

MARTINGALE SOLUTIONS FOR THE COMPRESSIBLE MHD SYSTEMS WITH STOCHASTIC EXTERNAL FORCES

HUAQIAO WANG

ABSTRACT. In this paper we consider the three-dimensional compressible MHD system with stochastic external forces in a bounded domain. We obtain the existence of martingale solution which is a weak solution for the fluid variables, the Brownian motion on a probability space. The construction of the solution is based on the Galerkin approximation method, stopping time, the compactness method and Jakubowski Skorokhod theorem, etc.

1. INTRODUCTION

In this paper, we consider the following three-dimensional stochastic compressible Magneto-hydrodynamic (MHD) system:

$$\begin{cases} d\rho + \operatorname{div}(\rho u)dt = 0, \\ d(\rho u) + [\operatorname{div}(\rho u \otimes u) + \nabla p - \mu\Delta u - (\lambda + \mu)\nabla \operatorname{div} u - (\nabla \times B) \times B]dt = \sum f_k(\rho, \rho u, x)d\beta_k^1(t), \\ dB - [\nabla \times (u \times B) + \nu\Delta B]dt = \sum g_k(B, x)d\beta_k^2(t), \quad \operatorname{div} B = 0, \end{cases} \quad (1.1)$$

where ρ is the density, u is the velocity field of the fluid, p is the scalar pressure and $p = a\rho^\gamma$ with a positive constant a and adiabatic index $\gamma \geq 1$, B is the magnetic field induced by the charged fluid, μ and λ are two viscosity constants satisfying $2\mu + 3\lambda \geq 0$, $\nu > 0$ is the magnetic diffusivity acting as a magnetic diffusion coefficient of the magnetic field, and all these kinetic coefficients and the magnetic diffusivity are independent of the magnitude and direction of the magnetic field, the symbol \otimes denotes the tensor product, $f_k(\rho, \rho u, x)d\beta_k^1$ and $g_k(B, x)d\beta_k^2$ are stochastic external forces, where $\beta_k^1, \beta_k^2, k = 1, 2, \dots$ are two sequences of independent one-dimensional \mathbb{R} -valued Brownian motions. Denote $\beta_k = (\beta_k^1, \beta_k^2)$.

System (1.1) models the interaction between a conductor fluid and the magnetic field in the presence of random perturbations. The inclusion of the stochastic terms in the governing equations is widely used to account for random fluctuations and past history of the system.

Notice that we have followed the usual convention of including the divergence-free condition on the magnetic field B . This is really a condition on the initial data, as this property is preserved by the dynamics of (1.1). To be more precise, we impose the assumptions on the initial data

$$\rho|_{t=0} = \rho_0, \quad \rho u|_{t=0} = m_0, \quad B|_{t=0} = B_0, \quad \operatorname{div} B_0 = 0, \quad (1.2)$$

and the boundary conditions

$$u|_{\partial D} = 0, \quad B|_{\partial D} = 0, \quad (1.3)$$

Date: January 15, 2022.

2010 Mathematics Subject Classification. 35Q35, 76D05, 76W05, 60H15, 76M35.

Key words and phrases. Stochastic compressible MHD systems, martingale solutions, stochastic compactness method.

where $D \subset \mathbb{R}^3$ is a bounded domain with smooth boundary ∂D , and \mathbf{n} is the unit outer normal at the boundary.

The deterministic version of system (1.1) was extensively studied in the literature. For instance, the existence of local strong solutions with large initial data has proved by Fan-Yu [11]; the existence of global weak and variational solution have obtained by Ducomet-Feireisl[10], Fan-Yu[12], Hu-Wang [17, 18]; the time decay of the compressible MHD systems with small initial data has established in [8, 25, 31, 36, 37]; the low Mach number limit problem was studied in [19, 21].

When $\rho = 1$, system (1.1) reduces to the stochastic incompressible MHD system, which has also received a lot of attention [1, 32, 35]. On the other hand, when $B = 0$ and ρ is not a constant, system (1.1) becomes the stochastic compressible Navier-Stokes equations, to which, to the best of our knowledge, much less is known. Feireisl-Maslowski-Novotný in [14] considered the three-dimensional compressible Navier-Stokes equations driven by the stochastic external forces in Sobolev spaces and obtained the existence of solutions by using the abstract measurability theorem ([4]) to prove that the weak solution generates a random variable. For the existence of martingale solutions to the three-dimensional compressible Navier-Stokes equations driven by the multiplicative noise, see [7, 34, 38]. Recently, Breit-Feireisl-Hofmanová considered the low Mach number limit problem in [6].

1.1. Definition of solutions and assumptions. We will seek a particular type of solution to system (1.1), namely the *martingale solution*. The definition of such solutions is given as follows.

Definition 1.1. A martingale solution of (1.1)-(1.3) is a system $((\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P}), \beta_k, \rho, u, B)$ with the following properties:

(1) $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$ is a stochastic basis, where \mathcal{F}_t is a filtration on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, i.e., a nondecreasing family $\{\mathcal{F}_t : t \geq 0\}$ of sub σ -fields of $\mathcal{F} : \mathcal{F}_s \subset \mathcal{F}_t \subset \mathcal{F}$ for $0 \leq s < t < \infty$;

(2) $\{\beta_k^1\}_{k \geq 1}, \{\beta_k^2\}_{k \geq 1}$ are two sequences of independent \mathbb{R} -valued Brownian motions;

(3) for any $\varphi, \psi \in C_0^\infty(D)$, $\langle \rho, \varphi \rangle, \langle \rho u, \psi \rangle$ and $\langle B, \psi \rangle$ are progressively measurable;

(4) for all $1 \leq p < \infty$, $\rho \geq 0$, $\rho \in L^p(\Omega, L^\infty(0, T; L^\gamma(D)))$, $u \in L^p(\Omega, L^2(0, T; H_0^1(D)))$, $B \in L^p(\Omega, L^\infty(0, T; L^2(D))) \cap L^p(\Omega, L^2(0, T; H_0^1(D)))$ such that for all $t \in [0, T]$, $\varphi \in C_0^\infty([0, T] \times D)$ and $\psi \in C_0^\infty([0, T] \times D)$, it holds that P-a.s.

$$\begin{aligned} & \int_D \rho(t) \varphi dx - \int_D \rho_0 \varphi dx = \int_0^t \int_D \rho u \cdot \nabla \varphi dx dt, \\ & \int_D (\rho u)(t) \cdot \psi dx - \int_0^t \int_D [\rho u u \cdot \nabla \psi - \mu \nabla u \cdot \nabla \psi - (\lambda + \mu) \operatorname{div} u \cdot \operatorname{div} \psi + a \rho^\gamma \cdot \operatorname{div} \psi] dx dt \\ & = \int_D \rho_0 u_0 \cdot \psi dx + \int_0^t \int_D (\nabla \times B) \times B \cdot \psi dx dt + \sum_{k=1}^{\infty} \int_0^t \int_D f_k(\rho, \rho u, x) \cdot \psi dx d\beta_k^1, \\ & \int_D B(t) \cdot \psi dx - \int_0^t \int_D [\nabla \times (u \times B) + \nu \Delta B] \cdot \psi dx dt = \int_D B_0 \cdot \psi dx + \sum_{k=1}^{\infty} \int_0^t \int_D g_k(B, x) \cdot \psi dx d\beta_k^2; \end{aligned}$$

(5) the equation (1.1)₁ is satisfied in sense of renormalized solutions, that is,

$$b(\rho)_t + \operatorname{div}(b(\rho)u) + (b'(\rho)\rho - b(\rho))\operatorname{div}u = 0 \quad (1.4)$$

holds in $\mathcal{D}'((0, T) \times \mathbb{R}^3)$ P-a.s. for any $b \in C^1(\mathbb{R})$ with $b'(z) \equiv 0$ for all $z \in \mathbb{R}$ large enough.

Notations. Throughout this paper, we drop the parameter $\omega \in \Omega$ for simplicity of notations. $\{\mathcal{F}_t\}$ is a right continuous filtration over the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that \mathcal{F}_0 contains all \mathbb{P} -negligible subsets of Ω . We use $\mathbb{E}X = \int_{\Omega} X d\mathbb{P}$ to denote the expectation of the random variable $X(\omega, t)$ for fixed t . All stochastic integrals are defined in the sense of Itô; see [9, 23]. Moreover, C denotes a generic constant which may vary in different estimates. For simplicity, we will write $A \lesssim B$ if $A \leq CB$.

We use $C([0, T]; V_w)$ to denote the subspace of $L^\infty([0, T]; V)$ consisting of those functions which are a.e. equal to weakly continuous functions with values in V . We denote $L_2(U, Y)$ the set of all Hilbert-Schmidt operators from U to Y .

Assumptions on the stochastic forces. We assume that the stochastic forces f_k, g_k satisfy the following conditions:

(A) $f_k(\rho, \rho u, x), g_k(B, x)$ are progressively measurable and C^1 continuous in $\rho, \rho u, x; B, x$, respectively, and

$$\left\{ \begin{array}{l} f_k(\rho, \rho u, x) = f_{k,1}(x, \rho) + f_{k,2}(x) \rho u, \\ \sum_{k=1}^{\infty} |f_{k,1}|^2 \leq C|\rho|^{\gamma+1}, \quad \sum_{k=1}^{\infty} |\partial_{\rho} f_{k,1}|^2 \leq C|\rho|^{\gamma-1}, \quad \sum_{k=1}^{\infty} |f_{k,2}|^2 \leq C, \\ \sum_{k=1}^{\infty} |g_k(B, x)|^2 \leq C|B|^2, \quad \sum_{k=1}^{\infty} |\partial_B g_k| \leq C. \end{array} \right. \quad (1.5)$$

The data ρ_0, m_0, B_0 are assumed to satisfy the following conditions:

$$\begin{aligned} \rho_0 &\in L^{\gamma p}(\Omega, L^{\gamma}(D)), \quad \rho_0 \geq 0; \quad m_0 = 0 \quad \text{if } \rho_0 = 0, \\ \frac{|m_0|^2}{\rho_0} &\in L^p(\Omega, L^1(D)); \quad B_0 \in L^{2p}(\Omega, L^2(D)), \quad \operatorname{div} B_0 = 0. \end{aligned} \quad (1.6)$$

Denote

$$F(\rho, \rho u) dW_1 := \sum_{k=1}^{\infty} f_k(\rho, \rho u, x) d\beta_k^1, \quad G(B) dW_2 := \sum_{k=1}^{\infty} g_k(B, x) d\beta_k^2.$$

By (1.5) and Hölder's inequality, we have for $\ell > \frac{3}{2}$,

$$\begin{aligned} \|F(\rho, \rho u)\|_{L_2(U, W^{-\ell, 2}(D))}^2 &= \sum_{k=1}^{\infty} \|f_k(\rho, \rho u, x)\|_{W^{-\ell, 2}(D)}^2 \lesssim \sum_{k=1}^{\infty} \|f_k(\rho, \rho u, x)\|_{L^1(D)}^2 \\ &\lesssim \left(\int_D \sqrt{\rho} \sum_{k=1}^{\infty} \frac{f_{k,1}(\rho, x)}{\sqrt{\rho}} + \sqrt{\rho} \sqrt{\rho u} \sum_{k=1}^{\infty} f_{k,2}(x) dx \right)^2 \\ &\lesssim \|\rho\|_{L^1} (\|\rho\|_{L^{\gamma}(D)}^{\gamma} + \|\sqrt{\rho u}\|_{L^2(D)}^2) \lesssim 1, \\ \|G(B)\|_{L_2(U, W^{-1, 2}(D))}^2 &= \sum_{k=1}^{\infty} \|g_k(B, x)\|_{W^{-1, 2}(D)}^2 \leq \sum_{k=1}^{\infty} \|g_k(B, x)\|_{L^2(D)}^2 \\ &\lesssim \|B\|_{L^2(D)}^2 \lesssim 1. \end{aligned}$$

Then the Itô's integrals $\int_0^T F(\rho, \rho u) dW_1$ and $\int_0^T G(B) dW_2$ are well-defined martingales in $W^{-\ell, 2}(D)$ and $W^{-1, 2}(D)$ respectively. Since W_1 and W_2 don't converge on U , we define $U_0 \supset U$ by $U_0 = \{v = \sum_{k \geq 1} \alpha_k e_k : \sum_{k \geq 1} \frac{\alpha_k^2}{k^2} < \infty\}$ with the norm $\|v\|_{U_0}^2 = \sum_{k \geq 1} \alpha_k^2 / k^2$,

$v = \sum_{k \geq 1} \alpha_k e_k$. Note that the embedding $U \hookrightarrow U_0$ is Hilbert-Schmidt and $\{e_k\}_{k \geq 1}$ is a complete orthonormal basis of U . Moreover, we have $W_1, W_2 \in C([0, T], U_0)$ \mathbb{P} -a.s..

1.2. Main results. Our main result is stated in the following theorem:

Theorem 1.2. Suppose that the initial data (ρ_0, m_0, B_0) satisfies (1.6) and (1.5) holds. Let $\gamma > 3/2$. If $D \subset \mathbb{R}^3$ is a bounded domain of class $C^{2+\alpha}$ with $\alpha > 0$, then there exists a martingale solution $((\Omega, \mathcal{F}, \mathbb{P}), \mathcal{F}_t, \beta_k, \rho, u, B)$ in the sense of Definition 1.1 with

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} \mathcal{E}(t) + \int_0^T \int_D (\mu |\nabla u|^2 + (\lambda + \mu) |\operatorname{div} u|^2 + \nu |\nabla B|^2) dx dt \right]^p \lesssim 1$$

to the problem (1.1)-(1.3) for any given $T > 0$ and for all $1 \leq p < \infty$.

The basic idea to prove Theorem 1.2 is to first construct the approximate schemes to the problem (1.1)-(1.3), and then establish tightness estimates of the approximate solutions and pass to the limits in the spirit of Feireisl [13, 15]. Of course, the apparent challenge stems from the stochastic term and the Itô integral, which is the primary focus of the work.

More precisely, as a first step we use Faedo-Galerkin method to construct the approximate system involving the artificial viscosity and artificial pressure, and apply the standard fixed point argument to obtain an n -dimensional truncated local solution $(\hat{\beta}_k, \rho_n, u_n, B_n)$ in the time interval $[0, \tau_{n,N})$. The existence of solutions in the whole time $[0, T]$ will be achieved by the energy estimates. Additional challenges arise due to the probabilistic nature of the system. Our method relies on a delicate stopping time argument with the use of the Burkholder-Davis-Gundy inequality. In particular, this approach leads to the convergence of the approximate solutions on $[0, T]$.

However the convergence is too weak to guarantee that the limit is a solution on $[0, T]$. In the two-dimensional case, it can be shown by using certain monotonicity principle that the nonlinear terms converge to the right limit and hence a global strong solution can be obtained [26]. But when the space dimension is three the monotonicity does not hold and to the best of our knowledge there is no result on the global strong solutions. This is why we pursue instead the Martingale solutions. As is explained, the main issue is the convergence of the nonlinear terms. To this end, we need to derive the tightness and show the convergence as $n \rightarrow \infty$ of the approximate finite-dimensional solutions in the Faedo-Galerkin method in three steps: (1) We apply the deterministic compactness criterion and Arzela-Ascoli's Theorem to get the tightness property of the approximate solution $(\hat{\beta}_k, \rho_n, u_n, B_n)$. Moreover in order to analyze the nonlinear term, we prove the tightness of $\rho_n u_n$ as well. Then from the Jakubowski-Skorokhod Theorem, there exist a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ and random variables $(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n) \rightarrow (\beta_k, \rho, u, h, B)$ $\tilde{\mathbb{P}}$ -a.s., (Denote $m_n = \rho_n u_n$, since they have same distribution, that is, for $A \in \mathcal{F}$, $\tilde{\mathbb{P}}(\tilde{m}_n \in A) = \tilde{\mathbb{P}}(\rho_n u_n \in A) = \tilde{\mathbb{P}}(\tilde{\rho}_n \tilde{u}_n \in A)$, we have $\tilde{m}_n = \tilde{\rho}_n \tilde{u}_n$ almost surely) with the property that the probability distribution of $(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n)$ is the same as that of $(\hat{\beta}_k, \rho_n, u_n, \rho_n u_n, B_n)$. By using a cut-off function we can also show that the random variable $(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n)$ satisfy the approximate equations in $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$; (2) In view of the uniformly integrable criterion and Vitali's convergence Theorem, together with the almost sure convergence on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, we can obtain the limit process of the stochastic parts, it's quadratic variation and cross variation are martingale; (3) Finally, by the new method in [3, 28] on the martingale problem of Stroock and Varadhan, we can prove that the system $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ is a martingale solution of the approximation scheme. After obtaining the foregoing solutions to the approximate

scheme by the Faedo-Galerkin method, we shall follow the idea of [15, 30] to deal with the artificial viscosity and artificial pressure using the similar convergence framework for $n \rightarrow \infty$ in the Faedo-Galerkin solutions. Due to insufficient integrability of the density, the effective viscous flux and the cut-off function technique will be used to obtain the strong convergence of the density. Then we pass to the limits of vanishing artificial viscosity and vanishing artificial pressure, and the limit of the solutions to the approximate scheme is a martingale solution with finite energy to the initial-boundary value problem of (1.1)-(1.3). Note that here we shall use the property of martingale.

The rest of the paper is organized as follows. We shall recall some basic theory of stochastic analysis in Section 2. In Section 3, we shall construct the solutions to an approximate scheme by the Faedo-Galerkin method. In Section 4, we shall pass to the limit as the artificial viscosity goes to zero, and in Section 5, we shall pass to the limit as the artificial pressure goes to zero.

2. PRELIMINARIES

In this section, we first introduce some function spaces. Let $D \subset \mathbb{R}^3$ be an open subset with smooth boundary ∂D . Let $L^p(D)$ ($1 \leq p < \infty$) denote the Banach space of Lebesgue measurable \mathbb{R}^3 valued integrable functions on the set D with the standard norm $\|u\|_{L^p(D)}$, and $L^\infty(D)$ the Banach space of Lebesgue measurable essentially bounded \mathbb{R}^3 valued functions defined on D with the standard norm $\|u\|_{L^\infty(D)}$. If $p = 2$, then $L^2(D)$ is a Hilbert space with the scalar product denoted by

$$\langle u, v \rangle_{L^2(D)} = \int_D u(x)v(x)dx, \quad \text{for } u, v \in L^2(D).$$

Let $H^1(D)$ stand for the Sobolev space of all $u \in L^2(D)$ for which there exist weak derivatives $\frac{\partial u}{\partial x_i} \in L^2(D)$, $i = 1, 2, 3$. It is a Hilbert space with the scalar product denoted by

$$\langle u, v \rangle_{H^1(D)} = \langle u, v \rangle_{L^2(D)} + \langle \nabla u, \nabla v \rangle_{L^2(D)}, \quad \text{for } u, v \in H^1(D).$$

For a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a Banach space X , denote by $L^p(\Omega, L^q(0, T; X))$ ($1 \leq p, q < \infty$) the space of random functions defined on Ω with value in $L^q(0, T; X)$, endowed with the norm:

$$\|u\|_{L^p(\Omega, L^q(0, T; X))} = (\mathbb{E}\|u\|_{L^q(0, T; X)}^p)^{\frac{1}{p}}.$$

If $q = \infty$, we write

$$\|u\|_{L^p(\Omega, L^\infty(0, T; L^q(X)))} = (\mathbb{E} \operatorname{ess\,sup}_{0 \leq t \leq T} \|u\|_{L^q(X)}^p)^{\frac{1}{p}}.$$

We now recall some preliminaries of stochastic analysis and useful tools for the sake of convenience and completeness. For details, we refer the reader to [9, 23] and the references therein.

Let us first recall the Itô formula for a local martingale.

Lemma 2.1. Suppose that $M_t = (M_t^1, M_t^2, \dots, M_t^n)$ is a vector valued continuous local martingale, that is (M_t^i, \mathcal{F}_t) is a local martingale for each $i = 1, 2, \dots, n$ and $t \in \mathbb{R}_+$. Let $A_t = (A_t^1, A_t^2, \dots, A_t^n)$ be a vector valued continuous processes adapted to the same filtration $\{\mathcal{F}_t\}$ such that the total variation of A_t^i on each finite interval is bounded almost surely, and

$A_0^i = 0$ almost surely. Let $X_t = (X_t^1, X_t^2, \dots, X_t^n)$ be a vector valued adapted processes such that $X_t = X_0 + M_t + A_t$, and let $\Phi \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^n)$. Then, for any $t \geq 0$, the equality

$$\begin{aligned} \Phi(t, X_t) &= \Phi(0, X_0) + \sum_{i=1}^n \int_0^t \frac{\partial}{\partial x_i} \Phi(s, X_s) dM_s^i + \sum_{i=1}^n \int_0^t \frac{\partial}{\partial x_i} \Phi(s, X_s) dA_s^i \\ &\quad + \frac{1}{2} \sum_{i,j=1}^n \int_0^t \frac{\partial^2}{\partial x_i \partial x_j} \Phi(s, X_s) d\langle M^i, M^j \rangle_s + \int_0^t \frac{\partial}{\partial s} \Phi(s, X_s) ds \end{aligned}$$

holds almost surely. Here $\langle \cdot, \cdot \rangle_t$ is the cross variation process defined by $\langle X, Y \rangle_t = \frac{1}{4} \{ \langle X + Y \rangle_t - \langle X - Y \rangle_t \}$, and $\langle X \rangle_t$ denotes the quadratic variation of X on $[0, t]$.

3. THE FAEDO-GALERKIN APPROXIMATION AND A PRIORI ESTIMATES

3.1. Approximation scheme and a priori estimates. For fixed $\varepsilon, \delta > 0$, we consider the following approximation problem with $\beta > \max\{4, \gamma\}$:

$$\begin{cases} d\rho + \operatorname{div}(\rho u) dt = \varepsilon \Delta \rho dt \\ d(\rho u) + [\operatorname{div}(\rho u \otimes u) + a \nabla \rho^\gamma + \delta \nabla \rho^\beta + \varepsilon \nabla u \cdot \nabla \rho - (\lambda + \mu) \nabla \operatorname{div} u - \mu \Delta u] dt \\ \quad = (\nabla \times B) \times B dt + \sum_{k \geq 1} f_k(\rho, \rho u, x) d\beta_k^1(t) \\ dB - [\nabla \times (u \times B) + \nu \Delta B] dt = \sum_{k \geq 1} g_k(B, x) d\beta_k^2(t), \end{cases} \quad (3.1)$$

with the boundary conditions:

$$\nabla \rho \cdot \mathbf{n}|_{\partial D} = 0, \quad u|_{\partial D} = 0, \quad B|_{\partial D} = 0 \quad (3.2)$$

and the initial data:

$$\begin{aligned} \rho|_{t=0} &= \rho_{0,\delta} \in C^{2+\alpha}(\bar{D}), \quad \nabla \rho_{0,\delta} \cdot \mathbf{n}|_{\partial D} = 0, \\ (\rho u)|_{t=0} &= m_{0,\delta} \in C^2(\bar{D}), \quad B|_{t=0} = B_{0,\delta} \in C^2(\bar{D}), \quad \operatorname{div} B_{0,\delta} = 0. \end{aligned} \quad (3.3)$$

Here we take the initial data that satisfies the following conditions:

$$\begin{aligned} \rho_{0,\delta} &\rightarrow \rho_0 \text{ in } L^\gamma(D), \quad B_{0,\delta} \rightarrow B_0 \text{ in } L^2(D) \text{ as } \delta \rightarrow 0; \\ 0 < \delta &\leq \rho_{0,\delta}(x) \leq \delta^{-\frac{1}{\beta}}, \quad m_{0,\delta} = h_\delta \sqrt{\rho_{0,\delta}}, \end{aligned}$$

almost surely and h_δ is defined as follows. First, take

$$\tilde{m}_{0,\delta}(x) = \begin{cases} m_0(x) \sqrt{\frac{\rho_{0,\delta}(x)}{\rho_0(x)}}, & \text{if } \rho_0(x) > 0, \\ 0, & \text{if } \rho_0(x) = 0. \end{cases}$$

By using (1.6), we have

$$\frac{|\tilde{m}_{0,\delta}|^2}{\rho_{0,\delta}} \text{ is bounded in } L^p(\Omega, L^1(D)) \text{ independently of } \delta > 0, \forall p \in [1, \infty).$$

Since $C^2(\bar{D})$ is dense in $L^2(D)$, then we can find $h_\delta \in C^2(\bar{D})$ such that

$$\left\| \frac{\tilde{m}_{0,\delta}}{\sqrt{\rho_{0,\delta}}} - h_\delta \right\|_{L^p(\Omega, L^2(D))} < \delta.$$

On the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with two given Brownian motions $\hat{\beta}_k^1$ and $\hat{\beta}_k^2$, $k \geq 1$, denote $\hat{\beta}_k = (\hat{\beta}_k^1, \hat{\beta}_k^2)_{k \geq 1}$.

In order to solve (3.1)-(3.3), we first consider a suitable orthogonal system formed by a family of smooth functions φ_n vanishing on ∂D . One can take the eigenfunctions of the Dirichlet problem for the Laplacian operator:

$$-\Delta\varphi_n = \lambda_n\varphi_n \text{ on } D, \quad \varphi_n|_{\partial D} = 0.$$

Now, we consider a sequence of finite dimensional spaces

$$X_n = \text{span}\{\varphi_j\}_{j=1}^n, \quad n = 1, 2, \dots$$

First, we introduce a family of operators

$$\mathcal{M}[\rho] : X_n \rightarrow X_n; \quad \langle \mathcal{M}[\rho]v, w \rangle = \int_D \rho v \cdot w dx, \quad \forall v, w \in X_n.$$

If $\inf_{x \in D} \rho > 0$, then these operators are invertible and we have

$$\|\mathcal{M}^{-1}[\rho]\|_{\mathcal{L}(X_n, X_n)} \leq \left(\inf_{x \in D} \rho(x) \right)^{-1},$$

where $\mathcal{M}^{-1}[\rho]$ is the inverse of $\mathcal{M}[\rho]$. Moreover, by using the following identity

$$\mathcal{M}^{-1}[\rho_1] - \mathcal{M}^{-1}[\rho_2] = \mathcal{M}^{-1}[\rho_2](\mathcal{M}[\rho_2] - \mathcal{M}[\rho_1])\mathcal{M}^{-1}[\rho_1],$$

we can deduce that the map

$$\rho \mapsto \mathcal{M}^{-1}[\rho] \text{ mapping } L^1(D) \text{ into } \mathcal{L}(X_n, X_n)$$

is well-defined and satisfies

$$\|\mathcal{M}^{-1}[\rho_1] - \mathcal{M}^{-1}[\rho_2]\|_{\mathcal{L}(X_n, X_n)} \leq C(n, \eta) \|\rho_1 - \rho_2\|_{L^1(D)} \quad (3.4)$$

for any ρ_1, ρ_2 belonging to the set

$$N_\eta = \left\{ \rho \in L^1 : \inf_{x \in D} \rho \geq \eta > 0 \right\}.$$

Now we will try to solve (3.1) on X_n . For this, let \mathbb{P} be the projection from $L^2(D)$ to X_n . \mathbb{P} is also a linear projection from $H^\ell(D)$ to X_n . In fact,

$$\mathbb{P}h = \sum_{i=1}^n \langle h, e_i \rangle_{H^\ell - H^{-\ell}} e_i, \quad h \in H^\ell(D), e_i \in X_n.$$

We shall look for the sequence of pairs $(u_n, B_n) \in C([0, T]; X_n)$ satisfying the integral equations:

$$\begin{aligned} & \int_D \rho(t) u_n(t) \cdot \varphi dx + \int_0^t \int_D [\text{div}(\rho u_n \otimes u_n) - \mu \Delta u_n - (\nabla \times B_n) \times B_n] \cdot \varphi dx ds \\ &= \int_D m_0 \cdot \varphi dx + \int_0^t \int_D \left[\nabla \left((\lambda + \mu) \text{div} u_n - a \rho^\gamma - \delta \rho^\beta \right) - \varepsilon \nabla u_n \cdot \nabla \rho \right] \cdot \varphi dx ds \\ & \quad + \int_0^t \left\langle \sum_{k \geq 1} f_k^n(\rho, \rho u_n, x), \varphi \right\rangle d\hat{\beta}_k^1, \\ & \int_D B_n(t) \cdot \varphi dx - \int_D B_0 \cdot \varphi dx = \int_0^t \int_D [\nabla \times (u_n \times B_n) + \nu \Delta B_n] \cdot \varphi dx ds \\ & \quad + \int_0^t \int_D \sum_{k \geq 1} g_k(B_n, x) \cdot \varphi dx d\hat{\beta}_k^2, \end{aligned}$$

for all $t \in [0, T]$ and any function $\varphi \in X_n$, where

$$\sum_{k \geq 1} f_k^n(\rho, \rho u_n, x) := \mathcal{M}^{\frac{1}{2}}[\rho] \mathbb{P} \left(\sum_{k \geq 1} f_k(\rho, \rho u_n, x) / \sqrt{\rho} \right), \quad \text{with } \langle \mathcal{M}^{\frac{1}{2}}[\rho] v, w \rangle = \int_D \sqrt{\rho} v \cdot w dx.$$

Then, we can rewrite the above integral equations as

$$(u_n(t), B_n(t)) = \left(\mathcal{M}^{-1}[\rho(t)] \left[m_0^* + \int_0^t \mathcal{N}_1[\rho(s), u_n(s), B_n(s)] ds + \int_0^t \sum_{k \geq 1} f_k^n(\rho, \rho u_n, x) d\hat{\beta}_k^1 \right], \right. \\ \left. B_0^* + \int_0^t \mathcal{N}_2[u_n(s), B_n(s)] ds + \int_0^t \sum_{k \geq 1} g_k(B_n, x) d\hat{\beta}_k^2 \right), \quad (3.5)$$

where

$$\begin{aligned} \langle m_0^*, \varphi \rangle &= \int_D m_0 \cdot \varphi dx, & \langle B_0^*, \varphi \rangle &= \int_D B_0 \cdot \varphi dx, \\ \langle \mathcal{N}_1[\rho, u_n, B_n], \varphi \rangle &= \int_D [\mu \Delta u_n - \operatorname{div}(\rho u_n \otimes u_n) \\ &\quad + \nabla \left((\lambda + \mu) \operatorname{div} u_n - a \rho^\gamma - \delta \rho^\beta \right) - \varepsilon \nabla u_n \cdot \nabla \rho + (\nabla \times B_n) \times B_n] \cdot \varphi dx, \\ \langle \mathcal{N}_2[u_n, B_n], \varphi \rangle &= \int_D [\nabla \times (u_n \times B_n) + \nu \Delta B_n] \cdot \varphi dx. \end{aligned}$$

We will take the following steps to solve (3.5).

Step 1: Solve ρ in terms of u_n and derive a fixed-point problem. Note that ρ is determined as the solution of the following Neumann initial-boundary value problem (for e.g., see Lemma 2.2 [13]):

$$\begin{cases} \rho_t + \operatorname{div}(\rho u) = \varepsilon \Delta \rho, \\ \nabla \rho \cdot \mathbf{n}|_{\partial D} = 0, \\ \rho|_{t=0} = \rho_{0,\delta}(x). \end{cases} \quad (3.6)$$

From [15, Lemma 2.1, Lemma 2.2] we know that there exists a mapping $\mathcal{S} : C([0, T]; C^2(\overline{D})) \rightarrow C([0, T]; C^{2+\alpha}(\overline{D}))$ such that

- (1) $\rho = \mathcal{S}[u]$ is the unique classical solution of (3.6);
- (2) $\underline{\rho} \exp \left(- \int_0^t \|\operatorname{div} u(s)\|_{L^\infty(D)} ds \right) \leq \mathcal{S}[u](t, x) \leq \overline{\rho} \exp \left(- \int_0^t \|\operatorname{div} u(s)\|_{L^\infty(D)} ds \right)$ for all $t \in [0, T]$, where $0 < \underline{\rho} \leq \rho_{0,\delta} \leq \overline{\rho}$;
- (3) $\|\mathcal{S}[u_1] - \mathcal{S}[u_2]\|_{C([0,T]; W^{1,2}(D))} \leq TC(K, T) \|u_1 - u_2\|_{C([0,T]; W^{1,2}(D))}$ for any u_1, u_2 in $M_K := \{u \in C([0, T]; W^{1,2}(D)) : \|u(t)\|_{L^\infty(D)} + \|\nabla u(t)\|_{L^\infty(D)} \leq K \text{ for all } t\}$.

Thus on X_n we can write $\rho = \mathcal{S}[u_n]$, and construct the approximate solutions for (3.1)-(3.3) by means of (3.5). That is, we can get the approximate solutions by solving the following integral equations:

$$(u_n(t), B_n(t)) = \left(\mathcal{M}^{-1}[\mathcal{S}[u_n](t)] \left[m_0^* + \int_0^t \mathcal{N}_1[\mathcal{S}[u_n](s), u_n(s), B_n(s)] ds + \int_0^t \sum_{k \geq 1} f_k^n(\mathcal{S}[u_n], \mathcal{S}[u_n] u_n, x) d\hat{\beta}_k^1(s) \right], \right.$$

$$B_0^* + \int_0^t \mathcal{N}_2[u_n(s), B_n(s)] ds + \int_0^t \sum_{k \geq 1} g_k(B_n, x) d\hat{\beta}_k^2(s) \Big).$$

Step 2. Solve a cut-off problem. For $N > 0$, choose a C^∞ smooth cut-off function $\theta_N : [0, \infty) \rightarrow [0, 1]$ such that

$$\theta_N(x) := \begin{cases} 1, & \text{for } |x| \leq N, \\ 0, & \text{for } |x| \geq N + 1. \end{cases}$$

Note that we consider the following cut-off problem of (3.5) for a fixed n :

$$\begin{aligned} (u_n^N(t), B_n^N(t)) = & \left(\mathcal{M}^{-1}[\mathcal{S}[u_n^N](t)] \left[m_0^* + \int_0^t \theta_N^{u_n^N, B_n^N}(s) \mathcal{N}_1[\mathcal{S}[u_n^N](s), u_n^N(s), B_n^N(s)] ds \right. \right. \\ & \left. \left. + \int_0^t \sum_{k \geq 1} \theta_N^{u_n^N, B_n^N}(s) f_k^n(\mathcal{S}[u_n^N], \mathcal{S}[u_n^N]u_n^N, x) d\hat{\beta}_k^1(s) \right], \right. \\ & \left. B_0^* + \int_0^t \theta_N^{u_n^N, B_n^N}(s) \mathcal{N}_2[u_n^N(s), B_n^N(s)] ds + \int_0^t \sum_{k \geq 1} \theta_N^{u_n^N, B_n^N}(s) g_k(B_n^N, x) d\hat{\beta}_k^2(s) \right), \end{aligned} \quad (3.7)$$

where $\theta_N^{u_n^N, B_n^N}(s) = \theta_N(\max\{\|u_n^N(s)\|_{W^{1,\infty}}, \|B_n^N(s)\|_{W^{1,\infty}}\})$. Let N and n be fixed. By means of the standard fixed point argument on the Banach space $C([0, T]; X_n)$, we can solve the integral equations (3.7), at least on a short time interval $[0, T_{n,N}]$, $T_{n,N} \leq T$. For simplicity, we denote $(u^N, B^N) := (u_n^N, B_n^N)$.

Proposition 3.1. Given a $T > 0$, for each fixed n and N , there exists a $T_{n,N} \in (0, T]$ such that the equation (3.7) admits a unique solution $(u^N, B^N) \in L^2(\Omega, C([0, T_{n,N}]; X_n))^2$.

Proof. Define

$$B_{N, T_{n,N}} = \left\{ U^N := (u^N, h^N) \in L^2(\Omega, C(I_{n,N}; X_n))^2 : \|(u^N, h^N)\|_{C(I_{n,N}; W^{1,\infty}(D))} \leq N \right\},$$

with the norm $\|U^N\|_{B_{N, T_{n,N}}}^2 = \mathbb{E} \sup_{I_{n,N}} \|U^N\|_{X_n}^2$, where $I_{n,N} = [0, T_{n,N}]$ and $\|U^N\|_{X_n}^2 = \|u^N\|_{X_n}^2 + \|h^N\|_{X_n}^2$. Introduce a map

$$\mathcal{T} : L^2(\Omega, C(I_{n,N}; X_n))^2 \rightarrow L^2(\Omega, C(I_{n,N}; X_n))^2,$$

defined by

$$\mathcal{T}(U^N) := q_0^* + \int_0^t \theta_N^{U^N}(s) \mathcal{N} ds + \int_0^t \sum_{k \geq 1} \theta_N^{U^N}(s) F_k d\hat{\beta}_k(s),$$

where

$$\begin{aligned} q_0^* &= (\mathcal{M}^{-1}[\mathcal{S}(u^N)(t)] m_0^*, B_0^*), \\ \mathcal{N} &= (\mathcal{M}^{-1}[\mathcal{S}(u^N)(t)] \mathcal{N}_1[\mathcal{S}(u^N), u^N, h^N], \mathcal{N}_2[u^N, h^N]), \\ F_k &= (\mathcal{M}^{-1}[\mathcal{S}(u^N)(t)] f_k^n(\mathcal{S}(u^N), \mathcal{S}(u^N)u^N, x), g_k(h^N, x)). \end{aligned}$$

Similar to [27], by using the properties of $\mathcal{M}[\rho]$, $\mathcal{S}[u]$, Hölder's inequality and the Burkholder-Davis-Gundy inequality, we can infer that \mathcal{T} maps $B_{N, T_{n,N}}$ into itself for some suitable $T_{n,N}$.

We will prove that \mathcal{T} is a contraction on $L^2(\Omega, C(I_{n,N}; X_n))^2$ for some (small) $T_{n,N} > 0$. Since the deterministic parts are similar to [27, Section 7], we only need to consider the stochastic parts

$$\mathcal{T}^s(U^N) := \int_0^t \sum_{k \geq 1} \theta_N^{U^N}(s) F_k d\hat{\beta}_k(s).$$

For $U = (u, h)$, $V = (v, \ell)$, one can write

$$\begin{aligned} \mathcal{T}^s(U) - \mathcal{T}^s(V) &= (\mathcal{M}^{-1}[\mathcal{S}(u)(t)] - \mathcal{M}^{-1}[\mathcal{S}(v)(t)]) \int_0^t \sum_{k \geq 1} \theta_N^U(s) f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) d\hat{\beta}_k^1 \\ &\quad + \mathcal{M}^{-1}[\mathcal{S}(v)(t)] \int_0^t \sum_{k \geq 1} (\theta_N^U(s) f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) - \theta_N^V(s) f_k^n(\mathcal{S}(v), \mathcal{S}(v)v, x)) d\hat{\beta}_k^1 \\ &\quad + \int_0^t \sum_{k \geq 1} (\theta_N^U(s) g_k(h, x) - \theta_N^V(s) g_k(\ell, x)) d\hat{\beta}_k^2. \end{aligned}$$

Then

$$\begin{aligned} &\mathbb{E} \|\mathcal{T}^s(U) - \mathcal{T}^s(V)\|_{X_n}^2 \\ &\leq \mathbb{E} \sup_{0 \leq t \leq T_{n,N}} \left\| \int_0^t (\mathcal{M}^{-1}[\mathcal{S}(u)(t)] - \mathcal{M}^{-1}[\mathcal{S}(v)(t)]) \sum_{k \geq 1} f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) d\hat{\beta}_k^1 \right\|_{X_n}^2 \\ &\quad + \mathbb{E} \sup_{0 \leq t \leq T_{n,N}} \left\| \int_0^t \mathcal{M}^{-1}[\mathcal{S}(v)(t)] \sum_{k \geq 1} [f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) - f_k^n(\mathcal{S}(v), \mathcal{S}(v)v, x)] d\hat{\beta}_k^1 \right\|_{X_n}^2 \\ &\quad + \mathbb{E} \sup_{0 \leq t \leq T_{n,N}} \left\| \int_0^t \sum_{k \geq 1} (g_k(h, x) - g_k(\ell, x)) d\hat{\beta}_k^2 \right\|_{X_n}^2 \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

For the term I_1 , using the Burkholder-Davis-Gundy inequality, (1.5), (3.4) and the fact that $\mathcal{S}(u)$ is a contraction map, we have

$$\begin{aligned} I_1 &\leq \mathbb{E} \left(\int_0^{T_{n,N}} \left\| (\mathcal{M}^{-1}[\mathcal{S}(u)(t)] - \mathcal{M}^{-1}[\mathcal{S}(v)(t)]) \sum_{k \geq 1} f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) \right\|_{X_n}^2 ds \right) \\ &\leq \mathbb{E} \left(\|\mathcal{M}^{-1}[\mathcal{S}(u)(t)] - \mathcal{M}^{-1}[\mathcal{S}(v)(t)]\|_{\mathcal{L}(X_n, X_n)}^2 \int_0^{T_{n,N}} \left\| \sum_{k \geq 1} f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) \right\|_{X_n}^2 ds \right) \\ &\leq \mathbb{E} \left(\|\mathcal{S}(u) - \mathcal{S}(v)\|_{L^1(D)}^2 \int_0^{T_{n,N}} \left\| \sum_{k \geq 1} f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) \right\|_{X_n}^2 ds \right) \\ &\leq CT_{n,N} \mathbb{E} (\|u - v\|_{X_n}^2) \leq CT_{n,N} \mathbb{E} (\|U - V\|_{X_n}^2). \end{aligned}$$

Similarly, for the terms I_2 and I_3 , by (1.5), we obtain

$$I_2 \leq \mathbb{E} \left(\int_0^{T_{n,N}} \|\mathcal{M}^{-1}[\mathcal{S}(u)(t)]\|_{\mathcal{L}(X_n, X_n)}^2 \left\| \sum_{k \geq 1} (f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) - f_k^n(\mathcal{S}(v), \mathcal{S}(v)v, x)) \right\|_{X_n}^2 ds \right)$$

$$\begin{aligned}
&\leq \mathbb{E} \left([\inf \mathcal{S}(u)(t)]^{-1} \int_0^{T_{n,N}} \left\| \sum_{k \geq 1} (f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) - f_k^n(\mathcal{S}(v), \mathcal{S}(v)v, x)) \right\|_{X_n}^2 ds \right) \\
&\leq [\inf \mathcal{S}(u)(t)]^{-1} \mathbb{E} \left(\int_0^{T_{n,N}} \left\| \sum_{k \geq 1} (f_k^n(\mathcal{S}(u), \mathcal{S}(u)u, x) - f_k^n(\mathcal{S}(v), \mathcal{S}(v)v, x)) \right\|_{X_n}^2 ds \right) \\
&\leq CT_{n,N} \mathbb{E} (\|\mathcal{S}(u) - \mathcal{S}(v)\|_{X_n}^2 + \|u - v\|_{X_n}^2) \\
&\leq CT_{n,N} \mathbb{E} (\|u - v\|_{X_n}^2 + \|h - \ell\|_{X_n}^2) \leq CT_{n,N} \mathbb{E} (\|U - V\|_{X_n}^2),
\end{aligned}$$

and

$$I_3 \leq \mathbb{E} \int_0^{T_{n,N}} \left\| \sum_{k \geq 1} (g_k(h, x) - g_k(\ell, x)) \right\|_{X_n}^2 dt \leq CT_{n,N} \mathbb{E} (\|U - V\|_{X_n}^2).$$

Here we have used the fact that $\|\mathcal{M}^{\frac{1}{2}}[\mathcal{S}(u)] - \mathcal{M}^{\frac{1}{2}}[\mathcal{S}(v)]\|_{\mathcal{L}(X_n, X_n)} \leq C\|\mathcal{S}(u) - \mathcal{S}(v)\|_{L^2}$. This can be proved by using the definition of \mathcal{M} and the mean value theorem. Thus, for sufficiently small $T_{n,N}$ such that $\kappa = CT_{n,N} < 1$,

$$\mathbb{E} \|\mathcal{T}(U) - \mathcal{T}(V)\|_{X_n}^2 \leq \kappa \mathbb{E} (\|U - V\|_{X_n}^2).$$

Then \mathcal{T} is a contraction. Similar arguments indicate that with a further refined choice of $T_{n,N}$, \mathcal{T} maps $B_{N, T_{n,N}}$ into itself. Therefore \mathcal{T} has a unique fixed point (u^N, B^N) , proving the proposition. \square

Remark 3.2. The cut-off function $\theta_N(x)$ is introduced here to guarantee the global Lipschitz continuity of the map.

Step 3. Derive uniform a priori estimates to extend $\tau_{n,N}$ to T . Set $U_n^N = (u_n^N, B_n^N)$ be the solution to equation (3.5) on $[0, \tau_{n,N})$ obtained from Proposition 3.1. Let us introduce the following stopping times:

$$\tau_{n,N} = \begin{cases} \inf \{t \geq 0 : \|U_n^N(t)\|_{L^2(D)} \geq N\} \wedge \inf \left\{ t \geq 0 : \left\| \int_0^t F_n^N dW \right\|_{L^2(D)} ds \geq N \right\}, \\ T, \text{ otherwise.} \end{cases}$$

We denote $(u_N, B_N) := (u_n^N, B_n^N)$. Let $\rho_N := \mathcal{S}(u^N)$.

Our next goal is to prove

Lemma 3.3. For any fixed n ,

$$\lim_{N \rightarrow \infty} \mathbb{P}(\tau_{n,N} = T) = 1. \quad (3.8)$$

To prove the above lemma, we need the following energy estimates.

Proposition 3.4. For any (u_N, B_N) solving equation (3.5), for $1 \leq p < \infty$, we have

$$\begin{aligned}
&\mathbb{E} \left[\sup_{0 \leq t \leq T} \varepsilon_\delta + \mathbb{E} \int_0^T \left(\mu \|\nabla u_N(t)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(t)\|_{L^2(D)}^2 + \nu \|\nabla B_N(t)\|_{L^2(D)}^2 \right) dt \right. \\
&\quad \left. + \varepsilon \mathbb{E} \int_0^T \int_D \left(a\gamma \rho_N^{\gamma-2} + \delta \beta \rho_N^{\beta-2} \right) |\nabla \rho_N(t)|^2 dx dt \right]^p \leq C \mathbb{E} (\varepsilon_{0,\delta})^p.
\end{aligned} \quad (3.9)$$

Proof. Define the function $\Phi(\rho, m) := \int_D \frac{|m|^2}{\rho} dx$. Note that

$$\nabla_m \Phi(\rho, m) = \int_D \frac{2m}{\rho} dx, \quad \nabla_m^2 \Phi(\rho, m) = \int_D \frac{2}{\rho} \mathbb{I} dx, \quad \partial_\rho \Phi(\rho, m) = - \int_D \frac{|m|^2}{\rho^2} dx.$$

Here \mathbb{I} is the identity matrix.

Apply Itô's formula in Lemma 2.1 to the above function Φ with $(\rho, m) = (\rho_n, \rho_n u_N)$, and then apply Itô's formula again with $\Phi(B) := \int_D |B|^2 dx$. From equation (3.1), one deduces that

$$\begin{aligned} & d \int_D \left(\frac{1}{2} |\sqrt{\rho_N} u_N|^2 + \frac{a}{\gamma-1} \rho_N^\gamma + \frac{\delta}{\beta-1} \rho_N^\beta + \frac{1}{2} |B_N|^2 \right) dx + \int_D (\mu |\nabla u_N|^2 + (\lambda + \mu) |\operatorname{div} u_N|^2) dx ds \\ & \quad + \int_D \nu |\nabla B_N|^2 dx ds + \varepsilon \int_D (a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2}) |\nabla \rho_N|^2 dx ds \\ & = \int_D \sum_{k=1}^n f_k^n(\rho_N, \rho_N u_N, x) \cdot u_N dx d\hat{\beta}_k^1 + \frac{1}{2} \int_D \sum_{k \geq 1} |f_k(\rho_N, \rho_N u_N, x) / \sqrt{\rho_N}|^2 dx ds \\ & \quad + \int_D \sum_{k \geq 1} g_k(B_N, x) \cdot B_N dx d\hat{\beta}_k^2 + \frac{1}{2} \int_D \sum_{k \geq 1} |g_k(B_N, x)|^2 dx ds, \end{aligned} \quad (3.10)$$

where $s \in [0, t \wedge \tau_{n,N}]$, $t \wedge \tau_{n,N} = \min\{t, \tau_{n,N}\}$ and $t \in [0, T]$. Let

$$\mathcal{E}_\delta = \int_D \left(\frac{1}{2} \rho_N |u_N|^2 + \frac{a}{\gamma-1} \rho_N^\gamma + \frac{\delta}{\beta-1} \rho_N^\beta + \frac{1}{2} |B_N|^2 \right) dx,$$

$$\mathcal{E}_{0,\delta} = \int_D \left(\frac{1}{2} \rho_{0,\delta} |u_{0,\delta}|^2 + \frac{a}{\gamma-1} \rho_{0,\delta}^\gamma + \frac{\delta}{\beta-1} \rho_{0,\delta}^\beta + \frac{1}{2} |B_{0,\delta}|^2 \right) dx,$$

where $m_N = \rho_N u_N$. Integrating (3.10) on $[0, s]$ for all $s \in [0, t \wedge \tau_{n,N}]$ yields

$$\begin{aligned} & \mathcal{E}_\delta + \int_0^s \left[\mu \|\nabla u_N(r)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(r)\|_{L^2(D)}^2 + \nu \|\nabla B_N(r)\|_{L^2(D)}^2 \right] dr \\ & \quad + \varepsilon \int_0^s \int_D (a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2}) |\nabla \rho_N|^2 dx dr \\ & \leq \mathcal{E}_{0,\delta} + \frac{1}{2} \int_0^s \sum_{k \geq 1} \left\| \frac{f_k^n(\rho_N, \rho_N u_N, x)}{\sqrt{\rho_N(r)}} \right\|_{L^2(D)}^2 dr + \left| \int_0^s \langle u_N(r), \sum_{k \geq 1} f_k^n(\rho_N, \rho_N u_N, x) \rangle d\hat{\beta}_k^1 \right| \\ & \quad + \frac{1}{2} \int_0^s \sum_{k \geq 1} \|g_k(B_N, x)\|_{L^2(D)}^2 dr + \left| \int_0^s \langle B_N(r), \sum_{k \geq 1} g_k(B_N, x) \rangle d\hat{\beta}_k^2 \right|, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ is the inner product in $L^2(D)$. Taking the sup over $[0, t \wedge \tau_{n,N}]$ and the mathematical expectation in the above inequality, one easily deduces that

$$\begin{aligned} & \mathbf{E} \sup_{0 \leq s \leq t \wedge \tau_{n,N}} \mathcal{E}_\delta + \mathbf{E} \int_0^{t \wedge \tau_{n,N}} \left(\mu \|\nabla u_N(s)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(s)\|_{L^2(D)}^2 + \nu \|\nabla B_N(s)\|_{L^2(D)}^2 \right) ds \\ & \quad + \varepsilon \mathbf{E} \int_0^{t \wedge \tau_{n,N}} \int_D (a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2}) |\nabla \rho_N(s)|^2 dx ds \\ & \leq \mathbf{E} \mathcal{E}_{0,\delta} + \frac{1}{2} \mathbf{E} \int_0^{t \wedge \tau_{n,N}} \sum_{k \geq 1} \left\| \frac{f_k^n(\rho_N, \rho_N u_N, x)}{\sqrt{\rho_N(s)}} \right\|_{L^2(D)}^2 ds + \mathbf{E} \sup_s \left| \int_0^s \langle u_N, \sum_{k \geq 1} f_k^n(\rho_N, \rho_N u_N, x) \rangle d\hat{\beta}_k^1 \right| \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \sum_{k \geq 1} \|g_k(B_N, x)\|_{L^2(D)}^2 + \mathbb{E} \sup_s \left| \int_0^s \langle B_N, \sum_{k \geq 1} g_k(B_N, x) \rangle d\hat{\beta}_k^2 \right| \\
& =: \mathbb{E} \mathcal{E}_{0,\delta} + I_1 + I_2 + I_3 + I_4.
\end{aligned} \tag{3.11}$$

We now estimate the terms $I_i, i = 1, 2, 3, 4$. For the first term I_1 , the assumptions (1.5) on f and the Cauchy-Schwarz inequality imply that

$$\begin{aligned}
I_1 & \lesssim \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \int_D (|\sqrt{\rho_N} u_N|^2 + |\rho_N|^\gamma) dx ds \\
& \lesssim \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \left(\|\sqrt{\rho_N} u_N\|_{L^2(D)}^2 + \|\rho_N\|_{L^\gamma(D)}^\gamma \right) ds,
\end{aligned}$$

For I_2 , by the Burkholder-Davis-Gundy, Hölder and Young inequalities, and (1.5), for small $\eta > 0$, we have

$$\begin{aligned}
I_2 & \lesssim \mathbb{E} \left[\int_0^{t \wedge \tau_{n,N}} \sum_{k \geq 1} \langle f_k^n(\rho_N, \rho_N u_N, x), u_N \rangle^2 ds \right]^{\frac{1}{2}} \\
& \lesssim \mathbb{E} \left[\int_0^{t \wedge \tau_{n,N}} \|\sqrt{\rho_N} u_N\|_{L^2(D)}^2 \left(\|\sqrt{\rho_N} u_N\|_{L^2(D)}^2 + \|\rho_N\|_{L^\gamma(D)}^\gamma \right) ds \right]^{\frac{1}{2}} \\
& \leq \eta \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_{n,N}} \|\sqrt{\rho_N} u_N(s)\|_{L^2(D)}^2 + C_\eta \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \int_D (|\sqrt{\rho_N} u_N|^2 + |\rho_N|^\gamma) dx ds.
\end{aligned}$$

Similarly, from (1.5), one has

$$I_3 \leq C \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \|B_N\|_{L^2(D)}^2 ds,$$

and

$$I_4 \leq \eta \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_{n,N}} \|B_N(s)\|_{L^2(D)}^2 + C_\eta \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \|B_N\|_{L^2(D)}^2 ds.$$

When $p > 2$, we obtain as I_2 that

$$\begin{aligned}
|I_2|^p & \leq C \mathbb{E} \left[\int_0^{t \wedge \tau_{n,N}} \sum_{k \geq 1} \langle f_k(\rho_N, \rho_N u_N, x), u_N \rangle^2 ds \right]^{\frac{p}{2}} \\
& \leq \eta \mathbb{E} \left(\sup_{0 \leq s \leq t \wedge \tau_{n,N}} \|\sqrt{\rho_N} u_N(s)\|_{L^2(D)}^{2p} \right) + C_\eta \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \left(\int_D |\sqrt{\rho_N} u_N|^2 + |\rho_N|^\gamma dx \right)^p ds,
\end{aligned}$$

and

$$\begin{aligned}
|I_4|^p & \leq C \mathbb{E} \left[\int_0^{t \wedge \tau_{n,N}} \sum_{k \geq 1} \langle g_k(B_N, x), B_N \rangle^2 ds \right]^{\frac{p}{2}} \\
& \leq \eta \mathbb{E} \left(\sup_{0 \leq s \leq t \wedge \tau_{n,N}} \|B_N(s)\|_{L^2(D)}^{2p} \right) + C_\eta \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \left(\int_D |B_N|^2 dx \right)^p ds.
\end{aligned}$$

Plugging the estimates on $I_1 - I_4$ into (3.11), one has, for small enough $\eta > 0$,

$$\begin{aligned} & \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_{n,N}} \mathcal{E}_\delta + \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \mu \|\nabla u_N(s)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(s)\|_{L^2(D)}^2 + \nu \|\nabla B_N(s)\|_{L^2(D)}^2 ds \\ & \quad + \varepsilon \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \int_D \left(a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2} \right) |\nabla \rho_N(s)|^2 dx ds \\ & \leq \mathbb{E} \mathcal{E}_{0,\delta} + C_\eta \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \int_D (|\sqrt{\rho_N} u_N|^2 + |B_N|^2) dx ds. \end{aligned}$$

Then by Gronwall's inequality, we have

$$\begin{aligned} & \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_{n,N}} \mathcal{E}_\delta + \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \left(\mu \|\nabla u_N(s)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(s)\|_{L^2(D)}^2 + \nu \|\nabla B_N(s)\|_{L^2(D)}^2 \right) ds \\ & \quad + \varepsilon \mathbb{E} \int_0^{t \wedge \tau_{n,N}} \int_D \left(a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2} \right) |\nabla \rho_N(s)|^2 dx ds \leq CE \mathcal{E}_{0,\delta}, \end{aligned}$$

from which we get the following L^2 energy estimates for any $t \in [0, T]$:

$$\begin{aligned} & \mathbb{E} \sup_{0 \leq s \leq t} \mathcal{E}_\delta + \mathbb{E} \int_0^t \left(\mu \|\nabla u_N(s)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(s)\|_{L^2(D)}^2 + \nu \|\nabla B_N(s)\|_{L^2(D)}^2 \right) ds \\ & \quad + \varepsilon \mathbb{E} \int_0^t \int_D \left(a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2} \right) |\nabla \rho_N(s)|^2 dx ds \leq CE \mathcal{E}_{0,\delta}. \end{aligned} \tag{3.12}$$

In the same way, for $p > 2$, by virtue of the estimates on $|I_2|^p$ and $|I_4|^p$, we obtain the L^p energy estimates for $t \in [0, T]$

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq t} \mathcal{E}_\delta + \mathbb{E} \int_0^t \left(\mu \|\nabla u_N(s)\|_{L^2(D)}^2 + (\lambda + \mu) \|\operatorname{div} u_N(s)\|_{L^2(D)}^2 + \nu \|\nabla B_N(s)\|_{L^2(D)}^2 \right) ds \right. \\ & \quad \left. + \varepsilon \mathbb{E} \int_0^t \int_D \left(a\gamma \rho_N^{\gamma-2} + \delta\beta \rho_N^{\beta-2} \right) |\nabla \rho_N(s)|^2 dx ds \right]^p \leq CE (\mathcal{E}_{0,\delta})^p, \end{aligned}$$

which completes the proof of the proposition. \square

Now, we are ready to prove Lemma 3.3.

Proof of Lemma 3.3. It follows from (3.12) that

$$\mathbb{E} \int_0^T \|\nabla u_N\|_{L^2(D)}^2 dt \leq CE \mathcal{E}_{0,\delta}, \quad \mathbb{E} \sup_{t \in [0, T]} \|\sqrt{\rho_N} u_N(t)\|_{L^2(D)}^2 \leq CE \mathcal{E}_{0,\delta}.$$

Since $\dim X_n$ is finite, the L^∞ , C^2 and L^2 norms are equivalent on X_n . Moreover, $\rho = \mathcal{S}[u]$ is bounded. It follows that

$$\underline{\rho} \exp \left(- \int_0^T \|\nabla u_N(s)\|_{L^2(D)}^2 ds \right) \lesssim \rho_N(t, x) \lesssim \bar{\rho} \exp \left(\int_0^T \|\nabla u_N(s)\|_{L^2(D)}^2 ds \right), \tag{3.13}$$

which yields that

$$\mathbb{E} \left[\exp \left(- \int_0^T \|\nabla u_N(s)\|_{L^2(D)}^2 ds \right) \sup_{t \in [0, T]} \|u_N\|_{L^2(D)} \right] \leq C, \tag{3.14}$$

where C is independent of N . It follows from (3.9), (3.13) and (3.14) that $\lim_{N \rightarrow \infty} \mathbb{P}(\tau_{n,N} = T) = 1$. \square

Moreover, it follows from (3.9) that

$$\sqrt{\varepsilon\delta}\rho_N^{\frac{\beta}{2}} \text{ is bounded in } L^p(\Omega, L^2(0, T; H^1(D))).$$

The Sobolev embedding $H^1(D) \hookrightarrow L^6(D)$ yields that

$$\rho_N^{\frac{\beta}{2}} \in L^p(\Omega, L^2(0, T; L^6(D))),$$

that is,

$$\mathbb{E} \left(\|\rho_N^\beta\|_{L^1(0, T; L^3(D))} \right)^{2p} \leq C, \text{ where } C \text{ is independent of } N.$$

In view of (3.9), we deduce that

$$\mathbb{E} \left(\sup_{t \in [0, T]} \|\rho_N^\beta\|_{L^1(D)} \right)^p \leq C,$$

this together with the interpolation inequality $\|\rho_N^\beta\|_{L^2(D)} \lesssim \|\rho_N^\beta\|_{L^1(D)}^{\frac{1}{4}} \|\rho_N^\beta\|_{L^3(D)}^{\frac{3}{4}}$ yields that

$$\begin{aligned} \mathbb{E} \left(\int_0^T \|\rho_N^\beta\|_{L^2(D)}^{\frac{4}{3}} dt \right)^p &\lesssim \mathbb{E} \left(\int_0^T \|\rho_N^\beta\|_{L^1(D)}^{\frac{1}{3}} \|\rho_N^\beta\|_{L^3(D)} dt \right)^p \\ &\lesssim \mathbb{E} \left(\sup_{t \in [0, T]} \|\rho_N^\beta(t)\|_{L^1(D)} \right)^{\frac{2p}{3}} + \mathbb{E} \left(\int_0^T \|\rho_N^\beta\|_{L^3(D)} dt \right)^{2p} \\ &\leq C. \end{aligned}$$

By the Hölder inequality, we have

$$\mathbb{E} \left(\int_0^T \int_D \rho_N^{\frac{4\beta}{3}} dx dt \right)^p \leq \mathbb{E} \left(\int_0^T \left(\int_D \rho_N^{2\beta} dx \right)^{\frac{2}{3}} \left(\int_D dx \right)^{\frac{1}{3}} dt \right) \leq C.$$

Thus

$$\mathbb{E} \left(\|\rho_N\|_{L^{\beta+1}((0, T) \times D)} \right)^p \leq C \text{ if } \beta \geq 3.$$

If $\beta \geq 4$, multiply (3.1)₁ by ρ_N and integrate by parts to get

$$\begin{aligned} \mathbb{E} \left(\varepsilon \int_0^T \|\nabla \rho_N\|_{L^2(D)}^2 dt \right)^p &\lesssim \mathbb{E} \left(\|\rho_N(0)\|_{L^2(D)}^2 + \int_0^T \int_D \rho_N^2 \operatorname{div} u_N dx dt \right)^p \\ &\lesssim \mathbb{E} \left(\|\rho_N(0)\|_{L^2(D)}^2 + \sup_{t \in [0, T]} \|\rho_N\|_{L^4(D)}^4 + \|\nabla u_N\|_{L^2([0, T] \times D)}^2 \right)^p \\ &\lesssim C. \end{aligned}$$

For each $t \in [0, T]$, we define $(\rho_n, u_n, B_n) := \lim_{N \rightarrow \infty} (\rho_n^N, u_n^N, B_n^N)$. By the same argument as above, we can infer that (ρ_n, u_n, B_n) satisfies the corresponding a priori estimate uniformly in n . More precisely, for any $1 \leq p < \infty$, we have the following lemma:

Lemma 3.5. Let (ρ_n, u_n, B_n) be the solution of (3.1)–(3.2) on $\Omega \times (0, T) \times D$ constructed above. Then, for $\beta \geq 4$, we have,

$$\begin{aligned}
& \mathbb{E} \left(\sup_{t \in [0, T]} \|\rho_n(t)\|_{L^\gamma(D)}^\gamma \right)^p \leq C, \quad \mathbb{E} \left(\delta \sup_{t \in [0, T]} \|\rho_n(t)\|_{L^\beta(D)}^\beta \right)^p \leq C, \\
& \mathbb{E} \left(\sup_{t \in [0, T]} \|\sqrt{\rho_n(t)} u_n(t)\|_{L^2(D)}^2 \right)^p \leq C, \quad \mathbb{E} \left(\|u_n(t)\|_{L^2([0, T]; H^1(D))}^2 \right)^p \leq C, \\
& \mathbb{E} \left(\sup_{t \in [0, T]} \|B_n(t)\|_{L^2(D)}^2 \right)^p \leq C, \quad \mathbb{E} \left(\|B_n(t)\|_{L^2([0, T]; H^1(D))}^2 \right)^p \leq C, \\
& \mathbb{E} \left(\varepsilon \|\nabla \rho_n(t)\|_{L^2((0, T) \times D)}^2 \right)^p \leq C, \quad \mathbb{E} \left(\|\rho_n\|_{L^{\beta+1}((0, T) \times D)} \right)^p \leq C,
\end{aligned} \tag{3.15}$$

where the constant C is independent of n .

3.2. Tightness Property. In this subsection, we shall show the tightness property for the approximation solution in the following lemma.

Lemma 3.6. Define

$$\begin{aligned}
S := & C(0, T; \mathbb{R}) \times \left(C([0, T]; L_w^\beta(D)) \cap L^2(0, T; L^2(D)) \cap L^2(0, T; H_w^1(D)) \right) \\
& \times L^2(0, T; H_w^1(D)) \times C([0, T]; L_w^{\frac{2\beta}{\beta+1}}(D)) \times (L^2(0, T; H_w^1(D)) \cap L^2(0, T; L^2(D)))
\end{aligned}$$

equipped with its Borel σ -algebra. Let Π_n be the probability on S which is the image of \mathbb{P} on Ω by the map: $\omega \mapsto (\hat{\beta}_k(\omega, \cdot), \rho_n(\omega, \cdot), u_n(\omega, \cdot), \rho_n u_n(\omega, \cdot), B_n(\omega, \cdot))$, that is, for any $A \subseteq S$,

$$\Pi_n(A) = \mathbb{P} \left\{ \omega \in \Omega : (\hat{\beta}_k(\omega, \cdot), \rho_n(\omega, \cdot), u_n(\omega, \cdot), \rho_n u_n(\omega, \cdot), B_n(\omega, \cdot)) \in A \right\}.$$

Then the family Π_n is tight.

Proof. We want to check the tightness of the family of Π_n in the following five steps:

1° (the tightness of $\hat{\beta}_k$) First, we will check the tightness of $\hat{\beta}_k$, that is, for $\varepsilon > 0$, we now need to find the compact subset $\Sigma_\varepsilon \subset C(0, T; \mathbb{R})$ such that $\mathbb{P}(\hat{\beta}_k \notin \Sigma_\varepsilon) \leq \frac{\varepsilon}{5}$. For Σ_ε we rely on classical results concerning the Brownian motion. For a constant L_ε to be chosen later, we consider the set

$$\Sigma_\varepsilon = \left\{ \hat{\beta}_k(\cdot) \in C(0, T; \mathbb{R}) : \sup_{\substack{t_1, t_2 \in [0, T], \\ |t_1 - t_2| < \frac{1}{m^6}}} m |\hat{\beta}_k(t_2) - \hat{\beta}_k(t_1)| \leq L_\varepsilon, \forall m \in \mathbb{N} \right\}.$$

Σ_ε is relatively compact in $C(0, T; \mathbb{R})$ by Arzela-Ascoli's Theorem. Furthermore Σ_ε is closed in $C(0, T; \mathbb{R})$. Therefore Σ_ε is a compact subset of $C(0, T; \mathbb{R})$. We can show that $\mathbb{P}(\hat{\beta}_k \notin \Sigma_\varepsilon) \leq \frac{C}{L_\varepsilon^4}$. In fact, by Chebyshev's inequality $\mathbb{P}\{\omega : \xi(\omega) \geq r\} \leq \frac{1}{r^q} \mathbb{E}[|\xi(\omega)|^q]$, one has

$$\begin{aligned}
& \mathbb{P}\{\omega : \hat{\beta}_k(\omega, \cdot) \notin \Sigma_\varepsilon\} \\
& \leq \mathbb{P} \left[\bigcup_{m=1}^{\infty} \left\{ \omega : \sup_{t_1, t_2 \in [0, T], |t_1 - t_2| < m^{-6}} |\hat{\beta}_k(t_1) - \hat{\beta}_k(t_2)| > \frac{L_\varepsilon}{m} \right\} \right] \\
& \leq \sum_{m=1}^{\infty} \sum_{i=0}^{m^6-1} \left(\frac{m}{L_\varepsilon} \right)^4 \mathbb{E} \left[\sup_{iTm^{-6} \leq t \leq (i+1)Tm^{-6}} |\hat{\beta}_k(t) - \hat{\beta}_k(iTm^{-6})|^4 \right]
\end{aligned}$$

$$\leq C \sum_{m=1}^{\infty} \left(\frac{m}{L_\varepsilon} \right)^4 (Tm^{-6})^2 m^6 = \frac{C}{L_\varepsilon^4} \sum_{m=1}^{\infty} \frac{1}{m^2}.$$

We choose $L_\varepsilon^4 = \frac{1}{5C\varepsilon} \left(\sum_{m=1}^{\infty} \frac{1}{m^2} \right)^{-1}$ to obtain $\mathbb{P}(\hat{\beta}_k \notin \Sigma_\varepsilon) \leq \frac{\varepsilon}{5}$.

2° (the tightness of ρ_n) In this step, we want to find $X_\varepsilon \subset C([0, T]; L_w^\beta(D)) \cap L^2(0, T; L^2(D)) \cap L^2(0, T; H_w^1(D))$ such that $\mathbb{P}(\rho_n \notin X_\varepsilon) \leq \frac{\varepsilon}{5}$. For this, we define a function space \mathcal{X} with the norm

$$\begin{aligned} \|f\|_{\mathcal{X}} &= \sup_{0 \leq t \leq T} \|f(t)\|_{L^\beta(D)} + \|f(t)\|_{L^2([0, T]; H^1(D))} \\ &\quad + \sup_{0 \leq t \leq T} \|\partial_t f\|_{W^{-1, \frac{2\beta}{\beta+1}}(D)} + \|\partial_t f\|_{L^2([0, T]; H^{-1}(D))}. \end{aligned}$$

Choose X_ε to be a closed ball of radius r_ε centered at 0 in \mathcal{X} . By Aubin-Lions Lemma, we know that X_ε is compact in $C(0, T; L_w^\beta(D)) \cap L^2(0, T; H_w^1(D)) \cap L^2(0, T; L^2(D))$. It follows from (3.15) in Lemma 3.5 that

$$\mathbb{P}(\rho_n \notin X_\varepsilon) = \mathbb{P}(\|\rho_n\|_{\mathcal{X}} > r_\varepsilon) \leq \frac{1}{r_\varepsilon} \mathbb{E}(\|\rho_n\|_{\mathcal{X}}) \leq \frac{C}{r_\varepsilon}.$$

Choosing $r_\varepsilon = 5C\varepsilon^{-1}$, we have $\mathbb{P}(\rho_n \notin X_\varepsilon) \leq \frac{\varepsilon}{5}$. Then $\mathbb{P}\{\omega : \rho_n(\omega, \cdot) \in X_\varepsilon\} \geq 1 - \frac{\varepsilon}{5}$.

3° (the tightness of u_n) In this step, we find $Y_\varepsilon \subset L^2(0, T; H_w^1(D))$ such that $\mathbb{P}(u_n \notin Y_\varepsilon) \leq \frac{\varepsilon}{5}$. To this end, we choose Y_ε as a closed ball of radius \tilde{r}_ε centered at 0 in $L^2(0, T; H_w^1(D))$. Then Y_ε is compact in $L^2(0, T; H_w^1(D))$. (3.15) in Lemma 3.5 implies that

$$\mathbb{P}(u_n \notin Y_\varepsilon) = \mathbb{P}(\|u_n\|_{L^2(0, T; H^1(D))} > \tilde{r}_\varepsilon) \leq \frac{1}{\tilde{r}_\varepsilon} \mathbb{E}(\|u_n\|_{L^2(0, T; H^1(D))}) \leq \frac{C}{\tilde{r}_\varepsilon}.$$

Choosing $\tilde{r}_\varepsilon = 5C\varepsilon^{-1}$, we have $\mathbb{P}(u_n \notin Y_\varepsilon) \leq \frac{\varepsilon}{5}$. Then $\mathbb{P}\{\omega : u_n(\omega, \cdot) \in Y_\varepsilon\} \geq 1 - \frac{\varepsilon}{5}$.

4° (the tightness of $\rho_n u_n$) In this step, we find $Z_\varepsilon \subset C([0, T]; L_w^{\frac{2\beta}{\beta+1}}(D))$ such that $\mathbb{P}(\rho_n u_n \notin Z_\varepsilon) \leq \frac{\varepsilon}{5}$. For this, define a function space $\mathcal{Z} = L^\infty([0, T]; L^{\frac{2\beta}{\beta+1}}(D)) \cap C^{0, \alpha}([0, T]; W^{-\ell, 2}(D))$, $\ell > 3$, $0 < \alpha < \frac{1}{2}$ with the norm

$$\|f\|_{\mathcal{Z}} = \sup_{0 \leq t \leq T} \|f(t)\|_{L^{\frac{2\beta}{\beta+1}}(D)} + \|f(t)\|_{C^{0, \alpha}([0, T]; W^{-\ell, 2}(D))}.$$

Because the Brownian motion is not differentiable in time, we can not estimate $\partial_t(\rho_n u_n)$ directly. To overcome the difficulty, we consider the momentum equation directly. From (3.1), we have

$$\begin{aligned} \mathbb{P}[\rho_n u_n(t)] &= \mathbb{P}[\rho_0 u_0] - \int_0^t \mathbb{P} \left[\operatorname{div}(\rho_n u_n \otimes u_n) - \mu \Delta u_n - (\lambda + \mu) \nabla \operatorname{div} u_n - a \nabla \rho_n^\gamma - \delta \nabla \rho_n^\beta \right] ds \\ &\quad + \int_0^t \mathbb{P} [(\nabla \times B_n) \times B_n - \varepsilon \nabla u_n \cdot \nabla \rho_n] ds - \int_0^t \sum_{k \geq 1} f_k^n(\rho_n, \rho_n u_n, x) d\hat{\beta}_k^1, \end{aligned} \quad (3.16)$$

where $\mathbb{P} : L^2(D) \rightarrow X_n$ is the projection onto X_n . First, from (3.15), we have

$$\mathbb{E} \int_0^T \|\mathbb{P}[\operatorname{div}(\rho_n u_n \otimes u_n)]\|_{W^{-1, \frac{6\beta}{4\beta+3}}(D)}^2 ds \leq \mathbb{E} \int_0^T \|\rho_n u_n \otimes u_n\|_{L^{\frac{6\beta}{4\beta+3}}(D)}^2 ds \leq C. \quad (3.17)$$

Similarly, from Lemma 3.5, we can handle the other deterministic terms as:

$$\begin{aligned}
& \mathbb{E} \int_0^T \|\mathbb{P}[\mu\Delta u_n + (\lambda + \mu)\nabla\operatorname{div}u_n]\|_{W^{-1,2}(D)}^2 ds \leq C, \\
& \mathbb{E} \int_0^T \|\mathbb{P}(\varepsilon\nabla u_n \cdot \nabla\rho_n)\|_{W^{-\ell,2}(D)}^{\frac{2q}{q+2}} ds \leq C, \\
& \mathbb{E} \int_0^T \|\mathbb{P}(a\nabla\rho_n^\gamma + \delta\nabla\rho_n^\beta)\|_{W^{-1, \frac{\beta+1}{\beta}}(D)}^{\frac{\beta+1}{\beta}} ds \leq C, \\
& \mathbb{E} \int_0^T \|\mathbb{P}[(\nabla \times B_n) \times B_n]\|_{W^{-\ell,2}(D)}^2 ds \leq C.
\end{aligned} \tag{3.18}$$

Here we have used $\nabla\rho_n \in L^q([0, T]; L^2(D))$, $q > 2$ (see [14, Lemma 2.4]). For the stochastic terms, by Lemma 3.5, one has for $\sigma > 1$

$$\begin{aligned}
& \mathbb{E} \left(\int_s^t \sum_{k \geq 1} \|f_k^n(\rho_n, \rho_n u_n, x)\|_{W^{-\ell,2}(D)} d\hat{\beta}_k^1(s) \right)^{2\sigma} \\
& \leq \mathbb{E} \left[\int_s^t \sum_{k=1} \|f_k^n(\rho_n, \rho_n u_n, x)\|_{W^{-\ell,2}(D)}^2 ds \right]^\sigma \\
& \leq \mathbb{E} \left[\int_s^t \sum_{k=1} \|\mathbb{P}(f_k(\rho_n, \rho_n u_n, x)/\sqrt{\rho_n})\|_{L^2(D)}^2 \sup_{\|\varphi\|_{W^{\ell,2}(D)}=1} \|\mathcal{M}^{\frac{1}{2}}[\rho_n]\varphi\|_{L^2(D)}^2 ds \right]^\sigma \\
& \leq C(t-s)^\sigma.
\end{aligned} \tag{3.19}$$

Thanks to the Kolmogorov continuity Theorem, we can infer that the stochastic term is almost surely α -Hölder continuous for every $0 < \alpha < \frac{1}{2} - \frac{1}{2\sigma}$. Note that when $\ell > 3$, we have $W^{-1, \frac{6\beta}{4\beta+3}}(D) \hookrightarrow W^{-\ell,2}(D)$ and $W^{-1, \frac{\beta+1}{\beta}}(D) \hookrightarrow W^{-\ell,2}(D)$.

Choose Z_ε to be a closed ball of radius r'_ε centered at 0 in \mathcal{Z} . By [28, Corollary B.2] and [33], we know that Z_ε is compact in $C([0, T]; L_w^{\frac{2\beta}{\beta+1}}(D))$. Moreover, from (3.16)-(3.19), by the Chebyshev inequality, we have

$$\mathbb{P}(\rho_n u_n \notin Z_\varepsilon) = \mathbb{P}(\|\rho_n u_n\|_{\mathcal{Z}} > r'_\varepsilon) \leq \frac{1}{r'_\varepsilon} \mathbb{E}(\|\rho_n u_n\|_{\mathcal{Z}}) \leq \frac{C}{r'_\varepsilon}. \tag{3.20}$$

Let $r'_\varepsilon = 5C\varepsilon^{-1}$ in (3.20), one deduces that

$$\mathbb{P}(\rho_n u_n \notin Z_\varepsilon) \leq \frac{\varepsilon}{5}.$$

Then $\mathbb{P}\{\omega : \rho_n u_n(\omega, \cdot) \in Z_\varepsilon\} \geq 1 - \frac{\varepsilon}{5}$.

Remark 3.7. Here we can also use the following tightness criterion: Let $\{f_n\}_{n=1}^\infty$ be a collection of $C([0, T], L_w^p(D))$ -valued random variables defined on a probability space $(\Omega_n, \mathcal{F}_n, \mathbb{P}_n)$ such that $\sup_n \mathbb{E}(\|f_n\|_{L^\infty([0, T], L^p(D))}) < \infty$ and for any $\varphi \in C_0^\infty(D)$, there exist an integer $k, \sigma > 0, q > \frac{1}{\sigma}$ such that $\sup_n \mathbb{E}[|\langle f_n(t) - f_n(s), \varphi \rangle|^q] \leq \|\varphi\|_{C^k}^q |t - s|^{\sigma q}$ for all $0 \leq s, t \leq T$. Then the sequence of the induced measures $\mathbb{P} \circ f_n^{-1}$ are tight on $C([0, T], L_w^p(D))$.

For example, by Hölder's inequality and Lemma 3.5, we have

$$\mathbb{E} \left| \int_s^t \langle \operatorname{div}(\rho_n u_n \otimes u_n), \varphi \rangle dr \right|$$

$$\begin{aligned}
&\leq \mathbb{E} \int_s^t \|\nabla \varphi\|_{L^\infty} \|\sqrt{\rho_n} u_n\|_{L^2(D)} \|\nabla u_n\|_{L^2(D)} \|\rho_n\|_{L^\gamma(D)}^{\frac{1}{2}} dr \\
&\leq C(t-s)^{\frac{1}{2}} \mathbb{E} \left(\|\sqrt{\rho_n} u_n\|_{L^\infty(0,T;L^2(D))} \|\nabla u_n\|_{L^2(0,T;L^2(D))} \|\rho_n\|_{L^\infty(0,T;L^\gamma(D))}^{\frac{1}{2}} \right) \\
&\leq C(t-s)^{\frac{1}{2}}, \\
&\mathbb{E} \left| \int_s^t \langle (\nabla \times B_n) \times B_n, \varphi \rangle dr \right| \\
&\leq C(t-s)^{\frac{1}{2}} \|\varphi\|_{L^\infty} \mathbb{E} (\|\nabla B_n\|_{L^2([0,T],L^2(D))} \|B_n\|_{L^\infty([0,T],L^2(D))}) \leq C(t-s)^{\frac{1}{2}}, \\
&\mathbb{E} \left| \int_s^t \langle \mu \Delta u_n + (\lambda + \mu) \nabla \operatorname{div} u_n, \varphi \rangle dr \right| \\
&\leq C(t-s)^{\frac{1}{2}} \|\nabla \varphi\|_{L^\infty} \mathbb{E} (\|\nabla u_n\|_{L^2([0,T],L^2(D))}) \leq C(t-s)^{\frac{1}{2}}, \\
&\mathbb{E} \left| \int_s^t \langle a \nabla \rho_n^\gamma - \delta \nabla \rho_n^\beta, \varphi \rangle dr \right| \\
&\leq C \mathbb{E} \int_s^t \|\nabla \varphi\|_{L^\infty} \|\rho_n\|_{L^\gamma(D)}^\gamma \|\rho_n\|_{L^\beta(D)}^\beta dr \leq C(t-s)^{\frac{1}{2}}, \\
&\mathbb{E} \left| \int_s^t \langle \varepsilon \nabla u_n \cdot \nabla \rho_n, \varphi \rangle dr \right| \\
&\leq \mathbb{E} \int_s^t \varepsilon \|\nabla u_n\|_{L^2(D)} \|\nabla \rho_n\|_{L^2(D)} \|\varphi\|_{L^\infty(D)} dr \\
&\leq (t-s)^{\frac{1}{2}-\frac{1}{q}} \mathbb{E} (\|\nabla u_n\|_{L^2([0,T],L^2(D))} \|\nabla \rho_n\|_{L^q([0,T],L^2(D))}) \leq C(t-s)^{\frac{1}{2}-\frac{1}{q}}.
\end{aligned}$$

Here we have used $\nabla \rho_n \in L^q([0, T]; L^2(D))$, $q > 2$. Then we can get the tightness of $\rho_n u_n$.

5° (the tightness of B_n) First, we will find $R_\varepsilon \subset L^2(0, T; H_w^1(D))$ such that $\mathbb{P}(B_n \notin R_\varepsilon) \leq \frac{\varepsilon}{5}$. First, we choose R_ε as a closed ball of radius \mathring{r}_ε centered at 0 in $L^2(0, T; H^1(D))$. Then R_ε is compact in $L^2(0, T; H_w^1(D))$. From (3.15), we have

$$\mathbb{P}(B_n \notin R_\varepsilon) = \mathbb{P}(\|B_n\|_{L^2(0,T;H^1(D))} > \mathring{r}_\varepsilon) \leq \frac{1}{\mathring{r}_\varepsilon} \mathbb{E} (\|B_n\|_{L^2(0,T;H^1(D))}) \leq \frac{C}{\mathring{r}_\varepsilon}.$$

Choosing $\mathring{r}_\varepsilon = 5C\varepsilon^{-1}$, we have $\mathbb{P}(B_n \notin R_\varepsilon) \leq \frac{\varepsilon}{5}$. Then $\mathbb{P}\{\omega : B_n(\omega, \cdot) \in R_\varepsilon\} \geq 1 - \frac{\varepsilon}{5}$.

On the other hand, it follows from (3.1) that

$$\begin{aligned}
\mathbb{E} \int_0^{T-\theta} \|B_n(t+\theta) - B_n(t)\|_{H^{-1}(D)}^2 dt &= \mathbb{E} \int_0^{T-\theta} \left\| \int_t^{t+\theta} dB_n(s) \right\|_{H^{-1}(D)}^2 dt \\
&\leq \mathbb{E} \int_0^{T-\theta} (J_1 + J_2 + J_3) dt,
\end{aligned} \tag{3.21}$$

where

$$J_1(t) = \left\| \int_t^{t+\theta} \nabla \times (u_n \times B_n) ds \right\|_{H^{-1}(D)}^2, \quad J_2(t) = \left\| \int_t^{t+\theta} \nu \Delta B_n ds \right\|_{H^{-1}(D)}^2,$$

$$J_3(t) = \left\| \int_t^{t+\theta} \sum_{k \geq 1} g_k(B_n, x) d\hat{\beta}_k^2 \right\|_{H^{-1}(D)}^2.$$

For the term $J_1(t)$, one has

$$\begin{aligned} J_1^{1/2} &= \sup_{\varphi \in H_0^1(D), \|\varphi\|_{H^1(D)}=1} \left\{ \int_D \left(\int_t^{t+\theta} \nabla \times (u_n \times B_n) ds \right) \varphi(x) dx \right\} \\ &\leq C \int_t^{t+\theta} \|u_n\|_{L^4(D)} \|B_n\|_{L^4(D)} ds. \end{aligned}$$

By Hölder's inequality and Lemma 3.5, then

$$\mathbb{E} \int_0^{T-\theta} J_1(t) dt \lesssim \theta^{\frac{1}{4}} \mathbb{E} \left(\|u_n\|_{L^2(0,T;H^1(D))}^2 \|B_n\|_{L^{\frac{8}{3}}(0,T;L^4(D))}^2 \right) \lesssim \theta^{\frac{1}{4}}. \quad (3.22)$$

For the term $J_2(t)$, similar to (3.22), we have

$$\mathbb{E} \int_0^{T-\theta} J_2(t) dt \lesssim \mathbb{E} \int_0^{T-\theta} \left(\int_t^{t+\theta} \|\nabla B_n\|_{L^2(D)} ds \right)^2 dt \lesssim \theta \mathbb{E} \int_0^T \|\nabla B_n\|_{L^2(D)}^2 dt \lesssim \theta. \quad (3.23)$$

For the term J_3 , using the Burkholder-Davis-Gundy inequality, Hölder's inequality and (1.5), we obtain

$$\begin{aligned} \mathbb{E} \int_0^{T-\theta} J_3(t) dt &\leq \int_0^T \mathbb{E} \left(\sup_{\varphi \in H_0^1(D), \|\varphi\|_{H^1(D)}=1} \int_t^{t+\theta} \int_D \sum_{k \geq 1} g_k(B_n, x) \varphi dx d\hat{\beta}_k^2 \right)^2 dt \\ &\leq \int_0^T \mathbb{E} \left(\sup_{\varphi \in H_0^1(D), \|\varphi\|_{H^1(D)}=1} \int_t^{t+\theta} \left(\sum_{k \geq 1} \int_D g_k(B_n, x) \varphi dx \right)^2 ds \right) dt \\ &\leq \int_0^T \left(\mathbb{E} \int_t^{t+\theta} \|\varphi\|_{L^2(D)}^2 \sum_{k=1} \|\hat{g}_k(B_n, x)\|_{L^2(D)}^2 ds \right) dt \\ &\leq \int_0^T \left(\mathbb{E} \int_t^{t+\theta} \|B_n\|_{L^2(D)}^2 ds \right) dt \\ &\lesssim \theta. \end{aligned} \quad (3.24)$$

Combining (3.21)-(3.24), and then by Aubin-Simon Lemma in [33], we know that B_n is compact in $L^2(0, T; L^2(D))$.

From the above argument, it follows from [24, pp.16, Corollary1.3] that the distribution of the joint processes $(\hat{\beta}_k, \rho_n, u_n, \rho_n u_n, B_n)$ are tight. Hence, the proof of the lemma is completed. \square

3.3. Application of Jakubowski-Skorokhod Theorem. From the tightness property and Jakubowski-Skorokhod's Theorem, there exists a subsequence, still indexed by n , such that $\Pi_n \rightarrow \Pi$ weakly, where Π is a probability on S , a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$; and there exist random variables $(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n)$ with distribution Π_n , (β_k, ρ, u, h, B) with values in S such that

$$(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n) \rightarrow (\beta_k, \rho, u, h, B) \text{ in } S \tilde{\mathbb{P}} - \text{a.s.} \quad (3.25)$$

We set

$$\tilde{\mathcal{F}}_t = \sigma\{\beta_k(s), \rho(s), u(s), B(s)\}_{s \in [0, t]},$$

which is the union σ -algebra generated by random variables $(\beta_k(s), \rho(s), u(s), B(s))$ for $s \in [0, t]$. It's easy to see that $\beta_k(t)$ is a $\tilde{\mathcal{F}}_t$ standard Brownian motion.

Now, we need to prove that $(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{B}_n)$ satisfies the equation (3.1), that is,

$$\mathbb{P}[\tilde{\rho}_n(t) - \tilde{\rho}_n(0)] = \int_0^t \mathbb{P}[\varepsilon \Delta \tilde{\rho}_n - \operatorname{div}(\tilde{\rho}_n \tilde{u}_n)] ds,$$

$$\begin{aligned} & \mathbb{P}[\tilde{\rho}_n \tilde{u}_n(t)] + \int_0^t \mathbb{P} \left[\operatorname{div}(\tilde{\rho}_n \tilde{u}_n \otimes \tilde{u}_n) - \mu \Delta \tilde{u}_n - (\lambda + \mu) \nabla \operatorname{div} \tilde{u}_n + a \nabla \tilde{\rho}_n^\gamma + \delta \nabla \tilde{\rho}_n^\beta \right] ds \\ &= \mathbb{P}[\tilde{\rho}_n \tilde{u}_n(0)] + \int_0^t \mathbb{P} \left((\nabla \times \tilde{B}_n) \times \tilde{B}_n - \varepsilon \nabla \tilde{u}_n \cdot \nabla \tilde{\rho}_n \right) ds + \int_0^t \sum_{k \geq 1} f_k^n(\tilde{\rho}_n, \tilde{\rho}_n \tilde{u}_n, x) d\tilde{\beta}_{k,n}^1, \end{aligned}$$

and

$$\mathbb{P}[\tilde{B}_n(t)] - \int_0^t \mathbb{P} \left[\nabla \times (\tilde{u}_n \times \tilde{B}_n) + \nu \Delta \tilde{B}_n \right] ds = \mathbb{P}[\tilde{B}_n(0)] + \int_0^t \sum_{k \geq 1} \mathbb{P} \left[g_k(\tilde{B}_n, x) \right] d\tilde{\beta}_{k,n}^2.$$

Here \mathbb{P} is a projection from L^2 to X_n . To this end, we define

$$\varpi_n(t) = \mathbb{P}[\rho_n(t) - \rho(0)] - \int_0^t \mathbb{P}[\varepsilon \Delta \rho_n - \operatorname{div}(\rho_n u_n)] ds,$$

$$\begin{aligned} \xi_n(t) &= \mathbb{P}[\rho_n u_n(t)] + \int_0^t \mathbb{P} \left[\operatorname{div}(\rho_n u_n \otimes u_n) - \mu \Delta u_n - (\lambda + \mu) \nabla \operatorname{div} u_n + a \nabla \rho_n^\gamma + \delta \nabla \rho_n^\beta \right] ds \\ &\quad - \mathbb{P}[\rho_n u_n(0)] - \int_0^t \mathbb{P} \left((\nabla \times B_n) \times B_n - \varepsilon \nabla u_n \cdot \nabla \rho_n \right) ds + \int_0^t \sum_{k \geq 1} f_k^n(\rho_n, \rho_n u_n) d\hat{\beta}_k^1, \end{aligned}$$

$$\zeta_n(t) = \mathbb{P}[B_n(t) - B_n(0)] - \int_0^t \mathbb{P} \left[\nabla \times (u_n \times B_n) + \nu \Delta B_n \right] dt - \int_0^t \sum_{k \geq 1} \mathbb{P} \left[g_k(B_n, x) \right] d\hat{\beta}_k^2,$$

$$G_n = \int_0^T \|\varpi_n(t)\|_{H^{-1}(D)}^2 dt, \quad Z_n = \int_0^T \|\xi_n(t)\|_{W^{-\ell, 2}(D)}^2 dt, \quad \ell \geq 3, \quad H_n = \int_0^T \|\zeta_n(t)\|_{H^{-1}(D)}^2 dt.$$

Since (ρ_n, u_n, B_n) is a solution, we have

$$G_n = 0, \quad Z_n = 0, \quad H_n = 0, \quad \mathbb{P} - \text{a.s.} \quad (3.26)$$

Let

$$\tilde{\varpi}_n(t) = \mathbb{P}[\tilde{\rho}_n(t) - \tilde{\rho}_n(0)] - \int_0^t \mathbb{P}[\varepsilon \Delta \tilde{\rho}_n - \operatorname{div}(\tilde{\rho}_n \tilde{u}_n)] ds,$$

$$\begin{aligned} \tilde{\xi}_n(t) &= \mathbb{P}[\tilde{\rho}_n \tilde{u}_n(t)] - \int_0^t \mathbb{P} \left[\operatorname{div}(\tilde{\rho}_n \tilde{u}_n \otimes \tilde{u}_n) - \mu \Delta \tilde{u}_n - (\lambda + \mu) \nabla \operatorname{div} \tilde{u}_n + a \nabla \tilde{\rho}_n^\gamma + \delta \nabla \tilde{\rho}_n^\beta \right] ds \\ &\quad - \mathbb{P}[\tilde{\rho}_n \tilde{u}_n(0)] + \int_0^t \mathbb{P} \left(\varepsilon \nabla \tilde{u}_n \cdot \nabla \tilde{\rho}_n + (\nabla \times \tilde{B}_n) \times \tilde{B}_n \right) ds - \int_0^t \sum_{k \geq 1} f_k^n(\tilde{\rho}_n, \tilde{\rho}_n \tilde{u}_n, x) d\tilde{\beta}_{k,n}^1, \end{aligned}$$

$$\tilde{\zeta}_n(t) = \mathbb{P}[\tilde{B}_n(t) - \tilde{B}_n(0)] - \int_0^t \mathbb{P}[\nabla \times (\tilde{u}_n \times \tilde{B}_n) + \nu \Delta \tilde{B}_n] dt - \int_0^t \sum_{k \geq 1} \mathbb{P}[g_k(\tilde{B}_n, x)] d\tilde{\beta}_{k,n}^2,$$

and

$$\tilde{G}_n = \int_0^T \|\tilde{\omega}_n(t)\|_{H^{-1}(D)}^2 dt, \quad \tilde{Z}_n = \int_0^T \|\tilde{\xi}_n(t)\|_{W^{-\ell,2}(D)}^2 dt, \quad \tilde{H}_n = \int_0^T \|\tilde{\zeta}_n(t)\|_{H^{-1}(D)}^2 dt.$$

We want to verify that

$$\tilde{\mathbb{E}}\tilde{G}_n = 0, \quad \tilde{\mathbb{E}}\tilde{Z}_n = 0, \quad \tilde{\mathbb{E}}\tilde{H}_n = 0.$$

Here $\tilde{\mathbb{E}}$ denotes the mathematical expectation with respect to the probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$. To this end, we have the following proposition:

Proposition 3.8. $\tilde{G}_n = 0, \tilde{Z}_n = 0, \tilde{H}_n = 0$ $\tilde{\mathbb{P}}$ - a.s., that is, $(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{B}_n)$ solves the equation (3.1).

Proof. The difficulty comes from the fact that (Z_n, H_n) is not expressed as a deterministic function of $(\hat{\beta}_k, \rho_n, u_n, B_n)$ due to the presence of the stochastic integrals. By Theorem 2.4 and [2, Corollary 2.5], we can infer that

$$\mathcal{L}(\hat{\beta}_k, \rho_n, u_n, \rho_n u_n, B_n, \xi_n, \zeta_n) = \mathcal{L}(\tilde{\beta}_{k,n}, \tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n, \tilde{\xi}_n, \tilde{\zeta}_n). \quad (3.27)$$

Here $\mathcal{L}(f)$ is the probability distribution of f . Note that $(\tilde{Z}_n, \tilde{H}_n)$ is continuous as a function of $(\tilde{\xi}_n, \tilde{\zeta}_n)$. From (3.27), one deduces that the distribution of $(\tilde{Z}_n, \tilde{H}_n)$ is equal to the distribution of (Z_n, H_n) on \mathbb{R}_+ , which yields that

$$\tilde{\mathbb{E}}\phi(\tilde{Z}_n) = \mathbb{E}\phi(Z_n), \quad \tilde{\mathbb{E}}\phi(\tilde{H}_n) = \mathbb{E}\phi(H_n), \quad \text{for any } \phi \in C_b(\mathbb{R}_+), \quad (3.28)$$

where $C_b(X)$ is the space of continuous bounded functions defined on X . Now, define $\phi_\varepsilon \in C_b(\mathbb{R}_+)$ by

$$\phi_\varepsilon = \begin{cases} \frac{y}{\varepsilon}, & 0 \leq y < \varepsilon; \\ 1, & y \geq \varepsilon. \end{cases}$$

One can check that

$$\begin{aligned} \tilde{\mathbb{P}}(\tilde{Z}_n \geq \varepsilon) &= \int_{\tilde{\Omega}} 1_{[\varepsilon, \infty]} \tilde{Z}_n d\tilde{\mathbb{P}} \leq \int_{\tilde{\Omega}} 1_{[0, \varepsilon]} \frac{\tilde{Z}_n}{\varepsilon} d\tilde{\mathbb{P}} + \int_{\tilde{\Omega}} 1_{[\varepsilon, \infty]} \tilde{Z}_n d\tilde{\mathbb{P}}, \\ \tilde{\mathbb{P}}(\tilde{H}_n \geq \varepsilon) &= \int_{\tilde{\Omega}} 1_{[\varepsilon, \infty]} \tilde{H}_n d\tilde{\mathbb{P}} \leq \int_{\tilde{\Omega}} 1_{[0, \varepsilon]} \frac{\tilde{H}_n}{\varepsilon} d\tilde{\mathbb{P}} + \int_{\tilde{\Omega}} 1_{[\varepsilon, \infty]} \tilde{H}_n d\tilde{\mathbb{P}}, \end{aligned}$$

Hence by the definition of $\tilde{\mathbb{E}}\phi_\varepsilon(\tilde{Z}_n)$ and $\tilde{\mathbb{E}}\phi_\varepsilon(\tilde{H}_n)$, we can infer that

$$\tilde{\mathbb{P}}(\tilde{Z}_n \geq \varepsilon) \leq \tilde{\mathbb{E}}\phi_\varepsilon(\tilde{Z}_n), \quad \tilde{\mathbb{P}}(\tilde{H}_n \geq \varepsilon) \leq \tilde{\mathbb{E}}\phi_\varepsilon(\tilde{H}_n),$$

which together with (3.28) imply that

$$\tilde{\mathbb{P}}(\tilde{Z}_n \geq \varepsilon) \leq \mathbb{E}\phi_\varepsilon(Z_n), \quad \tilde{\mathbb{P}}(\tilde{H}_n \geq \varepsilon) \leq \mathbb{E}\phi_\varepsilon(H_n).$$

From (3.26), it holds that

$$\tilde{\mathbb{P}}(\tilde{Z}_n \geq \varepsilon) \leq \mathbb{E}\phi_\varepsilon(Z_n) = 0, \quad \tilde{\mathbb{P}}(\tilde{H}_n \geq \varepsilon) \leq \mathbb{E}\phi_\varepsilon(H_n) = 0. \quad (3.29)$$

Since $\varepsilon > 0$ is arbitrary, from (3.29), we can infer that

$$\tilde{Z}_n = 0, \quad \tilde{H}_n = 0 \quad \tilde{\mathbb{P}} - \text{a.s.}$$

Similarly, we can prove that $\tilde{G}_n = 0$, $\tilde{\mathbb{P}}$ - a.s. □

3.4. Passage to the limit for $n \rightarrow \infty$. By Proposition 3.8, we know that $(\tilde{\rho}_n, \tilde{u}_n, \tilde{\rho}_n \tilde{u}_n, \tilde{B}_n)$ satisfies the same estimates as $(\rho_n, u_n, \rho_n u_n, B_n)$. In this subsection, to simplify notations, we will drop the tildes on the random variables. In order to obtain a solution of the problem (3.1)-(3.3), we will use the estimates obtained in Lemma 3.5 to pass to the limit for $n \rightarrow \infty$ in the sequence (ρ_n, u_n, B_n) in the sample space. For this, we need the following proposition and lemma in [22, Chapter 3].

Proposition 3.9 (Uniformly integrability). If there exists a nonnegative measurable function f in \mathbb{R}^+ , such that $\lim_{x \rightarrow \infty} \frac{f(x)}{x} = \infty$ and $\sup_{t \in T} \mathbb{E}[f(|X_t|)] < \infty$. Then X_t is a set of uniformly integrable.

Lemma 3.10 (Vitali's convergence Theorem). Suppose $p \in [1, \infty)$, $X_n \in L^p$ and X_n converges to X in probability. Then the following are equivalent:

- (1) $X_n \xrightarrow{L^p} X$;
- (2) $|X_n|^p$ is uniformly integrable;
- (3) $\mathbb{E}(|X_n|^p) \rightarrow \mathbb{E}(|X|^p)$.

The estimate (3.15) along with the fact that $\partial_t \rho_n$ satisfies (3.1) makes it possible to use (3.25) to infer that there exists a function ρ such that

$$\rho_n \rightarrow \rho \text{ in } L^4((0, T) \times D) \quad \tilde{\mathbb{P}} - \text{a.s.} \quad (3.30)$$

In fact, (3.25) implies that $\rho_n \rightarrow \rho$ in $L^2((0, T) \times D)$ $\tilde{\mathbb{P}}$ -a.s.. Since $\rho_n \in L^\infty(0, T; L^\beta(D))$, $\beta > 4$, using the interpolation of $L^2((0, T) \times D)$ and $L^\infty(0, T; L^\beta(D))$, we have (3.30).

On the other hand, similar to (3.30), in view of (3.15), and $\beta > \gamma$, one has

$$\rho_n^\gamma \rightarrow \rho^\gamma, \quad \rho_n^\beta \rightarrow \rho^\beta \text{ in } L^1((0, T) \times D) \quad \tilde{\mathbb{P}} - \text{a.s.},$$

which together with integration by parts yields that for $\phi \in C_0^\infty([0, T] \times D)$

$$\langle \nabla \rho_n^\gamma, \phi \rangle \rightarrow \langle \nabla \rho^\gamma, \phi \rangle, \quad \langle \nabla \rho_n^\beta, \phi \rangle \rightarrow \langle \nabla \rho^\beta, \phi \rangle \quad \tilde{\mathbb{P}} - \text{a.s. as } n \rightarrow \infty. \quad (3.31)$$

Now, we pass to the limit in the products $\rho_n u_n$ and $\rho_n u_n \otimes u_n$. From (3.25), we know that

$$\rho_n u_n \rightharpoonup h \text{ in } C([0, T]; L^{\frac{2\beta}{\beta+1}}(D)) \quad \tilde{\mathbb{P}} - \text{a.s.}$$

Then, by (3.25) ($\rho_n \rightarrow \rho$ in $L^2(0, T; L^2(D))$, $u_n \rightharpoonup u$ in $L^2(0, T; H^1(D))$) $\tilde{\mathbb{P}}$ - a.s.), one has

$$\rho_n u_n \rightharpoonup \rho u \text{ in } L^1(0, T; L^1(D)) \quad \tilde{\mathbb{P}} - \text{a.s.}$$

which together with [15, Lemma 2.4] yields that $h = \rho u$. Then $\rho_n u_n \rightharpoonup \rho u$ in $C([0, T]; L^{\frac{2\beta}{\beta+1}}(D))$ $\tilde{\mathbb{P}}$ -a.s.. Since $L^{\frac{2\beta}{\beta+1}}(D) \hookrightarrow H^{-1}(D)$, by Aubin-Lions Lemma, one has

$$\rho_n u_n \rightarrow \rho u \text{ in } C([0, T]; H^{-1}(D)) \quad \tilde{\mathbb{P}} - \text{a.s.} \quad (3.32)$$

Therefore, from (3.25) ($u_n \rightharpoonup u$ in $L^2(0, T; H^1(D))$) $\tilde{\mathbb{P}}$ - a.s.) and (3.32), we can deduce that

$$\rho_n u_n \otimes u_n \rightarrow \rho u \otimes u \text{ in } \mathcal{D}'([0, T] \times D) \quad \tilde{\mathbb{P}} - \text{a.s.} \quad (3.33)$$

Now, we turn to the magnetic terms. By (3.25), we know that

$$B_n \rightarrow B \text{ in } L^2(0, T; L^2(D)) \text{ and } B_n \rightharpoonup B \text{ in } L^2(0, T; H^1(D)) \quad \tilde{\mathbb{P}} - \text{a.s.} \quad (3.34)$$

Moreover, from (3.1)₃ and (3.15), by Aubin-Lions Lemma, one has

$$B_n \rightarrow B \text{ in } C([0, T]; H^{-1}(D)) \quad \tilde{\mathbb{P}} - \text{a.s.}$$

Then we have

$$(\nabla \times B_n) \times B_n \rightarrow (\nabla \times B) \times B \text{ in } \mathcal{D}'([0, T] \times D) \quad \tilde{\mathbb{P}} - \text{a.s.} \quad (3.35)$$

Similarly, it follows from (3.25) ($u_n \rightharpoonup u$ in $L^2(0, T; H^1(D))$) $\tilde{\mathbb{P}} - \text{a.s.}$) and (3.34) that

$$\nabla \times (u_n \times B_n) \rightarrow \nabla \times (u \times B) \text{ in } \mathcal{D}'([0, T] \times D) \quad \tilde{\mathbb{P}} - \text{a.s.} \quad (3.36)$$

Finally, we will treat the force terms. We claim that

$$\begin{aligned} \langle f_k^n(\rho_n, \rho_n u_n, x), \mathbb{P}\phi \rangle &\rightarrow \langle f_k(\rho, \rho u, x), \phi \rangle \text{ in } L^1([0, T]) \quad \tilde{\mathbb{P}} - \text{a.s.}, \\ \left\langle \sum_{k \geq 1} f_k^n(\rho_n, \rho_n u_n, x), \mathbb{P}\phi \right\rangle^2 &\rightarrow \left\langle \sum_{k \geq 1} f_k(\rho, \rho u, x), \phi \right\rangle^2 \text{ in } L^1([0, T]) \quad \tilde{\mathbb{P}} - \text{a.s.} \end{aligned} \quad (3.37)$$

Proof of the claim. By definition and the symmetry of $\mathcal{M}[\rho_n]$ we have

$$\langle f_k^n(\rho_n, \rho_n u_n, x), \phi \rangle = \left\langle \mathcal{M}^{\frac{1}{2}}[\rho_n] \mathbb{P} \left(\frac{f_k(\rho_n, \rho_n u_n, x)}{\sqrt{\rho_n}} \right), \mathbb{P}\phi \right\rangle = \left\langle \mathbb{P} \left(\frac{f_k(\rho_n, \rho_n u_n, x)}{\sqrt{\rho_n}} \right), \mathcal{M}^{\frac{1}{2}}[\rho_n] \mathbb{P}\phi \right\rangle.$$

By using (1.5), (3.15) and the strong convergence in (3.25), (3.30), (3.32), we have

$$\frac{f_k(\rho_n, \rho_n u_n, x)}{\sqrt{\rho_n}} \rightarrow \frac{f_k(\rho, \rho u, x)}{\sqrt{\rho}} \text{ in } L^2(D) \quad \tilde{\mathbb{P}} \otimes \tilde{\mathbb{L}} - \text{a.e.}, \quad (3.38)$$

where $\tilde{\mathbb{L}}$ is the Lebesgue measure in time.

Next we extend the operator $\mathcal{M}[\rho]$ to

$$\tilde{\mathcal{M}}[\rho] : H^1(D) \rightarrow L^2(D), \quad \langle \tilde{\mathcal{M}}[\rho]v, w \rangle = \langle \mathcal{M}[\rho]\mathbb{P}v, \mathbb{P}w \rangle$$

for $v \in H^1(D)$, $w \in L^2(D)$. It is easy to see that as $n \rightarrow \infty$,

$$\begin{aligned} \left| \langle \tilde{\mathcal{M}}[\rho_n]v, w \rangle - \langle \rho v, w \rangle \right| &\leq \|\rho_n\|_{L^\infty} \|v - \mathbb{P}v\|_{L^2} \|w - \mathbb{P}w\|_{L^2} + \|\rho_n\|_{L^\infty} \|v\|_{L^2} \|w - \mathbb{P}w\|_{L^2} \\ &\quad + \|\rho_n - \rho\|_{L^4} \|v\|_{L^4} \|w\|_{L^2} + \|\rho_n\|_{L^\infty} \|w\|_{L^2} \|v - \mathbb{P}v\|_{L^2} \rightarrow 0, \end{aligned}$$

which implies that

$$\langle \tilde{\mathcal{M}}[\rho_n]v, \cdot \rangle \rightarrow \langle \rho v, \cdot \rangle_2 \quad \tilde{\mathbb{P}} \otimes \tilde{\mathbb{L}} - \text{a.e.},$$

Next we look at $\tilde{\mathcal{M}}^{\frac{1}{2}}[\rho_n]$. Note that $\tilde{\mathcal{M}}\rho_n$ and ρ are both symmetric positive definite operators with a lower bound (independent of n) strictly away from 0, and hence they are both invertible. Moreover we know that they also commute. This way we have

$$\tilde{\mathcal{M}}^{\frac{1}{2}}[\rho_n] - \sqrt{\rho} = \left(\tilde{\mathcal{M}}^{\frac{1}{2}}[\rho_n] + \sqrt{\rho} \right)^{-1} \left(\tilde{\mathcal{M}}[\rho_n] - \rho \right),$$

and hence

$$\langle \tilde{\mathcal{M}}^{\frac{1}{2}}[\rho_n]v, \cdot \rangle \rightarrow \langle \sqrt{\rho}v, \cdot \rangle_2 \quad \tilde{\mathbb{P}} \otimes \tilde{\mathbb{L}} - \text{a.e.} \quad (3.39)$$

This way (3.37)₁ follows from (3.38) and (3.39).

Next, we prove (3.37)₂. Direct computation yields that

$$\begin{aligned} &\left| \sum_{k \geq 1} \langle f_k^n(\rho_n, \rho_n u_n, x), \mathbb{P}\phi \rangle^2 - \sum_{k \geq 1} \langle f_k(\rho_n, \rho_n u_n, x), \phi \rangle^2 \right| \\ &\leq \sum_{k \geq 1} \left| \langle f_k^n(\rho_n, \rho_n u_n, x), \mathbb{P}\phi \rangle - \langle f_k(\rho_n, \rho_n u_n, x), \phi \rangle \right| \left| \langle f_k^n(\rho_n, \rho_n u_n, x), \mathbb{P}\phi \rangle + \langle f_k(\rho_n, \rho_n u_n, x), \phi \rangle \right|. \end{aligned}$$

By (1.5) and Lemma 3.5, we can bound

$$\sum_{k \geq 1} \left| \langle f_k(\rho_n, \rho_n u_n, x), \phi \rangle \right| \lesssim \sum_{k \geq 1} \|f_k\|_{L^2} \lesssim \|\rho_n\|_{L^{\gamma+1}}^{(\gamma+1)/2} + \|\rho_n u_n\|_{L^2} \leq C.$$

Similarly we can bound $\sum_{k \geq 1} \left| \langle f_k^n(\rho_n, \rho_n u_n, x), \mathbb{P}\phi \rangle \right|$. Then from (3.37)₁ we get (3.37)₂. \square

Thanks to (3.34) and (1.5), we can similarly obtain

$$\begin{aligned} \langle g_k(B_n, x), \phi \rangle &\rightarrow \langle g_k(B, x), \phi \rangle \quad \text{in } L^1([0, T]) \quad \tilde{\mathbb{P}} - \text{a.s.}, \\ \left\langle \sum_{k \geq 1} g_k(B_n, x), \phi \right\rangle^2 &\rightarrow \left\langle \sum_{k \geq 1} g_k(B, x), \phi \right\rangle^2 \quad \text{in } L^1([0, T]) \quad \tilde{\mathbb{P}} - \text{a.s.} \end{aligned} \quad (3.40)$$

Consequently, by (3.25), (3.30)-(3.36), Lemma 3.5, Proposition 3.9 and Lemma 3.10, we can pass to the limit in the continuity equation (3.1)₁ and have the following propositions.

Proposition 3.11. $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ is a weak solution of (3.1)₁. Moreover, $\nabla \rho_n \cdot \nabla u_n \rightarrow \nabla \rho \cdot \nabla u$ in $\mathcal{D}'((0, T) \times D)$, $\tilde{\mathbb{P}}$ -a.s..

Proof. For all $t \in [0, T]$ and any $\varphi \in C_0^\infty(D)$, denote

$$m_n(t) = \langle \rho_n(t), \varphi \rangle - \langle \rho_0, \varphi \rangle + \int_0^t \langle \rho_n u_n, \nabla \varphi \rangle ds - \int_0^t \langle \varepsilon \nabla \rho_n, \nabla \varphi \rangle ds.$$

By Lemma 3.5, we know that $\sup_{0 \leq t \leq T} \tilde{\mathbb{E}}(|m_n(t)|^2) \leq C$. Then $m_n(t)$ is uniformly integrable. By (3.25), (3.32), and Lemma 3.5 we know that $m_n(t) \rightarrow m(t)$ $\tilde{\mathbb{P}}$ -a.s.. Hence applying Proposition 3.9 and Lemma 3.10, we have $\tilde{\mathbb{E}}(|m(t)|^2) = \lim_{n \rightarrow \infty} \tilde{\mathbb{E}}(|m_n(t)|^2) = 0$.

Next, we shall show the convergence of the term $\nabla \rho_n \cdot \nabla u_n$. Multiplying (3.1)₁ by ρ_n , integrating by parts, we obtain $\tilde{\mathbb{P}}$ -a.s.

$$\|\rho_n(t)\|_{L^2(D)}^2 + 2\varepsilon \int_0^t \|\nabla \rho_n\|_{L^2(D)}^2 dt = - \int_0^t \int_D \rho_n^2 \operatorname{div} u_n dx dt + \|\rho_0\|_{L^2(D)}^2. \quad (3.41)$$

Sending $n \rightarrow \infty$ in equation (3.1)₁ and then multiplying the result by ρ (since from [15, Lemma 2.4] we know ρ enjoys enough regularity due to parabolic smoothing), one has $\tilde{\mathbb{P}}$ -a.s.,

$$\|\rho(t)\|_{L^2(D)}^2 + 2\varepsilon \int_0^t \|\nabla \rho\|_{L^2(D)}^2 dt = - \int_0^t \int_D \rho^2 \operatorname{div} u dx dt + \|\rho_0\|_{L^2(D)}^2. \quad (3.42)$$

From [15, Lemma 2.4], (3.25), (3.30), (3.41), and (3.42), for any t , we deduce that $\tilde{\mathbb{P}}$ -a.s.

$$\|\nabla \rho_n\|_{L^2((0, T) \times D)}^2 \rightarrow \|\nabla \rho\|_{L^2((0, T) \times D)}^2, \quad \|\rho_n(t)\|_{L^2(D)}^2 \rightarrow \|\rho(t)\|_{L^2(D)}^2,$$

which together with (3.25) imply $\nabla \rho_n \rightarrow \nabla \rho$ in $L^2(0, T; L^2(D))$ $\tilde{\mathbb{P}}$ -a.s.. Then, by the fact $u_n \rightarrow u \in L^2(0, T; H^1(D))$ $\tilde{\mathbb{P}}$ -a.s.((3.25)), one has

$$\nabla \rho_n \cdot \nabla u_n^i \rightarrow \nabla \rho \cdot \nabla u^i \quad \text{in } \mathcal{D}'((0, T) \times D) \quad \tilde{\mathbb{P}} - \text{a.s.}, \quad i = 1, 2, 3.$$

\square

Proposition 3.12. The system $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ is a martingale solution of (3.1)₂ and (3.1)₃.

Proof. It follows from Proposition 3.8 that

$$\begin{aligned}
\langle M_n(t), \phi \rangle &= \langle \rho_n u_n(t) - \rho_n u_n(0), \phi \rangle + \int_0^t \langle \operatorname{div}(\rho_n u_n \otimes u_n) + \delta \nabla \rho_n^\beta - \mu \Delta u_n, \phi \rangle ds \\
&\quad + \int_0^t \langle (\nabla \times B_n) \times B_n + \varepsilon \nabla \rho_n \cdot \nabla u_n - (\lambda + \mu) \nabla \operatorname{div} u_n + a \nabla \rho_n^\gamma, \phi \rangle ds \\
&:= \langle \rho_n u_n(t) - \rho_n u_n(0), \phi \rangle + \int_0^t \langle \Lambda_n, \phi \rangle ds, \\
\langle \tilde{M}_n(t), \phi \rangle &= \langle B_n(t) - B_n(0), \phi \rangle + \int_0^t \langle \nabla \times (u_n \times B_n) + \nu \Delta B_n, \phi \rangle ds \\
&:= \langle B_n(t) - B_n(0), \phi \rangle + \int_0^t \langle \tilde{\Lambda}_n, \phi \rangle ds,
\end{aligned}$$

where $\langle M_n(t), \phi \rangle = \int_0^t \sum_{k \geq 1} \langle f_k(\rho_n, \rho_n u_n, x), \phi \rangle d\beta_{k,n}^1$ and $\langle \tilde{M}_n(t), \phi \rangle = \int_0^t \sum_{k \geq 1} \langle g_k(B_n, x), \phi \rangle d\beta_{k,n}^2$ are martingales under $\tilde{\mathbb{P}}$. Therefore, with $0 \leq s < t < \infty$ and a bounded, continuous \mathcal{F}_s -measurable function $\varphi_n = \varphi(\beta_{k,n}, \rho_n, u_n, B_n)$, we have

$$\begin{aligned}
\tilde{\mathbb{E}}[\langle M_n(t) - M_n(s), \phi \rangle \varphi_n] &= 0, \quad \tilde{\mathbb{E}}[\langle \tilde{M}_n(t) - \tilde{M}_n(s), \phi \rangle \varphi_n] = 0, \\
\tilde{\mathbb{E}}\left[\left(\langle M_n(t) - M_n(s), \phi \rangle \beta_{k,n}^1 - \int_s^t \langle f_k^n(\rho_n, \rho_n u_n, x), \phi \rangle d\tau\right) \varphi_n\right] &= 0, \\
\tilde{\mathbb{E}}\left[\left(\langle \tilde{M}_n(t) - \tilde{M}_n(s), \phi \rangle \beta_{k,n}^2 - \int_s^t \langle g_k(B_n, x), \phi \rangle d\tau\right) \varphi_n\right] &= 0, \\
\tilde{\mathbb{E}}\left[\left(\langle M_n(t), \phi \rangle^2 - \langle M_n(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle f_k^n(\rho_n, \rho_n u_n, x), \phi \rangle^2 d\tau\right) \varphi_n\right] &= 0, \\
\tilde{\mathbb{E}}\left[\left(\langle \tilde{M}_n(t), \phi \rangle^2 - \langle \tilde{M}_n(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle g_k(B_n, x), \phi \rangle^2 d\tau\right) \varphi_n\right] &= 0.
\end{aligned} \tag{3.43}$$

By (3.25), (3.30)-(3.36), one deduces that

$$\langle \rho_n u_n(t) - \rho_n u_n(s), \phi \rangle + \int_s^t \langle \Lambda_n, \phi \rangle d\tau, \quad \langle B_n(t) - B_n(s), \phi \rangle + \int_s^t \langle \tilde{\Lambda}_n, \phi \rangle d\tau$$

converges to

$$\langle \rho u(t) - \rho u(s), \phi \rangle - \int_s^t \langle \Lambda, \phi \rangle d\tau, \quad \langle B(t) - B(s), \phi \rangle + \int_s^t \langle \tilde{\Lambda}, \phi \rangle d\tau \quad \text{almost surely on } \tilde{\Omega},$$

where

$$\begin{aligned}
\langle \Lambda, \phi \rangle &= \langle \operatorname{div}(\rho u \otimes u) - \mu \Delta u - (\lambda + \mu) \operatorname{div} \nabla u + a \nabla \rho^\gamma + \delta \nabla \rho^\beta + \varepsilon \nabla u \cdot \nabla \rho + (\nabla \times B) \times B, \phi \rangle, \\
\langle \tilde{\Lambda}, \phi \rangle &= \langle \nabla \times (u \times B) + \nu \Delta B, \phi \rangle.
\end{aligned}$$

Next, we prove that the functions $\overline{f_n} := \langle M_n(t) - M_n(s), \phi \rangle \varphi(\beta_{k,n}, \rho_n, u_n, B_n)$ are uniformly integrable. Indeed, by the Schwarz inequality and the Burkholder-Davis-Gundy inequality, for each n , we have

$$\tilde{\mathbb{E}}(|\overline{f_n}|^2) \lesssim \tilde{\mathbb{E}}\left(\|M_n(t)\|_{W^{-2,2}(D)}^2 + \|M_n(s)\|_{W^{-2,2}(D)}^2\right)$$

$$\begin{aligned}
&\lesssim \tilde{\mathbb{E}} \left(\int_0^T \sum_{k=1}^n \|f_k^n(\rho_n, \rho_n u_n, x)\|_{L^1(D)}^2 dt \right) \\
&\lesssim \tilde{\mathbb{E}} \left(\int_0^T \|\rho_n\|_{L^1(D)} (\|\rho_n\|_{L^\gamma(D)}^\gamma + \|\sqrt{\rho_n} u_n\|_{L^2(D)}^2) dt \right) \leq C.
\end{aligned} \tag{3.44}$$

Then the sequence $\overline{f_n}$ is uniformly integrable. For some $r > 1$, we obtain as (3.44)

$$\tilde{\mathbb{E}} \left(\int_0^T \sum_{k=1}^n \|f_k^n(\rho_n, \rho_n u_n, x)\|_{L^1(D)}^2 dt \right)^r \lesssim C. \tag{3.45}$$

Similarly, we have

$$\tilde{\mathbb{E}} \left(\int_0^T \sum_{k=1}^n \|g_k(B_n, x)\|_{H^{-1}(D)}^2 dt \right)^r \leq C. \tag{3.46}$$

In view of (3.37), (3.40), (3.43), (3.45)-(3.46), together with Proposition 3.9 and Lemma 3.10, one has

$$\begin{aligned}
&\tilde{\mathbb{E}} [\langle M(t) - M(s), \phi \rangle \varphi(\beta_k, \rho, u, B)] = \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} [\langle M_n(t) - M_n(s), \phi \rangle \varphi_n] = 0, \\
&\tilde{\mathbb{E}} [\langle \tilde{M}(t) - \tilde{M}(s), \phi \rangle \varphi(\beta_k, \rho, u, B)] = \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} [\langle \tilde{M}_n(t) - \tilde{M}_n(s), \phi \rangle \varphi_n] = 0, \\
&\tilde{\mathbb{E}} \left[\left(\langle M(t) - M(s), \phi \rangle \beta_k^1 - \int_s^t \langle f_k(\rho, \rho u, x), \phi \rangle d\tau \right) \varphi(\beta_k, \rho, u, B) \right] \\
&= \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} \left[\left(\langle M_n(t) - M_n(s), \phi \rangle \beta_{k,n}^1 - \int_s^t \langle f_k^n(\rho_n, \rho_n u_n, x), \phi \rangle d\tau \right) \varphi_n \right] = 0, \\
&\tilde{\mathbb{E}} \left[\left(\langle \tilde{M}(t) - \tilde{M}(s), \phi \rangle \beta_k^2 - \int_s^t \langle g_k(B, x), \phi \rangle d\tau \right) \varphi(\beta_k, \rho, u, B) \right] \\
&= \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} \left[\left(\langle \tilde{M}_n(t) - \tilde{M}_n(s), \phi \rangle \beta_{k,n}^2 - \int_s^t \langle g_k(B_n, x), \phi \rangle d\tau \right) \varphi_n \right] = 0,
\end{aligned}$$

and

$$\begin{aligned}
&\tilde{\mathbb{E}} \left[\left(\langle M(t), \phi \rangle^2 - \langle M(s), \phi \rangle^2 - \int_s^t \sum_{k=1}^n \langle f_k(\rho, \rho u, x), \phi \rangle^2 d\tau \right) \varphi(\beta_k, \rho, u, B) \right] \\
&= \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} \left[\left(\langle M_n(t), \phi \rangle^2 - \langle M_n(s), \phi \rangle^2 - \int_s^t \sum_{k=1}^n \langle f_k^n(\rho_n, \rho_n u_n, x), \phi \rangle^2 d\tau \right) \varphi_n \right] = 0, \\
&\tilde{\mathbb{E}} \left[\left(\langle \tilde{M}(t), \phi \rangle^2 - \langle \tilde{M}(s), \phi \rangle^2 - \int_s^t \sum_{k=1}^n \langle g_k(B, x), \phi \rangle^2 d\tau \right) \varphi(\beta_k, \rho, u, B) \right] \\
&= \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} \left[\left(\langle \tilde{M}_n(t), \phi \rangle^2 - \langle \tilde{M}_n(s), \phi \rangle^2 - \int_s^t \sum_{k=1}^n \langle g_k(B_n, x), \phi \rangle^2 d\tau \right) \varphi_n \right] = 0.
\end{aligned}$$

Then we can deduce that $\langle M(t), \phi \rangle$, $\langle M(t) - M(s), \phi \rangle \beta_k^1 - \int_s^t \langle f_k(\rho, \rho u, x), \phi \rangle d\tau$, $\langle M(t), \phi \rangle^2 - \int_0^t \sum_{k=1}^n \langle f_k(\rho, \rho u, x), \phi \rangle^2 ds$; $\langle \tilde{M}(t), \phi \rangle$, $\langle \tilde{M}(t) - \tilde{M}(s), \phi \rangle \beta_k^2 - \int_s^t \langle g_k(B, x), \phi \rangle d\tau$ and $\langle \tilde{M}(t), \phi \rangle^2 - \int_0^t \sum_{k=1}^n \langle g_k(B, x), \phi \rangle^2 ds$ are martingales.

Applying the method in [3, 28], we can infer that

$$\langle M(t), \phi \rangle = \int_0^t \langle f_k(\rho, \rho u, x), \phi \rangle d\beta_k^1, \quad \langle \tilde{M}(t), \phi \rangle = \int_0^t \langle g_k(B, x), \phi \rangle d\beta_k^2.$$

Therefore, $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ is a martingale solution of (3.1)-(3.3). \square

4. THE VANISHING VISCOSITY LIMIT

In this section, we shall consider the limit of (3.1)-(3.3) as $\varepsilon \rightarrow 0$ following the idea of [13, 15, 30]. Similar to Lemma 3.5, we can get the following estimates:

$$\begin{aligned} \rho_\varepsilon &\in L^p(\Omega, L^\infty(0, T; L^\gamma(D))) \cap L^p(\Omega, L^\infty(0, T; L^\beta(D))), \\ \sqrt{\rho_\varepsilon} u_\varepsilon &\in L^p(\Omega, L^\infty(0, T; L^2(D))), \quad \rho_\varepsilon u_\varepsilon \in L^p(\Omega, L^\infty(0, T; L^{\frac{2\beta}{\beta+1}}(D))), \\ u_\varepsilon &\in L^p(\Omega, L^2(0, T; H^1(D))), \\ \sqrt{\varepsilon} \nabla \rho_\varepsilon &\in L^p(\Omega, L^2(0, T; L^2(D))), \quad \rho_\varepsilon u_\varepsilon^i u_\varepsilon^j \in L^p(\Omega, L^2(0, T; L^{\frac{6\beta}{4\beta+3}}(D))), \\ B_\varepsilon &\in L^p(\Omega, L^2(0, T; H^1(D))) \cap L^p(\Omega, L^\infty(0, T; L^2(D))), \end{aligned} \tag{4.1}$$

for $1 \leq p < \infty$, $\beta > \max\{4, \gamma\}$. It follows from (4.1) that $\rho_\varepsilon \in L^\infty(0, T; L^\beta(D))$, i.e., $\rho_\varepsilon^\beta \in L^\infty(0, T; L^1(D))$, and hence we can only conclude that ρ_ε^β converges to a Radon measure. To gain some better convergence of this term we need to improve the integrability of the density ρ as follows.

4.1. On the equation $\operatorname{div} v = f$. We introduce a linear operator

$$\mathcal{B} : \left\{ f \in L^p(D) : \int_D f dx = 0 \right\} \rightarrow [W_0^{1,p}(D)]^3,$$

which is a bounded linear operator, that is,

$$\|\mathcal{B}[f]\|_{W^{1,p}(D)} \leq C(p) \|f\|_{L^p(D)}, \quad 1 < p < \infty. \tag{4.2}$$

and $\mathcal{B}(f)$ solves

$$\operatorname{div} \mathcal{B}(f) = f \text{ in } D, \quad \mathcal{B}(f)|_{\partial D} = 0.$$

Moreover, if $g \in L^r(D)$ and $g \cdot \mathbf{n}|_{\partial D} = 0$, then

$$\|\mathcal{B}[\operatorname{div} g]\|_{L^r(D)} \leq C(p) \|g\|_{L^p(D)} \text{ for } 1 < p < \infty.$$

The operator \mathcal{B} was first constructed by Bogovskii [5]. The existence and the above properties of the operator \mathcal{B} have been proved in many papers, and we refer the reader to [16] and the references therein for details.

4.2. Uniform (on viscosity) estimates of density. Next, we shall use the operator \mathcal{B} to improve the integrability of the density. First, let $\psi \in \mathcal{D}(0, T)$, $0 \leq \psi \leq 1$ and $\bar{m} = \frac{1}{|D|} \int_D \rho(t) dx$, we note that \bar{m} is conserved. Taking $\psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}]$ as a test function for (3.1)₂, integrating over $\Omega \times (0, T) \times D$, we arrive at

$$\begin{aligned} \mathbb{E} \int_0^T \int_D [\psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}] (\rho_\varepsilon u_\varepsilon)_t + \psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}] \operatorname{div}(\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon)] dx dt \\ + a \mathbb{E} \int_0^T \int_D \psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}] \nabla \rho_\varepsilon^\gamma dx dt + \delta \mathbb{E} \int_0^T \int_D \psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}] \nabla \rho_\varepsilon^\beta dx dt \end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \int_0^T \int_D \psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}] [\varepsilon \nabla \rho_\varepsilon \cdot \nabla u_\varepsilon - (\nabla \times B_\varepsilon) \times B_\varepsilon] dx dt \\
& = \mathbb{E} \int_0^T \int_D [\mu \psi(t) \mathcal{B}_i[\rho_\varepsilon - \bar{m}] \Delta u_\varepsilon + (\lambda + \mu) \psi(t) \mathcal{B}[\rho_\varepsilon - \bar{m}] \nabla \operatorname{div} u_\varepsilon] dx dt \\
& + \mathbb{E} \left(\int_0^T \int_D \psi(s) \mathcal{B}[\rho_\varepsilon - \bar{m}] \sum_{k=1} f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x) dx d\beta_{k,\varepsilon}^1 \right).
\end{aligned}$$

For convenience, we write the above equation as

$$I_1 + I_2 + I_3 + I_4 + I_5 + I_6 = J_1 + J_2 + J_3.$$

We will just discuss the term I_6 . Other terms can be handled as in [34, 38]. For the term I_6 , using Hölder's inequality and the fact that $\beta > 4$, from (4.1) and (4.2), we obtain

$$\begin{aligned}
I_6 & = -\mathbb{E} \int_0^T \psi \int_D \mathcal{B}[\rho_\varepsilon - \bar{m}] (\nabla \times B_\varepsilon) \times B_\varepsilon dx dt \\
& \lesssim \mathbb{E} \left(\|B_\varepsilon\|_{L^2(0,T;L^2(D))} \|\nabla B_\varepsilon\|_{L^2(0,T;L^2(D))} \|\mathcal{B}[\rho_\varepsilon - \bar{m}]\|_{L^\infty(0,T;L^\infty(D))} \right) \\
& \lesssim \mathbb{E} \left(\|\nabla B_\varepsilon\|_{L^2(0,T;L^2(D))}^2 \|\rho_\varepsilon\|_{L^\infty(0,T;L^\beta(D))} \right) \\
& \lesssim \left(\mathbb{E} \|\nabla B_\varepsilon\|_{L^2(0,T;L^2(D))}^4 \right)^{\frac{1}{2}} \left(\mathbb{E} \|\rho_\varepsilon\|_{L^\infty(0,T;L^\beta(D))}^2 \right)^{\frac{1}{2}} \leq C.
\end{aligned}$$

Summing up the estimates for I_1, I_2, \dots, I_6 and J_1, J_2, J_3 , we obtain the following results:

Lemma 4.1. Let $(\beta_{k,\varepsilon}, \rho_\varepsilon, u_\varepsilon, B_\varepsilon)$ be a weak solution of the problem (3.1)-(3.3). Then

$$\mathbb{E} \left(\|\rho_\varepsilon\|_{L^{\gamma+1}((0,T) \times D)} + \|\rho_\varepsilon\|_{L^{\beta+1}((0,T) \times D)} \right) \leq C, \quad (4.3)$$

where $C = C(\delta, \rho_{0,\delta}, m_{0,\delta}, B_{0,\delta})$ is a constant independent of ε .

4.3. Tightness property.

Lemma 4.2. Define

$$\begin{aligned}
S & = C(0, T; \mathbb{R}) \times C([0, T]; L_w^\beta(D)) \times L^2(0, T; H_w^1(D)) \\
& \quad \times C([0, T]; L_w^{\frac{2\beta}{\beta+1}}(D)) \times (L^2(0, T; H_w^1(D)) \cap L^2(0, T; L^2(D)))
\end{aligned}$$

equipped with its Borel σ -algebra. Let Π_ε be the probability on S which is the image of \mathbb{P} on Ω by the map: $\omega \mapsto (\beta_{k,\varepsilon}(\omega, \cdot), \rho_\varepsilon(\omega, \cdot), u_\varepsilon(\omega, \cdot), \rho_\varepsilon u_\varepsilon(\omega, \cdot), B_\varepsilon(\omega, \cdot))$, that is, for any $A \subseteq S$,

$$\Pi_\varepsilon(A) = \mathbb{P} \{ \omega \in \Omega : (\beta_{k,\varepsilon}(\omega, \cdot), \rho_\varepsilon(\omega, \cdot), u_\varepsilon(\omega, \cdot), \rho_\varepsilon u_\varepsilon(\omega, \cdot), B_\varepsilon(\omega, \cdot)) \in A \}.$$

Then the family Π_ε is tight.

Proof. Let Π_ε^1 be the probability on

$$S_1 = C(0, T; \mathbb{R}) \times C([0, T]; L_w^\beta(D)) \times L^2(0, T; H_w^1(D)) \times (L^2(0, T; H_w^1(D)) \cap L^2(0, T; L^2(D)))$$

which is the image of \mathbb{P} on Ω by the map $\omega \mapsto (\beta_{k,\varepsilon}(\omega, \cdot), \rho_\varepsilon(\omega, \cdot), u_\varepsilon(\omega, \cdot), B_\varepsilon(\omega, \cdot))$. Similar to Section 3.2, we can obtain the tightness of the family of Π_ε^1 :

$$\Pi_\varepsilon^1(\Sigma_\eta \times X_\eta \times Y_\eta \times R_\eta) \geq \left(1 - \frac{\eta}{5}\right)^4. \quad (4.4)$$

Hence we only need to check the tightness of Π_ε^2 , which is defined as: for any $A \subseteq S_2 = C([0, T]; L_w^{\frac{2\beta}{\beta+1}}(D))$,

$$\Pi_\varepsilon^2(A) = \mathbb{P}(\omega \in \Omega : \rho_\varepsilon u_\varepsilon(\omega, \cdot) \in A).$$

From (3.1), we have

$$\begin{aligned} \rho_\varepsilon u_\varepsilon(t) = & \rho_0 u_0 - \int_0^t \left[\operatorname{div}(\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon) - \mu \Delta u_\varepsilon - (\lambda + \mu) \operatorname{div} \nabla u_\varepsilon + a \nabla \rho_\varepsilon^\gamma + \delta \nabla \rho_\varepsilon^\beta \right] ds \\ & - \int_0^t [\varepsilon \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon - (\nabla \times B_\varepsilon) \times B_\varepsilon] ds + \int_0^t f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x) d\beta_{k,\varepsilon}^1. \end{aligned}$$

For $\phi \in C_0^\infty(D)$, denote $\Gamma_\varepsilon := \langle \rho_\varepsilon u_\varepsilon, \phi \rangle + \varepsilon \int_0^t \langle \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon, \phi \rangle ds$. By virtue of (4.1), for some $\alpha > 1$, one has P-a.s.

$$\begin{aligned} \partial_t \Gamma_\varepsilon = & \langle \partial_t(\rho_\varepsilon u_\varepsilon) + \varepsilon \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon, \phi \rangle \\ = & \langle -\operatorname{div}(\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon) - a \nabla \rho_\varepsilon^\gamma - \delta \nabla \rho_\varepsilon^\beta + \mu \Delta u_\varepsilon + (\lambda + \mu) \nabla \operatorname{div} u_\varepsilon, \phi \rangle \\ & + \langle (\nabla \times B_\varepsilon) \times B_\varepsilon, \phi \rangle \in L^\alpha(0, T). \end{aligned}$$

On the other hand, Γ_ε is uniformly bounded in $L^\alpha(0, T)$. Hence Γ_ε is equi-continuous in $C[0, T]$. We can infer that $\int_0^T \varepsilon \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon dt \rightarrow 0$ in $C([0, T]; L^1(D))$ almost surely on Ω . Since $L^1(D) \hookrightarrow H^{-\ell}(D)$ ($\ell \geq 3$) is compact, by [22, Lemma 3.9], the distribution of $\int_0^T \varepsilon \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon dt$ is tight on $H^{-\ell}(D)$. That is, for any $\eta > 0$, there exists a compact set $K \subset C([0, T]; H^{-\ell}(D))$ such that $\mathbb{P}(\omega \in \Omega : \int_0^T \varepsilon \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon dt \notin K) \leq \frac{\eta}{10}$. Set

$$\mathcal{H} = L^\infty([0, T]; L_w^{\frac{2\beta}{\beta+1}}(D)) \cap C^{0,\alpha'}([0, T]; H^{-\ell}(D)), \quad \ell \geq 3, \quad 0 < \alpha' < \frac{1}{2} - \frac{1}{2\sigma},$$

with the norm $\|f\|_{\mathcal{H}} = \sup_{0 \leq t \leq T} \|f(t)\|_{L_w^{\frac{2\beta}{\beta+1}}(D)} + \|f(t)\|_{C^{0,\alpha'}([0, T]; H^{-\ell}(D))}$.

We choose Z_η as a closed ball of radius r_η centered at 0 in $\mathcal{H} \cap K$. Similar to Section 3.2, choosing $r_\eta = 10C\eta^{-1}$, we have

$$\begin{aligned} \Pi_\varepsilon^2(Z_\eta) &= 1 - \Pi_\varepsilon^2(Z_\eta^c) = 1 - \mathbb{P}(\omega \in \Omega : \rho_\varepsilon u_\varepsilon(\omega, \cdot) \in Z_\eta^c) \\ &= 1 - \mathbb{P}(\omega \in \Omega : \rho_\varepsilon u_\varepsilon(\omega, \cdot) \in \mathcal{H}^c \cup K^c) \\ &\geq 1 - \mathbb{P}(\omega \in \Omega : \rho_\varepsilon u_\varepsilon(\omega, \cdot) \notin \mathcal{H}) - \mathbb{P}\left(\omega \in \Omega : \int_0^T \varepsilon \nabla \rho_\varepsilon \cdot \nabla u_\varepsilon(\omega, \cdot) dt \notin K\right) \quad (4.5) \\ &\geq 1 - \frac{1}{r_\eta} \mathbb{E}(\|\rho_\varepsilon u_\varepsilon\|_{\mathcal{H}}) - \frac{\eta}{10} \\ &\geq 1 - \frac{C}{r_\eta} - \frac{\eta}{10} = 1 - \frac{\eta}{5}, \end{aligned}$$

where Z^c is the complement of Z in $\mathcal{H} \cap K$. Therefore, (4.4) and (4.5) imply that

$$\Pi_\varepsilon(\Sigma_\eta \times X_\eta \times Y_\eta \times Z_\eta \times R_\eta) = \Pi_\varepsilon^1(\Sigma_\eta \times X_\eta \times Y_\eta \times R_\eta) \times \Pi_\varepsilon^2(Z_\eta) \geq 1 - \eta.$$

Then Π_ε is tight. \square

According to Jakubowski-Skorohod Theorem, there exists a subsequence such that $\Pi_\varepsilon \rightarrow \Pi$ weakly, where Π is a probability on S . Moreover, there exist a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$

and two random variables $(\tilde{\beta}_{k,\varepsilon}, \tilde{\rho}_\varepsilon, \tilde{u}_\varepsilon, \tilde{\rho}_\varepsilon \tilde{u}_\varepsilon, \tilde{B}_\varepsilon)$ with distribution Π_ε , (β_k, ρ, u, h, B) with values in S such that as $\varepsilon \rightarrow 0$

$$(\tilde{\beta}_{k,\varepsilon}, \tilde{\rho}_\varepsilon, \tilde{u}_\varepsilon, \tilde{\rho}_\varepsilon \tilde{u}_\varepsilon, \tilde{B}_\varepsilon) \rightarrow (\beta_k, \rho, u, h, B) \text{ in } S \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.6)$$

4.4. The vanishing viscosity limit. In this subsection, we shall pass to the limit as $\varepsilon \rightarrow 0$ in (3.1)-(3.3). Note that the parameter δ is kept fixed throughout this section and then we can use the previously derived estimates. Similarly, from Proposition 3.8, we can deduce that $(\tilde{\rho}_\varepsilon, \tilde{u}_\varepsilon, \tilde{\rho}_\varepsilon \tilde{u}_\varepsilon, \tilde{B}_\varepsilon)$ satisfies the same estimates as $(\rho_\varepsilon, u_\varepsilon, \rho_\varepsilon u_\varepsilon, B_\varepsilon)$. For simplicity of notations, we will again drop the tildes on the random variables. It follows from (4.1) that

$$\varepsilon \nabla u_\varepsilon \cdot \nabla \rho_\varepsilon \rightarrow 0 \text{ in } L^1(\tilde{\Omega} \times (0, T) \times D). \quad (4.7)$$

Similarly, we have $\varepsilon \Delta \rho_\varepsilon \rightarrow 0$ in $L^2(\tilde{\Omega}, L^2(0, T; H^{-1}(D)))$. From (4.3), we have

$$a \rho_\varepsilon^\gamma + \delta \rho_\varepsilon^\beta \rightarrow \bar{P} \text{ weakly in } L^{\frac{\beta+1}{\beta}}(\tilde{\Omega} \times (0, T) \times D). \quad (4.8)$$

We can obtain as in Section 3.4

$$\rho_\varepsilon \rightarrow \rho \text{ in } C(0, T; H^{-1}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.9)$$

Furthermore, (4.6) implies that

$$u_\varepsilon \rightharpoonup u \text{ in } L^2(0, T; H^1(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.10)$$

It follow from (4.10) and (4.9) that

$$\rho_\varepsilon u_\varepsilon \rightarrow \rho u \text{ in } \mathcal{D}'((0, T) \times D) \text{ } \tilde{\mathbb{P}} - \text{a.s.}$$

Since $C_0^\infty(D)$ is density in $L^{\frac{2\gamma}{\gamma-1}}(D)$, we know that

$$\rho_\varepsilon u_\varepsilon \rightarrow \rho u \text{ in } C([0, T]; L_w^{\frac{2\gamma}{\gamma+1}}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.},$$

which together with the fact that $\frac{2\gamma}{\gamma+1} > \frac{6}{5}$, $L^{\frac{2\gamma}{\gamma+1}}(D) \hookrightarrow H^{-1}(D)$ and Aubin-Lions Lemma implies that

$$\rho_\varepsilon u_\varepsilon \rightarrow \rho u \text{ in } C([0, T]; H^{-1}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.11)$$

Therefore, by (4.10) and (4.11) one has

$$\rho_\varepsilon u_\varepsilon^i u_\varepsilon^j \rightarrow \rho u^i u^j \text{ in } \mathcal{D}'((0, T) \times D), i, j = 1, 2, 3 \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.12)$$

By (4.6), we know that

$$B_\varepsilon \rightarrow B \text{ in } L^2(0, T; L^2(D)) \text{ and } B_\varepsilon \rightharpoonup B \text{ in } L^2(0, T; H^1(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.13)$$

Moreover, from (3.1)₃ and (4.2), by Aubin-Lions Lemma, one has

$$B_\varepsilon \rightarrow B \text{ in } C([0, T]; H^{-1}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.}$$

which together with (4.13) yields

$$(\nabla \times B_\varepsilon) \times B_\varepsilon \rightarrow (\nabla \times B) \times B \text{ in } \mathcal{D}'([0, T] \times D) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.14)$$

Similarly, it follows from (4.10) and (4.13) that

$$\nabla \times (u_\varepsilon \times B_\varepsilon) \rightarrow \nabla \times (u \times B) \text{ in } \mathcal{D}'([0, T] \times D) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (4.15)$$

By (1.5), Hölder's inequality, (4.6), (4.11), (4.13), we have

$$\begin{aligned} \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle &\rightarrow \langle f_k(\rho, \rho u, x), \phi \rangle \text{ in } L^1([0, T]) \text{ } \tilde{\mathbb{P}} - \text{a.s.}, \\ \langle g_k(B_\varepsilon, x), \phi \rangle &\rightarrow \langle g_k(B, x), \phi \rangle \text{ in } L^1([0, T]) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \end{aligned} \quad (4.16)$$

For $\phi \in C_0^\infty(D)$, we set $\langle M_\varepsilon(t), \phi \rangle := \int_0^t \sum_{k \geq 1} \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle d\beta_{k,\varepsilon}^1$ and $\langle \tilde{M}_\varepsilon(t), \phi \rangle := \int_0^t \sum_{k \geq 1} \langle g_k(B_\varepsilon, x), \phi \rangle d\beta_{k,\varepsilon}^2$, which are martingales under $\tilde{\mathbb{P}}$. It follows from Proposition 3.8 that

$$\begin{aligned} \langle M_\varepsilon(t), \phi \rangle &= \langle \rho_\varepsilon u_\varepsilon(t) - \rho_\varepsilon u_\varepsilon(0), \phi \rangle + \int_0^t \langle \operatorname{div}(\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon) + \delta \nabla \rho_\varepsilon^\beta - \mu \Delta u_\varepsilon, \phi \rangle ds \\ &\quad + \int_0^t \langle (\nabla \times B_\varepsilon) \times B_\varepsilon + \varepsilon \nabla \rho_\varepsilon \cdot \nabla u_\varepsilon - (\lambda + \mu) \nabla \operatorname{div} u_\varepsilon + a \nabla \rho_\varepsilon^\gamma, \phi \rangle ds \\ &:= \langle \rho_\varepsilon u_\varepsilon(t) - \rho_\varepsilon u_\varepsilon(0), \phi \rangle + \int_0^t \langle \Lambda_\varepsilon, \phi \rangle ds, \\ \langle \tilde{M}_\varepsilon(t), \phi \rangle &= \langle B_\varepsilon(t) - B_\varepsilon(0), \phi \rangle + \int_0^t \langle \nabla \times (u_\varepsilon \times B_\varepsilon) + \nu \Delta B_\varepsilon, \phi \rangle ds \\ &:= \langle B_\varepsilon(t) - B_\varepsilon(0), \phi \rangle + \int_0^t \langle \tilde{\Lambda}_\varepsilon, \phi \rangle ds, \end{aligned}$$

We define $\langle M(t), \phi \rangle$ and $\langle \tilde{M}(t), \phi \rangle$ by dropping the subscript ε on the right-hand side of the above two equalities.

Proposition 4.3. The system $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ is a martingale solution of the following equations:

$$\begin{cases} d\rho + \operatorname{div}(\rho u) dt = 0, \\ d(\rho u) + [\operatorname{div}(\rho u \otimes u) + \nabla \bar{P} - \mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u - (\nabla \times B) \times B] dt = dM, \\ dB - [\nabla \times (u \times B) + \nu \Delta B] dt = d\tilde{M}, \end{cases} \quad (4.17)$$

in $\mathcal{D}'((0, T) \times D)$ $\tilde{\mathbb{P}}$ -a.s..

Proof. Similar to Proposition 3.11, we can easily prove that $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ satisfies the equation (4.17)₁. Now, we only need to prove that $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ satisfies the equations (4.17)₂ and (4.17)₃.

For $0 \leq s < t < \infty$ and a bounded, continuous \mathcal{F}_s -measurable function $\varphi_\varepsilon = \varphi(\beta_{k,\varepsilon}, \rho_\varepsilon, u_\varepsilon, B_\varepsilon)$, we have

$$\begin{aligned} \tilde{\mathbb{E}}[\langle M_\varepsilon(t) - M_\varepsilon(s), \phi \rangle \varphi_\varepsilon] &= 0, \quad \tilde{\mathbb{E}}[\langle \tilde{M}_\varepsilon(t) - \tilde{M}_\varepsilon(s), \phi \rangle \varphi_\varepsilon] = 0, \\ \tilde{\mathbb{E}}\left[\left(\langle M_\varepsilon(t) - M_\varepsilon(s), \phi \rangle \beta_{k,\varepsilon}^1 - \int_s^t \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle d\tau\right) \varphi_\varepsilon\right] &= 0, \\ \tilde{\mathbb{E}}\left[\left(\langle \tilde{M}_\varepsilon(t) - \tilde{M}_\varepsilon(s), \phi \rangle \beta_{k,\varepsilon}^2 - \int_s^t \langle g_k(B_\varepsilon, x), \phi \rangle d\tau\right) \varphi_\varepsilon\right] &= 0. \end{aligned} \quad (4.18)$$

By (4.6)-(4.8), (4.10)-(4.16), we can infer that

$$\langle \rho_\varepsilon u_\varepsilon(t) - \rho_\varepsilon u_\varepsilon(s), \phi \rangle + \int_s^t \langle \Lambda_\varepsilon, \phi \rangle d\tau, \quad \langle B_\varepsilon(t) - B_\varepsilon(s), \phi \rangle + \int_s^t \langle \tilde{\Lambda}_\varepsilon, \phi \rangle d\tau$$

converges to

$$\langle \rho u(t) - \rho u(s), \phi \rangle - \int_s^t \langle \Lambda, \phi \rangle d\tau, \quad \langle B(t) - B(s), \phi \rangle + \int_s^t \langle \tilde{\Lambda}, \phi \rangle d\tau \quad \text{almost surely on } \tilde{\Omega},$$

where

$$\begin{aligned}\langle \Lambda, \phi \rangle &= \langle \operatorname{div}(\rho u \otimes u) - \mu \Delta u - (\lambda + \mu) \operatorname{div} \nabla u + (\nabla \times B) \times B + \overline{P}, \phi \rangle \\ \langle \tilde{\Lambda}, \phi \rangle &= \langle \nabla \times (u \times B) + \nu \Delta B, \phi \rangle.\end{aligned}$$

We can prove that the functions $f_\varepsilon := \langle M_\varepsilon(t) - M_\varepsilon(s), \phi \rangle \varphi(\beta_{k,\varepsilon}, \rho_\varepsilon, u_\varepsilon, B_\varepsilon)$ are uniformly integrable as (3.44), (3.45) and (3.46). By the uniform integrability, (4.18), and Lemma 3.10, we have

$$\begin{aligned}\tilde{\mathbb{E}}[\langle M(t) - M(s), \phi \rangle \varphi(\beta_k, \rho, u, B)] &= \lim_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}}[\langle M_\varepsilon(t) - M_\varepsilon(s), \phi \rangle \varphi_\varepsilon] = 0, \\ \tilde{\mathbb{E}}[\langle \tilde{M}(t) - \tilde{M}(s), \phi \rangle \varphi(\beta_k, \rho, u, B)] &= \lim_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}}[\langle \tilde{M}_\varepsilon(t) - \tilde{M}_\varepsilon(s), \phi \rangle \varphi_\varepsilon] = 0, \\ \tilde{\mathbb{E}}\left[\left(\langle M(t) - M(s), \phi \rangle \beta_k^1 - \int_s^t \langle f_k(\rho, \rho u, x), \phi \rangle d\tau\right) \varphi(\beta_k, \rho, u, B)\right] \\ &= \lim_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}}\left[\left(\langle M_\varepsilon(t) - M_\varepsilon(s), \phi \rangle \beta_{k,\varepsilon}^1 - \int_s^t \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle d\tau\right) \varphi_\varepsilon\right] = 0, \\ \tilde{\mathbb{E}}\left[\left(\langle \tilde{M}(t) - \tilde{M}(s), \phi \rangle \beta_k^2 - \int_s^t \langle g_k(B, x), \phi \rangle d\tau\right) \varphi(\beta_k, \rho, u, B)\right] \\ &= \lim_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}}\left[\left(\langle \tilde{M}_\varepsilon(t) - \tilde{M}_\varepsilon(s), \phi \rangle \beta_{k,\varepsilon}^2 - \int_s^t \langle g_k(B_\varepsilon, x), \phi \rangle d\tau\right) \varphi_\varepsilon\right] = 0.\end{aligned}$$

□

Our next goal is to prove

$$\overline{P} = \alpha \rho^\gamma + \delta \rho^\beta \quad (\overline{P} \text{ appears in the limit in (4.8)})$$

which is equivalent to the strong convergence of ρ_ε in $L^1(\tilde{\Omega} \times (0, T) \times D)$.

4.5. The effective viscous flux. We introduce the quantity $a\rho^\gamma + \delta\rho^\beta - (\lambda + 2\mu)\operatorname{div}u$ that is usually called the effective viscous flux. This quantity satisfies many properties for which we refer to [15, 30] for details. As in [34, 38], we can obtain the following lemma:

Lemma 4.4. Assume that $\beta > \max\{4, \gamma, \frac{4\gamma}{2\gamma-3}\}$. Let $(\rho_\varepsilon, u_\varepsilon, B_\varepsilon)$ be the sequence of approximate solutions obtained in Lemma 3.5, and let ρ, u, \overline{P} be the limits appearing in (4.9), (4.10), and (4.8) respectively. Then we have

$$\begin{aligned}\lim_{\varepsilon \rightarrow 0^+} \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \left[a\rho_\varepsilon^\gamma + \delta\rho_\varepsilon^\beta - (\lambda + 2\mu)\operatorname{div}u_\varepsilon \right] \rho_\varepsilon dx dt \\ = \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \left[\overline{P} - (\lambda + 2\mu)\operatorname{div}u \right] \rho dx dt,\end{aligned}$$

for any $\psi \in \mathcal{D}(0, T)$ and $\phi \in \mathcal{D}(D)$.

The idea of the proof of the above lemma is based on the Div-Curl Lemma. First, we introduce the operator \mathcal{A} by

$$\mathcal{A}_i v = \Delta^{-1}[\partial_{x_i} v], \quad i = 1, 2, 3.$$

Here Δ^{-1} stands for the inverse of the Laplace operator in \mathbb{R}^3 . Furthermore

$$\mathcal{A}_j(\xi) = \frac{-i\xi_j}{|\xi|^2}, \quad \partial_{x_i} \mathcal{A}_i[v] = v.$$

The classical Mihklin multiplier theorem implies that

$$\|\mathcal{A}_i v\|_{W^{1,s}(D)} \leq C(s, D)\|v\|_{L^s(\mathbb{R}^3)}, \quad 1 < s < \infty,$$

$$\|\mathcal{A}_i v\|_{L^q(D)} \leq C(q, s, D)\|\mathcal{A}_i v\|_{W^{1,s}(D)} \leq C(q, s, D)\|v\|_{L^s(\mathbb{R}^3)}, \quad \frac{1}{q} \geq \frac{1}{s} - \frac{1}{3},$$

and

$$\|\mathcal{A}_i v\|_{L^\infty(D)} \leq C(s, D)\|v\|_{L^s(\mathbb{R}^3)}, \quad s > 3.$$

Now, extending ρ_ε to be zero outside D . Consider

$$\varphi(t, x) = \psi(t)\phi(x)\mathcal{A}[\rho_\varepsilon], \quad \psi \in \mathcal{D}(0, T), \quad \phi \in \mathcal{D}(D).$$

Note that (4.1) ensures that φ can be used as a test function for (3.1). Note that $\operatorname{div}\mathcal{A}[\rho_\varepsilon] = \rho_\varepsilon$ and $\Delta\mathcal{A}_j = \partial_j$. Similar to Section 4.2, we have

$$\begin{aligned} & \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \left[a\rho_\varepsilon^\gamma + \delta\rho_\varepsilon^\beta - (\lambda + 2\mu)\operatorname{div}u_\varepsilon \right] \rho_\varepsilon dxdt \\ &= (\lambda + \mu)\tilde{\mathbb{E}} \int_0^T \psi \int_D \operatorname{div}u_\varepsilon \partial_{x_i} \phi \mathcal{A}_i[\rho_\varepsilon] dxdt - \tilde{\mathbb{E}} \int_0^T \psi \int_D (a\rho_\varepsilon^\gamma + \delta\rho_\varepsilon^\beta) \partial_{x_i} \phi \mathcal{A}_i[\rho_\varepsilon] dxdt \\ &+ \mu\tilde{\mathbb{E}} \int_0^T \psi \int_D \partial_{x_j} u_\varepsilon^i \partial_{x_j} \phi \mathcal{A}_i[\rho_\varepsilon] dxdt - \tilde{\mathbb{E}} \int_0^T \psi \int_D \rho_\varepsilon u_\varepsilon^i u_\varepsilon^j \partial_{x_j} \phi \mathcal{A}_i[\rho_\varepsilon] dxdt \\ &- \tilde{\mathbb{E}} \int_0^T \psi_t \int_D \phi \rho_\varepsilon u_\varepsilon^i \mathcal{A}_i[\rho_\varepsilon] dxdt - \varepsilon\tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \rho_\varepsilon u_\varepsilon \mathcal{A}_i[\operatorname{div}(1_D \nabla \rho_\varepsilon)] dxdt \\ &+ \varepsilon\tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \partial_{x_j} \rho_\varepsilon \partial_{x_j} u_\varepsilon^i \mathcal{A}_i[\rho_\varepsilon] dxdt + \mu\tilde{\mathbb{E}} \int_0^T \psi \int_D u_\varepsilon \partial_{x_i} \phi \rho_\varepsilon dxdt \\ &- \mu\tilde{\mathbb{E}} \int_0^T \psi \int_D u_\varepsilon^i \partial_{x_j} \phi \partial_{x_j} \mathcal{A}_i[\rho_\varepsilon] dxdt + \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \mathcal{A}[\rho_\varepsilon] (\nabla \times B_\varepsilon) \times B_\varepsilon dxdt \\ &+ \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi u_\varepsilon^i (\rho_\varepsilon \mathcal{R}_{ij}[\rho_\varepsilon u_\varepsilon^j] - \rho_\varepsilon u_\varepsilon^j \mathcal{R}_{ij}[\rho_\varepsilon]) dxdt, \end{aligned} \tag{4.19}$$

where \mathcal{R}_{ij} is the Rize operators, which is defined by

$$\mathcal{R}_{i,j}[v] = \partial_{x_j} \mathcal{A}_i[v] \quad \text{or} \quad \hat{\mathcal{R}}_{i,j}(\xi) = \frac{\xi_i \xi_j}{|\xi|^2}.$$

Here we can extend the function ρ_ε by zero outside D and

$$\partial_t \rho_\varepsilon = \begin{cases} \varepsilon \Delta \rho_\varepsilon - \operatorname{div}(\rho_\varepsilon u_\varepsilon) & \text{in } D, \\ 0 & \text{in } \mathbb{R}^3 \setminus D. \end{cases}$$

Since $u_\varepsilon|_{\partial D} = 0$, then we have

$$\operatorname{div}(\rho_\varepsilon u_\varepsilon) = 0 \quad \text{on } \mathbb{R}^3 \setminus D.$$

Similarly, for $\nabla \rho_\varepsilon$, we also have

$$\operatorname{div}(1_D \nabla \rho_\varepsilon) = \begin{cases} \Delta \rho_\varepsilon & \text{in } D, \\ 0 & \text{on } \mathbb{R}^3 \setminus D. \end{cases}$$

We have the following lemma:

Lemma 4.5. Assume that $\rho \in L^p(\tilde{\Omega}, L^2((0, T) \times D))$ and $u \in L^p(\tilde{\Omega}, L^2(0, T; H_0^1(D)))$ solve (4.17) in $\mathcal{D}'((0, T) \times D)$ almost surely. Then we can extend (ρ, u) to be zero on $\mathbb{R}^3 \setminus D$. Moreover, the equation (4.17) holds in $\mathcal{D}'((0, T) \times \mathbb{R}^3)$ almost surely.

Next, we consider the test functions

$$\varphi_i(t, x) = \psi(t)\phi(x)\mathcal{A}_i[\rho], \quad i = 1, 2, 3,$$

where ρ is zero outside D . Similar to (4.19), taking φ as test functions for (4.17)₂, integrating over $\tilde{\Omega} \times (0, T) \times D$, we obtain

$$\begin{aligned} & \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi [\overline{P} - (\lambda + 2\mu)\operatorname{div}u] \rho dx dt \\ &= (\lambda + \mu) \tilde{\mathbb{E}} \int_0^T \psi \int_D \operatorname{div}u \partial_{x_i} \phi \mathcal{A}_i[\rho] dx dt - \tilde{\mathbb{E}} \int \psi \int_D \overline{P} \partial_{x_i} \phi \mathcal{A}_i[\rho] dx dt \\ &+ \mu \tilde{\mathbb{E}} \int_0^T \psi \int_D \partial_{x_j} u_i \partial_{x_j} \phi \mathcal{A}_i[\rho] dx dt - \tilde{\mathbb{E}} \int_0^T \psi \int_D \rho u_i u_j \partial_{x_j} \phi \mathcal{A}_i[\rho] dx dt \\ &- \tilde{\mathbb{E}} \int_0^T \psi_t \int_D \phi \rho u_i \mathcal{A}_i[\rho] dx dt + \mu \tilde{\mathbb{E}} \int_0^T \psi \int_D u_i \partial_{x_i} \phi \rho dx dt \\ &- \mu \tilde{\mathbb{E}} \int_0^T \psi \int_D u_i \partial_{x_j} \phi \partial_{x_j} \mathcal{A}_i[\rho] dx dt + \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi \mathcal{A}[\rho] (\nabla \times B) \times B dx dt \\ &+ \tilde{\mathbb{E}} \int_0^T \psi \int_D \phi u_i (\rho \mathcal{R}_{i,j}[\rho u_j] - \rho u_j \mathcal{R}_{i,j}[\rho]) dx dt. \end{aligned} \tag{4.20}$$

Here we have used the following property: Recall that $M(t) = \rho u(t) - \rho u(0) + \int_0^t [\operatorname{div}(\rho u \otimes u) + \nabla \overline{P} - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div}u - (\nabla \times B) \times B] ds$ and M is a martingale. So we can prove that $\int_0^t \psi(t)\phi(x)\mathcal{A}_i[\rho] dM = 0$, where $\psi \in C_0^\infty([0, T])$, $\phi(x) \in C_0^\infty(D)$. In fact, integrating by parts, we have

$$\int_0^t \psi(t)\phi(x)\mathcal{A}_i[\rho] dM = \psi(t)\phi(x)\mathcal{A}_i[\rho]M - \int_0^t M d(\psi(s)\phi(x)\mathcal{A}_i[\rho]).$$

We denote the term $\int_0^t M d(\psi(s)\phi(x)\mathcal{A}_i[\rho])$ by $\int_0^t M dN(s)$. Let $\Delta_n := \{0 = t_0 \leq t_1 \leq t_2 \leq \dots \leq t_{k-1} \leq t_k \leq \dots \leq t_n = t\}$ be a subdivision of the time interval and denote $\|\Delta_n\| := \max_{1 \leq k \leq n} |t_k - t_{k-1}|$. Then we have

$$\begin{aligned} & \int_0^t M d(\psi(s)\phi(x)\mathcal{A}_i[\rho]) = \sum_{k=1}^n M(t_k)[N(t_k) - N(t_{k-1})] \\ &= \sum_{k=1}^n N(t_k)M(t_k) - \sum_{k=1}^{n-1} N(t_k)M(t_k) + \sum_{k=0}^{n-1} N(t_k)M(t_k) - \sum_{k=0}^{n-1} M(t_{k+1})N(t_k) \\ &= N(t)M(t) - \sum_{k=0}^{n-1} N(t_k)[M(t_{k+1}) - M(t_k)]. \end{aligned}$$

Let $\|\Delta_n\| \rightarrow 0$, since $N(t_k)$ is independent on $M(t_{k+1}) - M(t_k)$ and M is a martingale, we have $\tilde{\mathbb{E}} \int_0^t M d(\psi(s)\phi(x)\mathcal{A}_i[\rho]) = \tilde{\mathbb{E}} N(t)M(t)$. Then $\tilde{\mathbb{E}} \int_0^t \psi(t)\phi(x)\mathcal{A}_i[\rho] dM = 0$.

Now following [34, 38] we can prove that the right-hand side of (4.19) converges to the right-hand side of (4.20) as $\varepsilon \rightarrow 0$, which proves Lemma 4.4.

4.6. Strong convergence of the density. In this subsection, we will follow the idea of [15, 30] to prove the strong convergence of the sequence ρ_ε . Our goal is to show $\bar{P} = a\rho^\gamma + \delta\rho^\beta$.

Lemma 4.5 implies that we can extend (ρ, u) to be zero on $\mathbb{R}^3 \setminus D$, and

$$\rho \in L^p(\tilde{\Omega}, L^2(0, T; L^2(D))), \quad u \in L^p(\tilde{\Omega}, L^2(0, T; H_0^1(D)))$$

solve the continuity equation (4.17)₁ in $\mathcal{D}'((0, T) \times \mathbb{R}^3)$ almost surely. Taking a regularizing sequence $\vartheta_m = \vartheta_m(x)$ and then using S_m to test (4.17)₁, we have

$$\partial_t S_m[\rho] + \operatorname{div}[S_m(\rho)u] = r_m \quad \text{on } (0, T) \times \mathbb{R}^3. \quad (4.21)$$

Here $S_m[v] = \vartheta_m * v$ is the standard smoothing operator. We note that [29, Lemma 2.3] implies that the remainder term $r_m \rightarrow 0$ in $L^1(\tilde{\Omega} \times (0, T) \times \mathbb{R}^3)$ as $m \rightarrow \infty$.

For any b satisfying the conditions in Definition 1.1, multiplying (4.21) by $b'(S_m[\rho])$ and passing to the limit as $m \rightarrow \infty$, we can infer that (ρ, u) solves (4.17)₁ in the sense of renormalized solutions. Furthermore, taking $b(z) = z \ln z$ and then integrating (1.4) over $\tilde{\Omega} \times (0, T) \times D$, one has

$$\tilde{\mathbb{E}} \int_0^T \int_D \rho \operatorname{div} u \, dx \, dt = \tilde{\mathbb{E}} \int_D \rho_0 \ln \rho_0 \, dx - \tilde{\mathbb{E}} \int_0^T \rho(T) \ln \rho(T) \, dx. \quad (4.22)$$

Since $(\rho_\varepsilon, u_\varepsilon)$ satisfies the equation (3.1)₁ a.e. on $\tilde{\Omega} \times (0, T) \times D$, multiplying (3.1)₁ with $b'(\rho_\varepsilon)$, then for any b convex and globally Lipschitz on \mathbb{R}^+ , we have

$$\partial_t b(\rho_\varepsilon) + \operatorname{div}(b(\rho_\varepsilon)u_\varepsilon) + [b'(\rho_\varepsilon)\rho_\varepsilon - b(\rho_\varepsilon)] \operatorname{div} u_\varepsilon - \varepsilon \Delta b(\rho_\varepsilon) = -\varepsilon b''(\rho_\varepsilon) |\nabla \rho_\varepsilon|^2 \leq 0.$$

Integrating the above on $\tilde{\Omega} \times (0, T) \times D$, the boundary condition yields

$$\tilde{\mathbb{E}} \int_0^T \int_D [b'(\rho_\varepsilon)\rho_\varepsilon - b(\rho_\varepsilon)] \operatorname{div} u_\varepsilon \, dx \, dt \leq \tilde{\mathbb{E}} \int_D b(\rho_0) \, dx - \tilde{\mathbb{E}} \int_D b(\rho_\varepsilon(T)) \, dx.$$

Let $b(z) = z \ln z$ in the above inequality, then

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_0^T \int_D \rho_\varepsilon \operatorname{div} u_\varepsilon \, dx \, dt &\leq \tilde{\mathbb{E}} \int_D \rho_0 \ln \rho_0 \, dx - \underline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_D \rho_\varepsilon(T) \ln \rho_\varepsilon(T) \, dx \\ &\leq \tilde{\mathbb{E}} \int_0^T \int_D \rho \operatorname{div} u \, dx \, dt. \end{aligned} \quad (4.23)$$

Taking two nondecreasing sequences $0 \leq \psi_n \in \mathcal{D}(0, T)$, $0 \leq \phi_n \in \mathcal{D}(D)$ satisfying

$$\begin{aligned} \psi_n &\rightarrow 1, \quad \phi_n \rightarrow 1 \quad \text{as } n \rightarrow \infty; \quad \psi_n(t) = 1 \quad \text{for } t \geq \frac{1}{n} \quad \text{or } t \leq T - \frac{1}{n}; \\ \phi_n(x) &= 1 \quad \text{for } x \in D, \quad \operatorname{dist}(x, \partial D) \geq \frac{1}{n}. \end{aligned}$$

Combining Lemma 4.4, (4.22) and (4.23), for all $m \leq n$, we obtain

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_0^T \psi_m \int_D \phi_m (a\rho_\varepsilon^\gamma + \delta\rho_\varepsilon^\beta) \rho_\varepsilon \, dx \, dt &\leq \overline{\lim}_{\varepsilon \rightarrow 0^+} \tilde{\mathbb{E}} \int_0^T \psi_n \int_D \phi_n (a\rho_\varepsilon^\gamma + \delta\rho_\varepsilon^\beta) \rho_\varepsilon \, dx \, dt \\ &\leq \overline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_0^T \psi_n \int_D \phi_n [a\rho_\varepsilon^\gamma + \delta\rho_\varepsilon^\beta - (\lambda + 2\mu) \operatorname{div} u_\varepsilon] \rho_\varepsilon \, dx \, dt \\ &\quad + (\lambda + 2\mu) \overline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_0^T \psi_n \int_D \phi_n \rho_\varepsilon \operatorname{div} u_\varepsilon \, dx \, dt \end{aligned}$$

$$\begin{aligned}
&\leq \tilde{\mathbb{E}} \int_0^T \psi_n \int_D \phi_n [\bar{P} - (\lambda + 2\mu) \operatorname{div} u] \rho dx dt + (\lambda + 2\mu) \overline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_0^T \int_D \rho_\varepsilon |1 - \psi_n \phi_n| |\operatorname{div} u_\varepsilon| dx dt \\
&\quad + (\lambda + 2\mu) \overline{\lim}_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \int_0^T \int_D \rho_\varepsilon \operatorname{div} u_\varepsilon dx dt \\
&\leq \tilde{\mathbb{E}} \int_0^T \int_D \bar{P} \rho dx dt + (\lambda + 2\mu) \tilde{\mathbb{E}} \int_0^T \int_D \rho \operatorname{div} u dx dt - (\lambda + 2\mu) \tilde{\mathbb{E}} \int_0^T \int_D \rho \operatorname{div} u dx dt + \eta(n) \\
&\leq \tilde{\mathbb{E}} \int_0^T \int_D \bar{P} \rho dx dt + \eta(n),
\end{aligned}$$

where $\eta(n) = \tilde{\mathbb{E}} \int_0^T \int_D (\psi_n \phi_n - 1) [\bar{P} - (\lambda + 2\mu) \operatorname{div} u] \rho dx dt + (\lambda + 2\mu) \tilde{\mathbb{E}} \int_0^T \int_D \rho_\varepsilon |1 - \psi_n \phi_n| |\operatorname{div} u_\varepsilon| dx dt \rightarrow 0$ as $n \rightarrow \infty$. Passing to the limit in the above inequality as $n \rightarrow \infty$, one deduces that

$$\overline{\lim}_{\varepsilon \rightarrow 0^+} \tilde{\mathbb{E}} \int_0^T \psi_m \int_D \phi_m (a \rho_\varepsilon^\gamma + \delta \rho_\varepsilon^\beta) \rho_\varepsilon dx dt \leq \tilde{\mathbb{E}} \int_0^T \int_D \bar{P} \rho dx dt \quad \text{for all } m = 1, 2, \dots$$

Since $P(z) = az^\gamma + \delta z^\beta$ is monotone, then we have

$$\tilde{\mathbb{E}} \int_0^T \psi_m \int_D \phi_m [P(\rho_\varepsilon) - P(v)] (\rho_\varepsilon - v) dx dt \geq 0.$$

Furthermore, the facts $\rho_\varepsilon \rightarrow \rho$ and $P(\rho_\varepsilon) \rightarrow \bar{P}$ imply that

$$\tilde{\mathbb{E}} \int_0^T \psi_m \int_D \phi_m \bar{P} \rho dx dt + \tilde{\mathbb{E}} \int_0^T \psi_m \int_D \phi_m P(v) v dx dt - \tilde{\mathbb{E}} \int_0^T \psi_m \int_D \phi_m [\bar{P} v + P(v) \rho] dx dt \geq 0.$$

Now, letting $m \rightarrow \infty$ in the above inequality, one has

$$\tilde{\mathbb{E}} \int_0^T \int_D [\bar{P} - P(v)] (\rho - v) dx dt \geq 0.$$

Choosing $v = \rho + \alpha \varphi$, for an arbitrary φ , and then letting $\alpha \rightarrow 0$, we have

$$\bar{P} = a \rho^\gamma + \delta \rho^\beta.$$

That is,

$$a \rho_\varepsilon^\gamma + \delta \rho_\varepsilon^\beta \rightarrow a \rho^\gamma + \delta \rho^\beta \quad \text{in } L^{\frac{\beta+1}{\beta}}(\tilde{\Omega} \times (0, T) \times D).$$

Then we have

$$\rho_\varepsilon \rightarrow \rho \quad \text{in } L^s(\tilde{\Omega} \times (0, T) \times D), \quad \forall 1 \leq s < \beta + 1. \quad (4.24)$$

Similar to the proof of (3.40), by (1.5), Hölder's inequality, (4.11), (4.13), (4.24), we can deduce that

$$\begin{aligned}
\left\langle \sum_{k \geq 1} f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \right\rangle^2 &\rightarrow \left\langle \sum_{k \geq 1} f_k(\rho, \rho u, x), \phi \right\rangle^2 \quad \text{in } L^1([0, T]) \quad \tilde{\mathbb{P}} - \text{a.s.}, \\
\left\langle \sum_{k \geq 1} g_k(B_\varepsilon, x), \phi \right\rangle^2 &\rightarrow \left\langle \sum_{k \geq 1} g_k(B, x), \phi \right\rangle^2 \quad \text{in } L^1([0, T]) \quad \tilde{\mathbb{P}} - \text{a.s.}
\end{aligned} \quad (4.25)$$

Note that

$$\begin{aligned} \tilde{\mathbb{E}} \left[\left(\langle M_\varepsilon(t), \phi \rangle^2 - \langle M_\varepsilon(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle^2 d\tau \right) \varphi_\varepsilon \right] &= 0, \\ \tilde{\mathbb{E}} \left[\left(\langle \tilde{M}_\varepsilon(t), \phi \rangle^2 - \langle \tilde{M}_\varepsilon(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle g_k(B_\varepsilon, x), \phi \rangle^2 d\tau \right) \varphi_\varepsilon \right] &= 0. \end{aligned} \quad (4.26)$$

Then, it follows from (4.25), (4.26), the uniform integrability, Proposition 3.9 and Lemma 3.10 that

$$\begin{aligned} &\tilde{\mathbb{E}} \left[\left(\langle M(t), \phi \rangle^2 - \langle M(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle f_k(\rho, \rho u, x), \phi \rangle^2 d\tau \right) \varphi(\beta_k, \rho, u, B) \right] \\ &= \lim_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \left[\left(\langle M_\varepsilon(t), \phi \rangle^2 - \langle M_\varepsilon(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle^2 d\tau \right) \varphi_\varepsilon \right] = 0, \\ &\tilde{\mathbb{E}} \left[\left(\langle \tilde{M}(t), \phi \rangle^2 - \langle \tilde{M}(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle g_k(B, x), \phi \rangle^2 d\tau \right) \varphi(\beta_k, \rho, u, B) \right] \\ &= \lim_{\varepsilon \rightarrow 0} \tilde{\mathbb{E}} \left[\left(\langle \tilde{M}_\varepsilon(t), \phi \rangle^2 - \langle \tilde{M}_\varepsilon(s), \phi \rangle^2 - \int_s^t \sum_{k \geq 1} \langle g_k(B_\varepsilon, x), \phi \rangle^2 d\tau \right) \varphi_\varepsilon \right] = 0. \end{aligned}$$

Hence we deduce that $\langle M(t), \phi \rangle^2 - \int_0^t \sum_{k \geq 1} \langle f_k(\rho, \rho u, x), \phi \rangle^2 ds$; $\langle \tilde{M}(t), \phi \rangle$ and $\langle \tilde{M}(t), \phi \rangle^2 - \int_0^t \sum_{k \geq 1} \langle g_k(B, x), \phi \rangle^2 ds$ are continuous martingales. Using again the method in [3, 28] we can also infer that

$$\langle M(t), \phi \rangle = \int_0^t \left\langle \sum_{k \geq 1} f_k(\rho, \rho u, x), \phi \right\rangle d\beta_k^1, \quad \langle \tilde{M}(t), \phi \rangle = \int_0^t \left\langle \sum_{k \geq 1} g_k(B, x), \phi \right\rangle d\beta_k^2.$$

Summing up the above results, we have the following proposition:

Proposition 4.6. Assume that $D \in C^{2+\alpha}$ is a bounded domain. If $\beta > \max\{\frac{6\gamma}{2\gamma-3}, \gamma, 4\}$, then there exists a finite energy martingale solution $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ of the problem:

$$\begin{cases} d\rho + \operatorname{div}(\rho u)dt = 0, \\ d(\rho u) + [\operatorname{div}(\rho u \otimes u) + a\nabla\rho^\gamma + \delta\nabla\rho^\beta - \mu\Delta u + (\lambda + \mu)\nabla\operatorname{div}u] dt \\ \quad = (\nabla \times B) \times B dt + \sum_{k \geq 1} f_k(\rho, \rho u, x) d\beta_k^1, \\ dB - [\nabla \times (u \times B) + \nu\Delta B]dt = \sum_{k \geq 1} g_k(B, x) d\beta_k^2, \end{cases} \quad (4.27)$$

$\tilde{\mathbb{P}}$ -a.s. and (ρ, u, B) satisfies the initial-boundary valued conditions (3.2), (3.3). Moreover, if (ρ, u) is prolonged to be zero on $\mathbb{R}^3 \setminus D$, then the equation (4.27)₁ holds in sense of renormalized solutions on $\mathcal{D}'((0, T) \times \mathbb{R}^3)$ $\tilde{\mathbb{P}}$ -a.s.. Finally, (ρ, u, B) satisfies the following estimates:

$$\tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\rho(t)\|_{L^\gamma(D)}^\gamma \right)^P \leq C, \quad \tilde{\mathbb{E}} \left(\delta \sup_{t \in [0, T]} \|\rho(t)\|_{L^\beta(D)}^\beta \right)^P \leq C,$$

$$\begin{aligned} \tilde{\mathbb{E}}\left(\sup_{t \in [0, T]} \|\sqrt{\rho}u\|_{L^2(D)}^2\right)^p &\leq C, \quad \tilde{\mathbb{E}}\left(\|\rho\|_{L^{\beta+1}((0, T) \times D)}\right)^p \leq C, \\ \tilde{\mathbb{E}}\left(\|u\|_{L^2(0, T; H_0^1(D))}^2\right)^p &\leq C, \quad \tilde{\mathbb{E}}\left(\|B\|_{L^\infty(0, T; L^2(D))} + \|B\|_{L^2(0, T; H_0^1(D))}^2\right)^p \leq C, \end{aligned}$$

where the constant C is independent of $\delta > 0$.

5. PASSING TO THE LIMIT IN THE ARTIFICIAL PRESSURE TERM

In this section, we shall follow the idea of [15, 30] to pass to the limit in (4.27) as $\delta \rightarrow 0$ for the artificial pressure term and relax the hypotheses on the initial data $(\rho_{0, \delta}, m_{0, \delta}, B_{0, \delta})$, that is, (3.2) and (3.3). First, from (3.1), we have

$$\rho_{0, \delta} \rightarrow \rho_0 \text{ in } L^\gamma(D), \quad m_{0, \delta} \rightarrow m_0 \text{ in } L^1(D), \quad B_{0, \delta} \rightarrow B_0 \text{ in } L^2(D) \quad \tilde{\mathbb{P}} - \text{a.s. as } \delta \rightarrow 0.$$

Next, we consider the problem (4.27) with the initial data $(\rho_{0, \delta}, m_{0, \delta}, B_{0, \delta})$. From Proposition 4.6, we obtain the existence of martingale solution, denoted by $(\beta_{k, \delta}, \rho_\delta, u_\delta, B_\delta)$. From (3.1), $\mathcal{E}_{0, \delta}$ is bounded uniformly in δ . Hence, the estimates in Proposition 4.6 hold independently of δ .

5.1. On integrability of the density. First, we shall estimate the density ρ_δ , uniformly in $\delta > 0$. Recall

$$\partial_t S_m[b(\rho_\delta)] + \operatorname{div}(S_m[b(\rho_\delta)]u_\delta) + S_m[(b'(\rho_\delta)\rho_\delta - b(\rho_\delta))\operatorname{div}u_\delta] = r_m, \quad (5.1)$$

where

$$r_m \rightarrow 0 \text{ in } L^2(\Omega, L^2(0, T; L^2(\mathbb{R}^3))) \text{ as } m \rightarrow \infty, \quad (5.2)$$

and b is uniformly bounded.

Denote $\int_D f dx = \frac{1}{|D|} \int_D f dx$. Similar to Section 4.2, we use the operator \mathcal{B} introduced in Section 4.1 to construct the following test functions:

$$\varphi_i(t, x) = \psi(t)\mathcal{B}_i \left[S_m[b(\rho_\delta)] - \int_D S_m[b(\rho_\delta)] dx \right], \quad i = 1, 2, 3, \quad \psi \in \mathcal{D}(0, T).$$

Note that $\varphi_i|_{\partial D} = 0$, $\varphi_i \in L^\infty(0, T; H_0^1(D))$, $\partial_t \varphi_i \in L^2(0, T; H_0^1(D))$. Thus, we can use φ_i as test functions for the equation (4.27)₂. Using (5.1), we obtain

$$\begin{aligned} &\mathbb{E} \int_0^T \psi \int_D (a\rho_\delta^\gamma + \delta\rho_\delta^\beta) S_m[b(\rho_\delta)] dx dt \\ &= \mathbb{E} \int_0^T \psi \int_D (a\rho_\delta^\gamma + \delta\rho_\delta^\beta) dx \int_D S_m[b(\rho_\delta)] dx dt + (\lambda + \mu) \mathbb{E} \int_0^T \psi \int_D S_m[b(\rho_\delta)] \operatorname{div}u_\delta dx dt \\ &\quad - \mathbb{E} \int_0^T \psi_t \int_D \rho_\delta u_\delta^i \mathcal{B}_i \left[S_m[b(\rho_\delta)] - \int_D S_m[b(\rho_\delta)] dx \right] dx dt \\ &\quad + \mathbb{E} \int_0^T \psi \int_D (\mu \partial_{x_j} u_\delta^i - \rho_\delta u_\delta^i u_\delta^j) \partial_{x_j} \mathcal{B}_i \left[S_m[b(\rho_\delta)] - \int_D S_m[b(\rho_\delta)] dx \right] dx dt \\ &\quad + \mathbb{E} \int_0^T \psi \int_D \rho_\delta u_\delta^i \mathcal{B}_i \left[S_m[(b(\rho_\delta) - b'(\rho_\delta)\rho_\delta)\operatorname{div}u_\delta] - \int_D S_m[(b(\rho_\delta) - b'(\rho_\delta)\rho_\delta)\operatorname{div}u_\delta] dx \right] dx dt \\ &\quad - \mathbb{E} \int_0^T \psi \int_D \rho_\delta u_\delta^i \mathcal{B}_i \left[r_m - \int_D r_m dx \right] + \mathbb{E} \int_0^T \psi \int_D \rho_\delta u_\delta^i \mathcal{B}_i [\operatorname{div}(S_m[b(\rho_\delta)]u_\delta)] dx dt \end{aligned}$$

$$+ \mathbb{E} \int_0^T \psi \int_D \mathcal{B}_i \left[S_m[b(\rho_\delta)] - \int_D S_m[b(\rho_\delta)] dx \right] (\nabla \times B_\delta) \times B_\delta dx dt. \quad (5.3)$$

Using (5.2), we can pass to the limit in (5.3) for $m \rightarrow \infty$. Moreover, we can apply the function $z^\theta \approx b(z)$ to obtain

$$\mathbb{E} \int_0^T \psi \int_D \left(a\rho_\delta^{\gamma+\theta} + \delta\rho_\delta^{\beta+\theta} \right) dx dt := \sum_{i=1}^8 J_i,$$

for some $\theta > 0$ to be determined below. Now, we just estimate the term J_8 on the right hand side of the above identity. Other terms can be bounded as in [38]. For the term J_8 , if $\theta < \frac{\gamma}{3}$, using Proposition 4.6 and (4.2), together with the embedding $W^{1,p}(D) \subset L^\infty(D)$ for $p > 3$, we have

$$\begin{aligned} J_8 &\leq \mathbb{E} \left| \int_0^T \int_D \psi \mathcal{B}_i \left[\rho_\delta^\theta - \int_D \rho_\delta^\theta dx \right] (\nabla \times B_\delta) \times B_\delta dx dt \right| \\ &\lesssim \mathbb{E} \int_0^T |\psi| \|\nabla B_\delta\|_{L^2(D)} \|B_\delta\|_{L^2(D)} \left\| \mathcal{B}_i \left[\rho_\delta^\theta - \int_D \rho_\delta^\theta dx \right] \right\|_{L^\infty(D)} dt \\ &\lesssim \mathbb{E} \int_0^T |\psi| \|\nabla B_\delta\|_{L^2(D)} \|B_\delta\|_{L^2(D)} \left\| \mathcal{B}_i \left[\rho_\delta^\theta - \int_D \rho_\delta^\theta dx \right] \right\|_{W^{1,\frac{\gamma}{\theta}}(D)} dt \\ &\lesssim \mathbb{E} \int_0^T |\psi| \|\nabla B_\delta\|_{L^2(D)} \|B_\delta\|_{L^2(D)} \left\| \rho_\delta^\theta \right\|_{L^{\frac{\gamma}{\theta}}(D)} dt \\ &\lesssim \mathbb{E} \left(\|\nabla B_\delta\|_{L^2(0,T;L^2(D))} \|B_\delta\|_{L^2(0,T;L^2(D))} \|\rho_\delta\|_{L^\gamma(D)}^\theta \right) \\ &\lesssim \left(\mathbb{E} \|\nabla B_\delta\|_{L^2(0,T;L^2(D))}^3 \right)^{\frac{1}{3}} \left(\mathbb{E} \|B_\delta\|_{L^2(0,T;L^2(D))}^3 \right)^{\frac{1}{3}} \left(\mathbb{E} \|\rho_\delta\|_{L^\gamma(D)}^{3\theta} \right)^{\frac{1}{3}} \\ &\leq C. \end{aligned}$$

Here the constant C is independent of δ .

Summing up the estimates for (5.3), we have

$$\mathbb{E} \int_0^T \psi \int_D \left(a\rho_\delta^{\gamma+\theta} + \delta\rho_\delta^{\beta+\theta} \right) dx dt \leq C. \quad (5.4)$$

Taking $\psi_n \in \mathcal{D}(0,T)$, $0 \leq \psi_n \leq 1$ and $\int_0^T |\partial_t \psi_n| dt \leq C$. By approximation and Lebesgue convergence theorem, we can take $\psi = 1$ in (5.4). Then we have proved the following result:

Lemma 5.1. For $\gamma > \frac{3}{2}$ and $0 < \theta < \min \{1, \frac{\gamma}{3}, \frac{2}{3}\gamma - 1\}$, then there exists a constant C independently of $\delta > 0$, such that

$$\mathbb{E} \int_0^T \int_D \left(a\rho_\delta^{\gamma+\theta} + \delta\rho_\delta^{\beta+\theta} \right) dx dt \leq C.$$

5.2. Tightness Property. Similar to Section 4.3, we can prove the following lemma:

Lemma 5.2. Define

$$\begin{aligned} S &= C(0, T; \mathbb{R}) \times C([0, T]; L_w^\gamma(D)) \times L^2(0, T; H_w^1(D)) \times C([0, T]; L_w^{\frac{2\gamma}{\gamma+1}}(D)) \\ &\quad \times (L^2(0, T; H_w^1(D)) \cap L^2(0, T; L^2(D))) \end{aligned}$$

equipped with its Borel σ -algebra. Let Π_δ be the probability on S which is the image of \mathbb{P} on Ω by the map: $\omega \mapsto (\beta_{k,\delta}(\omega, \cdot), \rho_\delta(\omega, \cdot), u_\delta(\omega, \cdot), \rho_\delta u_\delta(\omega, \cdot), B_\delta(\omega, \cdot))$, that is, for any $A \subseteq S$,

$$\Pi_\delta(A) = \mathbb{P} \{ \omega \in \Omega : (\beta_{k,\delta}(\omega, \cdot), \rho_\delta(\omega, \cdot), u_\delta(\omega, \cdot), \rho_\delta u_\delta(\omega, \cdot), B_\delta(\omega, \cdot)) \in A \}.$$

Then the family Π_δ is tight.

5.3. The limit passage. In this subsection, we shall pass to the limit for $\delta \rightarrow 0$ following the idea of [15, 30]. According to Jakubowski-Skorohod Theorem, there exists a subsequence such that $\Pi_\delta \rightarrow \Pi$ weakly, where Π is a probability on S . Moreover, there exist a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ and random variables $(\tilde{\beta}_{k,\delta}, \tilde{\rho}_\delta, \tilde{u}_\delta, \tilde{\rho}_\delta \tilde{u}_\delta, \tilde{B}_\delta)$ with distribution Π_δ , (β_k, ρ, u, h, B) with values in S such that

$$(\tilde{\beta}_{k,\delta}, \tilde{\rho}_\delta, \tilde{u}_\delta, \tilde{\rho}_\delta \tilde{u}_\delta, \tilde{B}_\delta) \rightarrow (\beta_k, \rho, u, h, B) \text{ in } S \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (5.5)$$

Since the distribution of $(\tilde{\beta}_{k,\delta}, \tilde{\rho}_\delta, \tilde{u}_\delta, \tilde{\rho}_\delta \tilde{u}_\delta, \tilde{B}_\delta)$ is Π_δ , we can deduce that $(\tilde{\rho}_\delta, \tilde{u}_\delta, \tilde{\rho}_\delta \tilde{u}_\delta, \tilde{B}_\delta)$ satisfies the same estimate as $(\rho_\delta, u_\delta, \rho_\delta u_\delta, B_\delta)$. To simplify notations, we drop the tilde on the random variables. First, we can apply Lemma 5.1 to obtain

$$\begin{aligned} \delta \rho_\delta^\beta &\rightarrow 0 \text{ in } L^{\frac{\beta+\theta}{\beta}}(\tilde{\Omega} \times (0, T) \times D), \\ \rho_\delta^\gamma &\rightarrow \overline{\rho^\gamma} \text{ weakly in } L^{\frac{\gamma+\theta}{\gamma}}(\tilde{\Omega} \times (0, T) \times D). \end{aligned} \quad (5.6)$$

Similar to Section 4.4, (5.5) implies that

$$\begin{aligned} \rho_\delta &\rightarrow \rho \text{ in } C([0, T]; L_w^\gamma(D)) \cap C([0, T]; H^{-1}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.}, \\ u_\delta &\rightarrow u \text{ in } L^2(0, T; H^1(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.}, \\ \rho_\delta u_\delta &\rightarrow \rho u \text{ in } C([0, T]; L_w^{\frac{2\gamma}{\gamma+1}}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \end{aligned} \quad (5.7)$$

Since $\gamma > \frac{3}{2}$, then $\frac{2\gamma}{\gamma+1} > \frac{6}{5}$. By (5.7), one has

$$\rho_\delta u_\delta^i u_\delta^j \rightarrow \rho u^i u^j \text{ in } \mathcal{D}'((0, T) \times D) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (5.8)$$

By (5.5), we know that

$$B_\delta \rightarrow B \text{ in } L^2(0, T; L^2(D)) \text{ and } B_\delta \rightharpoonup B \text{ in } L^2(0, T; H^1(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (5.9)$$

Moreover, from (3.1)₃ and Proposition 4.6, by Aubin-Lions Lemma, one has

$$B_\delta \rightarrow B \text{ in } C([0, T]; H^{-1}(D)) \text{ } \tilde{\mathbb{P}} - \text{a.s.}$$

Then we have

$$(\nabla \times B_\delta) \times B_\delta \rightarrow (\nabla \times B) \times B \text{ in } \mathcal{D}'([0, T] \times D) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (5.10)$$

Similarly, it follows from (5.7), and (5.9) that

$$\nabla \times (u_\delta \times B_\delta) \rightarrow \nabla \times (u \times B) \text{ in } \mathcal{D}'([0, T] \times D) \text{ } \tilde{\mathbb{P}} - \text{a.s.} \quad (5.11)$$

Now, it remains to show the strong convergence of ρ_δ in $L^1(\tilde{\Omega} \times (0, T) \times D)$, i.e., $\overline{\rho^\gamma} = \rho^\gamma$ for proving Theorem 1.2. First, we introduce the cut off functions:

$$T_k(z) = kT \left(\frac{z}{k} \right), k = 1, 2, 3, \dots,$$

where

$$T(z) = \begin{cases} z, & z \leq 1 \\ 2, & z \geq 3 \end{cases} \in C^\infty(\mathbb{R}^3), \text{ concave in } z \in \mathbb{R}.$$

Since (ρ_δ, u_δ) is a renormalized solution of (4.27)₁ in $\mathcal{D}'((0, T) \times \mathbb{R}^3)$ $\tilde{\text{P}}$ -a.s., choosing $b(z) = T_k(z)$, one deduces that in $\mathcal{D}'((0, T) \times \mathbb{R}^3)$ $\tilde{\text{P}}$ -a.s.

$$\partial_t T_k(\rho_\delta) + \operatorname{div}(T_k(\rho_\delta)u_\delta) + [T_k'(\rho_\delta)\rho_\delta - T_k(\rho_\delta)]\operatorname{div}u_\delta = 0. \quad (5.12)$$

Noting that for any fixed k , $T_k(\rho_\delta) \in L^p(\tilde{\Omega}, L^\infty((0, T) \times D))$ and $T_k(\rho_\delta) \rightarrow \overline{T_k(\rho)}$ weakly star in $L^p(\tilde{\Omega}, L^\infty((0, T) \times D))$ and $\partial_t T_k(\rho_\delta)$ satisfies the equation (5.12), by Aubin-Lions Lemma, as $\delta \rightarrow 0$, for all $1 \leq p < \infty$, we can infer that $\tilde{\text{P}}$ -a.s.

$$T_k(\rho_\delta) \rightarrow \overline{T_k(\rho)} \quad \text{in } C_w([0, T]; L^p(D)) \cap C([0, T]; H^{-1}(D)). \quad (5.13)$$

Letting $\delta \rightarrow 0+$ in (5.12) and by using (5.13), we have $\tilde{\text{P}}$ -a.s.

$$\partial_t \overline{T_k(\rho)} + \operatorname{div} \left(\overline{T_k(\rho)u} \right) + \overline{[T_k'(\rho)\rho - T_k(\rho)]\operatorname{div}u} = 0, \quad (5.14)$$

in $\mathcal{D}'((0, T) \times \mathbb{R}^3)$, where

$$[T_k'(\rho_\delta)\rho_\delta - T_k(\rho_\delta)]\operatorname{div}u_\delta \rightharpoonup \overline{[T_k'(\rho)\rho - T_k(\rho)]\operatorname{div}u} \quad \text{in } L^2(\tilde{\Omega}, L^2((0, T) \times D)).$$

Here $\overline{h(\rho)}$ is the weak limit of $h(\rho_\delta)$.

5.4. The effective viscous flux. Similar to the Section 4.5, we have the following lemma:

Lemma 5.3. Let $(\rho_\delta, u_\delta, B_\delta)$ be the sequence of approximate solutions constructed in Proposition 4.6. Then for any $\psi \in \mathcal{D}(0, T)$, $\phi \in \mathcal{D}(D)$,

$$\begin{aligned} & \lim_{\delta \rightarrow 0+} \tilde{\text{E}} \int_0^T \psi \int_D \phi [a\rho_\delta^\gamma - (\lambda + 2\mu)\operatorname{div}u_\delta] T_k(\rho_\delta) dx dt \\ &= \tilde{\text{E}} \int_0^T \psi \int_D \phi [a\overline{\rho^\gamma} - (\lambda + 2\mu)\operatorname{div}u] \overline{T_k(\rho)} dx dt. \end{aligned}$$

Here we have used (5.14) and the property under (4.20).

5.5. The amplitude of oscillations. In this subsection, we have the following lemma in the sense of expectation as in [38]:

Lemma 5.4. There exists a constant C independent of $k \geq 1$ such that

$$\overline{\lim}_{\delta \rightarrow 0+} \tilde{\text{E}} \|T_k(\rho_\delta) - T_k(\rho)\|_{L^{\gamma+1}((0, T) \times D)}^{\gamma+1} \leq C.$$

5.6. The renormalized solutions. In this subsection, we shall use Lemma 5.4 and the cut-off function technique to show the following crucial lemma as in [38]:

Lemma 5.5. Suppose (ρ, u) be zero outside D , then (ρ, u) solves the continuity equation in the sense of renormalized solutions, that is,

$$\partial_t b(\rho) + \operatorname{div}(b(\rho)u) + [b'(\rho)\rho - b(\rho)] \operatorname{div}u = 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^3),$$

$\tilde{\text{P}}$ -a.s. for any b satisfying the conditions on b .

5.7. Strong convergence of the density. This subsection is devoted to completing the proof of Theorem 1.2. First, we introduce a family of functions L_k as

$$L_k(z) = \begin{cases} z \ln z, & \text{for } 0 \leq z < k, \\ z \ln k + z \int_k^z \frac{T_k(s)}{s^2} ds, & \text{for } z \geq k, \end{cases} \in C^1(\mathbb{R}_+) \cap C^0[0, \infty).$$

Then, for large enough z , $L_k(z)$ is a linear function. More precisely, for $z \geq 3k$, $L_k(z) = \beta_k z - 2k$. Here

$$\beta_k = \ln k + \int_k^{3k} \frac{T_k(s)}{s^2} ds + \frac{2}{3}.$$

Denote $b_k(z) = L_k(z) - \beta_k z$. Then $b_k(z) \in C^1(\mathbb{R}_+) \cap C^0[0, \infty)$, and $b'_k(z) = 0$ when z is large enough. Meanwhile, $b'_k(z) - b_k(z) = T_k(z)$. Taking $b(z) = b_k(z)$, the fact that (ρ_δ, u_δ) is renormalized solution of (4.27)₁ yields that

$$\partial_t L_k(\rho_\delta) + \operatorname{div}(L_k(\rho_\delta)u_\delta) + T_k(\rho_\delta)\operatorname{div}u_\delta = 0. \quad (5.15)$$

As in (5.15), it follows from the continuity equation and Lemma 5.5 that

$$\partial_t L_k(\rho) + \operatorname{div}(L_k(\rho)u) + T_k(\rho)\operatorname{div}u = 0 \quad \text{in } \mathcal{D}'((0, T) \times D). \quad (5.16)$$

Since $L_k(z)$ is a linear function when z is large enough, by (5.15) and Aubin-Lions Lemma, then it holds that $\tilde{\mathbb{P}}$ -a.s.

$$L_k(\rho_\delta) \rightarrow \overline{L_k(\rho)} \quad \text{in } C_w([0, T]; L^\gamma(D)) \cap C^0([0, T]; H^{-1}(D)), \quad (5.17)$$

as $\delta \rightarrow 0$. Taking the difference of (5.15) and (5.16) and then taking $\phi \in \mathcal{D}(D)$ as a test function, we have

$$\begin{aligned} \tilde{\mathbb{E}} \int_D (L_k(\rho_\delta) - L_k(\rho))(t)\phi dx &= \tilde{\mathbb{E}} \int_D (L_k(\rho_{0,\delta}) - L_k(\rho_0))(t)\phi dx \\ &+ \tilde{\mathbb{E}} \int_0^t \int_D (L_k(\rho_\delta)u_\delta - L_k(\rho)u) \cdot \nabla \phi + (T_k(\rho)\operatorname{div}u - T_k(\rho_\delta)\operatorname{div}u_\delta)\phi dx dt. \end{aligned} \quad (5.18)$$

Letting $\delta \rightarrow 0$ in (5.18), noting that $L_k(\rho_{0,\delta}) - L_k(\rho_0) \rightarrow 0$ as $\delta \rightarrow 0$, by (5.17), one has

$$\begin{aligned} \tilde{\mathbb{E}} \int_D (\overline{L_k(\rho)} - L_k(\rho))(t)\phi dx &= \tilde{\mathbb{E}} \int_0^t \int_D (\overline{L_k(\rho)}u - L_k(\rho)u) \cdot \nabla \phi dx dt \\ &+ \lim_{\delta \rightarrow 0^+} \tilde{\mathbb{E}} \int_0^t \int_D [T_k(\rho)\operatorname{div}u - T_k(\rho_\delta)\operatorname{div}u_\delta]\phi dx ds, \quad \forall \phi \in \mathcal{D}(D). \end{aligned} \quad (5.19)$$

By approximating ϕ in the above inequality, similar to Lemma 4.5, taking $\phi = 1$ in (5.19), $u|_{\partial D} = 0$, Lemma 5.3 and Lemma 5.4 imply

$$\begin{aligned} &\tilde{\mathbb{E}} \int_D (\overline{L_k(\rho)} - L_k(\rho))(t) dx \\ &= \tilde{\mathbb{E}} \int_0^t \int_D T_k(\rho)\operatorname{div}u dx dt - \lim_{\delta \rightarrow 0} \tilde{\mathbb{E}} \int_0^t \int_D T_k(\rho_\delta)\operatorname{div}u_\delta dx dt \\ &\leq \tilde{\mathbb{E}} \int_0^T \int_D (T_k(\rho) - \overline{T_k(\rho)})\operatorname{div}u dx dt \\ &\lesssim \tilde{\mathbb{E}} \left(\|\operatorname{div}u\|_{L^2(\{\rho \geq k\})} + \|T_k(\rho) - \overline{T_k(\rho)}\|_{L^2(\{\rho \leq k\})} \right). \end{aligned} \quad (5.20)$$

On the one hand, since $T_k(\rho) \geq \overline{T_k(\rho)}$ and $T_k(\rho) \leq \rho$, by (5.20) in [38], we have

$$\begin{aligned} \tilde{\mathbb{E}} \|T_k(\rho) - \overline{T_k(\rho)}\|_{L^1(\{\rho \leq k\})} &= \tilde{\mathbb{E}} \|\rho - \overline{T_k(\rho)}\|_{L^1(\{\rho \leq k\})} \\ &\leq \tilde{\mathbb{E}} \|\rho - \overline{T_k(\rho)}\|_{L^1(0,T;D)} \leq 2Ck^{-(\gamma-1)} \rightarrow 0, \end{aligned}$$

as $k \rightarrow \infty$. By interpolation and Lemma 5.4, we have

$$\begin{aligned} &\tilde{\mathbb{E}} \|T_k(\rho) - \overline{T_k(\rho)}\|_{L^2(\{\rho \leq k\})} \\ &\leq \left(\tilde{\mathbb{E}} \|T_k(\rho) - \overline{T_k(\rho)}\|_{L^1(\{\rho \leq k\})} \right)^{\frac{\gamma-1}{2\gamma}} \left(\tilde{\mathbb{E}} \|T_k(\rho) - \overline{T_k(\rho)}\|_{L^{\gamma+1}(\{\rho \leq k\})}^{\gamma+1} \right)^{\frac{1}{2\gamma}} \rightarrow 0. \end{aligned}$$

This and $u \in L^2(\tilde{\Omega}, L^2(0, T; H_0^1(D)))$ imply that the right-hand side of (5.20) goes to zero as $k \rightarrow \infty$. Therefore,

$$\lim_{k \rightarrow \infty} \tilde{\mathbb{E}} \int_D \left(\overline{L_k(\rho)} - L_k(\rho) \right) dx \leq 0, \quad t \in [0, T]. \quad (5.21)$$

On the other hand, since $L_k(\rho)$ is linear when ρ is large enough, we have

$$\begin{aligned} \tilde{\mathbb{E}} \|L_k(\rho) - \rho \ln \rho\|_{L^1(0,T;L^1(D))} &\leq \tilde{\mathbb{E}} \int_0^T \int_{\{\rho \geq k\}} |L_k(\rho) - \rho \ln \rho| dx dt \\ &\leq \tilde{\mathbb{E}} \int_0^T \int_{\{\rho \geq k\}} (|L_k(\rho)| + |\rho \ln \rho|) dx dt \lesssim \tilde{\mathbb{E}} \int_0^T \int_{\{\rho \geq k\}} |\rho \ln \rho| \rightarrow 0, \end{aligned} \quad (5.22)$$

as $k \rightarrow \infty$.

Similarly, for small enough ε , one has

$$\begin{aligned} \tilde{\mathbb{E}} \|L_k(\rho_\delta) - \rho_\delta \ln \rho_\delta\|_{L^1(0,T;L^1(D))} &\leq \tilde{\mathbb{E}} \int_0^T \int_{\{\rho_\delta \geq k\}} |L_k(\rho_\delta) - \rho_\delta \ln \rho_\delta| dx dt \\ &\leq \tilde{\mathbb{E}} \int_0^T \int_{\{\rho_\delta \geq k\}} \frac{|L_k(\rho_\delta)| + |\rho_\delta \ln \rho_\delta|}{\rho_\delta^\gamma} \rho_\delta^\gamma dx dt \leq C(\varepsilon) \tilde{\mathbb{E}} \int_0^T \int_{\{\rho_\delta \geq k\}} \frac{\rho_\delta^\gamma}{\rho_\delta^{\gamma-1-\varepsilon}} dx dt \\ &\leq C(\varepsilon) k^{1-\gamma+\varepsilon} \tilde{\mathbb{E}} \int_0^T \int_{\{\rho_\delta \geq k\}} \rho_\delta^\gamma dx dt \leq C(\varepsilon) k^{1-\gamma+\varepsilon} \rightarrow 0 \quad \text{as } k \rightarrow \infty, \end{aligned}$$

uniformly in δ . Let $k \rightarrow \infty$ in the above inequality, then the weak lower semi-continuity of norm implies that

$$\tilde{\mathbb{E}} \|\overline{L_k(\rho)} - \overline{\rho \ln \rho}\|_{L^1(0,T;L^1(D))} \rightarrow 0, \quad \text{uniformly in } \delta. \quad (5.23)$$

Combining (5.21)-(5.23), we have

$$\tilde{\mathbb{E}} \int_D (\overline{\rho \ln \rho} - \rho \ln \rho)(x, t) dx \leq 0,$$

this together with $\rho \ln \rho \leq \overline{\rho \ln \rho}$ imply that

$$\overline{\rho \ln \rho}(t) = \rho \ln \rho(t) \quad \text{for all } t \in [0, T] \text{ a.e..}$$

From this, by [13, Theorem 2.11], we can infer that

$$\rho_\delta \rightarrow \rho \quad \text{almost everywhere in } \tilde{\Omega} \times [0, T] \times D.$$

It follows from Proposition 4.6 and [13, Proposition 2.1] that

$$\rho_\delta \rightarrow \rho \quad \text{weakly in } L^1(\tilde{\Omega} \times (0, T) \times D),$$

By [13, Theorem 2.10], for any $\eta > 0$ and all $\delta > 0$, there exists $\sigma > 0$ such that $\int_F \rho_\delta(t, x) dx dt d\tilde{\mathbb{P}} < \eta$ for any measurable set $F \subset \tilde{\Omega} \times (0, T) \times D$ with $\text{meas}\{F\} < \sigma$. On the other hand, by Egorov's Theorem, there exists a measurable set $F_\sigma \subset \tilde{\Omega} \times (0, T) \times D$ such that $\text{meas}\{F_\sigma\} < \sigma$ and $\rho_\delta \rightarrow \rho$ uniformly in $\tilde{\Omega} \times (0, T) \times D - F_\sigma$. Note that

$$\begin{aligned} \tilde{\mathbb{E}} \int_0^T \int_D |\rho_\delta - \rho| dx dt &\leq \iiint_{F_\sigma} |\rho_\delta - \rho| dx dt d\tilde{\mathbb{P}} + \iiint_{\tilde{\Omega} \times (0, T) \times D - F_\sigma} |\rho_\delta - \rho| dx dt d\tilde{\mathbb{P}} \\ &\leq 2\eta + T|D| \sup_{\tilde{\Omega} \times (0, T) \times D - F_\sigma} |\rho_\delta - \rho|. \end{aligned} \quad (5.24)$$

Then (5.24) tends to 0 as $\delta \rightarrow 0$ and $\eta \rightarrow 0$. This implies the strong convergence of the sequence ρ_δ in $L^1(\tilde{\Omega} \times (0, T) \times D)$, that is,

$$\overline{\rho^\gamma} = \rho^\gamma. \quad (5.25)$$

Similar to the proof of (3.37) and (3.40) in Section 3.4, by using (1.5), (5.7), (5.9), (5.25) and Hölder's inequality, we have

$$\begin{aligned} \langle f_k(\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, x), \phi \rangle &\rightarrow \langle f_k(\rho, \rho u, x), \phi \rangle \text{ in } L^1([0, T]) \tilde{\mathbb{P}} - \text{a.s.}, \\ \langle g_k(B_\varepsilon, x), \phi \rangle &\rightarrow \langle g_k(B, x), \phi \rangle \text{ in } L^1([0, T]) \tilde{\mathbb{P}} - \text{a.s.}, \end{aligned} \quad (5.26)$$

and

$$\begin{aligned} \sum_{k \geq 1} \langle f_k(\rho_\delta, \rho_\delta u_\delta, x), \phi \rangle^2 &\rightarrow \sum_{k \geq 1} \langle f_k(\rho, \rho u, x), \phi \rangle^2 \text{ in } L^1([0, T]) \tilde{\mathbb{P}} - \text{a.s.}, \\ \sum_{k \geq 1} \langle g_k(B_\delta, x), \phi \rangle^2 &\rightarrow \sum_{k \geq 1} \langle g_k(B, x), \phi \rangle^2 \text{ in } L^1([0, T]) \tilde{\mathbb{P}} - \text{a.s.} \end{aligned} \quad (5.27)$$

As in Section 4, by using (5.6)-(5.11), (5.25)-(5.27), we deduce that $((\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}), \beta_k, \rho, u, B)$ satisfies the following equations:

$$\begin{cases} d\rho + \text{div}(\rho u) dt = 0, \\ (\rho u)_t + \text{div}(\rho u \otimes u) + a \nabla \rho^\gamma - \mu \Delta u - (\lambda + \mu) \nabla \text{div} u = (\nabla \times B) \times B + \sum_{k \geq 1} f_k(\rho, \rho u, x) \dot{\beta}_k^1, \\ B_t - (\nabla \times u) \times B - \nu \Delta B = \sum_{k \geq 1} g_k(B, x) \dot{\beta}_k^2, \end{cases}$$

in $\mathcal{D}'((0, T) \times D)$ $\tilde{\mathbb{P}}$ -a.s.. Here we formally denote $\dot{W} := dW/dt$. The proof of Theorem 1.2 is thus completed.

ACKNOWLEDGEMENT

H. Wang is supported by the National Natural Science Foundation of China (No. 11901066), the Natural Science Foundation of Chongqing (No. cstc2019jcyj-msxmX0167) and Project No. 2019CDXYST0015 and No. 2020CDJQY-A040 supported by the Fundamental Research Funds for the Central Universities.

REFERENCES

- [1] V. Barbu and G. Da Prato, *Existence and ergodicity for the 2D stochastic MHD equations*, Preprint di Matematica.
- [2] Z. Brzeźniak and E. Hausenblas, *Uniqueness in law of the Itô integral with respect to Lévy Noise*, In: Seminar on Stochastic Analysis, Random Fields and Applications VI Progress in Probability, 63 (2011), 37-57.

- [3] Z. Brzeźniak and M. Ondreját, *Stochastic geometric wave equations with values in compact Riemannian homogeneous space*, The Annals of Probability, 41 (2013), 1938-1977.
- [4] A. Bensoussan and R. Temam, *Équations stochastiques du type Navier-Stokes*, J. Functional Analysis, 13 (1973), 195-222.
- [5] M.E. Bogovskii, *Solution of some vector analysis problems connected with operators div and grad*, (in Russian), Trudy Sem. S. L. Sobolev, 80 (1980), 5-40.
- [6] D. Breit, E. Feireisl and M. Hofmanová, *Incompressible limit for compressible fluids with stochastic forcing*, Arch. Ration. Mech. Anal., 222(2) (2016), 895-926.
- [7] D. Breit and M. Hofmanová, *Stochastic Navier-Stokes equations for compressible fluids*, Indiana Univ. Math. J., 65(4) (2016), 1183-1250.
- [8] Q. Chen and Z. Tan, *Global existence and convergence rates of smooth solutions for the compressible magnetohydrodynamical equations*, 72 (2010), 4438-4451.
- [9] G. Da Prato and J. Zabczyk, *Stochastic Equations in Infinite Dimensions*, Cambridge University Press, Cambridge, 1992.
- [10] B. Ducomet and E. Feireisl, *The equations of magnetohydrodynamics: On the interaction between matter and radiation in the evolution of gaseous stars*, Commun. Math. Phys, 266 (2006), 595-629.
- [11] J. Fan and W. Yu, *Strong solution to the compressible magnetohydrodynamic equations with vacuum*, Nonlinear Analysis: Real World Applications, 10 (2009), 392-409.
- [12] J. Fan and W. Yu, *Global variational solutions to the compressible magnetohydrodynamic equations*, Nonlinear Anal., 69 (2008), 3637-3660.
- [13] E. Feireisl, *Dynamics of viscous compressible fluids*, Oxford University Press, Oxford, 2004.
- [14] E. Feireisl, B. Maslowski and A. Novotný, *Compressible fluid flows driven by stochastic forcing*, J. Differential Equations, 254 (2013), 1342-1358.
- [15] E. Feireisl, A. Novotný, and H. Petzeltová, *On the existence of globally defined weak solutions to the Navier-Stokes equations*, J. Math. Fluid Mech., 3 (2001), 358-392.
- [16] G.P. Galdi, *An introduction to the mathematical theory of the Navier-Stokes equations*, Springer-Verlag, New York, 1994.
- [17] X. Hu and D. Wang, *Global solutions to the three-dimensional full compressible magnetohydrodynamic flows*, Commun. Math. Phys., 283 (2008), 255-284.
- [18] X. Hu and D. Wang, *Global existence and large-time behavior of solutions to the three-dimensional equations of compressible magnetohydrodynamic flows*, Arch. Ration. Mech. Anal., 197 (2010), 203-238.
- [19] X. Hu and D. Wang, *Low Mach number limit of viscous compressible magnetohydrodynamic flows*, SIAM J. Math. Anal., 41 (2009), 1272-1294.
- [20] A. Jakubowski, *The a.s. Skorokhod representation for subsequences in nonmetric spaces*, Teor. Veroyatnost. i Primenen., 42 (1997), 209-216.
- [21] S. Jiang, Q. Ju and F. Li, *Incompressible limit of the compressible magnetohydrodynamic equations with periodic boundary conditions*, Commun. Math. Phys, 297 (2010), 371-400.
- [22] O. Kallenberg, *Foundations of Modern Probability*, in: *Probability and its Applications*, Springer-Verlag, New York, 1997.
- [23] I. Karatzas and S.E. Shreve, *Brownian Motion and Stochastic Calculus*, Springer-Verlag, New York, 1991.
- [24] H.J. Kushner, *Numerical Methods for Controlled Stochastic Delay Systems*, Birkhäuser, Boston, 2008.
- [25] F. Li and H. Yu, *Optimal decay rate of classical solutions to the compressible magnetohydrodynamic equations*, Proc. Royal Soc. Edinburgh, 141A (2011), 109-126.
- [26] J.L. Menaldi and S.S. Sritharan, *Stochastic 2-D Navier-Stokes Equation*, Appl Math Optim, 46 (2002), 31-53.
- [27] A. Novotný and I. Straškraba, *Introduction to the Mathematical Theory of Compressible Flow*, Oxford University Press Inc., New York, 2004.
- [28] Martin Ondreját, *Stochastic nonlinear wave equations in local Sobolev spaces*, Electronic Journal of Probability, 15 (2010), 1041-1091.
- [29] P.L. Lions, *Mathematical topics in fluid dynamics, Vol. 1, Incompressible models*, Oxford Science Publication, Oxford, 1996.
- [30] P.L. Lions, *Mathematical topics in fluid dynamics, Vol. 2, Compressible models*, Oxford Science Publication, Oxford, 1998.

- [31] X. Pu and B. Guo, *Global existence and convergence rates of smooth solutions for the full compressible MHD equations*, *Z. Angew. Math. Phys.*, 64 (2013), 519-538.
- [32] M. Sango, *Magnetohydrodynamic turbulent flows: Existence results*, *Physica D*, 239 (2010), 912-923.
- [33] J. Simon, *Compact sets in the space $L^p(0, T; B)$* , *Ann. Mat. Pura Appl.*, 146 (1987), 65-96.
- [34] S. A. Smith, *Random Perturbations of Viscous, Compressible Fluids: Global Existence of Weak Solutions*, *SIAM J. Math. Anal.* 49 (2017), 4521-4578.
- [35] P. Sundar, *Stochastic Magnetohydrodynamic system perturbed by general noise*, *Communications on Stochastic Analysis*, 365 (2016), 1219-1256.
- [36] Z. Tan, H.Q. Wang, *Optimal decay rates of the compressible MHD equations*, *Nonlinear Analysis: Real World Applications*, 14 (2013), 188-201.
- [37] Z. Tan, Q.J. Xu and H.Q. Wang, *Global existence and convergence rates for the compressible magnetohydrodynamic equations without heat conductivity*, *Discrete Continuous Dynam. Systems-A*, 35(10) (2015), 5083-5105.
- [38] D.H. Wang and H.Q. Wang, *Global existence of martingale solutions to the three-dimensional stochastic compressible Navier-Stokes equations*, *Differential and Integral Equations*, 28(11-12) (2015), 1105-1154.

COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, CHONGQING 400044, CHINA.
Email address: wanghuaqiao@cqu.edu.cn