

Quantitative unique continuation and observability on an equidistributed set for the diffusion equation in \mathbb{R}^N

Yueliang Duan ^{*}, Huaiqiang Yu [†], Can Zhang [‡]

December 10, 2021

Abstract

In this paper, we obtain a quantitative estimate of unique continuation and an observability inequality from an equidistributed set for solutions of the diffusion equation in the whole space \mathbb{R}^N . This kind of observability indicates that the total energy of solutions can be controlled by the energy localized in a measurable subset, which is equidistributed over the whole space. The proof of our results is based on an interesting reduction method [18, 22], as well as the propagation of smallness for the gradient of solutions to elliptic equations [24].

Keywords. Unique continuation, Observability, Equidistributed set, Diffusion equation.

2010 AMS Subject Classifications. 35B60, 35K10, 93C20.

Contents

1	Introduction	2
1.1	Main results	2
1.2	Motivation and novelty	3
1.3	Plan of this paper	4
2	Some auxiliary lemmas	4
2.1	Local energy estimates	4
2.2	Stability estimate for elliptic equations	11
3	Proof of Theorem 1.1	13
4	Proof of Theorem 1.2	19
5	Appendix: Proof of (2.23)	21

^{*}School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China; e-mail: ylduan@whu.edu.cn.

[†]School of Mathematics, Tianjin University, Tianjin 300354, China; e-mail: huaiqiangyu@tju.edu.cn.

[‡]School of Mathematics and Statistics; Computational Science Hubei Key Laboratory, Wuhan University, Wuhan 430072, China; e-mail: zhangcansx@163.com

1 Introduction

Let N be a positive integer, $\mathbb{N}^+ := \{1, 2, \dots\}$ and $\mathbb{R}^+ := (0, +\infty)$. Consider the linear diffusion equation over \mathbb{R}^N :

$$\begin{cases} \partial_t u(x, t) - \sum_{j,k=1}^N \partial_j (a_{jk}(x) \partial_k u(x, t)) = 0 & \text{in } \mathbb{R}^N \times \mathbb{R}^+, \\ u(x, 0) = u_0(x) & \text{in } \mathbb{R}^N, \end{cases} \quad (1.1)$$

where $u_0 \in L^2(\mathbb{R}^N)$, $a_{jk}(x) = a_{kj}(x)$ for all $j, k = 1, \dots, N$, and all $x \in \mathbb{R}^N$. We assume the matrix-valued function $A(\cdot) = (a_{jk}(\cdot))_{j,k=1}^N$ is Lipschitz continuous and satisfies the uniform ellipticity condition, i.e., there is a constant $\lambda \geq 1$ such that

$$|a_{jk}(x) - a_{jk}(y)| \leq \lambda |x - y| \quad \text{and} \quad \lambda^{-1} |\xi|^2 \leq \sum_{j,k=1}^N a_{jk}(x) \xi_j \xi_k \leq \lambda |\xi|^2 \quad (1.2)$$

for all $j, k = 1, \dots, N$, $x, y \in \mathbb{R}^N$, and all $\xi = (\xi_1, \xi_2, \dots, \xi_N) \in \mathbb{R}^N$.

According to [4, Theorem 10.9], (1.1) has a unique solution $u \in L^2(\mathbb{R}^+; H^1(\mathbb{R}^N)) \cap C([0, +\infty); L^2(\mathbb{R}^N))$. Furthermore, if $u_0 \in H^1(\mathbb{R}^N)$, then $u \in C([0, +\infty); H^1(\mathbb{R}^N))$ and $\partial_t u \in L^2(\mathbb{R}^+; L^2(\mathbb{R}^N))$.

Throughout the paper, $B_R(x_0)$ stands for an open ball in \mathbb{R}^N with the center x_0 and of radius $R > 0$, while $B_R(x_0, 0)$ stands for an open ball in \mathbb{R}^{N+1} with the center $(x_0, 0) \in \mathbb{R}^N \times \mathbb{R}$ and of radius $R > 0$. We denote the usual inner product and norm in $L^2(B_R(x_0))$ by $\langle \cdot, \cdot \rangle_{L^2(B_R(x_0))}$ and $\|\cdot\|_{L^2(B_R(x_0))}$, respectively. Define the cube with center $x_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,N})$ and side length $R > 0$ as $Q_R(x_0) := \{v = (v_1, v_2, \dots, v_N) : |v_i - x_{0,i}| \leq R, \forall i = 1, 2, \dots, N\}$. Denote by $\text{int}(Q_R(x_0))$ the interior of $Q_R(x_0)$, and by $\partial B_R(x_0)$ the boundary of $B_R(x_0)$. Write \bar{z} for the conjugate of a complex number $z \in \mathbb{C}$, and $|\omega|$ for the Lebesgue measure for a measurable set $\omega \subset \mathbb{R}^N$.

1.1 Main results

The main results of this paper concerning the quantitative estimate of unique continuation and the observability inequality for solutions of (1.1) are stated as follows.

Theorem 1.1. *Let $T > 0$, $\rho > 0$ and $0 < r < +\infty$. Let $\{x_i\}_{i \in \mathbb{N}^+} \subset \mathbb{R}^N$ be arbitrarily fixed so that*

$$\mathbb{R}^N = \bigcup_{i \in \mathbb{N}^+} Q_r(x_i) \quad \text{with} \quad \text{int}(Q_r(x_i)) \cap \text{int}(Q_r(x_j)) = \emptyset \quad \text{for each } i \neq j \in \mathbb{N}^+.$$

Let $\{\omega_i\}_{i \in \mathbb{N}^+}$ be some N -dimensional Lebesgue measurable sets in \mathbb{R}^N so that

$$\omega_i \subset B_{\frac{r}{2}}(x_i) \quad \text{and} \quad \frac{|\omega_i|}{|B_r(x_i)|} \geq \rho \quad \text{for each } i \in \mathbb{N}^+.$$

Then there exist constants $C = C(r, N, \lambda, \rho) > 0$ and $\sigma = \sigma(r, N, \lambda, \rho) \in (0, 1)$ such that for any $u_0 \in L^2(\mathbb{R}^N)$, the corresponding solution u of (1.1) satisfies

$$\int_{\mathbb{R}^N} |u(x, T)|^2 dx \leq C e^{C(T+\frac{1}{T})} \left(\int_{\omega} |u(x, T)|^2 dx \right)^{\sigma} \left(\int_{\mathbb{R}^N} |u_0(x)|^2 dx \right)^{1-\sigma},$$

where $\omega := \bigcup_{i \in \mathbb{N}^+} \omega_i$.

Theorem 1.2. *Let E be a subset of positive Lebesgue measure in $(0, T)$. Under the same assumptions in Theorem 1.1, then there exist constants $C = C(r, N, \lambda, \rho) > 0$ and $\tilde{C} = \tilde{C}(E, r, N, \lambda, \rho) > 0$ such that for any $u_0 \in L^2(\mathbb{R}^N)$, the corresponding solution u of (1.1) satisfies*

$$\int_{\mathbb{R}^N} |u(x, T)|^2 dx \leq e^{\tilde{C} + CT} \int_{\omega \times E} |u(x, t)|^2 dx dt.$$

Several remarks are given below.

Remark 1.3. When $E = (0, T)$, the constant $\tilde{C}(E, r, N, \lambda, \rho)$ in Theorem 1.2 can take the form $\tilde{C}(r, N, \lambda, \rho)/T$.

Remark 1.4. When $A(\cdot) \equiv I$ (the $N \times N$ order identity matrix) in the equation (1.1), the above two theorems have been already established in [34] for solutions of the heat equation in \mathbb{R}^N , when the observation set is thick. The proofs there are based on quantitative estimates from measurable sets for real analytic functions, and uncertainty principle. To the best of our knowledge, however, this approach cannot be extended to the case of variable coefficients. In the present paper, we built a new approach to obtain the observability inequality for the diffusion equation (1.1).

Remark 1.5. Recall that a subset $\mathcal{O} \subset \mathbb{R}^N$ is called thick with parameters $L > 0$ and $\gamma > 0$ means that in each cube $Q_L \subset \mathbb{R}^N$ with the length L , the N -dimensional Lebesgue measure of $\mathcal{O} \cap Q_L$ is bigger than or equals to γL^N . As the thick set \mathcal{O} is almost dense in the whole space, one may imagine that such kind of thick set has a close relationship with the equidistributed set ω as defined in Theorem 1.1. Indeed, an equidistributed set ω is clearly a thick set. On the other hand, if \mathcal{O} is a thick set, then one could construct an equidistributed set ω to be such that $\omega \subset \mathcal{O}$.

Remark 1.6. By [34, Lemma 2.5], one can obtain that the equidistributed observation set is also a necessary condition such that the observability inequality holds for the diffusion equation (1.1).

Remark 1.7. It is also worth mentioning that the authors in [34, 9] have proved that the observability inequality for the heat equation holds if and only if the observation set is thick. Combined with Remarks 1.5 and 1.6 above, one can finally conclude that the observability inequality for the diffusion equation (1.1) holds if and only if the observation set is thick or equidistributed in \mathbb{R}^N .

1.2 Motivation and novelty

The observability inequality for linear parabolic equations on bounded domains are previously studied in a large number of publications. When E is the whole time interval and the observation region ω is a nonempty open subset, we refer the readers to [15, 14] and a vast number of references therein for the observability inequality for parabolic equations. Their approach is mainly based on the Carleman inequality method. When E is only a subset of positive Lebesgue measure in the time interval and the observation region ω is a non-empty open subset, we refer the readers to [2, 32, 33, 35] for the observability inequality for parabolic equations. More generally, when the observation subdomain is a measurable subset of positive measure in the space and time variables, we refer the readers to [12, 13, 1] for the observability inequality for analytic parabolic equations. The proofs of those inequalities are mainly based on the parabolic frequency function method, and the propagation of smallness estimate for real analytic functions. Note that the solution to (1.1) is obviously not spatial real-analyticity any more, and thus we cannot derive the observability inequality from measurable sets by the real analytic technique.

The studies on the observability inequality for parabolic equations on unbounded domains are rather few in last decades. The observability inequality may not be true when the heat equation is evolving in the whole space and the observation subdomain is only a bounded and open subset (see, e.g., [27, 28, 29]). We would like to mention the work [5] for sufficient conditions so that the observability inequalities hold true for heat equations in unbounded domains. It showed that, for some parabolic equations in an unbounded domain $\Omega \subset \mathbb{R}^N$, the observability inequality holds when observations are made over an open subset $\omega \subset \Omega$, with $\Omega \setminus \omega$ bounded. The proofs are based on Carleman estimates. For other similar results, we refer the reader to [3, 6, 17, 19, 36].

Recently, [34] and [9] independently obtained the observability inequality for the pure heat equation on the whole space, where the observation is a thick set (see Remark 1.5). This could be extended to the time-independent parabolic equation associated to the Schrödinger operator with analytic coefficients (see [10, 20, 30]). The methods utilized in these papers are all based on the spectral inequality (or uncertainty principle). Unfortunately, they are not valid any more for the case that the coefficients in parabolic equations are non-analytic.

A Hölder type propagation of smallness for solutions, as well as for their gradients, on a possibly lower dimensional Hausdorff subset to general second order elliptic equations has been established in [24]. The approach in [24] does not base on the analyticity of solutions. This result stimulates our motivation and opens a possible way to establish the observability inequality from measurable sets for solutions of general parabolic equations, getting rid of the above-mentioned technique of analyticity.

Recently, quantitative estimates of unique continuation for solutions of second order parabolic equations, such as the doubling property, as well as the two-ball and one-cylinder inequality, have been well understood (see, e.g., [2], [7], [8], [11], [18], [22], [31], [32]).

The main effort of this paper is to prove a locally quantitative estimate of strong unique continuation for solutions to the diffusion equation, by utilizing the propagation of smallness estimate for gradients of solutions to elliptic equations. Combing with the above local result and the geometry of the equidistributed observation, we obtain a globally quantitative estimate at one time point for solutions of the diffusion equation. We finally apply the telescoping series method to prove the desired observability inequality.

1.3 Plan of this paper

The rest of this paper is organized as follows. In Section 2, we give several auxiliary lemmas. Sections 3 and 4 prove Theorem 1.1 and Theorem 1.2, respectively.

2 Some auxiliary lemmas

In this section, we first show two exponential estimates (see Lemmas 2.1 and 2.2 below) in Section 2.1, and then give a quantitative estimate of Cauchy uniqueness (see Lemma 2.6 below) in Section 2.2.

2.1 Local energy estimates

Let $T > 0$ and $\tau \in (0, T)$ be arbitrarily fixed. Let $\eta \in C^\infty(\mathbb{R}^+; [0, 1])$ be a cutoff function satisfying

$$\begin{cases} \eta = 1 & \text{in } [0, \tau], \\ \eta = 0 & \text{in } [T, +\infty), \\ |\eta_t| \leq \frac{C_1}{T-\tau} & \text{in } (\tau, T), \end{cases} \quad (2.1)$$

where the positive constant C_1 is independent of τ and T .

Let $R > 0$ and $x_0 \in \mathbb{R}^N$. Let v be the solution to

$$\begin{cases} v_t - \operatorname{div}(A(x)\nabla v) = 0 & \text{in } B_R(x_0) \times \mathbb{R}^+, \\ v = \eta u & \text{on } \partial B_R(x_0) \times \mathbb{R}^+, \\ v(\cdot, 0) = 0 & \text{in } B_R(x_0), \end{cases} \quad (2.2)$$

where u satisfies (1.1) with the initial value $u_0 \in H^1(\mathbb{R}^N)$ and η verifies (2.1).

Then, we have the following exponential decay result.

Lemma 2.1. *There exist positive constants $\hat{C}_1 = \hat{C}_1(\lambda)$ and $\hat{C}_2 = \hat{C}_2(\lambda, N)$ such that*

$$\|v(\cdot, t)\|_{H^1(B_R(x_0))} \leq \hat{C}_1 \left(1 + \frac{1}{T}\right)^{\frac{1}{2}} e^{\frac{\hat{C}_1 T}{T-\tau} - \hat{C}_2 R^{-2}(t-T)^+} F(R) \quad \text{for all } t \in \mathbb{R}^+, \quad (2.3)$$

where $(t-T)^+ := \max\{0, t-T\}$ and $F(R) := \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_R(x_0))}$.

Proof. We proceed the proof into two steps as follows.

Step 1. To prove (2.3) when $t \in [0, T]$.

Setting $w = v - \eta u$ in $B_R(x_0) \times \mathbb{R}^+$, we see that w verifies

$$\begin{cases} w_s - \operatorname{div}(A(x)\nabla w) = -\eta_s u & \text{in } B_R(x_0) \times \mathbb{R}^+, \\ w = 0 & \text{on } \partial B_R(x_0) \times \mathbb{R}^+, \\ w(\cdot, 0) = -u_0 & \text{in } B_R(x_0). \end{cases} \quad (2.4)$$

We next prove that for each $t \in [0, T]$,

$$\|w(\cdot, t)\|_{L^2(B_R(x_0))} \leq e^{\frac{C_1 T}{T-\tau}} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}. \quad (2.5)$$

Indeed, multiplying first (2.4) by w and then integrating by parts over $B_R(x_0) \times (0, t)$ lead to

$$\begin{aligned} & \|w(\cdot, t)\|_{L^2(B_R(x_0))}^2 + 2 \int_0^t \int_{B_R(x_0)} A \nabla w \cdot \nabla w \, dx ds \\ & \leq \|u_0\|_{L^2(B_R(x_0))}^2 + \frac{C_1 t}{T-\tau} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}^2 + \frac{C_1}{T-\tau} \int_0^t \int_{B_R(x_0)} |w|^2 \, dx ds \\ & \leq \left(1 + \frac{C_1 T}{T-\tau}\right) \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}^2 + \frac{C_1}{T-\tau} \int_0^t \int_{B_R(x_0)} |w|^2 \, dx ds. \end{aligned}$$

Here, we used (2.1). This, together with the uniform ellipticity condition (see (1.2)), implies that

$$\|w(\cdot, t)\|_{L^2(B_R(x_0))}^2 \leq \left(1 + \frac{C_1 T}{T-\tau}\right) \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}^2 + \frac{C_1}{T-\tau} \int_0^t \int_{B_R(x_0)} |w|^2 \, dx ds.$$

By the Gronwall inequality, we get (2.5) immediately. From (2.5) and the definition of w , we know that for any $t \in [0, T]$,

$$\|v(\cdot, t)\|_{L^2(B_R(x_0))} \leq 2e^{\frac{C_1 T}{T-\tau}} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}. \quad (2.6)$$

We next claim that

$$\|\nabla w(\cdot, t)\|_{L^2(B_R(x_0))}^2 \leq \frac{4C_1^2\lambda T}{(T-\tau)^2} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}^2 + \lambda^2 \|\nabla u_0\|_{L^2(B_R(x_0))}^2. \quad (2.7)$$

If it holds true, along with the definition of w , we have that, for each $t \in [0, T]$,

$$\begin{aligned} \|\nabla v(\cdot, t)\|_{L^2(B_R(x_0))}^2 &\leq 2\|\nabla w(\cdot, t)\|_{L^2(B_R(x_0))}^2 + 2\|\nabla u(\cdot, t)\|_{L^2(B_R(x_0))}^2 \\ &\leq 2\lambda^2\|\nabla u_0\|_{L^2(B_R(x_0))}^2 + 2\|\nabla u(\cdot, t)\|_{L^2(B_R(x_0))}^2 \\ &\quad + \frac{8C_1^2\lambda T}{(T-\tau)^2} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}^2 \\ &\leq \left[2 + 2\lambda^2 + \frac{8C_1^2\lambda T}{(T-\tau)^2}\right] \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_R(x_0))}^2. \end{aligned} \quad (2.8)$$

Hence, the desired estimate (2.3) follows from $\lambda \geq 1$, (2.6) and (2.8) when $t \in [0, T]$ with $\hat{C}_1 = 2(3 + \lambda^2)(1 + 2C_1)$.

The rest is devoted to prove the claim (2.7). In fact, multiplying first (2.4) by w_s and then integrating by parts over $B_R(x_0) \times (0, t)$, we find

$$\begin{aligned} &\int_0^t \int_{B_R(x_0)} |w_s|^2 dx ds + \frac{1}{2} \int_0^t \int_{B_R(x_0)} [\nabla w \cdot (A\nabla w)]_s dx ds \\ &= \int_0^t \int_{B_R(x_0)} |w_s|^2 dx ds + \frac{1}{2} \int_0^t \int_{B_R(x_0)} [(\nabla w)_s \cdot (A\nabla w) + \nabla w \cdot (A\nabla w)_s] dx ds \\ &= \int_0^t \int_{B_R(x_0)} |w_s|^2 dx ds + \int_0^t \int_{B_R(x_0)} \nabla w_s \cdot (A\nabla w) dx ds \\ &= \int_0^t \int_{B_R(x_0)} |w_s|^2 dx ds - \int_0^t \int_{B_R(x_0)} w_s \operatorname{div}(A\nabla w) dx ds \\ &= - \int_0^t \int_{B_R(x_0)} w_s \eta_s u dx ds \\ &\leq \frac{C_1}{\epsilon_1(T-\tau)} \int_0^t \int_{B_R(x_0)} |w_s|^2 dx ds + \frac{C_1\epsilon_1}{T-\tau} \int_0^t \int_{B_R(x_0)} |u|^2 dx ds \quad \text{for any } \epsilon_1 > 0. \end{aligned}$$

Here, we used (2.1). Letting $\epsilon_1 = 2C_1/(T-\tau)$ in the inequality above, combined with the uniform ellipticity condition (1.2), we get that

$$\begin{aligned} &\int_{B_R(x_0)} \nabla w(x, t) \cdot (A(x)\nabla w(x, t)) dx \\ &\leq \int_{B_R(x_0)} \nabla u_0 \cdot (A(x)\nabla u_0) dx + \frac{4C_1^2}{(T-\tau)^2} \int_0^t \int_{B_R(x_0)} |u|^2 dx ds \\ &\leq \lambda \int_{B_R(x_0)} |\nabla u_0|^2 dx + \frac{4C_1^2}{(T-\tau)^2} \int_0^t \int_{B_R(x_0)} |u|^2 dx ds. \end{aligned}$$

By the uniform ellipticity condition (1.2) again, this means that

$$\|\nabla w(\cdot, t)\|_{L^2(B_R(x_0))}^2 \leq \lambda^2 \int_{B_R(x_0)} |\nabla u_0|^2 dx + \frac{4\lambda C_1^2}{(T-\tau)^2} \int_0^t \int_{B_R(x_0)} |u|^2 dx ds$$

$$\leq \lambda^2 \int_{B_R(x_0)} |\nabla u_0|^2 dx + \frac{4C_1^2 \lambda T}{(T-\tau)^2} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))}^2.$$

Thus, (2.7) is true.

Step 2. To prove (2.3) when $t \geq T$.

Denoting $\mathcal{A}(\cdot) := -\operatorname{div}(A(\cdot)\nabla)$ with domain $D(\mathcal{A}) := H_0^1(B_R(x_0)) \cap H^2(B_R(x_0))$, we claim that there is a generic constant $C_2 = C_2(\lambda, N) > 0$ such that

$$\langle \mathcal{A}f, f \rangle_{L^2(B_R(x_0))} \geq C_2 R^{-2} \|f\|_{L^2(B_R(x_0))}^2 \quad \text{for each } f \in D(\mathcal{A}). \quad (2.9)$$

In fact, by the Poincaré inequality

$$\int_{B_R(x_0)} |f|^2 dx \leq \left(\frac{2R}{N}\right)^2 \int_{B_R(x_0)} |\nabla f|^2 dx \quad \text{for each } f \in H_0^1(B_R(x_0)),$$

we derive

$$\langle \mathcal{A}f, f \rangle_{L^2(B_R(x_0))} \geq \lambda^{-1} \int_{B_R(x_0)} |\nabla f|^2 dx \geq \lambda^{-1} \left(\frac{N}{2R}\right)^2 \int_{B_R(x_0)} |f|^2 dx.$$

We can conclude the claim (2.9).

As a consequence of (2.9), we see that the inverse of $(\mathcal{A}, D(\mathcal{A}))$ is positive, self-adjoint and compact in $L^2(B_R(x_0))$. By the spectral theorem for compact self-adjoint operators, there are eigenvalues $\{\mu_i\}_{i \in \mathbb{N}^+} \subset \mathbb{R}^+$ and eigenfunctions $\{f_i\}_{i \in \mathbb{N}^+} \subset H_0^1(B_R(x_0))$, which make up a complete orthogonal basis of $L^2(B_R(x_0))$, such that

$$\begin{cases} -\mathcal{A}f_i = \mu_i f_i \quad \text{and} \quad \|f_i\|_{L^2(B_R(x_0))} = 1 & \text{for each } i \in \mathbb{N}^+, \\ C_2 R^{-2} \leq \mu_1 \leq \mu_2 \leq \dots \leq \mu_i \rightarrow +\infty & \text{as } i \rightarrow +\infty. \end{cases} \quad (2.10)$$

Then, by the formula of Fourier decomposition, the solution w of (2.4) in $[T, +\infty)$ is given by

$$w(\cdot, t) = \sum_{i=1}^{\infty} \langle w(\cdot, T), f_i \rangle_{L^2(B_R(x_0))} e^{-\mu_i(t-T)} f_i \quad \text{in } B_R(x_0) \quad \text{for each } t \in [T, +\infty).$$

Hence, we deduce that for each $t \in [T, +\infty)$,

$$\|w(\cdot, t)\|_{L^2(B_R(x_0))}^2 \leq e^{-C_2 R^{-2}(t-T)} \|w(\cdot, T)\|_{L^2(B_R(x_0))}^2 \quad (2.11)$$

and

$$w_t(\cdot, t) = - \sum_{i=1}^{\infty} \mu_i \langle w(\cdot, T), f_i \rangle_{L^2(B_R(x_0))} e^{-\mu_i(t-T)} f_i.$$

It follows that

$$-\langle w(\cdot, t), w_t(\cdot, t) \rangle_{L^2(B_R(x_0))} = \sum_{i=1}^{\infty} \mu_i |\langle w(\cdot, T), f_i \rangle_{L^2(B_R(x_0))}|^2 e^{-2\mu_i(t-T)}, \quad (2.12)$$

for each $t \in [T, +\infty)$. In particular, by taking $t = T$ in the above identify, we have

$$-\langle w(\cdot, T), w_t(\cdot, T) \rangle_{L^2(B_R(x_0))} = \sum_{i=1}^{\infty} \mu_i |\langle w(\cdot, T), f_i \rangle_{L^2(B_R(x_0))}|^2. \quad (2.13)$$

Meanwhile, it follows from (2.4), (2.1) and (1.2) that

$$-\langle w(\cdot, T), w_t(\cdot, T) \rangle_{L^2(B_R(x_0))} \leq \lambda \|\nabla w(\cdot, T)\|_{L^2(B_R(x_0))}^2. \quad (2.14)$$

From (2.13) and (2.14), we have

$$\sum_{i=1}^{\infty} \mu_i |\langle w(\cdot, T), f_i \rangle_{L^2(B_R(x_0))}|^2 \leq \lambda \|\nabla w(\cdot, T)\|_{L^2(B_R(x_0))}^2.$$

This, together with (2.12) and (2.10), gives

$$-\langle w(\cdot, t), w_t(\cdot, t) \rangle_{L^2(B_R(x_0))} \leq \lambda e^{-2C_2 R^{-2}(t-T)} \|\nabla w(\cdot, T)\|_{L^2(B_R(x_0))}^2, \quad (2.15)$$

for each $t \in [T, +\infty)$. On the other hand, by (2.1), (2.4) and (1.2), we see that for each $t \in [T, +\infty)$,

$$-\langle w(\cdot, t), w_t(\cdot, t) \rangle_{L^2(B_R(x_0))} = \langle w(\cdot, t), \mathcal{A}w(\cdot, t) \rangle_{L^2(B_R(x_0))} \geq \lambda^{-1} \|\nabla w(\cdot, t)\|_{L^2(B_R(x_0))}^2. \quad (2.16)$$

By (2.15) and (2.16), we find that for each $t \in [T, \infty)$,

$$\|\nabla w(\cdot, t)\|_{L^2(B_R(x_0))}^2 \leq \lambda^2 e^{-2C_2 R^{-2}(t-T)} \|\nabla w(\cdot, T)\|_{L^2(B_R(x_0))}^2.$$

This, together with (2.11) and $\lambda \geq 1$, means that

$$\|w(\cdot, t)\|_{H^1(B_R(x_0))}^2 \leq \lambda^2 e^{-C_2 R^{-2}(t-T)} \|w(\cdot, T)\|_{H^1(B_R(x_0))}^2.$$

By (2.1), we have that $w(\cdot, t) = v(\cdot, t)$ for each $t \geq T$. Thus, by (2.3) for the case $t \in [0, T]$, we conclude the desired result with $\hat{C}_1 = 2\lambda^2(3 + \lambda^2)(1 + 2C_1)$ and $\hat{C}_2 = C_2/2$. \square

Define

$$\tilde{v}(\cdot, t) := \begin{cases} v(\cdot, t) & \text{if } t \geq 0, \\ 0 & \text{if } t < 0, \end{cases}$$

where v is the solution of (2.2). By Lemma 2.1, we see the Fourier transform of \tilde{v} with respect to the time variable $t \in \mathbb{R}$ is meaningful

$$\hat{v}(x, \mu) = \int_{\mathbb{R}} e^{-i\mu t} \tilde{v}(x, t) dt \quad \text{for } (x, \mu) \in B_R(x_0) \times \mathbb{R}.$$

We further have

Lemma 2.2. *There exist positive constants $\hat{C}_3 = \hat{C}_3(\lambda, N)$ and $\hat{C}_4 = \hat{C}_4(\lambda, N)$ such that for each $\mu \in \mathbb{R}$, the following two estimates hold:*

$$\|\nabla \hat{v}(\cdot, \mu)\|_{L^2(B_r(x_0))}^2 \leq \frac{\hat{C}_3}{(R - 2r)^2} \|\hat{v}(\cdot, \mu)\|_{L^2(B_{\frac{R}{2}}(x_0))}^2 \quad \text{for all } 0 < r < \frac{R}{2} \quad (2.17)$$

and

$$\|\hat{v}(\cdot, \mu)\|_{L^2(B_{\frac{R}{2}}(x_0))} \leq \hat{C}_4 (T + R^2) e^{\frac{\hat{C}_1 T}{T - \tau} - \frac{\sqrt{|\mu|} R}{4e\hat{C}_5}} F(R), \quad (2.18)$$

where $\hat{C}_5 = \sqrt{8\pi^2 \sqrt{2 + \lambda^6}}$, $\hat{C}_1 > 0$ and $F(R)$ are given by Lemma 2.1.

Proof. By (2.2), we have that for each $\mu \in \mathbb{R}$,

$$i\mu\hat{v}(x, \mu) - \operatorname{div}(A(x)\nabla\hat{v}(x, \mu)) = 0 \text{ in } B_R(x_0). \quad (2.19)$$

Take arbitrarily $r \in (0, R/2)$ and define a cutoff function $\psi \in C^\infty(\mathbb{R}^N; [0, 1])$ verifying

$$\begin{cases} \psi = 1 & \text{in } \overline{B_r(x_0)}, \\ \psi = 0 & \text{in } \mathbb{R}^N \setminus B_{\frac{R}{2}}(x_0), \\ |\nabla\psi| \leq \frac{C_3}{R-2r} & \text{in } \mathbb{R}^N, \end{cases} \quad (2.20)$$

where $C_3 = C_3(N)$ is a positive constant. The rest proof is divided into two steps.

Step 1. The proof of (2.17).

Multiplying first (2.19) by $\bar{v}\psi^2$ and then integrating by parts over $B_{\frac{R}{2}}(x_0)$, we have

$$\int_{B_{\frac{R}{2}}(x_0)} \nabla\bar{v} \cdot (A\nabla\hat{v})\psi^2 dx + 2 \int_{B_{\frac{R}{2}}(x_0)} \nabla\psi \cdot (A\nabla\hat{v})\bar{v}\psi dx = -i \int_{B_{\frac{R}{2}}(x_0)} \mu|\hat{v}|^2\psi^2 dx.$$

By (1.2) and the Young inequality, we derive that for each $\epsilon_2 > 0$,

$$\begin{aligned} & \lambda^{-1} \int_{B_{\frac{R}{2}}(x_0)} |\nabla\hat{v}|^2\psi^2 dx = \lambda^{-1} \operatorname{Re} \int_{B_{\frac{R}{2}}(x_0)} |\nabla\hat{v}|^2\psi^2 dx \\ & \leq -2\operatorname{Re} \int_{B_{\frac{R}{2}}(x_0)} \nabla\psi \cdot (A\nabla\hat{v})\bar{v}\psi dx \leq 2\lambda \int_{B_{\frac{R}{2}}(x_0)} |\nabla\psi| |\nabla\hat{v}| |\bar{v}| |\psi| dx \\ & \leq \epsilon_2 \int_{B_{\frac{R}{2}}(x_0)} |\nabla\hat{v}|^2\psi^2 dx + \frac{\lambda^2}{\epsilon_2} \int_{B_{\frac{R}{2}}(x_0)} |\hat{v}|^2 |\nabla\psi|^2 dx. \end{aligned} \quad (2.21)$$

Taking $\epsilon_2 = 1/(2\lambda)$ in the above inequality, by (2.20) and (2.21), we derive (2.17).

Step 2. The proof of (2.18).

Note that when $\mu = 0$, by (2.6), (2.11) and the definitions of \hat{v}, \tilde{v}, w , we have

$$\begin{aligned} & \|\hat{v}(\cdot, 0)\|_{L^2(B_R(x_0))} \\ & = \left\| \int_{\mathbb{R}} \tilde{v}(x, t) dt \right\|_{L^2(B_R(x_0))} = \left\| \int_0^{+\infty} v(x, t) dt \right\|_{L^2(B_R(x_0))} \\ & \leq \int_0^T \|v(x, t)\|_{L^2(B_R(x_0))} dt + \int_T^{+\infty} \|v(x, t)\|_{L^2(B_R(x_0))} dt \\ & \leq 2Te^{\frac{C_1 T}{T-\tau}} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{L^2(B_R(x_0))} + \int_T^{+\infty} e^{-\frac{C_2(t-T)}{2R^2}} \|v(T)\|_{L^2(B_R(x_0))} dt \\ & \leq 2e^{\frac{C_1 T}{T-\tau}} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_R(x_0))} \left[T + \int_T^{+\infty} e^{-\frac{C_2(t-T)}{2R^2}} dt \right] \\ & \leq 2e^{\frac{C_1 T}{T-\tau}} \left[T + \frac{2R^2}{C_2} \right] \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_R(x_0))}. \end{aligned} \quad (2.22)$$

This yields that (2.18) is true when $\mu = 0$. Thus it suffices to prove (2.18) in the case that $\mu \neq 0$. To this end, define for each $\mu \in \mathbb{R} \setminus \{0\}$,

$$p(x, \xi, \mu) = e^{i\sqrt{|\mu|\xi}} \hat{v}(x, \mu) \text{ for a.e. } (x, \xi) \in B_R(x_0) \times \mathbb{R}.$$

Then, by (2.19), one can check easily that, for each fixed $\mu \in \mathbb{R} \setminus \{0\}$, $p(\cdot, \cdot, \mu)$ verifies

$$\operatorname{div}(A(x)\nabla p(x, \xi, \mu)) + i\operatorname{sign}(\mu)\partial_{\xi\xi}p(x, \xi, \mu) = 0 \text{ in } B_R(x_0) \times \mathbb{R}.$$

Here

$$\operatorname{sign}(\mu) = \begin{cases} 1 & \text{if } \mu > 0, \\ -1 & \text{if } \mu < 0. \end{cases}$$

As for the equation verified by $p(\cdot, \cdot, \mu)$, it is elliptic with complex coefficients and its coefficients are independent of the ξ -variable. These facts imply by standard energy methods (m times localized Cacciopoli's inequalities, see also [7]) that

$$\int_{B_{\frac{R}{2}}(x_0) \times (-\frac{R}{2}, \frac{R}{2})} |p^{(m)}|^2 dx d\xi \leq 2R \left[\frac{C_4 m^2}{R^2} \right]^m \int_{B_R(x_0)} |\hat{v}(x, \mu)|^2 dx \quad (2.23)$$

with $p^{(m)} = \partial_{\xi}^m p$ and $C_4 = 8\pi^2 \sqrt{2 + \lambda^6}$ (For the sake of completion, a detail proof for (2.23) is presented in Appendix). By the similar arguments in (2.22) and the definition of \hat{v} , we get that for each $\mu \in \mathbb{R}$,

$$\begin{aligned} \|\hat{v}(\cdot, \mu)\|_{L^2(B_R(x_0))} &\leq \int_0^{+\infty} \|v(x, t)\|_{L^2(B_R(x_0))} dt \\ &\leq 2e^{\frac{C_1 T}{T-\tau}} \left[T + \frac{2R^2}{C_2} \right] \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_R(x_0))}. \end{aligned} \quad (2.24)$$

Therefore, by (2.23), (2.24) and the definition of $F(R)$, we get that for each $m \in \mathbb{N}^+$,

$$\int_{B_{\frac{R}{2}}(x_0) \times (-\frac{R}{2}, \frac{R}{2})} |p^{(m)}|^2 dx d\xi \leq 8R \left[\frac{C_4 m^2}{R^2} \right]^m e^{\frac{2C_1 T}{T-\tau}} \left[T + \frac{2R^2}{C_2} \right]^2 F^2(R). \quad (2.25)$$

For any $\varphi \in L^2(B_{R/2}(x_0); \mathbb{C})$, we define

$$P_{\mu}(\xi) := \int_{B_{\frac{R}{2}}(x_0)} p(x, \xi, \mu) \bar{\varphi}(x) dx, \quad \xi \in \left(-\frac{R}{2}, \frac{R}{2} \right).$$

It is well known that the following interpolation inequality holds

$$\|f\|_{L^\infty(I)} \leq C_5 \left(|I| \|f'\|_{L^2(I)}^2 + \frac{1}{|I|} \|f\|_{L^2(I)}^2 \right)^{\frac{1}{2}} \text{ for each } f \in H^1(I), \quad (2.26)$$

where $C_5 > 0$, I is a bounded nonempty interval of \mathbb{R} and $|I|$ is the length. Therefore, by (2.25) and (2.26), we have that for any $\xi \in (-R/2, R/2)$ and $m \in \mathbb{N}^+$,

$$\begin{aligned} &|P_{\mu}^{(m)}(\xi)| \\ &\leq C_5 \left(R \int_{-\frac{R}{2}}^{\frac{R}{2}} |P_{\mu}^{(m+1)}(\xi)|^2 d\xi + \frac{1}{R} \int_{-\frac{R}{2}}^{\frac{R}{2}} |P_{\mu}^{(m)}(\xi)|^2 d\xi \right)^{\frac{1}{2}} \\ &= C_5 \left(R \int_{-\frac{R}{2}}^{\frac{R}{2}} \left| \int_{B_{\frac{R}{2}}(x_0)} p^{(m+1)}(x, \xi, \mu) \bar{\varphi}(x) dx \right|^2 d\xi + \frac{1}{R} \int_{-\frac{R}{2}}^{\frac{R}{2}} \left| \int_{B_{\frac{R}{2}}(x_0)} p^{(m)}(x, \xi, \mu) \bar{\varphi}(x) dx \right|^2 d\xi \right)^{\frac{1}{2}} \end{aligned}$$

$$\begin{aligned}
&\leq C_5 \left(R \int_{B_{\frac{R}{2}}(x_0) \times (-\frac{R}{2}, \frac{R}{2})} |p^{(m+1)}|^2 dx d\xi + \frac{1}{R} \int_{B_{\frac{R}{2}}(x_0) \times (-\frac{R}{2}, \frac{R}{2})} |p^{(m)}|^2 dx d\xi \right)^{\frac{1}{2}} \|\varphi\|_{L^2(B_{\frac{R}{2}}(x_0))} \\
&\leq 4C_5 e^{\frac{C_1 T}{T-\tau}} \left[T + \frac{2R^2}{C_2} \right] F(R) \frac{C_4^{\frac{m+1}{2}} (m+1)^{m+1}}{R^m} \|\varphi\|_{L^2(B_{\frac{R}{2}}(x_0))}. \tag{2.27}
\end{aligned}$$

This implies that $P_\mu(\cdot)$ can be analytically extended to the complex ball (still denoted by the same notation)

$$E_0 := \left\{ \xi \in \mathbb{C} : |\xi| < \frac{R}{2\sqrt{C_4 e}} \right\}.$$

Then,

$$|P_\mu(\xi)| \leq \sum_{m=0}^{\infty} \frac{|P_\mu^{(m)}(0)|}{m!} |\xi|^m,$$

when $\xi \in i\mathbb{R} \cap E_0$. Taking $\xi_0 = -iR/(4\sqrt{C_4 e})$, by (2.27), we get that

$$|P_\mu(\xi_0)| \leq 4C_5 \sqrt{C_4} e^{\frac{C_1 T}{T-\tau}} \left[T + \frac{2R^2}{C_2} \right] F(R) \sum_{m=0}^{\infty} \frac{(m+1)^{m+1}}{m!(4e)^m} \|\varphi\|_{L^2(B_{\frac{R}{2}}(x_0))}. \tag{2.28}$$

While, by the definitions of $P_\mu(\cdot)$ and ξ_0 ,

$$P_\mu(\xi_0) = e^{\frac{\sqrt{|\mu|R}}{4\sqrt{C_4 e}}} \int_{B_{\frac{R}{2}}(x_0)} \hat{v}(x, \mu) \bar{\varphi}(x) dx.$$

This, together with (2.28) and the arbitrariness of φ , means that,

$$\|\hat{v}(\cdot, \mu)\|_{L^2(B_{\frac{R}{2}}(x_0))} \leq 4C_5 \sqrt{C_4} \left[T + \frac{2R^2}{C_2} \right] e^{\frac{C_1 T}{T-\tau} - \frac{\sqrt{|\mu|R}}{4\sqrt{C_4 e}}} \sum_{m=0}^{\infty} \frac{(m+1)^{m+1}}{m!(4e)^m} F(R). \tag{2.29}$$

By (2.29) and (2.22), we derive (2.18) and complete the proof. \square

2.2 Stability estimate for elliptic equations

For our purpose and the convenience of the reader, we first quote from [24, Theorem 5.1] the following propagation estimate of smallness (for sets on hyperplanes) for gradients of solutions to elliptic equations in the way we understood.

Lemma 2.3. *Let $0 < r < r' < R < +\infty$ and $\rho > 0$. Suppose a positive measurable set $\omega \subset B_{r/2}(x_0)$ satisfy $|\omega|/|B_r(x_0)| > \rho$. If g is a solution to $\operatorname{div}(A(x)\nabla g) + g_{x_{N+1}x_{N+1}} = 0$ in $B_R(x_0, 0)$, with $A(\cdot)$ satisfying the condition (1.2), then there exists $\hat{C}_6 = \hat{C}_6(r, r', \rho, \lambda) > 0$ and $\alpha_1 = \alpha_1(r, r', \rho, \lambda) \in (0, 1)$ such that*

$$\sup_{B_r(x_0, 0)} |(\nabla g, \partial_{N+1}g)| \leq \hat{C}_6 \left(\sup_{\omega} |(\nabla g(\cdot, 0), \partial_{N+1}g(\cdot, 0))| \right)^{\alpha_1} \left(\sup_{B_{r'}(x_0, 0)} |(\nabla g, \partial_{N+1}g)| \right)^{1-\alpha_1}. \tag{2.30}$$

Remark 2.4. We refer the interesting reader to [24] for more general statements on propagation of smallness on possibly lower dimensional subsets for solutions to elliptic equations.

Remark 2.5. Regarding the dependence of ω , we here emphasize that the constants \hat{C}_6 and α_1 in (2.30) depend only on the r, r' and N -dimensional Lebesgue measure $|\omega|$, but not on the shape or position of ω . This uniform dependence is very important in our approach to prove the main result of this paper (see also [2, 12, 13]).

Let $0 < 2r < R < +\infty$ and $\rho > 0$. Suppose a positive measurable set $\omega \subset B_{r/2}(x_0)$ satisfy $|\omega|/|B_r(x_0)| > \rho$. Let $g \in H^1(B_R(x_0, 0))$ be a solution of

$$\begin{cases} \operatorname{div}(A(x)\nabla g) + g_{x_{N+1}x_{N+1}} = 0 & \text{in } B_R(x_0, 0), \\ g(x, 0) = 0 & \text{in } B_R(x_0), \\ g_{x_{N+1}}(x, 0) = z(x) & \text{in } B_R(x_0), \end{cases} \quad (2.31)$$

where $z \in L^2(B_R(x_0))$.

Lemma 2.6. *There exist constants $\hat{C}_7 = \hat{C}_7(N, \rho, r, \lambda) > 0$ and $\alpha_2 = \alpha_2(N, \rho, r, \lambda) \in (0, 1)$ such that*

$$\|g\|_{H^1(B_r(x_0, 0))} \leq \hat{C}_7 \|g\|_{H^1(B_{\frac{3r}{2}}(x_0, 0))}^{\alpha_2} \|z\|_{L^2(\omega)}^{1-\alpha_2}. \quad (2.32)$$

Proof. We quote from [23, Lemma 4.3], [21, Lemma A] and [25, Theorem 5.1] that the following stability estimate holds: there exist constants $C_6 = C_6(\lambda, N) > 0$ and $\gamma = \gamma(r, \lambda, N) \in (0, 1)$ such that

$$\|g\|_{H^1(B_r(x_0, 0))} \leq C_6 r^{-2} \|g\|_{H^1(B_{\frac{7r}{6}}(x_0, 0))}^{\gamma} \|z\|_{L^2(B_{\frac{7r}{6}}(x_0))}^{1-\gamma}. \quad (2.33)$$

We first assume $z \in C(\overline{B_R(x_0)})$. Applying Lemma 2.3 to gradients of solutions for (2.31), we have

$$\|(\nabla g, \partial_{N+1}g)\|_{L^\infty(B_{\frac{7r}{6}}(x_0, 0))} \leq C_7 \|(\nabla g(\cdot, 0), \partial_{N+1}g(\cdot, 0))\|_{L^\infty(\omega)}^\theta \|(\nabla g, \partial_{N+1}g)\|_{L^\infty(B_{\frac{4r}{3}}(x_0, 0))}^{1-\theta} \quad (2.34)$$

with $C_7 = C_7(\rho, r, \lambda) > 0$ and $\theta = \theta(\rho, r, \lambda) \in (0, 1)$.

Next, we observe from the second line in (2.31) that $\nabla g(\cdot, 0) = 0$ in $B_R(x_0)$. By (2.34) and the standard elliptic regularity (see, e.g., [16, Theorem 8.32])

$$\|(\nabla g, \partial_{N+1}g)\|_{L^\infty(B_{\frac{4r}{3}}(x_0, 0))} \leq C_8 \|g\|_{H^1(B_{\frac{3r}{2}}(x_0, 0))}$$

with $C_8 = C_8(r, \lambda, N) > 0$, we have

$$\|z\|_{L^\infty(B_{\frac{7r}{6}}(x_0))} \leq \|(\nabla g, \partial_{N+1}g)\|_{L^\infty(B_{\frac{7r}{6}}(x_0, 0))} \leq C_7 C_8^{1-\theta} \|z\|_{L^\infty(\omega)}^\theta \|g\|_{H^1(B_{\frac{3r}{2}}(x_0, 0))}^{1-\theta}.$$

This, combined with (2.33), leads to

$$\begin{aligned} \|g\|_{H^1(B_r(x_0, 0))} &\leq C_6 r^{-2} \|g\|_{H^1(B_{\frac{7r}{6}}(x_0, 0))}^{\gamma} \|z\|_{L^2(B_{\frac{7r}{6}}(x_0))}^{1-\gamma} \\ &\leq C_6 r^{-2} |B_{\frac{7r}{6}}|^{1-\gamma} \|g\|_{H^1(B_{\frac{7r}{6}}(x_0, 0))}^{\gamma} \|z\|_{L^\infty(B_{\frac{7r}{6}}(x_0))}^{1-\gamma} \\ &\leq C_6 r^{-2} |B_{\frac{7r}{6}}|^{1-\gamma} \|g\|_{H^1(B_{\frac{4}{3}r}(x_0, 0))}^{\gamma} \left[C_7 C_8^{1-\theta} \|z\|_{L^\infty(\omega)}^\theta \|g\|_{H^1(B_{\frac{3r}{2}}(x_0, 0))}^{1-\theta} \right]^{1-\gamma} \end{aligned}$$

$$\leq C_9 \|z\|_{L^\infty(\omega)}^\beta \|g\|_{H^1(B_{\frac{3r}{2}}(x_0,0))}^{1-\beta} \quad (2.35)$$

with $C_9 = C_9(r, \rho, N, \lambda) > 0$ and $\beta = \beta(r, \rho, N, \lambda) \in (0, 1)$.

Finally, we complete the proof by replacing the L^∞ -norm in (2.35) with L^2 -norm. To this end, we define a new Lebesgue measurable set

$$\tilde{\omega} := \left\{ x \in \omega : |z(x)| \leq \left(\frac{2}{|\omega|} \int_\omega |z|^2 dx \right)^{\frac{1}{2}} \right\}. \quad (2.36)$$

It is clear that

$$\int_\omega |z|^2 dx \geq \int_{\omega \setminus \tilde{\omega}} |z|^2 dx \geq \frac{2|\omega \setminus \tilde{\omega}|}{|\omega|} \int_\omega |z|^2 dx,$$

where $\omega \setminus \tilde{\omega} = \omega \cap \tilde{\omega}^c$. This implies that $|\omega \setminus \tilde{\omega}| \leq |\omega|/2$, and hence $\tilde{\omega} \geq |\omega|/2$. Applying (2.35) with ω replaced by $\tilde{\omega}$ leads to

$$\|g\|_{H^1(B_r(x_0,0))} \leq C_{10} \|g\|_{H^1(B_{\frac{3r}{2}}(x_0,0))}^{\beta_1} \|z\|_{L^\infty(\tilde{\omega})}^{1-\beta_1}$$

for some new constant $C_{10} = C_{10}(r, \rho, N, \lambda) > 0$ and $\beta_1 = \beta_1(r, \rho, N, \lambda) \in (0, 1)$. This, together with (2.36), indicates the desired estimate (2.32) for the case $z \in C(\overline{B_R(x_0)})$. This, together with (2.33) and the fact $C(\overline{B_R(x_0)})$ is dense in $L^2(B_R(x_0))$, yields that (2.32) holds for any $z \in L^2(B_R(x_0))$. This ends the proof. \square

3 Proof of Theorem 1.1

In order to give the proof of Theorem 1.1, we need the following the interpolation inequality (i.e., Lemma 3.1), whose proof below is based on a reduction method [18] (see also [22]). It is worth mentioning that the interpolation inequality established in Lemma 3.1 is related to the so-called two spheres and one cylinder inequality obtained in [7, Theorem 3.1.1'] (see also [11, Theorem 2]) which has some applications in the study of inverse problem (see, for instance, [7]). Comparing with them, however, there are two main differences: (i) the radius r of the observation sphere is here allowed to be independent of the observation time τ ; (ii) we specify the dependence of the observability constant on the observation time τ .

Lemma 3.1. *Let $T > 0$, $0 < r < +\infty$ and $\rho > 0$. Suppose a positive measurable set $\omega \subset B_{r/2}(x_0)$ satisfy $|\omega|/|B_r(x_0)| > \rho$. There exist constants $\hat{C}_8 = \hat{C}_8(r, N, \lambda, \rho) > 0$ and $\sigma = \sigma(r, N, \lambda, \rho) \in (0, 1)$ such that for each $\tau \in (0, T/2)$, the corresponding solution u of (1.1) with the initial value $u_0 \in H^1(\mathbb{R}^N)$ satisfies*

$$\|u(\cdot, \tau)\|_{L^2(B_r(x_0))} \leq \hat{C}_8 \left[T^2 e^{\frac{\hat{C}_1 T}{T-\tau}} + e^{\frac{\hat{C}_8}{\tau}} \right] \|u(\cdot, \tau)\|_{L^2(\omega)}^\sigma \left(\sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{gr}(x_0))}^2 \right)^{\frac{1-\sigma}{2}},$$

where $\varrho := 16\hat{C}_5 e$, the constant $\hat{C}_1 > 0$ is given by Lemma 2.1 and the constant $\hat{C}_5 = \sqrt{8\pi^2 \sqrt{2} + \lambda^6}$ is given by Lemma 2.2.

Proof. Taking $u_0 \in H^1(\mathbb{R}^N)$ arbitrarily. Let u be the solution to the equation (1.1) with the initial value $u_0 \in H^1(\mathbb{R}^N)$, and $R := \varrho r$. Let u_1 and u_2 be accordingly the solutions to

$$\begin{cases} \partial_t u_1 - \operatorname{div}(A(x)\nabla u_1) = 0 & \text{in } B_R(x_0) \times (0, 2T), \\ u_1 = u & \text{on } \partial B_R(x_0) \times (0, 2T), \\ u_1(\cdot, 0) = 0 & \text{in } B_R(x_0) \end{cases}$$

and

$$\begin{cases} \partial_t u_2 - \operatorname{div}(A(x)\nabla u_2) = 0 & \text{in } B_R(x_0) \times (0, 2T), \\ u_2 = 0 & \text{on } \partial B_R(x_0) \times (0, 2T), \\ u_2(\cdot, 0) = u_0 & \text{in } B_R(x_0). \end{cases}$$

It is clear that $u = u_1 + u_2$ in $B_R(x_0) \times [0, 2T]$. By the standard energy estimate for solutions of parabolic equations, we have

$$\sup_{t \in [0, T]} \|u_2(\cdot, t)\|_{H^1(B_R(x_0))} \leq C_{11} \|u_0\|_{H^1(B_R(x_0))} \quad (3.1)$$

with $C_{11} = C_{11}(N, \lambda) > 0$.

Arbitrarily fix $\tau \in (0, T/2)$. Let v_1 be the solution to

$$\begin{cases} \partial_t v_1 - \operatorname{div}(A(x)\nabla v_1) = 0 & \text{in } B_R(x_0) \times \mathbb{R}^+, \\ v_1 = \eta u & \text{on } \partial B_R(x_0) \times \mathbb{R}^+, \\ v_1(\cdot, 0) = 0 & \text{in } B_R(x_0), \end{cases}$$

where η is given by (2.1). It is clear that $u = v_1 + u_2$ in $B_R(x_0) \times [0, \tau]$. We extend v_1 to $\mathbb{R}^- \times B_R(x_0)$ by zero, and still denote it by the same way. Define

$$\hat{v}_1(x, \mu) = \int_{\mathbb{R}} e^{-i\mu t} v_1(x, t) dt \quad \text{for } (x, \mu) \in B_R(x_0) \times \mathbb{R}.$$

Note from Lemma 2.1 that \hat{v}_1 is well defined.

Let

$$\kappa := \frac{\sqrt{2}}{4e\hat{C}_5} \quad \text{with } \hat{C}_5 = \sqrt{8\pi^2\sqrt{2} + \lambda^6} \quad \text{given in Lemma 2.2.}$$

We define

$$V = V_1 + V_2 \quad \text{in } B_R(x_0) \times (-\kappa R, \kappa R)$$

with

$$V_1(x, y) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\tau\mu} \hat{v}_1(x, \mu) \frac{\sinh(\sqrt{-i\mu}y)}{\sqrt{-i\mu}} d\mu \quad \text{in } B_R(x_0) \times (-\kappa R, \kappa R), \quad (3.2)$$

$$V_2(x, y) = \sum_{k=1}^{\infty} \alpha_k e^{-\mu_k \tau} f_k(x) \frac{\sinh(\sqrt{\mu_k}y)}{\sqrt{\mu_k}} \quad \text{in } B_R(x_0) \times (-\kappa R, \kappa R) \quad (3.3)$$

where $\{\mu_k\}_{k=1}^{\infty}$, $\{f_k\}_{k=1}^{\infty}$ are given by (2.10), and $\alpha_k = \langle u_2(\cdot, 0), f_k \rangle_{L^2(B_R(x_0))}$. Note from Lemma 2.2 that V_1 is also well defined. One can readily check that

$$\begin{cases} \operatorname{div}(A(x)\nabla V(x, y)) + V_{yy}(x, y) = 0 & \text{in } B_{\frac{R}{2}}(x_0) \times (-\kappa R, \kappa R), \\ V(x, 0) = 0 & \text{in } B_{\frac{R}{2}}(x_0), \\ V_y(x, 0) = u(x, \tau) & \text{in } B_{\frac{R}{2}}(x_0). \end{cases} \quad (3.4)$$

Let $r \in (0, \kappa R/2)$. By the three-ball inequality for elliptic operators (cf., e.g., [26, Theorem 3.1]), there exist constants $C_{13} = C_{13}(\lambda, N, r) > 0$ and $\beta_2 = \beta_2(\lambda, N, r) \in (0, 1)$ such that

$$\|V\|_{L^2(B_{\frac{r}{4}}(x_0, 0))} \leq C_{13} \|V\|_{L^2(B_r(x_0, 0))}^{\beta_2} \|V\|_{L^2(B_{2r}(x_0, 0))}^{1-\beta_2}. \quad (3.5)$$

Since V_y satisfies the first equation of (3.4), by the interior estimate of elliptic equations we find

$$\begin{aligned}
\int_{B_{\frac{7r}{4}}(x_0,0)} |V|^2 dx dy &\geq C_{14} r^2 \int_{B_{\frac{3r}{2}}(x_0,0)} (|\nabla V|^2 + |V_y|^2) dx dy \\
&\geq \frac{C_{14} r^2}{2} \left(\int_{B_{\frac{3r}{2}}(x_0,0)} |V_y|^2 dx dy + \int_{B_{\frac{3r}{2}}(x_0,0)} |V_y|^2 dx dy \right) \\
&\geq C_{14} r^2 \left(\int_{B_{\frac{3r}{2}}(x_0,0)} |V_y|^2 dx dy + r^2 \int_{B_{\frac{5r}{4}}(x_0,0)} (|\nabla V_y|^2 + |V_{yy}|^2) dx dy \right) \\
&\geq C_{14} r^3 \left(\frac{1}{r} \int_{B_{\frac{5r}{4}}(x_0,0)} |V_y|^2 dx dy + r \int_{B_{\frac{5r}{4}}(x_0,0)} |V_{yy}|^2 dx dy \right) \tag{3.6}
\end{aligned}$$

with $C_{14} = C_{14}(\lambda, N) > 0$.

As a simple corollary of [4, Lemma 9.9, Page 315], we have the following trace theorem

$$\int_{B_r(x_0)} |f(x, 0)|^2 dx \leq C_{15}(N) \left(\frac{1}{r} \int_{B_{\frac{5r}{4}}(x_0,0)} |f|^2 dx dy + r \int_{B_{\frac{5r}{4}}(x_0,0)} |f_y|^2 dx dy \right)$$

for any $f \in H^1(B_{5r/4}(x_0, 0))$. Hence, by (3.4) and (3.6) we have

$$C_{16} r^3 \int_{B_r(x_0)} |u(x, \tau)|^2 dx \leq \int_{B_{\frac{7r}{4}}(x_0,0)} |V|^2 dx dy \tag{3.7}$$

with $C_{16} = C_{16}(\lambda, N) > 0$.

By Lemma 2.6, we obtain that there is $C_{17} = C_{17}(\rho, N, r, \lambda) > 0$ and $\beta_3 = \beta_3(\rho, N, r, \lambda) \in (0, 1)$ such that

$$\|V\|_{L^2(B_r(x_0,0))} \leq C_{17} \|V\|_{H^1(B_{\frac{3r}{2}}(x_0,0))}^{\beta_3} \|u(\cdot, \tau)\|_{L^2(\omega)}^{1-\beta_3}. \tag{3.8}$$

Again, by the interior estimate, there is a constant $C_{18} = C_{18}(N) > 0$ such that

$$\|V\|_{H^1(B_{\frac{3r}{2}}(x_0,0))} \leq C_{18} r^{-1} \|V\|_{L^2(B_{2r}(x_0,0))}. \tag{3.9}$$

Hence, it follows from (3.8) and (3.9) that

$$\|V\|_{L^2(B_r(x_0,0))} \leq C_{19} \|V\|_{L^2(B_{2r}(x_0,0))}^{\beta_3} \|u(\cdot, \tau)\|_{L^2(\omega)}^{1-\beta_3} \tag{3.10}$$

with $C_{19} = C_{19}(\rho, N, r, \lambda) > 0$. It follows from (3.7), (3.5) and (3.10) that

$$\|u(\cdot, \tau)\|_{L^2(B_r(x_0))} \leq C_{20} \|u(\cdot, \tau)\|_{L^2(\omega)}^{(1-\beta_3)\beta_2} \|V\|_{L^2(B_{2r}(x_0,0))}^{1-(1-\beta_3)\beta_2} \tag{3.11}$$

with $C_{20} = C_{20}(\rho, N, r, \lambda) > 0$.

To finish the proof, it suffices to bound the term $\|V\|_{L^2(B_{2r}(x_0,0))}$. Recall that $V = V_1 + V_2$, we will treat V_1 and V_2 separately.

In fact, we derive from (3.2) that for each $x \in B_{2r}(x_0) \subset B_R(x_0)$ and $|y| < \kappa R/(4\sqrt{2})$,

$$\begin{aligned}
|V_1(x, y)| &= \left| \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\tau\mu} \hat{v}_1(x, \mu) \int_{-y}^y e^{\sqrt{-i}\mu s} ds d\mu \right| \leq \frac{1}{2\pi} \int_{\mathbb{R}} |\hat{v}_1(x, \mu)| \int_{-y}^y |e^{\sqrt{-i}\mu s}| ds d\mu \\
&\leq \frac{\kappa R}{4\sqrt{2}\pi} \int_{\mathbb{R}} |\hat{v}_1(x, \mu)| e^{\frac{1}{4\sqrt{2}}\kappa\sqrt{|\mu|}R} d\mu \\
&\leq \frac{\kappa R}{4\sqrt{2}\pi} \left(\int_{\mathbb{R}} |\hat{v}_1(x, \mu)|^2 e^{\frac{1}{2}\kappa\sqrt{|\mu|}R} d\mu \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}} e^{-\frac{1}{2\sqrt{2}}\kappa\sqrt{|\mu|}R} d\mu \right)^{\frac{1}{2}} \\
&= \frac{1}{2\pi} \left(\int_{\mathbb{R}} |\hat{v}_1(x, \mu)|^2 e^{\frac{1}{2}\kappa\sqrt{|\mu|}R} d\mu \right)^{\frac{1}{2}}.
\end{aligned}$$

Hence, by Lemma 2.2, we have for each $r < \kappa R/(4\sqrt{2})$,

$$\begin{aligned}
\int_{B_{2r}(x_0, 0)} |V_1|^2 dx dy &\leq \frac{1}{2\pi^2} \hat{C}_4^2 R e^{\frac{2\hat{C}_1 T}{T-\tau}} [T + R^2]^2 F^2(R) \int_{\mathbb{R}} e^{-\frac{1}{2\sqrt{2}}\kappa\sqrt{|\mu|}R} d\mu \\
&\leq \frac{32e}{\pi^2} \hat{C}_5^2 \hat{C}_4^2 R^{-1} e^{\frac{2\hat{C}_1 T}{T-\tau}} [T + R^2]^2 F^2(R).
\end{aligned} \tag{3.12}$$

While, by (3.3) and (3.1) we obtain

$$\begin{aligned}
\int_{B_{2r}(x_0, 0)} |V_2|^2 dx dy &\leq \int_{-2r}^{2r} \int_{B_R(x_0)} |V_2|^2 dx dy \leq \int_{-2r}^{2r} \sum_{k=1}^{\infty} \alpha_k^2 e^{-2\mu_k \tau} \left| \frac{\sinh(\sqrt{\mu_k} y)}{\sqrt{\mu_k}} \right|^2 dy \\
&\leq \frac{e^{\frac{2r^2}{\tau}}}{4\hat{C}_2^{3/2}} R^3 \sum_{k=1}^{\infty} \alpha_k^2 = \frac{e^{\frac{2r^2}{\tau}}}{4\hat{C}_2^{3/2}} R^3 \int_{B_R(x_0)} |u(x, 0)|^2 dx \\
&\leq C_{21} R^3 e^{\frac{C_{21}}{\tau}} F^2(R)
\end{aligned} \tag{3.13}$$

with $C_{21} = C_{21}(N, r, \lambda) > 0$. Therefore, by (3.12) and (3.13), we conclude that

$$\|V\|_{L^2(B_{2r}(x_0, 0))} \leq C_{22} \left[e^{\frac{\hat{C}_1 T}{T-\tau}} R^{-1} (T + R^2)^2 + e^{\frac{C_{21}}{\tau}} R^3 \right]^{\frac{1}{2}} F(R)$$

with $C_{22} = C_{22}(N, r, \lambda) > 0$. This, together with (3.11), means that

$$\begin{aligned}
\|u(\cdot, \tau)\|_{L^2(B_r(x_0))} &\leq C_{20} \|u(\cdot, \tau)\|_{L^2(\omega)}^{(1-\beta_3)\beta_2} \|V\|_{L^2(B_{2r}(x_0, 0))}^{1-(1-\beta_3)\beta_2} \\
&\leq C_{20} C_{22}^{1-(1-\beta_3)\beta_2} \left[e^{\frac{\hat{C}_1 T}{T-\tau}} R^{-1} (T + R^2)^2 + e^{\frac{C_{21}}{\tau}} R^3 \right]^{\frac{1-(1-\beta_3)\beta_2}{2}} \\
&\quad \times \|u(\cdot, \tau)\|_{L^2(\omega)}^{(1-\beta_3)\beta_2} F(R)^{1-(1-\beta_3)\beta_2}.
\end{aligned}$$

Taking $\sigma = (1 - \beta_3)\beta_2$, the proof is immediately achieved by the arbitrariness of u_0 . \square

Lemma 3.2. *Let $T > 0$, $0 < r < +\infty$ and $\rho > 0$. Assume that there is a sequence $\{x_i\}_{i \in \mathbb{N}^+} \subset \mathbb{R}^N$ so that*

$$\mathbb{R}^N = \bigcup_{i \in \mathbb{N}^+} Q_r(x_i) \quad \text{with} \quad \text{int}(Q_r(x_i)) \cap \text{int}(Q_r(x_j)) = \emptyset \quad \text{for each} \quad i \neq j \in \mathbb{N}^+.$$

Let

$$\tilde{\omega} \triangleq \bigcup_{i \in \mathbb{N}^+} \tilde{\omega}_i \quad \text{with} \quad \tilde{\omega}_i \subset B_{r/2}(x_i) \quad \text{satisfy} \quad \frac{|\tilde{\omega}_i|}{|B_r(x_i)|} \geq \rho \quad \text{for each} \quad i \in \mathbb{N}^+$$

where, for each $i \in \mathbb{N}^+$, $\tilde{\omega}_i$ is a N -dimensional Lebesgue measurable set of positive measure. There exist constants $C = C(r, N, \lambda, \rho) > 0$ and $\sigma = \sigma(r, N, \lambda, \rho) \in (0, 1)$ such that for any $u_0 \in H^1(\mathbb{R}^N)$, the corresponding solution u of (1.1) satisfies

$$\|u(\cdot, T)\|_{L^2(\mathbb{R}^N)} \leq C \left(T^3 + e^{\frac{C}{T}} \right) \|u(\cdot, T)\|_{L^2(\tilde{\omega})}^\sigma \left(\sup_{s \in [0, 2T]} \|u(\cdot, s)\|_{H^1(\mathbb{R}^N)} \right)^{1-\sigma}.$$

Proof. By Lemma 3.1 (where r and ω are replaced by $\sqrt{N}r$ and $\tilde{\omega}_i$ ($i \in \mathbb{N}^+$), respectively), we get that

$$\begin{aligned} \int_{Q_r(x_i)} |u(x, \tau)|^2 dx &\leq \int_{B_{\sqrt{N}r}(x_i)} |u(x, \tau)|^2 dx \\ &\leq \hat{C}_9 \left(T^2 e^{\frac{\hat{C}_1 T}{T-\tau}} + e^{\frac{\hat{C}_9}{\tau}} \right)^2 \left(\int_{\tilde{\omega}_i} |u(x, \tau)|^2 dx \right)^\theta \\ &\quad \times \left(\sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{\sqrt{N}r}(x_i))}^2 \right)^{1-\theta}, \end{aligned}$$

where $\hat{C}_9 = \hat{C}_9(r, N, \lambda, \rho) > 0$ and $\theta = \theta(r, N, \lambda, \rho) \in (0, 1)$. This, along with Young's inequality, implies that for each $\varepsilon > 0$,

$$\begin{aligned} \int_{Q_r(x_i)} |u(x, \tau)|^2 dx &\leq \hat{C}_9^2 \left(T^2 e^{\frac{\hat{C}_1 T}{T-\tau}} + e^{\frac{\hat{C}_9}{\tau}} \right)^2 \left(\varepsilon \theta \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{\sqrt{N}r}(x_i))}^2 \right. \\ &\quad \left. + \varepsilon^{-\frac{\theta}{1-\theta}} (1-\theta) \int_{\tilde{\omega}_i} |u(x, \tau)|^2 dx \right). \end{aligned}$$

Then

$$\begin{aligned} \int_{\mathbb{R}^N} |u(x, \tau)|^2 dx &= \sum_{i \in \mathbb{N}^+} \int_{Q_r(x_i)} |u(x, \tau)|^2 dx \\ &\leq \hat{C}_9^2 \left(T^2 e^{\frac{\hat{C}_1 T}{T-\tau}} + e^{\frac{\hat{C}_9}{\tau}} \right)^2 \left(\varepsilon \theta \sum_{i \in \mathbb{N}^+} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{\sqrt{N}r}(x_i))}^2 + \varepsilon^{-\frac{\theta}{1-\theta}} (1-\theta) \int_{\tilde{\omega}} |u(x, \tau)|^2 dx \right). \end{aligned} \tag{3.14}$$

Next, let $R := \varrho r$, we show that

$$\sum_{i \in \mathbb{N}^+} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{\sqrt{N}R}(x_i))}^2 \leq \hat{C}_{10}(1+T) \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(\mathbb{R}^N)}^2, \tag{3.15}$$

where $\hat{C}_{10}(\lambda, R, N) > 0$. Indeed, we take a cut-off function $\psi \in C_0^\infty(\mathbb{R}^N; [0, 1])$ so that $\text{supp} \psi \Subset Q_{2\sqrt{N}R}(0)$ and $\psi \equiv 1$ in $B_{\sqrt{N}R}(0)$. For each $i \in \mathbb{N}^+$, we let $\psi_i(x) := \psi(x - x_i)$ for any $x \in \mathbb{R}^+$. It is clear that $\text{supp} \psi_i \Subset Q_{2\sqrt{N}R}(x_i)$ and $\psi_i \equiv 1$ in $B_{\sqrt{N}R}(x_i)$. We suppose that

$$\begin{cases} \text{Card}\{j \in \mathbb{N}^+ : Q_r(x_1) \cap Q_{2\sqrt{N}R}(x_j) \neq \emptyset\} = m^*, \\ \text{Card}\{j \in \mathbb{N}^+ : Q_{2\sqrt{N}R}(x_1) \cap Q_r(x_j) \neq \emptyset\} = n^*. \end{cases}$$

This means that, for each $k \in \mathbb{N}^+$,

$$\begin{cases} \text{Card}\{j \in \mathbb{N}^+ : Q_r(x_k) \cap Q_{2\sqrt{N}R}(x_j) \neq \emptyset\} = m^*, \\ \text{Card}\{j \in \mathbb{N}^+ : Q_{2\sqrt{N}R}(x_k) \cap Q_r(x_j) \neq \emptyset\} = n^*. \end{cases}$$

For each k , we suppose $\{y_{k,i}\}_{i=1}^{n^*} = \{x_i | Q_{2\sqrt{N}R}(x_k) \cap Q_r(x_i) \neq \emptyset\}$. Thus, for any $f \in H^1(\mathbb{R}^N)$,

$$\begin{aligned} \sum_{k \in \mathbb{N}^+} \|f\|_{H^1(Q_{2\sqrt{N}R}(x_k))}^2 &\leq \sum_{k \in \mathbb{N}^+} \sum_{i=1}^{n^*} \|f\|_{H^1(Q_r(y_{k,i}))}^2 \leq \sum_{i=1}^{n^*} \sum_{k \in \mathbb{N}^+} \|f\|_{H^1(Q_r(y_{k,i}))}^2 \\ &\leq m^* n^* \sum_{k \in \mathbb{N}^+} \|f\|_{H^1(Q_r(x_k))}^2 = m^* n^* \|f\|_{H^1(\mathbb{R}^N)}^2. \end{aligned} \quad (3.16)$$

For each $i \in \mathbb{N}^+$, we let $v_i = \psi_i u$. It is obvious that v_i verifies

$$\begin{cases} (v_i)_t - \text{div}(A(x)\nabla v_i) = -2\nabla\psi_i \cdot (A(x)\nabla u) - u \text{div}(A(x)\nabla\psi_i) & \text{in } Q_{2\sqrt{N}R}(x_i) \times [0, T], \\ v_i = 0 & \text{on } \partial Q_{2\sqrt{N}R}(x_i) \times [0, T], \\ v_i(0) = \psi_i u_0 & \text{in } Q_{2\sqrt{N}R}(x_i). \end{cases} \quad (3.17)$$

By the standard energy estimate for (3.17), one can easily check that

$$\begin{aligned} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{\sqrt{N}R}(x_i))}^2 &\leq \sup_{s \in [0, T]} \|v_i(\cdot, s)\|_{H^1(Q_{2\sqrt{N}R}(x_i))}^2 \\ &\leq C(\lambda, N)(1 + R^{-4}) \left(\|u_0\|_{H^1(Q_{2\sqrt{N}R}(x_i))}^2 + \int_0^T \|u(\cdot, s)\|_{H^1(Q_{2\sqrt{N}R}(x_i))}^2 ds \right), \end{aligned}$$

where $C(\lambda, N) > 0$. This, along with (3.16), yields that

$$\begin{aligned} \sum_{i \in \mathbb{N}^+} \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(B_{\sqrt{N}R}(x_i))}^2 &\leq C(\lambda, N)(1 + R^{-4}) \sum_{i \in \mathbb{N}^+} \left(\|u_0\|_{H^1(Q_{2\sqrt{N}R}(x_i))}^2 + \int_0^T \|u(\cdot, s)\|_{H^1(Q_{2\sqrt{N}R}(x_i))}^2 ds \right) \\ &\leq m^* n^* C(\lambda, N)(1 + R^{-4})(1 + T) \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(\mathbb{R}^N)}^2. \end{aligned}$$

Thus, (3.15) is true. By (3.15) and (3.14), we obtain that

$$\begin{aligned} \int_{\mathbb{R}^N} |u(x, \tau)|^2 dx &\leq \hat{C}_9^2 \left(T^2 e^{\frac{\hat{C}_1 T}{T-\tau}} + e^{\frac{\hat{C}_9}{\tau}} \right)^2 \left(\varepsilon \theta \hat{C}_{10} (1 + T) \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(\mathbb{R}^N)}^2 \right. \\ &\quad \left. + \varepsilon^{-\frac{\theta}{1-\theta}} (1 - \theta) \int_{\tilde{\omega}} |u(x, \tau)|^2 dx \right) \end{aligned}$$

for each $\varepsilon > 0$. This implies that

$$\begin{aligned} \int_{\mathbb{R}^N} |u(x, \tau)|^2 dx &\leq \hat{C}_9^2 \left(T^2 e^{\frac{\hat{C}_1 T}{T-\tau}} + e^{\frac{\hat{C}_9}{\tau}} \right)^2 \left(\hat{C}_{10} (1 + T) \sup_{s \in [0, T]} \|u(\cdot, s)\|_{H^1(\mathbb{R}^N)}^2 \right)^\theta \\ &\quad \times \left(\int_{\tilde{\omega}} |u(x, \tau)|^2 dx \right)^{1-\theta}. \end{aligned}$$

By Lemma 3.1 (where τ and T are replaced by T and $2T$, respectively), we finish the proof of Lemma 3.2. \square

Proof of Theorem 1.1. First, by standard energy estimates of solutions to (1.1) we have

$$\|u(\cdot, t)\|_{H^1(\mathbb{R}^N)} \leq \frac{C_{23}e^{C_{23}t}}{\sqrt{t}} \|u_0\|_{L^2(\mathbb{R}^N)} \quad (3.18)$$

with $C_{23} = C_{23}(N, \lambda) > 0$, for each $t \in (0, 6T]$. Moreover, if $u_0 \in H^1(\mathbb{R}^N)$, then

$$\|u(\cdot, t)\|_{H^1(\mathbb{R}^N)} \leq C_{24}e^{C_{24}t} \|u_0\|_{H^1(\mathbb{R}^N)} \quad \text{for each } t \in [0, 6T], \quad (3.19)$$

where $C_{24} = C_{24}(N, \lambda) > 0$.

Second, we consider the following equation

$$\begin{cases} v_t - \operatorname{div}(A(x)\nabla v) = 0 & \text{in } \mathbb{R}^N \times (0, 4T), \\ v(\cdot, 0) = u(\cdot, \frac{T}{2}) & \text{in } \mathbb{R}^N. \end{cases}$$

It is obvious that $v(\cdot, t) = u(\cdot, t+T/2)$ when $t \in [0, 4T]$. Moreover, by (3.18) we have $u(\cdot, T/2) \in H^1(\mathbb{R}^N)$, which means that $v \in C([0, 4T]; H^1(\mathbb{R}^N))$. From Lemma 3.2 (where T and $\tilde{\omega}$ are replaced by $T/2$ and ω , respectively), it follows that there are $C > 0$ and $\sigma \in (0, 1)$ such that

$$\left\| v\left(\cdot, \frac{T}{2}\right) \right\|_{L^2(\mathbb{R}^N)} \leq C \left(T^3 + e^{\frac{C}{T}}\right) \left\| v\left(\cdot, \frac{T}{2}\right) \right\|_{L^2(\omega)}^\sigma \left(\sup_{s \in [0, T]} \|v(\cdot, s)\|_{H^1(\mathbb{R}^N)} \right)^{1-\sigma}.$$

This, along with (3.19), gives that

$$\left\| v\left(\cdot, \frac{T}{2}\right) \right\|_{L^2(\mathbb{R}^N)} \leq CC_{24}e^{C_{24}T} \left(T^3 + e^{\frac{C}{T}}\right) \left\| v\left(\cdot, \frac{T}{2}\right) \right\|_{L^2(\omega)}^\sigma \|v(\cdot, 0)\|_{H^1(\mathbb{R}^N)}^{1-\sigma}.$$

Which is

$$\|u(\cdot, T)\|_{L^2(\mathbb{R}^N)} \leq CC_{24}e^{C_{24}T} \left(T^3 + e^{\frac{C}{T}}\right) \|u(\cdot, T)\|_{L^2(\omega)}^\sigma \left\| u\left(\cdot, \frac{T}{2}\right) \right\|_{H^1(\mathbb{R}^N)}^{1-\sigma}.$$

This, together with (3.18), completes the proof. \square

4 Proof of Theorem 1.2

Now, we are able to present the proof of Theorem 1.2.

Proof of Theorem 1.2. By Theorem 1.1 (where r, x_i and w_i are replaced by r, x_i and w_i , respectively) and Young's inequality, for any $0 \leq t_1 < t_2 \leq T$, we see that

$$\|u(t_2)\|_{L^2(\mathbb{R}^N)}^2 \leq \varepsilon \|u(t_1)\|_{L^2(\mathbb{R}^N)}^2 + \frac{C_{25}}{\varepsilon^\alpha} e^{\frac{C_{26}}{t_2 - t_1}} \|u(t_2)\|_{L^2(\omega)}^2 \quad \text{for each } \varepsilon > 0, \quad (4.1)$$

where $C_{25} := (Ce^{CT})^{2/(1-\sigma)}$, $C_{26} := 2C/(1-\sigma)$ and $\alpha := \sigma/(1-\sigma)$. Let l be a density point of E . According to Proposition 2.1 in [32], for each $\kappa > 1$, there exists $l_1 \in (l, T)$, depending on κ and E , so that the sequence $\{l_m\}_{m \in \mathbb{N}^+}$, given by

$$l_{m+1} = l + \frac{1}{\kappa^m} (l_1 - l) \quad \text{for each } m \in \mathbb{N}^+,$$

satisfies that

$$l_m - l_{m+1} \leq 3|E \cap (l_{m+1}, l_m)|. \quad (4.2)$$

Next, let $0 < l_{m+2} < l_{m+1} \leq t < l_m < l_1 < T$. It follows from (4.1) that

$$\|u(t)\|_{L^2(\mathbb{R}^N)}^2 \leq \varepsilon \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 + \frac{C_{25}}{\varepsilon^\alpha} e^{\frac{C_{26}}{t-l_{m+2}}} \|u(t)\|_{L^2(\omega)}^2 \text{ for each } \varepsilon > 0. \quad (4.3)$$

By a standard energy estimate, we have that

$$\|u(l_m)\|_{L^2(\mathbb{R}^N)} \leq C_{27} \|u(t)\|_{L^2(\mathbb{R}^N)},$$

where $C_{27} = C_{27}(\lambda, N) \geq 1$. This, along with (4.3), implies that

$$\|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 \leq C_{27}^2 \left(\varepsilon \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 + \frac{C_{25}}{\varepsilon^\alpha} e^{\frac{C_{26}}{t-l_{m+2}}} \|u(t)\|_{L^2(\omega)}^2 \right) \text{ for each } \varepsilon > 0,$$

which indicates that

$$\|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 \leq \varepsilon \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 + \frac{C_{28}}{\varepsilon^\alpha} e^{\frac{C_{26}}{t-l_{m+2}}} \|u(t)\|_{L^2(\omega)}^2 \text{ for each } \varepsilon > 0,$$

where $C_{28} = C_{27}^{2(1+\alpha)} C_{25}$. Integrating the latter inequality over $E \cap (l_{m+1}, l_m)$, we get that

$$\begin{aligned} |E \cap (l_{m+1}, l_m)| \|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 &\leq \varepsilon |E \cap (l_{m+1}, l_m)| \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 \\ &\quad + \frac{C_{28}}{\varepsilon^\alpha} e^{\frac{C_{26}}{l_{m+1}-l_{m+2}}} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt \text{ for each } \varepsilon > 0. \end{aligned} \quad (4.4)$$

Here and throughout the proof of Theorem 1.2, χ_E denotes the characteristic function of E . Since $l_m - l_{m+1} = (\kappa - 1)(l_1 - l)/\kappa^m$, by (4.4) and (4.2), we obtain that

$$\begin{aligned} \|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 &\leq \varepsilon \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 + \frac{1}{|E \cap (l_{m+1}, l_m)|} \frac{C_{28}}{\varepsilon^\alpha} e^{\frac{C_{26}}{l_{m+1}-l_{m+2}}} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt \\ &\leq \frac{3\kappa^m}{(l_1 - l)(\kappa - 1)} \frac{C_{28}}{\varepsilon^\alpha} e^{2C_{26} \left(\frac{1}{l_1 - l} \frac{\kappa^{m+1}}{\kappa - 1} \right)} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt + \varepsilon \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 \end{aligned}$$

for each $\varepsilon > 0$. This yields that

$$\|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 \leq \frac{1}{\varepsilon^\alpha} \frac{3}{\kappa} \frac{C_{28}}{C_{26}} e^{2C_{26} \left(\frac{1}{l_1 - l} \frac{\kappa^{m+1}}{\kappa - 1} \right)} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt + \varepsilon \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 \quad (4.5)$$

for each $\varepsilon > 0$. Denote $d := 2C_{26}/[\kappa(l_1 - l)(\kappa - 1)]$. It follows from (4.5) that

$$\varepsilon^\alpha e^{-d\kappa^{m+2}} \|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 - \varepsilon^{1+\alpha} e^{-d\kappa^{m+2}} \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 \leq \frac{3}{\kappa} \frac{C_{28}}{C_{26}} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt$$

for each $\varepsilon > 0$. Choosing $\varepsilon = e^{-d\kappa^{m+2}}$ in the latter inequality, we observe that

$$e^{-(1+\alpha)d\kappa^{m+2}} \|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 - e^{-(2+\alpha)d\kappa^{m+2}} \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 \leq \frac{3}{\kappa} \frac{C_{28}}{C_{26}} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt. \quad (4.6)$$

Take $\kappa = \sqrt{(\alpha + 2)/(\alpha + 1)}$ in (4.6). Then we have that

$$e^{-(2+\alpha)d\kappa^m} \|u(l_m)\|_{L^2(\mathbb{R}^N)}^2 - e^{-(2+\alpha)d\kappa^{m+2}} \|u(l_{m+2})\|_{L^2(\mathbb{R}^N)}^2 \leq \frac{3}{\kappa} \frac{C_{28}}{C_{26}} \int_{l_{m+1}}^{l_m} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt.$$

Changing m to $2m'$ and summing the above inequality from $m' = 1$ to infinity give the desired result. Indeed,

$$\begin{aligned} & C_{27}^{-2} e^{-(2+\alpha)d\kappa^2} \|u(T)\|_{L^2(\mathbb{R}^N)}^2 \leq e^{-(2+\alpha)d\kappa^2} \|u(l_2)\|_{L^2(\mathbb{R}^N)}^2 \\ & \leq \sum_{m'=1}^{+\infty} \left(e^{-(2+\alpha)d\kappa^{2m'}} \|u(l_{2m'})\|_{L^2(\mathbb{R}^N)}^2 - e^{-(2+\alpha)d\kappa^{2m'+2}} \|u(l_{2m'+2})\|_{L^2(\mathbb{R}^N)}^2 \right) \\ & \leq \frac{3}{\kappa} \frac{C_{28}}{C_{26}} \sum_{m'=1}^{+\infty} \int_{l_{2m'+1}}^{l_{2m'}} \chi_E \|u(t)\|_{L^2(\omega)}^2 dt \leq \frac{3}{\kappa} \frac{C_{28}}{C_{26}} \int_0^T \chi_E \|u(t)\|_{L^2(\omega)}^2 dt. \end{aligned}$$

In summary, we finish the proof of Theorem 1.2. \square

5 Appendix: Proof of (2.23)

Let $m \in \mathbb{N}^+$ and $a_j = 1 - j/(2m)$ for $j = 0, 1, \dots, m+1$. For each $j \in \{0, 1, \dots, m\}$, we define a cutoff function

$$h_j(s) := \begin{cases} 0 & \text{if } |s| > a_j, \\ \frac{1}{2} \left[1 + \cos \left(\frac{\pi(a_{j+1}-|s|)}{a_{j+1}-a_j} \right) \right] & \text{if } a_{j+1} \leq |s| \leq a_j, \\ 1 & \text{if } |s| < a_{j+1}. \end{cases}$$

Clearly,

$$|h'_j(s)| \leq m\pi \text{ for any } s \in \mathbb{R}.$$

Denote $p_j = \partial^j p / \partial \xi^j$, $j = 0, 1, \dots, m$. Then p_j verifies

$$\operatorname{div}(A \nabla p_j(\cdot, \cdot, \mu)) + i \operatorname{sign}(\mu) p_{j+2}(\cdot, \cdot, \mu) = 0 \text{ in } B_R(x_0) \times \mathbb{R}. \quad (5.1)$$

Let

$$\eta_j(x, \xi) = h_j \left(\frac{|x - x_0|}{R} \right) h_j \left(\frac{\xi}{R} \right) \text{ for } (x, \xi) \in B_R(x_0) \times \mathbb{R}.$$

Multiplying first (5.1) by $\bar{p}_j \eta_j^2$ and then integrating by parts over $D_j = B_{a_j R}(x_0) \times (-a_j R, a_j R)$, we obtain

$$\begin{aligned} & - \int_{D_j} \nabla \bar{p}_j \cdot (A \nabla p_j) \eta_j^2 dx d\xi - i \operatorname{sign}(\mu) \int_{D_j} |p_{j+1}|^2 \eta_j^2 dx d\xi \\ & = \int_{D_j} \nabla \eta_j^2 \cdot (A \nabla p_j) \bar{p}_j dx d\xi + i \operatorname{sign}(\mu) \int_{D_j} p_{j+1} \partial_\xi \eta_j^2 \bar{p}_j dx d\xi. \end{aligned} \quad (5.2)$$

Since $\nabla \bar{p}_j \cdot (A \nabla p_j)$ and $|p_{j+1}|^2$ are real-valued, by (5.2) we get

$$\left(\int_{D_j} \nabla \bar{p}_j \cdot (A \nabla p_j) \eta_j^2 dx d\xi \right)^2 + |\operatorname{sign}(\mu)|^2 \left(\int_{D_j} |p_{j+1}|^2 \eta_j^2 dx d\xi \right)^2$$

$$\leq 2 \left(\int_{D_j} |\nabla \eta_j^2 \cdot (A \nabla p_j)| |p_j| dx d\xi \right)^2 + 2 \left(\int_{D_j} |p_{j+1}| |p_j| |\partial_\xi \eta_j^2| dx d\xi \right)^2 := 2 \sum_{i=1}^2 I_i. \quad (5.3)$$

Next, we will estimate I_i ($i = 1, 2$) one by one. By Young's inequality, we have

$$\begin{aligned} & \int_{D_j} |\nabla \eta_j^2 \cdot (A \nabla p_j)| |p_j| dx d\xi \leq 2\lambda \int_{D_j} |\nabla \eta_j| |\eta_j| |\nabla p_j| |p_j| dx d\xi \\ & \leq \epsilon_1 \lambda \int_{D_j} |\nabla p_j|^2 \eta_j^2 dx d\xi + \frac{\lambda}{\epsilon_1} \int_{D_j} |p_j|^2 |\nabla \eta_j|^2 dx d\xi \\ & \leq \epsilon_1 \lambda \int_{D_j} |\nabla p_j|^2 \eta_j^2 dx d\xi + \frac{\lambda \pi^2 m^2}{R^2 \epsilon_1} \int_{D_j} |p_j|^2 dx d\xi. \end{aligned}$$

Thus, we get that

$$I_1 \leq 2\lambda^2 \epsilon_1^2 \left(\int_{D_j} |\nabla p_j|^2 \eta_j^2 dx d\xi \right)^2 + \frac{2\lambda^2 \pi^4 m^4}{R^4 \epsilon_1^2} \left(\int_{D_j} |p_j|^2 dx d\xi \right)^2, \quad \forall \epsilon_1 > 0. \quad (5.4)$$

Furthermore,

$$\begin{aligned} \int_{D_j} |p_{j+1}| |p_j| |\partial_\xi \eta_j^2| dx d\xi & \leq \epsilon_2 \int_{D_j} |p_{j+1}|^2 \eta_j^2 dx d\xi + \frac{1}{\epsilon_2} \int_{D_j} |p_j|^2 |\partial_\xi \eta_j|^2 dx d\xi \\ & \leq \epsilon_2 \int_{D_j} |p_{j+1}|^2 \eta_j^2 dx d\xi + \frac{m^2 \pi^2}{R^2 \epsilon_2} \int_{D_j} |p_j|^2 dx d\xi. \end{aligned}$$

Thus, we have that

$$I_2 \leq 2\epsilon_2^2 \left(\int_{D_j} |p_{j+1}|^2 \eta_j^2 dx d\xi \right)^2 + \frac{2\pi^4 m^4}{R^4 \epsilon_2^2} \left(\int_{D_j} |p_j|^2 dx d\xi \right)^2, \quad \forall \epsilon_2 > 0. \quad (5.5)$$

Taking $\epsilon_1 = \sqrt{2}/(4\lambda^2)$, $\epsilon_2 = 1/4$ in (5.4) and (5.5), respectively, we derive that

$$\begin{aligned} \sum_{i=1}^2 I_i & \leq \frac{\lambda^{-2}}{4} \left(\int_{D_j} |\nabla p_j|^2 \eta_j^2 dx d\xi \right)^2 + \frac{1}{8} \left(\int_{D_j} |p_{j+1}|^2 \eta_j^2 dx d\xi \right)^2 \\ & \quad + \frac{(2 + \lambda^6) 4^2 \pi^4 m^4}{R^4} \left(\int_{D_j} |p_j|^2 dx d\xi \right)^2. \end{aligned} \quad (5.6)$$

On the other hand, by the uniform ellipticity condition (1.2), we find that

$$\left(\int_{D_j} \nabla \bar{p}_j \cdot (A \nabla p_j) \eta_j^2 dx d\xi \right)^2 \geq \lambda^{-2} \left(\int_{D_j} |\nabla p_j|^2 \eta_j^2 dx d\xi \right)^2.$$

This, together with (5.3) and (5.6), gives that for each $j \in \{0, 1, \dots, m-1\}$,

$$\int_{D_{j+1}} |p_{j+1}|^2 dx d\xi = \int_{D_{j+1}} |p_{j+1}|^2 \eta_j^2 dx d\xi \leq \sqrt{\frac{4^3 (2 + \lambda^6) \pi^4 m^4}{3R^4}} \int_{D_j} |p_j|^2 dx d\xi$$

$$\leq \frac{C_4 m^2}{R^2} \int_{D_j} |p_j|^2 dx d\xi, \quad (5.7)$$

where $C_4 = 8\pi^2 \sqrt{2 + \lambda^6}$. Here, we used the definition of D_j . Iterating (5.7) for each $j \in \{0, 1, \dots, m-1\}$, by the fact that $p_0 = p = \hat{v}$ we obtain (2.23) and complete the proof. \square

Acknowledgement. The authors thank the financial support by the National Natural Science Foundation of China under grants 11971363.

References

- [1] J. Apraiz, L. Escauriaza, Null-control and measurable sets, *ESAIM Control Optim. Calc. Var.* 19 (2013), 239-254.
- [2] J. Apraiz, L. Escauriaza, G. Wang, C. Zhang, Observability inequalities and measurable sets, *J. Eur. Math. Soc.* 16 (2014), 2433-2475.
- [3] V. Barbu, Exact null internal controllability for the heat equation on unbounded convex domains, *ESAIM: COCV* 20 (2014), 222-235.
- [4] H. Brezis, *Functional Analysis, Sobolev Space and Partial Differential Equations*, Springer, New York, 2011.
- [5] V. R. Cabanillas, S. B. de Menezes, E. Zuazua, Null controllability in unbounded domains for the semilinear heat equation with nonlinearities involving gradient terms, *J. Optim. Theory App.* 110 (2001), 245-264.
- [6] P. Cannarsa, P. Martinez, J. Vancostenoble, Null controllability of the heat equation in unbounded domains by a finite measure control region, *ESAIM: COCV* 10 (2004), 381-408.
- [7] B. Canuto, E. Rosset, S. Vessella, Quantitative estimates of unique continuation for parabolic equations and inverse initial-boundary value problems with unknown boundaries, *Trans. Amer. Math. Soc.* 354 (2002), 491-535.
- [8] Y. Duan, L. Wang, C. Zhang, Observability inequalities for the heat equation with bounded potentials on the whole space, *SIAM J. Control Optim.* 58 (4) (2020) 1939-1960.
- [9] M. Egidi, I. Veselić, Sharp geometric condition for null-controllability of the heat equation on \mathbb{R}^d and consistent estimates on the control cost, *Arch. Math. (Basel)* 111 (2018), 85-99.
- [10] M. Egidi, A. Seelmann, The reflection principle in the control problem of the heat equation, arXiv preprint, arXiv:1902.08141, (2019).
- [11] L. Escauriaza, F. J. Fernández, S. Vessella, Doubling properties of Caloric functions, *Appl. Anal.* 85 (2006), 205-223.
- [12] L. Escauriaza, S. Montaner, C. Zhang, Observation from measurable sets for parabolic analytic evolutions and applications, *J. Math. Pures Appl.* 9 (5) (2015), 837-867.
- [13] L. Escauriaza, S. Montaner, C. Zhang, Analyticity of solutions to parabolic evolutions and applications, *SIAM J. Math. Anal.* 49 (2017), 4064-4092.

- [14] E. Fernández-Cara, E. Zuazua, Null and approximate controllability for weakly blowing up semilinear heat equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 17 (2000), 583-616.
- [15] A. V. Fursikov, O. Y. Imanuvilov, *Controllability of Evolution Equations*, Seoul National University, 1996.
- [16] D. Gilbarg, N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1983.
- [17] M. González-Burgos, L. de Teresa, Some results on controllability for linear and nonlinear heat equations in unbounded domains, *Adv. Diff. Equ.*, 12 (2007), 1201-1240.
- [18] E. M. Landis, O. A. Oleinik, Generalized analyticity and some related properties of solutions of elliptic and parabolic equations, *Russian Math. Surveys* 29 (1974), 195-212.
- [19] J. Le Rousseau, I. Moyano, Null-controllability of the Kolmogorov equation in the whole phase space, *J. Differ. Equ.* 260 (2016), 3193-3233.
- [20] G. Lebeau, I. Moyano, Spectral inequalities for the Schrödinger operator, arXiv preprint, arXiv:1901.03513,(2019).
- [21] G. Lebeau, E. Zuazua, Null-controllability of a system of linear thermoelasticity, *Arch. Rational Mech. Anal.* 141 (1998), 297-329.
- [22] F. H. Lin, A uniqueness theorem for parabolic equations, *Comm. Pure Appl. Math.* 43 (1990), 127-136.
- [23] F. H. Lin, Nodal sets of solutions of elliptic and parabolic equations, *Comm. Pure Appl. Math.* 44 (1991), 287-308.
- [24] A. Logunov, E. Malinnikova, Quantitative propagation of smallness for solutions of elliptic equations, *Proceedings of the International Congress of Mathematicians-Rio de Janeiro 2018. Vol. III. Invited lectures*, 2391-2411, World Sci. Publ., Hackensack, NJ, 2018.
- [25] Q. Lü, A lower bound on local energy of partial sum of eigenfunctions for Laplace-Beltrami operators, *ESAIM: COCV* 19 (1) (2013), 255-273.
- [26] E. Malinnikova, S. Vessella, Quantitative uniqueness for elliptic equations with singular lower order terms, *Math. Ann.* 353 (2012), 1157-1181.
- [27] S. Micu, E. Zuazua, On the lack of null-controllability of the heat equation on the half-line, *Trans. Amer. Math. Soc.*, 353 (2001), 1635-1659.
- [28] S. Micu, E. Zuazua, On the lack of null-controllability of the heat equation on the half space, *Port. Math.* 58 (2001), 1-24.
- [29] L. Miller, On the null-controllability of the heat equation in unbounded domains, *Bull. Sci. math.* 129 (2005), 175-185.
- [30] I. Nakić, M. Täufer, M. Tautenhahn, I. Veselić, Sharp estimates and homogenization of the control cost of the heat equation on large domains, *ESAIM: COCV* 26 (2020) 26-54.
- [31] K. D. Phung, G. Wang, Quantitative unique continuation for the semilinear heat equation in a convex domain, *J. Funct. Anal.* 259 (2010), 1230-1247.

- [32] K. D. Phung, G. Wang, An observability estimate for parabolic equations from a measurable set in time and its applications, *J. Eur. Math. Soc.* 15 (2013), 681-703.
- [33] K. D. Phung, L. Wang, C. Zhang, Bang-bang property for time optimal control of semilinear heat equation, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 31 (2014), 477-499.
- [34] G. Wang, M. Wang, C. Zhang, Y. Zhang, Observable set, observability, interpolation inequality and spectral inequality for the heat equation in \mathbb{R}^N , *J. Math. Pures Appl.* 126 (2019), 144-194.
- [35] G. Wang, C. Zhang, Observability inequalities from measurable sets for some evolution equations, *SIAM J. Control Optim.* 55 (2017), 1862-1886.
- [36] Y. Zhang, Unique continuation estimates for the Kolmogorov equation in the whole space, *C. R. Math. Acad. Sci. Paris.*, 354 (2016), 389-393.