

CO-ROTATING AND TRAVELING VORTEX SHEETS FOR THE 2D INCOMPRESSIBLE EULER EQUATION

DAOMIN CAO, GUOLIN QIN, CHANGJUN ZOU

ABSTRACT. We construct co-rotating and traveling vortex sheets for 2D incompressible Euler equation, which are supported on several small closed curves. These examples represent a new type of vortex sheet solutions other than two known classes. The construction is based on Birkhoff-Rott operator, and accomplished by using implicit function theorem at point vortex solutions with suitably chosen function spaces.

Keywords: The Euler equation, vortex sheets, Birkhoff-Rott operator, implicit function theorem

2020 MSC Primary: 76B47; **Secondary:** 35Q31, 76B03.

1. INTRODUCTION AND MAIN RESULTS

The vorticity formulation of 2D incompressible Euler equation in the whole space reads as follows

$$\begin{cases} \partial_t \vartheta + \mathbf{v} \cdot \nabla \vartheta = 0 & \text{in } \mathbb{R}^2 \times (0, T), \\ \vartheta|_{t=0} = \vartheta_0 & \text{in } \mathbb{R}^2, \end{cases} \quad (1.1)$$

where $\mathbf{v} = (v_1, v_2)$ is the velocity field, and $\vartheta = \text{curl} \mathbf{v} := \partial_1 v_2 - \partial_2 v_1$ is the vorticity function describing the rotating of fluid. Using the incompressible condition $\nabla \cdot \mathbf{v} = 0$, the velocity can be recovered by the Biot-Savart law from ϑ :

$$\mathbf{v} = \nabla^\perp (-\Delta)^{-1} \vartheta,$$

where $(x_1, x_2)^\perp = (x_2, -x_1)$ and the operator $(-\Delta)^{-1}$ is given by

$$(-\Delta)^{-1} \vartheta(x) = \int_{\mathbb{R}^2} G(x-y) \vartheta(y) dy.$$

Here $G(x, y) = \frac{1}{2\pi} \ln \frac{1}{|x-y|}$ is the fundamental solution of $-\Delta$ in \mathbb{R}^2 .

In this paper, we are concerned with the existence of co-rotating and traveling vortex sheet solutions to (1.1). Generally speaking, a vortex sheet is a weak solution in $\mathcal{M}(\mathbb{R}^2) \cap H^{-1}(\mathbb{R}^2)$ concentrated on simple closed curves $\Gamma_i := \{\mathbf{z}_i(t, x) \in \mathbb{R}^2 : x \in [0, 2\pi)\}$ with

This work was supported by NNSF of China Grant 11831009 and Chinese Academy of Sciences (No. QYZDJ-SSW-SYS021).

vortex strength ω_i , namely

$$\vartheta(t, \mathbf{x}) = \sum_{i=0}^{m-1} \omega_i(t, x) \delta_{\Gamma_i(t, \mathbf{z})},$$

where $\omega_i(t, x)$ is the vorticity strength with respect to the parametrization \mathbf{z}_i , and δ_Γ is the Dirac measure on the curve Γ . The above form of vorticity is understood in the sense that

$$\int_{\mathbb{R}^2} \varphi(x) d\vartheta(t, \mathbf{x}) = \sum_{i=0}^{m-1} \int_0^{2\pi} \varphi(\mathbf{z}_i(t, x)) \omega_i(t, x) dx, \quad \forall \varphi(x) \in C_c^\infty(\mathbb{R}^2).$$

In the formulation of a vortex sheet solution $\vartheta(t, \mathbf{x})$, we take $\gamma(t, x) = \omega(t, x)/|\partial_x \mathbf{z}(t, x)|$ be the regularized vortex strength. According to the Birkhoff-Rott operator, the velocity field generated by the vortex sheet Γ is given by

$$\mathbf{v}(\mathbf{z}(t, x), t) = -\mathcal{BR}(\mathbf{z}, \gamma) := -\frac{1}{2\pi} PV \int_0^{2\pi} \frac{(\mathbf{z}(t, x) - \mathbf{z}(t, y))^\perp}{|\mathbf{z}(t, x) - \mathbf{z}(t, y)|^2} \gamma(t, y) dy, \quad (1.2)$$

with PV the notation of principle value. In the fluid near a sheet, the respective limits of velocity on one side of Γ_i is denoted by \mathbf{v}_i^+ , while on the other is denoted by \mathbf{v}_i^- . Although the velocity field has a ‘‘jump’’ across Γ_i which causes some kind of singularity, we can define the velocity \mathbf{v}_{Γ_i} on Γ_i by imposing $\mathbf{v}_{\Gamma_i} = (\mathbf{v}_i^+ + \mathbf{v}_i^-)/2$. Using above notations, the evolution of $\mathbf{z}(t, x)$ and $\gamma(t, x)$ can be described by

$$\partial_t \mathbf{z}(t, x) = \mathbf{v}(\mathbf{z}, t) + c(t, x) \partial_x \mathbf{z}(t, x), \quad (1.3)$$

$$\partial_t \gamma(t, x) = \partial_x (c(t, x) \gamma(t, x)), \quad (1.4)$$

where $c(t, x)$ is the reparametrization freedom of the sheet curve.

The study of vortex sheets arises from the phenomenon of discontinuity for velocity field across an interface, which is caused by small viscosity flow passing rigid walls or sharp corners. It is observed that although the viscosity plays an important role in the generation of vortex sheets, the later transportation and evolution of vortices can be modeled by the incompressible Euler equation [2, 21]. However, the vortex sheet structure is proved to be extremely unstable, which leads to coherent vortex pattern often found in mixing layers, jets and wakes.

Before discussing our main topic of constructing special vortex sheets, let us first take some time to review the effort on existence of solutions to (1.1) in different settings: In 1963, Yudovich [29] proved the global well-posedness of (1.1) with the initial data in $L^1 \cap L^\infty$. This result covers the ‘‘vortex patch’’ initial data solution, but is not applicable to more singular cases. For an L_{loc}^2 initial data whose vorticity has a definite sign, Delort [11] and Majda [20] proved the global existence of a weak solution separately. When the vorticity does not have a definite sign, by imposing reflection symmetry on the initial vorticity, the global existence was proved by Lopes Filho et al. in [18]. In the case that initial data $\mathbf{z}(0, x)$, $\gamma(0, x)$ for a vortex sheet are analytic, Sulem et al. [26] obtained local existence for the analytic solution, which corresponds to the cases we discuss in this paper. On the other hand, Birkhoff et al. [3, 4] conjectured that curvature may blow up in finite time for

vortex sheets with analytic initial data. This was verified by asymptotic analysis of Moore [23] and numerical simulations of Krasny [19], Meiron et al. [22]. It is notable that the system (1.3) and (1.4) was proved to be ill-posed in H^s for $s > 3/2$ in [5]. For more results on the well-posedness theory, we refer to [21, 25, 28] and references therein.

Concerning the existence of global solutions (also known as relative equilibria) to (1.1), many vivid examples whose vorticity is supported on compact areas are known, which can be roughly divided into two classes: one contains solutions with relatively large support, which are bifurcated from known solutions including disks, annuli, Kirchhoff ellipses or Lamb dipole, see [8, 10, 16]; the other contains solutions with highly concentrated vorticity, which can be regarded as a regularization of point vortex solutions, see [6, 7, 17]. Different approaches were developed for construction of this type, such as study of contour dynamic equation, variational method, Lyapunov-Schmidt reduction, etc.

However, for equation (1.1), there are only two known classes of relative equilibria composed of vortex sheets other than circles and lines before our results which will be given later in this paper (see Theorem 1.1 and Theorem 1.2). The first one is Protas-Sakajo class [24], which is made out of segments rotating about a common center of rotation with endpoints at the vertices of a regular polygon. Duchon and Robert [12] discussed the global existence when the initial data near one example constructed in [3], where the vorticity is supported on a segment of length $2l$ with density

$$\gamma(x) = \Omega \sqrt{l^2 - x^2}, \quad \text{for } x \in [-l, l]$$

and angular velocity Ω . ($\Omega > 0$ for counterclockwise)

Recently, Gómez-Serrano et al. [13] gave another vortex sheet supported on a closed curve. By Lyapunov-Schmidt reduction and degenerate bifurcation, they obtained two curves of solution bifurcated from one trivial solution, which is a unit circle at specific angular velocity. This remarkable result indicates that solutions bifurcated from concentric circles can be constructed in the same way, and constitute the second class of vortex sheet solutions. In their follow-up work [14], they have further showed that one can not expect vortex sheets other than concentric circles when $\omega \geq 0$ and $\Omega \leq 0$.

In the present paper, using the Birkhoff-Rott operator (1.1), we will give a new class of vortex sheet solutions to (1.1), whose support is on several small closed curves. Our strategy is to transform (1.3) and (1.4) into an integro-differential system, where well-located point vortices are observed to be trivial solutions. By extending the range of parameter ε to a neighborhood of 0, we hope that there exist nontrivial vortex sheet solutions near the trivial ones. However, having done the local linearization, we find the Gateaux derivatives do not give an isomorphism, which makes the implicit function theorem invalid. To obtain local curves of nontrivial vortex sheet solutions, a few techniques of subtle analysis are adopted. We will introduce a series of quotient function spaces for this problem, and let the angular velocity Ω and traveling speed W vary with the unknown pair (z, γ) . Then the isomorphism condition for linearized operator will prove to be equivalent to a dynamic relationship, which exactly corroborates our physical intuition. From this point of view, the existence of co-rotating and traveling vortex sheets can be established by the Implicit function theorem in a refined version. Since Dirac measure at one point is in $H^{-1-\tau}(\mathbb{R}^2)$

for any $\tau > 0$, which is more singular than a vortex sheet, our result can also be regarded as regularization of point vortices in a special way.

To transform the construction of vortex sheets into a nonlinear problem of our preference, we introduce some necessary notations. For $\varepsilon \in (0, 1/2)$, we denote

$$\Gamma^\varepsilon := \{\mathbf{z}(t, x) \in \mathbb{R}^2 : x \in [0, 2\pi)\}$$

as a simple closed curve encircling the origin, which is close to the circle with radius ε centered at the origin. In the rest of this paper, the unit vector in x_i direction ($i = 1, 2$) is denoted by \mathbf{e}_i . We also use Q_θ to denote the counterclockwise rotation operator of angle θ with respect to the origin, and use

$$\oint g(\tau) d\tau := \frac{1}{2\pi} \int_0^{2\pi} g(\tau) d\tau$$

to denote the mean value of integral on the unit circle. For simplicity, we will omit the principal value notation PV for possible singularities in the integral.

The first kind of relative equilibria we are looking for are co-rotating sheet solutions. Suppose the solution ϑ_ε is composed of m -fold symmetric sheets rotating in a uniform angular velocity Ω centered at $(d, 0)$, that is

$$\vartheta_\varepsilon(t, \mathbf{x}) = \vartheta_{0,\varepsilon}(Q_{-\Omega t}(\mathbf{x} - d\mathbf{e}_1) + d\mathbf{e}_1) = \frac{1}{\varepsilon} \sum_{i=0}^{m-1} \omega(t, x) \delta_{\Gamma_i^\varepsilon(t, \mathbf{z})}, \quad (1.5)$$

where $\Gamma_i^\varepsilon(t, \mathbf{z}) \subset \mathbb{R}^2$ are closed curves satisfying

$$\Gamma_i^\varepsilon(t, \mathbf{z}) - d\mathbf{e}_1 = Q_{\frac{2\pi i}{m} - \Omega t}(\Gamma^\varepsilon(\mathbf{z}) - d\mathbf{e}_1), \quad i = 0, \dots, m-1$$

with some $d > 1$ fixed. In this rotating frame, the evolution equations (1.3) and (1.4) become

$$\partial_t \mathbf{z}(t, x) = \mathbf{v}(t, \mathbf{z}) + \Omega(\mathbf{z}(t, x) - d\mathbf{e}_1) + c(t, x) \partial_x \mathbf{z}(t, x), \quad (1.6)$$

and

$$\partial_t \gamma(t, x) = \partial_x (c(t, x) \gamma(t, x)), \quad (1.7)$$

with $c(t, x)$ being the reparameterization freedom introduced before. Since $\mathbf{z}(t, x)$ parameterizes the same curve Γ^ε as $\mathbf{z}(0, x)$, $\partial_t \mathbf{z}(t, x)$ is tangent to Γ^ε . Let $\mathbf{n}(\mathbf{z}(t, x))$ be the unit outward vector at $\mathbf{z}(t, x)$, and $\mathbf{s}(\mathbf{z}(t, x))$ be the unit tangent vector counterclockwise. Multiplying (1.6) by $\mathbf{n}(\mathbf{z}(t, x))$ and using $\mathbf{n}(\mathbf{z}(t, x)) \cdot \partial_x \mathbf{z}(t, x) = 0$, we obtain

$$(\mathbf{v}(t, \mathbf{z}) + \Omega(\mathbf{z}(t, x) - d\mathbf{e}_1)^\perp) \cdot \mathbf{n}(\mathbf{z}(t, x)) = 0, \quad \forall \mathbf{z}(t, x) \in \Gamma^\varepsilon. \quad (1.8)$$

To eliminate time parameter and derive another necessary relationship, we choose

$$c(t, x) = -\frac{(\mathbf{v}(t, \mathbf{z}) + \Omega(\mathbf{z}(t, x) - d\mathbf{e}_1)^\perp) \cdot \mathbf{s}(\mathbf{z}(t, x))}{|\partial_x \mathbf{z}(t, x)|},$$

and hence obtain $\partial_t \mathbf{z}(t, x) \cdot \mathbf{s}(\mathbf{z}(t, x)) = 0$ by multiplying $\mathbf{s}(\mathbf{z}(t, x))$ to (1.6). Note that $\mathbf{s}(\mathbf{z}(t, x))$ is parallel to $\partial_t \mathbf{z}(t, x)$ and has unit length. So it holds $\partial_t \mathbf{z}(t, x) = 0$, which implies that $\mathbf{z}(t, x) = \mathbf{z}(x)$ remains fixed in time. On the other hand, ϑ_ε is stationary in

the rotating frame. Since $\mathbf{z}(t, x) = \mathbf{z}(x)$ is invariant, the regularized vortex strength will not evolve with time, or namely $\gamma(t, x) = \gamma(x)$. From (1.7), we derive

$$c(t, x)\gamma(t, x) = C.$$

According to the definition of $c(t, x)$ given above, we have

$$\frac{(\mathbf{v}(\mathbf{z}) + \Omega(\mathbf{z}(x) - d\mathbf{e}_1)^\perp) \cdot \mathbf{s}(\mathbf{z}(x))\gamma(x)}{|\partial_x \mathbf{z}(x)|} = -C, \quad \forall x \in [0, 2\pi). \quad (1.9)$$

Now we can assume that $\mathbf{z}(x)$ is parameterized as

$$\mathbf{z}(x) = (\varepsilon R(x) \cos(x), \varepsilon R(x) \sin(x))$$

for some function $R(x)$ close to 1. From (1.8) (1.9) and the definition of Birkhoff-Rott operator (1.2), if we let

$$\begin{aligned} F_1(\varepsilon, \Omega, R, \gamma) &= \Omega(\varepsilon R(x)R'(x) - dR'(x) \cos(x) + dR(x) \sin(x)) \\ &+ \frac{1}{\varepsilon} \int \frac{R(x)R'(x) - R'(x)R(y) \cos(x-y) + R(x)R(y) \sin(x-y)}{(R(x) - R(y))^2 + 4R(x)R(y) \sin^2(\frac{x-y}{2})} \times \gamma(y) dy \\ &+ \sum_{i=1}^{m-1} \int \frac{R(x)R'(x) - R'(x)R(y) \cos(x-y - \frac{2\pi i}{m}) + R(x)R(y) \sin(x-y - \frac{2\pi i}{m})}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(\mathbf{z}(y) - (d, 0)) \right|^2} \times \gamma(y) dy \\ &- \sum_{i=1}^{m-1} \int \frac{dR(x) (1 - \cos(\frac{2\pi i}{m})) \sin(x) - dR'(x) (1 - \cos(\frac{2\pi i}{m})) \cos(x)}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(\mathbf{z}(y) - (d, 0)) \right|^2} \times \gamma(y) dy, \end{aligned}$$

and

$$\begin{aligned} \tilde{F}_2(\varepsilon, \Omega, R, \gamma) &= \frac{\gamma(x)\Omega}{R'(x)^2 + R(x)^2} \cdot (-\varepsilon R(x)^2 + dR(x) \cos(x) + dR'(x) \sin(x)) \\ &+ \frac{1}{\varepsilon} \frac{\gamma(x)}{R'(x)^2 + R(x)^2} \int \frac{-R(x)^2 + R(x)R(y) \cos(x-y) + R'(x)R(y) \sin(x-y)}{(R(x) - R(y))^2 + 4R(x)R(y) \sin^2(\frac{x-y}{2})} \times \gamma(y) dy \\ &+ \sum_{i=1}^{m-1} \frac{\gamma(x)}{R'(x)^2 + R(x)^2} \int \frac{-R(x)^2 + R(x)R(y) \cos(x-y - \frac{2\pi i}{m}) + R'(x)R(y) \sin(x-y - \frac{2\pi i}{m})}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(\mathbf{z}(y) - (d, 0)) \right|^2} \times \gamma(y) dy \\ &- \sum_{i=1}^{m-1} \frac{\gamma(x)}{R'(x)^2 + R(x)^2} \int \frac{dR(x) (1 - \cos(\frac{2\pi i}{m})) \cos(x) + dR'(x) (1 - \cos(\frac{2\pi i}{m})) \sin(x)}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(\mathbf{z}(y) - (d, 0)) \right|^2} \times \gamma(y) dy, \end{aligned}$$

then the construction of co-rotating solutions is transformed to finding $R(x)$ and $\gamma(x)$ such that

$$\begin{cases} F_1(\varepsilon, \Omega, R, \gamma) = 0, \\ F_2(\varepsilon, \Omega, R, \gamma) = (I - \mathcal{P})\tilde{F}_2 = 0, \end{cases} \quad (1.10)$$

where \mathcal{P} is the operator of projection to the mean, namely, $\mathcal{P}g := \int g(x) dx$.

By choosing suitable quotient function spaces, we will show that the structure of F_1 and F_2 allows us to use implicit function theorem at $(\varepsilon, \Omega, R, \gamma) = (0, \Omega^*, 1, 1)$ with

$$\Omega^* := \frac{m-1}{2d^2}, \quad (1.11)$$

and thus establish the existence of a family of co-rotating vortex sheet solutions to (1.1) generated from $(0, \Omega^*, 1, 1)$. One can also verify that $(0, \Omega^*, 1, 1)$ corresponds to co-rotating m -fold point vortices solutions to (1.1), where the intensity of every single point vortex is 2π , the perimeter of unit circle. Now we can state our first result.

Theorem 1.1. *Suppose that $m \geq 2$. Then there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$, (1.1) has a global co-rotating vortex sheet solution $\vartheta_\varepsilon(\mathbf{x}, t) = \vartheta_{0,\varepsilon}(Q_{-\Omega t}(\mathbf{x} - d\mathbf{e}_1) + d\mathbf{e}_1)$ defined in (1.5), and the uniform angular velocity Ω_ε satisfies*

$$\Omega = \Omega^* + O(\varepsilon)$$

with Ω^* given in (1.11). Moreover, each component of the solution is supported on a C^∞ closed curve with convex interior.

Next we turn to the existence of traveling sheet solutions. In this case, the solution ϑ_ε is composed of two mirror symmetric sheets with opposite sign traveling at a uniform speed $W > 0$ in x_2 -direction, namely

$$\vartheta_\varepsilon(t, \mathbf{x}) = \vartheta_{0,\varepsilon}(\mathbf{x} - tW\mathbf{e}_2) = \frac{1}{\varepsilon}\omega(t, x)\delta_{\Gamma_O^\varepsilon(t, \mathbf{z})} - \frac{1}{\varepsilon}\omega(t, x)\delta_{\Gamma_T^\varepsilon(t, \mathbf{z})}, \quad (1.12)$$

where $\Gamma_O^\varepsilon(t, \mathbf{z}), \Gamma_T^\varepsilon(t, \mathbf{z}) \subset \mathbb{R}^2$ satisfy

$$\Gamma_O^\varepsilon(t, \mathbf{z}) = \Gamma^\varepsilon(\mathbf{z}) + tW\mathbf{e}_2,$$

and

$$\Gamma_T^\varepsilon(t, \mathbf{z}) = -\Gamma^\varepsilon(\mathbf{z}) + 2d\mathbf{e}_1 + tW\mathbf{e}_2$$

with $d > 1$ the half distance of two sheets center. By a similar method as we investigate co-rotating sheet solutions, we can make ϑ_ε stationary in the translation frame, and $\mathbf{z}(t, x), \gamma(t, x)$ invariant of time. Hence the two dynamic equations for traveling sheet solutions are

$$(\mathbf{v}(\mathbf{z}) - W\mathbf{e}_2) \cdot \mathbf{n}(\mathbf{z}(x)) = 0, \quad \forall x \in [0, 2\pi), \quad (1.13)$$

and

$$\frac{(\mathbf{v}(\mathbf{z}) + \Omega(\mathbf{z}(x) - d\mathbf{e}_1)^\perp) \cdot \mathbf{s}(\mathbf{z}(x))\gamma(x)}{|\partial_x \mathbf{z}(x)|} = -C, \quad \forall x \in [0, 2\pi). \quad (1.14)$$

By doing a same parameterization for $\mathbf{z}(x)$ as before and using Birkhoff-Rott operator (1.2), we see that (1.13) and (1.14) are equivalent to

$$\begin{cases} G_1(\varepsilon, W, R, \gamma) = 0, \\ G_2(\varepsilon, W, R, \gamma) = (I - \mathcal{P})\tilde{G}_2 = 0, \end{cases} \quad (1.15)$$

where

$$\begin{aligned}
G_1(\varepsilon, W, R, \gamma) &= -W(R(x) \sin(x) - R'(x) \cos(x)) \\
&+ \frac{1}{\varepsilon} \int \frac{R(x)R'(x) - R'(x)R(y) \cos(x-y) + R(x)R(y) \sin(x-y)}{(R(x) - R(y))^2 + 4R(x)R(y) \sin^2(\frac{x-y}{2})} \times \gamma(y) dy \\
&- \int \frac{R(x)R'(x) + R'(x)R(y) \cos(x-y) - R(x)R(y) \sin(x-y)}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \times \gamma(y) dy \\
&+ \int \frac{2dR(x) \sin(x) - 2dR'(x) \cos(x)}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \times \gamma(y) dy,
\end{aligned}$$

and

$$\begin{aligned}
\tilde{G}_2(\varepsilon, W, R, \gamma) &= -\frac{\gamma(x)W}{R'(x)^2 + R(x)^2} \cdot (R'(x) \sin(x) + R(x) \cos(x)) \\
&+ \frac{1}{\varepsilon} \frac{\gamma(x)}{R'(x)^2 + R(x)^2} \int \frac{-R(x)^2 + R(x)R(y) \cos(x-y) + R'(x)R(y) \sin(x-y)}{(R(x) - R(y))^2 + 4R(x)R(y) \sin^2(\frac{x-y}{2})} \times \gamma(y) dy \\
&+ \frac{\gamma(x)}{R'(x)^2 + R(x)^2} \int \frac{R(x)^2 + R(x)R(y) \cos(x-y) + R'(x)R(y) \sin(x-y)}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \gamma(y) dy \\
&+ \frac{\gamma(x)}{R'(x)^2 + R(x)^2} \int \frac{dR(x) (1 - \cos(\frac{2\pi i}{m})) \cos(x) + dR'(x) (1 - \cos(\frac{2\pi i}{m})) \sin(x)}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \gamma(y) dy.
\end{aligned}$$

We can use implicit function theorem at $(\varepsilon, W, R, \gamma) = (0, W^*, 1, 1)$ with

$$W^* := \frac{1}{2d} \tag{1.16}$$

as before. Careful analysis gives a curve of nontrivial traveling vortex sheets passing the trivial one $(0, W^*, 1, 1)$, which represents the traveling point vortex pair with opposite intensity $\pm 2\pi$. This fact leads to our second theorem:

Theorem 1.2. *There exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$, (1.1) has a global traveling vortex sheet solution $\vartheta_\varepsilon(\mathbf{x}, t) = \vartheta_{0,\varepsilon}(\mathbf{x} - tW\mathbf{e}_2)$ defined in (1.12), and the uniform traveling speed W satisfies*

$$W = W^* + O(\varepsilon).$$

with W^* given in (1.16). Moreover, each component of the solution is supported on a C^∞ closed curve with convex interior.

Remark 1.3. Since $R(x)$ and $\gamma(x)$ will be chosen in analytic spaces, the support curves for vortex sheets obtained in Theorem 1.1 and 1.2 are not only C^∞ smooth, but also real analytic. For vortex patch in 2D incompressible Euler flow, the regularity of boundary is an interesting topic. Chemin [9] and Bertozzi et al. [1] showed that the patch boundary will persist $C^{1,\alpha}$ regularity. As a special case, Hmidi et al. [15] proved the C^∞ boundary regularity for relative equilibria bifurcated from disks. Different from the patch case, the regularity will not be preserved for a general vortex sheet initial data, see [2, 3].

It is notable that since there is no semilinear elliptic problem corresponding to vortex sheet, traditional variational method for vortex patch does not work here. Our method for concentrative vortex sheets is novel and widely applicable: various solutions can be constructed in this way, where each local cluster of sheet is supported on small concentric circles or has other different locations. However, the system will be more complicated in foresaid situations and it takes much computation time. We also bring our readers' attention to the fact that besides the existence, the local uniqueness for relative equilibria of such type remains another open problem.

This paper is organized, as follows: In Section 2, we introduce the functional settings and a special form of Taylor's formula. Then we will prove the existence of co-rotating vortex sheets in Section 3. The complete proof contains three essential elements: the verification of regularity for the integro-differential system, the linearization of system, and the application of implicit function theorem. In Section 4, the existence of traveling vortex sheets will be given by a brief discussion, since it is similar to that of the co-rotating case.

2. FUNCTIONAL SETTINGS AND TAYLOR'S FORMULA

For some small constant $a > 0$, let $C_w(a)$ denote the space of analytic functions in the strip $|\mathbf{Im}(z)| \leq a$. To apply the implicit function theorem, we define the even function spaces

$$X^k = \left\{ u \in C_w(a), u(x) = \sum_{j=1}^{\infty} a_j \cos(jx), \sum_{\pm} \int_0^{2\pi} |u(x \pm a\mathbf{i})|^2 + |\partial^k u(x \pm a\mathbf{i})|^2 dx \leq +\infty \right\},$$

and the odd function spaces

$$Y^k = \left\{ u \in C_w(a), u(x) = \sum_{j=1}^{\infty} a_j \sin(jx), \sum_{\pm} \int_0^{2\pi} |u(x \pm a\mathbf{i})|^2 + |\partial^k u(x \pm a\mathbf{i})|^2 dx \leq +\infty \right\},$$

where the norm of X^k, Y^k is the standard H^k norm on the strip $|\mathbf{Im}(z)| \leq a$, namely, for some $u_1 \in X^k$ and $u_2 \in Y^k$

$$\|u_1\|_{X^k} = \|u_1(x \pm a\mathbf{i})\|_{H^k} \quad \text{and} \quad \|u_2\|_{Y^k} = \|u_2(x \pm a\mathbf{i})\|_{H^k}.$$

Our construction has a distinguished feature: due to the spectral structure on Fourier series, the linearized operator is no longer an isomorphism on above spaces. To remediate to this weak point, we define the quotient function spaces

$$\mathcal{A}^k := \left\{ (u, v) \in X^{k+1} \times X^k \mid \int u(x) \cos(x) dx = - \int v(x) \cos(x) dx \right\},$$

$$\mathcal{B}^k := \left\{ (u, v) \in Y^k \times X^k \mid \int u(x) \sin(x) dx = - \int v(x) \cos(x) dx \right\},$$

which give a restriction on the first Fourier coefficient. On the other hand, in the first step we extend the range of ε from $(0, 1/2)$ to $(-1/2, 1/2)$, where the functionals should prove

to be well-defined. To eliminate possible singularity and derive asymptotic expansion for F_1, F_2 and G_1, G_2 , we will frequently use the following Taylor's formula in the proof of regularity part.

$$\frac{1}{(A+B)^\lambda} = \frac{1}{A^\lambda} - \lambda \int_0^1 \frac{B}{(A+tB)^{1+\lambda}} dt. \quad (2.1)$$

In the next two sections, we will consider the existence of co-rotating and traveling vortex sheets solutions separately. Since the problem is transformed to solving an integro-differential system, we will focus on dealing with relevant functionals.

3. EXISTENCE OF CO-ROTATING VORTEX SHEETS

To normalize our problem, we extend the range of ε to $(-\frac{1}{2}, \frac{1}{2})$. Suppose that $R(x) = 1 + \varepsilon f(x)$, $\gamma(x) = 1 + \varepsilon g(x)$ with $x \in [0, 2\pi)$, and f, g being two C^1 functions. From discussion in the previous section, we will consider following functionals

$$F_1(\varepsilon, \Omega, f, g) = F_{11} + F_{12} + F_{13} + F_{14},$$

where

$$F_{11} = \Omega \left(\varepsilon^2 (1 + \varepsilon f(x)) f'(x) - \varepsilon f'(x) d \cos(x) + (1 + \varepsilon f(x)) d \sin(x) \right), \quad (3.1)$$

$$\begin{aligned} F_{12} &= \frac{1}{\varepsilon} \int \frac{\varepsilon (1 + \varepsilon f(x)) f'(x) (1 - \cos(x-y)) (1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\ &\quad + \frac{1}{\varepsilon} \int \frac{\varepsilon^2 f'(x) (f(x) - f(y)) \cos(x-y) (1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\ &\quad - \frac{1}{\varepsilon} \int \frac{\varepsilon (1 + \varepsilon f(x)) (f(x) - f(y)) \sin(x-y) (1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\ &\quad + \frac{1}{\varepsilon} \int \frac{(1 + \varepsilon f(x))^2 \sin(x-y) (1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\ &= F_{121} + F_{122} + F_{123} + F_{124}, \end{aligned} \quad (3.2)$$

$$\begin{aligned} F_{13} &= \sum_{i=1}^{m-1} \int \frac{\varepsilon (1 + \varepsilon f(x)) f'(y) (1 + \varepsilon g(y))}{\left| (z(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (z(y) - (d, 0)) \right|^2} dy \\ &\quad - \sum_{i=1}^{m-1} \int \frac{\varepsilon f'(x) (1 + \varepsilon f(y)) \cos(x - y - \frac{2\pi i}{m}) (1 + \varepsilon g(y))}{\left| (z(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (z(y) - (d, 0)) \right|^2} dy \\ &\quad + \sum_{i=1}^{m-1} \int \frac{(1 + \varepsilon f(x)) (1 + \varepsilon f(y)) \sin(x - y - \frac{2\pi i}{m}) (1 + \varepsilon g(y))}{\left| (z(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (z(y) - (d, 0)) \right|^2} dy, \end{aligned} \quad (3.3)$$

$$\begin{aligned}
F_{14} &= \sum_{i=1}^{m-1} \int \frac{d(1 + \varepsilon f(x)) (1 - \cos(\frac{2\pi i}{m})) \sin(x)(1 + \varepsilon g(y))}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (\mathbf{z}(y) - (d, 0)) \right|^2} dy \\
&\quad - \sum_{i=1}^{m-1} \int \frac{\varepsilon df'(x) (1 - \cos(\frac{2\pi i}{m})) \cos(x)(1 + \varepsilon g(y))}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (\mathbf{z}(y) - (d, 0)) \right|^2} dy,
\end{aligned} \tag{3.4}$$

and

$$F_2(\varepsilon, \Omega, f, g) = (I - \mathcal{P})\tilde{F}_2 = (I - \mathcal{P})(\tilde{F}_{21} + \tilde{F}_{22} + \tilde{F}_{23} + \tilde{F}_{24}),$$

where

$$\tilde{F}_{21} = \frac{\Omega(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \cdot (-\varepsilon(1 + \varepsilon f(x))^2 + (1 + \varepsilon f(x))d \cos(x) + \varepsilon f'(x)d \sin(x)), \tag{3.5}$$

$$\begin{aligned}
\tilde{F}_{22} &= \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{(1 + \varepsilon f(x))(1 + \varepsilon f(y))(\cos(x - y) - 1)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2(\frac{x-y}{2})} dy \\
&\quad - \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon^2 f'(x)(f(x) - f(y)) \sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2(\frac{x-y}{2})} dy \\
&\quad - \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon(1 + \varepsilon f(x))(f(x) - f(y))(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2(\frac{x-y}{2})} dy \\
&\quad + \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon f'(x)(1 + \varepsilon f(x)) \sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2(\frac{x-y}{2})} dy \\
&= F_{221} + F_{222} + F_{223} + F_{224},
\end{aligned} \tag{3.6}$$

$$\begin{aligned}
\tilde{F}_{23} &= \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \sum_{i=1}^{m-1} \int \frac{(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \cos(x - y - \frac{2\pi i}{m})(1 + \varepsilon g(y))}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (\mathbf{z}(y) - (d, 0)) \right|^2} dy \\
&\quad + \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \sum_{i=1}^{m-1} \int \frac{\varepsilon f'(x)(1 + \varepsilon f(y)) \sin(x - y - \frac{2\pi i}{m})(1 + \varepsilon g(y))}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (\mathbf{z}(y) - (d, 0)) \right|^2} dy, \\
&\quad - \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \sum_{i=1}^{m-1} \int \frac{(1 + \varepsilon f(x))^2(1 + \varepsilon g(y))}{\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}} (\mathbf{z}(y) - (d, 0)) \right|^2} dy,
\end{aligned} \tag{3.7}$$

$$\begin{aligned}
\tilde{F}_{24} = & -\frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \sum_{i=1}^{m-1} \int \frac{d(1 + \varepsilon f(x)) (1 - \cos(\frac{2\pi i}{m})) \cos(x)(1 + \varepsilon g(y))}{\left| (z(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(z(y) - (d, 0)) \right|^2} dy \\
& -\frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \sum_{i=1}^{m-1} \int \frac{\varepsilon df'(x) (1 - \cos(\frac{2\pi i}{m})) \sin(x)(1 + \varepsilon g(y))}{\left| (z(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(z(y) - (d, 0)) \right|^2} dy.
\end{aligned} \tag{3.8}$$

We also denote $F_{2i} = (I - \mathcal{P})\tilde{F}_{2i}$ for $i = 1, 2, 3, 4$. The proof will be split into several lemmas, each of which is associated with checking the hypotheses of the implicit function theorem.

Let V^r be the open neighborhood of zero in \mathcal{A}^k

$$V^r := \{(f, g) \in \mathcal{A}^k : \|f\|_{X^{k+1}} < r, \|g\|_{X^k} < r\}$$

with $0 < r < 1$ and $k \geq 3$. In the following two lemmas, we investigate the regularity of $F_1(\varepsilon, \Omega, f, g)$, $F_2(\varepsilon, \Omega, f, g)$, and show the extension for ε to $(-\frac{1}{2}, \frac{1}{2})$ is reasonable.

Lemma 3.1. $F_1(\varepsilon, \Omega, f, g) : (-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R} \times V^r \rightarrow Y^k$ and $F_2(\varepsilon, \Omega, f, g) : (-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R} \times V^r \rightarrow X^k$ are both continuous.

Proof. Let us start with F_{11} . From (2.1), it is obvious that $F_{11} : (-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R} \times V^r \rightarrow X^k$ is continuous. We can rewrite F_{11} as

$$F_{11} = \Omega(d \sin(x) + \varepsilon \mathcal{R}_{11}(\varepsilon, f, g)), \tag{3.9}$$

where $\mathcal{R}_{11}(\varepsilon, f) : (-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R} \times V^r \rightarrow X^k$ is continuous.

The next step is to study F_{12} . To begin with, we show F_{12} is odd. Notice that $f(x) \in X^{k+1}$, $g(x) \in X^k$ are both even functions and $f'(x) \in Y^{k-1}$ is odd. By substituting $-y$ for y in F_{12} , we can deduce that $F_{12}(\varepsilon, f, g)$ is odd with respect to x .

Then we are going to prove that the range of F_{12} is in Y^k . Since $R(x) = 1 + \varepsilon f(x)$ and $\gamma(x) = 1 + \varepsilon g(x)$, the possible singularity for $\varepsilon = 0$ may occur only when we take zeroth derivative of F_{12} . More precisely, it may occur only in F_{124} . To prove this case can not happen, we use Taylor's formula (2.1) to do following decomposition

$$\frac{1}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} = \frac{1}{4 \sin^2\left(\frac{x-y}{2}\right)} + \varepsilon K_\varepsilon(x, y), \tag{3.10}$$

where we let $A = 4 \sin^2\left(\frac{x-y}{2}\right)$ and B the remaining terms in the denominator. As for this decomposition, an important fact is that $K_\varepsilon(x, y)$ is not singular at $x = y$. Since $\sin(\cdot)$ is odd, it holds

$$F_{124} = \frac{1}{\varepsilon} \int \frac{\sin(x-y)}{4 \sin^2\left(\frac{x-y}{2}\right)} dy + \varepsilon \mathcal{R}_{124}(\varepsilon, f, g) = \varepsilon \mathcal{R}_{124}(\varepsilon, f, g),$$

where $\mathcal{R}_{124}(\varepsilon, f, g)$ is not singular at $\varepsilon = 0$.

Make the change of variable $x - y \rightarrow y$, and take k -th partial derivatives of F_{12} with respect to x . Since we have $1 - \cos(y) = 2 \sin\left(\frac{y}{2}\right)$, for $\partial^k F_{121}$ it holds

$$\partial^k F_{121} = \frac{1}{2} \int (1 + \varepsilon f(x)) \partial^{k+1} f(x) (1 + \varepsilon g(x-y)) dy + \frac{1}{2} \int \varepsilon (1 + \varepsilon f(x)) f'(x) \partial^k g(x-y) dy + l.o.t.,$$

where we have used decomposition (3.10) and *l.o.t.* represents all lower order terms. Since $f(x) \in X^{k+1}$, $g(x) \in X^k$ and $k \geq 3$, we have $\|\partial^i f(x \pm \mathbf{a}\mathbf{i})\|_{L^\infty} \leq C\|f\|_{X^{k+1}} < \infty$ and $\|\partial^i g(x \pm \mathbf{a}\mathbf{i})\|_{L^\infty} \leq C\|g\|_{X^k} < \infty$ for $i = 0, 1, 2$. Using Hölder inequality and mean value theorem, we can conclude that

$$\|\partial^k F_{121}(x \pm \mathbf{a}\mathbf{i})\|_{L^2} \leq C\|f\|_{X^{k+1}} + C\|g\|_{X^k} < \infty.$$

We now turn to the most singular term $\partial^k F_{122}$, which has an expansion as

$$\begin{aligned} \partial^k F_{122} &= \int \frac{\varepsilon \partial^{k+1} f(x) (f(x) - f(x-y)) \cos(y) (1 + \varepsilon g(x-y))}{\varepsilon^2 (f(x) - f(x-y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(x-y)) \sin^2\left(\frac{y}{2}\right)} dy \\ &\quad + \int \frac{\varepsilon f'(x) (\partial^k f(x) - \partial^k f(x-y)) \cos(y) (1 + \varepsilon g(x-y))}{\varepsilon^2 (f(x) - f(x-y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(x-y)) \sin^2\left(\frac{y}{2}\right)} dy \\ &\quad - \int \frac{2\varepsilon^2 \partial^{k+1} f(x) (f(x) - f(x-y)) \cos(y) (1 + \varepsilon g(x-y))}{(\varepsilon^2 (f(x) - f(x-y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(x-y)) \sin^2\left(\frac{y}{2}\right))^2} \times \left(\varepsilon (f(x) - f(x-y)) \right. \\ &\quad \left. (f'(x) - f'(x-y)) + 2((1 + \varepsilon f(x))f'(x-y) + f'(x)(1 + \varepsilon f(x-y))) \sin^2\left(\frac{y}{2}\right) \right) dy \\ &\quad + \int \frac{\varepsilon f'(x) (f(x) - f(x-y)) \cos(y) (1 + \varepsilon \partial^k g(x-y))}{\varepsilon^2 (f(x) - f(x-y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(x-y)) \sin^2\left(\frac{y}{2}\right)} dy \\ &\quad + l.o.t. \\ &= \partial F_{1221} + \partial F_{1222} + \partial F_{1223} + \partial F_{1224} + l.o.t.. \end{aligned}$$

We first deal with ∂F_{1221} . Using the decomposition (3.10) with $\|K_\varepsilon(x, y)\|_{L^\infty} \leq C(\varepsilon, \|f\|_{x^{k+1}}) < \infty$ and the fact $|y|^2$ is the equivalent infinitesimal for $4 \sin^2\left(\frac{y}{2}\right)$ as $|y| \rightarrow 0$, it holds

$$\begin{aligned} \|\partial F_{1221}(x \pm \mathbf{a}\mathbf{i})\|_{L^2} &\leq C \left\| \int \frac{\varepsilon \partial^{k+1} f(x \pm \mathbf{a}\mathbf{i}) (f(x \pm \mathbf{a}\mathbf{i}) - f(x \pm \mathbf{a}\mathbf{i} - y)) \cos(y) (1 + \varepsilon g(x \pm \mathbf{a}\mathbf{i} - y))}{4 \sin^2\left(\frac{y}{2}\right)} dy \right\|_{L^2} \\ &\leq C\|f\|_{X^{k+1}} (1 + \|g(x \pm \mathbf{a}\mathbf{i})\|_{L^\infty}) \left| \int \frac{f(x \pm \mathbf{a}\mathbf{i}) - f(x \pm \mathbf{a}\mathbf{i} - y)}{4 \sin^2\left(\frac{y}{2}\right)} dy \right| \\ &\leq C\|f\|_{X^{k+1}} (1 + \|g\|_{X^k}) \left| \int \frac{f'(x \pm \mathbf{a}\mathbf{i}) + f''(x \pm \mathbf{a}\mathbf{i}) \cdot |y|}{2 \sin\left(\frac{y}{2}\right)} dy \right| \\ &\leq C\|f\|_{X^{k+1}} (1 + \|g\|_{X^k}) \int f''(x \pm \mathbf{a}\mathbf{i}) dy \\ &\leq C\|f\|_{X^{k+1}}^2 (1 + \|g\|_{X^k}), \end{aligned}$$

where we have used the identity $\int \frac{1}{\sin(\frac{y}{2})} dy = 0$. For the term ∂F_{1222} , one can use a similar method to obtain the upper bound

$$\|\partial F_{1222}(x \pm a\mathbf{i})\|_{L^2} \leq C\|f\|_{X^{k+1}}^2(1 + \|g\|_{X^k}).$$

Hence in the next we will focus on ∂F_{1223} . By Hölder inequality and mean value theorem, we deduce that

$$\begin{aligned} \|\partial F_{1223}(x \pm a\mathbf{i})\|_{L^2} &\leq C\|f\|_{X^{k+1}}(1 + \|g\|_{X^k}) \left| \int \frac{f(x \pm a\mathbf{i}) - f(x \pm a\mathbf{i} - y)}{4 \sin^2(\frac{y}{2})} dy \right| \\ &\quad \times (\|f'(x \pm a\mathbf{i})\|_{L^\infty} \|f''(x \pm a\mathbf{i})\|_{L^\infty} + (1 + \|f(x \pm a\mathbf{i})\|_{L^\infty}) \|f'(x \pm a\mathbf{i})\|_{L^\infty}) \\ &\leq C\|f\|_{X^{k+1}}^3(1 + \|f\|_{X^{k+1}})(1 + \|g\|_{X^k}) \end{aligned}$$

through a direct calculation. In the last, it is easy to obtain

$$\|\partial F_{1224}(x \pm a\mathbf{i})\|_{L^2} \leq C\|f\|_{X^{k+1}}^2(1 + \|g\|_{X^k}).$$

Since F_{122} is the most singular term in F_{12} , it always holds $\|F_{12}\|_{Y^k} \leq C\|F_{122}\|_{Y^k} \leq \infty$ from above discussion, and we can conclude that the range of F_{12} is in Y^k .

To prove the continuity of F_{12} , we also consider the most singular term F_{122} . We will use following notations: for a general function u , we denote

$$\Delta u = u(x) - u(y), \quad u = u(x), \quad \tilde{u} = u(y),$$

and

$$D(u) = \varepsilon^2 \Delta u^2 + 4(1 + \varepsilon u)(1 + \varepsilon \tilde{u}) \sin^2\left(\frac{x-y}{2}\right).$$

To show the continuity with respect to f , letting $(f_1, g), (f_2, g) \in V^r$, then we can calculate the difference

$$\begin{aligned} F_{122}(\varepsilon, f_1, g) - F_{122}(\varepsilon, f_2, g) &= (f'_1 - f'_2) \int \frac{\varepsilon \Delta f_1 \cos(x-y)(1 + \varepsilon \tilde{g}) dy}{D(f_1)} \\ &\quad + f'_2 \int \frac{\varepsilon (\Delta f_1 - \Delta f_2) \cos(x-y)(1 + \varepsilon \tilde{g}) dy}{D(f_1)} \\ &\quad + \left(\int \frac{f'_2 \Delta f_2 \cos(x-y)(1 + \varepsilon \tilde{g}) dy}{D(f_1)} - \int \frac{f'_2 \Delta f_2 \cos(x-y)(1 + \varepsilon \tilde{g}) dy}{D(f_2)} \right) \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

For the first two terms I_1 and I_2 , by the technique we have used before it is easy to prove

$$\|I_1\|_{Y^k} \leq C\|f_1 - f_2\|_{X^{k+1}}$$

and

$$\|I_2\|_{Y^k} \leq C\|f_1 - f_2\|_{X^{k+1}}.$$

For the last term I_3 , since

$$\begin{aligned} & \frac{1}{D(f_1)} - \frac{1}{D(f_2)} \\ &= \frac{\varepsilon^2(\Delta f_2^2 - \Delta f_1^2) + 4\varepsilon((f_2 - f_1)(1 + \varepsilon \tilde{f}_2) + (\tilde{f}_2 - \tilde{f}_1)(1 + \varepsilon f_1)) \sin^2(\frac{x-y}{2})}{D(f_1)D(f_2)}, \end{aligned} \quad (3.11)$$

it holds

$$\begin{aligned} \|\partial^k I_3(x \pm \mathbf{a}i)\|_{L^2} &\leq C \|f_2\|_{X^{k+1}}^2 (\|f_1\|_{X^{k+1}} + \|f_2\|_{X^{k+1}} + C) \|f_1 - f_2\|_{X^{k+1}} (1 + \|g\|_{X^k}) \\ &\leq C \|f_1 - f_2\|_{X^{k+1}}. \end{aligned}$$

On the other hand, using the fact that F_{122} is linear with respect to g , we can verify the continuity of F_{122} on g directly. Since F_{122} is the most singular term in F_{12} , we have actually shown that $F_{12}(\varepsilon, f, g) : (-\frac{1}{2}, \frac{1}{2}) \times V^r \rightarrow Y^k$ is continuous. For further use, we write F_{12} as

$$F_{12} = \frac{1}{2} f'(x) + \int \frac{g(y) \sin(x-y)}{4 \sin^2(\frac{x-y}{2})} dy + \varepsilon \mathcal{R}_{12}(\varepsilon, f, g), \quad (3.12)$$

where $\mathcal{R}_{12}(\varepsilon, f, g) : (-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R} \times V^r \rightarrow X^k$ is regular.

Having dealt with F_{12} , we are to study F_{13} and F_{14} . Notice that for $\mathbf{z} \in \Gamma^\varepsilon$ and $i \geq 1$, $\left| (\mathbf{z}(x) - (d, 0)) - Q_{\frac{2\pi i}{m}}(\mathbf{z}(y) - (d, 0)) \right|$ has positive lower bounds. So F_{13} and F_{14} are less singular than F_{12} , and $F_{14}(\varepsilon, f, g), F_{13}(\varepsilon, f, g) : (-\frac{1}{2}, \frac{1}{2}) \times V^r \rightarrow Y^k$ are both continuous. Using the Taylor's formula (2.1), we also rewrite F_{13} and F_{14} as

$$F_{13} = \varepsilon \mathcal{R}_{13}(\varepsilon, f, g), \quad (3.13)$$

and

$$\begin{aligned} F_{14} &= - \sum_{i=1}^{m-1} \frac{(1 - \cos(\frac{2\pi i}{m})) \sin(x)}{((-1 + \cos(\frac{2\pi i}{m}))^2 + \sin^2(\frac{2\pi i}{m})) d} + \varepsilon \mathcal{R}_{14}(\varepsilon, f, g) \\ &= - \frac{m-1}{2d} \sin(x) + \varepsilon \mathcal{R}_{14}(\varepsilon, f, g) \end{aligned} \quad (3.14)$$

with both $\mathcal{R}_{13}, \mathcal{R}_{14}$ being good terms.

The continuity for $F_2(\varepsilon, \Omega, f, g) : (-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R} \times V^r \rightarrow X^k$ can be verified in a similar way. Notice that projection operator $I - \mathcal{P}$ (defined in (1.10)) eliminates all constant terms in \tilde{F}_2 , and hence there is no singularity in $F_2 = (I - \mathcal{P})\tilde{F}_2$. According to (2.1), the four terms in F_2 can be rewritten into the following form respectively

$$F_{21} = \Omega(d \cos x + \mathcal{R}_{21}(\varepsilon, f, g)), \quad (3.15)$$

$$F_{22} = f(x) - \int \frac{f(x) - f(y)}{4 \sin^2(\frac{x-y}{2})} dy - \frac{1}{2} g(x) + \varepsilon \mathcal{R}_{22}(\varepsilon, f, g), \quad (3.16)$$

$$F_{23} = \varepsilon \mathcal{R}_{23}(\varepsilon, f, g), \quad (3.17)$$

$$F_{24} = - \frac{m-1}{2d} \cos(x) + \varepsilon \mathcal{R}_{24}(\varepsilon, f, g), \quad (3.18)$$

where \mathcal{R}_{2i} ($1 \leq i \leq 4$) are all continuous from $(-\frac{1}{2}, \frac{1}{2}) \times V^r$ to X^k .

Summing up the above, we complete the proof of this lemma. \square

For $(\varepsilon, \Omega, f, g) \in \left(-\frac{1}{2}, \frac{1}{2}\right) \times \mathbb{R} \times V^r$ and $(h_1, h_2) \in X^{k+1} \times X^k$, let

$$\partial_f F_i(\varepsilon, \Omega, f, g)h_1 := \lim_{t \rightarrow 0} \frac{1}{t} (F_i(\varepsilon, \Omega, f + th_1, g) - F_i(\varepsilon, \Omega, f, g))$$

and

$$\partial_g F_i(\varepsilon, \Omega, f, g)h_2 := \lim_{t \rightarrow 0} \frac{1}{t} (F_i(\varepsilon, \Omega, f, g + th_2) - F_i(\varepsilon, \Omega, f, g))$$

be the Gateaux derivatives of $F_i(\varepsilon, \Omega, f, g)$ with $i = 1, 2$. Then we have following lemma:

Lemma 3.2. *For each $(\varepsilon, \Omega, f, g) \in \left(-\frac{1}{2}, \frac{1}{2}\right) \times \mathbb{R} \times V^r$ and $(h_1, h_2) \in X^{k+1} \times X^k$,*

$$\partial_f F_1(\varepsilon, \Omega, f, g)h_1 : X^{k+1} \rightarrow Y^k, \quad \partial_g F_1(\varepsilon, \Omega, f, g)h_2 : X^k \rightarrow Y^k$$

and

$$\partial_f F_2(\varepsilon, \Omega, f, g)h_1 : X^{k+1} \rightarrow X^k, \quad \partial_g F_2(\varepsilon, \Omega, f, g)h_2 : X^k \rightarrow X^k$$

are continuous.

Proof. We only verify that $\partial_f F_1(\varepsilon, \Omega, f, g)h_1 : X^{k+1} \rightarrow Y^k$ is continuous, since the continuity of other three Gateaux derivatives can be shown in the same way.

According to Lemma 3.1, it is obvious that

$$\partial_f F_{11}(\varepsilon, \Omega, f)h = \Omega \varepsilon |\varepsilon| \partial_f \mathcal{R}_1(\varepsilon, f)h$$

is continuous, where $\mathcal{R}_{11}(\varepsilon, f)$ is given in (3.3).

Then we claim $\partial_f F_{12}(\varepsilon, f, g)h_1 = \partial_f F_{121} + \partial_f F_{122} + \partial_f F_{123} + \partial_f F_{124}$ is continuous (one can observe that $\partial_f F_{12}(\varepsilon, f, g)h_1$ is the main source of $\partial_f F_1(\varepsilon, f, g)h_1$ as $\varepsilon \rightarrow 0$), where

$$\begin{aligned} \partial_f F_{121} = & \int \frac{\varepsilon h_1(x) f'(x) (1 - \cos(x - y)) (1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\ & + \int \frac{(1 + \varepsilon f(x)) h_1'(x) (1 - \cos(x - y)) (1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\ & - \int \frac{2\varepsilon (1 + \varepsilon f(x)) f'(x) (1 - \cos(x - y)) (1 + \varepsilon g(y))}{(\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right))^2} \\ & \times \left(\varepsilon (f(x) - f(y)) (h_1(x) - h_1(y)) + 2((1 + \varepsilon f(x)) h_1(y) + h_1(x) (1 + \varepsilon f(y))) \sin^2\left(\frac{x-y}{2}\right) \right) dy, \end{aligned}$$

$$\begin{aligned}
\partial_f F_{122} &= \int \frac{\varepsilon h_1'(x)(f(x) - f(y)) \cos(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
&\quad + \int \frac{\varepsilon f'(x)(h_1(x) - h_1(y)) \cos(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
&\quad - \int \frac{2\varepsilon^2 f'(x)(f(x) - f(y)) \cos(x - y)(1 + \varepsilon g(y))}{\left(\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)\right)^2} \\
&\quad \times \left(\varepsilon(f(x) - f(y))(h_1(x) - h_1(y)) + 2((1 + \varepsilon f(x))h_1(y) + h_1(x)(1 + \varepsilon f(y))) \sin^2\left(\frac{x-y}{2}\right) \right) dy,
\end{aligned}$$

$$\begin{aligned}
\partial_f F_{123} &= - \int \frac{\varepsilon h_1(x)(f(x) - f(y)) \sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
&\quad - \int \frac{(1 + \varepsilon f(x))(h_1(x) - h_1(y)) \sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
&\quad + \int \frac{2\varepsilon(1 + \varepsilon f(x))(f(x) - f(y)) \sin(x - y)(1 + \varepsilon g(y))}{\left(\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)\right)^2} \\
&\quad \times \left(\varepsilon(f(x) - f(y))(h_1(x) - h_1(y)) + 2((1 + \varepsilon f(x))h_1(y) + h_1(x)(1 + \varepsilon f(y))) \sin^2\left(\frac{x-y}{2}\right) \right) dy,
\end{aligned}$$

$$\begin{aligned}
\partial_f F_{124} &= \int \frac{2(1 + \varepsilon f(x))h_1(x) \sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
&\quad - \int \frac{2(1 + \varepsilon f(x))^2 \sin(x - y)(1 + \varepsilon g(y))}{\left(\varepsilon^2 (f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)\right)^2} \\
&\quad \times \left(\varepsilon(f(x) - f(y))(h_1(x) - h_1(y)) + 2((1 + \varepsilon f(x))h_1(y) + h_1(x)(1 + \varepsilon f(y))) \sin^2\left(\frac{x-y}{2}\right) \right) dy.
\end{aligned}$$

First we are to show that these Gateaux derivatives are well-defined, that is

$$\lim_{t \rightarrow 0} \left\| \frac{F_{12i}(\varepsilon, f + th_1, g) - F_{12i}(\varepsilon, f, g)}{t} - \partial_f F_{12i} \right\|_{Y^k} = 0, \quad \text{for } i = 1, 2, 3, 4.$$

To simplify our proof, we will deal with the most singular case $i = 2$. Using the notations given in Lemma 3.1, we deduce that

$$\begin{aligned}
& \frac{F_{122}(\varepsilon, f + th_1, g) - F_{122}(\varepsilon, f, g)}{t} - \partial_f F_{122} \\
&= \varepsilon h_1'(x) \int (f(x) - f(y)) \cos(x - y) (1 + \varepsilon g(y)) \left(\frac{1}{D(f + th_1)} - \frac{1}{D(f)} \right) dy \\
&+ \frac{\varepsilon f'(x)}{t} \int (f(x) - f(y)) \cos(x - y) (1 + \varepsilon g(y)) \\
&\quad \times \left(\frac{1}{D(f + th_1)} - \frac{1}{D(f)} + t \frac{2\varepsilon^2 \Delta f \Delta h_1 + 4((1 + \varepsilon \tilde{f})\varepsilon h_1 + \varepsilon \tilde{h}_1(1 + \varepsilon f)) \sin^2(\frac{x-y}{2})}{D(f)^2} \right) dy \\
&+ \varepsilon f'(x) \int (h_1(x) - h_1(y)) \cos(x - y) (1 + \varepsilon g(y)) \left(\frac{1}{D(f + th_1)} - \frac{1}{D(f)} \right) dy \\
&+ t\varepsilon h_1'(x) \int \frac{(h_1(x) - h_1(y)) \cos(x - y) (1 + \varepsilon g(y))}{D(f + th_1)} dy \\
&= \partial_f^2 F_{1221} + \partial_f^2 F_{1222} + \partial_f^2 F_{1223} + \partial_f^2 F_{1224}.
\end{aligned}$$

From (3.11), for $\partial_f^2 F_{1221}$ and $\partial_f^2 F_{1223}$ we have

$$\|\partial^k \partial_f^2 F_{1221}(x \pm a\mathbf{i})\|_{L^2} \leq tC \|h_1\|_{x^{k+1}} \|f\|_{X^{k+1}} (1 + \|g\|_{X^k}) (1 + \|f\|_{X^{k+1}}) \|h_1\|_{X^k},$$

and

$$\|\partial^k \partial_f^2 F_{1223}(x \pm a\mathbf{i})\|_{L^2} \leq tC \|f\|_{x^{k+1}} \|f\|_{X^{k+1}} (1 + \|g\|_{X^k}) (1 + \|f\|_{X^{k+1}}) \|h_1\|_{X^k}.$$

For $\partial_f^2 F_{1224}$, the computation will be a little tedious. However, using mean value theorem, we are able to verify that

$$\frac{1}{D(f + th_1)} - \frac{1}{D(f)} + t \frac{2\varepsilon^2 \Delta f \Delta h_1 + 4(\varepsilon \tilde{R} h_1 + \varepsilon \tilde{h}_1 R) \sin^2(\frac{x-y}{2})}{D(f)^2} \leq \frac{t^2}{4 \sin^2(\frac{x-y}{2})} \xi(\varepsilon, f, h_1),$$

where $\|\xi(\varepsilon, f, h_1)(x \pm a\mathbf{i})\|_{L^\infty} \leq C(\varepsilon, \|f\|_{x^{k+1}}, \|h_1\|_{x^{k+1}}) < \infty$. As a result, it holds

$$\|\partial^k \partial_f^2 F_{1224}(x \pm a\mathbf{i})\|_{L^2} \leq tC \|f\|_{x^{k+1}} \|f\|_{X^{k+1}} (1 + \|g\|_{X^k}) \|\xi(\varepsilon, f, h_1)(x \pm a\mathbf{i})\|_{L^\infty}.$$

For the fourth term $\partial_f^2 F_{1224}$, it is obvious that

$$\|\partial^k \partial_f^2 F_{1224}(x \pm a\mathbf{i})\|_{L^2} \leq tC \|f\|_{x^{k+1}} \|h_1\|_{X^{k+1}} (1 + \|g\|_{X^k}),$$

and we have achieved our first target. In the next step, the continuity of $\partial_f F_{12}(\varepsilon, f, g)h_1$ should be shown. One can proceed as the proof Lemma 3.1 and hence we omit the proof.

To finish our discussion on F_1 , we have to verify the continuity of $\partial_f F_{13}(\varepsilon, f, g)h_1$ and $\partial_f F_{14}(\varepsilon, f, g)h_1$. For this purpose, a similar method can be applied as above. Moreover, due to the positive lower bound of denominator, the proof of this part is much simpler.

In the last, noting that F_1 and F_2 are almost linear dependent on g , it is easy to compute their Gateaux derivatives with respect to g , and we leave them to our readers. On the other hand, since $\partial_f F_{22}(\varepsilon, f, g)h_1 = \partial_f F_{221} + \partial_f F_{222} + \partial_f F_{223} + \partial_f F_{224}$ is a little bit complicated

and is the main source of $\partial_f F_2(\varepsilon, f, g)h_1$, we also write the four terms down here for readers' convenience.

$$\begin{aligned}
\partial_f F_{221} = & -\frac{2\varepsilon(1+\varepsilon g(x))f'(x)h'_1(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{(1+\varepsilon f(x))(1+\varepsilon f(y))(\cos(x-y)-1)(1+\varepsilon g(y))}{\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
& -\frac{2(1+\varepsilon g(x))(1+\varepsilon f(x))h_1(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{(1+\varepsilon f(x))(1+\varepsilon f(y))(\cos(x-y)-1)(1+\varepsilon g(y))}{\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
& +\frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{(1+\varepsilon f(x))h_1(y)(\cos(x-y)-1)(1+\varepsilon g(y))}{\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
& +\frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{h_1(x)(1+\varepsilon f(y))(\cos(x-y)-1)(1+\varepsilon g(y))}{\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
& -\frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{2(1+\varepsilon f(x))(1+\varepsilon f(y))(\cos(x-y)-1)(1+\varepsilon g(y))}{(\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right))^2} \\
& \times \left(\varepsilon(f(x)-f(y))(h_1(x)-h_1(y)) + 2((1+\varepsilon f(x))h_1(y) + h_1(x)(1+\varepsilon f(y))) \sin^2\left(\frac{x-y}{2}\right) \right) dy,
\end{aligned}$$

$$\begin{aligned}
\partial_f F_{222} = & \frac{2\varepsilon(1+\varepsilon g(x))f'(x)h'_1(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{\varepsilon^2 f'(x)(f(x)-f(y)) \sin(x-y)(1+\varepsilon g(y))}{(\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right))^2} dy \\
& +\frac{2(1+\varepsilon g(x))(1+\varepsilon f(x))h_1(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{\varepsilon^2 f'(x)(f(x)-f(y)) \sin(x-y)(1+\varepsilon g(y))}{(\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right))^2} dy \\
& -\frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{\varepsilon^2 h'_1(x)(f(x)-f(y)) \sin(x-y)(1+\varepsilon g(y))}{\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
& -\frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{\varepsilon^2 f'(x)(h_1(x)-h_1(y)) \sin(x-y)(1+\varepsilon g(y))}{\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right)} dy \\
& +\frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{2\varepsilon^2 f'(x)(f(x)-f(y)) \sin(x-y)(1+\varepsilon g(y))}{(\varepsilon^2 (f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y)) \sin^2\left(\frac{x-y}{2}\right))^2} \\
& \times \left(\varepsilon(f(x)-f(y))(h_1(x)-h_1(y)) + 2((1+\varepsilon f(x))h_1(y) + h_1(x)(1+\varepsilon f(y))) \sin^2\left(\frac{x-y}{2}\right) \right) dy,
\end{aligned}$$

$$\begin{aligned}
\partial_f F_{223} &= \frac{2\varepsilon(1+\varepsilon g(x))f'(x)h_1'(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{\varepsilon(1+\varepsilon f(x))(f(x)-f(y))(1+\varepsilon g(y))}{(\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2}))^2} dy \\
&+ \frac{2(1+\varepsilon g(x))(1+\varepsilon f(x))h_1(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{\varepsilon(1+\varepsilon f(x))(f(x)-f(y))(1+\varepsilon g(y))}{(\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2}))^2} dy \\
&- \frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{\varepsilon h_1(x)(f(x)-f(y))(1+\varepsilon g(y))}{\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2})} dy \\
&- \frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{(1+\varepsilon f(x))(h_1(x)-h_1(y))(1+\varepsilon g(y))}{\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2})} dy \\
&+ \frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{2\varepsilon(1+\varepsilon f(x))(f(x)-f(y))(1+\varepsilon g(y))}{(\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2}))^2} \\
&\times \left(\varepsilon(f(x)-f(y))(h_1(x)-h_1(y)) + 2((1+\varepsilon f(x))h_1(y) + h_1(x)(1+\varepsilon f(y)))\sin^2(\frac{x-y}{2}) \right) dy, \\
\partial_f F_{224} &= -\frac{2\varepsilon(1+\varepsilon g(x))f'(x)h_1'(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{\varepsilon f'(x)(1+\varepsilon f(x))\sin(x-y)(1+\varepsilon g(y))}{(\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2}))^2} dy \\
&- \frac{2(1+\varepsilon g(x))(1+\varepsilon f(x))h_1(x)}{(\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2)^2} \int \frac{\varepsilon f'(x)(1+\varepsilon f(x))\sin(x-y)(1+\varepsilon g(y))}{(\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2}))^2} dy \\
&+ \frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{h_1'(x)(1+\varepsilon f(x))\sin(x-y)(1+\varepsilon g(y))}{\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2})} dy \\
&+ \frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{\varepsilon f'(x)h_1(x)\sin(x-y)(1+\varepsilon g(y))}{\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2})} dy \\
&- \frac{(1+\varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1+\varepsilon f(x))^2} \int \frac{2\varepsilon f'(x)(1+\varepsilon f(x))\sin(x-y)(1+\varepsilon g(y))}{(\varepsilon^2(f(x)-f(y))^2 + 4(1+\varepsilon f(x))(1+\varepsilon f(y))\sin^2(\frac{x-y}{2}))^2} \\
&\times \left(\varepsilon(f(x)-f(y))(h_1(x)-h_1(y)) + 2((1+\varepsilon f(x))h_1(y) + h_1(x)(1+\varepsilon f(y)))\sin^2(\frac{x-y}{2}) \right) dy.
\end{aligned}$$

Concluding all the facts above, we have completed the proof. \square

According to Lemma 3.1 and Lemma 3.2, when $\varepsilon = 0$ and $f, g \equiv 0$, the Gateaux derivatives are

$$\begin{aligned}
\partial_f F_1(\varepsilon, \Omega, f, g)h_1 &= \frac{1}{2}h_1'(x), \\
\partial_g F_1(\varepsilon, \Omega, f, g)h_2 &= \int \frac{h_2(y)\sin(x-y)}{4\sin^2(\frac{x-y}{2})} dy, \\
\partial_f F_2(\varepsilon, \Omega, f, g)h_1 &= h_1(x) - \int \frac{h_1(x)-h_1(y)}{4\sin^2(\frac{x-y}{2})} dy, \\
\partial_g F_1(\varepsilon, \Omega, f, g)h_2 &= -\frac{1}{2}h_2(x).
\end{aligned}$$

Take $(h_1, h_2) \in X^{k+1} \times X^k$, where

$$h_1(x) = \sum_{j=1}^{\infty} a_j \cos(jx) \quad \text{and} \quad h_2(x) = \sum_{j=1}^{\infty} b_j \cos(jx). \quad (3.19)$$

If we denote $\mathbf{F} = (F_1, F_2)(\varepsilon, \Omega, f, g)$, the linearization of \mathbf{F} at $(0, \Omega, 0, 0)$ has the following Fourier series form

$$D\mathbf{F}(0, \Omega, 0, 0)(h_1, h_2) = \begin{pmatrix} \partial_f F_1(0, \Omega, 0, 0)h_1 + \partial_g F_1(0, \Omega, 0, 0)h_2 \\ \partial_f F_2(0, \Omega, 0, 0)h_1 + \partial_g F_2(0, \Omega, 0, 0)h_2 \end{pmatrix} = \sum_{j=1}^{\infty} \begin{pmatrix} \hat{a}_j \sin(jx) \\ \hat{b}_j \cos(jx) \end{pmatrix}, \quad (3.20)$$

where

$$\begin{pmatrix} \hat{a}_j \\ \hat{b}_j \end{pmatrix} = M_j \begin{pmatrix} a_j \\ b_j \end{pmatrix}$$

with M_j a 2×2 matrix and a_j, b_j given in (3.19). In the following lemma we will calculate the matrix M_j , and claim that the linearization of $\mathbf{F}(\varepsilon, \Omega, f)$ at $(0, \Omega, 0, 0)$ is an isomorphism from \mathcal{A}^k to \mathcal{B}^k . Notice that M_1 is not invertible. This is the reason why we give \mathcal{A}^k and \mathcal{B}^k a special restriction on the first Fourier coefficient.

Lemma 3.3. *One has*

$$M_j = \begin{pmatrix} -j/2 & 1/2 \\ (2-j)/2 & -1/2 \end{pmatrix}.$$

Moreover, $D\mathbf{F}(0, \Omega, 0, 0)(h_1, h_2)$ is an isomorphism from \mathcal{A}^k to \mathcal{B}^k .

Proof. We first show

$$\int \frac{\cos(jx) \sin(x-y)}{4 \sin^2\left(\frac{x-y}{2}\right)} dy = \frac{1}{2} \sin(jx) \quad (3.21)$$

holds for all $j \geq 1$ and $j \in \mathbb{N}^*$. It can be deduced from the identity

$$\int \frac{\cos(jy) \sin(x-y)}{4 \sin^2\left(\frac{x-y}{2}\right)} dy = \frac{1}{2} \int \cos(jy) \cot\left(\frac{x-y}{2}\right) dy = \frac{1}{2} \mathcal{H}(\cos(jx))(x),$$

where $\mathcal{H}(\cdot)$ is the Hilbert transform on torus. This fact implies that

$$\partial_f F_1(\varepsilon, \Omega, f, g)h_1 = - \sum_{j=1}^{\infty} \frac{j}{2} a_j \sin(jx),$$

and

$$\partial_g F_1(\varepsilon, \Omega, f, g)h_2 = \frac{1}{2} \sum_{j=1}^{\infty} b_j \sin(jx).$$

On the other hand, for all $j \geq 1$ and $j \in \mathbb{N}^*$ it holds

$$\int \frac{\cos(jx) - \cos(jy)}{4 \sin^2\left(\frac{x-y}{2}\right)} dy = \frac{1}{2} (-\Delta)^{\frac{1}{2}} \cos(jx) = \frac{j}{2} \cos(jx).$$

Then we obtain

$$\partial_f F_2(\varepsilon, \Omega, f, g)h_1 = \sum_{j=1}^{\infty} \frac{2-j}{2} a_j \sin(jx),$$

and

$$\partial_g F_2(\varepsilon, \Omega, f, g)h_2 = -\frac{1}{2} \sum_{j=1}^{\infty} b_j \sin(jx).$$

Thus we have verified the first part of this lemma.

Now we are going to prove $D\mathbf{F}(0, \Omega, 0, 0)(h_1, h_2)$ is an isomorphism from \mathcal{A}^k to \mathcal{B}^k . Recall the definition of \mathcal{A}^k and \mathcal{B}^k given in the beginning of this section. From Lemma (3.1) and $M_1 = \begin{pmatrix} -1/2 & 1/2 \\ 1/2 & -1/2 \end{pmatrix}$, it is obvious that $D\mathbf{F}$ maps \mathcal{A}^k into \mathcal{B}^k . Hence only the invertibility need to be considered.

For $j \geq 2$, $\det(M_j) = \frac{j-1}{2}$ which implies M_j is invertible at this time. We can compute the inverse of M_j as

$$M_j^{-1} = \begin{pmatrix} -1/(j-1) & -1/(j-1) \\ (j-2)/(j-1) & -j/(j-1) \end{pmatrix}, \quad \forall j \geq 2. \quad (3.22)$$

Thus for any $(u, v) \in \mathcal{B}^k$ with the formulation

$$u = \sum_{j=1}^{\infty} p_j \sin(jx) \quad \text{and} \quad g = -p_1 \cos(x) + \sum_{j=1}^{\infty} q_j \cos(jx),$$

we can write $D\mathbf{F}^{-1}(0, \Omega, 0, 0)(u, v)$ as

$$D\mathbf{F}^{-1}(0, \Omega, 0, 0)(u, v) = \begin{pmatrix} -p_1 \cos(jx) \\ p_1 \cos(jx) \end{pmatrix} + \sum_{j=2}^{\infty} \begin{pmatrix} \tilde{p}_j \cos(jx) \\ \tilde{q}_j \cos(jx) \end{pmatrix},$$

where

$$\begin{pmatrix} \tilde{p}_j \\ \tilde{q}_j \end{pmatrix} = M_j^{-1} \begin{pmatrix} p_j \\ q_j \end{pmatrix}, \quad \forall j \geq 2.$$

From (3.22), we see $\tilde{p}_j \sim -j^{-1}(p_j + q_j)$ and $\tilde{q}_j \sim p_j - q_j$ as $j \rightarrow +\infty$. Hence $D\mathbf{F}^{-1}(0, \Omega, 0, 0)(u, v)$ does belong to \mathcal{A}^k and the proof is finished. \square

One can easily verify that for

$$\Omega^* := \frac{m-1}{2d^2}, \quad (3.23)$$

the four-tuple $(0, \Omega^*, 0, 0)$ is a trivial solution to $\mathbf{F}(\varepsilon, \Omega, f, g) = (F_1, F_2)(\varepsilon, \Omega, f, g) = 0$, which corresponds to co-rotating m -fold point vortex solution. To apply the implicit function theorem and obtain concentrative vortex sheet solutions, we need to choose suitable $\Omega = (\varepsilon, f, g)$ such that $\mathbf{F}(\varepsilon, \Omega, f, g)$ maps $V^r \subset \mathcal{A}^k$ into \mathcal{B}^k . This is achieved in the following lemma.

Lemma 3.4. *Let Ω^* be given by (3.23). Then there exists some continuous function $\Omega_\varepsilon(\varepsilon, f, g) : (-1/2, 1/2) \times V^r \rightarrow \mathbb{R}$, such that if*

$$\Omega(\varepsilon, f, g) := \Omega^* + \varepsilon \cdot \Omega_\varepsilon(\varepsilon, f, g),$$

then $\mathbf{F}^(\varepsilon, f, g) := \mathbf{F}(\varepsilon, \Omega(\varepsilon, f, g), f, g)$ maps $(-1/2, 1/2) \times V^r$ into \mathcal{B}^k . Moreover, for any $(\varepsilon, f, g) \in (-1/2, 1/2) \times V^r$, $\partial_f \Omega_\varepsilon(\varepsilon, f, g)(\cdot) : X^{k+1} \rightarrow \mathbb{R}$ and $\partial_g \Omega_\varepsilon(\varepsilon, f, g)(\cdot) : X^k \rightarrow \mathbb{R}$ are both continuous.*

Proof. According to the definition of \mathcal{B}^k , it is sufficient to find $\Omega = \Omega(\varepsilon, f, g)$ such that

$$\int F_1(\varepsilon, \Omega, f, g) \sin(x) dx = - \int F_2(\varepsilon, \Omega, f, g) \cos(x) dx \quad (3.24)$$

for ε sufficiently small and $(f, g) \in V^r$. Let

$$f = \sum_{j=1}^{\infty} a_j \cos(jx) \quad \text{and} \quad g = -a_1 \cos(x) + \sum_{j=1}^{\infty} b_j \cos(jx).$$

Then by Lemma 3.1, Lemma 3.3 and (3.23), identity (3.24) is equivalent to

$$\begin{aligned} \Omega d - a_1 + \int \sum_{i=1}^4 \varepsilon \mathcal{R}_{1i}(\varepsilon, f, g) \sin(x) dx - \Omega^* d \\ = -(\Omega d + a_1 - \int \sum_{i=1}^4 \varepsilon \mathcal{R}_{2i}(\varepsilon, f, g) \cos(x) dx - \Omega^* d). \end{aligned}$$

So we deduce that

$$\Omega(\varepsilon, f, g) = \Omega^* - \frac{\varepsilon}{2d} \int \sum_{i=1}^4 \mathcal{R}_{1i}(\varepsilon, f, g) \sin(x) dx + \frac{\varepsilon}{2d} \int \sum_{i=1}^4 \mathcal{R}_{2i}(\varepsilon, f, g) \cos(x) dx.$$

Let

$$\Omega_\varepsilon(\varepsilon, f, g) := -\frac{1}{2d} \int \sum_{i=1}^4 \mathcal{R}_{1i}(\varepsilon, f, g) \sin(x) dx + \frac{1}{2d} \int \sum_{i=1}^4 \mathcal{R}_{2i}(\varepsilon, f, g) \cos(x) dx.$$

From the discussion in Lemma 3.2, it is easy to see that $\partial_f \Omega_\varepsilon(\varepsilon, f, g)(\cdot) : X^{k+1} \rightarrow \mathbb{R}$ and $\partial_g \Omega_\varepsilon(\varepsilon, f, g)(\cdot) : X^k \rightarrow \mathbb{R}$ are continuous. Thus the proof is therefore completed. \square

Having made all the preparations in the previous Lemmas, we are now able to prove Theorem 1.1.

Proof of Theorem 1.1: We first show that $D\mathbf{F}^*(\varepsilon, f, g)(\cdot, \cdot) : \mathcal{A}^k \rightarrow \mathcal{B}^k$ is an isomorphism. By the chain rule, we have for $i = 1, 2$

$$\partial_f F_i^*(0, 0, 0)h_1 = \partial_\Omega F_i(0, \Omega^*, 0, 0)\partial_f \Omega(0, 0, 0)h_1 + \partial_f F_1(0, \Omega^*, 0, 0)h_1, \quad \forall h_1 \in X^{k+1},$$

and

$$\partial_g F_i^*(0, 0, 0)h_2 = \partial_\Omega F_i(0, \Omega^*, 0, 0)\partial_g \Omega(0, 0, 0)h_2 + \partial_g F_1(0, \Omega^*, 0, 0)h_2, \quad \forall h_2 \in X^k.$$

On the other hand, using the formulation of angular velocity Ω in Lemma 3.4, we deduce that $\partial_f \Omega(0, 0, 0)h_1 = \partial_g \Omega(0, 0, 0)h_2 \equiv 0$ for any $h_1 \in X^{k+1}$ and $h_2 \in X^k$. Hence for $i = 1, 2$, it should be

$$\partial_f F_i^*(0, 0, 0)(\cdot) = \partial_f F_i(0, \Omega^*, 0, 0)(\cdot) \quad \text{and} \quad \partial_g F_i^*(0, 0, 0)(\cdot) = \partial_g F_i(0, \Omega^*, 0, 0)(\cdot),$$

and we have achieved our target by Lemma 3.3.

According to Lemma 3.1 – Lemma 3.4, we can apply implicit function theorem to derive that there exists $\varepsilon_0 > 0$ such that

$$\{(\varepsilon, f, g) \in [-\varepsilon_0, \varepsilon_0] \times V^r : \mathbf{F}^*(\varepsilon, f, g) = 0\}$$

is parameterized by one-dimensional curve $\varepsilon \in [-\varepsilon_0, \varepsilon_0] \rightarrow (\varepsilon, f_\varepsilon, g_\varepsilon)$. Moreover, it is not difficult to verify that for $\varepsilon \in [-\varepsilon_0, \varepsilon_0]$,

$$-\frac{1}{2d} \int \sum_{i=1}^4 \mathcal{R}_{1i}(\varepsilon, f_\varepsilon, g_\varepsilon) \sin(jx) dx + \frac{1}{2d} \int \sum_{i=1}^4 \mathcal{R}_{2i}(\varepsilon, f_\varepsilon, g_\varepsilon) \cos(jx) dx \neq 0, \quad \exists j \in \mathbb{N}^*.$$

So we have actually obtained a family of nontrivial vortex sheet solutions.

To show the convexity of the area surrounded by the closed curve corresponding to the vortex sheet, we use direct computation. Indeed,

$$\varepsilon \kappa(x) = \frac{R(x)^2 + 2R'(x)^2 - R(x)R''(x)}{(R(x)^2 + R'(x)^2)^{\frac{3}{2}}} = \frac{1 + O(\varepsilon)}{1 + O(\varepsilon)} > 0,$$

for ε small and each $x \in [0, 2\pi)$. Hence we claim that every component of vorticity is supported on a closed curve with convex interior. Since \mathcal{A}^k and \mathcal{B}^k are analytic spaces, this curve is also of C^∞ .

Let $\bar{f}(x) = f(-x)$ and $\bar{g}(x) = g(-x)$. The last thing to be verified is that if (ε, f, g) is a solution to $\mathbf{F}^*(\varepsilon, f, g) = 0$, then $(-\varepsilon, \bar{f}, \bar{g})$ is also a solution. By replacing y with $-y$ in the integral representation of $\mathbf{F}(\varepsilon, \Omega, f, g)$ and using the fact that f, g are both even functions, we derive $\Omega_\varepsilon(\varepsilon, f) = \Omega_\varepsilon(-\varepsilon, \bar{f}, \bar{g})$. Then we can insert this relationship into $\mathbf{F}(\varepsilon, \Omega, f, g)$ and obtain $\mathbf{F}^*(-\varepsilon, \bar{f}, \bar{g}) = 0$ by a similar substitution of variables as before.

Thus we have finished the proof of Theorem 1.1. \square

4. EXISTENCE OF TRAVELING VORTEX SHEETS

As before, let $\varepsilon \in (-\frac{1}{2}, \frac{1}{2})$, $R(x) = 1 + \varepsilon f(x)$, $\gamma(x) = 1 + \varepsilon g(x)$ with $x \in [0, 2\pi)$, and f, g be two C^1 functions. By (1.15), we need to find the zero points of following two functionals

$$G_1(\varepsilon, W, f, g) = G_{11} + G_{12} + G_{13} + G_{14},$$

where

$$G_{11} = -W ((1 + \varepsilon f(x)) \sin(x) - \varepsilon f'(x) \cos(x)),$$

$$\begin{aligned}
G_{12} &= \frac{1}{\varepsilon} \int \frac{\varepsilon(1 + \varepsilon f(x))f'(x)(1 - \cos(x - y))(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy \\
&+ \frac{1}{\varepsilon} \int \frac{\varepsilon^2 f'(x)(f(x) - f(y))\cos(x - y)(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy \\
&- \frac{1}{\varepsilon} \int \frac{\varepsilon(1 + \varepsilon f(x))(f(x) - f(y))\sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy \\
&+ \frac{1}{\varepsilon} \int \frac{(1 + \varepsilon f(x))^2 \sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy,
\end{aligned}$$

$$\begin{aligned}
G_{13} &= - \int \frac{\varepsilon(1 + \varepsilon f(x))f'(y)(1 + \varepsilon g(y))}{|(\varepsilon R(x)\cos(x) + \varepsilon R(y)\cos(y) - 2d)^2 + (\varepsilon R(x)\sin(x) + \varepsilon R(y)\sin(y))^2|^2} dy \\
&- \int \frac{\varepsilon f'(x)(1 + \varepsilon f(y))\cos(x - y)(1 + \varepsilon g(y))}{|(\varepsilon R(x)\cos(x) + \varepsilon R(y)\cos(y) - 2d)^2 + (\varepsilon R(x)\sin(x) + \varepsilon R(y)\sin(y))^2|^2} dy \\
&+ \int \frac{(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin(x - y)(1 + \varepsilon g(y))}{|(\varepsilon R(x)\cos(x) + \varepsilon R(y)\cos(y) - 2d)^2 + (\varepsilon R(x)\sin(x) + \varepsilon R(y)\sin(y))^2|^2} dy,
\end{aligned}$$

$$\begin{aligned}
G_{14} &= - \int \frac{2d(1 + \varepsilon f(x))\sin(x)(1 + \varepsilon g(y))}{|(\varepsilon R(x)\cos(x) + \varepsilon R(y)\cos(y) - 2d)^2 + (\varepsilon R(x)\sin(x) + \varepsilon R(y)\sin(y))^2|^2} dy \\
&+ \int \frac{2\varepsilon d f'(x)\cos(x)(1 + \varepsilon g(y))}{|(\varepsilon R(x)\cos(x) + \varepsilon R(y)\cos(y) - 2d)^2 + (\varepsilon R(x)\sin(x) + \varepsilon R(y)\sin(y))^2|^2} dy,
\end{aligned}$$

and

$$G_2(\varepsilon, W, f, g) = (I - \mathcal{P})\tilde{G}_2 = (I - \mathcal{P})(\tilde{G}_{21} + \tilde{G}_{22} + \tilde{G}_{23} + \tilde{G}_{24}),$$

where

$$\tilde{G}_{21} = -\frac{W(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \cdot ((1 + \varepsilon f(x))\cos(x) + \varepsilon f'(x)\sin(x)),$$

$$\begin{aligned}
\tilde{G}_{22} &= \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{(1 + \varepsilon f(x))(1 + \varepsilon f(y))(\cos(x - y) - 1)(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy \\
&- \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon^2 f'(x)(f(x) - f(y))\sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy \\
&- \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon(1 + \varepsilon f(x))(f(x) - f(y))(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy \\
&+ \frac{1}{\varepsilon} \cdot \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon f'(x)(1 + \varepsilon f(x))\sin(x - y)(1 + \varepsilon g(y))}{\varepsilon^2(f(x) - f(y))^2 + 4(1 + \varepsilon f(x))(1 + \varepsilon f(y))\sin^2\left(\frac{x-y}{2}\right)} dy,
\end{aligned}$$

$$\begin{aligned}
\tilde{G}_{23} &= \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{(1 + \varepsilon f(x))(1 + \varepsilon f(y)) \cos(x - y)(1 + \varepsilon g(y)) dy}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \\
&+ \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{\varepsilon f'(x)(1 + \varepsilon f(y)) \sin(x - y)(1 + \varepsilon g(y)) dy}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \\
&+ \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{(1 + \varepsilon f(x))^2(1 + \varepsilon g(y)) dy}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2}, \\
\tilde{G}_{24} &= \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{2d(1 + \varepsilon f(x)) \cos(x)(1 + \varepsilon g(y)) dy}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2} \\
&+ \frac{(1 + \varepsilon g(x))}{\varepsilon^2 f'(x)^2 + (1 + \varepsilon f(x))^2} \int \frac{2\varepsilon d f'(x) \sin(x)(1 + \varepsilon g(y)) dy}{|(\varepsilon R(x) \cos(x) + \varepsilon R(y) \cos(y) - 2d)^2 + (\varepsilon R(x) \sin(x) + \varepsilon R(y) \sin(y))^2|^2}.
\end{aligned}$$

Proof of Theorem 1.2: Comparing G_1, G_2 with F_1, F_2 for co-rotating sheet solutions, we observe that G_{12}, \tilde{G}_{22} equal F_{12}, \tilde{F}_{22} given in (3.2) and (3.6) separately. Moreover, for $i = 3, 4$, G_{1i}, \tilde{G}_{2i} equal $-F_{1i}, -\tilde{F}_{2i}$ in the case $m = 2$. Hence by a same procedure as seeking for the regularity of F_1, F_2 , we can prove the continuity of $\mathbf{G} = (G_1, G_2)$ and their Gateaux derivatives $D\mathbf{G}$ near $(0, W, 0, 0)$. From Lemma 3.3, we can obtain the exact matrix form for $D\mathbf{G}(0, W, 0, 0)$, which is an isomorphism from \mathcal{A}^k to \mathcal{B}^k .

It remains to choose W appropriately such that for $(\varepsilon, W, f, g) \in [-\varepsilon_0, \varepsilon_0] \times \mathbb{R} \times V^r$ with $\varepsilon_0, r > 0$ small enough, the range of $\mathbf{G} = (G_1, G_2)$ is in \mathcal{B}^k . This can be achieved by letting

$$W = W^* + \varepsilon \cdot W_\varepsilon(\varepsilon, f, g)$$

with $W^* = \frac{1}{2d}$ and some regular perturbation $W_\varepsilon(\varepsilon, f, g)$. Then we can apply implicit function theorem on $\mathbf{G}^*(\varepsilon, f, g)$ at $(0, 0, 0)$ to derive the existence and properties for a local curve of nontrivial traveling vortex sheet solutions. \square

REFERENCES

- [1] A. L. Bertozzi and P. Constantin, Global regularity for vortex patches, *Comm. Math. Phys.*, 152(1)(1993), 19–28. <https://doi.org/10.1007/BF02097055>
- [2] G. Birkhoff, Hydrodynamics: A study in logic, fact and similitude. Revised ed. Princeton Univ. Press, Princeton, N.J. 1960 xi+184 pp.
- [3] G. Birkhoff, Helmholtz and Taylor instability, *Proc. Symp. Appl. Math*, 13(1962), 55–76.
- [4] G. Birkhoff and J. Fisher, Do vortex sheets roll up? *Rendiconti del Circolo matematico di Palermo*, 8(1959), 77–90. <https://doi.org/10.1007/BF02843773>
- [5] R.E. Caffisch and O.F. Orellana, Singular solutions and ill-posedness for the evolution of vortex sheets, *SIAM J. Math. Anal.*, 20(1989)(2), 293–307. <https://doi.org/10.1137/0520020>
- [6] D. Cao, Z. Liu and J. Wei, Regularization of point vortices for the Euler equation in dimension two, *Arch. Rational Mech. Anal.*, 212 (2014), 179–217. <https://doi.org/10.1007/s00205-013-0692-y>
- [7] D. Cao, S. Peng and S. Yan, Planar vortex patch problem in incompressible steady flow, *Adv. Math.*, 270 (2015), 263–301. <https://doi.org/10.1016/j.aim.2014.09.027>

- [8] A. Castro, D. Córdoba and J. Gómez-Serrano, Uniformly rotating smooth solutions for the incompressible 2D Euler equations, *Arch. Ration. Mech. Anal.*, 231(2019), no. 2, 719–785. <https://doi.org/10.1007/s00205-018-1288-3>
- [9] J.-Y. Chemin, *Fluides Parfaits Incompressibles*, Astérisque 230, 1995 (Perfect Incompressible Fluids translated by I. Gallagher and D. Iftimie, Oxford Lecture Series in Mathematics and Its Applications, Vol. 14, Clarendon Press-Oxford University Press, New York, 1998).
- [10] F. de la Hoz, T. Hmidi, J. Mateu and J. Verdera, Doubly connected V-states for the planar Euler equations, *SIAM J. Math. Anal.*, 48(3)(2016), 1892–1928. <https://doi.org/10.1137/140992801>
- [11] J.-M., Delort, Existence de nappes de tourbillon en dimension deux, *J. Amer. Math. Soc.*, 4(1991), no. 3, 553–586. <https://doi.org/10.1090/S0894-0347-1991-1102579-6>
- [12] J. Duchon and R. Robert, Global vortex sheet solutions of Euler equations in the plane, *J. Differential Equations*, 73(1988), no. 2, 215–224. [https://doi.org/10.1016/0022-0396\(88\)90105-2](https://doi.org/10.1016/0022-0396(88)90105-2)
- [13] J. Gómez-Serrano, J. Park, J. Shi and Y. Yao, Remarks on stationary and uniformly-rotating vortex sheets: Flexibility results, *Comm. Math. Phys.*, (2021). <https://doi.org/10.1007/s00220-021-04146-3>
- [14] J. Gómez-Serrano, J. Park, J. Shi and Y. Yao, Remarks on stationary and uniformly-rotating vortex sheets: Rigidity result, to appear in *Phil. Trans. R. Soc. A.*, (2021).
- [15] T. Hmidi, J. Mateu and J. Verdera, Boundary regularity of rotating vortex patches, *Arch. Ration. Mech. Anal.*, 209(1)(2013), 171–208. <https://doi.org/10.1007/s00205-013-0618-8>
- [16] T. Hmidi and J. Mateu, Bifurcation of rotating patches from Kirchhoff vortices, *Discret. Contin. Dyn. Syst.*, 36(10)(2016), 5401–5422. [https://doi.org/10.1016/0021-9991\(84\)90051-2](https://doi.org/10.1016/0021-9991(84)90051-2)
- [17] T. Hmidi and J. Mateu, Existence of corotating and counter-rotating vortex pairs for active scalar equations, *Comm. Math. Phys.*, 350(2017), 699–747. <https://doi.org/10.1007/s00220-016-2784-7>
- [18] M.C. Lopes Filho, H.J. Nussenzweig Lopes and Z. Xin. Existence of vortex sheets with reflection symmetry in two space dimensions, *Arch. Ration. Mech. Anal.*, 158(2007)(3), 235–257. <https://doi.org/10.1007/s002050100145>
- [19] R. Krasny, A study of singularity formation in a vortex sheet by the point-vortex approximation, *Journal of Fluid Mechanics*, 167(1986), 65–93. <https://doi.org/10.1017/S0022112086002732>
- [20] A. Majda, Remarks on weak solutions for vortex sheets with a distinguished sign, *Indiana Univ. Math. J.*, 42(1993)(3), 921–939.
- [21] A. J. Majda and A. L. Bertozzi, Vorticity and incompressible flow, *Cambridge Texts in Applied Mathematics, Vol. 27.*, Cambridge University Press, 2002.
- [22] D. I. Meiron, G. R. Baker and S.A. Orszag, Analytic structure of vortex sheet dynamics. Part 1. Kelvin-Helmholtz instability, *Journal of Fluid Mechanics*, 114(1982), 283–298. <https://doi.org/10.1017/S0022112082000159>
- [23] D.W. Moore, The spontaneous appearance of a singularity in the shape of an evolving vortex sheet, *Proc. Roy. Soc. London Ser. A*, 365(1979)(1720), 105–119. <https://doi.org/10.1098/rspa.1979.0009>
- [24] B. Protas and T. Sakajo, Rotating equilibria of vortex sheets, *Phys. D*, 403(2020), 132268, 9. <https://doi.org/10.1016/j.physd.2019.132268>
- [25] P. G. Saffman, *Vortex dynamics*. Cambridge Monographs on Mechanics and Applied Mathematics, Cambridge University Press, New York, 1992.
- [26] C. Sulem and P.-L. Sulem, Finite time analyticity for the two- and three-dimensional Rayleigh-Taylor instability, *Trans. Amer. Math. Soc.*, 287(1985)(1), 127–160. <https://doi.org/10.1090/S0002-9947-1985-0766210-5>
- [27] B. Turkington, Corotating steady vortex flows with N -fold symmetry, *Nonlinear Anal.*, 9(1985), no. 4, 351–369. [https://doi.org/10.1016/0362-546X\(85\)90059-8](https://doi.org/10.1016/0362-546X(85)90059-8)
- [28] S. Wu, Mathematical analysis of vortex sheets, *Comm. Pure Appl. Math.*, 59(2006)(8), 1065–1206. <https://doi.org/10.1002/cpa.20110>
- [29] V.I. Yudovich, Non-stationary flows of an ideal incompressible fluid, *Zhurnal Vych Matematika*, 3(1963), 1032–1106.

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND
UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA

Email address: dmcao@amt.ac.cn

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND
UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA

Email address: qinguolin18@mailsucas.edu.cn

INSTITUTE OF APPLIED MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, AND
UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P.R. CHINA

Email address: zouchangjun17@mailsucas.ac.cn