

The local characterizations of the singularity formation for the MHD equations

Wenke Tan* Fan Wu†

Key Laboratory of Computing and Stochastic Mathematics (Ministry of Education),
School of Mathematics and Statistics, Hunan Normal University,
Changsha, Hunan 410081, China

Abstract: This paper characterizes the possible blow-up of solutions for the 3D magneto-hydrodynamics (MHD for short) equations. We first establish some ϵ -regularity criteria in $L^{q,\infty}$ spaces for suitable weak solutions, and then together with an embedding theorem from $L^{p,\infty}$ space into a Morrey type space to characterize the local behaviors of solutions near a potential singular point. More precisely, we show that if $z_0 = (t_0, x_0)$ is a singular point, then for any $r > 0$ it holds that

$$\limsup_{t \rightarrow t_0^-} \left(\|u(t, x) - u(t)_{x_0, r}\|_{L^{3,\infty}(B_r(x_0))} + \|b(t, x) - b(t)_{x_0, r}\|_{L^{3,\infty}(B_r(x_0))} \right) > \delta^*;$$

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu}} r^{\frac{2}{\nu} - \frac{3}{p}} \|(u, b)(t)\|_{L^{p,\infty}(B_r(x_0))} > \delta^* \text{ for } \frac{1}{\mu} + \frac{1}{\nu} = \frac{1}{2}, 2 \leq \nu \leq \frac{2p}{3}, 3 < p \leq \infty;$$

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu}} r^{\frac{2}{\nu} - \frac{3}{p} + 1} \|(\nabla u, \nabla b)(t)\|_{L^p(B_r(x_0))} > \delta^* \text{ for } \frac{1}{\mu} + \frac{1}{\nu} = \frac{1}{2}, \nu \in \begin{cases} [2, \infty], & p \geq 3 \\ [2, \frac{2p}{3-p}], & \frac{3}{2} \leq p < 3 \end{cases}$$

where δ^* is a positive constant independent on ν and p .

Mathematics Subject Classification (2010): 76D05, 35B65, 35Q30.

Keywords: MHD equations; Weak solutions; Singularities; Concentration rate

1 Introduction

This paper is concerned with the local behaviors near a possible singularity of suitable weak solutions to the incompressible MHD equations in three spatial dimensions with unit viscosity and zero external force

$$\begin{cases} \partial_t u + (u \cdot \nabla)u - \Delta u + \nabla P = (b \cdot \nabla)b, \\ \partial_t b + (u \cdot \nabla)b - \Delta b = (b \cdot \nabla)u, \\ \nabla \cdot u = \nabla \cdot b = 0, \\ u(x, 0) = u_0(x), b(x, 0) = b_0(x), \end{cases} \quad (1.1)$$

where u , P and b are quantities corresponding to the velocity of the fluid, its pressure and the magnetic field, and u_0, b_0 are the prescribed initial data satisfying the compatibility condition $\nabla \cdot u_0 = \nabla \cdot b_0 = 0$ in the distributional sense. Physically, equations (1.1) governs the dynamics of the velocity and magnetic

*tanwenkeybfq@163.com

†wufan0319@yeah.net

fields in electrically conducting fluids, such as plasmas, liquid metals, and salt water. For more details on the related background, we refer the reader to [3].

While many mathematicians have made a lot of important contributions to theoretical research of the incompressible 3D MHD equations (1.1), many questions are left open. It is well known that the global existence of weak solutions, local existence, and uniqueness of smooth solutions to equations (1.1) were established in [11, 30]. In the absence of magnetic field (i.e., $b = 0$), (1.1) reduces to the incompressible Navier-Stokes (NS) equations. The problem of the global regularity of solutions to the NS equations in three and higher space dimensions is a fundamental question in fluid dynamics and is still widely open. In a seminal paper [22], Leray shown that for $3 < p \leq \infty$, there exists c_p such that the conditions

$$\|u(t)\|_{L^p} < \frac{c_p}{(T-t)^{\frac{p-3}{2p}}}$$

imply the regularity of weak solutions on $[0, T]$. It is also possible to guarantee that a solution must remain smooth as long as there is control on some scale critical quantity. The Prodi-Serrin and Escauriaza-Seregin-Sverak regularity criterion [12, 29, 34] states that if

$$u \in L^q(0, T; L^p(\Omega)) \quad \text{with} \quad \frac{2}{q} + \frac{3}{p} \leq 1 \quad \text{and} \quad 3 \leq p \leq \infty, \quad (1.2)$$

then the solution is smooth and can be continued to a smooth solution for some time $T' > T$. Recently, the endpoint case $p = 3$, $q = +\infty$ was further improved by Tao [38], who proved a triply logarithmic lower bound on the rate of blowup of the L^3 norm. Somewhat more precisely, Tao showed that for an absolute constant $c > 0$, if $T < +\infty$,

$$\limsup_{t \rightarrow T} \frac{\|u(\cdot, t)\|_{L^3}}{\left(\log \log \log \frac{1}{T-t}\right)^c} = +\infty. \quad (1.3)$$

In [20], Kim-Kozono proved the local boundedness of a weak solution u under the assumption that $\|u\|_{L^{r,\infty}([0,T]; L^{s,\infty}(\mathbb{R}^3))}$ is sufficiently small for some (r, s) with $\frac{2}{r} + \frac{3}{s} = 1$ and $3 \leq s < \infty$. The limiting case of the regularity criteria obtained by Kim and Kozono was proved by He and Wang [13], i.e., any weak solution for the NS or MHD equations is regular under the assumption that $\|u\|_{L^{2,\infty}([0,T]; L^\infty(\mathbb{R}^3))}$ is sufficiently small. These results involving the NS equations were improved by Wang-Zhang [39], they showed that $\|u_3\|_{L^{r,\infty}([0,T]; L^{s,\infty}(\mathbb{R}^3))} \leq M$ and $\|u_h\|_{L^{r,\infty}([0,T]; L^{s,\infty}(\mathbb{R}^3))} \leq \epsilon_M$ with $\frac{2}{r} + \frac{3}{s} = 1$ and $3 < s \leq \infty$ imply the regularity of the suitable weak solution, where $u_h = (u_1, u_2)$, $u = (u_h, u_3)$ and ϵ_M is a small constant depending on M . More recently, Tan-Yin [36] extended previous results, they proved that u is regular in $[-1, 0] \times \mathbb{R}^3$ provided that u satisfies

$$\|u_3\|_{L^{2,\infty}([-1,0]; BMO(\mathbb{R}^3))} = M < \infty$$

and

$$\|u_h\|_{L^{2,\infty}([-1,0]; BMO(\mathbb{R}^3))} \leq \epsilon.$$

Another important step towards a better understanding of the Navier-Stokes equations is the partial regularity theory. Scheffer introduced the idea of suitable weak solutions and generalized energy inequality in a series of papers [31–33]. Scheffer's results were further generalized by Caffarelli, Kohn and Nirenberg [4], where a criterion of ϵ -regularity theorem was established and the nullity of one-dimensional parabolic Hausdorff measure of the singular set was provided. Lin [24] gave a greatly simplified new proof by an indirect argument. Later, Choe and Lewis [7] improved the parabolic Hausdorff dimension by a logarithmic factor.

For incompressible MHD equations, we would like to mention here the paper of He-Xin [14] that first extended the Serrin type criteria of NS equations to the MHD equations only in terms of velocity, these

results merit attention, especially, which indicates that the velocity field plays a more dominant role than the magnetic field on the regularity of solutions. For conditional regularity or blow-up criteria of NS and MHD equations, please refer to [5, 6, 10, 19, 23, 27, 41, 42] for a more comprehensive discussions. On the other hand, He and Xin [15] introduced the definition of suitable weak solutions and obtained partial regularity theorems of suitable weak solutions. Kang and Lee [18] found regularity criteria under the smallness of some scaling invariant quantities for velocity field without a smallness condition for the magnetic field. But they assumed uniform bounds of some scaling invariant quantities for magnetic field. And then, Wang-Zhang [40] presented a new interior regularity criterion for suitable weak solutions of the MHD equations that the scaling invariant quantity of velocity is involved only.

Investigation of singular weak Leray-Hopf solutions was first performed by Leray in [22]. In particular, he proved that if $[0, T)$ is the maximal existence interval of a smooth solution u for the 3D Navier-Stokes equations, then for any $3 < p \leq \infty$, there exists a constant c_p depending only on p such that

$$\|u(\cdot, t)\|_{L^p(\mathbb{R}^3)} \geq \frac{c_p}{(T-t)^{\frac{p-3}{2p}}}.$$

In general, except for the case of $p = 3$ (see [12, 21]), if the solution u satisfies

$$\|u(\cdot, t)\|_{L^p(\mathbb{R}^3)} \leq \frac{C}{(T-t)^{\frac{p-3}{2p}}}$$

with some constant $C > 0$, the regularity of u at time $t = T$ is still open. For the axi-symmetric Navier-Stokes equations, Chen-Strain-Yau-Tsai [8, 9] and Koh-Nadirashvili-Seregin-Sverák [21] made some important progresses, and established that the weak solution does not develop type-I singularities on time $t = T$, i.e.,

$$\|u(x, t)\|_{L^\infty(\mathbb{R}^3)} \leq \frac{C}{\sqrt{T-t}}.$$

Behaviour of the L^3 norm is more subtle. In a breakthrough paper, Escauriaza, Seregin and Sverák showed that if (\bar{x}, T^*) is a singular point then

$$\limsup_{t \rightarrow T^*} \|u(\cdot, t)\|_{L^3(B_\delta(\bar{x}))} = \infty \quad (1.4)$$

for any fixed $\delta > 0$. Later in [35], Seregin improved (1.4):

$$\lim_{t \rightarrow T^*} \|u(\cdot, t)\|_{L^3(\mathbb{R}^3)} = \infty. \quad (1.5)$$

Recently in [1], Albritton and Barker refined (1.4) and (1.5) to show that if Ω is a bounded domain with C^2 boundary one has

$$\lim_{t \rightarrow T^*} \|u(\cdot, t)\|_{L^3(B_\delta(\bar{x}) \cap \Omega)} = \infty \quad (1.6)$$

for any fixed $\delta > 0$.

We may now go back to the issue of the ‘‘concentration of solutions’’. Such phenomenon was investigated for other equations e.g. nonlinear Schrödinger in the wake of the pioneering work [26], also see [16, 17]. In [25], for the NS equations, an interesting concentration result is proven for a weak Leray-Hopf solution u which first blows up at $T^* > 0$. In particular, they showed that there exists $t_n \rightarrow T^*$ and $x_n \in \mathbb{R}^3$ such that

$$\|u(\cdot, t_n)\|_{L^m(B_{\sqrt{C(m)(T^*-t_n)}}(x_n))} \geq \frac{C(m)}{(T^* - t_n)^{\frac{1}{2}(1-\frac{3}{m})}}, \quad 3 \leq m \leq \infty. \quad (1.7)$$

Later, Maekawa-Miura-Prange in [28] improved (1.7) and showed that for every $t \in (0, T^*)$ (not just a sequence $t_n \rightarrow T^*$) there exists $x(t) \in \mathbb{R}^3$ such that

$$\|u(\cdot, t)\|_{L^p(B_{\sqrt{C(m)(T^*-t)}}(x(t)))} \geq \frac{C(m)}{(T^* - t)^{\frac{1}{2}(1 - \frac{3}{m})}} \quad 3 \leq m \leq \infty. \quad (1.8)$$

However, in (1.7) and (1.8) no information is provided on x_n and $x(t)$. It is natural to ask whether the concentration phenomenon occurs on balls $B(x, R)$ with $R = O(\sqrt{T^* - t})$ and with (x, T^*) being a singular point. Recently, Barker-Prange in [2] gave this problem an affirmative answer for the L^3 for Leray-Hopf solutions which first blow-up at time T^* and which satisfy the Type I bound:

$$\sup_{\bar{x} \in \mathbb{R}^3} \sup_{r \in (0, r_0)} \sup_{T^* - r^2 < t < T^*} \left(\int_{B_r(\bar{x})} |u(x, t)|^2 dx \right)^{\frac{1}{2}} \leq M \quad (1.9)$$

for a fixed radius $r_0 \in (0, \infty]$ and $M, T^* \in (0, \infty)$. Then it holds

$$\|u(\cdot, t)\|_{L^3(B_R(x))} \geq \gamma_{\text{univ}}, \quad R = O(\sqrt{T - t}).$$

In conclusion, the existed works characterized the singularity formation accurately by the critical L^3 norm. But, if we consider the local characterization of singularity by L^3 norm, there is an unpleasant problem. On the one hand, It is well-known that if $|u(T, x)| \leq \frac{c}{|x|}$ with small enough c then $(T, 0)$ can not be a singular point. On the other hand, it is clear that $\|u(T, \cdot)\|_{L^3(B_r)} = \infty$ for any $r > 0$. This means that one can not exclude such point from the singular set by using L^3 norm. Noticing that $\|u(T, \cdot)\|_{L^{3, \infty}(B_r)} = (\frac{4\pi}{3})^{\frac{1}{3}} c$, it is more natural to characterize the singularity formation for the Navier-Stokes equations by $L^{3, \infty}$ norm. Very recently, the first author in [37] established the concentration rate for the $L^{p, \infty}$ norm of u with $3 \leq p \leq \infty$ for the Navier-Stokes equations. Precisely, it showed that if $z_0 = (t_0, x_0)$ is a singular point, then for any $r > 0$ it holds that

$$\limsup_{t \rightarrow t_0^-} \|u(t, x) - u(t)_{x_0, r}\|_{L^{3, \infty}(B_r(x_0))} > \delta^* \quad (1.10)$$

or

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu}} r^{\frac{2}{\nu} - \frac{3}{p}} \|u(t)\|_{L^{p, \infty}(B_r(x_0))} > \delta^* \quad \text{for } \frac{1}{\mu} + \frac{1}{\nu} = \frac{1}{2}, \quad 2 \leq \nu \leq \frac{2p}{3}, \quad 3 < p \leq \infty, \quad (1.11)$$

where δ^* is a positive constant independent on p and ν .

Concerning the collective behavior of the MHD equations (1.1), to the best of our knowledge, there is no available result in this topic. In this paper, we shall consider the local characterization of singularity by the $L^{p, \infty}$ norm of (u, b) with $3 \leq p \leq \infty$ and the L^p norm of $(\nabla u, \nabla b)$ with $\frac{3}{2} \leq p \leq \infty$.

Before stating our main results, we first introduce the definition of suitable weak solutions and the notations of some scaled dimensionless quantities.

Definition 1.1 *We say that a triple (u, b, P) is a suitable weak solution of (1.1) in $\mathbb{R}^3 \times (0, T)$ if the following conditions hold:*

1. $u, b \in L^\infty(0, T; L^2(\mathbb{R}^3)) \cap L^2(0, T; H^1(\mathbb{R}^3))$;
2. (1.1)_{1,2} holds in the sense of distributions, that is,

$$\begin{aligned} \int_0^t (u, \partial_t \phi + \Delta \phi + (u \cdot \nabla) \phi) ds + (u_0, \phi(0)) &= \int_0^t ((b \cdot \nabla) \phi, b) ds, \\ \int_0^t (b, \partial_t \phi + \Delta \phi + (u \cdot \nabla) \phi) ds + (u_0, \phi(0)) &= \int_0^t ((b \cdot \nabla) \phi, u) ds, \end{aligned}$$

for all $\phi \in C_c^\infty([0, T] \times \mathbb{R}^3)$ with $\nabla \cdot \phi = 0$, where (\cdot, \cdot) is the scalar product in $L^2(\Omega)$;

3. (u, b) satisfies the energy inequality, that is,

$$\|u(t)\|_{L^2}^2 + \|b(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla u(s)\|_{L^2}^2 + \|\nabla b(s)\|_{L^2}^2 ds \leq \|u_0\|_{L^2}^2 + \|b_0\|_{L^2}^2, \quad 0 \leq t \leq T.$$

4. the pressure $P \in L_{loc}^{\frac{3}{2}}(\mathbb{R}^3 \times (0, T))$ and the following local energy inequality holds

$$\begin{aligned} & \int_{\mathbb{R}^3} (|u(x, t)|^2 + |b(x, t)|^2) \phi dx + 2 \int_0^t \int_{\mathbb{R}^3} (|\nabla u(x, t)|^2 + |\nabla b(x, t)|^2) \phi dx dt \\ & \leq \int_0^t \int_{\mathbb{R}^3} [(|u|^2 + |b|^2)(\Delta \phi + \partial_t \phi) + u \cdot \nabla \phi (|u|^2 + |b|^2 + 2P) - (b \cdot u)(b \cdot \nabla \phi)] dx dt, \end{aligned} \quad (1.12)$$

for any $\phi \in C_c^\infty((0, T] \times \mathbb{R}^3)$.

We say a point $z_0 = (t_0, x_0) \in (0, T] \times \mathbb{R}^3$ is a regular point of solution (u, b) to (1.1) if there exists a non-empty neighborhood $\mathcal{O}_{z_0} \subset (0, T] \times \mathbb{R}^3$ of z_0 such that $u, b \in L^\infty(\mathcal{O}_{z_0})$. The complement of the set of regular points will be called the singular set.

For the sake of completeness, we introduce some invariant quantities: Let (u, b, P) be a solution of the MHD equations (1.1), let

$$u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t), \quad b_\lambda(x, t) = \lambda b(\lambda x, \lambda^2 t), \quad P_\lambda(x, t) = \lambda^2 P(\lambda x, \lambda^2 t) \quad (1.13)$$

for any $\lambda > 0$, then the family $(u_\lambda, b_\lambda, P_\lambda)$ is also a solution of (1.1). Define

$$\begin{aligned} A(u, b, r, z) &= \sup_{t-r^2 \leq s < t} \frac{1}{r} \int_{B_r(x) \times \{s\}} (|u|^2 + |b|^2) dx, \\ B(u, b, r, z) &= \frac{1}{r} \iint_{Q_r(z)} (|\nabla u|^2 + |\nabla b|^2) dx dt, \\ C(u, b, r, z) &= \frac{1}{r^2} \iint_{Q_r(z)} (|u|^3 + |b|^3) dx dt, \\ D(P, r, z) &= \frac{1}{r^2} \iint_{Q_r(z)} |P|^{\frac{3}{2}} dx dt, \end{aligned}$$

where

$$\begin{aligned} B_r(x_0) &= \{x \in \mathbb{R}^3 : |x - x_0| < r\}, \quad B_r = B_r(0), \quad B = B_1; \\ Q_r(x_0) &= B_r(x_0) \times (t_0 - r^2, t_0), \quad Q_r = Q_r(0), \quad Q = Q_1. \end{aligned}$$

For simplicity, we denote

$$A(u, b, r) = A(u, b, r, 0), \quad B(u, b, r) = B(u, b, r, 0), \quad C(u, b, r) = C(u, b, r, 0), \quad D(P, r) = D(P, r, 0).$$

Next, we recall the following function spaces:

We use $L^q((0, T]; L^p(\mathbb{R}^3))$ to denote the space of measurable functions with the following norm

$$\|f\|_{L^q((0, T]; L^p(\mathbb{R}^3))} = \begin{cases} \left(\int_0^T \left(\int_{\mathbb{R}^3} |f(t, x)|^p dx \right)^{\frac{q}{p}} dt \right)^{\frac{1}{q}}, & 1 \leq q < \infty \\ \text{ess sup}_{t \in (0, T]} \|f(t, \cdot)\|_{L^p(\mathbb{R}^3)}, & q = \infty \end{cases}$$

The Lorentz space $L^{r,s}([0, T])$ is the space of measurable functions with the following norm:

$$\|f\|_{L^{r,s}([0, T])} = \begin{cases} \left(\int_0^\infty \sigma^{s-1} |\{x \in [0, T] : |f(x)| > \sigma\}|^{\frac{s}{r}} d\sigma \right)^{\frac{1}{s}}, & 1 \leq s < \infty \\ \sup_{\sigma > 0} \sigma |\{x \in [0, T] : |f(x)| > \sigma\}|^{\frac{1}{r}}, & s = \infty \end{cases}$$

Throughout this paper, $u_{x_0, \rho} \doteq \frac{1}{|B_\rho|} \int_{B_\rho(x_0)} u \, dx$ and C denotes an absolute positive number which can change from line to line.

With weak solutions and the relevant function spaces defined, we can now state the main results of this paper.

Theorem 1.1 *Let $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$ with $2 \leq p \leq \infty$. Assume (u, b, P) be a suitable weak solution to the MHD equations (1.1) on $Q_1(z_0)$, there exists a absolute constant $\delta > 0$ such that if*

$$\left\| \left(\sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t) - u_{x_0, \eta}|^p \, dx \right)^{\frac{1}{p}}, \sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t) - b_{x_0, \eta}|^p \, dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} \leq \delta \quad (1.14)$$

or

$$\left\| \left(\sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t)|^p \, dx \right)^{\frac{1}{p}}, \sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t)|^p \, dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} \leq \delta, \quad (1.15)$$

then z_0 is a regular point.

Let us now state the characterization for singularity formation in MHD equations (1.1).

Theorem 1.2 *Let (u, b, P) be a suitable weak solution of equations (1.1) in $Q_1(z_0)$. Assume z_0 be a singular point, then for any given $r \in (0, 1)$, it holds that*

$$\limsup_{t \rightarrow t_0^-} \left(\|u(t, x) - u(t)_{x_0, r}\|_{L^{3, \infty}(B_r(x_0))} + \|b(t, x) - b(t)_{x_0, r}\|_{L^{3, \infty}(B_r(x_0))} \right) > \delta^*; \quad (1.16)$$

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu}} r^{\frac{2}{\nu} - \frac{3}{p}} \|(u, b)(t)\|_{L^{p, \infty}(B_r(x_0))} > \delta^*, \frac{1}{\mu} + \frac{1}{\nu} = \frac{1}{2}, 2 \leq \nu \leq \frac{2p}{3}, 3 < p \leq \infty; \quad (1.17)$$

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu}} r^{\frac{2}{\nu} - \frac{3}{p} + 1} \|(\nabla u, \nabla b)(t)\|_{L^p(B_r(x_0))} > \delta^* \text{ for } \frac{1}{\mu} + \frac{1}{\nu} = \frac{1}{2}, \nu \in \begin{cases} [2, \infty], & p \geq 3 \\ [2, \frac{2p}{3-p}], & \frac{3}{2} \leq p < 3 \end{cases} \quad (1.18)$$

where $\delta^* > 0$ is independent on μ, ν, p and r .

2 Several auxiliary lemmas

In this section, we shall give several auxiliary lemmas, which present some uniform estimates of invariant quantities and a crucial embedding theorem from the Lorentz space $L^{p, \infty}$ to a Morrey type space.

Lemma 2.1 *Let (u, b, P) be a suitable weak solution of equations (1.1) on $Q_1(z_0)$ satisfying*

$$\left\| \left(\sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(x, t) - u_{x_0, \rho}|^p \, dx \right)^{\frac{1}{p}}, \sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(x, t) - b_{x_0, \rho}|^p \, dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} = M < \infty \quad (2.1)$$

or

$$\left\| \left(\sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(x, t)|^p dx \right)^{\frac{1}{p}}, \sup_{\eta \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(x, t)|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} = M < \infty, \quad (2.2)$$

where p and q satisfy $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$, then for $0 < r < \rho \leq 1$, we have following estimates

$$C(u, b, r, z_0) \leq C \frac{r}{\rho} C(u, b, \rho, z_0) + C \left(\frac{\rho}{r} \right)^2 B^{\frac{9-3p}{6-p}}(u, b, \rho, z_0) M^{\frac{3p}{6-p}}, \quad 2 \leq p < 3; \quad (2.3)$$

$$C(u, b, r, z_0) \leq C \frac{r}{\rho} C(u, b, \rho, z_0) + C \left(\frac{\rho}{r} \right) A^{\frac{p-3}{p-2}}(u, b, \rho, z_0) M^{\frac{p}{p-2}}, \quad 3 \leq p \leq 6; \quad (2.4)$$

$$C(u, b, r, z_0) \leq C \frac{r}{\rho} C(u, b, \rho, z_0) + C \left(\frac{\rho}{r} \right)^{\frac{3}{2}} A^{\frac{3}{4}}(u, b, \rho, z_0) M^{\frac{3}{2}}, \quad 6 < p \leq \infty. \quad (2.5)$$

Proof. First, we consider the assumption (2.1), and the proof of conclusions will be divided into three cases:

case 1 : $2 \leq p < 3$;

case 2 : $3 \leq p \leq 6$;

case 3 : $6 < p \leq \infty$.

We define

$$f_p(t) = \left(\sup_{\rho \leq 1} \frac{1}{\rho} \int_{B_\rho(x_0)} |u(t, x) - u_{x_0, \rho}|^p dx \right)^{\frac{1}{p}}$$

and

$$g_p(t) = \left(\sup_{\rho \leq 1} \frac{1}{\rho} \int_{B_\rho(x_0)} |b(t, x) - b_{x_0, \rho}|^p dx \right)^{\frac{1}{p}}.$$

For almost every time $t \in (t_0 - \rho^2, t_0]$, we have

$$\begin{aligned} \int_{B_r(x_0)} (|u|^3 + |b|^3) dx &\leq \int_{B_r(x_0)} (|u - u_{x_0, \rho}|^3 + |b - b_{x_0, \rho}|^3) dx + C|B_r| (|u_{x_0, \rho}|^3 + |b_{x_0, \rho}|^3) \\ &= J_1 + J_2. \end{aligned} \quad (2.6)$$

For case 1 : $2 \leq p < 3$, J_1 can be estimated as

$$\begin{aligned} J_1 &\leq C \|u - u_{x_0, \rho}\|_{L^p}^{\frac{3p}{6-p}} \|u - u_{x_0, \rho}\|_{L^6}^{\frac{6(3-p)}{6-p}} + C \|b - b_{x_0, \rho}\|_{L^p}^{\frac{3p}{6-p}} \|b - b_{x_0, \rho}\|_{L^6}^{\frac{6(3-p)}{6-p}} \\ &\leq C \left(\|u - u_{x_0, \rho}\|_{L^6}^{\frac{6(3-p)}{6-p}} + \|b - b_{x_0, \rho}\|_{L^6}^{\frac{6(3-p)}{6-p}} \right) \left(\|u - u_{x_0, \rho}\|_{L^p}^{\frac{3p}{6-p}} + \|b - b_{x_0, \rho}\|_{L^p}^{\frac{3p}{6-p}} \right). \end{aligned} \quad (2.7)$$

Integrating with respect to time from $t_0 - r^2$ to t_0 and using Holder's inequality, one has

$$\begin{aligned} &\int_{t_0-r^2}^{t_0} \int_{B_r(x_0)} J_1 dx ds \\ &\leq C \left(\int_{t_0-r^2}^{t_0} \|u - u_{x_0, \rho}\|_{L^6(B_\rho(x_0))}^2 + \|b - b_{x_0, \rho}\|_{L^6(B_\rho(x_0))}^2 ds \right)^{\frac{9-3p}{6-p}} \\ &\quad \times \left(\int_{t_0-r^2}^{t_0} \|u - u_{x_0, \rho}\|_{L^p(B_\rho(x_0))}^{\frac{3p}{2p-3}} + \|b - b_{x_0, \rho}\|_{L^p(B_\rho(x_0))}^{\frac{3p}{2p-3}} ds \right)^{\frac{2p-3}{6-p}} \\ &\leq C \left(\int_{t_0-r^2}^{t_0} \int_{B_\rho(x_0)} (|\nabla u|^2 + |\nabla b|^2) dx ds \right)^{\frac{9-3p}{6-p}} \rho^{\frac{3}{6-p}} \left(\int_{t_0-r^2}^{t_0} f_p^{\frac{3p}{2p-3}} + g_p^{\frac{3p}{2p-3}} ds \right)^{\frac{2p-3}{6-p}}. \end{aligned} \quad (2.8)$$

Using the assumption (2.1), and if $2 < p < 3$, we see that

$$\begin{aligned}
\int_{t_0-r^2}^{t_0} f_p^{\frac{3p}{2p-3}} + g_p^{\frac{3p}{2p-3}} ds &\leq \int_{t_0-r^2}^{t_0} (f_p + g_p)^{\frac{3p}{2p-3}} ds \\
&= \frac{3p}{2p-3} \int_0^\infty \sigma^{\frac{3+p}{2p-3}} |\{s \in [t-r^2, t]; (f_p + g_p) > \sigma\}| d\sigma \\
&\leq \frac{3p}{2p-3} \left\{ \int_0^R \sigma^{\frac{3+p}{2p-3}} r^2 d\sigma + M^{\frac{2p}{p-2}} \int_R^\infty \sigma^{\frac{3+p}{2p-3} - \frac{2p}{p-2}} d\sigma \right\} \\
&\leq R^{\frac{3p}{2p-3}} r^2 + \frac{3(p-2)}{p} R^{\frac{3p}{2p-3} - \frac{2p}{p-2}} M^{\frac{2p}{p-2}} \\
&\leq \left(\frac{4p-6}{p} \right) r^{\frac{p}{2p-3}} M^{\frac{3p}{2p-3}},
\end{aligned} \tag{2.9}$$

where we take $R = r^{-\frac{p-2}{p}} M$. When $p = 2$, we take $R = M$ such that

$$\mu\{s \in [t-r^2, t]; (f_p + g_p) > M\} = 0.$$

Indeed,

$$\begin{aligned}
\int_{t_0-r^2}^{t_0} f_p^{\frac{3p}{2p-3}} + g_p^{\frac{3p}{2p-3}} ds &\leq \int_{t_0-r^2}^{t_0} (f_p + g_p)^{\frac{3p}{2p-3}} ds \\
&= \frac{3p}{2p-3} \int_0^\infty \sigma^{\frac{3+p}{2p-3}} |\{s \in [t-r^2, t]; (f_p + g_p) > \sigma\}| d\sigma \\
&\leq \frac{3p}{2p-3} \int_0^M \sigma^{\frac{3+p}{2p-3}} r^2 d\sigma \leq \left(\frac{4p-6}{p} \right) r^{\frac{p}{2p-3}} M^{\frac{3p}{2p-3}},
\end{aligned} \tag{2.10}$$

where we used the fact $r < \rho \leq 1$. The estimate of (2.9)-(2.10) together with (2.8) yield that

$$\begin{aligned}
&\int_{t_0-r^2}^{t_0} \int_{B_r(x_0)} J_1 dx ds \\
&\leq C \left(\frac{4p-6}{p} \right)^{\frac{2p-3}{6-p}} \rho^{\frac{3}{6-p}} r^{\frac{p}{6-p}} \left(\int_{t_0-r^2}^{t_0} \int_{B_\rho(x_0)} (|\nabla u|^2 + |\nabla b|^2) dx ds \right)^{\frac{9-3p}{6-p}} M^{\frac{3p}{6-p}}.
\end{aligned} \tag{2.11}$$

For J_2 , we have

$$J_2 \leq C \left(\frac{r}{\rho} \right)^3 \int_{B_\rho(x_0)} (|u|^3 + |b|^3) dx \tag{2.12}$$

Combining (2.6), (2.11) and (2.12), one get

$$\begin{aligned}
\frac{1}{r^2} \int_{Q_r(z_0)} (|u|^3 + |b|^3) dx dt &\leq C \rho^{\frac{3}{6-p}} r^{\frac{3p-12}{6-p}} \left(\int_{t-r^2}^t \int_{Q_\rho(z_0)} (|\nabla u|^2 + |\nabla b|^2) dx ds \right)^{\frac{9-3p}{6-p}} M^{\frac{3p}{6-p}} \\
&\quad + C \frac{r}{\rho} \frac{1}{\rho^2} \int_{\mathbb{P}_\rho(z)} (|u|^3 + |b|^3) dx dt.
\end{aligned} \tag{2.13}$$

That is to say

$$C(u, b, r, z_0) \leq C \frac{r}{\rho} C(u, b, \rho, z_0) + C \left(\frac{\rho}{r} \right)^2 B(u, b, \rho, z_0)^{\frac{9-3p}{6-p}} M^{\frac{3p}{6-p}}. \tag{2.14}$$

For case 2 : $3 \leq p \leq 6$, J_1 can be showed as

$$\begin{aligned}
J_1 &\leq C \|u - u_{x,\rho}\|_{L^p}^{\frac{p}{p-2}} \|u - u_{x_0,\rho}\|_{L^2}^{\frac{2(p-3)}{p-2}} + C \|b - b_{x_0,\rho}\|_{L^p}^{\frac{p}{p-2}} \|b - b_{x_0,\rho}\|_{L^2}^{\frac{2(p-3)}{p-2}} \\
&\leq C \left(\|u - u_{x_0,\rho}\|_{L^2}^{\frac{2(p-3)}{p-2}} + \|b - b_{x_0,\rho}\|_{L^2}^{\frac{2(p-3)}{p-2}} \right) \left(\|u - u_{x_0,\rho}\|_{L^p}^{\frac{p}{p-2}} + \|b - b_{x_0,\rho}\|_{L^p}^{\frac{p}{p-2}} \right) \\
&\leq C \rho A(\rho, z)^{\frac{p-3}{p-2}} \left(\frac{1}{\rho} \right)^{\frac{1}{p-2}} \left(\|u - u_{x_0,\rho}\|_{L^p}^{\frac{p}{p-2}} + \|b - b_{x_0,\rho}\|_{L^p}^{\frac{p}{p-2}} \right).
\end{aligned} \tag{2.15}$$

Together with estimates J_1 and J_2 and integrating with respect to time from $t_0 - r^2$ to t_0 , we get

$$\begin{aligned}
&\int_{t_0-r^2}^{t_0} \int_{B_r(x_0)} |u|^3 + |b|^3 dx ds \\
&\leq C \left(\frac{r}{\rho} \right)^3 \int_{t_0-\rho^2}^{t_0} \int_{B_\rho(x_0)} |u|^3 + |b|^3 dx ds + C \rho A(u, b, \rho, z_0)^{\frac{p-3}{p-2}} \int_{t_0-r^2}^{t_0} f_p^{\frac{p}{p-2}} + g_p^{\frac{p}{p-2}} ds,
\end{aligned} \tag{2.16}$$

under the assumption (2.1), we infer that

$$\begin{aligned}
\int_{t_0-r^2}^{t_0} f_p^{\frac{p}{p-2}} + g_p^{\frac{p}{p-2}} ds &\leq \int_{t-r^2}^t (f_p + g_p)^{\frac{p}{p-2}} ds \\
&= \frac{p}{p-2} \int_0^\infty \sigma^{\frac{2}{p-2}} |\{s \in [t-r^2, t] : (f_p + g_p)(s) > \sigma\}| d\sigma \\
&= \frac{p}{p-2} \left\{ \int_0^R \sigma^{\frac{2}{p-2}} |\{s \in [t-r^2, t] : (f_p + g_p)(s) > \sigma\}| d\sigma \right. \\
&\quad \left. + \int_R^\infty \sigma^{\frac{2}{p-2}} |\{s \in [t-r^2, t] : (f_p + g_p)(s) > \sigma\}| d\sigma \right\} \\
&\leq \frac{p}{p-2} \int_0^R \sigma^{\frac{2}{p-2}} r^2 d\sigma + \frac{p}{p-2} \int_R^\infty \sigma^{\frac{2}{p-2} - \frac{2p}{p-2}} d\sigma M^{\frac{2p}{p-2}} \\
&= r^2 R^{\frac{p}{p-2}} + \frac{p^2}{(p-2)^2} R^{-\frac{p}{p-2}} M^{\frac{2p}{p-2}} \\
&= \left(1 + \frac{p^2}{(p-2)^2} \right) r M^{\frac{p}{p-2}},
\end{aligned} \tag{2.17}$$

where we have take $R = r^{-\frac{p-2}{p}} M$. Combining (2.7) and (2.17) it follows that

$$\begin{aligned}
&\int_{t_0-r^2}^{t_0} \int_{B_r(x_0)} |u|^3 + |b|^3 dx ds \\
&\leq C \left(\frac{r}{\rho} \right)^3 \int_{t_0-\rho^2}^{t_0} \int_{B_\rho(x_0)} |u|^3 + |b|^3 dx ds + C \left(1 + \frac{p^2}{(p-2)^2} \right) \rho A(u, b, \rho, z_0)^{\frac{p-3}{p-2}} M^{\frac{p}{p-2}}.
\end{aligned} \tag{2.18}$$

Therefore, we arrive at

$$C(u, b, r, z_0) \leq C \frac{r}{\rho} C(u, b, \rho, z_0) + C \left(\frac{\rho}{r} \right) A(u, b, \rho, z_0)^{\frac{p-3}{p-2}} M^{\frac{p}{p-2}}. \tag{2.19}$$

For case 3 : $6 < p \leq \infty$, J_1 can be showed as

$$\begin{aligned}
J_1 &\leq C \|u - u_{x_0, \rho}\|_{L^p}^{\frac{3}{2}} \|u - u_{x_0, \rho}\|_{L^2}^{\frac{3}{2}} \rho^{\frac{3(p-6)}{4p}} + C \|b - b_{x_0, \rho}\|_{L^p}^{\frac{3}{2}} \|b - b_{x_0, \rho}\|_{L^2}^{\frac{3}{2}} \rho^{\frac{3(p-6)}{4p}} \\
&\leq C \left(\|u - u_{x_0, \rho}\|_{L^2}^{\frac{3}{2}} + \|b - b_{x_0, \rho}\|_{L^2}^{\frac{3}{2}} \right) \left(\|u - u_{x_0, \rho}\|_{L^p}^{\frac{3}{2}} + \|b - b_{x_0, \rho}\|_{L^p}^{\frac{3}{2}} \right) \rho^{\frac{3(p-6)}{4p}} \\
&\leq C \rho A^{\frac{3}{4}}(u, b, \rho, z_0) \rho^{\frac{3(p-6)}{4p} + \frac{3}{4}} \left(\|u - u_{x_0, \rho}\|_{L^p}^{\frac{3}{2}} + \|b - b_{x_0, \rho}\|_{L^p}^{\frac{3}{2}} \right) \\
&\leq C \rho^{\frac{3}{4} + \frac{3}{2p} + \frac{3(p-6)}{4p}} A^{\frac{3}{4}}(u, b, \rho, z_0) (f_p + g_p)^{\frac{3}{2}}.
\end{aligned} \tag{2.20}$$

Together with estimates J_1 and J_2 and integrating with respect to time from $t_0 - r^2$ to t_0 , we get

$$\begin{aligned}
&\int_{t_0 - r^2}^{t_0} \int_{B_r(x_0)} |u|^3 + |b|^3 dx ds \\
&\leq C \left(\frac{r}{\rho} \right)^3 \int_{t_0 - \rho^2}^{t_0} \int_{B_\rho(x_0)} |u|^3 + |b|^3 dx ds + C \rho^{\frac{3p-6}{2p}} A(u, b, \rho, z_0)^{\frac{3}{4}} \int_{t_0 - r^2}^{t_0} (f_p + g_p)^{\frac{3}{2}}(s) ds,
\end{aligned} \tag{2.21}$$

under the assumption (2.1), it follows that

$$\begin{aligned}
&\int_{t_0 - r^2}^{t_0} (f_p + g_p)^{\frac{3}{2}}(s) ds \\
&= \frac{3}{2} \int_0^\infty \sigma^{\frac{1}{2}} \left| \left\{ s \in [t - r^2, t] : (f_p + g_p)^{\frac{3}{2}}(s) > \sigma \right\} \right| d\sigma \\
&= \frac{3}{2} \left\{ \int_0^R \sigma^{\frac{1}{2}} \left| \left\{ s \in [t - r^2, t] : (f_p + g_p)^{\frac{3}{2}}(s) > \sigma \right\} \right| d\sigma \right. \\
&\quad \left. + \int_R^\infty \sigma^{\frac{1}{2}} \left| \left\{ s \in [t - r^2, t] : (f_p + g_p)^{\frac{3}{2}}(s) > \sigma \right\} \right| d\sigma \right. \\
&\leq \frac{3}{2} \int_0^R \sigma^{\frac{1}{2}} r^2 d\sigma + \frac{3}{2} \int_R^\infty \sigma^{\frac{1}{2} - \frac{2p}{p-2}} d\sigma M^{\frac{2p}{p-2}} \\
&= r^2 R^{\frac{3}{2}} - \frac{3(p-2)}{p-6} R^{\frac{3}{2} - \frac{2p}{p-2}} M^{\frac{2p}{p-2}} \\
&= \left(1 + \frac{3(p-2)}{6+p} \right) r^{2 - \frac{3p-6}{2p}} M^{\frac{3}{2}},
\end{aligned} \tag{2.22}$$

where we have take $R = r^{\frac{2-p}{p}} M$. Substituting (2.22) into (2.21), one has

$$\begin{aligned}
&\int_{t_0 - r^2}^{t_0} \int_{B_r(x_0)} |u|^3 + |\nabla Q|^3 dx ds \\
&\leq C \left(\frac{r}{\rho} \right)^3 \int_{t_0 - \rho^2}^{t_0} \int_{B_\rho(x_0)} |u|^3 + |b|^3 dx ds + C \rho^{\frac{3p-6}{2p}} r^{2 - \frac{3p-6}{2p}} A(u, b, \rho, z_0)^{\frac{3}{4}} M^{\frac{3}{2}},
\end{aligned} \tag{2.23}$$

which allows us to get

$$C(u, b, r, z_0) \leq C \frac{r}{\rho} C(u, b, \rho, z_0) + C \left(\frac{\rho}{r} \right)^{\frac{3}{2}} A(u, b, \rho, z_0)^{\frac{3}{4}} M^{\frac{3}{2}}. \tag{2.24}$$

We complete the proof of Lemma 2.1 under the assumption (2.1).

Now, we consider the assumption (2.2) is true, we denote

$$f_p(t) = \sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(t, x)|^p dx \right)^{\frac{1}{p}}$$

and

$$g_p(t) = \sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(t, x)|^p dx \right)^{\frac{1}{p}},$$

and we can modify the processes of proofs as follows:

In the case $2 \leq p < 3$, we can replace $\|u - u_\rho\|_{L^p(B_\rho)}$ and $\|b - b_\rho\|_{L^p(B_\rho)}$ by $C_p \|u\|_{L^p(B_\rho)}$ and $C_p \|b\|_{L^p(B_\rho)}$ in (2.8) and repeat the processes of proofs for (2.9)-(2.13) to get (2.3). The difference is that in this case, the constant C_p is depended on p . Noticing $2 \leq p < 3$, we can choose a large enough constant C to get rid of the dependence on p .

If $3 \leq p \leq 6$ or $6 < p \leq \infty$, we just need to replace $\int_{B_r(x_0)} (|u - u_{x_0, \rho}|^3 + |b - b_{x_0, \rho}|^3) dx$ by $C \int_{B_\rho(x_0)} (|u|^3 + |b|^3) dx$ and repeat the processes of proofs step by step to get (2.4) and (2.5). Thus the proof of Lemma 2.1 is completed under the assumption (2.2).

Lemma 2.2 *Let (u, b, P) be a suitable weak solution of equations (1.1) on $Q_1(z_0)$ satisfying*

$$\left\| \left(\sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(x, t) - u_{x_0, \rho}|^p dx \right)^{\frac{1}{p}}, \sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(x, t) - b_{x_0, \rho}|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} = M < \infty \quad (2.25)$$

or

$$\left\| \left(\sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(x, t)|^p dx \right)^{\frac{1}{p}}, \sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(x, t)|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} = M < \infty, \quad (2.26)$$

where p and q satisfy $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$, then there exists a constant $\rho_0 > 0$ only depended on $A(u, b, 1, z_0)$, $B(u, b, 1, z_0)$, $C(u, b, 1, z_0)$ and $D(P, 1, z_0)$, such that for $r \leq \rho_0$, it follows

$$A(u, b, r, z_0) + B(u, b, r, z_0) + C(u, b, r, z_0) + D(P, r, z_0) \leq C(M).$$

Proof. Without loss of generality, we set $z_0 = 0$ and for any $2r < \rho < 1$, let ζ be a cutoff function, which vanishes outside of Q_ρ and equals 1 in $Q_{\frac{\rho}{2}}$, and satisfies

$$|\nabla \zeta| \leq C\rho^{-1}, \quad |\partial_t \zeta|, |\Delta \zeta| \leq C\rho^{-2}.$$

Define the backward heat kernel as

$$\Gamma(x, t) = \frac{1}{4\pi(r^2 - t)^{\frac{3}{2}}} e^{-\frac{|x|^2}{4(r^2 - t)}}.$$

Taking the test function $\phi = \Gamma \zeta$ in the local energy inequality, and noting that

$$(\partial_t + \Delta)\Gamma = 0,$$

we have

$$\begin{aligned}
& \sup_t \int_{B_r} (|u(x,t)|^2 + |b(x,t)|^2) \phi dx + \int_{Q_r} (|\nabla u(x,t)|^2 + |\nabla b(x,t)|^2) \phi dx dt \\
& \leq \int_{Q_\rho} [(|u|^2 + |b|^2) (\Delta \phi + \partial_t \phi) + u \cdot \nabla \phi (|u|^2 + |b|^2 + 2p) - (b \cdot u) (b \cdot \nabla \phi)] dx dt \\
& \leq \int_{Q_\rho} [(|u|^2 + |b|^2) (\Gamma \Delta \zeta + \Gamma \partial_t \zeta + 2 \nabla \Gamma \cdot \nabla \zeta) + |\nabla \phi| |u| (|u|^2 + |b|^2 + 2P)] dx dt.
\end{aligned} \tag{2.27}$$

Direct computation shows that

$$\begin{aligned}
Cr^{-3} & \geq \Gamma(x,t) \geq C^{-1}r^{-3} \quad \text{in } Q_r; \\
|\nabla \phi| & \leq |\nabla \Gamma| \zeta + \Gamma |\nabla \zeta| \leq Cr^{-4}; \\
|\Gamma \Delta \zeta| + |\Gamma \partial_t \zeta| + 2|\nabla \Gamma \cdot \nabla \zeta| & \leq C\rho^{-5}.
\end{aligned}$$

From these inequalities and the Hölder's inequality, it follows that

$$\begin{aligned}
& A(u, b, r) + B(u, b, r) \\
& \leq r^2 \int_{Q_\rho} [(|u|^2 + |b|^2) (\Gamma \Delta \zeta + \Gamma \partial_t \zeta + 2 \nabla \Gamma \cdot \nabla \zeta) + |\nabla \phi| |u| (|u|^2 + |b|^2 + 2P)] dx dt \\
& \leq C \left(\frac{r}{\rho} \right)^2 A(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 \rho^{-2} \int_{Q_\rho} |u|^3 + |b|^3 + 2|u||p| dx dt \\
& = C \left(\frac{r}{\rho} \right)^2 A(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 C(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 \rho^{-2} \int_{Q_\rho} 2|u||P| dx dt \\
& \leq C \left(\frac{r}{\rho} \right)^2 A(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 C(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 C^{\frac{1}{3}}(u, b, \rho) D^{\frac{2}{3}}(P, \rho) \\
& \leq C \left(\frac{r}{\rho} \right)^2 A(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 C(u, b, \rho) + C \left(\frac{\rho}{r} \right)^2 D(P, \rho).
\end{aligned} \tag{2.28}$$

Now, we turn to show bounds on $D(P, r)$. Note that p satisfies the following equation in distribution sense:

$$-\Delta P = \partial_i \partial_j (u_i u_j - b_i b_j),$$

which in turn can be derived by calculating the divergence of (1.1)₁. Let $\eta(x)$ be a cut-off function, which equals 1 in $B_{\frac{\rho}{2}}$ and vanishes outside of B_ρ . Set $P = P_1 + P_2$ with

$$P_1 = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{1}{|x-y|} \partial_i \partial_j [(u_i u_j - b_i b_j) \eta] dy$$

and P_2 is harmonic in $B_{\frac{\rho}{2}}$. Using the Calderón-Zygmund inequality, one know that

$$\int_{B_\rho} |P_1|^{\frac{3}{2}} dx \leq C \int_{B_\rho} |u|^3 + |b|^3 dx.$$

Since p_2 is harmonic in $B_{\frac{\rho}{2}}$, we conclude that

$$\int_{B_r} |P_2|^{\frac{3}{2}} dx \leq Cr^3 \sup_{x \in B_r} |P_2(x)|^{\frac{3}{2}} \leq C \left(\frac{r}{\rho} \right)^3 \int_{B_\rho} |P_2(x)|^{\frac{3}{2}} dx$$

for any $r \leq \frac{\rho}{2}$. Hence, we infer that

$$\begin{aligned} \int_{B_r} |P|^{\frac{3}{2}} dx &\leq C \int_{B_\rho} |u|^3 + |b|^3 dx + C \left(\frac{r}{\rho}\right)^3 \int_{B_\rho} |P - P_1|^{\frac{3}{2}} dx \\ &\leq C \int_{B_\rho} |u|^3 + |b|^3 dx + C \left(\frac{r}{\rho}\right)^3 \int_{B_\rho} |P|^{\frac{3}{2}} dx. \end{aligned}$$

Integrating above inequality with respect to t from $-r^2$ to 0, we get

$$\int_{Q_r} |P|^{\frac{3}{2}} dx dt \leq C \int_{Q_\rho} |u|^3 + |b|^3 dx dt + C \left(\frac{r}{\rho}\right)^3 \int_{Q_\rho} |P|^{\frac{3}{2}} dx dt,$$

which implies

$$\begin{aligned} D(P, r) &\leq C \frac{1}{r^2} \int_{Q_\rho} (|u|^3 + |b|^3) dx dt + C \frac{1}{r^2} \left(\frac{r}{\rho}\right)^3 \int_{Q_\rho} |P|^{\frac{3}{2}} dx dt \\ &\leq C \left(\frac{\rho}{r}\right)^2 C(u, b, \rho) + C \frac{r}{\rho} D(P, \rho). \end{aligned} \quad (2.29)$$

By using Young's inequality, we calculate

$$\begin{aligned} D^{\frac{8}{7}}(P, r) &\leq C \left(\frac{\rho}{r}\right)^{\frac{16}{7}} C^{\frac{8}{7}}(u, b, \rho) + C \left(\frac{r}{\rho}\right)^{\frac{8}{7}} D^{\frac{8}{7}}(P, \rho) \\ &\leq C \left(\frac{\rho}{r}\right)^{\frac{16}{7}} \left(\frac{\rho}{r}\right)^{\frac{8}{7}} \left(\frac{r}{\rho}\right)^{\frac{8}{7}} C^{\frac{8}{7}}(u, b, \rho) + C \left(\frac{r}{\rho}\right)^{\frac{8}{7}} D^{\frac{8}{7}}(P, \rho) \\ &\leq \left(\frac{r}{\rho}\right)^{\frac{7}{6}} C^{\frac{7}{6}}(u, b, \rho) + C \left(\frac{r}{\rho}\right)^{\frac{8}{7}} D^{\frac{8}{7}}(P, \rho) + C \left(\frac{\rho}{r}\right)^{168}. \end{aligned} \quad (2.30)$$

Similarly, we show some crucial bounds for $C(u, b, r)$ under the assumptions of Lemma 2.1. First, from Lemma 2.1, we know that

$$\begin{cases} C(u, b, r) \leq C \frac{r}{\rho} C(u, b, \rho) + C \left(\frac{\rho}{r}\right)^2 B(u, b, \rho)^{\frac{9-3p}{6-p}} M^{\frac{3p}{6-p}}, & 2 \leq p < 3; \\ C(u, b, r) \leq C \frac{r}{\rho} C(u, b, \rho) + C \left(\frac{\rho}{r}\right)^{\frac{p-3}{p-2}} A(u, b, \rho)^{\frac{p}{p-2}} M^{\frac{p}{p-2}}, & 3 \leq p \leq 6; \\ C(u, b, r) \leq C \frac{r}{\rho} C(u, b, \rho) + C \left(\frac{\rho}{r}\right)^{\frac{3}{2}} A(u, b, \rho)^{\frac{3}{4}} M^{\frac{3}{2}}, & 6 < p \leq \infty. \end{cases}$$

Clearly, we have

$$\begin{cases} C^{\frac{7}{6}}(u, b, r) \leq C \left(\frac{r}{\rho}\right)^{\frac{7}{6}} C^{\frac{7}{6}}(u, b, \rho) + C \left(\frac{r}{\rho}\right)^{\frac{7}{6}} B(u, b, \rho) + C \left(\frac{\rho}{r}\right)^{40} M^{\frac{7p}{5p-9}}, & 2 \leq p < 3; \\ C^{\frac{7}{6}}(u, b, r) \leq C \left(\frac{r}{\rho}\right)^{\frac{7}{6}} C^{\frac{7}{6}}(u, b, \rho) + \left(\frac{r}{\rho}\right) A(u, b, \rho) + C \left(\frac{\rho}{r}\right)^4 M^{\frac{7p}{9-p}}, & 3 \leq p \leq 6; \\ C^{\frac{7}{6}}(u, b, r) \leq C \left(\frac{r}{\rho}\right)^{\frac{7}{6}} C^{\frac{7}{6}}(u, b, \rho) + \left(\frac{r}{\rho}\right)^{\frac{7}{6}} A(u, b, \rho) + C \left(\frac{\rho}{r}\right)^{23} M^{14}, & 6 < p \leq \infty. \end{cases}$$

From (2.28) and combine with above estimates of $C^{\frac{7}{6}}(r)$ and $D^{\frac{8}{7}}(r)$, we discover that

$$\begin{aligned} &A(r) + B(r) + C^{\frac{7}{6}}(r) + D^{\frac{8}{7}}(r) \\ &\leq C \left(\frac{r}{\rho}\right) (A(\rho) + B(\rho)) + C \left(\frac{r}{\rho}\right)^{\frac{7}{6}} C^{\frac{7}{6}}(\rho) + C \left(\frac{r}{\rho}\right)^{\frac{8}{7}} D^{\frac{8}{7}}(\rho) + C \left(\frac{\rho}{r}\right)^{40} M^{\frac{7p}{5p-9}} \\ &\quad + C \left(\left(\frac{\rho}{r}\right)^{14} + \left(\frac{\rho}{r}\right)^{16} + \left(\frac{\rho}{r}\right)^{168} \right), \end{aligned} \quad (2.31)$$

$$\begin{aligned}
& A(r) + B(r) + C^{\frac{7}{6}}(r) + D^{\frac{8}{7}}(r) \\
& \leq C \left(\frac{r}{\rho} \right) (A(\rho) + B(\rho)) + C \left(\frac{r}{\rho} \right)^{\frac{7}{6}} C^{\frac{7}{6}}(\rho) + C \left(\frac{r}{\rho} \right)^{\frac{8}{7}} D^{\frac{8}{7}}(\rho) + C \left(\frac{\rho}{r} \right)^4 M^{\frac{7p}{9-p}} \\
& \quad + C \left(\left(\frac{\rho}{r} \right)^{14} + \left(\frac{\rho}{r} \right)^{16} + \left(\frac{\rho}{r} \right)^{168} \right),
\end{aligned} \tag{2.32}$$

$$\begin{aligned}
& A(r) + B(r) + C^{\frac{7}{6}}(r) + D^{\frac{8}{7}}(r) \\
& \leq C \left(\frac{r}{\rho} \right) (A(\rho) + B(\rho)) + C \left(\frac{r}{\rho} \right)^{\frac{7}{6}} C^{\frac{7}{6}}(\rho) + C \left(\frac{r}{\rho} \right)^{\frac{8}{7}} D^{\frac{8}{7}}(\rho) + C \left(\frac{\rho}{r} \right)^{23} M^{14} \\
& \quad + C \left(\left(\frac{\rho}{r} \right)^{14} + \left(\frac{\rho}{r} \right)^{16} + \left(\frac{\rho}{r} \right)^{168} \right).
\end{aligned} \tag{2.33}$$

These estimates combined with the fact that $\frac{r}{\rho} < 1$, one get

$$\begin{cases} G(r) \leq C \left(\frac{r}{\rho} \right) G(\rho) + C \left(1 + M^{\frac{7p}{5p-9}} \right) \left(\frac{\rho}{r} \right)^{168}, \\ G(r) \leq C \left(\frac{r}{\rho} \right) G(\rho) + C \left(1 + M^{\frac{7p}{9-p}} \right) \left(\frac{\rho}{r} \right)^{168}, \\ G(r) \leq C \left(\frac{r}{\rho} \right) G(\rho) + C \left(1 + M^{14} \right) \left(\frac{\rho}{r} \right)^{168}. \end{cases}$$

Let $r = \theta\rho$ with $\theta < \frac{1}{2}$ such that $C \cdot \frac{r}{\rho} < \frac{1}{2}$, i.e.,

$$\begin{cases} G(\theta\rho) \leq \frac{1}{2}G(\rho) + C \left(1 + M^{\frac{7p}{5p-9}} \right) \theta^{-168}, \\ G(\theta\rho) \leq \frac{1}{2}G(\rho) + C \left(1 + M^{\frac{7p}{9-p}} \right) \theta^{-168}, \\ G(\theta\rho) \leq \frac{1}{2}G(\rho) + C \left(1 + M^{14} \right) \theta^{-168}. \end{cases} \tag{2.34}$$

Iterating the inequality (2.34) k times yields

$$\begin{cases} G(\theta^k \rho) \leq \frac{1}{2^k} G(\rho) + C \left(1 + M^{\frac{7p}{5p-9}} \right) \theta^{-168k}, \\ G(\theta^k \rho) \leq \frac{1}{2^k} G(\rho) + C \left(1 + M^{\frac{7p}{9-p}} \right) \theta^{-168k}, \\ G(\theta^k \rho) \leq \frac{1}{2^k} G(\rho) + C \left(1 + M^{14} \right) \theta^{-168k}. \end{cases} \tag{2.35}$$

These complete the proof of Lemma 2.2.

Lemma 2.3 *Let (u, b, P) be a suitable weak solution of equations (1.1) on $Q_1(z_0)$. For any fixed $\epsilon > 0$, there exists two constants δ and r^* depended ϵ on such that if*

$$\begin{aligned}
& \left\| \left(\sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(x, t) - u_{x_0, \rho}|^p dx \right)^{\frac{1}{p}}, \sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(x, t) - b_{x_0, \rho}|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} \\
& \leq \delta \ll 1
\end{aligned} \tag{2.36}$$

or

$$\left\| \left(\sup_{\rho \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |u(x, t)|^p dx \right)^{\frac{1}{p}}, \sup_{\eta \leq 1} \left(\frac{1}{\rho} \int_{B_\rho(x_0)} |b(x, t)|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} \leq \delta \ll 1, \tag{2.37}$$

where p and q satisfy $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$, then it follows

$$C(u, b, r, z_0) + D(P, r, z_0) \leq \epsilon, \quad 0 < r \leq r^* < 1.$$

Proof. Without loss of generality, we assume $z_0 = 0$ and $\delta \ll 1$. From Lemma 2.2, we know that if $\rho \leq \rho_0$, then

$$A(u, b, \rho) + B(u, b, \rho) + C(u, b, \rho) + D(P, \rho) \leq C. \quad (2.38)$$

Therefore, we choosing $\rho \leq \rho_0$ in (2.3) ($2 \leq p < 3$) and using (2.38) to obtain

$$\begin{aligned} C(u, b, r) &\leq C \frac{r}{\rho} C(u, b, \rho) + C \left(\frac{\rho}{r}\right)^2 B^{\frac{9-3p}{6-p}}(u, b, \rho) \delta^{\frac{3p}{6-p}} \\ &\leq C \frac{r}{\rho} + C \left(\frac{\rho}{r}\right)^2 \delta^{\frac{3}{2}}. \end{aligned} \quad (2.39)$$

We take $r = 2^{-k} \rho_0$ in (2.39) and obtain by a standard iterative argument that

$$\begin{aligned} C(u, b, 2^{-k} \rho_0) &\leq \frac{1}{2} C(u, b, 2^{-k+1} \rho_0) + C 2^2 \delta^{\frac{3}{2}} \\ &\leq \left(\frac{1}{2}\right)^k C(u, b, \rho_0) + \sum_{j=0}^{k-1} \left(\frac{1}{2}\right)^j C(2^2 \delta^{\frac{3}{2}}) \\ &\leq \left(\frac{1}{2}\right)^k C(u, b, \rho_0) + C 2^2 \delta^{\frac{3}{2}}. \end{aligned} \quad (2.40)$$

Similarly, using the same trick to see that

$$\begin{aligned} D(P, 2^{-k_0 k} \rho_0) &\leq \frac{1}{2} D(P, 2^{-k_0(k-1)} \rho_0) + C 2^{2k_0} \left(\left(\frac{1}{2}\right)^{k_0(k-1)} C(u, b, \rho_0) + C 2^2 \delta^{\frac{3}{2}} \right) \\ &\leq \left(\frac{1}{2}\right)^k D(P, 2^{-k_0} \rho_0) + C 2^{2k_0} \left(\left(\frac{1}{2}\right)^{k_0(k-1)} C(u, b, \rho_0) + C 2^2 \delta^{\frac{3}{2}} \right), \end{aligned} \quad (2.41)$$

where we choose k_0 such that $C 2^{-k_0} \leq \frac{1}{2}$. Summing up (2.40) and (2.41), we have

$$\begin{aligned} &C(u, b, 2^{-k_0 k} \rho_0) + D(P, 2^{-k_0 k} \rho_0) \\ &\leq \left(\left(\frac{1}{2}\right)^k + \left(\frac{1}{2}\right)^{k_0 k} \right) (C(u, b, \rho_0) + D(P, 2^{-k_0} \rho_0)) + C 2^{2k_0} \left(\left(\frac{1}{2}\right)^{k_0(k-1)} C(u, b, \rho_0) + C 2^2 \delta^{\frac{3}{2}} \right) \\ &\leq C \left(\left(\frac{1}{2}\right)^k + \left(\frac{1}{2}\right)^{k_0 k} \right) + C 2^{2k_0} \left(\left(\frac{1}{2}\right)^{k_0(k-1)} + C 2^2 \delta^{\frac{3}{2}} \right), \end{aligned} \quad (2.42)$$

and then we take k large enough such that $C(2^{-k} + 2^{-k k_0} + 2^{-k k_0 + 3k_0}) \leq \frac{\epsilon}{2}$ and $r^* = 2^{-k k_0} \rho_0$, then take $\delta \leq \left(\frac{\epsilon}{2^{2k_0+3} C}\right)^{\frac{2}{3}}$, we discover that

$$C(u, b, r) + D(P, r) \leq \epsilon, \quad 0 < r \leq r^* < 1.$$

Next, by choosing $\rho \leq \rho_0$ in (2.4) ($3 \leq p \leq 6$) and using (2.38), we infer that

$$\begin{aligned} C(u, b, r) &\leq C \frac{r}{\rho} C(u, b, \rho) + C \left(\frac{\rho}{r}\right)^2 A^{\frac{p-3}{p-2}}(u, b, \rho) \delta^{\frac{p}{p-2}} \\ &\leq C \frac{r}{\rho} + C \left(\frac{\rho}{r}\right)^2 \delta^{\frac{3}{2}} \\ &\leq C \frac{r}{\rho} + C \left(\frac{\rho}{r}\right)^2 \delta^{\frac{3}{2}}, \end{aligned} \quad (2.43)$$

where we have used $\frac{p-3}{p-2} \leq 1$ and $\frac{p}{p-2} \geq \frac{3}{2}$. By choosing $\rho \leq \rho_0$ in (2.5) ($6 < p \leq \infty$) and using (2.38), we obtain by similar computations that

$$C(u, b, r) \leq C \frac{r}{\rho} + C \left(\frac{\rho}{r} \right)^2 \delta^{\frac{3}{2}}. \quad (2.44)$$

Therefore, repeating the process of (2.40)-(2.42), we also have

$$C(u, b, r) + D(P, r) \leq \epsilon, \quad 0 < r \leq r^* < 1.$$

These imply that the Lemma 2.3 holds true.

Lemma 2.4 [37] *For any given $r > 0$ and $2 \leq p \leq \infty$, it follows that*

$$\left(\sup_{\eta \leq r} \frac{1}{\eta} \int_{B_\eta} |f|^p dx \right)^{\frac{1}{p}} \leq C \|f\|_{L^{\frac{3p}{2}, \infty}(B_r)}, \quad (2.45)$$

where $C > 0$ is a constant independent on p .

Lemma 2.5 [15] *Let (u, b, P) be a suitable weak solution of the MHD equations (1.1) in $Q_1(z_0)$. There exists an $\epsilon > 0$ such that if*

$$C(u, b, r, z_0) + D(P, r, z_0) \leq \epsilon$$

for some $r > 0$, then (u, b) is regular at z_0 .

3 Proof of main results

This section is devoted to the proof of the main theorems. We first prove the local in space regularity near initial time, i.e. Theorem 1.1. Then, we prove the concentration result, i.e. Theorem 1.2.

3.1 Proof of Theorem 1.1

From Lemma 2.3, there exists two constants δ and r^* depended ϵ on such that if

$$\left\| \left(\sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t) - u_{x_0, \rho}|^p dx \right)^{\frac{1}{p}}, \sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t) - b_{x_0, \eta}|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} \leq \delta \quad (3.1)$$

or

$$\left\| \left(\sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t)|^p dx \right)^{\frac{1}{p}}, \sup_{\eta \leq 1} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t)|^p dx \right)^{\frac{1}{p}} \right) \right\|_{L^{q, \infty}[t_0-1, t_0]} \leq \delta, \quad (3.2)$$

where p and q satisfy $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$, then it follows that

$$C(u, b, r, z_0) + D(P, r, z_0) \leq \epsilon, \quad 0 < r \leq r^* < 1.$$

By Lemma 2.5, we deduce that z_0 is a regular point. This completes the proof of Theorem 1.1.

3.2 Proof of Theorem 1.2

The proof of Theorem 1.2 is by contradiction. If Theorem 1.2 is false, then there exists some $r_0 \in (0, 1)$ such that it holds that

$$\limsup_{t \rightarrow t_0^-} \left(\|u(t, x) - u(t)_{x_0, r_0}\|_{L^{3, \infty}(B_{r_0}(x_0))} + \|b(t, x) - b(t)_{x_0, r_0}\|_{L^{3, \infty}(B_{r_0}(x_0))} \right) \leq \delta^* \quad (3.3)$$

or

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu_0}} r_0^{\frac{2}{\nu_0} - \frac{3}{p_0}} \|(u, b)(t)\|_{L^{p_0, \infty}(B_{r_0}(x_0))} \leq \delta^*, \quad \frac{1}{\mu_0} + \frac{1}{\nu_0} = \frac{1}{2}, \quad 2 \leq \nu_0 \leq \frac{2p_0}{3}, \quad 3 < p_0 \leq \infty \quad (3.4)$$

or

$$\limsup_{t \rightarrow t_0^-} (t_0 - t)^{\frac{1}{\mu_0}} r_0^{\frac{2}{\nu_0} - \frac{3}{p_0} + 1} \|(\nabla u, \nabla b)(t)\|_{L^{p_0}(B_{r_0}(x_0))} \leq \delta^* \text{ for } \frac{1}{\mu_0} + \frac{1}{\nu_0} = \frac{1}{2}, \quad \nu_0 \in \begin{cases} [2, \infty], & p_0 \geq 3 \\ [2, \frac{2p_0}{3-2p_0}], & \frac{3}{2} \leq p_0 < 3 \end{cases} \quad (3.5)$$

If (3.3) holds, by using the fact

$$\min_{c \in \mathbb{R}} \int_{B_\eta(x_0)} |u - c|^2 dx = \int_{B_\eta(x_0)} |u - u_{x_0, \eta}|^2 dx, \quad \min_{c \in \mathbb{R}} \int_{B_\eta(x_0)} |b - c|^2 dx = \int_{B_\eta(x_0)} |b - b_{x_0, \eta}|^2 dx$$

and Lemma 2.4, we immediately get

$$\begin{aligned} & \limsup_{t \rightarrow t_0} \sup_{\eta < r_0} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u - u_{x_0, \eta}|^2 dx \right)^{\frac{1}{2}} + \limsup_{t \rightarrow t_0} \sup_{\eta < r_0} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b - b_{x_0, \eta}|^2 dx \right)^{\frac{1}{2}} \\ & \leq C \limsup_{t \rightarrow t_0} \sup_{\eta < r_0} \left(\eta^{-1} \left(\int_{B_\eta(x_0)} |u - u_{x_0, r_0}|^2 dx \right)^{\frac{1}{2}} \right) + C \limsup_{t \rightarrow t_0} \sup_{\eta < r_0} \left(\eta^{-1} \left(\int_{B_\eta(x_0)} |b - b_{x_0, r_0}|^2 dx \right)^{\frac{1}{2}} \right) \\ & \leq C \limsup_{t \rightarrow t_0} \|u(t) - u(t)_{x_0, r_0}\|_{L^{3, \infty}(B_{r_0}(x_0))} + C \limsup_{t \rightarrow t_0} \|b(t) - b(t)_{x_0, r_0}\|_{L^{3, \infty}(B_{r_0}(x_0))} \\ & \leq C\delta^* = \delta. \end{aligned} \quad (3.6)$$

This implies that

$$\left\| \left(\sup_{\eta \leq r_0^*} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t) - u_{x_0, \eta}|^2 dx \right)^{\frac{1}{2}}, \sup_{\eta \leq r_0^*} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t) - b_{x_0, \eta}|^2 dx \right)^{\frac{1}{2}} \right) \right\|_{L^\infty[t_0 - (r_0^*)^2, t_0]} \leq \delta \quad (3.7)$$

for some $r_0^* \leq r_0$. Due to Theorem 1.1, we know that z_0 is a regular point, this is a contradiction. If (3.4) holds, by using Hölder's inequality and Lemma 2.4, we find that

$$\begin{aligned}
& \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} \left(\sup_{\eta < r_0} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u|^{\nu_0} dx \right)^{\frac{1}{\nu_0}} + \sup_{\eta < r_0} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b|^{\nu_0} dx \right)^{\frac{1}{\nu_0}} \right) \\
& \leq \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} \sup_{\eta < r_0} \left(\left(\eta^{2 - \frac{3\nu_0}{p_0}} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u|^{\frac{2p_0}{3}} dx \right)^{\frac{3\nu_0}{2p_0}} \right)^{\frac{1}{\nu_0}} + \left(\eta^{2 - \frac{3\nu_0}{p_0}} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b|^{\frac{2p_0}{3}} dx \right)^{\frac{3\nu_0}{2p_0}} \right)^{\frac{1}{\nu_0}} \right) \\
& \leq \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} \sup_{\eta < r_0} \left(\eta^{\frac{2}{\nu_0} - \frac{3}{p_0}} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u|^{\frac{2p_0}{3}} dx \right)^{\frac{3}{2p_0}} \right) + \sup_{\eta < r_0} \left(\eta^{\frac{2}{\nu_0} - \frac{3}{p_0}} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b|^{\frac{2p_0}{3}} dx \right)^{\frac{3}{2p_0}} \right) \\
& \leq \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} \sup_{\eta < r_0} \left(\eta^{\frac{2}{\nu_0} - \frac{3}{p_0}} \|(u, b)(t)\|_{L^{p_0, \infty}(B_\eta(x_0))} \right) \\
& \leq C \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} r_0^{\frac{2}{\nu_0} - \frac{3}{p_0}} \|(u, b)(t)\|_{L^{p_0, \infty}(B_\eta(x_0))} \\
& \leq C \delta^* = \delta,
\end{aligned} \tag{3.8}$$

where δ is the same constant in Theorem 1.1 and we choose $\delta^* = \frac{\delta}{C}$. This implies

$$\left\| \left(\sup_{\eta \leq r_0^*} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t)|^{\nu_0} dx \right)^{\frac{1}{\nu_0}}, \sup_{\eta \leq r_0^*} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t)|^{\nu_0} dx \right)^{\frac{1}{\nu_0}} \right) \right\|_{L^{\mu_0, \infty}[t_0 - (r_0^*)^2, t_0]} \leq \delta \tag{3.9}$$

for some $r_0^* \leq r_0$. Due to facts of Theorem 1.1, we know that z_0 is a regular point, which is a contradiction. If (3.5) is true, by using Poincaré's inequality, we have

$$\begin{aligned}
& \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} \left(\sup_{\eta < r_0} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u - u_{x_0, \eta}|^{\nu_0} dx \right)^{\frac{1}{\nu_0}} + \sup_{\eta < r_0} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b - b_{x_0, \eta}|^{\nu_0} dx \right)^{\frac{1}{\nu_0}} \right) \\
& \leq C \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} \sup_{\eta < r_0} \left(\left(\eta^{2 - \frac{3\nu_0}{p_0} + \nu_0} \left(\int_{B_\eta(x_0)} |\nabla u|^{p_0} dx \right)^{\frac{\nu_0}{p_0}} \right)^{\frac{1}{\nu_0}} + \left(\eta^{2 - \frac{3\nu_0}{p_0} + \nu_0} \left(\int_{B_\eta(x_0)} |\nabla b|^{p_0} dx \right)^{\frac{\nu_0}{p_0}} \right)^{\frac{1}{\nu_0}} \right) \\
& \leq C \limsup_{t \rightarrow t_0} (t_0 - t)^{\frac{1}{\mu_0}} r_0^{\frac{2}{\nu_0} - \frac{3}{p_0} + 1} \|(\nabla u, \nabla b)\|_{L^{p_0}(B_{r_0}(x_0))} \\
& \leq C \delta^* = \delta,
\end{aligned} \tag{3.10}$$

which implies

$$\left\| \left(\sup_{\eta \leq r_0^*} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |u(x, t) - u_{x_0, \eta}|^{\nu_0} dx \right)^{\frac{1}{\nu_0}}, \sup_{\eta \leq r_0^*} \left(\frac{1}{\eta} \int_{B_\eta(x_0)} |b(x, t) - b_{x_0, \eta}|^{\nu_0} dx \right)^{\frac{1}{\nu_0}} \right) \right\|_{L^{\mu_0, \infty}[t_0 - (r_0^*)^2, t_0]} \leq \delta \tag{3.11}$$

for some $r_0^* \leq r_0$ and $\nu_0 \in \left\{ \begin{array}{l} [2, \infty], \quad p_0 \geq 3 \\ [2, \frac{2p_0}{3-p_0}], \quad \frac{3}{2} \leq p_0 < 3 \end{array} \right.$. By using Theorem 1.1, we deduce z_0 is a regular point, this is also a contradiction, hence concludes the proof of Theorem 1.2.

Acknowledgments

The authors are supported by the Construct Program of the Key Discipline in Hunan Province and NSFC Grant No. 11871209.

References

- [1] Albritton D, Barker T. Localised necessary conditions for singularity formation in the Navier-Stokes equations with curved boundary. *Journal of Differential Equations*, 2020, 269(9): 7529-7573.
- [2] Barker T, Prange C. Localized smoothing for the Navier–Stokes equations and concentration of critical norms near singularities. *Archive for Rational Mechanics and Analysis*, 2020: 1-55.
- [3] Cabannes H. *Theoretical Magnetofluidynamics*. Academic Press, New York and London, 1970.
- [4] Caffarelli L, Kohn R, Nirenberg L. Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Communications on Pure and Applied Mathematics*, 1982, 35(6): 771-831.
- [5] Cao C, Titi E. Global regularity criterion for the 3D Navier-Stokes equations involving one entry of the velocity gradient tensor. *Archive for Rational Mechanics and Analysis*, 2011, 202(3): 919-932.
- [6] Cao C, Wu J. Two regularity criteria for the 3D MHD equations. *Journal of Differential Equations*, 2010, 248(9): 2263-2274.
- [7] Choe H J, Lewis J L. On the singular set in the Navier-Stokes equations. *Journal of Functional Analysis*, 2000, 175(2): 348-369.
- [8] Chen C, Strain R, Yau H, et al. Lower bound on the blow-up rate of the axisymmetric Navier-Stokes equations. *International Mathematics Research Notices*, 2008.
- [9] Chen C, Strain R, Tsai T, et al. Lower bounds on the blow-up rate of the axisymmetric Navier-Stokes equations II. *Communications in Partial Differential Equations*, 2009, 34(3): 203-232.
- [10] Da Veiga H B. A new regularity class for the Navier-Stokes equations in \mathbb{R}^n . *Chinese Ann. Math. Ser. B*, 1995, 16(4): 407-412.
- [11] Duvaut G, Lions J L. Inéquations en thermoélasticité et magnétohydrodynamique. *Archive for Rational Mechanics and Analysis*, 1972, 46(4): 241-279.
- [12] Escauriaza L, Seregin G, Sverak V. $L_{3,\infty}$ -solutions of the Navier-Stokes equations and backward uniqueness. *Russian Mathematical Surveys*, 2003, 58(2): 211-250.
- [13] He C, Wang Y. Limiting case for the regularity criterion of the Navier-Stokes equations and the magnetohydrodynamic equations. *Science China Mathematics*, 2010, 53(7): 1767-1774.
- [14] He C, Xin Z. On the regularity of weak solutions to the magnetohydrodynamic equations. *Journal of Differential Equations*, 2005, 213(2): 235-254.
- [15] He C, Xin Z. Partial regularity of suitable weak solutions to the incompressible magnetohydrodynamic equations. *Journal of Functional Analysis*, 2005, 227(1): 113-152.
- [16] Hmidi T, Keraani S. Remarks on the blowup for the L^2 -critical nonlinear Schrödinger equations. *Siam Journal on Mathematical Analysis*, 2006, 38(4): 1035-1047.

- [17] Holmer J, Roudenko S. On blow-up solutions to the 3D cubic nonlinear Schrödinger equation. Applied Mathematics Research eXpress, 2007, 2007.
- [18] Kang K, Lee J. Interior regularity criteria for suitable weak solutions of the magnetohydrodynamic equations. Journal of Differential Equations, 2009, 247(8): 2310-2330.
- [19] Kang K, Kim J. Regularity criteria of the magnetohydrodynamic equations in bounded domains or a half space. Journal of Differential Equations, 2012, 253(2): 764-794.
- [20] Kim H, Kozono H. Interior regularity criteria in weak spaces for the Navier-Stokes equations. Manuscripta Mathematica, 2004, 115(1): 85-100.
- [21] Koch G, Nadirashvili N, Seregin G, et al. Liouville theorems for the Navier-Stokes equations and applications. Acta Mathematica, 2009, 203(1): 83-105.
- [22] Leray J. Sur le mouvement d'un liquide visqueux emplissant l'espace. Acta mathematica, 1934, 63: 193-248.
- [23] Lin H, Du L. Regularity criteria for incompressible magnetohydrodynamics equations in three dimensions. Nonlinearity, 2012, 26(1): 219.
- [24] Lin F. A new proof of the Caffarelli-Kohn-Nirenberg theorem. Communications on Pure and Applied Mathematics, 1998, 51(3): 241-257.
- [25] Li K, Ozawa T, Wang B. Dynamical behavior for the solutions of the Navier-Stokes equation. arXiv preprint arXiv:1608.06680, 2016.
- [26] Merle F, Tsutsumi Y. L^2 concentration of blow-up solutions for the nonlinear Schrödinger equation with critical power nonlinearity. Journal of Differential Equations, 1990, 84(2): 205-214.
- [27] Miller E. A regularity criterion for the Navier-Stokes equation involving only the middle eigenvalue of the strain tensor. Archive for Rational Mechanics and Analysis, 2020, 235(1): 99-139.
- [28] Maekawa Y, Miura H, Prange C. Estimates for the Navier-Stokes equations in the half-space for nonlocalized data. Analysis & PDE, 2020, 13(4): 945-1010.
- [29] Prodi G. Un teorema di unicità per le equazioni di Navier-Stokes. Annali di Matematica Pura ed Applicata, 1959, 48(1): 173-182.
- [30] Sermange M, Temam R. Some mathematical questions related to the MHD equations. Communications on Pure and Applied Mathematics, 1983, 36(5): 635-664.
- [31] Scheffer V. Hausdorff measure and the Navier-Stokes equations. Communications in Mathematical Physics, 1977, 55(2): 97-112.
- [32] Scheffer V. The Navier-Stokes equations in space dimension four. Communications in Mathematical Physics, 1978, 61(1): 41-68.
- [33] Scheffer V. The Navier-Stokes equations on a bounded domain. Communications in Mathematical Physics, 1980, 73(1): 1-42.
- [34] Serrin J. On the interior regularity of weak solutions of the Navier-Stokes equations. Archive for Rational Mechanics and Analysis, 1962, 9(1): 187-195.
- [35] G. Seregin. A certain necessary condition of potential blow up for Navier-Stokes equations. Communications in Mathematical Physics, 2012, 312(3):833-845.

- [36] Tan W, Yin Z. The energy conservation and regularity for the Navier-Stokes equations. arXiv preprint arXiv:2107.04157, 2021.
- [37] Tan W. The localized characterization for the singularity formation in the Navier-Stokes equations. arXiv preprint arXiv:2107.04597, 2021.
- [38] Tao T. Quantitative bounds for critically bounded solutions to the Navier-Stokes equations. arXiv preprint arXiv:1908.04958, 2019.
- [39] Wang W, Zhang Z. On the interior regularity criteria and the number of singular points to the Navier-Stokes equations. *Journal d'Analyse Mathématique*, 2014, 123(1): 139-170.
- [40] Wang W, Zhang Z. On the interior regularity criteria for suitable weak solutions of the magnetohydrodynamics equations. *SIAM Journal on Mathematical Analysis*, 2013, 45(5): 2666-2677.
- [41] Wu F. Conditional regularity for the 3D Navier-Stokes equations in terms of the middle eigenvalue of the strain tensor. *Evolution Equations & Control Theory*, 2021, 10(3): 511-518.
- [42] Zhang Z, Chen Q. Regularity criterion via two components of vorticity on weak solutions to the Navier-Stokes equations in \mathbb{R}^3 . *Journal of Differential Equations*, 2005, 216(2): 470-481.