

THE SECOND BOGOLYUBOV THEOREM AND GLOBAL AVERAGING PRINCIPLE FOR SPDES WITH MONOTONE COEFFICIENTS

MENGYU CHENG AND ZHENXIN LIU

ABSTRACT. In this paper, we establish the second Bogolyubov theorem and global averaging principle for stochastic partial differential equations (in short, SPDEs) with monotone coefficients. Firstly, we prove that there exists a unique L^2 -bounded solution to SPDEs with monotone coefficients and this bounded solution is globally asymptotically stable in square-mean sense. Then we show that the L^2 -bounded solution possesses the same recurrent properties (e.g. periodic, quasi-periodic, almost periodic, almost automorphic, Birkhoff recurrent, Levitan almost periodic, etc.) in distribution sense as the coefficients. Thirdly, we prove that the recurrent solution of the original equation converges to the stationary solution of averaged equation under the compact-open topology as the time scale goes to zero — in other words, there exists a unique recurrent solution to the original equation in a neighborhood of the stationary solution of averaged equation when the time scale is small. Finally, we establish the global averaging principle in weak sense, i.e. we show that the attractor of original system tends to that of the averaged equation in probability measure space as the time scale goes to zero. For illustration of our results, we give two applications, including stochastic reaction diffusion equations and stochastic generalized porous media equations.

1. INTRODUCTION

Averaging principle is an effective method for studying dynamical systems with highly oscillating components. Under suitable conditions, the highly oscillating components can be “averaged out” to produce an averaged system. The averaged system is easier for analysis and governs the evolution of the original system over long time scales.

Consider the following deterministic systems in \mathbb{R}^n , $n \in \mathbb{N}$:

$$(1.1) \quad \dot{X}^\varepsilon = F\left(\frac{t}{\varepsilon}, X^\varepsilon\right)$$

and

$$(1.2) \quad \dot{X} = \bar{F}(X)$$

for small parameter $0 < \varepsilon \ll 1$, where $F \in C(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$, and $\bar{F}(X) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T F(t, X) dt$.

It is a basic problem of averaging principle to determine in what sense the behavior of solutions to the averaged system (1.2) approximates the behavior of solutions to the non-autonomous system (1.1) as the time scale ε goes to zero. For the connotation of approximation, there are three natural types of interpretation. One is the so-called *first Bogolyubov theorem*, i.e. the convergence of the solution of the original Cauchy problem (1.1) to that of the averaged equation (1.2) on a finite interval $[0, T]$ when the initial data are such that $X^\varepsilon(0) = X(0)$. And another one is to request that the approximation be valid on the entire real axis, which is the so-called *second Bogolyubov theorem* (sometimes called “theorem

Date: August 31, 2021.

2010 Mathematics Subject Classification. 70K65, 60H15, 37B20, 37L15.

Key words and phrases. Second Bogolyubov theorem; Global averaging principle; Monotone SPDEs; Periodic solutions; Quasi-periodic solutions; Almost periodic solutions; Birkhoff recurrent solutions; Poisson stable solutions; Stochastic reaction diffusion equations; Stochastic generalized porous media equations.

for periodic solution by averaging"). In addition, it is meaningful to determine whether the attractor of the averaged equation (1.2) approximates the attractor of the original equation (1.1). One calls this result the *global averaging principle*.

The idea of averaging dates back to the perturbation theory which was proposed by Clairaut, Laplace and Lagrange in the 18th century. Then fairly rigorous averaging method for nonlinear oscillations was presented by Krylov, Bogolyubov and Mitropolsky [26, 2], which is called the Krylov-Bogolyubov method nowadays. After that, there is a lot of works on averaging for deterministic finite and infinite dimensional systems, which we will not mention here.

Meanwhile, Stratonovich firstly proposed the stochastic averaging method on the basis of physical considerations, which was later proved mathematically by Khasminskii. Then extensive investigations concerning averaging principle for stochastic differential equations were conducted, following Khasminskii's mathematically pioneering work [23]; see, e.g. [1, 4, 5, 6, 13, 15, 16, 18, 24, 28, 30, 32, 39, 40, 41, 42, 43] and the references therein. Note that the above existing results are concerned with the first Bogolyubov theorem.

Despite considerable advances in this direction, there are few works on stochastic averaging concerning with the second Bogolyubov theorem, which states: there exists a unique periodic solution to the original equation in a neighborhood of the stationary solution of averaged equation. Consider the following SPDEs on a separable Hilbert space $(H, \langle \cdot, \cdot \rangle)$

$$(1.3) \quad dX_\varepsilon(t) = \left(A(X_\varepsilon(t)) + F\left(\frac{t}{\varepsilon}, X_\varepsilon(t)\right) \right) dt + G\left(\frac{t}{\varepsilon}, X_\varepsilon(t)\right) dW(t),$$

where A satisfies some monotone condition, F and G are Lipschitz in second variable. Here W is a two-sided cylindrical Wiener process defined on another separable Hilbert space U and $0 < \varepsilon \leq 1$. Recall that the second Bogolyubov theorem for stochastic differential equations with almost periodic coefficients was studied in [22], and [9] investigated the averaging principle for stochastic ordinary differential equations with general recurrent coefficients. As discussed in [22] and [9], equations are semilinear with globally Lipschitz and linear growth nonlinear terms, which cannot cover the monotone case. However, the coefficients of many interesting practical models just satisfy monotone conditions. Some typical examples are reaction diffusion equations and porous media equations.

Generally, reaction diffusion equations can be used to describe the growth of biological population and the spatial spread of epidemic diseases, which are largely affected by time-varying environment. In particular, the recurrent phenomenon has been found in the growth of population and the spread of diseases, since some regular environmental changes such as seasonal changes. And porous media equations appear in the description of different natural phenomenon related to diffusion, filtration or heat propagation. Since the noise models the small irregular fluctuations generated by microscopic effects, it is more practical to consider the above systems perturbed by white noise.

From the perspective of theoretical and practical value, we establish the second Bogolyubov theorem for SPDEs with monotone coefficients in this paper. More specifically, consider equation (1.3), we assume that A is strongly monotone. Compared with the assumption that A is a linear bounded operator in [9], this condition admits wider applications. It includes unbounded linear operators and quasi-linear operators.

Denoting by $F_\varepsilon(t, x) := F(\frac{t}{\varepsilon}, x)$ and $G_\varepsilon(t, x) := G(\frac{t}{\varepsilon}, x)$, we transform equation (1.3) to

$$(1.4) \quad dX_\varepsilon(t) = (A(X_\varepsilon(t)) + F_\varepsilon(t, X_\varepsilon(t))) dt + G_\varepsilon(t, X_\varepsilon(t)) dW(t).$$

In this paper, we firstly show that there exists a unique L^2 -bounded solution $X_\varepsilon(t), t \in \mathbb{R}$ of (1.4) which shares the same recurrent properties (in particular, periodic, quasi-periodic, almost periodic, almost automorphic, Birkhoff recurrent, Levitan almost periodic, almost recurrent, pseudo-periodic, pseudo-recurrent, Poisson stable) in distribution sense as the

coefficients for each $0 < \varepsilon \leq 1$. And we prove that this L^2 -bounded solution $X_\varepsilon(t), t \in \mathbb{R}$ to (1.4) is globally asymptotically stable in square-mean sense. Without loss of generality, we assume $\varepsilon = 1$ in this part, then the L^2 -bounded solution is denoted by $X(t), t \in \mathbb{R}$. Note that coefficient F in (1.4) need not to be Lipschitz in this part. This result is interesting on its own rights. To our knowledge, there are only a few works on general recurrent solutions to SPDEs in a unified framework, see [8, 29]. As discussed in [8] and [29], they dealt with recurrent solutions to semilinear SPDEs with Lipschitz continuous and globally linear growth nonlinearities according to Shcherbakov's comparability method by character of recurrence. B. A. Shcherbakov gave the existence condition of at least one (or exactly one) solution to deterministic equation with the same character of recurrence as the coefficient. This solution is said to be compatible (respectively, uniformly compatible). Comparing to [8] and [29], we consider SPDEs with monotone coefficients.

Let X_ε be the recurrent solution to equation (1.4). Then one of the major aims of this paper is to prove that

$$(1.5) \quad \lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X})) = 0 \quad \text{in } Pr(C(\mathbb{R}, H))$$

(see Theorem 4.5 and Corollary 4.6), where \bar{X} is the unique stationary solution of the following averaged equation

$$(1.6) \quad dX(t) = (A(X(t)) + \bar{F}(X(t))) dt + \bar{G}(X(t))dW(t).$$

Here $\bar{F} \in C(H, H), \bar{G} \in C(H, L_2(U, H)), \bar{F}$ and \bar{G} satisfy

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} F(s, x) ds = \bar{F}(x), \quad \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \|G(s, x) - \bar{G}(x)\|_{L_2(U, H)}^2 ds = 0$$

uniformly with respect to $t \in \mathbb{R}$.

This averaging principle is also applicable to the following system

$$(1.7) \quad dX(t) = \varepsilon (A(X(t)) + F(t, X(t))) dt + \sqrt{\varepsilon} G(t, X(t))dW(t),$$

where $0 < \varepsilon \leq 1$. With the time scaling $t \mapsto \frac{t}{\varepsilon}$, denote by $\Phi_\varepsilon(t) := X(\frac{t}{\varepsilon})$ and $W_\varepsilon(t) := \sqrt{\varepsilon} W(\frac{t}{\varepsilon})$ for all $t \in \mathbb{R}$, we transform equation (1.7) to

$$(1.8) \quad d\Phi_\varepsilon(t) = (A(\Phi_\varepsilon(t)) + F_\varepsilon(t, \Phi_\varepsilon(t))) dt + G_\varepsilon(t, \Phi_\varepsilon(t))dW_\varepsilon(t).$$

Then we can consider the following equation

$$(1.9) \quad d\tilde{X}_\varepsilon(t) = (A(\tilde{X}_\varepsilon(t)) + F_\varepsilon(t, \tilde{X}_\varepsilon(t))) dt + G_\varepsilon(t, \tilde{X}_\varepsilon(t))dW(t).$$

It is obvious that $\mathcal{L}(\tilde{X}_\varepsilon(t)) = \mathcal{L}(\Phi_\varepsilon(t))$ for any $t \in \mathbb{R}$.

In contrast to the first Bogolyubov averaging principle for Cauchy problem of stochastic differential equations on finite intervals, we prove that there exists a unique recurrent solution in a small neighborhood of the stationary solution to the averaged equation when the time scale is small. Note that it is non-initial value problem, and this recurrent solution is more general than the classical second Bogolyubov theorem which only treats the periodic case.

Since the SPDEs we concern in this paper are not semilinear, the semigroup framework in [22] and [9] is not applicable to our problem. Therefore, a difficulty that we face is how to deal with the monotone SPDEs. Firstly, under some suitable conditions, employing the technique of truncation which is used in [4, 5, 6, 28], we show that

$$\lim_{\varepsilon \rightarrow 0} E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 = 0$$

for all $s \in \mathbb{R}$ and $T > 0$ provided $\lim_{\varepsilon \rightarrow 0} E \|\zeta_s^\varepsilon - \zeta_s\|^2 = 0$, where Y_ε is the solution of (1.4) with the initial condition $Y_\varepsilon(s) = \zeta_s^\varepsilon$ and \bar{Y} is the solution of (1.6) with initial condition $\bar{Y}(s) = \zeta_s$

(see Theorem 4.3). In fact, this is the first Bogolyubov theorem, which is new despite that there have already been many results in this direction mentioned above.

In view of Theorem 4.3, tightness of family of measures $\{\mathbb{P} \circ [X_\varepsilon(t)]^{-1}\}_{\varepsilon \in (0,1]}$ for any $t \in \mathbb{R}$ plays an important role in establishing the second Bogolyubov theorem. Although the tightness of $\{\mathbb{P} \circ [X_\varepsilon(t)]^{-1}\}_{\varepsilon \in (0,1]}$ on $C([0, T]; H)$ was proved by using Ascoli-Arzelà theorem and the Garcia-Rademich-Rumsey theorem in [4, 5, 6], it is different from the technique used in our paper. We find that

$$(1.10) \quad \sup_{t \in \mathbb{R}} E \|X_\varepsilon(t)\|_S^2 < \infty$$

uniformly with respect to $\varepsilon \in (0, 1]$. Therefore, for any $t \in \mathbb{R}$, the tightness of $\{\mathbb{P} \circ [X_\varepsilon(t)]^{-1}\}_{\varepsilon \in (0,1]}$ is a consequence of the compactness of the inclusion $S \subset H$.

Another major result in present paper is to establish the global averaging principle in weak sense. Namely, we prove that uniform attractor of original system tends to uniform attractor of averaged equation in probability measure space $Pr(H)$. Global averaging of deterministic systems was conducted, see e.g. [19, 20, 21, 44] among others. But to our knowledge, there is no work so far on global averaging for stochastic equations. Notice that it is still an open problem whether solutions of general SPDEs can generate random dynamical systems (in short, RDS). So we consider attractors in the probability measure space $Pr(H)$ instead of pullback attractors in the framework of RDS. That is why we call it “in weak sense”.

Let $Pr_2(H)$ be a subspace of $Pr(H)$ such that

$$\int_H \|z\|^2 \mu(dz) < +\infty$$

for any $\mu \in Pr_2(H)$. With the transition probability $P_{\mathbb{F}}(s, x, t, dy) := \mathbb{P} \circ (X(t, s, x))^{-1}(dy)$ to equation (1.4) when $\varepsilon = 1$, we associate a mapping $P^*(t, \mathbb{F}, \cdot) : Pr(H) \rightarrow Pr(H)$ defined by

$$P^*(t, \mathbb{F}, \mu)(B) := \int_H P_{\mathbb{F}}(0, x, t, B) \mu(dx)$$

for all $\mu \in Pr(H)$, $B \in \mathcal{B}(H)$ and $\mathbb{F} := (F, G)$. Firstly, we show that P^* is a cocycle over $(H(\mathbb{F}), \mathbb{R}, \sigma)$ with fiber $Pr_2(H)$, where $(H(\mathbb{F}), \mathbb{R}, \sigma)$ is a shift dynamical system (see Section 5 for details). Suppose that $H(\mathbb{F})$ is compact. Then we prove that P_ε^* associated with (1.4) has a uniform attractor \mathcal{A}^ε in $Pr_2(H)$ for any $0 < \varepsilon \leq 1$, and

$$\lim_{\varepsilon \rightarrow 0} \text{dist}_{Pr_2(H)}(\mathcal{A}^\varepsilon, \bar{\mathcal{A}}) = 0$$

(see Theorem 5.14), where $\text{dist}_{Pr_2(H)}$ is the Hausdorff semi-metric and $\bar{\mathcal{A}} := \{\mathcal{L}(\bar{X}(0))\}$ is the attractor of \bar{P}^* to the averaged equation (1.6). Note that $H(\mathbb{F})$ is compact provided \mathbb{F} is Birkhoff recurrent.

The remainder of this paper is organized as follows. In the next section, we recall some definitions and facts concerning dynamical systems, Poisson stable (or recurrent) functions, Shcherbakov’s comparability method by character of recurrence and variational approach. In the third section, we show that there exists a unique L^2 -bounded solution which possesses the same recurrent properties in distribution sense as the coefficients and this bounded solution is globally asymptotically stable in square-mean sense. In section 4, we establish the second Bogolyubov theorem for SPDEs with monotone coefficients. In section 5, we prove the global averaging principle for these SPDEs. In the last section, we illustrate our theoretical results by stochastic reaction diffusion equations and stochastic generalized porous media equations.

2. PRELIMINARIES

In this section, we introduce some useful preliminaries, including dynamical systems, poisson stable functions, Shcherbakov's comparability method by character of recurrence, variational approach.

2.1. Shift dynamical systems. In this subsection, let (\mathcal{X}, ρ) be a complete metric space and $(\mathcal{X}, \mathbb{R}, \pi)$ be a dynamical system (flow) on \mathcal{X} , i.e. the mapping $\pi : \mathbb{R} \times \mathcal{X} \rightarrow \mathcal{X}$ is continuous, $\pi(0, x) = x$ and $\pi(t + s, x) = \pi(t, \pi(s, x))$ for any $x \in \mathcal{X}$ and $t, s \in \mathbb{R}$. We write $C(\mathbb{R}, \mathcal{X})$ to mean the space of all continuous functions $\varphi : \mathbb{R} \rightarrow \mathcal{X}$ equipped with the distance

$$d(\varphi_1, \varphi_2) := \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{d_k(\varphi_1, \varphi_2)}{1 + d_k(\varphi_1, \varphi_2)},$$

where

$$d_k(\varphi_1, \varphi_2) := \sup_{|t| \leq k} \rho(\varphi_1(t), \varphi_2(t)),$$

which generates the compact-open topology on $C(\mathbb{R}, \mathcal{X})$. The space $(C(\mathbb{R}, \mathcal{X}), d)$ is a complete metric space (see, e.g. [33, 35, 37, 38]).

Remark 2.1. Let $\{\varphi_n\}_{n=1}^{\infty}, \varphi \in C(\mathbb{R}, \mathcal{X})$. Then the following statements are equivalent.

- (i) $\lim_{n \rightarrow \infty} d(\varphi_n, \varphi) = 0$.
- (ii) $\lim_{n \rightarrow \infty} \max_{|t| \leq l} \rho(\varphi_n(t), \varphi(t)) = 0$ for any $l > 0$.
- (iii) There exists a sequence $l_n \rightarrow +\infty$ such that $\lim_{n \rightarrow \infty} \max_{|t| \leq l_n} \rho(\varphi_n(t), \varphi(t)) = 0$.

Let us now consider two examples of shift dynamical systems which we will use in this paper.

Example 2.2. We say φ^τ is the τ -translation of φ if $\varphi^\tau(t) := \varphi(t + \tau)$ for any $t \in \mathbb{R}$ and $\varphi \in C(\mathbb{R}, \mathcal{X})$. For any $(\tau, \varphi) \in \mathbb{R} \times C(\mathbb{R}, \mathcal{X})$, the mapping $\sigma : \mathbb{R} \times C(\mathbb{R}, \mathcal{X}) \rightarrow C(\mathbb{R}, \mathcal{X})$ is defined by $\sigma(\tau, \varphi) := \varphi^\tau$. Then the triplet $(C(\mathbb{R}, \mathcal{X}), \mathbb{R}, \sigma)$ is a dynamical system which is called *shift dynamical system* or *Bebutov's dynamical system*. Indeed, it is easy to check that $\sigma(0, \varphi) = \varphi$ and $\sigma(\tau_1 + \tau_2, \varphi) = \sigma(\tau_2, \sigma(\tau_1, \varphi))$ for any $\varphi \in C(\mathbb{R}, \mathcal{X})$ and $\tau_1, \tau_2 \in \mathbb{R}$. And it can be proved that the mapping $\sigma : \mathbb{R} \times C(\mathbb{R}, \mathcal{X}) \rightarrow C(\mathbb{R}, \mathcal{X})$ is continuous, see, e.g. [7, 33, 35, 38].

In what follows, let (\mathcal{Y}, ρ_1) be a complete metric space. We employ $H(\varphi)$ to denote the hull of φ , which is the set of all the limits of φ^{τ_n} in $C(\mathbb{R}, \mathcal{X})$, i.e.

$$H(\varphi) := \{\psi \in C(\mathbb{R}, \mathcal{X}) : \psi = \lim_{n \rightarrow \infty} \varphi^{\tau_n} \text{ for some sequence } \{\tau_n\} \subset \mathbb{R}\}.$$

Notice that the set $H(\varphi) \subset C(\mathbb{R}, \mathcal{X})$ is closed and translation invariant. Consequently, it naturally defines on $H(\varphi)$ a shift dynamical system $(H(\varphi), \mathbb{R}, \sigma)$. Now we give the second example, which is similar to Section 2.4 in [8].

Example 2.3. We write $BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ to mean the space of all continuous functions $f : \mathbb{R} \times \mathcal{X} \rightarrow \mathcal{Y}$ which satisfy the following conditions:

- (i) f is bounded on every bounded subset from $\mathbb{R} \times \mathcal{X}$;
- (ii) f is continuous in $t \in \mathbb{R}$ uniformly with respect to x on each bounded subset $Q \subset \mathcal{X}$.

We endow $BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ with the following d metric

$$(2.1) \quad d(f, g) := \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{d_k(f, g)}{1 + d_k(f, g)},$$

where $d_k(f, g) := \sup_{|t| \leq k, x \in Q_k} \rho_1(f(t, x), g(t, x))$. Here $Q_k \subset \mathcal{X}$ is bounded, $Q_k \subset Q_{k+1}$ and $\cup_{k \in \mathbb{N}} Q_k = \mathcal{X}$. Note that d generates the topology of uniform convergence on bounded subsets on $BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ and $(BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y}), d)$ is a complete metric space.

Given $f \in BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ and $\tau \in \mathbb{R}$. We write f^τ to mean the τ -translation of f if $f^\tau(t, x) := f(t + \tau, x)$ for all $(t, x) \in \mathbb{R} \times \mathcal{X}$. It is proved that $BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ is invariant with respect to translations. Like in Example 2.2, we define a mapping $\sigma : \mathbb{R} \times BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y}) \rightarrow BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$, $(\tau, f) \mapsto f^\tau$. Then it can be proved that the triplet $(BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y}), \mathbb{R}, \sigma)$ is a dynamical system. See Chapter I in [7] for details. Given $f \in BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$, $H(f) \subset BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ is closed and translation invariant. Consequently, it naturally defines on $H(f)$ a shift dynamical system $(H(f), \mathbb{R}, \sigma)$.

We employ $BC(\mathcal{X}, \mathcal{Y})$ to denote the space of all continuous functions $f : \mathcal{X} \rightarrow \mathcal{Y}$ which are bounded on every bounded subset of \mathcal{X} and equipped with the distance

$$d(f, g) := \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{d_k(f, g)}{1 + d_k(f, g)},$$

where $d_k(f, g) := \sup_{x \in Q_k} \rho_1(f(x), g(x))$, Q_k is similar to that in Example 2.3. Note that $(BC(\mathcal{X}, \mathcal{Y}), d)$ is a complete metric space. For any $F \in BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$, the mapping $\mathcal{F} : \mathbb{R} \rightarrow BC(\mathcal{X}, \mathcal{Y})$ is defined by $\mathcal{F}(t) := F(t, \cdot) : \mathcal{X} \rightarrow \mathcal{Y}$. Clearly, $\mathcal{F} \in C(\mathbb{R}, BC(\mathcal{X}, \mathcal{Y}))$.

Remark 2.4. It can be proved that the following statements are true.

- (i) The mapping $h : BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y}) \rightarrow C(\mathbb{R}, BC(\mathcal{X}, \mathcal{Y}))$ defined by equality $h(F) := \mathcal{F}$ establishes an isometry between $BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ and $C(\mathbb{R}, BC(\mathcal{X}, \mathcal{Y}))$.
- (ii) $h(F^\tau) = \mathcal{F}^\tau$ for any $\tau \in \mathbb{R}$ and $F \in BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$, i.e. the shift dynamical systems $(BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y}), \mathbb{R}, \sigma)$ and $(C(\mathbb{R}, BC(\mathcal{X}, \mathcal{Y})), \mathbb{R}, \sigma)$ are (dynamically) homeomorphic.

2.2. Poisson stable functions. Let us recall the types of Poisson stable (or recurrent) functions to be studied in this paper; For further details and the relations among these types of functions, see [33, 35, 37, 38].

Definition 2.5. We say that a function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *T-periodic*, if there exists a constant $T \in \mathbb{R}$ such that $\varphi(t + T) = \varphi(t)$ for all $t \in \mathbb{R}$. In particular, φ is called *stationary* provided $\varphi(t) = \varphi(0)$ for all $t \in \mathbb{R}$.

Definition 2.6. A function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is called *Bohr almost periodic* if the set $\mathcal{T}(\varphi, \varepsilon)$ of ε -almost periods of φ is relatively dense for each $\varepsilon > 0$, i.e. for each $\varepsilon > 0$ there exists a constant $l = l(\varepsilon) > 0$ such that $\mathcal{T}(\varphi, \varepsilon) \cap [a, a + l] \neq \emptyset$ for all $a \in \mathbb{R}$, where

$$\mathcal{T}(\varphi, \varepsilon) := \left\{ \tau \in \mathbb{R} : \sup_{t \in \mathbb{R}} \rho(\varphi(t + \tau), \varphi(t)) < \varepsilon \right\},$$

and $\tau \in \mathcal{T}(\varphi, \varepsilon)$ is called ε -almost period of φ .

Definition 2.7. We say that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *pseudo-periodic in the positive* (respectively, *negative*) *direction* if for each $\varepsilon > 0$ and $l > 0$ there exists a ε -almost period $\tau > l$ (respectively, $\tau < -l$) of the function φ . The function φ is called *pseudo-periodic* if it is pseudo-periodic in both directions.

Definition 2.8. A function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is called *almost recurrent (in the sense of Bebutov)* if the set $\mathfrak{T}(\varphi, \varepsilon)$ is relatively dense for every $\varepsilon > 0$, where $\mathfrak{T}(\varphi, \varepsilon) := \{\tau \in \mathbb{R} : d(\varphi^\tau, \varphi) < \varepsilon\}$. And $\tau \in \mathfrak{T}(\varphi, \varepsilon)$ is said to be ε -shift for φ .

Definition 2.9. (i) We say that a function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *Lagrange stable* provided $\{\varphi^h : h \in \mathbb{R}\}$ is a relatively compact subset of $C(\mathbb{R}, \mathcal{X})$.

- (ii) We say that a function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *Birkhoff recurrent* if it is almost recurrent and Lagrange stable.

Definition 2.10. We say that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *Poisson stable in the positive* (respectively, *negative*) *direction* if for every $\varepsilon > 0$ and $l > 0$ there exists $\tau > l$ (respectively, $\tau < -l$) such that $d(\varphi^\tau, \varphi) < \varepsilon$. The function φ is called *Poisson stable* if it is Poisson stable in both directions.

Definition 2.11. We say that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *Levitan almost periodic* if there exists an almost periodic function $\psi \in C(\mathbb{R}, \mathcal{Y})$ such that for any $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon) > 0$ such that $\mathcal{T}(\psi, \delta) \subseteq \mathfrak{T}(\varphi, \varepsilon)$.

Definition 2.12. We say that a function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *almost automorphic* if it is Levitan almost periodic and Lagrange stable.

Definition 2.13. We say that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is *quasi-periodic with the spectrum of frequencies* $\nu_1, \nu_2, \dots, \nu_k$ if it satisfies the following conditions:

- (i) the numbers $\nu_1, \nu_2, \dots, \nu_k$ are rationally independent;
- (ii) there exists a continuous function $\Phi : \mathbb{R}^k \rightarrow \mathcal{X}$ such that $\Phi(t_1 + 2\pi, t_2 + 2\pi, \dots, t_k + 2\pi) = \Phi(t_1, t_2, \dots, t_k)$ for all $(t_1, t_2, \dots, t_k) \in \mathbb{R}^k$;
- (iii) $\varphi(t) = \Phi(\nu_1 t, \nu_2 t, \dots, \nu_k t)$ for $t \in \mathbb{R}$.

Let $\varphi \in C(\mathbb{R}, \mathcal{X})$. We employ \mathfrak{N}_φ (respectively, \mathfrak{M}_φ) to denote the family of all sequences $\{t_n\} \subset \mathbb{R}$ such that $\varphi^{t_n} \rightarrow \varphi$ (respectively, $\{\varphi^{t_n}\}$ converges) in $C(\mathbb{R}, \mathcal{X})$ as $n \rightarrow \infty$.

Definition 2.14. ([34, 35, 37]) A function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is called *pseudo-recurrent* if for any $\varepsilon > 0$ and $l \in \mathbb{R}$ there exists $L \geq l$ such that for any $\tau_0 \in \mathbb{R}$ we can find a number $\tau \in [l, L]$ satisfying

$$\sup_{|t| \leq 1/\varepsilon} \rho(\varphi(t + \tau_0 + \tau), \varphi(t + \tau_0)) \leq \varepsilon.$$

Remark 2.15. ([34, 35, 37, 38])

- (i) Every Birkhoff recurrent function is pseudo-recurrent, but not vice versa.
- (ii) Suppose that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is pseudo-recurrent, then every function $\psi \in H(\varphi)$ is pseudo-recurrent.
- (iii) Suppose that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is Lagrange stable and every function $\psi \in H(\varphi)$ is Poisson stable, then φ is pseudo-recurrent.

Finally, we remark that a Lagrange stable function is not Poisson stable in general, but all other types of functions introduced above are Poisson stable.

Definition 2.16. (i) We say that a function $\varphi \in C(\mathbb{R}, \mathcal{X})$ *possesses the property A* if the motion $\sigma(\cdot, \varphi)$ through φ with respect to the Bebutov dynamical system $(C(\mathbb{R} \times \mathcal{X}), \mathbb{R}, \sigma)$ possesses the property A.
(ii) Similarly, we say that $F \in BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ *possesses the property A in $t \in \mathbb{R}$ uniformly with respect to x on each bounded subset $Q \subset \mathcal{X}$* , if the motion $\sigma(\cdot, F) : \mathbb{R} \rightarrow BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y})$ through F with respect to the Bebutov dynamical system $(BUC(\mathbb{R} \times \mathcal{X}, \mathcal{Y}), \mathbb{R}, \sigma)$ possesses the property A.

Here the property A may be stationary, periodic, Bohr/Levitan almost periodic, etc.

2.3. Shcherbakov's comparability method by character of recurrence.

Definition 2.17. A function $\varphi \in C(\mathbb{R}, \mathcal{X})$ is called *comparable* (respectively, *strongly comparable*) *by character of recurrence* with $\psi \in C(\mathbb{R}, \mathcal{Y})$ provided $\mathfrak{N}_\psi \subseteq \mathfrak{N}_\varphi$ (respectively, $\mathfrak{M}_\psi \subseteq \mathfrak{M}_\varphi$).

Theorem 2.18. ([35, ChII], [36])

- (i) $\mathfrak{M}_\psi \subseteq \mathfrak{M}_\varphi$ implies $\mathfrak{N}_\psi \subseteq \mathfrak{N}_\varphi$, and hence strong comparability implies comparability.
- (ii) Assume that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is comparable by character of recurrence with $\psi \in C(\mathbb{R}, \mathcal{Y})$. If the function ψ is stationary (respectively, T -periodic, Levitan almost periodic, almost recurrent, Poisson stable), then so is φ .
- (iii) Assume that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is strongly comparable by character of recurrence with $\psi \in C(\mathbb{R}, \mathcal{Y})$. If the function ψ is quasi-periodic with the spectrum of frequencies $\nu_1, \nu_2, \dots, \nu_k$ (respectively, almost periodic, almost automorphic, Birkhoff recurrent, Lagrange stable), then so is φ .
- (iv) Assume that $\varphi \in C(\mathbb{R}, \mathcal{X})$ is strongly comparable by character of recurrence with $\psi \in C(\mathbb{R}, \mathcal{Y})$. And suppose further that ψ is Lagrange stable. If ψ is pseudo-periodic (respectively, pseudo-recurrent), then so is φ .

2.4. Variational approach. Recall that H is a separable Hilbert space with norm $\|\cdot\|_H$ and inner product $\langle \cdot, \cdot \rangle_H$, and that H^* is the dual space of H . Let $(V, \|\cdot\|_V)$ be a reflexive Banach space such that $V \subset H$ continuously and densely. So we have $H^* \subset V^*$ continuously and densely. Identifying H with its dual H^* via the Riesz isomorphism, then we have

$$V \subset H \subset V^*$$

continuously and densely. We write ${}_{V^*}\langle \cdot, \cdot \rangle_V$ to denote the pairing between V^* and V . It follows that

$${}_{V^*}\langle h, v \rangle_V = \langle h, v \rangle_H$$

for all $h \in H, v \in V$. (V, H, V^*) is called *Gelfand triple*. Since $H \subset V^*$ continuously and densely, we deduce that V^* is separable, hence so is V .

Assume that $(V_1, \|\cdot\|_{V_1})$ and $(V_2, \|\cdot\|_{V_2})$ are reflexive Banach spaces and embedded in H continuously and densely. Then we get two triples:

$$V_1 \subset H \simeq H^* \subset V_1^* \quad \text{and} \quad V_2 \subset H \simeq H^* \subset V_2^*.$$

We define the norm $\|v\|_V := \|v\|_{V_1} + \|v\|_{V_2}$ on the space $V := V_1 \cap V_2$. Note that $(V, \|\cdot\|_V)$ is also a Banach space. Since V_1^* and V_2^* can be thought as subspaces of V^* , we get a Banach space $W := V_1^* + V_2^* \subset V^*$ with norm

$$\|f\|_W := \inf \{ \|f_1\|_{V_1^*} + \|f_2\|_{V_2^*} : f = f_1 + f_2, f_i \in V_i^*, i = 1, 2 \}.$$

Similarly, we write ${}_{V_i^*}\langle \cdot, \cdot \rangle_{V_i}$ to denote the pairing between V_i^* and $V_i, i = 1, 2$. Then, for all $v \in V$ and $f = f_1 + f_2 \in W \subset V^*$ we have

$${}_{V^*}\langle f, v \rangle_V = {}_{V_1^*}\langle f_1, v \rangle_{V_1} + {}_{V_2^*}\langle f_2, v \rangle_{V_2}.$$

Note carefully that if $f \in H$ and $v \in V$, then we obtain

$${}_{V^*}\langle f, v \rangle_V = {}_{V_1^*}\langle f, v \rangle_{V_1} = {}_{V_2^*}\langle f, v \rangle_{V_2} = \langle f, v \rangle_H.$$

We write $Pr(H)$ to mean the set of all Borel probability measures on H . Denote by $C_b(H)$ the space of all continuous functions $\varphi : H \rightarrow \mathbb{R}$ for which the norm $\|\varphi\|_\infty := \sup_{x \in H} |\varphi(x)|$ is

finite. Let $\{\mu_n\} := \{\mu_n\}_{n=1}^\infty \subset Pr(H)$ and $\mu \in Pr(H)$. We say μ_n converges weakly to μ in $Pr(H)$ provided $\int \varphi d\mu_n$ converges to $\int \varphi d\mu$ for all $\varphi \in C_b(H)$. Let $\varphi \in C_b(H)$ be Lipschitz continuous, we define

$$\|\varphi\|_{BL} := Lip(\varphi) + \|\varphi\|_\infty,$$

where $Lip(\varphi) = \sup_{x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\|x - y\|_H}$. Then $Pr(H)$ is a separable complete metric space with the following metric

$$d_{BL}(\mu, \nu) := \sup \left\{ \left| \int \varphi d\mu - \int \varphi d\nu \right| : \|\varphi\|_{BL} \leq 1 \right\}$$

for all $\mu, \nu \in Pr(H)$. It is well known that d_{BL} generates the weak topology on $Pr(H)$, i.e. $\mu_n \rightarrow \mu$ weakly in $Pr(H)$ if and only if $d_{BL}(\mu_n, \mu) \rightarrow 0$ as $n \rightarrow \infty$. See Chapter 11 in [14] for this metric d_{BL} (denoted by β there) and its related properties.

We assume in the following exposition that $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space. The space $L^2(\Omega, \mathbb{P}; H)$ consists of all H -valued random variables ζ such that $E\|\zeta\|_H^2 = \int_{\Omega} \|\zeta\|_H^2 dP < \infty$. An H -valued stochastic process $X = X(t)$, $t \in \mathbb{R}$ is called L^2 -bounded provided $\sup_{t \in \mathbb{R}} E\|X(t)\|_H^2 < \infty$. Throughout the paper, we denote by $\mathcal{L}(\zeta) \in Pr(H)$ the law or distribution of H -valued random variable ζ . A sequence of H -valued continuous stochastic processes $\{X_n\}$ is said to *converge in distribution to X (on $C(\mathbb{R}, H)$)* provided $\mathcal{L}(X_n)$ weakly converges to $\mathcal{L}(X)$ in $Pr(C(\mathbb{R}, H))$, where $\mathcal{L}(X)$ is the law or distribution of X on $C(\mathbb{R}, H)$. If $d_{BL}(\mathcal{L}(X_n(t)), \mathcal{L}(X(t))) \rightarrow 0$ as $n \rightarrow \infty$ for each $t \in \mathbb{R}$, we simply say that X_n *converges in distribution to X on H* .

3. COMPATIBLE SOLUTIONS

Let $W(t)$, $t \in \mathbb{R}$ be a two-sided cylindrical Q -Wiener process with $Q = I$ on a separable Hilbert space $(U, \langle \cdot, \cdot \rangle_U)$ with respect to a complete filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$. Denote by $L_2(U, H)$ the space of all Hilbert-Schmidt operators from U into H .

In this section, coefficient F in (1.3) need not to be Lipschitz. Therefore, instead of explicitly writing F in (1.3), we consider the following stochastic partial differential equation on H

$$(3.1) \quad dX(t) = A(t, X(t))dt + G(t, X(t))dW(t),$$

where $A(t, x) = A_1(x) + A_2(t, x)$, $A_i : \mathbb{R} \times V_i \rightarrow V_i^*$, $i = 1, 2$ and $G : \mathbb{R} \times V \rightarrow L_2(U, H)$.

Consider equation (3.1). Let us introduce the following conditions.

(H1) (Continuity) For all $u, v, w \in V$ and $t \in \mathbb{R}$ the map

$$(3.2) \quad \mathbb{R} \ni \theta \mapsto V_1^* \langle A_1(u + \theta v), w \rangle_{V_1}$$

is continuous. $A_2 : \mathbb{R} \times V_2 \rightarrow V_2^*$ and $G : \mathbb{R} \times V \rightarrow L_2(U, H)$ are continuous. Here A_1 is called *hemicontinuity* provided (3.2) hold.

(H2) (Strong monotonicity) There exist constants $\lambda \geq 0$, $r > 2$ and $\lambda' \geq 0$ such that for all $u, v \in V$, $t \in \mathbb{R}$

$$V^* \langle A(t, u) - A(t, v), u - v \rangle_V \leq -\lambda \|u - v\|_H^2 - \lambda' \|u - v\|_H^r$$

and

$$\|G(t, u) - G(t, v)\|_{L_2(U, H)}^2 \leq L_G^2 \|u - v\|_H^2.$$

(H3) (Coercivity) There exist constants $\alpha_1, \alpha_2 \in (1, \infty)$, $c_1 \in \mathbb{R}$, $c_2, c_2' \in (0, \infty)$ and $M_0 \in (0, \infty)$ such that for all $v \in V$, $t \in \mathbb{R}$

$$V^* \langle A(t, v), v \rangle_V \leq c_1 \|v\|_H^2 - c_2 \|v\|_{V_1}^{\alpha_1} - c_2' \|v\|_{V_2}^{\alpha_2} + M_0.$$

(H4) (Boundedness) There exist constants $c_3, c_3' \in (0, \infty)$ such that for all $v \in V$, $t \in \mathbb{R}$

$$\|A_1(v)\|_{V_1^*} \leq c_3 \|v\|_{V_1}^{\alpha_1 - 1} + M_0, \quad \|A_2(t, v)\|_{V_2^*} \leq c_3' \|v\|_{V_2}^{\alpha_2 - 1} + M_0$$

and

$$\|G(t, 0)\|_{L_2(U, H)} \leq M_0,$$

where α_i and M_0 are as in (H3).

(H5) A_2 and G are continuous in $t \in \mathbb{R}$ uniformly with respect to v on each bounded subset $Q \subset V$.

Remark 3.1. Since we consider compatible solutions (see Definition 3.10) by the method of dynamical systems, we assume that A_2 and G satisfy (H1) and (H5) that are different from the usual situation (i.e. we request stronger continuity conditions here). Under conditions of (H1)–(H2) and (H4)–(H5), $(H(A_2), \mathbb{R}, \sigma)$ and $(H(G), \mathbb{R}, \sigma)$ are dynamical systems, where $\sigma : \mathbb{R} \times H(A_2) \rightarrow H(A_2)$, $(\tau, \tilde{A}_2) \mapsto \tilde{A}_2^\tau$ and similarly for the action σ on $H(G)$. Note that we only need hemicontinuity of A_2 and do not need (H5), as usual, when we consider estimates of solutions, such as Lemmas 3.3–3.5, Theorems 3.6 and 3.8, Proposition 3.11, Lemma 3.12.

Definition 3.2 (see, e.g. [31, 45]). We say continuous H -valued (\mathcal{F}_t) -adapted process $X(t)$, $t \in [0, T]$ is a *solution* to equation (3.1), if $X \in \cap_{i=1,2} L^{\alpha_i}([0, T] \times \Omega, dt \otimes \mathbb{P}; V_i) \cap L^2([0, T] \times \Omega, dt \otimes \mathbb{P}; H)$ with α_i as in (H3) and \mathbb{P} -a.s.

$$(3.3) \quad X(t) = X(s) + \int_s^t A(\sigma, X(\sigma))d\sigma + \int_s^t G(\sigma, X(\sigma))dW(\sigma), \quad 0 \leq s \leq t \leq T.$$

In this paper, we write C_α to mean some positive constant which depends on α . Here α is one or more than one parameter and C_α may change from line to line. Now we discuss the L^2 -bounded solution to equation (3.1) by employing the classical pullback attraction method in random and non-autonomous dynamics (see, e.g. [10, 12] etc). For this we need three lemmas.

Lemma 3.3. *Assume that (H1)–(H4) hold. Let $\zeta_s \in L^2(\Omega, \mathcal{F}_s, \mathbb{P}; H)$ and $X(t, s, \zeta_s)$, $t \geq s$ be the solution to the following Cauchy problem*

$$\begin{cases} dX(t) = A(t, X(t))dt + G(t, X(t))dW(t) \\ X(s) = \zeta_s. \end{cases}$$

- (i) *If $2\lambda > L_G^2$, let $\eta \in (0, 2\lambda - L_G^2)$. Then there exist constants $1 \leq p < \frac{\eta}{2L_G^2} + 1$ and $\kappa, M_1 > 0$ such that*

$$(3.4) \quad E\|X(t, s, \zeta_s)\|_H^{2p} \leq e^{-\kappa(t-s)} E\|\zeta_s\|_H^{2p} + M_1,$$

where M_1 depends only on $\eta, c_2, c_3, c'_2, c'_3, \alpha_1, \alpha_2, \kappa, p, r$.

- (ii) *If $\lambda' > 0$ then estimate (3.4) hold for any $p \in [1, +\infty)$ and $\kappa > 0$.*

Proof. By (H2)–(H4) and Young's inequality, we have

$$(3.5) \quad 2_{V^*} \langle A(t, u), u \rangle_V + \|G(t, u)\|_{L_2(U, H)}^2 \leq \begin{cases} -\eta\|u\|_H^2 + C_{\alpha_1, \alpha_2, c_2, c'_2, M_0}, & \text{if } 2\lambda > L_G^2 \\ -\lambda'\|u\|_H^2 + (c_1 + 2L_G^2 - \lambda)\|u\|_H^2 + C_{\alpha_1, \alpha_2, c_2, c'_2, M_0}, & \text{if } \lambda' > 0. \end{cases}$$

Given $\kappa > 0$ and $p \geq 1$, in view of Itô's formula, we get

$$(3.6) \quad \begin{aligned} & E \left(e^{\kappa(t-s)} \|X(t, s, \zeta_s)\|_H^{2p} \right) \\ &= E \|\zeta_s\|_H^{2p} + \int_s^t \kappa e^{\kappa(\sigma-s)} E \|X(\sigma, s, \zeta_s)\|_H^{2p} d\sigma \\ & \quad + pE \int_s^t \|X(\sigma, s, \zeta_s)\|_H^{2p-2} e^{\kappa(\sigma-s)} \left(2_{V^*} \langle A(\sigma, X(\sigma, s, \zeta_s)), X(\sigma, s, \zeta_s) \rangle_V \right. \\ & \quad \left. + \|G(\sigma, X(\sigma, s, \zeta_s))\|_{L_2(U, H)}^2 \right) d\sigma \\ & \quad + 2p(p-1)E \int_s^t e^{\kappa(\sigma-s)} \|X(\sigma, s, \zeta_s)\|_H^{2p-4} \|(G(\sigma, X(\sigma, s, \zeta_s)))^* X(\sigma, s, \zeta_s)\|_U^2 d\sigma. \end{aligned}$$

If $2\lambda > L_G^2$, according to (3.5)–(3.6), (H2) and Young's inequality, we obtain

$$\begin{aligned}
& E \left(e^{\kappa(t-s)} \|X(t, s, \zeta_s)\|_H^{2p} \right) \\
& \leq E \|\zeta_s\|_H^{2p} + \int_s^t \kappa e^{\kappa(\sigma-s)} E \|X(\sigma, s, \zeta_s)\|_H^{2p} d\sigma \\
& \quad + pE \int_s^t \|X(\sigma, s, \zeta_s)\|_H^{2p-2} e^{\kappa(\sigma-s)} \left(-\eta \|X(\sigma, s, \zeta_s)\|_H^2 + C_{\alpha_1, \alpha_2, c_2, c'_2, M_0} \right) d\sigma \\
& \quad + 2p(p-1)E \int_s^t e^{\kappa(\sigma-s)} \|X(\sigma, s, \zeta_s)\|_H^{2p-2} \left(L_G^2 \|X(\sigma, s, \zeta_s)\|_H^2 + \varepsilon L_G^2 \|X(\sigma, s, \zeta_s)\|_H^2 + C_{\varepsilon, M_0} \right) d\sigma \\
& \leq E \|\zeta_s\|_H^{2p} + E \int_s^t e^{\kappa(\sigma-s)} \left(\kappa - \eta p + 2p(p-1)\varepsilon L_G^2 + 2p(p-1)L_G^2 \right) \|X(\sigma, s, \zeta_s)\|_H^{2p} d\sigma \\
& \quad + E \int_s^t e^{\kappa(\sigma-s)} \left(C_{\alpha_1, \alpha_2, c_2, c'_2, M_0, \varepsilon p} + C_{\varepsilon p(p-1)} \right) \|X(\sigma, s, \zeta_s)\|_H^{2p-2} d\sigma.
\end{aligned}$$

Let $1 \leq p < \frac{\eta}{2L_G^2} + 1$ and $\kappa \in (0, \eta p - 2p(p-1)L_G^2)$. Employing Young's inequality and taking ε small enough, we obtain

$$E \|X(t, s, \zeta_s)\|_H^{2p} \leq e^{-\kappa(t-s)} E \|\zeta_s\|_H^{2p} + M_1.$$

If $\lambda' > 0$, for any $\kappa > 0$, by (3.5)–(3.6), (H2) and Young's inequality we have

$$\begin{aligned}
& E \left(e^{\kappa(t-s)} \|X(t, s, \zeta_s)\|_H^{2p} \right) \\
& \leq E \|\zeta_s\|_H^{2p} + \int_s^t \kappa e^{\kappa(\sigma-s)} E \|X(\sigma, s, \zeta_s)\|_H^{2p} d\sigma + pE \int_s^t \|X(\sigma, s, \zeta_s)\|_H^{2p-2} e^{\kappa(\sigma-s)} \left(C_{\alpha_1, \alpha_2, c_2, c'_2, M_0} \right. \\
& \quad \left. + (c_1 + 2L_G^2 - \lambda) \|X(\sigma, s, \zeta_s)\|_H^2 - \lambda' \|X(\sigma, s, \zeta_s)\|_H^r \right) d\sigma \\
& \quad + 2p(p-1)E \int_s^t e^{\kappa(\sigma-s)} \|X(\sigma, s, \zeta_s)\|_H^{2p-2} \left(2L_G^2 \|X(\sigma, s, \zeta_s)\|_H^2 + 2M_0^2 \right) d\sigma \\
& \leq E \|\zeta_s\|_H^{2p} + E \int_s^t e^{\kappa(\sigma-s)} \left[-\lambda' p \|X(\sigma, s, \zeta_s)\|_H^{r+2p-2} + \left(\kappa + p(c_1 + 2L_G^2 - \lambda) \right. \right. \\
& \quad \left. \left. + 4p(p-1)L_G^2 \right) \|X(\sigma, s, \zeta_s)\|_H^{2p} + \left(pC_{\alpha_1, \alpha_2, c_2, c'_2, M_0} + 4p(p-1)M_0^2 \right) \|X(\sigma, s, \zeta_s)\|_H^{2p-2} \right] d\sigma \\
& \leq E \|\zeta_s\|_H^{2p} + E \int_s^t e^{\kappa(\sigma-s)} C_{\alpha_1, \alpha_2, c_2, c'_2, M_0, p, r} d\sigma.
\end{aligned}$$

Therefore, for any $p \in [1, +\infty)$ and $\kappa > 0$

$$E \|X(t, s, \zeta_s)\|_H^{2p} \leq e^{-\kappa(t-s)} E \|\zeta_s\|_H^{2p} + M_1.$$

□

Lemma 3.4. *Consider equation (3.1). Assume that $2\lambda - L_G^2 \geq 0$ and (H1)–(H4) hold. Let X and Y be two solutions of equation (3.1). If $\lambda' > 0$ or $2\lambda > L_G^2$, then for any $s \leq t$ we have*

$$\begin{aligned}
(3.7) \quad & E \|X(t, s, X(s)) - Y(t, s, Y(s))\|_H^2 \\
& \leq \begin{cases} E \|X(s) - Y(s)\|_H^2 \wedge \{\lambda'(r-2)(t-s)\}^{-\frac{2}{r-2}}, & \text{if } \lambda' > 0 \\ e^{-(2\lambda - L_G^2)(t-s)} E \|X(s) - Y(s)\|_H^2, & \text{if } 2\lambda > L_G^2. \end{cases}
\end{aligned}$$

In particular, for any $t \in \mathbb{R}$ there exists some random variable $X(t)$ such that

$$(3.8) \quad X(t, -n, 0) \rightarrow X(t) \quad \text{in } L^2(\Omega, \mathbb{P}; H) \text{ as } n \rightarrow \infty.$$

Proof. If $2\lambda > L_G^2$, by Itô's formula and (H2) we get

$$\begin{aligned} & E\|X(t, s, X(s)) - Y(t, s, Y(s))\|_H^2 \\ & \leq E\|X(s) - Y(s)\|_H^2 + E \int_s^t (-2\lambda + L_G^2) \|X(\sigma, s, X(s)) - Y(\sigma, s, Y(s))\|_H^2 d\sigma. \end{aligned}$$

It follows from Gronwall's lemma that

$$E\|X(t, s, X(s)) - Y(t, s, Y(s))\|_H^2 \leq e^{-(2\lambda - L_G^2)(t-s)} E\|X(s) - Y(s)\|_H^2.$$

If $\lambda' > 0$ and $2\lambda \geq L_G^2$, in view of Itô's formula and (H2), we have

$$\begin{aligned} & E\|X(t, s, X(s)) - Y(t, s, Y(s))\|_H^2 \\ & \leq E\|X(s) - Y(s)\|_H^2 + E \int_s^t -2\lambda' \|X(\sigma, s, X(s)) - Y(\sigma, s, Y(s))\|_H^r d\sigma \\ & \leq E\|X(s) - Y(s)\|_H^2 - 2\lambda' \int_s^t (E\|X(\sigma, s, X(s)) - Y(\sigma, s, Y(s))\|_H^2)^{\frac{r}{2}} d\sigma. \end{aligned}$$

Employing comparison theorem, we obtain

$$E\|X(t, s, X(s)) - Y(t, s, Y(s))\|_H^2 \leq E\|X(s) - Y(s)\|_H^2 \wedge \{\lambda'(r-2)(t-s)\}^{-\frac{2}{r-2}}.$$

□

Lemma 3.5. *Suppose that (H1)–(H4) hold. Let $X(t, s, \zeta_s)$ be a solution to equation (3.1) with initial value $X(s, s, \zeta_s) = \zeta_s$. We have*

$$(3.9) \quad \begin{aligned} & E \left(\sup_{t \in [s, s+T]} \|X(t, s, \zeta_s)\|_H^2 \right) + E \int_s^{s+T} \left(\|X(t, s, \zeta_s)\|_{V_1}^{\alpha_1} + \|X(t, s, \zeta_s)\|_{V_2}^{\alpha_2} \right) dt \\ & + E \int_s^{s+T} \left(\|A_1(X(t, s, \zeta_s))\|_{V_1^*}^{\frac{\alpha_1}{\alpha_1-1}} + \|A_2(t, X(t, s, \zeta_s))\|_{V_2^*}^{\frac{\alpha_2}{\alpha_2-1}} \right) dt \\ & \leq C_{c_1, L_G, M_0, T} (1 + E\|\zeta_s\|_H^2) \end{aligned}$$

for any $s \in \mathbb{R}, T > 0$.

Proof. By Itô's formula, (H2) and (H3), we have

$$(3.10) \quad \begin{aligned} & \|X(t, s, \zeta_s)\|_H^2 \\ & = \|\zeta_s\|_H^2 + \int_s^t 2V^* \langle A(\sigma, X(\sigma, s, \zeta_s)), X(\sigma, s, \zeta_s) \rangle_V + \|G(\sigma, X(\sigma, s, \zeta_s))\|_{L_2(U, H)}^2 d\sigma \\ & + 2 \int_s^t \langle X(\sigma, s, \zeta_s), G(\sigma, X(\sigma, s, \zeta_s)) dW(\sigma) \rangle_H \\ & \leq \|\zeta_s\|_H^2 + \int_s^t \left(2c_1 \|X(\sigma, s, \zeta_s)\|_H^2 - 2c_2 \|X(\sigma, s, \zeta_s)\|_{V_1}^{\alpha_1} - 2c_2' \|X(\sigma, s, \zeta_s)\|_{V_2}^{\alpha_2} + 2M_0 \right. \\ & \quad \left. + 2L_G^2 \|X(\sigma, s, \zeta_s)\|_H^2 + 2M_0^2 \right) d\sigma + 2 \int_s^t \langle X(\sigma, s, \zeta_s), G(\sigma, X(\sigma, s, \zeta_s)) dW(\sigma) \rangle_H. \end{aligned}$$

Dropping negative terms on the right of the above inequality, according to Burkholder-Davis-Gundy inequality (see, e.g. [31]) and Young's inequality, we get

$$(3.11) \quad E \sup_{t \in [s, s+T]} \|X(t, s, \zeta_s)\|_H^2$$

$$\begin{aligned}
&\leq E\|\zeta_s\|_H^2 + E \int_s^{s+T} (2c_1 + 2L_G^2) \|X(\sigma, s, \zeta_s)\|_H^2 + 2M_0^2 + 2M_0 d\sigma \\
&\quad + 6E \left(\int_s^{s+T} \|G(\sigma, X(\sigma, s, \zeta_s))\|_{L_2(U, H)}^2 \|X(\sigma, s, \zeta_s)\|_H^2 d\sigma \right)^{\frac{1}{2}} \\
&\leq E\|\zeta_s\|_H^2 + E \int_s^{s+T} (C_{c_1, L_G} \|X(\sigma, s, \zeta_s)\|_H^2 + C_{M_0}) d\sigma \\
&\quad + \frac{1}{2} E \sup_{t \in [s, s+T]} \|X(t, s, \zeta_s)\|_H^2.
\end{aligned}$$

By Gronwall's lemma, we obtain

$$(3.12) \quad E \sup_{t \in [s, s+T]} \|X(t, s, \zeta_s)\|_H^2 \leq C_{c_1, L_G, T, M_0} (1 + E\|\zeta_s\|_H^2).$$

Take expectations on both sides of (3.10) and let $t = s + T$, then by (3.12) we have

$$E \int_s^{s+T} \left(\|X(t, s, \zeta_s)\|_{V_1}^{\alpha_1} + \|X(t, s, \zeta_s)\|_{V_2}^{\alpha_2} \right) dt \leq C_{c_1, L_G, T, M_0} (1 + E\|\zeta_s\|_H^2).$$

In view of (H4), we complete the proof. \square

Theorem 3.6. *Consider equation (3.1). Suppose that $2\lambda - L_G^2 \geq 0$ and (H1)–(H4) hold. If $\lambda' > 0$ or $2\lambda > L_G^2$, then there exists a unique L^2 -bounded continuous H -valued solution $X(t)$, $t \in \mathbb{R}$ to equation (3.1). Moreover, the mapping $\hat{\mu} : \mathbb{R} \rightarrow Pr(H)$, defined by $\hat{\mu}(t) := \mathbb{P} \circ [X(t)]^{-1}$, is unique with the following properties:*

- (i) L^2 -boundedness: $\sup_{t \in \mathbb{R}} \int_H \|x\|_H^2 \hat{\mu}(t)(dx) < +\infty$;
- (ii) Flow property: $\mu(t, s, \hat{\mu}(s)) = \hat{\mu}(t)$ for all $t \geq s$.

Here $\mu(t, s, \mu_0)$ denotes the distribution of $X(t, s, \zeta_s)$ on H , with $\mu_0 = \mathbb{P} \circ \zeta_s^{-1}$.

Proof. For any fixed interval $[a, b] \subset \mathbb{R}$, we denote

$$\begin{aligned}
J &:= L^2([a, b] \times \Omega, dt \otimes \mathbb{P}; L_2(U, H)), \quad K_i := L^{\alpha_i}([a, b] \times \Omega, dt \otimes \mathbb{P}; V_i), \\
K_i^* &:= L^{\frac{\alpha_i}{\alpha_i-1}}([a, b] \times \Omega, dt \otimes \mathbb{P}; V_i^*), \quad i = 1, 2.
\end{aligned}$$

According to the reflexivity of K_i , $i = 1, 2$, (3.7) and (3.9), we may assume, going if necessary to a subsequence, that

- (1) $X(\cdot, -n, 0) \rightarrow X(\cdot)$ in $L^2([a, b] \times \Omega, dt \otimes \mathbb{P}; H)$ and $X(\cdot, -n, 0) \rightarrow X(\cdot)$ weakly in K_1 and K_2 ;
- (2) $A_i(\cdot, X(\cdot, -n, 0)) \rightarrow Y_i(\cdot)$ weakly in K_i^* , $i = 1, 2$;
- (3) $G(\cdot, X(\cdot, -n, 0)) \rightarrow Z(\cdot)$ weakly in J and hence

$$\int_a^t G(\sigma, X(\sigma, -n, 0)) dW(\sigma) \rightarrow \int_a^t Z(\sigma) dW(\sigma)$$

weakly* in $L^\infty([a, b], dt; L^2(\Omega, \mathbb{P}; H))$.

Thus for all $v \in V$, $\varphi \in L^\infty([a, b] \times \Omega)$ by Fubini's theorem we get

$$\begin{aligned}
&E \int_a^b v^* \langle X(t), \varphi(t)v \rangle_V dt \\
&= \lim_{n \rightarrow \infty} E \int_a^b v^* \langle X(t, -n, 0), \varphi(t)v \rangle_V dt \\
&= \lim_{n \rightarrow \infty} E \int_a^b v^* \langle X(a, -n, 0) + \int_a^t A(\sigma, X(\sigma, -n, 0)) d\sigma, \varphi(t)v \rangle_V dt
\end{aligned}$$

$$\begin{aligned}
& + \lim_{n \rightarrow \infty} E \left(\int_a^b \left\langle \int_a^t B(\sigma, X(\sigma, -n, 0)) dW(\sigma), \varphi(t)v \right\rangle_H dt \right) \\
& = E \int_a^b {}_{V^*} \langle X(a) + \int_a^t (Y_1(\sigma) + Y_2(\sigma)) d\sigma, \varphi(t)v \rangle_V dt \\
& \quad + E \left(\int_a^b \left\langle \int_a^t Z(\sigma) dW(\sigma), \varphi(t)v \right\rangle_H dt \right).
\end{aligned}$$

Let $Y(\sigma) := Y_1(\sigma) + Y_2(\sigma) \in W \subset V^*$, we have

$$X(t) = X(a) + \int_a^t Y(\sigma) d\sigma + \int_a^t Z(\sigma) dW(\sigma), \quad dt \otimes \mathbb{P}\text{-a.e.}$$

Thus, it remains to verify that

$$Y = A(\cdot, X), \quad Z = G(\cdot, X), \quad dt \otimes \mathbb{P}\text{-a.e.}$$

To this end, for any $\phi \in K_1 \cap K_2 \cap L^2([a, b] \times \Omega, dt \otimes \mathbb{P}; H)$, we have

$$\begin{aligned}
(3.13) \quad & E \|X(t, -n, 0)\|_H^2 - E \|X(a, -n, 0)\|_H^2 \\
& = E \int_a^t \left(2_{V^*} \langle A(\sigma, X(\sigma, -n, 0)), X(\sigma, -n, 0) \rangle + \|G(\sigma, X(\sigma, -n, 0))\|_{L_2(U, H)}^2 \right) d\sigma \\
& \leq E \int_a^t \left[2_{V^*} \langle A(\sigma, X(\sigma, -n, 0)) - A(\sigma, \phi(\sigma)), X(\sigma, -n, 0) - \phi(\sigma) \rangle_V \right. \\
& \quad + \|G(\sigma, X(\sigma, -n, 0)) - G(\sigma, \phi(\sigma))\|_{L_2(U, H)}^2 + 2_{V^*} \langle A(\sigma, \phi(\sigma)), X(\sigma, -n, 0) \rangle_V \\
& \quad + 2_{V^*} \langle A(\sigma, X(\sigma, -n, 0)) - A(\sigma, \phi(\sigma)), \phi(\sigma) \rangle_V \\
& \quad \left. + 2 \langle G(\sigma, X(\sigma, -n, 0)), G(\sigma, \phi(\sigma)) \rangle_{L_2(U, H)} - \|G(\sigma, \phi(\sigma))\|_{L_2(U, H)}^2 \right] d\sigma.
\end{aligned}$$

For every nonnegative $\psi \in L^\infty([a, b], dt; \mathbb{R})$, first multiplying $\psi(t)$ on both sides of (3.13), then integrating with respect to t from a to b and letting $n \rightarrow \infty$, it follows from (H2) and $2\lambda - L_G^2 > 0$ that

$$\begin{aligned}
(3.14) \quad & E \int_a^b \psi(t) (\|X(t)\|_H^2 - \|X(a)\|_H^2) dt \\
& \leq E \left(\int_a^b \psi(t) \int_a^t \left(2_{V_1^*} \langle Y_1(\sigma) - A_1(\phi(\sigma)), \phi(\sigma) \rangle_{V_1} + 2_{V_1^*} \langle A_1(\phi(\sigma)), X(\sigma) \rangle_{V_1} \right. \right. \\
& \quad + 2_{V_2^*} \langle Y_2(\sigma) - A_2(\sigma, \phi(\sigma)), \phi(\sigma) \rangle_{V_2} + 2_{V_2^*} \langle A_2(\sigma, \phi(\sigma)), X(\sigma) \rangle_{V_2} \\
& \quad \left. \left. + 2 \langle Z(\sigma), G(\sigma, \phi(\sigma)) \rangle_{L_2(U, H)} - \|G(\sigma, \phi(\sigma))\|_{L_2(U, H)}^2 \right) d\sigma dt \right).
\end{aligned}$$

Applying Itô's formula to $\|X(t)\|_H^2 - \|X(a)\|_H^2$ in (3.14), we get

$$\begin{aligned}
(3.15) \quad & 0 \geq E \left(\int_a^b \psi(t) \int_a^t \left(2_{V^*} \langle Y(\sigma) - A(\sigma, \phi(\sigma)), X(\sigma) - \phi(\sigma) \rangle_V \right. \right. \\
& \quad \left. \left. + \|G(\sigma, \phi(\sigma)) - Z(\sigma)\|_{L_2(U, H)}^2 \right) d\sigma dt \right).
\end{aligned}$$

Taking $\phi = X$ in (3.15), we have $Z = G(\cdot, X)$, $dt \otimes \mathbb{P}$ -a.e. Then, applying (3.15) to $\phi = X - \epsilon \tilde{\phi} v$ for $\epsilon > 0$ and $\tilde{\phi} \in L^\infty([a, b] \times \Omega, dt \otimes \mathbb{P}; \mathbb{R})$, $v \in V$, we have

$$E \left(\int_a^b \psi(t) \int_a^t 2_{V^*} \langle Y(\sigma) - A(\sigma, X(\sigma) - \epsilon \tilde{\phi}(\sigma)v), \epsilon \tilde{\phi}(\sigma)v \rangle_V d\sigma dt \right) \leq 0.$$

Dividing both sides by ϵ and letting $\epsilon \rightarrow 0$, according to Lebesgue's dominated convergence theorem, (H1) and (H4), we obtain

$$E \left(\int_a^b \psi(t) \int_a^t \tilde{\phi}(\sigma)_{V^*} \langle Y(\sigma) - A(\sigma, X(\sigma)), v \rangle_V d\sigma dt \right) \leq 0.$$

In view of the arbitrariness of ψ , $\tilde{\phi}$ and v , we conclude that $Y = A(\cdot, X)$, $dt \otimes \mathbb{P}$ -a.e. This completes the existence proof, i.e.

$$X(t) = X(a) + \int_a^t A(\sigma, X(\sigma)) d\sigma + \int_a^t G(\sigma, X(\sigma)) dW(\sigma), \quad dt \otimes \mathbb{P}\text{-a.e.}$$

By the arbitrariness of interval $[a, b] \subset \mathbb{R}$, we conclude that $X(\cdot)$ is a solution on \mathbb{R} . It follows from (3.4) that $\sup_{t \in \mathbb{R}} E \|X(t)\|_H^2 < \infty$. The uniqueness of L^2 -bounded solution is a consequence of (3.7).

The goal next is to prove that $\hat{\mu}$ is unique with the properties (i) and (ii). Note that

$$\sup_{t \in \mathbb{R}} \int_H \|x\|_H^2 \hat{\mu}(t)(dx) = \sup_{t \in \mathbb{R}} E \|X(t)\|_H^2 < \infty.$$

In view of the Chapman-Kolmogorov equation, we have $\mu(t, s, \mathcal{L}(X(s, -n, 0))) = \mathcal{L}(X(t, -n, 0))$. Then according to the Feller property, we get

$$\mu(t, s, \hat{\mu}(s)) = \hat{\mu}(t).$$

Suppose that μ_1 and μ_2 satisfy properties (i) and (ii), let $\zeta_{n,1}$ and $\zeta_{n,2}$ be random variables with the distributions $\mu_1(-n)$ and $\mu_2(-n)$ respectively. Then consider the solutions $X(t, -n, \zeta_{n,1})$ and $X(t, -n, \zeta_{n,2})$ on $[-n, \infty)$, we have

$$\begin{aligned} d_{BL}(\mu_1(t), \mu_2(t)) &= \sup_{\|f\|_{BL} \leq 1} \left| \int_H f(x) d(\mu(t, -n, \mu_1(-n)) - \mu(t, -n, \mu_2(-n))) \right| \\ &\leq (E \|X(t, -n, \zeta_{n,1}) - X(t, -n, \zeta_{n,2})\|_H^2)^{1/2}. \end{aligned}$$

Thus (3.7) yield that $\mu_1(t) = \mu_2(t)$ for all $t \in \mathbb{R}$. \square

Definition 3.7 (See [17]). We say that a solution $X(\cdot)$ of equation (3.1) is *stable in square-mean sense*, if for each $\epsilon > 0$, there exists $\delta > 0$ such that for all $t \geq 0$

$$E \|X(t, 0, \zeta_0) - X(t)\|_H^2 < \epsilon,$$

whenever $E \|\zeta_0 - X(0)\|_H^2 < \delta$. The solution $X(\cdot)$ is said to be *asymptotically stable in square-mean sense* if it is stable in square-mean sense and

$$(3.16) \quad \lim_{t \rightarrow \infty} E \|X(t, 0, \zeta_0) - X(t)\|_H^2 = 0.$$

We say $X(\cdot)$ is *globally asymptotically stable in square-mean sense* provided (3.16) holds for any $\zeta_0 \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$.

Applying Lemma 3.4 we obtain the following result:

Theorem 3.8. Consider equation (3.1). Suppose that $2\lambda - L_G^2 \geq 0$ and (H1)–(H4) hold. If $\lambda' > 0$ or $2\lambda > L_G^2$, then the unique L^2 -bounded solution of equation (3.1) is globally asymptotically stable in square-mean sense. Moreover,

$$(3.17) \quad E\|X(t, s, \zeta_s) - X(t)\|_H^2 \leq \begin{cases} E\|\zeta_s - X(s)\|_H^2 \wedge \{\lambda'(r-2)(t-s)\}^{-\frac{2}{r-2}}, & \text{if } \lambda' > 0 \\ e^{-(2\lambda - L_G^2)(t-s)} E\|\zeta_s - X(s)\|_H^2, & \text{if } 2\lambda > L_G^2 \end{cases}$$

for any $t \geq s$ and $\zeta_s \in L^2(\Omega, \mathcal{F}_s, \mathbb{P}; H)$.

Remark 3.9. (i) If A_2 and G satisfy (H2) and (H3), then every pair of functions $(\tilde{A}_2, \tilde{G}) \in H(A_2, G)$ possess the same property with the same constants, where

$$H(A_2, G) := \overline{\{(A_2^\tau, G^\tau) : \tau \in \mathbb{R}\}}.$$

Here $\overline{\{(A_2^\tau, G^\tau) : \tau \in \mathbb{R}\}}$ means the closure of $\{(A_2^\tau, G^\tau) : \tau \in \mathbb{R}\}$.

- (ii) If A_2 and G satisfy the conditions (H1), (H2), (H4) and (H5), then $A_2 \in BUC(\mathbb{R} \times V, V_2^*)$, $G \in BUC(\mathbb{R} \times V, L_2(U, H))$ and $H(A_2, G) \subset BUC(\mathbb{R} \times V, V_2^*) \times BUC(\mathbb{R} \times V, L_2(U, H))$.

Definition 3.10. Let $\{\varphi(t)\}_{t \in \mathbb{R}}$ be a solution of equation (3.1). Then φ is called *compatible* (respectively, *strongly compatible*) in distribution if the following conditions are fulfilled:

- (i) there exists a bounded closed subset $\mathcal{Q} \subset L^2(\Omega, \mathbb{P}; H)$ such that $\varphi(\mathbb{R}) \subseteq \mathcal{Q}$;
(ii) $\mathfrak{N}_{(F, G)} \subseteq \tilde{\mathfrak{N}}_\varphi$ (respectively, $\mathfrak{M}_{(F, G)} \subseteq \tilde{\mathfrak{M}}_\varphi$), where $\tilde{\mathfrak{N}}_\varphi$ (respectively, $\tilde{\mathfrak{M}}_\varphi$) means the set of all sequences $\{t_n\} \subset \mathbb{R}$ such that the sequence $\{\varphi(\cdot + t_n)\}$ converges to $\varphi(\cdot)$ (respectively, $\{\varphi(\cdot + t_n)\}$ converges) in distribution uniformly on any compact interval.

Now we show that the L^2 -bounded solution $X(t), t \in \mathbb{R}$ for equation (3.1) is strongly compatible in distribution. To this end, we need the tightness of the family of distributions $\{\mathbb{P} \circ [X(t)]^{-1}\}_{t \in \mathbb{R}}$. Therefore, we need the following condition (H6) which is used by many works (see, e.g. [27]).

(H6) Assume that there exists a closed subset $S \subset H$ equipped with the norm $\|\cdot\|_S$ such that $V \subset S$ is continuous and $S \subset H$ is compact. Let T_n be a sequence of positive definite self-adjoint operators on H such that for each $n \geq 1$,

$$\langle x, y \rangle_n := \langle x, T_n y \rangle_H, \quad x, y \in H,$$

defines a new inner product on H . Assume further that the norms $\|\cdot\|_n$ generated by $\langle \cdot, \cdot \rangle_n$ are all equivalent to $\|\cdot\|_H$ and for all $x \in S$ we have

$$\|x\|_n \uparrow \|x\|_S \quad \text{as } n \rightarrow \infty.$$

Furthermore, we suppose that for each $n \geq 1$, $T_n : V \rightarrow V$ is continuous and there exist constants $c_4 > 0$, $M_0 > 0$ such that for all $v \in V$, $t \in \mathbb{R}$

$$2_{V^*} \langle A(t, v), T_n v \rangle_V + \|G(t, v)\|_{L_2(U, H_n)}^2 \leq -c_4 \|v\|_n^2 + M_0.$$

Proposition 3.11. Consider equation (3.1). Suppose that conditions of Theorem 3.8 hold. If (H6) hold then the L^2 -bounded solution $X(\cdot)$ satisfies

$$(3.18) \quad \sup_{t \in \mathbb{R}} E\|X(t)\|_S^2 < \infty.$$

In particular, the family of distributions $\{P \circ [X(t)]^{-1}\}_{t \in \mathbb{R}}$ is tight.

Proof. Similar to the proof of Proposition 1 in [10], (3.18) can be obtained by Itô's formula, (H6) and Gronwall's lemma. \square

The following lemma is a direct corollary of Theorem 3.1 in [10].

Lemma 3.12. *Suppose that A_n, A, G_n, G satisfy (H1)–(H4) with the same constants $c, c_1, c_2, c_3, c'_2, c'_3, M_0, \alpha_i, i = 1, 2$ and L_G . Let X_n be the solution of the Cauchy problem*

$$(3.19) \quad \begin{cases} dX(t) = A_n(t, X(t))dt + G_n(t, X(t))dW(t) \\ X(s) = \zeta_n^s \end{cases}$$

and X be the solution to the Cauchy problem

$$(3.20) \quad \begin{cases} dX(t) = A(t, X(t))dt + G(t, X(t))dW(t) \\ X(s) = \zeta^s. \end{cases}$$

Assume further that

- (1) $\lim_{n \rightarrow \infty} A_{i,n}(t, x) = A_i(t, x)$ in V_i^* for all $t \in \mathbb{R}, x \in V, i = 1, 2$;
- (2) $\lim_{n \rightarrow \infty} G_n(t, x) = G(t, x)$ in $L_2(U, H)$ for all $t \in \mathbb{R}, x \in V$.

Then we have the following conclusions:

- (i) If $\lim_{n \rightarrow \infty} E \|\zeta_n^s - \zeta^s\|_H^2 = 0$, then $\lim_{n \rightarrow \infty} E \sup_{s \leq \tau \leq t} \|X_n(\tau) - X(\tau)\|_H^2 = 0$ for any $t > s$;
- (ii) If $\lim_{n \rightarrow \infty} \zeta_n^s = \zeta^s$ in probability, then $\lim_{n \rightarrow \infty} \sup_{\tau \in [s, t]} \|X_n(\tau) - X(\tau)\|_H = 0$ in probability;
- (iii) If $\lim_{n \rightarrow \infty} d_{BL}(\mathcal{L}(\zeta_n^s), \mathcal{L}(\zeta^s)) = 0$ in $Pr(H)$, then

$$\lim_{n \rightarrow \infty} d_{BL}(\mathcal{L}(X_n), \mathcal{L}(X)) = 0 \quad \text{in } Pr(C([s, \infty), H)).$$

Theorem 3.13. *Consider equation (3.1). Suppose that $2\lambda - L_G^2 \geq 0$ and (H1)–(H6) hold. If $\lambda' > 0$ or $2\lambda > L_G^2$, then the unique L^2 -bounded solution is strongly compatible in distribution.*

Proof. It follows from Remark 3.9 that $H(A_2, G) \subset BUC(\mathbb{R} \times V, V_2^*) \times BUC(\mathbb{R} \times V, L_2(U, H))$. Let $\{t_n\} \in \mathfrak{M}_{(A_2, G)}$, then there exists $(\tilde{A}_2, \tilde{G}) \in H(A_2, G)$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{|t| \leq l, \|x\|_V \leq r} \|A_2(t + t_n, x) - \tilde{A}_2(t, x)\|_{V_2^*} &= 0, \\ \lim_{n \rightarrow \infty} \sup_{|t| \leq l, \|x\|_V \leq r} \|G(t + t_n, x) - \tilde{G}(t, x)\|_{L_2(U, H)} &= 0, \end{aligned}$$

for any $l > 0$ and $r > 0$. Let X_n be the unique L^2 -bounded solution of

$$dX(t) = (A_1(X(t)) + A_2(t + t_n, X(t))) dt + G(t + t_n, X(t))dW(t)$$

and \tilde{X} be the unique L^2 -bounded solution of

$$(3.21) \quad dX(t) = \left(A_1(X(t)) + \tilde{A}_2(t, X(t)) \right) dt + \tilde{G}(t, X(t))dW(t).$$

We now prove that for any $[a, b] \subset \mathbb{R}$, $\lim_{n \rightarrow \infty} \sup_{t \in [a, b]} d_{BL}(\mathcal{L}(X_n(t)), \mathcal{L}(\tilde{X}(t))) = 0$. According to

Lemma 3.12, we only need to prove that $\lim_{n \rightarrow \infty} d_{BL}(\mathcal{L}(X_n(t)), \mathcal{L}(\tilde{X}(t))) = 0$ in $Pr(H)$ for every $t \in \mathbb{R}$. To this end, it suffices to show that for every sequence $\{\gamma'_k\} := \{\gamma'_k\}_{k=1}^\infty \subset \mathbb{N}$, there exists a subsequence $\{\gamma_k\}$ of $\{\gamma'_k\}$ such that $\lim_{k \rightarrow \infty} d_{BL}(\mathcal{L}(X_{\gamma_k}(t)), \mathcal{L}(\tilde{X}(t))) = 0$ in $Pr(H)$ for every $t \in \mathbb{R}$.

Given $r \geq 1$, according to the tightness of $\{\mathcal{L}(X_{\gamma'_k}(-r))\}$, there exists a subsequence $\{\gamma_k\} \subset \{\gamma'_k\}$ such that $\mathcal{L}(X_{\gamma_k}(-r))$ converges weakly to some probability measure μ_r in $Pr(H)$. Let ξ_r be a random variable with distribution μ_r . Define $Y_r(t) := X(t, -r, \xi_r)$, where $X(t, -r, \xi_r), t \in [-r, +\infty)$ is a solution to the following Cauchy problem

$$\begin{cases} dX(t) = \left(A_1(X(t)) + \tilde{A}_2(t, X(t)) \right) dt + \tilde{G}(t, X(t))dW(t) \\ X(-r) = \xi_r. \end{cases}$$

In view of Lemma 3.12, we have

$$\lim_{k \rightarrow \infty} d_{BL}(\mathcal{L}(X_{\gamma_k}), \mathcal{L}(Y_r)) = 0 \quad \text{in } Pr(C([-r, +\infty), H)).$$

Since $\{\mathcal{L}(X_{\gamma_k}(-r-1))\}$ is tight, going if necessary to a subsequence, we can assume that $\mathcal{L}(X_{\gamma_k}(-r-1))$ converges weakly to some probability measure μ_{r+1} in $Pr(H)$. Let ξ_{r+1} be a random variable with distribution μ_{r+1} . In light of Lemma 3.12, we have

$$\lim_{k \rightarrow \infty} d_{BL}(\mathcal{L}(X_{\gamma_k}), \mathcal{L}(Y_{r+1})) = 0 \quad \text{in } Pr(C([-r-1, +\infty), H)),$$

where $Y_{r+1}(t) := X(t, -r-1, \xi_{r+1})$, $t \in [-r-1, +\infty)$. Therefore, we have $d_{BL}(\mathcal{L}(Y_r), \mathcal{L}(Y_{r+1})) = 0$ in $Pr(C([-r, +\infty), H))$. In particular, $\mathcal{L}(Y_r(t)) = \mathcal{L}(Y_{r+1}(t))$ for all $t \geq -r$.

Define $\nu(t) := \mathcal{L}(Y_r(t))$, $t \geq -r$. We use a standard diagonal argument to extract a subsequence which we still denote by $\{X_{\gamma_k}\}$ satisfying

$$\lim_{k \rightarrow \infty} d_{BL}(\mathcal{L}(X_{\gamma_k}(t)), \nu(t)) = 0 \quad \text{in } Pr(H)$$

for every $t \in \mathbb{R}$. Note that $\sup_{t \in \mathbb{R}} \int_H \|x\|^2 \nu(t)(dx) < +\infty$. And we have \mathbb{P} -a.s.

$$Y_r(t) = Y_r(s) + \int_s^t \left(A_1(Y_r(\sigma)) + \tilde{A}_2(\sigma, Y_r(\sigma)) \right) d\sigma + \int_s^t \tilde{G}(\sigma, Y_r(\sigma)) dW(\sigma),$$

where $t \geq s \geq -r$. By the uniqueness in law of the solutions for equation (3.21), we have $\mathcal{L}(Y_r(t)) = \mu(t, s, \mathcal{L}(Y_r(s)))$, $t \geq s \geq -r$, i.e. $\nu(t) = \mu(t, s, \nu(s))$, $t \geq s$. In view of Theorem 3.6, we obtain $\nu = \mathcal{L}(\tilde{X})$. Therefore, we have

$$\lim_{k \rightarrow \infty} d_{BL}(\mathcal{L}(X_{\gamma_k}(t)), \mathcal{L}(\tilde{X}(t))) = 0 \quad \text{in } Pr(H)$$

for every $t \in \mathbb{R}$.

Note that $X(\cdot + t_n)$ and $X_n(\cdot)$ share the same distribution. It follows from Definition 3.10 and Lemma 3.12 that $X(\cdot)$ is strongly compatible in distribution. \square

By Theorems 2.18 and 3.13, we have the following result.

Corollary 3.14. *Under the conditions of Theorem 3.13 the following statements hold:*

- (i) *If $A_2 \in C(\mathbb{R} \times V, V_2^*)$ and $G \in C(\mathbb{R} \times V, L_2(U, H))$ are jointly stationary (respectively, T -periodic, quasi-periodic with the spectrum of frequencies ν_1, \dots, ν_k , almost periodic, almost automorphic, Birkhoff recurrent, Lagrange stable, Levitan almost periodic, almost recurrent, Poisson stable) in $t \in \mathbb{R}$ uniformly with respect to x on each bounded subset, then so is the unique solution $\varphi \in C_b(\mathbb{R}, L^2(\Omega, \mathbb{P}; H))$ of equation (3.1) in distribution;*
- (ii) *If $A_2 \in C(\mathbb{R} \times V, V_2^*)$ and $G \in C(\mathbb{R} \times V, L_2(U, H))$ are Lagrange stable and jointly pseudo-periodic (respectively, pseudo-recurrent) in $t \in \mathbb{R}$ uniformly with respect to x on each bounded subset, then equation (3.1) has a unique solution $\varphi \in C_b(\mathbb{R}, L^2(\mathbb{P}, H))$ which is pseudo-periodic (respectively, pseudo-recurrent) in distribution.*

4. THE SECOND BOGOLYUBOV THEOREM

Consider the following stochastic partial differential equation

$$(4.1) \quad dX_\varepsilon(t) = \left(A(X_\varepsilon(t)) + F\left(\frac{t}{\varepsilon}, X_\varepsilon(t)\right) \right) dt + G\left(\frac{t}{\varepsilon}, X_\varepsilon(t)\right) dW(t),$$

where $A(x) = A_1(x) + A_2(x)$, $A_i : V_i \rightarrow V_i^*$, $i = 1, 2$, $F \in C(\mathbb{R} \times H, H)$, $G \in C(\mathbb{R} \times H, L_2(U, H))$ and $0 < \varepsilon \leq 1$. Here W is a U -valued two-sided cylindrical Wiener process with the identity covariance operator with respect to a filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{F}_t)$,

where $\mathcal{F}_t := \sigma\{W(u) - W(v) : u, v \leq t\}$. In this section, we will omit the index H of $\|\cdot\|_H$ and $\langle \cdot, \cdot \rangle_H$ if it does not cause confusion.

We employ Ψ to denote the space of all decreasing, positive bounded functions $\omega : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\lim_{t \rightarrow +\infty} \omega(t) = 0$. Below we need additional conditions.

(H2') There exist constants $\lambda, \lambda' \geq 0$, $L_F, M_0 > 0$, $r > 2$ and $\lambda_F \in \mathbb{R}$ such that

$$\begin{aligned} v^* \langle A(u) - A(v), u - v \rangle_V &\leq -\lambda \|u - v\|^2 - \lambda' \|u - v\|^r, \\ \langle F(t, u) - F(t, v), u - v \rangle &\leq \lambda_F \|u - v\|^2, \quad \|F(t, 0)\| \leq M_0 \\ \|F(t, u) - F(t, v)\| &\leq L_F \|u - v\|, \quad \|G(t, u) - G(t, v)\|_{L_2(U, H)} \leq L_G \|u - v\| \end{aligned}$$

for any $t \in \mathbb{R}$ and $x, y \in H$;

(G1) There exist functions $\omega_1 \in \Psi$ and $\bar{F} \in C(H, H)$ such that

$$\frac{1}{T} \left\| \int_t^{t+T} [F(s, x) - \bar{F}(x)] ds \right\| \leq \omega_1(T)(1 + \|x\|)$$

for any $T > 0$, $x \in H$ and $t \in \mathbb{R}$;

(G2) There exist functions $\omega_2 \in \Psi$ and $\bar{G} \in C(H, L_2(U, H))$ such that

$$\frac{1}{T} \int_t^{t+T} \|G(s, x) - \bar{G}(x)\|_{L_2(U, H)}^2 ds \leq \omega_2(T)(1 + \|x\|^2)$$

for any $T > 0$, $x \in H$ and $t \in \mathbb{R}$.

Remark 4.1. (i) Note that we can just assume A_2 is hemicontinuous when A_2 is independent of time t . In the following, we still say that equation (4.1) satisfies (H5) (respectively, (H6)), if F and G are continuous in $t \in \mathbb{R}$ uniformly with respect to u on each bounded $Q \subset H$ (respectively, there exist constants $c_4, M_0 > 0$ such that $2v^* \langle A(v), T_n v \rangle_V + 2 \langle F(t, v), T_n v \rangle + \|G(t, v)\|_{L_2(U, H)}^2 \leq -c_4 \|v\|_n^2 + M_0$ for all $v \in V$ and $t \in \mathbb{R}$).

(ii) If F and G satisfy (H2'), (H4) and (G1)–(G2), then \bar{F} and \bar{G} also satisfy (H2') and (H4) with the same constants. Therefore, under the same conditions, estimates (3.4), (3.9), (3.17), (3.18) uniformly hold for $\varepsilon \in (0, 1]$, \bar{F} and \bar{G} .

(iii) It can be verified that (G1) (respectively, (G2)) holds if and only if

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} F(s, x) ds = \bar{F}(x)$$

(respectively, $\lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \|G(s, x) - \bar{G}(x)\|_{L_2(U, H)}^2 ds = 0$) uniformly with respect to $t \in \mathbb{R}$.

Denote by $F_\varepsilon(t, x) := F(\frac{t}{\varepsilon}, x)$ and $G_\varepsilon(t, x) := G(\frac{t}{\varepsilon}, x)$ for any $t \in \mathbb{R}$, $x \in H$ and $\varepsilon \in (0, 1]$. Equation (4.1) can be written as

$$(4.2) \quad dX_\varepsilon(t) = (A(X_\varepsilon(t)) + F_\varepsilon(t, X_\varepsilon(t)))dt + G_\varepsilon(t, X_\varepsilon(t))dW(t).$$

Along with equations (4.1)–(4.2) we consider the following averaged equation

$$(4.3) \quad dX(t) = (A(X(t)) + \bar{F}(X(t)))dt + \bar{G}(X(t))dW(t).$$

Recall that C_α mean some positive constant which depends on α . For simplicity, we just write C when C depends on some parameters of $\lambda, \lambda', r, \lambda_F, L_F, L_G, c_1, c_2, c'_2, \alpha_1, \alpha_2, M_0, c_3, c'_3$ in (H1), (H2') and (H3). Let $\delta > 0$ be a fixed constant depending on ε . Denote a step process $\tilde{\phi}$ such that $\tilde{\phi}(\sigma) = \phi(s + k\delta)$ for any $\sigma \in [s + k\delta, s + (k+1)\delta)$. Employing the technique of time discretization, we have the following estimates.

Lemma 4.2. *Assume that (H1), (H2'), (H3)–(H4) and (G1)–(G2) hold. Let Y_ε be the solution of (4.2) with the initial value $Y_\varepsilon(s) = \zeta_s^\varepsilon$ and \bar{Y} be the solution of (4.3) with the initial value $\bar{Y}(s) = \zeta_s$. Then we have*

$$(4.4) \quad E \int_s^{s+T} \|Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma)\|^2 d\sigma \leq C_T(1 + E\|\zeta_s^\varepsilon\|^2)\delta^{\frac{1}{2}}$$

and

$$(4.5) \quad E \int_s^{s+T} \|\bar{Y}(\sigma) - \tilde{Y}(\sigma)\|^2 d\sigma \leq C_T(1 + E\|\zeta_s\|^2)\delta^{\frac{1}{2}}$$

for any $s \in \mathbb{R}$ and $T > 0$.

Proof. By Lemma 3.5 we have

$$(4.6) \quad \begin{aligned} & E \int_s^{s+T} \|Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma)\|^2 d\sigma \\ &= E \int_s^{s+\delta} \|Y_\varepsilon(\sigma) - \zeta_s^\varepsilon\|^2 d\sigma + E \sum_{k=1}^{T(\delta)-1} \int_{s+k\delta}^{s+(k+1)\delta} \|Y_\varepsilon(\sigma) - Y_\varepsilon(s+k\delta)\|^2 d\sigma \\ &\quad + E \int_{s+T(\delta)\delta}^{s+T} \|Y_\varepsilon(\sigma) - Y_\varepsilon(s+T(\delta)\delta)\|^2 d\sigma \\ &\leq C_T(1 + E\|\zeta_s^\varepsilon\|^2)\delta + 2E \sum_{k=1}^{T(\delta)-1} \int_{s+k\delta}^{s+(k+1)\delta} \|Y_\varepsilon(\sigma) - Y_\varepsilon(\sigma-\delta)\|^2 d\sigma \\ &\quad + 2E \sum_{k=1}^{T(\delta)-1} \int_{s+k\delta}^{s+(k+1)\delta} \|Y_\varepsilon(\sigma-\delta) - Y_\varepsilon(s+k\delta)\|^2 d\sigma \\ &=: C_T(1 + E\|\zeta_s^\varepsilon\|^2)\delta + 2 \sum_{k=1}^{T(\delta)-1} I_k + 2 \sum_{k=1}^{T(\delta)-1} J_k. \end{aligned}$$

Given $k \in [1, T(\delta) - 1]$, for any $\sigma \in [s+k\delta, s+(k+1)\delta)$, by Itô's formula, (H2'), (H3)–(H4) and Young's inequality we get

$$(4.7) \quad \begin{aligned} & \|Y_\varepsilon(\sigma) - Y_\varepsilon(\sigma-\delta)\|^2 \\ &= \int_{\sigma-\delta}^{\sigma} 2V^* \langle A(Y_\varepsilon(u)), Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta) \rangle_V + 2 \langle F_\varepsilon(u, Y_\varepsilon(u)), Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta) \rangle du \\ &\quad + \int_{\sigma-\delta}^{\sigma} \|G_\varepsilon(u, Y_\varepsilon(u))\|_{L_2(U, H)}^2 du + 2 \int_{\sigma-\delta}^{\sigma} \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta), G_\varepsilon(u, Y_\varepsilon(u)) \rangle dW(u) \\ &\leq \int_{\sigma-\delta}^{\sigma} \left(2\|A_1(Y_\varepsilon(u))\|_{V_1^*} \|Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta)\|_{V_1} + 2\|A_2(Y_\varepsilon(u))\|_{V_2^*} \|Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta)\|_{V_2} \right. \\ &\quad \left. + 2\|F_\varepsilon(u, Y_\varepsilon(u))\| \|Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta)\| + 2L_G^2 \|Y_\varepsilon(u)\|^2 + 2M_0^2 \right) du \\ &\quad + 2 \int_{\sigma-\delta}^{\sigma} \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta), G_\varepsilon(u, Y_\varepsilon(u)) \rangle dW(u) \\ &\leq \int_{\sigma-\delta}^{\sigma} \left[2 \left(\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1-1} + M_0 \right) \|Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta)\|_{V_1} \right. \\ &\quad \left. + 2 \left(\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2-1} + M_0 \right) \|Y_\varepsilon(u) - Y_\varepsilon(\sigma-\delta)\|_{V_2} \right] du \end{aligned}$$

$$\begin{aligned}
& + 2(L_F \|Y_\varepsilon(u)\| + M_0) \|Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta)\| + 2L_G^2 \|Y_\varepsilon(u)\|^2 + 2M_0^2 \Big] du \\
& + 2 \int_{\sigma-\delta}^\sigma \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta), G_\varepsilon(u, Y_\varepsilon(u)) dW(u) \rangle \\
\leq & \int_{\sigma-\delta}^\sigma \left[2\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + 2\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1-1} \|Y_\varepsilon(\sigma - \delta)\|_{V_1} + 2M_0 \|Y_\varepsilon(u)\|_{V_1} + 2M_0 \|Y_\varepsilon(\sigma - \delta)\|_{V_1} \right. \\
& + 2\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + 2\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2-1} \|Y_\varepsilon(\sigma - \delta)\|_{V_2} + 2M_0 \|Y_\varepsilon(u)\|_{V_2} + 2M_0 \|Y_\varepsilon(\sigma - \delta)\|_{V_2} \\
& \left. + (2L_F + L_F^2 + 2L_G^2 + 1) \|Y_\varepsilon(u)\|^2 + 2\|Y_\varepsilon(\sigma - \delta)\|^2 + 4M_0^2 \right] du \\
& + 2 \int_{\sigma-\delta}^\sigma \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta), G_\varepsilon(u, Y_\varepsilon(u)) dW(u) \rangle \\
\leq & \int_{\sigma-\delta}^\sigma \left[4\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + \frac{4}{\alpha_1} \|Y_\varepsilon(\sigma - \delta)\|_{V_1}^{\alpha_1} + 4\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + \frac{4}{\alpha_2} \|Y_\varepsilon(\sigma - \delta)\|_{V_2}^{\alpha_2} + \frac{4(\alpha_1 - 1)}{\alpha_1} M_0^{\frac{\alpha_1}{\alpha_1-1}} \right. \\
& \left. + \frac{4(\alpha_2 - 1)}{\alpha_2} M_0^{\frac{\alpha_2}{\alpha_2-1}} + (2L_F + L_F^2 + 2L_G^2 + 1) \|Y_\varepsilon(u)\|^2 + 2\|Y_\varepsilon(\sigma - \delta)\|^2 + 4M_0^2 \right] du \\
& + 2 \int_{\sigma-\delta}^\sigma \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta), G_\varepsilon(u, Y_\varepsilon(u)) dW(u) \rangle.
\end{aligned}$$

Then we have

(4.8)

$$\begin{aligned}
I_k & = E \int_{s+k\delta}^{s+(k+1)\delta} \|Y_\varepsilon(\sigma) - Y_\varepsilon(\sigma - \delta)\|^2 d\sigma \\
& \leq E \int_{s+k\delta}^{s+(k+1)\delta} \left\{ \int_{\sigma-\delta}^\sigma \left[4\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + \frac{4}{\alpha_1} \|Y_\varepsilon(\sigma - \delta)\|_{V_1}^{\alpha_1} + 4\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + \frac{4}{\alpha_2} \|Y_\varepsilon(\sigma - \delta)\|_{V_2}^{\alpha_2} \right. \right. \\
& \quad \left. \left. + \frac{4(\alpha_1 - 1)}{\alpha_1} M_0^{\frac{\alpha_1}{\alpha_1-1}} + \frac{4(\alpha_2 - 1)}{\alpha_2} M_0^{\frac{\alpha_2}{\alpha_2-1}} + (2L_F + L_F^2 + 2L_G^2 + 1) \|Y_\varepsilon(u)\|^2 \right. \right. \\
& \quad \left. \left. + 2\|Y_\varepsilon(\sigma - \delta)\|^2 + 4M_0^2 \right] du + 2 \int_{\sigma-\delta}^\sigma \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta), G_\varepsilon(u, Y_\varepsilon(u)) dW(u) \rangle \right\} d\sigma \\
& \leq E \int_{s+(k-1)\delta}^{s+(k+1)\delta} \int_u^{u+\delta} \left[4\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + 4\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + C_{\alpha_1, \alpha_2, M_0} + C_{L_F, L_G} \|Y_\varepsilon(u)\|^2 \right] d\sigma du \\
& \quad + E \int_{s+k\delta}^{s+(k+1)\delta} \delta \left(\frac{4}{\alpha_1} \|Y_\varepsilon(\sigma - \delta)\|_{V_1}^{\alpha_1} + \frac{4}{\alpha_2} \|Y_\varepsilon(\sigma - \delta)\|_{V_2}^{\alpha_2} + 2\|Y_\varepsilon(\sigma - \delta)\|^2 \right) d\sigma + I_k^2 \\
& \leq E \int_{s+(k-1)\delta}^{s+(k+1)\delta} \delta \left[4\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + 4\|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + C_{\alpha_1, \alpha_2, M_0} + C_{L_F, L_G} \|Y_\varepsilon(u)\|^2 \right] du \\
& \quad + E \int_{s+k\delta}^{s+(k+1)\delta} \delta \left(\frac{4}{\alpha_1} \|Y_\varepsilon(\sigma - \delta)\|_{V_1}^{\alpha_1} + \frac{4}{\alpha_2} \|Y_\varepsilon(\sigma - \delta)\|_{V_2}^{\alpha_2} + 2\|Y_\varepsilon(\sigma - \delta)\|^2 \right) d\sigma + I_k^2 \\
& \leq \delta CE \int_{s+(k-1)\delta}^{s+(k+1)\delta} \left(\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + \|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + \|Y_\varepsilon(u)\|^2 + 1 \right) du + I_k^2.
\end{aligned}$$

Now we estimate I_k^2 . In view of Burkholder-Davis-Gundy inequality, (H2'), (H4) and Young's inequality, we obtain

$$\begin{aligned}
(4.9) \quad I_k^2 &= 2E \int_{s+k\delta}^{s+(k+1)\delta} \int_{\sigma-\delta}^{\sigma} \langle Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta), G_\varepsilon(u, Y_\varepsilon(u)) dW(u) \rangle d\sigma \\
&\leq 6 \int_{s+k\delta}^{s+(k+1)\delta} E \left(\int_{\sigma-\delta}^{\sigma} \|G_\varepsilon(u, Y_\varepsilon(u))\|_{L_2(U, H)}^2 \|Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta)\|^2 du \right)^{\frac{1}{2}} d\sigma \\
&\leq 6 \int_{s+k\delta}^{s+(k+1)\delta} E \left(\int_{\sigma-\delta}^{\sigma} (2L_G^2 \|Y_\varepsilon(u)\|^2 + 2M_0^2) \|Y_\varepsilon(u) - Y_\varepsilon(\sigma - \delta)\|^2 du \right)^{\frac{1}{2}} d\sigma \\
&\leq \delta^{\frac{1}{2}} C_{L_G, M_0} \left[\int_{s+k\delta}^{s+(k+1)\delta} E \int_{\sigma-\delta}^{\sigma} (\|Y_\varepsilon(u)\|^4 + \|Y_\varepsilon(\sigma - \delta)\|^4 + 1) dud\sigma \right]^{\frac{1}{2}} \\
&\leq \delta^{\frac{1}{2}} C_{L_G, M_0} \left(E \int_{s+(k-1)\delta}^{s+(k+1)\delta} \delta \|Y_\varepsilon(u)\|^4 du + \delta^2 \right)^{\frac{1}{2}} \\
&\leq \delta^{\frac{3}{2}} C_{L_G, M_0} \left[\left(E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(u)\|^2 \right) + 1 \right].
\end{aligned}$$

Therefore (4.8) and (4.9) yield

$$\begin{aligned}
(4.10) \quad I_k &\leq \delta C E \int_{s+(k-1)\delta}^{s+(k+1)\delta} \left(\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + \|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + \|Y_\varepsilon(u)\|^2 + 1 \right) du \\
&\quad + \delta^{\frac{3}{2}} C \left(E \sup_{s \leq t \leq t+T} \|Y_\varepsilon(t)\|^2 + 1 \right).
\end{aligned}$$

By (3.9) and Remark 4.1 (ii) we get

$$\begin{aligned}
(4.11) \quad 2 \sum_{k=1}^{T(\delta)-1} I_k &\leq \delta C E \int_s^{s+T} \left(\|Y_\varepsilon(u)\|_{V_1}^{\alpha_1} + \|Y_\varepsilon(u)\|_{V_2}^{\alpha_2} + \|Y_\varepsilon(u)\|^2 + 1 \right) du \\
&\quad + \delta^{\frac{1}{2}} C \left(E \sup_{s \leq t \leq t+T} \|Y_\varepsilon(t)\|^2 + 1 \right) \\
&\leq C_T (1 + E \|\zeta_s^\varepsilon\|^2) \delta^{\frac{1}{2}}.
\end{aligned}$$

Similarly, we have

$$(4.12) \quad 2 \sum_{k=1}^{T(\delta)-1} J_k \leq C_T (1 + E \|\zeta_s^\varepsilon\|^2) \delta^{\frac{1}{2}}.$$

Combining (4.6), (4.11) and (4.12), we obtain

$$E \int_s^{s+T} \|Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma)\|^2 d\sigma \leq C_T (1 + E \|\zeta_s^\varepsilon\|^2) \delta^{\frac{1}{2}}.$$

It follows from the same steps as in the proof of (4.4) that

$$E \int_s^{s+T} \|\bar{Y}(\sigma) - \tilde{Y}(\sigma)\|^2 d\sigma \leq C_T (1 + E \|\zeta_s\|^2) \delta^{\frac{1}{2}}.$$

□

Now we establish the following first Bogolyubov theorem.

Theorem 4.3. *Suppose that (G1)–(G2), (H1), (H2') and (H3)–(H4) hold. For any $s \in \mathbb{R}$, let Y_ε be the solution of the following Cauchy problem*

$$\begin{cases} dX(t) = (A(X(t)) + F_\varepsilon(t, X(t))) dt + G_\varepsilon(t, X(t))dW(t) \\ X(s) = \zeta_s^\varepsilon \end{cases}$$

and \bar{Y} be the solution of the following Cauchy problem

$$\begin{cases} dX(t) = (A(X(t)) + \bar{F}(X(t))) dt + \bar{G}(X(t))dW(t) \\ X(s) = \zeta_s. \end{cases}$$

Assume further that $\lim_{\varepsilon \rightarrow 0} E \|\zeta_s^\varepsilon - \zeta_s\|^2 = 0$. Then

$$\lim_{\varepsilon \rightarrow 0} E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 = 0$$

for any $T > 0$.

Proof. In view of Itô's formula and (H2'), we have

(4.13)

$$\begin{aligned} & \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 \\ &= \|\zeta_s^\varepsilon - \zeta_s\|^2 + \int_s^t \left(2_{V^*} \langle A(Y_\varepsilon(\sigma)) - A(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle_V \right. \\ &\quad \left. + 2 \langle F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle + \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U, H)}^2 \right) d\sigma \\ &\quad + 2 \int_s^t \langle Y_\varepsilon(\sigma) - \bar{Y}(\sigma), (G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))) dW(\sigma) \rangle \\ &\leq \|\zeta_s^\varepsilon - \zeta_s\|^2 + 2 \int_s^t \langle Y_\varepsilon(\sigma) - \bar{Y}(\sigma), (G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))) dW(\sigma) \rangle \\ &\quad + \int_s^t \left(2 \langle F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle + \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U, H)}^2 \right) d\sigma. \end{aligned}$$

Therefore, by Burkholder-Davis-Gundy inequality and Young's inequality we get

$$\begin{aligned} (4.14) \quad & E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 \\ &\leq E \|\zeta_s^\varepsilon - \zeta_s\|^2 + E \sup_{s \leq t \leq s+T} \int_s^t 2 \langle F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\ &\quad + E \int_s^{s+T} \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U, H)}^2 d\sigma \\ &\quad + 6E \left(\int_s^{s+T} \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U, H)}^2 \|Y_\varepsilon(\sigma) - \bar{Y}(\sigma)\|^2 d\sigma \right)^{\frac{1}{2}} \\ &\leq E \|\zeta_s^\varepsilon - \zeta_s\|^2 + E \sup_{s \leq t \leq s+T} \int_s^t 2 \langle F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\ &\quad + \frac{1}{2} E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 + 19E \int_s^{s+T} \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U, H)}^2 d\sigma. \end{aligned}$$

Then we obtain

$$(4.15) \quad E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2$$

$$\begin{aligned}
&\leq 2E\|\zeta_s^\varepsilon - \zeta_s\|^2 + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\
&\quad + 38E \int_s^{s+T} \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U, H)}^2 d\sigma \\
&=: 2E\|\zeta_s^\varepsilon - \zeta_s\|^2 + \mathbb{I}_1 + \mathbb{I}_2.
\end{aligned}$$

For \mathbb{I}_1 ,

$$\begin{aligned}
(4.16) \quad \mathbb{I}_1 &= 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\
&\leq 4E \int_s^{s+T} \|F_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - F_\varepsilon(\sigma, \bar{Y}(\sigma))\| \|Y_\varepsilon(\sigma) - \bar{Y}(\sigma)\| d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\
&\leq 4E \int_s^{s+T} L_F \|Y_\varepsilon(\sigma) - \bar{Y}(\sigma)\|^2 d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma) \rangle d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), \tilde{Y}(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\
&\leq 4E \int_s^{s+T} L_F \|Y_\varepsilon(\sigma) - \bar{Y}(\sigma)\|^2 d\sigma + \mathbb{I}_1^2 + \mathbb{I}_1^3 + \mathbb{I}_1^4.
\end{aligned}$$

For \mathbb{I}_1^2 , by (H2'), Hölder inequality, (3.9) and (4.4) we have

$$\begin{aligned}
(4.17) \quad \mathbb{I}_1^2 &:= 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma) \rangle d\sigma \\
&\leq 4E \int_s^{s+T} \|F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma))\| \|Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma)\| d\sigma \\
&\leq 4 \left[E \int_s^{s+T} (2L_F \|\bar{Y}(\sigma)\| + 2M_0)^2 d\sigma \right]^{\frac{1}{2}} \left(E \int_s^{s+T} \|Y_\varepsilon(\sigma) - \tilde{Y}_\varepsilon(\sigma)\|^2 d\sigma \right)^{\frac{1}{2}} \\
&\leq C_T (1 + E\|\zeta_s\|^2) \delta^{\frac{1}{4}}.
\end{aligned}$$

For \mathbb{I}_1^4 , similar to \mathbb{I}_1^2 , we get

$$\begin{aligned}
(4.18) \quad \mathbb{I}_1^4 &:= 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), \tilde{Y}(\sigma) - \bar{Y}(\sigma) \rangle d\sigma \\
&\leq C_T (1 + E\|\zeta_s\|^2) \delta^{\frac{1}{4}}.
\end{aligned}$$

For \mathbb{I}_1^3 , by (H2') and Hölder inequality we have

$$(4.19) \quad \mathbb{I}_1^3 := 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma$$

$$\begin{aligned}
&= 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \bar{Y}(\sigma)) - F_\varepsilon(\sigma, \tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{F}(\tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle \bar{F}(\tilde{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&\leq 4E \int_s^{s+T} \|F_\varepsilon(\sigma, \bar{Y}(\sigma)) - F_\varepsilon(\sigma, \tilde{Y}(\sigma))\| \|\tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma)\| d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{F}(\tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&\quad + 4E \int_s^{s+T} \|\bar{F}(\tilde{Y}(\sigma)) - \bar{F}(\bar{Y}(\sigma))\| \|\tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma)\| d\sigma \\
&\leq 8E \int_s^{s+T} L_F \|\tilde{Y}(\sigma) - \bar{Y}(\sigma)\| \|\tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma)\| d\sigma \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{F}(\tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&\leq 8L_F \left(E \int_s^{s+T} \|\tilde{Y}(\sigma) - \bar{Y}(\sigma)\|^2 d\sigma \right)^{\frac{1}{2}} \left(E \int_s^{s+T} \|\tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma)\|^2 d\sigma \right)^{\frac{1}{2}} \\
&\quad + 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{F}(\tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&\leq C_T (1 + E \|\zeta_s\|^2) \delta^{\frac{1}{4}} + \mathbb{I}_1^{3,2}.
\end{aligned}$$

Denote by $t(s, \delta) := s + \lceil \frac{t-s}{\delta} \rceil \delta$, where $\lceil \frac{t-s}{\delta} \rceil$ is the integer part of $\frac{t-s}{\delta}$. Now we estimate $\mathbb{I}_1^{3,2} := 4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{F}(\tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma$.

(4.20)

$$\begin{aligned}
&4E \sup_{s \leq t \leq s+T} \int_s^t \langle F_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{F}(\tilde{Y}(\sigma)), \tilde{Y}_\varepsilon(\sigma) - \tilde{Y}(\sigma) \rangle d\sigma \\
&= 4E \sup_{s \leq t \leq s+T} \left\{ \sum_{k=0}^{\lceil \frac{t-s}{\delta} \rceil - 1} \int_{s+k\delta}^{s+(k+1)\delta} \langle F_\varepsilon(\sigma, \bar{Y}(s+k\delta)) - \bar{F}(\bar{Y}(s+k\delta)), Y_\varepsilon(s+k\delta) - \bar{Y}(s+k\delta) \rangle d\sigma \right. \\
&\quad \left. + \int_{t(s,\delta)}^t \langle F_\varepsilon(\sigma, \bar{Y}(t(s,\delta))) - \bar{F}(\bar{Y}(t(s,\delta))), Y_\varepsilon(t(s,\delta)) - \bar{Y}(t(s,\delta)) \rangle d\sigma \right\} \\
&\leq 4E \sup_{s \leq t \leq s+T} \left\{ \sum_{k=0}^{\lceil \frac{t-s}{\delta} \rceil - 1} \left\langle \int_{s+k\delta}^{s+(k+1)\delta} F_\varepsilon(\sigma, \bar{Y}(s+k\delta)) - \bar{F}(\bar{Y}(s+k\delta)) d\sigma, Y_\varepsilon(s+k\delta) - \bar{Y}(s+k\delta) \right\rangle \right. \\
&\quad \left. + \int_{t(s,\delta)}^t \|F_\varepsilon(\sigma, \bar{Y}(t(s,\delta))) - \bar{F}(\bar{Y}(t(s,\delta)))\| \|Y_\varepsilon(t(s,\delta)) - \bar{Y}(t(s,\delta))\| d\sigma \right\} \\
&\leq 4E \sup_{s \leq t \leq s+T} \left\{ \sum_{k=0}^{\lceil \frac{t-s}{\delta} \rceil - 1} \left\| \int_{s+k\delta}^{s+(k+1)\delta} F_\varepsilon(\sigma, \bar{Y}(s+k\delta)) - \bar{F}(\bar{Y}(s+k\delta)) d\sigma \right\| \right.
\end{aligned}$$

$$\begin{aligned}
& \times \|Y_\varepsilon(s+k\delta) - \bar{Y}(s+k\delta)\| \Big\} + C_T (1 + E\|\zeta_s\|^2) \delta \\
& \leq \frac{4T}{\delta} \max_{0 \leq k \leq T(\delta)-1} \left(E \left\| \int_{s+k\delta}^{s+(k+1)\delta} F_\varepsilon(\sigma, \bar{Y}(s+k\delta)) - \bar{F}(\bar{Y}(s+k\delta)) d\sigma \right\|^2 \right)^{\frac{1}{2}} C_T (1 + E\|\zeta_s\|^2) \\
& \quad + C_T (1 + E\|\zeta_s\|^2) \delta \\
& \leq \frac{4T}{\delta} \max_{0 \leq k \leq T(\delta)-1} \delta \omega_1 \left(\frac{\delta}{\varepsilon} \right) \left(E (1 + \|\bar{Y}(s+k\delta)\|)^2 \right)^{\frac{1}{2}} C_T (1 + E\|\zeta_s\|^2) + C_T (1 + E\|\zeta_s\|^2) \delta \\
& \leq C_T (1 + E\|\zeta_s\|^2) \left(\omega_1 \left(\frac{\delta}{\varepsilon} \right) + \delta \right).
\end{aligned}$$

Combining (4.19) and (4.20), we deduce

$$(4.21) \quad \mathbb{I}_1^3 \leq C_T (1 + E\|\zeta_s\|^2) \left(\delta^{\frac{1}{4}} + \delta + \omega_1 \left(\frac{\delta}{\varepsilon} \right) \right).$$

Therefore, (4.16)–(4.18) and (4.21) yield

$$(4.22) \quad \mathbb{I}_1 \leq 4L_F \int_s^{s+T} E \sup_{s \leq u \leq \sigma} \|Y_\varepsilon(u) - \bar{Y}(u)\|^2 d\sigma + C_T (1 + E\|\zeta_s\|^2) \left(\delta^{\frac{1}{4}} + \delta + \omega_1 \left(\frac{\delta}{\varepsilon} \right) \right).$$

Now we estimate \mathbb{I}_2 .

$$\begin{aligned}
(4.23) \quad \mathbb{I}_2 & := 38E \int_s^{s+T} \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \leq 76E \int_s^{s+T} \|G_\varepsilon(\sigma, Y_\varepsilon(\sigma)) - G_\varepsilon(\sigma, \bar{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \quad + 76E \int_s^{s+T} \|G_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \leq 76L_G^2 E \int_s^{s+T} \|Y_\varepsilon(\sigma) - \bar{Y}(\sigma)\|^2 d\sigma \\
& \quad + 76E \int_s^{s+T} \|G_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \leq 76L_G^2 \int_s^{s+T} E \sup_{s \leq u \leq \sigma} \|Y_\varepsilon(u) - \bar{Y}(u)\|^2 d\sigma + \mathbb{I}_2^2.
\end{aligned}$$

Denote by $T(\delta) := \lceil \frac{T}{\delta} \rceil$. For \mathbb{I}_2^2 , it follows from (H2'), (3.9) and Remark 4.1 that

$$(4.24)$$

$$\begin{aligned}
\mathbb{I}_2^2 & := 76E \int_s^{s+T} \|G_\varepsilon(\sigma, \bar{Y}(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \leq 228E \int_s^{s+T} \|G_\varepsilon(\sigma, \bar{Y}(\sigma)) - G_\varepsilon(\sigma, \tilde{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \quad + 228E \int_s^{s+T} \|G_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{G}(\tilde{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
& \quad + 228E \int_s^{s+T} \|\bar{G}(\tilde{Y}(\sigma)) - \bar{G}(\bar{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma
\end{aligned}$$

$$\begin{aligned}
&\leq 456E \int_s^{s+T} L_G^2 \|\bar{Y}(\sigma) - \tilde{Y}(\sigma)\|^2 d\sigma + 228E \int_s^{s+T} \|G_\varepsilon(\sigma, \tilde{Y}(\sigma)) - \bar{G}(\tilde{Y}(\sigma))\|_{L_2(U,H)}^2 d\sigma \\
&\leq C_T (1 + E\|\zeta_s\|^2) \delta^{\frac{1}{2}} + 228E \sum_{k=0}^{T(\delta)-1} \int_{s+k\delta}^{s+(k+1)\delta} \|G_\varepsilon(\sigma, \bar{Y}(s+k\delta)) - \bar{G}(\bar{Y}(s+k\delta))\|_{L_2(U,H)}^2 d\sigma \\
&\quad + 228E \int_{s+T(\delta)\delta}^{s+T} \|G_\varepsilon(\sigma, \bar{Y}(s+T(\delta)\delta)) - \bar{G}(\bar{Y}(s+T(\delta)\delta))\|_{L_2(U,H)}^2 d\sigma \\
&\leq C_T (1 + E\|\zeta_s\|^2) \delta^{\frac{1}{2}} + 228 \sum_{k=0}^{T(\delta)-1} \delta \omega_2 \left(\frac{\delta}{\varepsilon} \right) E (1 + \|\bar{Y}(s+k\delta)\|^2) \\
&\quad + C_{L_G, M_0} E \int_{s+T(\delta)\delta}^{s+T} (\|\bar{Y}(s+T(\delta)\delta)\|^2 + 1) d\sigma \\
&\leq C_T (1 + E\|\zeta_s\|^2) \left(\delta^{\frac{1}{2}} + \delta + \omega_2 \left(\frac{\delta}{\varepsilon} \right) \right).
\end{aligned}$$

Therefore,

(4.25)

$$\mathbb{I}_2 \leq 76L_G^2 \int_s^{s+T} E \sup_{s \leq u \leq \sigma} \|Y_\varepsilon(u) - \bar{Y}(u)\|^2 d\sigma + C_T (1 + E\|\zeta_s\|^2) \left(\delta^{\frac{1}{2}} + \delta + \omega_2 \left(\frac{\delta}{\varepsilon} \right) \right).$$

Combining (4.15), (4.22) and (4.25), we get

$$\begin{aligned}
(4.26) \quad &E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 \\
&\leq 2E\|\zeta_s^\varepsilon - \zeta_s\|^2 + (4L_F + 76L_G^2) \int_s^{s+T} E \sup_{s \leq u \leq \sigma} \|Y_\varepsilon(u) - \bar{Y}(u)\|^2 d\sigma \\
&\quad + C_T (1 + E\|\zeta_s\|^2) \left(\delta^{\frac{1}{4}} + \delta^{\frac{1}{2}} + \delta + \omega_1 \left(\frac{\delta}{\varepsilon} \right) + \omega_2 \left(\frac{\delta}{\varepsilon} \right) \right).
\end{aligned}$$

It follows from Gronwall's lemma that

$$\begin{aligned}
(4.27) \quad &E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 \leq \left[2E\|\zeta_s^\varepsilon - \zeta_s\|^2 + C_T (1 + E\|\zeta_s\|^2) \left(\delta^{\frac{1}{4}} + \delta^{\frac{1}{2}} + \delta \right. \right. \\
&\quad \left. \left. + \omega_1 \left(\frac{\delta}{\varepsilon} \right) + \omega_2 \left(\frac{\delta}{\varepsilon} \right) \right) \right] \exp \{ (4L_F + 76L_G^2) T \}.
\end{aligned}$$

Taking $\delta = \sqrt{\varepsilon}$ and letting $\varepsilon \rightarrow 0$ in (4.27), we have

$$\lim_{\varepsilon \rightarrow 0} E \sup_{s \leq t \leq s+T} \|Y_\varepsilon(t) - \bar{Y}(t)\|^2 = 0.$$

□

Remark 4.4. As mentioned in Introduction, there are three types of averaging principle and most existing works (except for [9, 22]) on *stochastic* averaging focus on the first Bogolyubov theorem. But to the best of our knowledge, the above result is new and hence interesting on its own rights. Meanwhile, it is helpful for us to establish the second Bogolyubov theorem and global averaging principle in what follows.

With the help of Theorem 4.3, we can now establish the second Bogolyubov theorem.

Theorem 4.5. *Suppose that conditions (G1)–(G2), (H1), (H2') and (H3)–(H6) hold. Assume further that $2\lambda - 2\lambda_F - L_G^2 \geq 0$. If $\lambda' > 0$ or $2\lambda - 2\lambda_F - L_G^2 > 0$ then for any $0 < \varepsilon \leq 1$*

- (i) *equation (4.2) has a unique solution $X_\varepsilon \in C_b(\mathbb{R}, L^2(\Omega, \mathbb{P}; H))$;*

(ii) the L^2 -bounded solution X_ε of (4.2) is strongly compatible in distribution, i.e. $\mathfrak{M}_{(F_\varepsilon, G_\varepsilon)} \subseteq \tilde{\mathfrak{M}}_{X_\varepsilon}$, and

$$\lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X})) = 0 \quad \text{in } Pr(C(\mathbb{R}, H)),$$

where \bar{X} is the unique stationary solution of averaged equation (4.3).

Proof. (i) follows from Theorem 3.6.

(ii) By Theorem 3.13 the bounded solution X_ε of equation (4.2) is strongly compatible in distribution, i.e. $\mathfrak{M}_{(F_\varepsilon, G_\varepsilon)} \subseteq \tilde{\mathfrak{M}}_{X_\varepsilon}$, for any $0 < \varepsilon \leq 1$.

Now we prove that $\lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon(t)), \mathcal{L}(\bar{X}(t))) = 0$ in $Pr(H)$ for any $t \in \mathbb{R}$. Take a sequence $\{\varepsilon_n\}_{n=1}^\infty \subset (0, 1]$ such that $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. Similar to Proposition 3.11, we have $\sup_{t \in \mathbb{R}, \varepsilon \in (0, 1]} E \|X_\varepsilon(t)\|_S^2 < \infty$. It follows from Chebychev's inequality and the compact imbedding $S \subset H$ that $\{\mathcal{L}(X_{\varepsilon_n}(t))\}_{n=1}^\infty$ is tight for all $t \in \mathbb{R}$. For every $r \geq 1$, according to the tightness of $\{\mathcal{L}(X_{\varepsilon_n}(-r))\}_{n=1}^\infty$, there exists a subsequence $\{\varepsilon_{n_k}\} \subset \{\varepsilon_n\}$ such that $\mathcal{L}(X_{\varepsilon_{n_k}}(-r))$ weakly converges to μ_r in $Pr(H)$. Due to the Skorohod representation theorem, there exists a sequence of random variables $\hat{\psi}^k(-r)$ and $\hat{\zeta}_r$ with laws of $\mathcal{L}(X_{\varepsilon_{n_k}}(-r))$ and μ_r respectively, defined on another probability space $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$, such that $\hat{\psi}^k(-r) \rightarrow \hat{\zeta}_r$ $\hat{\mathbb{P}}$ -a.s. It follows from (3.4) and Remark 4.1 that there exists $p > 1$ such that

$$\hat{E} \left\| \hat{\psi}^k(-r) \right\|^{2p} = \int_H \|x\|^{2p} \mathcal{L}(\hat{\psi}^k(-r))(dx) = \int_H \|x\|^{2p} \mathcal{L}(X_{\varepsilon_{n_k}}(-r))(dx) < \infty.$$

By the Vitali L^p convergence criterion, we have $\lim_{k \rightarrow \infty} \hat{E} \left\| \hat{\psi}^k(-r) - \hat{\zeta}_r \right\|^2 = 0$.

Let $\hat{\psi}^k$ be the solution of the following Cauchy problem

$$\begin{cases} dX(t) = \left(A(X(t)) + F_{\varepsilon_{n_k}}(t, X(t)) \right) dt + G_{\varepsilon_{n_k}}(t, X(t)) d\hat{W}(t) \\ X(-r) = \hat{\psi}^k(-r) \end{cases}$$

and \hat{Y}_r be the solution of the following Cauchy problem

$$\begin{cases} dX(t) = \left(A(X(t)) + \bar{F}(X(t)) \right) dt + \bar{G}(X(t)) d\hat{W}(t) \\ X(-r) = \hat{\zeta}_r, \end{cases}$$

where \hat{W} is a cylindrical Wiener process with the identity covariance operator on $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$. In view of Theorem 4.3, we get

$$\lim_{k \rightarrow \infty} \hat{E} \sup_{-r \leq s \leq t} \left\| \hat{\psi}^k(s) - \hat{Y}_r(s) \right\|^2 = 0$$

for any $t \geq -r$.

Let ζ_r be a random variable defined on $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\mathcal{L}(\zeta_r) = \mu_r$, and Y_r be the solution of

$$dX(t) = \left(A(X(t)) + \bar{F}(X(t)) \right) dt + \bar{G}(X(t)) dW(t)$$

with initial value $Y_r(-r) = \zeta_r$. Since the law of the solutions for equation (4.2) (respectively, equation (4.3)) is unique, $\mathcal{L}(\hat{\psi}^k) = \mathcal{L}(X_{\varepsilon_{n_k}})$ and $\mathcal{L}(\hat{Y}_r) = \mathcal{L}(Y_r)$ in $Pr(C([-r, +\infty), H])$. Then we have

$$(4.28) \quad \lim_{k \rightarrow \infty} d_{BL}(\mathcal{L}(X_{\varepsilon_{n_k}}), \mathcal{L}(Y_r)) = 0 \quad \text{in } Pr(C([-r, +\infty), H]).$$

It follows from the tightness of $\left\{ \mathcal{L} \left(X_{\varepsilon_{n_k}}(-r-1) \right) \right\}$ that there exists a subsequence $\{\varepsilon_{n_{k_j}}\} \subset \{\varepsilon_{n_k}\}$ such that $\mathcal{L} \left(X_{\varepsilon_{n_{k_j}}}(-r-1) \right)$ weakly converges to μ_{r+1} . We can find a random variable ζ_{r+1} on $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\mathcal{L}(\zeta_{r+1}) = \mu_{r+1}$. Let Y_{r+1} be the solution of

$$dX(t) = (A(X(t)) + \bar{F}(X(t))) dt + \bar{G}(X(t))dW(t)$$

with initial value $Y_{r+1}(-r-1) = \zeta_{r+1}$. Similar to the procedure of calculating (4.28), we get

$$\lim_{j \rightarrow \infty} d_{BL} \left(\mathcal{L}(X_{\varepsilon_{n_{k_j}}}), \mathcal{L}(Y_{r+1}) \right) = 0 \quad \text{in } Pr(C([-r-1, +\infty), H)).$$

Therefore, we have $d_{BL}(\mathcal{L}(Y_r), \mathcal{L}(Y_{r+1})) = 0$ in $Pr(C([-r, +\infty), H))$. In particular, $\mathcal{L}(Y_r(t)) = \mathcal{L}(Y_{r+1}(t))$ for all $t \geq -r$.

Define $\nu(t) := \mathcal{L}(Y_r(t))$, $t \geq -r$. We can extract a subsequence which we still denote by $\{X_{\varepsilon_{n_{k_j}}}\}$ satisfying $\lim_{j \rightarrow \infty} d_{BL} \left(\mathcal{L}(X_{\varepsilon_{n_{k_j}}}(t)), \nu(t) \right) = 0$ in $Pr(H)$ for every $t \in \mathbb{R}$. In view of Theorem 3.6, we obtain that ν is the law of the L^2 -bounded solution of (4.3). Therefore we have

$$\lim_{j \rightarrow \infty} d_{BL} \left(\mathcal{L}(X_{\varepsilon_{n_{k_j}}}(t)), \mathcal{L}(\bar{X}(t)) \right) = 0 \quad \text{in } Pr(H)$$

for every $t \in \mathbb{R}$. By the arbitrariness of $\{\varepsilon_n\}_{n=1}^\infty \subset (0, 1]$, we have

$$\lim_{\varepsilon \rightarrow 0} d_{BL} \left(\mathcal{L}(X_\varepsilon(t)), \mathcal{L}(\bar{X}(t)) \right) = 0 \quad \text{in } Pr(H).$$

Now we show that $\lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X})) = 0$ in $Pr(C(\mathbb{R}, H))$. For any $[a, b] \subset \mathbb{R}$, we have $\mathcal{L}(X_\varepsilon(a))$ converges weakly to $\mathcal{L}(X(a))$ in $Pr(H)$. In light of Skorohod representation theorem, there exist random variables $\hat{\psi}_\varepsilon(a)$ and $\hat{\psi}(a)$ defined on another probability space $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ satisfying $\lim_{\varepsilon \rightarrow 0} \hat{\psi}_\varepsilon(a) = \hat{\psi}(a)$ $\hat{\mathbb{P}}$ -a.s., where $\mathcal{L}(\hat{\psi}_\varepsilon(a)) = \mathcal{L}(X_\varepsilon(a))$ and $\mathcal{L}(\hat{\psi}(a)) = \mathcal{L}(\bar{X}(a))$. Similar to the procedure of calculating (4.28), we have

$$(4.29) \quad \lim_{\varepsilon \rightarrow 0} d_{BL} \left(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X}) \right) = 0 \quad \text{in } Pr(C([a, b], H)).$$

The proof is complete. \square

Corollary 4.6. *Under the conditions of Theorem 4.5 the following statements hold:*

- (i) *If $F \in C(\mathbb{R} \times H, H)$ and $G \in C(\mathbb{R} \times H, L_2(U, H))$ are jointly stationary (respectively, T -periodic, quasi-periodic with the spectrum of frequencies ν_1, \dots, ν_k , almost periodic, almost automorphic, Birkhoff recurrent, Lagrange stable, Levitan almost periodic, almost recurrent, Poisson stable) in t uniformly with respect to x on each bounded subset, then equation (4.2) has a unique solution $X_\varepsilon \in C_b(\mathbb{R}, L^2(\Omega, \mathbb{P}; H))$ which is stationary (respectively, T -periodic, quasi-periodic with the spectrum of frequencies ν_1, \dots, ν_k , almost periodic, almost automorphic, Birkhoff recurrent, Lagrange stable, Levitan almost periodic, almost recurrent, Poisson stable) in distribution;*
- (ii) *If $F \in C(\mathbb{R} \times H, H)$ and $G \in C(\mathbb{R} \times H, L_2(U, H))$ are Lagrange stable and jointly pseudo-periodic (respectively, pseudo-recurrent) in t uniformly with respect to x on each bounded subset, then equation (4.2) has a unique solution $X_\varepsilon \in C_b(\mathbb{R}, L^2(\Omega, \mathbb{P}; H))$ which is pseudo-periodic (respectively, pseudo-recurrent) in distribution;*
- (iii)

$$\lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X})) = 0 \quad \text{in } Pr(C(\mathbb{R}, H)),$$

where \bar{X} is the unique stationary solution of averaged equation (4.3).

5. GLOBAL AVERAGING IN WEAK SENSE

We recall firstly that some known definitions and lemma in dynamical systems (see, e.g. [3, 7, 11, 25, 33] for more details). Let (X, d_X) and (P, d_P) be metric spaces.

Definition 5.1. A *nonautonomous dynamical system* (σ, φ) (in short, φ) consists of two ingredients:

- (i) A *dynamical system* σ on P with time set $\mathbb{T} = \mathbb{Z}$ or \mathbb{R} , i.e.
 - (1) $\sigma_0(\cdot) = Id_P$,
 - (2) $\sigma_{t+s}(p) = \sigma_t(\sigma_s(p))$ for all $t, s \in \mathbb{T}$ and $p \in P$,
 - (3) the mapping $(t, p) \mapsto \sigma_t(p)$ is continuous.
 If $\mathbb{T} = \mathbb{R}$, σ is called *flow* on P ; if $\mathbb{T} = \mathbb{R}^+$, σ is called *semiflow* on P .
- (ii) A *cocycle* $\varphi : \mathbb{T}^+ \times P \times X \rightarrow X$ satisfies
 - (1) $\varphi(0, p, x) = x$ for all $(p, x) \in P \times X$,
 - (2) $\varphi(t + s, p, x) = \varphi(t, \sigma_s(p), \varphi(s, p, x))$ for all $s, t \in \mathbb{T}^+$ and $(p, x) \in P \times X$,
 - (3) the mapping $(t, p, x) \mapsto \varphi(t, p, x)$ is continuous.

P is called the *base* or *parameter space* and X is the *fiber* or *state space*. For convenience, we also write $\sigma_t(p)$ as $\sigma_t p$.

Definition 5.2. Let (σ, φ) be a nonautonomous dynamical system with base space P and state space X . The *skew product semiflow* $\Pi : \mathbb{T}^+ \times P \times X \rightarrow P \times X$ is a semiflow of the form:

$$\Pi(t, (p, x)) := (\sigma_t p, \varphi(t, p, x)).$$

Definition 5.3. Define $\mathfrak{X} := P \times X$. A nonempty compact subset \mathfrak{A} of \mathfrak{X} is called *global attractor* for skew product semiflow Π , if

- (i) $\Pi(t, \mathfrak{A}) = \mathfrak{A}$ for all $t \in \mathbb{T}^+$,
- (ii) $\lim_{t \rightarrow +\infty} \text{dist}_{\mathfrak{X}}(\Pi(t, D), \mathfrak{A}) = 0$ for every nonempty bounded subset D of \mathfrak{X} ,

where $\text{dist}_{\mathfrak{X}}(A, B)$ is the Hausdorff semi-metric between sets A and B , i.e. $\text{dist}_{\mathfrak{X}}(A, B) := \sup_{x \in A} d(x, B)$ with $d(x, B) := \inf_{y \in B} d_{\mathfrak{X}}(x, y)$. Here $d_{\mathfrak{X}}((p_1, x_1), (p_2, x_2)) = d_P(p_1, p_2) + d_X(x_1, x_2)$ for all $(p_1, x_1), (p_2, x_2) \in P \times X$.

Lemma 5.4 (see, e.g. [11]). *Let $\{S(t)\}_{t \geq 0}$ be a semiflow in a complete metric space \mathcal{X} having a compact attracting set $K \subset \mathcal{X}$, i.e.*

$$\lim_{t \rightarrow +\infty} \text{dist}_{\mathcal{X}}(S(t)B, K) = 0$$

for all bounded set $B \subset \mathcal{X}$. Then $\{S(t)\}_{t \geq 0}$ has a global attractor $\mathcal{A} := \omega(K)$. Where $\omega(K)$ is the ω -limit set of K , i.e. $\omega(K) := \bigcap_{t \geq 0} \overline{\bigcup_{s \geq t} S(s)K}$.

Definition 5.5. A family $D := \{D_p : p \in P\}$ of subsets of X is called a *non-autonomous set*. If every fiber D_p is compact, then $D = \{D_p : p \in P\}$ is called *non-autonomous compact set*.

Definition 5.6 (see, e.g. [11]). A compact set $\mathcal{A} \subset X$ is called the *uniform attractor* (with respect to $p \in P$) of cocycle φ if the following conditions are fulfilled:

- (i) The set \mathcal{A} is uniformly attracting, i.e.

$$\lim_{t \rightarrow +\infty} \sup_{p \in P} \text{dist}_X(\varphi(t, p, B), \mathcal{A}) = 0$$

for every bounded set $B \subset X$.

- (ii) If \mathcal{A}_1 is another closed uniformly attracting set, then $\mathcal{A} \subset \mathcal{A}_1$.

Remark 5.7. It follows from Definition 5.6 (ii) that the uniform attractor is unique.

Denote by $\mathbb{F} := (F, G) \in BUC(\mathbb{R} \times H, H) \times BUC(\mathbb{R} \times H, L_2(U, H))$. Recall that $F^\tau(t, x) = F(t + \tau, x)$ for all $(t, x) \in \mathbb{R} \times H$,

$$H(\mathbb{F}) = \overline{\{\mathbb{F}^\tau = (F^\tau, G^\tau) : \tau \in \mathbb{R}\}} \subset BUC(\mathbb{R} \times H, H) \times BUC(\mathbb{R} \times H, L_2(U, H)),$$

and $(H(\mathbb{F}), \mathbb{R}, \sigma)$ is a shift dynamical system. Here $\sigma : \mathbb{R} \times H(\mathbb{F}) \rightarrow H(\mathbb{F}), (\tau, \mathbb{F}) \mapsto \mathbb{F}^\tau$.

Let $X(t, s, x), t \geq s$ be the solution of equation

$$(5.1) \quad dX(t) = (A(X(t)) + F(t, X(t))) dt + G(t, X(t))dW(t)$$

with initial condition $X(s, s, x) = x$. Define $P_{\mathbb{F}}(s, x, t, dy) := \mathbb{P} \circ (X(t, s, x))^{-1}(dy)$. Then we can associate a mapping $P^*(t, \mathbb{F}, \cdot) : Pr(H) \rightarrow Pr(H)$ defined by

$$P^*(t, \mathbb{F}, \mu)(B) := \int_H P_{\mathbb{F}}(0, x, t, B) \mu(dx)$$

for all $\mu \in Pr(H)$ and $B \in \mathcal{B}(H)$. We write $Pr_2(H)$ to mean the space of probability measures $\mu \in Pr(H)$ such that $\int_H \|z\|^2 \mu(dz) < \infty$. We say that $B \subset Pr_2(H)$ is *bounded* if there exists a constant $r > 0$ such that $\int_H \|z\|^2 \mu(dz) \leq r^2$ for all $\mu \in B$. In the following, we define

$$B_r := \left\{ \mu \in Pr_2(H) : \int_H \|z\|^2 \mu(dz) \leq r^2 \right\}$$

and

$$\mathcal{O}_\rho(B) := \{\mu \in Pr_2(H) : d(\mu, B) < \rho\}$$

for all $r, \rho > 0$, where $d(\mu, B) := \inf_{\nu \in B} d_{BL}(\mu, \nu)$.

Lemma 5.8. *Consider equation (5.1). Assume that conditions (H1), (H2') and (H3)–(H4) hold. Then P^* is a cocycle on $(H(\mathbb{F}), \mathbb{R}, \sigma)$ with fiber $Pr_2(H)$.*

Proof. It follows from Lemma 3.12 that P^* is a continuous mapping from $\mathbb{R}^+ \times H(\mathbb{F}) \times Pr_2(H)$ into $Pr_2(H)$. For any $\mu \in Pr_2(H)$, $t, \tau \in \mathbb{R}^+$ and $\tilde{\mathbb{F}} \in H(\mathbb{F})$, according to the uniqueness in law of the solutions for equation (5.1), we have $P^*(t + \tau, \tilde{\mathbb{F}}, \mu) = P^*(t, \sigma_\tau \tilde{\mathbb{F}}, P^*(\tau, \tilde{\mathbb{F}}, \mu))$. And by the definition of P^* we have $P^*(0, \tilde{\mathbb{F}}, \cdot) = Id_{Pr_2(H)}$ for all $\tilde{\mathbb{F}} \in H(\mathbb{F})$. \square

Corollary 5.9. *Under conditions of Lemma 5.8, the mapping given by*

$$\Pi : \mathbb{R}^+ \times H(\mathbb{F}) \times Pr_2(H) \rightarrow H(\mathbb{F}) \times Pr_2(H),$$

$$\Pi(t, (\tilde{\mathbb{F}}, \mu)) := \left(\sigma_t \tilde{\mathbb{F}}, P^*(t, \tilde{\mathbb{F}}, \mu) \right)$$

is a continuous skew-product semiflow.

For any given $\tilde{\mathbb{F}} \in H(\mathbb{F})$, suppose that (H1), (H2'), (H3)–(H4) hold and $2\lambda - 2\lambda_F - L_G^2 \geq 0$. If $\lambda' > 0$ or $2\lambda - 2\lambda_F - L_G^2 > 0$, then equation (5.1) has a unique L^2 -bounded solution $X_{\tilde{\mathbb{F}}}$ with the distribution $\mathcal{L}(X_{\tilde{\mathbb{F}}}(t)) =: \mu_{\tilde{\mathbb{F}}}(t), t \in \mathbb{R}$. In the following, we denote by $\mathfrak{X} := H(\mathbb{F}) \times Pr_2(H)$.

Lemma 5.10. *Let $\bar{F}(x) := \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} F(s, x) ds$ and $\lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \|G(s, x) - \bar{G}(x)\|^2 ds = 0$ uniformly with respect to $t \in \mathbb{R}$. If $H(\mathbb{F})$ is compact, then for any $\tilde{\mathbb{F}} = (\tilde{F}, \tilde{G}) \in H(\mathbb{F})$ we have*

$$(5.2) \quad \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \tilde{F}(s, x) ds = \bar{F}(x)$$

and

$$(5.3) \quad \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \|\tilde{G}(s, x) - \bar{G}(x)\|^2 ds = 0$$

uniformly with respect to $t \in \mathbb{R}$.

Proof. By Remark 4.1 (iii), there exists $\omega_1 \in \Psi$ such that

$$\frac{1}{T} \left\| \int_t^{t+T} F(s, x) - \bar{F}(x) ds \right\| \leq \omega_1(T) (1 + \|x\|)$$

for all $x \in H$ and $t \in \mathbb{R}$. Here $\lim_{T \rightarrow +\infty} \omega_1(T) = 0$. Given $\tilde{F} \in H(F)$, there exists $\{t_n\} \subset \mathbb{R}$ such that

$$\lim_{n \rightarrow \infty} \sup_{|t| \leq l, \|x\| \leq r} \|\tilde{F}(t, x) - F(t + t_n, x)\| = 0$$

for all $l, r > 0$. Then by Lebesgue dominated convergence theorem, we have

$$\begin{aligned} \frac{1}{T} \left\| \int_t^{t+T} \tilde{F}(s, x) - \bar{F}(x) ds \right\| &= \frac{1}{T} \left\| \int_t^{t+T} \lim_{n \rightarrow \infty} F(s + t_n, x) - \bar{F}(x) ds \right\| \\ &= \lim_{n \rightarrow \infty} \frac{1}{T} \left\| \int_t^{t+T} F(s + t_n, x) - \bar{F}(x) ds \right\| \\ &\leq \omega_1(T) (1 + \|x\|). \end{aligned}$$

Therefore, (5.2) follows from Remark 4.1 (iii). The proof of (5.3) is similar. \square

Remark 5.11. It follows from Remark 3.9 that estimates (3.4), (3.7), (3.9), (3.17), (3.18) and (4.27) hold uniformly for all $\tilde{\mathbb{F}} \in H(\mathbb{F})$ and $\varepsilon \in (0, 1]$.

Proposition 5.12. *Consider equation (5.1). Assume that conditions (H1), (H2'), (H3)–(H4), (H6) hold, and $2\lambda - 2\lambda_F - L_G^2 \geq 0$. Suppose further that $\lambda' > 0$ or $2\lambda - 2\lambda_F - L_G^2 > 0$. Then we have the following results.*

(i) Denote $\mathfrak{A}_{\tilde{\mathbb{F}}} := \overline{\{\mu_{\tilde{\mathbb{F}}}(t) \in Pr_2(H) : t \in \mathbb{R}\}}$. Then

$$P^*(t, \tilde{\mathbb{F}}, \mathfrak{A}_{\tilde{\mathbb{F}}}) = \mathfrak{A}_{\sigma_t \tilde{\mathbb{F}}}$$

for all $t \in \mathbb{R}^+$ and $\tilde{\mathbb{F}} \in H(\mathbb{F})$.

(ii) If $H(\mathbb{F})$ is compact, then the skew product semiflow Π admits a global attractor $\mathfrak{A} := \omega\left(H(\mathbb{F}) \times \overline{\cup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}}}\right)$. Moreover, $\Pi_2 \mathfrak{A}$ is the uniform attractor of cocycle P^* .

Here $\Pi_2(\tilde{\mathbb{F}}, \mu) := \mu$ for all $(\tilde{\mathbb{F}}, \mu) \in H(\mathbb{F}) \times Pr_2(H)$.

Proof. (i) Given $t \in \mathbb{R}^+$ and $\tilde{\mathbb{F}} \in H(\mathbb{F})$, let $X_{\sigma_t \tilde{\mathbb{F}}}$ be the unique L^2 -bounded solution of equation

$$dX(s) = \left(A(X(s)) + \tilde{F}(s + t, X(s)) \right) ds + \tilde{G}(s + t, X(s)) dW(s).$$

Note that $\mathcal{L}(X_{\tilde{\mathbb{F}}}(s + t)) = \mathcal{L}(X_{\sigma_t \tilde{\mathbb{F}}}(s))$ for all $s \in \mathbb{R}$. Consequently, $P^*(t, \tilde{\mathbb{F}}, \mathfrak{A}_{\tilde{\mathbb{F}}}) = \mathfrak{A}_{\sigma_t \tilde{\mathbb{F}}}$.

(ii) Employing Proposition 3.11 and Remark 5.11, there exists a constant $R > 0$ such that

$$\bigcup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}} \subset \left\{ \mu \in Pr_2(H) : \int_S \|z\|^2 \mu(dz) < R^2 \right\}.$$

According to the Chebychev's inequality and the compactness of the inclusion $S \subset H$, $\bigcup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}}$ is compact in $Pr(H)$.

Let $r > 0$ be an arbitrary constant. For any $\mu \in B_r$, take a random variable ξ such that $\mathcal{L}(\xi) = \mu$. Let $Y(t, \xi), t \geq 0$ be the solution to

$$Y(t, \xi) = \xi + \int_0^t \left(A(Y(s, \xi)) + \tilde{F}(s, Y(s, \xi)) \right) ds + \int_0^t \tilde{G}(s, Y(s, \xi)) dW(s).$$

In view of Theorem 3.8, we have

$$E\|Y(t, \xi) - X_{\tilde{\mathbb{F}}}(t)\|^2 \leq \begin{cases} E\|\xi - X_{\tilde{\mathbb{F}}}(s)\|_H^2 \wedge \{\lambda'(r-2)(t-s)\}^{-\frac{2}{r-2}}, & \text{if } \lambda' > 0 \\ e^{-(2\lambda-2\lambda_F-L_G^2)(t-s)} E\|\xi - X_{\tilde{\mathbb{F}}}(s)\|_H^2, & \text{if } 2\lambda - 2\lambda_F - L_G^2 > 0. \end{cases}$$

Therefore, $\lim_{t \rightarrow +\infty} \sup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \text{dist}_{Pr_2(H)} \left(P^*(t, \tilde{\mathbb{F}}, \mu), \overline{\cup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}}} \right) = 0$ uniformly with respect to $\mu \in B_r$, i.e. $\overline{\cup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}}}$ is a compact uniformly attracting set. Obviously, $H(\mathbb{F}) \times \overline{\cup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}}}$ is a compact attracting set for Π . By Lemma 5.4, Π admits a global attractor $\mathfrak{A} := \omega \left(H(\mathbb{F}) \times \overline{\cup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \mathfrak{A}_{\tilde{\mathbb{F}}}} \right)$.

Let us now prove that $\Pi_2 \mathfrak{A}$ is the uniform attractor of cocycle P^* . Let $B \subset Pr_2(H)$ be bounded, then $H(\mathbb{F}) \times B$ is bounded in $H(\mathbb{F}) \times Pr_2(H)$. Therefore,

$$\begin{aligned} \text{dist}_{Pr_2(H)}(P^*(t, \tilde{\mathbb{F}}, B), \Pi_2 \mathfrak{A}) &\leq \text{dist}_{\mathfrak{X}}(H(\mathbb{F}) \times P^*(t, \tilde{\mathbb{F}}, B), \mathfrak{A}) \\ &= \text{dist}_{\mathfrak{X}}(\Pi(t, H(\mathbb{F}) \times B), \mathfrak{A}) \rightarrow 0 \quad \text{as } t \rightarrow +\infty. \end{aligned}$$

Next we verify the minimality property. Denote by $\omega_{H(\mathbb{F})}(B) := \overline{\cap_{t \geq 0} \cup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \cup_{s \geq t} P^*(s, \tilde{\mathbb{F}}, B)}$. Then $\mu \in \omega_{H(\mathbb{F})}(B)$ if and only if there exist $\{\nu_n\} \subset B$, $\{\mathbb{F}_n\} \subset H(\mathbb{F})$ and $\{t_n\} \subset \mathbb{R}_+$ such that $t_n \rightarrow +\infty$ and $P^*(t_n, \mathbb{F}_n, \nu_n) \rightarrow \mu$ as $n \rightarrow +\infty$. Let \mathcal{A}_1 be a closed uniformly attracting set. Then we show that $\omega_{H(\mathbb{F})}(\Pi_2 \mathfrak{A}) \subset \mathcal{A}_1$. Indeed, if this is false, i.e. $\omega_{H(\mathbb{F})}(\Pi_2 \mathfrak{A}) \not\subset \mathcal{A}_1$. Take $\mu \in \omega_{H(\mathbb{F})}(\Pi_2 \mathfrak{A}) \setminus \mathcal{A}_1$, there exist $\{\nu_n\} \subset \Pi_2 \mathfrak{A}$, $\{\mathbb{F}_n\} \subset H(\mathbb{F})$ and $\{t_n\} \subset \mathbb{R}_+$ such that $t_n \rightarrow +\infty$ and $P^*(t_n, \mathbb{F}_n, \nu_n) \rightarrow \mu$ as $n \rightarrow +\infty$. Hence we have

$$\begin{aligned} 0 < d(\mu, \mathcal{A}_1) &\leq \lim_{n \rightarrow +\infty} d(P^*(t_n, \mathbb{F}_n, \nu_n), \mathcal{A}_1) \\ &\leq \lim_{n \rightarrow +\infty} \text{dist}_{Pr_2(H)}(P^*(t_n, \mathbb{F}_n, \Pi_2 \mathfrak{A}), \mathcal{A}_1) \\ &\leq \lim_{n \rightarrow +\infty} \sup_{\tilde{\mathbb{F}} \in H(\mathbb{F})} \text{dist}_{Pr_2(H)}(P^*(t_n, \tilde{\mathbb{F}}, \Pi_2 \mathfrak{A}), \mathcal{A}_1) = 0, \end{aligned}$$

a contradiction. On the other hand, for any $(\tilde{\mathbb{F}}, \mu) \in \omega(H(\mathbb{F}) \times \Pi_2 \mathfrak{A}) = \mathfrak{A}$, there exist $\{\nu_n\} \subset \Pi_2 \mathfrak{A}$, $\{\mathbb{F}_n\} \subset H(\mathbb{F})$, $\{t_n\} \subset \mathbb{R}_+$ such that $P^*(t_n, \mathbb{F}_n, \nu_n) \rightarrow \mu$ and $\sigma_{t_n} \mathbb{F}_n \rightarrow \tilde{\mathbb{F}}$ as $n \rightarrow +\infty$. Then $\mu \in \omega_{H(\mathbb{F})}(\Pi_2 \mathfrak{A})$. Therefore, $\Pi_2 \mathfrak{A} \subset \mathcal{A}_1$.

The proof is complete. \square

Remark 5.13. It is known that $H(\mathbb{F})$ is compact provided \mathbb{F} is Birkhoff recurrent, which includes periodic, quasi-periodic, almost periodic, almost automorphic as special cases.

Next we prove the global averaging principle for strongly monotone SPDEs.

Theorem 5.14. *Suppose that $2\lambda - 2\lambda_F - L_G^2 \geq 0$, (G1)–(G2), (H1), (H2'), (H3)–(H4) and (H6) hold. Assume further that $\lambda' > 0$ or $2\lambda - 2\lambda_F - L_G^2 > 0$. If $H(\mathbb{F})$ is compact, then*

- (i) *the cocycle P_ε^* associated with SPDE (4.2) has a uniform attractor \mathcal{A}^ε for any $0 < \varepsilon \leq 1$;*
- (ii) *the cocycle \bar{P}^* associated with SPDE (4.3) has a uniform attractor $\bar{\mathcal{A}}$, which is a singleton set;*
- (iii) *for arbitrary large R_1 and small $\rho > 0$ there exist $\varepsilon_0 = \varepsilon_0(R_1, \rho)$ and $T = T(R_1, \rho)$ such that for all $\varepsilon \leq \varepsilon_0$, $t \geq T$*

$$(5.4) \quad P_\varepsilon^*(t, \mathbb{F}, B_{R_1}) \subset \mathcal{O}_\rho(\bar{\mathcal{A}}).$$

In particular,

$$(5.5) \quad \lim_{\varepsilon \rightarrow 0} \text{dist}_{Pr_2(H)}(\mathcal{A}^\varepsilon, \bar{\mathcal{A}}) = 0.$$

Proof. (i)–(ii) It follows from Proposition 5.12 that P_ε^* and \bar{P}^* admit uniform attractors, and $\bar{\mathcal{A}} = \{\mathcal{L}(\bar{X}(0))\} \in Pr_2(H)$. Here $\bar{X}(t), t \in \mathbb{R}$ is the unique stationary solution to averaged equation (4.3).

(iii) It follows from Theorem 3.8 that there exists $\delta, 0 < \delta < \frac{\rho}{2}$ such that

$$(5.6) \quad \bar{P}^*(t, \mathcal{O}_\delta(\bar{\mathcal{A}})) \subset \mathcal{O}_{\frac{\rho}{2}}(\bar{\mathcal{A}})$$

for all $t \geq 0$. Fix R_1 large enough. In view of (3.4), there exists $T_0 > 0$ such that

$$(5.7) \quad P_\varepsilon^*(t, \mathbb{F}, B_{R_1}) \subset B_{R_1}$$

for all $t \geq T_0$. Since $\bar{\mathcal{A}}$ is attractor, we can choose $T_1 = T_1(R_1, \rho)$ so large such that

$$(5.8) \quad \bar{P}^*(t, B_{R_1}) \subset \mathcal{O}_{\frac{\delta}{2}}(\bar{\mathcal{A}})$$

for all $t \geq T_1$. Denote by $T := \max\{T_0, T_1\}$. Employing (4.27), we have

$$(5.9) \quad \sup_{0 \leq t \leq T} d\left(P_\varepsilon^*(t, \tilde{\mathbb{F}}, \mu), \bar{P}^*(t, \mu)\right) < \eta(T, R_1)(\varepsilon)$$

for all $\mu \in B_{R_1}$ and $\tilde{\mathbb{F}} \in H(\mathbb{F})$, where $\eta(T, R_1)(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. Then, there exists $\varepsilon_0 = \varepsilon_0(T, R_1)$ such that $\eta(T, R_1)(\varepsilon) < \frac{\delta}{2}$ for all $\varepsilon \leq \varepsilon_0$.

For any $\mu \in B_{R_1}$, in view of (5.7)–(5.9), we have

$$P_\varepsilon^*(T, \mathbb{F}, \mu) \in \mathcal{O}_\delta(\bar{\mathcal{A}}) \cap B_{R_1}$$

for all $\varepsilon \leq \varepsilon_0$. It can be verified that $P_\varepsilon^*(t, \mathbb{F}, \mu) \in \mathcal{O}_\rho(\bar{\mathcal{A}})$ for all $t \geq T$ and $\varepsilon \leq \varepsilon_0$. To this end, define $\mu_1^\varepsilon := P_\varepsilon^*(T, \mathbb{F}, \mu)$. Then $\bar{P}^*(t, \mu_1^\varepsilon) \in \mathcal{O}_{\frac{\rho}{2}}(\bar{\mathcal{A}})$ and $P_\varepsilon^*(t+T, \mathbb{F}, \mu) = P_\varepsilon^*(t, \sigma_T \mathbb{F}, \mu_1^\varepsilon)$ for all $t \geq 0$. Therefore, according to (5.8)–(5.9), we get

$$P_\varepsilon^*(2T, \mathbb{F}, \mu) \in \mathcal{O}_\delta(\bar{\mathcal{A}}) \cap B_{R_1}$$

and

$$P_\varepsilon^*(t+T, \mathbb{F}, \mu) \in \mathcal{O}_{\frac{\rho}{2}+\frac{\delta}{2}}(\bar{\mathcal{A}}) \subset \mathcal{O}_\rho(\bar{\mathcal{A}})$$

for all $t \in [0, T]$. Repeating the above procedure, we have

$$P_\varepsilon^*(t, \mathbb{F}, \mu) \in \mathcal{O}_\rho(\bar{\mathcal{A}})$$

for all $t \geq T$ and $\varepsilon \leq \varepsilon_0$.

Take R_1 large enough so that $\mathcal{A}^\varepsilon \subset B_{R_1}$, then (5.5) follows from (5.4) and Definition 5.6. \square

6. APPLICATIONS

In this section, we illustrate our theoretical results by two examples. We mainly consider the additive or linear multiplicative noise in these examples for brevity. Let $\Lambda \subset \mathbb{R}^n, n \in \mathbb{N}$ be an open bounded subset and $0 < \varepsilon \leq 1$. Denote by $f^+(t) := \max\{f(t), 0\}$ for all $t \in \mathbb{R}$ and λ_* the first eigenvalue of $-\Delta$ with the Dirichlet boundary condition.

6.1. Stochastic reaction diffusion equations. Consider the equation

$$(6.1) \quad du = (\Delta u - au|u|^{p-2} + \phi(t/\varepsilon)u + g(t/\varepsilon)) dt + \kappa u dW(t),$$

where $W(\cdot)$ is a two-sided standard real-valued Wiener process, $p \in [2, +\infty)$ and $g \in C_b(\mathbb{R}, H_0^{1,2}(\Lambda))$. Here $a > 0$ and $\kappa \in \mathbb{R}$ are constants. We define $V_1 := H_0^{1,2}(\Lambda)$, $V_2 := L^p(\Lambda)$, $H := L^2(\Lambda)$, $V := V_1 \cap V_2$ and

$$A_1(u) := \Delta u, \quad A_2(u) := -au|u|^{p-2}, \quad F(t, u) := \phi(t)u + g(t), \quad G(t, u) = \kappa u.$$

Assume that $\lambda_* - |\phi^+|_\infty - \frac{\kappa^2}{2} > 0$, then we have the following theorem.

- Theorem 6.1.** (1) *There exists a unique L^2 -bounded solution $X_\varepsilon(\cdot)$ to equation (6.1) which is globally asymptotically stable in square-mean sense for any $0 < \varepsilon \leq 1$.*
- (2) *If ϕ is almost automorphic and g is almost periodic, then the L^2 -bounded solution $X_\varepsilon(\cdot)$ is almost automorphic in distribution.*
- (3) *Let \bar{X} be the unique stationary solution of the following averaged equation*

$$(6.2) \quad du = (\Delta u - a|u|^{p-2}u + \bar{\phi}u + \bar{g}) dt + \kappa u dW(t),$$

where $\bar{\phi} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \phi(s) ds$ and $\bar{g} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} g(s) ds$ uniformly for all $t \in \mathbb{R}$.

Then

$$\lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X})) = 0 \quad \text{in } Pr(C(\mathbb{R}, L^2(\Lambda))).$$

- (4) *The cocycle P_ε^* generated by equation (6.1) has a uniform attractor \mathcal{A}^ε , and*

$$\lim_{\varepsilon \rightarrow 0} \text{dist}_{Pr_2(H)}(\mathcal{A}^\varepsilon, \bar{\mathcal{A}}) = 0.$$

Here $\bar{\mathcal{A}} := \mathcal{L}(\bar{X}(0))$ is the attractor for \bar{P}^* and $H := L^2(\Lambda)$.

Proof. (1)–(2) It suffices to show that conditions of Theorem 3.8 and Corollary 3.14 hold.

(H1) A_1 is obviously hemicontinuous. We now prove that A_2 is hemicontinuous. Let $u, v, w \in V$. For $\theta \in \mathbb{R}$, without loss of generality, we assume $|\theta| \leq 1$, then we have

$$(6.3) \quad \begin{aligned} & V_2^* \langle A_2(u + \theta v) - A_2(u), w \rangle_{V_2} \\ &= \int_{\Lambda} (-(u(\xi) + \theta v(\xi)) |u(\xi) + \theta v(\xi)|^{p-2} w(\xi) + u(\xi) |u(\xi)|^{p-2} w(\xi)) d\xi \\ &\leq \int_{\Lambda} (4(|u(\xi)|^{p-1} + |v(\xi)|^{p-1}) |w(\xi)| + |u(\xi)|^{p-1} |w(\xi)|) d\xi < \infty. \end{aligned}$$

The last inequality holds since $u, v, w \in L^p(\Lambda)$. Then $V_2^* \langle A_2(u + \theta v) - A_2(u), w \rangle_{V_2}$ converges to zero as $\theta \rightarrow 0$ by Lebesgue's dominated convergence theorem. So, (H1) holds.

(H2') For all $u, v \in V$ and $t \in \mathbb{R}$

$$V_1^* \langle A_1(u) - A_1(v), u - v \rangle_{V_1} \leq -\lambda_* \|u - v\|_H^2,$$

$$V_2^* \langle A_2(u) - A_2(v), u - v \rangle_{V_2} = -a \int_{\Lambda} (u(\xi) |u(\xi)|^{p-2} - v(\xi) |v(\xi)|^{p-2}) (u(\xi) - v(\xi)) d\xi \leq 0,$$

$$\langle F(t, u) - F(t, v), u - v \rangle \leq |\phi^+|_{\infty} \|u - v\|_H^2, \quad \|F(t, 0)\|_H \leq \sup_{t \in \mathbb{R}} \|g(t)\|_{H_0^{1,2}(\Lambda)},$$

$$\|F(t, u) - F(t, v)\|_H \leq |\phi|_{\infty} \|u - v\|_H \quad \text{and} \quad \|\kappa u - \kappa v\|^2 \leq \kappa^2 \|u - v\|^2.$$

So (H2') holds with $\lambda = \lambda_*$, $\lambda' = 0$, $\lambda_F = |\phi^+|_{\infty}$, $L_F = |\phi|_{\infty}$ and $L_G = |\kappa|$.

(H3) For all $v \in V, t \in \mathbb{R}$ we have

$$V_1^* \langle A_1(v), v \rangle_{V_1} = - \int_{\Lambda} |\nabla v(\xi)|^2 d\xi = \|v\|_H^2 - \|v\|_{V_1}^2,$$

$$V_2^* \langle A_2(v), v \rangle_{V_2} = -a \int_{\Lambda} |v(\xi)|^p d\xi = -a \|v\|_{V_2}^p.$$

Then (H3) holds with $\alpha_1 = 2, \alpha_2 = p$.

(H4) For all $u, v \in V, t \in \mathbb{R}$ we have

$$|V_1^* \langle A_1(u), v \rangle_{V_1}| \leq \|\nabla u\|_H \|\nabla v\|_H \leq \|u\|_{V_1} \|v\|_{V_1},$$

$$|V_2^* \langle A_2(u), v \rangle_{V_2}| = \left| a \int_{\Lambda} -u(\xi) |u(\xi)|^{p-2} v(\xi) d\xi \right| \leq a \|u\|_{V_2}^{p-1} \|v\|_{V_2}.$$

Therefore, we get $\|A_1(u)\|_{V_1^*} \leq \|u\|_{V_1}$ and $\|A_2(u)\|_{V_2^*} \leq a \|u\|_{V_2}^{p-1}$.

(2) In order to prove the almost automorphic property of the L^2 -bounded solution, it suffices to show that (H5)–(H6) holds. To this end, let $S := H_0^{1,2}(\Lambda)$, we define $T_n = -\Delta \left(I - \frac{\Delta}{n}\right)^{-1} = n \left(I - \left(I - \frac{\Delta}{n}\right)^{-1}\right)$. Note that T_n are continuous on $W_0^{1,2}(\Lambda)$. Since the heat semigroup $\{P_t\}_{t \geq 0}$ (generated by Δ) is contractive on $L^p(\Lambda)$, $p > 1$ and $\left(I - \frac{\Delta}{n}\right)^{-1}u = \int_0^\infty e^{-t} P_{\frac{t}{n}} u dt$, T_n are continuous on $L^p(\Lambda)$.

For all $u \in V$, $t \in \mathbb{R}$ we have

$$V_1^* \langle \Delta u, T_n u \rangle_{V_1} \leq -\lambda_* \|u\|_n^2 \quad \text{and} \quad \phi(t) \langle u, T_n u \rangle_H = \phi(t) \|u\|_n^2 \leq |\phi^+|_\infty \|u\|_n^2.$$

In view of the contractivity of $\{P_t\}_{t \geq 0}$ on $L^p(\Lambda)$, we have

$$\begin{aligned} V_2^* \langle A_2(u), T_n u \rangle_{V_2} &= \langle -a|u|^{p-2}u, nu - n \left(I - \frac{\Delta}{n}\right)^{-1} u \rangle \\ &= n \int_0^\infty e^{-t} \left(\int_\Lambda -a u(\xi) |u(\xi)|^{p-2} \left(u(\xi) - P_{\frac{t}{n}} u(\xi)\right) d\xi \right) dt \leq 0. \end{aligned}$$

Then we obtain

$$2V^* \langle A(t, u), T_n u \rangle_V + 2 \langle F(t, u), T_n u \rangle \leq -2(\lambda_* - |\phi^+|_\infty - \varepsilon) \|u\|_n^2 + C_\varepsilon \sup_{t \in \mathbb{R}} \|g(t)\|_S^2.$$

That is, (H6) holds. And (H5) is obvious.

(3)–(4) follows from Corollary 4.6 and Theorem 5.14. \square

Remark 6.2. Note that equation (6.1) is the real Ginzburg-Landau equation when $p = 4$.

6.2. Stochastic generalized porous media equations. Consider the equation

$$(6.4) \quad du = (\Delta(|u|^{p-2}u + au) + \phi(t/\varepsilon)u) dt + GdW(t),$$

where $W(\cdot)$ is a two-sided cylindrical Q -Wiener process with $Q = I$ on $L^p(\Lambda)$, $p > 2$ and $a \geq 0$, $G \in L_2(L^p(\Lambda))$. And there exists a constant $C_1 > 0$ such that $\phi(t) < -C_1$ for all $t \in \mathbb{R}$. We define

$$V := L^p(\Lambda) \subset H := W_0^{-1,2}(\Lambda) \subset V^*.$$

Theorem 6.3. (1) *There exists a unique L^2 -bounded solution $X_\varepsilon(\cdot)$ to equation (6.4), which is globally asymptotically stable in square-mean sense.*

(2) *The L^2 -bounded solution $X_\varepsilon(\cdot)$ is almost periodic in distribution provided ϕ is almost periodic.*

(3) *Let \bar{X} be the unique stationary solution of averaged equation*

$$(6.5) \quad du = (-\Delta(|u|^{p-2}u + au) + \bar{\phi}u) dt + GdW(t),$$

where $\bar{\phi} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} \phi(s) ds$ uniformly for all $t \in \mathbb{R}$. Then

$$\lim_{\varepsilon \rightarrow 0} d_{BL}(\mathcal{L}(X_\varepsilon), \mathcal{L}(\bar{X})) = 0 \quad \text{in } Pr(C(\mathbb{R}, H)).$$

(4) *The cocycle P_ε^* generated by equation (6.4) has a uniform attractor \mathcal{A}^ε , and*

$$\lim_{\varepsilon \rightarrow 0} \text{dist}_{Pr_2(H)}(\mathcal{A}^\varepsilon, \bar{\mathcal{A}}) = 0.$$

Here $\bar{\mathcal{A}} := \mathcal{L}(\bar{X}(0))$ is the attractor for \bar{P}^* .

Proof. Let $A(u) := \Delta(|u|^{p-2}u + au)$, $F(t, u) := \phi(t)u$ for all $u \in V$ and $t \in \mathbb{R}$. Fix $u \in V$, for all $v \in V$ we denote

$$V^* \langle A(u), v \rangle_V := - \int_\Lambda u(\xi) |u(\xi)|^{p-2} v(\xi) d\xi - a \int_\Lambda u(\xi) v(\xi) d\xi.$$

We first show that $A : V \rightarrow V^*$ is well-defined. Indeed, employing Hölder inequality and Young's inequality, we get

$$|v^* \langle A(u), v \rangle_V| \leq \|u\|_{L^p}^{p-1} \|v\|_{L^p} + a \left(C_p \|u\|_{L^p}^{p-1} + C_p (|\Lambda|)^{\frac{p-1}{p}} \right) \|v\|_{L^p}$$

for all $u, v \in V$, where C_p is a constant depending only on p . Therefore, $A : V \rightarrow V^*$ is well-defined and

$$(6.6) \quad \|A(u)\|_{V^*} \leq (1 + aC_p) \|u\|_{L^p}^{p-1} + C_p (|\Lambda|)^{\frac{p-1}{p}} a.$$

Similar to the proof of Theorem 6.1, it suffices to show that (H1), (H2') and (H3)–(H6) hold. Note that (H1), (H5) hold and $\lambda_F = 0$, $L_F = |\phi|_\infty$ in (H2').

(H2') For all $u, v \in V$, $t \in \mathbb{R}$ we have

$$\begin{aligned} v^* \langle A(u) - A(v), u - v \rangle_V &= -\langle u|u|^{p-2} - v|v|^{p-2}, u - v \rangle_{L^2} - a \|u - v\|_{L^2}^2 \\ &\leq -2^{2-p} \|u - v\|_{L^2}^p - a \|u - v\|_{L^2}^2 \\ &\leq -2^{2-p} \|u - v\|_H^p - a \|u - v\|_H^2. \end{aligned}$$

Therefore, (H2) holds with $r = p$, $\lambda' = 2^{2-p}$ and $\lambda = a$.

(H3) Note that for all $u \in V$, $t \in \mathbb{R}$

$$v^* \langle A(u), u \rangle_V = - \int_\Lambda u(\xi) |u(\xi)|^{p-2} u(\xi) d\xi - a \int_\Lambda u(\xi) u(\xi) d\xi \leq -\|u\|_V^p.$$

That is, (H3) holds with $\alpha = p$.

(H4) holds by (6.6) with $\alpha = p$.

(H6) Let $S = L^2(\Lambda)$ and Δ be the Laplace operator on $L^2(\Lambda)$ with the Dirichlet boundary condition. Define $T_n = -\Delta (I - \frac{\Delta}{n})^{-1} = n (I - (I - \frac{\Delta}{n})^{-1})$. Then we obtain

$$\begin{aligned} v^* \langle \Delta(u|u|^{p-2} + au) + \phi(t)u, -\Delta(I - \frac{\Delta}{n})^{-1}u \rangle_V \\ = -\langle u|u|^{p-2} + au, nu - n \int_0^\infty e^{-t} P_{\frac{t}{n}} u dt \rangle_{L^2} + \phi(t) \|u\|_n^2 \\ \leq -C_1 \|u\|_n^2. \end{aligned}$$

That is, (H6) holds. □

ACKNOWLEDGEMENTS

This work is partially supported by NSFC Grants 11522104, 11871132, 11925102, Dalian High-level Talent Innovation Project (Grant 2020RD09), and Xinghai Jieqing and DUT19TD14 funds from Dalian University of Technology.

REFERENCES

- [1] V. Bakhtin and Y. Kifer, Diffusion approximation for slow motion in fully coupled averaging. *Probab. Theory Related Fields* **129** (2004), 157–181.
- [2] N. N. Bogolyubov and Y. A. Mitropolsky, *Asymptotic Methods in the Theory of Non-linear Oscillations*. Translated from the second revised Russian edition. International Monographs on Advanced Mathematics and Physics Hindustan Publishing Corp., Delhi, Gordon and Breach Science Publishers, New York 1961, x+537 pp.
- [3] T. Caraballo, J. C. Jara, J. A. Langa and Z. Liu, Morse decomposition of attractors for non-autonomous dynamical systems. *Adv. Nonlinear Stud.* **13** (2013), 309–329.
- [4] S. Cerrai, A Khasminskii type averaging principle for stochastic reaction–diffusion equations. *Ann. Appl. Probab.* **19** (2009), 899–948.
- [5] S. Cerrai and M. Freidlin, Averaging principle for a class of stochastic reaction-diffusion equations. *Probab. Theory Related Fields* **144** (2009), 137–177.

- [6] S. Cerrai and A. Lunardi, Averaging principle for nonautonomous slow-fast systems of stochastic reaction-diffusion equations: the almost periodic case. *SIAM J. Math. Anal.* **49** (2017), 2843–2884.
- [7] D. Cheban, *Global Attractors of Nonautonomous Dynamical and Control Systems. 2nd Edition*. Interdisciplinary Mathematical Sciences, vol.18, River Edge, NJ: World Scientific, 2015, xxv+589 pp.
- [8] D. Cheban and Z. Liu, Periodic, quasi-periodic, almost periodic, almost automorphic, Birkhoff recurrent and Poisson stable solutions for stochastic differential equations. *J. Differential Equations* **269** (2020), 3652–3685.
- [9] D. Cheban and Z. Liu, Averaging principle on infinite intervals for stochastic ordinary differential equations. *Electron. Res. Arch.* **29** (2021), 2791–2817.
- [10] M. Cheng and Z. Liu, Periodic, almost periodic and almost automorphic solutions for SPDEs with monotone coefficients. *Discrete Contin. Dyn. Syst. Ser. B* doi: 10.3934/dcdsb.2021026.
- [11] V. V. Chepyzhov and M. I. Vishik, *Attractors for Equations of Mathematical Physics*. American Mathematical Society Colloquium Publications, 49. American Mathematical Society, Providence, RI, 2002, xii+363 pp.
- [12] G. Da Prato and C. Tudor, Periodic and almost periodic solutions for semilinear stochastic equations. *Stochastic Anal. Appl.* **13** (1995), 13–33.
- [13] J. Duan and W. Wang, *Effective Dynamics of Stochastic Partial Differential Equations*. Elsevier Insights. Elsevier, Amsterdam, 2014, xii+270 pp.
- [14] R. M. Dudley, *Real Analysis and Probability*. Revised reprint of the 1989 original. Cambridge Studies in Advanced Mathematics, 74. Cambridge University Press, Cambridge, 2002, x+555 pp.
- [15] M. I. Freidlin and A. D. Wentzell, Long-time behavior of weakly coupled oscillators. *J. Stat. Phys.* **123** (2006), 1311–1337.
- [16] M. I. Freidlin and A. D. Wentzell, *Random Perturbations of Dynamical Systems*. Translated from the 1979 Russian original by Joseph Szücs. Third edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 260. Springer, Heidelberg, 2012, xxviii+458 pp.
- [17] M. Fu and Z. Liu, Square-mean almost automorphic solutions for some stochastic differential equations. *Proc. Amer. Math. Soc.* **138** (2010), 3689–3701.
- [18] P. Gao, Averaging principle for complex Ginzburg-Landau equation perturbed by mixing random forces. *SIAM J. Math. Anal.* **53** (2021), 32–61.
- [19] J. K. Hale and S. M. Verduyn Lunel, Averaging in infinite dimensions. *J. Integral Equations Appl.* **2** (1990), 463–494.
- [20] A. A. Ilyin, Averaging of dissipative dynamical systems with rapidly oscillating right-hand sides. (in Russian) *Mat. Sb.* **187** (1996), 15–58; translation in *Sb. Math.* **187** (1996), 635–677.
- [21] A. A. Ilyin, Global averaging of dissipative dynamical systems. *Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5)* **22** (1998), 165–191.
- [22] M. Kamenskii, O. Mellah and P. Raynaud de Fitte, Weak averaging of semilinear stochastic differential equations with almost periodic coefficients. *J. Math. Anal. Appl.* **427** (2015), 336–364.
- [23] R. Khasminskii, On the principle of averaging the Itô’s stochastic differential equations. *Kybernetika (Prague)* **4** (1968), 260–279. (in Russian)
- [24] Y. Kifer, Some recent advances in averaging. *Modern dynamical systems and applications*, 385–403, Cambridge Univ. Press, Cambridge, 2004.
- [25] P. E. Kloeden and M. Rasmussen, *Nonautonomous Dynamical Systems*. Mathematical Surveys and Monographs, 176. American Mathematical Society, Providence, RI, 2011, viii+264 pp.
- [26] N. Krylov and N. N. Bogolyubov, *Introduction to Non-Linear Mechanics*. Annals of Mathematics Studies, no. 11. Princeton University Press, Princeton, N. J., 1943, iii+105 pp.
- [27] W. Liu, Invariance of subspaces under the solution flow of SPDE. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **13** (2010), 87–98.
- [28] W. Liu, M. Röckner, X. Sun and Y. Xie, Averaging principle for slow-fast stochastic differential equations with time dependent locally Lipschitz coefficients. *J. Differential Equations* **268** (2020), 2910–2948.
- [29] X. Liu and Z. Liu, Poisson stable solutions for stochastic differential equations with Lévy noise, *Acta Math. Sin. (Engl. Ser.)* doi: 10.1007/s10114-021-0107-1.
- [30] B. Maslowski, J. Seidler and I. Vrkoč, An averaging principle for stochastic evolution equations. II. *Math. Bohem.* **116** (1991), 191–224.
- [31] C. Prévôt and M. Röckner, *A Concise Course on Stochastic Partial Differential Equations*. Lecture Notes in Mathematics, 1905. Springer, Berlin, 2007. vi+144 pp.
- [32] M. Röckner and L. Xie, Averaging principle and normal deviations for multiscale stochastic systems. *Comm. Math. Phys.* **383** (2021), 1889–1937.
- [33] G. R. Sell, *Topological Dynamics and Ordinary Differential Equations*. Van Nostrand-Reinhold, 1971.

- [34] B. A. Shcherbakov, A certain class of Poisson stable solutions of differential equations, *Differentsial'nye Uravneniya* **4** (1968), 238–243. (in Russian)
- [35] B. A. Shcherbakov, *Topologic Dynamics and Poisson Stability of Solutions of Differential Equations*. Știința, Chișinău, 1972, 231 pp. (in Russian)
- [36] B. A. Shcherbakov, The comparability of the motions of dynamical systems with regard to the nature of their recurrence. *Differentsial'nye Uravneniya* **11** (1975), 1246–1255. (in Russian) [English translation: *Differential Equations* **11** (1975), 937–943]
- [37] B. A. Shcherbakov, *Poisson Stability of Motions of Dynamical Systems and Solutions of Differential Equations*. Știința, Chișinău, 1985, 147 pp. (in Russian)
- [38] K. S. Sibirsky, *Introduction to Topological Dynamics*. Kishinev, RIA AN MSSR, 1970, 144 pp. (in Russian) [English translation: *Introduction to Topological Dynamics*. Noordhoff, Leyden, 1975]
- [39] A. V. Skorokhod, *Asymptotic Methods in the Theory of Stochastic Differential Equations*. Translated from the Russian by H. H. McFaden. Translations of Mathematical Monographs, 78. American Mathematical Society, Providence, RI, 1989, xvi+339 pp.
- [40] X. Sun, L. Xie, Y. Xie, Averaging principle for slow-fast stochastic partial differential equations with Hölder continuous coefficients. *J. Differential Equations* **270** (2021), 476–504.
- [41] A. Yu. Veretennikov, On an averaging principle for systems of stochastic differential equations. *Mat. Sb.* **181** (1990), 256–268. (in Russian) [English translation: *Math. USSR-Sb.* **69** (1991), 271–284]
- [42] I. Vrkoč, Weak averaging of stochastic evolution equations. *Math. Bohem.* **120** (1995), 91–111.
- [43] W. Wang and A. J. Roberts, Average and deviation for slow-fast stochastic partial differential equations. *J. Differential Equations* **253** (2012), 1265–1286.
- [44] S. Zelik, Global averaging and parametric resonances in damped semilinear wave equations. *Proc. Roy. Soc. Edinburgh Sect. A* **136** (2006), 1053–1097.
- [45] X. Zhang, On stochastic evolution equations with non-Lipschitz coefficients. *Stoch. Dyn.* **9** (2009), 549–595.

M. CHENG: SCHOOL OF MATHEMATICAL SCIENCES, DALIAN UNIVERSITY OF TECHNOLOGY, DALIAN 116024, P. R. CHINA

Email address: mengyucheng@mail.dlut.edu.cn; mengyu.cheng@hotmail.com

Z. LIU (CORRESPONDING AUTHOR): SCHOOL OF MATHEMATICAL SCIENCES, DALIAN UNIVERSITY OF TECHNOLOGY, DALIAN 116024, P. R. CHINA

Email address: zxliu@dlut.edu.cn