

Physics of \mathcal{PT} -Symmetric Quantum Systems at Finite Temperature

Qian Du,^{1,*} Kui Cao,^{1,*} and Su-Peng Kou^{1,†}

¹*Center for Advanced Quantum Studies, Department of Physics,
Beijing Normal University, Beijing 100875, China*

Parity-time (\mathcal{PT}) non-Hermitian (NH) system has significant effects on observable in a great variety of physical phenomena in NH physics. However, the \mathcal{PT} -symmetric NH quantum system at finite temperature (the so-called thermal \mathcal{PT} system) has never been addressed. In this letter, based on a controlled open quantum system coupling two separated environments, we proposed a design to realize a thermal \mathcal{PT} system. After solving quantum master equation in the Gorini-Kossakowski-Sudarshan-Lindblad form, the unexpected, abnormal, universal properties of NH thermal states (the unique final states under time evolution) are explored, for example, the non-Boltzmann/Gibbs distribution, high-temperature non-thermalization effect, etc. To understand the anomalous behaviours in thermal \mathcal{PT} system, we developed the quantum Liouvillian statistical theory – the generalization of usual quantum statistical theory to finite-temperature NH systems. With its help, we derived the analytical results of thermodynamic properties. In addition, we found that at EP a continuous thermodynamic phase transition occurs, of which there exists zero temperature anomaly. This discovery will open a door to novel physics for NH systems at finite temperature.

Recently, the non-Hermitian (NH) problem in controlled open quantum systems has become the frontier of physics and attracted the attentions of many physicists. For a class of NH quantum systems under specific conditions, their Hamiltonians may have entirely real eigenvalues like Hermitian quantum systems. As an example, parity-time (\mathcal{PT})-symmetric NH quantum models were put forward by Bender and Boettcher[1–3] in the 1990s. \mathcal{PT} -symmetry spontaneous breaking (\mathcal{PT} -SSB) occurs accompanied by a transition from real spectra to complex. At the critical point (the so called “exceptional point”), the energy levels coalesce and eigenvectors merge with each other. It had attracted a lot of researches in different fields in recent years[4–8] and many platforms were proposed to realize \mathcal{PT} -symmetric NH models[9–38].

Now, the theory about \mathcal{PT} -symmetric NH quantum system and its \mathcal{PT} -SSB becomes mature. However, people are in the dark about the properties of a \mathcal{PT} -symmetric NH quantum system at finite temperature (the so-called *thermal \mathcal{PT} systems*) and know nothing of the thermodynamic behaviour of a thermal \mathcal{PT} system with Hamiltonian at EPs. In this letter, to explore this new research field, we are going to face a lot of difficult problems: *When a NH system coupling thermal bath, do there exist thermal states? For these possible thermal states, does the Boltzmann/Gibbs distribution still work? Does there exist thermodynamic phase transition for a thermal \mathcal{PT} system with Hamiltonian at EPs?...* In this letter, firstly, by taking a designed thermal \mathcal{PT} system (an open system coupling two separated environments) as an example, we solved the quantum master equation to answered above questions and found possible new physical phenomena. Then, we tried to develop

a systematic theory for them (we call it quantum Liouvillian statistical theory). Finally, we applied the theory to study physics of \mathcal{PT} -symmetric quantum systems at finite temperature.

A controlled open quantum system coupling two separated environments – physical model for thermal \mathcal{PT} system. At the beginning, to design a thermal \mathcal{PT} system, we introduce a controlled open quantum system S coupling two separated environments B, and E, respectively. The two-level system S is a tight-binding model of two lattice sites with single fermionic particle. B denotes a thermal bath with finite- $T = \frac{1}{k_B \beta_T}$ that will thermalize S to its temperature (we call it *thermal environment*), and E denotes the environment that leads to a relevant NH term for S under post-selection (we call it *NH environment*).

The Hamiltonian of the total system is given by

$$\hat{H} = \hat{H}_S \otimes \hat{I}_E \otimes \hat{I}_B + \hat{I}_S \otimes \hat{H}_E \otimes \hat{I}_B + \hat{I}_S \otimes \hat{I}_E \otimes \hat{H}_B + \hat{H}_{ES} \otimes \hat{I}_B + \hat{I}_E \otimes \hat{H}_{BS}, \quad (1)$$

where $\hat{H}_S = h(c_1^\dagger c_2 + c_2^\dagger c_1)$ is the Hamiltonian of S, \hat{H}_E is the Hamiltonian of E, and \hat{H}_B is the Hamiltonian of B. $\hat{H}_{ES} = \lambda[a^\dagger(c_1 - ic_2) + h.c.]$ represents the coupling between S and E that leads to a Lindblad operator $\hat{L}_{ES} = \sqrt{2\gamma}(c_1 + ic_2) \otimes \hat{I}_B$. $\hat{H}_{BS} = \gamma_1 c_1^\dagger \otimes \hat{B}_1 + \gamma_2 c_2^\dagger \otimes \hat{B}_2$ represents the coupling between S and B, where \hat{B}_1, \hat{B}_2 are the operators of system B. Here, we assume that $\gamma_1, \gamma_2 \ll \gamma, h$. In particular, there is *no directly coupling* between the two environments B, and E.

Steady states for the thermal \mathcal{PT} system. In general, under the Markovian approximation, the non-unitary dynamics of an open quantum system S is described by quantum master equation in the Gorini-Kossakowski-Sudarshan-Lindblad (GKSL) form[39–41], $\frac{d\rho^S(t)}{dt} \equiv \mathcal{L}\rho^S(t)$ where \mathcal{L} is a Liouville super-operator acting on the (reduced) density matrix ρ^S of the subsystem S. Then, to learn the properties of the thermal

*Who has same contribution to this work

†Corresponding author; Electronic address: spkou@bnu.edu.cn

\mathcal{PT} system, the key point is to find the possible thermal steady states by solving the GKSL equation.

First, we trace out E from the total system and consider its effect on the combined sub-system S+B. If we do post-selection by continuous particle number measurement on the sub-systems S, the continuous particle number measurement guarantees the conservation of particle number in S[42, 43]. When such post-selection process is considered, there appears effective NH Hamiltonian.

The GKSL equation of the reduced density matrix ρ^{S+B} for the combined sub-system S+B becomes

$$\frac{d\rho^{S+B}}{dt} = -i[\hat{H}_{S+B}, \rho^{S+B}] - \frac{1}{2}\{\hat{L}_{ES}^\dagger \hat{L}_{ES}, \rho^{S+B}\} + \hat{L}_{ES} \rho^{S+B} \hat{L}_{ES}^\dagger, \quad (2)$$

where the Lindblad operator $\hat{L}_{ES} = \sqrt{2\gamma}(c_1 + ic_2) \otimes \hat{I}_B$ describes loss from the coupling between E and S. After post-selection from particle-number measurement $(c_1^\dagger c_1 + c_2^\dagger c_2) \otimes \hat{I}_B$, the quantum jumping term $\hat{L}_{ES} \rho^{S+B} \hat{L}_{ES}^\dagger$ will be projected out[44, 45]. Then, the reduced density matrix of S+B satisfies $\frac{d\rho^{S+B}}{dt} = -i(\hat{H}_{S+B, \text{eff}} \rho^{S+B} - \rho^{S+B} \hat{H}_{S+B, \text{eff}}^\dagger)$ where $\hat{H}_{S+B, \text{eff}} = \hat{H}_{NH} \otimes \hat{I}_B + \hat{I}_S \otimes \hat{H}_B + \hat{H}_{BS}$ and $\hat{H}_{NH} = (h + \gamma)c_1^\dagger c_2 + (h - \gamma)c_2^\dagger c_1 - i\gamma$. Due to the post-selections, the density matrix at each moment is always normalized. Then, we can ignore the term $-i\gamma$ after normalization. During the post-selections of the system that is performed in a way so that the particle number of the system is fixed, the density matrix at each moment needs to be normalized. Then, owing to normalization of ρ_i^S , we can obtain a "usual" steady state with zero eigenvalue of \mathcal{L} by globally shifting certain imaginary part of the original NH Hamiltonian.

Therefore, we can effectively describe the sub-system S by a NH Hamiltonian \hat{H}_{NH} . An eigenstate of the NH Hamiltonian \hat{H}_{NH} can be labeled by the number of particles. Let $|N, i\rangle$ be a right eigenstate of \hat{H}_{NH} with $N = 1$ particle: $\hat{H}_{NH}|1, i\rangle = E_N|1, i\rangle$, where $i = 1$ or 2 distinguishes the eigenstates with $N = 1$. \hat{H}_{NH} can be regarded as a \mathcal{PT} -symmetric two-level NH model on the Hilbert space $\begin{pmatrix} |\uparrow\rangle \\ |\downarrow\rangle \end{pmatrix}$, of which the Hamiltonian \hat{H}_{NH} is transformed into $\hat{H}_{NH} = h\sigma_x + i\gamma\sigma_y$. Here, $|\uparrow\rangle$ and $|\downarrow\rangle$ denote the quantum states of the single fermion on site 1 and site 2, respectively. In this paper, h is set be unit. For \hat{H}_{NH} , at $h = \gamma$ a usual \mathcal{PT} -SSB occurs that is really transition from real spectra to complex: for the case $h > \gamma$, the energy levels are $E_\pm = \pm\sqrt{h^2 - \gamma^2}$; for the case $h < \gamma$, the two energy levels are $E_\pm = \pm i\sqrt{\gamma^2 - h^2}$; for the case of $h = \gamma$, the system is at exceptional point (EP) with state coalescing and energy degeneracy.

In Hermitian system at finite temperature, the theory is based on thermodynamic properties of thermal state. To reach it, one can prepare a unique final state

under time-evolution (from arbitrary initial states). For \hat{H}_{NH} at finite temperature, an important concept is "non-Hermitian thermal state" (NHTS) that is the generalization of equilibrium thermal state in Hermitian system. Using similar logic, a NHTS is the unique final state under time-evolution in a NH system at finite temperature. Let us show the rigorous mathematical definition:

Definition – Non-Hermitian thermal state (NHTS) in thermal \mathcal{PT} systems: In thermal \mathcal{PT} systems, if the spectrum of Liouville super-operator \mathcal{L} is λ_i , i.e., $\mathcal{L}\rho_i^S = \lambda_i\rho_i^S$, the NHTS ρ_{TSS}^S is the eigenstate of \mathcal{L} corresponding to the eigenvalue with maximum real part $\lambda_{TSS} = \max(\lambda_i)$. We need to point out that the NHTS is different from the usual nonequilibrium steady state, which is characterized by the eigenstates with zero eigenvalue of \mathcal{L} . In experiments, according to the uniqueness, one can firstly prepare an arbitrary initial state $\rho^S(t=0)$. Then, let the system naturally evolve. After a long enough time evolution, the initial state $\rho^S(t=0)$ will eventually turn into the only NHTS $\rho^S(t \rightarrow \infty) = \rho_{NHTS}^S$.

We first study the possible NHTSs ρ_{NHTS}^S in the phase with \mathcal{PT} -symmetry ($h > \gamma$). We trace out B and consider the effect of thermal environment B on the NH sub-system S. This is an issue that had never been studied before. We assume that the NH sub-system S has little influence on the environment B and the relaxation time of the environment E is much smaller than S, and then average out the high frequency part of quantum transition processes. Under these conditions, the temperature of a possible NHTS of the NH system is assumed to be same to that of the thermal environment B.

To illustrate the thermalization effect from coupling the environment B, we re-write the reduced density matrix of B+S in the interaction picture, i.e., $\rho_I^{S+B} = e^{iH_{\text{eff},0}t} \rho^{S+B} e^{-iH_{\text{eff},0}^\dagger t}$, where $H_{\text{eff},0}$ is the first two terms of $\hat{H}_{S+B, \text{eff}}$. The GKSL equation for reduced density matrix of S in the interaction picture is then obtained as

$$\frac{d}{dt}\rho_I^S(t) = \sum_{a,b} \sum_{\omega} \{\Gamma_{ab}(\omega) [A_b(\omega)\rho_I^S(t)A_a^\dagger(\omega) - A_a(-\omega)A_b(\omega)\rho_I^S(t)] + h.c.\}, \quad (3)$$

where $A_{a,b}(\omega)$ is a NH operator with $a, b = 1$ or 2, and its forms is $A_{a,b}(\omega) = \sum_m |m\rangle_R \langle m|_L \gamma_{a,b} c_{a,b}^\dagger c_{a,b} |m+\omega\rangle_R \langle m+\omega|_L$. Here, $|m\rangle_{R,L}$ are the right/left eigenstates of \hat{H}_{NH} with eigenvalues E_m , and $|m+\omega\rangle_{R,L}$ are the eigenvalues \hat{H}_{NH} with eigenvalues $E_m + \omega$ and they satisfy ${}_L \langle m|m\rangle_R = 1$. The reservoir correlation functions $\Gamma_{ab}(\omega)$ are defined as $\Gamma_{ab}(\omega) = \int_0^\infty dt e^{i\omega t} \langle B_a^\dagger(t)B_b(0) \rangle$, where $\langle B_a^\dagger(t)B_b(0) \rangle = \text{Tr}_B[B_a^\dagger(t)B_b(0)\rho^B]$. Because the imaginary part of Γ_{ab} only provides a shift of energy level and does not affect the steady state that we will focus on, we ignore them in the calculation below and

get $\Gamma_{ab} = \frac{1}{2}\gamma_{ab}$, where $\gamma_{ab} = \int_{-\infty}^{\infty} dt e^{i\omega t} \langle B_a^\dagger(t) B_b(0) \rangle$ is the real part of Γ_{ab} . Using the Kubo-Martin-Schwinger (KMS) condition $\langle B_a^\dagger(t) B_b(0) \rangle = \langle B_b(0) B_a^\dagger(t + i\frac{1}{T}) \rangle$, we derive temperature dependent behavior of γ_{ab} , i.e., $\gamma_{ab}(-\omega) = e^{-\omega/k_B T} \gamma_{ba}(\omega)$ [41, 46]. For simplicity, we ignore the field correlation between two lattice sites a

and b , i.e. $\gamma_{ab} = \gamma_{aa}\delta_{ab}$.

Calculating the expression of the operator $A_{a,b}(\omega)$ under base $\{|\uparrow\rangle, |\downarrow\rangle\}$ and substituting it into the GKSL master equation, we get $\frac{d}{dt}\rho_I^S = \mathcal{L}\rho_I^S$ where \mathcal{L} in the interaction picture is equal to

$$\begin{pmatrix} -\gamma_0(1 + e^{\omega_0/k_B T}) & 0 & 0 & \gamma_0(1 + e^{\omega_0/k_B T})\frac{h+\gamma}{h-\gamma} \\ -2\gamma_0(e^{\omega_0/k_B T} - 1)\sqrt{\frac{h-\gamma}{h+\gamma}} & -3\gamma_0(1 + e^{\omega_0/k_B T}) & -\gamma_0(1 + e^{\omega_0/k_B T}) & -2\gamma_0(e^{\omega_0/k_B T} - 1)\sqrt{\frac{h+\gamma}{h-\gamma}} \\ -2\gamma_0(e^{\omega_0/k_B T} - 1)\sqrt{\frac{h-\gamma}{h+\gamma}} & -\gamma_0(1 + e^{\omega_0/k_B T}) & -3\gamma_0(1 + e^{\omega_0/k_B T}) & -2\gamma_0(e^{\omega_0/k_B T} - 1)\sqrt{\frac{h+\gamma}{h-\gamma}} \\ \gamma_0(1 + e^{\omega_0/k_B T})\frac{h-\gamma}{h+\gamma} & 0 & 0 & -\gamma_0(1 + e^{\omega_0/k_B T}) \end{pmatrix} \quad (4)$$

where $\gamma_0 = \frac{1}{16}[\gamma_a\gamma_{aa}(-\omega_0) + \gamma_b\gamma_{bb}(-\omega_0)]$, $\omega_0 = 2\sqrt{h^2 - \gamma^2}$. The eigenvalues of super-operator \mathcal{L} are $-4\gamma_0(1 + e^{\omega_0\beta T})$, $-2\gamma_0(1 + e^{\omega_0\beta T})$, $-2\gamma_0(1 + e^{\omega_0\beta T})$, and 0, respectively. The first three will dissipate over time—the reciprocal of the first eigenvalue is characteristic time to reach thermal equilibrium and the second, third eigenvalue is related to the characteristic time of decoherence. The last eigenvalue 0 with maximum real part indicates a steady state in the interaction picture and it is turn out that this state is also a steady state in the Schrodinger picture. As a result, for the thermal \mathcal{PT} system, the NHTS ρ_{NHTS}^S is also a usual nonequilibrium steady state. More detailed discussion is provided in supplementary materials.

On the other hand, we study the possible NHTSs ρ_{NHTS}^S in the phase with \mathcal{PT} -symmetry breaking ($h < \gamma$). Now, the two energy levels of NH Hamiltonian are pure imaginary, $\pm i\sqrt{\gamma^2 - h^2}$. Under the conditions $\gamma_1, \gamma_2 \ll \gamma, h$, the dissipation term in Liouville super operator only has little effect, i.e. $\mathcal{L}\rho^S \approx -i(\hat{H}_{NH}\rho^S - \rho^S\hat{H}_{NH}^\dagger)$. The eigenvalues of super-operator \mathcal{L} are $-2\sqrt{\gamma^2 - h^2}$, 0, 0, $2\sqrt{\gamma^2 - h^2}$, respectively. The eigenstate with maximum real part $2\sqrt{\gamma^2 - h^2}$ corresponds to a NHTS $\rho_{NHTS}^S = |\Psi_+^R\rangle\langle\Psi_+^R|$, i.e., the system reaches a pure state $|\Psi_+^R\rangle$ which corresponding to eigenvalue with largest imaginary part $+i\sqrt{\gamma^2 - h^2}$ under a long time evolution $t \rightarrow \infty$. This can be understand as the fact that the other ($|\Psi_-^R\rangle$) will decay much more faster than these states.

To directly and strictly verify the existence of NHTSs ρ_{NHTS}^S in the thermal \mathcal{PT} system, we numerically solved the GKSL equation. The temperature T of thermal environment B is set to be $10h$. The numerical results are shown in Fig.1. The results of the expectative values of σ_i ($i = x, y, z$), i.e., $n_i = \langle\sigma_i\rangle$ for the controlled open system coupling two separated environments are represented by a point on Bloch sphere in Fig.1. Fig.1(a) indicates the existence of a *unique* steady state ρ_{NHTS}^S -

for \hat{H}_{NH} with fixed γ ($\gamma = 0.2h$), in the limit of $t \rightarrow \infty$, different initial states will eventually evolve into the same NHTS. That means under time evolution the final state $\rho^S(t \rightarrow \infty)$ is NHTS ρ_{NHTS}^S ; Fig.1(b) indicates the *abnormality* of the NHTS ρ_{NHTS}^S - for \hat{H}_{NH} with different γ ($\gamma = 0.3, \gamma = 0.9, \gamma = 1.5$), from a same initial state, for example, a fully thermalized state described by $\rho^S(t = 0) = \frac{1}{2}I_{2 \times 2}$, the system will eventually evolve into the different NHTSs $\rho_{NHTS}^S(\gamma)$. This result indicates that the NH system is no more thermalized at high temperature. Consequently, the usual Boltzmann/Gibbs distribution of equilibrium states in Hermitian systems doesn't work! What's the underlying theory to explain these unexpected results? *Is there an analytical formula to unify these calculation results?*

Quantum Liouvillian statistics theory for thermal \mathcal{PT} systems. In this section, to answer above questions, we develop a quantum Liouvillian statistics theory to systematically, and analytically characterize the thermal \mathcal{PT} systems.

We focus on the case with real energy levels ($h > \gamma$). By doing similarity transformation (ST) $\hat{S} = e^{\beta_{NH}\hat{H}'}$, \hat{H}_{NH} can be transformed into a Hermitian Hamiltonian $\hat{H}_0 = \sqrt{h^2 - \gamma^2}\sigma_x$, i.e., $\hat{H}_{NH} = \hat{S}\hat{H}_0\hat{S}^{-1}$. Here, the real number $\beta_{NH} = \frac{1}{2} \ln \frac{h+\gamma}{h-\gamma}$ characterizes the strength of NH terms and the Hermitian operator $\hat{H}' = \frac{1}{2}\sigma_z$ determines the form of the NH terms. Under NH ST \hat{S} , the energy levels E_\pm of \hat{H}_{NH} are same to those of the Hermitian model \hat{H}_0 . The corresponding eigenstates of \hat{H}_{NH} become $|\Psi_n^R\rangle = \hat{S}|\Psi_{0,n}\rangle$, $n = \pm$. Here $|\Psi_{0,n}\rangle$ denotes eigenstates of \hat{H}_0 .

The key point is to obtain the *analytical* and *correct* formula of the density matrix ρ_{NHTS}^S . Here, the word "correct" means the equivalence to the results from numerical calculations on GKSL equation. Let's show the analytical and correct formula:

The density matrix for the thermal state of a NH system at temperature $T = \frac{1}{k_B\beta T}$ described by $\hat{H}_{NH} =$




FIG. 1: (a) Time-evolution of NH systems for fixed γ ($\gamma = 0.2h$) from the different initial states: the red line corresponding to $\rho^S(t=0) = (1, 0; 0, 0)$, the blue line corresponding to $\rho^S(t=0) = (0, 0; 0, 1)$, the black line corresponding to $\rho^S(t=0) = (\frac{1}{2}, \frac{1}{2}; \frac{1}{2}, \frac{1}{2})$. The single final state indicates the uniqueness of the non-Hermitian thermal states, i.e., $\rho^S(t \rightarrow \infty) = \rho_{TSS}^S$, where $\gamma_0 = 0.05h$; (b) Time-evolution of NH systems for different γ ($\gamma = 0.3, \gamma = 0.9, \gamma = 1.5$) from the same fully thermalized initial state, where $\gamma_0 = 0.05h$. Each point on Bloch sphere denotes the expectative values of σ_i ($i = x, y, z$), i.e., $n_i = \langle \sigma_i \rangle$ for a given thermal steady state (or the final state under time-evolution) of the two-level NH system at temperature $T = 10h$. The arrows denote the direction of time-evolution.

$\hat{S}\hat{H}_0\hat{S}^{-1}$ ($\hat{S} = e^{\beta_{NH}\hat{H}'}$) is given as

$$\rho_{NHTS}^S = \sum_{n=\pm} |\Psi_n^R\rangle e^{-\beta_T E_n} \langle \Psi_n^R| = \hat{S}\rho_0\hat{S}^\dagger \quad (5)$$

where $|\Psi_n^R\rangle = \hat{S}|\Psi_{0,n}\rangle$ and $\langle \Psi_n^R| = \langle \Psi_{0,n}|\hat{S}^\dagger$. ρ_0 is the density matrix for Hermitian model \hat{H}_0 . The partition function Z_{NHTS}^S for the NHST is defined as $Z_{NHTS}^S = \text{Tr}(\rho_{NHTS}^S)$.

To simplify the Liouvillian physics for NHTSs, we introduce an effective Hamiltonian \hat{H}_L as $e^{-\beta_T \hat{H}_L} = \hat{S}(e^{-\beta_T \hat{H}_0})\hat{S}^\dagger$ or $\hat{H}_L = -\frac{1}{\beta_T} \ln[\hat{S}(e^{-\beta_T \hat{H}_0})\hat{S}^\dagger]$. In this paper, we call the temperature-dependent Hamiltonian \hat{H}_L to be *Liouvillian Hamiltonian* (LH) and the corresponding eigenvalues E_n^L to be *Liouvillian energy level*. According to $\hat{H}_L = \hat{H}_L^\dagger$, E_n^L must be real.

It was known that for Hermitian systems, the weights of different equilibrium states obey Boltzmann/Gibbs distribution, i.e., $P_n^T = \frac{1}{Z_0} e^{-\beta_T E_n}$ where E_n are energy levels. For NHTSs in this NH system, although the energy levels don't change under STs, the weights of NHTSs are deformed. As a result, usual Boltzmann/Gibbs distribution is deformed into non-Boltzmann/Gibbs distribution[47, 48], i.e., $P_n^T = \frac{1}{Z_{NHTS}^S} e^{-\beta_T E_n^L}$ which represents the weights of the different eigenstates of the Liouvillian Hamiltonian and E_n^L are Liouvillian energy levels. To distinguish the phenomenon of weights in the thermal \mathcal{PT} system to that in Hermitian thermal systems, we call the corresponding weight P_n^T the *Liouvillian weight*.

In particular, when the temperature is high, $\beta_T \rightarrow 0$, the density matrix $\rho_{NHTS}^S = e^{-\beta_T \hat{H}_L}$ for a NH sys-




FIG. 2: (a) The average spin operator from $n_i = \langle \sigma_i \rangle$ for thermal state ρ_{TSS}^S of the thermal \mathcal{PT} system. (b), (c) and (d) are n_x, n_y, n_z at temperature $T = 10h$, respectively. Red triangle symbols denote the two-level NH model with different γ ($\gamma = 0.3, \gamma = 0.9, \gamma = 1.5$) that correspond to the three different NH steady states in Fig.1(b).

tem with real spectra is reduced into $\rho_{NHTS}^S \sim \hat{S}\hat{S}^\dagger = e^{2\beta_{NH}\hat{H}'}$ and the Liouvillian weight turns into $P_n^T = \frac{1}{Z_{NHTS}^S} e^{-\beta_{NH}(-2E_n')}$ where E_n' is the “energy levels” of \hat{H}' . Such non-thermalization effect in high temperature limit ($T \rightarrow \infty$) is quite different from that for usual Hermitian model. In particular, the theoretical prediction from $\rho_{NHTS}^S \sim e^{2\beta_{NH}\hat{H}'}$ is exactly consistent to the numerical results in Fig.1(b).

Thermodynamic properties for the thermal \mathcal{PT} system: With the help of the quantum Liouvillian statistical theory, we study thermodynamic properties for the NHTSs ρ_{NHTS}^S of the thermal \mathcal{PT} system.

Firstly, we consider the case in the region of $h \geq \gamma$. The physical properties of the original NH system \hat{H}_{NH} at finite- T corresponds to that for a Hermitian system \hat{H}_L . According to $e^{-\beta_T \hat{H}_L} = \hat{S}(e^{-\beta_T \hat{H}_0})\hat{S}^\dagger$, we obtain the analytical result as $e^{-\beta_T \hat{H}_L} = \sigma_x A + \sigma_z B + IC$ where $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, $A = -\sinh(\beta_T \sqrt{h^2 - \gamma^2})$, $B = \sinh(\beta_{NH}) \cdot \cosh(\beta_T \sqrt{h^2 - \gamma^2})$ and $C = \cosh(\beta_{NH}) \cdot \cosh(\beta_T \sqrt{h^2 - \gamma^2})$. As a result, the Liouvillian Hamiltonian is derived as $\hat{H}_L = -\frac{1}{\beta_T} \frac{\cosh^{-1} C}{|\vec{r}_{\text{eff}}|} (\vec{\sigma} \cdot \vec{r}_{\text{eff}})$ where $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$, $\vec{r}_{\text{eff}} = (A, 0, B)$ and $|\vec{r}_{\text{eff}}| = \sqrt{A^2 + B^2}$.

We then calculate the expectative values of σ_i ($i = x, y, z$), i.e., $n_i = \langle \sigma_i \rangle$. The expected value of physical operator σ_i is defined as $\langle \sigma_i \rangle = \frac{1}{Z_{NHTS}^S} \text{Tr}(\sigma_i \cdot e^{-\beta_T \hat{H}_L})$. After straightforwardly calculations, we have $\vec{n} = (n_x, n_y, n_z) = (\frac{A}{C}, 0, \frac{B}{C})$. Fig.2(a) shows the results of \vec{n} . From Fig.2(a), when β_{NH} increases from zero to infinite, the spin direction changes from x -direction to z -direction together with amplitude increasing.

Approaching to EP, at finite- T , the Liouvillian Hamil-

tonian \hat{H}_L becomes an effective Hamiltonian with *infinite* external field, $\hat{H}_L \rightarrow \frac{\beta_{\text{NH}}}{\beta_T} \sigma_z$ with $\beta_{\text{NH}} \rightarrow \infty$. The expectative values of σ_i ($i = x, y, z$) is derived as $\vec{n} = (0, 0, \text{sgn}(\gamma))$. As a result, \vec{n} is fixed to be polarized along z -direction with saturated amplitude 1 (See the magenta arrows in Fig.2(a)) marked by a magenta line).

Next, we consider the case in the region of $h < \gamma$. The NHTS becomes a pure state described by $|\Psi_+^R\rangle$. We have $\vec{n} = (0, -\sqrt{1 - (\frac{h}{\gamma})^2}, \frac{h}{\gamma})$. As shown in Fig.2(a), with increasing γ , the red arrows (the expectative values of σ_i) swerve from z -direction to y -direction.

Fig.2(b)-(d) separately show the behavior of $n_i = \langle \sigma_i \rangle$ at a high temperature $T = 10h$. The dotted magenta lines mark the position of EP. The non-zero results of n_i at high temperature once again implies non-Boltzmann/Gibbs distribution. In addition, we point out that the red triangle symbols come from the numerical results in Fig.1(b) by solving GKSL equation for the two-level NH model with different γ ($\gamma = 0.3, \gamma = 0.9, \gamma = 1.5$). The consistence between the results from quantum Liouvillian statistical theory and those by directly solving GKSL equation verifies the correctness of both calculations.

Thermodynamic phase transition at EP. Finally, we discuss the possible thermodynamic phase transition at EP $h = \gamma$.

From above discussion, we found that the average spin operator $\langle \sigma_i \rangle$ is always continuous crossing over EP. Furthermore, we calculate the derivatives of spin average values $\frac{\partial \vec{n}}{\partial \gamma}$. Because $h = \gamma$ is a transition from real spectra to complex, the steady states of both sides ($h > \gamma$ and $h < \gamma$) are described by different functions. After straightforward calculation, we have $\frac{\partial \vec{n}}{\partial \gamma} = \frac{1}{h} (\{ \tanh(\beta_T \sqrt{h^2 - \gamma^2}) + \sqrt{h^2 - \gamma^2} [1 - [\tanh(\beta_T \sqrt{h^2 - \gamma^2})]^2] \beta_T \} \frac{\gamma}{\sqrt{h^2 - \gamma^2}}, 0, 1)$ for

$$h > \gamma \text{ and } \frac{\partial \vec{n}}{\partial \gamma} = (0, -\frac{1}{\sqrt{1 - (\frac{h}{\gamma})^2}} \frac{h^2}{\gamma^3}, -\frac{h}{\gamma^2}) \text{ for } h < \gamma.$$

The discontinuity of $\frac{\partial \vec{n}}{\partial \gamma}$ indicates a "second-order" phase transition: At finite temperature $T \neq 0$, there exists a $\nu = \frac{1}{2}$ critical rule for $\frac{\partial \vec{n}}{\partial \gamma}$: $\frac{\partial \vec{n}}{\partial \gamma} = (2\beta_T, 0, h^{-1})$ for $h \rightarrow \gamma + 0^+$ and $\frac{\partial \vec{n}}{\partial \gamma} = (0, -(2h)^{-\frac{1}{2}} (\gamma - h)^{-\nu}, -\gamma^{-1})$ for $\gamma \rightarrow h + 0^+$; At zero temperature $T = 0$, $\frac{\partial \vec{n}}{\partial \gamma} = ((2h)^{-\frac{1}{2}} (h - \gamma)^{-\nu}, 0, h^{-1})$ for $h \rightarrow \gamma + 0^+$ and $\frac{\partial \vec{n}}{\partial \gamma} = (0, -(2h)^{-\frac{1}{2}} (\gamma - h)^{-\nu}, -\gamma^{-1})$ for $\gamma \rightarrow h + 0^+$. The results obviously illustrate a *zero temperature anomaly* for the thermodynamic phase transition at EP! See the detailed discussion in supplementary materials.

Conclusion and summary. In this paper, we developed a quantum Liouvillian statistics theory to characterize thermodynamic properties for a typical thermal \mathcal{PT} system. With the help of the theory, we derived the analytical results of non-Hermitian thermal state ρ_{NHTS}^S that accurately describe their unexpected, abnormal, universal properties, for example, the non-Boltzmann/Gibbs distribution, high-temperature non-thermalization effect, continuous thermodynamic phase transition with zero temperature anomaly occurs at EP, etc. In the future, we will study the thermal state of more complex NH models and explore the possible exotic physics for NH systems at finite-T.

Acknowledgments

This work is supported by NSFC Grant No. 11974053, 1217040237. We thank the helpful discussion with Wei Yi, Shu Chen, Gao-Yong Sun.

-
- [1] C. M. Bender, and S. Boettcher, Phys. Rev. Lett. **80**, 5243 (1998).
 [2] C. M. Bender, D. C. Brody, and H. F. Jones, Phys. Rev. Lett. **89**, 270401 (2002).
 [3] C. M. Bender, Rep. Prog. Phys. **70**, 947 (2007).
 [4] G. L. Giorgi, Phys. Rev. B **82**, 052404 (2010).
 [5] L. Ge and H. E. Türeci, Phys. Rev. A **88**, 053810 (2013).
 [6] C. Hang, G. Huang, and V. V. Konotop, Phys. Rev. Lett. **110**, 083604 (2013).
 [7] C. Korff and R. Weston, J. Phys. A **40**, 8845 (2007).
 [8] C. Korff and R. Weston, J. Phys. A **41**, 295206 (2008).
 [9] C. E. Rüter, K. G. Makris, R. El-Ganainy, D. N. Christodoulides, M. Segev, and D. Kip, Nat. Phys. **6**, 192 (2010).
 [10] C. Hang, G. Huang, and V. V. Konotop, Phys. Rev. Lett. **110**, 083604 (2013).
 [11] A. Guo, G. J. Salamo, D. Duchesne and R. Morandotti, M. Volatier-Ravat and V. Aimez, G. A. Siviloglou and D. N. Christodoulides, Phys. Rev. Lett. **103**, 093902 (2009).
 [12] P. Peng W. Cao, C. Shen, W. Qu, J. Wen, L. Jiang and Y. Xiao, Nat. Phys. **12**, 1139 (2016).
 [13] Z. Zhang, Y. Zhang, J. Sheng, L. Yang, M.-A. Miri, D. N. Christodoulides, B. He, Y. Zhang, and M. Xiao, Phys. Rev. Lett. **117**, 123601 (2016).
 [14] Y. D. Chong, Li Ge, and A. Douglas Stone, Phys. Rev. Lett. **106**, 093902 (2011).
 [15] A. Regensburger, C. Bersch, M.-A. Miri, G. Onishchukov, D. N. Christodoulides, and U. Peschel, Nature (London) **488**, 167 (2012).
 [16] L. Feng, Y.-L. Xu, W. S. Fegadolli, M.-H. Lu, J. E. B. Oliveira, V. R. Almeida, Y.-F. Chen, and A. Scherer, Nat. Mater. **12**, 108 (2013).
 [17] M. Wimmer, M.-A. Miri, D. Christodoulides, and U. Peschel, Sci. Rep. **5**, 17760 (2015).
 [18] S. Weimann, M. Kremer, Y. Lumer, S. Nolte, K. G. Makris, M. Segev, M. C. Rechtsman and A. Szameit, Nat. Mater. **16**, 433 (2017).
 [19] H. Hodaei, A. U. Hassan, S. Wittek, H. Garcia-Gracia, R. El-Ganainy, D. N. Christodoulides, and M. Khajavikhan, Nature **548**, 187 (2017).

- [20] S. K. Özdemir, S. Rotter, F. Nori, and L. Yang, *Nat. Mater.* **18**, 783 (2019).
- [21] L. Feng, Y. L. Xu, W. S. Fegadolli, M. H. Lu, J. E. B. Oliveira, V. R. Almeida, Y. F. Chen and A. Scherer, *Nat. Mater.* **12**, 108 (2013).
- [22] J. Li, A. K. Harter, J. Liu, L. de Melo, Y. N. Joglekar and L. Luo, *Nat. Commun.* **10**, 855 (2019).
- [23] N. Bender, S. Factor, J. D. Bodyfelt, H. Ramezani, D. N. Christodoulides, F. M. Ellis, and T. Kottos, *Phys. Rev. Lett.* **110**, 234101 (2013).
- [24] J. Schindler, A. Li, M. C. Zheng, F. M. Ellis, and T. Kottos, *Phys. Rev. A* **84**, 040101(R) (2011).
- [25] S. Assaworarith, X. Yu, S. Fan, *Nature* **546**, 387 (2017).
- [26] Y. Choi, C. Hahn, J. W. Yoon, and S. H. Song, *Nat. Commun.* **9**, 2182 (2018).
- [27] S. Bittner, B. Dietz, U. Günther, H. L. Harney, M. Miski-Oglu, A. Richter, and F. Schäfer, *Phys. Rev. Lett.* **108**, 024101 (2012).
- [28] B. Peng, S. K. Özdemir, F. Lei, F. Monifi, M. Gianfreda, G. L. Long, S. Fan, F. Nori, C. M. Bender, and L. Yang, *Nat. Phys.* **10**, 394 (2014).
- [29] C. Poli, M. Bellec, U. Kuhl, F. Mortessagne, and H. Schomerus, *Nat. Commun.* **6**, 6710 (2015).
- [30] Z.-P. Liu, J. Zhang, S. K. Özdemir, B. Peng, H. Jing, X.-Y. Lü, C.-W. Li, L. Yang, F. Nori, and Y.-X. Liu, *Phys. Rev. Lett.* **117**, 110802 (2016).
- [31] L. Feng, Z. J. Wong, R. M. Ma, Y. Wang and X. Zhang, *Science*, **346** 972 (2014).
- [32] B. Peng, S. K. Özdemir, F. Lei, F. Monifi, M. Giandreda, G. L. Long S. Fan, F. Nori, C. M. Bender and L. Yang, *Nat. Phys.* **10**, 394 (2014).
- [33] H. Hodaei, M. A. Miri, M. Heinrich, D. N. Christodoulides, M. Khajavikhan, *Science*, **346**, 975 (2014).
- [34] X. Zhu, H. Ramezani, C. Shi, J. Zhu, and X. Zhang, *Phys. Rev. X* **4**, 031042 (2014).
- [35] B.-I. Popa, and S. A. Cummer, *Nat. Commun.* **5**, 3398 (2014).
- [36] R. Fleury, D. Sounas, and A. Alù, *Nat. Commun.* **6**, 5905 (2015).
- [37] K. Ding, G. Ma, M. Xiao, Z. Q. Zhang, and C. T. Chan, *Phys. Rev. X* **6**, 021007 (2016).
- [38] Y. Wu, W. Liu, J. Geng, X. Song, X. Ye, C. K. Duan, X. Rong, and J. Du, *Science*, **364**, 878 (2019).
- [39] G. Lindblad, *Commun. Math. Phys.*, **48**, 119 (1976).
- [40] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, *J. Math. Phys.*, **17**, 821 (1976).
- [41] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
- [42] Y. Aharonov, D. Z. Albert, and L. Vaidman, *Phys. Rev. Lett.* **60**, 1351 (1988); Y. Aharonov and L. Vaidman, *Phys. Rev. Lett.* **62**, 2327 (1989); A. J. Leggett, *Phys. Rev. Lett.* **62**, 2325 (1989); A. Peres, *Phys. Rev. Lett.* **62**, 2326 (1989); Y. Aharonov and L. Vaidman, *Phys. Rev. A* **41**, 11 (1990).
- [43] T. A. Brun, *Am. J. Phys.* **70**, 719 (2002).
- [44] N. Matsumoto, K. Kawabata, Y. Ashida, S. Furukawa, and M. Ueda, *Phys. Rev. Lett.* **125**, 260601 (2020).
- [45] Y. Ashida and M. Ueda, *Phys. Rev. Lett.* **120**, 185301 (2018).
- [46] R. Haag, N. Hugenholtz, and M. Winnink, *Commun. Math. Phys.* **5**, 215 (1967).
- [47] B. Gardas, S. Deffner and A. Saxena, *Scientific Reports* **6**, 23408 (2016).
- [48] P. N. Meisinger and M. C. Ogilvie, *Phil. Trans. R. Soc. A* **371**, 20120058 (2013).