

First indication on self-similarity of strangeness production in $Au + Au$ collisions at RHIC: Search for signature of phase transition in nuclear matter

M. V. Tokarev^{1,2*}, I. Zborovský^{3**}

1 Joint Institute for Nuclear Research, Dubna, 141980, Russia

2 Dubna State University, Dubna, 141980, Russia

3 The Czech Academy of Science, Nuclear Physics Institute,
Řež, 250 68, Czech Republic

* tokarev@jinr.ru

** zborovsky@ujf.cas.cz

November 17, 2021

*50th International Symposium on Multiparticle Dynamics
(ISMD2021)*

12-16 July 2021

doi:10.21468/SciPostPhysProc.?

Abstract

New results of analysis of K_S^0 -meson spectra measured over a wide range of energy $\sqrt{s_{NN}} = 7.7 - 200$ GeV and centrality in $Au + Au$ collisions by the STAR Collaboration at RHIC using the z -scaling approach are presented. Indication on self-similarity of fractal structure of nuclei and fragmentation processes with K_S^0 probe is demonstrated. The energy loss as a function of the collision energy, centrality and transverse momentum of the inclusive strange meson is estimated.

1 Introduction

The idea of self-similarity of hadron interactions is a fruitful concept to study collective phenomena in hadron matter. Important manifestation of such a concept is existence of scaling itself. Some of the scaling features constitute pillars of modern critical phenomena. Other category of scaling laws (self-similarity in point explosion, laminar and turbulent fluid flow, super-fluidity far from phase boundary and critical point, etc.) reflects features not related to phase transitions. The notions "scaling" and "universality" have special importance in critical phenomena. The universality hypothesis reduces the great variety of critical phenomena to a small number of equivalence classes, the so-called "universality classes", which depend only on few parameters (critical exponents). The universality has its origin in the long range character of interactions. Close to the transition point, the behavior of the cooperative phenomena becomes independent of the microscopic details of the considered system. The fundamental parameters determining the universality class are the symmetry of the order parameter and the dimensionality of space.

2 General concept of z -scaling

The concept of the z -scaling is based on the fundamental principles of self-similarity, locality, and fractality of particle production in $p+p$ and $A+A$ interactions at a constituent level [1–4]. At sufficiently high energies, the collision of hadrons or nuclei is considered as an ensemble of individual interactions of their constituents. Structures of the colliding objects are characterized by fractal dimension δ_1 and δ_2 . The interacting constituents carry the fractions x_1 and x_2 of the momenta P_1 and P_2 of the colliding hadrons (or nuclei). The fragmentation of the scattered constituent and fragmentation of the produced recoil is described by fractal dimensions ϵ_a and ϵ_b , and by the momentum fractions y_a and y_b , respectively. The gross features of the momentum distributions of produced inclusive particles can be described in terms of the kinematic characteristics of the corresponding sub-processes. The sub-process is considered as a binary collision of constituents with masses x_1M_1 and x_2M_2 , which results in the scattered and recoil object in the final state with masses m_a/y_a and $M_X = x_1M_1 + x_2M_2 + m_b/y_b$, respectively. The associate production of the particle with mass m_b ensures conservation of additive quantum numbers. The momentum conservation in the sub-process is connected with the recoil mass as follows $(x_1P_1 + x_2P_2 - p/y_a)^2 = M_X^2$. A function of the momentum fractions, $\Omega = (1 - x_1)^{\delta_1}(1 - x_2)^{\delta_2}(1 - y_a)^{\epsilon_a}(1 - y_b)^{\epsilon_b}$, is introduced, which is proportional to the number of all such constituent configurations, which contain the configuration defined by the fractions x_1 , x_2 , y_a and y_b . It plays the role of a relative volume which occupy these configurations in space of the momentum fractions. The self-similarity variable z and the scaling function $\psi(z)$ are expressed via model parameters, momentum fractions, multiplicity density, total inelastic and inclusive cross section of produced hadron, and the corresponding Jacobian J [1–4]: $z = z_0\Omega^{-1}$, $\psi(z) = \pi/[(dN/d\eta)\sigma_{inel}J] \cdot Ed^3\sigma/dp^3$.

3 Self-similarity of K_S^0 production in $Au + Au$ collisions

Figure 1(a) shows the dependence of the scaling function $\psi(z)$ on the variable z for K_S^0 mesons produced in the (0–5)% central $Au + Au$ collisions for different energies. The symbols correspond to p_T -distributions measured in the $Au+Au$ system and the solid line is the z -scaling curve for $p+p$ interactions. Reasonable coincidence of the symbols and the solid curve in the interval $z = 0.1 - 7$ was found provided specific dependencies of model parameters on energy and multiplicity. There are no irregularities in the behavior of $\psi(z)$ over a wide range of collision energy $\sqrt{s_{NN}} = 7.7 - 200$ GeV. An indication on a flattening of the scaling function at low z and a power-law at high z is clearly observed. Similar results were obtained for other collision centralities. This result is interpreted as a self-similar modification of the constituent sub-processes by the created medium.

The energy loss is characterized by the fraction y_a in the z -scaling scheme. The quantity is sensitive characteristic of the nuclear medium. Figure 1(b) shows the dependence of the momentum fraction y_a on the transverse momentum p_T of K_S^0 mesons produced in the most central (0–5)% $Au + Au$ collisions at different energies. A monotonic growth of y_a with p_T is found for all energies and all collision centralities. This means that the relative energy dissipation $\Delta E_q/E_q = (1 - y_a)$ associated with a high- p_T particle is smaller than for processes with lower transverse momenta. The energy loss becomes larger as the collision energy increases. It was found that the energy loss is greater in the most central $Au + Au$ collisions than in




Figure 1: The scaling function $\psi(z)$ (a) and momentum fraction y_a (b) for K_S^0 mesons produced in (0–5)% central $Au + Au$ collisions at $\sqrt{s_{NN}} = 7.7, 11.5, 19.6, 27, 39, 62.4, 200$ GeV in the rapidity interval $|y| < 0.5$. The symbols correspond to data obtained by the STAR Collaboration at RHIC. The solid line is a reference curve for $p + p$ interactions.

the peripheral ones. The energy losses alone, which have been estimated for the production of K_S^0 mesons in $Au + Au$ collisions over a wide range of collisions energy, centrality and p_T , show no sign of a phase transition.

4 Conclusion

We analyzed data on transverse momentum spectra of K_S^0 mesons measured in $Au + Au$ collisions at mid-rapidity by the STAR Collaboration in BES-I program at RHIC in the z -scaling approach. Self-similarity of K_S^0 -meson production in the gold-gold collisions over a wide kinematic and centrality range was found. Constituent energy loss as a function of the collision energy, centrality and transverse momentum of K_S^0 mesons was estimated. The method of analysis is extended for systematic description of $A + A$ collisions with production of identified hadrons.

Acknowledgements

This work was partially supported by Project funded by the MEYS of the Czech Republic under the contract LTT18021.

References

- [1] M. V. Tokarev and I. Zborovský, *New indication on scaling properties of strangeness production in pp collisions at RHIC*, Int. J. Mod. Phys. A **32**, 1750029 (2017), doi:10.1142/S0217751X17500294.
- [2] M. Tokarev, A. Kechechyan, and I. Zborovský, *Validation of z-scaling for negative particle*

- production in Au+Au collisions from BES-I at STAR*, Nucl. Phys. A **993**, 121646 (2020), doi:10.1016/j.nuclphysa.2019.121646.
- [3] M. V. Tokarev, I. Zborovský, A. O. Kechechyan, and T. G. Dedovich, *Verification of z -scaling in $p+p$, $\bar{p}+p$ and $Au+Au$ collisions at RHIC, Tevaron and LHC*, Phys. Part. Nucl. **51**, 141 (2020), doi:10.1134/S1063779620020045.
- [4] I. Zborovský and M. Tokarev, *Self-similarity, fractality, and entropy principle in collisions of hadrons and nuclei at RHIC, Tevaron and LHC*, Phys. Part. Nucl. Lett. **18**, 302 (2021), doi:10.1134/S1547477121030110.