

The scaling of the entropy production rate in a φ^4 model of Active Matter

Matteo Paoluzzi^{1, *}

¹*Departament de Física de la Matèria Condensada,
Universitat de Barcelona, C. Martí Franquès 1, 08028 Barcelona, Spain.*

(Dated: December 13, 2021)

Active φ^4 theories consider non-equilibrium terms that play an important role in describing active phase separation but are irrelevant, in the Renormalization Group sense, at the critical point. Their irrelevance makes the critical exponents the same of Ising universality class. Despite their irrelevance, they contribute with a non-trivial scaling of the entropy production rate at the critical point. However, active baths also constitute a continuous source of correlated noise. We consider the non-equilibrium dynamics of a non-conserved scalar field φ (Model A) driven out-of-equilibrium by a persistent noise that is correlated on a finite time scale τ . We perform the computation of the density of entropy production rate σ and we study its scaling near the critical point. We found that similar to the case of Active Model A, and although the non-linearities responsible for non-vanishing entropy production rates in the two models are quite different, the irrelevant parameter τ makes the critical dynamics irreversible.

INTRODUCTION

Self-propelled particles can undergo a second-order phase transition whose numerical estimate of the critical exponents are consistent with those of the Ising universality class [1, 2]. This happens despite the microscopic mechanism responsible for this non-equilibrium phase transition, i. e., the so-called Motility-Induced Phase Separation, does not have any equilibrium counterpart [3, 4]. Coarse-grained models of active particles suggest that non-equilibrium gradient terms have to be taken into account for describing the active system on large scale, e. g., the so-called Active Model A, Active Model B, etc... [5]. However, once we study the system around the critical point, the non-equilibrium terms of active φ^4 field theories are irrelevant in the Renormalization Group (RG) sense and thus they run to zero under RG transformations [6]. It has been shown recently that, even though irrelevant, the non-equilibrium gradient term $(\nabla\varphi)^2$ of Active Model A/B causes an anomalous scaling of the density of entropy production rate σ that signals a non-vanishing entropy production at criticality, i. e., the critical dynamics remains non-equilibrium like at the critical point [6]. This result implies that non-equilibrium irrelevant terms might make the critical dynamics irreversible. We can speculate that this fact might open the way to the definition of new non-equilibrium universality classes.

On the other hand, one can expect that the non-equilibrium terms considered in the Active Model A/B are not the only ingredients that one should take into account for building an active φ^4 field theory. And thus, a natural question is whether or not other non-equilibrium terms make the critical dynamics effectively non-equilibrium. In particular, usually one assumes that the noise acting on the coarse-grained fields has no mem-

ory, not in time, neither in space. This assumption seems reasonable because, in order to describe the system on large scales, one has to maintain only the relevant degrees of freedom, e. g., the order parameters, that are slow-varying variables. In this framework, the noise represents the effect of the fast degrees of freedom on the slow ones [7].

In particular, in the case of active systems, the noise is not necessarily delta-correlated, as proved by experiments of passive beads in active baths [8–10]. Moreover, not only the noise generated by the active bath is correlated on a finite time-scale τ , but it violates the Fluctuation-Dissipation Theorem in a peculiar way [10], i. e., although the active bath exerts time-correlated forces, linear response reveals an instantaneous friction kernel.

If we interpret the passive bead as a probe that captures the emerging properties of the active bath, it looks reasonable to assume that fluctuations are not delta-correlated but they are characterized by a finite correlation time. Moreover, recent numerical experiments suggest that even approaching the MIPS critical point, i. e., in a situation where the critical slowing down dominates over any other time scale, field theories driven by an exponentially-correlated stochastic force well capture the non-thermal *active* fluctuations [11]. It is known that, once we consider an exponentially correlated noise in a system composed of discrete degrees of freedom, a non-vanishing entropy production rate can be due only to non-linear interactions [12, 13]. A natural question is what happens in the case of a field theory.

In this work, using the standard machinery of dynamical critical phenomena and stochastic thermodynamics, we shall compute the entropy production rate in the case of a Model A driven by a persistent noise on a finite time scale τ . We focus our attention on the behavior of the system at the critical point and we consider the situation where the transition is approached from the symmetric phase, i. e., the system is not phase separated and thus there are not reasons for including other non-equilibrium

* matteopaoluzzi@ub.edu

active terms. The computation does not require any perturbation expansion, neither in the persistence time τ nor in the non-linear couplings of the field theory.

We obtain that all the non-linear terms (in principle infinite) appearing in the effective hamiltonian that describes the critical system contribute to the entropy production. Moreover, focusing our attention on a φ^4 theory, we show that, in agreement with that observed in Active Model A [6], the irrelevant parameter τ , on one hand, puts the system in the Ising universality class, on the other hand, maintains the relaxation dynamics out-of-equilibrium even at the critical point.

MODEL A DRIVEN BY A PERSISTENT NOISE

As it has been recently shown in Ref. [2] (where one can find details about the RG of Model B driven by a persistent noise), field theories with exponentially correlated noise well reproduce the critical dynamics of MIPS and are consistent with the Ising universality class. Here we perform the computation of the entropy production rate to understand the impact of the persistent noise on the critical dynamics. Following Ref. [7], we will focus our attention on the simplest case of Model A. We consider the non-equilibrium dynamics of a non-conserved scalar field $\varphi = \varphi(x, t)$ in d spatial dimensions (with $x \equiv \mathbf{x} \in \mathbb{R}^d$) that is driven out-of-equilibrium by a persistent stochastic force $\eta = \eta(x, t)$ that is correlated on a finite time scale τ , the stochastic dynamics of the two fields are

$$\dot{\varphi} = -\Gamma_0 \frac{\delta H_{LG}}{\delta \varphi} + \eta \quad (1)$$

$$\dot{\eta} = -\frac{1}{\tau} \eta + \zeta, \quad (2)$$

with the noise $\zeta = \zeta(x, t)$ satisfying

$$\langle \zeta(x, t) \rangle = 0, \quad \langle \zeta(x, t) \zeta(y, s) \rangle = \frac{2D\Gamma_0}{\tau^2} \delta(t-s) \delta^{(d)}(x-y).$$

In the present work we set $\Gamma_0 = 1$. We now consider the most general expression for the Landau-Ginzburg hamiltonian H_{LG} that is

$$H_{LG}[\varphi] = \int dx (H_0 + H_I) \quad (3)$$

$$H_0 \equiv \frac{1}{2} [\mu(\nabla\varphi)^2 + r\varphi^2]$$

$$H_I \equiv \sum_{p=2}^{\infty} \frac{u_p}{(2p)!} \varphi^{2p},$$

where we adopt the notation $dx \equiv d^d x$. In eq. (3) H_0 is the Gaussian contribute and H_I contains the non-linear interactions. The coupling constant r sets the distance from the critical point that is located at $r=0$. We work in a condition such that the critical point is always approached from above, i. e., from the disordered phase

where $r \geq 0$ and thus $\langle \varphi \rangle = 0$. We stress that, since we are interested in studying only the contribution of the persistent noise on the entropy production rate around the critical point, we neglect other non-equilibrium terms like those in the Active Model A and B [5]. These terms play surely an important role on active phase separation and thus have to be taken into account for describing the spinodal decomposed active phases [14, 15]. In order to establish the relevant parameters of the theory, we can perform the scaling analysis of eq. (1) that is obtained by considering Kadanoff transformations with scaling parameter b , i. e., we replace $x \rightarrow bx$, $t \rightarrow b^z t$, and $\varphi \rightarrow b^\chi \varphi$ (more details about the scaling analysis of the dynamical action can be found in Ref. [11]). We obtain the following scaling transformations for the parameters of the theory

$$\begin{aligned} r' &= rb^z & (4) \\ \mu' &= \mu b^{z-2} \\ u'_p &= u_p b^{2(p-1)\chi+z} \\ D' &= Db^{z-2\chi-d} \\ \tau' &= \tau b^{-z}. \end{aligned}$$

Where z is the dynamic exponent and χ the dimension of the scalar field φ . Once we impose $\mu = \mu'$, we get $z=2$, i. e., the usual z exponent of Model A [7]. Because of that, τ is an irrelevant parameter that runs to zero under RG transformations [11, 16] and thus the critical exponents are the same of Ising universality class. However, non-universal quantities as the location of the critical point might depend on τ , as shown in Ref. [17]. From the scaling of the noise strength D we obtain that the natural dimension of the field is $\chi = 1-d/2$, i. e., that is obtained by setting $D=D'$. The scaling of the coupling constants u_p sets the (upper) critical dimensions of each non-linear interaction, i. e., $d_c = 4$ for u_2 (that we will name as u), $d_c = 3$ for u_3 , etc... meaning that the φ^4 interaction can be neglected above 4 and φ^6 interactions above $d = 3$ [18]. As discussed in Ref. [6], non-equilibrium control parameters might play a role at the critical point even if they are irrelevant in the RG sense. In order to study the impact of τ on critical dynamics, we have to compute the entropy production rate of the model.

Following stochastic thermodynamics [19–21], we are going to compute the steady-state entropy production rate \mathcal{S} that is

$$\mathcal{S} = \lim_{t_F \rightarrow \infty} \frac{1}{t_F} \left\langle \log \frac{P[\varphi]}{P_R[\varphi]} \right\rangle. \quad (5)$$

with $P[\varphi]$ indicating the probability of a path $\{\varphi(x, t)\}_{t_0 \leq t \leq t_F}$, with t_0 and t_F the initial and final time, respectively. In Eq. (5), $P_R[\varphi]$ is the probability for the time-reversed trajectory that can be computed considering the transformations $t \rightarrow t_F - t$ and $\varphi \rightarrow \varphi_R(x, t) = \varphi(x, t_F - t)$. The average $\langle \cdot \rangle$ is performed with respect to the noise realizations. Under suitable ergodicity assumptions that we make in the present work,

when it is necessary, we replace averages over independent noise realizations with average over a long trajectory.

As we shall see, the probability $P[\varphi]$ can be written in terms of a dynamical action $A[\varphi]$, i. e., the Onsager–Machlup action, so that

$$P[\varphi] \propto \exp(-A[\varphi]) \quad (6)$$

$$A[\varphi] = \int dx dt \mathcal{L}[\varphi]. \quad (7)$$

Here we have written the dynamical action $A[\varphi]$ in terms of the "density" $\mathcal{L}[\varphi]$. Similarly for $P_R[\varphi]$ one has $P_R[\varphi] \propto \exp(-A_R[\varphi])$ so that

$$\mathcal{S} = \lim_{t_F \rightarrow \infty} \frac{1}{t_F} \int_{t_0}^{t_0+t_F} dt dx \{ \mathcal{L}_R[\varphi] - \mathcal{L}[\varphi] \}. \quad (8)$$

ONSAGER-MACHLUP DYNAMICAL ACTION

In order to proceed with the program illustrated above, it is convenient to write the dynamics in the Fourier space. We consider the system closed in a d -dimensional box of side L and volume $V = L^d$ where periodic boundary conditions are employed. We can thus write $\varphi(x, t)$ and $\eta(x, t)$ in terms of their space-Fourier components as follows [18, 22, 23]

$$\varphi(x, t) = \frac{1}{L^{d/2}} \sum_{|k| \in (0, \Lambda)} e^{ikx} \varphi_k(t) \quad (9)$$

$$\eta(x, t) = \frac{1}{L^{d/2}} \sum_{|k| \in (0, \Lambda)} e^{ikx} \eta_k(t)$$

where we have introduced the ultraviolet cutoff $\Lambda = \frac{2\pi}{a}$ and a the lattice spacing (or the typical particle size) used as a regulator. In the Fourier space, the hamiltonian is

$$H_{LG} = H_0 + H_I \quad (10)$$

$$H_0 = \frac{1}{2} \sum_k (\mu k^2 + r) |\varphi_k|^2$$

$$H_I = \sum_{p=2}^{\infty} \frac{u_p}{(2p)!} \frac{1}{V^{-p+1}} \sum_{k_1, \dots, k_{2p}} \varphi_{k_1} \varphi_{k_2} \dots \varphi_{k_{2p}} \delta_{k_1 + \dots + k_{2p}}$$

where H_I contains all the non-linear interactions. In this way, the dynamics is

$$\dot{\varphi}_k = -\frac{\partial H_{LG}}{\partial \varphi_k} + \eta_k \quad (11)$$

$$\dot{\eta}_k = -\frac{1}{\tau} \eta_k + \zeta_k \quad (12)$$

with

$$\langle \zeta_k(t) \rangle = 0, \quad \langle \zeta_k(t) \zeta_q(s) \rangle = \frac{2D}{\tau^2} \delta_{k, -q} \delta(t - s).$$

For making progresses, we perform the time derivative of Eq. (11) obtaining

$$\ddot{\varphi}_k + \mathbb{H}_{kq} \dot{\varphi}_q + \frac{1}{\tau} [\dot{\varphi}_k + F_k] = \zeta_k \quad (13)$$

$$F_k \equiv \frac{\partial H_{LG}}{\partial \varphi_k}$$

$$\mathbb{H}_{kq} \equiv \frac{\partial^2 H_{LG}}{\partial \varphi_k \partial \varphi_q},$$

where we have adopted the Einstein summation convention. In order to obtain the expression of the probability distribution $P[\varphi]$, we start with introducing the expectation value $\langle \mathcal{O} \rangle$ of a generic observable $\mathcal{O}[\varphi]$ that is

$$\langle \mathcal{O} \rangle = \int \mathcal{D}[\zeta_k] \mathcal{D}[\varphi_k] \mathcal{P}[\zeta] \mathcal{O}[\varphi] \delta[\varphi - \varphi_\zeta] \quad (14)$$

with φ_ζ indicating a solution of the Langevin equation (13) and $\mathcal{P}[\zeta]$ the Gaussian distribution of the noise ξ_k . We now perform some standard manipulations [24–27]. First, we perform the change of variable

$$\delta[\varphi - \varphi_\zeta] = \mathcal{J}[\varphi] \delta \left[\ddot{\varphi}_k + \mathbb{H}_{kq} \dot{\varphi}_q + \frac{1}{\tau} [\dot{\varphi}_k + F_k] - \zeta_k \right]$$

$$\mathcal{J}[\varphi] \equiv \left| \det \frac{\delta \xi_k(t)}{\delta \varphi_q(t')} \right|.$$

Second, we represent the delta-functional through a set of response fields $\hat{\varphi}_k$. Third, we perform the Gaussian integral over the noise ζ_k . In the following, we neglect the contribution of $\mathcal{J}[\varphi]$ since it does not contribute to the entropy production rate [28, 29]. These standard manipulations [25–27] lead to

$$\langle \mathcal{O} \rangle = \int \mathcal{D}[\varphi_k] \mathcal{D}[\hat{\varphi}_k] e^{-S[\hat{\varphi}, \varphi]} \mathcal{O}[\varphi], \quad (15)$$

where we $S[\hat{\varphi}, \varphi]$ is the so-called Jensen-De Dominicis response functional [24, 30, 31]

$$S[\hat{\varphi}, \varphi] \equiv \frac{D}{2\tau^2} \sum_k \int_{t_0}^{t_F} dt \hat{\varphi}_k^2$$

$$- \frac{1}{\tau} \sum_k \int_{t_0}^{t_F} dt \hat{\varphi}_k [\tau \dot{\varphi}_k + M_{kq} \dot{\varphi}_q + F_k]. \quad (16)$$

After performing the Gaussian integration over $\hat{\varphi}_k$, we finally obtain that the probability $P[\varphi]$ of a path $\{\varphi_k(t)\}_{t_0 \leq t \leq t_F}$ can be expressed in terms of the following Onsager-Machlup action (the proportional symbol indicates that we are neglecting factors that does not contribute to the computation of the entropy production rate)

$$P[\varphi] \propto \exp(-A[\varphi]) \quad (17)$$

$$A[\varphi] = \frac{1}{2D} \sum_k \int_{t_0}^{t_F} dt [\tau \ddot{\varphi}_k + M_{kq} \dot{\varphi}_q + F_k]^2.$$

Since we are interested in the critical behavior of the system by approaching the critical point *from above*, i. e., from the *disordered phase* where $\langle \varphi \rangle = 0$, the dynamical action $A[\varphi]$ is well-defined for every value of τ . In such a situation the Hessian matrix \mathbb{H}_{kq} is positive-definite and so M_{kq} is positive-definite as well.

The probability of the time-reversed path can be obtained by employing the time-reversal operator $t \rightarrow t_F - t$ that brings to the following expressions

$$P_R[\varphi] \propto \exp(-A_R[\varphi]) \quad (18)$$

$$A_R[\varphi] = \frac{1}{2D} \sum_k \int_{t_0}^{t_F - t_0} dt [\tau \ddot{\varphi}_k - M_{kq} \dot{\varphi}_q + F_k]^2.$$

ENTROPY PRODUCTION RATE

Now we can perform the computation of the entropy production rate using the expression

$$S = \lim_{t_F \rightarrow \infty} \frac{1}{2Dt_F} \sum_k \int_{t_0}^{t_F - t_0} dt \left\{ [\tau \ddot{\varphi}_k + M_{kq} \dot{\varphi}_q + F_k]^2 - [\tau \ddot{\varphi}_k - M_{kq} \dot{\varphi}_q + F_k]^2 \right\}, \quad (19)$$

that brings to

$$S = \lim_{t_F \rightarrow \infty} \frac{1}{Dt_F} \sum_q \int_{t_0}^{t_F - t_0} dt [A_q + B_q] \quad (20)$$

$$A_q \equiv \dot{\varphi}_q [\tau \ddot{\varphi}_q + F_q + \tau \mathbb{H}_{kq} F_k]$$

$$B_q \equiv \tau^2 \ddot{\varphi}_k \mathbb{H}_{kq} \dot{\varphi}_q.$$

The term A_q can be written as

$$A_q = \frac{d}{dt} Q_q \quad (21)$$

$$Q_q = \frac{\tau}{2} \dot{\varphi}_q^2 + H_{LG} + \tau F_q$$

whose time integral is a border term and thus vanishes in the limit $t_F \rightarrow \infty$ because of the factor t_F^{-1} . After integrating by parts B_q , we finally get

$$S = - \lim_{t_F \rightarrow \infty} \frac{\tau^2}{Dt_F} \int_{t_0}^{t_F - t_0} dt \mathbb{H}_{kq} \ddot{\varphi}_k \dot{\varphi}_q. \quad (22)$$

Performing an integration by parts and neglecting vanishing boundary terms, we obtain the following expression for the entropy production rate

$$S = \frac{\tau^2}{2D} \langle \dot{\varphi}_k \dot{\varphi}_q \dot{\varphi}_l G_{kql} \rangle \quad (23)$$

$$G_{kql} \equiv \frac{\partial^3 H_{LG}}{\partial \varphi_k \partial \varphi_q \partial \varphi_l} = \frac{\partial^3 H_I}{\partial \varphi_k \partial \varphi_q \partial \varphi_l}. \quad (24)$$

This is a central result of this paper: From Eqs. (23)-(24) we immediately realize that, in principle, all the non-linear terms in a Landau-Ginzburg model contribute to the entropy production. Moreover, the expression

obtained here generalized to field theories the entropy production rate formula obtained in the case of Active Ornstein-Uhlenbeck particles [12, 13]. Finally, no matter how big is τ , we notice that for a Gaussian model one has $S = 0$. Using Eqs. (23)-(24), S can be computed numerically by solving the equations of motion. This can be done, for instance, by discretizing the dynamics on a d -dimensional grid.

Once we perform the inverse Fourier transform, we get the following expression

$$S = \int dx \sigma(x) \quad (25)$$

$$\sigma \equiv \frac{\tau^2}{2D} \left\langle \dot{\varphi}(x)^3 \frac{\delta^3 H_I}{\delta \varphi(x)^3} \right\rangle,$$

where σ is the density of the entropy production rate.

The critical scaling

We now study the scaling of σ close to the critical point in the case of the φ^4 theory. For doing that, since φ satisfies the equation of motion (1), we can write

$$\sigma = \frac{\tau^2}{2D} \left\langle \left(-\frac{\delta H_{LG}}{\delta \varphi} + \eta \right)^3 \frac{\delta^3 H_{LG}}{\delta \varphi^3} \right\rangle_{\eta}. \quad (26)$$

Differently from Active Model A, where averages involve stochastic trajectories of φ driven by a delta-correlated noise, here the stochastic force η is time-correlated. Because of that we do not have, in general, a way for evaluating analytically the scaling of σ . However, since we aim to study the system in the vicinity of the critical point, we consider some reasonable assumptions for making the problem tractable. For making progress, we notice that, since τ is an irrelevant parameter, its renormalized value goes to zero on the critical surface. Once we approach the critical point we can thus replace the correlated noise over τ with a white noise that is delta-correlated. Moreover, as well as in the case of AOUPs [13], we replace the average over trajectories with an average over the stationary distribution $P_s[\varphi]$, i. e., $\langle \mathcal{O}[\varphi] \rangle_{\eta} = \int \mathcal{D}[\varphi] P_s[\varphi] \mathcal{O}[\varphi]$. Again, since τ is an irrelevant parameter these averages can be done by replacing the stationary distribution with the Boltzmann distribution, i. e., $P_s[\varphi] \simeq e^{-\beta H_{LG}}$, with β the inverse temperature sets to 1 in the following. Once we perform these approximations, we obtain that σ results from four contributions (in the following, the proportional symbol indicates that we are neglecting numer-

ical factors that are inessential for our purpose)

$$\begin{aligned}
\sigma &\propto \sigma_1 + \sigma_2 + \sigma_3 + \sigma_4 & (27) \\
\sigma_1 &\equiv \frac{\tau^2 u r^3}{D} \langle \varphi^4 \rangle \\
\sigma_2 &\equiv \frac{\tau^2 u \mu^3}{D} \langle (\nabla^2 \varphi)^3 \varphi \rangle \\
\sigma_3 &\equiv \frac{\tau^2 r^2 \mu}{D} \langle \varphi^3 \nabla^2 \varphi \rangle \\
\sigma_4 &\equiv \frac{\tau^2 u r \mu^2}{D} \langle (\nabla^2 \varphi)^2 \varphi^2 \rangle.
\end{aligned}$$

We will first consider the four contributions above at the Gaussian level, i.e., by performing averages using the Gaussian measure given by $P_s = e^{-H_0}$. In the Gaussian model, we obtain that the only non vanishing contribute in the limit $k \rightarrow 0$ is provided by σ_1 . In particular we get

$$\sigma_1 \propto \frac{\tau^2 u}{D} r^{1+d/2} \quad (28)$$

indicating that, at the mean-field level, i. e., above the upper critical dimension, the density of the entropy production rate tends to zero at the critical point, i. e., $\sigma \rightarrow 0$ for $r \rightarrow 0$. More in general, by performing the naive Gaussian scaling analysis of σ , we obtain that it follows the natural scaling, i. e., $\sigma \rightarrow \sigma b^{-(d+z)}$, where the arrow indicates how σ scales under Kadanoff transformations. As discussed in Ref. [6] in the case of Active Model A, the fact that σ goes to zero at the critical point does not guaranty that the dynamics is *effectively reversible* at criticality because what is really matter is the scaling of the singular part of the observable we are interested in. The suitable observable for doing that in the case of σ is the density of entropy production rate per spacetime correlation volume ψ defined as $\psi \equiv \xi^{d+z} \sigma$ [6]. Since $\xi \sim r^{-\nu}$, and the mean-field values are $\nu_{MF} = 1/2$ and $z_{MF} = 2$, we obtain that $\psi = \xi^{d+2} \xi^{-\frac{1}{\nu_{MF}} \frac{d+z}{2}} = 1$ meaning that, if we look for a singular part of ψ that behaves as $\psi \sim r^{-\theta_\sigma}$, we get the mean-field value $\theta_\sigma = 0$, that is in agreement also with Active Models A/B [6]. As discussed in details in [6], a non-negative value of θ_σ , i. e., $\theta_\sigma \geq 0$, indicates already that, although the universality class of the model remains untouched, the dynamics remains irreversible even at the critical point.

We shall now provide arguments that support an exponent θ_σ of order ϵ around the upper critical dimension. First of all, we notice that the only contribution to σ that depends on a relevant operator is σ_1 . In particular, the other composite operators in $\sigma_{2,3,4}$ have naive dimension $m+n > 4$, with m the number of fields and n the number of gradients. In order to understand the scaling of σ_4 , we consider the following perturbative RG transformation (at the lowest order in u). Using perturbation theory, one has $\langle \mathcal{O}[\varphi] \rangle = \int \mathcal{D}[\varphi] (1 - H_I) e^{-H_0} \mathcal{O}[\varphi]$ [18]. Since the observable we are going to consider is $\mathcal{O} = \varphi^4$, at the lowest order, one has $\sigma_1 = \frac{\tau^2 u r^3}{D} \langle \varphi^4 \rangle_{H_0}$. Once we consider the scaling transformations (4), the standard one-loop

RG equations bring to the following scaling [24, 32] (here we are using the fact that, under RG transformations, $r = r' b^{-(z+\epsilon/3)}$, $r\tau = r'\tau' = 1$ and $D = D' = 1$)

$$\sigma_1 = b^{\epsilon/3} b^{-(z+d)} r' u' \langle (\varphi')^4 \rangle \quad (29)$$

with $\epsilon = 4 - d$. As a result, in this framework, the scaling of ψ is $\psi \sim r^{-\theta_\sigma}$ with $\theta_\sigma = \frac{\nu\epsilon}{3}$ that is positive below the upper critical dimension.

DISCUSSION AND CONCLUSIONS

In this work, motivated by the experimental and numerical evidence that active baths always develop correlations on finite time scales [9–11], we have computed the entropy production rate in the case of a scalar field theory (with non-conserved order parameter) driven out-of-equilibrium by a persistent noise. The computation predicts non-vanishing entropy production rate only when non-linear interactions are taken into account (with a vanishing entropy production rate at equilibrium, i. e., that is recovered for $\tau = 0$). We have performed a scaling analysis of the density of entropy production rate σ near the critical point in a static picture by replacing the averages over the persistent dynamics with averages over equilibrium configurations. The computation provides evidence in favor of an exponent θ_σ that is positive below the upper critical dimension and negative above it. However, even in this simplified framework, for having a better estimate of the critical exponent finest computations involving the scaling of composite operators are required [22]. Moreover, it will be crucial to test those predictions against numerical data in three dimensions. As it has been shown in Ref. [6], a positive value of θ_σ indicates that, although the system share the same critical exponents of the Ising universality class, its dynamics at the critical point remains *effectively* a non-equilibrium dynamics and thus irreversible. Since the non-linear terms responsible for a non-vanishing σ in the model presented here are different from those in Active Model A, it is quite natural to obtain different critical scaling of σ . In Active Model A/B the leading non-equilibrium contribute is provided by a non-integrable gradient term, while the stochastic force at the coarse-grained level is considered equilibrium-like. Here we neglected non-equilibrium gradient terms and we have focused our attention on the role of a non-equilibrium noise (as a step forward, it might be interesting to consider both contributions). It is remarkable that, despite this qualitative difference in the non-equilibrium dynamics, the phenomenology of the two models are somehow related, in the sense that their non-equilibrium nature can be described by an additional critical exponent that might be the smoking gun for defining new non-equilibrium universality classes that are characterized by Ising universality class and positive exponent θ_σ .

As a future direction, the computation of σ presented here might be extended to the case of Model B and in

the presence of noise that is also correlated in space [11]. It might be also interesting to compare the phenomenological model presented here with recent coarse-graining description of active systems obtained from microscopic models [33].

ACKNOWLEDGMENTS

I am deeply grateful to Mattia Scandolo (as well as to all the CoBBS group in Rome) for his comments and his critical reading of the manuscript. I also thank Claudio Maggi and Andrea Puglisi for illuminating discussions and for their critical reading of the early version of the manuscript. This work has received funding from the European Union's Horizon 2020 research and innovation programme under the MSCA grant agreement No 801370 and by the Secretary of Universities and Research of the Government of Catalonia through Beatriu de Pinós program Grant No. BP 00088 (2018).

-
- [1] B. Partridge and C. F. Lee, *Phys. Rev. Lett.*, 2019, **123**, 068002.
- [2] C. Maggi, M. Paoluzzi, A. Crisanti, E. Zaccarelli and N. Gnan, *Soft Matter*, 2021, **17**, 3807–3812.
- [3] J. Tailleur and M. E. Cates, *Phys. Rev. Lett.*, 2008, **100**, 218103.
- [4] M. E. Cates and J. Tailleur, *Annu. Rev. Condens. Matter Phys.*, 2015, **6**, 219–244.
- [5] M. E. Cates, *arXiv preprint arXiv:1904.01330*, 2019.
- [6] F. Caballero and M. E. Cates, *Phys. Rev. Lett.*, 2020, **124**, 240604.
- [7] P. C. Hohenberg and B. I. Halperin, *Reviews of Modern Physics*, 1977, **49**, 435.
- [8] X.-L. Wu and A. Libchaber, *Phys. Rev. Lett.*, 2000, **84**, 3017–3020.
- [9] C. Maggi, M. Paoluzzi, N. Pellicciotta, A. Lepore, L. Angelani and R. Di Leonardo, *Phys. Rev. Lett.*, 2014, **113**, 238303.
- [10] C. Maggi, M. Paoluzzi, L. Angelani and R. Di Leonardo, *Scientific reports*, 2017, **7**, 17588.
- [11] C. Maggi, N. Gnan, M. Paoluzzi, A. Crisanti and E. Zaccarelli, *arXiv:2108.13971*, 2021.
- [12] L. Caprini, U. M. B. Marconi, A. Puglisi and A. Vulpiani, *Journal of Statistical Mechanics: Theory and Experiment*, 2019, **2019**, 053203.
- [13] E. Fodor, C. Nardini, M. E. Cates, J. Tailleur, P. Visco and F. van Wijland, *Phys. Rev. Lett.*, 2016, **117**, 038103.
- [14] R. Wittkowski, A. Tiribocchi, J. Stenhammar, R. J. Allen, D. Marenduzzo and M. E. Cates, *Nature communications*, 2014, **5**, 4351.
- [15] J. Stenhammar, A. Tiribocchi, R. J. Allen, D. Marenduzzo and M. E. Cates, *Physical review letters*, 2013, **111**, 145702.
- [16] J. Sancho, J. Garcia-Ojalvo and H. Guo, *Physica D: Non-linear Phenomena*, 1998, **113**, 331–337.
- [17] M. Paoluzzi, C. Maggi, U. Marini Bettolo Marconi and N. Gnan, *Phys. Rev. E*, 2016, **94**, 052602.
- [18] S.-K. Ma, *Modern theory of critical phenomena*, Routledge, 2018.
- [19] U. Seifert, *Reports on progress in physics*, 2012, **75**, 126001.
- [20] J. L. Lebowitz and H. Spohn, *Journal of Statistical Physics*, 1999, **95**, 333–365.
- [21] C. Nardini, É. Fodor, E. Tjhung, F. Van Wijland, J. Tailleur and M. E. Cates, *Physical Review X*, 2017, **7**, 021007.
- [22] D. J. Amit and V. Martin-Mayor, *Field theory, the renormalization group, and critical phenomena: graphs to computers*, World Scientific Publishing Company, 2005.
- [23] S.-k. Ma, *Reviews of Modern Physics*, 1973, **45**, 589.
- [24] U. C. Täuber, *Critical dynamics: a field theory approach to equilibrium and non-equilibrium scaling behavior*, Cambridge University Press, 2014.
- [25] P. C. Martin, E. D. Siggia and H. A. Rose, *Phys. Rev. A*, 1973, **8**, 423–437.
- [26] C. De Dominicis and L. Peliti, *Phys. Rev. B*, 1978, **18**, 353–376.
- [27] R. V. Jensen, *Journal of Statistical Physics*, 1981, **25**, 183–210.
- [28] R. E. Spinney and I. J. Ford, *Phys. Rev. Lett.*, 2012, **108**, 170603.
- [29] R. E. Spinney and I. J. Ford, *Phys. Rev. E*, 2012, **85**, 051113.
- [30] C. d. Dominicis, *J. Phys.(Paris)*, Colloq, 1976, pp. C1–247.
- [31] H.-K. Janssen, *Zeitschrift für Physik B Condensed Matter*, 1976, **23**, 377–380.
- [32] M. Le Bellac, *Quantum and statistical field theory*, Clarendon Press, 1991.
- [33] U. M. B. Marconi, A. Puglisi and L. Caprini, *arXiv preprint arXiv:2105.13832*, 2021.