

EXPONENTIAL APPROXIMATION IN VARIABLE EXPONENT LEBESGUE SPACES ON THE REAL LINE

RAMAZAN AKGÜN

Abstract Present work contains a method to obtain Jackson and Stechkin type inequalities of approximation by integral functions of finite degree (IFFD) in some variable exponent Lebesgue space of real functions defined on $\mathbf{R} := (-\infty, +\infty)$. To do this we employ a transference theorem which produce norm inequalities starting from norm inequalities in $\mathcal{C}(\mathbf{R})$, the class of bounded uniformly continuous functions defined on \mathbf{R} . Let $B \subseteq \mathbf{R}$ be a measurable set, $p(x) : B \rightarrow [1, \infty)$ be a measurable function. For the class of functions f belonging to variable exponent Lebesgue spaces $L_{p(x)}(B)$ we consider difference operator $(I - T_\delta)^r f(\cdot)$ under the condition that $p(x)$ satisfies the Log Hölder continuity condition and $1 \leq \text{ess inf}_{x \in B} p(x)$, $\text{ess sup}_{x \in B} p(x) < \infty$ where I is the identity operator, $r \in \mathbf{N} := \{1, 2, 3, \dots\}$, $\delta \geq 0$ and

$$T_\delta f(x) = \frac{1}{\delta} \int_0^\delta f(x+t) dt, \quad x \in \mathbf{R}, \quad T_0 \equiv I, \quad (*)$$

is the forward Steklov operator. It is proved that

$$\|(I - T_\delta)^r f\|_{p(\cdot)}, \quad (**)$$

is a suitable measure of smoothness for functions in $L_{p(x)}(B)$ where $\|\cdot\|_{p(\cdot)}$ is Luxemburg norm in $L_{p(x)}(B)$. We obtain main properties of difference operator $\|(I - T_\delta)^r f\|_{p(\cdot)}$ in $L_{p(x)}(B)$. We give proof of direct and inverse theorems of approximation by IFFD in $L_{p(x)}(\mathbf{R})$.

Key Words Variable exponent Lebesgue space, One sided Steklov operator, Integral functions of finite degree, Best approximation, Direct theorem, Inverse theorem, Modulus of smoothness, Marchaud inequality, K-functional.

2010 Mathematics Subject Classifications 41A10; 41A25; 41A27; 41A65.

1. INTRODUCTION

Some inequalities of Approximation Theory in a Homogenous Banach Spaces (HBS) can be obtained their uniform-norm counterparts. This information is known for a long time, (see e.g., [21] for definition of HBS). This elegant method was generalized to some variable exponent Lebesgue spaces functions defined on \mathbf{R} . (see Theorem 21 of [9]). Generally, these scale of function classes are non-translation invariant with respect to the ordinary translation $x \rightarrow f(x+a)$. Here we give several uniform-norm

inequalities on $C(\mathbf{R})$ and apply them to obtain several inequalities of approximation by IFFD in some variable exponent Lebesgue spaces $L_{p(x)}(\mathbf{R})$. Under some condition on $p(x)$ of $L_{p(x)}(\mathbf{R})$ we obtain main inequalities of exponential approximation by IFFD such as Jackson-Stechkin-Timan type estimates and equivalence of K -functional with suitable modulus of smoothness (***) given in abstract for functions of $L_{p(x)}(\mathbf{R})$. Note that many results of approximation by IFFD can be obtained easily their uniform-norm counterparts in $C(\mathbf{R})$.

Consider an entire function $f(z)$ and put $M(r) = \max_{|z|=r} |f(z)|$ for $z = x + iy$. We say that an entire function f is of exponential type σ if $\limsup_{r \rightarrow \infty} r^{-1} \ln M(r) \leq \sigma$, $\sigma < \infty$.

The approximation by entire function of finite degree in the real line was originated in the beginning of twentieth century by Serge Bernstein [15] and became a separate branch of analysis due to the efforts of many mathematicians such as N. Wiener and R. Paley [46], N.I. Ahiezer [4], S.M. Nikolskii [43], I.I. Ibragimov [30], A. F. Timan [53], M. F. Timan [54], R. Taberski [55, 56], F.G. Nasibov [42], V. Yu. Popov [47], A. A. Ligun [44], and others.

Studying function spaces with variable exponent is now an extensively developed field after their applications in elasticity theory [59], fluid mechanics [48, 49], differential operators [20, 49], nonlinear Dirichlet boundary value problems [41], nonstandard growth [59], and variational calculus. See the books [16, 19, 52] for more references. Nowadays many mathematician solved many problems for the approximation of function in these type spaces defined on $[0, 2\pi] \subset \mathbf{R}$ (see e.g., [7, 8, 27, 31, 32, 35], [1, 2, 3, 11, 12], [5, 6, 9, 13, 14], [23, 25, 26, 29, 33, 34, 37], [38, 39, 45, 50, 51, 57]). In this paper we propose generalized our last results in [10] which we obtained a direct and inverse theorems for approximation by entire functions of finite degree in variable exponent Lebesgue spaces on the whole real axis \mathbf{R} with

$$(1.1) \quad \sup_{0 < h \leq \delta} \|(I - T_h)f\|_{p(\cdot)}$$

as modulus of continuity $\Omega_1(f, \delta)_{p(\cdot)}$. Instead of (1.1), here we will use

$$(1.2) \quad \|(I - T_\delta)^r f\|_{p(\cdot)}$$

as modulus smoothness $\Omega_r(f, \delta)_{p(\cdot)}$ and we obtain stronger Jackson inequality than obtained in [10].

Let $B \subseteq \mathbf{R}$ be a measurable set and $p(x) : B \rightarrow [1, \infty)$ be a measurable function. We define $\tilde{P}(B)$ as the class of measurable functions $p(x)$ satisfying the conditions

$$(1.3) \quad 1 \leq p_B^- := \text{ess inf}_{x \in B} p(x), \quad p_B^+ := \text{ess sup}_{x \in B} p(x) < \infty.$$

We also set $p^- := p_B^-$ and $p^+ := p_B^+$. We define the $L_{p(\cdot)}(B)$ as the set of all functions $f : B \rightarrow \mathbf{R}$ such that

$$(1.4) \quad I_{p(\cdot), B} \left(\frac{f}{\lambda} \right) := \int_B \left| \frac{f(y)}{\lambda} \right|^{p(y)} dy < \infty$$

for some $\lambda > 0$. We set $I_{p(\cdot)}(f) := I_{p(\cdot), \mathbf{R}}(f)$. The set of functions $L_{p(\cdot)}(B)$, with norm

$$\|f\|_{p(\cdot), B} := \inf \left\{ \eta > 0 : I_{p(\cdot), B} \left(\frac{f}{\eta} \right) < 1 \right\}$$

is Banach space. We set $L_{p(\cdot)} := L_{p(\cdot)}(\mathbf{R})$.

For $i \in \mathbb{N}$, all constants $c_i(x, y, \dots)$ will be some positive number such that they depend on the parameters x, y, \dots given in the brackets. Also constants $c_i(x, y, \dots)$ can be change only when the parameters x, y, \dots change. Absolute constants $\mathbf{c}_1, \mathbf{c}_2, \dots$ will not change in each occurrence.

Definition 1.1. For a measurable set $B \subseteq \mathbf{R}$, a measurable function $p(\cdot) : B \rightarrow \mathbf{R}$ is said to locally log-Hölder continuous on B if there is a positive constant $c_1(p)$ such that

$$(1.5) \quad |p(x) - p(y)| \log(e + 1/|x - y|) \leq c_1(p) < \infty$$

for any $x, y \in B$. We say that p satisfies log-Hölder decay condition if there is a constant $c_2(p) > 0$ and $p_\infty > 1$ such that

$$(1.6) \quad |p(x) - p_\infty| \log(e + |x|) \leq c_2(p) < \infty$$

for any $x \in B$.

Define the class $P^{Log}(B) := \left\{ p \in \tilde{P}(B) : \frac{1}{p} \text{ is satisfy (1.5)-(1.6)} \right\}$. We set $c_3(p) := \max\{c_1(p), c_2(p)\}$.

Definition 1.2. ([28, p.96]) Let $\mathbb{N} := \{1, 2, 3, \dots\}$ be natural numbers and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

(a) A family Q of measurable sets $E \subset \mathbf{R}$ is called locally N -finite ($N \in \mathbb{N}$) if

$$\sum_{E \in Q} \chi_E(x) \leq N$$

almost everywhere in \mathbf{R} where χ_U is the characteristic function of the set U .

(b) A family Q of open bounded sets $U \subset \mathbf{R}$ is locally 1-finite if and only if the sets $U \in Q$ are pairwise disjoint.

(c) Let $U \subset \mathbf{R}$ be a measurable set and

$$A_U f := \frac{1}{|U|} \int_U |f(t)| dt.$$

(d) For a family Q of open sets $U \subset \mathbf{R}$ we define averaging operator by

$$T_Q : L_{loc}^1 \rightarrow L^0,$$

$$T_Q f(x) := \sum_{U \in Q} \chi_U(x) A_U f = \sum_{U \in Q} \frac{\chi_U(x)}{|U|} \int_U |f(y)| dy, \quad x \in \mathbf{R},$$

where L^0 is the set of measurable functions on \mathbf{R} .

For a measurable set $A \subset \mathbf{R}$, symbol $|A|$ will represent the Lebesgue measure of A .

We consider Transference Result.

Definition 1.3. For $0 < \delta < \infty$, $\tau \in \mathbf{R}$, we define family of Steklov operators

$$(1.7) \quad S_\delta f(x) := \frac{1}{\delta} \int_{x-\delta/2}^{x+\delta/2} f(t) dt = \frac{1}{\delta} \int_{-\delta/2}^{\delta/2} f(x+t) dt, \quad x \in \mathbf{R},$$

where f is a locally integrable function, defined on \mathbf{R} .

The following result was obtained by Drihem for every cubes or balls in \mathbf{R}^d . We write below its restricted version with constants. The proof of this is the same with Theorem 2 of [24].

Proposition 1.4. ([24]) *Suppose that $p \in P^{Log}(\mathbf{R})$ and Q is a bounded interval of \mathbf{R} having Lebesgue measure greater than 1. For every $m > 0$ there is $c_4(m, c_3(p)) := \exp(-4mc_3(p)) \in (0, 1)$ such that*

$$\begin{aligned} \left(\frac{c_4(m, c_3(p))}{|Q|} \int_Q |f(y+\tau)| dy \right)^{p(x)} &\leq \frac{3^{p^+}}{|Q|} \int_Q |f(y+\tau)|^{p(y+\tau)} dy + \frac{3^{p^+-1}}{(e+|x|)^m} + \\ &+ 3^{p^+-1} \int_Q \frac{dy}{(e+|y+\tau|)^m} \end{aligned}$$

holds for all $x \in Q$, $\tau \in \mathbf{R}$ and all $f \in L_{p(\cdot)} + L_\infty(\mathbf{R})$ with $\|f\|_{p(\cdot)} + \|f\|_\infty \leq 1$.

Theorem 1.5. *Suppose that $p \in P^{Log}(\mathbf{R})$. Then, the family of operators $\{\mathcal{U}_\tau f\}_{\tau \in \mathbf{R}}$, defined by*

$$\mathcal{U}_\tau f(x) := \mathbf{S}_1 f(x+\tau) = \int_{-1/2}^{+1/2} f(x+\tau+t) dt, \quad x \in \mathbf{R}, \quad \tau \in \mathbf{R}$$

is uniformly bounded (in τ) in $L_{p(\cdot)}$, namely,

$$\|\mathcal{U}_\tau f\|_{p(\cdot)} \leq c_5(p^+, c_3(p)) \|f\|_{p(\cdot)},$$

holds with $c_5(p^+, c_3(p)) := 2^{p^++1} 3^{p^+} \left(1 + 2 \cdot 3^{p^+} \left[\sum_{k=2}^\infty 2^{-k} + 2\right]\right) \exp(8c_3(p))$.

Proof of Theorem 1.5. Let us consider $f \in L_{p(\cdot)}$ with $\|f\|_{p(\cdot)} \leq 1/2$. Suppose that $Q := \{U \subset \mathbf{R} : U \text{ open interval and } |U| = 1\}$ be a locally 1-finite family of partition of \mathbf{R} . Choose $m = 2 > 1$ (constant $c_6(p^+)$ below becomes a finite number)

$$c_6(p^+) = 2^{p^+} 3^{p^+} \left(1 + 2 \cdot 3^{p^+} \left[\sum_{k=2}^\infty 2^{-k} + 2\right]\right) < \infty.$$

We can select $c_4(2, c_3(p)) = \exp(-8c_3(p)) \in (0, 1)$ as in Proposition 1.4. Then using Corollary 2.2.2 of [28, p.20] we obtain

$$\rho_{p(\cdot)} \left(\frac{c_4(2, c_3(p))}{c_6(p^+)} \mathcal{U}_\tau f \right) = \frac{1}{c_6(p^+)} \int_{\mathbf{R}} \left| c_4(2, c_3(p)) \int_{-1/2}^{+1/2} f(x+\tau+t) dt \right|^{p(x)} dx$$

$$\begin{aligned}
&\leq \frac{1}{c_6(p^+)} \sum_{U \in Q} \int_U \left| c_4(2, c_3(p)) \int_{-1/2}^{+1/2} f(x + \tau + t) dt \right|^{p(x)} dx \\
&\leq \frac{2^{p^+}}{c_6(p^+)} \sum_{U \in Q} \int_U \left| \frac{c_4(2, c_3(p))}{|2U|} \int_{2U} \chi_{2U}(y) f(y + \tau) dy \right|^{p(x)} dx \\
&\leq \frac{2^{p^+}}{c_6(p^+)} \sum_{U \in Q} \int_U \left[\frac{3^{p^+} \chi_{2U}(y)}{|2U|} \int_{2U} |f(y + \tau)|^{p(y+\tau)} dy + \right. \\
&\quad \left. + \frac{3^{p^+-1}}{(e + |x|)^2} + \frac{\chi_{2U}(y)}{|2U|} \int_{2U} \frac{3^{p^+-1} dy}{(e + |y + \tau|)^2} \right] dx \\
&\leq \frac{2^{p^+-1} 3^{p^+}}{c_6(p^+)} \sum_{U \in Q} \int_U \left[\chi_{2U}(y) \int_{2U+\tau} |f(s)|^{p(s)} ds + \frac{3^{p^+-1} 2}{(e + |x|)^2} + \int_{2U+\tau} \frac{3^{p^+-1} ds}{(e + |s|)^2} \right] dx \\
&\leq \frac{2^{p^+-1} 3^{p^+}}{c_6(p^+)} \left(\sum_{U \in Q} \chi_{2U} \right) \left(1 + 3^{p^+} \int_{\mathbf{R}} \frac{ds}{(e + |s|)^2} \right) \\
&= \frac{2^{p^+} 3^{p^+}}{c_6(p^+)} \left(1 + 3^{p^+} \int_{\mathbf{R}} \frac{ds}{(e + |s|)^2} \right) \\
&\leq \frac{2^{p^+} 3^{p^+}}{c_6(p^+)} \left(1 + 2 \cdot 3^{p^+} \left[\sum_{k=2}^{\infty} \frac{1}{2^k} + 2 \right] \right) = 1
\end{aligned}$$

and hence

$$\|\mathcal{U}_\tau f\|_{p(\cdot)} \leq 2^{-1} c_5(p^+, c_3(p)).$$

General case $f \in L_{p(\cdot)}$ can be obtained easily by rescaling:

$$\|\mathcal{U}_\tau f\|_{p(\cdot)} \leq c_5(p^+, c_3(p)) \|f\|_{p(\cdot)}.$$

□

Theorem 1.6. ([19, Theorem 4.4.8]) *Suppose that $p \in P^{Log}(\mathbf{R})$ and $f \in L_{p(\cdot)}$. If Q is locally 1-finite family of open bounded subintervals of \mathbf{R} having Lebesgue measure 1, then, the averaging operator T_Q is uniformly bounded in $L_{p(\cdot)}$, namely,*

$$\|T_Q f\|_{p(\cdot)} \leq c_7(c_{\log}(p)) \|f\|_{p(\cdot)}$$

holds with $c_7(c_3(p)) := 2 \exp(8c_3(p))$.

Let $C(\mathbf{R})$ be the class of continuous functions defined on \mathbf{R} . For $r \in \mathbf{N}$, we define C^r consisting of every member $f \in C(\mathbf{R})$ such that the derivative $f^{(k)}$ exists and is continuous on \mathbf{R} for $k = 1, \dots, r$. We set $C^\infty := \{f \in C^r \text{ for any } r \in \mathbf{N}\}$. We denote by $C_c(\mathbf{R})$, the collection of real valued continuous functions on \mathbf{R} and support of f is compact set in \mathbf{R} . We define $C_c^r := C^r \cap C_c(\mathbf{R})$ for $r \in \mathbf{N}$ and $C_c^\infty := C^\infty \cap C_c(\mathbf{R})$. Let $L_p(\mathbf{R})$, $1 \leq p \leq \infty$ be the classical Lebesgue space of functions on \mathbf{R} .

Theorem 1.7. [19, Corollary 4.6.6] *Let $p \in P^{Log}(\mathbf{R})$ and $f \in L_{p(\cdot)}$. Then*

$$(1.8) \quad \frac{\|f\|_{p(\cdot)}}{12c_7(c_3(p))} \leq \sup_{g \in L_{p'(\cdot)} \cap C_c^\infty: \|g\|_{p'(\cdot)} \leq 1} \int_{\mathbf{R}} |f(x)g(x)| dx \leq 2\|f\|_{p(\cdot)}$$

Definition 1.8. Let $p \in P^{Log}(\mathbf{R})$. For an $f \in L_{p(\cdot)}$, we define

$$(1.9) \quad F_f(u) := \int_{\mathbf{R}} (\mathbf{S}_1 f)(x+u) |G(x)| dx, \quad u \in \mathbf{R},$$

where $G \in L_{p'(\cdot)} \cap C_c^\infty$ and $\|G\|_{p'(\cdot)} \leq 1$.

Let $W_{p(\cdot)}^r$, $r \in \mathbf{N}$, be the class of functions $f \in L_{p(\cdot)}$ such that derivatives $f^{(k)}$ exist for $k = 1, \dots, r-1$, $f^{(r-1)}$ absolutely continuous and $f^{(r)} \in L_{p(\cdot)}$.

Some properties of the function $F_f(\cdot)$ is given in the following theorem.

Theorem 1.9. *Let $p \in P^{Log}(\mathbf{R})$, $0 < \delta < \infty$, and $f \in L_{p(\cdot)}$. Then,*

- (a) *the function $F_f(\cdot)$ defined in (1.9) is a bounded, uniformly continuous on \mathbf{R} ,*
- (b) *$(\mathbf{S}_\delta f)' = \mathbf{S}_\delta(f')$ on \mathbf{R} for $f \in W_{p(\cdot)}^1$.*

Main theorem of this section is as follows.

Theorem 1.10. *Let $p \in P^{Log}(\mathbf{R})$. If $f, g \in L_{p(\cdot)}$ and*

$$\|F_f\|_{C(\mathbf{R})} \leq \mathbf{c}_1 \|F_g\|_{C(\mathbf{R})},$$

holds with an absolute constant $\mathbf{c}_1 > 0$, then, norm inequality

$$(1.10) \quad \|f\|_{p(\cdot)} \leq c_8(\mathbf{c}_1, p^+, c_3(p)) \|g\|_{p(\cdot)}$$

also holds with $c_8(\mathbf{c}_1, p^+, c_3(p)) := 48c_7(c_3(p)) \mathbf{c}_1 c_5(p^+, c_3(p))$.

For the proof of these results we will need the following Propositions.

Proposition 1.11. (a) $C_c(\mathbf{R})$ and C_c^∞ are dense subsets of $L_p(\mathbf{R})$, $1 \leq p < \infty$. (Theorems 17.10 and 23.59 of [58, p. 415 and p. 575]).

(b) $C_c(\mathbf{R})$ contained $L_\infty(\mathbf{R})$ but not dense (Remark 17.11 of [58, p.416]) in $L_\infty(\mathbf{R})$.

(c) if $r \in \mathbf{N}$ and $f \in C_c^r$, then, $\mathbf{S}_\delta(f) \in C_c^r$.

Proposition 1.12. ([19, Theorem 2.26]) *Let $B \subseteq \mathbf{R}$ be a measurable set. If $1 \leq p(x) < p_B^+ < \infty$, $p'(x) = p(x)/(p(x) - 1)$, $f \in L_{p(\cdot)}(B)$, and $g \in L_{p'(\cdot)}(B)$, then, Hölder's inequality*

$$(1.11) \quad \int_B f(x)g(x)dx \leq 2\|f\|_{p(\cdot),B} \|g\|_{p'(\cdot),B}$$

holds.

1.0.1. *Proofs of Section 1.*

Proof of Proposition 1.11. (a) and (b) are known. (c) is follows from definitions. \square

Proof of Theorem 1.9. (a) Since $C_c(\mathbf{R})$ is a dense subset ([40, Theorem 4.1 (I)]) of $L_{p(\cdot)}$, we consider functions $H \in C_c(\mathbf{R})$ and prove that $F_H(\cdot) = \int_{\mathbf{R}} (S_1 H)(x + u_1) |G(x)| dx$ is bounded and uniformly continuous on \mathbf{R} , where $G \in L_{p'(\cdot)} \cap C_c^\infty$ and $\|G\|_{p'(\cdot)} \leq 1$. Boundedness of $F_H(\cdot)$ is easy consequence of the Hölder's inequality (1.11) and Theorem 1.5. On the other hand, note that H is uniformly continuous on \mathbf{R} , see e.g. Lemma 23.42 of [58, pp.557-558]. Take $\varepsilon > 0$ and $u_1, u_2, x \in \mathbf{R}$. Then, there exists a $\delta := \delta(\varepsilon) > 0$ such that

$$|H(x + u_1) - H(x + u_2)| \leq \frac{\varepsilon}{2(1 + |\text{supp}(G)|)}$$

for $|u_1 - u_2| < \delta$. Then, for $|u_1 - u_2| < \delta$, $u_1, u_2 \in \mathbf{R}$ we have

$$\begin{aligned} |F_H(u_1) - F_H(u_2)| &= \left| \int_{\mathbf{R}} S_1(H(x + u_1) - H(x + u_2)) |G(x)| dx \right| \\ &\leq \frac{1}{2(1 + |\text{supp}(G)|)} \int_{\mathbf{R}} |S_1(\varepsilon)| |G(x)| dx = \frac{\varepsilon}{2(1 + |\text{supp}(G)|)} \int_{\mathbf{R}} |G(x)| dx \\ &\leq \frac{\varepsilon}{(1 + |\text{supp}(G)|)} (1 + |\text{supp}(G)|) \|G\|_{p'(\cdot)} \leq \varepsilon. \end{aligned}$$

Now, the conclusion of Theorem 1.9 follows for the class $C_c(\mathbf{R})$. For the general case $f \in L_{p(\cdot)}$ there exists an $H \in C_c(\mathbf{R})$ so that

$$\|f - H\|_{p(\cdot)} < \xi/8c$$

for any $\xi > 0$. Then, for this ξ ,

$$\begin{aligned} |F_f(u_1) - F_f(u_2)| &\leq \left| \int_{\mathbf{R}} S_1(f - H)(x + u_1) |G(x)| dx \right| + \\ &\quad + \left| \int_{\mathbf{R}} S_1(H(x + u_1) - H(x + u_2)) |G(x)| dx \right| \\ &\quad + \left| \int_{\mathbf{R}} S_1(H - f)(x + u_2) |G(x)| dx \right| \\ &\leq 2 \|S_1(f - H)(\cdot + u_1)\|_{p(\cdot)} + \left| \int_{\mathbf{R}} S_1(H(x + u_1) - H(x + u_2)) |G(x)| dx \right| \\ &\quad + 2 \|S_1(f - H)(\cdot + u_2)\|_{p(\cdot)} \\ &\leq 4c \|f - H\|_{p(\cdot)} + \xi/2 \leq \xi/2 + \xi/2 = \xi \end{aligned}$$

As a result F_f is bounded, uniformly continuous function defined on \mathbf{R} .

(b) can be obtained easily from definition. \square

Proof of Theorem 1.10. Let $f \in L_{p(\cdot)}$ be nonnegative. If $\|f\|_{p(\cdot)} = 0$, then, the result (1.10) is obvious. So we assume that $\infty > \|f\|_{p(\cdot)} > 0$. In this case

$$\begin{aligned} \|F_f\|_{C(\mathbf{R})} &\leq \mathbf{c}_1 \|F_g\|_{C(\mathbf{R})} = \mathbf{c}_1 \left\| \int_{\mathbf{R}} \mathbf{S}_1(g)(u+x) |G(x)| dx \right\|_{C(\mathbf{R})} \\ &= \mathbf{c}_1 \max_{u \in \mathbf{R}} \left| \int_{\mathbf{R}} \mathbf{S}_1(g)(u+x) |G(x)| dx \right| \\ &\leq 2\mathbf{c}_1 \max_{u \in \mathbf{R}} \|\mathbf{S}_1(g)(u+\cdot)\|_{p(\cdot)} \leq 2c_5(p^+, c_3(p)) \mathbf{c}_1 \|g\|_{p(\cdot)} \end{aligned}$$

where we used hypothesis, Hölder's inequality and Theorem 1.5 respectively. On the other hand, for any $\varepsilon \in \left(0, \frac{\|f\|_{p(\cdot)}}{12c_7(c_3(p))}\right)$ and appropriately chosen $\tilde{G}_\varepsilon \in L_{p'(\cdot)}$ with $\|\tilde{G}_\varepsilon\|_{X'} \leq 1$ (see e.g. Theorem 1.7)

$$\int_{\mathbf{R}} |g(x)| |\tilde{G}_\varepsilon(x)| dx \geq \frac{1}{12c_7(c_3(p))} \|g\|_{p(\cdot)} - \varepsilon,$$

one can find

$$\begin{aligned} \|F_f\|_{C(\mathbf{R})} &\geq |F_f(0)| \geq \int_{\mathbf{R}} \mathbf{S}_1(f)(x) |G(x)| dx \\ &= \mathbf{S}_1\left(\int_{\mathbf{R}} f(x) |G(x)| dx\right) \geq \mathbf{S}_1\left(\frac{1}{12c_7(c_3(p))} \|f\|_{p(\cdot)} - \varepsilon\right) \\ &= \frac{1}{12c_7(c_3(p))} \|f\|_{p(\cdot)} - \varepsilon. \end{aligned}$$

In the last inequality we take as $\varepsilon \rightarrow 0+$ and obtain

$$\|F_f\|_{C(\mathbf{R})} \geq \frac{1}{12c_7(c_3(p))} \|f\|_{p(\cdot)}.$$

Then for $f \in L_{p(\cdot)}$ we get

$$\begin{aligned} \|f\|_{p(\cdot)} &\leq 24c_7(c_3(p)) \|F_f\|_{C(\mathbf{R})} \leq 24c_7(c_3(p)) \mathbf{c}_1 \|F_g\|_{C(\mathbf{R})} \\ &\leq 48c_7(c_3(p)) \mathbf{c}_1 c_5(p^+, c_3(p)) \|g\|_{p(\cdot)}. \end{aligned}$$

□

Remark 1.13. Theorem 1.10 is a powerful tool to obtain norm inequalities in $L_{p(\cdot)}$ (and other non-translation invariant Banach spaces of functions) for $p \in P^{Log}(\mathbf{R})$. In this work we will use it frequently. See for example the following result.

As a corollaries of Theorem 1.10 we get the following two results:

Theorem 1.14. *Suppose that $p \in P^{Log}(\mathbf{R})$, $0 < \delta < \infty$ and $\tau \in \mathbf{R}$. Then, the family of operators $\{\mathcal{S}_{\delta,\tau}f\}$, defined by*

$$\mathcal{S}_{\delta,\tau}f(x) := \mathbf{S}_\delta f(\cdot + \tau) = \frac{1}{\delta} \int_{x+\tau-\delta/2}^{x+\tau+\delta/2} f(s) ds, \quad x \in \mathbf{R},$$

is uniformly bounded (in δ and τ) in $L_{p(\cdot)}$, namely,

$$\|\mathcal{S}_{\delta,\tau}f\|_{p(\cdot)} \leq 48c_7(c_3(p)) c_5(p^+, c_3(p)) \|f\|_{p(\cdot)}$$

holds.

Corollary 1.15. *Let $p \in P^{Log}(\mathbf{R})$, $0 < \delta < \infty$, and $f \in L_{p(\cdot)}$. If $\tau = \delta/2$ then,*

$$\mathcal{S}_{\delta, \delta/2} f(x) = \frac{1}{\delta} \int_0^\delta f(x+t) dt = T_\delta f(x),$$

$$(1.12) \quad \begin{aligned} \|T_\delta f\|_{p(\cdot)} &\leq 48c_7(c_3(p)) c_5(p^+, c_3(p)) \|f\|_{p(\cdot)} \quad \text{and} \\ \|(I - T_\delta)^r f\|_{p(\cdot)} &\leq (1 + 48c_7(c_3(p)) c_5(p^+, c_3(p)))^r \|f\|_{p(\cdot)}. \end{aligned}$$

Definition 1.16. For $p \in P^{Log}(\mathbf{R})$, $f \in L_{p(\cdot)}$, $0 < \delta < \infty$, $r \in \mathbf{N}_0$, we can define modulus of smoothness as

$$\begin{aligned} \Omega_r(f, \delta)_{p(\cdot)} &= \|(I - T_\delta)^r f\|_{p(\cdot)} \\ \Omega_0(f, \delta)_{p(\cdot)} &:= \|f\|_{p(\cdot)} =: \Omega_r(f, 0)_{p(\cdot)}. \end{aligned}$$

2. UNIFORM NORM ESTIMATES

In this section, let $\Omega \subseteq \mathbf{R}$ be a measurable set and $C(\Omega)$ be the collection of functions continuous on Ω . If $\Omega \neq \mathbf{R}$ and $f \in C(\Omega)$, we will extend f to whole \mathbf{R} by " $f(s) \equiv 0$ whenever $s \notin \Omega$." when necessary. For $f \in C(\Omega)$ and $\delta \geq 0$, we define the modulus of smoothness as

$$(2.1) \quad \begin{aligned} \Omega_r(f, \delta)_{C(\Omega)} &:= \|(I - T_\delta)^r f\|_{C(\Omega)}, \quad r \in \mathbf{N}, \\ \Omega_0(f, \cdot)_{C(\Omega)} &:= \|f\|_{C(\Omega)} \end{aligned}$$

with $T_\delta f$ of (*).

Lemma 2.1. *Let $0 \leq \delta < \infty$, $r \in \mathbf{N}$ and $f \in C^r(\Omega)$. Then*

$$(2.2) \quad \frac{d^r}{dx^r} T_\delta f(x) = T_\delta \frac{d^r}{dx^r} f(x) \quad \text{on } \Omega.$$

The following theorem states the main properties of (2.1).

Theorem 2.2. *For $f \in C(\Omega)$, $0 \leq \delta < \infty$, and $r \in \mathbf{N}$, the following properties hold.*

- (1) $\Omega_r(f, \delta)_{C(\Omega)}$ is non-negative, non-decreasing function of δ ,
- (2) $\Omega_r(f, \delta)_{C(\Omega)}$ is subadditive with respect to f ,
- (3) $\|T_\delta f\|_{C(\Omega)} \leq \|f\|_{C(\Omega)}$,
- (4) $\Omega_r(f, \delta)_{C(\Omega)} \leq 2\Omega_{r-1}(f, \delta)_{C(\Omega)} \leq \cdots \leq 2^{r-1}\Omega_1(f, \delta)_{C(\Omega)} \leq 2^r \|f\|_{C(\Omega)}$, (***)
- (5) $\Omega_r(f, \delta)_{C(\Omega)} \leq 2^{-1}\delta\Omega_{r-1}(f', \delta)_{C(\Omega)} \leq \cdots \leq 2^{-r}\delta^r \|f^{(r)}\|_{C(\Omega)}$, if $f \in C^r(\Omega)$.

Let X be a Banach space with a norm $\|\cdot\|_X$ and $r \in \mathbf{N}$. We define Peetre's K -functional for the pair X and W_X^r as follows :

$$K_r(f, \delta, X)_X := \inf_{g \in W_X^r} \{ \|f - g\|_X + \delta^r \|g^{(r)}\|_X \}, \quad \delta > 0.$$

We set $T_\delta^r f := (T_\delta f)^r$.

Lemma 2.3. *Let $0 \leq \delta < \infty$, $r - 1 \in \mathbb{N}$, and $f \in C^r(\Omega)$ be given. Then*

$$(2.3) \quad \frac{d^r}{dx^r} T_\delta^r f(x) = \frac{d}{dx} T_\delta \frac{d^{r-1}}{dx^{r-1}} T_\delta^{r-1} f(x) \quad \text{on } \Omega.$$

Lemma 2.4. *(see e.g. [18, p.177]) Let $\Omega \subseteq \mathbf{R}$ be a measurable set, $\delta > 0$, $f \in C(\Omega)$ and $\tilde{T}_\delta f(\cdot) = f(\cdot + \delta)$. Then, for any $r \in \mathbb{N}$, there holds*

$$\frac{1}{r^r + 2^r} \leq \frac{\sup_{|h| \leq \delta} \left\| \left(I - \tilde{T}_h \right)^r f \right\|_{C(\Omega)}}{K_r(f, \delta, L_p(\Omega))_{C(\Omega)}} \leq 2^r.$$

Main result of this section is the following theorem.

Theorem 2.5. *Let $\Omega \subseteq \mathbf{R}$ be a measurable set, $0 < \delta < \infty$, $f \in C(\Omega)$, $r \in \mathbb{N}$ and $g \in C^2(\Omega)$. Then, the following inequalities*

$$\begin{aligned} \left\| \frac{d}{dx} T_\delta f(x) \right\|_{C(\Omega)} &\leq \frac{2}{\delta} \|f\|_{C(\Omega)}, \\ \left\| \frac{d^2}{dx^2} T_\delta f(x) \right\|_{C(\Omega)} &\leq \frac{2}{\delta} \left\| \frac{d}{dx} T_\delta f \right\|_{C(\Omega)}, \\ \left\| g(x) - T_\delta g(x) + \frac{\delta}{2} \frac{d}{dx} g(x) \right\|_{C(\Omega)} &\leq \frac{\delta^2}{6} \left\| \frac{d^2}{dx^2} g \right\|_{C(\Omega)}, \end{aligned}$$

$$(2.4) \quad (c_8(r))^{-1} K_r(f, \delta, C(\Omega))_{C(\Omega)} \leq \|(I - T_\delta)^r f\|_{C(\Omega)} \leq 2^r K_r(f, \delta, C(\Omega))_{C(\Omega)},$$

are hold with $c_8(1) = 36$, $c_8(r) = 2^r (r^r + (34)^r)$ for $r > 1$.

As a corollary of Theorem 2.5 we can state the following result.

Proposition 2.6. *If $0 < h \leq \delta < \infty$ and $f \in C(\Omega)$, then*

$$(2.5) \quad \|(I - T_h) f\|_{C(\Omega)} \leq 72 \|(I - T_\delta) f\|_{C(\Omega)}.$$

As a corollary of (2.4) and Lemma 2.4 we can write

Corollary 2.7. *Let $\Omega \subseteq \mathbf{R}$ be a measurable set, $\delta > 0$, $f \in C(\Omega)$ and $r \in \mathbb{N}$. Then,*

(i) *there holds*

$$1 + 2^{-r} r^r \leq \frac{\sup_{|h| \leq \delta} \left\| \left(I - \tilde{T}_h \right)^r f \right\|_{C(\Omega)}}{\|(I - T_\delta)^r f\|_{C(\Omega)}} \leq 2^r c_8(r),$$

(ii) *for $0 < \delta_1 \leq \delta_2$, there holds*

$$(1 + 2^{-r} r^r) \Omega_r(f, \delta_1)_{C(\Omega)} \leq c_8(r) 2^r \Omega_r(f, \delta_2)_{C(\Omega)}.$$

Remark 2.8. From Theorem 23.62 of [58, p.579] we have

$$(2.6) \quad \lim_{\delta \searrow 0} \Omega_1(f, \delta)_{C(\mathbf{R})} = \lim_{\delta \searrow 0} \|(I - T_\delta) f\|_{C(\mathbf{R})} = 0.$$

Corollary 2.9. *If $f \in C(\mathbf{R})$, $0 < \delta < \infty$, and $r \in \mathbf{N}$, then, by (2.6) and (***)*,

$$\lim_{\delta \searrow 0} \Omega_r(f, \delta)_{C(\mathbf{R})} = \lim_{\delta \searrow 0} \|(I - T_\delta)^r f\|_{C(\mathbf{R})} = 0$$

holds.

Let $\mathcal{G}_\sigma(X)$ be the subspace of entire function of exponential type σ that belonging to a Banach space X . The quantity

$$(2.7) \quad A_\sigma(f)_X := \inf_g \{\|f - g\|_X : g \in \mathcal{G}_\sigma(X)\}$$

is called the deviation of the function $f \in X$ from $\mathcal{G}_\sigma(X)$.

Let $\mathcal{G}_{\sigma, p(\cdot)} := \mathcal{G}_\sigma(L_{p(\cdot)})$ be the subspace of integral function f of exponential type σ that belonging to $L_{p(\cdot)}$. The quantity

$$A_\sigma(f)_{p(\cdot)} := \inf_g \{\|f - g\|_{p(\cdot)} : g \in \mathcal{G}_{\sigma, p(\cdot)}\}$$

is the deviation of the function $f \in L_{p(\cdot)}$ from \mathcal{G}_σ .

Remark 2.10. Let $\sigma > 0$, $1 \leq p \leq \infty$, $f \in L_p(\mathbf{R})$,

$$\vartheta(x) := \frac{2 \sin(x/2) \sin(3x/2)}{\pi x^2}$$

and

$$J(f, \sigma) = \sigma \int_{\mathbf{R}} f(x-u) \vartheta(\sigma u) du$$

be the de la Valèe Poussin operator ([13, definition given in (5.3)]). It is known (see (5.4)-(5.5) of [13]) that, if $f \in L_p(\mathbf{R})$, $1 \leq p \leq \infty$, then,

- (i) $J(f, \sigma) \in \mathcal{G}_{2\sigma}(L_p(\mathbf{R}))$,
- (ii) $J(g_\sigma, \sigma) = g_\sigma$ for any $g_\sigma \in \mathcal{G}_\sigma(L_p(\mathbf{R}))$,
- (iii) $\|J(f, \sigma)\|_{L_p(\mathbf{R})} \leq \frac{3}{2} \|f\|_{L_p(\mathbf{R})}$.
- (iv) $(J(f, \sigma))^{(r)} = J(f^{(r)}, \sigma)$ for any $r \in \mathbf{N}$ and $f \in (L_p(\mathbf{R}))^r$.
- (v) $\|J(f, \frac{\sigma}{2}) - f\|_{L_p(\mathbf{R})} \rightarrow 0$ (as $\sigma \rightarrow \infty$) and hence

$$\left\| \left(J\left(f, \frac{\sigma}{2}\right) \right)^{(k)} - f^{(k)} \right\|_{L_p(\mathbf{R})} \rightarrow 0 \text{ as } \sigma \rightarrow \infty,$$

for $f \in W_{L_p(\mathbf{R})}^r$ and $1 \leq k \leq r$.

Corollary 2.11. *Let $0 < \sigma < \infty$.*

- (i) *If $1 \leq p \leq \infty$, $f \in L_p(\mathbf{R})$. Then, using (v) of the last remark, we conclude*

$$\lim_{\sigma \rightarrow \infty} A_\sigma(f)_{L_p(\mathbf{R})} = 0.$$

- (ii) *Let $g : \mathbf{R} \rightarrow \mathbb{C}$ be bounded on the real axis \mathbf{R} . Then (see [14])*

$$\lim_{\sigma \rightarrow \infty} A_\sigma(g)_{C(\mathbf{R})} = 0$$

if and only if g is uniformly continuous on \mathbf{R} .

Theorem 2.12. *Let $r \in \mathbb{N}$, $\sigma > 0$, $\delta \in (0, 1)$ and $f \in \mathcal{C}(\mathbf{R})$. Then, the following Jackson type inequality*

$$(2.8) \quad A_\sigma(f)_{\mathcal{C}(\mathbf{R})} \leq 5\pi 4^{r-1} c_8(r) \Omega_r(f, 1/\sigma)_{\mathcal{C}(\mathbf{R})}, \text{ and}$$

its weak inverse

$$(2.9) \quad \Omega_r(f, \delta)_{\mathcal{C}(\mathbf{R})} \leq (1 + 2^{2r-1}) 2^{r-1} \delta^r \left(A_0(f)_{\mathcal{C}(\mathbf{R})} + \int_{1/2}^{1/\delta} u^{r-1} A_u(f)_{\mathcal{C}(\mathbf{R})} du \right)$$

are hold.

We set $[\sigma] := \max\{n \in \mathbb{Z} : n \leq \sigma\}$.

Theorem 2.13. *Let $r \in \mathbb{N}$, $f \in X_{\mathcal{C}(\mathbf{R})}^r$ and $\sigma > 0$. Then,*

(a) (i) *there exists (see [13, Proposition 25]) a $g_\sigma \in \mathcal{G}_\sigma(\mathcal{C}(\mathbf{R}))$ such that*

$$A_\sigma(f)_{\mathcal{C}(\mathbf{R})} \leq \|f - g_\sigma\|_{\mathcal{C}(\mathbf{R})} \leq \frac{5\pi}{4} \frac{4^r}{\sigma^r} \|f^{(r)}\|_{\mathcal{C}(\mathbf{R})}, \text{ and}$$

(ii) *its weak inverse*

$$\|f^{(k)}\|_{\mathcal{C}(\mathbf{R})} \leq (1 + 2^{2k-1}) 2^{k+2} \pi^k c_8(k) \sum_{\nu=0}^{\infty} \frac{(\nu+1)^r}{\nu+1} A_\nu(f)_{\mathcal{C}(\mathbf{R})},$$

holds whenever $k = 1, 2, \dots, r$ and $\sum_{\nu=0}^{\infty} (\nu+1)^{r-1} A_\nu(f)_{\mathcal{C}(\mathbf{R})} < \infty$.

(b) (i) *the following inequality (see [30, p.397])*

$$A_\sigma(f)_{\mathcal{C}(\mathbf{R})} \leq \frac{(5\pi)^r}{\sigma^r} A_\sigma(f^{(r)})_{\mathcal{C}(\mathbf{R})}, \text{ and}$$

(ii) *its weak inverse*

$$\begin{aligned} A_\sigma(f^{(r)})_{\mathcal{C}(\mathbf{R})} &\leq \left\| f^{(r)} - \left(J \left(f^{(r)}, \frac{\sigma}{2} \right) \right) \right\|_{\mathcal{C}(\mathbf{R})} \leq \\ &\leq (1 + 2^{2r-1}) 2^{r+2} \pi^r c_8(r) \left(A_\sigma(f)_{\mathcal{C}(\mathbf{R})} \sum_{k=0}^{[\sigma]} \frac{k^r}{k} + \sum_{\nu=[\sigma]+1}^{\infty} \frac{(\nu+1)^r}{\nu+1} A_\nu(f)_{\mathcal{C}(\mathbf{R})} \right) \end{aligned}$$

hold when $\sum_{\nu=0}^{\infty} (\nu+1)^{r-1} A_\nu(f)_{\mathcal{C}(\mathbf{R})} < \infty$.

Theorem 2.14. *Let $r, k \in \mathbb{N}$, $0 < t \leq 1/2$, $0 \leq \delta < \infty$ and $f \in \mathcal{C}(\mathbf{R})$. Then*

(i) *there holds*

$$\Omega_{r+k}(f, \delta)_{\mathcal{C}(\mathbf{R})} \leq 2^k \Omega_r(f, \delta)_{\mathcal{C}(\mathbf{R})}, \text{ and}$$

(ii) *its weak inverse (Marchaud inequality)*

$$\Omega_r(f, t)_{\mathcal{C}(\mathbf{R})} \leq C_9(r, k) t^r \int_t^1 \frac{\Omega_{r+k}(f, u)_{\mathcal{C}(\mathbf{R})}}{u^{r+1}} du$$

with $C_9(r, k) = 10\pi (1 + 2^{2r-1}) 2^{2r+3k} c_8(r+k)$.

Theorem 2.15. *Let $\sigma > 0$ and $f \in C(\mathbf{R})$. If $\sum_{\nu=0}^{\infty} (\nu+1)^{k-1} A_{\nu}(f)_{C(\mathbf{R})} < \infty$, holds for some $k \in \mathbf{N}$, then,*

(i) *the following Jackson type inequality for derivatives*

$$A_{\sigma}(f)_{C(\mathbf{R})} \leq (5\pi)^{k+1} c_8(r) \sigma^{-k} \Omega_r(f^{(k)}, \sigma^{-1})_{C(\mathbf{R})}, \quad \text{and}$$

(ii) *its weak inverse (see Theorem 6.3.4 of [30, p.343])*

$$\Omega_r\left(f^{(k)}, \frac{1}{\sigma}\right)_{C(\mathbf{R})} \leq 2^{2k+r+1} \left(\frac{1}{\sigma^r} \sum_{\nu=0}^{[\sigma]} \frac{(\nu+1)^{r+k}}{\nu+1} A_{\nu}(f)_{C(\mathbf{R})} + \sum_{\nu=[\sigma]+1}^{\infty} \frac{\nu^k}{\nu} A_{\nu}(f)_{C(\mathbf{R})} \right)$$

are hold.

2.1. Proofs of the results of section 2.

Proof of Lemma 2.1. For $\delta = 0$ (2.2) is obvious. For $0 < \delta < \infty$ and $r = 1$, one can find

$$\begin{aligned} (2.10) \quad \frac{d}{dx} T_{\delta} f(x) &= \frac{d}{dx} \left(\frac{1}{\delta} \int_0^{\delta} f(x+t) dt \right) = \frac{d}{dx} \left(\frac{1}{\delta} \int_x^{x+\delta} f(\tau) d\tau \right) \\ &= \frac{1}{\delta} \int_x^{x+\delta} \frac{d}{dx} f(\tau) d\tau = T_{\delta} \frac{d}{dx} f(x). \end{aligned}$$

For $r > 1$, (2.2) follows from (2.10). □

Proof of Theorem 2.2. (1)-(3) is known. (4) is seen from binomial expansion. To prove (5) it is sufficient to note inequality (see [10])

$$\|(I - T_{\delta}) f\|_{C(\Omega)} \leq 2^{-1} \delta \|f'\|_{C(\Omega)}, \quad \delta > 0$$

for $f \in C^1(\Omega)$. Then

$$\|(I - T_{\delta})^r f\|_{C(\Omega)} \leq 2^{-1} \delta \|(I - T_{\delta})^{r-1} f'\|_{C(\Omega)} \leq \dots \leq 2^{-r} \delta^r \|f^{(r)}\|_{C(\Omega)}$$

for $f \in C^r(\Omega)$, because

$$[(I - T_{\delta})^r f]' = (I - T_{\delta})^r f'. \quad \square$$

Proof of Lemma 2.3. For $r = 2$, by Lemma 2.1,

$$\begin{aligned} \frac{d^2}{dx^2} T_{\delta}^2 f &= \frac{d}{dx} \frac{d}{dx} T_{\delta} T_{\delta} f = \frac{d}{dx} \frac{d}{dx} T_{\delta} \Psi \quad [\Psi := T_{\delta} f] \\ &= \frac{d}{dx} T_{\delta} \frac{d}{dx} \Psi = \frac{d}{dx} T_{\delta} \frac{d}{dx} T_{\delta} f \end{aligned}$$

and the result (2.3) follows. For $r = 3$, by Lemma 2.1,

$$\begin{aligned} \frac{d^3}{dx^3} T_{\delta}^3 f &= \frac{d}{dx} \frac{d^2}{dx^2} T_{\delta}^2 R_{\nu} f = \frac{d}{dx} \frac{d^2}{dx^2} T_{\delta}^2 \Psi = \frac{d}{dx} \frac{d}{dx} T_{\delta} \frac{d}{dx} T_{\delta} \Psi \\ &= \frac{d}{dx} \frac{d}{dx} T_{\delta} \frac{d}{dx} T_{\delta}^2 f = \frac{d}{dx} T_{\delta} \frac{d}{dx} \frac{d}{dx} T_{\delta}^2 f = \frac{d}{dx} T_{\delta} \frac{d^2}{dx^2} T_{\delta}^2 f \end{aligned}$$

and (2.3) holds. Let (2.3) holds for $k \in \mathbb{N}$:

$$(2.11) \quad \frac{d^k}{dx^k} T_\delta^k f = \frac{d}{dx} T_\delta \frac{d^{k-1}}{dx^{k-1}} T_\delta^{k-1} f.$$

Then, for $k + 1$, (2.11) and Lemma 2.1 implies that

$$\begin{aligned} \frac{d^{k+1}}{dx^{k+1}} T_\delta^{k+1} f &= \frac{d}{dx} \frac{d^k}{dx^k} T_\delta^k T_\delta f = \frac{d}{dx} \frac{d^k}{dx^k} T_\delta^k \Psi = \frac{d}{dx} \frac{d}{dx} T_\delta \frac{d^{k-1}}{dx^{k-1}} T_\delta^{k-1} \Psi \\ &= \frac{d}{dx} \frac{d}{dx} T_\delta \frac{d^{k-1}}{dx^{k-1}} T_\delta^k f = \frac{d}{dx} T_\delta \frac{d}{dx} \frac{d^{k-1}}{dx^{k-1}} T_\delta^k f = \frac{d}{dx} T_\delta \frac{d^k}{dx^k} T_\delta^k f. \end{aligned}$$

□

Proof of Theorem 2.5. For $f \in C(\Omega)$ we have

$$(2.12) \quad \begin{aligned} \left\| \frac{d}{dx} T_\delta f(x) \right\|_{C(\Omega)} &= \left\| \frac{d}{dx} \frac{1}{\delta} \int_0^\delta f(x+t) dt \right\|_{C(\Omega)} = \\ \left\| \frac{1}{\delta} \frac{d}{dx} \int_x^{x+\delta} f(\tau) d\tau \right\|_{C(\Omega)} &= \left\| \frac{1}{\delta} (f(x+\delta) - f(x)) \right\|_{C(\Omega)} \leq \frac{2}{\delta} \|f\|_{C(\Omega)}. \end{aligned}$$

Inequality (2.12) also implies

$$\left\| \left(\frac{d}{dx} \right)^2 T_\delta f(x) \right\|_{C(\Omega)} \leq \frac{2}{\delta} \left\| \frac{d}{dx} T_\delta f \right\|_{C(\Omega)}$$

for $f \in C(\Omega)$. If $f \in C^2(\Omega)$ one can get

$$(2.13) \quad \left\| f(x) - T_\delta f(x) + \frac{\delta}{2} \frac{d}{dx} f(x) \right\|_{C(\Omega)} \leq \frac{\delta^2}{6} \left\| \frac{d^2}{dx^2} f \right\|_{C(\Omega)}.$$

To obtain (2.13) we will use the Taylor formula

$$f(x+t) = f(x) + t \frac{d}{dx} f(x) + \frac{t^2}{2} \frac{d^2}{dx^2} f(\xi)$$

for some $\xi \in [x, x+t]$. Then integrating the last equation with respect to t

$$\begin{aligned} \frac{1}{\delta} \int_0^\delta f(x+t) dt &= f(x) + \frac{1}{\delta} \int_0^\delta t dt \frac{d}{dx} f(x) + \frac{1}{2} \frac{1}{\delta} \int_0^\delta t^2 dt \frac{d^2}{dx^2} f(\xi), \\ T_\delta f(x) &= f(x) + \frac{\delta}{2} \frac{d}{dx} f(x) + \frac{\delta^2}{6} \frac{d^2}{dx^2} f(\xi) \end{aligned}$$

and (2.13) holds.

Now (2.12) and (2.13) imply that

$$(2.14) \quad (1/36) K_1(f, \delta, C(\Omega))_{C(\Omega)} \leq \|(\mathbb{I} - T_\delta) f\|_{C(\Omega)} \leq 2K_1(f, \delta, C(\Omega))_{C(\Omega)}.$$

Firstly, let us prove the right hand side of (2.14). For any $g \in C^1(\Omega)$

$$\begin{aligned} \|f - T_\delta f\|_{C(\Omega)} &\leq \|f - g\|_{C(\Omega)} + \|g - T_\delta g\|_{C(\Omega)} + \|T_\delta(g - f)\|_{C(\Omega)} \\ &\leq 2\|f - g\|_{C(\Omega)} + \frac{\delta}{2} \|g'\|_{C(\Omega)} \leq 2K_1(f, \delta, C(\Omega))_{C(\Omega)}. \end{aligned}$$

For the left hand side of inequality (2.14) we need inequalities

$$(2.15) \quad \|f - T_\delta^2 f\|_{C(\mathbf{R})} \leq 2 \|f - T_\delta f\|_{C(\mathbf{R})},$$

$$(2.16) \quad \delta \left\| \left(\frac{d}{dx} \right)^2 T_\delta^2 f \right\|_{C(\mathbf{R})} \leq 34 \|f - T_\delta f\|_{C(\mathbf{R})}.$$

First we prove (2.15). Then

$$\begin{aligned} \|f - T_\delta^2 f\|_{C(\Omega)} &\leq \|f - T_\delta f\|_{C(\Omega)} + \|T_\delta f - T_\delta T_\delta f\|_{C(\Omega)} \\ &\leq 2 \|f - T_\delta f\|_{C(\Omega)}. \end{aligned}$$

Now we consider inequality (2.16). In (2.13) we replace f by $T_\delta^2 f$ and obtain

$$\left\| T_\delta^2 f(x) - T_\delta T_\delta^2 f(x) + \frac{\delta}{2} \frac{d}{dx} T_\delta^2 f(x) \right\|_{C(\Omega)} \leq \frac{\delta^2}{6} \left\| \frac{d^2}{dx^2} T_\delta^2 f \right\|_{C(\Omega)}.$$

On the other hand, by (2.12),

$$\begin{aligned} \left\| \frac{d^2}{dx^2} T_\delta^2 f \right\|_{C(\Omega)} &\leq \frac{2}{\delta} \left\| \frac{d}{dx} T_\delta f \right\|_{C(\Omega)} \\ &\leq \frac{2}{\delta} \left\{ \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} + \left\| \frac{d}{dx} T_\delta (T_\delta f - f) \right\|_{C(\Omega)} \right\} \\ &\leq \frac{2}{\delta} \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} + \frac{4}{\delta^2} \|T_\delta f - f\|_{C(\Omega)}. \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\delta}{2} \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} &\leq \left\| T_\delta^2 f - T_\delta T_\delta^2 f - \frac{\delta}{2} \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} + \|T_\delta^2 f - T_\delta T_\delta^2 f\|_{C(\Omega)} \\ &\leq \frac{\delta^2}{6} \left\| \frac{d^2}{dx^2} T_\delta^2 f \right\|_{C(\Omega)} + \|T_\delta^2 f - T_\delta T_\delta^2 f\|_{C(\Omega)} \\ &\leq \frac{\delta^2}{6} \frac{2}{\delta} \left\{ \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} + \frac{2}{\delta} \|T_\delta f - f\|_{C(\Omega)} \right\} + \|T_\delta^2 f - f\|_{C(\Omega)} \\ &\quad + \|T_\delta (T_\delta^2 f - f)\|_{C(\Omega)} + \|T_\delta f - f\|_{C(\Omega)}. \end{aligned}$$

Then

$$\begin{aligned} \frac{\delta}{6} \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} &\leq \frac{17}{3} \|T_\delta f - f\|_{C(\Omega)}, \\ \delta \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} &\leq 34 \|T_\delta f - f\|_{C(\Omega)}. \end{aligned}$$

To finish proof of the left hand side of inequality (2.4) with $r = 1$, we proceed as

$$\begin{aligned} K_1(f, \delta, C(\Omega))_{C(\Omega)} &\leq \|f - T_\delta^2 f\|_{C(\Omega)} + \delta \left\| \frac{d}{dx} T_\delta^2 f \right\|_{C(\Omega)} \\ &\leq 36 \|T_\delta f - f\|_{C(\Omega)}. \end{aligned}$$

The proof of (2.4) with $r = 1$ now completed.

Let $r > 1$ be a natural number and we define

$$g(\cdot) = \sum_{l=1}^r (-1)^{l-1} \binom{r}{l} T_\delta^{2rl} f(\cdot).$$

Then,

$$\|f - g\|_{C(\Omega)} = \|(I - T_\delta^{2r})^r f\|_{C(\Omega)} \leq (2r)^r \|(I - T_\delta)^r f\|_{C(\Omega)}.$$

On the other hand,

$$\begin{aligned} \delta^r \left\| \frac{d^r}{dx^r} T_\delta^{2r} f \right\|_{C(\Omega)} &= \delta^{r-1} \delta \left\| \frac{d}{dx} T_\delta^2 \left(\frac{d^{r-1}}{dx^{r-1}} \right) T_\delta^{2r-2} f \right\|_{C(\Omega)} \\ &\leq 34 \delta^{r-1} \left\| (I - T_\delta) \frac{d^{r-1}}{dx^{r-1}} T_\delta^{2r-2} f \right\|_{C(\Omega)} \\ &\leq (34)^2 \delta^{r-2} \left\| (I - T_\delta)^2 \frac{d^{r-2}}{dx^{r-2}} T_\delta^{2r-4} f \right\|_{C(\Omega)} \\ &\leq \dots \leq (34)^r \|(I - T_\delta)^r f\|_{C(\Omega)}. \end{aligned}$$

Then

$$\begin{aligned} \delta^r \left\| \frac{d^r}{dx^r} T_\delta^{2rl} f \right\|_{C(\Omega)} &\leq (34)^r \left\| (I - T_\delta)^r T_\delta^{2r(l-1)} f \right\|_{C(\Omega)} \\ &= (34)^r \left\| T_\delta^{2r(l-1)} (I - T_\delta)^r f \right\|_{C(\Omega)} \leq (34)^r \|(I - T_\delta)^r f\|_{C(\Omega)}. \end{aligned}$$

Using the last inequality we find

$$\begin{aligned} \delta^r \left\| \frac{d^r}{dx^r} g \right\|_{C(\Omega)} &= \delta^r \left\| \frac{d^r}{dx^r} \sum_{l=1}^r (-1)^{l-1} \binom{r}{l} T_\delta^{2rl} f \right\|_{C(\Omega)} \\ &= \delta^r \left\| \sum_{l=1}^r (-1)^{l-1} \binom{r}{l} \frac{d^r}{dx^r} T_\delta^{2rl} f \right\|_{C(\Omega)} \\ &\leq \sum_{l=1}^r \left| \binom{r}{l} \right| \delta^r \left\| \frac{d^r}{dx^r} T_\delta^{2rl} f \right\|_{C(\Omega)} \leq 2^r (34)^r \|(I - T_\delta)^r f\|_{C(\Omega)} \end{aligned}$$

and

$$\begin{aligned} K_r(f, \delta, C(\Omega))_{C(\Omega)} &\leq \|f - g\|_{C(\Omega)} + \delta^r \left\| \frac{d^r}{dx^r} g \right\|_{C(\Omega)} \\ &\leq 2^r (r^r + (34)^r) \|(I - T_\delta)^r f\|_{C(\Omega)}. \end{aligned}$$

For the opposite direction of the last inequality, when $g \in W_{p(\cdot)}^r$,

$$(2.17) \quad \begin{aligned} \Omega_r(f, \delta)_{C(\Omega)} &\leq 2^r \|f - g\|_{C(\Omega)} + \Omega_r(g, \delta)_{C(\Omega)} \\ &\leq 2^r \|f - g\|_{C(\Omega)} + 2^{-r} \delta^r \|g^{(r)}\|_{C(\Omega)}, \end{aligned}$$

and taking infimum on $g \in W_{p(\cdot)}^r$ in (2.17) we get

$$\Omega_r(f, \delta)_{C(\Omega)} \leq 2^r K_r(f, \delta, C(\Omega))_{C(\Omega)}.$$

□

Proof of Proposition 2.6. Let $f \in C(\Omega)$. Then

$$\begin{aligned} \|(I - T_h) f\|_{C(\Omega)} &\leq 2K_1(f, h, C(\Omega))_{C(\Omega)} \\ &\leq 2K_1(f, \delta, C(\Omega))_{C(\Omega)} \leq 72 \|(I - T_\delta) f\|_{C(\Omega)}. \end{aligned}$$

□

Proof of Theorem 2.12. (i) We consider Jackson type inequality (2.8). For any $g \in X_{\mathcal{C}(\mathbf{R})}^r$ we have

$$\begin{aligned} A_\sigma(f)_{\mathcal{C}(\mathbf{R})} &\leq A_\sigma(f - g)_{\mathcal{C}(\mathbf{R})} + A_\sigma(g)_{\mathcal{C}(\mathbf{R})} \\ &\leq \|f - g\|_{\mathcal{C}(\mathbf{R})} + \frac{5\pi}{4} \frac{4^r}{\sigma^r} \left\| \frac{d^r}{dx^r} g \right\|_{\mathcal{C}(\mathbf{R})}. \end{aligned}$$

Taking infimum on $g \in X_{\mathcal{C}(\mathbf{R})}^r$ in the last inequality we have

$$A_\sigma(f)_{\mathcal{C}(\mathbf{R})} \leq \frac{5\pi 4^r}{4} K_r\left(f, \frac{1}{\sigma}, \mathcal{C}(\mathbf{R})\right)_{\mathcal{C}(\mathbf{R})} \leq \frac{5\pi}{4} c_8(r) 4^r \left\| \left(I - T_{\frac{1}{\sigma}}\right)^r f \right\|_{\mathcal{C}(\mathbf{R})}.$$

(ii) We give the proof of inverse estimate (2.9). Let $\sigma > 0$ and $g_\sigma \in \mathcal{G}_\sigma(\mathcal{C}(\mathbf{R}))$ be the best approximating IFFD of $f \in \mathcal{C}(\mathbf{R})$. Suppose that $r \in \mathbf{N}$, $0 < \delta < 1$. Then, there exists a $m \in \mathbf{N}$ such that $\lfloor 1/\delta \rfloor = 2^{m-1}$. Hence, $2^{m-1} \leq 1/\delta < 2^m$. Now we have

$$\begin{aligned} \Omega_r(f, \delta)_{\mathcal{C}(\mathbf{R})} &\leq \Omega_r(f - g_{2^m}, \delta)_{\mathcal{C}(\mathbf{R})} + \Omega_r(g_{2^m}, \delta)_{\mathcal{C}(\mathbf{R})} \\ &\leq 2^r A_{2^m}(f)_{\mathcal{C}(\mathbf{R})} + 2^{-r} \delta^r \left\| \frac{d^r}{dx^r} g_{2^m} \right\|_{\mathcal{C}(\mathbf{R})}. \end{aligned}$$

On the other hand

$$\begin{aligned} \left\| \frac{d^r}{dx^r} g_{2^m} \right\|_{\mathcal{C}(\mathbf{R})} &= \left\| \sum_{\gamma=1}^m \left(\frac{d^r}{dx^r} g_{2^\gamma} - \frac{d^r}{dx^r} g_{2^{\gamma-1}} \right) + \left(\frac{d^r}{dx^r} g_1 - \frac{d^r}{dx^r} g_0 \right) \right\|_{\mathcal{C}(\mathbf{R})} \\ &\leq \sum_{\gamma=1}^m 2^{\gamma r} \|g_{2^\gamma} - g_{2^{\gamma-1}}\|_{\mathcal{C}(\mathbf{R})} + \|g_1 - g_0\|_{\mathcal{C}(\mathbf{R})} \\ &\leq A_0(f)_{\mathcal{C}(\mathbf{R})} + A_1(f)_{\mathcal{C}(\mathbf{R})} + \sum_{\gamma=1}^m 2^{\gamma r} \left(A_{2^\gamma}(f)_{\mathcal{C}(\mathbf{R})} + A_{2^{\gamma-1}}(f)_{\mathcal{C}(\mathbf{R})} \right) \\ &\leq A_0(f)_{\mathcal{C}(\mathbf{R})} + 2^r A_1(f)_{\mathcal{C}(\mathbf{R})} + 2 \sum_{\gamma=1}^m 2^{\gamma r} A_{2^{\gamma-1}}(f)_{\mathcal{C}(\mathbf{R})} \end{aligned}$$

$$\leq 2 \left(A_0(f)_{C(\mathbf{R})} + \sum_{\gamma=1}^m 2^{\gamma r} A_{2^{\gamma-1}}(f)_{C(\mathbf{R})} \right).$$

Then

$$\frac{\delta^r}{2^r} \left\| \frac{d^r}{dx^r} g_{2^m} \right\|_{C(\mathbf{R})} \leq \frac{2}{2^r} \delta^r \left(A_0(f)_{C(\mathbf{R})} + \sum_{\gamma=1}^m 2^{\gamma r} A_{q^{\gamma-1}}(f)_{C(\mathbf{R})} \right).$$

Hence

$$\begin{aligned} \Omega_r(f, \delta)_{C(\mathbf{R})} &\leq \frac{2^{(m+1)r}}{2^{mr}} A_{2^m}(f)_{C(\mathbf{R})} + \frac{2}{2^r} \delta^r \left(A_0(f)_{C(\mathbf{R})} + \sum_{\gamma=1}^m 2^{\gamma r} A_{q^{\gamma-1}}(f)_{C(\mathbf{R})} \right) \\ &\leq (1 + 2^{2r-1}) 2^{1-r} 2^{2r} \delta^r \left(A_0(f)_{C(\mathbf{R})} + \sum_{\gamma=1}^m \int_{2^{\gamma-2}}^{2^{\gamma-1}} u^{r-1} A_u(f)_{C(\mathbf{R})} du \right) \\ &\leq (1 + 2^{2r-1}) 2^{r-1} \delta^r \left(A_0(f)_{C(\mathbf{R})} + \int_{1/2}^{2^{m-1}} u^{r-1} A_u(f)_{C(\mathbf{R})} du \right) \\ &\leq (1 + 2^{2r-1}) 2^{r-1} \delta^r \left(A_0(f)_{C(\mathbf{R})} + \int_{1/2}^{1/\delta} u^{r-1} A_u(f)_{C(\mathbf{R})} du \right). \end{aligned}$$

□

Proof of Theorem 2.13. Results a) (i) and b) (i) are known. Let us consider a) (ii). Suppose that $\sum_{\nu=0}^{\infty} \frac{(\nu+1)^r}{\nu+1} A_{\nu}(f)_{C(\mathbf{R})} < \infty$ and $k \in \{1, 2, \dots, r\}$. Then, using Nikolskii inequality, one gets

$$\begin{aligned} \|f^{(k)}\|_{C(\mathbf{R})} &= \lim_{\sigma \rightarrow \infty} \|J\left(f^{(k)}, \frac{\sigma}{2}\right)\|_{C(\mathbf{R})} = \lim_{\sigma \rightarrow \infty} \left\| \left(J\left(f, \frac{\sigma}{2}\right) \right)^{(k)} \right\|_{C(\mathbf{R})} \\ &\leq \frac{\pi^k \sup_{|h| \leq \delta} \left\| \left(I - \tilde{T}_h \right)^k \left(J\left(f, \frac{\sigma}{2}\right) \right) \right\|_{C(\mathbf{R})}}{\delta^k} \leq \frac{\pi^k 2^k c_8(k) \Omega_k\left(J\left(f, \frac{\sigma}{2}\right), \delta\right)_{C(\mathbf{R})}}{\delta^k} \\ &\leq (1 + 2^{2k-1}) 2^{k+2} \pi^k c_8(k) \sum_{\nu=0}^{\lfloor 1/\delta \rfloor} \frac{(\nu+1)^k}{\nu+1} A_{\nu}\left(J\left(f, \frac{\sigma}{2}\right)\right)_{C(\mathbf{R})} \\ &\leq (1 + 2^{2k-1}) 2^{k+2} \pi^k c_8(k) \sum_{\nu=0}^{\infty} \frac{(\nu+1)^r}{\nu+1} A_{\nu}(f)_{C(\mathbf{R})}. \end{aligned}$$

Note that (ii) b) is follow from (i) b). □

Proof of Theorem 2.14. (i) follows from properties of modulus of smoothness. We consider Marchaud type inequality (ii). Let $0 < t < 1/2$. Assume that $2^{m-1} \leq \frac{1}{t} < 2^m$ for some $m \in \mathbb{N}$. Then

$$\begin{aligned}
 \Omega_r(f, t)_{C(\mathbf{R})} &\leq (1 + 2^{2r-1}) 2^{1-r} t^r \left(\sum_{\nu=1}^m 2^{\nu r} A_{2^{\nu-1}}(f)_{C(\mathbf{R})} + A_0(f)_{C(\mathbf{R})} \right) \\
 &\leq \frac{5\pi}{2} (1 + 2^{2r-1}) 2^{r+2k} c_8(r+k) t^r \left(A_0(f)_{C(\mathbf{R})} + \sum_{\nu=1}^m 2^{\nu r} \Omega_{k+r}(f, \frac{1}{2^\nu})_{C(\mathbf{R})} \right) \\
 &\leq \frac{5\pi}{2} (1 + 2^{2r-1}) 2^{2r+3k} c_8(r+k) t^r \left(\Omega_{k+r}(f, \frac{1}{2})_{C(\mathbf{R})} + \sum_{\nu=1}^m \int_{2^{-\nu}}^{2^{-\nu+1}} \frac{\Omega_{k+r}(f, u)_{C(\mathbf{R})} du}{u^{r+1}} \right) \\
 &\leq \frac{5\pi}{2} (1 + 2^{2r-1}) 2^{2r+3k} c_8(r+k) t^r \left(\Omega_{k+r}(f, \frac{1}{2})_{C(\mathbf{R})} + \int_{2^{-1}}^{2^{-m+1}} \frac{\Omega_{k+r}(f, u)_{C(\mathbf{R})} du}{u^{r+1}} \right) \\
 &\leq 5\pi (1 + 2^{2r-1}) 2^{2r+3k} c_8(r+k) t^r \left(\int_{1/2}^1 \frac{\Omega_{k+r}(f, u)_{C(\mathbf{R})} du}{u^{r+1}} + \int_t^1 \frac{\Omega_{k+r}(f, u)_{C(\mathbf{R})} du}{u^{r+1}} \right) \\
 &\leq 10\pi (1 + 2^{2r-1}) 2^{2r+3k} c_8(r+k) t^k \int_t^1 \frac{\Omega_{k+r}(f, u)_{C(\mathbf{R})} du}{u^{r+1}}
 \end{aligned}$$

□

Using this section's estimates and Transference Result Theorem 1.10, in the next section we will give several results on difference operator $\|(I - T_\delta)^r f\|_{p(\cdot)}$ and approximation by IFFD in $L_{p(\cdot)}$.

3. APPLICATIONS ON DIFFERENCE OPERATOR AND APPROXIMATION

Notation . Since the $48c_7(c_3(p))c_5(p^+, c_3(p))$ of (1.12) will be used very frequently in the next parts we will set $c_{10} := c_{10}(p^+, c_3(p)) := 48c_7(c_3(p))c_5(p^+, c_3(p))$.

Lemma 3.1. *Let $p \in P^{Log}(\mathbf{R})$, $r \in \mathbf{N}$, and $0 < \delta < \infty$. Then*

$$\|(I - T_\delta)^r f\|_{p(\cdot)} \leq c_{10}^r 2^{-r} \delta^r \|f^{(r)}\|_{p(\cdot)}, \quad f \in W_{L_{p(\cdot)}}^r$$

hold.

We will use notation $K_r(f, \delta, p(\cdot)) := K_r(f, \delta, L_{p(\cdot)})_{L_{p(\cdot)}}$ for $r \in \mathbf{N}$, $p \in P^{Log}(B)$, $\delta > 0$ and $f \in L_{p(\cdot)}(B)$.

As a corollary of Transference Result we can obtain the following lemma.

Lemma 3.2. *Let $0 < h \leq \delta < \infty$, $p \in P^{Log}(\mathbf{R})$ and $f \in L_{p(\cdot)}$. Then*

$$(3.1) \quad \|(I - T_h) f\|_{p(\cdot)} \leq c_8(72, p^+, c_3(p)) \|(I - T_\delta) f\|_{p(\cdot)}$$

holds.

In the following theorem we show that K -functional $K_r(f, \delta, p(\cdot))_{p(\cdot)}$ and $\Omega_r(f, \delta)_{p(\cdot)}$ are equivalent.

Theorem 3.3. *Let $p(\cdot) \in P^{Log}(\mathbf{R})$. If $L_{p(\cdot)}$, then the K -functional $K_r(f, \delta, p(\cdot))_{p(\cdot)}$ and the modulus $\Omega_r(f, \delta)_{p(\cdot)}$, are equivalent, namely,*

$$\begin{aligned} \frac{1}{48c_7(c_3(p))2^r c_5(p^+, c_3(p))} &\leq \frac{K_r(f, \delta, p(\cdot))_{p(\cdot)}}{\Omega_r(f, \delta)_{p(\cdot)}} \\ &\leq 48c_7(c_3(p))\{(2r)^r + 2^r(34)^r\}c_5(p^+, c_3(p)). \end{aligned}$$

Theorem 3.4. *For $p(\cdot) \in P^{Log}(\mathbf{R})$, $f, g \in L_{p(\cdot)}$ and $\delta > 0$, the modulus of smoothness $\Omega_r(f, \delta)_{p(\cdot)}$, has the following properties:*

- (1) $\Omega_r(f, \delta)_{p(\cdot)}$ is non-negative; non-decreasing function of δ ;
- (2) For $f, g \in L_{p(\cdot)}$ and $\delta > 0$,

$$(3.2) \quad \Omega_r(f + g, \delta)_{p(\cdot)} \leq \Omega_r(f, \delta)_{p(\cdot)} + \Omega_r(g, \delta)_{p(\cdot)}.$$

- (3) For $f \in L_{p(\cdot)}$,

$$(3.3) \quad \lim_{\delta \rightarrow 0} \Omega_r(f, \delta)_{p(\cdot)} = 0.$$

As a corollary of Theorem 3.3

Corollary 3.5. *Let $p(\cdot) \in P^{Log}(\mathbf{R})$. If $\delta, \lambda \in (0, 1)$, $f \in L_{p(\cdot)}$, then,*

$$\frac{\Omega_r(f, \lambda\delta)_{p(\cdot)}}{(1 + [\lambda])^r \Omega_r(f, \delta)_{p(\cdot)}} \leq (48)^2 c_7^2(c_3(p)) 2^r c_5^2(p^+, c_3(p)) ((2r)^r + 2^r(34)^r)$$

holds.

Theorem 3.6. *Let $p(\cdot) \in P^{Log}(\mathbf{R})$, $r \in \mathbf{N}$, $\sigma > 0$ and $f \in L_{p(\cdot)}$. Then,*

$$(3.4) \quad A_\sigma(f)_{p(\cdot)} \leq c_{11} \|(I - T_{1/\sigma})^r f\|_{p(\cdot)}$$

with $c_{11} := c_{11}(r, p^+, c_3(p)) := 30\pi 8^r c_5(p^+, c_3(p)) c_7(c_3(p)) c_8(r)$.

Now we present the inverse theorem.

Theorem 3.7. *Let $p(\cdot) \in P^{Log}(\mathbf{R})$, $r \in \mathbf{N}$, $\delta \in (0, 1)$ and $f \in L_{p(\cdot)}$. Then,*

$$\Omega_r(f, \delta)_{p(\cdot)} \leq c_{12} \delta^r \left(A_0(f)_{p(\cdot)} + \int_{1/2}^{1/\delta} u^{r-1} A_{u/2}(f)_{p(\cdot)} du \right)$$

holds with $c_{12} := c_{12}(r, p^+, c_3(p)) := c_{13} 12c_7(c_3(p)) (1 + 2^{2r-1}) 2^r$ where $c_{13} := c_{13}(p^+, c_3(p)) := 2c_5(p^+, c_3(p)) (1 + 72c_7(c_3(p)) c_5(p^+, c_3(p)))$.

In this section we obtain Marchaud inequality.

Theorem 3.8. *Let $r, k \in \mathbf{N}$, $p \in P^{Log}(\mathbf{R})$, $f \in L_{p(\cdot)}$ and $t \in (0, 1/2)$. Then,*

$$\Omega_r(f, t)_{p(\cdot)} \leq c_{14} t^r \int_t^1 \frac{\Omega_{r+k}(f, u)_{p(\cdot)}}{u^{r+1}} du$$

holds with $c_{14} := c_{14}(r, k, p^+, c_3(p)) := 48c_7(c_3(p)) C_9(r, k) c_5(p^+, c_3(p))$.

Theorem 3.9. *Let $p \in P^{Log}(\mathbf{R})$, $r \in \mathbb{N}$ and $f \in L_{p(\cdot)}$. If*

$$\sum_{\nu=0}^{\infty} \nu^{k-1} A_{\nu/2}(f)_{p(\cdot)} < \infty$$

holds for some $k \in \mathbb{N}$, then $f^{(k)} \in L_{p(\cdot)}$ and

$$(3.5) \quad \Omega_r \left(f^{(k)}, \frac{1}{\sigma} \right)_{p(\cdot)} \leq c_{14} \left(\frac{1}{\sigma^r} \sum_{\nu=0}^{\lfloor \sigma \rfloor} (\nu+1)^{r+k-1} A_{\nu/2}(f)_{p(\cdot)} + \sum_{\nu=\lfloor \sigma \rfloor+1}^{\infty} \nu^{k-1} A_{\nu/2}(f)_{p(\cdot)} \right)$$

with $c_{14} := c_{14}(r, k, p^+, c_3(p)) := 48c_7(c_3(p)) c_5(p^+, c_3(p)) 2^{2k+r+2}$.

3.1. Proofs of the results of section 3.

Proof of Lemma 3.1. We note that (see [10]) the following inequality

$$(3.6) \quad \|(I - T_\delta) f\|_{p(\cdot)} \leq 2^{-1} c_{10} \delta \|f'\|_{p(\cdot)}, \quad \delta > 0$$

holds for $f \in L_{p(\cdot)}$. Then

$$\Omega_r(f, \delta)_{p(\cdot)} = \|(I - T_\delta)^r f\|_{p(\cdot)} \leq \dots \leq 2^{-r} c_{10}^r \delta^r \|f^{(r)}\|_{p(\cdot)}, \quad \delta > 0$$

for $f \in W_{L_{p(\cdot)}}^r$. □

Proof of Theorem 3.3. For any $g \in W_{L_{p(\cdot)}}^r(\Omega)$ we have $F_g \in C^r(\Omega)$. Since F_f is linear in f ,

$$(I - T_\delta)^r F_f = F_{(I - T_\delta)^r f} \quad \text{and} \quad (F_g)^{(r)} = F_{g^{(r)}}$$

using Theorem 1.10 we obtain

$$\begin{aligned} \|(I - T_\delta)^r f\|_{p(\cdot)} &\leq 24c_7(c_3(p)) \|F_{(I - T_\delta)^r f}\|_{C(\Omega)} \\ &= 24c_7(c_3(p)) \|(I - T_\delta)^r F_f\|_{C(\Omega)} \\ &\leq 24c_7(c_3(p)) 2^r K_r(F_f, \delta, C(\Omega))_{C(\Omega)} \\ &\leq 24c_7(c_3(p)) 2^r \left\{ \|F_f - F_g\|_{C(\Omega)} + \delta^r \|(F_g)^{(r)}\|_{C(\Omega)} \right\} \\ &= 24c_7(c_3(p)) 2^r \left\{ \|F_{(f-g)}\|_{C(\Omega)} + \delta^r \|F_{g^{(r)}}\|_{C(\Omega)} \right\} \\ &\leq 48c_7(c_3(p)) 2^r c_5(p^+, c_3(p)) \left\{ \|f - g\|_{p(\cdot)} + \delta^r \|g^{(r)}\|_{p(\cdot)} \right\}. \end{aligned}$$

Taking infimum and considering definition of K -functional one gets

$$\|(I - T_\delta)^r f\|_{p(\cdot)} \leq 48c_7(c_3(p)) 2^r c_5(p^+, c_3(p)) K_r(f, \delta, p(\cdot))_{p(\cdot)}.$$

Now we consider the opposite direction of the last inequality. For

$$g(\cdot) = \sum_{l=1}^r (-1)^{l-1} \binom{r}{l} T_\delta^{2rl} f(\cdot)$$

we have

$$\begin{aligned}
K_r(f, \delta, p(\cdot))_{p(\cdot)} &\leq \|f - g\|_{p(\cdot)} + \delta^r \left\| \frac{d^r}{dx^r} g \right\|_{p(\cdot)} \\
&\leq 24c_7(c_3(p)) \left\{ \|F_{(f-g)}\|_{C(\Omega)} + \delta^r \|F_{g^{(r)}}\|_{C(\Omega)} \right\} \\
&= 24c_7(c_3(p)) \left\{ \|F_f - F_g\|_{C(\Omega)} + \delta^r \|(F_g)^{(r)}\|_{C(\Omega)} \right\} \\
&\leq 24c_7(c_3(p)) \left\{ \|(I - T_\delta^{2r})^r F_f\|_{C(\Omega)} + \delta^r \left\| \left(\sum_{l=1}^r (-1)^{l-1} \binom{r}{l} T_\delta^{2rl} F_f \right)^{(r)} \right\|_{C(\Omega)} \right\} \\
&= 24c_7(c_3(p)) \left\{ \|(I - T_\delta^{2r})^r F_f\|_{C(\Omega)} + \sum_{l=1}^r \left| \binom{r}{l} \right| \delta^r \|(T_\delta^{2rl} F_f)^{(r)}\|_{C(\Omega)} \right\} \\
&\leq 24c_7(c_3(p)) \left\{ (2r)^r \|(I - T_\delta)^r F_f\|_{C(\Omega)} + 2^r (34)^r \|(I - T_\delta)^r F_f\|_{C(\Omega)} \right\} \\
&= 24c_7(c_3(p)) \{(2r)^r + 2^r (34)^r\} \|F_{(I-T_\delta)^r f}\|_{C(\Omega)} \\
&\leq 48c_7(c_3(p)) \{(2r)^r + 2^r (34)^r\} c_5(p^+, c_3(p)) \|(I - T_\delta)^r f\|_{p(\cdot)}.
\end{aligned}$$

□

Proof of Theorem 3.4. Properties (1) and (2), by definition of $\Omega_r(f, \delta)_{p(\cdot)}$ and the triangle inequality of $L_{p(\cdot)}$ are clearly valid. By using [22, Theorem 10.1] and [36, Lemma 2], the relation (3.3) is satisfied. □

Proof of Corollary 3.5. We have

$$\begin{aligned}
\frac{\Omega_r(f, \lambda\delta)_{p(\cdot)}}{(1 + \lfloor \lambda \rfloor)^r \Omega_r(f, \delta)_{p(\cdot)}} &\leq \frac{48c_7(c_3(p)) 2^r c_5(p^+, c_3(p)) K_r(f, \lambda\delta, p(\cdot))_{p(\cdot)}}{(1 + \lfloor \lambda \rfloor)^r \Omega_r(f, \delta)_{p(\cdot)}} \\
&\leq \frac{(48)^2 c_7^2(c_3(p)) 2^r c_5^2(p^+, c_3(p)) (1 + \lfloor \lambda \rfloor)^r}{(1 + \lfloor \lambda \rfloor)^r} \frac{\{(2r)^r + 2^r (34)^r\}}{1} \\
&= (48)^2 c_7^2(c_3(p)) 2^r c_5^2(p^+, c_3(p)) \{(2r)^r + 2^r (34)^r\}.
\end{aligned}$$

□

Proof of Theorem 3.6. First we obtain

$$(3.7) \quad A_{2\sigma}(f)_{p(\cdot)} \leq 30\pi 8^r c_5(p^+, c_3(p)) c_7(c_3(p)) c_8(r) \|(I - T_{1/(2\sigma)})^r f\|_{p(\cdot)}$$

and (3.4) follows from (3.7). Let g_σ be an exponential type entire function of degree $\leq \sigma$, belonging to $\mathcal{C}(\mathbf{R})$, as best approximation of $F_f \in \mathcal{C}(\mathbf{R})$. Since $F_{V_\sigma f} = V_\sigma F_f$ and $V_\sigma g_\sigma = g_\sigma$, there holds

$$\begin{aligned}
A_{2\sigma}(f)_{p(\cdot)} &\leq \|f - V_\sigma f\|_{p(\cdot)} \leq 24c_7(c_3(p)) \|F_{f-V_\sigma f}\|_{C(\mathbf{R})} \\
&= 24c_7(c_3(p)) \|F_f - V_\sigma F_f\|_{C(\mathbf{R})} \\
&= 24c_7(c_3(p)) \|F_f - g_\sigma + g_\sigma - V_\sigma F_f\|_{C(\mathbf{R})}
\end{aligned}$$

$$\begin{aligned}
&= 24c_7(c_3(p)) \|F_f - g_\sigma + V_\sigma g_\sigma - V_\sigma F_f\|_{\mathcal{C}(\mathbf{R})} \\
&\leq 24c_7(c_3(p)) (A_\sigma(F_f)_{\mathcal{C}(\mathbf{R})} + \frac{3}{2}A_\sigma(F_f)_{\mathcal{C}(\mathbf{R})}) = 12c_7(c_3(p)) A_\sigma(F_f)_{\mathcal{C}(\mathbf{R})}.
\end{aligned}$$

For any $g \in W_{\mathcal{C}(\mathbf{R})}^r$

$$\begin{aligned}
A_\sigma(u)_{\mathcal{C}(\mathbf{R})} &\leq A_\sigma(u - g)_{\mathcal{C}(\mathbf{R})} + A_\sigma(g)_{\mathcal{C}(\mathbf{R})} \\
&\leq \|u - g\|_{\mathcal{C}(\mathbf{R})} + \frac{5\pi 4^r}{4 \sigma^r} \left\| \frac{d^r}{dx^r} g \right\|_{\mathcal{C}(\mathbf{R})} \\
&\leq \frac{5\pi 4^r}{4} K_r \left(u, \frac{1}{\sigma}, \mathcal{C}(\mathbf{R}) \right)_{\mathcal{C}(\mathbf{R})} \leq \frac{5\pi 8^r}{4} K_r \left(u, \frac{1}{2\sigma}, \mathcal{C}(\mathbf{R}) \right)_{\mathcal{C}(\mathbf{R})} \\
&\leq \frac{5\pi 8^r}{4} c_8(r) \left\| \left(I - T_{\frac{1}{2\sigma}} \right)^r u \right\|_{\mathcal{C}(\mathbf{R})}.
\end{aligned}$$

Therefore

$$\begin{aligned}
A_{2\sigma}(f)_{p(\cdot)} &\leq 12c_7(c_3(p)) A_\sigma(F_f)_{\mathcal{C}(\mathbf{R})} \\
&\leq 15\pi 8^r c_7(c_3(p)) c_8(r) \left\| \left(I - T_{\frac{1}{2\sigma}} \right)^r F_f \right\|_{\mathcal{C}(\mathbf{R})} \\
&= 15\pi 8^r c_7(c_3(p)) c_8(r) \left\| F_{(I - T_{1/(2\sigma)})^r} f \right\|_{\mathcal{C}(\mathbf{R})} \\
&\leq 30\pi 8^r c_5(p^+, c_3(p)) c_7(c_3(p)) c_8(r) \left\| (I - T_{1/(2\sigma)})^r f \right\|_{p(\cdot)}.
\end{aligned}$$

□

Proof of Theorem 3.7. Let g_σ be an exponential type entire function of degree $\leq \sigma$, belonging to $L^{p(\cdot)}$, as best approximation of $f \in L^{p(\cdot)}$. Then

$$\begin{aligned}
\Omega_r(f, \delta)_{p(\cdot)} &= \|(I - T_\delta)^r f\|_{p(\cdot)} \leq 24c_7(c_3(p)) \|F_{(I - T_\delta)^r} f\|_{\mathcal{C}(\mathbf{R})} \\
&= 24c_7(c_3(p)) \|(I - T_\delta)^r F_f\|_{\mathcal{C}(\mathbf{R})} \\
&\leq 12c_7(c_3(p)) (1 + 2^{2r-1}) 2^r \delta^r \left(A_0(F_f)_{\mathcal{C}(\mathbf{R})} + \int_{1/2}^{1/\delta} u^{r-1} A_u(F_f)_{\mathcal{C}(\mathbf{R})} du \right) \\
&\leq c_{13} 12c_7(c_3(p)) (1 + 2^{2r-1}) 2^r \delta^r \left(A_0(f)_{p(\cdot)} + \int_{1/2}^{1/\delta} u^{r-1} A_{u/2}(f)_{p(\cdot)} du \right)
\end{aligned}$$

because

$$\begin{aligned}
A_{2\sigma}(F_f)_{\mathcal{C}(\mathbf{R})} &\leq \|F_f - V_\sigma F_f\|_{\mathcal{C}(\mathbf{R})} = \|F_f - V_\sigma f\|_{\mathcal{C}(\mathbf{R})} \leq 2c_5(p^+, c_3(p)) \|f - V_\sigma f\|_{p(\cdot)} \\
&= 2c_5(p^+, c_3(p)) \|f - g_\sigma + g_\sigma - V_\sigma f\|_{p(\cdot)} \\
&\leq 2c_5(p^+, c_3(p)) \left(\|f - g_\sigma\|_{p(\cdot)} + \|V_\sigma g_\sigma - V_\sigma f\|_{p(\cdot)} \right) \\
&\leq 2c_5(p^+, c_3(p)) \left(\|f - g_\sigma\|_{p(\cdot)} + 72c_7(c_3(p)) c_5(p^+, c_3(p)) \|g_\sigma - f\|_{p(\cdot)} \right) \\
&= 2c_5(p^+, c_3(p)) (1 + 72c_7(c_3(p)) c_5(p^+, c_3(p))) A_\sigma(f)_{p(\cdot)}.
\end{aligned}$$

□

Proof of Theorem 3.8. Let g_σ be an exponential type entire function of degree $\leq \sigma$, belonging to $L^{p(\cdot)}$, as best approximation of $f \in L_{p(\cdot)}$. Then

$$\begin{aligned} \Omega_r(f, t)_{p(\cdot)} &= \|(I - T_t)^r f\|_{p(\cdot)} \leq 24c_7(c_3(p)) \|F_{(I-T_t)^r f}\|_{\mathcal{C}(\mathbf{R})} \\ &= 24c_7(c_3(p)) \|(I - T_t)^r F_f\|_{\mathcal{C}(\mathbf{R})} \\ &\leq 24c_7(c_3(p)) C_9(r, k) t^r \int_t^1 \frac{\|(I - T_t)^{r+k} F_f\|_{\mathcal{C}(\mathbf{R})}}{u^{r+1}} du \\ &= 24c_7(c_3(p)) C_9(r, k) t^r \int_t^1 \frac{\|F_{(I-T_t)^{r+k} f}\|_{\mathcal{C}(\mathbf{R})}}{u^{r+1}} du \\ &\leq 48c_7(c_3(p)) C_9(r, k) c_5(p^+, c_3(p)) t^r \int_t^1 \frac{\|(I - T_t)^{r+k} f\|_{p(\cdot)}}{u^{r+1}} du \\ &= 48c_7(c_3(p)) C_9(r, k) c_5(p^+, c_3(p)) t^r \int_t^1 \frac{\Omega_{r+k}(f, t)_{p(\cdot)}}{u^{r+1}} du. \end{aligned}$$

□

Proof of Theorem 3.9. Proof of (3.5) is similar to that of proof of Theorem 3.8. □

REFERENCES

- [1] F. Abdullaev, S. Chaichenko, M. Imashgizi, and A. Shidlich, Direct and inverse approximation theorems in the weighted Orlicz-type spaces with a variable exponent, Turk. J. Math., 44 (2020), 284-299.
- [2] F. Abdullaev, A. Shidlich and S. Chaichenko, Direct and inverse approximation theorems of functions in the Orlicz type spaces, Math. Slovaca, 69 (2019), 1367-1380.
- [3] F. Abdullaev, N. Özkaratepe, V. Savchuk and A. Shidlich, Exact constants in direct and inverse approximation theorems for functions of several variables in the spaces S_p , FILOMAT, 33 (2019), 1471-1484.
- [4] N. I. Akhiezer, Theory of approximation, Fizmatlit, Moscow, 1965; English transl. of 2nd ed. Frederick Ungar, New York, 1956.
- [5] R. Akgün, Approximation of Functions of Weighted Lebesgue and Smirnov Spaces, Mathematica (Cluj), Tome 54 (77), No: Special (2012), pp. 25-36.
- [6] R. Akgün Sharp Jackson and converse theorems of trigonometric approximation in weighted Lebesgue spaces, Proc. A. Razmadze Math. Inst., 152 (2010), pp. 1-18.
- [7] R. Akgün, Inequalities for one sided approximation in Orlicz spaces, Hacet. J. Math. Stat., Volume: 40 Issue: 2 (2011), pp. 231-240.
- [8] R. Akgün, Some convolution inequalities in Musielak Orlicz spaces, Proc. Inst. Math. Mech., NAS Azerbaijan, 42 (2016), No: 2, 279-291.
- [9] R. Akgün, Approximation properties of Bernstein's singular integrals in variable exponent Lebesgue spaces on the real axis, Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, In press, arXiv:1210.4714v3 [math.CA], <https://doi.org/10.48550/arXiv.1210.4714>
- [10] R. Akgün; A. Ghorbanalizadeh, "Approximation by integral functions of finite degree in variable exponent Lebesgue spaces on the real axis." Turkish Journal of Mathematics 42, (2018), no. 4, 1887-1903.
- [11] A.H. Avşar and H. Koç, Jackson and Stechkin type inequalities of trigonometric approximation in $A_{-p, q(\cdot)}^w, \theta$, Turk J Math (2018) 42: 2979-2993.

- [12] A.H. Avşar and Y.E. Yildirim, On the trigonometric approximation of functions in weighted Lorentz spaces using Cesaro submethod, *Novi Sad J. Math.* Vol. 48, No. 2, 2018, 41-54.
- [13] C. Bardaro, P.L. Butzer, R.L. Stens and G. Vinti, Approximation error of the Whittaker cardinal series in terms of an averaged modulus of smoothness covering discontinuous signals, *J. Math. Anal. Appl.* 316 (2006) 269-306.
- [14] S.N. Bernstein, Sur la meilleure approximation sur tout l'axe reel des fonctions continues par des fonctions entieres de degre n. I, *C.R. (Doklady) Acad. Sci. URSS (N.S.)* 51 (1946), 331-334.
- [15] S. N. Bernstein, *Collected works, M. Vol. I*, Izdat. Akad. Nauk SSSR, Moscow, 1952., 11-104.
- [16] D. Cruz-Uribe, A. Fiorenza, *Variable Lebesgue Spaces, Foundations and Harmonic Analysis*, Birkhauser, Applied and Numerical Harmonic Analysis, 2013.
- [17] D. Cruz-Uribe, A. Fiorenza, Approximate identities in variable L_p spaces, *Mathematische Nachrichten*, 280 (2007), No:3, 256-270.
- [18] R. A. DeVore, G. G. Lorentz, *Constructive Approximation*, Springer-Verlag, (1993).
- [19] L. Diening, P. Harjulehto, P. Hasto and M. Ružička, *Lebesgue and Sobolev spaces with variable exponents*, *Lecture Notes in Math.*, vol. 2017, Springer, Berlin, Heidelberg, 2011.
- [20] L. Diening and M. Ružička, Calderon–Zygmund operators on generalized Lebesgue spaces $L^{p(x)}$ and problems related to fluid dynamics, preprint, *Mathematische Fakultät, Albert-Ludwigs-Universität Freiburg*, 21/2002, 04.07.2002, 1-20, 2002.
- [21] Z. Ditzian, Inverse theorems for functions in L^p and other spaces, *Proc. Amer. Math. Soc.* 54 (1976), 80-82.
- [22] Z. Ditzian and K. G. Ivanov, Strong converse inequalities, *Journal D'analyse mathematique* **61**:1, 1993, 61-111.
- [23] A. Dogu, A.H. Avsar and Y.E. Yildirim, Some inequalities about convolution and trigonometric approximation in weighted Orlicz spaces, *Proceedings of the Institute of Mathematics and Mechanics, National Academy of Sciences of Azerbaijan*, Volume 44, Number 1, 2018, Pages 107-115.
- [24] D. Drihem, Restricted boundedness of translation operators on variable Lebesgue spaces, <https://doi.org/10.48550/arXiv.1507.08089>
- [25] D. P. Dryanov, M. A. Qazi, and Q. I. Rahman, Entire functions of exponential type in Approximation Theory, In: *Constructive Theory of Functions*, Varna 2002 (B. Bojanov, Ed.), DARBA, Sofia, 2003, pp. 86-135.
- [26] X. Fan and D. Zhao, On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$, *J. Math. Anal. Appl.* **263**:2 (2001), 424–446.
- [27] A. Guven and D.M. Israfilov, Trigonometric approximation in generalized Lebesgue spaces $L^{p(x)}$, *J. Math. Inequal.* 4 (2010), no. 2, 285–299.
- [28] P. Harjulehto and P. Hästö, *Orlicz Spaces and Generalized Orlicz Spaces*, *Lecture Notes in Mathematics*, vol. 2236, Springer, 2019, X+168.
- [29] H. Hudzik, On generalized Orlicz–Sobolev space, *Funct. Approximatio Comment. Math.* **4** (1976), 37–51.
- [30] I.I. Ibragimov, *Teoriya priblizheniya tselymi funktsiyami. (Russian) [The theory of approximation by entire functions] "Elm"*, Baku, 1979. 468 pp.
- [31] S.Z. Jafarov, Linear Methods for Summing Fourier Series and Approximation in Weighted Lebesgue Spaces with Variable Exponents. *Ukrainian Mathematical Journal.* 2015; 66(10): 1509–1518.
- [32] S.Z. Jafarov, Approximation by trigonometric polynomials in subspace of variable exponent grand Lebesgue spaces. *Global Journal of Mathematics*, 2016; 8(2): 836–843.
- [33] S. Z. Jafarov, Ul'yanov type inequalities for moduli of smoothness, *Applied Mathematics E-Notes*, 12 (2012), 221-227.
- [34] S. Z. Jafarov, S. M. Nikolskii type inequality and estimation between the best approximations of a function in norms of different spaces. *Math. Balkanica (N.S.)* 21 (2007), no. 1-2, 173-182
- [35] D. M. Israfilov and R. Akgün, Approximation by polynomials and rational functions in weighted rearrangement invariant spaces, *J. Math. Anal. Appl.* 346 (2008), 489-500.
- [36] D. M. Israfilov and A. Testici, Approximation problems in the Lebesgue spaces with variable exponent, *Journal of Mathematical Analysis and Applications* 459:1, 2018, 112–123.

- [37] D. M. Israfilov and A. Testici, Approximation by Faber–Laurent rational functions in Lebesgue spaces with variable exponent, *Indag. Mat.* 27 (2016), No:4, 914-922.
- [38] D. M. Israfilov and E. Yirtici, Convolutions and best approximations in variable exponent Lebesgue spaces, *Math. Reports* 18(68) (2016), No:4 , 497-508.
- [39] H. Koc, Simultaneous approximation by polynomials in Orlicz spaces generated by quasiconvex Young functions, *Kuwait J. Sci.* 43 (2016); No:4, 18-31.
- [40] V. Kokilashvili and S. Samko, Singular integrals in weighted Lebesgue spaces with variable exponent, *Georgian Math. J.*, 10 (2003), N0: 1, 145-156.
- [41] Z. O. Kováčik and J. Rákosník, On spaces $L^{p(x)}$ and $W^{k,p(x)}$, *Czechoslovak Math. J.* **41(116)**:4 (1991), 592–618.
- [42] F.G. Nasibov, Approximation in L_2 by entire functions.(Russian) *Akad. Nauk Azerbaidzhan. SSR Dokl.* 1986; 42(4): 3–6.
- [43] S. M. Nikolskii, Inequalities for entire functions of finite degree and their application to the theory of differentiable functions of several variables, *Amer. Math. Soc. Transl. Ser. 2*, 80 (1969), 1-38, (*Trudy Mat. Inst. Steklov* 38 (1951), 211-278).
- [44] A.A. Ligun and V.G. Doronin, Exact constants in Jackson-type inequalities for the L_2 -approximation on a straight line. (Russian) *Ukrain. Mat. Zh.* 2009; 61(1): 92–98; translation in *Ukrainian Math. J.* 2009; 61(1): 112–120.
- [45] W. Orlicz, Über konjugierte Exponentenfolgen, *Studia Math.* **3** (1931), 200-212.
- [46] R. Paley and N. Wiener, *Fourier Transforms in the Complex Domain*, Amer. Math. Soc., 1934.
- [47] V. Yu. Popov, Best mean square approximations by entire functions of exponential type. (Russian) *Izv. Vysš. Uceb. Zaved. Matematika.* 1972; 121(6): 65—73.
- [48] K. R. Rajagopal and M. Ružička, On the modeling elektoreological materials, *Mech. Res. Commun.* **23**:4 (1996), 401–407.
- [49] M. Ružička, *Electrorheological Fluids: Modeling and Mathematical Theory*, Lecture Notes in Mathematics, 1748. Springer-Verlag, Berlin, 2000.
- [50] S. Samko, Differentiation and integration of variable order and the spaces $L^{p(x)}$, in: *Operator theory for complex and hypercomplex analysis* (Mexico City, 1994), 203–219, *Contemp. Math.*, 212, Amer. Math. Soc., Providence, RI, 1998.
- [51] I. I. Sharapudinov, The topology of the space $L^{p(t)}([0, 1])$, (Russian) *Mat. Zametki* **26**:4 (1979), 613–632.
- [52] I. I. Sharapudinov, Some questions in the theory of approximation in Lebesgue spaces with variable exponent, *Itogi Nauki. Yug Rossii. Mat. Monografiya*, vol. 5, Southern Institute of Mathematics of the Vladikavkaz Science Centre of the Russian Academy of Sciences and the Government of the Republic of North Ossetia-Alania, Vladikavkaz 2012, 267 pp. Russian.
- [53] A.F. Timan, *Theory of approximation of functions of a real variable*. Translated from the Russian by J. Berry. English translation edited and editorial preface by J. Cossar. *International Series of Monographs in Pure and Applied Mathematics*, Vol. 34, The Macmillan Co., New York: A Pergamon Press Book. 1963.
- [54] M.F. Timan, The approximation of functions defined on the whole real axis by entire functions of exponential type. *Izv. Vyssh. Uchebn. Zaved. Mat.* 1968; 2: 89—101.
- [55] R. Taberski, Approximation by entire functions of exponential type, 1981, *Demonstr. Math.* 14, 151-181 (1981).
- [56] R. Taberski, Contributions to fractional calculus and exponential approximation, 1986, *Funct. Approximatio, Comment. Math.* 15, 81-106 (1986).
- [57] S. S. Volosivets, Approximation of functions and their conjugates in variable Lebesgue spaces, *Sbornik: Mathematics*, 208 (2017), No:1, 44-59.
- [58] J. Yeh, *Real analysis: theory of measure and integration*, 2nd ed., 2006.
- [59] V. V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR Ser. Mat.* **50**:4 (1986), 675–710 (in Russian).
- [60] V. V. Zhuk, *Approximation of periodic functions*, LGU Press, 1982.