

# SIMPLICITY OF TWISTED C\*-ALGEBRAS OF DEACONU–RENAULT GROUPOIDS

BECKY ARMSTRONG, NATHAN BROWNLOWE, AND AIDAN SIMS

**ABSTRACT.** We consider Deaconu–Renault groupoids associated to actions of finite-rank free abelian monoids by local homeomorphisms of locally compact Hausdorff spaces. We study simplicity of the twisted C\*-algebra of such a groupoid determined by a continuous circle-valued groupoid 2-cocycle. When the groupoid is not minimal, this C\*-algebra is never simple, so we focus on minimal groupoids. We describe an action of the quotient of the groupoid by the interior of its isotropy on the spectrum of the twisted C\*-algebra of the interior of the isotropy. We prove that the twisted groupoid C\*-algebra is simple if and only if this action is minimal. We describe applications to crossed products of topological-graph C\*-algebras by quasi-free actions.

## 1. INTRODUCTION

The purpose of this paper is to characterise simplicity of twisted C\*-algebras arising from continuous 2-cocycles on Deaconu–Renault groupoids of actions of  $\mathbb{N}^k$  on second-countable locally compact Hausdorff spaces. The study of twisted C\*-algebras associated to continuous groupoid 2-cocycles dates back to Renault’s seminal work [31]. They serve both as a very flexible C\*-algebraic framework for modelling dynamical systems, and as a source of tractable models for classifiable C\*-algebras [29, 16, 8, 27]. So it is important to be able to determine when a given twisted groupoid C\*-algebra is simple; but this is in general a complicated question.

Deaconu–Renault groupoids encode actions of submonoids of abelian groups by local homeomorphisms of locally compact Hausdorff spaces. In hindsight, the first example of such a groupoid was the one associated to the one-sided full shift on  $n$  letters, introduced by Renault in [31] as a model for the Cuntz algebra. However groupoids of this type for generic local homeomorphisms (that is, actions of  $\mathbb{N}$ ) were first studied by Deaconu [11], and have come to be known as (rank-1) Deaconu–Renault groupoids. Shortly afterwards they were used as models for graph C\*-algebras in [20, 19], and later still, Yeend [38] showed that rank-1 Deaconu–Renault groupoids provide models for the topological-graph C\*-algebras of Katsura [17].

For the dual reasons that most of the key examples studied had been related to 0-dimensional spaces, and that  $\mathbb{N}$  embeds in  $\mathbb{Z}$ , which has trivial cohomology, no work was done on twisted C\*-algebras associated to Deaconu–Renault groupoids for many years. However, in 2000, Kumjian and Pask [18] introduced higher-rank graphs (or  $k$ -graphs) and demonstrated that the associated C\*-algebras can be described as the C\*-algebras of Deaconu–Renault groupoids of actions of  $\mathbb{N}^k$ . This led to the development [21, 22, 23] of twisted  $k$ -graph C\*-algebras. Kumjian, Pask, and Sims showed that from a

---

*Date:* 28th December 2021.

*2020 Mathematics Subject Classification.* 46L05 (primary).

*Key words and phrases.* C\*-algebra, groupoid, Deaconu–Renault, twist, simplicity.

The first author was supported by an Australian Government Research Training Program Stipend Scholarship. The second and third authors were supported by the Australian Research Council grant DP200100155. The first author would like to thank the second and third authors for being awesome PhD super(hero)visors.

2-cocycle on a  $k$ -graph, one can construct a 2-cocycle on the associated Deaconu–Renault groupoid so that the twisted  $C^*$ -algebras coincide, and they used this model to characterise simplicity of twisted  $k$ -graph  $C^*$ -algebras [24], as well as to describe applications of this characterisation to the study of crossed products of graph algebras by quasi-free actions.

Here we build substantially on elements of the analysis of [24] to describe precisely when the twisted  $C^*$ -algebra of a Deaconu–Renault groupoid for an action of  $\mathbb{N}^k$  by local homeomorphisms is simple (Theorem 6.1). To demonstrate the applicability of our main theorem, we use this result to investigate simplicity of crossed products of  $C^*$ -algebras associated to rank-1 Deaconu–Renault groupoids by actions of  $\mathbb{Z}$  induced by  $\mathbb{T}$ -valued 1-cocycles (Theorem 7.3), and we specialise to the Deaconu–Renault groupoids of topological graphs to characterise simplicity of crossed products of topological-graph  $C^*$ -algebras by quasi-free automorphisms (Corollaries 7.5 and 7.6).

The paper is organised as follows. In Section 2 we establish background and notation. In Section 3 we describe the periodicity group  $P_T$  of a minimal action  $T$  of  $\mathbb{N}^k$  on a second-countable locally compact Hausdorff space  $X$ , and we show that the interior of the isotropy of  $\mathcal{G}_T$  is isomorphic to the group bundle  $X \times P_T$ . In Section 4 we show that every 2-cocycle on  $\mathcal{G}_T$  is cohomologous to one whose restriction to  $X \times P_T$  is determined by a fixed bicharacter  $\omega$  of  $P_T$  that vanishes on its own centre  $Z_\omega$ , and we use this to give a concrete description of the spectral action  $\theta$  of  $\mathcal{G}_T/\mathcal{I}_T$  on  $X \times \widehat{Z}_\omega$ . Then in Section 6 we state and prove our main theorem. We finish in Section 7 by describing an application to crossed products of rank-1 Deaconu–Renault groupoid  $C^*$ -algebras by automorphisms induced by continuous 1-cocycles. We provide two appendices—one on group cohomology and one on twisted group  $C^*$ -algebras—to provide a handy reference to some key results on these two topics that we need in the body of the paper, and have found difficult to locate explicitly in the literature.

## 2. BACKGROUND

In this section we present the necessary background on Hausdorff étale groupoids, 2-cocycles, twisted groupoid  $C^*$ -algebras, and Deaconu–Renault groupoids.

We refer to a topological groupoid  $\mathcal{G}$  with a locally compact Hausdorff topology under which multiplication and inversion are continuous as a *Hausdorff groupoid*. We write  $\mathcal{G}^{(0)}$  for the unit space of  $\mathcal{G}$ , and  $\mathcal{G}^{(2)}$  for the set of composable pairs in  $\mathcal{G}$ . Given subsets  $A, B \subseteq \mathcal{G}$ , we write  $AB := \{\alpha\beta : (\alpha, \beta) \in (A \times B) \cap \mathcal{G}^{(2)}\}$  and  $A^{-1} := \{\alpha^{-1} : \alpha \in A\}$ , and for  $\gamma \in \mathcal{G}$ , we write  $\gamma A := \{\gamma\}A$  and  $A\gamma := A\{\gamma\}$ . We say that  $\mathcal{G}$  is *étale* if the range and source maps  $r, s : \mathcal{G} \rightarrow \mathcal{G}^{(0)}$  are local homeomorphisms. We call a subset  $B$  of  $\mathcal{G}$  a *bisection* if  $B$  is contained in an open subset  $U$  of  $\mathcal{G}$  such that  $r|_U$  and  $s|_U$  are homeomorphisms onto open subsets of  $\mathcal{G}^{(0)}$ . Every second-countable Hausdorff étale groupoid has a countable basis of open bisections. For each  $x \in \mathcal{G}^{(0)}$ , we define  $\mathcal{G}_x := s^{-1}(x)$  and  $\mathcal{G}^x := r^{-1}(x)$ ; and  $\mathcal{G}_x^x := \mathcal{G}_x \cap \mathcal{G}^x$ . We say that  $\mathcal{G}$  is *minimal* if  $r(\mathcal{G}_x)$  is dense in  $\mathcal{G}^{(0)}$  for every  $x \in \mathcal{G}^{(0)}$ . The *isotropy subgroupoid* of  $\mathcal{G}$  is the groupoid  $\text{Iso}(\mathcal{G}) := \bigcup_{x \in \mathcal{G}^{(0)}} \mathcal{G}_x^x = \{\gamma \in \mathcal{G} : r(\gamma) = s(\gamma)\}$ . The interior  $\mathcal{I}$  of the isotropy of a Hausdorff étale groupoid  $\mathcal{G}$  is itself a Hausdorff étale groupoid with unit space  $\mathcal{I}^{(0)} = \mathcal{G}^{(0)}$ . We say that  $\mathcal{G}$  is *effective* if  $\mathcal{I} = \mathcal{G}^{(0)}$ , and we say that  $\mathcal{G}$  is *topologically principal* if  $\{x \in \mathcal{G}^{(0)} : \mathcal{G}_x^x = \{x\}\}$  is dense in  $\mathcal{G}^{(0)}$ . By [7, Lemma 3.1], every topologically principal Hausdorff étale groupoid is effective, and every effective second-countable Hausdorff étale groupoid is topologically principal.

The following definition of a groupoid action comes from [15, Definition 1.60].

**Definition 2.1.** Suppose that  $\mathcal{G}$  is a topological groupoid and  $X$  is a topological space. We say that  $\mathcal{G}$  *acts continuously on (the left of)  $X$* , and that  $X$  is a *continuous (left)*

$\mathcal{G}$ -space, if there is a continuous surjective map  $R: X \rightarrow \mathcal{G}^{(0)}$  and a continuous map  $\theta: (\gamma, x) \mapsto \gamma \cdot x$  from  $\mathcal{G} \star X := \{(\gamma, x) \in \mathcal{G} \times X : s(\gamma) = R(x)\}$  to  $X$ , satisfying

(A1) if  $(\alpha, \beta) \in \mathcal{G}^{(2)}$  and  $(\beta, x) \in \mathcal{G} \star X$ , then  $(\alpha\beta, x)$ ,  $(\alpha, \beta \cdot x) \in \mathcal{G} \star X$ , and we have  $\alpha \cdot (\beta \cdot x) = (\alpha\beta) \cdot x$ ; and

(A2) for all  $x \in X$ , we have  $(R(x), x) \in \mathcal{G} \star X$ , and  $R(x) \cdot x = x$ .

We refer to the map  $\theta$  as a *continuous (left) action* of  $\mathcal{G}$  on  $X$ . For each  $x \in X$ , the *orbit* of  $x$  under  $\theta$  is the set

$$[x]_\theta := \{\gamma \cdot x : (\gamma, x) \in \mathcal{G} \star X\}.$$

We now recall the relevant cohomology theory for groupoids from [31, Section I.1]. See Appendix A for the relevant cohomology theory for groups.

**Definition 2.2.** Let  $\mathcal{G}$  be a topological groupoid, and let  $A$  be a topological abelian group with identity  $e_A$ .

- (i) A *continuous  $A$ -valued 1-cochain* on  $\mathcal{G}$  is a continuous map  $b: \mathcal{G} \rightarrow A$ . We say that  $b$  is *normalised* if  $b(r(\gamma)) = b(s(\gamma)) = e_A$  for all  $\gamma \in \mathcal{G}$ .
- (ii) A *continuous  $A$ -valued 1-cocycle* on  $\mathcal{G}$  is a continuous 1-cochain  $c: \mathcal{G} \rightarrow A$  satisfying  $c(\alpha\beta) = c(\alpha)c(\beta)$  for all  $(\alpha, \beta) \in \mathcal{G}^{(2)}$ .
- (iii) A *continuous  $A$ -valued 2-cocycle* on  $\mathcal{G}$  is a continuous map  $\sigma: \mathcal{G}^{(2)} \rightarrow A$  that satisfies the *2-cocycle identity*:  $\sigma(\alpha, \beta)\sigma(\alpha\beta, \gamma) = \sigma(\alpha, \beta\gamma)\sigma(\beta, \gamma)$  for all  $\alpha, \beta, \gamma \in \mathcal{G}$  such that  $s(\alpha) = r(\beta)$  and  $s(\beta) = r(\gamma)$ , and is *normalised*, in the sense that  $\sigma(r(\gamma), \gamma) = \sigma(\gamma, s(\gamma)) = e_A$  for all  $\gamma \in \mathcal{G}$ . We write  $Z^2(\mathcal{G}, A)$  for the group of continuous  $A$ -valued 2-cocycles on  $\mathcal{G}$ .
- (iv) The *continuous 2-coboundary* associated to a continuous normalised  $A$ -valued 1-cochain  $b: \mathcal{G} \rightarrow A$  is the map  $\delta^1 b: \mathcal{G}^{(2)} \rightarrow A$  given by  $\delta^1 b(\alpha, \beta) := b(\alpha)b(\beta)b(\alpha\beta)^{-1}$ .
- (v) We say that two continuous 2-cocycles  $\sigma, \tau: \mathcal{G}^{(2)} \rightarrow A$  are *cohomologous* if there exists a continuous normalised 1-cochain such that  $\delta^1 b(\alpha, \beta) = \sigma(\alpha, \beta)^{-1}\tau(\alpha, \beta)$  for all  $(\alpha, \beta) \in \mathcal{G}^{(2)}$ .

We write  $\mathbb{T}$  for the multiplicative group of complex numbers of modulus 1. Suppose that  $\mathcal{G}$  is a Hausdorff groupoid and take  $\sigma \in Z^2(\mathcal{G}, \mathbb{T})$ . Let  $\mathcal{G} \times_\sigma \mathbb{T}$  be the set  $\mathcal{G} \times \mathbb{T}$  endowed with the product topology, and equipped with the multiplication operation

$$(\alpha, w)(\beta, z) := (\alpha\beta, \sigma(\alpha, \beta)wz), \quad (2.1)$$

defined for all  $(\alpha, \beta) \in \mathcal{G}^{(2)}$  and  $w, z \in \mathbb{T}$ , and the inversion operation

$$(\alpha, w)^{-1} := (\alpha^{-1}, \overline{\sigma(\alpha, \alpha^{-1})w}), \quad (2.2)$$

defined for all  $(\alpha, w) \in \mathcal{G} \times \mathbb{T}$ . Then  $\mathcal{G} \times_\sigma \mathbb{T}$  is a Hausdorff groupoid.

We now recall Renault's construction of the full twisted groupoid C\*-algebra  $C^*(\mathcal{G}, \sigma)$  associated to a Hausdorff étale groupoid  $\mathcal{G}$  and a continuous  $\mathbb{T}$ -valued 2-cocycle  $\sigma$  on  $\mathcal{G}$ . Note that Renault gives this construction for groupoids that are not necessarily étale, but we specialise to the étale case since we will primarily be dealing with Deaconu–Renault groupoids, which are étale. Renault also defines reduced twisted groupoid C\*-algebras, but we will only be working with amenable groupoids, and in this setting, the full and reduced C\*-algebras coincide. Let  $C_c(\mathcal{G}, \sigma)$  denote the complex vector space of continuous compactly supported complex-valued functions on  $\mathcal{G}$ , equipped with multiplication given by the *twisted convolution* formula

$$(f * g)(\gamma) := \sum_{\substack{(\alpha, \beta) \in \mathcal{G}^{(2)}, \\ \alpha\beta = \gamma}} \sigma(\alpha, \beta) f(\alpha) g(\beta) = \sum_{\zeta \in \mathcal{G}^r(\gamma)} \sigma(\zeta, \zeta^{-1}\gamma) f(\zeta) g(\zeta^{-1}\gamma),$$

and involution given by

$$f^*(\gamma) := \overline{\sigma(\gamma, \gamma^{-1}) f(\gamma^{-1})}.$$

Then  $C_c(\mathcal{G}, \sigma)$  is a  $*$ -algebra. We write  $fg$  for the twisted convolution product  $f * g$  when the intended meaning is clear. The *full twisted groupoid  $C^*$ -algebra*  $C^*(\mathcal{G}, \sigma)$  is the completion of  $C_c(\mathcal{G}, \sigma)$  with respect to the *full  $C^*$ -norm*, which is given by

$$\|f\| := \sup \{ \|\pi(f)\| : \pi \text{ is a } *\text{-representation of } C_c(\mathcal{G}, \sigma) \}.$$

Given a locally compact Hausdorff space  $Y$  and a function  $f \in C_c(Y)$ , we define the *open support* of  $f$  to be the set  $\text{osupp}(f) := f^{-1}(\mathbb{C} \setminus \{0\})$ , and the *support* of  $f$  to be the set  $\text{supp}(f) := \overline{\text{osupp}(f)}$ .

We now recall the definition of the Deaconu–Renault groupoid associated to an action of  $\mathbb{N}^k$  by local homeomorphisms. Details appear in [35, Proposition 3.1].

Fix  $k \in \mathbb{N} \setminus \{0\}$ . Let  $T: n \mapsto T^n$  be an action of  $\mathbb{N}^k$  on a locally compact Hausdorff space  $X$  by local homeomorphisms. We call the pair  $(X, T)$  a *rank- $k$  Deaconu–Renault system*. Define

$$\mathcal{G}_T := \{(x, m - n, y) \in X \times \mathbb{Z}^k \times X : m, n \in \mathbb{N}^k, T^m(x) = T^n(y)\},$$

and

$$\mathcal{G}_T^{(2)} := \{((x, m, y), (w, n, z)) \in \mathcal{G}_T \times \mathcal{G}_T : y = w\}.$$

If  $((x, m, y), (y, n, z)) \in \mathcal{G}_T^{(2)}$ , then  $(x, m + n, z), (y, -m, x) \in \mathcal{G}_T$ . We define multiplication from  $\mathcal{G}_T^{(2)}$  to  $\mathcal{G}_T$  by  $(x, m, y)(y, n, z) := (x, m + n, z)$ , and inversion on  $\mathcal{G}_T$  by  $(x, m, y)^{-1} := (y, -m, x)$ . Then  $\mathcal{G}_T$  is a groupoid, called a *Deaconu–Renault groupoid*. The unit space of  $\mathcal{G}_T$  is  $\mathcal{G}_T^{(0)} = \{(x, 0, x) : x \in X\}$ , and we identify it with  $X$ . The range and source maps of  $\mathcal{G}_T$  are given by  $r(x, m, y) := x$  and  $s(x, m, y) := y$ . For open sets  $U, V \subseteq X$  and for  $m, n \in \mathbb{N}^k$ , we define

$$Z(U, m, n, V) := \{(x, m - n, y) : x \in U, y \in V, \text{ and } T^m(x) = T^n(y)\}.$$

The collection  $\{Z(U, m, n, V) : U, V \subseteq X \text{ are open, and } m, n \in \mathbb{N}^k\}$  is a basis for a locally compact Hausdorff topology on  $\mathcal{G}_T$ . The sets  $Z(U, m, n, V)$  such that  $T^m|_U$  and  $T^n|_V$  are homeomorphisms onto their ranges and  $T^m(U) = T^n(V)$  form a basis for the same topology. Under this topology,  $\mathcal{G}_T$  is a locally compact Hausdorff étale groupoid. If  $X$  is second-countable, then  $\mathcal{G}_T$  is also second-countable.

**Lemma 2.3.** *Let  $(X, T)$  be a rank- $k$  Deaconu–Renault system. The map  $c: (x, n, y) \mapsto n$  is a continuous  $\mathbb{Z}^k$ -valued 1-cocycle on  $\mathcal{G}_T$ , and for each  $x \in X$ , the restriction of  $c$  to  $(\mathcal{G}_T)_x$  is injective.*

*Proof.* Fix  $\alpha = (x, p, y)$  and  $\beta = (y, q, z) \in \mathcal{G}_T$ . Then  $c(\alpha\beta) = p + q = c(\alpha) + c(\beta)$ , and so  $c$  is a 1-cocycle. Since each  $c|_{Z(U, m, n, V)}$  is constant,  $c$  is locally constant and hence continuous.  $\square$

**Definition 2.4.** Let  $(X, T)$  be a rank- $k$  Deaconu–Renault system. The *orbit* under  $T$  of  $x \in X$  is

$$[x]_T := \bigcup_{m, n \in \mathbb{N}^k} (T^m)^{-1}(T^n(x)) = \{y \in X : T^m(y) = T^n(x) \text{ for some } m, n \in \mathbb{N}^k\}.$$

We say that  $(X, T)$  is *minimal* if  $[x]_T$  is dense in  $X$  for each  $x \in X$ . We frequently just write  $[x]$  for  $[x]_T$ .

*Remark 2.5.* We have  $[x] = r(s^{-1}(x)) \subseteq \mathcal{G}_T^{(0)}$ , and so  $\mathcal{G}_T$  is minimal if and only if  $(X, T)$  is minimal.

*Remark 2.6.* By [35, Lemma 3.5], every Deaconu–Renault groupoid is amenable, and so we can discuss the twisted C\*-algebra associated to a Deaconu–Renault groupoid and a continuous 2-cocycle without any ambiguity as to whether we mean the full or reduced C\*-algebra.

*Remark 2.7.* The C\*-algebras studied here are related to previous work. Suppose that  $\Lambda$  is a proper, source-free topological  $k$ -graph with infinite-path space  $\Lambda^\infty$  (as defined in [5, Section 3]). For each  $n \in \mathbb{N}^k$ , let  $T^n: \Lambda^\infty \rightarrow \Lambda^\infty$  be the shift map. Then  $(\Lambda^\infty, T)$  is a rank- $k$  Deaconu–Renault system, and the associated Deaconu–Renault groupoid  $\mathcal{G}_\Lambda := \mathcal{G}_T$  is called the *boundary-path groupoid* of the topological  $k$ -graph. The twisted C\*-algebras  $C^*(\mathcal{G}_\Lambda, \sigma)$  associated to continuous 2-cocycles  $\sigma \in Z^2(\mathcal{G}_\Lambda, \mathbb{T})$  on proper, source-free topological  $k$ -graphs generalise the twisted C\*-algebras of discrete  $k$ -graphs studied in [21, 22, 23, 24], and are studied in the first-named author’s PhD thesis [3]. In [5], the first- and second-named authors study an alternative notion of a twisted C\*-algebra of a topological  $k$ -graph associated to a continuous 2-cocycle on the topological  $k$ -graph itself, which is constructed using a product system of Hilbert bimodules. In the case where  $\Lambda$  is a discrete  $k$ -graph, it is known (see [2, Theorem 7.2.2]) that these two constructions give the same C\*-algebra, but in the more general topological setting, the relationship is unknown.

### 3. THE INTERIOR OF THE ISOTROPY OF A DEACONU–RENAULT GROUPOID

In this section we introduce the periodicity group  $P_T$  of a minimal Deaconu–Renault groupoid  $(X, T)$ , and we show that the interior  $\mathcal{I}_T$  of the isotropy of  $\mathcal{G}_T$  can be identified with  $X \times P_T$ .

**Definition 3.1.** Let  $(X, T)$  be a rank- $k$  Deaconu–Renault system. For each nonempty precompact open set  $U \subseteq X$ , we define

$$P_T(U) := \{m - n : m, n \in \mathbb{N}^k, \text{ and } T^m|_{\bar{U}} = T^n|_{\bar{U}} \text{ is injective}\}.$$

We define

$$P_T := \bigcup_{\emptyset \neq U \subseteq X \text{ precompact open}} P_T(U).$$

*Remark 3.2.* When  $k = 1$ , the set  $P_T(U)$  is related to the group  $\text{Stab}^{\text{ess}}(x)$  from [10, Page 29]. Specifically,  $\text{Stab}^{\text{ess}}(x)$  contains  $P_T(U)$  for any precompact open set  $U$  containing  $x$ ; but also, since  $\text{Stab}^{\text{ess}}(x)$  is a subgroup of  $\mathbb{Z}^k$ , and hence finitely generated, it is not too hard to check that there is an open cover of  $X$  by sets  $U$  such that  $P_T(U) = \text{Stab}^{\text{ess}}(x)$  for each  $x$  in  $U$ .

In addition to being needed for our own arguments, our next result, [Proposition 3.5](#), plugs a gap in the literature—it is mentioned without proof in [10, Page 30].

**Proposition 3.3.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system. Then*

$$P_T = \{p \in \mathbb{Z}^k : (x, p, x) \in \mathcal{G}_T \text{ for all } x \in X\},$$

*and  $P_T$  is a subgroup of  $\mathbb{Z}^k$ .*

In order to prove [Proposition 3.3](#), we need the following lemma.

**Lemma 3.4.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system. Suppose that  $m, n \in \mathbb{N}^k$  and  $y \in X$  satisfy  $T^m(y) \neq T^n(y)$ . Then there exists an open neighbourhood  $W \subseteq X$  of  $y$  such that  $T^m|_W$  and  $T^n|_W$  are injective and  $T^m(W) \cap T^n(W) = \emptyset$ .*



*Proof.* Since  $X$  is Hausdorff, we can choose open neighbourhoods  $U \subseteq X$  of  $T^m(y)$  and  $V \subseteq X$  of  $T^n(y)$  such that  $U \cap V = \emptyset$ . Define  $A := (T^m)^{-1}(U) \cap (T^n)^{-1}(V)$ . Then  $y \in A$ . Since  $T^m$  and  $T^n$  are local homeomorphisms, there is an open neighbourhood  $W \subseteq A$  of  $y$  such that  $T^m|_W$  and  $T^n|_W$  are injective, and we have  $T^m(W) \cap T^n(W) \subseteq U \cap V = \emptyset$ .  $\square$

*Proof of Proposition 3.3.* Fix  $p \in P_T$ . Then there exist  $m, n \in \mathbb{N}^k$  and a nonempty open set  $U \subseteq X$  such that  $p = m - n$ , and  $T^m|_U = T^n|_U$  is injective. Fix  $z \in X$ . We claim that  $(z, p, z) \in \mathcal{G}_T$ . Since  $[z]$  is dense in  $X$ , we have  $U \cap [z] \neq \emptyset$ , and so there exist  $y \in U$  and  $a, b \in \mathbb{N}^k$  such that  $T^a(y) = T^b(z)$ . Thus,

$$T^{b+m}(z) = T^m(T^b(z)) = T^m(T^a(y)) = T^a(T^m(y)),$$

and

$$T^{b+n}(z) = T^n(T^b(z)) = T^n(T^a(y)) = T^a(T^n(y)).$$

Since  $y \in U$ , we have  $T^m(y) = T^n(y)$ , and hence  $T^{b+m}(z) = T^{b+n}(z)$ . Therefore,  $(z, p, z) = (z, (b+m) - (b+n), z) \in \mathcal{G}_T$ , and so

$$P_T \subseteq \{p \in \mathbb{Z}^k : (x, p, x) \in \mathcal{G}_T \text{ for all } x \in X\}.$$

We now show that  $\mathbb{Z}^k \setminus P_T \subseteq \{p \in \mathbb{Z}^k : (x, p, x) \notin \mathcal{G}_T \text{ for some } x \in X\}$ . To see this, fix  $p \in \mathbb{Z}^k \setminus P_T$ . Let  $(m_i, n_i)_{i=1}^\infty$  be an enumeration of  $\{(m, n) \in \mathbb{N}^k \times \mathbb{N}^k : m - n = p\}$ . We must find  $x \in X$  such that  $T^{m_i}(x) \neq T^{n_i}(x)$  for all  $i \geq 1$ . We claim that there exist nonempty precompact open subsets  $V_0, V_1, V_2, \dots$  of  $X$  satisfying

- (1)  $\overline{V_i} \subseteq V_{i-1}$  for all  $i \geq 1$ ,
- (2)  $T^{m_i}|_{\overline{V_{i-1}}}$  and  $T^{n_i}|_{\overline{V_{i-1}}}$  are injective for all  $i \geq 1$ , and
- (3)  $T^{m_i}(\overline{V_i}) \cap T^{n_i}(\overline{V_i}) = \emptyset$  for all  $i \geq 1$ .

To start, let  $V_0$  be a nonempty precompact open subset of  $X$  such that  $T^{m_1}|_{\overline{V_0}}$  and  $T^{n_1}|_{\overline{V_0}}$  are injective. Now fix  $i \geq 1$  and suppose that  $V_0, \dots, V_{i-1}$  satisfy (1)–(3). Since  $m_i - n_i = p \notin P_T$ , we have  $T^{m_i}|_{V_{i-1}} \neq T^{n_i}|_{V_{i-1}}$ , and so there exists  $y \in V_{i-1}$  such that  $T^{m_i}(y) \neq T^{n_i}(y)$ . Thus, by Lemma 3.4, there exists an open neighbourhood  $W \subseteq V_{i-1}$  of  $y$  such that  $T^{m_i}(W) \cap T^{n_i}(W) = \emptyset$ . Since  $X$  is locally compact and Hausdorff and  $T^{m_{i+1}}$  and  $T^{n_{i+1}}$  are local homeomorphisms, there is an open neighbourhood  $V_i$  of  $y$  such that  $\overline{V_i} \subseteq W$  and  $T^{m_{i+1}}|_{\overline{V_i}}$  and  $T^{n_{i+1}}|_{\overline{V_i}}$  are injective. So induction gives the desired sets  $V_i$ . Each  $V_i$  is contained in the compact set  $\overline{V_0}$ , and so the descending intersection  $\bigcap_{i=1}^\infty \overline{V_i}$  is nonempty. Any  $x \in \bigcap_{i=1}^\infty \overline{V_i}$  satisfies  $T^{m_i}(x) \neq T^{n_i}(x)$  for all  $i \geq 1$ .

We conclude by showing that  $P_T$  is a subgroup of  $\mathbb{Z}^k$ . For all  $x \in X$ , we have  $(x, 0, x) \in \mathcal{G}_T^{(0)} \subseteq \mathcal{G}_T$ , and so  $0 \in P_T$ . Suppose that  $p, q \in P_T$ . For all  $x \in X$ , we have  $(x, p, x), (x, q, x) \in \mathcal{G}_T$ , and hence  $(x, p - q, x) = (x, p, x)(x, q, x)^{-1} \in \mathcal{G}_T$ . Thus  $p - q \in P_T$ , and so  $P_T$  is a subgroup of  $\mathbb{Z}^k$ .  $\square$

Given a rank- $k$  Deaconu–Renault system  $(X, T)$ , we write  $\mathcal{I}_T$  for the topological interior of  $\text{Iso}(\mathcal{G}_T)$ . Since  $\mathcal{G}_T$  is a locally compact Hausdorff étale groupoid, so is  $\mathcal{I}_T$ . From this point forward, we will assume that  $X$  is second-countable (and hence so are  $\mathcal{G}_T$  and  $\mathcal{I}_T$ ). We know from [35, Lemma 3.5] that  $\mathcal{G}_T$  is amenable, and hence [1, Proposition 5.1.1] implies that  $\mathcal{I}_T$  is amenable.

**Proposition 3.5.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Let  $P_T$  be as in Definition 3.1. Then*

$$\mathcal{I}_T = \{(x, p, x) : x \in X, p \in P_T\} \cong X \times P_T.$$

*Proof.* For  $\subseteq$ , fix  $\gamma \in \mathcal{I}_T$ . Let  $c: \mathcal{G}_T \rightarrow \mathbb{Z}^k$  be the continuous 1-cocycle defined in Lemma 2.3. Let  $p := c(\gamma)$  so that  $\gamma = (x, p, x)$  for some  $x \in X$ . We claim that  $p \in P_T$ .

By [Remark 2.5](#),  $\mathcal{G}_T$  is minimal, and hence [\[24, Proposition 2.1\]](#) implies that for all  $y \in X$ ,

$$c(\mathcal{I}_T \cap (\mathcal{G}_T)_y^y) = c(\mathcal{I}_T \cap (\mathcal{G}_T)_x^x),$$

and thus

$$p = c(x, p, x) \in c(\mathcal{I}_T \cap (\mathcal{G}_T)_x^x) = c(\mathcal{I}_T \cap (\mathcal{G}_T)_y^y).$$

So [Proposition 3.3](#) gives  $p \in P_T$ .

For  $\supseteq$ , fix  $x \in X$  and  $p \in P_T$ . By the definition of  $P_T$ , there exist  $m, n \in \mathbb{N}^k$  and a nonempty precompact open set  $U \subseteq X$  such that  $p = m - n$  and  $T^m|_{\bar{U}} = T^n|_{\bar{U}}$  is injective. This injectivity forces  $Z(U, m, n, U) = \{(y, p, y) : y \in U\} \subseteq \mathcal{I}_T$ . Fix  $y \in U$ . Then  $(y, p, y) \in \mathcal{I}_T$ , and so [\[24, Proposition 2.1\]](#) implies that

$$p = c(y, p, y) \in c(\mathcal{I}_T \cap (\mathcal{G}_T)_y^y) = c(\mathcal{I}_T \cap (\mathcal{G}_T)_x^x),$$

and hence  $(x, p, x) \in \mathcal{I}_T$ .  $\square$

*Remark 3.6.* [Proposition 3.5](#) is related to the sets  $\Sigma_X$  and  $H(T)$  of [\[35\]](#) as follows. Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. In the notation of [\[35, Section 3\]](#), suppose that  $\Sigma = \Sigma_X$ . Then  $T$  is an irreducible action of  $\mathbb{N}^k$  on  $X$ , and [\[35, Proposition 3.10\]](#) implies that

$$\mathcal{I}_T = \{(x, p, x) : x \in X, p \in H(T)\} \cong X \times H(T).$$

Thus [Proposition 3.5](#) implies that  $P_T = H(T)$ .

We now present two corollaries of [Proposition 3.5](#).

**Corollary 3.7.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Then  $\mathcal{G}_T$  is effective if and only if  $P_T = \{0\}$ .*

*Proof.* By [Proposition 3.5](#),  $\mathcal{I}_T = \{(x, p, x) : x \in X, p \in P_T\}$ . Hence

$$\mathcal{G}_T \text{ is effective} \iff \mathcal{I}_T = \mathcal{G}_T^{(0)} \iff P_T = \{0\}. \quad \square$$

**Corollary 3.8.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Let  $c : \mathcal{G}_T \rightarrow \mathbb{Z}^k$  be as in [Lemma 2.3](#). Then for each  $p \in P_T$ , we have  $(c|_{\mathcal{I}_T})^{-1}(p) = \{(x, p, x) : x \in X\}$ , and  $\{(c|_{\mathcal{I}_T})^{-1}(p) : p \in P_T\}$  is a collection of mutually disjoint clopen bisections whose union is  $\mathcal{I}_T$ .*

*Proof.* Fix  $p \in P_T$ . Since  $c$  is continuous and  $\mathbb{Z}^k$  is discrete,  $(c|_{\mathcal{I}_T})^{-1}(p)$  is clopen. Fix  $x \in X$ . If  $\alpha, \beta \in (c|_{\mathcal{I}_T})^{-1}(p)$  and  $r(\alpha) = x = r(\beta)$ , then  $s(\alpha) = x = s(\beta)$  because  $\alpha, \beta \in \text{Iso}(\mathcal{G}_T)$ , and hence  $\alpha = (x, p, x) = \beta$ . So  $r|_{(c|_{\mathcal{I}_T})^{-1}(p)}$  is injective, and a similar argument shows that  $s|_{(c|_{\mathcal{I}_T})^{-1}(p)}$  is also injective. Hence  $(c|_{\mathcal{I}_T})^{-1}(p)$  is a bisection. By [Proposition 3.5](#), we have  $c(\mathcal{I}_T) = P_T$ , and the result follows.  $\square$

We now prove that if  $\mathcal{G}_T$  is minimal, then we can form the quotient groupoid  $\mathcal{G}_T/\mathcal{I}_T$ .

**Proposition 3.9.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Then  $\mathcal{I}_T$  is a closed subgroupoid of  $\mathcal{G}_T$  and acts freely and properly on  $\mathcal{G}_T$  by right-multiplication. The set  $\mathcal{H}_T := \mathcal{G}_T/\mathcal{I}_T$  is a locally compact Hausdorff étale groupoid, with multiplication given by  $[\alpha][\beta] := [\alpha\beta]$  for  $(\alpha, \beta) \in \mathcal{G}_T^{(2)}$ , inversion given by  $[\gamma]^{-1} := [\gamma^{-1}]$  for  $\gamma \in \mathcal{G}_T$ , and range and source maps given by  $r([\gamma]) = [r(\gamma)]$  and  $s([\gamma]) = [s(\gamma)]$ .*

*Proof.* Together, [Remark 2.5](#) and [Lemma 2.3](#) allow us to apply [\[24, Proposition 2.1\]](#) to see that  $\mathcal{I}_T$  is a closed subgroupoid of  $\mathcal{G}_T$ . Therefore, [\[35, Proposition 2.5\(d\)\]](#) implies that  $\mathcal{H}_T$  is a locally compact Hausdorff étale groupoid under the given operations.  $\square$

We conclude this section with two technical lemmas that we use in the proof of our characterisation of simplicity of  $C^*(\mathcal{G}_T, \sigma)$  in [Section 6](#).

**Lemma 3.10.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Let  $c: \mathcal{G}_T \rightarrow \mathbb{Z}^k$  be as in [Lemma 2.3](#). Fix  $m, n \in \mathbb{Z}^k$ , and let  $U$  and  $V$  be precompact open bisections of  $\mathcal{G}_T$  such that  $U \subseteq c^{-1}(m)$  and  $V \subseteq c^{-1}(n)$ . Then  $\overline{\mathcal{I}_T U \cap V} \subseteq \overline{\mathcal{I}_T U}$  and  $\overline{\mathcal{I}_T V \cap U} \subseteq \overline{\mathcal{I}_T V}$ .*

*Proof.* Define  $K := r(\overline{U}) \times \{n - m\} \times r(\overline{U})$  and  $W := K \cap \mathcal{I}_T$ . Since  $r$  is continuous,  $K$  is compact, and hence closed. Since  $\mathcal{I}_T$  is closed by [Proposition 3.9](#),  $W$  is closed, and hence is a compact subset of  $K$  and of  $\mathcal{I}_T$ . We claim that  $\mathcal{I}_T U \cap V \subseteq W\overline{U}$ . For this, suppose that  $\gamma \in \mathcal{I}_T U \cap V$ . Then there exist  $\xi \in \mathcal{I}_T$  and  $\eta \in U \subseteq c^{-1}(m)$  such that  $\gamma = \xi\eta \in V \subseteq c^{-1}(n)$ . Hence  $\xi = \gamma\eta^{-1} \subseteq c^{-1}(n - m)$ . We also have  $r(\xi) = s(\xi) = r(\eta) \in r(\overline{U})$ , and so  $\xi \in K \cap \mathcal{I}_T = W$ . Hence  $\gamma = \xi\eta \in W\overline{U}$ , and so  $\mathcal{I}_T U \cap V \subseteq W\overline{U}$ . Since  $W$  and  $\overline{U}$  are compact,  $W\overline{U}$  is compact, and hence closed. Thus  $\overline{\mathcal{I}_T U \cap V} \subseteq \overline{\mathcal{I}_T U}$ . A symmetric argument shows that  $\overline{\mathcal{I}_T V \cap U} \subseteq \overline{\mathcal{I}_T V}$ .  $\square$

**Lemma 3.11.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Let  $c: \mathcal{G}_T \rightarrow \mathbb{Z}^k$  be as in [Lemma 2.3](#). Fix  $m, n \in \mathbb{Z}^k$ , and let  $U$  and  $V$  be precompact open bisections of  $\mathcal{G}_T$  such that  $U \subseteq c^{-1}(m)$  and  $V \subseteq c^{-1}(n)$ . Then  $s(\mathcal{I}_T U \cap V) = s(\mathcal{I}_T V \cap U)$ . Moreover, if  $(V^{-1}\mathcal{I}_T U) \cap \mathcal{I}_T \neq \emptyset$ , then  $n - m \in P_T$ , and for each  $\gamma \in (V^{-1}\mathcal{I}_T U) \cap \mathcal{I}_T$ , we have  $s(\gamma) \in s(\mathcal{I}_T U \cap V)$ .*

*Proof.* We first show that  $s(\mathcal{I}_T U \cap V) = s(\mathcal{I}_T V \cap U)$ . By symmetry, it suffices to show that  $s(\mathcal{I}_T U \cap V) \subseteq s(\mathcal{I}_T V \cap U)$ . Suppose that  $x \in s(\mathcal{I}_T U \cap V)$ . Then there exist  $\zeta \in \mathcal{I}_T$  and  $\eta \in U$  such that  $\zeta\eta \in V$  and  $x = s(\zeta\eta) = s(\eta)$ . Since  $\zeta^{-1} \in \mathcal{I}_T$ , we have  $\eta = \zeta^{-1}(\zeta\eta) \in \mathcal{I}_T V \cap U$ , and hence  $x = s(\eta) \in s(\mathcal{I}_T V \cap U)$ . Thus  $s(\mathcal{I}_T U \cap V) \subseteq s(\mathcal{I}_T V \cap U)$ , as required.

For the second statement, suppose that  $\gamma \in (V^{-1}\mathcal{I}_T U) \cap \mathcal{I}_T$ . Then there exist  $\alpha \in U$ ,  $\beta \in V$ , and  $\xi \in \mathcal{I}_T$  such that  $\gamma = \beta^{-1}\xi\alpha$ , and hence  $c(\gamma) = -n + c(\xi) + m$ . Since  $\gamma, \xi \in \mathcal{I}_T$ , we have  $c(\gamma), c(\xi) \in P_T$  by [Proposition 3.5](#), and hence  $n - m = c(\xi) - c(\gamma) \in P_T$ , because  $P_T$  is a group by [Proposition 3.3](#). Since  $\gamma \in \mathcal{I}_T$ , we have  $s(\beta) = r(\gamma) = s(\gamma) = r(\alpha^{-1})$ , and hence  $(\beta, \alpha^{-1}) \in \mathcal{G}_T^{(2)}$ . Since  $\xi \in \mathcal{I}_T$ , we have  $r(\beta\alpha^{-1}) = s(\beta^{-1}) = r(\xi) = s(\xi) = r(\alpha) = s(\beta\alpha^{-1})$ . We also have  $c(\beta\alpha^{-1}) = n - m \in P_T$ , and thus [Proposition 3.5](#) implies that  $\beta\alpha^{-1} \in \mathcal{I}_T$ . Hence  $\beta = (\beta\alpha^{-1})\alpha \in \mathcal{I}_T U \cap V$ , and so  $s(\gamma) = r(\gamma) = s(\beta) \in s(\mathcal{I}_T U \cap V)$ .  $\square$

#### 4. COHOMOLOGY OF DEACONU–RENAULT GROUPOIDS

In this section we show that every continuous  $\mathbb{T}$ -valued 2-cocycle on a minimal Deaconu–Renault groupoid  $\mathcal{G}_T$  is cohomologous to a continuous  $\mathbb{T}$ -valued 2-cocycle  $\sigma$  on  $\mathcal{G}_T$  that is constant on  $\mathcal{I}_T$  (in the sense of [Definition 4.1](#)). We also introduce the spectral action  $\theta$  of  $\mathcal{H}_T := \mathcal{G}_T/\mathcal{I}_T$ , analogous to [\[24, Lemma 3.6\]](#).

**Definition 4.1.** Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Suppose that  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . We say that  $\sigma$  is *constant on  $\mathcal{I}_T$*  if

$$\sigma((x, m, x), (x, n, x)) = \sigma((y, m, y), (y, n, y)) \quad \text{for all } x, y \in X \text{ and } m, n \in P_T.$$

If  $\omega \in Z^2(P_T, \mathbb{T})$  is the 2-cocycle satisfying  $\sigma((x, m, x), (x, n, x)) = \omega(m, n)$  for all  $x \in X$  and  $m, n \in P_T$ , then we say that  $\sigma$  is  *$\omega$ -constant on  $\mathcal{I}_T$* , and we write  $\sigma|_{\mathcal{I}_T^{(2)}} = 1_X \times \omega$ .

Recall that if  $A$  is a countable discrete abelian group, then a *bicharacter* of  $A$  is a function  $\omega: A \times A \rightarrow \mathbb{T}$  such that for each fixed  $a \in A$ , the maps  $\omega(\cdot, a): A \rightarrow \mathbb{T}$  and  $\omega(a, \cdot): A \rightarrow \mathbb{T}$  are group homomorphisms. (See [Appendix A](#) for some additional



background on bicharacters and cohomology of groups.) The following proposition and the lemmas used in its proof are extensions of cohomological results from [24, Section 3] about boundary-path groupoids of cofinal, row-finite, source-free  $k$ -graphs to the more general setting of Deaconu–Renault groupoids. **Proposition 4.2** is a generalisation of [24, Proposition 3.1], but we have adapted it slightly to prove that the bicharacter  $\omega \in Z^2(P_T, \mathbb{T})$  can be chosen in such a way that it vanishes on its centre  $Z_\omega$  in each coordinate (in the sense that  $\omega(Z_\omega, P_T) \cup \omega(P_T, Z_\omega) = \{1\}$ ; see **Definition A.5**), and hence descends to a bicharacter  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$ .

**Proposition 4.2.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Suppose that  $\rho \in Z^2(\mathcal{G}_T, \mathbb{T})$ . For each  $x \in X$ , define  $\rho_x: P_T \times P_T \rightarrow \mathbb{T}$  by*

$$\rho_x(m, n) := \rho((x, m, x), (x, n, x)).$$

*Then  $\rho_x \in Z^2(P_T, \mathbb{T})$ . There exists a bicharacter  $\omega \in Z^2(P_T, \mathbb{T})$  such that  $\omega$  vanishes on  $Z_\omega$  in each coordinate, and  $\omega$  is cohomologous to  $\rho_x$  for every  $x \in X$ . For any such bicharacter  $\omega$ , there exists  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$  such that  $\sigma$  is cohomologous to  $\rho$  and is  $\omega$ -constant on  $\mathcal{I}_T$  (in the sense of **Definition 4.1**), and there exists a bicharacter  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$  such that*

$$\tilde{\omega}(p + Z_\omega, q + Z_\omega) = \omega(p, q) \quad \text{for all } p, q \in P_T.$$

In order to prove **Proposition 4.2**, we need the following two results. The first of these results is an extension of [24, Lemma 3.2] to the setting of Deaconu–Renault groupoids.

**Lemma 4.3.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . For each  $x \in X$ , define  $\sigma_x: P_T \times P_T \rightarrow \mathbb{T}$  by*

$$\sigma_x(m, n) := \sigma((x, m, x), (x, n, x)).$$

*Then  $\sigma_x \in Z^2(P_T, \mathbb{T})$ . For  $\gamma \in \mathcal{G}_T$  and  $y := s(\gamma) \in X$ , define  $\tau_\gamma^\sigma: P_T \rightarrow \mathbb{T}$  by*

$$\tau_\gamma^\sigma(p) := \sigma(\gamma, (y, p, y)) \sigma(\gamma(y, p, y), \gamma^{-1}) \overline{\sigma(\gamma, \gamma^{-1})}. \quad (4.1)$$

- (a) *For all  $m, n \in P_T$ , the map  $x \mapsto \sigma_x(m, n)$  from  $X$  to  $\mathbb{T}$  is continuous.*
- (b) *For each  $p \in P_T$ , the map  $\gamma \mapsto \tau_\gamma^\sigma(p)$  from  $\mathcal{G}_T$  to  $\mathbb{T}$  is continuous.*
- (c) *Fix  $\gamma = (x, m, y) \in \mathcal{G}_T$ ,  $p \in P_T$ , and  $w, z \in \mathbb{T}$ . Under the multiplication and inversion operations on  $\mathcal{G}_T \times_\sigma \mathbb{T}$  (as defined in **Equations (2.1) and (2.2)**), we have*

$$(\gamma, w)((y, p, y), z)(\gamma, w)^{-1} = ((x, p, x), \tau_\gamma^\sigma(p)z). \quad (4.2)$$

- (d) *For all  $\gamma \in \mathcal{G}_T$  and  $p, q \in P_T$ , we have*

$$\tau_\gamma^\sigma(p + q) = \sigma_{r(\gamma)}(p, q) \overline{\sigma_{s(\gamma)}(p, q)} \tau_\gamma^\sigma(p) \tau_\gamma^\sigma(q). \quad (4.3)$$

- (e) *If  $\omega$  is a bicharacter of  $P_T$  such that  $\sigma$  is  $\omega$ -constant on  $\mathcal{I}_T$ , then  $\sigma_x = \sigma_y$  for all  $x, y \in X$ , and  $\gamma \mapsto \tau_\gamma^\sigma$  is a continuous  $\widehat{P}_T$ -valued 1-cocycle on  $\mathcal{G}_T$ .*

*Proof.* Routine calculations show that since  $\sigma$  is normalised and satisfies the 2-cocycle identity, we have  $\sigma_x \in Z^2(P_T, \mathbb{T})$  for each  $x \in X$ .

For part (a), note that for each  $m, n \in P_T$ , the map  $x \mapsto \sigma_x(m, n)$  is the composition of the continuous maps  $x \mapsto ((x, m, x), (x, n, x))$  and  $\sigma$ .

For part (b), fix  $p \in P_T$ . For  $\gamma \in \mathcal{G}_T$ ,

$$\tau_\gamma^\sigma(p) = \sigma(\gamma, (s(\gamma), p, s(\gamma))) \sigma(\gamma(s(\gamma), p, s(\gamma)), \gamma^{-1}) \overline{\sigma(\gamma, \gamma^{-1})}.$$

Thus the map  $\gamma \mapsto \tau_\gamma^\sigma(p)$  from  $\mathcal{G}_T$  to  $\mathbb{T}$  is continuous because it is a product of continuous functions.

For part (c), fix  $\gamma = (x, m, y) \in \mathcal{G}_T$ ,  $p \in P_T$ , and  $w, z \in \mathbb{T}$ . We have

$$\gamma(y, p, y)\gamma^{-1} = (x, m, y)(y, p, y)(y, -m, x) = (x, p, x),$$

and hence

$$\begin{aligned} (\gamma, w)((y, p, y), z)(\gamma, w)^{-1} &= (\gamma(y, p, y), \sigma(\gamma, (y, p, y)) wz)(\gamma^{-1}, \overline{\sigma(\gamma, \gamma^{-1}) \overline{w}}) \\ &= (\gamma(y, p, y)\gamma^{-1}, \sigma(\gamma(y, p, y), \gamma^{-1}) \sigma(\gamma, (y, p, y)) \overline{\sigma(\gamma, \gamma^{-1}) z}) \\ &= ((x, p, x), \tau_\gamma^\sigma(p)z). \end{aligned}$$

For part (d), fix  $\gamma = (x, m, y) \in \mathcal{G}_T$  and  $p, q \in P_T$ . For all  $z \in \mathbb{T}$ , we have

$$\begin{aligned} ((y, p, y), 1)((y, q, y), z) &= ((y, p, y)(y, q, y), \sigma((y, p, y), (y, q, y))z) \\ &= ((y, p+q, y), \sigma_y(p, q)z), \end{aligned}$$

and so, taking  $z = \overline{\sigma_y(p, q)}$ , we see that

$$((y, p, y), 1)((y, q, y), \overline{\sigma_y(p, q)}) = ((y, p+q, y), 1). \quad (4.4)$$

Together, Equations (4.2) and (4.4) imply that

$$\begin{aligned} ((x, p+q, x), \tau_\gamma^\sigma(p+q)) &= (\gamma, 1)((y, p+q, y), 1)(\gamma, 1)^{-1} \\ &= (\gamma, 1)((y, p, y), 1)((y, q, y), \overline{\sigma_y(p, q)})(\gamma, 1)^{-1} \\ &= (\gamma, 1)((y, p, y), 1)(\gamma, 1)^{-1}(\gamma, 1)((y, q, y), \overline{\sigma_y(p, q)})(\gamma, 1)^{-1} \\ &= ((x, p, x), \tau_\gamma^\sigma(p))((x, q, x), \tau_\gamma^\sigma(q) \overline{\sigma_y(p, q)}) \\ &= ((x, p+q, x), \sigma_x(p, q) \overline{\sigma_y(p, q)} \tau_\gamma^\sigma(p) \tau_\gamma^\sigma(q)), \end{aligned}$$

and hence

$$\tau_\gamma^\sigma(p+q) = \sigma_{r(\gamma)}(p, q) \overline{\sigma_{s(\gamma)}(p, q)} \tau_\gamma^\sigma(p) \tau_\gamma^\sigma(q).$$

For part (e), since  $\sigma$  is  $\omega$ -constant on  $\mathcal{I}_T$ , for all  $x, y \in X$  and  $p, q \in P_T$ , we have

$$\sigma_x(p, q) = \sigma((x, p, x), (x, q, x)) = \omega(p, q) = \sigma((y, p, y), (y, q, y)) = \sigma_y(p, q).$$

So  $\sigma_x = \sigma_y$ , and for each  $\gamma \in \mathcal{G}_T$ , Equation (4.3) reduces to

$$\tau_\gamma^\sigma(p+q) = \tau_\gamma^\sigma(p) \tau_\gamma^\sigma(q).$$

Thus  $\tau_\gamma^\sigma: P_T \rightarrow \mathbb{T}$  is a homomorphism, and so  $\tau_\gamma^\sigma \in \widehat{P}_T$  for each  $\gamma \in \mathcal{G}_T$ .

We now show that the map  $\gamma \mapsto \tau_\gamma^\sigma$  is multiplicative. Fix  $\alpha = (x, m, u)$ ,  $\beta = (u, n, y) \in \mathcal{G}_T$ , and  $p \in P_T$ . Using Equation (4.2), we compute

$$\begin{aligned} ((x, p, x), \tau_\alpha^\sigma(p) \tau_\beta^\sigma(p)) &= (\alpha, 1)((u, p, u), \tau_\beta^\sigma(p))(\alpha, 1)^{-1} \\ &= (\alpha, 1)(\beta, 1)((y, p, y), 1)(\beta, 1)^{-1}(\alpha, 1)^{-1} \\ &= (\alpha\beta, \sigma(\alpha, \beta))((y, p, y), 1)(\alpha\beta, \sigma(\alpha, \beta))^{-1} \\ &= ((x, p, x), \tau_{\alpha\beta}^\sigma(p)). \end{aligned}$$

Hence  $\tau_\alpha^\sigma \tau_\beta^\sigma = \tau_{\alpha\beta}^\sigma$ , and so  $\gamma \mapsto \tau_\gamma^\sigma$  is a  $\widehat{P}_T$ -valued 1-cocycle on  $\mathcal{G}_T$ .

We conclude by showing that the map  $\gamma \mapsto \tau_\gamma^\sigma$  is continuous. Fix a finite subset  $F \subseteq P_T$  and an open subset  $U \subseteq \mathbb{T}$ . The set

$$S_{\widehat{P}_T}(F, U) = \{\phi \in \widehat{P}_T : \phi(F) \subseteq U\}$$

is a typical subbasis element for the compact-open topology on  $\widehat{P}_T$ , and so it suffices to show that  $\{\gamma \in \mathcal{G}_T : \tau_\gamma^\sigma \in S_{\widehat{P}_T}(F, U)\}$  is an open subset of  $\mathcal{G}_T$ . We have

$$\{\gamma \in \mathcal{G}_T : \tau_\gamma^\sigma(F) \subseteq U\} = \bigcap_{p \in F} \{\gamma \in \mathcal{G}_T : \tau_\gamma^\sigma(p) \in U\},$$

which is open by part (b).  $\square$

The following lemma is an extension of [24, Lemma 3.3] to the setting of Deaconu–Renault groupoids.

**Lemma 4.4.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . As in Lemma 4.3, for each  $x \in X$ , define  $\sigma_x \in Z^2(P_T, \mathbb{T})$  by*

$$\sigma_x(m, n) := \sigma((x, m, x), (x, n, x)).$$

*Then the cohomology class of  $\sigma_x$  does not depend on  $x$ .*

*Proof.* By Theorem A.3, it suffices to show that  $\sigma_x \sigma_x^* = \sigma_y \sigma_y^*$  for all  $x, y \in X$ . By Proposition 3.3,  $P_T$  is a subgroup of the finitely generated free abelian group  $\mathbb{Z}^k$ , and so  $P_T \cong \mathbb{Z}^l$  for some  $l \leq k$ . Fix free abelian generators  $g_1, \dots, g_l$  of  $P_T$ . Since each  $\sigma_x \sigma_x^*$  is a bicharacter (by Theorem A.3), it suffices to show that  $(\sigma_x \sigma_x^*)(g_i, g_j) = (\sigma_y \sigma_y^*)(g_i, g_j)$  for all  $i, j \in \{1, \dots, l\}$  and  $x, y \in X$ . To see this, we first show that  $\sigma_{r(\gamma)} \sigma_{r(\gamma)}^*(g_i, g_j) = \sigma_{s(\gamma)} \sigma_{s(\gamma)}^*(g_i, g_j)$ , for all  $\gamma \in \mathcal{G}_T$  and  $i, j \in \{1, \dots, l\}$ .

Let  $P_T \times_\sigma \mathbb{T} := P_T \times \mathbb{T}$  be the semidirect product group, which is equal to  $P_T \times \mathbb{T}$  as a set, but has group operation

$$(p, w)(q, z) = (p + q, \sigma(p, q) w z).$$

Define  $i_\sigma : \mathbb{T} \rightarrow P_T \times_\sigma \mathbb{T}$  by  $i_\sigma(z) = (0, z)$  and  $q_\sigma : P_T \times_\sigma \mathbb{T} \rightarrow P_T$  by  $q_\sigma(p, z) = p$ . Consider the bijection  $M : H^2(P_T, \mathbb{T}) \rightarrow \text{Ext}(P_T, \mathbb{T})$  of Theorem A.1 that maps the cohomology class of a 2-cocycle  $\sigma \in Z^2(P_T, \mathbb{T})$  to the congruence class of the central extension

$$1 \rightarrow \mathbb{T} \xrightarrow{i_\sigma} P_T \times_\sigma \mathbb{T} \xrightarrow{q_\sigma} P_T \rightarrow 0.$$

Fix  $\gamma \in \mathcal{G}_T$ . We aim to prove that  $\sigma_{s(\gamma)}$  and  $\sigma_{r(\gamma)}$  are cohomologous by showing that their cohomology classes have the same image under  $M$ . So we must find a homomorphism

$$\varphi_\gamma : P_T \times_{\sigma_{s(\gamma)}} \mathbb{T} \rightarrow P_T \times_{\sigma_{r(\gamma)}} \mathbb{T}$$

that makes the diagram

$$\begin{array}{ccc} & P_T \times_{\sigma_{s(\gamma)}} \mathbb{T} & \\ i_{\sigma_{s(\gamma)}} \nearrow & \downarrow \varphi_\gamma & \searrow q_{\sigma_{s(\gamma)}} \\ \mathbb{T} & & P_T \\ i_{\sigma_{r(\gamma)}} \searrow & \downarrow q_{\sigma_{r(\gamma)}} & \nearrow \\ & P_T \times_{\sigma_{r(\gamma)}} \mathbb{T} & \end{array} \quad (4.5)$$

commute. Let  $\tau_\gamma^\sigma : P_T \rightarrow \mathbb{T}$  be the map of Lemma 4.3, and define  $\varphi_\gamma : P_T \times_{\sigma_{s(\gamma)}} \mathbb{T} \rightarrow P_T \times_{\sigma_{r(\gamma)}} \mathbb{T}$  by  $\varphi_\gamma(m, z) := (m, \tau_\gamma^\sigma(m)z)$ . Fix  $(m, z), (n, w) \in P_T \times_{\sigma_{s(\gamma)}} \mathbb{T}$ . Recalling from Lemma 4.3(d) that

$$\tau_\gamma^\sigma(m + n) = \sigma_{r(\gamma)}(m, n) \overline{\sigma_{s(\gamma)}(m, n)} \tau_\gamma^\sigma(m) \tau_\gamma^\sigma(n),$$

we obtain

$$\begin{aligned}
\varphi_\gamma((m, z)(n, w)) &= \varphi_\gamma(m + n, \sigma_{s(\gamma)}(m, n) zw) \\
&= (m + n, \tau_\gamma^\sigma(m + n) \sigma_{s(\gamma)}(m, n) zw) \\
&= (m + n, \sigma_{r(\gamma)}(m, n) \tau_\gamma^\sigma(m) \tau_\gamma^\sigma(n) zw) \\
&= (m, \tau_\gamma^\sigma(m) z)(n, \tau_\gamma^\sigma(n) w) \\
&= \varphi_\gamma(m, z) \varphi_\gamma(n, w),
\end{aligned}$$

and thus  $\varphi_\gamma$  is a homomorphism. Since  $\sigma$  is normalised, the formula (4.1) from Lemma 4.3 gives  $\tau_\gamma^\sigma(0) = 1$ , and it follows that the diagram (4.5) commutes. Therefore,  $\sigma_{r(\gamma)}$  is cohomologous to  $\sigma_{s(\gamma)}$ , and so Theorem A.3 implies that

$$(\sigma_{r(\gamma)} \sigma_{r(\gamma)}^*)(g_i, g_j) = (\sigma_{s(\gamma)} \sigma_{s(\gamma)}^*)(g_i, g_j) \quad \text{for all } i, j \in \{1, \dots, l\}, \quad (4.6)$$

as claimed.

Now fix  $x, y \in X$ . Since  $(X, T)$  is minimal, there is a sequence  $(\gamma_n)_{n \in \mathbb{N}}$  in  $\mathcal{G}_T$  such that  $s(\gamma_n) = x$  for all  $n \in \mathbb{N}$ , and  $r(\gamma_n) \rightarrow y$  as  $n \rightarrow \infty$ . Fix  $i, j \in \{1, \dots, l\}$ . By Lemma 4.3(a), the map  $u \mapsto \sigma_u(g_i, g_j)$  is continuous, and hence the map  $u \mapsto (\sigma_u \sigma_u^*)(g_i, g_j)$  is continuous. So  $(\sigma_y \sigma_y^*)(g_i, g_j) = \lim_{n \rightarrow \infty} (\sigma_{r(\gamma_n)} \sigma_{r(\gamma_n)}^*)(g_i, g_j)$ . Equation (4.6) gives  $(\sigma_{r(\gamma_n)} \sigma_{r(\gamma_n)}^*)(g_i, g_j) = (\sigma_x \sigma_x^*)(g_i, g_j)$  for each  $n \in \mathbb{N}$ , and so  $(\sigma_y \sigma_y^*)(g_i, g_j) = (\sigma_x \sigma_x^*)(g_i, g_j)$ .  $\square$

*Proof of Proposition 4.2.* Lemma 4.4 shows that  $\rho_x$  is a  $\mathbb{T}$ -valued 2-cocycle on  $P_T$  whose cohomology class is independent of  $x$ . So there exists a 2-cocycle  $\omega \in Z^2(P_T, \mathbb{T})$  whose cohomology class agrees with that of each  $\rho_x$ . By Theorem A.6, we may assume that  $\omega$  is a bicharacter that vanishes on  $Z_\omega$  in each coordinate, and that there is a bicharacter  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$  such that

$$\tilde{\omega}(p + Z_\omega, q + Z_\omega) = \omega(p, q) \quad \text{for all } p, q \in P_T.$$

We now construct  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$  such that  $\sigma$  is cohomologous to  $\rho$ , and  $\sigma$  is  $\omega$ -constant on  $\mathcal{I}_T$ . For each  $x \in X$ , the 2-cocycles  $\rho_x$  and  $\omega$  are cohomologous, and so the map  $\tilde{c}_x: P_T \times P_T \rightarrow \mathbb{T}$  defined by

$$\tilde{c}_x(p, q) := \overline{\omega(p, q)} \rho_x(p, q)$$

is a 2-coboundary on  $P_T$ . Since  $P_T$  is a subgroup of  $\mathbb{Z}^k$  (by Proposition 3.3), there is an integer  $l \in \{1, \dots, k\}$  such that  $P_T \cong \mathbb{Z}^l$ . Fix free abelian generators  $g_1, \dots, g_l$  for  $P_T$ . For  $m \in P_T$ , let  $m_1, \dots, m_l$  be the unique integers such that  $m = \sum_{i=1}^l m_i g_i$ . For each  $i \in \{1, \dots, l\}$ , we write  $\langle g_j : j \leq i \rangle$  for the group generated by the set  $\{g_j : 1 \leq j \leq i\}$ . We claim that there are maps  $b_x: P_T \rightarrow \mathbb{T}$ , indexed by  $x \in X$ , such that  $x \mapsto b_x(m)$  is continuous for each  $m \in P_T$ , and for each  $i \in \{1, \dots, l\}$ , we have

$$b_x(m) \overline{b_x(m + g_i)} = \tilde{c}_x(g_i, m), \quad \text{whenever } m \in \langle g_j : j \leq i \rangle. \quad (4.7)$$

To see this, for each  $x \in X$  define  $b_x(0) := 1 \in \mathbb{T}$ . The map  $x \mapsto b_x(0)$  is trivially continuous. Fix  $i \in \{1, \dots, l\}$ . Suppose inductively that the maps  $b_x$  have been defined on  $\langle g_j : j < i \rangle$ , and that  $x \mapsto b_x(m)$  is continuous for each  $m \in \langle g_j : j < i \rangle$ . To extend  $b_x$  to  $\langle g_j : j \leq i \rangle$ , first observe that  $b_x(m)$  is already defined when  $m = \sum_{j=1}^i m_j g_j$  and  $m_i = 0$ . Now suppose inductively that  $b_x(m)$  is defined and  $x \mapsto b_x(m)$  is continuous whenever  $|m_i| \leq a$  for some  $a \in \mathbb{N}$ , and that  $b_x$  satisfies Equation (4.7) whenever  $|m_i|, |m_i + 1| \leq a$ . Fix  $m \in \langle g_j : j \leq i \rangle$  such that  $|m_i| = a + 1$ . Define

$$b_x(m) := \begin{cases} b_x(m - g_i) \overline{\tilde{c}_x(g_i, m - g_i)} & \text{if } m_i > 0 \\ b_x(m + g_i) \tilde{c}_x(g_i, m) & \text{if } m_i < 0. \end{cases}$$

Since Lemma 4.3(a) implies that the maps  $x \mapsto \tilde{c}_x(p, q)$  are continuous for all  $p, q \in P_T$ , the inductive hypothesis guarantees that  $x \mapsto b_x(m)$  is continuous. Moreover, rearranging each of the cases in the definition of  $b_x(m)$  shows that Equation (4.7) is satisfied. So the claim follows by induction.

Recall the coboundary map  $\delta^1$  of Definition 2.2(iv). We claim that  $\delta^1 b_x = \tilde{c}_x$ . To see this, first choose a normalised 1-cochain  $\tilde{b}_x: P_T \rightarrow \mathbb{T}$  such that  $\delta^1 \tilde{b}_x = \tilde{c}_x$ . (This is possible because  $\tilde{c}_x$  is a 2-coboundary on  $P_T$ .) Define  $a_x: P_T \rightarrow \mathbb{T}$  by

$$a_x(m) := \prod_{i=1}^l \overline{\tilde{b}_x(g_i)^{m_i}}.$$

A straightforward calculation shows that  $a_x$  is a 1-cocycle, and so  $\delta^1 a_x$  is trivial. Hence  $\delta^1(a_x \tilde{b}_x) = \delta^1 \tilde{b}_x = \tilde{c}_x$ . Putting  $m = 0$  in Equation (4.7), we see that for each  $i \in \{1, \dots, l\}$ ,  $b_x(g_i) = 1$ . Hence

$$(a_x \tilde{b}_x)(0) = \left( \prod_{i=1}^l \overline{\tilde{b}_x(g_i)^0} \right) \tilde{b}_x(0) = 1 = b_x(0),$$

and for each  $i \in \{1, \dots, l\}$ ,

$$(a_x \tilde{b}_x)(g_i) = \left( \prod_{\substack{j=1, \\ j \neq i}}^l \overline{\tilde{b}_x(g_j)^0} \right) \left( \overline{\tilde{b}_x(g_i)^1} \right) \tilde{b}_x(g_i) = 1 = b_x(g_i).$$

Thus, for all  $i \in \{1, \dots, l\}$  and  $m \in \langle g_j : j \leq i \rangle$ , we have

$$\begin{aligned} (a_x \tilde{b}_x)(m) \overline{(a_x \tilde{b}_x)(m + g_i)} &= (a_x \tilde{b}_x)(g_i) (a_x \tilde{b}_x)(m) \overline{(a_x \tilde{b}_x)(g_i + m)} \\ &= \delta^1(a_x \tilde{b}_x)(g_i, m) \\ &= \tilde{c}_x(g_i, m) \\ &= b_x(m) \overline{b_x(m + g_i)}. \end{aligned}$$

So  $b_x$  and  $a_x \tilde{b}_x$  both map 0 and each generator  $g_i$  to 1, and they also both satisfy Equation (4.7). Hence  $a_x \tilde{b}_x = b_x$ , and thus  $\delta^1 b_x = \delta^1(a_x \tilde{b}_x) = \tilde{c}_x$ , as claimed.

Since the maps  $(x, p, x) \mapsto x$  and  $x \mapsto b_x(p)$  are both continuous for each fixed  $p \in P_T$ , the map  $\tilde{b}: \mathcal{I}_T \rightarrow \mathbb{T}$  given by  $\tilde{b}(x, p, x) := b_x(p)$  is a continuous 1-cochain on  $\mathcal{I}_T$ . We extend  $\tilde{b}$  to a map  $b: \mathcal{G}_T \rightarrow \mathbb{T}$  by setting  $b(\gamma) := 1$  for all  $\gamma \in \mathcal{G}_T \setminus \mathcal{I}_T$ . Since  $\mathcal{I}_T$  is a clopen subset of  $\mathcal{G}_T$  (by Proposition 3.9), this map  $b$  is a continuous 1-cochain on  $\mathcal{G}_T$ . We have  $b(x, 0, x) = b_x(0) = 1$  for all  $x \in X$ , and so  $b$  is normalised. Thus the map  $\delta^1 b: \mathcal{G}_T^{(2)} \rightarrow \mathbb{T}$  given by  $\delta^1 b(\alpha, \beta) := b(\alpha) b(\beta) \overline{b(\alpha\beta)^{-1}}$  is a continuous 2-coboundary on  $\mathcal{G}_T$ . Define  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$  by  $\sigma(\alpha, \beta) := \rho(\alpha, \beta) \overline{\delta^1 b(\alpha, \beta)}$ . Since  $\sigma$  and  $\rho$  differ by the 2-coboundary  $\delta^1 b$ , they are cohomologous, and so [31, Proposition II.1.2] implies that  $C^*(\mathcal{G}_T, \rho) \cong C^*(\mathcal{G}_T, \sigma)$ . Finally, fix  $x \in X$  and  $p, q \in P_T$ . Since  $\delta^1 b_x = \tilde{c}_x = \rho_x \omega$ , we have

$$\sigma((x, p, x), (x, q, x)) = \rho_x(p, q) \overline{\delta^1 b_x(p, q)} = \omega(p, q),$$

and so  $\sigma$  is  $\omega$ -constant on  $\mathcal{I}_T$ .  $\square$

The following result is an extension of [24, Lemma 3.6] to the setting of Deaconu–Renault groupoids.

**Proposition 4.5.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Suppose that  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ , and that  $\omega \in Z^2(P_T, \mathbb{T})$  is a bicharacter that vanishes on  $Z_\omega$  in each coordinate such that  $\sigma$  is  $\omega$ -constant on  $\mathcal{I}_T$ , as in Proposition 4.2. Let  $\gamma \mapsto \tau_\gamma^\sigma$  be the continuous  $\widehat{P}_T$ -valued 1-cocycle on  $\mathcal{G}_T$  defined in*



*Lemma 4.3(e).* For all  $\gamma \in \mathcal{I}_T$  and  $p \in Z_\omega$ , we have  $\tau_\gamma^\sigma(p) = 1$ . Let  $\mathcal{H}_T = \mathcal{G}_T/\mathcal{I}_T$  be the quotient groupoid of [Proposition 3.9](#). There is a continuous  $\widehat{Z}_\omega$ -valued 1-cocycle  $[\gamma] \mapsto \tilde{\tau}_{[\gamma]}^\sigma$  on  $\mathcal{H}_T$  such that  $\tilde{\tau}_{[\gamma]}^\sigma(p) = \tau_\gamma^\sigma(p)$  for all  $\gamma \in \mathcal{G}_T$  and  $p \in Z_\omega$ . There is a continuous action  $\theta$  of  $\mathcal{H}_T$  on  $X \times \widehat{Z}_\omega$  such that

$$\theta_{[\gamma]}(s(\gamma), \chi) = (r(\gamma), \tilde{\tau}_{[\gamma]}^\sigma \chi) \quad \text{for all } \gamma \in \mathcal{G}_T \text{ and } \chi \in \widehat{Z}_\omega.$$

We call the action  $\theta$  of [Proposition 4.5](#) the *spectral action* associated to  $(T, \sigma)$ . We denote the orbit of  $(x, \chi) \in X \times \widehat{Z}_\omega$  under  $\theta$  by  $[x, \chi]_\theta$ .

*Proof of Proposition 4.5.* Fix  $\gamma \in \mathcal{I}_T$ . [Proposition 3.5](#) implies that there exist  $y \in X$  and  $m \in P_T$  such that  $\gamma = (y, m, y)$ . We claim that  $\tau_\gamma^\sigma(Z_\omega) = \{1\}$ . Fix  $p \in Z_\omega$ . Using the formula (4.1) from [Lemma 4.3](#), and that  $\omega$  is a bicharacter satisfying  $\sigma|_{\mathcal{I}_T^{(2)}} = 1_X \times \omega$ , and that  $\omega\omega^*$  is an antisymmetric bicharacter, we see that

$$\begin{aligned} \tau_\gamma^\sigma(p) &= \sigma(\gamma, (y, p, y)) \sigma(\gamma(y, p, y), \gamma^{-1}) \overline{\sigma(\gamma, \gamma^{-1})} \\ &= \sigma((y, m, y), (y, p, y)) \sigma((y, m + p, y), (y, -m, y)) \overline{\sigma((y, m, y), (y, -m, y))} \\ &= \omega(m, p) \omega(m + p, -m) \overline{\omega(m, -m)} \\ &= \omega(m, p) \overline{\omega(p, m)} \\ &= \overline{(\omega\omega^*)(p, m)}, \end{aligned}$$

which is 1 because  $p \in Z_\omega$ . Thus  $\tau_\gamma^\sigma(Z_\omega) = \{1\}$ , as claimed.

For any  $\gamma \in \mathcal{G}_T$ , we have  $\tau_\gamma^\sigma \in \widehat{P}_T$  by [Lemma 4.3\(e\)](#), and so  $\tau_\gamma^\sigma|_{Z_\omega} \in \widehat{Z}_\omega$ . Suppose that  $\alpha, \beta \in \mathcal{G}_T$  satisfy  $[\alpha] = [\beta]$ . Then  $\eta = \beta^{-1}\alpha \in \mathcal{I}_T$  satisfies  $\alpha = \beta\eta$ . For  $p \in Z_\omega$ , we have  $\tau_\eta^\sigma(p) = 1$ , and thus, since  $\gamma \mapsto \tau_\gamma^\sigma$  is a 1-cocycle,

$$\tau_\alpha^\sigma(p) = \tau_{\beta\eta}^\sigma(p) = \tau_\beta^\sigma(p) \tau_\eta^\sigma(p) = \tau_\beta^\sigma(p).$$

Therefore, there is a map  $[\gamma] \mapsto \tilde{\tau}_{[\gamma]}^\sigma$  from  $\mathcal{H}_T$  to  $\widehat{Z}_\omega$  such that  $\tilde{\tau}_{[\gamma]}^\sigma(p) = \tau_\gamma^\sigma(p)$  for all  $\gamma \in \mathcal{G}_T$  and  $p \in Z_\omega$ . For  $\alpha, \beta \in \mathcal{G}_T$  and  $p \in Z_\omega$ ,

$$\tilde{\tau}_{[\alpha][\beta]}^\sigma(p) = \tilde{\tau}_{[\alpha\beta]}^\sigma(p) = \tau_{\alpha\beta}^\sigma(p) = \tau_\alpha^\sigma(p) \tau_\beta^\sigma(p) = \tilde{\tau}_{[\alpha]}^\sigma(p) \tilde{\tau}_{[\beta]}^\sigma(p),$$

and so  $\tilde{\tau}_{[\alpha][\beta]}^\sigma = \tilde{\tau}_{[\alpha]}^\sigma \tilde{\tau}_{[\beta]}^\sigma$ . Thus  $[\gamma] \mapsto \tilde{\tau}_{[\gamma]}^\sigma$  is a  $\widehat{Z}_\omega$ -valued 1-cocycle on  $\mathcal{H}_T$ .

We claim that  $[\gamma] \mapsto \tilde{\tau}_{[\gamma]}^\sigma$  is continuous on  $\mathcal{H}_T$ . Fix a finite subset  $F \subseteq Z_\omega$  and an open subset  $U \subseteq \mathbb{T}$ , so that  $S_{\widehat{Z}_\omega}(F, U) := \{\chi \in \widehat{Z}_\omega : \chi(F) \subseteq U\}$  is a typical subsbasis element for the topology on  $\widehat{Z}_\omega$ . It suffices to show that  $\{[\gamma] \in \mathcal{H}_T : \tilde{\tau}_{[\gamma]}^\sigma \in S_{\widehat{Z}_\omega}(F, U)\}$  is open in  $\mathcal{H}_T$ . Since  $F$  is finite,

$$S_{\widehat{P}_T}(F, U) := \{\chi \in \widehat{P}_T : \chi(F) \subseteq U\}$$

is open in  $\widehat{P}_T$ . By [Lemma 4.3\(e\)](#), the map  $\gamma \mapsto \tau_\gamma^\sigma$  is continuous on  $\mathcal{G}_T$ , and hence  $\{\gamma \in \mathcal{G}_T : \tau_\gamma^\sigma \in S_{\widehat{P}_T}(F, U)\}$  is open in  $\mathcal{G}_T$ . Let  $\pi_T: \mathcal{G}_T \rightarrow \mathcal{H}_T$  denote the quotient map  $\gamma \mapsto [\gamma]$ . Then  $\pi_T^{-1}(\{[\gamma] : \tilde{\tau}_{[\gamma]}^\sigma \in S_{\widehat{Z}_\omega}(F, U)\}) = \{\gamma \in \mathcal{G}_T : \tau_\gamma^\sigma(F) \subseteq U\}$  is open. Thus, by the definition of the quotient topology,  $\{[\gamma] : \tilde{\tau}_{[\gamma]}^\sigma \in S_{\widehat{Z}_\omega}(F, U)\}$  is open in  $\mathcal{H}_T$ .

It remains to show that  $\theta$  is a continuous action of  $\mathcal{H}_T$  on  $X \times \widehat{Z}_\omega$ . For  $\alpha, \beta \in \mathcal{G}_T$  such that  $[\alpha] = [\beta]$ , we have  $\alpha\beta^{-1} \in \mathcal{I}_T$ , and hence  $r(\alpha) = r(\beta)$  and  $s(\alpha) = s(\beta)$ . Define  $R: X \times \widehat{Z}_\omega \rightarrow \mathcal{H}_T^{(0)}$  by  $R(x, \chi) := [x]$ . Then  $R$  is continuous and surjective. Recall from [Definition 2.1](#) that the fibred product  $\mathcal{H}_T \star (X \times \widehat{Z}_\omega)$  is defined by

$$\mathcal{H}_T \star (X \times \widehat{Z}_\omega) = \{([\gamma], (x, \chi)) : x \in X, \chi \in \widehat{Z}_\omega, \gamma \in (\mathcal{G}_T)_x\}.$$

Since  $[\gamma] \mapsto \tilde{\tau}_{[\gamma]}^\sigma$  is a continuous map from  $\mathcal{H}_T$  to  $\widehat{Z}_\omega$  and  $r: \mathcal{H}_T \rightarrow X$  is continuous, the map  $([\gamma], (s(\gamma), \chi)) \mapsto \theta_{[\gamma]}(s(\gamma), \chi) = (r(\gamma), \tilde{\tau}_{[\gamma]}^\sigma \chi)$  from  $\mathcal{H}_T \star (X \times \widehat{Z}_\omega)$  to  $X \times \widehat{Z}_\omega$  is continuous. To see that  $\theta$  is an action, we must show that conditions (A1) and (A2) of Definition 2.1 are satisfied.

For (A1), fix  $(x, \chi) \in X \times \widehat{Z}_\omega$  and  $([\alpha], [\beta]) \in \mathcal{H}_T^{(2)}$  such that  $([\beta], (x, \chi)) \in \mathcal{H}_T \star (X \times \widehat{Z}_\omega)$ . Then  $s([\alpha]) = r([\beta])$ , and  $s([\beta]) = R(x, \chi) = [x]$ . Hence  $s([\alpha][\beta]) = s([\beta]) = R(x, \chi)$ , and so  $([\alpha][\beta], (x, \chi)) \in \mathcal{H}_T \star (X \times \widehat{Z}_\omega)$ . Since  $s(\beta) = x$ ,

$$\theta_{[\beta]}(x, \chi) = \theta_{[\beta]}(s(\beta), \chi) = (r(\beta), \tilde{\tau}_{[\beta]}^\sigma \chi).$$

Thus

$$R(\theta_{[\beta]}(x, \chi)) = R(r(\beta), \tilde{\tau}_{[\beta]}^\sigma \chi) = [r(\beta)] = r([\beta]) = s([\alpha]),$$

and so  $([\alpha], \theta_{[\beta]}(x, \chi)) \in \mathcal{H}_T \star (X \times \widehat{Z}_\omega)$ . Finally, since  $[\gamma] \mapsto \tilde{\tau}_{[\gamma]}^\sigma$  is a  $\widehat{Z}_\omega$ -valued 1-cocycle on  $\mathcal{H}_T$ , we have  $\tilde{\tau}_{[\alpha]}^\sigma \tilde{\tau}_{[\beta]}^\sigma = \tilde{\tau}_{[\alpha][\beta]}^\sigma = \tilde{\tau}_{[\alpha\beta]}^\sigma$ , and hence

$$\begin{aligned} \theta_{[\alpha]}(\theta_{[\beta]}(x, \chi)) &= \theta_{[\alpha]}(\theta_{[\beta]}(s(\beta), \chi)) = \theta_{[\alpha]}(r(\beta), \tilde{\tau}_{[\beta]}^\sigma \chi) = \theta_{[\alpha]}(s(\alpha), \tilde{\tau}_{[\beta]}^\sigma \chi) \\ &= (r(\alpha), \tilde{\tau}_{[\alpha]}^\sigma (\tilde{\tau}_{[\beta]}^\sigma \chi)) = (r(\alpha\beta), \tilde{\tau}_{[\alpha\beta]}^\sigma \chi) = \theta_{[\alpha\beta]}(s(\alpha\beta), \chi) = \theta_{[\alpha][\beta]}(x, \chi). \end{aligned}$$

Thus, (A1) is satisfied.

For (A2), fix  $(x, \chi) \in X \times \widehat{Z}_\omega$ . Then  $s(R(x, \chi)) = s([x]) = [x] = R(x, \chi)$ , and so  $(R(x, \chi), (x, \chi)) \in \mathcal{H}_T \star (X \times \widehat{Z}_\omega)$ . Since  $x \in \mathcal{I}_T$ , we have  $\tau_x^\sigma(Z_\omega) = \{1\}$ . Thus, for all  $p \in Z_\omega$ , we have  $\tilde{\tau}_{[x]}^\sigma(p) = \tau_x^\sigma(p) = 1$ , and so  $\tilde{\tau}_{[x]}^\sigma \chi = \chi$ . Hence

$$\theta_{R(x, \chi)}(x, \chi) = \theta_{[x]}(s(x), \chi) = (r(x), \tilde{\tau}_{[x]}^\sigma \chi) = (x, \chi). \quad \square$$

## 5. REALISING $C^*(\mathcal{I}_T, \sigma)$ AS AN INDUCED ALGEBRA

In this section we realise the twisted C\*-algebra associated to the interior  $\mathcal{I}_T$  of the isotropy of a Deaconu–Renault groupoid  $\mathcal{G}_T$  and a continuous 2-cocycle  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$  as an induced algebra. We then describe the ideals of this induced algebra. We begin by introducing a spanning set  $\mathcal{B}_T$  for  $C_c(\mathcal{G}_T)$  and then giving a tensor-product decomposition of  $C^*(\mathcal{I}_T, \sigma)$ .

**Lemma 5.1.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Let  $c: \mathcal{G}_T \rightarrow \mathbb{Z}^k$  be as in Lemma 2.3. Let*

$$\mathcal{B}_T := \{f \in C_c(\mathcal{G}_T) : \text{supp}(f) \text{ is a bisection contained in } c^{-1}(n), \text{ for some } n \in \mathbb{Z}^k\}.$$

Then  $C_c(\mathcal{G}_T) = \text{span } \mathcal{B}_T$ .

*Proof.* Fix  $f \in C_c(\mathcal{G}_T)$ . Since  $\text{supp}(f)$  is compact, there is a finite set  $\mathcal{F}$  of precompact open bisections that cover  $\text{supp}(f)$ . Since each  $U \in \mathcal{F}$  is precompact, there are only finitely many  $n \in \mathbb{Z}^k$  such that  $U \cap c^{-1}(n) \neq \emptyset$ . Since each  $c^{-1}(n) \cap U$  is open, it is a precompact open bisection, so we can assume that  $c$  is constant on each  $U \in \mathcal{F}$ . Now, as in the proof of [33, Lemma 9.1.3], fix a partition of unity  $\{g_U : U \in \mathcal{F}\}$  on  $\text{supp}(f)$  subordinate to  $\mathcal{F}$ . By the Tietze extension theorem, each  $g_U$  extends to an element  $\tilde{g}_U$  of  $C_c(\mathcal{G}_T)$ . Now the pointwise products  $f_U := \tilde{g}_U \cdot f$  satisfy  $\text{supp}(f_U) \subseteq U$ , and  $\sum_{U \in \mathcal{F}} f_U = f$ .  $\square$

**Lemma 5.2.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. For each  $h \in C_c(X)$  and  $p \in P_T$ , define  $h \cdot 1_p: \mathcal{I}_T \rightarrow \mathbb{C}$  by*

$$(h \cdot 1_p)(x, m, x) := \delta_{p, m} h(x).$$

Then  $h \cdot 1_p \in C_c(\mathcal{I}_T)$  for each  $p \in P_T$ , and  $C_c(\mathcal{I}_T) = \text{span}\{h \cdot 1_p : h \in C_c(X), p \in P_T\}$ .

*Proof.* For each  $h \in C_c(X)$  and  $p \in P_T$ , we have

$$\text{osupp}(h \cdot 1_p) = (\text{osupp}(h) \times \{p\} \times \text{osupp}(h)) \cap \mathcal{I}_T,$$

and hence  $h \cdot 1_p \in C_c(\mathcal{I}_T)$ . Fix  $f \in C_c(\mathcal{I}_T)$ . Since  $\text{supp}(f)$  is compact, there is a finite set  $F \subseteq P_T$  such that  $\text{supp}(f) \subseteq \bigcup_{p \in F} c|_{\mathcal{I}_T}^{-1}(p)$ . For  $p \in F$ , define  $h_p: X \rightarrow \mathbb{C}$  by  $h_p(x) := f(x, p, x)$ . Then  $\text{osupp}(h_p) = r(c|_{\mathcal{I}_T}^{-1}(p) \cap \text{osupp}(f))$ , and hence  $h_p \in C_c(X)$ . Moreover,  $f = \sum_{p \in F} h_p \cdot 1_p$ .  $\square$

**Proposition 5.3.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Suppose that  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ , and that  $\omega \in Z^2(P_T, \mathbb{T})$  is a bicharacter that vanishes on  $Z_\omega$  in each coordinate and satisfies  $\sigma|_{\mathcal{I}_T^{(2)}} = 1_X \times \omega$ , as in Proposition 4.2.*

*Let  $\{u_p : p \in P_T\}$  be the canonical family of generating unitaries for the twisted group  $C^*$ -algebra  $C^*(P_T, \omega)$ . There is an isomorphism  $\Upsilon: C^*(\mathcal{I}_T, \sigma) \rightarrow C_0(X) \otimes C^*(P_T, \omega)$  such that  $\Upsilon(h \cdot 1_p) = h \otimes u_p$  for all  $h \in C_c(X)$  and  $p \in P_T$ .*

*Proof.* The argument used to prove [24, Lemma 4.1] works here—for more detail and an alternative approach to proving injectivity, see [3, Proposition 8.1.3].  $\square$

Before stating the next theorem, we recall the following facts relating to twisted group  $C^*$ -algebras. Define  $B := P_T/Z_\omega$ . There is a right action of  $\widehat{B}$  on  $\widehat{P}_T$  such that

$$(\phi \cdot \chi)(p) = \phi(p) \chi(p + Z_\omega) \quad \text{for all } \phi \in \widehat{P}_T, \chi \in \widehat{B}, \text{ and } p \in P_T.$$

This action induces a continuous, free, proper, right action of  $\widehat{B}$  on  $X \times \widehat{P}_T$  given by  $(x, \phi) \cdot \chi := (x, \phi \cdot \chi)$ . By [14, Theorem 4.40], the map  $\phi \cdot \widehat{B} \mapsto \phi|_{Z_\omega}$  is an isomorphism  $\widehat{P}_T/\widehat{B} \cong \widehat{Z}_\omega$ . Thus  $\phi \mapsto \phi|_{Z_\omega}$  is a quotient map from  $\widehat{P}_T$  to  $\widehat{Z}_\omega$ , and so [12, Theorem 3.3.17] implies that  $Q: (x, \phi) \mapsto (x, \phi|_{Z_\omega})$  is a quotient map from  $X \times \widehat{P}_T$  to  $X \times \widehat{Z}_\omega$ .

Let  $\{U_{p+Z_\omega} : p+Z_\omega \in B\}$  be the canonical family of generating unitaries for the twisted group  $C^*$ -algebra  $C^*(B, \tilde{\omega})$ . By the universal property of  $C^*(B, \tilde{\omega})$ , there is a strongly continuous action  $\beta^B$  of  $\widehat{B}$  on  $C^*(B, \tilde{\omega})$  such that

$$\beta_\chi^B(U_{p+Z_\omega}) = \chi(p + Z_\omega) U_{p+Z_\omega} \quad \text{for all } \chi \in \widehat{B} \text{ and } p \in P_T.$$

(See [3, Theorem 4.3.1] for proofs of the existence of these two actions of  $\widehat{B}$ .) Recall from Definition B.1 the definition of the induced algebra  $\text{Ind}_{\widehat{B}}^{X \times \widehat{P}_T}(C^*(B, \tilde{\omega}), \beta^B)$  associated to the dynamical system  $(C^*(B, \tilde{\omega}), \widehat{B}, \beta^B)$ .

**Theorem 5.4.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ , and let  $\omega \in Z^2(P_T, \mathbb{T})$  and  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$  be bicharacters chosen as in Proposition 4.2. Define*

$$\begin{aligned} \mathcal{X}_T^\omega &:= \text{Ind}_{\widehat{B}}^{X \times \widehat{P}_T}(C^*(B, \tilde{\omega}), \beta^B) \\ &= \left\{ f \in C_0(X \times \widehat{P}_T, C^*(B, \tilde{\omega})) : \begin{array}{l} f(x, \phi \cdot \chi) = (\beta_\chi^B)^{-1}(f(x, \phi)) \text{ for} \\ \text{all } (x, \phi) \in X \times \widehat{P}_T \text{ and } \chi \in \widehat{B} \end{array} \right\}. \end{aligned}$$

*There is an isomorphism  $\psi_T: C^*(\mathcal{I}_T, \sigma) \rightarrow \mathcal{X}_T^\omega$  such that*

$$\psi_T(h \cdot 1_p)(x, \phi) = h(x) \overline{\phi(p)} U_{p+Z_\omega}$$

*for all  $h \in C_c(X)$ ,  $p \in P_T$ , and  $(x, \phi) \in X \times \widehat{P}_T$ .*

*Proof.* Define  $\mathcal{Y}_T^\omega := \text{Ind}_{\widehat{B}}^{\widehat{P}_T}(C^*(B, \tilde{\omega}), \beta^B)$ . Recall from Proposition 5.3 and Theorem B.2 the definitions of the isomorphisms

$$\Upsilon: C^*(\mathcal{I}_T, \sigma) \rightarrow C_0(X) \otimes C^*(P_T, \omega) \quad \text{and} \quad \Omega: C^*(P_T, \omega) \rightarrow \mathcal{Y}_T^\omega.$$

By [30, Propositions B.13 and B.16], there is an isomorphism

$$\Gamma: C_0(X) \otimes C^*(P_T, \omega) \rightarrow C_0(X, \mathcal{Y}_T^\omega)$$

such that  $\Gamma(f \otimes a)(x) = f(x) \Omega(a)$  for all  $f \in C_0(X)$ ,  $a \in C^*(P_T, \omega)$ , and  $x \in X$ . Hence

$$\Gamma(\Upsilon(h \cdot 1_p))(x) = \Gamma(h \otimes u_p)(x) = h(x) \Omega(u_p), \quad (5.1)$$

for all  $h \in C_0(X)$ ,  $p \in P_T$ , and  $x \in X$ . Applications of [30, Propositions B.13, B.15(b), and B.16, and Corollary B.17] show that there is an isomorphism

$$\Lambda: C_0(X, C(\widehat{P}_T, C^*(B, \tilde{\omega}))) \rightarrow C_0(X \times \widehat{P}_T, C^*(B, \tilde{\omega}))$$

given by  $\Lambda(g)(x, \phi) = g(x)(\phi)$ . (See the proof of [3, Proposition 8.2.2] for details.) We claim that for each  $g \in C_0(X, C(\widehat{P}_T, C^*(B, \tilde{\omega})))$ ,

$$\Lambda(g) \in \mathcal{X}_{T, \omega} \quad \text{if and only if} \quad g(x) \in \mathcal{Y}_{T, \omega} \text{ for all } x \in X. \quad (5.2)$$

To see this, fix  $g \in C_0(X, C(\widehat{P}_T, C^*(B, \tilde{\omega})))$ . For all  $x \in X$ ,  $\phi \in \widehat{P}_T$ , and  $\chi \in \widehat{B}$ , we have

$$\Lambda(g)(x, \phi \cdot \chi) = g(x)(\phi \cdot \chi) \quad \text{and} \quad (\beta_\chi^B)^{-1}(\Lambda(g)(x, \phi)) = (\beta_\chi^B)^{-1}(g(x)(\phi)),$$

and hence

$$\Lambda(g)(x, \phi \cdot \chi) = (\beta_\chi^B)^{-1}(\Lambda(g)(x, \phi)) \quad \text{if and only if} \quad g(x)(\phi \cdot \chi) = (\beta_\chi^B)^{-1}(g(x)(\phi)).$$

It is now clear from the definitions of  $\mathcal{X}_{T, \omega}$  and  $\mathcal{Y}_{T, \omega}$  that [Claim \(5.2\)](#) holds. Therefore,  $\Lambda$  restricts to an isomorphism  $\tilde{\Lambda}: C_0(X, \mathcal{Y}_{T, \omega}) \rightarrow \mathcal{X}_{T, \omega}$ , and so

$$\psi_T := \tilde{\Lambda} \circ \Gamma \circ \Upsilon: C^*(\mathcal{I}_T, \sigma) \rightarrow \mathcal{X}_{T, \omega}$$

is an isomorphism. Using [Equation \(5.1\)](#) and the definitions of  $\tilde{\Lambda}$  and  $\Omega$ , we see that for all  $h \in C_c(X)$ ,  $p \in P_T$ , and  $(x, \phi) \in X \times \widehat{P}_T$ ,

$$\psi_T(h \cdot 1_p)(x, \phi) = \Gamma(\Upsilon(h \cdot 1_p))(x)(\phi) = h(x) \Omega(u_p)(\phi) = h(x) \overline{\phi(p)} U_{p+Z_\omega}. \quad \square$$

We now give a useful description of the ideals of the induced algebra  $\mathcal{X}_T^\omega$ .

**Proposition 5.5.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ , and let  $\omega \in Z^2(P_T, \mathbb{T})$  and  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$  be bicharacters chosen as in [Proposition 4.2](#). Define  $\mathcal{X}_T^\omega := \text{Ind}_{\widehat{B}}^{X \times \widehat{P}_T}(C^*(B, \tilde{\omega}), \beta^B)$ . If  $I$  is an ideal of  $\mathcal{X}_T^\omega$ , then*

$$K_I := \{(x, \phi) \in X \times \widehat{P}_T : f(x, \phi) = 0 \text{ for all } f \in I\}$$

is a closed subset of  $X \times \widehat{P}_T$ , and we have

$$I = \{f \in \mathcal{X}_T^\omega : f|_{K_I} \equiv 0\}.$$

In order to prove [Proposition 5.5](#), we need the following lemma.

**Lemma 5.6** ([3, Proposition 8.2.4]). *Let  $G$  be a countable discrete abelian group with identity  $e$ , and let  $\varsigma \in Z^2(G, \mathbb{T})$  be a bicharacter. Let  $\{u_g : g \in G\}$  be the canonical family of generating unitaries for the twisted group C\*-algebra  $C^*(G, \varsigma)$ . Suppose that for all  $g \in G$ , we have  $(\varsigma \varsigma^*)(\{g\} \times G) = \{1\}$  if and only if  $g = e$ . Then  $C^*(G, \varsigma)$  is a simple C\*-algebra with a unique trace  $\tau_e: C^*(G, \varsigma) \rightarrow \mathbb{C}$ , which satisfies  $\tau_e(u_g) = \delta_{g, e}$  for all  $g \in G$ .*

*Proof.* A routine argument shows that there is a state  $\tau_e: C^*(G, \varsigma) \rightarrow \mathbb{C}$  satisfying  $\tau_e(u_g) = \delta_{g,e}$  for all  $g \in G$ . For all  $g, h \in G$ , we have

$$\tau_e(u_g u_h) = \varsigma(g, h) \tau_e(u_{g+h}) = \varsigma(g, h) \delta_{g+h,e} = \varsigma(g, -g) \delta_{h,-g} = \varsigma(-g, g) \delta_{h,-g} = \tau_e(u_h u_g),$$

and it follows from bilinearity and continuity of multiplication that  $\tau_e$  is a trace.

For uniqueness, suppose that  $\tau: C^*(G, \varsigma) \rightarrow \mathbb{C}$  is a trace. Then  $\tau(u_e) = 1 = \tau_e(u_e)$ . We claim that  $\tau(u_g) = 0 = \tau_e(u_g)$  for all  $g \in G \setminus \{e\}$ . To see this, suppose that  $\tau(u_g) \neq 0$  for some  $g \in G$ . We must show that  $g = e$ . Fix  $h \in G$ . Since  $\tau$  is a trace, we have

$$\varsigma(g-h, h) \tau(u_g) = \varsigma(g-h, h) \tau(u_{(g-h)+h}) = \tau(u_{g-h} u_h) = \tau(u_h u_{g-h}) = \varsigma(h, g-h) \tau(u_g).$$

Since  $\tau(u_g) \neq 0$  by assumption, it follows that  $\varsigma(g-h, h) = \varsigma(h, g-h)$ . Thus, since  $\varsigma$  is a bicharacter, we have

$$\varsigma(g, h) \varsigma(-h, h) = \varsigma(g-h, h) = \varsigma(h, g-h) = \varsigma(h, -h) \varsigma(h, g).$$

Since  $\varsigma(-h, h) = \overline{\varsigma(h, h)} = \varsigma(h, -h)$ , we obtain  $\varsigma(g, h) = \varsigma(h, g)$ . Hence  $(\varsigma\varsigma^*)(g, h) = \varsigma(g, h) \varsigma(h, g) = 1$ , and so  $(\varsigma\varsigma^*)(\{g\} \times G) = \{1\}$ . Therefore,  $g = e$  by our hypothesis, and it follows that  $\tau = \tau_e$ .

Now, since  $G$  is abelian and  $C^*(G, \varsigma)$  has a unique trace, [6, Corollary 2.3] implies that  $C^*(G, \varsigma)$  is simple.  $\square$

*Proof of Proposition 5.5.* We have  $K_I = \bigcap_{f \in I} f^{-1}(0)$ , which is closed because each  $f \in I$  is continuous. It is well known that if  $C^*(B, \tilde{\omega})$  is simple, then  $I = \{f \in \mathcal{X}_T^\omega : f|_{K_I} \equiv 0\}$  (see [3, Proposition 4.2.1] for a proof). We will use Lemma 5.6 to show that  $C^*(B, \tilde{\omega})$  is simple. Fix  $p \in P_T$ . Proposition 4.2 implies that for all  $q \in P_T$ , we have

$$(\tilde{\omega}\tilde{\omega}^*)(p + Z_\omega, q + Z_\omega) = \omega(p, q) \overline{\omega(q, p)} = (\omega\omega^*)(p, q). \quad (5.3)$$

By the definition of  $Z_\omega$ , we have  $p \in Z_\omega$  if and only if  $(\omega\omega^*)(p, q) = 1$  for all  $q \in P_T$ . Thus, Equation (5.3) implies that  $p + Z_\omega$  is the identity element of  $B$  if and only if  $(\tilde{\omega}\tilde{\omega}^*)(\{p + Z_\omega\} \times B) = \{1\}$ , and so Lemma 5.6 implies that  $C^*(B, \tilde{\omega})$  is simple.  $\square$

## 6. SIMPLICITY OF TWISTED $C^*$ -ALGEBRAS OF DEACONU–RENAULT GROUPOIDS

In this section we characterise simplicity of twisted  $C^*$ -algebras of Deaconu–Renault groupoids in terms of the underlying data, using the spectral action defined in Proposition 4.5.

**Theorem 6.1.** *Let  $(X, T)$  be a rank- $k$  Deaconu–Renault system such that  $X$  is second-countable. Fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ .*

- (a) *If  $(X, T)$  is not minimal, then  $C^*(\mathcal{G}_T, \sigma)$  is not simple.*
- (b) *Suppose that  $(X, T)$  is minimal. Let  $\omega \in Z^2(P_T, \mathbb{T})$  and  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$  be bicharacters chosen as in Proposition 4.2. Let  $\theta$  be the spectral action associated to  $(T, \sigma)$  as in Proposition 4.5. Then  $C^*(\mathcal{G}_T, \sigma)$  is simple if and only if  $\theta$  is minimal.*

*Proof of Theorem 6.1(a).* This follows from [32, Corollary 4.9] applied to the groupoid dynamical system  $(\mathcal{G}_T, \mathcal{G}_T \times_\sigma \mathbb{T}, C_0(\mathcal{G}_T^{(0)}))$ , but it is easy to provide a short direct proof. Since  $(X, T)$  is not minimal, there exists  $x \in X$  such that  $\overline{[x]}$  is a proper closed invariant set. Let  $\mathcal{H} := \mathcal{G}_T|_{\overline{[x]}} = \{\gamma \in \mathcal{G}_T : s(\gamma) \in \overline{[x]}\}$ , and let  $\tau$  be the restriction of  $\sigma$  to  $\mathcal{H}^{(2)}$ . Then the restriction map  $f \mapsto f|_{\mathcal{H}}$  is a  $*$ -homomorphism from  $C_c(\mathcal{G}_T, \sigma)$  to  $C^*(\mathcal{H}, \tau)$ , and so it extends to a homomorphism  $R: C^*(\mathcal{G}_T, \sigma) \rightarrow C^*(\mathcal{H}, \tau)$ . Since  $\ker(R) \cap C_0(\mathcal{G}_T^{(0)}) = C_0(X \setminus \overline{[x]})$  is neither  $\{0\}$  nor all of  $C_0(\mathcal{G}_T^{(0)})$ , we see that  $\ker(R)$  is a nonzero proper ideal of  $C^*(\mathcal{G}_T, \sigma)$ .  $\square$



To prove part (b) of [Theorem 6.1](#), we need several preliminary results. Let  $\omega \in Z^2(P_T, \mathbb{T})$  and  $\tilde{\omega} \in Z^2(P_T/Z_\omega, \mathbb{T})$  be bicharacters chosen as in [Proposition 4.2](#). Define  $B := P_T/Z_\omega$ , and recall from [Theorem 5.4](#) the definition of the isomorphism

$$\psi_T: C^*(\mathcal{I}_T, \sigma) \rightarrow \mathcal{X}_T^\omega = \text{Ind}_B^{X \times \widehat{P}_T}(C^*(B, \tilde{\omega}), \beta^B).$$

Let  $\iota: C^*(\mathcal{I}_T, \sigma) \rightarrow C^*(\mathcal{G}_T, \sigma)$  be the homomorphism of [\[4, Proposition 6.1\]](#), so

$$\iota(f)(\gamma) = \begin{cases} f(\gamma) & \text{if } \gamma \in \mathcal{I}_T \\ 0 & \text{if } \gamma \notin \mathcal{I}_T \end{cases} \quad \text{for all } f \in C_c(\mathcal{I}_T, \sigma) \text{ and } \gamma \in \mathcal{G}_T.$$

Since  $\mathcal{I}_T$  is amenable (by [\[35, Lemma 3.5\]](#) and [\[1, Proposition 5.1.1\]](#)),  $\iota$  is injective by [\[4, Proposition 6.1\]](#). Define  $M_T^\sigma := \iota(C^*(\mathcal{I}_T, \sigma)) \subseteq C^*(\mathcal{G}_T, \sigma)$ .

We begin by showing that there is a bounded linear map on  $M_T^\sigma$  given by conjugation in  $C^*(\mathcal{G}_T, \sigma)$  by a fixed element of  $C_c(\mathcal{G}_T, [0, 1])$  that is supported on a bisection.

**Lemma 6.2.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Let  $U$  be an open bisection of  $\mathcal{G}_T$ . Suppose that  $g \in C_c(\mathcal{G}_T, [0, 1])$  satisfies  $\text{supp}(g) \subseteq U$ . For all  $f \in C_c(\mathcal{I}_T, \sigma)$ , we have  $g^* \iota(f) g \in \iota(C_c(\mathcal{I}_T, \sigma))$ . There is a linear contraction  $\Xi_g: M_T^\sigma \rightarrow M_T^\sigma$  given by  $\Xi_g(a) := g^* a g$ .*

*Proof.* Fix  $f \in C_c(\mathcal{I}_T, \sigma)$ . Since  $U$  is a bisection containing  $\text{supp}(g)$ , we have

$$\text{supp}(g^* \iota(f) g) \subseteq U^{-1} \mathcal{I}_T U \subseteq \mathcal{I}_T,$$

and hence  $g^* \iota(f) g \in \iota(C_c(\mathcal{I}_T, \sigma))$ . Since  $g$  has range in  $[0, 1]$  and is supported on a bisection,  $\|g\| = \|g\|_\infty \leq 1$ , and thus

$$\|g^* \iota(f) g\| \leq \|g^*\| \|\iota(f)\| \|g\| \leq \|\iota(f)\|.$$

Therefore,  $\iota(f) \mapsto g^* \iota(f) g$  extends to a linear contraction  $\Xi_g: M_T^\sigma \rightarrow M_T^\sigma$ .  $\square$

In the next lemma we introduce a bounded linear map  $\Theta_{U,g}$  on the induced algebra  $\mathcal{X}_{T,\omega}$  that is reminiscent of the spectral action  $\theta$  associated to the pair  $(T, \sigma)$ . This map  $\Theta_{U,g}$  is defined in terms of a fixed element  $g$  of  $C_c(\mathcal{G}_T, [0, 1])$  that is supported on an open bisection  $U$  of  $\mathcal{G}_T$ , and as we show in [Proposition 6.5\(a\)](#), it simply amounts to conjugation of elements of  $M_T^\sigma \cong \mathcal{X}_{T,\omega}$  by  $g$ .

**Lemma 6.3.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Let  $U$  be an open bisection of  $\mathcal{G}_T$ . Suppose that  $g \in C_c(\mathcal{G}_T, \sigma)$  satisfies  $\text{supp}(g) \subseteq U$  and that  $g(U) \subseteq [0, 1]$ . For each  $x \in s(U)$ , let  $\alpha_{U,x}$  denote the unique element of  $U$  with source  $x$ . Let  $\gamma \mapsto \tau_\gamma^\sigma$  be the continuous  $\widehat{P}_T$ -valued 1-cocycle of [Lemma 4.3\(e\)](#). For  $f \in \mathcal{X}_T^\omega$ , define  $\Theta_{U,g}(f): X \times \widehat{P}_T \rightarrow C^*(B, \tilde{\omega})$  by*

$$\Theta_{U,g}(f)(x, \phi) := \begin{cases} |g(\alpha_{U,x})|^2 f(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) & \text{if } x \in s(U) \\ 0 & \text{if } x \notin s(U). \end{cases}$$

Then  $\Theta_{U,g}(f) \in \mathcal{X}_T^\omega$ , and  $\Theta_{U,g}: \mathcal{X}_T^\omega \rightarrow \mathcal{X}_T^\omega$  is a bounded linear map.

*Proof.* Fix  $f \in \mathcal{X}_T^\omega$ . Then

$$f(x, \phi \cdot \chi) = (\beta_\chi^B)^{-1}(f(x, \phi)) \quad \text{for all } (x, \phi) \in X \times \widehat{P}_T \text{ and } \chi \in \widehat{B}. \quad (6.1)$$

We first show that  $\Theta_{U,g}(f) \in C_0(X \times \widehat{P}_T, C^*(B, \tilde{\omega}))$ . The map  $\Theta_{U,g}(f)$  is continuous because  $x \mapsto \alpha_{U,x}$  is continuous. We have  $\text{supp}(\Theta_{U,g}(f)) \subseteq s(\text{supp}(g)) \times \widehat{P}_T$ , and so

$\Theta_{U,g}(f)$  has compact support. Hence  $\Theta_{U,g}(f) \in C_0(X \times \widehat{P}_T, C^*(B, \tilde{\omega}))$ . We must show that  $\Theta_{U,g}(f)$  satisfies Equation (6.1). Fix  $(x, \phi) \in X \times \widehat{P}_T$  and  $\chi \in \widehat{B}$ . If  $x \notin s(U)$ , then

$$\Theta_{U,g}(f)(x, \phi \cdot \chi) = 0 = (\beta_\chi^B)^{-1}(0) = (\beta_\chi^B)^{-1}(\Theta_{U,g}(f)(x, \phi)).$$

Suppose that  $x \in s(U)$ . Since  $f \in \mathcal{X}_T^\omega$ , Equation (6.1) implies that

$$\begin{aligned} \Theta_{U,g}(f)(x, \phi \cdot \chi) &= |g(\alpha_{U,x})|^2 f(r(\alpha_{U,x}), (\tau_{\alpha_{U,x}}^\sigma \phi) \cdot \chi) \\ &= |g(\alpha_{U,x})|^2 (\beta_\chi^B)^{-1}(f(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi)) = (\beta_\chi^B)^{-1}(\Theta_{U,g}(f)(x, \phi)). \end{aligned}$$

Therefore,  $\Theta_{U,g}(f) \in \mathcal{X}_T^\omega$ . Since the range of  $g$  is contained in  $[0, 1]$ , routine calculations show that  $\Theta_{U,g}: \mathcal{X}_T^\omega \rightarrow \mathcal{X}_T^\omega$  is a bounded linear map.  $\square$

In the next lemma we show that the set of functions of the form  $\iota(h \cdot 1_p)$  (as defined in Lemma 5.2) is invariant under conjugation in  $C^*(\mathcal{G}_T, \sigma)$  by a fixed element of  $C_c(\mathcal{G}_T, [0, 1])$  that is supported on a bisection.

**Lemma 6.4.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Let  $U$  be an open bisection of  $\mathcal{G}_T$ . Suppose that  $g \in C_c(\mathcal{G}_T, \sigma)$  satisfies  $\text{supp}(g) \subseteq U$  and that  $g(U) \subseteq [0, 1]$ . For each  $x \in s(U)$ , let  $\alpha_{U,x}$  denote the unique element of  $U$  with source  $x$ . Let  $\Xi_g: M_T^\sigma \rightarrow M_T^\sigma$  and  $\gamma \mapsto \tau_\gamma^\sigma$  be as in Lemmas 6.2 and 4.3(e). For each  $x \in s(U)$ , let  $\alpha_{U,x}$  denote the unique element of  $U$  with source  $x$ . Fix  $h \in C_c(X)$  and  $p \in P_T$ , and define  $H_{g,p}: X \rightarrow \mathbb{C}$  by*

$$H_{g,p}(x) := \begin{cases} |g(\alpha_{U,x})|^2 \overline{\tau_{\alpha_{U,x}}^\sigma(p)} h(r(\alpha_{U,x})) & \text{if } x \in s(U) \\ 0 & \text{if } x \notin s(U). \end{cases}$$

Then  $H_{g,p} \in C_c(X)$ , and we have  $\Xi_g(\iota(h \cdot 1_p)) = \iota(H_{g,p} \cdot 1_p)$ .

*Proof.* Since  $x \mapsto \alpha_{U,x}$  is continuous on  $s(U)$ , and since  $\gamma \mapsto \tau_\gamma^\sigma(p)$  is continuous by Lemma 4.3(b), the map  $H_{g,p}$  is continuous. Since  $\text{supp}(H_{g,p}) \subseteq s(\text{supp}(g))$ , we have  $H_{g,p} \in C_c(X)$ .

By Lemma 6.2, we have  $\Xi_g(\iota(h \cdot 1_p)) \in \iota(C_c(\mathcal{I}_T, \sigma))$ . Thus, for all  $\gamma \in \mathcal{G}_T \setminus \mathcal{I}_T$ , we have

$$\Xi_g(\iota(h \cdot 1_p))(\gamma) = 0 = \iota(H_{g,p} \cdot 1_p)(\gamma).$$

Suppose that  $\gamma \in \mathcal{I}_T$ . Then by Proposition 3.5, there exist  $x \in X$  and  $m \in P_T$  such that  $\gamma = (x, m, x)$ . We have

$$\text{supp}(\Xi_g(\iota(h \cdot 1_p))) \subseteq \text{supp}(g^*) \text{supp}(\iota(h \cdot 1_p)) \text{supp}(g) \subseteq U^{-1} \mathcal{I}_T U.$$

Thus, if  $x \notin s(U)$ , then  $\gamma \notin \text{supp}(\Xi_g(\iota(h \cdot 1_p)))$  and  $H_{g,p}(x) = 0$ , and hence

$$\Xi_g(\iota(h \cdot 1_p))(x, m, x) = 0 = \iota(H_{g,p} \cdot 1_p)(x, m, x).$$

Suppose that  $x \in s(U)$ . Since  $g$  is supported on the bisection  $U$ ,

$$\begin{aligned} \Xi_g(\iota(h \cdot 1_p))(x, m, x) &= \tau_{(\alpha_{U,x})^{-1}}^\sigma(m) g^*(\alpha_{U,x}) \iota(h \cdot 1_p)(r(\alpha_{U,x}), m, r(\alpha_{U,x})) g(\alpha_{U,x}) \\ &= |g(\alpha_{U,x})|^2 \overline{\tau_{\alpha_{U,x}}^\sigma(m)} \delta_{p,m} h(r(\alpha_{U,x})) \\ &= \delta_{p,m} H_{g,p}(x) \\ &= \iota(H_{g,p} \cdot 1_p)(x, m, x). \end{aligned}$$

Therefore,  $\Xi_g(\iota(h \cdot 1_p)) = \iota(H_{g,p} \cdot 1_p)$ .  $\square$

In the following proposition we describe exactly how the map  $\Theta_{U,g}$  defined in Lemma 6.3 relates to the conjugation map  $\Xi_g$  defined in Lemma 6.2. We also show that ideals of  $\mathcal{X}_{T,\omega}$  induced by ideals of  $C^*(\mathcal{G}_T, \sigma)$  are invariant under  $\Theta_{U,g}$ , which is a key result used in the proof of Theorem 6.1(b).

**Proposition 6.5.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Let  $U$  be an open bisection of  $\mathcal{G}_T$ . Suppose that  $g \in C_c(\mathcal{G}_T, [0, 1])$  satisfies  $\text{supp}(g) \subseteq U$ . Recall the definitions of the bounded linear maps  $\Xi_g: M_T^\sigma \rightarrow M_T^\sigma$  from Lemma 6.2 and  $\Theta_{U,g}: \mathcal{X}_T^\omega \rightarrow \mathcal{X}_T^\omega$  from Lemma 6.3.*

(a) *For all  $a \in C^*(\mathcal{I}_T, \sigma)$ , we have*

$$\Theta_{U,g}(\psi_T(a)) = \psi_T(\iota^{-1}(\Xi_g(\iota(a)))).$$

(b) *Suppose that  $I$  is an ideal of  $C^*(\mathcal{G}_T, \sigma)$ , and that  $J$  is an ideal of  $C^*(\mathcal{I}_T, \sigma)$  such that  $\iota(J) = I \cap M_T^\sigma$ . Then the ideal  $\psi_T(J)$  is invariant under  $\Theta_{U,g}$ .*

*Proof.* For part (a), fix  $h \in C_c(X)$  and  $p \in P_T$ . Since all the maps involved are bounded and linear, Lemma 5.2 implies that it suffices to show that

$$\Theta_{U,g}(\psi_T(h \cdot 1_p)) = \psi_T(\iota^{-1}(\Xi_g(\iota(h \cdot 1_p)))).$$

Recall from Lemma 6.4 that there is a function  $H_{g,p} \in C_c(X)$  given by

$$H_{g,p}(x) = \begin{cases} |g(\alpha_{U,x})|^2 \overline{\tau_{\alpha_{U,x}}^\sigma(p)} h(r(\alpha_{U,x})) & \text{if } x \in s(U) \\ 0 & \text{if } x \notin s(U), \end{cases}$$

which satisfies  $\Xi_g(\iota(h \cdot 1_p)) = \iota(H_{g,p} \cdot 1_p)$ . Thus, for all  $(x, \phi) \in X \times P_T$ , we have

$$\begin{aligned} \Theta_{U,g}(\psi_T(h \cdot 1_p))(x, \phi) &= \begin{cases} |g(\alpha_{U,x})|^2 \psi_T(h \cdot 1_p)(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) & \text{if } x \in s(U) \\ 0 & \text{if } x \notin s(U) \end{cases} \\ &= \begin{cases} |g(\alpha_{U,x})|^2 h(r(\alpha_{U,x})) \overline{\tau_{\alpha_{U,x}}^\sigma(p)} \overline{\phi(p)} U_{p+Z_\omega} & \text{if } x \in s(U) \\ 0 & \text{if } x \notin s(U) \end{cases} \\ &= H_{g,p}(x) \overline{\phi(p)} U_{p+Z_\omega} \\ &= \psi_T(H_{g,p} \cdot 1_p)(x, \phi) \\ &= \psi_T(\iota^{-1}(\Xi_g(\iota(h \cdot 1_p))))(x, \phi). \end{aligned}$$

For part (b), fix  $a \in J$ . Then  $\iota(a) \in I \cap M_T^\sigma$ . Since  $I$  is an ideal of  $C^*(\mathcal{G}_T, \sigma)$  and the range of  $\Xi_g$  is contained in  $M_T^\sigma$ , we have  $\Xi_g(\iota(a)) = g^* \iota(a) g \in I \cap M_T^\sigma = \iota(J)$ , and so  $\iota^{-1}(\Xi_g(\iota(a))) \in J$ . Hence part (a) implies that

$$\Theta_{U,g}(\psi_T(a)) = \psi_T(\iota^{-1}(\Xi_g(\iota(a)))) \in \psi_T(J),$$

and thus  $\Theta_{U,g}(\psi_T(J)) \subseteq \psi_T(J)$ .  $\square$

We now use Proposition 6.5(b) to show that the closed subsets of  $X \times \widehat{P}_T$  characterising the ideals of the induced algebra  $\mathcal{X}_{T,\omega}$  are invariant under the spectral action  $\theta$  associated to the pair  $(T, \sigma)$ .

**Proposition 6.6.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Suppose that  $I$  is an ideal of  $C^*(\mathcal{G}_T, \sigma)$ , and that  $J$  is an ideal of  $C^*(\mathcal{I}_T, \sigma)$  such that  $\iota(J) = I \cap M_T^\sigma$ . Define*

$$K_J := \{(x, \phi) \in X \times \widehat{P}_T : f(x, \phi) = 0 \text{ for all } f \in \psi_T(J)\}.$$

*Let  $Q: X \times \widehat{P}_T \rightarrow X \times \widehat{Z}_\omega$  be the quotient map  $(x, \phi) \mapsto (x, \phi|_{Z_\omega})$ . Then  $Q^{-1}(Q(K_J)) = K_J$ , and  $Q(K_J)$  is closed and invariant under the spectral action  $\theta$  of Proposition 4.5.*

*Proof.* We trivially have  $K_J \subseteq Q^{-1}(Q(K_J))$ . We must show that  $Q^{-1}(Q(K_J)) \subseteq K_J$ . Fix  $(x, \phi) \in Q^{-1}(Q(K_J))$ . Then  $(x, \phi|_{Z_\omega}) = Q(x, \phi) \in Q(K_J)$ , and so there exists  $(y, \rho) \in K_J$  such that  $(x, \phi|_{Z_\omega}) = Q(y, \rho) = (y, \rho|_{Z_\omega})$ . We have  $x = y$  and  $\phi|_{Z_\omega} = \rho|_{Z_\omega}$ , and hence [14, Theorem 4.40] implies that  $\phi \cdot \widehat{B} = \rho \cdot \widehat{B}$ . So there exists  $\chi \in \widehat{B}$  such that  $\phi \cdot \chi = \rho \cdot 1_{\widehat{B}} = \rho$ .

Since  $(x, \phi \cdot \chi) = (y, \rho) \in K_J$ , we have  $f(x, \phi \cdot \chi) = 0$  for all  $f \in \psi_T(J)$ . Thus, since  $\psi_T(J) \subseteq \mathcal{X}_T^\omega$ , we have  $f(x, \phi) = \beta_\chi^B(f(x, \phi \cdot \chi)) = 0$  for all  $f \in \psi_T(J)$ . Hence  $(x, \phi) \in K_J$ , and so  $Q^{-1}(Q(K_J)) = K_J$ . Since  $Q$  is a quotient map, [12, Proposition 2.4.3] implies that  $C \subseteq X \times \widehat{Z}_\omega$  is closed if and only if  $Q^{-1}(C) \subseteq X \times \widehat{P}_T$  is closed. Since  $Q^{-1}(Q(K_J)) = K_J$  is closed in  $X \times \widehat{P}_T$  (by Proposition 5.5), we deduce that  $Q(K_J)$  is closed.

We now show that  $Q(K_J)$  is invariant under  $\theta$ . Fix  $(x, \zeta) \in Q(K_J)$  and  $\gamma \in (\mathcal{G}_T)_x$ . Then there exists  $\phi \in \widehat{P}_T$  such that  $(x, \phi) \in K_J$  and  $\phi|_{Z_\omega} = \zeta$ . We must show that  $\theta_{[\gamma]}(x, \zeta) \in Q(K_J)$ . Proposition 4.5 implies that  $\tau_\gamma^\sigma|_{Z_\omega} = \tilde{\tau}_{[\gamma]}^\sigma$ , and so

$$\theta_{[\gamma]}(x, \zeta) = (r(\gamma), \tilde{\tau}_{[\gamma]}^\sigma \zeta) = Q(r(\gamma), \tau_\gamma^\sigma \phi).$$

Hence it suffices to show that  $(r(\gamma), \tau_\gamma^\sigma \phi) \in K_J$ . Fix  $f \in \psi_T(J)$ . We must show that  $f(r(\gamma), \tau_\gamma^\sigma \phi) = 0$ . Let  $U \subseteq \mathcal{G}_T$  be an open bisection containing  $\gamma$ . By Urysohn's lemma there exists  $g \in C_c(\mathcal{G}_T, [0, 1])$  such that  $\text{supp}(g) \subseteq U$  and  $g(\gamma) = 1$ . Let  $\Theta_{U,g}: \mathcal{X}_T^\omega \rightarrow \mathcal{X}_T^\omega$  be as in Lemma 6.3. Since  $s|_U^{-1}(x) = \gamma$  and  $g(\gamma) = 1$ ,

$$\Theta_{U,g}(f)(x, \phi) = f(r(\gamma), \tau_\gamma^\sigma \phi). \quad (6.2)$$

Since  $f \in \psi_T(J)$ , Proposition 6.5(b) implies that  $\Theta_{U,g}(f) \in \psi_T(J)$ . Since  $\psi_T(J)$  is an ideal of  $\mathcal{X}_T^\omega$ , Proposition 5.5 implies that

$$\psi_T(J) = \{f \in \mathcal{X}_T^\omega : f|_{K_J} \equiv 0\}.$$

Thus, since  $(x, \phi) \in K_J$  and  $\Theta_{U,g}(f) \in \psi_T(J)$ , we have

$$\Theta_{U,g}(f)(x, \phi) = 0. \quad (6.3)$$

Together, Equations (6.2) and (6.3) imply that  $f(r(\gamma), \tau_\gamma^\sigma \phi) = 0$ , as required.  $\square$

We now prove several technical results that we use in the proof of Theorem 6.1(b) to show that when the spectral action  $\theta$  is not minimal, the twisted groupoid C\*-algebra  $C^*(\mathcal{G}_T, \sigma)$  is not simple. We first show that, given an element  $(x, \phi) \in X \times \widehat{P}_T$  with non-dense orbit under  $\theta$ , there is a nonzero element of  $\mathcal{X}_{T,\omega} \cong M_T^\sigma$  that is supported off the orbit of  $(x, \phi)$ .

**Lemma 6.7.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Let  $Q: X \times \widehat{P}_T \rightarrow X \times \widehat{Z}_\omega$  be the quotient map  $(x, \phi) \mapsto (x, \phi|_{Z_\omega})$ . Suppose that  $(x, \phi) \in X \times \widehat{P}_T$  satisfies  $\overline{[x, \phi|_{Z_\omega}]_\theta} \neq X \times \widehat{Z}_\omega$ . Then  $Q^{-1}(\overline{[x, \phi|_{Z_\omega}]_\theta})$  is a proper closed subset of  $X \times \widehat{P}_T$ , and there exists  $f \in M_T^\sigma \setminus \{0\}$  such that*

$$\text{supp}((\psi_T \circ \iota^{-1})(f)) \subseteq (X \times \widehat{P}_T) \setminus Q^{-1}(\overline{[x, \phi|_{Z_\omega}]_\theta}).$$

*Proof.* Let  $C_{(x,\phi)} := Q^{-1}(\overline{[x, \phi|_{Z_\omega}]_\theta})$ . Since  $\overline{[x, \phi|_{Z_\omega}]_\theta} \neq X \times \widehat{Z}_\omega$  and  $Q$  is surjective,  $C_{(x,\phi)} \neq X \times \widehat{P}_T$ . Since  $Q$  is continuous,  $C_{(x,\phi)}$  is closed. By Urysohn's lemma there exists  $h \in C_c(X \times \widehat{P}_T, [0, 1]) \setminus \{0\}$  such that  $\text{supp}(h) \subseteq (X \times \widehat{P}_T) \setminus C_{(x,\phi)}$ . Define  $g: X \times \widehat{P}_T \rightarrow C^*(B, \tilde{\omega})$  by

$$g(y, \rho) := \int_{\widehat{B}} h(y, \rho \cdot \chi) \beta_\chi^B(U_{0+Z_\omega}) d\chi = \int_{\widehat{B}} h(y, \rho \cdot \chi) U_{0+Z_\omega} d\chi.$$

By [30, Lemma 6.17], we have  $g \in \mathcal{X}_T^\omega$ . Since  $h \neq 0$  and  $h(y, \rho) \geq 0$  for all  $(y, \rho) \in \text{supp}(h)$ , we have  $g \neq 0$ . We claim that  $\text{supp}(g) \subseteq (X \times \widehat{P}_T) \setminus C_{(x,\phi)}$ . Fix  $(y, \rho) \in C_{(x,\phi)}$ . Then  $Q(y, \rho) \in \overline{[x, \phi|_{Z_\omega}]_\theta}$ . It suffices to show that  $g(y, \rho) = 0$ . Fix  $\chi \in \widehat{B}$ . For all  $m \in Z_\omega$ , we have  $\chi(m + Z_\omega) = 1$ , and hence  $(\rho \cdot \chi)(m) = \rho(m) \chi(m + Z_\omega) = \rho(m)$ . Thus

$$Q(y, \rho \cdot \chi) = (y, (\rho \cdot \chi)|_{Z_\omega}) = (y, \rho|_{Z_\omega}) = Q(y, \rho) \in \overline{[x, \phi|_{Z_\omega}]_\theta},$$

and hence  $(y, \rho \cdot \chi) \in C_{(x, \phi)}$ . Since  $\text{supp}(h) \subseteq (X \times \widehat{P}_T) \setminus C_{(x, \phi)}$ , we have  $h(y, \rho \cdot \chi) = 0$  for all  $\chi \in \widehat{B}$ , and therefore,

$$g(y, \rho) = \int_{\widehat{B}} h(y, \rho \cdot \chi) U_{0+Z_\omega} d\chi = 0.$$

Define  $f := (\iota \circ \psi_T^{-1})(g) \in M_T^\sigma$ . Since  $g \neq 0$  and  $\iota \circ \psi_T^{-1}$  is injective, we have  $f \neq 0$ . Since  $(\psi_T \circ \iota^{-1})(f) = g$ , we have

$$\text{supp}((\psi_T \circ \iota^{-1})(f)) = \text{supp}(g) \subseteq (X \times \widehat{P}_T) \setminus C_{(x, \phi)}. \quad \square$$

Recall from [4, Lemma 6.3(b)] that since  $\mathcal{I}_T$  is closed in  $\mathcal{G}_T$  (by Proposition 3.9) and amenable, there is a conditional expectation  $\Phi: C^*(\mathcal{G}_T, \sigma) \rightarrow M_T^\sigma$  satisfying  $\Phi \circ \iota = \iota$  and  $\Phi(f) = \iota(f|_{\mathcal{I}_T})$  for all  $f \in C_c(\mathcal{G}_T, \sigma)$ .

**Lemma 6.8.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Recall from Lemma 5.1 the definition of the spanning set  $\mathcal{B}_T$  for  $C_c(\mathcal{G}_T, \sigma)$ . Given  $a, b \in \mathcal{B}_T$  and  $f \in M_T^\sigma$ , there exist  $p, q, g \in \mathcal{B}_T$  such that  $gq^*, pg^* \in \iota(C_c(\mathcal{I}_T, \sigma))$ , the range of  $g$  is contained in  $[0, 1]$ , and the map  $\Xi_g$  of Lemma 6.2 satisfies  $\Phi(b^*fa) = \Xi_g(gq^*fpq^*)$ .*

*Proof.* Define  $U := \text{osupp}(a)$  and  $V := \text{osupp}(b)$ . Since  $a, b \in \mathcal{B}_T$ , both  $\overline{U}$  and  $\overline{V}$  are compact bisections, and there exist  $m, n \in \mathbb{Z}^k$  such that  $\overline{U} \subseteq c^{-1}(m)$  and  $\overline{V} \subseteq c^{-1}(n)$ . Define  $X := \mathcal{I}_T V \cap U$  and  $Y := \mathcal{I}_T U \cap V$ . Define  $p, q: \mathcal{G}_T \rightarrow \mathbb{C}$  by

$$p(\gamma) := \begin{cases} a(\gamma) & \text{if } \gamma \in \overline{X} \\ 0 & \text{if } \gamma \notin \overline{X}, \end{cases} \quad \text{and} \quad q(\gamma) := \begin{cases} b(\gamma) & \text{if } \gamma \in \overline{Y} \\ 0 & \text{if } \gamma \notin \overline{Y}. \end{cases}$$

Since  $\text{supp}(p) \subseteq \overline{X} \subseteq \overline{U}$  and  $\text{supp}(q) \subseteq \overline{Y} \subseteq \overline{V}$ , we have  $p, q \in \mathcal{B}_T$ . Let  $W$  be an open bisection of  $\mathcal{G}_T$  such that  $\overline{V} \subseteq W \subseteq c^{-1}(n)$ . By Urysohn's lemma there exists  $g \in C_c(\mathcal{G}_T, [0, 1])$  such that  $\text{supp}(g) \subseteq W$  and  $g|_{\overline{V}} \equiv 1$ . Then  $g \in \mathcal{B}_T$ , and  $gq^*, pg^* \in C_c(\mathcal{G}_T, \sigma)$ . We claim that  $gq^*, pg^* \in \iota(C_c(\mathcal{I}_T, \sigma))$ . To see this, it suffices to show that  $\text{osupp}(gq^*) \cup \text{osupp}(pg^*) \subseteq \mathcal{I}_T$ . Since  $q$  and  $g$  are supported on bisections, we have

$$\text{osupp}(gq^*) = \text{osupp}(g)(\text{osupp}(q))^{-1} \subseteq WV^{-1} \subseteq WW^{-1} = r(W) \subseteq \mathcal{I}_T.$$

By Lemma 3.10,  $\overline{X} = \overline{\mathcal{I}_T V \cap U} \subseteq \mathcal{I}_T \overline{V} \subseteq \mathcal{I}_T W$ , and since  $p$  and  $g$  are supported on bisections, we deduce that

$$\text{osupp}(pg^*) = \text{osupp}(p)(\text{osupp}(g))^{-1} \subseteq \overline{X}W^{-1} \subseteq \mathcal{I}_T WW^{-1} = \mathcal{I}_T r(W) \subseteq \mathcal{I}_T.$$

Therefore,  $gq^*, pg^* \in \iota(C_c(\mathcal{I}_T, \sigma))$ , and Lemma 6.2 implies that

$$\text{osupp}(\Xi_g(gq^*fpq^*)) \subseteq \mathcal{I}_T. \quad (6.4)$$

We conclude by showing that  $\Phi(b^*fa) = \Xi_g(gq^*fpq^*)$ . Since  $\iota, \Phi$ , and  $\Xi_g$  are bounded linear maps, Lemma 5.2 implies that it suffices to consider  $f = \iota(h \cdot 1_p)$  for some  $h \in C_c(X)$  and  $p \in P_T$ . Define  $D := \text{osupp}(f) \subseteq c|_{\mathcal{I}_T}^{-1}(p)$ . Then  $\text{osupp}(g^*g) \subseteq s(W)$ , and so

$$\text{osupp}(\Xi_g(gq^*fpq^*)) = \text{osupp}(g^*gq^*fpq^*g) \subseteq s(W)(V^{-1}DU)s(W) \subseteq V^{-1}DU. \quad (6.5)$$

Together, Equations (6.4) and (6.5) imply that

$$\text{osupp}(\Xi_g(gq^*fpq^*)) \subseteq (V^{-1}DU) \cap \mathcal{I}_T = \text{osupp}(b^*fa) \cap \mathcal{I}_T = \text{osupp}(\Phi(b^*fa)).$$

Thus, if  $\Phi(b^*fa)(\gamma) = 0$  for some  $\gamma \in \mathcal{G}_T$ , then  $\Xi_g(gq^*fpq^*)(\gamma) = 0$ . Suppose that  $\gamma \in \mathcal{G}_T$  satisfies  $\Phi(b^*fa)(\gamma) \neq 0$ . Then  $\gamma \in \mathcal{I}_T$ , and Equation (6.5) implies that there exist  $\alpha \in U$ ,  $\beta \in V$ , and  $\xi \in D \subseteq \mathcal{I}_T$  such that  $\gamma = \beta^{-1}\xi\alpha \in \mathcal{I}_T$ . A routine calculation gives

$$\Phi(b^*fa)(\gamma) = (b^*fa)(\beta^{-1}\xi\alpha) = \sigma(\beta^{-1}\xi, \alpha) \sigma(\beta^{-1}, \xi) \overline{\sigma(\beta^{-1}, \beta)} \overline{b(\beta)} f(\xi) a(\alpha). \quad (6.6)$$



Define  $y := s(\gamma)$ . Since  $\gamma \in \mathcal{I}_T$ , we have  $s(\beta) = r(\gamma) = y$ . Since  $\beta \in V$  and  $g|_V \equiv 1$ ,

$$(g^*g)(y) = (g^*g)(s(\beta)) = |g(\beta)|^2 = 1. \quad (6.7)$$

A routine calculation using Equation (6.7) and that  $\sigma$  is normalised gives

$$\begin{aligned} \Xi_g(gq^*fpg^*)(\gamma) &= (g^*gq^*fpg^*g)(y\gamma y) \\ &= \sigma(y, \gamma y) \sigma(\gamma, y) (g^*g)(y) (q^*fp)(\gamma) (g^*g)(y) \\ &= (q^*fp)(\beta^{-1}\xi\alpha) \\ &= \sigma(\beta^{-1}\xi, \alpha) \sigma(\beta^{-1}, \xi) \overline{\sigma(\beta^{-1}, \beta)} \overline{q(\beta)} f(\xi) p(\alpha). \end{aligned} \quad (6.8)$$

We claim that  $p(\alpha) = a(\alpha)$  and  $q(\beta) = b(\beta)$ . Since  $\gamma \in (V^{-1}\mathcal{I}_T U) \cap \mathcal{I}_T$ , Lemma 3.11 implies that

$$y = s(\gamma) \in s(\mathcal{I}_T U \cap V) = s(\mathcal{I}_T V \cap U).$$

So there exist  $\eta \in \mathcal{I}_T V \cap U = X \subseteq U$  and  $\zeta \in \mathcal{I}_T U \cap V = Y \subseteq V$  such that  $s(\eta) = y = s(\zeta)$ . Since  $s|_U$  and  $s|_V$  are homeomorphisms onto their ranges and

$$s(\eta) = s(\alpha) = y = s(\beta) = s(\zeta),$$

we deduce that  $\alpha = \eta \in X$  and  $\beta = \zeta \in Y$ . Hence  $p(\alpha) = a(\alpha)$  and  $q(\beta) = b(\beta)$ . Together, Equations (6.6) and (6.8) now give

$$\Phi(b^*fa)(\gamma) = \sigma(\beta^{-1}\xi, \alpha) \sigma(\beta^{-1}, \xi) \overline{\sigma(\beta^{-1}, \beta)} \overline{b(\beta)} f(\xi) a(\alpha) = \Xi_g(gq^*fpg^*)(\gamma). \quad \square$$

**Proposition 6.9.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Let  $Q: X \times \widehat{P}_T \rightarrow X \times \widehat{Z}_\omega$  be the quotient map  $(x, \phi) \mapsto (x, \phi|_{Z_\omega})$ , and let  $\Phi: C^*(\mathcal{G}_T, \sigma) \rightarrow M_T^\sigma$  be the conditional expectation of [4, Lemma 6.3(b)] that extends restriction of functions to  $\mathcal{I}_T$ . Fix  $(x, \phi) \in X \times \widehat{P}_T$ . Suppose that  $f \in M_T^\sigma$  satisfies*

$$\text{supp}((\psi_T \circ \iota^{-1})(f)) \subseteq (X \times \widehat{P}_T) \setminus Q^{-1}(\overline{[x, \phi|_{Z_\omega}]_\theta}).$$

Then for all  $a, b \in C^*(\mathcal{G}_T, \sigma)$ , we have

$$(\psi_T \circ \iota^{-1} \circ \Phi)(b^*fa)(x, \phi) = 0.$$

*Proof.* Let  $\text{ev}_{(x, \phi)}: \mathcal{X}_T^\omega \rightarrow C^*(B, \tilde{\omega})$  denote the evaluation map  $f \mapsto f(x, \phi)$ . Recall from Lemma 5.1 the definition of the spanning set  $\mathcal{B}_T$  for  $C_c(\mathcal{G}_T, \sigma)$ . Let  $C_{(x, \phi)} := Q^{-1}(\overline{[x, \phi|_{Z_\omega}]_\theta})$ . Fix  $a, b \in \mathcal{B}_T$ , and suppose that  $f \in M_T^\sigma$  satisfies

$$\text{supp}((\psi_T \circ \iota^{-1})(f)) \subseteq (X \times \widehat{P}_T) \setminus C_{(x, \phi)}.$$

Since  $\text{ev}_{(x, \phi)}$ ,  $\psi_T$ ,  $\iota^{-1}$ , and  $\Phi$  are all bounded linear maps, it suffices to show that

$$(\text{ev}_{(x, \phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi)(b^*fa) = 0.$$

Let  $\Xi_g$  be the bounded linear map defined in Lemma 6.2. By Lemma 6.8 there exist  $p, q, g \in \mathcal{B}_T$  such that  $gq^*, pg^* \in \iota(C_c(\mathcal{I}_T, \sigma))$ , the range of  $g$  is contained in  $[0, 1]$ , and

$$\Phi(b^*fa) = \Xi_g(gq^*fpg^*). \quad (6.9)$$

Let  $U$  be an open bisection of  $\mathcal{G}_T$  containing  $\text{supp}(g)$ . For  $y \in s(U)$ , let  $\alpha_{U, y}$  denote the unique element of  $U$  with source  $y$ . Define  $h_q := \psi_T(\iota^{-1}(gq^*))$  and  $h_p := \psi_T(\iota^{-1}(pg^*))$ . Then

$$\psi_T(\iota^{-1}(gq^*fpg^*)) = h_q \psi_T(\iota^{-1}(f)) h_p. \quad (6.10)$$

By Proposition 6.5(a),

$$\psi_T(\iota^{-1}(\Xi_g(gq^*fpg^*))) = \Theta_{U, g}(\psi_T(\iota^{-1}(gq^*fpg^*))). \quad (6.11)$$

Together, [Equations \(6.9\)](#) and [\(6.11\)](#) imply that

$$\begin{aligned}
& (\text{ev}_{(x,\phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi)(b^*fa) \\
&= \psi_T(\iota^{-1}(\Xi_g(gq^*fpq^*)))(x, \phi) \\
&= \Theta_{U,g}(\psi_T(\iota^{-1}(gq^*fpq^*)))(x, \phi) \\
&= \begin{cases} |g(\alpha_{U,x})|^2 \psi_T(\iota^{-1}(gq^*fpq^*))(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) & \text{if } x \in s(U) \\ 0 & \text{if } x \notin s(U). \end{cases}
\end{aligned}$$

Thus, to see that  $(\text{ev}_{(x,\phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi)(b^*fa) = 0$ , it suffices to show that if  $x \in s(U)$ , then

$$\psi_T(\iota^{-1}(gq^*fpq^*))(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) = 0.$$

If  $x \in s(U)$ , then

$$Q(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) = (r(\alpha_{U,x}), \tilde{\tau}_{[\alpha_{U,x}]}^\sigma \phi|_{Z_\omega}) = \theta_{[\alpha_{U,x}]}(x, \phi|_{Z_\omega}),$$

and hence

$$(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) \in Q^{-1}(\theta_{[\alpha_{U,x}]}(x, \phi|_{Z_\omega})) \subseteq C_{(x,\phi)}.$$

Since  $\text{supp}((\psi_T \circ \iota^{-1})(f)) \subseteq (X \times \widehat{P}_T) \setminus C_{(x,\phi)}$ , we obtain  $\psi_T(\iota^{-1}(f))(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) = 0$ . Combining this with [Equation \(6.10\)](#) gives

$$\psi_T(\iota^{-1}(gq^*fpq^*))(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) = (h_q \psi_T(\iota^{-1}(f)) h_p)(r(\alpha_{U,x}), \tau_{\alpha_{U,x}}^\sigma \phi) = 0. \quad \square$$

We now construct a state  $\kappa_{(x,\phi)}$  of  $C^*(\mathcal{G}_T, \sigma)$  defined in terms of a fixed element  $(x, \phi) \in X \times \widehat{P}_T$ . In the proof of [Theorem 6.1\(b\)](#), we show that if  $(x, \phi)$  has non-dense orbit under  $\theta$ , then the GNS representation associated to  $\kappa_{(x,\phi)}$  is nonzero and has nontrivial kernel, and thus  $C^*(\mathcal{G}_T, \sigma)$  is simple.

**Lemma 6.10.** *Let  $(X, T)$  be a minimal rank- $k$  Deaconu–Renault system such that  $X$  is second-countable, and fix  $\sigma \in Z^2(\mathcal{G}_T, \mathbb{T})$ . Fix  $(x, \phi) \in X \times \widehat{P}_T$ . Let  $\Phi: C^*(\mathcal{G}_T, \sigma) \rightarrow M_T^\sigma$  be the conditional expectation of [[4](#), Lemma 6.3(b)] that extends restriction of functions to  $\mathcal{I}_T$ , and let  $\text{ev}_{(x,\phi)}: \mathcal{X}_T^\omega \rightarrow C^*(B, \tilde{\omega})$  be the evaluation map  $f \mapsto f(x, \phi)$ . Let  $\text{Tr}_T^\omega$  denote the canonical trace on  $C^*(B, \tilde{\omega})$  (as defined in [Lemma 5.6](#)). Let*

$$\kappa_{(x,\phi)} := \text{Tr}_T^\omega \circ \text{ev}_{(x,\phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi: C^*(\mathcal{G}_T, \sigma) \rightarrow \mathbb{C}.$$

For all  $h \in C_c(X)$  such that  $h(x) = 1$ , we have  $\kappa_{(x,\phi)}(\iota(h \cdot 1_0)) = 1$ . Moreover,  $\kappa_{(x,\phi)}$  is a state of  $C^*(\mathcal{G}_T, \sigma)$ .

*Proof.* Suppose that  $h \in C_c(X)$  satisfies  $h(x) = 1$ . Since  $\Phi \circ \iota = \iota$ , we have

$$(\text{ev}_{(x,\phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi)(\iota(h \cdot 1_0)) = \psi_T(h \cdot 1_0)(x, \phi) = h(x) \overline{\phi(0)} U_{0+Z_\omega} = U_{0+Z_\omega},$$

and hence

$$\kappa_{(x,\phi)}(\iota(h \cdot 1_0)) = \text{Tr}_T^\omega(U_{0+Z_\omega}) = 1. \quad (6.12)$$

Since  $\Phi$ ,  $\iota^{-1}$ ,  $\psi_T$ ,  $\text{ev}_{(x,\phi)}$ , and  $\text{Tr}_T^\omega$  are all positive norm-decreasing linear maps,  $\kappa_{(x,\phi)}$  is a positive linear functional, and  $\|\kappa_{(x,\phi)}\| \leq 1$ . By Urysohn's lemma there exists  $h \in C_c(X)$  such that  $h(x) = 1$ . Then [Equation \(6.12\)](#) implies that  $\|\kappa_{(x,\phi)}\| \geq 1$ , and thus  $\kappa_{(x,\phi)}$  is a state of  $C^*(\mathcal{G}_T, \sigma)$ .  $\square$

We conclude this section by proving [Theorem 6.1\(b\)](#), which says that if  $(X, T)$  is minimal, then  $C^*(\mathcal{G}_T, \sigma)$  is simple if and only if the spectral action  $\theta$  is minimal.

*Proof of Theorem 6.1(b).* Suppose that  $\theta$  is minimal. Let  $I$  be a nontrivial ideal of  $C^*(\mathcal{G}_T, \sigma)$ . Then there exists a non-injective homomorphism  $\Psi$  of  $C^*(\mathcal{G}_T, \sigma)$  such that  $I = \ker(\Psi)$ . By [4, Theorem 6.4],  $J := \ker(\Psi \circ \iota)$  is a nontrivial ideal of  $C^*(\mathcal{I}_T, \sigma)$ . We have

$$\iota(J) = \{\iota(a) : a \in C^*(\mathcal{I}_T, \sigma), \Psi(\iota(a)) = 0\} = \{b \in M_T^\sigma : \Psi(b) = 0\} = I \cap M_T^\sigma \subseteq I.$$

Thus, to see that  $C^*(\mathcal{G}_T, \sigma)$  is simple, it suffices to show that  $J = C^*(\mathcal{I}_T, \sigma)$ , because then  $\iota(C_0(X)) \subseteq \iota(J) \subseteq I$ , and (as argued in [3, Theorem 5.3.13]) [13, Proposition 3.18] implies that  $I = C^*(\mathcal{G}_T, \sigma)$ . Define

$$K_J := \{(x, \phi) \in X \times \widehat{P}_T : f(x, \phi) = 0 \text{ for all } f \in \psi_T(J)\}.$$

Since  $\psi_T(J)$  is an ideal of  $\mathcal{X}_T^\omega$ , Proposition 5.5 implies that  $K_J$  is a closed subset of  $X \times \widehat{P}_T$ , and

$$\psi_T(J) = \{f \in \mathcal{X}_T^\omega : f|_{K_J} \equiv 0\}.$$

Let  $Q_T: X \times \widehat{P}_T \rightarrow X \times \widehat{Z}_\omega$  be the quotient map  $(x, \phi) \mapsto (x, \phi|_{Z_\omega})$ . Suppose that  $Q(K_J)$  is nonempty, and fix  $(x, \zeta) \in Q(K_J)$ . By Proposition 6.6,  $Q(K_J)$  is closed and invariant under  $\theta$ , and hence

$$\overline{[x, \zeta]_\theta} = \overline{\{\theta_{[\gamma]}(x, \zeta) : \gamma \in (\mathcal{G}_T)_x\}} \subseteq Q(K_J).$$

Since  $\theta$  is minimal by assumption,  $Q(K_J) = X \times \widehat{Z}_\omega$ . Thus, Proposition 6.6 implies that

$$K_J = Q^{-1}(Q(K_J)) = Q^{-1}(X \times \widehat{Z}_\omega) = X \times \widehat{P}_T.$$

Hence

$$\psi_T(J) = \{f \in \mathcal{X}_T^\omega : f|_{K_J} \equiv 0\} = \{0\},$$

which contradicts that  $J$  is nontrivial, because  $\psi_T$  is injective. Therefore,  $Q(K_J) = \emptyset$ , forcing  $K_J = \emptyset$ , and hence  $\psi_T(J) = \mathcal{X}_T^\omega$ . Since  $\psi_T$  is an isomorphism,  $J = C^*(\mathcal{I}_T, \sigma)$ , and hence  $C^*(\mathcal{G}_T, \sigma)$  is simple.

For the converse, we prove the contrapositive. Suppose that  $\theta$  is not minimal. Then there exists  $(x, \phi) \in X \times \widehat{P}_T$  such that

$$\overline{[x, \phi|_{Z_\omega}]_\theta} = \overline{\{\theta_{[\gamma]}(x, \phi|_{Z_\omega}) : \gamma \in (\mathcal{G}_T)_x\}} \neq X \times \widehat{Z}_\omega.$$

Let

$$\kappa_{(x, \phi)} := \text{Tr}_T^\omega \circ \text{ev}_{(x, \phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi : C^*(\mathcal{G}_T, \sigma) \rightarrow \mathbb{C}$$

be the state of  $C^*(\mathcal{G}_T, \sigma)$  defined in Lemma 6.10. Let  $\kappa := \kappa_{(x, \phi)}$ , let

$$N_\kappa := \{f \in C^*(\mathcal{G}_T, \sigma) : \kappa(f^*f) = 0\}$$

be the null space for  $\kappa$ , and let  $\pi_\kappa : C^*(\mathcal{G}_T, \sigma) \rightarrow B(\mathcal{H}_\kappa)$  be the GNS representation associated to  $\kappa$ . To see that  $C^*(\mathcal{G}_T, \sigma)$  is not simple, it suffices to prove that

$$\{0\} \neq \ker(\pi_\kappa) \neq C^*(\mathcal{G}_T, \sigma).$$

Since  $\kappa \neq 0$ , we have  $\mathcal{H}_\kappa \neq \{0\}$ . So since  $\pi_\kappa$  is nondegenerate,  $\ker(\pi_\kappa) \neq C^*(\mathcal{G}_T, \sigma)$ . We now show that  $\ker(\pi_\kappa) \neq \{0\}$ . Define  $C_{(x, \phi)} := Q^{-1}(\overline{[x, \phi|_{Z_\omega}]_\theta})$ . Since  $\overline{[x, \phi|_{Z_\omega}]_\theta} \neq X \times \widehat{Z}_\omega$ , Lemma 6.7 shows that  $C_{(x, \phi)}$  is a proper closed subset of  $X \times \widehat{P}_T$ , and there exists  $f \in M_T^\sigma \setminus \{0\}$  such that

$$\text{supp}((\psi_T \circ \iota^{-1})(f)) \subseteq (X \times \widehat{P}_T) \setminus C_{(x, \phi)}.$$

Fix  $a, b \in C^*(\mathcal{G}_T, \sigma)$ . To see that  $\pi_\kappa(f) = 0$ , it suffices to show that

$$(\pi_\kappa(f)(a + N_\kappa) \mid b + N_\kappa) = 0.$$

Since  $\pi_\kappa$  is the GNS representation associated to  $\kappa$ , we have

$$(\pi_\kappa(f)(a + N_\kappa) \mid b + N_\kappa) = (fa + N_\kappa \mid b + N_\kappa) = \kappa(b^*fa).$$

By [Proposition 6.9](#), we have  $(\psi_T \circ \iota^{-1} \circ \Phi)(b^*fa)(x, \phi) = 0$ , and hence

$$\kappa(b^*fa) = (\mathrm{Tr}_T^\omega \circ \mathrm{ev}_{(x, \phi)} \circ \psi_T \circ \iota^{-1} \circ \Phi)(b^*fa) = \mathrm{Tr}_T^\omega(0) = 0.$$

Hence  $(\pi_\kappa(f)(a + N_k) \mid b + N_k) = 0$ , giving  $\ker(\pi_\kappa) \neq \{0\}$ .  $\square$

*Remark 6.11.* If  $X$  is the infinite-path space of a cofinal, row-finite  $k$ -graph with no sources, and each  $T^n$  is the degree- $n$  shift map, then [Theorem 6.1](#) coincides with the simplicity characterisation given in [\[24, Corollary 4.8\]](#).

*Remark 6.12.* [Theorem 5.1](#) of [\[7\]](#) shows that  $C^*(\mathcal{G}_T)$  is simple if and only if  $\mathcal{G}_T$  is minimal and effective. We claim that [\[7, Theorem 5.1\]](#) is equivalent to [Theorem 6.1](#) when  $\sigma$  is trivial. In this case,  $\omega$  and each  $\tilde{\tau}_\gamma^\sigma$  are also trivial, and  $Z_\omega = P_T$ . So [Theorem 6.1](#) says that  $C^*(\mathcal{G}_T, \sigma) = C^*(\mathcal{G}_T)$  is simple if and only if the set

$$[x, \phi]_\theta = \{(r(\gamma), \tilde{\tau}_{[\gamma]}^\sigma \phi) : \gamma \in (\mathcal{G}_T)_x\} = r((\mathcal{G}_T)_x) \times \{\phi\}$$

is dense in  $X \times \widehat{P}_T$  for all  $(x, \phi) \in X \times \widehat{P}_T$ . Since  $r((\mathcal{G}_T)_x)$  is dense in  $X$  and  $X \times \{\phi\}$  is closed, we deduce that  $\theta$  is minimal if and only if  $P_T = \{0\}$ . By [Corollary 3.7](#), this occurs precisely when  $\mathcal{G}_T$  is effective.

## 7. AN APPLICATION TO SOME CROSSED PRODUCTS BY $\mathbb{Z}$

In this section we apply our theorem to characterise simplicity of crossed products of C\*-algebras of rank-1 Deaconu–Renault groupoids arising from continuous  $\mathbb{T}$ -valued functions on the underlying spaces. We then specialise this to the analogue of quasi-free actions on topological-graph C\*-algebras.

To set up our example, we will need the following folklore result about multipliers of the C\*-algebras of Hausdorff étale groupoids. We write  $C_b(Y)$  for the set of continuous, bounded, complex-valued functions on a locally compact Hausdorff space  $Y$ .

**Lemma 7.1.** *Let  $\mathcal{G}$  be a Hausdorff étale groupoid, and fix  $\sigma \in Z^2(\mathcal{G}, \mathbb{T})$ . Suppose that  $B \subseteq \mathcal{G}$  is a clopen bisection of  $\mathcal{G}$  such that  $s(B)$  and  $r(B)$  are closed, and fix  $f \in C_b(B)$ . For  $g \in C_c(\mathcal{G}, \sigma)$ , the convolution product  $f * g$  given by  $(f * g)(\gamma) = \sum_{\alpha \in \mathcal{G}^r(\gamma)} \sigma(\alpha, \alpha^{-1}\gamma) f(\alpha) g(\alpha^{-1}\gamma)$  belongs to  $C_c(\mathcal{G}, \sigma)$ . There is a multiplier  $M_f$  of  $C^*(\mathcal{G}, \sigma)$  such that for  $g \in C_c(\mathcal{G}, \sigma)$  we have  $M_f(g) = f * g$ . If  $s(B) = r(B) = \mathcal{G}^{(0)}$  and  $f(B) \subseteq \mathbb{T}$ , then  $M_f$  is a unitary multiplier of  $C^*(\mathcal{G}, \sigma)$ . If  $B_1$  and  $B_2$  are two clopen bisections such that  $r(B_i)$  and  $s(B_i)$  are closed for each  $i$ , and  $f_i \in C_b(B_i)$  for each  $i$ , then the convolution product  $f_1 * f_2$  belongs to  $C_b(B_1 B_2)$ , and we have  $M_{f_1} \circ M_{f_2} = M_{f_1 * f_2}$ ; likewise,  $f_1^* \in C_b(B_1^{-1})$  and  $M_{f_1}^* = M_{f_1^*}$ .*

*Proof.* Fix  $g \in C_c(\mathcal{G}, \sigma)$ . Since  $\mathrm{supp}(g)$  is compact, its image  $r(\mathrm{supp}(g))$  under the continuous range map is also compact. Fix any compactly supported function  $h \in C_c(\mathcal{G}^{(0)}, [0, 1])$  such that  $h|_{r(\mathrm{supp}(g))} \equiv 1$ . Then  $f * g = f * (h * g) = (f * h) * g$ . The function  $f * h$  is given by  $(f * h)(\gamma) = f(\gamma)h(s(\gamma))$ , and since  $\mathrm{supp}(f * h) \subseteq s|_B^{-1}(\mathrm{supp}(h) \cap s(B)) \subseteq B$ , it follows that  $f * h \in C_c(\mathcal{G}, \sigma)$ . So  $f * g = (f * h) * g \in C_c(\mathcal{G}, \sigma)$ .

Using the same function  $h$  as above, we see that

$$\|f * g\|^2 = \|((f * h) * g)^* * ((f * h) * g)\| = \|g^* * (h * f^* * f * h) * g\|.$$

In  $C^*(\mathcal{G}, \sigma)$ , we have  $g^* * (h * f^* * f * h) * g \leq \|h * f^* * f * h\|_\infty g^* g \leq \|f\|_\infty^2 g^* g$ , and so we deduce that  $\|f * g\| \leq \|f\|_\infty \|g\|$ . So the map  $g \mapsto f * g$  on  $C_c(\mathcal{G}, \sigma)$  extends to a bounded linear map  $M_f$  of norm at most  $\|f\|_\infty$  on  $C^*(\mathcal{G}, \sigma)$ . Defining  $f^*: B^{-1} \rightarrow \mathbb{C}$  by  $f^*(\gamma) = \overline{\sigma(\gamma^{-1}, \gamma)} f(\gamma^{-1})$  as usual, associativity of multiplication shows that for  $g, h \in C_c(\mathcal{G}, \sigma)$ , we have  $M_f(g)^* * h = g^* * f^* * h = g^* * M_{f^*}(h)$ . Thus  $M_f$  is adjointable

with respect to the standard inner product on  $C^*(\mathcal{G}, \sigma)$ , with adjoint  $M_{f^*}$ . So  $M_f$  is a multiplier, as claimed. If  $s(B) = r(B) = \mathcal{G}^{(0)}$  and  $f(B) \subseteq \mathbb{T}$ , then for  $g \in C_c(\mathcal{G}, \sigma)$  we have  $M_f^*(M_f(g)) = f^* * f * g = 1_{\mathcal{G}^{(0)}} * g = g$ , and similarly,  $M_f(M_f^*(g)) = g$ . So continuity gives  $M_f^* M_f = M_f M_f^* = 1_{\mathcal{M}(C^*(\mathcal{G}, \sigma))}$ , and thus  $M_f$  is a unitary.

For the final statement, we already saw that  $M_{f^*} = M_f^*$  for all  $f$ , so we just have to establish the multiplicativity. If  $B_1$  and  $B_2$  are clopen bisections and  $f_i \in C_b(B_i)$ , then  $B_1 B_2$  is an open bisection because multiplication is open. To see that  $B_1 B_2$  is also closed, suppose that  $\gamma_i \rightarrow \gamma$  and each  $\gamma_i \in B_1 B_2$ . Then each  $\gamma_i$  can be written as  $\alpha_i \beta_i$  with each  $\alpha_i$  in  $B_1$  and each  $\beta_i$  in  $B_2$ . Since  $\gamma_i \rightarrow \gamma$ , we have  $r(\alpha_i) = r(\gamma_i) \rightarrow r(\gamma)$ , and then since  $r|_{B_1}$  is a homeomorphism,  $\alpha_i \rightarrow \alpha$  for some  $\alpha \in B_1$ . Similarly (using  $s$  in place of  $r$ ), we have  $\beta_i \rightarrow \beta$  for some  $\beta \in B_2$ . Since each  $s(\alpha_i) = r(\beta_i)$ , continuity gives  $s(\alpha) = r(\beta)$ , and since  $\alpha_i \beta_i = \gamma_i \rightarrow \gamma$ , continuity also gives  $\alpha \beta = \gamma$ . So  $\gamma \in B_1 B_2$ . The convolution formula shows that  $\text{supp}(f_1 * f_2) \subseteq B_1 B_2$  and that  $\|f_1 * f_2\|_\infty \leq \|f_1\|_\infty \|f_2\|_\infty$ . For  $g \in C_c(\mathcal{G}, \sigma)$  we have  $M_{f_1}(M_{f_2}(g)) = f_1 * (f_2 * g) = (f_1 * f_2) * g = M_{f_1 * f_2}(g)$ , and then continuity gives  $M_{f_1} M_{f_2} = M_{f_1 * f_2}$ .  $\square$

**Lemma 7.2.** *Let  $(X, T)$  be a rank-1 Deaconu–Renault system such that  $X$  is second-countable, and let  $h: X \rightarrow \mathbb{T}$  be a continuous function.*

(a) *For each  $(x, p, y) \in \mathcal{G}_T \subseteq X \times \mathbb{Z} \times X$ , the sequence*

$$\left( \prod_{i=0}^N h(T^i(x)) \prod_{i=0}^{N-p} \overline{h(T^i(y))} \right)_{N=|p|}^\infty$$

*is eventually constant.*

(b) *There is a continuous 1-cocycle  $\tilde{h}: \mathcal{G}_T \rightarrow \mathbb{T}$  such that*

$$\tilde{h}(x, p, y) = \prod_{i=0}^N h(T^i(x)) \prod_{i=0}^{N-p} \overline{h(T^i(y))}$$

*for large  $N \in \mathbb{N}$ .*

(c) *There is an action  $\alpha^h: \mathbb{Z} \rightarrow \text{Aut}(C^*(\mathcal{G}_T))$  such that  $\alpha_n^h(f)(\gamma) = \tilde{h}(\gamma)^n f(\gamma)$  for all  $f \in C_c(\mathcal{G}_T)$  and  $\gamma \in \mathcal{G}_T$ .*

(d) *There is a continuous 2-cocycle  $c_h: (\mathcal{G}_T \times \mathbb{Z})^{(2)} \rightarrow \mathbb{T}$  given by*

$$c_h((\alpha, m), (\beta, n)) := \tilde{h}(\beta)^m,$$

*and there is an isomorphism  $\phi: C^*(\mathcal{G}_T) \rtimes_{\alpha^h} \mathbb{Z} \rightarrow C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  such that*

$$\phi(i_{C^*(\mathcal{G}_T)}(f) i_{\mathbb{Z}}(n))(\gamma, p) = \delta_{n,p} f(\gamma).$$

*Proof.* (a) Fix  $(x, p, y) \in \mathcal{G}_T$ . By definition there exist  $m, n \in \mathbb{N}$  such that  $m - n = p$  and  $T^m(x) = T^n(y)$ . Let  $N := \max\{m, n\}$ . For  $M \geq N$ , we have  $T^{M+1}(x) = T^{M-m+1}(T^m(x)) = T^{M-m+1}(T^n(y)) = T^{M+1-p}(y)$ . Hence

$$\prod_{i=0}^{M+1} h(T^i(x)) \prod_{i=0}^{M+1-p} \overline{h(T^i(y))} = \prod_{i=0}^M h(T^i(x)) \prod_{i=0}^{M-p} \overline{h(T^i(y))}.$$

Now a simple induction proves (a).

(b) For fixed  $N \in \mathbb{N}$ , the map  $x \mapsto \prod_{i=0}^N h(T^i(x))$  is continuous, and it follows that  $\tilde{h}$  is continuous on each basic open set  $Z(U, m, n, V)$ . So  $\tilde{h}$  is continuous. To see that  $\tilde{h}$  is a 1-cocycle, fix  $((x, p, y), (y, q, z)) \in \mathcal{G}_T^{(2)}$ . Then for sufficiently large  $M, N \in \mathbb{N}$ , we have

$$\tilde{h}(x, p, y) \tilde{h}(y, q, z) = \prod_{i=0}^M h(T^i(x)) \prod_{i=0}^{M-p} \overline{h(T^i(y))} \prod_{i=0}^N h(T^i(y)) \prod_{i=0}^{N-q} \overline{h(T^i(z))}.$$

In particular, by choosing  $M$  large enough, we can take  $N = M - p$ , so that  $N - q = M - (p + q)$ , and then  $\prod_{i=0}^{M-p} \overline{h(T^i(y))} \prod_{i=0}^N h(T^i(y)) = 1$ , giving

$$\tilde{h}(x, p, y) \tilde{h}(y, q, z) = \prod_{i=0}^M h(T^i(x)) \prod_{i=0}^{M-(p+q)} \overline{h(T^i(z))} = \tilde{h}(x, p + q, z).$$

Thus  $\tilde{h}$  is a 1-cocycle.

(c) It is standard that a continuous 1-cocycle on  $\mathcal{G}_T$  determines an automorphism of its C\*-algebra: the given formula for  $\alpha_n^h$  determines a linear transformation of  $C_c(\mathcal{G}_T)$ . For  $f, g \in C_c(\mathcal{G}_T)$  and  $\gamma \in \mathcal{G}_T$ , using that  $\tilde{h}$  is a 1-cocycle at the third equality, we calculate

$$\begin{aligned} (\alpha_n^h(f) * \alpha_n^h(g))(\gamma) &= \sum_{\beta \in \mathcal{G}_T^{r(\gamma)}} \alpha_n^h(f)(\beta) \alpha_n^h(g)(\beta^{-1}\gamma) \\ &= \sum_{\beta \in \mathcal{G}_T^{r(\gamma)}} \tilde{h}(\beta)^n \tilde{h}(\beta^{-1}\gamma)^n f(\beta) g(\beta^{-1}\gamma) \\ &= \tilde{h}(\gamma)^n \sum_{\beta \in \mathcal{G}_T^{r(\gamma)}} f(\beta) g(\beta^{-1}\gamma) = \alpha_n^h(f * g)(\gamma). \end{aligned}$$

Similarly, we have  $\alpha_n^h(f)^*(\gamma) = \overline{\tilde{h}(\gamma^{-1})^n f(\gamma^{-1})} = \tilde{h}(\gamma)^n \overline{f(\gamma^{-1})} = \alpha_n^h(f^*)(\gamma)$ . Hence  $\alpha_n^h$  is a \*-homomorphism from  $C_c(\mathcal{G}_T)$  to  $C_c(\mathcal{G}_T) \subseteq C^*(\mathcal{G}_T)$ . The universal property of  $C^*(\mathcal{G}_T)$  then implies that  $\alpha_n^h$  extends to a homomorphism  $C^*(\mathcal{G}_T) \rightarrow C^*(\mathcal{G}_T)$ , and since  $\alpha_{-n}^h$  is an inverse for it on the dense subspace  $C_c(\mathcal{G}_T)$ , we see that  $\alpha^h$  is an automorphism. The map  $n \mapsto \alpha_n^h$  is a homomorphism because  $\alpha_n^h(\alpha_m^h(f))(\gamma) = \tilde{h}(\gamma)^n \tilde{h}(\gamma)^m f(\gamma) = \alpha_{n+m}^h(f)(\gamma)$ .

(d) The map  $c_h$  is continuous because  $\tilde{h}$  is continuous. To see that  $c_h$  satisfies the 2-cocycle identity, fix a composable triple  $((\alpha, m), (\beta, n), (\gamma, p))$  and calculate

$$\begin{aligned} c_h((\alpha, m), (\beta, n)) c_h((\alpha, m)(\beta, n), (\gamma, p)) &= \tilde{h}(\beta)^m \tilde{h}(\gamma)^{m+n} = \tilde{h}(\beta\gamma)^m \tilde{h}(\gamma)^n \\ &= c_h((\alpha, m), (\beta, n)(\gamma, p)) c_h((\beta, n), (\gamma, p)). \end{aligned}$$

To see that  $c_h$  is normalised, observe that since  $\tilde{h}$  is a 1-cocycle, it vanishes on  $\mathcal{G}_T^{(0)}$ , and so  $c_h((r(\gamma), 0), (\gamma, n)) = \tilde{h}(\gamma)^0 = 1 = \tilde{h}(s(\gamma))^n = c_h((\gamma, n), (s(\gamma), 0))$ .

For the final statement, first note that for  $n \in \mathbb{Z}$ , the set  $\mathcal{G}_T^{(0)} \times \{n\}$  is a clopen bisection of  $\mathcal{G}_T \times \mathbb{Z}$  with range and source equal to  $(\mathcal{G}_T \times \mathbb{Z})^{(0)}$ . Hence Lemma 7.1 shows that there is a unitary multiplier  $U_n$  of  $C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  that acts on  $C_c(\mathcal{G}_T \times \mathbb{Z}, c_h)$  by convolution with the characteristic function  $1_{\mathcal{G}_T^{(0)} \times \{n\}}$ . Since  $\tilde{h}$  vanishes on  $\mathcal{G}_T^{(0)}$ , the final statement of Lemma 7.1 shows that  $n \mapsto U_n$  is a unitary representation of  $\mathbb{Z}$ .

Since  $\mathcal{G}_T \times \{0\}$  is isomorphic to  $\mathcal{G}_T$  and  $c_h$  is trivial on  $\mathcal{G}_T \times \{0\}$ , the universal property of  $C^*(\mathcal{G}_T)$  yields a homomorphism  $\pi: C^*(\mathcal{G}_T) \rightarrow C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  such that  $\pi(f)(\gamma, m) = \delta_{m,0} f(\gamma)$  for all  $f \in C_c(\mathcal{G}_T)$  and  $(\gamma, m) \in \mathcal{G}_T \times \mathbb{Z}$ . We claim that  $\pi$  is nondegenerate. To see this, fix  $g \in C_c(\mathcal{G}_T \times \mathbb{Z}, c_h)$ , and use Urysohn's lemma to choose  $f \in C_c(\mathcal{G}_T)$  such that  $\text{supp}(f) \subseteq \mathcal{G}_T^{(0)}$  and  $f|_{\pi_1(r(\text{supp}(g)))} \equiv 1$ , where  $\pi_1$  is the projection of  $\mathcal{G}_T \times \mathbb{Z}$  onto the first coordinate. A routine calculation shows that  $\pi(f)g = g$ , and hence  $\pi$  is nondegenerate.



We claim that  $(\pi, U)$  is a covariant representation of  $(C^*(\mathcal{G}_T), \mathbb{Z}, \alpha^h)$ . To see this, fix  $f \in C_c(\mathcal{G}_T)$  and  $n \in \mathbb{Z}$ . For all  $(\gamma, m) \in \mathcal{G}_T \times \mathbb{Z}$ , we have

$$\begin{aligned} (U_n \pi(f) U_n^*)(\gamma, m) &= (U_n \pi(f) U_{-n})(\gamma, m) \\ &= \sum_{(\alpha, p)(\beta, q)(\lambda, l) = (\gamma, m)} \tilde{h}(\beta)^p \tilde{h}(\lambda)^{p+q} 1_{\mathcal{G}_T^{(0)} \times \{n\}}(\alpha, p) \pi(f)(\beta, q) 1_{\mathcal{G}_T^{(0)} \times \{-n\}}(\lambda, l). \end{aligned}$$

If  $(\alpha, p)(\beta, q)(\lambda, l) = (\gamma, m)$  contributes a nonzero term, then  $(\alpha, p) \in \mathcal{G}_T^{(0)} \times \{n\}$  and  $(\lambda, l) \in \mathcal{G}_T^{(0)} \times \{-n\}$ ; thus  $(\alpha, p) = (r(\gamma), n)$  and  $(\lambda, l) = (s(\gamma), -n)$ , and hence  $(\beta, q) = (\gamma, m)$ . So we obtain

$$(U_n \pi(f) U_n^*)(\gamma, m) = \tilde{h}(\gamma)^n \pi(f)(\gamma, m) = \delta_{m,0} \tilde{h}(\gamma)^n f(\gamma) = \pi(\alpha_n^h(f))(\gamma, m).$$

Therefore,  $(\pi, U)$  is a nondegenerate covariant representation of  $(C^*(\mathcal{G}_T), \mathbb{Z}, \alpha^h)$ , and so the universal property of the crossed product gives a homomorphism  $\phi: C^*(\mathcal{G}_T) \rtimes_{\alpha^h} \mathbb{Z} \rightarrow C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  such that  $\phi(i_{C^*(\mathcal{G}_T)}(f) i_{\mathbb{Z}}(n))(\gamma, p) = (\pi(f) U_n)(\gamma, p) = \delta_{n,p} f(\gamma)$ .

To see that  $\phi$  is injective, it suffices by [17, Theorem 4.5] to show that  $\pi$  is injective and that there is a strongly continuous action  $\beta$  of  $\mathbb{T}$  on  $C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  such that for each  $z \in \mathbb{T}$ , we have  $\beta_z(\pi(f)) = \pi(f)$  for all  $f \in C_c(\mathcal{G}_T)$ , and the extension  $\bar{\beta}_z$  of  $\beta_z$  to the multiplier algebra  $\mathcal{M}(C^*(\mathcal{G}_T \times \mathbb{Z}, c_h))$  satisfies  $\bar{\beta}_z(U_n) = z^n U_n$  for all  $n \in \mathbb{Z}$ .

The homomorphism  $\pi$  is clearly injective on  $C_0(\mathcal{G}_T^{(0)})$  and intertwines the canonical expectations  $C^*(\mathcal{G}_T) \rightarrow C_0(\mathcal{G}_T^{(0)})$  and  $C^*(\mathcal{G}_T \times \mathbb{Z}, c_h) \rightarrow C_0((\mathcal{G}_T \times \mathbb{Z})^{(0)})$  extending restriction of  $C_c$ -functions to  $\mathcal{G}_T^{(0)}$  and  $(\mathcal{G}_T \times \mathbb{Z})^{(0)}$ , respectively. Since  $\mathcal{G}_T$  is amenable [35, Lemma 3.5], the expectation  $C^*(\mathcal{G}_T) \rightarrow C_0(\mathcal{G}_T^{(0)})$  is faithful, and so a standard argument (see, for example, [34, Lemma 3.14]) shows that  $\pi$  is injective.

So we just need to construct the action  $\beta$ . For  $z \in \mathbb{T}$ , the map  $\beta_z: C_c(\mathcal{G}_T \times \mathbb{Z}, c_h) \rightarrow C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  given by  $\beta_z(f)(\gamma, n) = z^n f(\gamma, n)$  is a  $*$ -homomorphism, so the universal property of  $C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$  implies that it extends to an endomorphism  $\beta_z$  of  $C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$ . Since  $\beta_{\bar{z}} \circ \beta_z$  is the identity on  $C_c(\mathcal{G}_T \times \mathbb{Z}, c_h)$ , we deduce that  $\beta_z$  is an automorphism, and since  $\beta_z \circ \beta_w$  agrees with  $\beta_{zw}$  on  $C_c(\mathcal{G}_T \times \mathbb{Z}, c_h)$  we see that  $z \mapsto \beta_z$  is a homomorphism. For  $f \in C_c(\mathcal{G}_T \times \{n\})$ , the map  $z \mapsto \beta_z(f)$  is clearly continuous. So an  $\varepsilon$ -argument shows that  $\beta$  is a strongly continuous action of  $\mathbb{T}$ .

We claim that the extension of each  $\beta_z$  to the multiplier algebra  $\mathcal{M}(C^*(\mathcal{G}_T \times \mathbb{Z}, c_h))$  satisfies  $\bar{\beta}_z(U_n) = z^n U_n$  for each  $n \in \mathbb{Z}$ . To see this, fix  $n \in \mathbb{Z}$  and an increasing sequence  $K_i \subseteq \mathcal{G}_T^{(0)}$  of compact sets with  $\bigcup_{i \in \mathbb{N}} K_i = \mathcal{G}_T^{(0)}$ , and for each  $i \in \mathbb{N}$ , fix  $h_i \in C_c(\mathcal{G}_T^{(0)} \times \{n\}, [0, 1])$  such that  $h_i|_{K_i \times \{n\}} \equiv 1$ . For  $f \in C_c(\mathcal{G}_T \times \mathbb{Z})$  there exists  $N \in \mathbb{N}$  large enough so that  $r(\text{supp}(f)) \subseteq K_N \times \{0\}$ , and then  $h_i * f = U_n f$  for all  $i \geq N$ . So the  $h_i$  converge strictly to  $U_n$ , and since  $\beta_z(h_i) = z^n h_i$  for all  $z \in \mathbb{T}$ , this establishes the claim.

It remains only to prove that  $\phi$  is surjective. For this, fix an open bisection  $B$  of  $\mathcal{G}_T \times \mathbb{Z}$  and distinct points  $\beta, \gamma \in B$ . Then  $B = B' \times \{n\}$  for some open bisection  $B'$  of  $\mathcal{G}_T$  and some  $n \in \mathbb{Z}$ , and so  $\beta = (\beta', n)$  and  $\gamma = (\gamma', n)$  for distinct  $\beta', \gamma' \in \mathcal{G}_T$ . Fix  $f \in C_c(\mathcal{G}_T)$  such that  $\text{supp}(f) \subseteq B'$ ,  $f(\beta') = 1$ , and  $f(\gamma') = 0$ . Then  $\phi(i_{C^*(\mathcal{G}_T)}(f) i_{\mathbb{Z}}(n))(\beta', n) = (\pi(f) U_n)(\beta', n) = 1$  and  $\phi(i_{C^*(\mathcal{G}_T)}(f) i_{\mathbb{Z}}(n))(\gamma', n) = (\pi(f) U_n)(\gamma', n) = 0$ . So [33, Corollary 9.3.5] shows that  $\phi$  is surjective.  $\square$

**Theorem 7.3.** *Let  $(X, T)$  be a rank-1 Deaconu–Renault system such that  $X$  is second-countable, and let  $h: X \rightarrow \mathbb{T}$  be a continuous function. Let  $\tilde{h}: \mathcal{G}_T \rightarrow \mathbb{T}$  be the 1-cocycle of Lemma 7.2(b), and let  $\alpha^h: \mathbb{Z} \rightarrow \text{Aut}(C^*(\mathcal{G}_T))$  be the action of Lemma 7.2(c). Write  $\rho$*

for the action of  $\mathcal{G}_T$  on  $X \times \mathbb{T}$  given by  $\rho_\gamma(s(\gamma), z) := (r(\gamma), \tilde{h}(\gamma)z)$ . Suppose that  $X$  is an uncountable space. Then the crossed product  $C^*(\mathcal{G}_T) \rtimes_{\alpha^h} \mathbb{Z}$  is simple if and only if  $\rho$  is minimal.

In order to prove [Theorem 7.3](#), we need the following lemma.

**Lemma 7.4.** *Let  $(X, T)$  be a minimal rank-1 Deaconu–Renault system such that  $X$  is second-countable. If  $X$  is uncountable, then  $\mathcal{G}_T$  is topologically principal.*

*Proof.* Since  $\mathcal{G}_T$  is second-countable, it suffices by [[7](#), Lemma 3.1] to show that  $\mathcal{I}_T = \mathcal{G}_T^{(0)}$ . To see this, we suppose that  $\mathcal{I}_T \neq \mathcal{G}_T^{(0)}$  and derive a contradiction. Recall from [Proposition 3.5](#) that  $\mathcal{I}_T = \{(x, p, x) : p \in P_T\}$ . Since  $\mathcal{I}_T$  is nontrivial, there exists  $p \in \mathbb{Z} \setminus \{0\}$  such that  $(x, p, x) \in \mathcal{G}_T$  for all  $x$ . By definition of the topology on  $\mathcal{G}_T$ , it follows that for each  $x \in X$  there is an open neighbourhood  $U$  of  $x$  and a pair  $m > n \in \mathbb{N}$  such that  $T^m(x) = T^n(x)$  for all  $x \in U$ . Since the pairs  $m > n \in \mathbb{N}$  are countable and  $X$  is not countable, that  $\mathcal{G}_T$  is second-countable implies that there exist  $x, U, m, n$  as above so that  $U$  is not countable. Since  $X$  is second-countable and  $T^n$  is a local homeomorphism,  $(T^n)^{-1}(x)$  is countable for every  $x \in X$ , and so  $V = T^n(U)$  is an uncountable open set and  $p = m - n > 0$  satisfies  $T^p(x) = x$  for all  $x \in V$ . Fix  $x \in V$ . Since  $V$  is uncountable, there exists  $y \in V$  such that

$$T^q(x) \neq y \quad \text{for all } q \in \mathbb{N}. \quad (7.1)$$

Since  $\mathcal{G}_T$  is minimal, there is a sequence  $(z_i, m_i, x)_{i=1}^\infty$  in  $\mathcal{G}_T$  such that  $z_i \rightarrow y$ . Write each  $m_i = a_i - b_i$  with  $a_i, b_i \geq 0$  so that  $T^{a_i}(z_i) = T^{b_i}(x)$ . For each  $i \in \mathbb{N}$ , there exists  $k > 0$  such that  $kp > b_i$ ; and then  $T^{a_i+(kp-b_i)}(z_i) = T^{kp}(x) = x$ . So we can assume that each  $m_i > 0$  and that  $T^{m_i}(z_i) = x$  for all  $i \in \mathbb{N}$ . By passing to a subsequence, we may assume that each  $m_i - m_1$  is divisible by  $p$ . Fix  $l > 0$  such that  $lp > m_1$ , let  $d := lp - m_1$ , let  $z := T^d(x)$ , and let  $n_i := m_i + d$  for all  $i \in \mathbb{N}$ . Then  $T^{n_i}(z_i) = z$  for all  $i \in \mathbb{N}$ , and each  $n_i$  is divisible by  $p$ . Since  $z_i \rightarrow y$ , we eventually have  $z_i \in V$ , and so we eventually have  $z = T^{n_i}(z_i) = z_i$ . But this forces  $y = z = T^d(x)$ , which contradicts [Condition \(7.1\)](#). Thus  $\mathcal{I}_T$  is trivial, as claimed.  $\square$

*Proof of [Theorem 7.3](#).* By [Lemma 7.2\(d\)](#), the crossed product  $C^*(\mathcal{G}_T) \rtimes_{\alpha^h} \mathbb{Z}$  is isomorphic to the twisted groupoid C\*-algebra  $C^*(\mathcal{G}_T \times \mathbb{Z}, c_h)$ , to which we aim to apply [Theorem 6.1](#).

For this, observe first that if  $\bar{T}$  is the action of  $\mathbb{N}^2$  given by  $\bar{T}^{(m,n)} = T^m$ , then  $\mathcal{G}_T \times \mathbb{Z} \cong \mathcal{G}_{\bar{T}}$

First suppose that  $(X, T)$  is not minimal. Then  $\mathcal{G}_{\bar{T}}$  is also not minimal, and the action  $\rho$  is not minimal. So  $(X, \bar{T})$  is not minimal, and hence [Theorem 6.1\(a\)](#) implies that  $C^*(\mathcal{G}_T) \rtimes_{\alpha^h} \mathbb{Z} \cong C^*(\mathcal{G}_{\bar{T}}, c_h)$  is not simple. So it suffices to prove the result when  $(X, T)$  is minimal.

Now suppose that  $(X, T)$  is minimal. Since  $X$  is uncountable, [Lemma 7.4](#) implies that  $\mathcal{I}_T = \mathcal{G}_T^{(0)}$ . The isomorphism  $\mathcal{G}_T \times \mathbb{Z} \rightarrow \mathcal{G}_{\bar{T}}$  is given by  $((x, m, y), n) \mapsto (x, (m, n), y)$ . So the interior  $\mathcal{I}_{\bar{T}}$  of the isotropy of  $\mathcal{G}_{\bar{T}}$  is precisely  $\{(x, (m, n), x) : (x, m, x) \in \mathcal{I}_T, n \in \mathbb{Z}\}$ . So the preceding paragraph implies that  $\mathcal{I}_{\bar{T}} = \{(x, (0, n), x) : x \in X, n \in \mathbb{Z}\}$ . The isomorphism  $\mathcal{G}_T \times \mathbb{Z} \rightarrow \mathcal{G}_{\bar{T}}$  intertwines  $c_h$  with the 2-cocycle  $\sigma \in Z^2(\mathcal{G}_{\bar{T}}, \mathbb{T})$  given by  $\sigma((x, (m, n), y), (y, (p, q), z)) := \tilde{h}(y, p, z)^n$ . The restriction of this  $\sigma$  to  $\mathcal{I}_{\bar{T}}^{(2)}$  satisfies

$$\sigma((x, (0, m), x), (x, (0, n), x)) = \tilde{h}(x, 0, x)^m = 1.$$

Hence  $\sigma$  is  $\omega$ -constant on  $\mathcal{I}_{\bar{T}}$  with  $\omega = 1$ . It follows that  $Z_\omega = P_{\bar{T}} = \{0\} \times \mathbb{Z} \cong \mathbb{Z}$ .

The  $\widehat{\mathbb{Z}}$ -valued 1-cocycle  $\tau^\sigma$  obtained from [Lemma 4.3\(e\)](#) satisfies

$$\begin{aligned} \tau_{(x,(m,n),y)}^\sigma(p) &= \sigma((x, (m, n), y), (y, (0, p), y)) \\ &\quad \cdot \sigma((x, (m, n+p), y), (y, (-m, -n), x)) \\ &\quad \cdot \overline{\sigma((x, (m, n), y), (y, (-m, -n), x))} \\ &= \widetilde{h}(y, 0, y)^n \widetilde{h}(y, -m, x)^{n+p} \overline{\widetilde{h}(y, -m, x)^n} \\ &= \widetilde{h}(y, -m, x)^p. \end{aligned}$$

So the isomorphism  $\chi \mapsto \chi(1)$  from  $\widehat{\mathbb{Z}}$  to  $\mathbb{T}$  carries  $\tau_{(x,(m,n),y)}^\sigma$  to  $\widetilde{h}(y, -m, x) \in \mathbb{T}$ .

We have  $\mathcal{H}_{\overline{T}} := \mathcal{G}_{\overline{T}}/\mathcal{I}_{\overline{T}} \cong (\mathcal{G}_T \times \mathbb{Z})/(\mathcal{G}_T^{(0)} \times \mathbb{Z}) \cong \mathcal{G}_T$  and the isomorphism is the map  $[(x, (m, n), y)] \mapsto (x, m, y)$ . So the spectral action  $\theta$  of  $\mathcal{H}_{\overline{T}}$  on  $X \times \widehat{\mathbb{Z}}_\omega$  of [Proposition 4.5](#) is identified with the action of  $\mathcal{G}_T$  on  $X \times \mathbb{T}$  given by  $\theta_{(x,m,y)}(y, z) := (x, \widetilde{h}(y, -m, x)z)$ , which is precisely the action  $\rho$ . So [Theorem 6.1](#) shows that  $C^*(\mathcal{G}_{\overline{T}}, c_h)$  is simple if and only if  $\rho$  is minimal.  $\square$

In the following result, we write  $t$  and  $o$  for the terminus (range) and origin (source) map in a topological graph, so as to avoid confusion with the range and source maps  $r$  and  $s$  in the associated groupoid. We write  $X(E)$  for the graph correspondence associated to a topological graph  $E = (E^0, E^1, t, o)$ , and we write  $\mathcal{O}_{X(E)}$  for the associated Cuntz–Pimsner algebra. We write  $(j_{C_0(E^0)}, j_{X(E)})$  for the universal Cuntz–Pimsner-covariant representation of  $X(E)$  that generates  $\mathcal{O}_{X(E)}$ . See [\[17, 37, 26\]](#) for background on topological graphs and their C\*-algebras.

**Corollary 7.5.** *Let  $E = (E^0, E^1, t, o)$  be a second-countable topological graph such that the terminus map  $t: E^1 \rightarrow E^0$  is proper and surjective, and the infinite-path space  $E^\infty$  is uncountable. Suppose that  $\ell: E^1 \rightarrow \mathbb{T}$  is a continuous function. There is an action  $\beta^\ell: \mathbb{Z} \curvearrowright C^*(E)$  such that  $\beta_n^\ell(j_{X(E)}(\xi)) = j_{X(E)}(\ell^n \cdot \xi)$  for all  $\xi \in C_c(E^1)$ . Extend  $\ell$  to a continuous function  $\ell: E^* \rightarrow \mathbb{T}$  by defining  $\ell(e_1 \cdots e_n) := \prod_{i=1}^n \ell(e_i)$  and  $\ell|_{E^0} \equiv 1$ , and let  $T: E^\infty \rightarrow E^\infty$  be the shift map  $T(x_1 x_2 x_3 \cdots) = x_2 x_3 \cdots$ . Then  $C^*(E) \rtimes_{\beta^\ell} \mathbb{Z}$  is simple if and only if for every infinite path  $x \in E^\infty$ , the set*

$$\{(\lambda T^n(x), \ell(\lambda) \overline{\ell(x(0, n))}) : n \in \mathbb{N}, \lambda \in E^* t(T^n(x))\} \quad (7.2)$$

is dense in  $E^\infty \times \mathbb{T}$ .

*Proof.* The map  $\xi \mapsto \ell \cdot \xi$  is a unitary operator  $U_\ell$  on the graph correspondence  $X(E)$ . If  $\xi \in C_c(E^1)$  is a positive-valued function such that  $o$  is injective on  $\text{supp}(\xi)$ , then a quick calculation shows that conjugation by  $U_\ell$  fixes the rank-1 operator  $\Theta_{\xi, \xi}$ . Using this, it is routine to see that if  $(\psi, \pi)$  is a covariant Toeplitz representation of  $E$  as in [\[26, Definitions 2.2 and 2.10\]](#), then so is  $(\psi \circ U_\ell, \pi)$ . So the universal property of  $C^*(E) \cong \mathcal{O}_{X(E)}$  described by [\[26, Theorem 2.13\]](#) yields a unique automorphism  $\beta^\ell$  that fixes  $j_{C_0(E^0)}(C_0(E^0))$  and satisfies  $\beta_n^\ell(j_{X(E)}(\xi)) = j_{X(E)}(\ell \cdot \xi)$  for  $\xi \in C_c(E^1)$ . The formula  $\beta_n^\ell := (\beta^\ell)^n$  then gives the desired action.<sup>1</sup>

Since  $t: E^1 \rightarrow E^0$  is proper, [\[5, Propositions 3.11 and 3.16\]](#) show that  $E^\infty$  is a locally compact Hausdorff space and  $T$  is a local homeomorphism. By [\[37, Theorem 5.2\]](#), there is an isomorphism  $\phi: C^*(E) \rightarrow C^*(\mathcal{G}_T)$  such that  $\phi(j_{C_0(E^0)}(f))(x, m, y) = \delta_{x,y} \delta_{m,0} f(t(x))$  for  $f \in C_0(E^0)$  and  $\phi(j_{X(E)}(\xi))(x, m, y) = \delta_{T(x),y} \delta_{m,1} \xi(x_1)$  for  $\xi \in C_c(E^1)$ .

Define  $h: E^\infty \rightarrow \mathbb{T}$  by  $h(x) := \ell(x_1)$ . Then  $h$  is continuous. Let  $\alpha^h \in \text{Aut}(C^*(\mathcal{G}_T))$  be the automorphism  $\alpha_1^h$  of [Lemma 7.2\(c\)](#). A routine calculation shows that  $\alpha^h \circ \phi$  agrees with

<sup>1</sup>We could also appeal to the fourth paragraph of [\[25, Page 462\]](#).

$\phi \circ \beta^\ell$  on  $j_{C_0(E^0)}(C_0(E^0)) \cup j_{X(E)}(C_c(E^1))$ , and hence the uniqueness of the automorphism  $\beta^\ell$  discussed in the first paragraph shows that  $\alpha^h \circ \phi = \phi \circ \beta^\ell$ . It therefore suffices to show that  $C^*(\mathcal{G}_T) \rtimes_{\alpha^h} \mathbb{Z}$  is simple if and only if the set described in Equation (7.2) is dense for each  $x \in E^\infty$ .

Fix  $x \in E^\infty$ . We claim that the set described in Equation (7.2) is precisely the orbit of  $(x, 1)$  under the action  $\rho$  of Theorem 7.3. We have

$$(\mathcal{G}_T)_x = \{(\lambda T^n(x), |\lambda| - n, x) : n \in \mathbb{N}, \lambda \in E^*t(T^n(x))\},$$

and so for each  $\gamma \in (\mathcal{G}_T)_x$ , we have

$$\rho_\gamma(x, 1) = (\lambda T^n(x), \tilde{h}(\lambda T^n(x), |\lambda| - n, x)), \quad (7.3)$$

for some  $n \in \mathbb{N}$  and  $\lambda \in E^*t(T^n(x))$ . Direct calculation shows that

$$\tilde{h}(\mu x, |\mu| - |\nu|, \nu x) = \ell(\mu) \overline{\ell(\nu)}, \quad (7.4)$$

for all  $x \in E^\infty$  and  $\mu, \nu \in E^*t(x)$ . Together, Equations (7.3) and (7.4) imply that the set described in Equation (7.2) is the orbit of  $(x, 1)$  under  $\rho$ . Since  $\rho$  commutes with the action of  $\mathbb{T}$  on  $E^\infty \times \mathbb{T}$  by translation in the second coordinate, the orbit of  $(x, 1)$  is dense if and only if the orbit of  $(x, z)$  is dense for every  $z \in \mathbb{T}$ . That is, the set described in Equation (7.2) is dense for each  $x \in E^\infty$  if and only if every  $\rho$ -orbit is dense. So the result follows from Theorem 7.3.  $\square$

To conclude, for the class of topological graphs appearing in Corollary 7.5, we give a sufficient condition phrased purely in terms of the graph without reference to the shift map on its infinite-path space, for simplicity of the crossed product described there. (The hypothesis that  $E^\infty$  is uncountable is quite weak, and follows from a number of elementary conditions on the graph: for example, that  $E^0$  is uncountable, or that  $E$  has at least one vertex that supports at least two distinct cycles.)

**Corollary 7.6.** *Let  $E = (E^0, E^1, t, o)$  be a second-countable topological graph such that the terminus map  $t: E^1 \rightarrow E^0$  is proper and surjective, and the infinite-path space  $E^\infty$  is uncountable. Let  $\ell: E^1 \rightarrow \mathbb{T}$  be a continuous function. Extend  $\ell$  to  $E^*$  by defining  $\ell(e_1 \cdots e_n) := \prod_{i=1}^n \ell(e_i)$  and  $\ell|_{E^0} \equiv 1$ . For each  $v \in E^0$ , define*

$$\text{Orb}^+(v) := \bigcup_{\mu \in E^*v} (t(\mu), \ell(\mu)) \subseteq E^0 \times \mathbb{T}.$$

*If  $\text{Orb}^+(v)$  is dense in  $E^0 \times \mathbb{T}$  for each  $v \in E^0$ , then the crossed product  $C^*(E) \rtimes_{\beta^\ell} \mathbb{Z}$  of Corollary 7.5 is simple.*

*Proof.* Suppose that  $\text{Orb}^+(v)$  is dense in  $E^0 \times \mathbb{T}$  for each  $v \in E^0$ . We aim to invoke Theorem 7.3. Fix  $(x, w), (y, z) \in E^\infty \times \mathbb{T}$ . Recall from [5, Proposition 3.11 and Lemma 3.13] that for  $n \in \mathbb{N}$  and an open neighbourhood  $U \subseteq E^n$  of  $y(0, n)$  such that  $o|_U$  is injective, the set  $Z(U) = \{y' \in E^\infty : y'(0, n) \in U\}$  is a basic open neighbourhood of  $y$ . Let  $d$  be the metric on  $\mathbb{T}$  induced by the usual metric on  $\mathbb{R}$  via the local homeomorphism  $t \mapsto e^{it}$  from  $\mathbb{R}$  to  $\mathbb{T}$ . Let  $\rho$  be the action of  $\mathcal{G}_T$  on  $E^\infty \times \mathbb{T}$  from Theorem 7.3. It suffices to fix a neighbourhood  $U$  as above and an  $\varepsilon > 0$  and show that there exists  $\gamma \in (\mathcal{G}_T)_x$  such that  $\rho_\gamma(x, w) \in Z(U) \times B_d(z; \varepsilon)$ . Let  $\mu_y := y(0, n) \in U$ . Since  $\ell$  is continuous, by shrinking  $U$  if necessary, we may assume that

$$d(\ell(\mu), \ell(\mu_y)) < \frac{\varepsilon}{2} \quad \text{for all } \mu \in U. \quad (7.5)$$

Since  $o: E^1 \rightarrow E^0$  is a local homeomorphism, it is an open map, and so  $o(U)$  is open. Since  $\text{Orb}^+(t(x))$  is dense in  $E^0 \times \mathbb{T}$ , we can find  $\lambda \in E^*t(x)$  such that

$$(t(\lambda), \ell(\lambda)) \in o(U) \times B_d(z \overline{\ell(\mu_y)} \bar{w}; \frac{\varepsilon}{2}).$$

Let  $\mu_{t(\lambda)}$  be the unique element of  $U$  such that  $o(\mu_{t(\lambda)}) = t(\lambda)$ . Since  $d$  is rotation-invariant, [Condition \(7.5\)](#) implies that

$$d(\ell(\mu_{t(\lambda)}\lambda)w, \ell(\mu_y)\ell(\lambda)w) = d(\ell(\mu_{t(\lambda)})\ell(\lambda)w, \ell(\mu_y)\ell(\lambda)w) = d(\ell(\mu_{t(\lambda)}), \ell(\mu_y)) < \frac{\varepsilon}{2}. \quad (7.6)$$

Moreover, since  $d$  is rotation-invariant and  $\ell(\lambda) \in B_d(z\overline{\ell(\mu_y)\bar{w}}; \frac{\varepsilon}{2})$ , we have

$$d(\ell(\mu_y)\ell(\lambda)w, z) = d(\ell(\lambda), z\overline{\ell(\mu_y)\bar{w}}) < \frac{\varepsilon}{2}. \quad (7.7)$$

Together, [Equations \(7.6\)](#) and [\(7.7\)](#) imply that

$$d(\ell(\mu_{t(\lambda)}\lambda)w, z) \leq d(\ell(\mu_{t(\lambda)}\lambda)w, \ell(\mu_y)\ell(\lambda)w) + d(\ell(\mu_y)\ell(\lambda)w, z) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \quad (7.8)$$

Now [Equations \(7.4\)](#) and [\(7.8\)](#) imply that

$$\rho_{(\mu_{t(\lambda)}\lambda x, n+|\lambda|, x)}(x, w) = (\mu_{t(\lambda)}\lambda x, \ell(\mu_{t(\lambda)}\lambda)w) \in Z(U) \times B_d(z; \varepsilon),$$

as required.  $\square$

## APPENDIX A. COHOMOLOGY OF GROUPS

In this appendix we provide some background on basic ideas on group cohomology used in [Section 4](#). A much more detailed summary appears in [[3](#), Chapter 2], so we give relatively few details.

Let  $G$  be a countable discrete group with identity  $e_G$ , and let  $A$  be an abelian group with identity  $1_A$  (we write the group operation in  $A$  multiplicatively because our primary example is  $A = \mathbb{T}$ ).

An  $A$ -valued *2-cocycle* on  $G$  is a map  $\sigma: G \times G \rightarrow A$  that satisfies the *2-cocycle identity*:  $\sigma(g, h)\sigma(gh, k) = \sigma(g, hk)\sigma(h, k)$  for all  $g, h, k \in G$ , and is *normalised*, in the sense that  $\sigma(e_G, g) = 1_A = \sigma(g, e_G)$  for all  $g \in G$ . The collection of  $A$ -valued 2-cocycles on  $G$  forms a group under multiplication of functions, which we denote by  $Z^2(G, A)$ .

For every 2-cocycle  $\sigma$  on  $G$ , we have

$$\begin{aligned} \sigma(g, g^{-1}) &= \sigma(g, g^{-1})\sigma(e_G, g) = \sigma(g, g^{-1})\sigma(gg^{-1}, g) \\ &= \sigma(g, g^{-1}g)\sigma(g^{-1}, g) = \sigma(g, e_G)\sigma(g^{-1}, g) = \sigma(g^{-1}, g), \end{aligned}$$

for all  $g \in G$ .

If  $\sigma: G \times G \rightarrow A$  is a 2-cocycle, then a quick calculation shows that the formula

$$\sigma^*(g, h) := \sigma(h, g)^{-1} \quad (\text{A.1})$$

also defines an  $A$ -valued 2-cocycle on  $G$ . If  $\sigma = \sigma^*$ , then we say that  $\sigma$  is *antisymmetric*.

If  $b: G \rightarrow A$  is a function such that  $b(e_G) = 1_A$  (sometimes called a *normalised 1-cochain*), then the formula  $\delta^1 b(g, h) := b(g)b(h)b(gh)^{-1}$  defines a 2-cocycle, called the *2-coboundary* associated to  $b$ . The set  $B^2(G, A)$  of such coboundaries is a subgroup of  $Z^2(G, A)$ .

The quotient  $H^2(G, A) := Z^2(G, A)/B^2(G, A)$  is called the *second cohomology group* of  $G$  with values in  $A$ . Two 2-cocycles  $\sigma, \tau \in Z^2(G, A)$  are said to be *cohomologous* if they represent the same class in  $H^2(G, A)$ ; that is, if  $\sigma^{-1}\tau \in B^2(G, A)$ . Given  $\sigma \in Z^2(G, A)$ , we write  $[\sigma]$  for its coset in  $H^2(G, A)$ .

A *central extension* of  $G$  by  $A$  is an exact sequence  $1_A \rightarrow A \xrightarrow{i} E \xrightarrow{q} G \rightarrow e_G$  such that  $i(A)$  is contained in the centre of  $E$ . A *normalised section* for  $q$  is a map  $s: G \rightarrow E$  such that  $s(e_G) = e_E$  and  $q \circ s = \text{id}_G$ . We say that two central extensions

$$1_A \rightarrow A \xrightarrow{i} E \xrightarrow{q} G \rightarrow e_G \quad \text{and} \quad 1_A \rightarrow A \xrightarrow{i'} E' \xrightarrow{q'} G \rightarrow e_G$$



are *congruent* (or *equivalent*) if there is a homomorphism  $\varphi: E \rightarrow E'$  such that  $\varphi \circ i = i'$  and  $q' \circ \varphi = q$ ; by the Five Lemma, this  $\varphi$  must be an isomorphism of groups. We write  $\text{Ext}(G, A)$  for the collection of congruence classes of central extensions of  $G$  by  $A$ .

Cocycles and central extensions are connected as follows. Given  $\sigma \in Z^2(G, A)$ , we can define a group operation on  $G \times A$  by  $(g, a)(h, b) := (gh, \sigma(g, h)ab)$ . We write  $G \times_\sigma A$  for the group  $G \times A$  under this operation. The formulae  $i(a) := (e_G, a)$  and  $q(g, a) := g$  determine a central extension  $1_A \rightarrow A \xrightarrow{i} G \times_\sigma A \xrightarrow{q} G \rightarrow e_G$ . Conversely, given any normalised section  $s: G \rightarrow E$  and given any  $g, h \in G$ , we have  $q(s(g)s(h)s(gh)^{-1}) = e_G$ , and hence there exists a unique element  $\tau(g, h) \in A$  such that  $i(\tau(g, h)) = s(g)s(h)s(gh)^{-1}$ . The map  $\tau$  is an  $A$ -valued 2-cocycle on  $G$ , and we have the following result.

**Theorem A.1** ([9, Theorem IV.3.12]). *Let  $G$  be a countable discrete group, and let  $A$  be an abelian group. If  $\sigma, \tau \in Z^2(G, A)$  are cohomologous 2-cocycles and  $b: G \rightarrow A$  is a normalised 1-cochain such that  $\sigma = \delta^1 b \tau$ , then the map  $(g, a) \mapsto (g, b(g)a)$  is an isomorphism from  $G \times_\sigma A$  to  $G \times_\tau A$  that induces a congruence of central extensions. There is a bijection  $M: H^2(G, A) \rightarrow \text{Ext}(G, A)$  that sends  $[\sigma]$  to the congruence class of  $G \times_\sigma A$ .*

There is a natural pullback construction (the Baer sum) that describes the group operation on  $\text{Ext}(G, A)$  that makes  $M$  an isomorphism of groups, but the details are not needed here.

We now focus on the situation  $A = \mathbb{T}$  and single out a particular type of 2-cocycle.

**Definition A.2.** Let  $G$  be a countable discrete group. A *bicharacter* of  $G$  is a function  $\omega: G \times G \rightarrow \mathbb{T}$  such that for each fixed  $g \in G$ , the maps  $\omega(\cdot, g): G \rightarrow \mathbb{T}$  and  $\omega(g, \cdot): G \rightarrow \mathbb{T}$  are group homomorphisms.

Every bicharacter of  $G$  is a  $\mathbb{T}$ -valued 2-cocycle:

$$\omega(g, h)\omega(gh, k) = \omega(g, h)\omega(g, k)\omega(h, k) = \omega(g, hk)\omega(h, k), \quad \text{for all } g, h, k \in G.$$

**Theorem A.3** ([28, Proposition 3.2]). *Suppose that  $A$  is a discrete abelian group and that  $\sigma \in Z^2(A, \mathbb{T})$ . With  $\sigma^*$  defined as in Equation (A.1), the 2-cocycle  $\sigma\sigma^*$  is an antisymmetric bicharacter of  $A$ . Two 2-cocycles  $\sigma, \tau \in Z^2(A, \mathbb{T})$  are cohomologous if and only if  $\sigma\sigma^* = \tau\tau^*$ , and the map  $[\sigma] \mapsto \sigma\sigma^*$  is an isomorphism of  $H^2(A, \mathbb{T})$  onto the group of antisymmetric bicharacters of  $A$ .*

It follows that every 2-cocycle on a finitely generated abelian group is cohomologous to a bicharacter. For a detailed proof, see [3, Proposition 2.2.5].

**Corollary A.4.** *Suppose that  $A$  is a finitely generated discrete abelian group and that  $\sigma \in Z^2(A, \mathbb{T})$ . Then there is a bicharacter  $\omega$  of  $A$  such that  $[\sigma] = [\omega]$  in  $H^2(A, \mathbb{T})$ .*

*Proof.* Choose generators  $e_1, \dots, e_l$  for  $A$  such that  $A = \bigoplus_{1 \leq i \leq l} \mathbb{Z}e_i$ . For  $i, j \leq l$ , define

$$\omega(e_i, e_j) := \begin{cases} (\sigma\sigma^*)(e_i, e_j) & \text{if } i \geq j \\ 1 & \text{otherwise.} \end{cases}$$

This formula extends uniquely to a bicharacter of  $A$ . We have  $(\omega\omega^*)(e_i, e_j) = (\sigma\sigma^*)(e_i, e_j)$  for all  $i, j \in \{1, \dots, l\}$  by construction. Since both  $\omega\omega^*$  and  $\sigma\sigma^*$  are bicharacters, it follows that  $\omega\omega^* = \sigma\sigma^*$ . Thus  $[\omega] = [\sigma]$  by Theorem A.3.  $\square$

**Definition A.5.** Let  $A$  be a countable discrete abelian group. For  $\sigma \in Z^2(A, \mathbb{T})$ , we define the *centre* of  $\sigma$  to be the set

$$Z_\sigma := \{a \in A : (\sigma\sigma^*)(a, b) = 1 \text{ for all } b \in A\}.$$

If  $\sigma(a, b) = 1 = \sigma(b, a)$  for all  $a \in Z_\sigma$  and  $b \in A$ , then we say that  $\sigma$  *vanishes on its centre*.



We finish the appendix by showing that every cohomology class for a finitely generated abelian group is represented by a bicharacter that vanishes on its centre. For a detailed proof, see [3, Theorem 2.2.8].

**Theorem A.6.** *Let  $A$  be a finitely generated discrete abelian group. For each  $\sigma \in Z^2(A, \mathbb{T})$ , there exists a bicharacter  $\omega$  of  $A$  such that  $[\sigma] = [\omega]$  in  $H^2(A, \mathbb{T})$  and  $\omega$  vanishes on its centre. There is a bicharacter  $\tilde{\omega} \in Z^2(A/Z_\omega, \mathbb{T})$  such that  $\omega(a, b) = \tilde{\omega}(a + Z_\omega, b + Z_\omega)$  for all  $a, b \in A$ .*

*Proof.* Fix  $\sigma \in Z^2(A, \mathbb{T})$ . By Corollary A.4, there is a bicharacter  $\rho$  of  $A$  that is cohomologous to  $\sigma$ . Theorem A.3 implies that there is an antisymmetric bicharacter  $\tau$  of  $A/Z_\rho$  given by  $\tau(a + Z_\rho, b + Z_\rho) = (\rho\rho^*)(a, b)$  for all  $a, b \in A$ . By the final statement of Theorem A.3, there is a 2-cocycle  $\tilde{\omega}$  of  $A/Z_\rho$  such that  $\tilde{\omega}\tilde{\omega}^* = \tau$ , and by another application of Corollary A.4, we can take  $\tilde{\omega}$  to be a bicharacter. Now  $\omega(a, b) := \tilde{\omega}(a + Z_\rho, b + Z_\rho)$  defines a bicharacter of  $A$ . A quick calculation shows that  $\omega\omega^* = \rho\rho^*$ , and so  $[\omega] = [\rho]$  by Theorem A.3. Moreover,  $Z_\omega = Z_\rho$ , and so  $\omega$  vanishes on its centre by definition.  $\square$

## APPENDIX B. REALISING TWISTED GROUP $C^*$ -ALGEBRAS AS INDUCED ALGEBRAS

In this appendix we describe how to realise twisted group  $C^*$ -algebras as induced algebras, which is a key step in the proof of Theorem 5.4. These results are fairly well known and a detailed treatment is given in [3, Theorem 4.3.1], so we give relatively little detail here. We assume knowledge of  $C(X)$ -algebras (or, more generally,  $C_0(X)$ -algebras). See [36, Section C.1] for the definition and relevant results.

We first recall the definition of the induced algebra of a dynamical system. (See [30, Section 6.3] for more details.)

**Definition B.1.** Let  $G$  be a compact Hausdorff group acting continuously on the right of a locally compact Hausdorff space  $X$ , and let  $\alpha$  be a strongly continuous action of  $G$  on a  $C^*$ -algebra  $D$ . The *induced algebra* of the dynamical system  $(D, G, \alpha)$  is defined by

$$\text{Ind}_G^X(D, \alpha) := \{f \in C_0(X, D) : f(x \cdot g) = \alpha_g^{-1}(f(x)) \text{ for all } x \in X \text{ and } g \in G\}.$$

**Theorem B.2.** *Let  $A$  be a countable discrete abelian group. Suppose that  $\omega \in Z^2(A, \mathbb{T})$  is a bicharacter that vanishes on  $Z_\omega$ , in the sense that  $\omega(Z_\omega, A) \cup \omega(A, Z_\omega) = \{1\}$ . Define  $B := A/Z_\omega$ , and let  $\tilde{\omega} \in Z^2(B, \mathbb{T})$  be the bicharacter satisfying  $\tilde{\omega}(p + Z_\omega, q + Z_\omega) = \omega(p, q)$  for all  $p, q \in A$ . Let  $\{u_p : p \in A\}$  be the canonical family of generating unitaries for the twisted group  $C^*$ -algebra  $C^*(A, \omega)$ , and let  $\{U_{p+Z_\omega} : p + Z_\omega \in B\}$  be the canonical family of generating unitaries for the twisted group  $C^*$ -algebra  $C^*(B, \tilde{\omega})$ .*

(a) *There is a continuous, free, proper right action of  $\widehat{B}$  on  $\widehat{A}$  given by*

$$(\phi \cdot \chi)(p) := \phi(p) \chi(p + Z_\omega) \text{ for all } \phi \in \widehat{A}, \chi \in \widehat{B}, \text{ and } p \in A.$$

*The orbit space  $\widehat{A}/\widehat{B}$  is compact.*

(b) *There is a strongly continuous action  $\beta^B$  of  $\widehat{B}$  on  $C^*(B, \tilde{\omega})$  such that*

$$\beta_\chi^B(U_{p+Z_\omega}) = \chi(p + Z_\omega) U_{p+Z_\omega} \text{ for all } \chi \in \widehat{B} \text{ and } p \in A.$$

(c) *There is an isomorphism  $\Omega : C^*(A, \omega) \rightarrow \text{Ind}_{\widehat{B}}^{\widehat{A}}(C^*(B, \tilde{\omega}), \beta^B)$  such that*

$$\Omega(u_p)(\phi) = \overline{\phi(p)} U_{p+Z_\omega} \text{ for all } p \in A \text{ and } \phi \in \widehat{A}.$$

*In particular,  $C^*(A, \omega)$  is a  $C(\widehat{A}/\widehat{B})$ -algebra.*

A detailed proof of [Theorem B.2](#) can be found in [[3](#), Theorem 4.3.1]. Parts (a) and (b) are routine, but we reproduce some of the details of part (c) below. For this, we need the following preliminary result.

**Lemma B.3.** *Let  $A$  be a countable discrete abelian group, and let  $\omega \in Z^2(A, \mathbb{T})$  be a bicharacter.*

- (a) *There is a strongly continuous action  $\beta^A$  of  $\widehat{A}$  on  $C^*(A, \omega)$  such that  $\beta_\phi^A(u_p) = \phi(p)u_p$  for all  $\phi \in \widehat{A}$  and  $p \in A$ .*
- (b) *There is a faithful conditional expectation  $\Phi^A: C^*(A, \omega) \rightarrow \mathbb{C}1_{C^*(A, \omega)}$  such that*

$$\Phi^A(x) = \int_{\widehat{A}} \beta_\phi^A(x) d\phi \quad \text{for all } x \in C^*(A, \omega).$$

- (c) *Suppose that  $Y$  is a nonzero unital C\*-algebra and  $\Psi: C^*(A, \omega) \rightarrow Y$  is a unital homomorphism. If  $x \in C^*(A, \omega)$  satisfies  $\Psi(\beta_\phi^A(x)) = 0$  for all  $\phi \in \widehat{A}$ , then  $x = 0$ .*

*Proof.* Parts (a) and (b) follow from standard arguments (see [[3](#), Lemmas 4.3.2 and 4.3.4]).

For part (c), fix  $x \in C^*(A, \omega)$  such that  $\Psi(\beta_\phi^A(x)) = 0$  for all  $\phi \in \widehat{A}$ . Then

$$\Psi(\beta_\phi^A(x^*x)) = \Psi(\beta_\phi^A(x))^* \Psi(\beta_\phi^A(x)) = 0,$$

and [[30](#), Lemma C.3] implies that

$$\Psi(\Phi^A(x^*x)) = \Psi\left(\int_{\widehat{A}} \beta_\phi^A(x^*x) d\phi\right) = \int_{\widehat{A}} \Psi(\beta_\phi^A(x^*x)) d\phi = 0. \quad (\text{B.1})$$

Since  $\Psi$  is unital, it is injective on  $\mathbb{C}1_{C^*(A, \omega)}$ , and so we deduce that  $\Phi^A(x^*x) = 0$ . Hence  $x = 0$ , because  $\Phi^A$  is faithful.  $\square$

*Proof of [Theorem B.2\(c\)](#).* Let  $\mathcal{Y}_{A, \omega} := \text{Ind}_{\widehat{B}}^{\widehat{A}}(C^*(B, \tilde{\omega}), \beta^B)$ . We aim to use the universal property of  $C^*(A, \omega)$  to find a homomorphism  $\Omega: C^*(A, \omega) \rightarrow \mathcal{Y}_{A, \omega}$  such that  $\Omega(u_p)(\phi) = \overline{\phi(p)} U_{p+Z_\omega}$  for all  $p \in A$  and  $\phi \in \widehat{A}$ . For each  $p \in A$ , define  $v_p: \widehat{A} \rightarrow C^*(B, \tilde{\omega})$  by  $v_p(\phi) := \overline{\phi(p)} U_{p+Z_\omega}$ . A routine argument shows that each  $v_p$  is continuous and that  $v_p(\phi \cdot \chi) = (\beta_\chi^B)^{-1}(v_p(\phi))$  for all  $\phi \in \widehat{A}$  and  $\chi \in \widehat{B}$ , and hence  $v_p \in \mathcal{Y}_{A, \omega}$ . It is clear that each  $v_p$  is a unitary. For all  $p, q \in A$ , we have

$$U_{p+Z_\omega} U_{q+Z_\omega} = \tilde{\omega}(p+Z_\omega, q+Z_\omega) U_{p+q+Z_\omega} = \omega(p, q) U_{p+q+Z_\omega},$$

and hence for all  $\phi \in \widehat{A}$ , we have

$$(v_p v_q)(\phi) = \overline{\phi(p)} U_{p+Z_\omega} \overline{\phi(q)} U_{q+Z_\omega} = \omega(p, q) \overline{\phi(p+q)} U_{p+q+Z_\omega} = \omega(p, q) v_{p+q}(\phi).$$

Therefore,  $v_p v_q = \omega(p, q) v_{p+q}$ , and so the universal property of  $C^*(A, \omega)$  implies that there is a homomorphism  $\Omega: C^*(A, \omega) \rightarrow \mathcal{Y}_{A, \omega}$  such that  $\Omega(u_p) = v_p$  for each  $p \in A$ .

We first show that  $\Omega$  is surjective. Let  $Z(\mathcal{Y}_{A, \omega})$  denote the centre of  $\mathcal{Y}_{A, \omega}$ . By [[36](#), Proposition 3.49], the unital C\*-algebra  $\mathcal{Y}_{A, \omega}$  is a  $C(\widehat{A}/\widehat{B})$ -algebra with respect to the nondegenerate homomorphism  $\Phi_{\mathcal{Y}_{A, \omega}}: C(\widehat{A}/\widehat{B}) \rightarrow Z(\mathcal{Y}_{A, \omega})$  given by

$$\Phi_{\mathcal{Y}_{A, \omega}}(f)(\phi) = f(\phi \cdot \widehat{B}) 1_{C^*(B, \tilde{\omega})}.$$

For each  $\phi \in \widehat{A}$ , the set

$$I_{\phi \cdot \widehat{B}} := \overline{\text{span}}\{\Phi_{\mathcal{Y}_{A, \omega}}(f)g : f \in C(\widehat{A}/\widehat{B}), g \in \mathcal{Y}_{A, \omega}, f(\phi \cdot \widehat{B}) = 0\}$$

is an ideal of  $\mathcal{Y}_{A, \omega}$ . Define

$$\mathcal{A} := \bigsqcup_{\phi \cdot \widehat{B} \in \widehat{A}/\widehat{B}} \mathcal{Y}_{A, \omega} / I_{\phi \cdot \widehat{B}},$$

and let  $\rho: \mathcal{A} \rightarrow \widehat{A}/\widehat{B}$  be the surjective map given by  $\rho(g + I_{\phi \cdot \widehat{B}}) := \phi \cdot \widehat{B}$ . An application of [36, Proposition C.10(a) and Theorem C.25] shows that there is a unique topology on  $\mathcal{A}$  such that  $(\mathcal{A}, \rho, \widehat{A}/\widehat{B})$  is an upper semicontinuous  $C^*$ -bundle, and that for each  $g \in \mathcal{Y}_{A,\omega}$ , the section  $\phi \cdot \widehat{B} \mapsto g + I_{\phi \cdot \widehat{B}}$  is continuous. Define

$$\Gamma(\mathcal{A}) := \{h: \widehat{A}/\widehat{B} \rightarrow \mathcal{A} : h \text{ is continuous, and } \rho(h(\phi \cdot \widehat{B})) = \phi \cdot \widehat{B} \text{ for all } \phi \in \widehat{A}\},$$

and let  $F: \mathcal{Y}_{A,\omega} \rightarrow \Gamma(\mathcal{A})$  be the map given by  $F(g)(\phi \cdot \widehat{B}) := g + I_{\phi \cdot \widehat{B}}$ , for all  $g \in \mathcal{Y}_{A,\omega}$  and  $\phi \in \widehat{A}$ . By [36, Theorem C.26],  $F$  is a  $C(\widehat{A}/\widehat{B})$ -linear isomorphism of  $\mathcal{Y}_{A,\omega}$  onto the  $C(\widehat{A}/\widehat{B})$ -algebra  $\Gamma(\mathcal{A})$ . An application of [36, Proposition C.24] shows that  $F(\Omega(C^*(A, \omega)))$  is a dense subspace of  $\Gamma(\mathcal{A})$ . Since  $F$  is an isomorphism, it follows that  $\Omega(C^*(A, \omega))$  is dense in  $\mathcal{Y}_{A,\omega}$ , and hence  $\Omega$  is surjective.

To see that  $\Omega$  is injective, let  $\beta^A$  be the strongly continuous action of Lemma B.3(a). For each  $\phi \in \widehat{A}$ , the map  $v_p \mapsto \phi(p)v_p$  extends to an automorphism  $\alpha_\phi^A$  of  $\mathcal{Y}_{A,\omega}$  satisfying  $\alpha_\phi^A \circ \Omega = \Omega \circ \beta_\phi^A$ . If  $x \in C^*(A, \omega)$  satisfies  $\Omega(x) = 0$ , then for all  $\phi \in \widehat{A}$ , we have  $\Omega(\beta_\phi^A(x)) = \alpha_\phi^A(\Omega(x)) = 0$ , and then Lemma B.3(c) gives  $x = 0$ . Hence  $\Omega$  is injective. Thus  $\Omega$  is an isomorphism, and  $C^*(A, \omega)$  is a  $C(\widehat{A}/\widehat{B})$ -algebra with respect to the homomorphism  $\Omega^{-1} \circ \Phi_{\mathcal{Y}_{A,\omega}}$ .  $\square$

## REFERENCES

- [1] C. Anantharaman-Delaroche and J. Renault, *Amenable Groupoids (with a foreword by Georges Skandalis and Appendix B by E. Germain)*, Monographies de L'Enseignement Mathématique, vol. 36, L'Enseignement Mathématique, Geneva, 2000.
- [2] B. Armstrong, *Twisted topological higher-rank graphs and their  $C^*$ -algebras*, Honours thesis, University of Wollongong, Australia, 2014.
- [3] ———, *Simplicity of twisted  $C^*$ -algebras of topological higher-rank graphs*, Ph.D. thesis, The University of Sydney, Australia, August 2019.
- [4] ———, *A uniqueness theorem for twisted groupoid  $C^*$ -algebras*, preprint, 2021, [arXiv:2103.03063v2](https://arxiv.org/abs/2103.03063v2) [[math.OA](https://arxiv.org/abs/2103.03063v2)].
- [5] B. Armstrong and N. Brownlowe, *Product-system models for twisted  $C^*$ -algebras of topological higher-rank graphs*, J. Math. Anal. Appl. **466** (2018), 1443–1475, DOI: [10.1016/j.jmaa.2018.06.052](https://doi.org/10.1016/j.jmaa.2018.06.052).
- [6] E. Bédos and T. Omland, *On twisted group  $C^*$ -algebras associated with FC-hypercentral groups and other related groups*, Ergodic Theory Dynam. Systems **36** (2016), 1743–1756, DOI: [10.1017/etds.2015.9](https://doi.org/10.1017/etds.2015.9).
- [7] J.H. Brown, L.O. Clark, C. Farthing, and A. Sims, *Simplicity of algebras associated to étale groupoids*, Semigroup Forum **88** (2014), 433–452, DOI: [10.1007/s00233-013-9546-z](https://doi.org/10.1007/s00233-013-9546-z).
- [8] J.H. Brown, L.O. Clark, A. Sierakowski, and A. Sims, *Purely infinite simple  $C^*$ -algebras that are principal groupoid  $C^*$ -algebras*, J. Math. Anal. Appl. **439** (2016), 213–234, DOI: [10.1016/j.jmaa.2016.02.055](https://doi.org/10.1016/j.jmaa.2016.02.055).
- [9] K.S. Brown, *Cohomology of Groups*, Graduate Texts in Mathematics, vol. 87, Springer, New York, 1982, DOI: [10.1007/978-1-4684-9327-6](https://doi.org/10.1007/978-1-4684-9327-6).
- [10] T.M. Carlsen, E. Ruiz, A. Sims, and M. Tomforde, *Reconstruction of groupoids and  $C^*$ -rigidity of dynamical systems*, Adv. Math. **390** (2021), 1–55, DOI: [10.1016/j.aim.2021.107923](https://doi.org/10.1016/j.aim.2021.107923).
- [11] V. Deaconu, *Groupoids associated with endomorphisms*, Trans. Amer. Math. Soc. **347** (1995), 1779–1786, DOI: [10.1090/s0002-9947-1995-1233967-5](https://doi.org/10.1090/s0002-9947-1995-1233967-5).
- [12] R. Engelking, *General Topology*, 2nd ed., Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989.
- [13] R. Exel, *Inverse semigroups and combinatorial  $C^*$ -algebras*, Bull. Braz. Math. Soc. (N.S.) **39** (2008), 191–313, DOI: [10.1007/s00574-008-0080-7](https://doi.org/10.1007/s00574-008-0080-7).
- [14] G.B. Folland, *A Course in Abstract Harmonic Analysis*, 2nd ed., Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2016, DOI: [10.1201/b19172](https://doi.org/10.1201/b19172).

- [15] G. Goehle, *Groupoid crossed products*, Ph.D. thesis, Dartmouth College, Hanover, NH, May 2009, [arXiv:0905.4681v1](https://arxiv.org/abs/0905.4681v1) [[math.OA](https://arxiv.org/abs/0905.4681v1)].
- [16] A. an Huef, A. Kumjian, and A. Sims, *A Dixmier–Douady theorem for Fell algebras*, *J. Funct. Anal.* **260** (2011), 1543–1581, DOI: [10.1016/j.jfa.2010.11.011](https://doi.org/10.1016/j.jfa.2010.11.011).
- [17] T. Katsura, *A class of  $C^*$ -algebras generalizing both graph algebras and homeomorphism  $C^*$ -algebras I, fundamental results*, *Trans. Amer. Math. Soc.* **356** (2004), 4287–4322, DOI: [10.1090/s0002-9947-04-03636-0](https://doi.org/10.1090/s0002-9947-04-03636-0).
- [18] A. Kumjian and D. Pask, *Higher rank graph  $C^*$ -algebras*, *New York J. Math.* **6** (2000), 1–20.
- [19] A. Kumjian, D. Pask, and I. Raeburn, *Cuntz–Krieger algebras of directed graphs*, *Pacific J. Math.* **184** (1998), 161–174, DOI: [10.2140/pjm.1998.184.161](https://doi.org/10.2140/pjm.1998.184.161).
- [20] A. Kumjian, D. Pask, I. Raeburn, and J. Renault, *Graphs, groupoids, and Cuntz–Krieger algebras*, *J. Funct. Anal.* **144** (1997), 505–541, DOI: [10.1006/jfan.1996.3001](https://doi.org/10.1006/jfan.1996.3001).
- [21] A. Kumjian, D. Pask, and A. Sims, *Homology for higher-rank graphs and twisted  $C^*$ -algebras*, *J. Funct. Anal.* **263** (2012), 1539–1574, DOI: [10.1016/j.jfa.2012.05.023](https://doi.org/10.1016/j.jfa.2012.05.023).
- [22] ———, *On the  $K$ -theory of twisted higher-rank-graph  $C^*$ -algebras*, *J. Math. Anal. Appl.* **401** (2013), 104–113, DOI: [10.1016/j.jmaa.2012.11.030](https://doi.org/10.1016/j.jmaa.2012.11.030).
- [23] ———, *On twisted higher-rank graph  $C^*$ -algebras*, *Trans. Amer. Math. Soc.* **367** (2015), 5177–5216, DOI: [10.1090/s0002-9947-2015-06209-6](https://doi.org/10.1090/s0002-9947-2015-06209-6).
- [24] ———, *Simplicity of twisted  $C^*$ -algebras of higher-rank graphs and crossed products by quasifree actions*, *J. Noncommut. Geom.* **10** (2016), 515–549, DOI: [10.4171/jncg/241](https://doi.org/10.4171/jncg/241).
- [25] M. Laca and S. Neshveyev, *KMS states of quasi-free dynamics on Pimsner algebras*, *J. Funct. Anal.* **211** (2004), 457–482, DOI: [10.1016/j.jfa.2003.08.008](https://doi.org/10.1016/j.jfa.2003.08.008).
- [26] H. Li, D. Pask, and A. Sims, *An elementary approach to  $C^*$ -algebras associated to topological graphs*, *New York J. Math.* **20** (2014), 447–469.
- [27] X. Li, *Every classifiable simple  $C^*$ -algebra has a Cartan subalgebra*, *Invent. Math.* **219** (2020), 653–699, DOI: [10.1007/s00222-019-00914-0](https://doi.org/10.1007/s00222-019-00914-0).
- [28] D. Olesen, G.K. Pedersen, and M. Takesaki, *Ergodic actions of compact abelian groups*, *J. Operator Theory* **3** (1980), 237–269.
- [29] I. Raeburn and J.L. Taylor, *Continuous trace  $C^*$ -algebras with given Dixmier–Douady class*, *J. Austral. Math. Soc. Ser. A* **38** (1985), 394–407, DOI: [10.1017/s1446788700023661](https://doi.org/10.1017/s1446788700023661).
- [30] I. Raeburn and D.P. Williams, *Morita Equivalence and Continuous-Trace  $C^*$ -Algebras*, *Math. Surveys and Monographs*, vol. 60, Amer. Math. Soc., Providence, RI, 1998, DOI: [10.1090/surv/060](https://doi.org/10.1090/surv/060).
- [31] J. Renault, *A Groupoid Approach to  $C^*$ -Algebras*, *Lecture Notes in Math.*, vol. 793, Springer, Berlin, 1980, DOI: [10.1007/bfb0091072](https://doi.org/10.1007/bfb0091072).
- [32] ———, *The ideal structure of groupoid crossed product  $C^*$ -algebras (with an appendix by Georges Skandalis)*, *J. Operator Theory* **25** (1991), 3–36.
- [33] A. Sims, *Hausdorff étale groupoids and their  $C^*$ -algebras*, *Operator algebras and dynamics: groupoids, crossed products, and Rokhlin dimension* (F. Perera, ed.), *Advanced Courses in Mathematics*, CRM Barcelona, Birkhäuser, 2020, DOI: [10.1007/978-3-030-39713-5](https://doi.org/10.1007/978-3-030-39713-5).
- [34] A. Sims, B. Whitehead, and M.F. Whittaker, *Twisted  $C^*$ -algebras associated to finitely aligned higher-rank graphs*, *Documenta Math.* **19** (2014), 831–866.
- [35] A. Sims and D.P. Williams, *The primitive ideals of some étale groupoid  $C^*$ -algebras*, *Algebr. Represent. Theor.* **19** (2016), 255–276, DOI: [10.1007/s10468-015-9573-4](https://doi.org/10.1007/s10468-015-9573-4).
- [36] D.P. Williams, *Crossed Products of  $C^*$ -Algebras*, *Math. Surveys and Monographs*, vol. 134, Amer. Math. Soc., Providence, RI, 2007, DOI: [10.1090/surv/134](https://doi.org/10.1090/surv/134).
- [37] T. Yeend, *Topological higher-rank graphs and the  $C^*$ -algebras of topological 1-graphs*, *Operator theory, operator algebras, and applications*, *Contemp. Math.*, vol. 414, Amer. Math. Soc., Providence, RI, 2006, DOI: [10.1090/conm/414/07812](https://doi.org/10.1090/conm/414/07812), pp. 231–244.
- [38] ———, *Groupoid models for the  $C^*$ -algebras of topological higher-rank graphs*, *J. Operator Theory* **57** (2007), 95–120.

(B. Armstrong) MATHEMATICAL INSTITUTE, WWU MÜNSTER, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY

*Email address:* [becky.armstrong@uni-muenster.de](mailto:becky.armstrong@uni-muenster.de)

(N. Brownlowe) SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA

*Email address:* [nathan.brownlowe@sydney.edu.au](mailto:nathan.brownlowe@sydney.edu.au)

(A. Sims) SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOLLONGONG, NSW 2522, AUSTRALIA

*Email address:* [asims@uow.edu.au](mailto:asims@uow.edu.au)