

The essential p -dimension of finite simple groups of Lie type

Hannah Knight*
 Mathematics, UCI, Irvine, CA, USA
hknight1@uci.edu

Abstract

In this paper, we compute the essential p -dimension of the finite simple groups of classical Lie type at the defining prime, specifically the simple groups arising from the linear groups, the symplectic groups, the orthogonal groups, and the unitary groups.

Contents

1	Introduction	2
1.1	General Outline for Proofs	5
2	Essential p-Dimension Background	6
3	Representation Theory Background	7
3.1	The Wigner-Mackey Little Group Method	8
4	PSLn	9
4.1	Finding the p -Sylows and identifying the center	10
4.2	Identifying the generalized Heisenberg subgroup	10
4.3	Calculating $\text{ed}_k(H_n)$	10
4.3.1	Maximizing L_s	11
4.3.2	Proof	11
4.4	Showing that $\text{ed}_k(\text{Up}_n(\mathbb{F}_{p^r})) = \text{ed}_k(H_n(\mathbb{F}_{p^r}))$	12
4.4.1	Proof	12
5	The Symplectic Groups	13
5.1	Definitions	13
5.2	The p -Sylows	13
5.3	Calculating the center	14
5.3.1	$p \neq 2$	14
5.3.2	$p = 2$	16
5.4	Maximizing L_s	18
5.4.1	$p \neq 2$	19
5.4.2	$p = 2$	20
5.5	Proof	25

*This work was supported in part by NSF Grant Nos. DMS-1811846 and DMS-1944862.

6	The Orthogonal Groups	26
6.1	Definitions	26
6.1.1	$n = 2m, p \neq 2$	26
6.1.2	$n = 2m, p = 2$	27
6.1.3	$n = 2m + 1$	28
6.2	The p -Sylows	28
6.2.1	$n = 2m$	29
6.2.2	$n = 2m + 1, p \neq 2$	31
6.3	Calculating the center	32
6.3.1	$n = 4, \text{ any } p$	32
6.3.2	$n = 2m, n > 4, \text{ any } p$	33
6.3.3	$n = 2m + 1, p \neq 2$	36
6.4	Maximizing L_s	38
6.4.1	$n = 2m$	38
6.4.2	$n = 2m + 1, p \neq 2$	42
6.5	Proof	45
7	The Unitary Groups	47
7.1	Definitions	47
7.2	The p -Sylows	48
7.2.1	$n = 2m$	48
7.2.2	$n = 2m + 1$	49
7.3	Calculating the center	51
7.3.1	$n = 2m$	51
7.3.2	$n = 2m + 1$	52
7.4	Maximizing L_s	53
7.4.1	$n = 2m$	53
7.4.2	$n = 2m + 1$	57
7.5	Proof	62

1 Introduction

The goal of this paper is to compute the essential p -dimension at the defining prime for the finite groups of classical Lie type. Fix a field k . The essential dimension of a finite group G is the smallest number of algebraically independent parameters needed to define a Galois G -algebra over any field extension of k . In other words, the essential dimension of a finite group G is the supremum taken over all field extensions F/k of the smallest number of algebraically independent parameters needed to define a Galois G -algebra over F . The essential p -dimension of a finite group is similar, except that before taking the supremum, we first allow finite extensions of F of degree prime to p and take the minimum of the number of parameters needed. In other words, the essential p -dimension of a finite group is the supremum taken over all fields F/k of the smallest number of algebraically independent parameters needed to define a Galois G -algebra over a field extension of F of degree prime to p . See the Essential p -Dimension Background Section for more formal definitions. See also [5] and [12] for more detailed discussions.

Question 1. What is the essential dimension of the finite simple groups?

This question is quite difficult to answer. A few results for small groups (not necessarily simple) have been proven. For example, it is known that $\text{ed}_k(S_5) = 2$, $\text{ed}_k(S_6) = 3$ for k of characteristic not 2 [3], and $\text{ed}_k(A_7) = \text{ed}_k(S_7) = 4$ in characteristic 0 [6]. It is also known that for k a field of characteristic 0 containing all roots of unity, $\text{ed}_k(G) = 1$ if and only if G is isomorphic to a cyclic group $\mathbb{Z}/n\mathbb{Z}$ or a dihedral

group D_m where m is odd ([5], Theorem 6.2). Various bounds have also been proven. See [5], [15], [23],[17], among others. For a nice summary of the results known in 2010, see [23].

We can find a lower bound to this question by considering the corresponding question for essential p -dimension.

Theorem 1.1. *Let p be a prime, k a field with $\text{char } k \neq p$. Then*

(1) (Theorem 4.1)

$$\text{ed}_k(PSL_n(\mathbb{F}_{p^r}), p) = \text{ed}_k(GL_n(\mathbb{F}_{p^r})) = rp^{r(n-2)}.$$

(2) (Theorem 5.1)

$$\text{ed}_k(PSp(2n, p^r)', p) = \text{ed}_k(Sp(2n, p^r)) = \begin{cases} rp^{r(n-1)}, & p \neq 2 \text{ or } p = 2, n = 2 \\ r2^{r(n-1)-1}(2^{r(n-2)} + 1), & p = 2, n > 2 \end{cases}$$

(3) (Theorem 6.1)

$$\text{ed}_k(P\Omega^\epsilon(n, p^r), p) = \text{ed}_k(O^\epsilon(n, p^r), p) = \begin{cases} 2r, & n = 4, \text{ any } p \\ rp^{2r(m-2)}, & n = 2m, n > 4, \text{ any } p \\ rp^{r(m-1)} + rp^{r(m-1)(m-2)}, & n = 2m + 1, p \neq 2 \end{cases}$$

(4) (Theorem 7.1)

$$\text{ed}_k(PSU(n, p^{2r}), p) = \text{ed}_k(U(n, p^{2r}), p) = \begin{cases} rp^{2r(m-1)} & n = 2m \\ 3rp^{2r(m-1)} & n = 2m + 1 \end{cases}$$

Remark 1. For $p = 2, n = 2, r = 1$, we have $PSp(4, 2)' \cong A_6$, and so $\text{ed}_k(PSp(4, 2)', 2) = \text{ed}_k(A_6, 2) = 2$. Note that $\text{ed}_k(Sp(2n, p^r), p) = \text{ed}_k(PSp(2n, p^r), p)$ (see [9], Lemma 2.2.7). Except for $p = 2, n = 2, r = 1$, $PSp(2n, p^r) = PSp(2n, p^r)'$ is simple. This methods of this paper can recover the proof that $\text{ed}_k(PSp(4, 2), 2) = \text{ed}_k(S_6, 2) = 3$ and that $\text{ed}_k(PSp(4, 2)', 2) = \text{ed}_k(A_6, 2) = 2$, but for brevity, because these are known theorems, we will omit the proofs here.

Remark 2. Note that if $\text{char } k = p$, then $\text{ed}_k(G, p) = 1$ unless $p \nmid |G|$, in which case $\text{ed}_k(G, p) = 0$ [25]. So we have a full description of $\text{ed}_k(G, p)$ for the finite simple groups of Lie type for any field k and any p .

Remark 3. Dave Benson independently proved $\text{ed}_{\mathbb{C}}(Sp(2n, p), p) = p^{n-1}$ for p odd ([4], Appendix A).

Remark 4. The following results were known prior to this paper:

1. $\text{ed}_{\mathbb{C}}(PSL_n(\mathbb{F}_{p^r}), p) = \text{ed}_{\mathbb{C}}(GL_n(\mathbb{F}_{p^r})) = rp^{r(n-2)}$ for $p \neq 2$ ([1], Theorems 1.1 and 1.2).
2. Duncan and Reichstein calculated the essential p -dimension of the pseudo-reflection groups: For G a pseudo-reflection group with $k[V]^G = k[f_1, \dots, f_n]$, $d_i = \deg(f_i)$, $\text{ed}_k(G, p) = a(p) = |\{i : d_i \text{ is divisible by } p\}|$ ([7], Theorem 1.1). These groups overlap with the groups above in a few small cases (The values of d_i are in [27], Table VII):

(i) Group 12 in the Shephard-Todd classification, $Z_2.O \cong GL_2(\mathbb{F}_3)$: d_1, d_2 are 6, 8; so

$$\text{ed}_k(Z_2.O, 3) = 1 = \text{ed}_k(GL_2(\mathbb{F}_3), 3).$$

(ii) Group 23 in the Shephard-Todd classification, $W(H_3) \cong \mathbb{Z}/2\mathbb{Z} \times PSL_2(\mathbb{F}_5)$: d_1, \dots, d_3 are 2, 6, 10; so

$$\text{ed}_k(W(H_3), 5) = 1 = \text{ed}_k(PSL_2(\mathbb{F}_5), 5).$$

- (iii) Group 32 in the Shephard-Todd classification, $W(L_4) \cong \mathbb{Z}/3\mathbb{Z} \times Sp(4, 3)$: d_1, \dots, d_4 are 12, 18, 24, 30; so

$$\text{ed}_k(W(L_4), 3) = 4 = 1 + \text{ed}_k(Sp(4, 3), 3).$$

- (iv) Group 33 in the Shephard-Todd classification, $W(K_5) \cong \mathbb{Z}/2\mathbb{Z} \times PSp(4, 3) \cong \mathbb{Z}/2\mathbb{Z} \times PSU(4, 2)$: d_1, \dots, d_5 are 4, 6, 10, 12, 18; so

$$\text{ed}_k(W(K_5), 3) = 3 = \text{ed}_k(PSp(4, 3), 3)$$

and

$$\text{ed}_k(W(K_5), 2) = 5 = 1 + \text{ed}_k(PSU(4, 2)).$$

- (v) Group 35 in the Shephard-Todd classification, $W(E_6) \cong O^-(6, 2)$: d_1, \dots, d_6 are 2, 5, 6, 8, 9, 12; so

$$\text{ed}_k(W(E_6), 2) = 4 = \text{ed}_k(O^-(6, 2), 2).$$

- (vi) Group 36 in the Shephard-Todd classification, $W(E_7) \cong \mathbb{Z}/2\mathbb{Z} \times Sp(6, 2)$: d_1, \dots, d_7 are 2, 6, 8, 10, 12, 14, 18; so

$$\text{ed}_k(W(E_7), 2) = 7 = 1 + \text{ed}_k(Sp(6, 2), 2).$$

3. Reichstein and Shukla calculated the essential 2-dimension of double covers of the symmetric and alternating groups in characteristic $\neq 2$: Write $n = 2^{a_1} + \dots + 2^{a_s}$, where $a_1 > a_2 > \dots > a_s \geq 0$. For \tilde{S}_n a double cover of S_n , $\text{ed}_k(\tilde{S}_n, 2) = 2^{\lfloor (n-s)/2 \rfloor}$. and for \tilde{A}_n a double cover of A_n , $\text{ed}_k(\tilde{A}_n, 2) = 2^{\lfloor (n-s-1)/2 \rfloor}$. ([24], Theorem 1.2). These groups overlap with the groups above in a few small cases: $\tilde{A}_4 \cong SL_2(3)$, $\tilde{A}_5 \cong SL_2(5)$, $\tilde{A}_6 \cong SL_2(9)$, $\tilde{S}_4^+ \cong GL_2(3)$. However, in each case the defining prime is not 2, and as I only calculate the essential dimension at the defining prime, there is no overlap with the results of Reichstein and Shukla.

The key tools in the proofs of Theorem 1.1 are the Karpenko-Merkurjev Theorem (Theorem 2.9), a lemma of Meyer and Reichstein (Lemma 3.4), and Wigner Mackey Theory.

Theorem 1.2 (Karpenko-Merkurjev [12], Theorem 4.1, See Theorem 2.9 below). *Let G be a p -group, k a field with $\text{char } k \neq p$ containing a primitive p th root of unity. Then $\text{ed}_k(G, p) = \text{ed}_k(G)$ and $\text{ed}_k(G, p)$ coincides with the least dimension of a faithful representation of G over k .*

The Karpenko-Merkurjev Theorem allows us to translate the question formulated in terms of extensions and transcendence degree into a question of representation theory of p -Sylow subgroups.

Lemma 1.3 ([16], Lemma 2.3; See Lemma 3.4 below). *Let H be a finite p -group and let ρ be a faithful representation of H of minimal dimension. Then ρ decomposes as a direct sum of exactly $r = \text{rank}(Z(H))$ irreducible representations*

$$\rho = \rho_1 \oplus \dots \oplus \rho_r.$$

and if χ_i are the central characters of ρ_i , then $\{\chi_i|_{\Omega_1(Z(H))}\}$ is a basis for $\widehat{\Omega_1(Z(H))}$. ($\Omega_1(Z(H))$ is defined to be the largest elementary abelian p -group contained in $Z(H)$; see Definition 3.1.)

This lemma allows us to translate a question of analyzing faithful representations into a question of analyzing irreducible representations. Our main tool for the case at hand is Wigner-Mackey Theory. This method from representation theory allows us to classify the irreducible representations for groups of the form $\Delta \rtimes L$ with Δ abelian. (See section 3.1.)

1.1 General Outline for Proofs

By the Karpenko-Merkurjev Theorem (Theorem 2.9), we need to find the minimal dimension of a faithful representation of the p -Sylows. By Lemma 3.4, if the center of the p -Sylow has rank s , a faithful representation ρ of the p -Sylow of minimal dimension decomposes as a direct sum

$$\rho = \rho_1 \oplus \cdots \oplus \rho_s$$

of exactly s irreducibles, and if χ_i are the central characters of ρ_i , then $\{\chi_i|_{\Omega_1(Z(G))}\}$ is a basis for $\widehat{\Omega_1(Z(G))}$ (where $\Omega_1(Z(G))$ is the largest elementary abelian p -group contained in $Z(G)$; see Definition 3.1). If for some i , ρ_i has a trivial central character χ_i , then $\chi_i|_{\Omega_1(Z(G))}$ would also be trivial, which would contradict the previous sentence. Therefore, each of the ρ_i must have non-trivial central character. Let χ_i be the central characters of ρ_i . Denote the center by Z . In each of the following cases, Z is a finite elementary abelian group, and so $\widehat{\Omega_1(Z)} = Z$. So we have that $\{\chi_i|_Z\}$ is a basis for \widehat{Z} .

Our proofs will follow the following steps:

- Step 1: Find the p -Sylows and their centers

By inspection, we can write each of the following p -Sylows in the form required for Wigner-Mackey: $\Delta \rtimes L$. By Wigner-Mackey theory, we get that irreducible representations come from extending characters of Δ . We will calculate the minimum dimensions of these irreducible representations.

Recall that $L_s = \text{Stab}_L(\chi_s)$ and $G_s = \Delta \cdot L_s$. χ_s is extended to G_s by setting

$$\chi_s(al) = \chi_s(a), \text{ for } a \in \Delta, l \in L_s.$$

And for λ an irreducible representation of L_s , we can extend to G_s by

$$\lambda(al) = \lambda(l), \text{ for } a \in \Delta, l \in L_s.$$

And $\theta_{s,\lambda} = \text{Ind}_{G_s}^G(\chi_s \otimes \lambda)$.

The dimension is given by

$$\begin{aligned} \dim(\theta_{s,\lambda}) &= \dim(\text{Ind}_{G_s}^G \chi_s \otimes \lambda) \\ &= [G : G_s] \dim(\chi_s \otimes \lambda) \\ &= \frac{|L|}{|L_s|} \dim(\lambda), \text{ since } \dim(\chi_s) = 1 \end{aligned}$$

If we pick λ to be the trivial representation, then this will minimize the dimension of the representation. And for λ trivial, we have $\dim(\lambda) = 1$. And so we have that for λ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|}.$$

So the dimension of the representation will be minimized when $|L_s|$ is maximized.

- Step 2: Maximize L_s .
- Step 3: Proof: We proceed to construct upper and lower bounds and prove they are equal.

Find the Lower Bound: By Corollary 3.4, for minimal dimensional faithful representations with central characters χ_i , we must have that $\{\chi_i|_Z\}$ is a basis for \widehat{Z} . We use this fact and the classification in Step 2 to get a lower bound for the dimension of a minimal dimensional faithful representation, and thus a lower bound for the essential p -dimension.

Construct the Upper Bound: We then construct a faithful representation of the p -Sylow to get an upper bound equal to the lower bound. We do this in the following manner: We will create a basis for the characters of the center and extend these to representations of the p -Sylow via Wigner-Mackey. We pick them via the classification in Step 2 such that when we take the direct sum of these representations, we will get a representation of the desired dimension. We then use Lemma 3.3 to show that the direct sum is a faithful representation. And thus we get the desired upper bound.

Remark 5. A more recent paper of Bardestani, Mallahi-Karai, and Salmasian [2] gives a way of calculating the essential dimension by means of commutator matrices. We do not pursue this method here, but rather pursue the method outlined above.

Acknowledgements: I would like to thank Jesse Wolfson for his kind mentorship and generous support. I am also grateful to Vladimir Chernousev, Alexander Duncan, Najmuddin Fakhruddin, Nate Harman, Keivan Mallahi-Karia, Mark MacDonald, Zinovy Reichstein, Hadi Salmasian, Federico Scavia, Jean-Pierre Serre, Alex Sutherland, and Burt Totaro for very helpful comments on a draft.

2 Essential p -Dimension Background

Fix a field k . In the introduction, we defined the following:

Definition 2.1. Let $T : \text{Fields}/k \rightarrow \text{Sets}$ be a functor. Let F/k be a field extension, and $t \in T(F)$. The *essential dimension of t* is

$$\text{ed}_k(t) = \min_{F' \subset F \text{ s.t. } t \in \text{Im}(T(F') \rightarrow T(F))} \text{trdeg}_k(F').$$

Definition 2.2. Let $T : \text{Fields}/k \rightarrow \text{Sets}$ be a functor. The *essential dimension of T* is

$$\text{ed}_k(T) = \sup_{t \in T(F), F/k \in \text{Fields}/k} \text{ed}_k(t).$$

Definition 2.3. Let G be a finite group. Let $H'(-; G) : \text{Fields}/k \rightarrow \text{Sets}$ be defined by $H'(-; G)(F/k) = \{\text{the isomorphism classes of } G\text{-torsors over } \text{Spec}F\}$. The *essential dimension of G* is

$$\text{ed}_k(G) = \text{ed}_k(H'(-; G)).$$

Definition 2.4. Let $T : \text{Fields}/k \rightarrow \text{Sets}$ be a functor. Let F/k be a field extension, and $t \in T(F)$. The *essential p -dimension of t* is

$$\text{ed}_k(t, p) = \min_{F'' \subset F', F \subset F' \text{ s.t. } p \nmid [F':F] \text{ and the image of } t \text{ in } T(F') \text{ is in } \text{Im}(T(F'') \rightarrow T(F'))} \text{trdeg}_k(F'').$$

Note. $\text{ed}_k(t, p) = \min_{F \subset F', p \nmid [F':F]} \text{ed}_k(t|_{F'})$.

Definition 2.5. Let $T : \text{Fields}/k \rightarrow \text{Sets}$ be a functor. The *essential p -dimension of T* is

$$\text{ed}_k(T, p) = \sup_{t \in T(F), F/k \in \text{Fields}/k} \text{ed}_k(t, p).$$

Definition 2.6. Let G be a finite group, p a prime. Let $H'(-; G) : \text{Fields}/k \rightarrow \text{Sets}$ be defined by $H'(-; G)(F/k) = \{\text{the isomorphism classes of } G\text{-torsors over } \text{Spec}F\}$. The *essential p -dimension of G* is

$$\text{ed}_k(G, p) = \text{ed}_k(H'(-; G), p).$$

Let G be a finite group, p a prime. The statements in the following lemma follows directly from the definitions:

Lemma 2.7. (1) $ed_k(G, p) \leq ed_k(G)$.

(2) If $H \subset G$, then $ed_k(H, p) \leq ed_k(G, p)$.

(3) If $H \subset G$ is a subgroup of index prime to p , then $ed_k(G, p) = ed_k(H, p)$.

(4) Let G be a finite group, p a prime, $Syl_p(G)$ a p -Sylow of G . Then

$$ed_k(G, p) = ed_k(Syl_p(G), p).$$

Lemma 2.8 ([12], Remark 4.8). Let G be a finite p -group, k a field of characteristic $\neq p$, k_1/k a finite field extension of degree prime to p . Then

$$ed_k(G, p) = ed_{k_1}(G, p).$$

Remark 6. The idea for this lemma was brought to my attention by Federico Scavia and Zinovy Reichstein.

Note. Let ζ be a primitive p -th root of unity. Then ζ is a root of the polynomial of the form $x^p - 1 = (x - 1)(1 + \dots + x^{p-1})$. Then the minimal polynomial over a field of characteristic prime to p divides $1 + \dots + x^{p-1}$ and so has degree prime to p . So for F a field, we have that $p \nmid [F(\zeta) : F]$. Therefore, in calculating the essential p -dimension over a field k of characteristic $\neq p$, we may assume that k contains a primitive p -th root of unity. Thus to use the result of Karpenko-Merkurjev below, we need only assume that the characteristic of k is not p .

The key theorem allowing us to work with representation theory is the following:

Theorem 2.9 (Karpenko-Merkurjev [12], Theorem 4.1). Let G be a p -group, k a field with char $k \neq p$ containing a primitive p th root of unity. Then $ed_k(G, p) = ed_k(G)$ and $ed_k(G, p)$ coincides with the least dimension of a faithful representation of G over k .

Putting this together with Lemma 2.8, we get

Corollary 2.10. Let G be a finite group, p a prime, $Syl_p(G)$ a p -Sylow of G , k a field with char $k \neq p$. Then

$$ed_k(G, p) = ed_k(Syl_p(G), p) = ed_k(Syl_p(G)).$$

We will also use the following theorem of Karpenko-Merkurjev

Theorem 2.11 (Karpenko-Merkurjev [12], Theorem 5.1). Let G_1 and G_2 be two p -groups, k a field with char $k \neq p$, then

$$ed_k(G_1 \times G_2) = ed_k(G_1) + ed_k(G_2).$$

Corollary 2.12. For k a field with char $k \neq p$, $ed_k((\mathbb{Z}/p\mathbb{Z})^r, p) = r$

Corollary 2.13. Let G be a finite abelian p -group, k a field with char $k \neq p$. Then $ed_k(G) = \text{rank}(G) = ed_k(G, p)$.

3 Representation Theory Background

We will need the following representation theory background:

Definition 3.1. Let H be a p -group. Define $\Omega_1(Z(H))$ (also called the socle of H) to be the largest elementary abelian p -group contained in $Z(H)$, i.e. $\Omega_1(Z(H)) = \{z \in Z(H) : z^p = 1\}$.

Definition 3.2. For G an abelian group, let \widehat{G} denote the group of characters of G (homomorphisms from G to k^\times). We will use the notation $\widehat{\Omega}_1(Z(H))$ for the character group of $\Omega_1(Z(H))$.

The next two lemmas are due to Meyer-Reichstein [16] and reproduced in [1].

Lemma 3.3 ([16], Lemma 2.3). *Let H be a finite p -group and let $(\rho_i, V_i)_{1 \leq i \leq n}$ be a family of irreducible representations of H with central characters χ_i . Suppose that $\{\chi_i|_{\Omega_1(Z(H))} : 1 \leq i \leq n\}$ spans $\widehat{\Omega}_1(Z(H))$. Then $\bigoplus_i \rho_i$ is a faithful representation of H .*

Lemma 3.4 ([16], Lemma 2.3). *Assume that $\text{char } k \neq p$. Let H be a finite p -group and let ρ be a faithful representation of H over k of minimal dimension. Then ρ decomposes as a direct sum of exactly $r = \text{rank}(Z(H))$ irreducible representations*

$$\rho = \rho_1 \oplus \cdots \oplus \rho_r.$$

and if χ_i are the central characters of ρ_i , then $\{\chi_i|_{\Omega_1(Z(H))}\}$ is a basis for $\widehat{\Omega}_1(Z(H))$

Lemma 3.5 ([13], Theorem 9.1). *For G a finite abelian group, $\widehat{G} \cong G$, and for G_1 and G_2 finite abelian groups,*

$$\widehat{G_1 \times G_2} \cong \widehat{G_1} \times \widehat{G_2}.$$

Definition 3.6. Fix a nontrivial character ψ of \mathbb{F}_p^+ . Define $\psi_b(x) = \psi(bx)$.

Remark 7. The map given by $b \mapsto \psi_b$ is an isomorphism between \mathbb{F}_p^+ and $\widehat{\mathbb{F}_p^+}$.

Definition 3.7. Fix a nontrivial character ψ of \mathbb{F}_p^+ . Fix m . For $\mathbf{b} = (b_j) \in (\mathbb{F}_p^+)^m$, define

$$\psi_{\mathbf{b}}(\mathbf{d}) = \prod_j (\psi_{b_j}(d_j)) \in (\widehat{\mathbb{F}_p^+})^m$$

Lemma 3.8. *Fix a nontrivial character ψ of \mathbb{F}_p^+ . The elements of $(\widehat{\mathbb{F}_p^+})^m$ are of the form $\mathbf{d} \mapsto \psi(\mathbf{b}\mathbf{d}^T)$ for fixed $\mathbf{b} \in \mathbb{F}_p^m$.*

Proof. By Remark 7 and Lemma 3.5, $(b_j) \mapsto (\psi_{b_j}) = \psi_{\mathbf{b}}$ is an isomorphism between $(\mathbb{F}_p^+)^m$ and $(\widehat{\mathbb{F}_p^+})^m$. And

$$\psi_{\mathbf{b}}(\mathbf{d}) = \prod_j \psi(b_j d_j) = \psi\left(\sum_j b_j d_j\right) = \psi(\mathbf{b}\mathbf{d}^T).$$

□

3.1 The Wigner-Mackey Little Group Method

I will first describe the Wigner-Mackey little group method over \mathbb{C} . The following exposition of Wigner-Mackey Theory is in [26] Section 8.2 and is also reproduced in [1] page 7: Let $G = \Delta \rtimes L$ with Δ abelian. Since Δ is abelian, its irreducible characters are of degree 1 and form a group $\widehat{\Delta} = \text{Hom}(\Delta, \mathbb{C}^\times)$. The group G acts on $\widehat{\Delta}$ by

$$(g\chi)(a) = \chi(g^{-1}ag), \text{ for } g \in G, \chi \in \widehat{\Delta}, a \in \Delta.$$

Let $(\chi_s)_{\chi_s \in \widehat{\Delta}/L}$ be a system of representatives for the orbits of L in $\widehat{\Delta}$. For each χ_s , let L_s be the subgroup of L consisting of those elements such that $l\chi_s = \chi_s$, that is $L_s = \text{Stab}_L(\chi_s)$. Let $G_s = \Delta \cdot L_s$ be the corresponding subgroup of G . Extend the function χ_s to G_s by setting

$$\chi_s(al) = \chi_s(a), \text{ for } a \in \Delta, l \in L_s.$$

Then since $l\chi_s = \chi_s$ for all $l \in L_s$, we see that χ_s is a one-dimensional representation of G_s . Now let λ be an irreducible representation of L_s ; by composing λ with the canonical projection $G_s \rightarrow L_s$ we obtain an irreducible representation λ of G_s , i.e

$$\lambda(al) = \lambda(l), \text{ for } a \in \Delta, l \in L_s.$$

Finally, by taking the tensor product of χ_s and λ , we obtain an irreducible representation $\chi_s \otimes \lambda$ of G_s . Let $\theta_{s,\lambda}$ be the corresponding induced representation of G , i.e. $\theta_{s,\lambda} := \text{Ind}_{G_s}^G(\chi_s \otimes \lambda)$. The following is Proposition 25 in Chapter 8 of [26], it is called ‘‘Wigner-Mackey theory’’ in [1] (Theorem 4.2):

Theorem 3.9 ([26], Proposition 25). *Under the above assumptions, we have*

- (i) $\theta_{s,\lambda}$ is irreducible.
- (ii) Every irreducible representation of G is isomorphic to one of the $\theta_{s,\lambda}$.

Remark 8. Serre also proves a uniqueness statement: If $\theta_{s,\lambda}$ and $\theta_{s',\lambda'}$ are isomorphic, then $\chi_s = \chi'_s$ and λ is isomorphic to λ' . But we do not care about the uniqueness of the irreducible representations. In what follows, we will consider characters χ_s with $\chi_s \in \widehat{\Delta}$ rather than $\chi_s \in \widehat{\Delta}/L$. The two points above still hold.

Geetha Venkataraman proves a generalization of the Wigner-Mackey little group method [28] with the following assumptions: Let k be a field whose characteristic does not divide $|G|$. Suppose that $G = \Delta \rtimes L$ with Δ abelian and that all irreducible representations of Δ over k have degree 1.

Remark 9. Note that in the cases considered in this paper, these conditions hold so long as $\text{char } k \neq p$. Since we are considering the p -Sylows, this takes care of the first condition. All of our p -Sylows have the form $\Delta \rtimes L$. In each case, $\Delta \cong (\mathbb{Z}/p\mathbb{Z})^N$ for some N . By the note following Lemma 2.8, we may assume that k contains a primitive p -th root of unity. Thus we can conclude that all irreducible representations of Δ over k have degree 1.

Theorem 3.10 ([28], Theorem 4.1). *Let G be a finite group which is a semi-direct product of an abelian group Δ by a subgroup L . Let k be a field such that the characteristic of k does not divide $|G|$, and assume that all irreducible representation of Δ over k are of degree 1. Let O_1, \dots, O_t be the distinct orbits under the action of G on $\widehat{\Delta}$ and let χ_s be a representative of the orbit O_s . Let G_s be the stabilizer of χ_s and let $L_s = I_s \cap L$ (i.e. $L_s = \text{Stab}_L(\chi_s)$). For any irreducible representation λ of L_s , let $\theta_{s,\lambda}$ denote the representation of G induced from the irreducible representation $\chi_s \otimes \lambda$ of G_s . Then*

- (i) $\theta_{s,\lambda}$ is irreducible.
- (ii) Every irreducible representation of G is isomorphic to one of the $\theta_{s,\lambda}$.

Remark 10. Ventakaraman also proves a uniqueness statement, but again we do not care about the uniqueness of the irreducible representations. In what follows, we will consider characters χ_s with $\chi_s \in \widehat{\Delta}$ rather than $\chi_s \in \widehat{\Delta}/L$. The two points above still hold.

4 PSL_n

In this section, we will prove that

Theorem 4.1. *For any p, k a field such that $\text{char } k \neq p$,*

$$ed_k(\text{PSL}_n(\mathbb{F}_{p^r}), p) = ed_k(\text{GL}_n(\mathbb{F}_{p^r})) = rp^{r(n-2)}.$$

Remark 11. For $p \neq 2$, this is a theorem of [1] (Theorems 1.1 and 1.2).

For PSL_n , we will actually identify a subgroup of the p -SyLOW (the generalized Heisenberg group) to which we will apply our general (Wigner-Mackey) proof method. This will give a lower bound for the essential p -dimension. We will find an upper bound by constructing a specific faithful representation (we will extend the minimal dimensional representation of the generalized Heisenberg to a representation of the same dimension).

4.1 Finding the p -Sylows and identifying the center

For any p , $|\text{Syl}_p(PSL_n(\mathbb{F}_{p^r}))| = (p^r)^{n(n-1)/2} = |\text{Syl}_p(GL_n(\mathbb{F}_{p^r}))|$. So $\text{Syl}_p(PSL_n(\mathbb{F}_{p^r}))$ maps injectively onto $\text{Syl}_p(GL_n(\mathbb{F}_{p^r}))$; thus their p -sylows are isomorphic, and hence they have the same essential p -dimension.

Definition 4.2. Define $\text{Up}_n(\mathbb{F}_{p^r})$ to be the unitriangular $n \times n$ matrices over \mathbb{F}_{p^r} (upper triangular matrices with 1's on the diagonal.)

Then since this is a subgroup of $PSL_n(\mathbb{F}_{p^r})$ of order $(p^r)^{\sum_{k=1}^{n-1} k} = (p^r)^{(\sum_{k=1}^{n-1} k)}$,

Lemma 4.3. $\text{Syl}_p(PSL_n(\mathbb{F}_{p^r})) = \text{Up}_n(\mathbb{F}_{p^r})$.

Also,

Lemma 4.4.

$$Z(\text{Up}_n(\mathbb{F}_{p^r})) = \left\{ \begin{pmatrix} 1 & 0 & \cdots & 0 & a_{1,n} \\ 0 & 1 & 0 & \cdots & 0 \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix} \right\} \cong (\mathbb{Z}/p\mathbb{Z})^r$$

Proof. This follows by direct computation: Given a matrix A in the center, pick B with $b_{i,j} = 0$ for $i < j$ except for b_{i_0, j_0} . Then since $AB = BA$, we can see by comparing the entries of AB and BA that we must have $a_{i,j} = 0$ except for $i = 1, j = n$. \square

4.2 Identifying the generalized Heisenberg subgroup

Definition 4.5. Define

$$H_n(\mathbb{F}_{p^r}) = \left\{ \begin{pmatrix} 1 & \mathbf{a} & x \\ \mathbf{0} & \text{Id}_{n-2} & \mathbf{b}^T \\ 0 & \mathbf{0} & 1 \end{pmatrix} : x \in \mathbb{F}_{p^r}, \mathbf{a}, \mathbf{b} \in (\mathbb{F}_{p^r})^{n-2} \right\}$$

This is a subgroup of $\text{Up}_n(\mathbb{F}_{p^r})$. A straightforward calculations shows that

$$\mathbf{Lemma 4.6.} \quad Z(H_n(\mathbb{F}_{p^r})) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & x \\ \mathbf{0} & \text{Id}_{n-2} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \right\} = Z(\text{Up}_n(\mathbb{F}_{p^r})).$$

Another straightforward calculation shows that

Lemma 4.7. $H_n(\mathbb{F}_{p^r})$ is isomorphic to the the group with underlying set $(\mathbb{F}_{p^r})^{2(n-2)} \times \mathbb{F}_{p^r}$ and multiplication given by

$$(\mathbf{a}, \mathbf{b}, x) \cdot (\mathbf{c}, \mathbf{d}, y) = (\mathbf{a} + \mathbf{c}, \mathbf{b} + \mathbf{d}, x + y + \mathbf{ad}^T),$$

where $\mathbf{a}, \mathbf{b} \in \mathbb{F}_{p^r}^{n-2}, x \in \mathbb{F}_{p^r}$.

4.3 Calculating $\text{ed}_k(H_n)$

In this section, we will show that

Theorem 4.8 ([1], Theorem 1.1).

$$\text{ed}_k(H_n(\mathbb{F}_{p^r})) = rp^{r(n-2)}$$

Bardestani et. al. proved this for all p using Wigner-Mackey. Their proof is for a more general definition of the generalized Heisenberg groups, and so it is more complicated than we need. Here, we provide a simplified version of their proof for our particular generalized Heisenberg group. This provides a warmup for our proofs for the other finite simple groups.

4.3.1 Maximizing L_s

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element in $\widehat{\Delta} \cong (\widehat{\mathbb{F}_{p^r}^{n-1}})^+$ is of the form $\psi_{\mathbf{a},b} : (\mathbf{v}, 0, x) \mapsto \psi(\mathbf{v}\mathbf{a}^T + bx)$ for $(\mathbf{a}, b) \in \mathbb{F}_{p^r}^{n-1}$.

Proposition 4.9. *The characters extending a non-trivial central character are $\psi_{\mathbf{a},b}$ with $b \neq 0$.*

Proof. $\psi_{\mathbf{a},b}$ restricted to the center is ψ_b . This is non-trivial if and only if $b \neq 0$. \square

Define

$$\Delta = \{(\mathbf{v}, 0, k) : \mathbf{v} \in \mathbb{F}_{p^r}^{n-2}, k \in \mathbb{F}_{p^r}\}, \quad L = \{(0, \mathbf{w}, 0) : \mathbf{w} \in \mathbb{F}_{p^r}^{n-2}\}.$$

Then $H_n(\mathbb{F}_{p^r}) = \Delta \rtimes L$ with Δ is an abelian subgroup of $H_n(\mathbb{F}_{p^r})$. The action of L on Δ given by

$${}^{(0, \mathbf{w}, 0)}(\mathbf{v}, 0, x) = (\mathbf{v}, 0, x + \mathbf{v}\mathbf{w}^T).$$

For $h = (0, \mathbf{w}, 0)$, the action on $\psi_{\mathbf{a},b}$ is given by

$${}^h(\psi_{\mathbf{a},b}(\mathbf{v}, \mathbf{0}, x)) = \psi_{\mathbf{a},b}(\mathbf{v}, \mathbf{0}, x + \mathbf{v}\mathbf{w}^T).$$

Proposition 4.10. *For $s = (\mathbf{a}, b)$ with $b \neq 0$*

$$L_s = \text{Stab}_L(\psi_s) = \{\mathbf{0}\}.$$

Corollary 4.11. *For $s = (\mathbf{a}, b) \neq (\mathbf{0}, 0)$, since L_s is trivial, λ is trivial, and*

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = |L| = (p^r)^{(n-2)}$$

Proof of Proposition 4.10. $(0, \mathbf{w}, 0) \in L_s$ if and only if for all $(\mathbf{v}, \mathbf{0}, x) \in \Delta$,

$$\begin{aligned} \psi_{\mathbf{a},b}(\mathbf{v}, \mathbf{0}, x + \mathbf{v}\mathbf{w}^T) &= \psi_{\mathbf{a},b}(\mathbf{v}, \mathbf{0}, x) \\ \Leftrightarrow \psi_{\mathbf{a},b}(\mathbf{0}, \mathbf{0}, \mathbf{v}\mathbf{w}^T) &= 1 \\ \Leftrightarrow \psi(b\mathbf{v}\mathbf{w}^T) &= 1 \end{aligned}$$

If $b\mathbf{w} \neq \mathbf{0}$, $\mathbf{v}\mathbf{w}^T$ will run through all values of \mathbb{F}_{p^r} , and since ψ is nontrivial, this is not possible. Thus we must have $b\mathbf{w}^T = \mathbf{0}$. And since $b \neq 0$, this means that $\mathbf{w}^T = \mathbf{0}$. So $L_s = \{(0, 0, 0)\}$. \square

4.3.2 Proof

Proof of Theorem 4.8. By Theorem 2.9, we need to find the minimal dimension of a faithful representation of $H_n(\mathbb{F}_{p^r})$. By Lemma 3.4, since $Z(H_n(\mathbb{F}_{p^r})) \cong (\mathbb{Z}/p\mathbb{Z})^r$ has rank r , a faithful representation ρ of $H_n(\mathbb{F}_{p^r})$ of minimal dimension decomposes as a direct sum

$$\rho = \rho_1 \oplus \cdots \oplus \rho_r$$

of exactly r irreducibles. By Wigner-Mackey each of the ρ_i is isomorphic to $\theta_{s,\lambda}$ for some χ_s, λ where $\chi_s = \psi_{\mathbf{a},b} \in \widehat{\Delta}$. And since the ρ_i have non-trivial central characters, we can conclude that $b \neq 0$, and hence ρ_i has dimension $p^{r(n-2)}$ by Corollary 4.11. Thus ρ has dimension $rp^{r(n-2)}$. Therefore,

$$\text{ed}_k(H_n(\mathbb{F}_{p^r})) = rp^{r(n-2)}.$$

\square

Remark 12. Since we classified the irreps with non-trivial central characters completely and they all have the same dimension, we get both the upper and lower bound from the classification. In the other groups, we will only find the irreps which give representations of minimal dimension, which will give the lower bound. The upper bound is found by construction.

4.4 Showing that $\text{ed}_k(\text{Up}_n(\mathbb{F}_{p^r})) = \text{ed}_k(H_n(\mathbb{F}_{p^r}))$

In this section, we will show that

Theorem 4.12. $\text{ed}_k(\text{Up}_n(\mathbb{F}_{p^r})) = \text{ed}_k(H_n(\mathbb{F}_{p^r}))$

Remark 13. For $p \neq 2$, this is a theorem of [1] (Theorem 1.2).

Since $H_n(\mathbb{F}_{p^r}) \subset \text{Up}_n(\mathbb{F}_{p^r})$, we already have that $\text{ed}_k(H_n(\mathbb{F}_{p^r})) \leq \text{ed}_k(\text{Up}_n(\mathbb{F}_{p^r}))$. So it suffices to prove

$$\text{ed}_k(\text{Up}_n(\mathbb{F}_{p^r})) \leq \text{ed}_k(H_n(\mathbb{F}_{p^r})).$$

We will do this by constructing a faithful representation of $\text{Up}_n(\mathbb{F}_{p^r})$ of dimension $rp^{r(n-2)}$.

Proposition 4.13. $\text{Up}_n(\mathbb{F}_{p^r})$ is isomorphic to $H_n(\mathbb{F}_{p^r}) \rtimes \text{Up}_{n-2}(\mathbb{F}_{p^r})$, where the action of $\text{Up}_{n-2}(\mathbb{F}_{p^r})$ on $H_n(\mathbb{F}_{p^r})$ is given by $A(\mathbf{a}, \mathbf{b}, x) = (\mathbf{a}A^{-1}, \mathbf{b}A^T, x)$.

Proof. Note that

$$\text{Up}_n(\mathbb{F}_{p^r}) = \left\{ \begin{pmatrix} 1 & \mathbf{a} & x \\ 0 & A & \mathbf{b}^T \\ 0 & \mathbf{0} & 1 \end{pmatrix} : \mathbf{a}, \mathbf{b} \in \mathbb{F}_{p^r}^2, x \in \mathbb{F}_{p^r}, A \in \text{Up}_{n-2}(\mathbb{F}_{p^r}) \right\},$$

and

$$\begin{pmatrix} 1 & \mathbf{a} & x \\ 0 & A & \mathbf{b}^T \\ 0 & \mathbf{0} & 1 \end{pmatrix} \begin{pmatrix} 1 & \mathbf{c} & y \\ 0 & B & \mathbf{d}^T \\ 0 & \mathbf{0} & 1 \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{a}B + \mathbf{c} & x + y + \mathbf{a}\mathbf{d}^T \\ 0 & AB & \mathbf{b}^T + A\mathbf{d}^T \\ 0 & \mathbf{0} & 1 \end{pmatrix}$$

So $\text{Up}_n(\mathbb{F}_{p^r})$ is isomorphic to the group with underlying set $H_n(\mathbb{F}_{p^r}) \times \text{Up}_{n-2}(\mathbb{F}_{p^r})$ and multiplication given by

$$(\mathbf{a}, \mathbf{b}, x, A)(\mathbf{c}, \mathbf{d}, y, B) = (\mathbf{a}B + \mathbf{c}, \mathbf{b} + \mathbf{d}A^T, x + y + \mathbf{a}\mathbf{d}^T, AB).$$

$H_n(\mathbb{F}_{p^r})$ is normal in the group described above. Therefore, we actually have that the group described above is a semi-direct product $H_n(\mathbb{F}_{p^r}) \rtimes \text{Up}_{n-2}(\mathbb{F}_{p^r})$. And since

$$(\mathbf{0}, \mathbf{0}, 0, A)(\mathbf{a}, \mathbf{b}, x, \text{Id}_{n-2})(\mathbf{0}, \mathbf{0}, 0, A)^{-1} = (\mathbf{a}A^{-1}, \mathbf{b}A^T, x, \text{Id}_{n-2}),$$

the action of $\text{Up}_{n-2}(\mathbb{F}_{p^r})$ on $H_n(\mathbb{F}_{p^r})$ is given by

$$A(\mathbf{a}, \mathbf{b}, x) = (\mathbf{a}A^{-1}, \mathbf{b}A^T, x)$$

□

4.4.1 Proof

We will construct a faithful representation of $\text{Up}_n(\mathbb{F}_{p^r})$ of dimension $rp^{r(n-2)}$ by extending the irreducible representations used above. Define

$$\tilde{\Delta} = \Delta \rtimes \text{Up}_{n-2}(\mathbb{F}_{p^r}).$$

For $s = (\mathbf{a}, b) \in (\mathbb{F}_{p^r}^+)^{n-2}$, define

$$\tilde{\psi}_s(\mathbf{v}, k, A) = \psi_s(\mathbf{v}, k).$$

Define

$$\rho_s = \text{Ind}_{\tilde{\Delta}}^{H_n(\mathbb{F}_{p^r}) \rtimes \text{Up}_{n-2}(\mathbb{F}_{p^r})} \tilde{\psi}_s.$$

Then

$$\dim(\rho_s) = [H_n(\mathbb{F}_{p^r}) \rtimes \text{Up}_{n-2}(\mathbb{F}_{p^r}) : \tilde{\Delta}] \dim(\tilde{\psi}_s)$$

$$= p^{r(n-2)}$$

Let $\rho = \oplus_i \rho_{e_i}$. Then ρ is a representation of $H_n(\mathbb{F}_{p^r}) \rtimes \text{Up}_{n-2}(\mathbb{F}_{p^r})$ of dimension $rp^{r(n-2)}$. And since

$$\rho_{e_i}|_{Z(\text{Up}_n(\mathbb{F}_{p^r}))} = \rho|_{Z(H_n(\mathbb{F}_{p^r}))} = \psi_{e_i}$$

form a basis for $\widehat{\mathbb{F}_{p^r}^+}$, we can conclude by Lemma 3.3 that $\oplus_i \rho_{e_i}$ is a faithful representation of $\text{Up}_n(\mathbb{F}_{p^r})$.

Remark 14. A potential alternate construction of the upper bound brought to my attention by Nate Harman: Let $q = p^r$ and let A denote the copy of \mathbb{F}_q^{n-1} in $U\mathcal{P}_n(\mathbb{F}_q)$ where everything but the diagonal is zero except the last column which can be anything. $\text{Up}_n(\mathbb{F}_q)$ acts naturally on the set of vectors $(a_1, a_2, \dots, a_{n-1}, 1)^T$ in \mathbb{F}_q^n , and moreover this action is simply transitive for A . If we linearize this action over \mathbb{C} , we get a q^{n-1} dimensional $\text{Up}_n(\mathbb{F}_q)$ representation, which restricts to the regular representation of A . Now we can split this up according to the action of the center of $Z(\text{Up}_n(\mathbb{F}_q))$ to decompose this as q summands each of dimension q^{n-2} . Note that when we restrict to $Z(\text{Up}_n(\mathbb{F}_q))$, we have the regular representation of $Z(\text{Up}_n(\mathbb{F}_q))$. And every irreducible representation is contained in the regular representation ([26], Section 2.4, Corollary 1). Since a character of $Z(\text{Up}_n(\mathbb{F}_q))$ is an irreducible representation, this means every character must be contained in the regular representation of $Z(\text{Up}_n(\mathbb{F}_q))$. Thus for each character ψ_b of $Z(\text{Up}_n(\mathbb{F}_q))$, there must exist a summand that restricts to ψ_b on $Z(\text{Up}_n(\mathbb{F}_q))$. Choose a basis for the characters of \mathbb{F}_q as a \mathbb{F}_p vector space. Then take ρ to be the direct sum of the corresponding summands. Then the restriction of ρ to the center is faithful by Lemma 3.3. Therefore, ρ is faithful as well, and its dimension is rq^{n-2} , as desired.

5 The Symplectic Groups

In this section, we will show that

Theorem 5.1. *For k a field such that $\text{char } k \neq p$,*

$$\text{ed}_k(\text{PSp}(2n, p^r)', p) = \text{ed}_k(\text{Sp}(2n, p^r)) = \begin{cases} rp^{r(n-1)}, & p \neq 2 \text{ or } p = 2, n = 2 \\ r2^{r(n-1)-1}(2^{r(n-2)} + 1), & p = 2, n > 2 \end{cases}$$

Remark 15. We do not prove the case $p = 2, n = 2, r = 1$, since it is already known that $\text{ed}_k(\text{PSp}(4, 2)', 2) = \text{ed}_k(A_6, 2) = 2$. In any other case, $\text{PSp}(2n, p^r)' = \text{PSp}(n, p^r)$, so it suffices to prove the theorem for $\text{PSp}(2n, p^r)$.

5.1 Definitions

Definition 5.2. Let $S = \begin{pmatrix} 0 & \text{Id}_n \\ -\text{Id}_n & 0 \end{pmatrix}$. $\text{Sp}(2n, p^r)$ is defined as $\{M \in \text{GL}_{2n}(\mathbb{F}_{p^r}) : M^T S M = S\}$.

Remark 16. A matrix $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is symplectic if and only if $A^T C, B^T D$ are symmetric and $A^T D - C^T B = \text{Id}_n$.

Definition 5.3. $\text{PSp}(2n, p^r)$ is defined as $\text{Sp}(2n, p^r)/Z(\text{Sp}(2n, p^r))$.

5.2 The p -Sylows

For any p , $|\text{Syl}_p(\text{PSp}(2n, p^r))| = (p^r)^{n^2} = |\text{Syl}_p(\text{Sp}(2n, p^r))|$. So $\text{Syl}_p(\text{Sp}(2n, p^r))$ maps injectively onto $\text{Syl}_p(\text{PSp}(2n, p^r))$; thus their p -sylows are isomorphic, and hence they have the same essential p -dimension.

Recall that $\text{Up}_n(\mathbb{F}_{p^r})$ is the unitriangular $n \times n$ matrices over \mathbb{F}_{p^r} (upper triangular matrices with 1's on the diagonal) and $\text{Syl}_p(\text{PSL}(n, p^r)) = \text{Up}_n(\mathbb{F}_{p^r})$.

Definition 5.4. For any p , define $\text{Sym}(n, p^r)$ as the group of $n \times n$ symmetric matrices under addition (with entries from \mathbb{F}_{p^r}).

Lemma 5.5 (See [19], [20], [22],[11]). *For any p ,*

$$\text{Syl}_p(\text{PSp}(2n, p^r)) = \left\{ \begin{pmatrix} A & 0_n \\ 0_n & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \text{Id}_n & B \\ 0_n & \text{Id}_n \end{pmatrix} : A \in \text{Up}_n(\mathbb{F}_{p^r}), B \in \text{Sym}(n, p^r) \right\}.$$

Proof. Let

$$N = \left\{ \begin{pmatrix} \text{Id}_n & B \\ 0_n & \text{Id}_n \end{pmatrix} : B \text{ is symmetric} \right\} \subset \text{PSp}(2n, p^r).$$

Then $N \cong \text{Sym}(n, p^r)$ and $|N| = (p^r)^{\frac{n(n+1)}{2}}$. Let

$$O = \left\{ \begin{pmatrix} A & 0_n \\ 0_n & (A^{-1})^T \end{pmatrix} : A \in \text{Syl}_p(\text{PSL}(n, p^r)) = \text{Up}_n(\mathbb{F}_{p^r}) \right\} \subset \text{PSp}(2n, p^r).$$

Then $O \cong \text{Up}_n(\mathbb{F}_{p^r})$ and $|O| = (p^r)^{n(n-1)/2}$. A direct computation shows that $N \cdot O = N \rtimes O$ and

$$|N \rtimes O| = (p^r)^{n^2} = |\text{Syl}_p(\text{PSp}(2n, p^r))|.$$

Thus

$$N \rtimes O = \left\{ \begin{pmatrix} A & 0_n \\ 0_n & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \text{Id}_n & B \\ 0_n & \text{Id}_n \end{pmatrix} : A \in \text{Up}_n(\mathbb{F}_{p^r}), B \in \text{Sym}(n, p^r) \right\}.$$

is a p -Sylow of $\text{PSp}(2n, p^r)$. □

Corollary 5.6. [See [20]] *For any p , $\text{Syl}_p(\text{PSp}(2n, p^r)) \cong \text{Sym}(n, p^r) \rtimes \text{Up}_n(\mathbb{F}_{p^r})$, where the action is given by $A(B) = ABA^T$.*

5.3 Calculating the center

5.3.1 $p \neq 2$

Lemma 5.7. *For $p \neq 2$,*

$$Z(\text{Syl}_p(\text{PSp}(2n, p^r))) = \left\{ \begin{pmatrix} \text{Id}_n & D \\ 0_n & \text{Id}_n \end{pmatrix} : D = \begin{pmatrix} d & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix} \right\} \cong \mathbb{F}_{p^r}^+ \cong (\mathbb{Z}/p\mathbb{Z})^r$$

For the proof of Lemma 5.7, we need the following lemma:

Lemma 5.8. *For $p \neq 2$, $D \in \text{Sym}(n, p^r)$, $AD = D(A^{-1})^T$ for all $A \in \text{Up}_n(\mathbb{F}_{p^r})$ if and only if $D = \begin{pmatrix} d & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix}$.*

Granting this lemma, we can prove Lemma 5.7:

Proof of Lemma 5.7.

$$\text{Syl}_p(\text{PSp}(2n, p^r)) = \left\{ \begin{pmatrix} A & AB \\ 0_n & (A^{-1})^T \end{pmatrix} : A \in \text{Up}_n(\mathbb{F}_{p^r}), B \in \text{Sym}(n, p^r) \right\}.$$

Note that

$$\begin{pmatrix} A & AB \\ 0_n & (A^{-1})^T \end{pmatrix}^{-1} \begin{pmatrix} C & CD \\ 0_n & (C^{-1})^T \end{pmatrix} \begin{pmatrix} A & AB \\ 0_n & (A^{-1})^T \end{pmatrix} = \begin{pmatrix} A^{-1}CA & A^{-1}CAB + A^{-1}CD(A^{-1})^T - B((A^{-1}CA)^{-1})^T \\ 0_n & ((A^{-1}CA)^{-1})^T \end{pmatrix}$$

So $\begin{pmatrix} C & CD \\ 0_n & (C^{-1})^T \end{pmatrix} \in Z(\text{Syl}_p(\text{PSp}(n, p^r)))$ if and only if $C \in Z(\text{Syl}_p(\text{PSL}(n, p^r)))$ and

$$CD = CB + CA^{-1}D(A^{-1})^T - B(C^{-1})^T, \quad \text{for all } A \in \text{Up}_n(\mathbb{F}_{p^r}), B \in \text{Sym}(n, p^r).$$

Choosing $A, B = \text{Id}_n$, we need $CD = C + CD - (C^{-1})^T$. So $C = (C^{-1})^T$ and thus $C = \text{Id}_n$. So the other requirement above becomes

$$D = A^{-1}D(A^{-1})^T \Leftrightarrow AD = D(A^{-1})^T, \quad \text{for all } A \in \text{Up}_n(\mathbb{F}_{p^r}).$$

By Lemma 5.8, we get that

$$Z(\text{Syl}_p(\text{PSp}(2n, p^r))) = \left\{ \begin{pmatrix} \text{Id}_n & D \\ 0_n & \text{Id}_n \end{pmatrix} : D = \begin{pmatrix} d & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix} \right\}$$

□

Proof of Lemma 5.8.

⇐: This is a straightforward calculation.

⇒: We will prove this by induction.

Base Case: When $n = 2$, we can write $A = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ and $D = \begin{pmatrix} x & y \\ y & z \end{pmatrix}$.

$$AD = \begin{pmatrix} x + ay & y + az \\ y & z \end{pmatrix},$$

and

$$D(A^{-1})^T = \begin{pmatrix} x - ay & y - az \\ y & z \end{pmatrix}.$$

So the condition that $AD = D(A^{-1})^T$ for all A implies that $y = 0$ and $z = 0$.

Induction Step: Write

$$D = \begin{pmatrix} d_{1,1} & d_{1,2} & d_{1,3} & \cdots & d_{1,n} \\ d_{1,2} & d_{2,2} & d_{2,3} & \cdots & d_{2,n} \\ \vdots & & \ddots & & \vdots \\ d_{1,n-1} & d_{2,n-1} & \cdots & d_{n-1,n-1} & d_{n-1,n} \\ d_{1,n} & d_{2,n} & \cdots & d_{n-1,n} & d_{n,n} \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & a_{n-1,n} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Then

$$AD = \begin{pmatrix} d_{1,1} & \cdots & d_{1,n-1} & d_{1,n} \\ d_{2,2} & \cdots & d_{2,n-1} & d_{2,n} \\ \vdots & \ddots & \vdots & \vdots \\ d_{1,n-1} + a_{n-1,n}d_{1,n} & \cdots & d_{n-1,n-1} + a_{n-1,n}d_{n-1,n} & d_{n-1,n} + a_{n-1,n}d_{n,n} \\ d_{1,n} & \cdots & d_{n-1,n} & d_{n,n} \end{pmatrix}$$

and

$$D(A^{-1})^T = \begin{pmatrix} d_{1,1} & \cdots & d_{1,n-2} & d_{1,n-1} - a_{n-1,n}d_{1,n} & d_{1,n} \\ d_{1,2} & \cdots & d_{2,n-2} & d_{2,n-1} - a_{n-1,n}d_{2,n} & d_{2,n} \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ d_{1,n-1} & \cdots & d_{n-1,n-2} & d_{n-1,n-1} - a_{n-1,n}d_{n-1,n} & d_{n-1,n} \\ d_{1,n} & \cdots & d_{n,n-2} & d_{n-1,n} - a_{n-1,n}d_{n,n} & d_{n,n} \end{pmatrix}$$

In order for these to be equal for all $a_{n-1,n}$, we must have $d_{k,n} = 0$ for all k . So the matrix

$$D' = \begin{pmatrix} d_{1,1} & d_{1,2} & d_{1,3} & \cdots & d_{1,n-1} \\ d_{1,2} & d_{2,2} & d_{2,3} & \cdots & d_{2,n-1} \\ \vdots & & \ddots & & \vdots \\ d_{1,n-2} & d_{2,n-2} & \cdots & d_{n-2,n-2} & d_{n-2,n} \\ d_{1,n-1} & d_{2,n-1} & \cdots & d_{n-2,n-1} & d_{n-1,n-1} \end{pmatrix}$$

satisfies the condition $A'D' = D'(A'^{-1})^T$ for all $A' \in \text{Up}_{n-1}(\mathbb{F}_{p^r})$. By induction, we conclude that

$$D' = \begin{pmatrix} d & \mathbf{0} \\ \mathbf{0} & 0_{n-2} \end{pmatrix},$$

and hence

$$D = \begin{pmatrix} d & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix}.$$

□

5.3.2 $p = 2$

Lemma 5.9.

$$Z(\text{Syl}_p(\text{PSp}(2n, 2^r))) = \left\{ \begin{pmatrix} \text{Id}_n & D \\ 0_n & \text{Id}_n \end{pmatrix} : D_{i,j} = 0, \text{ for all } (i,j) \notin \{(1,1), (1,2), (2,1)\} \right\} \cong (\mathbb{F}_{p^r}^+)^2 \cong (\mathbb{Z}/p\mathbb{Z})^{2r}$$

For the proof of Lemma 5.9, we need the following lemma:

Lemma 5.10. *For $p = 2$, $D \in \text{Sym}(n, 2^r)$, $AD = D(A^{-1})^T$ for all $A \in \text{Up}_n(\mathbb{F}_{2^r})$ if and only if $D_{i,j} = 0$, for all $(i,j) \notin \{(1,1), (1,2), (2,1)\}$.*

Granting this lemma, we can prove Lemma 5.9:

Proof of Lemma 5.9.

$$\text{Syl}_2(\text{PSp}(2n, 2^r)) = \left\{ \begin{pmatrix} A & AB \\ 0_n & (A^{-1})^T \end{pmatrix} : A \in \text{Up}_n(\mathbb{F}_{2^r}), B \in \text{Sym}(n, 2^r) \right\}.$$

Just as for $p \neq 2$, $\begin{pmatrix} C & CD \\ 0_n & (C^{-1})^T \end{pmatrix} \in Z(\text{Syl}_p(\text{PSp}(n, 2^r)))$ if and only if $C = \text{Id}_n$ and

$$D = A^{-1}D(A^{-1})^T \Leftrightarrow AD = D(A^{-1})^T, \quad \text{for all } A \in \text{Up}_n(\mathbb{F}_{p^r}).$$

By Lemma 5.10, then we have that

$$Z(\text{Syl}_2(\text{PSp}(2n, 2^r))) = \left\{ \begin{pmatrix} \text{Id}_n & D \\ 0_n & \text{Id}_n \end{pmatrix} : D_{i,j} = 0, \text{ for all } (i,j) \notin \{(1,1), (1,2), (2,1)\} \right\}$$

□

Proof of Lemma 5.10.

⇐: This is a straightforward calculation.

⇒: We will prove this by induction.

Base Case: When $n = 2$, we can write $A = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ and $D = \begin{pmatrix} x & y \\ y & z \end{pmatrix}$.

$$AD = \begin{pmatrix} x + ay & y + az \\ y & z \end{pmatrix},$$

and

$$D(A^{-1})^T = \begin{pmatrix} x + ay & y \\ y + az & z \end{pmatrix}.$$

So the condition that $AD = D(A^{-1})^T$ for all A implies that $z = 0$.

Remark 17. This calculation is the key difference between odd and even characteristic.

Induction Step: Assume that $n > 2$. Write

$$D = \begin{pmatrix} d_{1,1} & d_{1,2} & d_{1,3} & \cdots & d_{1,n} \\ d_{1,2} & d_{2,2} & d_{2,3} & \cdots & d_{2,n} \\ \vdots & & \ddots & & \vdots \\ d_{1,n-1} & d_{2,n-1} & \cdots & d_{n-1,n-1} & d_{n-1,n} \\ d_{1,n} & d_{2,n} & \cdots & d_{n-1,n} & d_{n,n} \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & a_{n-1,n} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Then

$$AD = \begin{pmatrix} d_{1,1} & \cdots & d_{1,n-1} & d_{1,n} \\ d_{1,2} & \cdots & d_{2,n-1} & d_{2,n} \\ \vdots & \ddots & \vdots & \vdots \\ d_{1,n-1} + a_{n-1,n}d_{1,n} & \cdots & d_{n-1,n-1} + a_{n-1,n}d_{n-1,n} & d_{n-1,n} + a_{n-1,n}d_{n,n} \\ d_{1,n} & \cdots & d_{n-1,n} & d_{n,n} \end{pmatrix}$$

and

$$D(A^{-1})^T = \begin{pmatrix} d_{1,1} & \cdots & d_{1,n-2} & d_{1,n-1} + a_{n-1,n}d_{1,n} & d_{1,n} \\ d_{1,2} & \cdots & d_{2,n-2} & d_{2,n-1} + a_{n-1,n}d_{2,n} & d_{2,n} \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ d_{1,n-1} & \cdots & d_{n-1,n-2} & d_{n-1,n-1} + a_{n-1,n}d_{n-1,n} & d_{n-1,n} \\ d_{1,n} & \cdots & d_{n,n-2} & d_{n-1,n} + a_{n-1,n}d_{n,n} & d_{n,n} \end{pmatrix}$$

In order for these to be equal for all $a_{n-1,n}$, we must have $d_{k,n} = 0$ for all k except $k = n - 1$. Since $n > 2$, we can pick

$$A = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ & & \ddots & & \vdots \\ 0 & \cdots & 1 & a_{n-2,n-1} & 0 \\ 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

By comparing the entries of AD and $D(A^{-1})^T$, we see that in order to have $AD = D(A^{-1})^T$ for all $a_{n-2,n-1}$, we must have $d_{k,n-1} = 0$ for all k except $k = n - 2$. In particular, we get that $d_{n,n-1} = d_{n-1,n} = 0$. Thus $d_{k,n} = 0$ for all k . So the matrix

$$D' = \begin{pmatrix} d_{1,1} & d_{1,2} & d_{1,3} & \cdots & d_{1,n-1} \\ d_{1,2} & d_{2,2} & d_{2,3} & \cdots & d_{2,n-1} \\ \vdots & & \ddots & & \vdots \\ d_{1,n-2} & d_{2,n-2} & \cdots & d_{n-2,n-2} & d_{n-2,n} \\ d_{1,n-1} & d_{2,n-1} & \cdots & d_{n-2,n-1} & d_{n-1,n-1} \end{pmatrix}$$

satisfies the condition $A'D' = D'(A'^{-1})^T$ for all $A' \in \text{Up}_{n-1}(\mathbb{F}_{p^r})$. By induction, we conclude that

$$D_{i,j} = 0, \text{ for all } (i,j) \notin \{(1,1), (1,2), (2,1)\}.$$

□

5.4 Maximizing L_s

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element of $\widehat{\text{Sym}}(n, p^r) \cong (\mathbb{F}_{p^r}^+)^{n(n+1)/2}$ is of the form $\mathbf{d} \mapsto \psi(\mathbf{b}\mathbf{d}^T)$, where $\mathbf{d}, \mathbf{b} \in (\mathbb{F}_{p^r})^{n(n+1)/2}$.

Proposition 5.11. *For $p \neq 2$, the characters extending a non-trivial central character are $\psi_{(b_j)}$ with $b_1 \neq 0$.*

Proof. $\psi_{(b_j)}$ restricted to the center is ψ_{b_1} . This is non-trivial if and only if $b_1 \neq 0$. \square

Proposition 5.12. *For $p = 2$, the characters extending a non-trivial central character are $\psi_{(b_j)}$ with $(b_1, b_2) \neq (0, 0)$, that is $b_1 \neq 0$ or $b_2 \neq 0$.*

Proof. $\psi_{(b_j)}$ restricted to the center is $\psi_{(b_1, b_2)}$. This is non-trivial if and only if $(b_1, b_2) \neq (0, 0)$. \square

We will use Wigner-Mackey Theory with $G = \text{Syl}_p(\text{PSp}(2n, p^r)) = \text{Sym}(n, p^r) \rtimes \text{Up}_n(\mathbb{F}_{p^r})$ to compute the minimum dimension of an irreducible representation of G with non-trivial central character. So we have $\Delta = \text{Sym}(n, p^r) \times \{\text{Id}_n\}$, $L = \{0_n\} \rtimes \text{Up}_n(\mathbb{F}_{p^r})$. Recall that $L_s = \text{Stab}_L(\psi_s)$. For any p , let

$$B = \begin{pmatrix} b_1 & b_2 & \cdots & & b_n \\ b_2 & b_{n+1} & \cdots & & b_{2n-1} \\ \vdots & & \ddots & & \vdots \\ b_{n-1} & \cdots & & b_{n(n+1)/2-2} & b_{n(n+1)/2-1} \\ b_n & \cdots & & b_{n(n+1)/2-1} & b_{n(n+1)/2} \end{pmatrix}$$

be the matrix corresponding to $s = (b_1, \dots, b_{n(n+1)/2})$.

Remark 18. In all of the following, we view $\psi_{(b_j)}$ as a map on $\Delta \cong \text{Sym}(n, p^r) \cong \mathbb{F}_{p^r}^{n(n+1)/2}$. So $\psi_{(b_j)}(D, \text{Id}) = \psi_{(b_j)}(D) = \psi(\mathbf{b} \cdot \mathbf{d})$, where $\mathbf{b} = (b_j)$ and \mathbf{d} is the vector corresponding to the matrix D .

Note that $(0_n, H^{-1}) \in L_s$ if and only if for all $\mathbf{d} \in (\mathbb{F}_{p^r})^{n(n+1)/2} = D \in \text{Sym}(n, p^r)$,

$$\psi_{(b_j)}((0_n, H)(D, \text{Id}_n)(0_n, H^{-1})) = \psi_{(b_j)}(D, \text{Id}_n).$$

Let \mathbf{hdh}^T denote the vector corresponding to HDH^T . Then since

$$\psi_{(b_j)}((0_n, H)(D, \text{Id}_n)(0_n, H^{-1})) = \psi(\mathbf{b} \cdot \mathbf{hdh}^T),$$

and

$$\psi_{(b_j)}(D, \text{Id}_n) = \psi(\mathbf{b} \cdot \mathbf{d}),$$

we get that $(0_n, H^{-1}) \in L_s$ if and only if for all $\mathbf{d} \in (\mathbb{F}_{p^r})^{n(n+1)/2} = D \in \text{Sym}(n, p^r)$,

$$\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1.$$

Write

$$H = \begin{pmatrix} 1 & h_{1,2} & h_{1,3} & \cdots & h_{1,n} \\ 0 & 1 & h_{2,3} & \cdots & h_{2,n} \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & h_{n-1,n} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}, \quad D = \begin{pmatrix} d_{1,1} & d_{1,2} & d_{1,3} & \cdots & d_{1,n} \\ d_{1,2} & d_{2,2} & d_{2,3} & \cdots & d_{2,n} \\ \vdots & & \ddots & & \vdots \\ d_{1,n-1} & d_{2,n-1} & \cdots & d_{n-1,n-1} & d_{n-1,n} \\ d_{1,n} & d_{2,n} & \cdots & d_{n-1,n} & d_{n,n} \end{pmatrix}.$$

Then

$$\begin{aligned}
& HDH^T - D \\
&= \begin{pmatrix}
\left[\sum_{l=1}^n (h_{1,l} \sum_{k=1}^n d_{l,k} h_{1,k}) \right] - d_{1,1} & \left[\sum_{l=2}^n (h_{2,l} \sum_{k=1}^n d_{l,k} h_{1,k}) \right] - d_{1,2} & \cdots & \left(\sum_{k=1}^n d_{k,n} h_{1,k} \right) - d_{1,n} \\
\left[\sum_{l=1}^n (h_{1,l} \sum_{k=2}^n d_{l,k} h_{2,k}) \right] - d_{1,2} & \left[\sum_{l=2}^n (h_{2,l} \sum_{k=2}^n d_{l,k} h_{2,k}) \right] - d_{2,2} & \cdots & \left(\sum_{k=2}^n d_{k,n} h_{2,k} \right) - d_{2,n} \\
\vdots & \vdots & \ddots & \vdots \\
\left(\sum_{l=1}^n h_{1,l} d_{l,n} \right) - d_{1,n} & \left(\sum_{l=2}^n h_{2,l} d_{l,n} \right) - d_{2,n} & \cdots & 0
\end{pmatrix}
\end{aligned}$$

5.4.1 $p \neq 2$

Proposition 5.13. For $p \neq 2$, $s = (b_i)$, $b_1 \neq 0$, $\max_s |L_s| = |Up_{n-1}(\mathbb{F}_{p^r})| = p^{r(n-1)(n-2)/2}$. And this maximum is achieved when $s = (b, 0, \dots, 0)$.

Corollary 5.14. For $p \neq 2$, $s = (b, 0, \dots, 0)$, λ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|Up_n(\mathbb{F}_{p^r})|}{|Up_{n-1}(\mathbb{F}_{p^r})|} = |\mathbb{F}_{p^r}^{n-1}| = p^{r(n-1)}.$$

We will prove Proposition 5.13 in two steps:

Claim 5.15. For $p \neq 2$, for $s = (b_i)$, $b_1 \neq 0$, $|L_s| \leq |Up_{n-1}(\mathbb{F}_{p^r})| = p^{r(n-1)(n-2)/2}$.

Claim 5.16. For $p \neq 2$, $s = (b, 0, \dots, 0)$ with $b \neq 0$,

$$L_s = \text{Stab}_L(\psi_s) = \{H : H_{1,j} = 0, \forall j \neq 1\} \cong Up_{n-1}(\mathbb{F}_{p^r})$$

Proof of Claim 5.15. Pick $j_0 \neq 1$ and choose D with $d_{i,j} = 0$ except for $d_{1,j_0} = d_{j_0,1}$. Then we get that

$$HDH^T - D = \begin{pmatrix}
2d_{1,j_0} h_{1,j_0} & h_{2,j_0} d_{1,j_0} & \cdots & h_{j_0-1,j_0} d_{1,j_0} & 0 & \cdots & 0 \\
h_{2,j_0} d_{1,j_0} & 0 & & & \cdots & & 0 \\
\vdots & & & \ddots & & & \vdots \\
h_{j_0-1,j_0} d_{1,j_0} & 0 & & & \cdots & & 0 \\
0 & & & & \cdots & & 0 \\
\vdots & & & \ddots & & & \vdots \\
0 & 0 & & & \cdots & & 0
\end{pmatrix}$$

Thus we have

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = 2d_{1,j_0} h_{1,j_0} B_{1,1} + \sum_{i=2}^{j_0-1} h_{i,j_0} d_{1,j_0} B_{1,i} = d_{1,j_0} \left(2h_{1,j_0} B_{1,1} + \sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} \right)$$

If $\left(2h_{1,j_0} B_{1,1} + \sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} \right) \neq 0$, then as we run through all the values for d_{1,j_0} , we will get that $\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})$ runs through all the values of \mathbb{F}_{p^r} . And since ψ is non-trivial, this means that $\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}))$ cannot always equal 1. This is a contradiction. So we must have

$$2h_{1,j_0} B_{1,1} + \sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} = 0$$

for all choices of $j_0 \neq 1$. Recall that $B_{1,1} = b_1 \neq 0$. So, for all j_0 , given h_{i,j_0} for $i > 1$, the above dictates h_{1,j_0} :

$$h_{1,j_0} = \frac{-1}{2B_{1,1}} \sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i}.$$

Thus we can conclude that for all $s = (b_i)$ with $b_1 \neq 0$,

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed } \forall j \neq 1\}| = |\text{Up}_{n-1}(\mathbb{F}_{p^r})| = p^{r(n-1)(n-2)/2}$$

□

Proof of Claim 5.16. Let B be the matrix corresponding to $s = (b, 0, \dots, 0)$. Since the only nonzero entry of B is $B_{1,1} = b$, we have that

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = b(\mathbf{hdh}^T - \mathbf{d})_1 = b \left(\left[\sum_{l=1}^n (h_{1,l} \sum_{k=1}^n d_{l,k} h_{1,k}) \right] - d_{1,1} \right).$$

By the proof of Claim 5.15, if $H \in L_s$, then $\forall j_0 \neq 1$, we must have

$$h_{1,j_0} = \frac{-1}{2B_{1,1}} \sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} = 0.$$

And if $h_{1,j_0} = 0 \forall j \neq 1$, then we have

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = b(h_{1,1}d_{1,1}h_{1,1} - d_{1,1}) = 0, \text{ since } h_{1,1} = 1$$

Thus we have shown that $(0_n, H^{-1}) \in L_s$ if and only if $h_{1,j} = 0, \forall j \neq 1$. Therefore,

$$L_s = \{(0_n, H^{-1}) : H_{1,j} = 0, \forall j \neq 1\} \cong \text{Up}_{n-1}(\mathbb{F}_{p^r}).$$

□

5.4.2 $p = 2$

Case 1: $n = 2$

Proposition 5.17. For $p = 2, n = 2, s = (b_i), (b_1, b_2) \neq (0, 0)$:

For $b_1, b_2 \neq 0$, $\max_s |L_s| = 2$ and this maximum is achieved when $s = (b_1, b_2, 0)$ with $b_1, b_2 \neq 0$.
If $b_1 = 0$ or $b_2 = 0$, then $|L_s| = 1$.

Corollary 5.18. For $p = 2, n = 2, s = (b_1, b_2, 0), b_1, b_2 \neq 0, \lambda$ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|\text{Up}_2(\mathbb{F}_{2^r})|}{2} = \frac{2^r}{2} = 2^{r-1}.$$

For $s = (b_1, b_2, b_3)$, with $b_1 = 0$ or $b_2 = 0$,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|\text{Up}_2(\mathbb{F}_{2^r})|}{1} = 2^r$$

Proof. We will prove Proposition 5.17 in two steps:

Step 1: Proving that for $p = 2, n = 2, s = (b_i), (b_1, b_2) \neq (0, 0)$: if $b_1, b_2 \neq 0$, then $|L_s| \leq 2$, and otherwise $|L_s| = 1$.

$$HDH^T - D = \begin{pmatrix} \left[\sum_{l=1}^2 (h_{1,l} \sum_{k=1}^2 d_{l,k} h_{1,k}) \right] - d_{1,1} & \left(\sum_{k=1}^2 d_{k,2} h_{1,k} \right) - d_{1,2} \\ \left(\sum_{l=1}^2 h_{1,l} d_{l,2} \right) - d_{1,2} & 0 \end{pmatrix}$$

Let $p = 2, s = (b_i)$ with $(b_1, b_2) \neq (0, 0)$.

Calculation 1. Choose $d_{i,j} = 0$ except for $d_{2,2}$.

Then we get that

$$HDH^T - D = \begin{pmatrix} h_{1,2}^2 d_{2,2} & h_{1,2} d_{2,2} \\ h_{1,2} d_{2,2} & 0 \end{pmatrix}$$

Thus we have

$$\begin{aligned} \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) &= B_{1,1} h_{1,2}^2 d_{2,2} + B_{1,2} h_{1,2} d_{2,2} \\ &= d_{2,2} h_{1,2} (B_{1,1} h_{1,2} + B_{1,2}) \end{aligned}$$

Then since ψ is non-trivial, we must have $h_{1,2}(B_{1,1}h_{1,2} + B_{1,2}) = 0$. Thus either $h_{1,2} = 0$ or $B_{1,1}h_{1,2} + B_{1,2} = 0$. If $B_{1,1} \neq 0, B_{1,2} \neq 0$, then either $h_{1,2} = 0$ or $h_{1,2} = \frac{B_{1,2}}{B_{1,1}}$. If $B_{1,1} \neq 0, B_{1,2} = 0$ or $B_{1,1} = 0, B_{1,2} \neq 0$, then $h_{1,2} = 0$. Our findings can be summarized in a chart as follows (we only care when $(B_{1,1}, B_{1,2}) \neq (0, 0)$):

Case:	result	options
$B_{1,1} \neq 0, B_{1,2} \neq 0$	$h_{1,2} = 0$ or $h_{1,2} = \frac{B_{1,2}}{B_{1,1}}$	2
$B_{1,1} \neq 0, B_{1,2} = 0$	$h_{1,2} = 0$	1
$B_{1,1} = 0, B_{1,2} \neq 0$	$h_{1,2} = 0$	1

Thus we can conclude that for all $s = (b_i)$ with $(b_1, b_2) \neq (0, 0)$, then for $b_1, b_2 \neq 0, |L_s| \leq 2$ and otherwise $|L_s| = 1$.

Step 2: Showing that when $s = (b_1, b_2, 0)$ with $b_1, b_2 \neq 0, |L_s| = 2$.

For $s = (b_1, b_2, b_3)$,

$$\begin{aligned} \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) &= b_1 \left(\left[\sum_{l=1}^2 (h_{1,l} \sum_{k=1}^2 d_{l,k} h_{1,k}) \right] - d_{1,1} \right) + b_2 \left(\left[\sum_{k=1}^2 d_{k,2} h_{1,k} \right] - d_{1,2} \right) \\ &= b_1 h_{1,2}^2 d_{2,2} + b_2 d_{2,2} h_{1,2} \quad \text{since we are working in char 2} \\ &= d_{2,2} h_{1,2} (b_1 h_1 + b_2) \end{aligned}$$

If $b_1 \neq 0, b_2 \neq 0$, then either $h_{1,2} = 0$ or $h_1 = \frac{b_2}{b_1}$. In either case, the above is identically zero. Thus $|L_s| = 2$. □

Case 2: $n > 2$

Proposition 5.19. For $p = 2, n > 2, s = (b_i), (b_1, b_2) \neq (0, 0)$:

For $b_2 \neq 0, \max_s |L_s| = 2 |Up_{n-2}(\mathbb{F}_{2^r})| = 2^{r(n-2)(n-3)/2+1}$ and this maximum is achieved when $s = (b_i) = (b_1, b_2, 0, \dots, 0)$ with $b_1, b_2 \neq 0$.

For $b_1 \neq 0, \max_s |L_s| = 2 |Up_{n-1}(\mathbb{F}_{2^r})| = 2^{r(n-1)(n-2)/2+1}$, and this maximum is achieved when $s = (b_i) = (b_1, 0, b_3, \dots, 0)$ with $b_1, b_3 \neq 0$.

Corollary 5.20. For $p = 2, n > 2$,

For $s = (b_1, b_2, 0, \dots, 0)$ with $b_1, b_2 \neq 0, \lambda$ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|Up_n(\mathbb{F}_{2^r})|}{2 |Up_{n-2}(\mathbb{F}_{2^r})|} = \frac{|(\mathbb{F}_{2^r})^{n-1+n-2}|}{2} = 2^{r(2n-3)-1}$$

For $s = (b_1, 0, b_3, \dots, 0)$ with $b_1, b_3 \neq 0, \lambda$ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|Up_n(\mathbb{F}_{2^r})|}{2 |Up_{n-1}(\mathbb{F}_{2^r})|} = \frac{|(\mathbb{F}_{2^r})^{n-1}|}{2} = 2^{r(n-1)-1}$$

Remark 19. For any $n > 2$ and any $r, 2^{r(2n-3)-1} > 2^{r(n-1)}$.

Proof. Again, we will prove this in two steps:

Step 1: Proving that for $\mathbf{p} = \mathbf{2}, \mathbf{n} > \mathbf{2}, \mathbf{s} = (\mathbf{b}_i), (\mathbf{b}_1, \mathbf{b}_2) \neq (\mathbf{0}, \mathbf{0})$: If $\mathbf{b}_2 \neq \mathbf{0}$, then $|\mathbf{L}_s| \leq 2^{r(n-2)(n-3)/2+1}$, and if $\mathbf{b}_2 = \mathbf{0} (\Rightarrow \mathbf{b}_1 \neq \mathbf{0})$, then $|\mathbf{L}_s| \leq 2^{r(n-1)(n-2)/2+1}$.

Calculation 1. For $j_0 > 2$, choose $d_{i,j} = 0$ except for $d_{1,j_0} = d_{j_0,1}$.

Then

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = \sum_{i=2}^{j_0-1} h_{i,j_0} d_{1,j_0} B_{1,i} = d_{1,j_0} \sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i}$$

So for all $j_0 > 2$, we must have

$$\sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} = 0.$$

For $j_0 = 3$, this gives $h_{2,j_0} B_{1,2} = 0$, and thus if $B_{1,2} \neq 0$, we must have $h_{2,j_0} = 0$. For $2 \leq k \leq n$, if $B_{1,k} \neq 0$, then for all $j_0 > 3$, given h_{i,j_0} for $i \neq 1, k$, the above dictates h_{k,j_0} :

$$h_{k,j_0} = \frac{-1}{B_{1,k}} \sum_{i=2, i \neq k}^{j_0-1} h_{i,j_0} B_{1,i}.$$

Calculation 2. Now for $j_0 > 1$, choose $d_{i,j} = 0$ except for d_{j_0,j_0} .

Then

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = d_{j_0,j_0} \left(\sum_{l=1}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0} \right)$$

So for all $j_0 \neq 1$, we must have

$$\sum_{l=1}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0} = 0.$$

Thus we have that for all $j_0 \neq 1$,

$$h_{1,j_0} \left(\sum_{k=1}^{j_0} B_{1,k} h_{k,j_0} \right) + \sum_{l=2}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0} = 0$$

For $j_0 = 2$, this tells us $0 = h_{1,2}(B_{1,1}h_{1,2} + B_{1,2})$. If $B_{1,2} = 0 (\Rightarrow B_{1,1} \neq 0)$ or $B_{1,1} = 0 (\Rightarrow B_{1,2} \neq 0)$, then this implies that $h_{1,2} = 0$. If $B_{1,2} \neq 0$ and $B_{1,1} \neq 0$, then we have two options for $h_{1,2}$: $h_{1,2} = 0$ and $h_{1,2} = \frac{B_{1,2}}{B_{1,1}}$. For $j_0 > 2$, this is a quadratic expression for h_{1,j_0} in terms of $B_{i,j}$ and h_{k,j_0} for $k > 1$, namely

$$B_{1,1} h_{1,j_0}^2 + \left(\sum_{k=2}^{j_0} B_{1,k} h_{k,j_0} \right) h_{1,j_0} + \sum_{l=2}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0} = 0$$

Thus for $j_0 > 2$, given h_{i,j_0} for $i > 1$, there are up to two options for h_{1,j_0} .

Calculation 3. Now for $j_0 > 2$, choose $d_{i,j} = 0$ except for $d_{2,j_0} = d_{j_0,2}$.

Then

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = d_{2,j_0} \left(B_{1,2} h_{1,j_0} + \sum_{i=2}^{j_0} B_{1,i} h_{i,j_0} h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i} h_{i,j_0} \right)$$

So for all $j_0 > 2$, we must have

$$B_{1,2} h_{1,j_0} + \sum_{i=2}^{j_0} B_{1,i} h_{i,j_0} h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i} h_{i,j_0} = 0.$$

If $B_{1,2} \neq 0$, then for all $j_0 > 2$, given h_{i,j_0} for $i > 2$, the above dictates h_{1,j_0} :

$$h_{1,j_0} = \frac{-1}{B_{1,2}} \left(\sum_{i=2}^{j_0} B_{1,i} h_{i,j_0} h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,k} h_{i,j_0} \right)$$

Case 1. $b_2 \neq 0$

If $B_{1,2} = b_2 \neq 0$, then we have from the first calculation that for all $j_0 > 1$, given h_{i,j_0} for $i > 2$, h_{2,j_0} are dictated. By the second calculation we have that there are at most two options for $h_{1,2}$. And by the third calculation, h_{1,j_0} is dictated for $j_0 > 2$. Thus for $b_2 \neq 0$, we can conclude that

$$\begin{aligned} |L_s| &\leq |\{H : \text{two options for } H_{1,2}, \text{ and } \forall j > 2, H_{1,j}, H_{2,j} \text{ fixed}, \}| \\ &= 2|\text{Up}_{n-2}(\mathbb{F}_{p^r})| \\ &= 2^{r(n-2)(n-3)/2+1}. \end{aligned}$$

Case 2. $b_2 = 0, b_3 \neq 0$

If $B_{1,2} = b_2 = 0 (\Rightarrow B_{1,1} \neq 0)$: We have by the second calculation that for $j_0 > 2$,

$$0 = B_{1,1} h_{1,j_0}^2 + \left(\sum_{k=3}^{j_0} B_{1,k} h_{k,j_0} \right) h_{1,j_0} + \sum_{l=2}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0}$$

For $j_0 = 2$, we get $B_{1,1} h_{1,2}^2 = 0$. Thus we must have $h_{1,2} = 0$. For $j_0 = 3$, we get $0 = B_{1,1} h_{1,3}^2 + B_{1,3} h_{1,3} = h_{1,3}(B_{1,1} h_{1,3} + B_{1,3})$. Thus either $h_{1,3} = 0$ or $h_{1,3} = \frac{B_{1,3}}{B_{1,1}}$. For $j_0 > 3$, we have from the first calculation that $\sum_{i=3}^{j_0-1} h_{i,j_0} B_{1,i} = 0$, so the equality from the second calculation becomes

$$0 = B_{1,1} h_{1,j_0}^2 + B_{1,j_0} h_{1,j_0} + \sum_{l=2}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0}$$

We will use the following proposition:

Proposition 5.21 ([18], Proposition 1). *In a finite field of order 2^r , for $f(x) = ax^2 + bx + c$, we have the following:*

- (i) f has exactly one root $\Leftrightarrow b = 0$.
- (ii) f has exactly two roots $\Leftrightarrow b \neq 0$ and $\text{Tr}(\frac{ac}{b^2}) = 0$.
- (iii) f has no root $\Leftrightarrow b \neq 0$ and $\text{Tr}(\frac{ac}{b^2}) = 1$,

where $\text{Tr}(x) = x + x^2 + \dots + x^{2^r-1}$.

So, for $j_0 > 3$, if $B_{1,j_0} = 0$, then there is only one option for h_{1,j_0} . Otherwise, it might have two options or no options. Thus we have the following for $j_0 > 3$: If $B_{1,j_0} = 0$, then there is one option for h_{1,j_0} , but h_{k,j_0} can be anything for $k > 1$. And if $B_{1,j_0} \neq 0$, then there is only one option for h_{j_0,k_0} for all $k_0 > 2$ (by the first calculation with $k = j_0, j_0 = k_0$), but h_{1,j_0} might have two options. So we can obtain an upper bound for L_s by choosing $B_{1,j} = 0$ for all $j > 3$ and assuming all the options are in L_s . In this case $h_{2,j}$ can be anything, but $h_{1,j}$ is fixed for all j except $j = 3$, and there are two options for $h_{1,3}$. So we get that

$$\begin{aligned} |L_s| &\leq |\{H : H_{1,j} \text{ fixed } \forall j \neq 3, H_{1,3} = 0 \text{ or } \frac{B_{1,3}}{B_{1,1}}\}| \\ &= 2|\text{Up}_{n-1}(\mathbb{F}_{2^r})| \end{aligned}$$

$$= 2^{r(n-1)(n-2)/2+1}$$

Step 2: Showing that for $p = 2, n > 2$: When $\mathbf{s} = (\mathbf{b}_1, \mathbf{b}_2, \mathbf{0}, \dots, \mathbf{0})$ with $\mathbf{b}_1, \mathbf{b}_2 \neq \mathbf{0}$, $|\mathbf{L}_s| = 2^{r(n-2)(n-3)/2+1}$, and when $\mathbf{s} = (\mathbf{b}_1, \mathbf{0}, \mathbf{b}_3, \dots, \mathbf{0})$ with $\mathbf{b}_1, \mathbf{b}_3 \neq \mathbf{0}$, $|\mathbf{L}_s| = 2^{r(n-1)(n-2)/2+1}$.

Let $p = 2, s = (b_1, b_2, \dots, b_n, 0, \dots, 0)$. And let B be the corresponding matrix. Then

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = b_1 \sum_{k=2}^n h_{1,k}^2 d_{k,k} + \sum_{j=2}^n b_j \left(\left[\sum_{l=j}^n (h_{j,l} \sum_{k=1}^n d_{l,k} h_{1,k}) \right] - d_{1,j} \right)$$

Case 1. $b_1, b_2 \neq 0, b_3, \dots, b_n = 0$.

Since $B_{1,2} = b_2 \neq 0$, then we have from the first calculation in Step 1 that for all $j_0 > 2$,

$$h_{2,j_0} = \frac{-1}{B_{1,2}} \sum_{i=3}^{j_0-1} h_{i,j_0} B_{1,i} = 0.$$

By the second calculation we have that there are two options for $h_{1,2}$: $h_{1,2} = 0$ and $h_{1,2} = \frac{B_{1,2}}{B_{1,1}}$. And by the third calculation, for $j_0 > 2$,

$$h_{1,j_0} = \frac{-1}{B_{1,2}} \left(\sum_{i=2}^{j_0} B_{1,i} h_{i,j_0} h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i} h_{i,j_0} \right) = \frac{-1}{B_{1,2}} B_{1,2} h_{2,j_0} h_{1,2} = 0$$

Thus we have

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = d_{2,2} h_{1,2} (B_{1,1} h_{1,2} + B_{1,2})$$

So whether $h_{1,2} = 0$ or $h_{1,2} = \frac{B_{1,2}}{B_{1,1}}$, this is identically 0. Therefore

$$|L_s| = |\{H : H_{1,2} = 0 \text{ or } H_{1,2} = \frac{B_{1,2}}{B_{1,1}}, H_{1,j} = 0 = H_{2,j} \ \forall j > 0\}| = 2|\text{UP}_{n-2}(\mathbb{F}_{2^r})| = 2^{r(n-2)(n-3)/2+1}$$

Case 2. $b_1 \neq 0, b_2 = \dots = b_n = 0$.

If $B_{1,k} = b_k = 0$ for $2 \leq k \leq n$: We have the following by the work in Step 1:

$h_{1,2} = 0$. By the second calculation we have that there are two options for $h_{1,3}$: $h_{1,2} = 0$ and $h_{1,3} = \frac{B_{1,3}}{B_{1,1}}$. And for $j_0 > 3$,

$$0 = B_{1,1} h_{1,j_0}^2 + B_{1,j_0} h_{1,j_0} + \sum_{l=2}^{j_0-1} \sum_{k=l}^{j_0} B_{l,k} h_{l,j_0} h_{k,j_0} = B_{1,1} h_{1,j_0}^2$$

So we have $h_{1,j_0} = 0$ for $j_0 \neq 1$. Thus

$$\begin{aligned} \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) &= b_1 \sum_{k=2}^n h_{1,k}^2 d_{k,k} && \text{since } b_i = 0 \text{ for } i > 1 \\ &= 0 && \text{since } h_{1,j_0} = 0 \text{ for } j_0 \neq 1 \end{aligned}$$

Therefore

$$|L_s| = |\{H : H_{1,3} = 0 \text{ or } H_{1,3} = \frac{B_{1,3}}{B_{1,1}}, H_{1,j_0} = 0 \text{ for } j_0 \neq 1, 3\}| = 2|\text{UP}_{n-1}(\mathbb{F}_{2^r})| = 2^{r(n-1)(n-2)/2+1}$$

□

5.5 Proof

Proof of Theorem 5.1. Case 1: $p \neq 2$

Step 1: Find the lower bound

For $p \neq 2$, the minimum dimension of an irreducible representation of $\text{Syl}_p(\text{PSp}(2n, p^r))$ with non-trivial central character is given by $\theta_{s, \lambda}$ with λ trivial and $s = (b, 0, \dots, 0)$. So the minimum dimension is $p^{r(n-1)}$. Thus the minimum dimension of a faithful representation is $rp^{r(n-1)}$.

Step 2: Find the upper bound

Let $\{e_i\}$ be a basis for $\mathbb{F}_{p^r}^+$ over \mathbb{F}_p . Then by Lemma 7, $\{\psi_{e_i}\}$ form a basis for the characters of $\mathbb{F}_{p^r}^+$. Thus $\{\psi_{e_i}|_{\Omega_1(\mathbb{F}_{p^r})}\}$ form a basis for $\widehat{\Omega}_1(\mathbb{F}_{p^r})$. Let $s_i = (e_i, 0, \dots, 0)$. Then by Corollary 5.14, for λ trivial, $\dim(\theta_{s_i, \lambda}) = p^{r(n-1)}$. Let $\rho = \bigoplus_i \theta_{s_i, \lambda}$. Then

$$\dim(\rho) = \sum_{i=1}^r \dim(\theta_{s_i, \lambda}) = \sum_{i=1}^r p^{r(n-1)} = rp^{r(n-1)}.$$

And by Lemma 3.3, since $\psi_{e_i}|_{\Omega_1(\mathbb{F}_{p^r})}$ form a basis for $\widehat{\Omega}_1(\mathbb{F}_{p^r})$, we have that $\bigoplus_i \theta_{s_i, \lambda}$ is a faithful representation of $\text{Syl}_p(\text{PSp}(2n, p^r))$. Thus ρ is the desired faithful representation of $\text{Syl}_p(\text{PSp}(2n, p^r))$ of dimension $rp^{r(n-1)}$.

Case 2: $p = 2$

Step 1: Find the lower bound

For $p = 2$, the minimum dimension of an irreducible representation of $\text{Syl}_2(\text{PSp}(2n, 2^r))$ with non-trivial central character is given by $\theta_{s, \lambda}$ with λ trivial and $s = (b_1, b_2, \dots, 0)$ with $(b_1, b_2) \neq (0, 0)$.

Subcase 1: $n = 2$: The minimum dimension is 2^{r-1} . So we have a lower bound of $\text{ed}_k(\text{PSp}(4, 2^r)) \geq 2r2^{r-1} = r2^r$.

Subcase 2: $n > 2$: Since $\{\chi_i|_{Z(\text{Syl}_2(\text{PSp}(2n, 2^r)))}\}$ a basis for $Z(\widehat{\text{PSp}}(2n, 2^r))$, we must have $b_2 \neq 0$ for at least r of the ρ_i . So the minimum dimension is $2^{r(2n-3)-1}$. The other r may have $b_2 = 0$, so the minimum dimension is $2^{r(n-1)-1}$. Thus we have $\text{ed}_k(\text{Syl}_2(\text{PSp}(2n, 2^r))) \geq r2^{r(2n-3)-1} + r2^{r(n-1)-1} = r2^{r(n-1)-1}(2^{r(n-2)} + 1)$.

Step 2: Find the upper bound

Let e_i be a basis for $\mathbb{F}_{2^r}^+$ over \mathbb{F}_2 . Let x be a nonzero element of \mathbb{F}_{2^r} . For $n = 2$, let

$$S = \{(e_i, e_i, 0), (x, e_i, 0)\}.$$

And for $n > 2$, let

$$S = \{(e_i, e_i, 0, \dots, 0), (e_i, 0, x, 0, \dots, 0)\}.$$

For both cases, for $s \in S$,

$$\{\psi_s|_{Z(\text{Syl}_2(\text{PSp}(2n, 2^r)))}\}$$

forms a basis for

$$Z(\widehat{\text{PSp}}(2n, 2^r))$$

[and $\Omega_1(Z(\text{Syl}_2(\text{PSp}(2n, 2^r)))) = Z(\text{Syl}_2(\text{PSp}(2n, 2^r)))$]. Let $\rho = \bigoplus_{s \in S} \theta_{s, \lambda}$. Then by Lemma 3.3,

$$\rho = \bigoplus_{s \in S} \theta_{s, \lambda}$$

is a faithful representation of $\text{Syl}_2(\text{PSp}(2n, 2^r))$. By Corollaries 5.18 and 5.20, for $s = (b_i)$, λ trivial,

$$\dim(\theta_{s, \lambda}) = \begin{cases} 2^{r-1}, & n = 2, b_1, b_2 \neq 0, b_3 = 0 \\ 2^r, & n = 2, b_1 = 0 \text{ or } b_2 = 0 \\ 2^{r(2n-3)-1}, & n > 2, b_1, b_2 \neq 0, b_i = 0 \forall i > 2 \\ 2^{r(n-1)-1}, & n > 2, b_1 \neq 0, b_i = 0 \forall i > 1 \end{cases}.$$

So

$$\dim(\rho) = \sum_{s \in S} \dim(\theta_{s,\lambda}) = \begin{cases} 2r2^{r-1} = r2^r, & n = 2, r > 1 \\ r2^{r(2n-3)-1} + r2^{r(n-1)-1} = r2^{r(n-1)-1}(2^{r(n-2)} + 1), & n > 2 \end{cases}$$

Thus ρ is the faithful representation of $\text{Syl}_2(\text{PSp}(2n, 2^r))$ of desired dimension. □

6 The Orthogonal Groups

In this section, we will show the following theorem:

Theorem 6.1. *For $\epsilon = +, -$ in the notation in the following section, k a field such that $\text{char } k \neq p$,*

$$\begin{aligned} \text{ed}_k(P\Omega^\epsilon(n, p^r), p) &= \text{ed}_k(\Omega^\epsilon(n, p^r), p) \\ &= \text{ed}_k(O^\epsilon(n, p^r), p) \\ &= \begin{cases} 2r, & n = 4, \text{ any } p \\ rp^{2r(m-2)}, & n = 2m, n > 4, \text{ any } p \\ rp^{r(m-1)} + rp^{r(m-1)(m-2)}, & n = 2m + 1, p \neq 2 \end{cases} \end{aligned}$$

Remark 20. We do not need to consider the case $n = 2m + 1, p = 2$ since $O^\epsilon(2m + 1, 2^r) \cong \text{Sp}(2m, 2^r)$ ([10], Theorem 14.2), so this case is taken care of in the work on the symplectic groups.

6.1 Definitions

6.1.1 $n = 2m, p \neq 2$

Let

$$A^+ = \begin{pmatrix} 0_m & \text{Id}_m \\ \text{Id}_m & 0_m \end{pmatrix}.$$

Let $\eta \in \mathbb{F}_{p^r}^\times$ be a non-square and let Id_m^η be the $m \times m$ identity matrix with the first entry replaced by η . And let

$$A^- = \begin{pmatrix} 0_m & \text{Id}_m \\ \text{Id}_m^\eta & 0_m \end{pmatrix}.$$

Definition 6.2. Define $O^+(2m, p^r)$ as $\{M \in \text{GL}(2m, \mathbb{F}_{p^r}) : M^T A^+ M = A^+\}$.

Definition 6.3. Define $O^-(2m, p^r)$ as $\{M \in \text{GL}(2m, \mathbb{F}_{p^r}) : M^T A^- M = A^-\}$.

Definition 6.4. For $\epsilon = +, -$, define $\text{SO}^\epsilon(2m, p^r) = \{M \in O^\epsilon(2m, p^r) : \det(M) = 1\}$.

Definition 6.5. For $\epsilon = +, -$, define $\Omega^\epsilon(2m, p^r) = (\text{SO}^\epsilon)'(2m, p^r)$ (the commutator subgroup).

Definition 6.6. For $\epsilon = +, -$, define $P\Omega^\epsilon(2m, p^r) = \Omega^\epsilon(2m, p^r) / (\Omega^\epsilon(2m, p^r) \cap \{\pm \text{Id}\})$.

Remark 21. $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in O^+(2m, p^r)$ iff $A^T C, B^T D$ are anti-symmetric and $A^T D + C^T B = \text{Id}_m$.

And $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in O^-(2m, p^r)$ iff $A^T C = -C^T A^\eta, B^T D = -D^T B^\eta, A^T D + C^T B^\eta = \text{Id}_m$, and $B^T C + D^T A^\eta = \text{Id}_m^\eta$, where A^η, B^η are A and B with the first row multiplied by η .

6.1.2 $n = 2m, p = 2$

Let $Q^+(x_i) = \sum_{i=1}^m x_i x_{i+m}$, and let

$$A_m^+ = \begin{pmatrix} 0_m & \text{Id}_m \\ 0_m & 0_m \end{pmatrix}.$$

Then $Q^+(x) = xA_m^+x^T$. Let η be such that $z^2 + z + \eta$ is irreducible in $\mathbb{F}_{2^r}[z]$. And let

$$Q_m^-(x_i) = \sum_{i=1}^m x_i x_{i+m} + x_m^2 + x_m x_{2m} + \eta x_{2m}^2$$

and define A_n^- to be

$$A_m^- = \begin{pmatrix} 0_m^1 & \text{Id}_m \\ 0_m & 0_m^\eta \end{pmatrix},$$

where

$$0_m^1 = \begin{pmatrix} 0_{m-1} & \mathbf{0} \\ \mathbf{0} & 1 \end{pmatrix} \text{ and } 0_m^\eta = \begin{pmatrix} 0_{m-1} & \mathbf{0} \\ \mathbf{0} & \eta \end{pmatrix}.$$

Then $Q_m^-(x) = xA_m^-x^T$. So if we write $x = (\mathbf{a}, b, \mathbf{c}, e)$, then

$$Q_m^-(x) = Q_{m-1}^+(\mathbf{a}, \mathbf{c}) + b^2 + be + \eta e^2 = \mathbf{a}\mathbf{d}^T + b^2 + be + \eta e^2.$$

Or if we write $x = (y, z)$, then

$$Q_m^-(x) = \mathbf{y}\mathbf{z}^T + y_m^2 + \eta z_m^2.$$

Definition 6.7. Define $O^\epsilon(2m, 2^r)$ as

$$\begin{aligned} O^\epsilon(2m, 2^r) &= \{M \in GL(2m, \mathbb{F}_{2^r}) : Q^+(Mx) = Q^+(x) \text{ for all } x \in \mathbb{F}_{2^r}^{2m}\} \\ &= \{M \in GL(2m, \mathbb{F}_{2^r}) : x(M^T A_n^\epsilon M)x^T = xA_m^\epsilon x^T, \forall x \in \mathbb{F}_{2^r}^{2m}\}. \end{aligned}$$

Definition 6.8. Define $B^\epsilon(x, y) = Q^\epsilon(x + y) + Q^\epsilon(x) + Q^\epsilon(y)$, the bilinear form corresponding to Q^ϵ .

Note that $B^+(x, y) = \sum_{i=1}^m x_i y_{i+m} + \sum_{i=1}^m y_i x_{i+m}$. So the corresponding matrix is

$$S = \begin{pmatrix} 0 & \text{Id}_m \\ \text{Id}_m & 0 \end{pmatrix}.$$

That is, $B^+(x, y) = xSy^T$. And $B^-(x, y) = \sum_{i=1}^{m-1} x_i y_{i+m} + y_i x_{i+m} + x_m y_{2m} + y_m x_{2m}$. So the corresponding matrix is also

$$S = \begin{pmatrix} 0 & \text{Id}_m \\ \text{Id}_m & 0 \end{pmatrix}.$$

That is, we have $B^-(x, y) = xSy^T = B^+(x, y)$, the same bilinear form as for A^+ . Note that this is a nondegenerate alternating form and we have

$$O^\epsilon(2m, 2^r) \subset Sp(2m, 2^r),$$

where $Sp(2m, 2^r)$ is the symplectic group corresponding to S .

Definition 6.9. Define $\Omega^\epsilon(2m, 2^r) = (O^\epsilon)^\epsilon(2m, 2^r)$ (the commutator subgroup).

For the sake of combination with $\Omega^\epsilon(2m, p^r)$ for $p \neq 2$, we make the following definition:

Definition 6.10. Define $P\Omega^\epsilon(2m, 2^r) = \Omega^\epsilon(2m, 2^r) / (\Omega^\epsilon(2m, 2^r) \cap \{\pm \text{Id}\}) = \Omega^\epsilon(2m, 2^r)$.

Definition 6.11. The *Dickson invariant*, $\delta_{2m,2^r}^\epsilon$, is a homomorphism from $O^\epsilon(2m, 2^r)$ to $\mathbb{Z}/2\mathbb{Z}$ taking the value of its fixed space mod 2. Algebraically, this is $\delta_{2m,2^r}^\epsilon(M) = \text{rank}(\text{Id}_{2m} - M) \pmod{2}$.

Definition 6.12. Define $SO^\epsilon(2m, 2^r) = \ker \delta_{2m,2^r}^\epsilon$.

Definition 6.13. Given $\epsilon = +, -$, the Witt index is defined to be the dimension of a maximal totally isotropic subspace of \mathbb{F}_{2^r} (this depends on ϵ). For $\epsilon = +, -$, let w_ϵ denote the Witt index.

Remark 22. Grove shows ([10], Proposition 14.41) that for Witt index $w_\epsilon > 0$, and $n \geq 2$,

$$\Omega^\epsilon(2m, 2^r) = (O^\epsilon)'(2m, 2^r) = (SO^\epsilon)'(2m, 2^r),$$

where $(O^\epsilon)'(2m, 2^r)$ is the commutator subgroup. He also shows ([10], Theorem 14.43) that if $m \geq 2$ and $(m, w_\epsilon) \neq (2, 2)$, then $\Omega^\epsilon(2m, 2^r)$ is simple.

Remark 23. If we write $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ and $x = (y, z)$ where $y, z \in \mathbb{F}_{2^r}^n$, then we get

$$Q(Mx) = yA^T C y^T + yA^T D z^T + zB^T C y^T + zB^T D z^T$$

And $M \in O^+(2m, 2^r)$ iff this is equal to $Q^+(x) = yz^T = zy^T$ for all $x = (y, z)$.

6.1.3 $n = 2m + 1$

Let

$$L = \begin{pmatrix} -1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & 0_m & \text{Id}_m \\ \mathbf{0} & \text{Id}_m & 0_m \end{pmatrix}.$$

Definition 6.14. Define $O(2m + 1, p^r) = \{M \in GL(2m + 1, \mathbb{F}_{p^r}) : M^T L M = L\}$.

Definition 6.15. Define $SO(2m + 1, p^r) = \{M \in O(2m + 1, p^r) : \det(M) = 1\}$.

Definition 6.16. Define $\Omega(2m + 1, p^r) = SO'(2m + 1, p^r)$ (the commutator subgroup).

Remark 24.

$$M = \begin{pmatrix} a & \mathbf{w} & \mathbf{x} \\ \mathbf{y} & A & B \\ \mathbf{z} & C & D \end{pmatrix}$$

is in $O(2m + 1, p^r)$ if and only if

$$a^2 - 2\mathbf{y} \cdot \mathbf{z} = 1,$$

$$0 = -a\mathbf{w} + \mathbf{y}C + \mathbf{z}A = -a\mathbf{x} + \mathbf{y}D + \mathbf{z}B = -a\mathbf{w} + A^T \mathbf{z} + C^T \mathbf{y} = -a\mathbf{x} + B^T \mathbf{z} + D^T \mathbf{y}$$

$A^T C, B^T D$ are anti-symmetric

and

$$A^T D + C^T B = \text{Id}_m.$$

6.2 The p -Sylows

Recall that for any p , we defined Up_m to be the unitriangular matrices over \mathbb{F}_{p^r} and we defined $\text{Sym}(m, p^r)$ to be the group of symmetric matrices under addition (with entries from \mathbb{F}_{p^r}).

Definition 6.17. For any p , define $\text{Antisym}(m, p^r)$ as the group of $m \times m$ anti-symmetric matrices under addition (with entries from \mathbb{F}_{p^r}).

Definition 6.18. For $p = 2$, define $\text{Antisym}0(m, 2^r) \subset \text{Antisym}(m, 2^r) = \text{Sym}(m, 2^r)$ as the subgroup of symmetric/antisymmetric matrices with 0's on the diagonal. That is,

$$\text{Antisym}0(m, 2^r) = \{B \in \text{Sym}(m, 2^r) = \text{Antisym}(m, 2^r) : B_{i,i} = 0, \forall i\}.$$

Definition 6.19. Recall that η is a fixed non-square in $\mathbb{F}_{p^r}^\times$. Let B^η denote B with the first row multiplied by η .

6.2.1 $n = 2m$

For $p \neq 2$,

$$|\mathrm{Syl}_p(P\Omega^\epsilon(2m, p^r))| = |\mathrm{Syl}_p(\Omega^\epsilon(2m, p^r))| = |\mathrm{Syl}_p(O^\epsilon(2m, p^r))| = p^{rm(m-1)}.$$

So $\mathrm{Syl}_p(O^\epsilon(2m, p^r))$ maps injectively onto $\mathrm{Syl}_p(\Omega^\epsilon(2m, p^r))$; thus their p -sylows are isomorphic, and hence they have the same essential p -dimension. So it suffices to consider $\Omega^\epsilon(2m, p^r)$. (We do this for notational purposes so we can combine the arguments with the case $p = 2$.)

A direct computation shows that

Lemma 6.20. [See [19], [14]] For $p \neq 2$, $\epsilon = +$,

$$\mathrm{Syl}_p(\Omega^+(2m, p^r)) = \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \mathrm{Id}_m & B \\ 0_m & \mathrm{Id}_m \end{pmatrix} : A \in \mathrm{Up}_m(\mathbb{F}_{p^r}), B \in \mathrm{Antisym}(m, p^r) \right\}.$$

and for $p \neq 2$, $\epsilon = -$,

$$\mathrm{Syl}_p(\Omega^-(2m, p^r)) = \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \mathrm{Id}_m & 0_m \\ C & \mathrm{Id}_m \end{pmatrix} : A \in \mathrm{Up}_m(\mathbb{F}_{p^r}), B \in \mathrm{Antisym}(m, p^r) \right\}.$$

Corollary 6.21. For $p \neq 2$, $\epsilon = +, -$,

$$\mathrm{Syl}_p(\Omega^\pm(2m, p^r)) \cong \mathrm{Antisym}(m, p^r) \rtimes \mathrm{Up}_m(\mathbb{F}_{p^r}),$$

where the action is given by

$$A(B) = ABA^T.$$

Since $\mathrm{Syl}_p(O^+(2m, p^r)) \cong \mathrm{Syl}_p(O^-(2m, p^r))$, we need only prove Theorem 6.1 for $\epsilon = +$.

Lemma 6.22. For $\epsilon = +, -$,

$$\mathrm{Syl}_2(\Omega^\epsilon(2m, 2^r)) = \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \mathrm{Id}_m & B \\ 0_m & \mathrm{Id}_m \end{pmatrix} : A \in \mathrm{Up}_m(\mathbb{F}_{2^r}), B \in \mathrm{Antisym}0(m, 2^r) \right\}.$$

Proof. Since $\Omega^\epsilon(2m, 2^r) \subset O^\epsilon(2m, 2^r) \subset \mathrm{Sp}(2m, 2^r)$, we must have

$\mathrm{Syl}_2(\Omega^\epsilon(2m, 2^r)) \subset \mathrm{Syl}_2(O^\epsilon(2m, 2^r)) \subset \mathrm{Syl}_2(\mathrm{Sp}(2m, 2^r))$. And we showed in our document on $P\mathrm{Sp}(2m, 2^r)$ that $\mathrm{Syl}_2(\mathrm{Sp}(2m, 2^r)) = N \rtimes O$ where

$$N = \left\{ \begin{pmatrix} \mathrm{Id}_m & B \\ 0_m & \mathrm{Id}_m \end{pmatrix} : B \in \mathrm{Sym}(m, p^r) \right\} \cong \mathrm{Sym}(m, p^r)$$

and

$$O = \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} : A \in \mathrm{Up}_m(\mathbb{F}_{2^r}) \right\} \cong \mathrm{Up}_m(\mathbb{F}_{2^r})$$

Note O is a subgroup of both $\Omega^+(2m, 2^r)$ and Ω^- . O is isomorphic to $\mathrm{Up}_m(\mathbb{F}_{2^r})$. So $|O| = (2^r)^{m(m-1)/2}$. Let

$$N' = \left\{ \begin{pmatrix} \mathrm{Id}_m & B \\ 0_m & \mathrm{Id}_m \end{pmatrix} : B \in \mathrm{Antisym}0(m, 2^r) \right\} \subset N$$

Then $N' \cong \mathrm{Antisym}0(m, 2^r)$. And for $M \in N'$,

$$M^T A_m^+ M = \begin{pmatrix} 0_m & \mathrm{Id}_m \\ 0_m & B^T \end{pmatrix}$$

and for $x = (y, z)$,

$$Q(Mx) = y^T z + z^T B^T z$$

And

$$\begin{aligned}
z^T B^T z &= \sum_{i,j} B_{i,j} z_i z_j \\
&= \sum_{i < j} 2B_{i,j} z_i z_j + \sum_{i=1}^n B_{i,i} z_i^2 \text{ since } B \in \text{Antisym}0(m, 2^r) \subset \text{Sym}(m, 2^r) \\
&= 0 \text{ since we are in characteristic 2 and } B_{i,i} = 0, \forall i
\end{aligned}$$

Therefore, $Q^+(Mx) = y^T z = Q^+(x)$ for all $x = (y, z)$. So $N'^+ \subset O^+(2n, p^r)$. Also, for $M = \begin{pmatrix} \text{Id}_m & B \\ 0_n & \text{Id}_m \end{pmatrix} \in N'$,

$$\begin{aligned}
M^T A_n^- M &= \begin{pmatrix} \text{Id}_n & 0_n \\ B^T & \text{Id}_m \end{pmatrix} \begin{pmatrix} 0_m^1 & \text{Id}_m \\ 0_m & 0_m^d \end{pmatrix} \begin{pmatrix} \text{Id}_m & B \\ 0_m & \text{Id}_m \end{pmatrix} \\
&= \begin{pmatrix} \text{Id}_m & 0_m \\ B^T & \text{Id}_m \end{pmatrix} \begin{pmatrix} 0_m^1 & \text{Id}_m \\ 0_m & 0_m^d \end{pmatrix}, \text{ since } B_{m,m} = 0 \\
&= \begin{pmatrix} 0_m^1 & \text{Id}_m \\ 0_m & B^T + 0_m^d \end{pmatrix}
\end{aligned}$$

So for $x = (y, z)$,

$$\begin{aligned}
Q^-(Mx) &= \mathbf{y}\mathbf{z}^T + y_m^2 + dz_m^2 + \mathbf{z}B^T\mathbf{z}^T \\
&= \mathbf{y}\mathbf{z}^T + y_m^2 + dz_m^2 \text{ since } \mathbf{z}B^T\mathbf{z}^T = 0 \text{ by the work shown above} \\
&= Q^-(x)
\end{aligned}$$

Therefore $N' \subset O^-(2n, p^r)$ as well. And

$$|N'| = (p^r)^{\sum_{k=1}^{m-1} k} = (p^r)^{m(m-1)/2}.$$

Then consider $N' \times O \subset \Omega^\epsilon(2m, 2^r)$ for both $\epsilon = +$ and $\epsilon = -$ (the operation is inherited from $N \times O$). Then we have

$$\begin{aligned}
|N' \times O| &= |N'| \cdot |O| \\
&= (2^r)^{n(n-1)/2} \cdot (2^r)^{m(m-1)/2} \\
&= 2^{rn(n-1)}
\end{aligned}$$

We learned the following argument from an early draft of [8]:

Note that for $M = \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \in O$,

$$\begin{aligned}
\delta_{2m, 2^r}^+(M) &= \text{rank}(\text{Id}_{2m} - M) \pmod{2} \\
&= \text{rank} \begin{pmatrix} \text{Id}_m + A & 0_m \\ 0_m & \text{Id}_m + (A^{-1})^T \end{pmatrix} \pmod{2} \\
&= 2 \text{rank}(A) \pmod{2} \\
&= 0
\end{aligned}$$

And for $M = \begin{pmatrix} \text{Id}_m & B \\ 0_m & \text{Id}_m \end{pmatrix} \in N'$,

$$\begin{aligned}
\delta_{2m,2^r}^+(M) &= \text{rank}(\text{Id}_{2m} - M) \pmod 2 \\
&= \text{rank} \begin{pmatrix} 0_m & B \\ 0_m & 0_m \end{pmatrix} \pmod 2 \\
&= \text{rank}(B) \pmod 2
\end{aligned}$$

And since B is symmetric with $B_{i,i} = 0$, $\forall i$, B determines an alternating symmetric bilinear form, and thus has even rank.

Thus, $\delta_{2m,2^r}^+(M) = 0$ for $M \in N'$ as well. Hence we have that both N' and O are in $\Omega^+(2m, 2^r) = SO^+(2m, 2^r) = \ker(\delta_{2m,2^r}^+)$. Therefore, $N' \rtimes O \subset \Omega^+(2n, 2^r)$. And

$$|N' \rtimes O| = 2^{2m(m-1)} = |\text{Syl}_2(\Omega^\epsilon(2m, 2^r))|.$$

Thus we can conclude that for $\epsilon = +, -$,

$$\text{Syl}_2(\Omega^\epsilon(2m, 2^r)) = N' \rtimes O$$

□

So since the p -Sylows of Ω^+ and Ω^- are the same, to prove Theorem 6.1 it suffices to consider $\epsilon = +$.

Remark 25. Recall that the action of O on N is given by

$$\begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \text{Id}_m & B \\ 0_m & \text{Id}_m \end{pmatrix} = \begin{pmatrix} \text{Id}_m & ABA^T \\ 0_m & \text{Id}_m \end{pmatrix}.$$

6.2.2 $n = 2m + 1, p \neq 2$

Since

$$\det L = \det \begin{pmatrix} -1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & 0_m & \text{Id}_m \\ \mathbf{0} & \text{Id}_m & 0_m \end{pmatrix} = (-1)^{m+1},$$

the form corresponding to L is equivalent to the form corresponding to

$$\text{diag}(1, -1, 1, -1, \dots, 1, -1, -1),$$

used in Grove (the diagonal matrix with entries $1, -1, 1, -1, \dots, 1, -1, -1$ on the diagonal). By Grove ([10], Theorem 9.11 and Corollary 9.12), for $p \neq 2$,

$$|\text{Syl}_p(O(2m + 1, p^r))| = p^{rm^2} = |\text{Syl}_p(\Omega(2m + 1, p^r))|.$$

So $\text{Syl}_p(O(2m + 1, p^r))$ maps injectively onto $\text{Syl}_p(\Omega(2m + 1, p^r))$; thus their p -Sylows must be isomorphic, and hence they have the same essential p -dimension.

Lemma 6.23. For $p \neq 2$

$$\begin{aligned}
&\text{Syl}_p(O(2n + 1, p^r)) \\
&= \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & 0_m \\ \mathbf{0} & 0_m & \text{Id}_n \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A & 0_m \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \text{Id}_m & B \\ \mathbf{0} & \mathbf{0} & \text{Id}_m \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^r}^m, A \in U_m(\mathbb{F}_{p^r}), B \in \text{Antisym}(m, p^r) \right\}.
\end{aligned}$$

Proof. Let

$$P = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & 0_m \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^r}^m \right\}.$$

This is a subgroup of $O(2m+1, p^r)$ isomorphic to $(\mathbb{F}_{p^r}^+)^m$. So $|P| = |(\mathbb{F}_{p^r}^+)^m| = p^{rm}$. Let

$$N = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \text{Id}_m & B \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} : B \text{ is anti-symmetric} \right\}.$$

This is a subgroup of $O(2m+1, p^r)$ isomorphic to $\text{Antisym}(m, p^r)$. So $|N| = |\text{Antisym}(m, p^r)| = (p^r)^{\frac{m(m-1)}{2}}$.
Let

$$O = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A & 0_m \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^r}) \right\}.$$

This is a group of $O(2m+1, p^r)$ isomorphic to $\text{Up}_m(\mathbb{F}_{p^r})$.

$$|O| = |\text{Up}_m(\mathbb{F}_{p^r})| = (p^r)^{\sum_{k=1}^{m-1} k} = (p^r)^{m(m-1)/2}.$$

A direct computation shows that $P \cdot O \cdot N = P \cdot N \cdot O = (P \times N) \rtimes O$ and

$$|(P \times N) \rtimes O| = p^{rm^2} = |\text{Syl}_p(O(2m+1, p^r))|.$$

Thus

$$\begin{aligned} & (P \times N) \rtimes O \\ &= \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & 0_m \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A & 0_m \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \text{Id}_m & B \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^r}^m, A \in \text{Up}_m(\mathbb{F}_{p^r}), B \in \text{Sym}(m, p^r) \right\}. \end{aligned}$$

is a p -Sylow of $O(2m+1, p^r)$. □

Corollary 6.24. For $p \neq 2$,

$$\text{Syl}_p(O(2m+1, p^r)) \cong ((\mathbb{F}_{p^r}^+)^m \times \text{Antisym}(m, p^r)) \rtimes \text{Up}_m(\mathbb{F}_{p^r}),$$

where the action of $\text{Up}_m(\mathbb{F}_{p^r})$ on $\text{Antisym}(m, p^r)$ is given by $A(B) = ABA^T$. and the action of $\text{Up}_m(\mathbb{F}_{p^r})$ on $(\mathbb{F}_{p^r}^+)^m$ is given by $A(\mathbf{x}) = \mathbf{x}A^T$.

6.3 Calculating the center

6.3.1 $n = 4$, any p

For $n = 4$, the action of $\text{Up}_2(\mathbb{F}_{p^r}) \cong \mathbb{F}_{p^r}$ on $\text{Antisym}(2, p^r) \cong \mathbb{F}_{p^r}$ is trivial and so $\text{Syl}_p(\Omega^+(2n, p^r)) \cong \mathbb{F}_{p^r} \times \mathbb{F}_{p^r}$. Thus the p -Sylow is abelian.

6.3.2 $n = 2m, n > 4$, any p

Lemma 6.25. For any $p, n > 2$,

$$Z(\text{Syl}_p(\Omega^+(2m, p^r))) = \left\{ \begin{pmatrix} \text{Id}_m & D \\ 0_m & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\} \cong \mathbb{F}_{p^r}^+ \cong (\mathbb{Z}/p\mathbb{Z})^r$$

For the proof of Lemma 6.25, we need the following lemma:

Lemma 6.26. Given $D \in \begin{cases} \text{Antisym}(m, p^r) & p \neq 2 \\ \text{Antisym}0(m, 2^r) & p = 2 \end{cases}$,

$$AD = D(A^{-1})^T \quad \forall A \in \text{Up}_m(\mathbb{F}_{p^r}) \Leftrightarrow D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix}.$$

Remark 26. This lemma is true for any $m \geq 2$.

Granting this lemma, we can prove Lemma 6.25:

Proof of Lemma 6.25. For $p \neq 2$,

$$\begin{aligned} \text{Syl}_p(\Omega^+(2m, p^r)) &= \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \text{Id}_m & B \\ 0_m & \text{Id}_m \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^r}), B \in \text{Antisym}(m, p^r) \right\} \\ &= \left\{ \begin{pmatrix} A & AB \\ 0_m & (A^{-1})^T \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^r}), B \in \text{Antisym}(m, p^r) \right\}. \end{aligned}$$

and

$$\begin{aligned} \text{Syl}_2(\Omega^+(2m, 2^r)) &= \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (A^{-1})^T \end{pmatrix} \begin{pmatrix} \text{Id}_m & B \\ 0_m & \text{Id}_m \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{2^r}), B \in \text{Antisym}0(m, 2^r) \right\} \\ &= \left\{ \begin{pmatrix} A & AB \\ 0_n & (A^{-1})^T \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{2^r}), B \in \text{Antisym}0(m, 2^r) \right\}. \end{aligned}$$

Note that for any p , given

$$\begin{pmatrix} A & AB \\ 0_m & (A^{-1})^T \end{pmatrix}, \begin{pmatrix} C & CD \\ 0_m & (C^{-1})^T \end{pmatrix} \in \begin{cases} O^+(2m, p^r) & p \neq 2 \\ \Omega^+(2m, 2^r) & p = 2 \end{cases},$$

we have

$$\begin{pmatrix} A & AB \\ 0_m & (A^{-1})^T \end{pmatrix}^{-1} \begin{pmatrix} C & CD \\ 0_m & (C^{-1})^T \end{pmatrix} \begin{pmatrix} A & AB \\ 0_m & (A^{-1})^T \end{pmatrix} = \begin{pmatrix} A^{-1}CA & A^{-1}CAB + A^{-1}CD(A^{-1})^T - B((A^{-1}CA)^{-1})^T \\ 0_m & ((A^{-1}CA)^{-1})^T \end{pmatrix}.$$

So

$$\begin{pmatrix} C & CD \\ 0_m & (C^{-1})^T \end{pmatrix} \in \begin{cases} Z(\text{Syl}_p(O^+(2m, p^r))) & p \neq 2 \\ Z(\text{Syl}_2(\Omega^+(2m, 2^r))) & p = 2 \end{cases}$$

if and only if

$$C \in Z(\text{Up}_m(\mathbb{F}_{p^r})) = \left\{ \begin{pmatrix} 1 & 0 & x \\ \mathbf{0} & \text{Id}_{m-2} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \right\}$$

and

$$CD = CB + CA^{-1}D(A^{-1})^T - B(C^{-1})^T, \text{ for all } A \in \text{Up}_m(\mathbb{F}_{p^r}), B \in \begin{cases} \text{Antisym}(m, p^r) & p \neq 2 \\ \text{Antisym}0(m, 2^r) & p = 2 \end{cases}.$$

Remark 27. For the remainder of this proof p can be any prime. (When $p = 2$, the negatives will go away, but the argument is the same.)

Choosing $A = \text{Id}_m$, we need

$$CD = CB + CD - B(C^{-1})^T.$$

So we must have

$$CB = B(C^{-1})^T$$

for all

$$B \in \begin{cases} \text{Antisym}(m, p^r) & p \neq 2 \\ \text{Antisym0}(m, 2^r) & p = 2 \end{cases}.$$

Write

$$C = \begin{pmatrix} 1 & \mathbf{0} & x \\ \mathbf{0} & \text{Id}_{m-2} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \in Z(\text{Up}_m(\mathbb{F}_{p^r})) = Z(\text{Syl}_p(\text{PSL}(m, p^r))).$$

(We calculated this center in our document on $\text{PSL}(n, p^r)$.) Then

$$(C^{-1})^T = \begin{pmatrix} 1 & \mathbf{0} & -x \\ \mathbf{0} & \text{Id}_m & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix}^T = \begin{pmatrix} 1 & \mathbf{0} & 0 \\ \mathbf{0} & \text{Id}_m & \mathbf{0} \\ -x & \mathbf{0} & 1 \end{pmatrix}.$$

Then for

$$B = (b_{i,j}) \in \begin{cases} \text{Antisym}(m, p^r) & p \neq 2 \\ \text{Antisym0}(m, 2^r) & p = 2 \end{cases},$$

we get

$$CB = \begin{pmatrix} -xb_{1,m} & b_{1,2} - xb_{2,m} & \cdots & b_{1,m-1} - xb_{m-1,m} & b_{1,m} \\ -b_{1,2} & 0 & b_{2,3} & \cdots & b_{2,m} \\ \vdots & & \ddots & & \vdots \\ -b_{1,m-1} & & \cdots & & b_{m-1,m} \\ -b_{1,m} & & \cdots & -b_{m-1,m} & 0 \end{pmatrix}$$

and

$$B(C^{-1})^T = \begin{pmatrix} -xb_{1,m} & b_{1,2} & \cdots & b_{1,m} \\ -b_{1,2} - xb_{2,m} & 0 & b_{2,3} & \cdots & b_{2,m} \\ \vdots & & \ddots & & \vdots \\ -b_{1,m-1} - xb_{m-1,m} & -b_{2,m-1} & \cdots & b_{m-1,m} \\ -b_{1,m} & -b_{2,m} & \cdots & -b_{m-1,m} & 0 \end{pmatrix}$$

So if $m > 2$, we must have $x = 0$, and hence $C = \text{Id}_m$.

Remark 28. This is where I need $m > 2$

So the other requirement above becomes

$$D = A^{-1}D(A^{-1})^T \Leftrightarrow AD = D(A^{-1})^T$$

for all $A \in \text{Up}_m(\mathbb{F}_{p^r})$. Then by Lemma 6.26, we get that

$$Z(\text{Syl}_p(O^+(2m, p^r))) = \left\{ \begin{pmatrix} \text{Id}_m & D \\ 0_m & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\}$$

□

Proof of Lemma 6.26.

\Leftarrow : This is a straightforward calculation.

\Rightarrow : We will prove this by induction.

Base Case: When $m = 2$, we can write $A = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ and $D = \begin{pmatrix} 0 & x \\ -x & 0 \end{pmatrix}$.

$$AD = \begin{pmatrix} -ax & x \\ -x & 0 \end{pmatrix} = D(A^{-1})^T.$$

So the condition that $AD = D(A^{-1})^T$ always holds. When $n = 3$, we can write $A = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}$ and

$$D = \begin{pmatrix} 0 & x & y \\ -x & 0 & z \\ -y & -z & 0 \end{pmatrix}.$$

$$AD = \begin{pmatrix} -ax - by & x - bz & y + az \\ -x - cy & -cz & z \\ -y & -z & 0 \end{pmatrix},$$

and

$$D(A^{-1})^T = \begin{pmatrix} -ax + acy - by & x - cy & y \\ -x + acy - bz & -cz & z \\ -y + az - acz + bz & -z & 0 \end{pmatrix}.$$

So in order for these to be equal for all A , we must have $y = 0$ and $z = 0$.

Induction Step: Write

$$D = \begin{pmatrix} 0 & d_{1,2} & d_{1,3} & \cdots & d_{1,m} \\ -d_{1,2} & 0 & d_{2,3} & \cdots & d_{2,m} \\ \vdots & & \ddots & & \vdots \\ -d_{1,m-1} & -d_{2,m-1} & \cdots & 0 & d_{m-1,m} \\ -d_{1,m} & -d_{2,m} & \cdots & -d_{m-1,m} & 0 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & a_{m-1,m} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix},$$

then

$$AD = \begin{pmatrix} 0 & d_{1,2} & d_{1,3} & \cdots & d_{1,m-1} & d_{1,m} \\ -d_{1,2} & 0 & d_{2,3} & \cdots & d_{2,m-1} & d_{2,m} \\ \vdots & & \ddots & & \vdots & \vdots \\ -d_{1,m-1} - a_{m-1,m}d_{1,m} & \cdots & -d_{m-2,m-1} - a_{m-1,m}d_{m-1,m} & -a_{m-1,m}d_{m-1,m} & d_{m-1,m} \\ -d_{1,m} & \cdots & \cdots & -d_{m-1,m} & 0 \end{pmatrix}$$

And

$$D(A^{-1})^T = \begin{pmatrix} 0 & d_{1,2} & d_{1,3} & \cdots & d_{1,m-2} & d_{1,m-1} - a_{m-1,m}d_{1,m} & d_{1,m} \\ -d_{1,2} & 0 & d_{2,3} & \cdots & d_{2,m-2} & d_{2,m-1} - a_{m-1,m}d_{2,m} & d_{2,m} \\ \vdots & & \ddots & & \vdots & \vdots & \vdots \\ -d_{1,m-1} & \cdots & d_{m-1,m-2} & -a_{m-1,m}d_{m-1,m} & d_{m-1,m} \\ -d_{1,m} & \cdots & -d_{m,m-2} & -d_{m-1,m} & 0 \end{pmatrix}$$

In order for these to be equal for all $a_{m-1,m}$, we must have $d_{k,m} = 0$ for all $k \neq m - 1$. Since $m > 2$, we

can pick

$$A = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ & & \ddots & & \vdots \\ 0 & \cdots & 1 & a_{m-2,m-1} & 0 \\ 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix},$$

so we get

$$AD = \begin{pmatrix} 0 & d_{1,2} & d_{1,3} & \cdots & d_{1,m-1} & d_{1,m} \\ -d_{1,2} & 0 & d_{2,3} & \cdots & d_{2,m-1} & d_{2,m} \\ \vdots & & \ddots & & \vdots & \vdots \\ -d_{1,m-2} - a_{m-2,m-1}d_{1,m-1} & \cdots & -a_{m-2,m-1}d_{m-2,m-1} & d_{m-2,m-1} & d_{m-2,m} + a_{m-2,m-1}d_{m-1,m} \\ -d_{1,m-1} & \cdots & -d_{m-2,m-1} & 0 & d_{m-1,m} \\ -d_{1,m} & \cdots & \cdots & -d_{m-1,m} & 0 \end{pmatrix}$$

And

$$D(A^{-1})^T = \begin{pmatrix} 0 & \cdots & d_{1,m-3} & d_{1,m-2} - a_{m-2,m-1}d_{1,m-1} & d_{1,m-1} & d_{1,m} \\ -d_{1,2} & \cdots & d_{2,m-3} & d_{2,m-2} - a_{m-2,m-1}d_{2,m-1} & d_{2,m-1} & d_{2,m} \\ \vdots & & \ddots & & \vdots & \vdots \\ -d_{1,m-2} & \cdots & -d_{m-2,m-3} & -a_{m-2,m-1}d_{m-2,m-1} & d_{m-2,m-1} & d_{m-2,m} \\ -d_{1,m-1} & \cdots & -d_{m-1,m-3} & -d_{m-2,m-1} & 0 & d_{m-1,m} \\ -d_{1,m} & \cdots & -d_{m,m-3} & -d_{m-2,m} + a_{m-2,m-1}d_{m-1,m} & -d_{m-1,m} & 0 \end{pmatrix}$$

In order for these to be equal for all $a_{m-2,m}$, we must have $d_{k,m-1} = 0$ for all $k \neq m-2$. In particular, we get that $d_{mm,m-1} = d_{m-1,m} = 0$. Thus $d_{k,m} = 0$ for all k . So the matrix

$$D' = \begin{pmatrix} d_{1,1} & d_{1,2} & d_{1,3} & \cdots & d_{1,m-1} \\ -d_{1,2} & d_{2,2} & d_{2,3} & \cdots & d_{2,m-1} \\ \vdots & & \ddots & & \vdots \\ -d_{1,m-2} & -d_{2,m-2} & \cdots & d_{m-2,m-2} & d_{m-2,m} \\ -d_{1,m-1} & -d_{2,m-1} & \cdots & -d_{m-2,m-1} & d_{m-1,m-1} \end{pmatrix}$$

satisfies the condition $A'D' = D'(A'^{-1})^T$ for all $A' \in U_{m-1}(\mathbb{F}_{p^r})$. By induction, we conclude that

$$D' = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-3} \end{pmatrix},$$

and hence

$$D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix}.$$

□

6.3.3 $n = 2m + 1, p \neq 2$

Lemma 6.27. *For any p ,*

$$Z(\text{Syl}_p(\Omega(2m+1, p^r))) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & Id_m & D \\ \mathbf{0} & 0_m & Id_m \end{pmatrix} : \mathbf{x} = (x_1, 0, \dots, 0), D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\} \cong (\mathbb{F}_{p^r}^+)^2$$

Proof.

Case 1: $n = 5$

The proof for $n > 5$ uses the result for $Z(\text{Syl}_p(O^+(2m, p^r)))$, which we only calculated for $m > 2$. So we must prove the case $m = 2$ separately:

For $m = 2$, the action of $\text{Up}_2(\mathbb{F}_{p^r}) \cong \mathbb{F}_{p^r}$ on $\text{Antisym}(2, p^r) \cong \mathbb{F}_{p^r}$ is trivial. And the action on $\mathbb{F}_{p^r}^2$ is given by $a(x, y) = (x + ay, y)$. So we have $\text{Syl}_p(\Omega(5, p^r)) \cong \mathbb{F}_{p^r}^2 \rtimes \mathbb{F}_{p^r}^2$, where the action of $\mathbb{F}_{p^r}^2$ (2nd copy) on $\mathbb{F}_{p^r}^2$ (1st copy) is given by $(b, a)((x, y)) = (x + ay, y)$. An element $((x, y), (a, b))$ is in the center if and only if for all $((w, z), (d, c))$ we have

$$((w, z), (d, c))((x, y), (b, a)) = ((x, y), (b, a))((w, z), (d, c))$$

Note that

$$((w, z), (d, c))((x, y), (b, a)) = ((x + w + cy, y + z), (b + d, a + c))$$

and

$$((x, y), (b, a))((w, z), (d, c)) = ((x + w + az, y + z), (b + d, a + c))$$

These will be equal for all $((w, z), (d, c))$ if and only if $a = 0 = y$. Therefore the center is given by

$$\{((x, 0), (b, 0)) : x, b \in \mathbb{F}_{p^r}\}.$$

Translating this back into the original form in a matrix, we get that the center is

$$Z(\text{Syl}_p(\Omega(5, p^r))) = \left\{ \begin{pmatrix} 1 & 0 & 0 & w & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ w & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & b \\ 0 & 0 & 1 & -b & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & w & 0 \\ 0 & 1 & 0 & 0 & b \\ 0 & 0 & 1 & -b & 0 \\ w & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \right\} \cong (\mathbb{F}_{p^r})^2$$

Case 2: $n > 5$

Since

$$\text{Syl}_p(\Omega(2m + 1, p^r)) \cong (\mathbb{F}_{p^r}^+)^m \rtimes (\text{Antisym}(m, p^r) \rtimes \text{Up}_m(\mathbb{F}_{p^r})) \cong (\mathbb{F}_{p^r}^+)^m \rtimes \text{Syl}_p(\Omega^+(2m, p^r)),$$

we can conclude that

$$Z(\text{Syl}_p(\Omega(2m + 1, p^r))) \cap (\{\mathbf{0}\} \times \text{Syl}_p(\Omega^+(2m, p^r)))$$

must be a subset of $Z(\text{Syl}_p(O^+(2m, p^r)))$, which we proved in the previous section to be

$$Z(\text{Syl}_p(\Omega^+(2m, p^r))) = \left\{ \begin{pmatrix} \text{Id}_m & D \\ 0_m & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\} \cong \mathbb{F}_{p^r}^+ \cong (\mathbb{Z}/p\mathbb{Z})^r \text{ (for } n > 2\text{)}.$$

Thus the center of $(\mathbb{F}_{p^r}^+)^m \rtimes (\text{Antisym}(m, p^r) \rtimes \text{Up}_m(\mathbb{F}_{p^r}))$ is a subset of

$$(\mathbb{F}_{p^r}^+)^m \rtimes \left\{ \begin{pmatrix} \text{Id}_m & D \\ 0_m & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\} = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^r}^m, D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\}$$

Given

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} \\ \mathbf{y}^T & A & AB \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix} \in \text{Syl}_p(O(2m + 1, p^r)),$$

we have that

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} \\ \mathbf{y}^T & A & AB \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} + \mathbf{x}(A^{-1})^T \\ \mathbf{x}^T + \mathbf{y}^T & A & \mathbf{x}^T \mathbf{y} + AB + D(A^{-1})^T \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix}$$

and

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} \\ \mathbf{y}^T & A & AB \\ \mathbf{0} & 0_n & (A^{-1})^T \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} + \mathbf{y} \\ \mathbf{y}^T + A\mathbf{x}^T & A & \mathbf{x}\mathbf{y}^T + AD + AB \\ \mathbf{0} & 0_m & (A^{-1})^T \end{pmatrix}$$

So in order for

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix}$$

to be in the center, we need $\mathbf{x}^T = A\mathbf{x}^T$, $\mathbf{x} = \mathbf{x}(A^{-1})^T$, and $AD = D(A^{-1})^T$ for all choices of A . By the work on even orthogonal groups, $AD = D(A^{-1})^T$ is satisfied if and only if

$$D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix}.$$

Note that the k th entry of $\mathbf{x} = A\mathbf{x}^T$ is given by $x_k + \sum_{i=k+1}^m x_i a_{k,i}$. In order for this to be equal to x_k for all $a_{k,i}$, must have $x_i = 0$ for all $i > 1$. So $\mathbf{x} = (x_1, 0, \dots, 0)$. In this case $\mathbf{x} = \mathbf{x}(A^{-1})^T$ will be satisfied as well. Therefore the center is

$$Z(\text{Syl}_p(\Omega(2m+1, p^r))) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} : \mathbf{x} = (x_1, 0, \dots, 0), D = \begin{pmatrix} 0 & x & \mathbf{0} \\ -x & 0 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & 0_{m-2} \end{pmatrix} \right\} \cong (\mathbb{F}_{p^r}^+)^2$$

□

6.4 Maximizing L_s

6.4.1 $n = 2m$

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element of

$$\begin{cases} \widehat{\text{Antisym}}(n, p^r) \cong (\mathbb{F}_{p^r}^+)^{\widehat{m(m-1)/2}} & p \neq 2 \\ \widehat{\text{Antisym0}}(m, 2^r) \cong (\mathbb{F}_{2^r}^+)^{\widehat{m(m-1)/2}} & p = 2 \end{cases}$$

is of the form

$$\mathbf{d} \mapsto \psi(\mathbf{bd}^T),$$

where $\mathbf{d}, \mathbf{b} \in (\mathbb{F}_{p^r})^{m(m-1)/2}$.

Proposition 6.28. *For any p , the characters extending a non-trivial central character are $\psi_{(b_j)}$ with $b_1 \neq 0$.*

Proof. $\psi_{(b_j)}$ restricted to the center of the p -Sylow is ψ_{b_1} . This is non-trivial if and only if $b_1 \neq 0$. □

We will use Wigner-Mackey Theory with

$$G = \text{Syl}_p(\Omega^+(2m, p^r)) = \begin{cases} \text{Antisym}(m, p^r) \rtimes \text{Up}_m(\mathbb{F}_{p^r}) & p \neq 2 \\ \text{Antisym0}(m, 2^r) \rtimes \text{Up}_m(\mathbb{F}_{2^r}) & p = 2 \end{cases}$$

to see what is the minimum dimension of an irreducible representation of G with central character ψ_{b_1} with $b_1 = b \neq 0$. So we have

$$\Delta = \begin{cases} \text{Antisym}(m, p^r) \rtimes \{\text{Id}_m\} & p \neq 2 \\ \text{Antisym}0(m, 2^r) \rtimes \{\text{Id}_m\} & p = 2 \end{cases}$$

and

$$L = \{0_n\} \rtimes \text{Up}_m(\mathbb{F}_{p^r}).$$

Let

$$B = \begin{pmatrix} 0 & b_1 & \cdots & & b_{m-1} \\ -b_1 & 0 & b_m & \cdots & b_{2m-3} \\ \vdots & & \ddots & & \vdots \\ -b_{m-2} & \cdots & & 0 & b_{m(m-1)/2} \\ -b_{m-1} & \cdots & & -b_{m(m-1)/2} & 0 \end{pmatrix}$$

be the matrix corresponding to $s = (b_1, \dots, b_{m(m-1)/2})$ with $b_1 \neq 0$. (When $p = 2$, the negatives go away.)

Remark 29. In all of the following, we view $\psi_{(b_j)}$ as a map on

$$\begin{cases} \Delta \cong \text{Antisym}(m, p^r) \cong \mathbb{F}_{p^r}^{m(m-1)/2} & p \neq 2 \\ \Delta \cong \text{Antisym}0(m, 2^r) \cong \mathbb{F}_{2^r}^{m(m-1)/2} & p = 2 \end{cases}.$$

So $\psi_{(b_j)}(D, \text{Id}) = \psi_{(b_j)}(D) = \psi(\mathbf{b} \cdot \mathbf{d})$, where $\mathbf{b} = (b_j)$ and \mathbf{d} is the vector corresponding to the matrix D .

The action of $h \in \text{Syl}_p(\Omega^+(2m, p^r))$ on $\widehat{\Delta}$ is given by

$${}^h\psi(D, \text{Id}_m) = \psi(h^{-1}(D, \text{Id}_m)h).$$

So for $h = (0_m, H^{-1})$, the action on $\psi_{(b_j)}$ is given by

$${}^h\psi_{(b_j)}(D, \text{Id}_m) = \psi_{(b_j)}((0_m, H)(D, \text{Id}_m)(0_m, H^{-1})).$$

So $(0_m, H^{-1}) \in L_s$ if and only if

$$\psi_{(b_j)}((0_m, H)(D, \text{Id}_m)(0_m, H^{-1})) = \psi_{(b_j)}(D, \text{Id}_m)$$

for all

$$\mathbf{d} \in (\mathbb{F}_{p^r}^+)^{m(m-1)/2} \text{ corresponding to } D \in \begin{cases} \text{Antisym}(m, p^r) & p \neq 2 \\ \text{Antisym}0(m, 2^r) & p = 2 \end{cases}.$$

Let \mathbf{hdh}^T be the vector corresponding to HDH^T . Then since

$$\psi_{(b_j)}((0_m, H)(D, \text{Id}_m)(0_m, H^{-1})) = \psi(\mathbf{b} \cdot \mathbf{hdh}^T),$$

and

$$\psi_{(b_j)}(D, \text{Id}_m) = \psi(\mathbf{b} \cdot \mathbf{d}).$$

we get that $(0_m, H^{-1}) \in L_s$ if and only if

$$\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1$$

for all

$$\mathbf{d} \in (\mathbb{F}_{p^r})^{m(m-1)/2} \text{ corresponding to } D \in \begin{cases} \text{Antisym}(m, p^r) & p \neq 2 \\ \text{Antisym}0(m, 2^r) & p = 2 \end{cases}.$$

Write

$$H = \begin{pmatrix} 1 & h_{1,2} & h_{1,3} & \cdots & h_{1,n} \\ 0 & 1 & h_{2,3} & \cdots & h_{2,m} \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & h_{m-1,m} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 0 & d_{1,2} & d_{1,3} & \cdots & d_{1,m} \\ -d_{1,2} & 0 & d_{2,3} & \cdots & d_{2,m} \\ \vdots & & \ddots & & \vdots \\ -d_{1,m-1} & -d_{2,m-1} & \cdots & 0 & d_{m-1,m} \\ -d_{1,m} & -d_{2,m} & \cdots & -d_{m-1,m} & 0 \end{pmatrix}.$$

Proposition 6.29. For any p : for $s = (b_i)$, $b_1 \neq 0$, the maximum of $|L_s|$ is

$$|L_s| = |\mathbb{F}_{p^r}| \cdot |U_{m-2}(\mathbb{F}_{p^r})| = p^{r[(m-2)(m-3)/2+1]}.$$

And this maximum is achieved when $s = (b, 0, \dots, 0)$.

Corollary 6.30. For $s = (b, 0, \dots, 0)$, λ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|Up_m(\mathbb{F}_{p^r})|}{|\mathbb{F}_{p^r}| \cdot |U_{m-2}(\mathbb{F}_{p^r})|} = |(\mathbb{F}_{p^r}^{m-2})^2| = p^{2r(m-2)}.$$

Proof of Proposition 6.29.

Step 1: Proving that for any $\mathbf{s} = (b_i)$, $b_1 \neq 0$, $|L_s| \leq |\mathbb{F}_{p^r}| \cdot |U_{m-2}(\mathbb{F}_{p^r})| = p^{2r(m-2)}$.

In all the following, in characteristic 2, the negatives will go away, but the argument is the same.

Calculation 1. For $j_0 > 2$, choose $d_{i,j} = 0$ except for $d_{1,j_0} = -d_{j_0,1}$.

Then

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = \sum_{i=2}^{j_0-1} h_{i,j_0} d_{1,j_0} B_{1,i} = d_{1,j_0} \left(\sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} \right)$$

If $\sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} \neq 0$, then as we run through all the values for d_{1,j_0} , we will get that $\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})$ runs through all the values of \mathbb{F}_{p^r} . And since ψ is non-trivial, this means that $\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}))$ cannot always equal 1. This is a contradiction. So we must have

$$\sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} = 0$$

for all choices of $j_0 > 2$. Recall that $B_{1,2} = b_1 \neq 0$. So, for all $j_0 > 2$, given h_{i,j_0} for $i > 2$, the above dictates h_{2,j_0} : If we know h_{i,j_0} for $i > 1$, then we have

$$\sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} = 0 \Rightarrow h_{2,j_0} = \frac{-1}{B_{1,2}} \sum_{i=3}^{j_0-1} h_{i,j_0} B_{1,i}.$$

(In particular, note $h_{2,3} = 0$.) For $3 \leq k \leq n$, if $B_{1,k} \neq 0$, then for all $j_0 > 2$, given h_{i,j_0} for $i \neq 1, k$, the above dictates h_{k,j_0} : If we know h_{i,j_0} for $i \neq 1, k$, then we have

$$\sum_{i=2}^{j_0-1} h_{i,j_0} B_{1,i} = 0 \Rightarrow h_{k,j_0} = \frac{-1}{B_{1,k}} \sum_{i=2, i \neq k}^{j_0-1} h_{i,j_0} B_{1,i}.$$

Calculation 2. Now for $j_0 > 2$, choose $d_{i,j} = 0$ except for $d_{2,j_0} = -d_{j_0,2}$.

Then

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = d_{2,j_0} \left(-B_{1,2}h_{1,j_0} + \sum_{i=2}^{j_0} B_{1,i}h_{i,j_0}h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i}h_{i,j_0} \right)$$

By the same reasoning as before, we must have

$$-B_{1,2}h_{1,j_0} + \sum_{i=2}^{j_0} B_{1,i}h_{i,j_0}h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i}h_{i,j_0} = 0$$

for all choices of $j_0 > 2$. Recall that $B_{1,2} = b_1 \neq 0$. So for all $j_0 > 2$, given h_{i,j_0} for $i > 2$, the above dictates h_{1,j_0} : If we know $h_{1,2}$ and h_{i,j_0} for $i > 1$, then we have

$$-B_{1,2}h_{1,j_0} + \sum_{i=2}^{j_0} B_{1,i}h_{i,j_0}h_{1,2} + \sum_{k=3}^{j_0-1} B_{2,k}h_{k,j_0} = 0 \Rightarrow h_{1,j_0} = \frac{1}{B_{1,2}} \left(\sum_{i=2}^{j_0} B_{1,i}h_{i,j_0}h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i}h_{i,j_0} \right)$$

Thus we can conclude that for all $s = (b_i)$ with $b_1 \neq 0$,

$$|L_s| \leq |\{H : H_{2,j} \text{ fixed}, \forall j > 2, H_{1,j} \text{ fixed}, \forall j > 2\}| = |\mathbb{F}_{p^r}| \cdot |U_{m-2}(\mathbb{F}_{p^r})| = p^{r[(m-2)(m-3)/2+1]}.$$

Step 2: Exhibiting that the max is achieved when $\mathbf{s} = (b, \mathbf{0}, \dots, \mathbf{0})$ with $\mathbf{b} \neq \mathbf{0}$.

Let B be the matrix corresponding to $s = (b, \mathbf{0}, \dots, \mathbf{0})$. So since the only nonzero entry of B is $B_{1,2} = b$, we have that

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = b(HDH^T - D)_{1,2} = b \left(\left[\sum_{l=2}^n [h_{2,l} \left(\sum_{k=1}^l d_{k,l}h_{1,k} - \sum_{k=l+1}^{m-1} d_{l,k}h_{1,k} \right)] - d_{1,2} \right] \right).$$

By the first calculation above, we have that for $j_0 > 2$,

$$h_{2,j_0} = \frac{-1}{B_{1,2}} \sum_{i=3}^{j_0-1} h_{i,j_0} B_{1,i} = 0.$$

So we have

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = b \left(\left[\sum_{k=1}^2 d_{k,2}h_{1,k} - \sum_{k=3}^{m-1} d_{2,k}h_{1,k} \right] - d_{1,2} \right)$$

By the second calculation above, we have that for $j_0 > 2$,

$$\begin{aligned} h_{1,j_0} &= \frac{1}{B_{1,2}} \left(\sum_{i=2}^{j_0} B_{1,i}h_{i,j_0}h_{1,2} + \sum_{i=3}^{j_0-1} B_{2,i}h_{i,j_0} \right) \\ &= h_{2,j_0}h_{1,2} \\ &= 0 \end{aligned}$$

So we have

$$\begin{aligned} \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) &= b \left(\left[\sum_{k=1}^2 d_{k,2}h_{1,k} \right] - d_{1,2} \right) \\ &= b(d_{1,2}h_{1,1} + d_{2,2}h_{1,2} - d_{1,2}) \\ &= 0 \text{ since } h_{1,1} = 0, d_{2,2} = 0 \end{aligned}$$

Thus we have shown that $(0_m, H^{-1}) \in L_s$ if and only if $h_{2,j} = 0, \forall j > 2$ and $h_{1,j} = 0, \forall j > 2$. Therefore,

$$L_s = \{(0_m, H^{-1}) : H_{1,j} = 0, \forall j > 2, H_{2,j} = 0, \forall j > 2\}.$$

So $|L_s| = |\mathbb{F}_{p^r}| \cdot |U_{m-2}(\mathbb{F}_{p^r})| = p^{r[(m-2)(m-3)/2+1]}$.

□

6.4.2 $n = 2m + 1, p \neq 2$

Case 1: $n = 5$

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element of

$$(\mathbb{F}_{p^r}^+)^2 \times \widehat{\text{Antisym}}(2, p^r) \cong \widehat{\text{Antisym}}(2, p^r) \times (\mathbb{F}_{p^r}^+)^2 \cong \widehat{(\mathbb{F}_{p^r}^+)^3}$$

is of the form

$$\mathbf{d} \mapsto \psi(\mathbf{bd}^T),$$

where $\mathbf{d}, \mathbf{b} \in (\mathbb{F}_{p^r})^3$.

Proposition 6.31. *For $p \neq 2$, the characters of $(\mathbb{F}_{p^r}^+)^3$ extending a non-trivial central character of the p -Sylow are $\psi_{(b_j)}$ with $b_1 \neq 0$ or $b_3 \neq 0$.*

Proof. $\psi_{(b_j)}$ restricted to the center is $\psi_{b_1} \times \psi_{b_3}$. This is non-trivial if and only if $b_1 \neq 0$ or $b_3 \neq 0$. \square

We will use Wigner-Mackey Theory with

$$G = \text{Syl}_p(\Omega(5, p^r)) = (\text{Antisym}(2, p^r) \times (\mathbb{F}_{p^r}^+)^2) \rtimes \text{Up}_2(\mathbb{F}_{p^r})$$

to see what is the minimum dimension of an irreducible representation of G with non-trivial central character. So we have $\Delta = (\text{Antisym}(2, p^r) \times (\mathbb{F}_{p^r}^+)^2) \rtimes \{\text{Id}_2\}$, $L = (\{0_2\} \times \{\mathbf{0}\} \rtimes \text{Up}_2(\mathbb{F}_{p^r}))$. Recall that $L_s = \text{Stab}_L(\psi_s)$.

Proposition 6.32. *For $p \neq 2$, $s = (b_1, b_2, b_3)$, $L_s \cong \begin{cases} \text{Up}_2(\mathbb{F}_{p^r}), & b_3 = 0 \\ \{0\}, & b_3 \neq 0 \end{cases}$.*

Corollary 6.33. *For $s = (b_1, b_2, b_3)$, λ trivial, $\dim(\theta_{s, \lambda}) = \begin{cases} 1, & b_3 = 0 \\ p^r, & b_3 \neq 0 \end{cases}$.*

Proof of Proposition 6.32. Just like before, we have that $(\mathbf{0}, 0_n, H^{-1}) \in L_s$ if and only if for all $(\mathbf{d}, \mathbf{x}) \in (\mathbb{F}_{p^r})^3$ corresponding to $(D, \mathbf{x}) \in \text{Antisym}(2, p^r) \times (\mathbb{F}_{p^r}^+)^2$,

$$\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}, \mathbf{x}H^T - \mathbf{x})) = 1.$$

Given $H = \begin{pmatrix} 1 & h \\ 0 & 1 \end{pmatrix} \in \text{Up}_2(\mathbb{F}_{p^r})$, $D = \begin{pmatrix} 0 & d \\ -d & 0 \end{pmatrix} \in \text{Antisym}(2, p^r)$, $\mathbf{x} = (x, y) \in (\mathbb{F}_{p^r})^2$,

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}, \mathbf{x}H^T - \mathbf{x}) = b_3(\mathbf{x}H^T - \mathbf{x}) = b_3(h, 0)$$

Thus if $b_3 = 0$, then we can have any h . And if $b_3 \neq 0$, we must have $h = 0$. Hence we have

$$L_s \cong \begin{cases} \text{Up}_2(\mathbb{F}_{p^r}), & b_3 = 0 \\ \{0\}, & b_3 \neq 0 \end{cases}.$$

\square

Case 2: $n > 5$

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element of

$$(\mathbb{F}_{p^r}^+)^m \times \widehat{\text{Antisym}}(m, p^r) \cong (\mathbb{F}_{p^r}^+)^{m+m(m-1)/2}$$

is of the form $\mathbf{d} \mapsto \psi(\mathbf{bd}^T)$, where $\mathbf{d}, \mathbf{b} \in (\mathbb{Z}/p\mathbb{Z})^{m+m(m-1)/2}$.

Proposition 6.34. For $p \neq 2$, the characters of $(\mathbb{F}_{p^r})^{m+m(m-1)/2}$ extending a non-trivial central character of the p -Sylow are $\psi_{(a_j, b_j)}$ with $(a_1, b_1) \neq (0, 0)$.

Proof. $\psi_{(a_j, b_j)}$ restricted to the center is ψ_{a_1, b_1} . This is non-trivial if and only if $(a_1, b_1) \neq (0, 0)$. \square

We will use Wigner-Mackey Theory with

$$G = \text{Syl}_p(\Omega(2m+1, p^r)) = ((\mathbb{F}_{p^r}^+)^m \times \text{Antisym}(m, p^r)) \rtimes \text{Up}_m(\mathbb{F}_{p^r})$$

to see what is the minimum dimension of an irreducible representation of G with non-trivial central character. So we have

$$\Delta = ((\mathbb{F}_{p^r}^+)^m \times \text{Antisym}(m, p^r)) \rtimes \{\text{Id}_m\}, \quad L = (\{\mathbf{0} \times \{\mathbf{0}_m\}\}) \rtimes \text{Up}_m(\mathbb{F}_{p^r}).$$

Recall that $(\mathbf{0}, 0_m, H^{-1}) \in L_s$ if and only if for all $(\mathbf{x}, \mathbf{d}) \in (\mathbb{F}_{p^r})^{m+m(m-1)/2} = (\mathbf{x}, D) \in (\mathbb{F}_{p^r}^+)^m \times \text{Antisym}(m, p^r)$,

$$\psi_{(a_j, b_j)}((\mathbf{0}, 0_m, H)(\mathbf{x}, D, \text{Id}_m)(\mathbf{0}, 0_m, H^{-1})) = \psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}_m)$$

Remark 30. In all of the following, we view $\psi_{(a_j, b_j)}$ as a map on

$$\Delta \cong ((\mathbb{F}_{p^r}^+)^m \times \text{Antisym}(m, p^r)) \cong (\mathbb{F}_{p^r}^+)^{m+m(m-1)/2}.$$

So $\psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}) = \psi_{(a_j, b_j)}(\mathbf{x}, D) = \psi((\mathbf{a}, \mathbf{b}) \cdot (\mathbf{x}, \mathbf{d}))$, where $(\mathbf{a}, \mathbf{b}) = (a_j, b_j)$ and (\mathbf{x}, \mathbf{d}) is the vector corresponding to (\mathbf{x}, D) .

Let \mathbf{hdh}^T denote the vector corresponding to HDH^T . Then since

$$\psi_{(a_j, b_j)}((\mathbf{0}, 0_m, H)(\mathbf{x}, D, \text{Id}_m)(\mathbf{0}, 0_m, H^{-1})) = \psi((\mathbf{a}, \mathbf{b}) \cdot (\mathbf{x}H^T, \mathbf{hdh}^T))$$

and

$$\psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}_m) = \psi((\mathbf{a}, \mathbf{b}) \cdot (\mathbf{x}, \mathbf{d})),$$

We get that $(\mathbf{0}, 0_m, H^{-1}) \in L_s$ if and only if for all $(\mathbf{x}, \mathbf{d}) \in (\mathbb{F}_{p^r})^{m+m(m-1)/2}$ corresponding to $(\mathbf{x}, D) \in \text{Antisym}(m, p^r) \times (\mathbb{F}_{p^r}^+)^m$,

$$\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1.$$

Proposition 6.35. For $m > 2$, $p \neq 2$, $s = (a_i, b_i)$,

$$\max_{s: b_1 \neq 0} |L_s| = |\text{Up}_{m-2}(\mathbb{F}_{p^r})| = p^{r(m-2)(m-3)/2}.$$

And this max is achieved when $s = (0, \dots, 0, b_1, 0, \dots, 0)$.

Corollary 6.36. For $n > 2$, $p \neq 2$, $s = (0, \dots, 0, b_1, 0, \dots, 0)$, λ trivial,

$$\dim(\theta_{s, \lambda}) = \frac{|L|}{|L_s|} = \frac{|\text{Up}_n(\mathbb{F}_{p^r})|}{|\text{Up}_{m-2}(\mathbb{F}_{p^r})|} = |\mathbb{F}_{p^r}^{(m-1)(m-2)}| = p^{r(m-1)(m-2)}.$$

Proof of Proposition 6.35. If we take $\mathbf{x} = 0$, then $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1$ reduces to the condition for $\Omega^+(2m, p^r)$. So if $b_1 \neq 0$, we must have L_s is a subset of that calculated in the document for even orthogonal groups. Thus

$$|L_s| \leq |\{H : H_{2,j} \text{ fixed}, \forall j > 2, H_{1,j} \text{ fixed}, \forall j > 2\}| = p^{r[(m-2)(m-3)/2+1]}.$$

If $b_i = 0$ for $i \neq 1$, then we get

$$L_s \subset \{H \in \text{Up}_m(\mathbb{F}_{p^r}) : H_{1,j} = 0, \forall j \neq 2, H_{2,j} = 0, \forall j > 2\}.$$

Given H of this form, we have $\mathbf{b} \cdot (\mathbf{h}\mathbf{d}\mathbf{h}^T - \mathbf{d}) = 0$. Then for $\mathbf{a} = 0$,

$$\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{h}\mathbf{d}\mathbf{h}^T - \mathbf{d}) = 0.$$

So for $(0, \dots, 0, b_1, 0, \dots, 0)$,

$$L_s = \{H \in \text{Up}_m(\mathbb{F}_{p^r}) : H_{1,j} = 0, \forall j \neq 2, H_{2,j} = 0, \forall j > 2\}.$$

□

Proposition 6.37. For $m > 2$, $p \neq 2$, $s = (a_i, b_i)$,

$$\max_{s: a_1 \neq 0} |L_s| = |\text{Up}_{m-1}(\mathbb{F}_{p^r})| = p^{r(m-1)(m-2)/2}.$$

And this max is achieved when $s = (a_1, 0, \dots, 0)$.

Corollary 6.38. For $m > 2$, $p \neq 2$, $s = (a_1, 0, \dots, 0)$, λ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|\text{Up}_m(\mathbb{F}_{p^r})|}{|\text{Up}_{m-1}(\mathbb{F}_{p^r})|} = |\mathbb{F}_{p^r}^{m-1}| = p^{r(m-1)}.$$

Proof of Proposition 6.37. If we take $\mathbf{d} = \mathbf{0}$ then $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{h}\mathbf{d}\mathbf{h}^T - \mathbf{d})) = 1$ reduces to $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x})) = 1$. Write

$$H = \begin{pmatrix} 1 & h_{1,2} & h_{1,3} & \cdots & h_{1,m} \\ 0 & 1 & h_{2,3} & \cdots & h_{2,m} \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & h_{m-1,m} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Then

$$\begin{aligned} \mathbf{x}H^T &= (x_1, \dots, x_m) \begin{pmatrix} 1 & 0 & \cdots & 0 \\ h_{1,2} & 1 & 0 & \cdots & 0 \\ h_{1,3} & h_{2,3} & 1 & & 0 \\ \vdots & & \ddots & \ddots & \vdots \\ h_{1,m} & h_{2,m} & \cdots & h_{m-1,m} & 1 \end{pmatrix} \\ &= \left(\sum_{k=1}^m x_k h_{1,k}, \dots, x_{m-1} + x_m h_{m-1,m}, x_m \right) \end{aligned}$$

So

$$\mathbf{x}H^T - \mathbf{x} = \left(\sum_{k=2}^m x_k h_{1,k}, \dots, x_m h_{m-1,m}, 0 \right).$$

Thus

$$\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) = \sum_{k=1}^{m-1} a_k \cdot \left(\sum_{j=k+1}^m x_j h_{k,j} \right)$$

Calculation 1. For $j_0 > 1$, $\mathbf{x} = (x_i)$ with $x_i = 0$ except for x_{j_0} .

Then we get

$$\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) = \sum_{k=1}^{j_0-1} a_k \cdot x_{j_0} h_{k,j_0} = x_{j_0} \left(\sum_{k=1}^{j_0-1} a_k h_{k,j_0} \right)$$

So for all $j_0 > 1$, we must have

$$\sum_{k=1}^{j_0-1} a_k h_{k,j_0} = 0.$$

So if $a_1 \neq 0$, given h_{i,j_0} for $i \neq 1, k$, the above dictates h_{1,j_0} :

$$h_{1,j_0} = \frac{-1}{a_1} \sum_{k=2}^{j_0-1} a_k h_{k,j_0}.$$

Therefore,

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed } \forall j \neq 1\}| = |\text{Up}_{n-1}(\mathbb{F}_{p^r})| = p^{r(n-1)(n-2)/2}.$$

If $a_i = 0$ for $i \neq 0$, then we get from the calculation above that

$$h_{1,j_0} = \frac{-1}{a_1} \sum_{k=2}^{j_0-1} a_k h_{k,j_0} = 0.$$

So

$$\begin{aligned} \mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) &= \sum_{k=1}^{m-1} a_k \cdot \left(\sum_{j=k+1}^m x_j h_{k,j} \right) \\ &= a_1 \cdot \left(\sum_{j=2}^m x_j h_{1,j} \right), && \text{since } a_i = 0, i > 1 \\ &= 0 && \text{since } h_{1,j} = 0, j > 1 \end{aligned}$$

So we get that for $s = (a_1, 0, \dots, 0)$ with $a_1 \neq 0$,

$$L_s = \{H : H_{1,j} = 0, \forall j \neq 1\}.$$

□

6.5 Proof

Proof of Theorem 6.1.

Case 1 ($n = 4$).

Proof of Theorem 6.1 for $n = 2$. For $p \neq 2$, the action of $\text{Up}_2(\mathbb{F}_{p^r}) \cong \mathbb{F}_{p^r}$ on $\text{Antisym}(2, p^r) \cong \mathbb{F}_{p^r}$ is trivial and so $\text{Syl}_p(\Omega^+(2n, p^r)) \cong \mathbb{F}_{p^r} \times \mathbb{F}_{p^r}$. So $\text{ed}_k(\text{Syl}_p(\Omega^+(4, p^r))) = \text{ed}_k(\mathbb{F}_{p^r} \times \mathbb{F}_{p^r}) = 2\text{ed}_k(\mathbb{F}_{p^r}) = 2r$. This proves Theorem 6.1 for $n = 4$.

Similarly for $n = 4, p = 2$, the action of $\text{Up}_2(\mathbb{F}_{2^r}) \cong \mathbb{F}_{2^r}$ on $\text{Antisym}_0(2, 2^r) \cong \mathbb{F}_{2^r}$ is trivial and so $\text{Syl}_2(\Omega^+(2n, 2^r)) \cong \mathbb{F}_{2^r} \times \mathbb{F}_{2^r}$. So $\text{ed}_k(\text{Syl}_2(\Omega^+(4, 2^r))) = \text{ed}_k(\mathbb{F}_{2^r} \times \mathbb{F}_{2^r}) = 2\text{ed}_k(\mathbb{F}_{2^r}) = 2r$. This proves Theorem 6.1 for $n = 4$.

□

Remark 31. The work in the previous section is valid, though unnecessary, for $n = 4$. It gives us that the minimum dimension of an irrep is 1. And then since the center has rank $2r$, we will get an essential dimension of $2r$.

Remark 32. Alternate proof:

$$\begin{aligned}\Omega^+(4, p^r) &\cong (SL_2(\mathbb{F}_{p^r}) \times SL_2(\mathbb{F}_{p^r})) / \pm 1 \\ \Rightarrow \text{ed}_k(\Omega^+(4, p^r), p) &= \text{ed}_k(SL_2(\mathbb{F}_{p^r}), p) + \text{ed}_p(SL_2(\mathbb{F}_{p^r}), p) = 2r\end{aligned}$$

Case 2 ($n = 5$).

Step 1: Find the lower bound

Since $\{\chi_i|_{Z(\text{Syl}_p(\Omega(5, p^r)))}\}$ is a basis for $Z(\widehat{\text{Syl}_p(\Omega(5, p^r))})$. We can conclude that for at least r of the $\chi_i = \psi_{(b_i)}$, we have $b_1 \neq 0$ and so the dimension is at least p^r . The others might have $b_1 = 0$ and so could have dimension 1. Thus we must have that $\dim(\rho) \geq rp^r + r$. So

$$\text{ed}_k(\text{Syl}_p(\Omega(5, p^r))) \geq rp^r + r.$$

Step 2: Construct the upper bound

Let e_i be a basis for $\mathbb{F}_{p^r}^+$ over \mathbb{F}_p . Then $T = \{(e_i, 0), (0, e_i)\}$ forms a basis for $(\mathbb{F}_{p^r}^+)^2$ over \mathbb{F}_p . Then by Lemma 7, $\{\psi_{(e_i, 0)}, \psi_{(0, e_i)}\}$ forms a basis for the characters of $(\mathbb{F}_{p^r}^+)^2 = \Omega_1((\mathbb{F}_{p^r})^2)$. Let

$$S = \{(e_i, 0, 0), (0, 0, e_i)\}.$$

Then for $s \in S$, $\theta_{s, \lambda}$ have central character ψ_t where $t = (s_1, s_3)$. By Corollary 6.33, for λ trivial,

$$\dim(\theta_{s, \lambda}) = \begin{cases} p^r, & \text{if } s = (e_i, 0, 0) \\ 1, & \text{if } s = (0, 0, e_i) \end{cases}.$$

Let $\rho = \bigoplus_{s \in S} \theta_{s, \lambda}$. Then

$$\dim(\rho) = \sum_{s \in S} \dim(\theta_{s, \lambda}) = rp^r + r$$

And by Lemma 3.3, since $\{\psi_{(e_i, 0)}, \psi_{(0, e_i)}\}$ forms a basis for the characters of $(\mathbb{F}_{p^r}^+)^2 = \Omega_1((\mathbb{F}_{p^r})^2)$, we have that $\rho = \bigoplus_{s \in S} \theta_{s, \lambda}$ is a faithful representation of $\text{Syl}_p(\Omega(5, p^r))$. Thus ρ is the desired faithful representation of $\text{Syl}_p(\Omega(5, p^r))$ of dimension $rp^r + r$.

Case 3 ($n = 2m, m > 2$).

Step 1: Find the lower bound

The minimum dimension of an irreducible representation of $\text{Syl}_p(\Omega^+(2n, p^r))$ with non-trivial central character is given by $\theta_{s, \lambda}$ with λ trivial and $s = (b, 0, \dots, 0)$. So the minimum dimension is $p^{2r(m-2)}$. Thus the minimum dimension of a faithful representation $rp^{2r(m-2)}$ since the center has rank r .

Step 2: Construct the upper bound

Let $\{e_i\}$ be a basis for $\mathbb{F}_{p^r}^+$ over \mathbb{F}_p . Then by Lemma 7, $\{\psi_{e_i}\}$ form a basis for the characters of $\mathbb{F}_{p^r}^+$. Let $s_i = (e_i, 0, \dots, 0)$. Then by Corollary 6.30, for λ trivial,

$$\dim(\theta_{s_i, \lambda}) = p^{2r(n-2)}.$$

Let $\rho = \bigoplus_i \theta_{s_i, \lambda}$. Then

$$\dim(\rho) = \sum_{i=1}^r \dim(\theta_{s_i, \lambda}) = \sum_{i=1}^r p^{2r(m-2)} = rp^{2r(m-2)}.$$

And by Lemma 3.3, since $\psi_{e_i}|_{\mathbb{F}_{p^r}}$ form a basis for $\widehat{\mathbb{F}_{p^r}}$, we have that $\bigoplus_i \theta_{s_i, \lambda}$ is a faithful representation of $\text{Syl}_p(\Omega^+(2m, p^r))$. Thus ρ is the desired faithful representation of $\text{Syl}_p(\Omega^+(2m, p^r))$ of dimension $rp^{2r(m-2)}$.

Case 4 ($n = 2m + 1, m > 2$).

Step 1: Find the lower bound

Since $\{\chi_i|_{Z(\text{Syl}_p(\Omega(2m+1, p^r)))}\}$ a basis for $Z(\widehat{\text{Syl}_p(\Omega(2m+1, p^r))})$, we must have $b_1 \neq 0$ for at least r of the ρ_i . So the minimum dimension is $p^{r(m-1)(m-2)}$. The other r may have $b_1 = 0$ but then we must have $a_1 \neq 0$, so the minimum dimension is $p^{r(m-1)}$. Thus we have

$$\text{ed}_k(\text{Syl}_p(\Omega(2m+1, p^r))) \geq rp^{r(m-1)(m-2)} + rp^{r(m-1)} = rp^{r(m-1)}(p^{r(m-2)} + 1).$$

Step 2: Construct the upper bound

Let $\{e_i\}$ be a basis for $\mathbb{F}_{p^r}^+$ over \mathbb{F}_p . Then by Lemma 7, $\{\psi_{e_i}\}$ form a basis for the characters of $\mathbb{F}_{p^r}^+$. Let

$$S = \{(e_i, 0, \dots, 0), (0, \dots, 0, e_i, 0, \dots, 0)\}.$$

For $s \in S$, $\{\psi_s|_{Z(\widehat{\text{Syl}_p(\Omega(2m+1, p^r))})}\}$ forms a basis for $Z(\widehat{\text{Syl}_p(\Omega(2m+1, p^r))})$. Let

$$\rho = \bigoplus_{s \in S} \theta_{s, \lambda}.$$

Then by Lemma 3.3, $\rho = \bigoplus_{s \in S} \theta_{s, \lambda}$ is a faithful representation of $\text{Syl}_p(\Omega(2m+1, p^r))$. By Corollaries 6.36 and 6.38, for $s \in S$, λ trivial,

$$\dim(\theta_{s, \lambda}) = \begin{cases} p^{r(m-1)}, & s = (e_i, 0, \dots, 0) \\ p^{r(m-1)(m-2)}, & s = (0, \dots, 0, e_i, 0, \dots, 0) \end{cases}.$$

Thus

$$\dim(\rho) = \sum_{s \in S} \dim(\theta_{s, \lambda}) = rp^{r(m-1)} + rp^{r(m-1)(m-2)} = rp^{r(m-1)}(p^{r(m-2)} + 1).$$

Thus ρ is the faithful representation of $\text{Syl}_p(\Omega(2m+1, p^r))$ of desired dimension. \square

7 The Unitary Groups

In this section, we will show the following theorem:

Theorem 7.1. *For any p, k a field such that $\text{char } k \neq p$,*

$$\text{ed}_k(\text{PSU}(n, p^{2r}), p) = \text{ed}_k(\text{U}(n, p^{2r}), p) = \begin{cases} rp^{2r(m-1)} & n = 2m \\ 3rp^{2r(m-1)} & n = 2m + 1 \end{cases}$$

7.1 Definitions

For any p , by the Artin-Schreier Theorem ([13], Theorem 6.4) we can write $\mathbb{F}_{p^{2r}} = \mathbb{F}_{p^r}[\alpha]$ where α is a root of $x^2 - x - \eta$. Then $\mathbb{F}_{p^{2r}} = \mathbb{F}_{p^r} + \alpha\mathbb{F}_{p^r}$.

Definition 7.2. For $p \neq 2$, we can change $x^2 - x - \eta$ by a linear change of variables to the form $x^2 + \eta'$. Let α be a root of $x^2 + \eta'$. Then we still have $\mathbb{F}_{p^{2r}} = \mathbb{F}_{p^r} + \alpha\mathbb{F}_{p^r}$. We will write elements in $\mathbb{F}_{p^{2r}}$ as $a = \dot{a} + \ddot{a}\alpha$, where $\dot{a}, \ddot{a} \in \mathbb{F}_{p^r}$. For $a = \dot{a} + \ddot{a}\alpha$, define $\bar{a} = \dot{a} - \ddot{a}\alpha$.

Definition 7.3. For $p = 2$, let α be a root of $x^2 + x + \eta$. Note that $(\alpha + 1)^2 + (\alpha + 1) + \eta = \alpha^2 + \alpha + \eta = 0$. So $\alpha + 1$ gives the other root of $x^2 + x + \eta$. We will write elements in $\mathbb{F}_{p^{2r}}$ as $a = \dot{a} + \ddot{a}\alpha$, where $\dot{a}, \ddot{a} \in \mathbb{F}_{p^r}$. For $a = \dot{a} + \ddot{a}\alpha$, define $\bar{a} = \dot{a} + \ddot{a}(\alpha + 1) = (\dot{a} + \ddot{a}) + \ddot{a}\alpha$.

Remark 33. For any p , we have that for all $a \in \mathbb{F}_{p^{2r}}$, $\bar{\bar{a}} = a$.

Definition 7.4. For $\mathbf{x} = (x_i) \in (\mathbb{F}_{p^{2r}})^n$, let $\bar{\mathbf{x}} = (\bar{x}_i)$.

Definition 7.5. For $M = (M_{ij}) \in GL(n, \mathbb{F}_{p^{2r}})$, let $\overline{M} = (\overline{M_{ij}})$.

Definition 7.6. A matrix β is called **Hermitian** if $\beta^T = \overline{\beta}$.

Definition 7.7. Given a Hermitian matrix β , define

$$U_\beta(n, p^{2r}) = \{M \in GL(n, \mathbb{F}_{p^{2r}}) : M^T \beta \overline{M} = \beta\}.$$

By Grove ([10], Corollary 10.4), for any p, r, n, β, β' , $U_\beta(n, p^{2r}) \cong U_{\beta'}(n, p^{2r})$, so we may work with whichever choice of β we like. So for the remainder of this paper, we will drop the β in our notation.

Definition 7.8. Define $SU(n, p^{2r})$ as $SU(n, p^{2r}) = \{M \in U(n, p^{2r}) : \det(M) = 1\}$.

Definition 7.9. Define $PSU(n, p^{2r})$ as $PSU(n, p^{2r}) = SU(n, p^{2r}) / (Z(SU(n, p^{2r})))$.

Definition 7.10. Just as for the orthogonal groups the Witt index is defined to be the dimension of a maximal totally isotropic subspace of \mathbb{F}_{p^r} . Let w denote the Witt index.

Remark 34. Grove shows ([10], Theorem 11.26) that for $n \geq 2$ and Witt index $w \geq 1$, $PSU(n, p^r)$ is simple except for $(n, p^r) = (2, 4), (2, 9)$, or $(3, 4)$. Grove also shows ([10], Proposition 11.17) that

$$Z(SU(n, p^{2r})) = \{a \text{Id}_n : a \in \mathbb{F}_{p^{2r}}, a\overline{a} = 1, \text{ and } a^n = 1\}.$$

7.2 The p -Sylows

By Grove ([10], Theorem 11.28 and Cor 11.29), $|\text{Syl}_p(U(n, p^{2r}))| = p^{rn(n-1)/2} = |\text{Syl}_p(PSU(n, p^{2r}))|$. So $\text{Syl}_p(U(n, p^r))$ maps injectively onto $\text{Syl}_p(PSU(n, p^r))$; thus their p -Sylows must be isomorphic, and hence they have the same essential p -dimension.

7.2.1 $n = 2m$

If $n = 2m$, then we can choose

$$\beta = \begin{pmatrix} 0_m & \text{Id}_m \\ \text{Id}_m & 0_m \end{pmatrix}.$$

Remark 35.

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \in U_\beta(n, p^{2r}) \Leftrightarrow A^T \overline{C} + C^T \overline{A} = 0 = B^T \overline{D} + D^T \overline{B} \text{ and } A^T \overline{D} + C^T \overline{B} = \text{Id}_m = B^T \overline{C} + D^T \overline{A}.$$

Definition 7.11. For any p , define

$$G_{p^r} = \{a \in \mathbb{F}_{p^{2r}} : \overline{a} = -a\}.$$

Remark 36.

$$G_{p^r} = \begin{cases} \{\dot{a} + \ddot{a}\alpha : \dot{a} = 0\} = \alpha \mathbb{F}_{p^r} \cong (\mathbb{Z}/p\mathbb{Z})^r & p \neq 2 \\ \{\dot{a} + \ddot{a}\alpha : \ddot{a} = 0\} = \mathbb{F}_{p^r} \cong (\mathbb{Z}/p\mathbb{Z})^r & p = 2 \end{cases}$$

Definition 7.12. Let $\text{antisym}^*(m, p^{2r}) = \{B \in GL(n, p^{2r}) : B^T = -\overline{B}\}$

To construct a matrix in $\text{antisym}^*(m, p^{2r})$, we can choose the entries in the upper right to be anything, and this will dictate the entries in the lower right. On the diagonal, we must have $B_{i,i} = -\overline{B_{i,i}}$. So

$$|\text{antisym}^*(m, p^{2r})| = |\text{Up}_m(\mathbb{F}_{p^{2r}})| \cdot |G_{p^r}|^m = (p^{2r})^{\frac{m(m-1)}{2}} \cdot (p^r)^m = p^{rm(m-1)} \cdot p^{rm} = p^{rm^2 - rm + rm} = p^{rm^2}$$

Proposition 7.13. [See [21]] For $n = 2m$, any p ,

$$\text{Syl}_p(U(n, p^{2r})) = \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (\overline{A^{-1}})^T \end{pmatrix} \begin{pmatrix} \text{Id}_m & B \\ 0_m & \text{Id}_m \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(m, p^{2r}) \right\}.$$

Proof. Let

$$O = \left\{ \begin{pmatrix} A & 0_m \\ 0_m & (\overline{A^{-1}})^T \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^{2r}}) \right\}.$$

This is a subgroup of $U(2m, p^{2r})$ isomorphic to $\text{Up}_m(\mathbb{F}_{p^{2r}})$. So $|O| = |\text{Up}_m(\mathbb{F}_{p^{2r}})| = p^{rm(m-1)}$. Let

$$N = \left\{ \begin{pmatrix} \text{Id}_n & B \\ 0_n & \text{Id}_n \end{pmatrix} : B \in \text{antisym}^*(m, p^{2r}) \right\}.$$

This is a subgroup of $U(2m, p^{2r})$ isomorphic to $\text{antisym}^*(n, p^{2r})$. So $|N| = |\text{antisym}^*(m, p^{2r})| = p^{rm^2}$. A direct computation shows that $N \cdot O = N \rtimes O$ and

$$|N \rtimes O| = p^{rn(n-1)/2} = |\text{Syl}_p(U(n, p^{2r}))|.$$

Thus

$$N \rtimes O = \left\{ \begin{pmatrix} A & 0_n \\ 0_n & (\overline{A^{-1}})^T \end{pmatrix} \begin{pmatrix} \text{Id}_n & B \\ 0_n & \text{Id}_n \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(n, p^{2r}) \right\}$$

is a p -Sylow of $U(2m, p^{2r})$. □

Corollary 7.14. For $n = 2m$, any p ,

$$\text{Syl}_p(U(n, p^{2r})) \cong \text{antisym}^*(m, p^{2r}) \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}}),$$

where the action is given by

$$A(B) = AB\overline{A}^T.$$

7.2.2 $n = 2m + 1$

If $n = 2m + 1$, then we can choose

$$\beta = \begin{pmatrix} -1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & 0_m & \text{Id}_m \\ \mathbf{0} & \text{Id}_m & 0_m \end{pmatrix}$$

since $\overline{-1} = -1$ when p is odd and $-1 = 1$ when p is even.

Remark 37.

$$M = \begin{pmatrix} a & \mathbf{w} & \mathbf{x} \\ \mathbf{y} & A & B \\ \mathbf{z} & C & D \end{pmatrix}$$

is in $U(2m + 1, p^{2r})$ if and only if

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U(2m, p^{2r})$$

and

$$a\overline{a} - \mathbf{y} \cdot \overline{\mathbf{z}} - \mathbf{z} \cdot \overline{\mathbf{y}} = 1$$

and

$$0 = -a\overline{\mathbf{w}} + \mathbf{y}\overline{C} + \mathbf{z}\overline{A} = -a\overline{\mathbf{x}} + \mathbf{y}\overline{D} + \mathbf{z}\overline{B} = -a\overline{\mathbf{w}} + A^T\overline{\mathbf{z}} + C^T\overline{\mathbf{y}} = -a\overline{\mathbf{x}} + B^T\overline{\mathbf{z}} + D^T\overline{\mathbf{y}}$$

If $A = D = \text{Id}_m$ and $B = C = 0_m$, $a = 1$, this becomes

$$\mathbf{y} \cdot \overline{\mathbf{z}} = -\mathbf{z} \cdot \overline{\mathbf{y}}, \quad 0 = -\overline{\mathbf{w}} + \mathbf{z} = -\overline{\mathbf{x}} + \mathbf{y}.$$

Note this is satisfied when $\mathbf{z} = \mathbf{w} = \mathbf{0}$, $\mathbf{x} \in \mathbb{F}_{p^{2r}}^m$, $\mathbf{y} = \overline{\mathbf{x}}$.

Lemma 7.15. For $n = 2m + 1$, any p ,

$$\begin{aligned} & \text{Syl}_p(U(n, p^{2r})) \\ &= \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \bar{\mathbf{x}}^T & \text{Id}_n & 0_n \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A & 0_n \\ \mathbf{0} & 0_n & (A^{-1})^T \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \text{Id}_n & B \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^{2r}}^m, A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(m, p^{2r}) \right\}. \end{aligned}$$

Proof. Let

$$P = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \bar{\mathbf{x}}^T & \text{Id}_n & 0_n \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} : \mathbf{y} \in \mathbb{F}_{p^{2r}}^m \right\}.$$

This is a subgroup of $U(2n + 1, p^{2r})$ isomorphic to $(\mathbb{F}_{p^{2r}}^+)^m$. So $|P| = |(\mathbb{F}_{p^{2r}}^+)^m| = p^{2rm}$. Let

$$N = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \text{Id}_n & B \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} : B \in \text{antisym}^*(m, p^{2r}) \right\}.$$

This is a subgroup of $U(2n + 1, p^{2r})$ isomorphic to $\text{antisym}^*(m, p^{2r})$. So $|N| = |\text{antisym}^*(m, p^{2r})| = p^{rm^2}$. Let

$$O = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A & 0_n \\ \mathbf{0} & 0_n & (A^{-1})^T \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^{2r}}) \right\}.$$

This is a subgroup of $U(2n + 1, p^{2r})$ isomorphic to $\text{Up}_m(\mathbb{F}_{p^{2r}})$. So $|O| = |\text{Up}_m(\mathbb{F}_{p^{2r}})| = p^{rm(m-1)}$. A direct computation shows that $P \cdot N \cdot O = (P \times N) \rtimes O$ and

$$|(P \times N) \rtimes O| = p^{rn(n-1)/2} = |\text{Syl}_p(U(n, p^{2r}))|$$

Thus

$$\begin{aligned} & (P \times N) \rtimes O \\ &= \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \bar{\mathbf{x}}^T & \text{Id}_n & 0_n \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A & 0_n \\ \mathbf{0} & 0_n & (A^{-1})^T \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \text{Id}_n & B \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^{2r}}^m, A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(m, p^{2r}) \right\}. \end{aligned}$$

is a p -Sylow of $U(2m + 1, p^{2r})$. □

Corollary 7.16. For $n = 2m + 1$, any p ,

$$\text{Syl}_p(U(n, p^{2r})) \cong \left((\mathbb{F}_{p^{2r}}^+)^m \times \text{antisym}^*(m, p^{2r}) \right) \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}}),$$

where the action of $\text{Up}_m(\mathbb{F}_{p^{2r}})$ on $\text{antisym}^*(m, p^{2r})$ is given by

$$A(B) = AB\bar{A}^T.$$

and the action of $\text{Up}_m(\mathbb{F}_{p^{2r}})$ on $(\mathbb{F}_{p^{2r}}^+)^m$ is given by

$$A(\mathbf{x}) = \mathbf{x}\bar{A}^T.$$

7.3 Calculating the center

7.3.1 $n = 2m$

Lemma 7.17. For $n = 2m$, any p ,

$$Z(\text{Syl}_p(U(n, p^{2r}))) = \left\{ \begin{pmatrix} \text{Id}_m & D \\ 0_n & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix}, y = -\bar{y} \right\} \cong G_{p^r} \cong (\mathbb{Z}/p\mathbb{Z})^r$$

For the proof of Lemma 7.17, we need the following lemma:

Lemma 7.18. Given $D \in \text{antisym}^*(m, p^{2r})$,

$$AD = D(\overline{A^{-1}})^T \forall A \in \text{Up}_m(\mathbb{F}_{p^{2r}}) \Leftrightarrow D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{m-1} \end{pmatrix}$$

Granting this lemma, we can prove Lemma 7.17:

*Proof of Lemma 7.17.*¹

$$\begin{aligned} \text{Syl}_p(U(2m, p^{2r})) &= \left\{ \begin{pmatrix} A & 0_n \\ 0_n & (\overline{A^{-1}})^T \end{pmatrix} \begin{pmatrix} \text{Id}_n & B \\ 0_n & \text{Id}_n \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(m, p^{2r}) \right\} \\ &= \left\{ \begin{pmatrix} A & AB \\ 0_n & (\overline{A^{-1}})^T \end{pmatrix} : A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(m, p^{2r}) \right\}. \end{aligned}$$

Note that

$$\begin{aligned} &\begin{pmatrix} A & AB \\ 0_n & (\overline{A^{-1}})^T \end{pmatrix}^{-1} \begin{pmatrix} C & CD \\ 0_n & (\overline{C^{-1}})^T \end{pmatrix} \begin{pmatrix} A & AB \\ 0_n & (\overline{A^{-1}})^T \end{pmatrix} \\ &= \begin{pmatrix} A^{-1}CA & A^{-1}CAB + A^{-1}CD(\overline{A^{-1}})^T - B((\overline{A^{-1}CA})^{-1})^T \\ 0_n & ((\overline{A^{-1}CA})^{-1})^T \end{pmatrix} \end{aligned}$$

So $\begin{pmatrix} C & CD \\ 0_n & (\overline{C^{-1}})^T \end{pmatrix} \in Z(\text{Syl}_p(U(2m, p^{2r})))$ if and only if

$$C \in Z(\text{Up}_m(\mathbb{F}_{p^{2r}})) = \left\{ \begin{pmatrix} 1 & 0 & x \\ \mathbf{0} & \text{Id}_{n-2} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \right\}$$

and

$$CD = CB + CA^{-1}D(\overline{A^{-1}})^T - B(\overline{C^{-1}})^T$$

for all $A \in \text{Up}_m(\mathbb{F}_{p^{2r}}), B \in \text{antisym}^*(m, p^{2r})$. Choosing $A = \text{Id}_n$, we need $CD = CB + CD - B(C^{-1})^T$. So we must have $CB = B(\overline{C^{-1}})^T$ for all $B \in \text{antisym}^*(m, p^{2r})$. Write

$$C = \begin{pmatrix} 1 & \mathbf{0} & x \\ \mathbf{0} & \text{Id}_{n-2} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \in Z(\text{Up}_m(\mathbb{F}_{p^{2r}})) = Z(\text{Syl}_p(\text{PSL}(m, p^{2r}))).$$

(We calculated this center in the section on $\text{PSL}(n, p^r)$.) Then

$$(\overline{C^{-1}})^T = \begin{pmatrix} 1 & \mathbf{0} & -\bar{x} \\ \mathbf{0} & \text{Id}_{n-2} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix}^T = \begin{pmatrix} 1 & \mathbf{0} & 0 \\ \mathbf{0} & \text{Id}_{n-2} & \mathbf{0} \\ -\bar{x} & \mathbf{0} & 1 \end{pmatrix}.$$

¹When $p = 2$, the negatives will go away, but the arguments will still hold.

Then for $B = (b_{i,j}) \in \text{antisym}^*(m, p^{2r})$,

$$CB = \begin{pmatrix} b_{1,1} - x\overline{b_{1,n}} & b_{1,2} - x\overline{b_{2,n}} & \cdots & b_{1,n-1} - x\overline{b_{n-1,n}} & b_{1,n} + xb_{n,n} \\ -\overline{b_{1,2}} & b_{2,2} & b_{2,3} & \cdots & b_{2,n} \\ \vdots & & \ddots & & \vdots \\ -\overline{b_{1,n-1}} & & \cdots & & b_{n-1,n} \\ -\overline{b_{1,n}} & & \cdots & -\overline{b_{n-1,n}} & b_{n,n} \end{pmatrix}$$

and

$$B(\overline{C^{-1}})^T = \begin{pmatrix} b_{1,1} - \overline{xb_{1,n}} & b_{1,2} & \cdots & b_{1,n} \\ -\overline{b_{1,2}} - \overline{xb_{2,n}} & b_{2,2} & b_{2,3} & \cdots & b_{2,n} \\ \vdots & & \ddots & & \vdots \\ -\overline{b_{1,n-1}} - \overline{xb_{n-1,n}} & -\overline{b_{2,n-1}} & \cdots & \cdots & b_{n-1,n} \\ -\overline{b_{1,n}} - \overline{xb_{n,n}} & -\overline{b_{2,n}} & \cdots & -\overline{b_{n-1,n}} & b_{n,n} \end{pmatrix}$$

Remark 38. I do not need $m > 2$ here like I did for the orthogonal groups.

So we must have $x = 0$, and hence $C = \text{Id}_n$. So the other requirement above becomes

$$D = A^{-1}D(\overline{A^{-1}})^T \Leftrightarrow AD = D(\overline{A^{-1}})^T$$

for all $A \in \text{Up}_m(\mathbb{F}_{p^{2r}})$. Then by Lemma 7.18, we get that

$$Z(\text{Syl}_p(U(2m, p^{2r}))) = \left\{ \begin{pmatrix} \text{Id}_m & D \\ 0_n & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix} \right\}$$

□

Proof of Lemma 7.18. \Leftarrow : This is a straightforward calculation.

\Rightarrow : We will prove this by induction.

Base Case: When $n = 2$, we can write $A = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ and $D = \begin{pmatrix} y & x \\ -\overline{x} & z \end{pmatrix}$.

$$AD = \begin{pmatrix} y - a\overline{x} & x + az \\ -\overline{x} & z \end{pmatrix},$$

and

$$D(\overline{A^{-1}})^T = \begin{pmatrix} y - \overline{ax} & x \\ -\overline{x} - \overline{az} & z \end{pmatrix}.$$

So in order for these to be equal for all A , we must have $z = 0$ and $x = 0$.

Induction Step: The induction step is basically identical to that in the proof of Lemma 5.8. □

7.3.2 $n = 2m + 1$

Lemma 7.19. [See [19]] For $n = 2m + 1$, any p ,

$$\begin{aligned} Z(\text{Syl}_p(U(n, p^{2r}))) &= \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_n & D \\ \mathbf{0} & 0_n & \text{Id}_n \end{pmatrix} : \mathbf{x} = (x_1, 0, \dots, 0) : D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix}, y = -\overline{y} \right\} \\ &\cong \mathbb{F}_{p^{2r}} \times G_{p^r} \cong (\mathbb{Z}/p\mathbb{Z})^{3r} \end{aligned}$$

Proof. Since $\text{Syl}_p(U(2m + 1, p^{2r})) \cong \left((\mathbb{F}_{p^{2r}}^+)^m \times (\text{antisym}^*(m, p^{2r})) \times \text{Up}_m(\mathbb{F}_{p^{2r}}) \right) \cong (\mathbb{F}_{p^{2r}}^+)^n \times \text{Syl}_p(U(2m, p^{2r}))$,

$$Z(\text{Syl}_p(U(2m + 1, p^{2r}))) \cap \{\mathbf{0}\} \times \text{Syl}_p(U(2m, p^{2r}))$$

must be a subset of $Z(\text{Syl}_p(U(2m, p^{2r})))$, which we proved above to be

$$Z(\text{Syl}_p(U(2m, p^{2r}))) = \left\{ \begin{pmatrix} \text{Id}_m & D \\ \mathbf{0} & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{m-1} \end{pmatrix}, y = -\bar{y} \right\}$$

Thus the center of $Z(\text{Syl}_p(U(2m+1, p^r)))$ is a subset of

$$(\mathbb{F}_{p^{2r}}^+)^m \rtimes \left\{ \begin{pmatrix} \text{Id}_m & D \\ \mathbf{0} & \text{Id}_m \end{pmatrix} : D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{m-1} \end{pmatrix} \right\} = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & \mathbf{0}_m & \text{Id}_m \end{pmatrix} : \mathbf{x} \in \mathbb{F}_{p^{2r}}^m, D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{m-1} \end{pmatrix} \right\}$$

Given $\begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} \\ \mathbf{y}^T & A & AB \\ \mathbf{0} & 0_n & (\overline{A^{-1}})^T \end{pmatrix} \in \text{Syl}_p(U(2m+1, p^{2r}))$, we have that

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} \\ \mathbf{y}^T & A & AB \\ \mathbf{0} & 0_m & (\overline{A^{-1}})^T \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} + \mathbf{x}(\overline{A^{-1}})^T \\ \mathbf{x}^T + \mathbf{y}^T & A & \mathbf{x}^T \mathbf{y} + AB + D(\overline{A^{-1}})^T \\ \mathbf{0} & 0_m & (\overline{A^{-1}})^T \end{pmatrix}$$

and

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{y} \\ \mathbf{y}^T & A & AB \\ \mathbf{0} & 0_m & (\overline{A^{-1}})^T \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} + \mathbf{y} \\ \mathbf{y}^T + A\mathbf{x}^T & A & \mathbf{x}\mathbf{y}^T + AD + AB \\ \mathbf{0} & 0_m & (\overline{A^{-1}})^T \end{pmatrix}$$

So in order for

$$\begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_n & \text{Id}_m \end{pmatrix}$$

to be in the center, we need $\mathbf{x}^T = A\mathbf{x}^T$, $\mathbf{x} = \mathbf{x}(\overline{A^{-1}})^T$, and $AD = D(\overline{A^{-1}})^T$ for all choices of A . By Lemma 7.18, $AD = D(\overline{A^{-1}})^T$ is satisfied if and only if

$$D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{n-1} \end{pmatrix}.$$

Note that the k th entry of $A\mathbf{x}^T$ is given by $x_k + \sum_{i=k+1}^n x_i a_{k,i}$. In order for this to be equal to x_k (the k th entry of \mathbf{x}^T) for all $a_{k,i}$, must have $x_i = 0$ for all $i > 1$. So $\mathbf{x} = (x_1, 0, \dots, 0)$. In this case $\mathbf{x} = \mathbf{x}(\overline{A^{-1}})^T$ will be satisfied as well. Therefore the center is

$$Z(\text{Syl}_p(U(2m+1, p^{2r}))) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & \mathbf{x} \\ \mathbf{x}^T & \text{Id}_m & D \\ \mathbf{0} & 0_m & \text{Id}_m \end{pmatrix} : \mathbf{x} = (x_1, 0, \dots, 0) : D = \begin{pmatrix} y & \mathbf{0} \\ \mathbf{0} & 0_{m-1} \end{pmatrix} \right\}$$

□

7.4 Maximizing L_s

7.4.1 $n = 2m$

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element of

$$\widehat{\text{antisym}^*(m, p^{2r})} \cong (\mathbb{F}_{p^{2r}}^+)^{\widehat{m(m-1)/2}} \times \mathbb{F}_{p^r}^m \cong (\mathbb{F}_{p^r}^+)^{\widehat{m^2}}$$

is of the form

$$\mathbf{d} \mapsto \psi(\mathbf{bd}^T),$$

where $\mathbf{d}, \mathbf{b} \in (\mathbb{F}_{p^r}^+)^{m^2}$. Let the isomorphism $\mathbb{F}_{p^{2r}}^+ \rightarrow (\mathbb{F}_{p^r}^+)^2$ be defined as

$$\dot{b} + \ddot{b}\alpha \mapsto (\dot{b}, \ddot{b}).$$

Note that

$$\overline{(\dot{b}, \ddot{b})} = \begin{cases} (\dot{b}, -\ddot{b}) & p \neq 2 \\ (\dot{b} + \ddot{b}, \ddot{b}) & p = 2 \end{cases}.$$

And for $b = (\dot{b}, \ddot{b})$,

$$b + \bar{b} = \begin{cases} (2\dot{b}, 0) & p \neq 2 \\ (\dot{b}, 0) & p = 2 \end{cases}.$$

Note that for a matrix $(b_{i,j})$ in $\text{Antisym}(m, p^{2r})$, $b_{i,i} \in G_{p^r}$, so

$$b_{i,i} = \begin{cases} b_{i,i}\alpha & p \neq 2 \\ b_{i,i} & p = 2 \end{cases}.$$

Let the isomorphism $\text{antisym}^*(m, p^{2r}) \cong (\mathbb{F}_{p^r}^+)^{m^2}$ be given by

$$\begin{pmatrix} b_{1,1} & b_{1,2} & b_{1,3} & \cdots & b_{1,n} \\ -\overline{b_{1,2}} & b_{2,2} & b_{2,3} & \cdots & b_{2,n} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ -\overline{b_{1,n-1}} & -\overline{b_{2,n-1}} & \cdots & b_{n-1,n-1} & b_{n-1,n} \\ -\overline{b_{1,n}} & -\overline{b_{2,n}} & \cdots & -\overline{b_{n-1,n}} & b_{n,n} \end{pmatrix} \mapsto \begin{cases} (b_{1,1}, b_{1,2}, b_{1,2}, \cdots, b_{1,n}, b_{1,n}, b_{2,2}, b_{2,3}, b_{2,3}, \cdots, \cdots, b_{n,n}) & p \neq 2 \\ (b_{1,1}, b_{1,2}, b_{1,2}, \cdots, b_{1,n}, b_{1,n}, b_{2,2}, b_{2,3}, b_{2,3}, \cdots, \cdots, b_{n,n}) & p = 2 \end{cases}$$

Proposition 7.20. *For any p , the characters extending a non-trivial central character are $\psi_{(b_j)}$ with $b_1 \neq 0$.*

Proof. $\psi_{(b_j)}$ restricted to the center of the p -Sylow is ψ_{b_1} . This is non-trivial if and only if $b_1 \neq 0$. \square

We will use Wigner-Mackey Theory with $G = \text{Syl}_p(U(2m, p^{2r})) = \text{antisym}^*(m, p^{2r}) \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}})$ to see what is the minimum dimension of an irreducible representation of G with non-trivial central character. So we have $\Delta = \text{antisym}^*(m, p^{2r}) \rtimes \{\text{Id}_m\}$, $L = \{0_m\} \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}})$. Recall that $L_s = \text{Stab}_L(\psi_s)$.

Remark 39. We will view $\psi_{(b_j)}$ as a map on $\Delta \cong \text{antisym}^*(m, p^{2r}) \cong (\mathbb{F}_{p^r}^+)^{m^2}$. So $\psi_{(b_j)}(D, \text{Id}) = \psi_{(b_j)}(D) = \psi(\mathbf{b} \cdot \mathbf{d})$, where $\mathbf{b} = (b_j)$ and \mathbf{d} is the vector corresponding to the matrix D .

The action of $g \in \text{Syl}_p(U(2m, p^r))$ on $\hat{\Delta}$ is given by

$${}^g\psi(D, \text{Id}_m) = \psi(g^{-1}(D, \text{Id}_m)g).$$

So for $h = (0_n, H^{-1}) \in L = \{0_m\} \times \text{Up}_m(\mathbb{F}_{p^{2r}})$, the action on $\psi_{(b_j)}$ is given by

$${}^h\psi_{(b_j)}(D, \text{Id}_m) = \psi_{(b_j)}((0_n, H)(D, \text{Id}_m)(0_n, H^{-1})).$$

So $(0_n, H^{-1}) \in L_{(b_j)}$ if and only if for all $\mathbf{d} \in ((\mathbb{Z}/p\mathbb{Z})^r)^{m^2} = D \in \text{antisym}^*(m, p^{2r})$,

$$\psi_{(b_j)}((0_n, H)(D, \text{Id}_n)(0_n, H^{-1})) = \psi_{(b_j)}(D, \text{Id}_n)$$

Let \mathbf{hdh}^T be the vector corresponding to $H D H^T$. Then since

$$\psi_{(b_j)}((0_m, H)(D, \text{Id}_m)(0_m, H^{-1})) = \psi(\mathbf{b} \cdot \mathbf{hdh}^T),$$

and

$$\psi_{(b_j)}(D, \text{Id}_m) = \psi(\mathbf{b} \cdot \mathbf{d}).$$

we get that $(0_m, H^{-1}) \in L_s$ if and only if

$$\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1$$

for all $\mathbf{d} \in (\mathbb{F}_{p^r})^{m^2}$ corresponding to $D \in \text{antisym}^*(m, p^{2r})$.

Write

$$H = \begin{pmatrix} 1 & h_{1,2} & h_{1,3} & \cdots & h_{1,m} \\ 0 & 1 & h_{2,3} & \cdots & h_{2,m} \\ & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & h_{m-1,m} \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}, \quad D = \begin{pmatrix} \overline{d_{1,1}} & d_{1,2} & d_{1,3} & \cdots & d_{1,m} \\ -\overline{d_{1,2}} & d_{2,2} & d_{2,3} & \cdots & d_{2,m} \\ \vdots & & \ddots & & \vdots \\ -\overline{d_{1,m-1}} & -\overline{d_{2,m-1}} & \cdots & d_{m-1,m-1} & d_{m-1,m} \\ -\overline{d_{1,m}} & -\overline{d_{2,m}} & \cdots & -\overline{d_{m-1,m}} & d_{m,m} \end{pmatrix}.$$

Proposition 7.21. For any p , $s = (b_j)$, with $b_1 \neq 0$, $\max_s |L_s| = |Up_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}$, and this maximum is achieved when $s = (b_j) = (b_1, 0, \dots, 0)$ with $b_1 \neq 0$.

Corollary 7.22. For any p , for $s = (b_1, 0, 0, \dots)$ with $b_1 \neq 0$, λ trivial,

$$\dim(\theta_{s,\lambda}) = \frac{|L|}{|L_s|} = \frac{|Up_m(\mathbb{F}_{p^{2r}})|}{|Up_{m-1}(\mathbb{F}_{p^{2r}})|} = |\mathbb{F}_{p^{2r}}^{(m-1)}| = p^{2r(m-1)}.$$

Proof of Proposition 7.21.

Step 1: Proving that for any s , $|L_s| \leq p^{r(m-1)(m-2)}$.

For $j_0 > 2$, choose $d_{i,j} = 0$ except for $d_{1,j_0} = -\overline{d_{j_0,1}}$. Then we get that

$$HD\overline{H}^T - D = \begin{pmatrix} \overline{h_{1,j_0}d_{1,j_0}} - \overline{h_{1,j_0}d_{1,j_0}} & \overline{h_{2,j_0}d_{1,j_0}} & \cdots & \overline{h_{j_0-1,j_0}d_{1,j_0}} & 0 & \cdots & 0 \\ -h_{2,j_0}d_{1,j_0} & 0 & & & \cdots & \cdots & 0 \\ \vdots & & & \ddots & & & \vdots \\ -h_{j_0-1,j_0}d_{1,j_0} & 0 & & & \cdots & \cdots & 0 \\ 0 & & & & \cdots & \cdots & 0 \\ \vdots & & & \ddots & & & \vdots \\ 0 & & & & \cdots & \cdots & 0 \end{pmatrix}$$

Case 1: $p \neq 2$

Note that

$$\overline{(h_{1,j_0} + h_{1,j_0}\alpha)(d_{1,j_0} + d_{1,j_0}\alpha)} = (h_{1,j_0}d_{1,j_0} + h_{1,j_0}d_{1,j_0}\eta') + (h_{1,j_0}d_{1,j_0} - h_{1,j_0}d_{1,j_0})\alpha$$

So we have

$$\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = d_{1,j_0} \left(2h_{1,j_0}b_{1,1} + \sum_{i=2}^{j_0-1} (h_{i,j_0}b_{1,i} - h_{i,j_0}b_{1,i}) \right) + d_{1,j_0} \left(2h_{1,j_0}b_{1,1} + \sum_{i=2}^{j_0-1} (h_{i,j_0}\eta'b_{1,i} + h_{i,j_0}b_{1,i}) \right)$$

If $2h_{1,j_0}b_{1,1} + \sum_{i=2}^{j_0-1} (h_{i,j_0}b_{1,i} - h_{i,j_0}b_{1,i}) \neq 0$, then as we run through all the values for d_{1,j_0} with $d_{1,j_0} = 0$, we will get that $\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})$ runs through all the values of \mathbb{F}_{p^r} . And since ψ is non-trivial, this means that $\psi(\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}))$ cannot always equal 1. This is a contradiction. So we must have

$$2h_{1,j_0}b_{1,1} + \sum_{i=2}^{j_0-1} (h_{i,j_0}b_{1,i} - h_{i,j_0}b_{1,i}) = 0$$

for all choices of $j_0 > 1$. Similarly, by running through all values for $d_{1,j_0}^{\ddot{\cdot}}$ with $d_{1,j_0}^{\dot{\cdot}} = 0$, we see that we must have

$$2h_{1,j_0}^{\ddot{\cdot}} b_{1,1}^{\ddot{\cdot}} + \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta' b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0$$

for all choices of $j_0 > 1$. If $b_{1,1} = b_{1,1}^{\ddot{\cdot}} \alpha \neq 0$, then given h_{i,j_0} for $i > 1$, $h_{1,j_0}^{\dot{\cdot}}$ and $h_{1,j_0}^{\ddot{\cdot}}$ are dictated by the equations above:

$$h_{1,j_0}^{\dot{\cdot}} = \frac{-1}{2b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\dot{\cdot}} - h_{i,j_0}^{\ddot{\cdot}} b_{1,i}^{\ddot{\cdot}})$$

and

$$h_{1,j_0}^{\ddot{\cdot}} = \frac{-1}{2b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta' b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}).$$

Thus for $b_{1,1} \neq 0$, $p \neq 2$,

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed } \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}$$

This completes Step 1 for $p \neq 2$.

Case 2: $p = 2$

Note that

$$\overline{(h_{1,j_0}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} \alpha)} (d_{1,j_0}^{\dot{\cdot}} + d_{1,j_0}^{\ddot{\cdot}} \alpha) = (h_{1,j_0}^{\dot{\cdot}} d_{1,j_0}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} d_{1,j_0}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} d_{1,j_0}^{\ddot{\cdot}} \eta) + (h_{1,j_0}^{\dot{\cdot}} d_{1,j_0}^{\ddot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} d_{1,j_0}^{\dot{\cdot}}) \alpha$$

So we have

$$\begin{aligned} & \mathbf{b} \cdot (\mathbf{hd}\overline{\mathbf{h}}^{\mathbf{T}} - \mathbf{d}) \\ &= d_{1,j_0}^{\dot{\cdot}} \left(h_{1,j_0}^{\ddot{\cdot}} b_{1,1}^{\ddot{\cdot}} + \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\ddot{\cdot}} b_{1,i}^{\ddot{\cdot}}) \right) + d_{1,j_0}^{\ddot{\cdot}} \left(h_{1,j_0}^{\dot{\cdot}} b_{1,1}^{\ddot{\cdot}} + \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}) \right) \end{aligned}$$

By the same reasoning as before, we must have

$$h_{1,j_0}^{\ddot{\cdot}} b_{1,1}^{\ddot{\cdot}} + \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\ddot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0$$

and

$$h_{1,j_0}^{\dot{\cdot}} b_{1,1}^{\ddot{\cdot}} + \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0$$

for all choices of $j_0 > 1$. If $b_{1,1} = b_{1,1}^{\ddot{\cdot}} \alpha \neq 0$, then given h_{i,j_0} for $i > 1$, $h_{1,j_0}^{\dot{\cdot}}$ and $h_{1,j_0}^{\ddot{\cdot}}$ are dictated by the equations above:

$$h_{1,j_0}^{\dot{\cdot}} = \frac{1}{b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\ddot{\cdot}} b_{1,i}^{\ddot{\cdot}})$$

and

$$h_{1,j_0}^{\ddot{\cdot}} = \frac{1}{b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}).$$

Thus for $b_{1,1} \neq 0$,

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed } \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}$$

This completes Step 1 for $p = 2$.

Step 2: Exhibiting that the max is achieved when $\mathbf{s} = (\mathbf{b}_1, \mathbf{0}, \cdot, \mathbf{0})$ with $\mathbf{b}_1 \neq \mathbf{0}$.

Let $s = (b_1, 0, 0, \dots, 0)$. And let B be the corresponding matrix. By the calculation above, we must have that for $j_0 > 1$:

For $p \neq 2$,

$$h_{1,j_0}^{\dot{\cdot}} = \frac{1}{2b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\dot{\cdot}} - h_{i,j_0}^{\ddot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0$$

and

$$h_{1,j_0}^{\ddot{\cdot}} = \frac{1}{2b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta' b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0.$$

And for $p = 2$,

$$h_{1,j_0}^{\dot{\cdot}} = \frac{1}{b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{1,j_0}^{\ddot{\cdot}} b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\ddot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0$$

and

$$h_{1,j_0}^{\ddot{\cdot}} = \frac{1}{b_{1,1}^{\ddot{\cdot}}} \sum_{i=2}^{j_0-1} (h_{i,j_0}^{\ddot{\cdot}} \eta b_{1,i}^{\dot{\cdot}} + h_{i,j_0}^{\dot{\cdot}} b_{1,i}^{\ddot{\cdot}}) = 0.$$

So, for any p we have that for $j_0 > 0$, $h_{1,j_0}^{\dot{\cdot}} = 0 = h_{1,j_0}^{\ddot{\cdot}}$, and hence $h_{1,j_0} = 0$. So

$$\begin{aligned} \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) &= b_1 \left(\sum_{l=1}^n \overline{h_{1,l}} \left(\sum_{k=1}^l d_{k,l} h_{1,k} - \sum_{k=l+1}^n \overline{d_{l,k}} h_{1,k} \right) - d_{1,1} \right) && \text{since } b_i = 0, i > 1 \\ &= b_1 (d_{1,1} - d_{1,1}) && \text{since } h_{1,j} = 0, j > 1 \\ &= 0 \end{aligned}$$

Therefore,

$$L_s = \{H : H_{1,j} = 0 \forall j > 1\}.$$

□

7.4.2 $n = 2m + 1$

Fix ψ a non-trivial character of $\mathbb{F}_{p^r}^+$. By Lemma 3.8, every element of

$$(\mathbb{F}_{p^{2r}}^+)^m \times \widehat{\text{antisym}^*(m, p^{2r})} \cong (\mathbb{F}_{p^{2r}}^+)^m \times (\mathbb{F}_{p^{2r}}^+)^{m^2} \cong (\mathbb{F}_{p^r}^+)^{2m+m^2}$$

is of the form

$$\mathbf{d} \mapsto \psi(\mathbf{bd}^T),$$

where $\mathbf{d}, \mathbf{b} \in (\mathbb{F}_{p^r}^+)^{2m+m^2}$. Let the isomorphism $\text{antisym}^*(m, p^{2r}) \cong (\mathbb{F}_{p^r}^+)^{m^2}$ be as it was for $n = 2m$:

$$\left(\begin{array}{ccccc} \overline{b_{1,1}} & b_{1,2} & b_{1,3} & \cdots & b_{1,n} \\ -\overline{b_{1,2}} & b_{2,2} & b_{2,3} & \cdots & b_{2,n} \\ \vdots & & \ddots & & \vdots \\ -\overline{b_{1,n-1}} & -\overline{b_{2,n-1}} & \cdots & b_{n-1,n-1} & b_{n-1,n} \\ -\overline{b_{1,n}} & -\overline{b_{2,n}} & \cdots & -\overline{b_{n-1,n}} & b_{n,n} \end{array} \right) \mapsto \begin{cases} (b_{1,1}^{\ddot{\cdot}}, b_{1,2}^{\dot{\cdot}}, b_{1,2}^{\ddot{\cdot}}, \dots, b_{1,n}^{\dot{\cdot}}, b_{1,n}^{\ddot{\cdot}}, b_{2,2}^{\dot{\cdot}}, b_{2,3}^{\dot{\cdot}}, b_{2,3}^{\ddot{\cdot}}, \dots, \dots, b_{n,n}^{\ddot{\cdot}}) & p \neq 2 \\ (b_{1,1}, b_{1,2}, b_{1,2}, \dots, b_{1,n}, b_{1,n}, b_{2,2}, b_{2,3}, b_{2,3}, \dots, \dots, b_{n,n}) & p = 2 \end{cases}$$

Proposition 7.23. *For any p , the characters of $(\mathbb{F}_{p^r}^+)^{2m+m^2}$ extending a non-trivial central character of the p -Sylow are $\psi_{(a_j, b_j)}$ with $(\dot{a}_1, \ddot{a}_1, b_1) \neq (0, 0, 0)$.*

Proof. $\psi_{(a_j, b_j)}$ restricted to the center is $\psi_{\hat{a}_1, \hat{a}_1, b_1}$. This is non-trivial if and only if $(\hat{a}_1, \hat{a}_1, b_1) \neq (0, 0, 0)$. \square

We will use Wigner-Mackey Theory with $G = \text{Syl}_p(U(2m+1, p^r)) = ((\mathbb{F}_{p^{2r}}^+)^m \times \text{antisym}^*(m, p^{2r})) \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}})$ to see what is the minimum dimension of an irreducible representation of G with non-trivial central character. So we have $\Delta = ((\mathbb{F}_{p^{2r}}^+)^m \times \text{antisym}^*(m, p^{2r})) \rtimes \{\text{Id}_n\}$, $L = (\{\mathbf{0}\} \times \{0_m\}) \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}})$. Recall that $L_s = \text{Stab}_L(\psi_s)$.

Remark 40. We will view $\psi_{(a_j, b_j)}$ as a map on $\Delta \cong ((\mathbb{F}_{p^{2r}}^+)^m \times \text{antisym}^*(n, p^r)) \cong (\mathbb{Z}/p\mathbb{Z})^{2rm+rm^2}$. So $\psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}) = \psi_{(a_j, b_j)}(\mathbf{x}, D) = \psi((\mathbf{a}, \mathbf{b}) \cdot (\mathbf{d}, \mathbf{x}))$, where $(\mathbf{a}, \mathbf{b}) = (a_j, b_j)$ and (\mathbf{x}, \mathbf{d}) is the vector corresponding to (\mathbf{x}, D) .

The action of $g \in \text{Syl}_p(U(2m+1, p^{2r}))$ on $\hat{\Delta}$ is given by

$${}^g\psi(\mathbf{x}, D, \text{Id}_m) = \psi(g^{-1}(\mathbf{x}, D, \text{Id}_m)g).$$

So for $h = (\mathbf{0}, 0_m, H^{-1}) \in L = (\{\mathbf{0}\} \times \{0_m\}) \rtimes \text{Up}_m(\mathbb{F}_{p^{2r}})$, the action on $\psi_{(a_j, b_j)}$ is given by

$${}^h\psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}_m) = \psi_{(a_j, b_j)}((\mathbf{0}, 0_m, H)(\mathbf{x}, D, \text{Id}_m)(\mathbf{0}, 0_m, H^{-1})).$$

So $(\mathbf{0}, 0_m, H^{-1}) \in L_s$ if and only if for all $(\mathbf{x}, \mathbf{d}) \in (\mathbb{F}_{p^r}^+)^{2m+m^2} = (\mathbf{x}, D) \in (\mathbb{F}_{p^{2r}}^+)^m \times \text{antisym}^*(m, p^{2r})$,

$$\psi_{(a_j, b_j)}((\mathbf{0}, 0_n, H)(\mathbf{x}, D, \text{Id}_m)(\mathbf{0}, 0_n, H^{-1})) = \psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}_m)$$

Let \mathbf{hdh}^T be the vector corresponding to HDH^T . Then since

$$\psi_{(a_j, b_j)}((\mathbf{0}, 0_n, H)(\mathbf{x}, D, \text{Id}_m)(\mathbf{0}, 0_n, H^{-1})) = \psi((\mathbf{a}, \mathbf{b}) \cdot (\mathbf{x}H^T, \mathbf{hdh}^T))$$

and

$$\psi_{(a_j, b_j)}(\mathbf{x}, D, \text{Id}_n) = \psi((\mathbf{a}, \mathbf{b}) \cdot (\mathbf{x}, \mathbf{d}))$$

we get that $(\mathbf{0}, 0_n, H^{-1}) \in L_s$ if and only if

$$\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1$$

for all $(\mathbf{x}, \mathbf{d}) \in (\mathbb{F}_{p^r}^+)^{2m+m^2}$ corresponding to $(\mathbf{x}, D) \in (\mathbb{F}_{p^{2r}}^+)^m \times \text{antisym}^*(m, p^{2r})$.

Proposition 7.24. For any p , $s = (a_j, b_j)$, $\max_s |L_s| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}$. And this max is achieved when $s = (0, \dots, 0, b_1, 0, \dots, 0)$ and when $s = (a_1, 0, \dots, 0)$ with $\hat{a}_1 \neq 0, \hat{a}_1 = 0$ or $\hat{a}_1 = 0, \hat{a}_1 \neq 0$.

Corollary 7.25. For any p , λ trivial, if $s = (0, \dots, 0, b_1, 0, \dots, 0)$ or $s = (a_1, 0, \dots, 0)$ with $\hat{a}_1 \neq 0, \hat{a}_1 = 0$ or $\hat{a}_1 = 0, \hat{a}_1 \neq 0$, then

$$\dim(\theta_{s, \lambda}) = \frac{|L|}{|L_s|} = \frac{|\text{Up}_m(\mathbb{F}_{p^{2r}})|}{|\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})|} = |\mathbb{F}_{p^{2r}}^{(m-1)}| = p^{2r(m-1)}.$$

Proof of Proposition 7.24.

Step 1: Proving that for any s , $|L_s| \leq p^{r(m-1)(m-2)}$.

Case 1: $b_1 \neq 0$

If we take $\mathbf{x} = \mathbf{0}$, then $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d})) = 1$ reduces to the condition for $U(2m, p^{2r})$. So if $b_1 \neq 0$ we must have $L_s = L_{(a_j, b_j)}$ a subset of the $L_{(b_j)}$ calculated in the proof of Proposition 7.21. I.e.

$$L_s \subset \{H : H_{1,j} \text{ fixed } \forall j \neq 1\} = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Thus $|L_s| \leq p^{r(m-1)(m-2)}$.

Case 2: $\mathbf{a}_1 \neq \mathbf{0}$ and $\mathbf{p} \neq \mathbf{2}$

If we take $\mathbf{d} = \mathbf{0}$, then $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{h}\mathbf{d}\mathbf{h}^T - \mathbf{d})) = 1$ reduces to $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x})) = 1$. Note that

$$x_{j_0} h_{k,j_0} = \dot{x}_{j_0} \dot{h}_{k,j_0} - \ddot{x}_{j_0} \ddot{h}_{k,j_0} \eta' + (x_{j_0} \ddot{h}_{k,j_0} + \dot{x}_{j_0} \dot{h}_{k,j_0}) \alpha$$

So we have

$$\begin{aligned} \mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) &= \sum_{k=1}^{j_0-1} \dot{a}_k x_{j_0} \dot{h}_{k,j_0} + \ddot{a}_k x_{j_0} \ddot{h}_{k,j_0} \\ &= \sum_{k=1}^{j_0-1} \dot{a}_k (\dot{x}_{j_0} \dot{h}_{k,j_0} - \ddot{x}_{j_0} \ddot{h}_{k,j_0} \eta') + \ddot{a}_k (\dot{x}_{j_0} \ddot{h}_{k,j_0} + \dot{x}_{j_0} \dot{h}_{k,j_0}) \\ &= \dot{x}_{j_0} \left(\sum_{k=1}^{j_0-1} \dot{a}_k \dot{h}_{k,j_0} + \ddot{a}_k \ddot{h}_{k,j_0} \right) + \ddot{x}_{j_0} \left(\sum_{k=1}^{j_0-1} \ddot{a}_k \dot{h}_{k,j_0} - \dot{a}_k \ddot{h}_{k,j_0} \eta' \right) \end{aligned}$$

If $\sum_{k=1}^{j_0-1} \dot{a}_k \dot{h}_{k,j_0} + \ddot{a}_k \ddot{h}_{k,j_0} \neq 0$, then as we run through all the values for x_{j_0} with $\ddot{x}_{j_0} = 0$, we will get that $\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x})$ runs through all the values of \mathbb{F}_{p^r} . And since ψ is non-trivial, this means that $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}))$ cannot always equal 1. This is a contradiction. So we must have

$$0 = \sum_{k=1}^{j_0-1} \dot{a}_k \dot{h}_{k,j_0} + \ddot{a}_k \ddot{h}_{k,j_0} = \dot{a}_1 \dot{h}_{1,j_0} + \ddot{a}_1 \ddot{h}_{1,j_0} + \sum_{k=2}^{j_0-1} \dot{a}_k \dot{h}_{k,j_0} + \ddot{a}_k \ddot{h}_{k,j_0}$$

Similarly, we must have

$$0 = \sum_{k=1}^{j_0-1} \ddot{a}_k \dot{h}_{k,j_0} - \dot{a}_k \ddot{h}_{k,j_0} \eta' = -\dot{a}_1 \ddot{h}_{1,j_0} \eta' + \ddot{a}_1 \dot{h}_{1,j_0} + \sum_{k=2}^{j_0-1} \ddot{a}_k \dot{h}_{k,j_0} - \dot{a}_k \ddot{h}_{k,j_0} \eta'$$

Subcase 2a: $\dot{\mathbf{a}}_1 \neq \mathbf{0}, \ddot{\mathbf{a}}_1 = \mathbf{0}$ or $\dot{\mathbf{a}}_1 = \mathbf{0}, \ddot{\mathbf{a}}_1 \neq \mathbf{0}$

If $\dot{a}_1 \neq 0, \ddot{a}_1 = 0$, then given h_{k,j_0} for $k > 1$, h_{1,j_0} and \ddot{h}_{1,j_0} are dictated by the above equations:

$$h_{1,j_0} = \frac{-1}{\dot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k \dot{h}_{k,j_0} + \ddot{a}_k \ddot{h}_{k,j_0} \right)$$

and

$$\ddot{h}_{1,j_0} = \frac{1}{\dot{a}_1 \eta'} \left(\sum_{k=2}^{j_0-1} \ddot{a}_k \dot{h}_{k,j_0} - \dot{a}_k \ddot{h}_{k,j_0} \eta' \right)$$

Similarly, if $\ddot{a}_1 \neq 0, \dot{a}_1 = 0$, then given h_{k,j_0} for $k > 1$, \dot{h}_{1,j_0} and \ddot{h}_{1,j_0} are still dictated by the above equations:

$$\ddot{h}_{1,j_0} = \frac{-1}{\ddot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k \dot{h}_{k,j_0} + \ddot{a}_k \ddot{h}_{k,j_0} \right)$$

and

$$\dot{h}_{1,j_0} = \frac{-1}{\ddot{a}_1} \left(\sum_{k=2}^{j_0-1} \ddot{a}_k \dot{h}_{k,j_0} - \dot{a}_k \ddot{h}_{k,j_0} \eta' \right)$$

Thus we have that for $\dot{a}_1 \neq 0, \ddot{a}_1 = 0$ or $\dot{a}_1 = 0, \ddot{a}_1 \neq 0$,

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed, } \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Subcase 2b: $\dot{\mathbf{a}}_1, \ddot{\mathbf{a}}_1 \neq \mathbf{0}$

If $\dot{a}_1, \ddot{a}_1 \neq 0$, then given h_{k,j_0} for $k > 1$, and $h_{1,j_0}^{\ddot{\cdot}}, h_{1,j_0}^{\dot{\cdot}}$ is dictated:

$$h_{1,j_0}^{\dot{\cdot}} = \frac{-1}{\dot{a}_1} \left(\ddot{a}_1 h_{1,j_0}^{\ddot{\cdot}} + \sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}} \right)$$

and we must also have

$$h_{1,j_0}^{\dot{\cdot}} = \frac{-1}{\dot{a}_1} \left(-\dot{a}_1 h_{1,j_0}^{\ddot{\cdot}} \eta' + \sum_{k=2}^{j_0-1} \ddot{a}_k h_{k,j_0}^{\dot{\cdot}} - \dot{a}_k h_{k,j_0}^{\ddot{\cdot}} \eta' \right)$$

Subtracting the first equation from the second we also get

$$0 = h_{1,j_0}^{\ddot{\cdot}} \left(\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta'}{\ddot{a}_1} \right) + \frac{1}{\dot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}} \right) - \frac{1}{\dot{a}_1} \left(\sum_{k=2}^{j_0-1} \ddot{a}_k h_{k,j_0}^{\dot{\cdot}} - \dot{a}_k h_{k,j_0}^{\ddot{\cdot}} \eta' \right)$$

If $\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta'}{\ddot{a}_1} \neq 0$, we can solve for $h_{1,j_0}^{\ddot{\cdot}}$. And hence we would get that $h_{1,j_0}^{\ddot{\cdot}}$ is dictated by the above equations as well. So if $\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta'}{\ddot{a}_1} \neq 0$, we again get that

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed, } \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Note that

$$\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta'}{\ddot{a}_1} = 0 \Leftrightarrow \left(\frac{\ddot{a}_1}{\dot{a}_1} \right)^2 + \eta' = 0$$

This will have a solution if and only if $f(x) = x^2 + \eta'$ has a solution in \mathbb{F}_{p^r} . But we chose η' such that this was irreducible over \mathbb{F}_{p^r} and thus does not have any root in \mathbb{F}_{p^r} . Therefore we can conclude that for any $\dot{a}_1, \ddot{a}_2 \neq 0$, we also have

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed, } \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Case 3: $\mathbf{a}_1 \neq \mathbf{0}$ and $\mathbf{p} = 2$

As in Case 2, if we take $\mathbf{d} = \mathbf{0}$, then $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{h}\mathbf{d}H^T - \mathbf{d})) = 1$ reduces to $\psi(\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x})) = 1$. Note that

$$x_{j_0} h_{k,j_0} = \dot{x}_{j_0} h_{k,j_0}^{\dot{\cdot}} + \ddot{x}_{j_0} h_{k,j_0}^{\ddot{\cdot}} \eta + (\dot{x}_{j_0} h_{k,j_0}^{\ddot{\cdot}} + \ddot{x}_{j_0} h_{k,j_0}^{\dot{\cdot}} + \ddot{x}_{j_0} h_{k,j_0}^{\ddot{\cdot}}) \alpha$$

So we have

$$\begin{aligned} \mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) &= \sum_{k=1}^{j_0-1} \dot{a}_k x_{j_0} \dot{h}_{k,j_0} + \ddot{a}_k x_{j_0} \ddot{h}_{k,j_0} \\ &= \sum_{k=1}^{j_0-1} \dot{a}_k (\dot{x}_{j_0} h_{k,j_0}^{\dot{\cdot}} + \ddot{x}_{j_0} h_{k,j_0}^{\ddot{\cdot}} \eta) + \ddot{a}_k (\dot{x}_{j_0} h_{k,j_0}^{\ddot{\cdot}} + \ddot{x}_{j_0} h_{k,j_0}^{\dot{\cdot}} + \ddot{x}_{j_0} h_{k,j_0}^{\ddot{\cdot}}) \\ &= \dot{x}_{j_0} \left(\sum_{k=1}^{j_0-1} \dot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}} \right) + \ddot{x}_{j_0} \left(\sum_{k=1}^{j_0-1} \dot{a}_k h_{k,j_0}^{\ddot{\cdot}} \eta + \ddot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}} \right) \end{aligned}$$

By the same reasoning as before, we must have

$$0 = \sum_{k=1}^{j_0-1} \dot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}} = \dot{a}_1 h_{1,j_0}^{\dot{\cdot}} + \ddot{a}_1 h_{1,j_0}^{\ddot{\cdot}} + \sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}}$$

and

$$0 = \sum_{k=1}^{j_0-1} \dot{a}_k h_{k,j_0}^{\ddot{\cdot}} \eta + \ddot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}} = \dot{a}_1 h_{1,j_0}^{\ddot{\cdot}} \eta + \ddot{a}_1 h_{1,j_0}^{\dot{\cdot}} + \ddot{a}_1 h_{1,j_0}^{\ddot{\cdot}} + \sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0}^{\ddot{\cdot}} \eta + \ddot{a}_k h_{k,j_0}^{\dot{\cdot}} + \ddot{a}_k h_{k,j_0}^{\ddot{\cdot}}$$

Subcase 3a: $\dot{\mathbf{a}}_1 \neq \mathbf{0}, \ddot{\mathbf{a}}_1 = \mathbf{0}$ or $\dot{\mathbf{a}}_1 = \mathbf{0}, \ddot{\mathbf{a}}_1 \neq \mathbf{0}$

If $\dot{a}_1 \neq 0, \ddot{a}_1 = 0$, then given h_{k,j_0} for $k > 1, h_{1,j_0}$ and \dot{h}_{1,j_0} are dictated by the above equations:

$$\dot{h}_{1,j_0} = \frac{1}{\dot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0} + \ddot{a}_k h_{k,j_0} \right)$$

and

$$\ddot{h}_{1,j_0} = \frac{1}{\dot{a}_1 \eta} \left(\sum_{k=2}^{j_0-1} \ddot{a}_k h_{k,j_0} - \dot{a}_k h_{k,j_0} \eta \right)$$

Similarly, if $\ddot{a}_1 \neq 0, \dot{a}_1 = 0$, then given h_{k,j_0} for $k > 1, \dot{h}_{1,j_0}$ and \ddot{h}_{1,j_0} are still dictated by the above equations:

$$\ddot{h}_{1,j_0} = \frac{1}{\ddot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0} + \ddot{a}_k h_{k,j_0} \right)$$

and

$$\dot{h}_{1,j_0} = \frac{1}{\ddot{a}_1} \left(\ddot{a}_1 h_{1,j_0} + \sum_{k=2}^{j_0-1} \ddot{a}_k h_{k,j_0} - \dot{a}_k h_{k,j_0} \eta \right)$$

Thus we have that for $\dot{a}_1 \neq 0, \ddot{a}_1 = 0$ or $\dot{a}_1 = 0, \ddot{a}_1 \neq 0$,

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed}, \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Subcase 3b: $\dot{\mathbf{a}}_1, \ddot{\mathbf{a}}_1 \neq \mathbf{0}$

If $\dot{a}_1, \ddot{a}_1 \neq 0$, then given h_{k,j_0} for $k > 1$ and $\dot{h}_{1,j_0}, \ddot{h}_{1,j_0}$ is dictated:

$$\dot{h}_{1,j_0} = \frac{1}{\dot{a}_1} \left(\ddot{a}_1 h_{1,j_0} + \sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0} + \ddot{a}_k h_{k,j_0} \right)$$

and we must also have

$$\ddot{h}_{1,j_0} = \frac{1}{\ddot{a}_1} \left(\dot{a}_1 h_{1,j_0} \eta + \ddot{a}_1 h_{1,j_0} + \sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0} \eta + \ddot{a}_k h_{k,j_0} + \ddot{a}_k h_{k,j_0} \right)$$

Adding these two equations we also get

$$0 = \ddot{h}_{1,j_0} \left(\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta}{\ddot{a}_1} + 1 \right) + \frac{1}{\dot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0} + \ddot{a}_k h_{k,j_0} \right) + \frac{1}{\ddot{a}_1} \left(\sum_{k=2}^{j_0-1} \dot{a}_k h_{k,j_0} \eta + \ddot{a}_k h_{k,j_0} + \ddot{a}_k h_{k,j_0} \right)$$

If $\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta}{\ddot{a}_1} + 1 \neq 0$, we can solve for \ddot{h}_{1,j_0} . And hence we would get that \dot{h}_{1,j_0} is dictated by the above equations as well. So if $\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta}{\ddot{a}_1} + 1 = 0$, we again get that

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed}, \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Note that

$$\frac{\ddot{a}_1}{\dot{a}_1} + \frac{\dot{a}_1 \eta}{\ddot{a}_1} + 1 = 0 \Leftrightarrow \ddot{a}_1^2 + \dot{a}_1 \ddot{a}_1 + \dot{a}_1^2 \eta = 0.$$

Recall that for $p = 2$, we have

Proposition 7.26 ([18], Proposition 1). *In a finite field of order 2^r , for $f(x) = ax^2 + bx + c$, we have the following:*

(i) f has exactly one root $\Leftrightarrow b = 0$.

(ii) f has exactly two roots $\Leftrightarrow b \neq 0$ and $\text{Tr}(\frac{ac}{b^2}) = 0$.

(iii) f has no root $\Leftrightarrow b \neq 0$ and $\text{Tr}(\frac{ac}{b^2}) = 1$,

where $\text{Tr}(x) = x + x^2 + \dots + x^{2^r-1}$.

So $\ddot{a}_1^2 + \dot{a}_1\ddot{a}_1 + \dot{a}_1^2\eta = 0$ has a root if and only if $0 = \text{Tr}(\frac{\dot{a}_1^2\eta}{\ddot{a}_1^2}) = \text{Tr}(\eta)$. Recall that η was chosen such that $x^2 + x + \eta$ was irreducible, and thus $x^2 + x + \eta$ has no root. Hence by Proposition 7.26, we must have that $\text{Tr}(\eta) = 1$. Thus $\ddot{a}_1^2 + \dot{a}_1\ddot{a}_1 + \dot{a}_1^2\eta = 0$ has no root. Therefore we can conclude that $\dot{a}_1, \ddot{a}_2 \neq 0$, we also have

$$|L_s| \leq |\{H : H_{1,j} \text{ fixed, } \forall j \neq 1\}| = |\text{Up}_{m-1}(\mathbb{F}_{p^{2r}})| = p^{r(m-1)(m-2)}.$$

Step 2: Exhibiting that the max is achieved when $s = (0, \dots, 0, b_1, 0, \dots, 0)$ and when $s = (a_1, 0, \dots, 0)$ with $\dot{a}_1 \neq 0, \ddot{a}_1 = 0$ or $\dot{a}_1 = 0, \ddot{a}_1 \neq 0$

Case 1: $\mathbf{b}_1 \neq \mathbf{0}$

We showed in Step 1 that we must have $L_{(a_j, b_j)}$ a subset of the $L_{(b_j)}$ calculated for $n = 2m$. Thus by Proposition 7.21 if $(b_j) = (b_1, 0, \dots, 0)$, then we have $L_s \subset \{H \in \text{Up}_m(\mathbb{F}_{p^{2r}}) : H_{1,j} = 0 \forall j > 1\}$ and for H of this form, we have that $\mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = 0$. Thus for H of this form, if $\mathbf{a} = \mathbf{0}$,

$$\mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) + \mathbf{b} \cdot (\mathbf{hdh}^T - \mathbf{d}) = 0.$$

So for $s = (0, \dots, 0, b_1, 0, \dots, 0)$,

$$L_s = \{H \in \text{Up}_m(\mathbb{F}_{p^{2r}}) : H_{1,j} = 0 \forall j > 1\}.$$

Case 2: $\dot{\mathbf{a}}_1 \neq \mathbf{0}, \ddot{\mathbf{a}}_1 = \mathbf{0}$ or $\dot{\mathbf{a}}_1 = \mathbf{0}, \ddot{\mathbf{a}}_1 \neq \mathbf{0}$

If $a_i = 0$ for $i \neq 1$, then by Subcase 2a and Subcase 3a in Step 1, we get that for $j_0 > 1$, $h_{1,j_0} = 0 = h_{1,j_0}^{\ddot{\cdot}}$, and hence $h_{1,j_0} = 0$. So

$$\begin{aligned} \mathbf{a} \cdot (\mathbf{x}H^T - \mathbf{x}) &= \sum_{k=1}^{m-1} a_k \cdot \left(\sum_{j=k+1}^m x_j h_{k,j} \right) \\ &= a_1 \cdot \left(\sum_{j=2}^m x_j h_{1,j} \right), && \text{since } a_i = 0, i > 1 \\ &= 0 && \text{since } h_{1,j} = 0, j > 1 \end{aligned}$$

So we get that for any p , $s = (a_1, 0, \dots, 0)$ with $\dot{a}_1 \neq 0, \ddot{a}_1 = 0$ or $\dot{a}_1 = 0, \ddot{a}_1 \neq 0$,

$$L_s = \{H : H_{1,j} = 0, \forall j \neq 1\}.$$

□

7.5 Proof

Proof of Theorem 7.1.

Case 1 ($n = 2$).

$\text{Up}_1(\mathbb{F}_{p^{2r}})$ is trivial and so $\text{Syl}_p(U(2, p^{2r})) \cong \text{antisym}^*(1, p^{2r}) \cong \mathbb{F}_{p^r}^+$.

So $\text{ed}_k(\text{Syl}_p(U(2, p^{2r}))) = \text{ed}_k(\mathbb{F}_{p^r}^+) = r$. This proves Theorem 7.1 for $n = 2$.

Case 2 ($n = 3$).

$\text{Up}_1(\mathbb{F}_{p^{2r}})$ is trivial and so $\text{Syl}_p(U(3, p^{2r})) \cong \mathbb{F}_{p^{2r}}^+ \times \text{antisym}^*(1, p^{2r}) \cong \mathbb{F}_{p^{2r}}^+ \times \mathbb{F}_{p^r}^+ \cong (\mathbb{Z}/p\mathbb{Z})^{3r}$.

So $\text{ed}_k(\text{Syl}_p(U(3, p^{2r}))) = \text{ed}_k((\mathbb{Z}/p\mathbb{Z})^{3r}) = 3r$. This proves Theorem 7.1 for $n = 3$.

Case 3 ($n = 2m, m > 1$).

Step 1: Find the lower bound

For any p , the minimum dimension of an irreducible representation of $\text{Syl}_p(U(2m, p^r))$ with non-trivial central character is given by $\theta_{s, \lambda}$ with λ trivial and $s = (b, 0, \dots, 0)$. So the minimum dimension is $p^{r(n-1)}$. Thus the minimum dimension of a faithful representation is $rp^{2r(m-1)}$ since the center has rank r .

Step 2: Construct the upper bound

Let $\{e_i\}$ be a basis for $Z(U(2m, p^{2r})) \cong (\mathbb{Z}/p\mathbb{Z})^r$ over \mathbb{F}_p . Then by Lemma 7, $\{\psi_{e_i}\}$ form a basis for the characters of $Z(U(2m, p^{2r}))$. Thus $\{\psi_{e_i}|_{Z(U(2m, p^{2r}))}\}$ form a basis for $(Z(\widehat{U(2m, p^{2r})}))$. Let

$$s_i = (e_i, 0, \dots, 0).$$

Then by Corollary 7.22, for λ trivial, $\dim(\theta_{s_i, \lambda}) = p^{2r(m-1)}$. Let

$$\rho = \oplus_i \theta_{s_i, \lambda}.$$

Then

$$\dim(\rho) = \sum_{i=1}^r \dim(\theta_{s_i, \lambda}) = \sum_{i=1}^r p^{2r(m-1)} = rp^{2r(m-1)}.$$

And by Lemma 3.3, since $\psi_{e_i}|_{Z(U(2m, p^{2r}))}$ form a basis for $(Z(\widehat{U(2m, p^{2r})}))$, we have that $\oplus_i \theta_{s_i, \lambda}$ is a faithful representation of $\text{Syl}_p(U(2m, p^{2r}))$. Thus ρ is the desired faithful representation of $\text{Syl}_p(U(2m, p^{2r}))$ of dimension $rp^{2r(m-1)}$.

Case 4 ($n = 2m + 1, m > 1$).

Step 1: Find the lower bound

By Proposition 7.24, the minimum dimension of an irreducible representation with non-trivial central character is $p^{2r(m-1)}$. Thus the minimum dimension of a faithful representation is $3rp^{2r(m-1)}$ since the center has rank $3r$.

Step 2: Construct the upper bound

Let $\{e_i\}$ be a basis for $(\mathbb{Z}/p\mathbb{Z})^r$ over \mathbb{F}_p . Let

$$S = \{(0, \dots, 0, e_i, 0, \dots, 0), (e_i, 0, \dots, 0), (e_i \alpha, 0, \dots, 0)\}.$$

Then $\{\psi_s\}_{s \in S}$ form a basis for the characters of $Z(U(2m+1, p^{2r}))$. Thus $\{\psi_s|_{\Omega_1(Z(U(2m+1, p^{2r})))}\}$ form a basis for $\widehat{\Omega_1(Z(U(2m+1, p^{2r})))}$. By Corollary 7.25, for $s \in S$, λ trivial,

$$\dim(\theta_{s, \lambda}) = p^{2r(m-1)}$$

Let $\rho = \oplus_{s \in S} \theta_{s, \lambda}$. Then

$$\dim(\rho) = \sum_s \dim(\theta_{s, \lambda}) = 3rp^{2r(m-1)}$$

And by Lemma 3.3, since $\psi_s|_{\Omega_1(Z(U(2m+1, p^{2r})))}$ form a basis for $\widehat{\Omega_1(Z(U(2m+1, p^{2r})))}$, we have that $\oplus_{s \in S} \theta_{s, \lambda}$ is a faithful representation of $\text{Syl}_p(U(2m+1, p^{2r}))$. Thus ρ is the desired faithful representation of $\text{Syl}_p(U(2m+1, p^{2r}))$ of dimension $3rp^{2r(m-1)}$. □

References

- [1] M Bardestani, K. Mallahi-Karai, and H. Salmasian. Minimal dimension of faithful representations for p -groups. *Journal of Group Theory*, 19(4):589–608. <https://arxiv.org/abs/1505.00626>.
- [2] M Bardestani, K. Mallahi-Karai, and H. Salmasian. Kirillov’s orbit method and polynomiality of the faithful dimension of p -groups. *Compositio Mathematica*, 155(8):1618–1654, 2019. <https://arxiv.org/abs/1712.02019>.
- [3] G. Berhuy and G. Favi. Essential dimension: A functorial point of view (after a. merkurjev). *Doc. Math.*, 8:279–330, 2003. <https://www.math.uni-bielefeld.de/documenta/vol-08/11.pdf>.
- [4] P. Brosnan and N. Fakhruddin. Fixed points, local monodromy, and incompressibility of congruence covers, 2020. <https://arxiv.org/abs/2007.01752>.
- [5] J. Buhler and Z. Reichstein. On the essential dimension of a finite group. *Composito Mathematica*, 106:159–179, 1997. <https://www.cambridge.org/core/journals/compositio-mathematica/article/on-the-essential-dimen>
- [6] A. Duncan. Essential dimensions of a_7 and s_7 . *Math. Res. Lett.*, 17(2):263–266, 2010. <https://arxiv.org/abs/0908.3220v1>.
- [7] A. Duncan and Z. Reichstein. Pseudo-reflection groups and essential dimension, 2014. <https://arxiv.org/abs/1307.5724>.
- [8] B. Farb, M. Kisin, and J. Wolfson. Modular functions and resolvent problems. <https://arxiv.org/abs/1912.12536>, 2018.
- [9] B. Farb, M. Kisin, and J. Wolfson. The essential dimension of congruence covers. *Compos. Math.*, 157(11):2407–2432, 2021.
- [10] L. Grove. *Classical Groups and Geometric Algebra*. American Mathematical Society, 2000. Graduate Studies in Mathematics Volume 39.
- [11] I.M. Isaacs. Characters of groups associated with finite algebras. *Journal of Algebra*, 177:708–730, 1995. <https://www.sciencedirect.com/science/article/pii/S0021869385713257>.
- [12] N. Karpenko and A. Merkurjev. Essential dimension of finite p -groups. *Inventiones mathematicae*, 172:491–508, 2008. <https://www.math.ucla.edu/~merkurev/papers/esdim-final.pdf>.
- [13] S. Lang. *Algebra*, volume 211. Springer, 2002.
- [14] M. Marjoram. Irreducible characters of a Sylow p -subgroup of the orthogonal group. *Commun. Algebra*, 27(3):1171–1195, 1999. https://www.researchgate.net/publication/28238784_Irreducible_characters_of_a_Sylow_p-subgroup
- [15] A. Merkurjev. Essential dimension. *Séminaire BOURBAKI*, 1101, 2015. <https://www.math.ucla.edu/~merkurev/papers/Exp1102.A.Merkurjev.2.pdf>.
- [16] A. Meyer and Z. Reichstein. Some consequences of the karpenko-merkurjev theorem. *Doc. Math.*, Extra Vol: Andrei A. Suslin sixtieth birthday:445–457, 2010.
- [17] A. Moretó. On the minimal dimension of a faithful linear representation of a finite group, 2021. <https://arxiv.org/abs/2102.01463>.
- [18] K. Pommerening. Quadratic equations in finite fields of characteristic 2. *unpublished manuscript*, 2000, English version 2012. <https://www.staff.uni-mainz.de/pommeren/MathMisc/QuG1Char2.pdf>.

- [19] A. Previtali. On a conjecture concerning character degrees of some p -groups. *Arch. Math.*, 65(5):375–378, 1995. https://www.researchgate.net/publication/226465705_On_a_conjecture_concerning_characters_of_some_p-groups
- [20] A. Previtali. Orbit lengths and character degrees in p -sylow subgroups of some classical lie groups. *Journal of Algebra*, 177(3):658–675, 1995. <https://www.sciencedirect.com/science/article/pii/S0021869385713221>.
- [21] A. Previtali. Maps behaving like exponentials and maximal unipotent subgroups of groups of Lie type. *Commun. Algebra*, 27(5):2511–2519, 1999. <https://www.semanticscholar.org/paper/Maps-behaving-like-exponentials-and-maximal-of-of-Previtali>
- [22] M. Marjoram R. Gow and A. Previtali. On the irreducible characters of a sylow 2-subgroup of the finite symplectic group in characteristic 2. *J. Algebra*, 2001. <https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.613.8582>.
- [23] Z. Reichstein. Essential dimension. *Proceedings of the International Congress of Mathematicians*, II:162–188, 2010.
- [24] Z. Reichstein and A. Shukla. Essential dimension of double covers of symmetric and alternating groups. *J. Ramanujan Math. Soc.*, 35(4):357–372, 2020. <https://arxiv.org/abs/1906.03698>.
- [25] Zinovy Reichstein and Angelo Vistoli. Dimension essentielle des groupes finis en caractéristique positive. *C. R., Math., Acad. Sci. Paris*, 356(5):463–467, 2018.
- [26] J.P. Serre. *Linear Representations of Finite Groups*. Springer, 1977. Translated from the French by Leonard L. Scott.
- [27] G.C. Shephard and J.A. Todd. Finite unitary reflection groups. *Canadian J. Math.*, pages 274–304, 1954. <http://www.math.ucsd.edu/~nwallach/shephard-todd.pdf>.
- [28] G. Venkataraman. On irreducibility of induced modules and an adaptation of the wigner-mackey method of little groups. *J. Korean Math. Soc.*, 50(6):1213–1222, 2013. <https://arxiv.org/abs/0908.0026>.