

Rigorous justification for the space-split sensitivity algorithm to compute linear response in Anosov systems

Nisha Chandramoorthy*, Malo Jézéquel†

Abstract

Ruelle [Rue97] first proved a rigorous formula for linear response of smooth, transitive Anosov diffeomorphisms. Recently, practically computable realizations of Ruelle’s formula have emerged that potentially enable sensitivity analysis of the high-dimensional chaotic numerical simulations such as encountered in climate studies. In this paper, we provide a rigorous convergence proof of one such efficient computation, the space-split sensitivity, or S3, algorithm [CW21] for one-dimensional unstable manifolds. In S3, Ruelle’s formula is computed as a sum of two terms obtained by decomposing the perturbation vector field into a coboundary and a remainder that is parallel to the unstable direction. Such a decomposition results in a splitting of Ruelle’s formula that is amenable to efficient computation. We prove the existence of the S3 decomposition and the convergence of the computations of both resulting components of Ruelle’s formula.

1 Introduction

In the *linear response* regime, statistical outputs of a dynamical system vary proportionally to infinitesimal input perturbations applied to the dynamics. In idealized chaotic maps, known as Anosov diffeomorphisms, Ruelle first proved the existence of linear response and proposed a linear response formula [Rue97, Rue03] (see §1.1 for details). Linear response studies are of fundamental scientific and engineering interest, especially in climate studies [LBH⁺14, RLL16, BLL20], because linear response quantifies the influence of infinitesimal parameter perturbations, such as anthropogenic forcings, on the ensemble behavior, or equivalently in ergodic systems, the long-term behavior of the system. Furthermore, the derivatives of statistics to parameters are crucial objects in scientific computing: they are used to solve design

*Center for Computational Science and Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139. Email: nishac@mit.edu

†Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139. Email: mpjez@mit.edu

optimization, parameter selection and uncertainty quantification problems (see, e.g., [JS18, DIX19, SBH⁺16] for surveys of such applications in aerodynamic turbulent flows). But despite their importance, the lack of an efficient and rigorous computation of Ruelle’s linear response formula [Rue97, Rue03] has hindered downstream applications of linear response in high-dimensional chaotic models encountered in practice. Recently, some new methods have been proposed to evaluate decomposed and regularized versions of Ruelle’s formula [CW21, Ni21, CW20]. The purpose of this paper is the rigorous proof of one such method, the space-split sensitivity (S3) algorithm [CW21].

Before recalling the steps of the S3 algorithm in §1.2 and stating our main result, Theorem 2, let us recall a few facts about statistical properties of Anosov systems.

1.1 Linear response of Anosov diffeomorphisms

Let T denote a \mathcal{C}^r transitive Anosov diffeomorphism, with $r > 2$, on a compact \mathcal{C}^∞ Riemannian manifold M . We recall that Anosov means that there is a splitting $TM = E_u \oplus E_s$ of the tangent bundle, invariant by the derivative of T , such that E_u is uniformly expanded and E_s is uniformly contracted by the derivative of T (see [KH95, Definition 6.4.2] for a detailed definition). The fundamental *stable/unstable manifold theorem* [Yoc95, §3.6] asserts that the subbundle E_u and E_s are uniquely integrable: there are Hölder-continuous foliations W_u and W_s with uniformly \mathcal{C}^r leaves (called respectively unstable and stable manifolds) that are tangent to E_u and E_s respectively.

Owing to their idealized setting, Anosov diffeomorphisms enjoy a neat statistical description. In particular, it is well-known [Sin68] that T admits a unique SRB measure μ , that is a unique invariant probability measure with smooth conditionals on unstable manifolds (see [You02] for an introduction and survey of results on SRB measures). The SRB measure is *physical* in the sense that for Lebesgue almost every $x \in M$ and every continuous function f on M ,

$$\frac{1}{n} \sum_{k=0}^{n-1} f(T^k x) \xrightarrow{n \rightarrow \infty} \int_M f d\mu. \quad (1)$$

Even though the measure μ is in general not absolutely continuous [Sin73, Corollary 1], its properties given above imply that μ governs the statistical properties of T at the scale of Lebesgue measure, hence their importance in applied sciences. Many strong properties of the SRB measure μ are now well-known: central limit theorem [Den89], large deviation principle [OP89], full asymptotics for the correlations in term of resonances [GL06], etc.

Of much importance for applications is the regularity of the dependence of μ on the dynamics T . To put it in more concrete terms, let $t \mapsto T_t$ be

a smooth family of Anosov diffeomorphisms on M with $T_0 = T$. For each t near 0, the Anosov diffeomorphism T_t admits a unique SRB measure μ_t and it is then natural to wonder what is the regularity of the map $t \mapsto \mu_t$. When this map is differentiable at 0, we say that *linear response* holds. In that case, the value of the derivative at 0 of the map $t \mapsto \mu_t$ is sometimes called the linear response, as it describes the first order response of statistical properties of a physical system under perturbations.

In the context of transitive Anosov diffeomorphisms, linear response has been established first by Ruelle in [Rue97, Rue03]. In [GL06], Gouëzel and Liverani used the spectral properties of transfer operators to give a much more detailed description of the variation of the statistical properties of the SRB measure under perturbation. See also [Bal18, §5.3] for a textbook treatment of this problem. These references give a very detailed picture of the situation, let us just mention the following result that ensures that linear response holds.

Theorem 1 ([GL06, Theorem 2.8], [Bal18, Theorem 5.26]). *Let $t \mapsto T_t$ be a \mathcal{C}^2 family of \mathcal{C}^r diffeomorphisms, with $r > 2$ and $T_0 = T$. Then, for t close enough to 0 the map T_t has a unique SRB measure μ_t . Moreover, for every $\epsilon > 0$, there is $\eta > 0$ such that the map $t \mapsto \mu_t \in (\mathcal{C}^{2+\epsilon})'$ is $\mathcal{C}^{1+\eta}$ near 0.*

Although the above assumptions are rather restrictive, and cannot be verified with certainty in chaotic models encountered in practice, linear response theory has been applied successfully for a variety of systems including climate models, turbulent flows and biological systems [HM20, Ni19, BLL20, Ces19]. The justification for this robustness of linear response in noisy real-life systems has been studied from the statistical physics standpoint [WG19, WG18]. Meanwhile, in the dynamical systems literature, rigorous proofs of existence of linear response have been found for a wider class of systems than Anosov systems, including certain partially hyperbolic systems [Dol04], some stochastic systems [HM10][GG19], certain nonuniformly hyperbolic systems [BCV16] and even for some systems with intermittencies [BS12][BT16][BS16]. See the survey [Bal14] and references therein for further information on linear response.

The subject of our paper is the S3 algorithm, which computes the derivative at 0 of the map $t \mapsto \mu_t$. To do so, we will rely on the following formula that has been proposed, and first proved, by Ruelle [Rue97]: for a smooth observable f , we expect that

$$\frac{d}{dt} \left(\int_M f d\mu_t \right)_{|t=0} = \sum_{n \geq 0} \mu(X(f \circ T^n)). \quad (2)$$

where the vector field X describes the infinitesimal parameter perturbation:

$$X = \frac{d}{dt} (T_t)_{|t=0} \circ T^{-1},$$

so that the perturbed dynamics are given by $T_t = T + tX \circ T + \mathcal{O}(t^2)$. Under the assumption of Theorem 1, Ruelle’s formula (2) holds as soon as f is $\mathcal{C}^{3+\epsilon}$ (see for instance [Bal18, §5.3]).

At first glance, the series in the right-hand side of (2) does not seem to converge, due to the exponential growth of the derivatives of T^n when n tends to $+\infty$. However, spectral investigation of a transfer operator associated to T shows that, thanks to the integration against μ , the series on the right-hand side is actually converging exponentially fast [Rue97, Rue03]. We will prove a slightly more general statement in Lemma 5.2. Nevertheless, it is inefficient to practically approximate the integrals against μ in (2) using a Monte Carlo method, i.e., approximating each integral as a finite sample average. The variance of the sample average also grows exponentially with n , making such a direct evaluation of Ruelle’s formula (2) computationally impractical for high-dimensional systems encountered in practice [EHL04, CFTW19].

1.2 The S3 algorithm

The central object of this paper is the rigorous justification for the use of the space-split sensitivity, or S3, algorithm proposed by Chandramoorthy and Wang in [CW20, CW21]. The S3 algorithm is one method to efficiently compute Ruelle’s formula (2). Ni’s Linear Response Algorithm [Ni20] is another alternative that converges to (2), wherein the computations of the non-intrusive least squares shadowing algorithm [NW17] and a particular characterization of the unstable divergence are leveraged to provide a fast computation of (2). Ni’s Linear Response Algorithm proposes to split Ruelle’s formula into shadowing and an unstable contribution while we propose a different decomposition in the S3 algorithm (see [CW21, Appendix A], [Ni20] for a more detailed comparison of these two algorithms). Abramov and Majda [AM07] developed a blended response technique for (2) whereby an ensemble sensitivity estimate [LHAH02, EHL04] – a direct numerical approximation of (2) – is used for responses to stable components of the perturbations. Since, as mentioned above, Monte Carlo evaluation is infeasible for long-term response (large n in (2)) to the unstable component of the perturbation, Ruelle’s formula is rewritten, using integration by parts, as a sum of correlations between the observable f and an unstable divergence (see Ruelle’s original work [Rue97] and [Jia12] for details) with respect to the conditional measures of the SRB measure on the unstable manifold. In [AM07], this unstable divergence is computed under the assumption (based on a fluctuation-dissipation argument) that the conditional densities of the SRB measure on the unstable manifold are Gaussian.

Other computational approaches to linear response are available, such as [BGNN17], where using suitable discretization schemes, the authors approximate the transfer operator associated to a one-dimensional dynamics,

to deduce an estimate of linear response. Even if transfer operators appear in this paper, our approach is quite different: we rely on formula (2) rather than on a functional analytic characterization of the linear response.

We recall that $r > 2$ and T is a transitive C^r Anosov diffeomorphism on a compact C^∞ Riemannian manifold M . In addition, we assume that the unstable direction of T is 1-dimensional and orientable. Consequently, there is a unit Hölder-continuous vector field X_u spanning the unstable direction (unique up to sign). Here, the orientability condition could be removed by replacing M by a twofold cover.

Considering the formula (2), the S3 algorithm gives an approximate value of the sum

$$\Psi_T(X, f) := \sum_{n \geq 0} \mu(X(f \circ T^n)) \quad (3)$$

for any C^{1+} vector field X and C^{1+} observable f .

The S3 algorithm is as follows. The input data are the map T , the vector field X and the observable f . Further implementation details are discussed in Remark 1.1.

Step 1: Find the unstable direction for T . To do so, start with any vector field X_u^0 and then set $X_u^{k+1} = \frac{T_* X_u^k}{|T_* X_u^k|}$ for $k \geq 0$. At most points, X_u^n converges to $\pm X_u$ exponentially fast (if we choose X_u^0 in an unstable cone for T , then the convergence holds everywhere and uniformly).

Step 2: Compute a vector field V so as to decompose X as

$$X = \underbrace{Y}_{\text{unstable term}} + \underbrace{V - T_* V}_{\text{coboundary term}}, \quad (4)$$

where Y is parallel to the unstable direction. Moreover, we require that Y and V are Hölder-continuous and that they have one Hölder-continuous derivative in the unstable direction (the precise meaning of this regularity assumption is detailed in §2). The existence of the above decomposition is proved in Proposition 3.1, and a way to compute Y and V explicitly is explained in Remark 3.2. Notice that we can write

$$\Psi_T(X, f) = \Psi_T(Y, f) + \Psi_T(V - T_* V, f)$$

and compute the contribution of Y and $V - T_* V$ separately. Moreover, Lemma 5.1 asserts that $\Psi_T(V - T_* V, f) = \mu(Vf)$.

Step 3: Find a Hölder-continuous function ρ such that for any C^{1+} function g we have $\mu(Yg) = \mu(g\rho)$. The existence of ρ is obtained by integration by parts using the fact that μ has smooth conditionals on unstable manifolds (see Proposition 4.1). In Lemma 4.2, we give a characterization of ρ that allows us to compute it explicitly (see also Remark 4.3). Using ρ , we can obtain the unstable contribution as

$$\Psi_T(Y, f) = \sum_{n \geq 0} \mu(\rho f \circ T^n), \quad (5)$$

where the sum converges exponentially fast due to the exponential decay of correlations (see for [Liv95]) for T (notice that ρ is Hölder and that $\mu(\rho) = \mu(Y1) = 0$). The advantage of (5) over the initial definition of $\Psi_T(Y, f)$ is that we do not need to compute exponentially growing derivatives anymore. Due to the exponential convergence of the series in (5), we get a good approximation of $\Psi_T(Y, f)$ by truncating the series at some $L \gg 1$:

$$\Psi_T(Y, f) \simeq \sum_{n=0}^{L-1} \mu(\rho f \circ T^n). \quad (6)$$

Step 4: Compute $\mu(f\rho), \dots, \mu(f \circ T^{L-1}\rho)$ and $\mu(Vf)$ using either of the two methods (probabilistic or deterministic) proposed in §6, and take as an approximate value for $\Psi_T(X, f)$:

$$\Psi_T(X, f) \simeq \mu(Vf) + \sum_{k=0}^{L-1} \mu(f \circ T^k \rho). \quad (7)$$

Our main result is then the following:

Theorem 2. *The S3 algorithm converges: with the notations above, we have*

$$\Psi_T(X, f) = \mu(Vf) + \sum_{n \geq 0} \mu(f \circ T^n \rho), \quad (8)$$

where we recall that X and f are C^{1+} , while T is C^{2+} . Moreover, there is a $\gamma > 0$ such that the error made after a computation of time τ when estimating $\Psi_T(X, f)$ using the S3 algorithm with the deterministic method in Step 4 is $\mathcal{O}(\tau^{-\gamma})$. If one uses instead the probabilistic method, then γ may be chosen arbitrarily close to $\frac{1}{2}$ but the convergence only holds almost surely.

The identity (8) is an immediate consequence of Propositions 3.1 and 4.1 and Lemma 5.2. The speed of convergence is discussed in Remarks 6.2 and 6.3.

Remark 1.1. Let us describe briefly how the S3 algorithm is actually implemented (see [CW21] for details). In order to approximate the integrals $\mu(f\rho), \dots, \mu(f \circ T^{L-1}\rho)$ and $\mu(Vf)$ in Step 4, we only need to know ρ and V at a finite number of points. To compute ρ and V at a point x , we can proceed as follows.

We start by computing a large segment $T^{-n}x, \dots, T^{-1}x, x$ of the backward orbit of x . Then, we cut this orbit in 3 parts of comparable length $O_1 = \{T^{-n}x, \dots, T^{-m-1}x\}$, $O_2 = \{T^{-m}x, \dots, T^{-\ell-1}x\}$ and $O_3 = \{T^{-\ell}x, \dots, x\}$. We use the first orbit segment O_1 to perform Step 1: we choose any vector $X_u^0(T^{-n}x)$ over $T^{-n}x$ (if possible near the unstable direction), we can then recursively compute $X_u^1(T^{-n+1}x), \dots, X_u^k(T^{-n+k}x)$. If O_1 is large enough,

this process gives us a good approximation of X_u at the points of O_2 and O_3 (the error is exponentially small in the size of O_1 , and the convergence holds provided the initial vector has not been chosen in the stable direction).

We use the orbit segment O_2 to perform Step 2. The method given in Remark 3.2 (see also Remark 2.7) to compute Y and V at a point $y \in M$ only uses values of X, X_u and of the derivative of T at points in the backward orbit of y . This procedure converges to V and Y exponentially fast in the length of the orbit used. Hence, using the points of O_2 (with the value of X_u computed at Step 1), we get an approximation of Y and V at the points of O_3 .

Finally, we proceed at Step 3 using the orbit segment O_3 : using a similar method as in Step 2, we can compute ρ at x using the values of Y and X_u at the points of O_3 (see Remark 4.3), and the error made is exponentially small in the length of O_3 .

Now that we know how to compute the values of V and ρ at a given point, we can apply the methods described in §6 to approximate the integrals $\mu(f\rho), \dots, \mu(f \circ T^{L-1}\rho)$ and $\mu(Vf)$. In practice, these methods do not need the value of V and ρ at any specific x , but rather along any μ -typical orbit of T . As a result, one need not invert the map T , but only compute forward orbits (and use the forward orbit of x as a backward orbit for $T^n x$).

Remark 1.2. Instead of (4), one could naturally imagine decomposing X as $X = V_u + V_s$ where V_u and V_s are respectively in the unstable and stable direction. Then, the terms in the sum (3) defining $\Psi_T(V_s, f)$ would be exponentially decaying. However, one could not use integration by parts in the unstable direction to compute $\Psi_T(V_u, f)$ as we do for $\Psi_T(Y, f)$ in Step 3 of the S3 algorithm. Indeed, in general the stable and unstable directions of T are only Hölder-continuous, and so will be V_u and V_s , and this is not enough regularity to perform an integration by parts in the unstable direction (one needs at least one derivative in the unstable direction). See §4 for details.

The remainder of this article proves the main result (Theorem 2). The organization is as follows. In §2, we give a precise meaning to the expression “having a Hölder-continuous derivative in the unstable direction”. We also give a bound on the spectral radii of certain transfer operators, Proposition 2.6, that will be the main tool in order to construct Y, V and ρ in Steps 2 and 3 of the S3 algorithm. This bound is also what ensures that the Y, V and ρ may be computed with exponential precision. In §3, we explain how the vector fields Y and V from Step 2 are constructed (Proposition 3.1). The section §4 is devoted to the existence, Proposition 4.1, and computation, Lemma 4.2, of the function ρ from Step 3. In §5, we prove the formula $\Psi_T(V - T_*V, f) = \mu(Vf)$ which is essential to the validity of the S3 algorithm (Lemma 5.1). This point was missing in [CW21] and is the main novelty from our work. The proof of this crucial result is based on the spectral

decomposition of a Koopman operator acting on the spaces of anisotropic distributions from [GL06]. Finally, in §6, we explain how the integrals in Step 4 of the algorithm may be approximated, and we estimate the precision of the S3 algorithm (see Remarks 6.2 and 6.3).

Acknowledgements

Both authors thank Semyon Dyatlov for introducing them to each other and for useful discussions about this work. N.C. is supported by the Air Force Office of Scientific Research Grant No. FA8650-19-C-2207 and Department of Energy Research Grant No. DE-FOA-0002068-0018. During the elaboration of this work, M.J. was supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 787304). Most of this work was done while MJ was working at LPSM¹.

2 Spectral radii of transfer operators on vector bundles

In this section, we define precisely what we mean by having a Hölder-continuous derivative in the unstable direction (as in Step 2 of the S3 algorithm) and give the basic properties of this regularity condition (see Lemma 2.12). The main point of this section is Proposition 2.6 that gives a bound on the spectral radii of certain transfer operators associated to T . Since we will need to consider transfer operators acting both on functions and on vector fields, it is convenient to work here in the context of a general vector bundle.

Let $p : E \mapsto M$ be a C^r complex vector bundle over M . We denote by $E_x := p^{-1}(\{x\})$ the fiber of E over x . Let \mathbb{T} be a lift of T , that is a map from E to itself that satisfies $p \circ \mathbb{T} = T \circ p$, and whose restriction to each fiber is linear. To \mathbb{T} , we associate a transfer operator acting on sections of E , defined by

$$\mathcal{L}_{\mathbb{T}}v(x) = \mathbb{T}(v(T^{-1}x)), \quad (9)$$

for $x \in M$ and v a section of E .

Before studying the properties of operators of the form $\mathcal{L}_{\mathbb{T}}$, let us give the main examples of such operators that appear in the S3 algorithm.

Example 2.1. If $E = M \times \mathbb{C}$ is the trivial line bundle over M and $h : M \rightarrow \mathbb{C}$ is a scalar function, then we can define a lift \mathbb{T} for T by

$$\mathbb{T}(x, z) = (Tx, h(Tx)z) \quad \text{for } (x, z) \in M \times \mathbb{C}$$

¹Laboratoire de Probabilités, Statistique et Modélisation (LPSM), CNRS, Sorbonne Université, Université de Paris, 4, Place Jussieu, 75005 Paris, France

In this case, we will use an abbreviated notation, \mathcal{L}_h , for the transfer operator $\mathcal{L}_{\mathbb{T}}$ which is defined by

$$\mathcal{L}_h u(x) = h(x)u(T^{-1}x),$$

for $x \in M$ and u a function on M .

Example 2.2. Another basic example is $E = TM$, the tangent vector bundle of M , we can then define the lift dT of T by

$$\mathbb{T}(x, v) = d_x T \cdot v \quad \text{for } v \in T_x M.$$

This example will be crucial in the decomposition of X . Indeed, we see that in this case we retrieve $\mathcal{L}_{\mathbb{T}} = T_*$, defined by

$$T_* V(x) = d_{T^{-1}x} T \cdot V(T^{-1}x),$$

for $x \in M$ and V a vector field on M .

Let us fix a \mathcal{C}^r covariant derivative ∇ on E . By this, we just mean that if X is a \mathcal{C}^r vector field on M and v a \mathcal{C}^r section of E then $\nabla_X v$ is a \mathcal{C}^{r-1} section of E , and we require in addition that if f is a \mathcal{C}^r function on M then $\nabla_{fX} v = f \nabla_X v$ and $\nabla_X(fv) = Xf \cdot v + f \nabla_X v$. Notice that if we work in a trivialization of E that identifies locally the fibers of E with \mathbb{R}^m , with basis e_1, \dots, e_m , and the tangent space of M with \mathbb{R}^n , with basis $\partial_{x_1}, \dots, \partial_{x_m}$, then, writing $X = \sum_{j=1}^n X^j \partial_{x_j}$ and $v = \sum_{k=1}^m v^k e_k$, we have

$$\nabla_X v = dv \cdot X + \sum_{\substack{i,k=1,\dots,n \\ j=1,\dots,m}} \Gamma_{j,k}^i X^j v^k e_i, \quad (10)$$

where the Christoffel symbols $\Gamma_{j,k}^i$ are \mathcal{C}^{r-1} functions. If E is as in Example 2.1, we can just take ∇ to be the trivial connection, and in Example 2.2 it is natural to choose the Levi-Civita connection associated to the Riemannian metric on M .

We will study the operator $\mathcal{L}_{\mathbb{T}}$ acting on spaces that we introduce now. The first space is the space $\mathcal{C}^0(M; E)$ of continuous sections of E . In order to define the topology on $\mathcal{C}^0(M; E)$, we choose a smooth Riemannian metric on E and, letting $|\cdot|$ denote the associated norm on the fibers of E , we define the norm on $\mathcal{C}^0(M; E)$ by

$$|v|_{\infty} = \sup_{x \in M} |v(x)| \quad \text{for } v \in \mathcal{C}^0(M; E).$$

One can then check that the norm $|\cdot|_{\infty}$ induces a structure of Banach space on $\mathcal{C}^0(M; E)$. Moreover, another choice of Riemannian metric on E leads to an equivalent norm.

We will also need for $0 < \alpha < 1$ the space $\mathcal{C}^\alpha(M; E)$ of α -Hölder continuous sections of E . In order to define the topology on this space, we recall that we fixed a smooth Riemannian metric on M and choose $\epsilon > 0$ such that for $x, y \in M$ with $d(x, y) \leq 2\epsilon$ there is a unique geodesic $c_{y,x}$ from y to x in the ball of center x and radius 2ϵ . For such x and y , we let $S_{x,y} : E_y \rightarrow E_x$ denote the parallel transport for ∇ along $c_{y,x}$ (which is well-defined since $r > 1$). We can then define for $v \in \mathcal{C}^\alpha(M; E)$ the best Hölder constant for v as

$$|v|_\alpha := \sup_{\substack{x,y \in M \\ d(x,y) \leq \epsilon}} \frac{|v(x) - S_{x,y}v(y)|}{d(x,y)^\alpha}. \quad (11)$$

A norm on $\mathcal{C}^\alpha(M; E)$ is then defined by

$$\|v\|_\alpha := |v|_\infty + |v|_\alpha \quad \text{for } v \in \mathcal{C}^\alpha(M; E).$$

The norm $\|\cdot\|_\alpha$ defines a structure of Banach space on $\mathcal{C}^\alpha(M; E)$. Other choices of Riemannian metric on M and of covariant derivative on E lead to equivalent norms. Notice that one could also define the best Hölder constant (11) by using a finite number of trivialization for E : this procedure also gives an equivalent norm.

We will also need a slightly less classical space of sections of E : the space $\mathcal{C}^{\alpha,k}(M; E)$ of sections of E that admits k derivatives in the unstable direction that are α -Hölder. Similar classes of regularity have already been introduced in the literature on hyperbolic dynamics, see [DLL01]. The point of introducing the spaces $\mathcal{C}^{\alpha,k}(M; E)$ is that on one hand there are large enough so that we can perform the decomposition (4) in these spaces (see §3), and on the other hand having derivatives in the unstable direction is essential in Steps 3 and 4 of the S3 algorithm (see §4 and §5). The class of differentiable function is in general too large to get the decomposition (4), while the Hölder regularity is not enough for the arguments in §4 and §5 (that involve integration by parts for instance). To define precisely the space $\mathcal{C}^{\alpha,k}(M; E)$, we recall the vector field X_u on M which is such that for every $x \in M$ the vector $X_u(x)$ spans the unstable direction at x and has size 1 for the Riemannian metric on M .

We say that a section v of E is differentiable in the unstable direction if its restriction to any unstable manifold is differentiable. In that case, we can define $\nabla_{X_u} v$ by applying the pull-back of ∇ onto the unstable manifold to the restriction of v . Let us detail this definition. Let x_0 be a point in M and W be a local unstable manifold for T at x_0 . Since W is a \mathcal{C}^r manifold and the restriction of v to W is differentiable, we can find a differentiable section \tilde{v} of E that coincides with v on W . Then, it makes sense to consider $\nabla_{X_u} \tilde{v}$ (even if X_u is not \mathcal{C}^r , see (10)), and since X_u is tangent to W , we see that the value of $\nabla_{X_u} \tilde{v}$ in the interior of W does not depend on the choice of \tilde{v} . In particular it makes sense to set

$$\nabla_{X_u} v(x_0) := \nabla_{X_u} \tilde{v}(x_0).$$

We can then define the space $\mathcal{C}^{\alpha,k}(M; E)$ by induction on k , setting $\mathcal{C}^{\alpha,0}(M; E) = \mathcal{C}^\alpha(M; E)$. Then for $k \in \mathbb{N}$, we say that $v \in \mathcal{C}^{\alpha,k+1}(M; E)$ if $v \in \mathcal{C}^\alpha(M; E)$, the section v is differentiable in the unstable direction and $\nabla_{X_u} v \in \mathcal{C}^{\alpha,k}(M; E)$. The space $\mathcal{C}^{\alpha,k}(M; E)$ will be endowed with the norm:

$$\|v\|_{\alpha,k} = \sum_{\ell=0}^k \|\nabla_{X_u}^\ell v\|_\alpha \quad \text{for } v \in \mathcal{C}^{\alpha,k}(M; E).$$

If $v \in \mathcal{C}^{\alpha,k}(M; E)$, we will sometimes say for short that v is $\mathcal{C}^{\alpha,k}$. We will also write $\mathcal{C}^{\alpha,k}$ instead of $\mathcal{C}^{\alpha,k}(M; M \times \mathbb{C})$.

Remark 2.3. We gave a definition of the space $\mathcal{C}^{\alpha,k}(M; E)$ for any $k \in \mathbb{N}$ and $0 \leq \alpha < 1$. However, when $k \geq 1$, it is not clear *a priori* that this definition does not depend on the choices of Riemannian metric on M and of covariant derivative on E . Since the vector field X_u is not smooth, it is not even clear that the space $\mathcal{C}^{\alpha,k}(M; E)$ is non-trivial. It follows from Lemma 2.4 below that the space $\mathcal{C}^{\alpha,k}(M; E)$ is well-behaved as soon as $k + \alpha \leq r$ and X_u is $\mathcal{C}^{\alpha,k-1}(M; E)$. It follows from standard results in hyperbolic dynamical systems that this condition is satisfied if $k < r$ and α is small enough (see for instance [HP70, §6] and [DLL01, Theorem 2]). However, we will retrieve this regularity of X_u by a bootstrap argument using Proposition 2.6, see Lemma 2.12.

Let us mention here some elementary properties of these spaces that will be useful later.

Lemma 2.4. *Let $k \in \mathbb{N}$ and $0 \leq \alpha < 1$ be such that $k + \alpha \leq r$. If $k \geq 1$, assume in addition that $X_u \in \mathcal{C}^{\alpha,k-1}(M; TM)$. Then the following properties hold:*

- (i) *if v is a $\mathcal{C}^{k+\alpha}$ section of E , then $v \in \mathcal{C}^{\alpha,k}(M; E)$;*
- (ii) *the space $\mathcal{C}^{\alpha,k}(M; E)$ depends neither on the choice of the Riemannian metric on M nor on the choice of the covariant derivative ∇ (up to taking an equivalent norm);*
- (iii) *$\mathcal{C}^{\alpha,k}(M; E)$ is a Banach space;*
- (iv) *if $k \geq 1$ and $Y \in \mathcal{C}^{\alpha,k-1}(M; TM)$ is parallel to the unstable direction, then there is $a \in \mathcal{C}^{\alpha,k-1}$ such that $Y = aX_u$.*

Proof. (i) The restriction of v to any unstable manifold is \mathcal{C}^k , we can consequently define the $\nabla_{X_u}^\ell v$ for $\ell = 0, \dots, k$. Using the expression (10) for ∇ in local coordinates, we see that $\nabla_{X_u}^\ell v$ may be expressed in these coordinates as

$$\begin{aligned} \nabla_{X_u}^\ell v &= d^\ell v \cdot (X_u, \dots, X_u) \\ &\quad + \Gamma \left(X_u, \nabla_{X_u} X_u, \dots, \nabla_{X_u}^{\ell-1} X_u, v, \nabla_{X_u} v, \dots, \nabla_{X_u}^{\ell-1} v \right), \end{aligned}$$

where Γ is a polynomial, of order at most $\ell + 1$ in each entry, with $\mathcal{C}^{r-\ell}$ coefficients (to actually prove this formula when $k \geq 1$, consider the restriction of X_u to an unstable curve and extends it to a \mathcal{C}^{k-1} vector field near that curve). Hence, we see, by induction on ℓ , that $\nabla_{X_u}^\ell v$ is indeed α -Hölder for $\ell = 0, \dots, k$.

(ii) The proof is an induction on k , the case $k = 0$ being well-known. Assume that $k \geq 1$ and that the result holds for $k - 1$. Changing the Riemannian metric on M amounts to replacing X_u with $\tilde{X}_u = aX_u$, where the normalizing factor $a > 0$ is in $\mathcal{C}^{\alpha, k-1}$ using our assumption on X_u . Let $\tilde{\nabla}$ be another \mathcal{C}^r covariant derivative on E . Then, there is a \mathcal{C}^{r-1} section γ of the vector bundle $T^*M \otimes E^* \otimes E$ such that for X a vector field and v a section of E , we have

$$\tilde{\nabla}_X v = \nabla_X v + \gamma(X, v).$$

Consequently, if v belongs to the space $\mathcal{C}^{\alpha, k}(M; E)$ defined using the original metric on M and the covariant derivative ∇ we find that

$$\tilde{\nabla}_{\tilde{X}_u} v = a\nabla_{X_u} v + a\gamma(X_u, v).$$

By assumption on v , we know that $\nabla_{X_u} v$ and v belongs to $\mathcal{C}^{\alpha, k-1}(M; E)$, whose definition does not depend on the choice of Riemannian metric on M nor on ∇ by induction hypothesis. Applying Leibniz rule and point (i) to γ , we see that $\tilde{\nabla}_{\tilde{X}_u} v$ belongs to $\mathcal{C}^{\alpha, k-1}(M; E)$. Consequently, v also belongs to the space $\mathcal{C}^{\alpha, k}(M; E)$ defined using the new metric on M and the covariant derivative $\tilde{\nabla}$. It follows that the space $\mathcal{C}^{\alpha, k}(M; E)$ does not depend on these choices.

(iii) Let $(v_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in $\mathcal{C}^{\alpha, k}(M; E)$. Since $\mathcal{C}^\alpha(M; E)$ is a Banach space, we see that for $\ell = 0, \dots, k$, the sequence $(\nabla_{X_u}^\ell v_n)_{n \in \mathbb{N}}$ converges to an element u_ℓ of $\mathcal{C}^\alpha(M; E)$. Hence, we only need to prove that the restriction of u_0 to any unstable submanifold is \mathcal{C}^k and that $\nabla_{X_u}^\ell u_0 = u_\ell$. This is immediate by restricting to a piece of submanifolds parametrized by arclength and applying the fact that the space of \mathcal{C}^k sections of a \mathcal{C}^k vector bundle is a Banach space.

(iv) Just set $a = \langle X_u, Y \rangle$ and apply Leibniz formula to find that a is $\mathcal{C}^{k-1, \alpha}$. \square

In order to bound the spectral radius of the transfer operator $\mathcal{L}_\mathbb{T}$ defined by (9) acting on $\mathcal{C}^{\alpha, k}(M; E)$, let us introduce the quantity

$$\theta = \theta(\mathbb{T}) := \limsup_{n \rightarrow +\infty} \left(\sup_{x \in M} \sup_{v \in E_x \setminus \{0\}} \frac{|\mathbb{T}^n v|}{|v|} \right)^{\frac{1}{n}}.$$

We will also need the maximal expansion rate for T^{-1} :

$$\Lambda := \limsup_{n \rightarrow +\infty} \left(\sup_{x \in M} \sup_{v \in T_x M \setminus \{0\}} \frac{|d_x T^{-n} v|}{|v|} \right)^{\frac{1}{n}}.$$

Remark 2.5. Let us notice that the lift \mathbb{T} of T identifies with a section of the bundle of linear application from E to T^*E (the reciprocal image of E by T). Consequently, it makes sense to say that \mathbb{T} is $\mathcal{C}^{\alpha,k}$.

We can now state the main result of this section.

Proposition 2.6. *Let $0 \leq \alpha < 1$ and $k \leq r$. Assume that \mathbb{T} is $\mathcal{C}^{\alpha,k}$ (in the sense of Remark 2.5), that $k + \alpha \leq r$ and that $X_u \in \mathcal{C}^{\alpha, \min(k-1, 0)}(M; TM)$. Then $\mathcal{L}_{\mathbb{T}}$ induces a bounded operator on $\mathcal{C}^{\alpha,k}(M; E)$ with spectral radius less than $\Lambda^\alpha \theta$.*

To prove Proposition 2.6, we will need the function $h = \langle T_* X_u, X_u \rangle^{-1}$ that satisfies

$$h.T_* X_u = X_u. \quad (12)$$

Notice that if $0 \leq \alpha < 1$ and $k < r - 1$ are such that X_u is $\mathcal{C}^{\alpha,k}$, then so is h by Leibniz formula. It follows from the hyperbolicity of T that

$$\limsup_{n \rightarrow +\infty} \sup_{x \in M} \left(\prod_{\ell=1}^n |h \circ T^\ell| \right)^{\frac{1}{n}} < 1. \quad (13)$$

Before proving Proposition 2.6, let us discuss how we will use it in the context of the S3 algorithm.

Remark 2.7. Before proving Proposition 2.6, let us explain how we will use it in the context of the S3 algorithm. In order to compute Y, V and ρ in Steps 2 and 3 of the algorithm, we will need to solve for v equations of the form

$$(I - \mathcal{L}_{\mathbb{T}})v = w, \quad (14)$$

where w is a section of a vector bundle E over M and $\mathbb{T} : E \rightarrow E$ a lift of T , see Remarks 3.2 and 4.3. If \mathbb{T} and w are continuous and $\theta(\mathbb{T}) < 1$, then it follows from Proposition 2.6 that (14) has a unique continuous solution v which is given by

$$v = (I - \mathcal{L}_{\mathbb{T}})^{-1}w = \sum_{n \geq 0} \mathcal{L}_{\mathbb{T}}^n w. \quad (15)$$

It will be useful to notice that if \mathbb{T} and w are Hölder-continuous, then so is v , since we can always find a small $\alpha > 0$ such that $\theta(\mathbb{T})\Lambda^\alpha < 1$. Similarly, if \mathbb{T} and w are $\mathcal{C}^{\alpha,k}$ then so is v .

When actually implementing the S3 algorithm, equations of the form (14) are solved approximately by choosing a large N and using as a solution

$$v_N = \sum_{n=0}^{N-1} \mathcal{L}_{\mathbb{T}}^n w \simeq v.$$

Since the series in the right hand side of (15) converges exponentially fast, we see that the error we make when approximating v by v_N is exponentially

small in N . Moreover, we see from the definition (9) of the operator $\mathcal{L}_{\mathbb{T}}$ that, in order to compute the value of v_N at a point $x \in M$, one only needs to know the value of w at $x, T^{-1}x, \dots, T^{-N+1}x$, which makes the computation doable.

Remark 2.8. When the solution v to (14) is $\mathcal{C}^{\alpha,1}$, we will sometimes need to compute the derivative $\nabla_{X_u}v$, see Remark 4.3. To do so, one can work as in the proof of Proposition 2.6 in the case $k \geq 1$ below. One can find a lift $\tilde{\mathbb{T}}$ of T on the vector bundle $E \oplus E$ such that

$$(I - \mathcal{L}_{\tilde{\mathbb{T}}})(v \oplus \nabla_{X_u}v) = w \oplus \nabla_{X_u}w. \quad (16)$$

Indeed, working in a trivialization for E , the lift \mathbb{T} becomes a matrix-valued function $\mathbb{T}(x)$ and from (10) and Leibniz formula, we get

$$\begin{aligned} \nabla_{X_u}(\mathcal{L}_{\mathbb{T}}v)(x) &= d_x(\mathcal{L}_{\mathbb{T}}v) \cdot X_u(x) + \Gamma_1(X_u(x), v(T^{-1}x)) \\ &= \mathbb{T}(T^{-1}x)(d_{T^{-1}x}v \cdot d_x T^{-1} \cdot X_u(x)) + \Gamma_2(X_u(x), v(T^{-1}x)) \\ &= h(x)\mathbb{T}(T^{-1}x)(d_{T^{-1}x}v \cdot X_u(T^{-1}x)) + \Gamma_3(X_u(x), v(T^{-1}x)) \\ &= h(x)\mathbb{T}(T^{-1}x)(\nabla_{X_u}v(T^{-1}x)) + \Gamma_4(X_u(x), X_u(T^{-1}x), v(T^{-1}x)), \end{aligned}$$

where $\Gamma_1, \dots, \Gamma_4$ are polynomials with \mathcal{C}^{r-1} coefficients and h is defined by (12). Hence, we can define the lift $\tilde{\mathbb{T}}$ of \mathbb{T} by

$$\tilde{\mathbb{T}}(v_1 \oplus v_2) = \mathbb{T}v_1 \oplus (h(Tx)\mathbb{T}v_2 + \Gamma_4(X_u(Tx), X_u(x), v_1))$$

for $v_1, v_2 \in E_x$.

It follows from (13) that we have $\theta(\tilde{\mathbb{T}}) = \theta(\mathbb{T})$, and hence we can use the same methods to solve approximately (14) and (16). This trick allows us to effectively compute approximate values of the derivative $\nabla_{X_u}v$, where v is a solution of (14).

Example 2.9. Let us illustrate the method from Remarks 2.7 and 2.8 through a very basic example. Assume that T is a CAT map of the two-dimensional torus (a linear example of Anosov diffeomorphism) and that we try to solve for v the equation

$$v - \frac{1}{2}v \circ T^{-1} = w, \quad (17)$$

where w is a smooth function. This is just the equation (14) in the context of Example 2.1 with h identically equal to $1/2$. By a von Neumann series argument, we find a continuous solution v to (17) given by

$$v = \sum_{n \geq 0} 2^{-n}w \circ T^{-n}.$$

We want now to investigate the regularity of v . Let $\lambda_u > 1$ and $\lambda_s = \lambda_u^{-1}$ be the eigenvalues of T . Differentiating in the stable direction, we see that

each term is multiplied by a factor λ_u^n , so that v will not be differentiable (unless λ_u is less than 2). This can be mitigated by choosing $\alpha > 0$ such that $\lambda_u^\alpha < 2$ and noticing that v is α -Hölder. Differentiating in the unstable direction, the situation is much better since each term is multiplied by a factor λ_s^n , so that we can see that v is indeed differentiable in the unstable direction. The proof of Proposition 2.6 is an adaptation of this observation.

Remark 2.10. A natural question regarding the implementation of the S3 algorithm is whether the inevitable computational error will affect the quality of the approximation of the solution to an equation of the form (14). A small error when computing the action of \mathbb{T} on the fiber amounts to replacing the operator $\mathcal{L}_{\mathbb{T}}$ by an operator which is close in the norm topology of operators from $\mathcal{C}^{\alpha,k}(M; E)$ to itself. Thus, numerical errors made at an $n < N$ are not amplified with n in (14).

An error when computing the action of T on a point x is more serious. Indeed, such a perturbation of the dynamics induces a perturbation of the operator $\mathcal{L}_{\mathbb{T}}$ which may not be small in the norm topology of operators from $\mathcal{C}^{\alpha,k}(M; E)$ to itself (the perturbation is small in the strong operator topology but this is not enough to get spectral stability). However, if we see $\mathcal{L}_{\mathbb{T}}$ as an operator from $\mathcal{C}^{\alpha,k}$ to $\mathcal{C}^{\beta,k}$ with $\beta < \alpha$, then the perturbation becomes small. One can then use the machinery developed by Gouëzel, Keller and Liverani in the context of linear response to get some kind of spectral stability [KL99, GL06] (see [Bal18, A.3] for a textbook presentation). In particular, since we are only working away from the spectrum, the resolvent of $\mathcal{L}_{\mathbb{T}}$ is well approximated by the resolvent of its perturbation, and so we can expect that the computational errors do not affect the result of the S3 algorithm too much.

We will prove Proposition 2.6 first in the case $k = 0$.

Proof of Proposition 2.6 in the case $k = 0$. The case $\alpha = 0$ is elementary. It is also a consequence of the much more general statement [CL99, Theorem 8.15]. Let us consequently assume that $0 < \alpha < 1$.

Let $\tilde{\Lambda} > \Lambda$ and $\tilde{\theta} > \theta$. Choose some number $\delta \ll 1$, and let $v \in \mathcal{C}^{\alpha,0}(M; E)$ and $n \in \mathbb{N}$. From the definition of r , we see that there is a constant $C > 0$, that does not depend on n , such that, for every $x \in M$, the operator norm of \mathbb{T}^n from E_x to $E_{T^n x}$ is less than $C\tilde{\theta}^n$. It follows that

$$|\mathcal{L}_{\mathbb{T}}^n v|_\infty \leq C\tilde{\theta}^n |v|_\infty. \quad (18)$$

We want to estimate the semi-norm $|\mathcal{L}_{\mathbb{T}}^n v|_\alpha$. Let $x, y \in M$ be such that $d(x, y) \leq \epsilon$. If $d(x, y) \geq \tilde{\Lambda}^{-n}\delta$, then we have, for some $C > 0$ that may vary from one line to another (but does not depend on n nor v),

$$\begin{aligned} |\mathbb{T}^n v(T^{-n}x) - S_{x,y} \mathbb{T}^n v(T^{-n}y)| &\leq C (|\mathbb{T}^n v(T^{-n}x)| + |\mathbb{T}^n v(T^{-n}y)|) \\ &\leq C\tilde{\theta}^n |v|_\infty \leq C\delta^{-\alpha} (\tilde{\theta}\tilde{\Lambda}^\alpha)^n d(x, y)^\alpha. \end{aligned}$$

Let us now consider the case $d(x, y) \leq \tilde{\Lambda}^{-n}\delta$. Notice that there is a constant $C > 0$ such that the derivative of T^{-n} is bounded by $C\tilde{\Lambda}^n$. Hence we have

$$d(T^{-n}x, T^{-n}y) \leq C\tilde{\Lambda}^n d(x, y) \leq C\delta.$$

In particular, we have $d(T^{-n}x, T^{-n}y) \leq \epsilon$, provided δ is small enough, and we can estimate

$$\begin{aligned} & |\mathbb{T}^n v(T^{-n}x) - S_{x,y} \mathbb{T}^n v(T^{-n}y)| \\ & \leq |\mathbb{T}^n (v(T^{-n}y) - S_{T^{-n}x, T^{-n}y} v(T^{-n}x))| \\ & \quad + |\mathbb{T}^n S_{T^{-n}x, T^{-n}y} v(T^{-n}x) - S_{x,y} \mathbb{T}^n v(T^{-n}x)| \\ & \leq C\tilde{\theta}^n |v|_\alpha d(T^{-n}x, T^{-n}y)^\alpha \\ & \quad + |\mathbb{T}^n S_{T^{-n}x, T^{-n}y} v(T^{-n}x) - S_{x,y} \mathbb{T}^n v(T^{-n}x)| \\ & \leq C |v|_\alpha (\tilde{\theta}\tilde{\Lambda}^\alpha)^n d(x, y)^\alpha + |\mathbb{T}^n S_{T^{-n}x, T^{-n}y} v(T^{-n}x) - S_{x,y} \mathbb{T}^n v(T^{-n}x)|. \end{aligned}$$

We want then to estimate the operator norm of $\mathbb{T}^n S_{T^{-n}x, T^{-n}y} - S_{x,y} \mathbb{T}^n$, which is a linear application from $E_{T^{-n}y}$ to E_x . To do so, we write

$$\begin{aligned} & \mathbb{T}^n S_{T^{-n}x, T^{-n}y} - S_{x,y} \mathbb{T}^n \\ & = \sum_{\ell=0}^{n-1} \mathbb{T}^\ell S_{T^{-\ell}x, T^{-\ell}y} (S_{T^{-\ell}y, T^{-\ell}x} \mathbb{T} S_{T^{-\ell-1}x, T^{-\ell-1}y} - \mathbb{T}) \mathbb{T}^{n-\ell-1}. \end{aligned}$$

In each term, we may bound the operator norm of \mathbb{T}^ℓ by $C\tilde{\theta}^\ell$, the operator norm of $\mathbb{T}^{n-\ell-1}$ by $C\tilde{\theta}^{n-\ell}$ and the operator norm of $S_{T^{-\ell}y, T^{-\ell}x}$ by a constant. Notice then that, for $\ell = 0, \dots, n-1$, the distance between $T^{-\ell-1}x$ and $T^{-\ell-1}y$ is less than $C\tilde{\Lambda}^\ell d(x, y)$, which is less than ϵ , provided δ is small enough. Consequently, it follows from our hypothesis on \mathbb{T} that the operator norm of

$$S_{T^{-\ell}y, T^{-\ell}x} \mathbb{T} S_{T^{-\ell-1}x, T^{-\ell-1}y} - \mathbb{T}$$

is less than $C\tilde{\Lambda}^{\alpha\ell} d(x, y)^\alpha$. The operator norm of $\mathbb{T}^n S_{T^{-n}x, T^{-n}y} - S_{x,y} \mathbb{T}^n$ is consequently less than $C(\tilde{\theta}\tilde{\Lambda}^\alpha)^n d(x, y)^\alpha$ with C that does not depend on n .

Consequently, we find that for some $C > 0$ that does not depend on n nor v , we have

$$|\mathcal{L}_{\mathbb{T}}^n v|_\alpha \leq C(\tilde{\theta}\tilde{\Lambda}^\alpha)^n \|v\|_\alpha.$$

Recalling (18), it follows that the spectral radius of $\mathcal{L}_{\mathbb{T}}$ acting on $\mathcal{C}^\alpha(M; E)$ is less than $\tilde{\theta}\tilde{\Lambda}^\alpha$. Since $\tilde{\theta} > \theta$ and $\tilde{\Lambda} > \Lambda$ are arbitrary, the result follows. \square

Now, using the same trick as in Remark 2.8, we deduce the general case from the case $k = 0$.

Proof of Proposition 2.6 in full generality. The plan is to reduce to the previous case. Let us consider a new vector bundle over M :

$$F = \underbrace{E \oplus \cdots \oplus E}_{k+1 \text{ times}}.$$

We define then the map

$$\begin{aligned} \Phi : \mathcal{C}^{\alpha,k}(M; E) &\rightarrow \mathcal{C}^\alpha(M; F) \\ v &\mapsto v \oplus \nabla_{X_u} v \oplus \cdots \oplus \nabla_{X_u}^k v. \end{aligned}$$

Due to our assumption on X_u , the map Φ induces a linear isomorphism between $\mathcal{C}^{\alpha,k}(M; E)$ and its (closed) image in $\mathcal{C}^\alpha(M; F)$. Working as in Remark 2.8 inductively, we find a \mathcal{C}^α lift $\tilde{\mathbb{T}} : F \rightarrow F$ of T such that $\Phi \circ \mathcal{L}_{\mathbb{T}} = \mathcal{L}_{\tilde{\mathbb{T}}} \circ \Phi$. Moreover, if $x \in M$, then the linear operator from F_x to F_{T_x} induced by $\tilde{\mathbb{T}}$ has the upper block triangular form

$$\begin{pmatrix} \mathbb{T}|_{E_x} & 0 & 0 & 0 \\ * & h\mathbb{T}|_{E_x} & 0 & 0 \\ * & * & \dots & 0 \\ * & * & * & h^k \mathbb{T}|_{E_x} \end{pmatrix}.$$

Notice that in the lower part of this matrix, there may be derivatives of X_u or h along the X_u direction, but of order at most $k-1$, so that our assumption on X_u ensures that $\tilde{\mathbb{T}}$ is α -Hölder. It follows from (13) that $\theta(\tilde{\mathbb{T}}) = \theta(\mathbb{T})$. Consequently, the spectral radius of $\mathcal{L}_{\tilde{\mathbb{T}}}$ acting on $\mathcal{C}^\alpha(M; F)$ is less than $\theta\Lambda^\alpha$. Since Φ is an isomorphism on its image and $\Phi \circ \mathcal{L}_{\mathbb{T}} = \mathcal{L}_{\tilde{\mathbb{T}}} \circ \Phi$, the spectral radius of $\mathcal{L}_{\mathbb{T}}$ is smaller than the one of $\mathcal{L}_{\tilde{\mathbb{T}}}$, hence the result. \square

There is a situation in which Proposition 2.6 will be particularly useful:

Lemma 2.11. *Let $\mathcal{P} : TM \rightarrow TM$ be the map that acts on each fiber by orthogonal projection on the orthogonal of X_u . Then $\mathcal{P}dT : TM \rightarrow TM$ is a lift of T and $\theta(\mathcal{P}dT) < 1$.*

Proof. Let $\pi : TM \rightarrow TM$ acts on each fiber by projection on E_s along E_u . Since π and \mathcal{P} are projectors with the same kernel in each fiber, we have that

$$\mathcal{P}\pi = \mathcal{P} \quad \text{and} \quad \pi\mathcal{P} = \pi.$$

We also have $\pi dT = dT\pi$. Hence, for $n \geq 1$, we have

$$\begin{aligned} (\mathcal{P}dT)^n &= (\mathcal{P}\pi dT)^n = (\mathcal{P}dT\pi)^n \\ &= \mathcal{P} (dT\pi\mathcal{P})^{n-1} dT\pi = \mathcal{P} (dT\pi)^n. \end{aligned}$$

From the hyperbolicity of T , we know that operator norm of $(dT\pi)^n$ in the fibers decays exponentially fast with n , so that we have $\theta(\mathcal{P}dT) < 1$. \square

Lemma 2.12. *Let $k < r - 1$. Then there is $0 < \alpha < 1$ such that we have $X_u \in \mathcal{C}^{\alpha,k}(M; TM)$.*

Proof. We will use Proposition 2.6 to bootstrap the regularity of X_u . The proof is consequently an induction on k , the case $k = 0$ being well-known. Let $1 \leq k < r - 1$ and assume that $X_u \in \mathcal{C}^{\alpha,k-1}(M; TM)$. We will prove that $X_u \in \mathcal{C}^{\alpha_0,k}(M; TM)$ for some $0 < \alpha_0 < 1$.

Since $X_u \in \mathcal{C}^{\alpha,k-1}(M; TM)$ by assumption, we may use any covariant derivative on TM to define the space $\mathcal{C}^{\alpha,k}(M; TM)$. We will use the Levi-Civita connection associated to the Riemannian metric on M . This choice ensures that $\nabla_{X_u} X_u$ is orthogonal to X_u , since X_u is normalized. Notice that $\nabla_{X_u} X_u$ is well-defined and continuous thanks to the unstable manifold theorem [Yoc95, §3.6]. Computing in coordinates with (10), we see that there is a \mathcal{C}^{r-2} section γ of $T^*M \otimes T^*M \otimes M$ such that, for two vector fields X and Y , we have

$$T_*^{-1} \nabla_{T_* X} T_* Y = \nabla_X Y + \gamma(X, Y).$$

Consequently, the equality $X_u = h.T_* X_u$ gives

$$\nabla_{X_u} X_u = \frac{X_u h}{h} X_u + h^2 T_* \nabla_{X_u} X_u + h^2 T_* \gamma(X_u, X_u).$$

Now, let $\mathcal{P} : TM \rightarrow TM$ be as in Lemma 2.11, and notice that:

$$\nabla_{X_u} X_u = h^2 \mathcal{P} T_* \nabla_{X_u} X_u + h^2 \mathcal{P} T_* \gamma(X_u, X_u). \quad (19)$$

Define then a lift $\mathbb{T} : TM \rightarrow TM$ for T by

$$\mathbb{T}v = h(Tx)^2 \mathcal{P} d_x T \cdot v,$$

for $x \in M$ and $v \in T_x M$. It follows from our assumption on X_u that \mathbb{T} is $\mathcal{C}^{\alpha,k-1}$. Moreover, (19) rewrites

$$(I - \mathcal{L}_{\mathbb{T}}) \nabla_{X_u} X_u = h^2 \mathcal{P} T_* \gamma(X_u, X_u). \quad (20)$$

From (13) and Lemma 2.11, we see that $\theta(\mathbb{T}) < 1$.

We can consequently apply Proposition 2.6 to see that $\nabla_{X_u} X_u$ is the unique solution of (20) in $\mathcal{C}^0(M; TM)$. On the other hand, if $0 < \alpha_0 < 1$ is small enough we have $\theta(\mathbb{T}) \Lambda^{\alpha_0} < 1$, and by Proposition 2.6, the equation (20) also has a unique solution v in $\mathcal{C}^{\alpha_0,k-1}(M; TM)$. Here, we use the fact that $r - 1 > k$ and Lemma 2.4 to see that the right-hand side of (20) is in $\mathcal{C}^{\alpha_0,k-1}$, since γ is \mathcal{C}^{r-2} . The space $\mathcal{C}^{\alpha_0,k}(M; TM)$ is smaller than $\mathcal{C}^0(M; TM)$, thus v is nothing else than $\nabla_{X_u} X_u$. Consequently, $\nabla_{X_u} X_u \in \mathcal{C}^{\alpha_0,k-1}(M; TM)$, and thus $X_u \in \mathcal{C}^{\alpha_0,k}(M; TM)$. \square

3 S3 algorithm decomposition

In this section, we explain how the vector fields Y and V are constructed in the second step of the S3 algorithm.

Proposition 3.1. *There is $\alpha_0 > 0$ such that, if $0 \leq \alpha \leq \alpha_0, k < r - 1$ and $X \in \mathcal{C}^{\alpha,k}(M; TM)$, then there are Y and V in $\mathcal{C}^{\alpha,k}(M; TM)$ such that Y is parallel to the unstable direction and*

$$X = Y + V - T_*V.$$

Proof. Let $\mathbb{T} = \mathcal{P}dT$ be the lift of T from Lemma 2.11. It follows from Lemma 2.12 that \mathbb{T} is $\mathcal{C}^{\alpha,k}$ for $\alpha > 0$ small. According to Lemma 2.11, we have $\theta(\mathbb{T}) < 1$.

Let then α_0 be such that $\theta(\mathbb{T})\Lambda^{\alpha_0} < 1$ and $X_u \in \mathcal{C}^{\alpha_0,k}(M; TM)$. Then, if $\alpha \leq \alpha_0$, the spectral radius of \mathcal{L}_T acting on $\mathcal{C}^{\alpha,k}(M; TM)$ is strictly less than 1. Consequently, if $X \in \mathcal{C}^{\alpha,k}(M; TM)$, we can set

$$V = (I - \mathcal{L}_{\mathbb{T}})^{-1}X \quad \text{and} \quad Y = X - (V - T_*V). \quad (21)$$

It follows then from the definition of \mathbb{T} that Y is parallel to the unstable direction. \square

Remark 3.2. Proposition 3.1 implies that there exist vector fields Y and V as in the second step of the S3 algorithm. Moreover, the proof of Proposition 3.1 gives a way to compute Y and V . Indeed, if we look at the expression (21) for V and Y , we see that their values at a point $x \in M$ may be approximated from the knowledge of X and X_u on the backward orbit of x , as explained in Remark 2.7. In addition, the quality of the approximation will be exponentially small in the number of points in the backward orbit of x that we use. Here, we recall that X is a data and that X_u has been computed in the first step of the algorithm.

Remark 3.3. Notice that in Proposition 3.1, even if X is $\mathcal{C}^{k+\alpha}$, then Y and V will *a priori* only be $\mathcal{C}^{\alpha,k}$. This is due to the phenomenon highlighted in Example 2.9 that causes loss of regularity in the stable direction. However, this is not a problem since, even if we cannot work with mere Hölder regularity (because of the use of integration by parts for instance), having a Hölder-continuous derivative in the unstable direction is enough for the arguments from §4 and §5.

4 Unstable contribution

In this section, we give a method to compute the quantity $\Psi_T(Y, f)$, defined by (3), when Y is parallel to the unstable direction, justifying the third step in the S3 algorithm.

Proposition 4.1. *Let $0 \leq \alpha < 1$ and $1 \leq k < r - 1$. Let $Y \in \mathcal{C}^{\alpha,k}(M; TM)$ be parallel to the unstable direction. Then, there is a continuous function $\rho : M \rightarrow \mathbb{C}$ such that if we set $Y^* = -Y + \rho$ then, for every $\mathcal{C}^{1+\alpha}$ functions f, g , we have*

$$\mu(Yf.g) = \mu(f.Y^*g).$$

Proof. Cover M by open sets $(U_\omega)_{\omega \in \Omega}$ foliated by local unstable manifolds. Then, let $(\varphi_\omega)_{\omega \in \Omega}$ be a partition of unity subordinated to the open cover $(U_\omega)_{\omega \in \Omega}$. We can then split

$$\mu(Yf.g) = \sum_{\omega \in \Omega} \mu(Yf.\varphi_\omega g)$$

and consider each term separately. Fix then $\omega \in \Omega$ and let W_ω denote a transverse to the unstable foliation in U_ω . We can then disintegrate the restriction of μ to U_ω and write

$$\mu(Yf.\varphi_\omega g) = \int_{W_\omega} \left(\int_{W_{u,\text{loc}}(y)} Yf(x).\varphi_\omega(x)g(x)dm_y(x) \right) d\nu_\omega(y),$$

where ν_ω is a measure on W_ω and, for every $y \in W_\omega$, the measure m_y is absolutely continuous and fully supported on the local unstable manifold $W_{u,\text{loc}}(y)$. Consequently, for $y \in W_\omega$, we can integrate by parts in the integral on $W_{u,\text{loc}}(y)$ to find

$$\begin{aligned} & \int_{W_{u,\text{loc}}(y)} Yf(x).\varphi_\omega(x)g(x)dm_y(x) \\ &= \int_{W_{u,\text{loc}}(y)} f(x) (-\varphi_\omega(x)Yg(x) + \rho_\omega(x)g(x)) dm_y(x), \end{aligned}$$

for some bounded measurable function ρ_ω . Here, the integration by part is licit since the measure dm_y has a differentiable density, and there is no boundary term since φ_ω vanishes near the boundary of the local unstable manifold. Moreover, using the explicit formula for the density of dm_y given for instance in [LY85, Corollary 6.1.4], we see that ρ_ω is actually continuous. Summing over ω , the lemma is proven. \square

Lemma 4.2. *Under the hypotheses of Proposition 4.1, assume in addition that $X_u \in \mathcal{C}^{\alpha,k}(M; TM)$ and write $Y = aX_u$ with $a \in \mathcal{C}^{\alpha,k}$. Then, the function ρ from Proposition 4.1 is given by the formula*

$$\rho = a\rho_0 - X_u a,$$

where ρ_0 is given by

$$\rho_0 = -(I - \mathcal{L}_h)^{-1} \left(\frac{X_u h}{h} \right),$$

where h is defined by (12). In particular, if $\beta > 0$ is small enough then ρ is $\mathcal{C}^{\beta,k-1}$.

Proof. Notice first that, by taking $g = 1$ in Proposition 4.1, we see that for every f smooth enough we have

$$\mu(f\rho) = \mu(Yf),$$

and this relation completely characterizes the function ρ , since μ is fully supported.

1. Let us start with the case $Y = X_u$, in which we write $\rho_0 = \rho$. It follows from the definition of h that for smooth f , we have

$$h \circ T.X_u(f \circ T) = X_u f \circ T,$$

and that h does not vanish on M . Hence, using that μ is T -invariant we find that for smooth f we have

$$\begin{aligned} \mu(f\rho_0) &= \mu(X_u f) = \mu(X_u f \circ T) = \mu(h \circ T.X_u(f \circ T)) \\ &= \mu(f \circ T.(-X_u(h \circ T) + \rho_0 h \circ T)) \\ &= \mu(f.(-X_u(h \circ T) + \rho_0 h \circ T) \circ T^{-1}). \end{aligned}$$

Thus, we find that

$$\rho_0 = -X_u(h \circ T) \circ T^{-1} + \rho_0 \circ T^{-1}h = -\frac{X_u h}{h} + \rho_0 \circ T^{-1}h.$$

Now, notice that it follows from (13), Proposition 2.6 and Lemma 2.12 that the operator \mathcal{L}_h has spectral radius strictly less than 1 on $\mathcal{C}^{\beta, k-1}$, if β is small enough, so that we can write

$$\rho_0 = -(I - \mathcal{L}_h)^{-1} \left(\frac{X_u h}{h} \right).$$

Indeed ρ_0 is the unique continuous solution of the equation $(I - \mathcal{L}_h)\rho_0 = -X_u h/h$.

2. Let us now deal with the general case, where $Y = aX_u$. Then, we have for smooth f

$$\mu(f\rho) = \mu(Yf) = \mu(X_u f.a) = \mu(f.(-X_u a + a\rho_0)),$$

and the result follows. \square

Remark 4.3. As explained in the introduction, the knowledge of ρ allows us to approximate the contribution $\Psi_T(Y, f)$ of Y to the linear response using the formula (5). Notice that the characterization of ρ given in Lemma 4.2 enables to use the method explained in Remark 2.7 to compute ρ from the knowledge of $h, X_u h$ and $X_u a$ (or equivalently $\nabla_{X_u} Y$).

All these quantities are deduced from previous computations. The value of h is deduced from X_u by its definition $h = \langle T_* X_u, X_u \rangle^{-1}$. By differentiation of this equality, we deduce the derivative $X_u h$ from $\nabla_{X_u} X_u$ (that

can be computed using the method from Remark 2.7 and the formula (20)). Finally, we explained in Remark 2.8 how to compute the derivative $\nabla_{X_u} Y$ using the equation (21).

As usual, the value of ρ at x is obtained with a precision which is exponential in the number of points in the backward orbit of x that we use for the computation.

Remark 4.4. Notice that if W is a local unstable manifold for T and Υ denotes the density (with respect to the arclength) of the conditional of μ on W (which is well-defined up to a multiplicative constant), then one has $\rho_0 = -X_u \log \Upsilon$.

5 Coboundary contribution

In order to end the proof of Theorem 2, we want now to evaluate the contribution to $\Psi_T(f, X)$ of the coboundary term in the decomposition (4). To do so, we use the following result.

Lemma 5.1. *Let $0 < \alpha < 1$, $V \in \mathcal{C}^{\alpha,1}(M; TM)$ and f be a $\mathcal{C}^{1+\alpha}$ function on M . Then the series defining $\Psi_T(f, V - T_*V)$ converges and*

$$\Psi_T(f, V - T_*V) = \mu(Vf).$$

Lemma 5.1 follows from the estimate:

Lemma 5.2. *Let $\alpha > 0$, $V \in \mathcal{C}^{\alpha,1}(M; TM)$ and $f \in \mathcal{C}^{1+\alpha}$. Then the integral $\mu(V(f \circ T^n))$ tends to 0 (exponentially fast) as n tends to $+\infty$.*

Proof of Lemma 5.1. Just notice that for $n \in \mathbb{N}$, we have

$$\begin{aligned} \mu((V - T_*V)(f \circ T^n)) &= \mu(V(f \circ T^n)) - \mu(V(f \circ T^{n+1}) \circ T^{-1}) \\ &= \mu(V(f \circ T^n)) - \mu(V(f \circ T^{n+1})). \end{aligned}$$

Hence, $\Psi_T(f, V - T_*V)$ becomes a telescopic sum, and we get thanks to Lemma 5.1:

$$\Psi_T(f, V - T_*V) = \mu(Vf).$$

□

The rest of this section is dedicated to the proof of Lemma 5.2. We will use the machinery from [GL06, GL08]. Letting $0 < q < \min(\alpha, r - 2)$, we recall the space $\mathcal{B}^{1,q}$ from [GL06, GL08]. For convenience, we will reverse time with respect to [GL06, GL08] and consider the Koopman operator

$$\mathcal{K} : u \mapsto u \circ T \tag{22}$$

instead of the transfer operator. In particular, we will use admissible unstable leaves instead of admissible stable leaves to define the space $\mathcal{B}^{1,q}$. On

a technical level, it amounts to replace T by T^{-1} and consider a weighted transfer operator. Very general weighted transfer operators are considered in [GL08], and we could use the results from this reference. However, since we do not need all the subtleties from [GL08], we rather rely on the simpler exposition from [GL06], letting the reader check that the results from [GL06] also applies to the Koopman operator (22) with the required modifications.

The space $\mathcal{B}^{1,q}$ is defined as the completion of $\mathcal{C}^{1+\alpha}$ with respect to the norm

$$\|u\|_{\text{GL}} := \max \left(|u|_{0,q}, |u|_{1,1+q} \right),$$

where

$$|u|_{0,q} = \sup_{W \in \Sigma} \sup_{\substack{\varphi \in \mathcal{C}_0^q(W, \mathbb{R}) \\ |\varphi|_{\mathcal{C}^q} \leq 1}} \int_W u \varphi dm_W$$

and

$$|u|_{1,1+q} = \sup_{W \in \Sigma} \sup_{\substack{v \in \mathcal{V}^{1+\alpha}(W) \\ |v|_{\mathcal{C}^{1+\alpha}} \leq 1}} \sup_{\substack{\varphi \in \mathcal{C}_0^{1+q}(W, \mathbb{R}) \\ |\varphi|_{\mathcal{C}^{1+q}} \leq 1}} \int_W (vu) \varphi dm_W.$$

Here, Σ denotes a set of admissible unstable leaves (that is a set of admissible stable leaves for T^{-1} in the words of [GL06]). We will only need to know that the elements of Σ are small curves in M and that there is $\delta > 0$ such that any piece of unstable manifolds of size δ is an element of Σ . For $W \in \Sigma$, we write dm_W for the Lebesgue measure on W , we denote by $\mathcal{V}^{1+\alpha}(W)$ the set of $\mathcal{C}^{1+\alpha}$ vector fields on a neighbourhood of W and by $\mathcal{C}_0^q(W)$ the space of \mathcal{C}^q functions on W supported away from the boundary. Notice that we recall this definition of the space $\mathcal{B}^{1,q}$ mostly for educational purpose, as we will only need the local description of the space $\mathcal{B}^{1,q}$ given by [GL06, Lemma 3.2].

From [GL06, Theorem 2.3], or rather its adaption to the case of Koopman operator instead of transfer operator, we know that the essential spectral radius of \mathcal{K} acting on $\mathcal{B}^{1,q}$ is strictly less than 1, and that 1 is a simple eigenvalue for \mathcal{K} . Moreover, this is the only eigenvalue of \mathcal{K} on the unit circle, the associated right eigenspace consists of constant functions, while the left eigenspace is generated by the SRB measure μ of T . Consequently, we may decompose \mathcal{K} as

$$\mathcal{K} = P + Q, \tag{23}$$

where P is the rank 1 operator defined by $P(u) = \mu(u) \mathbb{1}$ and Q is a bounded operator on $\mathcal{B}^{1,q}$ with spectral radius strictly less than 1 that satisfies $PQ = QP = 0$. Consequently, for $n \geq 1$, we have

$$\mathcal{K}^n = P + Q^n. \tag{24}$$

With these tools from [GL06], Lemma 5.2 follows from:

Lemma 5.3. *Let $\alpha > 0$ and $V \in \mathcal{C}^{\alpha,1}(M; TM)$. Then the map*

$$f \mapsto \mu(Vf)$$

extends to a bounded linear form on $\mathcal{B}^{1,q}$.

Let us explain how Lemma 5.3 allows us to prove Lemma 5.2.

Proof of Lemma 5.2. Since $f \in \mathcal{C}^{1+\alpha}$, we see that f belongs to $\mathcal{B}^{1,q}$. Consequently, we have for $n \in \mathbb{N}$

$$f \circ T^n = \mathcal{K}^n f = \mu(f) \mathbb{1} + Q^n f$$

and thus

$$\mu(V(f \circ T^n)) = \mu(V(f \circ T^n - \mu(f) \mathbb{1})) = \mu(VQ^n f).$$

Here, the expression $\mu(VQ^n f)$ makes sense due to Lemma 5.3, and we have for some $C > 0$ that does not depend on n :

$$|\mu(VQ^n f)| \leq C \|Q^n f\|_{\text{GL}}.$$

However, since the spectral radius of Q is strictly less than 1, we see that $\|Q^n f\|_{\text{GL}}$ decays exponentially fast when n tends to $+\infty$. \square

We need now to prove Lemma 5.3.

Proof of Lemma 5.3. Let $f \in \mathcal{C}^{1+\alpha}$. Let χ be a \mathcal{C}^∞ function on M with small support. We will estimate $\mu(\chi Vf)$ using the $\mathcal{B}^{1,q}$ norm of f under the assumption that the support of χ is small enough. The result then follows by a partition of unity argument.

By assuming that the support of χ is small enough, we may work in coordinates. In these coordinates, we can write

$$V = \sum_{j=1}^d v_j \frac{\partial}{\partial x_j}.$$

Here, the v_j 's are $\mathcal{C}^{\alpha,1}$. In particular, the restriction of the v_j 's to unstable leaves are $\mathcal{C}^{1+\alpha}$ with uniform $\mathcal{C}^{1+\alpha}$ estimates. Then, up to taking the support of χ smaller, we can cover the patch of coordinates we are working in by a family $(W_z)_{z \in Z}$ of local (exact) unstable leaves of uniform size. We endow Z with a measurable structure by identifying it with a manifold transverse to the unstable direction. Moreover, we may assume that the W_z 's belong to Σ and that each W_z fully intersects that support of χ (i.e. the support of the restriction of χ to W_z does not intersect the boundary of W_z). Then, we

disintegrate μ (restricted to the coordinate patch) with respect to the W_z 's and find that, for $j = 1, \dots, d$, we have

$$\mu \left(\chi v_j \frac{\partial f}{\partial x_j} \right) = \int_Z \left(\int_{W_z} \chi v_j \frac{\partial f}{\partial x_j} \rho_z dm_{W_z} \right) d\nu(z),$$

where ν is a measure with finite mass on Z and the ρ_z 's are uniformly \mathcal{C}^{r-1} (and hence \mathcal{C}^{1+q}). Hence, we see that, for $z \in Z$, the function $\chi v_j \rho_z$ on W_z is \mathcal{C}^{1+q} and supported away from the boundary of W_z , so that we have by [GL06, Lemma 3.2]

$$\left| \int_{W_z} \chi v_j \frac{\partial f}{\partial x_j} \rho_z dm_{W_z} \right| \leq C \|f\|_{\text{GL}},$$

for some constant $C > 0$ which is uniform in $z \in Z$ (because all data are). Integrating this estimate with respect to ν , we find that

$$\left| \mu \left(\chi v_j \frac{\partial f}{\partial x_j} \right) \right| \leq C \|f\|_{\text{GL}}.$$

Summing over j , we find that $f \mapsto \mu(\chi V f)$ extends to a continuous linear form on $\mathcal{B}^{1,q}$, and then the lemma follows by a partition of unity argument. \square

Remark 5.4. The two main elements in the proof of Lemma 5.2 are Lemma 5.3 and the spectral decomposition (23) of the Koopman operator, valid on the space $\mathcal{B}^{1,q}$. Such a decomposition for the Koopman operator has been established on many other spaces, see for instance [BT07, BT08, FRS08]. One can consequently wonder if Lemma 5.3 also holds on these spaces (it would give an alternative proof of Lemma 5.2). It seems that the answer is positive for the spaces from [BT07], but the proof is much more technically involved. The advantage of the space $\mathcal{B}^{1,q}$ from [GL06] here is that it is easy to relate the linear form from Lemma 5.3 with the definition of these spaces.

6 Computing integrals with respect to μ

In order to achieve Step 4 of the S3 algorithm, we need to find a way to approximate the integral $\mu(g)$ of a Hölder observable g against the SRB measure μ for T . We propose two simple methods to do so: a probabilistic one that relies on the physicality of μ (this is the method that is used in [CW21]) and a deterministic one that uses a classical construction of the SRB measure. One could also use more advanced techniques to compute integrals against μ , based on approximations of transfer operators. See for instance the approach from [CF20] based on Fourier decomposition.

6.1 Probabilistic method

We describe here a ‘‘Monte Carlo-like’’ method to approximate integrals against μ . This approach is quite standard (see for instance [GWW16] for another application of this method). In view of the property (1) of the measure μ , it seems natural to approximate $\mu(g)$ by

$$\mu(g) \simeq \frac{1}{N} \sum_{k=0}^{N-1} g(T^k x), \quad (25)$$

where x is a point in M and N a large integer. This strategy works for Lebesgue almost every choice of x . The error in this approximation can be bounded almost surely using the following corollary of the almost sure invariance principle. We expect that non-asymptotic bound with high probability could be derived using a concentration inequality in the spirit of [CG12].

Lemma 6.1. *Let g be a Hölder-continuous function on M . Then, for Lebesgue almost every $x \in M$, there is a constant C such that, for $N \geq 3$, we have*

$$\left| \frac{1}{N} \sum_{k=0}^{N-1} g(T^k x) - \mu(g) \right| \leq C \sqrt{\frac{\log \log N}{N}}.$$

Proof. Replacing g by $g - \mu(g)$, we can assume that $\mu(g) = 0$. If M is endowed with any smooth measure, then the sequence of functions $(g, g \circ T, g \circ T^2, \dots)$ becomes a stochastic process and it follows from [Gou10, Theorem 2.1], using the functional analytic tools from [GL06, GL08] or [BT07, BT08] as explained in [Gou10], that this stochastic process satisfies an almost sure principle. Consequently, there is a probability space Ω , a process (A_0, A_1, \dots) on Ω with the same law as $(g, g \circ T, \dots)$ and a Brownian motion $(B_t)_{t \geq 0}$ on Ω such that, almost surely,

$$\left| \sum_{k=0}^{N-1} A_k - B_N \right| \underset{N \rightarrow +\infty}{=} o(N^{\frac{1}{2}}).$$

From the law of the iterated logarithm, we know that almost surely there is a constant C such that

$$|B_N| \leq C \sqrt{N \log \log N},$$

so that

$$\left| \frac{1}{N} \sum_{k=0}^{N-1} A_k \right| \leq C' \sqrt{\frac{\log \log N}{N}},$$

for some constant $C' > C$ that depends on the alea. Since (A_0, A_1, \dots) has the same law as $(g, g \circ T, \dots)$, the result follows. \square

Remark 6.2. Let us go back to the S3 algorithm. If we use (7) as an approximation for $\Psi_T(X, f)$, the integrals being computed by the probabilistic method we just described, then we make three kinds of error:

- if we use orbits of length n to compute approximate values of ρ and V , then we make an error of size $\mathcal{O}(\eta^n)$, for some $\eta < 1$, when approximating them (see Remarks 2.7, 3.2 and 4.3);
- the truncation after the L th term in the series in the right hand side of (7) produces an error of size $\mathcal{O}(\eta^L)$;
- the errors when approximating the integrals are almost surely of size $\mathcal{O}((\log \log N/N)^{1/2})$ by Lemma 6.1.

This suggest to take n and L of the order of $C \log N$ with $C \gg 1$ in order to get an error which is $\mathcal{O}(N^{-\frac{1}{2}-\epsilon})$ for any $\epsilon > 0$. The N points that we use in the approximation (25) lie on the same orbits, so that this computation requires to compute only one orbit of length $n + N$, rather than N orbits of length n . We need to compute $1 + L$ such integrals, so that the total computation time τ is proportional to $L(n + N) = \mathcal{O}(N \log N)$, and we find as announced that the error made by the S3 algorithm is $\mathcal{O}(\tau^{-\frac{1}{2}+\epsilon})$.

6.2 Deterministic method

The integral $\mu(g)$ may also be approximated by a deterministic method. The advantage is that we do not rely on chance to get an accurate approximation. However, the convergence will be slower in general. Choose W a one-dimensional submanifold of M whose tangent line never aligns with the stable direction of T . For simplicity, we assume that W belongs to Σ , the space of admissible unstable manifolds used to define the space $\mathcal{B}^{1,q}$ from [GL06] in §5. Choose then a smooth function $\rho : W \rightarrow \mathbb{R}_+$ supported away from the boundary of W and such that

$$\int_W \rho dm_W = 1.$$

Then, if n is a very large integer, we can approximate $\mu(g)$ by

$$\mu(g) \simeq \int_W g \circ T^n \rho dm_W.$$

Indeed, the linear form

$$u \mapsto \int_W u \rho dm_W$$

is bounded on $\mathcal{B}^{1,q}$ (by the very definition of the space), and recalling the decomposition (24) for the Koopman operator we find that

$$\int_W g \circ T^n \rho dm_W = \mu(g) + \int_W Q^n u \rho dm_W.$$

Here, the last term decays exponentially fast with n since the spectral radius of Q is strictly less than 1.

The integral $\int_W g \circ T^n \rho dm_W$ can be approximated by Riemann sums. In the context of the S3 algorithm, the observable g is only Hölder, so that it does not make sense to try to use a more sophisticated method to compute the integral. Moreover, when n tends to $+\infty$ the Hölder constant of $g \circ T^n$ grows exponentially fast, so that even the approximation by Riemann sum must be dealt with carefully. Let $RS_{N,n}(g)$ denote the approximation of $\int_W g \circ T^n \rho dm_W$ by a Riemann sum with N terms. If g is α -Hölder, then $g \circ T^n$ is α -Hölder with a Hölder constant $\mathcal{O}(\kappa^{\alpha n})$, where $\kappa > 1$ denotes the maximal expansion rate of T . Hence, we have

$$\begin{aligned} & |RS_{N,n}(g) - \mu(g)| \\ & \leq \left| RS_{N,n}(g) - \int_W g \circ T^n \rho dm_W \right| + \left| \int_W g \circ T^n \rho dm_W - \mu(g) \right| \\ & \lesssim N^{-\alpha} \kappa^{\alpha n} + \eta^n, \end{aligned} \quad (26)$$

with $\eta < 1$. We see that the best approximation here is obtained by taking $n \simeq \frac{\alpha}{\log(\kappa^\alpha/\eta)} \log N$. With this value of n , the error when approximating $\mu(g)$ by $RS_{N,n}(g)$ is $\mathcal{O}(N^{-\beta})$ with $\beta = -\frac{\alpha \log \eta}{\log(\kappa^\alpha/\eta)}$.

Remark 6.3. If g is one of the observables for which we need to compute $\mu(g)$ in the fourth step of the S3 algorithm, then the value of g at a point may be computed with an error $\mathcal{O}(\eta^n)$ in a time proportional to n (one only needs to use an orbit of length n , see Remarks 2.7, 3.2 and 4.3). Hence, this error may be considered as part of the exponentially decaying terms in (26). Computing $RS_{N,n}(g)$ requires the value of g at N points, hence the computation time τ_0 of one such integral will be proportional to $nN = \mathcal{O}(N \log N)$. Consequently, we compute one integral with a precision $\mathcal{O}(\tau_0^{-\gamma})$, for some $\gamma > 0$, in a time τ_0 .

We also need to take into account the number of integrals that we want to compute: taking L proportional to $\log \tau_0$ in (5) gives an approximation of $\Psi_T(Y, f)$ with a precision $\mathcal{O}(\tau_0^{-\gamma})$. The total computation time τ will consequently be bounded by $\tau_0 \log \tau_0$. Up to taking $\gamma > 0$ smaller, we see that the error made by the S3 algorithm when estimating $\Psi_T(X, f)$ after a computation of time τ is $\mathcal{O}(\tau^{-\gamma})$.

Remark 6.4. Instead of using a piece of unstable manifold W , one could integrate directly against a smooth measure on the manifold M . The convergence also holds, but the quality of the approximation of an integral by Riemann sums deteriorates when the dimension gets bigger. Hence, we still get a polynomial precision, but *a priori* with a worst exponent.

References

- [AM07] Rafail V. Abramov and Andrew J. Majda. Blended response algorithms for linear fluctuation-dissipation for complex nonlinear dynamical systems. *Nonlinearity*, 20(12):2793–2821, 2007.
- [Bal14] Viviane Baladi. Linear response, or else. In *Proceedings of the International Congress of Mathematicians (ICM 2014), Seoul, Korea, August 13–21, 2014. Vol. III: Invited Lectures*, pages 525–545. KM Kyung Moon Sa, Seoul, 2014.
- [Bal18] Viviane Baladi. *Dynamical Zeta Functions and Dynamical Determinants for Hyperbolic Maps*, volume 68 of *Ergebnisse*. Springer, 2018.
- [BCV16] Thiago Bomfim, Armando Castro, and Paulo Varandas. Differentiability of thermodynamical quantities in non-uniformly expanding dynamics. *Advances in Mathematics*, 292:478–528, 2016.
- [BGNN17] Wael Bahsoun, Stefano Galatolo, Isaia Nisoli, and Xiaolong Niu. A Rigorous Computational Approach to Linear Response. *arXiv:1506.08661*, 2017.
- [BLL20] Tamás Bódai, Valerio Lucarini, and Frank Lunkeit. Can we use linear response theory to assess geoengineering strategies? *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 30(2):023124, 2020.
- [BS12] Viviane Baladi and Daniel Smania. Linear response for smooth deformations of generic nonuniformly hyperbolic unimodal maps. *Annales scientifiques de l'ENS*, 45(6):861–926, 2012.
- [BS16] Wael Bahsoun and Benoît Saussol. Linear response in the intermittent family: Differentiation in a weighted c^0 -norm. *Discrete & Continuous Dynamical Systems*, 36(12):6657–6668, 2016.
- [BT07] Viviane Baladi and Masato Tsujii. Anisotropic Hölder and Sobolev spaces for hyperbolic diffeomorphisms. *Ann. Inst. Fourier (Grenoble)*, 57(1):127–154, 2007.
- [BT08] Viviane Baladi and Masato Tsujii. Dynamical determinants and spectrum for hyperbolic diffeomorphisms. *Geometric and probabilistic structures in dynamics*, Amer. Math. Soc., Providence, RI(469):29–68, 2008.

- [BT16] Viviane Baladi and Mike Todd. Linear response for intermittent maps. *Commun. Math. Phys.*, 347:857–874, 2016.
- [Ces19] Bruno Cessac. Linear response in neuronal networks: From neurons dynamics to collective response. *Chaos*, 29:103105, 2019.
- [CF20] Harry Crimmins and Gary Froyland. Fourier approximation of the statistical properties of Anosov maps on tori. *Nonlinearity*, 33(11):6244–6296, 2020.
- [CFTW19] Nisha Chandramoorthy, Pablo Fernandez, Chaitanya Talnikar, and Qiqi Wang. Feasibility analysis of ensemble sensitivity computation in turbulent flows. *AIAA Journal*, 57(10):4514–4526, 2019.
- [CG12] Jean-René Chazottes and Sébastien Gouëzel. Optimal Concentration Inequalities for Dynamical Systems. *Communications in Mathematical Physics*, 316(3):843–889, 2012.
- [CL99] Carmen Chicone and Yuri Latushkin. *Evolution Semigroups in Dynamical Systems and Differential Equations*, volume 70 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, Rhode Island, 1999.
- [CW20] Nisha Chandramoorthy and Qiqi Wang. A computable realization of Ruelle’s formula for linear response of statistics in chaotic systems. *arXiv:2002.04117*, 2020.
- [CW21] Nisha Chandramoorthy and Qiqi Wang. Efficient computation of linear response of chaotic attractors with one-dimensional unstable manifolds. *arXiv:2103.08816*, 2021.
- [Den89] Manfred Denker. *The Central Limit Theorem for Dynamical Systems*. 1989. Dynamical systems and ergodic theory, 28th Sem. St. Banach Int. Math. Cent., Warsaw/Pol. 1986, Banach Cent. Publ. 23, 33-62 (1989).
- [DIX19] Karthik Duraisamy, Gianluca Iaccarino, and Heng Xiao. Turbulence modeling in the age of data. *Annual Review of Fluid Mechanics*, 51(1):357–377, 2019.
- [DLL01] Rafael De La Llave. Remarks on Sobolev regularity in Anosov systems. *Ergodic Theory and Dynamical Systems*, 21(04), 2001.
- [Dol04] Dmitry Dolgopyat. On differentiability of SRB states for partially hyperbolic systems. *Inventiones Mathematicae*, 155(2):389–449, 2004.

- [EHL04] Gregory Eyink, Thomas Haine, and D. J. Lea. Ruelle's linear response formula, ensemble adjoint schemes and Lévy flights. *Nonlinearity*, 17(5):1867–1889, 2004.
- [FRS08] Frédéric Faure, Nicolas Roy, and Johannes Sjöstrand. Semi-classical approach for Anosov diffeomorphisms and Ruelle resonances. *Open Mathematics Journal*, 1:35–81, 2008.
- [GG19] Stefano Galatolo and Paolo Giulietti. A linear response for dynamical systems with additive noise. *Nonlinearity*, 32:2269–2301, 2019.
- [GL06] Sebastien Gouëzel and Carlangelo Liverani. Banach spaces adapted to Anosov systems. *Ergod. Th. and Dyn. Sys.*, 26:189–217, 2006.
- [GL08] Sébastien Gouëzel and Carlangelo Liverani. Compact locally maximal hyperbolic sets for smooth maps: Fine statistical properties. *J. Differential Geom.*, 79(3):433–477, 2008.
- [Gou10] Sébastien Gouëzel. Almost sure invariance principle for dynamical systems by spectral methods. *The Annals of Probability*, 38(4):1639–1671, 2010.
- [GWW16] Georg A. Gottwald, J.P. Wormell, and Jeroen Wouters. On spurious detection of linear response and misuse of the fluctuation–dissipation theorem in finite time series. *Physica D: Nonlinear Phenomena*, 331:89–101, 2016.
- [HM10] Martin Hairer and Andrew J. Majda. A simple framework to justify linear response theory. *Nonlinearity*, 23(4):909–922, 2010.
- [HM20] Francisco Huhn and Luca Magri. Stability, sensitivity and optimisation of chaotic acoustic oscillations. *Journal of Fluid Mechanics*, 882:A24, 2020.
- [HP70] Morris W. Hirsch and Charles C. Pugh. Stable manifolds and hyperbolic sets. In *Global Analysis (Proc. Sympos. Pure Math., Vol. XIV, Berkeley, Calif., 1968)*, pages 133–163. Amer. Math. Soc., Providence, R.I., 1970.
- [Jia12] Miaohua Jiang. Differentiating potential functions of srb measures on hyperbolic attractors. *Ergodic Theory and Dynamical Systems*, 32(4):1350–1369, 2012.
- [JS18] Matthew P. Juniper and R.I. Sujith. Sensitivity and nonlinearity of thermoacoustic oscillations. *Annual Review of Fluid Mechanics*, 50(1):661–689, 2018.

- [KH95] Anatole Katok and Boris Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*, volume 54 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, Cambridge, 1995.
- [KL99] Gerhard Keller and Carlangelo Liverani. Stability of the spectrum for transfer operators. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, 28(1):141–152, 1999.
- [LBH⁺14] Valerio Lucarini, Richard Blender, Corentin Herbert, Salvatore Pascale, Francesco Ragone, and Jeroen Wouters. Mathematical and physical ideas for climate science. *Rev. Geophys.*, 52:809–859, 2014.
- [LHAH02] Daniel J. Lea, Thomas W. N. Haine, Myles R. Allen, and James A. Hansen. Sensitivity analysis of the climate of a chaotic ocean circulation model. *Quarterly Journal of the Royal Meteorological Society*, 128(586):2587–2605, 2002.
- [Liv95] Carlangelo Liverani. Decay of correlations. *Annals of Mathematics*, 142(2):239–301, 1995.
- [LY85] Francois Ledrappier and Lai-Sang Young. The Metric Entropy of Diffeomorphisms: Part I: Characterization of Measures Satisfying Pesin’s Entropy Formula. *The Annals of Mathematics*, 122(3):509, 1985.
- [Ni19] Angxiu Ni. Hyperbolicity, shadowing directions and sensitivity analysis of a turbulent three-dimensional flow. *Journal of Fluid Mechanics*, 863:644–669, 2019.
- [Ni20] Angxiu Ni. Linear response algorithm for differentiating stationary measures of chaos. *arXiv:2009.00595*, 2020.
- [Ni21] Angxiu Ni. Fast linear response algorithm for differentiating stationary measures of chaos. *arXiv:2009.00595*, 2021.
- [NW17] Angxiu Ni and Qiqi Wang. Sensitivity analysis on chaotic dynamical systems by non-intrusive least squares shadowing (nilss). *Journal of Computational Physics*, 347:56–77, 2017.
- [OP89] Steven Orey and Stephan Pelikan. Deviations of trajectory averages and the defect in Pesin’s formula for Anosov diffeomorphisms. *Trans. Am. Math. Soc.*, 315(2):741–753, 1989.
- [RLL16] Francesco Ragone, Valerio Lucarini, and Frank Lunkeit. A new framework for climate sensitivity and prediction: a modelling perspective. *Climate Dynamics*, 46:1459–1471, 2016.

- [Rue97] David Ruelle. Differentiation of SRB States. *Communications in Mathematical Physics*, 187(1):227–241, 1997.
- [Rue03] David Ruelle. Differentiation of SRB states: Correction and complements. *Communications in Mathematical Physics*, 234(1):185–190, 2003.
- [SBH⁺16] D. Stammer, M. Balmaseda, P. Heimbach, A. Köhl, and A. Weaver. Ocean data assimilation in support of climate applications: Status and perspectives. *Annual Review of Marine Science*, 8(1):491–518, 2016. PMID: 26473335.
- [Sin68] Yakov G. Sinai. Markov partitions and C^∞ -diffeomorphisms. *Functional Analysis and its Applications*, 2:61–82, 1968.
- [Sin73] Yakov G. Sinai. Gibbs measures in ergodic theory. *Russ. Math. Surv.*, 27(4):21–69, 1973.
- [WG18] Caroline L. Wormell and Georg A. Gottwald. On the validity of linear response theory in high-dimensional deterministic dynamical systems. *J Stat Phys*, 172:1479—1498, 2018.
- [WG19] Caroline L. Wormell and Georg A. Gottwald. Linear response for macroscopic observables in high-dimensional systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 29(11):113127, 2019.
- [Yoc95] Jean-Christophe Yoccoz. Introduction to hyperbolic dynamics. In *Real and Complex Dynamical Systems. Proceedings of the NATO Advanced Study Institute Held in Hillerød, Denmark, June 20-July 2, 1993*, pages 265–291. Kluwer Academic Publishers, Dordrecht, 1995.
- [You02] Lai-Sang Young. What are SRB measures, and which dynamical systems have them? In *Journal of Statistical Physics*, volume 108, pages 733–754. 2002.