

# APPROXIMATE FACTOR MODELS WITH WEAKER LOADINGS

Jushan Bai\*      Serena Ng†

December 6, 2021

## Abstract

Pervasive cross-section dependence is increasingly recognized as an appropriate characteristic of economic data and the approximate factor model provides a useful framework for analysis. Assuming a strong factor structure, early work established convergence of the principal component estimates of the factors and loadings to a rotation matrix. This paper shows that the estimates are still consistent and asymptotically normal for a broad range of weaker factor loadings, albeit at slower rates and under additional assumptions on the sample size. Standard inference procedures can be used except in the case of extremely weak loadings which has encouraging implications for empirical work. The simplified proofs are of independent interest.

JEL Classification: C30, C31

Keywords: asymptotic principal components, low rank decomposition. factor augmented regressions.

---

\*Columbia University, 420 W. 118 St. MC 3308, New York, NY 10027. Email: jb3064@columbia.edu

†Columbia University and NBER, 420 W. 118 St. MC 3308, New York, NY 10027. Email: serena.ng@columbia.edu. This work is supported by the National Science Foundation SES-2018369 (Ng).

# 1 Introduction

Starting with Forni et al. (2000) and Stock and Watson (1998, 2002), a large body of research has been developed to estimate the latent common variations in large panels in which the  $N$  units observed over  $T$  periods are cross-sectionally correlated. A fundamental result shown in Bai and Ng (2002) is that the space spanned by the factors can be consistently estimated by the method of static principal components (APC) at rate  $\min(\sqrt{N}, \sqrt{T})$ . Bai (2003) then establishes  $\sqrt{N}$  asymptotic normality of the estimated factors  $\tilde{\mathbf{F}}$  up to a rotation matrix  $\mathbf{H}$ . The maintained assumption is that the factor structure is strong, meaning that if  $\mathbf{F}$  and  $\mathbf{\Lambda}$  are the latent factors and loadings, the matrices  $\mathbf{F}'\mathbf{F}/T$  and  $\mathbf{\Lambda}'\mathbf{\Lambda}/N$  are both positive definite in the limit. However, Onatski (2012) shows that the APC estimates are inconsistent when  $\mathbf{\Lambda}'\mathbf{\Lambda}$  (without dividing by  $N$ ) has a positive definite limit. This has generated a good deal of interest in determining the number of less pervasive factors. Some assume large idiosyncratic variances, some assume that the entries of  $\mathbf{\Lambda}$  are non-zero but small, while others assume a sparse  $\mathbf{\Lambda}$  with many zero entries. See, for example, DeMol et al. (2008), Lettau and Pelger (2020), Uematsu and Yamagata (2019), Freyaldenhoven (2021). Though the term ‘weak factors’ is used in different ways, there is a presumption that the APC estimator has undesirable properties when the strong factor assumption fails. However, to our knowledge, there does not exist a clear statement of what those properties are.

In this paper, we consider the weaker condition that  $\mathbf{\Lambda}'\mathbf{\Lambda}/N^\alpha$  has a positive definite limit with  $\alpha \in (0, 1]$ . Since it is the strength of the loadings that is being weakened and positive definiteness of  $\mathbf{F}'\mathbf{F}/T$  is maintained throughout, we use the terminology of *weaker loadings*. Our main result is that while the strong factor assumption of  $\alpha = 1$  yields the fastest convergence rates possible, and the estimates are inconsistent when the loadings are extremely weak in the case of  $\alpha = 0$ , the principal component estimator continues to be consistent when  $\alpha \in (0, 1]$ . For the factor space, we establish that  $\frac{1}{T} \sum_{t=1}^T \|\tilde{\mathbf{F}}_t - \mathbf{H}'\mathbf{F}_t^0\|^2 = O_p(\frac{1}{N^\alpha}) + O_p((\frac{N^{1-\alpha}}{T})^2)$ , which depends on  $\alpha$ . However, for the common component, we obtain  $\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \|\tilde{C}_{it} - C_{it}^0\|^2 = O_p(\frac{1}{N}) + O_p(\frac{1}{T})$  as in the strong factor case. In terms of distribution theory, we show that  $\sqrt{N^\alpha}(\tilde{\mathbf{F}}_t - \mathbf{H}'\mathbf{F}_t^0)$ ,  $\sqrt{T}(\tilde{\Lambda}_i - \mathbf{H}^{-1}\Lambda_i^0)$ , and  $\min(\sqrt{N^\alpha}, \sqrt{T})(\tilde{C}_{it} - C_{it}^0)$  are all asymptotically normal when  $\alpha > 1/2$ . Though some additional assumptions on  $N$  and  $T$  are needed,  $N$  is not required to grow at the same rate as  $T$ , and knowledge of  $\alpha$  is not necessary for inference. The results have implications for factor augmented regressions, and more generally, for panel data models with cross-section dependence.

The convergence rate of the factor space we now obtain for  $\alpha = 1$  is  $\min(N, T^2)$ , which is faster than previously derived for the strong factor case. This is made possible by a

different proof technique that also leads to significant simplifications, hence of independent interest.<sup>1</sup> The simplifications come partly from using higher level assumptions, and partly from using approximations to the original rotation matrix  $\mathbf{H}$  which also make it possible to conduct inference using a representation of the asymptotic variance that the user deems most convenient.

The paper proceeds as follows. Section 2 sets up the econometric framework and presents three useful preliminary results. Section 3 studies consistent estimation of the factors, the loadings, and introduces four asymptotically equivalent rotation matrices. The distribution theory is given in Section 4. Implications of weaker loadings for factor augmented regressions are discussed.

Throughout, matrices are written in bold-face to distinguish them from vectors. As a matter of notation,  $\|\mathbf{A}\|^2 = \sum_{i=1}^m \sum_{j=1}^n |A_{ij}|^2 = \text{Tr}(\mathbf{A}\mathbf{A}')$  is the squared Frobenius norm of a  $m \times n$  matrix  $\mathbf{A}$ ,  $\|\mathbf{A}\|_{sp}^2 = \rho_{\max}(\mathbf{A}'\mathbf{A})$  denotes the squared spectral norm of  $\mathbf{A}$ , where  $\rho_{\max}(\mathbf{B})$  denotes the largest eigenvalue for a positive semi-definite matrix  $\mathbf{B}$ . Note that  $\|\mathbf{A}\|_{sp} \leq \|\mathbf{A}\| \leq \sqrt{q}\|\mathbf{A}\|_{sp}$ , where  $q = \text{rank}(\mathbf{A})$ . Thus when the rank  $q$  is fixed, the two norms are equivalent in terms of asymptotic behavior.

## 2 The Econometric Setup

We use  $i = 1, \dots, N$  to index cross-section units and  $t = 1, \dots, T$  to index time series observations. Let  $X_i = (X_{i1}, \dots, X_{iT})'$  be a  $T \times 1$  vector of random variables and  $\mathbf{X} = (X_1, X_2, \dots, X_N)$  be a  $T \times N$  matrix. The normalized data  $\mathbf{Z} = \frac{\mathbf{X}}{\sqrt{NT}}$  admit singular value decomposition (SVD)

$$\mathbf{Z} = \frac{\mathbf{X}}{\sqrt{NT}} = \mathbf{U}_{NT} \mathbf{D}_{NT} \mathbf{V}'_{NT} = \mathbf{U}_{NT,k} \mathbf{D}_{NT,k} \mathbf{V}'_{NT,k} + \mathbf{U}_{NT,N-k} \mathbf{D}_{NT,N-k} \mathbf{V}'_{NT,N-k}$$

where  $\mathbf{U}'_{NT} \mathbf{U}_{NT} = \mathbf{I}_T$  and  $\mathbf{V}'_{NT} \mathbf{V}_{NT} = \mathbf{I}_N$ . In the above,  $\mathbf{D}_{NT,k}$  is a diagonal matrix of  $k$  singular values  $d_{NT,1}, \dots, d_{NT,k}$  arranged in descending order,  $\mathbf{U}_{NT,k}, \mathbf{V}_{NT,k}$  are the corresponding left and right singular vectors respectively. By the Eckart and Young (1936) theorem, the best rank  $k$  approximation of  $\mathbf{Z}$  is  $\mathbf{U}_{NT,k} \mathbf{D}_{NT,k} \mathbf{V}'_{NT,k}$ . This is obtained without imposing probabilistic assumptions on the data.

We represent the data using a static factor model with  $r$  factors. In matrix form,

$$\mathbf{X} = \mathbf{F}\mathbf{\Lambda}' + \mathbf{e}. \tag{1}$$

---

<sup>1</sup>An earlier version of the paper circulated as *Simpler proofs for approximate factor models of large dimensions* considers  $\alpha = 1$  only.

To simplify notation, the subscripts indicating that  $\mathbf{F}$  is  $T \times r$  and  $\mathbf{\Lambda}$  is  $N \times r$  will be suppressed when the context is clear. The common component  $\mathbf{C} = \mathbf{F}\mathbf{\Lambda}'$  has reduced rank  $r$  because  $\mathbf{F}$  and  $\mathbf{\Lambda}$  both have rank  $r$ . The  $N \times N$  covariance matrix of  $\mathbf{X}$  takes the form

$$\mathbf{\Sigma}_X = \mathbf{\Lambda}\mathbf{\Sigma}_F\mathbf{\Lambda}' + \mathbf{\Sigma}_e = \mathbf{\Sigma}_C + \mathbf{\Sigma}_e.$$

A strict factor model obtains when the errors  $e_{it}$  are cross-sectionally and serially uncorrelated so that  $\mathbf{\Sigma}_e$  is a diagonal matrix. The classical factor model studied in Anderson and Rubin (1956) uses the stronger assumption that  $e_{it}$  is iid and normally distributed. For economic analysis, this error structure is overly restrictive. We work with the approximate factor model formulated in Chamberlain and Rothschild (1983) which allows the idiosyncratic errors to be weakly correlated in both the cross-section and time series dimensions. In such a case,  $\mathbf{\Sigma}_e$  need not be a diagonal matrix.

## 2.1 The APC Estimator and Assumptions

Let  $\mathbf{F}^0$  and  $\mathbf{\Lambda}^0$  be the true values of  $\mathbf{F}$  and  $\mathbf{\Lambda}$ . The model for unit  $i$  at time  $t$  as

$$x_{it} = \Lambda_i^{0'} F_t^0 + e_{it}.$$

Letting  $e_i' = (e_{i1}, e_{i2}, \dots, e_{iT})$  and  $e_t' = (e_{1t}, e_{2t}, \dots, e_{Nt})$ , the model for unit  $i$  is

$$X_i = \mathbf{F}^0 \Lambda_i^{0'} + e_i$$

Estimation of  $\mathbf{F}$  and  $\mathbf{\Lambda}$  in an approximate factor model with  $r$  factors proceeds by minimizing the sum of squared residuals:

$$\begin{aligned} \min_{\mathbf{F}, \mathbf{\Lambda}} \text{SSR}(\mathbf{F}, \mathbf{\Lambda}; r) &= \min_{\mathbf{F}, \mathbf{\Lambda}} \frac{1}{NT} \|\mathbf{X} - \mathbf{F}\mathbf{\Lambda}'\|^2 \\ &= \min_{\mathbf{F}, \mathbf{\Lambda}} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (x_{it} - \Lambda_i' F_t)^2. \end{aligned}$$

As  $\mathbf{F}$  and  $\mathbf{\Lambda}$  are not separately identified, we impose the normalization restrictions

$$\frac{\mathbf{F}'\mathbf{F}}{T} = I_r, \quad \mathbf{\Lambda}'\mathbf{\Lambda} \text{ is diagonal.} \quad (2)$$

The solution is the (static) APC estimator defined as:

$$(\tilde{\mathbf{F}}, \tilde{\mathbf{\Lambda}}) = (\sqrt{T}\mathbf{U}_{NT,r}, \sqrt{N}\mathbf{V}_{NT,r}\mathbf{D}_{NT,r}). \quad (3)$$

APC estimation of large dimensional approximate factor models must overcome two challenges not present in the classical factor analysis of Anderson and Rubin (1956). The first pertains to the fact that the errors are now allowed to be cross-sectionally correlated. The second issue arises because the  $T \times T$  covariance matrix of  $\mathbf{X}$  and the  $N \times N$  covariance of  $\mathbf{X}'$  are of infinite dimensions when  $N$  and  $T$  are large. To study the properties of the APC estimates, we use  $\mathbf{X} = \mathbf{F}^0 \boldsymbol{\Lambda}^{0'} + \mathbf{e}$  to obtain:

$$\frac{1}{NT} \mathbf{X} \mathbf{X}' = \frac{\mathbf{F}^0 (\boldsymbol{\Lambda}^{0'} \boldsymbol{\Lambda}^0) \mathbf{F}^{0'}}{N} + \frac{\mathbf{F}^0 \boldsymbol{\Lambda}^{0'} \mathbf{e}'}{NT} + \frac{\mathbf{e} \boldsymbol{\Lambda}^0 \mathbf{F}^{0'}}{NT} + \frac{\mathbf{e} \mathbf{e}'}{NT}. \quad (4)$$

But  $\frac{1}{NT} \mathbf{X} \mathbf{X}' = \mathbf{U}_{NT} \mathbf{D}_{NT}^2 \mathbf{U}_{NT}'$  and thus  $\frac{1}{NT} \mathbf{X} \mathbf{X}' \tilde{\mathbf{F}} = \tilde{\mathbf{F}} \mathbf{D}_{NT,r}^2$ . It follows that

$$\frac{\mathbf{F}^0 (\boldsymbol{\Lambda}^{0'} \boldsymbol{\Lambda}^0) \mathbf{F}^{0'} \tilde{\mathbf{F}}}{N} + \frac{\mathbf{F}^0 \boldsymbol{\Lambda}^{0'} \mathbf{e}' \tilde{\mathbf{F}}}{NT} + \frac{\mathbf{e} \boldsymbol{\Lambda}^0 \mathbf{F}^{0'} \tilde{\mathbf{F}}}{NT} + \frac{\mathbf{e} \mathbf{e}' \tilde{\mathbf{F}}}{NT} = \tilde{\mathbf{F}} \mathbf{D}_{NT,r}^2. \quad (5)$$

Rearranging terms yields

$$\tilde{\mathbf{F}}_t - \tilde{\mathbf{H}}'_{NT,0} \mathbf{F}_t^0 = \tilde{\mathbf{D}}_{NT}^{-2} \left( \frac{1}{T} \sum_{t=1}^T \tilde{\mathbf{F}}_s \gamma_{st} + \frac{1}{T} \sum_{s=1}^T \tilde{\mathbf{F}}_s \zeta_{st} + \frac{1}{T} \sum_{s=1}^T \tilde{\mathbf{F}}_s \eta_{st} + \frac{1}{T} \sum_{s=1}^T \tilde{\mathbf{F}}_s \xi_{st} \right)$$

where  $\tilde{\mathbf{H}}_{NT,0} = \left( \frac{\boldsymbol{\Lambda}^{0'} \boldsymbol{\Lambda}^0}{N} \right) \left( \frac{\mathbf{F}^{0'} \tilde{\mathbf{F}}}{T} \right) \mathbf{D}_{NT,r}^{-2}$ ,  $\gamma_{st} = E(\frac{1}{N} \mathbf{e}'_s \mathbf{e}_t) = E(\frac{1}{N} \sum_{i=1}^N e_{is} e_{it})$ ,  $\zeta_{st} = \frac{1}{N} \mathbf{e}'_s \mathbf{e}_t - \gamma_{st}$ ,  $\eta_{st} = \mathbf{F}_s^{0T} \boldsymbol{\Lambda}^{0T} \mathbf{e}_t / N$ , and  $\xi_{st} = \mathbf{F}_t^{0T} \boldsymbol{\Lambda}^{0T} \mathbf{e}_s / N$ . Stock and Watson (2002); Bai and Ng (2002); Bai (2003) established properties of the APC estimator by analyzing the four terms under certain assumptions, and this is by and large the approach that the literature has taken. In what follows, we work directly with the matrix norms of the terms in (5). This makes it possible to obtain simpler proofs under more general assumptions.

**Assumption A1:** Let  $M < \infty$  not depending on  $N$  and  $T$  and define

$$\delta_{NT} = \min(\sqrt{N}, \sqrt{T}).$$

i Mean independence:  $E(e_{it} | \Lambda_i^0, \mathbf{F}_t^0) = 0$ .

ii Weak (cross-sectional and serial) correlation in the errors.

(a)  $E \left[ \frac{1}{\sqrt{N}} \sum_{i=1}^N [e_{it} e_{is} - E(e_{it} e_{is})] \right]^2 \leq M$ ,

(b) For all  $i$ ,  $\frac{1}{T} \sum_{t=1}^T \sum_{s=1}^T |E(e_{it} e_{is})| \leq M$ , and for all  $t$ ,  $\frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N |E(e_{it} e_{jt})| \leq M$

(c) For all  $t$ ,  $\frac{1}{N\sqrt{T}} \|\mathbf{e}'_t \mathbf{e}\| = O_p(\delta_{NT}^{-1})$  and for all  $i$ ,  $\frac{1}{T\sqrt{N}} \|\mathbf{e}'_i \mathbf{e}\| = O_p(\delta_{NT}^{-1})$ .

(d)  $\|\mathbf{e}\|_{sp}^2 = \rho_{\max}(\mathbf{e}' \mathbf{e}) = O_p(\max\{N, T\})$ .

**Assumption A2:** (i)  $E\|F_t\|^4 \leq M$ ,  $\text{plim}_{T \rightarrow \infty} \frac{\mathbf{F}^{0'} \mathbf{F}^0}{T} = \boldsymbol{\Sigma}_F > 0$ ;  
(ii)  $\|\Lambda_i\| \leq M$ ,  $\lim_{N \rightarrow \infty} \frac{\Lambda^{0'} \Lambda^0}{N^\alpha} = \boldsymbol{\Sigma}_\Lambda > 0$ , for some  $\alpha > 0$  with  $\alpha \in (0, 1]$ ;  
(iii) the eigenvalues of  $\boldsymbol{\Sigma}_\Lambda \boldsymbol{\Sigma}_F$  are distinct.

**Assumption A3:** For each  $t$ , (i)  $E\|N^{-\alpha/2} \sum_i \Lambda_i^0 e_{it}\|^2 \leq M$ , (ii)  $\frac{1}{NT} e'_t e' \mathbf{F}^0 = O_p(\delta_{NT}^{-2})$ ; for each  $i$ , (iii)  $E\|T^{-1/2} \sum_t F_t^0 e_{it}\|^2 \leq M$ , (iv)  $\frac{1}{N^\alpha T} e'_i e \Lambda^0 = O_p(\frac{1}{N^\alpha}) + O_p(\frac{1}{\sqrt{TN^\alpha}})$ ; (v)  $\Lambda^{0'} e' \mathbf{F}^0 = \sum_{i=1}^N \sum_{t=1}^T \Lambda_i^0 F_t^{0'} e_{it} = O_p(\sqrt{N^\alpha T})$ .

**Assumption A4:** As  $N, T \rightarrow \infty$ ,  $(\frac{N}{N^\alpha})^{\frac{1}{T}} \rightarrow 0$  for the same  $\alpha$  in Assumption A2.

Assumption A1.i uses mean independence in place of moment conditions on  $e_{it}$  as in previous work. Assumption A1.ii assumes weak time and cross-section dependence. Assumption A.1(d) is a bound on the maximum eigenvalues of  $e'e$ . For iid data with uniformly bounded fourth moments, the rate is implied by random matrix theory. Moon and Weidner (2017) extends the case to data that are weakly correlated across  $i$  and  $t$ . We use it to obtain simpler proofs. Assumption A2 implies  $\|\mathbf{F}^0\|^2/T = O_p(1)$  and  $\|\Lambda^0\|^2/N^\alpha = O_p(1)$ . Parts of Assumption A3 also appear in Bai and Ng (2002). When the errors  $e_{it}$  are independent, Assumptions A1 and A2 are enough to validate A3. The assumption should hold under weak cross-sectional and serial correlations. Allowing for weaker factors comes at a cost. As stated in Assumption A4, which is new, a small  $\alpha$  must be compensated by a larger  $T$ .

The defining characteristic of an approximate factor model is that the first  $r$  largest population eigenvalues of  $\boldsymbol{\Sigma}_C$  diverge with  $N$  while all remaining eigenvalues of  $\boldsymbol{\Sigma}_C$  are zero, and all eigenvalues of  $\boldsymbol{\Sigma}_e$  are bounded. Previous works model the ‘diverge with  $N$ ’ feature by assuming that  $\mathbf{F}^{0'} \mathbf{F}^0/T > 0$  and  $\Lambda^{0'} \Lambda^0/N$  are positive definite in the limit. These two conditions have come to be known as the strong factor structure. Onatski (2012) considers the other extreme of  $\alpha = 0$ . We accommodate weaker loadings but require that  $\alpha > 0$  which nests the strong factor model as a special case. Note that the strength of the factor loadings affects the normalization of  $\Lambda^{0'} \Lambda^0$  but not  $\mathbf{F}^{0'} \mathbf{F}^0$ . Assumption A2.(ii) allows the  $r$  eigenvalues of  $\Lambda^{0'} \Lambda^0$  to diverge at a rate of  $N^\alpha$  with  $1 \geq \alpha > 0$ . Assumptions A3.(b) and (d) reflect the more general setup. Assumption A3 implies

$$\frac{\mathbf{F}^{0'} e e' \mathbf{F}^0}{NT} = \frac{1}{N} \sum_{i=1}^N \left[ \left( \frac{1}{\sqrt{T}} \sum_t F_t^0 e_{it} \right) \left( \frac{1}{\sqrt{T}} \sum_t F_t^0 e_{it} \right)' \right] = O_p(1) \quad (6)$$

$$\frac{\Lambda^{0'} e' e \Lambda^0}{N^\alpha T} = \frac{1}{T} \sum_{t=1}^T \left[ \left( \frac{1}{\sqrt{N^\alpha}} \sum_i \Lambda_i^0 e_{it} \right) \left( \frac{1}{\sqrt{N^\alpha}} \sum_i \Lambda_i^0 e_{it} \right)' \right] = O_p(1). \quad (7)$$

DeMol et al. (2008) considers a setting in which the eigenvalues of  $\Sigma_e$  are large relative to those of  $\Sigma_C$ . As discussed in Onatski (2012), such a setup can be rewritten in terms of weaker loadings with  $1 \geq \alpha > 0$ . The focus of DeMol et al. (2008) is shrinkage priors in Bayesian estimation and the implications for forecasting. Lettau and Pelger (2020) considers a penalized APC estimator to account for pricing errors in expected returns. Uematsu and Yamagata (2019, 2020) assume in our notation that  $\Lambda^0 \mathbf{H}^{-1} \mathbf{H}^{-1} \Lambda^0$  is a diagonal matrix with  $d_j^2 N^{\alpha_j}$  in the  $j$ -th diagonal, where  $\alpha_j$  reflects the strength of factor  $j$ , and for an unspecified choice of  $\mathbf{H}$ . Their main interest is in a new regularized estimator that captures sparsity of the loadings. Freyaldenhoven (2021) develops criteria to determine the number of factors of differing strength (local factors) and provides some results for the APC estimates when  $\alpha < 1$ . We consider standard APC estimation without knowledge of  $\alpha$  and assume that it is the same for all factors, but we provide the complete distribution theory for  $\tilde{F}_t$ ,  $\tilde{\Lambda}_i$  and  $\tilde{C}_{it}$ , making explicit that  $\mathbf{H}$  is the limit of a matrix that depends on the data generating process. As we will now proceed to show,  $\mathbf{H}$  plays an important role in the asymptotic theory.

## 2.2 Useful Identities and Matrices

In the strong factor case,  $\mathbf{D}_{NT,r}^2$  is a diagonal matrix of the  $r$  largest eigenvalues of  $\mathbf{Z}'\mathbf{Z} = \frac{1}{NT} \mathbf{X}\mathbf{X}'$ . The singular values of  $\mathbf{Z}$  are those of  $\mathbf{X}$  divided by  $\sqrt{NT}$ . In practice, each column of  $\mathbf{Z}$  is transformed to have unit variance so  $d_{NT,j}^2$  is the fraction of variation in  $\mathbf{Z}$  explained by factor  $j$ . The following lemma shows that to accommodate weaker factors,  $\mathbf{D}_{NT,r}^2$  must be scaled up by  $N^{1-\alpha}$  to have a limit matrix  $\mathbf{D}_r^2$  that is full rank.

**Lemma 1** *Let  $\mathbf{D}_r^2$  be a diagonal matrix consisting of the the ordered eigenvalues of  $\Sigma_\Lambda \Sigma_F$ . Under Assumption A, we have*

$$\left(\frac{N}{N^\alpha}\right) \mathbf{D}_{NT,r}^2 \xrightarrow{p} \mathbf{D}_r^2 > 0, \quad 1 \geq \alpha > 0.$$

**Proof:** Proper normalization of  $\mathbf{D}_{NT,r}^2$  is key to accommodating  $1 \geq \alpha > 0$ . The diagonal matrix  $\frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2$  consists of the  $r$  largest eigenvalues of  $\frac{1}{N^\alpha T} \mathbf{X}\mathbf{X}'$ , and

$$\frac{1}{N^\alpha T} \mathbf{X}\mathbf{X}' = \frac{\mathbf{F}^0 (\Lambda^0 \Lambda^0) \mathbf{F}^{0'}}{N^\alpha} \frac{\mathbf{F}^{0'}}{T} + \frac{\mathbf{F}^0 \Lambda^0 \mathbf{e}'}{N^\alpha T} + \frac{\mathbf{e} \Lambda^0 \mathbf{F}^{0'}}{N^\alpha T} + \frac{\mathbf{e} \mathbf{e}'}{N^\alpha T}.$$

Since the largest eigenvalue of  $\mathbf{e} \mathbf{e}'$  is of order  $\max\{N, T\}$ , the largest eigenvalue of the last matrix is bounded by  $O_p\left(\frac{N}{N^\alpha} \frac{1}{T}\right) + O_p\left(\frac{1}{N^\alpha}\right) \xrightarrow{p} 0$ . Furthermore,  $\|\mathbf{e}\|_{sp} = O_p(\sqrt{\max\{N, T\}})$ ,

$\|\mathbf{F}^0\|_{sp} = O_p(T^{1/2})$  and  $\|\mathbf{\Lambda}^0\|_{sp} = O_p(N^{\alpha/2})$ . A bound in spectral norm for the second matrix on the right hand side is

$$\frac{\|e\|_{sp}\|\mathbf{F}^0\|_{sp}\|\mathbf{\Lambda}^0\|_{sp}}{TN^\alpha} \leq O_p\left(\sqrt{\frac{N}{N^\alpha T}}\right) + O_p\left(\frac{1}{N^{\alpha/2}}\right) \xrightarrow{p} 0.$$

The third matrix is the transpose of the second. Thus, the largest eigenvalues of the last three matrices converges to zero. By the matrix perturbation theorem, the  $r$  largest eigenvalues of  $\frac{1}{N^\alpha T}\mathbf{X}\mathbf{X}'$  are determined by the first matrix on the right hand side. The eigenvalues of this matrix are the same as those of

$$\left(\frac{\mathbf{\Lambda}^{0'}\mathbf{\Lambda}^0}{N^\alpha}\right)\left(\frac{\mathbf{F}^{0'}\mathbf{F}^0}{T}\right).$$

This matrix converges to  $\mathbf{\Sigma}_\Lambda\mathbf{\Sigma}_F$  whose eigenvalues are  $\mathbf{D}_r^2$ , proving the lemma.  $\square$

Next, we turn to two matrices that will play important roles subsequently. The first is the covariance matrix  $\tilde{\mathbf{F}}'\mathbf{F}^0/T$ . To obtain its limit, we multiply  $(N/N^\alpha)\tilde{\mathbf{F}}'$  on each side of (5) and use the fact that  $\tilde{\mathbf{F}}'\tilde{\mathbf{F}} = T$  to obtain

$$\left(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right)\frac{\mathbf{\Lambda}^{0'}\mathbf{\Lambda}^0}{N^\alpha}\left(\frac{\mathbf{F}^{0'}\tilde{\mathbf{F}}}{T}\right) + \frac{\tilde{\mathbf{F}}'\mathbf{F}^0\mathbf{\Lambda}^{0'}e'\tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\tilde{\mathbf{F}}'e\mathbf{\Lambda}^0\mathbf{F}^{0'}\tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\tilde{\mathbf{F}}'ee'\tilde{\mathbf{F}}}{N^\alpha T^2} = \frac{N}{N^\alpha}\mathbf{D}_{NT,r}^2. \quad (8)$$

The right hand side converges to a positive definite matrix (thus invertible) by Lemma 1. The last three matrices on the left hand side converges in probability to zero. In particular,

$$\left\|\frac{\tilde{\mathbf{F}}'\mathbf{F}^0\mathbf{\Lambda}^{0'}e'\tilde{\mathbf{F}}}{N^\alpha T^2}\right\| \leq \left\|\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right\|\|\mathbf{\Lambda}^{0'}e'\|\|\tilde{\mathbf{F}}\|\frac{1}{N^\alpha T} = O_p\left(\frac{1}{N^{\alpha/2}}\right) = o_p(1)$$

and

$$\left\|\frac{\tilde{\mathbf{F}}'ee'\tilde{\mathbf{F}}}{N^\alpha T^2}\right\| \leq \rho_{\max}(ee')\frac{\|\tilde{\mathbf{F}}\|^2}{T}\frac{1}{N^\alpha T} \leq \frac{\max\{N, T\}}{N^\alpha T}O_p(1) = o_p(1). \quad (9)$$

The limit on the left hand side is thus determined by the first matrix, ie.

$$\left(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right)\frac{\mathbf{\Lambda}^{0'}\mathbf{\Lambda}^0}{N^\alpha}\left(\frac{\mathbf{F}^{0'}\tilde{\mathbf{F}}}{T}\right) + o_p(1) = \frac{N}{N^\alpha}\mathbf{D}_{NT,r}^2. \quad (10)$$

The limit of  $\tilde{\mathbf{F}}'\mathbf{F}^0/T$  can be obtained from this representatino.

**Lemma 2** *Under Assumption A,*

i.  $\tilde{\mathbf{F}}'\mathbf{F}^0/T \xrightarrow{p} \mathbf{Q} := \mathbf{D}_r\mathbf{\Upsilon}\mathbf{\Sigma}_\Lambda^{-1/2}$ , where  $\mathbf{\Upsilon}$  consists of the eigenvectors of the matrix  $\mathbf{\Sigma}_F^{1/2}\mathbf{\Sigma}_\Lambda\mathbf{\Sigma}_F^{1/2}$  with  $\mathbf{\Upsilon}'\mathbf{\Upsilon} = \mathbf{I}_r$ .

ii. For  $\mathbf{H}_{NT,0} = \left(\frac{\mathbf{\Lambda}^{0'}\mathbf{\Lambda}^0}{N}\right)\left(\frac{\mathbf{F}^{0'}\tilde{\mathbf{F}}}{T}\right)\mathbf{D}_{NT,r}^{-2}$ ,  $\mathbf{H}_{NT,0} \xrightarrow{p} \mathbf{Q}^{-1}$ .



Part (i) is obtained by taking limit on each side of (10) and to yield  $\mathbf{Q}\Sigma_{\Lambda}\mathbf{Q}' = \mathbf{D}_r^2$ . Since  $\mathbf{D}_r^2$  is diagonal with distinct elements, we can solve for  $\mathbf{Q}$  as  $\mathbf{Q} = \mathbf{D}_r\Upsilon\Sigma_{\Lambda}^{-1/2}$  where  $\Upsilon$  consists of the orthonormal eigenvectors of the matrix  $\Sigma_F^{1/2}\Sigma_{\Lambda}\Sigma_F^{1/2}$ . The  $\mathbf{Q}$  matrix is full rank and invertible. The solution holds up to a column sign change, just like  $\tilde{\mathbf{F}}$  is determined up to a column sign change.

The rotation matrix  $\mathbf{H}_{NT,0}$ , first derived in Stock and Watson (1998), has been used to evaluate the precision of  $\tilde{\mathbf{F}}$ . Bai (2003) shows that  $\mathbf{H}_{NT,0} \xrightarrow{p} \mathbf{Q}^{-1}$  when  $\alpha = 1$ . To accommodate weaker loadings, we consider

$$\mathbf{H}_{NT,0} = \left( \frac{\Lambda^{0'}\Lambda^0}{N^\alpha} \right) \left( \frac{\mathbf{F}^{0'}\tilde{\mathbf{F}}}{T} \right) \left( \frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2 \right)^{-1}.$$

By assumption, the first matrix on the right hand side is invertible while the last two matrices are invertible by the previous lemmas. Hence  $\mathbf{H}_{NT,0} \xrightarrow{p} \Sigma_{\Lambda}\mathbf{Q}'\mathbf{D}_r^{-2} \equiv \mathbf{Q}^{-1}$ . The matrix  $\mathbf{Q}$  and its relation to  $\mathbf{H}_{NT,0}$  are fundamental to the asymptotic theory in the strong factor case. Lemma 2 shows that the relations are unaffected when weaker loadings are allowed.

### 3 Consistent Estimation of the Factor Space

This section has three parts. Subsection 1 presents consistent estimation of the factors up to rotation by  $\mathbf{H}_{NT,0}$ . Subsection 2 introduces four new rotation matrices. Subsection 3 uses these new matrices to show consistent estimation of the loadings.

#### 3.1 The Factors

To establish consistent estimation of  $\tilde{\mathbf{F}}$  for  $\mathbf{F}^0$  up to rotation by  $\mathbf{H}_{NT,0}$ , we multiply  $\mathbf{D}_{NT,r}^{-2}$  to both sides of (5) and use the definition of  $\mathbf{H}_{NT,0}$  to obtain

$$\begin{aligned} \tilde{\mathbf{F}} - \mathbf{F}^0\mathbf{H}_{NT,0} &= \left( \frac{\mathbf{F}^0\Lambda^{0'}\mathbf{e}'\tilde{\mathbf{F}}}{NT} + \frac{\mathbf{e}\Lambda^0\mathbf{F}^{0'}\tilde{\mathbf{F}}}{NT} + \frac{\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}}{NT} \right) \mathbf{D}_{NT,r}^{-2} \\ &= \left( \frac{\mathbf{F}^0\Lambda^{0'}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T} + \frac{\mathbf{e}\Lambda^0\mathbf{F}^{0'}\tilde{\mathbf{F}}}{N^\alpha T} + \frac{\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T} \right) \left( \frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2 \right)^{-1}. \end{aligned} \quad (11)$$

This implies

$$\begin{aligned} \frac{1}{\sqrt{T}} \|\tilde{\mathbf{F}} - \mathbf{F}^0\mathbf{H}_{NT,0}\| &\leq \left\{ 2 \left( \frac{\|\mathbf{F}^0\| \|\tilde{\mathbf{F}}\|}{T} \right) \left( \frac{1}{\sqrt{T}N^\alpha} \|\Lambda^{0'}\mathbf{e}'\| \right) + \frac{\|\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}\|}{N^\alpha T^{3/2}} \right\} \left\| \left( \frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2 \right)^{-1} \right\| \\ &= O_p \left( \frac{1}{\sqrt{T}N^\alpha} \|\Lambda^{0'}\mathbf{e}'\| \right) + O_p \left( \frac{\|\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}\|}{N^\alpha T^{3/2}} \right). \end{aligned}$$

But  $\frac{1}{\sqrt{T}N^\alpha}\|\boldsymbol{\Lambda}^{0'}\mathbf{e}'\| = O_p(\frac{1}{\sqrt{N^\alpha}})$  by (7) and  $\frac{\|\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}\|}{N^\alpha T^{3/2}} \leq \frac{\rho_{\max}(\mathbf{e}\mathbf{e}')\|\tilde{\mathbf{F}}\|}{N^\alpha T^{3/2}} = O_p(\frac{1}{N^\alpha}) + \frac{N}{N^\alpha} \frac{1}{T} O_p(1)$ . Thus

$$\frac{1}{\sqrt{T}}\|\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0}\| = O_p(\frac{1}{\sqrt{N^\alpha}}) + \frac{1}{T} \frac{N}{N^\alpha} O_p(1).$$

Squaring it gives the following proposition.

**Proposition 1** *Under Assumption A, the following holds:*

$$\frac{1}{T}\|\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0}\|^2 = \frac{1}{T} \sum_{t=1}^T \|\tilde{F}_t - \mathbf{H}'_{NT,0} F_t^0\|^2 = O_p\left(\frac{1}{N^\alpha}\right) + \frac{1}{T^2} \left(\frac{N}{N^\alpha}\right)^2 O_p(1).$$

The result is stated in in squared Frobenius norm. For  $\alpha = 1$ , Theorem 1 of Bai and Ng (2002) gives a convergence rate for the same quantity of  $O_p(\delta_{NT}^{-2})$ . The proposition here uses a different proof to obtain a faster convergence rate of  $O_p(\frac{1}{N}) + \frac{1}{T^2} O_p(1)$  for the strong factor case of  $\alpha = 1$ . Implications of the proposition will be discussed subsequently.

## 3.2 Equivalent Rotation Matrices

The rotation matrix  $\mathbf{H}_{NT,0}$  is a product of three  $r \times r$  matrices and it is not easy to interpret. However, we can rewrite (8) as

$$\left(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right)\mathbf{H}_{NT,0} = \mathbf{I}_r - \left\{ \frac{\tilde{\mathbf{F}}'\mathbf{F}^0\boldsymbol{\Lambda}^{0'}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\tilde{\mathbf{F}}'\mathbf{e}\boldsymbol{\Lambda}^0\mathbf{F}^{0'}\tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\tilde{\mathbf{F}}'\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T^2} \right\} \left(\frac{N}{N^\alpha}\mathbf{D}_{NT,r}^2\right)^{-1}. \quad (12)$$

As  $\tilde{\mathbf{F}}'\mathbf{F}^0/T = O_p(1)$ , the product of  $\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}$  and  $\mathbf{H}_{NT,0}$  is an identity matrix up to an negligible term if it can be shown that the three terms inside the bracket is small. The next Lemma formalizes this result and shows that it also holds for four other rotation.

**Lemma 3** *Under Assumption A,*

$$i. \mathbf{H}_{NT,0} = \left(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right)^{-1} + O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + \left[ O_p\left(\frac{1}{N^\alpha}\right) + \left(\frac{N}{N^\alpha}\right)\frac{1}{T} O_p(1) \right].$$

ii. For  $\ell = 1, 2, 3, 4$ ,  $\mathbf{H}_{NT,\ell} = \mathbf{H}_{NT,0} + O_p(\frac{1}{\sqrt{N^\alpha T}}) + O_p(\frac{1}{N^\alpha}) + (\frac{N}{N^\alpha})\frac{1}{T} O_p(1)$ , where

$$\mathbf{H}_{NT,1} = (\boldsymbol{\Lambda}^{0'}\boldsymbol{\Lambda}^0)(\tilde{\boldsymbol{\Lambda}}'\boldsymbol{\Lambda}^0)^{-1},$$

$$\mathbf{H}_{NT,2} = (\mathbf{F}^{0'}\mathbf{F}^0)^{-1}(\mathbf{F}^{0'}\tilde{\mathbf{F}}),$$

$$\mathbf{H}_{NT,3} = (\tilde{\mathbf{F}}'\mathbf{F}^0)^{-1}(\tilde{\mathbf{F}}'\tilde{\mathbf{F}}) = (\tilde{\mathbf{F}}'\mathbf{F}^0/T)^{-1}, \text{ and}$$

$$\mathbf{H}_{NT,4} = (\boldsymbol{\Lambda}^{0'}\tilde{\boldsymbol{\Lambda}})(\tilde{\boldsymbol{\Lambda}}'\tilde{\boldsymbol{\Lambda}})^{-1} = (\boldsymbol{\Lambda}^{0'}\tilde{\boldsymbol{\Lambda}}/N)\mathbf{D}_{NT,r}^{-2}.$$

iii.  $\mathbf{H}_{NT,\ell} \xrightarrow{p} \mathbf{Q}^{-1}$ .

Part (i), shown in the Appendix, establishes the error in approximating  $\mathbf{H}_{NT,0}$  by  $(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T})^{-1}$  while part (ii) considers four additional approximations that provide an intuitive interpretation of  $\mathbf{H}_{NT}F_t^0$ . For example,  $\mathbf{H}_{NT,2}$  is coefficient matrix from projecting  $\tilde{\mathbf{F}}$  on the space spanned by  $\mathbf{F}^0$  and  $\mathbf{H}'_{NT,2}F_t^0$  is asymptotically the fit from the projection. These alternative rotation matrices were used in Bai and Ng (2019) for  $\alpha = 1$ . The above Lemma shows that they can still be used in place of  $\mathbf{H}_{NT,0}$  when  $\alpha < 1$ , but the adequacy of approximation will depend on  $\alpha$ .

### 3.3 The Loadings and the Common Component

The APC estimator satisfies  $\frac{1}{N}\tilde{\Lambda}'\tilde{\Lambda} = \mathbf{D}_{NT,r}^2$  and we already have  $\frac{1}{N^\alpha}\tilde{\Lambda}'\tilde{\Lambda} = \frac{N}{N^\alpha}\mathbf{D}_{NT,r}^2 \xrightarrow{p} \mathbf{D}_r^2$ . We can now provide a simple consistency proof for  $\tilde{\Lambda}$ . Multiply  $\frac{1}{T}\tilde{\mathbf{F}}'$  to both sides of  $\mathbf{X} = \mathbf{F}^{0'}\Lambda^{0'} + \mathbf{e}$  to obtain  $\frac{1}{T}\tilde{\mathbf{F}}'\mathbf{X} = (\tilde{\mathbf{F}}'\mathbf{F}^0/T)\Lambda^{0'} + \tilde{\mathbf{F}}'\mathbf{e}/T$ . We have

$$\begin{aligned}\tilde{\Lambda}' &= \mathbf{H}_{NT,3}^{-1}\Lambda^{0'} + \tilde{\mathbf{F}}'\mathbf{e}/T \\ \tilde{\Lambda}' - \mathbf{H}_{NT,3}^{-1}\Lambda^{0'} &= \mathbf{H}'_{NT,0}\mathbf{F}^{0'}\mathbf{e}/T + (\tilde{\mathbf{F}} - \mathbf{F}^0\mathbf{H}_{NT,0})'\mathbf{e}/T\end{aligned}\quad (13)$$

and thus

$$\frac{1}{\sqrt{N}}\|\tilde{\Lambda}' - \mathbf{H}_{NT,3}^{-1}\Lambda^{0'}\| \leq \|H_{NT,0}\| \frac{\|\mathbf{F}^{0'}\mathbf{e}\|}{T\sqrt{N}} + \frac{\|(\tilde{\mathbf{F}} - \mathbf{F}^0\mathbf{H}_{NT,0})'\mathbf{e}\|}{T\sqrt{N}}$$

The first term  $\|\mathbf{F}^0\mathbf{e}\|/(T\sqrt{N}) = O_p(1/\sqrt{T})$  by equation (6). The second term is  $O_p(\frac{1}{\sqrt{N^{1+\alpha}}})$ , shown in the Appendix. Combining results and ignoring terms dominated by  $O_p(T^{-1/2})$ , we have

$$\frac{1}{\sqrt{N}}\|\tilde{\Lambda}' - \mathbf{H}_{NT,3}^{-1}\Lambda^{0'}\| = O_p\left(\frac{1}{\sqrt{T}}\right) + O_p\left(\frac{1}{\sqrt{N^{1+\alpha}}}\right).$$

Squaring gives the next proposition:

**Proposition 2** *Under Assumption A, the following holds*

$$\frac{1}{N}\|\tilde{\Lambda} - \Lambda^0(\mathbf{H}'_{NT,0})^{-1}\|^2 = \frac{1}{N}\sum_{i=1}^N \|\tilde{\Lambda}_i - \mathbf{H}_{NT,0}^{-1}\Lambda_i^0\|^2 = O_p\left(\frac{1}{T}\right) + O_p\left(\frac{1}{N^{1+\alpha}}\right).$$

Note that here we replace  $\mathbf{H}_{NT,3}$  by  $\mathbf{H}_{NT,0}$ , but this does not affect the rate analysis. In fact, we can use other rotation matrices to gain intuition. For example,  $\mathbf{H}_{NT,1}^{-1}$  is obtained by regressing  $\tilde{\Lambda}$  on  $\Lambda_0$ . Hence  $\Lambda_0(\mathbf{H}'_{NT,1})^{-1}$  is asymptotically the fit from projecting  $\tilde{\Lambda}$  on the space spanned by  $\Lambda_0$ , and  $\tilde{\Lambda} - \Lambda^0(\mathbf{H}'_{NT,0})^{-1}$  is the error from that projection.

While  $\tilde{\mathbf{F}}$  and  $\tilde{\Lambda}$  only estimate  $\mathbf{F}^0$  and  $\Lambda^0$  up to a rotation matrix,  $\tilde{\mathbf{C}}$  does not depend on rotations and is directly comparable to  $\mathbf{C}^0$ .

**Proposition 3** *Under Assumption A,*

$$\frac{1}{NT} \|\tilde{\mathbf{C}} - \mathbf{C}^0\|^2 = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \|\tilde{C}_{it} - C_{it}^0\|^2 = O_p(\delta_{NT}^{-2}).$$

**Proof:** From  $\tilde{\mathbf{C}} - \mathbf{C}_0 = \tilde{\mathbf{F}}\tilde{\Lambda}' - \mathbf{F}^0\Lambda^{0'} = (\tilde{\mathbf{F}} - \mathbf{F}^0\mathbf{H}')\tilde{\Lambda}' + \mathbf{F}^0\mathbf{H}'\tilde{\Lambda}' - \mathbf{F}^0\Lambda^{0'}$ , we have

$$\begin{aligned} \frac{1}{NT} \|\tilde{\mathbf{C}} - \mathbf{C}^0\|^2 &\leq \frac{\|\tilde{\mathbf{F}} - \mathbf{F}^0\mathbf{H}'\|^2 \|\tilde{\Lambda}'\|^2}{T N} + \frac{\|\mathbf{F}^0\mathbf{H}'\|^2 \|\tilde{\Lambda}' - \mathbf{H}^{-1}\Lambda^{0'}\|^2}{T N} \\ &\leq \left[ O_p\left(\frac{1}{N^\alpha}\right) + \frac{1}{T^2} \left(\frac{N}{N^\alpha}\right)^2 O_p(1) \right] O_p\left(\frac{N^\alpha}{N}\right) + O_p\left(\frac{1}{T}\right) + O_p\left(\frac{1}{N^{1+\alpha}}\right) \\ &= O_p\left(\frac{1}{N}\right) + O_p\left(\frac{1}{T}\right) + \frac{1}{T^2} \frac{N}{N^\alpha} O_p(1) = O_p(\delta_{NT}^{-2}) \end{aligned}$$

where the second inequality follows from Propositions 1 and 2. The term  $\frac{1}{T^2} \frac{N}{N^\alpha} O_p(1)$  is dominated by  $O_p(1/T)$  since  $\frac{N}{N^\alpha} \frac{1}{T} \rightarrow 0$  by Assumption A.4. Note this is the same rate as the strong factor case.  $\square$

## 4 Distribution Theory

As we do not observe  $\mathbf{F}^0$  or  $\Lambda^0$ , we need an inferential theory for  $\tilde{F}_t$ ,  $\tilde{\Lambda}_i$ , and  $\tilde{C}_{it} = \tilde{F}_t \tilde{\Lambda}_i'$ . Assuming that  $\alpha = 1$ , Theorems 1 and 2 of Bai (2003) establish that

$$\sqrt{N}(\tilde{F}_t - \mathbf{H}'_{NT,0} F_t^0) \xrightarrow{d} \mathcal{N}(\mathbf{D}_r^{-2} \mathbf{Q} \Gamma_t \mathbf{Q}' \mathbf{D}_r^{-2}) \quad (14a)$$

$$\sqrt{T}(\tilde{\Lambda}_i - \mathbf{H}_{NT,0}^{-1} \Lambda_i^0) \xrightarrow{d} \mathcal{N}(0, \mathbf{Q}'^{-1} \Phi_i \mathbf{Q}^{-1}). \quad (14b)$$

by positing appropriate central limit theorems appropriate for  $\alpha = 1$ . Assumption B accommodates weaker loadings.

**Assumption B.** The following holds for each  $i$  and  $t$  as  $N, T \rightarrow \infty$ :

$$\frac{1}{\sqrt{N^\alpha}} \sum_{i=1}^N \Lambda_i^0 e_{it} \xrightarrow{d} \mathcal{N}(0, \mathbf{\Gamma}_t), \quad \frac{1}{\sqrt{T}} \sum_{t=1}^T F_t^0 e_{it} \xrightarrow{d} \mathcal{N}(0, \mathbf{\Phi}_i).$$

To obtain comparable asymptotic normality results when  $1 \geq \alpha > 0$  requires additional assumptions. To see why, first consider the limiting distribution  $\tilde{\Lambda}$ . From (13),

$$\sqrt{T}(\tilde{\Lambda}_i - \mathbf{H}_{NT,3}^{-1} \Lambda_i^0) = \mathbf{H}'_{NT,3} \frac{1}{\sqrt{T}} \sum_{t=1}^T F_t^0 e_{it} + \frac{(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,3})' e_i}{\sqrt{T}}.$$

The first term is asymptotically normal by Assumption B and  $\mathbf{H}'_{NT,3} \xrightarrow{p} \mathbf{Q}'^{-1}$  by Lemma 3. For the second term, we show in Lemma 6 part (iii) that for each  $i$ ,

$$\frac{1}{T} e'_i(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,\ell}) = O_p\left(\frac{1}{N^\alpha}\right) + O_p\left(\frac{1}{\sqrt{T} N^\alpha}\right) + O_p\left(\frac{N}{N^\alpha T}\right).$$

For  $\sqrt{T}$  consistency of  $\tilde{\mathbf{\Lambda}}$ , we need  $\sqrt{T}$  times the above three terms to be negligible. We thus require  $\frac{\sqrt{T}}{N^\alpha} \rightarrow 0$  and  $\frac{1}{\sqrt{T}} \frac{N}{N^\alpha} \rightarrow 0$ . A necessary condition for both to hold is  $\alpha > 1/2$ . These are listed as Assumption C(i)-(iii) below.

Similarly, for the distribution of  $\tilde{F}_t$ , we multiply  $\tilde{\mathbf{\Lambda}}(\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1}$  to both sides of  $\mathbf{X} = \mathbf{F}^0 \mathbf{\Lambda}^{0'} + \mathbf{e}$  to obtain  $\mathbf{X} \tilde{\mathbf{\Lambda}}(\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1} = \mathbf{F}^0 \mathbf{\Lambda}^{0'} \tilde{\mathbf{\Lambda}}(\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1} + \mathbf{e} \tilde{\mathbf{\Lambda}}(\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1}$ . Using the definition of  $\mathbf{H}_{NT,4}$ ,

$$\begin{aligned} \tilde{\mathbf{F}} &= \mathbf{F}^0 \mathbf{H}_{NT,4} + \mathbf{e} \tilde{\mathbf{\Lambda}}(\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1} \\ &= \mathbf{F}^0 \mathbf{H}_{NT,4} + \mathbf{e} \mathbf{\Lambda}^0 \mathbf{H}'_{NT,4} (\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1} + \mathbf{e}(\tilde{\mathbf{\Lambda}} - \mathbf{\Lambda}^0 \mathbf{H}'_{NT,4}) (\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}})^{-1}. \end{aligned} \quad (15)$$

Thus

$$\sqrt{N^\alpha}(\tilde{F}_t - \mathbf{H}'_{NT,4} F_t^0) = \left(\frac{\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}}}{N^\alpha}\right)^{-1} \mathbf{H}_{NT,4}^{-1} \frac{1}{\sqrt{N^\alpha}} \sum_{i=1}^N \Lambda_i^0 e_{it} + \left(\frac{\tilde{\mathbf{\Lambda}}'\tilde{\mathbf{\Lambda}}}{N^\alpha}\right)^{-1} \frac{(\tilde{\mathbf{\Lambda}} - \mathbf{\Lambda}^0 \mathbf{H}'_{NT,4})' e_t}{\sqrt{N^\alpha}}$$

The first term is asymptotically normal by Assumption B. Now Lemma 6 part (iv) shows that for each  $t$ , the average correlation between  $e_t$  and the error from estimating  $\tilde{\mathbf{\Lambda}}$  is

$$\frac{1}{N} e'_t(\tilde{\mathbf{\Lambda}} - \mathbf{\Lambda}^0 \mathbf{H}'_{NT,\ell}) = \frac{1}{T} \sqrt{\frac{N}{N^\alpha}} O_p(1) + \frac{1}{\sqrt{N^{1+\alpha}}} O_p(1) + \frac{1}{T^{3/2}} \left(\frac{N}{N^\alpha}\right) O_p(1) + \frac{1}{N^\alpha \sqrt{T}} O_p(1).$$

In the strong factor case when  $\alpha = 1$ , having the three terms vanish as  $N, T \rightarrow \infty$  suffice for  $\sqrt{N}$  consistency of  $\tilde{F}_t$ . With weaker loadings, we can only get  $\sqrt{N^\alpha}$  consistency of  $\tilde{F}_t$ , and this requires  $N/\sqrt{N^\alpha}$  times each of the four terms to be negligible. That is,

$$\frac{1}{T} \frac{N^{3/2}}{N^\alpha} \rightarrow 0, \quad N^{\frac{1}{2}-\alpha} \rightarrow 0, \quad \frac{1}{T^{3/2}} \frac{N^2}{N^{3\alpha/2}} \rightarrow 0, \quad \frac{N}{N^{3\alpha/2}} \frac{1}{\sqrt{T}} \rightarrow 0.$$

The third and the fourth conditions hold if the first two along with C(iii) are satisfied. Thus, in addition to  $\alpha > 1/2$ , we also need C(iv),  $\frac{N}{T} N^{1/2-\alpha} \rightarrow 0$  to restrict the relation between  $N$  and  $T$ .

We collect the required conditions for  $\sqrt{T}$  asymptotic normality of  $\tilde{\mathbf{\Lambda}}$  and  $\sqrt{N^\alpha}$  asymptotic normality of  $\tilde{\mathbf{F}}$  into the following:

**Assumption C:** (i)  $\alpha > \frac{1}{2}$ , (ii)  $\frac{\sqrt{T}}{N^\alpha} \rightarrow 0$ , (iii)  $\frac{1}{\sqrt{T}} \frac{N}{N^\alpha} \rightarrow 0$ , and (iv)  $\frac{1}{T} \frac{N^{3/2}}{N^\alpha} \rightarrow 0$ .

The above conditions reduce to  $\sqrt{T}/N \rightarrow 0$  and  $\sqrt{N}/T \rightarrow 0$  when  $\alpha = 1$ . In general, they are distinctive restrictions, though some conditions may be redundant depending on the value of  $\alpha$ .

Under Assumption C, the preceding analysis implies

$$\sqrt{N^\alpha}(\tilde{F}_t - \mathbf{H}'_{NT,4}F_t^0) = \left(\frac{\tilde{\Lambda}'\tilde{\Lambda}}{N^\alpha}\right)^{-1} \mathbf{H}_{NT,4}^{-1} \frac{1}{\sqrt{N^\alpha}} \sum_{i=1}^N \Lambda_i^0 e_{it} + o_p(1) \quad (16a)$$

$$\sqrt{T}(\tilde{\Lambda}_i - \mathbf{H}_{3,NT}^{-1}\Lambda_i^0) = \mathbf{H}'_{NT,3} \frac{1}{\sqrt{T}} \sum_{t=1}^T F_t^0 e_{it} + o_p(1). \quad (16b)$$

**Proposition 4** *Under Assumptions A, B, and C and the normalization that  $\mathbf{F}'\mathbf{F}/T = I_r$  and  $\Lambda'\Lambda$  is diagonal, we have, as  $N, T \rightarrow \infty$ ,*

$$\begin{aligned} \sqrt{N^\alpha}(\tilde{F}_t - \mathbf{H}'_{NT,4}F_t^0) &\xrightarrow{d} \mathcal{N}(\mathbf{D}_r^{-2}\mathbf{Q}\Gamma_t\mathbf{Q}'\mathbf{D}_r^{-2}), \\ \sqrt{T}(\tilde{\Lambda}_i - \mathbf{H}_{NT,3}^{-1}\Lambda_i^0) &\xrightarrow{d} \mathcal{N}(0, \mathbf{Q}'^{-1}\Phi_i\mathbf{Q}^{-1}). \end{aligned}$$

The Proposition establishes  $(\sqrt{N^\alpha}, \sqrt{T})$  asymptotic normality of  $(\tilde{F}_t, \tilde{\Lambda}_i)$ . Though the conditions are more restrictive than the strong factor case, the results are more general than previously understood. Conditions weaker than Assumption C may be possible. For example, if  $T/N \rightarrow c \in (0, \infty)$ , then  $\alpha > 1/2$  is sufficient for asymptotic normality.

For hypothesis testing, there is no need to know  $\alpha$  as  $\frac{1}{N^\alpha}\mathbf{D}_r^{-2}\mathbf{Q}\Gamma_t\mathbf{Q}'\mathbf{D}_r^{-2}$  can be consistently estimated by  $\mathbf{D}_{NT,r}^{-2}(\sum_{i=1}^N \tilde{\Lambda}_i\tilde{\Lambda}_i'\tilde{e}_{it}^2)\mathbf{D}_{NT,r}^{-2}$  (assuming no cross-section correlation in  $e_{it}$ ). The square root of the  $k$ -th diagonal element of this matrix gives the standard error of the  $k$ -th component of  $\tilde{F}_t$ . Similarly,  $\frac{1}{T}\mathbf{Q}'^{-1}\Phi_i\mathbf{Q}^{-1}$  can be consistently estimated by  $\frac{1}{T}\sum_{s=1}^T \tilde{F}_s\tilde{F}_s'\tilde{e}_{is}^2$  (assuming no serial correlation in  $e_{it}$ ). These formulae are identical to the case of strong factors.

Although the limiting covariance matrices look different from those given in (14a) and (14b), they are mathematically identical because of the different ways to represent  $\mathbf{H}$  and thus  $\mathbf{Q}$ . Indeed, other representations of the asymptotic variances can be used. For example, since  $(\tilde{\Lambda}'\tilde{\Lambda}/N^\alpha)^{-1}\mathbf{H}_{NT,4}^{-1} = (\Lambda^0'\Lambda^0/N^\alpha)^{-1} = \mathbf{H}'_{1,NT}(\Lambda^0'\Lambda^0/N^\alpha)^{-1}$  and  $\mathbf{H}_{NT} \xrightarrow{p} \mathbf{Q}^{-1}$ , we also have

$$\begin{aligned} \sqrt{N^\alpha}(\tilde{F}_t - \mathbf{H}'_{NT,4}F_t^0) &= \mathbf{H}'_{NT,1} \left(\frac{\Lambda^0'\Lambda^0}{N^\alpha}\right)^{-1} \frac{1}{\sqrt{N^\alpha}} \sum_{i=1}^N \Lambda_i^0 e_{it} + o_p(1) \\ &\xrightarrow{d} \mathcal{N}(0, \mathbf{Q}^{-1'}\Sigma_\Lambda^{-1}\Gamma_t\Sigma_\Lambda^{-1}\mathbf{Q}). \end{aligned} \quad (17)$$

From  $\mathbf{H}'_{NT,3} = \mathbf{H}_{NT,2}^{-1}(\mathbf{F}^{0'}\mathbf{F}^0/T)^{-1}$  and using (16b), it also holds that

$$\begin{aligned}\sqrt{T}(\tilde{\Lambda}_i - \mathbf{H}_{NT,3}^{-1}\Lambda_i^0) &= \mathbf{H}_{NT,2}^{-1}\left(\frac{\mathbf{F}^{0'}\mathbf{F}^0}{T}\right)^{-1}\frac{1}{\sqrt{T}}\sum_{t=1}^T F_t^0 e_{it} + o_p(1) \\ &\xrightarrow{d} \mathcal{N}(0, \mathbf{Q}\Sigma_F^{-1}\Phi_i\Sigma_F^{-1}\mathbf{Q}').\end{aligned}\quad (18)$$

As the factor estimates are asymptotically normal regardless of the choice of the rotation matrix, one can use the most convenient representation for inference.

While there are many ways to represent the sampling error of  $\tilde{F}_t$  and  $\tilde{\Lambda}_i$ , the properties of  $\tilde{C}_{it}$  are invariant to the choice of  $\mathbf{H}_{NT,\ell}$ , so we can simply write  $\mathbf{H}_{NT}$ . By definition,  $C_{it}^0 = \Lambda_i^{0'}F_t^0$  and  $\tilde{C}_{it} = \tilde{\Lambda}_i'\tilde{F}_t^0$ . Adding and subtracting terms

$$\begin{aligned}\tilde{C}_{it} - C_{it}^0 &= \Lambda_i^{0'}\mathbf{H}'_{NT}{}^{-1}(\tilde{F}_t - \mathbf{H}'_{NT}F_t^0)' + (\tilde{\Lambda}_i - \mathbf{H}_{NT}^{-1}\Lambda_i^0)'\tilde{F}_t \\ &= \Lambda_i^{0'}\mathbf{H}'_{NT}{}^{-1}(\tilde{F}_t - \mathbf{H}'_{NT}F_t^0)' + F_t^{0'}\mathbf{H}_{NT}(\tilde{\Lambda}_i - \mathbf{H}_{NT}^{-1}\Lambda_i^0) + O_p\left(\frac{1}{\sqrt{TN^\alpha}}\right).\end{aligned}$$

Using (17) and (18), we have

$$\begin{aligned}(\tilde{C}_{it} - C_{it}^0) &= \frac{1}{\sqrt{N^\alpha}}\Lambda_i^{0'}\left(\frac{\Lambda^{0'}\Lambda^0}{N^\alpha}\right)^{-1}\frac{1}{\sqrt{N^\alpha}}\sum_{i=1}^N \Lambda_i^0 e_{it} \\ &\quad + \frac{1}{\sqrt{T}}F_t^{0'}\left(\frac{\mathbf{F}^{0'}\mathbf{F}^0}{T}\right)^{-1}\frac{1}{\sqrt{T}}\sum_{t=1}^T F_t^0 e_{it} + o_p(N^{-\alpha/2}) + o_p(T^{-1/2})\end{aligned}$$

This leads to the distribution theory for the estimated common components.

**Proposition 5** *Under Assumptions A, B and C, we have, as  $N, T \rightarrow \infty$ ,*

$$\frac{\tilde{C}_{it} - C_{it}^0}{\sqrt{\frac{1}{N^\alpha}W_{NT,it}^\Lambda + \frac{1}{T}W_{NT,it}^F}} \xrightarrow{d} \mathcal{N}(0, 1)$$

where  $W_{it}^\Lambda = \Lambda_i^{0'}\Sigma_\Lambda^{-1}\Gamma_t\Sigma_\Lambda^{-1}\Lambda_i^0$  and  $W_{it}^F = F_t^{0'}\Sigma_F^{-1}\Phi_i\Sigma_F^{-1}F_t^0$ .

The proposition implies a convergence rate of  $\min\{\sqrt{N^\alpha}, \sqrt{T}\}$  for the estimated common components. To estimate the standard error of  $\tilde{C}_{it}$ , there is also no need to know  $\alpha$ . The term  $\frac{1}{N^\alpha}W_{NT,it}^\Lambda$  is consistently estimated by  $\tilde{\Lambda}_i'(\sum_{i=1}^N \tilde{\Lambda}_i\tilde{\Lambda}_i')^{-1}(\sum_{i=1}^N \tilde{\Lambda}_i\tilde{\Lambda}_i'\tilde{e}_{it}^2)(\sum_{i=1}^N \tilde{\Lambda}_i\tilde{\Lambda}_i')^{-1}\tilde{\Lambda}_i$  (assuming no cross-sectional correlation in  $e_{it}$ ). Furthermore,  $\frac{1}{T}W_{NT,it}^F$  is consistently estimated by  $\frac{1}{T}\tilde{F}_t'(\sum_{s=1}^T \tilde{F}_s\tilde{F}_s'\tilde{e}_{is}^2)\tilde{F}_t$  (assuming no serial correlation in  $e_{it}$ ).

## 4.1 Implications for Factor-Augmented Regressions

The results have implications for empirical work. Consider the infeasible regression

$$y_{t+h} = \gamma' F_t^0 + \beta' W_t + \epsilon_{t+h}$$

where  $W_t$  is observed but  $F_t^0$  is not,  $\mathbb{E}(W_t \epsilon_{t+h}) = 0$  and  $\mathbb{E}(F_t^0 \epsilon_{t+h}) = 0$ . The feasible regression upon replacing  $F_t^0$  with  $\tilde{F}$  is

$$y_{t+h} = \gamma' \mathbf{H}_{NT}'^{-1} \tilde{F}_t + \beta' W_t + \epsilon_{t+h} + \gamma' \mathbf{H}_{NT}'^{-1} (\mathbf{H}_{NT}' F_t^0 - \tilde{F}_t). \quad (19)$$

Bai and Ng (2006) shows under the strong factor assumption  $\tilde{F}$  can be used in a second step regression without the need for standard error adjustments if  $\sqrt{T}/N \rightarrow 0$ . To establish the comparable conditions when  $1 \geq \alpha > 0$ , we need to analyze the correlation between the regressors  $(\tilde{F}_t, W_t)$  and the errors  $\epsilon_{t+h}$  as well as with the first step estimation error  $(\mathbf{H}_{NT}' F_t^0 - \tilde{F}_t)$ .

**Lemma 4** *Let  $\hat{z}_t = (\hat{F}_t', W_t')$  be used in place of  $z_t = (F_t', W_t')$  in the factor-augmented regression (19), and denote  $\delta^0 = (\gamma' \mathbf{H}_{NT}'^{-1}, \beta)'$ . Suppose  $\mathbf{W}' \mathbf{e} \Lambda^0 = \sum_{i=1}^N \sum_{t=1}^T W_t \Lambda_i^0 e_{it} = O_p(\sqrt{N^\alpha T})$  and Assumptions A, B, and C hold. Then*

$$\sqrt{T}(\hat{\delta} - \delta^0) \xrightarrow{d} N(0, \mathbf{J}'^{-1} \Sigma_{zz}^{-1} \Sigma_{zz, \epsilon} \Sigma_{zz}^{-1} \mathbf{J}^{-1})$$

where  $\mathbf{J}$  is the limit of  $\mathbf{J}_{NT} = \text{diag}(\mathbf{H}_{NT}', \mathbf{I}_{\dim(W)})$ .

To understand the result, notice that

$$\frac{1}{\sqrt{T}} \sum_{t=1}^T \tilde{F}_t \epsilon_{t+h} = \mathbf{H}_{NT}' \frac{1}{\sqrt{T}} \sum_{t=1}^T F_t^0 \epsilon_{t+h} + \frac{1}{\sqrt{T}} \sum_{t=1}^T (\tilde{F}_t - \mathbf{H}_{NT}' F_t^0) \epsilon_{t+h}.$$

We need to show that the second term is  $o_p(1)$  and that the first step error is negligible. More precisely we need to verify the properties of the following three terms:

$$(i.) \frac{1}{\sqrt{T}} \sum_{t=1}^T (\tilde{F}_t - \mathbf{H}_{NT}' F_t^0) \epsilon_{t+h}, \quad (ii.) \frac{1}{\sqrt{T}} \sum_{t=1}^T \tilde{F}_t (F_t^{0'} \mathbf{H}_{NT} - \tilde{F}_t'), \quad (iii.) \frac{1}{\sqrt{T}} \sum_{t=1}^T W_t (F_t^{0'} \mathbf{H}_{NT} - \tilde{F}_t').$$

But  $\frac{1}{\sqrt{T}} \epsilon' (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT, \ell}) = o_p(1)$  upon replacing  $e_i = (e_{i1}, \dots, e_{iT})'$  by  $\epsilon = (\epsilon_{1+h}, \dots, \epsilon_{T+h})$  in Lemma 6(iii). Furthermore,  $\frac{1}{\sqrt{T}} \mathbf{W}' (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT, \ell}) = o_p(1)$  upon replacing  $\mathbf{F}^0$  by  $\mathbf{W} = (W_1, \dots, W_T)'$  in Lemma 6(i). Hence all three terms are  $o_p(1)$  under the assumptions of the analysis, and  $\tilde{\mathbf{F}}$  can be used in the augmented regression as though it were  $\mathbf{F}$ . However, the conditions  $\alpha > 1/2$  and  $\sqrt{T}/N^\alpha \rightarrow 0$  are stronger than for the  $\alpha = 1$  case.



## 5 Conclusion

A sizable literature has emerged since Onatski (2012) shows that the APC estimates are inconsistent when the loadings are extremely weak in the sense that  $\frac{\Lambda'\Lambda}{N^\alpha} \rightarrow \Sigma_\Lambda$  with  $\alpha = 0$ . This note establishes conditions under which the APC estimates are consistent and asymptotically normal in more moderate cases when  $1 \geq \alpha > 0$ .

Allowing  $\alpha$  to take on a range of values naturally raises questions about the different criteria available to determine the number of factors. If we want to estimate the number of factors with  $\alpha > 0$ , the criteria in Bai and Ng (2002) remain useful. While the criteria of Onatski (2010) and Ahn and Horenstein (2013) try to better separate the bounded from the diverging eigenvalues of  $\mathbf{X}\mathbf{X}'$ , Bai and Ng (2019) seeks to isolate the factors with a tolerated level of explanatory power by singular value thresholding. This criteria will return an estimate of the ‘minimum rank’, a concept of long standing interest in classical factor analysis, see, eg. ten Berge and Kiers (1991). For documenting the number of factors with different strength, methods of Bailey et al. (2016) and Uematsu and Yamagata (2020) are available. The criteria in Freyaldenhoven (2021) determine the number factors with  $\alpha > 1/2$ . But as seen above, the  $\alpha$  required for consistent estimation of the factor space can be smaller than the one needed for asymptotic normality. Ultimately, the desired number of factors depends on the objective of the exercise and the assumptions that the researcher find defensible. It seems difficult to avoid taking a stand on what is meant by weak in practice.

## Appendix

**Lemma 5** *The following holds under Assumption A:*

$$\|\mathbf{e}'\mathbf{F}^0\|^2 \leq r\|\mathbf{F}^{0'}\mathbf{e}\mathbf{e}'\mathbf{F}^0\| = O_p(NT) \quad (20a)$$

$$\|\mathbf{e}\mathbf{\Lambda}^0\|^2 \leq r\|\mathbf{\Lambda}^{0'}\mathbf{e}'\mathbf{e}\mathbf{\Lambda}^0\| = O_p(N^\alpha T) \quad (20b)$$

$$\|\mathbf{e}\mathbf{e}'\mathbf{F}^0\| \leq \rho_{\max}(\mathbf{e}\mathbf{e}')\|\mathbf{F}^0\| = O_p(\max\{N, T\}\sqrt{T}) \quad (20c)$$

$$\|\mathbf{e}'\mathbf{e}\mathbf{\Lambda}^0\| \leq \rho_{\max}(\mathbf{e}\mathbf{e}')\|\mathbf{\Lambda}^0\| = O_p(\max\{N, T\}\sqrt{N^\alpha}) \quad (20d)$$

The matrix norms here are Frobenius norms. Since the matrices are rank  $r$ , spectral norms would give the same bounds.

**Proof of Lemma 5:** (20a) and (20b) follow from (6) and (7), respectively. For (20c),

$$\|\mathbf{e}\mathbf{e}'\mathbf{F}^0\| \leq \|\mathbf{e}\mathbf{e}'\|_{sp}\|\mathbf{F}^0\| = \rho_{\max}(\mathbf{e}\mathbf{e}')\|\mathbf{F}^0\| = O_p(\max\{N, T\}\sqrt{T})$$

here we used  $\|\mathbf{A}\mathbf{B}\| \leq \|\mathbf{A}\|_{sp}\|\mathbf{B}\|$ . The argument for (20d) is the same.

**Proof of Lemma 3 part (i)** We start with (12). We derive the order of magnitude for each of the three terms in braces.

$$\begin{aligned} \frac{\mathbf{\Lambda}^{0'}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T} &= \frac{\mathbf{\Lambda}^{0'}\mathbf{e}'\mathbf{F}^0}{N^\alpha T} + \frac{\mathbf{\Lambda}^{0'}\mathbf{e}'(\tilde{\mathbf{F}} - \mathbf{F}^0 H_{NT,0})}{N^\alpha T}, \\ \left\| \frac{\mathbf{\Lambda}^{0'}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T} \right\| &\leq \left\| \frac{\mathbf{\Lambda}^{0'}\mathbf{e}'\mathbf{F}^0}{N^\alpha T} \right\| + \left\| \frac{\mathbf{\Lambda}^{0'}\mathbf{e}'}{N^\alpha\sqrt{T}} \right\| \frac{\|(\tilde{\mathbf{F}} - \mathbf{F}^0 H_{NT,0})\|}{\sqrt{T}} \\ &= O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + O_p\left(\frac{1}{\sqrt{N^\alpha}}\right) \left[ O_p\left(\frac{1}{\sqrt{N^\alpha}}\right) + \frac{1}{T} \frac{N}{N^\alpha} O_p(1) \right] \\ &= O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + O_p\left(\frac{1}{N^\alpha}\right) + \frac{1}{T} \frac{N}{N^{3\alpha/2}} O_p(1), \end{aligned}$$

where we have used Assumption A3(v), (20b), and Proposition 1. Next (9) implies

$$\left\| \frac{\tilde{\mathbf{F}}'\mathbf{e}\mathbf{e}'\tilde{\mathbf{F}}}{N^\alpha T^2} \right\| = \left(\frac{N}{N^\alpha}\right) \frac{1}{T} O_p(1) + \frac{1}{N^\alpha} O_p(1).$$

Collecting the non-dominated terms, we obtain

$$\left(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right)\mathbf{H}_{NT,0} = \mathbf{I}_r + O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + O_p\left(\frac{1}{N^\alpha}\right) + \left(\frac{N}{N^\alpha}\right) \frac{1}{T} O_p(1).$$

Equivalently,

$$\mathbf{H}_{NT,0} = \left(\frac{\tilde{\mathbf{F}}'\mathbf{F}^0}{T}\right)^{-1} + O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + O_p\left(\frac{1}{N^\alpha}\right) + \left(\frac{N}{N^\alpha}\right) \frac{1}{T} O_p(1).$$

**Proof of Lemma 3 part (ii):** Part (i) makes clear that the approximations for  $\mathbf{H}_{NT,0}$  depends on  $\alpha$ . Part (ii) follows from part (i) and  $\mathbf{H}_{NT,0} \xrightarrow{p} \mathbf{Q}^{-1}$ . For part (ii), we already proved the case of  $\ell = 3$ . It remains to show asymptotic equivalence of the three remaining matrices to  $\mathbf{H}_{NT,0}$ .

We begin with  $\ell = 1$  where  $\mathbf{H}_{NT,1} = (\mathbf{\Lambda}^{0'} \mathbf{\Lambda}^0 / N^\alpha) (\tilde{\mathbf{\Lambda}}' \mathbf{\Lambda}^0 / N^\alpha)^{-1}$ . From  $\tilde{\mathbf{\Lambda}}' = \tilde{\mathbf{F}}' \mathbf{X} / T = \tilde{\mathbf{F}}' (\mathbf{F}^0 \mathbf{\Lambda}^{0'} + \mathbf{e}) / T$ , we have

$$\frac{\tilde{\mathbf{\Lambda}}' \mathbf{\Lambda}^0}{N^\alpha} = \frac{\tilde{\mathbf{F}}' \mathbf{F}^0 \mathbf{\Lambda}^{0'} \mathbf{\Lambda}^0}{T N^\alpha} + \frac{\tilde{\mathbf{F}}' \mathbf{e} \mathbf{\Lambda}^0}{T N^\alpha} = \frac{\tilde{\mathbf{F}}' \mathbf{F}^0 \mathbf{\Lambda}^{0'} \mathbf{\Lambda}^0}{T N^\alpha} + \frac{1}{\sqrt{T N^\alpha}} O_p(1). \quad (21)$$

Thus  $\left( \frac{\tilde{\mathbf{\Lambda}}' \mathbf{\Lambda}^0}{N^\alpha} \right)^{-1} = \left( \frac{\mathbf{\Lambda}^{0'} \mathbf{\Lambda}^0}{N^\alpha} \right)^{-1} \left( \frac{\tilde{\mathbf{F}}' \mathbf{F}^0}{T} \right)^{-1} + \frac{1}{\sqrt{T N^\alpha}} O_p(1)$ , implying

$$\mathbf{H}_{NT,1} = \left( \frac{\tilde{\mathbf{F}}' \mathbf{F}^0}{T} \right)^{-1} + \frac{1}{\sqrt{T N^\alpha}} O_p(1).$$

For  $\ell = 4$ , taking the transpose of (21) gives

$$\begin{aligned} \mathbf{H}_{NT,4} &= \frac{\tilde{\mathbf{\Lambda}}' \mathbf{\Lambda}^0}{N^\alpha} \frac{N^\alpha}{N} \mathbf{D}_{NT,r}^{-2} = \left( \frac{\mathbf{\Lambda}^{0'} \mathbf{\Lambda}^0}{N^\alpha} \right) \left( \frac{\mathbf{F}^0 \tilde{\mathbf{F}}}{T} \right) \frac{N^\alpha}{N} \mathbf{D}_{NT,r}^{-2} + \frac{1}{\sqrt{T N^\alpha}} O_p(1) \\ &= \mathbf{H}_{NT,0} + \frac{1}{\sqrt{T N^\alpha}} O_p(1). \end{aligned}$$

Finally, for  $\ell = 2$ , we use the expression for  $\tilde{\mathbf{F}}$  in (11) to obtain

$$\mathbf{H}_{NT,2} - \mathbf{H}_{NT,0} \equiv \left( \frac{\mathbf{F}^{0'} \mathbf{F}^0}{T} \right)^{-1} \left( \frac{\mathbf{F}^{0'} \mathbf{F}^0 \mathbf{\Lambda}^{0'} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\mathbf{F}^{0'} \mathbf{e} \mathbf{\Lambda}^0 \mathbf{F}^{0'} \tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\mathbf{F}^{0'} \mathbf{e} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} \right) \left( \frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2 \right)^{-1}$$

The right hand side is bounded by  $O_p(\frac{1}{\sqrt{N^\alpha T}}) + O_p(\frac{1}{N^\alpha}) + (\frac{N}{N^\alpha}) \frac{1}{T} O_p(1)$ .  $\square$

**Proof of Proposition 2.** As argued in the text, it remains to analyze the term  $\frac{\|(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0})' \mathbf{e}\|}{T \sqrt{N}}$ .

Notice

$$\frac{\mathbf{e}' (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0})}{T} = \left( \frac{\mathbf{e}' \mathbf{F}^0 \mathbf{\Lambda}^{0'} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\mathbf{e}' \mathbf{e} \mathbf{\Lambda}^0 \mathbf{F}^{0'} \tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\mathbf{e}' \mathbf{e} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} \right) \left( \frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2 \right)^{-1}$$

Now

$$\begin{aligned} \left\| \frac{\mathbf{e}' \mathbf{F}^0 \mathbf{\Lambda}^{0'} \mathbf{e}' \mathbf{F}^0}{N^\alpha T^2} \right\| &\leq \frac{1}{N^\alpha T^2} \|\mathbf{e}' \mathbf{F}^0\| \cdot \|\mathbf{\Lambda}^{0'} \mathbf{e}' \mathbf{F}^0\| = \frac{1}{T} \sqrt{\frac{N}{N^\alpha}} O_p(1) \\ \left\| \frac{\mathbf{e}' \mathbf{e} \mathbf{\Lambda}^0}{N^\alpha T} \right\| &= \frac{\max\{N, T\} \sqrt{N^\alpha}}{N^\alpha T} O_p(1) \leq \frac{N}{\sqrt{N^\alpha} T} O_p(1) + \frac{1}{\sqrt{N^\alpha}} O_p(1) \\ \left\| \frac{\mathbf{e}' \mathbf{e} \mathbf{e}' \mathbf{F}^0}{N^\alpha T^2} \right\| &\leq \frac{1}{N^\alpha T^2} O_p(\max\{N, T\}) \|\mathbf{e}' \mathbf{F}^0\| = \frac{1}{N^\alpha T^2} O_p(\max\{N, T\}) O_p(\sqrt{NT}) \\ &= \frac{N^{3/2-\alpha}}{T^{3/2}} O_p(1) + \sqrt{\frac{N}{T}} \frac{1}{N^\alpha} O_p(1). \end{aligned}$$

Collecting the non-dominating terms, we obtain

$$\frac{\|(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0})' \mathbf{e}\|}{T} \leq \frac{N}{\sqrt{N^\alpha}} \frac{1}{T} O_p(1) + \frac{1}{\sqrt{N^\alpha}} O_p(1) + \frac{N^{3/2-\alpha}}{T^{3/2}} O_p(1) + \sqrt{\frac{N}{T}} \frac{1}{N^\alpha} O_p(1)$$

So

$$\frac{\|(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0})' \mathbf{e}\|}{T\sqrt{N}} \leq \frac{1}{T} \sqrt{\frac{N}{N^\alpha}} O_p(1) + \frac{1}{\sqrt{N^{1+\alpha}}} + \frac{1}{T^{3/2}} \left(\frac{N}{N^\alpha}\right) O_p(1) + \frac{1}{N^\alpha \sqrt{T}} O_p(1).$$

**Lemma 6** *Suppose that Assumption A holds. We have, for  $\ell = 0, 1, 2, 3, 4$ ,*

- i  $\frac{1}{T} \mathbf{F}^{0'} (\tilde{\mathbf{F}} - \mathbf{F}^{0'} \mathbf{H}_{NT,\ell}) = O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + O_p\left(\frac{N}{N^\alpha T}\right) + O_p\left(\frac{1}{N^\alpha}\right) + \frac{1}{N^{2\alpha}} \left(\frac{N}{T}\right)^{3/2} O_p(1) + \frac{1}{N^{2\alpha}} \left(\frac{N}{T}\right)^{1/2} O_p(1)$
- ii  $\frac{1}{N^\alpha} \boldsymbol{\Lambda}^{0'} (\tilde{\boldsymbol{\Lambda}} - \boldsymbol{\Lambda}^0 \mathbf{H}'_{NT,\ell}) = O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right) + O_p\left(\frac{N}{N^\alpha T}\right) + O_p\left(\frac{1}{N^\alpha}\right) + \frac{1}{N^{2\alpha}} \left(\frac{N}{T}\right)^{3/2} O_p(1) + \frac{1}{N^{2\alpha}} \left(\frac{N}{T}\right)^{1/2} O_p(1)$
- iii  $\frac{1}{T} e'_i (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,\ell}) = O_p\left(\frac{1}{N^\alpha}\right) + O_p\left(\frac{1}{\sqrt{TN^\alpha}}\right) + O_p\left(\frac{N}{N^\alpha T}\right)$ , for each  $i$ ,
- iv  $\frac{1}{N} e'_t (\tilde{\boldsymbol{\Lambda}} - \boldsymbol{\Lambda}^0 \mathbf{H}'_{NT,\ell}) = \frac{1}{T} \sqrt{\frac{N}{N^\alpha}} O_p(1) + \frac{1}{\sqrt{N^{1+\alpha}}} O_p(1) + \frac{1}{T^{3/2}} \left(\frac{N}{N^\alpha}\right) O_p(1) + \frac{1}{N^\alpha \sqrt{T}} O_p(1)$ , for each  $t$ .

The same bounds in parts (i) and (ii) hold for  $\frac{1}{T} \tilde{\mathbf{F}}' (\tilde{\mathbf{F}} - \mathbf{F}^{0'} \mathbf{H}_{NT,\ell})$  and  $\frac{1}{N^\alpha} \tilde{\boldsymbol{\Lambda}}' (\tilde{\boldsymbol{\Lambda}} - \boldsymbol{\Lambda}^0 \mathbf{H}'_{NT,\ell})$ . This follows from, for example,  $\frac{1}{T} \tilde{\mathbf{F}}' (\tilde{\mathbf{F}} - \mathbf{F}^{0'} \mathbf{H}_{NT,\ell}) = \mathbf{H}'_{NT} \frac{1}{T} \mathbf{F}^{0'} (\tilde{\mathbf{F}} - \mathbf{F}^{0'} \mathbf{H}_{NT,\ell}) + \frac{1}{T} \|\tilde{\mathbf{F}} - \mathbf{F}^{0'} \mathbf{H}_{NT,\ell}\|^2$  and Proposition 1. In the strong factor case, all four quantities are  $O_p(\delta_{NT}^{-2})$ . The rates now depend on  $\alpha$  and affect the convergence rates.

**Proof of Lemma 6.** Consider (i).

$$\frac{1}{T} \mathbf{F}^{0'} (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0}) = \left[ \left( \frac{\mathbf{F}^{0'} \mathbf{F}^0}{T} \right) \left( \frac{\boldsymbol{\Lambda}^{0'} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T} \right) + \left( \frac{\mathbf{F}^{0'} \mathbf{e} \boldsymbol{\Lambda}^0}{N^\alpha T} \right) \frac{\mathbf{F}^{0'} \tilde{\mathbf{F}}}{T} + \frac{\mathbf{F}^{0'} \mathbf{e} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} \right] O_p(1)$$

Consider the first term in the bracket

$$\frac{\boldsymbol{\Lambda}^{0'} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T} = \frac{\boldsymbol{\Lambda}^{0'} \mathbf{e}' \mathbf{F}^0 \mathbf{H}}{N^\alpha T} + \frac{\boldsymbol{\Lambda}^{0'} \mathbf{e}' (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H})}{N^\alpha T}$$

the first term is  $O_p\left(\frac{1}{\sqrt{N^\alpha T}}\right)$ . The second term is bounded by

$$\begin{aligned} \left\| \frac{\boldsymbol{\Lambda}^{0'} \mathbf{e}' (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H})}{N^\alpha T} \right\| &\leq \frac{1}{\sqrt{N^\alpha}} \left( \frac{\|\boldsymbol{\Lambda}^0\|}{\sqrt{N^\alpha}} \right) \left\| \frac{\mathbf{e}' (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H})}{T} \right\| \\ &= O_p \left( \left( \frac{N}{N^\alpha} \frac{1}{T} \right) + \frac{1}{N^\alpha} + \left( \frac{N}{T} \right)^{3/2} \frac{1}{N^{2\alpha}} + \left( \frac{N}{T} \right)^{1/2} \frac{1}{N^{2\alpha}} \right) \end{aligned} \quad (22)$$

Next

$$\begin{aligned}
\left\| \frac{\mathbf{F}^{0'} \mathbf{e} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} \right\| &\leq \left\| \frac{\mathbf{F}^{0'} \mathbf{e} \mathbf{e}' \mathbf{F}^0}{N^\alpha T^2} \right\| + \left\| \frac{\mathbf{F}^{0'} \mathbf{e}}{N^\alpha T} \right\| \left\| \frac{\mathbf{e}'(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H})}{T} \right\| \\
&= \frac{1}{T} O_p\left(\frac{N}{N^\alpha}\right) + \frac{1}{N^\alpha} O_p\left(\sqrt{\frac{N}{T}}\right) \left\| \frac{\mathbf{e}'(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H})}{T} \right\| \\
&= \frac{1}{T} O_p\left(\frac{N}{N^\alpha}\right) + \frac{1}{N^\alpha} O_p\left(\sqrt{\frac{N}{N^\alpha T}}\right)
\end{aligned} \tag{23}$$

Collecting the non-dominated terms, we obtain (i). None of these terms can be dominated by others, all depending on the relative magnitude of  $N$  and  $T$ , and when  $\alpha = 1$ , they simplify to  $O_p(\delta_{NT}^{-2})$ .

Consider (ii).

$$\frac{1}{N^\alpha} (\tilde{\mathbf{\Lambda}} - \mathbf{\Lambda}^0 \mathbf{H}'_{NT,3})' \mathbf{\Lambda}^0 = \mathbf{H}'_{3,NT} \mathbf{F}^{0'} \mathbf{e} \mathbf{\Lambda}^0 / (N^\alpha T) + (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,3})' \mathbf{e} \mathbf{\Lambda}^0 / (N^\alpha T)$$

the first term is  $O_p(\frac{1}{\sqrt{N^\alpha T}})$  by Assumption A3(v), the second term is analyzed in (22). This proves (ii).

Consider (iii)

$$\frac{\mathbf{e}'_i (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,0})}{T} = \left( \frac{\mathbf{e}'_i \mathbf{F}^0 \mathbf{\Lambda}^0' \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\mathbf{e}'_i \mathbf{e} \mathbf{\Lambda}^0 \mathbf{F}^{0'} \tilde{\mathbf{F}}}{N^\alpha T^2} + \frac{\mathbf{e}'_i \mathbf{e} \mathbf{e}' \tilde{\mathbf{F}}}{N^\alpha T^2} \right) \left( \frac{N}{N^\alpha} \mathbf{D}_{NT,r}^2 \right)^{-1}$$

The first term can be ignored. The second term is determined by

$$\frac{\mathbf{e}'_i \mathbf{e} \mathbf{\Lambda}^0}{N^\alpha T} = O_p\left(\frac{1}{N^\alpha}\right) + O_p\left(\frac{1}{\sqrt{TN^\alpha}}\right)$$

by Assumption A3(ii). The third term is  $O_p(\frac{1}{N^\alpha} + \frac{N}{N^\alpha T})$ . Collecting terms gives (iii).

Consider (iv).

$$\frac{1}{N} (\tilde{\mathbf{\Lambda}} - \mathbf{\Lambda}^0 \mathbf{H}'_{NT,3})' \mathbf{e}_t = \mathbf{H}'_{3,NT} \mathbf{F}^{0'} \mathbf{e} \mathbf{e}_t / (NT) + (\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,3})' \mathbf{e} \mathbf{e}_t / (NT)$$

First term is  $O_p(\delta_{NT}^{-2})$  by Assumption A3. The second term is bounded by

$$\frac{\|(\tilde{\mathbf{F}} - \mathbf{F}^0 \mathbf{H}_{NT,3})' \mathbf{e}\| \|e_t\|}{T \sqrt{N}} = \frac{1}{T} \sqrt{\frac{N}{N^\alpha}} O_p(1) + \frac{1}{\sqrt{N^{1+\alpha}}} O_p(1) + \frac{1}{T^{3/2}} \left(\frac{N}{N^\alpha}\right) O_p(1) + \frac{1}{N^\alpha \sqrt{T}} O_p(1)$$

This proves (iv).  $\square$

## References

- Ahn, S. and Horenstein, R. 2013, Eigenvalue Ratio Test for the Number of Factors, *Econometrica* **81:3**, 1203–1227.
- Anderson, T. W. and Rubin, H. 1956, Statistical Inference in Factor Analysis, in J. Neyman (ed.), *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*, Vol. V, Berkeley: University of California Press, pp. 114–150.
- Bai, J. 2003, Inferential Theory for Factor Models of Large Dimensions, *Econometrica* **71:1**, 135–172.
- Bai, J. and Ng, S. 2002, Determining the Number of Factors in Approximate Factor Models, *Econometrica* **70:1**, 191–221.
- Bai, J. and Ng, S. 2006, Confidence Intervals for Diffusion Index Forecasts and Inference with Factor-Augmented Regressions, *Econometrica* **74:4**, 1133–1150.
- Bai, J. and Ng, S. 2019, Rank Regularized Estimation of Approximate Factor Models, *Journal of Econometrics* **78-96**, 212:1.
- Bailey, N., Kapetanios, G. and Pesaran, H. 2016, Exponent of Cross-Section Dependence: Estimation and Inference, *Journal of Applied Econometrics* **31**, 929–960.
- Chamberlain, G. and Rothschild, M. 1983, Arbitrage, Factor Structure and Mean-Variance Analysis in Large Asset Markets, *Econometrica* **51**, 1281–2304.
- DeMol, C., Giannone, D. and Reichlin, L. 2008, Forecasting Using a Large Number of Predictors: Is Bayesian Regression a Valid Alternative to Principal Components, *Journal of Econometrics* **146**, 318–328.
- Eckart, C. and Young, G. 1936, The Approximation of One Matrix by Another of Lower Rank, *Psychometrika*.
- Forni, M., Hallin, M., Lippi, M. and Reichlin, L. 2000, The Generalized Dynamic Factor Model: Identification and Estimation, *Review of Economics and Statistics* **82:4**, 540–554.
- Freyaldenhoven, S. 2021, Factor Models with Local Factors: Determining the Number of Relevant Factors, *Journal of Econometrics*.

- Lettau, M. and Pelger, M. 2020, Estimating Latent Asset Pricing Factors, *Journal of Econometrics* **218:1**, 1–31.
- Moon, R. and Weidner, M. 2017, Dynamic Linear Panel Regression Models with Interactive Fixed Effects, *Econometric Theory* **33**, 158–195.
- Onatski, A. 2010, Determining the Number of Factors From Empirical Distribution of Eigenvalues, *Review of Economics and Statistics* **92:4**, 1004–1016.
- Onatski, A. 2012, Asymptotics of the Principal Components Estimator of Large Factor Models with Weakly Influential Factors, *Journal of Econometrics* **168**, 244–258. manuscript under revision.
- Stock, J. H. and Watson, M. W. 1998, Diffusion Indexes, NBER Working Paper 6702.
- Stock, J. H. and Watson, M. W. 2002, Forecasting Using Principle Components from a Large Number of Predictors, *Journal of American Statistical Association* **97**(460), 1167–1179.
- ten Berge, J. and Kiers, H. 1991, A Numerical Approach to the Exact and the Approximate Minimum Rank of a Covariance Matrix, *Psychometrika* **56**, 309–315.
- Uematsu, Y. and Yamagata, T. 2019, Estimation of Weak Factors, ISER Discussion Paper 1053.
- Uematsu, Y. and Yamagata, T. 2020, Inference of Weak Factors, ISER Discussion Paper 1080.