

SOME IMPOSSIBILITY RESULTS FOR INFERENCE WITH CLUSTER DEPENDENCE WITH LARGE CLUSTERS

Denis Kojevnikov and Kyungchul Song
Tilburg University and University of British Columbia

September 9, 2022

ABSTRACT. This paper focuses on a setting with observations having a cluster dependence structure, and presents two main impossibility results. First, we show that when there is only one large cluster, i.e., the researcher does not have any knowledge on the dependence structure of the observations, it is not possible to consistently discriminate the mean. Focusing on a Gaussian experiment with large clusters where the within-cluster long run variances are nearly identical, we also show that the necessary and sufficient condition for consistent \sqrt{n} -discrimination of the mean is that we have at least two clusters. This result shows some limitations for inference when we lack information on the dependence structure, i.e., information on which groups of observations exhibit strong within-group dependence. Our second result provides a necessary and sufficient condition for the cluster structure that the long run variance is consistently estimable. Our result implies that when there is at least one large cluster, the long run variance is not consistently estimable.

KEY WORDS. Consistent Discrimination; Local Dependence; Unknown Dependence Structure; Consistent Estimation of Long-Run Variance; Cluster Dependence; Log Likelihood Process

JEL CLASSIFICATION: C01, C12, C13

We thank Michael Leung and James MacKinnon for valuable comments. We thank the Associate Editor and three anonymous referees for their constructive comments which helped us improve our paper's exposition and results. All errors are ours. Song acknowledges financial support from Social Sciences and Humanities Research Council of Canada. Corresponding Address: Kyungchul Song, Vancouver School of Economics, University of British Columbia, 6000 Iona Drive, Vancouver, V6T 1L4, Canada.

1. Introduction

Statistical inference from data usually begins by imposing a form of a dependence structure on the data, by specifying which groups of observations exhibit strong within-group dependence. Various tools of asymptotic inference such as law of large numbers and central limit theorem are available under the imposed dependence structure. A standard case is the independence assumption or an assumption on time series dependence. However, it is well known that in the case of cross-sectional dependence, a researcher is often less confident about the correctness of the dependence structure used, despite its crucial role for inference.

A popular way to deal with this challenge is to use cluster dependence modeling, where the dependence structure among observations within each cluster is left unspecified, while independence is imposed between observations from different clusters. The inference procedures when there are many clusters are well known and can be analyzed using standard methods of asymptotic inference. However, less is known about the case where there are large clusters, and the dependence structure within such a cluster is unknown. [Cameron, Gelbach, and Miller \(2008\)](#) proposed a wild bootstrap procedure and showed by simulations that their tests perform well even when there are a small number of clusters. The robustness of this result was confirmed by [MacKinnon and Webb \(2017\)](#) even when the sizes of the clusters are highly heterogeneous. This large cluster issue has also drawn interest in the literature of difference-in-differences when there are only few treated clusters. (See [Conley and Taber \(2011\)](#), [Hagemann \(2019\)](#), and [MacKinnon and Webb \(2020\)](#), and references therein.) [Djogbenou, MacKinnon, and Nielsen \(2019\)](#) studied inference on regression models with clustered errors. They provided conditions for the cluster sizes so that asymptotic and bootstrap inferences are asymptotically valid. They showed that their conditions exclude the presence of a large cluster.

There are several methods proposed to deal with the problem of inference with large clusters. [Donald and Lang \(2007\)](#) and [Bester, Conley, and Hansen \(2011\)](#) considered linear models and proposed inference where the asymptotic distribution of the long run variance estimator is fully known. This approach is related to the HAR (Heteroskedasticity-Autocorrelation Robust) inference of [Kiefer and Vogelsang \(2002\)](#) and [Sun \(2014\)](#) in time series, which uses a normalization by an inconsistent long run variance estimator that has a stochastic limit.

[Ibragimov and Müller \(2010\)](#) proposed a t -test approach based on within-cluster estimators together with a t -distribution, where the degree of freedom in the t distribution is determined by the number of clusters. They used the result of [Bakirov and Szekely \(2005\)](#) and showed that their approach is asymptotically valid, even if the variances of the cluster specific estimators are different across the clusters. [Ibragimov and Müller \(2016\)](#) extended these results to the problem of two-sample comparison and developed a testing procedure for the level of clustering.

Some studies adopted the approach of randomized testing to deal with cluster dependence with large clusters. [Canay, Romano, and Shaikh \(2017\)](#) developed asymptotic inference procedures when the inference involves statistics whose limiting distribution satisfy symmetry properties. [Hagemann \(2019\)](#) proposed randomized tests for treatment effects when there are only a small number of clusters. Like [Ibragimov and Müller \(2010\)](#), both proposals assumed large sample properties for within-cluster statistics. A recent work by [Canay, Santos, and Shaikh \(2021\)](#) use the analogue between wild bootstrap and randomized tests, and provided conditions under which the wild bootstrap for cluster-dependent regression models is asymptotically valid when there are only a small number of clusters.

Our paper focuses on observations with a cluster dependence structure and explores implications on statistical inference when there are large clusters. First, we show that when the sample consists of large clusters, the mean cannot be consistently discriminated if there is only one cluster, i.e., the researcher does not have any knowledge on the dependence structure of the data. Furthermore, when the observations form large clusters, a necessary and sufficient condition for the mean to be consistently discriminated at the rate of \sqrt{n} is that the sample consists of at least two large clusters.

This impossibility result has a significant implication in a setting where the researcher does not know the dependence structure of observations. Note that [Song \(2016\)](#) proposed a randomized subsampling approach, and [Leung \(2021\)](#) provided a set of general conditions for the approach to produce asymptotically valid inference, which does not require any knowledge on the dependence structure. Our result shows that part of those conditions implies consistent discrimination of the mean, and hence imposes more conditions than those permitted via lack of knowledge on the dependence structure.

Our second result is concerned with consistent estimation of long run variances. More specifically, suppose that $X_n = [X_{n,1}, \dots, X_{n,n}]^\top$ is a given random vector of dimension n , where each observation $X_{n,i}$ has the same mean μ . Let us define the long-run variance of X_n as¹

$$(1.1) \quad \sigma_{LR}^2 = \text{Var} \left(\frac{1}{\sqrt{n}} \sum_{i=1}^n X_{n,i} \right).$$

Recently, [Hansen and Lee \(2019\)](#) derived an asymptotic distribution theory for clustered data, including a law of large numbers and a central limit theorem. One of their results presents a condition for the cluster structure that is necessary and sufficient for the weak law of large numbers to hold for the sample average of the clustered observations. Our paper shows that the same condition is in fact necessary and sufficient for the consistent estimability of the long

¹Note that when there is a common shock, say, C_n , such as cluster-specific fixed effects with few clusters, the analysis in this paper carries over to this case with σ_{LR}^2 replaced by the conditional variance given common shock C_n . Our impossibility results do not depend on whether there is a common shock of this form in the data or not. For simplicity, we consider a setting without such cluster-specific fixed effects.

run variance of the clustered observations as well. Our condition for the cluster structure also implies that when there is at least one large cluster, i.e., the researcher does not know the dependence structure on a nonnegligible portion of the data, the long-run variance is not consistently estimable. It is not hard to show that the existing cluster-robust variance estimators are inconsistent when the cluster structure is severely misspecified. However, to the best of our knowledge, it has not been known whether there exists any consistent estimator of the long run variance when there is a lack of knowledge on the dependence structure on a nonnegligible portion of the data. Our result gives a negative answer to this question.

There has long been a strand of literature that studies impossibility of estimation and inference. (e.g., [Bahadur and Savage \(1956\)](#), [Dufour \(1997\)](#), [Pötscher \(2002\)](#), [Bertanha and Moreira \(2020\)](#).) Our impossibility of consistent estimation of long run variance is related to [Pötscher \(2002\)](#) who established a minimax risk lower bound for a general estimation problem. His result implies impossibility of consistent estimation uniform in P , and unbounded confidence sets in various estimation settings in practice. Similarly as in [Pötscher \(2002\)](#), our impossibility result comes from the parameter of interest (i.e., the long-run variance here) behaving “discontinuously” in response to a contiguous change of the joint distribution of random variables. However, unlike [Pötscher \(2002\)](#), our probability model is not indexed by a set of parameters fixed independently of the sample size, such as \mathcal{H} in his paper, because here we have to deal with the *joint distribution of the entire sample* whose dependence structure varies in the model, not a representative marginal probability as in settings with i.i.d. or stationary random variables. [Bertanha and Moreira \(2020\)](#) studied impossibility results of two types: nondistinguishability of the null hypothesis from the alternative hypothesis and unbounded confidence sets. Their study of impossibility of the first type is related to impossibility of consistent discrimination of the mean in our paper. However, in our setting, while consistent discrimination of the mean is not possible in the absence of knowledge on the dependence structure, it is still possible to have a test with maximal power greater than the size. Hence, our result does not fall into their framework. [Menzel \(2021\)](#) recently developed and verified the validity of a bootstrap procedure in multi-way clustered observations with two or more dimensions. Part of his results shows that it is not possible to consistently estimate the distribution of the cluster dependent observations. Our results are not the special case of his results, because our impossibility result holds for models that exclude the counterexample that he used to prove the impossibility result. In particular, our cluster dependence accommodates within-cluster heterogeneity in terms of marginal distributions and dependence structures.

The rest of the paper is organized as follows. The next section studies the consistent discrimination of the mean. Section 3 is devoted to presenting the result of the impossibility of consistent estimation of the long run variance. In Section 4, we illustrate the implication of

our results for the case of difference-in-difference models. In Section 5, we conclude. The mathematical proofs are found in the appendix.

2. Gaussian Experiments

Let $X_n = [X_{n,1}, \dots, X_{n,n}]^\top$ be a random vector with a joint distribution P_n which belongs to the class of distributions \mathcal{P}_n . Throughout the paper, we assume that for each $P_n \in \mathcal{P}_n$,

$$\mathbf{E}[X_{n,1}] = \mathbf{E}[X_{n,2}] = \dots = \mathbf{E}[X_{n,n}].$$

In many situations, the dependence structure is partially observed. Here we consider cluster dependence, where the dependence structure is entirely unknown within each cluster, and observations are independent between clusters. Let $N_{n,m}, m = 1, \dots, M_n$, be a partition of $N_n = \{1, \dots, n\}$ such that $|N_{n,m}| = n_m$ for each $m = 1, \dots, M_n$, so that $\sum_{m=1}^{M_n} n_m = n$. Define $\mathcal{M}_n = \{N_{n,m} : m = 1, \dots, M_n\}$ and call it a **cluster structure**. Throughout the paper we assume that $(X_{n,i})_{i \in N_{n,m}}$ are independent across m 's under all $P_n \in \mathcal{P}_n$, i.e., the joint distribution of X_n has a cluster dependence structure.

Our impossibility results rely on the assumption that the probability model, \mathcal{P}_n , includes Gaussian experiments. It is convenient for us to use Gaussian experiments because we have an explicit finite sample form of the log likelihood ratio of the random vector X_n . Since we do not assume independence or time series dependence, we cannot rely on the standard procedure based on the local asymptotic normality or local asymptotic mixed normality for the log likelihood process for X_n .

We say that an $n \times n$ matrix is **block diagonal along the cluster structure** \mathcal{M}_n , if it is block diagonal and the m -th block of the matrix for each $m = 1, \dots, M_n$ is an $n_m \times n_m$ matrix and has row and column indices in $N_{n,m}$. Let \mathcal{S}_n^0 be the set of symmetric matrices with zero diagonals, and let \mathcal{S}_n^+ be the set of symmetric and positive definite matrices, where matrices in \mathcal{S}_n^0 and \mathcal{S}_n^+ are all block diagonal along \mathcal{M}_n .

For each $\Sigma_n \in \mathcal{S}_n^+$, we let $\Phi(\mu_n, \Sigma_n)$ be the multivariate normal distribution with mean $\mu_n \in \mathbf{R}^n$, and covariance matrix Σ_n . Define

$$\mathcal{P}_{n,\mathcal{N}} = \left\{ \Phi(\bar{\mu}_n \mathbf{1}_n, \Sigma_n) : \Sigma_n = I_n + \Delta_n, \Delta_n \in \mathcal{S}_n^0, \Sigma_n \in \mathcal{S}_n^+, \bar{\mu}_n \in \mathbf{R} \right\},$$

where $\mathbf{1}_n$ denotes the n -dimensional vector of ones. The set $\mathcal{P}_{n,\mathcal{N}}$ is the set of multivariate normal distributions with a common mean and a covariance matrix which is a deviation from I_n by a symmetric matrix with zero diagonals. As we focus on the role of knowledge of the dependence structure, we do not consider variations in the marginal variances. Thus, we take Δ_n to have zero diagonal entries.

Throughout the paper, we assume that $\mathcal{P}_{n,\mathcal{N}} \subset \mathcal{P}_n$. That is, the researcher knows that the joint distribution of X_n belongs to \mathcal{P}_n and does not exclude the possibility that it is a Gaussian distribution.

3. Consistent Discrimination of the Mean

3.1. Consistent Discrimination of the Mean

Let us explore the consistent discrimination of the mean under the general cluster dependence structure. We introduce the notion of consistent discrimination formally. Let \mathcal{P}_n be a set of the distributions of $X_n \in \mathbf{R}^n$ such that $\mathbf{E}[X_{n,i}]$ is identical across i for each $n \geq 1$. Let $\mathcal{P}_{n,0} = \{P_n \in \mathcal{P}_n : \mathbf{E}[X_{n,i}] = 0\}$, i.e., the set of probabilities under the null hypothesis of $\mathbf{E}[X_{n,i}] = 0$.

Definition 3.1. The mean of $X_{n,i}$, $n \geq 1$, is **consistently discriminated at level $\alpha \in (0, 1)$ in model \mathcal{P}_n** , if there is a sequence of (potentially randomized) tests $\{\varphi_n(X_n)\}_{n \geq 1}$ such that $\limsup_{n \rightarrow \infty} \mathbf{E}[\varphi_n(X_n)] \leq \alpha$, along any sequence $P_{n,0}$, and $\liminf_{n \rightarrow \infty} \mathbf{E}[\varphi_n(X_n)] = 1$, along any sequence $P_n \in \mathcal{P}_n$ such that $\liminf_{n \geq 1} \mathbf{E}[X_{n,i}] > 0$.

The following theorem shows that when the sample consists of nonnegligible clusters, a necessary condition for the consistent discrimination of the mean at the parametric rate is that there exist at least two clusters.

Theorem 3.1. *Suppose that $\mathcal{P}_{n,\mathcal{N}} \subset \mathcal{P}_n$. Suppose further that $\alpha \in (0, 1/2)$, and $M_n = 1$. Then, the mean cannot be consistently discriminated at level α .*

The theorem implies that when we do not know the local dependence structure of the random variables (i.e., $M_n = 1$), it is not possible to consistently discriminate the mean. In fact, as the proof of the theorem shows, the maximum power against any alternative hypothesis in $\mathcal{P}_{n,\mathcal{N}}$ is bounded by 2α . Hence, it is necessary that $M_n \geq 2$ for there to be a consistent test for the mean.

3.2. Consistent \sqrt{n} -Discrimination of the Mean

We introduce the following notion of consistent \sqrt{n} -discrimination.

Definition 3.2. The mean of $X_{n,i}$, $n \geq 1$, is **consistently \sqrt{n} -discriminated at level $\alpha \in (0, 1)$ in model \mathcal{P}_n** , if there is a sequence of (potentially randomized) tests $\{\varphi_n(X_n)\}_{n \geq 1}$ such that

$$\limsup_{n \rightarrow \infty} \mathbf{E}[\varphi_n(X_n)] \leq \alpha,$$

along any sequence $P_n \in \mathcal{P}_{n,0}$, and

$$\liminf_{n \rightarrow \infty} \mathbf{E}[\varphi_n(X_n)] = 1,$$

along any sequence $P_n \in \mathcal{P}_n$ such that $\lim_{n \rightarrow \infty} \sqrt{n} \mathbf{E}[X_{n,i}] / \sigma_{LR} = \infty$.

We consider consistent discrimination against alternatives after normalizing by σ_{LR} (which depends on n), so that when σ_{LR}^2 is larger, we focus on the alternative hypothesis that is farther away from the null hypothesis. Hence, if it is not possible to consistently \sqrt{n} -discriminate the mean, it is not necessarily due to the long run variance increasing to infinity fast.

The consistent \sqrt{n} -discrimination is often obtained when the parameter is in a finite dimensional space, and one knows the local dependence structure of the observations. To illustrate this point, suppose that $X_{n,1}, \dots, X_{n,n}$'s are i.i.d. Then, often we have

$$\frac{\sqrt{n}(\bar{X}_n - \mathbf{E}[X_{n,i}])}{\hat{\sigma}_n} \rightarrow_d \mathcal{N}(0, 1),$$

where $\hat{\sigma}_n^2 \rightarrow_p \sigma^2 = \text{Var}(X_{n,1}) > 0$, and $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_{n,i}$. Let us consider the usual t -test as follows:

$$\varphi_n(X_n) = 1 \left\{ \frac{\sqrt{n}\bar{X}_n}{\hat{\sigma}_n} > z_{1-\alpha} \right\}.$$

Under the Pitman local alternatives such that $\mathbf{E}[X_{n,i}] / \sigma = \bar{\mu} / \sqrt{n}$, $\bar{\mu} > 0$, we have

$$\liminf_{n \rightarrow \infty} \mathbf{E}[\varphi_n(X_n)] = 1 - \Phi(z_{1-\alpha} - \bar{\mu}),$$

where Φ denotes the CDF of $\mathcal{N}(0, 1)$. The last term converges to 1 as $\bar{\mu} \rightarrow \infty$. Hence, the mean is consistently \sqrt{n} -discriminated. The discrimination results extend to the case with locally dependent observations where we know the local dependence structure and the long run variance is consistently estimable.

Theorem 3.2. *Suppose that there exists $\mathcal{M}'_n \subset \mathcal{M}_n$ such that $|\mathcal{M}'_n| \geq \min\{|\mathcal{M}_n|, 2\}$ and*

$$(3.1) \quad \liminf_{n \rightarrow \infty} \min_{N_n \in \mathcal{M}'_n} |N_n| / n > 0.$$

Suppose further that the set \mathcal{M}'_n satisfies that for each $t \in \mathbf{R}$ and any sequence $P_n \in \mathcal{P}_n$,

$$(3.2) \quad \max_{m \in \mathcal{M}'_n} \left| P_n \left\{ \frac{\sqrt{n_m}(\bar{X}_m - \mathbf{E}[X_{n,i}])}{\sigma_{n,m}} \leq t \right\} - \Phi(t) \right| \rightarrow 0,$$

as $n \rightarrow \infty$, where $\sigma_{n,m}^2 = \text{Var}(\sqrt{n_m}\bar{X}_m)$ and $\bar{X}_m = \frac{1}{n_m} \sum_{i \in N_m} X_{n,i}$.

Then the mean is consistently \sqrt{n} -discriminated at level $\alpha \in (0, 1/2)$ if and only if $|\mathcal{M}_n| \geq 2$ for all but a finite number of n 's.

The theorem says that when the cluster structure consists of large clusters, the mean is \sqrt{n} -consistently discriminated if and only if there are at least two clusters. The necessity of $|\mathcal{M}_n| \geq 2$ already comes from Theorem 3.1. For the sufficiency, we construct a t -test statistic as in Ibragimov and Müller (2010), and show that using the test, we can consistently \sqrt{n} -discriminate the mean.

4. Consistent Estimation of Variance

Recently, Hansen and Lee (2019) showed that it is necessary and sufficient for the weak law of large numbers to hold for the sample average of the clustered observations that

$$\lim_{n \rightarrow \infty} \sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m}{n} \right)^2 = 0.$$

In this section, we show that this condition is necessary and sufficient for consistent estimability of the long run variance. This implies that when there is a large cluster (i.e., which takes up an asymptotically nonnegligible fraction of observations), the long run variance is not consistently estimable. This is a consequence of lack of knowledge of the dependence structure within the large cluster. It means that the usual asymptotic inference based on the asymptotic normal approximation of statistics is generally not applicable in this situation.

4.1. Consistent Estimability

Let us introduce a notion of consistent estimability of a parameter. Let \mathcal{P}_n be the set of joint distributions of observed random variables, say, $\{X_1, \dots, X_n\}$. Given a parameter space $\Theta \subset \mathbf{R}^d$, we define our object of interest to be a map $\theta_n : \mathcal{P}_n \rightarrow \Theta$. Let us introduce the definition of consistent estimability of the parameter θ_n as follows.

Definition 4.1. For any sequence of subsets $\mathcal{P}'_n \subset \mathcal{P}_n$, we say that θ_n is *consistently estimable in* \mathcal{P}'_n , if there exists an estimator $\hat{\theta}$ such that along any sequence $P_n \in \mathcal{P}'_n$,

$$P_n \{ \|\hat{\theta} - \theta_n(P_n)\| > \epsilon \} \rightarrow 0,$$

as $n \rightarrow \infty$, for each $\epsilon > 0$.

One can find a similar definition of consistent estimability in LeCam and Schwartz (1960). They provide necessary and sufficient conditions for a parameter to be consistently estimable when the data are i.i.d. See also Section 1.4 of Ibragimov and Has'minskii (1981) and Section 6.2 of Pfanzagl (1994).

Our setting is somewhat nonstandard, requiring a different technique to prove impossibility of consistent estimation. It is usually assumed that the probability model is indexed by a

certain set, i.e., $\mathcal{P}_n = \{P_{n,h} : h \in \mathcal{H}\}$, where each $P_{n,h}$ is a probability measure indexed by h in some topological space \mathcal{H} that is independent of the sample size n . One can then redefine the parameter $\psi_n(h) = \theta_n(P_{n,h})$, $h \in \mathcal{H}$, i.e., as a map on \mathcal{H} . As long as ψ_n behaves “continuously” on \mathcal{H} , the parameter ψ can be shown to be consistently estimable. (See, e.g., Theorem 4.1 of [Ibragimov and Has’minskii \(1981\)](#) and Theorem 6.2.11 of [Pfanzagl \(1994\)](#).) Then the impossibility of consistent estimation stems from the discontinuity of ψ as a map on \mathcal{H} , which yields “non-identifiability” of the parameter ([Pötscher, 2002](#)).

However, we cannot apply this standard approach in our setting, because there is no natural space \mathcal{H} that indexes \mathcal{P}_n independently of n . The main reason is that we need to deal with a situation potentially with a large cluster with an unknown within-cluster dependence structure. This means that we need to require our probability model to accommodate a wide range of dependence structures for the entire sample. For example, suppose that there is only one large cluster, so that one does not know the dependence structure at all. This means, among other things, that our model needs to include various network dependence structures (such as those studied in [Kojevnikov, Marmor, and Song, 2021](#)) for the joint distribution of the entire random vector $[X_1, \dots, X_n]$ whose dimension grows with the sample size n . One might consider parametrizing the probabilities in terms of the networks governing the dependence structure, but each network depends on the sample size n . To the best of our knowledge, there is no obvious way to topologize such a probability model and to define the continuity of the parameter θ_n on the probabilities, independently of sample size n .

Our approach relies on the following simple lemma that uses contiguity of probabilities at a primitive level. For any two sequences of probabilities P_n and P'_n , we say that P_n is **contiguous with respect to** P'_n if $P'_n(A_n) \rightarrow 0$ implies $P_n(A_n) \rightarrow 0$ for any sequence of Borel sets A_n , and write $P_n \triangleleft P'_n$. When $P_n \triangleleft P'_n$ and $P'_n \triangleleft P_n$, we say that P_n and P'_n are **mutually contiguous**, and write $P_n \triangleleft \triangleright P'_n$. Contiguity between probabilities was introduced by [LeCam \(1960\)](#) and is widely used, especially for deriving the limiting distribution of a test statistic under local alternatives. By tracing out the limiting distribution along a range of local alternatives, one obtains a limiting experiment which one can use to compute the asymptotic risk lower bound in statistical decision theory. (See, e.g., Chapter 6 of [van der Vaart \(1998\)](#).)

The following lemma summarizes our scheme of proving the impossibility of consistent estimability of σ_{LR}^2 .

Lemma 4.1. *Suppose that there exists $P_{n,0} \in \mathcal{P}_n$ such that $P_{n,1} \triangleleft P_{n,0}$ for every sequence $P_{n,1} \in \mathcal{P}_n$. Then θ_n is consistently estimable in \mathcal{P}_n if and only if $\lim_{n \rightarrow \infty} \|\theta_n(P_{n,1}) - \theta_n(P_{n,0})\| = 0$ for any sequence $P_{n,1} \in \mathcal{P}_n$.*

Condition (i) is quite weak and satisfied in many applications. For this condition, it suffices to choose a sequence of probabilities $P_{n,0}$ such that it is easy to build an estimator for θ_n which is

consistent along $P_{n,0}$. When θ_n is σ_{LR}^2 , this condition is immediately satisfied, if the model does not exclude the case of independent observations, because in this case, one can simply take $P_{n,0}$ to be the sequence under which observations are independent, and take the estimator to be the sample variance of the observations. Later we use Lemma 4.1 to prove the impossibility of consistent estimation of the long run variance, by selecting two Gaussian probabilities, $P_{n,0}$ and $P_{n,1}$, such that (i) and (ii) in Lemma 4.1 are satisfied, but the values of the long run variance stay apart as $n \rightarrow \infty$. (See the discussion below Theorem 4.1.)

The notion of consistent estimability in Definition 4.1 coincides with consistent estimability *uniform in P* , i.e., the existence of an estimator $\hat{\theta}$ such that for each $\epsilon > 0$, as $n \rightarrow \infty$,

$$\sup_{P_n \in \mathcal{P}_n} P_n \{ \|\hat{\theta} - \theta_n(P_n)\| > \epsilon \} \rightarrow 0.$$

(See Ibragimov and Has'minskii (1981), p.31. See also Pötscher (2002) for discussion on asymptotics uniform in P .) When $\mathcal{P}_n = \{P_{n,h} : h \in \mathcal{H}\}$ for some index set \mathcal{H} which does not depend on the sample size n , uniform consistent estimability is stronger than pointwise consistent estimability which assumes the existence of an estimator $\hat{\theta}$ such that for each $h \in \mathcal{H}$, and for each $\epsilon > 0$,

$$P_{n,h} \{ \|\hat{\theta} - \theta_n(P_{n,h})\| > \epsilon \} \rightarrow 0,$$

as $n \rightarrow \infty$. However, as explained above, in our setting, there is no space \mathcal{H} that indexes \mathcal{P}_n and is independent of n . Hence there is no natural notion of pointwise consistency in P in our set-up.

4.2. Consistent Estimation of Variance in Gaussian Experiments

4.2.1. A Necessary and Sufficient Condition for Consistent Estimation of Variance. In our context, a major challenge is to show the contiguity condition (ii) of Lemma 4.1. A standard argument proving contiguity utilizes the local asymptotic normality or local asymptotic mixed normality results for a log-likelihood process. However, these latter results often use an i.i.d. or time-series set-up where the researcher knows the dependence structure, and hence are not useful for our purpose here. For this reason, we focus on Gaussian experiments, where we can explicitly compute the log-likelihood process in finite samples and investigate its asymptotic behavior as the dependence structure varies. As long as the probability model in practice does not exclude Gaussian distributions, our impossibility results have direct implications for a more general setting.

Theorem 4.1. Suppose that $\mathcal{P}_{n,\mathcal{N}} \subset \mathcal{P}_n$, and there exists $c > 0$ such that $\sigma_{LR}^2 \leq c$ for all $P_n \in \mathcal{P}_n$ and all $n \geq 1$. Then, the long run variance σ_{LR}^2 is consistently estimable in \mathcal{P}_n , if and only if

$$(4.1) \quad \lim_{n \rightarrow \infty} \sum_{m=1}^{M_n} \frac{n_m^2}{n^2} \rightarrow 0,$$

as $n \rightarrow \infty$.

The sufficiency part of the theorem is straightforward. To see this, suppose for simplicity that there is no singleton cluster in the data. If (4.1) is satisfied, it means that the number of clusters M_n grows to infinity as $n \rightarrow \infty$. Then, we consider the following estimator.

$$\hat{\sigma}_{LR}^2 \equiv \frac{1}{n} \sum_{m=1}^{M_n} \sum_{i,j \in N_{n,m}} (X_{n,i} - \bar{X}_n)(X_{n,j} - \bar{X}_n).$$

The nontrivial part of the theorem is to show that the condition (4.1) is *necessary* for the consistent estimability of σ_{LR}^2 in $\mathcal{P}_{n,\mathcal{N}}$. Suppose that the condition of (4.1) fails, which implies that one has at least one nonnegligible cluster. Then, we show that σ_{LR}^2 is not consistently estimable. For this, we employ Lemma 4.1 after computing the log-likelihood process under cluster dependence. More specifically, suppose first that the observations consist of only a single large cluster. Then, we note that the model \mathcal{P}_n , due to the lack of knowledge on the dependence structure, does not exclude what we call the *local-to-independence common shock models*: $\Phi(0, \Sigma_n)$, with

$$\Sigma_n = I_n + \delta_n \mathbf{1}_n \mathbf{1}_n^\top - \delta_n I_n,$$

where $\mathbf{1}_n$ denotes the n -dimensional column vector of ones. When we take $\delta_n = \bar{\delta}/n$ for some $\bar{\delta} \neq 0$, this model represents the following form of the cross-sectional dependence structure of $X_{n,i}$'s generated as follows:

$$X_{n,i} = \varepsilon_i + \frac{\eta}{\sqrt{n}},$$

where ε_i 's are i.i.d. standard normal random variables, and η is a ‘‘common shock’’ which creates the global dependence among $X_{n,i}$'s, and η and $\{\varepsilon_i\}_{i=1}^n$ are independent. With this choice of Σ_n , we show that $\Phi(0, I_n) \triangleleft \Phi(0, \Sigma_n)$, whereas

$$\sigma_{LR}^2(\Phi(0, I_n)) - \sigma_{LR}^2(\Phi(0, \Sigma_n)) \rightarrow \delta \neq 0.$$

Hence, by Lemma 4.1, σ_{LR}^2 cannot be consistently estimated in any probability model that includes $\Phi(0, \Sigma_n)$, $\Sigma_n \in \mathcal{S}_n^+$. It is not hard to extend the same arguments to a setting where there are potentially multiple large clusters.

Theorem 4.1 then implies that if a nonnegligible portion of the sample belongs to non-singleton clusters, the long run variance σ_{LR}^2 is consistently estimable in $\mathcal{P}_{n,\mathcal{N}}$ if and only if the cluster structure consists of negligible clusters. We formalize this in the following corollary.

Corollary 4.1. *Suppose that the conditions of Theorem 4.1 hold, and $\liminf_{n \rightarrow \infty} n^*/n > 0$, where $n^* = \sum_{m=1: n_m \geq 2}^{M_n} n_m$. Then, the long run variance σ_{LR}^2 is consistently estimable in \mathcal{P}_n if and only if the cluster structure \mathcal{M}_n consists of negligible clusters, i.e.,*

$$\lim_{n \rightarrow \infty} \max_{1 \leq m \leq M_n} \frac{n_m}{n} = 0.$$

The condition $\liminf_{n \rightarrow \infty} n^*/n > 0$ requires that the fraction of random variables $X_{n,i}$ that do not belong to a singleton cluster is asymptotically nonnegligible. In this case, if the probability model in practice includes the Gaussian model $\mathcal{P}_{n,\mathcal{N}}$ as a subclass and there is at least one nonnegligible cluster, it is not possible to consistently estimate the long run variance. Certainly, this impossibility result carries over to a model where the long run variance σ_{LR}^2 is allowed to increase with the sample size n .

Hence, by combining Theorem 3.2 with Theorem 4.1, we find that when we have several large clusters, the long run variance is not consistently estimable because the sample contains large clusters, but the mean can still be consistently \sqrt{n} -discriminated.

4.2.2. Implications for Network Dependent Observations. One might wonder whether the result extends to the case where the observations exhibit a dependence structure other than cluster dependence. Below we give a partial answer for the case of a dependency graph. Dependency graphs were introduced by Stein (1986), and have been studied and used in statistics and econometrics. (See, e.g., Aronow and Samii (2017), Song (2018), Leung (2020) and Canen, Schwartz, and Song (2020) and references therein.)

A **graph** (or network) is a pair $G_n = (N_n, E_n)$, where $N_n = \{1, \dots, n\}$ denotes the set of vertices and E_n the set of edges, where we denote $N(i) = \{j : ij \in E_n\}$ to mean the neighborhood of vertex i . (Here, we consider only simple, undirected graphs, i.e., $ii \notin E_n$, for all $i \in N_n$, and $ij \in E_n$ if and only if $ji \in E_n$.) We define

$$d_{mx} = \max_{i \in N_n} |N(i)|, \text{ and } d_{av} = \frac{1}{n} \sum_{i \in N_n} |N(i)|,$$

where d_{mx} is called the **maximum degree**, and d_{av} the **average degree** of the graph G_n . The maximum and average degrees are often used to capture the denseness of the graph. A subset of vertices in graph G_n is called a **clique** if any two distinct vertices are adjacent in G_n , and the number of vertices in the clique is called the size of the clique. The maximum clique size refers to the size of the clique that is largest in the graph G_n .

Recall that a graph $G_n = (N_n, E_n)$ on $N_n = \{1, \dots, n\}$ is called a **dependency graph** for $X_n = (X_{n,i})_{i \in N_n}$, if for any subset $A \subset N_n$, $(X_{n,i})_{i \in A}$ and $(X_{n,i})_{i \in N_n \setminus \overline{N}_n(A)}$ is independent, where $\overline{N}_n(A) = \{j : ij \in E_n, \text{ for some } i \in A\} \cup \{i\}$. It is important to note that while the dependency graph imposes independence between $X_{n,i}$ and $X_{n,j}$, when they are not adjacent in the graph, it says nothing about dependence between them when they are adjacent. Therefore, as the dependency graph becomes denser, this reflects our limited knowledge on the dependence structure, similarly to large clusters in the cluster dependence case.

Corollary 4.2. *Suppose that for each $n \geq 1$, there exists a graph $G_n = (N_n, E_n)$ which has maximum degree d_{mx} , average degree d_{av} , and maximum clique size n_C , and each distribution $P_n \in \mathcal{P}_n$ of X_n has G_n as a dependency graph. Suppose further that there exists $c > 0$ such that $\sigma_{LR}^2 \leq c$ for all $P_n \in \mathcal{P}_n$ and all $n \geq 1$. Then, the following holds.*

- (i) *If $\limsup_{n \rightarrow \infty} n_C/n > 0$, the long run variance σ_{LR}^2 is not consistently estimable.*
- (ii) *If $\lim_{n \rightarrow \infty} d_{mx}^2 d_{av}/n = 0$, the long run variance σ_{LR}^2 is consistently estimable.*

The impossibility result in (i) has an important implication in many models with network dependent observations. As in the case of a dependency graph, many models of network dependence do not specify the strength of dependence between observations that are adjacent in the network (e.g., [Kojevnikov, Marmer, and Song, 2021](#)). Weak dependence is usually imposed between observations that are far from each other in terms of the shortest path in the network. Hence, when there is a large clique in the network which constitutes a nonnegligible fraction of the entire sample in the limit as $n \rightarrow \infty$, Corollary 4.2(i) implies that the long run variance of the network dependent observations is not consistently estimable.

It is interesting to note that one cannot characterize a necessary and sufficient condition for the network solely in terms of its maximum degree. For example, if X_n is a multivariate normal random vector such that each component has a bounded variance and has a star graph as a dependency graph, the long run variance is consistently estimable. To see this, let $X_n = [X_{n,1}, \dots, X_{n,n}]^\top$ be a centered multivariate random vector which has a graph G_n as a dependency graph. Let the graph G_n be a star graph with the unit 1 being its center.² In the context of multivariate normality, we can write

$$X_{n,1} = \sum_{i \in N_n \setminus \{1\}} \theta_i X_{n,i} + \varepsilon,$$

²The star graph as a dependency graph is different from an additive common shock model such as $X_{n,i} = C_n + \varepsilon_i$, where C_n is a common shock, and ε_i 's are cross-sectionally independent idiosyncratic shock. In this case, the dependency graph is a *complete graph*, because every pair of random variables is correlated through the common shock. Hence, the center in the star graph as a dependency graph cannot be a source like a common shock. It is more plausible to imagine the center to be an aggregated outcome of independent sources. In this case, by simply eliminating the star, one obtains independent random variables.

where the leading sum is the best linear projection, so that ε is independent of $X_{n,i}$'s, $i \in N_n \setminus \{1\}$, which are independent from each other (due to the dependency graph being a star graph). Since the variance of $X_{n,1}$ is bounded, we should have $\sum_{i \in N_n \setminus \{1\}} \theta_i^2 < C$, for all $n \geq 1$, for some $C > 0$. Note that we can identify

$$\theta_i = \text{Cov}(X_{n,i}, X_{n,1}) / \text{Var}(X_{n,i}).$$

Now, we can write

$$\sigma_{LR}^2 = \frac{1}{n} \sum_{i=1}^n \mathbf{E}[X_{n,i}^2] + \frac{1}{n} \sum_{i=1}^n \sum_{j=1, i \neq j}^n \mathbf{E}[X_{n,i} X_{n,j}].$$

The second term is written as

$$\frac{2}{n} \sum_{i=1, i \neq 1}^n \mathbf{E}[X_{n,i} X_{n,1}] = 2\mathbf{E} \left[\left(\frac{1}{n} \sum_{i=1, i \neq 1}^n X_{n,i} \right) X_{n,1} \right] = o(1),$$

as $n \rightarrow \infty$, because the normalized sum in the parenthesis converges to zero in moments. Hence, we can simply take

$$\hat{\sigma}_{LR}^2 = \frac{1}{n} \sum_{i=1}^n X_{n,i}^2$$

to be an estimator of the long run variance. It is not hard to see that $\hat{\sigma}_{LR}^2$ is consistent for σ_{LR}^2 . This example shows that one cannot express the condition for the consistent estimability solely in terms of the maximum degree of the dependency graph.

5. Implications

5.1. A Linear Regression Model with Cluster-Dependent Errors

Let us consider the following regression model with cluster-dependent errors (see, e.g., [Cameron, Gelbach, and Miller \(2008\)](#), [Djogbenou, MacKinnon, and Nielsen \(2019\)](#) and [Hansen and Lee \(2019\)](#) and references therein):

$$y = X\beta + u,$$

where $y = [y'_1, \dots, y'_{M_n}]'$, $X = [X'_1, \dots, X'_{M_n}]'$ and $u = [u'_1, \dots, u'_{M_n}]'$, with $\mathbf{E}[u_m | X] = 0$ for each $m = 1, \dots, M_n$, and each cluster m has n_m observations (so that y_m and u_m are n_m dimensional column vectors and X_m is an $n_m \times k$ matrix.) We assume that u_1, \dots, u_{M_n} are independent, but for each m , the dependence structure of u_m is not known. The OLS estimator of β is given by

$$\hat{\beta} = (X'X)^{-1}X'y.$$

The sandwich form of the variance matrix of $\hat{\beta}$ is given by

$$V = (X'X)^{-1} \left(\sum_{m=1}^{M_n} X'_m \mathbf{E}[u_m u'_m | X] X_m \right) (X'X)^{-1}.$$

For asymptotic normal inference, one needs to obtain a consistent estimator of V . As part of their results, [Djogbenou, MacKinnon, and Nielsen \(2019\)](#) considered the following estimator:

$$\hat{V} = d(X'X)^{-1} \left(\sum_{m=1}^{M_n} X'_m \hat{u}_m \hat{u}'_m X_m \right) (X'X)^{-1},$$

where $\hat{u}_m = y_m - X'_m \hat{\beta}$, and $d \rightarrow 1$ whenever $M_n \rightarrow \infty$, as $n \rightarrow \infty$, and established the consistency of this estimator under a set of conditions. They showed that their conditions are not compatible with a setting in which one of the clusters is large, i.e., its size is proportional to the entire sample.

In this setting, one might wonder whether there is any estimator that consistently estimates V . Our impossibility result implies that as long as we do not exclude the possibility that the error term follows a normal distribution, it is not possible to consistently estimate V when there is at least one large cluster, even if the researcher knows a constant $c > 0$ such that $\|V\| \leq c$.

5.2. Difference-in-Differences with Spillovers

Let us explore the implications of the impossibility results in the context of a difference-in-differences approach to causal inference. (See Section 6.5 of [Imbens and Wooldridge \(2009\)](#) for an overview of this method. See also [Roth, Sant'Anna, Bilinski, and Poe \(2022\)](#) for an overview including recent advances in the literature.) Suppose that there are n individuals who are subject to a treatment and the researcher observes their outcomes before and after the treatment. We let $Y_{i,1}(1)$ and $Y_{i,1}(0)$ denote the potential outcomes at time 1 for the treated state and the control state respectively. The observed outcome at $Y_{i,1}$ at time 1 is defined by

$$Y_{i,1} = D_i Y_{i,1}(1) + (1 - D_i) Y_{i,1}(0),$$

where D_i is the indicator of treatment for i that happens between times 0 and 1. Our parameter of interest is on the average treatment effect on the treated:

$$\text{ATT} = \mathbf{E}[Y_{i,1}(1) - Y_{i,1}(0) | D_i = 1].$$

Suppose that we have observations $\{(Y_{i,1}, Y_{i,0}, D_i)\}_{i=1}^n$, where $Y_{i,0}$ is the outcome for person i in time 0. Furthermore, we assume that the researcher knows the probability $p = P\{D_i = 1\}$. (The impossibility result we mention below carries over to the case where p is not known.)

Let us introduce the standard parallel trend assumption used in the literature:

$$\mathbf{E}[Y_{i,1}(1) - Y_{i,1}(0) \mid D_i = 1] = \mathbf{E}[Y_{i,1}(1) - Y_{i,1}(0) \mid D_i = 0].$$

Under this assumption, we can identify

$$\text{ATT} = \mathbf{E}[\Delta Y_i \mid D_i = 1] - \mathbf{E}[\Delta Y_i \mid D_i = 0],$$

where $\Delta Y_i = Y_{i,1} - Y_{i,0}$. Below we consider two situations where dependence along the observed outcomes can arise: treatment spillover and spillover of treatment effects. We explore implications of our impossibility results in these situations.

5.2.1. Treatment Spillover. Suppose that there is a spillover of the treatments so that D_i 's are correlated across i , along some network among people. For example, one can think of a situation in a social program where two people i and j are neighbors and participating in the program by i can induce the participation by j . Suppose that the researcher does not have information on the neighborhoods among the people. Then, this creates dependence among $Y_{i,1}$'s along a dependence structure that is unknown to the researcher. Then, our impossibility result shows that ATT cannot be consistently discriminated.

In practice, the treatment assignment is often done at the cluster level, where the potential outcomes $Y_{i,1}(1)$ and $Y_{i,1}(0)$ may exhibit arbitrary dependence within each cluster. (See Section 5 of [Roth, Sant'Anna, Bilinski, and Poe \(2022\)](#) for examples and references studying such a setting.) Suppose that we have at least two large clusters such that $(Y_{i,1}(1), Y_{i,0}(0), D_i)$ are independent across the clusters but arbitrarily correlated within each cluster. Then, our impossibility result implies that as long as each within-cluster sum of $(Y_{i,1}(1), Y_{i,0}(0), D_i)$ satisfies the central limit theorem, the ATT is consistently \sqrt{n} -discriminated. The sample analogue of the ATT is given by

$$\widehat{\text{ATT}} = \frac{1}{n} \sum_{i \in N} \left(\frac{\Delta Y_i D_i}{p} - \frac{\Delta Y_i (1 - D_i)}{1 - p} \right).$$

Under mild conditions that do not require the knowledge of the within-cluster dependence structure, one can show that this estimator is consistent. However, our impossibility result implies that it is not possible to consistently estimate the variance of $\sqrt{n}(\widehat{\text{ATT}} - \text{ATT})$.

5.2.2. Spillover of Treatment Effects. Suppose that the treatments D_i themselves do not exhibit any spillover, but the cross-sectional dependence of $(Y_{i,1}(1), Y_{i,0}(0))$ arises due to the spillover of the treatment effects, for example, the treatment of a person i influences the outcome of the person j in the next period. Such a setting has been studied in the recent literature (see [Aronow and Samii \(2017\)](#), [Leung \(2020\)](#), [He and Song \(2022\)](#) and references therein.)

Suppose that the spillover of the treatment effects arises along some network among people, and yet the researcher does not have any information on the network. Then, our impossibility result implies that we cannot consistently discriminate ATT in such a situation. However, the researcher may observe a group structure where the spillover does not arise between groups, so that $(Y_{i,1}(1), Y_{i,0}(0), D_i)$ are independent across groups. If each within-group sum of $(Y_{i,1}(1), Y_{i,0}(0), D_i)$ satisfies the central limit theorem, our result shows that the ATT can be consistently \sqrt{n} -discriminated. However, when there is at least one large group, there does not exist a consistent estimator of the variance of $\sqrt{n}(\widehat{\text{ATT}} - \text{ATT})$.

6. Conclusion

In this paper, we show two impossibility results on the estimation of the mean when the dependence structure is not known. The first result is the impossibility of consistent estimation of the long run variance. The second result is the impossibility of the consistent \sqrt{n} -discrimination of the mean. We make an attempt to accommodate partial knowledge of the dependence structure through cluster dependence, and has obtained some necessary and sufficient conditions for the cluster structure for the impossibility results.

While cluster dependence is a popularly used specification of the cross-sectional dependence structure, it is not general enough to accommodate other forms of a dependence structure such as dependency graphs, Markov graphs, and network dependence. It would be interesting to investigate the implications of partial knowledge of a dependence structure for a more general setting. We leave this for future research.

7. Appendix: Mathematical Proofs

7.1. Preliminary Results

For the proof of the main results, we first prove auxiliary lemmas. As a first step, we provide an explicit form of a log-likelihood process in Gaussian experiments which is Lemma 7.2. The following lemma is an auxiliary result for the lemma.

Lemma 7.1. *Let $\Sigma_0 = USU^\top$ be the spectral decomposition of $\Sigma_0 \in \mathcal{S}^+$ and let $\Sigma_1 \in \mathcal{S}^+$ be such that*

$$\Sigma_1 = \Sigma_0 + UAU^\top,$$

for some positive semidefinite matrix A . Let BAB^\top be the spectral decomposition of $S^{-1/2}AS^{-1/2}$. Suppose that $|\lambda_i| < 1$ for all $i = 1, \dots, n$, where λ_i denote the i -th diagonal entry of Λ .

Then the following results hold.

(i)

$$(7.1) \quad \log(|\Sigma_1|^{-1/2}) - \log(|\Sigma_0|^{-1/2}) = -\frac{1}{2} \sum_{i=1}^n \log(1 + \lambda_i).$$

(ii) For any vectors a and b ,

$$(7.2) \quad a^\top (\Sigma_1^{-1} - \Sigma_0^{-1}) b = - \sum_{i=1}^n \frac{\lambda_i}{1 + \lambda_i} \tilde{a}_i \tilde{b}_i,$$

where \tilde{a}_i and \tilde{b}_i are the i -th entries of \tilde{a} and \tilde{b} , with

$$(7.3) \quad \tilde{a} = B^\top S^{-1/2} U^\top a \quad \text{and} \quad \tilde{b} = B^\top S^{-1/2} U^\top b.$$

Proof: Let $Q = US^{1/2}$ such that $\Sigma_0 = QBB^\top Q^\top$ and $\Sigma_{n,1} = QB(I + \Lambda)B^\top Q^\top$. Thus,

$$|\Sigma_1| = |\Sigma_0| \cdot |I + \Lambda| = |\Sigma_0| \cdot \prod_{i=1}^n (1 + \lambda_i),$$

and

$$\Sigma_0^{-1} - \Sigma_1^{-1} = (B^\top Q^\top)^{-1} (I - (I + \Lambda)^{-1}) (QB)^{-1} = US^{-1/2} B \tilde{\Lambda} B^\top S^{-1/2} U^\top,$$

where

$$\tilde{\Lambda} = \text{diag} \left(\frac{\lambda_1}{1 + \lambda_1}, \dots, \frac{\lambda_n}{1 + \lambda_n} \right).$$

■

The following lemma gives a general log-likelihood process for Gaussian measures. Recall that $\Phi(\mu, \Sigma)$ denotes the multivariate normal distribution with mean vector μ and covariance matrix Σ .

Lemma 7.2. Let $\Sigma_0, \Sigma_1, \Lambda, B, S$ and U be the matrices in Lemma 7.1, and let X be an n -dimensional random vector that follows either $\Phi(\mu_0, \Sigma_0)$ or $\Phi(\mu_1, \Sigma_1)$.

Then

$$\log \frac{d\Phi(\mu_1, \Sigma_1)}{d\Phi(\mu_0, \Sigma_0)}(X) = - \sum_{i=1}^n \log q_i + \frac{1}{2} \sum_{i=1}^n \frac{1}{q_i^2} (Z_i(q_i + 1) - \tilde{\mu}_i)(Z_i(q_i - 1) + \tilde{\mu}_i),$$

where $q_i = \sqrt{1 + \lambda_i}$, λ_i is the i -th diagonal entry of Λ , Z_i is the i -th entry of Z and $\tilde{\mu}_i$ is the i -th entry of $\tilde{\mu}$ with

$$(7.4) \quad Z = B^\top S^{-1/2} U^\top (X - \mu_0) \quad \text{and} \quad \tilde{\mu} = B^\top S^{-1/2} U^\top (\mu_1 - \mu_0).$$

Proof: We write

$$(7.5) \quad \log \frac{d\Phi(\mu_1, \Sigma_1)}{d\Phi(\mu_0, \Sigma_0)}(X) = \log(|\Sigma_1|^{-1/2}) - \log(|\Sigma_0|^{-1/2}) \\ - \frac{1}{2} \left((X - \mu_1)^\top \Sigma_1^{-1} (X - \mu_1) - (X - \mu_0)^\top \Sigma_0^{-1} (X - \mu_0) \right).$$

We apply Lemma 7.1(i) to the first term on the right hand side. As for the last term we let $\mu_\Delta = \mu_1 - \mu_0$, and $X_* = X - \mu_0$. Note that

$$\begin{aligned} \mu_\Delta^\top \Sigma_0^{-1} \mu_\Delta &= \mu_\Delta^\top U S^{-1} U \mu_\Delta = \mu_\Delta^\top U S^{-1/2} S^{-1/2} U^\top \mu_\Delta \\ &= \mu_\Delta^\top U S^{-1/2} B B^\top S^{-1/2} U^\top \mu_\Delta = \tilde{\mu}^\top \tilde{\mu}. \end{aligned}$$

Similarly, $\mu_\Delta^\top \Sigma_0^{-1} X_* = \tilde{\mu}^\top Z$. We rewrite the last term in (7.5) as

$$\begin{aligned} & - \frac{1}{2} X_*^\top (\Sigma_1^{-1} - \Sigma_0^{-1}) X_* - \frac{1}{2} \left(\mu_\Delta^\top (\Sigma_1^{-1} - \Sigma_0^{-1}) \mu_\Delta - 2 \mu_\Delta^\top (\Sigma_1^{-1} - \Sigma_0^{-1}) X_* \right) \\ & \quad - \frac{1}{2} \left(\mu_\Delta^\top \Sigma_0^{-1} \mu_\Delta - 2 \mu_\Delta^\top \Sigma_0^{-1} X_* \right) \\ & = \frac{1}{2} \sum_{i=1}^n \frac{\lambda_i}{1 + \lambda_i} Z_i^2 + \frac{1}{2} \sum_{i=1}^n \frac{\lambda_i}{1 + \lambda_i} (\tilde{\mu}_i^2 - 2 \tilde{\mu}_i Z_i) - \frac{1}{2} \sum_{i=1}^n (\tilde{\mu}_i^2 - 2 \tilde{\mu}_i Z_i) \\ & = \frac{1}{2} \sum_{i=1}^n \left(\left(\frac{\lambda_i}{1 + \lambda_i} - 1 \right) (Z_i - \tilde{\mu}_i)^2 + Z_i^2 \right), \end{aligned}$$

(by applying Lemma 7.1(ii)). By rearranging terms, we rewrite the last sum as

$$(7.6) \quad \frac{1}{2} \sum_{i=1}^n \frac{1}{q_i^2} (Z_i(q_i + 1) - \tilde{\mu}_i)(Z_i(q_i - 1) + \tilde{\mu}_i).$$

Combining this with an earlier result, we obtain the desired result. ■

Lemma 7.3. *Let*

$$(7.7) \quad \Sigma = I + \delta \mathbf{1} \mathbf{1}^\top - \delta I,$$

where δ is such that $n\delta \in (-1, 1)$, and $\mathbf{1}$ denotes the n -dimensional vector of ones.

Then, for any $\bar{\mu} \in \mathbf{R}$,

$$\begin{aligned} \log \frac{d\Phi(\bar{\mu} \mathbf{1}, \Sigma)}{d\Phi(0, I)}(X) &= -\log \sqrt{1 + (n-1)\delta} - (n-1) \log \sqrt{1 - \delta} \\ & \quad + \frac{(n-1)\delta Z_1^2}{2(1 + (n-1)\delta)} - \frac{\delta}{2(1 - \delta)} \sum_{k=2}^n Z_k^2 \\ & \quad + \frac{\sqrt{n\bar{\mu}}}{(1 + (n-1)\delta)} Z_1 - \frac{n\bar{\mu}^2}{2(1 + (n-1)\delta)}, \end{aligned}$$

where Z_k is the k -th entry of $Z \equiv B^\top X$, and $B = [b_1, \dots, b_n]$ is an $n \times n$ orthogonal matrix such that $b_1 = n^{-1/2} \mathbf{1}$, $b_k^\top \mathbf{1} = 0$ for all $k = 2, \dots, n$, $b_k^\top b_\ell = 0$ for all $k \neq \ell = 2, \dots, n$.

Proof: We apply Lemma 7.2 with $S = U = I$, and $A = \delta \mathbf{1}\mathbf{1}^\top - \delta I$. Note that the spectral decomposition of A is given as $B\Lambda B^\top$, where Λ is the diagonal matrix with the diagonal elements $\lambda_1, \dots, \lambda_n$ given as $\lambda_1 = (n-1)\delta$, $\lambda_2 = \dots = \lambda_n = -\delta$, and the orthogonal matrix B as given the lemma. The desired result follows from Lemma 7.2. ■

Lemma 7.3 yields the following result for the case with cluster dependence. From here on, we make the dimension of the matrices and vectors explicit. Let I_{n_m} be the n_m -dimensional identity matrix and $\mathbf{1}_{n_m}$ denote the n_m -dimensional column vector of ones.

Corollary 7.1. Let Σ_n be the block diagonal matrix whose m -th block, $m = 1, \dots, M_n$, is given by

$$(7.8) \quad \Sigma_{n,m} = I_{n_m} + \delta_{n,m} \mathbf{1}_{n_m} \mathbf{1}_{n_m}^\top - \delta_{n,m} I_{n_m},$$

for some $\delta_{n,m} \in \mathbf{R}$ such that $n_m \delta_{n,m} \in (-1, 1)$.

Then, for any $\bar{\mu}_n \in \mathbf{R}$,

$$\begin{aligned} \log \frac{d\Phi(\bar{\mu}_n \mathbf{1}_n, \Sigma_n)}{d\Phi(0, I_n)}(X_n) &= - \sum_{m=1}^{M_n} \log \sqrt{1 + (n_m - 1)\delta_{n,m}} - \sum_{m=1}^{M_n} (n_m - 1) \log \sqrt{1 - \delta_{n,m}} \\ &\quad + \sum_{m=1}^{M_n} \frac{(n_m - 1)\delta_{n,m} Z_{m,i_m}^2}{2(1 + (n_m - 1)\delta_{n,m})} - \sum_{m=1}^{M_n} \frac{\delta_{n,m}}{2(1 - \delta_{n,m})} \sum_{i \in N_{n,m} \setminus \{i_m\}} Z_{m,i}^2 \\ &\quad + \sum_{m=1}^{M_n} \frac{n\bar{\mu}_n / \sqrt{n_m}}{(1 + (n_m - 1)\delta_{n,m})} Z_{n,i_m} - \frac{1}{2} \sum_{m=1}^{M_n} \frac{n^2 \bar{\mu}_n^2 / \sqrt{n_m}}{1 + (n_m - 1)\delta_{n,m}}, \end{aligned}$$

where $Z_{m,i}$, $i \in N_{n,m}$, are the entries of $Z_m = B_m^\top X_{m,n}$, B_m is an $n_m \times n_m$ orthogonal matrix, i_m denotes the first index in $N_{n,m}$, and $X_{m,n} = [X_{n,i}]_{i \in N_{n,m}}$.

Lemma 7.4. Suppose that $f : \mathbf{R} \rightarrow \mathbf{R}^+$ is a continuously differentiable function such that for some $\delta > 0$,

$$\sup_{x \in [-\delta, \delta]} \left| \frac{d \log f(x)}{dx} \right| |\bar{x}| < 1,$$

where $\bar{x} \in \mathbf{R}$ is such that

$$\sup_{x \in [-\delta, \delta]} f(x) = f(\bar{x}).$$

Then,

$$f(\bar{x}) \leq \left(1 - \sup_{x \in [-\delta, \delta]} \left| \frac{d \log f(x)}{dx} \right| |\bar{x}| \right)^{-1} f(0).$$

Proof: Using the Mean Value Theorem,

$$\begin{aligned} \exp \log f(x) &= \exp \log f(0) + \frac{d \log f(x^*(x))}{dx} (\exp \log f(x^*(x))) x \\ &\leq \exp \log f(0) + \sup_{\tilde{x} \in [-\delta, \delta]} \left| \frac{d \log f(\tilde{x})}{dx} \right| (\exp \log f(\bar{x})) |x|, \end{aligned}$$

where $x^*(x)$ is a point on the line segment between 0 and x . Evaluating the inequality at $x = \bar{x}$ gives us the desired result. ■

To accommodate the case with singleton-clusters (so that the case with independent observations is included as a special case), let us introduce the notation:

$$(7.9) \quad n^* = \sum_{m=1: n_m \geq 2}^{M_n} n_m,$$

where the summation is over $m = 1, \dots, M_n$ such that $n_m \geq 2$. (When the summation is over m in the empty set, the sum is taken to be zero.) The number $n - n^*$ represents the number of random variables, $X_{n,i}$, that are known to be mutually independent. Each variable outside this set belongs to a non-singleton cluster.

Lemma 7.5. $\lim_{n \rightarrow \infty} \sum_{m=1}^{M_n} (n_m/n)^2 = 0$, if and only if

- (a) $\lim_{n \rightarrow \infty} n^*/n = 0$, or
- (b) $\lim_{n \rightarrow \infty} \sum_{m=1: n_m \geq 2}^{M_n} (n_m/n^*)^2 = 0$.

Proof: For each $n \geq 1$, we have either

$$\sum_{m=1}^{M_n} \frac{n_m^2}{n^2} = \begin{cases} \frac{1}{n}, & \text{if } n^* = 0, \\ \left(\frac{n^*}{n}\right)^2 \sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*}\right)^2 + \left(\frac{1}{n}\right)^2 (n - n^*), & \text{if } n^* > 0. \end{cases}$$

Since $(1/n)^2(n - n^*) = o(1)$, $\lim_{n \rightarrow \infty} \sum_{m=1}^{M_n} (n_m/n)^2 = 0$ if and only if (a) or (b) holds. ■

Lemma 7.6. Suppose that $n \geq 2$, and Σ_n is a block diagonal matrix along a cluster structure \mathcal{M}_n , where the m -th block, denoted by $\Sigma_{n,m}$ is given by

$$\Sigma_{n,m} = I_{n_m} + \delta_{n,m} \mathbf{1}_{n_m}^\top \mathbf{1}_{n_m} - \delta_{n,m} I_{n_m},$$

with

$$(7.10) \quad \delta_{n,m} = \frac{\bar{\delta}}{n^*},$$

for some $\bar{\delta} \in [-a, a]$, with $0 < a < 1/2$, if $n_m \geq 2$.

Then, the following holds.

- (i) $(d\Phi(0, \Sigma_n)/d\Phi(0, I_n))(X_n)$ is uniformly integrable under $\Phi(0, I_n)$.
(ii) $\log((d\Phi(0, \Sigma_n)/d\Phi(0, I_n))(X_n))$ is uniformly tight under $\Phi(0, I_n)$.

Proof: By Corollary 7.1,

$$\log \frac{d\Phi(0, \Sigma_n)}{d\Phi(0, I_n)}(X_n) = A_n + R_n,$$

where

$$A_n = - \sum_{m=1}^{M_n} \log \sqrt{1 + (n_m - 1)\delta_{n,m}} + \sum_{m=1}^{M_n} \frac{(n_m - 1)\delta_{n,m} Z_{m,i_m}^2}{2(1 + (n_m - 1)\delta_{n,m})}, \text{ and}$$

$$R_n = - \sum_{m=1}^{M_n} (n_m - 1) \log \sqrt{1 - \delta_{n,m}} - \sum_{m=1}^{M_n} \frac{\delta_{n,m}}{2(1 - \delta_{n,m})} \sum_{i \in N_{n,m} \setminus \{i_m\}} Z_{m,i}^2,$$

and i_m denotes the first i in block m .

(i) Let us take small $\epsilon > 0$ such that

$$(7.11) \quad a < \frac{2}{(1 + \epsilon)(4 + \epsilon)}.$$

We write (under $\Phi(0, I_n)$)

$$\mathbf{E} \left[\left(\frac{d\Phi(0, \Sigma_n)}{d\Phi(0, I_n)} \right)^{(1+\epsilon)} \right] = \mathbf{E}[\exp((1 + \epsilon)A_n)] \mathbf{E}[\exp((1 + \epsilon)R_n)],$$

since A_n and R_n are independent. Let $t_m = (n_m - 1)/n^*$, and write

$$\begin{aligned} \mathbf{E}[\exp((1 + \epsilon)A_n)] &= \prod_{m=1}^{M_n} \left(1 + t_m \bar{\delta}\right)^{-\frac{1+\epsilon}{2}} \mathbf{E} \left[\exp \left(\frac{t_m(1 + \epsilon)\bar{\delta}}{2(1 + t_m \bar{\delta})} Z_{m,1}^2 \right) \right] \\ &\leq \prod_{m=1}^{M_n} \left(1 + t_m \bar{\delta}\right)^{-\frac{1+\epsilon}{2}} \left(1 - t_m(1 + \epsilon)\bar{\delta}\right)^{-1/2} \equiv f_n(\bar{\delta}), \text{ say.} \end{aligned}$$

After some algebra, we find that

$$\frac{d \log f_n(\bar{\delta})}{d \bar{\delta}} = \frac{(1 + \epsilon)(2 + \epsilon)\bar{\delta}}{2} \sum_{m=1}^{M_n} \frac{t_m^2}{(1 + t_m \bar{\delta})(1 - t_m \bar{\delta}(1 + \epsilon))} > 0,$$

because $t_m \leq 1$ and $\bar{\delta}(1 + \epsilon) < 1$ by (7.11). This means that $f_n(\bar{\delta})$ is increasing in $\bar{\delta} \in [-a, a]$ and achieves its maximum at $\bar{\delta} = a$. Hence,

$$\begin{aligned} \left| \frac{d \log f_n(\bar{\delta})}{d \bar{\delta}} \right|_a &\leq \left| \frac{d \log f_n(\bar{\delta})}{d \bar{\delta}} \right| \leq \frac{(1 + \epsilon)(2 + \epsilon)\bar{\delta}}{2(1 - (1 + \epsilon)\bar{\delta})} \sum_{m=1}^{M_n} t_m^2 \\ &\leq \frac{(1 + \epsilon)(2 + \epsilon)a}{2(1 - (1 + \epsilon)a)} < 1, \end{aligned}$$

because $\sum_{m=1}^{M_n} t_m^2 \leq 1$ and we chose ϵ such that (7.11) holds. By Lemma 7.4, we have

$$f_n(\bar{\delta}) \leq f_n(a) \leq \left(1 - \frac{(1+\epsilon)(2+\epsilon)a}{2(1-(1+\epsilon)a)}\right)^{-1}.$$

The bound does not depend on n , and hence

$$\sup_{n \geq 1} \mathbf{E}[\exp((1+\epsilon)A_n)] < \infty.$$

Now, we turn to $\mathbf{E}[\exp((1+\epsilon)R_n)]$. We can write

$$(7.12) \quad R_n = -\frac{1}{2}n^* \log\left(1 - \frac{\bar{\delta}}{n^*}\right) - \frac{1}{2} \frac{\bar{\delta}/n^*}{1 - \bar{\delta}/n^*} \sum_{m=1: n_m \neq 1}^{M_n} \sum_{i \in N_{n,m} \setminus \{i_m\}} Z_{m,i}^2.$$

Using this expression, we rewrite

$$\begin{aligned} \mathbf{E}[\exp((1+\epsilon)R_n)] &= \left(1 - \frac{\bar{\delta}}{n^*}\right)^{-\frac{n^*(1+\epsilon)}{2}} \prod_{m=1: n_m \neq 1}^{M_n} \prod_{i \in N_{n,m} \setminus \{i_m\}} \mathbf{E}\left[\exp\left(-\frac{1}{2} \frac{\bar{\delta}(1+\epsilon)/n^*}{1 - \bar{\delta}/n^*} Z_{m,i}^2\right)\right] \\ &= \left(1 - \frac{\bar{\delta}}{n^*}\right)^{-\frac{n^*(1+\epsilon)}{2}} \prod_{m=1: n_m \neq 1}^{M_n} \prod_{i \in N_{n,m} \setminus \{i_m\}} \left(1 + \frac{\bar{\delta}(1+\epsilon)/n^*}{1 - \bar{\delta}/n^*}\right)^{-\frac{1}{2}} \\ &\leq \left(1 - \frac{\bar{\delta}}{n^*}\right)^{-\frac{n^*(1+\epsilon)}{2}} \leq \left(1 - \frac{a}{n^*}\right)^{-\frac{n^*(1+\epsilon)}{2}}. \end{aligned}$$

The last bound is a sequence converging to $\exp(a(1+\epsilon)/2)$ as $n^* \rightarrow \infty$, and hence is a bounded sequence. Thus, we conclude that

$$\sup_{n \geq 1} \mathbf{E}[\exp((1+\epsilon)R_n)] < \infty.$$

This proves that

$$\sup_{n \geq 1} \mathbf{E}\left[\left(\frac{d\Phi(0, \Sigma_n)}{d\Phi(0, I_n)}\right)^{(1+\epsilon)} (X_n)\right] < \infty.$$

Hence, the proof of (i) is complete.

(ii) We rewrite

$$(7.13) \quad A_n = -\frac{1}{2} \sum_{m=1}^{M_n} \left(\log(1 + (n_m - 1)\delta_{n,m}) - \frac{(n_m - 1)\delta_{n,m}}{1 + (n_m - 1)\delta_{n,m}} \right) + \sum_{m=1}^{M_n} \frac{(n_m - 1)\delta_{n,m}(Z_{m,i_m}^2 - 1)}{2(1 + (n_m - 1)\delta_{n,m})}.$$

For any $x \in [0, 1]$, we have

$$\frac{x^2}{8} \leq \frac{1}{2} \frac{x^2}{(1+x)^2} \leq \log(1+x) - \frac{x}{1+x} \leq \frac{1}{2} \frac{x^2}{1+x} \leq \frac{x^2}{2}.$$

Hence,

$$\left| -\frac{1}{2} \sum_{m=1}^{M_n} \left(\log(1 + (n_m - 1)\delta_{n,m}) - \frac{(n_m - 1)\delta_{n,m}}{1 + (n_m - 1)\delta_{n,m}} \right) \right| \leq \frac{\bar{\delta}^2}{2} \sum_{m=1}^{M_n} \frac{(n_m - 1)^2}{2n^{*2}} \leq \frac{\bar{\delta}^2}{2}.$$

It suffices to show the uniform tightness of the second sum in (7.13). Under $\Phi(0, I_n)$, it has mean zero, and

$$\begin{aligned} \text{Var} \left(\sum_{m=1: n_m \geq 2}^{M_n} \frac{\bar{\delta}(n_m - 1)/n^*}{1 + (\bar{\delta}(n_m - 1)/n^*)} (Z_{m, i_m}^2 - 1) \right) &\leq \sum_{m=1}^{M_n} \frac{(\bar{\delta}(n_m - 1)/n^*)^2}{(1 + (\bar{\delta}(n_m - 1)/n^*))^2} \text{Var}(Z_{m, i_m}^2) \\ &\leq 2\bar{\delta}^2 \sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m - 1}{n^*} \right)^2 \leq 2\bar{\delta}^2, \end{aligned}$$

because $\text{Var}(Z_{m, i_m}^2) = 2$. Therefore, A_n is uniformly tight.

As for R_n , we recall (7.12), and can follow similar arguments to show that R_n is uniformly tight as well. ■

7.2. Consistent Discrimination of the Mean

We let

$$(7.14) \quad \Sigma(\delta) = I_n + \delta \mathbf{1}_n \mathbf{1}_n^\top - \delta I_n,$$

and for any $\mu \in \mathbf{R}$, we write $\Phi(\mu \mathbf{1}_n, \Sigma(\delta))$ simply as $\Phi(\mu, \delta)$. Let us recall some basic notions of optimality of tests (Lehmann and Romano, 2005). Given a model \mathcal{P}_n which is partitioned as $\mathcal{P}_{n,0} \cup \mathcal{P}_{n,1}$, a test ϕ_n is said to be a **UMP (uniformly most powerful) test** of $\mathcal{P}_{n,0}$ against $\mathcal{P}_{n,1}$ at level $\alpha \in (0, 1)$, if under any $P_{n,0} \in \mathcal{P}_{n,0}$,

$$\mathbf{E}[\phi_n] \leq \alpha,$$

and for any alternative test ϕ'_n such that $\mathbf{E}[\phi'_n] \leq \alpha$ under any sequence $P_{n,0} \in \mathcal{P}_{n,0}$, we have

$$\mathbf{E}[\phi'_n] - \mathbf{E}[\phi_n] \leq 0,$$

under any $P_{n,1} \in \mathcal{P}_{n,1}$.

A sequence of tests ϕ_n is said to be an **AUMP (asymptotically uniformly most powerful) test** of $\mathcal{P}_{n,0}$ against $\mathcal{P}_{n,1}$ at level $\alpha \in (0, 1)$, if under any sequence $P_{n,0} \in \mathcal{P}_{n,0}$,

$$\limsup_{n \rightarrow \infty} \mathbf{E}[\phi_n] \leq \alpha,$$

and for any alternative test ϕ'_n such that $\limsup_{n \rightarrow \infty} \mathbf{E}[\phi'_n] \leq \alpha$ under any sequence $P_{n,0} \in \mathcal{P}_{n,0}$, we have

$$\limsup_{n \rightarrow \infty} \mathbf{E}[\phi'_n] - \mathbf{E}[\phi_n] \leq 0,$$

under any sequence $P_{n,1} \in \mathcal{P}_{n,1}$.

Lemma 7.7. *Suppose that $A \subset \mathbf{R}$ is an open interval, and $\{\varphi_\alpha\}_{\alpha \in A}$ is a class of tests of $\mathcal{P}_{n,0}$ against $\mathcal{P}_{n,1}$, such that for each $\alpha \in A$, the test φ_α is UMP at level α , and for any $\tilde{\alpha} = \alpha + o(1)$ as $n \rightarrow \infty$,*

$$(7.15) \quad \mathbf{E}[\varphi_\alpha] = \mathbf{E}[\varphi_{\tilde{\alpha}}] + o(1),$$

under any sequence $P_n \in \mathcal{P}_{n,0} \cup \mathcal{P}_{n,1}$. Then φ_α is AUMP at level α .

Proof: Choose any test $\tilde{\varphi}$ such that under any $P_n \in \mathcal{P}_{n,0}$,

$$\mathbf{E}[\tilde{\varphi}] \leq \alpha + \epsilon_n,$$

for some sequence $\epsilon_n \rightarrow 0$, as $n \rightarrow \infty$. Fix one such sequence $P_n \in \mathcal{P}_{n,0}$, together with the sequence ϵ_n , and let $\tilde{\alpha} = \alpha + \epsilon_n$. Now, select large enough n such that $\tilde{\alpha} \in A$, and choose any $P'_n \in \mathcal{P}_{n,1}$. Then since $\varphi_{\tilde{\alpha}}$ is UMP at level $\tilde{\alpha}$, we have

$$\mathbf{E}_{P'_n}[\varphi_{\tilde{\alpha}}] \geq \mathbf{E}_{P'_n}[\tilde{\varphi}]$$

under P'_n . By (7.15), we can see that φ_α is AUMP at level α . ■

Proof of Theorem 3.1: Suppose that $M_n = 1$. Let us consider the following model:

$$\mathcal{P}_n = \{\Phi(\mu, \delta) : (n-1)\delta \in (0, 1), \mu \geq 0\}.$$

Also, define $\mathcal{P}_{n,0} = \{\Phi(0, \delta) : \delta > 0\}$ and let $\mathcal{P}_{n,1} = \mathcal{P}_n \setminus \mathcal{P}_{n,0}$. In light of Lemma 7.7, it suffices to construct a class of tests $\{\varphi_\alpha\}_{\alpha \in (0, 1/2)}$ of $\mathcal{P}_{n,0}$ against $\mathcal{P}_{n,1}$ such that

- (a) it satisfies (7.15) in Lemma 7.7,
- (b) the test φ_α has power bounded by a constant below 1 uniformly over n , and
- (c) each test φ_α is a UMP test of $\mathcal{P}_{n,0}$ against $\mathcal{P}_{n,1}$.

Let us first construct such a test and show that (a)-(c) are satisfied. Define

$$V_n = \frac{Z_{n,1}}{|Z_{n,1}|} = \text{sgn}(Z_{n,1}).$$

For each $\alpha \in (0, 1/2)$, let

$$\phi_\alpha(V_n) = \begin{cases} 1, & \text{if } V_n > C_0 \\ \gamma_0, & \text{if } V_n = C_0, \\ 0, & \text{if } V_n < C_0, \end{cases}$$

for some $C_0 > 0$ and $\gamma_0 \in [0, 1]$. Let $Z = (Z_{n,1} - \mathbf{E}[Z_{n,1}]) / \sqrt{\text{Var}(Z_{n,1})}$. Then the size control requires that under the null hypothesis,

$$(7.16) \quad \mathbf{E}[\phi_\alpha(V_n)] = P\{Z > C_0 | Z|\} + P\{Z = C_0 | Z|\} \gamma_0 = \alpha.$$

Since $\alpha \in (0, 1/2)$, we must have $C_0 = 1$ and $\gamma_0 = 2\alpha$.

Let us first show that this test satisfies the condition (a). For any $\tilde{\alpha}$ such that $\tilde{\alpha} = \alpha + o(1)$, and under any sequence $P_n \in \mathcal{P}_n$,

$$(7.17) \quad \mathbf{E}[\phi_{\tilde{\alpha}}(V_n)] = 2\tilde{\alpha}P_n\{Z_{n,1} \geq 0\} = 2\alpha P_n\{Z_{n,1} \geq 0\} + o(1) = \mathbf{E}[\phi_\alpha(V_n)] + o(1).$$

Hence, the class of tests $\{\phi_\alpha(V_n)\}_{\alpha \in (0, 1/2)}$ satisfies the condition (7.15).

As for the condition (b), note that under any alternative hypothesis in $\mathcal{P}_{n,1}$, we have

$$\mathbf{E}[\varphi_\alpha(V_n)] = P_n\{Z_{n,1} > |Z_{n,1}|\} + 2\alpha P_n\{Z_{n,1} = |Z_{n,1}|\} = 2\alpha P_n\{Z_{n,1} \geq 0\} \leq 2\alpha.$$

Hence, the test does not have power exceeding $2\alpha < 1$.

Finally, we show that the condition (c) is satisfied. Let

$$(7.18) \quad \begin{aligned} \mathcal{L}(X_n; \mu, \delta_n) = & -\log \sqrt{1 + (n-1)\delta_n} - (n-1) \log \sqrt{1 - \delta_n} + \frac{(n-1)\delta_n Z_{n,1}^2}{2(1 + (n-1)\delta_n)} \\ & - \frac{\delta_n}{2(1 - \delta_n)} \sum_{k=2}^n Z_{n,k}^2 + \sqrt{n}\mu Z_{n,1} - \frac{n\mu^2(1 + (n-1)\delta_n)}{2}. \end{aligned}$$

Hence, $\mathcal{L}(X_n; \mu, \delta_n)$ is the same as $\log(d\Phi(\mu_n \mathbf{1}_n, \Sigma_n)/d\Phi(0, I_n))(X_n)$ in Lemma 7.3, except that the coefficient of $Z_{n,1}$ is $\sqrt{n}\mu$ and the last term is different. Define a probability measure $P_n(\mu, \delta_n)$ as follows: for any Borel B ,

$$P_n(\mu, \delta_n)(B) = \int_B \exp(\mathcal{L}(x; \mu, \delta_n)) d\Phi(0, I_n)(x).$$

Similarly as before, we define

$$\tilde{\mathcal{P}}_n = \{P_n(\mu, \delta) : (n-1)\delta \in (0, 1), \mu \geq 0\}, \tilde{\mathcal{P}}_{n,0} = \{P_n(0, \delta) : (n-1)\delta \in (0, 1)\},$$

and let $\tilde{\mathcal{P}}_{n,1} = \tilde{\mathcal{P}}_n \setminus \tilde{\mathcal{P}}_{n,0}$. It is not hard to see that

$$\tilde{\mathcal{P}}_{n,0} = \mathcal{P}_{n,0}, \text{ and } \tilde{\mathcal{P}}_{n,1} = \mathcal{P}_{n,1}.$$

Therefore, a UMP test of $\tilde{\mathcal{P}}_{n,0}$ against $\tilde{\mathcal{P}}_{n,1}$ is also a UMP test of $\mathcal{P}_{n,0}$ against $\mathcal{P}_{n,1}$. It suffices for condition (c) to show that the test $\varphi_\alpha(V_n)$ is a UMP test of $\tilde{\mathcal{P}}_{n,0}$ against $\tilde{\mathcal{P}}_{n,1}$. From (7.18), the sufficient statistics for $\tilde{\mathcal{P}}_n$ in the case of $M_n = 1$ are given by

$$\left(Z_{n,1}, Z_{n,1}^2, \sum_{k=2}^n Z_{n,k}^2 \right),$$

where $Z_{n,k}$'s are as in Lemma 7.3. For any $t \geq 0$,

$$\begin{aligned} P\{V_n = 1, Z_{n,1}^2 \leq t\} &= P\{V_n = 1, -\sqrt{t} \leq Z_{n,1} \leq \sqrt{t}\} = P\{Z_{n,1} > 0, -\sqrt{t} \leq Z_{n,1} \leq \sqrt{t}\} \\ &= P\{0 \leq Z_{n,1} \leq \sqrt{t}\} = \frac{1}{2}P\{-\sqrt{t} \leq Z_{n,1} \leq \sqrt{t}\} = P\{V_n = 1\}P\{Z_{n,1}^2 \leq t\}, \end{aligned}$$

under the null hypothesis. Hence, V_n and $Z_{n,1}^2$ are independent under any probability in $\tilde{\mathcal{P}}_{n,0}$. Furthermore, under any probability in $\tilde{\mathcal{P}}_{n,0}$,

$$\begin{aligned} \mathbf{E}[Z_n Z_n^\top] &= B_n^\top \mathbf{E}[Z_n Z_n^\top] B_n \\ &= B_n^\top \Sigma_n B_n = B_n^\top (I_n + \delta_n \mathbf{1}_n \mathbf{1}_n^\top - \delta_n I_n) B_n = (1 - \delta_n) I_n + B_n^\top \mathbf{1}_n \mathbf{1}_n B_n. \end{aligned}$$

Note that $B_n^\top \mathbf{1}_n \mathbf{1}_n B_n$ is a matrix whose $(1, 1)$ -th entry is $b_1^\top \mathbf{1}_n \mathbf{1}_n^\top b_1 = 1$ and all the other entries are zeros. Hence, $Z_{n,k}$'s are independent across k 's under any probability in $\tilde{\mathcal{P}}_{n,0}$. Therefore, V_n and $(Z_{n,1}^2, \sum_{k=2}^n Z_{n,k}^2)$ are independent under any probability in $\tilde{\mathcal{P}}_{n,0}$. By Theorem 5.1.1 of [Lehmann and Romano \(2005\)](#), the randomized test $\varphi_\alpha(V_n)$ is an α -level UMP test. ■

The following lemma is used for the proof of Theorem 3.2. For $w \in [0, 1]$, define $\xi_1 = wZ_1$ and $\xi_2 = (1-w)Z_2$, where $Z_i \sim N(0, 1)$, independent across $i = 1, 2$. Define

$$T(w) = \frac{(\xi_1 + \xi_2)/\sqrt{2}}{\sqrt{(\xi_1 - \bar{\xi})^2 + (\xi_2 - \bar{\xi})^2}},$$

where $\bar{\xi} = (\xi_1 + \xi_2)/2$. Let us take $t \geq 0$, and define

$$p(w; t) = P\{T(w) \leq t\}.$$

Lemma 7.8. (i) For all $t \geq 1$ and all $w \in [0, 1]$, $p(w; t) \geq p(1/2; t)$.

(ii) For all $0 \leq t < 1$ and all $w \in [0, 1]$, $p(w; t) \leq p(1/2; t)$.

Proof: First, we write

$$T(w) = \frac{\xi_1 + \xi_2}{\sqrt{(\xi_1 - \xi_2)^2}}.$$

For (i) and (ii), since $[\xi_1, \xi_2]$ is symmetrically distributed around the origin, it suffices to show that $p(\cdot; t)$ is increasing on $[1/2, 1]$ for all $t \geq 1$, and $p(\cdot; t)$ is decreasing on $[1/2, 1]$ for all $0 \leq t < 1$. Let A denote the event that $\xi_1 + \xi_2 < 0$. Then, on the event A , for all $w \geq 0$ and all $t \geq 0$, we have $T(w) \leq t$. Hence

$$P(\{T(w) \leq t\} \cap A) = P(A) = 0.5.$$

On the event A^c , $T(w) \leq t$ if and only if

$$w^2 Z_1^2 + (1-w)^2 Z_2^2 + 2w(1-w)Z_1 Z_2 \leq (w^2 Z_1^2 + (1-w)^2 Z_2^2 - 2w(1-w)Z_1 Z_2) t^2$$

if and only if

$$0 \leq (w^2 Z_1^2 + (1-w)^2 Z_2^2)(t^2 - 1) - 2(t^2 + 1)w(1-w)Z_1 Z_2 = f(w; Z_1, Z_2),$$

where

$$f(w; Z_1, Z_2) = (wZ_1 + (1-w)Z_2)^2(t^2 - 1) - 4t^2 w(1-w)Z_1 Z_2.$$

Take $t \geq 1$. Let B be the event $Z_1 Z_2 \leq 0$. Certainly, if $Z_1 Z_2 \leq 0$, $f(w; Z_1, Z_2) \geq 0$ for all $w \in [1/2, 1]$. Hence

$$\begin{aligned} P(\{T(w) \leq t\} \cap A^c) &= P(\{T(w) \leq t\} \cap A^c \cap B) + P(\{T(w) \leq t\} \cap A^c \cap B^c) \\ &= P(A^c \cap B) + P(\{T(w) \leq t\} \cap A^c \cap B^c). \end{aligned}$$

We show that $f(w; Z_1, Z_2)$ is increasing in w on the event A^c . We take the derivative $f'(w; Z_1, Z_2)$ with respect to w :

$$f'(w; Z_1, Z_2) = 2(wZ_1 - (1-w)Z_2)(t^2 - 1)(Z_1 - Z_2) - 4t^2(1-2w)Z_1 Z_2.$$

The function is linear in w . First, we take $w = 1$. Then, on the event B^c ,

$$\begin{aligned} f'(1; Z_1, Z_2) &= 2(t^2 - 1)Z_1(Z_1 - Z_2) + 4t^2 Z_1 Z_2 = 2(t^2 - 1)Z_1^2 + (4t^2 - 2(t^2 - 1))Z_1 Z_2 \\ &= 2(t^2 - 1)Z_1^2 + (2t^2 + 2)Z_1 Z_2 \geq 0. \end{aligned}$$

Second, we take $w = 1/2$. Then,

$$f'(1/2; Z_1, Z_2) = (t^2 - 1)(Z_1 - Z_2)^2.$$

Hence, for all Z_1, Z_2 such that $Z_1 Z_2 > 0$, $f(w; Z_1, Z_2)$ is increasing on $[1/2, 1]$ for all $t \geq 1$. Therefore, whenever $t \geq 1$, $P\{T(w) \leq t\}$ is increasing on $[1/2, 1]$.

Take $0 \leq t < 1$. If $Z_1 Z_2 > 0$, then $f(w; Z_1, Z_2) < 0$. Hence

$$\begin{aligned} P(\{T(w) \leq t\} \cap A) &= P(\{T(w) \leq t\} \cap A^c \cap B) + P(\{T(w) \leq t\} \cap A^c \cap B^c) \\ &= P(\{T(w) \leq t\} \cap A^c \cap B). \end{aligned}$$

On the event B , when $w = 1$,

$$f'(1; Z_1, Z_2) = 2(t^2 - 1)Z_1^2 + (2t^2 + 2)Z_1 Z_2 \leq 0,$$

and when $w = 1/2$,

$$(7.19) \quad f'(1/2; Z_1, Z_2) = (t^2 - 1)(Z_1 - Z_2)^2 \leq 0.$$

Hence, for all Z_1, Z_2 such that $Z_1 Z_2 \leq 0$, $f(w; Z_1, Z_2)$ is decreasing on $[1/2, 1]$ for all $0 \leq t < 1$. Therefore, whenever $0 \leq t < 1$, $P\{T(w) \leq t\}$ is decreasing on $[1/2, 1]$. ■

Proof of Theorem 3.2: Since $\mathcal{P}_{n,\mathcal{N}} \subset \mathcal{P}_n$, when $M_n = 1$ for infinitely many n 's, the mean is not consistently \sqrt{n} -discriminated by Theorem 3.1. Suppose that we have at least two non-negligible clusters, i.e., $M_n \geq 2$ for all but finite n 's. Consider testing the null hypothesis of $\mathbf{E}[X_{n,i}] = 0$ against $\mathbf{E}[X_{n,i}] > 0$. Without loss of generality, we enumerate $\mathcal{M}'_n = \{N_1, \dots, N_{M'_n}\}$, $M'_n \geq 2$. Now we construct a test that consistently \sqrt{n} -discriminates the mean. Define

$$\xi_m = \frac{1}{\sqrt{n_m}} \sum_{i \in N_{n,m}} X_{n,i},$$

and

$$U'_n = \frac{1}{2} \sum_{m=1}^2 \xi_m \quad \text{and} \quad T'_n = \sum_{m=1}^2 \left(\xi_m - \frac{1}{2} \sum_{m=1}^2 \xi_m \right)^2.$$

We take

$$V'_n = \frac{\sqrt{2}U'_n}{\sqrt{T'_n}}.$$

Let $c_{1-\alpha}$ be the $1 - \alpha$ quantile of the t -distribution with degree of freedom 1. Define

$$\varphi_n = 1\{V'_n > \max\{c_{0.75}, c_{1-\alpha}\}\}.$$

Note that $c_{0.75} = 1$.

We first show that this test controls the size of the test under α asymptotically under the null hypothesis. Define an infeasible test statistic

$$V''_n = \frac{\sqrt{2}U''_n}{\sqrt{T''_n}},$$

where

$$U''_n = \frac{1}{2} \sum_{m=1}^2 \frac{\xi_m}{\sigma_{n,m}} \quad \text{and} \quad T''_n = \sum_{m=1}^2 \left(\frac{\xi_m}{\sigma_{n,m}} - \frac{1}{2} \sum_{m=1}^2 \frac{\xi_m}{\sigma_{n,m}} \right)^2.$$

Here $\sigma_{n,m}^2$ is the variance of ξ_m under the probability under the null hypothesis. Then V''_n converges in distribution to the t -distribution with the degree of freedom equal to 1 under the null hypothesis. By Lemma 7.8, if $0 \leq \alpha < 0.25$ so that $c_{1-\alpha} > 1$,

$$(7.20) \quad \mathbf{E}[\varphi_n] = P\{V'_n > c_{1-\alpha}\} \leq P\{V''_n > c_{1-\alpha}\} = \alpha + o(1),$$

and if $\alpha \geq 0.25$ so that $0 \leq c_{1-\alpha} \leq 1$,

$$(7.21) \quad \mathbf{E}[\varphi_n] = 1 - P\{V'_n \leq c_{0.75}\} \leq 1 - P\{V''_n \leq c_{0.75}\} = 0.25 \leq \alpha + o(1),$$

as $n \rightarrow \infty$. Hence, the size of the test φ_n is bounded by α asymptotically.

Suppose that we are under the local alternatives such that $\mathbf{E}[X_n] = \bar{\mu}_n \sigma_{LR} \mathbf{1}_n / \sqrt{n}$, for some sequence $\bar{\mu}_n \rightarrow \infty$. Define

$$\bar{U}_n = U'_n - \frac{1}{2} \sum_{m=1}^2 \sqrt{\frac{n_m}{n}} \bar{\mu}_n \sigma_{LR}.$$

Note that

$$(7.22) \quad \begin{aligned} P\{V'_n > c_{1-\alpha}\} &= P\left\{ \frac{\sqrt{2}\bar{U}_n}{\sigma_{LR}} > c_{1-\alpha} \sqrt{\frac{T'_n}{\sigma_{LR}^2}} - \frac{1}{\sqrt{2}} \sum_{m=1}^2 \sqrt{\frac{n_m}{n}} \bar{\mu}_n \right\} \\ &\geq P\left\{ \frac{\sqrt{2}\bar{U}_n}{\sigma_{LR}} > c_{1-\alpha} \sqrt{\frac{T'_n}{\sigma_{LR}^2}} - \frac{1}{\sqrt{2}} \sum_{m=1}^2 \frac{n_m}{n} \bar{\mu}_n \right\}, \end{aligned}$$

because $n_m/n \leq 1$. Note that

$$(7.23) \quad \sigma_{LR}^2 = \sum_{m=1}^{M_n} \frac{n_m}{n} \sigma_{n,m}^2 \geq \sum_{m=1}^2 \frac{n_m}{n} \sigma_{n,m}^2.$$

Since

$$0 < \liminf_{n \rightarrow \infty} \min_{m=1,2} \frac{n_m}{n} \leq \limsup_{n \rightarrow \infty} \min_{m=1,2} \frac{n_m}{n} \leq 1,$$

there exist $\epsilon > 0$ and $n_0 > 0$ such that for all $n \geq n_0$,

$$\sigma_{LR}^2 \geq \epsilon \sum_{m=1}^2 \sigma_{n,m}^2 \geq \epsilon \max_{m=1,2} \sigma_{n,m}^2.$$

Therefore,

$$\frac{|\bar{U}_n|}{\sigma_{LR}} \leq \frac{\max_{m=1,2} \sigma_{n,m}}{2\sigma_{LR}} \sum_{m=1}^2 \frac{|\xi_m - \mathbf{E}[\xi_m]|}{\sigma_{n,m}} \leq \frac{1}{2\sqrt{\epsilon}} \sum_{m=1}^2 \frac{|\xi_m - \mathbf{E}[\xi_m]|}{\sigma_{n,m}}.$$

Since $(\xi_m - \mathbf{E}\xi_m)/\sigma_{n,m}$ converges in distribution to $N(0, 1)$ under any sequence $P_n \in \mathcal{P}_n$ as $n \rightarrow \infty$ by (3.2), we have $\bar{U}_n/\sigma_{LR} = O_p(1)$. Similarly, we can show that $T'_n/\sigma_{LR}^2 = O_p(1)$. Hence, the last probability in (7.22) converges to one as $\bar{\mu}_n \rightarrow \infty$, proving that the mean is consistently \sqrt{n} -discriminated. ■

7.3. Impossibility of Consistent Estimation of Long Run Variance

Proof of Lemma 4.1: We first show sufficiency. Suppose that $\|\theta_n(P_{n,1}) - \theta_n(P_{n,0})\| = o(1)$ for all sequence $P_{n,1} \in \mathcal{P}_n$. Then we take $\hat{\theta} = \theta_n(P_{n,0})$, so that $\|\hat{\theta} - \theta_n(P_{n,1})\| \rightarrow 0$, along $P_{n,1} \in \mathcal{P}_n$, as $n \rightarrow \infty$. Hence sufficiency follows.

Conversely, suppose that θ_n is consistently estimable in \mathcal{P}_n , so that there exists an estimator, say, $\tilde{\theta}$, such that $\tilde{\theta} - \theta_n(P_{n,1}) = o_p(1)$ along any $P_{n,1} \in \mathcal{P}_n$. Since $P_{n,0} \in \mathcal{P}_n$, this means that

$\tilde{\theta} - \theta_n(P_{n,0}) = o_p(1)$ under $P_{n,0}$. Since $P_{n,1} \triangleleft P_{n,0}$, $\tilde{\theta} - \theta_n(P_{n,0}) = o_p(1)$ under any $P_{n,1} \in \mathcal{P}_n$. We choose any $P_{n,1} \in \mathcal{P}_n$ and write

$$(7.24) \quad \tilde{\theta} - \theta_n(P_{n,1}) = \tilde{\theta} - \theta_n(P_{n,0}) + \theta_n(P_{n,0}) - \theta_n(P_{n,1}).$$

The difference on the left hand side and the first difference on the right hand side are $o_p(1)$ under $P_{n,1}$. This implies that $\|\theta_n(P_{n,0}) - \theta_n(P_{n,1})\| = o(1)$. ■

The lemma below essentially shows that the long run variance is not possible to consistently estimate in local-to-independence common shock models.

Lemma 7.9. *It is not possible to consistently estimate σ_{LR}^2 along $\Phi(0, I_n + \delta_n \mathbf{1}_n \mathbf{1}_n^\top - \delta_n I_n)$ with a sequence δ_n such that $(n-1)|\delta_n| < 1$ for all $n \geq 1$, and*

$$(7.25) \quad n\delta_n \rightarrow \delta,$$

for some $\delta \in [-1, 1] \setminus \{0\}$.

Proof: By Lemma 7.3, we have (under the sequence $\{\Phi(0, I_n)\}$)

$$\begin{aligned} \log \frac{d\Phi(0, \Sigma_n)}{d\Phi(0, I_n)}(X_n) &= -\frac{1}{2} \log(1 + (n-1)\delta_n) - \frac{1}{2} \log(1 - \delta_n)^{n-1} \\ &\quad + \frac{(n-1)\delta_n Z_{n,1}^2}{2(1 + (n-1)\delta_n)} - \frac{(n-1)\delta_n}{2(1 - \delta_n)} - \frac{(n-1)\delta_n}{2(1 - \delta_n)} \frac{1}{n-1} \sum_{k=2}^n (Z_{n,k}^2 - 1) \\ &= -\frac{1}{2} \log(1 + (n-1)\delta_n) \\ &\quad + \frac{(n-1)\delta_n Z_{n,1}^2}{2(1 + (n-1)\delta_n)} - \frac{(n-1)\delta_n}{2(1 - \delta_n)} \frac{1}{n-1} \sum_{k=2}^n (Z_{n,k}^2 - 1) + o(1) \\ &\rightarrow_d -\log \sqrt{1 + \delta} + \frac{\delta \mathbb{Z}^2}{2(1 + \delta)}, \end{aligned}$$

as $(n-1)\delta_n \rightarrow \delta$, where $\mathbb{Z} \sim \mathcal{N}(0, 1)$. It is not hard to check that

$$\mathbb{E} \left[\exp \left(-\frac{1}{2} \log(1 + \delta) + \frac{1}{2} \frac{\delta \mathbb{Z}^2}{1 + \delta} \right) \right] = 1.$$

Hence, by Le Cam's First Lemma (e.g., [van der Vaart, 1998](#), Lemma 6.4), it follows that

$$\Phi(0, \Sigma_n) \triangleleft \Phi(0, I_n).$$

However,

$$\sigma_{LR}^2 = 1 + (n-1)\delta_n \rightarrow 1 + \delta \neq 1.$$

By Lemma 4.1, it is impossible to consistently estimate σ_{LR}^2 along $\Phi(0, I_n + \delta_n \mathbf{1}_n \mathbf{1}_n^\top - \delta_n I_n)$. ■

Proof of Theorem 4.1: Let us first show sufficiency. Suppose that either (a) or (b) in Lemma 7.5 holds. Let us take

$$\hat{\sigma}_{LR}^2 = \frac{1}{n} \sum_{m=1}^{M_n} \sum_{i,j \in N_{n,m}} (X_{n,i} - \bar{X}_n)(X_{n,j} - \bar{X}_n).$$

Since $\sigma_{LR}^2 \leq c$ for all $n \geq 1$, we have

$$\bar{X}_n = \mathbf{E}[X_{n,i}] + o_p(1).$$

Hence

$$\hat{\sigma}_{LR}^2 = \frac{1}{n} \sum_{m=1}^{M_n} \sum_{i,j \in N_{n,m}} (X_{n,i} - \mathbf{E}[X_{n,i}])(X_{n,j} - \mathbf{E}[X_{n,i}]) + o_p(1).$$

Note that

$$(7.26) \quad \sum_{m=1}^{M_n} \left(\frac{n_m}{n}\right)^2 \leq \sum_{m=1}^{M_n} \frac{n_m}{n} \leq 1.$$

Choose $P_n \in \mathcal{P}_{n,\mathcal{N}}$. Then, under P_n ,

$$(7.27) \quad \begin{aligned} \mathbf{E}\left[(\hat{\sigma}_{LR}^2 - \sigma_{LR}^2)^2\right] &= \frac{1}{n^2} \sum_{m=1}^{M_n} \sum_{i,j \in N_{n,m}} \sum_{i',j' \in N_{n,m}} \text{Cov}(X_{n,i}X_{n,j}, X_{n,i'}X_{n,j'}) \\ &\leq \frac{C'}{n^2} \sum_{m=1}^{M_n} n_m^2 = \frac{C'}{n^2} \left((n - n^*) + \sum_{m=1:n_m \geq 2}^{M_n} n_m^2 \right) \\ &= C' \left(\frac{n - n^*}{n^2} + C' \left(\frac{n^*}{n}\right)^2 \sum_{m=1:n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*}\right)^2 \right), \end{aligned}$$

where $C' > 0$ is a constant that does not depend on n . By (7.26), the last term is $o(1)$, if either of the conditions (a) and (b) in Lemma 7.5. Therefore, σ_{LR}^2 is consistently estimable in \mathcal{P}_n .

Now, let us show necessity. Suppose that both (a) and (b) in Lemma 7.5 are violated. That is,

$$(7.28) \quad \limsup_{n \rightarrow \infty} \frac{n^*}{n} > 0 \quad \text{and} \quad \limsup_{n \rightarrow \infty} \sum_{m \geq 1:n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*}\right)^2 > 0.$$

Let $\Pi_{\mathcal{M}_n}$ be the set of permutations π_n on $\{1, \dots, n\}$ such that their matrix representation is block diagonal along the partition \mathcal{M}_n of $\{1, \dots, n\}$. Let $F_{n,\mathcal{M}_n}(\cdot)$ is the uniform distribution over Π_{n,\mathcal{M}_n} . Let us fix $c > 0$ and let $\mathcal{S}_n^0(c) \subset \mathcal{S}_n^0$ be the set of the matrices in \mathcal{S}_n^0 such that each diagonal block $\Delta_{n,m}$ has eigenvalues bounded by $c - 1$. Define

$$\mathcal{P}'_{n,\mathcal{N}}(c) \equiv \left\{ \Phi(0, \Sigma_n) : \Sigma_n = I_n + \int \pi_n \Delta_n \pi_n^\top dF_{n,\mathcal{M}_n}(\pi_n), \Delta_n \in \mathcal{S}_n^0(c), \Sigma_n \in \mathcal{S}_n^+ \right\}.$$

Note that for any $\Delta_n \in \mathcal{S}_n^0(c)$, for each $m = 1, \dots, M_n$, if $n_m = 1$, its m -th block $\Delta_{n,m}$ is zero, and if $n_m \geq 2$, we have

$$(7.29) \quad \frac{\mathbf{1}_{n_m}^\top \Delta_{n,m} \mathbf{1}_{n_m}}{n_m - 1} = \frac{\text{tr}(\Delta_{n,m} \mathbf{1}_{n_m} \mathbf{1}_{n_m}^\top)}{n_m - 1} \leq \frac{\lambda_{\max}(\Delta_{n,m}) \text{tr}(\mathbf{1}_{n_m} \mathbf{1}_{n_m}^\top)}{n_m - 1} \leq \frac{cn_m}{n_m - 1}.$$

Hence,

$$\frac{1}{n} \mathbf{1}' \left(I_n + \int \pi_n \Delta_n \pi_n^\top dF_n(\pi_n) \right) \mathbf{1} \leq 1 + \frac{\mathbf{1}' \Delta_n \mathbf{1}}{n} \leq 1 + \frac{\lambda_{\max}(\Delta_n) \text{tr}(\mathbf{1}_n \mathbf{1}_n^\top)}{n} \leq c,$$

because $\lambda_{\max}(\Delta_n) \leq \max_{1 \leq m \leq M_n} \lambda_{\max}(\Delta_{n,m}) \leq c$. This means that the long run variance under any $P_n \in \mathcal{P}'_{n,\mathcal{N}}(c)$ is bounded by c . Hence

$$(7.30) \quad \mathcal{P}'_{n,\mathcal{N}}(c) \subset \mathcal{P}_{n,\mathcal{N}}.$$

It suffices to show that it is impossible to consistently estimate the long run variance in $\mathcal{P}'_{n,\mathcal{N}}(c)$.

By (7.26), there exists a subsequence $\{n_k\} \subset \{n\}$ such that

$$(7.31) \quad \sum_{m=1}^{M_{n_k}} \left(\frac{n_{m,n_k}}{n_k} \right)^2 \rightarrow \tilde{c}_1 > 0 \quad \text{and} \quad \frac{n_k^*}{n_k} \rightarrow \tilde{c}_2,$$

for some constants $\tilde{c}_1, \tilde{c}_2 \in (0, 1]$. For simplicity of notation, we fix this subsequence, and identify it as $\{n\}$. For each $m = 1, \dots, M_n$ such that $n_m \geq 2$, we let

$$(7.32) \quad \delta_{n,m} = \frac{\mathbf{1}'_{n_m} \Delta_{n,m} \mathbf{1}_{n_m}}{n_m(n_m - 1)}.$$

In particular, we consider a sequence of block matrices $\Delta_n \in \mathcal{S}_n^0(c)$ such that (7.10) holds for some $\bar{\delta} \in [-a, a] \setminus \{0\}$, $a < 1/2$, if $n_m \geq 2$. Such a sequence Δ_n exists in $\mathcal{S}_n^0(c)$ by (7.29). Note that the matrix

$$\int \pi_n \Delta_n \pi_n^\top dF_{n,\mathcal{M}_n}(\pi_n)$$

is a block diagonal matrix whose m -th block is given by

$$\delta_{n,m} \mathbf{1}_{n_m} \mathbf{1}_{n_m}^\top - \delta_{n,m} I_{n_m}.$$

Now we choose

$$\Sigma_n = I_n + \int \pi_n \Delta_n \pi_n^\top dF_{n,\mathcal{M}_n}(\pi_n),$$

and show that $\Phi(0, \Sigma_n) \triangleleft \Phi(0, I_n)$. By Lemma 7.6(ii), $\log(d\Phi(0, \Sigma_n)/d\Phi(0, I_n))(X_n)$ is uniformly tight under $\Phi(0, I_n)$. Furthermore, by Prohorov's Theorem, there exists a subsequence $\{n_k\}$ of $\{n\}$ such that $\log(d\Phi(0, \Sigma_{n_k})/d\Phi(0, I_{n_k}))(X_n)$ weakly converges. Let W be a random variable whose distribution is identical to the weak limit. By the Continuous Mapping Theorem, we

have

$$\frac{d\Phi(0, \Sigma_{n_k})}{d\Phi(0, I_{n_k})}(X_{n_k}) \rightarrow_d e^W,$$

along the subsequence $\{n_k\}$. Note that $\mathbf{E}[(d\Phi(0, \Sigma_{n_k})/d\Phi(0, I_{n_k}))(X_{n_k})] = 1$, where the expectation is under $\Phi(0, I_{n_k})$. By Lemma 7.6(i), $(d\Phi(0, \Sigma_{n_k})/d\Phi(0, I_{n_k}))(X_{n_k})$ is uniformly integrable under $\Phi(0, I_{n_k})$. Hence, we find that $\mathbf{E}e^W = 1$. By Le Cam's First Lemma (e.g., van der Vaart, 1998, Lemma 6.4), we conclude that $\Phi(0, \Sigma_{n_k}) \triangleleft \Phi(0, I_{n_k})$.

On the other hand, note that

$$\begin{aligned} \frac{\mathbf{1}_n^\top \int \pi_n \Delta_n \pi_n^\top dF_{n, \mathcal{M}_n}(\pi_n) \mathbf{1}_n}{n} &= \frac{\mathbf{1}_n^\top \Delta_n \mathbf{1}_n}{n} \\ &= \frac{1}{n} \sum_{m=1: n_m \geq 2}^{M_n} \mathbf{1}_{n_m}^\top \Delta_{n,m} \mathbf{1}_{n_m} \\ &= \bar{\delta} \sum_{m=1: n_m \geq 2}^{M_n} \frac{n_m(n_m-1)}{nn^*} = \bar{\delta} \sum_{m=1}^{M_n} \frac{n_m^2}{nn^*} - \bar{\delta} \sum_{m=1: n_m \geq 2}^{M_n} \frac{n_m}{nn^*}. \end{aligned}$$

We rewrite the last term as

$$\begin{aligned} \bar{\delta} \sum_{m=1: n_m \geq 2}^{M_n} \frac{n_m^2}{nn^*} - \frac{\bar{\delta}}{n} &= \bar{\delta} \sum_{m=1: n_m \geq 2}^{M_n} \frac{n_m^2}{nn^*} + o(1) \\ &= \frac{\bar{\delta} n^*}{n} \sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*}\right)^2 + o(1) \rightarrow \bar{\delta} \tilde{c}_1 \tilde{c}_2 \neq 0, \end{aligned}$$

as $n \rightarrow \infty$, where the last convergence is due to (7.31). Certainly, σ_{LR}^2 is consistently estimable along $\Phi(0, I_n)$. By Lemma 4.1, we conclude that σ_{LR}^2 is not consistently estimable in $\mathcal{P}'_{n, \mathcal{N}}(c)$. By (7.30), it is not consistently estimable in $\mathcal{P}_{n, \mathcal{N}}$ either. ■

Proof of Corollary 4.1: Let us show sufficiency. First suppose that \mathcal{M}_n consists of negligible clusters, so that $\max_{1 \leq m \leq M_n} n_m/n \rightarrow 0$, as $n \rightarrow \infty$. From (7.26), this implies that

$$\sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*}\right)^2 \leq \left(\frac{n}{n^*}\right) \max_{1 \leq m \leq M_n} \frac{n_m}{n} \rightarrow 0,$$

as $n \rightarrow \infty$. Therefore, σ_{LR}^2 is consistently estimable in \mathcal{P}_n by Theorem 4.1.

Conversely, suppose that for some $\epsilon > 0$,

$$\limsup_{n \rightarrow \infty} \max_{1 \leq m \leq M_n} \frac{n_m}{n} > \epsilon.$$

Then there exist subsequences $\{n_k\} \subset \{n\}$ and $\{n_{m(n_k)}\} \subset \{n_{m(n)}\}_{n \geq 1}$, $m(n) \in \{1, \dots, M_n\}$, such that

$$\lim_{k \rightarrow \infty} \frac{n_{m(n_k)}}{n_k} > \epsilon.$$

This implies that

$$\limsup_{n \rightarrow \infty} \sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*} \right)^2 > 0.$$

Hence, σ_{LR}^2 is not consistently estimable in \mathcal{P}_n by Theorem 4.1. ■

Proof of Corollary 4.2: (i) Let $N_o \subset \{1, \dots, n\}$ be the set of nodes in the clique with size n_c . Fix $c > 0$ and let $\mathcal{S}'_n(c)$ be the set of symmetric $n \times n$ matrices with zero diagonals, whose eigenvalues are bounded by $c - 1$. Take \mathcal{M}_n to be the cluster structure such that there is only one non-singleton cluster that is N_o and let \mathcal{S}_n^0 be the set of symmetric $n \times n$ matrices with zero diagonals and block diagonal along \mathcal{M}_n . Let

$$\mathcal{S}_n(c; G_n) = \{\Delta_n \in \mathcal{S}'_n(c) : \Delta_{n,ij} = 0, \text{ if and only if } ij \notin E_n\},$$

where $\Delta_{n,ij}$ denotes the (i, j) -th entry of Δ_n . Then,

$$\mathcal{S}_n^0(c) \subset \mathcal{S}_n(c; G_n).$$

Therefore, it suffices to show that σ_{LR}^2 is not consistently estimable in $\mathcal{P}_{n, \mathcal{N}}(c)$ with the cluster structure \mathcal{M}_n . Note that

$$\sum_{m=1: n_m \geq 2}^{M_n} \left(\frac{n_m}{n^*} \right)^2 = 1,$$

because we have only one non-singleton cluster in \mathcal{M}_n . By Theorem 4.1, σ_{LR}^2 is not consistently estimable in $\mathcal{P}_{n, \mathcal{N}}(c)$.

(ii) Let us define $\bar{N}(i) = \{j \in N_n : ij \in \bar{E}_n\}$, where $\bar{E}_n = E_n \cup \{ii : i \in N_n\}$, and consider

$$\hat{\sigma}_{LR}^2 = \frac{1}{n} \sum_{i=1}^n \sum_{j \in \bar{N}(i)} (X_{n,i} - \bar{X}_n)(X_{n,j} - \bar{X}_n).$$

By rearranging terms, we can write

$$(7.33) \quad \begin{aligned} \hat{\sigma}_{LR}^2 - \sigma_{LR}^2 &= \frac{1}{n} \sum_{i=1}^n \sum_{j: ij \in \bar{E}_n} (X_{n,i} X_{n,j} - \mathbf{E}[X_{n,i} X_{n,j}]) - 2(\bar{X}_n - \mathbf{E}[X_{n,i}]) \frac{1}{n} \sum_{j \in \bar{N}(i)} X_{n,j} \\ &\quad - 2\mathbf{E}[X_{n,i}] \frac{1}{n} \sum_{i=1}^n \sum_{j \in \bar{N}(i)} (X_{n,j} - \mathbf{E}X_{n,j}) + \bar{X}_n^2 - (\mathbf{E}[X_{n,i}])^2 \frac{1}{n} \sum_{i=1}^n |\bar{N}(i)|. \end{aligned}$$

As for the squared L^2 norm of the leading term on the right hand side, we can write it as

$$\frac{1}{n^2} \sum_{i_1=1}^n \sum_{i_2=1}^n \sum_{j_1: i_1 j_1 \in \bar{E}_n} \sum_{j_2: i_2 j_2 \in \bar{E}_n} \text{Cov}(X_{n,i_1} X_{n,j_1}, X_{n,i_2} X_{n,j_2}).$$

If $\{i_1, j_1\}$ and $\{i_2, j_2\}$ are not adjacent in G_n , the covariance above is zero by the dependency graph assumption. The number of the terms in the above sum such that $\{i_1, j_1\}$ and $\{i_2, j_2\}$ are adjacent in G_n is of the order $O(nd_{mx}^2 d_{av}) = o(n^2)$. The last rate comes from our assumption that $\lim_{n \rightarrow \infty} d_{mx}^2 d_{av}/n = 0$. Therefore, the leading term on the right hand side of (7.33) is $o_p(1)$. Similarly, we can show that the remainder terms are $o_p(1)$. Hence, $\hat{\sigma}_{LR}^2$ is a consistent estimator of σ_{LR}^2 . ■

References

- ARONOW, P., AND C. SAMII (2017): “Estimating Average Causal Effects Under General Interference, with Application to a Social Network Experiment,” *Annals of Applied Statistics*, 11(4), 1912–1947.
- BAHADUR, R. R., AND L. J. SAVAGE (1956): “The Nonexistence of Certain Statistical Procedures in Nonparametric Problems,” *Annals of Mathematical Statistics*, 27, 1115–1122.
- BAKIROV, N. K., AND G. J. SZEKELY (2005): “Student’s t -Test for Gaussian Scale Mixtures,” *Zapiski Nauchnyh Seminarov POMI*, 328, 5–19.
- BERTANHA, M., AND M. J. MOREIRA (2020): “Impossible Inference in Econometrics: Theory and Applications,” *Journal of Econometrics*, 218, 247–270.
- BESTER, A. C., T. G. CONLEY, AND C. B. HANSEN (2011): “Inference with Dependent Data Using Cluster Covariance Estimators,” *Journal of Econometrics*, 165, 137–151.
- CAMERON, A. C., J. B. GELBACH, AND D. L. MILLER (2008): “Bootstrap Based Improvements for Inference with Clustered Errors,” *Review of Economics and Statistics*, 90, 414–427.
- CANAY, I. A., J. P. ROMANO, AND A. M. SHAIKH (2017): “Randomization Tests Under an Approximate Symmetry Assumption,” *Econometrica*, 85, 1013–1030.
- CANAY, I. A., A. SANTOS, AND A. M. SHAIKH (2021): “The Wild Bootstrap with a “Small” Number of “Large” Clusters,” *Review of Economics and Statistics*, 103, 346–363.
- CANEN, N., J. SCHWARTZ, AND K. SONG (2020): “Estimating Local Interactions among Many Agents Who Observe Their Neighbors,” *Quantitative Economics*, 11, 346–363.
- CONLEY, T. G., AND C. R. TABER (2011): “Inference with “Difference in Differences” with a Small Number of Policy Changes,” *Review of Economics and Statistics*, 93, 113–125.
- DJOGBENOU, A. A., J. G. MACKINNON, AND M. O. NIELSEN (2019): “Asymptotic Theory and Wild Bootstrap Inference with Clustered Errors,” *Journal of Econometrics*, 212, 393–412.

- DONALD, S. G., AND K. LANG (2007): “Inference with Difference-in-Difference and Other Panel Data,” *Review of Economics and Statistics*, 89, 221–233.
- DUFOUR, J. M. (1997): “Some Impossibility Theorems in Econometrics with Applications to Structural and Dynamic Models,” *Econometrica*, 65, 1365–1387.
- HAGEMANN, A. (2019): “Placebo Inference on Treatment Effects When the Number of Clusters is Small,” *Journal of Econometrics*, 213, 190–209.
- HANSEN, B. E., AND S. LEE (2019): “Asymptotic Theory for Clustered Samples,” *Journal of Econometrics*, 210, 268–290.
- HE, X., AND K. SONG (2022): “Measuring Diffusion over a Large Network,” *arXiv: 1812.04195v4 [stat.ME]*.
- IBRAGIMOV, I. A., AND R. Z. HAS’MINSKII (1981): *Statistical Estimation: Asymptotic Theory*. Springer Science+Business Media, New York.
- IBRAGIMOV, R., AND U. K. MÜLLER (2010): “t-Statistic Based Correlation and Heterogeneity Robust Inference,” *Journal of Business and Economic Statistics*, 28, 453–468.
- (2016): “Inference With Few Heterogeneous Clusters,” *Review of Economics and Statistics*, 98, 83–96.
- IMBENS, G. W., AND J. M. WOOLDRIDGE (2009): “Recent Developments in the Econometrics of Program Evaluation,” *Journal of Economic Literature*, 47, 5–86.
- KIEFER, N. M., AND T. J. VOGELSANG (2002): “Heteroskedasticity-Autocorrelation Robust Standard Errors Using the Bartlett Kernel without Truncation,” *Econometrica*, 70, 2093–2095.
- KOJEVNIKOV, D., V. MARMER, AND K. SONG (2021): “Limit theorems for Network Dependent Random Variables,” *Journal of Econometrics*, 222, 882–908.
- LECAM, L. (1960): “Locally Asymptotically Normal Families of Distributions,” *University of California Publications in Statistics*, 3, 37–98.
- LECAM, L., AND L. SCHWARTZ (1960): “A Necessary and Sufficient Condition for the Existence of Consistent Estimates,” *Annals of Mathematical Statistics*, 31, 140–150.
- LEHMANN, E. L., AND J. P. ROMANO (2005): *Testing Statistical Hypotheses*. Springer, New York.
- LEUNG, M. P. (2020): “Treatment and Spillover Effects Under Network Interference,” 102, 368–380, *Review of Economics and Statistics*, forthcoming.
- (2021): “Dependence-Robust Inference Using Resampled Statistics,” *Journal of Applied Econometrics*, 37, 270–285.
- MACKINNON, J. G., AND M. D. WEBB (2017): “Wild bootstrap Inference for Wildly Different Cluster Sizes,” *Journal of Applied Econometrics*, 32, 233–254.
- (2020): “Randomization Inference for Difference-in-Differences with Few Treated Clusters,” *Journal of Econometrics*, 218, 435–450.
- MENZEL, K. (2021): “Bootstrap with Cluster-Dependence in Two or More Dimensions,” *Econometrica*, 89, 2143–2188.

- PFANZAGL, J. (1994): *Parametric Statistical Theory*. De Gruyter, Berlin.
- PÖTSCHER, B. M. (2002): “Lower Risk Bounds and Properties of Confidence Sets for Ill-Posed Estimation Problems with Applications to Spectral Density and Persistence Estimation, Unit Roots, and Estimation of Long Memory Parameters,” *Econometrica*, 70, 1035–1065.
- ROTH, J., P. H. C. SANT’ANNA, A. BILINSKI, AND J. POE (2022): “What’s Trending in Difference-in-Differences? A Synthesis of the Recent Econometrics Literature,” *arXiv:2201.01194v2 [econ.EM]*.
- SONG, K. (2016): “Ordering-Free Inference from Locally Dependent Data,” *arXiv:1604.00447v1 [stat.ME]*.
- (2018): “Measuring the Graph Concordance of Locally Dependent Observations,” *Review of Economics and Statistics*, 100, 535–549.
- STEIN, C. (1986): “Approximate Computation of Expectations,” *Lecture Notes-Monograph Series*, 7, i–164.
- SUN, Y. (2014): “Fixed-Smoothing Asymptotics in a Two-Step Generalized Method of Moments Framework,” *Econometrica*, 82, 2327–2370.
- VAN DER VAART, A. W. (1998): *Asymptotic Statistics*. Cambridge University Press, New York, USA.