

CONTINUUM LIMITS FOR DISCRETE DIRAC OPERATORS ON 2D SQUARE LATTICES

KARL MICHAEL SCHMIDT AND TOMIO UMEDA

2020 Mathematics Subject Classification: Primary 47A10; Secondary 47B37, 47B93

Keywords: discrete Dirac operators, Dirac operators on square lattices, discrete Fourier transform, continuum limits, spectrum, complex potentials.

ABSTRACT. We discuss the continuum limit of discrete Dirac operators on the square lattice in \mathbb{R}^2 as the mesh size tends to zero. To this end, we propose a natural and simple embedding of $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$ that enables us to compare the discrete Dirac operators with the continuum Dirac operators in the same Hilbert space $L^2(\mathbb{R}^2)^2$. In particular, we prove strong resolvent convergence. Potentials are assumed to be bounded and uniformly continuous functions on \mathbb{R}^2 and allowed to be complex matrix-valued.

1. INTRODUCTION

This paper is concerned with the discrete Dirac operator $\mathbb{D}_{m,h} + V_h$ defined by

$$(1.1) \quad \mathbb{D}_{m,h} + V_h = \begin{pmatrix} m & i\partial_{1,h}^* + \partial_{2,h}^* \\ -i\partial_{1,h} + \partial_{2,h} & -m \end{pmatrix} + V_h \quad \text{in } \ell^2(\mathbb{Z}_h^2)^2,$$

which is a discrete analogue of the two-dimensional Dirac operator defined by

$$(1.2) \quad \mathbb{D}_m + V = -i\sigma_1 \frac{\partial}{\partial x_1} - i\sigma_2 \frac{\partial}{\partial x_2} + m\sigma_3 + V(x) \quad \text{in } L^2(\mathbb{R}^2)^2,$$

where $m \geq 0$ and $\sigma_1, \sigma_2, \sigma_3$ are the Pauli matrices,

$$(1.3) \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

and V is a complex matrix valued function. For the definition of $\ell^2(\mathbb{Z}_h^2)^2$, see (2.1) in section 2; the finite difference operators $\partial_{j,h}$ and $\partial_{j,h}^*$ ($j \in \{1, 2\}$) are defined in (2.3) and (2.4) in section 2, respectively; for V_h , see (5.3) in section 5.

We remark that both operators $\mathbb{D}_{m,h}$ in (1.1) and \mathbb{D}_m in (1.2) possess supersymmetry structure (see [25, Chapter 5], [26, Chapter 3]). The discrete Dirac operator (1.1) can be rewritten in a form analogous to (1.2),

$$\mathbb{D}_{m,h} = -i\sigma_1 \begin{pmatrix} \partial_{1,h} & 0 \\ 0 & -\partial_{1,h}^* \end{pmatrix} - i\sigma_2 \begin{pmatrix} \partial_{2,h} & 0 \\ 0 & -\partial_{2,h}^* \end{pmatrix} + m\sigma_3 + V_h.$$

It is widely recognized that 2D Dirac operators, especially in the massless case, have been the object of extensive research in the context of graphene since its

Date: February 9, 2022.

T.Umeda was partially supported by the Japan Society for the Promotion of Science ‘‘Grant-in-Aid for Scientific Research’’ (C) No. 26400175.

discovery in 2004, see [5] or [18] for an exposition. In particular, we would like to mention the work [17], which reported that electron transport is essentially governed by a massless Dirac equation and that a variety of unusual phenomena are characteristic of two-dimensional Dirac fermions. These are the main reasons why we focus on the two-dimensional case, although it is apparent that the methods and ideas developed in the present paper are directly applicable to the one-dimensional and the three-dimensional cases.

It is quite natural to make an attempt to show that the discrete operator (1.1) converges to the continuum operator (1.2) as the mesh size h of the lattice \mathbb{Z}_h^2 tends to 0. However, there is a difficulty in that these two operators work in completely different Hilbert spaces. For example, it is not immediately obvious how one can make sense of the expression $(\mathbb{D}_{m,h} + V_h) - (\mathbb{D}_m + V)$. For this reason, it is necessary to embed $\ell^2(\mathbb{Z}_h^2)^2$ into an appropriate function space on \mathbb{R}^2 .

In this paper, we propose a simple and natural embedding of $\ell^2(\mathbb{Z}_h^2)^2$ into $L^2(\mathbb{R}^2)^2$ by assigning to each element in $\ell^2(\mathbb{Z}_h^2)^2$ a step function in $L^2(\mathbb{R}^2)^2$ (see subsection 2.1 for a precise description). This idea of embedding $\ell^2(\mathbb{Z}_h^2)^2$ into $L^2(\mathbb{R}^2)^2$ induces a subspace $L^2(\mathbb{Z}_h^2)^2$ of $L^2(\mathbb{R}^2)^2$ where one can naturally define the difference operators $\partial_{j,h}$ and $\partial_{j,h}^*$. We emphasize that the induced difference operators are the exact counterparts of the difference operators in $\ell^2(\mathbb{Z}_h^2)^2$.

When we apply our idea to the discrete Dirac operator (1.1) to discuss the continuum limit as the mesh size h tends to 0, we require a convergence theorem for the orthogonal projection P_h onto the closed subspace $L^2(\mathbb{Z}_h^2)^2$ of $L^2(\mathbb{R}^2)^2$. In connection with this, we also introduce a discrete Fourier transform in $L^2(\mathbb{R}^d)$ (related, but not identical to the d -dimensional Fourier series with coefficients in $\ell^2(\mathbb{Z}_h^d)$), enabled by the natural embedding of $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$, and prove a convergence theorem for the discrete Fourier transform as $h \rightarrow 0$. Specifically, we establish both convergence theorems in the sense of strong convergence in $L^2(\mathbb{R}^d)$. Precise descriptions are given in subsection 2.2 and section 3. With these strong convergence theorems, we can prove that the resolvents of the discrete Dirac operator (1.1) strongly converge to those of the Dirac operator (1.2) in $L^2(\mathbb{R}^2)^2$.

In the literature, there have been few papers studying spectral properties of discrete Dirac operators on 2D or 3D lattices, while there have been many working on 1D lattices, see for example the recent works [1], [4], [6], [7], [13], [14], [15], [19], [22], [23], [24]. We mention in passing that the discrete Dirac operator on a 1D lattice, when written in matrix form, is a tri-diagonal matrix and indeed unitarily equivalent to a discrete Schrödinger-type operator; for example, with the unitary operator $U : \ell^2(\mathbb{Z}_h) \rightarrow \ell^2(\mathbb{Z}_h)^2$ given as $(Uf)(nh) = (-1)^n \begin{pmatrix} f(2nh) \\ f((2n+1)h) \end{pmatrix}$,

$$U^* \begin{pmatrix} m & \partial_h^* \\ \partial_h & -m \end{pmatrix} U = h(-\partial_h^* \partial_h + q),$$

where $q(nh) = (-1)^n m/h - 2/h^2$.

To our knowledge, the only paper working in dimension 2 or 3 is [20], where dynamical localization for random Dirac operators on d -dimensional lattices with $d \in \{1, 2, 3\}$ was discussed. The lack of works on the continuum limit of discrete Dirac operators in the literature, as far as we know, is hardly surprising in view of the fact that research on the continuum limit of discrete analogs of quantum Hamiltonians began rather recently; see [16] and [12].

It is worth mentioning that a nice embedding of $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$ was introduced in [16], where the embedded discrete Schrödinger operators, in contrast to the present paper, were proved to converge to the continuum Schrödinger operator in the norm resolvent sense as the mesh size h tends to 0. (However, the expression “norm resolvent sense” here is a slight abuse of terminology because the Hilbert space for the discrete operators in [16] is different from that for the continuum operators; see [21, p.284] for the definition of the convergence the norm resolvent sense.) What allows the work [16] to derive the convergence in the norm resolvent sense is the identification operator used for the embedding of $\ell^2(\mathbb{Z}_h^d)$.

In [12] a natural embedding of $\ell^2(\mathbb{Z}_h^d)$ into $\mathcal{S}'(\mathbb{R}^d)$, the space of tempered distributions, was introduced and continuum limits of lattice Schrödinger operators for various models were investigated. In particular, lattice Laplacians satisfying suitable assumptions were shown to converge to the 2D Dirac operators (1.2) with $m = 0$ and $V = 0$. Specifically for the hexagonal (graphene) lattice, see also [8].

The present paper is organised as follows. Section 2 illustrates the idea of embedding $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$ and shows how the finite difference operators in the embedded space can naturally be defined. It also describes how the discrete Fourier transform can be extended as an operator in $L^2(\mathbb{R}^d)$. In section 3, convergence of discrete Fourier transform in $L^2(\mathbb{R}^d)$ is discussed. Resolvent convergence of the discrete Dirac operator without potential is discussed in section 4, based on the results obtained in the previous sections. Strong resolvent convergence of the discrete Dirac operators with potentials is discussed in section 5.

2. EMBEDDING OF $\ell^2(\mathbb{Z}_h^d)$ INTO $L^2(\mathbb{R}^d)$ AND DISCRETE FOURIER TRANSFORM IN $L^2(\mathbb{R}^d)$

In applications to the Dirac operator (1.1), the underlying Hilbert space is $\ell^2(\mathbb{Z}_h^d)^2 := \ell^2(\mathbb{Z}_h^d) \otimes \mathbb{C}^2$, which consists of \mathbb{C}^2 -valued functions on the 2 dimensional lattice. In this section, we focus on the space of complex-valued functions on the d -dimensional lattice, $\ell^2(\mathbb{Z}_h^d)$, and related spaces of functions on \mathbb{R}^d ; the results naturally extend to the corresponding spaces of \mathbb{C}^2 -valued functions.

The d -dimensional square lattice with the mesh size $h > 0$ is denoted by

$$\mathbb{Z}_h^d := \{ hn \mid n \in \mathbb{Z}^d \}.$$

The Hilbert space

$$(2.1) \quad \ell^2(\mathbb{Z}_h^d) := \left\{ f \mid f : \mathbb{Z}_h^d \rightarrow \mathbb{C}, \sum_{n \in \mathbb{Z}^d} |f(hn)|^2 < \infty \right\},$$

has the standard inner product

$$(2.2) \quad (f, g)_{\ell^2(\mathbb{Z}_h^d)} = \sum_{n \in \mathbb{Z}^d} f(hn) \overline{g(hn)}.$$

For $f \in \ell^2(\mathbb{Z}_h^d)$, define

$$(2.3) \quad [\partial_{j,h} f](n) := \frac{1}{h} \left\{ f(h(n + e_j)) - f(hn) \right\} \quad (j \in \{1, \dots, d\}),$$

where $e_1 = (1, 0, \dots, 0), \dots, e_d = (0, \dots, 0, 1)$. The adjoint of $\partial_{j,h}$ is given by

$$(2.4) \quad [\partial_{j,h}^* f](n) := \frac{1}{h} \left\{ f(h(n - e_j)) - f(hn) \right\} \quad (j \in \{1, \dots, d\}),$$

so

$$(2.5) \quad (\partial_{j,h} f, g)_{\ell^2(\mathbb{Z}_h^d)} = (f, \partial_{j,h}^* g)_{\ell^2(\mathbb{Z}_h^d)}, \quad \forall f, g \in \ell^2(\mathbb{Z}_h^d).$$

2.1. Embedding of $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$. We introduce an embedding of $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$ by assigning to $f \in \ell^2(\mathbb{Z}_h^d)$ the step function

$$[J_h f](x) := \sum_{n \in \mathbb{Z}^d} f(hn) \chi_{I_{n,h}}(x) \in L^2(\mathbb{R}^d),$$

where $\chi_{I_{n,h}}$ is a characteristic function of the set

$$(2.6) \quad I_{n,h} := \{x \mid hn_j \leq x_j < h(n_j + 1) \ (j \in \{1, \dots, d\})\};$$

clearly, $\sum_{n \in \mathbb{Z}^d} \chi_{I_{n,h}} \equiv 1$. Note that

$$(2.7) \quad \|J_h f\|_{L^2(\mathbb{R}^d)} = h^{d/2} \|f\|_{\ell^2(\mathbb{Z}_h^d)},$$

so $h^{-d/2} J_h$ is an isometry from $\ell^2(\mathbb{Z}_h^d)$ into $L^2(\mathbb{R}^d)$. Since $\ell^2(\mathbb{Z}_h^d)$ is a Hilbert space, the image $J_h(\ell^2(\mathbb{Z}_h^d))$ is a closed subspace of the Hilbert space $L^2(\mathbb{R}^d)$. We thus have an orthogonal decomposition

$$(2.8) \quad L^2(\mathbb{R}^d) = L^2(\mathbb{Z}_h^d) \oplus L^2(\mathbb{Z}_h^d)^\perp, \quad L^2(\mathbb{Z}_h^d) := J_h(\ell^2(\mathbb{Z}_h^d)).$$

(The notation $L^2(\mathbb{Z}_h^d)$ already appeared in [12], but there it essentially denotes $\ell^2(\mathbb{Z}_h^d)$; for details, see [12, Subsection 2.1].) The orthogonal projection P_h of $L^2(\mathbb{R}^d)$ onto $L^2(\mathbb{Z}_h^d)$ can be described, for general $\varphi \in L^2(\mathbb{R}^d)$, as $P_h \varphi = J_h \tilde{\varphi}$, where

$$\tilde{\varphi}(hn) = \frac{1}{h^d} \int_{I_{n,h}} \varphi(x) dx \quad (n \in \mathbb{Z}^d).$$

Remark 2.1. *The following example illustrates the action of the projection P_h . Let $\varphi \in L^2(\mathbb{R})$ be defined by $\varphi(x) = 0$ ($|x| \geq 2h$), $\varphi(x) = x + 2h$ ($-2h < x < -h$), $\varphi(x) = h$ ($|x| \leq h$), and $\varphi(x) = -x + 2h$ ($h < x < 2h$). This function can be decomposed as $\varphi = f_h + g_h$ with*

$$f_h(x) = \begin{cases} h & \text{if } -h \leq x < h, \\ \frac{h}{2} & \text{if } -2h \leq x < -h \text{ or } h \leq x < 2h, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$g_h(x) = \begin{cases} x + \frac{3h}{2} & \text{if } -2h \leq x < -h, \\ -x + \frac{3h}{2} & \text{if } h \leq x < 2h, \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that $f_h \in L^2(\mathbb{Z}_h)$ and $g_h \in L^2(\mathbb{Z}_h)^\perp$, so $P_h \varphi = f_h$.

As the above embedding gives a one-to-one relationship between the elements of $\ell^2(\mathbb{Z}_h^d)$ and of $L^2(\mathbb{Z}_h^d)$, we can define the finite difference operators $\partial_{j,h}$ and $\partial_{j,h}^*$ on $L^2(\mathbb{Z}_h^d)$ by applying them, as defined in (2.3) and (2.4), to the corresponding element of $\ell^2(\mathbb{Z}_h^d)$, such that

$$(2.9) \quad \partial_{j,h} J_h[f] := J_h[\partial_{j,h} f] \quad \partial_{j,h}^* J_h[f] := J_h[\partial_{j,h}^* f] \quad (f \in \ell^2(\mathbb{Z}_h^d)).$$

Then we again have

$$(2.10) \quad (\partial_{j,h} f, g)_{L^2(\mathbb{Z}_h^d)} = (f, \partial_{j,h}^* g)_{L^2(\mathbb{Z}_h^d)} \quad (f, g \in L^2(\mathbb{Z}_h^d)).$$

in analogy to (2.5).

2.2. Discrete Fourier transform. Let

$$(2.11) \quad \mathbb{T}_{1/h}^d = [-\pi/h, \pi/h]^d = \{\xi \in \mathbb{R}^d \mid |\xi|_\infty \leq \pi/h\},$$

where $|\xi|_\infty = \max\{|\xi_1|, \dots, |\xi_d|\}$. (Although we use a notation alluding to the interpretation, natural in the following, of this set as a flat d -dimensional torus of side length $2\pi/h$, we emphasize that it is a bounded interval in \mathbb{R}^d .) As the functions

$$e_n(\xi) = \left(\frac{h}{2\pi}\right)^{d/2} e^{-ihn \cdot \xi} \quad (\xi \in \mathbb{T}_{1/h}^d; n \in \mathbb{Z}^d)$$

form an orthonormal basis of $L^2(\mathbb{T}_{1/h}^d)$, any function $f \in \ell^2(\mathbb{Z}_h^d)$ serves as a collection of Fourier coefficients for a d -dimensional Fourier series in $L^2(\mathbb{T}_{1/h}^d)$,

$$\sum_{n \in \mathbb{Z}^d} f(hn) e_n(\xi) \quad (\xi \in \mathbb{T}_{1/h}^d).$$

In view of the bijection between $\ell^2(\mathbb{Z}_h^d)$ and $L^2(\mathbb{Z}_h^d)$, this motivates the following definition of a *discrete Fourier transform* $\mathcal{F}_h : L^2(\mathbb{Z}_h^d) \rightarrow L^2(\mathbb{T}_{1/h}^d)$,

$$(2.12) \quad \begin{aligned} [\mathcal{F}_h J_h f](\xi) &:= h^{d/2} \sum_{n \in \mathbb{Z}^d} f(hn) e_n(\xi) \\ &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \sum_{n \in \mathbb{Z}^d} e^{-ihn \cdot \xi} f(hn) \chi_{I_{n,h}}(x) dx \quad (\xi \in \mathbb{T}_{1/h}^d) \end{aligned}$$

for $f \in \ell^2(\mathbb{Z}_h^d)$. By Parseval's identity for the orthonormal basis $\{e_n \mid n \in \mathbb{Z}^d\}$,

$$(2.13) \quad \|\mathcal{F}_h J_h f\|_{L^2(\mathbb{T}_{1/h}^d)}^2 = h^d \sum_{n \in \mathbb{Z}^d} |f(hn)|^2 = h^d \|f\|_{\ell^2(\mathbb{Z}_h^d)}^2 = \|J_h f\|_{L^2(\mathbb{Z}_h^d)}^2$$

for any $f \in \ell^2(\mathbb{Z}_h^d)$, so \mathcal{F}_h is a unitary operator.

Its inverse is the operator $\overline{\mathcal{F}}_h : L^2(\mathbb{T}_{1/h}^d) \rightarrow L^2(\mathbb{Z}_h^d)$,

$$(2.14) \quad [\overline{\mathcal{F}}_h u](x) = \sum_{n \in \mathbb{Z}^d} \left\{ (2\pi)^{-d/2} \int_{\mathbb{T}_{1/h}^d} e^{ihn \cdot \xi} u(\xi) d\xi \right\} \chi_{I_{n,h}}(x) \quad (x \in \mathbb{R}^d).$$

for $u \in L^2(\mathbb{T}_{1/h}^d)$.

By direct computations, we have

$$(2.15) \quad [\mathcal{F}_h(\partial_{j,h} f)](\xi) = \frac{1}{h} (e^{ih\xi_j} - 1) [\mathcal{F}_h f](\xi) \quad (f \in L^2(\mathbb{Z}_h^d))$$

and

$$(2.16) \quad [\mathcal{F}_h(\partial_{j,h}^* f)](\xi) = \frac{1}{h} (e^{-ih\xi_j} - 1) [\mathcal{F}_h f](\xi) \quad (f \in L^2(\mathbb{Z}_h^d)).$$

Extending functions by 0 outside $\mathbb{T}_{1/h}^d$, the space $L^2(\mathbb{T}_{1/h}^d)$ naturally forms a closed subspace of $L^2(\mathbb{R}^d)$; it is the range of the orthogonal projection $Q_{1/h}$ defined as the operator of multiplication with the characteristic function of $\mathbb{T}_{1/h}^d$.

Using the projections $P_h \in \mathbf{B}(L^2(\mathbb{R}^d))$ and $Q_{1/h} \in \mathbf{B}(L^2(\mathbb{R}^d))$, we can extend \mathcal{F}_h and its inverse $\overline{\mathcal{F}}_h$ to become elements of $\mathbf{B}(L^2(\mathbb{R}^d))$ by setting

$$(2.17) \quad \mathcal{F}_h := Q_{1/h} \mathcal{F}_h P_h, \quad \overline{\mathcal{F}}_h := P_h \overline{\mathcal{F}}_h Q_{1/h}.$$

Here $\mathbf{B}(L^2(\mathbb{R}^d))$ denotes the Banach space of all bounded linear operators in $L^2(\mathbb{R}^d)$, equipped with the uniform operator topology. Note that \mathcal{F}_h is a partial isometry from $L^2(\mathbb{R}^d)$ to $L^2(\mathbb{R}^d)$ with the initial set $L^2(\mathbb{Z}_h^d)$ and the final set $L^2(\mathbb{T}_{1/h}^d)$, and that $\overline{\mathcal{F}}_h$ is a partial isometry from $L^2(\mathbb{R}^d)$ to $L^2(\mathbb{R}^d)$ with the initial set $L^2(\mathbb{T}_{1/h}^d)$ and the final set $L^2(\mathbb{Z}_h^d)$. Clearly,

$$(2.18) \quad \overline{\mathcal{F}}_h \mathcal{F}_h = P_h, \quad \mathcal{F}_h \overline{\mathcal{F}}_h = Q_{1/h},$$

and

$$(2.19) \quad P_h^\perp \overline{\mathcal{F}}_h = \mathbf{0}, \quad Q_{1/h}^\perp \mathcal{F}_h = \mathbf{0}.$$

3. CONVERGENCE OF DISCRETE FOURIER TRANSFORMS

As in the previous section, we will work in \mathbb{C} -valued functions on d dimensional Euclidean space \mathbb{R}^d .

For $\varphi \in \mathcal{S}(\mathbb{R}^d)$, the Schwartz space of rapidly decreasing functions, we define

$$(3.1) \quad \varphi_h(x) = \sum_{n \in \mathbb{Z}^d} \varphi(hn) \chi_{I_{n,h}}(x).$$

It is clear that $\varphi_h \in L^2(\mathbb{Z}_h^d)$ and $P_h \varphi_h = \varphi_h$. However, we emphasize that $P_h \varphi \neq \varphi_h$ in general. In fact, for each $h > 0$ one can easily choose a function $\varphi \in \mathcal{S}(\mathbb{R}^d)$ such that $(\varphi_h, \varphi - \varphi_h)_{L^2(\mathbb{R}^d)} \neq 0$.

Remark 3.1. *As was noted in (2.7), we have $\|\varphi_h\|_{L^2(\mathbb{R}^d)} = h^{d/2} \|\varphi_h\|_{\ell^2(\mathbb{Z}_h^d)}$.*

Lemma 3.1. *Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and $k \in \mathbb{N}$.*

(i) *There exists a constant $C_{\varphi k}$, depending only on φ and k , such that*

$$(3.2) \quad |\varphi_h(x) - \varphi(x)| \leq C_{\varphi k} \langle x \rangle^{-k} h \quad (x \in \mathbb{R}^d, 0 < h < \frac{1}{\sqrt{d}}),$$

where $\langle x \rangle = \sqrt{1 + |x|^2}$. In particular, φ_h is rapidly decreasing, i.e., $|\varphi_h(x)| \leq C_{\varphi k} \langle x \rangle^{-k}$ for any $k \in \mathbb{N}$.

(ii) *There exists a constant C_φ , depending only on φ , such that*

$$(3.3) \quad \|\varphi_h - \varphi\|_{L^2} \leq C_\varphi h \quad (0 < h < \frac{1}{\sqrt{d}}).$$

Proof. We prove statement (i); then statement (ii) follows as a straightforward consequence, taking $k > d/2$. Let $n \in \mathbb{Z}^d$. Then, for $x \in I_{n,h}$,

$$\varphi(hn) - \varphi(x) = (hn - x) \cdot \int_0^1 (\nabla \varphi)(t(hn - x) + x) dt.$$

Consequently,

$$\begin{aligned} & |\varphi(hn) - \varphi(x)| \langle x \rangle^k \\ & \leq |hn - x| \int_0^1 \frac{\langle x \rangle^k}{\langle t(hn - x) + x \rangle^k} |(\nabla \varphi)(t(hn - x) + x)| \langle t(hn - x) + x \rangle^k dt \\ & \leq \left(\sup_{y \in \mathbb{R}^d} |(\nabla \varphi)(y)| \langle y \rangle^k \right) |hn - x| \int_0^1 \frac{\langle x \rangle^k}{\langle t(hn - x) + x \rangle^k} dt. \end{aligned}$$

Now if $|x| \geq 2\sqrt{dh}$, then

$$|t(hn - x) + x| \geq |x| - |hn - x| \geq |x| - \sqrt{dh} \geq \frac{|x|}{2},$$

so

$$\langle t(hn - x) + x \rangle \geq \sqrt{1 + \frac{|x|^2}{4}} = \sqrt{1 + \frac{\langle x \rangle^2 - 1}{4}} \geq \frac{\langle x \rangle}{2};$$

if $|x| < 2\sqrt{dh}$, then (trivially) $\langle t(hn - x) + x \rangle \geq 1$ and $\langle x \rangle < \sqrt{1 + 4dh^2} < \sqrt{5}$. In either case,

$$\int_0^1 \frac{\langle x \rangle^k}{\langle t(hn - x) + x \rangle^k} dt \leq \sqrt{5}^k,$$

and, noting that $|hn - x| \leq \sqrt{dh}$, the inequality (3.2) follows. ■

The statement of Lemma 3.1 (i) has the following immediate consequence.

Corollary 3.1. *Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and $k \in \mathbb{N}$. There exists a constant $C_{\varphi k}$, depending only on φ and k , such that*

$$(3.4) \quad \sum_{n \in \mathbb{Z}^d} |\varphi(hn)| \chi_{I_{n,h}}(x) \leq |\varphi(x)| + C_{\varphi k} h \langle x \rangle^{-k} \quad (x \in \mathbb{R}^d)$$

for $0 < h < 1/\sqrt{d}$.

In what follows, we shall use the notation

$$(3.5) \quad \mathcal{S}_h^{\text{step}}(\mathbb{R}^d) = \left\{ \varphi_h = \sum_{n \in \mathbb{Z}^d} \varphi(hn) \chi_{I_{n,h}} \mid \varphi \in \mathcal{S}(\mathbb{R}^d) \right\},$$

$$(3.6) \quad \mathcal{S}_{0+}^{\text{step}}(\mathbb{R}^d) = \bigcup_{h>0} \mathcal{S}_h^{\text{step}}(\mathbb{R}^d).$$

Note that, unlike $\mathcal{S}_h^{\text{step}}(\mathbb{R}^d)$, the set of functions $\mathcal{S}_{0+}^{\text{step}}(\mathbb{R}^d)$ is not a vector space. As we shall see in Lemma 3.4 in subsection 3.1, each $\varphi_h \in \mathcal{S}_{0+}^{\text{step}}(\mathbb{R}^d)$ allows an explicit expression of its Fourier transform in a certain sense.

The following lemma is a direct consequence of Lemma 3.1(ii).

Lemma 3.2. $\mathcal{S}_{0+}^{\text{step}}(\mathbb{R}^d)$ is a dense subset of $L^2(\mathbb{R}^d)$.

We can now prove the strong convergence of the orthogonal projectors P_h and $Q_{1/h}$ to the identity.

Lemma 3.3. *For any $u \in L^2(\mathbb{R}^d)$, $\|P_h u - u\|_{L^2(\mathbb{R}^d)} \rightarrow 0$ and $\|Q_{1/h} u - u\|_{L^2(\mathbb{R}^d)} \rightarrow 0$ as $h \rightarrow 0$.*

Proof. Since $\|P_h\|_{\mathbf{B}(L^2(\mathbb{R}^d))} = 1$ for all $h > 0$ and $\mathcal{S}(\mathbb{R}^d)$ is dense in $L^2(\mathbb{R}^d)$, it is sufficient to prove that for $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $\|P_h\varphi - \varphi\|_{L^2(\mathbb{R}^d)} \rightarrow 0$ as $h \rightarrow 0$.

Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and, for each $h > 0$, let $\varphi_h \in L^2(\mathbb{Z}_h^d)$ be the step function defined in (3.1). Then $P_h\varphi_h = \varphi_h$, so

$$\begin{aligned} \|P_h\varphi - \varphi\|_{L^2(\mathbb{R}^d)} &\leq \|P_h\varphi - P_h\varphi_h\|_{L^2(\mathbb{R}^d)} + \|\varphi_h - \varphi\|_{L^2(\mathbb{R}^d)} \\ &\leq (\|P_h\|_{\mathbf{B}(L^2(\mathbb{R}^d))} + 1) \|\varphi_h - \varphi\|_{L^2(\mathbb{R}^d)} \rightarrow 0 \quad (h \rightarrow 0) \end{aligned}$$

by Lemma 3.1(ii).

Furthermore, for any $u \in L^2(\mathbb{R}^d)$

$$\|Q_{1/h}u - u\|_{L^2(\mathbb{R}^d)}^2 = \int_{\mathbb{R}^d \setminus \mathbb{T}_{1/h}^d} |u|^2 \rightarrow 0 \quad (h \rightarrow 0).$$

■

3.1. Convergence of \mathcal{F}_h and $\overline{\mathcal{F}}_h$. As usual, the Fourier transform \mathcal{F} on $L^2(\mathbb{R}^d)$ and its inverse $\overline{\mathcal{F}}$ arise by extension of the integral operators

$$(3.7) \quad [\mathcal{F}\varphi](\xi) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{-ix \cdot \xi} \varphi(x) dx \quad (\xi \in \mathbb{R}^d; \varphi \in \mathcal{S}(\mathbb{R}^d))$$

and

$$(3.8) \quad [\overline{\mathcal{F}}\varphi](x) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \varphi(\xi) d\xi \quad (x \in \mathbb{R}^d; \varphi \in \mathcal{S}(\mathbb{R}^d)),$$

respectively. We emphasize that we can compare \mathcal{F}_h with \mathcal{F} and $\overline{\mathcal{F}}_h$ with $\overline{\mathcal{F}}$ on $L^2(\mathbb{R}^d)$, as we have extended the discrete Fourier transform and its inverse to all of $L^2(\mathbb{R}^d)$ in (2.17).

Lemma 3.4. *Let $\varphi_h \in \mathcal{S}_{0+}^{\text{step}}(\mathbb{R}^d)$. Then*

$$[\mathcal{F}\varphi_h](\xi) = \left\{ (2\pi)^{-d/2} \sum_{n \in \mathbb{Z}^d} \varphi(hn) h^d e^{-ihn \cdot \xi} \right\} \prod_{j=1}^d a(h\xi_j) \quad (\xi \in \mathbb{R}^d),$$

where

$$a(\theta) = \frac{1 - e^{-i\theta}}{i\theta} \quad (\theta \in \mathbb{R}).$$

Proof. Let $n \in \mathbb{Z}^d$ and $h > 0$. A direct computation shows that

$$(3.9) \quad \int_{I_{n,h}} e^{-ix \cdot \xi} dx = \prod_{j=1}^d e^{-ihn_j \xi_j} \frac{e^{-ih\xi_j} - 1}{-i\xi_j} = h^d e^{-ihn \cdot \xi} \prod_{j=1}^d a(h\xi_j).$$

We then have

$$\begin{aligned} (3.10) \quad [\mathcal{F}\varphi_h](\xi) &= (2\pi)^{-d/2} \sum_{n \in \mathbb{Z}^d} \varphi(hn) \int_{I_{n,h}} e^{-ix \cdot \xi} dx \\ &= (2\pi)^{-d/2} \sum_{n \in \mathbb{Z}^d} \varphi(hn) h^d e^{-ihn \cdot \xi} \prod_{j=1}^d a(h\xi_j). \end{aligned}$$

This completes the proof. ■

An immediate consequence of (2.12) and Lemma 3.4 is the following

Corollary 3.2. *Let $\varphi_h \in \mathcal{S}_{0+}^{\text{step}}(\mathbb{R}^d)$. Then*

$$(3.11) \quad [\mathcal{F}_h \varphi_h - \mathcal{F} \varphi_h](\xi) = \left(1 - \prod_{j=1}^d a(h\xi_j)\right) [\mathcal{F}_h \varphi_h](\xi) \quad (\xi \in \mathbb{T}_{1/h}^d).$$

Remark 3.2. *Note that $a(\theta) \rightarrow 1$ as $\theta \rightarrow 0$, and that $|a(\theta)| \leq 1$ for all θ .*

Remark 3.3. *One can deduce that $\mathcal{F}_h \varphi$ converges locally in L^2 to $\mathcal{F} \varphi$ for any $\varphi \in L^2(\mathbb{R}^d)$ in the following manner.*

Since $\|\mathcal{F}_h\|_{\mathbf{B}(L^2(\mathbb{R}^d))} = 1$ for all $h > 0$, and since $\mathcal{S}(\mathbb{R}^d)$ is dense in $L^2(\mathbb{R}^d)$, it is sufficient to prove the local convergence in L^2 for $\varphi \in \mathcal{S}(\mathbb{R}^d)$.

Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and let φ_h be given by (3.1). Then, by (2.12) and (3.4),

$$(3.12) \quad \begin{aligned} |[\mathcal{F}_h \varphi_h](\xi)| &= (2\pi)^{-d/2} \left| \int_{\mathbb{R}^d} \sum_{n \in \mathbb{Z}^d} e^{-ihn \cdot \xi} \varphi(hn) \chi_{I_{n,h}}(x) dx \right| \\ &\leq (2\pi)^{-d/2} \int_{\mathbb{R}^d} \sum_{n \in \mathbb{Z}^d} |\varphi(hn)| \chi_{I_{n,h}}(x) dx \\ &\leq (2\pi)^{-d/2} \int_{\mathbb{R}^d} (|\varphi(x)| + \text{const}_\varphi \langle x \rangle^{-d-1}) dx =: C_\varphi < \infty. \end{aligned}$$

Taken together with Corollary 3.2, this estimate implies that

$$\|[\mathcal{F}_h - \mathcal{F}] \varphi_h\|_{L^2(K)} \leq C_\varphi \sup_{\xi \in K} \left| 1 - \prod_{j=1}^d a(h\xi_j) \right| \rightarrow 0 \quad \text{as } h \rightarrow 0$$

for any compact subset K of \mathbb{R}^d . The decomposition

$$(3.13) \quad [\mathcal{F}_h - \mathcal{F}] \varphi = \mathcal{F}_h(\varphi - \varphi_h) + [\mathcal{F}_h - \mathcal{F}] \varphi_h + \mathcal{F}(\varphi_h - \varphi),$$

together with Lemma 3.1 (ii), gives the local convergence in L^2 . However, the following lemma shows a stronger convergence of \mathcal{F}_h .

Lemma 3.5. *Let $\varphi \in L^2(\mathbb{R}^d)$ and $s > 0$. Then*

$$(3.14) \quad \|\langle \cdot \rangle^{-s} [\mathcal{F}_h - \mathcal{F}] \varphi\|_{L^2(\mathbb{R}^d)} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

Proof. It is sufficient to show (3.14) for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$, since

$$\|\langle \cdot \rangle^{-s} \mathcal{F}_h\|_{\mathbf{B}(L^2(\mathbb{R}^d))} \leq 1$$

for all $h > 0$.

Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and, for each $h > 0$, let φ_h be given by (3.1). We first write

$$(3.15) \quad \|\langle \cdot \rangle^{-s} [\mathcal{F}_h - \mathcal{F}] \varphi_h\|_{L^2(\mathbb{R}^d)}^2 = \int_{\mathbb{R}^d} |\langle \xi \rangle^{-s} ([\mathcal{F}_h - \mathcal{F}] \varphi_h)(\xi)|^2 d\xi = I_h + II_h,$$

where

$$(3.16) \quad \begin{aligned} I_h &= \int_{\mathbb{T}_{1/h}^d} |\langle \xi \rangle^{-s} ([\mathcal{F}_h - \mathcal{F}] \varphi_h)(\xi)|^2 d\xi, \\ II_h &= \int_{\mathbb{R}^d \setminus \mathbb{T}_{1/h}^d} |\langle \xi \rangle^{-s} [\mathcal{F} \varphi_h](\xi)|^2 d\xi. \end{aligned}$$

The term II_h is easier to deal with. In fact, we see that

$$(3.17) \quad II_h \leq 2 \left(\int_{\mathbb{R}^d \setminus \mathbb{T}_{1/h}^d} |\mathcal{F} \varphi(\xi)|^2 d\xi + \int_{\mathbb{R}^d \setminus \mathbb{T}_{1/h}^d} |\mathcal{F}(\varphi - \varphi_h)(\xi)|^2 d\xi \right).$$

Here the second term is bounded by $\|\mathcal{F}(\varphi - \varphi_h)\|_{L^2(\mathbb{R}^d)}^2$, which tends to 0 as $h \rightarrow 0$ by (3.3). Hence $I_h \rightarrow 0$ as $h \rightarrow 0$.

To estimate the term I_h , let $\varepsilon \in (0, \pi)$ and divide $\mathbb{T}_{1/h}^d$ into two subsets,

$$(3.18) \quad \mathbb{T}_{1/h}^d = A_{\varepsilon, h} + B_{\varepsilon, h}$$

$$(3.19) \quad A_{\varepsilon, h} = \left\{ \xi \in \mathbb{T}_{1/h}^d \mid |\xi|_\infty < \frac{\varepsilon}{h} \right\}, \quad B_{\varepsilon, h} = \left\{ \xi \in \mathbb{T}_{1/h}^d \mid |\xi|_\infty \geq \frac{\varepsilon}{h} \right\}.$$

It follows from (3.16), (3.11) and (2.12) that

$$(3.20) \quad \begin{aligned} I_h &= h^{2d} (2\pi)^{-d} \int_{A_{\varepsilon, h} + B_{\varepsilon, h}} \langle \xi \rangle^{-2s} \left| 1 - \prod_{j=1}^d a(h\xi_j) \right|^2 \left| \sum_{n \in \mathbb{Z}^d} \varphi(hn) e^{-ihn \cdot \xi} \right|^2 d\xi \\ &\leq h^{2d} (2\pi)^{-d} \sup_{|\eta|_\infty < \varepsilon} \left\{ \left| 1 - \prod_{j=1}^d a(\eta_j) \right|^2 \right\} \int_{A_{\varepsilon, h}} \left| \sum_{n \in \mathbb{Z}^d} \varphi(hn) e^{-ihn \cdot \xi} \right|^2 d\xi \end{aligned}$$

$$(3.21) \quad + h^{2d} (2\pi)^{-d} (1 + \{\sup_{\theta \in \mathbb{R}} |a(\theta)|\}^d)^2 \varepsilon^{-2s} h^{2s} \int_{B_{\varepsilon, h}} \left| \sum_{n \in \mathbb{Z}^d} \varphi(hn) e^{-ihn \cdot \xi} \right|^2 d\xi$$

where we have used, in (3.21), the fact that $\langle \xi \rangle \geq \varepsilon/h$ for $\xi \in B_{\varepsilon, h}$. Both of the integrals in (3.20) and (3.21) are estimated by

$$\begin{aligned} \int_{\mathbb{T}_{1/h}^d} \left| \sum_{n \in \mathbb{Z}^d} \varphi(hn) e^{-ihn \cdot \xi} \right|^2 d\xi &= \left(\frac{2\pi}{h} \right)^d \sum_{n \in \mathbb{Z}^d} |\varphi(hn)|^2 = (2\pi)^d h^{-d} \|\varphi_h\|_{\ell^2(\mathbb{Z}_h^d)}^2 \\ &= (2\pi)^d h^{-2d} \|\varphi_h\|_{L^2(\mathbb{R}^d)}^2, \end{aligned}$$

using Parseval's identity for the orthonormal basis $\{e_n \mid n \in \mathbb{Z}^d\}$ of $L^2(\mathbb{T}_{1/h}^d)$ and Remark 3.1. Thus we obtain the estimate

$$(3.22) \quad \begin{aligned} I_h &\leq \left[\sup_{|\eta|_\infty < \varepsilon} \left\{ \left| 1 - \prod_{j=1}^d a(\eta_j) \right|^2 \right\} \right. \\ &\quad \left. + (1 + \{\sup_{\theta \in \mathbb{R}} |a(\theta)|\}^d)^2 \varepsilon^{-2s} h^{2s} \right] \|\varphi_h\|_{L^2(\mathbb{R}^d)}^2 \end{aligned}$$

and, by (3.3), arrive at

$$(3.23) \quad \limsup_{h \rightarrow 0} I_h \leq \left[\sup_{|\eta|_\infty < \varepsilon} \left\{ \left| 1 - \prod_{j=1}^d a(\eta_j) \right|^2 \right\} \right] \|\varphi\|_{L^2(\mathbb{R}^d)}^2.$$

Since $\varepsilon \in (0, \pi)$ is arbitrary and $a(\theta) \rightarrow 1$ as $\theta \rightarrow 0$, we can conclude that I_h converges to 0 as $h \rightarrow 0$. ■

Remark 3.4. Note that $\langle \cdot \rangle^{-s} \mathcal{F} = \mathcal{F} \langle D \rangle^{-s}$, where $D = -i\nabla$. However, one cannot expect in general that $\langle \Xi \rangle^{-s} \mathcal{F}_h = \mathcal{F}_h \langle D \rangle^{-s}$. The reason for this is illustrated by the following example. Let φ be the function given in Remark 2.1, which is absolutely continuous on \mathbb{R} . Hence the distributional derivative of φ coincides almost everywhere with its pointwise derivative $\varphi' = \chi_{I_{-2, h}} - \chi_{I_{1, h}}$. This expression of φ' implies that $[\mathcal{F}_h(-i\varphi')](\xi) = -ih(2\pi)^{-1/2}(e^{2ih\xi} - e^{-ih\xi})$. On the other hand, $[\mathcal{F}_h\varphi](\xi) = [\mathcal{F}_h P_h \varphi](\xi) = [\mathcal{F}_h f_h](\xi) = h^2(2\pi)^{-1/2}(e^{ih\xi} + 1)$. Therefore $\xi \mathcal{F}_h \varphi \neq \mathcal{F}_h(-i\varphi')$.

We close section 3 with the (fairly straightforward) proof of convergence of $\overline{\mathcal{F}}_h$.

Lemma 3.6. *For any $u \in L^2(\mathbb{R}^d)$, $\|\overline{\mathcal{F}}_h u - \overline{\mathcal{F}}u\|_{L^2(\mathbb{R}^d)} \rightarrow 0$ as $h \rightarrow 0$.*

Proof. Since $\|\overline{\mathcal{F}}_h\|_{\mathbf{B}(L^2(\mathbb{R}^d))} = 1$ for all $h > 0$, and since $C_0^\infty(\mathbb{R}^d)$ is dense in $L^2(\mathbb{R}^d)$, it is sufficient to prove the assertion for $u \in C_0^\infty(\mathbb{R}^d)$.

Let $u \in C_0^\infty(\mathbb{R}^d)$ and choose $h_* > 0$ so that $\mathbb{T}_{1/h}^d \supset \text{supp}[u]$ for all $h \in (0, h_*)$. Then we have, by (2.14) and by the fact that $\mathbb{T}_{1/h}^d \supset \text{supp}[u]$,

$$\begin{aligned} \overline{\mathcal{F}}_h u &= \sum_{n \in \mathbb{Z}^d} \left\{ (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{ihn \cdot \xi} u(\xi) d\xi \right\} \chi_{I_{n,h}} \\ (3.24) \quad &= \sum_{n \in \mathbb{Z}^d} \varphi(hn) \chi_{I_{n,h}} \end{aligned}$$

for all $h \in (0, h_*)$, where we set $\varphi := \overline{\mathcal{F}}u$. This equality implies that

$$(3.25) \quad |\overline{\mathcal{F}}_h u(x) - \overline{\mathcal{F}}u(x)|^2 = \sum_{n \in \mathbb{Z}^d} |\varphi(hn) - \varphi(x)|^2 \chi_{I_{n,h}}(x) \quad (x \in \mathbb{R}^d).$$

In view of the fact that $\varphi \in \mathcal{S}(\mathbb{R}^d)$, it follows from (3.25) and Lemma 3.1(i) that $\|\overline{\mathcal{F}}_h u - \overline{\mathcal{F}}u\|_{L^2(\mathbb{R}^d)} \rightarrow 0$ as $h \rightarrow 0$. ■

4. RESOLVENT CONVERGENCES OF $\mathbb{D}_{m,h}$

The continuum Dirac operator we shall consider in this section is

$$(4.1) \quad \mathbb{D}_m = -i\sigma_1 \frac{\partial}{\partial x_1} - i\sigma_2 \frac{\partial}{\partial x_2} + m\sigma_3 \quad \text{in } L^2(\mathbb{R}^2)^2,$$

where $m \geq 0$ and

$$(4.2) \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

It is well-known that \mathbb{D}_m is a self-adjoint operator in $L^2(\mathbb{R}^2)^2$ with domain $H^1(\mathbb{R}^2)^2$, the Sobolev space of order 1 of \mathbb{C}^2 -valued functions.

The discrete Dirac operator $\mathbb{D}_{m,h}$ we shall consider is

$$(4.3) \quad \mathbb{D}_{m,h} = \begin{pmatrix} m & i\partial_{1,h}^* + \partial_{2,h}^* \\ -i\partial_{1,h} + \partial_{2,h} & -m \end{pmatrix} \quad \text{in } L^2(\mathbb{Z}_h^2)^2,$$

where difference operators $\partial_{j,h}$ and $\partial_{j,h}^*$ ($j \in \{1, 2\}$) are as defined in (2.9). It is evident that $\mathbb{D}_{m,h}$ is a bounded self-adjoint operator in $L^2(\mathbb{Z}_h^2)^2$. We mention in passing that (4.3) is a Dirac operator with supersymmetry in the abstract sense (see [25, Chapter 5]). It can be rewritten in the form

$$\mathbb{D}_{m,h} = -i\sigma_1 \begin{pmatrix} \partial_{1,h} & 0 \\ 0 & -\partial_{1,h}^* \end{pmatrix} - i\sigma_2 \begin{pmatrix} \partial_{2,h} & 0 \\ 0 & -\partial_{2,h}^* \end{pmatrix} + m\sigma_3,$$

which is comparable with (1.2).

In accordance with the decomposition (2.8) of $L^2(\mathbb{R}^d)$, we can compare \mathbb{D}_m and $\mathbb{D}_{m,h} \oplus \mathbf{0}$ in the same Hilbert space $L^2(\mathbb{R}^2)^2$, and in particular investigate the difference

$$(4.4) \quad (\mathbb{D}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\mathbb{D}_m - z)^{-1} \quad \text{as } h \rightarrow 0.$$

We define $\widehat{\mathbb{D}}_m := \mathcal{F}\mathbb{D}_m\overline{\mathcal{F}}$, which is the operator of multiplication with the matrix-valued function

$$(4.5) \quad \widehat{\mathbb{D}}_m(\xi) = \begin{pmatrix} m & \xi_1 - i\xi_2 \\ \xi_1 + i\xi_2 & -m \end{pmatrix} \quad (\xi \in \mathbb{R}^2)$$

in $L^2(\mathbb{R}^2)^2$.

With help of (2.15), (2.16) and (4.3), we also define $\widehat{\mathbb{D}}_{m,h} := \mathcal{F}_h\mathbb{D}_{m,h}\overline{\mathcal{F}}_h$, where we abbreviate $\mathcal{F}_h := \mathcal{F}_h \otimes \mathbf{1}_{\mathbb{C}^2} \in \mathbf{B}(L^2(\mathbb{Z}_h^2)^2, L^2(\mathbb{T}_{1/h}^2)^2)$ and $\overline{\mathcal{F}}_h := \overline{\mathcal{F}}_h \otimes \mathbf{1}_{\mathbb{C}^2} \in \mathbf{B}(L^2(\mathbb{T}_{1/h}^2)^2, L^2(\mathbb{Z}_h^2)^2)$. The operator $\widehat{\mathbb{D}}_{m,h} \in \mathbf{B}(L^2(\mathbb{T}_{1/h}^2)^2)$ is the operator of multiplication with the matrix-valued function

$$(4.6) \quad \widehat{\mathbb{D}}_{m,h}(\xi) = \begin{pmatrix} m & \frac{i(e^{-ih\xi_1} - 1) + (e^{-ih\xi_2} - 1)}{h} \\ \frac{-i(e^{ih\xi_1} - 1) + (e^{ih\xi_2} - 1)}{h} & -m \end{pmatrix}$$

($\xi \in \mathbb{T}_{1/h}^2$). With the notation of (2.17), we see that

$$(4.7) \quad \mathbb{D}_{m,h} \oplus \mathbf{0} = \overline{\mathcal{F}}_h(\widehat{\mathbb{D}}_{m,h} \oplus \mathbf{0})\mathcal{F}_h \quad \text{in } L^2(\mathbb{R}^2)^2.$$

Therefore, the difference in (4.4) can be written as

$$(4.8) \quad \begin{aligned} & (\mathbb{D}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\mathbb{D}_m - z)^{-1} \\ &= \overline{\mathcal{F}}_h((\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0})\mathcal{F}_h - \overline{\mathcal{F}}(\widehat{\mathbb{D}}_m - z)^{-1}\mathcal{F}. \end{aligned}$$

To investigate the matrices $\widehat{\mathbb{D}}_m(\xi)$ and $\widehat{\mathbb{D}}_{m,h}(\xi)$, we start by noting the following.

Lemma 4.1. *Let $m \geq 0$ and $\zeta \in \mathbb{C}$, and assume that either $m > 0$ or $\zeta \neq 0$. Then, with the unitary matrix*

$$(4.9) \quad U_m(\zeta) = \frac{1}{\sqrt{2}\sqrt{\mu_m(\zeta)^2 + m\mu_m(\zeta)}} \begin{pmatrix} \mu_m(\zeta) + m & -\bar{\zeta} \\ \zeta & \mu_m(\zeta) + m \end{pmatrix},$$

where $\mu_m(\zeta) = \sqrt{|\zeta|^2 + m^2}$, we have

$$(4.10) \quad U_m(\zeta)^* \begin{pmatrix} m & \bar{\zeta} \\ \zeta & -m \end{pmatrix} U_m(\zeta) = \begin{pmatrix} \mu_m(\zeta) & 0 \\ 0 & -\mu_m(\zeta) \end{pmatrix}.$$

Note that the columns of the matrix on the right hand side of (4.9) are the eigenvectors of the matrix

$$(4.11) \quad \begin{pmatrix} m & \bar{\zeta} \\ \zeta & -m \end{pmatrix}.$$

Applying Lemma 4.1 to the matrix $\widehat{\mathbb{D}}_m(\xi)$ in (4.5) with $\zeta = \xi_1 + i\xi_2$, we see that the eigenvalues of the matrix $\widehat{\mathbb{D}}_m(\xi)$ are given by

$$(4.12) \quad \pm \lambda_m(\xi) := \pm \sqrt{|\xi|^2 + m^2},$$

and that the unitary transformation defined by $U_m(\xi_1 + i\xi_2)^*[\mathcal{F}\varphi](\xi)$ ($\varphi \in L^2(\mathbb{R}^2)^2$) brings the Dirac operator \mathbb{D}_m in (1.2) into the form of the operator of multiplication with the diagonal matrix-valued function

$$(4.13) \quad \begin{pmatrix} \lambda_m(\xi) & 0 \\ 0 & -\lambda_m(\xi) \end{pmatrix}$$

in $L^2(\mathbb{R}^2)^2$. This unitary transformation is a two-dimensional version of the Foldy-Wouthuysen-Tani transformation; see [9], [25, Section 1.4], [3, Section 2.1]. As is well-known, one can infer from (4.13) that the Dirac operator \mathbb{D}_m is absolutely continuous and that its spectrum is given by $\sigma(\mathbb{D}_m) = (-\infty, -m] \cup [m, \infty)$.

In a similar manner, applying Lemma 4.1 to the matrix $\widehat{\mathbb{D}}_{m,h}(\xi)$ in (4.6) with

$$(4.14) \quad \zeta = \frac{-i(e^{ih\xi_1} - 1) + (e^{ih\xi_2} - 1)}{h},$$

we see that the eigenvalues of the matrix $\widehat{\mathbb{D}}_{m,h}(\xi)$ are given by

$$(4.15) \quad \pm \lambda_{m,h}(\xi) := \pm \sqrt{h^{-2}\omega(h\xi) + m^2},$$

where

$$(4.16) \quad \begin{aligned} \omega(\xi) &= 4 + 2\sin(\xi_1 - \xi_2) - 2(\sin \xi_1 + \cos \xi_1) + 2(\sin \xi_2 - \cos \xi_2) \\ &= 2(1 - \cos \xi_1)(1 + \sin \xi_2) + 2(1 - \cos \xi_2)(1 - \sin \xi_1). \end{aligned}$$

Then the unitary transformation defined by $U_m(\zeta)^*[\mathcal{F}_h f](\xi)$ ($f \in L^2(\mathbb{Z}_h^2)^2$), with ζ as specified in 4.14, brings the discrete Dirac operator $\mathbb{D}_{m,h}$ in (4.3) into the form of the operator of multiplication with the diagonal matrix-valued function

$$(4.17) \quad \begin{pmatrix} \lambda_{m,h}(\xi) & 0 \\ 0 & -\lambda_{m,h}(\xi) \end{pmatrix}$$

in $L^2(\mathbb{T}_{1/h}^2)^2$.

Remark 4.1. *It is interesting that the Laplacian on the hexagonal lattice considered in [12] has a similar form to the discrete Dirac operator (4.3) in the massless case $m = 0$. As a result, a function similar to $\omega(\xi)$ appears in [12, subsection 8.1].*

Simple calculations show that for $\xi \in \mathbb{T}_1^2$

$$\nabla\omega(\xi) = 0 \iff$$

$$\xi \in \left\{ (0, 0), \left(\frac{\pi}{2}, -\frac{\pi}{2}\right), \left(-\frac{3\pi}{4}, \frac{3\pi}{4}\right), \left(\frac{\pi}{4}, -\frac{\pi}{4}\right), \left(\frac{\pi}{4}, \frac{3\pi}{4}\right), \left(-\frac{3\pi}{4}, -\frac{\pi}{4}\right) \right\},$$

and that the function $\omega : \mathbb{T}_1^2 \rightarrow \mathbb{R}$ has the following properties:

- (1) ω attains its minimum value 0 at $(0, 0)$ and at $\left(\frac{\pi}{2}, -\frac{\pi}{2}\right)$;
- (2) ω attains its unique maximum $6 + 4\sqrt{2}$ at $\left(-\frac{3\pi}{4}, \frac{3\pi}{4}\right)$;
- (3) The saddle points of ω are $\left(\frac{\pi}{4}, -\frac{\pi}{4}\right)$, $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$, $\left(-\frac{3\pi}{4}, -\frac{\pi}{4}\right)$.

Summing up, we have shown the following.

Theorem 4.1. *The discrete Dirac operator $\mathbb{D}_{m,h}$ is a bounded self-adjoint operator in $L^2(\mathbb{Z}_h^2)^2$ with purely absolutely continuous spectrum*

$$\sigma(\mathbb{D}_{m,h}) = \left[-\sqrt{h^{-2}(6 + 4\sqrt{2}) + m^2}, -m \right] \cup \left[m, \sqrt{h^{-2}(6 + 4\sqrt{2}) + m^2} \right].$$

Remark 4.2. *As was mentioned in the introduction, some spectral properties of 2D and 3D discrete Dirac operators with a fixed mesh size were already discussed in [20, Theorem 2.1], where the statement of Theorem 4.1 is shown in the special case $h = 1$.*

To prove the convergence theorems on $\mathbb{D}_{m,h}$, we need to examine the function ω in more detail. It follows from (4.16) that

$$(4.18) \quad 0 \leq \omega(\xi) \leq 2|\xi|^2 \quad (\xi \in \mathbb{R}^2)$$

and that

$$(4.19) \quad \omega(\xi) \geq \frac{2-\sqrt{2}}{8}|\xi|^2 \quad (|\xi|_\infty \leq \frac{\pi}{4})$$

(Obviously, the number $(2-\sqrt{2})/8$ can be replaced by any smaller positive constant, but we fix this value for the sake of definiteness.) Also, it follows from (4.16) that, setting $\alpha = (\frac{\pi}{2}, -\frac{\pi}{2})$,

$$(4.20) \quad \begin{aligned} \omega(\alpha + \xi) &= 4 - 2\sin(\xi_1 - \xi_2) + 2(\sin \xi_1 - \cos \xi_1) - 2(\sin \xi_2 + \cos \xi_2) \\ &= 2(1 - \cos \xi_2)(1 + \sin \xi_1) + 2(1 - \cos \xi_1)(1 - \sin \xi_2), \end{aligned}$$

which, together with (4.18), implies that

$$(4.21) \quad 0 \leq \omega(\alpha + \xi) \leq 2|\xi|^2 \quad (\xi \in \mathbb{R}^2)$$

and that

$$(4.22) \quad \omega(\alpha + \xi) \geq \frac{2-\sqrt{2}}{8}|\xi|^2 \quad (|\xi|_\infty \leq \frac{\pi}{4}).$$

Now, for any $\varepsilon \in (0, \pi^2/128)$ we divide \mathbb{T}_1^2 into two disjoint subsets,

$$(4.23) \quad \begin{aligned} \mathbb{T}_1^2 &= E(\varepsilon) \cup F(\varepsilon), \\ E(\varepsilon) &:= \{ \xi \in \mathbb{T}_1^2 \mid \omega(\xi) \geq \varepsilon \}, \quad F(\varepsilon) := \{ \xi \in \mathbb{T}_1^2 \mid \omega(\xi) < \varepsilon \}. \end{aligned}$$

In view of (4.18), (4.19), (4.21) and (4.22), one can see that $F(\varepsilon)$ consists of two disjoint components $F_0(\varepsilon)$ and $F_1(\varepsilon)$ satisfying

$$(4.24) \quad B\left(0, \sqrt{\frac{\varepsilon}{2}}\right) \subset F_0(\varepsilon) \subset B(0, 4\sqrt{\varepsilon}),$$

and

$$(4.25) \quad B\left(\alpha, \sqrt{\frac{\varepsilon}{2}}\right) \subset F_1(\varepsilon) \subset B(\alpha, 4\sqrt{\varepsilon}),$$

where $B(a, r) = \{ \xi \in \mathbb{R}^2 \mid |a - \xi| < r \}$ is the ball with center a and radius $r > 0$. Hence we have a disjoint decomposition of \mathbb{T}_1^2 :

$$(4.26) \quad \mathbb{T}_1^2 = E(\varepsilon) \cup F_0(\varepsilon) \cup F_1(\varepsilon).$$

Accordingly, we have a disjoint decomposition of $\mathbb{T}_{1/h}^2$:

$$(4.27) \quad \mathbb{T}_{1/h}^2 = E(\varepsilon, 1/h) \cup F_0(\varepsilon, 1/h) \cup F_1(\varepsilon, 1/h),$$

where

$$(4.28) \quad \begin{aligned} E(\varepsilon, 1/h) &:= \{ \xi \in \mathbb{T}_{1/h}^2 \mid h\xi \in E(\varepsilon) \}, \\ F_j(\varepsilon, 1/h) &:= \{ \xi \in \mathbb{T}_{1/h}^2 \mid h\xi \in F_j(\varepsilon) \} \quad (j = 0, 1). \end{aligned}$$

In the following proposition, we refer to the decomposition $L^2(\mathbb{R}^2)^2 = L^2(\mathbb{T}_{1/h}^2)^2 \oplus L^2(\mathbb{R}^2 \setminus \mathbb{T}_{1/h}^2)^2$; remember that $Q_{1/h}$, the operator of multiplication with $\chi_{\mathbb{T}_{1/h}^2}$, is the orthogonal projection onto the first direct summand.

Proposition 4.1. *Let $z \in \mathbb{C} \setminus \mathbb{R}$. Then*

- (i) $\lim_{h \rightarrow 0} \|[(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}]u\|_{L^2(\mathbb{R}^2)^2} = 0$ for any $u \in L^2(\mathbb{R}^2)^2$;
- (ii) $\lim_{h \rightarrow 0} \|(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}\|_{\mathbf{B}(L^{2,s}(\mathbb{R}^2)^2; L^2(\mathbb{R}^2)^2)} = 0$ for any $s > 0$, where $L^{2,s}(\mathbb{R}^2)^2$ denotes the weighted L^2 space defined by

$$(4.29) \quad L^{2,s}(\mathbb{R}^2)^2 = \left\{ u : \mathbb{R}^2 \rightarrow \mathbb{C}^2 \mid \|u\|_{L^{2,s}(\mathbb{R}^2)^2}^2 = \int_{\mathbb{R}^2} \langle \xi \rangle^{2s} |u(\xi)|_{\mathbb{C}^2}^2 d\xi < \infty \right\}$$

Lemma 4.2. *Let $h > 0$. Then we have*

$$(4.30) \quad \|\widehat{\mathbb{D}}_{m,h}(\xi) - \widehat{\mathbb{D}}_m(\xi)\|_{\mathbf{B}(\mathbb{C}^2)} \leq \frac{h}{2} |\xi|^2 \quad (\xi \in \mathbb{T}_{1/h}^2),$$

where $\|M\|_{\mathbf{B}(\mathbb{C}^2)}$ denotes the operator norm of a 2×2 matrix M as a linear operator in \mathbb{C}^2 .

Proof. By using the inequality

$$(4.31) \quad |e^{i\theta} - 1 - i\theta| \leq \frac{\theta^2}{2} \quad (\theta \in \mathbb{R}),$$

we see that

$$(4.32) \quad \left| \frac{-i(e^{ih\xi_1} - 1) + (e^{ih\xi_2} - 1)}{h} - (\xi_1 + i\xi_2) \right| \leq \frac{1}{h} \left\{ \frac{(h\xi_1)^2}{2} + \frac{(h\xi_2)^2}{2} \right\} = \frac{h}{2} |\xi|^2 \quad (\xi \in \mathbb{R}^2),$$

which, together with (4.5) and (4.6), implies the lemma. ■

Since $h|\xi| \leq \sqrt{2}\pi$ for $\xi \in \mathbb{T}_{1/h}^2$, the above lemma immediately gives the following estimate.

Lemma 4.3. *Let $h > 0$. Then we have*

$$(4.33) \quad \|\widehat{\mathbb{D}}_{m,h}(\xi) - \widehat{\mathbb{D}}_m(\xi)\|_{\mathbf{B}(\mathbb{C}^2)} \leq \frac{\sqrt{2}\pi}{2} |\xi| \quad (\xi \in \mathbb{T}_{1/h}^2).$$

Lemma 4.4. *For any $z \in \mathbb{C} \setminus \mathbb{R}$, there exists a constant C_z such that*

$$(4.34) \quad \|(\widehat{\mathbb{D}}_m(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)} \leq C_z (|\xi|^2 + m^2 + |\Im z|^2)^{-1/2} \quad (\xi \in \mathbb{R}^2).$$

Proof. We first note that the matrix $\widehat{\mathbb{D}}_m(\xi) - z$ is unitarily equivalent to the matrix

$$(4.35) \quad \begin{pmatrix} \lambda_m(\xi) - z & 0 \\ 0 & -\lambda_m(\xi) - z \end{pmatrix},$$

as was discussed after Lemma 4.1. Hence it is clear that

$$(4.36) \quad \|(\widehat{\mathbb{D}}_m(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)} = \max \left\{ \frac{1}{|\lambda_m(\xi) - z|}, \frac{1}{|-\lambda_m(\xi) - z|} \right\}.$$

Now let $z_0 = i \Im z$. Then

$$\frac{|z_0 - t|^2}{|z - t|^2} = \frac{(\Im z)^2 + t^2}{(\Im z)^2 + (t - \Re z)^2}$$

is a continuous function of $t \in \mathbb{R}$ that tends to 1 as $t \rightarrow \pm\infty$. It is therefore bounded, so there exists a constant C_z such that $|z_0 - t| \leq C_z |z - t|$ for all $t \in \mathbb{R}$, which together with (4.36) gives (4.34). ■

Lemma 4.5. *Let $z \in \mathbb{C} \setminus \mathbb{R}$, and let $\varepsilon > 0$. Then for any $h \in \left(0, \frac{\sqrt{\varepsilon}}{2|\Re z|}\right)$ we have*

$$(4.37) \quad \|(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)} \leq \min\left(\frac{1}{|\Im z|}, \frac{2h}{\sqrt{\varepsilon}}\right) \quad (\xi \in E(\varepsilon, 1/h)).$$

Proof. As a consequence of the spectral theorem for self-adjoint operators,

$$(4.38) \quad \|(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)} \leq \frac{1}{|\Im z|} \quad (\xi \in \mathbb{T}_{1/h}^2)$$

for any $z \in \mathbb{C} \setminus \mathbb{R}$. Therefore, it is sufficient to prove the inequality

$$(4.39) \quad \|(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)} \leq \frac{2h}{\sqrt{\varepsilon}} \quad (\xi \in E(\varepsilon, 1/h)).$$

To this end, we note the fact that the matrix $\widehat{\mathbb{D}}_{m,h}(\xi) - z$ is unitarily equivalent to the matrix

$$(4.40) \quad \begin{pmatrix} \lambda_{m,h}(\xi) - z & 0 \\ 0 & -\lambda_{m,h}(\xi) - z \end{pmatrix},$$

as was discussed after Lemma 4.1. By the reverse triangle inequality,

$$(4.41) \quad \begin{aligned} |\lambda_{m,h}(\xi) \pm z| &> \sqrt{h^{-2}\omega(h\xi) + m^2} - |\Re z| \\ &> h^{-1}\sqrt{\varepsilon} - |\Re z| \quad (\text{for } \xi \in E(\varepsilon, 1/h)). \end{aligned}$$

If h satisfies the inequality $0 < h < \sqrt{\varepsilon}/(2|\Re z|)$, then the inequality (4.39) follows from (4.41). ■

Proof of Proposition 4.1. (i) Let $u \in L^2(\mathbb{R}^2)^2$. Then

$$\begin{aligned} &\left\| \left((\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1} \right) u \right\|_{L^2(\mathbb{R}^2)^2}^2 \\ &= \int_{\mathbb{T}_{1/h}^2} \left| \left((\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1} - (\widehat{\mathbb{D}}_m - z)^{-1} \right) u(\xi) \right|_{\mathbb{C}^2}^2 d\xi \\ &\quad + \int_{\mathbb{R}^2 \setminus \mathbb{T}_{1/h}^2} \left| (\widehat{\mathbb{D}}_m(\xi) - z)^{-1} u(\xi) \right|_{\mathbb{C}^2}^2 d\xi. \end{aligned}$$

As the operator norm of $(\widehat{\mathbb{D}}_m(\xi) - z)^{-1}$ can be estimated by $|\Im z|^{-1}$, the second integral tends to 0 as $h \rightarrow 0$. Let $\varepsilon \in (0, \pi^2/128)$. In view of (4.27), we divide the first integral into three terms,

$$(4.42) \quad \begin{aligned} &\left(\int_{E(\varepsilon, 1/h)} + \int_{F_0(\varepsilon, 1/h)} + \int_{F_1(\varepsilon, 1/h)} \right) \left| \left((\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1} - (\widehat{\mathbb{D}}_m(\xi) - z)^{-1} \right) u(\xi) \right|_{\mathbb{C}^2}^2 d\xi \\ &=: I(\varepsilon, h) + II_0(\varepsilon, h) + II_1(\varepsilon, h). \end{aligned}$$

Since

$$(4.43) \quad \begin{aligned} &(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1} - (\widehat{\mathbb{D}}_m(\xi) - z)^{-1} \\ &= (\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1} (\widehat{\mathbb{D}}_m(\xi) - \widehat{\mathbb{D}}_{m,h}(\xi)) (\widehat{\mathbb{D}}_m(\xi) - z)^{-1}, \end{aligned}$$

we get

$$\begin{aligned}
(4.44) \quad I(\varepsilon, h) &\leq \int_{E(\varepsilon, 1/h)} \|(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)}^2 \\
&\quad \times \|\widehat{\mathbb{D}}_m(\xi) - \widehat{\mathbb{D}}_{m,h}(\xi)\|_{\mathbf{B}(\mathbb{C}^2)}^2 \|\widehat{\mathbb{D}}_m(\xi) - z\|_{\mathbf{B}(\mathbb{C}^2)}^{-1} |u(\xi)|_{\mathbb{C}^2}^2 d\xi \\
&\leq \int_{E(\varepsilon, 1/h)} \left\{ \min\left(\frac{1}{|\mathfrak{Jm} z|}, \frac{2h}{\sqrt{\varepsilon}}\right) \right\}^2 \\
&\quad \times \left(\frac{\sqrt{2}\pi}{2}|\xi|\right)^2 C_z^2 (|\xi|^2 + m^2 + |\mathfrak{Jm} z|^2)^{-1} |u(\xi)|_{\mathbb{C}^2}^2 d\xi \\
&\leq \left\{ \min\left(\frac{1}{|\mathfrak{Jm} z|}, \frac{2h}{\sqrt{\varepsilon}}\right) \right\}^2 \left(\frac{\sqrt{2}\pi}{2}\right)^2 C_z^2 \int_{\mathbb{R}^2} |u(\xi)|_{\mathbb{C}^2}^2 d\xi,
\end{aligned}$$

where we have used Lemmas 4.3, 4.4, 4.5. It follows from (4.44) that

$$(4.45) \quad \lim_{h \rightarrow 0} I(\varepsilon, h) = 0.$$

The second term in (4.42) can be estimated in a similar manner:

$$\begin{aligned}
(4.46) \quad II_0(\varepsilon, h) &\leq \int_{F_0(\varepsilon, 1/h)} \|(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)}^2 \\
&\quad \times \|\widehat{\mathbb{D}}_m(\xi) - \widehat{\mathbb{D}}_{m,h}(\xi)\|_{\mathbf{B}(\mathbb{C}^2)}^2 \|\widehat{\mathbb{D}}_m(\xi) - z\|_{\mathbf{B}(\mathbb{C}^2)}^{-1} |u(\xi)|_{\mathbb{C}^2}^2 d\xi \\
&\leq \int_{F_0(\varepsilon, 1/h)} \left(\frac{1}{|\mathfrak{Jm} z|}\right)^2 \\
&\quad \times \left(\frac{h}{2}|\xi|^2\right)^2 C_z^2 (|\xi|^2 + m^2 + |\mathfrak{Jm} z|^2)^{-1} |u(\xi)|_{\mathbb{C}^2}^2 d\xi,
\end{aligned}$$

where we have used (4.38), Lemmas 4.2 and 4.4. If $\xi \in F_0(\varepsilon, 1/h)$, then by (4.24) we see that $h|\xi| < 4\sqrt{\varepsilon}$, hence that

$$(4.47) \quad \left(\frac{h}{2}|\xi|^2\right)^2 C_z^2 (|\xi|^2 + m^2 + |\mathfrak{Jm} z|^2)^{-1} < 4\varepsilon C_z^2.$$

It follows from (4.45) and (4.47) that

$$(4.48) \quad \limsup_{h \rightarrow 0} II_0(\varepsilon, h) \leq \left(\frac{2C_z}{|\mathfrak{Jm} z|} \|u\|_{L_\xi^2}\right)^2 \varepsilon.$$

Also, the third term in (4.42) can be estimated in a similar manner:

$$\begin{aligned}
(4.49) \quad II_1(\varepsilon, h) &\leq \int_{F_1(\varepsilon, 1/h)} \|(\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1}\|_{\mathbf{B}(\mathbb{C}^2)}^2 \\
&\quad \times \|\widehat{\mathbb{D}}_m(\xi) - \widehat{\mathbb{D}}_{m,h}(\xi)\|_{\mathbf{B}(\mathbb{C}^2)}^2 \|\widehat{\mathbb{D}}_m(\xi) - z\|_{\mathbf{B}(\mathbb{C}^2)}^{-1} |u(\xi)|_{\mathbb{C}^2}^2 d\xi \\
&\leq \int_{F_1(\varepsilon, 1/h)} \left(\frac{1}{|\mathfrak{Jm} z|}\right)^2 \\
&\quad \times \left(\frac{\sqrt{2}\pi}{2}|\xi|\right)^2 C_z^2 (|\xi|^2 + m^2 + |\mathfrak{Jm} z|^2)^{-1} |u(\xi)|_{\mathbb{C}^2}^2 d\xi \\
&\leq \left(\frac{\sqrt{2}\pi C_z}{2|\mathfrak{Jm} z|}\right)^2 \int_{F_1(\varepsilon, 1/h)} |u(\xi)|_{\mathbb{C}^2}^2 d\xi
\end{aligned}$$

where we have used (4.38), Lemmas 4.3 and 4.4. Since, by (4.25),

$$(4.50) \quad F_1(\varepsilon, 1/h) \subset \{ \xi \in \mathbb{R}^2 \mid |\xi| \geq h^{-1}(|\alpha| - 4\sqrt{\varepsilon}) \},$$

we get

$$(4.51) \quad \lim_{h \rightarrow 0} II_1(\varepsilon, h) = 0.$$

We can deduce from (4.42), together with (4.45), (4.48) and (4.51), that

$$(4.52) \quad \limsup_{h \rightarrow 0} \int_{\mathbb{T}_{1/h}^2} | \{ (\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1} - (\widehat{\mathbb{D}}_m(\xi) - z)^{-1} \} u(\xi) |_{\mathbb{C}^2}^2 d\xi \leq \left(\frac{2C_z}{|\mathfrak{Jm} z|} \|u\|_{L_\xi^2} \right)^2 \varepsilon.$$

This completes the proof of conclusion (i) of Proposition 4.1, since $\varepsilon > 0$ was arbitrarily small.

(ii) Let $u \in L^{2,s}(\mathbb{R}^2)^2$ and $\varepsilon \in (0, \pi/128)$. Similarly to the proof of part (i), we start with the equality

$$(4.53) \quad \begin{aligned} & \|Q_{1/h} \{ (\widehat{\mathbb{D}}_{m,h} - z)^{-1} - (\widehat{\mathbb{D}}_m - z)^{-1} \} Q_{1/h} u \|_{L^2(\mathbb{R}^2)^2}^2 \\ &= \int_{\mathbb{T}_{1/h}^2} | \{ (\widehat{\mathbb{D}}_{m,h}(\xi) - z)^{-1} - (\widehat{\mathbb{D}}_m(\xi) - z)^{-1} \} u(\xi) |_{\mathbb{C}^2}^2 d\xi \\ &= I(\varepsilon, h) + II_0(\varepsilon, h) + II_1(\varepsilon, h). \end{aligned}$$

Since $s > 0$, we get, with the help of (4.44), (4.46) and (4.47), that

$$(4.54) \quad \begin{aligned} & I(\varepsilon, h) + II_0(\varepsilon, h) \\ & \leq \left[\left\{ \min \left(\frac{1}{|\mathfrak{Jm} z|}, \frac{2h}{\sqrt{\varepsilon}} \right) \right\}^2 \left(\frac{\sqrt{2}\pi}{2} \right)^2 + \left(\frac{2}{|\mathfrak{Jm} z|} \right)^2 \varepsilon \right] C_z^2 \times \|u\|_{L^{2,s}(\mathbb{R}^2)^2}^2 \end{aligned}$$

In a similar manner, we get, with the help of (4.49) and (4.50), that

$$(4.55) \quad II_1(\varepsilon, h) \leq \left(\frac{\sqrt{2}\pi C_z}{2|\mathfrak{Jm} z|} \right)^2 \left(\frac{h}{|\alpha| - 4\sqrt{\varepsilon}} \right)^{2s} \times \|u\|_{L^{2,s}(\mathbb{R}^2)^2}^2,$$

where we have used the inequality

$$(4.56) \quad \sup_{|\xi| \geq h^{-1}(|\alpha| - 4\sqrt{\varepsilon})} \langle \xi \rangle^{-2s} \leq \left(\frac{h}{|\alpha| - 4\sqrt{\varepsilon}} \right)^{2s}.$$

Combining (4.53), (4.54) and (4.55) leads to the inequality

$$(4.57) \quad \begin{aligned} & \|Q_{1/h} \{ (\widehat{\mathbb{D}}_{m,h} - z)^{-1} - (\widehat{\mathbb{D}}_m - z)^{-1} \} Q_{1/h} u \|_{\mathbf{B}(L^{2,s}(\mathbb{R}^2)^2; L^2(\mathbb{R}^2)^2)}^2 \\ & \leq \left[\left\{ \min \left(\frac{1}{|\mathfrak{Jm} z|}, \frac{2h}{\sqrt{\varepsilon}} \right) \right\}^2 \left(\frac{\sqrt{2}\pi}{2} \right)^2 + \left(\frac{2}{|\mathfrak{Jm} z|} \right)^2 \varepsilon \right] C_z^2 \\ & \quad + \left(\frac{\sqrt{2}\pi C_z}{2|\mathfrak{Jm} z|} \right)^2 \left(\frac{h}{|\alpha| - 4\sqrt{\varepsilon}} \right)^{2s}. \end{aligned}$$

It follows from (4.57) that

$$(4.58) \quad \begin{aligned} & \limsup_{h \rightarrow 0} \|Q_{1/h} \{ (\widehat{\mathbb{D}}_{m,h} - z)^{-1} - (\widehat{\mathbb{D}}_m - z)^{-1} \} Q_{1/h} u \|_{\mathbf{B}(L^{2,s}(\mathbb{R}^2)^2; L^2(\mathbb{R}^2)^2)}^2 \\ & \leq \frac{2C_z}{|\mathfrak{Jm} z|} \sqrt{\varepsilon}. \end{aligned}$$

This completes the proof of conclusion (ii) of Proposition 4.1, since $\varepsilon > 0$ was arbitrarily small. ■

Theorem 4.2. *Let $z \in \mathbb{C} \setminus \mathbb{R}$. Then*

$$(4.59) \quad \text{s-lim}_{h \rightarrow 0} \{(\mathbb{D}_{m,h} - z)^{-1} \oplus \mathbf{0}\} = (\mathbb{D}_m - z)^{-1} \text{ in } L^2(\mathbb{R}^2)^2.$$

Proof. Let $\varphi \in L^2(\mathbb{R}^2)^2$. It follows from (4.8) that

$$(4.60) \quad \begin{aligned} & \{(\mathbb{D}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\mathbb{D}_m - z)^{-1}\} \varphi \\ &= \overline{\mathcal{F}}_h \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0}\} \mathcal{F}_h \varphi - \overline{\mathcal{F}} \{(\widehat{\mathbb{D}}_m - z)^{-1}\} \mathcal{F} \varphi \\ &= \overline{\mathcal{F}}_h \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0}\} (\mathcal{F}_h - \mathcal{F}) \varphi \end{aligned}$$

$$(4.61) \quad + (\overline{\mathcal{F}}_h - \overline{\mathcal{F}}) \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0}\} \mathcal{F} \varphi$$

$$(4.62) \quad + \overline{\mathcal{F}} \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}\} \mathcal{F} \varphi.$$

We write the term in (4.60) as

$$(4.63) \quad \overline{\mathcal{F}}_h \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0}\} (\mathcal{F}_h - \mathcal{F}) \varphi$$

$$(4.64) \quad = \overline{\mathcal{F}}_h \{(\widehat{\mathbb{D}}_m - z)^{-1}\} (\mathcal{F}_h - \mathcal{F}) \varphi$$

$$(4.65) \quad + \overline{\mathcal{F}}_h \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}\} (\mathcal{F}_h - \mathcal{F}) \varphi.$$

The L^2 norm of the term in (4.64) is estimated by

$$(4.66) \quad \|(\widehat{\mathbb{D}}_m - z)^{-1} \langle \cdot \rangle^s\|_{\mathbf{B}(L^2(\mathbb{R}^2)^2)} \|\langle \cdot \rangle^{-s} (\mathcal{F}_h - \mathcal{F}) \varphi\|_{L^2(\mathbb{R}^2)^2},$$

which, by Lemma 3.5, tends to 0 as $h \rightarrow 0$ for any $s \in (0, 1]$. The L^2 norm of the term in (4.65) is bounded by

$$(4.67) \quad \|(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}\|_{\mathbf{B}(L^{2,s}(\mathbb{R}^2)^2; L^2(\mathbb{R}^2)^2)} \|\langle \cdot \rangle^{-s} (\mathcal{F}_h - \mathcal{F}) \varphi\|_{L^2(\mathbb{R}^2)^2},$$

which, by Proposition 4.1(ii) and Lemma 3.5, tends to 0 as $h \rightarrow 0$ for any $s > 0$. Hence the L^2 norm of the term in (4.60) tends to 0 as $h \rightarrow 0$ for any $s > 0$.

Similarly, we write the term in (4.61) as

$$(4.68) \quad (\overline{\mathcal{F}}_h - \overline{\mathcal{F}}) \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0}\} \mathcal{F} \varphi$$

$$(4.69) \quad = (\overline{\mathcal{F}}_h - \overline{\mathcal{F}}) \{(\widehat{\mathbb{D}}_m - z)^{-1} \oplus \mathbf{0}\} \mathcal{F} \varphi$$

$$(4.70) \quad + (\overline{\mathcal{F}}_h - \overline{\mathcal{F}}) \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}\} \mathcal{F} \varphi,$$

where, by Lemma 3.6, the L^2 norm of the term (4.69) tends to 0 as $h \rightarrow 0$, and the L^2 norm of the term (4.70) is bounded by

$$(4.71) \quad 2\| \{(\widehat{\mathbb{D}}_{m,h} - z)^{-1} \oplus \mathbf{0} - (\widehat{\mathbb{D}}_m - z)^{-1}\} \mathcal{F} \varphi \|_{L^2(\mathbb{R}^2)^2},$$

because $\|\overline{\mathcal{F}}_h\|_{\mathbf{B}(L^2(\mathbb{R}^2)^2, L^2(\mathbb{R}^2)^2)} = 1$. Proposition 4.1(i) implies that L^2 norm of the term in (4.71) tends to 0 as $h \rightarrow 0$. Therefore, the L^2 norm of the term in (4.61) tends to 0 as $h \rightarrow 0$ for any $s > 0$.

In the same manner as above, Proposition 4.1(i) implies that L^2 norm of the term in (4.62) tends to 0 as $h \rightarrow 0$. ■

5. STRONG RESOLVENT CONVERGENCE OF $\mathbb{D}_{m,h} + V_h$

In this section, we shall discuss the continuum Dirac operators

$$(5.1) \quad \mathbb{D}_m + V = -i\sigma_1 \frac{\partial}{\partial x_1} - i\sigma_2 \frac{\partial}{\partial x_2} + m\sigma_3 + V(x) \text{ in } L^2(\mathbb{R}^2)^2,$$

where V is a complex 2×2 matrix-valued potential. More precisely, we make the following

Assumption (V). $V : \mathbb{R}^2 \rightarrow \mathbb{C}^{2 \times 2}$ is a matrix-valued function each element of which is a bounded and uniformly continuous function.

We note that V can be decomposed into its Hermitian and skew-Hermitian parts, $V = V_{\mathfrak{R}} + iV_{\mathfrak{J}}$, where

$$(5.2) \quad V_{\mathfrak{R}}(x) = \frac{V(x) + V(x)^*}{2}, \quad iV_{\mathfrak{J}}(x) = \frac{V(x) - V(x)^*}{2} \quad (x \in \mathbb{R}^2).$$

It is evident that under the assumption (V), the operator V of multiplication with the matrix-valued function V is a bounded operator in $L^2(\mathbb{R}^2)^2$ and that the operator $\mathbb{D}_m + V$ is well-defined. In particular, if $V_{\mathfrak{J}} = 0$, then $\mathbb{D}_m + V$ is a self-adjoint operator in $L^2(\mathbb{R}^2)^2$ with domain $H^1(\mathbb{R}^2)^2$.

In analogy to (5.1), we consider the discrete Dirac operator $\mathbb{D}_{m,h} + V_h$ in $L^2(\mathbb{Z}_h^2)^2$, where $\mathbb{D}_{m,h}$ is the operator introduced in (4.3) and V_h is the operator of multiplication by

$$(5.3) \quad V_h(x) = \sum_{n \in \mathbb{Z}^2} V(hn) \chi_{I_{n,h}}(x) \quad (x \in \mathbb{R}^2).$$

It is clear that $\mathbb{D}_{m,h} + V_h$ is a bounded operator in $L^2(\mathbb{Z}_h^2)^2$. In the same manner as in (5.2), we split $V_h = V_{\mathfrak{R},h} + iV_{\mathfrak{J},h}$.

Theorem 5.1. *Under the assumption (V),*

(i) *both the spectra $\sigma(\mathbb{D}_{m,h} + V_h)$ and $\sigma(\mathbb{D}_m + V)$ are subsets of the strip*

$$\left\{ z \in \mathbb{C} \mid |\Im z| \leq \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)} \right\};$$

(ii) *for z with $|\Im z| > \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}$*

$$(5.4) \quad \text{s-lim}_{h \rightarrow 0} \left\{ (\mathbb{D}_{m,h} + V_h - z)^{-1} \oplus \mathbf{0} \right\} = (\mathbb{D}_m + V - z)^{-1} \text{ in } L^2(\mathbb{R}^2)^2.$$

Note that in the self-adjoint case $V_{\mathfrak{J}} = 0$, (5.4) holds for $z \in \mathbb{C} \setminus \mathbb{R}$.

We prepare the proof of Theorem 5.1 by providing the following auxiliary statements.

Lemma 5.1. *Suppose that (V) holds. Then*

$$\sigma(\mathbb{D}_m + V) \subset \left\{ z \in \mathbb{C} \mid |\Im z| \leq \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)} \right\}.$$

Moreover,

$$(5.5) \quad (\mathbb{D}_m + V - z)^{-1} = (\mathbb{D}_m + V_{\mathfrak{R}} - z)^{-1} \left\{ I_{L^2(\mathbb{R}^2)^2} + iV_{\mathfrak{J}} (\mathbb{D}_m + V_{\mathfrak{R}} - z)^{-1} \right\}^{-1}$$

for all z with $|\Im z| > \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}$.

Proof. Let $|\Im z| > \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}$. Since $\mathbb{D}_m + V_{\mathfrak{R}}$ is self-adjoint, it follows that $z \in \rho(\mathbb{D}_m + V_{\mathfrak{R}})$, the resolvent set of $\mathbb{D}_m + V_{\mathfrak{R}}$. This enables us to write

$$(5.6) \quad \mathbb{D}_m + V - z = \{I_{L^2(\mathbb{R}^2)} + iV_{\mathfrak{J}}(\mathbb{D}_m + V_{\mathfrak{R}} - z)^{-1}\}(\mathbb{D}_m + V_{\mathfrak{R}} - z).$$

Since $\|V_{\mathfrak{J}}\|_{\mathbf{B}(L^2(\mathbb{R}^2)^2)} \leq \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}$, we see that

$$\|V_{\mathfrak{J}}(\mathbb{D}_m + V_{\mathfrak{R}} - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{R}^2)^2)} < 1,$$

so the operator on the right hand side of (5.6) is invertible in $L^2(\mathbb{R}^2)^2$, and therefore $z \in \rho(\mathbb{D}_m + V)$. ■

In the same manner as in the proof of Lemma 5.1, one can prove a similar statement for $\mathbb{D}_{m,h} + V_h$. Recall that $\mathbb{D}_{m,h} + V_h$ is a bounded operator acting in $L^2(\mathbb{Z}_h^2)^2$.

Lemma 5.2. *Under assumption (V),*

$$\sigma(\mathbb{D}_{m,h} + V_h) \subset \{z \in \mathbb{C} \mid |\Im z| \leq \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}\}.$$

Moreover,

$$(5.7) \quad (\mathbb{D}_{m,h} + V_h - z)^{-1} = (\mathbb{D}_{m,h} + V_{\mathfrak{R},h} - z)^{-1} \{I_{L^2(\mathbb{Z}_h^2)^2} + iV_{\mathfrak{J},h}(\mathbb{D}_{m,h} + V_{\mathfrak{R},h} - z)^{-1}\}^{-1}$$

for all z with $|\Im z| > \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}$.

Lemma 5.3. *Under the assumption (V), $V_h \oplus \mathbf{0} \rightarrow V$ strongly in $L^2(\mathbb{R}^2)^2$.*

Proof. In this proof, we distinguish the multiplication operator V_h in $L^2(\mathbb{R}^2)^2$ from the embedded version of the multiplication operator V_h in $L^2(\mathbb{Z}_h^2)^2$, denoted by $V_h \oplus \mathbf{0}$.

Let $\varphi \in L^2(\mathbb{R}^2)^2$. In view of the fact that $\{V_h \oplus \mathbf{0}\}\varphi = V_h P_h \varphi$, we infer that

$$\begin{aligned} \|\{V_h \oplus \mathbf{0}\} - V\}\varphi\|_{L^2(\mathbb{R}^2)^2} &\leq \|V_h(P_h \varphi - \varphi)\|_{L^2(\mathbb{R}^2)^2} + \|(V_h - V)\varphi\|_{L^2(\mathbb{R}^2)^2} \\ &\leq \left\{ \sup_{x \in \mathbb{R}^2} \|V(x)\|_{\mathbf{B}(\mathbb{C}^2)} \right\} \|P_h \varphi - \varphi\|_{L^2(\mathbb{R}^2)^2} \\ &\quad + \sup_{n \in \mathbb{Z}^2} \left\{ \sup_{x \in I_{n,h}} \|V(x) - V(hn)\|_{\mathbf{B}(\mathbb{C}^2)} \right\} \|\varphi\|_{L^2(\mathbb{R}^2)^2}. \end{aligned}$$

By virtue of Lemma 3.3 and the assumption that the function V is bounded and uniformly continuous, it follows that $V_h \oplus \mathbf{0} \rightarrow V$ strongly in $L^2(\mathbb{R}^2)^2$ as $h \rightarrow 0$. ■

Applying the above lemma to V and to V^* and using (5.2), we obtain the following convergence results for the Hermitian and skew-Hermitian parts separately.

Corollary 5.1. *Suppose that (V) is verified. Then $V_{\mathfrak{R},h} \oplus \mathbf{0} \rightarrow V_{\mathfrak{R}}$ strongly in $L^2(\mathbb{R}^2)^2$ and $V_{\mathfrak{J},h} \oplus \mathbf{0} \rightarrow V_{\mathfrak{J}}$ strongly in $L^2(\mathbb{R}^2)^2$.*

Furthermore, we shall use the following two abstract lemmas. We omit the quite straightforward proof of the first of them.

Lemma 5.4. *Let \mathcal{H} be a Hilbert space. Let S_h and T_h belong to $\mathbf{B}(\mathcal{H})$ for each $h > 0$, and suppose that S_h and T_h strongly converge to S and $T \in \mathbf{B}(\mathcal{H})$ respectively as $h \rightarrow 0$. If $\sup_{h>0} \|S_h\|_{\mathbf{B}(\mathcal{H})} < \infty$, then $S_h T_h$ strongly converges to ST as $h \rightarrow 0$.*

Lemma 5.5. *Let \mathcal{H} be a Hilbert space. Suppose that \mathcal{H} has an orthogonal decomposition $\mathcal{H} = \mathcal{X}_h \oplus \mathcal{X}_h^\perp$ for each $h > 0$ and that the orthogonal projection P_h onto \mathcal{X}_h strongly converges to $I_{\mathcal{H}}$ as $h \rightarrow 0$. Let A_h , for each $h > 0$, and A be invertible operators in \mathcal{X}_h and in \mathcal{H} respectively such that $A_h \oplus 0$ strongly converges to A as $h \rightarrow 0$. If $\sup_{h>0} \|A_h^{-1}\|_{\mathbf{B}(\mathcal{X}_h)} < \infty$, then $A_h^{-1} \oplus 0$ strongly converges to A^{-1} as $h \rightarrow 0$.*

Proof. Let $\varphi \in \mathcal{H}$. By hypothesis, we see that

$$(A_h \oplus \mathbf{0})A^{-1}\varphi \rightarrow AA^{-1}\varphi = \varphi \text{ in } \mathcal{H}$$

as $h \rightarrow 0$. Hence we obtain

$$\begin{aligned} \|A^{-1}\varphi - (A_h^{-1} \oplus \mathbf{0})\varphi\|_{\mathcal{H}} \\ \leq \|(I_{\mathcal{H}} - P_h)A^{-1}\varphi\|_{\mathcal{H}} + \|(A_h^{-1} \oplus \mathbf{0})((A_h \oplus \mathbf{0})A^{-1}\varphi - \varphi)\|_{\mathcal{H}} \rightarrow 0 \quad (h \rightarrow 0) \end{aligned}$$

by the uniform boundedness of A_h^{-1} . ■

Proof of Theorem 5.1. Statement (i) was shown in Lemma 5.1 and Lemma 5.2. For (ii), let $z \in \mathbb{C}$, $|\Im z| > \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}\|_{\mathbf{B}(\mathbb{C}^2)}$. Then Lemmas 5.3, 5.4, and Theorem 4.2, together with the fact that

$$(5.8) \quad \|V_h \oplus \mathbf{0}\|_{\mathbf{B}(L^2(\mathbb{R}^2)^2)} \leq \sup_{x \in \mathbb{R}^2} \|V(x)\|_{\mathbf{B}(\mathbb{C}^2)} < \infty$$

imply that

$$(5.9) \quad \{V_h(\mathbb{D}_{m,h} - z)^{-1}\} \oplus \mathbf{0} = \{V_h \oplus \mathbf{0}\} \{(\mathbb{D}_{m,h} - z)^{-1} \oplus \mathbf{0}\} \rightarrow V(\mathbb{D}_m - z)^{-1}$$

strongly in $L^2(\mathbb{R}^2)^2$ as $h \rightarrow 0$, so also

$$(5.10) \quad \{I_{L^2(\mathbb{Z}_h^2)} + V_h(\mathbb{D}_{m,h} - z)^{-1}\} \oplus \mathbf{0} \rightarrow I_{L^2(\mathbb{R}^2)} + V(\mathbb{D}_m - z)^{-1}$$

strongly in $L^2(\mathbb{R}^2)^2$ as $h \rightarrow 0$.

Since z lies in the resolvent set of both $\mathbb{D}_{m,h}$ and $\mathbb{D}_{m,h} + V_h$ (see Lemma 5.2), we see that the right hand (and therefore the left hand) side of

$$(5.11) \quad I_{L^2(\mathbb{Z}_h^2)^2} + V_h(\mathbb{D}_{m,h} - z)^{-1} = (\mathbb{D}_{m,h} + V_h - z)(\mathbb{D}_{m,h} - z)^{-1}$$

is invertible in $L^2(\mathbb{Z}_h^2)^2$, so

$$(\mathbb{D}_{m,h} + V_h - z)^{-1} = (\mathbb{D}_{m,h} - z)^{-1} \{I_{L^2(\mathbb{Z}_h^2)^2} + V_h(\mathbb{D}_{m,h} - z)^{-1}\}^{-1}.$$

Now, in order to apply Lemma 5.5 with

$$A_h = I_{L^2(\mathbb{Z}_h^2)^2} + V_h(\mathbb{D}_{m,h} - z)^{-1} \text{ in } L^2(\mathbb{Z}_h^2)^2$$

and

$$A = I_{L^2(\mathbb{R}^2)^2} + V(\mathbb{D}_m - z)^{-1} \text{ in } L^2(\mathbb{R}^2)^2,$$

we require uniform boundedness of A_h^{-1} . To this end, we note that one can write

$$(5.12) \quad A_h^{-1} = I_{L^2(\mathbb{Z}_h^2)^2} - V_h(\mathbb{D}_{m,h} + V_h - z)^{-1},$$

and analogously

$$\left(I_{L^2(\mathbb{Z}_h^2)^2} + iV_{\mathfrak{J},h}(\mathbb{D}_{m,h} + V_{\mathfrak{A},h} - z)^{-1} \right)^{-1} = I_{L^2(\mathbb{Z}_h^2)^2} - iV_{\mathfrak{J},h}(\mathbb{D}_{m,h} + V_h - z)^{-1};$$

so we find, using (5.7),

$$\begin{aligned}
& \|(\mathbb{D}_{m,h} + V_h - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \\
&= \left\| (\mathbb{D}_{m,h} + V_{\mathfrak{R},h} - z)^{-1} \left(I_{L^2(\mathbb{Z}_h^2)^2} + iV_{\mathfrak{J},h}(\mathbb{D}_{m,h} + V_{\mathfrak{R},h} - z)^{-1} \right)^{-1} \right\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \\
&\leq \|(\mathbb{D}_{m,h} + V_{\mathfrak{R},h} - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \|I_{L^2(\mathbb{Z}_h^2)^2} - iV_{\mathfrak{J},h}(\mathbb{D}_{m,h} + V_h - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \\
&\leq \frac{1}{|\mathfrak{Im} z|} \left(1 + \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)} \|(\mathbb{D}_{m,h} + V_h - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \right),
\end{aligned}$$

giving

$$\|(\mathbb{D}_{m,h} + V_h - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \leq \frac{1}{|\mathfrak{Im} z| - \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}}$$

and further by (5.12)

$$\begin{aligned}
\|A_h^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} &\leq 1 + \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)} \|(\mathbb{D}_{m,h} + V_h - z)^{-1}\|_{\mathbf{B}(L^2(\mathbb{Z}_h^2)^2)} \\
&\leq \frac{|\mathfrak{Im} z|}{|\mathfrak{Im} z| - \sup_{x \in \mathbb{R}^2} \|V_{\mathfrak{J}}(x)\|_{\mathbf{B}(\mathbb{C}^2)}}
\end{aligned}$$

for all $h > 0$. Thus we can conclude, with the help of (5.10) and Lemma 5.5, that

$$(5.13) \quad \{I_{L^2(\mathbb{Z}_h^2)^2} + V_h(\mathbb{D}_{m,h} - z)^{-1}\}^{-1} \oplus \mathbf{0} \rightarrow \{I_{L^2(\mathbb{R}^2)^2} + V(\mathbb{D}_m - z)^{-1}\}^{-1}$$

strongly in $L^2(\mathbb{R}^2)^2$. Noting that

$$\|(\mathbb{D}_{m,h} - z)^{-1} \oplus \mathbf{0}\|_{L^2(\mathbb{R}^2)^2} \leq \frac{1}{|\mathfrak{Im} z|}$$

for all $h > 0$, we see from Lemma 5.4, Theorem 4.2 and (5.13) that

$$(\mathbb{D}_{m,h} + V_h - z)^{-1} \oplus \mathbf{0} \rightarrow (\mathbb{D}_m - z)^{-1} \{I_{L^2(\mathbb{R}^2)} + V(\mathbb{D}_m - z)^{-1}\}^{-1}$$

strongly in $L^2(\mathbb{R}^2)^2$. This completes the proof. ■

REFERENCES

- [1] R.I. Aleskerov, *An application of the inverse scattering problem for the discrete Dirac operator*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **46** (2020), no.1, 94-101.
- [2] E. Bairamov and S. Solmaz, *Spectrum and scattering function of the impulsive discrete Dirac systems*, Turkish J. Math. **42**(2018), no.6, 3182-3194.
- [3] M. Ben-Artzi and T. Umeda, *Spectral theory of first-order systems: From crystals to Dirac operators*, Rev. Math. Phys. **33** (2021), no.5, 2150014, 52pp.
- [4] O. Bourget, G. R. Moreno Flores and A. Taarabt, *One-dimensional discrete Dirac operators in a decaying random potential I: Spectrum and dynamics*, Math. Phys. Anal. Geom. **23**(2020), no. 2., Paper No.20, 51pp.
- [5] A.H. Castro Neto, F. Guinea, N.M.R. Peres, K.S. Novoselov and A.K. Geim, *The electron properties of graphene*, Rev. Mod. Phys. **81** (2009), no.1, 109-162.
- [6] B. Cassano, O.O. Ibrogimov, D. Krejčířík and F. Štampach, *Location of eigenvalues of non-selfadjoint discrete Dirac operators*, Ann. Henri Poincaré **21**(2020), no.7, 2193-2217.
- [7] N. Coskun and N. Yokus, *The spectrum of discrete Dirac operator with a general boundary condition*, Adv. Difference Equ. **2020** (2020), Paper No. 409, 9pp.
- [8] C. L. Fefferman, M. I. Weinstein, *Honeycomb lattice potentials and Dirac points*, J. Amer. Math. Soc. **25** (2012), no.4, 1169-1220.
- [9] L. L. Foldy and S. A. Wouthuysen, *On the Dirac theory of spin 1/2 particles and its non-relativistic limit*, Phys. Rev. **78** (1950), 29-35.

- [10] B. Fritzsche, B. Kirstein, I. Ya. Roitberg and A.L. Sakhnovich, *Discrete Dirac systems on the semiaxis: rational reflection coefficients and Weyl functions*, J. Difference Equ. Appl. **25** (2019), no.2, 294-304.
- [11] H. A. Hulko, *On the number of eigenvalues of the discrete one-dimensional Dirac operator with a complex potential*, Anal. Math. Phys. **9** (2019), no.1, 639-654.
- [12] H. Isozaki and A. Jensen, *Continuum limit for lattice Schrödinger operators*, arXiv:2006.00854v1(2020).
- [13] S. Klimek, M. McBride, S. Rathnayake and K. Sakai, *A value region problem for continued fractions and discrete Dirac equations*, Hokkaido Math. J. **49** (2020), no.2, 333-348.
- [14] T. Koprubasi, *The cubic eigenparameter dependent discrete Dirac equations with principal functions*, Commun. Fac. Sci. Univ. Ank. Ser.A1, Math. Stat. **68** (2019), no.2, 1742-1760.
- [15] E. Kopylova and G. Teschl, *Scattering properties and dispersion estimates for a one-dimensional discrete Dirac equation*, arXiv:2001.08445v1(2020).
- [16] S. Nakamura and Y. Tadano, *On a continuum limit of discrete Schrödinger operators on square lattice*, J. Spectr. Theory **11**(2021) no.1, 355-367.
- [17] K.S. Novoselov, A.K. Geim, S.V. Morozov, D. Jiang, M.I. Katsnelson, I.V. Grigorieva, S.V. Dubonos and A.A. Firsov, *Two-dimensional gas of massless Dirac fermions in graphene*, Nature **438** (2005), no.10, 197-200.
- [18] N.M.R. Peres, *Colloquium: The transport properties of graphene: An introduction*, Rev. Mod. Phys. **82** (2010), no.3, 2673-2700.
- [19] R.A. Prado and R.C. Charão, *Invariants of the trace map and uniform spectral properties for discrete Strumian Dirac operators*, Osaka J. Math. **56** (2019), no.2, 391-416.
- [20] R.A. Prado, C.R. de Oliveira and S.L. Carvalho, *Dynamical localization for discrete Anderson Dirac operators*, J. Stat. Phys. **167** (2017), 260-296.
- [21] M. Reed and B. Simon, *Methods of Modern Mathematical Physics I*, Functional Analysis, Revised and enlarged edition, Academic Press, Orlando, 1980.
- [22] I. Roitberg and A. Sakhnovich, *The discrete self-adjoint Dirac systems of general type: explicit solutions of direct and inverse problems, asymptotic of Verblunsky-type coefficients and the stability of solving of the inverse problem*, Zh. Mat. Fiz. Anal. Geom. **14** (2018), no.4, 532-548.
- [23] A.L. Sakhnovich, *GBDT of discrete skew-selfadjoint Dirac systems and explicit solutions of the corresponding non-stationary problems*, Oper. Theory Adv. Appl. **271**, Birkhäuser/Springer, Cham, 2018.
- [24] A.L. Sakhnovich, *Discrete Self-adjoint Dirac Systems: Asymptotic Relations, Weyl Functions and Toeplitz Matrices*, to appear in Constr. Approx. (2021).
- [25] B. Thaller, *The Dirac equation*, Springer-Verlag, Berlin, 1992.
- [26] C. Tretter, *Spectral theory of block operator matrices and applications*, Imperial College Press, London, 2008.

KARL MICHAEL SCHMIDT: SCHOOL OF MATHEMATICS, CARDIFF UNIVERSITY, SENGHENNYDD ROAD, CARDIFF CF24 4AG, WALES, UK
 Email address: schmidtkm@cardiff.ac.uk

TOMIO UMEDA: DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF HYOGO, HIMEJI 671-2201, JAPAN
 Email address: umeda@sci.u-hyogo.ac.jp