

# KEISLER'S THEOREM AND CARDINAL INVARIANTS

TATSUYA GOTO

ABSTRACT. We consider several variants of Keisler's isomorphism theorem. We separated these variants by showing implications between them and cardinal invariants hypotheses.

## 1. INTRODUCTION

The method of ultrapowers is one of the most important way to construct models. We consider the problem of when there exists an ultrafilter  $U$  on  $\omega$  such that for two models  $\mathcal{A}, \mathcal{B}$  in a countable language  $\mathcal{L}$ , the respective ultrapowers  $\mathcal{A}^\omega/U, \mathcal{B}^\omega/U$  are isomorphic. Since ultrapowers are elementarily extension of original models, if  $\mathcal{A}^\omega/U$  and  $\mathcal{B}^\omega/U$  are isomorphic, then  $\mathcal{A}$  and  $\mathcal{B}$  must be elementarily equivalent. Keisler showed, under CH, conversely if  $\mathcal{A}$  and  $\mathcal{B}$  are elementarily equivalent and of size  $\leq \mathfrak{c}$ , then for every ultrafilter  $U$  over  $\omega$ ,  $\mathcal{A}^\omega/U$  and  $\mathcal{B}^\omega/U$  are isomorphic. The purpose of this paper is to give necessary conditions and sufficient conditions for when the variants of Keisler's theorem hold in a model where CH does not hold, and to separate their variants using those conditions.

**Definition 1.1.** Let  $\kappa$  be a cardinal.

- (1) We say  $\text{KT}(\kappa)$  holds if for every countable language  $\mathcal{L}$  and  $\mathcal{L}$ -structures  $\mathcal{A}, \mathcal{B}$  of size  $\leq \kappa$  with  $\mathcal{A} \equiv \mathcal{B}$ , there exists an ultrafilter  $U$  over  $\omega$  such that  $\mathcal{A}^\omega/U \simeq \mathcal{B}^\omega/U$ .
- (2) We say  $\text{SAT}(\kappa)$  holds if there exists an ultrafilter  $U$  over  $\omega$  such that for every countable language  $\mathcal{L}$  and every sequence of  $\mathcal{L}$ -structures  $(\mathcal{A}_i)_{i \in \omega}$  with each  $\mathcal{A}_i$  of size  $\leq \kappa$ ,  $\prod_{i \in \omega} \mathcal{A}_i/U$  is saturated.

$\text{SAT}(\kappa)$  implies  $\text{KT}(\kappa)$  for every  $\kappa$ . Keisler [Kei61] proved  $\text{CH} \Rightarrow \text{SAT}(\mathfrak{c})$ . Recently, Golshani and Shelah [GS21] proved the converse  $\text{KT}(\mathfrak{c}) \Rightarrow \text{CH}$ . In fact, they showed  $\neg \text{KT}(\aleph_2)$ . They also proved that  $\text{cov}(\text{meager}) = \mathfrak{c} \wedge \text{cf}(\mathfrak{c}) = \aleph_1$  implies  $\text{KT}(\aleph_1)$ . Another classical result is the theorem by Ellentuck–Rucker [ER72] which shows that  $\text{MA}(\sigma\text{-centered})$  implies  $\text{SAT}(\aleph_0)$ . Moreover, Shelah [She92] showed  $\text{Con}(\neg \text{KT}(\aleph_0))$  by showing that  $\mathfrak{d} < \mathfrak{v}^\forall$  implies  $\neg \text{KT}(\aleph_0)$  and that the former is consistent.

In this paper, we prove the implications indicated by thick lines in Figure 1.

In the rest of this section, we recall basic notions related to cardinal invariants.

- Definition 1.2.**
- (1) If  $X, Y$  are sets and  $R$  is a subset of  $X \times Y$ , we call a triple  $(X, Y, R)$  a relational system.
  - (2) For a relational system  $\mathcal{A} = (X, Y, R)$ , define  $\mathcal{A}^\perp = (Y, X, \hat{R})$ , where  $\hat{R} = \{(y, x) \in Y \times X : \neg(x R y)\}$ .

---

*Date:* November 21, 2021.




FIGURE 1. Implications

- (3) For a relational system  $\mathcal{A} = (X, Y, R)$ , define  $\|\mathcal{A}\| = \min\{|B| : B \subseteq Y \wedge (\forall x \in X)(\exists y \in B)(x R y)\}$ .
- (4) For relational systems  $\mathcal{A} = (X, Y, R), \mathcal{B} = (X', Y', S)$ , we call a pair  $(\varphi, \psi)$  a Galois–Tukey morphism from  $\mathcal{A}$  to  $\mathcal{B}$  if  $\varphi: X \rightarrow X', \psi: Y' \rightarrow Y$  and  $(\forall x \in X)(\forall y \in Y')(\varphi(x) S y \implies x R \psi(y))$ .

**Fact 1.3** ([Bla10, Theorem 4.9]). If there is a Galois–Tukey morphism  $(\varphi, \psi)$  from  $\mathcal{A}$  to  $\mathcal{B}$ , then  $\|\mathcal{A}\| \leq \|\mathcal{B}\|$  and  $\|\mathcal{B}^\perp\| \leq \|\mathcal{A}^\perp\|$ .

- Definition 1.4.**
- (1) For  $f, g \in \omega^\omega$ , define  $f <^* g$  iff  $(\forall^\infty n)(f(n) < g(n))$  and  $f \leq^\infty g$  iff  $\neg(g <^* f)$  that is  $(\exists^\infty n)(f(n) \leq g(n))$ .
  - (2) For  $c \in (\omega+1)^\omega, h \in \omega^\omega$ , define  $\prod c = \prod_{n \in \omega} c(n)$  and  $S(c, h) = \prod_{n \in \omega} [c(n)]^{\leq h(n)}$ .
  - (3) For  $x \in \prod c$  and  $\varphi \in S(c, h)$ , define  $x \in^* \varphi$  iff  $(\forall^\infty n)(x(n) \in \varphi(n))$  and define  $x \in^\infty \varphi$  iff  $(\exists^\infty n)(x(n) \in \varphi(n))$ .

- Definition 1.5.**
- (1) Define  $\mathbf{D} = (\omega^\omega, \omega^\omega, <^*)$  and  $\mathfrak{d} = \|\mathbf{D}\|$ .
  - (2) Define  $\mathfrak{b} = \|\mathbf{D}^\perp\|$ .
  - (3) For a poset  $(P, <)$ , define a relational system  $\mathbf{Cof}(P, <)$  by  $\mathbf{Cof}(P, <) = (P, P, <)$ . Then we have  $\text{cf}(P, <) = \|\mathbf{Cof}(P, <)\|$ .
  - (4) For  $c \in (\omega+1)^\omega, h \in \omega^\omega$ , define  $\mathbf{Lc}(c, h) = (\prod c, S(c, h), \in^*)$ ,  $\mathfrak{c}_{c,h}^\forall = \|\mathbf{Lc}(c, h)\|$  and  $\mathfrak{v}_{c,h}^\forall = \|\mathbf{Lc}(c, h)\|^\perp$ .
  - (5) Define  $\mathbf{wLc}(c, h) = (\prod c, S(c, h), \in^\infty)$ ,  $\mathfrak{c}_{c,h}^\exists = \|\mathbf{wLc}(c, h)\|$  and  $\mathfrak{v}_{c,h}^\exists = \|\mathbf{wLc}(c, h)\|^\perp$ .
  - (6) For an ideal  $I$  on  $X$ , define  $\mathbf{Cov}(I) = (X, I, \in)$  and  $\text{cov}(I) = \|\mathbf{Cov}(I)\|$ .

- Definition 1.6.**
- (1) Define  $\mathfrak{v}^\forall = \min\{\mathfrak{v}_{c,h}^\forall : c, h \in \omega^\omega, \lim h = \infty\}$ .
  - (2) Define  $\mathfrak{c}^\exists = \min\{\mathfrak{c}_{c,h}^\exists : c, h \in \omega^\omega, \sum h/c < \infty\}$ .

**Fact 1.7** ([Gar16, Proposition 1.1.5]). Let  $\langle \mathcal{A}_i : i \in \omega \rangle$  be a sequence of structures in a language  $\mathcal{L}$  such that each  $\mathcal{A}_i$  has size  $\leq \mathfrak{c}$ . Let  $U$  be an ultrafilter over  $\omega$ . Then the ultraproduct  $\prod_{i \in \omega} \mathcal{A}_i / U$  has size either finite or  $\mathfrak{c}$ .

2. SAT( $\aleph_1$ ) AND KT( $\aleph_1$ )

**Theorem 2.1.** SAT( $\aleph_1$ ) implies CH.

*Proof.* Assume SAT( $\aleph_1$ ) and  $\neg$ CH. Take an ultrafilter  $U$  over  $\omega$  that witnesses SAT( $\aleph_1$ ). Let  $\mathcal{A}_* = (\omega_1, <)^{\omega}/U$ . For  $\alpha < \omega_1$ , put  $\alpha_* = [\langle \alpha, \alpha, \alpha, \dots \rangle]$ . Define a set  $p$  of formulas with a free variable  $x$  by

$$p = \{\alpha_* < x : \alpha < \omega_1\}.$$

This  $p$  is finitely satisfiable and the number of parameters occurring in  $p$  is  $\aleph_1 < \mathfrak{c} = |\mathcal{A}_*|$  by  $\neg$ CH. Thus, by SAT( $\aleph_1$ ), we can take  $f: \omega \rightarrow \omega_1$  such that  $[f]$  realizes  $p$ . Put  $\beta = \sup_{n \in \omega} f(n)$ . Now we have  $\{n \in \omega : \beta < f(n)\} \in U$  and this contradicts to the definition of  $\beta$ .  $\square$

**Definition 2.2.** Let  $\mathfrak{mcf} = \min\{\text{cf}(\omega^{\omega}/U) : U \text{ an ultrafilter over } \omega\}$ .

**Lemma 2.3** ([GS21, Claim 2.2]). Let  $\mathcal{A}$  be a structure in a language  $\mathcal{L} = \{<\}$ . Suppose that  $a \in \mathcal{A}$  has cofinality  $\omega_1$ . Let  $U$  be an ultrafilter over  $\omega$ . Then  $a_* = [\langle a, a, a, \dots \rangle]$  have cofinality  $\omega_1$  in  $\mathcal{A}^{\omega}/U$ .

*Proof.* Take an increasing cofinal sequence  $\langle x_{\alpha} : \alpha < \omega_1 \rangle$  of points in  $\mathcal{A}$  below  $a$ . Then  $\langle x_{\alpha}^* : \alpha < \omega_1 \rangle$  is an increasing cofinal sequence in  $\mathcal{A}_*$ , where  $x_{\alpha}^* = [\langle x_{\alpha}, x_{\alpha}, x_{\alpha}, \dots \rangle]$  for each  $\alpha < \omega_1$ . This can be showed by regularity of  $\omega_1$ .  $\square$

**Lemma 2.4** ([GS21, Claim 2.4]). Let  $U$  be an ultrafilter over  $\omega$  and  $\mathcal{B}_* = (\mathbb{Q}, <)^{\omega}/U$ . Then for every  $a, b \in \mathcal{B}_*$ , there is an automorphism on  $\mathcal{B}_*$  that sends  $a$  to  $b$ .

*Proof.* Consider a map  $F: \mathbb{Q}^3 \rightarrow \mathbb{Q}$  defined by  $F(x, y, z) = x - y + z$ . Then we have  $\forall (y, z \in \mathbb{Q})(\text{the map } x \mapsto F(x, y, z) \text{ is an automorphism on } (\mathbb{Q}, <) \text{ that sends } y \text{ to } z)$ .

This statement can be written by a first-order formula in the language  $\mathcal{L}' = \{<, F\}$ . Thus the same statement is true in  $(\mathbb{Q}, <, F)^{\omega}/U$ . The map  $F_*: \mathcal{B}_*^3 \rightarrow \mathcal{B}_*$  induced by  $F$  satisfies that

$\forall (y, z \in \mathcal{B}_*)(\text{the map } x \mapsto F(x, y, z) \text{ is an automorphism on } (\mathcal{B}_*, <) \text{ that sends } y \text{ to } z)$ .  $\square$

**Theorem 2.5.** KT( $\aleph_1$ ) implies  $\mathfrak{mcf} = \aleph_1$ .

*Proof.* This proof is based on [GS21, Theorem 2.1]. Assume that  $\mathfrak{mcf} \geq \aleph_2$ . We shall show  $\neg$ KT( $\aleph_1$ ).

Let  $\mathcal{L} = \{<\}$ ,  $\mathcal{A} = (\mathbb{Q}, <)$  and  $\mathcal{B} = (\mathbb{Q} + (\omega_1 + 1) \times \mathbb{Q}_{\geq 0}, <_{\mathcal{B}})$ . Here  $<_{\mathcal{B}}$  is defined by a lexicographical order and a disjoint union order.  $\mathcal{A}$  and  $\mathcal{B}$  are dense linear ordered sets, so by completeness of DLO, we have  $\mathcal{A} \equiv \mathcal{B}$ . Take an ultrafilter  $U$  over  $\omega$ . Put  $\mathcal{A}_* = \mathcal{A}^{\omega}/U$ ,  $\mathcal{B}_* = \mathcal{B}^{\omega}/U$ .

There is a point  $a$  in  $\mathcal{B}$  such that  $\text{cf}(\mathcal{B}_a) = \aleph_1$ , where  $\mathcal{B}_a = \{x \in \mathcal{B} : x < a\}$ . Then  $a_* \in \mathcal{B}_*$  has cofinality  $\aleph_1$  by Lemma 2.3. Here  $a_* = [\langle a, a, a, \dots \rangle]$ . On the other hand, we shall show every point in  $\mathcal{A}_*$  has cofinality  $\geq \mathfrak{mcf}$ . If we do this, since we assumed  $\mathfrak{mcf} \geq \aleph_2$ , we will have  $\mathcal{A}_* \not\equiv \mathcal{B}_*$ .

By Lemma 2.4, it suffices to consider the point  $0_* = [\langle 0, 0, 0, \dots \rangle]$ . Since  $\mathbb{Q}$  is vertically symmetrical, we consider  $\text{cf}((\mathbb{Q}_{>0})^{\omega}/U, >_U)$ .

Now we construct a Galois–Tukey morphism  $(\varphi, \psi) : \mathbf{Cof}(\omega^\omega/U) \rightarrow \mathbf{Cof}((\mathbb{Q}_{>0})^\omega/U, >_U)$  by

$$\begin{aligned}\varphi : \omega^\omega/U &\rightarrow (\mathbb{Q}_{>0})^\omega/U; [f] \mapsto [\langle 1/(f(n) + 1) : n \in \omega \rangle], \\ \psi : (\mathbb{Q}_{>0})^\omega/U &\rightarrow \omega^\omega/U; [g] \mapsto [\langle \lfloor 1/g(n) - 1 \rfloor : n \in \omega \rangle].\end{aligned}$$

So  $\text{cf}((\mathbb{Q}_{>0})^\omega/U, >_U) \geq \text{cf}(\omega^\omega/U, <_U)$ .

Thus we have  $\text{cf}((\mathbb{Q}_{>0})^\omega/U, >_U) \geq \mathbf{mcf}$ . We are done.  $\square$

**Corollary 2.6.**  $\text{KT}(\aleph_1)$  implies  $\mathfrak{b} = \aleph_1$ .

*Proof.* This follows from Theorem 2.5 and the fact that  $\mathfrak{b} \leq \mathbf{mcf}$ .  $\square$

**Remark 2.7.** The combination of Theorem 2.5 and [GS21, Theorem 3.2] gives the implication  $(\text{cov}(\text{meager}) = \mathfrak{c} \wedge \text{cf}(\mathfrak{c}) = \aleph_1) \implies \mathfrak{b} = \aleph_1$ . This is well-known. Because  $\text{cov}(\text{meager}) = \mathfrak{c}$  implies  $\mathfrak{d} = \mathfrak{c}$  and then  $\text{cf}(\mathfrak{d}) = \text{cf}(\mathfrak{c})$ . So by  $\text{cf}(\mathfrak{c}) = \aleph_1$ , we have  $\text{cf}(\mathfrak{d}) = \aleph_1$ , but we know  $\mathfrak{b} \leq \text{cf}(\mathfrak{d})$ . Thus it follows that  $\mathfrak{b} = \aleph_1$ .

### 3. $\text{SAT}(\aleph_0)$ AND $\text{KT}(\aleph_0)$

**Theorem 3.1.**  $\text{cov}(\text{meager}) = \mathfrak{c}$  implies  $\text{KT}(\aleph_0)$ .

*Proof.* By the same proof as [GS21, Theorem 3.2].  $\square$

**Corollary 3.2.**  $\text{Con}(\text{KT}(\aleph_0) \wedge \neg \text{KT}(\aleph_1))$ .

*Proof.*  $\text{MA} + \neg \text{CH}$  implies  $\text{KT}(\aleph_0) \wedge \neg \text{KT}(\aleph_1)$  by Theorem 2.5 and 3.1.  $\square$

**Fact 3.3** ([BJ95, Lemma 2.4.2]).  $\text{cov}(\text{meager}) = \mathfrak{v}_{\langle \omega : n \in \omega \rangle, \text{id}}^\exists$ . In other words,  $\text{cov}(\text{meager}) \geq \kappa$  iff  $(\forall X \subseteq \omega^\omega \text{ of size } < \kappa)(\exists S \in \prod_{i \in \omega} [\omega]^{< i})(\forall x \in X)(\exists^\infty n)(x(n) \in S(n))$ .

**Theorem 3.4.**  $\text{SAT}(\aleph_0)$  implies  $\text{cov}(\text{meager}) = \mathfrak{c}$ .

*Proof.* Take an ultrafilter  $U$  that witnesses  $\text{SAT}$ . Fix  $X \subseteq \omega^\omega$  of size  $< \mathfrak{c}$ . Define a language  $\mathcal{L}$  by  $\mathcal{L} = \{\subseteq\}$  and for each  $i \in \omega$ , define a  $\mathcal{L}$ -structure  $\mathcal{A}_i$  by  $\mathcal{A}_i = ([\omega]^{< i}, \subseteq)$ . For each  $x \in \omega^\omega$ , let  $S_x = \langle \{x(i) : i \in \omega\} \rangle$ . In the ultraproduct  $\mathcal{A}_* = \prod_{i \in \omega} \mathcal{A}_i / U$ , define a set  $p$  of formulas of one free variable  $S$  by

$$p = \{[S_x] \subseteq S : x \in X\}.$$

This  $p$  is finitely satisfiable. In order to check this, let  $x_0, x_1, \dots, x_n$  be finitely many members of  $X$ . Define  $S$  by  $S(m) = \{x_1(m), \dots, x_n(m)\}$  for  $m \geq n$ . We don't need to care about  $S(m)$  for  $m < n$ . Then this  $S$  satisfies  $[S_{x_i}] \subseteq [S]$  for all  $i \leq n$ . Moreover, the number of parameters of  $p$  is  $< \mathfrak{c}$ .

So by  $\text{SAT}(\aleph_0)$ , we can take  $[S] \in \mathcal{A}_*$  that realizes  $p$ . Then  $S$  fulfills  $(\forall x \in X)(\{n \in \omega : x(n) \in S(n)\} \in U)$ . Thus  $(\forall x \in X)(\exists^\infty n)(x(n) \in S(n))$ .  $\square$

**Theorem 3.5.**  $\text{SAT}(\aleph_0)$  implies  $2^{< \mathfrak{c}} = \mathfrak{c}$ .

*Proof.* Take an ultrafilter  $U$  over  $\omega$  that witnesses  $\text{SAT}(\aleph_0)$ . Fix  $\kappa < \mathfrak{c}$ .

Put  $\mathcal{L} = \{\subseteq\}$  and define an  $\mathcal{L}$ -structure  $\mathcal{A}$  by  $\mathcal{A} = ([\omega]^{< \omega}, \subseteq)$ . Put  $\mathcal{A}^* = \mathcal{A}^\omega / U$ .

Define a map  $\iota : \omega^\omega / U \rightarrow \mathcal{A}^*$  by  $\iota([x]) = [\langle \{x(n)\} : n \in \omega \rangle]$ . By Fact 1.7, we have  $|\omega^\omega / U| = \mathfrak{c}$ . Take a subset  $F$  of  $\omega^\omega / U$  of size  $\kappa$ .

For each  $X \subseteq F$ , let  $p_X$  be a set of formulas with a free variable  $z$  defined by

$$p_X = \{\iota(y) \subseteq z : y \in X\} \cup \{\iota(y) \not\subseteq z : y \in F \setminus X\}$$

Each  $p_X$  is finitely satisfiable. In order to check this, take  $[x_0], \dots, [x_n] \in X$  and  $[y_0], \dots, [y_m] \in F \setminus X$ . Put  $z(i) = \{x_0(i), \dots, x_n(i)\}$ . Then  $\iota([x_0]), \dots, \iota([x_n]) \subseteq_U [z]$ . In order to prove  $\iota([y_j]) \not\subseteq_U [z]$  for each  $j \leq m$ , suppose that  $\{i \in \omega : y_j(i) \in z(i)\} \in U$ . Then for each  $i \in \omega$ , there is a  $k_i \leq n$  such that  $\{i \in \omega : y_j(i) = x_{k_i}(i)\} \in U$ . Then there is a  $k \leq n$  such that  $\{i \in \omega : y_j(i) = x_k(i)\} \in U$ . This implies  $[y_j] = [x_k]$ , which is a contradiction.

By  $\text{SAT}(\aleph_0)$ , for each  $X \subseteq F$ , take  $[z_X] \in \mathcal{A}^*$  that realizes  $p_X$ . For  $X, Y \subseteq F$  with  $X \neq Y$ , we have  $[z_X] \neq [z_Y]$ . So  $2^{\aleph_0} = |\{[z_X] : X \subseteq F\}| \leq |\mathcal{A}^*| = \mathfrak{c}$ . Therefore we have proved  $2^{<\mathfrak{c}} = \mathfrak{c}$ .  $\square$

**Theorem 3.6.**  $\text{cov}(\text{meager}) = \mathfrak{c} \wedge 2^{<\mathfrak{c}} = \mathfrak{c}$  implies  $\text{SAT}(\aleph_0)$ .

*Proof.* This proof is based on [ER72, Theorem 1].

Let  $\langle b_\alpha : \alpha < \mathfrak{c} \rangle$  be an enumeration of  $\omega^\omega$ . Let  $\langle (\mathcal{L}_\xi, \mathcal{B}_\xi, \Delta_\xi) : \xi < \mathfrak{c} \rangle$  be an enumeration of triples  $(\mathcal{L}, \mathcal{B}, \Delta)$  such that  $\mathcal{L}$  is a countable language,  $\mathcal{B} = \langle \mathcal{A}_i : i \in \omega \rangle$  is a sequence of  $\mathcal{L}$ -structures with universe  $\omega$  and  $\Delta$  is a subset of  $\text{Fml}(\mathcal{L}^+)$  with  $|\Delta| < \mathfrak{c}$ . Here  $\mathcal{L}^+ = \mathcal{L} \cup \{c_\alpha : \alpha < \mathfrak{c}\}$  where the  $c_\alpha$ 's are new constant symbols and  $\text{Fml}(\mathcal{L}^+)$  is the set of all  $\mathcal{L}^+$  formulas with one free variable. Here we used the assumption  $2^{<\mathfrak{c}} = \mathfrak{c}$ . And ensure each  $(\mathcal{L}, \mathcal{B}, \Delta)$  occurs cofinally in this sequence.

For  $\mathcal{B}_\xi = \langle \mathcal{A}_i^\xi : i \in \omega \rangle$ , put  $\mathcal{B}_\xi(i) = (\mathcal{A}_i^\xi, b_0(i), b_1(i), \dots)$ , which is a  $\mathcal{L}^+$ -structure.

Let  $\langle X_\xi : \xi < \mathfrak{c} \rangle$  be an enumeration of  $\mathcal{P}(\omega)$ .

We construct a sequence  $\langle F_\xi : \xi < \mathfrak{c} \rangle$  of filters inductively so that the following properties hold:

- (1)  $F_0$  is the filter consisting of all cofinite subsets of  $\omega$ .
- (2)  $F_\xi \subseteq F_{\xi+1}$  and  $F_\xi = \bigcup_{\alpha < \xi} F_\alpha$  for  $\xi$  limit.
- (3)  $X_\xi \in F_{\xi+1}$  or  $\omega \setminus X_\xi \in F_{\xi+1}$ .
- (4)  $F_\xi$  is generated by  $< \mathfrak{c}$  members.
- (5) If
  - (\*) for all  $\Gamma \subseteq \Delta_\xi$  finite,  $\{i \in \omega : \Gamma \text{ is satisfiable in } \mathcal{B}_\xi(i)\} \in F_\xi$ ,
 then there is a  $f \in \omega^\omega$  such that for all  $\varphi \in \Delta_\xi$ ,  $\{i \in \omega : f(i) \text{ satisfies } \varphi \text{ in } \mathcal{B}_\xi(i)\} \in F_{\xi+1}$ .

Suppose we have constructed  $F_\xi$ . We construct  $F_{\xi+1}$ . Let  $F'_\xi$  be a generating subset of  $F_\xi$  with  $|F'_\xi| < \mathfrak{c}$ . If (\*) is false, let  $F_{\xi+1}$  be the filter generated by  $F'_\xi \cup \{X_\xi\}$  or  $F'_\xi \cup \{\omega \setminus X_\xi\}$ . Suppose (\*).

Put  $\mathbb{P} = \text{Fn}(\omega, \omega)$ . For  $n \in \omega$ , put

$$D_n = \{p \in \mathbb{P} : n \in \text{dom } p\}.$$

For  $A \in F'_\xi$  and  $\varphi_1, \dots, \varphi_n \in \Delta_\xi$ , put

$$E_{A, \varphi_1, \dots, \varphi_n} = \{p \in \mathbb{P} : (\exists i \in \text{dom } p \cap A)(p(i) \text{ satisfies } \varphi_1, \dots, \varphi_n \text{ in } \mathcal{B}_\xi(i))\}.$$

Each  $D_n$  is clearly dense. In order to show that each  $E_{A, \varphi_1, \dots, \varphi_n}$  is dense, take  $p \in \mathbb{P}$ . By (\*) and the property  $A \in F_\xi$ , we can take  $i \in A \setminus \text{dom } p$  and  $k \in \omega$  such that  $k$  satisfies  $\varphi_1, \dots, \varphi_n$  in  $\mathcal{B}_\xi(i)$ . Put  $q = p \cup \{(i, k)\}$ . This is an extension of  $p$  in  $E_{A, \varphi_1, \dots, \varphi_n}$ .

By using MA(Cohen), take a generic filter  $G \subseteq \mathbb{P}$  with respect to above dense sets. Put  $f = \bigcup G$ . Then  $F'_\xi := F'_\xi \cup \{Y_\varphi : \varphi \in \Delta_\xi\}$  satisfies finite intersection property, where  $Y_\varphi = \{i \in \omega : f(i) \text{ satisfies } \varphi \text{ in } \mathcal{B}_\xi(i)\}$ . In order to check this, let  $A \in F'_\xi$  and  $\varphi_1, \dots, \varphi_n \in \Delta_\xi$ . Then by genericity, we can take  $p \in G \cap E_{A, \varphi_1, \dots, \varphi_n}$ .

So we can take  $i \in \text{dom } p \cap A$  such that  $p(i)$  satisfies  $\varphi_1, \dots, \varphi_n$  in  $\mathcal{B}_\xi(i)$ . Then we have  $i \in A \cap Y_{\varphi_1} \cap \dots \cap Y_{\varphi_n}$ .

Let  $F_{\xi+1}$  be the filter generated by  $F_\xi'' \cup \{X_\xi\}$  or  $F_\xi'' \cup \{\omega \setminus X_\xi\}$ .

We have constructed  $\langle F_\xi : \xi < \mathfrak{c} \rangle$ . In order to check that the resulting ultrafilter  $F = \bigcup_{\xi < \mathfrak{c}} F_\xi$  witnesses  $\text{SAT}(\aleph_0)$ , let  $\mathcal{L}$  and  $\mathcal{B} = \langle \mathcal{A}_i : i \in \omega \rangle$  satisfy the assumption of the theorem. Let  $\Delta$  be a subset of  $\text{Fml}(\mathcal{L}^+)$  with  $|\Delta| < \mathfrak{c}$ . Assume that for all  $\Gamma \subseteq \Delta$  finite,  $X_\Gamma := \{i \in \omega : \Gamma \text{ is satisfiable in } \mathcal{B}_\xi(i)\} \in F$ . By the regularity of  $\mathfrak{c}$ , we have  $\alpha < \mathfrak{c}$  such that for all  $\Gamma \subseteq \Delta$  finite,  $X_\Gamma \in F_\alpha$ . Let  $\xi \geq \alpha$  be satisfying  $(\mathcal{L}_\xi, \mathcal{B}_\xi, \Delta_\xi) = (L, \mathcal{B}, \Delta)$ . Then by (5), there is a  $f \in \omega$  such that for all  $\varphi \in \Delta$ ,  $\{i \in \omega : f(i) \text{ satisfies } \varphi \text{ in } \mathcal{B}(i)\} \in F$ . Thus  $\prod_{i \in \omega} \mathcal{A}_i / F$  is saturated.  $\square$

#### 4. $\text{KT}(\aleph_0)$ IMPLIES $\mathfrak{c}^\exists \leq \mathfrak{d}$

In this section, we will show the following theorem. This proof is based on [She92] and [Abr10].

**Theorem 4.1.**  $\text{KT}(\aleph_0)$  implies  $\mathfrak{c}^\exists \leq \mathfrak{d}$ .

**Definition 4.2.** Define a language  $\mathcal{L}$  by  $\mathcal{L} = \{E, U, V\}$ , where  $E$  is a binary predicate and  $U, V$  are unary predicates. We say a  $\mathcal{L}$ -structure  $M = (|M|, E^M, U^M, V^M)$  is a **bipartite graph** if the following conditions hold:

- (1)  $U^M \cup V^M = |M|$ ,
- (2)  $U^M \cap V^M = \emptyset$ ,
- (3)  $(\forall x, y \in |M|)(x E^M y \rightarrow (x \in U^M \text{ and } y \in V^M))$ .

**Definition 4.3.** For  $n, k \in \omega$ , define a bipartite graph  $\Delta_{n,k}$  as follows:

- (1)  $U^{\Delta_{n,k}} = \{1, 2, 3, \dots, n\}$
- (2)  $V^{\Delta_{n,k}} = [\{1, 2, 3, \dots, n\}]^{\leq k} \setminus \{\emptyset\}$
- (3) For  $u \in U^{\Delta_{n,k}}, v \in V^{\Delta_{n,k}}$ ,  $u E^{\Delta_{n,k}} v$  iff  $u \in v$ .

**Definition 4.4.** For  $n \in \omega$ , Let  $G_n = \Delta_{n^3, n}$ . Let  $\Gamma$  be the disjoint union of  $(G_n : n \geq 2)$ .

We define a natural order  $\triangleleft$  on  $\Gamma$  by  $x \triangleleft y$  if  $m < n$  for  $x \in G_m, y \in G_n$ . Then  $\Gamma$  is a bipartite graph with an order  $\triangleleft$ . Put  $\mathcal{L}' = \mathcal{L} \cup \{\triangleleft\}$ . From now on, we consider  $\mathcal{L}'$ -structures which are elementarily equivalent to  $\Gamma$ .

**Definition 4.5.** Let  $\Gamma_{\text{NS}}$  be a countable non-standard elementary extension of  $\Gamma$ .

When we say connected components, we mean the connected components when we ignore the orientation of the edges.

**Lemma 4.6.** Let  $M$  be an  $\mathcal{L}'$ -structure that is elementarily equivalent to  $\Gamma$ . Then the connected components of  $M$  are precisely the maximal antichains of  $M$  with respect to  $\triangleleft$ .

*Proof.* Use elementarity and the fact that any two connected vertexes in  $\Gamma$  have path of length at most 4.  $\square$

**Lemma 4.7.** Every infinite connected component  $C$  of  $\Gamma_{\text{NS}}$  satisfies the following:

$$(\forall F \subseteq C \cap U \text{ finite})(\exists v \in C \cap V)(v \text{ has an edge to each point in } F).$$

*Proof.* Let  $F = \{u_1, \dots, u_n\}$ . Observe that

$$\begin{aligned} \Gamma \models (\forall x_1) \dots (\forall x_n) [x_1, \dots, x_n \text{ are points in } U \text{ and belong to} \\ \text{the same connected component and} \\ \text{the index of this connected component is } \geq n \\ \rightarrow (\exists y)[y \text{ belongs to this component, } y \in V \text{ and } x_1, \dots, x_n E y]]. \end{aligned}$$

By elementarity,  $\Gamma_{\text{NS}}$  satisfies the same formula.  $\square$

**Lemma 4.8.** Let  $\langle \Delta_n : n \in \omega \rangle$  be a sequence of bipartite graphs with  $|U^{\Delta_n}| = |V^{\Delta_n}| = \aleph_0$ . Suppose that for each  $n \in \omega$ ,

$$(\forall F \subseteq U^{\Delta_n} \text{ finite})(\exists v \in V^{\Delta_n})(v \text{ has an edge to each point in } F).$$

Then for every ultraproduct  $R := \prod_{n \in \omega} \Delta_n / q$ , we have

$$\langle \exists v_i : i < \mathfrak{d} \rangle \text{ with each } v_i \in V^R (\forall u \in U^R)(\exists i < \mathfrak{d})(u E^R v_i).$$

*Proof.* We may assume that each  $U^{\Delta_n} = \omega$ . Let  $\{f_i : i < \mathfrak{d}\}$  be a cofinal subset of  $(\omega^\omega, <^*)$ . For each  $n, m \in \omega$ , take  $v_{n,m} \in V^{\Delta_n}$  that is connected with first  $m$  points in  $U^{\Delta_n}$ . For  $i < \mathfrak{d}$ , put

$$v_i = \langle v_{n, f_i(n)} : n \in \omega \rangle.$$

Let  $[u] \in U^R$ . Consider  $u$  as an element of  $\omega^\omega$ . Take  $f_i$  that dominates  $u$ . Then we have

$$\{n \in \omega : u E^{\Delta_n} v_{n, f_i(n)}\} \in q.$$

Therefore  $[u] E^R v_i$ .  $\square$

**Lemma 4.9.** Let  $q$  an ultrafilter over  $\omega$  and put  $Q = (\Gamma_{\text{NS}})^\omega / q$ . Then there exist connected components  $C$  cofinally with respect to  $\triangleleft$  such that

$$\langle \exists v_i : i < \mathfrak{d} \rangle \text{ with each } v_i \in C \cap V^Q (\forall u \in C \cap U^Q)(\exists i < \mathfrak{d})(u E^Q v_i).$$

*Proof.* Fix connected component  $C_0$  of  $Q$  and  $[x_0] \in C_0$ . Then for each  $n \in \omega$ , there is an infinite component  $C_n$  above  $x_0(n)$ . Now

$$C = \{[x] \in Q : x \in (\Gamma_{\text{NS}})^\omega \text{ and } (\forall n \in \omega)(x(n) \in C_n)\}.$$

is a connected component of  $Q$  above  $C_0$ . Since  $C$  can be viewed as  $C = \prod_{n \in \omega} C_n / q$ , the conclusion of the lemma holds for  $C$  by Lemma 4.7 and Lemma 4.8.  $\square$

**Lemma 4.10.** Let  $\kappa < \mathfrak{c}^\exists$  and  $p$  an ultrafilter over  $\omega$  and put  $P = \Gamma^\omega / p$ . Then for every  $C$  in a final set of connected components of  $P$ , we have

$$\langle \forall v_i : i < \kappa \rangle \text{ with each } v_i \in C \cap V^P (\exists u \in C \cap U^P)(\forall i < \kappa)(u \not E^P v_i).$$

*Proof.* Let  $f : \omega \rightarrow \Gamma$  satisfy  $f(n) \in G_n$  for all  $n$ . Let  $C_0$  be the connected component that  $[f]$  belongs to. Take a connected component  $C$  such that  $C_0 \triangleleft C$  and an element  $g \in C$ . Take a function  $h : \omega \rightarrow \omega$  such that  $\{n \in \omega : g(n) \in G_{h(n)}\} \in q$ . Then  $A := \{n \in \omega : h(n) \geq n\} \in q$ . Put  $h'(n) = \max\{h(n), n\}$ .

Take  $\langle [v_i] : i < \kappa \rangle$  with each  $[v_i] \in C \cap V^P$ . Then

$$B_i := \{n \in \omega : v_i(n) \in G_{h(n)} \cap V^\Gamma\} \in q.$$

Take  $v'_i$  such that  $v'_i(n) = v_i(n)$  for  $n \in A_i$  and  $v'_i(n) \in [h'(n)^3]^{\leq h'(n)}$  for  $n \in \omega$ . The assumption  $\kappa < \mathfrak{c}^\exists$  and the calculation

$$\sum_{n \geq 1} \frac{h'(n)}{h'(n)^3} = \sum_{n \geq 1} \frac{1}{h'(n)^2} \leq \sum_{n \geq 1} \frac{1}{n^2} < \infty$$

give a  $x \in \prod h'$  such that for all  $i < \kappa$ ,  $(\forall^\infty n)(x(n) \notin v'_i(n))$ . For each  $i < \kappa$ , take  $n_i$  such that  $(\forall n \geq n_i)(x(n) \notin v'_i(n))$ .

Take a point  $[u] \in U^P$  such that  $u(n) = x(n)$  for all  $n \in A$ . Then for all  $i < \kappa$  we have

$$\{n \in \omega : u(n) \notin v_i(n)\} \supseteq A \cap B_i \cap [n_i, \omega) \in q.$$

Therefore  $[u] \notin^P [v_i]$  for all  $i < \kappa$ .  $\square$

Assume that  $\mathfrak{d} < \mathfrak{c}^\exists$ . Then by Lemma 4.10 and Lemma 4.9, for any two ultrafilters  $p, q$  over  $\omega$ , we have  $\Gamma^\omega/p \not\equiv (\Gamma_{\text{NS}})^\omega/q$ . So  $\neg \text{KT}(\aleph_0)$  holds. We have proved Theorem 4.1.

**Fact 4.11** ([KM19, Lemma 2.3]).  $\text{cov}(\text{null}) \leq \mathfrak{c}^\exists$ .

**Corollary 4.12.** In the random model,  $\neg \text{KT}(\aleph_0)$  holds.  $\square$

**Remark 4.13.**  $\mathfrak{v}^\forall \leq \mathfrak{c}^\exists$  follows from [KM19, Lemma 2.6]. So the implication  $\text{KT}(\aleph_0) \implies \mathfrak{d} \geq \mathfrak{c}^\exists$  strengthens the implication  $\text{KT}(\aleph_0) \implies \mathfrak{d} \geq \mathfrak{v}^\forall$ .

**Remark 4.14.** In [She92], Shelah constructed a creature forcing that forces the following statements:

- (1) There are a finite language  $\mathcal{L}$  and countable  $\mathcal{L}$ -structures  $\mathcal{A}, \mathcal{B}$  with  $\mathcal{A} \equiv \mathcal{B}$  such that for all ultrafilters  $p, q$  over  $\omega$ , we have  $\mathcal{A}^\omega/p \not\equiv \mathcal{B}^\omega/q$ .
- (2) There is an ultrafilter  $p$  over  $\omega$  such that for every countable language  $\mathcal{L}$  and any sequence  $\langle (\mathcal{A}_n, \mathcal{B}_n) : n \in \omega \rangle$  of pairs of finite  $\mathcal{L}$ -structures, if  $\prod_{n \in \omega} \mathcal{A}_n/p \equiv \prod_{n \in \omega} \mathcal{B}_n/p$ , then these ultraproducts are isomorphic.

Shelah himself pointed out in [She92, Remark 2.2] item 2 holds in the random model. On the other hand, we have proved item 1 also holds in the random model. Therefore both of above two statements hold in the random model.

## 5. OPEN QUESTIONS

- Question 5.1.**
- (1) Can  $\text{KT}(\aleph_0)$  and  $\text{cov}(\text{meager}) = \mathfrak{c}$  be separated?
  - (2) Does  $\text{KT}(\aleph_1)$  imply a stronger hypothesis than  $\mathfrak{mcf} = \aleph_1$ ? Especially does  $\text{KT}(\aleph_1)$  imply  $\text{non}(\text{meager}) = \aleph_1$ ?
  - (3) Does  $\text{KT}(\aleph_1)$  imply a hypothesis that some cardinal invariant is large? Especially does  $\text{KT}(\aleph_1)$  imply  $\text{cov}(\text{meager}) = \mathfrak{c}$ ?

## ACKNOWLEDGEMENT

The author is grateful to his current supervisor Yasuo Yoshinobu and his future supervisor Jörg Brendle. Both of them gave him helpful comments.



## REFERENCES

- [Abr10] Uri Abraham. “Proper forcing”. In: *Handbook of set theory*. Springer, 2010, pp. 333–394.
- [BJ95] Tomek Bartoszyński and Haim Judah. *Set Theory: on the structure of the real line*. CRC Press, 1995.
- [Bla10] Andreas Blass. “Combinatorial cardinal characteristics of the continuum”. In: *Handbook of set theory*. Springer, 2010, pp. 395–489.
- [ER72] Erik Ellentuck and R v B Rucker. “Martin’s Axiom and saturated models”. In: *Proceedings of the American Mathematical Society* 34.1 (1972), pp. 243–249.
- [Gar16] Darío García. *Model theory of pseudofinite structures*. 2016.
- [GS21] Mohammad Golshani and Saharon Shelah. *The Keisler-Shelah isomorphism theorem and the continuum hypothesis*. 2021. arXiv: 2108.03977 [math.LO].
- [Kei61] H Jerome Keisler. “Ultraproducts and elementary classes”. PhD thesis. University of California, Berkeley, 1961.
- [KM19] Lukas Daniel Klausner and Diego Alejandro Mejía. *Many Different Uniformity Numbers of Yorioka Ideals*. 2019. arXiv: 1805.11005 [math.LO].
- [She92] Saharon Shelah. “Vive la différence I: Nonisomorphism of ultrapowers of countable models”. In: *Set theory of the continuum*. Springer, 1992, pp. 357–405.

GRADUATE SCHOOL OF INFORMATION SCIENCE  
NAGOYA UNIVERSITY  
FURO-CHO, CHIKUSA-KU, NAGOYA 464-8601  
JAPAN

*Email address:* goto.tatsuya@k.mbox.nagoya-u.ac.jp