

SLOPE GAP DISTRIBUTION OF SADDLE CONNECTIONS ON THE $2n$ -GON

JONAH BERMAN, TAYLOR MCADAM, ANANTH MILLER-MURTHY, CAGLAR UYANIK, AND HAMILTON WAN

ABSTRACT. We explicitly compute the limiting slope gap distribution for saddle connections on any $2n$ -gon for $n \geq 3$. Our calculations show that the slope gap distribution for a translation surface is not always unimodal, answering a question of Athreya. We also give linear lower and upper bounds for number of non-differentiability points as n grows. The latter result exhibits the first example of a non-trivial bound on an infinite family of translation surfaces and answers a question by Kumanduri-Wang.

1. INTRODUCTION

Consider, for $n \geq 3$, a regular $2n$ -gon O_{2n} in the plane $\mathbb{R}^2 \cong \mathbb{C}$ where the opposite sides of O_{2n} are glued by Euclidean translations. The resulting surface X_{2n} is topologically a closed surface. If $n = 2k$, then X_{2n} is a surface of genus k and has a single cone point of angle $2\pi(n-1)$. If $n = 2k+1$, then X_{2n} is a surface of genus k , with two cone points where the cone angles are both $\pi(n-1)$ [14]. For example, gluing the opposite sides of the regular decagon yields a genus 2 surface with two cone points (marked with black and red), where at each of the cone points the total angle is 4π . See section 2.1 for details.




FIGURE 1. The regular decagon, O_{10} . Black vertices are identified together and red vertices are identified together.

Outside of the finitely many cone points, the surface O_{2n} is locally Euclidean and hence one can talk about straight lines. A *saddle connection* γ on O_{2n} is a straight line trajectory that connects a cone point to another cone point without passing through a cone point in the interior. To each saddle connection γ one can assign a holonomy vector v_γ which records how far γ travels in \mathbb{C} .

In this article we are interested in fine statistical properties of the set of holonomy vectors on X_{2n} . Veech proved that the number of saddle connections (rather, the set of holonomy vectors associated to saddle connections) grow quadratically with length, and saddle connection directions equidistribute on S^1 with respect to Lebesgue measure [18, 19], which suggests that saddle connection directions appear “randomly.” As a finer notion of randomness, one can study the *gaps* between saddle connections [1]. Since the set of saddle connections are symmetric with respect to the coordinate axes, we consider the set of holonomy vectors that lie in the first quadrant.

Let \mathbb{S}_{2n} denote the set of slopes of holonomy vectors v_γ with positive real component, and non-negative imaginary component. Write \mathbb{S}_{2n} as an increasing union of $\mathbb{S}_{2n}(k)$ for $k \rightarrow \infty$, where

$$\mathbb{S}_{2n}(k) = \{v_\gamma \mid 0 < \operatorname{Re}(v_\gamma) \leq k, \operatorname{Im}(v_\gamma) \geq 0\}$$

and write $\mathbb{S}_{2n}(k)$ as a set of increasing slopes

$$\mathbb{S}_{2n}(k) = \left\{ 0 = s_0^k < s_1^k < s_2^k < \dots \right\}.$$

We are interested in the distribution of gaps $s_i^k - s_{i-1}^k$ between consecutive slopes in $\mathbb{S}_{2n}(k)$ as $k \rightarrow \infty$. It can be shown that the gaps in $\mathbb{S}_{2n}(k)$ eventually repeat after an index $N(k)$ depending on k (for details, see Section 4). Moreover, Veech shows in [18] that $N(k)$ grows quadratically with k . Thus we define the set of renormalized slope gaps on S_{2n} as

$$\mathcal{G}_{2n}(k) = \left\{ k^2(s_i^k - s_{i-1}^k) \mid 1 \leq i \leq N(k), s_i^k \in \mathbb{S}_{2n}(k) \right\}.$$

We have the following theorem for the limiting distribution of these sets.

Theorem 1.1. *For any regular $2n$ -gon O_{2n} where $n \geq 3$, there exists a limiting probability distribution function $f : \mathbb{R} \rightarrow [0, \infty)$ such that*

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{G}_{2n}(k) \cap (a, b)|}{N(k)} = \int_a^b f(x) dx.$$

The distribution is a piecewise analytic function with finitely many domains of analyticity that are describable in terms of integrals of elementary functions. Moreover, the distribution has finitely many points of non-differentiability, has no support at 0, and has a quadratic tail.

Theorem 1.1 tells us that the slopes of the $2n$ -gon are not randomly distributed because if they were, we would expect the slope gap distribution to have small gaps and an exponential tail. For examples of the distributions for a variety of n , see Figure 11. One consequence of our calculations is the observation that the slope gap distribution of a translation surface is not always unimodal (that is, it can have multiple local maxima). To our knowledge, this is the first example of such a distribution, answering a question of Athreya.

Following the strategy in [3, 4] and [16], in order to prove this first theorem, we will translate our problem into a problem involving dynamics. In particular, we will exploit the slope gap preserving properties of the horocycle flow and study return times (under the horocycle flow) of translation surfaces to an appropriate Poincaré section.

The second theorem concerns a generalized description of the first return time function of the horocycle flow pertaining to the slope gaps of the regular unit $2n$ -gon.

Theorem 1.2. *There is a staircase surface in the $GL_2(\mathbb{R})$ orbit of O_{2n} for which a Poincaré section to the horocycle flow can be parametrized by two disjoint triangles. In these coordinates the return time function is a piecewise function which can be described by $n + 1$ elementary functions of the form*

$$R(x, y) = \frac{b}{x(ax + by)}$$

where a and b are constants with explicit formulas computed in this paper.

We use Theorem 1.2 to explicitly compute the slope gap distributions mentioned in Theorem 1.1. Furthermore, building on Theorems 1.1 and 1.2 we prove:

Theorem 1.3. *The number of non-differentiable points in the slope gap distribution for a regular $2n$ -gon has linear lower and upper bounds. In particular,*

$$\left\lfloor \frac{n}{2} \right\rfloor - 59 \leq \#(\text{Non-Differentiable Points}) \leq \left\lfloor \frac{n}{2} \right\rfloor + 3n - 6.$$

To the best of our knowledge this theorem gives the first infinite family of translation surfaces for which there is a non-trivial upper (and lower) bound on the number of non-differentiability points. Theorem 1.3 answers a question by [9]. Furthermore, in Section 5 we provide experimental evidence towards the precise asymptotics.

1.1. Some History on Slope Gap Distributions. There is a rich body of work on the statistics of gaps between saddle connections, combining aspects of the geometry of translation surfaces, homogeneous dynamics, and number theory.

Inspired by work of Eskin-Masur and Marklof-Strömbergsson [6, 10], Athreya-Chaika prove that a translation surface is not a lattice surface if and only if the smallest gap in directions between saddle connections of bounded length decays faster than quadratically in length [2]. Their results show that the limiting gap distribution exists and is the same for almost every surface, and that this distribution has a quadratic tail and has support at zero, in contrast to lattice surfaces which never have support at zero.

In [4], Athreya-Cheung use dynamical methods to rederive the Hall distribution for the limiting behavior of gaps between elements in the Farey sequence of level n (realized as slopes of saddle connections on the marked torus) and use their construction to obtain new fine statistical results in this setting. In [3], Athreya-Chaika-Lelièvre compute the slope gap distribution for saddle connections on the golden L, the first example of an explicit computation of a slope gap distribution for a surface which is not a branched cover of a torus. In [1], Athreya codifies the methods applied in the previous works into a series of meta-theorems that apply to gap sequences satisfying a set of quite general conditions, which in particular apply to studying the slopes of saddle connections on both lattice and non-lattice translation surfaces.

In [16], Uyanik-Work expand the list of surfaces for which the slope gap distribution has been explicitly calculated by finding the distribution for the regular octagon, which directly inspired the present work. They also provide an algorithm for finding the slope gap distribution for any lattice surface, and use this to show that the distribution of any such surface is piecewise real analytic. Their algorithm is improved upon by Kumanduri-Wang in [9] to further show that the distribution of a lattice surface always has a finite number of points of non-differentiability. In [15], Taha finds a Poincaré section for the geodesic and horocycle flows on quotients $SL_2(\mathbb{R})$ by Hecke triangle groups to study the statistics of generalized Farey sequences.

In [12], Sanchez provides the first explicit computation of a slope gap distribution for a non-lattice surface by finding the distribution for the class of surfaces known as double-slit tori. In a step toward computing the slope gap distribution for a generic surface (as described by [2]), Work finds a Poincaré section for the horocycle flow on $\mathcal{H}_1(2)$, the stratum of unit area translation surfaces with a single cone point of angle 6π , and provides bounds on the return time function to this section [20].

1.2. Acknowledgements. This work started as a project during 2020 SUMRY and the authors are grateful for the support from Yale University. The second named author is grateful for support from the NSF Postdoctoral Fellowship DMS-1903099. The authors thank Jayadev Athreya, Anthony Sanchez, Sunrose Shrestha, and Jane Wang for helpful conversations.

2. PRELIMINARIES

2.1. Translation Surfaces and the $GL_2(\mathbb{R})$ action. A translation surface is a pair (X, ω) where X is a Riemann surface, and ω is a holomorphic 1-form on X . Outside of a finite subset Σ of X , the 1-form ω induces an atlas of charts

$$\varphi_i : U_i \subset X \setminus \Sigma \rightarrow \mathbb{R}^2$$

where transition maps are Euclidean translations. The finite set of points in Σ are called the *singularities* and correspond to zeroes of the holomorphic one form [11]. At a zero of order $k - 1$ (which means that in local coordinates the chart looks like z^k and hence $\omega = d(z^k) = z^{k-1}dz$) the total angle is $2\pi k$.

Equivalently, a translation surface is given by a finite collection of polygons $\{P_1, \dots, P_n\}$ in the plane together with a preferred vertical direction and such that for each edge there exists a parallel edge of the same length, and these pairs are glued together by a translation. Since translations are holomorphic functions, and they preserve the standard 1-form dz on the plane, one gets a holomorphic 1-form on the glued together surface [11].

Two translation surfaces are *equivalent* if there is an orientation preserving isometry that preserves the preferred vertical direction. Equivalently, there is a *cut-translate-paste* transformation from one to another.

A translation surface (X, ω) comes equipped with topological data: the genus g , the number of cone points (number of zeroes of ω), and the excess angle at each cone point (order of zeroes). The Riemann–Roch theorem asserts that the sum of the order of zeroes is equal to $2g - 2$. Hence, we can record this topological data by a vector $\vec{\alpha} = (\alpha_1, \dots, \alpha_k)$ where α_i is the order of i^{th} zero. The set of equivalence classes of translation surfaces with fixed topological data $\vec{\alpha}$ is called a *stratum*, and denoted by $\mathcal{H}(\vec{\alpha})$.

A *saddle connection* γ on (X, ω) is a straight line trajectory that connects a cone point to another (not necessarily distinct) cone point without any other cone point in the interior.




FIGURE 2. A saddle connection γ on X_{10}

Integrating the 1-form along an (oriented) saddle connection γ determines a *holonomy vector*

$$v_\gamma = \int_\gamma \omega \in \mathbb{C}$$

which records how far and in what direction γ travels in \mathbb{C} . This paper is concerned with holonomy vectors corresponding to saddle connections on (X, ω) but for brevity we will call them also saddle connections following the convention in [3, 16] and denote the set of saddle connections on (X, ω) by $\Lambda_{sc}(X, \omega)$. For any translation surface (X, ω) the set $\Lambda_{sc}(X, \omega)$ is a discrete subset of \mathbb{C} , see [8, 21].

There is a natural action of $GL_2(\mathbb{R})$ on the space of translation surfaces coming from the linear action of $GL_2(\mathbb{R})$ on $\mathbb{R}^2 \cong \mathbb{C}$: for a matrix $A \in GL_2(\mathbb{R})$ and a translation surface (X, ω) given by a collection $\{P_1, \dots, P_n\}$ of polygons, $A \cdot (X, \omega)$ is the translation surface given by $\{A \cdot P_1, \dots, A \cdot P_n\}$. For $A \in SL_2(\mathbb{R})$, the action preserves the Euclidean area, hence induces an action on $\mathcal{H}_1(\vec{\alpha})$ of unit area translation surfaces of given topological type.

The stabilizer of (X, ω) under the $SL_2(\mathbb{R})$ action is called the *Veech group* of (X, ω) and denoted by $SL(X, \omega)$. If $SL(X, \omega)$ is a lattice, meaning that it has finite covolume in $SL_2(\mathbb{R})$ with respect to the Haar measure, then (X, ω) is called a *lattice surface* or a *Veech surface* [8].

2.2. Flat geometry of $2n$ -gons and Staircases. Let O_{2n} be the regular $2n$ -gon. In this section we prove that the matrix

$$M = \begin{pmatrix} 1 & -\cot(\pi/(2n)) \\ 0 & \sec\left(\frac{(n-2)\pi}{2n}\right) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & \sec\left(\frac{(n-2)\pi}{2n}\right) \end{pmatrix} \begin{pmatrix} 1 & -\cot(\pi/(2n)) \\ 0 & 1 \end{pmatrix}$$

takes O_{2n} to a staircase shape \mathcal{S}_{2n} that is easier to study. In other words, instead of studying the slope gap distribution on O_{2n} , we will study the slope gap distribution on \mathcal{S}_{2n} and relate it to that of O_{2n} . In what follows we will blur the distinction between the polygon \mathcal{S}_{2n} and the glued up surface, and denote both of them by \mathcal{S}_{2n} . Further, for the ease of notation we will denote \mathcal{S}_{2n} by \mathcal{S} .

For $0 \leq i \leq \lceil n/2 \rceil - 1$ and $0 \leq j \leq \lfloor n/2 \rfloor - 1$, we define the finite sequences h_i and v_j as follows:

$$h_i = 1 + \sum_{k=1}^i 2 \cos(k\pi/n) = \csc\left(\frac{\pi}{2n}\right) \sin\left(\frac{\pi(1+2i)}{2n}\right)$$

$$v_j = \cos\left(\frac{(n-2-2j)\pi}{2n}\right) \sec\left(\frac{(n-2)\pi}{2n}\right) = \csc\left(\frac{\pi}{n}\right) \sin\left(\frac{\pi(1+j)}{n}\right)$$

For $1 \leq i \leq \lceil n/2 \rceil - 1$, let H_i be the rectangle with height v_{i-1} and length h_i . Moreover, for $0 \leq j \leq \lfloor n/2 \rfloor - 1$, let V_j be the rectangle with height v_j and length h_j . Create the surface \mathcal{S} by gluing the left edge of H_1 to the right edge of the unit square (which is V_0), the top edge of H_i with the bottom edge of V_i , and the right edge of V_i with the left edge of H_{i+1} for all relevant i . Finally, identify opposite edges of the resulting staircase shape. This surface \mathcal{S} is a staircase shape whose edge lengths are in the sets of all h_i and v_j .




FIGURE 3. An example of \mathcal{S} for $n = 7$.

Proposition 2.1. *Left multiplication by the matrix $M = \begin{pmatrix} 1 & -\cot(\pi/(2n)) \\ 0 & \sec\left(\frac{(n-2)\pi}{2n}\right) \end{pmatrix}$ takes O_{2n} to \mathcal{S} .*

Proof. The surface O_{2n} has a cylinder decomposition in $\theta = \pi/2n$ direction. Hence by a cut and paste transformation while keeping track of the side identifications, we can obtain a *slanted* staircase representation of O_{2n} as below.

The matrix

$$\begin{pmatrix} 1 & -\cot(\pi/(2n)) \\ 0 & 1 \end{pmatrix}$$

is a shear that preserves the horizontal lines and distances and sends the lines with slope $\pi/2n$ to vertical lines. Hence we get a *staircase to the left*.

On the other hand, the matrix

$$\begin{pmatrix} 1 & 0 \\ 0 & \sec\left(\frac{(n-2)\pi}{2n}\right) \end{pmatrix}$$




FIGURE 4. Cylinder decomposition of O_{2n} in the $\theta = \pi/2n$ direction

only scales the vertical direction while keeping the x component constant, and it is only necessary to get the shortest vertical side to be of length 1. We can then apply a cut-and-paste transformation to orient our staircase from left to right. The claim about the edge lengths of the staircase follows from a straightforward computation.




FIGURE 5. A staircase representative of O_{12}

□

We finish this section by recording some geometric properties of \mathcal{S} that will be useful in the future. These result from the formulae for h_i and v_i and trigonometric identities.

Remark 2.2.

$$v_{i-1} + v_i = h_i$$

for $1 \leq i \leq \lfloor n/2 \rfloor - 1$. Equivalently, for all relevant i , the rectangle formed by $V_i \cup H_i$ is a square.

Remark 2.3.

$$2 + 2 \cos(\pi/n) = \frac{h_i + h_{i+1}}{v_i}$$

for $0 \leq i \leq \lfloor (n-1)/2 \rfloor - 1$. Equivalently, the rectangles formed by $V_i \cup H_{i+1}$ all have the same aspect ratio.

3. FROM SLOPE GAPS TO DYNAMICS

3.1. Dynamical Reframing. Let \mathcal{S} be the staircase representative of a regular $2n$ -gon. Recall from the introduction that

$$\mathbb{S}(k) = \{0 = s_0 < s_1 < s_2 < \dots\}$$

is the ordered set of slopes of holonomy vectors on \mathcal{S} with nonnegative imaginary component and positive real component $\leq k$. Recall that the gaps between consecutive elements of $\mathbb{S}(k)$ repeat with a period of $N(k)$, and that $N(k)$ grows like k^2 .

Let

$$\mathcal{G}(k) = \{k^2(s_i - s_{i-1}) \mid 1 \leq i \leq N(k), s_i \in \mathbb{S}(k)\}$$

denote the associated set of renormalized slope gaps on \mathcal{S} . Our goal is to compute the limit

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{G}(k) \cap (a, b)|}{N(k)}$$

for any interval $(a, b) \subseteq \mathbb{R}$. To compute this limit, we reframe the question about gaps between slopes in terms of return times for the horocycle flow to an appropriate Poincaré section.

In this article, the horocycle flow is given by the left action of the following one parameter subgroup:

$$\left\{ h_s = \begin{pmatrix} 1 & 0 \\ -s & 1 \end{pmatrix} \mid s \in \mathbb{R} \right\}.$$

Observe that given any vector v ,

$$\text{slope}(h_s v) = \text{slope}(v) - s,$$

so this definition of the horocycle flow acts on slopes by translation. Most importantly, the horocycle flow preserves *gaps* in slopes.

Let Γ denote the Veech group of \mathcal{S} . We will study the action of h_s on the space $\text{SL}_2(\mathbb{R})/\Gamma$, which we can identify with $\text{SL}_2(\mathbb{R}) \cdot \mathcal{S}$. If we denote the set of holonomy vectors on \mathcal{S} by $\Lambda_{sc}(\mathcal{S})$, then for any $g \in \text{GL}_2(\mathbb{R})$ we have

$$g \cdot \Lambda_{sc}(\mathcal{S}) = \Lambda_{sc}(g \cdot \mathcal{S}).$$

Consider the set

$$\Omega = \{g\Gamma \mid g \cdot \Lambda_{sc}(\mathcal{S}) \cap (0, 1] \neq \emptyset\}$$

in $\text{SL}_2(\mathbb{R})/\Gamma$, where the interval $(0, 1]$ is considered as a subset of the real axis inside of \mathbb{C} . In [1], Athreya proved that Ω is a Poincaré section to the horocycle flow h_s on $\text{SL}_2(\mathbb{R})/\Gamma$. Building on work of [3, 16] we parametrize this Poincaré section and relate the first return time function for the Poincaré section to the slope gap distribution of \mathcal{S} .

3.2. Description of the Poincaré Section. We now give an explicit description of the Poincaré section for the surface \mathcal{S} following the algorithm described in [16]. The parabolic and elliptic generators for $\Gamma_{O_{2n}}$, the Veech group of O_{2n} are respectively given by

$$S = \begin{pmatrix} 1 & -2 \cot(\pi/(2n)) \\ 0 & 1 \end{pmatrix}, \quad R = \begin{pmatrix} \cos(\pi/n) & \sin(\pi/n) \\ -\sin(\pi/n) & \cos(\pi/n) \end{pmatrix},$$

so the corresponding generators S' and R' for Γ , the Veech group of \mathcal{S} , are given by conjugating S and R by M :

$$S' = MSM^{-1} = \begin{pmatrix} 1 & -2(1 + \cos(\pi/n)) \\ 0 & 1 \end{pmatrix},$$

$$R' = MRM^{-1} = \begin{pmatrix} 1 + 2 \cos(\pi/n) & 2(1 + \cos(\pi/n)) \\ -1 & -1 \end{pmatrix}.$$

It has been shown by Veech, see [7], that the Veech group for O_{2n} (and hence \mathcal{S}) has two cusps. In fact, the image of $\Gamma_{O_{2n}}$ in $\mathrm{PSL}_2(\mathbb{R})$ is isomorphic to the triangle group $\Delta(n, \infty, \infty)$. Moreover, we know by [17] that given any Veech surface (X, ω) , the set $\Lambda_{sc}(X, \omega)$ of saddle connections is a finite disjoint union of $SL(X, \omega)$ orbits of some collection of saddle connections. Since our Veech group has two orbits, we will consider only the shorter orbits $\Gamma \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\Gamma \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

The decomposition of $\Lambda_{sc}(\mathcal{S})$ into disjoint orbits allows us to write

$$\Omega = \Omega_{w_1} \cup \Omega_{w_2},$$

where $w_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $w_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and $\Omega_{w_i} = \{g\Gamma \mid g\Lambda_{w_i} \cap (0, 1] \neq \emptyset\}$.

Let

$$M_{x,y} = \begin{pmatrix} x & y \\ 0 & x^{-1} \end{pmatrix},$$

$$\Omega_1 = \{(x, y) \in \mathbb{R}^2 \mid 0 < x \leq 1, 1 - 2(1 + \cos(\pi/n))x < y \leq 1\},$$

and

$$\Omega_2 = \{(x, y) \in \mathbb{R}^2 \mid 0 < x \leq 1, 1 - x < y \leq 1\}.$$

Proposition 3.1. *There are coordinates from the set Ω to $\Omega_1 \cup \Omega_2$. More precisely, the bijection between $\mathrm{SL}_2(\mathbb{R})/\Gamma$ and $\mathrm{SL}_2(\mathbb{R}) \cdot \mathcal{S}$ induces a bijection from Ω_{w_i} to $\{M_{x,y}C_i \cdot \mathcal{S} \mid (x, y) \in \Omega_i\}$ where the latter set is in bijection with Ω_i for a suitable matrix C_i .*

Proof. This statement follows from Theorem 1.2 in [16], but we illustrate the explicit computations below. The Veech group Γ has two cusps, and let S_1 and S_2 be the generators of the infinite cyclic factors corresponding to maximal parabolic subgroups. One can check that

$$S_1 = S' = \begin{pmatrix} 1 & -2(1 + \cos(\pi/n)) \\ 0 & 1 \end{pmatrix}$$

and

$$S_2 = ((R')^{n-1}S')^{-1} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}. \tag{1}$$

Note that S_1 and S_2 both have eigenvalue 1, and that their eigenvectors are in the direction of w_1 and w_2 , respectively. Now, pick $C_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, so that

$$C_1 S_1 C_1^{-1} = \begin{pmatrix} 1 & \alpha_1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -2(1 + \cos(\pi/n)) \\ 0 & 1 \end{pmatrix}$$

and $C_1 \cdot w_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. Thus, we can set

$$\Omega_1 = \{(x, y) \mid 0 < x \leq 1, 1 - 2(1 + \cos(\pi/n))x < y \leq 1\}.$$

Now, pick $C_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, so that

$$C_2 S_2 C_2^{-1} = \begin{pmatrix} 1 & \alpha_2 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

and $C_2 \cdot w_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. Thus, we can set

$$\Omega_2 = \{(x, y) \mid 0 < x \leq 1, 1 - x < y \leq 1\}.$$

□

3.3. **Return time and volume computations.** Recall that the Poincaré section Ω is defined as

$$\Omega = \{g\Gamma \mid g \cdot \Lambda_{sc}(\mathcal{S}) \cap (0, 1] \neq \emptyset\}.$$

Given a point X in Ω the return time function $R : \Omega \rightarrow \mathbb{R}^+$ gives the amount of time required for X to return to Ω under the horocycle flow h_s . Since h_s acts on slopes by translation, in the language of our parametrization, for any $(x, y) \in \Omega_i$, the return time $R(x, y)$ is the slope of the saddle connection (with horizontal component at most 1) on $M_{x,y}C_i \cdot \mathcal{S}$ with smallest positive slope among all saddle connections with horizontal component at most 1.

In this section, we prove that there are finitely many saddle connections of interest on $C_i \cdot \mathcal{S}$, i.e. for any given $(x, y) \in \Omega_i$, there is a finite set of *fixed* saddle connections on $C_i \mathcal{S}$ that are candidates for being the saddle connection on $M_{x,y}C_i \cdot \mathcal{S}$ of minimum positive slope.

Following the terminology introduced in [9], we say that a (holonomy) vector on $C_i \cdot \mathcal{S}$ is a *winner* or a *winning saddle connection* at the point $(x, y) \in \Omega_i$ if it is the vector whose image under $M_{x,y}$ has the smallest slope among all saddle connections with positive slope and horizontal length at most 1 on $M_{x,y}C_i \cdot \mathcal{S}$.

In Section 3.3.1, we show that the only winning saddle connection for Ω_2 is $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$. On the other hand, there are n saddle connections of interest for Ω_1 , given by $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and the diagonals of the rectangles H_i and V_i (see Figure 6). Each of these saddle connections wins at some point on Ω_1 and hence partition Ω_1 into n regions. We provide an explicit description of this partition in Section 3.3.2. These divisions and saddle connections therefore give us an explicit description of the return time function: if $(x, y) \in \Omega_i$ is arbitrary and $\begin{pmatrix} a \\ b \end{pmatrix}$ is the winning saddle connection on $M_{x,y}C_i \cdot \mathcal{S}$, then

$$R(x, y) = \frac{b}{x(ax + by)}.$$




FIGURE 6. The saddle connections of interest for Ω_1 (in magenta) on \mathcal{S} for $n = 7$.

3.3.1. *Description of $R(x, y)$ Over Ω_2 .* We describe $R(x, y)$ for $(x, y) \in \Omega_2$ by showing that $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ is the only winning vector on Ω_2 .

Proposition 3.2. *The vector $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ is the only winning saddle connection on Ω_2 .*




FIGURE 7. An illustration of Ω_2 with coordinates in \mathbb{R}^2 , for any n .

Proof. Since we will be studying Ω_2 , we will let $\mathcal{S}^R = C_2\mathcal{S}$ and study the saddle connections on \mathcal{S}^R . We claim that the image of the saddle connection $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ on \mathcal{S}^R is the only saddle connection whose image is a vector with positive slope and horizontal component at most 1.

Let $\begin{pmatrix} a \\ b \end{pmatrix}$ be some saddle connection on \mathcal{S}^R with an image under $M_{x,y}$ (for $(x,y) \in \Omega_2$) that has positive slope and horizontal component at most 1. Note that $M_{x,y} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} ax + by \\ b/x \end{pmatrix}$, $0 < ax + by \leq 1$.

We compute the slope of this image as

$$\frac{b}{x(ax + by)} = \frac{1}{x(x(a/b) + y)}.$$

Then, note that if $a/b < 0$ (recall that the slope is positive), then the slope would be greater than the case where $a/b = 0$, i.e. when $\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, so we need only consider the cases when $a \geq 0$.

Moreover, since $\begin{pmatrix} a \\ b \end{pmatrix}$ cannot be horizontal (or else its image would be horizontal), b must be positive (if $\begin{pmatrix} a \\ b \end{pmatrix}$ had negative slope, we can assume without loss of generality that $a < 0$ and $b > 0$). The shortest vertical distance on \mathcal{S}^R is 1, so $b \geq 1$. Suppose for the sake of contradiction that $a > 0$. Since the shortest horizontal distance on the surface \mathcal{S}^R is 1, we have $a \geq 1$. Combining this with the fact that $b \geq 1$, we have

$$ax + by \geq x + y.$$

Recalling the definition of b ,

$$ax + by \geq x + y > x + 1 - x = 1,$$

which is a contradiction. This forces $a = 0$, so our desired original saddle connection must be $\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ over all of Ω_2 . □

In turn, this tells us that

$$R(x, y) = \frac{1}{xy} \text{ for } (x, y) \in \Omega_2.$$

3.3.2. *Description of $R(x, y)$ Over Ω_1 .* To rephrase an earlier remark, given $(x, y) \in \Omega_1$, we say the saddle connection $v = \begin{pmatrix} a \\ b \end{pmatrix}$ is a winner at (x, y) if $M_{x,y}v$ has horizontal component $0 < ax + by \leq 1$ and positive slope, and if for all $v' = \begin{pmatrix} a' \\ b' \end{pmatrix}$ saddle connections such that $M_{x,y}v'$ has horizontal component $0 < a'x + b'y \leq 1$ and positive slope, we have $0 < \text{slope}(M_{x,y}v) \leq \text{slope}(M_{x,y}v')$.

Note that we consider vertical v to have positive slope ∞ . In this case, $M_{x,y}v$ will be the next saddle connection to reach the x -axis with horizontal length at most 1, and thus the slope of $M_{x,y}v$ is the return time at (x, y) . In Ω_1 , there are only finitely many winners and each winner has an associated convex polygonal region $P_i \subset \Omega_1$ as described below.

Proposition 3.3. *In the region*

$$P_1 = \{(x, y) \in \Omega_1 \mid x + y > 1\}$$

the winner is $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$, which forms the vertical edge of the rectangle V_0 , so the return time is given by

$$R(x, y) = \frac{1}{xy}.$$

For $2 \leq i \leq \lfloor n/2 \rfloor$, in the region

$$P_i = \{(x, y) \in \Omega_1 \mid xh_{i-2} + yv_{i-2} \leq 1, xh_{i-1} + yv_{i-1} > 1, xh_1 + y > 1\}$$

the winner is $\begin{pmatrix} h_{i-2} \\ v_{i-2} \end{pmatrix}$, which is the positive-sloped diagonal of the rectangle V_{i-2} , so the return time is given by

$$R(x, y) = \frac{v_{i-2}}{x(h_{i-2}x + v_{i-2}y)}.$$

In the region

$$P_{\lfloor n/2 \rfloor + 1} = \{(x, y) \in \Omega_1 \mid xh_{\lfloor n/2 \rfloor - 1} + yv_{\lfloor n/2 \rfloor - 1} \leq 1, xh_{\lfloor n/2 \rfloor - 1} + yv_{\lfloor n/2 \rfloor - 2} > 1, xh_1 + y > 1\}$$

the winner is $\begin{pmatrix} h_{\lfloor n/2 \rfloor - 1} \\ v_{\lfloor n/2 \rfloor - 1} \end{pmatrix}$, which is the positive-sloped diagonal of the rectangle $V_{\lfloor n/2 \rfloor - 2}$, so the return time is given by

$$R(x, y) = \frac{v_{\lfloor n/2 \rfloor - 1}}{x(h_{\lfloor n/2 \rfloor - 1}x + v_{\lfloor n/2 \rfloor - 1}y)}.$$

For $\lfloor n/2 \rfloor + 2 \leq i < n$, in the region

$$P_i = \{(x, y) \in \Omega_1 \mid xh_{n-i+1} + yv_{n-i} \leq 1, xh_{n-i} + yv_{n-i-1} > 1, xh_1 + y > 1\}$$

the winner is $\begin{pmatrix} h_{n+1-i} \\ v_{n-i} \end{pmatrix}$, which is the positive-sloped diagonal of the rectangle H_{n+1-i} , so the return time is given by

$$R(x, y) = \frac{v_{n-i}}{x(h_{n+1-i}x + v_{n-i}y)}.$$

In the region

$$P_n = \{(x, y) \in \Omega_1 \mid h_1x + y \leq 1\}$$

the winner is $\begin{pmatrix} h_1 \\ v_0 \end{pmatrix}$, which is the positive-sloped diagonal of the rectangle H_1 , so the return time is given by

$$R(x, y) = \frac{v_0}{x(h_1x + v_0y)}.$$




FIGURE 8. Color-coded relevant saddle connections for $n = 7$




FIGURE 9. The corresponding division of Ω_1 for $n = 7$ using coordinates in \mathbb{R}^2 . Each saddle connection is color coded to match the region it wins in. The purple region is P_1 , the blue region is P_2 , the green region is P_3 , the yellow region is P_4 , the gray region is P_5 , the orange region is P_6 , and the red region is P_7 .

The proof of Proposition 3.3 is quite long, and requires a case-by-case analysis. Hence it is postponed to the 24-page Appendix A. On the other hand, using hyperbolic geometry we can compute the volume of $\mathrm{SL}_2(\mathbb{R})/\Gamma_{O_{2n}}$ and compare it to our computation. More precisely, since $R(x, y)$ is a roof function over Ω in the suspension space $\mathrm{SL}_2(\mathbb{R})/\Gamma$, computing the volume under $R(x, y)$ should yield the volume of $\mathrm{SL}_2(\mathbb{R})/\Gamma$. The table in Figure 10 provides numerical volume computations for several values of n to verify our division of the transversal.

n	Computed Volume	Actual Volume	Error
2	4.9348	$\pi^2/2$	< 0.001%
3	6.5797	$2\pi^2/3$	< 0.001%
4	7.4022	$3\pi^2/4$	< 0.001%
5	7.8957	$4\pi^2/5$	< 0.001%
6	8.2247	$5\pi^2/6$	< 0.001%
7	8.4597	$6\pi^2/7$	< 0.001%
8	8.6359	$7\pi^2/8$	< 0.001%
9	8.7730	$8\pi^2/9$	< 0.001%
10	8.8826	$9\pi^2/10$	< 0.001%
50	9.7622	$49\pi^2/50$	< 0.001%
100	9.7709	$99\pi^2/100$	< 0.001%

FIGURE 10. Numerical computation of the volume of $\mathrm{SL}_2(\mathbb{R})/\Gamma_{O_{2n}}$ using our transversal compared to the exact volume. Within the limits of our computational software, there was essentially no error in the computed volume.

4. SLOPE GAP DISTRIBUTIONS

Let $k \in \mathbb{N}$. Denote by $V(k)$ the vertical strip $(0, k] \times [0, \infty)$ and let $\Lambda_{sc}^S(k)$ be the set of holonomy vectors of \mathcal{S} that lie inside $V(k)$. Let

$$\mathbb{S}(k) = \{0 = s_0 < s_1 < s_2 < \dots\}$$

denote the ordered set of slopes of vectors in $\Lambda_{sc}^S(k)$. Recall that the horocycle flow

$$\left\{ h_s = \begin{pmatrix} 1 & 0 \\ -s & 1 \end{pmatrix} \middle| s \in \mathbb{R} \right\}$$

acts on slopes by translation, so the slopes in $\mathbb{S}(k)$ are precisely the times when one of the vectors in $\Lambda_{sc}^S(k)$ hits the horizontal axis.

From (1) we see that the Veech group Γ of \mathcal{S} contains the parabolic element $S_2^{-1} = h_1$, which implies that \mathcal{S} is periodic under the horocycle flow with period 1. Hence $h_1 \Lambda_{sc}^S(k) = \Lambda_{sc}^S(k)$, i.e. after subtracting 1 from all of the slopes of holonomy vectors in the vertical strip $V(k)$ and removing the slopes that are negative, we get the same ordered set of slopes $\mathbb{S}(k)$ that we started with. In other words, there exists some $N(k) \in \mathbb{N}$ such that $s_{N(k)+i} - 1 = s_i$ for all $i \geq 0$. This in turn implies that the slope gaps $s_i - s_{i-1}$ repeat after $i = N(k)$, since

$$s_{N(k)+i} - s_{N(k)+i-1} = (s_i + 1) - (s_{i-1} + 1) = s_i - s_{i-1}.$$

Thus the gaps between the elements of $\mathbb{S}(k)$ are given by the (not renormalized) gap set

$$\overline{\mathcal{G}}(k) = \{s_i - s_{i-1} \mid s_i \in \mathbb{S}(k), 1 \leq i \leq N(k)\}.$$

Since $N(k)$ grows like k^2 , we may thus define our renormalized gap set to be

$$\mathcal{G}(k) = \{k^2(s_i - s_{i-1}) \mid s_i \in \mathbb{S}(k), 1 \leq i \leq N(k)\}.$$

Recall that our goal is to find the distribution of the renormalized gaps in $\mathcal{G}(k)$ as $k \rightarrow \infty$, i.e. we want to find a probability density function $f : \mathbb{R} \rightarrow [0, \infty)$ such that

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{G}(k) \cap (a, b)|}{N(k)} = \int_a^b f(x) dx$$

for any interval $(a, b) \subseteq \mathbb{R}$. We will do this by representing the quantity on the left as a limit of Birkhoff sums over longer and longer periodic orbits for the first return map to the transversal Ω and leveraging known ergodic theory results for such limits.

Let

$$g_k = \begin{pmatrix} 1/k & 0 \\ 0 & k \end{pmatrix}.$$

Observe that $g_k V(k) = V(1)$ and $\text{slope}(g_k v) = k^2 \text{slope}(v)$ for any $v \in \mathbb{R}^2$ and $k \in \mathbb{N}$. Moreover, one can verify that

$$\begin{aligned} g_k \cdot \Lambda_{sc}^S(k) &= \Lambda_{sc}^{g_k S}(1), \\ k^2 \mathbb{S}(k) &= \mathbb{S}^{g_k S}, \end{aligned}$$

and thus

$$\mathcal{G}(k) = \overline{\mathcal{G}}^{g_k S}$$

where $\mathbb{S}^{g_k S}$ denotes the ordered set of slopes of vectors in $\Lambda_{sc}^{g_k S}(1)$ and $\overline{\mathcal{G}}^{g_k S}$ is the associated non-renormalized gap set. In other words, the renormalized gaps between slopes of saddle connections on \mathcal{S} with horizontal length less than or equal to k is precisely equal to the set of gaps between slopes of saddle connections on $g_k \mathcal{S}$ with horizontal length less than or equal to 1.

Let $T : \Omega \rightarrow \Omega$ denote the first return map for the horocycle flow to the transversal Ω , i.e. for $x = g\Gamma \in \Omega$ (considered as a subset of $\text{SL}_2(\mathbb{R})/\Gamma$) we have

$$T(x) = h_{R(x)} x$$

where $R : \Omega \rightarrow \mathbb{R}^+$ is the usual return time function.

Observe that since $\Gamma \in \Omega$ is periodic under the horocycle flow with period 1, then $g_k \Gamma \in \Omega$ is periodic with period k^2 . Since the horocycle flow acts on slopes by translation, it preserves gaps between slopes. Then the slope gaps in

$$\mathcal{G}(k) = \overline{\mathcal{G}}^{g_k S} = \{s'_i - s'_{i-1} \mid s'_i \in \mathbb{S}^{g_k S}, 1 \leq i \leq N(k)\}$$

are precisely the times elapsed between consecutive intersections of the orbit of $g_k \Gamma$ with Ω under the horocycle flow, i.e.

$$s'_i - s'_{i-1} = R(T^i(g_k \Gamma))$$

for all $i \geq 0$. If we let

$$\chi(x) = \begin{cases} 1 & R(x) < t \\ 0 & R(x) \geq t \end{cases},$$

be the indicator function on $R^{-1}([0, t])$, then we can write a formula for the distribution of slope gaps as a limit of Birkhoff sums of the form

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{G}(k) \cap [0, t]|}{N(k)} = \lim_{k \rightarrow \infty} \frac{1}{N(k)} \sum_{i=0}^{N(k)-1} \chi(T^i(g_k \Gamma)). \quad (2)$$

Thus, we have reframed our geometric problem about slope gaps on \mathcal{S} into a dynamical problem concerning return times for a sequence of periodic points in Ω .

We may now apply the following theorem and its corollary, which are adapted from Theorems 4.1-4.2 in [3], or Theorem 2.5 in [1], which applies in a more general context. An analogous theorem

that applies to non-lattice surfaces also appears in [3, 1] and relies on the equidistribution of long horocycle orbits proved in [5], however we do not need this result for our setting.

Theorem 4.1 ([3], Theorem 4.1). *Let (X, ω) be a lattice surface with Veech group Γ , and let $\Omega \subset \mathrm{SL}_2(\mathbb{R})/\Gamma$ and T be the transversal and associated return map defined previously. If $x_k \in \Omega$ is a sequence of points that are periodic under the return map T with periods $N(k) \rightarrow \infty$ as $k \rightarrow \infty$, then for any bounded, measurable function $f : \Omega \rightarrow \mathbb{R}$, we have*

$$\lim_{k \rightarrow \infty} \frac{1}{N(k)} \sum_{i=0}^{N(k)-1} f(T^i(x_k)) = \int_{\Omega} f dm$$

where m is the unique ergodic probability measure for T supported on Ω .

Proof. We can realize $\mathrm{SL}_2(\mathbb{R})/\Gamma$ as a suspension space over the transversal Ω with roof function given by the return time function:

$$\mathrm{SL}_2(\mathbb{R})/\Gamma \cong \{(x, s) \mid x \in \Omega, s \in [0, R(x, y)]\} / \sim$$

where $(x, R(x)) \sim (T(x), 0)$. The probability Haar measure μ on $\mathrm{SL}_2(\mathbb{R})/\Gamma$ then decomposes as $d\mu = C ds dm$ for some constant C . A point $x_k \in \Omega$ is periodic for the return map T if and only if x_k is periodic under the continuous horocycle flow on $\mathrm{SL}_2(\mathbb{R})/\Gamma$. Let ρ_k denote the invariant probability measure on $\mathrm{SL}_2(\mathbb{R})/\Gamma$ supported on the periodic orbit of x_k under the horocycle flow, and let σ_k denote the invariant probability measure on Ω supported on the periodic orbit of x_k under T . Observe that for bounded, measurable $f : \mathrm{SL}_2(\mathbb{R})/\Gamma \rightarrow \mathbb{R}$,

$$\begin{aligned} \int f d\rho_k &= \frac{1}{k^2} \int_0^{k^2} f(h_s x_k) ds, \\ &= \frac{N(k)}{k^2} \frac{1}{N(k)} \sum_{i=0}^{N(k)-1} \int_0^{R(T^i(x_k))} f(h_s T^i(x_k)) ds, \\ &= \frac{N(k)}{k^2} \int_{\Omega} \int_0^{R(x)} f(h_s x) ds d\sigma_k(x), \end{aligned}$$

in other words, $d\rho_k = \frac{N(k)}{k^2} ds d\sigma_k$. Moreover, by a theorem of Sarnak [13], we know that long periodic horocycle orbits equidistribute with respect to the Haar measure, i.e. $d\rho_k \rightarrow \mu = ds dm$. Since we know that $\frac{N(k)}{k^2} \rightarrow C$ by [18], this implies that the corresponding long periodic orbits under T must equidistribute in Ω , i.e. $d\sigma_k \rightarrow dm$, which gives the result. \square

One can check that the measure dm on Ω is the appropriately scaled Lebesgue measure $dx dy$ on our parametrization of the transversal given above.

Corollary 4.2 ([3], Theorem 4.2). *Let the setting be as above, and let $\mathcal{G}(k)$ be the renormalized gap set for (X, ω) . Then*

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{G}(k) \cap [0, t]|}{N(k)} = m(\{(x, y) \in \Omega \mid R(x, y) \in [0, t]\}). \quad (3)$$

Proof. This is a direct application of Theorem 4.1 for the function χ and the sequence of points $x_k = g_k \Gamma$ with periods $N(k) = k^2$, along with the identity in (2). \square

With our parametrization of the transversal and formula for the return time function at every point in the transversal calculated in the last section, it is now a standard problem in multivariable calculus to compute the right hand side of (3). This gives us Theorem 1.1 from the introduction.

Since our limiting distribution has no small gaps, this indicates that the slope gap distribution for O_{2n} is not random (if this were the case, we would expect an exponential distribution *with* small

gaps). In Figure 11 we provide a few examples of the slope gap distributions on O_{2n} for several values of n .

An interesting conclusion that can be drawn from our calculation of the distribution for the $2n$ -gon is that the slope gap distribution of a translation surface is not always unimodal. For example, in the distribution for O_{14} one finds that there is a local maximum at $t \approx 0.715353$ of about 0.691264, followed by a local minimum at $t \approx 0.781831$ of about 0.681558, followed by another local maximum at $t \approx 0.870497$ of about 0.700232. This answers a question of Jayadev Athreya.




FIGURE 11. Gap distributions for the $2n$ -gon for several values of n . Note the changed scaling on the horizontal axis in the last two graphs.

5. BOUNDS ON THE NUMBER OF NON-DIFFERENTIABILITY POINTS

Now that we have an explicit description of the slope gap distribution, we can prove Theorem 1.3. In particular, since the slope gap distribution was computed by finding the area bounded by $R(x, y) = 0$ and $R(x, y) = t$ over $\Omega = \Omega_1 \cup \Omega_2$, we can see that non-differentiability points occur when the pieces of the return time function cross region boundaries on the transversal. Recall the theorem statement as follows.

Theorem. *The number of non-differentiable points in the slope gap distribution for a regular unit $2n$ -gon has linear lower and upper bounds. In particular,*

$$\left\lfloor \frac{n}{2} \right\rfloor - 59 \leq \#(\text{Non-Differentiable Points}) \leq \left\lceil \frac{n}{2} \right\rceil + 3n - 6.$$

Proof. Recall our division of Ω_1 into the regions P_1, P_2, \dots, P_n . Since the slope gap distribution is computed by finding the area bounded by $R(x, y) = 0$ and $R(x, y) = t$ over $\Omega = \Omega_1 \cup \Omega_2$, note that the distribution has a non-differentiable point at $t = k$ only if the return time function crosses a boundary of some P_i at $t = k$. We can disregard the boundary crossings on Ω_2 since they coincide with the boundary crossings for P_1 (they are the same region in \mathbb{R}^2 and have the same return time function as well). Thus, we can find an upper bound on the number of non-differentiable points by simply finding a bound on the number of boundary crossings on Ω_1 . The bulk of our proof will be in finding this upper bound. We will then use the information we gather for computing the upper bound to get a linear lower bound in the second half of this chapter. There are seven cases we must consider.

Case 1. We first count the number of boundary crossings in the region P_1 . Recall the definition of

$$P_1 = \{(x, y) \in \Omega_1 \mid x + y > 1\}$$

and that $R(x, y) = \frac{1}{xy}$ over P_1 . Thus, for any n , our region has the following shape:




FIGURE 12. An illustration of P_1 with coordinates in \mathbb{R}^2 , for any n , along with the hyperbola $R(x, y) = t$ (in black; here $t = 3$).

As $t \rightarrow \infty$, the function $\frac{1}{xy} = t$ moves towards the origin, but never crosses the points $(0, 1)$ and $(1, 0)$ (the upper left and lower right corners of P_1). Thus, $\frac{1}{xy}$ first enters P_1 through the point $(1, 1)$, which it crosses when

$$t = 1,$$

giving us a point of non-differentiability there. Moreover, when

$$t = 4,$$

$\frac{1}{xy} = t$ crosses the line $y = 1 - x$, giving us another possible point of non-differentiability. Since this function never passes through $(0, 1)$ or $(1, 0)$, there are no other possible boundary crossings in this case. Thus, we have **two** boundary crossings in this case.

Case 2. We now count the number of boundary crossings in the region P_2 . Recall that in this case,

$$P_2 = \{(x, y) \in \Omega_1 \mid xh_0 + yv_0 \leq 1, xh_1 + yv_1 > 1, xh_1 + y > 1\}$$

with $R(x, y) = \frac{v_0}{x(h_0x + v_0y)}$. The constraints on P_i in this case give us four sides: the line $xh_0 + yv_0 = 1$, the line $xh_1 + yv_1 = 1$, the line $xh_1 + y = 1$, and the line $x = 1$. Denote the intersection of $xh_0 + yv_0 = 1$ and $xh_1 + y = 1$ as A ; the intersection of $xh_0 + yv_0 = 1$ and $x = 1$ as B ; the intersection of $xh_1 + yv_1 = 1$ and $x = 1$ as C , and the intersection of $xh_1 + yv_1 = 1$ and $xh_1 + y = 1$ as D . For instance, we can refer to the following diagram to get a good idea of the general shape of these P_i and the labelling that we will be using:




FIGURE 13. In this example, $n = 7$. The black line is the hyperbola $R(x, y) = 3.6$.

The hyperbola enters the region from the right as $t \rightarrow \infty$, it moves to the left. Thus, the hyperbola always enters the region at the point B . Since $h_0 = v_0 = 1$ for all n , we see that the coordinates of B are always $(0, 1)$. Then,

$$R(x, y) = \frac{1}{x(x+y)} = t$$

intersects $(1, 0)$ at $t = 1$, which does not give us a new non-differentiability point (we had already accounted for $t = 1$ in case 1). Moreover, we find that the line CD (which is $xh_1 + yv_1 = 1$) intersects $x = 1$ when y satisfies $h_1 + yv_1 = 1$, or equivalently, $y = \frac{1-h_1}{v_1} = -1$. In particular, this means that the coordinates for C are $(1, -1)$. Plugging these values into $R(x, y)$ gives us an undefined value, meaning that $R(x, y)$ never crosses C . Similarly, plugging in $A := (0, 1)$ gives us an undefined value for $R(x, y)$, so $R(x, y)$ never crosses A .

Since the hyperbola never crosses A or C , we conclude that the first intersections of the hyperbola with the lines AD and CD either both occur inside the region P_2 or are both outside. We now solve for the first intersection of $\frac{1}{x(x+y)} = t$ and the line $AD := xh_1 + y = 1$. Doing so gives us a quadratic equation with discriminant $t^2 - 8 \cos(\pi/n)t$. This means that

$$t = 8 \cos(\pi/n)$$

gives us the first intersection between the hyperbola and AD .

Similarly, we solve for the intersection between $\frac{1}{x(x+y)} = t$ and the line $CD := xh_1 + yv_1 = 1$, giving us a quadratic equation with discriminant $t^2 - 8 \cos(\pi/n)t$ as well. In other words, the first intersections of the hyperbola with the lines AD and CD happen simultaneously.

Similarly, we see that the crossing of D happens at $t = (1 + 2 \cos(\pi/n))^2$.

Thus, this case gives us at most **two** new boundary crossings: the possible (they might not be in the region P_2) simultaneous crossing of AD and DC and the possible crossing of D .

Case 3. We now count the number of boundary crossings in the regions P_i for $2 < i < \lfloor n/2 \rfloor + 1$. Recall that in this case,

$$P_i = \{(x, y) \in \Omega_1 \mid xh_{i-2} + yv_{i-2} \leq 1, xh_{i-1} + yv_{i-1} > 1, xh_1 + y > 1\}$$

with $R(x, y) = \frac{v_{i-2}}{x(h_{i-2}x + v_{i-2}y)}$. The constraints on P_i in this case give us four sides: the line $xh_{i-2} + yv_{i-2} = 1$, the line $xh_{i-1} + yv_{i-1} = 1$, the line $xh_1 + y = 1$, and the line $x = 1$. Denote the intersection of $xh_{i-2} + yv_{i-2} = 1$ and $xh_1 + y = 1$ as A ; the intersection of $xh_{i-2} + yv_{i-2} = 1$ and $x = 1$ as B ; the intersection of $xh_{i-1} + yv_{i-1} = 1$ and $x = 1$ as C , and the intersection of $xh_{i-1} + yv_{i-1} = 1$ and $xh_1 + y = 1$ as D . For instance, we can refer to Figure 14 to get a good idea of the general shape of these P_i and the labelling that we will be using.




FIGURE 14. In this example, $n = 7$ and $i = 3$. The black curve is the hyperbola $R(x, y) = 3.6$.

The hyperbola enters the region at point B , which has coordinates $(1, \frac{1-h_{i-2}}{v_{i-2}})$. We can solve to see that this happens at

$$t = v_{i-2}.$$

This is an important fact that we can use for computing a lower bound, and we will refer back to this fact later.

We now note that since A is the intersection of $xh_{i-2} + yv_{i-2} = 1$ and $xh_1 + y = 1$, it has coordinates

$$\left(\frac{v_{i-2} - 1}{h_{i-2} - h_1 v_{i-2}}, 1 - \frac{h_1(v_{i-2} - 1)}{h_{i-2} - h_1 v_{i-2}} \right).$$

Similarly, we find that C has coordinates

$$\left(1, \frac{1}{v_{i-1}} - \frac{h_{i-1}}{v_{i-1}} \right).$$

Using computational assistance, we see that $R(x, y)$ crosses A and C at

$$t = \frac{\csc^2(\pi/n) \sin((i-1)\pi/n)}{\cot(\pi/n) - \cot(i\pi/(2n))},$$

i.e., the hyperbola crosses A and C simultaneously.

Since we know that the intersection with A and C happen simultaneously and since our hyperbola moves to the left as $t \rightarrow \infty$, we see that the hyperbola must make its first intersections with the downward sloping lines AD and CD inside the region P_i .

We claim that these intersections are simultaneous. Indeed, solving for the intersection of $R(x, y)$ and AD gives a quadratic with discriminant $t^2 - 4t(h_1 - h_{i-2}/v_{i-2})$, i.e., the first intersection happens at $t = 4(h_1 - h_{i-2}/v_{i-2})$. Moreover, solving for the intersection of $R(x, y)$ and CD gives a quadratic with discriminant $t^2/v_{i-1}^2 - 4t(h_{i-1}/v_{i-1} - h_{i-2}/v_{i-2})$, i.e., the first intersection happens at $t = 4v_{i-1}^2(h_{i-1}/v_{i-1} - h_{i-2}/v_{i-2})$. Using computational assistance, we verify that both of these quantities are equal to

$$t = 4 \sin(i\pi/n) \csc(\pi(i-1)/n),$$

meaning that the hyperbola crosses AD and CD at the same time.

We can similarly solve to see that the hyperbola may cross point D at

$$t = \frac{\csc(\pi/n) \sin(\pi(i-1)/n) \sin^2(\pi(i+1)/n)}{(\sin(i\pi/n) - \sin(\pi/n))^2}.$$

Thus, this gives us **four** total crossings for each i in this case: a crossing at B , a (possible) simultaneous crossing of AD and DC , a simultaneous crossing of A and C , and a (possible) crossing of D .

Case 4. We now count the number of boundary crossings in $P_{\lfloor n/2 \rfloor + 1}$.

Subcase 4.1. Let n be odd. For the sake of ease, we will note that $\lfloor n/2 \rfloor = \lceil n/2 \rceil - 1$ for odd n . Thus we can rewrite

$$P_{\lfloor n/2 \rfloor + 1} = \{(x, y) \in \Omega_1 \mid xh_{\lfloor n/2 \rfloor - 2} + yv_{\lfloor n/2 \rfloor - 2} \leq 1, xh_{\lfloor n/2 \rfloor - 1} + yv_{\lfloor n/2 \rfloor - 2} > 1, xh_1 + y > 1\}$$

with

$$R(x, y) = \frac{v_{\lfloor n/2 \rfloor - 2}}{x(h_{\lfloor n/2 \rfloor - 2}x + v_{\lfloor n/2 \rfloor - 2}y)}.$$

The constraints on P_i in this case give us four sides: the line $xh_{\lfloor n/2 \rfloor - 2} + yv_{\lfloor n/2 \rfloor - 2} = 1$, the line $xh_{\lfloor n/2 \rfloor - 1} + yv_{\lfloor n/2 \rfloor - 1} = 1$, the line $xh_1 + y = 1$, and the line $x = 1$. Denote the intersection of $xh_{\lfloor n/2 \rfloor - 2} + yv_{\lfloor n/2 \rfloor - 2} = 1$ and $xh_1 + y = 1$ as A ; the intersection of $xh_{\lfloor n/2 \rfloor - 2} + yv_{\lfloor n/2 \rfloor - 2} = 1$ and $x = 1$ as B ; the intersection of $xh_{\lfloor n/2 \rfloor - 1} + yv_{\lfloor n/2 \rfloor - 1} = 1$ and $x = 1$ as C , and the intersection of $xh_{\lfloor n/2 \rfloor - 1} + yv_{\lfloor n/2 \rfloor - 1} = 1$ and $xh_1 + y = 1$ as D . For instance, we can refer to the following diagram to get a good idea of the general shape of $P_{\lfloor n/2 \rfloor}$ and the labelling that we will be using:




FIGURE 15. In this example, $n = 7$. The black line is the equation $R(x, y) = 3.6$.

The hyperbola enters this region by crossing the point B . The computation for the time at which this crossing happens is identical to the analogous computation in Case 3, that is, the hyperbola crosses B at

$$t = v_{\lceil n/2 \rceil - 2}.$$

We can see that A is the intersection of $xh_{\lceil n/2 \rceil - 2} + yv_{\lceil n/2 \rceil - 2} = 1$ and $xh_1 + y = 1$, so it has coordinates

$$\left(\frac{v_{\lceil n/2 \rceil - 2} - 1}{h_{\lceil n/2 \rceil - 2} - h_1 v_{\lceil n/2 \rceil - 2}}, 1 - \frac{h_1 v_{\lceil n/2 \rceil - 2} - 1}{h_{\lceil n/2 \rceil - 2} - h_1 v_{\lceil n/2 \rceil - 2}} \right).$$

Similarly, we find that C has coordinates

$$\left(1, \frac{1}{v_{\lceil n/2 \rceil - 2}} - \frac{h_{\lceil n/2 \rceil - 1}}{v_{\lceil n/2 \rceil - 2}} \right).$$

Using computational assistance, we see that $R(x, y)$ passes through A when

$$t = \frac{\csc^2(\pi/2) \sin(\pi \lceil n/2 \rceil / n)}{\csc(\pi/n) + \csc((\pi - \pi \lceil n/2 \rceil) / n)}$$

and through C when

$$t = \frac{\csc(\pi/n) \sin(\pi(\lceil n/2 \rceil - 1) / n)}{1 - 2 \cos((\pi - \pi \lceil n/2 \rceil) / n)}.$$

Since n is odd, we can let $n = 2k + 1$ for some integer k so that $\lceil n/2 \rceil = k + 1$. Then, computational assistance tells us that these two quantities for t are indeed equal, i.e., $R(x, y)$ passes through A and C at the same time.

Since we know that the intersection with A and C happen simultaneously and since our hyperbola moves to the left as $t \rightarrow \infty$, we see that the hyperbola must make its first intersections with the downward sloping lines AD and CD inside the region P_i .

We claim that these intersections are simultaneous. Indeed, solving for the intersection of $R(x, y)$ and AD gives a quadratic with discriminant

$$t^2 - 4t(h_1 - h_{\lceil n/2 \rceil - 2} / v_{\lceil n/2 \rceil - 2}),$$

i.e., the first intersection happens at

$$t = 4(h_1 - h_{\lceil n/2 \rceil - 2} / v_{\lceil n/2 \rceil - 2}).$$

Moreover, solving for the intersection of $R(x, y)$ and CD gives a quadratic with discriminant

$$t^2 / v_{\lceil n/2 \rceil - 2}^2 - 4t(h_{\lceil n/2 \rceil - 1} / v_{\lceil n/2 \rceil - 2} - h_{\lceil n/2 \rceil - 2} / v_{\lceil n/2 \rceil - 2}),$$

i.e., the first intersection happens at

$$t = 4v_{\lceil n/2 \rceil - 2}^2 (h_{\lceil n/2 \rceil - 1} / v_{\lceil n/2 \rceil - 2} - h_{\lceil n/2 \rceil - 2} / v_{\lceil n/2 \rceil - 2}).$$

Since n is odd, let $n = 2k + 1$ for some integer k , so that $\lceil n/2 \rceil = k + 1$. Using computational assistance (more precisely, we were able to use Mathematica to show that the first quantity is equal to 4 and took the derivative of the second quantity with respect to k and found that the second quantity is constant, with value 4 at $k = 1$), we verify that both of these quantities for t are equal to 4, meaning that the hyperbola crosses AD and CD at the same time.

Thus, this gives us **four** total crossings in this case: a crossing at B , a (possible) simultaneous crossing of AD and DC , a simultaneous crossing of A and C , and a (possible) crossing of D .

Subcase 4.2. Let n be even. Our region is

$$P_{\lceil n/2 \rceil + 1} = \{(x, y) \in \Omega_1 \mid xh_{n/2-1} + yv_{n/2-1} \leq 1, xh_{n/2-1} + yv_{n/2-2} > 1, xh_1 + y > 1\}$$

with return time given by

$$R(x, y) = \frac{v_{n/2-1}}{x(h_{n/2-1}x + v_{n/2-1}y)}.$$

Adopt the same labelling system as in Subcase 4.1 (the general shape of the region is the same). Once again, the hyperbola enters this region by crossing the point B . The computation for the time at which this crossing happens is identical to the analogous computation in Case 3, that is, the hyperbola crosses B at

$$t = v_{n/2-1}.$$

We can see that A is the intersection of $xh_{n/2-1} + yv_{n/2-1} = 1$ and $xh_1 + y = 1$, so it has coordinates

$$\left(\frac{v_{n/2-1} - 1}{h_{n/2-1}} - h_1 v_{n/2-1}, 1 - \frac{h_1 v_{n/2-1} - 1}{h_{n/2-1} - h_1 v_{n/2-1}} \right).$$

Similarly, we find that C has coordinates

$$\left(1, \frac{1}{v_{n/2-2}} - \frac{h_{n/2-1}}{v_{n/2-2}} \right).$$

Using computational assistance, we see that $R(x, y)$ passes through A when

$$t = \frac{\cot(\pi/n)}{1 - \sin(\pi/n)}$$

and through C when

$$t = \frac{\csc(\pi/n)}{\sec(\pi/n) - \cot(\pi/(2n))(\sec(\pi/n) - 1)}.$$

Then, computational assistance tells us that these two quantities for t are indeed equal, i.e., $R(x, y)$ passes through A and C at the same time.

Since we know that the intersection with A and C happen simultaneously and since our hyperbola moves to the left as $t \rightarrow \infty$, we see that the hyperbola must make its first intersections with the downward sloping lines AD and CD *inside* the region P_i .

We claim that these intersections are simultaneous. Indeed, solving for the intersection of $R(x, y)$ and AD gives a quadratic with discriminant

$$t^2 - 4t(h_1 - h_{n/2-1}/v_{n/2-1}),$$

i.e., the first intersection happens at

$$t = 4(h_1 - h_{n/2-1}/v_{n/2-1}).$$

Moreover, solving for the intersection of $R(x, y)$ and CD gives a quadratic with discriminant

$$t^2/v_{n/2-2}^2 - 4t(h_{n/2-1}/v_{n/2-2} - h_{n/2-1}/v_{n/2-1}),$$

i.e., the first intersection happens at

$$t = 4v_{n/2-2}^2(h_{n/2-1}/v_{n/2-2} - h_{n/2-1}/v_{n/2-1}).$$

Using computational assistance, we verify that both of these quantities for t are equal to $\cos(\pi/n)$, meaning that the hyperbola crosses AD and CD at the same time.

Thus, this gives us **four** total crossings in this case: a crossing at B , a (possible) simultaneous crossing of AD and DC , a simultaneous crossing of A and C , and a (possible) crossing of D .

We conclude that regardless of whether n is even or odd, we get **four** crossings out of case 4.

Case 5. We now count the number of boundary crossings in P_i for $\lfloor n/2 \rfloor + 1 < i < n - 1$. Now, our region is

$$P_i = \{(x, y) \in \Omega_1 \mid xh_{n-i+1} + yv_{n-i} \leq 1, xh_{n-i} + yv_{n-i-1} > 1, xh_1 + y > 1\}$$

with return time given by

$$R(x, y) = \frac{v_{n-i}}{x(h_{n+1-i}x + v_{n-i}y)}.$$

The constraints on P_i in this case give us four sides: the line $xh_{n-i+1} + yv_{n-i} = 1$, the line $xh_{n-i} + yv_{n-i-1} = 1$, the line $xh_1 + y = 1$, and the line $x = 1$. Denote the intersection of $xh_{n-i+1} + yv_{n-i} = 1$ and $xh_1 + y = 1$ as A ; the intersection of $xh_{n-i+1} + yv_{n-i} = 1$ and $x = 1$ as B ; the intersection of $xh_{n-i} + yv_{n-i-1} = 1$ and $x = 1$ as C , and the intersection of $xh_{n-i} + yv_{n-i-1} = 1$ and $xh_1 + y = 1$ as D . For instance, we can refer to the following diagram to get a good idea of the general shape of these P_i and the labelling that we will be using:




FIGURE 16. Here, $n = 7$ and $i = 5$. The black curve is the plot of $R(x, y) = 3$.

The hyperbola enters the region at point B , which has coordinates $(1, \frac{1-h_{n-i+1}}{v_{n-i}})$. We can solve to see that this happens at

$$t = v_{n-i}.$$

Yet, these times coincide with the set of times where $R(x, y)$ crosses point B in the previous cases (in particular, these times are equal to the vertical lengths on the staircase), so this introduces **no** new non-differentiability points.

We now note that since A is the intersection of $xh_{n-i+1} + yv_{n-i} = 1$ and $xh_1 + y = 1$, it has coordinates

$$\left(\frac{v_{n-i} - 1}{h_{n-i+1} - h_1 v_{n-i}}, 1 - \frac{h_1(v_{n-i} - 1)}{h_{n-i+1} - h_1 v_{n-i}} \right).$$

Similarly, we find that C has coordinates

$$\left(1, \frac{1}{v_{n-i-1}} - \frac{h_{n-i}}{v_{n-i-1}} \right).$$

Using computational assistance, we see that $R(x, y)$ crosses A and C at

$$t = \frac{\csc^2(\pi/n) \sin((1-i+n)\pi/n)}{\cot(\pi/n) - \cot(i\pi/(2n))},$$

i.e., the hyperbola crosses A and C simultaneously.

Since we know that the intersection with A and C happen simultaneously and since our hyperbola moves to the left as $t \rightarrow \infty$, we see that the hyperbola must make its first intersections with the downward sloping lines AD and CD *inside* the region P_i .

We claim that these intersections are simultaneous. Indeed, solving for the intersection of $R(x, y)$ and AD gives a quadratic with discriminant

$$t^2 - 4t(h_1 - h_{n-i+1}/v_{n-i}),$$

i.e., the first intersection happens at $t = 4(h_1 - h_{n-i+1}/v_{n-i})$. Moreover, solving for the intersection of $R(x, y)$ and CD gives a quadratic with discriminant

$$t^2/v_{n-i-1}^2 - 4t(h_{n-i}/v_{n-i-1} - h_{n-i+1}/v_{n-i}),$$

i.e., the first intersection happens at $t = 4v_{n-i-1}^2(h_{n-i}/v_{n-i-1} - h_{n-i+1}/v_{n-i})$. Using computational assistance, we verify that both of these quantities are equal to

$$t = 4 \csc((\pi(i-1))/n) \sin((i\pi)/n),$$

meaning that the hyperbola crosses AD and CD at the same time.

We can similarly solve to see that the hyperbola may cross point D at

$$t = \frac{\csc(\pi/n) \sin(\pi(i-1)/n) \sin^2(\pi(i+1)/n)}{(\sin(i\pi/n) - \sin(\pi/n))^2}.$$

Thus, this gives us **three** new crossings for each i in this case: a (possible) simultaneous crossing of AD and DC , a simultaneous crossing of A and C , and a (possible) crossing of D .

Case 6. We now count the number of boundary crossings in P_{n-1} . Our region is

$$P_i = \{(x, y) \in \Omega_1 \mid xh_2 + yv_1 \leq 1, xh_1 + y > 1\}$$

with return time given by

$$R(x, y) = \frac{v_1}{x(h_2x + v_1y)}.$$

Our region is bounded by three lines: $xh_2 + yv_1 = 1$, $xh_1 + y = 1$, and $x = 1$. Label the intersection of $xh_2 + yv_1 = 1$ and $xh_1 + y = 1$ as A , the intersection of $xh_2 + yv_1 = 1$ and $x = 1$ as B , and the intersection of $xh_1 + y = 1$ and $x = 1$ as C . Our region has the following general shape:




FIGURE 17. This is the example of $n = 7$. The black curve is the equation $R(x, y) = 2$.

The hyperbola first enters the region at B . As in Case 5, we can find that this happens at $t = v_1$, which has already been accounted for. Moreover, the same argument we used to show that the hyperbola passes through A and C at the same time still applies in this case. Thus, this case gives us at most **two** new boundary crossings: a simultaneous crossing of A and C and a possible crossing of the line AC .

Case 7. We finally count the number of boundary crossings in P_n . Recall that

$$P_n = \{(x, y) \in \Omega_1 \mid xh_1 + y \leq 1\}$$

with $R(x, y) = \frac{v_0}{x(h_1x+v_0y)} = \frac{1}{x(h_1x+y)}$. Thus, our region has three boundaries: the line $xh_1 + y = 1$, the line $x = 1$, and the line $2(1 + \cos(\pi/n))x + y = 1$. Below is an example of what this region would look like:




FIGURE 18. The region P_n with $n = 7$ with the graph of the equation $R(x, y) = 3$.

We can easily see that the three corners of the region are $(1, 1-h_1)$, $(0, 1)$ and $(1, 1-2(1+\cos(\pi/n)))$. We can easily compute to see that the hyperbola enters the region by passing through $(1, 1-h_1)$ first at $t = 1$, which does not contribute a new time.

Moreover, we can verify that the hyperbola does not pass through $(0, 1)$ and $(1, 1-2(1+\cos(\pi/n)))$ because substituting these points in gives an undefined value for $R(x, y)$.

Thus, in this case, we have at most **one** new boundary crossing, that of the hyperbola passing through the line $2(1 + \cos(\pi/n))x + y = 1$.

Finishing Touches. We now add up all the boundaries we have counted. In Case 1, we found at most 2 boundary crossings. In Case 2, we found at most 2 boundary crossings as well. In Case 3, we found at most 4 crossings for each i , and there are $\lfloor n/2 \rfloor + 1 - 2 - 1 = \lfloor n/2 \rfloor - 2$ values of i satisfying the relevant conditions, giving us a total of at most $4\lfloor n/2 \rfloor - 8$ boundary crossings in this case. In Case 4, we found at most 4 boundary crossings. In Case 5, we found at most 3 boundary crossings for each value of i , and there are $(n-1) - (\lfloor n/2 \rfloor + 1) - 1 = n - \lfloor n/2 \rfloor - 3$ values of i in these cases, giving us at most $3n - 3\lfloor n/2 \rfloor - 9$ boundary crossings in these cases. In Case 6, we found at most 2 new boundary crossings. In Case 7, we found at most 1 new boundary crossing. Adding these values together, we have our desired bound of

$$\#(\text{Non-Differentiability Points}) \leq 2 + 2 + 4\lfloor n/2 \rfloor - 8 + 4 + 3n - 3\lfloor n/2 \rfloor - 9 + 2 + 1 = \left\lfloor \frac{n}{2} \right\rfloor + 3n - 6.$$

We can now compute a lower bound on the number of differentiability points. Let us recall the time stamps where the boundary crossings occur in each case in Figure 19.

We omit the explicit values for t_1, \dots, t_7 in the table above as they won't make a difference in obtaining a linear lower bound. In fact the values that show up outside of the third and fifth rows above can be neglected all together, as there are a total of 14 such values. We now focus on the time stamps coming from the third and fifth rows of the list.

We first note that, the first family of time stamps obtained by evaluating the functions k_1 and k_2 in the third and fifth regions give the same set as $k_1(i, n)$ and $k_2(i, n)$ run over the given indices. The remaining three families of time stamps $l(i, n), m(i, n), r(i, n)$ have exactly the same expression but they run over different indices, and there are $\lfloor n/2 \rfloor - 3$ or $\lfloor n/2 \rfloor - 2$ terms in each family, depending on the parity of n . Experimental evidence based on computations in Mathematica suggests that all of these times are distinct, and hence there is no potential cancellation of a point of non-differentiability. Therefore the conjectural lower bound for number of non-differentiability points is $7(\lfloor n/2 \rfloor - 3)$ and is comparable to the upper bound.

Region	Time Stamps
P_1	$t = 1, t = 4$
P_2	$t = 1, t = 8 \cos(\pi/n), t = (1 + 2 \cos(\pi/n))^2$
$P_i, 2 < i < \lfloor n/2 \rfloor + 1$	$t = k_1(i, n) := \csc\left(\frac{\pi}{n}\right) \sin\left(\frac{\pi(i-1)}{n}\right),$ $t = l(i, n) := \frac{\csc^2(\pi/n) \sin((i-1)\pi/n)}{\cot(\pi/n) - \cot(i\pi/(2n))},$ $t = m(i, n) := 4 \sin(i\pi/n) \csc(\pi(i-1)/n),$ $t = r(i, n) := \frac{\csc(\pi/n) \sin(\pi(i-1)/n) \sin^2(\pi(i+1)/n)}{(\sin(i\pi/n) - \sin(\pi/n))^2}$
$P_{\lfloor i \rfloor + 1}$	$t = t_1, t = t_2, t = t_3, t = t_4$
$P_i, \lfloor n/2 \rfloor + 1 < i < n - 1$	$t = k_2(i, n) := \csc\left(\frac{\pi}{n}\right) \sin\left(\frac{\pi(n-i+1)}{n}\right),$ $t = l(i, n) := \frac{\csc^2(\pi/n) \sin((i-1)\pi/n)}{\cot(\pi/n) - \cot(i\pi/(2n))},$ $t = m(i, n) := 4 \sin(i\pi/n) \csc(\pi(i-1)/n),$ $t = r(i, n) := \frac{\csc(\pi/n) \sin(\pi(i-1)/n) \sin^2(\pi(i+1)/n)}{(\sin(i\pi/n) - \sin(\pi/n))^2}$
P_{n-1}	$t = \csc(\pi/n) \sin(2\pi/n), t = t_5, \text{ and } t = t_6$
P_n	$t = 1, t = t_7$

FIGURE 19. Time Stamps

In what follows we will provide a reasoning for the more modest lower bound $\lfloor n/2 \rfloor - 59$ which is still linear in n . A straightforward computation in multivariable calculus shows that there is a value $m_0 \geq 6.07$ such that $k_1(i, n) \geq m_0$ when $9 \leq i \leq \lfloor n/2 \rfloor + 1$, and $n \geq 20$, and $l(i, n), r(i, n) \geq m_0$ when $9 \leq i \leq n - 10$, and $n \geq 20$.

On the other hand, for any $3 \leq i \leq n - 2$, $n \geq 5$ we have $m(i, n) \leq 6$. Furthermore, for any fixed n , the function m is one-to-one for $3 \leq i \leq n - 2$. This means that there at least $\lfloor n/2 \rfloor - 3$ distinct time stamps coming from the function $m(i, n)$ that are also distinct from the time stamps coming from k, l, r restricted to above domains. Since outside of those domains those functions each produce 14 time stamps, there are at most $14 * 3$ time stamps coming from k, l, r that could possibly agree with the time stamps from m . Together with the initial 14, that are outside of region 3 and region 5, we get $\lfloor n/2 \rfloor - 3 - 56 = \lfloor n/2 \rfloor - 59$ unique time stamps.

Combining these two bounds, we conclude:

$$\left\lfloor \frac{n}{2} \right\rfloor - 59 \leq \#(\text{Non-Differentiability Points}) \leq \left\lfloor \frac{n}{2} \right\rfloor + 3n - 6.$$

□

The plot in Figure 20 shows our upper bound, the actual number of boundary crossings, and the actual number of non-differentiable points on the distribution (computed using numerical methods on Mathematica).

Thus, from Figure 20 we can see that our bound is a reasonable estimate for the number of boundary crossings, and for small values of n is a reasonably good estimate of the number of non-differentiable points. In any case, we have shown that there is a linear upper bound on the number of non-differentiable points for the slope gap distribution of the $2n$ -gon. The table below depicts the same data as in Figure 20.




FIGURE 20. Plot of the bound from Theorem 1.3 (olive dashed line), the actual number of boundary crossings (cyan circles), and the actual number of non-differentiable points (pink squares) against n .

n	Upper Bound	# of Boundary Crossings	# of Non-Differentiable Points
4	8	7	7
5	11	9	9
6	15	13	13
7	18	16	15
8	22	21	18
9	25	23	20
10	29	28	23
11	32	31	25
12	36	35	–
13	39	38	–
14	43	42	–
15	46	45	–
16	50	49	–
17	53	52	–
18	57	56	–
19	60	59	–
20	64	63	–
21	67	65	–

6. APPENDIX A: RETURN TIME FUNCTION COMPUTATIONS

Proof of Proposition 3.3. We begin with an outline of how the proof proceeds. We must show that at every point in Ω_1 , the winning vector is the one we assert it to be in Proposition 3.3; the winning vector will dictate the return time at that point. We begin by finding the set of all saddle connection holonomy vectors which win at some $(x, y) \in \Omega_1$. By Lemma 6.1, this is exactly the set of vectors which win at points $(1, y) \in \Omega_1$, meaning that it suffices to find the winning vectors for points along the right edge ($x = 1$) of Ω_1 . We claim that the vectors specified in Proposition 3.3 win at these points (i.e. λ_i wins at points $(1, y) \in P_i$). To prove this, we establish conditions which any other vector must satisfy in order to win over our proposed vector in each region P_i , then we prove

by exhaustive casework that for each P_i there is no saddle connection holonomy vector satisfying these conditions. This proves that our candidate vectors (those specified in Proposition 3.3) are the only vectors which win at points $(1, y) \in \Omega_1$, and hence at any point in Ω_1 . Finally, we use the definition of a dominating vector to describe the regions where each member of this finite set of vectors (the λ_i vectors) wins, fully determining the return time function on Ω_1 .

We will first establish some useful lemmas, then we will address the 4 cases provided in the lemma separately.

First of all, notice that the restrictions $ax + by > 0$, $\text{slope}(M_{x,y}v) > 0$ and $x > 0$ together necessitate that $b > 0$, so we only need to consider saddle connections with positive horizontal component.

Also, for $v = \begin{pmatrix} a \\ b \end{pmatrix}$ and $v' = \begin{pmatrix} a' \\ b' \end{pmatrix}$, if $\text{slope}(v) \leq \text{slope}(v')$, then we have:

$$\begin{aligned} \text{slope}(M_{x,y}v) - \text{slope}(M_{x,y}v') &= \frac{bx^{-1}}{ax + by} - \frac{b'x^{-1}}{a'x + b'y} = \\ \frac{a'b + bb'yx^{-1} - ab' - bb'yx^{-1}}{(ax + by)(a'x + b'y)} &= \frac{a'b - ab'}{(ax + by)(a'x + b'y)} = \\ aa' \cdot \frac{b/a - b'/a'}{(ax + by)(a'x + b'y)} &\leq 0 \end{aligned}$$

This proves that $\text{slope}(v) \leq \text{slope}(v')$ implies $\text{slope}(M_{x,y}v) \leq \text{slope}(M_{x,y}v')$, so to determine if v' wins over v , we can directly compare slopes.

Lemma 6.1. *The set of saddle connections which win at some $(x, y) \in \Omega_1$ is exactly the set of saddle connections which win at some $(1, y) \in \Omega_1$.*

Proof. Suppose the saddle connection with holonomy vector $v_0 = \begin{pmatrix} a_0 \\ b_0 \end{pmatrix}$ is the saddle connection whose image under some $K = M_{x_0, y_0}$ with $(x_0, y_0) \in \Omega_1$ has the smallest positive slope of any image of a saddle connection with horizontal length at most 1. We claim v_0 is also the vector satisfying this property for the matrix

$$K' = M_{1, y_0 - (1 - x_0)a_0/b_0} = \begin{pmatrix} 1 & y_0 - (1 - x_0)a_0/b_0 \\ 0 & 1 \end{pmatrix}$$

First we must show that $(1, y_0 - (1 - x_0)a_0/b_0) \in \Omega_1$. In order for v_0 to satisfy the conditions stated above, it must be true that $a_0x_0 + b_0y_0 \leq 1$. We also know from the boundaries of Ω_1 that $y_0 > -2(1 + \cos(\pi/n))x_0 + 1$, and we know that because the smallest vertical edge in S' is 1, $b_0 \geq 1$. Using these inequalities, we find:

$$\begin{aligned} a_0x_0 - 2(1 + \cos(\pi/n))b_0x_0 &\leq \\ a_0x_0 - 2(1 + \cos(\pi/n))b_0x_0 + b_0 - 1 &< \\ a_0x_0 + b_0y_0 - 1 &\leq 0, \\ a_0/b_0 &> -2(1 + \cos(\pi/n)) \end{aligned}$$

We also know a_0/b_0 is positive, so $-a_0/b_0 < 0$. Thus, $-a_0/b_0$ is between the slopes of the lower and upper boundaries of Ω_1 , so the line of slope $-a_0/b_0$ passing through (x_0, y_0) will intersect the right edge of Ω_1 (which is included in Ω_1) at $x = 1$; this point is exactly $(1, y_0 - (1 - x_0)a_0/b_0)$.

Now we show that $K'v_0$ is the saddle connection of horizontal length at most 1 with the smallest positive slope. Since the image of v_0 under K has horizontal length at most 1, we know

$$x_0a_0 + y_0b_0 \leq 1$$

Well, the same must be true for the horizontal length of the image of v_0 under K' . Henceforth we will denote the horizontal component of Kv_0 as c , and we will denote the horizontal component of $K'v_0$ as c' .

$$c' = 1 \cdot a_0 + (y_0 - (1 - x_0)a_0/b_0) \cdot b_0 = x_0a_0 + y_0b_0 \leq 1$$

Now let us consider the set of vectors v whose image under K' may have a smaller positive slope than $K'v_0$. Both K and K' are shear matrices which preserve the order of slopes of vectors, so v must have smaller slope than v_0 , and Kv must have smaller slope than Kv_0 . In other words, for

$v = \begin{pmatrix} a \\ b \end{pmatrix}$, it must be true that

$$b/a < b_0/a_0 \rightarrow a_0b/b_0 < a$$

Now consider the horizontal component of $K'v$:

$$\begin{aligned} c' &= 1 \cdot a + (y_0 - (1 - x_0)a_0/b_0) \cdot b = (a - a_0b/b_0)(1 - x_0) + ax_0 + by_0 \\ &= |a - a_0b/b_0| \cdot |1 - x_0| + ax_0 + by_0 > ax_0 + by_0 = c \end{aligned}$$

Well, since v_0 is the saddle connection whose image under K has the smallest positive slope of any image of a saddle connection with horizontal length at most 1, and since Kv has smaller slope than Kv_0 , it must be true that the horizontal component of Kv is greater than 1, so

$$c' > c > 1$$

Thus there is no vector v such that $k'v$ has smaller slope than $K'v_0$ and $K'v$ has horizontal component at most 1. In other words, if v_0 is the saddle connection whose image under some $K = M_{x_0, y_0}$ with $(x_0, y_0) \in \Omega_1$ has the smallest positive slope of any image of a saddle connection with horizontal length at most 1, then v_0 also satisfies this property for some $K' = M_{1, y}$ with $(1, y) \in \Omega_1$. \square

For the sake of brevity, we introduce some new notation. Let σ_i for $1 \leq i \leq \lceil n/2 \rceil - 1$ represent the saddle connections joining the lower left and upper right vertices on the rectangles H_i and ν_i for $0 \leq i \leq \lfloor n/2 \rfloor - 1$ represent the saddle connections joining the lower left and upper right vertices on the rectangles V_i , as shown in the below diagram for $n = 7$.




FIGURE 21. Labelling of S' using σ 's and ν 's for $n = 7$




FIGURE 22. Labelling of S' with λ 's for $n = 7$

Note that as holonomy vectors, $\sigma_i = \begin{pmatrix} h_i \\ v_{i-1} \end{pmatrix}$ and $\nu_i = \begin{pmatrix} h_i \\ v_i \end{pmatrix}$. Thus, restating Proposition 3.3 in terms of this new notation, our goal is to show that the vector $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ wins on P_1 , that ν_{i-2} wins on P_i for $1 < i \leq \lceil n/2 \rceil$, that σ_{n-i} wins on P_i for $\lfloor n/2 \rfloor + 1 < i < n$, and that $\begin{pmatrix} h_1 \\ v_0 \end{pmatrix}$ wins on P_n . Let λ_i denote the vector which we claim wins on P_i as described above.

The above lemma implies that in order to find the full set of winning vectors for Ω_1 , we only need to consider the set of vectors which win at points in the intersection of Ω_1 and the line $x = 1$. We will first establish that this set of vectors is indeed the λ_i vectors described above. Then, knowing that one of the λ_i wins at every point in Ω_1 , we will show that each λ_i wins on P_i as compared to the other λ_j vectors; since $\cup_i P_i = \Omega_1$, this provides a full description of the return time function on Ω_1 .

For each i , in order to prove that λ_i wins on $P_i \cap \{x = 1\}$, we establish four conditions which must be satisfied by any vector $\begin{pmatrix} a \\ b \end{pmatrix}$ which wins over to λ_i in the region, and prove that all saddle connections on S' apart from λ_i violate at least one condition.

- **Condition 1** is a restriction on the horizontal length of the vector, i.e. there exists some $q > 0$ such that $a < q$ (we use an explicit q which differs between cases).
- **Condition 2** is the restriction that $b/a < \text{slope}(\lambda_i)$ (otherwise $\begin{pmatrix} a \\ b \end{pmatrix}$ would not have a smaller slope than λ_i).
- Now, for vector $\begin{pmatrix} a \\ b \end{pmatrix}$, define $f\left(\begin{pmatrix} a \\ b \end{pmatrix}\right) = \frac{1-a}{b}$. Notice that for $(1, y) \in \Omega_1$, in order for the image of $\begin{pmatrix} a \\ b \end{pmatrix}$ to have horizontal component at most 1, y must be at most $f\left(\begin{pmatrix} a \\ b \end{pmatrix}\right)$. Thus we can invoke another condition, **Condition 3**, that for a candidate vector v , $f(v)$ must be greater than the lower bound for y in the region $P_i \cap \{x = 1\}$, which is simply $f(\lambda_{i+1})$. So this condition can be simplified to $a - b \cdot f(\lambda_{i+1}) < 1$. Notice that the above constraint can be rephrased as follows, with $m = b/a$ being the slope of a given vector

$$a(1 - m \cdot f(\lambda_{i+1})) < 1$$

So, if v and v' are vectors such that v' has a larger horizontal length a and a smaller slope m than v , then if v does not satisfy the above constraint, neither will v' .

- **Condition 4** is the restriction that $a/b < 2 + 2 \cos(\pi/n)$ (which is incidentally the aspect ratio of the rectangle formed by merging H_{i+1} and V_i for all i). This is because for $(1, y) \in \Omega_1$, $y > 1 - (2 + 2 \cos(\pi/n))$, and for any saddle connection $\begin{pmatrix} a \\ b \end{pmatrix}$ of S' , $1/b \leq 1$, so we can rearrange the previously-mentioned constraint $a + by \leq 1$ as follows:

$$1 - a/b \geq 1/b - a/b \geq y > -1 - 2 \cos(\pi/n)$$

The inequality $a/b < 2 + 2 \cos(\pi/n)$ directly follows from the above inequality.

Lemma 6.2. $f(\lambda_i) < f(\lambda_{i+1})$

Proof. We first show $f(\nu_0) > f\begin{pmatrix} 0 \\ 1 \end{pmatrix}$:

$$f(\nu_0) - f\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \frac{h_0 - 1}{v_0} - \frac{0 - 1}{1} = \frac{1 - 1}{1} + 1 = 1 > 0$$

Now we show $f(\nu_{i+1}) > f(\nu_i)$:

$$\begin{aligned} f(\nu_{i+1}) - f(\nu_i) &= \frac{h_{i+1} - 1}{v_{i+1}} - \frac{h_i - 1}{v_i} = \\ &= \frac{v_i^2 + v_i v_{i+1} - v_i - v_i v_{i+1} - v_{i-1} v_{i+1} + v_{i+1}}{v_i v_{i+1}} = \frac{1 + v_{i+1} - v_i}{v_i v_{i+1}} > 0 \end{aligned}$$

Now we show $f(\sigma_{i+1}) > f(\nu_i)$:

$$f(\sigma_{i+1}) - f(\nu_i) = \frac{h_{i+1} - 1}{v_i} - \frac{h_i - 1}{v_i} = \frac{h_{i+1} - h_i}{v_i} > 0$$

Finally we show $f(\sigma_i) > f(\sigma_{i+1})$:

$$\begin{aligned} f(\sigma_i) - f(\sigma_{i+1}) &= \frac{h_i - 1}{v_{i-1}} - \frac{h_{i+1} - 1}{v_i} = \frac{v_i(v_{i-1} + v_i - 1) - v_{i-1}(v_i + v_{i+1} - 1)}{v_{i-1}v_i} \\ &= \frac{v_i^2 - v_{i-1}v_{i+1} + v_{i-1} - v_i}{v_{i-1}v_i} = \frac{1 + v_{i-1} - v_i}{v_{i-1}v_i} > 0 \end{aligned}$$

□

Lemma 6.3.

$$\text{Slope}(\lambda_2) > \text{Slope}(\lambda_3) > \dots > \text{Slope}(\lambda_n)$$

Proof. First we show that for $0 \leq i < \lfloor n/2 \rfloor - 1$, $\text{Slope}(\nu_i) > \text{Slope}(\nu_{i+1})$. By direct computation,

$$\text{Slope}(\nu_i) = \frac{1}{\sin\left(\frac{\pi i}{n}\right) \csc\left(\frac{\pi(i+1)}{n}\right) + 1},$$

so we show that this is a decreasing function of i . Indeed, computing the derivative:

$$\frac{d}{di} \text{Slope}(\nu_i) = -\frac{\pi \sin\left(\frac{\pi}{n}\right)}{n \left(\sin\left(\frac{\pi i}{n}\right) + \sin\left(\frac{\pi(i+1)}{n}\right) \right)^2} < 0.$$

Since $\lambda_{i+1} = \nu_i$ for $0 \leq i \leq \lfloor n/2 \rfloor - 1$, we've shown the inequality $\text{Slope}(\lambda_j) > \text{Slope}(\lambda_{j+1})$ for $j = 2, \dots, \lfloor n/2 \rfloor$.

Now we show that $\text{Slope}(\lambda_{\lfloor n/2 \rfloor + 1}) > \text{Slope}(\lambda_{\lfloor n/2 \rfloor + 2})$. For even n , we must show that $\text{Slope}(\nu_{n/2-1}) > \text{Slope}(\sigma_{n/2-1})$. The horizontal component of both of these vectors is $h_{n/2-1}$. However, the vertical component of $\nu_{n/2-1}$ is $v_{n/2-1}$, which is greater than $v_{n/2-2}$, the vertical component of $\sigma_{n/2-1}$. This proves the claim for even n . For odd n , we must show that $\text{Slope}(\nu_{(n-3)/2}) > \text{Slope}(\sigma_{(n-1)/2})$. The

vertical components of both of these vectors is $v_{(n-3)/2}$, so it suffices to compare the horizontal components. The horizontal component of $\nu_{(n-3)/2}$ is $h_{(n-3)/2}$, which is less than $h_{(n-1)/2}$, the horizontal component of $\sigma_{(n-1)/2}$. This proves the inequality for odd n .

Finally we show that for $1 \leq i < \lceil n/2 \rceil - 1$, $\text{Slope}(\sigma_{i+1}) > \text{Slope}(\sigma_i)$. By direct computation,

$$\text{Slope}(\sigma_i) = \frac{1}{\sin\left(\frac{\pi(i+1)}{n}\right) \csc\left(\frac{\pi i}{n}\right) + 1}$$

so we show that this is an increasing function of i . Indeed, computing the derivative:

$$\frac{d}{di} \text{Slope}(\sigma_i) = \frac{\pi \sin\left(\frac{\pi}{n}\right)}{n \left(\sin\left(\frac{\pi i}{n}\right) + \sin\left(\frac{\pi(i+1)}{n}\right) \right)^2} > 0.$$

Since $\lambda_{n+1-i} = \sigma_i$ for $1 \leq i < \lceil n/2 \rceil - 1$, we've shown the inequality $\text{Slope}(\lambda_j) > \text{Slope}(\lambda_{j+1})$ for $\lceil n/2 \rceil + 1 \leq j < n$.

Together, these inequalities are equivalent to the lemma. \square

Now, with the above conditions and lemmas in mind, we begin casework.

Case 1. First we will prove $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ wins on $x = 1, 0 < y \leq 1$.

Well, in this case, condition 3 stipulates that $a < 1$; since each horizontal length in S' is at least 1, the only vectors satisfying this constraint are vertical ones. These have the same slope, as do their images when multiplied by matrices $M_{x,y}$, so we simply choose $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ as the winning vector (any choice of vertical saddle connection on S' would be valid). Note that for $x = 1, 0 < y \leq 1$, $M_{x,y} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} y \\ 1 \end{pmatrix}$, so the image has positive horizontal length at most 1 as desired.

Before preceding to the other cases, we will label vertices as follows: the concave vertex on the upper left side where the sides of length v_k and h_{k-1} meet will be labeled L_k , and the concave vertex on the lower right side where the sides of length v_{k-1} and h_k meet will be labeled R_k , as demonstrated in the diagram below. Under this labeling, note that ν_i joins L_i and L_{i+1} .




FIGURE 23. An example of vertex labelling on S' for $n = 7$.

Case 2. We now prove that ν_i wins on $P_{i+2} \cap \{x = 1\}$ for $0 \leq i \leq \lfloor n/2 \rfloor - 1$.

We begin by fixing i . Recall that as a holonomy vector, $v_i = \begin{pmatrix} h_i \\ v_i \end{pmatrix}$. We first compute the bound q for condition 1 in this case. Suppose $\nu' = \begin{pmatrix} a \\ b \end{pmatrix}$ is a vector whose image has horizontal component at most 1 and slope less than the image of ν_i . In order for the image of ν' to have horizontal component at most 1, it must be true that $a + by \leq 1$. Because ν' is a saddle connection on the staircase shape, which has horizontal and vertical distances ≥ 1 , it must be the case that $a \geq 1$ and $b \geq 1$. Furthermore, since the matrices in the transversal preserve the ordering of slopes, in order for the image ν' to have smaller slope than the image of ν_i , it must be true that $b/a < v_i/h_i$ (where the latter quantity is the slope of ν_i). Combining these facts, we find

$$\begin{aligned} a \leq 1 - by < 1 - \frac{v_i}{h_i}ay < 1 + \frac{v_i(h_{i+1} - 1)}{h_i v_{i+1}}a &\implies \\ a \left(1 - \frac{v_i(h_{i+1} - 1)}{h_i v_{i+1}}\right) = a \left(\frac{v_i - 1}{v_i v_{i+1} + v_{i-1} v_{i+1}}\right) < 1 &\implies \\ a < \frac{v_i v_{i+1} + v_{i-1} v_{i+1}}{v_i - 1} = 1 + h_{i+1} + \frac{v_{i+1}}{v_i - 1}. \end{aligned}$$

Now, note that $\frac{v_{i+1}}{v_i - 1}$ varies inversely with i :

$$\frac{d}{di} \left(\frac{v_{i+1}}{v_i - 1} \right) = -\frac{\pi \sin(\pi/n)(1 + \cos((2+i)\pi/n))}{n(\sin((1+i)\pi/n) - \sin(\pi/n))^2} < 0.$$

Thus, since $\frac{v_{i+1}}{v_i - 1}$ decreases as i increases, we find

$$a < 1 + h_{i+1} + \frac{v_{i+1}}{v_i - 1} \leq 1 + h_{i+1} + \frac{v_2}{v_1 - 1} = 1 + h_1 + h_{i+1}$$

Thus, our bound q for **Condition 1** will be $1 + h_1 + h_{i+1}$.

Subcase 2.1. We first rule out all vectors that do not pass through an edge of the staircase; that is, we will show that any vector other than ν_i that does not pass through an edge of the staircase fails to meet at least one of the conditions. Recall the labeling of \mathcal{S}' with an example given in Figure 23. Well, all σ_j and the ν_j with $j > i$ violate condition 3 by the ordering of the $f(\lambda_j)$ s proven in an earlier lemma, and for $j < i$, ν_j has larger slope than ν_i by an earlier lemma, so condition 2 is violated.

Subsubcase 2.1.1. Now consider vectors linking L_j and L_k for some $k > j + 1$ (we don't need to consider $k = j + 1$ because the vectors linking L_j and L_{j+1} are precisely the set of ν_j). If $k \leq i + 1$, we note that $\begin{pmatrix} a \\ b \end{pmatrix}$ will be strictly steeper than ν_k and hence violates condition 2; this is because $\begin{pmatrix} a \\ b \end{pmatrix}$ would have to pass to the right of L_{k-1} and hence would be steeper than ν_{k-1} (since ν_{k-1} joins L_{k-1} and L_k). Since ν_{k-1} is at least as steep as ν_i for $k \leq i + 1$, condition 2 is violated.

Now suppose $k > i + 1$. It must be true that $j \geq k - 2$, otherwise condition 1 is violated:

$$a \geq h_k + h_{k-1} + h_{k-2} \geq h_{i+1} + h_1 + 1$$

If $k > i + 2$, then $j > i$, meaning $\begin{pmatrix} a \\ b \end{pmatrix}$ has larger a and smaller slope than ν_j , which we showed earlier violates condition 3, and hence $\begin{pmatrix} a \\ b \end{pmatrix}$ violates condition 3 as well. This leaves the case of

$k = i + 2$, meaning $j = i$ is forced. The holonomy vector here is $\begin{pmatrix} h_i + h_{i+1} \\ v_i + v_{i+1} \end{pmatrix}$ and can be shown




FIGURE 24. If fix $i = 2$ in this case, we see that any vector linking L_j and L_k (black vectors) with $k \leq i + 1$ while staying in the staircase is either steeper than ν_1 (green vector) or ν_2 (yellow vector).

(using some computational assistance) to violate condition 3:

$$\begin{aligned}
 h_i + h_{i+1} - (v_{i+1} + v_i) \left(\frac{h_{i+1} - 1}{v_{i+1}} \right) &= 1 + h_i - v_i \left(\frac{h_{i+1} - 1}{v_{i+1}} \right) = \\
 1 + v_{i-1} + v_i - v_i \left(\frac{v_i + v_{i+1} - 1}{v_{i+1}} \right) &= 1 + \frac{v_i - (v_i^2 - v_{i-1}v_{i+1})}{v_{i+1}} = \\
 &= 1 + \frac{v_i - 1}{v_{i+1}} \geq 1
 \end{aligned}$$

Thus, all vectors connecting an L_j to another L_k have been eliminated.

Subsubcase 2.1.2. Now consider the vectors linking some R_j to some L_k . If $k \leq i + 1$, then condition 1 is violated, for the same reasons that we mentioned in the $k \leq i + 1$ case for vectors connecting L_j and L_k .




FIGURE 25. If fix $i = 2$ in this case, we see that any vector linking L_j and R_k with $k \leq i + 1$ while staying in the staircase (black vectors) is either steeper than ν_1 (green vector) or ν_2 (yellow vector).

So suppose $k > i + 1$. If $j < k - 3$ then condition 1 is violated:

$$a \geq h_{k-1} + h_{k-2} + h_{k-3} \geq h_{i+1} + h_1 + 1.$$

If $j = k - 2$, then condition 2 is violated:

$$b/a = (v_{k-1} + v_{k-2})/h_{k-1} = 1 \geq v_i/h_i$$

Since it must be true that $j < k - 1$ for $\begin{pmatrix} a \\ b \end{pmatrix}$ to have positive slope, we are left with $j = k - 3$, $j \geq i - 1$. Any of these vectors violates condition 2:

$$\begin{aligned} b/a &= \frac{v_j + v_{j+1} + v_{j+2}}{h_{j+1} + h_{j+2}} > \frac{v_j + v_{j+1} + v_{j+2}}{h_j + h_{j+1} + h_{j+2}} = \\ &= \frac{(v_j + v_{j+1} + v_{j+2})(v_j + v_{j+1})}{(h_j + h_{j+1} + h_{j+2})h_{j+1}} = \frac{v_j^2 + 2v_jv_{j+1} + v_{j+1}^2 + v_jv_{j+2} + v_{j+1}v_{j+2}}{(h_j + h_{j+1} + h_{j+2})h_{j+1}} = \\ &= \frac{v_j^2 - v_{j-1}v_{j+1} - v_{j+1}^2 + v_jv_{j+2} + v_{j+1}^2 + 2v_jv_{j+1} + v_{j+1}^2 + v_{j-1}v_{j+1} + v_{j+1}v_{j+2}}{(h_j + h_{j+1} + h_{j+2})h_{j+1}} = \\ &= \frac{1 - 1 + v_{j-1}v_{j+1} + 2v_jv_{j+1} + 2v_{j+1}^2 + v_{j+1}v_{j+2}}{(h_j + h_{j+1} + h_{j+2})h_{j+1}} = \\ &= \frac{(h_j + h_{j+1} + h_{j+2})v_{j+1}}{(h_j + h_{j+1} + h_{j+2})h_{j+1}} = v_{j+1}/h_{j+1} \geq v_i/h_i. \end{aligned}$$

Thus all vectors connecting an R_j to an L_k violate at least one condition.

Subsubcase 2.1.3. Now consider a saddle connection staying in the staircase connecting an R_j or L_j to some R_k . We showed previously that all σ_i 's violate condition 3. Given any k , a vector $\begin{pmatrix} a \\ b \end{pmatrix}$ terminating at R_k has a at least as large as the horizontal component of σ_k and slope that is shallower than the slope of σ_k ($\begin{pmatrix} a \\ b \end{pmatrix}$ passes into H_{k-1} above the starting point of σ_k and terminates at the same place as σ_k), so any such $\begin{pmatrix} a \\ b \end{pmatrix}$ violates condition 3. This eliminates all vectors staying in the staircase (i.e., vectors that do not pass through an edge).




FIGURE 26. For the example of $k = 2$, we see that any vector connecting some R_j or L_j (in this case L_0) and R_2 will pass above R_1 and hence be longer and shallower than σ_2 .

Subcase 2.2. We now rule out vectors that pass through the edges of the staircase.

Subsubcase 2.2.1. If the first edge a vector $\begin{pmatrix} a \\ b \end{pmatrix}$ passes through is a vertical edge of length v_k , then $\begin{pmatrix} a \\ b \end{pmatrix}$ is longer and shallower than σ_{k+1} , and so $\begin{pmatrix} a \\ b \end{pmatrix}$ violates condition 3.




FIGURE 27. For the example of $k = 1$, we see that any vector passing through v_1 is longer and shallower than σ_2 .

Subsubcase 2.2.2. Thus we are left with vectors that first pass through a horizontal edge h_k . Note that in this case, the vector must travel a horizontal distance of at least h_k . If $k \leq i$, then $\begin{pmatrix} a \\ b \end{pmatrix}$ is steeper than ν_k , which we showed is at least as steep as ν_i , so condition 2 is violated.




FIGURE 28. If fix $k = 1$ and $i = 2$ in this case, we see that any vector passing through the side of length h_1 is steeper than ν_1 (green vector), which in turn is steeper than ν_2 .

Given that $k > i$, $\begin{pmatrix} a \\ b \end{pmatrix}$ can travel at most two horizontal distances (with a singular exception), otherwise condition 1 is violated for $i > 0$; more precisely, if $\begin{pmatrix} a \\ b \end{pmatrix}$ travelled three horizontal distances, note that a must be at least $h_k + h_{k+1} + h_l$ for some distance h_l . This is because the vector must traverse h_k if it crosses through a horizontal edge with length h_k , and moreover, since the vector must have slope < 1 to satisfy condition 2, it must pass through the vertical edge of length v_{k-1} after passing through the horizontal edge of length h_k , then traverse horizontal length h_{k-1} as well. In this case, condition 1 is violated:

$$a \geq h_k + h_{k-1} + h_l \geq h_{i+1} + h_i + h_l \geq h_{i+1} + h_1 + 1$$

Thus, we are left with vectors $\begin{pmatrix} a \\ b \end{pmatrix}$ passing through at most 2 horizontal distances, with a singular exception. In the special case of $i = 0$, there is one particular vector of this form that does not violate condition 1 that we must check.

This is the vector beginning at L_0 , passing through the horizontal face just to the left of L_2 , passing through the vertical face below R_1 , then ending at L_1 . This vector is $\begin{pmatrix} 2 + h_1 \\ 2 + v_1 \end{pmatrix}$, so condition 3 is violated:

$$a - b \left(\frac{h_1 - 1}{v_1} \right) = 2 + h_1 - (2 + v_1) = 1 \geq 1.$$

Now we eliminate all other vectors passing through a horizontal side of length h_k . Note that $\begin{pmatrix} a \\ b \end{pmatrix}$ cannot travel only h_k horizontally, because if $\begin{pmatrix} a \\ b \end{pmatrix}$ first passes through a horizontal side h_k , then it must travel at least $v_k + v_{k-1} = h_k$ vertically, so having horizontal length h_k would mean having a slope of at least 1, violating condition 2. Thus, $\begin{pmatrix} a \\ b \end{pmatrix}$ must then pass through the vertical side of length v_{k-1} and then horizontally travel length h_{k-1} . This is only possible if $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at L_k or R_{k-1} .

Suppose $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at R_{k-1} . As it travels the horizontal length of h_k , it must climb vertically at least $v_{k-1} + v_k = h_k$ in order to pass through the horizontal edge of length h_k . As we found before, this violates condition 2.

So we are left with $\begin{pmatrix} a \\ b \end{pmatrix}$ beginning at L_k and passing through the horizontal edge of length h_k for $k > i$. After passing through h_k , the vector must pass through the vertical edge of length v_{k-1} as stated earlier. Upon passing through v_{k-1} , $\begin{pmatrix} a \\ b \end{pmatrix}$ can only end at L_k or pass through the horizontal edge of length h_{k-1} and end at R_{k-1} without travelling horizontally more than the width of two rectangles or having slope at least 1. If the saddle connection ends at L_k , it is the vector $(h_{k-1} + h_k, v_{k-1} + v_k)$, the same vector as the saddle connection connecting L_{k-1} and L_{k+1} without passing through edges, which we have previously eliminated. If the saddle connection ends at R_{k-1} , it is the vector $(h_{k-1} + h_k, v_{k-1} + v_{k-1} + v_k)$, the same vector as the saddle connection connecting R_{k-2} and L_{k+1} without passing through edges, which we have previously eliminated. $\begin{pmatrix} a \\ b \end{pmatrix}$ cannot end at L_k .

Subcase 2.3. We have now eliminated all vectors passing through edges of the staircase, with a few exceptions.

The above proof assumes the staircase extends for an arbitrary number of rectangles in either direction from the rectangle containing v_i , so in order to complete this proof we must eliminate vectors that reach the ends of the staircase shape.

Subsubcase 2.3.1. The vectors that pass through the edge between L_0 and R_0 will first be eliminated. For $\begin{pmatrix} a \\ b \end{pmatrix}$ passing through the edge between L_0 and R_0 , consider where it intersects the edge of the staircase or a vertex. If $\begin{pmatrix} a \\ b \end{pmatrix}$ ends at some R_k or passes through a vertical edge below some R_k , then $\begin{pmatrix} a \\ b \end{pmatrix}$ is both longer and less steep than σ_k , which violates condition 3, so

$\begin{pmatrix} a \\ b \end{pmatrix}$ also violates condition 3. If $\begin{pmatrix} a \\ b \end{pmatrix}$ travels the length of the staircase to the edge at the opposite end of the staircase, then $a \geq 1 + h_1 + h_i$, so condition 1 is violated. So $\begin{pmatrix} a \\ b \end{pmatrix}$ must end at some L_k or pass through a horizontal edge to the left of some L_k . Well, if $k \leq i + 1$ then $\begin{pmatrix} a \\ b \end{pmatrix}$ is steeper than ν_{k-1} , which itself is at least as steep as ν_i , so condition 2 is violated. If $k > i + 1$ then $a \geq 1 + h_1 + h_{k-1} \geq 1 + h_1 + h_{i+1}$, so condition 1 is violated.

Subsubcase 2.3.2. For even n , we will eliminate the vectors passing through the vertical edge below $L_{n/2}$, which forms an edge of the staircase when n is even. Let $k = n/2 - 1$. Notice that such a vector necessarily crosses h_k at least twice. Thus, such a vector cannot travel more than 2 horizontal distances or else it would violate condition 1. In fact, such a vector must travel a horizontal distance of exactly $2h_k$.

We first suppose that $n \geq 6$. The distance travelled by this vector is $2h_k$, so it suffices to show that $h_k \geq 1 + h_1$ in order to violate condition 1. We compute

$$h_k - 1 - h_1 = -2 \cos\left(\frac{\pi}{n}\right) + \cot\left(\frac{\pi}{2n}\right) - 2.$$

The derivative of this function is

$$\frac{\pi \left(\csc^2\left(\frac{\pi}{2n}\right) - 4 \sin\left(\frac{\pi}{n}\right) \right)}{2n^2},$$

which is positive for $n \geq 3$. At $n = 6$, this difference evaluates to 0, meaning that the difference is indeed nonnegative for all $n \geq 6$. In other words, condition 1 is violated for $n \geq 6$.

Let $n = 4$. The only possible ν_i vector in this case is ν_0 , since we address the last ν_i as an entirely separate case from the other ν_i s. Thus condition 3 is

$$a - b \left(\frac{h_1 - 1}{v_1} \right) = a - b \leq 1$$

Knowing $a = 2h_1$, b can be v_1 , $1 + v_1$, or $2 + v_1$ without reaching a slope of at least that of ν_1 . So,

$$a - b \geq 2h_1 - (2 + v_1) = v_1 > 1$$

Thus condition 3 is violated. Recall that we do not consider the $n = 2$ case in this paper, so casework is complete for even n .

Subsubcase 2.3.3. Let n be odd. In this case we must eliminate the vectors passing through the horizontal edge to the right of $L_{\lfloor n/2 \rfloor}$. Let $k = (n - 3)/2$. This vector cannot travel more than 2 horizontal distances, otherwise we would have $b \geq h_{k+1} + h_k + h_{k-1} \geq 1 + h_1 + h_i$, which violates condition 1. So $\begin{pmatrix} a \\ b \end{pmatrix}$ must start at L_k , R_{k-1} or R_k . Any $\begin{pmatrix} a \\ b \end{pmatrix}$ that begins at L_k and passes through the horizontal face before any other faces will have larger a and smaller slope than ν_k , meaning it violates condition 3.

Otherwise, $\begin{pmatrix} a \\ b \end{pmatrix}$ must pass through the horizontal face to the left of L_{k+1} then travel above R_k , meaning that over a horizontal distance of h_k , it travels vertically at least $v_k + v_{k-1} = h_k$, so condition 2 is violated.

Now suppose $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at R_{k-1} . The only way for $\begin{pmatrix} a \\ b \end{pmatrix}$ to avoid violating condition 1 is by ending at R_{k+1} after passing through the horizontal edge once. For $n > 5$, this violates condition

1, as $h_1 + 1 \leq h_{k+1}$:

$$h_{k+1} - h_1 - 1 = \left(\sin\left(\frac{\pi}{2n}\right) + \sin\left(\frac{3\pi}{2n}\right) - 1 \right) \left(-\csc\left(\frac{\pi}{2n}\right) \right) > 0.$$

For $n = 5$, this violates condition 3:

$$f((h_1 + h_2, 1 + 2v_1)) = \frac{1 - h_1 - h_2}{1 + 2v_1} \frac{-3\phi}{1 + 2\phi} < -1 = \frac{1 - h_1}{v_1} = f(\nu_1)$$

For $n = 3$, this saddle connection does not exist.

In the last case, where $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at R_k , it can end at either R_{k+1} , R_k , or L_{k+1} after crossing the right edge precisely once, or else it would violate condition 1. If $\begin{pmatrix} a \\ b \end{pmatrix}$ ends at R_{k+1} , then from direct computation (the lower bound of 1 is the slope when the vector crosses the edge exactly once), this vector has slope at least 1, which is definitely too steep.

If $\begin{pmatrix} a \\ b \end{pmatrix}$ ends at L_{k+1} , then it has smaller slope and larger horizontal component than ν_k (since it would pass through the vertical edge above L_k) and thus violates condition 3.

The only other possible vector that does not travel horizontally across more than two rectangles is ν' starting at R_k , passing through the horizontal edge to the right of L_{k+1} , passing through the vertical edge below R_{k+1} , passing through the horizontal edge to the left of L_{k+1} , then ending at R_k . This is the same vector as the one beginning at R_{k-1} , passing through the horizontal edge to the left of L_{k+1} , and ending at R_{k+1} , which we have eliminated. This eliminates all vectors other than ν_i for **Case 2**.

Furthermore, ν_i has positive slope v_i/h_i and is such that $0 < a + by \leq 1$:

$$\begin{aligned} a + by &= h_i + v_i y > h_i + v_i \cdot \frac{1 - h_{i+1}}{v_{i+1}} = \frac{h_i v_{i+1} + v_i - h_{i+1} v_i}{v_{i+1}} = \\ &= \frac{v_{i-1} v_{i+1} + v_i v_{i+1} + v_i - v_i^2 - v_i v_{i+1}}{v_{i+1}} = \frac{v_{i-1} v_{i+1} + v_i - v_i^2}{v_{i+1}} = \frac{v_i - 1}{v_{i+1}} \geq 0 \\ a + by &= h_i + v_i y \leq h_i + v_i \cdot \frac{1 - h_i}{v_i} = 1 \end{aligned}$$

Thus we have shown that for ν_i with $0 \leq i < \lfloor n/2 \rfloor - 1$, ν_i is the saddle connection with the smallest slope whose image has horizontal component at most 1 for all $(1, y) \in \Omega_1$ with $\frac{1 - h_{i+1}}{v_{i+1}} < y \leq \frac{1 - h_i}{v_i}$, which is simply $P_{i+2} \cap \{x = 1\}$.

Case 3.

We now prove $\nu_{\lfloor n/2 \rfloor - 1}$ has the smallest slope of any saddle connection whose image has horizontal component at most 1 for all $(1, y) \in \Omega_1$ satisfying $\frac{1 - h_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 2}} < y \leq \frac{1 - h_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 1}}$, which is simply the region $P_{\lfloor n/2 \rfloor + 1} \cap \{x = 1\}$. Let $\begin{pmatrix} a \\ b \end{pmatrix}$ be a hypothetical better vector.

We start by computing the bound q for Condition 1.

We have two cases. First suppose that n is even. Note that $n \geq 4$, as we exclude the $n = 2$ case from this paper. The bounds on y demand that $y > \frac{1 - h_{\lfloor n/2 \rfloor - 1}}{v_{n/2 - 2}}$. Then, using the fact that $a \geq 1$ and

$a + by \leq 1$, we find:

$$a \leq 1 - by < 1 - \frac{v_{\lfloor n/2 \rfloor - 1}}{h_{\lfloor n/2 \rfloor - 1}} ay < 1 - \frac{v_{\lfloor n/2 \rfloor - 1}}{h_{\lfloor n/2 \rfloor - 1}} \left(\frac{1 - h_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 2}} \right) = 1 - \frac{v_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 2}} \left(\frac{1 - h_{\lfloor n/2 \rfloor - 1}}{h_{\lfloor n/2 \rfloor - 1}} \right)$$

$$a \left(1 + \frac{v_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 2}} \left(\frac{1 - h_{\lfloor n/2 \rfloor - 1}}{h_{\lfloor n/2 \rfloor - 1}} \right) \right) < 1$$

$$a \left(h_{\lfloor n/2 \rfloor - 1} + \frac{v_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 2}} (1 - h_{\lfloor n/2 \rfloor - 1}) \right) = a \left(\frac{\cos(\pi/(2n)) - \sin(\pi/(2n))}{\cos(\pi/(2n)) + \sin(\pi/(2n))} \right) < h_{\lfloor n/2 \rfloor - 1}$$

For $n \geq 4$, $\cos(\pi/(2n)) - \sin(\pi/(2n)) > 0$, so we can safely rearrange:

$$a < h_{\lfloor n/2 \rfloor - 1} \left(\frac{\cos(\pi/(2n)) + \sin(\pi/(2n))}{\cos(\pi/(2n)) - \sin(\pi/(2n))} \right) = h_{\lfloor n/2 \rfloor - 1} \left(1 + \frac{2}{\cot(\pi/(2n)) - 1} \right)$$

Now, we claim that

$$\frac{2h_{\lfloor n/2 \rfloor - 1}}{\cot(\pi/(2n)) - 1} \leq 1 + h_1.$$

We prove this by showing that

$$\frac{2h_{\lfloor n/2 \rfloor - 1}}{\cot(\pi/(2n)) - 1} - (1 + h_1) \leq 0.$$

To do so, we first see that when $n = 4$, this expression is 0. Moreover, taking the derivative with respect to n gives us

$$\frac{\pi \left(2 \sin\left(\frac{\pi}{n}\right) + \cos\left(\frac{2\pi}{n}\right) \right)}{n^2 \left(\sin\left(\frac{\pi}{n}\right) - 1 \right)},$$

which is indeed always negative for $n \geq 4$, meaning that the inequality is indeed true. This means that our bound becomes

$$a < h_{\lfloor n/2 \rfloor - 1} \left(1 + \frac{2}{\cot(\pi/(2n)) - 1} \right) \leq 1 + h_1 + h_{\lfloor n/2 \rfloor - 1}.$$

Now suppose that n is odd and greater than 3 (we will consider the case $n = 3$ later). Then, $y > \frac{1 - h_{\lfloor n/2 \rfloor}}{v_{\lfloor n/2 \rfloor - 1}}$. Combining this with previously-mentioned facts, we find:

$$a \leq 1 - by < 1 - \frac{v_{\lfloor n/2 \rfloor - 1}}{h_{\lfloor n/2 \rfloor - 1}} ay < 1 - \frac{v_{\lfloor n/2 \rfloor - 1}}{h_{\lfloor n/2 \rfloor - 1}} \left(\frac{1 - h_{\lfloor n/2 \rfloor}}{v_{\lfloor n/2 \rfloor - 1}} \right) a = 1 - a \left(\frac{1 - h_{\lfloor n/2 \rfloor}}{h_{\lfloor n/2 \rfloor - 1}} \right)$$

$$a \left(1 + \frac{1 - h_{\lfloor n/2 \rfloor}}{h_{\lfloor n/2 \rfloor - 1}} \right) = a \left(\frac{h_{\lfloor n/2 \rfloor - 1} - h_{\lfloor n/2 \rfloor} + 1}{h_{\lfloor n/2 \rfloor - 1}} \right) = a \left(\frac{1 - 2 \cos((\lfloor n/2 \rfloor)\pi/n)}{h_{\lfloor n/2 \rfloor - 1}} \right) < 1$$

$$a \left(\frac{1 - 2 \cos((n-1)\pi/(2n))}{h_{\lfloor n/2 \rfloor - 1}} \right) < 1.$$

We must now verify that

$$1 - 2 \cos((n-1)\pi/(2n)) > 0.$$

Well, for $n \geq 5$, $(n-1)/(2n) > 1/3$, so $2 \cos((n-1)\pi/(2n)) < 2 \cos(\pi/3) = 1$. This implies the desired inequality. We rearrange:

$$a < \frac{h_{\lfloor n/2 \rfloor - 1}}{1 - 2 \cos((n-1)\pi/(2n))} = \frac{h_{\lfloor n/2 \rfloor - 1}}{1 - 2 \sin(\pi/(2n))}.$$

We find that in fact,

$$(1 + h_1 + h_{\lfloor n/2 \rfloor}) - \frac{h_{\lfloor n/2 \rfloor - 1}}{1 - 2 \sin(\pi/(2n))} = \frac{1}{2 \sin\left(\frac{\pi}{2n}\right) - 1} + 2 \cos\left(\frac{\pi}{n}\right) + 1 \geq 0,$$

meaning that our inequality becomes

$$a < \frac{h_{\lfloor n/2 \rfloor - 1}}{1 - 2 \sin(\pi/(2n))} \leq 1 + h_1 + h_{\lfloor n/2 \rfloor}.$$

Thus for all $n > 3$, we have that $a < 1 + h_1 + h_{\lfloor n/2 \rfloor}$ for odd n , and $a < 1 + h_1 + h_{\lfloor n/2 \rfloor - 1}$ for even n ; this is our bound q for Condition 1 in this case.

By an earlier proposition on the ordering of the saddle connections by slope, we note that the vectors ν_j for all $j < \lfloor n/2 \rfloor - 1$ are steeper than $\nu_{\lfloor n/2 \rfloor - 1}$, meaning that they all violate Condition 2. Moreover, we know from an earlier lemma that all σ vectors fail condition 3, because we are comparing to a ν vector.

Subcase 3.1. We begin with saddle connections staying in the staircase.

Subsubcase 3.1.1. We now consider saddle connections staying in the staircase linking any vertex to a vertex L_j for any j . Since these saddle connections stay entirely within the staircase, we note that they are necessarily steeper than the vector ν_j joining L_{j-1} and L_j (they enter the rectangle containing L_{j-1} and L_j at a point farther to the right than L_{j-1}). This means they violate condition 2 since ν_j is at least as steep as $\nu_{\lfloor n/2 \rfloor - 1}$.

Subsubcase 3.1.2. We proceed to saddle connections ending at R_k for any m . Condition 3 is violated, as this vector necessarily has larger slope and lower a value than σ_k .

Subcase 3.2. This rules out all saddle connections that do not pass through the edges of the staircase, so we now proceed to saddle connections that do pass through the edges of the staircase.

Subsubcase 3.2.1. Any saddle connection which first passes through a horizontal edge to the left of some L_k is necessarily steeper than ν_{k-1} , violating condition 2.

Subsubcase 3.2.2. If a saddle connection first passes through the vertical edge below some R_k for $k > 0$, then this vector necessarily has larger slope and lower a value than σ_k , meaning that it violates condition 3.

Subsubcase 3.2.3. The saddle connections that first pass through the horizontal edge between L_0 and R_0 will now be eliminated. The only possible starting vertex for such a saddle connection is L_0 ; in this case, the slope of the vector is higher than 1, meaning condition 2 is violated. Note that all other saddle connections which pass through the edge between L_0 and R_0 first pass through a different edge, and these have been eliminated in previous cases.

Subsubcase 3.2.4. If n is even, we must eliminate the vectors passing through the vertical edge below $L_{n/2}$. Notice that such a vector necessarily crosses $h_{n/2-1}$ at least twice. Thus, such a vector cannot travel more than 2 horizontal distances or else it would violate condition 1. In fact, such a vector must travel a horizontal distance of exactly $2h_{n/2-1}$.

We first suppose that $n \geq 6$. If we show that $h_{\lfloor n/2 \rfloor - 1} \geq 1 + h_1$, then this vector would violate condition 1. We compute

$$h_{\lfloor n/2 \rfloor - 1} - 1 - h_1 = -2 \cos\left(\frac{\pi}{n}\right) + \cot\left(\frac{\pi}{2n}\right) - 2.$$

The derivative of this function is

$$\frac{\pi \left(\csc^2\left(\frac{\pi}{2n}\right) - 4 \sin\left(\frac{\pi}{n}\right) \right)}{2n^2},$$

which is positive for $n \geq 3$. At $n = 6$, this difference evaluates to 0, meaning that the difference is indeed nonnegative for all $n \geq 6$. In other words, condition 1 is violated for $n \geq 6$, leaving us with the case $n = 4$.

Now consider when $n = 4$. In this case $\begin{pmatrix} a \\ b \end{pmatrix}$ must begin at R_0 or L_1 and end at either R_1 or L_2 .

If $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at R_0 , it must pass through the vertical face below L_2 and either (a) end at L_2 or (b) pass through the horizontal face to the left of L_2 and end at R_1 . In case (a), the vector is $(2h_1, 1 + v_1)$, which violates condition 3:

$$a - b(h_1 - 1) = 2 + 2v_1 - (1 + v_1)v_1 = 2 + v_1 - v_1^2 = \sqrt{2} > 1$$

In case (b), the vector is $\begin{pmatrix} 2h_1 \\ 2 + v_1 \end{pmatrix}$, which has slope $\frac{1+v_1/2}{h_1} > \frac{v_1}{h_1}$, which violates condition 2. If $\begin{pmatrix} a \\ b \end{pmatrix}$ instead begins at L_1 , then it must pass through the vertical edge below L_2 , and either (c) end at L_2 or (d) pass through the horizontal edge to the left of L_2 and end at R_1 . In case (c), the vector is $\begin{pmatrix} 2h_1 \\ v_1 \end{pmatrix}$ which has smaller slope and larger a than $\sigma_1 = \begin{pmatrix} h_1 \\ 1 \end{pmatrix}$, so it violates condition 3. The vector in case (d) is $\begin{pmatrix} 2h_1 \\ 1 + v_1 \end{pmatrix}$, which we already eliminated. This concludes the proof for $n = 4$ and hence for the case where n is even.

Subsubcase 3.2.5. Finally, we will eliminate the saddle connections passing through the horizontal edge to the right of $L_{\lfloor n/2 \rfloor}$, which forms an edge of the staircase when n is odd and greater than 3.

Such a saddle connection can travel no more than 2 horizontal distances, otherwise it will violate condition 1 (since one of these horizontal distances must be $h_{\lfloor n/2 \rfloor}$). So, letting $k = \lfloor n/2 \rfloor - 1$, a candidate saddle connection must start at L_k, R_{k-1} or R_k .

If $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at L_k , we observe that in order to avoid violating condition 3, it must end at R_{k+1} . The vector must therefore be $(h_k + h_{k+1}, 2v_k)$. For $n > 5$, this violates condition 1, as $h_1 + 1 \leq h_{\lfloor n/2 \rfloor}$:

$$h_{\lfloor n/2 \rfloor} - h_1 - 1 = \left(\sin\left(\frac{\pi}{2n}\right) + \sin\left(\frac{3\pi}{2n}\right) - 1 \right) \left(-\csc\left(\frac{\pi}{2n}\right) \right) > 0.$$

For $n = 5$, this vector violates condition 3:

$$a - b\left(\frac{h_{k+1} - 1}{v_k}\right) = h_k + h_{k+1} - 2v_k\left(\frac{h_{k+1} - 1}{v_k}\right) = 2 + h_k - h_{k+1} \geq 1$$

Now suppose $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at R_{k-1} . The only way for $\begin{pmatrix} a \\ b \end{pmatrix}$ to avoid violating condition 1 is by ending at R_{k+1} after passing through the horizontal edge at least once. For $n > 5$, this violates condition 1, as $h_1 + 1 \leq h_{\lfloor n/2 \rfloor}$. For $n \leq 5$, consider the vector crossing the edge once, $(h_k + h_{k+1}, v_{k-1} + 2v_k)$. For $n = 5$ this vector has slope greater than v_k/h_k , violating condition 2.

Since any $\begin{pmatrix} a \\ b \end{pmatrix}$ crossing the edge more than once will have even larger slope, this case also violates condition 2.

Finally, suppose that $\begin{pmatrix} a \\ b \end{pmatrix}$ begins at R_k . Then, it must end at R_{k+1}, L_{k+1} , or R_k or else it would violate condition 1. If it ends at R_{k+1} , then because the lower bound of 1 is the slope when the vector crosses the edge exactly once, this vector has slope at least 1, which violates condition 2.

The only possible $\begin{pmatrix} a \\ b \end{pmatrix}$ ending at L_{k+1} in this case is the vector passing once through the horizontal edge to the left of R_{k+1} , then through the vertical edge below R_{k+1} , before ending at L_{k+1} . This is the same vector as the saddle connection beginning at $L_{\lfloor n/2 \rfloor - 1}$, then passing through the horizontal edge to the right of L_{k+1} , and finally ending at R_{k+1} , which we have eliminated earlier.

The only other possible vector that does not travel horizontally across more than two rectangles starts at R_k , passes through the horizontal edge to the right of L_{k+1} , then passes through the vertical edge below R_{k+1} , then passes through the horizontal edge to the left of L_{k+1} , and finally ends at R_k . This is the same vector as the one beginning at R_{k-1} and ending at R_{k+1} discussed earlier, which we have eliminated. This completes the proof for all $n > 3$.

Subcase 3.3. Finally, suppose $n = 3$. We directly check that no other vectors work in the region corresponding to $\nu_{\lfloor n/2 \rfloor - 1}$. In this case, $h_{\lfloor n/2 \rfloor - 1} = h_0 = 1$, $v_{\lfloor n/2 \rfloor - 1} = v_0 = 1$, $y > -1$, and $h_1 = 2$. Suppose that a vector $\begin{pmatrix} a \\ b \end{pmatrix}$ did work. Then, we must have that $a + by \leq 1$. As $y > -1$ in this region, this gives us $a - b < 1$, or equivalently, $a < b + 1$. We must also have that $a > b$, or else the slope of the vector wouldn't be less than that of $\nu_{\lfloor n/2 \rfloor - 1}$. Thus, $b < a < b + 1$, which is a contradiction since all lengths and heights in the staircase when $n = 3$ are integers.

Thus we have shown that $\nu_{\lfloor n/2 \rfloor - 1}$ wins for all $(1, y) \in \Omega_1$ with $\frac{1 - h_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 2}} < y \leq \frac{1 - h_{\lfloor n/2 \rfloor - 1}}{v_{\lfloor n/2 \rfloor - 1}}$, which is simply $P_{\lfloor n/2 \rfloor + 1} \cap \{x = 1\}$.

Case 4. We now repeat the procedure, showing that the vectors σ_i win over $P_{n+1-i} \cap \{x = 1\}$. Begin by fixing i .

We first compute a bound q for Condition 1 in this case. In order for the image of some $\begin{pmatrix} a \\ b \end{pmatrix}$ to have horizontal length at most 1, we must have $a + by \leq 1$. We also have $a \geq 1$ and $b \geq 1$, as these are the shortest horizontal and vertical distances on the S' , respectively. Moreover, we must have that $b/a < v_{i-1}/h_i$, as this is the slope of the candidate vector for P_i . Finally, we have that $y > \frac{1 - h_{i-1}}{v_{i-2}}$. Combining these facts, we find

$$\begin{aligned} a \leq 1 - by &< 1 - \frac{v_{i-1}}{h_i} ay < 1 - \frac{v_{i-1}}{h_i} \left(\frac{1 - h_{i-1}}{v_{i-2}} \right) a \implies \\ &a \left(1 + \frac{v_{i-1}(1 - h_{i-1})}{h_i v_{i-2}} \right) < 1 \implies \\ a \left(h_i + \frac{v_{i-1}(1 - h_{i-1})}{v_{i-2}} \right) &= a \left(\frac{v_{i-1} - 1}{v_{i-2}} \right) < h_i \implies \\ a < h_i \left(\frac{v_{i-2}}{v_{i-1} - 1} \right) &= 1 + h_{i-1} + \frac{v_{i-2}}{v_{i-1} - 1} < 1 + h_{i-1} + \frac{v_i}{v_{i-1} - 1}. \end{aligned}$$

Well, in case 2 we showed

$$\frac{v_i}{v_{i-1} - 1} < h_1$$

So we can conclude that

$$a < 1 + h_1 + h_{i-1}$$

This gives us the desired bound q for **Condition 1**.

By an earlier proposition on the ordering of the saddle connections by slope, we note that vectors ν_k for all k and σ_k with $k < i$ are steeper than σ_i , violating condition 2. Moreover, our earlier lemma concerning the ordering of the values of the function f on the vectors implies that all σ_k with $k > i$ violate condition 3.

Subcase 4.1. We begin with saddle connections staying in the staircase.

Subsubcase 4.1.1. First consider any saddle connection staying in the staircase that ends at a vertex L_k for any k . Since these saddle connections stay entirely within the staircase, we note that they are necessarily steeper than the vector ν_k joining L_{k-1} and L_k (they enter the rectangle

containing L_{k-1} and L_k at a point farther to the right than L_{k-1}). This means they violate condition 2 as ν_k violates condition 2. This can be seen in Figure 29.




FIGURE 29. Vectors ending at L_k for any k (shown in black here) are necessarily steeper than the ν vector that ends at L_k (shown in green or yellow), which is in turn steeper than every σ vector.

Subsubcase 4.1.2. We now eliminate vectors staying in the staircase linking R_j and R_k (with $k > j + 1$). If $k < i$, then condition 3 is violated, as this vector is necessarily shallower and longer than the vector σ_k which also ends at R_k as seen in Figure 30. On the other hand, if $j \geq i$, then this saddle connection is necessarily steeper than the saddle connection σ_{j+1} starting at R_j and ending at R_{j+1} , as seen in Figure 31, meaning that these vectors violate condition 2.




FIGURE 30. Let $n = 7$ and $i = 3$ (so the vector we are trying to show victory for is σ_3). Vectors ending at R_k for any $k < i$ (shown in black here) are necessarily shallower and longer than the σ vector that ends at R_k (in orange), which in turn violates condition 3.

Finally, we check the case where $j < i$ and $k \geq i$. If $k - j \geq 3$, then this saddle connection must travel through at least h_k and two other horizontal lengths, which violates condition 1 (as one of these lengths must be at least 1 and the other length, being distinct from the first length, must be at least h_1). This implies that $k - j = 2$, so the only cases we need to consider are $k = i, j = i - 2$ and $k = i + 1, j = i - 1$. In the latter case, the vector is steeper than σ_i (it begins at the same vertex as σ_i but travels higher than R_i), violating condition 2.




FIGURE 31. Let $n = 7$ and $i = 2$ (so the vector we are trying to show victory for is σ_2). Vectors starting at R_j for any $j \leq i$ (shown in black here) are necessarily steeper than the σ vector that starts at R_j (in red), which in turn violates condition 2.

This leaves us with the former case. We show that in this case, $f\left(\frac{a}{b}\right) > 1$, violating condition 3. We have that

$$\begin{aligned}
& h_{i-1} + h_i - (v_{i-1} + v_{i-2}) \left(\frac{h_{i-1} - 1}{v_{i-2}} \right) = \\
& v_{i-2} + 2v_{i-1} + v_i - (v_{i-1} + v_{i-2}) \left(\frac{v_{i-2} + v_{i-1} - 1}{v_{i-2}} \right) = \\
& v_{i-2} + 2v_{i-1} + v_i - v_{i-2} - v_{i-1} + 1 - \frac{v_{i-2}v_{i-1} + v_{i-1}^2 - v_{i-1}}{v_{i-2}} = \\
& \frac{v_{i-2}v_{i-1} + v_{i-2}v_i + v_{i-2} - v_{i-2}v_{i-1} - v_{i-1}^2 + v_{i-1}}{v_{i-2}} = \\
& \frac{-1 + v_{i-1} + v_{i-2}}{v_{i-2}} = 1 + \frac{v_{i-1} - 1}{v_{i-2}} > 1,
\end{aligned}$$

as we know that $i \geq 2$, meaning $v_{i-1} > 1$. Thus we have eliminated all vectors connecting some R_j to some R_k without passing through an edge of S' .

Subsubcase 4.1.3. We now eliminate the vectors starting at L_j and ending at R_k for $k \geq j + 1$. If $k < i$, then condition 3 is violated as this vector is necessarily shallower and longer than σ_k , which also ends at R_k (which we noted earlier violates condition 3), as seen in Figure 32.

Suppose that $k \geq i$. If $k - j \geq 2$, this saddle connection violates condition 3 (as it encompasses at least three horizontal distances on the staircase, of which one is at least as long as the length h_i , and the other two of which are distinct). Thus, we must have $k - j = 1$, in which case the slope of the saddle connection is precisely the aspect ratio $2 + 2 \cos(\pi/n)$ (it will be the diagonal of some rectangle, as seen in Figure 33), which fails condition 4. This rules out all vectors that do not pass through the edges of the staircase.

Subcase 4.2. We now proceed to vectors that do pass through the edges of the staircase.

Subsubcase 4.2.1. Any of these vectors that starts by passing through the horizontal edge containing L_k is necessarily steeper than the ν vector ending at L_k , meaning these vectors violate condition 2, as in Figure 34.

Subsubcase 4.2.2. Now consider a vector that starts by passing through a right wall. Suppose the vector started at L_j and passed the vertical edge below R_{j+1} . Then, this vector would be shallower than the aspect ratio of the rectangle and can thus be eliminated on account of failing condition 4. If the vector starts at R_k or L_k with $k > i$ and travels across more than three horizontal




FIGURE 32. Let $n = 7$ and $i = 3$ (so the vector we are trying to show victory for is σ_3). Vectors starting at some L_j and ending at R_k for any $k < i$ (shown in black here) are necessarily shallower and longer than the σ vector that ends at R_k (in orange), which in turn violates condition 3.




FIGURE 33. Let $n = 7$. Vectors starting at L_j and ending at R_{j+1} are precisely the diagonals of the large rectangles, which have aspect ratio equal to $2 + 2 \cos(\pi/n)$.




FIGURE 34. For instance, any vector passing through the side of length h_1 is steeper than ν_1 (green vector), which in turn is steeper than every σ vector.

distances, then it violates condition 1, as one of the horizontal distances it traverses is at least as long as h_i .

Now suppose that the saddle connection first crosses the right edge below R_k for some $k < i$. Then, this saddle connection is longer and shallower than the σ vector in that rectangle, meaning that it violates condition 3, as in Figure 35.




FIGURE 35. Fix $i = 3$. For instance, we see that any vector passing through v_1 is longer and shallower than σ_2 , which fails condition 3.

Now suppose that the saddle connection first crosses the right edge below R_k for some $i \leq k$. Then, if the vector originates from some L_j , we know that the vector must have originated at L_{k-2} or before, or else it would have slope shallower than the aspect ratio. However, this vector must cross at least three horizontal distances and therefore violates condition 1. Thus, in this case, the vector must originate from some R_j . In particular, to avoid violating condition 1, it must originate from R_{k-1} or R_{k-2} . However, in either case, the vector must pass through the horizontal edge to the left of L_k after passing through the vertical edge to avoid violating condition 4. Thus if the vector begins at R_{k-2} , we have

$$a \geq h_{k-1} + h_k + h_{k-1} > 1 + h_1 + h_{i-1}$$

Thus condition 1 is violated. So we are left with vectors beginning at R_{k-1} . Such a vector may only travel 2 horizontal distances without violating condition 1, so it must end at either L_k or R_{k-1} upon passing through the horizontal edge to the left of L_k . If it were to end at L_k it would have slope greater than 1, violating condition 2. The only remaining option, the vector ending at R_{k-1} , is the same as the vector connecting R_{k-2} and R_k lying within the staircase, which we have previously eliminated.

Subcase 4.3. Now let us address the special cases of vectors passing through edges at the ends of the staircase.

Subsubcase 4.3.1. The vectors that pass through the edge between L_0 and R_0 will first be eliminated. Consider where such a vector next intersects an edge of S' or terminates at a vertex after passing through the aforementioned horizontal edge. Such vectors which next intersect a horizontal edge to the left of some L_k or terminate at some L_k have larger slope than ν_{k-1} , meaning they violate condition 2. Such vectors which next intersect a vertical edge below some R_k or end at some R_k for $k < i$ have larger a value and smaller slope than σ_{i-1} , meaning they violate condition 3. This leaves vectors which, after passing through the horizontal edge between L_0 and R_0 , intersect a vertical edge below some R_k with $k \geq i$ or reach the opposite end of S' . In this case, we have:

$$a \geq 1 + h_1 + h_k > 1 + h_1 + h_{i-1}$$

Thus condition 1 is violated.

Subsubcase 4.3.2. When n is even, we must consider the special case of saddle connections crossing the vertical edge below $L_{n/2}$. Let $k = n/2 - 1$. Notice that such a vector necessarily crosses $h_k > h_{i-1}$ at least twice. Such a vector cannot travel more than 2 horizontal distances or else it would violate condition 1.

Thus, such a vector must either start at R_{k-1} or L_k . If the vector starts at R_{k-1} , then it must be steeper than σ_{k-1} , which is at least as steep as σ_i , meaning such vectors fail condition 2. Now, suppose that the vector starts at L_k . Then the vector must end at L_{k+1} or R_k after traversing h_k

twice (steeper vectors have slope greater than 1, violating condition 2). In the first case, the slope of the vector is at most the aspect ratio, which means it violates condition 4. In the second case, we have:

$$b/a = \frac{v_k + v_{k-1}}{2h_k} > \frac{v_{k-1}}{h_k}$$

Thus this vector has a higher slope than σ_k , which has slope at least as steep as σ_i , meaning condition 2 is violated.

Subsubcase 4.3.3. We must also consider a special case when n is odd.

In this case, we must eliminate saddle connections crossing the horizontal edge to the right of $L_{(n-1)/2}$. Let $k = (n-3)/2$. First of all, notice that any such vector crosses the horizontal distance $h_{k+1} > h_{i-1}$, meaning that any such vector which travels horizontally across at least 3 rectangles violates condition 2. Furthermore, vectors which start at R_k have larger slope than σ_k and thus fail condition 2. Thus we are left with vectors beginning at L_k and R_{k-1} , which must end at R_{k+1} to avoid violating condition 1. Both of these vectors have larger slope than σ_{k+1} , violating condition 2.

We have now eliminated every vector except the corresponding σ_i for **Case 4**. Furthermore, σ_i has positive slope v_{i-1}/h_i and is such that $0 < a + by \leq 1$:

$$a+by = h_i+v_{i-1}y > h_i+v_{i-1} \cdot \frac{1-h_{i-1}}{v_{i-2}} = \frac{v_{i-2}v_{i-1} + v_{i-2}v_i + v_{i-1} - v_{i-2}v_{i-1} - v_{i-1}^2}{v_{i-2}} = \frac{v_{i-1} - 1}{v_{i-2}} > 0$$

$$a + by = h_i + v_{i-1}y \leq h_i + v_{i-1} \cdot \frac{1-h_i}{v_{i-1}} = 1$$

Thus we have shown that for $1 < i \leq \lceil n/2 \rceil - 1$, σ_i is the saddle connection that wins at all $(1, y) \in \Omega_1$ with $\frac{1-h_{i-1}}{v_{i-2}} < y \leq \frac{1-h_i}{v_{i-1}}$, which is simply $P_{n-i+1} \cap \{x = 1\}$.

Case 5. We finally repeat the procedure on σ_1 , showing that it wins on P_n .

We first establish bounds on the horizontal length of a candidate vector $\begin{pmatrix} a \\ b \end{pmatrix}$ if it were to win over σ_1 . In order to satisfy condition 3, we must have that $\frac{1-a}{b} > -h_1$, or equivalently, $a < 1 + h_1b$. Condition 2 stipulates that $a/b > h_1$, so

$$h_1b < a < 1 + h_1b.$$

For any $\begin{pmatrix} a \\ b \end{pmatrix}$ satisfying the above conditions, the endpoint of the saddle connection must be a distance of between 0 and 1 to the right of a line of slope $1/h_1$ passing through the starting vertex of $\begin{pmatrix} a \\ b \end{pmatrix}$. So let us consider the lines of slope $1/h_1$ passing through each vertex of the staircase. The line of slope $1/h_1$ beginning at L_i travels within the staircase to R_{i+2} , as using trigonometric identities gives us that

$$\frac{(h_{i-1} + h_i + h_{i+1})}{h_i} = h_1.$$

The above equation also shows that the line of slope $1/h_1$ beginning at R_i with $i > 0$ passes through the vertical edge below R_{i+1} , then through the horizontal edge to the left of L_{i+1} , then through the vertical edge below R_i , before intersecting L_i , as this saddle connection has the same holonomy vector (and thus slope) as the one connecting L_i and R_{i+2} . For $i = 0$, the line of slope $1/h_1$ beginning at R_i simply connects directly to R_1 .

Now we observe that there are no vertices with horizontal distance between 0 and 1 to the right of the lines of slope $1/h_1$ described above. The vertex R_{i+1} is horizontally at least distance 1 to the right of the line from L_i described above, as shown below. We show that

$$v_i h_1 + 1 \leq h_i + h_{i+1}.$$

We first compute the function

$$h_i + h_{i+1} - v_i h_1 - 1 = \csc\left(\frac{\pi}{n}\right) \sin\left(\frac{\pi(i+1)}{n}\right) - 1.$$

Notice that for fixed n , when $i = 0$, this function evaluates to 0. Moreover, the partial derivative with respect to i is

$$\frac{\pi \csc\left(\frac{\pi}{n}\right) \cos\left(\frac{\pi(i+1)}{n}\right)}{n},$$

which is strictly positive for $i \leq n/2 - 1$ (which is indeed true for our purposes).

Therefore,

$$h_i + h_{i+1} - v_i h_1 - 1 \geq 0$$

for all relevant i , meaning that

$$v_i h_1 + 1 \leq h_i + h_{i+1}.$$

This calculation also shows the vertex L_{i+1} is horizontally at least distance 1 to the right of the line from R_i described above. All other vertices are horizontally at least some distance $h_j \geq 1$ to the right of the lines from each L_i or R_i described above.

The only special case we have failed to consider above is when the lines constructed above pass through the upper edge of the staircase shape. For even n , the calculation below shows that the line of slope $1/h_1$ from $L_{n/2-2}$ passes through the vertical edge below $L_{n/2}$ once before intersecting $L_{n/2}$, as through trigonometric identities, we can verify that

$$h_1(v_{n/2-2} + v_{n/2-1}) = h_{n/2-2} + 2h_{n/2-1}.$$

The above calculation also shows that the line of slope $1/h_1$ through $L_{n/2-1}$ passes through the vertical face below $L_{n/2}$, then passes through the horizontal face to the left of $L_{n/2}$, then passes through the vertical face below $R_{n/2-1}$, before ending at $L_{n/2-1}$, because this saddle connection has the same holonomy vector as the one described immediately before. Again, notice that there are no vertices with horizontal distance between 0 and 1 to the right of the lines described above. By the calculation below, the vertex $L_{n/2}$ is horizontally at least distance 1 to the right of the line from $L_{n/2-1}$ described above. We show that

$$h_1 v_{n/2-1} + 1 \leq 2h_{n/2-1}.$$

We will show the equivalent statement

$$2h_{n/2-1} - h_1 v_{n/2-1} - 1 \geq 0.$$

Using the definitions of h_i and v_i , we have

$$2h_{n/2-1} - h_1 v_{n/2-1} - 1 = \frac{1}{2} \left(\tan\left(\frac{\pi}{2n}\right) + \cot\left(\frac{\pi}{2n}\right) - 2 \right).$$

Note that we must have $n > 2$. Evaluating the above expression at $n = 2$ gives us

$$2h_{2/2-1} - h_1 v_{2/2-1} - 1 = 0.$$

Moreover, the derivative of this function is

$$\frac{\pi \left(\csc^2\left(\frac{\pi}{2n}\right) - \sec^2\left(\frac{\pi}{2n}\right) \right)}{4n^2},$$

and since $\csc\left(\frac{\pi}{2n}\right) \geq \sec\left(\frac{\pi}{2n}\right)$ whenever $n \geq 2$, the derivative is positive for all $n \geq 2$, meaning that we can conclude

$$2h_{n/2-1} - h_1 v_{n/2-1} - 1 \geq 0,$$

and hence the desired inequality. All other vertices are horizontally at least some distance $h_j \geq 1$ to the right of the lines from each L_i or R_i described above, or have been addressed in the earlier cases ignoring the edges at the upper end of S' .

For odd n , the calculation below shows that the line of slope $1/h_1$ from $L_{\lfloor n/2 \rfloor - 1}$ passes through the horizontal face to the right of $L_{\lfloor n/2 \rfloor}$, then intersects $L_{\lfloor n/2 \rfloor}$. Using trigonometric identities, we can verify that

$$2h_1v_{\lfloor n/2 \rfloor - 1} = 2h_{\lfloor n/2 \rfloor - 1} + h_{\lfloor n/2 \rfloor}.$$

All other vertices are horizontally at least some distance $h_j \geq 1$ to the right of the line from each L_i or R_i described above, or have been addressed in the earlier cases in which the edges at the upper end of S' were ignored.

Thus, for every vertex in the staircase shape, there is no $\begin{pmatrix} a \\ b \end{pmatrix}$ beginning at that vertex and ending at another vertex such that $h_1b < a < 1 + h_1b$. We conclude that in the region of Ω_1 with $x = 1$ and $-h_1 < y \leq 1 - h_1$, σ_1 is the saddle connection holonomy vector whose image has the smallest positive slope with horizontal component at most 1.

We have now shown that the λ_i vectors for $1 \leq i \leq n$ are exactly the set of saddle connection vectors which win at points $(1, y) \in \Omega_1$. By lemma 5.3, the λ_i vectors are therefore the complete set of vectors which win at any point in Ω_1 .

Now, in order to completely describe the return time on Ω_1 , we must simply determine which λ_i wins over the other λ_j vectors at each point in Ω_1 .

By an earlier lemma, we know that the λ_i vectors are in order of decreasing slope, so the vector which wins at $(x, y) \in \Omega_1$ is λ_i with maximal i such that $ax + by \leq 1$. Well, λ_n has the maximal i value among the all λ_i vectors, so $\lambda_n = \begin{pmatrix} h_1 \\ 1 \end{pmatrix}$ wins at every $(x, y) \in \Omega_1$ with $h_1x + y \leq 1$; this is exactly the region P_n . Thus the return time at (x, y) will be the slope of $M_{x,y}\lambda_n$, which is

$$\frac{v_0}{x(h_1x + v_0y)}.$$

For $\lfloor n/2 \rfloor + 2 \leq i < n$, $\lambda_i = \sigma_{n-i+1}$ wins at every point $(x, y) \in \Omega_1$ with $h_{n-i+1}x + yv_{n-i} \leq 1$ where λ_j for $j > i$ does not satisfy the same condition, or equivalently, $h_{n-j+1}x + yv_{n-j} > 1$ for $i < j \leq n$. Well, for $j < k < n$ and for all $(x, y) \in \Omega_1$, if $h_{n-j+1}x + yv_{n-j} > 1$ and $h_1x + y > 1$ then $h_{n-k+1}x + yv_{n-k} > 1$ as well, meaning that the above conditions can be simplified to $h_{n-i}x + yv_{n-i-1} > 1$ and $h_1x + y > 1$, which along with the aforementioned condition $h_{n-i+1}x + yv_{n-i} \leq 1$ exactly defines P_i . Thus the return time at (x, y) will be the slope of $M_{x,y}\lambda_i$, which is

$$\frac{v_{n-i}}{x(h_{n+1-i}x + v_{n-i}y)}.$$

For $i = \lfloor n/2 \rfloor + 1$, $\lambda_i = \nu_{\lfloor n/2 \rfloor - 1}$ wins at every point $(x, y) \in \Omega_1$ with $h_{\lfloor n/2 \rfloor - 1}x + yv_{\lfloor n/2 \rfloor - 1} \leq 1$ where λ_j for $j > i$ does not satisfy the same condition, or equivalently, $h_jx + yv_{j-1} > 1$ for all $1 \leq j < \lfloor n/2 \rfloor$. As noted earlier, this condition can be simplified to $h_{\lfloor n/2 \rfloor - 1}x + yv_{\lfloor n/2 \rfloor - 2} > 1$ and $h_1x + y > 1$, which along with the aforementioned condition $h_{\lfloor n/2 \rfloor - 1}x + yv_{\lfloor n/2 \rfloor - 1} \leq 1$ exactly defines P_i . Thus the return time for $(x, y) \in P_i$ will be the slope of $M_{x,y}\lambda_i$, which is

$$\frac{v_{\lfloor n/2 \rfloor - 1}}{x(h_{\lfloor n/2 \rfloor - 1}x + v_{\lfloor n/2 \rfloor - 1}y)}.$$

For $1 < i \leq \lfloor n/2 \rfloor$, $\lambda_i = \nu_{i-2}$ wins at every point $(x, y) \in \Omega_1$ with $h_{i-2}x + yv_{i-2} \leq 1$ where λ_j for $j > i$ does not satisfy the same condition, or equivalently, $h_jx + yv_j > 1$ for all $j > i$, $h_jx + yv_{j-1} > 1$ for all $1 \leq j < \lfloor n/2 \rfloor$. For $j < k$ and for all $(x, y) \in \Omega_1$, if $h_jx + yv_j > 1$ and $h_1x + y > 1$ then $h_kx + yv_k > 1$ as well, and if $h_jx + yv_j > 1$ for any j then $h_{k+1}x + yv_k > 1$ for all k , meaning that the above conditions can be simplified to $h_{i-2}x + yv_{i-2} \leq 1$, $h_{i-1}x + v_{i-1}y > 1$, and $h_1x + y > 1$.

These conditions together exactly describe the region P_i . Thus the return time for $(x, y) \in P_i$ will be the slope of $M_{x,y}\lambda_i$, which is

$$\frac{v_{i-2}}{x(h_{i-2}x + v_{i-2}y)}.$$

Now the only λ vector that remains is $\lambda_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Since this vector has higher slope than all other λ vectors, it can only win within the region $\Omega_1 - \cup_{i>1} P_i$, which is exactly P_1 . λ_1 wins throughout P_1 because $0 \cdot x + 1 \cdot y = y \leq 1$ at every point in P_1 . Thus the return time for $(x, y) \in P_1$ will be the slope of $M_{x,y}\lambda_1$, which is

$$\frac{1}{xy}.$$

□

REFERENCES

- [1] ATHREYA, J. S. Gap distributions and homogeneous dynamics. In *Geometry, topology, and dynamics in negative curvature*, vol. 425 of *London Math. Soc. Lecture Note Ser.* Cambridge Univ. Press, Cambridge, 2016, pp. 1–31.
- [2] ATHREYA, J. S., AND CHAIKA, J. The distribution of gaps for saddle connection directions. *Geom. Funct. Anal.* 22, 6 (2012), 1491–1516.
- [3] ATHREYA, J. S., CHAIKA, J., AND LELIÈVRE, S. The gap distribution of slopes on the golden L. In *Recent trends in ergodic theory and dynamical systems*, vol. 631 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 2015, pp. 47–62.
- [4] ATHREYA, J. S., AND CHEUNG, Y. A Poincaré section for the horocycle flow on the space of lattices. *Int. Math. Res. Not. IMRN*, 10 (2014), 2643–2690.
- [5] DANİ, S. G., AND SMİLLİE, J. Uniform distribution of horocycle orbits for Fuchsian groups. *Duke Mathematical Journal* 51, 1 (1984), 184–194.
- [6] ESKIN, A., AND MASUR, H. Asymptotic formulas on flat surfaces. *Ergodic Theory and Dynamical Systems* 21, 2 (2001), 443–478.
- [7] HUBERT, P., AND SCHMİDT, T. A. Invariants of translation surfaces. *Ann. Inst. Fourier (Grenoble)* 51, 2 (2001), 461–495.
- [8] HUBERT, P., AND SCHMİDT, T. A. An introduction to Veech surfaces. In *Handbook of dynamical systems. Vol. 1B*. Elsevier B. V., Amsterdam, 2006, pp. 501–526.
- [9] KUMANDURI, L., AND WANG, J. Slope gap distributions of Veech surfaces. Preprint, arXiv:math/2102.10069.
- [10] MARKLOF, J., AND STRÖMBERGSSON, A. The distribution of free path lengths in the periodic lorentz gas and related lattice point problems. *Annals of Mathematics* (2010), 1949–2033.
- [11] MASUR, H. Ergodic theory of translation surfaces. In *Handbook of dynamical systems. Vol. 1B*. Elsevier B. V., Amsterdam, 2006, pp. 527–547.
- [12] SANCHEZ, A. Gaps of saddle connection directions for some branched covers of tori. to appear, ETDS, 2021.
- [13] SARNAK, P. Asymptotic behavior of periodic orbits of the horocycle flow and eisenstein series. *Communications on Pure and Applied Mathematics* 34 (1981), 719–739.
- [14] SMİLLİE, J., AND ULCİGRAI, C. Beyond Sturmian sequences: coding linear trajectories in the regular octagon. *Proc. Lond. Math. Soc. (3)* 102, 2 (2011), 291–340.
- [15] TAHA, D. On cross sections to the geodesic and horocycle flows on quotients of $SL(2, \mathbb{R})$ by hecke triangle groups G_q . Preprint, arXiv:math/1906.07250.
- [16] UYANIK, C., AND WORK, G. The distribution of gaps for saddle connections on the octagon. *Int. Math. Res. Not. IMRN*, 18 (2016), 5569–5602.
- [17] VEECH, W. A. Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards. *Inventiones mathematicae* 97 (1989), 553–583.
- [18] VEECH, W. A. Siegel measures. *Annals of Mathematics* 148 (1998), 865–944.
- [19] VOROBETS, Y. B. Plane structures and billiards in rational polygons: the Veech alternative. *Uspekhi Mat. Nauk* 51, 5(311) (1996), 3–42.
- [20] WORK, G. A transversal for horocycle flow on $\mathcal{H}(\alpha)$. *Geom. Dedicata* 205 (2020), 21–49.
- [21] ZORICH, A. Flat surfaces. *Frontiers in number theory, physics, and geometry I* (2006), 439–585.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 10 HILLHOUSE AVENUE, NEW HAVEN, CT 06520
Email address: jonah.berman@yale.edu

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 10 HILLHOUSE AVENUE, NEW HAVEN, CT 06520
Email address: taylor.mcadam@yale.edu

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 10 HILLHOUSE AVENUE, NEW HAVEN, CT 06520
Email address: ananth.miller-murthy@yale.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, 480 LINCOLN DRIVE, MADISON, WI 53706
Email address: caglar@math.wisc.edu

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 10 HILLHOUSE AVENUE, NEW HAVEN, CT 06520
Email address: hamilton.wan@yale.edu