

# MODELS FOR SHORT SEQUENCES OF MEASURES IN THE COFINALITY- $\omega$ CONSTRUCTIBLE MODEL

UR YA'AR

**ABSTRACT.** We investigate the relation between  $C^*$ , the model of sets constructible using first order logic augmented with the “cofinality- $\omega$ ” quantifier, and “short” sequences of measures – sequences of measures of order 1, which are shorter than their minimum. We show that certain core models for short sequences of measures are contained in  $C^*$ ; we compute  $C^*$  in a model of the form  $L[\mathcal{U}]$  where  $\mathcal{U}$  is a short sequence of measures, and in models of the form  $L[\mathcal{U}][G]$  where  $G$  is generic for adding Prikry sequences to some of the measurables of  $\mathcal{U}$ ; and prove that if there is an inner model with a short sequence of measures of order type  $\chi$ , then there is such an inner model in  $C^*$ .

## 1. INTRODUCTION

The model  $C^*$ , introduced by Kennedy, Magidor and Väänänen in [1], is the model of sets constructible using the logic  $\mathcal{L}(Q_\omega^{\text{cf}})$  – first order logic augmented with the “cofinality- $\omega$ ” quantifier. As in the case of  $L$  – the model of sets constructible using first order logic – this is a model of ZFC. It is however in many cases much larger than  $L$ , for example in the presence of large cardinals. Specifically, Kennedy, Magidor and Väänänen prove in [1] the following:

- Theorem.**
- (1)  $C^*$  contains the Dodd-Jensen core model.
  - (2) If there is an inner model for a measurable cardinal, then  $C^*$  contains an inner model for a measurable cardinal.
  - (3) If  $V = L^\mu$  for some measure  $\mu$  then  $C^* = M_{\omega^2}[\langle \kappa_{\omega \cdot n} \mid n < \omega \rangle]$  where  $M_{\omega^2}$  is the  $\omega^2$ -iterate of  $V$  and  $\kappa_{\omega \cdot n}$  is the  $\omega \cdot n$  image of the measurable under the iteration embedding.

Our aim in this paper is to generalize this theorem to obtain inner models with more measurable cardinals. Our focus here is on “short” sequences of measures – a notion introduced by Koepke in [2, 3] – sequences of measures of order 1, which are shorter than their minimum (so the sequence doesn’t “stretch” when taking ultrapowers). In section 2 we define the notions of *short mice* and *short core models*, and present some basic facts concerning them; in section 3 we show that certain short core models are contained in  $C^*$ ; in section 4 we compute  $C^*$  in a model of the form  $L[\mathcal{U}]$  where  $\mathcal{U}$  is a short sequence

---

I would like to thank my advisor, Prof. Menachem Magidor, for his guidance and support without which this work would not have been possible.

of measures, and in models of the form  $L[\mathcal{U}][G]$  where  $G$  is generic for adding Prikry sequences to some of the measurables of  $\mathcal{U}$ ; and in section 5 we prove that if there is an inner model with a short sequence of measures of order type  $\chi$ , then there is such an inner model in  $C^*$ .

It is interesting to note that while we can get inner models with many measurable cardinals contained in  $C^*$ , it is yet unknown whether  $C^*$  can actually contain even a single measurable cardinal. By [1, theorem 5.7], if  $V = C^*$  then there is no measurable cardinal in  $V$ , however  $C^*$  does not necessarily satisfy  $V = C^*$ , so it might still be possible to have a measurable cardinal in  $C^*$ , which would simply imply that  $(C^*)^{C^*} \neq C^*$ .

## 2. SHORT MICE

First we cite some definitions from [3] (as they appear in [4]).

**Definition 1.** Let  $D$  be a class.  $D$  is simple if the following statements hold:

- (1) If  $x \in D$ , then there is  $\delta \in \mathbf{Ord}$  with  $x = \langle \delta, a \rangle$  with  $a \subseteq \delta$ .
- (2) If  $\langle \delta, a \rangle \in D$ , then  $\langle \delta, \delta \rangle \in D$

In the above situation, we define  $\text{Dom}(D) = \{\delta \mid \langle \delta, \delta \rangle \in D\}$  and  $D(\delta) = \{a \subseteq \delta \mid \langle \delta, a \rangle \in D\}$  for all  $\delta \in \text{Dom}(D)$ .  $D$  is called a *sequence of measures* if  $D$  is simple and  $D(\delta)$  is a normal ultrafilter on  $\delta$  for every  $\delta \in \text{Dom}(D)$

**Definition 2.** Let  $D$  be a simple set.

- (1) We say that  $M = \langle |M|, F_M \rangle$  is a *premouse over  $D$*  if the following statements hold:

- (a)  $|M|$  is a transitive set and  $F_M$  is a simple set with

$$\sup(\text{Dom}(D)) < \min(\text{Dom}(F_M)) \in |M|.$$

- (b)  $\langle |M|, \in, F_M \rangle \models \text{“}F_M \text{ is a sequence of measures”}$ .
- (c)  $|M| = J_{\alpha_M}[D, F_M]$  for some ordinal  $\alpha_M$  with  $\omega \cdot \alpha_M = |M| \cap \mathbf{Ord}$ .

In the above situation, we set  $\text{meas}(M) = \text{Dom}(F_M) \cap (\omega \cdot \alpha_M)$  (the measurables of  $M$ ) and  $\text{lp}(M) = \mathbf{H}(\min(\text{meas}(M)))^{|M|}$  (the lower part of  $M$ ).

- (2) Given a premouse  $M$  over  $D$  and  $\delta \in \text{meas}(M)$ , a premouse  $M_*$  over  $D$  is the *ultrapower* of  $M$  at  $\delta$  if there is a unique map  $j : |M| \rightarrow |N|$  with the following properties, and  $M_* = N$  holds for this  $N$ .

- (a)  $j : \langle |M|, \in, D, F_M \rangle \rightarrow \langle |N|, \in, D, F_N \rangle$  is  $\Sigma_1$ -elementary.
- (b)  $|N| = \left\{ j(f)(\delta) \mid f \in {}^\delta |M| \cap |M| \right\}$ .
- (c)  $F_M(\delta) \cap |M| = \{x \in \mathcal{P}(\delta) \cap |M| \mid \delta \in j(x)\}$ .

If such a premouse exists, then we denote it by  $\text{Ult}(M, F(\delta))$  and we call the corresponding map  $j$  the ultrapower embedding of  $M$  at  $\delta$ .

- (3) Given a premouse  $M$  over  $D$  and a function  $I : \lambda \rightarrow \mathbf{Ord}$  with  $\lambda \in \mathbf{Ord}$ , a system

$$\text{It}(M, I) = \langle \langle M_\alpha \mid \alpha \leq \lambda \rangle, \langle j_{\alpha, \beta} \mid \alpha \leq \beta \leq \lambda \rangle \rangle$$

is called the *iterated ultrapower of  $M$  by  $I$*  if the following statements hold for all  $\gamma \leq \lambda$ :

- (a)  $M = M_0$  and  $M_\gamma$  is a premouse over  $D$ .
  - (b) Given  $\alpha \leq \beta \leq \gamma$ ,  $j_{\beta, \gamma} : |M_\beta| \rightarrow |M_\gamma|$  is a function,  $j_{\gamma, \gamma} = \text{id}_{|M_\gamma|}$  and  $j_{\alpha, \gamma} = j_{\beta, \gamma} \circ j_{\alpha, \beta}$ .
  - (c) If  $\gamma < \lambda$  and  $I(\gamma) \in \text{meas}(M_\gamma)$ , then  $M_{\gamma+1} = \text{Ult}(M_\gamma, F_{M_\gamma}(I(\gamma)))$  and  $j_{\gamma, \gamma+1}$  is the ultrapower embedding of  $M_\gamma$  at  $F_{M_\gamma}(I(\gamma))$ . In the other case, if  $\gamma < \lambda$  and  $I(\gamma) \notin \text{meas}(M_\gamma)$ , then  $M_\gamma = M_{\gamma+1}$  and  $j_{\gamma, \gamma+1} = \text{id}_{|M_\gamma|}$ .
  - (d) If  $\gamma$  is a limit ordinal, then  $\langle \langle M_\beta, \in, D, F_{M_\beta} \rangle, \langle j_{\beta, \gamma} \mid \beta < \gamma \rangle \rangle$  is a direct limit of the directed system  $\langle \langle M_\beta, \in, D, F_{M_\beta} \mid \beta < \gamma \rangle, \langle j_{\alpha, \beta} \mid \alpha \leq \beta < \gamma \rangle \rangle$ .
- In this situation, we let  $M_I$  denote  $M_\lambda$  and let  $j_I$  denote  $j_{0, \lambda}$ .

- (4) A premouse  $M$  over  $D$  is *iterable* if the system  $\text{It}(M, I)$  exists for every function  $I : \lambda \rightarrow \mathbf{On}$  with  $\lambda \in \mathbf{On}$ .
- (5) A premouse  $M$  over  $D$  is *short* if one of the following statements holds:
  - (a)  $D = \emptyset$  and  $\text{otp}(\text{meas}(M) \cap \gamma) < \min(\text{meas}(M))$  for all  $\gamma \in |M| \cap \mathbf{On}$ .
  - (b)  $D \neq \emptyset$  and  $\text{otp}(\text{meas}(M)) \leq \min(\text{Dom}(D))$ .

At this stage we diverge from Koepke's terminology and pick up Mitchell's notion of a mouse from [5, §3], in order to be able to use a stronger comparison lemma. Note that for short sequences, the notion of coherence (definition 1.6 in [5]) is trivial.

- (6) [5, definition 3.1] A premouse  $M = J_\alpha[D, F]$  is called a (*short*)  $D$ -*mouse* if it is short, iterable and satisfies that there is some  $\rho$ ,  $\text{sup}(D) \leq \rho < \min(F)$  and a parameter  $p \in [\alpha \setminus \rho]^{<\omega}$  such that  $M = H_1^M(\rho \cup p)$  (the  $\Sigma_1$  Skolem hull in  $M$ ).

Denote by  $\rho_M$  the least  $\rho$  for which such a  $p$  exists and  $p_M$  the least such parameter (under the usual parameter well-order).

- (7) [5, definition 3.2] If  $M, N$  are (*short*)  $D$ -mice, denote  $M <_D N$  if there are iterated ultrapowers  $i : M \rightarrow J_\alpha[D, F]$ ,  $j : N \rightarrow J_\beta[D, G]$  such that  $i \upharpoonright \rho_M$  and  $j \upharpoonright \rho_N$  are the identity,  $\alpha \leq \beta$ ,  $G \upharpoonright \alpha = F$  and  $(\alpha, \rho_M, i(p_M)) < (\beta, \rho_N, j(p_N))$  lexicographically.

**Lemma 3** ([5, theorem 3.3]).  *$<_D$  is a well ordering of (*short*)  $D$  mice such that  $M <_D N$  implies  $M \in L(N)$ .*

*Remark 4.* The original theorem in [5] discusses general mice, and the proof uses comparison of mice by iterations. Since iterations of short mice preserves shortness it holds when restricting to short mice as well.

**Lemma 5.** *If  $M$  is a  $D$ -mouse then for every  $\eta \in (\rho_M, \alpha_M)$  such that  $M \models \eta$  is regular,  $\text{cf}^V(\eta) = \text{cf}^V(\alpha_M)$ .*

*Proof.*  $M = J_{\alpha_M}[D, F]$  for some  $F$ . Let  $\eta \in (\rho_M, \alpha_M)$  such that  $M \models \eta$  is regular. Let  $\beta \in (\eta, \alpha_M)$  regular in  $M$ .  $M$  has a Skolem function  $h_\beta$  for  $J_\beta[D, F]$ , and since  $M \models \eta$  regular, and  $\rho_M < \eta$ , the closure of  $\rho_M \cup p_M$  under  $h_\beta$  (denoted  $h_\beta^{\text{“}}(\rho_M \cup p_M)$ ), which can be computed in  $M$ , is bounded by some  $\alpha_\beta < \eta$ . But since  $M = \mathcal{H}_1^M(\rho_M \cup p_M)$ , each element of  $\eta$  is in  $h_\beta^{\text{“}}(\rho_M \cup p_M)$  for some  $\beta < \alpha_M$  (if an element satisfies some  $\Sigma_1$  formula in  $M$ , then it satisfies this formula in some initial segment of  $M$ ), so the sequence  $\langle \alpha_\beta \mid \beta < \alpha_M \rangle$  is cofinal in  $\eta$ , hence  $\text{cf}^V(\eta) = \text{cf}^V(\alpha_M)$ .  $\square$

*Notation 6.* Denote by  $\kappa(\gamma)$  be the  $\gamma$ th member of  $\text{Dom}(F)$  and  $\xi(\gamma) = (\kappa(\gamma)^+)^M$ . Note that for every  $\gamma$  we have  $\xi(\gamma) > \rho_M$  and  $M \models \xi(\gamma)$  is regular so by lemma 5  $\text{cf}^V(\xi(\gamma)) = \text{cf}^V(\alpha_M)$ . Denote this by  $\delta_M$ .

**Definition 7.** Define  $K[D]$  as the class of sets constructible from short  $D$ -mice.

**Theorem 8.**  $K[D] = \bigcup \{\text{lp}(M) \mid M \text{ is a short } D\text{-mouse}\}$ .

*Remark 9.* This is the way  $K[D]$  is defined by Koepke in [2, 3], so our slightly different definition yields the same result.

*Proof.* The inclusion  $\supseteq$  is clear, so we need to show that for every  $x \in K[D]$  there is some short  $D$ -mouse containing  $x$  in its lower part.  $x$  is constructed by using a set of mice,  $X$ . Let  $S$  be an ascending sequence of regular cardinals greater than  $\lambda^+$  where  $\lambda = \sup(\{|M| \mid M \in X\} \cup \{|\text{tr}(x)|, |X|\})$ , such that  $\text{otp } S = \sup\{\text{otp meas}(M) \mid M \in X\}$ . We iterate each  $M \in X$  according to  $S$  (i.e. for  $\gamma < \text{otp meas}(M)$ , we iterate  $M$  by its  $\gamma$ th measure  $S(\gamma)$  many times), so by [2, lemma 5.17] the result of the iteration will be of the form  $\bar{M} = J_{\alpha_{\bar{M}}}[D, \text{CU} \upharpoonright S]$  where  $\text{CU}$  is the closed unbounded simple predicate  $-\langle \kappa, z \rangle \in \text{CU}$  iff  $\kappa$  is a limit ordinal and  $z \subseteq \kappa$  contains a closed unbounded subset of  $\kappa$ . Note that these will all be short iterable pre-mice (as iterations of short mice). Let  $\alpha^* = \sup\{\alpha_{\bar{M}} \mid M \in X\}$ ,  $M^* = J_{\alpha^*}[D, \text{CU} \upharpoonright S]$ . We claim that  $X \subseteq M^*$ : if  $M \in X$ ,  $i : M \rightarrow \bar{M}$  the iteration embedding, then  $\bar{M}$  is an initial segment of  $M^*$ , and  $M \cong H_1^{\bar{M}}(\rho_M \cup i(p_M)) \in M^*$ , so  $M \in M^*$ .

Assume first that for all  $M \in X$ ,  $\alpha^* > \alpha_{\bar{M}}$  and let  $M^{**}$  be the transitive collapse of  $H_1^{M^*}(\lambda^+)$ . This is a short  $D$ -mouse, and we claim that already  $X \subseteq M^{**}$ . Let  $M \in X$ , then  $M = H_1^{\bar{M}}(a)$  for some  $a \in \bar{M}$ . Since  $\alpha^* > \alpha_{\bar{M}}$ ,  $\bar{M} \in M^*$ , so  $M^*$  satisfies the  $\Sigma_1$  statement “ $\exists y \exists z (M = H_1^y(z))$ ”, so such  $y, z$  will be in  $H_1^{M^*}(\lambda^+)$  ( $|M| < \lambda$  so it can be coded as a subset of  $\lambda^+$ ), so  $M \in M^{**}$ . So  $x$  can be constructed in  $M^{**}$ , which is a mouse with minimal measure above  $|\text{tr}(x)|$  so  $x$  is in  $\text{lp}(M^{**})$ .

Now assume  $\alpha^* = \alpha_{\bar{N}}$  for some  $N \in X$ . This means that there is some  $N \in X$  such that  $M \leq_D N$  for every  $M \in X$ . Hence by lemma 3,  $X \subseteq L[N]$  so  $x \in L[N]$ . Assume

$x \in J_\kappa[N]$  and iterate  $N$  to some  $\bar{N}$  with  $\min \text{meas}(\bar{N}) > \lambda' = \max\{\lambda, \kappa\}^+$ . Then  $x \in M^{**} := H_1^{\bar{N}}(\lambda')$  which is a mouse with minimal measure  $> |\text{tr}(X)|$ .  $\square$

We now establish some notation and terminology for iterations of short  $D$ -mice.

**Notation 10.** Let  $M = \langle |M|, D, F \rangle$  be a short  $D$ -mouse,  $\chi \leq \text{otp Dom}(F)$ , and  $\vec{\theta} = \langle \theta_\gamma \mid \gamma < \chi \rangle$  some sequence of ordinals. We define the *iteration of  $M$  according to  $\vec{\theta}$*  as follows. Set  $M_0^0 = M$ ,  $F_0^0 = F$  and define an iteration  $\langle M_\beta^\gamma, i_{\alpha,\beta}^\gamma, j^\gamma \mid \gamma < \chi, \alpha \leq \beta < \theta_\gamma \rangle$  inductively on  $\gamma < \chi$  such that:

- ◆  $i_{\alpha,\beta}^\gamma : M_\alpha^\gamma \rightarrow M_\beta^\gamma$  and  $j^\gamma : M_0^\gamma \rightarrow M_0^\gamma$  are  $\Sigma_1$ -elementary;
- ◆  $i_{\beta,\beta}^\gamma = \text{Id}_{M_\beta^\gamma}$  and for  $\alpha < \beta' \leq \beta$ ,  $i_{\alpha,\beta}^\gamma = i_{\beta',\beta}^\gamma \circ i_{\alpha,\beta'}^\gamma$ ;
- ◆ If  $M_\beta^\gamma = \langle |M_\beta^\gamma|, D, F_\beta^\gamma \rangle$  we set  $A_\beta^\gamma = \text{Dom}(F_\beta^\gamma)$ ;

Note that these are iterations of short mice so  $\text{otp Dom}(F_\beta^\gamma) = \text{otp Dom}(F)$ .

- ◆  $\kappa_0^\gamma$  is the  $\gamma$ th element of  $A_0^\gamma$ ,  $\xi_0^\gamma = \left( (\kappa_0^\gamma)^+ \right)^{M_0^\gamma}$  or (if this doesn't exist in  $M_0^\gamma$ )  $\xi_0^\gamma = M_0^\gamma \cap \mathbf{Ord}$ ,  $U_0^\gamma = F_0^\gamma(\kappa_0^\gamma)$  is the ultrafilter of  $M_0^\gamma$  on  $\kappa_0^\gamma$ .
- ◆ If  $M_0^\gamma, \kappa_0^\gamma, \xi_0^\gamma, U_0^\gamma$  are defined, iterate  $M_0^\gamma$  according to  $U_0^\gamma$  up to  $\theta_\gamma$  to get the iteration  $\langle M_\beta^\gamma, i_{\alpha,\beta}^\gamma \mid \alpha, \beta \leq \theta_\gamma \rangle$  and set  $\kappa_\beta^\gamma = i_{0,\beta}^\gamma(\kappa_0^\gamma)$ ,  $\xi_\beta^\gamma = i_{0,\beta}^\gamma(\xi_0^\gamma)$ ,  $U_\beta^\gamma = i_{0,\beta}^\gamma(U_0^\gamma)$ .
- ◆ Set  $M_0^{\gamma+1} = M_{\theta_\gamma}^\gamma$ ,  $j^{\gamma+1} = i_{0,\theta_\gamma}^\gamma \circ j^\gamma$ .
- ◆ For limit  $\gamma$ , set  $M_0^\gamma, j^\gamma$  as the directed limit of  $\langle M_0^\beta, j^\beta \mid \beta < \gamma \rangle$ .
- ◆ Let  $M^\chi$  be the directed limit of  $\langle M_0^\beta, j^\beta \mid \beta < \chi \rangle$  if  $\chi$  is limit or  $M_{\theta_\gamma}^\gamma$  if  $\chi = \gamma+1$ , with  $j^\chi : M_0^0 \rightarrow M^\chi$  the corresponding embedding.

**Lemma 11.** *In the above context, for every  $\gamma < \chi$  and  $\beta < \theta_\gamma$  we have  $\text{cf}^V(\xi_\beta^\gamma) = \text{cf}^V(\xi(\gamma)) = \delta_M$ .*

*Proof.* From the definitions, we can see that  $\xi_\beta^\gamma$  is in  $M_\beta^\gamma$  the image of  $\xi(\gamma)$  under an iteration embedding  $j := i_{0,\beta}^\gamma \circ j^\gamma$ . Thus every  $\eta < \xi_\beta^\gamma$  is of the form  $j(f)(\kappa_1, \dots, \kappa_n)$  for  $f \in M_0^0$ ,  $f : \kappa(\gamma)^n \rightarrow \xi(\gamma)$  and some iteration points  $\kappa_1 < \dots < \kappa_n < \kappa_\beta^\gamma$ .  $f$  must be bounded in  $\xi(\gamma)$  so if  $\langle \gamma_\alpha \mid \alpha < \text{cf}^V(\xi(\gamma)) \rangle$  is cofinal in  $\xi(\gamma)$ , there is some  $\alpha$  such that  $f(\zeta) < \gamma_\alpha$  for every  $\alpha$  so  $j(f)(\kappa_1, \dots, \kappa_n) < j(\gamma_\alpha)$  hence  $\langle j(\gamma_\alpha) \mid \alpha < \text{cf}^V(\xi(\gamma)) \rangle$  is cofinal in  $\xi_\beta^\gamma$ , and since  $\text{cf}^V(\xi(\gamma))$  is regular we have  $\text{cf}^V(\xi(\gamma)) = \text{cf}^V(\xi_\beta^\gamma)$ .  $\square$

### 3. SHORT MICE IN $C^*$

**Theorem 12.** *Let  $D \in C^*$  be simple. Then any short  $D$ -mouse is in  $C^*$ . As a result,  $(K[D])^{C^*} = K[D]$ .*

*Proof.* Assume towards contradiction this is not the case, and let  $M = \langle |M|, D, F \rangle$  be the  $<_D$ -minimal short  $D$ -mouse not in  $C^*$ . Our goal is to find an appropriate sequence  $\vec{\theta} = \langle \theta_\gamma \mid \gamma < \chi \rangle$  where  $\chi := \text{otp Dom}(F)$ , iterate  $M$  according to  $\vec{\theta}$  and show, following

the proof of [1, theorem 5.5], that in fact  $M^\chi \in C^*$ , and from that deduce that  $M \in C^*$ , by contradiction.

For any sequence  $\langle \theta_\gamma \mid \gamma < \chi \rangle$ , the final model in the iteration is  $M^\chi = J_\alpha [D, F_\chi]$  for some simple  $F_\chi$ , where by the construction  $F_\chi = j^\chi(F)$  and  $\text{otp Dom } F_\chi = \chi$ . For any  $\gamma < \chi$ , the  $\gamma$ th element of  $F_\chi$  is  $\kappa_{\theta_\gamma}^\gamma$ , and the corresponding  $M^\chi$ -ultrafilter  $U_\gamma^\chi$  satisfies that for  $X \in M^\chi \cap \mathcal{P}(\kappa_{\theta_\gamma}^\gamma)$ ,  $X \in U_\gamma^\chi$  iff  $X$  contains a final segment of  $E_\gamma := \{ \kappa_\beta^\gamma \mid \beta < \theta_\gamma \}$  iff  $X$  contains a cofinal subset of  $E_\gamma$ . So for any cofinal  $\bar{E} \subseteq E_\gamma$ , if  $F_{\bar{E}}$  is the filter generated by final segments of  $\bar{E}$ , then  $U_\gamma^\chi = F_{\bar{E}} \cap M^\chi$ . So we want to find a sequence  $\vec{\theta}$  for which we can show the following claim which will imply that  $M^\chi \in C^*$ :

*Claim 13.* There is a sequence  $\langle \bar{E}_\gamma \mid \gamma < \chi \rangle \in C^*$  such that for every  $\gamma$   $\bar{E}_\gamma$  is cofinal in  $E_\gamma$ .

To choose the right  $\vec{\theta}$  and prove the claim, we need to be able to identify in  $C^*$  unboundedly many critical points of the iteration. For this, we need the following crucial claims:

*Claim 14.* For every  $\gamma < \chi$  and  $\beta$ , if  $\eta$  is such that  $\kappa_0^\gamma < \eta < \kappa_\beta^\gamma$  and  $M_\beta^\gamma \models \eta$  is regular, then either there is  $\alpha < \beta$  such that  $\eta = \kappa_\alpha^\gamma$  or  $\text{cf}^V(\eta) = \delta_M$ .

*Claim 15.* For every  $\gamma < \chi$  and  $\beta \leq \theta_\gamma$ ,  $\eta \leq \kappa_\beta^\gamma$  is regular in  $M_\beta^\gamma$  iff it is regular in  $K^* := (K[D])^{C^*}$ . In particular every  $\kappa_\beta^\gamma$  is regular in  $K^*$ .

Let's first assume these claims and show that claim 13 (and then the theorem) follows. First we need to find the appropriate  $\langle \theta_\gamma \mid \gamma < \chi \rangle$ . We define in  $C^*$  a sequence  $\langle \lambda_\gamma \mid \gamma < \chi \rangle$  recursively such that  $\lambda_0 > |M|^V$  and for every  $\gamma < \chi$ :

- (1)  $C^* \models \lambda_\gamma$  is a regular cardinal;
- (2)  $\gamma' < \gamma$  implies  $\lambda_{\gamma'} < \lambda_\gamma$ ;
- (3)  $\forall \beta < \lambda_\gamma \exists \eta \in (\beta, \lambda_\gamma)$  such that  $K^* \models \eta$  is a regular cardinal, and  $\text{cf}^V(\eta) = \omega \leftrightarrow \delta_M \neq \omega$ .

Assume we've defined  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle \in C^*$  for some  $\gamma < \chi$ . If  $\lambda > \sup \langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$  is regular in  $V$  then it satisfies 1-3: regularity in  $V$  implies regularity in  $C^*$ , and 2 is clear. To show 3 iterate  $M$  according to  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle \frown \langle \lambda \rangle$  (where in the  $\gamma$ th stages we iterate up to  $\lambda$ ), and consider the iteration points  $\langle \kappa_\beta^\gamma \mid \beta < \lambda \rangle$ . Since  $\lambda$  is regular in  $V$  and we can assume  $> \omega_1$ , there are cofinally many iteration points of cofinality  $\omega$  and cofinally many of cofinality  $> \omega$ . Each such iteration point is regular in  $K^*$  by claim 15, so  $\lambda$  satisfies 3 (choosing the appropriate iteration points according to  $\delta_M$ ). So there are ordinals satisfying 1-3, and whether an ordinal satisfies 1-3 can be determined within  $C^*$ . So in  $C^*$  we can define  $\lambda_\gamma$  to be the first ordinal  $> \sup \langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$  (or  $> |M|$  if  $\gamma = 0$ ) satisfying 1-3. Note that this defines the sequence  $\langle \lambda_\gamma \mid \gamma < \chi \rangle$  recursively in  $C^*$  so we can in fact proceed through limit stages.

Consider now  $\lambda = \lambda_\gamma > \sup \langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$  which satisfies 1-3, and iterate  $M$  only according to  $\kappa(\gamma)$  such that the image of  $\kappa(\gamma)$  is greater than  $\lambda$ . By 3  $\lambda$  is the limit of ordinals  $\eta$  satisfying  $K^* \models \eta$  is a regular cardinal, and  $\text{cf}^V(\eta) = \omega \leftrightarrow \delta_M \neq \omega$ . By claims 14 and 15 each such  $\eta$  is an iteration point, so also  $\lambda$  must be an iteration point. So we have some iteration  $j : M \rightarrow N$  by the ultrafilter  $F(\gamma)$  of some length  $\theta$ , such that  $j(\kappa(\gamma)) = \lambda$ . Note that everything below  $\kappa(\gamma)$  is fixed so  $j(F) \upharpoonright \lambda = F \upharpoonright \kappa(\gamma)$ , and we can iterate  $N$  according to  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$ . Since in  $N$   $\lambda$  is strongly inaccessible and is above  $\sup \langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$ , it is not moved by this iteration. It is a known fact that the resulting iteration is isomorphic to the iteration of  $M$  according to  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle \widehat{\ } \langle \theta \rangle$ . So we get that in fact  $\lambda$  is the image of  $\kappa(\gamma)$  under the iteration according to  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle \widehat{\ } \langle \theta \rangle$ , i.e.  $\lambda = \kappa_{\theta}^\gamma$ . We set  $\theta_\gamma$  to be this  $\theta$ .

So to conclude, we've defined recursively  $\langle \lambda_\gamma \mid \gamma < \chi \rangle$  inside  $C^*$ , and we have (in  $V$ !) a sequence  $\vec{\theta} = \langle \theta_\gamma \mid \gamma < \chi \rangle$  such that when iterating  $M$  according to  $\vec{\theta}$  we have for every  $\gamma$ ,  $\lambda_\gamma = \kappa_{\theta_\gamma}^\gamma$ .

*Proof of claim 13.* We iterate  $M$  according to  $\vec{\theta}$  as above and obtain (in  $V$ ) the iteration points  $E_\gamma = \{ \kappa_\beta^\gamma \mid \beta < \theta_\gamma \}$ . For every  $\gamma < \chi$  set

$$\bar{E}_\gamma = \begin{cases} \left\{ \eta < \lambda_\gamma \mid \eta \text{ is regular in } K^* \text{ and } \text{cf}^V(\eta) = \omega \right\} & \delta_M \neq \omega \\ \left\{ \eta < \lambda_\gamma \mid \eta \text{ is regular in } K^* \text{ and } \text{cf}^V(\eta) \neq \omega \right\} & \delta_M = \omega \end{cases}$$

Each of these sets is in  $C^*$ , and by claims 14 and 15 and lemma 5,  $\bar{E}_\gamma \subseteq E_\gamma$  for every  $\gamma$ . By our choice of  $\lambda_\gamma$  and  $\theta_\gamma$ , they are cofinal in  $\lambda_\gamma = \kappa_{\theta_\gamma}^\gamma$  (by clause 3). Since  $\langle \lambda_\gamma \mid \gamma < \chi \rangle \in C^*$ , the sequence  $\langle \bar{E}_\gamma \mid \gamma < \chi \rangle$  is in  $C^*$  as well.

Claim 13  $\square$

Now as we claimed in the beginning of the proof, we can compute each ultrafilter  $U_\gamma^\chi$  from this sequence, hence in  $C^*$  we can compute  $M^\chi$ . Thus by lemma 3  $M \in C^*$ , by contradiction.

Theorem 12  $\square$

We now move on to prove claims 14 and 15.

*Proof of claim 14.* By examining the proof of claim 2 in the proof of [1, theorem 5.6], which corresponds to our claim, one can see that it can be stated as a general lemma:

**Lemma 16** (in [1, theorem 5.5]). *Assume that  $\langle N_\alpha, k_{\alpha,\beta} \mid \alpha \leq \beta < \theta \rangle$  is an ultrapower iteration of  $U_0 \in N_0$  which is (in  $N_0$ ) an ultrafilter on  $\lambda_0$ , i.e. for every  $\alpha < \theta$ ,  $N_{\alpha+1} = \text{Ult}(N_\alpha, i_{0,\alpha}(U_0))$  and for limit  $\alpha$   $N_\alpha$  is the directed limit. Let  $\zeta_0 = (\lambda_0^+)^{M_0}$  (or  $M_0 \cap \mathbf{Ord}$  if this doesn't exist) and  $\delta = \text{cf}^V(\zeta_0)$ . Set  $\lambda_\alpha = i_{0,\alpha}(\lambda_0)$  and assume that for every  $\alpha < \theta$ ,  $(\lambda_\alpha)^{\lambda_\alpha} \cap N_\alpha$  is the increasing union of  $\delta$  members of  $N_\alpha$ , each of cardinality  $\lambda_\alpha$  in  $N_\alpha$ . Then for every  $\beta < \theta$  and every  $\eta$  such that  $\lambda_0 < \eta < \lambda_\beta$ , if  $N_\beta \models \eta$  is regular, then either  $\exists \gamma < \beta$  such that  $\eta = \lambda_\gamma$  or  $\text{cf}^V(\eta) = \delta$ .*

Recall that  $\text{cf}^V(\xi(\gamma)) = \delta_M = \text{cf}^V(\alpha_M)$ , so to use this lemma, we need to show that for every  $\gamma$  and  $\beta < \lambda_\gamma$ ,  $(\kappa_\beta^\gamma)^{\kappa_\beta^\gamma} \cap M_\beta = \bigcup_{\psi < \delta_M} F_\psi$  such that  $F_\psi \in M_\beta^\gamma$  and  $|F_\psi| = \kappa_\beta^\gamma$ . Set for the moment  $N = M_\beta^\gamma$ ,  $\kappa = \kappa_\beta^\gamma$ ,  $\xi = \xi_\beta^\gamma = (\kappa^+)^N$ . Assume  $N = J_{\alpha_N}[D, F]$  and note that the image of the iteration embedding from  $M$  to  $N$  is cofinal, so  $\text{cf}^V(\alpha_N) = \text{cf}^V(\alpha_M)$ . Since  $N = H_1^N(\rho_N \cup p_N)$ , every  $h \in \kappa^\kappa \cap N$  is in some  $H_1^{J_\beta^{[D, F]}}(\rho_N \cup p_N)$  for some  $\beta < \alpha_N$  (as in the proof of lemma 5), and since  $\rho_M < \kappa$ , this is of size at most  $\kappa$ . So if we take  $\langle \beta_\psi \mid \psi < \delta_M \rangle$  cofinal in  $\alpha_N$ , then  $F_\psi = \kappa^\kappa \cap H_1^{J_{\beta_\psi}^{[D, F]}}(\rho_N \cup p_N)$  is as required. So we can apply lemma 16 and get our claim. Claim 14  $\square$

As for claim 15, it follows by the minimality of  $M$  and its iterates:

*Proof of claim 15.* Fix  $\gamma < \chi$  and  $\beta \leq \theta_\gamma$ , and we need to show that  $\eta \leq \kappa_\beta^\gamma$  is regular in  $M_\beta^\gamma$  iff it is regular in  $K^*$ . This will follow by showing that  $\mathcal{P}(\kappa_\beta^\gamma) \cap K^* = \mathcal{P}(\kappa_\beta^\gamma) \cap M_\beta^\gamma$ .

If  $X \in \mathcal{P}(\kappa_\beta^\gamma) \cap K^*$ , by theorem 8  $X \in \text{lp}(N)$  for some short  $D$ -mouse  $N \in K^*$ . By minimality of  $M_0^0$  and its iterates,  $N <_D M_\beta^\gamma$ , so there are iterates  $J_\delta[D, F^*]$  and  $J_{\delta'}[D, F^*]$  of  $N, M_\beta^\gamma$  respectively, such that  $\delta \leq \delta'$ .  $X \in \text{lp}(N)$  so  $X$  is fixed in the iteration, hence  $X \in J_\delta[D, F^*] \subseteq J_{\delta'}[D, F^*]$ . We can also assume that the iteration of  $M_\beta^\gamma$  is done with measurables  $\geq \kappa_\beta^\gamma$  so  $\mathcal{P}(\kappa_\beta^\gamma) \cap M_\beta^\gamma = \mathcal{P}(\kappa_\beta^\gamma) \cap J_{\delta'}[D, F^*]$  so  $X \in M_\beta^\gamma$ .

If  $X \in \mathcal{P}(\kappa_\beta^\gamma) \cap M_\beta^\gamma$ ,  $M_\beta^\gamma = J_\alpha[D, \tilde{F}]$ , then there is some  $\eta \in M_\beta^\gamma$  such that  $X$  is definable in  $J_\eta[D, \tilde{F}]$  from a finite set of parameters  $p$ , so  $X \in H_1^{M_\beta^\gamma}(p)$  i.e.  $X$  is in mouse which is smaller than  $M_0^0$ , so by minimality it is in  $K^*$ , so  $X \in \mathcal{P}(\kappa_\beta^\gamma) \cap K^*$ . Claim 15  $\square$

#### 4. THE $C^*$ OF A SHORT $L[\mathcal{U}]$

Our next goal is to generalize theorem 5.16 of [1] which states that if  $V = L^\mu$  then  $C^* = M_{\omega^2}[E]$  where  $M_{\omega^2}$  is the  $\omega^2$ th iterate of  $V$  and  $E = \{\kappa_{\omega \cdot n} \mid 1 \leq n < \omega\}$  is the sequence of the  $\omega \cdot n$  iteration points. As noted in [1], this can be generalized using the same proof to a model with finitely many measurable cardinals  $\{\kappa^0, \dots, \kappa^m\}$ , using the Prikry sequences  $E^i = \{\kappa_{\omega \cdot n}^i \mid 1 \leq n < \omega\}$  for  $i \leq m$ . However, if we want infinitely many measurables,  $\langle \kappa^i \mid i < \omega \rangle$ , we'd need to show that the sequence  $\langle E^i \mid i < \omega \rangle$  is in  $C^*$ , and not only every  $E^i$ , and at first sight it is not clear that this is the case – note for example that by [1, theorem 5.10], the sequence of measurables  $\langle \kappa^i \mid i < \omega \rangle$  is necessarily *not* in  $C^*$ . We will however show that using the previous theorem, we can construct the sequence  $\langle E^i \mid i < \omega \rangle$ , and even longer (though still short) sequences, in  $C^*$ .

**Theorem 17.** *Assume  $V = L[\mathcal{U}]$  where  $\mathcal{U} = \langle U_\gamma \mid \gamma < \chi \rangle$  is a short sequence of measures on the increasing measurables  $\langle \kappa^\gamma \mid \gamma < \chi \rangle$ . Iterate  $V$  according to  $\mathcal{U}$  where each measurable is iterated  $\omega^2$  many times, to obtain  $\langle M_\alpha^\gamma \mid \gamma < \chi, \alpha \leq \omega^2 \rangle$ , with iteration points  $\langle \kappa_\alpha^\gamma \mid \gamma < \chi, \alpha \leq \omega^2 \rangle$ , and set  $M^\chi$  as the directed limit of this iteration. Then  $C^* = M^\chi[\langle E^\gamma \mid \gamma < \chi \rangle]$  where  $E^\gamma = \langle \kappa_{\omega \cdot n}^\gamma \mid 1 \leq n < \omega \rangle$ .*



*Remark 18.* This theorem appears already in [6], where the claim is that the “arguments in [1] can be straightforwardly deployed for each cardinal in turn”. However, as we noted above, dealing with each measurable separately is *prima facie* not sufficient to show that the sequence of Prikry sequences  $\langle E^\gamma \mid \gamma < \chi \rangle$  is in  $C^*$ , which is crucial for the argument. Hence we include a detailed proof of this theorem here, which addresses this issue.

*Proof.* First we prove that  $\langle E^\gamma \mid \gamma < \chi \rangle \in C^*$  by giving a recursive definition for the sequence inside  $C^*$ . Denote for every  $\gamma$  the filter on  $\kappa_{\omega,2}^\gamma$  generated by end-segments of  $E^\gamma$  by  $\bar{U}_\gamma$ . Assume we’ve shown that  $\langle E^\gamma \mid \gamma < \eta \rangle \in C^*$ , so also  $D_\eta = \langle \bar{U}_\gamma \mid \gamma < \eta \rangle \in C^*$ , hence by theorem 12  $K[D_\eta] = (K[D_\eta])^{C^*}$ . Denote  $K^\eta = K[D_\eta]$  and  $N^\eta = L[D_\eta, U_\eta]$ . Let  $N_\alpha^\eta$  be the  $\alpha$ th iteration of  $N^\eta$  by  $U_\eta$ .

*Claim 19.* For every  $\alpha$ ,  $\mathcal{P}(\kappa_\alpha^\eta) \cap N_\alpha^\eta = \mathcal{P}(\kappa_\alpha^\eta) \cap K^\eta$ . In particular, for every  $\delta \leq \kappa_\alpha^\eta$   $N_\alpha^\eta \models \delta$  is regular iff  $K^\eta \models \delta$  is regular.

*Proof.* Let  $P$  be a  $D_\eta$ -mouse such that  $\mathcal{P}(\kappa_\alpha^\eta) \cap K^\eta \in \text{lp}(P)$ . We can assume that  $P = J_\mu[D_\eta, F]$  where  $F$  is a single measure in  $P$ . Let  $\theta$  be such that  $\mathcal{P}(\kappa_\alpha^\eta) \cap N_\alpha^\eta \in Q := J_\theta[D_\eta, U_\eta]$ . It is enough to show that  $\mathcal{P}(\kappa_\alpha^\eta) \cap P = \mathcal{P}(\kappa_\alpha^\eta) \cap Q$ . We iterate  $P, Q$  by  $F, U_\eta$  respectively  $\lambda$  many times for some regular  $\lambda > |P| + |Q|$  and get iterations up to  $\bar{P} = J_{\mu'}[D_\eta, \text{CU}(\lambda)]$  and  $\bar{Q} = J_{\theta'}[D_\eta, \text{CU}(\lambda)]$ . Since the iterations were done above  $\kappa_\alpha^\eta$ , and in fact above  $(2^{\kappa_\alpha^\eta})^P$  and  $(2^{\kappa_\alpha^\eta})^Q$ , we get that

$$\mathcal{P}(\kappa_\alpha^\eta) \cap P = \mathcal{P}(\kappa_\alpha^\eta) \cap \bar{P} = \mathcal{P}(\kappa_\alpha^\eta) \cap \bar{Q} = \mathcal{P}(\kappa_\alpha^\eta) \cap Q$$

as required. Claim 19□

*Claim 20.*  $E^\eta = \langle \kappa_{\omega \cdot n}^\eta \mid 1 \leq n < \omega \rangle$  is the sequence of the first  $\omega$  ordinals greater than  $\sup \{ \sup E^\gamma \mid \gamma < \eta \}$  which are regular in  $K^\eta$  but of cofinality  $\omega$  in  $V$ .

*Proof.* For every  $\eta$  and  $\alpha$ ,  $\kappa_\alpha^\eta$  is regular in  $M_\alpha^\eta$ , so by the previous claim it is regular in  $K^\eta$  as well, and for every  $n \geq 1$   $\kappa_{\omega \cdot n}^\eta = \sup \{ \kappa_{\omega \cdot (n-1) + k}^\eta \mid k < \omega \}$  so is of cofinality  $\omega$  in  $V$ . Now note that for every  $\gamma < \eta$ ,  $\sup E^\gamma = \kappa_{\omega,2}^\gamma < \kappa^\eta$  (by the properties of iterations), so also  $\sup \{ \sup E^\gamma \mid \gamma < \eta \} < \kappa^\eta = \kappa_0^\eta$ . Hence by claim 19 every  $\delta \in (\{ \sup E^\gamma \mid \gamma < \eta \}, \kappa^\eta)$  is regular in  $K^\eta$  iff it is regular in  $L[D_\eta, U_\eta]$ , but since  $V = L[\mathcal{U}]$ , this is iff  $\delta$  is regular in  $V$ . For  $\eta \in (\kappa_{\omega \cdot n}^\eta, \kappa_{\omega \cdot (n+1)}^\eta)$  ( $n \geq 0$ ), by claim 19 we get that  $\eta$  is regular in  $K^\eta$  iff it is regular in  $N_{\omega \cdot (n+1)}^\eta$ , in which case we get by the same proof as that of claim 14 that either  $\eta = \kappa_{\omega \cdot n + k}^\eta$  for some  $k > 0$ , or  $\text{cf}^V(\eta) = (\kappa^\eta)^+$ , and in either case  $\eta$  is not of cofinality  $\omega$  in  $V$ . So we get that indeed the first  $\omega$  ordinals  $> \sup \{ \sup E^\gamma \mid \gamma < \eta \}$  which are regular in  $K^\eta$  but of cofinality  $\omega$  in  $V$  are  $\langle \kappa_{\omega \cdot n}^\eta \mid 1 \leq n < \omega \rangle$ . Claim 20□

This claim allows us to define  $E^\eta$  in  $C^*$  by recursion, so indeed  $\langle E^\gamma \mid \gamma < \chi \rangle \in C^*$ . Now we claim that indeed  $C^* = M^\chi[\langle E^\gamma \mid \gamma < \chi \rangle]$ . First of all, if we again denote by  $\bar{U}_\gamma$

the filter on  $\kappa_{\omega,2}^\gamma$  generated by end-segments of  $E^\gamma$ , then  $\langle \bar{U}_\gamma \mid \gamma < \chi \rangle \in C^*$  and  $M^\chi = L[\langle \bar{U}_\gamma \mid \gamma < \chi \rangle]$ , so  $M^\chi[\langle E^\gamma \mid \gamma < \chi \rangle] \subseteq C^*$ . On the other hand, the only ordinals which are of cofinality  $\omega$  in  $V$  but not in  $M^\chi$  are those whose cofinality in  $M^\chi$  is in  $E^\gamma \cup \{\sup E^\gamma\}$  for some  $\gamma < \chi$ , so in  $M^\chi[\langle E^\gamma \mid \gamma < \chi \rangle]$  we can correctly compute  $C^*$  by replacing the quantifier  $Q_\omega^{\text{cf}}(\alpha)$  with  $\text{cf}(\alpha) \in \{\omega\} \cup \bigcup \{E^\gamma \mid \gamma < \chi\} \cup \{\sup E^\gamma \mid \gamma < \chi\}$ , so  $C^* \subseteq M^\chi[\langle E^\gamma \mid \gamma < \chi \rangle]$  as required. Theorem 17□

**Theorem 21.** *Assume  $V = L[\mathcal{U}]$  where  $\mathcal{U}$  is as above,  $M^\chi$  as above. Let  $A \in V$ ,  $A \subseteq \chi$ , and set for  $\gamma < \chi$*

$$\hat{E}^\gamma = \begin{cases} \langle \kappa_{\omega,n}^\gamma \mid 0 \leq n < \omega \rangle & \gamma \in A \\ \langle \kappa_{\omega,n}^\gamma \mid 1 \leq n < \omega \rangle & \gamma \notin A \end{cases}$$

and let  $G$  be generic for the forcing over  $V$  adding a Prikry sequence  $G_\alpha \subseteq \kappa^\alpha$  for every  $\alpha \in A$ . Then  $(C^*)^{V[G]} = M^\chi[\langle \hat{E}^\gamma \mid \gamma < \chi \rangle]$ .

*Proof.* We repeat the proof of theorem 17. For the  $\supseteq$  direction, we note that the only difference is that in claim 20, for  $\eta \in A$ , the sequence of the first  $\omega$  ordinals greater than  $\sup \{\sup E^\gamma \mid \gamma < \eta\}$  which are regular in  $K^\eta$  but of cofinality  $\omega$  in  $V[G]$  is now  $\langle \kappa_{\omega,n}^\eta \mid 0 \leq n < \omega \rangle$  instead of  $\langle \kappa_{\omega,n}^\eta \mid 1 \leq n < \omega \rangle$ , since also  $\kappa^\eta$  has cofinality  $\omega$  in  $V[G]$ , but otherwise there is no difference in cofinality  $\omega$  between  $V$  and  $V[G]$  in this interval of ordinals. For the  $\subseteq$  direction,  $V[G] \models \text{cf}(\eta) = \omega$  iff  $V \models \text{cf}(\eta) \in \{\omega\} \cup \{\kappa^\alpha \mid \alpha \in A\}$ , and by the same arguments we get that  $(C^*)^{V[G]}$  can be computed in  $M^\chi[\langle \hat{E}^\gamma \mid \gamma < \chi \rangle]$ . □

*Remark 22.* (1) This does not imply that  $A \in (C^*)^{V[G]}$  – it is clear from the proof that  $(C^*)^{V[G]}$  might not distinguish which option was used in defining  $\hat{E}^\gamma$ .  
 (2) If  $A$  is finite then  $(C^*)^{V[G]} = (C^*)^V$ .  
 (3) If for example  $A = \omega \leq \chi$ , then the sequence  $\langle \kappa^i \mid i < \omega \rangle$  can be computed from  $\langle \hat{E}^\gamma \mid \gamma < \chi \rangle$ , so is in  $(C^*)^{V[G]}$ , while as we already noted it is not in  $(C^*)^V$ , so  $(C^*)^{V[G]} \neq (C^*)^V$ .

## 5. INNER MODELS FOR SHORT SEQUENCES OF MEASURES IN $C^*$

Our last task is to generalize theorem 5.6 of [1] from the case of an inner model with one measurable to the case of an inner model with a short sequence of measurables.

**Lemma 23.** *Assume  $M = L[\mathcal{U}]$  where  $\mathcal{U} = \langle U_\gamma \mid \gamma < \chi \rangle$  is in  $M$  a short sequence of measures on the increasing sequence  $\langle \kappa^\gamma \mid \gamma < \chi \rangle$ . Denote for each  $\gamma < \chi$   $\xi^\gamma = \left[ (\kappa^\gamma)^+ \right]^M$  and let  $f_\mathcal{U} : \chi \rightarrow 2$  be defined by  $f_\mathcal{U}(\gamma) = 0 \iff \text{cf}^V(\xi^\gamma) = \omega$ . If  $f_\mathcal{U} \in C^*$  then  $C^*$  contains an iteration of  $M$  by  $\mathcal{U}$ .*

*Proof.* We combine the methods used in the previous sections – from section 3 we take the method of recursively defining the sequence  $\langle \lambda_\gamma \mid \gamma < \chi \rangle$  which gives the lengths of

the iterations, and from section 4 we take the method of recursively defining the sequence of iterated measures  $\langle \bar{U}_\gamma \mid \gamma < \chi \rangle$ .

Let  $\Theta$  be some regular cardinal (in  $V$ ) greater than  $\sup \{ (2^{\kappa^\gamma})^V \mid \gamma < \chi \}$ . We define simultaneously sequences  $\langle \lambda_\gamma \mid \gamma < \chi \rangle$  and  $\langle \bar{U}_\gamma \mid \gamma < \chi \rangle$  recursively such that  $\lambda_0 > \Theta$  and for every  $\gamma < \chi$ :

- (1)  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$  and  $\langle \bar{U}_{\gamma'} \mid \gamma' < \gamma \rangle$  are in  $C^*$ ;
- (2)  $K^\gamma := K [\langle \bar{U}_{\gamma'} \mid \gamma' < \gamma \rangle] = (K [\langle \bar{U}_{\gamma'} \mid \gamma' < \gamma \rangle])^{C^*}$ ;
- (3)  $C^* \models \lambda_\gamma$  is a regular cardinal;
- (4)  $\gamma' < \gamma$  implies  $\lambda_{\gamma'} < \lambda_\gamma$ ;
- (5)  $\forall \beta < \lambda_\gamma \exists \eta \in (\beta, \lambda_\gamma)$  such that  $K^\gamma \models \eta$  is a regular cardinal, and  $\text{cf}^V(\eta) = \omega \iff f_U(\gamma) = 1$  (so  $\text{cf}^V(\eta) = \omega \iff \text{cf}^V(\xi^\gamma) \neq \omega$ ).

Assume we've defined  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle, \langle \bar{U}_{\gamma'} \mid \gamma' < \gamma \rangle \in C^*$  for some  $\gamma < \chi$ . By essentially the same argument as in the proof of theorem 12 we show that every regular cardinal in  $V$  satisfies 3-5, where the main differences are that here in  $K^\gamma$  we take the iterations of the measures  $\langle \bar{U}_{\gamma'} \mid \gamma' < \gamma \rangle$  - instead of the original ones, and that we need to use  $f_U$  to know which iteration points to choose, instead of the single  $\delta_M$  we had there. So in  $C^*$  we can define  $\lambda_\gamma$  to be the first ordinal  $> \sup \langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle$  (or  $> \Theta$  if  $\gamma = 0$ ) satisfying 3-5. Again by the same argument as before we get that in fact  $\lambda_\gamma$  is the image of  $\kappa^\gamma$  under the iteration of  $M$  by the measures  $\langle U_{\gamma'} \mid \gamma' \leq \gamma \rangle$  according to  $\langle \lambda_{\gamma'} \mid \gamma' < \gamma \rangle \frown \langle \theta_\gamma \rangle$ , for some  $\theta_\gamma$  i.e.  $\lambda_\gamma = \kappa_{\theta_\gamma}^\gamma$ . Now we let

$$E_\gamma = \begin{cases} \left\{ \eta < \lambda_\gamma \mid \eta \text{ is regular in } K^\gamma \text{ and } \text{cf}^V(\eta) = \omega \right\} & f_U = 1 \\ \left\{ \eta < \lambda_\gamma \mid \eta \text{ is regular in } K^\gamma \text{ and } \text{cf}^V(\eta) \neq \omega \right\} & f_U = 0 \end{cases}$$

and get (using the appropriate variations of claims 14 and 15) that  $E_\gamma \in C^*$  generates the iterated measure  $\bar{U}_\gamma$  on  $\lambda_\gamma = \kappa_{\theta_\gamma}^\gamma$ .

Since all the definitions were done in a recursive way inside  $C^*$ , we can proceed through limit stages up to  $\chi$  to get that the sequences are in  $C^*$ , hence the iteration of  $M$  by  $\langle \theta_\gamma \mid \gamma < \chi \rangle$  (which is not in necessarily in  $C^*$ ) is in fact determined by  $\langle \lambda_\gamma \mid \gamma < \chi \rangle$  and  $\langle \bar{U}_\gamma \mid \gamma < \chi \rangle$  which are in  $C^*$ , so this iteration is in  $C^*$ .  $\square$

**Corollary 24.** *If there is an inner model with a short sequence of measures of order type  $\chi$  then there is such an inner model inside  $C^*$ .*

*Proof.* Let  $M = L[\mathcal{U}]$  where  $\mathcal{U}$  is a short sequence of measures of order type  $\chi$ , and let  $f_U : \chi \rightarrow 2$  be defined by  $f_U(\gamma) = 0 \iff \text{cf}^V(\xi^\gamma) = \omega$ . We will show that there is  $\tilde{\mathcal{U}}$  which is also of order type  $\chi$  such that  $f_{\tilde{\mathcal{U}}} \in C^*$ , so we'll be done by the previous lemma applied to  $L[\tilde{\mathcal{U}}]$ .

**Lemma 25.** *Let  $\alpha = \omega^\beta$ , and let  $f : \alpha \rightarrow 2$ . Then there is  $A \subseteq \alpha$ ,  $\text{otp } A = \alpha$  such that  $|f \upharpoonright A| = 1$ .*

*Proof.* Otherwise  $\alpha$  is the disjoint union of two sets of order-type  $< \alpha$ ,  $A_i = f^{-1}\{i\}$  for  $i = 0, 1$ . We prove that this is impossible. If  $\beta = 0$  then  $\alpha = 1$  so this is clear. If  $\beta = \gamma + 1$ , consider  $\omega^\beta = \omega^\gamma \cdot \omega$ . So  $\max\{\text{otp } A_0, \text{otp } A_1\} < \omega^\gamma \cdot n$  for some  $n$ , so  $\text{otp}(A_0 \cup A_1) < \omega^\gamma \cdot n + \omega^\gamma \cdot n = \omega^\gamma \cdot 2n < \omega^\gamma \cdot \omega = \omega^\beta$  by contradiction. For  $\beta$  limit,  $\omega^\beta = \sup\{\omega^\gamma \mid \gamma < \beta\}$ ,  $\max\{\text{otp } A_0, \text{otp } A_1\} < \omega^\gamma$  for some  $\gamma$ . So  $\text{otp}(A_0 \cup A_1) < \omega^\gamma + \omega^\gamma < \omega^{\gamma+1} < \omega^\beta$  by contradiction.  $\square$

Now by Cantor normal form, there are  $\beta_0 \geq \beta_1 \geq \dots \geq \beta_n \geq 0$  such that  $\chi = \omega^{\beta_0} + \dots + \omega^{\beta_n}$ , so  $\chi$  is partitioned into the finitely many successive segments  $I_0 = [0, \omega^{\beta_0})$ ,  $I_1 = [\omega^{\beta_0}, \omega^{\beta_0} + \omega^{\beta_1})$ ,  $\dots$ ,  $I_n = [\omega^{\beta_0} + \dots + \omega^{\beta_{n-1}}, \omega^{\beta_0} + \dots + \omega^{\beta_n})$ . Each segment  $I_k$  is order isomorphic to the corresponding  $\omega^{\beta_k}$ , so there is  $A_k \subseteq I_k$  such that  $\text{otp } A_k = \omega^{\beta_k}$  and  $|f_{\mathcal{U}} \upharpoonright A_k| = 1$ . Clearly  $\text{otp}(\bigcup_{i=0}^n A_k) = \chi$  so let  $\phi : \chi \rightarrow \bigcup_{i=0}^n A_k$  be the order isomorphism and define  $\tilde{\mathcal{U}}$  by  $\tilde{U}_\gamma = U_{\phi(\gamma)}$ . Then  $f_{\tilde{\mathcal{U}}} = (f_{\mathcal{U}} \upharpoonright \bigcup_{i=0}^n A_k) \circ \phi$  is simply the finite union of constant functions, so it is indeed in  $C^*$  as required.  $\square$

#### REFERENCES

1. Juliette Kennedy, Menachem Magidor, and Jouko Väänänen, *Inner models from extended logics: Part 1*, arXiv preprint (2020).
2. Peter Koepke, *A theory of short core models and applications*, Ph.D. thesis, University of Freiburg, 1983.
3. ———, *Some applications of short core models*, Annals of pure and applied logic **37** (1988), no. 2, 179–204.
4. Philipp Lücke and Philipp Schlicht, *Measurable cardinals and good  $\Sigma_1(\kappa)$ -wellorderings*, Mathematical Logic Quarterly **64** (2018), no. 3, 207–217.
5. William J. Mitchell, *The core model for sequences of measures. I*, Mathematical Proceedings of the Cambridge Philosophical Society, vol. 95, Cambridge University Press, 1984, pp. 229–260.
6. P.D Welch, *Closed and unbounded classes and the H\"artig quantifier model*, The Journal of Symbolic Logic, 1–22.

EINSTEIN INSTITUTE OF MATHEMATICS  
 UNIVERSITY OF JERUSALEM  
 EDMOND J. SAFRA CAMPUS, GIVAT RAM  
 JERUSALEM 91904, ISRAEL

*Email address:* ur.yaar@mail.huji.ac.il