

STRONG MORITA EQUIVALENCES FOR COMPLETELY POSITIVE LINEAR MAPS AND GNS- C^* -CORRESPONDENCES

KAZUNORI KODAKA

ABSTRACT. We will consider the set of all completely positive linear maps from a unital C^* -algebra to the C^* -algebra of all (bounded) adjointable right Hilbert C^* -module maps, which are automatically bounded, on a right Hilbert C^* -module and we will introduce strong Morita equivalence for elements in this set. In this paper, we will give the following result: If two classes of two unital C^* -algebras are strongly Morita equivalent, respectively, then we can construct a bijective correspondence between two sets of all strong Morita equivalence classes of completely positive linear maps given as above. Furthermore, we will discuss the relation between strong Morita equivalence for completely positive linear maps and strong Morita equivalence for GNS- C^* -correspondences.

1. INTRODUCTION

In the previous paper [8], we introduce the notion of strong Morita equivalence for completely positive linear maps from a C^* -algebra to the C^* -algebra of all bounded linear operators on a Hilbert space using their minimal Stinespring representations and [4, Definition 2.1] introduced by Echterhoff and Raeburn.

In this paper, we consider completely positive linear maps from a unital C^* -algebra C to the C^* -algebra of all adjointable right Hilbert B -module maps, which are automatically bounded by Blackadar [1, Proposition 13.2.2], on a right Hilbert B -module F , where B is a unital C^* -algebra. We denote by $\text{CP}_B(C)$, the set of all such completely positive linear maps. We introduce the notion of strong Morita equivalence for elements in $\text{CP}_B(C)$ using the KSGNS-representation given in Lance [9, Theorem 5.6]. In order to do this, we have to extend [4, Definition 2.1] to non-degenerate representations of a unital C^* -algebra on a right Hilbert C^* -module. This extension will be done in Section 2.

In Section 3, we will give a result, which is similar to [8, Corollary 6.6] in the following way: Let ψ be a completely positive linear map from a unital C^* -algebra D to $\mathbf{B}_B(F)$, where B is a unital C^* -algebra and $\mathbf{B}_B(F)$ is the C^* -algebra of all adjointable rightable right Hilbert B -module maps on a right Hilbert B -module F . Let C be a unital C^* -algebra which is strongly Morita equivalent to D . Then in the same way as in [8, Section 6], using ψ , we will construct a completely positive linear map ϕ from C to $\mathbf{B}_B(F \otimes \mathbf{C}^n)$, where n is some positive integer. We will show that this construction induces a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_B(C)/\sim$, where $\text{CP}_B(D)/\sim$ and $\text{CP}_B(C)/\sim$ denote the sets of all strong Morita equivalence classes in $\text{CP}_B(D)$ and $\text{CP}_B(C)$, respectively. Furthermore, we assume that A is strongly Morita equivalent to B . We will show that there exists a bijective correspondence between $\text{CP}_B(C)/\sim$ and $\text{CP}_A(C)/\sim$. Thus, we obtain the following result: If C and A are strongly Morita equivalent to D and B , respectively, then there exists a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_A(C)/\sim$.

2010 *Mathematics Subject Classification.* 46L05.

Key words and phrases. C^* -algebras, completely positive linear maps, equivalence bimodules, Hilbert C^* -modules, strong Morita equivalence.

In Section 4, we will discuss the relation between strong Morita equivalence for completely positive linear maps and strong Morita equivalence for GNS- C^* -correspondences induced by completely positive linear maps. It is natural to do it since GNS- C^* -correspondences induced by completely positive linear maps are constructed in the same way as in the KSGNS-construction.

Let A be a unital C^* -algebra and we denote by id_A and 1_A , the identity map on A and the unit element in A , respectively. If no confusion arises, we simply denote them by id and 1 , respectively.

For each $n \in \mathbf{N}$, let $M_n(A)$ be the $n \times n$ -matrix algebra over A . We identify $M_n(A)$ with $A \otimes M_n(\mathbf{C})$. Let I_n be the unit element in $M_n(\mathbf{C})$. For each $a \in M_n(A)$, we denote by a_{ij} the $i \times j$ -entry of a .

Let A and B be unital C^* -algebras and X an $A - B$ -equivalence bimodule. We denote its left A -action and right B -action on X by $a \cdot x$ and $x \cdot b$ for any $a \in A$, $b \in B$, $x \in X$, respectively. Let \tilde{X} be the dual $B - A$ -equivalence bimodule of X and let \tilde{x} denote the element in \tilde{X} associated to the element $x \in X$.

For right Hilbert A -modules E and F , let $\mathbf{B}_A(E, F)$ be the space of all bounded adjointable right A -module maps from E to F and if $E = F$, we denote $\mathbf{B}_A(E, F)$ by $\mathbf{B}_A(E)$. In this case, $\mathbf{B}_A(E)$ is a unital C^* -algebra.

2. REPRESENTATIONS OF A UNITAL C^* -ALGEBRA ON RIGHT HILBERT MODULES AND STRONG MORITA EQUIVALENCE

Let C and A be unital C^* -algebras.

Definition 2.1. We say that (π_C, E) is a *representation* of C on a right Hilbert A -module E if π_C is a homomorphism of C to $\mathbf{B}_A(E)$. Furthermore, we say that (π_C, E) is *non-degenerate* if $\overline{\pi_C(C)E} = E$.

Let B, C, D be unital C^* -algebras. Let (π_C, E) and (π_D, F) be non-degenerate representations of C and D on Hilbert B -modules E and F , respectively.

Definition 2.2. We say that (π_C, E) is *strongly Morita equivalent* to (π_D, F) by if there exist a $C - D$ -equivalence bimodule Y and a linear map π_Y from Y to $\mathbf{B}_B(F, E)$ satisfying the following conditions:

- (1) $\pi_Y(y)\pi_Y(z)^* = \pi_C(\langle y, z \rangle)$,
 - (2) $\pi_Y(y)^*\pi_Y(z) = \pi_D(\langle y, z \rangle_D)$,
 - (3) $\pi_Y(c \cdot y \cdot d) = \pi_C(c)\pi_Y(y)\pi_D(d)$ for any $y, z \in Y$, $c \in C$, $d \in D$.
- We denote it by $(\pi_C, E) \sim (\pi_D, F)$.

This definition is an extension of [4, Definition 2.1] in the case that C^* -algebras are unital.

Proposition 2.1. *Strong Morita equivalence for non-degenerate representations of unital C^* -algebras on right Hilbert C^* -modules is an equivalence relation.*

Proof. Let (π_C, E) be a non-degenerate representation of a unital C^* -algebra C on a right Hilbert B -module E , where B is a unital C^* -algebra. We regard C as the trivial $C - C$ -equivalence bimodule in the natural way. We denote it by Y_0 . Let π_{Y_0} be the linear map from Y_0 to $\mathbf{B}_B(E)$ defined by $\pi_{Y_0} = \pi_C$. Then π_{Y_0} satisfies Conditions (1)-(3) in Definition 2.2. Hence (π_C, E) is strongly Morita equivalent to itself.

Let (π_D, F) be a non-degenerate representation of a unital C^* -algebra D on a right Hilbert B -module F . We suppose that (π_C, E) is strongly Morita equivalent to (π_D, F) with respect to a $C - D$ -equivalence bimodule Y and a linear map π_Y from Y to $\mathbf{B}_B(F, E)$. Let $\pi_{\tilde{Y}}$ be the linear map from \tilde{Y} to $\mathbf{B}_B(E, F)$ defined by

$$\pi_{\tilde{Y}}(\tilde{y}) = \pi_Y(y)^*$$

for any $y \in Y$. Then by routine computations, we can see that $\pi_{\tilde{Y}}$ satisfies Conditions (1)-(3) in Definition 2.2. Thus (π_D, F) is strongly Morita equivalent to (π_C, E) .

Let (π_L, G) be a non-degenerate representation of a unital C^* -algebra L on a right Hilbert B -module G . We suppose that (π_C, E) is strongly Morita equivalent to (π_D, F) with respect to a $C - D$ -equivalence bimodule Y and a linear map π_Y from Y to $\mathbf{B}_B(F, E)$ satisfying Conditions (1)-(3) in Definition 2.2. Also, we suppose that (π_D, F) is strongly Morita equivalent to (π_L, G) with respect to a $D - L$ -equivalence bimodule W and a linear map π_W from W to $\mathbf{B}_B(G, F)$ satisfying Conditions (1)-(3) in Definition 2.2. We show that (π_C, E) is strongly Morita equivalent to (π_L, G) . Clearly C is strongly Morita equivalent to L with respect to the $C - L$ -equivalence bimodule $Y \otimes_D W$. Let $\pi_{Y \otimes_D W}$ be the linear map from $Y \otimes_D W$ to $\mathbf{B}_B(G, E)$ defined by

$$\pi_{Y \otimes_D W}(y \otimes w) = \pi_Y(y)\pi_W(w)$$

for any $y \in Y, w \in W$. Then by routine computations, we can see that $\pi_{Y \otimes_D W}$ satisfies Conditions (1)-(3) in Definition 2.2. Therefore, we obtain the conclusion. \square

Let C, A be unital C^* -algebras. For $i = 1, 2$, let (π_i, E_i) be a non-degenerate representation of C on a right Hilbert A -module E_i .

Lemma 2.2. *With the above notation, if (π_1, E_1) and (π_2, E_2) are unitarily equivalent, they are strongly Morita equivalent.*

Proof. Since (π_1, E_1) and (π_2, E_2) are unitarily equivalent, there is an isometry $u \in \mathbf{B}_B(E_1, E_2)$ such that $\pi_2 = \text{Ad}(u) \circ \pi_1$. Let $Y_0 (= C)$ be the trivial $C - C$ -equivalence bimodule. Let π_{Y_0} be the linear map from Y_0 to $\mathbf{B}_B(E_2, E_1)$ defined by

$$\pi_{Y_0}(y) = \pi_1(y)u^*$$

Then π_{Y_0} satisfies Conditions (1)-(3) in Definition 2.2. Thus, (π_1, E_1) and (π_2, E_2) are strongly Morita equivalent. \square

Let C, D and B be unital C^* -algebras and we suppose that C and D are strongly Morita equivalent with respect to a $C - D$ -equivalence bimodule Y . Let (π_D, F) be a non-degenerate representation of D on a right Hilbert B -module F . Modifying [4, Section 2], we construct a non-degenerate representation (π_C, E) of C on a right Hilbert B -module E , which is strongly Morita equivalent to (π_D, F) . We regard F as a Hilbert $D - B$ -bimodule using the representation (π_D, F) . Let E be a right Hilbert B -module by setting

$$E = Y \otimes_D F.$$

Let π_C be the map from C to $\mathbf{B}_B(E)$ defined by

$$\pi_C(c)(y \otimes f) = (c \cdot y) \otimes f$$

for any $c \in C, y \in Y, f \in F$. Then by [3, Proposition 1.7(i)] and easy computations, (π_C, E) is a non-degenerate representation of C . We will show that (π_C, E) is strongly Morita equivalent to (π_D, F) .

Lemma 2.3. *With the above notation, (π_C, E) is strongly Morita equivalent to (π_D, F) .*

Proof. It suffices to show that there is a linear map π_Y from Y to $\mathbf{B}_B(F, E)$ satisfying Conditions (1)-(3) in Definition 2.2. Let π_Y be the linear map from Y to the space of all bounded right B -module maps from F to E defined by

$$\pi_Y(y)f = y \otimes f$$

for any $y \in Y$, $f \in F$. We show that $\pi_Y(y) \in \mathbf{B}_B(F, E)$ for any $y \in Y$ and that π_Y satisfies Conditions (1)-(3) in Definition 2.2. For any $y, z \in Y$, $f, g \in F$.

$$\begin{aligned} \langle z \otimes f, \pi_Y(y)g \rangle_B &= \langle z \otimes f, y \otimes g \rangle_B = \langle f, \pi_D(\langle z, y \rangle_D)g \rangle_B \\ &= \langle \pi_D(\langle y, z \rangle_D)f, g \rangle_B. \end{aligned}$$

Thus $\pi(y) \in \mathbf{B}_B(F, E)$ for any $y \in Y$ and $\pi(y)^*(z \otimes f) = \pi_D(\langle y, z \rangle_D)f$ for any $y, z \in Y$, $f \in F$. Then for any $y, y_1, z \in Y$, $f \in F$,

$$\pi_Y(y)\pi_Y(y_1)^*(z \otimes f) = \pi_Y(y)\pi_D(\langle y_1, z \rangle_D)f = y \otimes \pi_D(\langle y_1, z \rangle_D)f.$$

On the other hand,

$$\pi_C(\langle y, y_1 \rangle)(z \otimes f) = (\langle y, y_1 \rangle \cdot z) \otimes f = y \cdot \langle y_1, z \rangle_D \otimes f = y \otimes \pi_D(\langle y_1, z \rangle_D)f.$$

Hence $\pi_Y(y)\pi_Y(y_1)^* = \pi_C(\langle y, y_1 \rangle)$. Also, for any $y, y_1 \in Y$, $f \in F$,

$$\pi_Y(y)^*\pi_Y(y_1)f = \pi_Y(y)^*(y_1 \otimes f) = \pi_D(\langle y, y_1 \rangle_D)f.$$

Hence $\pi_Y(y)^*\pi_Y(y_1) = \pi_D(\langle y, y_1 \rangle_D)$. Furthermore, for any $c \in C$, $d \in D$, $f \in F$, $y \in Y$,

$$\pi_Y(c \cdot y \cdot d)f = c \cdot y \cdot d \otimes f = \pi_C(c)(y \otimes \pi_D(d)f) = \pi_C(c)\pi_Y(y)\pi_D(d)f.$$

Thus $\pi_Y(c \cdot y \cdot d) = \pi_C(c)\pi_Y(y)\pi_D(d)$. Therefore, we obtain the conclusion. \square

We call the above (π_C, E) a non-degenerate representation induced by (π_D, F) and Y .

Let B, C and A be unital C^* -algebras. Let $\text{Rep}_B(C)$ be the set of all non-degenerate representations of C on right Hilbert B -modules. Let $\text{Rep}_B(C)/\sim$ be the set of all strong Morita equivalence classes of elements in $\text{Rep}_B(C)$. Similarly, we define $\text{Rep}_A(C)$ and $\text{Rep}_A(C)/\sim$, respectively. For any $(\pi, E) \in \text{Rep}_A(C)$, $[\pi, E]$ denotes the strong Morita equivalence class of (π, E) . We construct a bijective correspondence Φ between $\text{Rep}_B(C)/\sim$ and $\text{Rep}_A(C)/\sim$.

Let $(\pi, F) \in \text{Rep}_B(C)$ and let Z be a $B - A$ -equivalence bimodule. We consider the right Hilbert A -module $F \otimes_B Z$. Let π^Z be the map from C to $\mathbf{B}_A(F \otimes_B Z)$ defined by

$$\pi^Z(c)(f \otimes z) = \pi(c)f \otimes z$$

for any $f \in F$, $z \in Z$, $c \in C$. Then by easy computations, $(\pi^Z, F \otimes_B Z)$ is a non-degenerate representation of C on $F \otimes_B Z$. Thus $(\pi^Z, F \otimes_B Z) \in \text{Rep}_A(C)$. Let Φ be the map from $\text{Rep}_B(C)/\sim$ to $\text{Rep}_A(C)/\sim$ defined by

$$\Phi([\pi, F]) = [\pi^Z, F \otimes_B Z]$$

for any $[\pi, F] \in \text{Rep}_B(C)/\sim$.

Lemma 2.4. *With the above notation, Φ is well-defined and a bijective correspondence between $\text{Rep}_B(C)/\sim$ and $\text{Rep}_A(C)/\sim$.*

Proof. First, we show that Φ is well-defined. Let $(\pi, F), (\pi_1, F_1)$ be elements in $\text{Rep}_B(C)$ with $[\pi, F] = [\pi_1, F_1]$ in $\text{Rep}_B(C)/\sim$. Then there are a $C - C$ -equivalence bimodule Y and a linear map π_Y from Y to $\mathbf{B}_B(F, F)$ satisfying Conditions (1)-(3) in Definition 2.2. Let $(\pi^Z, F \otimes_B Z)$ and $(\pi_1^Z, F_1 \otimes_B Z)$ be non-degenerate representations of C induced by (π, F) , Y and (π_1, F_1) , Y respectively. Let π_Y^Z be the linear map from Y to $\mathbf{B}_A(F_1 \otimes_B Z, F \otimes_B Z)$ defined by

$$\pi_Y^Z(f \otimes z) = \pi_Y(f) \otimes z = (\pi_Y \otimes \text{id}_Z)(f \otimes z)$$

for any $f \in F$, $z \in Z$. By easy computations, π_Y^Z satisfies Conditions (1)-(3) in Definition 2.2. Thus, Φ is well-defined. Next we show that Φ is a bijective

correspondence between $\text{Rep}_B(C)/\sim$ and $\text{Rep}_A(C)/\sim$. Let Ψ be the map from $\text{Rep}_A(C)/\sim$ to $\text{Rep}_B(C)/\sim$ defined by

$$\Psi([\pi, E]) = [\pi^{\tilde{Z}}, E \otimes_A \tilde{Z}]$$

for any $(\pi, E) \in \text{Rep}_A(C)$. Then for any $(\pi, E) \in \text{Rep}_A(C)$,

$$(\Phi \circ \Psi)([\pi, E]) = \Phi([\pi^{\tilde{Z}}, E \otimes_A \tilde{Z}]) = [(\pi^{\tilde{Z}})^Z, Z \otimes_A \tilde{Z} \otimes_B Z].$$

We show that $[(\pi^{\tilde{Z}})^Z, Z \otimes_A \tilde{Z} \otimes_B Z] = [\pi, E]$ in $\text{Rep}_A(C)/\sim$. Let $Y_0 (= C)$ be the trivial $C - C$ -equivalence bimodule. Let π_{Y_0} be the linear map from Y_0 to the space of all bounded right A -module maps from $E \otimes_A \tilde{Z} \otimes_A Z$ to E defined by

$$\pi_{Y_0}(y)(e \otimes \tilde{z} \otimes z_1) = \pi(y)(e \cdot \langle z, z_1 \rangle_A)$$

for any $e \in E$, $z, z_1 \in Z$. First, we claim that $\pi_{Y_0}(y)$ is adjointable for any $y \in Y_0$. Indeed, for any $y \in Y_0$, $e, e_1 \in E$, $z, z_1 \in Z$,

$$\begin{aligned} \langle e, \pi_{Y_0}(y)(e_1 \otimes \tilde{z} \otimes z_1) \rangle_A &= \langle e, \pi(y)e_1 \cdot \langle z, z_1 \rangle_A \rangle_A = \langle \pi(y)^* e, e_1 \cdot \langle z, z_1 \rangle_A \rangle_A \\ &= \langle \pi(y)^* e, e_1 \rangle_A \langle z, z_1 \rangle_A. \end{aligned}$$

Since Z is full with the right A -valued inner product, there is a finite subset $\{u_i\}_{i=1}^n$ of Z such that $\sum_{i=1}^n \langle u_i, u_i \rangle_A = 1$ by Kajiwara and Watatani [6, Corollary 1.19]. Then

$$\begin{aligned} \langle \sum_{i=1}^n \pi(y)^* e \otimes \tilde{u}_i \otimes u_i, e_1 \otimes z \otimes z_1 \rangle_A &= \sum_{i=1}^n \langle \tilde{u}_i \otimes u_i, \langle \pi(y)^* e, e_1 \rangle_A \cdot \tilde{z} \otimes z \rangle_A \\ &= \sum_{i=1}^n \langle \tilde{u}_i \otimes u_i, [z \cdot \langle e_1, \pi(y)^* e \rangle_A] \tilde{\otimes} z_1 \rangle_A \\ &= \sum_{i=1}^n \langle u_i, \langle \tilde{u}_i, [z \cdot \langle e_1, \pi(y)^* e \rangle_A] \tilde{\otimes} z_1 \rangle_B \rangle_A \\ &= \sum_{i=1}^n \langle u_i, {}_B \langle u_i, [z \cdot \langle e_1, \pi(y)^* e \rangle_A] \rangle \cdot z_1 \rangle_A \\ &= \sum_{i=1}^n \langle {}_B \langle [z \cdot \langle e_1, \pi(y)^* e \rangle_A], u_i \rangle \cdot u_i, z_1 \rangle_A \\ &= \langle \pi(y)^* e, e_1 \rangle_A \langle z, z_1 \rangle_A. \end{aligned}$$

Thus $\pi_{Y_0}(y)$ is adjointable and $\pi_{Y_0}(y)^* = \sum_{i=1}^n \pi(y)^* e \otimes \tilde{u}_i \otimes u_i$ for any $y \in Y_0$. Hence $\pi_{Y_0}(y) \in \mathbf{B}_A(E \otimes_A \tilde{Z} \otimes_B Z, E)$ for any $y \in Y_0$ and π_{Y_0} is a linear map from Y_0 to $\mathbf{B}_A(E \otimes_A \tilde{Z} \otimes_B Z)$. Next, we show that π_{Y_0} satisfies Conditions (1)-(3) in Definition 2.2. For any $y, y_1 \in Y_0$, $e \in E$,

$$\begin{aligned} \pi_{Y_0}(y)\pi_{Y_0}(y_1)^* e &= \sum_{i=1}^n \pi_{Y_0}(y)\pi(y_1)^* \otimes \tilde{u}_i \otimes u_i = \sum_{i=1}^n \pi(y)\pi(y_1)^* e \cdot \langle u_i, u_i \rangle_A \\ &= \pi_C(y, y_1)e. \end{aligned}$$

Thus $\pi_{Y_0}(y)\pi_{Y_0}(y_1)^* = \pi_C(\langle y, y_1 \rangle)$. Also, for any $y, y_1 \in Y_0, e \in E, z, z_1 \in Z$,

$$\begin{aligned}
\pi_{Y_0}(y)^*\pi_{Y_0}(y_1)(e \otimes \tilde{z} \otimes z_1) &= \pi_{Y_0}(y)^*\pi(y_1)e \cdot \langle z, z_1 \rangle_A \\
&= \sum_{i=1}^n \pi(y)\pi(y_1)^*e \cdot \langle z, z_1 \rangle_A \otimes \tilde{u}_i \otimes u_i \\
&= \sum_{i=1}^n \pi(y)^*\pi(y_1)e \otimes [u_i \cdot \langle z_1, z \rangle_A] \tilde{\otimes} u_i \\
&= \sum_{i=1}^n \pi(y^*y_1)e \otimes [{}_B\langle u_i, z_1 \rangle \cdot z] \tilde{\otimes} u_i \\
&= \sum_{i=1}^n \pi(y^*y_1)e \otimes \tilde{z} \otimes {}_B\langle z_1, u_i \rangle \cdot u_i \\
&= \pi(y^*y_1)e \otimes \tilde{z} \otimes z_1 \\
&= (\pi^{\tilde{Z}})^Z(\langle y, y_1 \rangle_C).
\end{aligned}$$

Furthermore, for any $y \in Y_0, c, d \in C, e \in E, z, z_1 \in Z$,

$$\pi_{Y_0}(c \cdot y \cdot d)(e \otimes \tilde{z} \otimes z_1) = \pi(cyd)e \cdot \langle z, z_1 \rangle_A.$$

On the other hand,

$$\begin{aligned}
\pi(c)\pi_{Y_0}(y)(\pi^{\tilde{Z}})^Z(d)(e \otimes \tilde{z} \otimes z_1) &= \pi(c)\pi_{Y_0}(y)(\pi(d)e \otimes \tilde{z} \otimes z_1) \\
&= \pi(c)(\pi(yd)e \cdot \langle z, z_1 \rangle_A) \\
&= \pi(cyd)e \cdot \langle z, z_1 \rangle_A
\end{aligned}$$

since $\pi(c)$ is a right Hilbert A -module map. It follows that $[(\pi^{\tilde{Z}})^Z, E \otimes_A \tilde{Z} \otimes_B Z] = [\pi, E]$ in $\text{Rep}_A(C)/\sim$. Thus $\Phi \circ \Psi = \text{id}$ on $\text{Rep}_A(C)/\sim$. Similarly, $\Psi \circ \Phi = \text{id}$ on $\text{Rep}_B(C)/\sim$. Therefore, Φ is a bijective correspondence between $\text{Rep}_B(C)/\sim$ and $\text{Rep}_A(C)/\sim$. \square

3. A CORRESPONDENCE OF STRONG MORITA EQUIVALENCE CLASSES OF COMPLETELY POSITIVE LINEAR MAPS ON UNITAL C^* -ALGEBRAS

Following Lance [9, Chapter 5], we give the following theorem(KSGNS construction):

Theorem 3.1. ([9, Theorem 5.6]) *Let D nad B be unital C^* -algebras and let F be a right Hilbert B -module. Let ψ be a completely positive linear map from D to $\mathbf{B}_B(F)$.*

(1) *There exists a right Hilbert B -module F_ψ , a representation π_ψ of D on F_ψ and an element V_ψ in $\mathbf{B}_B(F, F_\psi)$ such that*

$$\psi(d) = V_\psi^* \pi_\psi(d) V_\psi$$

for any $d \in D$ and such that $\overline{\pi_\psi(D)V_\psi F} = F_\psi$.

(2) *Let G be a right Hilbert B -module, π a representation of D on G and $W \in \mathbf{B}_B(F, G)$. We suppose that $\psi(d) = W^* \pi(d) W$ for any $d \in D$ and that $\overline{\pi(D)WF} = G$. Then there is an isometry U from F_ψ onto G such that*

$$\pi(d) = U \pi_\psi(d) U^*$$

for any $d \in D$.

Proof. This is immediate by [9, Theorem 5.6]. \square

Remark 3.2. The representation (π_ψ, F_ψ) is non-degenerate since

$$F_\psi = \overline{\pi_\psi(D)V_\psi F} \subset \overline{\pi_\psi(D)F_\psi} \subset F_\psi.$$

Also, the representation (π, G) is non-degenerate.

We call the above $(\pi_\psi, V_\psi, F_\psi)$ a *minimal* KSGNS-representation for ψ .

Let C, D and B be unital C^* -algebras and let F be a right Hilbert B -module. Let ϕ and ψ be completely positive linear maps from C and D to $\mathbf{B}_B(F)$, respectively.

Definition 3.1. We say that ϕ is *strongly Morita equivalent* to ψ if a minimal KSGNS-representation for ϕ is strongly Morita equivalent to that for ψ . We denote it by $\phi \sim \psi$.

Proposition 3.3. *Strong Morita equivalence for completely positive linear maps from unital C^* -algebras to the C^* -algebras of all adjointable right Hilbert modules maps on right Hilbert modules is an equivalence relation.*

Proof. This is immediate by Proposition 2.1, Lemma 2.2 and Theorem 3.1. \square

Let C, D and B be unital C^* -algebras. We suppose that C and D are strongly Morita equivalent with respect to a $C - D$ -equivalence bimodule Y . Let F be a right Hilbert B -module and let ψ be a completely positive linear map from D to $\mathbf{B}_B(F)$. Let (π_C, E) be the non-degenerate representation of C on E induced by $(\pi_\psi, V_\psi, F_\psi)$ and Y . Let $\{u_i\}_{i=1}^n$ be a finite subset of Y . We construct a linear map ϕ from C to $\mathbf{B}_B(F) \otimes M_n(\mathbf{C})$ in the same way as in [8, Section 6], that is, let ϕ be the linear map from C to $\mathbf{B}_B(F) \otimes M_n(\mathbf{C})$ defined by

$$[\phi(c)_{ij}]_{i,j=1}^n = [\psi(\langle u_i, c \cdot u_j \rangle_D)]_{i,j=1}^n$$

for any $c \in C$. In the same way as in the proof of [8, Lemma 6.1], we can see that ϕ is a completely positive linear map from C to $\mathbf{B}_B(F) \otimes M_n(\mathbf{C})$. Let $F \otimes \mathbf{C}^n$ be the algebraic tensor product F and \mathbf{C}^n and we regard $F \otimes \mathbf{C}^n$ as a right Hilbert B -module in the natural way. We identify $\mathbf{B}_B(F) \otimes M_n(\mathbf{C})$ with $\mathbf{B}_B(F \otimes \mathbf{C}^n)$. Also, we identify $F \otimes \mathbf{C}^n$ with $\bigoplus_1^n F$ as right Hilbert B -modules. Let $(\pi_\phi, Y_\phi, E_\phi)$ be a minimal KSGNS-representation for ϕ . We show that (π_C, E) is unitarily equivalent to $(\pi_\phi, Y_\phi, E_\phi)$.

Let $C \odot (F \otimes \mathbf{C}^n)$ be the algebraic tensor product of C and $F \otimes \mathbf{C}^n$. We define a map U from $C \odot (F \otimes \mathbf{C}^n)$ to E by setting

$$U(c \otimes f \otimes \lambda) = \sum_{i=1}^n \lambda_i(c \cdot u_i) \otimes 1_D \otimes f$$

for any $c \in C, f \in F, \lambda = \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{bmatrix} \in \mathbf{C}^n$ and extending linearly. Then in the

same way as in the proof of [8, Lemma 6.2], we can see that U is an isometry from $C \odot (F \otimes \mathbf{C}^n)$ to E . Hence we can extend U to an isometry from E_ϕ to E . We denote it the same symbol U . From now on, we assume that $\{u_i\}_{i=1}^n$ is a left C -basis in Y . Such a finite subset $\{u_i\}_{i=1}^n$ of Y exists by Kajiwara and Watatani [6, Corollary 1.19]. Then we can see that U is surjective in the same way as in the proof of [8, Lemma 6.3]. Thus, U is an isometry from E_ϕ onto E . Furthermore, in the same way as in [8, Lemma 6.4], we can see that $\pi_\phi(c) = U^* \pi_C(c) U$ for any $c \in C$. Therefore, we obtain the following theorem:

Theorem 3.4. *Let C, D and B be unital C^* -algebras. We suppose that C and D are strongly Morita equivalent with respect to a $C - D$ -equivalence bimodule Y . Let $\{u_i\}_{i=1}^n$ be a left C -basis in Y . Let ψ be a completely positive linear map from D to*

$\mathbf{B}_B(F)$, where F is a right Hilbert B -module. Let ϕ be a mp from C to $\mathbf{B}_B(F \otimes \mathbf{C}^n)$ defined by

$$[\phi(c)_{ij}]_{i,j=1}^n = [\psi(\langle u_i, c \cdot u_j \rangle_D)]_{i,j=1}^n$$

for any $c \in C$. Then ϕ is a completely positive linear map from C to $\mathbf{B}_B(F \otimes \mathbf{C}^n)$, which is strongly Morita equivalent to ψ .

For any $\psi \in \text{CP}_B(D)$, we denote by $\mathcal{F}_0(\psi)$ the element in $\text{CP}_B(C)$ defined in the above and \mathcal{F}_0 denotes the map

$$\text{CP}_B(D) \rightarrow \text{CP}_B(C) : \psi \mapsto \mathcal{F}_0(\psi).$$

By the above theorem, we can obtain the map

$$\text{CP}_B(D)/\sim \rightarrow \text{CP}_B(C)/\sim : [\psi] \mapsto [\mathcal{F}_0(\psi)],$$

where $[\psi]$ and $[\mathcal{F}_0(\psi)]$ are the strong Morita equivalence classes of ψ and $\mathcal{F}_0(\psi)$, respectively. We denote by the symbol \mathcal{F} the map from $\text{CP}_B(D)/\sim$ to $\text{CP}_B(C)/\sim$. In the same as above, we can also obtain the map from $\text{CP}_B(C)/\sim$ to $\text{CP}_B(D)/\sim$, which is the inverse map of \mathcal{F} . Hence \mathcal{F} is a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_B(C)/\sim$. Therefore, we obtain the following:

Corollary 3.5. *Let C, D, B be unital C^* -algebras. Then \mathcal{F} is a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_B(C)/\sim$.*

Let C, A and B be unital C^* -algebras. We suppose that A and B are strongly Morita equivalent with respect to a $B - A$ -equivalence bimodule Z . Let ϕ be a completely positive linear map from C to $\mathbf{B}_B(F)$, where F is a right Hilbert B -module. Let ϕ^Z be the linear map from C to $\mathbf{B}_A(F \otimes_B Z)$ defined by

$$\phi^Z(c)(f \otimes z) = \phi(c)(f) \otimes z = (\phi \otimes \text{id}_Z)(c)(f \otimes z)$$

for any $c \in C$, $f \in F$, $z \in Z$. Also, let $(\pi_\phi, V_\phi, F_\phi)$ be a minimal KSGNC-representation for ϕ . Let $(\pi_\phi^Z, F_\phi \otimes_B Z)$ be the non-degenerate representation of C on the right Hilbert A -module $F_\phi \otimes_B Z$ defined in Section 2.

Lemma 3.6. *With the above notation, for any $c \in C$,*

$$\phi^Z(c) = (V_\phi^* \otimes \text{id}_Z) \pi_\phi^Z(c) (V_\phi \otimes \text{id}_Z)$$

and

$$\overline{\pi_\phi^Z(C)(V_\phi \otimes \text{id}_Z)(F \otimes_B Z)} = F_\phi \otimes_B Z.$$

Proof. For any $c \in C$, $f \in F$, $z \in Z$,

$$\begin{aligned} (V_\phi^* \otimes \text{id}_Z) \pi_\phi^Z(c) (V_\phi \otimes \text{id}_Z)(f \otimes z) &= (V_\phi^* \otimes \text{id}_Z) \pi_\phi^Z(c) (V_\phi f \otimes z) \\ &= (V_\phi^* \otimes \text{id}_Z) (\pi_\phi(c) V_\phi F \otimes z) \\ &= V_\phi^* \pi_\phi(c) V_\phi F \otimes z \\ &= \phi(c)(f) \otimes z \\ &= \phi^Z(c)(f \otimes z). \end{aligned}$$

Hence

$$\phi^Z(c) = (V_\phi^* \otimes \text{id}_Z) \pi_\phi^Z(c) (V_\phi \otimes \text{id}_Z)$$

for any $c \in C$ and clearly $V_\phi \otimes \text{id} \in \mathbf{B}_A(F \otimes_B Z, F_\phi \otimes_B Z)$. Thus ϕ^Z is completely positive by [9, Proposition 5.5]. Also,

$$\overline{\pi_\phi^Z(C)(V_\phi \otimes \text{id}_Z)(F \otimes_B Z)} = \overline{\pi_\phi(C) V_\phi F \otimes_B Z} = \overline{F_\phi \otimes_B Z}$$

since $\overline{\pi_\phi(C) V_\phi F} = F_\phi$. □

By Lemma 3.6, $(\pi_\phi^Z, V_\phi \otimes \text{id}_Z, F_\phi \otimes_B Z)$ is a minimal KSGNS-representation for ϕ^Z .

We denote by $\tilde{\Phi}$, the map $[\phi] \in \text{CP}_B(C)/\sim \mapsto [\phi^Z] \in \text{CP}_A(C)/\sim$, where $[\phi]$ and $[\phi^Z]$ are the strong Morita equivalence classes of $\phi \in \text{CP}_B(C)$ and $\phi^Z \in \text{CP}_A(C)$, respectively.

Lemma 3.7. *With the above notation, $\tilde{\Phi}$ is well-defined and $\tilde{\Phi}$ is a bijective correspondence between $\text{CP}_B(C)/\sim$ and $\text{CP}_A(C)/\sim$.*

Proof. This is immediate by Lemmas 2.4 and 3.6. \square

We give a similar result to [8, Corollary 6.6].

Theorem 3.8. *Let C, A and D, B be unital C^* -algebras. We suppose that C and A are strongly Morita equivalent to D and B , respectively. Then there exists a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_A(C)/\sim$.*

Proof. By Corollary 3.5, there is a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_B(C)/\sim$. Also, by Lemma 3.7, there is a bijective correspondence between $\text{CP}_B(C)/\sim$ and $\text{CP}_A(C)/\sim$. Thus, we obtain a bijective correspondence between $\text{CP}_B(D)/\sim$ and $\text{CP}_A(C)/\sim$. \square

4. GNS- C^* -CORRESPONDENCES AND STRONG MORITA EQUIVALENCE

Let C, D and A, B be unital C^* -algebras. Let ϕ be a completely positive linear map from C to A . In the same way as the KSGNS-construction, we can construct a $C - A$ -correspondence E_ϕ . Following Marrero and Muhly [10], we call E_ϕ a *GNS- C^* -correspondence* induced by ϕ . Also, let ψ be a completely positive linear map from D to B and let E_ψ be the GNS- C^* -correspondence induced by ψ , which is a $D - B$ -correspondence. Following Muhly and Solel [11], we give the following definition:

Definition 4.1. We say that E_ϕ is *strongly Morita equivalent* to E_ψ if there exist a $C - D$ -equivalence bimodule Y and an $A - B$ -equivalence bimodule X such that $E_\phi \otimes_A X$ is isomorphic to $Y \otimes_D E_\psi$ as $C - B$ -correspondences, that is, there is a linear map Φ from $E_\phi \otimes_A X$ onto $Y \otimes_D E_\psi$ satisfying the following:

- (1) $\Phi(c \cdot x \cdot b) = c \cdot \Phi(x) \cdot b$,
- (2) $\langle \Phi(x), \Phi(y) \rangle_B = \langle x, y \rangle_B$,

for any $x, y \in E_\phi \otimes_A X$, $c \in C$, $b \in B$.

Let ϕ be a completely positive linear map from C to A and E_ϕ the $C - A$ -correspondence induced by ϕ . We regard A as the trivial right Hilbert A -module in the natural way and ϕ can be regarded as a completely positive linear map from C to $\mathbf{B}_A(A)$. Also E_ϕ can be regarded as a right Hilbert A -module in a minimal KSGNS-representation for ϕ .

Let ψ be a completely positive linear map from D to B and E_ψ the $D - B$ -correspondence induced by ψ . We also regard E_ψ as a right Hilbert B -module in a minimal KSGNS-representation for ψ .

We suppose that there is an $A - B$ -equivalence bimodule X . Then $A \cong \mathbf{B}_B(X)$ as C^* -algebras by the isomorphism T of A onto $\mathbf{B}_B(X)$ defined by

$$T_a(x) = a \cdot x$$

for any $a \in A$, $x \in X$. Let $\phi_T = T \circ \phi$, a completely positive linear map from C to $\mathbf{B}_B(X)$. First, we show that if ϕ_T and ψ are strongly Morita equivalent as completely positive linear maps, then E_ϕ and E_ψ are strongly Morita equivalent

as GNS- C^* -correspondences. Since X is an $A - B$ -equivalence bimodule and B is unital, there is a finite subset $\{v_j\}_{j=1}^m$ of X such that

$$\sum_{j=1}^m \langle v_j, v_j \rangle_B = 1_B$$

by the proof of [6, Corollary 1.19]. Let $(\pi_{\phi_T}, V_{\phi_T}, E_{\phi_T})$ be a minimal KSGNS-representation for ϕ_T .

Lemma 4.1. *With the above notation, let Ψ be a linear map from $E_{\phi} \otimes_A X$ to E_{ϕ_T} defined by*

$$\Psi((c \otimes a) \otimes x) = c \otimes (a \cdot x)$$

for any $a \in A$, $c \in C$, $x \in X$. Then Ψ is a $C - B$ -correspondence isomorphism of $E_{\phi} \otimes_A X$ onto E_{ϕ_T} . Furthermore, Ψ^{-1} is given by

$$\Psi^{-1}(c \otimes x) = \sum_{j=1}^m c \otimes_A \langle x, v_j \rangle \otimes v_j$$

for any $c \in C$, $x \in X$, where E_{ϕ_T} is regarded as a $C - B$ -correspondence.

Proof. Let $c, c_1 \in C$, $a, a_1 \in A$, $x, x_1 \in X$. Then

$$\begin{aligned} \langle \Psi((c \otimes a) \otimes x), (c_1 \otimes a_1) \otimes x_1 \rangle_B &= \langle c \otimes a \cdot x, c_1 \otimes a_1 \cdot x_1 \rangle_B \\ &= \langle a \cdot x, \phi_T(c^* c_1)(a_1 \cdot x) \rangle_B \\ &= \langle x, a^* \phi(c^* c_1) a_1 \cdot x \rangle_B. \end{aligned}$$

On the other hand,

$$\begin{aligned} \langle (c \otimes a) \otimes x, (c_1 \otimes a_1) \otimes x_1 \rangle_B &= \langle x, \langle c \otimes a, c_1 \otimes a_1 \rangle_A \cdot x_1 \rangle_B \\ &= \langle x, \langle a, \phi(c^* c_1) a_1 \rangle_A \cdot x_1 \rangle_B \\ &= \langle x, a^* \phi(c^* c_1) a_1 \cdot x_1 \rangle_B. \end{aligned}$$

Hence Ψ preserves the right B -valued inner products. Also, let $c, c_1 \in C$, $a \in A$, $x \in X$, $b \in B$. Then

$$\begin{aligned} \Psi(c_1 \cdot (c \otimes a) \otimes x \cdot b) &= \Psi((c_1 c \otimes a) \otimes (x \cdot b)) \\ &= c_1 c \otimes (a \cdot x \cdot b) \\ &= c_1 \cdot (c \otimes a \cdot x) \cdot b \\ &= c_1 \cdot \Psi((c \otimes a) \otimes x) \cdot b. \end{aligned}$$

Thus, Ψ is a $C - B$ -bimodule map. Furthermore, by easy computations, $\Psi \circ \Psi^{-1} = \text{id}_{E_{\phi_T}}$ and $\Psi^{-1} \circ \Psi = \text{id}_{E_{\phi} \otimes_A X}$. Therefore, we obtain the conclusion. \square

Let $(\pi_{\psi}, V_{\psi}, E_{\psi})$ be a minimal KSGNS-representations for ψ . We also regard E_{ψ} as a $D - B$ -correspondence for ψ . We suppose that ϕ_T and ψ are strongly Morita equivalent as completely positive linear maps with respect to a $C - D$ -equivalence bimodule Y and a linear map π_Y from Y to $\mathbf{B}_B(E_{\psi}, E_{\phi_T})$ satisfying Conditions (1)-(3) in Definition 2.2.

Lemma 4.2. *With the above notation, let Φ be the linear map from $Y \otimes_D E_{\psi}$ to E_{ϕ_T} defined by*

$$\Phi(y \otimes (d \otimes b)) = \pi_Y(y)(d \otimes b)$$

for any $y \in Y$, $d \in D$, $b \in B$. Then Φ is a $C - B$ -correspondence isomorphism of $Y \otimes_D E_{\psi}$ onto E_{ϕ_T} .

Proof. Let $y, y_1 \in Y$, $d, d_1 \in D$, $b, b_1 \in B$. Then

$$\begin{aligned} \langle \Phi(y \otimes (d \otimes b)), \Phi(y_1 \otimes (d_1 \otimes b_1)) \rangle_B &= \langle \pi_Y(y)(d \otimes b), \pi_Y(y_1)(d_1 \otimes b_1) \rangle_B \\ &= \langle [\pi_Y(y_1)^* \pi_Y(y)](d \otimes b), d_1 \otimes b_1 \rangle_B \\ &= \langle \pi_\psi(\langle y_1, y \rangle_D)(d \otimes b), d_1 \otimes b_1 \rangle_B \\ &= \langle \langle y_1, y \rangle_D d \otimes b, d_1 \otimes b_1 \rangle_B. \end{aligned}$$

On the other hand,

$$\begin{aligned} \langle y \otimes (d \otimes b), y_1 \otimes (d_1 \otimes b_1) \rangle_B &= \langle d \otimes b, \langle y, y_1 \rangle_D d_1 \otimes b_1 \rangle_B \\ &= \langle \langle y_1, y \rangle_D d \otimes b, d_1 \otimes b_1 \rangle_B. \end{aligned}$$

Hence Φ preserves the right B -valued inner products. Let $c \in C$, $y \in Y$, $d \in D$, $b, b_1 \in B$. Then

$$\begin{aligned} \Phi(c \cdot y \otimes (d \otimes b) \cdot b_1) &= \Phi((c \otimes y) \otimes (d \otimes b b_1)) \\ &= \pi_Y(c \cdot y)(d \otimes b b_1) \\ &= (\pi_\phi(c) \pi_Y(y))(d \otimes b) \cdot b_1 \\ &= c \cdot \pi_Y(y)(d \otimes b) \cdot b_1 \\ &= c \cdot \Phi(y \otimes (d \otimes b)) \cdot b_1. \end{aligned}$$

Hence Φ is a $C - B$ -bimodule map from $Y \otimes_D E_\psi$ to E_{ϕ_T} . Furthermore, since Y is a $C - D$ -equivalence bimodule and C is unital, there is a finite subset $\{u_i\}_{i=1}^n$ of Y such that

$$\sum_{i=1}^n C \langle u_i, u_i \rangle = 1_C.$$

Let $c \in C$, $x \in X$. Then $\sum_{i=1}^n u_i \otimes \pi_Y(u_i)^*(c \otimes x) \in Y \otimes_D E_\psi$ and

$$\begin{aligned} \Phi\left(\sum_{i=1}^n u_i \otimes \pi_Y(u_i)^*(c \otimes x)\right) &= \sum_{i=1}^n \pi_Y(u_i)(\pi_Y(u_i)^*(c \otimes x)) \\ &= \sum_{i=1}^n (\pi_Y(u_i) \pi_Y(u_i)^*)(c \otimes x) \\ &= \sum_{i=1}^n \pi_\phi(C \langle u_i, u_i \rangle)(c \otimes x) \\ &= c \otimes x \end{aligned}$$

since $\pi_Y(u_i) \pi_Y(u_i)^* = \pi_\phi(C \langle u_i, u_i \rangle)$ for $i = 1, 2, \dots, n$. Thus Φ is surjective. Therefore, Φ is a $C - B$ -correspondence isomorphism of $Y \otimes_D E_\psi$ onto E_{ϕ_T} . \square

Next, we suppose that E_ϕ and E_ψ are strongly Morita equivalent as GNS- C^* -correspondences. Then there are a $C - D$ -equivalence bimodule Y , $A - B$ -equivalence bimodule X and a $C - B$ -correspondence isomorphism Φ_1 of $Y \otimes_D E_\psi$ onto $E_\phi \otimes_A X$. Let Ψ be the $C - B$ -correspondence isomorphism of $E_\phi \otimes_A X$ onto E_{ϕ_T} defined in Lemma 4.1. Let $\Phi = \Psi \circ \Phi_1$, a $C - B$ -correspondence isomorphism of $Y \otimes_D E_\psi$ onto E_{ϕ_T} .

Lemma 4.3. *With the above notation, let π_Y be the space of all right B -module maps from E_ψ to E_ϕ defined by*

$$\pi_Y(y)(d \otimes b) = \Phi(y \otimes (d \otimes b))$$

for any $y \in Y$, $d \in D$, $b \in B$. Then π_Y is a linear map from Y to $\mathbf{B}_B(E_\psi, E_{\phi_T})$ satisfying Conditions (1) – (3) in Definition 2.2.

Proof. First, we show that $\pi_Y(y) \in \mathbf{B}_B(E_\psi, E_{\phi_T})$ for any $y \in Y$. It suffices to show that $\pi_Y(y)$ is adjointable for any $y \in Y$. For any $y, y_1 \in Y, d, d_1 \in D, b, b_1 \in B$,

$$\begin{aligned} \langle \pi_Y(y)(d \otimes b), \pi_Y(y_1)(d_1 \otimes b_1) \rangle_B &= \langle \Phi(y \otimes (d \otimes b)), \Phi(y_1 \otimes (d_1 \otimes b_1)) \rangle_B \\ &= \langle y \otimes (d \otimes b), y_1 \otimes (d_1 \otimes b_1) \rangle_B \\ &= \langle d \otimes b, \langle y, y_1 \rangle_D d_1 \otimes b_1 \rangle_B. \end{aligned}$$

On the other hand,

$$\begin{aligned} \langle \pi_\psi(\langle y_1, y \rangle_D)(d \otimes b), d_1 \otimes b_1 \rangle_B &= \langle \langle y_1, y \rangle_D d \otimes b, d_1 \otimes b_1 \rangle_B \\ &= \langle d \otimes b, \langle y, y_1 \rangle_D d_1 \otimes b_1 \rangle_B. \end{aligned}$$

Since the linear span of the set

$$\{\Phi(y \otimes (d \otimes b)) \mid y \in Y, d \in D, b \in B\}$$

is dense in E_{ϕ_T} , $\pi_Y(y_1)$ is adjointable for any $y \in Y$ and

$$\pi_Y(y_1)^*(\Phi(y \otimes (d \otimes b))) = \pi_\psi(\langle y_1, y \rangle_D)(d \otimes b).$$

And,

$$\pi(y_1)^*\pi(y) = \pi_\psi(\langle y_1, y \rangle_D).$$

Furthermore, for any $y, y_1, z \in Y, d \in D, b \in B$,

$$\pi_Y(z)\pi_Y(y_1)^*\Phi(y \otimes (d \otimes b)) = \pi_Y(z)(\langle y_1, y \rangle_D d \otimes b) = \Phi(z \otimes \langle y_1, y \rangle_D d \otimes b).$$

On the other hand,

$$\begin{aligned} \pi_{\phi_T}(C\langle z, y_1 \rangle)\Phi(y \otimes (d \otimes b)) &= C\langle z, y_1 \rangle \cdot \Phi(y \otimes (d \otimes b)) \\ &= \Phi((C\langle z, y_1 \rangle \cdot y) \otimes (d \otimes b)) \\ &= \Phi(z \cdot \langle y_1, y \rangle_D \otimes (d \otimes b)) \\ &= \Phi(z \otimes (\langle y_1, y \rangle_D d \otimes b)). \end{aligned}$$

Hence $\pi_Y(z)\pi_Y(y_1)^* = \pi_{\phi_T}(C\langle z, y_1 \rangle)$ for any $z, y_1 \in Y$. Finally, for any $c \in C, y \in Y, b \in B, d, d_1 \in D$,

$$\begin{aligned} \pi_{\phi_T}(c)\pi_Y(y)\pi_\psi(d_1)(d \otimes b) &= \pi_{\phi_T}(c)\pi_Y(y)(d_1 d \otimes b) \\ &= \pi_{\phi_T}(c)\Phi(y \otimes (d_1 d \otimes b)) \\ &= c \cdot \Phi(y \otimes (d_1 d \otimes b)) \\ &= \Phi((c \cdot y \cdot d_1) \otimes (d \otimes b)) \\ &= \pi_Y(c \cdot y \cdot d_1)(d \otimes b). \end{aligned}$$

Thus $\pi_{\phi_T}(c)\pi_Y(y)\pi_\psi(d_1) = \pi_Y(c \cdot y \cdot d_1)$ for any $c \in C, y \in Y, d_1 \in D$. Therefore, we obtain the conclusion. \square

Combining Lemmas 4.1, 4.2 and 4.3, we obtain the following theorem:

Theorem 4.4. *Let C, D and A, B be unital C^* -algebras. Let ϕ and ψ be completely positive linear maps from C and D to A and B , respectively and E_ϕ and E_ψ the $C - A$ -correspondence and the $D - B$ -correspondence induced by ϕ and ψ , respectively. Then the following hold:*

(1) *If there is an $A - B$ -equivalence bimodule X and ϕ_T and ψ are strongly Morita equivalent as completely positive linear maps, then E_ϕ and E_ψ are strongly Morita equivalent as GNS- C^* -correspondences, where ϕ_T is a completely positive linear map from C to $\mathbf{B}_B(X)$ induced by ϕ and the isomorphism T of A onto $\mathbf{B}_B(X)$ defined by $T_a(x) = a \cdot x$ for any $a \in A, x \in X$.*

(2) *If E_ϕ and E_ψ are strongly Morita equivalent as GNS- C^* -correspondences, then there is an $A - B$ -equivalence bimodule X and ϕ_T and ψ are strongly Morita equivalent as completely positive linear maps, where ϕ_T is a completely positive linear*

map from C to $\mathbf{B}_B(X)$ induced by ϕ and the isomorphism T of A onto $\mathbf{B}_B(X)$ defined by $T_a(x) = a \cdot x$ for any $a \in A$, $x \in X$.

Proof. We can show (1) by Lemmas 4.1 and 4.2. We can also show (2) by Lemma 4.3. \square

Finally, we give an example on Theorem 4.4.

Example 4.5. Let C, D, B be unial C^* -algebras and ψ a completely positive linear map from D to B . We regard ψ as a completely positive linear map from D to $\mathbf{B}_B(B)$. We assume that C and D are strongly Morita equivalent. Then we can construct ϕ_T , a completely positive linear map from C to $\mathbf{B}_B(B \otimes \mathbf{C}^n)$ by Theorem 3.4, where n is some positive integer. Let T be the isomorphism of $B \otimes M_n(\mathbf{C})$ onto $\mathbf{B}_B(B \otimes \mathbf{C}^n)$ defined by

$$T_b(x) = b \cdot x$$

for any $b \in B \otimes M_n(\mathbf{C})$, $x \in B \otimes \mathbf{C}^n$, where we regard $B \otimes \mathbf{C}^n$ as a $B \otimes M_n(\mathbf{C})$ - B -equivalence bimodule in the natural way. Let $\phi = T^{-1} \circ \phi_T$. Then by Theorems 3.4 and 4.4, E_ϕ and E_ψ are strongly Morita equivalent as GNS- C^* -correspondences, where E_ϕ and E_ψ are a C - $B \otimes M_n(\mathbf{C})$ -correspondence and a D - B -correspondence, respectively.

REFERENCES

- [1] B. Blackadar, *K-theory for operator algebras*, M. S. R. I. Publications 5, 2nd Edition, Cambridge Univ. Press, Cambridge, 1998.
- [2] B. Blackadar, *Operator algebras. Theory of C^* -algebras and von Neumann algebras*, Encyclopaedia of Mathematical Sciences, **122**, Operator algebras and Non-commutative Geometry III, Springer-Verlag, Berlin, 2006.
- [3] L. G. Brown, J. Mingo and N-T. Shen, *Quasi-multipliers and embeddings of Hilbert C^* -bimodules*, Can. J. Math. **46** (1994), 1150–1174.
- [4] S. Echterhoff and I. Raeburn, *Multipliers of imprimitivity bimodules and Morita equivalence of crossed products*, Math. Scand. **76** (1995), 289–309.
- [5] K. K. Jensen and K. Thomsen, *Elements of KK -theory*, Birkhäuser, 1991.
- [6] T. Kajiwara and Y. Watatani, *Jones index theory by Hilbert C^* -bimodules and K -Theory*, Trans. Amer. Math. Soc. **352** (2000), 3429–3472.
- [7] G. G. Kasparov, *Hilbert C^* -modules: theorems of Stinespring and Voiculescu*, J. Operator Theory **4** (1980), 133–150.
- [8] K. Kodaka, *Strong Morita equivalence for completely positive linear maps on C^* -algebras*, preprint, arXiv: 2102.13317.
- [9] E. C. Lance, *Hilbert C^* -modules*, A toolkit for operator algebraists, London Math. Soc., Lecture Note Series, **210**, Cambridge Univ. Press, Cambridge, 1995.
- [10] A. E. Marrero and P. Muhly, *Cuntz-Pimsner algebras, completely positive maps and Morita equivalence*, Proc. Amer. Math. Soc. **134** (2006), 1133–1135.
- [11] P. Muhly and B. Solel, *On the Morita equivalence of tensor algebras*, Proc. London Math. Soc. **81** (2000), 113–168.
- [12] W. F. Stinespring, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc., **6** (1955), 211–216.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, RYUKYU UNIVERSITY,
NISHIHARA-CHO, OKINAWA, 903-0213, JAPAN

E-mail address: kodaka@math.u-ryukyu.ac.jp