

POINCARÉ SERIES FOR MODULAR GRAPH FORMS AT DEPTH TWO

II. Iterated integrals of cusp forms

Daniele Dorigoni¹, Axel Kleinschmidt^{2,3} and Oliver Schlotterer⁴

¹*Centre for Particle Theory & Department of Mathematical Sciences
Durham University, Lower Mountjoy, Stockton Road, Durham DH1 3LE, UK*

²*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut)
Am Mühlenberg 1, DE-14476 Potsdam, Germany*

³*International Solvay Institutes
ULB-Campus Plaine CP231, BE-1050 Brussels, Belgium*

⁴*Department of Physics and Astronomy, Uppsala University, 75108 Uppsala, Sweden*

We continue the analysis of modular invariant functions, subject to inhomogeneous Laplace eigenvalue equations, that were determined in terms of Poincaré series in a companion paper. The source term of the Laplace equation is a product of (derivatives of) two non-holomorphic Eisenstein series whence the modular invariants are assigned depth two. These modular invariant functions can sometimes be expressed in terms of single-valued iterated integrals of holomorphic Eisenstein series as they appear in generating series of modular graph forms. We show that the set of iterated integrals of Eisenstein series has to be extended to include also iterated integrals of holomorphic cusp forms to find expressions for all modular invariant functions of depth two. The coefficients of these cusp forms are identified as ratios of their L-values inside and outside the critical strip.

Contents

1	Introduction	1
2	Basics of iterated integrals	5
2.1	Iterated integrals of Eisenstein series	6
2.2	Iterated integrals of cusp forms	10
2.3	Properties of multiple modular values and the β^{sv}	15
3	Modular properties of solutions to the Laplace equations	21
3.1	Examples involving the Ramanujan cusp form	22
3.2	Examples with cusp forms of higher weight	25
3.3	An example involving the two weight 24 cusp forms	26
3.4	Structure for general weight	29
4	Selection rules on β^{sv} from Tsunogai's derivation algebra	30
4.1	Overview of ϵ_k relations at depth two	31
4.2	Modular graph forms and ϵ_k relations at depth two	36
5	Kloosterman sums and the Fourier expansion of the $F_{m,k}^{\pm(s)}$	41
5.1	Odd example	41
5.2	Even example	42
5.3	General comments	43
A	Recap of seed functions	44
B	Examples of cocycles	45
B.1	Weight 16	45
B.2	Weight 18	46

1 Introduction

In the companion Part I [1] to this paper we introduced the Laplace equations

$$(\Delta - s(s - 1))F_{m,k}^{+(s)} = E_m E_k, \tag{1.1a}$$

$$(\Delta - s(s - 1))F_{m,k}^{-(s)} = \frac{(\nabla E_m)(\overline{\nabla} E_k) - (\nabla E_k)(\overline{\nabla} E_m)}{2(\text{Im } \tau)^2}, \tag{1.1b}$$

with integers $s \geq 2$ and $2 \leq m \leq k$, and where E_k are non-holomorphic Eisenstein series

$$E_k = \frac{(\text{Im } \tau)^k}{\pi^k} \sum_{(m,n) \neq (0,0)} \frac{1}{|m\tau + n|^{2k}}. \tag{1.2}$$

The modular parameter τ is in the upper half-plane, and E_k is invariant under the modular transformations

$$\tau \rightarrow \gamma \cdot \tau = \frac{a\tau + b}{c\tau + d} \quad \text{for } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z}). \quad (1.3)$$

The Cauchy–Riemann derivatives $\nabla = 2i(\text{Im } \tau)^2 \partial_\tau$ of E_k are modular forms of weight $(0, -2)$ and the Laplacian $\Delta = 4(\text{Im } \tau)^2 \partial_\tau \partial_{\bar{\tau}} = \bar{\nabla}((\text{Im } \tau)^{-2} \nabla)$ is modular invariant. The superscripts \pm on $F_{m,k}^{\pm(s)}$ indicate that these functions are required to be even/odd under the involution $\tau \rightarrow -\bar{\tau}$ of the upper half-plane, in line with the respective right-hand sides of (1.1). The spectrum of eigenvalues appearing in (1.1) is

$$F_{m,k}^{+(s)} : \quad s \in \{k-m+2, k-m+4, \dots, k+m-4, k+m-2\}, \quad (1.4a)$$

$$F_{m,k}^{-(s)} : \quad s \in \{k-m+1, k-m+3, \dots, k+m-3, k+m-1\}. \quad (1.4b)$$

As all the objects in (1.1) are modular invariant, we focus on modular invariant solutions $F_{m,k}^{\pm(s)}$ to the Laplace problem. The transcendental weight of $F_{m,k}^{\pm(s)}$ is $m+k$ from (1.1) given that E_m and E_k have transcendental weight m and k , respectively.

In Part I, we constructed solutions to (1.1) in terms of absolutely convergent¹ Poincaré series

$$F_{m,k}^{\pm(s)}(\tau) = \sum_{\gamma \in B(\mathbb{Z}) \setminus \text{SL}(2, \mathbb{Z})} f_{m,k}^{\pm(s)}(\gamma \cdot \tau), \quad (1.5)$$

where the seed functions $f_{m,k}^{\pm(s)}$ are invariant under shifts $\tau \rightarrow \tau+n$ for $n \in \mathbb{Z}$ which form the stabiliser of the cusp $\tau \rightarrow i\infty$

$$B(\mathbb{Z}) = \left\{ \begin{pmatrix} \pm 1 & n \\ 0 & \pm 1 \end{pmatrix} \middle| n \in \mathbb{Z} \right\} \subset \text{SL}(2, \mathbb{Z}). \quad (1.6)$$

For the convenience of the reader, appendix A recaps the explicit form for our choice of representatives of these seeds $f_{m,k}^{\pm(s)}$.

Even though the solution (1.5) is fully explicit and has interesting structures analysed in Part I, extracting the complete Fourier expansion of $F_{m,k}^{\pm(s)}$ from the Poincaré-series representations is fairly involved. For the Fourier zero mode one can use the methods of [2–5] but the non-zero modes with respect to $\tau \rightarrow \tau+1$ are hard to obtain. For this reason it is desirable to find alternative expressions for the modular invariants $F_{m,k}^{\pm(s)}$.

A family of functions with well-defined modular transformation properties is provided by modular graph forms (MGFs) [6–8]. These arise in the α' -expansion of configuration-space integrals of genus-one closed-string amplitudes and have been studied from a physical perspective in [9–11, 6, 12, 13, 7, 8, 14–20, 2, 21–23, 3, 24–30, 5, 31–34] and a mathematical perspective in [35–43]. As they arise from string amplitudes, MGFs possess a lattice-sum description over discrete momenta of Feynman graphs drawn on the genus-one string world-sheet.

¹Absolute convergence is guaranteed for $m < k$ and for $m = k$ a suitable regularisation was described in Part I.

In particular, generating functions of closed-string integrals and their associated differential equations [28, 30] lead to expressions for MGFs in terms of real-analytic objects denoted by

$$\beta^{\text{sv}} \left[\begin{matrix} j_1 & j_2 & \dots & j_\ell \\ k_1 & k_2 & \dots & k_\ell \end{matrix}; \tau \right] \quad \text{with } k_i \in \{4, 6, 8, \dots\} \text{ and } 0 \leq j_i \leq k_i - 2, \quad (1.7)$$

where $\ell \in \mathbb{N}$ is called the depth of β^{sv} .² Depth serves as a filtration, and the highest-depth terms in the complex-conjugation and modular properties of the β^{sv} take a simple form. The β^{sv} are constructed from (single-valued) iterated integrals over holomorphic Eisenstein series and should be closely related to Brown's non-holomorphic modular forms [37, 38], although a precise dictionary between the two formalisms is still missing beyond depth one. Together with certain antiholomorphic integration constants determined in Part I, the complete Fourier expansion of the β^{sv} at depths one and two is known. Therefore it seems desirable to express the $F_{m,k}^{\pm(s)}$ in terms of the β^{sv} . A further advantage of using such a representation of a modular-invariant function in terms of iterated integrals is that it is unique [44], unlike lattice-sum representations that are more frequent for MGFs.

In many cases, the Poincaré-series representations in this work may be viewed as interpolating between double sums over lattice momenta and double integrals over holomorphic Eisenstein series: The seeds $f_{m,k}^{\pm(s)}$ in (1.5) are constructed from depth-one integrals, and the sum over $\text{SL}(2, \mathbb{Z})$ transformations is comparable to a single lattice momentum. However, the Poincaré sums in (1.5) often produce MGFs that require three and more lattice momenta (see Part I for details) or modular invariant functions without any known lattice-sum representation.

In Part I, we have presented a procedure for obtaining linear combinations $\check{F}_{m,k}^{\pm(s)}$ of β^{sv} of depths two and one, together with Laurent polynomial terms in $y = \pi \text{Im } \tau$ (that can be thought of as depth zero). These linear combinations were constructed by starting from depth-two terms that solve the Laplace equation (1.1) modulo terms of lower depth. The latter were fixed from certain requirements on the desired solutions concerning their Cauchy–Riemann derivatives and asymptotics at the cusp, see Part I for further details. However, this procedure was tailored towards solving the Laplace system in terms of the building blocks β^{sv} of MGFs and does not guarantee that the resulting expression is modular invariant.

By comparing the dimensions of the space of solutions to (1.1) and the space of MGFs at depth two we have seen in Part I that the MGFs do not suffice to span the space of $F_{m,k}^{\pm(s)}$. This is reflected in the fact that certain $\check{F}_{m,k}^{\pm(s)}$ fail to be modular invariant exactly in those cases when the dimensions of the functions spaces differ. In the present paper, we shall discuss how to augment the $\check{F}_{m,k}^{\pm(s)}$ so that they become modular invariant and therefore equal the corresponding Poincaré series $F_{m,k}^{\pm(s)}$ in (1.5). In other words, we illustrate through a variety of examples that MGFs do not exhaust the modular-invariant combinations of iterated integrals of holomorphic modular forms and their complex conjugates.

As we shall see, the missing ingredients beyond the β^{sv} are (real and imaginary parts of) iterated integrals of holomorphic cusp forms. From the Eichler–Shimura theorem [45, 46] and the work of Brown [35, 37, 38, 47] on iterated integrals of general holomorphic modular forms, it is not surprising that restricting to the β^{sv} , that only involve iterated integrals of holomorphic

²A more detailed review of the construction and properties of the β^{sv} can be found in section 2.

Eisenstein series, is insufficient to describe the full space of modular-invariant solutions to (1.1). The first discrepancy in the dimensions of the function spaces $F_{m,k}^{\pm(s)}$ and MGFs appears for eigenvalues $s = 6, 8, 9, 10, \dots$ which coincide exactly with half the modular weight of the first holomorphic cusp forms of $SL(2, \mathbb{Z})$. This can be seen as a hint that cusp forms are the missing piece of the puzzle.

A further indication for the relevance of holomorphic cusp forms stems from the appearance of conjectural matrix representations of Tsunogai's derivation algebra [48] in the generating series of MGFs [28, 30]. Relations in the derivation algebra are also tied to holomorphic cusp forms [49] and imply that, starting from depth two, there are combinations of the β^{sv} that are not contained in the generating series of MGFs. Completing these to modular invariants requires holomorphic cusp forms as we shall see. This follows from the S-modular transformations of the various β^{sv} which contain interesting so-called multiple modular values [35] that involve the values of completed L-functions of cusp forms at integers [50] and extend the set of single-valued multiple zeta values. In order to cancel these L-values from S-modular transformations in general one has to combine the β^{sv} with iterated integrals of cusp forms. We shall work out these ideas in detail in this paper and spell out a variety of examples.

As a byproduct of our analysis we derive that the series expansions in $q = \exp(2\pi i\tau)$ and $\bar{q} = \exp(-2\pi i\bar{\tau})$ of these non-holomorphic modular objects $F_{m,k}^{\pm(s)}$ display very interesting structures. Firstly, the leading terms in the expansion of the even functions $F_{m,k}^{+(s)}$ around the cusp $\text{Im } \tau \gg 1$ are Laurent polynomials in $y = \pi \text{Im } \tau$ that will also be referred to as ‘‘perturbative’’. These Laurent polynomials have a single term with a rational coefficient, a single term with a \mathbb{Q} -multiple of the product $\zeta_{2m-1}\zeta_{2k-1}$, while all other coefficients are \mathbb{Q} -multiples of odd zeta values $\zeta_{2m-1}, \zeta_{2k-1}, \zeta_{m+k+s-1}$, see Part I for further details. Secondly, the infinite tower of exponentially suppressed, non-perturbative terms of the form $q^n \bar{q}^m$, with both of $n, m > 0$, have Laurent polynomials in y with rational coefficients for both the even and odd $F_{m,k}^{\pm(s)}$. Finally, and perhaps more interestingly, due to the presence of iterated integrals of holomorphic cusp forms we find that the exponentially suppressed terms of the form $q^n \bar{q}^0$ (and their complex conjugates $q^0 \bar{q}^n$) with $n > 0$ are multiplied by particularly rich Laurent polynomials in y : Their coefficients are either rationals, or \mathbb{Q} -multiples of single odd zeta values or surprisingly rationals (or more general number-field extensions of \mathbb{Q}) times special ratios of completed L-values associated to whichever cusp form is at play.

These results allow us to make novel predictions regarding the non-zero Fourier-mode decomposition of the Poincaré series (1.5). In particular in Part I, we have thoroughly explained how, for all the constructed seed functions $f_{m,k}^{\pm(s)}(\tau)$, one can exploit the results of [3] to obtain the purely perturbative Laurent polynomials in y . To pass from the seed function $f_{m,k}^{\pm(s)}(\tau)$ to the actual associated Poincaré series (1.5) one needs to use a particular integral transform detailed for instance in appendix A of Part I. Such a mapping between seed and modular function can also be used to formally obtain the non-zero modes for the modular invariant Poincaré series. However, the computation of non-zero modes from this integral transform of the seed involves very complicated Kloosterman sums and the analogue of the analysis in [3] to this case is currently unknown. Despite this lack of full control over Kloosterman sums, our results imply that

these Kloosterman sums must contain completed L-values of holomorphic cusp forms. It would be extremely interesting to extend the results of [3] to the non-zero Fourier mode sectors, thus deriving directly from the seed functions the exponentially suppressed terms $q^n \bar{q}^0$ and $q^0 \bar{q}^n$ with $n > 0$ including their Laurent polynomials.

Outlook

The results of Part I and this work raise a variety of follow-up questions of relevance to string perturbation theory, algebraic geometry and number theory. Most obviously, the Fourier expansion of depth-two MGFs and their extension by iterated integrals of holomorphic cusp forms call for generalisations to higher depth. Among other things, (single-valued) multiple zeta values beyond depth one, iterated integrals that mix holomorphic Eisenstein series with cusp forms and generalisations of L-values [51] are expected to play a key role starting from depth three. The respective seed should have one unit of depth less than its modular invariant Poincaré sum, and it will be rewarding to study this kind of recursive structure at general depth.

Furthermore, a detailed connection with the recent mathematics literature promises powerful synergies. Various important properties of the β^{sv} at general depth will follow once their precise relation to Brown’s non-holomorphic modular forms is established. Moreover, we note that iterated integrals of cusp forms and their Poincaré sums have featured prominently in recent work [52] that also relates to so-called higher-order modular forms. Certain Laplace systems similar to (1.1) but at depth three have also been studied recently in [53]. These references can provide useful guidance when generalising our work.

Outline

In section 2, we review the basic properties of iterated integrals of holomorphic modular forms, with particular emphasis on their modular properties and certain $\text{SL}(2, \mathbb{Z})$ group cocycles that arise. In section 3, we then use these results in the analysis of the modular invariant solutions to the Laplace equations. We further show how to combine the β^{sv} with iterated integrals of cusp forms based on the vanishing of the cocycles thus restoring modularity. We explain the relation between Tsunogai’s derivation algebra and the modular invariant Laplace eigenfunctions in section 4. Further properties of the solutions to (1.1), such as connections to Kloosterman sums, are discussed in section 5. An ancillary file that accompanies the arXiv submission of this work contains many examples and explicit expressions related to the functions $F_{m,k}^{\pm(s)}$.

2 Basics of iterated integrals

This section is dedicated to the central aspects of iterated integrals as well as their differential and modular properties as they enter our analysis. Frequent use will be made of the Cauchy–Riemann derivatives

$$\nabla = 2i(\text{Im } \tau)^2 \partial_\tau, \quad \bar{\nabla} = -2i(\text{Im } \tau)^2 \partial_{\bar{\tau}} \quad (2.1)$$

and the Laplace operator

$$\Delta = 4(\operatorname{Im} \tau)^2 \partial_\tau \partial_{\bar{\tau}} = (\pi \bar{\nabla})(y^{-2} \pi \nabla). \quad (2.2)$$

As in the equation above, we often use the symbol $y = \pi \operatorname{Im} \tau$, and powers of y satisfy

$$(\pi \nabla)^k y^a = (\pi \bar{\nabla})^k y^a = \frac{\Gamma(a+k)}{\Gamma(a)} y^{a+k}, \quad \Delta y^a = a(a-1)y^a. \quad (2.3)$$

2.1 Iterated integrals of Eisenstein series

In the present work, we shall only require the depth-one and depth-two versions of the single-valued iterated Eisenstein integrals (1.7). These are defined by the integrals [30]³

$$\beta^{\text{sv}} \left[\begin{matrix} j \\ k \end{matrix} \right] = \frac{(2\pi i)^{-1}}{(4y)^{k-2-j}} \left\{ \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^{k-2-j} (\bar{\tau} - \tau_1)^j G_k(\tau_1) - \int_{\bar{\tau}}^{-i\infty} d\bar{\tau}_1 (\tau - \bar{\tau}_1)^{k-2-j} (\bar{\tau} - \bar{\tau}_1)^j \overline{G_k(\tau_1)} \right\} \quad (2.4a)$$

with $0 \leq j \leq k-2$ and

$$\begin{aligned} \beta^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix} \right] &= \sum_{p_1=0}^{k_1-2-j_1} \sum_{p_2=0}^{k_2-2-j_2} \frac{\binom{k_1-2-j_1}{p_1} \binom{k_2-2-j_2}{p_2}}{(4y)^{p_1+p_2}} \alpha \left[\begin{matrix} j_1+p_1 & j_2+p_2 \\ k_1 & k_2 \end{matrix} \right] + \frac{(2\pi i)^{-2}}{(4y)^{k_1+k_2-j_1-j_2-4}} \\ &\times \left\{ \int_{\tau}^{i\infty} d\tau_2 (\tau - \tau_2)^{k_2-j_2-2} (\bar{\tau} - \tau_2)^{j_2} G_{k_2}(\tau_2) \int_{\tau_2}^{i\infty} d\tau_1 (\tau - \tau_1)^{k_1-j_1-2} (\bar{\tau} - \tau_1)^{j_1} G_{k_1}(\tau_1) \right. \\ &- \int_{\tau}^{i\infty} d\tau_2 (\tau - \tau_2)^{k_2-j_2-2} (\bar{\tau} - \tau_2)^{j_2} G_{k_2}(\tau_2) \int_{\bar{\tau}}^{-i\infty} d\bar{\tau}_1 (\tau - \bar{\tau}_1)^{k_1-j_1-2} (\bar{\tau} - \bar{\tau}_1)^{j_1} \overline{G_{k_1}(\tau_1)} \\ &\left. + \int_{\bar{\tau}}^{-i\infty} d\bar{\tau}_1 (\tau - \bar{\tau}_1)^{k_1-j_1-2} (\bar{\tau} - \bar{\tau}_1)^{j_1} \overline{G_{k_1}(\tau_1)} \int_{\bar{\tau}_1}^{-i\infty} d\bar{\tau}_2 (\tau - \bar{\tau}_2)^{k_2-j_2-2} (\bar{\tau} - \bar{\tau}_2)^{j_2} \overline{G_{k_2}(\tau_2)} \right\} \end{aligned} \quad (2.4b)$$

with $0 \leq j_i \leq k_i-2$. The holomorphic Eisenstein series are normalised as

$$G_k(\tau) = \sum_{(m,n) \neq (0,0)} \frac{1}{(m\tau + n)^k} = 2\zeta_k + \frac{2(2\pi i)^k}{(k-1)!} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \quad (2.5)$$

with divisor sum $\sigma_s(n) = \sum_{d|n} d^s$. The integrals (2.4) have to be understood with tangential base-point regularisation [35] and satisfy the shuffle relations

$$\beta^{\text{sv}} \left[\begin{matrix} j_1 \\ k_1 \end{matrix} \right] \beta^{\text{sv}} \left[\begin{matrix} j_2 \\ k_2 \end{matrix} \right] = \beta^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix} \right] + \beta^{\text{sv}} \left[\begin{matrix} j_2 & j_1 \\ k_2 & k_1 \end{matrix} \right], \quad (2.6)$$

³We shall often suppress the argument τ of various functions to simplify the notation.

as well as the differential equations [30]

$$-4\pi\nabla\beta^{\text{sv}}\left[\begin{matrix} j \\ k \end{matrix}; \tau\right] = (k-2-j)\beta^{\text{sv}}\left[\begin{matrix} j+1 \\ k \end{matrix}; \tau\right] - \delta_{j,k-2}(\tau-\bar{\tau})^k G_k(\tau), \quad (2.7a)$$

$$\begin{aligned} -4\pi\nabla\beta^{\text{sv}}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau\right] &= (k_1-j_1-2)\beta^{\text{sv}}\left[\begin{matrix} j_1+1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau\right] + (k_2-j_2-2)\beta^{\text{sv}}\left[\begin{matrix} j_1 & j_2+1 \\ k_1 & k_2 \end{matrix}; \tau\right] \\ &\quad - \delta_{j_2,k_2-2}(\tau-\bar{\tau})^{k_2} G_{k_2}(\tau)\beta^{\text{sv}}\left[\begin{matrix} j_1 \\ k_1 \end{matrix}; \tau\right]. \end{aligned} \quad (2.7b)$$

The objects $\alpha\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right]$ appearing in (2.4b) are purely antiholomorphic functions and constrained by the shuffle relation (2.6). They are not fixed by the differential equation (2.7) and therefore referred to as integration constants — see [30] for a detailed discussion. A method to determine them from the reality properties of $F_{m,k}^{\pm(s)}$ is discussed in Part I, and a large number of examples can be found in an ancillary file.

2.1.1 Fourier expansions of iterated Eisenstein integrals

The compact definition (2.4) of the β^{sv} can be unpackaged to yield expressions in terms of other iterated integrals of the form [54, 20]

$$\mathcal{E}_0(k, 0^p; \tau) = \frac{(2\pi i)^{p+1-k}}{p!} \int_{\tau}^{i\infty} d\tau_1 (\tau-\tau_1)^p G_k^0(\tau_1), \quad (2.8a)$$

$$\mathcal{E}_0(k_1, 0^{p_1}, k_2, 0^{p_2}; \tau) = \frac{(2\pi i)^{p_1+p_2+2-k_1-k_2}}{p_1!p_2!} \int_{\tau}^{i\infty} d\tau_2 (\tau-\tau_2)^{p_2} G_{k_2}^0(\tau_2) \int_{\tau_2}^{i\infty} d\tau_1 (\tau_2-\tau_1)^{p_1} G_{k_1}^0(\tau_1) \quad (2.8b)$$

and their complex conjugates with integers $k, k_1, k_2 = 4, 6, 8, \dots$ and $p, p_1, p_2 \geq 0$. In the above expressions, 0^p is a placeholder for p successive zeroes (reminiscent of integration kernels $G_k^0 = -1$ [54]), and the G_k^0 are obtained from the holomorphic Eisenstein series G_k by removing the zero mode

$$G_k^0(\tau) = \begin{cases} G_k(\tau) - 2\zeta_k & : k > 0 \text{ even}, \\ 0 & : k > 0 \text{ odd}. \end{cases} \quad (2.9)$$

The removal of the zero mode destroys the good modular transformations of G_k but renders the integrals convergent without regularisation. Moreover, the integrals $\mathcal{E}_0(\dots)$ have fully explicit q -expansions, e.g.

$$\mathcal{E}_0(k, 0^p; \tau) = -\frac{2}{(k-1)!} \sum_{m=1}^{\infty} \frac{\sigma_{k-1}(m)}{m^{p+1}} q^m, \quad (2.10a)$$

$$\mathcal{E}_0(k_1, 0^{p_1}, k_2, 0^{p_2}; \tau) = \frac{4}{(k_1-1)!(k_2-1)!} \sum_{m,n=1}^{\infty} \frac{\sigma_{k_1-1}(m)\sigma_{k_2-1}(n)}{m^{p_1+1}(m+n)^{p_2+1}} q^{m+n}, \quad (2.10b)$$

see [20, Eq. (2.21)] for arbitrary depth. This can be used to obtain the full Fourier expansions of the β^{sv} . The rewriting of the β^{sv} in terms of the $\mathcal{E}_0(\dots)$ requires a number of steps that are well-understood and whose precise form can be found in section 3.3 and appendix D of [20] as well as appendix G of [30].

2.1.2 Differential equations and non-holomorphic Eisenstein series

The definition (2.8a) readily implies the Cauchy–Riemann derivative

$$\pi \nabla \mathcal{E}_0(k, 0^p) = -4y^2 \mathcal{E}_0(k, 0^{p-1}) \quad (2.11)$$

for $p \geq 0$, where we define $\mathcal{E}_0(k, 0^{-1}) = -(2\pi i)^{-k} G_k^0$, and the Laplace equation $\Delta \mathcal{E}_0(k, 0^p) = 0$.

The non-holomorphic Eisenstein series E_k defined in (1.2) can be decomposed in terms of iterated Eisenstein integrals as follows [55, 7]:

$$\begin{aligned} E_k(\tau) &= (-1)^{k-1} \frac{B_{2k}}{(2k)!} (4y)^k + \frac{4(2k-3)! \zeta_{2k-1}}{(k-2)!(k-1)!} (4y)^{1-k} \\ &\quad - 2 \frac{\Gamma(2k)}{\Gamma(k)} \sum_{\ell=0}^{k-1} (4y)^{-\ell} \frac{\Gamma(k+\ell)}{\ell! \Gamma(k-\ell)} \operatorname{Re} \mathcal{E}_0(2k, 0^{k-1+\ell}; \tau) \\ &= (-1)^{k-1} \frac{B_{2k}}{(2k)!} (4y)^k + \frac{4(2k-3)! \zeta_{2k-1}}{(k-2)!(k-1)!} (4y)^{1-k} \\ &\quad + \left[-\frac{1}{2\pi i} \frac{\Gamma(2k)}{[\Gamma(k)]^2} (4y)^{1-k} \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^{k-1} (\bar{\tau} - \tau_1)^{k-1} G_{2k}^0(\tau_1) + \text{c.c.} \right] \\ &= \frac{(2k-1)!}{[(k-1)!]^2} \left\{ -\beta^{\text{sv}} \left[\begin{matrix} k-1 \\ 2k \end{matrix} \right] + \frac{2\zeta_{2k-1}}{(2k-1)(4y)^{k-1}} \right\}, \end{aligned} \quad (2.12)$$

where +c.c. instructs us to add the complex conjugate, making E_k real-analytic and even under $\tau \rightarrow -\bar{\tau}$. We also used $y = \pi \operatorname{Im} \tau$ as we shall do frequently. This relation between E_k and the depth-one β^{sv} , already present in [30], comes directly from (2.4a) when using the relation between G_k and G_k^0 as well as tangential base-point regularisation.

From both the lattice-sum representation (1.2) and the final form of (2.12), one can show the well-known formula for the k -th Cauchy–Riemann derivative of E_k [8]:

$$(\pi \nabla)^k E_k(\tau) = \frac{\Gamma(2k)}{\Gamma(k)} (\operatorname{Im} \tau)^{2k} [2\zeta_{2k} + G_{2k}^0(\tau)] = \frac{\Gamma(2k)}{\Gamma(k)} (\operatorname{Im} \tau)^{2k} G_{2k}(\tau), \quad (2.13)$$

where we have used Euler’s formula relating the even Bernoulli numbers to the even Riemann zeta values

$$2\zeta_{2n} = (-1)^{n+1} \frac{4^n \pi^{2n}}{(2n)!} B_{2n}, \quad n = 1, 2, 3, \dots \quad (2.14)$$

We also record the following general formula

$$(\pi \nabla)^s \left[iy^{1-s} \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^{s-1} (\bar{\tau} - \tau_1)^{s-1} f(\tau_1) \right] = 2^{2s-1} \pi \Gamma(s) (\operatorname{Im} \tau)^{2s} f(\tau) \quad (2.15)$$

for any integer $s > 0$ and holomorphic function $f(\tau)$ irrespective of its modular properties. If $f(\tau)$ has a q -expansion in terms of positive powers of q only, the integral in (2.15) is well-defined without tangential base-point regularisation. With (2.15) and (2.3) it is easy to demonstrate (2.13).

For the Laplacian there is a similar lemma given by

$$(\Delta - s(s-1)) \left[iy^{1-s} \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^{s-1} (\bar{\tau} - \tau_1)^{s-1} f(\tau_1) \right] = 0. \quad (2.16)$$

Besides direct evaluation of the Laplacian on the integral, we can also consider (2.16) by Fourier expanding the integrand $f(\tau)$. Specialising to the case of a single Fourier mode $f(\tau) = e^{2\pi i n \tau}$ with $n > 0$, the integral can be evaluated in terms of Bessel functions $K_{s-1/2}$ giving

$$iy^{1-s} \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^{s-1} (\bar{\tau} - \tau_1)^{s-1} e^{2\pi i n \tau_1} = \frac{(-1)^s \Gamma(s)}{\pi^{2s-1} n^s} \sqrt{\frac{ny}{\pi}} K_{s-1/2}(2ny) e^{2\pi i n \operatorname{Re} \tau}, \quad (2.17)$$

which is the well-known solution to the Laplace equation (2.16) in the n^{th} Fourier mode sector.

From this and (2.3) one can also prove the classic Laplace equation

$$(\Delta - k(k-1))E_k(\tau) = 0, \quad (2.18)$$

since the Laurent monomials y^k and y^{1-k} in (2.12) are in the kernel of $(\Delta - k(k-1))$.

2.1.3 Multiple modular values

Besides the version of iterated Eisenstein integrals in (2.8), we shall also make use of

$$\mathcal{G} \left[\begin{matrix} j \\ k \end{matrix}; \tau \right] = \int_{\tau}^{i\infty} d\tau_1 \tau_1^j G_k(\tau_1), \quad (2.19a)$$

$$\mathcal{G} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau \right] = \int_{\tau}^{i\infty} d\tau_2 \tau_2^{j_2} G_{k_2}(\tau_2) \int_{\tau_2}^{i\infty} d\tau_1 \tau_1^{j_1} G_{k_1}(\tau_1), \quad (2.19b)$$

that are, up to normalisation conventions, Brown's holomorphic iterated Eisenstein integrals and require tangential base-point regularisation [35]. At depth one, this regularisation means

$$\mathcal{G} \left[\begin{matrix} j \\ k \end{matrix}; \tau \right] = \int_{\tau}^{i\infty} d\tau_1 \tau_1^j G_k^0(\tau_1) - 2\zeta_k \int_0^{\tau} d\tau_1 \tau_1^j, \quad (2.20)$$

treating the zero mode of (2.5) differently, while the depth-two generalisation can be found in [35, Eq. (4.13)]. A more general translation of (2.19) into the integrals (2.8) can be found in [20], and the depth-one instance of the dictionary is

$$\mathcal{E}_0(k, 0^p) = (2\pi i)^{p+1-k} \sum_{a=0}^p \frac{(-1)^a \tau^{p-a}}{a!(p-a)!} \mathcal{G} \left[\begin{matrix} a \\ k \end{matrix} \right] + 2\zeta_k (2\pi i)^{p+1-k} \frac{\tau^{p+1}}{(p+1)!}. \quad (2.21)$$

The extra term proportional to ζ_k is due to $\mathcal{E}_0(k, 0^p)$ being defined in terms of G_k^0 , thus lacking the zero mode ζ_k when compared to G_k appearing in (2.19), see (2.9).

The virtue of the definition (2.19) is that it is easier to describe the behaviour under S-modular transformations [35, 20]:

$$\mathcal{G} \left[\begin{matrix} j \\ k \end{matrix}; -\frac{1}{\tau} \right] = (-1)^j \mathcal{G} \left[\begin{matrix} k-2-j \\ k \end{matrix}; \tau \right] + \mathbf{m} \left[\begin{matrix} j \\ k \end{matrix} \right], \quad (2.22a)$$

$$\mathcal{G} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; -\frac{1}{\tau} \right] = (-1)^{j_1+j_2} \mathcal{G} \left[\begin{matrix} k_1-2-j_1 & k_2-2-j_2 \\ k_1 & k_2 \end{matrix}; \tau \right] + (-1)^{j_2} \mathcal{G} \left[\begin{matrix} k_2-2-j_2 \\ k_2 \end{matrix}; \tau \right] \mathbf{m} \left[\begin{matrix} j_1 \\ k_1 \end{matrix} \right] + \mathbf{m} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix} \right]. \quad (2.22b)$$

The objects $\mathbf{m}[\dots]$ appearing in this equation do not depend on τ – they are examples of multiple modular values [35] and correspond to period integrals

$$\mathbf{m}\left[\begin{smallmatrix} j \\ k \end{smallmatrix}\right] = \int_0^{i\infty} d\tau_1 \tau_1^j G_k(\tau_1), \quad (2.23a)$$

$$\mathbf{m}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}\right] = \int_0^{i\infty} d\tau_2 \tau_2^{j_2} G_{k_2}(\tau_2) \int_{\tau_2}^{i\infty} d\tau_1 \tau_1^{j_1} G_{k_1}(\tau_1), \quad (2.23b)$$

which are obtained formally as limits $\tau \rightarrow 0$ of (2.19). This limit is divergent and has to be treated again with tangential base-point regularisation. One way of doing this is to consider at depths one and two

$$\mathbf{m}\left[\begin{smallmatrix} j \\ k \end{smallmatrix}\right] = \mathcal{G}\left[\begin{smallmatrix} j \\ k \end{smallmatrix}; i\right] - (-1)^j \mathcal{G}\left[\begin{smallmatrix} k-2-j \\ k \end{smallmatrix}; i\right], \quad (2.24a)$$

$$\mathbf{m}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}\right] = \mathcal{G}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}; i\right] - (-1)^{j_1+j_2} \mathcal{G}\left[\begin{smallmatrix} k_1-2-j_1 & k_2-2-j_2 \\ k_1 & k_2 \end{smallmatrix}; i\right] - (-1)^{j_2} \mathcal{G}\left[\begin{smallmatrix} k_2-2-j_2 \\ k_2 \end{smallmatrix}; i\right] \mathbf{m}\left[\begin{smallmatrix} j_1 \\ k_1 \end{smallmatrix}\right], \quad (2.24b)$$

where we rewrote the S-modular behaviour (2.22) and evaluated this expression at the self-dual point $\tau = i$. We recall that the integrals (2.19) are well-defined for any finite τ , using tangential base-point regularisation at the upper integration boundary $\tau \rightarrow i\infty$. The choice of the self-dual point $\tau = i$ in (2.24) is arbitrary (any pair of S-dual points would do) but convenient for numerical evaluations.

For depth one we can work out the multiple modular values explicitly as

$$\mathbf{m}\left[\begin{smallmatrix} j \\ k \end{smallmatrix}\right] = \begin{cases} -\frac{2(2\pi i)^{k-j-1} j!}{(k-1)!} \zeta_{j+1} \zeta_{j+2-k} & \text{for } j > 0, \\ -\frac{2\pi i \zeta_{k-1}}{k-1} & \text{for } j = 0 \end{cases} \quad (2.25)$$

and these correspond to periods of the holomorphic Eisenstein series [56]. The $j = 0$ case can also be obtained as a limit after using the functional relations of the zeta function. Since $k \geq 4$ is an even integer, the multiple modular values of depth one vanish for even $0 < j < k-2$ as they involve the zeta function evaluated at a negative even integer.

For depth two, numbers beyond (multiple) zeta values can occur [35, 50]. We will discuss further properties of multiple modular values and how they arise directly in the S-modular transformation of the β^{sv} in section 2.3.

2.2 Iterated integrals of cusp forms

We now let $\Delta_{2s}(\tau)$ denote a holomorphic cusp form of weight $2s \in \{12, 16, 18, \dots\}$. Then we define the analogue of (2.8a) as

$$\begin{aligned} \mathcal{E}_0(\Delta_{2s}, 0^p; \tau) &= (-1)^p (2\pi i)^{p+1} \int_{\tau}^{i\infty} d\tau_1 \int_{\tau_1}^{i\infty} d\tau_2 \dots \int_{\tau_p}^{i\infty} d\tau_{p+1} \Delta_{2s}(\tau_{p+1}) \\ &= \frac{(2\pi i)^{p+1}}{p!} \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^p \Delta_{2s}(\tau_1). \end{aligned} \quad (2.26)$$

Since Δ_{2s} is a cusp form, this integral is well-defined for any $p \geq 0$; however, in everything that follows we shall only encounter the usual range of values $0 \leq p \leq 2s-2$. The cusp forms in this definition are Hecke normalised with $\Delta_{2s}(\tau) = q + O(q^2)$ such that the transcendentalty of the iterated integral (2.26) is given by $p+1$, just like for (2.8a).⁴ The objects that are on a similar footing are Δ_{2s} and $\frac{G_k}{(2\pi i)^k}$ since both have algebraic Fourier coefficients, for instance

$$\begin{aligned}\Delta_{12}(\tau) &= \frac{1}{1728} \left\{ \left(\frac{G_4(\tau)}{2\zeta_4} \right)^3 - \left(\frac{G_6(\tau)}{2\zeta_6} \right)^2 \right\} = q - 24q^2 + 252q^3 - 1472q^4 + O(q^5) \\ \Delta_{16}(\tau) &= \frac{1}{1728} \frac{G_4(\tau)}{2\zeta_4} \left\{ \left(\frac{G_4(\tau)}{2\zeta_4} \right)^3 - \left(\frac{G_6(\tau)}{2\zeta_6} \right)^2 \right\} = q + 216q^2 - 3348q^3 + 13888q^4 + O(q^5) \\ \Delta_{18}(\tau) &= \frac{1}{1728} \frac{G_6(\tau)}{2\zeta_6} \left\{ \left(\frac{G_4(\tau)}{2\zeta_4} \right)^3 - \left(\frac{G_6(\tau)}{2\zeta_6} \right)^2 \right\} = q - 528q^2 - 4284q^3 + 147712q^4 + O(q^5).\end{aligned}\tag{2.27}$$

If the cusp form has Fourier expansion $\Delta_{2s}(\tau) = \sum_{n=1}^{\infty} a(n)q^n$ then

$$\mathcal{E}_0(\Delta_{2s}, 0^p; \tau) = - \sum_{n=1}^{\infty} n^{-p-1} a(n)q^n.\tag{2.28}$$

While the seed functions of $F_{m,k}^{+(s)}$ and $F_{m,k}^{-(s)}$ in (1.5) determined in Part I are essentially⁵ constructed from real and imaginary parts of $\mathcal{E}_0(2m, 0^p; \tau)$ in (2.8a), respectively, Poincaré sums of $\mathcal{E}_0(\Delta_{2s}, 0^p; \tau)$ have been discussed in [52].

2.2.1 Real-analytic integrals of holomorphic cusp forms

From the fourth line of (2.12), we see that one can define even and odd analogues of the non-holomorphic Eisenstein series by trading G_k in the integration kernel for holomorphic cusp forms Δ_{2s} ,

$$\mathbf{H}_{\Delta_{2s}}^{\pm}(\tau) = (-1)^s \frac{\pi^{2s-1} i}{\Gamma(s)} y^{1-s} \int_{\tau}^{i\infty} d\tau_1 (\tau - \tau_1)^{s-1} (\bar{\tau} - \tau_1)^{s-1} \Delta_{2s}(\tau_1) \pm \text{c.c.},\tag{2.29}$$

where we have fixed a convenient normalisation. This function satisfies from (2.15) and (2.16)

$$(\pi \nabla)^s \mathbf{H}_{\Delta_{2s}}^{\pm} = \frac{1}{2} (2\pi i)^{2s} (\text{Im } \tau)^{2s} \Delta_{2s}(\tau),\tag{2.30a}$$

$$(\Delta - s(s-1)) \mathbf{H}_{\Delta_{2s}}^{\pm} = 0.\tag{2.30b}$$

Clearly, the even function $\mathbf{H}_{\Delta_{2s}}^{+}$ is obtained from the cusp form Δ_{2s} in the same way as E_s is obtained from G_{2s} . Moreover $\mathbf{H}_{\Delta_{2s}}^{-}$ is its odd cousin, and the appearance of an odd analogue of E_k (denoted by $E_k^{(-)}$) in the lower-depth terms of $F_{m,k}^{-(s)}$ is discussed in section 5.5 of Part I. Variants of (2.29) with more general exponents $(\tau - \tau_1)^j (\bar{\tau} - \tau_1)^{2s-2-j}$, $j = 0, 1, \dots, 2s-2$ have been studied in [52].

⁴The factor of $(-1)^p$ in the first line of (2.26) is due to the insertion of p copies of $G_0^0 = -1$ in the iterated integral which is the meaning of the notation 0^p [54, 20].

⁵The seed functions of $F_{m,k}^{+(s)}$ constructed in Part I additionally involve \mathbb{Q} -multiples of y^{m+k} and $\zeta_{2m-1} y^{1-m+k}$.

Following our discussion around (2.17) we expose the q -expansion of $H_{\Delta_{2s}}^{\pm}$ by rewriting them as a finite sum over the $\mathcal{E}_0(\Delta_{2s}, 0^p; \tau)$ in (2.26) and (2.28),

$$\begin{aligned} H_{\Delta_{2s}}^{\pm}(\tau) &= \frac{1}{2} \sum_{\ell=0}^{s-1} (4y)^{-\ell} \frac{\Gamma(s+\ell)}{\ell! \Gamma(s-\ell)} \sum_{n=1}^{\infty} \frac{a(n)}{n^{s+\ell}} (q^n \pm \bar{q}^n) \\ &= -\frac{1}{2} \sum_{\ell=0}^{s-1} (4y)^{-\ell} \frac{\Gamma(s+\ell)}{\ell! \Gamma(s-\ell)} \left[\mathcal{E}_0(\Delta_{2s}, 0^{s-1+\ell}; \tau) \pm \text{c.c.} \right], \end{aligned} \quad (2.31)$$

which is the direct analogue of the second line in (2.12) for E_s and the definition of $E_s^{(-)}$ in section 5.5 of Part I. The sum can also be recast in terms of Bessel functions using (2.17) as

$$H_{\Delta_{2s}}^{\pm}(\tau) = \sum_{n=1}^{\infty} a(n) n^{-s} \sqrt{\frac{ny}{\pi}} K_{s-1/2}(2ny) (e^{2\pi i n \operatorname{Re} \tau} \pm e^{-2\pi i n \operatorname{Re} \tau}). \quad (2.32)$$

Even though the functions $H_{\Delta_{2s}}^{\pm}$ and the modular invariant functions defined in [47] are both real-analytic and are both obtained from iterated integrals of holomorphic cusp forms Δ_{2s} , they differ crucially in their modular properties. In particular, as we shall show next, the $H_{\Delta_{2s}}^{\pm}$ are not invariant under S-modular transformations.

2.2.2 Modular properties

For studying modular transformations of functions $F(\tau)$ on the upper half-plane, it is convenient to introduce the following cocycles under the generating T- and S-transformations of $\mathrm{SL}(2, \mathbb{Z})$:

$$\delta_{\mathrm{T}} F(\tau) = F(\tau) - F(\tau+1), \quad \delta_{\mathrm{S}} F(\tau) = F(\tau) - F(-\frac{1}{\tau}). \quad (2.33)$$

When both of them vanish, F is invariant under modular transformations and in general there is a connection to the group cohomology of $\mathrm{SL}(2, \mathbb{Z})$ [57].

From the explicit q -series in (2.31), it is clear that the functions $H_{\Delta_{2s}}^{\pm}$ have a vanishing T-cocycle

$$\delta_{\mathrm{T}} H_{\Delta_{2s}}^{\pm}(\tau) = H_{\Delta_{2s}}^{\pm}(\tau) - H_{\Delta_{2s}}^{\pm}(\tau+1) = 0. \quad (2.34)$$

However, the functions $H_{\Delta_{2s}}^{\pm}$ have a non-trivial cocycle under the S-transformation

$$\begin{aligned} \delta_{\mathrm{S}} H_{\Delta_{2s}}^{\pm}(\tau) &= H_{\Delta_{2s}}^{\pm}(\tau) - H_{\Delta_{2s}}^{\pm}(-\frac{1}{\tau}) = (-1)^s \frac{i\pi^{2s-1}}{\Gamma(s)} y^{1-s} \int_0^{i\infty} d\tau_1 (\tau - \tau_1)^{s-1} (\bar{\tau} - \tau_1)^{s-1} \Delta_{2s}(\tau_1) \pm \text{c.c.} \\ &= (-1)^{s+1} \frac{\pi^{2s-1}}{\Gamma(s)} y^{1-s} \sum_{a,b=0}^{s-1} (-i)^{a+b} \binom{s-1}{a} \binom{s-1}{b} \tau^{s-1-a} \bar{\tau}^{s-1-b} \Lambda(\Delta_{2s}, a+b+1) \pm \text{c.c.} \\ &= \begin{cases} 2(-1)^{s+1} \frac{\pi^{2s-1}}{\Gamma(s)} y^{1-s} \sum_{\substack{a,b=0 \\ a+b \in 2\mathbb{Z}}}^{s-1} (-1)^{(a+b)/2} \binom{s-1}{a} \binom{s-1}{b} \tau^{s-1-a} \bar{\tau}^{s-1-b} \Lambda(\Delta_{2s}, a+b+1) \\ 2(-1)^s \frac{i\pi^{2s-1}}{\Gamma(s)} y^{1-s} \sum_{\substack{a,b=0 \\ a+b \in 2\mathbb{Z}+1}}^{s-1} (-1)^{(a+b-1)/2} \binom{s-1}{a} \binom{s-1}{b} \tau^{s-1-a} \bar{\tau}^{s-1-b} \Lambda(\Delta_{2s}, a+b+1), \end{cases} \end{aligned} \quad (2.35)$$

where in the last equality the upper line is for $H_{\Delta_{2s}}^+$ and the lower line for $H_{\Delta_{2s}}^-$. The function $\Lambda(\Delta_{2s}, t)$ appearing in the above expressions is real-valued for real t and corresponds to the completed L-function of the cusp form $\Delta_{2s}(\tau) = \sum_{n=1}^{\infty} a(n)q^n$ of weight $2s$ defined by

$$\Lambda(\Delta_{2s}, t) = (2\pi)^{-t} \Gamma(t) \sum_{n=1}^{\infty} \frac{a(n)}{n^t} = (-i)^t \int_0^{i\infty} d\tau_1 \tau_1^{t-1} \Delta_{2s}(\tau_1), \quad (2.36)$$

where the sum converges absolutely for $\text{Re}(t) > s + \frac{1}{2}$, using [58] and the improved growth of the Fourier coefficients of cusp forms following from [59].⁶ From its integral Mellin-form representation, it is well-known that the completed L-function enjoys an analytic continuation to the complex plane and satisfies the functional relation

$$\Lambda(\Delta_{2s}, t) = (-1)^s \Lambda(\Delta_{2s}, 2s-t). \quad (2.37)$$

The interval $t \in (0, 2s)$ is called the critical strip, and (2.35) implies that the failure of modularity of $H_{\Delta_{2s}}^{\pm}$ involves the completed L-function evaluated at integers inside the critical strip, with only odd integer arguments t contributing to $H_{\Delta_{2s}}^+$ and only even integers for $H_{\Delta_{2s}}^-$.

2.2.3 Integrals of Hecke normalised holomorphic cusp forms

If Δ_{2s} is a normalised eigenform of all Hecke operators T_n and of weight $2s$, i.e.,

$$[T_n \Delta_{2s}](\tau) = n^{2s-1} \sum_{d|n} d^{-2s} \sum_{b=0}^{d-1} \Delta_{2s}\left(\frac{n\tau + bd}{d^2}\right) = a(n) \Delta_{2s}(\tau) \quad (2.38)$$

for all $n > 0$, implying Hecke normalisation $a(1) = 1$, then it is moreover known that the values $\Lambda(\Delta_{2s}, t)$ for all even t inside the critical strip are related, as are all the values for odd t [45, 46, 60]. The ratios between the even (or the odd) values must belong to the number-field extension of \mathbb{Q} defined by the Fourier coefficients $\{a(n), n \in \mathbb{N}\}$ of Δ_{2s} . The first time a non-trivial extension arises is for cusp forms of weight $2s = 24$ where there are two linearly independent cusp forms and the number field is $\mathbb{Q}(\sqrt{144169})$. The non-trivial Galois automorphism of the number field exchanges the two independent Hecke eigenforms.

Therefore, for (normalised) Hecke eigenforms Δ_{2s} of weight $2s$ we find that we can rearrange (2.35) to

$$\delta_S H_{\Delta_{2s}}^+(\tau) = \frac{2(-1)^{s+1} \pi^{2s-1}}{\Gamma(s)} y^{1-s} \Lambda(\Delta_{2s}, 2s-1) \sum_{\substack{a,b=0 \\ a+b \in 2\mathbb{Z}}}^{s-1} (-1)^{\frac{a+b}{2}} \binom{s-1}{a} \binom{s-1}{b} \tau^{s-1-a} \bar{\tau}^{s-1-b} c_{a+b}^+, \quad (2.39a)$$

$$\delta_S H_{\Delta_{2s}}^-(\tau) = \frac{2(-1)^s i \pi^{2s-1}}{\Gamma(s)} y^{1-s} \Lambda(\Delta_{2s}, 2s-2) \sum_{\substack{a,b=0 \\ a+b \in 2\mathbb{Z}+1}}^{s-1} (-1)^{\frac{a+b-1}{2}} \binom{s-1}{a} \binom{s-1}{b} \tau^{s-1-a} \bar{\tau}^{s-1-b} c_{a+b}^-, \quad (2.39b)$$

⁶We thank Nils Matthes for correspondence on this point.

with coefficients c_ℓ^\pm from the number field associated with Δ_{2s} defined by

$$\Lambda(\Delta_{2s}, \ell+1) = \begin{cases} c_\ell^+ \Lambda(\Delta_{2s}, 2s-1) & : \ell \text{ even,} \\ c_\ell^- \Lambda(\Delta_{2s}, 2s-2) & : \ell \text{ odd,} \end{cases} \quad (2.40)$$

for $\ell = 0, 1, \dots, 2s-2$. The polynomials arising in (2.39) have also appeared in [35, 50] and are related to period polynomials as we also discuss in section 4.1.1.

We had argued above that $H_{\Delta_{2s}}^\pm$ should be assigned transcendental weight s and this is consistent with the fact that there is no transcendental weight carried by L-values inside the critical strip $(0, 2s)$ like $\Lambda(\Delta_{2s}, 2s-1)$ and $\Lambda(\Delta_{2s}, 2s-2)$. As we will see later on, the value $\Lambda(\Delta_{2s}, 2s+m)$ has transcendental weight $(m+1)$, so that just at the end of the critical strip we have $\Lambda(\Delta_{2s}, 2s)$ of transcendental weight one. This is analogous to the Riemann zeta function whose transcendental weight grows in the same way from the upper end of its critical strip $(0, 1)$. The transcendental weight of the cocycles (2.39) is therefore determined by the prefactor, including $y = \pi \operatorname{Im} \tau$, and is also given by s , consistent with that of $H_{\Delta_{2s}}^\pm$ itself.

Note that with our definition (2.29) for the iterated integral of a cusp form, the S-cocycles in (2.35) or (2.39) are two-variable generalisations of the classic period polynomial associated to the same cusp form [56], with $\delta_S H_{\Delta_{2s}}^\pm(\tau)$ playing the roles of the even/odd part of said period polynomial. Furthermore, along the same lines as Manin's original work [60], we have that $\delta_S H_{\Delta_{2s}}^\pm(\tau)$ satisfies the two cocycle conditions [57]

$$\left[\delta_S H_{\Delta_{2s}}^\pm(\tau) \right] \Big|_{1+S} = 0, \quad (2.41a)$$

$$\left[\delta_S H_{\Delta_{2s}}^\pm(\tau) \right] \Big|_{1+U+U^2} = 0, \quad (2.41b)$$

where $|$ denotes the $\operatorname{SL}(2, \mathbb{Z})$ -action on τ and $U = \operatorname{TS}$ is an order 3 generator of $\operatorname{SL}(2, \mathbb{Z})$.

When applied to (2.35), the first cocycle condition (2.41a) is equivalent to the reflection formula (2.37) for the even/odd values inside the critical strip. Similarly the second condition (2.41b), together with the Hecke condition (2.38), is equivalent to the statement that all the ratios between the even (or odd) critical values must be in the number field generated by the Fourier coefficients. In (2.39) we chose to factorise out $\Lambda(\Delta_{2s}, 2s-1)$ and $\Lambda(\Delta_{2s}, 2s-2)$, respectively, thus making a particular choice for what are usually called the (holomorphic) periods of the cusp form Δ_{2s} , sometimes denoted by $\omega_{\Delta_{2s}}^\pm$ [50].

2.2.4 Example with $s = 6$

As a concrete example we can study the cusp form of lowest weight $2s = 12$, i.e. the Ramanujan cusp form $\Delta_{2s} = \Delta_{12} = \sum_{n=1}^\infty \tau(n)q^n$ with $\tau(1) = 1$. Since the vector space of cusp forms at weight $2s = 12$ is one-dimensional we trivially have that Δ_{12} is a normalised Hecke eigenform and obviously the associated number field is simply \mathbb{Q} , i.e. $\tau(n) \in \mathbb{Q}$ for all $n > 0$. Following [60] we have the following number-field relations amongst the completed L-values, even and odd,

inside the critical strip:

$$\Lambda(\Delta_{12}, 6) : \Lambda(\Delta_{12}, 8) : \Lambda(\Delta_{12}, 10) = \frac{5}{12} : \frac{25}{48} : 1, \quad (2.42)$$

$$\Lambda(\Delta_{12}, 7) : \Lambda(\Delta_{12}, 9) : \Lambda(\Delta_{12}, 11) = \frac{691}{2520} : \frac{691}{1620} : 1,$$

where the remaining values can be obtained via the reflection formula $\Lambda(\Delta_{12}, 12-t) = \Lambda(\Delta_{12}, t)$.

If we compute the cocycles (2.39) defined above we obtain

$$\begin{aligned} H_{\Delta_{12}}^+(\tau) - H_{\Delta_{12}}^+\left(-\frac{1}{\tau}\right) &= \frac{\pi^{11}\Lambda(\Delta_{12}, 11)(1-\tau\bar{\tau})}{60y^5} \left\{ 1 + \tau^4\bar{\tau}^4 - \frac{691}{162}(\tau^2+\bar{\tau}^2)(1+\tau^2\bar{\tau}^2) \right. \\ &\quad \left. + \frac{691}{504}(\tau^4+\bar{\tau}^4) - \frac{3131\tau\bar{\tau}}{324}(1+\tau^2\bar{\tau}^2) + \frac{21421\tau\bar{\tau}}{2268}(\tau^2+\bar{\tau}^2) + \frac{40273\tau^2\bar{\tau}^2}{2268} \right\} \end{aligned} \quad (2.43a)$$

$$\begin{aligned} H_{\Delta_{12}}^-(\tau) - H_{\Delta_{12}}^-\left(-\frac{1}{\tau}\right) &= \frac{i\pi^{11}\Lambda(\Delta_{12}, 10)(\tau+\bar{\tau})}{12y^5} \left\{ 1 + \tau^4\bar{\tau}^4 - \frac{25}{24}(\tau^2+\bar{\tau}^2)(1+\tau^2\bar{\tau}^2) \right. \\ &\quad \left. + \frac{1}{12}(\tau^4+\bar{\tau}^4) - \frac{25}{6}\tau\bar{\tau}(1+\tau^2\bar{\tau}^2) + 2\tau\bar{\tau}(\tau^2+\bar{\tau}^2) + \frac{19\tau^2\bar{\tau}^2}{3} \right\}. \end{aligned} \quad (2.43b)$$

Similar expressions for the cocycles of $H_{\Delta_{16}}^\pm$ and $H_{\Delta_{18}}^\pm$ can be found in appendix B, while in the ancillary file the expressions are given up to modular weight $2s = 26$.

In summary, we can construct even and odd solutions $H_{\Delta_{2s}}^\pm$ of the homogeneous Laplace equation (1.1) whenever we have a holomorphic cusp form Δ_{2s} of weight $2s$. These homogeneous solutions are expressible through iterated integrals of Δ_{2s} of depth one. They are not modular invariant but their failure of modularity is characterised by a single number that is a value of the completed L-function inside the critical strip. This number is expected to be independent over \mathbb{Q} from the set of multiple zeta values.

2.3 Properties of multiple modular values and the β^{sv}

We now study the multiple modular values defined in (2.23) in more detail and also present expressions for the S-modular transformation of the β^{sv} introduced in (2.4).

From their definition (2.23) the multiple modular values inherit the shuffle relations of the (regularised) iterated integrals in (2.19)

$$\mathbf{m} \left[\begin{smallmatrix} j_1 \\ k_1 \end{smallmatrix} \right] \mathbf{m} \left[\begin{smallmatrix} j_2 \\ k_2 \end{smallmatrix} \right] = \mathbf{m} \left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix} \right] + \mathbf{m} \left[\begin{smallmatrix} j_2 & j_1 \\ k_2 & k_1 \end{smallmatrix} \right]. \quad (2.44)$$

By applying another S-transformation to (2.22) one can show the reflection properties

$$\mathbf{m} \left[\begin{smallmatrix} j \\ k \end{smallmatrix} \right] = -(-1)^j \mathbf{m} \left[\begin{smallmatrix} k-2-j \\ k \end{smallmatrix} \right], \quad (2.45a)$$

$$\begin{aligned} \mathbf{m} \left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix} \right] &= -(-1)^{j_1+j_2} \mathbf{m} \left[\begin{smallmatrix} k_1-2-j_1 & k_2-2-j_2 \\ k_1 & k_2 \end{smallmatrix} \right] + \mathbf{m} \left[\begin{smallmatrix} j_1 \\ k_1 \end{smallmatrix} \right] \mathbf{m} \left[\begin{smallmatrix} j_2 \\ k_2 \end{smallmatrix} \right] \\ &= (-1)^{j_1+j_2} \mathbf{m} \left[\begin{smallmatrix} k_2-2-j_2 & k_1-2-j_1 \\ k_2 & k_1 \end{smallmatrix} \right]. \end{aligned} \quad (2.45b)$$

Under complex conjugation they satisfy

$$\overline{\mathbf{m} \left[\begin{smallmatrix} j \\ k \end{smallmatrix} \right]} = (-1)^{j+1} \mathbf{m} \left[\begin{smallmatrix} j \\ k \end{smallmatrix} \right], \quad \overline{\mathbf{m} \left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix} \right]} = (-1)^{j_1+j_2+2} \mathbf{m} \left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix} \right]. \quad (2.46)$$

The transcendentality of the multiple modular values as defined in (2.23) is given by $\sum_i k_i$. For the depth-one case (2.25) this is evident from the fact that $(2\pi i)^{k-j-1}$ has transcendentality $k-j-1$, ζ_{j+1} has transcendentality $j+1$ and ζ_{j+2-k} has transcendentality zero.⁷ In the general case, this follows from the definition (2.19) by realising that G_k in our convention has transcendentality k , see (2.5). The iterated integrals (2.8a) and (2.8b) therefore have transcendentality $p+1$ and p_1+p_2+2 , respectively.

2.3.1 Reduced multiple modular values

While the multiple modular values (2.23) appear in the S-transformation of the holomorphic iterated integrals (2.19), the S-transformation of the β^{sv} only contains the following specific combinations of them:

$$\mathcal{M}\left[\begin{matrix} j \\ k \end{matrix}\right] = \mathfrak{m}\left[\begin{matrix} j \\ k \end{matrix}\right] - \overline{\mathfrak{m}\left[\begin{matrix} j \\ k \end{matrix}\right]} = (1 + (-1)^j) \mathfrak{m}\left[\begin{matrix} j \\ k \end{matrix}\right], \quad (2.47a)$$

$$\begin{aligned} \mathcal{M}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right] &= \mathfrak{m}\left[\begin{matrix} j_2 & j_1 \\ k_2 & k_1 \end{matrix}\right] - \overline{\mathfrak{m}\left[\begin{matrix} j_2 & j_1 \\ k_2 & k_1 \end{matrix}\right]} + \overline{\mathfrak{m}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right]} \\ &= (-1)^{j_1} (1 + (-1)^{j_2}) \mathfrak{m}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right] + (1 + (-1)^{j_1}) \mathfrak{m}\left[\begin{matrix} j_2 & j_1 \\ k_2 & k_1 \end{matrix}\right] \\ &= ((-1)^{j_1+j_2} - 1) \mathfrak{m}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right] + (1 + (-1)^{j_1}) \mathfrak{m}\left[\begin{matrix} j_1 \\ k_1 \end{matrix}\right] \mathfrak{m}\left[\begin{matrix} j_2 \\ k_2 \end{matrix}\right]. \end{aligned} \quad (2.47b)$$

We refer to the combinations $\mathcal{M}\left[\begin{matrix} j \\ k \end{matrix}\right]$ and $\mathcal{M}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right]$ as *reduced multiple modular values* of depth one and two, respectively. The simplifications at depth two are based on (2.44) and (2.46), and they show that reduced multiple modular values vanish if both of j_1 and j_2 are odd, whereas cases with both j_1 and j_2 even yield the product $2\mathfrak{m}\left[\begin{matrix} j_1 \\ k_1 \end{matrix}\right] \mathfrak{m}\left[\begin{matrix} j_2 \\ k_2 \end{matrix}\right]$. Moreover, reduced multiple modular values inherit the shuffle property and so satisfy

$$\mathcal{M}\left[\begin{matrix} j_1 \\ k_1 \end{matrix}\right] \mathcal{M}\left[\begin{matrix} j_2 \\ k_2 \end{matrix}\right] = \mathcal{M}\left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}\right] + \mathcal{M}\left[\begin{matrix} j_2 & j_1 \\ k_2 & k_1 \end{matrix}\right]. \quad (2.48)$$

2.3.2 Depth one reduced multiple modular values and β^{sv} modular transformations

At depth one, (2.25) leads to the following explicit expressions

$$\mathcal{M}\left[\begin{matrix} j \\ k \end{matrix}\right] = \begin{cases} -\frac{4\pi i \zeta_{k-1}}{k-1} & j = 0, \\ +\frac{4\pi i \zeta_{k-1}}{k-1} & j = k-2, \\ 0 & \text{otherwise,} \end{cases} \quad (2.49)$$

where the vanishing of all cases with $j = 1, 2, \dots, k-3$ is in agreement with their occurrence in certain coboundary polynomials [35, 38, 50]. They appear in the transformation of the depth-one

⁷The last statement is true since $j \leq k-2$ and zeta values at non-positive integers are either zero or given by rational numbers (expressible through Bernoulli numbers).

β^{sv} according to

$$\begin{aligned}
\beta^{\text{sv}} \left[\begin{matrix} j \\ k \end{matrix}; -\frac{1}{\bar{\tau}} \right] &= \bar{\tau}^{k-2-2j} \beta^{\text{sv}} \left[\begin{matrix} j \\ k \end{matrix}; \tau \right] - \frac{(\tau\bar{\tau})^{k-2-j}}{2\pi i (4y)^{k-2-j}} \\
&\times \sum_{A=0}^{k-2-j} \sum_{B=0}^j \binom{k-2-j}{A} \binom{j}{B} \left(-\frac{1}{\tau}\right)^A \left(-\frac{1}{\bar{\tau}}\right)^B \mathcal{M} \left[\begin{matrix} A+B \\ k \end{matrix} \right] \\
&= \bar{\tau}^{k-2-2j} \beta^{\text{sv}} \left[\begin{matrix} j \\ k \end{matrix}; \tau \right] + \frac{2\zeta_{k-1}}{(k-1)(4y)^{k-2-j}} \left\{ (\tau\bar{\tau})^{k-2-j} - \bar{\tau}^{k-2-2j} \right\}.
\end{aligned} \tag{2.50}$$

2.3.3 Modular transformation of β^{sv} at depth two

In the same way the depth-one reduced multiple modular values arise in the modular transformations of the depth-one β^{sv} , the depth-two $\mathcal{M} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix} \right]$ arise in the modular transformation of the β^{sv} at depth two. Performing the calculation based on the integral representation (2.4b) one can show

$$\begin{aligned}
\beta^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; -\frac{1}{\bar{\tau}} \right] &= \bar{\tau}^{k_1+k_2-4-2j_1-2j_2} \beta^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau \right] \\
&+ \bar{\tau}^{k_2-2-2j_2} \beta^{\text{sv}} \left[\begin{matrix} j_2 \\ k_2 \end{matrix}; \tau \right] \frac{2\zeta_{k_1-1} \left\{ (\tau\bar{\tau})^{k_1-2-j_1} - \bar{\tau}^{k_1-2-2j_1} \right\}}{(k_1-1)(4y)^{k_1-2-j_1}} \\
&+ \frac{(\tau\bar{\tau})^{k_1+k_2-4-j_1-j_2}}{(2\pi i)^2 (4y)^{k_1+k_2-4-j_1-j_2}} \sum_{A_1=0}^{k_1-2-j_1} \sum_{A_2=0}^{k_2-2-j_2} \sum_{B_1=0}^{j_1} \sum_{B_2=0}^{j_2} \binom{k_1-2-j_1}{A_1} \binom{k_2-2-j_2}{A_2} \\
&\times \binom{j_1}{B_1} \binom{j_2}{B_2} \left(-\frac{1}{\tau}\right)^{A_1+A_2} \left(-\frac{1}{\bar{\tau}}\right)^{B_1+B_2} \mathcal{M} \left[\begin{matrix} A_1+B_1 & A_2+B_2 \\ k_1 & k_2 \end{matrix} \right] \\
&+ \frac{\bar{\tau}^{k_1+k_2-4-2j_1-2j_2}}{(2\pi i)^2 (4y)^{k_1+k_2-4-j_1-j_2}} C \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau \right].
\end{aligned} \tag{2.51}$$

Here, the $C[\dots]$ are pure depth-zero terms, i.e. rational functions of τ and $\bar{\tau}$ multiplied by rational combinations of odd zeta values and powers of π , that can be traced back to the modular transformation of the $\overline{\alpha[\dots]}$. Their definition is most conveniently given in terms of the shorthand

$$\beta_{(\alpha)}^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau \right] = \sum_{p_1=0}^{k_1-2-j_1} \sum_{p_2=0}^{k_2-2-j_2} \frac{\binom{k_1-2-j_1}{p_1} \binom{k_2-2-j_2}{p_2}}{(4y)^{p_1+p_2}} \overline{\alpha \left[\begin{matrix} j_1+p_1 & j_2+p_2 \\ k_1 & k_2 \end{matrix}; \tau \right]} \tag{2.52}$$

for the contributions of the antiholomorphic $\overline{\alpha[\dots]}$ to (2.4b), namely

$$\begin{aligned}
C \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau \right] &= \frac{(2\pi i)^2 (4y)^{k_1+k_2-4-j_1-j_2}}{\bar{\tau}^{k_1+k_2-4-2j_1-2j_2}} \beta_{(\alpha)}^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; -\frac{1}{\bar{\tau}} \right] - (2\pi i)^2 (4y)^{k_1+k_2-4-j_1-j_2} \beta_{(\alpha)}^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix}; \tau \right] \\
&+ 2\pi i \left\{ \frac{2\zeta_{k_2-1}}{k_2-1} (1 - \tau^{k_2-j_2-2\bar{\tau}j_2}) \int_{\bar{\tau}}^{-i\infty} d\bar{\tau}_1 (\tau - \bar{\tau}_1)^{k_1-2-j_1} (\bar{\tau} - \bar{\tau}_1)^{j_1} \overline{G_{k_1}(\tau_1)} \right. \\
&\left. - \frac{2\zeta_{k_1-1}}{k_1-1} (1 - \tau^{k_1-j_1-2\bar{\tau}j_1}) \int_{\bar{\tau}}^{-i\infty} d\bar{\tau}_1 (\tau - \bar{\tau}_1)^{k_2-2-j_2} (\bar{\tau} - \bar{\tau}_1)^{j_2} \overline{G_{k_2}(\tau_1)} \right\}.
\end{aligned} \tag{2.53}$$

Then, the known expressions for $\overline{\alpha[\cdots]}$ [30, 1] and the modular transformations of the $\mathcal{E}_0(k, 0^p)$ (see for instance section 2.1.3) lead to representative examples such as⁸

$$\begin{aligned}
C \left[\begin{smallmatrix} 0 & 1 \\ 4 & 4 \end{smallmatrix} \right] &= \frac{32\bar{\tau}^2\pi^4 y\zeta_3}{405} - \frac{4\pi^4 y\zeta_3}{135} + \frac{128i\bar{\tau}\pi^3 y^2\zeta_3}{405} - \frac{16\pi^2 y^3\zeta_3}{81} - \frac{16\pi^2 y^3\zeta_3}{135\bar{\tau}^2} + \frac{16i\bar{\tau}\pi y\zeta_3^2}{9} - \frac{32y^2\zeta_3^2}{9}, \\
C \left[\begin{smallmatrix} 0 & 2 \\ 4 & 4 \end{smallmatrix} \right] &= -\frac{8\pi^4 y\zeta_3}{135} + \frac{64\bar{\tau}^2\pi^4 y\zeta_3}{405} + \frac{128i\bar{\tau}\pi^3 y^2\zeta_3}{405} + \frac{32i\bar{\tau}\pi y\zeta_3^2}{9} - \frac{32y^2\zeta_3^2}{9}, \\
C \left[\begin{smallmatrix} 1 & 2 \\ 4 & 4 \end{smallmatrix} \right] &= -\frac{4\pi^4 y\zeta_3}{135} + \frac{32\bar{\tau}^2\pi^4 y\zeta_3}{405} + \frac{16i\bar{\tau}\pi y\zeta_3^2}{9}, \\
C \left[\begin{smallmatrix} 4 & 2 \\ 6 & 4 \end{smallmatrix} \right] &= \frac{4i\bar{\tau}^3\pi^7\zeta_3}{2025} - \frac{8i\bar{\tau}\pi^7\zeta_3}{14175} + \frac{4i\bar{\tau}^5\pi^7\zeta_3}{2835} + \frac{8i\bar{\tau}^3\pi^5\zeta_5}{675} - \frac{4i\bar{\tau}^5\pi^5\zeta_5}{135} - \frac{8\bar{\tau}^2\pi^2\zeta_3\zeta_5}{15} + \frac{8\bar{\tau}^4\pi^2\zeta_3\zeta_5}{15}.
\end{aligned} \tag{2.54}$$

While the antisymmetry of $\beta_{(\alpha)}^{\text{sv}}$ in $(j_1, k_1) \leftrightarrow (j_2, k_2)$ clearly propagates to

$$C \left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix} \right] = -C \left[\begin{smallmatrix} j_2 & j_1 \\ k_2 & k_1 \end{smallmatrix} \right], \tag{2.55}$$

it is not immediately obvious from the definition (2.53) that the depth-one terms cancel. In fact, one may view the dropout of $\overline{\mathcal{E}_0(k, 0^p)}$ from (2.53) as a defining property of $\overline{\alpha[\cdots]}$.

2.3.4 Examples at depth two expressible via zeta values

For reduced multiple modular values at depth two, no analogue of the closed formula (2.49) is known. We begin with a few illustrative examples. In the (G_4, G_4) sector we have [50]

$$\begin{aligned}
\mathcal{M} \left[\begin{smallmatrix} 0 & 0 \\ 4 & 4 \end{smallmatrix} \right] &= -\frac{8\pi^2\zeta_3^2}{9}, & \mathcal{M} \left[\begin{smallmatrix} 0 & 1 \\ 4 & 4 \end{smallmatrix} \right] &= -\frac{16i\pi^5\zeta_3}{405} + \frac{10i\pi^3\zeta_5}{27}, & \mathcal{M} \left[\begin{smallmatrix} 1 & 0 \\ 4 & 4 \end{smallmatrix} \right] &= \frac{16i\pi^5\zeta_3}{405} - \frac{10i\pi^3\zeta_5}{27}, \\
\mathcal{M} \left[\begin{smallmatrix} 1 & 1 \\ 4 & 4 \end{smallmatrix} \right] &= 0, & \mathcal{M} \left[\begin{smallmatrix} 0 & 2 \\ 4 & 4 \end{smallmatrix} \right] &= \frac{8\pi^2\zeta_3^2}{9}, & \mathcal{M} \left[\begin{smallmatrix} 2 & 0 \\ 4 & 4 \end{smallmatrix} \right] &= \frac{8\pi^2\zeta_3^2}{9}, \\
\mathcal{M} \left[\begin{smallmatrix} 2 & 2 \\ 4 & 4 \end{smallmatrix} \right] &= -\frac{8\pi^2\zeta_3^2}{9}, & \mathcal{M} \left[\begin{smallmatrix} 1 & 2 \\ 4 & 4 \end{smallmatrix} \right] &= \frac{2i\pi^5\zeta_3}{135} - \frac{10i\pi^3\zeta_5}{27}, & \mathcal{M} \left[\begin{smallmatrix} 2 & 1 \\ 4 & 4 \end{smallmatrix} \right] &= -\frac{2i\pi^5\zeta_3}{135} + \frac{10i\pi^3\zeta_5}{27}.
\end{aligned} \tag{2.56}$$

We note that all rational multiples of π^8 present in the individual terms such as [50]

$$\mathfrak{m} \left[\begin{smallmatrix} 0 & 2 \\ 4 & 4 \end{smallmatrix} \right] = \frac{209\pi^8}{364500}, \quad \mathfrak{m} \left[\begin{smallmatrix} 2 & 0 \\ 4 & 4 \end{smallmatrix} \right] = -\frac{209\pi^8}{364500} + \frac{4\pi^2\zeta_3^2}{9}, \tag{2.57}$$

disappear in the combination (2.47).

⁸Here, we have chosen to replace τ by $\bar{\tau} + \frac{2i}{\pi}y$ for slightly more compact expressions.

More remarkable instances of such simplifications occur for higher weight, for instance in the reduced multiple modular values of the (G_4, G_6) sector [50]

$$\begin{aligned}
\mathcal{M}\left[\begin{smallmatrix} 0 & 0 \\ 4 & 6 \end{smallmatrix}\right] &= -\frac{8\pi^2\zeta_3\zeta_5}{15}, & \mathcal{M}\left[\begin{smallmatrix} 0 & 1 \\ 4 & 6 \end{smallmatrix}\right] &= -\frac{i\pi^7\zeta_3}{1575} + \frac{7i\pi^3\zeta_7}{90}, \\
\mathcal{M}\left[\begin{smallmatrix} 0 & 2 \\ 4 & 6 \end{smallmatrix}\right] &= 0, & \mathcal{M}\left[\begin{smallmatrix} 0 & 3 \\ 4 & 6 \end{smallmatrix}\right] &= \frac{i\pi^7\zeta_3}{4725} - \frac{4i\pi^5\zeta_5}{675}, \\
\mathcal{M}\left[\begin{smallmatrix} 0 & 4 \\ 4 & 6 \end{smallmatrix}\right] &= \frac{8\pi^2\zeta_3\zeta_5}{15}, & \mathcal{M}\left[\begin{smallmatrix} 1 & 0 \\ 4 & 6 \end{smallmatrix}\right] &= \frac{8i\pi^7\zeta_3}{14175} + \frac{2i\pi^5\zeta_5}{135} - \frac{7i\pi^3\zeta_7}{45}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 1 \\ 4 & 6 \end{smallmatrix}\right] &= 0, & \mathcal{M}\left[\begin{smallmatrix} 1 & 2 \\ 4 & 6 \end{smallmatrix}\right] &= -\frac{4i\pi^7\zeta_3}{8505} + \frac{2i\pi^5\zeta_5}{675}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 3 \\ 4 & 6 \end{smallmatrix}\right] &= 0, & \mathcal{M}\left[\begin{smallmatrix} 1 & 4 \\ 4 & 6 \end{smallmatrix}\right] &= \frac{8i\pi^7\zeta_3}{14175} - \frac{7i\pi^3\zeta_7}{45}, \\
\mathcal{M}\left[\begin{smallmatrix} 2 & 0 \\ 4 & 6 \end{smallmatrix}\right] &= \frac{8\pi^2\zeta_3\zeta_5}{15}, & \mathcal{M}\left[\begin{smallmatrix} 2 & 1 \\ 4 & 6 \end{smallmatrix}\right] &= \frac{2i\pi^7\zeta_3}{2835} - \frac{4i\pi^5\zeta_5}{675}, \\
\mathcal{M}\left[\begin{smallmatrix} 2 & 2 \\ 4 & 6 \end{smallmatrix}\right] &= 0, & \mathcal{M}\left[\begin{smallmatrix} 2 & 3 \\ 4 & 6 \end{smallmatrix}\right] &= -\frac{2i\pi^7\zeta_3}{14175} + \frac{7i\pi^3\zeta_7}{90}, \\
\mathcal{M}\left[\begin{smallmatrix} 2 & 4 \\ 4 & 6 \end{smallmatrix}\right] &= -\frac{8\pi^2\zeta_3\zeta_5}{15}.
\end{aligned} \tag{2.58}$$

Rational multiples of π^{10} and $\pi^2\zeta_{3,5}$ drop out from all the reduced counterparts $\mathcal{M}\left[\begin{smallmatrix} j_1 & j_2 \\ 4 & 6 \end{smallmatrix}\right]$ even though they appear in individual multiple modular values such as [50]

$$\begin{aligned}
\mathbf{m}\left[\begin{smallmatrix} 0 & 0 \\ 4 & 6 \end{smallmatrix}\right] &= -\frac{503\pi^{10}}{25515000} + \frac{4\pi^2}{75}\zeta_{3,5}, \\
\mathbf{m}\left[\begin{smallmatrix} 0 & 0 \\ 6 & 4 \end{smallmatrix}\right] &= \frac{503\pi^{10}}{25515000} - \frac{4\pi^2}{75}\zeta_{3,5} - \frac{4\pi^2}{15}\zeta_3\zeta_5.
\end{aligned} \tag{2.59}$$

We expect that more generally, the double zeta values ζ_{n_1, n_2} present in individual $\mathbf{m}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}\right]$ [61] will drop out in the combination to their reduced counterparts at arbitrary weight.

2.3.5 Examples at depth two involving L-values

Individual $\mathbf{m}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}\right]$ at weight $k_1+k_2 \geq 14$ involve certain “new numbers” [50] such as $c(\Delta; 12)$ and L-values of holomorphic cusp forms outside the critical strip. However, the reduced combinations (2.47) are conjectured to feature only single zeta values, L-values of cusp forms and powers of π . This can be checked from the $\mathcal{M}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}\right]$ provided in the ancillary file up to $k_1+k_2 \leq 28$ and the examples presented in this work.

The simplest examples of reduced multiple modular values involving non-critical L-values occur in the (G_4, G_{10}) and (G_6, G_8) sectors [50],⁹

$$\begin{aligned}
\mathcal{M}\left[\begin{smallmatrix} 0 & 1 \\ 4 & 10 \end{smallmatrix}\right] &= -\frac{43i\pi^{11}\zeta_3}{25259850} + \frac{11i\pi^3\zeta_{11}}{540} + \frac{256i\pi^{13}\Lambda(\Delta_{12}, 12)}{1913625}, \\
\mathcal{M}\left[\begin{smallmatrix} 0 & 3 \\ 4 & 10 \end{smallmatrix}\right] &= \frac{17i\pi^{11}\zeta_3}{265228425} - \frac{i\pi^5\zeta_9}{17010} - \frac{16i\pi^{13}\Lambda(\Delta_{12}, 12)}{229635}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 0 \\ 4 & 10 \end{smallmatrix}\right] &= \frac{8i\pi^{11}\zeta_3}{2525985} + \frac{2i\pi^5\zeta_9}{243} - \frac{11i\pi^3\zeta_{11}}{135} + \frac{256i\pi^{13}\Lambda(\Delta_{12}, 12)}{1913625}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 2 \\ 4 & 10 \end{smallmatrix}\right] &= -\frac{4i\pi^{11}\zeta_3}{10609137} + \frac{i\pi^5\zeta_9}{5670} - \frac{16i\pi^{13}\Lambda(\Delta_{12}, 12)}{229635}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 4 \\ 4 & 10 \end{smallmatrix}\right] &= -\frac{4i\pi^{11}\zeta_3}{37889775} + \frac{64i\pi^{13}\Lambda(\Delta_{12}, 12)}{1148175},
\end{aligned} \tag{2.60}$$

as well as

$$\begin{aligned}
\mathcal{M}\left[\begin{smallmatrix} 0 & 1 \\ 6 & 8 \end{smallmatrix}\right] &= -\frac{4i\pi^9\zeta_5}{297675} + \frac{22i\pi^3\zeta_{11}}{1575} - \frac{256i\pi^{13}\Lambda(\Delta_{12}, 12)}{826875}, \\
\mathcal{M}\left[\begin{smallmatrix} 0 & 3 \\ 6 & 8 \end{smallmatrix}\right] &= -\frac{2i\pi^9\zeta_5}{496125} + \frac{i\pi^5\zeta_9}{2250} + \frac{16i\pi^{13}\Lambda(\Delta_{12}, 12)}{99225}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 0 \\ 6 & 8 \end{smallmatrix}\right] &= \frac{i\pi^7\zeta_7}{4725} - \frac{11i\pi^3\zeta_{11}}{525} - \frac{256i\pi^{13}\Lambda(\Delta_{12}, 12)}{826875}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 2 \\ 6 & 8 \end{smallmatrix}\right] &= \frac{4i\pi^9\zeta_5}{2480625} - \frac{i\pi^5\zeta_9}{2250} + \frac{16i\pi^{13}\Lambda(\Delta_{12}, 12)}{99225}, \\
\mathcal{M}\left[\begin{smallmatrix} 2 & 1 \\ 6 & 8 \end{smallmatrix}\right] &= -\frac{2i\pi^9\zeta_5}{496125} + \frac{i\pi^5\zeta_9}{1350} + \frac{16i\pi^{13}\Lambda(\Delta_{12}, 12)}{99225}, \\
\mathcal{M}\left[\begin{smallmatrix} 2 & 3 \\ 6 & 8 \end{smallmatrix}\right] &= \frac{i\pi^9\zeta_5}{165375} - \frac{4i\pi^7\zeta_7}{99225} - \frac{64i\pi^{13}\Lambda(\Delta_{12}, 12)}{496125}.
\end{aligned} \tag{2.61}$$

All other cases with $(k_1, k_2) = (4, 10)$ or $(6, 8)$ are determined from (2.47), the depth-one results (2.25) and the reflection properties (2.45). Starting from $k_1+k_2 = 16$, we also find L-values $\Lambda(\Delta_{2s}, t)$ at odd t such as

$$\begin{aligned}
\mathcal{M}\left[\begin{smallmatrix} 1 & 6 \\ 4 & 12 \end{smallmatrix}\right] &= -\frac{2764i\pi^{13}\zeta_3}{1005657778125} + \frac{64i\pi^{14}\Lambda(\Delta_{12}, 13)}{245581875}, \\
\mathcal{M}\left[\begin{smallmatrix} 3 & 4 \\ 6 & 10 \end{smallmatrix}\right] &= -\frac{2i\pi^{11}\zeta_5}{21049875} + \frac{i\pi^7\zeta_9}{297675} + \frac{8i\pi^{14}\Lambda(\Delta_{12}, 13)}{2679075}, \\
\mathcal{M}\left[\begin{smallmatrix} 2 & 5 \\ 8 & 8 \end{smallmatrix}\right] &= -\frac{2i\pi^9\zeta_7}{496125} + \frac{4i\pi^7\zeta_9}{83349} + \frac{32i\pi^{14}\Lambda(\Delta_{12}, 13)}{3472875}, \\
\mathcal{M}\left[\begin{smallmatrix} 1 & 6 \\ 10 & 10 \end{smallmatrix}\right] &= \frac{i\pi^{11}\zeta_9}{5893965} - \frac{i\pi^9\zeta_{11}}{612360} - \frac{4i\pi^{16}\Lambda(\Delta_{12}, 15)}{104483925} - \frac{8i\pi^{18}\Lambda(\Delta_{16}, 17)}{1316497455}, \\
\mathcal{M}\left[\begin{smallmatrix} 0 & 1 \\ 4 & 16 \end{smallmatrix}\right] &= -\frac{31034i\pi^{17}\zeta_3}{51288546684375} + \frac{680i\pi^3\zeta_{17}}{97659} - \frac{256i\pi^{18}\Lambda(\Delta_{16}, 17)}{138175277625} - \frac{512i\pi^{19}\Lambda(\Delta_{18}, 18)}{28733079375}.
\end{aligned} \tag{2.62}$$

In the ancillary file accompanying the arXiv submission of this work and Part I, we present the complete list of reduced multiple modular values at depth two up to $k_1+k_2 = 28$. The values

⁹We are indebted to Francis Brown for correspondence on his work [62] and making many explicit expressions available to us, such as (2.60), (2.61) and the first three lines of (2.62).

there were fixed by knowing on which numbers to expand the reduced multiple modular value [50] and fitting the rational coefficients via numerical evaluation.¹⁰ Since we assign transcendental weight $m+1$ to $\Lambda(\Delta_{2s}, 2s+m)$, the explicit expressions are compatible with the transcendentality k_1+k_2 of $\mathcal{M}\begin{bmatrix} j_1 & j_2 \\ k_1 & k_2 \end{bmatrix}$.

3 Modular properties of solutions to the Laplace equations

The problem that motivates this work is to find modular invariant solutions $F_{m,k}^{\pm(s)}$ to the Laplace equations (1.1). In Part I, we showed how a leading-depth solution could be constructed in terms of the β^{sv} and how to complete it by including lower-depth β^{sv} terms. We already pointed out in Part I that the modular invariance of the resulting function $\check{F}_{m,k}^{\pm(s)}$ is not guaranteed by the construction that was only tailored to produce an exact solution of (1.1) in terms of various β^{sv} . Since the β^{sv} have the more involved modular properties presented above, this does not necessarily entail modular invariance of $\check{F}_{m,k}^{\pm(s)}$.

As we argued, failure of modular invariance can and will arise whenever the space of $F_{m,k}^{\pm(s)}$ is larger than the space of modular graph forms constructed from β^{sv} at depth ≤ 2 . The explicit counting done in section 3.6 of Part I showed that this can happen at Laplace eigenvalue $s(s-1)$ whenever there are holomorphic cusp forms at modular weight $2s$. This is not surprising since the generating series of MGFs [30] only contains combinations of β^{sv} that are compatible with the relations in Tsunogai's derivation algebra, see section 4 for a more detailed discussion of the corresponding 'dropouts' from the β^{sv} . As the relations in the derivation algebra are triggered by holomorphic cusp forms [49] we have a consistent picture that iterated integrals of such cusp forms should arise. They also feature naturally in the space of real-analytic modular functions studied in [37, 38, 47, 52, 53].

For every holomorphic cusp form Δ_{2s} of modular weight $2s$, we have constructed even and odd homogeneous solutions $H_{\Delta_{2s}}^{\pm}$ to the Laplace equation in (2.29) and we have also shown that they are not modular invariant, see (2.39). Therefore, if the combination $\check{F}_{m,k}^{\pm(s)}$ of β^{sv} is not modular invariant but solves the correct inhomogeneous Laplace equation, we can consider

$$F_{m,k}^{\pm(s)}(\tau) = \check{F}_{m,k}^{\pm(s)}(\tau) + \sum_{\Delta_{2s} \in \mathcal{S}_{2s}} a_{\Delta_{2s}, m, k}^{\pm} H_{\Delta_{2s}}^{\pm}(\tau), \quad (3.1)$$

where the sum runs over the space \mathcal{S}_{2s} of holomorphic cusp forms of weight $2s$. Then we can ask whether a suitable choice of constants $a_{\Delta_{2s}, m, k}^{\pm}$ renders this new solution $F_{m,k}^{\pm(s)}$ to the same Laplace equation modular invariant.

To answer this question we have to determine the modular transformation of $\check{F}_{m,k}^{\pm(s)}$. As this is a combination of β^{sv} , potentially multiplied by powers of y , we have to use the S-modular transformation of the β^{sv} discussed in section 2.3. As is evident from (2.50) and (2.51), the modular transformation generates special combinations of multiple modular values and additional

¹⁰Numerical approximations of the L-values can be obtained efficiently using PARI/GP [63, 64]. In the ancillary file, we have collected the numerical values of relevance to this work, meaning up to modular weight 26 and the first integer values outside critical strip.

depth-zero terms $C[\dots]$, which can be derived from the relevant $\overline{\alpha[\dots]}$ appearing in (2.4b). Ultimately, we obtain the explicit S-modular transformation of $\check{F}_{m,k}^{\pm(s)}$ in terms of these multiple modular values. In order for a failure of modularity of $\check{F}_{m,k}^{\pm(s)}$ to be cancelled by that of $H_{\Delta_{2s}}^{\pm}$, one must obtain very specific combinations of multiple modular values that ultimately are proportional to the same polynomial in τ and $\bar{\tau}$ given in (2.39). This is true in all examples and should follow from the general analysis in [35, 37]. We exemplify the mechanism for a variety of weights and Laplace eigenvalues.

The occurrence of these extra terms $H_{\Delta_{2s}}^{\pm}$ was also argued for on general grounds from the Cauchy–Riemann equation in Part I. We shall initially focus on the case when \mathcal{S}_{2s} is one-dimensional and defer more general cases to sections 3.3 and 3.4. We also note that the presence of $H_{\Delta_{2s}}^{\pm}$ in the modular-invariant solution $F_{m,k}^{\pm(s)}$ of the Laplace equation (1.1) implies that, inspecting Fourier by Fourier mode, there are non-zero coefficients for the homogeneous solutions (provided by Bessel functions) to the Laplace equations, see (2.32). This is in contrast to what was observed in examples in an $SL(2, \mathbb{Z})$ U-duality context in [65–67].

3.1 Examples involving the Ramanujan cusp form

The cusp form of lowest weight is the Ramanujan cusp form $\Delta_{12} = \sum_{n=1}^{\infty} \tau(n)q^n$. Since it has holomorphic modular weight 12, its iterated integrals can arise as the modular completion of $\check{F}_{m,k}^{\pm(s)}$ at eigenvalue $s = 6$. According to the spectrum (1.4) this happens first for $m+k = 7$ in the odd sector and for $m+k = 8$ in the even sector.

3.1.1 Odd functions for $(m, k) = (2, 5)$ and $(m, k) = (3, 4)$

As presented in Part I, we can carry out the procedure for obtaining a solution to the Laplace equations (1.1), and the simplest odd functions where modular invariance at the level of the β^{sv} breaks down are given by

$$\begin{aligned} \check{F}_{2,5}^{-(6)} &= -1890\beta^{\text{sv}}\left[\begin{smallmatrix} 1 & 4 \\ 4 & 10 \end{smallmatrix}\right] - 1512\beta^{\text{sv}}\left[\begin{smallmatrix} 2 & 3 \\ 4 & 10 \end{smallmatrix}\right] + 1890\beta^{\text{sv}}\left[\begin{smallmatrix} 4 & 1 \\ 10 & 4 \end{smallmatrix}\right] + 1512\beta^{\text{sv}}\left[\begin{smallmatrix} 5 & 0 \\ 10 & 4 \end{smallmatrix}\right] + 1008\zeta_3\beta^{\text{sv}}\left[\begin{smallmatrix} 3 \\ 10 \end{smallmatrix}\right] \\ &\quad + \frac{315\zeta_3}{y}\beta^{\text{sv}}\left[\begin{smallmatrix} 4 \\ 10 \end{smallmatrix}\right] - \frac{21\zeta_9}{4y^3}\beta^{\text{sv}}\left[\begin{smallmatrix} 0 \\ 4 \end{smallmatrix}\right] - \frac{105\zeta_9}{64y^4}\beta^{\text{sv}}\left[\begin{smallmatrix} 1 \\ 4 \end{smallmatrix}\right] + \frac{7\zeta_9}{1920y^2}, \\ \check{F}_{3,4}^{-(6)} &= -2100\beta^{\text{sv}}\left[\begin{smallmatrix} 2 & 3 \\ 6 & 8 \end{smallmatrix}\right] - 2100\beta^{\text{sv}}\left[\begin{smallmatrix} 3 & 2 \\ 6 & 8 \end{smallmatrix}\right] + 2100\beta^{\text{sv}}\left[\begin{smallmatrix} 3 & 2 \\ 8 & 6 \end{smallmatrix}\right] - 210\beta^{\text{sv}}\left[\begin{smallmatrix} 4 & 1 \\ 6 & 8 \end{smallmatrix}\right] + 2100\beta^{\text{sv}}\left[\begin{smallmatrix} 4 & 1 \\ 8 & 6 \end{smallmatrix}\right] \\ &\quad + 210\beta^{\text{sv}}\left[\begin{smallmatrix} 5 & 0 \\ 8 & 6 \end{smallmatrix}\right] + 84\zeta_5\beta^{\text{sv}}\left[\begin{smallmatrix} 1 \\ 8 \end{smallmatrix}\right] + \frac{210\zeta_5}{y}\beta^{\text{sv}}\left[\begin{smallmatrix} 2 \\ 8 \end{smallmatrix}\right] + \frac{105\zeta_5}{2y^2}\beta^{\text{sv}}\left[\begin{smallmatrix} 3 \\ 8 \end{smallmatrix}\right] - \frac{15\zeta_7}{y}\beta^{\text{sv}}\left[\begin{smallmatrix} 0 \\ 6 \end{smallmatrix}\right] \\ &\quad - \frac{75\zeta_7}{2y^2}\beta^{\text{sv}}\left[\begin{smallmatrix} 1 \\ 6 \end{smallmatrix}\right] - \frac{74\zeta_7}{8y^3}\beta^{\text{sv}}\left[\begin{smallmatrix} 2 \\ 6 \end{smallmatrix}\right] - \frac{\zeta_7}{15120}. \end{aligned} \tag{3.2}$$

The modular transformations of these expressions can be obtained by the methods of section 2.3 where we also use the knowledge of the $\overline{\alpha[\dots]}$ whose values were determined in Part I, building

on [30]. The modular transformation (2.33) of the functions above yields

$$\begin{aligned}\delta_S \check{F}_{2,5}^{-(6)}(\tau) &= \left(\frac{1701i}{8\pi^7} \mathcal{M} \begin{bmatrix} 0 & 1 \\ 4 & 10 \end{bmatrix} + \frac{1701i}{32\pi^7} \mathcal{M} \begin{bmatrix} 1 & 0 \\ 4 & 10 \end{bmatrix} - \frac{23\pi^4 \zeta_3}{118800} + \frac{7\zeta_9}{16\pi^2} \right) \frac{\tau + \bar{\tau}}{(\tau - \bar{\tau})^5} + \dots, \\ \delta_S \check{F}_{3,4}^{-(6)}(\tau) &= \left(\frac{6615i}{32\pi^7} \mathcal{M} \begin{bmatrix} 0 & 1 \\ 6 & 8 \end{bmatrix} + \frac{2205i}{16\pi^7} \mathcal{M} \begin{bmatrix} 1 & 0 \\ 6 & 8 \end{bmatrix} - \frac{\pi^2 \zeta_5}{360} + \frac{7\zeta_7}{240} \right) \frac{\tau + \bar{\tau}}{(\tau - \bar{\tau})^5} + \dots,\end{aligned}\quad (3.3)$$

where the dots denote similar terms that consist of multiple modular values and polynomials in τ and $\bar{\tau}$ over the same denominator $(\tau - \bar{\tau})^5$. Even without further knowledge of the exact multiple modular values one can check numerically, using (2.24), that neither of the above S-transformations vanishes. Hence, we are indeed in one of the cases where the combination $\check{F}_{2,5}^{-(6)}$ of β^{sv} that solves the Laplace equation fails to be modular invariant.

Fortunately, further knowledge of the multiple modular values is available through the beautiful work of Brown [35], and the examples relevant to (3.3) can be found in (2.60) and (2.61). We emphasise that the multiple modular values are not built out of multiple zeta values alone but also contain L-values of cusp forms. In general the individual multiple modular values also contain other ‘new numbers’ such as the object $c(\Delta; 12)$ in [50], but these do not enter in the examples above and cancel in all reduced multiple modular values (2.47) that we have encountered.

Using these and similar results one can show that

$$\begin{aligned}\delta_S \check{F}_{2,5}^{-(6)}(\tau) &= \frac{i\pi^{11} \Lambda(\Delta_{12}, 12)}{900y^5} p_{12}^-(\tau, \bar{\tau}) \\ \delta_S \check{F}_{3,4}^{-(6)}(\tau) &= -\frac{i\pi^{11} \Lambda(\Delta_{12}, 12)}{300y^5} p_{12}^-(\tau, \bar{\tau})\end{aligned}\quad (3.4)$$

with the same polynomial in $\tau, \bar{\tau}$ in both cases

$$\begin{aligned}p_{12}^-(\tau, \bar{\tau}) &= (\tau + \bar{\tau}) \left\{ 1 + \tau^4 \bar{\tau}^4 - \frac{25}{24} (\tau^2 + \bar{\tau}^2) (1 + \tau^2 \bar{\tau}^2) + \frac{1}{12} (\tau^4 + \bar{\tau}^4) \right. \\ &\quad \left. - \frac{25}{6} \tau \bar{\tau} (1 + \tau^2 \bar{\tau}^2) + 2\tau \bar{\tau} (\tau^2 + \bar{\tau}^2) + \frac{19}{3} \tau^2 \bar{\tau}^2 \right\}.\end{aligned}\quad (3.5)$$

These cocycles turn out to be proportional to those of $H_{\Delta_{12}}^-$ in (2.43b),

$$\begin{aligned}\check{F}_{2,5}^{-(6)}(\tau) - \check{F}_{2,5}^{-(6)}\left(-\frac{1}{\tau}\right) &= \frac{1}{75} \frac{\Lambda(\Delta_{12}, 12)}{\Lambda(\Delta_{12}, 10)} \left(H_{\Delta_{12}}^-(\tau) - H_{\Delta_{12}}^-\left(-\frac{1}{\tau}\right) \right), \\ \check{F}_{3,4}^{-(6)}(\tau) - \check{F}_{3,4}^{-(6)}\left(-\frac{1}{\tau}\right) &= -\frac{1}{25} \frac{\Lambda(\Delta_{12}, 12)}{\Lambda(\Delta_{12}, 10)} \left(H_{\Delta_{12}}^-(\tau) - H_{\Delta_{12}}^-\left(-\frac{1}{\tau}\right) \right).\end{aligned}\quad (3.6)$$

We can therefore form the linear combinations

$$\begin{aligned}F_{2,5}^{-(6)} &= \check{F}_{2,5}^{-(6)} - \frac{1}{75} \frac{\Lambda(\Delta_{12}, 12)}{\Lambda(\Delta_{12}, 10)} H_{\Delta_{12}}^-, \\ F_{3,4}^{-(6)} &= \check{F}_{3,4}^{-(6)} + \frac{1}{25} \frac{\Lambda(\Delta_{12}, 12)}{\Lambda(\Delta_{12}, 10)} H_{\Delta_{12}}^-, \end{aligned}\quad (3.7)$$

that are modular invariant thanks to (3.6) and solve the Laplace equation (1.1b). The ratio of L-values appearing here might appear a bit surprising since in all previous expressions of this work and Part I, (multiple) zeta values only appear in the numerator. The reason for the ratio showing up here lies in our normalisation of the homogeneous solution in (2.29) as a simple iterated integral. This definition leads to the explicit appearance of an L-value in the S-cocycle, see (2.43b), which needs to be cancelled by the denominator in (3.7). Choosing a different normalisation for $H_{\Delta_{2s}}^{\pm}$ one could turn the quotient in (3.7) into a multiplication by an L-value only.

We note that the combination

$$3F_{2,5}^{-(6)} + F_{3,4}^{-(6)} = 3\check{F}_{2,5}^{-(6)} + \check{F}_{3,4}^{-(6)}, \quad (3.8)$$

is an eigenfunction of the Laplacian with eigenvalue 30 which is perfectly modular invariant on its own without the need of adding any iterated integral of Δ_{12} . This is one of the examples of modular objects analysed in Part I that is expressible in terms of β^{sv} , y and odd zeta values.

3.1.2 Even functions for $(m, k) = (2, 6)$, $(m, k) = (3, 5)$ and $(m, k) = (4, 4)$

In the even sector, the first occurrence of the eigenvalue $s = 6$ is for $F_{2,6}^{+(6)}$, $F_{3,5}^{+(6)}$ and $F_{4,4}^{+(6)}$. The β^{sv} -representations of the associated $\check{F}_{m,8-m}^{+(6)}$ take a form similar to their odd counterparts in (3.2) and can be found in section 4.5 of Part I. Performing the S-transformations on the combinations constructed only out of the β^{sv} we find

$$\begin{aligned} \delta_S \check{F}_{2,6}^{+(6)}(\tau) &= \left(\frac{693}{64\pi^8} (\mathcal{M}[\begin{smallmatrix} 0 & 1 \\ 4 & 12 \end{smallmatrix}]] - \mathcal{M}[\begin{smallmatrix} 1 & 0 \\ 4 & 12 \end{smallmatrix}]] \right) + \frac{823i\pi^5\zeta_3}{202702500} + \frac{7i\zeta_{11}}{96\pi^3} - \frac{9555i\zeta_{13}}{11056\pi^5} \left) \frac{1}{(\tau - \bar{\tau})^5} + \dots \\ &= -\frac{\pi^{11}\Lambda(\Delta_{12}, 13)}{518250y^5} p_{12}^+(\tau, \bar{\tau}), \\ \delta_S \check{F}_{3,5}^{+(6)}(\tau) &= \left(\frac{945}{64\pi^8} (\mathcal{M}[\begin{smallmatrix} 0 & 1 \\ 6 & 10 \end{smallmatrix}]] - \mathcal{M}[\begin{smallmatrix} 1 & 0 \\ 6 & 10 \end{smallmatrix}]] \right) + \frac{i\pi^3\zeta_5}{86400} + \frac{7i\zeta_9}{2880\pi} - \frac{63063i\zeta_{13}}{176896\pi^5} \left) \frac{1}{(\tau - \bar{\tau})^5} + \dots \\ &= \frac{\pi^{11}\Lambda(\Delta_{12}, 13)}{132672y^5} p_{12}^+(\tau, \bar{\tau}), \\ \delta_S \check{F}_{4,4}^{+(6)}(\tau) &= \left(\frac{1029}{32\pi^8} \mathcal{M}[\begin{smallmatrix} 0 & 1 \\ 8 & 8 \end{smallmatrix}]] + \frac{i\pi\zeta_7}{3240} - \frac{5005i\zeta_{13}}{16584\pi^5} \right) \frac{1}{(\tau - \bar{\tau})^5} + \dots \\ &= -\frac{7\pi^{11}\Lambda(\Delta_{12}, 13)}{621900y^5} p_{12}^+(\tau, \bar{\tau}). \end{aligned} \quad (3.9)$$

In the second steps, we have inserted the relevant expressions for the multiple modular values and thereby arrived at the polynomial

$$\begin{aligned} p_{12}^+(\tau, \bar{\tau}) &= (1 - \tau\bar{\tau}) \left\{ 1 + \tau^4\bar{\tau}^4 - \frac{691}{162}(\tau^2 + \bar{\tau}^2)(1 + \tau^2\bar{\tau}^2) + \frac{691}{504}(\tau^4 + \bar{\tau}^4) \right. \\ &\quad \left. - \frac{3131}{324}\tau\bar{\tau}(1 + \tau^2\bar{\tau}^2) + \frac{21421}{2268}\tau\bar{\tau}(\tau^2 + \bar{\tau}^2) + \frac{40273}{2268}\tau^2\bar{\tau}^2 \right\} \end{aligned} \quad (3.10)$$

which determines the ellipses in the first steps of (3.9). Since the corresponding S-cocycle of the homogeneous solution $H_{\Delta_{12}}^+$ in (2.43a) is proportional to the same polynomial in (3.10), the following combinations are modular invariant:

$$\begin{aligned} F_{2,6}^{+(6)} &= \check{F}_{2,6}^{+(6)} + \frac{2}{17275} \frac{\Lambda(\Delta_{12}, 13)}{\Lambda(\Delta_{12}, 11)} H_{\Delta_{12}}^+, \\ F_{3,5}^{+(6)} &= \check{F}_{3,5}^{+(6)} - \frac{5}{11056} \frac{\Lambda(\Delta_{12}, 13)}{\Lambda(\Delta_{12}, 11)} H_{\Delta_{12}}^+, \\ F_{4,4}^{+(6)} &= \check{F}_{4,4}^{+(6)} + \frac{7}{10365} \frac{\Lambda(\Delta_{12}, 13)}{\Lambda(\Delta_{12}, 11)} H_{\Delta_{12}}^+. \end{aligned} \quad (3.11)$$

From this we also see that the following combinations are modular invariant without the inclusions of an iterated integral of a holomorphic cusp form,

$$\begin{aligned} F_{2,6}^{+(6)} - \frac{6}{35} F_{4,4}^{+(6)} &= \check{F}_{2,6}^{+(6)} - \frac{6}{35} \check{F}_{4,4}^{+(6)}, \\ F_{3,5}^{+(6)} + \frac{75}{112} F_{4,4}^{+(6)} &= \check{F}_{3,5}^{+(6)} + \frac{75}{112} \check{F}_{4,4}^{+(6)}, \end{aligned} \quad (3.12)$$

as already discussed in section 4.5.2 of Part I. There is a two-dimensional subspace of the three modular invariants $\{F_{2,6}^{+(6)}, F_{3,5}^{+(6)}, F_{4,4}^{+(6)}\}$ that does not require a cusp form – in agreement with our counting of MGFs in section 3.6 of Part I.

3.2 Examples with cusp forms of higher weight

We have performed the same analysis as in the previous section for all $F_{m,k}^{\pm(s)}$ with $m+k \leq 14$. The combinations $\check{F}_{m,k}^{\pm(s)}$ of β^{sv} require additional cusp forms for their modular invariant completion whenever s is half the weight of a holomorphic cusp form, i.e. $s \in \{6, 8, 9, \dots\}$. We only list those cases here, a full list of the $\check{F}_{m,k}^{\pm(s)}$ with $m+k \leq 14$ in terms of β^{sv} can be found in the ancillary file along with their modular completions.

Besides the examples of the previous section, the Ramanujan cusp form occurs in the following modular invariant functions at transcendental weight $m+k = 9, 10$,

$$\begin{aligned} F_{2,7}^{-(6)} &= \check{F}_{2,7}^{-(6)} + \frac{1}{51825} \frac{\Lambda(\Delta_{12}, 14)}{\Lambda(\Delta_{12}, 10)} H_{\Delta_{12}}^-, \\ F_{3,6}^{-(6)} &= \check{F}_{3,6}^{-(6)} - \frac{1}{4500} \frac{\Lambda(\Delta_{12}, 14)}{\Lambda(\Delta_{12}, 10)} H_{\Delta_{12}}^-, \\ F_{4,5}^{-(6)} &= \check{F}_{4,5}^{-(6)} + \frac{1}{1800} \frac{\Lambda(\Delta_{12}, 14)}{\Lambda(\Delta_{12}, 10)} H_{\Delta_{12}}^-, \\ F_{3,7}^{+(6)} &= \check{F}_{3,7}^{+(6)} + \frac{1}{725550} \frac{\Lambda(\Delta_{12}, 15)}{\Lambda(\Delta_{12}, 11)} H_{\Delta_{12}}^+, \\ F_{4,6}^{+(6)} &= \check{F}_{4,6}^{+(6)} - \frac{1}{232176} \frac{\Lambda(\Delta_{12}, 15)}{\Lambda(\Delta_{12}, 11)} H_{\Delta_{12}}^+, \\ F_{5,5}^{+(6)} &= \check{F}_{5,5}^{+(6)} + \frac{1}{165840} \frac{\Lambda(\Delta_{12}, 15)}{\Lambda(\Delta_{12}, 11)} H_{\Delta_{12}}^+. \end{aligned} \quad (3.13)$$

The Hecke normalised cusp form of weight 16 arises in (also see appendix B.1)

$$\begin{aligned}
F_{2,7}^{-(8)} &= \check{F}_{2,7}^{-(8)} - \frac{13}{7350} \frac{\Lambda(\Delta_{16}, 16)}{\Lambda(\Delta_{16}, 14)} H_{\Delta_{16}}^{-}, \\
F_{3,6}^{-(8)} &= \check{F}_{3,6}^{-(8)} + \frac{13}{2100} \frac{\Lambda(\Delta_{16}, 16)}{\Lambda(\Delta_{16}, 14)} H_{\Delta_{16}}^{-}, \\
F_{4,5}^{-(8)} &= \check{F}_{4,5}^{-(8)} - \frac{143}{14700} \frac{\Lambda(\Delta_{16}, 16)}{\Lambda(\Delta_{16}, 14)} H_{\Delta_{16}}^{-}, \\
F_{2,8}^{+(8)} &= \check{F}_{2,8}^{+(8)} + \frac{39}{3544660} \frac{\Lambda(\Delta_{16}, 17)}{\Lambda(\Delta_{16}, 15)} H_{\Delta_{16}}^{+}, \\
F_{3,7}^{+(8)} &= \check{F}_{3,7}^{+(8)} - \frac{91}{2083392} \frac{\Lambda(\Delta_{16}, 17)}{\Lambda(\Delta_{16}, 15)} H_{\Delta_{16}}^{+}, \\
F_{4,6}^{+(8)} &= \check{F}_{4,6}^{+(8)} + \frac{1001}{13021200} \frac{\Lambda(\Delta_{16}, 17)}{\Lambda(\Delta_{16}, 15)} H_{\Delta_{16}}^{+}, \\
F_{5,5}^{+(8)} &= \check{F}_{5,5}^{+(8)} - \frac{143}{1620416} \frac{\Lambda(\Delta_{16}, 17)}{\Lambda(\Delta_{16}, 15)} H_{\Delta_{16}}^{+}.
\end{aligned} \tag{3.14}$$

The Hecke normalised cusp form of weight 18 arises in (also see appendix B.2)

$$\begin{aligned}
F_{2,8}^{-(9)} &= \check{F}_{2,8}^{-(9)} + \frac{1}{1260} \frac{\Lambda(\Delta_{18}, 18)}{\Lambda(\Delta_{18}, 16)} H_{\Delta_{18}}^{-}, \\
F_{3,7}^{-(9)} &= \check{F}_{3,7}^{-(9)} - \frac{5}{2016} \frac{\Lambda(\Delta_{18}, 18)}{\Lambda(\Delta_{18}, 16)} H_{\Delta_{18}}^{-}, \\
F_{4,6}^{-(9)} &= \check{F}_{4,6}^{-(9)} + \frac{13}{5040} \frac{\Lambda(\Delta_{18}, 18)}{\Lambda(\Delta_{18}, 16)} H_{\Delta_{18}}^{-}.
\end{aligned} \tag{3.15}$$

Just as discussed in the previous examples, it is possible to find suitable rational linear combinations of these objects to produce modular invariant functions for which the homogeneous solutions $H_{\Delta_{2s}}^{\pm}$ cancel out, thus properly living in the realm of MGFs.

3.3 An example involving the two weight 24 cusp forms

As explained in Part I [1], as we increase the total transcendental weight $w = m+k$ we encounter higher and higher eigenvalues $s \leq k+m-1$ in the spectrum, see (1.4). This in turn means that the obstructions to finding modular solutions to the Laplace systems (1.1) are related to iterated integrals (2.29) of cusp forms Δ_{2s} of higher and higher modular weight $2s$. Denoting by \mathcal{S}_{2s} the vector space of holomorphic cusp forms for $\mathrm{SL}(2, \mathbb{Z})$ with even integer modular weight $2s$, we have the classic result [68]

$$\dim \mathcal{S}_{2s} = \begin{cases} \lfloor \frac{2s}{12} \rfloor - 1 & 2s \equiv 2 \pmod{12}, \\ \lfloor \frac{2s}{12} \rfloor & \text{otherwise.} \end{cases} \tag{3.16}$$

Hence, we see that starting with $2s = 24$ we should in general expect the space of obstructions to have dimension greater than one since $\dim \mathcal{S}_{24} = 2$. From (1.4) we know that the first instance for which the eigenvalue $s = 12$ appears is for the odd sector and with transcendental weight

$w = m+k = 13$. We are then led to expect that a modular invariant solution to (1.1) in this sector must take the form

$$F_{m,k}^{-(12)} = \check{F}_{m,k}^{-(12)} + a_{\Delta_{24,i},m,k}^- H_{\Delta_{24,i}}^- + a_{\Delta_{24,ii},m,k}^- H_{\Delta_{24,ii}}^-, \quad (3.17)$$

with $m+k = 13$ and $m, k \geq 2$, where we use a basis of Hecke eigenforms subject to (2.38) and denoted by $\Delta_{24,i}, \Delta_{24,ii}$.

Such a basis can be constructed by considering the linear combination $\alpha\Delta_{12}^2 + \beta\Delta_{12}G_{12}$, i.e. the most general holomorphic cusp form of weight $2s = 24$. Then, the real coefficients α, β for Hecke eigenforms are obtained by imposing the resulting Fourier coefficients to be multiplicative, i.e. $a(m)a(n) = a(m \cdot n)$ for m, n coprime: $\gcd(m, n) = 1$. This procedure constructs the two Hecke eigenforms¹¹

$$\Delta_{24,i}(\tau) = \left(\frac{324204}{691} - 12\sqrt{144169} \right) \Delta_{12}(\tau)^2 + \frac{G_{12}(\tau)}{2\zeta_{12}} \Delta_{12}(\tau), \quad (3.18a)$$

$$\Delta_{24,ii}(\tau) = \left(\frac{324204}{691} + 12\sqrt{144169} \right) \Delta_{12}(\tau)^2 + \frac{G_{12}(\tau)}{2\zeta_{12}} \Delta_{12}(\tau), \quad (3.18b)$$

whose Fourier coefficients lie in the number field $\mathbb{Q}(\sqrt{144169})$:

$$\Delta_{24,i}(\tau) = q^1 + (540 - 12\sqrt{144169})q^2 + (169740 + 576\sqrt{144169})q^3 + O(q^4), \quad (3.19a)$$

$$\Delta_{24,ii}(\tau) = q^1 + (540 + 12\sqrt{144169})q^2 + (169740 - 576\sqrt{144169})q^3 + O(q^4). \quad (3.19b)$$

The number field generated by the above Fourier coefficients has a non-trivial Galois automorphism $\sigma \in \text{Aut}_{\mathbb{Q}}(\mathbb{Q}(\sqrt{144169}))$, which acts as $\sigma : \sqrt{144169} \rightarrow -\sqrt{144169}$ and under which the two basis elements are exchanged, i.e. $\sigma(\Delta_{24,i}) = \Delta_{24,ii}$.

In the basis of Hecke eigenforms we have that all the even/odd completed L-values inside the critical strip are $\mathbb{Q}(\sqrt{144169})$ multiples of one another [60] and the S-cocycles for $\Delta_{24,i}$ and $\Delta_{24,ii}$ can be put in the form (2.39). The Galois automorphism exchanges the two cocycles, as well as the completed L-values.

Following the same types of arguments that led to (3.6), we see that in general the S-cocycle for $\check{F}_{m,k}^{-(12)}$ does not vanish and involves the completed L-values $\Lambda(\Delta_{24,i}, t), \Lambda(\Delta_{24,ii}, t)$ of the two

¹¹Alternative expressions in terms of the ring generators G_4, G_6 read

$$\begin{aligned} \Delta_{24,i}(\tau) = & \left(\frac{131}{248832} - \frac{\sqrt{144169}}{248832} \right) \left(\frac{G_4(\tau)}{2\zeta_4} \right)^6 - \left(\frac{13}{248832} + \frac{\sqrt{144169}}{248832} \right) \left(\frac{G_6(\tau)}{2\zeta_6} \right)^4 \\ & + \left(-\frac{59}{124416} + \frac{\sqrt{144169}}{124416} \right) \left(\frac{G_4(\tau)}{2\zeta_4} \right)^3 \left(\frac{G_6(\tau)}{2\zeta_6} \right)^2 \end{aligned}$$

and the analogous combination with $\sqrt{144169} \rightarrow -\sqrt{144169}$ for $\Delta_{24,ii}(\tau)$.

Hecke eigenforms through the reduced multiple modular values. For example, we find

$$\begin{aligned} \mathcal{M}\left[\begin{smallmatrix} 0 & 1 \\ 4 & 22 \end{smallmatrix}\right] &= -\frac{24438334i\pi^{23}\zeta_3}{77661373832214046875} + \frac{4016053i\pi^3\zeta_{23}}{1153592550} + \frac{4096i\pi^{24}\Lambda(\Delta_{22}, 23)}{7791418647627615} \\ &+ \left(\frac{39121664i\pi^{25}}{7428479236124821875} + \frac{1279322368i\pi^{25}\sqrt{144169}}{1070956422992879444896875} \right) \Lambda(\Delta_{24,i}, 24) \quad (3.20a) \\ &+ \left(\frac{39121664i\pi^{25}}{7428479236124821875} - \frac{1279322368i\pi^{25}\sqrt{144169}}{1070956422992879444896875} \right) \Lambda(\Delta_{24,ii}, 24), \end{aligned}$$

$$\begin{aligned} \mathcal{M}\left[\begin{smallmatrix} 1 & 0 \\ 4 & 22 \end{smallmatrix}\right] &= \frac{1242928i\pi^{23}\zeta_3}{847214987260516875} + \frac{2i\pi^5\zeta_{21}}{567} + \frac{4016053i\pi^3\zeta_{23}}{115359255} - \frac{8192i\pi^{24}\Lambda(\Delta_{22}, 23)}{1558283729525523} \\ &+ \left(\frac{39121664i\pi^{25}}{7428479236124821875} + \frac{1279322368i\pi^{25}\sqrt{144169}}{1070956422992879444896875} \right) \Lambda(\Delta_{24,i}, 24) \quad (3.20b) \\ &+ \left(\frac{39121664i\pi^{25}}{7428479236124821875} - \frac{1279322368i\pi^{25}\sqrt{144169}}{1070956422992879444896875} \right) \Lambda(\Delta_{24,ii}, 24). \end{aligned}$$

Using these and similar expressions for the other $\mathcal{M}\left[\begin{smallmatrix} j_1 & j_2 \\ k_1 & k_2 \end{smallmatrix}\right]$ with $k_1+k_2 = 26$, there is a unique choice of the constants $a_{\Delta_{24,i},m,k}^-, a_{\Delta_{24,ii},m,k}^-$ such that the combinations (3.17) are modular invariant. Explicitly,

$$\begin{aligned} a_{\Delta_{24,i},2,11}^- &= \frac{1}{7200} \left(-\frac{152819}{76230} - \frac{4997353\sqrt{144169}}{10990002870} \right) \frac{\Lambda(\Delta_{24,i}, 24)}{\Lambda(\Delta_{24,i}, 22)}, \\ a_{\Delta_{24,i},3,10}^- &= -\frac{19}{4320} \left(-\frac{18289}{87780} - \frac{554243\sqrt{144169}}{12655154820} \right) \frac{\Lambda(\Delta_{24,i}, 24)}{\Lambda(\Delta_{24,i}, 22)}, \\ a_{\Delta_{24,i},4,9}^- &= \frac{323}{7200} \left(-\frac{2117}{78540} - \frac{44479\sqrt{144169}}{11323033260} \right) \frac{\Lambda(\Delta_{24,i}, 24)}{\Lambda(\Delta_{24,i}, 22)}, \quad (3.21) \\ a_{\Delta_{24,i},5,8}^- &= -\frac{323}{1680} \left(-\frac{1}{220} + \frac{13\sqrt{144169}}{31717180} \right) \frac{\Lambda(\Delta_{24,i}, 24)}{\Lambda(\Delta_{24,i}, 22)}, \\ a_{\Delta_{24,i},6,7}^- &= \frac{4199}{10800} \left(-\frac{823}{660660} + \frac{82699\sqrt{144169}}{95246691540} \right) \frac{\Lambda(\Delta_{24,i}, 24)}{\Lambda(\Delta_{24,i}, 22)}, \end{aligned}$$

where we have split the result into three factors: The ratio of L-values carries transcendental weight one, the middle factor is valued in the number field $\mathbb{Q}(\sqrt{144169})$ and corresponds to the inverse of the Petersson–Haberland pairing [69] between two properly normalised polynomials associated with the cusp from Δ_{2s} . The first factor is a rational number multiplying the vector in the number field and we refer to [35] for why this splitting occurs. The constants $a_{\Delta_{24,ii},m,k}^-$ for the second cusp form can be directly obtained by the application of the Galois automorphism to $a_{\Delta_{24,i},m,k}^-$, i.e. $a_{\Delta_{24,ii},m,k}^- = \sigma(a_{\Delta_{24,i},m,k}^-)$ that acts on the number-field-valued middle factor by the Galois action and on the L-values by $\sigma(\Lambda(\Delta_{24,i}, t)) = \Lambda(\Delta_{24,ii}, t)$.

With the above values we obtain that the combinations (3.17) are then the unique modular invariant solutions to the Laplace system (1.1). Furthermore, we can easily check that, for

some particular rational linear combinations of $F_{m,k}^{-(12)}$, the iterated integrals of the cusp forms $H_{\Delta_{24,i}}^-(\tau)$ and $H_{\Delta_{24,ii}}^-(\tau)$ drop out, thus producing the modular objects discussed in Part I, for example:

$$\begin{aligned} \frac{1862}{103}F_{2,11}^{-(12)} - \frac{470}{103}F_{4,9}^{-(12)} + F_{6,7}^{-(12)} &= \frac{1862}{103}\check{F}_{2,11}^{-(12)} - \frac{470}{103}\check{F}_{4,9}^{-(12)} + \check{F}_{6,7}^{-(12)}, \\ \frac{171}{49}F_{3,10}^{-(12)} + \frac{165}{49}F_{4,9}^{-(12)} + F_{5,8}^{-(12)} &= \frac{171}{49}\check{F}_{3,10}^{-(12)} + \frac{165}{49}\check{F}_{4,9}^{-(12)} + \check{F}_{5,8}^{-(12)}, \\ -\frac{76}{11}F_{3,10}^{-(12)} - \frac{62}{11}F_{4,9}^{-(12)} + F_{6,7}^{-(12)} &= -\frac{76}{11}\check{F}_{3,10}^{-(12)} - \frac{62}{11}\check{F}_{4,9}^{-(12)} + \check{F}_{6,7}^{-(12)}. \end{aligned} \quad (3.22)$$

In section 4, we discuss the relation between linear combinations of this type and Tsunogai's derivation algebra.

The complete list of modular completions $\check{F}_{m,k}^{\pm(s)} \rightarrow F_{m,k}^{\pm(s)}$ with $m+k \leq 14$ can be found in the ancillary file.

3.4 Structure for general weight

For general transcendental weight $w = m+k$ and eigenvalue s in the spectrum (1.4), the modular invariant solutions $F_{m,k}^{\pm(s)}$ to the Laplace system (1.1) must take the form given in (3.1).

From their general form given in (2.39) we know that the cocycles $\delta_S H_{\Delta_{2s}}^{\pm}$ can be normalised so that we have factorised out the completed L-values $\Lambda(\Delta_{2s}, 2s-1)$ in the even case and $\Lambda(\Delta_{2s}, 2s-2)$ in the odd case, times a rational function in $\tau, \bar{\tau}$ with coefficients in $K_{\Delta_{2s}}$, the number field generated by the Fourier coefficients of Δ_{2s} .

To understand the generic structure of the coefficients $a_{\Delta_{2s},m,k}^{\pm}$ we can analyse in more depth their transcendental properties. Recalling here the expression (2.31) to rewrite $H_{\Delta_{2s}}^{\pm}$ in terms of iterated integrals

$$H_{\Delta_{2s}}^{\pm}(\tau) = -\frac{1}{2} \sum_{\ell=0}^{s-1} (4y)^{-\ell} \frac{\Gamma(s+\ell)}{\ell! \Gamma(s-\ell)} \left[\mathcal{E}_0(\Delta_{2s}, 0^{s-1+\ell}; \tau) \pm \text{c.c.} \right], \quad (3.23)$$

we notice that all these terms have uniform transcendental weight s , since in our convention the transcendental weight of any iterated integral $\mathcal{E}_0(\Delta_{2s}, 0^\ell; \tau)$ is $\ell+1$, and $y = \pi \text{Im } \tau$ has transcendental weight 1. Hence, we deduce that we can assign transcendental weight s to $H_{\Delta_{2s}}^{\pm}$. However, from (3.1) we see that the combination $a_{\Delta_{2s},m,k}^{\pm} H_{\Delta_{2s}}^{\pm}$ must have transcendental weight $w = m+k$ which assigns weight $m+k-s$ to $a_{\Delta_{2s},m,k}^{\pm}$. On these grounds, by the transcendental weight ℓ of $\Lambda(\Delta_{2s}, 2s+\ell-1)$ outside the critical strip $\ell \geq 1$, we are led to conclude that

$$a_{\Delta_{2s},m,k}^{\pm} = \begin{cases} q_{\Delta_{2s},m,k}^+ \kappa_{\Delta_{2s},m,k}^+ \frac{\Lambda(\Delta_{2s}, m+k+s-1)}{\Lambda(\Delta_{2s}, 2s-1)}, \\ q_{\Delta_{2s},m,k}^- \kappa_{\Delta_{2s},m,k}^- \frac{\Lambda(\Delta_{2s}, m+k+s-1)}{\Lambda(\Delta_{2s}, 2s-2)}, \end{cases} \quad (3.24)$$

where we have $q_{\Delta_{2s},m,k}^{\pm} \in \mathbb{Q}$, while $\kappa_{\Delta_{2s},m,k}^{\pm} \in K_{\Delta_{2s}}$ is given by the inverse of the Petersson-Haberland pairing between the two cocycles $\delta_S H_{\Delta_{2s}}^+(\tau)$ and $\delta_S H_{\Delta_{2s}}^-(\tau)$ properly normalised,

see [35]. Given that $m+k+s$ is even (odd) for even functions $F_{m,k}^{+(s)}$ (odd functions $F_{m,k}^{-(s)}$), the L-function $\Lambda(\Delta_{2s}, m+k+s-1)$ in the numerator of (3.24) is evaluated at odd integers in the modular invariant completion of $F_{m,k}^{+(s)}$ (at even integers in the case of $F_{m,k}^{-(s)}$).

Note that, as discussed previously, to determine the number field $K_{\Delta_{2s}}$, which contains the Fourier coefficient of $\Delta_{2s} \in \mathcal{S}_{2s}$, one has to diagonalise the Hecke operators (2.38) in \mathcal{S}_{2s} . The nature of this number field $K_{\Delta_{2s}}$ is clarified¹² by the *Maeda conjecture* [70], which states that the characteristic polynomials of the Hecke operator T_n are irreducible over \mathbb{Q} . In a certain sense the number field $K_{\Delta_{2s}}$ is “maximal” in that the Maeda conjecture suggests that the associated Galois group is the full symmetric group \mathfrak{S}_d with $d = \dim \mathcal{S}_{2s}$.

For example at weight $2s = 28$ with $d = \dim \mathcal{S}_{28} = 2$ we have $K_{\Delta_{28}} = \mathbb{Q}(\sqrt{18209})$, for $2s = 30$ we have once more $d = 2$ and now $K_{\Delta_{30}} = \mathbb{Q}(\sqrt{51349})$, while moving to weight $2s = 36$, the lowest weight for which $d = 3$, we have $K_{\Delta_{36}} = \mathbb{Q}[x]/(x^3 - 12422194x - 2645665785)$, which means that the algebraic number field extension of \mathbb{Q} contains all three roots of the polynomial. The Maeda conjecture, although still unproven, has been extensively tested for all modular weights up to $2s = 12000$, see [71] where strong evidence is presented to support its validity. More examples of such number fields can be found on the very comprehensive LMFDB database [72] of L-functions and modular forms.

As a final comment we stress that the action of a non-trivial element, σ , of the Galois automorphism group $\text{Aut}_{\mathbb{Q}}(K_{\Delta_{2s}})$ allows us to relate the constants $a_{\Delta_{2s}, m, k}^{\pm}$ for different Hecke eigenforms $\sigma(a_{\Delta_{2s}, m, k}^{\pm}) = a_{\sigma(\Delta_{2s}), m, k}^{\pm}$, hence if Maeda’s conjecture were to be true it would be enough to find one such number $a_{\Delta_{2s}, m, k}^{\pm}$ for a single Hecke cusp Δ_{2s} to deduce all the others.

4 Selection rules on β^{sv} from Tsunogai’s derivation algebra

In this section, we study the interplay between the generating series of MGFs introduced in [28], the modular invariant functions $F_{m,k}^{\pm(s)}$ and an abstract algebra on generating derivations ϵ_k introduced by Tsunogai [48] that is related to holomorphic cusp forms [49]. This connection will clarify why sometimes $\check{F}_{m,k}^{\pm(s)}$ are not modular invariant. The commutation relations among Tsunogai’s derivations ϵ_k with $k = 0, 4, 6, 8, \dots$ govern which linear combinations of the modular completions $F_{m,k}^{\pm(s)}$ appear as MGFs.

The generating series of MGFs introduced in [28] captures the structure of the α' -expansion of certain genus-one integrals in closed-string amplitudes that eventually comprise all MGFs when integrating over sufficiently many torus punctures. More specifically, the first-order differential equations in τ of this generating series is solved by

$$\begin{aligned} \Phi^\tau = 1 &+ \sum_{k=4}^{\infty} \sum_{j=0}^{k-2} \frac{(-1)^j (k-1)}{(k-2-j)!} \beta^{\text{sv}} \left[\begin{matrix} j \\ k \end{matrix} \right] \epsilon_k^{(k-j-2)} \\ &+ \sum_{k_1=4}^{\infty} \sum_{k_2=4}^{\infty} \sum_{j_1=0}^{k_1-2} \sum_{j_2=0}^{k_2-2} \frac{(-1)^{j_1+j_2} (k_1-1)(k_2-1)}{(k_1-2-j_1)!(k_2-2-j_2)!} \beta^{\text{sv}} \left[\begin{matrix} j_1 & j_2 \\ k_1 & k_2 \end{matrix} \right] \epsilon_{k_2}^{(k_2-j_2-2)} \epsilon_{k_1}^{(k_1-j_1-2)} + O(\epsilon_k^3) \end{aligned} \quad (4.1)$$

¹²We would like to thank Herbert Gangl for related discussions.

with conjectural matrix representations of ϵ_k acting on suitable initial values as $\tau \rightarrow i\infty$ that are series in zeta values [30].¹³ In [28, 30] the ϵ_k were explicit finite-dimensional matrix operators that were checked at low orders to obey the relations of Tsuonagai's derivation algebra, and this is conjectured to hold to all orders. Here, we think of the ϵ_k as the abstract generators of Tsuonagai's derivation algebra. In (4.1) and the following we make use of the convenient shorthand notation

$$\epsilon_k^{(j)} = (\text{ad}_{\epsilon_0})^j(\epsilon_k), \quad (4.2)$$

for the repeated adjoint actions $\text{ad}_{\epsilon_0}(\ast) = [\epsilon_0, \ast]$. The suppressed terms $O(\epsilon_k^3)$ in (4.1) are all multiplied by β^{sv} of depth ≥ 3 .

4.1 Overview of ϵ_k relations at depth two

Tsuonagai's derivations satisfy a wealth of commutator relations. First of all, $[\epsilon_2, \epsilon_k] = 0$ with $k = 0, 2, 4, \dots$ identifies ϵ_2 to be a central element which does not occur in the series (4.1). On the remaining derivations, ad_{ϵ_0} enjoys the nilpotency properties

$$\text{ad}_{\epsilon_0}^{k-1}(\epsilon_k) = \epsilon_k^{(k-1)} = 0, \quad k = 4, 6, 8, \dots \quad (4.3)$$

such that the $j_i = 0$ terms in (4.1) exhaust the maximal non-vanishing nested commutators $\epsilon_k^{(k-2)}$. Apart from the relations (4.3) that take a simple and universal form for all k , commutators of $\epsilon_{k_1}, \epsilon_{k_2}, \dots$ at weight $k_1 + k_2 + \dots \geq 14$ obey more involved identities starting with [73, 49, 54]

$$\begin{aligned} 0 &= [\epsilon_{10}, \epsilon_4] - 3[\epsilon_8, \epsilon_6], \\ 0 &= [\epsilon_{14}, \epsilon_4] - \frac{7}{2}[\epsilon_{12}, \epsilon_6] + \frac{11}{2}[\epsilon_{10}, \epsilon_8], \\ 0 &= [\epsilon_{16}, \epsilon_4] - \frac{25}{8}[\epsilon_{14}, \epsilon_6] + \frac{13}{4}[\epsilon_{12}, \epsilon_8]. \end{aligned} \quad (4.4)$$

Nested commutators of three and more derivations obey corollaries of these relations obtained from action of ad_{ϵ_k} . At the same time, there are infinite families of *indecomposable* relations among three or more derivations, i.e. relations that cannot be obtained by repeatedly acting with ad_{ϵ_k} on simpler ones. The simplest indecomposable relation trilinear in derivations reads

$$\begin{aligned} 0 &= 80[\epsilon_{12}, [\epsilon_4, \epsilon_0]] + 16[\epsilon_4, [\epsilon_{12}, \epsilon_0]] - 250[\epsilon_{10}, [\epsilon_6, \epsilon_0]] - 125[\epsilon_6, [\epsilon_{10}, \epsilon_0]] + 280[\epsilon_8, [\epsilon_8, \epsilon_0]] \\ &\quad - 462[\epsilon_4, [\epsilon_4, \epsilon_8]] - 1725[\epsilon_6, [\epsilon_6, \epsilon_4]] \end{aligned} \quad (4.5)$$

and already illustrates a generic feature: When referring to the number of $\epsilon_{k \neq 0}$ in a nested commutator as its *depth*,¹⁴ indecomposable relations involving more than two ϵ_k usually mix terms of different depth. The depth-two terms in the first line of (4.5) affect the appearance of

¹³The generating series of genus-one integrals are denoted by Y^τ in [30] and given by $\Phi^\tau \exp(-\frac{\epsilon_0}{4y}) \widehat{Y}^{i\infty}$, where the initial values $\widehat{Y}^{i\infty}$ comprise all the (conjecturally single-valued [36, 7]) MZVs in the expansion of the resulting MGFs around the cusp.

¹⁴This terminology differs from the work of Pollack [49] where also ϵ_0 is assigned depth one.

β^{sv} of depth two in (4.1) while the depth-three terms in the second line are related to β^{sv} of depth three that are part of the suppressed terms $O(\epsilon_k^3)$ in (4.1) and beyond the scope of this work.

Relations in the derivation algebra are also assigned a notion of depth according to the maximal depth of the nested commutator therein, e.g. (4.5) is said to have depth three. As will be reviewed in the remainder of this section, Pollack determined the depth-two terms in indecomposable relations of arbitrary depth in closed form. More precisely, their rational coefficients are determined from the period polynomials of holomorphic cusp forms in [49].

4.1.1 Cusp forms and depth-two relations

In order to concisely relate the coefficients in relations like (4.4) or (4.5) to holomorphic cusp forms Δ_{2s} of modular weight $2s$, we follow the conventions of [49] for period polynomials

$$\begin{aligned} r_{\Delta_{2s}}(X, Y) &= \int_0^{i\infty} d\tau \Delta_{2s}(\tau)(X - \tau Y)^{2s-2} \\ &= i \sum_{k=0}^{2s-2} \binom{2s-2}{k} X^{2s-2-k} (-iY)^k \Lambda(\Delta_{2s}, k+1), \end{aligned} \quad (4.6)$$

where the arguments of the L-function $\Lambda(\Delta_{2s}, t)$ are all within the critical strip $t \in (0, 2s)$. Moreover, we introduce the even and odd parts of the period polynomials (4.6) via

$$\begin{aligned} r_{\Delta_{2s}}^{\pm}(X, Y) &= \frac{1}{2} [r_{\Delta_{2s}}(X, Y) \pm r_{\Delta_{2s}}(X, -Y)] \\ &= \begin{cases} i \sum_{\substack{k=0 \\ k \text{ even}}}^{2s-2} \binom{2s-2}{k} X^{2s-2-k} (-iY)^k \Lambda(\Delta_{2s}, k+1) & : r_{\Delta_{2s}}^+ \\ i \sum_{\substack{k=0 \\ k \text{ odd}}}^{2s-2} \binom{2s-2}{k} X^{2s-2-k} (-iY)^k \Lambda(\Delta_{2s}, k+1) & : r_{\Delta_{2s}}^- \end{cases} \end{aligned} \quad (4.7)$$

The ratios of L-values at $2s = 12, 16$ and 18 noted in (2.42), (B.1) and (B.3) are equivalent to

$$\begin{aligned} r_{\Delta_{12}}^+(X, Y) &= -\frac{691i\Lambda(\Delta_{12}, 11)}{36} \left\{ X^8 Y^2 - X^2 Y^8 - 3(X^6 Y^4 - X^4 Y^6) - \frac{36}{691}(X^{10} - Y^{10}) \right\}, \\ r_{\Delta_{16}}^+(X, Y) &= -\frac{3617i\Lambda(\Delta_{16}, 15)}{180} \left\{ X^{12} Y^2 - X^2 Y^{12} - \frac{7}{2}(X^{10} Y^4 - X^4 Y^{10}) \right. \\ &\quad \left. + \frac{11}{2}(X^8 Y^6 - X^6 Y^8) - \frac{180}{3617}(X^{14} - Y^{14}) \right\}, \\ r_{\Delta_{18}}^+(X, Y) &= \frac{43867i\Lambda(\Delta_{18}, 17)}{2250} \left\{ X^{14} Y^2 - X^2 Y^{14} - \frac{25}{8}(X^{12} Y^4 - X^4 Y^{12}) \right. \\ &\quad \left. + \frac{13}{4}(X^{10} Y^6 - X^6 Y^{10}) - \frac{2250}{43867}(X^{16} - Y^{16}) \right\}. \end{aligned} \quad (4.8)$$

The rational coefficients $r_{\Delta_{2s}}^+(X, Y)|_{X^a Y^b}$ of the non-zero powers $X^a Y^b$ with $a, b > 0$ are easily seen to match those in the depth-two relations (4.4) among $[\epsilon_{k_1}, \epsilon_{k_2}]$ with $k_1 + k_2 = 2s + 2$. More

generally, the depth-two relations in the derivation algebra are given by

$$0 = \sum_{a+b=2s-2} r_{\Delta_{2s}}^+(X, Y) \Big|_{X^a Y^b} [\epsilon_{a+2}, \epsilon_{b+2}] \quad (4.9)$$

in terms of the even parts of period polynomials (4.6) of cusp forms [49]. The extremal terms $\sim (X^{2s-2} - Y^{2s-2})$ in (4.8) are mapped to coefficients of the vanishing commutators $[\epsilon_2, *]$ in (4.9), that is why their more involved coefficients $\frac{36}{691}, \frac{180}{3617}$ and $\frac{2250}{43867}$ do not enter the depth-two relations (4.4).

4.1.2 Cusp forms and higher-depth relations

Also for higher-depth relations such as (4.5), the coefficients of the depth-two terms $[\epsilon_{k_1}^{(j_1)}, \epsilon_{k_2}^{(j_2)}] = [\text{ad}_{\epsilon_0}^{j_1}(\epsilon_{k_1}), \text{ad}_{\epsilon_0}^{j_2}(\epsilon_{k_2})]$ are determined by the period polynomials (4.6). More specifically, relations of even (odd) depth are governed by the even part $r_{\Delta_{2s}}^+$ (the odd part $r_{\Delta_{2s}}^-$) in (4.7). At weight $2s = 12, 16, 18$, the coefficients in the odd counterpart of (4.8)

$$\begin{aligned} r_{\Delta_{12}}^-(X, Y) &= 10\Lambda(\Delta_{12}, 10) \left\{ X^9 Y + X Y^9 - \frac{25}{4}(X^7 Y^3 + X^3 Y^7) + \frac{21}{2} X^5 Y^5 \right\}, \\ r_{\Delta_{16}}^-(X, Y) &= 14\Lambda(\Delta_{16}, 14) \left\{ X^{13} Y + X Y^{13} - \frac{245}{36}(X^{11} Y^3 + X^3 Y^{11}) \right. \\ &\quad \left. + \frac{539}{36}(X^9 Y^5 + X^5 Y^9) - \frac{55}{3} X^7 Y^7 \right\}, \\ r_{\Delta_{18}}^-(X, Y) &= -16\Lambda(\Delta_{18}, 16) \left\{ X^{15} Y + X Y^{15} - \frac{77}{12}(X^{13} Y^3 + X^3 Y^{13}) \right. \\ &\quad \left. + \frac{91}{8}(X^{11} Y^5 + X^5 Y^{11}) - \frac{143}{24}(X^9 Y^7 + X^7 Y^9) \right\}, \end{aligned} \quad (4.10)$$

enter relations of depth three, five, \dots such as (4.5). By rewriting the first line of (4.5) as

$$-160 \left\{ \frac{[\epsilon_{12}, \epsilon_4^{(1)}]}{2} + \frac{[\epsilon_4, \epsilon_{12}^{(1)}]}{10} - \frac{25}{4} \left(\frac{[\epsilon_{10}, \epsilon_6^{(1)}]}{4} + \frac{[\epsilon_6, \epsilon_{10}^{(1)}]}{8} \right) + \frac{21}{2} \frac{[\epsilon_8, \epsilon_8^{(1)}]}{6} \right\} = 0 \text{ mod depth } 3, \quad (4.11)$$

one can identify the relative factors of $-\frac{25}{4}$ and $\frac{21}{2}$ inside the curly bracket with those in the expression (4.10) for the odd part $r_{\Delta_{12}}^-(X, Y)$. The additional denominators of the combinations $\frac{[\epsilon_{k_1}, \epsilon_{k_2}^{(1)}]}{k_2-2}$ line up with the $d = 3$ instance of the general depth-two combination¹⁵

$$t_{p,q}^d = (d-2)! \sum_{i=0}^{d-2} (-1)^i \frac{(p-2-i)!(q-d+i)!}{i!(p-2)!(d-2-i)!(q-2)!} [\epsilon_p^{(i)}, \epsilon_q^{(d-2-i)}], \quad (4.12)$$

with $d \geq 2$ subject to alternating symmetry properties $t_{p,q}^d = (-1)^{d-1} t_{q,p}^d$. The ratios of factorials in (4.12) are engineered such that $t_{p,q}^d$ is firstly annihilated by $p+q-2d+1$ powers of ad_{ϵ_0} , i.e.

$$\text{ad}_{\epsilon_0}^{p+q-2d+1}(t_{p,q}^d) = 0, \quad \text{ad}_{\epsilon_0}^{p+q-2d}(t_{p,q}^d) \neq 0. \quad (4.13)$$

¹⁵We depart from Pollack's conventions for the commutators (4.12) to ensure that the subscripts of $t_{p,q}^d$ line up with the ϵ_p, ϵ_q in their definition. The commutators $h_{p,q}^d$ in [49] are reproduced by $t_{p,q}^d = h_{p-2,q-2}^d$.

In the simplest instances of (4.12),

$$\begin{aligned}
t_{p,q}^2 &= [\epsilon_p, \epsilon_q], \\
t_{p,q}^3 &= \frac{[\epsilon_p, \epsilon_q^{(1)}]}{q-2} + \frac{[\epsilon_q, \epsilon_p^{(1)}]}{p-2}, \\
t_{p,q}^4 &= \frac{[\epsilon_p, \epsilon_q^{(2)}]}{(q-2)(q-3)} - \frac{2[\epsilon_p^{(1)}, \epsilon_q^{(1)}]}{(p-2)(q-2)} + \frac{[\epsilon_p^{(2)}, \epsilon_q]}{(p-2)(p-3)},
\end{aligned} \tag{4.14}$$

the relative factors ensure that $t_{p,q}^3$ and $t_{p,q}^4$ are annihilated by $p+q-5$ and $p+q-7$ powers of ad_{ϵ_0} , respectively, e.g. that terms of the form $[\epsilon_p^{(p-2)}, \epsilon_q^{(q-2)}]$ cancel from $\text{ad}_{\epsilon_0}^{p+q-5}(t_{p,q}^3)$. One can therefore view the $t_{p,q}^d$ at different values of d (together with their non-vanishing $\text{ad}_{\epsilon_0}^j$ actions at $j = 0, 1, \dots, p+q-2d$) as spanning different $\text{SL}(2)$ representations contained in the tensor product of $\{\epsilon_p^{(j)}, j = 0, 1, \dots, p-2\}$ and $\{\epsilon_q^{(j)}, j = 0, 1, \dots, q-2\}$.

With the combinations of commutators in (4.12) and the even or odd parts of period polynomials in (4.7), Pollack's result for the depth-two coefficients in indecomposable relations takes the compact form [49]

$$\begin{aligned}
0 &= \sum_{a+b=2s-2} r_{\Delta_{2s}}^+(X, Y) \Big|_{X^a Y^b} t_{a+d, b+d}^d \pmod{\text{depth} \geq 3} \quad : \quad d \text{ even}, \\
0 &= \sum_{a+b=2s-2} r_{\Delta_{2s}}^-(X, Y) \Big|_{X^a Y^b} t_{a+d, b+d}^d \pmod{\text{depth} \geq 3} \quad : \quad d \text{ odd}.
\end{aligned} \tag{4.15}$$

In other words, the ratios of odd L-values $\Lambda(\Delta_{2s}, 2n+1)$ (with $n \in \mathbb{N}$) in the even parts $r_{\Delta_{2s}}^+$ determine the coefficients in indecomposable relations among $t_{p,q}^d$ at even d . Conversely, $t_{p,q}^d$ at odd d are related by even L-values $\Lambda(\Delta_{2s}, 2n)$ (with $n \in \mathbb{N}$) in the odd parts $r_{\Delta_{2s}}^-$. In both cases, the L-values are within the critical strip, and the coefficients of higher-depth terms such as the second line of (4.5) remain undetermined by (4.15). The ϵ -weight of the $[\epsilon_{k_1}^{(j_1)}, \epsilon_{k_2}^{(j_2)}]$ in a relation (4.15) due to cusp forms of modular weight $2s$ is given by $k_1 + k_2 = 2s + 2d - 2$.

4.1.3 Examples

The depth-two relations (4.9) are the special case $d = 2$ of (4.15), i.e. the examples in (4.4) are reproduced by the first three lines of

$$\begin{aligned}
r_{\Delta_{12}}^+ &\Rightarrow 0 = -2t_{4,10}^2 + 6t_{6,8}^2, \\
r_{\Delta_{16}}^+ &\Rightarrow 0 = -2t_{4,14}^2 + 7t_{6,12}^2 - 11t_{8,10}^2, \\
r_{\Delta_{18}}^+ &\Rightarrow 0 = -2t_{4,16}^2 + \frac{25}{4}t_{6,14}^2 - \frac{13}{2}t_{8,12}^2, \\
r_{\Delta_{20}}^+ &\Rightarrow 0 = -2t_{4,18}^2 + \frac{20}{3}t_{6,16}^2 - \frac{28}{3}t_{8,14}^2 + \frac{26}{3}t_{10,12}^2, \\
r_{\Delta_{22}}^+ &\Rightarrow 0 = -2t_{4,20}^2 + \frac{105}{16}t_{6,18}^2 - \frac{17}{2}t_{8,16}^2 + \frac{85}{16}t_{10,14}^2,
\end{aligned} \tag{4.16}$$

depth	2	3	4	5
Δ_{12}	14	16	18	20
Δ_{16}	18	20	22	24
Δ_{18}	20	22	24	26
Δ_{20}	22	24	26	28
Δ_{22}	24	26	28	30
$\Delta_{24,i}, \Delta_{24,ii}$	2×26	2×28	2×30	2×32

Table 1: The number given corresponds to the ϵ -weight of the indecomposable relation of the given depth triggered by the cusp form Δ_{2s} , so that for instance Δ_{12} induces one indecomposable relation of depth 2 at weight 14. There are two indecomposable relations of depth 2 at weight 26 due to the two cusp forms of modular weight 24.

where $t_{2,q}^d = 0$ since ϵ_2 commutes with all the ϵ_k . Indecomposable relations at depth three due to (4.15) at $d = 3$ include

$$\begin{aligned}
r_{\Delta_{12}}^- &\Rightarrow 0 = 2t_{4,12}^3 - \frac{25}{2}t_{6,10}^3 + \frac{21}{2}t_{8,8}^3 \pmod{\text{depth} \geq 3}, \\
r_{\Delta_{16}}^- &\Rightarrow 0 = 2t_{4,16}^3 - \frac{245}{18}t_{6,14}^3 + \frac{539}{18}t_{8,12}^3 - \frac{55}{3}t_{10,10}^3 \pmod{\text{depth} \geq 3}, \\
r_{\Delta_{18}}^- &\Rightarrow 0 = 2t_{4,18}^3 - \frac{77}{6}t_{6,16}^3 + \frac{91}{4}t_{8,14}^3 - \frac{143}{12}t_{10,12}^3 \pmod{\text{depth} \geq 3}, \\
r_{\Delta_{20}}^- &\Rightarrow 0 = 2t_{4,20}^3 - \frac{715}{54}t_{6,18}^3 + \frac{242}{9}t_{8,16}^3 - \frac{260}{9}t_{10,14}^3 + \frac{715}{54}t_{12,12}^3 \pmod{\text{depth} \geq 3}, \\
r_{\Delta_{22}}^- &\Rightarrow 0 = 2t_{4,22}^3 - \frac{1183}{90}t_{6,20}^3 + \frac{1547}{60}t_{8,18}^3 - \frac{697}{30}t_{10,16}^3 + \frac{1547}{180}t_{12,14}^3 \pmod{\text{depth} \geq 3},
\end{aligned} \tag{4.17}$$

whose first line due to $r_{\Delta_{12}}^-$ is equivalent to (4.11). The relations at higher $d \geq 4$ start with

$$\begin{aligned}
r_{\Delta_{12}}^+ &\Rightarrow 0 = \frac{72}{691}t_{4,14}^4 - 2t_{6,12}^4 + 6t_{8,10}^4 \pmod{\text{depth} \geq 3}, \\
r_{\Delta_{16}}^+ &\Rightarrow 0 = \frac{360}{3617}t_{4,18}^4 - 2t_{6,16}^4 + 7t_{8,14}^4 - 11t_{10,12}^4 \pmod{\text{depth} \geq 3},
\end{aligned} \tag{4.18}$$

as well as

$$\begin{aligned}
r_{\Delta_{12}}^- &\Rightarrow 0 = 2t_{6,14}^5 - \frac{25}{2}t_{8,12}^5 + \frac{21}{2}t_{10,10}^5 \pmod{\text{depth} \geq 3}, \\
r_{\Delta_{16}}^- &\Rightarrow 0 = 2t_{6,18}^5 - \frac{245}{18}t_{8,16}^5 + \frac{539}{18}t_{10,14}^5 - \frac{55}{3}t_{12,12}^5 \pmod{\text{depth} \geq 3},
\end{aligned} \tag{4.19}$$

and feed into relations among $[[\dots [[\epsilon_{k_1}, \epsilon_{k_2}], \epsilon_{k_3}], \dots], \epsilon_{k_\ell}]$ of increasing ϵ -weight $k_1 + k_2 + \dots + k_\ell$. Table 1 summarises the counting of indecomposable higher-depth relations along with their ϵ_k -weight (which is $p+q$ in case of $t_{p,q}^d$ for arbitrary $d \geq 2$) due to given holomorphic cusp forms.

4.1.4 Comparison with the eMZV datamine

The explicit form of various indecomposable relations up to depth 5 can be downloaded from the datamine [74] of relations among elliptic multiple zeta values (eMZVs) [75]. The datamine

uses the notation $\mathbf{rRel}[w, r]$ for relations of ϵ -weight w and depth r . The depth-two parts of these relations $\mathbf{rRel}[w, r]$ are usually combinations of several relations (4.15) at different d . For instance, $\mathbf{rRel}[20, 3]$ at depth three mixes the specialisations of (4.15) to $(r_{\Delta_{16}}^-, d=3)$ with ad_{ϵ_0} acting on the relation from $(r_{\Delta_{18}}^+, d=2)$. Similarly, the depth-two terms of $\mathbf{rRel}[18, 4]$ can be reconstructed from $(r_{\Delta_{12}}^+, d=4)$ and $\text{ad}_{\epsilon_0}^2$ action on $(r_{\Delta_{16}}^+, d=2)$ whereas $\mathbf{rRel}[20, 5]$ mixes input from all of $(r_{\Delta_{12}}^-, d=5)$, $\text{ad}_{\epsilon_0}^2(r_{\Delta_{16}}^-, d=3)$ and $\text{ad}_{\epsilon_0}^3(r_{\Delta_{18}}^+, d=2)$.

4.2 Modular graph forms and ϵ_k relations at depth two

Based on the relations among depth-two commutators $[\epsilon_{k_1}^{(j_1)}, \epsilon_{k_2}^{(j_2)}]$ reviewed above, we shall now describe the dropouts of iterated Eisenstein integrals of depth two from the generating series (4.1) of MGFs. We will be interested in the modular invariant cases where the entries of $\beta^{\text{sv}} \begin{bmatrix} j_1 & j_2 \\ k_1 & k_2 \end{bmatrix}$ obey $2j_1+2j_2+4 = k_1+k_2$, see (2.51). These cases of β^{sv} are related to the $F_{m,k}^{\pm(s)}$ with $2m+2k = k_1+k_2$, modulo shuffles $\beta^{\text{sv}} \begin{bmatrix} j_1 \\ k_1 \end{bmatrix} \beta^{\text{sv}} \begin{bmatrix} j_2 \\ k_2 \end{bmatrix}$ and lower-depth terms. The derivations $[\epsilon_{k_1}^{(j_1)}, \epsilon_{k_2}^{(j_2)}]$ relevant to modular invariant terms in (4.1) involve a total of $j_1+j_2 = \frac{1}{2}(k_1+k_2)-2$ powers of ad_{ϵ_0} .

In the following, we shall rewrite the shuffle-irreducible modular invariants with β^{sv} at depth two in terms of $F_{m,k}^{\pm(s)}$. In this way, the $[\epsilon_{k_1}^{(j_1)}, \epsilon_{k_2}^{(j_2)}]$ conspire to the commutators $t_{p,q}^d$ defined in (4.12) and their images under $\text{ad}_{\epsilon_0}^N$ with $N \leq p+q-2d$,

$$\text{ad}_{\epsilon_0}^N[\epsilon_{k_1}^{(j_1)}, \epsilon_{k_2}^{(j_2)}] = \sum_{j=0}^N \binom{N}{j} [\epsilon_{k_1}^{(j_1+j)}, \epsilon_{k_2}^{(j_2+N-j)}]. \quad (4.20)$$

We will decompose the generating series (4.1) into depth-two sectors $\Phi^\tau(k_1, k_2)$ associated with double integrals of given (G_{k_1}, G_{k_2}) . The modular invariant contributions are isolated by means of the delta function in

$$\begin{aligned} \Phi^\tau(k_1, k_2) &= (k_1-1)(k_2-1) \sum_{j_1=0}^{k_1-2} \sum_{j_2=0}^{k_2-2} \frac{(-1)^{j_1+j_2} \delta(2j_1+2j_2+4-k_1-k_2)}{(k_1-2-j_1)!(k_2-2-j_2)!} \\ &\quad \times \beta^{\text{sv}} \begin{bmatrix} j_1 & j_2 \\ k_1 & k_2 \end{bmatrix} \epsilon_{k_2}^{(k_2-j_2-2)} \epsilon_{k_1}^{(k_1-j_1-2)}, \end{aligned} \quad (4.21)$$

and the depth-two modular-invariant part of the generating series (4.1) is obtained by summing over $k_1, k_2 \in 2\mathbb{N}+2$. The shuffle-irreducible modular invariants of the (G_4, G_6) sector are for instance encoded in

$$\begin{aligned} \Phi^\tau(4, 6) + \Phi^\tau(6, 4) &= -\frac{5}{2}(\beta^{\text{sv}} \begin{bmatrix} 1 & 2 \\ 6 & 4 \end{bmatrix} \epsilon_4 \epsilon_6^{(3)} + 3\beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 6 & 4 \end{bmatrix} \epsilon_4^{(1)} \epsilon_6^{(2)} + 3\beta^{\text{sv}} \begin{bmatrix} 0 & 3 \\ 4 & 6 \end{bmatrix} \epsilon_6^{(1)} \epsilon_4^{(2)} \\ &\quad + 3\beta^{\text{sv}} \begin{bmatrix} 3 & 0 \\ 6 & 4 \end{bmatrix} \epsilon_4^{(2)} \epsilon_6^{(1)} + 3\beta^{\text{sv}} \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix} \epsilon_6^{(2)} \epsilon_4^{(1)} + \beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 4 & 6 \end{bmatrix} \epsilon_6^{(3)} \epsilon_4). \end{aligned} \quad (4.22)$$

Using (2.6) we can rewrite $\beta^{\text{sv}} \begin{bmatrix} j_1 & j_2 \\ k_1 & k_2 \end{bmatrix} = -\beta^{\text{sv}} \begin{bmatrix} j_2 & j_1 \\ k_2 & k_1 \end{bmatrix} \pmod{\sqcup}$, where \sqcup refers to shuffle products (2.6) of depth-one β^{sv} . Picking the basis¹⁶

$$\begin{aligned} F_{2,3}^{+(3)} &= 30\beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 4 & 6 \end{bmatrix} + 30\beta^{\text{sv}} \begin{bmatrix} 3 & 0 \\ 6 & 4 \end{bmatrix} \pmod{\text{lower depth}}, \\ F_{2,3}^{-(2)} &= -90\beta^{\text{sv}} \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix} + 90\beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 4 & 6 \end{bmatrix} + 90\beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 6 & 4 \end{bmatrix} - 90\beta^{\text{sv}} \begin{bmatrix} 3 & 0 \\ 6 & 4 \end{bmatrix} \pmod{\text{lower depth}}, \\ F_{2,3}^{-(4)} &= -90\beta^{\text{sv}} \begin{bmatrix} 1 & 2 \\ 4 & 6 \end{bmatrix} - 60\beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 4 & 6 \end{bmatrix} + 90\beta^{\text{sv}} \begin{bmatrix} 2 & 1 \\ 6 & 4 \end{bmatrix} + 60\beta^{\text{sv}} \begin{bmatrix} 3 & 0 \\ 6 & 4 \end{bmatrix} \pmod{\text{lower depth}}, \end{aligned} \quad (4.23)$$

of shuffle irreducibles, the derivations in (4.22) conspire to the following ad_{ϵ_0} -images of the commutators $t_{p,q}^d$ in (4.12):

$$\begin{aligned} \Phi^\tau(4,6) + \Phi^\tau(6,4) &= \frac{1}{10} F_{2,3}^{-(2)} \text{ad}_{\epsilon_0}(t_{4,6}^4) - \frac{1}{120} F_{2,3}^{-(4)} \text{ad}_{\epsilon_0}^3(t_{4,6}^2) \\ &\quad + \frac{1}{6} F_{2,3}^{+(3)} \text{ad}_{\epsilon_0}^2(t_{4,6}^3) \pmod{\text{lower depth} \ \& \ \sqcup}. \end{aligned} \quad (4.24)$$

The analogous expressions at weights $m+k \geq 7$ will in the first place involve the combinations $\check{F}_{m,k}^{\pm(s)}$ of β^{sv} rather than the full modular invariants $F_{m,k}^{\pm(s)}$: The iterated integrals (2.29) of cusp forms are consistently absent from the generating series (4.1). In the matrix representations of (4.1) relevant to closed-string genus-one integrals [30], the combinations of $\beta^{\text{sv}} \begin{bmatrix} j \\ k \end{bmatrix}$ contributing to MGFs at depth two can be recovered by the initial conditions $\tau \rightarrow i\infty$ that the derivations in Φ^τ act on.

However, the initial conditions do not allow us to retrieve the iterated integrals of cusp forms in (2.29): They do not have any known realisation in closed-string integrals over torus punctures since Cauchy–Riemann derivatives of MGFs [8] or their generating series [28] do not introduce any holomorphic cusp forms. Many of the subsequent equations will hold modulo lower depth and shuffles as in (4.24), and we will indicate by using \cong in the place of $=$ that shuffles, β^{sv} of depth one and depth-zero terms have been dropped while depth-one integrals of holomorphic cusp forms are still tracked.

We will exemplify in the following sections that the depth-two terms (4.15) of ϵ_k -relations are sufficient to effectively replace all the $\check{F}_{m,k}^{\pm(s)}$ in Φ^τ by $F_{m,k}^{\pm(s)}$. In all cases up to and including $m+k = 14$, the coefficient of $\check{F}_{m,k}^{\pm(s)}$ is checked to be a \mathbb{Q} -multiple of $\text{ad}_{\epsilon_0}^{s-1}(t_{2m,2k}^{m+k-s+1})$. Given that $m+k-s$ is even for $\check{F}_{m,k}^{+(s)}$ and odd for $\check{F}_{m,k}^{-(s)}$, the transition to $F_{m,k}^{+(s)}$ (to $F_{m,k}^{-(s)}$) by adding cusp-form contributions is governed by $t_{p,q}^d$ at odd values of d (even values of d).

¹⁶The relation between (4.23) and MGFs in their lattice-sum representations is discussed in section 5.4 of Part I and section 5 of [30].

4.2.1 Weight 7

At weight $m+k=7$, the analogue of (4.24) for the shuffle-irreducible modular invariants reads

$$\begin{aligned}
\Phi^\tau(4, 10) + \Phi^\tau(10, 4) &\cong \frac{\check{F}_{2,5}^{-(4)}}{540} \text{ad}_{\epsilon_0}^3(t_{4,10}^4) - \frac{\check{F}_{2,5}^{-(6)}}{30240} \text{ad}_{\epsilon_0}^5(t_{4,10}^2) + \frac{\check{F}_{2,5}^{+(5)}}{420} \text{ad}_{\epsilon_0}^4(t_{4,10}^3), \\
\Phi^\tau(6, 8) + \Phi^\tau(8, 6) &\cong -\frac{3\check{F}_{3,4}^{-(2)}}{140} \text{ad}_{\epsilon_0}(t_{6,8}^6) + \frac{\check{F}_{3,4}^{-(4)}}{360} \text{ad}_{\epsilon_0}^3(t_{6,8}^4) - \frac{\check{F}_{3,4}^{-(6)}}{30240} \text{ad}_{\epsilon_0}^5(t_{6,8}^2) \\
&\quad - \frac{\check{F}_{3,4}^{+(3)}}{14} \text{ad}_{\epsilon_0}^2(t_{6,8}^5) + \frac{\check{F}_{3,4}^{+(5)}}{280} \text{ad}_{\epsilon_0}^4(t_{6,8}^3).
\end{aligned} \tag{4.25}$$

There are no cusp-form contributions to the even functions at this weight, $\check{F}_{m,7-m}^{+(s)} = F_{m,7-m}^{+(s)}$, but the individual differences $\check{F}_{m,7-m}^{-(6)} - F_{m,7-m}^{-(6)}$ in (3.7) involve the odd combination $H_{\Delta_{12}}^-$. This leads to the extra terms $\sim H_{\Delta_{12}}^-$ in the last line of

$$\begin{aligned}
\Phi^\tau(4, 10) + \Phi^\tau(10, 4) + \Phi^\tau(6, 8) + \Phi^\tau(8, 6) &\cong \frac{F_{2,5}^{-(4)}}{540} \text{ad}_{\epsilon_0}^3(t_{4,10}^4) - \frac{F_{2,5}^{-(6)}}{30240} \text{ad}_{\epsilon_0}^5(t_{4,10}^2) \\
&\quad + \frac{F_{2,5}^{+(5)}}{420} \text{ad}_{\epsilon_0}^4(t_{4,10}^3) - \frac{3F_{3,4}^{-(2)}}{140} \text{ad}_{\epsilon_0}(t_{6,8}^6) + \frac{F_{3,4}^{-(4)}}{360} \text{ad}_{\epsilon_0}^3(t_{6,8}^4) - \frac{F_{3,4}^{-(6)}}{30240} \text{ad}_{\epsilon_0}^5(t_{6,8}^2) \\
&\quad - \frac{F_{3,4}^{+(3)}}{14} \text{ad}_{\epsilon_0}^2(t_{6,8}^5) + \frac{F_{3,4}^{+(5)}}{280} \text{ad}_{\epsilon_0}^4(t_{6,8}^3) - \frac{\Lambda(\Delta_{12}, 12)H_{\Delta_{12}}^-}{2268000\Lambda(\Delta_{12}, 10)} \text{ad}_{\epsilon_0}^5(t_{4,10}^2 - 3t_{6,8}^2),
\end{aligned} \tag{4.26}$$

which vanish due to $\text{ad}_{\epsilon_0}^5(t_{4,10}^2 - 3t_{6,8}^2) = 0$, i.e. due to the ϵ_k -relation in the first line of (4.16). Using the same relation also for the coefficients of $F_{m,k}^{-(6)}$ shows that the linear combination $3F_{2,5}^{-(6)} + F_{3,4}^{-(6)}$ appears, consistent with (3.8).

4.2.2 Weight 8

At weight $m+k=8$, we can similarly start from

$$\begin{aligned}
\Phi^\tau(4, 12) + \Phi^\tau(12, 4) &\cong -\frac{\check{F}_{2,6}^{-(5)}}{7392} \text{ad}_{\epsilon_0}^4(t_{4,12}^4) + \frac{\check{F}_{2,6}^{-(7)}}{665280} \text{ad}_{\epsilon_0}^6(t_{4,12}^2) - \frac{\check{F}_{2,6}^{+(6)}}{6048} \text{ad}_{\epsilon_0}^5(t_{4,12}^3), \\
\Phi^\tau(6, 10) + \Phi^\tau(10, 6) &\cong \frac{\check{F}_{3,5}^{-(3)}}{252} \text{ad}_{\epsilon_0}^2(t_{6,10}^6) - \frac{\check{F}_{3,5}^{-(5)}}{4620} \text{ad}_{\epsilon_0}^4(t_{6,10}^4) + \frac{\check{F}_{3,5}^{-(7)}}{665280} \text{ad}_{\epsilon_0}^6(t_{6,10}^2) \\
&\quad + \frac{\check{F}_{3,5}^{+(4)}}{90} \text{ad}_{\epsilon_0}^3(t_{6,10}^5) - \frac{\check{F}_{3,5}^{+(6)}}{3780} \text{ad}_{\epsilon_0}^5(t_{6,10}^3), \\
\Phi^\tau(8, 8) &\cong -\frac{9\check{F}_{4,4}^{+(2)}}{280} \text{ad}_{\epsilon_0}(t_{8,8}^7) + \frac{\check{F}_{4,4}^{+(4)}}{120} \text{ad}_{\epsilon_0}^3(t_{8,8}^5) - \frac{\check{F}_{4,4}^{+(6)}}{6720} \text{ad}_{\epsilon_0}^5(t_{8,8}^3),
\end{aligned} \tag{4.27}$$

and immediately identify $\check{F}_{m,8-m}^{-(s)} = F_{m,8-m}^{-(s)}$ since all of these odd depth-two combinations are modular invariant without any need for cusp forms. However, relating $\check{F}_{m,8-m}^{+(6)}$ to $F_{m,8-m}^{+(6)}$

introduces the even combination $H_{\Delta_{12}}^+$ via (3.11), leading to extra terms in the last line of

$$\begin{aligned}
& \Phi^\tau(4, 12) + \Phi^\tau(12, 4) + \Phi^\tau(6, 10) + \Phi^\tau(10, 6) + \Phi^\tau(8, 8) \\
& \cong -\frac{F_{2,6}^{-(5)}}{7392} \text{ad}_{\epsilon_0}^4(t_{4,12}^4) + \frac{F_{2,6}^{-(7)}}{665280} \text{ad}_{\epsilon_0}^6(t_{4,12}^2) - \frac{F_{2,6}^{+(6)}}{6048} \text{ad}_{\epsilon_0}^5(t_{4,12}^3) + \frac{F_{3,5}^{-(3)}}{252} \text{ad}_{\epsilon_0}^2(t_{6,10}^6) \\
& - \frac{F_{3,5}^{-(5)}}{4620} \text{ad}_{\epsilon_0}^4(t_{6,10}^4) + \frac{F_{3,5}^{-(7)}}{665280} \text{ad}_{\epsilon_0}^6(t_{6,10}^2) + \frac{F_{3,5}^{+(4)}}{90} \text{ad}_{\epsilon_0}^3(t_{6,10}^5) - \frac{F_{3,5}^{+(6)}}{3780} \text{ad}_{\epsilon_0}^5(t_{6,10}^3) \quad (4.28) \\
& - \frac{9F_{4,4}^{+(2)}}{280} \text{ad}_{\epsilon_0}(t_{8,8}^7) + \frac{F_{4,4}^{+(4)}}{120} \text{ad}_{\epsilon_0}^3(t_{8,8}^5) - \frac{F_{4,4}^{+(6)}}{6720} \text{ad}_{\epsilon_0}^5(t_{8,8}^3) \\
& + \frac{\Lambda(\Delta_{12}, 13)H_{\Delta_{12}}^+}{208958400\Lambda(\Delta_{12}, 11)} \text{ad}_{\epsilon_0}^5(4t_{4,12}^3 - 25t_{6,10}^3 + 21t_{8,8}^3).
\end{aligned}$$

In this case, it is the depth-two terms in first line of (4.17) which imply the vanishing of $\text{ad}_{\epsilon_0}^5(4t_{4,12}^3 - 25t_{6,10}^3 + 21t_{8,8}^3)$ and therefore the dropout of $H_{\Delta_{12}}^+$ modulo higher-depth terms that are given in the second line of (4.5). The higher-depth commutators of ϵ_k in the second line of (4.5) will be associated with higher-depth β^{sv} , and their modular invariant completions must also contain the iterated integrals of $H_{\Delta_{12}}^+$ of depth one. Using the first line of (4.17) also for the coefficients of the functions $F_{m,k}^{+(6)}$ in (4.28) reproduces the linear combinations appearing in (3.12) (modulo higher depth).

4.2.3 Weight 9

Similar to (4.26) at weight seven, the even functions $\check{F}_{m,9-m}^{+(s)}$ at weight $m+k=9$ are all identical to the modular invariants $F_{m,9-m}^{+(s)}$. The odd functions in turn introduce iterated integrals of the cusp forms Δ_{12} and Δ_{16} from the Laplace eigenspaces with $s=6$ and $s=8$, respectively:

$$\begin{aligned}
& \Phi^\tau(4, 14) + \Phi^\tau(14, 4) + \Phi^\tau(6, 12) + \Phi^\tau(12, 6) + \Phi^\tau(8, 10) + \Phi^\tau(10, 8) \\
& \cong \frac{F_{2,7}^{-(6)}}{131040} \text{ad}_{\epsilon_0}^5(t_{4,14}^4) - \frac{F_{2,7}^{-(8)}}{17297280} \text{ad}_{\epsilon_0}^7(t_{4,14}^2) + \frac{F_{2,7}^{+(7)}}{110880} \text{ad}_{\epsilon_0}^6(t_{4,14}^3) - \frac{F_{3,6}^{-(4)}}{2376} \text{ad}_{\epsilon_0}^3(t_{6,12}^6) \\
& + \frac{F_{3,6}^{-(6)}}{78624} \text{ad}_{\epsilon_0}^5(t_{6,12}^4) - \frac{F_{3,6}^{-(8)}}{17297280} \text{ad}_{\epsilon_0}^7(t_{6,12}^2) - \frac{F_{3,6}^{+(5)}}{924} \text{ad}_{\epsilon_0}^4(t_{6,12}^5) + \frac{F_{3,6}^{+(7)}}{66528} \text{ad}_{\epsilon_0}^6(t_{6,12}^3) \\
& + \frac{F_{4,5}^{-(2)}}{210} \text{ad}_{\epsilon_0}(t_{8,10}^8) - \frac{F_{4,5}^{-(4)}}{1320} \text{ad}_{\epsilon_0}^3(t_{8,10}^6) + \frac{F_{4,5}^{-(6)}}{65520} \text{ad}_{\epsilon_0}^5(t_{8,10}^4) - \frac{F_{4,5}^{-(8)}}{17297280} \text{ad}_{\epsilon_0}^7(t_{8,10}^2) \quad (4.29) \\
& + \frac{F_{4,5}^{+(3)}}{42} \text{ad}_{\epsilon_0}^2(t_{8,10}^7) - \frac{3F_{4,5}^{+(5)}}{1540} \text{ad}_{\epsilon_0}^4(t_{8,10}^5) + \frac{F_{4,5}^{+(7)}}{55440} \text{ad}_{\epsilon_0}^6(t_{8,10}^3) \\
& - \frac{691\Lambda(\Delta_{12}, 14)H_{\Delta_{12}}^-}{244481328000\Lambda(\Delta_{12}, 10)} \text{ad}_{\epsilon_0}^5\left(\frac{36}{691}t_{4,14}^4 - t_{6,12}^4 + 3t_{8,10}^4\right) \\
& - \frac{\Lambda(\Delta_{16}, 16)H_{\Delta_{16}}^-}{19559232000\Lambda(\Delta_{16}, 14)} \text{ad}_{\epsilon_0}^7(2t_{4,14}^2 - 7t_{6,12}^2 + 11t_{8,10}^2).
\end{aligned}$$

Both combinations $\frac{36}{691}t_{4,14}^4 - t_{6,12}^4 + 3t_{8,10}^4$ and $2t_{4,14}^2 - 7t_{6,12}^2 + 11t_{8,10}^2$ of ϵ_k commutators vanish by (4.18) and (4.16), respectively (modulo higher depth in the case of the $t_{p,q}^4$).

4.2.4 Weight 10

Starting from weight 10, both even and odd functions $\check{F}_{m,k}^{\pm(s)}$ involve Laplace eigenvalues $s = 6, 8, 9, \dots$ where cusp forms are required in the modular completion to $F_{m,k}^{\pm(s)}$. All of Δ_{12}, Δ_{16} and Δ_{18} feature in the corrections in the lowest lines of the weight-10 terms at depth two,

$$\begin{aligned}
& \Phi^\tau(4, 16) + \Phi^\tau(16, 4) + \Phi^\tau(6, 14) + \Phi^\tau(14, 6) + \Phi^\tau(8, 12) + \Phi^\tau(12, 8) + \Phi^\tau(10, 10) \\
& \cong \frac{F_{5,5}^{+(2)}}{105} \text{ad}_{\epsilon_0}(t_{10,10}^9) - \frac{F_{5,5}^{+(4)}}{330} \text{ad}_{\epsilon_0}^3(t_{10,10}^7) + \frac{F_{5,5}^{+(6)}}{10920} \text{ad}_{\epsilon_0}^5(t_{10,10}^5) - \frac{F_{5,5}^{+(8)}}{2162160} \text{ad}_{\epsilon_0}^7(t_{10,10}^3) \\
& - \frac{F_{2,8}^{-(7)}}{2851200} \text{ad}_{\epsilon_0}^6(t_{4,16}^4) + \frac{F_{2,8}^{-(9)}}{518918400} \text{ad}_{\epsilon_0}^8(t_{4,16}^2) - \frac{F_{2,8}^{+(8)}}{2471040} \text{ad}_{\epsilon_0}^7(t_{4,16}^3) + \frac{F_{3,7}^{-(5)}}{32032} \text{ad}_{\epsilon_0}^4(t_{6,14}^6) \\
& - \frac{F_{3,7}^{-(7)}}{1663200} \text{ad}_{\epsilon_0}^6(t_{6,14}^4) + \frac{F_{3,7}^{-(9)}}{518918400} \text{ad}_{\epsilon_0}^8(t_{6,14}^2) + \frac{F_{3,7}^{+(6)}}{13104} \text{ad}_{\epsilon_0}^5(t_{6,14}^5) - \frac{F_{3,7}^{+(8)}}{1441440} \text{ad}_{\epsilon_0}^7(t_{6,14}^3) \\
& - \frac{5F_{4,6}^{-(3)}}{5544} \text{ad}_{\epsilon_0}^2(t_{8,12}^8) + \frac{F_{4,6}^{-(5)}}{16016} \text{ad}_{\epsilon_0}^4(t_{8,12}^6) - \frac{F_{4,6}^{-(7)}}{1330560} \text{ad}_{\epsilon_0}^6(t_{8,12}^4) + \frac{F_{4,6}^{-(9)}}{518918400} \text{ad}_{\epsilon_0}^8(t_{8,12}^2) \quad (4.30) \\
& - \frac{F_{4,6}^{+(4)}}{264} \text{ad}_{\epsilon_0}^3(t_{8,12}^7) + \frac{F_{4,6}^{+(6)}}{6552} \text{ad}_{\epsilon_0}^5(t_{8,12}^5) - \frac{F_{4,6}^{+(8)}}{1153152} \text{ad}_{\epsilon_0}^7(t_{8,12}^3) \\
& - \frac{\Lambda(\Delta_{12}, 15)H_{\Delta_{12}}^+}{38030428800\Lambda(\Delta_{12}, 11)} \text{ad}_{\epsilon_0}^5(4t_{6,14}^5 - 25t_{8,12}^5 + 21t_{10,10}^5) \\
& + \frac{\Lambda(\Delta_{16}, 17)H_{\Delta_{16}}^+}{8085227673600\Lambda(\Delta_{16}, 15)} \text{ad}_{\epsilon_0}^7(36t_{4,16}^3 - 245t_{6,14}^3 + 539t_{8,12}^3 - 330t_{10,10}^3) \\
& - \frac{\Lambda(\Delta_{18}, 18)H_{\Delta_{18}}^-}{5230697472000\Lambda(\Delta_{18}, 16)} \text{ad}_{\epsilon_0}^8(8t_{4,16}^2 - 25t_{6,14}^2 + 26t_{8,12}^2).
\end{aligned}$$

Again, all the $H_{\Delta_{2s}}^\pm$ are accompanied by combinations $4t_{6,14}^5 - 25t_{8,12}^5 + 21t_{10,10}^5$ as well as $36t_{4,16}^3 - 245t_{6,14}^3 + 539t_{8,12}^3 - 330t_{10,10}^3$ and $8t_{4,16}^2 - 25t_{6,14}^2 + 26t_{8,12}^2$ which vanish by (4.19), (4.17) and (4.16), respectively, modulo higher depth.

4.2.5 Summary

In this section, we have demonstrated in detail how (specific linear combinations of) the modular functions $F_{m,k}^{\pm(s)}$ appear in the generating series of MGFs. On the one hand, the integral- or lattice-sum representations of MGFs manifest that they are modular forms; on the other hand, their differential equations [8, 28] rule out iterated integrals of holomorphic cusp forms. These requirements have been explicitly confirmed for the β^{sv} -contributions (4.1) to modular invariant MGFs at depth two and a wide range of weights.

By reorganising the shuffle irreducible β^{sv} of depth two in terms of $\check{F}_{m,k}^{\pm(s)}$, the accompanying derivations in (4.1) conspire to specific combinations $t_{p,q}^d$ of commutators defined in (4.12) that are singled out by representation theory of $\text{SL}(2, \mathbb{Z})$. More importantly, these combinations $t_{p,q}^d$ were identified by Pollack [49] to streamline relations in the derivation algebra. By rewriting parts of the generating series (4.1) in terms of $\check{F}_{m,k}^{\pm(s)}$ and $t_{p,q}^d$, we have exhibited the interplay between Pollack's relations and the dropout of the modular completions $H_{\Delta_{2s}}^\pm$ from the generating series of MGFs at depth two. These findings are consistent with the main result of this

work that not all linear combinations of $F_{m,k}^{\pm(s)}$ can be represented by β^{sv} of depth one and two, powers of y and odd zeta values.

5 Kloosterman sums and the Fourier expansion of the $F_{m,k}^{\pm(s)}$

In this last section we want to highlight some of the “side-effects” due to the presence of iterated integrals of holomorphic cusp forms in the generic expression (3.1) for $F_{m,k}^{\pm(s)}$. One of the consequences will be to provide a connection between L-values and Kloosterman sums that come out of the Poincaré-series representation of the $F_{m,k}^{\pm(s)}$ as anticipated in Part I.

The addition of the iterated integrals over cusp forms has consequences for the Fourier expansion of the $F_{m,k}^{\pm(s)}$: In the following, we will compare different approaches to determining the coefficients of $q^a \bar{q}^b$ in the expansion around the cusp

$$F_{m,k}^{\pm(s)}(\tau) = \sum_{a,b=0}^{\infty} d_{a,b}(y) q^a \bar{q}^b. \quad (5.1)$$

The coefficients $d_{a,b}(y)$ are Laurent polynomials in $y = \pi \text{Im } \tau$ (with powers ranging from y^{m+k} to y^{-m-k+2}) which will be referred to as $F_{m,k}^{\pm(s)}|_{q^a \bar{q}^b}$ in the rest of this section. They are straightforwardly determined from the representation of $F_{m,k}^{\pm(s)}$ in terms of β^{sv} and $H_{\Delta_{2s}}^{\pm}$ and carry highly non-trivial information on (infinite combinations of) Kloosterman sums by comparing with the Poincaré-series representations (1.5) of $F_{m,k}^{\pm(s)}$. As detailed in Part I, the β^{sv} -part $\check{F}_{m,k}^{\pm(s)}$ only introduces single-valued zeta values and rational numbers into $d_{a,b}(y)$. However, the novelty is that now the addition of $H_{\Delta_{2s}}^{\pm}$ also introduces L-values into the Fourier coefficients in (5.1).

5.1 Odd example

As a first example we assemble the order $q^1 \bar{q}^0$ term in $F_{2,5}^{-(6)}$ from (3.2) and (3.7),

$$\begin{aligned} F_{2,5}^{-(6)}|_{q^1 \bar{q}^0} &= \frac{2y^5}{93555} - \frac{y^2}{1080} - \frac{y}{360} + \frac{19}{720} + \frac{2}{9y} + \frac{301}{384y^2} + \frac{301}{192y^3} + \frac{903}{512y^4} + \frac{903}{1024y^5} \\ &\quad - \zeta_3 \left(\frac{1}{24y} + \frac{1}{4y^2} + \frac{21}{32y^3} + \frac{7}{8y^4} + \frac{63}{128y^5} \right) + \zeta_9 \left(\frac{35}{64y^4} + \frac{63}{128y^5} \right) \\ &\quad - \frac{\Lambda(\Delta_{12}, 12)}{\Lambda(\Delta_{12}, 10)} \left(\frac{1}{150} + \frac{1}{20y} + \frac{7}{40y^2} + \frac{7}{20y^3} + \frac{63}{160y^4} + \frac{63}{320y^5} \right). \end{aligned} \quad (5.2)$$

Here, we can see clearly the separate contributions from $\check{F}_{2,5}^{-(6)}$ in the first two lines, containing only rational coefficients and single odd zetas, and $H_{\Delta_{12}}^-$ in the last line, which is instead multiplied by the ratio of L-values. Note that the complete Fourier mode $e^{2\pi i \text{Re } \tau}$ receives an infinite series of additional contributions beyond the $q^1 \bar{q}^0$ term in (5.2): All the exponentially suppressed corrections $q(q\bar{q})^n$ for $n > 0$ share the factor of $e^{2\pi i \text{Re } \tau}$ and are multiplied by Laurent polynomials in y with rational coefficients, see section 7.1 of Part I for their precise form.

At the same time, we have found a Poincaré-series representation of $F_{2,5}^{-(6)}$ in Part I, see section 5.2 there. Specialising the general seed formula, recapped in (A.3), to the case at hand

yields

$$\begin{aligned} F_{2,5}^{-(6)}(\tau) &= \sum_{\gamma \in B(\mathbb{Z}) \backslash \mathrm{SL}(2, \mathbb{Z})} \left[-\frac{4iy^5}{31185} \mathrm{Im} \mathcal{E}_0(4, 0; \tau) \right]_{\gamma} \\ &= \sum_{\gamma \in B(\mathbb{Z}) \backslash \mathrm{SL}(2, \mathbb{Z})} \left[\frac{2y^5}{93555} \sum_{n=1}^{\infty} n^{-2} \sigma_3(n) (q^n - \bar{q}^n) \right]_{\gamma}, \end{aligned} \quad (5.3)$$

where the $\mathrm{SL}(2, \mathbb{Z})$ element γ acts on the τ -dependence via both q, \bar{q} and y .

The general formula for obtaining the Fourier series of a Poincaré sum from the Fourier series of its seed function [76, 77], see also appendix A of Part I, then leads to the following identity

$$\begin{aligned} F_{2,5}^{-(6)}(\tau) \Big|_{q^1 \bar{q}^0} + \sum_{n=1}^{\infty} F_{2,5}^{-(6)}(\tau) \Big|_{q^{n+1} \bar{q}^n} &= \frac{2y^5}{93555} + e^{2y} \sum_{d=1}^{\infty} \sum_{\ell \in \mathbb{Z}} S(\ell, 1; d) \\ &\quad \times \int_{\mathbb{R}} e^{-2\pi i \omega - 2\pi i \ell \frac{\omega}{d^2((\mathrm{Im} \tau)^2 + \omega^2)}} c_{\ell} \left(\frac{\mathrm{Im} \tau}{d^2((\mathrm{Im} \tau)^2 + \omega^2)} \right) d\omega, \end{aligned} \quad (5.4)$$

where $c_{\ell}(\mathrm{Im} \tau)$ is the ℓ^{th} Fourier mode of the Poincaré seed in (5.3), i.e. $c_0(\mathrm{Im} \tau) = 0$ and

$$c_{\ell}(\mathrm{Im} \tau) = \frac{2 \mathrm{sign}(\ell)}{93555} |\ell|^{-2} \sigma_3(|\ell|) (\pi \mathrm{Im} \tau)^5 e^{-2\pi |\ell| \mathrm{Im} \tau} \quad \text{for } \ell \neq 0, \quad (5.5)$$

where the $\mathrm{sign}(\ell)$ arises since we are considering an odd modular invariant. Moreover, $S(\ell, 1; d)$ denotes a particular instance of the Kloosterman sum

$$S(\ell, n; d) = \sum_{r \in (\mathbb{Z}/d\mathbb{Z})^{\times}} \exp \left(\frac{2\pi i}{d} \left[\ell r + \frac{n}{r} \right] \right), \quad (5.6)$$

where $0 \leq r \leq d$ is coprime to d , such that r has the multiplicative inverse r^{-1} in $(\mathbb{Z}/d\mathbb{Z})^{\times}$. Note that the above expression (5.4) contains the full $e^{2\pi i \mathrm{Re} \tau}$ Fourier mode sector, i.e. it contains both the $q^1 \bar{q}^0$ term as well as the infinite tower of exponentially suppressed corrections $q(q\bar{q})^n$ for $n > 0$. Since $H_{\Delta_{2s}}^{\pm}$ does not have any $(q\bar{q})^n$ terms, one can restrict to $\check{F}_{m,k}^{\pm(s)}$, and the only sources of $(q\bar{q})^n$ are the depth-two β^{sv} , for which the full $q^{>0} \bar{q}^{>0}$ terms were given in section 7.1 of Part I. All of them are accompanied by Laurent polynomials with rational coefficients.

5.2 Even example

The same kind of analysis can be performed for the q, \bar{q} -expansion (5.1) of the even modular functions $F_{m,k}^{+(s)}$. As an example we can focus on $F_{2,6}^{+(6)}$, where the $q^1 \bar{q}^0$ coefficient is given by

$$\begin{aligned} F_{2,6}^{+(6)} \Big|_{q^1 \bar{q}^0} &= -\frac{691y^5}{3192564375} - \frac{y}{10800} - \frac{5123}{4914000} - \frac{7061}{1310400y} - \frac{6151}{374400y^2} - \frac{23239}{748800y^3} \\ &\quad - \frac{89}{2600y^4} - \frac{89}{5200y^5} + \zeta_3 \left(\frac{1}{480y^2} + \frac{7}{480y^3} + \frac{7}{160y^4} + \frac{21}{320y^5} + \frac{21}{512y^6} \right) \\ &\quad + \frac{21\zeta_{11}}{512y^6} + \frac{\Lambda(\Delta_{12}, 13)}{17275\Lambda(\Delta_{12}, 11)} \left(1 + \frac{15}{2y} + \frac{105}{4y^2} + \frac{105}{2y^3} + \frac{945}{16y^4} + \frac{945}{32y^5} \right) \end{aligned} \quad (5.7)$$

and strongly resembles its odd counterpart (5.2). These two examples (5.2) and (5.7) clearly illustrate the fact that the ratio of L-values $\frac{\Lambda(\Delta_{2s,t_1})}{\Lambda(\Delta_{2s,t_2})}$, appearing in the perturbative expansion of the non-zero Fourier modes, has t_1, t_2 odd for $F_{m,k}^{+(s)}$ as compared to t_1, t_2 even for $F_{m,k}^{-(s)}$.

While the seed functions of the odd $F_{m,k}^{-(s)}$ are series of $q^n - \bar{q}^n$ as in (5.3), the seed functions of the even $F_{m,k}^{+(s)}$ contain $q^n + \bar{q}^n$ and furthermore have zero modes involving \mathbb{Q} multiples of y^{m+k} and $\zeta_{2m-1} y^{k-m+1}$ such as

$$\begin{aligned} F_{2,6}^{+(6)} &= \sum_{\gamma \in B(\mathbb{Z}) \backslash \text{SL}(2, \mathbb{Z})} \left[\frac{691y^8}{373530031875} - \frac{691y^5 \zeta_3}{3192564375} + \frac{1382y^5}{1064188125} \text{Re } \mathcal{E}_0(4, 0^2) \right]_{\gamma} \quad (5.8) \\ &= \sum_{\gamma \in B(\mathbb{Z}) \backslash \text{SL}(2, \mathbb{Z})} \left[\frac{691y^8}{373530031875} - \frac{691y^5}{3192564375} \left(\zeta_3 + \sum_{n=1}^{\infty} n^{-3} \sigma_3(n) (q^n + \bar{q}^n) \right) \right]_{\gamma}, \end{aligned}$$

see section 3.3 of Part I for a construction at generic s, m, k , and (A.1) for a quick recap. We can again employ the relations between the Fourier expansion of seed functions and Poincaré series reviewed in appendix A of Part I to generate examples of how the ratios of L-values as in (5.2) and (5.7) arise from Kloosterman sums and integrals as in (5.4).

5.3 General comments

As we discussed in Part I, the seed functions for the various $F_{m,k}^{\pm(s)}$ can all be written as finite, rational linear combinations of building blocks whose Fourier coefficients of $e^{2\pi i \ell \text{Re } \tau}$ take the simple form

$$c_{\ell}(\text{Im } \tau) \sim \begin{cases} |\ell|^{-r} \sigma_{2m-1}(|\ell|) (\pi \text{Im } \tau)^{m+k-r} e^{-2\pi |\ell| \text{Im } \tau} & : \ell > 0, \\ \pm |\ell|^{-r} \sigma_{2m-1}(|\ell|) (\pi \text{Im } \tau)^{m+k-r} e^{-2\pi |\ell| \text{Im } \tau} & : \ell < 0, \end{cases} \quad (5.9)$$

for integers r in the range $m+1 \leq r \leq 2m-1$ and where the sign \pm for the negative Fourier modes $\ell < 0$ is adapted to the modular function $F_{m,k}^{\pm(s)}$ considered.

In [3] it was explained how to extract the asymptotic expansion for $\text{Im } \tau \rightarrow \infty$ for the Poincaré sum of a seed of the form (5.9) and derive its Laurent polynomial, discussed in full details in Part I, for the zero Fourier mode, i.e. the $(q\bar{q})^0$ term. It is furthermore possible to exploit the asymptotic nature of such an expansion to obtain, via resurgence analysis, the exponentially suppressed terms in the same Fourier mode sector, i.e. the terms $\sim (q\bar{q})^n$ with $n > 0$.

In a similar spirit, we believe that it should be possible, starting from the Poincaré sum of the general seed (5.9), to extract its asymptotic expansion for $\text{Im } \tau \rightarrow \infty$ in any Fourier mode sector. Unlike for the zero-mode sector, no such general expression is at the present time known for (5.9). For example, it would be extremely interesting to start from the expression (5.4) for the first Fourier mode $e^{2\pi i \text{Re } \tau}$ of $F_{2,5}^{-(6)}(\tau)$, or the analogous expression for $F_{2,6}^{+(6)}$ starting from (5.8), and to derive their asymptotic expansions for $\text{Im } \tau \rightarrow \infty$. Similar to what was done in [3] the integral in (5.4) could be done term-wise after expanding the τ -dependent part of the exponential in an absolutely convergent series. This yields multiple, partly divergent, infinite sums over Kloosterman sums. The analytic continuation of these sums is left for future work.

Irrespective of the explicit result, we can still make some predictions. Firstly, due to the presence of this novel, and extremely non-trivial, Kloosterman sum $S(\ell, 1; d)$ (or (5.6) for the generic Fourier mode), we should find that this asymptotic expansion truncates after finitely many terms. More importantly, this asymptotic expansion has either rational numbers, single odd zetas or L-values in its coefficients, reproducing all the $q^1 \bar{q}^0$ terms given in (5.2) and similarly (5.7). Secondly, the asymptotic nature of such an expansion should also hide and encode the presence of an infinite tower of exponentially suppressed corrections, i.e. the $q(q\bar{q})^n$ for $n > 0$, each one of them accompanied by a Laurent polynomial in y with rational coefficients.

Acknowledgements

We are grateful to Francis Brown for his generous help in understanding his results on multiple modular values, providing us with explicit examples beyond the ones published in [50] and valuable comments on an earlier version of this work. We would like to thank Eric D'Hoker, Nikolaos Diamantis, Mehregan Doroudiani, Herbert Gangl, Jan Gerken, Martijn Hidding, Kim Klinger-Logan, Nils Matthes, Stephen D. Miller, Bram Verbeek, and Federico Zerbini for valuable discussions and collaboration on related topics. DD and OS thank the Albert Einstein Institute Potsdam for kind hospitality during several stages of this project and creating a stimulating atmosphere. OS is supported by the European Research Council under ERC-STG-804286 UNISCAMP.

A Recap of seed functions

For the convenience of the reader, in this appendix we summarise the final results of Part I for the seed functions $f_{m,k}^{\pm(s)}$ associated with the modular invariant solutions (1.5) to the Laplace systems (1.1).

In the even case we have the seeds

$$\begin{aligned} f_{m,k}^{+(s)}(\tau) &= (-1)^{k+m} \frac{B_{2k} B_{2m} (4y)^{k+m}}{(2k)!(2m)!(\mu_{k+m} - \mu_s)} - (-1)^k \frac{4B_{2k} (2m-3)! \zeta_{2m-1} (4y)^{k+1-m}}{(2k)!(m-2)!(m-1)!(\mu_{k-m+1} - \mu_s)} \\ &\quad - (-1)^k \frac{2B_{2k} \Gamma(2m)}{(2k)! \Gamma(m)} \sum_{\ell=k-m+1}^{k-1} g_{m,k,\ell,s}^+ (4y)^\ell \operatorname{Re} \mathcal{E}_0(2m, 0^{k+m-\ell-1}), \end{aligned} \quad (\text{A.1})$$

with $\mu_s = s(s-1)$, iterated Eisenstein integrals $\mathcal{E}_0(2m, 0^p)$ defined in (2.8a) and rational coefficients $g_{m,k,\ell,s}^+$ given by

$$g_{m,k,\ell,s}^+ = \frac{\Gamma(\ell)}{\Gamma(\ell+s)} \sum_{i=\ell}^{k-1} \frac{(\ell+1-s)_{i-\ell} \Gamma(s+i) \Gamma(m+k-i-1)}{\Gamma(k-i) \Gamma(i+1) \Gamma(m-k+i+1)}. \quad (\text{A.2})$$

Similarly, in the odd case we have the seeds

$$f_{m,k}^{-(s)} = i(-1)^k \frac{B_{2k} (2m-1)!}{2\Gamma(2k) \Gamma(m)} \sum_{\ell=k-m+1}^k g_{m,k,\ell,s}^- (4y)^\ell \operatorname{Im} \mathcal{E}_0(2m, 0^{k+m-\ell-1}) \quad (\text{A.3})$$

with rational coefficients $g_{m,k,\ell,s}^-$ given by

$$g_{m,k,\ell,s}^- = \frac{\Gamma(\ell)}{\Gamma(\ell+s)} \sum_{i=\ell}^k \frac{(\ell+1-s)_{i-\ell} \Gamma(s+i) \Gamma(m+k-i)}{\Gamma(k-i+1) \Gamma(i+1) \Gamma(m-k+i)}, \quad (\text{A.4})$$

see sections 3.3 and 5.2 of Part I for the derivation of (A.1) and (A.3), respectively.

B Examples of cocycles

In this appendix, we spell out further examples of the cocycles (2.39) of the solutions $H_{\Delta_{2s}}^\pm$ to homogeneous Laplace equations. In the ancillary file, all cocycles up to $2s = 26$ are given along with the ratios of the completed L-values inside the critical strip.

B.1 Weight 16

For the cusp form of weight 16 in the Hecke normalisation $\Delta_{16} = q + O(q^2)$, the ratios of L-values

$$\begin{aligned} \Lambda(\Delta_{16}, 8) : \Lambda(\Delta_{16}, 10) : \Lambda(\Delta_{16}, 12) : \Lambda(\Delta_{16}, 14) &= \frac{35}{468} : \frac{49}{468} : \frac{245}{936} : 1, \\ \Lambda(\Delta_{16}, 9) : \Lambda(\Delta_{16}, 11) : \Lambda(\Delta_{16}, 13) : \Lambda(\Delta_{16}, 15) &= \frac{3617}{98280} : \frac{3617}{51480} : \frac{3617}{16380} : 1, \end{aligned} \quad (\text{B.1})$$

and similar ratios $\Lambda(\Delta_{16}, t) : \Lambda(\Delta_{16}, 14)$ and $\Lambda(\Delta_{16}, t) : \Lambda(\Delta_{16}, 15)$ with $t < 8$ following from the reflection formula (2.37) lead to the cocycles

$$\begin{aligned} H_{\Delta_{16}}^+(\tau) - H_{\Delta_{16}}^+\left(-\frac{1}{\tau}\right) &= \frac{\pi^{15} \Lambda(\Delta_{16}, 15) (1-\tau\bar{\tau})}{2520y^7} \left\{ 1 + \tau^6\bar{\tau}^6 - \frac{3617}{780} (\tau^2 + \bar{\tau}^2) (1 + \tau^4\bar{\tau}^4) \right. \\ &\quad + \frac{25319}{10296} (1 + \tau^2\bar{\tau}^2) (\tau^4 + \bar{\tau}^4) - \frac{3617}{14040} (\tau^6 + \bar{\tau}^6) - \frac{22979}{2340} \tau\bar{\tau} (1 + \tau^4\bar{\tau}^4) \\ &\quad + \frac{647443}{51480} \tau\bar{\tau} (\tau^2 + \bar{\tau}^2) (1 + \tau^2\bar{\tau}^2) - \frac{25319}{8580} \tau\bar{\tau} (\tau^4 + \bar{\tau}^4) - \frac{372551}{25740} \tau^2\bar{\tau}^2 (\tau^2 + \bar{\tau}^2) \\ &\quad \left. + \frac{1089559}{51480} \tau^2\bar{\tau}^2 (1 + \tau^2\bar{\tau}^2) - \frac{461756}{19305} \tau^3\bar{\tau}^3 \right\}, \end{aligned} \quad (\text{B.2a})$$

$$\begin{aligned} H_{\Delta_{16}}^-(\tau) - H_{\Delta_{16}}^-\left(-\frac{1}{\tau}\right) &= \frac{i\pi^{15} \Lambda(\Delta_{16}, 14) (\tau + \bar{\tau})}{360y^7} \left\{ 1 + \tau^6\bar{\tau}^6 - \frac{1225}{936} (\tau^2 + \bar{\tau}^2) (1 + \tau^4\bar{\tau}^4) \right. \\ &\quad + \frac{49}{156} (1 + \tau^2\bar{\tau}^2) (\tau^4 + \bar{\tau}^4) - \frac{5}{468} (\tau^6 + \bar{\tau}^6) - \frac{490}{117} \tau\bar{\tau} (1 + \tau^4\bar{\tau}^4) \\ &\quad + \frac{392}{117} \tau\bar{\tau} (\tau^2 + \bar{\tau}^2) (1 + \tau^2\bar{\tau}^2) - \frac{20}{39} (\tau^5\bar{\tau} + \tau\bar{\tau}^5) + \frac{3577}{468} \tau^2\bar{\tau}^2 (1 + \tau^2\bar{\tau}^2) \\ &\quad \left. - \frac{655}{156} \tau^2\bar{\tau}^2 (\tau^2 + \bar{\tau}^2) - \frac{80}{9} \tau^3\bar{\tau}^3 \right\}. \end{aligned} \quad (\text{B.2b})$$

B.2 Weight 18

For the cusp form of weight 18 in the Hecke normalisation $\Delta_{18} = q + O(q^2)$, the ratios of L-values

$$\begin{aligned} \Lambda(\Delta_{18}, 10) : \Lambda(\Delta_{18}, 12) : \Lambda(\Delta_{18}, 14) : \Lambda(\Delta_{18}, 16) &= \frac{1}{120} : \frac{1}{24} : \frac{11}{60} : 1, \\ \Lambda(\Delta_{18}, 11) : \Lambda(\Delta_{18}, 13) : \Lambda(\Delta_{18}, 15) : \Lambda(\Delta_{18}, 17) &= \frac{43867}{5544000} : \frac{43867}{1310400} : \frac{43867}{270000} : 1, \end{aligned} \quad (\text{B.3})$$

along with the remaining values, including $\Lambda(\Delta_{18}, 9) = 0$, due to (2.37) lead to the cocycles

$$\begin{aligned} H_{\Delta_{18}}^+(\tau) - H_{\Delta_{18}}^+\left(-\frac{1}{\tau}\right) &= \frac{\pi^{17}\Lambda(\Delta_{18}, 17)(1-\tau^2\bar{\tau}^2)}{20160y^8} \left\{ 1 + \tau^6\bar{\tau}^6 - \frac{307069}{67500}(\tau^2+\bar{\tau}^2)(1+\tau^4\bar{\tau}^4) \right. \\ &\quad + \frac{43867}{18720}(1+\tau^2\bar{\tau}^2)(\tau^4+\bar{\tau}^4) - \frac{43867}{198000}(\tau^6+\bar{\tau}^6) - \frac{175468}{16875}\tau\bar{\tau}(1+\tau^4\bar{\tau}^4) \\ &\quad + \frac{43867}{2925}\tau\bar{\tau}(\tau^2+\bar{\tau}^2)(1+\tau^2\bar{\tau}^2) - \frac{43867}{12375}\tau\bar{\tau}(\tau^4+\bar{\tau}^4) + \frac{318769}{11700}\tau^2\bar{\tau}^2(1+\tau^2\bar{\tau}^2) \\ &\quad \left. - \frac{29785693}{1485000}\tau^2\bar{\tau}^2(\tau^2+\bar{\tau}^2) - \frac{6536183}{185625}\tau^3\bar{\tau}^3 \right\}, \end{aligned} \quad (\text{B.4a})$$

$$\begin{aligned} H_{\Delta_{18}}^-(\tau) - H_{\Delta_{18}}^-\left(-\frac{1}{\tau}\right) &= \frac{i\pi^{17}\Lambda(\Delta_{18}, 16)(\tau+\bar{\tau})(1+\tau\bar{\tau})}{2520y^8} \left\{ 1 + \tau^6\bar{\tau}^6 - \frac{77}{60}(\tau^2+\bar{\tau}^2)(1+\tau^4\bar{\tau}^4) \right. \\ &\quad + \frac{7}{24}(1+\tau^2\bar{\tau}^2)(\tau^4+\bar{\tau}^4) - \frac{97}{20}\tau\bar{\tau}(1+\tau^4\bar{\tau}^4) + \frac{469}{120}\tau\bar{\tau}(\tau^2+\bar{\tau}^2)(1+\tau^2\bar{\tau}^2) \\ &\quad - \frac{31}{60}\tau\bar{\tau}(\tau^4+\bar{\tau}^4) + \frac{1247}{120}\tau^2\bar{\tau}^2(1+\tau^2\bar{\tau}^2) - \frac{319}{60}\tau^2\bar{\tau}^2(\tau^2+\bar{\tau}^2) \\ &\quad \left. - \frac{1}{120}(\tau^6+\bar{\tau}^6) - \frac{196}{15}\tau^3\bar{\tau}^3 \right\}. \end{aligned} \quad (\text{B.4b})$$

References

- [1] D. Dorigoni, A. Kleinschmidt, and O. Schlotterer, ‘‘Poincaré series for modular graph forms at depth two. I. Seeds and Laplace system,’’ [arXiv:2109.05017 \[hep-th\]](#).
- [2] O. Ahlén and A. Kleinschmidt, ‘‘ D^6R^4 curvature corrections, modular graph functions and Poincaré series,’’ *JHEP* **05** (2018) 194, [arXiv:1803.10250 \[hep-th\]](#).
- [3] D. Dorigoni and A. Kleinschmidt, ‘‘Modular graph functions and asymptotic expansions of Poincaré series,’’ *Commun. Num. Theor. Phys.* **13** (2019) no. 3, 569–617, [arXiv:1903.09250 \[hep-th\]](#).
- [4] D. Dorigoni and A. Kleinschmidt, ‘‘Resurgent expansion of Lambert series and iterated Eisenstein integrals,’’ *Commun. Num. Theor. Phys.* **15** (2021) no. 1, 1–57, [arXiv:2001.11035 \[hep-th\]](#).
- [5] A. Basu, ‘‘Zero mode of the Fourier series of some modular graphs from Poincaré series,’’ *Phys. Lett. B* **809** (2020) 135715, [arXiv:2005.07793 \[hep-th\]](#).

- [6] E. D'Hoker, M. B. Green, and P. Vanhove, "On the modular structure of the genus-one Type II superstring low energy expansion," *JHEP* **08** (2015) 041, [arXiv:1502.06698 \[hep-th\]](#).
- [7] E. D'Hoker, M. B. Green, Ö. Gürdoğan, and P. Vanhove, "Modular graph functions," *Commun. Num. Theor. Phys.* **11** (2017) no. 1, 165–218, [arXiv:1512.06779 \[hep-th\]](#).
- [8] E. D'Hoker and M. B. Green, "Identities between modular graph forms," *J. Number Theory* **189** (2018) 25–80, [arXiv:1603.00839 \[hep-th\]](#).
- [9] M. B. Green and P. Vanhove, "The Low-energy expansion of the one loop type II superstring amplitude," *Phys.Rev.* **D61** (2000) 104011, [arXiv:hep-th/9910056 \[hep-th\]](#).
- [10] M. B. Green, J. G. Russo, and P. Vanhove, "Low energy expansion of the four-particle genus-one amplitude in type II superstring theory," *JHEP* **02** (2008) 020, [arXiv:0801.0322 \[hep-th\]](#).
- [11] M. B. Green, C. R. Mafra, and O. Schlotterer, "Multiparticle one-loop amplitudes and S-duality in closed superstring theory," *JHEP* **10** (2013) 188, [arXiv:1307.3534 \[hep-th\]](#).
- [12] E. D'Hoker, M. B. Green, and P. Vanhove, "Proof of a modular relation between 1-, 2- and 3-loop Feynman diagrams on a torus," *J. Number Theory* (2018) 381, [arXiv:1509.00363 \[hep-th\]](#).
- [13] A. Basu, "Poisson equation for the Mercedes diagram in string theory at genus one," *Class. Quant. Grav.* **33** (2016) no. 5, 055005, [arXiv:1511.07455 \[hep-th\]](#).
- [14] A. Basu, "Poisson equation for the three loop ladder diagram in string theory at genus one," *Int. J. Mod. Phys. A* **31** (2016) no. 32, 1650169, [arXiv:1606.02203 \[hep-th\]](#).
- [15] A. Basu, "Proving relations between modular graph functions," *Class. Quant. Grav.* **33** (2016) no. 23, 235011, [arXiv:1606.07084 \[hep-th\]](#).
- [16] A. Basu, "Simplifying the one loop five graviton amplitude in type IIB string theory," *Int. J. Mod. Phys. A* **32** (2017) no. 14, 1750074, [arXiv:1608.02056 \[hep-th\]](#).
- [17] E. D'Hoker and J. Kaidi, "Hierarchy of Modular Graph Identities," *JHEP* **11** (2016) 051, [arXiv:1608.04393 \[hep-th\]](#).
- [18] A. Kleinschmidt and V. Verschinin, "Tetrahedral modular graph functions," *JHEP* **09** (2017) 155, [arXiv:1706.01889 \[hep-th\]](#).
- [19] A. Basu, "Low momentum expansion of one loop amplitudes in heterotic string theory," *JHEP* **11** (2017) 139, [arXiv:1708.08409 \[hep-th\]](#).

- [20] J. Broedel, O. Schlotterer, and F. Zerbini, “From elliptic multiple zeta values to modular graph functions: open and closed strings at one loop,” *JHEP* **01** (2019) 155, [arXiv:1803.00527 \[hep-th\]](#).
- [21] J. E. Gerken and J. Kaidi, “Holomorphic subgraph reduction of higher-point modular graph forms,” *JHEP* **01** (2019) 131, [arXiv:1809.05122 \[hep-th\]](#).
- [22] J. E. Gerken, A. Kleinschmidt, and O. Schlotterer, “Heterotic-string amplitudes at one loop: modular graph forms and relations to open strings,” *JHEP* **01** (2019) 052, [arXiv:1811.02548 \[hep-th\]](#).
- [23] E. D’Hoker and J. Kaidi, “Modular graph functions and odd cuspidal functions. Fourier and Poincaré series,” *JHEP* **04** (2019) 136, [arXiv:1902.04180 \[hep-th\]](#).
- [24] E. D’Hoker and M. B. Green, “Absence of irreducible multiple zeta-values in melon modular graph functions,” *Commun. Num. Theor. Phys.* **14** (2020) no. 2, 315–324, [arXiv:1904.06603 \[hep-th\]](#).
- [25] E. D’Hoker, “Integral of two-loop modular graph functions,” *JHEP* **06** (2019) 092, [arXiv:1905.06217 \[hep-th\]](#).
- [26] E. D’Hoker and M. B. Green, “Exploring transcendentality in superstring amplitudes,” *JHEP* **07** (2019) 149, [arXiv:1906.01652 \[hep-th\]](#).
- [27] A. Basu, “Eigenvalue equation for the modular graph $C_{a,b,c,d}$,” *JHEP* **07** (2019) 126, [arXiv:1906.02674 \[hep-th\]](#).
- [28] J. E. Gerken, A. Kleinschmidt, and O. Schlotterer, “All-order differential equations for one-loop closed-string integrals and modular graph forms,” *JHEP* **01** (2020) 064, [arXiv:1911.03476 \[hep-th\]](#).
- [29] S. Hohenegger, “From Little String Free Energies Towards Modular Graph Functions,” *JHEP* **03** (2020) 077, [arXiv:1911.08172 \[hep-th\]](#).
- [30] J. E. Gerken, A. Kleinschmidt, and O. Schlotterer, “Generating series of all modular graph forms from iterated Eisenstein integrals,” *JHEP* **07** (2020) 190, [arXiv:2004.05156 \[hep-th\]](#).
- [31] P. Vanhove and F. Zerbini, “Building blocks of closed and open string amplitudes,” in *MathemAmplitudes 2019: Intersection Theory and Feynman Integrals*. 7, 2020. [arXiv:2007.08981 \[hep-th\]](#).
- [32] A. Basu, “Poisson equations for elliptic modular graph functions,” *Phys. Lett. B* **814** (2021) 136086, [arXiv:2009.02221 \[hep-th\]](#).
- [33] A. Basu, “Relations between elliptic modular graphs,” *JHEP* **12** (2020) 195, [arXiv:2010.08331 \[hep-th\]](#). [Erratum: *JHEP* **03**, 061 (2021)].

- [34] S. Hohenegger, “Diagrammatic Expansion of Non-Perturbative Little String Free Energies,” *JHEP* **04** (2021) 275, [arXiv:2011.06323 \[hep-th\]](#).
- [35] F. Brown, “Multiple modular values and the relative completion of the fundamental group of $\mathcal{M}_{1,1}$,” [arXiv:1407.5167 \[math.NT\]](#).
- [36] F. Zerbini, “Single-valued multiple zeta values in genus 1 superstring amplitudes,” *Commun. Num. Theor. Phys.* **10** (2016) no. 4, 703–737, [arXiv:1512.05689 \[hep-th\]](#).
- [37] F. Brown, “A class of non-holomorphic modular forms I,” *Res. Math. Sci.* **5** (2018) no. 1, Paper No. 7, 40, [arXiv:1707.01230 \[math.NT\]](#).
- [38] F. Brown, “A class of nonholomorphic modular forms II: Equivariant iterated Eisenstein integrals,” *Forum Math. Sigma* **8** (2020) Paper No. e31, 62, [arXiv:1708.03354 \[math.NT\]](#).
- [39] E. D’Hoker and W. Duke, “Fourier series of modular graph functions,” *J. Number Theory* **192** (2018) 1–36, [arXiv:1708.07998 \[math.NT\]](#).
- [40] F. Zerbini, *Elliptic multiple zeta values, modular graph functions and genus 1 superstring scattering amplitudes*. PhD thesis, Bonn U., 2017. [arXiv:1804.07989 \[math-ph\]](#).
- [41] F. Zerbini, “Modular and holomorphic graph function from superstring amplitudes,” in *KMPB Conference: Elliptic Integrals, Elliptic Functions and Modular Forms in Quantum Field Theory Zeuthen, Germany, October 23-26, 2017*. 2018. [arXiv:1807.04506 \[math-ph\]](#).
- [42] D. Zagier and F. Zerbini, “Genus-zero and genus-one string amplitudes and special multiple zeta values,” *Commun. Num. Theor. Phys.* **14** (2020) no. 2, 413–452, [arXiv:1906.12339 \[math.NT\]](#).
- [43] M. Berg, K. Bringmann, and T. Gannon, “Massive deformations of Maass forms and Jacobi forms,” *Commun. Num. Theor. Phys.* **15** (2021) no. 3, 575–603, [arXiv:1910.02745 \[math.NT\]](#).
- [44] N. Matthes, “On the algebraic structure of iterated integrals of quasimodular forms,” *Algebra & Number Theory* **11-9** (2017) 2113–2130, [arXiv:1708.04561](#).
- [45] M. Eichler, “Eine Verallgemeinerung der Abelschen Integrale,” *Math. Z.* **67** (1957) 267–298.
- [46] G. Shimura, “Sur les intégrales attachées aux formes automorphes,” *J. Math. Soc. Japan* **11** (1959) 291–311.
- [47] F. Brown, “A class of non-holomorphic modular forms III: real analytic cusp forms for $SL_2(\mathbb{Z})$,” *Res. Math. Sci.* **5** (2018) no. 3, Paper No. 34, 36, [arXiv:1710.07912 \[math.NT\]](#).

- [48] H. Tsunogai, “On some derivations of Lie algebras related to Galois representations,” *Publ. Res. Inst. Math. Sci.* **31** (1995) no. 1, 113–134.
- [49] A. Pollack, “Relations between derivations arising from modular forms.” <https://dukespace.lib.duke.edu/dspace/handle/10161/1281>, 2009. Undergraduate thesis, Duke University.
- [50] F. Brown, “From the Deligne-Ihara conjecture to multiple modular values,” [arXiv:1904.00179](https://arxiv.org/abs/1904.00179) [math.AG].
- [51] F. Brown, “Talk *Zeta and L-functions, old and new* given at the workshop *Elliptics and beyond*.” <https://indico.cern.ch/event/927781/contributions/3926291/attachments/2101487/3533016/EllipticConf.pdf>, 2020.
- [52] N. Diamantis, “Modular iterated integrals associated with cusp forms,” [arXiv:2009.07128](https://arxiv.org/abs/2009.07128) [math.NT].
- [53] J. Drewitt, “Triple equivariant eisenstein integrals,” [arXiv:2104.09916](https://arxiv.org/abs/2104.09916) [math.NT].
- [54] J. Broedel, N. Matthes, and O. Schlotterer, “Relations between elliptic multiple zeta values and a special derivation algebra,” *J. Phys.* **A49** (2016) no. 15, 155203, [arXiv:1507.02254](https://arxiv.org/abs/1507.02254) [hep-th].
- [55] D. Zagier and H. Gangl, “Classical and elliptic polylogarithms and special values of L -series,” in *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, vol. 548 of *NATO Sci. Ser. C Math. Phys. Sci.*, pp. 561–615. Kluwer Acad. Publ., Dordrecht, 2000.
- [56] D. Zagier, “Periods of modular forms and Jacobi theta functions,” *Invent. Math.* **104** (1991) 449–465.
- [57] D. B. Zagier, “Quelques conséquences surprenantes de la cohomologie de $SL(2, \mathbb{Z})$,” *Leçons de Mathématiques d’aujourd’hui (Cassini, Paris)* (2000) 99–123. <http://people.mpim-bonn.mpg.de/zagier/files/tex/ConsequencesCohomologySL/fulltext.pdf>.
- [58] E. Hecke, “Über Modulfunktionen und die Dirichletschen Reihen mit Eulerscher Produktentwicklung. I,” *Math. Ann.* **114** (1937) no. 1, 1–28.
- [59] P. Deligne, “La conjecture de Weil. I,” *Inst. Hautes Études Sci. Publ. Math.* (1974) no. 43, 273–307.
- [60] J. I. Manin, “Periods of parabolic forms and p -adic hecke series,” *Mathematics of the USSR-Sbornik* **21** (1973) 371–393.
- [61] A. Saad, “Multiple zeta values and iterated Eisenstein integrals,” [arXiv:2009.09885](https://arxiv.org/abs/2009.09885) [math.NT].
- [62] F. Brown, private communication.

- [63] The PARI Group, Univ. Bordeaux, *PARI/GP version 2.13.1*, 2020.
<http://pari.math.u-bordeaux.fr/>.
- [64] T. Dokchitser, *ComputeL - PARI package v1.3.8*, 2018.
<https://people.maths.bris.ac.uk/~matyd/computel/>.
- [65] M. B. Green, S. D. Miller, and P. Vanhove, “ $SL(2, \mathbb{Z})$ -invariance and D-instanton contributions to the D^6R^4 interaction,” *Commun. Num. Theor. Phys.* **09** (2015) 307–344, [arXiv:1404.2192](https://arxiv.org/abs/1404.2192) [hep-th].
- [66] K. Klinger-Logan, “Differential equations in automorphic forms,” *Commun. Number Theory Phys.* **12** (2018) no. 4, 767–827, [arXiv:1801.00838](https://arxiv.org/abs/1801.00838) [math.NT].
- [67] K. Klinger-Logan, S. D. Miller, and D. Radchenko, “The $D^6\mathcal{R}^4$ coupling, revisited,” work in progress.
- [68] T. M. Apostol, *Modular functions and Dirichlet series in number theory*. Graduate texts in mathematics; 41. Springer-Verlag, New York, 1976.
- [69] K. Haberland, “Perioden von Modulformen einer Variabler and Gruppencohomologie. I, II, III,” *Math. Nachr.* **112** (1983) 245–282, 283–295, 297–315.
- [70] H. Hida and Y. Maeda, “Non-abelian base change for totally real fields,” *Pacific J. Math.* (1997) 189–217. (Special Issue in memoriam of Olga Taussky-Todd).
- [71] A. Ghitza and A. McAndrew, “Experimental evidence for Maeda’s conjecture on modular forms,” *Tbilisi Mathematical Journal* **5** (2012) no. 2, 55 – 69.
- [72] The LMFDB Collaboration, “The L-functions and modular forms database.”
<http://www.lmfdb.org>, 2021.
- [73] J.-G. Luque, J.-C. Novelli, and J.-Y. Thibon, “Period polynomials and Ihara brackets,” *J. Lie Theory* **17** (2007) 229–239, [arXiv:math/0606301](https://arxiv.org/abs/math/0606301) [math.CO,math.NT].
- [74] J. Broedel, N. Matthes, and O. Schlotterer. <https://tools.aei.mpg.de/emzv>.
- [75] B. Enriquez, “Analogues elliptiques des nombres multizétas,” *Bull. Soc. Math. France* **144** (2016) no. 3, 395–427, [arXiv:1301.3042](https://arxiv.org/abs/1301.3042) [math.NT].
- [76] H. Iwaniec, *Spectral methods of automorphic forms*, vol. 53 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI; Revista Matemática Iberoamericana, Madrid, second ed., 2002.
- [77] P. Fleig, H. P. A. Gustafsson, A. Kleinschmidt, and D. Persson, *Eisenstein series and automorphic representations, with applications in string theory*, vol. 176 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2018.
[arXiv:1511.04265](https://arxiv.org/abs/1511.04265) [math.NT].