

GAMES CHARACTERIZING CERTAIN FAMILIES OF FUNCTIONS

MAREK BALCERZAK, TOMASZ NATKANIEC, AND PIOTR SZUCA

ABSTRACT. We obtain several game characterizations of Baire 1 functions between Polish spaces X, Y which extends the recent result of V. Kiss. Then we propose similar characterizations for equi-Baire 1 families of functions. Also, using related ideas, we give game characterizations of Baire measurable and Lebesgue measurable functions.

1. INTRODUCTION

The game approach plays an important role in descriptive set theory. Let us recall Choquet games and the Banach-Mazur game in the studies of the Baire category problems [12, Sec. 8], and Wadge games with their influence on investigations in the Borel hierarchy [12, Sec. 21]. It is commonly known that Borel and projective determinacy provide a strong tool in set-theoretical investigations, cf. [12, Sec. 20, 38]. Note that various kinds of topological games make fruitful inspirations in topology and analysis, cf. [3], [10], [5]. They can distinguish new kinds of topological objects, cf. [10], [3].

In the recent decades, several nice characterizations for some classes of regular functions were obtained. Duparc [7] and Carroy [6] characterized Baire 1 functions from $\mathbb{N}^{\mathbb{N}}$ into itself by using the so-called eraser game (for more applications of this game, see [4]). Other significant results for different classes of functions between Polish zero-dimensional Polish spaces are due to Andretta [2] (a game characterization of Δ_2^0 -measurable functions), Semmes [18] (Borel functions), Nobrega [16] (Baire class ξ functions) and Ros [17] (piecewise defined functions).

Recently, Kiss [13] introduced a game characterizing Baire class 1 functions between arbitrary two Polish spaces. This improved the results by Duparc [7] and Carroy [6] that have been mentioned above. Another idea characterizing Baire 1, real-valued functions, has been presented in [8].

Our first aim in this paper is to extend the result by Kiss. We simplify the proof of a harder implication of his result by the use of ε - δ characterization of Baire 1 functions. Then we modify the game defined by Kiss in two other manners which again yield characterizations of Baire 1 class. One of these manners uses a point-open game. The both manners can be modified to obtain the respective characterizations of equi-Baire 1 families. Our second aim is to characterize, by a modification of point-open games, families of Baire measurable and Lebesgue measurable functions.

2010 *Mathematics Subject Classification*. Primary: 03E15; Secondary: 03E60, 26A21, 28A05, 54H05, 91A44.

Key words and phrases. topological game, Baire 1 functions, equi-Baire 1 families, ε -gauge, equi-continuity, measurable functions.

Assume that X and Y are Polish spaces. Through the paper, we assume that d_X and d_Y are the respective metrics in X and Y .

Let us state preliminary facts on Baire 1 functions. A function $f: X \rightarrow Y$ between Polish spaces X, Y is called *Baire class 1* whenever the preimage $f^{-1}[U]$ is F_σ in X for any open set U in Y . If $Y = \mathbb{R}$, this is equivalent to the property that f is the limit of a pointwise convergent sequence of continuous functions, see e.g. [12, Theorem 24.10].

In the literature, we encounter various equivalent definitions of Baire 1 functions. The classical characterization given by Baire says that f is Baire 1 if and only if $f \upharpoonright P$ has a point of continuity for every non-empty closed set $P \subseteq X$. This is the so-called *Pointwise Continuity Property*, in short (PCP). An ε - δ characterization of Baire 1 functions, obtained in [15], says the following. A function $f: X \rightarrow Y$ is Baire 1 whenever, for any positive number ε , there is a positive function $\delta_\varepsilon: X \rightarrow \mathbb{R}$ such that for any $x_0, x_1 \in X$,

$$(1) \quad d_X(x_0, x_1) < \min \{\delta_\varepsilon(x_0), \delta_\varepsilon(x_1)\} \text{ implies } d_Y(f(x_0), f(x_1)) < \varepsilon.$$

We will call such a δ_ε an ε -*gauge* for f .

We say that a family $\mathcal{F} \subseteq Y^X$ is equi-continuous at a point $x \in X$ whenever

$$\forall \varepsilon > 0 \exists \delta > 0 \forall f \in \mathcal{F} (d_X(x, x_0) < \delta \Rightarrow d_Y(f(x), f(x_0)) < \varepsilon).$$

\mathcal{F} is equi-continuous if it is equi-continuous at every $x \in X$.

A family $\mathcal{F} \subseteq Y^X$ is said to fulfil the *Point of Equicontinuity Property* (\mathcal{F} has (PECP), in short) if for every non-empty closed $P \subseteq X$, the family

$$\mathcal{F} \upharpoonright P := \{f \upharpoonright P: f \in \mathcal{F}\}$$

has a point of equicontinuity.

We say that a family $\mathcal{F} \subseteq Y^X$ is *equi-Baire 1* if for any positive number ε there is a positive function $\delta_\varepsilon: X \rightarrow \mathbb{R}_+$ such that for any $x_0, x_1 \in X$ and $f \in \mathcal{F}$ the condition (1) holds (i.e. all $f \in \mathcal{F}$ have a common family of ε -gauges). Clearly, every equi-continuous family is equi-Baire 1 and has (PECP), and the opposite implications do not hold.

The both definitions were introduced by D. Lecomte in [14]. He proved the following equivalence.

Theorem 1 ([14, Prop. 32]). *\mathcal{F} has (PECP) if and only if \mathcal{F} is equi-Baire 1.*

Let us mention that, in (PCP) and (PECP), a non-empty closed set can be equivalently replaced by a perfect set (that is, a non-empty closed set without isolated points).

Note that the definition of equi-Baire 1 family of functions was rediscovered later by A. Alikhani-Koopaei in [1]. The definition of families with (PECP) was used by E. Glasner and M. Megrelishvili in the context of dynamical systems in [9] (under the name “barely continuous family”).

2. GAME CHARACTERIZATIONS OF BAIRE 1 FUNCTIONS

Recall the game defined by Kiss [13]. Let X and Y be Polish spaces. Let $f: X \rightarrow Y$ be an arbitrary function. At the n th step of the game G_f , Player I plays x_n , then Player II plays y_n ,

$$\begin{array}{rcccccccc} \text{Player I} & x_0 & & x_1 & & x_2 & & \cdots \\ \text{Player II} & & y_0 & & y_1 & & y_2 & \cdots \end{array}$$

with the rules that for each $n \in \mathbb{N}$:

- $x_n \in X$ and $d_X(x_n, x_{n+1}) \leq 2^{-n}$;
- $y_n \in Y$.

Since X is complete, $x_n \rightarrow x$ for some $x \in X$. Player II wins if and only if $\langle y_n \rangle_{n \in \mathbb{N}}$ is convergent and $y_n \rightarrow f(x)$. Recall the main result of Kiss:

Theorem 2 ([13, Theorem 1]). *The game G_f is determined, and*

- *Player I has a winning strategy in G_f if and only if f is not of Baire class 1.*
- *Player II has a winning strategy in G_f if and only if f is of Baire class 1.*

The longest part of the original proof is the part “if f is of Baire class 1 then Player II has a winning strategy”. We show that it can be significantly shortened by the use of ε - δ characterization of Baire 1 functions. We describe it in Lemma 4 which will be preceded by the following fact.

Lemma 3. *A function $f: X \rightarrow Y$ is Baire 1 if and only if it possesses a family of gauges $\{\delta_\varepsilon: \varepsilon > 0\}$ such that for every $x \in X$ the map $\varepsilon \mapsto \delta_\varepsilon(x)$ is non-decreasing.*

Proof. Only the implication “ \Rightarrow ” has to be proved. Assume that f is Baire 1 and $\{\delta_\varepsilon: \varepsilon > 0\}$ is a family of gauges for f . For every $\varepsilon > 0$ fix $N_\varepsilon \in \mathbb{N}$ such that $N_\varepsilon = 1$ if $\varepsilon \geq 1$ and $\frac{1}{N_\varepsilon} \leq \varepsilon < \frac{1}{N_\varepsilon - 1}$ for $\varepsilon < 1$. For $x \in X$ define

$$\delta'_\varepsilon(x) := \min \left\{ \delta_{\frac{1}{n}}(x) : n \leq N_\varepsilon \right\}.$$

Clearly, if $\varepsilon \leq \varepsilon_1$ then $N_\varepsilon \geq N_{\varepsilon_1}$, hence for any $x \in X$ we have $\delta'_\varepsilon(x) \leq \delta'_{\varepsilon_1}(x)$. We will show that $\{\delta'_\varepsilon: \varepsilon > 0\}$ is a family of gauges for f . Indeed, assume that $d_X(x_0, x_1) < \min(\delta'_\varepsilon(x_0), \delta'_\varepsilon(x_1))$ for some $\varepsilon > 0$ and $x_0, x_1 \in X$. Then $\frac{1}{N_\varepsilon} \leq \varepsilon$ and $\delta'_\varepsilon(x_i) \leq \delta_{\frac{1}{N_\varepsilon}}(x_i)$ for $i = 0, 1$, so $d_X(x_0, x_1) < \min(\delta_{\frac{1}{N_\varepsilon}}(x_0), \delta_{\frac{1}{N_\varepsilon}}(x_1))$, hence $d_Y(f(x_0), f(x_1)) < \frac{1}{N_\varepsilon} \leq \varepsilon$. \square

Lemma 4. *Let $\Delta := \{\delta_\varepsilon: \varepsilon > 0\}$ be a family of positive functions from X into \mathbb{R} such that, for every $x \in X$, the map $\varepsilon \mapsto \delta_\varepsilon(x)$ is non-decreasing. Then there is a function $\mathcal{S}'_\Delta: X^{<\omega} \rightarrow X$ such that for each sequence $\langle x_n \rangle$ with $d_X(x_n, x_{n+1}) \leq 2^{-n}$ for each n , and for every $\varepsilon > 0$ there exists $N_\varepsilon \in \mathbb{N}$ with the property*

- *for every Baire 1 function $f: X \rightarrow Y$, if Δ is a family of ε -gauges for f , then*

$$\forall n > N_\varepsilon \quad d_Y \left(f(\mathcal{S}'(x_0, x_1, \dots, x_n)), f \left(\lim_{n \rightarrow \infty} x_n \right) \right) < \varepsilon.$$

In particular, $\lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n)$, so the function $\mathcal{S} := f \circ \mathcal{S}'_\Delta$ is a winning strategy for Player II in the game G_f .

Proof. Fix $\varepsilon > 0$ and a sequence $\langle x_n \rangle \subseteq X$ such that $d_X(x_n, x_{n+1}) \leq 2^{-n}$ for each n . We may assume that, for each $n \in \mathbb{N}$, Player I plays x_n in the n th move of the game G_f . For each $n \in \mathbb{N}$ let $K_n := \overline{B}(x_n, 2^{-n+1})$ be the closed ball around x_n . Note that this is the smallest closed ball around x_n which ensures that $x := \lim_{j \rightarrow \infty} x_j \in K_n$. Denote by M_n the greatest index $m < n$ for which there exists a point $x' \in K_n$ such that $K_n \subseteq B(x', \delta_{\frac{1}{m}}(x'))$; then pick one of them and call it x'_n . If such an index m does not exist, put $M_n := -\infty$. Define

$$\mathcal{S}'_\Delta(x_0, \dots, x_n) := \begin{cases} x'_n & \text{if } M_n > -\infty, \\ x_n & \text{otherwise.} \end{cases}$$

It is enough to show that $\lim_{n \rightarrow \infty} f(\mathcal{S}'_{\Delta}(x_0, \dots, x_n)) = f(x)$ for each Baire 1 function $f: X \rightarrow Y$ with the family of ε -gauges equal to Δ . Fix $\varepsilon > 0$ and find a positive integer M such that $1/M < \varepsilon$. There exists $N \in \mathbb{N}$ such that for each $n \geq N$,

$$(2) \quad x \in K_n \subseteq B\left(x, \delta_{\frac{1}{M}}(x)\right).$$

Since $x \in K_n$ for all n , it follows that $M_n \geq M > -\infty$ for all $n > \max\{N, M\}$. Then $\delta_{\frac{1}{M_n}} \leq \delta_{\frac{1}{M}}$, hence

$$(3) \quad \mathcal{S}'_{\Delta}(x_0, \dots, x_n) = x'_n \in K_n \subseteq B\left(x'_n, \delta_{\frac{1}{M_n}}(x'_n)\right) \subseteq B\left(x'_n, \delta_{\frac{1}{M}}(x'_n)\right).$$

From (2) and (3) we get

$$x, x'_n \in K_n \subseteq B\left(x, \delta_{\frac{1}{M}}(x)\right) \cap B\left(x'_n, \delta_{\frac{1}{M}}(x'_n)\right),$$

and so

$$d_X(x'_n, x) < \min\left\{\delta_{\frac{1}{M}}(x'_n), \delta_{\frac{1}{M}}(x)\right\}.$$

To finish the proof it is enough to observe that, since $\delta_{\frac{1}{M}}$ is an $\frac{1}{M}$ -gauge for f , so

$$d_Y(f(\mathcal{S}'_{\Delta}(x_0, \dots, x_n)), f(x)) < \frac{1}{M} < \varepsilon.$$

□

Remark. In the original proof, Kiss noted that “the idea of the proof is to pick y_n as the image of a point in $\overline{B}(x_n, 2^{-n+1})$ at which f behaves *badly*”. In fact, we are able to shorten his argument, since the family of ε -gauges encodes the “bad” behaviour of f .

2.1. The games G'_f and G''_f . From Lemma 4 (see also [13, Theorem 1]) it follows that Player II has a winning strategy in the game G_f if and only if he/she has a winning strategy of the form $\mathcal{S}(x_0, x_1, \dots, x_n) = f(\mathcal{S}'(x_0, x_1, \dots, x_n))$.

Let X and Y be Polish spaces, $f: X \rightarrow Y$ be an arbitrary function. At the n th step of the game G'_f , Player I plays x_n , then Player II plays x'_n ,

$$\begin{array}{rccccccc} \text{Player I} & x_0 & x_1 & x_2 & \cdots & & \\ \text{Player II} & & x'_0 & x'_1 & x'_2 & \cdots & \end{array}$$

with the rules that for each $n \in \mathbb{N}$:

- $x_n \in X$ and $d_X(x_n, x_{n+1}) \leq 2^{-n}$;
- $x'_n \in X$.

Since X is complete, $x_n \rightarrow x$ for some $x \in X$. Player II wins if $\langle f(x'_n) \rangle$ is convergent to $f(x)$. Otherwise, Player I wins.

As a consequence of Lemma 4, we obtain the following result.

Theorem 5. *The game G'_f is determined, and*

- *Player I has a winning strategy in G'_f if and only if f is not of Baire class 1.*
- *Player II has a winning strategy in G'_f if and only if f is of Baire class 1.*

Proof. It is enough to show that:

- (i) if f is Baire class 1 then Player II has a winning strategy, and
- (ii) if f is not of Baire class 1 then Player I has a winning strategy.

To prove (i) observe that the function $\$'_\Delta$ from Lemma 4, for Δ being a family of gauges of f , is a winning strategy for Player II.

To see (ii), observe that the winning strategy for Player I in G_f is also a winning strategy for I in G'_f . Thus (ii) follows from Theorem 2. \square

Now, we propose a further modification of the game to obtain a similar effect. This time, we will define a point-open game G''_f . Let X and Y be Polish spaces, $f: X \rightarrow Y$ be an arbitrary function. At the first step of the game G''_f , Player I plays $x_0 \in X$ and then Player II plays an open set $U_0 \ni x_0$. At the n th step of the game G''_f ($n > 0$), Player I plays $x_n \in U_{n-1}$, then Player II plays an open set $U_n \ni x_n$:

Player I	x_0	x_1	x_2	\dots	
Player II	U_0	U_1	U_2	\dots	

with the rules that for each $n \in \mathbb{N}$:

- $x_0 \in X$, and $x_n \in U_{n-1}$ for $n > 0$;
- $U_n \ni x_n$.

If $\langle x_n \rangle$ is convergent and $\lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n)$ then Player II wins. Otherwise, Player I wins.

Theorem 6. *The game G''_f is determined, and*

- *Player I has a winning strategy in G''_f if and only if f is not of Baire class 1.*
- *Player II has a winning strategy in G''_f if and only if f is of Baire class 1.*

Proof. It is enough to prove two implications: if f is Baire 1 then Player II has a winning strategy, and if f is not Baire 1 then Player I has a winning strategy.

To see the first implication, assume that f is Baire 1 and let $\{\delta_\varepsilon: \varepsilon > 0\}$ be a family of ε -gauges for f . Without loss of generality (see Lemma 3) we may assume that, for any fixed $x \in X$, the sequence $\langle \delta_{\frac{1}{n}}(x) \rangle$ is decreasing, and $\delta_{\frac{1}{n}}(x) < 2^{-n}$ for every $n > 0$. In the n th move, Player II plays $U_n := B(x_n, \delta_{\frac{1}{n}}(x_n)/2)$. This is a winning strategy for Player II. Indeed, since $x_{n+1} \in U_n$, so $d_X(x_n, x_{n+1}) < \text{diam}(U_n) \leq 2^{-n}$ for every n . Hence $\langle x_n \rangle$ is a Cauchy sequence in a complete space X , so it converges. Let $x := \lim_{n \rightarrow \infty} x_n$. Fix $\varepsilon > 0$ and $N > 1/\varepsilon$. Then $x \in B(x_n, \delta_{\frac{1}{N}}(x_n))$ for each $n \geq N$, and for all n with $d_X(x, x_n) < \delta_{\frac{1}{N}}(x)$, we have $d_Y(f(x), f(x_n)) < \varepsilon$. Thus $\langle f(x_n) \rangle$ is convergent to $f(x)$.

Now assume that f is not Baire class 1. Then there are a perfect set $P \subseteq X$, $y_0 \in Y$ and $\varepsilon > 0$ such that both sets $A := \{x \in P: d_Y(f(x), y_0) < \varepsilon\}$ and $B := \{x \in P: d_Y(f(x), y_0) > 2\varepsilon\}$ are dense in P . The winning strategy for Player I in the game G''_f consists in choosing $x_n \in A$ for odd n and $x_n \in B$ for even n . In fact, if Player I plays this strategy then the sequence $\langle f(x_n) \rangle$ is not a Cauchy sequence. \square

3. GAMES FOR EQUI-BAIRE 1 FAMILIES OF FUNCTIONS

In this section, we modify games G'_f and G''_f to obtain characterizations of equi-Baire 1 families of functions.

Let X and Y be Polish spaces, let $\mathcal{F} \subseteq Y^X$. At the n th step of the game $G'_\mathcal{F}$, Player I plays x_n , then Player II plays x'_n ,

Player I	x_0	x_1	x_2	\dots	
Player II	x'_0	x'_1	x'_2	\dots	

with the rules that for each $n \in \mathbb{N}$:

- $x_n \in X$ and $d_X(x_n, x_{n+1}) \leq 2^{-n}$;
- $x'_n \in X$.

Since X is complete, $x_n \rightarrow x$ for some $x \in X$. Player II wins if

$$(4) \quad \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N \forall f \in \mathcal{F} \quad d_Y(f(x'_n), f(x)) < \varepsilon,$$

i.e. the indexed family of sequences $\{\langle f(x'_n) \rangle : f \in \mathcal{F}\}$ is equi-convergent to the indexed family $\{f(x) : f \in \mathcal{F}\}$. Otherwise, Player I wins.

We will use the fact that (4) is equivalent to the following Cauchy-type condition. (An easy proof of this fact is left to the reader.)

$$(5) \quad \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n, m \geq N \forall f \in \mathcal{F} \quad d_Y(f(x'_n), f(x'_m)) < \varepsilon.$$

Theorem 7. *The game $G'_{\mathcal{F}}$ is determined, and*

- *Player I has a winning strategy in $G'_{\mathcal{F}}$ if and only if \mathcal{F} is not equi-Baire 1.*
- *Player II has a winning strategy in $G'_{\mathcal{F}}$ if and only if \mathcal{F} is equi-Baire 1.*

Proof. It is enough to show that:

- (i) if \mathcal{F} is equi-Baire 1 then Player II has a winning strategy, and
- (ii) if \mathcal{F} is not of equi-Baire 1 then Player I has a winning strategy.

To prove (i) observe that from Lemma 4 we obtain:

if \mathcal{F} is equi-Baire 1 then there is a function $\$: X^{<\omega} \rightarrow X$ which is a winning strategy for Player II in the game $G'_{\mathcal{F}}$.

To see (ii) note that, if \mathcal{F} is not equi-Baire 1, then there exists a non-empty perfect set $P \subseteq X$ such that $f \upharpoonright P$ has no point of equicontinuity (see Theorem 1).

For any non-empty set $U \subseteq P$ let

$$\omega_{\mathcal{F} \upharpoonright P}(U) := \sup \{d_Y(f(u), f(v)) : u, v \in U \text{ and } f \in \mathcal{F}\}.$$

For any $x \in P$ define the *equi-oscillation of $\mathcal{F} \upharpoonright P$ at x* as

$$\omega_{\mathcal{F} \upharpoonright P}(x) := \inf \{\omega_{\mathcal{F} \upharpoonright P}(B(x, h) \cap P) : h > 0\}.$$

It is easy to observe that, for any integer $n > 0$, the set

$$P_n := \left\{ x \in P : \omega_{\mathcal{F} \upharpoonright P}(x) \geq \frac{1}{n} \right\}$$

is closed, and x is a point of equicontinuity for \mathcal{F} if and only if $\omega_{\mathcal{F} \upharpoonright P}(x) = 0$. Since $f \upharpoonright P$ has no point of equicontinuity, $\bigcup_{n \in \mathbb{N}} P_n = P$. Since P is a Baire space and all P_n 's are closed, by the Baire Category Theorem there exists P_n with non-empty interior (in P). Thus, without loss of generality, we may assume that for some $\varepsilon > 0$, $\omega_{\mathcal{F} \upharpoonright P}(x) \geq \varepsilon$ for each $x \in P$.

We are ready to provide a strategy $\$$ for Player I. In the first move she/he picks $x_0 \in P$. For $n > 0$, in the n th move Player I takes $\$(x'_0, \dots, x'_{n-1}) = x_n \in P$ with

$$x_n := \begin{cases} x_{n-1} & \text{if there exists } f \in \mathcal{F} \text{ such that } d_Y(f(x'_{n-1}), f(x_{n-1})) \geq \varepsilon/3; \\ a & \text{otherwise,} \end{cases}$$

where $a \in P$ and

- (j) $d_X(x_{n-1}, a) < 1/2^n$;
- (jj) $d_Y(f(x'_{n-1}), f(a)) \geq 2\varepsilon/3$ for some $f \in \mathcal{F}$.

Such a choice is possible because the set P is dense-in-itself, $x_{n-1} \in P$, and $\omega_{\mathcal{F}|P}(x_{n-1}) \geq \varepsilon$.

If the family of sequences $\{f(x'_n) : f \in \mathcal{F}\}$ is not equi-convergent then Player I wins. Otherwise, we use (5). So, there exists $N \in \mathbb{N}$ such that for all $n, m \geq N$ and all $f \in \mathcal{F}$,

$$(6) \quad d_Y(f(x'_n), f(x'_m)) < \frac{\varepsilon}{6}.$$

We claim that there exists $M \geq N$ with $x_n = x_M$ for all $n \geq M$.

Indeed, it is nothing to prove if $x_m = x_{m-1}$ for all $m > N$. Otherwise, there exists $m_1 > N$ such that $x_{m_1} \neq x_{m_1-1}$, then (by the formula defining x_n)

$$d_Y(f(x'_{m_1-1}), f(x_{m_1-1})) < \frac{\varepsilon}{3} \text{ for each } f \in \mathcal{F}.$$

Then it follows from (jj) that

$$(7) \quad \text{there exists } f_1 \in \mathcal{F} \text{ such that } d_Y(f_1(x'_{m_1-1}), f_1(x_{m_1})) \geq \frac{2\varepsilon}{3}.$$

Thus, by (6) and (7), for all $m \geq N$ we have

$$d_Y(f_1(x'_m), f_1(x_{m_1})) \geq \frac{\varepsilon}{3}.$$

Hence, by the definition of x_n for $n = m_1 + 1$ we obtain the equality $x_n = x_{m_1}$, so $f(x_n) = f(x_{m_1})$. Therefore, $d_Y(f_1(x'_{n+1}), f_1(x_n)) \geq \frac{\varepsilon}{3}$, so by the definition of x_{n+1} we get $x_{n+1} = x_n = x_{m_1}$. In this way we show, by induction, that $x_n = x_{m_1}$ for all $n \geq m_1$. This finishes the proof of the claim.

Since the sequence constructed by Player I is almost constant, i.e. $x_m = x_M$ for all $m \geq M$, so $\lim_{n \rightarrow \infty} x_n = x_M$. Recall that by the formula defining x_n ,

$$\forall n \in \mathbb{N} \exists f \in \mathcal{F} \quad d_Y(f(x'_n), f(x_n)) \geq \frac{\varepsilon}{3}.$$

Therefore, since $x_m = x_M = \lim_{n \rightarrow \infty} x_n$ for all $m \geq M$,

$$\forall m \geq M \exists f \in \mathcal{F} \quad d_Y\left(f(x'_m), f\left(\lim_{n \rightarrow \infty} x_n\right)\right) \geq \frac{\varepsilon}{3}.$$

Fix $f_2 \in \mathcal{F}$ with

$$d_Y\left(f_2(x'_M), f_2\left(\lim_{n \rightarrow \infty} x_n\right)\right) \geq \frac{\varepsilon}{3}.$$

By (6), for every $n \geq N$ we have

$$d_Y\left(f_2(x'_n), f_2\left(\lim_{n \rightarrow \infty} x_n\right)\right) \geq \frac{\varepsilon}{6},$$

thus $\langle f_2(x'_n) \rangle$ does not converge to $f_2(\lim_{n \rightarrow \infty} x_n)$, so Player I wins. \square

Now, we will describe the game $G''_{\mathcal{F}}$ which is a modification of G''_f for equi-Baire 1 families.

Let X and Y be Polish spaces, let $\mathcal{F} \subseteq Y^X$. At the first step of the game $G''_{\mathcal{F}}$, Player I plays $x_0 \in X$ and then Player II plays an open set $U_0 \ni x_0$. At the n th step of the game $G''_{\mathcal{F}}$, Player I plays $x_n \in U_{n-1}$, then Player II plays an open set $U_n \ni x_n$:

$$\begin{array}{ccccccc} \text{Player I} & x_0 & x_1 & x_2 & \cdots & \cdots & \\ \text{Player II} & U_0 & U_1 & U_2 & & & \cdots \end{array}$$

with the rules that for each $n \in \mathbb{N}$:

- $x_0 \in X$, and $x_n \in U_{n-1}$ for $n > 0$;
- $U_n \ni x_n$.

Player II wins if the sequence $\langle x_n \rangle$ converges to some $x \in X$, and the indexed family $\{\langle f(x_n) \rangle : f \in \mathcal{F}\}$ is equi-convergent to $\{f(x) : f \in \mathcal{F}\}$. Otherwise, Player I wins.

Theorem 8. *The game $G''_{\mathcal{F}}$ is determined, and*

- *Player I has a winning strategy in $G''_{\mathcal{F}}$ if and only if \mathcal{F} is not equi-Baire 1.*
- *Player II has a winning strategy in $G''_{\mathcal{F}}$ if and only if \mathcal{F} is equi-Baire 1.*

Proof. Firstly, we show that, if \mathcal{F} is equi-Baire 1, then Player II has a winning strategy. This is a consequence of the respective part in the proof of Theorem 6. Indeed, let \mathcal{F} be equi-Baire 1. Then the respective gauges $\delta_\varepsilon(x)$ do not depend on $f \in \mathcal{F}$. So, we choose $U_n := B(x_n, \delta_{\frac{1}{n}}(x_n)/2)$ (with the respective modification of $\delta_{\frac{1}{n}}$). Then $x_n \rightarrow x$ and note that the index N such that $d_Y(f(x), f(x_n)) < \varepsilon$ for all $n > N$ does not depend on $f \in \mathcal{F}$. Hence the family of sequences $\{\langle f(x_n) \rangle : f \in \mathcal{F}\}$ is equi-convergent to $\{f(x) : f \in \mathcal{F}\}$ and we are done.

Secondly, assuming that \mathcal{F} is not equi-Baire 1, we will show that Player I has a winning strategy. We follow the respective part in the proof of Theorem 7. We can assume that there exist a perfect set $P \subseteq X$ and $\varepsilon > 0$ such that $\omega_{\mathcal{F}|P}(x) \geq \varepsilon$ for each $x \in P$. Initially, Player I picks $x_0 \in P$. Let $n > 0$. Since $x_{n-1} \in P$, we have $\omega_{\mathcal{F}|P}(x_{n-1}) \geq \varepsilon$. Thus, knowing that U_{n-1} is an open neighbourhood of x_{n-1} , Player I can choose $x_n \in U_{n-1}$ and $f_n \in \mathcal{F}$ such that

$$d_Y(f_n(x_{n-1}), f_n(x_n)) \geq \frac{\varepsilon}{3}.$$

This, by condition (5), shows that the family of sequences $\{\langle f(x_n) \rangle : f \in \mathcal{F}\}$ is not equi-convergent. So, we have a winning strategy for Player I. \square

4. GAME CHARACTERIZATION OF MEASURABLE FUNCTIONS

In this section, we propose another modification of the game G''_f to obtain characterizations of Baire measurable and Lebesgue measurable functions.

Let Σ be a σ -algebra of subsets of a set $Z \neq \emptyset$. A function $f: Z \rightarrow Y$, where Y denotes a metric space, is called Σ -measurable if the preimage $f^{-1}[U]$ of any open set U in Y belongs to Σ . If Y is separable, we can use only open balls in the role of U in this condition.

Let $H(\Sigma)$ be the σ -ideal given by $H(\Sigma) := \{A \subseteq Z : \forall B \subseteq A, B \in \Sigma\}$. Denote $\Sigma^+ := \Sigma \setminus H(\Sigma)$. The following lemma is a simple exercise.

Lemma 9. *Let Y be a separable metric space. A function $f: Z \rightarrow Y$ is not Σ -measurable if and only if there exist a set $W \in \Sigma^+$, a point $y \in Y$ and $\varepsilon > 0$ such that the sets $\{x \in W : d_Y(f(x), y) < \varepsilon\}$ and $\{x \in W : d_Y(f(x), y) \geq 2\varepsilon\}$ intersect every subset of W that belongs to Σ^+ .*

First, we are going to characterize Baire measurable functions by the respective game. By definition, Baire measurable functions are Σ -measurable where Σ is the σ -algebra of sets with the Baire property in a topological space, cf. [12, 8.21].

Let X be Polish space, Y be a separable metric space, and $f: X \rightarrow Y$ an arbitrary function. At the first step of the game $c-G''_f$, Player I plays $x_0 \in X$ and then Player II plays a non-meager set $U_0 \ni x_0$ with the Baire property. At the n th step of the game $c-G''_f$, where $n > 0$, Player I plays $x_n \in U_{n-1}$, then Player II plays

a non-meager $U_n \ni x_n$ with the Baire property.

Player I	x_0	x_1	x_2	\dots	
Player II	U_0	U_1	U_2	\dots	

with the rules that for each $n \in \mathbb{N}$:

- $x_0 \in X$, and $x_n \in U_{n-1}$ for $n > 0$;
- $U_n \ni x_n$.

Player II wins if $\langle x_n \rangle$ is convergent and $\lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n)$. Otherwise Player I wins.

Theorem 10. *The game $c-G_f''$ is determined, and*

- *Player I has a winning strategy in $c-G_f''$ if and only if f is not Baire measurable.*
- *Player II has a winning strategy in $c-G_f''$ if and only if f is Baire measurable.*

Proof. As usual, we should prove two implication:

- (i) if f is Baire measurable then Player II has a winning strategy in the game $c-G_f''$;
- (ii) if f is not Baire measurable then Player I has a winning strategy in $c-G_f''$.

To prove (i) assume that f is Baire measurable. We will describe a winning strategy for Player II in the game $c-G_f''$. Let $G \subseteq X$ be a dense G_δ set such that $f \upharpoonright G$ is continuous. (See [12, Theorem 8.38].) Fix any point $a \in G$. Let x_0 be a point chosen by Player I at the first move. Then Player II picks $U_0 := \{x_0\} \cup (B(a, 1) \cap G)$. At the $(n+1)$ -th move, Player I chooses $x_{n+1} \in U_n$. Then Player II plays $U_{n+1} := \{x_{n+1}\} \cup (B(a, \frac{1}{n+1}) \cap G)$. When the game is finished, one of the two cases is possible: either $x_n = x_N$ for some $N \in \mathbb{N}$ and all $n > N$, or $x_n \in G$ for some $N \in \mathbb{N}$ and all $n > N$ which implies $\lim_{n \rightarrow \infty} x_n = a$. In both cases, $\lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n)$.

To prove (ii) assume that f is not Baire measurable. We use Lemma 9. In our case, $H(\Sigma)$ consists exactly of meager sets in X . Then there exist a non-meager set $W \subseteq X$ with the Baire property, $y \in Y$ and $\varepsilon > 0$ such that both sets $A := \{x \in W : d_Y(f(x), y) < \varepsilon\}$ and $B := \{x \in W : d_Y(f(x), y) \geq 2\varepsilon\}$ intersect every non-meager subset of W with the Baire property. Let Player I play the following strategy. At the first step, he/she chooses a point $x_0 \in W$. If $n > 0$, then $U_{n-1} \subseteq W$ and he/she chooses $x_n \in A \cap U_{n-1}$ when n is even and $x_n \in B \cap U_{n-1}$ when n is odd. Then $d_Y(f(x_{2n}), y) \leq \varepsilon$ and $d_Y(f(x_{2n+1}), y) \geq 2\varepsilon$, so $d_Y(f(x_{2n}), f(x_{2n+1})) \geq \varepsilon$ for every n , and therefore $\langle f(x_n) \rangle$ is not convergent. \square

A similar idea can be used to characterize Lebesgue measurable functions from $X := \mathbb{R}^k$ to a separable metric space Y by a modified game $m-G_f''$ where non-meager sets with the Baire property are replaced by Lebesgue measurable sets of positive measure. Denote this game by $m-G_f''$. The rules remain the same with the assumption that U_0, U_1, \dots are measurable sets of positive measure. Then we obtain the following

Theorem 11. *The game $m-G_f''$ is determined, and*

- *Player I has a winning strategy in $m-G_f''$ if and only if f is not measurable.*
- *Player II has a winning strategy in $m-G_f''$ if and only if f is measurable.*

Proof. First assume that f is measurable. We will describe a winning strategy for Player II. By the Lusin theorem, there exists a closed set $F_0 \subseteq \mathbb{R}$ such that $\lambda(F_0) > 0$ and $f \upharpoonright F_0$ is continuous. (Here $\lambda(F_0)$ denotes the Lebesgue measure of F_0 .) Let a point x_0 be chosen at the first move by Player I. Then Player II picks $U_0 := \{x_0\} \cup F_0$. At the $(n + 1)$ -th move, Player I chooses $x_{n+1} \in U_n$. Then Player II plays $U_{n+1} := \{x_{n+1}\} \cup F_{n+1}$ where $F_{n+1} \subseteq F_n$ is a closed set of a positive measure, with the diameter $< \frac{1}{n+1}$. When the game is finished, we have $\bigcap_{n \geq 0} F_n = \{a\}$ for some $a \in X$. As in the previous proof, we infer that $\langle x_n \rangle$ is convergent and $\lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n)$.

Now assume that f is non-measurable. We use Lemma 9. In this case, $H(\Sigma)$ consists exactly of Lebesgue null sets. The further reasoning is analogous to that applied in the second part of the proof of Theorem 10. \square

Observe that we can extend this result to the case where a σ -finite Borel regular measure is given on a Polish space; see [19] and [12, Theorem 17.12].

REFERENCES

- [1] A. Alikhani-Koopaei, *Equi-Baire one family on metric spaces: A generalization of equi-continuity; and some applications*, Topology Appl. **277** (2020), 107170.
- [2] A. Andretta, *More on Wadge determinacy*, Ann. Pure Appl. Logic **144** (2006), 2–32.
- [3] L. F. Aurichi, R. R. Dias, *A minicourse on topological games*, Topology Appl. **258** (2018), 305–335.
- [4] R. Camerlo, J. Duparc, *Some remarks on Baire’s grand theorem*, Arch. Math. Logic **57** (2018), 195–201.
- [5] J. Cao, W. B. Moors, *A survey on topological games and their applications in analysis*, RACSAM Rev. R. Acad. Cien. Serie A. Mat. **100** (2006), 39–49.
- [6] R. Carroy, *Playing in the first Baire class*, Math. Log. Q. **60** (2014), 118–132.
- [7] J. Duparc, *Wadge hierarchy and Veblen hierarchy I. Borel sets of finite rank*, J. Symb. Logic **66** (2001), 56–86.
- [8] M. Elekes, J. Flesch, V. Kiss, D. Nagy, M. Poór, A. Predtetchinski, *Games characterizing limsup functions and Baire class 1 functions*, preprint, arXiv:2010.03327.
- [9] E. Glasner, M. Megrelishvili, *Hereditarily non-sensitive dynamical systems and linear representations*, Colloq. Math. **104** (2006), 223–283.
- [10] G. Gruenhage, *Infinite games and generalizations of first-countable spaces*, Gen. Topol. Appl. **6** (1976), 339–352.
- [11] G. Gruenhage, *The story of a topological game*, Rocky Mt. Math. J. **36** (2006), 1885–1914.
- [12] A. S. Kechris, *Classical Descriptive Set Theory*, Springer, New York 1995.
- [13] V. Kiss, *A game characterizing Baire class 1 functions*, J. Symb. Logic **85** (2020), 456–466.
- [14] D. Lecomte, *How can we recover Baire class one functions?* Mathematika **50** (2003), 171–198.
- [15] P. Y. Lee, W.-K. Tang, D. Zhao, *An equivalent definition of functions of the first Baire class*, Proc. Amer. Math. Soc., **129** (2001), 2273–2275.
- [16] H. Nobrega, *Games for functions Baire classes, Weihrauch degrees, transfinite computations, and ranks*, Ph.D. thesis, University of Amsterdam 2018.
- [17] L. M. Ros, *Game representations of classes of piecewise definable functions*, MLQ Math. Log. Q., **57** (2011), 95–112.
- [18] B. Semmes, *Games, trees, and Borel functions*, Ph.D. thesis, University of Amsterdam, 2009.
- [19] A. H. Stone, *Lusin’s Theorem*, Atti Sem. Mat. Fis. Univ. Modena **44** (1996), 201–207.

INSTITUTE OF MATHEMATICS, LODZ UNIVERSITY OF TECHNOLOGY, AL. POLITECHNIKI 10,
93-590 ŁÓDŹ, POLAND

Email address: `marek.balcerzak@p.lodz.pl`

INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS, UNI-
VERSITY OF GDAŃSK, 80-308 GDAŃSK, POLAND

Email address: `tomasz.natkaniec@ug.edu.pl`

INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS, UNI-
VERSITY OF GDAŃSK, 80-308 GDAŃSK, POLAND

Email address: `piotr.szuca@ug.edu.pl`