

STABLY DIFFEOMORPHIC MANIFOLDS AND THE REALISATION OF MODIFIED SURGERY OBSTRUCTIONS

ANTHONY CONWAY, DIARMUID CROWLEY, MARK POWELL, AND JOERG SIXT

ABSTRACT. For every $k \geq 2$ we construct infinitely many $4k$ -dimensional manifolds that are all stably diffeomorphic but pairwise not homotopy equivalent. Each of these manifolds has hyperbolic intersection form and is stably parallelisable. In fact we construct infinitely many such infinite sets. To achieve this we prove a realisation result for appropriate subsets of Kreck's modified surgery monoid $\ell_{2q+1}(\mathbb{Z}[\pi])$, analogous to Wall's realisation of the odd-dimensional surgery obstruction L -group $L_{2q+1}^s(\mathbb{Z}[\pi])$.

1. INTRODUCTION

Let $W_g := \#_g(S^q \times S^q)$ be the g -fold connected sum of $S^q \times S^q$ with itself. Two compact, connected, smooth $2q$ -manifolds M_0 and M_1 with the same Euler characteristic are *stably diffeomorphic* if there exists a nonnegative integer g and a diffeomorphism

$$M_0 \# W_g \rightarrow M_1 \# W_g.$$

We write $M_0 \cong_{\text{st}} M_1$. Note that $S^q \times S^q$ admits an orientation reversing diffeomorphism for every q , using the antipodal map on one factor for q even, and switching the factors for q odd. Hence the same is true of W_g and it follows that when the M_i are orientable the diffeomorphism type of the connected sum does not depend on a choice of orientations.

An effective procedure for classifying $2q$ -dimensional manifolds M , pioneered by Hambleton and Kreck in dimension 4 [HK88a, HK88b, HK93b, HK93a] and systematised by Kreck [Kre99] in dimensions $2q \geq 4$, is to first classify manifolds up to stable diffeomorphism and then for a fixed manifold M to understand its *stable class*, which is the set of diffeomorphism classes

$$\mathcal{S}^{\text{st}}(M) := \{N \mid N \cong_{\text{st}} M\} / \text{diffeomorphism}.$$

The literature on stably diffeomorphic manifolds contains work on the stable classification of 4-manifolds, that is enumerating and detecting the possible stable classes, including [CHR95, Spa03, KLPT17, KPT20]. There is also work showing that, under additional assumptions, a stable diffeomorphism type contains a unique diffeomorphism class [HKT09, CS11, Kha17]. In addition Davis gave conditions under which the homotopy type of a 4-manifold M contains a unique stable class [Dav05].

In this paper, for $k \geq 2$ we present new general methods to construct stably diffeomorphic manifolds, examples of $4k$ -manifolds with infinite stable classes, and new invariants which distinguish stably diffeomorphic manifolds. Before stating our main results we briefly review the history of the stable class.

2020 *Mathematics Subject Classification*. Primary 57R65, 57R67.

Key words and phrases. Stable diffeomorphism, ℓ -monoid, $4k$ -manifold.

If $w_1(M)$ is the orientation character of M , then classical surgery theory provides an action of the surgery obstruction group $L_{2q+1}^s(\mathbb{Z}[\pi_1(M)], w_1(M))$ on $\mathcal{S}^{\text{st}}(M)$. This action can give rise to infinite stable classes, such as for $L \times S^1$, where L is a $(2q-1)$ -dimensional lens space [Wei99], and for $\mathbb{R}\mathbb{P}^{2n} \# \mathbb{R}\mathbb{P}^{2n}$ [BDK07]. However the action of $L_{2q+1}^s(\mathbb{Z}[\pi_1(M)], w_1(M))$ necessarily preserves the homotopy type of M , and in the simply-connected case $L_{2q+1}(\mathbb{Z})$ is trivial. We will investigate the stable class of M beyond the action of the classical L -group and so we define the *homotopy stable class* of M ,

$$\mathcal{S}_h^{\text{st}}(M_0) = \{M_1 \mid M_1 \cong_{\text{st}} M_0\} / \text{homotopy equivalence},$$

which is the set of homotopy types within the stable class of M_0 , and is thus a quotient of $\mathcal{S}^{\text{st}}(M_0)$.

As Kreck and Schafer point out [KS84, I], examples of smooth $(4k-1)$ -connected $8k$ -manifolds M with arbitrarily large homotopy class have been implicit in the literature since Wall's classification of these manifolds up to the action of exotic sphere [Wal62]. These examples are distinguished by their intersection form $\lambda_M: H_{2k}(M) \times H_{2k}(M) \rightarrow \mathbb{Z}$ (which must be definite and have signature divisible by $8|bP_{4k}|$, where $|bP_{4k}|$ is the order of the group of homotopy $(4k-1)$ -spheres which bound parallelisable manifolds.)

Kreck and Schafer themselves provided examples of $4k$ -manifolds M with nontrivial finite fundamental groups where the homotopy stable class of M contained distinct elements with isometric intersection forms and even isometric equivariant intersection forms.

In [CCPS21] we gave the first examples of simply-connected $4k$ -manifolds, for $k \geq 2$, with hyperbolic intersection form and arbitrarily large stable class. In contrast to Kreck and Schafer, we exhibited these examples in infinitely many stable classes. However as far as we know all of these stable classes are finite.

1.1. Manifolds with infinite homotopy stable class. Our main theorem asserts the existence, for every $k \geq 2$, of infinitely many mutually non stably diffeomorphic $4k$ -manifolds, each of which has *infinite homotopy stable class*. All of the $4k$ -manifolds we construct, like the examples of [KS84], have equivalent intersection forms and equivalent equivariant intersection forms. Our $4k$ -manifolds are also *stably parallelisable*, meaning that their tangent bundles become trivial after Whitney sum with a trivial bundle of sufficiently high rank, so one cannot use stable characteristic classes to distinguish these manifolds.

Theorem 1.1. *Fix an integer $k \geq 2$. There is a closed oriented smooth $4k$ -manifold M with $\pi_1(M) \cong \mathbb{Z}$ and $|\mathcal{S}_h^{\text{st}}(M)| = \infty$. Indeed, there are infinitely many stable diffeomorphism classes of such manifolds, $\{[M_i]_{\text{st}}\}_{i=1}^{\infty}$, such that each $\mathcal{S}^{\text{st}}(M_i)$ contains an infinite subset $\{M_i^j\}_{j=1}^{\infty}$ where $M_i^1 = M_i$ and each M_i^j is stably parallelisable, has hyperbolic intersection form and has hyperbolic equivariant intersection form. The manifolds $M_i^j \# S^{2k} \times S^{2k}$ are pairwise diffeomorphic.*

To construct the manifolds in the theorem, we build initial $4k$ -manifolds M_i . We describe a new general surgery procedure, leading to Theorem 1.3 and Theorem 3.10. When applied to M_i this procedure constructs new elements of the stable class of M_i , which are related to M_i by a bordism with a given modified surgery obstruction lying in a certain subset of the modified surgery monoid $\ell_{4k+1}(\mathbb{Z}[\mathbb{Z}])$. This procedure is a generalisation of the Wall Realisation of surgery obstructions in $L_{4k+1}^s(\mathbb{Z}[\pi])$ classical surgery theory. We prove Theorem 1.1 by finding infinitely many elements of $\ell_{4k+1}(\mathbb{Z}[\mathbb{Z}])$ which produce infinitely many homotopy types.

One can see that the manifolds $\{M_i^j\}_{j=1}^\infty$ are not homotopy equivalent as follows. The cohomology groups are $H^*(M_i^j; \mathbb{Z}[\mathbb{Z}]) \cong \mathbb{Z}[\mathbb{Z}]$ for $*$ = $2, 4, \dots, 2k - 2, 2k + 2, \dots, 4k - 2$, with $H^{2k}(M_i^j; \mathbb{Z}[\mathbb{Z}]) \cong \mathbb{Z}[\mathbb{Z}]^2$, and all odd degree cohomology trivial. For a generator $z \in H_c^2(M_i^j; \mathbb{Z}) \cong H^2(M_i^j; \mathbb{Z}[\mathbb{Z}]) \cong \mathbb{Z}[\mathbb{Z}]$, our homotopy invariant then arises by considering $z^k \in H^{2k}(M_i^j; \mathbb{Z}[\mathbb{Z}])$ up to the isometries of the hyperbolic intersection pairing; we refer to Remark 9.9 for more details.

1.2. Realising modified surgery obstructions. To describe the strategy in more detail, we expand to the consideration of smooth manifolds of dimension $2q$, for $q \geq 3$, recall some classical surgery theory, and explain the tools we have developed in terms of the analogy between classical and modified surgery. We will also mention where our results extend to dimension $2q = 4$, in the topological category.

One aim of surgery theory is to decide whether two compact, smooth n -dimensional manifolds M_0 and M_1 are diffeomorphic. To describe the surgery program in the briefest terms, assume that $\partial M_0 \cong \partial M_1$, that $f: M_0 \rightarrow M_1$ is a homotopy equivalence, and that $(W^{n+1}; M_0, M_1)$ is an oriented cobordism rel. boundary. Classical surgery, as developed by Browder, Novikov, Sullivan, and Wall, associates an obstruction $\sigma(F) \in L_{n+1}^s(\mathbb{Z}[\pi_1(M_0)])$ to a degree one normal bordism $(F, \text{Id}, f): (W; M_0, M_1) \rightarrow (M_0 \times [0, 1]; M_0, M_0)$. This obstruction vanishes if, and for $n+1 \geq 5$ only if, (W, F) is normally bordant rel. boundary to an s -cobordism. If this holds, then for $n \geq 5$, by the s -cobordism theorem, M_0 and M_1 are diffeomorphic.

If M_0 and M_1 are $2q$ -dimensional, then the aforementioned surgery obstruction is (an equivalence class of) a $(-1)^q$ -quadratic formation, i.e. a triple $((P, \psi); F, G)$, where (P, ψ) is a $(-1)^q$ -quadratic form, and F, G are lagrangians of (P, ψ) . A classical result in surgery theory is *Wall Realisation* [Wal99, Theorem 6.5]. We restrict to the oriented case here, since we shall do so throughout this article.

Theorem 1.2 (Wall Realisation). *Let M_0 be a compact, connected, oriented, smooth $2q$ -dimensional manifold with $q \geq 3$. Given $x \in L_{2q+1}^s(\mathbb{Z}[\pi_1(M_0)])$, there exists a degree one normal bordism rel. boundary $(W^{2q+1}; M_0, M_1) \rightarrow (M_0 \times [0, 1]; M_0 \times \{0\}, M_0 \times \{1\})$ whose surgery obstruction equals x .*

The same holds for $q = 2$ for topological manifolds if $\pi_1(M_0)$ is a good group [FQ90]. We shall prove an analogue of Theorem 1.2 for Kreck's modified surgery theory [Kre99]. The starting point of modified surgery theory is to weaken the homotopy theoretic assumption that M_0^{2q} and M_1^{2q} are homotopy equivalent and to strengthen the bundle-theoretic input as follows. First we fix a fibration $\xi: B \rightarrow BO$. A *normal j -smoothing* over (B, ξ) is a pair $(M, \bar{\nu})$, where M is a compact, smooth manifold, and $\bar{\nu}: M \rightarrow B$ is a $(j+1)$ -connected map that lifts the classifying map $\nu_M: M \rightarrow BO$ of the stable normal bundle.

We assume that M_0 and M_1 are connected, oriented, and admit normal $(q-1)$ -smoothings into (B, ξ) , which together bound a normal (B, ξ) -bordism $(W, \bar{\nu})$. By [Kre99, Theorem 4] there is a *modified surgery obstruction*

$$\Theta(W, \bar{\nu}) \in \ell_{2q+1}(\mathbb{Z}[\pi_1(B)]),$$

where the ℓ -monoid $\ell_{2q+1}(\mathbb{Z}[\pi_1(B)])$ consists of certain equivalence classes of quasi-formation. As we recall in Section 2, a *quasi-formation* is a triple $((P, \psi); F, V)$, where for $\varepsilon = (-1)^q$,

(P, ψ) is an ε -quadratic form, F is a lagrangian of (P, ψ) and $V \subseteq P$ is a half rank direct summand. The *induced form* of a quasi-formation $((P, \psi); F, V)$ is the quadratic form (V, θ) obtained by restricting ψ to V .

When $\Theta(W, \bar{\nu}) = [(P, \psi); F, V]$, we now explain how the induced form of $((P, \psi); F, V)$ is related to the topology of $(W, \bar{\nu})$. For the normal $(q-1)$ -smoothing $(M_0, \bar{\nu}_0)$, intersections and self-intersections endow the finitely generated $\mathbb{Z}[\pi_1(B)]$ module $K\pi_q(M_0) := \ker(\bar{\nu}_* : \pi_q(M_0) \rightarrow \pi_q(B))$ with the structure of a quadratic form $(K\pi_q(M_0), \theta_{M_0})$ called the *Wall form* of $(M_0, \bar{\nu}_0)$. If S is the finitely generated free $\mathbb{Z}[\pi_1(B)]$ -module on a set of generators for $K\pi_q(M_0)$, then the *0-stabilised Wall form* of $(M_0, \bar{\nu}_0)$ is

$$[v(\bar{\nu}_0)]_0 := [S, \mu]_0,$$

where μ is the ε -quadratic form induced on S and $[S, \mu]_0$ indicates the 0-stable isometry class of a form, as defined in Section 2.3. By [Kre99, Proposition 8], $[v(\bar{\nu}_0)]_0$ is well-defined and $[v(\bar{\nu}_0)]_0 = [V, \theta]_0$.

We can now state our realisation result for modified surgery obstructions, which is the second main result of this paper; (see Theorem 3.10 for a more detailed version of Theorem 1.3).

Theorem 1.3 (Realisation of modified surgery obstructions). *For $q \geq 3$, let $(M_0^{2q}, \bar{\nu}_0)$ be a normal $(q-1)$ -smoothing and let $x = ((P, \psi); F, V)$ be a quasi-formation such that $[v(\bar{\nu}_0)]_0 = [V, \theta]_0$. Then there exists a normal (B, ξ) -cobordism $(W, \bar{\nu})$ with $\partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0)$ and surgery obstruction*

$$[\Theta(W, \bar{\nu})] = [x] \in \ell_{2q+1}(\mathbb{Z}[\pi_1(B)]).$$

Again, the same results hold for $q = 2$, provided we are in the topological category and $\pi_1(B)$ is a good group. Given a $(q-1)$ -smoothing $(M_0^{2q}, \bar{\nu}_0)$ and $[x] \in \ell_{2q+1}(\mathbb{Z}[\pi_1(B)])$ whose 0-stabilised induced form is isomorphic to the 0-stabilised Wall form of $(M_0, \bar{\nu}_0)$, Theorem 3.10 produces another $(q-1)$ -smoothing $(M_1, \bar{\nu}_1) = \partial_+(W, \bar{\nu})$. Since $(M_0, \bar{\nu}_0)$ and $(M_1, \bar{\nu}_1)$ are normal $(q-1)$ -smoothings over (B, ξ) , [Kre99, Theorem C] ensures that M_0 and M_1 are stably diffeomorphic.

With $q = 2k$, Theorem 1.3 gives rise to a procedure to realise elements of $\ell_{4k+1}(\mathbb{Z}[\pi_1(B)]) \cong \ell_{4k+1}(\mathbb{Z}[\pi_1(M_0)])$ by a bordism based on a $4k$ -manifold M_0 , as alluded to after the statement of Theorem 1.1. The manifold at the other end of such a bordism is an element in the same stable class as M_0 .

Remark 1.4. Theorem 1.3 can be re-stated as follows. We define a $(2q+1)$ -dimensional *modified surgery problem* (W, ξ) to be a normal (B, ξ) -bordism from a fixed $(q-1)$ -smoothing $\partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0)$ to some $(q-1)$ -smoothing $\partial_+(W, \bar{\nu})$ and let

$$\mathcal{W}_{2q+1, q}(M_0, \bar{\nu}_0) := \{ [W, \bar{\nu}] \mid \partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0), \partial_+(W, \bar{\nu}) \text{ is a } (q-1)\text{-smoothing} \}$$

be the set of rel. boundary bordism classes of $(2q+1)$ -dimensional modified surgery problems over (B, ξ) with incoming boundary $(M_0, \bar{\nu}_0)$ and outgoing boundary a $(q-1)$ -smoothing. We have seen that there is a well-defined surgery obstruction map

$$\begin{aligned} \Theta : \mathcal{W}_{2q+1, q}(M_0, \bar{\nu}_0) &\rightarrow \ell_{2q+1}([v(\bar{\nu}_0)]_0) \\ [W, \bar{\nu}] &\mapsto \Theta(W, \bar{\nu}), \end{aligned}$$

where $\ell_{2q+1}([v(\bar{\nu}_0)]_0) \subseteq \ell_{2q+1}(\Lambda)$ is the subset of elements $[(P, \psi); F, V]$, where $[V, \theta]_0 = [v(\bar{\nu}_0)]_0$. Theorem 1.3 states that the surgery obstruction map Θ is onto.

The analogous statement for Wall Realisation also contains that statement that the surgery obstruction is injective, in the appropriate sense, and this allows one to formulate Wall Realisation as an action of $L_{2q+1}^s(\mathbb{Z}[\pi_1(M)])$ on an appropriate structure set of M . Calculations of the fourth author lead us to expect that Θ is not in general injective and this limits our current ability to use modified surgery to provide upper bounds on the stable class. However, we expect for an appropriate refinement of the ℓ -monoid, that a refined surgery obstruction can be defined for which the surgery obstruction map is injective and we plan to pursue this in further work.

1.3. Primitive embeddings and boundary isomorphisms. Now we describe the algebraic structures we use to distinguish the manifolds of Theorem 1.1, as well as their relevance to modified surgery theory. First, we return to the ℓ -monoid $\ell_{2q+1}(\mathbb{Z}[\pi_1(B)])$, and recall a structural result from [CS11]. Given a quasi-formation $((P, \psi); F, V)$ over a ring R with involution, there are induced quadratic forms (V, θ) and (V^\perp, θ^\perp) , where θ is the restriction of ψ to V , and θ^\perp is the restriction of ψ to the orthogonal complement V^\perp of $V \subseteq P$. Under favourable conditions, given $(-1)^q$ -quadratic forms v, v' , [CS11, Theorem 5.11] fits the set $\ell_{2q+1}(v, v')$ of quasi-formations whose induced forms are (0-stably) v and v' into an “exact sequence” (of sets):

$$(1) \quad L_{2q+1}^s(R) \xrightarrow{\rho} \ell_{2q+1}(v, v') \xrightarrow{\delta} \text{bIso}(\partial v, \partial v') \xrightarrow{\kappa} L_{2q}^s(R).$$

This means that fibres of δ equal orbits under the action ρ , and $\text{Im}(\delta) = \kappa^{-1}(0)$. We focus on the *boundary isomorphism set* $\text{bIso}(\partial v, \partial v')$. Briefly, ∂v and $\partial v'$ denote (split) *boundary formations* associated to v and v' , and $\text{bIso}(\partial v, \partial v')$ consists of stable homotopy classes of stable isomorphisms between ∂v and $\partial v'$; we refer to Section 4.2 for details. The map κ assigns to a stable isomorphism $f: \partial v \rightarrow \partial v'$ the element of $L_{2q}^s(R)$ represented by the *union* $v \cup_f -v'$. This is a nonsingular quadratic form whose definition is recalled in Section 4.3.

Next, we describe the relevance of the exact sequence displayed in (1) to modified surgery theory. In classical surgery theory, if a surgery obstruction $\sigma(F)$ is trivial in $L_{2q+1}^s(\mathbb{Z}[\pi_1(M_0)])$, then the degree one normal map F is normal bordant to a homotopy equivalence. In modified surgery theory, if a modified surgery obstruction $\Theta(W, \overline{\nu})$ is “elementary”, then the (B, ξ) -cobordism $(W, \overline{\nu})$ is (B, ξ) -bordant rel. boundary to an s -cobordism [Kre99, Theorem 3]. A quasi-formation is *elementary* if it is stably isomorphic (after adding trivial formations $(H_{(-1)^q}(F); F, F^*)$, with $H_{(-1)^q}(F)$ the standard hyperbolic form on $F \oplus F^*$) to a quasi-formation of the form $(H_{(-1)^q}(P), P, \begin{pmatrix} 1 \\ \phi \end{pmatrix}(P))$ for some homomorphism $\phi: P \rightarrow P^*$.

We note that an elementary quasi-formation x need not be trivial in the modified surgery monoid $\ell_{2q+1}(\mathbb{Z}[\pi_1(B)])$. In addition to the above statements, exactness of (1) means that $[x]$ is elementary modulo the action of $L_{2q+1}^s(\mathbb{Z}[\pi_1(B)])$, represented by the dashed arrow labelled ρ in (1), if and only if $\delta([x]) = 1$; see [CS11, Corollary 5.12].

The exact sequence (1) measures the degree to which a quasi-formation fails to be elementary. Boundary isomorphism sets remain difficult to compute directly, although see [CS11, Section 6] for some progress. We provide a new definition of $\text{bIso}(\partial v, \partial v')$ in terms what we call *2-sided primitive embeddings*. Given quadratic forms v, v' and a nonsingular quadratic form m , a 2-sided primitive embedding is a pair

$$(2) \quad v \xleftarrow{j} m \xleftarrow{j'} v'$$

of split isometric injections such that $j(v)^\perp = j'(v')$. The set $\text{bPrim}(v, v')$ of isomorphism classes of 2-sided primitive embedding is introduced in Section 6.1. Given $[f] \in \text{bIso}(\partial v, \partial v')$, the union $v \cup_f -v'$ comes with maps $v \hookrightarrow (v \cup_f -v') \hookrightarrow -v'$ that form a 2-sided primitive embedding. As a consequence, we obtain a map

$$\text{Pr}: \text{bIso}(\partial v, \partial v') \rightarrow \text{bPrim}(v, -v').$$

We show that Pr is a bijection, and so give a new formulation of the boundary isomorphism set. A more precise statement can be found in Theorem 6.11.

Theorem 1.5. *Let v, v' and m be ε -quadratic forms where the latter is nonsingular. The map Pr defines a bijection between the sets $\text{bIso}(\partial v, \partial v')$ and $\text{bPrim}(v, -v')$.*

Theorem 1.5 has two main features. Firstly, from a theoretical point of view, it provides a new perspective on the modified surgery obstruction and its failure to be elementary. Secondly, as we shall see in Example 6.6 below, in some situations the set $\text{bPrim}_m(v, -v')$ is easier to study than the boundary isomorphism set $\text{bIso}_m(\partial v, \partial v')$.

Next we outline how 2-sided primitive embeddings and boundary isomorphisms are used to construct the manifolds of Theorem 1.1.

The first step is carried out in Section 7, where for $q = 2k$ even we associate a boundary isomorphism $\delta_{\bar{\nu}}$ to every ‘‘even, split-free’’ normal $(2k-1)$ -smoothing $(M^{4k}, \bar{\nu})$. Under some technical assumptions, Theorem 7.7 provides a realisation result for these boundary isomorphisms, allowing us to change $\delta_{\bar{\nu}}$ by an element coming from $\ell_{4k+1}(\mathbb{Z}[\pi_1(B)])$ under the map δ in (1).

Next we build $4k$ -manifolds $\{M_1, M_2, \dots\}$ that are pairwise not stably diffeomorphic. The manifolds N_i that we constructed in [CCPS21] are recalled in Proposition 9.3, and then we define

$$M_i := N_i \# (S^1 \times S^{4k-1}).$$

For each i we use Theorem 7.7 to construct sets of normal $(2k-1)$ -smoothings $\{(M_i^j, \bar{\nu}_i^j)\}_{j=1}^\infty$ with distinct associated boundary isomorphisms $\{\delta_{\bar{\nu}_i^j}\}_{j=1}^\infty$. Section 5 provides infinitely many nontrivial elements. Finally, Section 8 shows that for these manifolds the $\delta_{\bar{\nu}_i^j}$ are invariants of the homotopy type. The proof once again uses 2-sided primitive embeddings.

1.4. Extended symmetric forms. To close the introduction we define a natural invariant of oriented normal $(q-1)$ -smoothings, which in the simple cases we consider associates a primitive embedding to a normal smoothing. This gives an alternative perspective on the results of this article.

Let $\bar{\nu}: M \rightarrow B$ a normal $(q-1)$ -smoothing with $q \geq 2$, where as above M is a closed, connected, smooth, oriented $2q$ -dimensional manifold. Since $\bar{\nu}$ is a q -equivalence, we use the map induced by $\bar{\nu}$ to identify the fundamental groups of M and B . We write $\Lambda := \mathbb{Z}[\pi_1(B)]$. For simplicity, we assume that $q = 2k$ is even. The *extended symmetric form* of $\bar{\nu}: M \rightarrow B$ is the triple

$$\bar{\lambda}(M, \bar{\nu}) := (H_{2k}(M; \Lambda), \lambda_M, \bar{\nu}_*),$$

where $\lambda_M: H_{2k}(M; \Lambda) \times H_{2k}(M; \Lambda) \rightarrow \Lambda$ is the equivariant intersection form of the manifold M and $\bar{\nu}_*: H_{2k}(M; \Lambda) \rightarrow H_{2k}(B; \Lambda)$ is the map induced by $\bar{\nu}$ and we use $\nu_*: \pi_1(M) \rightarrow \pi_1(B)$ to identify fundamental groups.

More generally, fix a unital ring with involution R . An *extended symmetric form over R* is a triple (H, λ, ν) , where $\lambda: H \times H \rightarrow R$ is a sesquilinear form and $\nu: H \rightarrow Q$ is an R -module homomorphism. For example, for any Q the standard hyperbolic form $(H_g(\Lambda), \lambda_g)$ on Λ^{2g} and the zero homomorphism $H_g(\Lambda) \rightarrow Q$ define the extended symmetric form $(H_g(\Lambda), \lambda_g, 0)$. Two extended symmetric forms (H, λ, ν) and (H', λ', ν') are *isometric* if there is an isometry $f: (H, \lambda) \rightarrow (H', \lambda')$ and an isomorphism $h: Q \rightarrow Q'$ such that $\nu'f = h\nu$. A *stable isometry* is an isometry $(H, \lambda, \nu) \oplus (H_g(\Lambda), \lambda_g, 0) \xrightarrow{\cong} (H', \lambda', \nu') \oplus (H_g(\Lambda), \lambda_g, 0)$ for some $g \geq 0$. In this case we write $(H, \lambda, \nu) \cong_{\text{st}} (H', \lambda', \nu')$.

We define the *stable class* of an extended symmetric form (H, λ, ν) to be the set of isometry classes of extended symmetric forms over Λ that become isometric to (H, λ, ν) after stabilisation with $(H_g(\Lambda), \lambda_g, 0)$ for some $g \geq 0$:

$$\mathcal{S}^{\text{st}}(H, \lambda, \nu) := \{(H', \lambda', \nu') \mid (H', \lambda', \nu') \cong_{\text{st}} (H, \lambda, \nu)\} / \text{isometry}.$$

If we stabilise $(M, \bar{\nu})$ by adding $(W_g, \bar{\nu}_g)$, where $\bar{\nu}_g$ is the trivial B -structure on W_g , then the effect on the extended symmetric form is to stabilise by $(H_g(\Lambda), \lambda_g, 0)$:

$$\bar{\lambda}(M \# W_g, \bar{\nu} \# \bar{\nu}_g) = \bar{\lambda}(M, \bar{\nu}) \oplus (H_g(\Lambda), \lambda_g, 0).$$

Recall that there is a homotopy functor $P_{2k-1}(-)$ sending a CW -complex to a model for its $(2k-1)$ st Postnikov type obtained by attaching cells of dimension $2k+1$ and higher. Now we introduce some simplifying assumptions on the normal smoothings, which will all be satisfied for the examples we build in Section 9 for the proof of Theorems 1.1.

- (i) The Λ -modules $H_{2k}(M; \Lambda)$, $H_{2k}(B; \Lambda)$, and $H^{2k}(B; \Lambda)$ are all finitely generated and free.
- (ii) The intersection form on $H_{2k}(M; \Lambda)$ is even.
- (iii) There is a map $B \rightarrow P_{2k-1}(M)$ inducing an isomorphism $H_{2k}(B; \Lambda) \xrightarrow{\cong} H_{2k}(P_{2k-1}(M); \Lambda)$, such that the composition $M \xrightarrow{\bar{\nu}} B \rightarrow P_{2k-1}(M)$ agrees with the Postnikov slice $M \rightarrow P_{2k-1}(M)$.

Write $\mathcal{S}_+^{\text{st}}(M)$ and $\mathcal{S}_{h,+}^{\text{st}}(M)$ for the set of oriented manifolds stably diffeomorphic to M modulo orientation-preserving diffeomorphisms and modulo orientation-preserving homotopy equivalences respectively, and equipped with an identification of their fundamental groups with π . We obtain a map

$$\bar{\lambda}: \mathcal{S}_+^{\text{st}}(M) \rightarrow \mathcal{S}^{\text{st}}(\bar{\lambda}(M, \bar{\nu})),$$

where using the third assumption we replace $H_{2k}(B; \Lambda)$ with $H_{2k}(P_{2k-1}(M); \Lambda)$ in the rôle of Q . Since orientation-preserving homotopy equivalences induce an isometry of intersection forms and since $P_{2k-1}(-)$ is functorial, $\bar{\lambda}$ factors through the forgetful map to $\mathcal{S}_{h,+}^{\text{st}}(M)$. Moreover Wall Realisation produces homotopy equivalent, stably diffeomorphic manifolds with the same normal $(2k-1)$ -type, so we obtain a factorisation:

$$\bar{\lambda}: \mathcal{S}_+^{\text{st}}(M) \rightarrow \mathcal{S}_+^{\text{st}}(M) / L_{4k+1}(\Lambda) \rightarrow \mathcal{S}_{h,+}^{\text{st}}(M) \rightarrow \mathcal{S}^{\text{st}}(\bar{\lambda}(M, \bar{\nu})).$$

We have the following refinement of Theorem 1.1, which can be compared with [BCD⁺20, Problem 11].

Theorem 1.6. *Fix $k \geq 2$. For the $4k$ -dimensional manifolds $\{M_i\}_{i=1}^{\infty}$ with $\pi_1(M_i) \cong \mathbb{Z}$, constructed in Section 9 for the proof of Theorem 1.1, the composition*

$$\bar{\lambda}: \mathcal{S}_+^{\text{st}}(M_i) / L_{4k+1}(\Lambda) \rightarrow \mathcal{S}_{h,+}^{\text{st}}(M) \rightarrow \mathcal{S}^{\text{st}}(\bar{\lambda}(M_i, \bar{\nu}_i))$$

is surjective and has infinite image. Moreover $\text{Aut}(\mathbb{Z}) = \{\pm 1\}$ is finite, so after further identifying two extended symmetric forms related by changing the identification of the fundamental group with \mathbb{Z} , the image remains infinite.

We conjecture that the map $\bar{\lambda}: \mathcal{S}^{\text{st}}(M_i)/L_{4k+1}(\Lambda) \rightarrow \mathcal{S}^{\text{st}}(\bar{\lambda}(M_i, \bar{\nu}_i))$ of Theorem 1.6 is also injective. All of the manifolds we construct to realise this infinite image have isometric intersection forms, so in other words forgetting $\nu: H \rightarrow Q$ yields isomorphic (non-extended) symmetric forms. We give the proof of Theorem 1.6 in Section 10. The proof that the image is infinite consists of relating extended symmetric forms with 2-sided primitive embeddings, which uses the first two assumptions listed above. Since in Section 9 we use 2-sided primitive embeddings to distinguish homotopy types of manifolds in the same stable class, in Section 10 we use this to show that the image of $\bar{\lambda}$ is infinite.

Organisation. In Section 2, we fix our notations on forms and formations. Section 3 reviews some aspects of modified surgery and prove Theorem 1.3 on the realisation of suitable elements of $\ell_{2q+1}(\mathbb{Z}[\pi_1(B)])$. In Section 4, we review the definition of the boundary isomorphism set and the boundary automorphism set bAut . Section 5, exhibits quadratic forms over $\mathbb{Z}[\mathbb{Z}]$ with infinite bAut . In Section 6, we introduce 2-sided primitive embeddings and prove Theorem 1.5 relating boundary automorphisms and two-sided primitive embeddings. In Section 7, we associate a boundary isomorphism to a normal smoothing and prove a second realisation result in Theorem 7.7. Section 8 interprets primitive embeddings as a homotopy invariant of manifolds in the same stable class. In Section 9, we combine realisation, the algebraic theory of boundary automorphisms and primitive embeddings, and the resulting homotopy invariant, to prove Theorem 1.1. Finally Section 10 proves Theorem 1.6.

Conventions. Throughout the article, manifolds are by default assumed to be compact, oriented, and connected. In Sections 7 and 9, $2q$ -dimensional manifolds are additionally assumed to be closed and q is assumed to be even, equal to $2k$.

The symbol Λ will always denote a unital ring with involution. In topological settings $\Lambda = \mathbb{Z}[\pi]$, the group ring of a fundamental group. If M is a left Λ -module then M^* or \overline{M} denotes the dual of M , $\text{Hom}(M, \lambda)$, which is given a left R -module structure using the involution on R in the usual way.

Let $f: X \rightarrow Y$ be a morphism in a category \mathcal{C} , and let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a contravariant functor. Assume that $F(f)$ is invertible. We shall often write the induced morphism in \mathcal{D} as $f^* = F(f)$, and denote its inverse by $f^{-*} := (f^*)^{-1} = F(f)^{-1}$.

Acknowledgements. The first three authors thank the Max Planck Institute for Mathematics in Bonn, where they were all visitors at some point during the long life of this project. MP was partially supported by EPSRC New Investigator grant EP/T028335/1 and EPSRC New Horizons grant EP/V04821X/1.

CONTENTS

1. Introduction	1
1.1. Manifolds with infinite homotopy stable class	2
1.2. Realising modified surgery obstructions	3
1.3. Primitive embeddings and boundary isomorphisms	5
1.4. Extended symmetric forms	6
Organisation	8

Conventions	8
Acknowledgements	8
2. Forms and formations	9
2.1. Forms and formations	9
2.2. Quasi-formations and the ℓ -monoid.	10
2.3. 0-stabilised quadratic forms and the structure of $\ell_{2q+1}(R)$	11
3. Realising odd dimensional modified surgery obstructions	12
3.1. Normal smoothings, normal bordisms and Wall forms	13
3.2. The modified surgery obstruction	14
3.3. Realising modified surgery obstructions	16
4. Boundary isomorphisms	21
4.1. Stable isomorphisms of split formations	22
4.2. Boundary isomorphisms	23
4.3. Gluing of forms	25
5. Infinite boundary automorphism sets	29
6. Primitive embeddings	33
6.1. 2-sided primitive embeddings	33
6.2. Primitive embeddings and boundary isomorphisms.	36
7. Realising boundary isomorphisms and primitive embeddings	40
7.1. The boundary isomorphism of an even split-free normal smoothing	41
7.2. The realisation of boundary automorphisms	44
8. Homotopy invariance of certain two-sided primitive embeddings	52
8.1. Normal $(q-1)$ -types	52
8.2. The Postnikov two-sided primitive embedding	53
9. Manifolds with infinite homotopy stable class	59
9.1. The simply-connected $4k$ -manifolds $N_{a,b}$	60
9.2. The $4k$ -manifolds $M_{a,b}$ and their associated primitive embeddings	62
9.3. Proof of Theorem 9.1	68
10. Extended symmetric forms and the homotopy stable class	69
References	71

2. FORMS AND FORMATIONS

In this section, we review some background material on forms and formations. From now on, we fix a positive integer $q \geq 2$, we set $\varepsilon = (-1)^q$ and let R be a weakly finite unital ring with an involution $x \mapsto \bar{x}$. We refer to [Coh77, p.143] for the definition of a weakly finite ring, but simply note that finitely generated modules over such rings have a well defined rank [Coh77, Proposition 4.6]. We abbreviate “finitely generated” by “f.g.” throughout. In Section 2.1, we review forms and formations and in Section 2.2 we review quasi-formations and some of the structure of the ℓ -monoids.

2.1. Forms and formations. In this subsection, we briefly review forms and formations and fix our notational conventions. References include [Ran02, CS11, Wal99].

Let $\tilde{K}_1(R)$ be the reduced K -group of R . The involution on R induces an involution $*$: $\tilde{K}_1(R) \rightarrow \tilde{K}_1(R)$ and we fix $Z \subseteq \tilde{K}_1(R)$ a $*$ -invariant subgroup. In topological applications, $R = \mathbb{Z}[\pi]$ will be the integral group ring of a group π and $Z = \{g \mid [g] \in \pi\} \subseteq \tilde{K}_1(\mathbb{Z}[\pi])$. An

s -basis for a stably free f.g. left R -module M is a basis of some f.g. free left module $M \oplus R^m$. A *based module* is a stably f.g. free left module together with an equivalence class of s -bases, where two bases are *equivalent* if the change of basis matrix represents an element in Z . An isomorphism f of based modules is *simple* if, with respect to some s -bases, the torsion of $f \oplus \text{Id}$ lies in Z .

A basis of M induces a dual basis on the dual module M^* and for $\varepsilon = \pm 1$, we consider the involution

$$\begin{aligned} \varepsilon T: \text{Hom}_R(M, M^*) &\rightarrow \text{Hom}_R(M, M^*) \\ \phi &\mapsto (x \mapsto (y \mapsto \overline{\varepsilon \phi(y)(x)})). \end{aligned}$$

We then set $Q_\varepsilon(M) := \text{coker}(1 - \varepsilon T)$.

Definition 2.1. An ε -quadratic form is a pair (M, ψ) , where M is a based module and $\psi \in Q_\varepsilon(M)$.

An ε -quadratic form is *nonsingular* (resp. *simple*) if its *symmetrization* $(1 + \varepsilon T)\psi$ is an isomorphism (resp. a simple isomorphism). An *isometry* $h: (M, \psi) \rightarrow (M', \psi')$ between quadratic forms consists of an isomorphism $h: M \rightarrow M'$ such that $h^* \theta' h = \theta \in Q_\varepsilon(M)$. Unless stated otherwise, we shall assume that all isometries are simple. For a based module L , the standard *hyperbolic* ε -quadratic form is defined as follows:

$$H_\varepsilon(L) = \left(L \oplus L^*, \left[\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right] \right).$$

The module L^* is based using the dual basis, and using this we have a canonical basis for $L \oplus L^*$, with respect to which $(1 + \varepsilon T)\psi$ is the simple isomorphism $\begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}$. A form (M, ψ) is called *hyperbolic* if it is isometric to $H_\varepsilon(L)$ for some L .

A *sublagrangian* of an ε -quadratic form (M, ψ) is a direct summand $j: L \hookrightarrow M$ which is based, and satisfies $j^* \psi j = 0 \in Q_\varepsilon(L)$. A sublagrangian $j: L \hookrightarrow M$ satisfies $j(L) \subseteq L^\perp$, where the (unbased) module $L^\perp = \ker(j^*(1 + \varepsilon T)\psi)$ is called the *annihilator* of $j: L \hookrightarrow M$. A sublagrangian $j: L \hookrightarrow M$ is a *lagrangian* if it satisfies $L = L^\perp$, as unbased modules. Setting $\phi = (1 + \varepsilon T)\psi$, observe that a lagrangian gives rise to an exact sequence

$$(3) \quad 0 \rightarrow L \xrightarrow{j} M \xrightarrow{j^* \phi} L^* \rightarrow 0.$$

A lagrangian L for (M, ψ) is *simple* if the s -basis of M is represented by a basis such that the exact sequence in (3) is based exact. The second main definition of this subsection is the following.

Definition 2.2. An ε -quadratic formation is a triple $((M, \psi); F, G)$, where (M, ψ) is a simple ε -quadratic form and F, G are simple lagrangians of (M, ψ) .

In the literature, what we call a formation is often referred to as a nonsingular formation. For concision, we drop this adjective. Finally, note that if $((M, \psi); F, G)$ is a formation, then (M, ψ) is a *nonsingular* quadratic form; in fact (M, ψ) is hyperbolic [Ran02, Proposition 12.3].

2.2. Quasi-formations and the ℓ -monoid. In this subsection, we describe quasi-formations and the ℓ -monoid $\ell_{2q+1}(R)$. Recall that $\varepsilon = (-1)^q$. The monoid $\ell_{2q+1}(R)$ was first defined by Kreck [Kre99]. The definition we present here is a reformulation from [CS11].

Definition 2.3. An ε -quadratic *quasi-formation* is a triple $x = ((M, \psi); F, V)$, where (M, ψ) is a simple ε -quadratic form, F is a simple lagrangian of (M, ψ) , and V is a half rank simple direct summand of M . A quasi-formation x is called *elementary* if the canonical map $F \oplus V \rightarrow M$ is a simple isomorphism.

If $x = ((M, \psi); F, V)$ is a quasi-formation in which V is a simple lagrangian of (M, ψ) , then x is a formation. In particular, all formations are quasi-formations and a *trivial (quasi)-formation* is a formation of type $(H_\varepsilon(P); P, P^*)$, for some f.g. stably free R -module P . Two quasi-formations $((M, \psi); F, V)$ and $((M', \psi'); F', V')$ are *isomorphic* if there is an isometry $h: (M, \psi) \rightarrow (M', \psi')$ such that $h(F) = F'$ and $h(V) = V'$, and such that the induced isomorphisms $F \xrightarrow{\cong} F'$ and $V \xrightarrow{\cong} V'$ are simple. Next, two quasi-formations $((M, \psi); F, V)$ and $((M', \psi'); F', V')$ are *stably isomorphic* if they become isomorphic after the addition of trivial formations. We use \cong to denote the isomorphism relation and \cong_s to denote the stable isomorphism relation.

Definition 2.4. The ℓ -monoid $\ell_{2q+1}(R) = (\{[x]\}, \oplus)$ is the unital abelian monoid of stable isomorphism classes of ε -quadratic quasi-formations modulo the further relation generated by

$$((M, \psi); F, G) \oplus ((M, \psi); G, V) \sim ((M, \psi); F, V),$$

where F and G are both simple lagrangians. Addition is by direct sum and the zero element is the class of the trivial formations.

The *elementary submonoid* $\mathcal{E}\ell_{2q+1}(R) \subseteq \ell_{2q+1}(R)$ is the submonoid of elements represented by elementary quasi-formations.

Note that Kreck's original definition of $\ell_{2q+1}(R)$ does not involve quasi-formations [Kre99, page 733]: it makes use of classes of pairs $(H_\varepsilon(\Lambda^k), V)$, where $V \subseteq \Lambda^{2k}$ is a based free direct summand of rank k . The equivalence between Kreck's definition and Definition 2.4 was proven in [CS11, Lemma 3.11]. Finally, if one is working up to isomorphism of quasi-formations, then without loss of generality one can write quasi-formations as $(H_\varepsilon(F); F, V)$.

2.3. 0-stabilised quadratic forms and the structure of $\ell_{2q+1}(R)$. In this subsection we briefly recall some of the structure of $\ell_{2q+1}(R)$, as discussed in [CS11]. Let $x = ((M, \psi); F, V)$ be an ε -quadratic quasi-formation with symmetric form $\lambda = \psi + \varepsilon\psi^*$. The *orthogonal complement* of V is the summand $V^\perp \subseteq M$ of elements orthogonal to V :

$$V^\perp := \{m \in M \mid \lambda(m, v) = 0 \text{ for all } v \in V\}.$$

Note that we take V^\perp with respect to the symmetrisation λ . The ε -quadratic form ψ induces ε -quadratic forms on the summands V and V^\perp , which we denote by $v = (V, \theta)$ and $v^\perp = (V^\perp, \theta^\perp)$ respectively. Since stabilisation with trivial quasi-formations has the effect of adding a zero form on a free module to the induced form, we shall consider quadratic forms up to *0-stabilisation*, where two ε -quadratic forms (N, χ) and (N', χ') are called *0-stably equivalent* if there is an isometry

$$(N, \chi) \oplus (P, 0) \cong (N', \chi') \oplus (Q, 0)$$

for some free modules P and Q . The 0-stable equivalence class of a form (N, χ) is denoted $[N, \chi]_0$ and such an equivalence class is called a *0-stabilised ε -quadratic form*. Often we abbreviate (N, χ) by n and $[N, \chi]_0$ by $[n]_0$. The set of 0-stabilised ε -quadratic forms,

$$\mathcal{F}_\varepsilon^{\text{zs}}(R) := (\{[n]_0\}, \oplus),$$

forms an abelian monoid under orthogonal sum, with zero the 0-stable class of the 0-form. We write $-[n] = [-n]$. There are monoid maps [CS11, Definition 5.2], called *boundary maps*,

$$b_- : \ell_{2q+1}(R) \rightarrow \mathcal{F}_\varepsilon^{\text{zs}}(R), \quad [(M, \psi); F, V] \mapsto [v]_0$$

and

$$b_+ : \ell_{2q+1}(R) \rightarrow \mathcal{F}_\varepsilon^{\text{zs}}(R), \quad [(M, \psi); F, V] \mapsto -[v^\perp]_0.$$

To characterise the image of the monoid map $b := b_- \oplus b_+ : \ell_{2q+1}(R) \rightarrow \mathcal{F}_\varepsilon^{\text{zs}}(R) \times \mathcal{F}_\varepsilon^{\text{zs}}(R)$ we recall that two 0-stabilised quadratic forms $[N_0, \chi_0]_0$ and $[N_1, \chi_1]_0$ are called *stably equivalent* if they become equal after the addition of 0-stabilised hyperbolic forms, so that

$$[N_0, \chi_0]_0 \oplus [H_\varepsilon(\Lambda^r)]_0 = [N_1, \chi_1]_0 \oplus [H_\varepsilon(\Lambda^s)]_0$$

for some r, s . In this case we write $[N_0, \chi_0]_0 =_{\text{st}} [N_1, \chi_1]_0$ and we define the *stable diagonal*

$$\Delta_{\text{st}} := \{([n_0]_0, [n_1]_0) \mid [n_0]_0 =_{\text{st}} [n_1]_0\} \subseteq \mathcal{F}_\varepsilon^{\text{zs}}(R) \times \mathcal{F}_\varepsilon^{\text{zs}}(R).$$

Proposition 2.5 (cf. [CS11, Corollary 5.3]).

- (i) $b(\ell_{2q+1}(R)) = \Delta_{\text{st}}$;
- (ii) $b_-|_{\mathcal{E}\ell_{2q+1}} = -b_+|_{\mathcal{E}\ell_{2q+1}}$ and $b_-|_{\mathcal{E}\ell_{2q+1}} : \mathcal{E}\ell_{2q+1} \rightarrow \mathcal{F}_\varepsilon^{\text{zs}}(R)$ is an isomorphism. \square

For future use we make the following definitions. In light of Proposition 2.5 (ii), we let

$$[e]_{[n]_0} \in \mathcal{E}\ell_{2q+1}(R)$$

denote the unique elementary element with $b_-([e]_{[n]_0}) = [n]_0$. For a fixed 0-stabilised ε -quadratic form $[n]_0$ we define

$$\ell_{2q+1}([n]_0) := b_-^{-1}([n]_0) \subseteq \ell_{2q+1}(R)$$

to be the subset represented by quasi-formations $((M, \psi); L, V)$ with $[v]_0 = [n]_0$. Similarly we define

$$\ell_{2q+1}([n]_0, [m]_0) := b_-^{-1}([n]_0) \cap b_-^{-1}([m]_0) \subseteq \ell_{2q+1}(R).$$

This appears as $\ell_{2q+1}(v, v')$ in the sequence (1), where the 0-stable equivalence class was dropped from the notation.

3. REALISING ODD DIMENSIONAL MODIFIED SURGERY OBSTRUCTIONS

As discussed in the introduction, Wall showed that under suitable conditions, elements of the quadratic L -group $L_{2q+1}^s(\mathbb{Z}[\pi])$ can be realised as surgery obstructions of degree one normal maps [Wal99, Theorem 6.5]. The goal of this section is to prove an analogue of Wall's realisation result for Kreck's odd-dimensional modified surgery obstructions. Unless otherwise stated, manifolds are assumed to be compact, connected, oriented, and smooth. This section is organised as follows: in Section 3.1 we recall the definition of normal smoothings and normal bordism, in Section 3.2 we review the definition of the odd-dimensional modified surgery obstruction, and in Section 3.3 we prove our realisation result, Theorem 3.10.

3.1. Normal smoothings, normal bordisms and Wall forms. In this subsection, we briefly review the notions of normal smoothing and normal bordisms. The main reference is [Kre99].

We fix a fibration $\xi: B \rightarrow BO$. An n -dimensional (B, ξ) -manifold consists of a pair $(M, \bar{\nu})$, where M is a compact, connected, oriented, smooth n -dimensional manifold and $\bar{\nu}: M \rightarrow B$ is a lift of the stable normal bundle $\nu: M \rightarrow BO$ of M , meaning that $\nu = \xi \circ \bar{\nu}$.

A (B, ξ) -null-bordism for a closed n -dimensional (B, ξ) -manifold $(M_0, \bar{\nu}_0)$ is an $(n+1)$ -dimensional (B, ξ) -manifold $(W, \bar{\nu})$ for which $\partial(W, \bar{\nu}) = (M_0, \bar{\nu}_0)$. A (B, ξ) -cobordism between two n -dimensional (B, ξ) -manifolds $(M_0, \bar{\nu}_0)$ and $(M_1, \bar{\nu}_1)$ consists of a diffeomorphism $f: \partial M_0 \rightarrow \partial M_1$ which is compatible with the (B, ξ) -structures and an $(n+1)$ -dimensional (B, ξ) -null bordism of $(M_0 \cup_f -M_1, \bar{\nu}')$ where $\bar{\nu}'|_{M_0} = \bar{\nu}_0$ and $\bar{\nu}'|_{M_1} = -\bar{\nu}_1$. We write

$$\partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0) \quad \text{and} \quad \partial_+(W, \bar{\nu}) = (M_1, \bar{\nu}_1).$$

Modified surgery theory is a flexible theory in that one can use any fibration $\xi: B \rightarrow BO$, however for classification results one needs the maps $\bar{\nu}_i: M_i \rightarrow B$ to be roughly $n/2$ -connected.

Definition 3.1. Let $\xi: B \rightarrow BO$ be a fibration and let $(M, \bar{\nu})$ be a (B, ξ) -manifold.

- (i) If $\bar{\nu}$ is $(k+1)$ -connected then $(M, \bar{\nu})$ is called a *normal k -smoothing*.
- (ii) If $(M, \bar{\nu})$ is a normal k -smoothing and $\xi: B \rightarrow BO$ is $(k+1)$ -coconnected, then by [Kre99, p.711] the fibre homotopy type of ξ is uniquely determined and called a *normal k -type* of M . Moreover, the existence of the Moore-Postnikov factorisation [Bau77] of $\nu: M \rightarrow BO$ ensures that such fibrations exist for all $k \geq 0$.

Remark 3.2. While the normal k -types provide an important class of targets in modified surgery, the theory applies to any fibration $\xi: B \rightarrow BO$. However, the universal properties of the Moore-Postnikov factorisation entail that if $\bar{\nu}: M \rightarrow B$ is a normal $(k+1)$ -smoothing, then $\xi: B \rightarrow BO$ factors through the normal k -type of M , $\xi: B \rightarrow B^{q-1}(M) \rightarrow BO$.

Whenever $\bar{\nu}_*$ is 2-connected, we will use $\bar{\nu}_*: \pi_1(M) \rightarrow \pi_1(B)$ to identify $\pi_1(M) = \pi_1(B)$. For concision, we let

$$\Lambda := \mathbb{Z}[\pi_1(B)]$$

denote the group ring of $\pi_1(B)$ and we will use Λ and $\mathbb{Z}[\pi_1(B)]$ interchangeably, depending on whether we wish to emphasise topology or algebra.

Now suppose that M is a $2q$ -dimensional manifold, with a normal $(q-1)$ -smoothing $(M, \bar{\nu})$. We set

$$K\pi_q(M) := \ker(\bar{\nu}_*: \pi_q(M) \rightarrow \pi_q(B))$$

and $\varepsilon := (-1)^q$. As explained in [Kre99, Section 5], intersection and self-intersections of framed immersions of q -spheres in M equip $K\pi_q(M)$ with a quadratic refinement. For $q \neq 1, 3, 7$ or for $q = 3, 7$ and $w_{q+1}(\xi)(\pi_{q+1}(B)) = 0$, the quadratic refinement is given by an ε -quadratic form $(K\pi_q(M), \theta)$ over Λ . For $q = 3, 7$ and $w_{q+1}(\xi)(\pi_{q+1}(B)) \neq 0$, the quadratic refinement takes values in a quotient of $Q_-(\Lambda)$. Since the main examples of this paper are $4k$ -dimensional, so that $q = 2k$ is even, we shall leave aside the subtleties of the case $q = 3, 7$ and $w_{q+1}(\xi)(\pi_{q+1}(B)) \neq 0$ and make the following definition.

Definition 3.3. (Standard pair) A pair $((B, \xi), q)$, consisting of a fibration over BO and a dimension, is *standard* if $q \neq 3, 7$ or $w_{q+1}(\xi)(\pi_{q+1}(B)) = 0$. In addition we require that B has finite q -skeleton.

Henceforth we restrict our attention to standard pairs $((B, \xi), q)$. We call the ε -quadratic form $(K\pi_q(M), \theta)$ the *Wall form* of $\bar{\nu}: M \rightarrow B$. Standard surgery arguments ensure that one can perform surgery on $x \in K\pi_q(M)$ so that the normal (B, ξ) -structure on M extends over the trace and effect of the surgery if and only if $\theta(x) = 0$ [Kre99, Lemma 2, Propositions 6 and 7].

It turns out that the 0-stabilisation of the Wall form is the invariant which appears in the algebra of modified surgery. While $K\pi_q(M)$ need not be a free Λ -module, it is finitely generated.

Lemma 3.4. *The Λ -module $K\pi_q(M)$ is finitely generated.*

Proof. Since $\bar{\nu}: M \rightarrow B$ is q -connected, $\pi_{q+1}(B, M) \cong H_{q+1}(B, M; \Lambda)$ by the Hurewicz theorem. Then $K\pi_q(M)$ is a quotient of $\pi_{q+1}(B, M)$, so it suffices to see that $H_{q+1}(B, M; \Lambda)$ is finitely generated. The algebraic mapping cone of map $\bar{\nu}_*$ between the cellular chain complexes gives a chain complex D_* computing $H_*(B, M; \Lambda)$. Since M is compact and B has finite q -skeleton, in degrees $0 \leq i \leq q$ this chain complex consists of f.g. free Λ -modules. Since $H_i(D_*) = 0$ for $0 \leq i \leq q$, we have an exact sequence

$$0 \rightarrow \ker d_{q+1} \rightarrow D_{q+1} \xrightarrow{d_{q+1}} D_q \rightarrow \cdots \rightarrow D_0 \rightarrow 0.$$

It follows from Schanuel's lemma [Lam99, Corollary 5.5] that $\ker(d_{q+1})$ is f.g. projective. Then $H_{q+1}(B, M; \Lambda)$ is a quotient of $\ker(d_{q+1})$, so is finitely generated. \square

Thus there is a surjection $\pi: S \twoheadrightarrow K\pi_q(M)$ from a f.g. free Λ -module S . The pull-back of $(K\pi_q(M), \theta)$ along π is denoted (S, μ) and called a *free Wall form* of $(M, \bar{\nu})$. By [Kre99, Proposition 8] the 0-stable class of (S, μ) is well-defined.

Definition 3.5. The 0-stabilised *Wall form* of a $(q-1)$ -smoothing $(M^{2q}, \bar{\nu})$ is defined by

$$[v(\bar{\nu})]_0 := [S, \mu]_0.$$

3.2. The modified surgery obstruction. We recall the definition of the odd-dimensional modified surgery obstruction and give a unified presentation of foundational results of Kreck about this obstruction. From now on, all normal smoothings will be to a fixed fibration $\xi: B \rightarrow BO$. The main reference is [Kre99, Section 6]; see also [CS11, Section 2].

Recall that a $(2q+1)$ -dimensional *modified surgery problem* (W, ξ) is a normal (B, ξ) -bordism where $(M_0, \bar{\nu}_0) = \partial_-(W, \bar{\nu})$ and $(M_1, \bar{\nu}_1) = \partial_+(W, \bar{\nu})$ are $(q-1)$ -smoothings. We define

$$\mathcal{W}_{2q+1, q}(M, \bar{\nu}_0) := \{ [W, \bar{\nu}] \mid \partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0), \partial_+(W, \bar{\nu}) \text{ is a } (q-1)\text{-smoothing} \}$$

to be the set of rel. boundary bordism classes of $(2q+1)$ -dimensional modified surgery problems over (B, ξ) with incoming boundary $(M_0, \bar{\nu}_0)$ and outgoing boundary a $(q-1)$ -smoothing. An important problem in surgery is to determine when modified surgery problem $(W, \bar{\nu})$ is (B, ξ) -bordant rel. boundary to an s -cobordism and more generally to give an algebraic classification of $\mathcal{W}_{2q+1, q}(M_0, \bar{\nu}_0)$. The following theorem combines [Kre99, Theorem 4 & Proposition 8], two fundamental results of Kreck on this problem.

Theorem 3.6 (The modified surgery obstruction in odd dimensions). *If $((B, \xi), q)$ is standard, then the map*

$$\Theta: \mathcal{W}_{2q+1, q}(M_0, \bar{\nu}_0) \rightarrow \ell_{2q+1}([v(\bar{\nu}_0)]_0) \subseteq \ell_{2q+1}(\Lambda)$$

$$[W, \bar{\nu}] \mapsto \Theta(W, \bar{\nu}),$$

is well-defined and is such that $\Theta(W, \bar{\nu}) \in \mathcal{E}\ell_{2q+1}(\Lambda)$ if and only if $(W, \bar{\nu})$ is (B, ξ) -bordant rel. boundary to an s -cobordism. Moreover the boundary maps on $\Theta(W, \bar{\nu})$ satisfy the equations $b_-(\Theta(W, \bar{\nu})) = [v(\bar{\nu}_0)]_0$ and $b_+(\Theta(W, \bar{\nu})) = -[v(\bar{\nu}_1)]_0$.

Remark 3.7. This theorem has been used to prove a truly vast array of classification results, often by showing that $\ell_{2q+1}([v(\bar{\nu}_0)]_0) = \{[e]_{[v(\bar{\nu}_0)]_0}\} \subseteq \mathcal{E}\ell_{2q+1}(\Lambda)$, using the notation introduced just after Proposition 2.5. However, to the best of our knowledge the image of Θ has not been systematically discussed in the literature, nor have the sets $\Theta^{-1}([x])$, for $[x] \neq [e]_{[v(\bar{\nu}_0)]_0}$. In this paper, we shall show that Θ is surjective; see Theorem 3.10 below. We plan to investigate the pre-images $\Theta^{-1}([x])$ in future work.

Remark 3.8. The appropriate analogue of Theorem 3.6 holds in the non-standard cases. The proof again follows from [Kre99, Theorem 4 & Proposition 8], but some care is needed to interpret [Kre99, Proposition 8] in non-standard cases.

We now recall the definition of the modified surgery obstruction $\Theta(W, \bar{\nu})$ for a bordism $(W, \bar{\nu})$. By surgery below the middle dimension [Kre99, Proposition 4], we assume that $\bar{\nu}: W \rightarrow B$ is q -connected. It follows by the same argument as that in Lemma 3.4 that $K\pi_q(W)$ is finitely generated. Choose a set ω of generators for $K\pi_q(W)$ and disjoint embeddings $\varphi_i: S^q \times D^{q+1} \hookrightarrow W$ compatible with the (B, ξ) -structure representing these generators, and set

$$U := \bigcup_i \varphi_i(S^q \times D^{q+1}).$$

Let ψ denote the quadratic form on $H_q(\partial U; \mathbb{Z}[\pi_1(B)])$. Since $\partial U = \bigcup_i S^q \times S^q$, we deduce that $(H_q(\partial U; \mathbb{Z}[\pi_1(B)]), \psi)$ is a hyperbolic quadratic form. For concision, we refer to the pair (ω, φ) as a *set of embedded generators for $K\pi_q(W)$* , we set $\Lambda := \mathbb{Z}[\pi_1(B)]$, and consider the Λ -modules

$$F := H_{q+1}(U, \partial U; \Lambda) \text{ and } V := H_{q+1}(W \setminus \mathring{U}, M_0 \sqcup \partial U; \Lambda).$$

Via the boundary maps in the long exact sequences of the appropriate pairs, F and V can be identified with submodules of $P := H_q(\partial U; \Lambda)$. In addition, F is a lagrangian and V is a half rank direct summand [Kre99, p. 734]. The *kernel quasi-formation* is then defined as

$$\Sigma(W, \bar{\nu}, \omega, \varphi) := ((P, \psi); F, V).$$

The class of this quasi-formation in $\ell_{2q+1}(\Lambda)$ only depends on the (B, ξ) -bordism class rel. boundary of the normal (B, ξ) -null-bordism $(W, \bar{\nu})$ [Kre99, Theorem 4]. This leads to the following definition, which is due to Kreck [Kre99, p. 734]; see also [CS11, p. 494].

Definition 3.9. Let $(M_0^{2q}, \bar{\nu}_0), (M_1^{2q}, \bar{\nu}_1)$ be two $(q-1)$ -smoothings. Let $f: \partial M_0^{2q} \rightarrow \partial M_1^{2q}$ be a diffeomorphism that is compatible with the smoothings. Assume that $(W^{2q+1}, \bar{\nu})$ is a normal (B, ξ) -null-bordism for $M_0 \cup_f -M_1$. The *modified surgery obstruction* of $(W, \bar{\nu})$ is

$$\Theta(W, \bar{\nu}) := [(\Sigma(W', \bar{\nu}', \omega', \varphi'))] \in \ell_{2q+1}(\Lambda),$$

where $(W', \bar{\nu}')$ is a normal $(q-1)$ -smoothing over (B, ξ) , which is (B, ξ) -bordant rel. boundary to $(W, \bar{\nu})$ and (ω', φ') is a system of embedded generators for $K\pi_q(W')$.

As stated in Theorem 3.6, the class $\Theta(W, \bar{\nu}) \in \ell_{2q+1}(\Lambda)$ only depends on the (B, ξ) -bordism class rel. boundary of the (B, ξ) -null-bordism $(W, \bar{\nu})$: it is independent of the choice of the $(q-1)$ -smoothing $(W', \bar{\nu}')$ and of the subsequent choice of a system of embedded generators for $K\pi_q(W')$.

3.3. Realising modified surgery obstructions. We recall that $((B, \xi), q)$ is a standard pair consisting of a fibration (B, ξ) and a dimension q . In this subsection, we state and prove our realisation theorem, Theorem 3.10, for the odd-dimensional modified surgery obstruction. For a $(q-1)$ -smoothing $(M_0, \bar{\nu}_0)$, this theorem implies that the surgery obstruction map

$$\Theta: \mathcal{W}_{2q+1, q}(M_0, \bar{\nu}_0) \rightarrow \ell_{2q+1}([v(\bar{\nu}_0)]_0)$$

of Theorem 3.6 is onto. In fact we prove a realisation result for quasi-formations of sufficiently large rank.

Let $\pi: S \rightarrow K\pi_q(M_0)$ be a surjection from a free Λ -module with (S, μ) the associated free Wall form of $(M_0, \bar{\nu}_0)$. Given $[x] \in \ell_{2q+1}([v(\bar{\nu}_0)]_0)$ we may assume that it is represented by a quasi-formation $x = (H_\varepsilon(K); K, V)$ of large enough rank so that for some r there is an isometry of quadratic forms

$$(4) \quad h: (S, \mu) \oplus (\Lambda^r, 0) \xrightarrow{\cong} (V, \theta).$$

Theorem 3.10. *Let $q \geq 3$, let $(M_0^{2q}, \bar{\nu}_0)$ be a normal $(q-1)$ -smoothing and let $[x] \in \ell_{2q+1}([v(\bar{\nu}_0)]_0)$ be represented by $x = (H_\varepsilon(K); K, V)$, where (V, θ) is 0-stably isomorphic to a free Wall form (S, μ) of $(M_0, \bar{\nu}_0)$ as in (4). Then there is a (B, ξ) -cobordism $(W, \bar{\nu})$ with $\partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0)$ such that $\Sigma(W, \bar{\nu}, \omega, \varphi) = (H_\varepsilon(K); K, V)$ for certain choices of ω and φ . Hence*

$$\Theta(W, \bar{\nu}) = [x].$$

The same statement holds for $q = 2$ in the topological category if $\pi_1(B)$ is a good group.

The proof of Theorem 3.10 occupies the remainder of the section. First we use $(M_0, \bar{\nu}_0)$ and the data x, π , and h to construct a (B, ξ) -cobordism $(W, \bar{\nu})$ based on $(M_0, \bar{\nu}_0)$. Then we prove a lemma, then in the proof environment we show that $\Theta(W, \bar{\nu}) = [x]$.

Note that since the Λ -modules V and K have the same rank, we deduce that K also has rank $s+r$, where s is the rank of S . The first step is to perform trivial surgeries on the lagrangian K that appears in the quasi-formation $x = (H_\varepsilon(K); K, V)$. Namely, we let W_0 be the cobordism obtained from $M_0 \times [0, 1]$ by adding trivial q -handles along $M_0 \times \{1\}$ with trivial framing, one handle for each element of a basis for K . In other words, we consider the following boundary connected sum:

$$W_0 = (M_0 \times [0, 1]) \natural (\natural_{i=1}^{s+r} S^q \times D^{q+1}).$$

The boundary components of W_0 are $\partial_- W_0 = M_0$ and $\partial_+ W_0 = -M_0 \# (\natural_{i=1}^{s+r} S^q \times S^q)$. We turn W_0 into a (B, ξ) -manifold $(W_0, \bar{\nu}_{W_0})$ by taking the trivial extension of the normal structure $\bar{\nu}_0$ to W_0 (i.e. the (B, ξ) -structure obtained by restricting a null-homotopic map $D^{q+1} \times D^q \rightarrow B$ on the handles to the $S^q \times D^{q+1}$). This concludes the first step in the construction of W .

Next we wish to add $(q+1)$ -handles to W_0 along $\partial_+ W_0$ in order to complete the construction of the (B, ξ) -cobordism $(W, \bar{\nu})$. The attaching maps for these handles must lie in $K\pi_q(\partial_+ W_0)$, the domain of the Wall form of $\partial_+ W_0$. The identification $\partial_+ W_0 = -M_0 \# (\#_{i=1}^{s+r} S^q \times S^q)$ gives

$$(K\pi_q(\partial_+ W_0), \theta_{\partial_+ W_0}) \cong (K\pi_q(M_0), -\theta_{M_0}) \oplus H_\varepsilon(K).$$

Recall that the form μ on S was obtained by pulling back the Wall form on $K\pi_q(M_0)$ via the surjection $\pi: S \twoheadrightarrow K\pi_q(M_0)$. By pre-composing with the projection $S \oplus \Lambda^r \twoheadrightarrow S$ onto the first component, we obtain a surjection $(\pi, 0): (S \oplus \Lambda^r, \mu \oplus 0) \twoheadrightarrow (K\pi_q(M_0), \theta_{M_0})$ of quadratic forms. Recall furthermore that we fixed an isometry $h: (S, \mu) \oplus (\Lambda^r, 0) \rightarrow (V, \theta)$, where (V, θ) is the restriction of the hyperbolic form on $H_\varepsilon(K)$ to V ; in particular $V \subseteq K \oplus K^*$. We assert that we obtain a sublagrangian of $(K\pi_q(\partial_+ W_0), \theta_{\partial_+ W_0})$ by considering the image of $S \oplus \Lambda^r$ under the map

$$(5) \quad \begin{pmatrix} \pi \\ h \end{pmatrix} := \begin{pmatrix} (\pi, 0) \\ h \end{pmatrix} : S \oplus \Lambda^r \rightarrow K\pi_q(M_0) \oplus V \subseteq K\pi_q(M_0) \oplus (K \oplus K^*) = K\pi_q(\partial_+ W_0)$$

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \mapsto \begin{pmatrix} (\pi(y_1), 0) \\ h \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \end{pmatrix}.$$

The fact that $\text{im} \begin{pmatrix} \pi \\ h \end{pmatrix}$ is a summand follows from the fact that $V \subseteq K \oplus K^*$ is a summand. Note that $\pi^* \theta_{M_0} \pi = \mu$ (by definition of μ) and $h^* \theta h = \mu \oplus 0$ (since h is an isometry). Using these facts, the following computation establishes the assertion that the image of $S \oplus \Lambda^r$ is a sublagrangian:

$$\begin{aligned} \begin{pmatrix} \pi \\ h \end{pmatrix}^* (-\theta_{M_0} \oplus H_\varepsilon) \begin{pmatrix} \pi \\ h \end{pmatrix} &= -(\pi, 0)^* \theta_{M_0} (\pi, 0) + h^* H_\varepsilon h \\ &= -(\mu \oplus 0) + (\mu \oplus 0) = 0. \end{aligned}$$

We now use the sublagrangian $\text{im} \begin{pmatrix} \pi \\ h \end{pmatrix}$ to provide data with which to attach $(q+1)$ -handles along $\partial_+ W_0$. Fix a basis $\{\bar{x}_1, \dots, \bar{x}_{s+r}\}$ for $S \oplus \Lambda^r$. Since the elements $x_i := \begin{pmatrix} \pi \\ h \end{pmatrix} (\bar{x}_i) \in K\pi_q(\partial_+ W_0)$ lie in a sublagrangian of the Wall form of $(\partial_+ W_0, \bar{\nu}_{W_0}|_{\partial_+ W_0})$ for $q \geq 3$, it follows from [Kre99, Proposition 6.7], that these classes can be represented by disjoint framed embedded spheres. We discuss the case $q = 2$ in Remark 3.11 below.

For $i = 1, \dots, s+r$, we choose pairwise disjoint framed embeddings $\phi_i: S^q \times D^q \hookrightarrow \partial_+ W_0$ representing x_i and also lifts $\omega_i \in \pi_{q+1}(B, \partial_+ W_0)$ of the x_i . We then attach $(q+1)$ -handles (which we denote by h_i^{q+1}) along the set of disjoint, framed embeddings $\phi = (\phi_1, \dots, \phi_{s+r})$ to obtain the cobordism. Extend the (B, ξ) structure from $(W_0, \bar{\nu}_0)$ using the classes $\omega = (\omega_1, \dots, \omega_{r+s})$ to obtain the (B, ξ) -bordism

$$(6) \quad W = W_{x, \pi, h, \phi, \omega} := W_0 \cup \bigcup_{i=1}^{s+r} h_i^{q+1}.$$

Since we performed surgery along elements of $K\pi_q(\partial_+ W_0)$, there is a canonical normal (B, ξ) -structure $(W, \bar{\nu})$ which extends the normal (B, ξ) -structure on W_0 ; see the end of Section 3.1. A key property of this construction is that the $x_i \in K\pi_q(\partial_+ W_0) \cong K\pi_q(M_0) \oplus K \oplus K^*$ project to a basis for $V \subseteq K \oplus K^*$. This concludes the construction of the (B, ξ) -cobordism $(W, \bar{\nu})$ based on $\partial_-(W, \bar{\nu}) = (M_0, \bar{\nu}_0)$.

Remark 3.11. We discuss the additional requirements of the 4-dimensional case $q = 2$ in this remark. In this case we pass to the topological category, and assume that the fundamental group $\pi_1(B)$ is good. In order to deduce that the homology classes $\{x_i\}$ we wish to surger

can be represented by π_1 -negligible framed embedded 2-spheres, we need to use the *sphere embedding theorem* of Freedman-Quinn [FQ90, Theorem 5.1B]. Here π_1 -negligible means that removing the embedded spheres, and therefore the process of performing surgery on them, does not change the fundamental group. This is equivalent to the existence of a collection of geometrically dual 2-spheres $\{y_i\}$ immersed in $\partial_+ W_0$, intersecting the $\{x_i\}$ transversely, and such that $x_i \cap y_j$ is a single point if $i = j$ and is empty for $i \neq j$. The existence of such a collection is guaranteed by [PRT20].

The extra input needed in the 4-dimensional case, which is not needed in high dimensions, is that the classes $\{x_i\}$ must be represented by an immersed collection of 2-spheres that admit a collection of framed algebraically dual spheres. Then the sphere embedding theorem applies to realise the $\{x_i\}$ by π_1 -negligible framed embedded 2-spheres as desired. So to extend our argument to the case $q = 2$, we must show that such a dual collection exists. Since the hyperbolic form on $K \oplus K^*$ is even, if we find the dual spheres their self-intersection numbers will be even. A cusp homotopy of an immersed 2-sphere in a 4-manifold adds a self-intersection point and changes the Euler number of the normal bundle by ± 2 . So apply these to trivialise the normal bundle, and choose a trivialisation, to obtain framed dual spheres.

It remains to find dual spheres for the spheres $\{x_i\}$. Let $f_i \in V$ be the image of x_i under the projection $K\pi_q(M_0) \oplus K \oplus K^* \rightarrow K \oplus K^*$. By the key property of the $\{x_i\}$, the $\{f_i\}$ form a basis for the direct summand $V \subseteq K \oplus K^*$. We will use that the hyperbolic form on $K \oplus K^*$ is nonsingular, and we will find dual spheres $\{g_i\}$ to the $\{f_i\}$ in $K \oplus K^*$. Write $H := K \oplus K^*$ temporarily. Since $V \subseteq H$ is a direct summand, then the dual restriction map $H^* \rightarrow V^*$ is surjective. As V^* is free we may choose a splitting $s: V^* \rightarrow H^*$. Let $\lambda_H: H \rightarrow H^*$ be the symmetrised hyperbolic form on H , which is an isomorphism, and consider $\lambda_H^{-1}(s(V^*)) \subseteq H$. Choose a basis g_1, \dots, g_n for this submodule that maps to the duals $\lambda_H(g_i) = s(f_i^*) \in H^*$. The $\{g_i\}$ are represented by immersed framed spheres in $\partial_+ W_0$, algebraically dual to the x_i . As explained above we may therefore apply the sphere embedding theorem to represent the $\{x_i\}$ by locally flat embedded spheres, to which we attach handles for the completion of the construction of the cobordism W .

The next lemma shows that $(W, \bar{\nu})$ is a $(q-1)$ -smoothing and describes its q -th homotopy group. Recall that $x \in \ell_{2q+1}(\Lambda)$ is a fixed element represented by a quasi-formation $(H_\varepsilon(K); K, V)$ such that there is a surjection $\pi: S \rightarrow K\pi_q(M_0)$, where S is a free module of rank s . Let $\text{proj}_1: K \oplus K^* \rightarrow K$ denote the canonical projection, and use \hat{h} to denote the composition

$$\hat{h}: S \oplus \Lambda^r \xrightarrow{h} V \hookrightarrow K \oplus K^* \xrightarrow{\text{proj}_1} K.$$

Lemma 3.12. *Let $(M_0^{2q}, \bar{\nu}_0)$ be a normal $(q-1)$ -smoothing and let $[x] \in \ell_{2q+1}(\Lambda)$. If the induced form of $x = (H_\varepsilon(K); K, V)$ is 0-stably isomorphic to a free Wall form (S, μ) of $(M_0, \bar{\nu}_0)$ as in (4), then the (B, ξ) -cobordism $(W, \bar{\nu})$ constructed in (6) satisfies the following.*

- (i) *The (B, ξ) -cobordism $(W, \bar{\nu})$ is a normal $(q-1)$ -smoothing.*
- (ii) *There is an isomorphism $\varphi: \pi_q(W) \xrightarrow{\cong} (\pi_q(M_0) \oplus K) / \text{Im} \left(\frac{\pi}{\hat{h}} \right)$.*
- (iii) *The isomorphism φ induces a surjection $K \twoheadrightarrow K\pi_q(W) \subseteq \pi_q(W)$.*

Proof. We start by proving the first assertion, but along the way we will prove the second. First, we check that $\bar{\nu}: W \rightarrow B$ induces an isomorphism on the first $q-1$ homotopy groups. Since we defined W_0 as $W_0 = (M_0 \times [0, 1]) \natural (\natural_{i=1}^{r+s} S^q \times D^{q+1})$, we deduce that $\pi_i(W_0) = \pi_i(M_0)$

for $i \leq q-1$. Adding $(q+1)$ -handles does not affect the first $q-1$ homotopy groups, by the Hurewicz theorem applied to the universal covers, and therefore $\pi_i(W) = \pi_i(M_0)$ for $i \leq q-1$. Since $(M_0, \bar{\nu}_0)$ is a normal $(q-1)$ -smoothing, and since $\bar{\nu}: W \rightarrow B$ restricts to $\bar{\nu}_0$ on $M_0 = \partial_- W_0$, we deduce that $\bar{\nu}_*: \pi_i(W) = \pi_i(M_0) \rightarrow \pi_i(B)$ is an isomorphism for $i \leq q-1$.

Next, we check that $\bar{\nu}: \pi_q(W) \rightarrow \pi_q(B)$ is surjective. As $W_0 = (M_0 \times [0, 1]) \natural (\natural_{i=1}^{r+s} S^q \times D^{q+1})$, we deduce that $\pi_q(W_0) = \pi_q(M_0) \oplus K$. Since $(M_0, \bar{\nu}_0)$ is a normal $(q-1)$ -smoothing, we know that $\pi_q(M_0) \rightarrow \pi_q(B)$ is surjective and therefore so is $\pi_q(W_0) \rightarrow \pi_q(B)$. Next, we compute $\pi_q(W)$. Up to homotopy equivalence, adding $(q+1)$ -handles is the same as adding $(q+1)$ -cells and in order to compute the effect on π_q of attaching we must mod out by the image of the $(s+r)$ $(q+1)$ -cells along the maps $S^q \times \{0\} \hookrightarrow \partial_+ W_0$. The homotopy classes of these attaching maps were specified by

$$\begin{pmatrix} \pi \\ \widehat{h} \end{pmatrix} : S \oplus \Lambda^r \rightarrow K\pi_q(M_0) \oplus V \subseteq K\pi_q(M_0) \oplus (K \oplus K^*) = K\pi_q(\partial_+ W_0)$$

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \mapsto \begin{pmatrix} \pi(x_1) \\ \widehat{h}(x_2) \end{pmatrix}$$

Since the induced map $\pi_q(\partial_+ W_0) \rightarrow \pi_q(W_0)$ is $\text{Id} \oplus \text{proj}_1: \pi_q(M_0) \oplus K \oplus K^* \rightarrow \pi_q(M_0) \oplus K$, it follows that $\pi_q(W) \cong (\pi_q(M_0) \oplus K) / \text{Im} \left(\begin{pmatrix} \pi \\ \widehat{h} \end{pmatrix} \right)$ as claimed. This concludes the computation of $\pi_q(W)$ and thus the proof of the second assertion of the lemma. It remains to show that $\bar{\nu}_*: \pi_q(W) \rightarrow \pi_q(B)$ is surjective. The computation of $\pi_q(W)$ shows that every element of $\pi_q(W)$ is represented by a pair $(y_1, y_2) \in \pi_q(M_0) \oplus K$ and that $\bar{\nu}_*([y_1, y_2]) = \bar{\nu}_{0*}(y_1)$. Since the (B, ξ) -structure on W was obtained by first extending the one on M_0 and imposing the trivial (B, ξ) -structure on $S^q \times S^q$, it follows that for any element $[(y_1, y_2)] \in \pi_q(W)$ we have $\bar{\nu}_*[(y_1, y_2)] = \bar{\nu}_{0*}(y_1) \in \pi_q(B)$. Since $\bar{\nu}_{0*}: \pi_1(M_0) \rightarrow \pi_q(B)$ is surjective, so is $\bar{\nu}_*: \pi_q(W) \rightarrow \pi_q(B)$, as desired.

It remains to prove the last assertion, namely that

$$\text{proj}_2: K \rightarrow K\pi_q(W) \subseteq \pi_q(W) \cong (\pi_q(M_0) \oplus K) / \text{Im} \left(\begin{pmatrix} \pi \\ \widehat{h} \end{pmatrix} \right), \quad k \mapsto [(0, k)]$$

is a surjection. Let $[(x_1, x_2)] \in K\pi_q(W)$ with $x_1 \in \pi_q(M_0)$ and $x_2 \in K$. We saw above that $\bar{\nu}_*[(y_1, y_2)] = \bar{\nu}_{0*}(y_1)$ for all $[(y_1, y_2)] \in \pi_q(W)$. This implies both that proj_2 is defined (i.e. its image lies in $K\pi_q(W)$ as asserted) and that, for $[(x_1, x_2)] \in K\pi_q(W)$, we have $x_1 \in K\pi_q(M_0)$. Next, since $\pi: S \rightarrow K\pi_q(M_0)$ is surjective, there exists $z \in S \oplus \Lambda^r$ such that $x_1 = \pi(z)$. Using our computation of $\pi_q(W)$, we obtain $[(x_1, 0)] = [(\pi(z), 0)] = [(0, \widehat{h}(w))]$ for some $w \in S \oplus \Lambda^r$. We thus obtain that $[(x_1, x_2)] = [(0, x_2 + \widehat{h}(w))]$, so that $\text{proj}_2(x_2 + \widehat{h}(w)) = [(x_1, x_2)]$. Since $[(x_1, x_2)]$ was arbitrary, we are done. \square

Proof of Theorem 3.10. We must show that the modified surgery obstruction $\Theta(W, \bar{\nu})$ of the (B, ξ) -bordism $(W, \bar{\nu})$ of (6) lies in the equivalence class $[x] = [H_\varepsilon(K); K, V] \in \ell_{2q+1}(\mathbb{Z}[\pi_1(B)])$ of the quasi-formation with which we started.

The definition of $\Theta(W, \bar{\nu})$ requires that we choose a set of generators for $K\pi_q(W)$ and represent them by framed embedded q -spheres in W . We will use $U \subseteq W$ to denote the resulting handlebody, and ψ to denote the hyperbolic quadratic form on $H_q(\partial U; \Lambda)$. Recall from Section 3.2 that the modified surgery obstruction of a $(q-1)$ -normal smoothing $(W, \bar{\nu})$ is given by

$$(7) \quad \Theta(W, \bar{\nu}) = [((H_q(\partial U; \Lambda), \psi), H_{q+1}(U, \partial U; \Lambda), H_{q+1}(W \setminus \overset{\circ}{U}; M_0 \sqcup \partial U; \Lambda))].$$

Lemma 3.12 established the existence of surjection $K \twoheadrightarrow K\pi_q(W)$. Consequently, a basis for K provides a generating set for $K\pi_q(W)$. After representing these generators by framed embedded q -spheres, we obtain $r+s$ embeddings $S^q \times D^{q+1} \hookrightarrow W$ whose images we join by thickened paths. This leads to a solid handlebody U as in (7). We can and shall assume that $U \subseteq W_0$, since U represents geometrically a basis for K .

We will show that the surgery obstruction stemming from U is given by x . For the rest of this proof, homology and cohomology groups are understood to have Λ -coefficients. Since it is clear that $((H_q(\partial U), \psi); H_{q+1}(U, \partial U)) \cong (H_\varepsilon(K); K)$, it remains to show that the map

$$H_{q+1}(W \setminus \mathring{U}; M_0 \sqcup \partial U; \Lambda) \rightarrow H_q(\partial U) \xrightarrow{\cong} K \oplus K^*$$

has image V .

Claim. *The following assertions hold.*

- (i) *The inclusion induces an isomorphism $H_{q+1}(W_0 \setminus \mathring{U}) \xrightarrow{\cong} H_{q+1}(W_0)$.*
- (ii) *There is an isomorphism $H_q(W_0 \setminus \mathring{U}) \cong K \oplus K^* \oplus H_q(M_0)$.*
- (iii) *We have $H_i(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) = 0$ for $i = q, q+1$.*

Proof of Claim. In order to prove the first assertion, we will consider the long exact sequence of the pair $(W_0, W_0 \setminus \mathring{U})$. By excision, we have $H_{q+2}(W_0, W_0 \setminus \mathring{U}) = H_{q+2}(U, \partial U) = 0$. We deduce that the inclusion induced map $\iota: H_{q+1}(W_0 \setminus \mathring{U}) \rightarrow H_{q+1}(W_0)$ is injective. Since $W_0 = (M_0 \times [0, 1]) \natural (\natural_{i=1}^{r+s} S^q \times D^{q+1}) \simeq M_0 \vee^{r+s} S^q$, we deduce that $H_{q+1}(M_0) \cong H_{q+1}(W_0)$, again induced by inclusion. This factors as $H_{q+1}(M_0) \rightarrow H_{q+1}(W_0 \setminus \mathring{U}) \xrightarrow{\iota} H_{q+1}(W_0)$, so ι is also surjective. This concludes the proof of the first assertion. We note for later that also $H_{q+1}(M_0) \cong H_{q+1}(W_0 \setminus \mathring{U})$.

In order to prove the second assertion, we consider the Mayer-Vietoris sequence associated to the decomposition $W = (W \setminus \mathring{U}) \cup U$. The portion of interest is

$$(8) \quad \begin{aligned} \cdots \rightarrow H_{q+1}(W_0 \setminus \mathring{U}) \oplus H_{q+1}(U) &\rightarrow H_{q+1}(W_0) \xrightarrow{0} H_q(\partial U) \\ &\rightarrow H_q(W_0 \setminus \mathring{U}) \oplus H_q(U) \rightarrow H_q(W_0) \rightarrow H_{q-1}(\partial U) \rightarrow \cdots \end{aligned}$$

Since the map $H_{q+1}(W_0 \setminus \mathring{U}) \xrightarrow{\cong} H_{q+1}(W_0)$ is an isomorphism (by the first assertion), we deduce that $H_{q+1}(W_0) \rightarrow H_q(\partial U)$ is the zero map as shown. Next, we use that $H_{q-1}(\partial U) = 0$ as well as the fact that $H_q(\partial U) = K \oplus K^*$ surjects onto $H_q(U) = K$ and $H_q(W_0) \cong H_q(M_0) \oplus K$ to obtain that the sequence displayed in (8) determines the following short exact sequence:

$$(9) \quad 0 \rightarrow K^* \rightarrow H_q(W_0 \setminus \mathring{U}) \rightarrow H_q(M_0) \oplus K \rightarrow 0.$$

The inclusion $M_0 \sqcup \partial U \hookrightarrow W_0 \setminus \mathring{U}$ gives rise to a splitting of this short exact sequence. This concludes the proof of the second assertion. In order to prove the third assertion, we consider the long exact sequence of the pair $(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U)$:

$$(10) \quad \begin{aligned} \cdots \rightarrow H_{q+1}(M_0 \sqcup \partial U) &\rightarrow H_{q+1}(W_0 \setminus \mathring{U}) \xrightarrow{j} H_{q+1}(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) \\ &\rightarrow H_q(M_0 \sqcup \partial U) \rightarrow H_q(W_0 \setminus \mathring{U}) \rightarrow H_q(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) \rightarrow \cdots \end{aligned}$$

We show that $H_{q+1}(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) = 0$. Note that $H_{q+1}(M_0 \sqcup \partial U) = H_{q+1}(M_0)$ and, as we showed in the first assertion, the inclusion induced map $H_{q+1}(M_0) \rightarrow H_{q+1}(W_0 \setminus \mathring{U})$ is an isomorphism. We deduce that $H_{q+1}(W_0 \setminus \mathring{U}) \rightarrow H_{q+1}(W_0)$ is an isomorphism. It

follows that the second map in (10) is surjective, from which we deduce that $j: H_{q+1}(W_0 \setminus \mathring{U}) \rightarrow H_{q+1}(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U)$ is the zero map. We now show that j is surjective, from which it follows that $H_{q+1}(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) = 0$. That j is surjective follows because $H_q(M_0 \sqcup \partial U) \rightarrow H_q(W_0 \setminus \mathring{U})$ is an isomorphism. To see this latter fact, apply the 5-lemma to the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & K^* & \xrightarrow{\text{incl}} & H_q(M_0 \sqcup \partial U) & \xrightarrow{\text{incl}} & H_q(M_0) \oplus K \longrightarrow 0 \\ & & \downarrow \text{Id} & & \downarrow \text{incl} & & \cong \downarrow \text{incl} \\ 0 & \longrightarrow & K^* & \xrightarrow{\text{incl}} & H_q(W_0 \setminus \mathring{U}) & \xrightarrow{\text{proj}} & H_q(W_0) \oplus K \longrightarrow 0, \end{array}$$

where we recall that $H_q(M_0 \sqcup \partial U) = H_q(M_0) \oplus K \oplus K^*$, and the bottom (split) exact sequence was described in (9). We have therefore shown that j is both surjective and the zero map, proving that $H_{q+1}(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) = 0$, as desired. Along the way, we have established that (10) gives the following exact sequence:

$$(11) \quad 0 \rightarrow H_q(M_0 \sqcup \partial U) \xrightarrow{\cong} H_q(W_0 \setminus \mathring{U}) \xrightarrow{0} H_q(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) \rightarrow \dots$$

It remains to prove that $H_q(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) = 0$. The isomorphism displayed in (11) implies that the map $H_q(W_0 \setminus \mathring{U}) \rightarrow H_q(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U)$ is zero, as shown. It therefore suffices to show that the next map is also zero. Extending the exact sequence (11) to the right, this is equivalent to proving that $H_{q-1}(M_0) \cong H_{q-1}(M_0 \sqcup \partial U) \rightarrow H_{q-1}(W_0 \setminus \mathring{U})$ is an injection. A Mayer-Vietoris argument shows that the map $H_{q-1}(W_0 \setminus \mathring{U}) \rightarrow H_{q-1}(W_0)$ induced by the inclusion is an isomorphism. Since we have $H_{q-1}(W_0) \cong H_{q-1}(M_0)$, we deduce that $H_{q-1}(M_0 \sqcup \partial U) \rightarrow H_{q-1}(W_0 \setminus \mathring{U})$ is an isomorphism, as claimed. We have therefore shown that $H_i(W_0 \setminus \mathring{U}, M_0 \sqcup \partial U) = 0$ for $i = q, q + 1$ and *this concludes the proof of the Claim*.

Using the Claim, we can now show that $H_{q+1}(W \setminus \mathring{U}, M_0 \sqcup \partial U)$ is sent to $V \subseteq H_q(\partial U) \cong H_\varepsilon(K)$. Consider the long exact sequence of the triple $(W \setminus \mathring{U}, W_0 \setminus \mathring{U}, M_0 \sqcup \partial U)$. The third assertion of the Claim implies that $H_{q+1}(W \setminus \mathring{U}, M_0 \sqcup \partial U) \cong H_{q+1}(W \setminus \mathring{U}, W_0 \setminus \mathring{U})$. By excision, this latter module is isomorphic to $\bigoplus_{i=1}^{r+s} H_{q+1}(D^{q+1}, S^q)$. Here, the D^{q+1} correspond to the cores of the handles h_i^{q+1} . By definition of the attaching maps of the handles h_i^{q+1} , this module coincides with V under the composition

$$\bigoplus_{i=1}^{r+s} H_{q+1}(D^{q+1}, S^q) \xrightarrow{\cong} H_{q+1}(W \setminus \mathring{U}, M_0 \sqcup \partial U) \rightarrow H_q(M_0) \oplus H_q(\partial U) \rightarrow H_q(\partial U) \cong K \oplus K^*.$$

It follows that the surgery obstruction $(W, \bar{\nu})$ is represented by $x = (H_\varepsilon(K); K, V)$, which concludes the proof of Theorem 3.10. \square

4. BOUNDARY ISOMORPHISMS

This algebraic section recalls the notion of a boundary isomorphism from [CS11]. Informally, these can be thought of as an algebraic analogue of the group of diffeomorphisms of the $(2q - 1)$ -dimensional boundary of a compact $2q$ -manifold modulo those diffeomorphisms that are isotopic to the restriction of a diffeomorphism of the $2q$ -manifold.

It is perhaps surprising that such an algebraic phenomenon should arise for closed $2q$ -manifolds, but in fact a normal $(q-1)$ -smoothing $\bar{\nu}: M \rightarrow B$ of a $2q$ -manifold M , satisfying certain algebraic conditions, induces a splitting of the intersection quadratic form as a union over two forms defined in terms of B and $\bar{\nu}$; details are given in Section 7.

In Sections 4.1 and 4.2, we review stable isomorphisms of split quadratic formations and homotopies between them, culminating in the definition of the boundary isomorphism set, while the gluing process for forms is reviewed in Section 4.3.

We briefly recall our general strategy for the proof of Theorem 1.1: elements of the boundary isomorphism sets will be the invariants we use to distinguish $4k$ -manifolds. In Section 7 we will show, via the notion of primitive embeddings to appear in Section 6, that the realisation results from Section 3 allow us to obtain many infinitely many nontrivial values of the boundary isomorphism invariant.

4.1. Stable isomorphisms of split formations. In this subsection, we review the notion of a split quadratic formation. References include [Ran81, Section 1.6], [Ran02, Section 12.5], as well as [CS11, Section 4.1].

Definition 4.1. A *simple split ε -quadratic formation* $(F, \binom{\gamma}{\delta})(G, \theta)$ consists of a based R -module F , a $(-\varepsilon)$ -quadratic form (G, θ) , and morphisms $\gamma: G \rightarrow F$ and $\delta: G \rightarrow F^*$ such that

- (i) the *Hessian* θ satisfies $\gamma^*\delta = \theta - \varepsilon\theta^*: G \rightarrow G^*$;
- (ii) the map $\binom{\gamma}{\delta}: (G, 0) \rightarrow H_\varepsilon(F)$ is the inclusion of a simple lagrangian.

The first examples of split quadratic formations are *the trivial split quadratic formations*:

$$(F, F^*) := (F, \binom{0}{1}(F^*, 0)).$$

Next, observe that for a split quadratic formation $(F, \binom{\gamma}{\delta})(G, \theta)$, the triple $(H_\varepsilon(F); F, \binom{\gamma}{\delta}G)$ is a simple quadratic formation. In fact, quadratic formations also give rise to split quadratic formations: writing γ and δ for the inclusions of the lagrangians, the map $\gamma^*\delta: G \rightarrow G^*$ is a $(-\varepsilon)$ -symmetric form and admits a quadratic refinement [Ran02, Section 12.5]. The choice of this quadratic refinement is remembered in a split quadratic formation, but not in a formation.

Definition 4.2. An *isomorphism* (α, β, ν) between two split ε -quadratic formations $(F_1, \binom{\gamma_1}{\delta_1})(G_1, \theta_1)$ and $(F_2, \binom{\gamma_2}{\delta_2})(G_2, \theta_2)$ consists of isomorphisms $\alpha: F_1 \rightarrow F_2$ and $\beta: G_1 \rightarrow G_2$, as well as a $(-\varepsilon)$ -quadratic form (F_1^*, ν) such that

- (i) the equality $\theta_1 + \delta_1^*\nu\delta_1 = \beta_1^*\theta_2\beta_1$ holds in $Q_{-\varepsilon}(G_1)$;
- (ii) the map $f := \begin{pmatrix} \alpha & \alpha(\nu - \varepsilon\nu^*) \\ 0 & (\alpha^*)^{-1} \end{pmatrix}: H_\varepsilon(F_1) \rightarrow H_\varepsilon(F_2)$ defines an isomorphism of quadratic formations or, equivalently, the following diagram commutes:

$$\begin{array}{ccc} G_1 & \xrightarrow{\beta} & G_2 \\ \downarrow \binom{\gamma_1}{\delta_1} & & \downarrow \binom{\gamma_2}{\delta_2} \\ F_1 \oplus F_1^* & \xrightarrow{f, \cong} & F_2 \oplus F_2^* . \end{array}$$

A *stable isomorphism* of split ε -quadratic formations is an isomorphism

$$(\alpha, \beta, \nu): (F_1, \binom{\gamma_1}{\delta_1} (G_1, \theta_1)) \oplus (H_1, H_1^*) \xrightarrow{\cong} (F_2, \binom{\gamma_2}{\delta_2} (G_2, \theta_2)) \oplus (H_2, H_2^*)$$

for some f.g. free based R -modules H_1 and H_2 .

We use \cong to denote the isomorphism relation and \cong_s to denote stable isomorphism relation. A split ε -quadratic formation is *trivial* if it is isomorphic to $(F, F^*) = (F, \binom{0}{1} (F^*, 0))$. The *composition* of stable isomorphisms (α', β', ν') and (α, β, ν) is defined as

$$(12) \quad (\alpha', \beta', \nu') \circ (\alpha, \beta, \nu) := (\alpha' \alpha, \beta' \beta, \nu + \alpha^{-1} \nu' \alpha^{-*}).$$

The *inverse* of a stable isomorphism (α, β, ν) is defined as $(\alpha^{-1}, \beta^{-1}, -\alpha \nu \alpha^*)$. The *identity* on a split ε -quadratic formation x is $(1, 1, 0)$. The next example shows that isometries of quadratic forms give rise to isomorphisms of their so-called boundary split formations.

Example 4.3. The *boundary* of a $(-\varepsilon)$ -quadratic form (P, ψ) is the split ε -quadratic formation

$$\partial(P, \psi) := (P, \binom{1}{\psi - \varepsilon \psi^*} (P, \psi)).$$

The *boundary of an isometry* $h: (P, \psi) \rightarrow (P', \psi')$ of simple quadratic forms is the isomorphism

$$\partial h = (h, h, 0): \partial(P, \psi) \rightarrow \partial(P', \psi').$$

4.2. Boundary isomorphisms. In this subsection, we review homotopies between stable isomorphisms of formations. References include [CS11, Section 4.1] and [Ran81, Section 1.6].

Here is a short motivation for the definition. Namely, given two isomorphisms of split quadratic formations

$$(\alpha, \beta, \nu), (\alpha', \beta', \nu'): (F_1, \binom{\gamma_1}{\delta_1} (G_1, \theta_1)) \rightarrow (F_2, \binom{\gamma_2}{\delta_2} (G_2, \theta_2)),$$

we will define a homotopy $\Delta: (\alpha, \beta, \nu) \rightarrow (\alpha', \beta', \nu')$. There is a one to one correspondence between stable isomorphism classes of split quadratic formations and chain equivalence classes of 1-dimensional split quadratic complexes [Ran81, Proposition 1.6.4]. Regardless of the definition of a quadratic complex, a chain complex corresponding to the split quadratic formation $(F, \binom{\gamma}{\delta} (G, \theta))$ is $0 \rightarrow F \xrightarrow{\delta^*} G^* \rightarrow 0$. The second condition of Definition 4.2 implies that the isomorphisms (α, β, ν) and (α', β', ν') of split quadratic formations fit into the following diagram:

$$(13) \quad \begin{array}{ccccccc} 0 & \longrightarrow & F_1 & \xrightarrow{\delta_1^*} & G_1^* & \longrightarrow & 0 \\ & & \alpha \downarrow & \swarrow \Delta & \downarrow & \searrow \beta'^{-*} & \\ & & F_2 & \xrightarrow{\delta_2^*} & G_2^* & \longrightarrow & 0. \end{array}$$

In (13), the dashed map Δ suggests the following definition for a homotopy between two isomorphisms of split formations.

Definition 4.4. A *homotopy* of isomorphisms (α, β, ν) and (α', β', ν') between split ε -quadratic formations $(F_1, \binom{\gamma_1}{\delta_1} (G_1, \theta_1))$ and $(F_2, \binom{\gamma_2}{\delta_2} (G_2, \theta_2))$ consists of an R -linear map $\Delta: G_1^* \rightarrow F_2$ which satisfies

$$(i) \quad \beta'^{-*} - \beta^{-*} = \delta_2^* \Delta,$$

- (ii) $\alpha' - \alpha = \Delta\delta_1^*$,
- (iii) $\alpha'\nu'\alpha'^* - \alpha\nu\alpha^* = (\varepsilon\alpha'\gamma_1 + \Delta\theta)\Delta^* \in Q_{-\varepsilon}(F_2^*)$.

Next, we move on to stable homotopies. Briefly, two stable isomorphisms are stably homotopic if they become homotopic in a common stabilization. The formal definition reads as follows.

Definition 4.5. Given trivial split formations u_1, u_2, v_1, v_2 , two isomorphisms $f_1: x \oplus u_1 \rightarrow y \oplus v_1$ and $f_2: x \oplus u_2 \rightarrow y \oplus v_2$ are *stably homotopic* if the following conditions are satisfied:

- (i) there are based modules R_1, R_2, P, Q and isomorphisms

$$\begin{aligned} u_1 \oplus (R_1, R_1^*) &\xrightarrow{g_1, \cong} (P, P^*) \xrightarrow{g_2, \cong} u_2 \oplus (R_2, R_2^*), \\ v_1 \oplus (R_1, R_1^*) &\xrightarrow{h_1, \cong} (Q, Q^*) \xleftarrow{h_2, \cong} v_2 \oplus (R_2, R_2^*); \end{aligned}$$

- (ii) there is a homotopy

$$\tilde{f}_1 \simeq \tilde{f}_2: x \oplus (P, P^*) \rightarrow y \oplus (Q, Q^*),$$

where the isomorphisms \tilde{f}_1, \tilde{f}_2 are $\tilde{f}_i := (\text{Id}_y \oplus h_i) \circ (f_i \oplus \text{Id}_{(R_i, R_i^*)}) \circ (\text{Id}_x \oplus g_i)$ for $i = 1, 2$:

$$\begin{array}{ccc} x \oplus (P, P^*) & \xrightarrow{\text{Id}_x \oplus g_i} & x \oplus u_i \oplus (R_i, R_i^*) \\ \vdots \downarrow \tilde{f}_i & & \downarrow f_i \oplus \text{Id}_{(R_i, R_i^*)} \\ y \oplus (Q, Q^*) & \xleftarrow{\text{Id}_y \oplus h_i} & y \oplus v_i \oplus (R_i, R_i^*). \end{array}$$

Given simple split ε -quadratic formations y and z , and using $[f]$ to denote the stable homotopy class of a stable isomorphism $f: y \cong_s z$, we set

$$\text{Iso}(y, z) := \{[f] \mid f \text{ is a stable isomorphism from } y \text{ to } z\}.$$

For quadratic forms (V, θ) and (V', θ') , there are left actions of $\text{Aut}(V, \theta)$ and $\text{Aut}(V', \theta')$ on the set $\text{Iso}(\partial(V, \theta), \partial(V', \theta'))$, respectively given by $g \cdot [f] := f \circ \partial g^{-1}$ and $h \cdot [f] := \partial h \circ f$. Here, recall from Example 4.3 that ∂g^{-1} and ∂h refer to the boundaries of the isometries g^{-1} and h . Since these actions commute, we obtain an action of $\text{Aut}(V, \theta) \times \text{Aut}(V', \theta')$ by setting $(g, h) \cdot f = \partial h \circ f \circ \partial g^{-1}$. Modding out by these left actions leads to the main definition of this section.

Definition 4.6. Given quadratic forms (V, θ) and (V', θ') , the *one-sided boundary isomorphism sets* and *boundary isomorphism set* are respectively defined as

$$\begin{aligned} \ell\text{Iso}(\partial(V, \theta), \partial(V', \theta')) &:= \text{Iso}(\partial(V, \theta), \partial(V', \theta')) / \text{Aut}(V, \theta), \\ \text{rIso}(\partial(V, \theta), \partial(V', \theta')) &:= \text{Iso}(\partial(V, \theta), \partial(V', \theta')) / \text{Aut}(V', \theta'), \\ \text{bIso}(\partial(V, \theta), \partial(V', \theta')) &:= \text{Iso}(\partial(V, \theta), \partial(V', \theta')) / \text{Aut}(V, \theta) \times \text{Aut}(V', \theta'). \end{aligned}$$

Informally, non-trivial elements of $\text{bIso}(\partial(V, \theta), \partial(V', \theta'))$ consist of stable isomorphisms of split boundary formations that are not induced by isometries of forms. The boundary isomorphism set was introduced in [CS11, Section 5]. We discuss the reasons why we consider ℓIso and rIso .

Remark 4.7. As we saw before Example 4.3, the composition of boundary isomorphisms gives rise to a composition

$$\text{Iso}(\partial(V, \theta), \partial(V', \theta')) \times \text{Iso}(\partial(V', \theta'), \partial(V'', \theta'')) \rightarrow \text{Iso}(\partial(V, \theta), \partial(V'', \theta'')).$$

It follows that $\text{Aut}(\partial(V, \theta))$ is a group. On the other hand, since $\text{Aut}(V, \theta)$ need not be a normal subgroup of $\text{Aut}(\partial(V, \theta))$, the composition does *not* descend to $\text{bAut}(\partial(V, \theta))$ in general. Nevertheless, composition does induce maps

$$\begin{aligned} \ell\text{Iso}(\partial(V, \theta), \partial(V', \theta')) \times \text{Iso}(\partial(V', \theta'), \partial(V'', \theta'')) &\rightarrow \ell\text{Iso}(\partial(V, \theta), \partial(V'', \theta'')), \\ \text{rIso}(\partial(V, \theta), \partial(V', \theta')) \times \text{rIso}(\partial(V', \theta'), \partial(V'', \theta'')) &\rightarrow \text{rIso}(\partial(V, \theta), \partial(V'', \theta'')). \end{aligned}$$

Another more topological motivation for the appearance of ℓIso will feature in Section 7 below. We conclude with a remark on some identifications we will often make in the sequel.

Remark 4.8. As mentioned in [CS11, Remark 5.6], if there exist isometries $k: (V_1, \theta_1) \rightarrow (V_2, \theta_2)$ and $h: (V'_1, \theta'_1) \rightarrow (V'_2, \theta'_2)$, then the assignments $[f] \mapsto [f \circ \partial k^{-1}]$, $[f] \mapsto [\partial h \circ f]$, $[f] \mapsto [\partial h \circ f \circ \partial k^{-1}]$ respectively give rise to bijections

$$\begin{aligned} \ell\text{Iso}(\partial(V_1, \theta_1), \partial(V'_1, \theta'_1)) &\xrightarrow{\cong} \ell\text{Iso}(\partial(V_2, \theta_2), \partial(V'_1, \theta'_1)), \\ \text{rIso}(\partial(V_1, \theta_1), \partial(V'_1, \theta'_1)) &\xrightarrow{\cong} \text{rIso}(\partial(V_1, \theta_1), \partial(V'_2, \theta'_2)), \\ \text{bIso}(\partial(V_1, \theta_1), \partial(V'_1, \theta'_1)) &\xrightarrow{\cong} \text{bIso}(\partial(V_2, \theta_2), \partial(V'_2, \theta'_2)) \end{aligned}$$

that are independent of the choices of the isometries h and k . For instance, if $h': (V'_1, \theta'_1) \rightarrow (V'_2, \theta'_2)$ is another isometry, then $\partial h' \circ f = \partial(h' \circ h^{-1}) \circ \partial h \circ f$ agrees with $\partial h \circ f$ in $\text{rIso}(\partial(V_1, \theta_1), \partial(V'_2, \theta'_2))$.

Thus in this situation, given $f: \partial(V_1, \theta_1) \xrightarrow{\cong} \partial(V'_1, \theta'_1)$, in order to compare boundary isomorphisms, we will sometimes write $[f] \in \text{bIso}(\partial(V_2, \theta_2), \partial(V'_2, \theta'_2))$, in which case the class we are really referring to is $[\tilde{f}]$, where $\tilde{f} = \partial h \circ f \circ \partial k^{-1}$ for some isometries $h: (V'_1, \theta'_1) \rightarrow (V'_2, \theta'_2)$ and $k: (V_1, \theta_1) \rightarrow (V_2, \theta_2)$, and the choice of isometries is immaterial.

4.3. Gluing of forms. In this subsection, we recall how two quadratic forms can be glued along a stable isomorphism of their boundary formations. References for this construction include [Ran81, Section 1.7] and [CS11, Section 4.2].

Fix ε -quadratic forms (V, θ) and (V', θ') and let $\lambda = \theta + \varepsilon\theta^*$ and $\lambda' = \theta' + \varepsilon\theta'^*$ be the corresponding symmetric forms. The following lemma provides a characterisation of stable isomorphisms $\partial(V, \theta) \cong_s \partial(V', \theta')$; a proof can be found in [CS11, Lemma 4.3].

Lemma 4.9. *Let P and P' be based modules. Let $\alpha: V \oplus P \rightarrow V' \oplus P'$ and $\beta: V \oplus P^* \rightarrow V' \oplus P'^*$ be simple isomorphisms and let $\nu \in Q_\varepsilon(V'^* \oplus P'^*)$. The following statements are equivalent:*

(1) *the triple (α, β, ν) defines a stable isomorphism*

$$(\alpha, \beta, \nu): \partial(V, \theta) \oplus (P, P^*) \xrightarrow{\cong} \partial(V', \theta') \oplus (P', P'^*);$$

(2) *there are homomorphisms a, b, s and a_1, b_1, a_3 such that $\alpha = \begin{pmatrix} a & a_1 \\ \varepsilon b_1^* \lambda & a_3^* \end{pmatrix}$, $\beta^{-1} = \begin{pmatrix} b & b_1 \\ a_1^* \lambda' & a_3^* \end{pmatrix}$*

and $\alpha\nu\alpha^ = \begin{pmatrix} s & -\varepsilon ab_1 \\ 0 & -b_1^* \theta b_1 \end{pmatrix} \in Q_\varepsilon(V'^* \oplus P'^*)$ which satisfy*

$$(i) \ 1 = ab + (s^* + \varepsilon s)\lambda',$$

- (ii) $a^*\lambda' = \lambda b$,
- (iii) $\theta' = b^*\theta b + \lambda'^*s\lambda' \in Q_\varepsilon(V')$. □

From now on, we refer to the homomorphisms a, b, s as the a , b , and s components of the stable isomorphism f . We give some examples of these components.

Example 4.10. Given an isometry $h: (V, \theta) \rightarrow (V', \theta')$, the components of $\partial h = (h, h, 0)$ are given by $a_{\partial h} = h, b_{\partial h} = h^{-1}$ and $s_{\partial h} = 0$. Consequently, if f is a stable isomorphism with components a, b, s , and if $h_1: (V, \theta) \rightarrow (V, \theta), h_2: (V', \theta') \rightarrow (V', \theta')$ are isometries, then the stable isomorphism $(h_1, h_2) \cdot f = \partial h_2 \circ f \circ \partial h_1^{-1}$ has components $h_2 a h_1^{-1}, h_1 b h_2^{-1}$ and $h_2 s h_2^*$.

In the sequel, we will often use the following criterion for stable homotopy.

Remark 4.11. Two stable isomorphisms $(\alpha, \beta, \nu), (\alpha', \beta', \nu'): \partial(V_1, \theta_1) \xrightarrow{\cong_s} \partial(V_2, \theta_2)$ are stably homotopic if and only if there is a $\Delta \in \text{Hom}_R(V^*, V')$ such that $a' - a = \Delta\lambda, b' - b = \Delta^*\lambda'^*$, and $s' - s = (-\varepsilon a' \Delta^* - \Delta\theta)\Delta^*$ [CS11, Lemma 4.3 ii]. In particular, if two stable isomorphisms have the same a, b and s components, then they are stably homotopic via $\Delta = 0$.

The main definition of this subsection is the following.

Definition 4.12. The *union* of two quadratic forms (V, θ) and (V', θ') along a stable isomorphism $f: \partial(V, \theta) \oplus (P, P^*) \rightarrow \partial(V', \theta') \oplus (P', P'^*)$ is defined to be the quadratic form

$$(V, \theta) \cup_f (V', -\theta') := \left(V \oplus V'^*, \begin{pmatrix} \theta & 0 \\ \varepsilon a & -s \end{pmatrix} \right),$$

where a and s denote the components of f as described in Lemma 4.9.

Up to isometry, the union $(V, \theta) \cup_f (V', -\theta')$ only depends on the stable homotopy class of the stable isomorphism f [CS11, Lemma 4.6 iii]. In fact, more is known: $(V, \theta) \cup_f (V', -\theta')$ only depends on the class of f in $\text{bIso}(\partial(V, \theta), \partial(V', \theta'))$ [CS11, Lemma 4.6 iv]. The next lemma describes isometric inclusions of (V, θ) and $(V', -\theta')$ into $(V, \theta) \cup_f (V', -\theta')$. These maps will be used frequently in the upcoming subsections.

Lemma 4.13. *Let $f: \partial(V, \theta) \oplus (P, P^*) \rightarrow \partial(V', \theta') \oplus (P', P'^*)$ be an isomorphism of split quadratic formations, with components a, b, s . The following maps are split isometric injections:*

- (i) $j_f := \begin{pmatrix} 1 \\ 0 \end{pmatrix}: (V, \theta) \rightarrow (V, \theta) \cup_f (V', -\theta')$,
- (ii) $j'_f := \begin{pmatrix} b \\ -\lambda' \end{pmatrix}: (V', -\theta') \rightarrow (V, \theta) \cup_f (V', -\theta')$.

Furthermore, (V, θ) and $(V', -\theta')$ are complementary inside the union: $(V, \theta)^\perp = j'(V', -\theta')$.

The gluing construction described in Definition 4.12 produces a nonsingular quadratic form starting from a boundary isomorphism. Combining this with Lemma 4.13, we conclude that two split isometric injections can be extracted from any boundary isomorphism. The next proposition provides a converse of sorts: starting from a split isometric embedding, it constructs a boundary isomorphism. We refer to [CS11, Proposition 4.8] for a proof.

Proposition 4.14. *Given a split isometric injection $j: (V, \theta) \hookrightarrow (M, \psi)$ of quadratic forms (V, θ) and (M, ψ) , one can construct*

- (i) *an isomorphism $f_j: \partial(V, \theta) \oplus (M', M'^*) \rightarrow \partial(V^\perp, -\theta^\perp) \oplus (M, M^*)$ that is well defined up to homotopy, where $M' := V^\perp \oplus V^{\perp*}$;*

(ii) an isometry $r_j: (M, \psi) \xrightarrow{\cong} (V, \theta) \cup_{f_j} (V^\perp, \theta^\perp)$.

Furthermore, f_j and r_j are natural with respect to isometries of such pairs of forms.

For later use, we recall the explicit formula for the isomorphism f_j of Proposition 4.14. Following [CS11, proof of Proposition 4.8], we write $(M', \psi') := H_\varepsilon(V^\perp)$, and let ϕ' be the ε -symmetrization of ψ' . The formula for the stable isomorphism $f_j = (\alpha_{f_j}, \beta_{f_j}, \nu_{f_j})$ is given by

$$(14) \quad f_j: \partial(V, \theta) \oplus (M', M'^*) \rightarrow \partial(V^\perp, -\theta^\perp) \oplus (M, M^*)$$

$$f_j := \left(\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \oplus \begin{pmatrix} 1 \\ \phi \\ -\phi^{-*}\psi\phi^{-1} \end{pmatrix} \right) \circ \partial h \circ \left(\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \oplus \begin{pmatrix} 1 \\ \phi'^{-1} \\ \phi'^{-*}\psi'^*\phi'^{-1} \end{pmatrix} \right),$$

To decode this a little, recalling that $\partial h = (h, h, 0)$ and the composition law from (12), the notation implies that $f_j = (\alpha_{f_j}, \beta_{f_j}, \nu_{f_j})$ is given by

$$\alpha_{f_j} = ((1) \oplus (1)) \circ h \circ ((1) \oplus (1)) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \circ h \circ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

and similarly for the β and ν components. Here the isometry h is given by

$$(15) \quad h: (V, \theta) \oplus H_\varepsilon(V^\perp) \rightarrow (V^\perp, -\theta^\perp) \oplus (M, \psi)$$

$$h := \begin{pmatrix} 1 & 0 & 0 \\ j^\perp & \sigma & j \end{pmatrix} \begin{pmatrix} -\sigma^*\phi^*j & 1 & -\sigma^*\psi^*\sigma \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}.$$

In this formula, $j^\perp: V^\perp \rightarrow M$ is the inclusion of V^\perp , and $\sigma: (V^\perp)^* \rightarrow M$ is a splitting of the short exact sequence

$$(16) \quad 0 \rightarrow V \xrightarrow{j} M \xrightarrow{j^{\perp*}\phi} (V^\perp)^* \rightarrow 0$$

with $(j^{\perp*}\phi)\sigma = \text{Id}$. This implies that $\tilde{\sigma} := j^{-1}(1 - \sigma j^{\perp*}\phi)$ is a splitting for j , since $j^{\perp*}\phi j = 0$, and so $j^{-1}(1 - \sigma j^{\perp*}\phi)j = \text{Id}$. Here the notation j^{-1} is justified since the image of $(1 - \sigma j^{\perp*}\phi)$ is a subset of $\ker(j^{\perp*}\phi) = \text{Im}(j)$.

The next proposition describes the a, b, s components of f_j .

Proposition 4.15. *Let $j: (V, \theta) \hookrightarrow (M, \psi)$ be a split isometric injection. The a, b, s -components of the isomorphism f_j described in Proposition 4.14 are*

$$a = -\sigma^*\phi^*j,$$

$$b = -\tilde{\sigma}j^\perp,$$

$$s = -\sigma^*\psi\sigma.$$

Proof. By definition, the b component of a stable isomorphism (α, β, ν) can be read off from the $(1, 1)$ -entry of β^{-1} . Write f_j as $f_j := J \circ \partial h \circ K$, where $J = (\alpha_J, \beta_J, \nu_J)$ and $K = (\alpha_K, \beta_K, \nu_K)$ are the stable isomorphisms that appear in (14). Since $\beta_J = \begin{pmatrix} 1 & 0 \\ 0 & \phi \end{pmatrix}$ and $\beta_K = \begin{pmatrix} 1 & 0 \\ 0 & \phi'^{-1} \end{pmatrix}$, the b component of f_j is equal to the b component of ∂h . In order to understand the b component of $\partial h = (h, h, 0)$, we compute h^{-1} . To achieve this, recall from (16) that we have the split exact sequence

$$(17) \quad 0 \rightarrow (V, \theta) \xrightarrow{j} (M, \psi) \xrightarrow{j^{\perp*}\phi} (V^\perp, \theta^\perp)^* \rightarrow 0,$$

that we used σ to denote a splitting of $j^{\perp*}\phi$, and that $\tilde{\sigma} := j^{-1}(1 - \sigma j^{\perp*}\phi)$ is a splitting for j . Write the isometry h as $h = AB$, where A and B are the matrices that appear in (15). A direct computation shows that both A and B are invertible, and

$$\begin{aligned} h^{-1} = B^{-1}A^{-1} &= \begin{pmatrix} -\sigma^*\phi^*j & 1 & -\sigma^*\psi^*\sigma \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 0 & 0 \\ j^{\perp} & \sigma & j \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 0 & 0 & 1 \\ 1 & \sigma^*\psi^*j & \sigma^*\phi^*j \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\lambda^{\perp} & j^{\perp*}\phi \\ -j^{-1}(1 - \sigma j^{\perp*}\phi)j^{\perp} & j^{-1}(1 - \sigma j^{\perp*}\phi) \end{pmatrix}. \end{aligned}$$

Although

$$A = \begin{pmatrix} 1 & 0 & 0 \\ j^{\perp} & \sigma & j \end{pmatrix} : V^{\perp} \oplus (V^{\perp})^* \oplus V \rightarrow V^{\perp} \oplus M$$

is written as a non-square matrix, in fact $M \cong (V^{\perp})^* \oplus V$, so the domain and codomain of A coincide, and moreover A is an isomorphism. Recall that the b component of f_j is equal to the b component of ∂h . Since the b component of $\partial h = (h, h, 0)$ is equal to the $(1, 1)$ -entry h_{11}^{-1} of h^{-1} , we deduce that

$$(18) \quad b_{f_j} = b_{\partial h} = h_{11}^{-1} = -j^{-1}(1 - \sigma j^{\perp*}\phi)j^{\perp} = -\tilde{\sigma}j^{\perp},$$

where the last equality uses the definition of $\tilde{\sigma}$. Next, we compute the a component of f_j . We write f_j as $f_j := J \circ \partial h \circ K$, where the stable isomorphisms $J = (\alpha_J, \beta_J, \nu_J)$ and $K = (\alpha_K, \beta_K, \nu_K)$ are those which appear in (14). Note that α_J and α_K are both identity matrices. As $\partial h = (h, h, 0)$, the definition of the composition of stable isomorphisms gives $\alpha_{f_j} = \alpha_J \circ \alpha_{\partial h} \circ \alpha_K = \alpha_{\partial h} = h$. Thus the a component of f_j is given by the $(1, 1)$ entry h_{11} of h . The definition of h from (15) implies that $h_{11} = -\sigma^*\phi^*j$, and thus

$$a_{f_j} = h_{11} = -\sigma^*\phi^*j.$$

Finally, we compute the s component of f_j . Recalling the composition law $\nu' \circ \nu = \nu + \alpha^{-1}\nu'\alpha^{-*}$ from (12), we deduce that $\nu_{J \circ \partial h} = \nu_{\partial h} + \alpha_{\partial h}^{-1}\nu_J\alpha_{\partial h}^{-*}$. Apply $\nu_{\partial h} = 0$ and $\alpha_{\partial h} = h$ to obtain $\nu_{J \circ \partial h} = h^{-1}\nu_Jh^{-*}$. Use the composition law a second time, observing that $\alpha_K = \text{Id}$, to obtain

$$\nu_{f_j} = \nu_{J \circ \partial h \circ K} = \nu_K + \alpha_K^{-1}\nu_{J \circ \partial h}\alpha_K^{-*} = \nu_K + h^{-1}\nu_Jh^{-*}.$$

By definition, the s component of f_j is obtained as the $(1, 1)$ -entry of

$$\alpha_{f_j}\nu_{f_j}\alpha_{f_j}^* = h(\nu_K + h^{-1}\nu_Jh^{-*})h^* = h\nu_Kh^* + \nu_J.$$

A glance at (14) shows that the $(1, 1)$ -entry of ν_J is zero, so the s component of f_j coincides with the $(1, 1)$ -entry of $h\nu_Kh^*$. In order to compute ν_K , we use that $\psi' = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $\phi' = \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}$ to deduce that $\phi'^{-*}\psi'\phi'^{-1} = \begin{pmatrix} 0 & 0 \\ \varepsilon & 0 \end{pmatrix}$. It follows from (14) and this computation that

$$(19) \quad \nu_K = \begin{pmatrix} 0 & 0 \\ 0 & \phi'^{-*}\psi'\phi'^{-1} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \varepsilon & 0 \end{pmatrix}.$$

Writing the entries of h as h_{ij} , a brief computation using (15) and (19) now implies that

$$s_{f_j} = (h\nu_Kh^*)_{11} = \varepsilon h_{13}h_{12}^* = -\varepsilon(\sigma^*\psi^*\sigma) = -\sigma^*\psi\sigma.$$

We have therefore obtained the announced values for the a, b, s components of f_j . \square

From now on, we will often denote quadratic forms by v instead of (V, θ) . Next, we give a concrete example of Proposition 4.15 in the case where $m := (M, \psi)$ is a union.

Example 4.16. We apply Proposition 4.15 to the situation where $m = v \cup_f -v'$ for some stable isomorphism $f: \partial v \xrightarrow{\cong} \partial v'$, and where $j = \begin{pmatrix} 1 \\ 0 \end{pmatrix}: v \hookrightarrow v \cup_f -v'$. Using the notation of Proposition 4.15, we have $\psi := \begin{pmatrix} \theta & 0 \\ \varepsilon a & -s \end{pmatrix}$ with symmetrisation $\phi := \begin{pmatrix} \lambda & \varepsilon a^* \\ a & -(s + \varepsilon s^*) \end{pmatrix}$, and $j' := \begin{pmatrix} b \\ -\lambda' \end{pmatrix}$. We deduce that $j^\perp = \begin{pmatrix} b \\ -\lambda' \end{pmatrix}$ and $\lambda^\perp = j^{\perp*} \psi j^\perp = -\theta'$. Consequently, a splitting of $j^{\perp*} \phi = \begin{pmatrix} 0 & 1 \end{pmatrix}$ is $\sigma = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and the corresponding splitting of j is $\tilde{\sigma} = \begin{pmatrix} 1 & 0 \end{pmatrix}$. Therefore, we obtain

$$\begin{aligned} a_{f_j} &= -\sigma^* \phi^* j = -\begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} \lambda & \varepsilon a^* \\ a & -(s + \varepsilon s^*) \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = -a, \\ b_{f_j} &= -\tilde{\sigma} j^\perp = -\begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} b \\ -\lambda' \end{pmatrix} = -b, \\ s_{f_j} &= -\sigma^* \psi \sigma = -\begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} \theta & 0 \\ \varepsilon a & -s \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = s. \end{aligned}$$

Consider the isometries $(-1): v \rightarrow v$ and $(-1)': v' \rightarrow v'$ that are induced by multiplication by -1 . The boundaries of these isometries are $(-1, -1, 0)$. The stable isomorphism f_j has the same components as $(\partial(-1)' \oplus 1) \circ f$ and $f \circ (\partial(-1) \oplus 1)^{-1}$. By Remark 4.11, these stable isomorphisms are thus pairwise stably homotopic. In particular, while f and f_j might not agree in $\text{Iso}(\partial v, \partial v')$, they do agree in $\text{rIso}(\partial v, \partial v')$ and $\text{lIso}(\partial v, \partial v')$. In particular, they also agree in $\text{bIso}(\partial v, \partial v')$.

Given forms v, v' , we are interested in the set of $[f] \in \text{Iso}(\partial v, \partial v')$ such that the algebraic union $v \cup_f -v'$ is a fixed isometry type m :

$$\text{Iso}_m(\partial v, \partial v') := \{[f] \in \text{Iso}(\partial v, \partial v') \mid v \cup_f -v' \cong m\}.$$

Here, observe that if $m \cong m'$ then, by definition, we have $\text{Iso}_m(\partial v, \partial v') = \text{Iso}_{m'}(\partial v, \partial v')$: postcomposing the isomorphism $v \cup_f -v' \cong m$ with the isomorphism $m \cong m'$, we deduce that $v \cup_f -v' \cong m'$. The analogous definitions and remarks can be made for rIso , lIso and bIso . We conclude this subsection by recalling a concrete example over the integers.

Example 4.17. Given a product $q = p_1 \cdots p_k$ of k distinct odd primes, [CS11, Section 6.3] implies that $\text{bAut}(\partial(\mathbb{Z}, q))$ is a group and $\text{bAut}(\partial(\mathbb{Z}, q)) \cong (\mathbb{Z}/2\mathbb{Z})^{k-1}$. Since we are working over \mathbb{Z} , any twisted double $w \cup_f -w$ is hyperbolic [Ran98, Proposition 3.11 and Remark 29.18], and we obtain $\text{bAut}_{H_+(\mathbb{Z})}(\partial(\mathbb{Z}, q)) = \text{bAut}(\partial(\mathbb{Z}, q)) \cong (\mathbb{Z}/2\mathbb{Z})^{k-1}$. We will prove a generalisation of this result in Example 6.6 below.

5. INFINITE BOUNDARY AUTOMORPHISM SETS

The aim of this section, which is again entirely algebraic, is to provide examples of infinite boundary automorphism sets. For our topological applications, we must find infinitely many $(+1)$ -quadratic forms (V, θ) such that the set $\text{bAut}_{H_+(\mathbb{Z}[t^{\pm 1}])}(\partial(V, \theta))$ is infinite. Here recall that this set consists of those (equivalence classes of) boundary isomorphisms whose double is hyperbolic over $\mathbb{Z}[\mathbb{Z}] = \mathbb{Z}[t^{\pm 1}]$. By *doubling*, we mean the map $\kappa(f) = (V, \theta) \cup_f (V, -\theta)$ that takes a boundary isomorphism to the corresponding algebraic union.

Throughout this section we make the following assumptions:

- (i) π is an infinite abelian group (e.g. \mathbb{Z});
- (ii) $\Lambda := \mathbb{Z}[\pi]$ is its group ring;
- (iii) our forms (V, θ) are $(+1)$ -quadratic ($\varepsilon = 1$) over a free module of rank 1, so that we will have $V = \Lambda$, $\lambda = \bar{\lambda} = \theta + \bar{\theta} \in \Lambda$, $\theta \in \Lambda$.

Note that under these assumptions, we have $\text{bAut}(V, \theta) = \text{rAut}(V, \theta) = \text{lAut}(V, \theta)$. Furthermore, given an integer $p \in \mathbb{Z}$, we will often implicitly use the following notation:

- (i) $q = 1 - 4p^2$,
- (ii) $\theta = pq$,
- (iii) $\lambda = 2\theta = 2pq$.

Note that $(2p)(2p) + q = 1$ and therefore $2p$ and q are coprime. We use this fact below when decomposing \mathbb{Z}/λ . We first define the boundary automorphisms of interest.

Lemma 5.1. *Let π be an abelian group and let $p \in \mathbb{Z}$. Set $q := 1 - 4p^2$ and consider the quadratic form $(V, \theta) = (\Lambda, pq)$. Using the notation of Lemma 4.9, for every $g \in \pi$ there is a boundary isomorphism $t_g = (\alpha, \beta, \nu): \partial(V, \theta) \xrightarrow{\cong} \partial(V, \theta)$ given by*

$$\begin{aligned} s &= -g + p & b &= 2pg + q \\ b_1 &= -(s + \bar{s}) & a &= \bar{b} \\ a_1 &= 1 & a_3 &= b \\ \alpha &= \begin{pmatrix} \bar{b} & 1 \\ -(s + \bar{s})\lambda & b \end{pmatrix} & \beta^{-1} &= \begin{pmatrix} b & -(s + \bar{s}) \\ \lambda & \bar{b} \end{pmatrix} \\ \alpha\nu\alpha^* &= \begin{pmatrix} s & \bar{b}(s + \bar{s}) \\ 0 & -(s + \bar{s})\theta(s + \bar{s}) \end{pmatrix}. \end{aligned}$$

Proof. We need to go through the checklist of Lemma 4.9.

- (i) Since λ is an integer, we have $a^*\lambda = \lambda b$.
- (ii) Next, we calculate:

$$\begin{aligned} ab &= \bar{b}b \\ &= (2pg^{-1} + q)(2pg + q) = 4p^2 + 2pq(g + g^{-1}) + q^2 = (1 - q) + q^2 + 2pq(g + g^{-1}) \\ &= 1 + q(q-1) + \lambda(g + g^{-1}) = 1 + q(-4p^2) + \lambda(g + g^{-1}) = 1 - 2p\lambda + \lambda(g + g^{-1}) \\ &= 1 + \lambda(g + g^{-1} - 2p) = 1 - (s + \bar{s})\lambda. \end{aligned}$$

- (iii) The next calculation takes place in $Q_{+1}(\Lambda) = \Lambda/(x - \bar{x})$. Using the previous result we check

$$\begin{aligned} b^*\theta b &= \theta\bar{b}b = \theta(1 - (s + \bar{s})\lambda) = \theta - (s + \bar{s})\theta\lambda \\ &= \theta - (s + \bar{s})\theta\lambda + (\bar{s}\theta\lambda) - \overline{(\bar{s}\theta\lambda)} \\ &= \theta - 2s\theta\lambda = \theta - \lambda s\lambda = \theta - \bar{\lambda}s\lambda. \end{aligned}$$

- (iv) We need to show that α is a simple isomorphism. We compute the product of these two simple isomorphisms using the first calculation

$$\begin{pmatrix} 0 & 1 \\ -1 & b \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \bar{b} & 1 \end{pmatrix} = \begin{pmatrix} \bar{b} & 1 \\ -1 + b\bar{b} & b \end{pmatrix} = \begin{pmatrix} \bar{b} & 1 \\ -1 + 1 - (s + \bar{s})\lambda & b \end{pmatrix} = \begin{pmatrix} \bar{b} & 1 \\ -(s + \bar{s})\lambda & b \end{pmatrix} = \alpha.$$

(v) At last we need to show that the matrix labelled β^{-1} is indeed an isomorphism:

$$\begin{pmatrix} b & -(s+\bar{s}) \\ \lambda & \bar{b} \end{pmatrix} \begin{pmatrix} \bar{b} & (s+\bar{s}) \\ -\lambda & b \end{pmatrix} = \begin{pmatrix} b\bar{b} + (s+\bar{s})\lambda & b(s+\bar{s}) - (s+\bar{s})b \\ \lambda\bar{b} - \bar{b}\lambda & \lambda(s+\bar{s}) + \bar{b}b \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

This shows that $t_g = (\alpha, \beta, \nu)$ is a boundary isomorphism and concludes the proof of the lemma. \square

Definition 5.2. Given a ring R with involution, the group of *unitary units* is defined as

$$U(R) = \{x \in R \mid x\bar{x} = 1\}.$$

Lemma 5.3. For π an abelian group, we have $U(\Lambda) = \{\pm g \mid g \in \pi\}$.

Proof. Let $x = \sum_{g \in \pi} n_g g \in U(\Lambda)$ then

$$1 = x\bar{x} = \sum_{g \in \pi} n_g g \cdot \sum_{g \in \pi} n_g g^{-1}.$$

The component of the product for $1 \in \pi$ must fulfill $1 = \sum n_g^2 \in \mathbb{Z}$ which means that exactly one of the n_g is ± 1 and the others are zero. \square

Lemma 5.4. Let π be an abelian group and let $p \in \mathbb{Z}$. Set $q := 1 - 4p^2$ and consider the quadratic form $(V, \theta) = (\Lambda, pq)$. The following diagram commutes:

$$\begin{array}{ccccc} \text{Aut}(V, \theta) & \xrightarrow{f} & U(\Lambda) & \xrightarrow{\text{Id}} & U(\Lambda) \\ \downarrow \partial & & \downarrow j & & \downarrow j' \\ \pi \xrightarrow{t} \text{Aut } \partial(V, \theta) & \xrightarrow{h} & U(\Lambda/\lambda) & \xrightarrow{r} & U(\Lambda/2p) \times U(\Lambda/q) \\ \downarrow k & & \downarrow l & & \downarrow l' \\ \text{bAut}(V, \theta) & \xrightarrow{h'} & U(\Lambda/\lambda)/U(\Lambda) & \xrightarrow{r'} & (U(\Lambda/2p) \times U(\Lambda/q))/U(\Lambda), \end{array}$$

where the maps are defined as follows:

$$\begin{aligned} f: x &\mapsto x; \\ \partial: x &\mapsto \partial x = (x, x, 0) \quad (\text{see Remark 4.3}); \\ j: x &\mapsto [x]; \\ j': x &\mapsto ([x], [x]); \\ t: g &\mapsto t_g \quad \text{from Lemma 5.1}; \\ h: (\alpha, \beta, \nu) &\mapsto b \quad \text{using the notation of Lemma 4.9}; \\ r: [x] &\mapsto ([x], [x]) \quad \text{induced by canonical isomorphism } \mathbb{Z}/2pq \cong \mathbb{Z}/2p \times \mathbb{Z}/q; \\ k: t &\mapsto [t] \quad \text{the projection to the quotient}; \\ l: [x] &\mapsto [x] \quad \text{the projection to the quotient}; \\ l': ([x], [y]) &\mapsto [[x], [y]] \quad \text{the projection to the quotient}; \\ h': [(\alpha, \beta, \nu)] &\mapsto [b] \quad \text{the map induced on the quotients by } h; \\ r': [x] &\mapsto ([x], [x]) \quad \text{the map induced on the quotients by } r. \end{aligned}$$

Proof. By construction $2p$ and q are coprime, so the definition of r makes sense. The map h is well-defined by Remark 4.11. We verify it is a group homomorphism. If (α, β, ν) and (α', β', ν') are automorphisms of $\partial(V, \theta)$, then $h(\alpha, \beta, \nu)h(\alpha', \beta', \nu') = bb'$, by definition of h . In order to understand $h((\alpha, \beta, \nu)(\alpha', \beta', \nu'))$, we use Lemma 4.9 to compute the product

$$(\beta\beta')^{-1} = \beta'^{-1}\beta^{-1} = \begin{pmatrix} b' & b'_1 \\ a_1'^*\lambda & a_3'^* \end{pmatrix} \begin{pmatrix} b & b_1 \\ a_1^*\lambda & a_3^* \end{pmatrix} = \begin{pmatrix} b'b + b'_1 a_1^* \lambda & * \\ * & * \end{pmatrix}$$

and deduce that $h((\alpha, \beta, \nu)(\alpha', \beta', \nu')) = b'b \in \Lambda/\lambda$, proving that h is indeed a homomorphism.

The top left square of the diagram now commutes: $j(f(x)) = j(x) = [x] = h(x, x, 0) = h(\partial(x))$. The remainder of the verifications are straightforward and are left to the reader. \square

Lemma 5.5. *Using the same notation as in Lemma 5.5, the map $l' \circ r \circ h \circ t$ is injective.*

Proof. Given $g \in \pi$, observe that

$$\begin{aligned} r \circ h \circ t(g) &= r(2pg + q) = ([2pg + q], [2pg + q]) \\ &= ([q], [2pg]) = ([1 - 4p^2], [2pg]) = (1, [2pg]) \in U(\Lambda/2p) \times U(\Lambda/q). \end{aligned}$$

If there are $g, g' \in \pi$ such that $l' \circ r \circ h \circ t(g) = l' \circ r \circ h \circ t(g')$, then there is an $x \in U(\Lambda)$ with

$$(1, [2pg]) = ([x], [x])(1, [2pg']) \in U(\Lambda/2p) \times U(\Lambda/q).$$

From Lemma 5.3, we know that $x = s\gamma$ for $s \in \{\pm 1\}, \gamma \in \pi$, and therefore

$$\begin{aligned} 1 &= s\gamma \in \mathbb{Z}/2p[\pi], \\ 2pg &= s\gamma 2pg' \in \mathbb{Z}/q[\pi]. \end{aligned}$$

The first equation shows that γ is the identity element of π and therefore the second equation implies that $g = g'$. \square

Proposition 5.6. *Let π be an abelian group and $p \in \mathbb{Z}$. Set $q := 1 - 4p^2$ and consider the quadratic form $(V, \theta) = (\Lambda, pq)$. The map*

$$\begin{aligned} k \circ t: \pi &\longrightarrow \text{bAut}(V, \theta) \\ g &\longmapsto [t_g] = [(\alpha, \beta, \nu)] \end{aligned}$$

is injective, where (α, β, ν) is defined as in Lemma 5.1. The elements t_g represent an infinite subset of $\text{bAut}(V, \theta)$.

Proof. Lemma 5.5 showed that $l' \circ r \circ h \circ t$ is injective. According to the diagram in Lemma 5.4, this is the same as $r' \circ h' \circ k \circ t$. This implies that $k \circ t$ is already injective. \square

Proposition 5.7. *Let π be an abelian group and $p \in \mathbb{Z}$. Set $q := 1 - 4p^2$ and consider the quadratic form $(V, \theta) = (\Lambda, pq)$. Given $g \in \pi$, the quadratic form $(V, \theta) \cup_{t_g} (V, -\theta)$ is hyperbolic.*

Proof. Recall from Definition 4.12 that the gluing construction gives

$$\kappa(t_g) := (V, \theta) \cup_{t_g} (V, -\theta) = \left(\Lambda^2, \begin{pmatrix} \theta & 0 \\ \bar{b} & -s \end{pmatrix} \right) = \left(\Lambda^2, \begin{pmatrix} \theta & 0 \\ 2pg^{-1} + q & g - p \end{pmatrix} \right).$$

Consider the following simple isomorphism $u \in SL_2(\Lambda)$

$$u = \begin{pmatrix} 1 & 0 \\ -p & 1 \end{pmatrix} \begin{pmatrix} 1 & g^{-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2p & 1 \end{pmatrix} = \begin{pmatrix} 1 + 2pg^{-1} & g^{-1} \\ p - 2p^2g^{-1} & 1 - pg^{-1} \end{pmatrix}$$

and then compute in $Q_1(\Lambda^2)$:

$$\begin{aligned}
u^*Hu &= \begin{pmatrix} 1+2pg^{-1} & g^{-1} \\ p-2p^2g^{-1} & 1-pg^{-1} \end{pmatrix}^* \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1+2pg^{-1} & g^{-1} \\ p-2p^2g^{-1} & 1-pg^{-1} \end{pmatrix} \\
&= \begin{pmatrix} 1+2pg & p-2p^2g \\ g & 1-pg \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1+2pg^{-1} & g^{-1} \end{pmatrix} \\
&= \begin{pmatrix} (p-2p^2g)(1+2pg^{-1}) & (p-2p^2g)g^{-1} \\ (1-pg)(1+2pg^{-1}) & (1-pg)g^{-1} \end{pmatrix} \\
&= \begin{pmatrix} p-4p^3+2p^2g^{-1}-2p^2g & pg^{-1}-2p^2 \\ 1-2p^2+2pg^{-1}-pg & g^{-1}-p \end{pmatrix} \\
&\equiv \begin{pmatrix} p-4p^3+2p^2g^{-1}-2p^2g & pg^{-1}-2p^2 \\ 1-2p^2+2pg^{-1}-pg & g^{-1}-p \end{pmatrix} + (1-T) \begin{pmatrix} 2p^2g & -(pg^{-1}-2p^2) \\ 0 & -g^{-1} \end{pmatrix} \\
&= \begin{pmatrix} p-4p^3 & 0 \\ 2pg^{-1}+(1-4p^2) & g-p \end{pmatrix} = \begin{pmatrix} \theta & 0 \\ 2pg^{-1}+q & g-p \end{pmatrix}.
\end{aligned}$$

This shows that $\kappa(t_g)$ is hyperbolic and concludes the proof of the proposition. \square

Specialising to $\pi = \mathbb{Z}$, the following result will be used in our topological applications.

Theorem 5.8. *For $p \in \mathbb{Z}$ and $\theta = p(1-4p^2)$, the set $\text{bAut}_{H_+(\mathbb{Z}[t^{\pm 1}])}(\partial(\mathbb{Z}[t^{\pm 1}], \theta))$ is infinite.*

Proof. By Proposition 5.6, $\text{bAut}(\partial(\mathbb{Z}[t^{\pm 1}], \theta))$ contains the infinite set $\{t_g \mid g \in \mathbb{Z}\}$ where the t_g were constructed in Lemma 5.1. By Proposition 5.7 each of the boundary automorphisms t_g belongs to $\text{bAut}_{H_+(\mathbb{Z}[t^{\pm 1}])}(\partial(\mathbb{Z}[t^{\pm 1}], \theta))$, which establishes the result. \square

6. PRIMITIVE EMBEDDINGS

This section, the last that develops algebraic machinery, gives a new formulation of the boundary automorphism set. Section 6.1 introduces primitive embeddings and Section 6.2 relates these novel algebraic objects to boundary isomorphisms.

Not only are primitive embeddings simpler to define than boundary isomorphisms, they will also be used in Section 7 to describe the image $\delta(\Theta(W, \bar{v}))$ of the modified surgery obstruction for the normal (B, ξ) -cobordism (W, \bar{v}) constructed in Theorem 3.10.

6.1. 2-sided primitive embeddings. The aim of this subsection is to introduce 2-sided primitive embeddings. In what follows, we shall often write quadratic forms as $v = (V, \theta)$.

Motivated by Lemma 4.13, the main definition of this section is the following.

Definition 6.1. A *primitive embedding* is a split isometric injection $i: v \hookrightarrow m$ of ε -quadratic forms, where m is nonsingular. A *2-sided primitive embedding* is a pair $v \xrightarrow{i} m \xleftarrow{i'} v'$ of primitive embeddings such that $i(v)^\perp = i'(v')$.

We already encountered examples of 2-sided primitive embeddings in Section 4.3.

Example 6.2. Given a boundary isomorphism $f \in \text{Iso}(\partial v, \partial v')$, the inclusions described in Lemma 4.13 provide a 2-sided primitive embedding $v \hookrightarrow v \cup_f -v' \hookrightarrow -v'$.

Two 2-sided primitive embeddings $v \hookrightarrow m_0 \hookleftarrow v'$ and $v \hookrightarrow m_1 \hookleftarrow v'$ are *isomorphic* if there is an isometry $m_0 \rightarrow m_1$ that makes the following diagram commute:

$$(20) \quad \begin{array}{ccccc} v & \hookrightarrow & m_0 & \hookleftarrow & v' \\ \downarrow = & & \downarrow \cong & & \downarrow = \\ v & \hookrightarrow & m_1 & \hookleftarrow & v' \end{array}$$

The isomorphism class of a 2-sided primitive embedding (j, j') is denoted by $[(j, j')]$. For ε -quadratic forms v and v' , we use $\text{Prim}(v, v')$ to denote the set of isomorphism classes of 2-sided primitive embeddings between v and v' . By additionally fixing a nonsingular ε -quadratic form m , we can also consider the set

$$\text{Prim}_m(v, v') := \{[(j, j')] \mid v \xrightarrow{j} m \xleftarrow{j'} v' \text{ is a 2-sided primitive embedding}\}.$$

Next, we give a concrete examples of these notions over the ring of integers.

Example 6.3. Given an integer $q = y_1 y_2$ with y_1 and y_2 coprime integers, the following maps define a 2-sided primitive embedding into the hyperbolic form $H_+(\mathbb{Z})$:

$$(i_{y_1, y_2}, j_{y_1, y_2}) := \left((\mathbb{Z}, q) \xrightarrow{\begin{pmatrix} y_1 \\ y_2 \end{pmatrix}} \left(\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) \xleftarrow{\begin{pmatrix} -y_1 \\ y_2 \end{pmatrix}} (\mathbb{Z}, -q) \right).$$

Furthermore, one can check that every 2-sided primitive embedding $(\mathbb{Z}, q) \hookrightarrow H_+(\mathbb{Z}) \hookleftarrow (\mathbb{Z}, -q)$ is either of the form $(i_{y_1, y_2}, j_{y_1, y_2})$ or of the form $(i_{y_1, y_2}, -j_{y_1, y_2})$ for some y_1, y_2 with $q = y_1 y_2$.

For each factorisation $q = y_1 y_2$ (writing $q = (-y_1)(-y_2)$ leads to the same observation), one can construct eight 2-sided primitive embedding $(\mathbb{Z}, q) \hookrightarrow H_+(\mathbb{Z}) \hookleftarrow (\mathbb{Z}, -q)$, namely $\pm(i_{y_1, y_2}, j_{y_1, y_2})$, $\pm(i_{y_2, y_1}, -j_{y_2, y_1})$, $\pm(i_{y_1, y_2}, -j_{y_1, y_2})$, and $\pm(i_{y_2, y_1}, j_{y_2, y_1})$. The first four (resp. last four) of these 2-sided primitive embeddings are isomorphic; this is because isometries of $H_+(\mathbb{Z})$ are of the form

$$\begin{pmatrix} \varepsilon_1 & 0 \\ 0 & \varepsilon_2 \end{pmatrix}, \quad \begin{pmatrix} 0 & \varepsilon_1 \\ \varepsilon_2 & 0 \end{pmatrix}, \quad \varepsilon_1 \varepsilon_2 = 1.$$

Summarising, for each factorisation $q = y_1 y_2$, one can associate two isomorphism classes in $\text{Prim}_{H_+(\mathbb{Z})}((\mathbb{Z}, q), (\mathbb{Z}, -q))$, namely $[(i_{y_1, y_2}, j_{y_1, y_2})]$ and $[(i_{y_1, y_2}, -j_{y_1, y_2})]$. These two classes are unchanged if the signs of both y_1 and y_2 are reversed.

The groups $\text{Aut}(v)$ and $\text{Aut}(v')$ both act on the left on $\text{Prim}_m(v, v')$: given a 2-sided primitive embedding (j, j') , and isometries $g: v \rightarrow v$ and $h: v' \rightarrow v'$, one obtains a new 2-sided primitive embedding by setting $(g, h) \cdot [(j, j')] := [(j \circ g^{-1}, j' \circ h^{-1})]$. Modding out by these actions leads to the second main definition of this section.

Definition 6.4. Given quadratic forms v and v' , the *one-sided boundary primitive embedding sets* and *boundary primitive embedding set* are respectively defined as

$$\begin{aligned} \ell\text{Prim}_m(v, v') &:= \text{Prim}_m(v, v') / \text{Aut}(v), \\ r\text{Prim}_m(v, v') &:= \text{Prim}_m(v, v') / \text{Aut}(v'), \\ b\text{Prim}_m(v, v') &:= \text{Prim}_m(v, v') / \text{Aut}(v) \times \text{Aut}(v'). \end{aligned}$$

Thus, two 2-sided primitive embeddings agree in $\text{rPrim}_m(v, v')$ and $\ell\text{Prim}_m(v, v')$ (resp. $\text{bPrim}_m(v, v')$) if and only if they fit in a commutative diagram as the one displayed in (20), but where instead one equality (resp. both equalities) are allowed to be self-isomorphisms. For instance, two 2-sided primitive embeddings $v \hookrightarrow m_0 \hookleftarrow v'$ and $v \hookrightarrow m_1 \hookleftarrow v'$ agree in $\ell\text{Prim}(v, v')$ if and only if there are isometries $m_0 \rightarrow m_1$ and $v \rightarrow v'$ that makes the following diagram commute:

$$(21) \quad \begin{array}{ccccc} v & \hookrightarrow & m_0 & \hookleftarrow & v' \\ \downarrow \cong & & \downarrow \cong & & \downarrow = \\ v & \hookrightarrow & m_1 & \hookleftarrow & v'. \end{array}$$

If $m \cong m'$, then we saw in Section 4.3 that $\text{Iso}_m(\partial v, \partial v')$ is isomorphic to $\text{Iso}_{m'}(\partial v, \partial v')$. The same property holds for 2-sided primitive embeddings: if $m \cong m'$, then after post-composing with the isomorphism $m \cong m'$, a primitive embedding $v \hookrightarrow m$ gives rise to a primitive embedding $v \hookrightarrow m'$. In Remark 4.8, we also recalled that isometries $k: v \rightarrow w$ and $h: v' \rightarrow w'$ give rise to a canonical identification $\text{bIso}_m(\partial v, \partial v') \approx \text{bIso}_m(\partial w, \partial w')$ via $[f] \mapsto [\partial h \circ f \circ \partial k^{-1}]$. The analogous remark holds for 2-sided primitive embeddings.

Remark 6.5. Observe that isometries $k: v \xrightarrow{\cong} w$ and $h: v' \xrightarrow{\cong} w'$ give rise to a canonical identification $\text{bPrim}_m(v, v') \approx \text{bPrim}_m(w, w')$ via $[(j, j')] \mapsto [(j \circ h^{-1}, j \circ k^{-1})]$. Analogous remarks also hold for ℓIso and rIso . As a consequence, for instance, when comparing primitive embeddings it makes sense to say that $v_0 \hookrightarrow m_0 \hookleftarrow v'$ and $v_1 \hookrightarrow m_1 \hookleftarrow v'$ agree in $\text{rIso}_{m_0}(v_0, v')$: this means that these two 2-sided primitive embeddings fit into a commutative diagram of the following type:

$$(22) \quad \begin{array}{ccccc} v_0 & \hookrightarrow & m_0 & \hookleftarrow & v' \\ \downarrow \cong & & \downarrow \cong & & \downarrow = \\ v_1 & \hookrightarrow & m_1 & \hookleftarrow & v'. \end{array}$$

In particular, $\ell\text{Iso}_{m_0}(v_0, v)$ only depends on v and on the isometry type of v_0 and m . We emphasise that such identifications are not possible if one is merely working in Iso .

Building on Example 6.3, we now describe explicitly the elements of $\text{Prim}_{H_+(\mathbb{Z})}((\mathbb{Z}, q), (\mathbb{Z}, -q))$.

Example 6.6. For a product $q = \pm p_1^{x_1} p_2^{x_2} \cdots p_k^{x_k} \in \mathbb{Z}$ of k distinct prime powers we introduce some notation and define:

$$\begin{aligned} \mathcal{F}_q &:= \text{Prim}_{H_+(\mathbb{Z})}((\mathbb{Z}, q), (\mathbb{Z}, -q)), \\ r\mathcal{F}_q &:= \text{rPrim}_{H_+(\mathbb{Z})}((\mathbb{Z}, q), (\mathbb{Z}, -q)), \\ \ell\mathcal{F}_q &:= \ell\text{Prim}_{H_+(\mathbb{Z})}((\mathbb{Z}, q), (\mathbb{Z}, -q)), \\ b\mathcal{F}_q &:= \text{bPrim}_{H_+(\mathbb{Z})}((\mathbb{Z}, q), (\mathbb{Z}, -q)). \end{aligned}$$

We assert that $r\mathcal{F}_q = \ell\mathcal{F}_q = b\mathcal{F}_q$, and

$$|b\mathcal{F}_q| = |r\mathcal{F}_q| = |\ell\mathcal{F}_q| = \frac{1}{2}|\mathcal{F}_q| = 2^{k-1}.$$

There are 2^{k-1} ways to factorise q as a product $y_1 y_2$ with y_1 and y_2 coprime. Indeed, since $\text{gcd}(y_1, y_2) = 1$, for $i = 1, \dots, k$, it suffices to choose whether or not the prime p_i appears in the prime decomposition of y_1 . Also note that Example 6.3 ensures that each

of these factorisations determines two isomorphisms classes of 2-sided primitive embeddings; $[(i_{y_1, y_2}, j_{y_1, y_2})]$ and $[(i_{y_1, y_2}, -j_{y_1, y_2})]$. This establishes the claim about $|\mathcal{F}_q|$.

To conclude, notice that $(i_{y_1, y_2}, j_{y_1, y_2})$ and $(i_{y_1, y_2}, -j_{y_1, y_2})$ agree in $r\mathcal{F}_q$ (and therefore in $b\mathcal{F}_q$), as can be seen by glancing at the following commutative diagram:

$$\begin{array}{ccccc} (\mathbb{Z}, q) & \xrightarrow{\begin{pmatrix} y_1 \\ y_2 \end{pmatrix}} & \left(\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) & \xleftarrow{\begin{pmatrix} -y_1 \\ y_2 \end{pmatrix}} & (\mathbb{Z}, -q) \\ \downarrow = & & \downarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & & \downarrow (-1) \\ (\mathbb{Z}, q) & \xrightarrow{\begin{pmatrix} y_1 \\ y_2 \end{pmatrix}} & \left(\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) & \xleftarrow{\begin{pmatrix} y_1 \\ -y_2 \end{pmatrix}} & (\mathbb{Z}, -q). \end{array}$$

Replacing respectively the labels on the left, middle and right vertical arrows by (-1) , $-\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $=$ shows that $(j_{y_1, y_2}, \iota_{y_1, y_2})$ and $(j_{y_1, y_2}, -\iota_{y_1, y_2})$ also agree in $\ell\mathcal{F}_q$. We conclude that there are twice as many elements in \mathcal{F}_q as there are in $r\mathcal{F}_q$, $\ell\mathcal{F}_q$ and $b\mathcal{F}_q$. This establishes the assertion.

The primitive embeddings in Example 6.6 being distinct is the algebra underlying the constructions from our paper [CCPS21].

6.2. Primitive embeddings and boundary isomorphisms. In this subsection we relate 2-sided primitive embeddings and boundary isomorphisms.

Recall that for ε -quadratic forms v, v' and a nonsingular ε -quadratic form m , we considered

$$\text{Iso}_m(\partial v, \partial v') := \{[f] \in \text{Iso}(\partial v, \partial v') \mid v \cup_f -v' \cong m\}.$$

The sets $\text{rIso}_m(\partial v, \partial v')$, $\ell\text{Iso}_m(\partial v, \partial v')$ and $\text{bIso}_m(\partial v, \partial v')$ are defined as quotients of the set $\text{Iso}_m(\partial v, \partial v')$.

Construction 6.7. We construct a map $\text{Iso}_m(\partial v, \partial v') \rightarrow \text{Prim}_m(v, -v')$. Given a stable homotopy class of a stable isomorphism $f: \partial v \rightarrow \partial v'$ such that $v \cup_f -v' \cong m$, we choose an isometry $k: v \cup_f -v' \rightarrow m$, use λ' for the symmetrisation of the form v' , and consider

$$(23) \quad \text{Pr}: \text{Iso}_m(\partial v, \partial v') \rightarrow \text{Prim}_m(v, -v')$$

$$[f] \mapsto \left(v \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} v \cup_f -v' \xrightarrow{k, \cong} m \xleftarrow{k, \cong} v \cup_f -v' \xleftarrow{\begin{pmatrix} b \\ -\lambda' \end{pmatrix}} -v' \right).$$

We verify that the definition of the map Pr makes sense. Set $j := \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $j' := \begin{pmatrix} b \\ -\lambda' \end{pmatrix}$ for concision. It follows from Example 6.2 that $k \circ j(v)^\perp = k \circ j'(-v')$. Also the isometry type of the algebraic union $v \cup_f -v'$ only depends on the homotopy class of the stable isomorphism f , so for any choice of representative f , a choice of isometry k exists. Thus Pr can be defined, after making a choice of k . To see that Pr is well-defined, we need to show that Pr does not depend on the choice of the isometry $k: v \cup_f -v' \rightarrow m$. Assume that $h: v \cup_f -v' \rightarrow m$ is another isometry. We must show that $(k \circ j, k \circ j')$ and $(h \circ j, h \circ j')$ define isomorphic 2-sided primitive embeddings. This follows since $\varphi := h \circ k^{-1}: m \rightarrow m$ is an isometry that satisfies $\varphi \circ (k \circ j) = h \circ j$ and $\varphi \circ (k \circ j') = h \circ j'$. We conclude that Pr is indeed well defined.

Remark 6.8. Construction 6.7 also defines a map $\text{Pr}: \text{Iso}(\partial v, \partial v') \rightarrow \text{Prim}(v, -v')$: the nonsingular form m need not be fixed. The results of this section also hold for this “less refined” map.

We now construct an inverse to the map Pr .

Construction 6.9. Let $v \xrightarrow{i} m \xleftarrow{i'} -v'$ be a 2-sided primitive embedding. According to Proposition 4.14, the primitive embedding i gives rise to a stable isomorphism $[f_i] \in \text{Iso}_m(\partial v, \partial(-v^\perp))$. Since (i, i') is a 2-sided primitive embedding, we know that $-v^\perp = i'(v')$; recall Example 6.2. We deduce the existence of the map $i'^{-1}: -v^\perp = i'(v') \rightarrow v'$. Consider the assignment

$$\begin{aligned} \delta: \text{Prim}_m(v, -v') &\rightarrow \text{Iso}_m(\partial v, \partial v') \\ [v \xrightarrow{i} m \xleftarrow{i'} -v'] &\mapsto [(\partial i'^{-1} \oplus 1) \circ f_i]. \end{aligned}$$

We check that δ is well defined. Namely, we assume that (j, j') is a 2-sided primitive embedding that is isomorphic to (i, i') , and wish to show that $\delta(i, i') = \delta(j, j')$ as boundary isomorphisms. Since (i, i') and (j, j') are isomorphic, there is an isometry $g: m \rightarrow m$ such that $g \circ j = i$ and $g \circ j' = i'$. Using the naturality statement of Proposition 4.14, we deduce that $f_i = (\partial g \oplus 1) \circ f_j$. We can combine these facts to conclude the proof of the fact that δ is well defined:

$$\delta(j, j') = [(\partial j'^{-1} \oplus 1) \circ f_j] = [(\partial i'^{-1} \oplus 1) \circ (\partial g \oplus 1) \circ f_j] = [(\partial i'^{-1} \oplus 1) \circ f_i] = \delta(i, i').$$

The next result shows that Pr and δ descend to the boundary sets.

Proposition 6.10. *Let v, v' and m be ε -quadratic forms where the latter is nonsingular. The map Pr from Construction 6.7 descends to*

$$\begin{aligned} \text{Pr}: \ell\text{Iso}_m(\partial v, \partial v') &\rightarrow \ell\text{Prim}_m(v, -v'), \\ \text{Pr}: \text{rIso}_m(\partial v, \partial v') &\rightarrow \text{rPrim}_m(v, -v'), \\ \text{Pr}: \text{bIso}_m(\partial v, \partial v') &\rightarrow \text{bPrim}_m(v, -v'), \end{aligned}$$

and the map δ from Construction 6.9 descends to

$$\begin{aligned} \delta: \ell\text{Prim}_m(v, -v') &\rightarrow \ell\text{Iso}_m(\partial v, \partial v'), \\ \delta: \text{rPrim}_m(v, -v') &\rightarrow \text{rIso}_m(\partial v, \partial v'), \\ \delta: \text{bPrim}_m(v, -v') &\rightarrow \text{bIso}_m(\partial v, \partial v'). \end{aligned}$$

Proof. Let $f: \partial v \cong_s \partial v'$ be a stable isomorphism, and let $h: v \rightarrow v$ and $k: v' \rightarrow v'$ be isometries. To prove that Pr is defined on the quotients, we must show that $\text{Pr}(\partial k \circ f \circ \partial h^{-1}) = \text{Pr}(f)$. We will use [CS11, Lemma 4.6 iv)] which states that $\begin{pmatrix} h & 0 \\ 0 & k^{-*} \end{pmatrix}: v \cup_f -v' \rightarrow v \cup_{\partial k \circ f \circ \partial h^{-1}} -v'$ is an isometry. Let a, b, s be the components of the stable isomorphism $f = (\alpha, \beta, \nu)$. Recall that for an isometry $h: v \rightarrow v$, we have $\partial h := (h, h, 0)$, and recall from Example 4.10 that $\partial k \circ f \circ \partial h^{-1}$ has a -component $k \circ a \circ h^{-1}$ and b -component $h \circ b \circ k^{-1}$. Next, we use λ, λ' to denote the symmetrized forms on v, v' and consider the following diagram in

which the horizontal maps are the primitive embeddings described in Lemma 4.13:

$$(24) \quad \begin{array}{ccccc} v & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & v \cup_f -v' & \xleftarrow{\begin{pmatrix} b \\ -\lambda' \end{pmatrix}} & -v' \\ h \downarrow \cong & & \begin{pmatrix} h & 0 \\ 0 & k^{-*} \end{pmatrix} \downarrow \cong & & k \downarrow \cong \\ v & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & v \cup_{\partial k \circ f \circ \partial h^{-1}} -v' & \xleftarrow{\begin{pmatrix} hb k^{-1} \\ -\lambda' \end{pmatrix}} & -v' \end{array}$$

The left hand square of (24) clearly commutes. Note that (24) also shows that $\text{Pr}((h, k) \cdot f) = (h, k) \cdot \text{Pr}(f)$ so that Pr intertwines the $\text{Aut}(v) \times \text{Aut}(v')$ actions. The right hand square of (24) commutes because k is an isometry. As a consequence, we deduce that the 2-sided primitive embeddings $\text{Pr}(\partial k \circ f \circ \partial h^{-1})$ and $\text{Pr}(f)$ agree in $\text{bPrim}(v, v')$. This shows that Pr descends to $\text{bIso}_m(\partial v, \partial v')$. Taking $k = \text{Id}$ (resp. $h = \text{Id}$) in this proof gives the analogous statement for $\text{lIso}_m(\partial v, \partial v')$ (resp. $\text{rIso}_m(\partial v, \partial v')$).

We show that δ descends to $\text{bPrim}_m(v, -v')$. Let $v \xrightarrow{j} m \xleftarrow{j'} -v'$ be a 2-sided primitive embedding, and let $h: v \rightarrow v$ and $k: v' \rightarrow v'$ be isometries. To show that δ descends to the orbits sets, we prove $\delta(jh^{-1}, j'k^{-1}) = \partial k \circ \delta(j, j') \circ \partial h^{-1}$. Recall that $\delta(j, j') := (\partial j'^{-1} \oplus 1) \circ f_j$, where f_j denotes the stable isomorphism from Proposition 4.14. Use a, b, s to denote the components of the stable isomorphism $\delta(j, j') := (\partial j'^{-1} \oplus 1) \circ f_j$. By Example 4.10, $\partial k \circ \delta(j, j') \circ \partial h^{-1}$ has components $kah^{-1}, hb k^{-1}$ and ksk^* . By Remark 4.11, it suffices to show that $\delta(jh^{-1}, j'k^{-1})$ has the same components. By definition, we have $\delta(jh^{-1}, j'k^{-1}) = (\partial k \partial j'^{-1} \oplus 1) \circ f_{jh^{-1}}$. It therefore suffices to show that $(\partial j'^{-1} \oplus 1) f_{jh^{-1}}$ has components ah^{-1}, hb and s . Since h is an isometry, the orthogonal complement $jh^{-1}(v)^\perp$ of $jh^{-1}(v)$ in m coincides with the orthogonal complement $j(v)^\perp$ of v in m . In particular, the splitting σ of $j^{\perp*} \phi$ can also be used in the computation of $f_{jh^{-1}}$. Here recall that ϕ denotes the symmetrisation of the form on $v \cup_f -v'$. On the other hand, since $\tilde{\sigma} := j^{-1}(1 - \sigma j^{\perp*} \phi)$ splits j as in (16), we deduce that $h\tilde{\sigma}$ splits jh^{-1} . It now follows from Proposition 4.15 that the components of $f_{jh^{-1}}$ are indeed ah^{-1}, hb and s . This establishes that δ descends to $\text{bPrim}(v, -v')$ as well as to $\text{lPrim}(v, -v')$ and $\text{rPrim}(v, -v')$ (by respectively taking $k = \text{Id}$ and $h = \text{Id}$), concluding the proof the proposition. \square

We can now prove the main result of this section.

Theorem 6.11. *Let v, v' and m be ε -quadratic forms where the latter is nonsingular. The map Pr from Construction 6.7 descends to bijections*

$$\begin{aligned} \text{Pr}: \text{lIso}_m(\partial v, \partial v') &\rightarrow \text{lPrim}_m(v, -v'), \\ \text{Pr}: \text{rIso}_m(\partial v, \partial v') &\rightarrow \text{rPrim}_m(v, -v'), \\ \text{Pr}: \text{bIso}_m(\partial v, \partial v') &\rightarrow \text{bPrim}_m(v, -v'). \end{aligned}$$

Its inverse is the map δ induced by Construction 6.9.

Proof. Given a 2-sided primitive embedding $v \xrightarrow{i} m \xleftarrow{i'} v'$, we first show that $\text{Pr} \circ \delta(i, i') \cong (i, i')$ in $\text{rIso}(v, -v')$. Recall that $\delta(i, i') = [(\partial i'^{-1} \oplus 1) \circ f_i]$, where $i'^{-1}: -v^\perp = i'(v') \rightarrow v'$. Using $b := b_{f_i}$ to denote the b component of the stable isomorphism f_i , Example 4.10 shows that the b component of $(\partial i'^{-1} \oplus 1) \circ f_i$ is bi' . The 2-sided primitive embedding $\text{Pr} \circ \delta(i, i')$ is

therefore

$$\mathrm{Pr} \circ \delta(i, i') = \left(v \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} v \cup_{(\partial^{i'-1} \oplus 1) \circ f_i} -v' \xrightarrow{k, \cong} m \xleftarrow{k, \cong} v \cup_{(\partial^{i'-1} \oplus 1) \circ f_i} -v' \xleftarrow{\begin{pmatrix} bi' \\ -\lambda' \end{pmatrix}} -v' \right).$$

Proposition 4.14 provides an isometry $r_i: m \rightarrow v \cup_{f_i} v^\perp$. We describe an explicit formula for this isometry following [CS11, proof of Proposition 4.8]: $r_i = (i - \sigma)$, where $\sigma: v^{\perp*} \rightarrow m$ is a splitting of $i^{\perp*} \phi: m \rightarrow v^{\perp*}$, meaning that $i^{\perp*} \phi \sigma = \mathrm{Id}_{v^{\perp*}}$; here $\phi = \psi + \varepsilon \psi^*$ is the ε -symmetrization of the quadratic form $m = (M, \psi)$. One can check that r_i is indeed an isometry by using Proposition 4.15. Consider the following diagram:

$$(25) \quad \begin{array}{ccccc} v & \xrightarrow{i} & m & \xleftarrow{i'} & -v' \\ \cong \downarrow & & \cong \uparrow (i - \sigma) & & \cong \downarrow -i' \\ v & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & v \cup_{f_i} v^\perp & \xleftarrow{\begin{pmatrix} b \\ \lambda^\perp \end{pmatrix}} & v^\perp \cong \\ \cong \downarrow & & \cong \downarrow \begin{pmatrix} 1 & 0 \\ 0 & i'^* \end{pmatrix} & & \cong \uparrow i' \\ v & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & v \cup_{(\partial^{i'-1} \oplus 1) \circ f_i} -v' & \xleftarrow{\begin{pmatrix} bi' \\ -\lambda' \end{pmatrix}} & -v'. \end{array} \quad (-1).$$

Observe that each horizontal line of the diagram in (25) consists of 2-sided primitive embeddings, and the third line is $\mathrm{Pr} \circ \delta(i, i')$. The equation $\mathrm{Pr} \circ \delta(i, i') \cong (i, i')$ will therefore follow once we show that the diagram in (25) commutes. The top and bottom left squares of (25) clearly commute. The bottom right square of (25) commutes since $i'^* \lambda^\perp i' = -\lambda'$. In order to prove that the top right square of (25) commutes, we show that $-(ib - \sigma \lambda^\perp) i' = i'$. Recall from Proposition 4.15 the b component of the stable isomorphism $f_i: \partial v \cong_s \partial(-v^\perp)$ is $b = -i^{-1}(1 - \sigma i^{\perp*} \phi) i^\perp$. By using the equation $i^{\perp*} \phi i^\perp = \lambda^\perp$, note that $ib = \sigma \lambda^\perp - i^\perp$. Using this formula and the fact that $i^\perp: v^\perp \hookrightarrow m$ is the canonical inclusion (so that $v' \xrightarrow{i'} v^\perp \xrightarrow{i^\perp} m$ is also denoted by i'), we obtain the desired result:

$$-(ib - \sigma \lambda^\perp) i' = -(\sigma \lambda^\perp - i^\perp - \sigma \lambda^\perp) i' = i^\perp i' = i'.$$

This shows that the diagram in (25) commutes, proving that $\mathrm{Pr} \circ \delta(i, i') \cong (i, i')$ in $\mathrm{rPrim}(v, -v')$, and thus in $\mathrm{bPrim}(v, -v')$. To see that this isomorphism also holds in $\ell\mathrm{Prim}(v, -v')$, fix two 2-sided primitive embeddings $(j, j'), (k, k')$ and consider the two following diagrams:

$$\begin{array}{ccc} v \xrightarrow{j} m \xleftarrow{j'} -v' & & v \xrightarrow{j} m \xleftarrow{j'} -v' \\ \cong \downarrow & \cong \downarrow (-1) & \cong \downarrow (-1) \\ v \xrightarrow{k} m \xleftarrow{k'} -v' & \cong \downarrow (-1) & v \xrightarrow{k} m \xleftarrow{k'} -v' \end{array} \quad \begin{array}{ccc} v \xrightarrow{j} m \xleftarrow{j'} -v' & & v \xrightarrow{j} m \xleftarrow{j'} -v' \\ \cong \downarrow (-1) & \cong \downarrow -\varphi & \cong \downarrow (-1) \\ v \xrightarrow{k} m \xleftarrow{k'} -v' & & v \xrightarrow{k} m \xleftarrow{k'} -v' \end{array}$$

The commutativity of one of the diagram is equivalent to the commutativity of the other. Consequently (25) also proves that $\mathrm{Pr} \circ \delta(i, i') \cong (i, i')$ in $\ell\mathrm{Prim}(v, -v')$.

For $f \in \mathrm{Iso}(\partial v, \partial v')$, Example 4.16 proves that $\delta \circ \mathrm{Pr}(f)$ and f agree in rIso , $\ell\mathrm{Iso}$ and bIso (we observed that the components of $\delta(\mathrm{Pr}(f))$ coincide with the components of both $(\partial(-1) \oplus 1) \circ f$ and $f \circ (\partial(-1) \oplus 1)^{-1}$ and applied Remark 4.11), so this concludes the proof of Theorem 6.11. \square

Example 6.12. As a concrete example of the calculations that appear in the proof of Theorem 6.11, we describe a 2-sided primitive embedding representing $\Pr(\delta([(i_{y_1, y_2}, j_{y_1, y_2})]))$. Here, recall from Example 6.3 that for $q = y_1 y_2$, we set

$$(i_{y_1, y_2}, j_{y_1, y_2}) := \left((\mathbb{Z}, q) \xrightarrow{\begin{pmatrix} y_1 \\ y_2 \end{pmatrix}} (\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}) \xleftarrow{\begin{pmatrix} -y_1 \\ y_2 \end{pmatrix}} (\mathbb{Z}, -q) \right).$$

We compute the a, b, s components of $\delta(i_{y_1, y_2}, j_{y_1, y_2}) := f_{i_{y_1, y_2}}$ (the boundary automorphism $f_{i_{y_1, y_2}}$ was introduced in (14)), as this will be sufficient to calculate $\Pr(\delta([(i_{y_1, y_2}, j_{y_1, y_2})]))$.

We set $\phi := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $\psi = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and note that $i_{y_1, y_2}^\perp = j_{y_1, y_2}$. We also let $\sigma = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}$ be a splitting of $i_{y_1, y_2}^{\perp*}$ ($\phi = \begin{pmatrix} y_2 & -y_1 \\ 0 & 0 \end{pmatrix}$) (so that $y_2 z_1 - y_1 z_2 = 1$), and let $\tilde{\sigma} = \begin{pmatrix} -z_2 & z_1 \end{pmatrix}$ be the associated splitting of i_{y_1, y_2} . By Example 4.16, we deduce that the a, b, s -components $f_{i_{y_1, y_2}}$ are

$$\begin{aligned} a &= -\sigma^* \phi^* i_{y_1, y_2} = -\begin{pmatrix} z_1 & z_2 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = -(y_1 z_2 + y_2 z_1), \\ b &= -\tilde{\sigma} j_{y_1, y_2} = -\begin{pmatrix} -z_2 & z_1 \end{pmatrix} \begin{pmatrix} -y_1 \\ y_2 \end{pmatrix} = -(y_1 z_2 + y_2 z_1), \\ s &= -\sigma^* \psi \sigma = -\begin{pmatrix} z_1 & z_2 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} = -z_1 z_2. \end{aligned}$$

By definition of \Pr , we therefore deduce that $\Pr(\delta([(i_{y_1, y_2}, j_{y_1, y_2})]))$ is represented by

$$\left((\mathbb{Z}, q) \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} (\mathbb{Z}^2, \begin{pmatrix} q & 0 \\ a & -s \end{pmatrix}) \xleftarrow{\begin{pmatrix} b \\ -2q \end{pmatrix}} (\mathbb{Z}, -q) \right).$$

One can directly verify using the values of a, b, s and q above, together with $y_2 z_1 - y_1 z_2 = 1$, that this is a primitive embedding. As a reality check, we verify that that this 2-sided primitive embeddings agrees with $(i_{y_1, y_2}, j_{y_1, y_2})$ in $\text{bPrim}((\mathbb{Z}, q), (\mathbb{Z}, -q))$:

$$\begin{array}{ccc} (\mathbb{Z}, q) & \xrightarrow{\begin{pmatrix} y_1 \\ y_2 \end{pmatrix}} & (\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}) & \xleftarrow{\begin{pmatrix} -y_1 \\ y_2 \end{pmatrix}} & (\mathbb{Z}, -q) \\ \uparrow & & \uparrow & & \uparrow \\ (\mathbb{Z}, q) & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & (\mathbb{Z}^2, \begin{pmatrix} q & 0 \\ a & -s \end{pmatrix}) & \xleftarrow{\begin{pmatrix} b \\ -2q \end{pmatrix}} & (\mathbb{Z}, -q). \end{array}$$

Combining $b = -(y_1 z_2 + y_2 z_1)$, $q = y_1 y_2$, and $y_2 z_1 - y_1 z_2 = 1$ one can check that the diagram commutes.

7. REALISING BOUNDARY ISOMORPHISMS AND PRIMITIVE EMBEDDINGS

Now we start to apply the algebra of Sections 4, 5, and 6. We return to the setting of Section 3. That is, we fix a fibration $\xi: B \rightarrow BO$ and let $\bar{\nu}: M^{2q} \rightarrow B$ be a normal $(q-1)$ -smoothing over (B, ξ) . Throughout this section, we work with closed, connected, oriented, smooth manifolds M, M' or M_i , for $i = 0, 1, 2$, and we assume that $q > 3$ is even, or that $q = 2$, in which case we must switch to the topological category and assume that $\pi_1(B)$ is a good group. We set $\Lambda := \mathbb{Z}[\pi_1(B)]$ and consider the surgery kernel

$$K_q(M; \Lambda) := \ker(\bar{\nu}_*: H_q(M; \Lambda) \rightarrow H_q(B; \Lambda)),$$

where we use $\bar{\nu}_*: \pi_1(M) \rightarrow \pi_1(B)$ to identify fundamental groups. Since q is even and M is oriented, if the Λ -valued intersection form of M , $\lambda_M: H_q(M; \Lambda) \times H_q(M; \Lambda) \rightarrow \Lambda$ admits a quadratic refinement, then it is unique. In this case we denote it by θ_M .

In Section 7.1, under simplifying algebraic assumptions we associate a 2-sided primitive embedding

$$K_q(M; \Lambda) \xrightarrow{\iota_{\bar{\nu}}} H_q(M; \Lambda) \xleftarrow{j_{\bar{\nu}}} H^q(B; \Lambda)$$

to $(M, \bar{\nu})$ (Definition 7.5). By Construction 6.9, this gives rise to a boundary isomorphism

$$\delta_{\bar{\nu}} := \delta(\iota_{\bar{\nu}}, j_{\bar{\nu}}) \in \text{Iso}(\partial(K_q(M; \Lambda), \iota_{\bar{\nu}}^* \theta_M \iota_{\bar{\nu}}), \partial(H^q(B; \Lambda), -j_{\bar{\nu}}^* \theta_M j_{\bar{\nu}})).$$

In Section 7.2, we prove a realisation result for this boundary automorphism. Roughly, given a boundary automorphism $f \in \text{Aut}(\partial(K_q(M; \Lambda), \iota_{\bar{\nu}}^* \theta_M \iota_{\bar{\nu}}))$, Theorem 7.7 constructs another normal $(q-1)$ -smoothing $(M', \bar{\nu}')$ which is normally (B, ξ) -bordant (and thus stably diffeomorphic) to $(M, \bar{\nu})$ and whose associated boundary isomorphism δ' agrees with δf in rIso .

Then for the quadratic form $v(\bar{\nu}) = (K_q(M; \Lambda), \iota_{\bar{\nu}}^* \theta_M \iota_{\bar{\nu}})$ we relate this result to the realisation of elements of $\ell_{2q+1}([v(\bar{\nu})]_0)$ from Section 3. Given an appropriate quasi-formation x representing $[x] \in \ell_{2q+1}([v(\bar{\nu}_0)]_0)$, Theorem 3.10 constructed a (B, ξ) -bordism $(W, \bar{\nu})$ with $\Theta(W, \bar{\nu}) = [x]$. In Corollary 7.14, under simplifying algebraic assumptions, we show that the two realisation theorems are related via the exact sequence (1) from [CS11], as one would hope.

7.1. The boundary isomorphism of an even split-free normal smoothing. Under simplifying algebraic assumptions, we associate a 2-sided primitive embedding and a boundary isomorphism to a $(q-1)$ -smoothing $\bar{\nu}: M^{2q} \rightarrow B$. In Section 9, these objects will be used to distinguish certain stably diffeomorphic $4k$ -manifolds.

Consider a map $\bar{\nu}: M^{2q} \rightarrow B$, with B homotopy equivalent to a CW -complex with finite q -skeleton and M closed, connected, and oriented. The inclusion $K_q(M; \Lambda) \hookrightarrow H_q(M; \Lambda)$ and the composition of $\bar{\nu}^*: H^q(B; \Lambda) \rightarrow H^q(M; \Lambda)$ with the Poincaré duality isomorphism $\text{PD}: H^q(M; \Lambda) \xrightarrow{\cong} H_q(M; \Lambda)$ produce the following Λ -homomorphisms:

$$(26) \quad \begin{aligned} \iota_{\bar{\nu}}: K_q(M; \Lambda) &\rightarrow H_q(M; \Lambda), \\ j_{\bar{\nu}} := \text{PD} \circ \bar{\nu}^*: H^q(B; \Lambda) &\rightarrow H_q(M; \Lambda). \end{aligned}$$

Recall that when q is even and M is oriented, if the symmetric form $(H_q(M; \Lambda), \lambda_M)$ admits a quadratic refinement θ_M , then the refinement is uniquely determined by λ_M . The quadratic form θ_M could be singular, because the evaluation homomorphism $\text{ev}_M: H^q(M; \Lambda) \rightarrow \overline{\text{Hom}}_{\Lambda}(H_q(M; \Lambda), \Lambda)$ need not be an isomorphism. Next, use $\iota_{\bar{\nu}}$ and $j_{\bar{\nu}}$ to pull the quadratic form θ_M back from $H_q(M; \Lambda)$ to $K_q(M; \Lambda)$ and $H^q(B; \Lambda)$ respectively. This leads to the two following quadratic forms:

$$\begin{aligned} (K_q(M; \Lambda), \theta_M) &:= (K_q(M; \Lambda), \iota_{\bar{\nu}}^* \theta_M \iota_{\bar{\nu}}), \\ (H^q(B; \Lambda), \theta_M) &:= (H^q(B; \Lambda), j_{\bar{\nu}}^* \theta_M j_{\bar{\nu}}). \end{aligned}$$

In order for $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ to define a 2-sided primitive embedding, we impose some further restrictions on the map $\bar{\nu}: M \rightarrow B$.

Definition 7.1. Fix B , a space homotopy equivalent to a CW -complex with finite q -skeleton. A map $\bar{\nu}: M \rightarrow B$ is called q -surjective if it induces a surjection $\bar{\nu}_*: H_q(M; \Lambda) \rightarrow H_q(B; \Lambda)$. Let M be a closed, connected, and oriented $2q$ -dimensional manifold with $q \geq 2$ even.

- (i) We say that M is *even* if the Λ -valued intersection form on $H_q(M; \Lambda)$ admits a quadratic refinement θ_M , which since M is oriented and q is even is unique.
- (ii) We say that $(M, \bar{\nu})$ is *split-free* if the evaluation map

$$\text{ev}_B: H^q(B; \Lambda) \rightarrow \overline{\text{Hom}_\Lambda(H_q(B; \Lambda), \Lambda)}$$

is an isomorphism, and if the Λ -modules $K_q(M; \Lambda) := \ker(\bar{\nu}_*)$ and $H_q(B; \Lambda)$ are f.g. free. If in addition M is even, we shall say that $(M, \bar{\nu})$ is an *even, split-free map*.

We will describe the two main examples of q -surjective maps that we wish to keep in mind. For this we need a quick lemma.

Lemma 7.2. *For $q \geq 1$, if $\bar{\nu}: M \rightarrow B$ is q -connected, then it is q -surjective.*

Proof. Since $\bar{\nu}$ induces an isomorphism on fundamental groups, the pullback $\bar{\nu}^*(\tilde{B}) \rightarrow M$ of the universal covering of B is homeomorphic to the universal cover \tilde{M} , so $\bar{\nu}^*(\tilde{B})$ is simply-connected. Apply the relative Hurewicz theorem to $(\tilde{B}, \bar{\nu}^*(\tilde{B}))$ to see that $\bar{\nu}$ induces a surjection on $H_q(-; \Lambda)$ as asserted. \square

Example 7.3. Using Lemma 7.2 we note the two main examples of q -surjective maps.

- (i) Assume that $B = P_{q-1}(M)$ is the $(q-1)$ -st Postnikov stage of M , a model for which is obtained from M by adding cells of dimension of dimension $(q+1)$ and higher in order to kill $\pi_i(M)$ for $i \geq q$ [Hat02, Chapter 4.3]. Since the inclusion map $\bar{\nu} := p_{q-1}(M): M \rightarrow P_{q-1}(M)$ is q -connected, it is q -surjective.
- (ii) Given a fibration, $\xi: B \rightarrow BO$, a normal $(q-1)$ -smoothing $\bar{\nu}: M \rightarrow B$ is q -connected and therefore q -surjective.

Lemma 7.4. *Let $q \geq 2$ be even. If $\bar{\nu}: M^{2q} \rightarrow B$ is an even, split-free q -surjective map and $(H_q(M; \Lambda), \theta_M)$ is a nonsingular quadratic form, then $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$,*

$$(K_q(M; \Lambda), \theta_M) \xrightarrow{\iota_{\bar{\nu}}} (H_q(M; \Lambda), \theta_M) \xleftarrow{j_{\bar{\nu}}} (H^q(B; \Lambda), \theta_M),$$

is a 2-sided primitive embedding.

Proof. First, $\iota_{\bar{\nu}}$ is a split injection, because $H_q(B; \Lambda)$ free implies that the short exact sequence

$$0 \rightarrow K_q(M; \Lambda) \xrightarrow{\iota_{\bar{\nu}}} H_q(M; \Lambda) \xrightarrow{\bar{\nu}_*} H_q(B; \Lambda) \rightarrow 0$$

splits. Next we prove that $j_{\bar{\nu}} = \text{PD} \circ \bar{\nu}^*$ is a split injection. Since PD is an isomorphism, it suffices to show that $\bar{\nu}^*$ is a split injection. The map $\bar{\nu}$ is q -surjective, so the induced map $\bar{\nu}_*: H_q(M; \Lambda) \rightarrow H_q(B; \Lambda)$ is onto. It follows that its dual $\bar{\nu}_d^*$ is injective. Now consider the following commutative diagram:

(27)

$$\begin{array}{ccccccc} H^q(B; \Lambda) & \xrightarrow{\bar{\nu}^*} & H^q(M; \Lambda) & & & & \\ \cong \downarrow \text{ev}_B & & \cong \downarrow \text{ev}_M & & & & \\ 0 \longrightarrow & \overline{\text{Hom}_\Lambda(H_q(B; \Lambda), \Lambda)} & \xrightarrow{\bar{\nu}_d^*} & \overline{\text{Hom}_\Lambda(H_q(M; \Lambda), \Lambda)} & \xrightarrow{\iota_{\bar{\nu}}^*} & \overline{\text{Hom}_\Lambda(K_q(M; \Lambda), \Lambda)} & \longrightarrow 0. \end{array}$$

We argue that the bottom line is split exact. We already established that $\overline{\nu}_d^*$ is injective. Since $H_q(B; \Lambda)$ is free, there is no Ext term and so it follows that $\iota_{\overline{\nu}}^*$ is surjective. Next, since $K_q(M; \Lambda)$ is finitely generated and free, so is $\overline{\text{Hom}}_{\Lambda}(K_q(M; \Lambda), \Lambda)$, and the exact sequence splits as claimed. We can now show that $\overline{\nu}^*$ is a split injection. Since $(M, \overline{\nu})$ is even and split-free, the evaluation map ev_B is an isomorphism, while ev_M is an isomorphism because the quadratic form $(H_q(M; \Lambda), \theta_M)$ is nonsingular. Since $\overline{\nu}_d^*$ is a split injection, the commutativity of the diagram displayed in (27) together with these facts implies that $\overline{\nu}^*$ must also be a split injection, as claimed.

It remains to show that $K_q(M; \Lambda)^\perp = j_{\overline{\nu}}(H^q(B; \Lambda))$. We first show that $j_{\overline{\nu}}(H^q(B; \Lambda))$ is contained in $K_q(M; \Lambda)^\perp$. Use $\langle -, - \rangle_B$ to denote the Kronecker pairing that evaluates a cohomology class on a homology class, so that $\langle \varphi, x \rangle_B = \text{ev}_B(\varphi)(x)$. For $x \in K_q(M; \Lambda)$ and for $\varphi \in H^q(B; \Lambda)$, the definition of the Λ -valued intersection pairing $\lambda_M = \theta_M + \varepsilon\theta_M^*$ (using Poincaré duality and the Kronecker evaluation map) gives

$$(28) \quad \lambda_M(j_{\overline{\nu}}(\varphi), x) = \langle \overline{\nu}^* \varphi, x \rangle_B = \langle \varphi, \overline{\nu}_*(x) \rangle_B.$$

The inclusion $j_{\overline{\nu}}(H^q(B; \Lambda)) \subseteq K_q(M; \Lambda)^\perp$ now follows readily: if $x \in K_q(M; \Lambda) = \ker(\overline{\nu}_*)$, then

$$\lambda_M(j_{\overline{\nu}}(\varphi), x) = \langle \varphi, \overline{\nu}_*(x) \rangle_B = \langle \varphi, 0 \rangle_B = 0,$$

so $j_{\overline{\nu}}(\varphi) \in K_q(M; \Lambda)^\perp$ as desired. To prove the reverse inclusion, $K_q(M; \Lambda)^\perp \subseteq j_{\overline{\nu}}(H^q(B; \Lambda))$, assume that $y \in H_q(M; \Lambda)$ satisfies $\lambda_M(y, \iota_{\overline{\nu}}(x)) = 0$ for all $x \in K_q(M; \Lambda)$. Equivalently, we have $\text{ev}_M \text{PD}_M^{-1}(y) \in \ker(\iota_{\overline{\nu}}^*) = \text{Im}(\overline{\nu}_d^*)$, by the exactness of the sequence in (27). In particular, $\text{ev}_M \text{PD}_M^{-1}(y) = \overline{\nu}_d^*(z)$ for some $z \in \overline{\text{Hom}}_{\Lambda}(H_q(B; \Lambda), \Lambda)$. Since ev_B is an isomorphism, $z = \text{ev}_B(\varphi)$ for some $\varphi \in H^q(B; \Lambda)$. By commutativity of (27), we obtain $\text{ev}_M \text{PD}_M^{-1}(y) = \text{ev}_M(\overline{\nu}^*(\varphi))$. Since ev_M is an isomorphism, we deduce that $y = j_{\overline{\nu}}(\varphi)$. We thus proved that $K_q(M; \Lambda)^\perp \subseteq j_{\overline{\nu}}(H^q(B; \Lambda))$, and this concludes the proof of the proposition. \square

Using Lemma 7.4, we associate a 2-sided primitive embedding to an even, split-free q -surjective map $\overline{\nu}$. We can also associate a boundary isomorphism to $\overline{\nu}$ by recalling from Construction 6.9 that a 2-sided primitive embedding $(\iota_{\overline{\nu}}, j_{\overline{\nu}})$ determines a boundary isomorphism $\delta(\iota_{\overline{\nu}}, j_{\overline{\nu}})$.

Definition 7.5. Let $(M^{2q}, \overline{\nu})$ be an even, split-free q -surjective map and assume that the quadratic form $(H_q(M; \Lambda), \theta_M)$ is nonsingular. The 2-sided primitive embedding associated with $\overline{\nu}$ is the 2-sided primitive embedding $(\iota_{\overline{\nu}}, j_{\overline{\nu}})$ from (26), the boundary isomorphism associated to $\overline{\nu}$ is

$$\delta_{\overline{\nu}} := \delta(\iota_{\overline{\nu}}, j_{\overline{\nu}}) \in \text{Iso}(\partial(K_q(M; \Lambda), \theta_M), \partial(H^q(B; \Lambda), -\theta_M)).$$

Some remarks on this definition are in order.

Remark 7.6. Let $\overline{\nu}: M^{2q} \rightarrow B$ be an even, split-free q -surjective map.

- (i) The invariant $\delta_{\overline{\nu}}$ determines an element in the one-sided boundary automorphism set $\ell\text{Iso}_{(H_q(M; \Lambda), \theta_M)}(\partial(K_q(M; \Lambda), \theta_M), \partial(H^q(B; \Lambda), -\theta_M))$. This follows from Theorem 6.11, in particular (25). As a consequence, there is an isometry

$$(H_q(M; \Lambda), \theta_M) \cong (K_q(M; \Lambda), \theta_M) \cup_{\delta_{\overline{\nu}}} (H^q(B; \Lambda), \theta_M).$$

- (ii) Recall from Remarks 4.8 and 6.5 that the sets $\ell\text{Iso}_m(\partial v, \partial v')$ and $\ell\text{Prim}_m(v, v')$ only depend on v' and on the isometry type of m and v . As a consequence, we say that the 2-sided primitive embeddings associated with the q -surjective maps $(M_0, \bar{\nu}_0)$ and $(M_1, \bar{\nu}_1)$ into B agree in ℓPrim if they fit into the following diagram:

$$\begin{array}{ccccc} (K_q(M_0; \Lambda)) & \xrightarrow{\iota_{\bar{\nu}_0}} & (H_q(M_0; \Lambda), \theta_{M_0}) & \xleftarrow{j_{\bar{\nu}_0}} & (H^q(B; \Lambda), \theta_{M_0}) \\ \downarrow \cong & & \downarrow \cong & & \downarrow = \\ (K_q(M_1; \Lambda)) & \xrightarrow{\iota_{\bar{\nu}_1}} & (H_q(M_1; \Lambda), \theta_{M_1}) & \xleftarrow{j_{\bar{\nu}_1}} & (H^q(B; \Lambda), \theta_{M_0}). \end{array}$$

If the right equality in this diagram is merely an isomorphism, then we say that these two 2-sided primitive embeddings agree in bPrim . We also use the corresponding terminology for the boundary isomorphism associated to $\bar{\nu}$, replacing Prim by Iso .

- (iii) If $H_q(M; \Lambda)$ is free, then the data of the 2-sided primitive embedding associated with $\bar{\nu}$ is determined by the map $\bar{\nu}^*: H^q(B; \Lambda) \rightarrow H^q(M; \Lambda)$: indeed, using that both $H_q(M; \Lambda)$ and $H_q(B; \Lambda)$ are free, the map $\bar{\nu}^*$ determines $\bar{\nu}_*$.

7.2. The realisation of boundary automorphisms. The goal of this section is to prove a realisation result for the boundary isomorphisms associated to bordant normal smoothings. Throughout this subsection, we fix a fibration $\xi: B \rightarrow BO$ as above, and assume that all homology and cohomology groups have $\Lambda := \mathbb{Z}[\pi_1(B)]$ coefficients.

Given split quadratic formations x, y and $f \in \text{Iso}(x, y)$, we write $[f]_\ell$ for the class of f in $\ell\text{Iso}(x, y)$ and $[f]_b$ for its class in $\text{bIso}(x, y)$. Recall from Remark 4.7 that given quadratic forms v and b , the left action of $\text{Aut}(\partial v)$ on $\text{Iso}(\partial v, \partial b)$ does not descend to an action of $\text{Aut}(\partial v)$ on $\ell\text{Iso}(\partial v, \partial b)$. However, if $g \in \text{Aut}(\partial v)$ and $\delta \in \text{Iso}(\partial v, \partial b)$, it is possible to consider the class $[\delta \circ g]_\ell \in \ell\text{Iso}(\partial v, \partial b)$. Keeping this in mind, the main result of this section is the following.

Theorem 7.7. *Let $q \geq 4$ be even and let $\bar{\nu}_0: M_0^{2q} \rightarrow B$ be an even, split-free normal $(q-1)$ -smoothing. Set $(V, \theta) := (K_q(M_0), \theta_{M_0})$, and $r := \text{rk}(V)$. For every $f \in \text{Aut}_{H_\varepsilon(\Lambda^r)}(\partial(V, \theta))$ there is an even, split-free normal $(q-1)$ -smoothing $(M_2^{2q}, \bar{\nu}_2)$ such that*

- (i) $(M_0, \bar{\nu}_0)$ and $(M_2, \bar{\nu}_2)$ are stably diffeomorphic;
- (ii) there is an isometry $(K_q(M_2), \theta_{M_2}) \cong (V, \theta)$, and in particular

$$\ell\text{Iso}(\partial(K_q(M_2), \theta_{M_2}), \partial(H^q(B), -\theta_{M_0})) \cong \ell\text{Iso}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}));$$

- (iii) $[\delta_{\bar{\nu}_2}]_\ell$ and $[\delta_{\bar{\nu}_0} f]_\ell$ belong to $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$;
- (iv) $[\delta_{\bar{\nu}_2}]_\ell$ and $[\delta_{\bar{\nu}_0} f]_\ell$ agree in $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$.

The same holds in the topological category for $q = 2$, assuming that $\pi_1(B)$ is a good group.

Remark 7.8. The geometric construction in this proof is similar to that in the proof of Theorem 3.10. We perform a set of trivial surgeries, and then a second set of surgeries on a “diagonal Lagrangian”. The difference is that Theorem 7.7 tracks the algebra differently. In order to obtain a bordism realising the desired change on boundary isomorphisms/primitive embeddings, it is much easier to return to the geometric construction than to track the change through the map δ from the sequence (1). The reason is that the definition of δ involves two different types of stabilisation which are tricky to control in this setting. We will discuss this further after Corollary 7.12.

Proof. First we will construct a (B, ξ) -cobordism W based on $(M_0, \bar{\nu}_0)$. Since $\bar{\nu}_0$ is split-free, the Λ -module $K_q(M_0) \cong V$ is finitely generated and free. Using r to denote the rank of V , let W_0 be the cobordism obtained from $M_0 \times [0, 1]$ by adding trivial q -handles along $M_0 \times \{1\}$ with trivial framing, one handle for each element of a basis for V . In other words, we consider the boundary connected sum:

$$\begin{aligned} W_0 &:= (M_0 \times [0, 1]) \natural (\natural_{i=1}^r S^q \times D^{q+1}), \text{ with} \\ M_1 &:= M_0 \# (\#_{i=1}^r S^q \times S^q). \end{aligned}$$

Let $\bar{\nu}_1: M_1 \rightarrow B$ be the normal $(q-1)$ -smoothing on M_1 obtained from $\bar{\nu}_0$ by imposing the trivial (B, ξ) -structure on the $S^q \times S^q$ connected summands. In order to obtain the (B, ξ) -cobordism W , we will perform r additional q -surgeries on M_1 . To do so, we first prescribe a rank r sublagrangian of $K_q(M_1)$; we then argue why it is possible to perform surgery along generators of this subspace. Consider the following decomposition

$$(29) \quad (K_q(M_1), \theta_{M_1}) \cong H_\varepsilon(V) \oplus (K_q(M_0), \theta_{M_0}) \cong ((V, \theta) \cup_f (V, -\theta)) \oplus (V, \theta),$$

the primitive embedding $j'_f: (V, -\theta) \hookrightarrow (V, \theta) \cup_f (V, -\theta)$ described in Lemma 4.13, and the sublagrangian

$$(30) \quad D := \{(j'_f(x), -x) \mid x \in V\} \subseteq K_q(M_1).$$

Note that $(V, \theta) \cup_f (V, -\theta) \cong H_\varepsilon(V)$ because $f \in \text{Aut}_{H_\varepsilon(\Lambda^r)}(\partial(V, \theta))$. One way to see that D is a summand is to use item (c) of Lemma 7.9 below.

We assert that we can represent a basis of D by r framed embedded spheres, and perform q -surgeries on M_1 along these spheres. The relative Hurewicz theorem implies that the Hurewicz map $K\pi_q(M_0) \rightarrow K_q(M_0)$ is surjective. Consequently, every element of $K_q(M_0)$ comes from an element x in $K\pi_q(M_0)$. Since D is a sublagrangian of $K_q(M_0)$, we deduce that $\lambda_{M_0}(x, x) = 0$, where λ_{M_0} denotes the Λ -valued intersection form on M_0 . As q is even and M_0 is orientable, this implies that the Wall form vanishes on x . The r generators of D can therefore be represented by framed embedded q -spheres on which we can perform surgery. For the $q = 2$ case, using the same proof as in Remark 3.11, the generators of D can be represented by framed, topologically embedded, locally flat 2-spheres, on which we can perform surgery.

Taking the trace and effect of these surgeries, we obtain

$$W := (M_1 \times [0, 1]) \cup \bigcup_{i=1}^r h_i^{r+1} \text{ and } M_2 := \partial_+ W.$$

Since we performed surgeries along elements of $K\pi_q(M_1)$, the (B, ξ) -structure on M_1 extends to (B, ξ) -structures $\bar{\nu}_2$ on M_2 and $\bar{\nu}$ on W ; recall the end of Section 3.1. As M_0 and M_2 are (B, ξ) -bordant and have the same Euler characteristic, they are stably diffeomorphic [Kre99, Theorem C], establishing (i) of the theorem.

Next we prove properties (ii) and (iii) of $(M_2, \bar{\nu}_2)$, namely that $(K_q(M_2), \theta_{M_2}) \cong (V, \theta)$, and that both

$$\delta_{\bar{\nu}_2} \in \text{Iso}(\partial(K_q(M_2), \theta_{M_2}), \partial(H^q(B), -\theta_{M_0})) \text{ and } \delta_{\bar{\nu}_0} f \in \text{Iso}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$$

define classes in

$$\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0})).$$

Note that if we show $(K_q(M_2), \theta_{M_2}) \cong (V, \theta)$, it follows straight away that $\delta_{\bar{\nu}_2}$ determines a class in $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$, so half of (ii) follows immediately from (iii). From now on, we use the shorthand $\delta_i := \delta_{\bar{\nu}_i}$ for the boundary isomorphism associated to $(M_i, \bar{\nu}_i)$.

Lemma 7.9. *Let $(V, \theta), (V', \theta'), (V'', \theta'')$ be ε -quadratic forms, and let $f: \partial(V'', \theta'') \cong_s \partial(V, \theta)$ and $f': \partial(V, \theta) \cong_s \partial(V', \theta')$ be stable isomorphisms. There is an isometry*

$$(31) \quad \varphi: ((V'', \theta'') \cup_f (V, -\theta)) \oplus ((V, \theta) \cup_{f'} (V', -\theta')) \xrightarrow{\cong} H_\varepsilon(V) \oplus (V'', \theta'') \cup_{f' \circ f} (V', -\theta')$$

that satisfies the following properties:

(a) φ takes the isometric copy of $(V', -\theta')$ to itself:

$$\varphi \circ \begin{pmatrix} 0 \\ j'_{f'} \end{pmatrix} = \begin{pmatrix} 0 \\ j'_{f'f} \end{pmatrix};$$

(b) φ restricts to an isometry $\varphi|$ on the first three summands; namely the following diagram of isometries and isometric injections commutes:

$$\begin{array}{ccc} (V'', \theta'') \cup_f (V, -\theta) \oplus (V, \theta) \cup_{f'} (V', -\theta') & \xrightarrow{\varphi} & H_\varepsilon(V) \oplus (V'', \theta'') \cup_{f'f} (V', -\theta') \\ \text{Id} \oplus j'_{f'} \uparrow & & \text{Id} \oplus j_f \uparrow \\ (V'', \theta'') \cup_f (V, -\theta) \oplus (V, \theta) & \xrightarrow{\varphi|} & H_\varepsilon(V) \oplus (V'', \theta''); \end{array}$$

(c) φ takes the diagonal lagrangian of $(V, -\theta) \oplus (V, \theta)$ to the lagrangian $V \oplus 0$ of $H_\varepsilon(V)$:

$$\varphi \circ \begin{pmatrix} j'_f \\ -j'_{f'} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

Proof. We use the notation from Lemma 4.9 for the components of the stable isomorphisms f and f' . Namely, we write a, b, s, a_1, b_1, a_3 (respectively $a', b', s', a'_1, b'_1, a'_3$) for the components of f (respectively f'). A fairly long but straightforward computation shows that

$$\varphi = \begin{pmatrix} a & -\varepsilon s & -s^* \lambda & -s^* a'^* \\ 0 & 1 & -\lambda & -a'^* \\ 1 & 0 & b & -b_1 a'_1{}^* \\ 0 & 0 & 0 & 1 \end{pmatrix} : V'' \oplus V^* \oplus V \oplus V'^* \xrightarrow{\cong} V \oplus V^* \oplus V'' \oplus V'^*$$

provides the desired isometry. The three properties of φ are also established by a direct computation. \square

We continue with the proof of Theorem 7.7. In order to identify $(H_q(M_2), \theta_{M_2})$ and $(K_q(M_2), \theta_{M_2})$, we describe the corresponding quadratic forms, but for the intermediate manifold M_1 . Using (29) and Lemma 7.9 (b) (with $(V'', \theta'') = (V, \theta), (V', \theta') = (H^q(B), -\theta_{M_0})$ and $f' = \delta_0$), we obtain

$$(32) \quad \begin{aligned} (K_q(M_1), \theta_{M_1}) &= H_\varepsilon(V) \oplus (K_q(M_0), \theta_{M_0}) = ((V, \theta) \cup_f (V, -\theta)) \oplus (V, \theta) \\ &\xrightarrow{\varphi|, \cong} H_\varepsilon(V) \oplus (V, \theta). \end{aligned}$$

Next, we obtain a similar description for $(H_q(M_1), \theta_{M_1})$. First, identifying $(K_q(M_0), \theta_{M_0})$ with (V, θ) , recall from Remark 7.6 that $(H_q(M_0), \theta_{M_0}) = (V, \theta) \cup_{\delta_0} (H^q(B), \theta_{M_0})$. Using the

definition of M_1 , this decomposition, and the isomorphism from Lemma 7.9, we deduce that

$$(33) \quad \begin{aligned} (H_q(M_1), \theta_{M_1}) &= H_\varepsilon(V) \oplus (H_q(M_0), \theta_{M_0}) = ((V, \theta) \cup_f (V, -\theta)) \oplus ((V, \theta) \cup_{\delta_0} (H^q(B), \theta_{M_0})) \\ &\xrightarrow{\varphi, \cong} H_\varepsilon(V) \oplus (V, \theta) \cup_{\delta_0 \circ f} (H^q(B), \theta_{M_0}). \end{aligned}$$

Items (b) and (c) of Lemma 7.9 imply that the sublagrangian D is mapped by φ to the lagrangian $V \oplus 0$ of $H_\varepsilon(V)$. By definition of M_2 and looking at (32) and (33), the second set of surgeries kills the $H_\varepsilon(V)$ -hyperbolic summands in $(H_q(M_1), \theta_{M_1})$ and $(K_q(M_1), \theta_{M_1})$ and thus

$$(34) \quad \begin{aligned} (K_q(M_2), \theta_{M_2}) &= (V, \theta); \\ (H_q(M_2), \theta_{M_2}) &= (V, \theta) \cup_{\delta_0 \circ f} (H^q(B), \theta_{M_0}). \end{aligned}$$

This shows (ii) and that $\delta_{\bar{\nu}_2}$ determines a class in $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$. It also implies that $\iota_{\bar{\nu}_2}$ is given by $(\frac{1}{0})$. Note that (34) also shows that the normal smoothing $(M_2, \bar{\nu}_2)$ is even and split-free: the target fibration is unchanged and $K_q(M_0) \cong V$ is finitely generated, and free since the normal smoothing $(M_0, \bar{\nu}_0)$ is even and split-free. Consequently, the 2-sided primitive embedding $(\iota_2, j_2) := (\iota_{\bar{\nu}_2}, j_{\bar{\nu}_2})$ is defined. Remark 7.6 implies that $[\delta_2]_\ell$ belongs to the set $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$. This establishes the property (iii) of the normal smoothing $(M_2, \bar{\nu}_2)$ since (34) shows that $[\delta_0 \circ f]_\ell \in \ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$.

Next we verify property (iv) of $(M_2, \bar{\nu}_2)$. We use (34) to identify (V, θ) with $(K_q(M_2), \theta_{M_2})$. We must show that $[\delta_2]_\ell$ coincides with $[\delta_0 \circ f]_\ell$ in the one-sided boundary automorphism set $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$. We prove that the two following 2-sided primitive embeddings coincide:

$$\begin{aligned} (j_{\delta_0 f}, j'_{\delta_0 f}) &:= \text{Pr}([\delta_0 f]_\ell) \in \ell\text{Prim}_{(H_q(M_2), \theta_{M_2})}((V, \theta), (H^q(B), \theta_{M_0})), \text{ and} \\ ([\iota_2, j_2]_\ell) &\in \ell\text{Prim}_{(H_q(M_2), \theta_{M_2})}((V, \theta), (H^q(B), \theta_{M_0})). \end{aligned}$$

Since (34) established that $(H_q(M_2), \theta_{M_2}) = (V, \theta) \cup_{\delta_0 \circ f} (H^q(B), \theta_{M_0})$, the definition of Pr from Construction 6.7 implies that $j_{\delta_0 f} = (\frac{1}{0})$. Below (34), we already argued that $\iota_2 = (\frac{1}{0})$, and thus $\iota_2 = j_{\delta_0 f}$. It remains to show that $j_2 = j'_{\delta_0 f}$.

Recall that the (B, ξ) -cobordism W_1 , originally obtained from M_1 as the trace of the q -surgeries on the “diagonal sublagrangian”, could also be understood as the trace of trivial $(q-1)$ -surgeries on M_2 . Similarly, W_0 was defined as the trace of trivial $(q-1)$ -surgeries on M_0 . For $i = 0, 1$, use $\bar{\nu}_{W_i}: W_i \rightarrow B$ to denote the (B, ξ) -structure on W_i , and consider the composition

$$j_{W_i}: H^q(B) \xrightarrow{\bar{\nu}_{W_i}^*} H^q(W_i) \xrightarrow{PD} H_{q+1}(W_i, \partial W_i).$$

For $k = 0, 1$, we additionally consider the maps $p_k: H_{q+1}(W_0, \partial W_0) \rightarrow H_q(\partial W_0) \rightarrow H_q(M_k)$ obtained by composing the connecting homomorphism from the long exact sequence of the pair $(W_0, \partial W_0)$ with the projection $H_q(\partial W_0) = H_q(M_0) \oplus H_q(M_1) \rightarrow H_q(M_k)$. The exact same process gives rise to maps $p'_k: H_{q+1}(W_1, \partial W_1) \rightarrow H_q(\partial W_1) \rightarrow H_q(M_k)$ for $k = 1, 2$. Recall that for $i = 0, 1, 2$, we defined $j_i := j_{\bar{\nu}_i}$ as $PD_{M_i} \circ \bar{\nu}_i^*$. A short diagram chase involving

Poincaré duality and the definition of the j_i yields the following commutative diagram:

$$(35) \quad \begin{array}{ccccc} H_q(M_0) & \xleftarrow{p_0} & H_{q+1}(W_0, \partial W_0) & \xrightarrow{p_1} & H_q(M_1) \\ & \searrow^{j_0} & \uparrow^{j_{W_0}} & \nearrow^{j_1} & \downarrow = \\ & & H^q(B) & & \\ & \swarrow_{j_2} & \downarrow^{j_{W_1}} & \searrow_{j_1} & \\ H_q(M_2) & \xleftarrow{p'_2} & H_{q+1}(W_1, \partial W_1) & \xrightarrow{p'_1} & H_q(M_1). \end{array}$$

Since the cobordism W_1 was obtained as the trace of trivial $(q-1)$ -surgeries on M_2 along $V = \varphi(D)$, we know that

$$W_1 = (M_2 \times [0, 1]) \natural (\natural_{i=1}^r S^q \times D^{q+1}) \simeq M_2 \vee \bigvee_{i=1}^r S^q \text{ and } M_1 \cong M_2 \# (\#_{i=1}^r S^q \times S^q).$$

We deduce that inclusion induces an isomorphism $H_i(M_2) \cong H_i(W_1)$ for $i \neq q$. Also, we infer that the inclusion induced map $\iota: H_q(\partial W_1) \rightarrow H_q(W_1)$ is surjective. As W_1 is a cobordism between M_1 and M_2 , the inclusion induces an isomorphism $H_i(M_1) \oplus H_i(M_2) \cong H_i(\partial W_1)$. Combining these observations, the long exact sequence of the pair $(W_1, \partial W_1)$ now looks as follows:

$$\begin{array}{ccccc} H_{q+1}(W_1, \partial W_1) & \hookrightarrow & H_q(\partial W_1) & \longrightarrow & H_q(W_1) \\ \uparrow \cong & & \uparrow \cong & \searrow & \uparrow \cong \\ & & & H_q(M_1) & \\ & & & \cong \uparrow & \\ & & & V \oplus V^* \oplus H_q(M_2) & \\ & & \nearrow \text{pr}_1 & & \\ V^* \oplus H_q(M_2) & \hookrightarrow & (V \oplus V^* \oplus H_q(M_2)) \oplus H_q(M_2) & \longrightarrow & V \oplus H_q(M_2) \end{array}$$

Since M_1 is also obtained by trivial surgeries on M_0 , the same reasoning can be applied on the trace cobordism W_0 by replacing M_1, M_2 by M_1, M_0 . Using the resulting identifications, we deduce that $j_{W_0} = \begin{pmatrix} 0 \\ j_0 \end{pmatrix}$ (the normal $(q-1)$ -smoothing on M_0 was extended trivially over the cobordism W_0) and therefore the commutative diagram displayed in (35) takes the following form:

$$\begin{array}{ccccc} H_q(M_0) & \xleftarrow{(0 \ 1)} & V^* \oplus H_q(M_0) & \xrightarrow{\begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}} & H_\varepsilon(V) \oplus H_q(M_0) \\ & \searrow^{j_0} & \uparrow \begin{pmatrix} 0 \\ j_0 \end{pmatrix} & \nearrow^{j_1} & \downarrow \varphi \\ & & H^q(B) & & \\ & \swarrow_{j_2} & \downarrow^{j_{W_1}} & \searrow_{\varphi \circ j_1} & \\ H_q(M_2) & \xleftarrow{(0 \ 1)} & V^* \oplus H_q(M_2) & \xrightarrow{p'_1 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}} & H_\varepsilon(V) \oplus H_q(M_2). \end{array}$$

The commutativity of the upper right triangle now gives $j_1 = (0 \ 0 \ j_0)^T$. Lemma 7.9 (a) then implies that $\varphi j_1 = (0 \ 0 \ j_{\delta_0 f})^T$. The commutativity of the bottom right triangle yields $p'_1 j_{W_1} = \varphi j_1$. Applying the splitting $(\begin{smallmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{smallmatrix})$ of p'_1 to both sides of this equation, we deduce that $j_{W_1} = (\begin{smallmatrix} 0 \\ j_{\delta_0 f} \end{smallmatrix})$. We conclude that $j_2 = (0 \ 1) j_{W_1} = j_{\delta_0 f}$, as desired.

Summarising, we have shown that the 2-sided primitive embeddings $(j_{\delta_0 f}, j'_{\delta_0 f}) := \text{Pr}([\delta_0 f]_\ell)$ and $(\iota_2, j_2) := \text{Pr}([\delta_2]_\ell)$ agree in ℓPrim . By Theorem 6.11 this implies that the boundary isomorphisms $[\delta_0 \circ f]_\ell$ and $[\delta_2]_\ell$ agree in the set $\ell\text{Iso}_{(H_q(M_2), \theta_{M_2})}(\partial(V, \theta), \partial(H^q(B), -\theta_{M_0}))$. This concludes the proof of (iv) and therefore of Theorem 7.7. \square

Example 7.10. We describe the diagonal sublagrangian $D \subseteq K_q(M_1)$ from (30) in a concrete example with $\Lambda = \mathbb{Z}$. Namely, we assume that $(M_0, \bar{v}_0) = (N_{y_1, y_2}, \bar{v}_{y_1, y_2})$ is an even, split-free normal $(2k - 1)$ -smoothing whose associated 2-sided primitive embedding is the one from our running example (namely Example 6.3): we set $q = y_1 y_2$ and consider

$$(i_{y_1, y_2}, j_{y_1, y_2}) := \left((\mathbb{Z}, q) \xrightarrow{\begin{pmatrix} y_1 \\ y_2 \end{pmatrix}} \left(\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) \xleftarrow{\begin{pmatrix} -y_1 \\ y_2 \end{pmatrix}} (\mathbb{Z}, -q) \right).$$

Such smoothings exist, as described in Propositions 9.3 and 9.5 below. In the notation of the proof of Theorem 7.7, we have $(V, \theta) = K_q(M_0) = (\mathbb{Z}, q)$, so that $K_q(M_1) = \mathbb{Z}^3 \supset D$. In this case, in order to describe D , we first rewrite the 2-sided primitive embedding as a union, along $[f] := [\delta(i_{y_1, y_2}, j_{y_1, y_2})]$. This was done in Example 6.12 and the outcome was

$$\left((\mathbb{Z}, q) \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} \left(\mathbb{Z}^2, \begin{pmatrix} q & 0 \\ a & -s \end{pmatrix} \right) \xleftarrow{\begin{pmatrix} b \\ -2q \end{pmatrix}} (\mathbb{Z}, -q) \right),$$

with $a = b = -(y_1 z_2 + y_2 z_1)$ and $s = -z_1 z_2$. As a consequence, we deduce that in this case, we have $j'_f = (\begin{smallmatrix} b \\ -2q \end{smallmatrix})$ and therefore

$$D = \{(j'_f(x), -x) \mid x \in \mathbb{Z}\} = \mathbb{Z} \begin{pmatrix} -(y_1 z_2 + y_2 z_1) \\ -2y_1 y_2 \\ -1 \end{pmatrix} \subseteq \mathbb{Z}^3 \cong K_q(M_1).$$

As a reality check, a brief calculation shows that $\lambda_{M_1} = (\begin{smallmatrix} 2q & a \\ a & -2s \end{smallmatrix}) \oplus (2q)$ does indeed vanish on D (for this calculation, it is helpful once again to recall that $y_2 z_1 - y_1 z_2 = 1$). This diagonal lagrangian was used implicitly in the construction of the manifolds $N_{a, b}$ from our previous article [CCPS21].

It is worth noting that we do not claim that in general M_0 and M_2 have isometric intersection forms, even though their kernel forms are isometric. Corollary 7.12 will exhibit conditions under which this favourable situation occurs, but first a remark which introduces some notation.

Remark 7.11. Given split quadratic formations x, y and $f \in \text{Iso}(x, y)$, we again write $[f]_\ell$ for the class of f in $\ell\text{Iso}(x, y)$ and $[f]_b$ for the class of f in $b\text{Iso}(x, y)$. Given three isomorphic quadratic forms $b \cong v \cong v'$, recall from Remark 6.5 that a boundary isomorphism $\delta \in \text{Iso}(\partial v', \partial b)$ defines an element $[\delta]_\ell \in \ell\text{Iso}(\partial v, \partial b)$ and an element $[\delta]_b \in b\text{Aut}(\partial v)$, but does not determine an element in $\ell\text{Aut}(\partial v)$.

Keeping Remark 7.11 in mind, the next corollary provides a situation in which we can realise elements of bAut sets as boundary isomorphisms associated to normal smoothings.

Corollary 7.12. *Let $q \geq 4$ be even, let (M_0^{2q}, \bar{v}_0) be an even, split-free normal $(q-1)$ -smoothing, and use n to denote the rank of $(V, \theta) := (K_q(M_0), \theta_{M_0})$. If there is an isometry $(H^q(B), -\theta_{M_0}) \cong (V, \theta)$ and if $[\delta_{\bar{v}_0}]_b = \text{Id} \in \text{bAut}(\partial(V, \theta))$, then for any boundary automorphism $f \in \text{Aut}_{H_\varepsilon(\Lambda^n)}(\partial(V, \theta))$, there exists an even, split-free normal $(q-1)$ -smoothing (M_2, \bar{v}_2) such that*

- (i) (M_0, \bar{v}_0) and (M_2, \bar{v}_2) are stably diffeomorphic;
- (ii) M_0 and M_2 have intersection forms isometric to $H_\varepsilon(\Lambda^n)$;
- (iii) the boundary isomorphism associated to (M_2, \bar{v}_2) is given by

$$[\delta_{\bar{v}_2}]_b = [f]_b \in \text{bAut}_{H_\varepsilon(\Lambda^n)}(\partial(V, \theta)).$$

The same holds in the topological category for $q = 2$, assuming that $\pi_1(B)$ is a good group.

Proof. During this proof, we abbreviate $(H^q(B), -\theta_{M_0}), (K_q(M_0), \theta_{M_0}), (K_q(M_2), \theta_{M_2})$ by b, v, v' , and we write $\delta_0 \in \text{Iso}(\partial v, \partial b), \delta_2 \in \text{Iso}(\partial v', \partial b)$ for the associated boundary isomorphisms. The assumption that $[\delta_0]_b = \text{Id}$ ensures that the intersection form of M_0 is hyperbolic. Apply Theorem 7.7 to $f \in \text{Aut}_{H_\varepsilon(\Lambda^n)}(\partial(V, \theta))$ in order to obtain a $(q-1)$ -smoothing (M_2, \bar{v}_2) that is stably diffeomorphic to (M_0, \bar{v}_0) , whose associated boundary isomorphism is given by

$$(36) \quad [\delta_2]_\ell = [\delta_0 f]_\ell \in \ell\text{Iso}(\partial v, \partial b),$$

We argue that $[\delta_2]_b = [f]_b$. Recall from Remark 4.8 that (36) means that for some $\tilde{\delta}_2 \in \text{Iso}(\partial v, \partial b)$, we have $\tilde{\delta}_2 = (\delta_0 \circ f) \circ \partial h^{-1}$ for some isometry $h: v \rightarrow v'$. Similarly, $[\delta_0]_b = \text{Id}$ means that for some $\tilde{\delta}_0 \in \text{Aut}(\partial v)$, we have $[\tilde{\delta}_0]_b = \text{Id}$, where $\tilde{\delta}_0 = \partial h_2 \circ \delta_0$ for some isometry $h_2: b \rightarrow v$. In turn this implies $\tilde{\delta}_0 = \partial h_3 \circ \text{Id} \circ \partial h_4^{-1}$ for some isometries $h_3, h_4: v \rightarrow v'$. In order to study $[\delta_2]_b$, we choose $\tilde{\delta}'_2 \in \text{bAut}(\partial v)$ so that $\tilde{\delta}'_2 = \partial h_1 \circ \tilde{\delta}_2$ for some isometry $h_1: b \rightarrow v$. Summing up, we obtain

$$\begin{aligned} [\delta_2]_b &:= [\tilde{\delta}'_2]_b = [\partial h_1 \circ \tilde{\delta}_2]_b = [\partial h_1 \circ (\delta_0 \circ f \circ \partial h^{-1})]_b = [\partial h_1 \circ (\partial h_2^{-1} \circ \tilde{\delta}_0) \circ f \circ \partial h^{-1}]_b \\ &= [\partial h_1 \circ \partial h_2^{-1} \circ (\partial h_3 \circ \partial h_4^{-1}) \circ f \circ \partial h^{-1}]_b = [f]_b. \end{aligned}$$

Finally, we argue that M_0 and M_2 both have hyperbolic intersection forms. For M_0 , this follows from Remark 7.6 and from our assumption that $\delta_0 = [\text{Id}]_b$, and so we focus on M_2 . Recall from (34) that $(H_q(M_2), \theta_{M_2}) \cong (V, \theta) \cup_{\delta_0 f} (H^q(B), \theta_{M_0})$. Since we have $(H^q(B), \theta_{M_0}) \cong (V, -\theta)$ (by assumption), this implies that $(H_q(M_2), \theta_{M_2}) \cong (V, \theta) \cup_{[\delta_0 f]_b} (V, -\theta)$. Since we have just established that $[\delta_0 f]_b = [f]_b$ and since $f \in \text{Aut}_{H_\varepsilon(\Lambda^n)}(\partial(V, \theta))$, we deduce that $(H_q(M_2), \theta_{M_2})$ is indeed hyperbolic. This completes the proof of the corollary. \square

We conclude this section by describing the application of Theorem 7.7 discussed in Remark 7.8. We recall the exact sequence of [CS11, Theorem 5.11] that was displayed in (1). Given a quadratic form $v = (V, \theta)$ over R whose symmetrisation $\lambda: V \rightarrow V^*$ is injective (we call such a form *non-degenerate*), [CS11, Theorem 5.11 and Lemma 6.2] establish the existence of a sequence

$$(37) \quad L_{2q+1}^s(R) \xrightarrow{\rho} \ell_{2q+1}(v, v') \xrightarrow{\delta} \text{bIso}(\partial v, \partial v') \xrightarrow{\kappa} L_{2q}^s(R).$$

Here, $\ell_{2q+1}(v, v')$ consists of quasi-formations whose induced form and its orthogonal complement respectively coincide 0-stably with v and v' , the dashed arrow ρ denotes the action of $L_{2q+1}^s(R)$ on $\ell_{2q+1}(v, v')$ by direct sum, κ is the map induced by the assignment $f \mapsto v \cup_f v'$, and the map δ is (roughly) described as follows. For a quasi-formation $x = ((M, \psi); F, V)$, use $j: (V, \theta) \hookrightarrow (M, \psi)$ to denote the split isometric inclusion of the induced form into (M, ψ) . The map δ assigns to x the boundary isomorphism f_j constructed in Proposition 4.14.

Remark 7.13. Given split quadratic formations x, y and $f \in \text{Iso}(x, y)$, we again write $[f]_\ell$ for the class of f in $\ell\text{Iso}(x, y)$ and $[f]_b$ for the class of f in $\text{bIso}(x, y)$. Even though $\ell\text{Iso}(x, y)$ is not a group, given a third split quadratic formation z , we can consider the map

$$\begin{aligned} \ell\text{Iso}(x, y) \times \ell\text{Iso}(x, z) &\rightarrow \text{bIso}(y, z) \\ ([f]_\ell, [g]_\ell) &\mapsto [g^{-1} \circ f]_b. \end{aligned}$$

This map does not depend on the choice of representatives for $[f]_\ell$ and $[g]_\ell$: indeed for $h_1 \in \text{Aut}(y)$ and $h_2 \in \text{Aut}(z)$, we can use the definitions of the actions to obtain the following equalities:

$$[(g \circ \partial h_2^{-1})^{-1} \circ (f \circ \partial h_1^{-1})]_b = [\partial h_2 \circ (g^{-1} \circ f) \circ \partial h_1^{-1}]_b = [(h_1, h_2) \cdot (g \circ f^{-1})]_b = [g \circ f^{-1}]_b.$$

Now let b, v, v' be quadratic forms with an isometry $v \xrightarrow{\cong} v'$. As explained in Section 4.2, even though we cannot identify $\text{Iso}(\partial v, \partial b)$ with $\text{Iso}(\partial v', \partial b)$, we can identify $\ell\text{Iso}(\partial v, \partial b)$ with $\ell\text{Iso}(\partial v', \partial b)$. In particular, if $g \in \ell\text{Iso}(\partial v', \partial b)$, then there is an isometry $h: v' \rightarrow v$ and a $\tilde{g} \in \ell\text{Iso}(\partial v, \partial b)$ such that \tilde{g} and $g \circ \partial h^{-1}$ agree in $\ell\text{Iso}(\partial v, \partial b)$. In particular, there is a map

$$\begin{aligned} \ell\text{Iso}(\partial v, \partial b) \times \ell\text{Iso}(\partial v', \partial b) &\rightarrow \text{bAut}(\partial v) \\ ([f]_\ell, [g]_\ell) &\mapsto [\tilde{g}^{-1} \circ f]_b =: [g]_\ell^{-1} \circ [f]_\ell. \end{aligned}$$

From the discussion above, we already know that this assignment does not depend on the choice of representatives for $[f]_\ell$ and $[g]_\ell$. This map also does not depend on the choice of the isometry h : indeed if $h_1, h_2: v' \xrightarrow{\cong} v$ are two isometries, then

$$[\partial h_1 \circ g^{-1} \circ f]_b = [(\partial h_2 \circ \partial h_1^{-1}) \circ \partial h_1 \circ g^{-1} \circ f]_b = [\partial h_2 \circ g^{-1} \circ f]_b.$$

Using the notation of Theorem 7.7, we apply this construction to the case where the quadratic forms b, v and v' are $(H^q(B), -\theta_{M_0})$, $(K_q(M_0), \theta_{M_0})$, and $(K_q(M_2), \theta_{M_2})$ respectively. In this case, the boundary isomorphism δ_0 associated to (M_0, \bar{v}_0) belongs to $\text{Iso}(\partial v, \partial b)$, while the associated normal smoothing δ_2 associated to (M_2, \bar{v}_2) belongs to $\text{Iso}(\partial v', \partial b)$. The remarks above nevertheless allow us to make sense of the composition $[\delta_2]_\ell^{-1} \circ [\delta_0]_\ell$ as an element of $\text{bAut}(\partial v)$.

The next result describes the image under the map $\delta: \ell_{2q+1}(v, v) \rightarrow \text{bAut}(\partial v)$ of the modified surgery obstruction for the (B, ξ) -cobordism (W, \bar{v}) constructed in Theorems 3.10 and 7.7.

Corollary 7.14. *Let $q \geq 4$ be even and let (M_0^{2q}, \bar{v}_0) be an even, split-free normal $(q-1)$ -smoothing with non-degenerate kernel form $(V, \theta) := (K_q(M_0), \theta_{M_0})$. Fix a quasi-formation $x = (H_\varepsilon(K); K, V)$ whose induced form is (V, θ) and satisfies $(V, \theta) \cong (V^\perp, \theta^\perp)$. Then there exists a normal (B, ξ) -cobordism (W, \bar{v}) , with modified surgery obstruction $\Theta(W, \bar{v}) = [x]$, from (M_0, \bar{v}_0) to another even, split-free normal smoothing (M_2, \bar{v}_2) such that*

$$\delta(\Theta(W, \bar{v})) = \delta(x) = [\delta_2]_\ell^{-1} \circ [\delta_0]_\ell \in \text{bAut}(\partial(V, \theta)).$$

The same holds in the topological category for $q = 2$, assuming that $\pi_1(B)$ is a good group.

Proof. During this proof, we abbreviate $(H^q(B), -\theta_{M_0}), (K_q(M_0), \theta_{M_0}), (K_q(M_2), \theta_{M_2})$ by b, v, v' . Since we assumed that $v \cong v^\perp$, we know that $[f]_b := \delta(x)$ belongs to $\text{bAut}(\partial v)$. As v is non-degenerate, the exact sequence displayed in (1) ensures that the union $v \cup_f v'$ is hyperbolic. We apply Theorem 7.7 to a representative $f^{-1} \in \text{Aut}(\partial v)$ of $\delta(x)^{-1}$ to obtain a (B, ξ) -cobordism $(W, \bar{\nu})$ with $[\delta_2]_\ell = [\delta_0 \circ f^{-1}]_\ell \in \ell\text{Aut}(\partial v, \partial b)$. Now pick any $\tilde{\delta}_2 \in \text{Iso}(\partial v, \partial b)$ as in Remark 7.13. In particular, there is an $h \in \text{Aut}(v)$ such that $\tilde{\delta}_2 = h \cdot (\delta_0 \circ f^{-1}) = \delta_0 \circ f^{-1} \circ \partial h^{-1}$. It follows that

$$\delta(x) = [f]_b = [\partial h \circ f]_b = [(\partial h \circ f \circ \delta_0^{-1}) \circ \delta_0]_b = [\tilde{\delta}_2^{-1} \circ \delta_0]_b = [\delta_2]_\ell^{-1} \circ [\delta_0]_\ell.$$

It remains to show that $\Theta(W, \bar{\nu}) = x$. Recall that $(W, \bar{\nu})$ was constructed by first performing $\text{rk}(V)$ trivial $(q-1)$ -surgeries on M_0 (yielding M_1), and then performing additional q -surgeries on the diagonal sublagrangian of $K_q(M_1) = (V, \theta) \oplus ((V, \theta) \cup_f (V, -\theta))$. We assert that this produces the (B, ξ) -cobordism of Theorem 3.10 which was shown to have surgery obstruction x . As $(M_0, \bar{\nu}_0)$ is even and split-free, $(K_q(M_0), \theta_{M_0})$ is its own free Wall form. Since we assumed that $(K\pi_q(M_0), \theta_{M_0})$ is isometric to the induced form (V, θ) , the assertion follows from the construction in Theorem 3.10; see in particular (5). This proves the equality $\Theta(W, \bar{\nu}) = x$, thus establishing the corollary. \square

8. HOMOTOPY INVARIANCE OF CERTAIN TWO-SIDED PRIMITIVE EMBEDDINGS

Let $\bar{\nu}: M \rightarrow B$ be a normal $(q-1)$ -smoothing. Due to the extra data afforded by $\bar{\nu}$, it is in general not clear that the two-sided primitive embedding associated to $(M, \bar{\nu})$ is a homeomorphism invariant, or even invariant under homotopy equivalences. On the other hand, this hurdle is not present when considering the two-sided primitive embedding associated to the functorial map $M \rightarrow P_{q-1}(M)$. In this section, we therefore develop the notion of the Postnikov two-sided primitive embedding of a manifold, show that this object is invariant under orientation-preserving homotopy equivalences, and then relate it to the two-sided primitive embedding of a particular normal $(q-1)$ -smoothing. We hope that this will be of independent interest as a general formalism for homotopy invariants.

8.1. Normal $(q-1)$ -types. Given $m \geq q$, this subsection introduces a specific kind of normal $(q-1)$ -smoothing $\bar{\nu}: M \rightarrow P_{q-1}(M) \times BO\langle m \rangle$ that we will use in Section 9.

Recall that the j -connected cover $X\langle j \rangle$ of a CW -complex X is a CW -complex which, up to homotopy equivalence, is characterized by the property that $X\langle j \rangle$ is j -connected and there is a fibration $p_j: X\langle j \rangle \rightarrow X$ inducing isomorphisms on π_i for $i > j$. In our case, we take $X = BO$ leading to a fibration

$$\vartheta_k: BO\langle k \rangle \rightarrow BO.$$

In this subsection, we are concerned with a closed, connected, oriented manifold M^{2q} such that for some $m \geq q$, the stable normal bundle $\nu: M \rightarrow BO$ lifts to a map

$$\nu_m(M): M \rightarrow BO\langle m \rangle.$$

Such manifolds are called m -parallelisable. The next lemma describes the normal $(q-1)$ -type of m -parallelisable $2q$ -manifolds.

Lemma 8.1. *Let M be a closed, connected, oriented $2q$ -manifold whose stable normal bundle lifts to a map $\nu_m(M): M \rightarrow BO\langle m \rangle$ for some $m \geq q$. The following map is a normal $(q-1)$ -smoothing*

$$p_{q-1}(M) \times \nu_m(M): M \rightarrow P_{q-1}(M) \times BO\langle m \rangle.$$

Furthermore, the normal $(q-1)$ -type of M is represented by the fibration

$$\vartheta_q \circ \text{pr}_2: P_{q-1}(M) \times BO\langle q \rangle \rightarrow BO,$$

where pr_2 is the projection onto the second factor.

Proof. Note that $\vartheta_q: BO\langle q \rangle \rightarrow BO$ is q -coconnected, and $\pi_i(P_{q-1}(M)) = 0$ for $i \geq q$. It follows that $\vartheta_q \circ \text{pr}_2$ is q -coconnected. Also $p_{q-1}(M) \times \nu_m(M)$ lifts the stable normal bundle of M :

$$\vartheta_m \circ \text{pr}_2 \circ (p_{q-1}(M) \times \nu_m(M)) = \vartheta_m \circ \nu_m(M) = \nu: M \rightarrow BO$$

Finally $p_{q-1}(M) \times \nu_m(M)$ is q -connected since $m \geq q$ and $p_{q-1}(M)$ is q -connected. \square

8.2. The Postnikov two-sided primitive embedding. We modify Definition 7.5 in order to obtain an invariant of manifolds, instead of an invariant of q -surjective maps. More precisely, under some technical conditions, we use the Postnikov $(q-1)$ -type to associate a two-sided primitive embedding $PPE(M)$ to an oriented manifold M^{2q} . After proving the invariance of $PPE(M)$ under orientation-preserving homotopy equivalences, we then relate $PPE(M)$ to the two-sided primitive embedding associated to the normal smoothing from Section 8.1.

Construction 8.2. Let π be a group and set $\Lambda := \mathbb{Z}[\pi]$. Given an automorphism $\varphi \in \text{Aut}(\pi)$ and an ε -quadratic form (V, θ) , we write $\overline{(V, \theta)}^\varphi = (\overline{V}^\varphi, \overline{\theta}^\varphi)$ for the quadratic form where

- (i) the underlying module \overline{V}^φ agrees with V as an abelian group but has left Λ -module structure $x \cdot v := \varphi(x)v$ for $x \in \Lambda$ and $v \in V$;
- (ii) the quadratic form is defined as $\overline{\theta}^\varphi(x, y) = \varphi^{-1}(\theta(x, y)) \in Q_\varepsilon(\Lambda)$. Here it is understood that φ extends to a ring automorphism $\varphi: \Lambda \rightarrow \Lambda$ which itself descends to $Q_\varepsilon(\Lambda)$.

If $i: (V, \theta) \rightarrow (M, \psi)$ is Λ -linear and preserves quadratic forms, then so does $i: \overline{(V, \theta)}^\varphi \rightarrow \overline{(M, \psi)}^\varphi$. The action of $\text{Aut}(\pi)$ on a two-sided primitive $(i, j) := v \hookrightarrow m \hookleftarrow v'$ is defined as

$$\overline{(i, j)}^\varphi := \overline{v}^\varphi \xrightarrow{i} \overline{m}^\varphi \xleftarrow{j} \overline{v'}^\varphi.$$

Since \overline{v}^φ need not be isomorphic to v , this construction in general does not provide an action of $\text{Aut}(\pi)$ on $\text{bPrim}_m(v, v')$. It does however provide an action of $\text{Aut}(\pi)$ on the universal collection

$$\text{bPrim}^\pi := \bigcup_{[m, v, v']} \text{bPrim}_m(v, v'),$$

where $[m, v, v']$ denotes isometry classes of triples of forms. We then say that two 2-sided primitive embeddings $(i_0, j_0), (i_1, j_1)$ agree in $\text{bPrim}^\pi / \text{Aut}(\pi)$ if (i_0, j_0) and $\overline{(i_1, j_1)}^\varphi$ agree in bPrim for some automorphism $\varphi \in \text{Aut}(\pi)$, meaning that there is a commutative diagram

with vertical maps isomorphisms:

$$\begin{array}{ccccc}
 v_0 & \xrightarrow{i_0} & m_0 & \xleftarrow{j_0} & v'_0 \\
 \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\
 \overline{v_1}^\varphi & \xrightarrow{i_1} & \overline{m_1}^\varphi & \xleftarrow{j_1} & \overline{v'_1}^\varphi.
 \end{array}$$

For a ring homomorphism $u: \Gamma \rightarrow \Lambda$, we write Λ_u for the (Λ, Γ) -bimodule with underlying abelian group Λ , where the left Λ -module structure is given by multiplication and the right Γ -module structure is given by $p \cdot \gamma = pu(\gamma)$, for $p \in \Lambda$ and $\gamma \in \Gamma$.

Remark 8.3. Fix a group π and set $\Lambda := \mathbb{Z}[\pi]$. Let M be a closed, connected, oriented $2q$ -manifold. Given an isomorphism $\psi: \pi_1(M) \xrightarrow{\cong} \pi$, we have that Λ_ψ is a $(\Lambda, \mathbb{Z}[\pi_1(M)])$ -bimodule, and the Λ -module $H_q(M; \Lambda_\psi)$ is endowed with a Λ -valued intersection form $\lambda_{M, \psi}$, obtained by composing the Poincaré duality isomorphism with evaluation. Since Λ_ψ admits a length zero free resolution as a right $\mathbb{Z}[\pi_1(M)]$ -module, we have $H_q(M; \Lambda_\psi) \cong \Lambda_\psi \otimes_{\mathbb{Z}[\pi_1(M)]} H_q(M; \mathbb{Z}[\pi_1(M)])$, and similarly in cohomology. Under this identification, it follows that $\lambda_{M, \psi}$ is isometric to $(p \otimes x, q \otimes y) \mapsto p\psi(\lambda_M(x, y))\overline{q}$.

Convention 8.4. Given $k > 0$, we always identify $\pi_1(P_k(M))$ with $\pi_1(M)$. This is possible because $P_k(M)$ is obtained from M by adding cells of dimension $\geq k+2 > 2$. In particular, $p_k(M): M \rightarrow P_k(M)$ induces maps $p_{q-1}(M)_*$ and $p_{q-1}(M)^*$ on twisted homology and cohomology respectively.

Definition 8.5. Fix a group π , and set $\Lambda := \mathbb{Z}[\pi]$. Let M be a closed, connected, oriented, even $2q$ -manifold where $p_{q-1}(M): M \rightarrow P_{q-1}(M)$ is split-free. Given an isomorphism $\psi: \pi_1(M) \xrightarrow{\cong} \pi$, the *Postnikov 2-sided primitive embedding of M* is defined as

$$PPE^\pi(M) = (\ker(p_{q-1}(M)_*) \hookrightarrow H_q(M; \Lambda_\psi) \hookrightarrow H^q(P_{q-1}(M; \Lambda_\psi)) \in \text{bPrim}^\pi / \text{Aut}(\pi).$$

When no identification is fixed, we will write $PPE(M) \in \text{bPrim}^{\pi_1(M)}$ for the analogous two-sided primitive embedding, but where the homology groups have coefficients in $\mathbb{Z}[\pi_1(M)]$. In this case we omit the superscript π . Note that $PPE(M)$ and $PPE^\pi(M)$ are indeed two-sided primitive embeddings by Example 7.3 and Lemma 7.4.

We show that $PPE^\pi(M)$ does not depend on the choice of the identification ψ .

Lemma 8.6. *Fix a group π , and set $\Lambda := \mathbb{Z}[\pi]$. Let M be a closed, connected, oriented $2q$ -manifold with $p_{q-1}(M): M \rightarrow P_{q-1}(M)$ even and split-free. The Postnikov 2-sided primitive embedding $PPE^\pi(M)$ of M does not depend on the identification $\pi_1(M) \cong \pi$.*

Proof. Let $\psi_i: \pi_1(M) \xrightarrow{\cong} \pi$ be isomorphisms for $i = 1, 2$. For concision, throughout this proof, we set $\overline{p} := p_{q-1}(M)$ as well as $K_q(M; \Lambda_{\psi_i}) := \ker(\overline{p}_*: H_q(M; \Lambda_{\psi_i}) \rightarrow H_q(P_{q-1}(M_i); \Lambda_{\psi_i}))$. It can be verified that $\psi_2\psi_1^{-1} \in \text{Aut}(\pi)$ induces Λ -linear maps on twisted homology and

cohomology that make the following diagram commute:

$$(38) \quad \begin{array}{ccccccc} & & & \xleftarrow{j_{\bar{p}}} & & & \\ & & & \swarrow & & \searrow & \\ K_q(M; \Lambda_{\psi_1}) & \xrightarrow{\iota_{\bar{p}}} & H_q(M; \Lambda_{\psi_1}) & \xleftarrow{\text{PD}} & H^q(M; \Lambda_{\psi_1}) & \xleftarrow{\bar{p}^*} & H^q(P_{q-1}(M); \Lambda_{\psi_1}) \\ & \cong \downarrow (\psi_2 \psi_1^{-1})_* & \cong \downarrow (\psi_2 \psi_1^{-1})_* & & \cong \downarrow (\psi_2 \psi_1^{-1})_* & & \cong \downarrow (\psi_2 \psi_1^{-1})_* \\ \overline{K_q(M; \Lambda_{\psi_2})}^{\psi_2 \psi_1^{-1}} & \xrightarrow{\iota_{\bar{p}}} & \overline{H_q(M; \Lambda_{\psi_2})}^{\psi_2 \psi_1^{-1}} & \xleftarrow{\text{PD}} & \overline{H^q(M; \Lambda_{\psi_2})}^{\psi_2 \psi_1^{-1}} & \xleftarrow{\bar{p}^*} & \overline{H^{2k}(P_{q-1}(M); \Lambda_{\psi_2})}^{\psi_2 \psi_1^{-1}} \\ & & & \swarrow & & \searrow & \\ & & & \xleftarrow{j_{\bar{p}}} & & & \end{array}$$

The change of coefficients map $(\psi_2 \psi_1^{-1})_*$ induces an isometry $\lambda_{M, \psi_1} \cong \overline{\lambda_{M, \psi_2}}^{\psi_2 \psi_1^{-1}}$. Indeed, the twisted intersection forms on $H_q(M; \Lambda_{\psi_1})$ and $\overline{H_q(M; \Lambda_{\psi_2})}^{\psi_2 \psi_1^{-1}}$ are both defined by composing duality with the evaluation map, and changing coefficients commutes with these maps. Since the two-sided primitive embeddings are even, this implies that $(\psi_2 \psi_1^{-1})_*$ preserves the quadratic forms. This concludes the proof of the lemma. \square

Note that $PPE^\pi(M)$ is not in general an invariant of M : changing the orientation on M changes its equivariant intersection form and may therefore change $PPE^\pi(M)$. On the other hand the next proposition shows that PPE^π is invariant under orientation-preserving homotopy equivalences.

Proposition 8.7. *Let M_1^{2q} and M_2^{2q} be closed, connected, oriented manifolds with fundamental group π , such that $p_{q-1}(M_i)$ is even and split-free for $i = 1, 2$. If $f: M_1 \rightarrow M_2$ is an orientation preserving homotopy equivalence, then $PPE^\pi(M_1)$ and $PPE^\pi(M_2)$ agree in $\text{bPrim}^\pi / \text{Aut}(\pi)$.*

Proof. Choose arbitrary isomorphisms $\psi_i: \pi_1(M_i) \cong \pi$ for $i = 1, 2$. Write $\bar{p}_i := p_{q-1}(M_i)$ as well as $K_q(M_i; \Lambda_{\psi_i}) := \ker((\bar{p}_i)_*)$ for $i = 1, 2$, and $\varphi := \psi_2 f_* \psi_1^{-1} \in \text{Aut}(\pi)$. The homotopy equivalence f and the automorphism φ induce Λ -linear isomorphism on twisted homology that fit into the following diagram:

$$(39) \quad \begin{array}{ccccccc} & & & \xleftarrow{j_{\bar{p}_1}} & & & \\ & & & \swarrow & & \searrow & \\ K_q(M_1; \Lambda_{\psi_1}) & \xrightarrow{\iota_{\bar{p}_1}} & H_q(M_1; \Lambda_{\psi_1}) & \xleftarrow{\text{PD}} & H^q(M_1; \Lambda_{\psi_1}) & \xleftarrow{\bar{p}_1^*} & H^q(P_{q-1}(M_1); \Lambda_{\psi_1}) \\ & f_* \downarrow \cong & f_* \downarrow \cong & & \cong \downarrow f^{-*} & & \cong \downarrow f^{-*} \\ K_q(M_2; \Lambda_{\psi_1 f_*^{-1}}) & \xrightarrow{\iota_{\bar{p}_1}} & H_q(M_2; \Lambda_{\psi_1 f_*^{-1}}) & \xleftarrow{\text{PD}} & H^q(M_2; \Lambda_{\psi_1 f_*^{-1}}) & \xleftarrow{\bar{p}_2^*} & H^q(P_{q-1}(M_2); \Lambda_{\psi_1 f_*^{-1}}) \\ & \varphi_* \downarrow \cong & \varphi_* \downarrow \cong & & \cong \downarrow \varphi_* & & \cong \downarrow \varphi_* \\ \overline{K_q(M_2; \Lambda_{\psi_2})}^\varphi & \xrightarrow{\iota_{\bar{p}_2}} & \overline{H_q(M_2; \Lambda_{\psi_2})}^\varphi & \xleftarrow{\text{PD}} & \overline{H^q(M_2; \Lambda_{\psi_2})}^\varphi & \xleftarrow{\bar{p}_2^*} & \overline{H^{2k}(P_{q-1}(M_2); \Lambda_{\psi_2})}^\varphi \\ & & & \swarrow & & \searrow & \\ & & & \xleftarrow{j_{\bar{p}_2}} & & & \end{array}$$

This diagram commutes (for the top central square, we use that f is an orientation-preserving homotopy equivalence) and, in the second column, both f_* and φ_* induce isometries. This concludes the proof of the proposition. \square

Construction 8.8. Let M be a closed, connected, oriented $2q$ -manifold with fundamental group π such that $p_{q-1}(M)$ is even and split-free. We wish to compare $PPE(M) \in \text{bPrim}^{\pi_1(M)}$ with the two-sided primitive embedding associated to the normal smoothing from Section 8.1.

Set $B := P_{q-1}(M) \times BO\langle m \rangle$ and consider the fibration $\xi: B \xrightarrow{\text{pr}_2} BO\langle m \rangle \xrightarrow{\vartheta_m} BO$. Assume that the stable normal bundle $M \rightarrow BO$ lifts to a map $\nu_m(M): M \rightarrow BO\langle m \rangle$. For $m \geq q$, Lemma 8.1 showed that the following map is a normal $(q-1)$ -smoothing

$$\bar{\nu} := p_{q-1}(M) \times \nu_m(M): M \rightarrow P_{q-1}(M) \times BO\langle m \rangle =: B.$$

Since $BO\langle m \rangle$ is simply-connected, we have $\pi_1(B) = \pi_1(P_{q-1}(M))$, which we can canonically identify with $\pi_1(M)$ as we have been doing up to now; recall Convention 8.4. Assuming that $\bar{\nu}$ is even and split-free, we saw in Definition 7.5 how to associate a two-sided primitive embedding to $(M, \bar{\nu})$ and we obtain

$$PE(M, \bar{\nu}) \in \text{bPrim}_{H_q(M; \mathbb{Z}[\pi_1(M)])}(K_q(M; \mathbb{Z}[\pi_1(M)]), H^q(B; \mathbb{Z}[\pi_1(M)])) \subseteq \text{bPrim}^{\pi_1(M)}.$$

Next we show that $PPE(M)$ and $PE(M, \bar{\nu})$ agree in $\text{bPrim}^{\pi_1(M)}$.

Proposition 8.9. *Let M be a closed, connected, oriented $2q$ -manifold such that $p_{q-1}(M)$ is even and split-free. Assume that M admits a map $\nu_m(M): M \rightarrow BO\langle m \rangle$ with $m \geq q$, which lifts the stable normal bundle of M and consider the normal $(q-1)$ -smoothing*

$$\bar{\nu} := p_{q-1}(M) \times \nu_m(M): M \rightarrow P_{q-1}(M) \times BO\langle m \rangle =: B.$$

If $\bar{\nu}$ is also even and split-free, then $PPE(M)$ and $PE(M, \bar{\nu})$ agree in $\text{bPrim}^{\pi_1(M)}$.

Proof. Set $\Lambda := \mathbb{Z}[\pi_1(B)] = \mathbb{Z}[\pi_1(M)]$. Since $\bar{\nu} = p_{q-1}(M) \times \nu_m(M)$, it suffices to prove that the projection $\text{pr}_1: B \rightarrow P_{q-1}(M)$ induces an isomorphism $(\text{pr}_1)_*: H_q(B; \Lambda) \cong H_q(P_{q-1}(M); \Lambda)$, as the isomorphism $(\text{pr}_1)^*: H^q(P_{q-1}(M); \Lambda) \cong H^q(B; \Lambda)$ then follows because $\bar{\nu}$ and $p_{q-1}(M)$ are split-free. To see that $(\text{pr}_1)^*$ is an isomorphism, consider the fibration $BO\langle m \rangle \rightarrow BO\langle m \rangle \times P_{q-1}(M) \xrightarrow{\text{pr}_1} P_{q-1}(M)$ and its homotopy long exact sequence. We see that $\text{pr}_1: \pi_i(BO\langle m \rangle \times P_{q-1}(M)) \rightarrow \pi_i(P_{q-1}(M))$ is an isomorphism for $i \leq q$. Applying the relative Hurewicz theorem to the universal covers of these spaces then implies the assertion. \square

Construction 8.10. Let M be a closed, connected, oriented $2q$ -manifold such that $p_{q-1}(M)$ is even and split-free. Assume that M admits a map $\nu_m(M): M \rightarrow BO\langle m \rangle$ with $m \geq q$, which lifts the stable normal bundle of M , set $\bar{\nu} := p_{q-1}(M) \times \nu_m(M)$, and consider the normal $(q-1)$ -smoothing

$$\bar{\nu} := p_{q-1}(M) \times \nu_m(M): M \rightarrow P_{q-1}(M) \times BO\langle m \rangle =: B.$$

We let $\xi = \vartheta_m \circ \text{pr}_{BO\langle m \rangle}: B \rightarrow BO$ and perform a compatible r -surgery on $(M, \bar{\nu})$ leading to a trace (B, ξ) -cobordism (W, M, M') , and therefore to a normal smoothing $\bar{\nu}': M' \rightarrow B$. Assume that either (i) $r = q$ or (ii) $r = q-1$ but with a surgery on unknotted framed embeddings. Note that $(M', \bar{\nu}')$ is again even and split-free. Also, we can identify $\pi_1(M')$ with $\pi_1(M)$.

The following proposition generalises Proposition 8.9.

Proposition 8.11. *Let M be a closed, connected, oriented $2q$ -manifold such that $p_{q-1}(M)$ is even and split-free. Assume that M admits a map $\nu_m(M): M \rightarrow BO\langle m \rangle$ with $m \geq q$, which lifts the stable normal bundle of M . Assume further that the normal $(q-1)$ -smoothing*

$$\bar{\nu} := p_{q-1}(M) \times \nu_m(M): M \rightarrow P_{q-1}(M) \times BO\langle m \rangle =: B$$

is split-free. If (W, M, M') is a trace (B, ξ) -cobordism as in Construction 8.10, then $PPE(M')$ and $PE(M', \bar{\nu}')$ agree in $\mathbf{bPrim}^{\pi_1(M)}$.

Proof. Set $\Lambda := \mathbb{Z}[\pi_1(B)] = \mathbb{Z}[\pi_1(M')] = \mathbb{Z}[\pi_1(M)]$. Our goal is to define a map $\Phi: B \rightarrow P_{q-1}(M')$ such that there is a homotopy $\bar{\nu}' \circ \Phi \simeq p_{q-1}(M')$. Indeed, if we had such a map, then the following diagram would commute and, after showing that $\Phi^*: H^q(P_{q-1}(M'); \Lambda) \rightarrow H^q(B; \Lambda)$ is an isomorphism, the proposition would be proved:

$$(40) \quad \begin{array}{ccccccc} K_q(M'; \Lambda) & \xrightarrow{\iota_{p_{q-1}(M')}} & H_q(M'; \Lambda) & \xleftarrow{\text{PD}} & H^q(M'; \Lambda) & \xleftarrow{p_{q-1}(M')^*} & H^q(P_{q-1}(M'); \Lambda) \\ \cong \downarrow & & \cong \downarrow & & \downarrow = & & \cong \downarrow \Phi^* \\ K_q(M'; \Lambda) & \xrightarrow{\iota_{\bar{\nu}'}} & H_q(M'; \Lambda) & \xleftarrow{\text{PD}} & H^q(M'; \Lambda) & \xleftarrow{\bar{\nu}'^*} & H^q(B; \Lambda). \end{array}$$

Use $\iota: M \rightarrow W$ and $\iota': M' \rightarrow W$ to denote the inclusions, and let $\bar{\nu}_W$ and $\bar{\nu}' = \bar{\nu}_W \circ \iota'$ be the extensions of $\bar{\nu}$. By construction, the trace cobordism W is obtained by attaching an $(r+1)$ -handle to $M \times [0, 1]$. The extension $\bar{\nu}_W$ is null-homotopic on the $(r+1)$ -handle and coincides with $\bar{\nu}$ on M . Composing $\bar{\nu}_W$ and $\bar{\nu}'$ with the projection $\text{pr}_2: P_{q-1}(M) \times BO\langle m \rangle \rightarrow BO\langle m \rangle$ to the second coordinate gives maps $W \rightarrow BO\langle m \rangle$ and $M' \rightarrow BO\langle m \rangle$ that fit into a commutative diagram:

$$(41) \quad \begin{array}{ccccc} M & \xrightarrow{\iota} & W & \xleftarrow{\iota'} & M' \\ \downarrow \bar{\nu} & \swarrow \bar{\nu}_W & \downarrow \text{proj}_1 \circ \bar{\nu}_W & & \downarrow \text{proj}_1 \circ \bar{\nu}' \\ P_{q-1}(M) \times BO\langle m \rangle & & BO\langle m \rangle & \xleftarrow{=} & BO\langle m \rangle \\ \downarrow \text{pr}_1 & & \downarrow \text{pr}_1 & & \downarrow \text{pr}_1 \\ BO\langle m \rangle & \xrightarrow{=} & BO\langle m \rangle & \xleftarrow{=} & BO\langle m \rangle. \end{array}$$

Next, composing $\bar{\nu}_W$ and $\bar{\nu}'$ with the projection $\text{pr}_1: P_{q-1}(M) \times BO\langle m \rangle \rightarrow P_{q-1}(M)$ to the first coordinate gives maps $W \rightarrow P_{q-1}(M)$ and $M' \rightarrow P_{q-1}(M)$ which fit into the following diagram:

$$(42) \quad \begin{array}{ccccc} M & \xrightarrow{\iota} & W & \xleftarrow{\iota'} & M' \\ \downarrow \bar{\nu} & \swarrow \bar{\nu}_W & \downarrow p_{q-1}(W) & & \downarrow p_{q-1}(M') \\ P_{q-1}(M) \times BO\langle m \rangle & & P_{q-1}(W) & \xleftarrow{p_{q-1}(\iota'), \simeq} & P_{q-1}(M') \\ \downarrow \text{pr}_1 & & \downarrow \text{pr}_1 & & \downarrow \text{pr}_1 \\ P_{q-1}(M) & \xrightarrow{p_{q-1}(\iota), \simeq} & P_{q-1}(W) & \xleftarrow{p_{q-1}(\iota'), \simeq} & P_{q-1}(M'). \end{array}$$

The two triangles and the right square commutes. The bottom maps are homotopy equivalences because M' is obtained from M by trivial $(q-1)$ -surgeries or by q -surgeries, both of which do not affect the first $q-1$ homotopy groups. We claim that the left square commutes

up to homotopy. As recalled above, $\bar{\nu}_W$ restricts to $\bar{\nu}$ on M and is a null-homotopic map on the handle. As a consequence, $\text{pr}_1 \circ \bar{\nu}_W$ restricts to $p_{q-1}(M)$ on M and is a null-homotopic map on the handle. It follows that $p_{q-1}(\iota) \circ \text{pr}_1 \circ \bar{\nu}_W$ restricts to $p_{q-1}(\iota) \circ p_{q-1}(M) = p_{q-1}(W) \circ \iota$ on M and is a null-homotopic map on the handle. Up to homotopy, this is precisely the effect of $p_{q-1}(W)$. The claim is established.

Via the diagrams in (41) and (42), we obtain the following homotopy-commutative diagram:

$$\begin{array}{ccccc}
M & \xrightarrow{\iota} & W & \xleftarrow{\iota'} & M' \\
\downarrow \bar{\nu} & \nearrow \bar{\nu}_W & \downarrow \left(\begin{array}{c} p_{q-1}(W) \\ \text{proj}_1 \circ \bar{\nu}_W \end{array} \right) & & \downarrow \left(\begin{array}{c} p_{q-1}(M') \\ \text{proj}_1 \circ \bar{\nu}' \end{array} \right) \\
B & \xrightarrow{p_{q-1}(\iota) \times \text{Id}, \simeq} & P_{q-1}(W) \times BO\langle m \rangle & \xleftarrow{p_{q-1}(\iota') \times \text{Id}} & P_{q-1}(M') \times BO\langle m \rangle \xrightarrow{\text{proj}_1} P_{q-1}(M'). \\
& \searrow g & & \swarrow g' & \\
& & & &
\end{array}$$

Here, the maps g and g' respectively denote the homotopy inverses of the homotopy equivalences $p_{q-1}(\iota) \times \text{Id}$ and $p_{q-1}(\iota') \times \text{Id}$. Note that up to homotopy, $\bar{\nu}' = \bar{\nu}_W \circ \iota'$ can be expressed as

$$(43) \quad \bar{\nu}' = \bar{\nu}_W \circ \iota' \simeq g \circ \left(\begin{array}{c} p_{q-1}(W) \\ \text{proj}_1 \circ \bar{\nu}_W \end{array} \right) \circ \iota' = g \circ (p_{q-1}(\iota') \times \text{Id}) \circ \left(\begin{array}{c} p_{q-1}(M') \\ \text{proj}_1 \circ \bar{\nu}' \end{array} \right).$$

Finally, we set

$$\Phi := \text{proj}_1 \circ g' \circ (p_{q-1}(\iota) \times \text{Id}): B \rightarrow P_{q-1}(M').$$

Using (43), one verifies that $\bar{\nu}' \circ \Phi \simeq p_{q-1}(M')$. Thanks to our identification of fundamental groups, we may assume that Φ induces a map on twisted homology. Taking a look at the definition of Φ and using the same reasoning as in the proof of Proposition 8.9, one then notes that $\Phi^*: H^q(P_{q-1}(M); \Lambda) \rightarrow H^q(B; \Lambda)$ is an isomorphism. This concludes the explanation of the diagram in (40) and therefore concludes the proof of the proposition. \square

Finally, we prove that the two-sided primitive embeddings associated to the normal smoothings from Section 8.1 are invariant under orientation-preserving homotopy equivalences.

Corollary 8.12. *Let M be a closed, connected, oriented $2q$ -manifold such that $p_{q-1}(M)$ is even and split-free. Assume that M admits a map $\nu_m(M): M \rightarrow BO\langle m \rangle$ with $m \geq q$, which lifts the stable normal bundle of M . Assume further that the normal $(q-1)$ -smoothing*

$$\bar{\nu} := p_{q-1}(M) \times \nu_m(M): M \rightarrow P_{q-1}(M) \times BO\langle m \rangle =: B$$

is split-free. Assume that (W, M, M') is a bordism over B obtained by performing a sequence of trivial $(q-1)$ -surgeries followed by a sequence of compatible q -surgeries. If M and M' are orientation-preserving homotopy equivalent, then $PE(M, \bar{\nu})$ and $PE(M', \bar{\nu}')$ agree in $\text{bPrim}^{\pi_1(M)} / \text{Aut}(\pi_1(M))$.

Proof. Since M and M' are orientation-preserving homotopy equivalent, Proposition 8.9 ensures that $PPE(M)$ and $PPE(M')$ agree in $\text{bPrim}^{\pi_1(M)} / \text{Aut}(\pi_1(M))$. But now by Proposition 8.11, we know that $PPE(M)$ and $PE(M, \bar{\nu})$ agree in $\text{bPrim}^{\pi_1(M)}$, and similarly for $PPE(M')$ and $PE(M', \bar{\nu}')$. This concludes the proof of the corollary. \square

9. MANIFOLDS WITH INFINITE HOMOTOPY STABLE CLASS

In this section we prove Theorem 1.1 from the introduction. For the readers' convenience, we recall the necessary terminology. Firstly, two closed $2q$ -manifolds M_0 and M_1 with the same Euler characteristic are *stably diffeomorphic* if there exists a positive integer g and a diffeomorphism

$$M_0 \# W_g \cong M_1 \# W_g,$$

where $W_g := \#_g(S^q \times S^q)$. Secondly, recall that the *homotopy stable class* of a $2q$ -manifold M_0 is defined to be the set of homotopy classes of manifolds stably diffeomorphic to M_0 :

$$\mathcal{S}_h^{\text{st}}(M_0) = \{M_1^{2q} \mid M_1 \text{ is stably diffeomorphic to } M_0\} / \text{homotopy equivalence.}$$

In this section we show that there are manifolds with infinite homotopy stable class. A manifold M is *stably parallelisable* if its tangent bundle becomes trivial after Whitney sum with a trivial bundle. The dimension of our manifolds will henceforth be denoted $4k$ instead of $2q$, so that $q = 2k$. We also exclude dimension four. This is because we rely on the structure of the cohomology ring of the manifolds we construct, in a way that cannot occur for closed 4-manifolds.

The main result of this section is the following, restating Theorem 1.1.

Theorem 9.1. *Let $k \geq 2$. There exist infinitely many closed, oriented, connected $4k$ -manifolds M_1, M_2, \dots with $\pi_1(M_i) \cong \mathbb{Z}$ that are stably parallelisable, have hyperbolic equivariant intersection form, identical Euler characteristic, are pairwise not stably diffeomorphic, and moreover each of the M_i has infinite homotopy stable class:*

$$|\mathcal{S}_h^{\text{st}}(M_i)| = \infty.$$

In fact, for each i , there are infinitely many $4k$ -manifolds M_i^1, M_i^2, \dots with $\pi_1(M_i^k) \cong \mathbb{Z}$ that are stably parallelisable, have hyperbolic equivariant intersection form, and are all stably diffeomorphic to M_i , but as j varies the M_i^j are pairwise not homotopy equivalent.

We briefly describe the strategy of the proof of the theorem. In Proposition 9.3 we recall the simply-connected $4k$ -manifolds $N_{a,b}$ that we constructed in [CCPS21], for a pair of positive, coprime integers $\{a, b\}$. The cohomology rings depend on a and b . We define

$$M_{a,b} := N_{a,b} \# S^1 \times S^{4k-1}.$$

These manifolds satisfy $\pi_1(M_{a,b}) \cong \mathbb{Z}$, and will serve as base manifolds. We apply Corollary 7.7 to realise the infinitely many boundary automorphisms $\{f_i\}$ that we constructed in Section 5 by bordisms W_{f_i} . The $4k$ -manifolds $M_{a,b}^{f_i}$ arising as the new boundary components of these bordisms $\partial_+ W_{f_i}$ will be the infinite family of pairwise stably diffeomorphic manifolds promised by the theorem. We apply Corollary 8.12 to show that they are pairwise not homotopy equivalent.

In order to rule out orientation-reversing homotopy equivalences, we shall appeal to the following observation, which also appeared in [CCPS21]. We recall it here for the convenience of the reader.

Lemma 9.2. *Let N and N' be closed, connected, oriented $4k$ -manifolds. Suppose that a class z freely generates $H^2(N; \mathbb{Z})$ and $z^{2k} = n$ for some nonzero $n \in \mathbb{Z} = H^{4k}(N; \mathbb{Z})$, and*

similarly for (N', z') then any homotopy equivalence $f: N \rightarrow N'$ must be orientation preserving. Moreover, if N'' is stably diffeomorphic to such an N and $k \geq 2$, then $H^2(N''; \mathbb{Z})$ is freely generated by a class z'' with $(z'')^{2k} = n$ and so any homotopy equivalence $N'' \rightarrow N'$ is orientation preserving.

In order to apply this lemma, we shall orient our manifolds N , N' and N'' in such a way that e.g. $z^{2k} \cap [N] \in H_0(N; \mathbb{Z}) = \mathbb{Z}$ is positive, and similarly for N' and N'' . If we do this then we will be able to deduce that any homotopy equivalence between two of them is orientation preserving.

Proof of Lemma 9.2. Assume that f is of degree $\varepsilon = \pm 1$. Since f is a homotopy equivalence, N and N' have isomorphic cohomology rings. In particular $H^2(N'; \mathbb{Z}) \cong \mathbb{Z}$ is generated by $z' = \pm f^{-*}(z)$. Since $z'^{2k} = n$ in $H^{4k}(N'; \mathbb{Z}) \cong \mathbb{Z}$, and $f^*(z'^{2k}) = z^{2k}$, properties of the cap and cup products show that

$$n = f_*(z^{2k} \cap [N]) = f_*(f^*(z'^{2k}) \cap [N]) = z'^{2k} \cap f_*([N]) = z'^{2k} \cap \varepsilon[N'] = \varepsilon n.$$

Since $n \neq 0$, this implies that f must be orientation-preserving.

To prove the last assertion, use the naturality of the cup product together with the degree one map $N \# S^{2k} \times S^{2k} \rightarrow N$ that collapses $S^{2k} \times S^{2k}$ to a point, and induces isomorphisms on the second and $4k$ -th cohomology. This shows that N has an element $z \in H^2(N; \mathbb{Z})$ as in the statement of the lemma, if and only if $H^2(N \# S^{2k} \times S^{2k}; \mathbb{Z})$ does, and similarly for N'' . The last sentence of the lemma then follows from the argument of the previous paragraph. \square

9.1. The simply-connected $4k$ -manifolds $N_{a,b}$. Let us recall the next proposition from [CCPS21], which builds the simply-connected $4k$ -manifolds $N_{a,b}$. We will then proceed to discuss some of their properties.

Proposition 9.3. *Fix $k \geq 2$. Given an unordered pair $\{a, b\}$ of positive coprime integers such that $(2k)!$ divides $2ab$, there exists a closed, connected, oriented, smooth $4k$ -manifolds $N_{a,b}^{4k}$ with the following properties.*

- (i) *The manifold $N_{a,b}$ is simply-connected and stably parallelisable.*
- (ii) *The ring $H^*(N_{a,b}; \mathbb{Z})$ has generators w, x, y, z and 1 of degrees $2k+2, 2k, 2k, 2$ and 0 respectively, with $z^k = ax + by, x^2 = 0 = y^2, 2abw = z^{k+1}, xz = bw, yz = aw$ and xy generates $H^{4k}(N_{a,b}; \mathbb{Z})$.*
- (iii) *There is a $2k$ -connected map $\beta: N_{a,b} \rightarrow \mathbb{C}P^\infty$*

In particular, the intersection form of $N_{a,b}$ is hyperbolic and $z^{2k} = 2abxy$ is $2ab$ times a fundamental class of $N_{a,b}$. If $\{a, b\} \neq \{a', b'\}$ then $N_{a,b}$ and $N_{a',b'}$ have non-isomorphic integral cohomology rings and so are not homotopy equivalent. Moreover if $ab \neq a'b'$, then $N_{a,b}$ and $N_{a',b'}$ are not even stably diffeomorphic.

As mentioned at the start of the section, we will use $N_{a,b}$ to construct our base manifolds $M_{a,b} := N_{a,b} \# S^1 \times S^{4k-1}$. We will prove some facts purely about $N_{a,b}$, and then later obtain their analogues for $M_{a,b}$.

Next we describe a normal $(2k-1)$ -smoothing for $N_{a,b}$. Since $N_{a,b}$ is stably parallelisable, the stable normal bundle $\nu: N_{a,b} \rightarrow BO$ factors through a map $\nu_{2k}: N_{a,b} \rightarrow BO\langle 4k \rangle$ followed by $\vartheta_{2k}: BO\langle 4k \rangle \rightarrow BO$. This is a normal $(2k-1)$ -smoothing but it is not the normal $(2k-1)$ -type. We also use $p_{2k-1}(N_{a,b}): N_{a,b} \rightarrow P_{2k-1}(N_{a,b})$ to denote the $(2k-1)$ st Postnikov stage of $N_{a,b}$.

Lemma 9.4. *Let a and b be positive, coprime integers such that $(2k)!$ divides ab . The $(2k-1)$ st Postnikov stage $P_{2k-1}(N_{a,b})$ is homotopy equivalent to $\mathbb{C}P^\infty$, and a normal $(2k-1)$ -type of $N_{a,b}$ is given by the fibration*

$$\vartheta_{2k} \circ \text{pr}_2: \mathbb{C}P^\infty \times BO\langle 4k \rangle \rightarrow BO,$$

where pr_2 is the projection onto the second factor. The map

$$\bar{\nu}_{a,b} := p_{2k-1}(N_{a,b}) \times \nu_{2k}: N_{a,b} \rightarrow \mathbb{C}P^\infty \times BO\langle 4k \rangle =: B_{2k-1}(N_{a,b})$$

is a normal $(2k-1)$ -smoothing. Write $(\bar{\nu}: N \rightarrow B_{2k-1}(N)) := (\bar{\nu}_{a,b}: N_{a,b} \rightarrow B_{2k-1}(N_{a,b}))$ to ease notation. It satisfies the following properties.

- (i) The evaluation $\text{ev}_N: H^{2k}(N; \mathbb{Z}) \rightarrow \text{Hom}_{\mathbb{Z}}(H_{2k}(N; \mathbb{Z}), \mathbb{Z})$ is an isomorphism.
- (ii) The following evaluation maps are isomorphisms:

$$\text{ev}_{P_{2k-1}(N)}: H^{2k}(P_{2k-1}(N); \mathbb{Z}) \rightarrow \text{Hom}_{\mathbb{Z}}(H_{2k}(P_{2k-1}(N); \mathbb{Z}), \mathbb{Z}),$$

$$\text{ev}_{B_{2k-1}(N)}: H^{2k}(B_{2k-1}(N); \mathbb{Z}) \rightarrow \text{Hom}_{\mathbb{Z}}(H_{2k}(B_{2k-1}(N); \mathbb{Z}), \mathbb{Z}).$$

- (iii) The abelian groups $K_{2k}(N; \mathbb{Z})$, $\ker(p_{2k-1}(N)_*)$, $H_{2k}(B_{2k-1}(N); \mathbb{Z})$, and $H_{2k}(P_{2k-1}(N); \mathbb{Z})$ are finitely generated and free.

In particular, the normal $(2k-1)$ -smoothing $\bar{\nu}: N \rightarrow B_{2k-1}(N)$ is split-free.

Proof. Since there is a $2k$ -connected map $\beta: N_{a,b} \rightarrow \mathbb{C}P^\infty$ (by item (iii) of Proposition 9.3) and since $\mathbb{C}P^\infty \simeq K(\mathbb{Z}, 2)$, we see that $P_{2k-1}(N_{a,b}) \simeq \mathbb{C}P^\infty$. The first part of the lemma then follows from Lemma 8.1.

Now we check the claimed properties. For (i), since $N \rightarrow \mathbb{C}P^\infty$ is $2k$ -connected, $H_{2k-1}(N; \mathbb{Z}) = 0$ and thus the universal coefficient theorem ensures that ev_N is an isomorphism. For (ii), as $B_{2k-1}(N) \simeq \mathbb{C}P^\infty \times BO\langle 4k \rangle$, we deduce from the Künneth theorem that $H_{2k-1}(B_{2k-1}(N); \mathbb{Z}) = H_{2k-1}(P_{2k-1}(N); \mathbb{Z}) = 0$, and thus we also obtain that $\text{ev}_{B_{2k-1}(N)}$ and $\text{ev}_{P_{2k-1}(N)}$ are isomorphisms. Since $H_{2k}(N; \mathbb{Z})$ is free and \mathbb{Z} is a PID, we deduce that $K_{2k}(N; \mathbb{Z}) = \ker(\bar{\nu}_*)$ and $\ker(p_{2k-1}(N)_*)$ are also free. Also $H_{2k}(B_{2k-1}(N); \mathbb{Z}) \cong H_{2k}(P_{2k-1}(N); \mathbb{Z}) \cong H_{2k}(\mathbb{C}P^\infty; \mathbb{Z}) \cong \mathbb{Z}$, so this is f.g. and free, proving (iii). The second and third assertions imply, by definition, that $p_{2k-1}(N)$ and the normal $(2k-1)$ -smoothing $\bar{\nu}$ are split-free. \square

As in the statement of Lemma 9.4, when the integers a, b are fixed or clear from the context, we write N , $B_{2k-1}(N)$ and $\bar{\nu}$ instead of $N_{a,b}$, $B_{2k-1}(N_{a,b})$ and $\bar{\nu}_{a,b}$. By Proposition 9.3, the intersection form on $H_{2k}(N; \mathbb{Z})$ is hyperbolic and, in particular nonsingular. Since $4k = 2q$ implies q is even, and the intersection form on N is hyperbolic, it admits a unique quadratic refinement, so the normal $(2k-1)$ -smoothing $\bar{\nu}: N \rightarrow B_{2k-1}(N)$ is even. We showed in Lemma 9.4 that $\bar{\nu}: N \rightarrow B_{2k-1}(N)$ is a split-free normal $(2k-1)$ -smoothing. Therefore the 2-sided primitive embedding $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ associated with the normal $(2k-1)$ -smoothing $\bar{\nu}$ is defined. The next lemma describes it in detail.

Proposition 9.5. *Given positive, coprime integers a and b such that $(2k)!$ divides ab , the 2-sided primitive embedding $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ associated to $(\bar{\nu}: N \rightarrow B_{2k-1}(N)) := (\bar{\nu}_{a,b}: N_{a,b} \rightarrow B_{2k-1}(N_{a,b}))$ is isomorphic to*

$$\left((\mathbb{Z}, -ab) \xrightarrow{\begin{pmatrix} -a \\ b \end{pmatrix}} H_+(\mathbb{Z}) \xleftarrow{\begin{pmatrix} a \\ b \end{pmatrix}} (\mathbb{Z}, ab) \right).$$

Proof. In order to describe the primitive embedding $j_{\bar{\nu}}: H^{2k}(B_{2k-1}(N); \mathbb{Z}) \rightarrow H_{2k}(N; \mathbb{Z})$, we first choose bases for the \mathbb{Z} -modules $H^{2k}(B_{2k-1}(N); \mathbb{Z}) = \mathbb{Z}$ and $H_{2k}(N; \mathbb{Z}) = \mathbb{Z}^2$. Recall from Proposition 9.3 that $H^2(N; \mathbb{Z}) = \mathbb{Z}\langle z \rangle$ and $H^{2k}(N; \mathbb{Z}) = \mathbb{Z}\langle x \rangle \oplus \mathbb{Z}\langle y \rangle$, where $z^{2k} = ax + by$. Passing to homology, we endow $H_{2k}(N; \mathbb{Z})$ with the basis (\bar{x}, \bar{y}) Poincaré dual to (x, y) . In other words, we set $\bar{x} := \text{PD}(x)$ and $\bar{y} := \text{PD}(y)$. Since $BO\langle 4k \rangle$ is $4k$ -connected, we know that $H^2(B_{2k-1}(N); \mathbb{Z}) = H^2(\mathbb{C}P^\infty; \mathbb{Z}) = \mathbb{Z}$. Since $\bar{\nu}^*$ is $2k$ -connected, it induces an isomorphism between the second cohomology groups. We set $v := \bar{\nu}^{-*}(z)$, so that $H^2(B_{2k-1}(N); \mathbb{Z}) = \mathbb{Z}\langle v \rangle$. It follows that $H^{2k}(B_{2k-1}(N); \mathbb{Z}) = \mathbb{Z}\langle v^k \rangle$ and therefore the primitive embedding $j_{\bar{\nu}}$ is determined by its value on v^k . Since, by definition, we have $j_{\bar{\nu}} = \text{PD} \circ \bar{\nu}^*$, we deduce that

$$j_{\bar{\nu}}(v^k) = \text{PD} \circ \bar{\nu}^*(v^k) = \text{PD}(z^k) = \text{PD}(ax + by) = a\bar{x} + b\bar{y}.$$

In other words, with respect to the bases v^k and (\bar{x}, \bar{y}) , the map $j_{\bar{\nu}}$ is represented by $\begin{pmatrix} a \\ b \end{pmatrix}$. The quadratic form $j_{\bar{\nu}}^* \theta_N j_{\bar{\nu}}$ on $H^{2k}(B_{2k-1}(N); \mathbb{Z})$ is obtained by pulling back the (hyperbolic) form θ_N on $H_{2k}(N; \mathbb{Z})$. The description of $j_{\bar{\nu}}$ is now concluded by the following computation:

$$j_{\bar{\nu}}^* \theta_N j_{\bar{\nu}}(v^k, v^k) = \theta_N(j_{\bar{\nu}}(v^k), j_{\bar{\nu}}(v^k)) = \begin{pmatrix} a & b \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = ab.$$

In order to describe $\iota_{\bar{\nu}}: K_{2k}(N; \mathbb{Z}) \hookrightarrow H_{2k}(N; \mathbb{Z})$, we first determine the isometry type of the quadratic form $(K_{2k}(N; \mathbb{Z}), \theta_N)$. To start with, we base $H_{2k}(B_{2k-1}(N); \mathbb{Z}) = H_{2k}(\mathbb{C}P^\infty; \mathbb{Z}) = \mathbb{Z}$. Using Lemma 9.4 (ii), we know that

$$\mathbb{Z}\langle v^k \rangle = H^{2k}(B_{2k-1}(N); \mathbb{Z}) \cong \text{Hom}_{\mathbb{Z}}(H_{2k}(B_{2k-1}(N); \mathbb{Z}), \mathbb{Z}).$$

We base $\text{Hom}_{\mathbb{Z}}(H_{2k}(B_{2k-1}(N); \mathbb{Z}), \mathbb{Z}) \cong \mathbb{Z}$ with the image of v^k under this isomorphism and base the group $H_{2k}(B_{2k-1}(N); \mathbb{Z}) \cong \mathbb{Z}$ with the dual basis v_*^k . Using Lemma 9.4 (i), we can apply the same procedure to N , yielding basis elements $x_*, y_* \in H_{2k}(N; \mathbb{Z}) = \mathbb{Z}^2$ that are hom-dual to x, y . Observe that $x_* = \bar{y}$ and $y_* = \bar{x}$. With respect to the bases (x_*, y_*) and v_*^k , the map $\bar{\nu}_*$ is given by $\begin{pmatrix} a \\ b \end{pmatrix}^T = \begin{pmatrix} a & b \end{pmatrix}$. Passing to the basis (\bar{x}, \bar{y}) , we see that $\bar{\nu}_*$ is represented by $\begin{pmatrix} b & a \end{pmatrix}$. Taking the kernel of this map, we deduce that

$$K_{2k}(N; \mathbb{Z}) = \ker(\bar{\nu}_*) = \mathbb{Z}\langle \begin{pmatrix} -a \\ b \end{pmatrix} \rangle.$$

Pulling back the hyperbolic form from $H_{2k}(N; \mathbb{Z})$, we infer that the quadratic form on $K_{2k}(N; \mathbb{Z})$ is represented by $(-ab)$. This implies that $(K_{2k}(N; \mathbb{Z}), \theta_N) = (\mathbb{Z}, -ab)$, and the computation of $\iota_{\bar{\nu}}$ now follows readily. This concludes the proof of the proposition. \square

9.2. The $4k$ -manifolds $M_{a,b}$ and their associated primitive embeddings. Given two positive, coprime integers a and b such that $(2k)!$ divides ab , as above let $N_{a,b}$ be the simply-connected $4k$ -manifold built in the proof of Proposition 9.3 and studied in the previous subsection, and let

$$(44) \quad M_{a,b} := N_{a,b} \# (S^1 \times S^{4k-1}).$$

We need the analogous results for $M_{a,b}$ to those that we proved in Lemma 9.4 and Proposition 9.5 for $N_{a,b}$. We will use these two results to deduce their analogues.

Since both $N_{a,b}$ and $S^1 \times S^{4k-1}$ are stably parallelisable, their stable normal bundles lift to $BO\langle 4k \rangle$. The same assertion therefore holds for $M_{a,b}$. In particular, its stable normal bundle lifts to a map $\nu_{2k}: M_{a,b} \rightarrow BO\langle 4k \rangle$. Using $p_{2k-1}(M_{a,b}): M_{a,b} \rightarrow P_{2k-1}(M_{a,b})$ to

denote the $(2k-1)$ st Postnikov stage of $M_{a,b}$, Lemma 9.4 and Lemma 8.1 yields the following result.

Lemma 9.6. *Let a and b be positive, coprime integers such that $(2k)!$ divides ab . A normal $(2k-1)$ -type for the stably parallelisable $4k$ -manifold $M_{a,b}$ is given by the fibration*

$$\vartheta_{2k} \circ \text{pr}_2: P_{2k-1}(M_{a,b}) \times BO\langle 4k \rangle \rightarrow BO,$$

where pr_2 is the projection onto the second factor. The following map is a normal $(2k-1)$ -smoothing:

$$\bar{\nu}_{a,b} := p_{2k-1}(M_{a,b}) \times \nu_{2k}: M_{a,b} \rightarrow P_{2k-1}(M_{a,b}) \times BO\langle 4k \rangle =: B_{2k-1}(M_{a,b}).$$

When the integers a, b are clear from the context, as before we write $M, N, B_{2k-1}(M), B_{2k-1}(N), \bar{\nu}$ instead of $M_{a,b}, N_{a,b}, B_{2k-1}(M_{a,b}), B_{2k-1}(N_{a,b}), \bar{\nu}_{a,b}$. Our aim is to describe the 2-sided primitive embedding $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ associated with the normal $(2k-1)$ -smoothing $\bar{\nu}$. From now on, we set

$$\Lambda := \mathbb{Z}[\pi_1(B_{2k-1}(M))] = \mathbb{Z}[\pi_1(M)]$$

(recall Convention 8.4) and note that the intersection form on the Λ -module $H_{2k}(M; \Lambda)$ is hyperbolic, and therefore admits a quadratic refinement. In particular, the form is nonsingular and the normal $(2k-1)$ -smoothing is even. To ensure that $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ is defined, we must check that $\bar{\nu}: M \rightarrow B_{2k-1}(M)$ is a split-free normal $(2k-1)$ -smoothing.

Lemma 9.7. *Let a and b be positive, coprime integers such that $(2k)!$ divides ab . Writing $M := M_{a,b}$ as well as $N := N_{a,b}$, the normal $(2k-1)$ -smoothing $\bar{\nu}: M \rightarrow B_{2k-1}(M)$ satisfies the following properties.*

(i) *The evaluation $\text{ev}_M: H^{2k}(M; \Lambda) \xrightarrow{\cong} \overline{\text{Hom}}_{\Lambda}(H_{2k}(M; \Lambda), \Lambda)$ is an isomorphism.*

(ii) *The following evaluation maps are isomorphisms*

$$\text{ev}_{P_{2k-1}(M)}: H^{2k}(P_{2k-1}(M); \Lambda) \xrightarrow{\cong} \overline{\text{Hom}}_{\Lambda}(H_{2k}(P_{2k-1}(M); \Lambda), \Lambda),$$

$$\text{ev}_{B_{2k-1}(M)}: H^{2k}(B_{2k-1}(M); \Lambda) \xrightarrow{\cong} \overline{\text{Hom}}_{\Lambda}(H_{2k}(B_{2k-1}(M); \Lambda), \Lambda).$$

(iii) *The inclusion $\mathring{N} \rightarrow M$ induces an isomorphism $H_{2k}(N; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \xrightarrow{\cong} H_{2k}(M; \Lambda)$.*

(iv) *The inclusion $\mathring{N} \rightarrow M$ induces isomorphisms*

$$H_{2k}(P_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \xrightarrow{\cong} H_{2k}(P_{2k-1}(M); \Lambda),$$

$$H_{2k}(B_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \xrightarrow{\cong} H_{2k}(B_{2k-1}(M); \Lambda).$$

In particular, $H_{2k}(P_{2k-1}(M); \Lambda)$ and $H_{2k}(B_{2k-1}(M); \Lambda)$ are finitely generated and free.

(v) *The kernels $K_{2k}(M; \Lambda)$ and $\ker(p_{2k-1}(M))$ are finitely generated free Λ -modules.*

In particular, $p_{2k-1}(M)$ and the normal $(2k-1)$ -smoothing $\bar{\nu}: M \rightarrow B_{2k-1}(M)$ are split-free.

Proof. The first three assertions are proved using the universal coefficient spectral sequence, or UCSS for short [Lev77, Theorem 2.3]. For a space Y , this spectral sequence is given by

$$E_2^{p,q} = \overline{\text{Ext}}_{\Lambda}^q(H_p(Y; \Lambda), \Lambda) \Rightarrow H^{p+q}(Y; \Lambda).$$

To describe the second page of the UCSS for $Y = M$, we compute $H_i(M; \Lambda)$ for $0 < i \leq 2k$. As $M = N \# (S^1 \times S^{4k-1})$, the universal cover is homeomorphic to an infinite connected

sum of copies of N , so we obtain $H_i(M; \Lambda) \cong H_i(N; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda$ for $0 < i \leq 2k$. We deduce the Λ -homology of M , establishing in particular the third assertion (iii):

$$(45) \quad H_i(M; \Lambda) = \begin{cases} \mathbb{Z} & \text{if } i = 0, \\ 0 & \text{if } 0 < i \leq 2k, i \text{ odd}, \\ H_i(N; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda & \text{if } 0 < i \leq 2k, i \text{ even}. \end{cases}$$

We infer that the odd columns of the second page of the UCSS vanish. Then since the homology of N is f.g. free over \mathbb{Z} , the homology groups $H_i(M; \Lambda)$ for $0 < i \leq 2k$ and i even are also f.g. free over Λ . It follows that for $p \geq 1$ and $0 < i \leq 2k$ we have

$$\overline{\text{Ext}}_{\Lambda}^p(H_i(M; \Lambda), \Lambda) = 0.$$

A spectral sequence computation then shows that $\text{ev}_N: H^{2k}(M; \Lambda) \rightarrow \overline{\text{Hom}}_{\Lambda}(H_{2k}(M; \Lambda), \Lambda)$ is an isomorphism. This establishes (i).

We prove the second assertion (ii), namely the analogous result for $Y = B_{2k-1}(M)$ and $Y = P_{2k-1}(M)$. It suffices to show that for $i \leq 2k - 1$, we have $H_i(Y; \Lambda) = H_i(M; \Lambda)$. The computation of $H^{2k}(Y; \Lambda)$ then follows the exact same steps as for $Y = M$. Using Lemma 9.6 we have $B_{2k-1}(M) \simeq P_{2k-1}(M) \times BO\langle 4k \rangle$. Since $BO\langle 4k \rangle$ is $4k$ -connected and $P_{2k-1}(M)$ is obtained from M by adding cells of dimension $2k+1$ or greater, we obtain $H_i(B_{2k-1}(M); \Lambda) = H_i(P_{2k-1}(M); \Lambda) = H_i(M; \Lambda)$ for $i \leq 2k - 1$, as desired. This establishes (ii).

We move on to the fourth assertion (iv). First note that

$$\begin{aligned} H_{2k}(B_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda &\cong H_{2k}(P_{2k-1}(N) \times BO\langle 4k \rangle; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \cong H_{2k}(P_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \\ &\cong H_{2k}(\mathbb{C}P^{\infty}; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \cong \Lambda \end{aligned}$$

by the Künneth theorem and since $BO\langle 4k \rangle$ is $4k$ -connected. Similarly, we claim that

$$H_{2k}(B_{2k-1}(M); \Lambda) \cong H_{2k}(P_{2k-1}(M) \times BO\langle 4k \rangle; \Lambda) \cong H_{2k}(P_{2k-1}(M); \Lambda).$$

To see this, first note that the fibration $BO\langle 4k \rangle \rightarrow BO\langle 4k \rangle \times P_{2k-1}(M) \rightarrow P_{2k-1}(M)$ implies that $\pi_i(BO\langle 4k \rangle \times P_{2k-1}(M)) = \pi_i(P_{2k-1}(M))$ for $i \leq 4k$. Applying the relative Hurewicz theorem to the universal covers of these spaces then implies the claim.

The proof of (iv) therefore reduces to showing that the inclusion induced map $P_{2k-1}(N) = P_{2k-1}(\mathring{N}) \rightarrow P_{2k-1}(M)$ gives rise to an isomorphism

$$H_{2k}(\mathbb{C}P^{\infty}; \Lambda) \xrightarrow{\cong} H_{2k}(P_{2k-1}(M); \Lambda).$$

For later use, note that $H_{2k}(\mathbb{C}P^{\infty}; \Lambda) = \Lambda$ is generated by the $2k$ -cells of $\mathbb{C}P^{\infty}$.

Since $P_{2k-1}(M)$ only depends on the homotopy type of the $2k$ -skeleton of $M = N \# S^1 \times S^{4k-1}$, which is homotopy equivalent to $N^{(2k)} \vee S^1$, we obtain

$$(46) \quad \begin{aligned} P_{2k-1}(M) &\simeq P_{2k-1}(N \# S^1 \times S^{4k-1}) \simeq P_{2k-1}((N \# S^1 \times S^{4k-1})^{(2k)}) \\ &\simeq P_{2k-1}(N^{(2k)} \vee S^1) \simeq P_{2k-1}(\mathbb{C}P^{\infty} \vee S^1). \end{aligned}$$

Here we use that since $N \rightarrow \mathbb{C}P^{\infty}$ is $2k$ -connected, so is the composition $N^{(2k)} \rightarrow N \rightarrow \mathbb{C}P^{\infty}$, from which it follows that $N^{(2k)} \vee S^1 \rightarrow \mathbb{C}P^{\infty} \vee S^1$ is $2k$ -connected and therefore we have $P_{2k-1}(N^{(2k)} \vee S^1) \simeq P_{2k-1}(\mathbb{C}P^{\infty} \vee S^1)$ as asserted for the last equivalence of (46).

In the Claim below, we prove that $H_{2k}(P_{2k-1}(\mathbb{C}P^{\infty} \vee S^1); \Lambda) = \Lambda = H_{2k}(\mathbb{C}P^{\infty}; \Lambda)$, generated by the $2k$ -cells of $\mathbb{C}P^{\infty}$. Since this is also the case for $P_{2k-1}(N)$, combining the

Claim below with the equalities from (46) will prove (iv), that the inclusion $\mathring{N} \rightarrow M$ induces isomorphisms

$$\begin{aligned} H_{2k}(P_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda &\xrightarrow{\cong} H_{2k}(P_{2k-1}(M); \Lambda), \\ H_{2k}(B_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda &\xrightarrow{\cong} H_{2k}(B_{2k-1}(M); \Lambda). \end{aligned}$$

Claim. $H_{2k}(P_{2k-1}(\mathbb{C}P^\infty \vee S^1); \Lambda) \cong \Lambda$, generated by the $2k$ -cells of $\mathbb{C}P^\infty$.

Proof. Let $X := \mathbb{C}P^\infty \vee S^1$ and consider the map $p_{2k-1}: X \rightarrow P_{2k-1}(X)$. By definition, p_{2k-1} is a $2k$ -connected and hence

$$p_{2k-1*}: H_{2k}(X; \Lambda) \rightarrow H_{2k}(P_{2k-1}(X); \Lambda)$$

is surjective. The universal cover \tilde{X} of X is homeomorphic to \mathbb{R} with a copy of $\mathbb{C}P^\infty$ wedged on at each integer, and so

$$(47) \quad H_{2k}(X; \Lambda) = H_{2k}(\tilde{X}; \mathbb{Z}) \cong H_{2k}(X; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda \cong \Lambda,$$

generated by the $2k$ cell of $\mathbb{C}P^\infty$. Therefore to prove the claim it suffices to show that p_{2k-1*} is injective, which we now do.

It follows from $H_{2k}(X; \Lambda) \cong H_{2k}(X; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda$ that for every nonzero $u \in H_{2k}(X; \Lambda)$ there is a class $\alpha \in H^{2k}(X; \Lambda)$ such that $\langle \alpha, u \rangle \neq 0$. For every $\ell > 0$ we have

$$\phi: H^{2\ell}(X; \Lambda) \cong H^{2\ell}(\mathbb{C}P^\infty; \Lambda) \cong H^{2\ell}(\mathbb{C}P^\infty; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda,$$

which is isomorphic to Λ and generated by $(\phi^{-1}(z \otimes 1))^\ell$ for $z \in H^2(\mathbb{C}P^\infty; \mathbb{Z})$ a generator. Here, to make sense of this cup product, we think of $H^{2\ell}(X; \Lambda)$ as the compactly support cohomology group $H_c^{2\ell}(\tilde{X}; \mathbb{Z})$. Since p_{2k-1} is a $2k$ -connected map, we have that $p_{2k-1}^*: H^2(P_{2k-1}(X); \Lambda) \rightarrow H^2(X; \Lambda)$ is an isomorphism. We may therefore invert p_{2k-1}^* to define the element

$$d := p_{2k-1}^{-*}(\phi^{-1}(z \otimes 1)) \in H^2(P_{2k-1}(X); \Lambda).$$

We consider its k th power $d^k \in H^{2k}(P_{2k-1}(X); \Lambda)$. Note that

$$p_{2k-1}^*(d^k) = (p_{2k-1}^*(d))^k = (\phi^{-1}(z \otimes 1))^k = \phi^{-1}(z^k \otimes 1).$$

This shows that d^k maps under p_{2k-1}^* to the generator of $H^{2k}(X; \Lambda)$. The map p_{2k-1}^* is therefore surjective and hence $\alpha = p_{2k-1}^*(\beta) \in H^{2k}(X; \Lambda)$ for some $\beta \in H^{2k}(P_{2k-1}(X); \Lambda)$. It follows that

$$\langle \beta, p_{2k-1*}(u) \rangle = \langle p_{2k-1}^*(\beta), u \rangle = \langle \alpha, u \rangle \neq 0,$$

so in particular $p_{2k-1*}(u) \neq 0$. Thus $p_{2k-1*}: H_{2k}(X; \Lambda) \rightarrow H_{2k}(P_{2k-1}(X); \Lambda)$ is injective, and so is an isomorphism. This proves the Claim that $H_{2k}(P_{2k+1}(X); \Lambda) = H_{2k}(P_{2k-1}(\mathbb{C}P^\infty \vee S^1); \Lambda) \cong \Lambda$, generated by the $2k$ -cells of $\mathbb{C}P^\infty$. As explained above this completes the proof of (iv) of Lemma 9.7. \square

Having completed the proof of the first four assertions of Proposition 9.7, we prove (v). Since \bar{v} is $2k$ -connected, the induced map $\bar{v}_*: H_{2k}(M; \Lambda) \rightarrow H_{2k}(B_{2k-1}(M); \Lambda)$ is surjective. The fourth assertion implies that $H_{2k}(B_{2k-1}(M); \Lambda)$ is free. We deduce that the following short exact sequence splits:

$$0 \rightarrow K_{2k}(M; \Lambda) \rightarrow H_{2k}(M; \Lambda) \xrightarrow{\bar{v}_*} H_{2k}(B_{2k-1}(M); \Lambda) \rightarrow 0.$$

The third item ensures that $H_{2k}(M; \Lambda)$ is free. We therefore deduce that $K_{2k}(M; \Lambda)$ is finitely generated and projective. Since finitely generated, projective Λ -modules are free [Lam06, Chapter V, Corollary 4.12], $K_{2k}(M; \Lambda) = \ker(\bar{\nu}_*)$ is finitely generated and free as desired. The proof for $\ker(p_{2k-1}(M)_*)$ is identical. This concludes the proof (v) and therefore of Lemma 9.7. \square

Since Lemma 9.7 implies that $\bar{\nu}: M \rightarrow B_{2k-1}(M)$ is an even, split-free normal $(2k-1)$ -smoothing, the 2-sided primitive embedding $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ is defined. The next lemma describes it in more detail.

Proposition 9.8. *Given positive, coprime integers a and b such that $(2k)!$ divides ab , the 2-sided primitive embedding $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ associated to $(\bar{\nu}: M \rightarrow B_{2k-1}(M)) := (\bar{\nu}_{a,b}: M_{a,b} \rightarrow B_{2k-1}(M_{a,b}))$ is isomorphic to*

$$\left((\Lambda, -ab) \xrightarrow{\begin{pmatrix} -a \\ b \end{pmatrix}} H_+(\Lambda) \xleftarrow{\begin{pmatrix} a \\ b \end{pmatrix}} (\Lambda, ab) \right).$$

Proof. In Proposition 9.5, an analogous statement was proven for the simply-connected $N := N_{a,b}$ by performing an explicit computation using the bases $H^{2k}(N; \mathbb{Z}) = \mathbb{Z}^2 \langle x, y \rangle$ and $H_{2k}(N; \mathbb{Z}) = \mathbb{Z}^2 \langle \bar{x}, \bar{y} \rangle$ and $H^{2k}(B_{2k-1}(N); \mathbb{Z}) = \mathbb{Z} \langle v^2 \rangle$, where $v := \bar{\nu}^{-1}(z)$ with z the generator of $H^2(N; \mathbb{Z}) = \mathbb{Z}$. These bases led to the computation of $j_{\bar{\nu}}$, while taking the hom-dual bases led to the computation of $\iota_{\bar{\nu}}$. Our aim is to show that these statements can be “tensor over Λ ”.

We start with the computation of $j_{\bar{\nu}}$. Applying the third and fourth assertions of Lemma 9.7, we know that $H_{2k}(M; \Lambda) = H_{2k}(N; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda$ and $H_{2k}(B_{2k-1}(M); \Lambda) = H_{2k}(B_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda$. Additionally applying the first and second isomorphisms of Lemma 9.7 leads to the identifications

$$\begin{aligned} H^{2k}(B_{2k-1}(M); \Lambda) &\cong \overline{\text{Hom}_{\Lambda}(H_{2k}(B_{2k-1}(M); \Lambda), \Lambda)} = \overline{\text{Hom}_{\Lambda}(H_{2k}(B_{2k-1}(N); \mathbb{Z}), \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda} \\ &= H^{2k}(B_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda; \\ H^{2k}(M; \Lambda) &\cong \overline{\text{Hom}_{\Lambda}(H_{2k}(M; \Lambda), \Lambda)} = \overline{\text{Hom}_{\Lambda}(H_{2k}(N; \mathbb{Z}), \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda} \\ &= H^{2k}(N; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda. \end{aligned}$$

Base the Λ -modules $H^{2k}(M; \Lambda) = \Lambda^2$ and $H_{2k}(M; \Lambda) = \Lambda^2$ as well as $H^{2k}(B_{2k-1}(M); \Lambda) = \Lambda$ with the tensored up bases $(x \otimes 1, y \otimes 1)$, $(\bar{x} \otimes 1, \bar{y} \otimes 1)$ and $v^k \otimes 1$ respectively, so that $z^k \otimes 1 = a(x \otimes 1) + b(y \otimes 1)$. The computation of $j_{\bar{\nu}}$ now proceeds as in the simply-connected case:

$$j_{\bar{\nu}}(v^k \otimes 1) = \text{PD} \circ \bar{\nu}^*(v^k \otimes 1) = \text{PD}(z^k \otimes 1) = \text{PD}(ax \otimes 1 + by \otimes 1) = a\bar{x} \otimes 1 + b\bar{y} \otimes 1.$$

Note that in the second equality, we use that $\bar{\nu}(M)^* = \bar{\nu}(N)^* \otimes 1$: we have $H_{2k}(M; \Lambda) = H_{2k}(N; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda$ as well as $H_{2k}(B_{2k-1}(M); \Lambda) = H_{2k}(B_{2k-1}(N); \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda = H_{2k}(\mathbb{C}P^{\infty}; \mathbb{Z}) \otimes_{\mathbb{Z}} \Lambda$, and in both cases, the normal smoothings come from the Postnikov-tower induced map.

The simply-connected procedure can also be applied to compute $\iota_{\bar{\nu}}$. Indeed, we can use Lemma 9.7 to base $H_{2k}(B_{2k-1}(M); \Lambda)$ and $H_{2k}(M; \Lambda)$ with the bases $v_*^k \otimes 1$ and $(x_* \otimes 1, y_* \otimes 1)$ that are hom-dual to the aforementioned ones on the cohomology groups. With respect to these bases, $\bar{\nu}_*$ is represented by $\begin{pmatrix} b \\ a \end{pmatrix}$. Taking the kernel, we deduce that $K_{2k}(M; \Lambda) = \Lambda \langle \begin{pmatrix} -a \\ b \end{pmatrix} \rangle$, and the conclusion follows. \square

Remark 9.9. Let $\bar{\nu}: M \rightarrow B$ be as in Proposition 9.8 and let $\bar{\nu}': M' \rightarrow B$ be another normal $(2k-1)$ -smoothing with the same homology and intersection form as M . For instance, M' could be a manifold obtained by the realisation process of Corollary 7.12. Thinking of $H^*(M'; \Lambda)$ as the compactly supported cohomology group $H_c^*(\widetilde{M}'; \mathbb{Z})$, we argue that the data of $PE(M', \bar{\nu}')$ is determined by the cup product

$$\begin{aligned} \cup^k: H^2(M'; \Lambda) &\rightarrow H^{2k}(M'; \Lambda) \\ x &\mapsto x^k. \end{aligned}$$

To see this, note that item (iii) of Remark 7.6 asserts that the data of $PE(M', \bar{\nu}')$ is determined by the map $\bar{\nu}'^*: H^{2k}(B; \Lambda) \rightarrow H^{2k}(M'; \Lambda)$ and use the naturality of the cup product to obtain the following commutative diagram

$$\begin{array}{ccc} H^{2k}(B; \Lambda) & \xrightarrow{\bar{\nu}'^*} & H^{2k}(M'; \Lambda) \\ \cup^k \uparrow & & \cup^k \uparrow \\ H^2(B; \Lambda) & \xrightarrow{\bar{\nu}'^*, \cong} & H^2(M'; \Lambda). \end{array}$$

The left hand side \cup^k is determined by B , while the bottom isomorphism is determined by choosing once and for all a generator of $H^2(B; \Lambda) = H^2(\mathbb{C}P^\infty; \Lambda) \cong H^2(\mathbb{C}P^\infty) \otimes_{\mathbb{Z}} \Lambda = \Lambda$ and basing $H^2(M'; \Lambda) = \Lambda$ by the image z of that generator. This shows that the data of $PE(M', \bar{\nu}')$ is determined by \cup^k and thus by $z^k \in H^{2k}(M'; \Lambda) = \Lambda^2$.

For $b = 1$ we can explicitly compute the 2-sided primitive embedding and boundary automorphism associated to $\bar{\nu}_{a,1}$.

Corollary 9.10. *If $(2k)!$ divides a , then the 2-sided primitive embedding $(\iota_{\bar{\nu}}, j_{\bar{\nu}})$ associated to the $(2k-1)$ -smoothing $\bar{\nu}_{a,1}: M_{a,1} \rightarrow B_{2k-1}(M_{a,1})$ is isomorphic to*

$$(\Lambda, -a) \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} (\Lambda, -a) \cup_{\text{Id}} (\Lambda, a) \xleftarrow{\begin{pmatrix} 1 \\ 2a \end{pmatrix}} (\Lambda, a).$$

In particular, the boundary automorphism $\delta_{\bar{\nu}}$ is trivial in $\ell\text{Aut}(\partial(\Lambda, -a))$.

Proof. For $b = 1$, Proposition 9.8 implies that the 2-sided primitive embedding associated to $\bar{\nu}_{a,1}$ is

$$(j_{a,1}, \iota_{a,1}) = \left((\Lambda, -a) \xrightarrow{\begin{pmatrix} -a \\ 1 \end{pmatrix}} (\Lambda^2, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}) \xleftarrow{\begin{pmatrix} a \\ 1 \end{pmatrix}} (\Lambda, a) \right).$$

The quadratic form on the union $(\Lambda, -a) \cup_{\text{Id}} (\Lambda, a)$ is represented by $\begin{pmatrix} -a & 0 \\ 1 & 0 \end{pmatrix}$. A quick verification shows that the isometry $\begin{pmatrix} 0 & 1 \\ 1 & a \end{pmatrix}: H_+(\Lambda) \rightarrow (\Lambda, -a) \cup_{\text{Id}} (\Lambda, a)$ provides the desired isomorphism of 2-sided primitive embeddings:

$$\begin{array}{ccc} (\Lambda, -a) & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & (\Lambda, -a) \cup_{\text{Id}} (\Lambda, a) & \xleftarrow{\begin{pmatrix} 1 \\ 2a \end{pmatrix}} & (\Lambda, a) \\ \uparrow = & & \uparrow \begin{pmatrix} 0 & 1 \\ 1 & -a \end{pmatrix} & & \uparrow = \\ (\Lambda, -a) & \xrightarrow{\begin{pmatrix} -a \\ 1 \end{pmatrix}} & H_+(\Lambda) & \xleftarrow{\begin{pmatrix} a \\ 1 \end{pmatrix}} & (\Lambda, a). \end{array}$$

The assertion on boundary isomorphisms now follows from the correspondence between 2-sided primitive embeddings and boundary isomorphisms (Theorem 6.11). \square

9.3. Proof of Theorem 9.1. We can now prove Theorem 9.1, which posits the existence of infinitely many stably parallelisable $4k$ -manifolds M_1, M_2, \dots with $\pi_1(M_i) = \mathbb{Z}$ that are stably parallelisable, have hyperbolic intersection form, identical Euler characteristic, are pairwise not stably diffeomorphic, and satisfy the following: for each i , there are infinitely many $4k$ -manifolds M_i^1, M_i^2, \dots with $\pi_1(M_i^k) = \mathbb{Z}$ that are stably parallelisable, have hyperbolic intersection form, are all stably diffeomorphic to M_i , but are pairwise not homotopy equivalent.

Proof of Theorem 9.1. Fix $k \geq 2$. Given an integer $p < 0$ that is divisible by $(2k)!$, set $q := -p(1 - 4p^2) > 0$. This is also divisible by $(2k)!$. The smooth closed $4k$ -manifolds $M_{q,1} = N_{q,1} \# (S^1 \times S^{4k-1})$ are stably parallelisable and have fundamental group \mathbb{Z} . The description of the cohomology ring of $N_{q,1}$ from Proposition 9.3 implies that the $M_{q,1}$ are pairwise not stably diffeomorphic. Next, consider the $(2k-1)$ -smoothing

$$\bar{\nu}_{q,1}: M_{q,1} \rightarrow B_{2k-1}(M_{q,1}) = P_{2k-1}(M_{q,1}) \times BO\langle 4k \rangle$$

constructed in Lemma 9.6. Corollary 9.10 shows that the boundary automorphism associated to $(M_{q,1}, \bar{\nu}_{q,1})$ is trivial in $\text{Aut}(\partial(\Lambda, -q))$. As $q = p(1 - 4p^2)$, Theorem 5.8 shows that the boundary automorphism group $\text{bAut}_{H_+(\Lambda)}(\partial(\Lambda, -q))$ is infinite. Recall from Construction 8.2 that there is an action of $\text{Aut}(\pi_1(M_{q,1})) = \mathbb{Z}_2$ on $\text{bPrim}^{\mathbb{Z}} := \bigcup_{[m,v,v']} \text{bPrim}_m(v, v')$. Use the bijection from Theorem 6.11 to pull this action back to $\text{bIso}^{\mathbb{Z}} := \bigcup_{[m,v,v']} \text{bIso}_m(v, v')$. Using that $\text{bAut}_{H_+(\Lambda)}(\partial(\Lambda, -q))$ is infinite and applying our realisation result, we will construct infinitely many homotopically inequivalent manifolds.

Corollary 7.12 implies the following: for each $f \in \text{Aut}_{H_+(\Lambda)}(\partial(\Lambda, -q))$, there is a normal $(2k-1)$ -smoothing $(M_{q,1}^f, \bar{\nu}_{q,1}^f)$ such that the $4k$ -manifold $M_{q,1}^f$ has hyperbolic intersection form, is stably diffeomorphic to $M_{q,1}$, and the boundary isomorphism associated with $(M_{q,1}^f, \bar{\nu}_{q,1}^f)$ is

$$\delta_{q,1}^f = [f] \in \text{bAut}_{H_+(\Lambda)}(\partial(\Lambda, -q)).$$

The action by $\text{Aut}(\pi_1(M_{q,1})) = \text{Aut}(\mathbb{Z}) = \mathbb{Z}_2$ may lead to some of these boundary automorphisms being identified in $\text{bAut}^{\mathbb{Z}}/\mathbb{Z}_2$. However since \mathbb{Z}_2 is finite, we still have an infinite family $\mathcal{F} \subseteq \text{bAut}_{H_+(\Lambda)}(\partial(\Lambda, -q))$ of elements that are pairwise distinct in $\text{bAut}^{\mathbb{Z}}/\mathbb{Z}_2$. In particular, for any $f_1, f_2 \in \mathcal{F}$, the two-sided primitive embeddings $PE(M_{q,1}^{f_1}, \bar{\nu}_{q,1}^{f_1})$ and $PE(M_{q,1}^{f_2}, \bar{\nu}_{q,1}^{f_2})$ do not agree in $\text{bPrim}^{\mathbb{Z}}/\mathbb{Z}_2$.

For a contradiction, assume that some pair of manifolds among the $M_{q,1}^f$ is homotopy equivalent, say $M_{q,1}^{f_1}, M_{q,1}^{f_2}$. By Lemma 9.2, the homotopy equivalence must be orientation preserving: indeed each $M_{q,1}^f$ is stably diffeomorphic to $M_{q,1}$, a manifold with $H^2(M_{q,1}; \mathbb{Z}) = \mathbb{Z}\langle z \rangle$, $z^{2k} \neq 0$, and $z^{2k} \cap [M_{q,1}] > 0$. For an orientation preserving homotopy equivalence, Corollary 8.12 implies that $PE(M_{q,1}^{f_1}, \bar{\nu}_{q,1}^{f_1})$ and $PE(M_{q,1}^{f_2}, \bar{\nu}_{q,1}^{f_2})$ agree in $\text{bPrim}^{\mathbb{Z}}/\mathbb{Z}_2$. This is a contradiction.

Finally, note that each manifold $M_{q,1}^f$ is stably parallelisable since its stable normal bundle lifts to $BO\langle 4k \rangle$. This concludes the proof of Theorem 9.1, and therefore of Theorem 1.1. \square

10. EXTENDED SYMMETRIC FORMS AND THE HOMOTOPY STABLE CLASS

In this section we prove Theorem 1.6. For the convenience of the reader we recall the statement.

Theorem 10.1. *For the $4k$ -dimensional manifolds M_i with $\pi_1(M_i) \cong \mathbb{Z}$, $k \geq 2$, constructed in Section 9 for the proof of Theorem 1.1,*

$$\mu: \mathcal{S}_+^{\text{st}}(M_i)/L_{4k+1}(\Lambda) \rightarrow \mathcal{S}^{\text{st}}(\mu(M_i, \bar{\nu}_i))$$

is surjective and has infinite image. Moreover $\text{Aut}(\mathbb{Z}) = \{\pm 1\}$ is finite, so after further identifying two extended symmetric forms related by changing the identification of the fundamental group with \mathbb{Z} , the image remains infinite.

Proof. In this proof $\pi = \mathbb{Z}$ and $\Lambda := \mathbb{Z}[\pi] = \mathbb{Z}[\mathbb{Z}]$. We start by showing that the image is infinite. Consider extended symmetric forms $(H, \lambda, \nu: H \rightarrow Q)$ with H and Q f.g. free and λ even. We shall construct a map from isomorphism classes of extended symmetric forms satisfying these hypotheses to isomorphism classes of 2-sided-primitive embeddings, and then we shall use this map to distinguish the extended symmetric forms of the manifolds we constructed in Section 9. The inclusion $i_\nu: \ker(\nu) \subseteq H$ and the composition of $\nu^*: Q^* \rightarrow H^*$ with $\hat{\lambda}^{-1}: H^* \rightarrow H$ (the inverse of the adjoint of λ) produce Λ -homomorphisms that we claim fit into a 2-sided primitive embedding

$$\ker(\nu) \xrightarrow{i_\nu} H \xleftarrow{j_\nu} Q^*.$$

Here we use the assumption that λ is an even symmetric form, and that the orientation character is trivial to see that (H, λ) admits a uniquely determined quadratic refinement θ . Use i_ν and j_ν to pull the quadratic form θ back from H to $\ker(\nu)$ and Q^* respectively. This leads to the quadratic forms:

$$(\ker(\nu), \theta) := (\ker(\nu), i_\nu^* \theta i_\nu) \text{ and } (Q^*, \theta) := (Q^*, j_\nu^* \theta j_\nu).$$

Observe that both i_ν and j_ν are injective: i_ν is an inclusion, while for j_ν use that the dual of a surjective map is injective. The same verification as the one made in the proof of Lemma 7.4 implies that we have a 2-sided primitive embedding

$$t(H, \lambda, \nu) := \left((\ker(\nu), \theta) \xrightarrow{i_\nu} (H, \theta) \xleftarrow{j_\nu} (Q^*, \theta) \right)$$

as claimed. Next we claim we have defined a map

$$t: \text{bES}^{\mathbb{Z}} \rightarrow \text{bPrim}^{\mathbb{Z}}$$

from the set $\text{bES}^{\mathbb{Z}}$ of isomorphism classes of extended symmetric forms with H and Q f.g. free and λ even to the set of isomorphism classes of 2-sided primitive embeddings $\text{bPrim}^{\mathbb{Z}}$. This latter set can be thought of as the union of $\text{bPrim}_m(v, v')$ over all isomorphism classes of m , v , and v' .

Assume that (f, g) is an isometry between extended symmetric forms (H_1, λ_1, ν_1) and (H_2, λ_2, ν_2) in $\text{bES}^{\mathbb{Z}}$. To prove the claim we must show that $t(H_1, \lambda_1, \nu_1)$ and $t(H_2, \lambda_2, \nu_2)$ agree in $\text{bPrim}^{\mathbb{Z}}$. By assumption, we have an isometry $f: (H_1, \lambda_1) \rightarrow (H_2, \lambda_2)$ and an isomorphism $g: Q_1 \rightarrow Q_2$ such that $\nu_2 f = g \nu_1$. Since θ_1 and θ_2 are determined by λ_1 and λ_2 respectively, it follows that f_* also determines an isometry of the quadratic refinements. In particular, f restricts to an isometry $f|: (\ker(\nu_1), \theta_1) \rightarrow (\ker(\nu_2), \theta_2)$. Additionally, the

isomorphism $g: Q_1 \rightarrow Q_2$ induces an isomorphism $g^*: Q_2^* \rightarrow Q_1^*$ and consider the following diagram:

$$\begin{array}{ccccc} (\ker(\nu_1), \theta_1) & \xrightarrow{i_{\nu_1}} & (H_1, \theta_1) & \xleftarrow{j_{\nu_1}} & (Q_1^*, \theta_1), \\ \cong \downarrow f_1 & & \cong \downarrow f & & \cong \uparrow g^* \\ (\ker(\nu_2), \theta_2) & \xrightarrow{i_{\nu_2}} & (H_2, \theta_2) & \xleftarrow{j_{\nu_2}} & (Q_2^*, \theta_2). \end{array}$$

The left square commutes because i_{ν_1} and i_{ν_2} are inclusion maps. The right square is seen to commute by using successively the definition of j_{ν_1} , the fact that $\nu_2 f = g \nu_1$, the fact that f is an isometry, and the definition of j_{ν_2} :

$$f j_{\nu_1} g^* = f(\widehat{\lambda}_1^{-1} \nu_1^*) g^* = f \widehat{\lambda}_1^{-1} f^* \nu_2^* = \widehat{\lambda}_2^{-1} \nu_2^* = j_{\nu_2}.$$

One can then check that g^* is in fact an isometry of the quadratic forms. This completes the proof that the map t is well-defined on isomorphism classes.

By inspection of the construction, if $(H, \lambda, \nu: H \rightarrow Q) = \mu(M, \bar{\nu})$ then $t(H, \lambda, \nu)$ coincides with the 2-sided primitive embedding corresponding to $(M, \bar{\nu})$; recall Definition 7.5. By Theorem 1.1, we know that there are infinitely many manifolds M_i^1, M_i^2, \dots each of which is stably diffeomorphic to M_i but that are pairwise not homotopy equivalent. We detected that the M_i^j are not homotopy equivalent by using the Postnikov primitive embedding of M_i^j , which coincides with $t(\mu(M_i^j)) \in \text{bPrim}^{\mathbb{Z}}$; see Section 8.2. Since the $t(\mu(M_i^j))$ are pairwise distinct, so must the $\mu(M_i^j)$ be distinct in $\text{bES}^{\mathbb{Z}}$. This concludes the proof of the fact that μ has infinite image. The action of the automorphisms $\pi = \mathbb{Z}$ can at most identify these elements in pairs, so indeed we have infinitely many pairwise not homotopy equivalent manifolds.

Now we show that μ is surjective. Let (H, λ, ν) be an extended symmetric form representing an element of $\mathcal{S}^{\text{st}}(\mu(M_i, \bar{\nu}_i))$, where $\bar{\nu}_i: M_i \rightarrow B_i = P_{2k-1}(M_i) \times BO\langle 4k \rangle$ is a normal $(2k-1)$ -smoothing. Since $(H, \lambda, \nu) \in \mathcal{S}^{\text{st}}(\mu(M_i, \bar{\nu}_i))$, there is an isometry α and an isomorphism β that fit into the following commutative diagram

$$\begin{array}{ccc} (H_q(M_i; \Lambda), \lambda_{M_i}) \oplus H^+(\Lambda)^{\oplus r} & \xrightarrow{(\bar{\nu}_i \ 0)} & H_q(B_i; \Lambda) \\ \cong \downarrow \alpha & & \cong \downarrow \beta \\ (H, \lambda) \oplus H^+(\Lambda)^{\oplus r} & \xrightarrow{(\nu \ 0)} & Q. \end{array}$$

Perform r trivial $(2k-1)$ surgeries on M_i to obtain $M_i \#^r (S^{2k} \times S^{2k})$, with a cobordism over B_i to M_i . The intersection form of $M_i \#^r (S^{2k} \times S^{2k})$ is $(H_q(M_i; \Lambda), \lambda_{M_i}) \oplus H^+(\Lambda)^{\oplus r}$. Then consider

$$G := \alpha^{-1}(H^+(\Lambda)^{\oplus r}) \subseteq \lambda_{M_i} \oplus H^+(\Lambda)^{\oplus r}.$$

This is a hyperbolic summand, since α is an isometry. Surgery on a collection of framed embedded $2k$ -spheres in $M_i \#^r (S^{2k} \times S^{2k})$ representing a lagrangian for G produces a $4k$ -manifold M_i' with intersection pairing isomorphic to $G^\perp \cong (H, \lambda)$, via $\alpha|_{G^\perp}$. We may find such framed embedded spheres since we have a lagrangian that maps trivially to $H_{2k}(B_i; \Lambda)$, under the normal smoothing $\bar{\nu}_i$. Also note that the vanishing of the intersection form implies the

vanishing of the quadratic enhancement. We have that $\beta \circ (\bar{\nu}_i \oplus 0)(G) = \{0\}$ by commutativity of the diagram. Hence

$$\begin{array}{ccc} G^\perp & \xrightarrow{\alpha|_{G^\perp}} & (H, \lambda) \\ & \searrow \beta \circ (\bar{\nu}_i \oplus 0) & \swarrow \nu \\ & & Q \end{array}$$

commutes and the extended symmetric form of M'_i is isomorphic to (H, λ, ν) as desired. Since the second set of surgeries are on spheres in the kernel of $\bar{\nu}_i: H_{2k}(M_i \#^r (S^{2k} \times S^{2k}); \Lambda) \rightarrow H_{2k}(B_i; \Lambda)$, the trace of the surgeries is a cobordism over B_i . The map to B_i factors through the normal $(2k-1)$ -type $P_{2k-1}(M_i) \times BO\langle 2k \rangle$, so we deduce that M_i and M'_i are bordant over their $(2k-1)$ type and have the same Euler characteristic. It then follows from [Kre99, Theorem C] that M_i and M'_i are stably diffeomorphic, so $M'_i \in \mathcal{S}_+^{\text{st}}(M_i)$ and therefore $\mu(M'_i) \in \mathcal{S}^{\text{st}}(\mu(M_i, \bar{\nu}_i))$. Alternatively one can directly see from the construction of M'_i that it stabilises to $M_i \#^r (S^{2k} \times S^{2k})$. This completes the proof that $\mu: \mathcal{S}_+^{\text{st}}(M_i) \rightarrow \mathcal{S}^{\text{st}}(\mu(M_i, \bar{\nu}_i))$ is surjective. \square

REFERENCES

- [Bau77] Hans J. Baues. *Obstruction theory on homotopy classification of maps*, volume 628. Springer, Cham, 1977.
- [BCD⁺20] Jonathon Bowden, Diarmuid Crowley, Jim Davis, Stefan Friedl, Carmen Rovi, and Stephan Tillmann. Open problems in the topology of manifolds. *MATRIX Annals*, 2019-20. https://www.matrix-inst.org.au/wp_Matrix2016/wp-content/uploads/2019/01/Bowden.pdf.
- [BDK07] Jeremy Brookman, James F. Davis, and Qayum Khan. Manifolds homotopy equivalent to $P^n \# P^n$. *Math. Ann.*, 338(4):947–962, 2007.
- [CCPS21] Anthony Conway, Diarmuid Crowley, Mark Powell, and Joerg Sixt. Simply connected manifolds with large homotopy stable classes. *Preprint, available at arXiv:2109.00654*, 2021.
- [CHR95] Alberto Cavicchioli, Friedrich Hegenbarth, and Dušan Repovš. On the stable classification of certain 4-manifolds. *Bull. Austral. Math. Soc.*, 52(3):385–398, 1995.
- [Coh77] Paul Moritz Cohn. *Algebra. Vol. 2*. John Wiley & Sons, London-New York-Sydney, 1977. With errata to Vol. I.
- [CS11] Diarmuid Crowley and Joerg Sixt. Stably diffeomorphic manifolds and $l_{2q+1}(\mathbb{Z}[\pi])$. *Forum Math.*, 23(3):483–538, 2011.
- [Dav05] James Davis. The Borel/Novikov conjectures and stable diffeomorphisms of 4-manifolds. In *Geometry and topology of manifolds*, volume 47 of *Fields Inst. Commun.*, pages 63–76. Amer. Math. Soc., Providence, RI, 2005.
- [FQ90] Michael Freedman and Frank Quinn. *Topology of Four Manifolds*. Princeton Univ. Press, 1990.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [HK88a] Ian Hambleton and Matthias Kreck. On the classification of topological 4-manifolds with finite fundamental group. *Math. Ann.*, 280(1):85–104, 1988.
- [HK88b] Ian Hambleton and Matthias Kreck. Smooth structures on algebraic surfaces with cyclic fundamental group. *Invent. Math.*, 91(1):53–59, 1988.
- [HK93a] Ian Hambleton and Matthias Kreck. Cancellation, elliptic surfaces and the topology of certain four-manifolds. *J. Reine Angew. Math.*, 444:79–100, 1993.
- [HK93b] Ian Hambleton and Matthias Kreck. Cancellation of hyperbolic forms and topological four-manifolds. *J. Reine Angew. Math.*, 443:21–47, 1993.
- [HKT09] Ian Hambleton, Matthias Kreck, and Peter Teichner. Topological 4-manifolds with geometrically two-dimensional fundamental groups. *J. Topol. Anal.*, 1(2):123–151, 2009.
- [Kha17] Qayum Khan. Cancellation for 4-manifolds with virtually abelian fundamental group. *Topology Appl.*, 220:14–30, 2017.

- [KLPT17] Daniel Kasprowski, Markus Land, Mark Powell, and Peter Teichner. Stable classification of 4-manifolds with 3-manifold fundamental groups. *J. Topol.*, 10(3):827–881, 2017.
- [KPT20] Daniel Kasprowski, Mark Powell, and Peter Teichner. Algebraic criteria for stable diffeomorphism of spin 4-manifolds. Preprint: arXiv:2006.06127, 2020.
- [Kre99] Matthias Kreck. Surgery and duality. *Ann. of Math. (2)*, 149(3):707–754, 1999.
- [KS84] Matthias Kreck and James A. Schafer. Classification and stable classification of manifolds: some examples. *Comment. Math. Helv.*, 59:12–38, 1984.
- [Lam99] Tsit Yuen Lam. *Lectures on modules and rings*. Springer-Verlag, New York, 1999.
- [Lam06] Tsit Yuen Lam. *Serre’s problem on projective modules*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2006.
- [Lev77] Jerome Levine. Knot modules. *Trans. Amer. Math. Soc.*, 229:1–50, 1977.
- [PRT20] Mark Powell, Arunima Ray, and Peter Teichner. The 4-dimensional disc embedding theorem and dual spheres. Preprint, available at arXiv:2006.05209, 2020.
- [Ran81] Andrew Ranicki. *Exact sequences in the algebraic theory of surgery*. Princeton University Press, Princeton, N.J., 1981.
- [Ran98] Andrew Ranicki. *High-dimensional knot theory*. Springer-Verlag, New York, 1998. Algebraic surgery in codimension 2. With an appendix by Elmar Winkelkemper.
- [Ran02] Andrew Ranicki. *Algebraic and Geometric Surgery*. Oxford University Press, Oxford, UK, 2002.
- [Spa03] Fulvia Spaggiari. On the stable classification of Spin four-manifolds. *Osaka J. Math.*, 40(4):835–843, 2003.
- [Wal62] Charles Terence Clegg Wall. Classification of $(n - 1)$ -connected $2n$ -manifolds. *Ann. of Math. (2)*, 75:163–189, 1962.
- [Wal99] Charles Terence Clegg Wall. *Surgery on compact manifolds*. American Mathematical Society, Providence, RI, second edition, 1999. Edited and with a foreword by A. A. Ranicki.
- [Wei99] Shmuel Weinberger. Higher ρ -invariants. In *Tel Aviv Topology Conference: Rothenberg Festschrift (1998)*, volume 231 of *Contemp. Math.*, pages 315–320. Amer. Math. Soc., Providence, RI, 1999.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE MA 02139, UNITED STATES
Email address: anthonyconway@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, PARKVILLE,
 VIC, 3010, AUSTRALIA
Email address: dcrowley@unimelb.edu.au

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, UNITED KINGDOM
Email address: mark.a.powell@durham.ac.uk

Email address: sixtj@yahoo.de