

# OBSTRUCTIONS TO INTEGRABILITY OF NEARLY INTEGRABLE DYNAMICAL SYSTEMS NEAR REGULAR LEVEL SETS

SHOYA MOTONAGA AND KAZUYUKI YAGASAKI

ABSTRACT. We study the existence of real-analytic first integrals and real-analytic integrability for perturbations of integrable systems in the sense of Bogoyavlenskij including non-Hamiltonian ones. We especially assume that there exists a family of periodic orbits on a regular level set of the first integrals having a connected and compact component and give sufficient conditions for nonexistence of the same number of real-analytic first integrals in the perturbed systems as the unperturbed ones and for their real-analytic nonintegrability near the level set such that the first integrals and commutative vector fields depend analytically on the small parameter. We compare our results with classical results of Poincaré and Kozlov for systems written in action and angle coordinates and discuss their relationships with the subharmonic and homoclinic Melnikov methods for periodic perturbations of single-degree-of-freedom Hamiltonian systems. We illustrate our theory for three examples containing the periodically forced Duffing oscillator.

## 1. INTRODUCTION

In his famous memoir [22], which was related to a prize competition celebrating the 60th birthday of King Oscar II, Henri Poincaré studied two-degree-of-freedom Hamiltonian systems depending on a small parameter, say  $\varepsilon$  here although he used the letter  $\mu$  instead, such that they are integrable when  $\varepsilon = 0$ , and showed the nonexistence of first integrals which are analytic in the state variables and parameter  $\varepsilon$  and functionally independent of Hamiltonians, under some nondegenerate conditions. If there exists such a first integral, then the Hamiltonian systems are integrable for  $|\varepsilon| \geq 0$  sufficiently small in the sense of Liouville [2, 16]. The result was improved significantly in the first volume of his masterpieces [23] published two years later, so that more-degree-of-freedom Hamiltonian systems can be treated. Using these results, he discussed the nonexistence of such first integrals in the restricted planar and spacial three-body problem there. See also [7] for an account of his work from a mathematical and historical perspectives. Subsequently, his results were sophisticated and generalized to non-Hamiltonian systems [13, 14]. In particular, Kozlov [14] treated multi-dimensional systems of the form

$$\dot{I} = \varepsilon h(I, \theta; \varepsilon), \quad \dot{\theta} = \omega(I) + \varepsilon g(I, \theta; \varepsilon), \quad (I, \theta) \in \mathbb{R}^\ell \times \mathbb{T}^m, \quad (1.1)$$

---

*Date:* November 17, 2021.

*2020 Mathematics Subject Classification.* 37J30; 34A05; 34E10; 37C27; 34C25; 37C29; 34C37; 34C40; 37J40; 37J46.

*Key words and phrases.* Integrability; nearly integrable system; resonant periodic orbit; first integral; commutative vector field; Melnikov method.

where  $\varepsilon$  is a small parameter such that  $|\varepsilon| \ll 1$ ,  $\mathbb{T}^m = \prod_{j=1}^m \mathbb{S}^1$  with  $\mathbb{S}^1 = \mathbb{R}/2\pi\mathbb{Z}$  is an  $m$ -dimensional torus and  $h(I, \theta; \varepsilon)$ ,  $\omega(I)$  and  $g(I, \theta; \varepsilon)$  are analytic in  $(I, \theta, \varepsilon)$ . Note that the system (1.1) is Hamiltonian if  $\ell = m$  as well as  $\varepsilon = 0$  or

$$D_I h(I, \theta; \varepsilon) = -D_\theta g(I, \theta; \varepsilon),$$

and non-Hamiltonian if not. When  $\varepsilon = 0$ , Eq. (1.1) becomes

$$\dot{I} = 0, \quad \dot{\theta} = \omega(I) \tag{1.2}$$

which we refer to as the *unperturbed system* for (1.1). We often use this terminology for other systems below. Here we state some details of his result.

We expand  $h(I, \theta; 0)$  in Fourier series as

$$h(I, \theta; 0) = \sum_{r \in \mathbb{Z}^m} \hat{h}_r(I) \exp(ir \cdot \theta), \tag{1.3}$$

where  $\hat{h}_r(I)$ ,  $r \in \mathbb{Z}^m$ , are the Fourier coefficients and “ $\cdot$ ” represents the inner product. We assume the following for (1.1):

**(K1)** The system (1.1) has  $s$  first integrals  $F_j(I, \theta; \varepsilon)$ ,  $j = 1, \dots, s$ , which are analytic in  $(I, \theta, \varepsilon)$ ;

**(K2)** If  $r \in \mathbb{Z}^m$  and  $r \cdot \omega(I) = 0$  for any  $I \in \mathbb{R}^\ell$ , then  $r = 0$ .

If assumption (K2) holds, then we say that the unperturbed system (1.2) is *non-degenerate*. Under (K1) and (K2) we can show that  $F_j(I, \theta; 0)$ ,  $j = 1, \dots, s$ , are independent of  $\theta$  (see Lemma 1 in Section 1 of Chapter IV of [14]), and write  $F_{j0}(I) = F_j(I, \theta; 0)$  and  $F_0(I) = (F_{10}(I), \dots, F_{s0}(I))$ . We refer to  $\mathcal{P}_s \subset \mathbb{R}^\ell$  as a *Poincaré set* if for each  $I \in \mathcal{P}_s$  there exists linearly independent vectors  $r_j \in \mathbb{Z}^m$ ,  $j = 1, \dots, \ell - s$ , such that

- (i)  $r_j \cdot \omega(I) = 0$ ,  $j = 1, \dots, \ell - s$ ;
- (ii)  $\hat{h}_{r_j}(I)$ ,  $j = 1, \dots, \ell - s$ , are linearly independent.

Let  $U$  be a domain in  $\mathbb{R}^\ell$ . A set  $\Delta \subset U$  is called a *key set* (or *uniqueness set*) for  $C^\omega(U)$  if any analytic function vanishing on  $\Delta$  vanishes on  $U$ . For example, any dense set in  $U$  is a key set for  $C^\omega(U)$ . In this situation, we have the following theorem (see Section 1 of Chapter IV of [14] for its proof).

**Theorem 1.1** (Kozlov). *Suppose that assumptions (K1) and (K2) hold, the Jacobian matrix  $DF_0(I)$  has the maximum rank at a point  $I_0 \in \mathbb{R}^\ell$  and a Poincaré set  $\mathcal{P}_s \subset U$  is a key set for  $C^\omega(U)$ , where  $U$  is a neighborhood of  $I_0$ . Then the system (1.1) has no first integral which is real-analytic in  $(I, \theta, \varepsilon)$  and functionally independent of  $F_j(I, \theta; \varepsilon)$ ,  $j = 1, \dots, s$ , near  $\varepsilon = 0$ .*

A version of Theorem 1.1 for the Hamiltonian case  $\ell = m$  was given in [13] (see also Theorem 7.1 of [3]). The Hamiltonian case of  $s = 1$  with a dense Poincaré set in Theorem 1.1 was treated by Poincaré for  $\ell = m \geq 2$  in [23]. When  $s = 0$  in (K1), Theorem 1.1 means that under its hypotheses there exists no first integral which is analytic in  $(I, \theta, \varepsilon)$ . When  $s = 1$  in (K1), which always occurs if the system (1.1) is Hamiltonian, it means that under its hypotheses there exists no first integral which is analytic in  $(I, \theta) \in U \times \mathbb{T}^m$  and  $\varepsilon$  near  $\varepsilon = 0$  and functionally independent of  $F_1(I, \theta, \varepsilon)$ . When  $s = m - 1$  in (K1), if besides (K1) and (K2) there exists a key set  $\Delta$  for  $C^\omega(U)$  such that  $r \cdot \omega(I) = 0$  and  $h_r(I) \neq 0$  on  $\Delta$  for some  $r \in \mathbb{Z}^m \setminus \{0\}$ , there exists no additional first integral which is analytic in  $(I, \theta, \varepsilon)$ . For Hamiltonian systems, if they are not Liouville-integrable for any  $|\varepsilon| > 0$  sufficiently small, then

there does not exist such a first integral. For non-Hamiltonian systems this may be true in an appropriate meaning but some additional ingredients are needed as seen below.

In this paper we study more general dynamical systems of the form

$$\dot{x} = X_\varepsilon(x), \quad x \in \mathcal{M}, \tag{1.4}$$

where  $\varepsilon$  is a small parameter such that  $|\varepsilon| \ll 1$ ,  $\mathcal{M}$  is an  $n$ -dimensional analytic manifold for  $n \geq 2$  and the vector field  $X_\varepsilon$  is analytic in  $x$  and  $\varepsilon$ . Let  $X_\varepsilon = X^0 + \varepsilon X^1 + O(\varepsilon^2)$  for  $|\varepsilon| > 0$  sufficiently small. When  $\varepsilon = 0$ , the system (1.4) becomes

$$\dot{x} = X^0(x), \tag{1.5}$$

which is assumed to be *analytically*  $(q, n - q)$ -integrable in the following meaning of Bogoyavlenskij [8] for some positive integer  $q \leq n$ .

**Definition 1.2** (Bogoyavlenskij). *The system (1.5) is called  $(q, n - q)$ -integrable or simply integrable if there exist  $q$  vector fields  $Y_1(x) (:= X^0(x)), Y_2(x), \dots, Y_q(x)$  and  $n - q$  scalar-valued functions  $F_1(x), \dots, F_{n-q}(x)$  such that the following two conditions hold:*

- (i)  $Y_1(x), \dots, Y_q(x)$  are linearly independent almost everywhere and commute with each other, i.e.,  $[Y_j, Y_k](x) \equiv 0$  for  $j, k = 1, \dots, q$ , where  $[\cdot, \cdot]$  denotes the Lie bracket;
- (ii)  $F_1(x), \dots, F_{n-q}(x)$  are functionally independent, i.e.,  $dF_1(x), \dots, dF_{n-q}(x)$  are linearly independent almost everywhere, and  $F_1(x), \dots, F_{n-q}(x)$  are first integrals of  $Y_1, \dots, Y_q$ , i.e.,  $dF_k(Y_j) = 0$  for  $j = 1, \dots, q$  and  $k = 1, \dots, n - q$ .

If  $Y_1, Y_2, \dots, Y_q$  and  $F_1, \dots, F_{n-q}$  are analytic (resp. meromorphic), then Eq. (1.5) is said to be analytically (resp. meromorphic) integrable.

Definition 1.2 is considered as a generalization of Liouville-integrability for Hamiltonian systems [2, 16] since an  $m$ -degree-of-freedom Liouville-integrable Hamiltonian system with  $m \geq 1$  has not only  $m$  functionally independent first integrals but also  $m$  linearly independent commutative (Hamiltonian) vector fields generated by the first integrals. The  $(\ell + m)$ -dimensional system (1.2) is  $(m, \ell)$ -integrable in the Bogoyavlenskij sense.

The system (1.4) is regarded as a perturbation of the analytically integrable system (1.5). If there exist  $q'$  linearly independent, analytic commutative vector fields and  $n - q'$  functionally independent, analytic first integrals depending on  $\varepsilon$  analytically near  $\varepsilon = 0$  for some positive integer  $q' \leq n$ , then the perturbed system (1.4) is analytically  $(q', n - q')$ -integrable for  $|\varepsilon| \geq 0$  sufficiently small. We especially assume that there exists a  $q$ -parameter family of periodic orbits on a regular level set of the  $n - q$  first integrals having a connected and compact component (see Section 2 for our precise definitions containing this one) and give sufficient conditions for nonexistence of such  $n - q$  real-analytic first integrals and for real-analytic nonintegrability of the perturbed system (1.4) near the regular level set such that the first integrals and commutative vector fields depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ . The persistence of such first integrals and commutative vector fields in the perturbed system (1.4) along with that of periodic and homoclinic orbits was previously discussed in [20]. Our approach is different from those of Poincaré [22, 23] and Kozlov [13, 14] and based on the technique of [20]. Recently, another sufficient condition for nonintegrability of nearly integrable dynamical systems near

resonant periodic orbits was also obtained using a different approach in [30] and the theory was applied to prove the nonintegrability of the restricted three-body problem in [31], when the independent and state variables are extended to complex ones.

The unperturbed system (1.5) can be transformed to (1.2) under our assumptions, as stated in Proposition 2.1 below. However, Theorem 1.1 only says in our setting for (1.1) that there exists no such additional first integral even if its hypotheses hold. In particular, it does not allow us to determine the nonintegrability of (1.4) in the Bogoyavlenskij sense when it is non-Hamiltonian. We also describe a consequence of our results to (1.1) and show how it improves the results of Poincaré [23] and Kozlov [13, 14].

Moreover, we discuss a relationship between our results and the subharmonic and homoclinic Melnikov methods [11, 25, 27] for time-periodic perturbations of single-degree-of-freedom analytic Hamiltonian systems, which can be transformed to the form of (1.4) with  $\ell = 1$  and  $m = 2$ , i.e.  $(2, 1)$ -integrable, and have a one-parameter family of periodic orbits when  $\varepsilon = 0$ . As well known, if the subharmonic Melnikov functions have a simple zero, then the corresponding unperturbed periodic orbits persist in the perturbed systems. Morales-Ruiz [17] studied the Hamiltonian perturbation case in which the unperturbed systems have homoclinic orbits, and showed a relationship between their nonintegrability and a version due to Ziglin [33] of the homoclinic Melnikov method when the independent and state variables are extended to complex ones and the small parameter  $\varepsilon$  is also regarded as a state variable. More concretely, under some restrictive conditions, based on the generalized version due to Ayoul and Zung [5] of the Morales-Ramis theory [16, 18], which provides a sufficient condition for nonintegrability of autonomous dynamical systems, he essentially proved that they are meromorphically nonintegrable if the homoclinic Melnikov functions which are obtained as integrals along closed loops on the complex plane are not identically zero. Here the version of the Melnikov method enables us to detect transversal self-intersection of complex separatrices of periodic orbits and to prove its analytic nonintegrability, unlike the standard version [11, 15, 25]. Here we prove two variants of Morales-Ruiz [17] for periodic orbits: if the subharmonic Melnikov functions for a dense set of the unperturbed periodic orbits are not identically zero, then there exists no first integral depending analytically on  $\varepsilon$  near  $\varepsilon = 0$ ; and if the ‘standard’ homoclinic Melnikov functions [11, 15, 25] are not identically zero, then the perturbed systems are not Bogoyavlenskij-integrable such that the commutative vector fields and first integrals depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ .

We illustrate our theory for three examples: Simple pendulum with a constant torque, second-order coupled oscillators and the periodically forced Duffing oscillator [11, 12, 25]. Real-analytic nonintegrability is discussed in special cases of the second and third examples while existence of real-analytic first integrals in the rest. In particular, the special case of the third one is shown to be nonintegrable in the above meaning even if it does not have transverse homoclinic orbits to a periodic orbit.

This paper is organized as follows: In Section 2, we state our precise assumptions and main theorems. In Section 3, we give proofs of the main theorems. We describe a consequence of our results to (1.1) in Section 4 and discuss a relationship between our results and the subharmonic and homoclinic Melnikov methods

for time-periodic perturbations of single-degree-of-freedom Hamiltonian systems in Section 5. Finally, we give the three examples in Section 6.

## 2. MAIN RESULTS

In this section, we state our main results for (1.4). We first make the following assumptions on the unperturbed system (1.5):

- (A1)** For some positive integer  $q < n$ , the system (1.5) is analytically  $(q, n - q)$  Bogoyavlenskij-integrable, i.e., there exist  $q$  analytic vector fields  $Y_1(x) := X^0(x), \dots, Y_q(x)$  and  $n - q$  analytic scalar-valued functions  $F_1(x), \dots, F_{n-q}(x)$  such that conditions (i) and (ii) of Definition 1.2 hold.
- (A2)** Let  $F(x) = (F_1(x), \dots, F_{n-q}(x))$ . There exists a regular value  $c \in \mathbb{R}^{n-q}$  of  $F$ , i.e.,  $\text{rank } dF(x) = n - q$  when  $F(x) = c$ , such that  $F^{-1}(c)$  has a connected and compact component and  $Y_1(x), \dots, Y_q(x)$  are linearly independent on  $F^{-1}(c)$ .

Henceforth we assume without loss of generality that  $F^{-1}(c)$  is connected and compact itself in (A2), by reducing the domain of  $F$  if necessary. Under (A1) and (A2) we have the following result like a well-known theorem for Hamiltonian systems [2] (see [8, 34, 35] for the details).

**Proposition 2.1** (Liouville-Miuner-Arnold-Jost). *Suppose that assumptions (A1) and (A2) hold. Then we have the following:*

- (i) *The level set  $F^{-1}(c)$  is analytically diffeomorphic to the  $q$ -dimensional torus  $\mathbb{T}^q$ ;*
- (ii) *There exists an analytic diffeomorphism  $\varphi : U \times \mathbb{T}^q \rightarrow \mathcal{U}$ , where  $U$  and  $\mathcal{U}$  are, respectively, neighborhoods of  $I = I_0$  in  $\mathbb{R}^{n-q}$  and of  $F^{-1}(c)$  in  $\mathcal{M}$  for some  $I_0 \in \mathbb{R}^{n-q}$ , such that*
  - (iia)  $\varphi(\{I_0\} \times \mathbb{T}) = F^{-1}(c)$ ;
  - (iib)  $F \circ \varphi(I, \theta)$  depends only on  $I$ , where  $(I, \theta) \in U \times \mathbb{T}^q$ ;
  - (iic) *The flow of  $X^0$  on  $\mathcal{U}$  is analytically conjugate to that of (1.2) with  $\ell = n - q$  and  $m = q$  on  $U \times \mathbb{T}^q$ .*

The variables  $I$  and  $\theta$  are called the *action* and *angle variables* as in Hamiltonian systems, and  $\omega(I)$  is referred to as the *angular frequency vector*. Let  $\omega_j(I)$  be the  $j$ th component of  $\omega(I)$  for  $j = 1, \dots, q$ . Henceforth  $U$  denotes the neighborhood of  $I = I_0$  in Proposition 2.1. Moreover, we assume the following on (1.2) along with (K2):

- (A3)** There exists a key set  $D_R$  for  $C^\omega(U)$  such that for  $I \in D_R$  a resonance of multiplicity  $q - 1$ ,

$$\dim_{\mathbb{Q}} \langle \omega_1(I), \dots, \omega_q(I) \rangle = 1,$$

occurs with  $\omega(I) \neq 0$ , i.e., there exists a positive constant  $\omega_0(I)$  depending on  $I$  such that

$$\frac{\omega(I)}{\omega_0(I)} \in \mathbb{Z}^q \setminus \{0\}.$$

We choose  $\omega_0(I)$  as large as possible below.

We easily see that if  $\text{rank } D\omega(I^*) = q$  for some  $I^* \in U$ , then both (K2) and (A3) hold. Assumption (A3) also implies that if  $I \in D_R$ , then the system (1.2) has a  $q$ -parameter family of periodic orbits

$$(I, \theta) = (I, \omega(I)t + \tau), \quad \tau \in \mathbb{T}^q, \tag{2.1}$$

with the period  $T^I = 2\pi/\omega_0(I)$ . Note that the periodic orbits (2.1) is parameterized by  $(q-1)$  parameters essentially since two periodic orbits  $(I, \omega(I)t + \tau)$  and  $(I, \omega(I)t + \tau_0)$  represent the same orbit if  $\tau - \tau_0 = \omega(I)t_0$  for some  $t_0 \in \mathbb{R}$ . We also have a  $q$ -parameter (but essentially  $(q-1)$ -parameter) family of periodic orbits

$$\gamma_\tau^I(t) = \varphi(I, \omega(I)t + \tau), \quad (I, \tau) \in D_{\mathbb{R}} \times \mathbb{T}^q,$$

with the period  $T^I$  in the unperturbed system (1.5). Define the integrals

$$\mathcal{I}_{F_k}^I(\tau) := \int_0^{T^I} dF_j(X^1)_{\gamma_\tau^I(t)} dt, \quad k = 1, \dots, n-q, \quad (2.2)$$

for  $I \in D_{\mathbb{R}}$  and set  $\mathcal{I}_F^I(\tau) := (\mathcal{I}_{F_1}^I(\tau), \dots, \mathcal{I}_{F_{n-q}}^I(\tau))$ . Note that

$$\mathcal{I}_{F_k}^I(\tau + \omega(I)t) = \mathcal{I}_{F_k}^I(\tau)$$

for  $\tau \in \mathbb{T}^q$  and  $t \in \mathbb{R}$ . We state the first of our main results as follow.

**Theorem 2.2.** *Suppose that assumptions (A1)-(A3) and (K2) hold. If there exists a key set  $D \subset D_{\mathbb{R}}$  for  $C^\omega(U)$  such that  $\mathcal{I}_F^I(\tau)$  is not identically zero for any  $I \in D$ , then the perturbed system (1.4) does not have  $n-q$  real-analytic first integrals in a neighborhood of  $F^{-1}(c)$  near  $\varepsilon = 0$  such that they are functionally independent for  $|\varepsilon| \neq 0$  and depend analytically on  $\varepsilon$ .*

We prove Theorem 2.2 in Section 3.1.

We next consider a special case in which Eq. (1.4) is a two- or more-degree-of-freedom Hamiltonian system. For an integer  $m \geq 2$ , let  $(\mathcal{M}, \Omega)$  be a  $2m$ -dimensional analytic symplectic manifold with a symplectic form  $\Omega$ , and let  $H_\varepsilon(x) = H^0(x) + \varepsilon H^1(x) + O(\varepsilon^2)$  be an analytic Hamiltonian for (1.4) and depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ . We assume the following:

**(A1')** The unperturbed Hamiltonian system (1.5) with the Hamiltonian  $H^0(x)$  is real-analytically Liouville-integrable, i.e., there exist  $m$  functionally independent analytic first integrals  $F_1(x) (= H^0(x)), F_2(x), \dots, F_m(x)$  such that they are pairwise Poisson commutative, i.e.,  $\{F_j, F_k\} = 0$  for  $j, k = 1, \dots, m$ , where  $\{\cdot, \cdot\}$  denotes the Poisson bracket for the symplectic form  $\Omega$ .

Assumption (A1') means (A1) with  $q = m$  and  $Y_j = X_{F_j}$  for  $j = 1, \dots, m$ , where  $X_{F_j}$  denotes the Hamiltonian vector field for the Hamiltonian  $F_j(x)$ . So we immediately obtain the following from Theorem 2.2.

**Corollary 2.3.** *Suppose that (A1'), (A2), (A3) and (K2) hold. If there exists a key set  $D \subset D_{\mathbb{R}}$  for  $C^\omega(U)$  such that  $\mathcal{I}_F^I(\tau)$  with  $X_1 = X_{H^1}$  is not identically zero for any  $I \in D$ , then the perturbed Hamiltonian system for the Hamiltonian  $H_\varepsilon(x)$  is not real-analytically Liouville-integrable in a neighborhood of  $F^{-1}(c)$  near  $\varepsilon = 0$  such that  $m-1$  additional first integrals also depend analytically on  $\varepsilon$ .*

For integrability of non-Hamiltonian systems (1.4), we have to consider not only first integrals but also commutative vector fields. So we need the following assumption additionally:

**(A4)** For some  $I^* \in U$ , the Jacobian matrix  $D\omega(I^*)$  is injective, i.e.,

$$\text{rank } D\omega(I^*) = n - q.$$

Note that (A4) holds only when  $n - q \leq q$ . Finally, we state the second main result as follows.

**Theorem 2.4.** *Suppose that assumptions (A1)-(A4) and (K2) hold. If there exists a key set  $D \subset D_{\mathbb{R}}$  for  $C^\omega(U)$  such that  $\mathcal{I}_F^1(\tau)$  is not constant for any  $I \in D$ , then for  $|\varepsilon| \neq 0$  sufficiently small the perturbed system (1.4) is not real-analytically integrable in the Bogoyavlenskij sense near  $F^{-1}(c)$  such that the first integrals and commutative vector fields depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ : In particular there exists only such  $n - q - 1$  first integrals and such  $q - 1$  commutative vector fields at most.*

**Remark 2.5.**

- (i) *Theorems 2.2 and 2.4 exclude even the possibility that the first integrals are functionally independent for  $|\varepsilon| \neq 0$  sufficiently small but not for  $\varepsilon = 0$ .*
- (ii) *Theorem 2.4 does not exclude the possibility that the system (1.4) is real-analytically integrable in the Bogoyavlenskij sense near  $F^{-1}(c)$  such that in a punctured neighborhood of  $\varepsilon = 0$  there exist such  $n - q' (\geq 0)$  first integrals and  $q'$  commutative vector fields for some  $q' > q$ .*

We prove Theorem 2.4 in Section 3.2.

### 3. PROOFS OF THE MAIN THEOREMS

In this section, we prove Theorems 2.2 and 2.4. The proofs are based on the results of [20]. Henceforth we mean real-analyticity when saying analyticity.

**3.1. Proof of Theorem 2.2.** We first prove Theorem 2.2. Henceforth we assume that conditions (A1)-(A3) and (K2) hold. We begin with the following lemma.

**Lemma 3.1.** *Suppose that  $G_1(x), \dots, G_k(x)$  are analytic first integrals of the unperturbed system (1.5) near  $F^{-1}(c)$  for  $k \geq 1$  and they may be functionally dependent. Let  $G(x) = (G_1(x), \dots, G_k(x))$  and let  $\phi_t$  denote the flow generated by (1.5). Then the following statements hold:*

- (i) *There exists an analytic map  $\psi : F(\mathcal{U}) \rightarrow \mathbb{R}^k$  satisfying  $G = \psi \circ F$  in a neighborhood  $\mathcal{U}$  of  $F^{-1}(c)$  in  $\mathcal{M}$ .*
- (ii) *If  $dG_1, \dots, dG_k$  are linearly independent at  $x \in \mathcal{M}$ , then so are they at  $\phi_t(x)$ .*

*Proof.* For the proof of part (i) we first transform (1.5) to (1.2) with  $\ell = n - q$  and  $m = q$ , based on Proposition 2.1. In particular, by Proposition 2.1(iiia)  $F \circ \varphi(I, \theta)$  depends only on  $I$ . Let  $\tilde{F}(I) = F \circ \varphi(I, \theta)$ . On the other hand, we use Lemma 1 in Section 1 of Chapter IV of [14] to show that  $G(\varphi(I, \theta))$  depends only on  $I$ . Let  $\tilde{G}(I) = G(\varphi(I, \theta))$ . So we have

$$G = \tilde{G} \circ \varphi^{-1} = \tilde{G} \circ \tilde{F}^{-1} \circ \tilde{F} \circ \varphi^{-1}$$

and take  $\psi = \tilde{G} \circ \tilde{F}^{-1}$  to obtain part (i).

We turn to the proof of part (ii). Since the pull-back of  $G_j$  by  $\phi_t$  satisfies  $\phi_t^* G_j = G_j$  for  $j = 1, \dots, k$ , we have

$$\begin{aligned} \phi_t^*(dG_1 \wedge \dots \wedge dG_k)_x &= (\phi_t^* dG_1 \wedge \dots \wedge \phi_t^* dG_k)_x \\ &= (d(\phi_t^* G_1) \wedge \dots \wedge d(\phi_t^* G_k))_x = (dG_1 \wedge \dots \wedge dG_k)_x \neq 0, \end{aligned}$$

where ‘ $\wedge$ ’ represents the wedge product. On the other hand, by the definition of pull-back

$$\phi_t^*(dG_1 \wedge \dots \wedge dG_k)_x = (dG_1 \wedge \dots \wedge dG_k)_{\phi_t(x)} \circ (d\phi_t)_x.$$

Since  $\phi^t$  is a diffeomorphism, we see that  $(d\phi_t)_x$  is regular. Hence,

$$(dG_1 \wedge \dots \wedge dG_k)_{\phi_t(x)} \neq 0,$$

which yields part (ii).  $\square$

Using Theorem 2.2 of [20] on persistence of first integrals and Lemma 3.1, we obtain the following result.

**Lemma 3.2.** *Let  $I \in D_{\mathbb{R}}$ . Suppose that near  $\varepsilon = 0$  the perturbed system (1.4) has  $n - q$  analytic first integrals  $G_1^\varepsilon, \dots, G_{n-q}^\varepsilon$  near  $\mathcal{I}_I = \{\gamma_\tau^I \mid \tau \in \mathbb{T}^q\}$  in  $\mathcal{M}$  such that they are functionally independent on  $\mathcal{I}_I$  and depend analytically on  $\varepsilon$ . Then  $\mathcal{I}_F^I(\tau)$  must be identically zero.*

*Proof.* Assume that the hypothesis of the lemma holds. Using Theorem 2.2 of [20], we have

$$\mathcal{I}_{G^0}^I(\tau) = \int_0^{T^I} dG^0(X^1)_{\gamma_\tau^I(t)} dt = 0.$$

On the other hand, by Lemma 3.1,  $G^0 := (G_1^0, \dots, G_{n-q}^0)$  is expressed as  $G^0 = \psi \circ F$  for some analytic map  $\psi : F(\mathcal{U}) \rightarrow \mathbb{R}^{n-q}$ , where  $\mathcal{U}$  is a neighborhood of  $F^{-1}(c)$ . Since  $F$  is constant along the periodic orbit  $\gamma_\tau^I(t)$ , we have  $dG_{\gamma_\tau^I(t)}^0 = d\psi_{F(\gamma_\tau^I(0))} dF_{\gamma_\tau^I(t)}$ , so that

$$\mathcal{I}_{G^0}^I(\tau) = \int_0^{T^I} d\psi_{F(\gamma_\tau^I(0))} dF_{\gamma_\tau^I(t)}(X_{\gamma_\tau^I(t)}) dt = d\psi_{F(\gamma_\tau^I(0))} \mathcal{I}_F^I(\tau).$$

Since  $dG^0$  and  $dF$  have the maximum rank on  $\gamma_\tau^I(t)$  so that  $\text{rank } d\psi_{F(\gamma_\tau^I(0))} = n - q$ , we obtain  $\mathcal{I}_F^I(\tau) = 0$  for any  $\tau \in \mathbb{T}^q$ .  $\square$

For the proof of Theorem 2.2 we also need the following result (see Appendix A for its proof).

**Proposition 3.3.** *Let  $k \leq n - q$  be a positive integer. Suppose that in a neighborhood of  $F^{-1}(c)$  the perturbed system (1.4) has  $k$  first integrals that are analytic in  $(x, \varepsilon)$  near  $\varepsilon = 0$ . If they are functionally independent for  $\varepsilon \neq 0$ , then in a neighborhood of  $F^{-1}(c)$  there exist  $k$  first integrals that are analytic in  $(x, \varepsilon)$  and functionally independent near  $\varepsilon = 0$ .*

**Remark 3.4.** *From Proposition 3.3 we also see that the condition for  $DF_0(I)$  to have a maximum rank at a point  $I_0 \in \mathbb{R}^\ell$  was unnecessary in Theorem 1.1.*

*Proof of Theorem 2.2.* Suppose that  $\mathcal{I}_F^I(\tau)$  is not identically zero for any  $I \in D$ . Using Lemma 3.2 for each  $I \in D$ , we see that the perturbed system (1.4) does not have  $n - q$  analytic first integrals near  $\mathcal{I}_I$  such that they are functionally independent on  $\mathcal{I}_I$  and depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ .

Additionally, suppose that there are  $n - q$  analytic first integrals such that they are functionally independent for  $|\varepsilon| \neq 0$  sufficiently small but not at  $\varepsilon = 0$  and depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ . Then by Proposition 3.3 there exist  $n - q$  analytic first integrals  $G_1^\varepsilon, \dots, G_{n-q}^\varepsilon$  which are functionally independent and depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ . Hence,  $dG_1^0, \dots, dG_{n-q}^0$  are linearly dependent on  $\mathcal{I}_I$  for  $I \in D$ . As in the proof of Lemma 3.1 (i), we consider the transformed system (1.2) and write  $\tilde{G}_j(I) = G_j(\varphi(I, \theta))$  for  $j = 1, \dots, n - q$ . Let  $\tilde{G}(I) = (\tilde{G}_1(I), \dots, \tilde{G}_{n-q}(I))$ . We see that the determinant of the Jacobi matrix of  $\tilde{G}(I)$  is



zero for  $I \in D$ , so that it is identically zero on  $U$  since  $D$  is a key set for  $C^\omega(U)$ . This yields a contradiction.  $\square$

**3.2. Proof of Theorem 2.4.** We turn to the proof of Theorem 2.4. Henceforth, we assume that conditions (A1)-(A4) and (K2) hold.

We begin with the following lemma. Recall that  $Y_l(x)$ ,  $l = 1, \dots, q$ , are commutative vector fields for the unperturbed system (1.5).

**Lemma 3.5.** *An analytic vector field  $Z(x)$  commutes with the vector field  $X^0(x)$  if and only if it can be written as  $Z(x) = \sum_{l=1}^q \rho_l(F(x))Y_l(x)$ , where  $\rho_l : \mathbb{R}^{n-q} \rightarrow \mathbb{R}$ ,  $l = 1, \dots, n - q$ , are analytic. In particular, the vector field  $X^0(x)$  has only  $q$  commutative vector fields which are linearly independent almost everywhere.*

*Proof.* By Proposition 2.1, we may consider (1.2). So we prove that it is necessary and sufficient for an analytic vector field  $Z(I, \theta)$  to commute with the vector field of (1.2) that it can be written as

$$Z(I, \theta) = \sum_{l=1}^q \rho_l(I) \frac{\partial}{\partial \theta_l}. \quad (3.1)$$

The sufficiency is obvious. The necessity follows from assumptions (K2) and (A4) by Lemma 1 in Section 3 of Chapter IV of [14].  $\square$

Using Theorem 3.5 of [20] on persistence of commutative vector fields and Lemma 3.5, we obtain the following.

**Lemma 3.6.** *Let  $I \in D_{\mathbb{R}}$ . Suppose that near  $\varepsilon = 0$  the perturbed system (1.4) has  $q$  analytic commutative vector fields  $Z_1^\varepsilon(x), \dots, Z_q^\varepsilon(x)$  near  $\mathcal{T}_I = \{\gamma_\tau^I \mid \tau \in \mathbb{T}^q\}$  in  $\mathcal{M}$  such that they are linearly independent on  $\mathcal{T}_I$  and depend analytically on  $\varepsilon$ . Then  $\mathcal{S}_F^I(\tau)$  must be constant.*

*Proof.* We first transform (1.4) to (1.1) with  $\ell = n - q$  and  $m = q$ , as in the proof of Lemma 3.1. Assume that the hypothesis of the lemma holds. Then by Lemma 3.5,  $Z_j^0(I, \theta)$  has the form (3.1) with  $\rho_l(I) = \rho_{jl}(I)$ ,  $l = 1, \dots, q$ , for  $j = 1, \dots, q$ . Let

$$\tilde{X}^1(I, \theta) = \sum_{j=1}^{\ell} h_j(I, \theta; 0) \frac{\partial}{\partial I_j} + \sum_{k=1}^m g_k(I, \theta; 0) \frac{\partial}{\partial \theta_k}$$

where  $h_j(I, \theta; \varepsilon)$  (resp.  $g_k(I, \theta; \varepsilon)$ ) is  $j$ th (resp.  $k$ th) component of  $h(I, \theta; \varepsilon)$  (resp.  $g(I, \theta; \varepsilon)$ ) for  $j = 1, \dots, \ell$  (for  $k = 1, \dots, m$ ). Using Theorem 3.5 of [20], we have

$$\begin{aligned} 0 &= \int_0^{T^I} dI \left( \left[ Z_j^0(I, \theta), \tilde{X}^1(I, \theta) \right] \right)_{\theta=\omega(I)t+\tau} dt \\ &= \sum_{l=1}^q \rho_{jl}(I) \left( \int_0^{T^I} \frac{\partial h}{\partial \theta_l}(I, \omega(I)t + \tau; 0) dt \right) = \sum_{l=1}^q \rho_{jl}(I) \frac{d\mathcal{S}_F^I}{d\tau_l}(\tau). \end{aligned}$$

Since  $Z_1^0, \dots, Z_q^0$  are linearly independent on  $\mathcal{T}_I$ , the  $q \times q$  matrix  $(\rho_{jl})_{j,l=1,\dots,q}$  is invertible, so that by  $n - q \leq q$ ,  $(d\mathcal{S}_F^I/d\tau)(\tau) = O$ , i.e.,  $\mathcal{S}_F^I(\tau)$  is constant.

On the other hand, by Lemma 3.1, there exists an analytic map  $\psi : F(\mathcal{U}) \rightarrow \mathbb{R}^k$  such that  $I = \psi \circ F(x)$ . As in the proof of Lemma 3.2, we show that  $\text{rank } d\psi_{F(\gamma^I(0))} = n - q$  and  $\mathcal{S}_F^I(\tau) = d\psi_{F(\gamma^I(0))} \mathcal{S}_F^I(\tau)$ . Hence,  $\mathcal{S}_F^I(\tau)$  is also constant.  $\square$

*Proof of Theorem 2.4.* Suppose that  $\mathcal{S}_F^I(\tau)$  is not constant for any  $I \in D$ . From Theorem 2.2 we see that there exist only  $n - q - 1$  first integrals at most such that they are functionally independent and depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ .

Assume that there exist  $q$  analytic commutative vector fields  $Z_1^\varepsilon, \dots, Z_q^\varepsilon$  such that for  $\varepsilon = 0$  they are linearly independent almost everywhere and depend analytically on  $\varepsilon$ . Applying Lemma 3.6 for each  $I \in D$ , we see that the perturbed system (1.4) does not have  $q$  analytic commutative vector fields near  $\mathcal{T}_I$  such that they are linearly independent on  $\mathcal{T}_I$  and depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ . Hence,  $Z_1^\varepsilon, \dots, Z_q^\varepsilon$  are linearly dependent on  $\mathcal{T}_I$  for  $I \in D$  at  $\varepsilon = 0$  if they depend analytically on  $\varepsilon$  near  $\varepsilon = 0$ . As in the proof of Lemma 3.6, we consider the transformed system (1.2) and use Lemma 3.5 to write  $Z_j^0(I, \theta)$  in the form (3.1) with  $\rho_l(I) = \rho_{jl}(I)$ ,  $l = 1, \dots, q$ , for  $j = 1, \dots, q$ . We see that the determinant of the matrix  $(\rho_{jl}(I))_{j,l=1,\dots,q}$  is zero for  $I \in D$ , so that it is identically zero on  $U$  since  $D$  is a key set for  $C^\omega(U)$ . This yields a contradiction. So we obtain the desired result.  $\square$

#### 4. CONSEQUENCES OF THE THEORY TO (1.1)

In this section, we consider nearly integrable systems of the form (1.1) written in the action-angle coordinates and describe consequences of Theorems 2.2 and 2.4 to it. The unperturbed system (1.2) is  $(m, \ell)$ -integrable in the Bogoyavlenskij sense and has  $\ell$  first integrals  $I_1, \dots, I_\ell$  and  $m$  commutative vector fields  $\omega(I) \frac{\partial}{\partial \theta_1}, \frac{\partial}{\partial \theta_2}, \dots, \frac{\partial}{\partial \theta_m}$ . Thus, conditions (A1) and (A2) with  $n = \ell + m$  and  $q = m$  already hold. In particular, the level set of  $I = c$  given by  $\{c\} \times \mathbb{T}^m$  is connected and compact. Take some  $I_0 \in \mathbb{R}^\ell$  and let  $U$  be its neighborhood in  $\mathbb{R}^\ell$ , as in the preceding sections.

We first discuss consequences of Theorem 2.2 to (1.1) and assume that conditions (K2) and (A3) with  $n = \ell + m$  and  $q = m$  hold. For  $I \in D_{\mathbb{R}}$  the unperturbed system (1.2) has an  $m$ -parameter family of periodic orbits given by (2.1) with  $q = m$ . The integrals given by (2.2) for the  $\ell$  first integrals  $I = (I_1, \dots, I_\ell)$  become

$$\mathcal{S}_I^I(\tau) = \int_0^{T^I} h(I, \omega(I)t + \tau; 0) dt,$$

where  $\tau \in \mathbb{T}^m$ .

Assume that  $m > 1$ . Using the Fourier expansion of  $h(I, \theta; 0)$  given in (1.3), we rewrite the above integral as

$$\mathcal{S}_I^I(\tau) = \int_0^{T^I} \sum_{r \in \mathbb{Z}^m} \hat{h}_r(I) \exp(ir \cdot (\omega(I)t + \tau)) dt = T^I \sum_{r \in \Lambda_I} \hat{h}_r(I) e^{ir \cdot \tau}, \quad (4.1)$$

where  $\Lambda_I = \{r \in \mathbb{Z}^m \mid r \cdot \omega(I) = 0\}$ . Applying Theorem 2.2, we obtain the following result for (1.1). Recall that  $\hat{h}_r(I)$ ,  $r \in \mathbb{Z}^m$ , represent the Fourier coefficients of  $h(I, \theta; 0)$  (see Eq. (1.3)).

**Theorem 4.1.** *Let  $m > 1$ , and suppose that assumptions (K2) and (A3) with  $n = \ell + m$  and  $q = m$  hold. If there exists a key set  $D \subset D_{\mathbb{R}}$  for  $C^\omega(U)$  such that  $\hat{h}_r(I) \neq 0$  for some  $r \in \Lambda_I$  with  $I \in D$ , then the perturbed system (1.1) does not have  $\ell$  real-analytic first integrals in a neighborhood of the level set  $\{c\} \times \mathbb{T}^m$  near  $\varepsilon = 0$  such that they are functionally independent for  $|\varepsilon| \neq 0$  and depend analytically on  $\varepsilon$ .*

**Remark 4.2.**

- (i) From the proof given in [14] we see that the conclusion of Theorem 1.1 also holds even if the zero vector is taken as one of  $r_j \in \mathbb{Z}^m$ ,  $j = 1, \dots, \ell - s$ , in the definition of a Poincaré set. This fact was overlooked in [14].
- (ii) If a Poincaré set  $\mathcal{P}_{\ell-1} \subset U$  modified as stated in part (i) is a key set for  $C^\omega(U)$ , then condition (A3) holds. Moreover, there exists a key set  $D \subset D_{\mathbb{R}}$  for  $C^\omega(U)$  such that  $\hat{h}_r(I) \neq 0$  with some  $r \in \Lambda_I$  for  $I \in D$  if and only if such a Poincaré set  $\mathcal{P}_{\ell-1} \subset U$  is a key set for  $C^\omega(U)$ .
- (iii) The hypothesis of Theorem 4.1 holds if both of  $\omega(I)$  and  $\hat{h}_0(I)$  are not identically zero in  $U$ .
- (iv) If the system (1.1) is Hamiltonian, then  $\hat{h}_0(I) \equiv 0$ .
- (v) From Theorem 2.2 of [20] and Eq. (4.1) we see that the first integrals  $I_1, \dots, I_m$  do not persist in (1.1) near the resonant torus  $\{I\} \times \mathbb{T}^m$  if  $\hat{h}_r(I) \neq 0$  for some  $r \in \Lambda_I$ .

Let  $m = 1$  and assume that  $\omega(I) \neq 0$ . Then the integral (4.1) becomes

$$\mathcal{I}_I^I(\tau) = \int_0^{2\pi/\omega(I)} h(I, \omega(I)t + \tau; 0) dt = \frac{2\pi\hat{h}_0(I)}{\omega(I)} \quad (4.2)$$

since  $\Lambda_I = \{0\} \subset \mathbb{Z}$ . Noting that assumptions (K2) and (A3) hold if  $\omega(I) \neq 0$  for some  $I \in U$ , we obtain the following.

**Theorem 4.3.** *Let  $m = 1$ . If  $\omega(I) \neq 0$  and  $\hat{h}_0(I) \neq 0$  for some  $I \in U$ , then the perturbed system (1.1) does not have  $\ell$  real-analytic first integrals in a neighborhood of  $\{c\} \times \mathbb{S}^1$  near  $\varepsilon = 0$  such that they are functionally independent for  $|\varepsilon| \neq 0$  and depend analytically on  $\varepsilon$ .*

Assuming the existence of  $\ell - 1$  functionally independent first integrals in the perturbed system (1.1) and taking Remarks 4.2(i) and (ii) into account, we obtain the same result as Theorems 4.1 and 4.3 from Theorem 1.1 (see also Remark 3.4). Moreover, when the existence of such only  $s$  ( $< \ell - 1$ ) first integrals is assumed, Theorem 1.1 guarantees the nonexistence of no additional first integral if a Poincaré set  $\mathcal{P}_s$  modified in Remark 4.2(i) is a key set for  $C^\omega(U)$ , in particular  $\hat{h}_{r_j}(I)$ ,  $j = 1, \dots, \ell - s$  ( $> 1$ ), are linearly independent. Note that such a Poincaré set  $\mathcal{P}_s$  does not exist when  $m = 1$  and  $s < \ell - 1$ .

We next apply Theorem 2.4 to (1.1). When  $m = 1$ , the integral  $\mathcal{I}_I^I(\tau)$  is constant by (4.2), so that Theorem 2.4 does not apply. Thus, we obtain the following result.

**Theorem 4.4.** *Let  $m > 1$ , and suppose that assumptions (K2), (A3) and (A4) hold. If there exists a key set  $D \subset D_{\mathbb{R}}$  for  $C^\omega(U)$  such that  $\hat{h}_r(I) \neq 0$  for some  $r \in \Lambda_I \setminus \{0\}$  with  $I \in D$ , then for  $|\varepsilon| \neq 0$  sufficiently small the perturbed system (1.1) is not real-analytically integrable in the meaning of Theorem 2.4 near  $\{c\} \times \mathbb{T}^m$ .*

## 5. RELATIONSHIPS WITH THE MELNIKOV METHODS

In this section, we discuss relationships of our main results in Section 2 with the subharmonic and homoclinic Melnikov methods for time-periodic perturbations of single-degree-of-freedom Hamiltonian systems. See [11, 15, 25, 27] for the details of the Melnikov methods. A concise review of the methods was also given in Section 4.1 of [20].

Consider systems of the form

$$\dot{x} = JDH(x) + \varepsilon u(x, \nu t), \quad x \in \mathbb{R}^2, \quad (5.1)$$

where  $\varepsilon$  is a small parameter as in the preceding sections,  $\nu > 0$  is a constant,  $H : \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $u : \mathbb{R}^2 \times \mathbb{S} \rightarrow \mathbb{R}^2$  are analytic, and  $J$  is the  $2 \times 2$  symplectic matrix,

$$J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Equation (5.1) represents a time-periodic perturbation of the single-degree-of-freedom Hamiltonian system

$$\dot{x} = JDH(x), \quad (5.2)$$

with the Hamiltonian  $H(x)$ . Letting  $\phi = \nu t \pmod{2\pi}$  such that  $\phi \in \mathbb{S}^1$ , we rewrite (5.1) as an autonomous system,

$$\dot{x} = JDH(x) + \varepsilon u(x, \phi), \quad \dot{\phi} = \nu. \quad (5.3)$$

We easily see that assumptions (A1) and (A2) hold in (5.3) with  $\varepsilon = 0$ :  $H(x)$  is a first integral and  $(0, 1) \in \mathbb{R}^2 \times \mathbb{R}$  is a commutative vector field. We make the following assumptions on the unperturbed system (5.2):

**(M1)** There exists a one-parameter family of periodic orbits  $x^\alpha(t)$  with period  $\hat{T}^\alpha > 0$ ,  $\alpha \in (\alpha_1, \alpha_2)$ , for some  $\alpha_1 < \alpha_2$ . Moreover,  $\hat{T}^\alpha$  is not constant as a function of  $\alpha$ .

**(M2)**  $x^\alpha(t)$  is analytic with respect to  $\alpha \in (\alpha_1, \alpha_2)$ .

Note that in (M1)  $x^\alpha(t)$  is automatically analytic with respect to  $t$  since the vector field of (5.2) is analytic.

We assume that at  $\alpha = \alpha^{l/n}$

$$\frac{2\pi}{\hat{T}^\alpha} = \frac{n}{l}\nu, \quad (5.4)$$

where  $l$  and  $n$  are relatively prime integers. We define the *subharmonic Melnikov function* as

$$M^{l/n}(\phi) = \int_0^{2\pi l/\nu} DH(x^\alpha(t)) \cdot u(x^\alpha(t), \nu t + \phi) dt, \quad (5.5)$$

where  $\alpha = \alpha^{l/n}$ . Let  $T^\alpha = n\hat{T}^\alpha = 2\pi l/\nu$  for  $\alpha = \alpha^{l/n}$ . If  $M^{l/n}(\phi)$  has a simple zero at  $\phi = \phi_0$  and  $d\hat{T}^\alpha/d\alpha \neq 0$  at  $\alpha = \alpha^{l/n}$ , then for  $|\varepsilon| > 0$  sufficiently small there exists a  $T^\alpha$ -periodic orbit near  $(x, \phi) = (x^\alpha(t), \nu t + \phi_0)$  in (5.3). See Theorem 3.1 of [27]. A similar result is also found in [11, 25]. The stability of the periodic orbit can also be determined easily [27]. Moreover, several bifurcations of periodic orbits when  $d\hat{T}^\alpha/d\alpha \neq 0$  or not were discussed in [27–29].

On the other hand, since it is a single-degree-of-freedom Hamiltonian system, the unperturbed system (5.2) is integrable, so that it can be transformed into the form (1.2) with  $\ell, m = 1$ . So the perturbed system (5.3) is transformed into the form (1.1) with  $\ell = 1$  and  $m = 2$ . Here we take  $I = \alpha$  unlike [27, 30], and have  $\omega(I) = (\Omega(I), \nu)$ , where

$$\Omega(\alpha) = \frac{2\pi}{\hat{T}^\alpha}.$$

We remark that the transformed system is not Hamiltonian even when  $\varepsilon = 0$ , unlike [27, 30]. Choose a point  $\alpha = \alpha_0 \in (\alpha_1, \alpha_2)$  such that  $d\hat{T}^\alpha/d\alpha \neq 0$ , and let  $U$

be a neighborhood of  $\alpha_0$ . We see that assumptions (K2) and (A3) hold for

$$D_R = \{\alpha^{l/n} \mid l, n \in \mathbb{N}\} \cap U.$$

Let  $\alpha = \alpha^{l/n}$  and let  $\gamma_\tau^\alpha(t) = (x^\alpha(t + \tau_1), \nu(t + \tau_1) + \tau_2)$ . We see that  $\gamma_\tau^\alpha(t)$  is a  $T^\alpha$ -periodic orbit in (5.3) with  $\varepsilon = 0$ . Note that  $\gamma_\tau^\alpha(t)$  is essentially parameterized by a single parameter, say  $\phi := \nu\tau_1 + \tau_2$ . So we write  $\gamma_\phi^\alpha(t) = (x^\alpha(t), \nu t + \phi)$ . The integral (2.2) for  $H(x)$  along  $\gamma_\phi^\alpha(t)$  becomes

$$\mathcal{I}_H^\alpha(\phi) = \int_0^{2\pi l/\nu} DH(x^\alpha(t)) \cdot u(x^\alpha(t), \nu t + \phi) dt = M^{l/n}(\phi) \quad (5.6)$$

by (5.5). As stated above, if  $M^{l/n}(\phi)$  has a simple zero at  $\phi = \phi_0$ , then there exists a  $T^\alpha$ -periodic orbit near  $\gamma_{\phi_0}^\alpha(t)$ . Applying Theorems 2.2 and 2.4, we have the following two results.

**Theorem 5.1.** *Suppose that there exists a key set  $D \subset D_R$  for  $C^\omega(U)$  such that  $M^{l/n}(\phi)$  is not identically zero for  $\alpha^{l/n} \in D$ . Then for  $|\varepsilon| \neq 0$  sufficiently small the system (5.3) has no real-analytic first integral in a neighborhood of  $\{x^{\alpha_0}(t) \mid t \in [0, \hat{T}^{\alpha_0})\} \times \mathbb{S}^1$  such that it depends analytically on  $\varepsilon$  near  $\varepsilon = 0$ .*

**Theorem 5.2.** *Suppose that there exists a key set  $D \subset D_R$  for  $C^\omega(U)$  such that  $M^{l/n}(\phi)$  is not constant for  $\alpha^{l/n} \in D$ . Then for  $|\varepsilon| \neq 0$  sufficiently small the system (5.3) is not real-analytically integrable in the meaning of Theorem 2.4 in a neighborhood of  $\{x^{\alpha_0}(t) \mid t \in [0, \hat{T}^{\alpha_0})\} \times \mathbb{S}^1$ .*

**Remark 5.3.**

- (i) *If  $D$  has an accumulation point, then it becomes a key set for  $C^\omega(U)$ .*
- (ii) *If the system (5.1) is Hamiltonian, then the hypotheses of Theorems 5.1 and 5.2 are equivalent to the condition that  $M^{l/n}(\phi)$  has a simple zero. Actually, letting*

$$u(x, \phi) = JD_x H^1(x, \phi) = J \sum_{r \in \mathbb{Z}} D\hat{H}_r^1(x) e^{ir\phi},$$

we have

$$\begin{aligned} M^{l/n}(\phi) &= \int_0^{2\pi l/\nu} DH(x^\alpha(t)) \cdot JD_x H^1(x^\alpha(t), \nu t + \phi) dt \\ &= \sum_{r \in \mathbb{Z}} e^{ir\phi} \int_0^{2\pi l/\nu} DH(x^\alpha(t)) \cdot JD\hat{H}_r^1(x^\alpha(t)) e^{ir\nu t} dt \end{aligned}$$

and

$$\begin{aligned} &\int_0^{2\pi l/\nu} DH(x^\alpha(t)) \cdot JD\hat{H}_0^1(x^\alpha(t)) dt \\ &= - \int_0^{2\pi l/\nu} D\hat{H}_0^1(x^\alpha(t)) \cdot JDH(x^\alpha(t)) dt \\ &= - \int_0^{2\pi l/\nu} D\hat{H}_0^1(x^\alpha(t)) \cdot \dot{x}^\alpha(t) dt = 0, \end{aligned}$$

where  $\hat{H}_r^1(x)$ ,  $r \in \mathbb{Z}$ , represent the Fourier coefficients of  $H^1(x, \phi)$ . Thus, we obtain the claim.

We additionally assume the following on the unperturbed system (5.2):




FIGURE 1. Assumption (M3).

(M3) There exists a hyperbolic saddle  $x_0$  with a homoclinic orbit  $x^h(t)$  such that

$$\lim_{\alpha \rightarrow \alpha_2} \sup_{t \in \mathbb{R}} d(x^\alpha(t), \Gamma) = 0,$$

where  $\Gamma = \{x^h(t) \mid t \in \mathbb{R}\} \cup \{x_0\}$  and  $d(x, \Gamma) = \inf_{y \in \Gamma} |x - y|$ . See Fig. 1.

We define the *homoclinic Melnikov function* as

$$M(\phi) = \int_{-\infty}^{\infty} DH(x^h(t)) \cdot u(x^h(t), t + \phi) dt. \quad (5.7)$$

If  $M(\phi)$  has a simple zero, then for  $|\varepsilon| > 0$  sufficiently small there exist transverse homoclinic orbits to a periodic orbit near  $\{x_0\} \times \mathbb{S}^1$  in (5.3) [11, 15, 25]. The existence of such transverse homoclinic orbits implies that the system (5.3) exhibits chaotic motions by the Smale-Birkhoff theorem [11, 25] and has no real-analytic (additional) first integral (see, e.g., Chapter III of [19]). We easily show that

$$\lim_{l \rightarrow \infty} M^{l/1}(\phi) = M(\phi) \quad (5.8)$$

for each  $\phi \in \mathbb{S}^1$  (see Theorem 4.6.4 of [11]). Let  $U$  be a neighborhood of  $\alpha = \alpha_2$ . It follows from (5.8) that if  $M(\phi)$  is not identically zero or constant, then for  $l > 0$  sufficiently large neither is  $M^{l/1}(\phi)$ . Let  $\hat{U} \subset \mathbb{R}^2$  be a region such that  $\partial \hat{U} \supset \Gamma$  and  $\hat{U} \supset \{x^\alpha(t) \mid t \in [0, \hat{T}^\alpha]\}$  for some  $\alpha \in (\alpha_1, \alpha_2)$ . We obtain the following from Theorems 5.1 and 5.2.

**Theorem 5.4.** *Suppose that  $M(\phi)$  is not identically zero. Then for  $|\varepsilon| \neq 0$  sufficiently small the system (5.3) has no real-analytic first integral in  $\hat{U} \times \mathbb{S}^1$  such that it depends analytically on  $\varepsilon$  near  $\varepsilon = 0$ .*

**Theorem 5.5.** *Suppose that  $M(\phi)$  is not constant. Then for  $|\varepsilon| \neq 0$  sufficiently small the system (5.3) is not real-analytically integrable in the meaning of Theorem 2.4 in  $\hat{U} \times \mathbb{S}^1$ .*

**Remark 5.6.**

- (i) *Theorems 5.4 and 5.5, respectively, mean that the system (5.3) has no first integral and is nonintegrable even if the Melnikov function  $M(\phi)$  does not have a simple zero, i.e., there may exist no transverse homoclinic orbit to the periodic orbit in (5.3), but it is not identically zero and constant. See Section 6.3.*
- (ii) *As in Remark 5.3(ii), if the system (5.1) is Hamiltonian, then the hypotheses of Theorems 5.4 and 5.5 are equivalent to the condition that  $M^{l/n}(\phi)$  has a simple zero.*

- (iii) *In the statements of Theorems 5.4 and 5.5, the region  $\hat{U} \times \mathbb{S}^1$  may be replaced with a neighborhood of  $\Gamma \times \mathbb{S}^1$  although they are weakened.*

## 6. EXAMPLES

We now illustrate the above theory for four examples: Simple pendulum with a constant torque, second-order coupled oscillators and the periodically forced Duffing oscillator [11, 12, 25].

**6.1. Simple pendulum with a constant torque.** Consider a simple pendulum with a constant torque:

$$\dot{I} = \varepsilon(\beta \sin \theta + 1), \quad \dot{\theta} = I, \quad (I, \theta) \in \mathbb{R} \times \mathbb{T} \quad (6.1)$$

where  $\beta \in \mathbb{R}$  is a constant. Equation (6.1) is of the form (1.1) with  $m = \ell = 1$ . Using Theorem 4.3, we obtain the following.

**Proposition 6.1.** *The system (6.1) has no real-analytic first integral depending analytically on  $\varepsilon$  near  $\varepsilon = 0$ .*

**Remark 6.2.**

- (i) *The system (6.1) has the first integral*

$$F(I, \theta; \varepsilon) = \frac{1}{2}I^2 + \varepsilon(\beta \cos \theta - \theta)$$

*and is (1, 1)-integrable as a system on  $\mathbb{R} \times \mathbb{R}$ , although  $F(I, \theta; \varepsilon)$  is not even a function on  $\mathbb{R} \times \mathbb{S}^1$ .*

- (ii) *Let  $\beta = 0$ . Then the system (6.1) is (2, 0)-integrable when  $\varepsilon \neq 0$ , where the vector fields  $\varepsilon \frac{\partial}{\partial I} + I \frac{\partial}{\partial \theta}$  and  $\frac{\partial}{\partial \theta}$  are commutative and linearly independent. However, when  $\varepsilon = 0$ , the two vector fields are linearly dependent and Eq. (6.1) is not (2, 0)-integrable. See also Remark 2.5(ii).*

**6.2. Second-order coupled oscillators.** Consider

$$\begin{aligned} \dot{I}_j &= \varepsilon \left( -\delta I_j + \Omega_j + \sum_{i=1}^{\ell} \sum_{k \in \mathbb{N}^2} a_k \sin(k_1 \theta_j - k_2 \theta_i) \right), \\ \dot{\theta}_j &= I_j, \quad j = 1, \dots, \ell, \end{aligned} \quad (6.2)$$

where  $\delta, \Omega_j \geq 0$ ,  $j = 1, \dots, \ell$ , and  $a_k$ ,  $k = (k_1, k_2) \in \mathbb{N}^2$ , are constants such that  $|a_k| \leq M e^{-(k_1 + k_2)\delta}$  for some  $M, \delta > 0$ . We see by the remark after Lemma 2 in Section 12 of Chapter 3 in [1] that the vector field of (6.2) is analytic. Equation (6.2) has the form (1.1) with  $m = \ell$  and is rewritten in a system of second-order differential equations as

$$\ddot{\theta}_j + \varepsilon \delta \dot{\theta}_j = \varepsilon \left( \Omega_j + \sum_{i=1}^{\ell} \sum_{k \in \mathbb{N}^2} a_k \sin(k_1 \theta_j - k_2 \theta_i) \right), \quad j = 1, \dots, \ell,$$

which reduces to the *second-order Kuramoto model* [24] when  $a_k \neq 0$  for  $k = (1, 1)$  and  $a_k = 0$  for  $k \neq (1, 1)$ . Obviously, assumptions (K2), (A3) and (A4) hold. Using Theorems 4.1 and 4.4, we obtain the following.

**Proposition 6.3.** *The following statements hold for (6.2):*




FIGURE 2. Phase portraits of (6.3) with  $\varepsilon = 0$ : (a)  $a = 1$ ; (b)  $a = -1$ .

- (i) If one of  $\delta$  and  $\Omega_j$ ,  $j = 1, \dots, \ell$ , is nonzero at least, then the system (6.2) does not have  $\ell$  real-analytic first integrals near  $\varepsilon = 0$  such that they are functionally independent for  $|\varepsilon| \neq 0$  and depend analytically on  $\varepsilon$ ;
- (ii) If  $K_1 = \{k_1/k_2 \mid a_k, k_2 \neq 0\}$  or  $K_2 = \{k_2/k_1 \mid a_k, k_1 \neq 0\}$  has an accumulation point, then for  $|\varepsilon| \neq 0$  sufficiently small the system (6.2) is not real-analytically integrable in the meaning of Theorem 2.4.

*Proof.* Part (i) immediately follows from Theorem 4.1 and Remark 4.2(iii) since  $\hat{h}_0(I)$  is not identically zero if one of  $\delta$  and  $\Omega_j$ ,  $j = 1, \dots, \ell$ , is nonzero at least.

We turn to the proof of part (ii). Let  $D_1 = \{I \in \mathbb{R}^\ell \mid k_1 I_j - k_2 I_i = 0, i, j = 1, \dots, \ell, k_1/k_2 \in K_1\}$  and  $D_2 = \{I \in \mathbb{R}^\ell \mid k_1 I_j - k_2 I_i = 0, i, j = 1, \dots, \ell, k_2/k_1 \in K_2\}$ . If  $K_1$  (resp.  $K_2$ ) has an accumulation point, then  $D_1$  (resp.  $D_2$ ) is a key set for  $C^\omega(\mathbb{R}^\ell)$ . Their claim is shown as follows. Assume that  $K_1$  has an accumulation point. Let  $f(I) \in C^\omega(\mathbb{R}^\ell)$  be an analytic function which vanishes on  $D_1$ , and take a line  $L_b := \{I \in \mathbb{R}^\ell \mid (I_1, \dots, I_{\ell-1}) = b\}$  for  $b = (b_1, \dots, b_{\ell-1}) \in \mathbb{R}^{\ell-1}$  fixed. Then  $(b_1, \dots, b_{\ell-1}, k_1 b_1/k_2) \in L_b \cap D_1$  for all  $k_1/k_2 \in K_1$ , so that  $f(I)$  is identically zero on  $L_b$ . This means that  $f(I)$  is identically zero in  $\mathbb{R}^\ell$ , and consequently  $D_1$  is a key set for  $C^\omega(\mathbb{R}^\ell)$ . Similarly, we see that the claim is true for  $K_2$  and  $D_2$ . Applying Theorem 4.4, we obtain the desired result.  $\square$

**6.3. Periodically forced Duffing oscillator.** Consider the periodically forced Duffing oscillator

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = ax_1 - x_1^3 + \varepsilon(\beta \cos \nu t - \delta x_2), \quad (6.3)$$

where  $\nu > 0$  and  $\beta, \delta \geq 0$  are constants, and  $a = -1$  or  $1$ . The system (6.3) has the form (5.1) with

$$H = -\frac{1}{2}ax_1^2 + \frac{1}{4}x_1^4 + \frac{1}{2}x_2^2$$

and the autonomous system (5.3) becomes

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = ax_1 - x_1^3 + \varepsilon(\beta \cos \theta - \delta x_2), \quad \dot{\phi} = \nu, \quad (6.4)$$

where  $(x, \phi) \in \mathbb{R}^2 \times \mathbb{S}$ . See Fig. 2 for the phase portraits of (6.3) with  $\varepsilon = 0$ .



We begin with the case of  $a = 1$ . When  $\varepsilon = 0$ , in the phase plane there exist a pair of homoclinic orbits

$$x_{\pm}^h(t) = (\pm\sqrt{2} \operatorname{sech} t, \mp\sqrt{2} \operatorname{sech} t \tanh t),$$

a pair of one-parameter families of periodic orbits

$$x_{\pm}^k(t) = \left( \pm \frac{\sqrt{2}}{\sqrt{2-k^2}} \operatorname{dn} \left( \frac{t}{\sqrt{2-k^2}} \right), \right. \\ \left. \mp \frac{\sqrt{2}k^2}{2-k^2} \operatorname{sn} \left( \frac{t}{\sqrt{2-k^2}} \right) \operatorname{cn} \left( \frac{t}{\sqrt{2-k^2}} \right) \right), \quad k \in (0, 1),$$

inside each of them, and a one-parameter periodic orbits

$$\tilde{x}^k(t) = \left( \frac{\sqrt{2}k}{\sqrt{2k^2-1}} \operatorname{cn} \left( \frac{t}{\sqrt{2k^2-1}} \right), \right. \\ \left. - \frac{\sqrt{2}k}{2k^2-1} \operatorname{sn} \left( \frac{t}{\sqrt{2k^2-1}} \right) \operatorname{dn} \left( \frac{t}{\sqrt{2k^2-1}} \right) \right), \quad k \in (1/\sqrt{2}, 1),$$

outside of them, as shown in Fig. 2(a), where  $\operatorname{sn}$ ,  $\operatorname{cn}$  and  $\operatorname{dn}$  represent the Jacobi elliptic functions with the elliptic modulus  $k$ . The periods of  $x_{\pm}^k(t)$  and  $\tilde{x}^k(t)$  are given by  $\hat{T}^k = 2K(k)\sqrt{2-k^2}$  and  $\tilde{T}^k = 4K(k)\sqrt{2k^2-1}$ , respectively, where  $K(k)$  is the complete elliptic integral of the first kind. Note that  $x_{\pm}^k(t)$  approaches  $x_{\pm}^h(t)$  as  $k \rightarrow 1$ . See [11, 25]. See also [9] for general information on elliptic functions.

Assume that the resonance conditions

$$l\hat{T}^k = \frac{2\pi n}{\nu}, \quad \text{i.e.,} \quad \nu = \frac{2\pi n}{2lK(k)\sqrt{2-k^2}}, \quad (6.5)$$

and

$$l\tilde{T}^k = \frac{2\pi n}{\nu}, \quad \text{i.e.,} \quad \nu = \frac{2\pi n}{4lK(k)\sqrt{2k^2-1}}, \quad (6.6)$$

hold for  $x_{\pm}^k(t)$  and  $\tilde{x}^k(t)$ , respectively, with  $l, n > 0$  relatively prime integers. Then the subharmonic Melnikov function (5.5) for  $x_{\pm}^k(t)$  and  $\tilde{x}^k(t)$  are

$$M_{\pm}^{n/l}(\tau) = -\delta J_1(k, l) \pm \beta J_2(k, n, l) \sin \tau$$

and

$$\tilde{M}^{n/l}(\tau) = -\delta \tilde{J}_1(k, l) + \beta \tilde{J}_2(k, n, l) \sin \tau,$$

respectively, where

$$J_1(k, l) = \frac{4l[(2-k^2)E(k) - 2k'^2K(k)]}{3(2-k^2)^{3/2}}, \\ J_2(k, n, l) = \begin{cases} \sqrt{2}\pi\nu \operatorname{sech} \left( \frac{n\pi K(k')}{K(k)} \right) & (\text{for } l = 1); \\ 0 & (\text{for } l \neq 1), \end{cases} \\ \tilde{J}_1(k, l) = \frac{8l[(2k^2-1)E(k) + k'^2K(k)]}{3(2k^2-1)^{3/2}}, \\ \tilde{J}_2(k, n, l) = \begin{cases} 2\sqrt{2}\pi\nu \operatorname{sech} \left( \frac{n\pi K(k')}{2K(k)} \right) & (\text{for } l = 1 \text{ and } n \text{ odd}); \\ 0 & (\text{for } l \neq 1 \text{ or } n \text{ even}). \end{cases}$$

Here  $E(k)$  is the complete elliptic integral of the second kind and  $k' = \sqrt{1-k^2}$  is the complimentary elliptic modulus. When  $\delta \neq 0$ , the subharmonic Melnikov functions  $M_{\pm}^{n/l}(\tau)$  and  $\tilde{M}^{n/l}(\tau)$  are not identically zero for any relatively prime integers  $n, l > 0$  since  $J_1(k, l)$  and  $\tilde{J}_1(k, l)$  are not zero. Moreover, the homoclinic Melnikov function (5.7) for  $x_{\pm}^h(t)$  is

$$M_{\pm}(\tau) = -\frac{4}{3}\delta \pm \sqrt{2}\pi\nu\beta \operatorname{csch}\left(\frac{\pi\nu}{2}\right) \sin \tau,$$

which is not identically zero for  $\beta \neq 0$ . See [11, 25] for the computations of the Melnikov functions.

Let

$$R = \{k \in (0, 1) \mid k \text{ satisfies (6.5) for } n, l \in \mathbb{N}\},$$

$$\tilde{R} = \{k \in (1/\sqrt{2}, 1) \mid k \text{ satisfies (6.6) for } n, l \in \mathbb{N}\},$$

and let

$$S_{\pm}^k = \{(x_{\pm}^k(t), \theta) \in \mathbb{R}^2 \times \mathbb{S}^1 \mid t \in [0, \hat{T}^k], \theta \in \mathbb{S}^1\},$$

$$\tilde{S}^k = \{(\tilde{x}^k(t), \theta) \in \mathbb{R}^2 \times \mathbb{S}^1 \mid t \in [0, \tilde{T}^k], \theta \in \mathbb{S}^1\},$$

$$\Gamma_{\pm} = \{x_{\pm}^h(t) \in \mathbb{R}^2 \mid t \in \mathbb{R}\} \cup \{0\}.$$

Noting that

$$\lim_{n \rightarrow \infty} \tilde{M}^{2n+1/1}(\tau) = M_+(\tau) + M_-(\tau)$$

and applying Theorems 5.1, 5.4, 5.5 and their slight extensions, we have the following.

**Proposition 6.4.** *The system (6.4) with  $a = 1$  has no real-analytic first integral depending analytically on  $\varepsilon$  in neighborhoods of  $S_{\pm}^k$  for  $k \in R$ , of  $\tilde{S}^k$  for  $k \in \tilde{R}$ , and of  $S_{\pm}^h$  near  $\varepsilon = 0$  if  $\delta \neq 0$ .*

**Proposition 6.5.** *Let  $\hat{U}_{\pm}$  (resp.  $\tilde{U}$ ) be regions (resp. a region) in  $\mathbb{R}^2$  such that  $\partial\hat{U}_{\pm} \supset \Gamma_{\pm}$  (resp.  $\partial\tilde{U} \supset \Gamma_+ \cup \Gamma_-$ ) and  $\hat{U}_{\pm} \supset \{x_{\pm}^k(t) \mid t \in [0, \hat{T}^k]\}$  (resp.  $\tilde{U} \supset \{\tilde{x}^k(t) \mid t \in [0, \tilde{T}^k]\}$ ) for some  $k \in (0, 1)$  (resp.  $k \in (1/\sqrt{2}, 1)$ ). For  $|\varepsilon| \neq 0$  sufficiently small the system (6.4) with  $a = 1$  is not real-analytically integrable in the regions  $\hat{U}_{\pm} \times \mathbb{S}^1$  (resp.  $\tilde{U} \times \mathbb{S}^1$ ) in the meaning of Theorem 2.4 if  $\beta \neq 0$ .*

If  $\beta \neq 0$  and

$$\frac{\delta}{\beta} < \frac{3}{4}\sqrt{2}\pi\nu \operatorname{csch}\left(\frac{\pi\nu}{2}\right), \quad (6.7)$$

then  $M_{\pm}(\tau)$  has a simple zero, so that for  $|\varepsilon| > 0$  sufficiently small there exist transverse homoclinic orbits to a periodic orbit near the origin and chaotic dynamics may occur in (6.4) with  $a = 1$ , as stated in Section 5. From Proposition 6.5 we see that the system (6.4) is nonintegrable in the meaning of Theorem 2.4 even if condition (6.7) does not hold, i.e., there may exist no transverse homoclinic orbit to the periodic orbit, as stated in Remark 5.6(i). On the other hand, when the system (6.3) is Hamiltonian, i.e.,  $\delta = 0$ , condition (6.7) always holds and such inconsistency does not occur. See also Remark 5.6(ii).

We turn to the case of  $a = -1$ . When  $\varepsilon = 0$ , in the phase plane there exists a one-parameter family of periodic orbits

$$\hat{x}^k(t) = \left( \frac{\sqrt{2}k}{\sqrt{1-2k^2}} \operatorname{cn} \left( \frac{t}{\sqrt{1-2k^2}} \right), -\frac{\sqrt{2}k}{1-2k^2} \operatorname{sn} \left( \frac{t}{\sqrt{1-2k^2}} \right) \operatorname{dn} \left( \frac{t}{\sqrt{1-2k^2}} \right) \right), \quad k \in (0, 1/\sqrt{2}),$$

as shown in Fig. 2(b), and their period is given by  $\hat{T}^k = 4K(k)\sqrt{1-2k^2}$ . See [26, 27]. Assume that the resonance conditions

$$l\hat{T}^k = \frac{2\pi n}{\nu}, \quad \text{i.e.,} \quad \nu = \frac{\pi n}{2lK(k)\sqrt{1-2k^2}} \quad (6.8)$$

holds for  $l, n > 0$  relatively prime integers. We compute the subharmonic Melnikov function (5.5) for  $\hat{x}^k(t)$  as

$$\hat{M}^{n/l}(\tau) = -\delta \hat{J}_1(k, l) \pm \beta \hat{J}_2(k, n, l) \sin \tau,$$

where

$$\hat{J}_1(k, l) = \frac{8l[(2k^2-1)E(k) + k'^2K(k)]}{3(1-2k^2)^{3/2}},$$

$$\hat{J}_2(k, n, l) = \begin{cases} \frac{\sqrt{2}\pi^2 n}{K(k)\sqrt{1-2k^2}} \operatorname{sech} \left( \frac{\pi n K(k')}{2K(k)} \right) & (\text{for } l = 1 \text{ and } n \text{ odd}); \\ 0 & (\text{for } l \neq 1 \text{ or } n \text{ even}). \end{cases}$$

See also [26, 27] for the computations of the Melnikov function. When  $\delta \neq 0$ , the subharmonic Melnikov function  $\hat{M}^{n/l}(\tau)$  is not identically zero for any relatively prime integers  $n, l > 0$  since  $J_1(k, l)$  is not zero.

Let

$$\hat{R} = \{k \in (0, 1/\sqrt{2}) \mid k \text{ satisfies (6.8) for } n, l \in \mathbb{N}\},$$

and let

$$\hat{S}^k = \{(\hat{x}^k(t), \theta) \in \mathbb{R}^2 \times \mathbb{S}^1 \mid t \in [0, \hat{T}^k], \theta \in \mathbb{S}^1\}.$$

Applying Theorem 5.1, we obtain the following.

**Proposition 6.6.** *The system (6.4) with  $a = -1$  has no real-analytic first integral depending analytically on  $\varepsilon$  in a neighborhood of  $\hat{S}^k$  for  $k \in \hat{R}$  near  $\varepsilon = 0$  if  $\delta \neq 0$ .*

**Remark 6.7.**

- (i) *Since the subharmonic Melnikov function  $\hat{M}^{n/l}(\tau)$  is constant for  $l \neq 1$ , Theorem 5.2 is not applicable to (6.4) with  $a = -1$ . So we cannot exclude the possibility that the system (6.4) with  $a = -1$  is  $(3, 0)$ -integrable in the meaning of Theorem 2.4 when  $\beta, \delta \neq 0$ .*
- (ii) *It was shown in [30] that the system (6.4) with  $a = -1$  is meromorphically nonintegrable in a meaning similar to that of Theorem 2.4 when the independent and state variables are extended to complex ones.*

#### ACKNOWLEDGEMENT

This work was partially supported by the JSPS KAKENHI Grant Numbers JP17H02859 and JP19J22791.

## APPENDIX A. PROOF OF PROPOSITION 3.3

In this Appendix, we prove Proposition 3.3. We begin with the following lemma.

**Lemma A.1.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^k$  and let  $\chi_j : \Omega \rightarrow \mathbb{R}$ ,  $j = 1, \dots, m$ , be analytic, where  $k, m \in \mathbb{N}$ . Let  $\chi(x) = (\chi_1(x), \dots, \chi_m(x))$ . If  $\text{rank } d\chi$  is constant on  $\Omega$  and less than  $m$ , then for any  $x \in \Omega$  there exists a neighborhood  $V$  of  $x$  on which  $\chi_1, \dots, \chi_m$  are analytically dependent, i.e., there exist an open set  $\Omega' \subset \mathbb{R}^m$  and a non-constant analytic map  $\zeta : \Omega' \rightarrow \mathbb{R}$  such that  $\chi(V) \subset \Omega'$  and  $\zeta(\chi(y)) = 0$  for any  $y \in V$ .*

*Proof.* Using Theorem 1.3.14 of [21] and an argument in the proof of Theorem 1.4.15 of [21], we can immediately obtain the desired result as follows. The theorem says that there exist a neighborhood  $V$  (resp.  $V'$ ) of  $x$  (resp. of  $\chi(x)$ ), a cube  $Q$  (resp.  $Q'$ ) in  $\mathbb{R}^k$  (resp. in  $\mathbb{R}^m$ ) and analytic isomorphisms  $u : Q \rightarrow V$  and  $u' : V' \rightarrow Q'$  such that the composite map  $u' \circ \chi \circ u$  has the form  $(x_1, \dots, x_k) \rightarrow (x_1, \dots, x_{m'}, 0, \dots, 0)$ , where  $x_j$  is the  $j$ th element of  $x$  for  $j = 1, \dots, k$  and  $m' = \text{rank } d\chi < m$ . Here a cube in  $\mathbb{R}^k$  is an open set of the form

$$\{x \mid |x_j - a_j| < r_j, j = 1, \dots, k\}$$

for some  $a_j \in \mathbb{R}$  and  $r_j > 0$ ,  $j = 1, \dots, k$ . Letting  $u' = (u'_1, \dots, u'_m)$  and  $\zeta = u'_m$ , we have  $\zeta(\chi(y)) = 0$  for every  $y \in V$ .  $\square$

Let  $f_\varepsilon : \mathcal{M} \rightarrow \mathbb{R}$  be an analytic function such that it depends on  $\varepsilon$  analytically. We expand it near  $\varepsilon = 0$  as  $f_\varepsilon(x) = \sum_{j=0}^{\infty} f^j(x)\varepsilon^j$ , where  $f^j(x)$ ,  $j \in \mathbb{Z}_0 := \mathbb{N} \cup \{0\}$ , are analytic functions on  $\mathcal{M}$ . Define the order function  $\sigma(f_\varepsilon)$  by

$$\sigma(f_\varepsilon) := \min\{j \in \mathbb{Z}_0 \mid f^j(x) \not\equiv 0\}$$

if  $f_\varepsilon \not\equiv 0$  and  $\sigma(0) := +\infty$ , as in [6].

**Lemma A.2.** *Suppose that  $f_\varepsilon(x)$  is a nonconstant analytic first integral of (1.4) depending analytically on  $\varepsilon$  near  $\varepsilon = 0$ . Then there exists an analytic first integral  $\tilde{f}_\varepsilon(x) = \tilde{f}^0(x) + O(\varepsilon)$  depending analytically on  $\varepsilon$  near  $\varepsilon = 0$  such that  $\tilde{f}^0(x)$  is not constant.*

*Proof.* Since  $f_\varepsilon$  is not constant,  $\sigma(df_\varepsilon)$  takes a finite value. Let  $k = \sigma(df_\varepsilon)$  and  $f_\varepsilon(x) = \sum_{j=0}^{\infty} \varepsilon^j f^j(x)$ . Define

$$\tilde{f}_\varepsilon(x) := \frac{1}{\varepsilon^k} \left( f_\varepsilon(x) - \sum_{j=0}^{k-1} \varepsilon^j f^j(x) \right).$$

Then  $\tilde{f}_\varepsilon(x) = f^k(x) + O(\varepsilon)$  and  $\tilde{f}^0(x) = f^k(x)$  is not constant. Moreover,  $\tilde{f}_\varepsilon(x)$  is a first integral of (1.4) since  $X_\varepsilon(f_\varepsilon) = 0$  and  $\sum_{j=0}^{k-1} \varepsilon^j f^j$  is constant.  $\square$

*Proof of Proposition 3.3.* Modifying the proof of Ziglin's lemma [4, 10, 32] slightly, we prove this proposition. For  $k = 1$  the statement of the proposition holds by Lemma A.2. Let  $k > 1$  and suppose that it is true up to  $k-1$ . Let  $G_1^\varepsilon(x), \dots, G_k^\varepsilon(x)$  be analytic first integrals of (1.4) in a neighborhood of  $F^{-1}(c)$  near  $\varepsilon = 0$  such that they are functionally independent for  $\varepsilon \neq 0$  and depend analytically on  $\varepsilon$ . Without loss of generality, we assume that  $G_1^0(x), \dots, G_{k-1}^0(x)$  are functionally independent near  $F^{-1}(c)$ . Letting  $G^\varepsilon(x) = (G_1^\varepsilon(x), \dots, G_k^\varepsilon(x))$ , we see that  $G^0(\varphi(I, \theta))$  depends

only on  $I$ , as in the proof of Lemma 3.1(i), where  $\varphi$  denotes the analytic diffeomorphism in Proposition 2.1(ii). Let  $\tilde{G}_j(I) = G_j^0(\varphi(I, \theta))$  for  $j = 1, \dots, k$ . Note that, if  $d\tilde{G}_1(I), \dots, d\tilde{G}_k(I)$  are linearly independent at  $I = I_0 \in U$ , then so are  $dG_1(x), \dots, dG_k(x)$  on  $\varphi(\{I_0\} \times \mathbb{T}^q) \subset \mathcal{U}$ .

Assume that  $\tilde{G}_1(I), \dots, \tilde{G}_{k-1}(I), \tilde{G}_k(I)$  are functionally dependent in an open set  $U' \subset U$ . So  $\Omega := \{p \in U' \mid \text{rank } d_p \tilde{G} = k - 1\}$  contains a dense open set in  $U'$  since  $d\tilde{G}_1(I), \dots, d\tilde{G}_{k-1}(I)$  are functionally independent on  $U$ . By Lemma A.1, there exist an open set  $\Omega' \subset \mathbb{R}^k$  and a nonzero analytic function  $\zeta : \Omega' \rightarrow \mathbb{R}$  such that  $\tilde{G}(V) = (\tilde{G}_1(V), \dots, \tilde{G}_k(V)) \subset \Omega'$  and

$$\zeta(\tilde{G}_1(I), \dots, \tilde{G}_k(I)) = 0$$

in a neighborhood  $V$  of  $p \in \Omega$ . Moreover, there is a positive integer  $s$  such that  $(\partial^s \zeta / \partial y_k^s)(\tilde{G}(I)) \neq 0$ , since if not, then  $\zeta(\tilde{G}_1(I), \dots, \tilde{G}_{k-1}(I), y_k)$  depends on  $y_k$  near  $\tilde{G}(V)$  and consequently  $\tilde{G}_1(I), \dots, \tilde{G}_{k-1}(I)$  are functionally dependent. Let  $s$  be the smallest one of such integers and let  $\tilde{\zeta}(y) = (\partial^{s-1} \zeta / \partial y_k^{s-1})(y)$ . Then  $\tilde{\zeta}$  satisfies

$$\tilde{\zeta}(\tilde{G}_1(I), \dots, \tilde{G}_k(I)) = 0$$

and  $(\partial \tilde{\zeta} / \partial y_k)(\tilde{G}(I)) \neq 0$  on  $V$ . Hence,

$$\tilde{\zeta}(G^0(x)) = 0 \tag{A.1}$$

and  $(\partial \tilde{\zeta} / \partial y_k)(G^0(x)) \neq 0$  on  $\varphi(V \times \mathbb{T}^q)$ .

Let  $\hat{G}_k^\varepsilon(x) = \tilde{\zeta}(G^\varepsilon(x)) / \varepsilon$ . By (A.1)  $\hat{G}_k^\varepsilon$  is an analytic first integral depending analytically on  $\varepsilon$ . We have

$$d\hat{G}_k^\varepsilon = \varepsilon^{-1} d(\tilde{\zeta}(G^\varepsilon(x))) = \varepsilon^{-1} \sum_{j=1}^k \frac{\partial \tilde{\zeta}}{\partial y_j}(G^\varepsilon(x)) dG_j^\varepsilon$$

and

$$N(\hat{G}^\varepsilon) := dG_1^\varepsilon \wedge \dots \wedge dG_{k-1}^\varepsilon \wedge d\hat{G}_k^\varepsilon = \varepsilon^{-1} \frac{\partial \tilde{\zeta}}{\partial y_k}(G^\varepsilon(x)) dG_1^\varepsilon \wedge \dots \wedge dG_k^\varepsilon.$$

Since  $(\partial \tilde{\zeta} / \partial y_k)(G^0(x)) \neq 0$  on  $\varphi(V \times \mathbb{T}^q)$ , we have  $\sigma((\partial \tilde{\zeta} / \partial y_k)(G^0(x))) = 0$ , so that

$$\sigma(N(\hat{G}^\varepsilon)) = \sigma(\varepsilon^{-1} N(G^\varepsilon)) + \sigma\left(\frac{\partial \tilde{\zeta}}{\partial y_k}(G^\varepsilon(x))\right) = \sigma(N(G^\varepsilon)) - 1.$$

Repeating this procedure till  $\sigma(N(\hat{G}^\varepsilon)) = 0$ , we obtain

$$dG_1^0 \wedge \dots \wedge dG_{k-1}^0 \wedge d\hat{G}_k^0 \neq 0,$$

which means the desired result.  $\square$

## REFERENCES

- [1] V.I. Arnold, *Geometrical Methods in the Theory of Ordinary Differential Equations*, 2nd ed., Springer, New York, 1988.
- [2] V.I. Arnold, *Mathematical Methods of Classical Mechanics*, 2nd ed., Springer, New York, 1989.
- [3] V.I. Arnold, V.V. Kozlov and A.I. Neishtadt, *Dynamical Systems III: Mathematical Aspects of Classical and Celestial Mechanics*, 3rd ed., Springer, Berlin, 2006.
- [4] M. Audin, *Hamiltonian Systems and Their Integrability*, American Mathematical Society, Providence, RI, 2008.

- [5] M. Ayoul and N.T. Zung, Galoisian obstructions to non-Hamiltonian integrability, *C. R. Math. Acad. Sci. Paris*, **348** (2010), 1323–1326.
- [6] A. Baider, R.C. Churchill, D.L. Rod and M.F. Singer, On the infinitesimal geometry of integrable systems, *Mechanics Days*, W.F. Shadwick, P.S. Krishnaprasad and T.S. Ratiu (eds.), American Mathematical Society, Providence, RI, 1996.
- [7] J. Barrow-Green, *Poincaré and the Three-Body Problem*, American Mathematical Society, Providence, RI, 1996.
- [8] O.I. Bogoyavlenskij, Extended integrability and bi-hamiltonian systems, *Comm. Math. Phys.*, **196** (1998), 19–51.
- [9] P.F. Byrd and M.D. Friedman, *Handbook of Elliptic Integrals for Engineers and Physicists*, Springer, Berlin, 1954.
- [10] R. Churchill and D.L. Rod, Geometrical aspects of Ziglin’s nonintegrability theorem for complex Hamiltonian systems, *J. Differential Equations*, **76** (1988), 91–114.
- [11] J. Guckenheimer and P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer, New York, 1983.
- [12] P. Holmes, A nonlinear oscillator with a strange attractor, *Philos. Trans. Roy. Soc. London Ser. A*, **292** (1979), 419–448.
- [13] V.V. Kozlov, Integrability and non-integrability in Hamiltonian mechanics, *Russian Math. Surveys*, **38** (1983), 1–76.
- [14] V.V. Kozlov, *Symmetries, Topology and Resonances in Hamiltonian Mechanics*, Springer, Berlin, 1996.
- [15] V.K. Melnikov, On the stability of the center for time periodic perturbations, *Trans. Moscow Math. Soc.*, **12** (1963), 1–56.
- [16] J.J. Morales-Ruiz, *Differential Galois Theory and Non-Integrability of Hamiltonian Systems*, Birkhäuser, Basel, 1999.
- [17] J.J. Morales-Ruiz, A note on a connection between the Poincaré-Arnold-Melnikov integral and the Picard-Vessiot theory, in *Differential Galois theory*, T. Crespo and Z. Hajto (eds.), Banach Center Publ. 58, Polish Acad. Sci. Inst. Math., 2002, pp. 165–175.
- [18] J.J. Morales-Ruiz and J.P. Ramis, Galoisian obstructions to integrability of Hamiltonian systems, *Methods, Appl. Anal.*, **8** (2001), 33–96.
- [19] J. Moser, *Stable and Random Motions in Dynamical Systems: With Special Emphasis on Celestial Mechanics*, Princeton Univ. Press, Princeton, NJ, 1973.
- [20] S. Motonaga and K. Yagasaki, Persistence of periodic and homoclinic orbits, first integrals and commutative vector fields in dynamical systems, submitted for publication.
- [21] R. Narasimhan, *Analysis on Real and Complex Manifolds*, North-Holland, Amsterdam, 1968.
- [22] H. Poincaré, Sur le problème des trois corps et les équations de la dynamique, *Acta Math.*, **13** (1890), 1–270; English translation: *The Three-Body Problem and the Equations of Dynamics*, Translated by D. Popp, Springer, Cham, Switzerland, 2017.
- [23] H. Poincaré, *New Methods of Celestial Mechanics*, Vol. 1, AIP Press, New York, 1992 (original 1892).
- [24] F.A. Rodrigues, T.K.D. Peron, P. Ji and J. Kurths, The Kuramoto model in complex networks, *Phys. Rep.*, **610** (2016), 1–98.
- [25] S. Wiggins, *Introduction to Applied Nonlinear Dynamical Systems and Chaos*, Springer, New York, 1990.
- [26] K. Yagasaki, Homoclinic motions and chaos in the quasiperiodically forced van der Pol-Duffing oscillator with single well potential, *Proc. R. Soc. Lond. A*, **445** (1994), 597–617.
- [27] K. Yagasaki, The Melnikov theory for subharmonics and their bifurcations in forced oscillations, *SIAM J. Appl. Math.*, **56** (1996), 1720–1765.
- [28] K. Yagasaki, Melnikov’s method and codimension-two bifurcations in forced oscillations, *J. Differential Equations*, **185** (2002), 1–24.
- [29] K. Yagasaki, Degenerate resonances in forced oscillators, *Discrete Contin. Dyn. Syst. B*, **3** (2003), 423–438.
- [30] K. Yagasaki, Nonintegrability of nearly integrable dynamical systems near resonant periodic orbits, submitted for publication.
- [31] K. Yagasaki, Nonintegrability of the restricted three-body problem, submitted for publication.
- [32] S.L. Ziglin, Bifurcation of solutions and the nonexistence of first integrals in Hamiltonian mechanics. I. *Funktsional. Anal. i Prilozhen.*, **16** (1982), 30–41, 96.

- [33] S.L. Ziglin, Self-intersection of the complex separatrices and the non-existing of the integrals in the Hamiltonian systems with one-and-half degrees of freedom, *J. Appl. Math. Mech.*, **45** (1982), 411–413.
- [34] N.T. Zung, Torus actions and integrable systems, in *Topological Methods in the Theory of Integrable Systems*, A.V. Bolsinov, A.T. Fomenko and A.A. Oshemkov (eds.), Cambridge Scientific Publications, Cambridge, 2006, pp. 289–328.
- [35] N.T. Zung, A conceptual approach to the problem of action-angle variables, *Arch. Ration. Mech. Anal.*, **229** (2018), 789–833.

DEPARTMENT OF APPLIED MATHEMATICS AND PHYSICS, GRADUATE SCHOOL OF INFORMATICS,  
KYOTO UNIVERSITY, YOSHIDA-HONMACHI, SAKYO-KU, KYOTO 606-8501, JAPAN

*Email address:* `mnaga@amp.i.kyoto-u.ac.jp` (S. Motonaga)

*Email address:* `yagasaki@amp.i.kyoto-u.ac.jp` (K. Yagasaki)