

# ON THE PRIME GRAPH OF A FINITE GROUP WITH UNIQUE NONABELIAN COMPOSITION FACTOR

MARIA A. GRECHKOSEVA

*Sobolev Institute of Mathematics  
Koptiyuga 4, Novosibirsk 630090, Russia*

ANDREY V. VASIL'EV

*Sobolev Institute of Mathematics  
Koptiyuga 4, Novosibirsk 630090, Russia  
Novosibirsk State University  
Pirogova 1, Novosibirsk 630090, Russia*

**ABSTRACT.** We say that finite groups are isospectral if they have the same sets of orders of elements. It is known that every nonsolvable finite group  $G$  isospectral to a finite simple group has a unique nonabelian composition factor, that is, the quotient of  $G$  by the solvable radical of  $G$  is an almost simple group. The main goal of this paper is prove that this almost simple group is a cyclic extension of its socle.

To this end, we consider a general situation when  $G$  is an arbitrary group with unique nonabelian composition factor, not necessarily isospectral to a simple group, and study the prime graph of  $G$ , where the prime graph of  $G$  is the graph whose vertices are the prime numbers dividing the order of  $G$  and two such numbers  $r$  and  $s$  are adjacent if and only if  $r \neq s$  and  $G$  has an element of order  $rs$ . Namely, we establish some sufficient conditions for the prime graph of such a group to have a vertex adjacent to all other vertices. Besides proving the main result, this allows us to refine a recent result by P. Cameron and N. Maslova concerning finite groups almost recognizable by prime graph.

**Keywords:** almost simple group, group of Lie type, order of an element, recognition by spectrum, prime graph

## 1. INTRODUCTION

Given a finite group  $G$ , we denote the set of prime divisors of the order of  $G$  by  $\pi(G)$ . The set of element orders of  $G$  is called the spectrum of  $G$  and denoted by  $\omega(G)$ . If  $\omega(G) = \omega(H)$ , then  $G$  and  $H$  are said to be isospectral.

Suppose that  $G$  is a finite group isospectral to a finite nonabelian simple group  $L$ . Then  $G$  is either solvable, in which case  $L$  is one of  $L_3(3)$ ,  $U_3(3)$ ,  $S_4(3)$ , or has exactly

---

*E-mail addresses:* grechkoseeva@gmail.com, vasand@math.nsc.ru

This work was supported by the Program of Fundamental Scientific Researches of the Siberian Branch of Russian Academy of Sciences, I.1.1, project 0314-2019-0001.

one nonabelian composition factor (see [5, Theorem 2]). In what follows, we assume that  $G$  is not solvable, and so  $G$  has a normal series

$$(1.1) \quad 1 \leq K < H \leq G,$$

where  $K$  is the solvable radical of  $G$ ,  $H/K$  is a nonabelian simple group and  $G/K$  is an almost simple group with socle  $H/K$ . Denoting  $H/K$  by  $S$ , we may identify  $G/K$  with a subgroup of  $\text{Aut } S$ , and then  $G/H$  with a subgroup of  $\text{Out } S = \text{Aut } S/S$ . Observe that  $G/H$  is solvable.

If  $L$  is sufficiently 'large', more precisely, if  $L$  is a classical group of dimension at least 38 or a non-classical group other than  $\text{Alt}_6$ ,  $\text{Alt}_{10}$ ,  $J_2$ ,  ${}^3D_4(2)$ , then  $K = 1$  and  $H \simeq L$  (see [9, 13]). Furthermore, it follows that  $G/H$  is cyclic (see [8] and the references therein). In general case,  $K$  is not always trivial and  $H/K$  is not always isomorphic to  $L$  but in all known examples,  $G/H$  is cyclic. This observation suggests us to conjecture that  $G/H$  is always cyclic and the main goal of this paper is to prove this conjecture.

**Theorem 1.** *Let  $L$  be a finite nonabelian simple group and let  $G$  be a nonsolvable finite group with  $\omega(G) = \omega(L)$ . Suppose that  $1 \leq K < H \leq G$  is the normal series of  $G$  as in (1.1). Then  $G/H$  is cyclic. Furthermore, if  $H/K$  is a simple group of Lie type other than  $L_2(q)$ , then  $G/H$  does not contain diagonal automorphisms.*

If  $L$  is sporadic or alternating, Theorem 1 is a direct consequence of the known description of groups isospectral to  $L$ . If  $L$  is a group of Lie type, the proof has several ingredients. The first is the well-known property of spectra of groups of Lie type stated in Lemma 2.1 in Section 2. The second is the nilpotency of the solvable radical of  $G$  established in [18]. The third is the following Theorem 2 which concerns all finite groups of some specific structure, not only those isospectral to simple groups.

**Theorem 2.** *Suppose that a finite group  $G$  has a normal series  $1 \leq K < H \leq G$ , where  $K$  is the solvable radical of  $G$ ,  $S = H/K$  is a finite simple group of Lie type, and  $G/K \leq \text{Aut } S$ . Suppose also that  $K$  is nilpotent.*

- (i) *If  $S \neq L_2(q)$  and  $G/H$  contains a diagonal automorphism of  $S$  of prime order  $r$ , then  $rs \in \omega(G)$  for all  $s \in \pi(G) \setminus \{r\}$ .*
- (ii) *If  $G/H$  is not cyclic, then there is  $r \in \pi(G/H)$  such that  $rs \in \omega(G)$  for all  $s \in \pi(G) \setminus \{r\}$ .*

The set  $\omega(G)$  defines the prime graph of  $G$  as follows: the vertex set of this is  $\pi(G)$  and two primes  $r, s \in \pi(G)$  are adjacent if and only if  $r \neq s$  and  $rs \in \omega(G)$ . The prime graph is also known as the Gruenberg–Kegel graph and we denote it by  $GK(G)$ . It is not hard to see that Theorem 2 states a property of the graph  $GK(G)$  rather than of the whole set  $\omega(G)$ . This allows us to apply this theorem to the problem of recognition of simple groups by prime graph. Recently, P. Cameron and N. Maslova [1] proved several new results relating to this problem. In Theorem 3, we slightly refine Theorem 1.4 of [1].

**Theorem 3.** *There exists a function  $F(x) = O(x^5)$  such that for each labeled graph  $\Gamma$ , the following conditions are equivalent:*

- (i) *there exist infinitely many groups  $H$  such that  $GK(H) = \Gamma$ ;*
- (ii) *there exist more than  $F(|V(\Gamma)|)$  groups  $H$  such that  $GK(H) = \Gamma$ , where  $V(\Gamma)$  is the set of the vertices of  $\Gamma$ .*

In fact, Theorem 1.4 of [1] states exactly the same as Theorem 3 but with  $x^7$  in place of  $x^5$ .

## 2. PROOFS OF THEOREMS 1 AND 2

We begin this section with notation and preliminary results. We write  $L_n^\varepsilon(q)$  and  $E_6^\varepsilon(q)$  assuming that  $\varepsilon \in \{+, -\}$ ,  $L_n^+(q) = L_n(q)$ ,  $L_n^-(q) = U_n(q)$ ,  $E_6^+(q) = E_6(q)$ , and  $E_6^-(q) = {}^2E_6(q)$ . If  $r$  is a prime and  $a$  is an integer, then  $(a)_r$  is the highest power of  $r$  dividing  $a$ . If  $S$  is a group of Lie type, then  $\text{Inndiag } S$  is the subgroup of  $\text{Aut } S$  generated by inner and diagonal automorphisms, and  $\text{Outdiag } S$  is the image of  $\text{Inndiag } S$  in  $\text{Out } S$ . Also we use the terms ‘field automorphism’ and ‘graph automorphism’ of  $S$  according to [4, Definition 2.5.13].

**Lemma 2.1.** *If  $S$  is a finite simple group of Lie type, then for every  $r \in \pi(S)$  there is  $s \in \pi(S)$  such that  $r \neq s$  and  $rs \notin \omega(S)$ .*

**Proof.** This follows from [16, 17] (see, for example, [6, Lemma 2.2]).  $\square$

**Lemma 2.2.** *Let  $S$  be a finite simple group of Lie type in characteristic  $p$ . If  $r$  divides  $|\text{Outdiag } S|$  and  $rp \notin \omega(S)$ , then either  $S = L_2(q)$ , or  $S = L_3^\varepsilon(q)$  and  $(q - \varepsilon)_3 = 3$ .*

**Proof.** This follows, for example, from [16, Propositions 3.1 and 3.2].  $\square$

**Lemma 2.3.** *Let  $S$  be a finite simple group of Lie type in characteristic  $p$ . If  $r \in \pi(S)$ ,  $r$  is odd and  $2r \notin \omega(S)$ , then either a Sylow  $r$ -subgroup of  $S$  is cyclic, or  $S = L_2(q)$  and  $r = p$ , or  $S = L_3^\varepsilon(q)$ ,  $p = 2$ ,  $r = 3$  and  $(q - \varepsilon)_3 = 3$ .*

**Proof.** This follows from the results of [16, Sections 3 and 4] and the cross-characteristic Sylow structure of groups of Lie type [3, (10-2)].  $\square$

**Lemma 2.4.** *Let  $S = {}^t\Sigma(q)$  be a finite simple group of Lie type, not a Suzuki–Ree group, and let  $\varphi$  be a field automorphism of  $S$  of prime order  $r$ . Then  $r \cdot \omega({}^t\Sigma(q^{1/r})) \subseteq \omega(S \rtimes \langle \varphi \rangle)$ .*

**Proof.** This follows from the Lang–Steinberg theorem [14, Section 10] (see, for example, [7, Lemma 2.8]).  $\square$

**Lemma 2.5.** *Suppose that  $G$  is a finite group,  $K$  is a normal subgroup of  $G$  and every  $g \in G \setminus K$  acts fixed-point-freely on  $K$ . Then every odd order Sylow subgroup of  $G/K$  is cyclic and a Sylow 2-subgroup of  $G/K$  is cyclic or generalized quaternion.*

**Proof.** This is a well-known property of fixed-point-free automorphisms (see, for example, [10, Satz 8.7]).  $\square$

**Proof of Theorem 2.** Denote the defining characteristic of  $S$  by  $p$ ,  $G/K$  by  $\overline{G}$  and  $G/H$  by  $\widehat{G}$ . As we remarked in the introduction,  $\widehat{G}$  can be regarded as a subgroup of  $\text{Out } S$ .

Clearly, we may assume that either  $\text{Outdiag } S \neq 1$  or  $\text{Out } S$  is not cyclic, in particular, we may assume that  $S$  is not a Suzuki–Ree group and so  $3 \in \pi(S)$ .

(i) Suppose that  $r \in \pi(\widehat{G} \cap \text{Outdiag } S)$ . Observe that  $r \in \pi(S)$  and  $r \neq p$ . By Lemma 2.2, it follows that  $rp \in \omega(S)$  unless  $S = L_3^\varepsilon(q)$ ,  $r = 3$  and  $(q - \varepsilon)_3 = 3$ . In this case  $PGL_3^\varepsilon(q) \leq \overline{G}$ , and since  $PGL_3^\varepsilon(q)$  has an element of order  $p(q - \varepsilon)$ , we see that  $rp \in \omega(\overline{G})$ .

Suppose that  $s \in \pi(S)$  and  $s \neq p$ . If  $s \in \pi(\text{Outdiag } S)$ , then  $rs \in \omega(S)$  since  $\text{Outdiag } S$  is abelian. So we may assume that  $s \notin \pi(\text{Outdiag } S)$ . The maximal tori of  $\text{Inndiag } S$  are isomorphic to those of the universal version  $\tilde{S}_u$  of  $\tilde{S}$ , where  $\tilde{S} = S$  if  $S$  is not of type  $B_n$  or  $C_n$ , and  $\tilde{B}_n(q) = C_n(q)$ ,  $\tilde{C}_n(q) = B_n(q)$  (see [2, Section 4.4]). Since every maximal torus of  $\tilde{S}_u$  contains the center  $Z(\tilde{S}_u)$  of  $\tilde{S}_u$  and  $|Z(\tilde{S}_u)| = |\text{Outdiag } S|$ , we see that  $\text{Inndiag } S$  includes a maximal torus whose order is divisible by  $s|\text{Outdiag } S|$ . So  $\overline{G}$  contains an abelian subgroup of order  $sr$ .

Let  $s \in \pi(\overline{G}) \setminus \pi(S)$ . Since  $s \neq 2, 3$  and  $s \notin \pi(\text{Outdiag } S)$ , it follows that  $G/K$  contains a field automorphism of  $S$  of order  $s$ . By Lemma 2.4, we have  $s \cdot \omega(S_0) \subseteq \omega(G/K)$ , where  $S_0$  is a group of the same Lie type as  $S$ . If  $r = 2, 3$ , then it is clear that  $r \in \pi(S_0)$ . If  $r \neq 2, 3$ , then  $S = L_n^\varepsilon(q)$ ,  $r$  divides  $(n, q - \varepsilon)$  and  $S_0 = L_n^\varepsilon(q^{1/s})$ . Since  $r$  divides  $p^{r-1} - \varepsilon^{r-1}$  and  $r - 1 \leq n - 1$ , we see that  $r \in \pi(S_0)$ .

Let  $s \in \pi(K) \setminus \pi(\overline{G})$ . If  $r = 2$ , then  $s$  is adjacent to  $r$  in  $GK(G)$  by [15, Proposition 2]. So we may assume that  $r$  is odd. If  $S = E_6^\varepsilon(q)$  or  $S = L_n^\varepsilon(q)$  with  $n \geq 4$ , then  $S$  includes a torus of the form  $\mathbb{Z}_{q-\varepsilon} \times \mathbb{Z}_{q-\varepsilon}$ , and hence  $S$  includes an elementary abelian group of order  $r^2$ . If  $L = L_3^\varepsilon(q)$ , then  $PGL_3^\varepsilon(q) \leq \overline{G}$  and so  $\overline{G}$  includes an elementary abelian group of order  $r^2$ . Now we apply Lemma 2.5 to conclude that  $rs \in \omega(G)$ .

(ii) Let  $S \neq L_2(q)$ . By (i), we may assume that  $\widehat{G} \cap \text{Outdiag } S = 1$ . Then either  $\widehat{G}$  includes an elementary abelian group of order  $2^2$ , or  $S = O_8^+(q)$  and, up to conjugation in  $\text{Out } S$ ,  $\widehat{G}$  contains the image of the graph automorphism  $\gamma$  of  $S$  induced by the symmetry of the Dynkin diagram of order 3.

In the first case,  $S = L_n(q)$ ,  $O_{2n}^+(q)$ , or  $E_6(q)$ , and we claim that 2 is adjacent to all odd primes in  $GK(G)$ . By [15, Proposition 2], every  $s \in \pi(K) \cup \pi(\widehat{G})$  is adjacent to 2. Now let  $t \in \pi(S)$  and suppose that  $2t \notin \omega(S)$ . Excluding for a while the case when  $t = 3$ ,  $S = L_3(q)$ ,  $p = 2$ ,  $(q - 1)_3 = 3$  and applying Lemma 2.3, we conclude that a Sylow  $t$ -subgroup  $T$  of  $S$  is cyclic, and hence  $N_{\overline{G}}(T)/C_{\overline{G}}(T)$  is cyclic. On the other hand, by the Frattini argument,  $N_{\overline{G}}(T)/(N_{\overline{G}}(T) \cap S) \simeq \widehat{G}$ , and so a Sylow 2-subgroup of  $N_{\overline{G}}(T)$  is not cyclic. Thus  $2 \in C_{\overline{G}}(T)$ , and  $2t \in \omega(\overline{G})$ .

Suppose that  $t = 3$ ,  $S = L_3(q)$ ,  $p = 2$ , and  $(q - 1)_3 = 3$ . Since  $\widehat{G}$  includes an elementary abelian group of order  $2^2$ , it follows that  $\overline{G}$  contains a field automorphism of  $S$  of order 2, and so  $6 \in \omega(\overline{G})$ .

Now suppose that  $S = O_8^+(q)$  and  $\overline{G}$  contains the graph automorphism  $\gamma$ . The centralizer of  $\gamma$  in  $S$  is isomorphic to  $G_2(q)$  [3, (9-1)] and so  $3s \in \omega(G)$  for all  $3 \neq s \in G_2(q)$ . Since  $S$  includes an elementary abelian group of order 9, we conclude that  $3s \in \omega(G)$  for all  $s \in \pi(K) \setminus \{3\}$ . Also a 2'-Hall subgroup of  $\text{Out } S$  is abelian, and hence 3 is adjacent to every  $s \in \pi(\widehat{G}) \setminus \{2, 3\}$  in  $GK(\widehat{G})$ . Let  $s \in \pi(S) \setminus \{3\}$  and  $3s \notin \omega(S)$ . Then  $s$  divides  $q^2 + q + 1$  or  $q^2 - q + 1$ , therefore,  $s \in \pi(G_2(q))$  and, as we remarked,  $3s \in \omega(G)$ . Thus 3 is adjacent to all vertices in  $GK(G)$ .

Let  $S = L_2(q)$ , where  $q = p^l$ . We claim that 2 is adjacent to all odd primes in  $GK(G)$ . Since  $\text{Out } S$  is a direct product of cyclic groups of orders  $(2, q - 1)$  and  $l$ , it follows that  $p$  is odd,  $l$  is even and  $\overline{G} = PGL_2(q) \rtimes \langle \varphi \rangle$ , where  $\varphi$  is a field automorphism of  $S$  of even order. Since  $PGL_2(q)$  contains elements of orders  $q \pm 1$  and  $2p \in \omega(\overline{G})$  by Lemma 2.4, we see that 2 is adjacent to every odd  $s \in \pi(\overline{G})$ . Let  $s \in \pi(K)$  be odd. A Sylow 2-subgroup of  $PGL_2(q)$  is dihedral, and so it cannot act fixed-point-freely on a Sylow  $s$ -subgroup of  $K$  by Lemma 2.5. Hence  $2s \in \pi(G)$ , and the proof of Theorem 2 is complete.  $\square$

Now we are able to prove Theorem 1. Let  $S = H/K$ . Clearly, we may assume that  $\text{Out } S$  is not cyclic. In particular, we may assume that  $S$  is neither sporadic nor alternating with the following convention: if  $S = \text{Alt}_6 \simeq L_2(9)$ , we regard  $S$  as a group of Lie type.

If  $L$  is sporadic and  $L \neq J_2$ , or if  $L = \text{Alt}_n$  and  $n \neq 6, 10$ , then  $G \simeq L$  (see [11] and [5] respectively). If  $L = J_2$ , then  $G \simeq L$  or  $S = \text{Alt}_8$  by [11]. If  $L = \text{Alt}_{10}$ , then  $G \simeq L$  or  $S = \text{Alt}_5$  by [12]. If  $L = \text{Alt}_6$ , we regard  $L$  as a group of Lie type.

Let  $L$  be a group of Lie type. By [18, Theorem 1], it follows that  $K$  is nilpotent, and so  $G$  satisfies the hypothesis of Theorem 2. If  $G/H$  is not cyclic or if  $S \neq L_2(q)$  and

$G/H$  contains a diagonal automorphism of  $S$ , then there is  $r \in \pi(G)$  adjacent to all other vertices in  $GK(G)$ . But this is impossible by Lemma 2.1 since  $GK(G) = GK(L)$ . This contradiction completes the proof of Theorem 1.

### 3. GROUPS ALMOST RECOGNIZABLE BY PRIME GRAPH

Given a positive integer  $k$ , a finite group  $G$  is said to be  $k$ -recognizable by prime graph if there are exactly  $k$  pairwise nonisomorphic finite groups  $H$  with  $GK(H) = GK(G)$  and almost recognizable by prime graph if it is  $k$ -recognizable for some  $k$ .

By [1, Theorem 1.3], if  $G$  is almost recognizable by prime graph, then  $G$  is almost simple and each group  $H$  with  $GK(H) = GK(G)$  is almost simple. So if  $G$  is a  $k$ -recognizable group, then  $k$  is at most the number of almost simple groups  $H$  such that  $\pi(H) = \pi(G)$ . By [1, Proposition 4.2], this number is at most  $O(|\pi(G)|^7)$ . A direct corollary of this discussion is the following theorem.

**Theorem A** [1, Theorem 1.4]. *There exists a function  $F(x) = O(x^7)$  such that for each labeled graph  $\Gamma$ , the following conditions are equivalent:*

- (i) *there exist infinitely many groups  $H$  such that  $GK(H) = \Gamma$ ;*
- (ii) *there exist more than  $F(|V(\Gamma)|)$  groups  $H$  such that  $GK(H) = \Gamma$ , where  $V(\Gamma)$  is the set of the vertices of  $\Gamma$ .*

It is clear that estimating  $k$  we do not need to calculate all almost simple groups  $H$  such that  $\pi(H) = \pi(G)$ . It is sufficient to calculate those  $H$  whose prime graph satisfies some necessary conditions for  $H$  to be almost recognizable by prime graph. One of these conditions is stated in [1, Theorem 1.3]: 2 is nonadjacent to at least one odd prime in  $GK(H)$ . But in fact this condition can be strengthened: every  $r \in \pi(H)$  is nonadjacent to at least one prime  $s \neq r$  in  $GK(H)$ . Indeed, otherwise  $GK(H) = GK(H \times \mathbb{Z}_r^k)$  for all positive integers  $k$ . Applying Theorem 2, we see that it sufficient to calculate  $H$  such that  $H/S$  is cyclic, where  $S$  is the socle of  $H$ .

**Lemma 3.1.** *There is a function  $F(x) = O(x^2)$  such that if  $S$  is a finite simple group of Lie type, then there are at most  $F(|\pi(S)|)$  almost simple groups  $H$  with socle  $S$  such that  $H/S$  is cyclic.*

**Proof.** Let  $n$  be the Lie rank of  $S$  and  $q = p^l$  the order of the base field of  $S$ . Denote the number of divisors of  $l$  by  $d(l)$ . By [1, Lemma 2.7], we have  $n \leq 2|\pi(S)| + 3$  and  $d(l) \leq |\pi(S)| + 1$ .

Steinberg's theorem [4, Theorem 2.5.12] states that  $\text{Out } S = \text{Outdiag } S \rtimes \Phi_S \Gamma_S$ , where  $|\text{Outdiag } S| \leq n + 1$  and  $\Phi_S \Gamma_S$  is either a subgroup in  $\mathbb{Z}_l \times \text{Sym}_3$  or a cyclic group of order  $2l$  or  $3l$ . In any case the number of cyclic subgroups of  $\Phi_S \Gamma_S$  is at most  $6d(l)$ . Thus the number of cyclic subgroups of  $\text{Out } S$  is at most  $6(n + 1)d(l)$ , which is  $O(|\pi(S)|^2)$  by the preceding paragraph.  $\square$

Now we are ready to prove Theorem 3 (in fact we follow the lines of the proof of [1, Theorem 1.4] but Theorem 2 allows us to use the bound of Lemma 3.1 instead of that of [1, Proposition 4.1]). It is sufficient to show that there exists a function  $F(x) = O(x^5)$  such that for every finite group  $G$ , if  $G$  is almost recognizable by prime graph, then there are at most  $F(|\pi(G)|)$  pairwise nonisomorphic groups  $H$  with  $GK(H) = GK(G)$ .

Assume that  $G$  is  $k$ -recognizable by prime graph. By [1, Theorem 1.3], each group  $H$  with  $GK(H) = GK(G)$  is almost simple. Furthermore, as we remarked, every  $r \in \pi(H)$  is nonadjacent to at least one prime  $s \neq r$  in  $GK(H)$ . By Theorem 2, it follows that  $H$  is

a cyclic extension of its socle. By [1, Proposition 3.1], the number of nonabelian simple groups  $S$  such that  $\pi(S) \subseteq \pi(G)$  is bounded by  $F_1(|\pi(G)|)$  with  $F_1(x) = O(x^3)$ . Applying Lemma 3.1, we see that the number of almost simple groups  $H$  with socle  $S$  such that  $H/S$  is cyclic is at most  $F_2(|\pi(S)|)$ , where  $F_2(x) = O(x^3)$ . Thus  $k \leq F_1(|\pi(G)|)F_2(|\pi(G)|) = O(|\pi(G)|^5)$ , and this completes the proof of Theorem 3.

## REFERENCES

- [1] P. J. Cameron and N. V. Maslova, Criterion of unrecognizability of a finite group by its Gruenberg-Kegel graph, 2021, arXiv:2012.01482 [math.GR].
- [2] R. W. Carter, Finite groups of Lie type. Conjugacy classes and complex characters, John Wiley & Sons, New York, 1985.
- [3] D. Gorenstein and R. Lyons, Local structure of finite groups of characteristic 2 type, *Memoirs Amer. Math. Soc.* **276**, 1983.
- [4] D. Gorenstein, R. Lyons, and R. Solomon, The classification of the finite simple groups. Number 3, *Amer. Math. Soc. Surveys and Monographs* **40.3**, 1998.
- [5] I. B. Gorshkov, Recognizability of alternating groups by spectrum, *Algebra Logic* **52** (2013), no. 1, 41–45.
- [6] M. A. Grechkoseeva, On element orders in covers of finite simple groups of Lie type, *J. Algebra Appl.* **14** (2015), 1550056 [16 pages].
- [7] M. A. Grechkoseeva, On spectra of almost simple groups with symplectic or orthogonal socle, *Siberian Math. J.* **57** (2016), no. 4, 582–588.
- [8] M. A. Grechkoseeva, On spectra of almost simple extensions of even-dimensional orthogonal groups, *Siberian Math. J.* **59** (2018), no. 4, 623–640.
- [9] M. A. Grechkoseeva and A. V. Vasil'ev, On the structure of finite groups isospectral to finite simple groups, *J. Group Theory* **18** (2015), no. 5, 741–759.
- [10] B. Huppert, *Endliche Gruppen. I*, Springer-Verlag, Berlin, 1967.
- [11] V. D. Mazurov and W. J. Shi, A note to the characterization of sporadic simple groups, *Algebra Colloq.* **5** (1998), no. 3, 285–288.
- [12] A. M. Staroletov, Groups isospectral to the degree 10 alternating group, *Siberian Math. J.* **51** (2010), no. 3, 507–514.
- [13] A. Staroletov, On almost recognizability by spectrum of simple classical groups, *Int. J. Group Theory* **6** (2017), no. 4, 7–33.
- [14] R. Steinberg, Endomorphisms of linear algebraic groups, *Memoirs Amer. Math. Soc.* **80**, 1968.
- [15] A. V. Vasil'ev, On connection between the structure of a finite group and the properties of its prime graph, *Siberian Math. J.* **46** (2005), no. 3, 396–404.
- [16] A. V. Vasil'ev and E. P. Vdovin, An adjacency criterion for the prime graph of a finite simple group, *Algebra Logic* **44** (2005), no. 6, 381–406.
- [17] A. V. Vasil'ev and E. P. Vdovin, Cliques of maximal size in the prime graph of a finite simple group, *Algebra Logic* **50** (2011), no. 4, 291–322.
- [18] N. Yang, M. A. Grechkoseeva, and A. V. Vasil'ev, On the nilpotency of the solvable radical of a finite group isospectral to a simple group, *J. Group Theory* **23** (2020), no. 3, 447–470.