

# FINDING AND COMBINING INDICABLE SUBGROUPS OF BIG MAPPING CLASS GROUPS

CAROLYN R. ABBOTT, HANNAH HOGANSON, MARISSA LOVING, PRIYAM PATEL,  
AND RACHEL SKIPPER

ABSTRACT. In this paper, we prove a combination theorem for indicable subgroups of infinite-type (or big) mapping class groups. Importantly, all subgroups produced by the combination theorem, as well as those coming from the other results of the paper, can be constructed so that they do not lie in the closure of the compactly supported mapping class group and do not lie in the isometry group for any hyperbolic metric on the relevant infinite-type surface. Along the way, we prove an embedding theorem for indicable subgroups of mapping class groups, a corollary of which gives embeddings of big mapping class groups into other big mapping class groups that are not induced by embeddings of the underlying surfaces. We also give new constructions of free groups, wreath products with  $\mathbb{Z}$ , and Baumslag-Solitar groups in big mapping class groups that can be used as an input for the combination theorem. One application of our combination theorem is a new construction of right-angled Artin groups in big mapping class groups.

## 1. INTRODUCTION

A fundamental question in low-dimensional topology asks which groups can arise as subgroups of the diffeomorphism group, homeomorphism group, and mapping class group of a surface, denoted by  $\text{Homeo}(S)$ ,  $\text{Diffeo}(S)$ , and  $\text{Map}(S)$ , respectively. A classical approach to this problem is to show that a particular group  $G$  acts by orientation-preserving isometries on a surface  $S$ , which implies that  $G$  is a subgroup of the three relevant groups for  $S$  mentioned above. In this paper, we give an explicit construction for building subgroups of the mapping class group of certain infinite-type surfaces that do not arise from isometries and describe how these subgroups sit inside the mapping class group.

One of the original motivating questions in this line of inquiry was posed by Nielsen. He asked whether every finite subgroup of the mapping class group of a finite-type surface  $\Sigma$  can be realized as a subgroup of the isometry group of some hyperbolic metric on  $\Sigma$ . This is often referred to as the “Nielsen realization problem”. In celebrated work, Kerckhoff answered Nielsen’s question in the affirmative, [Ker83]. Recently this result was extended to the infinite-type setting by Afton–Calegari–Chen–Lyman, [ACCL20]. However, this result only holds for finite subgroups of  $\text{Map}(S)$ . Since the mapping class group of an infinite-type surface (called a big mapping class group) is uncountably infinite, it is natural to ask if a similar type of Nielsen realization holds for countable, and even uncountable, subgroups.

---

*Date:* October 6, 2021.

Aougab–Patel–Vlamis [APV] recently gave a characterization of what groups can be obtained as the isometry group of some hyperbolic metric on various classes of infinite-type surfaces. For example, they show that if an infinite-type surface  $S$  contains a non-displaceable subsurface, then a group  $G$  can be realized as the isometry group of some hyperbolic metric on  $S$  if and only if  $G$  is finite. Thus, for these surfaces, any infinite subgroup of the mapping class group of  $S$  cannot act via isometries on  $S$ . They also show that no uncountable group can be obtained as the isometry group of a hyperbolic metric on any infinite-type surface. These results make clear that if we are interested in explicit constructions of interesting subgroups of big mapping class groups, then focusing solely on building them from isometries will be too restrictive.

Below, we use a method inspired by work of Allcock [All06] to explicitly construct various subgroups of infinite-type mapping class groups that cannot be realized as subgroups of the isometry group for any hyperbolic metric on the surface. In particular, Allcock uses a Cayley graph for a group  $G$  to construct a hyperbolic surface whose isometry group is exactly  $G$ . We edit this construction and utilize shift maps on the surface  $S$  in such a way that the relevant group cannot act via isometries on  $S$ . Furthermore, we construct (uncountably many) distinct embeddings of such a group  $G$  in  $\text{Map}(S)$ . Lastly, all of the subgroups we produce are of intrinsically infinite type, meaning that they do not lie fully in the closure of the compactly supported mapping class group,  $\overline{\text{Map}_c(S)}$ . The only exceptions to this are when  $S$  is finite-type or the Loch Ness Monster surface, for which  $\overline{\text{Map}_c(S)} = \text{Map}(S)$ . Our main theorem is a combination theorem for indicable subgroups of  $\text{Map}(S)$ , i.e., subgroups which admit a surjection to  $\mathbb{Z}$ . The  $\star$ -product used in the Theorem 1.1 is known in the literature as a “free product with commuting subgroups,” a natural construction that has been well-studied in the literature. Some basic group theoretic properties of such groups can be found in [MKS04, Section 4.2, Problems 22–25]. In particular, they show that the  $\star$ -product can be written as the iterated amalgamated product

$$(G_1, H_1) \star (G_2, H_2) = G_1 *_{H_1} (H_1 \times H_2) *_{H_2} G_2.$$

The algorithmic properties for these groups (such as the word and conjugacy problem) have been well-studied [MS73, Hur76]. Residual and separability properties of such groups have also been extensively studied in [Log99, Tie05, TM10, TM08, Sok14]. This product provides a natural interpolation between free products and direct products and includes, for example, graph products of groups.

**Theorem 1.1.** *Let  $G_i$  be indicable groups that embed in  $\text{Map}(S_i)$ , for  $i = 1, \dots, n$ , where  $S_i$  is a surface with exactly one boundary component. For each  $i$ , fix a surjective map  $f_i: G_i \rightarrow \mathbb{Z}$ , and let  $H_i$  be the kernel of  $f_i$ . Let  $\Pi$  be the surface obtained from an  $(n + 1)$ -holed sphere by gluing each  $S_i$  to one of its boundary components. Let  $\Gamma$  be the standard Cayley graph of  $\mathbb{F}_n$ , and let  $S = S_\Gamma(\Pi)$ . Then the indicable group  $(G_1, H_1) \star \dots \star (G_n, H_n)$  embeds in  $\text{Map}(S)$ .*

We direct the reader to Section 3 for a definition of the surface  $S_\Gamma(\Pi)$ . Importantly, the support of the homeomorphisms defined in our construction is not all of  $S_\Gamma(\Pi)$ . In particular, we may change the topology outside the support of the homeomorphisms in any way we choose. In this way, Theorem 1.1 actually shows that  $(G_1, H_1) \star \dots \star (G_n, H_n)$  embeds in the mapping class group of a wide class of infinite-type surfaces. In particular, we can arrange for the edited surface to have

a non-displaceable subsurface so that the subgroups we construct could not arise from a construction using isometries.

By carefully choosing  $G_i$  and  $H_i$ , we obtain the following corollary of the Theorem 1.1.

**Corollary 1.2.** *For any surface  $\Pi$  of sufficient complexity, there exists an infinite family of right-angled Artin groups which embed into  $\text{Map}(S)$  and are not completely contained in  $\overline{\text{Map}_c(S)}$ , where  $S = S_\Gamma(\Pi)$ .*

Right-angled Artin groups are well studied in connection to mapping class groups of finite-type surfaces as they give plentiful examples of convex cocompact subgroups. Although the constructions of right-angled Artin groups in the finite-type setting (for example, the Clay-Leininger-Mangahas Embedding Theorem [CLM12]), port immediately to the infinite-type setting through subsurface inclusion, they necessarily do not produce examples of subgroups of intrinsically infinite type. This is in stark contrast to the construction given in this paper.

One of the ingredients needed to prove Theorem 1.1 is Theorem 4.2, which demonstrates how to construct many distinct embeddings of a group  $G$  into  $\text{Map}(S)$ , where  $G$  is an indicable group that arises as a subgroup of the mapping class group of another surface  $\Pi$  with one boundary component, and  $S$  is a surface that admits a shift map with domain  $D_\Pi$  (see Definition 2.5). An important corollary to this result is obtained when the surface  $\Pi$  is itself of infinite type and  $G$  is taken to be the pure mapping class group  $\text{PMap}(\Pi)$ .

**Corollary 1.3.** *Let  $\Pi$  be an infinite-type surface with at least two non-planar ends and exactly one boundary component. Given any surface  $S$  that admits a shift whose domain is  $D_\Pi$ , there exist uncountably many embeddings of  $\text{PMap}(\Pi)$  into  $\text{Map}(S)$  that are not induced by an embedding of  $\Pi$  into  $S$ .*

This corollary is in line with a body of work dedicated to understanding and constructing interesting homomorphisms between mapping class groups; see, for example, [ALS09, AS13, ALM21]. In fact, when  $\text{Map}(\Pi)$  is indicable, we can use the full mapping class group instead of  $\text{PMap}(\Pi)$  in this corollary (see Corollary 4.5). We give examples of such surfaces  $\Pi$  in Example 4.4.

For other applications and to illustrate the breadth of Theorem 1.1, we note that there are a variety of indicable groups that arise as subgroups of mapping class groups of surfaces with exactly one boundary component that can play the role of  $G_i$  in the statement of Theorem 1.1 (or the role of  $G$  in the statement of Theorem 4.2). In particular, one can let  $G_i$  be any indicable subgroup of  $\text{Map}(S)$  where  $S$  is a finite-type surface with exactly one boundary component (for example, free group constructions coming from pseudo-Anosov elements, right-angled Artin groups, and braid groups, to name a few). In Section 3, we provide examples of new constructions of indicable subgroups of big mapping class groups which can play the role of  $G_i$ , including solvable Baumslag-Solitar groups  $BS(1, n)$ , free groups  $\mathbb{F}_n$ , and wreath products  $G \wr \mathbb{Z}$  where  $G$  is any group known to be a subgroup of the mapping class group of a surface with boundary. Moreover, all of these constructions yield subgroups that are of intrinsically infinite-type. These results are summarized in the following two propositions.

**Proposition 1.4.** *There exist infinite-type surfaces with non-normal, non-abelian free subgroups of any rank in their mapping class groups, each of which is of intrinsically infinite type.*

This construction of free subgroups is distinct from classical ones both because it produces non-normal subgroups and because it does not rely on applying the ping-pong lemma to collections of elements acting loxodromically on a hyperbolic metric space, for example, on the curve graph or on a projection complex as introduced by Bestvina–Bromberg–Fujiwara [BBF15]. A construction of free subgroups of big mapping class groups using the projection complex machinery is given by Horbez–Qing–Rafi [HQR20].

**Proposition 1.5.** *Let  $\Pi$  be a surface with a single boundary component. If  $S$  is a surface which admits a shift with domain  $D_\Pi$ , then, with appropriate restrictions on  $\Pi$ , the groups  $BS(1, n)$  and  $G \wr \mathbb{Z}$  arise as subgroups of  $\text{Map}(S)$ .*

More precise statements of Proposition 1.5 for each particular subgroup, including the necessary restrictions on  $\Pi$ , can be found in Section 3 (see Theorem 3.9 and Proposition 3.8). We construct these subgroups in the mapping class group of various surfaces, some of which have nondisplaceable subsurfaces. In the case of surfaces containing a nondisplaceable subsurface, these subgroups can never arise within the isometry group with respect to any hyperbolic metric [APV]; so our construction produces the first embeddings of these groups inside such big mapping class groups. On the other hand, it was already known that  $BS(1, n)$  and  $G \wr \mathbb{Z}$  must arise as the isometry group of, for example, some hyperbolic metric on the Blooming Cantor Tree Surface<sup>1</sup>, by work of Aougab–Patel–Vlamis [APV]. However, even in this case, Proposition 1.5 gives an uncountable collection of isomorphic copies of these subgroups in the big mapping class group, none of which can be realized by isometries due to the nature of our construction.

**Acknowledgements.** The authors would like to thank Women in Groups, Geometry, and Dynamics (WiGGD) for sponsoring their collaboration, which was supported by NSF DMS–1552234, DMS–1651963, and DMS–1848346. The authors also thank Mladen Bestvina and Rylee Lyman for helpful conversations, as well as George Domat for productive discussions about surfaces with indicable mapping class groups.

In addition, the authors acknowledge support from NSF grants DMS–1803368 and DMS–2106906 (Abbott), DMS–1906095 and RTG DMS–1840190 (Hoganson), DMS–1902729 (Loving), DMS–1937969 and DMS–2046889 (Patel), and DMS–2005297 (Skipper).

## 2. PRELIMINARIES

**2.1. Ends of surfaces.** Essential to the classification of infinite-type surfaces is the notion of an end of a surface and the space of all such ends for an infinite-type surface  $S$ .

**Definition 2.1.** An *exiting sequence* in  $S$  is a sequence  $\{U_n\}_{n \in \mathbb{N}}$  of connected open subsets of  $\Sigma$  satisfying:

- (1)  $U_n \subset U_m$  whenever  $m < n$ ;
- (2)  $U_n$  is not relatively compact for any  $n \in \mathbb{N}$ , that is, the closure of  $U_n$  in  $S$  is not compact;
- (3) the boundary of  $U_n$  is compact for each  $n \in \mathbb{N}$ ; and

---

<sup>1</sup>In fact, this is true for any surface of infinite genus with no planar ends, no boundary, and self similar end space.

- (4) any relatively compact subset of  $S$  is disjoint from all but finitely many of the  $U_n$ 's.

Two exiting sequences  $\{U_n\}_{n \in \mathbb{N}}$  and  $\{V_n\}_{n \in \mathbb{N}}$  are equivalent if for every  $n \in \mathbb{N}$  there exists  $m \in \mathbb{N}$  such that  $U_m \subset V_n$  and  $V_m \subset U_n$ . An *end* of  $S$  is an equivalence class of exiting sequences.

The *space of ends* of  $S$ , denoted  $E(S)$  or simply  $E$ , is the set of ends of  $S$  equipped with a natural topology for which it is totally disconnected, Hausdorff, second countable, and compact. In particular,  $E(S)$  is homeomorphic to a closed subset of the Cantor set. The definition of the topology on the space of ends is not relevant to this paper and is, therefore, omitted.

Ends of  $S$  can be isolated or not and can be planar (if there exists an  $i$  such that  $U_i$  is homeomorphic to an open subset of the plane  $\mathbb{R}^2$ ) or nonplanar (if every  $U_i$  has infinite genus). The set of nonplanar ends of  $S$  is a closed subspace of  $E(S)$  and will be denoted by  $E^g(S)$  (these are frequently called the ends accumulated by genus). We have the following classification theorem of Kerékjártó [Ker23] and Richards [Ric63]:

**Theorem 2.2** (Classification of infinite-type surfaces). *The homeomorphism type of an orientable infinite-type surface  $S$  is determined by the quadruple*

$$(g, b, E^g(S), E(S))$$

where  $g \in \mathbb{Z}_{\geq 0} \cup \infty$  is the genus of  $S$  and  $b \in \mathbb{Z}_{\geq 0}$  is the number of (compact) boundary components of  $S$ .

**2.2. Mapping class group.** The mapping class group of  $S$  is the set of orientation preserving homeomorphisms of  $S$  which fix the boundary pointwise, up to isotopy, and is denoted by  $\text{Map}(S)$ . The natural topology on the set of homeomorphisms of  $S$  is the compact-open topology, and  $\text{Map}(S)$  is endowed with the induced quotient topology. Equipped with this topology,  $\text{Map}(S)$  is a topological group. When  $S$  is a finite-type surface, this topology on  $\text{Map}(S)$  agrees with the discrete topology, but when  $S$  is of infinite type, the two topologies are distinct. Relevant subgroups of  $\text{Map}(S)$  include the pure mapping class group  $\text{PMap}(S)$ , which is the subgroup that fixes the set of ends of  $S$  pointwise, and the subgroup of compactly supported mapping classes denoted by  $\text{Map}_c(S)$ .

**Definition 2.3.** A mapping class  $f \in \text{Map}(S)$  is of intrinsically infinite type if  $f \notin \overline{\text{Map}_c(S)}$ . A subgroup  $H \leq \text{Map}(S)$  is of intrinsically infinite type if  $H$  is not completely contained in  $\overline{\text{Map}_c(S)}$ .

The closure of  $\text{Map}_c(S)$  is taken with respect to the compact-open topology in the above definition. In this paper, all of the subgroups of  $\text{Map}(S)$  that we construct contain many intrinsically infinite-type homeomorphisms and, therefore, cannot be completely contained in  $\overline{\text{Map}_c(S)}$  (except when  $S$  is finite-type or the Loch Ness Monster, in which case  $\overline{\text{Map}_c(S)} = \text{Map}(S)$ ). Recall that the Loch Ness Monster surface is the unique (up to homeomorphism) infinite-genus surface with one end.

We are particularly interested in indicable groups and various ways of embedding them in mapping class groups of infinite-type surfaces. A group  $G$  is indicable if there exists a surjective homomorphism  $f: G \rightarrow \mathbb{Z}$ . We show with Lemma 4.1 that a group  $G$  is indicable if and only if there is a presentation for  $G$  where the relators all have total exponent sum zero in the generators of  $G$ . Importantly, many of our

constructions require, as an input, an indicable subgroup  $G$  of  $\text{Map}(S)$  where  $S$  is a surface with exactly one boundary component. There are many examples of such groups that were mentioned in Section 1, but there are also some restrictions on what groups  $G$  can arise in this way, as is evidenced by the following lemma, which generalizes the same result from the finite-type setting [FM12, Corollary 7.3].

**Lemma 2.4** (Corollary 3, [ACCL20]). *If  $S$  is an orientable infinite-type surface with nonempty compact boundary, the mapping class group fixing the boundary pointwise is torsion-free.*

**2.3. Push and shift maps.** In this section, we define shift maps and push maps, which are central to all of our constructions. The first definition of a shift map was for handle shifts, defined in [PV18]. This inspired the following definition from Abbott–Miller–Patel [AMP]. A similar definition of shift maps appears in [MR19] and [LL20].

**Definition 2.5.** Let  $D_\Pi$  be the surface defined by taking the strip  $\mathbb{R} \times [-1, 1]$ , removing an open disk of radius  $\frac{1}{4}$  with center  $(n, 0)$  for  $n \in \mathbb{Z}$ , and attaching any fixed topologically nontrivial surface  $\Pi$  with exactly one boundary component to the boundary of each such disk. A shift on  $D_\Pi$  is the homeomorphism that acts like a translation, sending  $(x, y)$  to  $(x+1, y)$  for  $y \in [-1+\epsilon, 1-\epsilon]$  and which tapers to the identity on  $\partial D_\Pi$ .

Given a surface  $S$  with a proper embedding of  $D_\Pi$  into  $S$  so that the two ends of the strip correspond to two different ends of  $S$ , the shift on  $D_\Pi$  induces a shift on  $S$ , where the homeomorphism acts as the identity on the complement of  $\overline{D_\Pi}$ . If instead we have a proper embedding of  $D_\Pi$  into  $S$  where the two ends of the strip correspond to the same end, we call the resulting homeomorphism on  $S$  a one-ended shift. Given a shift or one-ended shift  $h$  on  $S$ , the embedded copy of  $D_\Pi$  in  $S$  is called the domain of  $h$ . By abuse of notation, we will sometimes say that the domain of the shift or one-ended shift  $h$  is  $D_\Pi$  rather than referring to it as an embedded copy of  $D_\Pi$  in  $S$  (when it is clear from context to which embedded copy of  $D_\Pi$  we are referring).

**Remark 2.6.** Given a shift map  $h$  corresponding to an embedding of  $D_\Pi$  into a surface  $S$ , one can define a new and distinct shift map  $h'$  on  $S$  by omitting some of the surfaces  $\Pi_i$  from the domain, as long as infinitely many remain. This gives another embedding of  $D_\Pi$  into  $S$ . Since there are uncountably many infinite subsets of  $\mathbb{Z}$ , we can construct uncountably many distinct embeddings of  $D_\Pi$  into  $S$ , and thus uncountably many distinct shift maps on  $S$ , in this way. The same argument goes through for one-ended shifts as well.

**Remark 2.7.** If the surface  $\Pi$  in Definition 2.5 has a nontrivial end space, then a shift or one-ended shift  $h$  on  $S$  with domain  $\overline{D_\Pi}$  is not in  $\text{PMap}(S)$  since there are ends of  $S$  that are not fixed by  $h$ . Thus,  $h \notin \overline{\text{Map}_c(S)}$  and is of intrinsically infinite type. On the other hand, if  $h$  is a shift map and if  $\Pi$  is a finite-genus surface with no planar ends, then  $h$  is a power of a handle shift on  $S$ , and the proof of [PV18, Proposition 6.3] again tells us that  $h \notin \overline{\text{Map}_c(S)}$ . However, the second conclusion does not hold when  $h$  is a one-ended handle shift since, in that case, it follows from work in [PV18] that  $h \in \overline{\text{Map}_c(S)}$ .




FIGURE 1. A surface  $S$  that admits a shift whose domain is an embedded copy of  $D_\Pi$ .

We now use the construction of shift maps to introduce finite shifts. These are constructed in a completely analogous way, starting with an annulus instead of a biinfinite strip.

**Definition 2.8.** Let  $A_\Pi$  be the surface defined by taking the annulus

$$([0, n]/0 \sim n) \times [-1, 1],$$

removing an open disk of radius  $\frac{1}{4}$  centered at the integer points, and attaching any fixed topologically nontrivial surface  $\Pi$  with exactly one boundary component to the boundary of each disk. A finite shift on  $A_\Pi$  is the homeomorphism that acts like a translation, sending  $(x, y)$  to  $(x+1, y)$  (modulo  $n$ ) for  $y \in [-1+\epsilon, 1-\epsilon]$  and which tapers to the identity on  $\partial A_\Pi$ . Given a surface  $S$  with a proper embedding of  $A_\Pi$  into  $S$ , the finite shift on  $A_\Pi$  induces a finite shift on  $S$ , where the homeomorphism acts as the identity on the complement of  $A_\Pi$ . We call the embedded copy of  $A_\Pi$  the domain of the finite shift.

**Definition 2.9.** A push is any map that is a finite shift, a one-ended shift, or a shift map.

In Section 3, we will introduce the notion of a multipush once we have developed some further notation and language.

### 3. CONSTRUCTION OF SUBGROUPS OF BIG MAPPING CLASS GROUPS

In this section, we begin by constructing a class of surfaces based on an underlying graph. In section 3.2, we show how to use this construction to find free subgroups of many big mapping class groups. In sections 3.3, and 3.4, we construct certain wreath products, and solvable Baumslag-Solitar groups as subgroups of many big mapping class groups. In all cases, the subgroups  $G$  we construct are intrinsically infinite type (except when  $S$  is finite-type or the Loch Ness Monster) and there is no hyperbolic metric for which these isomorphic copies of  $G$  in  $\text{Map}(S)$  are subgroups of the isometry group of  $S$ .

#### 3.1. A construction of surfaces.

**Definition 3.1.** A  $d$ -leg pants is a surface which is homeomorphic to a  $(d+1)$ -holed sphere. Recall that the usual pair of pants is a 2-leg pants.

**Definition 3.2.** A set of seams on a  $d$ -leg pants is a collection of  $d + 1$  disjointly embedded arcs such that each boundary component of the pants intersects exactly two components of the seams at two distinct points and such that the seams cut each pants into two components. Call one component the front side and the other component the back side. Note that these conditions imply that each component is homeomorphic to a disk.

Now, starting from any graph  $\Gamma$  with a countable vertex set, we describe a procedure for building a surface  $S_\Gamma$  using  $\Gamma$  as a framework. This mirrors a construction of Allcock in [All06] using the Cayley graph of a given group  $G$ . Fix a surface  $\Pi$  with exactly one boundary component.

For each vertex  $u$  of valence  $d + 1$ , start with a  $d$ -leg pants. Remove a disk on the interior of the front side, and attach the surface  $\Pi$  along the boundary component. Call the resulting surface the vertex surface for  $u$ , which we denote by  $V_u$ . For each edge of the graph, define the edge surface  $E$  to be the 1-leg pants with seams (topologically this is an annulus).

For each vertex, we take its corresponding vertex surface. Whenever  $u$  and  $v$  are two vertices of  $\Gamma$  connected by an edge, then connect the vertex surfaces  $V_u$  and  $V_v$  with an edge surface  $E(u, v)$  by gluing the first boundary component of the edge surface to a boundary component of  $V_u$  and the second boundary component of the edge surface to a boundary component of  $V_v$  so that the gluing is compatible in the following sense: the union of the seams separates  $S_\Gamma$  into two disjoint connected components, the front and the back. Call the resulting surface  $S_\Gamma(\Pi)$ . In  $S_\Gamma(\Pi)$ , we let  $\Pi_v$  be the copy of  $\Pi$  on the vertex surface  $V_v$ . See Figure 2 for an example. Notice that the assumption that the vertex set  $V(\Gamma)$  of  $\Gamma$  is countable is necessary for this construction to yield a surface. In particular, if  $V(\Gamma)$  is uncountable, then  $S_\Gamma(\Pi)$  is not second countable, and therefore cannot be a surface.

In this paper, we will work with the surfaces  $S_\Gamma(\Pi)$  as well as a more general class of surfaces constructed by editing the back of  $S_\Gamma(\Pi)$  as follows. Fix a graph  $\Gamma$  with a countable vertex set and a surface with one boundary component  $\Pi$ , and let  $S = S_\Gamma(\Pi)$ . Given any collection of surfaces  $\{\Omega_v\}_{v \in V(\Gamma)}$ , only finitely many of which have boundary, we form the surface  $S \#_{v \in V(\Gamma)} \Omega_v$  as follows. For each  $v \in V(\Gamma)$ ,

take the connect sum of  $V_v$  and the corresponding  $\Omega_v$ , where we assume that the connect sum is done on the back of  $V_v$ . We note that if every  $\Omega_v$  is a sphere, then  $S \#_{v \in V(\Gamma)} \Omega_v$  is homeomorphic to  $S$ . On the other hand, by choosing the  $\Omega_v$

to be more complicated, we can change the homeomorphism type of the surface by changing the genus or the space of ends. Thus, even for a fixed surface  $\Pi$ , this construction will result in a large family of surfaces, formed by varying the  $\Omega_v$ .

Another way to modify the surfaces that our constructions produce is to take multiple surfaces  $S_{\Gamma_i}(\Pi_i)$ , all of which have a subgroup isomorphic to  $G$ , and take their connect sum, done on the backs of the surfaces. The resulting surface will have the individual copies of  $G$  in its mapping class group, as well as a copy of  $G$  with larger support coming from the diagonal action.

The underlying graph  $\Gamma$  used to build  $S_\Gamma(\Pi)$  throughout this paper will often be a Schreier graph, which is defined as follows. Let  $G$  be a finitely generated group,  $H$  a subgroup of  $G$ , and  $T$  a finite generating set for  $G$ . The Schreier graph  $\Gamma(G, T, H)$  is the graph whose vertices are the left cosets of  $H$  and in which, for each pair of a coset  $gH$  and a generator  $s \in T$ , there is an edge from  $gH$  to  $sgH$  labeled by  $s$ .




FIGURE 2. An example of the surface  $S_\Gamma(\Pi)$  for the group  $\mathbb{Z}^2 = \langle a, b \rangle$  acting on its Cayley graph  $\Gamma = \Gamma(\mathbb{Z}^2, \{a, b\})$ .

Note that if  $gH = sgH$ , there is a loop labeled  $s$  at the vertex corresponding to  $gH$ . Our assumption on the finiteness of  $T$  ensures that  $\Gamma(G, T, H)$  has a countable vertex set. When  $\Gamma$  is a Schreier graph, we let  $\Pi_{gH}$  be the copy of  $\Pi$  on the vertex surface corresponding to the coset  $gH$ . In the special case when  $H = \{1\}$ , the Schreier graph  $\Gamma(G, T, \{1\})$  is simply the Cayley graph of  $G$  with respect to the generating set  $T$ , which we denote  $\Gamma(G, T)$ . There is a natural action of  $G$  on its Schreier graph given by  $k(gH) = (kg)H$ , for any coset  $gH$  and  $k \in G$ .

We are now ready to define a multipush on an infinite-type surface  $S_\Gamma(\Pi) \#_{v \in V(\Gamma)} \Omega_v$ .

**Definition 3.3.** Let  $G$  be a group and  $\Gamma = \Gamma(G, T, H)$  be a Schreier graph. Fix a surface  $\Pi$  with exactly one boundary component. Let  $S = S_\Gamma(\Pi) \#_{v \in V(\Gamma)} \Omega_v$  for any collection of surfaces  $\Omega_v$ , all but finitely many of which are without boundary. For each generator  $s \in T$ , fix a transversal  $\mathcal{T}$  for the subgroup  $\langle s \rangle$  in  $G$ . For each element  $t$  in the transversal we define a push  $h_{\langle s \rangle t}$  which maps  $\Pi_{s^i t H}$  to  $\Pi_{s^{i+1} t H}$ . The support of  $h_{\langle s \rangle t}$  is contained in front of

$$\left( \bigcup_{i \in \mathbb{Z}} V_{s^i t H} \right) \cup \left( \bigcup_{i \in \mathbb{Z}} E(s^i t H, s^{i+1} t H) \right).$$

The multipush  $x_s$  associated to  $s$  is the element of  $\text{Map}(S)$  that acts simultaneously as the pushes  $h_{\langle s \rangle t}$  for each  $t \in \mathcal{T}$ . We let  $D_s$  denote the domain of the multipush  $x_s$ . See Figure 3.




FIGURE 3. A portion of the domain  $D_a$  (in blue) of the multipush  $x_a$  on the surface  $S_\Gamma(\Pi)$ , where the group  $\mathbb{F}_2 = \langle a, b \rangle$  acts on its Cayley graph  $\Gamma = \Gamma(\mathbb{F}_2, \{a, b\})$ .

**Remark 3.4.** Just as in Remark 2.6, one multipush  $x_s$  associated to a generator  $s$  on a surface  $S$  can be used to produce uncountably many distinct domains for multipushes associated to  $s$  by simply removing some of the copies of  $\Pi$  from the domain of the multipush. One can picture this by moving some of the copies of  $\Pi$  in  $S$  to the back of  $S$ .

**Remark 3.5.** One of the key features of our subgroup constructions below is that they utilize push and multipush maps. Notice that if the complement of the domain of a push or multipush map is topologically nontrivial, the map cannot act as an isometry for any hyperbolic metric on  $S$ .

Given Remark 2.6, we have that for any surface admitting one shift map there are uncountably many other shift maps such that the complement of their domains are topologically nontrivial since they, in particular, contain some copies of the surface  $\Pi$  in  $S$ . This argument also works for one-ended shifts, and using Remark 3.4 in place of Remark 2.6 gives the same result for multipushes. However, this argument does not apply to finite shifts, as the copies of  $\Pi$  in the domain are no longer in bijection with  $\mathbb{Z}$ .

Many results in the remainder of this section (and later sections) apply to the mapping class group of a surface which contains an embedded copy of  $D_\Pi$  for an appropriate surface with one boundary component  $\Pi$ . Such surfaces include  $S = S_\Gamma(\Pi) \#_{v \in V(\Gamma)} \Omega_v$  for any graph  $\Gamma$  which contains a bi-infinite path and any collection of surfaces  $\{\Omega_v\}$ , only finitely many of which have boundary. By choosing

the collection  $\{\Omega_v\}$  carefully, we can often ensure that the resulting surface  $S$  has a non-displaceable subsurface, and hence its isometry group (with respect to any hyperbolic metric) contains only finite groups [APV]. In particular, the groups we construct could not arise from a construction using isometries for any such surface.

**3.2. Free groups.** The construction of  $S_\Gamma(\Pi)$  from Section 3.1 and Theorem 3.7 below were motivated by the following construction of a free subgroup of intrinsically infinite type.

**Example 3.6.** Let  $\Gamma$  be the Cayley graph of the free group  $\mathbb{F}_2 = \langle a, b \rangle$ , which is the Schreier graph  $\Gamma(\mathbb{F}_2, \{a, b\}, \{1\})$ , and build  $S = S_\Gamma(\Pi)$  with  $\Pi$  a torus with one boundary component. Then the resulting surface is homeomorphic to the blooming Cantor tree surface, that is, the surface with no boundary components and a Cantor set of nonplanar ends. The multipushes  $x_a$  and  $x_b$  generate a copy of  $\mathbb{F}_2$  in  $\text{PMap}(S)$ . To see this, observe that for any  $g \in \langle a, b \rangle$ , the multipush  $x_a$  maps  $\Pi_g$  to  $\Pi_{ag}$ , and similarly for  $x_b$ . Thus, the only way for a word  $w \in \langle x_a, x_b \rangle$  to act trivially on the surface is if the corresponding word in  $\langle a, b \rangle$  is trivial. Moreover, Remark 2.7 shows that this copy of  $\mathbb{F}_2$  in  $\text{PMap}(S)$  is not contained in  $\overline{\text{Map}_c(S)}$ .

This example is simplified by the fact that  $\mathbb{F}_2$  has no relations and  $\Gamma$  is a tree, so we only need to track where  $\Pi_{\text{id}}$  is mapped. Generalizing this example to other groups, we have the following theorem, from which Proposition 1.4 follows immediately.

**Theorem 3.7.** *Let  $\Gamma$  be a Schreier graph for a triple  $(G, T, H)$ . Then the set  $\{x_\alpha \mid \alpha \in T\}$  generates a free group of rank  $|T|$  in  $\text{Map}(S)$ , where  $S = S_\Gamma(\Pi) \#_{v \in V(\Gamma)} \Omega_v$*

*for any topologically nontrivial surface  $\Pi$  with exactly one boundary component, and any collection of surfaces  $\{\Omega_v\}$ , only finitely many of which have boundary. If  $|T| = 1$ , then we require that at least one  $\Omega_v$  is not a sphere.*

*Moreover, if  $S = S_\Gamma(\Pi)$  is not a finite-type surface, there exist uncountably many copies of such a free group in  $\text{Map}(S)$ , none of which can lie entirely in the isometry group for any hyperbolic metric on  $S$ . If  $S$  is neither a finite-type surface nor the Loch Ness monster surface, none of these isomorphic copies of  $\mathbb{F}_{|T|}$  can be completely contained in  $\overline{\text{Map}_c(S)}$ .*

*Proof.* Let  $w = t_1 \dots t_k$  be a non-trivial reduced word in the generating set  $T$  and let  $x_w := x_{t_1} \dots x_{t_k}$ . We aim to show that  $x_w$  is nontrivial. If  $w \notin H$ , then as in Example 3.6, we see  $x_w \Pi_H = \Pi_{wH} \neq \Pi_H$  and therefore  $x_w \neq 1$ .

Now suppose  $w = t_1 \dots t_k$  represents an element in  $H$  so that  $x_w \Pi_H = \Pi_H$ . Let  $\gamma$  be a simple closed curve given by the intersection of the vertex surface  $V_H$  for the identity coset of  $H$ , and the edge surface  $E(H, t_k H)$  for the generator  $t_k$ . We claim that  $\gamma$  and  $x_w \gamma$  bound surfaces with nontrivial topology. In the case where  $|T| \geq 2$ , Figure 4 demonstrates this phenomenon, where the cycle in  $\Gamma$  is of length 4. Note that if  $G$  is one-generated and  $\Gamma$  is infinite, then  $\Gamma(G, T, H) = \Gamma(\mathbb{Z}, \{1\}, \{\text{id}\})$  is the Cayley graph of  $\mathbb{Z}$  with its standard generator, and so there are no nontrivial words in  $H = \{\text{id}\}$ . Thus, if  $G$  is one-generated then  $\Gamma$  is a finite cycle. In this case, the requirement that some  $\Omega_i$  is not a sphere guarantees that  $\gamma$  and  $x_w \gamma$  bound surfaces with non trivial topology. We see that  $\gamma$  and  $x_w \gamma$  are not homotopic, and conclude that  $x_w \neq 1$ .

Note that if we apply Remark 3.4 to the multipush  $x_t$  for some  $t \in T$ , then this actually removes those copies of  $\Pi$  from the domains of  $x_t$  for all  $t \in T$ , and so




FIGURE 4. A portion of a Schreier graph with the curve  $\gamma$  from the proof of Theorem 3.7 in blue. The four figures show the intermediate steps for computing  $x_w \gamma$ , which is shown in red in (D).

the construction is still well-defined. Thus Remark 3.4 yields uncountably many distinct multipush domains for the argument above, and therefore uncountably many distinct embeddings of  $\mathbb{F}_{|T|}$  in  $\text{Map}(S)$ , unless the surface  $S = S_\Gamma(\Pi)$  is finite-type. In the case where  $S$  is finite-type, the underlying graph  $\Gamma$  is finite, so the multipushes are disjoint unions of finite shift maps for which Remark 3.4 does not apply.

The fact that the copies of  $\mathbb{F}_{|T|}$  cannot lie in the isometry group for any hyperbolic metric on  $S$  follows from Remark 3.5. Additionally, when  $S$  is not finite-type or the Loch Ness Monster (in which case  $\overline{\text{Map}_c(S)} = \text{Map}(S)$ ), each multipush in the argument above is a collection of shift maps so that Remark 2.7 gives us the last conclusion of the theorem.  $\square$

The free groups resulting from Theorem 3.7 are not normal subgroups of the mapping class group of  $S$ . For example, when  $\Pi$  is not a punctured disk, we can see this by choosing a simple closed curve  $\delta$  on  $\Pi_H$  and considering the Dehn twist about it,  $T_\delta$ . Because  $T_\delta \circ x_\alpha \circ T_\delta^{-1}$  is not in the group  $\langle x_\alpha \rangle$  we see that the group is not normal.




FIGURE 5. The shifts  $h_a$  and  $h_b$  do not generate a free group.

3.2.1. *Shift Maps that do not generate a free group.* Our construction above uses a countable collection of intersecting shift maps to ensure the resulting group is free. The following example demonstrates why this is necessary by showing that the group generated by two shift maps with minimal intersection is not free. We use the convention that  $[x, y] = xyx^{-1}y^{-1}$ .

Let  $\Gamma$  be the four-ended tree with a single vertex of valence four and all other vertices of valence two. Identify  $\Gamma$  with the coordinate axes in  $\mathbb{R}^2$  to get a labeling of the vertices as integer coordinates. Let  $\Pi$  be any surface with one boundary component that is not a disk, and construct the surface  $S = S_\Gamma(\Pi)$ . There is a horizontal shift,  $h_a$ , corresponding to the  $+(1, 0)$  map on the  $x$ -axis and a vertical shift,  $h_b$  corresponding to  $+(0, 1)$  on the  $y$ -axis, as shown in Figure 5. The intersection of the supports of these shifts is contained in the front of  $V_{(0,0)}$ . It can be checked that the support of  $[h_a, h_b]$  is contained in the fronts of  $V_{(1,0)}$ ,  $V_{(0,0)}$ , and  $V_{(0,1)}$ . The word  $w = h_a^2 h_b^{-1} h_a$  maps  $\{\Pi_{(0,1)}, \Pi_{(0,0)}, \Pi_{(1,0)}\}$  to the collection  $\{\Pi_{(2,0)}, \Pi_{(3,0)}, \Pi_{(4,0)}\}$ . Thus, the elements  $[h_a, h_b]$  and  $w[h_a, h_b]w^{-1}$  have disjoint supports and commute. More generally, the words  $w_n = h_a^{3n+2} h_b^{-1} h_a$  map  $\{\Pi_{(0,1)}, \Pi_{(0,0)}, \Pi_{(1,0)}\}$  to  $\{\Pi_{(2+3n,0)}, \Pi_{(3+3n,0)}, \Pi_{(4+3n,0)}\}$ . In this way, not only do we see that  $H := \langle h_a, h_b \rangle$  is not a free group, but that it actually contains copies of  $\mathbb{Z}^n$  for all  $n$ .

In fact, more can be said about the group  $H$ : it is isomorphic to a 2-generated subgroup of an infinite strand braid group. To see this, note that the group structure of  $H$  is not dependent on the surface  $\Pi$  that we attach, so we may assume  $\Pi$  is a punctured disk. We can also realize the shift domains as a disk, where the punctures in each shift domain have two distinct accumulation points on the boundary. Because braid groups are mapping class groups of punctured disks, this view point allows us to realize  $H$  as a subgroup of the infinite strand braid group in which braids are allowed to have non-compact support. In particular,  $H$  is isomorphic to the subgroup of this braid group generated by the elements  $h_a$  and  $h_b$ , viewed as braids with non-compact support.

**3.3. Wreath products.** We now generalize the construction of  $\mathbb{Z} \wr \mathbb{Z}$  in Lanier–Loving [LL20] to construct wreath products in big mapping class groups. In particular, when  $G$  is chosen to be the infinite cyclic group generated by a single Dehn twist, we recover [LL20, Theorem 4].

**Proposition 3.8.** *Let  $G \leq \text{Map}(\Pi)$  where  $\Pi$  is a surface with a single boundary component. If  $S$  is a surface that admits a shift whose domain is an embedded copy of  $D_\Pi$ , then the restricted wreath product  $G \wr \mathbb{Z}$  is a subgroup of  $\text{Map}(S)$ . Moreover, there are uncountably many copies of  $G \wr \mathbb{Z}$  in  $\text{Map}(S)$  that are not contained in  $\overline{\text{Map}_c(S)}$  and cannot be realized as subgroups of the isometry group for any hyperbolic metric on  $S$ .*

*Proof.* Recall that

$$G \wr \mathbb{Z} = \bigoplus_{-\infty}^{\infty} G \rtimes_{\gamma} \mathbb{Z},$$

that is, the semidirect product of  $\mathbb{Z}$  with the direct sum of countably many copies of  $G$ , where the action of  $\mathbb{Z}$  is by shifting the coordinates. In particular, indexing the copies of  $G$  in the direct sum by  $i \in \mathbb{Z}$ , we have that  $\gamma: \mathbb{Z} = \langle t \rangle \rightarrow \text{Aut}(\bigoplus G)$  is defined by  $\gamma(t)(G_i) = tG_it^{-1} = G_{i+1}$ .

Let  $G_i$  be a copy of  $G$  whose support is contained in  $\Pi_i$ , the surface at the  $i$ -index of  $D_\Pi$ , and let  $h$  be the shift with domain  $D_\Pi$ . As  $h^k$  takes the surface  $\Pi_0$  to  $\Pi_k$  for each  $k \in \mathbb{Z}$ , we see  $h^k G_0 h^{-k} = G_k$ . For  $k \neq \ell$ , the groups  $G_k$  and  $G_\ell$  have disjoint support and therefore commute. Thus

$$\langle h^k G_0 h^{-k} \rangle \cong \bigoplus_{-\infty}^{\infty} G.$$

Moreover, we have that  $\langle h \rangle \cap \langle h^k G_0 h^{-k} \rangle = \langle h \rangle \cap \langle G_k \rangle = \{1\}$ . We conclude that  $\langle G_0, h \rangle = G \wr \mathbb{Z} \leq \text{Map}(S)$ . The fact that there are uncountably many copies of  $G \wr \mathbb{Z}$  in  $\text{Map}(S)$  follows from Remark 2.6, and that these copies cannot lie in the isometry group for any hyperbolic metric on  $S$  from Remark 3.5. The fact that none of these copies are contained in  $\overline{\text{Map}_c(S)}$  follows from Remark 2.7.  $\square$

**3.4. Solvable Baumslag-Solitar groups.** In our third and final construction, we focus on solvable Baumslag-Solitar groups. Fixing a positive integer  $n$ , recall that the Baumslag-Solitar group  $BS(1, n)$  is the group with presentation

$$BS(1, n) = \langle a, t \mid tat^{-1}a^{-n} \rangle.$$

**Theorem 3.9.** *Let  $\Pi$  be a Cantor tree surface with one boundary component. If  $S$  is a surface which admits a shift with domain  $D_\Pi$ , then  $BS(1, n) \leq \text{Map}(S)$  for all  $n > 0$ . Moreover, for each  $n$  there are uncountably many copies of  $BS(1, n)$  in  $\text{Map}(S)$  that are not contained in  $\overline{\text{Map}_c(S)}$  and cannot be realized as subgroups of the isometry group for any hyperbolic metric on  $S$ .*

*Proof.* We will first construct a collection of homeomorphisms of  $\Pi$ . We consider  $\Pi$  to be the sphere  $\mathbb{S}^2$  with a Cantor set  $\mathcal{C}$  of planar ends on the equator and a small disk removed around the north pole. For each  $k \in \mathbb{Z}$ , we will define a collection of simple closed curves which divide  $\mathcal{C}$  into clopen sets. When  $k = 0$ , define an arbitrary countable collection of disjoint clopen sets of  $\mathcal{C}$  using a collection of simple closed curves  $\{\alpha_i^0\}_{i \in \mathbb{Z}}$ . When  $k = 1$ , for each  $i$ , divide the punctures contained in




FIGURE 6. The most straightforward example of a surface  $S$  for which Theorem 3.9 shows that  $BS(1, n) \leq \text{Map}(S)$  for all  $n > 0$ .

$\alpha_i^0$  into  $n$  clopen sets using simple closed curves  $\alpha_{i,1}^1, \dots, \alpha_{i,n}^1$ . Continue in this manner for all  $k \geq 2$ . When  $k = -1$ , for each  $i \equiv 0 \pmod n$ , let  $l = i/n$  and let  $\alpha_l^{-1}$  be a simple closed curve such that  $\alpha_l^{-1}, \alpha_i^0, \alpha_{i+1}^0, \dots, \alpha_{i+n-1}^0$  cobound an  $(n+1)$ -holed sphere. Thus,  $\alpha_l^{-1}$  groups together the ends that are cut away by  $\alpha_i^0, \alpha_{i+1}^0, \dots, \alpha_{i+n-1}^0$ . Continue in this manner for all  $k \leq -2$ .

We now define an element of  $\text{Map}(\Pi)$  for each  $k \in \mathbb{Z}$ . The mapping class  $\phi_0$  is the shift that sends  $\alpha_i^0$  to  $\alpha_{i+1}^0$  for all  $i \in \mathbb{Z}$ . The mapping class  $\phi_1$  is the shift that sends  $\alpha_{i,j}^1$  to  $\alpha_{i+1,j+1}^1$  when  $1 \leq j < n$  and  $\alpha_{i,n}^1$  to  $\alpha_{i+1,1}^1$ . Define  $\phi_k$  when  $k \geq 2$  analogously. The mapping class  $\phi_{-1}$  is the shift that sends  $\alpha_l^{-1}$  to  $\alpha_{l+1}^{-1}$ . Define  $\phi_k$  for  $k \leq -2$  analogously.

Let  $\phi \in \text{Map}(S)$  be the element which simultaneously acts as  $\phi_k$  on  $\Pi_k$  for each  $k \in \mathbb{Z}$  and as the identity elsewhere. Let  $h \in \text{Map}(S)$  be the shift whose domain is  $D_\Pi$ . See Figure 6.

Let  $f: BS(1, n) \rightarrow \text{Map}(S)$  be the map defined by  $f(a) = \phi$  and  $f(t) = h$ . We will show that  $f$  is an isomorphism onto its image, i.e.,  $\text{Map}(S)$  contains an isomorphic copy of  $BS(1, n)$ .

For each  $k \in \mathbb{Z}$ , the mapping class  $f(tat^{-1}) = h\phi h^{-1}$  first shifts  $\Pi_k$  to the left, applies  $\phi$ , which now acts as  $\phi_{k-1}$  on  $\Pi_k$ , and then shifts  $\Pi_k$  back to the right. By construction,  $\phi_{k-1}$  applied to  $\Pi_i$  is equivalent to  $\phi_k^n$  applied to  $\Pi_i$ . It follows that

$$f(tat^{-1}) = h\phi h^{-1} = \phi^n = f(a^n).$$

Therefore,  $f$  is a well-defined homomorphism.

Suppose there exists  $g \in BS(1, n)$  such that  $f(g)$  is the identity of  $\text{Map}(S)$ . Using the relation in  $BS(1, n)$ , the element  $g$  can be written as  $g = t^i a^k t^{-j}$  for some  $k \in \mathbb{Z}$  and  $i, j \in \mathbb{Z}_{\geq 0}$ . Since  $f(g) = h^i \phi^k h^{-j}$  is the identity, it must fix each  $\Pi_i$ , and so we must have  $i = j$ . Consider the surface  $\Pi_0$ . Then,  $f(g)$  first shifts  $\Pi_0$  to the left  $j$  times, applies  $\phi^k$  (which acts as  $\phi_{-j}^k$  on  $\Pi_{-j}$ ), and then shifts back to  $\Pi_0$ . The result of this is that  $f(g)$  acts as the shift  $\phi_{-j}^k$  on  $\Pi_0$ . The only way that  $f(g)$  acts as the identity on  $\Pi_0$  is if  $k = 0$ . Thus,  $g = t^i a^0 t^{-i} = 1$ , and  $f$  is injective, as desired.

Editing the domain of  $h$  as in Remark 2.6 and applying Remark 2.7 and Remark 3.5 completes the proof.  $\square$

The construction above embeds solvable Baumslag-Solitar groups into mapping class groups of certain infinite-type surfaces. This is in contrast to the finite-type case, where  $BS(1, n)$  is never a subgroup of the mapping class group. This

follows from the Tits alternative for mapping class groups of finite-type surfaces [Iva84, McC85]: every subgroup of such a mapping class group either contains a free subgroup or is virtually abelian. Since  $BS(1, n)$  is solvable, it does not contain any free subgroups, but it is also not virtually abelian.

#### 4. INDICABLE GROUPS

In this section, we give a general construction for embedding any indicable group which arises as a subgroup of a mapping class group of a surface with one boundary component into another big mapping class group in uncountably many intrinsically infinite ways. Recall that a group  $G$  is indicable if there exists a surjective homomorphism  $f: G \rightarrow \mathbb{Z}$ . We will need the following lemma in our construction.

**Lemma 4.1.** *A group  $G$  is indicable if and only if there exists a presentation  $G = \langle T \mid R \rangle$  such that for each  $r \in R$ , the total exponent sum of  $r$  with respect to the generators  $T$  is zero.*

Before presenting the proof of the lemma, we give an example that motivates the argument. Consider the Baumslag-Solitar group  $BS(1, n)$  with its standard presentation  $BS(1, n) = \langle a, t \mid tat^{-1}a^{-n} \rangle$ . This presentation does not have the desired property since the total exponent sum of the relator in the generators  $a$  and  $t$  is  $1 - n$ . However, there exists a homomorphism  $f: BS(1, n) \rightarrow \mathbb{Z}$  defined by letting  $f(a) = 0$  and  $f(t) = 1$ , so the lemma tells us that there must be a presentation of  $BS(1, n)$  with the desired property. If we augment the generator  $a$  to be  $at$  instead, then

$$BS(1, n) = \left\langle at, t \mid (t \cdot at \cdot t^{-1} \cdot t^{-1}) \cdot \underbrace{t(at)^{-1} \cdots t(at)^{-1}}_{n \text{ times}} \right\rangle,$$

and the relator has zero total exponent sum in the generators  $at$  and  $t$ . In this presentation, the generators of  $BS(1, n)$  both map to 1 under the homomorphism  $f$ , and we will use this property in the proof of the lemma.

*Proof of Lemma 4.1.* Given a group  $G = \langle T \mid R \rangle$  with all relators having total exponent sum zero, there is a well defined homomorphism  $f: G \rightarrow \mathbb{Z}$  defined by sending each generator to  $1 \in \mathbb{Z}$ .

For the other direction, assume there exists a homomorphism  $f: G \rightarrow \mathbb{Z}$  and let  $N = \ker(f)$ . Let  $N = \langle V \mid W \rangle$  be a presentation for  $N$ , and let  $a \in G$  be such that  $f(a) = 1$ . Then since  $G/N \cong \mathbb{Z}$ ,  $G$  is generated by  $T' = \{a\} \cup V$ . If we augment the generators in  $V \subseteq T'$  by  $a$ , then  $T = \{a\} \cup \{av : v \in V\}$  is also a generating set for  $G$ . Importantly, the image of every one of these generators under  $f$  is  $1 \in \mathbb{Z}$ .

Let  $G = \langle T \mid R \rangle$  be the presentation of  $G$  for the generating set  $T$ . If  $r \in R$  is a relator in  $G$ , then  $r$  is a word in  $\langle T \rangle$  that is the identity in  $G$ . Thus,  $f(r) = 0$ , and given that every element of  $T$  maps to  $1 \in \mathbb{Z}$ , the total exponent sum of  $r$  with respect to  $T$  must be zero. Therefore,  $\langle T \mid R \rangle$  is the desired presentation for  $G$ .  $\square$

We can now begin our construction. Take any indicable group  $G$  that arises as a subgroup of  $\text{Map}(\Pi)$ , where  $\Pi$  is a surface with exactly one boundary component. For example,  $G$  could be any of the indicable groups mentioned in the introduction or constructed in Section 3, among many others. Let  $h$  be a shift map on an infinite-type surface  $S$  where the domain of the shift  $h$  is an embedded copy of  $D_\Pi$  in  $S$ . As discussed in Section 3, this includes a wide range of surfaces, including surfaces

of the type  $S = S_\Gamma(\Pi) \#_{v \in V(\Gamma)} \Omega_v$ , where  $\Gamma$  is any graph with countable vertex set and a biinfinite path, and  $\Omega_v$  is any collection of surfaces.

The most trivial way to embed  $G$  into  $\text{Map}(S)$  is to let  $G$  act on one copy of  $\Pi$  in  $S$ . Indexing the copies of  $\Pi$  in  $S$  by  $\mathbb{Z}$  and taking any subset of  $I$  of  $\mathbb{Z}$ ,  $G$  can also act simultaneously on the subsurfaces  $\Pi_i$  of  $S$  for  $i \in I$ . Varying over all subsets of  $\mathbb{Z}$  gives an uncountable collection of copies of  $G$  in  $\text{Map}(S)$ . Our construction gives a new, distinct uncountable collection of copies of  $G$ , embedded in  $S$  in a more interesting way.

**Theorem 4.2.** *Let  $G$  be an indicable group that arises as a subgroup of  $\text{Map}(\Pi)$ , where  $\Pi$  is a surface with exactly one boundary component. Given a surface  $S$  that admits a shift  $h$  whose domain is  $D_\Pi$ , there exists an uncountable collection of distinct embeddings of  $G$  into  $\text{Map}(S)$  such that no embedded copy is contained in  $\overline{\text{Map}_c(S)}$  and no embedded copy is contained in the isometry group for any hyperbolic metric on  $S$ .*

*Proof.* Let  $G$  be an indicable group. Fix a presentation  $\langle T \mid R \rangle$  of  $G$  such that each  $r \in R$  has total exponent sum zero with respect to  $T$ , which exists by Lemma 4.1. Note that since  $G$  is a subgroup of  $\text{Map}(\Pi)$ ,  $G$  acts by homeomorphisms on each  $\Pi_i$  in the domain  $D_\Pi$  for the shift  $h$ . For each  $g \in G$  we define an element  $\bar{g} \in \text{Map}(S)$ , where  $\bar{g}$  acts as  $g$  simultaneously on each  $\Pi_i$ . We claim that the group generated by  $\bar{T} = \{\bar{t}h : t \in T\}$  in  $\text{Map}(S)$  is isomorphic to  $G$ . In particular, we must show that the words in  $\langle \bar{T} \rangle$  that are the identity in  $\text{Map}(S)$  are exactly those that are the image of a relator  $r \in R$  under the map  $\phi : \langle T \rangle \rightarrow \langle \bar{T} \rangle$ , defined by  $t \mapsto \bar{t}h$  for all  $t \in T$ .

Notice that  $h$  and  $\bar{t}$  commute as elements of  $\text{Map}(S)$  so that for any word  $g \in \langle T \rangle$  with total exponent sum  $k \in \mathbb{Z}$ , the image  $\phi(g)$  can be written as  $\bar{g}h^k$ . Since any  $r \in R$  has total exponent sum zero with respect to  $T$ ,  $\phi(r) = \bar{r}$ . In particular,  $\phi(r)$  is supported on  $\bigcup_i \Pi_i$ . Since  $G$  is a subgroup of  $\text{Map}(\Pi)$ ,  $r$  must act as the identity on  $\Pi$  so that  $\bar{r}$  acts as the identity on each surface  $\Pi_i$ . Thus,  $\phi(r)$  is the identity in  $\text{Map}(S)$ .

On the other hand, let  $w$  be a nontrivial word in  $\langle \bar{T} \rangle$ . If the total exponent sum of  $w$  with respect to  $\bar{T}$  is not zero, then the subsurfaces  $\Pi_i$  of  $S$  are not fixed by  $w$  and  $w$  is not identity in  $\text{Map}(S)$ . If the total exponent sum of  $w$  with respect to  $\bar{T}$  is zero, then the support of  $w$  is exactly  $\bigcup_i \Pi_i$ . By how we defined the elements of  $\bar{T}$ ,  $w$  acts as the same homeomorphism on each of the surfaces  $\Pi_i$  so that  $w$  is the identity in  $\text{Map}(S)$  only if  $w = \bar{r}$  for some  $r \in R$ . Thus, the group  $G'$  generated by  $\bar{T}$  in  $\text{Map}(S)$  is isomorphic to  $G$ .

Any element of  $G'$  that does not have total exponent sum zero with respect to  $\bar{T}$  is not in  $\overline{\text{Map}_c(S)}$ , since it must shift the surfaces  $\Pi_i$ . We can replace  $h$  above with any of the other shift maps in the uncountable collection arising from Remark 2.6 so that there are uncountably many copies of  $G$  in  $\text{Map}(S)$ , each of which is not contained in  $\overline{\text{Map}_c(S)}$  by the construction above. The fact that these copies cannot lie in the isometry group for any hyperbolic metric on  $S$  follows from Remark 3.5.  $\square$

**Remark 4.3.** It was suggested to the authors by Mladen Bestvina that one can get around constructing the presentation in Lemma 4.1 for the indicable group  $G$  by working instead with the wreath product construction in Proposition 3.8. More

specifically, let  $f: G \rightarrow \mathbb{Z}$  be a surjection from the indicable group to  $\mathbb{Z}$ . Let  $\Pi$  be a surface with exactly one boundary component such that  $G$  arises as a subgroup of  $\text{Map}(\Pi)$  and let  $S$  be a surface which admits a shift  $h$  with domain  $D_\Pi$ . For  $g \in G$ , let  $\bar{g}$  be the element which acts as  $g$  on each  $\Pi_i$ . Then, for  $g \in G$ , define a new map  $\psi: G \rightarrow G \wr \mathbb{Z} \leq \text{Map}(S)$  via  $g \mapsto \bar{g}h^{f(g)}$ . One readily checks that this map is an injective homomorphism, observing that the restriction of the image of  $G$  to  $\bigoplus_{-\infty}^{\infty} G$  is the diagonal subgroup and so the action of  $\mathbb{Z}$  is trivial. The embedding in the proof of Theorem 4.2 is exactly this map.

This theorem applies to all subgroups constructed in Section 3. Another interesting class of examples is given by Corollary 1.3, which we restate here.

**Corollary 1.3.** *Let  $\Pi$  be an infinite-type surface with at least two non-planar ends and exactly one boundary component. Given any surface  $S$  that admits a shift whose domain is  $D_\Pi$ , there exist uncountably many embeddings of  $\text{PMap}(\Pi)$  into  $\text{Map}(S)$  that are not induced by an embedding of  $\Pi$  into  $S$ .*

The corollary is immediate from Theorem 4.2 and work of Aramayona, Vlamis and the fourth author, which shows that the pure mapping class group of any surface with at least two non-planar ends is indicable [APV20, Corollary 6]. Corollary 1.3 is in line with a body of work aiming to find interesting homomorphisms between big mapping class groups. It also gives a natural set of examples of uncountable groups  $G$  to which one can apply Theorem 4.2. We note that it is an important open question for both finite- and infinite-type surfaces to determine which full mapping class groups are indicable. We now give a few examples of such mapping class groups.

**Examples 4.4.** Mann and Rafi build continuous homomorphisms to  $\mathbb{Z}^k$  and  $\mathbb{Z}$  from finite index subgroups of mapping class groups in the proofs of [MR19, Lemma 6.7, Theorem 1.7] respectively. To find surfaces whose full mapping class groups are indicable we focus on the cases where the subgroup has index 1, a few of which we list below. We will define the homomorphism to  $\mathbb{Z}$  explicitly for example (1); the others are defined similarly.

- (1) Let  $\Sigma$  be the surface with infinite genus, no boundary components, and whose end space is homeomorphic to the two point compactification of  $\mathbb{Z}$ , that is,  $E(\Sigma) = \{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$ , where  $E^g(\Sigma) = \{\infty\}$ . Let  $A \subset E(\Sigma)$  be the subset of ends corresponding to  $-\mathbb{N}$ , and let  $B$  be the subset of ends corresponding to  $\{0\} \cup \mathbb{N}$ . This surface is colloquially called the bi-infinite flute with one end accumulated by genus, and it admits a shift domain  $D_\Pi$  where  $\Pi$  is the punctured disk. A homomorphism  $\ell: \text{Map}(\Sigma) \rightarrow \mathbb{Z}$  can be defined by

$$\ell(\phi) = |\{x \in E \mid x \in A, \phi(x) \in B\}| - |\{x \in E \mid x \in B, \phi(x) \in A\}|.$$

The map  $\ell$  counts the difference in the number of punctures mapped from negative to positive and punctures mapped from positive to negative. Note that the puncture shift mentioned above evaluates to 1 under  $\ell$ , so the map is surjective.

- (2) Let  $\Sigma$  be a surface with any number of genus, any number of boundary components, and whose end space consists of a Cantor set and  $\{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$ , where the end  $\{\infty\}$  is identified with a point in the Cantor set. The ends corresponding to  $\{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$  must all be planar or all non-planar;

the other Cantor set ends can be planar or not. The homomorphism to  $\mathbb{Z}$  is defined as above, with sets  $A = -\mathbb{N}$  and  $B = \{0\} \cup \mathbb{N}$ .

- (3) Let  $\Sigma$  be the surface with infinite genus, any number of boundary components and end space  $\mathbb{N} \cup \{\infty\}$ , where the ends corresponding to 1 and  $\infty$  are nonplanar. This surface can be visualized as the ladder surface with punctures accumulating to one end (and possibly some boundary components). Here we can similarly define a homomorphism to  $\mathbb{Z}$ , which instead counts the number of genus that are moved between halves of the surface.

The common thread in the examples above is that the two ends of the shift map are of different topological types so that no element of  $\text{Map}(S)$  can exchange the two ends. This is the key fact necessary to ensure that the map  $\ell$  above is a homomorphism of  $\text{Map}(S)$  and not of a proper subgroup of  $\text{Map}(S)$ .

The third example can be extended to uncountably many more examples by replacing one of the isolated planar ends with a disk punctured by any closed subset of the Cantor set, of which there are uncountably many. Each of the examples above can be modified to have exactly one boundary component. Thus, their full mapping class groups can be used as the input for Theorem 4.2, and we arrive at the following corollary.

**Corollary 4.5.** *Let  $\Pi$  be any of the uncountably many surfaces with exactly one boundary component for which  $\text{Map}(\Pi)$  is indicable. Given any surface  $S$  that admits a shift whose domain is  $D_\Pi$ , there exist uncountably many embeddings of  $\text{Map}(\Pi)$  into  $\text{Map}(S)$  that are not induced by an embedding of  $\Pi$  into  $S$ .*

## 5. COMBINATION THEOREM

In this section, we give a construction that takes as its input a set of indicable subgroups of mapping class groups of surfaces with one boundary component and outputs a new surface whose mapping class group contains a new indicable subgroup of intrinsically infinite type built from the original subgroups.

**Definition 5.1.** Given two subgroups  $H_1$  and  $H_2$  of groups  $G_1$  and  $G_2$ , respectively, the free product of  $G_1$  and  $G_2$  with commuting subgroups  $H_1$  and  $H_2$  is

$$(G_1, H_1) \star (G_2, H_2) := (G_1 * G_2) / \langle\langle [H_1, H_2] \rangle\rangle.$$

More generally, the free product of  $G_1, \dots, G_n$  with commuting subgroups  $H_1, \dots, H_n$  is

$$(G_1, H_1) \star \dots \star (G_n, H_n) := G_1 * \dots * G_n / \langle\langle [H_i, H_j] : i \neq j \rangle\rangle.$$

These groups are a natural interpolation between free products (where the  $H_i$  are trivial) and direct products (where  $H_i = G_i$  for all  $i$ ). Free products with commuting subgroups arise in many natural contexts; for example, graph products of groups are a special kind of free product with commuting subgroups (where  $H_i = G_i$  for some indices  $i$  and the remaining  $H_j$  are trivial).

We are interested in the case where the  $G_i$  are indicable groups and the  $H_i$  are the kernels of the maps to  $\mathbb{Z}$ .

**Lemma 5.2.** *Let  $G_1, \dots, G_n$  be indicable groups with surjective maps  $f_i: G_i \rightarrow \mathbb{Z}$ , and let  $H_i = \ker(f_i)$ . Then the group  $(G_1, H_1) \star \dots \star (G_n, H_n)$  is also indicable.*




FIGURE 7. The domains of the two multipushes  $x_a$  (red) and  $x_b$  (blue) in the proof of Theorem 1.1.

*Proof.* Let  $T_i$  be a generating set for  $G_i$ . Then there is a map  $\phi: (G_1, H_1) \star \cdots \star (G_n, H_n) \rightarrow G_1$  defined by  $\phi(t) = 1$  for each  $t \in T_i$  with  $i \neq 1$ , and  $\phi(t') = t'$  for each  $t' \in T_1$ . Here 1 is the identity element of  $G_1$ . This map  $\phi$  is a homomorphism which restricts to the identity on  $G_1$ . By post-composing  $\phi$  with  $f_1$ , we obtain the desired map  $(G_1, H_1) \star \cdots \star (G_n, H_n) \rightarrow \mathbb{Z}$ .  $\square$

We are now ready to prove Theorem 1.1, which we restate for the convenience of the reader.

**Theorem 1.1.** *Let  $G_i$  be indicable groups that embed in  $\text{Map}(S_i)$ , for  $i = 1, \dots, n$ , where  $S_i$  is a surface with exactly one boundary component. For each  $i$ , fix a surjective map  $f_i: G_i \rightarrow \mathbb{Z}$ , and let  $H_i$  be the kernel of  $f_i$ . Let  $\Pi$  be the surface obtained from an  $(n+1)$ -holed sphere by gluing each  $S_i$  to one of its boundary components. Let  $\Gamma$  be the standard Cayley graph of  $\mathbb{F}_n$ , and let  $S = S_\Gamma(\Pi)$ . Then the indicable group  $(G_1, H_1) \star \cdots \star (G_n, H_n)$  embeds in  $\text{Map}(S)$ .*

Note that  $G_i$  (and therefore  $H_i$ ) is a subgroup of  $\text{Map}(\Pi)$  for each  $i = 1, \dots, n$ . We prove the theorem for  $n = 2$  for simplicity of notation, but the same proof works for all  $n$ . Additionally, the proof of Theorem 1.1 will demonstrate that the theorem

actually applies to a much wider class of surfaces than simply  $S_\Gamma(\Pi)$  where  $\Gamma$  is the standard Cayley graph for  $\mathbb{F}_n$ .

**Corollary 5.3.** *Let  $G_i$ ,  $H_i$ , and  $\Pi$  be as in Theorem 1.1, but now let  $\Gamma$  be a Schreier graph for any  $n$ -generated group  $G$ . Then, the indicable group  $(G_1, H_1) \star \cdots \star (G_n, H_n)$  embeds in  $\text{Map}(S)$ , where  $S = S_\Gamma(\Pi)$ .*

*Proof of Theorem 1.1.* Let  $\mathbb{F}_2 = \langle a, b \rangle$ . By construction,  $S$  admits two multipushes  $x_a$  and  $x_b$ , where each acts simultaneously as shifts whose domains correspond to translates of the axes of  $a$  and  $b$ , respectively. See Figure 7.

Let  $G_i = \langle T_i \mid R_i \rangle$  be the presentation of  $G_i$  such that each  $r \in R_i$  has total exponent sum zero with respect to  $T_i$ , coming from Lemma 4.1 for  $i = 1, 2$ . Similarly to Theorem 4.2, for each  $g \in G_i$  we define an element  $\bar{g} \in \text{Map}(S)$ , where  $\bar{g}$  acts as  $g$  simultaneously on each copy of  $\Pi$  in  $S$ . Importantly, for  $i = 1, 2$ , elements  $g_i \in G_i$  act on the copies of  $S_i$  in  $\Pi$ , and the copies of  $S_1$  and  $S_2$  in each copy of  $\Pi$  are disjoint. Thus,  $\bar{g}_1$  and  $\bar{g}_2$  commute for any  $g_1 \in G_1$  and  $g_2 \in G_2$ . Let  $\bar{T}_1 = \{\bar{t}x_a : t \in T_1\}$  and let  $\bar{T}_2 = \{\bar{t}x_b : t \in T_2\}$ . We claim that the group generated by  $\bar{T}_1 \cup \bar{T}_2$  in  $\text{Map}(S)$  is isomorphic to  $(G_1, H_1) \star (G_2, H_2)$ .

Let  $\langle T_1 \cup T_2 \rangle$  denote the free group on the generators  $T_1 \cup T_2$ . Let  $\phi: \langle T_1 \cup T_2 \rangle \rightarrow \langle \bar{T}_1 \cup \bar{T}_2 \rangle \leq \text{Map}(S)$  be the surjective map defined by  $t \mapsto \bar{t}x_a$  for all  $t \in T_1$  and  $t \mapsto \bar{t}x_b$  for all  $t \in T_2$ . In order to show that  $\langle \bar{T}_1 \cup \bar{T}_2 \rangle \leq \text{Map}(S)$  is isomorphic to  $(G_1, H_1) \star (G_2, H_2)$ , we must show that the kernel of  $\phi$  is generated by all relators in  $R_1 \cup R_2$  and the commutator  $[H_1, H_2]$ .

By Theorem 4.2, if  $r \in R_i$  for  $i = 1, 2$ , then  $\phi(r)$  is the identity element in  $\text{Map}(S)$  so that  $R_1 \cup R_2 \subset \ker(\phi)$ . Given  $w_i \in H_i$ , its image  $\phi(w_i)$  will fix each copy of  $\Pi$  in  $S$  for each  $i = 1, 2$ . This follows from the fact that the subgroups  $H_i$  are chosen to be the kernel of  $f_i$ , so  $w_i$  has total exponent sum zero in the generators  $T_i$ . Therefore,  $\phi(w_i)$  has total exponent sum zero with respect to  $\bar{T}_i$  and so can be written as  $\phi(w_i) = \bar{w}_i$ . Since the supports of  $w_1$  and  $w_2$  as elements of  $\text{Map}(\Pi)$  are disjoint by the construction of  $\Pi$ , the supports of  $\bar{w}_1$  and  $\bar{w}_2$  as elements  $\text{Map}(S)$  are disjoint. Thus, these elements commute and the image  $\phi(w_1 w_2 w_1^{-1} w_2^{-1}) = \bar{w}_1 \bar{w}_2 \bar{w}_1^{-1} \bar{w}_2^{-1}$  is the identity in  $\text{Map}(S)$ . It follows that  $[H_1, H_2] \subset \ker(\phi)$ .

Lastly, we show that if  $w \in \langle T_1 \cup T_2 \rangle$  is in  $\ker \phi$ , then  $w$  is in the group generated by  $R_1 \cup R_2 \cup [H_1, H_2]$ . Fix any nontrivial  $w$  in  $\langle T_1 \cup T_2 \rangle$  such that  $\phi(w)$  acts as the identity on  $S$ . Since elements of  $\bar{T}_1$  and  $\bar{T}_2$  commute with both  $x_a$  and  $x_b$ , we can write  $\phi(w) = uv$  where  $u \in \langle x_a, x_b \rangle$  and  $v$  is a word in  $\{\bar{t} : t \in T_1 \cup T_2\}$ . By Theorem 3.7, the group  $\langle x_a, x_b \rangle$  is isomorphic to  $\mathbb{F}_2$ . If  $\phi(w)$  acts trivially on  $S$ , then in particular, it must fix each copy of  $\Pi$ , and so  $u$  must be the trivial word in this  $\mathbb{F}_2$  so that  $\phi(w) = v$ . Moreover,  $w$  must have total exponent sum zero in each of  $T_1$  and  $T_2$ . Here,  $v$  is a nontrivial word in the free group generated by  $\{\bar{t} : t \in T_1 \cup T_2\}$ , given the assumption that  $w$  is nontrivial. We will now show that since  $v$  acts trivially on each copy of  $\Pi$ , then  $w$  is a product of elements in  $[H_1, H_2]$  and  $R_1 \cup R_2$ .

Since  $w$  must have total exponent sum zero in each of  $T_1$  and  $T_2$ ,  $w$  is a word in the generators  $U_1$  and  $U_2$  of  $H_1$  and  $H_2$ , respectively. If  $w$  contains a subword that is in  $R_1 \cup R_2$ , we may delete this subword from  $w$  and obtain a new word that still has total exponent sum zero in each of  $T_1$  and  $T_2$ . We do this for each such subword in  $w$  and call the resulting word  $w'$ . Thus,  $w'$  is in  $\langle H_1, H_2 \rangle$ . Moreover, the natural projection of  $w'$  to each of  $H_i$  is trivial since  $\phi(w')$  still acts as the identity on the

copies of  $S_1$  and  $S_2$  contained in each  $\Pi$  in  $S$ . Thus,  $w' \in [H_1, H_2]$ . Therefore,  $w$  is in the group generated by  $R_1 \cup R_2 \cup [H_1, H_2]$ .  $\square$

One crucial aspect of the proof above, is that the multipushes  $x_a$  and  $x_b$  generate a free group, and Corollary 5.3 now follows from Theorem 3.7.

As in Section 4, the methods of Theorem 1.1 yield an intrinsically infinite-type embedded subgroup  $(G_1, H_1) \star \cdots \star (G_n, H_n)$  in  $\text{Map}(S)$  and can be extended to yield uncountably many distinct such embedded subgroups.

**Corollary 5.4.** *In the notation of Theorem 1.1, there exists an uncountable collection of distinct embeddings of  $(G_1, H_1) \star \cdots \star (G_n, H_n)$  in  $\text{Map}(S)$ , such that each embedded copy is of intrinsically infinite type and such that no embedded copy is contained in the isometry group of  $S$  with respect to any hyperbolic metric.*

*Proof.* It follows from Remark 2.7 that  $(G_1, H_1) \star \cdots \star (G_n, H_n) \notin \overline{\text{Map}_c(S)}$ . The distinct isomorphic copies of  $(G_1, H_1) \star \cdots \star (G_n, H_n)$  in  $\text{Map}(S)$  are constructed using Remark 3.4, noting that when a copy of  $\Pi$  is moved to the back of  $S$ , it is deleted from the domain of both multipushes in the proof. The non-containment in isometry groups follows from Remark 3.5.  $\square$

**5.1. Constructing right-angled Artin groups.** In general, the free product of  $G_1$  and  $G_2$  with commuting subgroups  $H_1$  and  $H_2$  will not be finitely presented, even when the groups  $G_i$  are finitely presented. For example, consider the indicable group  $\mathbb{F}_2 = \langle a, b \rangle$  with the map to  $\mathbb{Z}$  defined by  $a \mapsto 1$  and  $b \mapsto 0$ . The kernel  $K$  of this map is the subgroup normally generated by  $[a, b]$  and  $ab^{-1}$ . In particular,  $K$  is infinitely generated, because it contains the commutator subgroup  $[\mathbb{F}_2, \mathbb{F}_2]$ . Therefore, the free product of  $G_1$  and  $G_2$  with commuting subgroups  $H_1$  and  $H_2$  where  $G_1 = G_2 = \mathbb{F}_2$  and  $H_1 = H_2 = K$  is a finitely generated but infinitely presented group in this case.

However, there are times when the free product of indicable groups with commuting subgroups is a recognizable finitely presented group. In this subsection, we describe how to use Theorem 1.1 to produce certain right-angled Artin groups  $A_\Lambda$  which embed in big mapping class groups in interesting ways. In particular, by Corollary 5.4, these groups  $A_\Lambda$  are never completely contained in  $\text{Map}_c(S)$  and can never act by isometries on the infinite-type surface.

Recall that when  $m \leq 3g - 3$ , we can embed  $\mathbb{Z}^m$  as a subgroup of  $\text{Map}(\Sigma^g)$  generated by Dehn twists about simple closed curves in a pants decomposition. We can add punctures and a boundary component to  $\Sigma^g$  that avoid the pants decomposition to get the same  $\mathbb{Z}^m$  subgroup in  $\Sigma_{1,k}^g$ . Corollary 1.2 follows immediately from Theorem 1.1 and the following proposition.

**Proposition 5.5.** *Consider surfaces  $S_i = S_{1,k_i}^{g_i}$  for  $i = 1, \dots, n$  and  $g_i, k_i \in \mathbb{Z}_{>1}$ . Then for any  $2 \leq m_i \leq 3g_i - 3$  and  $i = 1, \dots, n$ , the group*

$$(\mathbb{Z}^{m_1}, \mathbb{Z}^{m_1-1}) \star (\mathbb{Z}^{m_2}, \mathbb{Z}^{m_2-1}) \star \cdots \star (\mathbb{Z}^{m_n}, \mathbb{Z}^{m_n-1})$$

*is the right-angled Artin group defined by the graph shown in Figure 8.*

We will prove the proposition for  $n = 2$  for simplicity of notation; the general case is analogous.

*Proof.* Fix surfaces  $S_1, S_2$  as in the statement of the theorem, and let  $m_1 = m$  and  $m_2 = n$ . Let  $\mathbb{Z}^m = \langle x_1, \dots, x_m \rangle$  and  $\mathbb{Z}^n = \langle y_1 \dots y_n \rangle$ . For each group,




FIGURE 8. The defining graphs for the family of right-angled Artin groups constructed in Proposition 5.5. Each vertex in the complete subgraph  $K_{m_i-1}$  is adjacent to a single vertex as well as every vertex in each other  $K_{m_j-1}$ .

there is a surjective homomorphism to  $\mathbb{Z}$  defined by mapping every generator to 1. We write the kernels of these maps as  $H_m = \langle x_1x_2^{-1}, \dots, x_1x_m^{-1} \rangle \simeq \mathbb{Z}^{m-1}$  and  $H_n = \langle y_1y_2^{-1}, \dots, y_1y_n^{-1} \rangle \simeq \mathbb{Z}^{n-1}$ , respectively. Observe that by changes of basis, we may write the generators of  $\mathbb{Z}^m$  as  $x_1, x_1x_2^{-1}, \dots, x_1x_m^{-1}$  and the generators of  $\mathbb{Z}^n$  as  $y_1, y_1y_2^{-1}, \dots, y_1y_n^{-1}$ . Thus, the free product of  $\mathbb{Z}^m$  and  $\mathbb{Z}^n$  with commuting subgroups  $\mathbb{Z}^{m-1}$  and  $\mathbb{Z}^{n-1}$ , corresponding to  $H_m$  and  $H_n$ , respectively, is the right-angled Artin group:

$$\left\langle a_1, \dots, a_m, b_1, \dots, b_n \left| \begin{array}{l} [a_i, a_j] \ \forall i, j, \\ [b_k, b_l] \ \forall k, l, \\ [a_q, b_r] \ \forall q = 1, \dots, m-1, r = 1, \dots, n-1 \end{array} \right. \right\rangle,$$

where  $a_i$  and  $b_i$  are identified with  $x_1x_{i+1}^{-1}$  and  $y_1y_{i+1}^{-1}$  for all  $1 \leq i \leq m-1$ , respectively, and  $a_m$  and  $b_m$  are identified with  $x_1$  and  $y_1$ , respectively.

Notice that  $x_1$  is the only generator of  $\mathbb{Z}^m$  that does not commute with any generators of  $\mathbb{Z}^n$ , and similarly,  $y_1$  is the only generator of  $\mathbb{Z}^n$  that does not commute with any generators of  $\mathbb{Z}^m$ .  $\square$

REFERENCES

[ACCL20] Santana Afton, Danny Calegari, Lvzhou Chen, and Rylee Alanza Lyman. Nielsen realization for infinite-type surfaces. 2020. Preprint. arXiv: 2002.09760.

[All06] Daniel Allcock. Hyperbolic surfaces with prescribed infinite symmetry groups. *Proc. Amer. Math. Soc.*, 134(10):3057–3059, 2006.

[ALM21] Javier Aramayona, Christopher J. Leininger, and Alan McLeay. Homomorphisms between mapping class groups. 2021. Preprint. arXiv:2101.07188.

[ALS09] Javier Aramayona, Christopher J Leininger, and Juan Souto. Injections of mapping class groups. *Geometry & Topology*, 13(5):2523–2541, 2009.

[AMP] Carolyn Abbott, Nicholas Miller, and Priyam Patel. Infinite-type loxodromic isometries of the relative arc graph. In preparation.

[APV] Tarik Aougab, Priyam Patel, and Nicholas G. Vlamis. Isometry groups of infinite genus hyperbolic surfaces. *Math. Ann.* To appear. arXiv:2007.01982.

[APV20] Javier Aramayona, Priyam Patel, and Nicholas G. Vlamis. The first integral cohomology of pure mapping class groups. *Int. Math. Res. Not. IMRN*, (22):8973–8996, 2020.

- [AS13] Javier Aramayona and Juan Souto. Homomorphisms between mapping class groups. *Geometry & Topology*, 16(4):2285–2341, 2013.
- [BBF15] Mladen Bestvina, Ken Bromberg, and Koji Fujiwara. Constructing group actions on quasi-trees and applications to mapping class groups. *Publications mathématiques de l’IHÉS*, 122(1):1–64, 2015.
- [CLM12] Matt T. Clay, Christopher J. Leininger, and Johanna Mangahas. The geometry of right-angled Artin subgroups of mapping class groups. *Groups Geom. Dyn.*, 6(2):249–278, 2012.
- [FM12] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [HQR20] Camille Horbez, Yulan Qing, and Kasra Rafi. Big mapping class groups with hyperbolic actions: classification and applications. 2020. Preprint. arXiv: 2005.00428.
- [Hur76] R. Daniel Hurwitz. On the conjugacy problem in a free product with commuting subgroups. *Math. Ann.*, 221(1):1–8, 1976.
- [Iva84] N. V. Ivanov. Algebraic properties of the Teichmüller modular group. *Dokl. Akad. Nauk SSSR*, 275(4):786–789, 1984.
- [Ker23] B. v. Kerékjártó. *Vorlesungen über Topologie. I*. Springer, Berlin, 1923.
- [Ker83] Steven P. Kerckhoff. The Nielsen realization problem. *Ann. of Math. (2)*, 117(2):235–265, 1983.
- [LL20] Justin Lanier and Marissa Loving. Centers of subgroups of big mapping class groups and the Tits alternative. *Glasnik Matematički*, 55:85–91, 06 2020.
- [Log99] E. D. Logina. Residual finiteness of the free product of two groups with commuting subgroups. *Sibirsk. Mat. Zh.*, 40(2):395–407, iii, 1999.
- [McC85] John McCarthy. A “Tits-alternative” for subgroups of surface mapping class groups. *Trans. Amer. Math. Soc.*, 291(2):583–612, 1985.
- [MKS04] Wilhelm Magnus, Abraham Karrass, and Donald Solitar. *Combinatorial group theory*. Dover Publications, Inc., Mineola, NY, second edition, 2004. Presentations of groups in terms of generators and relations.
- [MR19] Kathryn Mann and Kasra Rafi. Large scale geometry of big mapping class groups. 2019. Preprint. arXiv: 1912.10914.
- [MS73] Charles F. Miller, III and Paul E. Schupp. The geometry of Higman-Neumann-Neumann extensions. *Comm. Pure Appl. Math.*, 26:787–802, 1973. Collection of articles dedicated to Wilhelm Magnus.
- [PV18] Priyam Patel and Nicholas G. Vlamis. Algebraic and topological properties of big mapping class groups. *Algebr. Geom. Topol.*, 18(7):4109–4142, 2018.
- [Ric63] Ian Richards. On the classification of noncompact surfaces. *Trans. Amer. Math. Soc.*, 106:259–269, 1963.
- [Sok14] E. V. Sokolov. On the cyclic subgroup separability of the free product of two groups with commuting subgroups. *Internat. J. Algebra Comput.*, 24(5):741–756, 2014.
- [Tie05] D. Tiedjo. Abelian subgroup separability of some one-relator groups. *Int. J. Math. Math. Sci.*, (14):2287–2298, 2005.
- [TM08] D. Tiedjo and D. I. Moldavanskii. A generalization of residual finiteness. *Lobachevskii J. Math.*, 29(2):56–59, 2008.
- [TM10] D. Tiedjo and D. I. Moldavanskii. Conjugacy separability of some one-relator groups. *Int. J. Math. Math. Sci.*, pages Art. ID 803243, 14, 2010.

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY, WALTHAM, MA 02543  
*Email address:* [carolynabbott@brandeis.edu](mailto:carolynabbott@brandeis.edu)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112  
*Email address:* [hoganson@math.utah.edu](mailto:hoganson@math.utah.edu)

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332  
*Email address:* [mloving6@gatech.edu](mailto:mloving6@gatech.edu)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112  
*Email address:* [patelp@math.utah.edu](mailto:patelp@math.utah.edu)

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS. OH 43210  
*Email address:* [skipper.26@osu.edu](mailto:skipper.26@osu.edu)