

# ANTI-CLASSIFICATION RESULTS FOR THE KAKUTANI EQUIVALENCE RELATION

MARLIES GERBER<sup>1</sup> AND PHILIPP KUNDE<sup>2</sup>

ABSTRACT. We prove that the Kakutani equivalence relation of ergodic invertible measure-preserving transformations is not a Borel set. This shows in a precise way that classification up to Kakutani equivalence is impossible. We also obtain this anti-classification result for ergodic area-preserving smooth diffeomorphisms on compact surfaces admitting a non-trivial circle action as well as real-analytic diffeomorphisms on the 2-torus.

## 1. INTRODUCTION

A fundamental theme in ergodic theory, going back to J. von Neumann [Ne32], is the isomorphism problem: classify measure-preserving transformations (MPT's) up to isomorphism. By an MPT we mean a measure-preserving automorphism of a standard non-atomic probability space. We let  $\mathcal{X}$  denote the set of all MPT's of a fixed standard non-atomic probability space  $(\Omega, \mathcal{M}, \mu)$ . Two automorphisms  $S, T \in \mathcal{X}$  are said to be *isomorphic* (written  $S \cong T$ ) if there exists  $\varphi \in \mathcal{X}$  such that  $S \circ \varphi$  and  $\varphi \circ T$  agree  $\mu$ -almost everywhere. Since there is a decomposition of each MPT into ergodic components that is preserved under isomorphism (see, for example, Chapter 5 of [VO16]), we often consider just the set  $\mathcal{E}$  of ergodic elements of  $\mathcal{X}$ . Two great successes in the classification of MPT's are the Halmos-von Neumann classification of ergodic MPT's with pure point spectrum by countable subgroups of the unit circle [HN42] and D. Ornstein's classification of Bernoulli shifts by their metric entropy [Or70]. Nonetheless, the isomorphism problem remains unsolved for general ergodic MPT's.

In contrast to the positive results in [HN42] and [Or70], a series of *anti-classification* results have appeared, starting with the work of F. Beuzenay and M. Foreman [BF96]. Anti-classification results demonstrate the complexity of the isomorphism problem. To describe some of this work, we endow  $\mathcal{X}$  with the weak topology. (Recall that  $T_n \rightarrow T$  in the weak topology if and only if  $\mu(T_n(A) \Delta T(A)) \rightarrow 0$  for every  $A \in \mathcal{M}$ .) This topology is compatible with a complete separable metric and hence makes  $\mathcal{X}$  into a Polish space. Since  $\mathcal{E}$  is a (dense)  $G_\delta$ -set in  $\mathcal{X}$ , the induced topology on  $\mathcal{E}$  is Polish as well. In contrast to the Halmos-von Neumann classification result, G. Hjorth [Hj01] showed that there is no Borel way of associating algebraic invariants to ergodic MPT's that completely determines isomorphism. Moreover,

---

<sup>1</sup> Indiana University, Department of Mathematics, Bloomington, IN 47405, USA.

<sup>2</sup> University of Hamburg, Department of Mathematics, 20146 Hamburg, Germany. P.K. acknowledges financial support from a DFG Forschungsstipendium under Grant No. 405305501.

*2020 Mathematics Subject classification:* Primary: 37A20; Secondary: 37A35, 37A05, 37C40, 03E15

*Key words:* Kakutani equivalence, anti-classification, complete analytic, smooth ergodic theory

Foreman and B. Weiss [FW04] proved that there is no generic class (that is, a dense  $G_\delta$ -set) of ergodic MPT's for which there is a Borel way of associating a complete algebraic invariant. Hjorth also proved that the isomorphism relation  $\mathcal{R}$  is itself not a Borel set when viewed as a subset of  $\mathcal{X} \times \mathcal{X}$ . However, his proof used nonergodic transformations in an essential way. In the case of ergodic MPT's, Foreman, D. Rudolph, and Weiss [FRW11] proved that  $\mathcal{R} \cap (\mathcal{E} \times \mathcal{E})$  is not Borel. One way to interpret an equivalence relation not being a Borel set is to say that there is no inherently countable technique for determining whether or not two members of the set are equivalent. (See the survey article by M. Foreman [Fo18] for more details.) Thus [FRW11] gives a precise formulation of the unclassifiability of ergodic MPT's up to isomorphism.

Recently, Foreman and A. Gorodetski [FGpp] obtained anti-classification results in a different context. They considered topological equivalence of  $C^\infty$  diffeomorphisms of a compact manifold  $M$ . Two diffeomorphisms  $f$  and  $g$  of  $M$  are said to be topologically equivalent if there is a homeomorphism  $\varphi$  of  $M$  such that  $f \circ \varphi = \varphi \circ g$ . In the case  $\dim(M) \geq 2$ , Foreman and Gorodetski proved there is no Borel function from the  $C^\infty$  diffeomorphisms to the reals that is a complete invariant for topological equivalence. Moreover, if  $\dim(M) \geq 5$ , they proved the analog of the unclassifiability result of [FRW11], that is, the set of topologically equivalent pairs of diffeomorphisms is not Borel.

When the anti-classification results in [FRW11] became known, immediately the question arose whether these results are also true if “isomorphism” is replaced by “Kakutani equivalence”. (See [We14] for some history of this question.) In this paper we answer this question affirmatively. Our arguments build on the methods in [FRW11] and also imply the anti-classification result of [FRW11].

Kakutani equivalence is the best known and most natural equivalence relation on ergodic MPT's, besides isomorphism, for which the classification problem can be considered. (Orbit equivalence is not interesting in the present context, because according to Dye's theorem, all ergodic MPT's are orbit equivalent.) Two ergodic MPT's are said to be *Kakutani equivalent* if they are isomorphic to measurable cross-sections of the same ergodic flow. Since the Kakutani equivalence relation is weaker than isomorphism, one might expect classification to be simpler. In fact, until the work of J. Feldman [Fe76], it seemed that there might be only three Kakutani equivalence classes: zero entropy, finite positive entropy, and infinite entropy. However, Feldman found examples to show that each entropy class (zero, finite positive, and infinite) contains at least two non-Kakutani equivalent ergodic MPT's. A. Katok [Ka77], and Ornstein, Rudolph, and Weiss [ORW82], independently, extended this to obtain uncountably many non-Kakutani equivalent examples in each entropy class.

A precursor to the anti-classification results for isomorphism is the observation by Feldman [Fe74] that there does not exist a Borel function from  $\mathcal{E}$  to the reals such that two transformations are isomorphic if and only if the Borel function takes the same value at these transformations. His proof is an easy application of a 0-1 law to the uncountable family of non-Bernoulli K-automorphisms constructed by Ornstein and P. Shields [OS73]. It is interesting to note that the uncountable family of automorphisms constructed in [ORW82] can be substituted for the Ornstein-Shields examples to conclude that there is no Borel function from  $\mathcal{E}$  to the reals whose values are complete invariants for Kakutani equivalence. The key common

feature of the examples in [OS73] and the examples in [ORW82] is that they are indexed by sequences of zeros and ones in such a way that two of the examples are isomorphic (in the case of [OS73]) or Kakutani equivalent (in the case of [ORW82]) if and only if the two corresponding sequences of zeros and ones agree except for possibly finitely many terms.

In the context of Kakutani equivalence for flows, M. Ratner [Ra81] introduced the Kakutani invariant and used this to prove that for distinct integers  $k, \ell > 0$ , the product of  $k$  copies of the horocycle flow with itself is not Kakutani equivalent to the product of  $\ell$  copies. More recently, A. Kanigowski, K. Vinhage, and D. Wei [KVV21] obtained an explicit formula for the Kakutani invariant for unipotent flows acting on compact quotients of semisimple Lie groups. It is an open question whether the Kakutani invariant is a complete invariant in the class of unipotent flows.

We also note that the analog of the result in [FW04] fails for Kakutani equivalence in a trivial way, because the rank one transformations form a dense  $G_\delta$ -subset of  $\mathcal{E}$ , and all rank one transformations are zero entropy loosely Bernoulli, and therefore in the same Kakutani equivalence class. In case of isomorphism, Foreman, Rudolph and Weiss [FRW11, section 10] use a theorem of J. King [Ki86] to prove that the collection of pairs  $(S, T)$  of rank one ergodic transformations such that  $S$  and  $T$  are isomorphic is a Borel subset of  $\mathcal{E} \times \mathcal{E}$ . Nonetheless, no useful structure theorem to classify rank one transformations up to isomorphism is known.

We prove that the set  $\mathcal{S}$  of pairs  $(S, T)$  in  $\mathcal{E} \times \mathcal{E}$  such that  $S$  and  $T$  are Kakutani equivalent is not a Borel set. We follow the general scheme in [FRW11], but with some notable differences. As in [FRW11] we consider the set,  $Trees$ , which consists of countable rooted trees with arbitrarily long branches. Then, analogously to the approach in [FRW11], we construct a continuous function  $\Psi : Trees \rightarrow \mathcal{E}$  such that  $\Psi(\mathcal{T})$  is Kakutani equivalent to  $\Psi(\mathcal{T})^{-1}$  if and only if  $\mathcal{T}$  has an infinite branch. According to a classical result [Ke95], the collection of  $Trees$  with one or more infinite branches is a complete analytic subset of  $Trees$ . It follows that the set  $\mathcal{S}$  of Kakutani equivalent pairs is a complete analytic set and hence not Borel. In fact, by an argument similar to that in [FRW11],  $\Psi(\mathcal{T})$  is actually *isomorphic* to  $\Psi(\mathcal{T})^{-1}$  if  $\mathcal{T}$  has an infinite branch. Therefore any equivalence relation on  $\mathcal{E}$  that is weaker than (or equal to) isomorphism and stronger than (or equal to) Kakutani equivalence is complete analytic. This includes *even* Kakutani equivalence (see Definition 64 in Section 9.2), and  $\alpha$ -equivalence (defined in [FJR94]). Indeed, we first prove our result for even Kakutani equivalence and then use the methods of [ORW82] to prove that for any  $\mathcal{T} \in Trees$ ,  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  cannot be Kakutani equivalent unless they are evenly Kakutani equivalent.

Foreman, Rudolph, and Weiss use the elegant method of joinings to prove that for  $\mathcal{T} \in Trees$  such that  $\mathcal{T}$  has no infinite branch, the corresponding element of  $\mathcal{E}$  is not isomorphic to its inverse. However, as they announced in [FRW06], they could also have used a finite coding argument and approximations in the  $\bar{d}$ -metric for this proof. We use the finite coding approach and apply the result in [ORW82] that for evenly equivalent processes  $(S, P)$  and  $(T, Q)$ , there is a finite code defined on the  $(T, Q)$  process such that the  $\bar{f}$ -distance between the  $(S, P)$  process and the process coded from  $(T, Q)$  is small.

As in [FRW11], each level  $s$  of a tree  $\mathcal{T} \in Trees$  corresponds to an equivalence relation  $\mathcal{Q}_s$  on certain finite words, and  $\mathcal{Q}_{s+1}$  refines  $\mathcal{Q}_s$ . One of the main steps in

the construction of  $\Psi(\mathcal{T})$  is a procedure for substituting  $\mathcal{Q}_{s+1}$  equivalence classes into  $\mathcal{Q}_s$  equivalence classes. Foreman, Rudolph, and Weiss use a random substitution method, while we use Feldman patterns (defined in Section 6) to execute a deterministic substitution.

The classification problem (with respect to isomorphism or Kakutani equivalence) can also be restricted to the class of smooth ergodic diffeomorphisms of a compact manifold  $M$  that preserve a smooth measure. For dimensions greater than or equal to three, there are no known obstacles to realizing an arbitrary ergodic MPT as a diffeomorphism of a compact manifold except for the requirement, proved by A. Kushnirenko [Ku65], that the ergodic MPT have finite entropy. Thus it is not clear if the restriction to smooth ergodic diffeomorphisms changes the classification problem. But even with this restriction, Foreman and Weiss [FW3] proved that the isomorphism relation is not Borel. Moreover, S. Banerjee and P. Kunde [BKpp] showed that the isomorphism relation is still not Borel if “smooth” is replaced by “real analytic”. We obtain the analogous results for Kakutani equivalence both in the smooth and the real analytic settings.

To obtain their anti-classification results for smooth diffeomorphisms, Foreman and Weiss [FW19b] introduced a functor  $\mathcal{F}$  from certain odometer-based systems in [FRW11] to the so-called circular systems, which they showed in [FW19a] have a smooth realization on  $\mathbb{D}^2$  and  $\mathbb{T}^2$  via the untwisted Anosov-Katok method [AK70], while on  $\mathbb{T}^2$  Banerjee and Kunde [Ba17, BKpp] extended these results to the real analytic case. We also make use of this machinery to obtain smooth and real analytic models of the images under  $\mathcal{F}$  of the odometer-based systems that we construct using the Feldman patterns. The same argument as in [FW19b] shows that in our construction  $\mathcal{F}(\Psi(\mathcal{T}))$  and  $\mathcal{F}(\Psi(\mathcal{T})^{-1})$  are isomorphic if  $\mathcal{T}$  has an infinite branch. But the more difficult part, showing that  $\mathcal{F}(\Psi(\mathcal{T}))$  and  $\mathcal{F}(\Psi(\mathcal{T})^{-1})$  are *not* Kakutani equivalent if  $\mathcal{T}$  does *not* have an infinite branch, requires a different argument from that in [FW19b]. In fact, in [GKpp], we obtained an example of two *non-Kakutani* equivalent odometer-based systems  $S$  and  $T$  such that  $\mathcal{F}(S)$  and  $\mathcal{F}(T)$  are Kakutani equivalent. However, for the systems  $\Psi(\mathcal{T})$  that we construct in the present paper, if  $\Psi(\mathcal{T})$  is not Kakutani equivalent to  $\Psi(\mathcal{T})^{-1}$ , then  $\mathcal{F}(\Psi(\mathcal{T}))$  is not Kakutani equivalent to  $\mathcal{F}(\Psi(\mathcal{T})^{-1})$ . Thus, we can transfer the non-Kakutani equivalence between  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  to non-Kakutani equivalence between  $\mathcal{F}(\Psi(\mathcal{T}))$  and  $\mathcal{F}(\Psi(\mathcal{T})^{-1})$  if  $\mathcal{T}$  does not have an infinite branch. This yields smooth and real analytic anti-classification results for isomorphism (already proved in [FW3] and [BKpp]), Kakutani equivalence, and every equivalence relation in between these two.

Classification questions can also be considered in the context of group actions. For example, E. Gardella and M. Lupini [GLpp] showed that for every nonamenable countable discrete group  $\Gamma$ , the relations of isomorphism and orbit equivalence of free ergodic (or weak mixing) measure preserving actions of  $\Gamma$  on a standard probability space are not Borel. The analogous question for amenable countable discrete groups and the isomorphism relation is open.

## 2. PRELIMINARIES

**2.1. Some terminology from ergodic theory.** We call a separable non-atomic probability space  $(\Omega, \mathcal{M}, \mu)$  a *standard measure space*. Since every standard measure space is isomorphic to the unit interval with Lebesgue measure  $\lambda$  on the Borel

sets  $\mathcal{B}$ , every invertible MPT (also called an *automorphism*) of a standard measure space is isomorphic to an invertible Lebesgue measure-preserving transformation on  $[0, 1]$ . As in the introduction, we let  $\mathcal{X}$  denote the group of MPT's on  $[0, 1]$  endowed with the weak topology, where two MPT's are identified if they are equal on sets of full measure.

While in this model we fix the measure and look at transformations preserving that measure, there is also a dual viewpoint of fixing the transformation and considering the collection of its invariant measures. The transformation that we use is the left shift  $sh : \Sigma^{\mathbb{Z}} \rightarrow \Sigma^{\mathbb{Z}}$ , defined by

$$sh((x_n)_{n=-\infty}^{\infty}) = (x_{n+1})_{n=-\infty}^{\infty}$$

on  $(\Sigma^{\mathbb{Z}}, \mathcal{C})$ , where  $\Sigma$  is a countable set and  $\mathcal{C}$  is the  $\sigma$ -algebra of Borel sets in  $\Sigma^{\mathbb{Z}}$ . We endow the set of shift-invariant measures with the weak\* topology. In our paper,  $\Sigma$  is finite and for each shift-invariant measure  $\nu$  that we construct, there is also an explicit cutting-and-stacking procedure that yields an ergodic Lebesgue measure-preserving automorphism  $T$  of  $[0, 1]$  such that  $([0, 1], \mathcal{B}, \lambda, T)$  is isomorphic to  $(\Sigma^{\mathbb{Z}}, \mathcal{C}, \psi(T), sh)$ . This is a special case of a more general result of Foreman [Fo10]: there is a Borel bijection  $\psi$  between the aperiodic Lebesgue measure-preserving transformations on  $[0, 1]$  and nonatomic shift-invariant measures on  $\Sigma^{\mathbb{Z}}$  taking comeager sets to comeager sets such that  $([0, 1], \mathcal{B}, \lambda, T)$  is isomorphic to  $(\Sigma^{\mathbb{Z}}, \mathcal{C}, \psi(T), sh)$ . In particular, the choice of model for measure-preserving transformations is not important for the Borel or analytic distinction.

Another important viewpoint is to assign a process to a MPT. For this purpose, we recall that a *partition*  $P$  of a standard measure space  $(\Omega, \mathcal{M}, \mu)$  is a collection  $P = \{c_\sigma\}_{\sigma \in \Sigma}$  of subsets  $c_\sigma \in \mathcal{M}$  with  $\mu(c_\sigma \cap c_{\sigma'}) = 0$  for all  $\sigma \neq \sigma'$  and  $\mu(\bigcup_{\sigma \in \Sigma} c_\sigma) = 1$ , where  $\Sigma$  is a finite set of indices. Each  $c_\sigma$  is called an *atom* of the partition  $P$ . For two partitions  $P$  and  $Q$  we define the join of  $P$  and  $Q$  to be the partition  $P \vee Q = \{c \cap d : c \in P, d \in Q\}$ , and for a sequence of partitions  $\{P_n\}_{n=1}^{\infty}$  we let  $\bigvee_{n=1}^{\infty} P_n$  be the smallest  $\sigma$ -algebra containing  $\bigcup_{n=1}^{\infty} P_n$ . We say that a sequence of partitions  $\{P_n\}_{n=1}^{\infty}$  is a *generating sequence* if  $\bigvee_{n=1}^{\infty} P_n = \mathcal{M}$ . We also have a standard notion of a *distance between two ordered partitions*: If  $P = \{c_i\}_{i=1}^N$  and  $Q = \{d_i\}_{i=1}^N$  are two ordered partitions with the same number of atoms, then we define

$$d(P, Q) = \sum_{i=1}^N \mu(c_i \Delta d_i),$$

where  $\Delta$  denotes the symmetric difference.

For an automorphism  $T$  of  $(\Omega, \mathcal{M}, \mu)$  and a partition  $P = \{c_\sigma\}_{\sigma \in \Sigma}$  of  $\Omega$  we can define the  $(T, P)$ -name for almost every  $x \in \Omega$  by

$$(a_n)_{n=-\infty}^{\infty} \in \Sigma^{\mathbb{Z}} \text{ with } T^i(x) \in c_{a_i}.$$

Note that the  $(T, P)$ -name of  $T(x)$  is the left shift of the  $(T, P)$ -name of  $x$ . If the partition  $P$  is a *generator* (that is,  $\{T^n P\}_{n=-\infty}^{\infty}$  is a generating sequence), then  $(X, \mu, T)$  is isomorphic to  $(\Sigma^{\mathbb{Z}}, \nu, sh)$  with the measure  $\nu$  satisfying  $\nu\{(x_n)_{n=-\infty}^{\infty} : x_n = a_n \text{ for } n = j, j+1, \dots, k\} = \mu(T^{-j}(c_{a_j}) \cap T^{-(j+1)}(c_{a_{j+1}}) \cap \dots \cap T^{-k}(c_{a_k}))$ . The pair  $(T, P)$  is called a *process*.

Recall that the *Hamming distance* between two strings of symbols  $a_1 \dots a_n$  and  $b_1 \dots b_n$  is defined by  $\bar{d}(a_1 \dots a_n, b_1 \dots b_n) = \frac{1}{n} |\{i : a_i \neq b_i\}|$ . It can be extended to infinite words  $w = \dots a_{-2} a_{-1} a_0 a_1 a_2 \dots$  and  $w' = \dots b_{-2} b_{-1} b_0 b_1 b_2 \dots$  by

$\bar{d}(w, w') = \limsup_{n \rightarrow \infty} \bar{d}(w_n, w'_n)$ , where  $w_n = a_{-n}a_{-n+1} \dots a_{n-1}a_n$  and  $w'_n = b_{-n}b_{-n+1} \dots b_n$  are the truncated words.

**2.2. Symbolic systems.** An *alphabet* is a countable or finite collection of symbols. In the following, let  $\Sigma$  be a finite alphabet endowed with the discrete topology. Then  $\Sigma^{\mathbb{Z}}$  with the product topology is a separable, totally disconnected and compact space. A usual base of the product topology is given by the collection of cylinder sets of the form  $\langle u \rangle_k = \{f \in \Sigma^{\mathbb{Z}} : f \upharpoonright [k, k+n) = u\}$  for some  $k \in \mathbb{Z}$  and finite sequence  $u = \sigma_0 \dots \sigma_{n-1} \in \Sigma^n$ . For  $k = 0$  we abbreviate this by  $\langle u \rangle$ .

The shift map  $sh : \Sigma^{\mathbb{Z}} \rightarrow \Sigma^{\mathbb{Z}}$  is a homeomorphism. If  $\mu$  is a shift-invariant Borel measure, then the measure-preserving dynamical system  $(\Sigma^{\mathbb{Z}}, \mathcal{B}, \mu, sh)$  is called a *symbolic system*. The closed support of  $\mu$  is a shift-invariant subset of  $\Sigma^{\mathbb{Z}}$  called a *symbolic shift* or *sub-shift*.

The symbolic shifts that we use are described by specifying a collection of words. A word  $w$  in  $\Sigma$  is a finite sequence of elements of  $\Sigma$ , and we denote its length by  $|w|$ .

**Definition 1.** A sequence of collections of words  $(W_n)_{n \in \mathbb{N}}$ , where  $\mathbb{N} = \{0, 1, 2, \dots\}$ , satisfying the following properties is called a *construction sequence*:

- (1) for every  $n \in \mathbb{N}$  all words in  $W_n$  have the same length  $h_n$ ,
- (2) each  $w \in W_n$  occurs at least once as a subword of each  $w' \in W_{n+1}$ ,
- (3) there is a summable sequence  $(\varepsilon_n)_{n \in \mathbb{N}}$  of positive numbers such that for every  $n \in \mathbb{N}$ , every word  $w \in W_{n+1}$  can be uniquely parsed into segments  $u_0 w_1 u_1 w_1 \dots w_l u_l$  such that each  $w_i \in W_n$ , each  $u_i$  (called spacer or boundary) is a word in  $\Sigma$  of finite length and for this parsing

$$\frac{\sum_{i=0}^{l+1} |u_i|}{h_{n+1}} < \varepsilon_{n+1}.$$

We will often call words in  $W_n$  *n-words* or *n-blocks*, while a general concatenation of symbols from  $\Sigma$  is called a *string*. We also associate a symbolic shift with a construction sequence: Let  $\mathbb{K}$  be the collection of  $x \in \Sigma^{\mathbb{Z}}$  such that every finite contiguous substring of  $x$  occurs inside some  $w \in W_n$ . Then  $\mathbb{K}$  is a closed shift-invariant subset of  $\Sigma^{\mathbb{Z}}$  that is compact since  $\Sigma$  is finite.

*Remark.* The symbolic shifts built from construction sequences can be realized as transformations built by cutting-and-stacking constructions. Spacers are not used in our constructions until Section 10.

In order to be able to unambiguously parse elements of  $\mathbb{K}$  we will use construction sequences consisting of uniquely readable words.

**Definition 2.** Let  $\Sigma$  be an alphabet and  $W$  be a collection of finite words in  $\Sigma$ . Then  $W$  is *uniquely readable* if and only if whenever  $u, v, w \in W$  and  $uv = pws$  with  $p$  and  $s$  strings of symbols in  $\Sigma$ , then either  $p$  or  $s$  is the empty word.

Moreover, our  $(n+1)$ -words will be strongly uniform in the  $n$ -words as defined below.

**Definition 3.** We call a construction sequence  $(W_n)_{n \in \mathbb{N}}$  *uniform* if there is a summable sequence  $(\varepsilon_n)_{n \in \mathbb{N}}$  of positive numbers and for each  $n \in \mathbb{N}$  a map  $d_n : W_n \rightarrow (0, 1)$  such that for all words  $w' \in W_{n+1}$  and  $w \in W_n$  we have

$$\left| \frac{r(w, w')}{h_{n+1}/h_n} - d_n(w) \right| < \frac{\varepsilon_{n+1}}{h_n},$$

where  $r(w, w')$  is the number of occurrences of  $w$  in  $w'$ . Moreover, the construction sequence is called *strongly uniform* if for each  $n \in \mathbb{N}$  there is a constant  $c > 0$  such that for all words  $w' \in W_{n+1}$  and  $w \in W_n$  we have  $r(w, w') = c$ .

In the following we will identify  $\mathbb{K}$  with the symbolic system  $(\mathbb{K}, sh)$ . We introduce the following natural set  $S$  which will be of measure one for measures that we consider.

**Definition 4.** Suppose that  $(W_n)_{n \in \mathbb{N}}$  is a construction sequence for a symbolic system  $\mathbb{K}$  with each  $W_n$  uniquely readable. Let  $S$  be the collection of  $x \in \mathbb{K}$  such that there are sequences of natural numbers  $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}}$  going to infinity such that for all  $m \in \mathbb{N}$  there is  $n \in \mathbb{N}$  such that  $x \upharpoonright [-a_m, b_m] \in W_n$ .

We note that  $S$  is a dense shift-invariant  $\mathcal{G}_\delta$  subset of  $\mathbb{K}$  and we recall the following properties from [FW19a, Lemma 11] and [FW19b, Lemma 12].

**Fact 5.** Fix a construction sequence  $(W_n)_{n \in \mathbb{N}}$  for a symbolic system  $\mathbb{K}$  in a finite alphabet  $\Sigma$ . Then:

- (1)  $\mathbb{K}$  is the smallest shift-invariant closed subset of  $\Sigma^{\mathbb{Z}}$  such that for all  $n \in \mathbb{N}$  and  $w \in W_n$ ,  $\mathbb{K}$  has non-empty intersection with the basic open interval  $\langle w \rangle \subset \Sigma^{\mathbb{Z}}$ .
- (2) Suppose that  $(W_n)_{n \in \mathbb{N}}$  is a uniform construction sequence. Then there is a unique non-atomic shift-invariant measure  $\nu$  on  $\mathbb{K}$  concentrating on  $S$  and  $\nu$  is ergodic.
- (3) If  $\nu$  is a shift-invariant measure on  $\mathbb{K}$  concentrating on  $S$ , then for  $\nu$ -almost every  $s \in S$  there is  $N \in \mathbb{N}$  such that for all  $n > N$  there are  $a_n \leq 0 < b_n$  such that  $s \upharpoonright [a_n, b_n] \in W_n$ .

Since our symbolic systems  $\mathbb{K} \cong (\Sigma^{\mathbb{Z}}, \mathcal{B}, \nu, sh)$  will be built from a uniquely readable uniform construction sequence, they will automatically be ergodic. To each symbolic system we will also consider its inverse  $\mathbb{K}^{-1}$  which stands for  $(\mathbb{K}, sh^{-1})$ . Since it will often be convenient to have the shifts going in the same direction, we also introduce another convention.

**Definition 6.** If  $w$  is a finite or infinite string, we write  $rev(w)$  for the reverse string of  $w$ . In particular, if  $x$  is in  $\mathbb{K}$  we define  $rev(x)$  by setting  $rev(x)(k) = x(-k)$ . Then for  $A \subseteq \mathbb{K}$  we define  $rev(A) = \{rev(x) : x \in A\}$ .

If we explicitly view a finite word  $w$  positioned at a location interval  $[a, b]$ , then we take  $rev(w)$  to be positioned at the same interval  $[a, b]$  and we set  $rev(w)(k) = w(a + b - (k + 1))$ . For a collection  $W$  of words  $rev(W)$  is the collection of reverses of words in  $W$ .

Then we introduce the symbolic system  $(rev(\mathbb{K}), sh)$  as the one built from the construction sequence  $(rev(W_n))_{n \in \mathbb{N}}$ . Clearly, the map sending  $x$  to  $rev(x)$  is a canonical isomorphism between  $(\mathbb{K}, sh^{-1})$  and  $(rev(\mathbb{K}), sh)$ . We often abbreviate the symbolic system  $(rev(\mathbb{K}), sh)$  as  $rev(\mathbb{K})$ . If we have a cutting-and-stacking model of  $(rev(\mathbb{K}), sh)$  where each level (except the top level) of each column is mapped to the level above it, then the model for  $(\mathbb{K}, sh^{-1})$  would map each level (except the bottom level) to the level below it. In analogy with  $(rev(\mathbb{K}), sh)$ , we could instead reverse the order of the levels in each column and map the levels upward.

**2.3. Odometer-Based Systems.** The systems constructed in the first part of the paper to prove our anti-classification result for ergodic MPT's will turn out to be so-called odometer-based systems. In this subsection we review the definition.

Let  $(k_n)_{n \in \mathbb{N}}$  be a sequence of natural numbers  $k_n \geq 2$  and

$$O = \prod_{n \in \mathbb{N}} (\mathbb{Z}/k_n \mathbb{Z})$$

be the  $(k_n)_{n \in \mathbb{N}}$ -adic integers. Then  $O$  has a compact abelian group structure and hence carries a Haar measure  $\mu$ . We define a transformation  $T : O \rightarrow O$  to be addition by 1 in the  $(k_n)_{n \in \mathbb{N}}$ -adic integers (that is, the map that adds one in  $\mathbb{Z}/k_0 \mathbb{Z}$  and carries right). Then  $T$  is a  $\mu$ -preserving invertible transformation, called an *odometer transformation*, which is ergodic and has discrete spectrum.

We now define the collection of symbolic systems that have odometer systems as their timing mechanism to parse typical elements of the system.

**Definition 7.** Let  $(W_n)_{n \in \mathbb{N}}$  be a uniquely readable construction sequence with  $W_0 = \Sigma$  and  $W_{n+1} \subseteq (W_n)^{k_n}$  for every  $n \in \mathbb{N}$ . The associated symbolic shift will be called an *odometer-based system*.

Thus, odometer-based systems are those built from construction sequences  $(W_n)_{n \in \mathbb{N}}$  such that the words in  $W_{n+1}$  are concatenations of a fixed number  $k_n$  of words in  $W_n$ . Hence, the words in  $W_n$  have length

$$h_n = \prod_{i=0}^{n-1} k_i,$$

if  $n > 0$ , and  $h_0 = 1$ . Moreover, the spacers in part 3 of Definition 1 are all the empty words (that is, an odometer-based transformation can be built by a cutting-and-stacking construction using no spacers).

*Remark.* By [FW4], any finite entropy system that has an odometer factor can be represented as an odometer-based system

**2.4. The  $\bar{f}$  metric.** Feldman [Fe76] introduced a notion of distance, now called  $\bar{f}$ , between strings of symbols.

**Definition 8.** A *match* between two strings of symbols  $a_1 a_2 \dots a_n$  and  $b_1 b_2 \dots b_m$  from a given alphabet  $\Sigma$ , is a collection  $\mathcal{M}$  of pairs of indices  $(i_s, j_s)$ ,  $s = 1, \dots, r$  such that  $1 \leq i_1 < i_2 < \dots < i_r \leq n$ ,  $1 \leq j_1 < j_2 < \dots < j_r \leq m$  and  $a_{i_s} = b_{j_s}$  for  $s = 1, 2, \dots, r$ . A *pairing* is defined the same way as a match, except we drop the requirement that  $a_{i_s} = b_{j_s}$ . Then

$$(2.1) \quad \bar{f}(a_1 a_2 \dots a_n, b_1 b_2 \dots b_m) = \frac{1}{1 - \frac{2 \sup\{|\mathcal{M}| : \mathcal{M} \text{ is a match between } a_1 a_2 \dots a_n \text{ and } b_1 b_2 \dots b_m\}}{n + m}}.$$

We will refer to  $\bar{f}(a_1 a_2 \dots a_n, b_1 b_2 \dots b_m)$  as the “ $\bar{f}$ -distance” between  $a_1 a_2 \dots a_n$  and  $b_1 b_2 \dots b_m$ , even though  $\bar{f}$  does not satisfy the triangle inequality unless the strings are all of the same length. A match  $\mathcal{M}$  is called a *best possible match* if it realizes the supremum in the definition of  $\bar{f}$ . The extension of the definition of  $\bar{f}$ -distance from finite words to infinite words is analogous to the extension for  $\bar{d}$ -distance.



If  $x = x_1x_2 \cdots x_n$  and  $y = y_1y_2 \cdots y_n$  are decompositions of the strings of symbols  $x$  and  $y$  into substrings such that in a given match (or pairing) between  $x$  and  $y$ , all matches (or pairings) are between symbols in  $x_i$  and  $y_i$ , for  $i = 1, 2, \dots, n$ , then we say that  $x_i$  *corresponds* to  $y_i$  (and vice versa), under this match (or pairing). If we are given a match or pairing between  $x = x_1x_2 \cdots x_n$  and  $y$ , then we can decompose  $y$  into  $y_1y_2 \cdots y_n$  so that  $x_i$  corresponds to  $y_i$ , for  $i = 1, 2, \dots, n$ . The choice of  $y_1, y_2, \dots, y_n$  is not unique, but this does not matter in any of our arguments.

*Remark.* In the proof of Lemma 13, we will use certain pairings to give an upper bound on  $|\mathcal{M}|$ , for a best possible match  $\mathcal{M}$ .

*Remark.* Alternatively, one can view a match as an injective order-preserving function  $\pi : \mathcal{D}(\pi) \subseteq \{1, \dots, n\} \rightarrow \mathcal{R}(\pi) \subseteq \{1, \dots, m\}$  with  $a_i = b_{\pi(i)}$  for every  $i \in \mathcal{D}(\pi)$ . Then  $\bar{f}(a_1 \dots a_n, b_1 \dots b_m) = 1 - \max \left\{ \frac{2|\mathcal{D}(\pi)|}{n+m} : \pi \text{ is a match} \right\}$ .

The following simple properties of  $\bar{f}$  from [GKpp], that were already used in [Fe76] and [ORW82], will appear frequently in our arguments. These properties can be proved easily by considering the *fit*  $1 - \bar{f}(a, b)$  between two strings  $a$  and  $b$ .

**Fact 9.** *Suppose  $a$  and  $b$  are strings of symbols of length  $n$  and  $m$ , respectively, from an alphabet  $\Sigma$ . If  $\tilde{a}$  and  $\tilde{b}$  are strings of symbols obtained by deleting at most  $\lfloor \gamma(n+m) \rfloor$  terms from  $a$  and  $b$  altogether, where  $0 < \gamma < 1$ , then*

$$(2.2) \quad \bar{f}(a, b) \geq \bar{f}(\tilde{a}, \tilde{b}) - 2\gamma.$$

*Moreover, if there exists a best possible match between  $a$  and  $b$  such that no term that is deleted from  $a$  and  $b$  to form  $\tilde{a}$  and  $\tilde{b}$  is matched with a non-deleted term, then*

$$(2.3) \quad \bar{f}(a, b) \geq \bar{f}(\tilde{a}, \tilde{b}) - \gamma.$$

*Likewise, if  $\tilde{a}$  and  $\tilde{b}$  are obtained by adding at most  $\lfloor \gamma(n+m) \rfloor$  symbols to  $a$  and  $b$ , then (2.3) holds.*

**Fact 10.** *If  $x = x_1x_2 \cdots x_n$  and  $y = y_1y_2 \cdots y_n$  are decompositions of the strings of symbols  $x$  and  $y$  into corresponding substrings under a best possible  $\bar{f}$ -match between  $x$  and  $y$ , that is, one that achieves the minimum in the definition of  $\bar{f}$ , then*

$$\bar{f}(x, y) = \sum_{i=1}^n \bar{f}(x_i, y_i)v_i,$$

where

$$(2.4) \quad v_i = \frac{|x_i| + |y_i|}{|x| + |y|}.$$

**Fact 11.** *If  $x$  and  $y$  are strings of symbols such that  $\bar{f}(x, y) \leq \gamma$ , for some  $0 \leq \gamma < 1$ , then*

$$(2.5) \quad \left( \frac{1-\gamma}{1+\gamma} \right) |x| \leq |y| \leq \left( \frac{1+\gamma}{1-\gamma} \right) |x|.$$

We also recall the following elementary fact from [ORW82, Proposition 2.3, p.85].

**Fact 12.** *Suppose  $a$  and  $b$  are strings of symbols of length  $n$  and  $m$ , respectively, from an alphabet  $\Sigma$ . Let  $h$  be a symbol that is not an element in  $\Sigma$ . Suppose  $\tilde{a}$  and*

$\tilde{b}$  are strings of symbols obtained by inserting at most  $(\beta - 1)n$  symbols  $h$  in  $a$  and at most  $(\beta - 1)m$  symbols  $h$  in  $b$ . Then

$$\bar{f}(\tilde{a}, \tilde{b}) \geq \frac{\bar{f}(a, b)}{\beta}.$$

The next result provides lower bounds on the  $\bar{f}$  distance when symbols  $a_i$  and  $b_j$  are replaced by blocks  $A_{a_i}$  and  $A_{b_j}$ .

**Lemma 13** (Symbol by block replacement). *Suppose that  $A_{a_1}, A_{a_2}, \dots, A_{a_n}$  and  $A_{b_1}, A_{b_2}, \dots, A_{b_m}$  are blocks of symbols with each block of length  $L$ . Assume that  $\alpha \in (0, 1/7)$ ,  $R \geq 2$ , and for all substrings  $C$  and  $D$  consisting of consecutive symbols from  $A_{a_i}$  and  $A_{b_j}$ , respectively, with  $|C|, |D| \geq L/R$ , we have*

$$\bar{f}(C, D) \geq \alpha, \text{ if } a_i \neq b_j.$$

Then

$$\bar{f}(A_{a_1}A_{a_2} \cdots A_{a_n}, A_{b_1}A_{b_2} \cdots A_{b_m}) > \alpha \bar{f}(a_1a_2 \cdots a_n, b_1b_2 \cdots b_m) - \frac{1}{R}.$$

*Proof.* We may assume  $n \leq m$ . Suppose  $\mathcal{M}$  is a best possible match between  $A_{a_1}A_{a_2} \cdots A_{a_n}$  and  $A_{b_1}A_{b_2} \cdots A_{b_m}$ . Recall that  $|\mathcal{M}|$  denotes the number of pairs of symbols in  $A_{a_1}A_{a_2} \cdots A_{a_n}$  and  $A_{b_1}A_{b_2} \cdots A_{b_m}$  that are matched by  $\mathcal{M}$ . We use the analogous notation  $|\mathcal{N}|$  for a match  $\mathcal{N}$  between the strings  $a_1a_2 \cdots a_n$  and  $b_1b_2 \cdots b_m$ .

*Claim.* There exists a match  $\mathcal{N}$  (not necessarily a best possible match) between the strings  $a_1a_2 \cdots a_n$  and  $b_1b_2 \cdots b_m$  such that

$$(2.6) \quad L(n+m) - 2|\mathcal{M}| \geq \alpha \left\{ L(n+m) - 2L|\mathcal{N}| - \frac{3L(n+m)}{R} \right\}.$$

The lemma will follow from this claim, because

$$\bar{f}(A_{a_1}A_{a_2} \cdots A_{a_n}, A_{b_1}A_{b_2} \cdots A_{b_m}) = 1 - \frac{2|\mathcal{M}|}{L(n+m)} \text{ and}$$

$$\begin{aligned} \frac{\alpha}{L(n+m)} \left\{ L(n+m) - 2L|\mathcal{N}| - \frac{3L(n+m)}{R} \right\} &\geq \alpha \bar{f}(a_1a_2 \cdots a_n, b_1b_2 \cdots b_m) - \frac{3\alpha}{R} \\ &> \alpha \bar{f}(a_1a_2 \cdots a_n, b_1b_2 \cdots b_m) - \frac{1}{R}. \end{aligned}$$

*Proof of Claim.* We will modify  $\mathcal{M}$  in a series of  $i$  steps, for  $i = 1, 2, \dots, n$ , obtaining pairings  $\mathcal{M}_i$  between  $A_{a_1}A_{a_2} \cdots A_{a_n}$  and  $A_{b_1}A_{b_2} \cdots A_{b_m}$  such that  $|\mathcal{M}| \leq |\mathcal{M}_1| \leq \dots \leq |\mathcal{M}_{n-1}| \leq |\mathcal{M}_n|$ . The positions of the modifications will move from left to right with each step. We will simultaneously construct a sequence of matches  $\mathcal{N}_i$  between  $a_1a_2 \cdots a_n$  and  $b_1b_2 \cdots b_m$  such that  $\mathcal{N}_1 \subseteq \mathcal{N}_2 \subseteq \dots \subseteq \mathcal{N}_n$ . At the end of this construction, we will show that the inequality (2.6) holds with  $\mathcal{N} = \mathcal{N}_n$  and  $\mathcal{M}$  replaced by  $\mathcal{M}_n$ . This implies (2.6) as written.

We let  $\mathcal{M}_0 = \mathcal{M}$  and  $\mathcal{N}_0 = \emptyset$ . In the inductive construction of  $\mathcal{M}_i$  and  $\mathcal{N}_i$  (as described below), at most one pair of matching symbols is added to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ , and each time such an addition occurs, we allow up to  $L$  pairs to be in the part of  $\mathcal{M}_i$  that is newly specified at stage  $i$ .

Suppose  $1 \leq i \leq n$ , and  $\mathcal{M}_{i-1}$  and  $\mathcal{N}_{i-1}$  have been constructed. Assume that  $A_{b_1}A_{b_2} \cdots A_{b_m} = B_1B_2 \cdots B_n$ , where  $A_{a_j}$  corresponds to  $B_j$  under  $\mathcal{M}_{i-1}$ . If  $B_i$  starts with a symbol in  $A_{b_{j_i}}$ , we also assume that none of symbols in the matching pairs in  $\mathcal{N}_{i-1}$  come from  $a_i a_{i+1} \cdots a_n$  or  $b_{j_i} b_{j_i+1} \cdots b_m$ . (Therefore these symbols

are still available to add to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ .) We consider the following four cases (with various subcases).

**Case 1.**  $B_i$  contains two or more complete  $A_{b_j}$ 's. Then  $|B_i| \geq 2|A_{a_i}|$  and therefore  $\bar{f}(A_{a_i}, B_i) > 1/7 > \alpha$  by Fact 11. In this case we take  $\mathcal{M}_i = \mathcal{M}_{i-1}$  and  $\mathcal{N}_i = \mathcal{N}_{i-1}$ .

**Case 2.**  $B_i$  consists of one complete  $A_{b_j}$  plus possibly a final string  $A_{b_{j-1}}^{\text{fin}}$  in  $A_{b_{j-1}}$  and possibly an initial string  $A_{b_{j+1}}^{\text{int}}$  in  $A_{b_{j+1}}$ . By the induction hypothesis,  $b_j, b_{j+1}$ , and  $a_i$  are not in any of the pairs in  $\mathcal{N}_{i-1}$ , and neither is  $b_{j-1}$  if  $A_{b_{j-1}}^{\text{fin}} \neq \emptyset$ .

**Case 2(i).** If  $a_i = b_{j-1}$  or  $a_i = b_j$ , then we add one pair of equal symbols from these equations to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ . To construct  $\mathcal{M}_i$  we modify  $\mathcal{M}_{i-1}$  on  $A_{a_i}$  and  $B_i$  to create a perfect pairing between  $A_{a_i}$  and  $A_{b_j}$ . (That is, each symbol in  $A_{a_i}$  is paired with a symbol in  $A_{b_j}$ , even though they might not actually match.) The symbols in  $B_i \setminus A_{b_j}$  are not paired to anything by  $\mathcal{M}_i$  and they are eliminated from consideration for any future pairings  $\mathcal{M}_\ell$  for  $\ell > i$ . Outside of  $A_{a_i}$  and  $B_i$ ,  $\mathcal{M}_i$  agrees with  $\mathcal{M}_{i-1}$ . Note that  $|\mathcal{M}_i| \geq |\mathcal{M}_{i-1}|$  because  $\mathcal{M}_i$  has the maximum possible number of pairings between  $A_{a_i}$  and  $B_i$ .

**Case 2(ii).** If  $a_i \neq b_{j-1}$ ,  $a_i \neq b_j$ , and  $a_i = b_{j+1}$ , then we add  $a_i$  and  $b_{j+1}$  to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ . In this subcase and the next one, we let  $A_{a_i}^{(0)}, A_{a_i}^{(1)}$ , and  $A_{a_i}^{(2)}$  be the subdivision of  $A_{a_i}$  into the segments corresponding to  $A_{b_{j-1}}^{\text{fin}}, A_{b_j}$ , and  $A_{b_{j+1}}^{\text{int}}$ , respectively. We construct  $\mathcal{M}_i$  from  $\mathcal{M}_{i-1}$  by creating a perfect pairing between the extension of  $A_{a_i}^{(2)}$  to a total length of  $L$  symbols and  $A_{b_{j+1}}$ . (In case  $i = n$ , we only pair  $A_{a_n}^{(2)}$ , with no extension). Any pairing in  $\mathcal{M}_{i-1}$  involving a symbol in this extension of  $A_{a_i}^{(2)}$  or in  $A_{b_{j+1}}$  that is different from the newly created perfect pairing is not included in  $\mathcal{M}_i$ . Otherwise  $\mathcal{M}_i$  is the same as  $\mathcal{M}_{i-1}$ . On the part that was changed the pairing is as good as it can be for that number of symbols, and therefore  $|\mathcal{M}_i| \geq |\mathcal{M}_{i-1}|$ . On the strings  $A_{a_i}^{(0)}$  and  $A_{b_{j-1}}^{\text{fin}}$  the subscripts of the blocks do not match, and therefore either  $\bar{f}(A_{a_i}^{(0)}, A_{b_{j-1}}^{\text{fin}}) \geq \alpha$  by hypothesis or at least one of these two strings has length less than  $L/R$ . In the latter case, if the other string has length at least  $4L/(3R)$ , then by Fact 11,  $\bar{f}(A_{a_i}^{(0)}, A_{b_{j-1}}^{\text{fin}}) > 1/7 > \alpha$ . If one string has length less than  $L/R$  and the other string has length less than  $4L/(3R)$  we refer to this as the ‘‘short substring case’’ and the number of symbols coming from such cases will be at most  $14Ln/(3R) \leq 7L(n+m)/(3R) < 3L(n+m)/R$ . (As we will see in the next subcase there could be up to two pairs of short substrings for each  $i \in \{1, 2, \dots, n\}$ .) The strings  $A_{a_i}^{(1)}$  and  $A_{b_j}$  are also in blocks whose subscripts do not match. Since  $A_{b_j}$  is an entire block, the short substring case does not apply, and we obtain  $\bar{f}(A_{a_i}^{(1)}, A_{b_j}) > \alpha$ . For the next step,  $B_{i+1}$  starts with  $A_{b_{j+2}}$  and we modify  $A_{a_{i+1}}$  to be only the last  $|A_{a_i}^{(2)}|$  symbols in the original  $A_{a_{i+1}}$ .

**Case 2(iii).** If  $a_i \neq b_{j-1}$ ,  $a_i \neq b_j$ , and  $a_i \neq b_{j+1}$ , then we keep  $\mathcal{M}_i = \mathcal{M}_{i-1}$  and  $\mathcal{N}_i = \mathcal{N}_{i-1}$ . For each of the pairs  $A_{a_i}^{(0)}$  and  $A_{b_{j-1}}^{\text{fin}}, A_{a_i}^{(1)}$  and  $A_{b_j}, A_{a_i}^{(2)}$  and  $A_{b_{j+1}}^{\text{int}}$ , either we know that the  $\bar{f}$  distance is greater than  $\alpha$  or (possibly for the first and the third pair) we are in the short substring case.

**Case 3.**  $B_i$  contains a final segment of  $A_{b_{j-1}}$  and an initial segment of  $A_{b_j}$ .

**Case 3(i).** If  $a_i = b_{j-1}$ , then we add  $a_i$  and  $b_{j-1}$  to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ . We replace  $\mathcal{M}_{i-1}$  on  $A_{a_i}$  and  $B_i$  by a perfect pairing between  $A_{a_i}$  and  $L$  consecutive symbols in the  $A_{b_{j-1}}A_{b_j}$  string starting with the first symbol in  $B_i$ . If  $B_i$  contains fewer than  $L$  symbols, then we continue the pairing beyond the end of  $B_i$ . (This

may eliminate some previous pairings that were in  $\mathcal{M}_{i-1}$  but the net result is still  $|\mathcal{M}_i| \geq |\mathcal{M}_{i-1}|$ .) If  $B_i$  contains more than  $L$  symbols, then the remaining  $|B_i| - L$  symbols stay unpaired in  $\mathcal{M}_\ell$  for  $\ell \geq i$ .

**Case 3(ii).** If  $a_i \neq b_{j-1}$  and  $a_i = b_j$ , then we add  $a_i$  and  $b_j$  to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ . We keep  $\mathcal{M}_i$  the same as  $\mathcal{M}_{i-1}$  up to the symbol prior to the first one in  $A_{a_i}$  that is paired with a symbol in  $A_{b_j}$ . Then we pair the next  $L$  consecutive symbols in  $A_{a_i}A_{a_{i+1}}$  to  $A_{b_j}$ . Otherwise  $\mathcal{M}_i$  stays the same as  $\mathcal{M}_{i-1}$ .

**Case 3(iii).** If  $a_i \neq b_{j-1}$  and  $a_i \neq b_j$ , then we again have two pairs of strings, one for the part of  $A_i$  that corresponds to a final segment of  $A_{b_{j-1}}$  and the other for the part of  $A_i$  that corresponds to the initial segment of  $A_{b_j}$ . Each of these pairs is  $\bar{f}$  distance at least  $\alpha$  apart or consists of short substrings. We let  $\mathcal{M}_i = \mathcal{M}_{i-1}$  and  $\mathcal{N}_i = \mathcal{N}_{i-1}$ .

**Case 4.**  $B_i$  is a substring of a single  $A_{b_j}$ .

**Case 4(i).** If  $a_i = b_j$ , then we add  $a_i$  and  $b_j$  to  $\mathcal{N}_{i-1}$  to form  $\mathcal{N}_i$ . If  $j \neq m$ , we construct  $\mathcal{M}_i$  by extending  $B_i$  to the right for a total of  $L$  symbols and pairing the extended  $B_i$  exactly with  $A_{a_i}$ . If  $j = m$ , we pair all of  $B_i$  with a substring in  $A_{a_i}$ . Whatever pairings in  $\mathcal{M}_{i-1}$  are not affected by these changes are included in  $\mathcal{M}_i$ .

**Case 4(ii).** If  $a_i \neq b_j$ , then  $\bar{f}(A_{a_i}, B_j) \geq \alpha$ , and we let  $\mathcal{M}_i = \mathcal{M}_{i-1}$  and  $\mathcal{N}_i = \mathcal{N}_{i-1}$ .

In summary, after we apply this procedure for  $i = 1, 2, \dots, n$ , and we remove from  $A_{a_1}A_{a_2} \cdots A_{a_n}$  and  $A_{b_1}A_{b_2} \cdots A_{b_m}$  the symbols involved in the (at most  $L|\mathcal{N}_n|$ ) pairings obtained from Cases (2i),(2ii),(3i),(3ii), and (4i) in the formation of  $\mathcal{M}_i$  from  $\mathcal{M}_{i-1}$ , and we remove the pairs of corresponding strings in the short substring case, we are left with a total of at least  $L(n+m) - 2L|\mathcal{N}_n| - 3L((n+m)/R)$  symbols in pairs of strings in  $A_{a_1}A_{a_2} \cdots A_{a_n}$  and  $A_{b_1}A_{b_2} \cdots A_{b_m}$  that correspond under  $\mathcal{M}_n$ . On these corresponding strings the  $\mathcal{M}_n$  pairing is a subset of the original  $\mathcal{M}$ , and the symbols in each pair of these corresponding strings are contained in some  $A_{a_i}$  and  $A_{b_j}$  where  $a_i \neq b_j$ . By hypothesis, the  $\bar{f}$ -distance between two such corresponding strings is at least  $\alpha$ . Therefore

$$\begin{aligned} L(n+m) - 2|\mathcal{M}_0| &\geq L(n+m) - 2|\mathcal{M}_n| \\ &\geq \alpha \left\{ L(n+m) - 2L|\mathcal{N}_n| - \frac{3L(n+m)}{R} \right\}, \end{aligned}$$

where the second inequality follows from Fact 10.  $\square$

**2.5. Some background from descriptive set theory.** To state our results precisely we need some concepts explained below. The main tool is the idea of a reduction. See [Ke95] and [Fo18] for further information.

**Definition 14.** Let  $X$  and  $Y$  be Polish spaces and  $A \subseteq X$ ,  $B \subseteq Y$ . A function  $f : X \rightarrow Y$  reduces  $A$  to  $B$  if and only if for all  $x \in X$ :  $x \in A$  if and only if  $f(x) \in B$ .

Such a function  $f$  is called a Borel (respectively, continuous) reduction if  $f$  is a Borel (respectively, continuous) function.

A being reducible to  $B$  can be interpreted as saying that  $B$  is at least as complicated as  $A$ . We note that if  $B$  is Borel and  $f$  is a Borel reduction, then  $A$  is also Borel. Equivalently, if  $f$  is a Borel reduction of  $A$  to  $B$  and  $A$  is not Borel, then  $B$  is not Borel.

**Definition 15.** If  $\mathcal{S}$  is a collection of sets and  $B \in \mathcal{S}$ , then  $B$  is called  $\mathcal{S}$ -complete for Borel (respectively, continuous) reductions if and only if every  $A \in \mathcal{S}$  is Borel (respectively, continuously) reducible to  $B$ .

Following the interpretation above  $B$  is said to be at least as complicated as each set in  $\mathcal{S}$ .

**Definition 16.** If  $X$  is a Polish space and  $B \subseteq X$ , then  $B$  is *analytic* if and only if it is the continuous image of a Borel subset of a Polish space. Equivalently, there is a Polish space  $Y$  and a Borel set  $C \subseteq X \times Y$  such that  $B$  is the  $X$ -projection of  $C$ .

There are analytic sets that are not Borel (see, for example, [Ke95, section 14]). This implies that a complete analytic set is not Borel. The collection of ill-founded trees, which we describe below, is an example of a complete analytic set, and it is fundamental to our construction, as well as that in [FRW11].

To define the ill-founded trees, we first consider the set  $\mathbb{N}^{<\mathbb{N}}$  of finite sequences of natural numbers. For  $\tau \in \mathbb{N}^{<\mathbb{N}}$ , let  $lh(\tau)$  denote the length of  $\tau$ . A *tree* is a set  $\mathcal{T} \subseteq \mathbb{N}^{<\mathbb{N}}$  such that if  $\tau = (\tau_1, \dots, \tau_n) \in \mathcal{T}$  and  $\sigma = (\tau_1, \dots, \tau_s)$  with  $s \leq n$  is an initial segment of  $\tau$ , then  $\sigma \in \mathcal{T}$ . If  $\sigma$  is an initial segment of  $\tau$ , then  $\sigma$  is a *predecessor* of  $\tau$  and  $\tau$  is a *successor* of  $\sigma$ . We define the level  $s$  of a tree  $\mathcal{T}$  to be the collection of elements of  $\mathcal{T}$  that have length  $s$ .

A *branch* through  $\mathcal{T}$  is a function  $f$  into  $\mathcal{T}$  with domain either some  $\{0, \dots, n-1\}$  or  $\mathbb{N}$  itself such that  $lh(f(s)) = s$ , and if  $s+1$  is in the domain of  $f$ , then  $f(s+1)$  is an immediate successor of  $f(s)$ . If a tree has an infinite branch (that is, the  $f$  above can be taken with domain  $\mathbb{N}$ ), it is called *ill-founded*. If it does not have an infinite branch, it is called *well-founded*.

In the following, let  $\{\sigma_n : n \in \mathbb{N}\}$  be an enumeration of  $\mathbb{N}^{<\mathbb{N}}$  with the property that every proper predecessor of  $\sigma_n$  is some  $\sigma_m$  for  $m < n$ . Under this enumeration subsets  $S \subseteq \mathbb{N}^{<\mathbb{N}}$  can be identified with characteristic functions  $\chi_S : \mathbb{N} \rightarrow \{0, 1\}$ . The collection of such  $\chi_S$  can be viewed as the members of an infinite product space  $\{0, 1\}^{\mathbb{N}^{<\mathbb{N}}}$  homeomorphic to the Cantor space. Here, each function  $a : \{\sigma_m : m < n\} \rightarrow \{0, 1\}$  determines a basic open set

$$\langle a \rangle = \{\chi : \chi \upharpoonright \{\sigma_m : m < n\} = a\} \subseteq \{0, 1\}^{\mathbb{N}^{<\mathbb{N}}}$$

and the collection of all such  $\langle a \rangle$  forms a basis for the topology. The collection of trees is a closed (hence compact) subset of  $\{0, 1\}^{\mathbb{N}^{<\mathbb{N}}}$  in this topology. Moreover, the collection of trees containing arbitrarily long finite sequences is a dense  $\mathcal{G}_\delta$  subset. In particular, this collection is a Polish space. We will denote the space of trees containing arbitrarily long finite sequences by *Trees*.

Since the topology on the space of trees was introduced via basic open sets giving us a finite amount of information about the trees in it, we can characterize continuous maps defined on *Trees* as follows.

**Fact 17.** *Let  $Y$  be a topological space. Then a map  $f : \text{Trees} \rightarrow Y$  is continuous if and only if for all open sets  $O \subseteq Y$  and all  $\mathcal{T} \in \text{Trees}$  with  $f(\mathcal{T}) \in O$  there is  $M \in \mathbb{N}$  such that for all  $\mathcal{T}' \in \text{Trees}$  we have:*

$$\text{if } \mathcal{T} \cap \{\sigma_n : n \leq M\} = \mathcal{T}' \cap \{\sigma_n : n \leq M\}, \text{ then } f(\mathcal{T}') \in O.$$

As mentioned before, we have the following classical fact (see, for example, [Ke95, section 27]).

**Fact 18.** *The collection of ill-founded trees is a complete analytic subset of  $\mathcal{Trees}$ .*

As described in the next section, we will prove the main results of this paper by reducing the collection of ill-founded trees to pairs of Kakutani-equivalent ergodic transformations. Then Fact 18 will show that this set of Kakutani-equivalent pairs of transformations is also complete analytic.

During our construction the following maps will prove useful.

**Definition 19.** We define a map  $M : \mathcal{Trees} \rightarrow \mathbb{N}^{\mathbb{N}}$  by setting  $M(\mathcal{T})(s) = n$  if and only if  $n$  is the least number such that  $\sigma_n \in \mathcal{T}$  and  $lh(\sigma_n) = s$ . Dually, we also define a map  $s : \mathcal{Trees} \rightarrow \mathbb{N}^{\mathbb{N}}$  by setting  $s(\mathcal{T})(n)$  to be the length of the longest sequence  $\sigma_m \in \mathcal{T}$  with  $m \leq n$ .

*Remark.* When  $\mathcal{T}$  is clear from the context we write  $M(s)$  and  $s(n)$ . We also note that  $s(n) \leq n$  and that  $M$  as well as  $s$  is a continuous function when we endow  $\mathbb{N}$  with the discrete topology and  $\mathbb{N}^{\mathbb{N}}$  with the product topology.

### 3. MAIN RESULTS

**3.1. Statement of results.** We are ready to state the main result of this paper.

**Theorem 20.** *The collection*

$$\{(S, T) : S \text{ and } T \text{ are ergodic and Kakutani equivalent}\} \subseteq \mathcal{E} \times \mathcal{E}$$

*is a complete analytic set. In particular, it is not Borel.*

As we show below, this will follow from the subsequent theorem which is proved in Section 9.

**Theorem 21.** *There is a continuous one-to-one map*

$$\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$$

*such that for  $\mathcal{T} \in \mathcal{Trees}$  and  $T = \Psi(\mathcal{T})$ :*

*$\mathcal{T}$  has an infinite branch if and only if  $T$  and  $T^{-1}$  are Kakutani equivalent.*

Using the notions from descriptive set theory in Section 2.5, Theorem 21 can also be expressed as follows.

**Corollary 22.** *There is a continuous one-to-one map*

$$\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$$

*that reduces the collection of ill-founded trees to the collection of ergodic transformations  $T$  such that  $T$  and  $T^{-1}$  are Kakutani equivalent.*

*Proof of Theorem 20.* The map  $i(T) = (T, T^{-1})$  is a continuous mapping from  $\mathcal{E}$  to  $\mathcal{E} \times \mathcal{E}$  and reduces  $\{T \in \mathcal{E} : T \text{ is Kakutani equivalent to } T^{-1}\}$  to

$$\{(S, T) \in \mathcal{E} \times \mathcal{E} : S \text{ and } T \text{ are Kakutani equivalent}\}.$$

Then Theorem 20 follows from Fact 18 and Corollary 22. □

To prove Theorem 21 we actually show the following stronger result.

**Proposition 23.** *There is a continuous one-to-one map*

$$\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$$

*such that for  $\mathcal{T} \in \mathcal{Trees}$  and  $T = \Psi(\mathcal{T})$ :*

- (1) If  $\mathcal{T}$  has an infinite branch, then  $T$  and  $T^{-1}$  are isomorphic.
- (2) If  $T$  and  $T^{-1}$  are Kakutani equivalent, then  $\mathcal{T}$  has an infinite branch.

This stronger proposition also implies the anti-classification result for the isomorphism relation in [FRW11].

**Corollary.** *There is a continuous one-to-one map*

$$\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$$

*that reduces the collection of ill-founded trees to the collection of ergodic automorphisms  $T$  such that  $T$  is isomorphic to  $T^{-1}$ . Hence, the collection*

$$\{(S, T) : S \text{ and } T \text{ are ergodic and isomorphic}\} \subseteq \mathcal{E} \times \mathcal{E}$$

*is a complete analytic set.*

In Section 10 we are able to upgrade these results to the setting of diffeomorphisms.

**Theorem 24.** *Let  $M$  be the disk, annulus or torus with Lebesgue measure  $\lambda$ . Then the collection*

$$\{(S, T) : S \text{ and } T \text{ are ergodic diffeomorphisms and Kakutani equivalent}\}$$

*in  $\text{Diff}^\infty(M, \lambda) \times \text{Diff}^\infty(M, \lambda)$  is a complete analytic set and hence not a Borel set with respect to the  $C^\infty$  topology.*

In case of  $M = \mathbb{T}^2$  we also obtain the results in the real-analytic category (see Subsection 10.1 for the definition of the space  $\text{Diff}_\rho^\omega(M, \lambda)$ ).

**Theorem 25.** *Let  $\rho > 0$  and  $\lambda$  be the Lebesgue measure on  $\mathbb{T}^2$ . Then the collection*

$$\{(S, T) : S \text{ and } T \text{ are ergodic diffeomorphisms and Kakutani equivalent}\}$$

*in  $\text{Diff}_\rho^\omega(M, \lambda) \times \text{Diff}_\rho^\omega(M, \lambda)$  is a complete analytic set and hence not a Borel set with respect to the  $\text{Diff}_\rho^\omega$  topology.*

**3.2. Outline of the proof of Proposition 23.** To construct the reduction  $\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$  we build for every  $\mathcal{T} \in \mathcal{Trees}$  a transformation  $\Psi(\mathcal{T})$  as a symbolic system with a strongly uniform and uniquely readable construction sequence  $(\mathcal{W}_n(\mathcal{T}))_{n \in \mathbb{N}}$  which implies its ergodicity. This will be done in such a way that  $\mathcal{W}_n(\mathcal{T})$  is entirely determined by  $\mathcal{T} \cap \{\sigma_m : m \leq n\}$ , which, by Fact 17 guarantees continuity of  $\Psi$  (see Proposition 20 in [FRW11] for details). (Alternatively, in the cutting-and-stacking model, the  $n$ th tower is determined by  $\mathcal{T} \cap \{\sigma_m : m \leq n\}$ , and this implies that  $\Psi$  is continuous.) Related to the structure of the tree we also specify equivalence relations  $\mathcal{Q}_s^n(\mathcal{T})$  on the collections  $\mathcal{W}_n(\mathcal{T})$  of  $n$ -words and group actions on the equivalence classes in  $\mathcal{W}_n(\mathcal{T})/\mathcal{Q}_s^n(\mathcal{T})$  as in [FRW11]. These specifications will allow us to prove in Section 5 that  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  are isomorphic if the tree  $\mathcal{T}$  has an infinite branch.

The harder part is to show that  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  are not Kakutani equivalent if the tree  $\mathcal{T}$  does not have an infinite branch. For this purpose, we build the  $n$ -words in our construction sequence using specific patterns of blocks (that we call Feldman patterns) which originate from Feldman's first example of an ergodic zero-entropy automorphism that is not loosely Bernoulli [Fe76]. As observed in [ORW82], different Feldman patterns cannot be matched well in  $\bar{f}$  even after a finite coding. We introduce these patterns in detail in Section 6. In our construction of  $n$ -words

we apply them repeatedly by substituting Feldman patterns of finer equivalence classes into Feldman patterns of coarser classes. We present such a substitution step in general in Section 7, and then use it in an iterative way to construct the collection  $\mathcal{W}_n(\mathcal{T})$  of  $n$ -words in Section 8. There we also verify that  $\mathcal{W}_n(\mathcal{T})$  satisfies the aforementioned specifications. Finally, using techniques from [ORW82] based on the structure of Feldman patterns, we show in Section 9 that  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  cannot be Kakutani equivalent if the tree  $\mathcal{T}$  does not have an infinite branch. In the analysis of possible finite codes we face the additional challenge that we do not know the precise words being coded, only their equivalence classes.

We transfer the results to the setting of diffeomorphisms in Section 10 and give an outline of this proof at the beginning of that section.

#### 4. THE SETUP

We present a general setup of our construction. It is very similar to the one in [FRW11, Section 4] with a few minor changes (for example, specification (Q4) and our specifications on the group actions).

**4.1. General properties of our construction.** For each  $n \in \mathbb{N}$  with  $\sigma_n \in \mathcal{T}$  we construct a set of words  $\mathcal{W}_n = \mathcal{W}_n(\mathcal{T})$  in the basic alphabet  $\Sigma = \{1, \dots, 2^{12}\}$  which depend only on  $\mathcal{T} \cap \{\sigma_m : m \leq n\}$ . To start we set  $\mathcal{W}_0 = \Sigma$ .

- (E1) All words in  $\mathcal{W}_n$  have the same length  $h_n$  and the cardinality  $|\mathcal{W}_n|$  is a power of 2.
- (E2) If  $\sigma_m$  and  $\sigma_n$  are consecutive elements of  $\mathcal{T}$ , then every word in  $\mathcal{W}_n$  is built by concatenating words in  $\mathcal{W}_m$ . There is  $f_m \in \mathbb{Z}^+$  such that every word in  $\mathcal{W}_m$  occurs in each word of  $\mathcal{W}_n$  exactly  $f_m$  times. The number  $f_m$  will be a product of  $p_n^2$  and powers of 2, where  $p_n$  is a large prime number chosen when  $\sigma_n$  is considered.
- (E3) If  $\sigma_m$  and  $\sigma_n$  are consecutive elements of  $\mathcal{T}$ ,  $w = w_1 \dots w_{h_n/h_m} \in \mathcal{W}_n$  and  $w' = w'_1 \dots w'_{h_n/h_m} \in \mathcal{W}_n$ , where  $w_i, w'_i \in \mathcal{W}_m$ , then for any  $k \geq \lfloor \frac{h_n/h_m}{2} \rfloor$  and  $1 \leq i \leq \frac{h_n}{h_m} - k$ , we have  $w_{i+1} \dots w_{i+k} \neq w'_{i+1} \dots w'_{i+k}$ .

*Remark 26.* In particular, these specifications say that  $\{\mathcal{W}_n : \sigma_n \in \mathcal{T}\}$  is a uniquely readable and strongly uniform construction sequence for an odometer-based system. Hence, the corresponding symbolic system  $\mathbb{K}$  has a unique non-atomic ergodic shift-invariant measure  $\nu$  by Fact 5.

**4.2. Equivalence relations and canonical factors.** Related to the structure of the tree we will specify, for each  $s \leq s(n)$ , an equivalence relation  $\mathcal{Q}_s^n$  on our words  $\mathcal{W}_n$ . We write  $Q_s^n$  for the number of equivalence classes in  $\mathcal{Q}_s^n$  and enumerate the classes by  $\{c_j^{(n,s)} : j = 1, \dots, Q_s^n\}$ . In the construction of the factor  $\mathbb{K}_s$  described in 4.2.2, we will identify  $\mathcal{W}_n/\mathcal{Q}_s^n$  with an alphabet denoted by  $(\mathcal{W}_n/\mathcal{Q}_s^n)^*$  of  $Q_s^n$  symbols  $\{1, \dots, Q_s^n\}$ .

Each equivalence relation  $\mathcal{Q}_s^n$  will induce an equivalence relation on  $rev(\mathcal{W}_n)$ , which we will also call  $\mathcal{Q}_s^n$ , as follows:  $rev(w), rev(w') \in rev(\mathcal{W}_n)$  are equivalent with respect to  $\mathcal{Q}_s^n$  if and only if  $w, w' \in \mathcal{W}_n$  are equivalent with respect to  $\mathcal{Q}_s^n$ .

To state the specifications on our equivalence relations we recall some general notions.

**Definition 27.** Let  $X$  be a set and let  $\mathcal{Q}$  and  $\mathcal{R}$  be equivalence relations on  $X$ .



- We write  $\mathcal{R} \subseteq \mathcal{Q}$  and say that  $\mathcal{R}$  *refines*  $\mathcal{Q}$  if considered as sets of ordered pairs we have  $\mathcal{R} \subseteq \mathcal{Q}$ .
- We define the *product equivalence relation*  $\mathcal{Q}^n$  on  $X^n$  by setting  $x_0 \dots x_{n-1} \sim x'_0 \dots x'_{n-1}$  if and only if  $x_i \sim x'_i$  for all  $i = 0, \dots, n-1$ .

Our equivalence relations will satisfy the following specifications. Let  $(\epsilon_n)_{n \in \mathbb{N}}$  be a decreasing sequence of positive numbers such that

$$(4.1) \quad \sum_{n \in \mathbb{N}} \epsilon_n < \infty.$$

To start, we let  $\mathcal{Q}_0^0$  be the equivalence relation on  $\mathcal{W}_0 = \Sigma$  which has one equivalence class, that is, any two elements of  $\Sigma$  are equivalent.

- (Q4) Suppose that  $n = M(s)$ . There is a specific number  $J_{s(n),n} \in \mathbb{Z}^+$  such that each word in  $w_n \in \mathcal{W}_n$  is a concatenation  $w_n = w_{n,1} \dots w_{n,J}$  of  $J = J_{s(n),n}$  strings of equal length. Then any two words in the same  $\mathcal{Q}_s^n$  class agree with each other except possibly on initial or final strings of length at most  $\frac{\epsilon_n}{2} \frac{h_n}{J_{s(n),n}}$  on the segments  $w_{n,i}$  for  $i = 1, \dots, J_{s(n),n}$ .
- (Q5) For  $n \geq M(s) + 1$  we can consider words in  $\mathcal{W}_n$  as concatenations of words from  $\mathcal{W}_{M(s)}$  and define  $\mathcal{Q}_s^n$  as the product equivalence relation of  $\mathcal{Q}_s^{M(s)}$ .
- (Q6)  $\mathcal{Q}_{s+1}^n$  refines  $\mathcal{Q}_s^n$  and each  $\mathcal{Q}_s^n$  class contains  $2^{4e(n)}$  many  $\mathcal{Q}_{s+1}^n$  classes.

*Remark.* By (Q5) we can view  $\mathcal{W}_n/\mathcal{Q}_s^n$  as sequences of elements  $\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)}$  and similarly for  $rev(\mathcal{W}_n)/\mathcal{Q}_s^n$ . In particular, it follows that  $\mathcal{Q}_0^n$  is the equivalence relation on  $\mathcal{W}_n$  which has one equivalence class. Moreover, it allows us to regard elements in  $\mathcal{W}_n/\mathcal{Q}_s^n$  for  $n \geq M(s) + 1$  as sequences of symbols in the alphabet  $(\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)})^*$ .

*Remark.* In case that the exponent is not relevant we will refer to the  $\mathcal{Q}_s^n$  as  $\mathcal{Q}_s$ . For  $u \in \mathcal{W}_n$  we write  $[u]_s$  for its  $\mathcal{Q}_s^n$  class.

As in [FRW11, section 5] these specifications are sufficient to describe a canonical tower of factors. For this purpose, we consider the symbolic shift  $\mathbb{K} \subset \Sigma^{\mathbb{Z}}$  with measure  $\nu$  corresponding to  $\Psi(\mathcal{T})$  for a fixed  $\mathcal{T} \in \mathcal{Trees}$ .

4.2.1. *The odometer factor.* We start by describing an odometer factor of our transformation which will turn out to be the Kronecker factor and will determine a natural “timing mechanism” to detect blocks.

Suppose that  $\sigma_m$  and  $\sigma_n$  with  $m < n$  are consecutive members of  $\mathcal{T}$  and let  $x \in \mathbb{K}$ . By unique readability there are unique  $w, w' \in \mathcal{W}_n$  and  $k \in \{0, \dots, h_n - 1\}$  such that  $x \upharpoonright [-k, -k + 2h_n - 1] = ww'$ . We call the interval of integers  $[-k, -k + h_n - 1]$  the  $n$ -block of  $x$  containing 0.

If the  $m$ -block of  $x$  containing 0 is given by  $[-k', -k' + h_m - 1]$ , then there is a unique  $\pi(x, n)$  such that  $k' = -k + \pi(x, n)h_m$ . We can view  $\pi(x, n)$  as an element in  $\mathbb{Z}_{h_n/h_m}$  and we define

$$\pi_0 : \mathbb{K} \rightarrow \prod \{ \mathbb{Z}_{h_n/h_m} : \sigma_m \text{ and } \sigma_n \text{ are successive elements of } \mathcal{T} \},$$

by  $\pi_0(x) = (\pi(x, n))_{\sigma_n \in \mathcal{T}}$ . Since  $\pi(sh(x), \cdot)$  is obtained from  $\pi(x, \cdot)$  by adding  $\bar{1} = (1, 0, 0, \dots)$  in the odometer  $\mathcal{O}_{\mathcal{T}} := \prod \{ \mathbb{Z}_{h_n/h_m} : \sigma_m \text{ and } \sigma_n \text{ are successive elements of } \mathcal{T} \}$ , we denote the corresponding odometer transformation by  $\mathcal{O}_{\mathcal{T}}$  and collect the following properties (see [FRW11, Lemma 22]).

**Lemma 28.** *The map  $\pi_0$  is a factor map from  $\Psi(\mathcal{T})$  to  $\mathcal{O}_{\mathcal{T}}$ . If  $p > 2$  is a prime number, then  $e^{2\pi i/p}$  is an eigenvalue of the unitary operator associated with  $\mathcal{O}_{\mathcal{T}}$  if and only if  $p = p_n$  for some  $n$  with  $\sigma_n \in \mathcal{T}$ .*

By the analogous procedure one associates the same odometer transformation  $\mathcal{O}_{\mathcal{T}}$  to  $\mathbb{K}^{-1}$  and we let  $\pi_0^* : \mathbb{K}^{-1} \rightarrow \prod \mathbb{Z}_{h_n/h_m}$  be the analogous map. We note that  $\pi_0^*(\text{rev}(x)) = i(\pi_0(x))$  for the involution  $i$  of  $\mathcal{O}_{\mathcal{T}}$  given by  $i(x) = -x$ .

4.2.2. *The canonical factors.* Now we define a canonical sequence of invariant sub- $\sigma$ -algebras of the algebra  $\mathcal{B}(\mathbb{K})$  of measurable subsets of  $\mathbb{K}$ .

We let  $\mathbb{K}_0$  be the odometer factor  $\mathcal{O}_{\mathcal{T}}$  introduced above. For each  $s \geq 1$  a typical  $x \in \mathbb{K}$  gives a bi-infinite sequence of the classes  $\{c_j^{(M(s),s)} : j = 1, \dots, Q_s^{M(s)}\}$  in  $\mathcal{Q}_s^{M(s)}$  which yields a shift-equivariant map from  $\mathbb{K}$  to  $\{1, \dots, Q_s^{M(s)}\}^{\mathbb{Z}}$ . To be more precise, we define a map  $\tilde{\pi}_s : \mathbb{K} \rightarrow \{1, \dots, Q_s^{M(s)}\} = (\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)})^*$  which assigns to a  $x \in \mathbb{K}$  the letter  $j$  if the word  $w \in \mathcal{W}_{M(s)}$  on the  $M(s)$ -block of  $x$  containing the position 0 satisfies  $[w]_s = c_j^{(M(s),s)}$ . Hereby, we define a shift-equivariant map  $\pi_s : \mathbb{K} \rightarrow \mathbb{K}_s := \{1, \dots, Q_s^{M(s)}\}^{\mathbb{Z}}$  by letting  $\pi_s(x) = (\tilde{\pi}_s(sh^k(x)))_{k \in \mathbb{Z}}$ . There is an analogous map from  $\text{rev}(\mathbb{K})$  to  $\text{rev}(\mathbb{K}_s)$  that we also denote by  $\pi_s$ .

Next, we describe a convenient base for the topology on  $\mathbb{K}_s$ : For  $n \geq M(s)$ ,  $w \in \mathcal{W}_n$  and  $0 \leq k < h_n$ , we let  $\langle [w]_s, k \rangle$  be the collection of  $(x, y) \in \mathbb{K}_s$  such that the position 0 is at the  $k$ -th place in the  $n$ -block  $B$  of  $x$  containing 0 and if  $v \in \{1, \dots, Q_s^{M(s)}\}^{h_n}$  is the word in  $x$  at the block  $B$ , then  $v$  is the sequence of  $\mathcal{Q}_s^{M(s)}$  classes given by  $[w]_s$ . Then the collection of those  $\langle [w]_s, k \rangle$  for  $\sigma_n \in \mathcal{T}$ ,  $w \in \mathcal{W}_n$  and  $0 \leq k < h_n$  forms a basis for the topology of  $\mathbb{K}_s$  consisting of clopen sets. In particular, for  $m = M(s)$  a word  $w \in \mathcal{W}_m$  gives a word of length  $h_m$  in our alphabet  $(\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)})^*$  that is the repetition of the same letter. We denote the collection of these words by  $(\mathcal{W}_m)_s^*$ . Then for  $n > m = M(s)$  with  $\sigma_n \in \mathcal{T}$  each word in  $\mathcal{W}_n$  is a concatenation of  $\frac{h_n}{h_m}$  words in  $\mathcal{W}_m$  by specification (E2) and, thus, determines a sequence of  $\frac{h_n}{h_m}$  many elements of  $(\mathcal{W}_m)_s^*$ . We let  $(\mathcal{W}_n)_s^*$  be the collection of words in the alphabet  $(\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)})^*$  arising this way. Then the sequence  $((\mathcal{W}_n)_s^*)_{n \geq M(s)}$  gives a well-defined odometer-based construction sequence for  $\mathbb{K}_s$  in the alphabet  $(\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)})^*$ .

We also define the measure  $\nu_s := \pi_s^* \nu$  on  $\mathbb{K}_s$ . To be more explicit, with the aid of specifications (E2) and (Q6) we see that  $\nu_s(\langle [w]_s, k \rangle) = \frac{1}{h_n Q_s^n}$  for any  $w \in \mathcal{W}_n$  and  $0 \leq k < h_n$ .

Finally, we let  $\mathcal{H}_0$  be the shift-invariant sub- $\sigma$ -algebra of the  $\mathcal{B}(\mathbb{K})$  generated by the collection of  $\pi_0^{-1}(B)$ , where  $B$  is a basic open set in  $\mathcal{O}_{\mathcal{T}}$ , and we let  $\mathcal{H}_s$  be the shift-invariant sub- $\sigma$ -algebra of the  $\mathcal{B}(\mathbb{K})$  generated by the collection of  $\pi_s^{-1}(B)$ , where  $B$  is a basic open set in  $\mathbb{K}_s$ . Then  $\mathcal{H}_s$  is the sub- $\sigma$ -algebra determined by the factor map  $\pi_s$ .

Since the equivalence relation  $\mathcal{Q}_{s+1}^{M(s+1)}$  refines  $(\mathcal{Q}_s^{M(s)})^{h_{M(s+1)}/h_{M(s)}}$  by specifications (E2) and (Q6), we have  $\mathcal{H}_{s+1} \supseteq \mathcal{H}_s$  and a continuous factor map  $\pi_{s+1,s} :$

$\mathbb{K}_{s+1} \rightarrow \mathbb{K}_s$ . To express this one explicitly, we note that a  $\mathbb{Z}$ -sequence of  $\mathcal{Q}_{s+1}^{M(s+1)}$  classes determines a sequence of  $\mathcal{Q}_s^{M(s)}$  classes, because a  $\mathcal{Q}_{s+1}^{M(s+1)}$  class is contained in a  $\mathcal{Q}_s^{M(s+1)}$  class which is a  $h_{M(s+1)}/h_{M(s)}$ -tuple of  $\mathcal{Q}_s^{M(s)}$  classes. Analogously we define a map from  $rev(\mathbb{K}_{s+1})$  to  $rev(\mathbb{K}_s)$  that we also denote by  $\pi_{s+1,s}$ .

In analogy with [FRW11, Proposition 23] we can prove the following statement making use of specification (Q4).

**Lemma 29.**  $\mathcal{B}(\mathbb{K})$  is the smallest invariant  $\sigma$ -algebra that contains  $\bigcup_s \mathcal{H}_s$ .

*Proof.* It suffices to show that for any  $m \in \mathbb{N}$  and any  $u \in \mathcal{W}_m$  the basic open set  $\langle u \rangle \subseteq \mathbb{K}$  can be approximated arbitrarily well in measure by elements of  $\bigcup_s \mathcal{H}_s$ . For this purpose, we fix  $\varepsilon > 0$  and choose  $n > m$  such that  $n = M(s)$  for some  $s \in \mathbb{N}$  and  $\varepsilon_n + 2 \frac{h_m J_{s(n),n}}{h_n} < \varepsilon$ . Then we let  $G \subset \mathbb{K}$  be the collection of  $x$  such that within the  $n$ -word  $w_n \in \mathcal{W}_n$  located at the  $n$ -block of  $x$  containing the position 0,  $x(0)$  is not among the first or last  $\left(\frac{\varepsilon_n}{2} + \frac{h_m J_{s(n),n}}{h_n}\right) \frac{h_n}{J_{s(n),n}} = \frac{\varepsilon_n}{2} \frac{h_n}{J_{s(n),n}} + h_m$  letters of one of the  $J_{s(n),n}$  many segments  $w_{n,i}$  in specification (Q4), that is,  $x(0)$  belongs to one of the center segments. By the ergodic theorem and our assumption on  $n$  the measure of  $G$  is at least  $1 - \varepsilon$ .

Now let  $x \in G \cap \langle u \rangle$  and let  $w \in \mathcal{W}_n$  be located at the  $n$ -block  $B = [-k, -k + h_n]$  of  $x$  containing 0. If  $c_j^{(n,s)}$  is the  $\mathcal{Q}_s^{M(s)}$  class of  $w$ , then

$$sh^{-k}(x) \in \pi_s^{-1} \left( \left\langle c_j^{(n,s)}, 0 \right\rangle \right) \subseteq sh^{-k}(\langle u \rangle)$$

by specification (Q4). Hence,  $G \cap \langle u \rangle$  is a union of shifts of sets of the form  $\pi_s^{-1} \left( \left\langle c_j^{(n,s)}, 0 \right\rangle \right)$  lying in  $\mathcal{H}_s$ .  $\square$

As in the proof above, for  $n = M(s)$ , the set  $L_n$  of  $x$  such that within the  $n$ -word  $w_n \in \mathcal{W}_n$  located at the  $n$ -block of  $x$  containing the position 0 we have that  $x(0)$  is among the first or last  $\frac{\varepsilon_n}{2} \frac{h_n}{J_{s(n),n}}$  letters of one of the  $J_{s(n),n}$  many segments  $w_{n,i}$  in specification (Q4), has measure  $\varepsilon_n$ . Let  $L$  be the collection of  $x \in \mathbb{K}$  such that for all large  $s$ , if  $n = M(s)$  then  $x$  does not belong to  $L_n$ , that is,  $x(0)$  is not among the first or last  $\frac{\varepsilon_n}{2} \frac{h_n}{J_{s(n),n}}$  letters of one of the  $J_{s(n),n}$  many segments  $w_{n,i}$  within the  $n$ -word  $w_n \in \mathcal{W}_n$  located at the  $n$ -block of  $x$  containing the position 0. Then  $L$  has measure one by equation (4.1) and the Borel-Cantelli Lemma.

We collect the following properties as in Propositions 24 and 25 of [FRW11].

**Lemma 30.** (1) For all  $x \neq y$  belonging to  $L$ , there is an open set  $S \in \bigcup_s \mathcal{H}_s$  such that  $x \in S$  and  $y \notin S$ .

(2) For all  $s \geq 1$ ,  $\mathcal{H}_s$  is a strict subalgebra of  $\mathcal{H}_{s+1}$ . Moreover, if  $\{\nu_x : x \in \mathbb{K}_s\}$  is the disintegration of  $\nu_{s+1}$  over  $\nu_s$ , then for  $\nu_s$ -a.e.  $x$  the measure  $\nu_x$  is nonatomic.

**4.3. Groups of involutions.** To approximate conjugacies we will use groups associated to trees as in [FRW11]. These groups will be direct sums of  $\mathbb{Z}_2 = \mathbb{Z}/2\mathbb{Z}$  and are called *groups of involutions*.

If  $G$  is a group of involutions and  $B = \{r_i : i \in I\}$  is a distinguished basis (that is, a particular choice of basis for  $G$  considered as a vector space over  $\mathbb{Z}_2$ ), then we call the elements  $r_i \in B$  *canonical generators* and we have a well-defined notion of *parity* for elements in  $G$ : an element  $g \in G$  is called *even* if it can be written as

the sum of an even number of elements in  $B$ . Otherwise, it is called *odd*. Parity is preserved under homomorphisms sending the canonical generators of one group to the canonical generators of the other. Moreover, for an inverse limit system of groups of involutions  $\{G_s : s \in \mathbb{Z}^+\}$  (where each group has a set of canonical generators) with parity-preserving homomorphisms  $\rho_{t,s} : G_t \rightarrow G_s$  for  $0 < s < t$ , the elements of the inverse limit  $\varprojlim G_s$  have a well-defined parity.

For a tree  $\mathcal{T} \subset \mathbb{N}^{\mathbb{N}}$  we assign to each level  $s > 0$  a group of involutions  $G_s(\mathcal{T})$  by taking a sum of copies of  $\mathbb{Z}_2$  indexed by the nodes of  $\mathcal{T}$  at level  $s$ . Let

$$G_s(\mathcal{T}) = \sum_{\tau \in \mathcal{T}, lh(\tau)=s} (\mathbb{Z}_2)_\tau.$$

Each node of  $\mathcal{T}$  at level  $s$  corresponds to an element of  $G_s(\mathcal{T})$  that is 1 in the copy of  $\mathbb{Z}_2$  for that node and 0 in the other copies of  $\mathbb{Z}_2$ . We also define  $G_0(\mathcal{T})$  to be the trivial group. For levels  $0 < s < t$  of  $\mathcal{T}$  we have a canonical homomorphism  $\rho_{t,s} : G_t(\mathcal{T}) \rightarrow G_s(\mathcal{T})$  that sends a generator  $\tau$  of  $G_t(\mathcal{T})$  to the unique generator  $\sigma$  of  $G_s(\mathcal{T})$  that is an initial segment of  $\tau$ . The map  $\rho_{t,0}$  is the trivial homomorphism  $\rho_{t,0} : G_t(\mathcal{T}) \rightarrow G_0(\mathcal{T}) = \{0\}$ . We denote the inverse limit of  $\langle G_s(\mathcal{T}), \rho_{t,s} : s < t \rangle$  by  $G_\infty(\mathcal{T})$  and we let  $\rho_s : G_\infty(\mathcal{T}) \rightarrow G_s(\mathcal{T})$  be the projection map.

Since there is a one-to-one correspondence between the infinite branches of  $\mathcal{T}$  and infinite sequences  $(g_s)_{s \in \mathbb{Z}^+}$  of generators  $g_s \in G_s(\mathcal{T})$  with  $\rho_{t,s}(g_t) = g_s$  for  $t > s > 0$ , we obtain the following characterization.

**Fact 31.** *Let  $\mathcal{T} \subset \mathbb{N}^{\mathbb{N}}$  be a tree. Then  $G_\infty(\mathcal{T})$  has a nonidentity element of odd parity if and only if  $\mathcal{T}$  is ill-founded (that is,  $\mathcal{T}$  has an infinite branch).*

In order to make the elements of  $G_\infty(\mathcal{T})$  correspond to conjugacies, we will build symmetries into our construction using the following finite approximations to  $G_\infty(\mathcal{T})$ . We let  $G_0^n(\mathcal{T})$  be the trivial group and for  $s > 0$  we let

$$G_s^n(\mathcal{T}) = \sum (\mathbb{Z}_2)_\tau \text{ where the sum is taken over } \tau \in \mathcal{T} \cap \{\sigma_m : m \leq n\}, lh(\tau) = s.$$

When  $\mathcal{T}$  is clear from the context, we will frequently write  $G_s^n$ . We also introduce the finite approximations  $\rho_{t,s}^{(n)} : G_t^n(\mathcal{T}) \rightarrow G_s^n(\mathcal{T})$  to the canonical homomorphisms.

During the course of construction we will consider group actions of  $G_s^n$  on our quotient spaces  $\mathcal{W}_n/\mathcal{Q}_s^n$ . Here, we will need to control systems of such group actions on the refining equivalence relations. For that purpose, the following general definitions will prove useful.

**Definition 32.** Suppose

- $\mathcal{Q}$  and  $\mathcal{R}$  are equivalence relations on a set  $X$  with  $\mathcal{R}$  refining  $\mathcal{Q}$ ,
- $G$  and  $H$  are groups with  $G$  acting on  $X/\mathcal{Q}$  and  $H$  acting on  $X/\mathcal{R}$ ,
- $\rho : H \rightarrow G$  is a homomorphism.

Then we say that the  $H$  action is *subordinate* to the  $G$  action if for all  $x \in X$ , whenever  $[x]_{\mathcal{R}} \subset [x]_{\mathcal{Q}}$  we have  $h[x]_{\mathcal{R}} \subset \rho(h)[x]_{\mathcal{Q}}$ .

**Definition 33.** If  $G$  acts on  $X$ , then the canonical *diagonal action* of  $G$  on  $X^n$  is defined by

$$g(x_0x_1 \dots x_{n-1}) = gx_0gx_1 \dots gx_{n-1} \text{ for any } g \in G.$$

If  $G$  is a group of involutions with a collection of canonical generators, then we define the *skew diagonal action* on  $X^n$  by setting

$$g(x_0x_1\dots x_{n-1}) = \begin{cases} gx_0gx_1\dots gx_{n-1}, & \text{if } g \in G \text{ is of even parity,} \\ gx_{n-1}gx_{n-2}\dots gx_0, & \text{if } g \in G \text{ is of odd parity.} \end{cases}$$

*Remark.* Recalling the notion of a product equivalence relation from Definition 27 we can identify  $X^n/\mathcal{Q}^n$  with  $(X/\mathcal{Q})^n$  in an obvious way. Then we can also extend an action of  $G$  on  $X/\mathcal{Q}$  to the diagonal or skew diagonal actions on  $(X/\mathcal{Q})^n$  in a straightforward way.

To extend our group actions we will use the following extension lemma from [FRW11, Lemma 46].

**Lemma 34.** *Let  $X$  be a set and  $\mathcal{R} \subseteq \mathcal{Q}$  be equivalence relations on  $X$ . Suppose that*

- $\rho : H \rightarrow G \times \mathbb{Z}_2$  is a homomorphism,
- $G \times \mathbb{Z}_2$  acts on  $X/\mathcal{Q}$ ,
- $H$  acts on  $X/\mathcal{R}$  by a free action subordinate to the  $G \times \mathbb{Z}_2$  action on  $X/\mathcal{Q}$ ,
- if  $H_1 = \{h \in H : \rho(h) = (0, i), i \in \{0, 1\}\}$ , then every orbit of the  $\mathbb{Z}_2$  factor of  $G \times \mathbb{Z}_2$  contains an even number of  $H_1$  orbits.

Then there is a free action of  $H \times \mathbb{Z}_2$  on  $X/\mathcal{R}$  subordinate to the  $G \times \mathbb{Z}_2$  action via the map  $\rho'(h, i) = \rho(h) + (0, i)$ .

We now list specifications on the group actions.

- (A7)  $G_s^n$  acts freely on  $\mathcal{W}_n/\mathcal{Q}_s^n$  and the  $G_s^n$  action is subordinate to the  $G_{s-1}^n$  action on  $\mathcal{W}_n/\mathcal{Q}_{s-1}^n$  via the canonical homomorphism  $\rho_{s,s-1}^{(n)} : G_s^n \rightarrow G_{s-1}^n$ .
- (A8) Suppose  $M(s) < n$ ,  $\sigma_m$  and  $\sigma_n$  are consecutive elements of  $\mathcal{T}$  and we view  $G_s^n = G_s^m \oplus H$ . Then the action of  $G_s^m$  on  $\mathcal{W}_m/\mathcal{Q}_s^m$  is extended to an action on  $\mathcal{W}_n/\mathcal{Q}_s^n$  by the skew diagonal action.

*Remark 35.* In particular, in the above situation with  $M(s) < n$  and  $\sigma_m, \sigma_n$  consecutive elements of  $\mathcal{T}$  both specifications together yield that  $\mathcal{W}_n/\mathcal{Q}_s^n$  is closed under the skew diagonal action by  $G_s^m$ . Clearly, this also holds if we view each element in  $\mathcal{W}_n/\mathcal{Q}_s^n$  as a sequence in the alphabet  $(\mathcal{W}_{M(s)}/\mathcal{Q}_s^{M(s)})^*$ .

## 5. INFINITE BRANCHES GIVE ISOMORPHISMS

As in [FRW11] the specifications on our transformations stated in the previous section suffice to build an isomorphism between  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  in case that the tree  $\mathcal{T}$  has an infinite branch.

**Lemma 36.** *Let  $s \in \mathbb{N}$  and  $g \in G_s^m$  for some  $m \in \mathbb{N}$ . Suppose that  $g$  has odd parity. Then there is a shift-equivariant isomorphism  $\eta_g : \mathbb{K}_s \rightarrow \text{rev}(\mathbb{K}_s)$  canonically associated to  $g$  as in (5.1).*

Moreover, if  $s' > s$  and  $g' \in G_{s'}^n$  for some  $n \geq m$  with  $\rho_{s',s}(g') = g$ , then  $\pi_{s',s} \circ \eta_{g'} = \eta_g \circ \pi_{s',s}$ .

*Proof.* We recall from Section 4.2 that  $((\mathcal{W}_m)_s^*)_{m \geq M(s)}$  defines a construction sequence for  $\mathbb{K}_s$ . Similarly,  $(\text{rev}((\mathcal{W}_m)_s^*))_{m \geq M(s)}$  is a construction sequence for  $\text{rev}(\mathbb{K}_s)$ . For  $n > m$  there is  $K \in \mathbb{N}$  such that we can write any element  $[w]_s^* \in (\mathcal{W}_n)_s^*$  as  $[w]_s^* = [w_0]_s^*[w_1]_s^* \dots [w_{K-1}]_s^*$  with  $[w_i]_s^* \in (\mathcal{W}_m)_s^*$  for  $i = 0, \dots, K-1$ .

By specification (A8) the group action is given by the skew diagonal action and, hence, we have

$$g([w_0]_s^*[w_1]_s^* \dots [w_{K-1}]_s^*) = g[w_{K-1}]_s^* g[w_{K-2}]_s^* \dots g[w_0]_s^*.$$

As pointed out in Remark 35,  $(\mathcal{W}_n)_s^*$  is closed under the skew diagonal action. Thus  $g[w_{K-1}]_s^* g[w_{K-2}]_s^* \dots g[w_0]_s^* \in (\mathcal{W}_n)_s^*$ , which implies  $g[w_0]_s^* g[w_1]_s^* \dots g[w_{K-1}]_s^* \in \text{rev}((\mathcal{W}_n)_s^*)$ . Hence the map

$$(5.1) \quad [w_0]_s^*[w_1]_s^* \dots [w_{K-1}]_s^* \mapsto g[w_0]_s^* g[w_1]_s^* \dots g[w_{K-1}]_s^*$$

is an invertible map from the construction sequence for  $\mathbb{K}_s$  to the construction sequence for  $\text{rev}(\mathbb{K}_s)$ . It can also be interpreted as a shift-equivariant map from cylinder sets in  $\mathbb{K}_s$  to cylinder sets located in the same position in  $\text{rev}(\mathbb{K}_s)$ . This yields the isomorphism in the first assertion.

The second assertion follows from specification (A7), which says that the action by  $g'$  is subordinate to the action by  $g$  via the homomorphism  $\rho_{s',s}$ .  $\square$

**Definition 37.** Let  $s \in \mathbb{N}$  and  $g \in G_s^m$  for some  $m \in \mathbb{N}$ . Suppose that  $g$  has odd parity. Then  $g$  yields a shift-equivariant isomorphism  $\eta_g : \mathbb{K}_s \rightarrow \text{rev}(\mathbb{K}_s)$  as in Lemma 36. For every  $n \geq m$  it also induces a map  $\mathcal{W}_n/\mathcal{Q}_s^n \rightarrow \text{rev}(\mathcal{W}_n)/\mathcal{Q}_s^n$  that we denote by  $\eta_g$  as well.

In the following we call a sequence of isomorphisms  $\zeta_s$  between  $\mathbb{K}_s$  and  $\text{rev}(\mathbb{K}_s)$  *coherent* if  $\pi_{s+1,s} \circ \zeta_{s+1} = \zeta_s \circ \pi_{s+1,s}$  for every  $s \in \mathbb{N}$ .

**Lemma 38.** Let  $(\zeta_s)_{s \in \mathbb{N}}$  be a coherent sequence of isomorphisms between  $\mathbb{K}_s$  and  $\text{rev}(\mathbb{K}_s)$ . Then there is an isomorphism  $\zeta : \mathbb{K} \rightarrow \text{rev}(\mathbb{K})$  such that  $\pi_s \circ \zeta = \zeta_s \circ \pi_s$  for every  $s \in \mathbb{N}$ .

*Proof.* Since the isomorphisms  $\zeta_s$  cohere, their inverse limit defines a measure-preserving isomorphism between the subalgebra of  $\mathcal{B}(\mathbb{K})$  generated by  $\bigcup_s \mathcal{H}_s$  and the subalgebra of  $\mathcal{B}(\text{rev}(\mathbb{K}))$  generated by  $\bigcup_s \text{rev}(\mathcal{H}_s)$ . By Lemma 29 this extends uniquely to a measure-preserving isomorphism  $\tilde{\zeta}$  between  $\mathcal{B}(\mathbb{K})$  and  $\mathcal{B}(\text{rev}(\mathbb{K}))$ . Then by part (1) of Lemma 30 we can find sets  $D \subset \mathbb{K}$ ,  $D' \subset \text{rev}(\mathbb{K})$  of measure zero such that  $\tilde{\zeta}$  determines a shift-equivariant isomorphism  $\zeta$  between  $\mathbb{K} \setminus D$  and  $\text{rev}(\mathbb{K}) \setminus D'$ .  $\square$

Now we are ready to prove the first half of Proposition 23.

*Proof of part (1) in Proposition 23.* Suppose that  $\mathcal{T} \in \mathcal{Trees}$  has an infinite branch. Then  $G_\infty(\mathcal{T})$  has an element  $g$  of odd parity according to Fact 31. By Lemma 36 we obtain a coherent sequence of isomorphisms  $\zeta_s := \eta_{\rho_s(g)}$  between  $\mathbb{K}_s$  and  $\text{rev}(\mathbb{K}_s)$ . Hence, Lemma 38 yields an isomorphism between  $\mathbb{K}$  and  $\text{rev}(\mathbb{K})$ . Since  $\text{rev}(\mathbb{K})$  is isomorphic to  $\mathbb{K}^{-1}$ , we conclude that  $\mathbb{K} \cong \mathbb{K}^{-1}$ .  $\square$

## 6. FELDMAN PATTERNS

In [Fe76] Feldman constructed the first example of an ergodic zero-entropy automorphism that is not loosely Bernoulli. The construction is based on the idea that no pair of the following strings

*abababab*  
*aabbaabb*  
*aaaabbbb*

can be matched very well. In our construction we will concatenate variants of his blocks that we call *Feldman patterns*.

We start with an observation that follows along the lines of Lemma 6.5 and Remark 6.6 in [GKpp], whose proofs were inspired by [Fe76, Theorem 4] and [ORW82, Proposition 1.1, p.79].

**Lemma 39.** *Suppose  $a_1, a_2, \dots, a_N$  are distinct symbols in an alphabet  $\Sigma$ . Let  $M, S, T, j, k \in \mathbb{Z}^+$ ,  $S \geq T$ ,  $M \geq j > k$ , and*

$$B_k = \left( a_1^{TN^{2k}} a_2^{TN^{2k}} \dots a_N^{TN^{2k}} \right)^{N^{2(M+1-k)}}, \quad B_j = \left( a_1^{SN^{2j}} a_2^{SN^{2j}} \dots a_N^{SN^{2j}} \right)^{N^{2(M+1-j)}}.$$

*Suppose  $B$  and  $\bar{B}$  are strings of consecutive symbols in  $B_j$  and  $B_k$ , respectively, where  $|B| \geq SN^{2M+2}$ . Assume that  $N \geq 20$ . Then*

$$\bar{f}(B, \bar{B}) > 1 - \frac{4}{\sqrt{N}}.$$

*Proof.* By removing fewer than  $2SN^{2j}$  symbols from the beginning and end of  $B$ , we can decompose the remaining part of  $B$  into strings  $C_1, C_2, \dots, C_r$  each of the form  $a_i^{SN^{2j}}$ . Since  $2SN^{2j} \leq 2 \frac{|B|+|\bar{B}|}{N^2}$ , it follows from Fact 9 that removing these symbols increases the  $\bar{f}$  distance between  $B$  and  $\bar{B}$  by less than  $\frac{4}{N^2}$ . Let  $\bar{C}_1, \bar{C}_2, \dots, \bar{C}_r$  be the decomposition of  $\bar{B}$  into substrings corresponding to  $C_1, C_2, \dots, C_r$  under a best possible match between  $C_1 C_2 \dots C_r$  and  $\bar{B}$ .

Let  $i \in \{1, 2, \dots, r\}$ .

*Case 1.*  $|\bar{C}_i| < \frac{3}{2\sqrt{N}}|C_i|$ . Then by Fact 11,  $\bar{f}(C_i, \bar{C}_i) > 1 - \frac{3}{\sqrt{N}}$ .

*Case 2.*  $|\bar{C}_i| \geq \frac{3}{2\sqrt{N}}|C_i| = \frac{3}{2}SN^{2j-(1/2)}$ . The length of a cycle  $a_1^{TN^{2k}} a_2^{TN^{2k}} \dots a_N^{TN^{2k}}$  in  $B_k$  is at most  $TN^{2j-1}$ . Therefore  $\bar{C}_i$  contains at least  $\lfloor \frac{3\sqrt{N}}{2} \rfloor - 1 > \sqrt{N}$  complete cycles. Thus deleting any partial cycles at the beginning and end of  $\bar{C}_i$  would increase the  $\bar{f}$  distance between  $C_i$  and  $\bar{C}_i$  by less than  $\frac{2}{\sqrt{N}}$ . On the rest of  $\bar{C}_i$ , only  $\frac{1}{N}$  of the symbols can match the symbol in  $C_i$ . Thus  $\bar{f}(C_i, \bar{C}_i) > 1 - \frac{2}{\sqrt{N}} - \frac{2}{N} > 1 - \frac{3}{\sqrt{N}}$ .

Therefore, by Fact 10,  $\bar{f}(C_1 C_2 \dots C_r, \bar{C}_1, \bar{C}_2, \dots, \bar{C}_r) > 1 - \frac{3}{\sqrt{N}}$ . Hence  $\bar{f}(B, \bar{B}) > 1 - \frac{3}{\sqrt{N}} - \frac{4}{N^2} > 1 - \frac{4}{\sqrt{N}}$ .  $\square$

Let  $T, N, M \in \mathbb{Z}^+$ . A  $(T, N, M)$ -Feldman pattern in building blocks  $A_1, \dots, A_N$  of equal length  $L$  is one of the strings  $B_1, \dots, B_M$  that are defined by

$$\begin{aligned} B_1 &= \left( A_1^{TN^2} A_2^{TN^2} \dots A_N^{TN^2} \right)^{N^{2M}} \\ B_2 &= \left( A_1^{TN^4} A_2^{TN^4} \dots A_N^{TN^4} \right)^{N^{2M-2}} \\ &\vdots \\ B_M &= \left( A_1^{TN^{2M}} A_2^{TN^{2M}} \dots A_N^{TN^{2M}} \right)^{N^2} \end{aligned}$$

Thus  $N$  denotes the number of building blocks,  $M$  is the number of constructed patterns, and  $TN^2$  gives the minimum number of consecutive occurrences of a building block. We also note that each building block  $A_i$ ,  $1 \leq i \leq N$ , occurs  $TN^{2M+2}$  many times in each pattern. Every block  $B_j$ ,  $1 \leq j \leq M$ , has total length

$TN^{2M+3}L$ . Moreover, we notice that  $B_j$  is built with  $N^{2(M+1-j)}$  many so-called *cycles*: Each cycle winds through all the  $N$  building blocks.

We conclude this analysis with a statement on the  $\bar{f}$ -distance between different  $(T, N, M)$ -Feldman patterns.

**Proposition 40** (Distance between different Feldman patterns in  $\bar{f}$ ). *Let  $N \geq 20$ ,  $M \geq 2$ , and  $B_j$ ,  $1 \leq j \leq M$ , be the  $(T, N, M)$ -Feldman patterns in the building blocks  $A_1, \dots, A_N$  of equal length  $L$ . Assume that  $\alpha \in (0, \frac{1}{7})$ ,  $R \geq 2$ , and  $\bar{f}(C, D) > \alpha$ , for all substrings  $C$  and  $D$  consisting of consecutive symbols from  $A_{i_1}$  and  $A_{i_2}$ , respectively, where  $i_1 \neq i_2$ , with  $|C|, |D| \geq \frac{L}{R}$ .*

*Then for all  $j, k \in \{1, \dots, M\}$ ,  $j \neq k$ , and all sequences  $B$  and  $\bar{B}$  of at least  $TN^{2M+2}L$  consecutive symbols from  $B_j$  and  $B_k$ , respectively, we have*

$$(6.1) \quad \bar{f}(B, \bar{B}) \geq \alpha - \frac{4}{\sqrt{N}} - \frac{1}{R}.$$

*Proof.* Using Lemma 13 the statement follows from Lemma 39 with  $S = T$ .  $\square$

In our iterative construction process, that we present in Sections 7 and 8, we substitute Feldman patterns of finer equivalence classes of words into Feldman patterns of coarser classes. In Section 9 we need estimates on the  $\bar{f}$  distance between such different Feldman patterns of equivalence classes even under finite coding. Since we do not know the precise words being coded but just their equivalence classes, the following Coding Lemma will prove useful, especially in Lemma 62 and Lemma 71.

*Remark.* The Coding Lemma below can be applied in the case  $A = B_k$  and  $B = B_j$ , where  $B_k$  and  $B_j$  are as in Lemma 39, with  $k \neq j$ , except in the definition of  $B_j$ , the  $a_i$ 's are replaced by  $b_i$ 's, and we assume  $T = S$ . In case  $j > k$ , each  $\Lambda_{im}$  in the Coding Lemma is equal to  $N^{2(j-k)-1}$  cycles in  $B_k$ , and the permutations are the identity, while  $\Upsilon_{im} = b_m^{TN^{2j}}$ . If  $k > j$ , each substring  $\Lambda_{i_1}\Lambda_{i_2}\dots\Lambda_{i_N}$  of  $B_k$  actually consists of repetitions of a single symbol, and we again take  $\Upsilon_{im} = b_m^{TN^{2j}}$ . The permutations in the Coding Lemma allow for the more general situation that occurs in Lemma 45 and Remark 51.

**Lemma 41** (Coding Lemma). *Let  $B_{mj}^{(i)}$ ,  $1 \leq i \leq z, 1 \leq m \leq N, 1 \leq j \leq q$ , and  $A_{ij}$ ,  $1 \leq i \leq z, 1 \leq j \leq q$ , be blocks of symbols, with each block of length  $L$ . Assume  $q \geq N > 2^{16}$ . Suppose  $R_1 \geq 2$ ,  $\alpha \in (0, 1/7)$ , and for all substrings  $C$  and  $D$  consisting of consecutive symbols in  $B_{mj}^{(i)}$  and  $B_{m'j'}^{(i')}$ , respectively, with  $|C|, |D| \geq L/R_1$ , we have  $\bar{f}(C, D) \geq \alpha$  if  $m \neq m'$ . For  $1 \leq i \leq z, 1 \leq m \leq N$ , let  $\Lambda_{im}$  be a permutation of the strings  $A_{i1}, A_{i2}, \dots, A_{iq}$ , and let  $\Upsilon_{im} = B_{m1}^{(i)}B_{m2}^{(i)}\dots B_{mq}^{(i)}$ . Let*

$$A = \Lambda_{11}\Lambda_{12}\dots\Lambda_{1N}\Lambda_{21}\Lambda_{22}\dots\Lambda_{2N}\dots\Lambda_{z1}\Lambda_{z2}\dots\Lambda_{zN}$$

and

$$B = \Upsilon_{11}\Upsilon_{12}\dots\Upsilon_{1N}\Upsilon_{21}\Upsilon_{22}\dots\Upsilon_{2N}\dots\Upsilon_{z1}\Upsilon_{z2}\dots\Upsilon_{zN}.$$

*Suppose  $R_2 \geq 2$  and  $\bar{A}, \bar{B}$  are strings of consecutive symbols in  $A, B$ , respectively, each of length at least  $zqNL/R_2$ . Then*

$$\bar{f}(\bar{A}, \bar{B}) \geq \alpha \left[ \frac{1}{8} - \frac{2}{\sqrt[4]{N}} \right] - \frac{1}{R_1} - \frac{4R_2}{z}.$$



*Proof.* We begin by considering, instead of strings of symbols  $B_{mj}^{(\ell)}$  and  $A_{ij}$ , individual symbols,  $b_{mj}^{(\ell)}$ ,  $1 \leq \ell \leq z''$ ,  $1 \leq m \leq N$ ,  $1 \leq j \leq q$ , such that  $b_{mj}^{(\ell)} \neq b_{m'j'}^{(\ell')}$  for  $m \neq m'$ , and  $a_{ij}$ ,  $1 \leq i \leq z'$ ,  $1 \leq j \leq q$ . Let  $\mathcal{A}_{ij}$  be a permutation of  $a_{i1}, a_{i2}, \dots, a_{iq}$  and let  $\mathcal{B}_m^{(\ell)} = b_{m1}^{(\ell)} b_{m2}^{(\ell)} \dots b_{mq}^{(\ell)}$  for some  $1 \leq \ell \leq z''$ . Let  $R_3 = \sqrt[4]{N}$  and  $R_4 = \sqrt{N}$ . If a substring  $\bar{\mathcal{A}}_{i,j}$  of  $\mathcal{A}_{i,j}$  of length at least  $q/R_3$  and a substring  $\bar{\mathcal{B}}_m^{(\ell)}$  of  $\mathcal{B}_m^{(\ell)}$  satisfy  $\bar{f}(\bar{\mathcal{A}}_{i,j}, \bar{\mathcal{B}}_m^{(\ell)}) \leq 1/8$ , then at least  $7/9$  of the symbols in  $\bar{\mathcal{A}}_{i,j}$  are equal to a symbol in  $\{b_{m1}^{(\ell)}, b_{m2}^{(\ell)}, \dots, b_{mq}^{(\ell)}\}$ . Then at least  $7q/(9R_3)$  symbols among  $a_{i1}, a_{i2}, \dots, a_{iq}$  are elements of  $\{b_{m1}^{(\ell)}, b_{m2}^{(\ell)}, \dots, b_{mq}^{(\ell)}\}$ . This can happen for at most  $9R_3/7$  values of  $m$ . For the other values of  $m$ , and all values of  $j$  and  $\ell$ ,  $\bar{f}(\bar{\mathcal{A}}_{i,j}, \bar{\mathcal{B}}_m^{(\ell)}) > 1/8$ . Now suppose  $\mathcal{A}_i$  and  $\mathcal{B}^{(\ell)}$  are substrings consisting of consecutive symbols in  $\mathcal{A}_{i1}\mathcal{A}_{i2} \dots \mathcal{A}_{iN}$ , and  $\mathcal{B}_1^{(\ell)}\mathcal{B}_2^{(\ell)} \dots \mathcal{B}_N^{(\ell)}$ , respectively, both of length at least  $qN/R_4$ . Then there are at least  $(N/R_4) - 2$  complete  $\mathcal{A}_{ij}$ 's in  $\mathcal{A}_i$  and at least  $(N/R_4) - 2$  complete  $\mathcal{B}_m^{(\ell)}$ 's in  $\mathcal{B}$ . By Fact 9, if we remove the incomplete strings at the beginning and end of  $\mathcal{A}_i$  and  $\mathcal{B}^{(\ell)}$  we increase the  $\bar{f}$  distance by at most  $4R_4/N$ . We apply Lemma 13 with  $\alpha = 1/8$  to the resulting complete strings and then compensate by subtracting  $4R_4/N$ . Thus

$$\bar{f}(\mathcal{A}_i, \mathcal{B}^{(\ell)}) > \frac{1}{8} \left( 1 - \frac{9R_3/7}{(N/R_4) - 2} \right) - \frac{1}{R_3} - \frac{4R_4}{N} > \frac{1}{8} - \frac{7/6}{\sqrt[4]{N}} - \frac{4}{\sqrt{N}} =: \beta.$$

Therefore by Lemma 13,

$$(6.2) \quad \bar{f}(\mathcal{A}_1\mathcal{A}_2 \dots \mathcal{A}_{z'}, \mathcal{B}^{(1)}\mathcal{B}^{(2)} \dots \mathcal{B}^{(z'')}) > \beta - \frac{1}{R_4}.$$

Suppose  $\bar{A}$  and  $\bar{B}$  are as in the statement of the lemma. We will apply Lemma 13 again, now regarding  $\mathcal{A}_1\mathcal{A}_2 \dots \mathcal{A}_{z'}$  and  $\mathcal{B}^{(1)}\mathcal{B}^{(2)} \dots \mathcal{B}^{(z'')}$  as indices on the blocks in  $\bar{A}$  and  $\bar{B}$ . By removing any incomplete sequence of blocks at the beginning and end of  $\bar{A}$  and  $\bar{B}$ , we may assume that  $\bar{A}$  consists of complete sequences  $\Lambda_{i1}\Lambda_{i2} \dots \Lambda_{iN}$  and  $\bar{B}$  consists of complete sequences  $\Upsilon_{\ell 1}\Upsilon_{\ell 2} \dots \Upsilon_{\ell N}$ , thereby possibly increasing the  $\bar{f}$  distance by at most  $4R_2/z$ . By Lemma 13 and (6.2),  $\bar{f}(\bar{A}, \bar{B}) > \alpha(\beta - \frac{1}{\sqrt{N}}) - \frac{1}{R_1} - \frac{4R_2}{z} > \alpha(\frac{1}{8} - \frac{2}{\sqrt[4]{N}}) - \frac{1}{R_1} - \frac{4R_2}{z}$ .  $\square$

## 7. A GENERAL SUBSTITUTION STEP

In this section we describe a step in our iteration of substitutions in a very general framework. The substitution will have the following initial data:

- An alphabet  $\Sigma$  and a collection of words  $X \subset \Sigma^b$
- Equivalence relations  $\mathcal{P}$  and  $\mathcal{R}$  on  $X$  with  $\mathcal{R}$  refining  $\mathcal{P}$
- Groups of involutions  $G$  and  $H$  with distinguished generators
- A homomorphism  $\rho : H \rightarrow G$  that preserves the distinguished generators. We denote the range of  $\rho$  by  $G'$  and its kernel by  $H_0$  with cardinality  $|H_0| = 2^t$  for some  $t \in \mathbb{N}$ .
- A free  $G$  action on  $X/\mathcal{P}$  and a free  $H$  action on  $X/\mathcal{R}$  such that the  $H$  action is subordinate to the  $G$  action via  $\rho$ . In particular, for each  $k \in \mathbb{Z}^+$  the skew diagonal actions of  $G$  on  $(X/\mathcal{P})^k$  and  $H$  on  $(X/\mathcal{R})^k$  are defined.
- There are  $N$  different equivalence classes in  $X/\mathcal{P}$  denoted by  $[A_i]_{\mathcal{P}}$ ,  $i = 1, \dots, N$ , where  $N = 2^{\nu+N'}$  with  $N', \nu \in \mathbb{N}$ .
- Each equivalence class  $[A_i]_{\mathcal{P}}$  contains  $2^{4e}$  elements of  $X/\mathcal{R}$ , where  $e \in \mathbb{Z}^+$  with  $e \geq \max(2, t)$ . We subdivide these  $\mathcal{R}$  classes into tuples as follows.

Pick an arbitrary class  $[A_1]_{\mathcal{P}}$  and take a set  $\{[A_{1,1}]_{\mathcal{R}}, \dots, [A_{1,2^{4e-t}}]_{\mathcal{R}}\}$  of  $\mathcal{R}$  classes in  $[A_1]_{\mathcal{P}}$  that intersects each orbit of the  $H_0$  action exactly once. This yields the first tuple  $([A_{1,1}]_{\mathcal{R}}, \dots, [A_{1,2^{4e-t}}]_{\mathcal{R}})$ . Then we obtain  $2^t - 1$  further tuples as the images  $(h[A_{1,1}]_{\mathcal{R}}, \dots, h[A_{1,2^{4e-t}}]_{\mathcal{R}})$  for each  $h \in H_0 \setminus \{\text{id}\}$ . Hereby, we have divided the  $2^{4e}$  elements of  $X/\mathcal{R}$  in  $[A_1]_{\mathcal{P}}$  into  $2^t$  many tuples such that each tuple intersects each orbit of the  $H_0$  action exactly once and the tuples are images of each other under the action by  $H_0$ . In the next step, we choose one element  $h \in H$  in each of the  $H/H_0$  cosets and apply each such  $h$  to the tuples in  $[A_1]_{\mathcal{P}}$  to obtain tuples in  $\rho(h)[A_1]_{\mathcal{P}}$ . If there are any  $[A_j]_{\mathcal{P}}$ 's that are not in the  $G'$  orbit of  $[A_1]_{\mathcal{P}}$ , then we repeat the procedure for such a  $[A_j]_{\mathcal{P}}$ , etc. Our procedure guarantees that the choices of elements in the  $H_0$  orbits are consistent between different  $[A_i]_{\mathcal{P}}$  equivalence classes, so that the action of  $H$  sends tuples to tuples. Let the  $2^t$  tuples in  $[A_i]_{\mathcal{P}}$  be written as

$$([A_{i,u2^{4e-t}+1}]_{\mathcal{R}}, \dots, [A_{i,(u+1)2^{4e-t}}]_{\mathcal{R}}),$$

where  $u \in \{0, \dots, 2^t - 1\}$ .

- For some  $R \geq 2$  and some  $\alpha \in (0, \frac{1}{8})$  we have

$$(7.1) \quad \bar{f}(A, \bar{A}) \geq \alpha$$

for any substantial substrings  $A$  and  $\bar{A}$  of at least  $\frac{h}{R}$  consecutive symbols in any representatives of two different  $\mathcal{P}$ -equivalence classes, that is, representatives of  $[A_{i_1, j_1}]_{\mathcal{R}}$  and  $[A_{i_2, j_2}]_{\mathcal{R}}$  for  $i_1 \neq i_2$  and any  $j_1, j_2 \in \{1, \dots, 2^{4e}\}$ .

- For some  $\beta \in (0, \alpha]$  we have

$$(7.2) \quad \bar{f}(A, \bar{A}) \geq \beta$$

for any substantial substrings  $A$  and  $\bar{A}$  of at least  $\frac{h}{R}$  consecutive symbols in any representatives of two different  $\mathcal{R}$ -equivalence classes, that is, representatives of  $[A_{i, j_1}]_{\mathcal{R}}$  and  $[A_{i, j_2}]_{\mathcal{R}}$ , respectively, for  $j_1 \neq j_2$ .

Let  $K, T_2 \in \mathbb{Z}^+$  and  $\tilde{R} \geq 2$  be given. The number  $\tilde{R}$  will be used in Proposition 43 to quantify substantial  $\tilde{R}$  substrings of newly constructed concatenations. Then in Section 8 we specify the size of  $\tilde{R}$  needed for the application of Proposition 43. We also suppose that there are numbers  $M_1, P, U_1 \in \mathbb{Z}^+$ , where  $\nu \cdot (2M_1 + 3) \geq \nu + N'$  and  $U_1$  is a multiple of  $2^t$  such that

$$(7.3) \quad U_1 \geq 2\tilde{R}^2.$$

Finally, let  $M_2 \in \mathbb{Z}^+$  with

$$(7.4) \quad M_2 \geq K \cdot P \cdot U_1 \cdot 2^{\nu \cdot (2M_1 + 3)}.$$

Hereby, we define the numbers

$$(7.5) \quad T_1 := T_2 \cdot 2^{(4e-t) \cdot (2M_2 + 3)},$$

$$(7.6) \quad U_2 := U_1 \cdot 2^{\nu \cdot (2M_1 + 3)},$$

and

$$(7.7) \quad k = U_1 \cdot T_1 \cdot 2^{\nu \cdot (2M_1 + 3)}.$$

Note that  $k$  is a multiple of  $N$  by our assumption  $\nu \cdot (2M_1 + 3) \geq \nu + N'$ .

Suppose we have a collection  $\Omega \subset (X/\mathcal{P})^k$  of cardinality  $|\Omega| = P$  that satisfies the following properties:

- $\Omega$  is closed under the skew diagonal action of  $G$ ,
- each  $\omega \in \Omega$  is a different concatenation of  $U_1$  many different  $(T_1, 2^\nu, M_1)$ -Feldman patterns as described in Section 6, each of which is constructed out of a tuple consisting of  $2^\nu$  many  $[A_i]_{\mathcal{P}}$ ,
- each  $[A_i]_{\mathcal{P}} \in X/\mathcal{P}$  occurs exactly  $\frac{k}{N}$  times in each  $\omega \in \Omega$ .

Then we construct a collection  $S \subset (X/\mathcal{R})^k$  of substitution instances of  $\Omega$  as follows.

- (1) We start by choosing a set  $\Upsilon \subset \Omega$  that intersects each orbit of the action by the group  $G'$  exactly once.
- (2) We construct a collection of  $M_2$  many different  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns, where the tuple of building blocks is to be determined in step (6). Note that each such pattern is constructed as a concatenation of  $T_2 2^{(4e-t) \cdot (2M_2+3)}$  many building blocks in total which motivates the definition of the number  $T_1$  from above.
- (3) By assumption on  $\Omega$  we can subdivide each element  $r \in \Upsilon$  as a concatenation of  $U_1 2^{\nu \cdot (2M_1+3)} = U_2$  strings of the form  $[A_i]_{\mathcal{P}}^{T_1}$  and each  $i \in \{1, \dots, N\}$  occurs exactly  $V := \frac{1}{N} U_2$  many times in this decomposition.
- (4) For each  $r \in \Upsilon$  we choose  $K$  different sequences of  $U_2$  concatenations of different  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns in ascending order from our collection in the second step and enumerate the sequences by  $j \in \{1, \dots, K\}$ . We require that each of the constructed patterns appears at most once, that is, we needed to construct at least  $|\Upsilon| \cdot K \cdot U_2$  many different patterns in that step. This explains the condition on the number  $M_2$  in equation (7.4).
- (5) We define a sequence  $\psi = (\psi_1, \dots, \psi_V)$  with  $V = \frac{1}{N} U_1 2^{\nu \cdot (2M_1+3)}$ , by  $\psi_v = u \in \{0, \dots, 2^t - 1\}$ , where  $u \equiv v \pmod{2^t}$ . That is, the sequence  $\psi$  cycles through the symbols in  $\{0, \dots, 2^t - 1\}$ . Since  $U_1$  was chosen as a multiple of  $2^t$  and  $\frac{1}{N} 2^{\nu \cdot (2M_1+3)} \in \mathbb{Z}$  by the assumption  $\nu \cdot (2M_1 + 3) \geq \nu + N'$ , each symbol from  $\{0, \dots, 2^t - 1\}$  occurs the same number  $\frac{V}{2^t}$  of times in the sequence  $\psi$ .
- (6) Let  $r \in \Upsilon$  and  $j \in \{1, \dots, K\}$ , and write

$$r = [A_{i(1)}]_{\mathcal{P}}^{T_1} \cdots [A_{i(\ell)}]_{\mathcal{P}}^{T_1} \cdots [A_{i(U_2)}]_{\mathcal{P}}^{T_1},$$

where  $i(1), \dots, i(U_2) \in \{1, \dots, N\}$ . If  $i(\ell) = i_0$  and this is the  $m$ th occurrence of  $i_0$  in the sequence  $i(1), \dots, i(\ell)$  then we let  $u = \psi_m$  and substitute a Feldman pattern built with the tuple

$$([A_{i_0, u 2^{4e-t+1}}]_{\mathcal{R}}, \dots, [A_{i_0, (u+1) 2^{4e-t}}]_{\mathcal{R}})$$

into  $[A_{i_0}]_{\mathcal{P}}^{T_1}$ . The Feldman pattern that is used is the  $\ell$ th pattern among the  $U_2$  patterns previously chosen for the given  $r$  and  $j$ . We follow this procedure for each  $\ell = 1, \dots, U_2$  to obtain an element  $s \in (X/\mathcal{R})^k$ . Let  $S$  be the collection of such  $s \in (X/\mathcal{R})^k$  obtained for all  $r \in \Upsilon$  and  $j \in \{1, \dots, K\}$ .

Using this collection  $S \subset (X/\mathcal{R})^k$  we define

$$\Omega' = HS.$$

**Proposition 42.** *This collection  $\Omega' \subset (X/\mathcal{R})^k$  satisfies the following properties.*




FIGURE 7.1. Visualization of the substitution step.

- (1)  $\Omega'$  is closed under the skew diagonal action by  $H$ .
- (2) For each element in  $\omega \in \Omega$  there are  $K \cdot |H_0|$  many substitution instances in  $\Omega'$ .
- (3) Each element of  $\Omega'$  contains each  $[A_{i,j}]_{\mathcal{R}}$  the same number of times.

*Proof.* The first statement follows from the definition of  $\Omega'$  immediately. For part (2) we recall that  $\Upsilon \subset \Omega$  was chosen to intersect each orbit of the action by the group  $G'$  exactly once and that for each element in  $\Upsilon$  there are  $K$  many substitution instances in  $S$ . To see the third part we use that all the  $[A_i]_{\mathcal{P}} \in X/\mathcal{P}$  occurred uniformly in each  $\omega \in \Omega$ . By construction of the sequence  $\psi$  in step (5) each tuple  $([A_{i,a+1}]_{\mathcal{R}}, \dots, [A_{i,a+2^{4e-t}}]_{\mathcal{R}})$  occurs the same number of times as building blocks for  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns substituted into strings  $[A_i]_{\mathcal{P}}^{T_1}$ . Since Feldman patterns are uniform in their building blocks, we conclude the third statement.  $\square$

In the following, we estimate the  $\bar{f}$  distance of elements in  $\Omega'$  that are equivalent with respect to the  $\mathcal{P}$  product equivalence relation but are not  $\mathcal{R}$ -equivalent. We conclude that they are still approximately  $\beta$  apart from each other in  $\bar{f}$ , where  $\beta$  is as in (7.2). The  $\bar{f}$  distance for strings of symbols in  $\Sigma_{\mathcal{R}}$  will be denoted  $\bar{f}_{\Sigma_{\mathcal{R}}}$ . When there is no subscript on the  $\bar{f}$ , it is understood that the alphabet is  $\Sigma$ .

**Proposition 43.** *Let  $\omega'_1$  and  $\omega'_2$  be two different substitution instances in  $\Omega'$  of an element  $\omega \in \Omega$ . We consider  $\omega'_1, \omega'_2 \in \Omega' \subset (X/\mathcal{R})^k$  as strings of length  $k$  in the alphabet  $\Sigma_{\mathcal{R}} := \{[A_{i,j}]_{\mathcal{R}} : 1 \leq i \leq N, 1 \leq j \leq 2^{4e}\}$ . Then we have*

$$(7.8) \quad \bar{f}_{\Sigma_{\mathcal{R}}}(W', \bar{W}') > 1 - \frac{4}{2^e} - \frac{1}{\bar{R}}$$

for any substrings  $W'$  and  $\bar{W}'$  of at least  $k/\bar{R}$  consecutive  $\Sigma_{\mathcal{R}}$ -symbols from  $\omega'_1$  and  $\omega'_2$ , respectively.

Moreover, we have

$$(7.9) \quad \bar{f}(V, \bar{V}) > \beta - \frac{1}{\bar{R}} - \frac{4}{2^e} - \frac{1}{\bar{R}}$$

for any substrings  $V, \bar{V}$  of at least  $k\mathfrak{h}/\tilde{R}$  consecutive symbols in any representatives  $W_1$  and  $W_2$  of  $\omega'_1$  and  $\omega'_2$ , respectively, in the alphabet  $\Sigma$ .

*Proof.* Let  $W'$  and  $\bar{W}'$  be any substrings of at least  $k/\tilde{R}$  consecutive  $\Sigma_{\mathcal{R}}$ -symbols from  $\omega'_1$  and  $\omega'_2$ , respectively. We subdivide both strings into sequences coming from the  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns and we recall that their length is  $T_2 2^{(4e-t) \cdot (2M_2+3)} = T_1$ . By adding fewer than  $2T_1$  many  $\Sigma_{\mathcal{R}}$ -symbols to each of  $W'$  and  $\bar{W}'$  we can complete partial patterns at the beginning and end. Let  $W'_{aug}$  and  $\bar{W}'_{aug}$  be the augmented  $W'$  and  $\bar{W}'$  strings obtained in this way. By Fact 9 and condition (7.3) we have

$$(7.10) \quad \bar{f}_{\Sigma_{\mathcal{R}}}(W', \bar{W}') > \bar{f}_{\Sigma_{\mathcal{R}}}(W'_{aug}, \bar{W}'_{aug}) - \frac{2\tilde{R}}{U_1 2^{v \cdot (2M_1+3)}} > \bar{f}_{\Sigma_{\mathcal{R}}}(W'_{aug}, \bar{W}'_{aug}) - \frac{1}{\tilde{R}}.$$

We write  $W'_{aug}$  and  $\bar{W}'_{aug}$  as

$$W'_{aug} = P_1^{(p_1, a_1, k_1)} \dots P_s^{(p_s, a_s, k_s)}, \quad \bar{W}'_{aug} = \bar{P}_1^{(q_1, b_1, l_1)} \dots \bar{P}_t^{(q_t, b_t, l_t)},$$

where  $P_i^{(p_i, a_i, k_i)}$ ,  $1 \leq i \leq s$ , as well as  $\bar{P}_j^{(q_j, b_j, l_j)}$ ,  $1 \leq j \leq t$ , are complete  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns in the  $\Sigma_{\mathcal{R}}$ -alphabet and the superscripts  $p_i, q_j \in \{1, \dots, M_2\}$  indicate which pattern structure is used. Moreover, the other superscripts identify the building blocks. Here,  $a_i, b_j \in \{1, \dots, N\}$  refer to the  $\mathcal{P}$ -equivalence class of the building blocks and  $k_i, l_j \in \{1, \dots, 2^{4e-t}\}$  refer to the tuple of  $\mathcal{R}$ -equivalence classes of building blocks. If we apply Lemma 39 with  $S = T = T_2$  and  $N$  replaced by  $2^{4e-t}$ , we obtain

$$(7.11) \quad \bar{f}_{\Sigma_{\mathcal{R}}}(P, \bar{P}) \begin{cases} = 1, & \text{if } (a_i, k_i) \neq (b_j, l_j) \\ > 1 - \frac{4}{2^{2e-0.5t}}, & \text{if } (a_i, k_i) = (b_j, l_j) \text{ and } p_i \neq q_j \end{cases}$$

for any sequences  $P$  and  $\bar{P}$  of at least  $\frac{T_1}{2^{4e-t}} = T_2 2^{(4e-t) \cdot (2M_2+2)}$  consecutive  $\Sigma_{\mathcal{R}}$ -symbols in  $P_i^{(p_i, a_i, k_i)}$  and  $\bar{P}_j^{(q_j, b_j, l_j)}$ , respectively.

We examine the two possible cases:  $\omega'_1$  and  $\omega'_2$  have disjoint  $H$  orbits (case 1) or  $\omega'_1 = h\omega'_2$  for some  $h \in H_0$  (case 2). From the construction we recall that in the first case all occurring  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns in  $\omega'_1$  and  $\omega'_2$  are different from each other. To investigate the second case we recall that a tuple of building blocks is mapped to a different tuple under the action of an element  $h \in H_0$ . Hence, identical Feldman patterns in  $\omega'_1 = h\omega'_2$  and  $\omega'_2$  have different tuples of building blocks.

In both cases we consider a best possible  $\bar{f}$  match between  $W'_{aug}$  and  $\bar{W}'_{aug}$ . To simplify notation we ignore the superscripts of the occurring  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns. We denote the string in  $\bar{W}'_{aug}$  matched with  $P_i$  by  $\tilde{P}_i$ . If  $|\tilde{P}_i| < \frac{T_1}{2^e}$  or  $|\tilde{P}_i| > 2^e T_1$ , then

$$(7.12) \quad \bar{f}_{\Sigma_{\mathcal{R}}}(P_i, \tilde{P}_i) > 1 - \frac{2}{2^e}.$$

Otherwise, we subdivide  $\tilde{P}_i$  into the strings coming from the Feldman patterns in  $\bar{W}'_{aug}$  and we denote the part of  $\tilde{P}_i$  belonging to  $\bar{P}_j$  by  $\tilde{P}_{i,j}$ . Note that there are at most  $2^e + 2$  many  $j$ 's with nonempty  $\tilde{P}_{i,j}$  due to  $|\tilde{P}_i| \leq 2^e T_1$ . In the following we ignore all  $\tilde{P}_{i,j}$  of length  $|\tilde{P}_{i,j}| < \frac{T_1}{2^{2e}}$ . Hereby, we omit at most  $\frac{2T_1}{2^{2e}}$  many symbols

from  $\tilde{P}_i$ . We denote the remaining  $\tilde{P}_i$  string by  $\tilde{P}_{i,red}$  and we obtain from Fact 9 that

$$(7.13) \quad \bar{f}_{\Sigma\mathcal{R}}(P_i, \tilde{P}_i) > \bar{f}_{\Sigma\mathcal{R}}(P_i, \tilde{P}_{i,red}) - \frac{2}{2^{2e}}.$$

For all the remaining  $j$ 's with  $|\tilde{P}_{i,j}| \geq \frac{T_1}{2^{2e}}$  the corresponding string in  $P_i$  under the matching is called  $P_{i,j}$ . If  $|P_{i,j}| < \frac{T_1}{2^{4e-t}}$ , then

$$\bar{f}_{\Sigma\mathcal{R}}(P_{i,j}, \tilde{P}_{i,j}) > 1 - \frac{2|P_{i,j}|}{|\tilde{P}_{i,j}| + |P_{i,j}|} = 1 - \frac{2}{\frac{|\tilde{P}_{i,j}|}{|P_{i,j}|} + 1} > 1 - \frac{2}{2^e}.$$

Otherwise, we use the estimates from formula (7.11). This yields

$$\bar{f}_{\Sigma\mathcal{R}}(P_i, \tilde{P}_i) > 1 - \frac{2}{2^e} - \frac{2}{2^{2e}} \geq 1 - \frac{3}{2^e}$$

by equation (7.13). Combining this with equation (7.12), we obtain

$$(7.14) \quad \bar{f}_{\Sigma\mathcal{R}}(W'_{aug}, \bar{W}'_{aug}) > 1 - \frac{4}{2^e}.$$

With the aid of equation (7.10) we finish the proof of (7.8).

Now we turn to the proof of (7.9) in the alphabet  $\Sigma$ . We treat  $V_{aug}$  and  $\bar{V}_{aug}$  as concatenations of complete  $(T_2, 2^{4e-t}, M_2)$ -Feldman patterns by adding fewer than  $2T_1h$  many  $\Sigma$ -symbols to complete partial patterns at the beginning and end of  $V$  and  $\bar{V}$ . Note that  $V_{aug}$  and  $\bar{V}_{aug}$  correspond to some  $W'_{aug}$  and  $\bar{W}'_{aug}$  in the  $\Sigma\mathcal{R}$ -alphabet of the same type as above. By Lemma 13, with  $\bar{f} \geq 1 - \frac{4}{2^e}$  coming from equation (7.14), and our assumptions from equations (7.1) and (7.2), we obtain

$$\bar{f}(V_{aug}, \bar{V}_{aug}) > \left(1 - \frac{4}{2^e}\right) \cdot \beta - \frac{1}{R} > \beta - \frac{1}{R} - \frac{4}{2^e}.$$

Finally, we conclude (7.9) with the aid of Fact 9 and equation (7.3).  $\square$

## 8. THE CONSTRUCTION PROCESS

To construct the map  $\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$  we must build a construction sequence  $(\mathcal{W}_n(\mathcal{T}))_{n \in \mathbb{N}}$  satisfying our specifications for each  $\mathcal{T} \in \mathcal{Trees}$ . Moreover, this has to be done in such a way that  $\mathcal{W}_n(\mathcal{T})$  is entirely determined by  $\mathcal{T} \cap \{\sigma_m : m \leq n\}$ , that is,

$$(8.1) \quad \begin{aligned} &\text{if } \mathcal{T} \cap \{\sigma_m : m \leq n\} = \mathcal{T}' \cap \{\sigma_m : m \leq n\}, \\ &\text{then for all } m \leq n : \mathcal{W}_m(\mathcal{T}) = \mathcal{W}_m(\mathcal{T}'), \end{aligned}$$

and we guarantee the continuity of our map  $\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$  by Fact 17. Therefore, we follow [FRW11] to organize our construction: For each  $n$  and for each subtree  $\mathcal{S} \subseteq \{\sigma_m : m \leq n\}$  and each  $\sigma_m \in \mathcal{S}$  we build  $\mathcal{W}_m(\mathcal{S})$ . By induction we want to pass from stage  $n-1$  to stage  $n$ . So, we assume that we have constructed  $(\mathcal{W}_m(\mathcal{S}))_{\sigma_m \in \mathcal{S}}$ ,  $(\mathcal{Q}_m(\mathcal{S}))_{\sigma_m \in \mathcal{S}}$ , and the action of  $(G_s^m(\mathcal{S}))_{\sigma_m \in \mathcal{S}}$  for each subtree  $\mathcal{S} \subseteq \{\sigma_m : m \leq n-1\}$ . In the inductive step, we now have to construct  $(\mathcal{W}_m(\mathcal{S}))_{\sigma_m \in \mathcal{S}}$  for each subtree  $\mathcal{S} \subseteq \{\sigma_m : m \leq n\}$ .

Obviously, for those subtrees  $\mathcal{S}$  with  $\sigma_n \notin \mathcal{S}$  there is nothing to do. So, we have to work on the finitely many subtrees  $\mathcal{S}$  with  $\sigma_n \in \mathcal{S}$ . We list those in arbitrary manner as  $\{\mathcal{S}_1, \dots, \mathcal{S}_E\}$ . Assume that  $\mathcal{T}$  is the  $e$ -th subtree on this list and that we have constructed the collections  $\mathcal{W}_n(\mathcal{S}_i)$  of  $n$ -words for all  $i \in \{1, \dots, e-1\}$ . Let

$\mathfrak{P}$  be the collection of prime numbers occurring in the prime factorization of any of the lengths of the words that we have constructed so far.<sup>1</sup>

We want to construct  $\mathcal{W}_n(\mathcal{T})$ ,  $\mathcal{Q}_s^n(\mathcal{T})$ , and the action of  $G_s^n(\mathcal{T})$ , which we abbreviate by  $\mathcal{W}_n$ ,  $\mathcal{Q}_s^n$ , and  $G_s^n$ , respectively. For that purpose, let  $m$  be the largest number less than  $n$  such that  $\sigma_m \in \mathcal{T}$ . If  $m = 0$  we have  $\mathcal{W}_0 = \Sigma = \{1, \dots, 2^{12}\}$  (see Remark 54 for comments on this base case) and for  $m > 0$  our induction assumption says that we have  $\mathcal{W}_m = \mathcal{W}_m(\mathcal{T})$ ,  $\mathcal{Q}_s^m = \mathcal{Q}_s^m(\mathcal{T})$ , and  $G_s^m = G_s^m(\mathcal{T})$  satisfying our specifications. Furthermore, there are  $\frac{1}{8} > \alpha_{1,m} > \dots > \alpha_{s(m),m} > \beta_m > 0$  and a positive integer  $R_m$  with

$$(8.2) \quad R_m \geq \frac{9}{\beta_m}$$

such that the following assumptions on  $\bar{f}$  distances hold:

- For every  $s \in \{1, \dots, s(m)\}$  we have

$$(8.3) \quad \bar{f}(W, W') > \alpha_{s,m}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{h_m}{R_m}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_m$  with  $[w]_s \neq [w']_s$ .

- We have

$$(8.4) \quad \bar{f}(W, W') > \beta_m$$

on any substrings  $W$  and  $W'$  of at least  $\frac{h_m}{R_m}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_m$  with  $w \neq w'$ .

We start the construction in stage  $n$  by choosing a prime number  $p_n$  larger than the maximum of  $\mathfrak{P}$  and satisfying

$$(8.5) \quad p_n > \max\left(4R_m, 2^n, \frac{1}{\epsilon_n}\right),$$

where we recall the definition of the sequence  $(\epsilon_n)_{n \in \mathbb{N}}$  from (4.1). Then we choose an integer  $R_n$  such that

$$(8.6) \quad R_n \geq \frac{40p_n}{\beta_m}$$

Moreover, we choose the integer  $e(n) > e(m)$  sufficiently large such that

$$(8.7) \quad 2^{e(n)} > \max\left(10R_n, \max_{s \leq s(n)} |G_s^m|\right).$$

In the two possible cases  $s(n) = s(m)$  and  $s(n) = s(m) + 1$  we now present the construction of  $n$ -words out of  $m$ -words as well as the extension of  $G_s^m$  actions to  $G_s^n$  actions. In case of  $s(n) = s(m) + 1$  we also define the new equivalence relation  $\mathcal{Q}_{s(n)}^n$ .

### 8.1. Case 1: $s(n)=s(m)$ .

---

<sup>1</sup>In our construction of  $\mathcal{W}_n(\mathcal{T})$  we will choose a prime number  $p_n$  larger than the maximum of  $\mathfrak{P}$ . By Lemma 28 we see that our map  $\Psi : \mathcal{Trees} \rightarrow \mathcal{E}$  takes distinct trees to transformations with nonisomorphic Kronecker factors. Hence, for  $\mathcal{T} \neq \mathcal{T}'$  we get that  $\Psi(\mathcal{T})$  is not isomorphic to  $\Psi(\mathcal{T}')$  or  $\Psi(\mathcal{T}')^{-1}$  (compare with [FRW11, Corollary 34]). As in [FRW11] this is actually not necessary for the proof of the main theorem but we keep this order of induction to parallel the strategy in [FRW11] and to allow possible extensions in future work.

8.1.1. *Construction of  $\mathcal{W}_n$ .* We apply the substitution step described in Section 7  $s(m) + 1$  many times to construct  $\mathcal{W}_n/\mathcal{Q}_s^n$  for  $s = 0, \dots, s(m)$  and eventually  $\mathcal{W}_n$ . Adapting to the notation in the general substitution step we set  $X = \mathcal{W}_m$  and

$$P_i = \mathcal{Q}_{i-1}^m, \mathcal{R}_i = \mathcal{Q}_i^m, \quad G_i = G_{i-1}^m, H_i = G_i^m$$

for the  $i$ -th application, where  $i = 1, \dots, s(m)$ . In the final  $(s(m) + 1)$ -th application we let  $\mathcal{P}_{s(m)+1} = \mathcal{Q}_{s(m)}^m$ ,  $\mathcal{R}_{s(m)+1} = \{(w, w) : w \in \mathcal{W}_m\}$ , that is,  $X/\mathcal{R}_{s(m)+1} = \mathcal{W}_m$ ,  $G_{s(m)+1} = G_{s(m)}^m$ , and we let  $H_{s(m)+1}$  be the trivial group acting trivially on  $\mathcal{W}_m$ .

To carry out the successive application of the substitution step we always set  $\tilde{R} = R_n$  and introduce  $t_i \in \mathbb{N}$  for  $i = 1, \dots, s(m) + 1$  such that  $t_{s(m)+1} = 0$  and

$$2^{t_i} = |\ker(\rho_{i,i-1}^{(m)})| \text{ for } i = 1, \dots, s(m).$$

Here, we recall that  $\rho_{i,i-1}^{(m)} : G_i^m \rightarrow G_{i-1}^m$  is the canonical homomorphism defined in Subsection 4.3. Moreover, we define parameters as follows

$$P_i = 2^{(i-1) \cdot 4e(n)}, \quad \text{for } i = 1, \dots, s(m) + 1, \\ N_i = 2^{(i-1) \cdot 4e(m)}, \quad \text{for } i = 1, \dots, s(m) + 1.$$

Then we have  $N_i = |\mathcal{W}_m/\mathcal{Q}_{i-1}^m|$  and the number  $P_i$  will describe  $|\mathcal{W}_n/\mathcal{Q}_{i-1}^n|$ . We also set  $K_{s(m)+1} = 2^{4e(n)}$ , and choose numbers  $K_i$ ,  $i = 1, \dots, s(m)$ , such that  $K_i \cdot |\ker(\rho_{i,i-1}^{(m)})| = 2^{4e(n)}$  for  $i = 1, \dots, s(m)$ . We point out that these numbers are chosen in this way to always obtain exactly  $2^{4e(n)}$  substitution instances of each coarser equivalence class (see part (2) of Proposition 42) in order to satisfy specification property (Q6). Moreover, we choose a number  $U_1 = 2^{U'_1}$  with

$$U'_1 \geq \max_{i=1, \dots, s(m)} t_i \quad \text{and} \quad U_1 \geq \max \left( 2R_n^2, 2^{8e(m)} \right).$$

In particular, condition (7.3) is satisfied in all construction steps.

By forward recursion we define the parameters  $M_i$ ,  $i = 1, \dots, s(m) + 2$ , as well as  $U_i$ ,  $i = 2, \dots, s(m) + 1$ :

$$M_1 = 1, \quad M_2 = K_1 P_1 U_1, \quad M_i = K_{i-1} P_{i-1} U_{i-1} 2^{(4e(m)-t_{i-2}) \cdot (2M_{i-1}+3)} \text{ for } i \geq 3, \\ U_2 = U_1, \quad U_i = U_{i-1} 2^{(4e(m)-t_{i-2}) \cdot (2M_{i-1}+3)} \text{ for } i \geq 3.$$

Hereby, we define the parameters  $T_i$ ,  $i = 1, \dots, s(m) + 2$ , by backward recursion:

$$T_{s(m)+2} = p_n^2, \quad T_i = T_{i+1} \cdot 2^{(4e(m)-t_i) \cdot (2M_{i+1}+3)} \text{ for } i \leq s(m) + 1.$$

We stress that the definitions of  $T_i$  and  $M_i$  are directly linked to the corresponding conditions in equations (7.4) and (7.5) from the general substitution step. We also conclude from equation (7.7) that the number  $k$  of concatenations working for all substitutions steps is given by

$$(8.8) \quad k = U_{s(m)+1} \cdot T_{s(m)+1} \cdot 2^{(4e(m)-t_{s(m)}) \cdot (2M_{s(m)+1}+3)} = U_{s(m)+1} \cdot T_{s(m)} = U_2 \cdot T_1,$$

which implies that the length of  $n$ -words is given by

$$h_n = U_{s(m)+1} \cdot T_{s(m)+1} \cdot 2^{(4e(m)-t_{s(m)}) \cdot (2M_{s(m)+1}+3)} \cdot h_m = U_1 \cdot T_1 \cdot h_m.$$

After determining all these parameters we describe the successive application of the general substitution step. We start by concatenating  $k = U_1 \cdot T_1$  many times the single element of  $X/\mathcal{P}_1 = \mathcal{W}_m/\mathcal{Q}_0^m$  which yields  $\Omega_1 = \mathcal{W}_n/\mathcal{Q}_0^n \subseteq (\mathcal{W}_m/\mathcal{Q}_0^m)^k = (X/\mathcal{Q}_0)^k$  of cardinality  $|\Omega_1| = P_1 = 1$ . Note that this element can be considered



as a repetition of  $U_1$  many  $(T_1, 2^0, 1)$ -Feldman patterns. This viewpoint allows the application of our first substitution step  $i = 1$  using the notation from above. More explicitly, the first substitution step produces a collection  $\Omega'_1 \subset (X/\mathcal{R}_1)^k = (\mathcal{W}_m/\mathcal{Q}_1^m)^k$  of  $K_1 \cdot |\ker(\rho_{1,0}^{(m)})| = 2^{4e(n)} = P_2$  substitution instances, which are concatenations of  $U_2$  many  $(T_2, 2^{4e(m)-t_1}, M_2)$ -Feldman patterns. On the one hand,  $\Omega'_1$  will serve as our collection  $\Omega_2 \subset (X/\mathcal{P}_2)^k = (\mathcal{W}_m/\mathcal{Q}_1^m)^k$  for the application of the substitution step  $i = 2$ . On the other hand,  $\Omega'_1$  will turn out to be our collection of equivalence classes in  $\mathcal{W}_n/\mathcal{Q}_1^n$ . Successively, we apply the substitution step for  $i = 2, \dots, s(m) + 1$ .

*Remark 44.* In correspondence with specification (Q5),  $\mathcal{Q}_s^n$  is defined as the product equivalence relation on  $\mathcal{W}_n$  for each  $s = 1, \dots, s(m)$ . Then we see from the procedure of the substitution step that for each  $w \in \mathcal{W}_n$  its class  $[w]_s \in \mathcal{W}_n/\mathcal{Q}_s^n$  is a concatenation of  $U_{s+1} \geq 2^{8e(m)}$  different  $(T_{s+1}, 2^{4e(m)-t_s}, M_{s+1})$ -Feldman patterns (with building blocks out of  $\mathcal{W}_m/\mathcal{Q}_s^m$ ). We call these the *s-Feldman patterns* and refer to this sequence of Feldman patterns as the *s-pattern type* of  $w$ . Using the same enumeration of  $(T_{s+1}, 2^{4e(m)-t_s}, M_{s+1})$ -Feldman patterns, we also define the *s-pattern types* for words in  $\text{rev}(\mathcal{W}_n)$ .

For such a *s-Feldman pattern*  $P_m$  we make the following observation.

**Lemma 45.** *Let  $w \in \mathcal{W}_n$  and  $[P_m]_s$  be a  $(T_{s+1}, 2^{4e(m)-t_s}, M_{s+1})$ -Feldman pattern in  $[w]_s$  with building blocks from the tuple  $([A_1]_s, \dots, [A_{2^{4e(m)-t_s}}]_s)$  in  $\mathcal{W}_m/\mathcal{Q}_s^m$ . Then for every  $i = 1, \dots, 2^{4e(m)-t_s}$  the repetitions  $(a_{i,j})^{p_n^2}$  of each substitution instance  $a_{i,j}$  in  $\mathcal{W}_m$  of  $[A_i]_s$  are substituted the same number of times into each occurrence of  $[A_i]_s^{T_{s+1} \cdot 2^{t_{s+1}}}$  in  $[P_m]_s$ . This also yields that the repetitions  $(a_{i,j})^{p_n^2}$  of each substitution instance  $a_{i,j}$  of each  $[A_i]_s$ ,  $i = 1, \dots, 2^{4e(m)-t_s}$ , occur the same number of times in each cycle of  $P_m$ .*

*Proof.* By definition of  $(T_{s+1}, 2^{4e(m)-t_s}, M_{s+1})$ -Feldman patterns we have in each complete cycle that for every  $i \in \{1, \dots, 2^{4e(m)-t_s}\}$  the maximum repetition

$$[A_i]_s^{T_{s+1} \cdot 2^{(4e(m)-t_s) \cdot 2j}}$$

occurs exactly once, where  $j \in \{1, \dots, M_{s+1}\}$  labels our Feldman pattern. Into each  $[A_i]_s^{T_{s+1}}$  we substitute a  $(T_{s+2}, 2^{4e(m)-t_{s+1}}, M_{s+2})$ -Feldman pattern built out of one of the  $2^{t_{s+1}}$  many tuples  $([A_{i,1}]_{s+1}, \dots, [A_{i,2^{4e(m)-t_{s+1}}}]_{s+1})$ . According to our choice of the tuples in step (5) of the general substitution step each of these tuples is used  $2^{(4e(m)-t_s) \cdot 2j - t_{s+1}}$  many times within one maximum repetition  $[A_i]_s^{T_{s+1} \cdot 2^{(4e(m)-t_s) \cdot 2j}}$ . Then we continue inductively for each of the obtained  $(T_{s+2}, 2^{4e(m)-t_{s+1}}, M_{s+2})$ -Feldman patterns until we finally insert repetitions of the form  $(a_{i,j})^{p_n^2}$  (recall that  $T_{s(m)+2} = p_n^2$ ).  $\square$

**8.1.2. Group actions.** We also need to define the group actions for  $G_s^n$  for  $s \leq s(n)$ . Except for  $s_0 = lh(\sigma_n)$  we have  $G_s^n = G_s^m$  and the group actions are extended via the skew-diagonal actions. In the remaining case we have  $G_{s_0}^n = G_{s_0}^m \oplus \mathbb{Z}_2$  and we use Lemma 34 to extend the  $G_{s_0}^m$  action to a  $G_{s_0}^n$  action subordinate to the  $G_{s_0-1}^n$  action. To be more precise, we set  $G \times \mathbb{Z}_2 := G_{s_0-1}^n = G_{s_0-1}^m$ ,  $H := G_{s_0}^m$ ,  $X/\mathcal{Q} := \mathcal{W}_n/\mathcal{Q}_{s_0-1}^n$ ,  $X/\mathcal{R} := \mathcal{W}_n/\mathcal{Q}_{s_0}^n$ , and  $\rho := \rho_{s_0, s_0-1}^{(m)}$  in the notation of the lemma.

8.1.3. *Properties.* By construction the specifications are satisfied. In particular, specification (E3) holds since we obtain words by concatenating different Feldman patterns.

We also make the following observations regarding  $\bar{f}$  distances.

**Lemma 46.** *For every  $s \in \{1, \dots, s(m)\}$  we have*

$$(8.9) \quad \bar{f}(W, W') > \alpha_{s,m} - \frac{2}{R_m} - \frac{1}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{h_n}{R_n}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_s \neq [w']_s$ .

Moreover, we have

$$(8.10) \quad \bar{f}(W, W') > \beta_m - \frac{2}{R_m} - \frac{1}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{h_n}{R_n}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_n$  with  $w \neq w'$ .

*Proof.* For each  $s \in \{1, \dots, s(m)\}$  we use Proposition 43 and assumption (8.3) to obtain the following estimate

$$\bar{f}(W, W') > \alpha_{s,m} - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{R_n} > \alpha_{s,m} - \frac{2}{R_m} - \frac{1}{R_n}$$

by  $2^{e(m)} > 10R_m$ , which follows from equation (8.7) on stage  $m$  of the induction. Similarly, we use Proposition 43 and assumption (8.4) to conclude equation (8.10).  $\square$

The expression on the right hand side in equation (8.9) will give us the definition

$$(8.11) \quad \alpha_{s,n} := \alpha_{s,m} - \frac{2}{R_m} - \frac{1}{R_n}$$

in this case of the construction. Similarly, equation (8.10) yields the definition

$$\beta_n := \beta_m - \frac{2}{R_m} - \frac{1}{R_n}.$$

We point out that

$$\beta_n > \frac{\beta_m}{2} > \frac{10}{R_n} > 0$$

by the conditions (8.2) and (8.6). This also implies  $R_n > \frac{10}{\beta_n}$ , that is, the induction assumption (8.2) for the next step is satisfied.

*Remark 47.* To prepare the transfer to the setting of diffeomorphisms, we consider the constructed words in  $\mathcal{W}_n$  as concatenations of  $k_m$  symbols in the alphabets  $\Sigma_{m,s}$  for  $s \leq s(m)$ . Here,  $\Sigma_{m,s}$  is the alphabet where we identify each class in  $\mathcal{W}_m/Q_s^m$  of length  $h_m$  with one symbol in  $\{1, \dots, Q_s^{M(s)}\}$ . As in the proof of Lemma 46 we use Proposition 43 to obtain

$$\bar{f}_{\Sigma_{m,s}}(W, W') > 1 - \frac{4}{2^{e(m)}} - \frac{1}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{k_m}{R_n}$  consecutive  $\Sigma_{m,s}$ -symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_s \neq [w']_s$ . Similarly, we consider words in  $\mathcal{W}_n$  as concatenations

of  $k_m$  symbols in the alphabets  $\Sigma_m$ , that is, each  $m$ -word represents one symbol. As above, we get

$$\bar{f}_{\Sigma_m}(W, W') > 1 - \frac{4}{2^{e(m)}} - \frac{1}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{k_m}{R_n}$  consecutive  $\Sigma_m$ -symbols in any words  $w, w' \in \mathcal{W}_n$  with  $w \neq w'$ .

**8.2. Case 2:  $\mathbf{s(n)=s(m)+1}$ .** In this case  $\sigma_n$  is the only sequence in  $\mathcal{T} \cap \{\sigma_k : k \leq n\}$  of length  $s(n)$ . Hence, we have  $G_{s(n)}^n = \mathbb{Z}_2$  and we additionally need to define  $\mathcal{Q}_{s(n)}^n$  as well as the action of  $\mathbb{Z}_2$  on  $\mathcal{W}_n / \mathcal{Q}_{s(n)}^n$ .

Recall that the specification (Q4) on  $\mathcal{Q}_{s(n)}^n$  is given in terms of a number  $J_{s(n),n}$ . We start the construction by choosing the number  $J_{s(n),n} = 2^{J'_{s(n),n}}$  sufficiently large such that

$$(8.12) \quad J_{s(n),n} > 2R_n^2.$$

This time the construction of  $\mathcal{W}_n$  uses two collections  $\mathcal{W}^\dagger$  and  $\mathcal{W}^{\dagger\dagger}$  of concatenation of  $m$ -words. Here, both collections  $\mathcal{W}^\dagger$  and  $\mathcal{W}^{\dagger\dagger}$  are obtained by a successive application of  $s(m) + 1$  substitution steps. Their construction parallels the one in Case 1 but with different choices of parameters.

This time we use  $\tilde{R} = 2J_{s(n),n}^3$  in each substitution step and we repeat the definition of parameters  $t_i$ ,  $i = 1, \dots, s(m) + 1$ , and

$$N_i = 2^{(i-1) \cdot 4e(m)}, \quad \text{for } i = 1, \dots, s(m) + 1.$$

Moreover, we choose a number  $U_1 = 2^{U'_1}$  with  $U'_1 \geq \max_{i=1, \dots, s(m)} t_i$  and

$$(8.13) \quad U_1 \geq \max \left( 8J_{s(n),n}^6, 2^{8e(m)} \right).$$

By forward recursion we define the parameters  $M_i$ ,  $i = 1, \dots, s(m) + 2$ , as well as  $U_i$ ,  $i = 2, \dots, s(m) + 1$ :

$$M_1 = 1, \quad M_2 = 2^{4e(n)+1}U_1, \quad M_i = 2^{(i+2) \cdot 4e(n)}U_{i-1}2^{(4e(m)-t_{i-2}) \cdot (2M_{i-1}+3)} \text{ for } i \geq 3,$$

$$U_2 = U_1, \quad U_i = U_{i-1}2^{(4e(m)-t_{i-2}) \cdot (2M_{i-1}+3)} \text{ for } i \geq 3.$$

Then for each  $i = 1, \dots, s(m) + 1$  we construct  $M_{i+1}$  many  $(T, 2^{4e(m)-t_i}, M_{i+1})$ -Feldman patterns, where the parameter  $T$  will be chosen later and will differ between the two cases  $\mathcal{W}^\dagger$  and  $\mathcal{W}^{\dagger\dagger}$ .

**8.2.1. Construction of  $\mathcal{W}^{\dagger\dagger}$ .** To construct the collection  $\mathcal{W}^{\dagger\dagger}$  we choose the remaining parameters as follows: We set  $K_i^{\dagger\dagger} = 1$  for  $i = 1, \dots, s(m)$  and

$$(8.14) \quad K_{s(m)+1}^{\dagger\dagger} = 2^{8e(n)}.$$

Moreover, we define the parameters  $T_i^{\dagger\dagger}$ ,  $i = 1, \dots, s(m) + 2$ , by backward recursion:

$$T_{s(m)+2}^{\dagger\dagger} = p_n, \quad T_i^{\dagger\dagger} = T_{i+1}^{\dagger\dagger} \cdot 2^{(4e(m)-t_i) \cdot (2M_{i+1}+3)} \text{ for } i \leq s(m) + 1.$$

Then we apply the general substitution step  $s(m) + 1$  many times as in Case 1, where  $P_1^{\dagger\dagger} = 1$ ,  $P_2^{\dagger\dagger} = |G_1^m|$ , and  $P_i^{\dagger\dagger} = |G_1^m| \cdot \prod_{j=2}^{i-1} |\ker(\rho_{j,j-1}^{(m)})|$  for  $i = 3, \dots, s(m) + 1$ . In particular, during the  $i$ -th application we use the first at most  $K_i^{\dagger\dagger} P_i^{\dagger\dagger} U_i$  many

different  $(T_{i+1}^{\dagger\dagger}, 2^{4e(m)-t_i}, M_{i+1})$ -Feldman patterns from our collection above. At the end, we obtain a collection  $\mathcal{W}^{\dagger\dagger} \subset (\mathcal{W}_m)^{k^{\dagger\dagger}}$  with

$$\begin{aligned} k^{\dagger\dagger} &= U_{s(m)+1} \cdot T_{s(m)+1}^{\dagger\dagger} \cdot 2^{(4e(m)-t_{s(m)}) \cdot (2M_{s(m)+1}+3)} \\ &= U_{s(m)+1} \cdot T_{s(m)+2}^{\dagger\dagger} \cdot 2^{8e(m) \cdot (M_{s(m)+2}+M_{s(m)+1}+3)-t_{s(m)} \cdot (2M_{s(m)+1}+3)} \end{aligned}$$

and  $|\mathcal{W}^{\dagger\dagger}/\mathcal{Q}_s| = P_{s+1}^{\dagger\dagger}$  for  $s = 1, \dots, s(m)$  as well as  $|\mathcal{W}^{\dagger\dagger}| = P_{s(m)+1}^{\dagger\dagger} 2^{8e(n)}$ .

*Remark 48.* We recall that the choice of  $U_1$  in (8.13) is made in this way to satisfy condition (7.3) from the general substitution step for  $\tilde{R} = 2J_{s(n),n}^3$ . Thus, for the collection  $\mathcal{W}^{\dagger\dagger}$  we obtain the same statements as in Lemma 46 on substantial substrings of length at least  $\frac{k^{\dagger\dagger} h_m}{2J_{s(n),n}^3}$ .

8.2.2. *Construction of  $\mathcal{W}^\dagger$ .* We continue by constructing the collection  $\mathcal{W}^\dagger$ . This time we choose the parameters  $K_i^\dagger \in \mathbb{Z}^+$ ,  $i = 1, \dots, s(m) + 1$ , as follows:

$$K_i^\dagger \cdot |\ker(\rho_{i,i-1}^{(m)})| = 2^{4e(n)} \text{ for } i = 2, \dots, s(m), \quad K_{s(m)+1}^\dagger = 2^{4e(n)}.$$

The parameters  $T_i$ ,  $i = 1, \dots, s(m) + 2$ , are defined by backward recursion:

$$T_{s(m)+2}^\dagger = p_n^2 - p_n, \quad T_i^\dagger = T_{i+1}^\dagger \cdot 2^{(4e(m)-t_i) \cdot (2M_{i+1}+3)} \text{ for } i \leq s(m) + 1.$$

Once again, we apply the general substitution step  $s(m) + 1$  many times as in Case 1. Here, we define  $P_i^\dagger = 2^{(i-1) \cdot 4e(n)}$  for  $i = 1, \dots, s(m) + 1$ . In the  $i$ -th application with  $i \in \{1, \dots, s(m) + 1\}$  we use at most  $K_i^\dagger P_i^\dagger U_i$  many different  $(T_{i+1}^\dagger, 2^{4e(m)-t_i}, M_{i+1})$ -Feldman patterns from our collection above that have not been used in the construction of  $\mathcal{W}^{\dagger\dagger}$ . In particular, in the enumeration of patterns they come after the patterns used for  $\mathcal{W}^{\dagger\dagger}$ . Note that there are enough patterns since

$$K_i^\dagger P_i^\dagger U_i + K_i^{\dagger\dagger} P_i^{\dagger\dagger} U_i \leq (2^{4e(n)} + K_i^{\dagger\dagger}) \cdot 2^{(i-1) \cdot 4e(n)} \cdot U_i \leq M_{i+1}.$$

Then we obtain a collection  $\mathcal{W}^\dagger \subset (\mathcal{W}_m)^{k^\dagger}$  with

$$\begin{aligned} k^\dagger &= U_{s(m)+1} \cdot T_{s(m)+1}^\dagger \cdot 2^{(4e(m)-t_{s(m)}) \cdot (2M_{s(m)+1}+3)} \\ &= U_{s(m)+1} \cdot T_{s(m)+2}^\dagger \cdot 2^{8e(m) \cdot (M_{s(m)+2}+M_{s(m)+1}+3)-t_{s(m)} \cdot (2M_{s(m)+1}+3)} \end{aligned}$$

and  $|\mathcal{W}^\dagger| = 2^{(s(m)+1) \cdot e(n)}$  as well as  $|\mathcal{W}^\dagger/\mathcal{Q}_s| = P_{s+1}^\dagger$  for  $s = 1, \dots, s(m)$ .

*Remark 49.* By the same reasoning as in Remark 48 we obtain for the collection  $\mathcal{W}^{\dagger\dagger}$  the same statements as in Lemma 46 on substantial substrings of length at least  $\frac{k^\dagger h_m}{2J_{s(n),n}^3}$ .

We also make the following observation on the  $\bar{f}$  distance between strings from  $\mathcal{W}^\dagger$  and  $\mathcal{W}^{\dagger\dagger}$ .

**Lemma 50.** *Let  $w \in \mathcal{W}^\dagger$  and  $w' \in \mathcal{W}^{\dagger\dagger}$ . For any substrings  $W$  and  $W'$  of  $w$  and  $w'$ , respectively, of lengths at least  $\frac{k^{\dagger\dagger} h_m}{2J_{s(n),n}^3}$  we have*

$$\bar{f}(W, W') > \alpha_{1,m} - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3}.$$

*Proof.* By construction, already in the first application of the substitution step the Feldman patterns on the level of the 1-equivalence classes of  $w$  and  $w'$  differ. Here, in the enumeration of  $(T, 2^{4e(m)-t_1}, M_2)$ -Feldman patterns the patterns used in the construction of  $\mathcal{W}^{\dagger\dagger}$  come before the ones in  $\mathcal{W}^\dagger$ . Since we also have  $T_2^{\dagger\dagger} < T_2^\dagger$ , we can apply Lemma 39. As in Lemma 46 we obtain from Proposition 43 that

$$\bar{f}(\tilde{W}, \tilde{W}') > \alpha_{1,m} - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3}$$

on all substrings  $\tilde{W}$  and  $\tilde{W}'$  of  $w$  and  $w'$ , respectively, of lengths at least  $\frac{k^\dagger h_m}{2J_{s(n),n}^3}$ .

Since  $\frac{k^\dagger}{k^{\dagger\dagger}} = p_n - 1 < J_{s(n),n}$  our strings  $W$  and  $W'$  are long enough to conclude the statement.  $\square$

*Remark 51.* As in Remark 44 we introduce the notion of  $s$ -pattern type for strings in  $\mathcal{W}^\dagger$  and  $rev(\mathcal{W}^\dagger)$ . As in Lemma 45 we make the following observation. Let  $w \in \mathcal{W}^\dagger$  and  $[P_m]_s$  be a  $(T_{s+1}^\dagger, 2^{4e(m)-t_s}, M_{s+1})$ -Feldman pattern in  $[w]_s$  with building blocks from the tuple  $([A_1]_s, \dots, [A_{2^{4e(m)-t_s}}]_s)$  in  $\mathcal{W}_m/\mathcal{Q}_s^m$ . Then in each complete cycle of  $P_m$  and for each substitution instance  $a_{i,j}$  in  $\mathcal{W}_m$  of each  $[A_i]_s$ ,  $i = 1, \dots, 2^{4e(m)-t_s}$ , its repetition  $(a_{i,j})^{p_n^2 - p_n}$  occurs the same number of times.

8.2.3. *Construction of  $\mathcal{W}_n$ .* To construct  $\mathcal{W}_n$  we recall that  $U_1$  (and, hence,  $U_{s(m)+1}$ ) is a multiple of  $2J_{s(n),n}$ . Thus, we can divide each element  $v_1 \in \mathcal{W}^\dagger$  and  $v_2 \in \mathcal{W}^{\dagger\dagger}$  into  $J = J_{s(n),n}$  or  $2J$  many pieces of equal length:

$$v_1 = v_{1,1}v_{1,2} \dots v_{1,J} \text{ and } v_2 = v_{2,1}v_{2,2} \dots v_{2,2J}.$$

Then we can concatenate their substrings in the following way:

$$(8.15) \quad v_1 * v_2 := v_{2,1}v_{1,1}v_{2,2}v_{2,3}v_{1,2}v_{2,4} \dots v_{2,2J-1}v_{1,J}v_{2,2J}$$

Using this notation we build  $\mathcal{W}_n/\mathcal{Q}_s^n$  for  $0 \leq s \leq s(m)$  inductively: Trivially, for  $s = 0$  we set  $\mathcal{W}_n/\mathcal{Q}_0^n = \{[v_1]_0 * [v_2]_0\}$ , where  $[v_1]_0$  and  $[v_2]_0$  are the single elements in  $\mathcal{W}^\dagger/\mathcal{Q}_0$  and  $\mathcal{W}^{\dagger\dagger}/\mathcal{Q}_0$ , respectively. For  $0 \leq s < s(m)$  we choose a collection  $\Upsilon_s \subseteq \mathcal{W}_n/\mathcal{Q}_s^n$  that intersects each orbit of the action by the group  $\rho_{s+1,s}^m(G_{s+1}^m)$  exactly once. For any element  $C * D \in \Upsilon_s \subseteq \mathcal{W}_n/\mathcal{Q}_s^n$  with  $C \in \mathcal{W}^\dagger/\mathcal{Q}_s$ ,  $D \in \mathcal{W}^{\dagger\dagger}/\mathcal{Q}_s$  we take a collection  $\mathcal{C}$  of substitution instances of  $C$  in  $\mathcal{W}^\dagger/\mathcal{Q}_{s+1}$  that intersects each orbit of  $ker(\rho_{s+1,s}^m)$  exactly once. We note that  $|\mathcal{C}| = K_{s+1}^\dagger$ . Moreover, we choose one substitution instance  $d \in \mathcal{W}^{\dagger\dagger}/\mathcal{Q}_{s+1}$  of  $D$ . For every  $c \in \mathcal{C}$  we let  $c * d$  and its images under the action by  $G_{s+1}^m$  belong to  $\mathcal{W}_n/\mathcal{Q}_{s+1}^n$ . Notice that for any  $g \in G_{s+1}^m$  and  $[w]_s = c * d \in \mathcal{W}_n/\mathcal{Q}_{s+1}^n$  we have  $g[w]_s = (gc) * (gd)$ . Since  $\mathcal{W}^\dagger/\mathcal{Q}_{s+1}$  as well as  $\mathcal{W}^{\dagger\dagger}/\mathcal{Q}_{s+1}$  is closed under the skew diagonal action by construction, we have  $gc \in \mathcal{W}^\dagger/\mathcal{Q}_{s+1}$  and  $gd \in \mathcal{W}^{\dagger\dagger}/\mathcal{Q}_{s+1}$ . Altogether there are  $2^{4e(n)}$  substitution instances of each element of  $\mathcal{W}_n/\mathcal{Q}_s^n$  in  $\mathcal{W}_n/\mathcal{Q}_{s+1}^n$  by our condition  $K_{s+1}^\dagger \cdot |ker(\rho_{s+1,s}^m)| = 2^{4e(n)}$  on  $K_{s+1}^\dagger$ .

For any element  $C * D \in \mathcal{W}_n/\mathcal{Q}_{s(m)}^n$  with  $C \in \mathcal{W}^\dagger/\mathcal{Q}_{s(m)}$ ,  $D \in \mathcal{W}^{\dagger\dagger}/\mathcal{Q}_{s(m)}$  we pair each of the  $2^{4e(n)}$  many substitution instances  $v_i \in \mathcal{W}^\dagger$  of  $C$  with  $2^{4e(n)}$  many substitution instances  $d_1^{(i)}, \dots, d_{2^{4e(n)}}^{(i)} \in \mathcal{W}^{\dagger\dagger}$  of  $D$ . This is consistent with the  $2^{8e(n)}$  substitution instances of  $D$  in  $\mathcal{W}^{\dagger\dagger}$  that we have according to our choice

$K_{s(m)+1}^{\dagger\dagger} = 2^{8e(n)}$  in (8.14). Then we define the new equivalence classes

$$[v_i]_{s(m)+1} = \left\{ v_i * d_j^{(i)} : j = 1, \dots, 2^{4e(n)} \right\}.$$

Moreover, we set

$$\mathcal{W}_n = \left\{ v_i * d_j^{(i)} : v_i \in \mathcal{W}^\dagger, j = 1, \dots, 2^{4e(n)} \right\}.$$

We note that these definitions fulfill specification (Q6).

8.2.4. *Equivalence relations on  $\mathcal{W}_n$ .* The equivalence classes  $\mathcal{W}_n/\mathcal{Q}_s^n$  for  $s \leq s(m)$  are given by the product equivalence relation. The new equivalence relation  $\mathcal{Q}_{s(n)}^n$  satisfies

$$v_1 * v_2 \sim v'_1 * v'_2 \text{ if and only if } v_1 = v'_1 \text{ and } [v_2]_{s(m)} = [v'_2]_{s(m)},$$

that is, two words in  $\mathcal{W}_n$  are equivalent with respect to  $\mathcal{Q}_{s(n)}^n$  if and only if when considering each as a concatenation of  $J_{s(n),n}$  strings of equal length all their central strings (whose concatenation forms an element from  $\mathcal{W}^\dagger$ ) coincide and their initial and terminal strings (whose concatenation forms an element from  $\mathcal{W}^{\dagger\dagger}$ ) are equivalent with respect to the product equivalence relation induced from  $\mathcal{Q}_{s(m)}^m$ . Hence, for  $w = v_1 * v_2 \in \mathcal{W}_n$  we have

$$[w]_{s(n)} = \left\{ v_1 * v'_2 : [v_2]_{s(m)} = [v'_2]_{s(m)} \right\}.$$

We also observe that

$$(8.16) \quad \frac{k^\dagger}{k} = \frac{k^\dagger}{k^\dagger + k^{\dagger\dagger}} = \frac{T_{s(m)+2}^\dagger}{T_{s(m)+2}^\dagger + T_{s(m)+2}^{\dagger\dagger}} = 1 - \frac{1}{p_n} > 1 - \epsilon_n$$

by our condition on  $p_n$  in equation (8.5). In this way, specification (Q4) is satisfied and the equivalence relation  $\mathcal{Q}_{s(n)}^n$  refines  $\mathcal{Q}_{s(m)}^m$ .

8.2.5. *Group actions.* For all  $s \leq s(m)$  we have  $G_s^n = G_s^m$  and the group actions are defined via the skew-diagonal action. According to our construction in Section 8.2.3,  $\mathcal{W}_n/\mathcal{Q}_s^n$  is closed under the skew diagonal action by  $G_s^n$ .

To define the action by  $G_{s(n)}^n = \mathbb{Z}_2$  we apply Lemma 34 with  $G \times \mathbb{Z}_2 := G_{s(m)}^n = G_{s(m)}^m$ ,  $H$  the trivial group,  $H \times \mathbb{Z}_2 := G_{s(n)}^n$ ,  $X/\mathcal{Q} := \mathcal{W}_n/\mathcal{Q}_{s(m)}^m$ ,  $X/\mathcal{R} := \mathcal{W}_n/\mathcal{Q}_{s(n)}^n$ , and  $\rho := \rho_{s(n),s(m)}$  in the notation of the Lemma. Hence, we obtain an action of  $G_{s(n)}^n = \mathbb{Z}_2$  on  $\mathcal{W}_n/\mathcal{Q}_{s(n)}^n$  subordinate to the action of  $G_{s(m)}^m$ .

8.2.6. *Further properties.* Also in this case we make observations regarding  $\bar{f}$  distances.

**Lemma 52.** (1) *For every  $s \in \{1, \dots, s(m)\}$  we have*

$$(8.17) \quad \bar{f}(W, W') > \alpha_{s,m} - \frac{3}{R_m} - \frac{2}{R_n}$$

*on any substrings  $W$  and  $W'$  of at least  $\frac{h_n}{R_n}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_s \neq [w']_s$ .*

(2) In case of  $s = s(n)$  we have

$$(8.18) \quad \bar{f}(W, W') > \beta_m - \frac{3}{R_m} - \frac{2}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{h_n}{R_n}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_{s(n)} \neq [w']_{s(n)}$ .

(3) Moreover, we have

$$(8.19) \quad \bar{f}(W, W') > \frac{1}{p_n} \cdot \left( \beta_m - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3} \right) - \frac{2}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{h_n}{R_n}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_n$  with  $w \neq w'$ .

*Proof.* We decompose the strings  $W$  and  $W'$  into the substrings of length  $\frac{h_n}{J_{s(n),n}}$  according to equation (8.15). To complete partial ones at the beginning and end we add fewer than  $4\frac{h_n}{J_{s(n),n}}$  symbols in total which corresponds to a proportion of less than

$$\frac{2R_n}{J_{s(n),n}} < \frac{1}{R_n}$$

of the total length by equation (8.12). This might increase the  $\bar{f}$  distance by at most  $\frac{1}{R_n}$  by Fact 9.

- (1) On each of these substrings of length  $\frac{h_n}{J_{s(n),n}}$  the part belonging to  $\mathcal{W}^\dagger$  has length  $\frac{k^\dagger h_m}{J_{s(n),n}}$  while the part belonging to  $\mathcal{W}^{\dagger\dagger}$  represents a proportion of  $\frac{1}{p_n}$  of the length by equation (8.16). We delete them for the consideration of this subcase increasing the  $\bar{f}$  distance by at most  $\frac{2}{p_n}$  according to Fact 9. On the remaining part we use Remark 49 on the level of the equivalence class  $s \in \{1, \dots, s(m)\}$ . With the aid of conditions (8.5) on  $p_n$  and (8.7) applied on  $e(m)$  we obtain the following estimate:

$$\bar{f}(W, W') > \alpha_{s,m} - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3} - \frac{2}{p_n} - \frac{1}{R_n} > \alpha_{s,m} - \frac{3}{R_m} - \frac{2}{R_n}.$$

- (2) We follow the argument from the previous subcase but apply Remark 49 on the level of the actual blocks in  $\mathcal{W}^\dagger$ .
- (3) While the strings might coincide on the central parts belonging to  $\mathcal{W}^\dagger$ , this time the  $\bar{f}$  distance comes from the initial and terminal parts of length  $\frac{k^{\dagger\dagger} h_m}{2J_{s(n),n}}$  belonging to  $\mathcal{W}^{\dagger\dagger}$ . To see this, we further subdivide the strings of length  $\frac{kh_m}{J_{s(n),n}}$  into pieces of length  $\frac{k^{\dagger\dagger} h_m}{2J_{s(n),n}}$ . Lemma 50 allows us to estimate the  $\bar{f}$  distance if one of those pieces lies in  $\mathcal{W}^\dagger$  and the other one lies in  $\mathcal{W}^{\dagger\dagger}$ . For both pieces lying in  $\mathcal{W}^{\dagger\dagger}$  we apply Remark 48 for blocks in  $\mathcal{W}^{\dagger\dagger}$ . Hence, we obtain by applying Lemma 13 with  $\bar{f} \geq \frac{1}{p_n}$  that

$$\begin{aligned} \bar{f}(W, W') &> \frac{1}{p_n} \cdot \left( \beta_m - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3} \right) - \frac{1}{J_{s(n),n}} - \frac{1}{R_n} \\ &> \frac{1}{p_n} \cdot \left( \beta_m - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3} \right) - \frac{2}{R_n} \end{aligned}$$

□

The expressions on the right hand side in Lemma 52 motivate the following definitions. For  $s \in \{1, \dots, s(m)\}$  we set

$$(8.20) \quad \alpha_{s,n} := \alpha_{s,m} - \frac{3}{R_m} - \frac{2}{R_n},$$

while for  $s = s(n)$  we introduce

$$(8.21) \quad \alpha_{s(n),n} := \beta_m - \frac{3}{R_m} - \frac{2}{R_n}.$$

We notice that

$$\alpha_{s(n),n} > \beta_m - \frac{1}{3}\beta_m - \frac{1}{20p_n}\beta_m \geq \frac{\beta_m}{2} > 0$$

by equations (8.2) and (8.6). Hence, we get

$$(8.22) \quad \frac{1}{R_n} < \frac{\alpha_{s(n),n}}{20p_n}$$

by condition (8.6). Finally, we set

$$\beta_n := \frac{1}{p_n} \cdot \left( \beta_m - \frac{1}{R_m} - \frac{4}{2^{e(m)}} - \frac{1}{2J_{s(n),n}^3} \right) - \frac{2}{R_n},$$

and we note with the aid of equation (8.6) again that

$$\beta_n > \frac{1}{p_n} \cdot \frac{\beta_m}{2} - \frac{2}{R_n} \geq \frac{9}{20} \cdot \frac{\beta_m}{p_n} \geq \frac{9}{R_n},$$

which not only yields  $\beta_n > 0$ , but also  $R_n > \frac{9}{\beta_n}$ . That is, the induction assumption (8.2) for the next step is satisfied.

*Remark 53.* As in Remark 47 we prepare the transfer to the setting of diffeomorphisms by considering the words in  $\mathcal{W}_n$  as concatenations of  $k_m$  symbols in the alphabets  $\Sigma_m$  and  $\Sigma_{m,s}$  for  $s \leq s(m)$ . As in the proof of Lemma 52 we use Proposition 43 to obtain

$$\bar{f}_{\Sigma_{m,s}}(W, W') > 1 - \frac{4}{2^{e(m)}} - \frac{1}{R_n} - \frac{1}{R_n} - \frac{2}{p_n} > 1 - \frac{1}{R_m} - \frac{2}{R_n}.$$

on any substrings  $W$  and  $W'$  of at least  $\frac{k_m}{R_n}$  consecutive  $\Sigma_{m,s}$ -symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_s \neq [w']_s$  for  $s \leq s(m)$ . Similarly, the proof of part (3) in Lemma 52 yields

$$(8.23) \quad \bar{f}_{\Sigma_m}(W, W') > \frac{1}{p_n} \cdot \left( 1 - \frac{4}{2^{e(m)}} - \frac{1}{R_n} \right) - \frac{2}{R_n} > \frac{1}{2p_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{k_m}{R_n}$  consecutive  $\Sigma_m$ -symbols in any words  $w, w' \in \mathcal{W}_n$  with  $w \neq w'$ . Following the proof of part (2) in Lemma 52 we can also express the  $\bar{f}$ -distance on substantial substrings of different classes of the new equivalence relation  $\mathcal{Q}_{s(n)}^n$  in the alphabet  $\Sigma_m$ : We have

$$(8.24) \quad \bar{f}_{\Sigma_m}(W, W') > 1 - \frac{4}{2^{e(m)}} - \frac{1}{R_n} - \frac{1}{R_n} - \frac{2}{p_n} > 1 - \frac{1}{R_m} - \frac{2}{R_n}$$

on any substrings  $W$  and  $W'$  of at least  $\frac{k_m}{R_n}$  consecutive  $\Sigma_m$ -symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_{s(n)} \neq [w']_{s(n)}$ .



*Remark 54.* In the base case  $m = 0$  of our inductive process we have  $\mathcal{W}_0 = \{1, \dots, 2^{12}\}$ , that is, the  $m$ -blocks are actual symbols and  $e(0) = 3$ . Thus, by (8.23) and (8.24), we can set

$$\alpha_{1,n} = \min\left(\frac{1}{2} - \frac{2}{R_n} - \frac{2}{p_n}, \frac{1}{9}\right), \quad \beta_n = \min\left(\frac{1}{p_n} \cdot \left(\frac{1}{2} - \frac{1}{R_n}\right) - \frac{2}{R_n}, \alpha_{1,n}\right).$$

**8.3. Conclusion of the construction.** We recall the definition of the function  $M(s)$  from Definition 19. For every  $s \in \mathbb{Z}^+$  we use equations (8.21) and (8.20) as well as (8.11) to conclude for every  $\ell > M(s)$  with  $\sigma_\ell \in \mathcal{T}$  that

$$\alpha_{s,\ell} > \alpha_{s,M(s)} - \frac{3}{R_{M(s)}} - \sum_{k>M(s):\sigma_k \in \mathcal{T}} \frac{5}{R_k}.$$

Together with condition (8.6) and estimate (8.22) this implies

$$\alpha_{s,\ell} > \alpha_{s,M(s)} - \frac{6\alpha_{s,M(s)}}{40p_{M(s)}} - \frac{5}{40}\alpha_{s,M(s)} > \frac{\alpha_{s,M(s)}}{2}.$$

Hereby, we obtain that for every  $s \in \mathbb{Z}^+$  the decreasing sequence  $(\alpha_{s,\ell})_{\ell \geq M(s):\sigma_\ell \in \mathcal{T}}$  is bounded from below by  $\frac{\alpha_{s,M(s)}}{2}$  and we denote its limit by  $\alpha_s$ . This yields the following proposition.

**Proposition 55.** *We have for every  $n \geq M(s)$  with  $\sigma_n \in \mathcal{T}$  that*

$$(8.25) \quad \bar{f}(W, W') > \alpha_s$$

*on any substrings  $W$  and  $W'$  of at least  $\frac{hn}{R_n}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_n$  with  $[w]_s \neq [w']_s$ .*

## 9. PROOF OF NON-KAKUTANI EQUIVALENCE

At first, we prove in Proposition 56 that for any tree  $\mathcal{T} \in \mathcal{Trees}$  the constructed transformation  $T = \Psi(\mathcal{T})$  is not isomorphic to a special transformation  $T^f$  for a nontrivial roof function  $f : X \rightarrow \mathbb{Z}^+$ . Then, in Subsection 9.2, we will use Proposition 56 to show that  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  can only be Kakutani equivalent if they are actually evenly Kakutani equivalent. In Subsection 9.3 we finish the proof by showing that for a tree  $\mathcal{T} \in \mathcal{Trees}$  without an infinite branch  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  are not evenly Kakutani equivalent.

**9.1.  $T^f$  is not isomorphic to  $T$  for nontrivial  $f$ .** We recall that for an ergodic invertible measure-preserving transformation  $T : (X, \mu) \rightarrow (X, \mu)$  and an integrable function  $f : X \rightarrow \mathbb{Z}^+$  one defines the *special transformation* (or sometimes called *integral transformation*)  $T^f$  on the set  $X^f = \{(x, j) : x \in X, 1 \leq j \leq f(x)\}$  by

$$T^f(x, j) = \begin{cases} (x, j+1) & \text{if } j < f(x), \\ (Tx, 1) & \text{if } j = f(x). \end{cases}$$

The map  $T^f$  preserves the probability measure  $\mu^f$  that is the restriction to  $X^f$  of  $(\mu \times \lambda) / (\int f d\mu)$  on  $X \times \mathbb{Z}^+$ , where  $\lambda$  is the counting measure on  $\mathbb{Z}^+$ . Moreover,  $T^f$  is ergodic with respect to  $\mu^f$  because  $T$  is ergodic.

We denote  $\{(x, j) \in X^f : j > 1\}$  by  $H$ . For a measurable partition  $P = \{C_1, \dots, C_N\}$  of  $X$  we define  $P^f = \{C_1, \dots, C_N, H\}$  as the corresponding measurable partition of  $X^f$ . Accordingly, if  $(T^f)^i(x) \in H$ , we let  $h$  be the letter in the  $i$ th position of the

$(T^f, P^f)$ -name of  $x$ . Hence the  $(T^f, P^f)$ -name is obtained from the  $(T, P)$ -name by inserting  $h$ 's.

In this section we consider a transformation of the form  $T = \Psi(\mathcal{T})$  for any  $\mathcal{T} \in \text{Trees}$ . We recall that  $\Psi(\mathcal{T})$  was built with a uniquely readable and strongly uniform construction sequence  $\{\mathcal{W}_n(\mathcal{T}) : \sigma_n \in \mathcal{T}\}$ . To simplify notation we enumerate it as  $\{\mathcal{W}_n\}_{n \in \mathbb{N}}$ , that is,  $(n+1)$ -words are built by concatenating  $n$ -words.

Following the approach in [ORW82, chapter 11] (see also [Be15, section 4] for another exposition) we show that for any nontrivial integrable function  $f : [0, 1] \rightarrow \mathbb{Z}^+$  the special transformation  $T^f$  is not isomorphic to  $T$ . In other words,  $T^f \cong T$  implies that  $f$  is the constant function equal to 1.

**Proposition 56.** *Suppose that  $\mathcal{T} \in \text{Trees}$ ,  $T = \Psi(\mathcal{T})$ , and  $f : X \rightarrow \mathbb{Z}^+$  is integrable. If  $T^f \cong T$ , then  $\int f d\mu = 1$ . Equivalently, if  $A$  is a measurable subset of  $X$  and  $T_A \cong T$ , then  $\mu(A) = 1$ .*

If  $T$  and  $T^f$  were isomorphic by an isomorphism  $\Phi : X \rightarrow X^f$ , then for every  $\tau > 0$  there would be  $K(\tau) \in \mathbb{N}$  and a partition  $P(\tau) \subset \bigvee_{i=-K(\tau)}^{K(\tau)} T^i P$  such that  $|P(\tau)| = |P^f|$  and  $d(P(\tau), \Phi^{-1}(P^f)) < \tau$ . For the remainder of Subsection 9.1, we assume that such an isomorphism  $\Phi$  exists and that  $\int f d\mu > 1$ . We obtain a contradiction, which proves Proposition 56.

*Remark 57.* Such a partition  $P(\tau)$  defines an equivalence relation on  $\bigvee_{i=-K(\tau)}^{K(\tau)} T^i P$  whereby two elements  $Q_1, Q_2$  of this partition are equivalent if they are contained in the same element of  $P(\tau)$ . For  $Q \in \bigvee_{i=-K(\tau)}^{K(\tau)} T^i P$ , denote the equivalence class containing  $Q$  by  $\overline{Q}$ . From the  $(T, P)$ -trajectory  $\dots a_{-1} a_0 a_1 \dots$  of a point  $x \in X$  one can derive a  $(T, \bigvee_{i=-K(\tau)}^{K(\tau)} T^i P)$  trajectory

$$\dots (a_{-K(\tau)-1} \dots a_{-1} \dots a_{K(\tau)-1}) (a_{-K(\tau)} \dots a_0 \dots a_{K(\tau)}) (a_{-K(\tau)+1} \dots a_1 \dots a_{K(\tau)+1}) \dots$$

with  $(a_{-K(\tau)+i} \dots a_i \dots a_{K(\tau)+i})$  such that

$$x \in T^{-(K(\tau)+i)}(C_{a_{-K(\tau)+i}}) \cap \dots \cap T^{-i}(C_{a_i}) \cap \dots \cap T^{-(K(\tau)+i)}(C_{a_{K(\tau)+i}}).$$

By taking equivalence classes we obtain the  $(T, P(\tau))$ -trajectory

$$\dots \overline{(a_{-K(\tau)-1} \dots a_{-1} \dots a_{K(\tau)-1})} \overline{(a_{-K(\tau)} \dots a_0 \dots a_{K(\tau)})} \overline{(a_{-K(\tau)+1} \dots a_1 \dots a_{K(\tau)+1})} \dots$$

with  $\overline{(a_{-K(\tau)+i} \dots a_i \dots a_{K(\tau)+i})}$  such that

$$x \in \overline{T^{-(K(\tau)+i)}(C_{a_{-K(\tau)+i}}) \cap \dots \cap T^{-i}(C_{a_i}) \cap \dots \cap T^{-(K(\tau)+i)}(C_{a_{K(\tau)+i}})}.$$

In this way the partition  $P(\tau)$  can be identified with a *finite code* and we use this term synonymously.

**Definition 58.** A *code* of length  $2K+1$  is a function  $\phi : \Sigma^{\mathbb{Z} \cap [-K, K]} \rightarrow \Sigma$ . We denote the length of the code by  $|\phi|$ . Given a code  $\phi$  of length  $2K+1$  the *stationary code*  $\overline{\phi}$  on  $\Sigma^{\mathbb{Z}}$  determined by  $\phi$  is defined as  $\overline{\phi}(s)$  for any  $s \in \Sigma^{\mathbb{Z}}$ , where for any  $l \in \mathbb{Z}$

$$\overline{\phi}(s)(l) = \phi(s \upharpoonright [l-K, l+K]).$$

In this setting  $\overline{\phi}(s) \upharpoonright [-N, N]$  denotes the string of symbols

$$\overline{\phi}(s)(-N) \overline{\phi}(s)(-N+1) \dots \overline{\phi}(s)(N-1) \overline{\phi}(s)(N)$$

in  $\Sigma^{2N+1}$ .

*Remark 59.* If we consider the symbolic systems  $\mathbb{K}$  and  $\mathbb{K}^{-1}$  as processes with generators  $P = Q = \{(\sigma) : \sigma \in \Sigma\}$ , then a code  $P(\varepsilon) \subset \bigvee_{i=-K}^K T^i Q$  as in Remark 57 corresponds to a code  $\phi$  as described in Definition 58.

*Remark 60.* There is an ambiguity in applying a code  $\phi$  of length  $2K + 1$  to blocks of the form  $s \upharpoonright [a, b]$ : It does not make sense to apply it to the initial string  $s \upharpoonright [a, a + K - 1]$  and end string  $s \upharpoonright [b - K + 1, b]$ . However, if  $b - a$  is large with respect to the code length  $2K + 1$ , we can fill in  $\phi(s) \upharpoonright [a, a + K - 1]$  and  $\phi(s) \upharpoonright [b - K + 1, b]$  arbitrarily and it makes a negligible difference to the  $\bar{d}$  or  $\bar{f}$  distance. In particular, if  $\bar{f}(\phi(s), \bar{s}) < \varepsilon$ , then for all large enough  $a, b \in \mathbb{N}$ , we have

$$\bar{f}(\phi(s) \upharpoonright [-a, b], \bar{s} \upharpoonright [-a, b]) < 2\varepsilon$$

independently of how we fill in the ambiguous portion. The general phenomenon of ambiguity or disagreement at the beginning and end of large strings is referred to by the phrase *end effects*.

We show that for a nontrivial function  $f$ , there exists  $\tau > 0$  sufficiently small so that there cannot be a finite code from  $(T, P)$  to  $(T^f, P^f)$  that approximates  $P^f$  within distance  $\tau$ . First we investigate some properties of  $(T^f, P^f)$ -names. We divide such names into  $n$ -blocks as follows: A  $(T^f, P^f)$ - $n$ -block consists of all the symbols from the leftmost symbol of a  $(T, P)$ - $n$ -block up to (but not including) the leftmost symbol in the next  $(T, P)$ - $n$ -block. In the following we denote the  $(T^f, P^f)$ - $n$ -block corresponding to a  $(T, P)$ - $n$ -block,  $w_n$ , by  $\tilde{w}_n$ . In general, the  $(T^f, P^f)$ - $n$ -blocks have varying lengths. The following fact from [ORW82, Proposition 2.1, p.84] based on the ergodic theorem shows that a large proportion of them have almost the same length.

**Fact 61.** *Let  $T : (X, \mu) \rightarrow (X, \mu)$  be an ergodic transformation and  $f : X \rightarrow \mathbb{Z}^+$  be integrable. For a measurable subset  $F$  of  $X$ , let  $R_F, R_F^f : F \rightarrow \mathbb{Z}^+$  denote, respectively, the return time under  $T$  to  $F$ , and the return time under  $T^f$  to  $F$  considered as a subset of the base of the tower  $X^f$ . Then for every  $\varepsilon > 0$  there is  $N(\varepsilon) \in \mathbb{N}$  such that for every measurable subset  $F$  of  $X$  with  $R_F \geq N(\varepsilon)$  there is a measurable subset  $F'$  of  $F$  such that*

$$\int_{F'} R_F^f d\mu \geq \int f d\mu - \varepsilon$$

and  $R_F(x) \cdot \left( \int f d\mu - \varepsilon \right) \leq R_F^f(x) \leq R_F(x) \cdot \left( \int f d\mu + \varepsilon \right)$  for all  $x \in F'$ .

In particular, the last line of Fact 61 says that if we think of  $F$  as the initial points in  $n$ -blocks, then for  $n$  large enough,  $n$ -blocks are expanded nearly uniformly. The inequality  $\int_{F'} R_F^f d\mu \geq \int f d\mu - \varepsilon$  implies that the set of indices in a typical  $(T^f, P^f)$ -trajectory in blocks not expanded within those bounds is at most  $\varepsilon / (\int f d\mu) \leq \varepsilon$ .

Now for almost every point  $x$  we consider its  $(T, P)$ -name  $a(T, P, x)$ , its  $(T, P(\tau))$ -name  $a(T, P(\tau), x)$ , and its  $(T, \Phi^{-1}(P^f))$ -name  $a(T, \Phi^{-1}(P^f), x)$ , which differs from its  $(T, P(\tau))$ -name on a set of indices with density at most  $\tau$ . Note that the  $(T, \Phi^{-1}(P^f))$ -name of  $x$  is identified with the  $(T^f, P^f)$ -name of  $\Phi(x)$ . Thus, for any sequence of integers (considered as indices labeling positions in names) we can consider the three names defined on it.




FIGURE 9.1. Feldman patterns  $F_{n,i}$  of  $n$ -blocks in the  $P$ -name of  $x$ , their corresponding strings  $\tilde{F}_{n,i}$  in the  $\Phi^{-1}(P^f)$ -name of  $x$ , and the decomposition of the set  $\mathbb{Z}$  of indices into  $I_i = F_{n,l} \cap \tilde{F}_{n,\tilde{l}}$ .

We recall from our constructions in Section 8 that the  $(n+1)$ -words in  $\mathcal{W}_{n+1}$  are built as substitution instances of the different 1-equivalence classes in  $\mathcal{W}_{n+1}/\mathcal{Q}_1^{n+1}$ . Considered on the level of the 1-equivalence relation these elements in  $\mathcal{W}_{n+1}/\mathcal{Q}_1^{n+1}$  are concatenations of different Feldman patterns with building blocks in  $\mathcal{W}_n/\mathcal{Q}_1^n$ . In the following we denote these Feldman patterns of 1-equivalence classes of  $n$ -blocks in the  $P$ -name of  $x$  by  $F_{n,i}$ , labeled so that  $F_{n,0}$  contains the position 0. Similarly the strings in the  $\Phi^{-1}(P^f)$ -name are denoted by  $\tilde{F}_{n,i}$ , where  $\tilde{F}_{n,0}$  contains the position 0. Thus

$$a(T, P, x) = \dots F_{n,-2} F_{n,-1} F_{n,0} F_{n,1} F_{n,2} \dots$$

$$a(T^f, P^f, \Phi(x)) = a(T, \Phi^{-1}(P^f), x) = \dots \tilde{F}_{n,-2} \tilde{F}_{n,-1} \tilde{F}_{n,0} \tilde{F}_{n,1} \tilde{F}_{n,2} \dots$$

We decompose the set  $\mathbb{Z}$  of indices of trajectories into sets  $I_i = F_{n,l} \cap \tilde{F}_{n,\tilde{l}}$ , that is,  $I_i$  is the intersection of the set of indices of a 1-pattern  $F_{n,l}$  in the  $(T, P)$ -trajectory and of the set of indices of a 1-pattern  $\tilde{F}_{n,\tilde{l}}$  in the  $(T^f, P^f)$ -trajectory (see Figure 9.1). Moreover,  $I_i(T, P)$  denotes the  $(T, P)$ -trajectory of  $x$  on the indices in  $I_i$ .

If we are in the second case  $s(n+1) = s(n) + 1$  of the construction (see Subsection 8.2), then we delete those parts of the  $(n+1)$ -words coming from the collection  $\mathcal{W}^{\dagger\dagger}$ . We also delete the corresponding strings in the  $(T^f, P^f)$ -trajectory. Hereby, we neglect a set of indices with density at most  $2\epsilon_{n+1}$ . After this, each element in  $\mathcal{W}_{n+1}/\mathcal{Q}_1^{n+1}$  is a concatenation of  $U_2^{(n+1)}$  many different  $(T_2^{(n+1)}, 2^{4e(n)-t_1}, M_2^{(n+1)})$ -Feldman patterns with building blocks out of  $\mathcal{W}_n/\mathcal{Q}_1^n$ , where  $U_2^{(n+1)}$ ,  $T_2^{(n+1)}$ , and  $M_2^{(n+1)}$  are the corresponding parameters chosen at our stage  $n+1$  of the construction in Subsections 8.1 and 8.2. Note that we had to remove the parts coming from  $\mathcal{W}^{\dagger\dagger}$  to have unique values of  $T_2^{(n+1)}$  and to guarantee that all Feldman patterns have the same length, which we denote by  $f_n$ . All of this automatically holds in the case  $s(n+1) = s(n)$  of the construction (see Subsection 8.1).

Recall the definition of  $\alpha_1$  from Proposition 55. We let

$$0 < \delta < \min \left( \frac{\int f \, d\mu - 1}{3 \int f \, d\mu}, \frac{\alpha_1}{160 \int f \, d\mu} \right),$$

and let  $n$  be sufficiently large such that

- (n1)  $\frac{2}{2^{\epsilon(n)}} < \delta$ ,  $\epsilon_{n+1} < \delta$ , and  $\frac{20}{R_n} < \alpha_1$ ,
- (n2)  $2 \int f \, d\mu < \min \left( 2^{\epsilon(n)}, U_2^{(n+1)}, \frac{\alpha_1 p_{n+1}}{80} \right)$ ,
- (n3) the code length  $2K(\tau) + 1$  is less than  $h_n$ ,
- (n4)  $h_n$  is larger than the number  $N(\delta^3)$  from Fact 61 for  $\delta^3$ .

In particular,  $h_n > N(\delta^3)$  implies that the set of indices lying in  $(T, \Phi^{-1}(P^f))$ - $n$ -words  $\tilde{w}_n$ , that are too long (i.e.  $|\tilde{w}_n| > (\int f \, d\mu + \delta^3) h_n$ ) or too short (i.e.  $|\tilde{w}_n| < (\int f \, d\mu - \delta^3) h_n$ ), has density at most  $\delta^3$ . Then at most  $\delta$  of the indices are lying in a 1-pattern  $\tilde{F}_{n,j}$  that contains a proportion of more than  $\delta^2$  of indices lying in too long or too short  $n$ -words. Moreover, condition (n4) implies that all but a set of indices of density at most  $\delta^3$  lie in 1-patterns  $\tilde{F}_{n,j}$  with  $(\int f \, d\mu - \delta^3) f_n \leq |\tilde{F}_{n,j}| \leq (\int f \, d\mu + \delta^3) f_n$ . Hence, at most  $\delta + \delta^3 < 2\delta$  of indices belong to those ‘‘bad’’ 1-patterns  $\tilde{F}_{n,j}$  and we neglect them until the end of the proof of Proposition 56.

In the first step, we give a lower bound for the  $\bar{d}$ -distance between the strings on  $I_i$  from the  $(T, P(\tau))$ -trajectory and the  $(T, \Phi^{-1}(P^f))$ -trajectory of  $x$ , if the 1-pattern types are different from each other. In fact, in (9.5) we even obtain a lower bound on the  $\bar{f}$  distance. The proof uses the Coding Lemma 41 and relies on the different periodicities of maximum repetitions of the same 1-equivalence class of  $n$ -blocks.

**Lemma 62.** *Suppose for  $I_i = F_{n,l} \cap \tilde{F}_{n,\tilde{l}}$  that  $|I_i| \geq \delta f_n$  and that  $F_{n,l}$  and  $\tilde{F}_{n,\tilde{l}}$  have different 1-pattern types. Then we have*

$$\bar{d}(I_i(T, \Phi^{-1}(P^f)), I_i(T, P(\tau))) \geq \frac{\alpha_1}{20 \cdot \int f \, d\mu},$$

where  $\alpha_1$  is defined in Proposition 55.

*Proof.* By the properties of the ‘‘good’’ 1-pattern  $\tilde{F}_{n,\tilde{l}}$  as described above,  $I_i(T, \Phi^{-1}(P^f))$  is  $2\delta$ -close in  $\bar{f}$  to a string whose  $(T, \Phi^{-1}(P^f))$ - $n$ -words have length  $\lfloor \int f \, d\mu \cdot h_n \rfloor$ . Until the end of the proof we identify  $I_i(T, \Phi^{-1}(P^f))$  with this string.

We denote the tuple of building blocks for  $[F_{n,l}]_1$  by  $([A_1]_1, \dots, [A_{2^{4e(n)-t_1}}]_1)$  and the one for  $[\tilde{F}_{n,\tilde{l}}]_1$  by  $([\tilde{A}_1]_1, \dots, [\tilde{A}_{2^{4e(n)-t_1}}]_1)$ . Furthermore, let  $r$  and  $\tilde{r}$  be the 1-pattern types of  $F_{n,l}$  and  $\tilde{F}_{n,\tilde{l}}$ , respectively.

Recalling  $p'_{n+1} = p_{n+1}^2$  in Case 1 ( $s(n+1) = s(n)$ ) or  $p'_{n+1} = p_{n+1}^2 - p_{n+1}$  in Case 2 ( $s(n+1) = s(n) + 1$ ) of the construction, we let

$$\begin{aligned} L &= p'_{n+1} \cdot h_n, \\ N &= 2^{4e(n)-t_1} \end{aligned}$$

in the notation of the Coding Lemma 41. Moreover, we subdivide  $\tilde{F}_{n,\tilde{l}}$  into substitution instances  $\tilde{\Upsilon}_{i,m}$  of a maximal repetition  $[\tilde{A}_m]_1^{T_2 N^{2\tilde{r}}}$ . Then we further subdivide

$\tilde{\Upsilon}_{i,m}$  into

$$q := \left\lfloor \frac{|\tilde{\Upsilon}_{i,m}|}{T_2 \cdot N^{2\tilde{r}-1} \cdot h_n} \right\rfloor \cdot \frac{T_2}{p'_{n+1}} \cdot N^{2\tilde{r}-1} = \left\lfloor N \cdot \int f \, d\mu \right\rfloor \cdot \frac{T_2}{p'_{n+1}} \cdot N^{2\tilde{r}-1}$$

many strings of  $L$  consecutive symbols, where we recall that  $T_2$  is a multiple of  $p'_{n+1}$ . In the notation of the Coding Lemma each of these  $L$ -strings corresponds to a  $B_{m,j}^{(i)}$  and altogether they form  $\Upsilon_{i,m}$ . Then we also have

$$z = \left\lfloor \frac{|F_{n,l}|}{NqL} \right\rfloor = \left\lfloor \frac{N^{2(M_2-\tilde{r})+3}}{[N \cdot \int f \, d\mu]} \right\rfloor.$$

Note that

$$(9.1) \quad 0 \leq \frac{|\tilde{\Upsilon}_{i,m}| - |\Upsilon_{i,m}|}{|\tilde{\Upsilon}_{i,m}|} \leq \frac{T_2 \cdot N^{2\tilde{r}-1} \cdot h_n}{T_2 \cdot N^{2\tilde{r}} \cdot \int f \, d\mu \cdot h_n} < \frac{1}{N}.$$

Combining Fact 12, Lemma 13, and Proposition 55 we have

$$(9.2) \quad \bar{f}(B, B') > \frac{\alpha_1 - \frac{1}{R_n}}{\int f \, d\mu} - \frac{2}{p_{n+1}} > \frac{9\alpha_1}{10 \cdot \int f \, d\mu}$$

on any substrings  $B$  and  $B'$  of at least  $\frac{L}{p_{n+1}}$  consecutive symbols in any  $B_{m,j}^{(i)}$  and  $B_{m',j'}^{(i')}$ , respectively, with  $m \neq m'$ . Here we used conditions (n1) and (n2) in the last estimate. Hence, the assumption of the Coding Lemma is satisfied. Now we examine both possible cases  $r < \tilde{r}$  and  $r > \tilde{r}$ .

In the case  $r < \tilde{r}$  the number  $T_2 N^{2r}$  of maximum repetitions of  $n$ -blocks of the same 1-class in  $F_{n,l}$  is smaller than in  $\tilde{F}_{n,\tilde{l}}$ . We subdivide  $F_{n,l}(T, P)$  into substitution instances  $\tilde{\Lambda}_{i,m}$  of

$$u_1 := \frac{|\Upsilon_{i,m}|}{T_2 N^{2r+1} h_n} = \frac{qL}{T_2 N^{2r+1} h_n} = \left\lfloor N \cdot \int f \, d\mu \right\rfloor \cdot N^{2(\tilde{r}-r-1)}$$

many consecutive complete cycles in  $[F_{n,l}]_s$ . Note that

$$(9.3) \quad 0 \leq \frac{|F_{n,l}| - zNu_1 \cdot T_2 N^{2r+1} h_n}{|F_{n,l}|} \leq \frac{zT_2 N^{2r+2} h_n}{T_2 N^{2M_2+3} h_n} < \frac{1}{N}.$$

Lemma 45 and Remark 51 yield that for each substitution instance  $a_{x,y}$  of each  $[A_x]_1$ ,  $x = 1, \dots, 2^{4e(n)-t_1}$ , the repetition  $(a_{x,y})^{p'_{n+1}}$  occurs the same number of times in each  $\tilde{\Lambda}_{i,m}$ . Since the code length  $2K(\tau) + 1 < h_n$  and there are  $p'_{n+1}$  repetitions, we obtain that the coded images of all occurrences of  $(a_{x,y})^{p'_{n+1}}$  are  $\frac{2}{p'_{n+1}}$  close to each other in  $\bar{d}$ . Hence, by making an  $\bar{f}$ -approximation error of at most  $\frac{2}{p'_{n+1}}$  we can decompose  $F_{n,l}(T, P(\tau))$  into permutations  $\Lambda_{i,m}$  as in the Coding Lemma.

In the second case  $r > \tilde{r}$  we decompose  $F_{n,l}(T, P)$  into substitution instances  $\tilde{\Lambda}_{i,m}$  of repetitions of the form

$$[A_x]_1^{qp'_{n+1}}, \quad x = 1, \dots, 2^{4e(n)-t_1}.$$

Within the maximum repetition  $[A_x]_1^{T_2 N^{2r}}$  in  $F_{n,l}(T, P)$ , there are

$$u_2 := \left\lfloor \frac{T_2 N^{2r}}{N q p'_{n+1}} \right\rfloor = \left\lfloor \frac{N^{2(r-\tilde{r})}}{[N \cdot \int f d\mu]} \right\rfloor$$

many such  $\tilde{\Lambda}_{i,1} \dots \tilde{\Lambda}_{i,N}$ . We note that

$$(9.4) \quad 0 \leq \frac{T_2 N^{2r} h_n - u_2 N q p'_{n+1} h_n}{T_2 N^{2r} h_n} \leq \frac{N q p'_{n+1} h_n}{T_2 N^{2r} h_n} \leq \frac{\int f d\mu}{N^{2(r-\tilde{r})-1}} \leq \frac{\int f d\mu}{N}.$$

Lemma 45 and Remark 51 yield again that for each substitution instance  $a_{x,y}$  of  $[A_x]_s$  the repetition  $(a_{x,y})^{p'_{n+1}}$  occurs the same number of times in each  $\tilde{\Lambda}_{i,m}$ . Again, we can decompose  $F_{n,l}(T, P(\tau))$  into permutations  $\Lambda_{i,m}$  as in the Coding Lemma by making an  $\bar{f}$ -approximation error of at most  $\frac{2}{p'_{n+1}}$ .

In both cases, we let  $A = \Lambda_{1,1} \dots \Lambda_{1,N} \Lambda_{2,1} \dots \Lambda_{2,N} \dots \Lambda_{z,1} \dots \Lambda_{z,N}$  and  $B = \Upsilon_{1,1} \dots \Upsilon_{1,N} \Upsilon_{2,1} \dots \Upsilon_{2,N} \dots \Upsilon_{z,1} \dots \Upsilon_{z,N}$  be the strings as in the Coding Lemma. Then the Coding Lemma yields with the aid of (9.2) that

$$\bar{f}(\bar{A}, \bar{B}) > \frac{9\alpha_1}{10 \cdot \int f d\mu} \cdot \left( \frac{1}{8} - \frac{2}{2^{e(n)-0.25t_1}} \right) - \frac{1}{p_{n+1}} - \frac{4 \cdot \int f d\mu}{N}$$

on any substrings  $\bar{A}$  and  $\bar{B}$  of at least  $zqL$  consecutive symbols in  $A$  and  $B$ , respectively. Since  $zqL \leq \delta f_n$  by condition (n1), we conclude

$$(9.5) \quad \begin{aligned} & \bar{f}(I_i(T, \Phi^{-1}(P^f)), I_i(T, P(\tau))) \\ & > \frac{\alpha_1}{10 \cdot \int f d\mu} - \frac{1}{p_{n+1}} - \frac{6 \cdot \int f d\mu}{N} - \frac{2}{N} - \frac{2}{p'_{n+1}} - 4\delta \\ & > \frac{\alpha_1}{20 \cdot \int f d\mu}, \end{aligned}$$

where we used conditions (n1) and (n2) in the last step.  $\square$

In the second step, we show that the 1-pattern types do not agree with each other very often. The proof uses the fact that on average patterns in the  $T^f$  trajectory are expanded by a factor close to  $\int f d\mu$  and thus only about  $\frac{1}{\int f d\mu}$  of the patterns are of the same type.

**Lemma 63.** *For almost every point  $x$  the set of indices in  $I_i = F_{n,l} \cap \tilde{F}_{n,\tilde{l}}$ , for which the 1-pattern types of  $F_{n,l}$  and  $\tilde{F}_{n,\tilde{l}}$  are the same or  $|I_i| < \delta f_n$ , has density at most*

$$\frac{1}{\int f d\mu} + 4\delta.$$

*Proof.* We decompose  $a(T, \Phi^{-1}(P^f), x)$  into its  $(T, \Phi^{-1}(P^f))$ - $(n+1)$ -blocks

$$a(T, \Phi^{-1}(P^f), x) = \dots \tilde{w}_{n+1,-2} \tilde{w}_{n+1,-1} \tilde{w}_{n+1,0} \tilde{w}_{n+1,1} \tilde{w}_{n+1,2} \dots$$

We consider its  $j$ -th word  $\tilde{w}_{n+1,j}$  and denote the corresponding sets of indices by  $\tilde{B}_{n+1,j}$  (that is,  $\tilde{B}_{n+1,j}(T, \Phi^{-1}(P^f), x) = \tilde{w}_{n+1,j}$ ). Then we further subdivide  $\tilde{w}_{n+1,j}$  into its 1-patterns. We denote those by  $\tilde{F}_{n,j,k}$  and their set of indices by  $\tilde{B}_{n+1,j,k}$ . By condition (n4) for almost every  $x$  for all but a set of indices of density at most  $\delta$  the occurrent word  $\tilde{w}_{n+1,j}$  is such that the subset of its indices  $\tilde{B}_{n+1,j}$  lying in a too long 1-pattern (that is,  $|\tilde{B}_{n+1,j,k}| > (\int f d\mu + \delta^3) \cdot |f_n|$ ) or in a too short 1-pattern ( $|\tilde{B}_{n+1,j,k}| < (\int f d\mu - \delta^3) \cdot |f_n|$ ) has density at most  $\delta$ .

Hence, neglecting a set of indices with density at most  $2\delta$  we can assume that  $(\int f d\mu - \delta^3) \cdot |f_n| < |\tilde{B}_{n+1,j,k}| < (\int f d\mu + \delta^3) \cdot |f_n|$ . Since  $U_2^{(n+1)} > \int f d\mu + \delta^3$  by our assumption (n2), there is at most one 1-pattern  $F_{n,l}$  in  $\tilde{B}_{n+1,j,k}(T, P)$  that has the same 1-pattern type as  $\tilde{F}_{n,j,k}$ . Thus, in  $\tilde{B}_{n+1,j,k}(T, P)$  the set of indices that lie in the same 1-pattern type as  $\tilde{F}_{n,j,k}$  has density at most

$$\frac{|f_n|}{|\tilde{B}_{n+1,j,k}|} < \frac{1}{\int f d\mu - \delta^3} < \frac{1}{\int f d\mu} + \delta^3.$$

Taking the neglected indices into account the lemma follows.  $\square$

Combining both results yields the proof of Proposition 56.

*Proof of Proposition 56.* For almost every  $x \in X$  we obtain from Lemma 62 and Lemma 63 that

$$\begin{aligned} & \bar{d}(a(T, \Phi^{-1}(P^f), x), a(T, P(\tau), x)) \\ & \geq \frac{\alpha_1}{20 \cdot \int f d\mu} \cdot \left(1 - \frac{1}{\int f d\mu} - 10\delta\right), \end{aligned}$$

where we also took into account the indices of density  $\delta$  and  $2\epsilon_{n+1} < 2\delta$  that we neglected at the very beginning. Letting  $\delta \rightarrow 0$  and  $n \rightarrow \infty$  we conclude

$$\bar{d}(a(T, \Phi^{-1}(P^f), x), a(T, P(\tau), x)) \geq \frac{\alpha_1}{20 \cdot \int f d\mu} \cdot \frac{\int f d\mu - 1}{\int f d\mu}.$$

For nontrivial  $f$  and  $\tau < \frac{\alpha_1}{20 \cdot \int f d\mu} \cdot \frac{\int f d\mu - 1}{\int f d\mu}$  this contradicts the assumption that  $P(\tau)$  is a finite code with  $d(\Phi^{-1}(P^f), P(\tau)) < \tau$ .  $\square$

**9.2. On even Kakutani equivalence.** In this subsection we state several properties of even Kakutani equivalence. We start by recalling its definition.

**Definition 64.** Two transformations  $S : (X, \mu) \rightarrow (X, \mu)$  and  $T : (Y, \nu) \rightarrow (Y, \nu)$  are said to be *evenly equivalent* if there are subsets  $A \subseteq X$  and  $B \subseteq Y$  of equal measure  $\mu(A) = \nu(B) > 0$  such that  $S_A$  and  $T_B$  are isomorphic to each other. Equivalently, there is  $A \subseteq X$  and  $f : A \rightarrow \mathbb{Z}^+$  with  $\mu(A) \cdot \int_A f d\mu_A = 1$  such that  $(S_A)^f \cong T$ .

*Remark.* We also present another equivalent definition from [FR92]. For this purpose, we define  $\mathcal{O}_S = \{(x_1, x_2) \in X \times X : \exists n \in \mathbb{Z} \text{ such that } S^n(x_1) = x_2\}$  and a map  $v_S : \mathcal{O}_S \rightarrow \mathbb{Z}$  by  $v_S(x, S^n(x)) = n$ . Then  $S$  and  $T$  are evenly equivalent if there is a measure-preserving orbit equivalence  $\phi : X \rightarrow Y$  and a set  $A \subseteq X$  of positive measure such that for all  $x_1 \neq x_2$  with  $(x_1, x_2) \in (A \times A) \cap \mathcal{O}_S$ ,  $v_S(x_1, x_2)$  and  $v_T(\phi(x_1), \phi(x_2))$  have the same sign.

The following lemma from [ORW82] is related to the second characterization in Definition 64.

**Lemma 65.** *Suppose  $S : (X, \mu) \rightarrow (X, \mu)$  is an ergodic measure-preserving automorphism.*

- (a) *If  $f, g : X \rightarrow \mathbb{Z}^+$  such that  $\int g d\mu < \int f d\mu$ , then there exists  $h : X^g \rightarrow \mathbb{Z}^+$  such that  $\int_{X^g} h d\mu^g = (\int f d\mu) / (\int g d\mu)$  and  $(S^g)^h \cong S^f$ .*
- (b) *If  $f : A \rightarrow \mathbb{Z}^+$  and  $\mu(A) \int_A f d\mu_A < 1$ , then there exists  $A' \subset X$  with  $\mu(A') = \mu(A) \int_A f d\mu_A$  such that  $(S_A)^f \cong S_{A'}$ .*



(c) If  $f : A \rightarrow \mathbb{Z}^+$  and  $\mu(A) \int_A f d\mu_A > 1$ , then there exists  $h : X \rightarrow \mathbb{Z}^+$  with  $\int h d\mu = \mu(A) \int_A f d\mu_A$  such that  $(S_A)^f \cong S^h$ .

*Proof.* Part (a) follows from [ORW82, Lemma 1.3, p. 3] and its proof. Parts (b) and (c) are stated on p. 91 of [ORW82], and they follow easily from (a) and the observation that  $S \cong (S_A)^{r_A}$  and  $\int_A r_A d\mu_A = 1/\mu(A)$ .  $\square$

**Lemma 66.** *If  $U, V : (X, \mu) \rightarrow (X, \mu)$  are ergodic measure-preserving automorphisms and  $U_A \cong V_B$  with  $0 < \mu(A) < \mu(B) \leq 1$ , then there exists  $C$  with  $0 < \mu(C) < 1$  such that  $U_C \cong V$ .*

*Proof.* Since  $\mu(A) < \mu(B)$ ,  $\int_A r_A d\mu_A > \int_B r_B d\mu_B$ . Then  $U \cong (U_A)^{r_A} \cong (V_B)^{\tilde{r}}$  where  $\int_B \tilde{r} d\mu_B = \int_A r_A d\mu_A > \int_B r_B d\mu_B = 1/\mu(B)$ . Therefore by Lemma 65(c),  $(V_B)^{\tilde{r}} \cong V^h$  for some  $h : X \rightarrow \mathbb{Z}^+$  with  $\int h d\mu > 1$ . Therefore  $U \cong V^h$  and consequently there exists  $C$  with  $\mu(C) = 1/(\int h d\mu)$  such that  $U_C \cong V$ .  $\square$

For the following proposition we use the same proof as on p. 90 of [ORW82] for the analogous result with  $T$  replaced by Feldman's example  $J$ .

**Proposition 67.** *Let  $\mathcal{T} \in \mathcal{Trees}$  and  $T = F(\mathcal{T})$ . If  $(T_A)^{-1} \cong T_B$ , then  $\mu(A) = \mu(B)$ . That is, if  $T$  and  $T^{-1}$  are Kakutani equivalent, then they are evenly equivalent.*

*Proof.* Suppose  $(T_A)^{-1} \cong T_B$  where  $\mu(A) \neq \mu(B)$ .

*Case 1.* Suppose  $\mu(A) < \mu(B)$ . Then by Lemma 66, there exists  $C$  with  $0 < \mu(C) < 1$  such that  $(T_C)^{-1} = (T^{-1})_C \cong T$ . This implies  $T_C \cong T^{-1}$  and there exists  $\tilde{C} \subset C$  with  $\mu(\tilde{C})/\mu(C) = \mu(C)$  such that

$$T_{\tilde{C}} = (T_C)_{\tilde{C}} \cong (T^{-1})_C \cong T.$$

Since  $0 < \mu(\tilde{C}) < 1$ , this contradicts Proposition 56.

*Case 2.* Suppose  $\mu(A) > \mu(B)$ . Then  $(T_B)^{-1} \cong T_A$ , which is the same as Case 1 with  $A$  and  $B$  switched. Again we have a contradiction to Proposition 56.  $\square$

The first inequality in Lemma 68 below is Proposition 3.2 in [ORW82, p. 92]. The second inequality will serve as an approximate ‘‘consistency property’’ for a sequence of finite codes between two evenly equivalent transformations. It follows from the proof of Lemma 1.3 in [ORW82, p. 3].

If we are given a partition  $P$  of the space  $X$  and  $A \subset X$  with  $\mu(A) > 0$ , we use the notation  $P_A^f$  to denote the partition  $\{A^f \setminus A\} \cup \{c \cap A : c \in P\}$  of  $A^f$ . Moreover, if we consider names in the  $(S_A^f, P_A^f)$  and  $(S_A^g, P_A^g)$  processes, we regard  $A^f \setminus A$  and  $A^g \setminus A$  as having different symbols associated to them. In other words, we have a match between the  $x$ - $S_A^f$ - $P_A^f$  name at time  $i$  and the  $x'$ - $S_A^g$ - $P_A^g$  name at time  $j$  if and only if  $T^i x, T^j x' \in c \cap A$  for some  $c \in P$ .

**Lemma 68.** *Let  $S : (X, \mu) \rightarrow (X, \mu)$  and  $T : (Y, \nu) \rightarrow (Y, \nu)$  be ergodic measure-preserving automorphisms,  $Q$  a finite generating partition for  $T$ , and  $P$  a finite partition of  $X$ . Suppose that  $S$  and  $T$  are evenly equivalent. Let  $(\varepsilon_\ell)_{\ell=1}^\infty$  be a decreasing sequence of positive numbers such that  $0 < \varepsilon_\ell < 1$  and  $\lim_{\ell \rightarrow \infty} \varepsilon_\ell = 0$ . Then there exist increasing sequences  $(m_\ell)_{\ell=1}^\infty$  and  $(K_\ell)_{\ell=1}^\infty$  of positive integers, sets  $F_\ell \subset Y$  with  $\nu(F_\ell) > 1 - \varepsilon_\ell$ , and codes  $\phi_\ell$  of length  $2K_\ell + 1$  from  $(T, Q)$ -names to*

$(S, P)$ -names such that for  $y \in F_\ell$  we have the following properties: There exists  $x = x_\ell(y) \in X$  such that for all  $m \geq m_\ell$ , the  $(S$ - $P$ -name of  $x)_0^{m-1}$  occurs with positive frequency in the  $(S, P)$  process,

$$\bar{f}(\phi_\ell((T\text{-}Q\text{-name of } y)_0^{m-1}), (S\text{-}P\text{-name of } x)_0^{m-1}) < \varepsilon_\ell, \text{ for } m \geq m_\ell,$$

and

$$\bar{f}(\phi_\ell((T\text{-}Q\text{-name of } y)_0^{m-1}), \phi_{\ell+1}((T\text{-}Q\text{-name of } y)_0^{m-1})) < \varepsilon_\ell, \text{ for } m \geq m_{\ell+1}.$$

*Proof.* Since  $S$  and  $T$  are evenly equivalent, there are  $\tilde{A}_1 \subseteq X$  and  $\tilde{f}_1 : \tilde{A}_1 \rightarrow \mathbb{Z}^+$  with  $\mu(\tilde{A}_1) \cdot \int_{\tilde{A}_1} \tilde{f}_1 d\mu_{\tilde{A}_1} = 1$  such that  $(S_{\tilde{A}_1})^{\tilde{f}_1} \cong T$ . We choose a set  $A_1$  with  $X \supset A_1 \supset \tilde{A}_1$  and  $\mu(A_1) > 1 - \frac{\varepsilon_1}{10}$ . Since  $((S_{A_1})_{\tilde{A}_1})^{\tilde{f}_1} \cong T$ ,  $S_{A_1}$  and  $T$  are equivalent. We note that

$$\mu_{A_1}(\tilde{A}_1) \cdot \int_{\tilde{A}_1} \tilde{f}_1 d\mu_{\tilde{A}_1} = \frac{\mu(\tilde{A}_1)}{\mu(A_1)} \cdot \int_{\tilde{A}_1} \tilde{f}_1 d\mu_{\tilde{A}_1} = \frac{1}{\mu(A_1)} > 1.$$

Then Lemma 65 (c) yields the existence of  $f_1 : A_1 \rightarrow \mathbb{Z}^+$  with  $(S_{A_1})^{f_1} \cong T$  and  $\int_{A_1} f_1 d\mu_{A_1} = \frac{1}{\mu(A_1)}$ . Let  $\theta_1 : Y \rightarrow (A_1)^{f_1}$  be an isomorphism between  $T$  and  $(S_{A_1})^{f_1}$ .

We let  $(A_\ell)_{\ell=1}^\infty$  be a sequence of sets such that  $A_1 \subset A_2 \subset A_3 \subset \dots \subset X$  with  $\mu(A_\ell) > 1 - \frac{\varepsilon_\ell}{10}$ . As in the above argument for  $A_1$ , there exist functions  $f_\ell : A_\ell \rightarrow \mathbb{Z}^+$  such that  $(S_{A_\ell})^{f_\ell} \cong T$  and  $\int_{A_\ell} f_\ell d\mu_{A_\ell} = \frac{1}{\mu(A_\ell)}$ . We also have  $(S_{A_\ell})^{r_\ell} \cong S$ , where  $r_\ell = r_{A_\ell, S}$  is the return time to  $A_\ell$  under  $S$  and  $\int_{A_\ell} r_\ell d\mu_{A_\ell} = \frac{1}{\mu(A_\ell)}$ .

For  $x \in A_\ell$ , the  $P_{A_\ell}^{f_\ell}$ -name of  $x$  at the  $i$ th time the orbit under  $(S_{A_\ell})^{f_\ell}$  returns to  $A_\ell$  agrees with the  $P_{A_\ell}^{r_\ell}$ -name of  $x$  at the  $i$ th time the orbit under  $(S_{A_\ell})^{r_\ell}$  returns to  $A_\ell$ . There exist  $k_\ell$  and a set  $C_\ell \subset A_\ell$  with  $\mu(C_\ell) > \mu(A_\ell) - \frac{\varepsilon_\ell}{10} > 1 - \frac{\varepsilon_\ell}{5}$  such that for  $x \in C_\ell$  and  $m \geq k_\ell$ , both the  $(S_{A_\ell})^{f_\ell}$  and the  $(S_{A_\ell})^{r_\ell}$  orbits of  $x$  are in  $A_\ell$  for more than  $1 - \frac{\varepsilon_\ell}{5}$  of the times  $0, 1, \dots, m-1$ . Thus, for  $x \in C_\ell$  and  $m \geq k_\ell$ ,

$$(9.6) \quad \bar{f}(((S_{A_\ell})^{f_\ell}\text{-}P_{A_\ell}^{f_\ell}\text{-name of } x)_0^{m-1}, (S\text{-}P\text{-name of } x)_0^{m-1}) < \frac{\varepsilon_\ell}{5}.$$

We may assume that any such  $S$ - $P$ -name of length  $m$  occurs with positive probability in the  $(S, P)$ -process.

Although we already know that  $(S_{A_\ell})^{f_\ell}$  and  $(S_{A_{\ell+1}})^{f_{\ell+1}}$  are isomorphic, since they are both isomorphic to  $T$ , the proof of the second  $\bar{f}$  inequality in the lemma will be based on a particular isomorphism  $\theta_{\ell+1} : (A_\ell)^{f_\ell} \rightarrow (A_{\ell+1})^{f_{\ell+1}}$  between  $(S_{A_\ell})^{f_\ell}$  and  $(S_{A_{\ell+1}})^{f_{\ell+1}}$ . To simplify notation for this part of the argument, we let  $A = A_\ell$ ,  $A' = A_{\ell+1}$ ,  $f = f_\ell$ ,  $\varepsilon = \varepsilon_\ell$ , and  $\varepsilon' = \varepsilon_{\ell+1}$ . Then  $A \subset A' \subset X$ ,  $\mu(A) > 1 - \frac{\varepsilon}{10}$ , and  $\mu(A') > 1 - \frac{\varepsilon'}{10}$ . The function  $f'$  constructed below is the function that we take as  $f_{\ell+1}$ .

We can write  $S_{A'} = (S_A)^g$ , where  $g : A \rightarrow \mathbb{Z}^+$  is the return time to  $A$  under  $S_{A'}$ . Then  $g$  satisfies  $\int_A g d\mu_A = (\int_A g d\mu) / (\mu(A)) = \mu(A') / \mu(A)$ . Since  $\int_A f d\mu_A = 1 / \mu(A) > \mu(A') / \mu(A)$ , we have  $\int_A f d\mu_A > \int_A g d\mu_A$ . By the ergodic theorem, for almost every  $x$  there exists  $N(x)$  such that

$$\sum_{i=0}^{n-1} f((S_A)^i(x)) > \sum_{i=0}^{n-1} g((S_A)^i(x)), \text{ for all } n \geq N(x).$$

Thus, we can choose a positive measure set  $B \subset A$  and  $N \in \mathbb{Z}^+$  such that  $N(x) \leq N$  for all  $x \in B$ , and the return time  $r_{B,S_A}$  to  $B$  under  $S_A$  satisfies  $r_{B,S_A}(x) \geq N$  for all  $x \in B$ . It follows that

$$\sum_{i=0}^{r_{B,S_A}(x)-1} f((S_A)^i(x)) > \sum_{i=0}^{r_{B,S_A}(x)-1} g((S_A)^i(x)) \text{ for all } x \in B.$$

Now  $(S_A)^f = (S_B^{r_{B,S_A}})^f$ , which can be visualized as a three-dimensional tower, with the two-dimensional base,  $A$ , being represented as a tower for  $S_A$  over  $B$  with roof function  $r_{B,S_A}$ , and then, in the third direction, using the roof function  $f : A \rightarrow \mathbb{Z}$  with base function  $S_A$ . This can also be represented as a two-dimensional tower over  $B$  with height function  $\sum_{i=0}^{r_{B,S_A}(x)-1} f((S_A)^i(x))$ . In the three-dimensional tower, we will denote points by  $(y, \ell)$ , where  $y \in A$  and  $\ell$  is the level in the third direction. That is, the orbit under  $(S_A)^f$  for  $x \in B$  until the time right before it returns to  $B$  is:

$$(9.7) \quad \left( \left( (S_A)^j(x), i \right)_{i=1}^{f((S_A)^j(x))} \right)_{j=0}^{r_{B,S_A}(x)-1}.$$

The part of this orbit that corresponds to points in  $A$  is

$$(9.8) \quad (x, 1), (S_A(x), 1), ((S_A)^2(x), 1), \dots, ((S_A)^{r_{B,S_A}(x)-1}(x), 1).$$

Similarly,  $S_{A'} = (S_A)^g = (S_B^{r_{B,S_A}})^g$  can be represented as a three-dimensional tower with the same two-dimensional base,  $A$ . In the third direction, the roof function is  $g$ . We use the same coordinates  $(y, \ell)$  as before. We define  $f' : A' \rightarrow \mathbb{Z}$  by  $f'(y, \ell) = 1 + \sum_{i=0}^{r_{B,S_A}(x)-1} f((S_A)^i(x)) - \sum_{i=0}^{r_{B,S_A}(x)-1} g((S_A)^i(x))$  if  $y = (S_A)^{r_{B,S_A}(x)-1}(x)$  for some  $x \in B$  and  $\ell = g((S_A)^{r_{B,S_A}(x)-1}(x))$ ;  $f'(y, \ell) = 1$  otherwise. Then  $(S_{A'})^{f'} = ((S_A)^g)^{f'} = (S_A)^f$ . The orbit under  $((S_A)^g)^{f'}$  for  $x \in B$  until the time right before it returns to  $B$  is:

$$(9.9) \quad \left( \left( (S_A)^j(x), i \right)_{i=1}^{g((S_A)^j(x))} \right)_{j=0}^{r_{B,S_A}(x)-1};$$

$$\left( (S_A)^{r_{B,S_A}(x)-1}(x), g \left( (S_A)^{r_{B,S_A}(x)-1}(x) \right) + j \right)_{j=1}^{f'(y,\ell)-1},$$

where  $y = (S_A)^{r_{B,S_A}(x)-1}(x)$  and  $\ell = g((S_A)^{r_{B,S_A}(x)-1}(x))$ . The isomorphism  $\theta$  is defined to be the identity on  $B$  and maps orbits of  $(S_A)^f$  to orbits of  $((S_A)^g)^{f'}$ , with points of the orbits remaining in consecutive order. Note that the part of the orbit in (9.9) that corresponds to points in  $A$  is also given by (9.8). At the points in  $A$ , the  $P_A^f$ -name agrees with the  $P_{A'}^{f'}$ -name. Thus, the number of  $\bar{f}$  matches between the  $P_A^f$ -names of the points in (9.7) and the  $P_{A'}^{f'}$ -names of the points in (9.9) is at least the number of terms in (9.8). Therefore, to estimate the  $\bar{f}$  distance between the  $P_A^f$ -name of  $x$  and the  $P_{A'}^{f'}$ -name of  $\theta(x)$  from time 0 to  $m-1$  we divide the orbits into segments that start with a point in  $B$  until right before the next return to  $B$ . The number of  $\bar{f}$  matches that we obtain is at least the number of times that points in the segments are in  $A$ . If  $\ell_0$  is sufficiently large, then the waiting time until the first entry to the set  $B$  is less than  $\ell_0$  with high probability, and the last entry to the set  $B$  occurs after  $m - \ell_0$  with high probability. Thus there exist  $m_0 \gg \ell_0$ , and a set  $D \subset A$  with  $\mu(D) > 1 - \frac{\varepsilon}{5}$  such that for  $x \in D$  and  $m \geq m_0$  the  $\bar{f}$  distance between the  $P_A^f$ -name of  $x$  and the  $P_{A'}^{f'}$ -name of  $\theta(x)$  from time 0 to  $m-1$  is less than  $1 - \mu(A) + \frac{\varepsilon}{10} < \frac{\varepsilon}{5}$ .

In our inductive argument, if we are given  $f_1, \dots, f_\ell$  and isomorphisms  $\theta_{j+1} : (A_j)^{f_j} \rightarrow (A_{j+1})^{f_{j+1}}$  between  $(S_{A_j})^{f_j}$  and  $(S_{A_{j+1}})^{f_{j+1}}$ , for  $j = 1, \dots, \ell - 1$ , then we let  $f_{\ell+1} = f'$ ,  $\theta_{\ell+1} = \theta$ ,  $u_\ell = m_0$ , and  $D_\ell = D \subset A_\ell$  be as constructed above. Thus  $\mu(D_\ell) > 1 - \frac{\varepsilon_\ell}{5}$  and for  $x \in D_\ell$  and  $m \geq u_\ell$  we have

$$(9.10) \quad \bar{f}_m(((S_{A_\ell})^{f_\ell} - P_{A_\ell}^{f_\ell}\text{-name of } x)_0^{m-1}, ((S_{A_{\ell+1}})^{f_{\ell+1}} - P_{A_{\ell+1}}^{f_{\ell+1}}\text{-name of } \theta_{\ell+1}(x))_0^{m-1}) < \frac{\varepsilon_\ell}{5}.$$

Since  $Q$  is a generator for  $T$  there exists  $K_\ell$  such that

$$(\theta_n \circ \dots \circ \theta_1)^{-1}(P_{A_\ell}^{f_\ell}) \subset_{-K_\ell}^{\varepsilon_\ell/10} \sqrt{-K_\ell} T^j Q,$$

and this defines a code  $\phi_\ell$  from  $(T, Q)$ -names to  $P_{A_\ell}^{f_\ell}$ -names. If  $m_\ell$  is sufficiently large, in particular,  $m_\ell \geq \max(k_\ell, u_\ell)$ , then there exists a set  $E_\ell \subset ((\theta_\ell \circ \dots \circ \theta_1)^{-1}(A_\ell))$  such that for  $y \in E_\ell$  the coded name  $\phi_\ell(y)$  at time  $j$  agrees with the actual  $(\theta_\ell \circ \dots \circ \theta_1)^{-1}(P_{A_\ell}^{f_\ell})$ -name of  $y$ , or equivalently, the  $(P_{A_\ell}^{f_\ell})$ -name of  $(\theta_\ell \circ \dots \circ \theta_1)(y)$ , at time  $j$ , with frequency greater than  $1 - \frac{\varepsilon_\ell}{5}$  among the times  $j = 0, 1, \dots, m - 1$ , for  $m \geq m_\ell$ . That is,

$$(9.11) \quad \bar{d}_m((\phi_\ell(y))_0^{m-1}, (P_{A_\ell}^{f_\ell}\text{-name of } (\theta_\ell \circ \dots \circ \theta_1)(y))_0^{m-1}) < \frac{\varepsilon_\ell}{5}.$$

Finally, we let  $F_\ell = E_\ell \cap E_{\ell+1} \cap (\theta_\ell \circ \dots \circ \theta_1)^{-1}(C_\ell \cap D_\ell) \cap (\theta_{\ell+1} \circ \dots \circ \theta_1)^{-1}(C_{\ell+1})$ . The first  $\bar{f}$  inequality in the lemma follows from (9.6) and (9.11), while the second  $\bar{f}$  inequality follows from (9.6) and (9.11) for  $\ell$  and  $\ell + 1$ , and (9.10) for  $\ell$ .  $\square$

*Remark 69.* We call such a sequence of codes as in Lemma 68 a sequence of  $(\varepsilon_\ell, K_\ell)$ -finite  $\bar{f}$  codes from  $(T, Q)$  to  $(S, P)$ .

**Corollary 70.** *Let  $0 < \gamma < 1/2$  and assume the set-up of Lemma 68, where  $T = \Phi(\mathcal{T})$  for some  $\mathcal{T} \in \mathcal{Trees}$  and  $Q$  is the partition of  $Y = \Sigma^{\mathbb{Z}}$  according to the symbol of  $\Sigma$  located at position 0. We also let  $X = \Sigma^{\mathbb{Z}}$  and  $P = Q$ , but the measures  $\mu$  and  $\nu$  may be different. Assume that the  $\varepsilon_\ell$  in Lemma 68 are chosen so that  $\sum_{i=\ell+1}^{\infty} \varepsilon_i < \varepsilon_\ell$ . Then there exists  $N(\gamma) \in \mathbb{Z}^+$  such that for  $N \geq N(\gamma)$  and  $k \in \mathbb{Z}^+$  and all sufficiently large  $n$  (how large depends on  $N, \gamma$ , and  $k$ ) there is a collection  $\mathcal{W}'_n \subset \mathcal{W}_n$  that includes at least  $1 - \gamma$  of the words in  $\mathcal{W}_n$  and the following condition holds: For  $w \in \mathcal{W}'_n$  there exists  $z = z(w) \in X$  such that the  $(S\text{-}Q\text{-name of } z)_0^{h_n-1}$  occurs with positive frequency in the  $(S, Q)$  process and we have*

$$\bar{f}((\phi_N(w))_0^{h_n-1}, (S\text{-}Q\text{-name of } z)_0^{h_n-1}) < \gamma,$$

and

$$\bar{f}((\phi_N(w))_0^{h_n-1}, (\phi_{N+k}(w))_0^{h_n-1}) < \gamma.$$

*Proof.* Choose  $N(\gamma)$  sufficiently large that  $\varepsilon_{N(\gamma)-1} < \gamma^2/2$ . Let  $N \geq N(\gamma)$ . Then choose  $n$  sufficiently large so  $m_{N+k} < h_n/2$ . Let  $F = F_N \cap \dots \cap F_{N+k-1}$  where  $F_N, \dots, F_{N+k-1}$  are as in Lemma 68. Then  $\nu(F) > 1 - \varepsilon_N - \dots - \varepsilon_{N+k-1} > 1 - \varepsilon_{N-1} > 1 - \gamma^2/2$ . Thus there is a collection  $\mathcal{W}'_n \subset \mathcal{W}_n$  that includes at least  $1 - \gamma$  of the words in  $\mathcal{W}_n$  such that for  $w \in \mathcal{W}'_n$ , for more than  $1 - \gamma/2$  of the indices  $i = 1, 2, \dots, h_n$ , the  $i$ th level of column in the tower representation of the word  $w$  intersects  $F$ . For  $w = a_1 a_2 \dots a_{h_n} \in \mathcal{W}'_n$ , there exists  $i_0 = i_0(w)$  with  $1 \leq i_0 \leq (\gamma/2)h_n$  such that for some  $y = y(w) \in F$  we have  $a_{i_0} \dots a_{h_n} = (T\text{-}Q\text{-name of } y)_0^{h_n-i_0-1}$ . Let  $x = x(y)$  be as in Lemma 68. Let  $z = z(y) \in X$  be such

that the  $(S\text{-}Q\text{-name of } z)_0^{h_n-1}$  occurs with positive frequency in the  $(S, Q)$  process and the  $(S\text{-}Q\text{-name of } z)_{i_0}^{h_n-1}$  is the same as the  $(S\text{-}Q\text{-name of } x)_0^{h_n-i_0-1}$ . Since

$$\bar{f}(\phi_N(a_{i_0} \cdots a_{h_n}), (S\text{-}Q\text{-name of } x)_0^{h_n-i_0-1}) < \varepsilon_N < \gamma^2$$

and  $i_0 \leq (\gamma/2)h_n$ , it follows that

$$\bar{f}(\phi_N(w), (S\text{-}Q\text{-name of } z)_0^{h_n-1}) < (\gamma/2) + \gamma^2 < \gamma.$$

By repeated application of the second  $\bar{f}$  inequality in Lemma 68, we obtain

$$\bar{f}(\phi_N(a_{i_0} \cdots a_{h_n}), \phi_{N+k}(a_{i_0} \cdots a_{h_n})) < \varepsilon_N + \cdots + \varepsilon_{N+k-1} < \varepsilon_{N-1} < \gamma^2/2.$$

Therefore  $\bar{f}(\phi_N(w), \phi_{N+k}(w)) < (\gamma/2) + \gamma^2 < \gamma$ .  $\square$

**9.3. Proof of part (2) in Proposition 23.** We have seen in Proposition 67 that if  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{T})^{-1}$  are Kakutani equivalent, then it can only be by an even equivalence. Thus, to prove part (2) in Proposition 23 it remains to show that under our assumption that the tree  $\mathcal{T} \in \text{Trees}$  does not have an infinite branch,  $\mathbb{K} := \Psi(\mathcal{T})$  is not evenly equivalent to  $\mathbb{K}^{-1} = \Psi(\mathcal{T})^{-1}$ .

As in Section 9.1 we enumerate the uniquely readable and strongly uniform construction sequence of  $\mathbb{K} = \Psi(\mathcal{T})$  as  $\{\mathcal{W}_n\}_{n \in \mathbb{N}}$ , that is,  $n$ -words are built by concatenating  $(n-1)$ -words. We also remind the reader of the phrase  $s$ -pattern type from Remark 44 as well as Remark 51 and that we can decompose an  $n$ -word  $w \in \mathcal{W}_n$  into its  $s$ -Feldman patterns (if we are in Case 2 of the construction we ignore the strings coming from  $\mathcal{W}^{\dagger\dagger}$ ):  $w = P_{n-1,1} \dots P_{n-1,U_{s+1}}$ . In the subsequent lemma we prove that even under finite coding substantial strings of Feldman patterns of different type cannot be matched well.

**Lemma 71.** *Let  $s \in \mathbb{N}$  and  $\phi$  be a finite code from  $\mathbb{K}$  to  $\text{rev}(\mathbb{K})$ . Then for sufficiently large  $n \in \mathbb{N}$  and any pair of  $s$ -Feldman patterns  $P_{n-1,\ell}$  in  $\mathbb{K}$  and  $\bar{P}_{n-1,\bar{\ell}}$  in  $\text{rev}(\mathbb{K})$  of different pattern type we have*

$$\bar{f}(\phi(E), \bar{E}) > \frac{\alpha_s}{10},$$

for any substrings  $E, \bar{E}$  of at least  $\frac{|P_{n-1,\ell}|}{2^{2e(n-1)}} = \frac{|\bar{P}_{n-1,\bar{\ell}}|}{2^{2e(n-1)}}$  consecutive symbols in  $P_{n-1,\ell}$  and  $\bar{P}_{n-1,\bar{\ell}}$ , respectively.

*Proof.* We recall the definition of  $M(s)$  from Definition 19 and choose  $n > M(s) + 1$  sufficiently large such that  $R_{n-1} > \frac{110}{\alpha_s}$  and  $h_{n-1} > 2^{4e(n-1)} \cdot |\phi|$ . In particular, end effects are negligible and we can assume  $\phi(E)$  and  $E$  to have the same length.

We denote the tuple of building blocks for  $[P_{n-1,\ell}]_s$  by  $([A_1]_s, \dots, [A_{2^{4e(n-1)-t_s}}]_s)$  and the tuple of building blocks for  $[\bar{P}_{n-1,\bar{\ell}}]_s$  by  $([\bar{A}_1]_s, \dots, [\bar{A}_{2^{4e(n-1)-t_s}}]_s)$ . Moreover, let  $r$  and  $\bar{r}$  denote the types of  $s$ -Feldman pattern of  $P_{n-1,\ell}$  and  $\bar{P}_{n-1,\bar{\ell}}$ , respectively. In the notation of the Coding Lemma 41 we let

$$\begin{aligned} L &= h_{n-1}, \\ N &= 2^{4e(n-1)-t_s}, \\ z &= N^{2 \cdot (M_{s+1} + 1 - \bar{r})}, \\ q &= T_{s+1} \cdot 2^{(4e(n-1)-t_s) \cdot 2\bar{r}} = T_{s+1} \cdot N^{2\bar{r}}. \end{aligned}$$

We emphasize that  $T_{s+1}$  and thus  $q$  are multiples of  $p'_n$ , where  $p'_n = p_n^2$  in Case 1 ( $s(n) = s(n-1)$ ) or  $p'_n = p_n^2 - p_n$  in Case 2 ( $s(n) = s(n-1) + 1$ ) of the construction.

Furthermore, each  $\Upsilon_{i,j}$  in the Coding Lemma 41 corresponds to a substitution instance of a maximal repetition  $[\overline{A}_j]_s^q$  in  $\overline{P}_{n-1,\bar{\ell}}$ . Since we have  $\overline{f}(W, W') > \alpha_s$  on any substrings  $W$  and  $W'$  of at least  $\frac{h_{n-1}}{R_{n-1}}$  consecutive symbols in any words  $w, w' \in \mathcal{W}_{n-1}$  with  $[w]_s \neq [w']_s$  by Proposition 55, the assumption of the Coding Lemma is fulfilled. Now we distinguish between the two possible cases.

In the first instance, let  $r < \bar{r}$ , that is, the number  $T_{s+1}2^{(4e(n-1)-t_s)\cdot 2r} = T_{s+1}N^{2r}$  of consecutive repetitions of the same  $s$ -block in  $P_{n-1,\ell}$  is smaller than in  $\overline{P}_{n-1,\bar{\ell}}$ . We subdivide  $P_{n-1,\ell}$  into substitution instances  $\tilde{\Lambda}_{i,j}$  of  $N^{2(\bar{r}-r)-1}$  consecutive complete cycles in  $[P_{n-1,\ell}]_s$ . Note that  $|\tilde{\Lambda}_{i,j}| = |\Lambda_{i,j}|$ . Lemma 45 and Remark 51 yield that for each substitution instance  $a_{x,y}$  of each  $[A_x]_s$ ,  $x = 1, \dots, 2^{4e(n-1)-t_s}$ , the repetition  $(a_{x,y})^{p'_n}$  occurs the same number of times in each  $\tilde{\Lambda}_{i,j}$ . Since  $h_{n-1} > 2^{4e(n-1)} \cdot |\phi|$ , we also see that the images under code  $\phi$  of all occurrences of  $(a_{x,y})^{p'_n}$  are  $\frac{2}{p'_n}$  close to each other in  $\bar{d}$ . Hence, by making a  $\bar{d}$ -approximation error of at most  $\frac{2}{p'_n}$  we can decompose  $\phi(P_{n-1,\ell})$  into permutations  $\Lambda_{i,j}$  as in the Coding Lemma.

In case of  $r > \bar{r}$  we decompose  $P_{n-1,\ell}$  into substitution instances  $\tilde{\Lambda}_{i,j}$  of repetitions of the form  $[A_x]_s^q$ ,  $x = 1, \dots, 2^{4e(n-1)-t_s}$ . Lemma 45 and Remark 51 yield again that for each substitution instance  $a_{x,y}$  of  $[A_x]_s$  the repetition  $(a_{x,y})^{p'_n}$  occurs the same number of times in each  $\tilde{\Lambda}_{i,j}$ . As above, we can decompose  $\phi(P_{n-1,\ell})$  into permutations  $\Lambda_{i,j}$  as in the Coding Lemma by making a  $\bar{d}$ -approximation error of at most  $\frac{2}{p'_n}$ .

In both cases, the Coding Lemma gives

$$\begin{aligned} \overline{f}(\phi(E), \overline{E}) &> \alpha_s \cdot \left( \frac{1}{8} - \frac{2}{2^{e(n-1)-0.25t_s}} \right) - \frac{1}{R_{n-1}} - \frac{4}{N} - \frac{2}{p'_n} \\ &> \frac{\alpha_s - \frac{11}{R_{n-1}}}{9} > \frac{\alpha_s}{10} \end{aligned}$$

for any substrings  $E, \overline{E}$  of at least  $zqL = \frac{|P_{n-1,\ell}|}{N}$  consecutive symbols in  $P_{n-1,\ell}$  and  $\overline{P}_{n-1,\bar{\ell}}$ , respectively. Here, we used conditions (8.5) and (8.7) as well as our assumption  $R_{n-1} > \frac{110}{\alpha_s}$ .  $\square$

**Lemma 72.** *Let  $s \in \mathbb{N}$  and  $\phi$  be a finite code from  $\mathbb{K}$  to  $\text{rev}(\mathbb{K})$ . Then for sufficiently large  $n \in \mathbb{N}$  and any pair  $w \in \mathcal{W}_n$ ,  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  of different  $s$ -pattern type we have*

$$\overline{f}(\phi(A), \overline{A}) > \frac{\alpha_s}{16}$$

for any substrings  $A, \overline{A}$  of at least  $\frac{h_n}{2^{e(n-1)}}$  consecutive symbols in  $w$  and  $\bar{w}$ , respectively.

*Proof.* As before, we choose  $n > M(s) + 1$  sufficiently large such that  $h_{n-1} > 2^{4e(n-1)} \cdot |\phi|$  and end effects are negligible, that is, we can assume  $\phi(A)$  and  $A$  to have the same length.

If  $\overline{f}(\phi(A), \overline{A}) > \frac{\alpha_s}{4}$ , we do not have to show anything. Otherwise, we have  $\overline{f}(\phi(A), \overline{A}) \leq \frac{1}{32}$  (recall that  $\alpha_s < \frac{1}{8}$ ) and let  $\pi$  be a best possible  $\overline{f}$  matching between  $\phi(A)$  and  $\overline{A}$ . Since  $\pi$  is a map on the indices, we can also view it as a map between  $A$  and  $\overline{A}$ . Since  $n > M(s) + 1$  and  $A$  as well as  $\overline{A}$  has length at least  $\frac{h_n}{2^{e(n-1)}}$ , Remark 44 yields that there are at least  $2^{6e(n-1)}$  complete  $s$ -Feldman




FIGURE 9.2. Illustration of the decompositions into  $E_i \subseteq P_{n-1,j(i)}$  and  $\bar{E}_i \subseteq \bar{P}_{n-1,\bar{j}(i)}$  corresponding to each other under  $\pi$ .

patterns (out of building blocks in  $\mathcal{W}_{n-1}/\mathcal{Q}_s^{n-1}$ ) in  $A$  and  $\bar{A}$ . Moreover, all of them have the same length, (If we are in Case 2 of the construction we ignore the strings coming from  $\mathcal{W}^{\dagger\dagger}$ , which might increase the  $\bar{f}$  distance by at most  $\frac{2}{p_n}$  according to (8.16) and Fact 9.) We complete partial  $s$ -patterns at the beginning and end of  $A$  and  $\bar{A}$ , which might increase the  $\bar{f}$  distance by at most  $\frac{2}{2^{6e(n-1)}}$  according to Fact 9 as well. The augmented strings obtained in this way are denoted by  $A_{aug}$  and  $\bar{A}_{aug}$ , respectively, and we decompose them into the  $s$ -Feldman patterns:

$$A_{aug} = P_{n-1,1} \dots P_{n-1,r} \quad \bar{A}_{aug} = \bar{P}_{n-1,1} \dots \bar{P}_{n-1,\bar{r}}.$$

Since  $w$  and  $\bar{w}$  are of different  $s$ -pattern type, the occurring Feldman patterns are either disjoint or traversed in opposite directions.

In the next step, we write  $A_{aug}$  and  $\bar{A}_{aug}$  as

$$A_{aug} = E_1 \dots E_v \quad \bar{A}_{aug} = \bar{E}_1 \dots \bar{E}_v,$$

where  $E_i$  and  $\bar{E}_i$  are maximal substrings such that  $E_i \subseteq P_{n-1,j(i)}$  and  $\bar{E}_i \subseteq \bar{P}_{n-1,\bar{j}(i)}$  correspond to each other under  $\pi$  (see Figure 9.2 for an illustration of this decomposition). In particular, we have  $v \geq 2^{6e(n-1)}$ .

We note that for at most one pair of  $E_i$  and  $\bar{E}_i$  the type of Feldman pattern coincides. We delete this possible pair. This might increase the  $\bar{f}$  distance by at most  $\frac{2}{2^{6e(n-1)}}$  according to Fact 9 because we have  $v \geq 2^{6e(n-1)}$ . Thus, in the following we can assume that all pairs  $E_i$  and  $\bar{E}_i$  lie in Feldman patterns of different type.

If  $\bar{f}(\phi(E_i), \bar{E}_i) > \frac{\alpha_s}{4}$ , we stop the investigation of these strings. Otherwise, we have

$$(9.12) \quad 1 - \frac{1}{2^4} < \frac{1 - \frac{1}{2^5}}{1 + \frac{1}{2^5}} \leq \frac{|E_i|}{|\bar{E}_i|} \leq \frac{1 + \frac{1}{2^5}}{1 - \frac{1}{2^5}} < 1 + \frac{1}{2^3}$$

by Fact 11. In each  $P_{n-1,l}$  there are at most two segments  $E_i$  with  $|E_i| < \frac{|P_{n-1,l}|}{2^{2e(n-1)}}$  because a string well-matched with  $P_{n-1,l}$  has to lie within at most three consecutive patterns in  $\bar{A}_{aug}$ . For their corresponding segments we also get  $|\bar{E}_i| < \frac{|P_{n-1,l}|}{2^{2e(n-1)-1}}$  by the estimate in (9.12). By the same reasons, there are at most two segments

$\bar{E}_i$  with  $|\bar{E}_i| < \frac{|P_{n-1,l}|}{2^{2\varepsilon(n-1)}}$  in each  $\bar{P}_{n-1,l}$  and we can also bound the lengths of their corresponding strings. Altogether, the density of symbols in those situations is at most  $\frac{1}{2^{2\varepsilon(n-1)-4}}$  and we ignore them in the following consideration.

Hence, we consider the case that both  $|E_i|$  and  $|\bar{E}_i|$  are at least  $\frac{|P_{n-1,l}|}{2^{2\varepsilon(n-1)}}$ . This is large enough such that we can apply Lemma 71 and conclude the statement.  $\square$

As a consequence, we deduce that a well-approximating finite code can be identified with an element of the group action, where we recall the notation  $\eta_g$  from Definition 37.

**Lemma 73.** *Let  $s \in \mathbb{N}$ ,  $0 < \delta < \frac{1}{4}$ ,  $0 < \varepsilon < \frac{\alpha_s}{200}\delta$ , and  $\phi$  be a finite code from  $\mathbb{K}$  to  $\text{rev}(\mathbb{K})$ . Then for  $n$  sufficiently large and  $w \in \mathcal{W}_n$  we have that for any string  $\bar{A}$  in  $\text{rev}(\mathbb{K})$  with  $\bar{f}(\phi(w), \bar{A}) < \varepsilon$  there must be exactly one  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  with  $|\bar{A} \cap \bar{w}| \geq (1 - \delta)|\bar{w}|$  and  $\bar{w}$  must be of the form  $[\bar{w}]_s = \eta_g[w]_s$  for a unique  $g \in G_s$ , which is necessarily of odd parity.*

*Proof.* For a start, we choose  $n$  large enough such that end effects are negligible with respect to the length  $h_n$  of  $n$ -words (that is,  $h_n$  has to be much larger than the length of the code). By Fact 11 we know from  $\bar{f}(\phi(w), \bar{A}) < \varepsilon$  that

$$1 - 3\varepsilon \leq \frac{1 - \varepsilon}{1 + \varepsilon} \leq \frac{|\bar{A}|}{|\bar{w}|} \leq \frac{1 + \varepsilon}{1 - \varepsilon} \leq 1 + 3\varepsilon.$$

Hence, there can be at most one  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  with  $|\bar{A} \cap \bar{w}| \geq (1 - \delta)|\bar{w}|$ . Suppose that there is no such  $\bar{w} \in \text{rev}(\mathcal{W}_n)$ . Then there are two words  $\bar{w}_1, \bar{w}_2 \in \text{rev}(\mathcal{W}_n)$  such that  $\bar{A}$  is a substring of  $\bar{w}_1\bar{w}_2$  and

$$|\bar{A} \cap \bar{w}_j| \geq (\delta - 3\varepsilon) \cdot |\bar{w}_j| > \frac{180}{\alpha_s} \varepsilon h_n \text{ for } j = 1, 2.$$

Then the beginning of  $\phi(w)$  is matched with the end of  $\bar{w}_1$ , while the end of  $\phi(w)$  is matched with the beginning of  $\bar{w}_2$ . There are two possible cases: Either at least one of  $\bar{w}_1, \bar{w}_2$  has a different  $s$ -pattern type from that of  $w$ , or  $\bar{w}_1$  and  $\bar{w}_2$  have the same  $s$ -pattern type as  $w$ . Here, we point out that the  $s$ -pattern structure of the beginning and end of  $w$  are different from each other by construction (recall from the general substitution step in Section 7 that substitution instances were constructed as a concatenation of different Feldman patterns). To investigate both cases we denote  $\bar{A}_j := \bar{A} \cap \bar{w}_j$ . Let  $\pi$  be a best possible match in  $\bar{f}$  between  $\phi(w)$  and  $\bar{A}$ . Then we denote the parts in  $\phi(w)$  corresponding to  $\bar{A}_j$  under  $\pi$  by  $A_j$ . If  $\bar{f}(A_j, \bar{A}_j) > \frac{\alpha_s}{4}$ , we stop. Otherwise, we have  $\frac{8}{9} \leq \frac{|A_j|}{|\bar{A}_j|} \leq \frac{9}{8}$ , that is,  $|A_j| \geq \frac{8}{9}|\bar{A}_j| \geq \frac{160}{\alpha_s} \varepsilon h_n$ . Hence, for sufficiently large  $n$  the strings are long enough that we can apply Lemmas 71 and 72. In both aforementioned cases this yields, with the aid of Fact 10, that

$$\bar{f}(\phi(w), \bar{A}) \geq \frac{80}{\alpha_s} \varepsilon \cdot \frac{\alpha_s}{16} = 5\varepsilon,$$

which contradicts the assumption  $\bar{f}(\phi(w), \bar{A}) < \varepsilon$ . Hence, there must be exactly one  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  with  $|\bar{A} \cap \bar{w}| \geq (1 - \delta)|\bar{w}|$ .

Lemma 72 also implies that  $w$  and  $\bar{w}$  need to have the same  $s$ -pattern structure. By construction this only happens if their equivalence classes  $[w]_s$  and  $[\bar{w}]_s$  lie on the same orbit of the group action by  $G_s^n$  (recall from the general substitution step that substitution instances not lying on the same  $G_s^n$  orbit were constructed as



a different concatenation of different Feldman patterns). By the freeness of the group action and Proposition 55, we conclude that there is a unique  $g \in G_s$  with  $[\bar{w}]_s = \eta_g[w]_s$ .  $\square$

Assume  $\mathbb{K}$  and  $\mathbb{K}^{-1}$  are evenly equivalent. Let  $(\varepsilon_\ell)_{\ell \in \mathbb{N}}$  be a sequence of positive reals satisfying  $\sum_{i=\ell+1}^{\infty} \varepsilon_i < \varepsilon_\ell$  for every  $\ell \in \mathbb{N}$ . Then there exists a sequence of  $(\varepsilon_\ell, K_\ell)$ -finite  $\bar{f}$  codes  $\phi_\ell$  from  $\mathbb{K}$  to  $\mathbb{K}^{-1}$  satisfying the properties in Lemma 68. We fix such a sequence of  $(\varepsilon_\ell, K_\ell)$ -finite  $\bar{f}$  codes  $\phi_\ell$  between  $\mathbb{K}$  and  $\mathbb{K}^{-1}$ . By applying Corollary 70 with  $\gamma_s = \frac{\alpha_s^4}{5 \cdot 10^9}$ , for each  $s \in \mathbb{Z}^+$  there is  $N(s) \in \mathbb{Z}^+$  such that for every  $N \geq N(s)$  and  $k \in \mathbb{Z}^+$  there is  $n(s, N, k) \in \mathbb{Z}^+$  sufficiently large so that for all  $n \geq n(s, N, k)$  Lemma 73 with  $\delta_s = \frac{\alpha_s^3}{2 \cdot 10^7}$  and  $\phi_N$  holds. Moreover, there exists a collection  $\mathcal{W}'_n$  of  $n$ -words (that includes at least  $1 - \gamma_s$  of the  $n$ -words) satisfying the following properties:

- (C1) For  $w \in \mathcal{W}'_n$  there exists  $z = z(w) \in \text{rev}(\mathbb{K})$  such that  $z \upharpoonright [0, h_n - 1]$  occurs with positive frequency in  $\text{rev}(\mathbb{K})$  and

$$\bar{f}(\phi_N(w), z \upharpoonright [0, h_n - 1]) < \frac{\alpha_s^4}{5 \cdot 10^9}.$$

- (C2) For  $w \in \mathcal{W}'_n$  we have

$$\bar{f}(\phi_N(w), \phi_{N+k}(w)) < \frac{\alpha_s^4}{5 \cdot 10^9}.$$

**Lemma 74.** *Suppose that  $\mathbb{K}$  and  $\mathbb{K}^{-1}$  are evenly equivalent and let  $s \in \mathbb{N}$ . There is a unique  $g_s \in G_s$  such that for every  $N \geq N(s)$  and sufficiently large  $n \in \mathbb{N}$  we have for every  $w \in \mathcal{W}'_n$  that there is  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  with  $[\bar{w}]_s = \eta_{g_s}[w]_s$  and*

$$\bar{f}(\phi_N(w), \bar{w}) < \frac{\alpha_s}{4},$$

where  $(\phi_\ell)_{\ell \in \mathbb{N}}$  is a sequence of  $(\varepsilon_\ell, K_\ell)$ -finite  $\bar{f}$  codes as described above and  $\mathcal{W}'_n$  is the associated collection of  $n$ -words satisfying properties (C1) and (C2).

Moreover,  $g_s \in G_s$  is of odd parity and the sequence  $(g_s)_{s \in \mathbb{N}}$  satisfies  $g_s = \rho_{s+1, s}(g_{s+1})$  for all  $s \in \mathbb{N}$ .

*Proof.* For  $n \geq n(s, N, 1)$  we see from property (C1) that for any  $w \in \mathcal{W}'_n$  there exists  $z \in \text{rev}(\mathbb{K})$  with

$$\bar{f}(\phi_N(w), z \upharpoonright [0, h_n - 1]) < \gamma_s = \frac{\alpha_s^4}{5 \cdot 10^9} < \frac{\alpha_s}{200} \cdot \delta_s.$$

We denote  $\bar{A} := z \upharpoonright [0, h_n - 1]$  in  $\text{rev}(\mathbb{K})$ . By Lemma 73 there is exactly one  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  with  $|\bar{A} \cap \bar{w}| \geq (1 - \delta_s)|\bar{w}|$  and  $\bar{w}$  must be of the form  $[\bar{w}]_s = \eta_g[w]_s$  for a unique  $g \in G_s$ , which is of odd parity. By Fact 9 we conclude  $\bar{f}(\phi_N(w), \bar{w}) < \gamma_s + \delta_s < 10^{-7}\alpha_s^3$ .

To see that one group element of  $G_s$  is supposed to work for  $\phi_N$  and all words in  $\mathcal{W}'_{n+1}$  we repeat the argument for a word  $w \in \mathcal{W}'_{n+1}$ , which is a concatenation of  $n$ -words by construction. Hence, we obtain a unique  $g_s \in G_s$  and  $\bar{w} \in \text{rev}(\mathcal{W}_{n+1})$  with  $[\bar{w}]_s = \eta_{g_s}[w]_s$  such that  $\bar{f}(\phi_N(w), \bar{w}) < 10^{-7}\alpha_s^3$ . We decompose  $w$  into its  $s$ -Feldman patterns:  $w = P_1 \dots P_r$ . Moreover, let  $\bar{w} = \bar{P}_1 \dots \bar{P}_r$  be the corresponding decomposition under a best possible  $\bar{f}$ -match between  $\phi_N(w)$  and  $\bar{w}$ . According to Fact 10 the proportion of such substrings with  $\bar{f}(\phi_N(P_i), \bar{P}_i) > 10^{-6}\alpha_s^3$  could be at most  $\frac{1}{10}$ . On the other substrings with  $\bar{f}(\phi_N(P_i), \bar{P}_i) \leq 10^{-6}\alpha_s^3$  we need to have

$\left(1 - \frac{2\alpha_s^3}{10^6}\right) |P_i| \leq |\bar{P}_i| \leq \left(1 + \frac{3\alpha_s^3}{10^6}\right) |P_i|$  by Fact 11. Then we obtain from Lemma 71 that there is a substring  $\bar{P}'_i \subseteq \bar{P}_i$  with  $|\bar{P}'_i| \geq \left(1 - \frac{\alpha_s^2}{5 \cdot 10^4}\right) |\bar{P}_i|$  such that  $\bar{P}'_i$  and  $P_i$  have the same  $s$ -pattern structure, that is,  $[\bar{P}'_i]_s$  must lie in  $\eta_{g_s}[P_i]_s$ . We deduce from  $\bar{f}(\phi_N(P_i), \bar{P}'_i) < \frac{\alpha_s^2}{4 \cdot 10^4}$  that a proportion of at least  $\frac{9}{10}$  of the  $n$ -blocks  $w_l$  occurring as substitution instances in  $P_i$  have to satisfy that there is a string  $\bar{A}_l$  in  $\bar{P}'_i$  such that  $\bar{f}(\phi_N(w_l), \bar{A}_l) < \frac{\alpha_s^2}{2000} = \frac{\alpha_s}{200} \cdot \frac{\alpha_s}{10}$ . As in the first paragraph of the proof, Lemma 73 implies that there is exactly one  $\bar{w}_l \in \text{rev}(\mathcal{W}_n)$  with  $|\bar{A}_l \cap \bar{w}_l| \geq (1 - \frac{\alpha_s}{10})|\bar{w}_l|$  and  $\bar{w}_l$  must be of the form  $[\bar{w}_l]_s = \eta_{h_l}[w_l]_s$  for a unique  $h_l \in G_s$ , which is of odd parity. As above, we also get  $\bar{f}(\phi_N(w_l), \bar{w}_l) < \frac{\alpha_s}{4}$ . Since the classes within the building tuple of  $P_i$  have disjoint  $G_s$  orbits, we obtain that a proportion of at least  $\frac{9}{10}$  of the  $n$ -blocks  $w_l$  occurring as substitution instances in  $P_i$  have to satisfy that there is  $\bar{w}_l \in \text{rev}(\mathcal{W}_n)$  with  $[\bar{w}_l]_s = \eta_{g_s}[w_l]_s$  and  $\bar{f}(\phi_N(w_l), \bar{w}_l) < \frac{\alpha_s}{4}$ . Altogether, we conclude that a proportion of at least  $\frac{8}{10}$  of all  $n$ -blocks  $w_l$  have to satisfy that there is  $\bar{w}_l \in \text{rev}(\mathcal{W}_n)$  with  $[\bar{w}_l]_s = \eta_{g_s}[w_l]_s$  and  $\bar{f}(\phi_N(w_l), \bar{w}_l) < \frac{\alpha_s}{4}$ . Suppose now that there is a word  $w' \in \mathcal{W}'_{n+1}$  such that the argument above gives a different element  $h \in G_s$  and  $\bar{w}' \in \text{rev}(\mathcal{W}_{n+1})$  with  $[\bar{w}']_s = \eta_h[w']_s$  and  $\bar{f}(\phi_N(w'), \bar{w}') < \frac{\alpha_s}{4}$ . Then a proportion of at least  $\frac{8}{10}$  of all  $n$ -blocks  $w_l$  would have to satisfy that there is  $\bar{w}_l \in \text{rev}(\mathcal{W}_n)$  with  $[\bar{w}_l]_s = \eta_h[w_l]_s$  and  $\bar{f}(\phi_N(w_l), \bar{w}_l) < \frac{\alpha_s}{4}$ . This contradicts  $\eta_{g_s}[w_l]_s \neq \eta_h[w_l]_s$  and Proposition 55. Hence, there is a unique group element  $g_s \in G_s$  working for all  $(n+1)$ -words in  $\mathcal{W}'_{n+1}$ .

In the next step, we check that the same group element  $g_s \in G_s$  works for all  $\phi_N$ ,  $N \geq N(s)$ . Let  $k \in \mathbb{Z}^+$ . For  $n \geq n(s, N, k)$  property (C2) guarantees  $\bar{f}(\phi_N(w), \phi_{N+k}(w)) < \alpha_s^4$  for every  $w \in \mathcal{W}'_n$ . On the one hand, let  $g_s$  be the element in  $G_s$  such that for every  $w \in \mathcal{W}'_n$  there is  $\bar{w} \in \text{rev}(\mathcal{W}_n)$  with  $[\bar{w}]_s = \eta_{g_s}[w]_s$  and  $\bar{f}(\phi_N(w), \bar{w}) < \frac{\alpha_s}{4}$ . On the other hand, let  $h_s \in G_s$  such that for every  $w \in \mathcal{W}'_n$  there is  $w' \in \text{rev}(\mathcal{W}_n)$  with  $[w']_s = \eta_{h_s}[w]_s$  and  $\bar{f}(\phi_{N+k}(w), w') < \frac{\alpha_s}{4}$ . Assume  $g_s \neq h_s$ . Then we have  $[w']_s = \eta_{h_s}[w]_s \neq \eta_{g_s}[w]_s = [\bar{w}]_s$  but

$$\bar{f}(\bar{w}, w') \leq \bar{f}(\bar{w}, \phi_N(w)) + \bar{f}(\phi_N(w), \phi_{N+k}(w)) + \bar{f}(\phi_{N+k}(w), w') < \frac{2}{3}\alpha_s,$$

which contradicts Proposition 55. Thus, we conclude that  $g_s \in G_s$  works for all  $\phi_N$ , for  $N \geq N(s)$ .

Suppose that for some  $s \in \mathbb{N}$  we have that  $g'_s := \rho_{s+1, s}(g_{s+1}) \neq g_s$ . Let  $N \geq N(s+1)$ . Then for  $n$  sufficiently large there is  $w \in \mathcal{W}'_n$  with  $\eta_{g'_s}[w]_s \neq \eta_{g_s}[w]_s = [\bar{w}_1]_s$ , where  $\bar{w}_1 \in \text{rev}(\mathcal{W}_n)$  is the element corresponding to  $w$  under the code  $\phi_N$  with  $\bar{f}(\phi_N(w), \bar{w}_1) < \frac{\alpha_s}{4}$ . Similarly, let  $\bar{w}_2 \in \text{rev}(\mathcal{W}_n)$  be the element corresponding to  $w$  under the code  $\phi_N$  with  $\bar{f}(\phi_N(w), \bar{w}_2) < \frac{\alpha_{s+1}}{4}$ . Since the  $G_{s+1}^n$  action is subordinate to the  $G_s^n$  action on  $\mathcal{W}_n/\mathcal{Q}_s^n$  by specification (A7), we see that  $[\bar{w}_2]_{s+1} = \eta_{g_{s+1}}[w]_{s+1}$  lies in  $\eta_{g'_s}[w]_s$ . Hence,  $[\bar{w}_2]_s \neq [\bar{w}_1]_s$ , which implies  $\bar{f}(\bar{w}_1, \bar{w}_2) > \alpha_s$  by Proposition 55. On the other hand, we have

$$\bar{f}(\bar{w}_1, \bar{w}_2) \leq \bar{f}(\bar{w}_1, \phi_N(w)) + \bar{f}(\phi_N(w), \bar{w}_2) < \frac{\alpha_s}{4} + \frac{\alpha_{s+1}}{4} < \frac{\alpha_s}{2}.$$

This contradiction shows that  $(g_s)_{s \in \mathbb{N}}$  must satisfy  $g_s = \rho_{s+1, s}(g_{s+1})$  for all  $s$ .  $\square$

*Proof of part (2) in Proposition 23.* If  $\mathbb{K} = \Psi(\mathcal{T})$  and  $\mathbb{K}^{-1}$  are Kakutani equivalent, then they have to be evenly equivalent by Proposition 67. Then the conclusion of Lemma 74 implies that the tree  $\mathcal{T}$  has an infinite branch.  $\square$

## 10. TRANSFER TO THE SMOOTH AND REAL-ANALYTIC CATEGORIES

In this final section we transfer our anti-classification result for Kakutani equivalence to the smooth and real-analytic categories. For this purpose, we review the definition of circular systems. These symbolic systems can be realized by the Approximation by Conjugation method as  $C^\infty$  diffeomorphisms on compact surfaces admitting a nontrivial circle action [FW19a] and as real-analytic diffeomorphisms on  $\mathbb{T}^2$  ([Ba17], [BKpp]). We also review the definition of the functor  $\mathcal{F}$  between odometer-based and circular systems introduced in [FW19b]. To simplify notation we enumerate construction sequences of our odometer-based systems as  $\{\mathcal{W}_n\}_{n \in \mathbb{N}}$ , that is,  $(n+1)$ -words are built by concatenating  $n$ -words. This functor  $\mathcal{F}$  preserves so-called synchronous and anti-synchronous isomorphisms [FW19b] but it does not necessarily preserve Kakutani equivalence as shown by examples in [GKpp]. In the setting of our constructions from Section 8 we can prove using the techniques from Section 9 that if  $\mathcal{T} \in \mathcal{Trees}$  does not have an infinite branch, then  $\mathcal{F}(\Psi(\mathcal{T}))$  and  $\mathcal{F}(\Psi(\mathcal{T})^{-1})$  are not Kakutani equivalent to each other. This will allow us to conclude the proofs of Theorems 24 and 25 in Subsection 10.6.

**10.1. Diffeomorphism spaces.** Let  $M$  be a smooth compact manifold of finite dimension equipped with a standard measure  $\mu$  determined by a smooth volume element. We denote the space of  $\mu$ -preserving  $C^\infty$  diffeomorphisms on  $M$  by  $\text{Diff}^\infty(M, \mu)$ , which inherits a Polish topology.

Following [BKpp, section 2.3] we give a more detailed description of the space of Lebesgue measure-preserving real-analytic diffeomorphisms on  $\mathbb{T}^2 := \mathbb{R}^2/\mathbb{Z}^2$  that are homotopic to the identity. Here,  $\lambda$  denotes the standard Lebesgue measure on  $\mathbb{T}^2$ .

Any real-analytic diffeomorphism on  $\mathbb{T}^2$  homotopic to the identity admits a lift to a map from  $\mathbb{R}^2$  to  $\mathbb{R}^2$  which has the form

$$F(x_1, x_2) = (x_1 + f_1(x_1, x_2), x_2 + f_2(x_1, x_2)),$$

where  $f_i : \mathbb{R}^2 \rightarrow \mathbb{R}$  are  $\mathbb{Z}^2$ -periodic real-analytic functions. Any real-analytic  $\mathbb{Z}^2$ -periodic function on  $\mathbb{R}^2$  can be extended as a holomorphic function defined on some open complex neighborhood of  $\mathbb{R}^2$  in  $\mathbb{C}^2$ , where we identify  $\mathbb{R}^2$  inside  $\mathbb{C}^2$  via the natural embedding  $(x_1, x_2) \mapsto (x_1 + i0, x_2 + i0)$ . For a fixed  $\rho > 0$  we define the neighborhood

$$\Omega_\rho := \{(z_1, z_2) \in \mathbb{C}^2 : |\text{Im}(z_1)| < \rho \text{ and } |\text{Im}(z_2)| < \rho\},$$

and for a function  $f$  defined on this set we let

$$\|f\|_\rho := \sup_{(z_1, z_2) \in \Omega_\rho} |f((z_1, z_2))|.$$

We define  $C_\rho^\omega(\mathbb{T}^2)$  to be the space of all  $\mathbb{Z}^2$ -periodic real-analytic functions  $f$  on  $\mathbb{R}^2$  that extend to a holomorphic function on  $\Omega_\rho$  and satisfy  $\|f\|_\rho < \infty$ .

Hereby, we define  $\text{Diff}_\rho^\omega(\mathbb{T}^2, \lambda)$  to be the set of all Lebesgue measure-preserving real-analytic diffeomorphisms of  $\mathbb{T}^2$  homotopic to the identity, whose lift  $F$  to  $\mathbb{R}^2$  satisfies  $f_i \in C_\rho^\omega(\mathbb{T}^2)$ , and we also require that the lift  $\tilde{F}(x) = (x_1 + \tilde{f}_1(x), x_2 + \tilde{f}_2(x))$

of its inverse to  $\mathbb{R}^2$  satisfies  $\tilde{f}_i \in C_\rho^\omega(\mathbb{T}^2)$ . Then the metric in  $\text{Diff}_\rho^\omega(\mathbb{T}^2, \lambda)$  is defined by

$$d_\rho(f, g) = \max\{\tilde{d}_\rho(f, g), \tilde{d}_\rho(f^{-1}, g^{-1})\}, \text{ where } \tilde{d}_\rho(f, g) = \max_{i=1,2} \left\{ \inf_{n \in \mathbb{Z}} \|f_i - g_i + n\|_\rho \right\}.$$

We note that if  $\{f_n\}_{n=1}^\infty \subset \text{Diff}_\rho^\omega(\mathbb{T}^2, \lambda)$  is a Cauchy sequence in the  $d_\rho$  metric, then  $f_n$  converges to some  $f \in \text{Diff}_\rho^\omega(\mathbb{T}^2, \lambda)$ . Thus, this space is Polish.

**10.2. Circular systems.** A *circular coefficient sequence* is a sequence of pairs of positive integers  $(k_n, l_n)_{n \in \mathbb{N}}$  such that  $k_n \geq 2$  and  $\sum_{n \in \mathbb{N}} \frac{1}{l_n} < \infty$ . From these numbers we inductively define numbers

$$q_{n+1} = k_n l_n q_n^2$$

and

$$p_{n+1} = p_n k_n l_n q_n + 1,$$

where we set  $p_0 = 0$  and  $q_0 = 1$ . Obviously,  $p_{n+1}$  and  $q_{n+1}$  are relatively prime. Moreover, let  $\Sigma$  be a non-empty finite alphabet (in our case  $\Sigma = \{1, \dots, 2^{12}\}$ ) and  $b, e$  be two additional symbols. Then given a circular coefficient sequence  $(k_n, l_n)_{n \in \mathbb{N}}$  we build collections of words  $\mathcal{W}_n^c$  in the alphabet  $\Sigma \cup \{b, e\}$  by induction as follows:

- Set  $\mathcal{W}_0^c = \Sigma$ .
- Having built  $\mathcal{W}_n^c$ , we choose a set  $P_{n+1} \subseteq (\mathcal{W}_n^c)^{k_n}$  of so-called *prewords* and build  $\mathcal{W}_{n+1}^c$  by taking all words of the form

$$\mathcal{C}_n(w_0, w_1, \dots, w_{k_n-1}) = \prod_{i=0}^{q_n-1} \prod_{j=0}^{k_n-1} \left( b^{q_n-j_i} w_j^{l_n-1} e^{j_i} \right)$$

with  $w_0 \dots w_{k_n-1} \in P_{n+1}$ . If  $n = 0$  we take  $j_0 = 0$ , and for  $n > 0$  we let  $j_i \in \{0, \dots, q_n - 1\}$  be such that

$$j_i \equiv (p_n)^{-1} i \pmod{q_n}.$$

We note that each word in  $\mathcal{W}_{n+1}^c$  has length  $k_n l_n q_n^2 = q_{n+1}$ .

**Definition 75.** A construction sequence  $(\mathcal{W}_n^c)_{n \in \mathbb{N}}$  will be called *circular* if it is built in this manner using the  $\mathcal{C}$ -operators and a circular coefficient sequence, and each  $P_{n+1}$  is uniquely readable in the alphabet with the words from  $\mathcal{W}_n^c$  as letters. (This last property is called the *strong readability assumption*.)

*Remark.* By [FW19b, Lemma 45] each  $\mathcal{W}_n^c$  in a circular construction sequence is uniquely readable even if the prewords are not uniquely readable. However, the definition of a circular construction sequence requires this stronger readability assumption.

**Definition 76.** A symbolic shift  $\mathbb{K}$  built from a circular construction sequence is called a *circular system*. For emphasis we will often denote it by  $\mathbb{K}^c$ .

As described in [FW3, section 4.3] one can give an explicit construction sequence  $\{\text{rev}(\mathcal{W}_n^c)\}_{n \in \mathbb{N}}$  of  $(\mathbb{K}^c)^{-1}$  as

$$\text{rev}(\mathcal{W}_{n+1}^c) = \{\mathcal{C}_n^r(\text{rev}(w_0), \text{rev}(w_1), \dots, \text{rev}(w_{k_n-1})) : w_0 w_1 \dots w_{k_n-1} \in P_{n+1}\}$$

using the operator

$$(10.1) \quad \mathcal{C}_n^r(w_0, w_1, \dots, w_{k_n-1}) = \prod_{i=0}^{q_n-1} \prod_{j=0}^{k_n-1} \left( e^{q_n-j_{i+1}} w_{k_n-j-1}^{l_n-1} b^{j_{i+1}} \right).$$

For a circular system  $\mathbb{K}^c$  with circular coefficients  $(k_n, l_n)_{n \in \mathbb{N}}$  we also construct a second circular system: Let  $\Sigma_0 = \{*\}$  and we define a construction sequence

$$\mathcal{W}_0^c := \Sigma_0, \quad \text{and if} \quad \mathcal{W}_n^c := \{w_n\} \quad \text{then} \quad \mathcal{W}_{n+1}^c = \{\mathcal{C}_n(w_n, \dots, w_n)\}.$$

We denote the resulting circular system by  $\mathcal{K}$ . As shown in [FW19b, section 3.4]  $\mathcal{K}$  is a factor of  $\mathbb{K}^c$  measure theoretically isomorphic to a rotation of the circle. We call it the *canonical circle factor*. In [FW19b, section 4.3] a specific isomorphism  $\natural : \mathcal{K} \rightarrow \text{rev}(\mathcal{K})$  is introduced. It is called the *natural map* and will serve as a benchmark for understanding maps from  $\mathbb{K}^c$  to  $\text{rev}(\mathbb{K}^c)$  (see, for example, Definition 79).

By [FW19a, Theorem 60] circular systems can be realized as smooth diffeomorphisms via the untwisted AbC method provided that the sequence  $(l_n)_{n \in \mathbb{N}}$  grows sufficiently fast.

**Lemma 77.** *Let  $M$  be  $\mathbb{D}^2$ ,  $\mathbb{A}$ , or  $\mathbb{T}^2$  equipped with Lebesgue measure  $\lambda$ , and let  $\{\mathcal{W}_n^c\}_{n \in \mathbb{N}}$  be a strongly uniform circular construction sequence with circular coefficients  $(k_n, l_n)_{n \in \mathbb{N}}$ . We suppose that  $|\mathcal{W}_n^c| \rightarrow \infty$  as  $n \rightarrow \infty$  and that  $|\mathcal{W}_n^c|$  divides  $|\mathcal{W}_{n+1}^c|$ . If the sequence  $(l_n)_{n \in \mathbb{N}}$  grows sufficiently fast, then there is an area-preserving  $C^\infty$  diffeomorphism  $T$  on  $M$  such that the system  $(M, \lambda, T)$  is isomorphic to  $(\mathbb{K}^c, \nu, sh)$ .*

The following real-analytic counterpart is proven in [BKpp, Theorem G].

**Lemma 78.** *Let  $\rho > 0$ , and let  $\lambda$  be the Lebesgue measure on  $\mathbb{T}^2$ . Moreover, let  $\{\mathcal{W}_n^c\}_{n \in \mathbb{N}}$  be a strongly uniform circular construction sequence with circular coefficients  $(k_n, l_n)_{n \in \mathbb{N}}$ . We suppose that  $|\mathcal{W}_n^c| \rightarrow \infty$  as  $n \rightarrow \infty$  and that  $|\mathcal{W}_n^c|$  divides  $|\mathcal{W}_{n+1}^c|$ . If the sequence  $(l_n)_{n \in \mathbb{N}}$  grows sufficiently fast, then there is  $T \in \text{Diff}_\rho^\omega(\mathbb{T}^2, \lambda)$  such that the system  $(\mathbb{T}^2, \lambda, T)$  is isomorphic to  $(\mathbb{K}^c, \nu, sh)$ .*

We end this subsection by introducing the following subscales for a word  $w \in \mathcal{W}_{n+1}^c$  as in [FW19b, Subsection 3.3]:

- Subscale 0 is the scale of the individual powers of  $w_j \in \mathcal{W}_n^c$  of the form  $w_j^{l_j-1}$  and each such occurrence of a  $w_j^{l_j-1}$  is called a *0-subsection*.
- Subscale 1 is the scale of each term in the product  $\prod_{j=0}^{k_n-1} (b^{q_n-j_i} w_j^{l_n-1} e^{j_i})$  that has the form  $(b^{q_n-j_i} w_j^{l_n-1} e^{j_i})$  and these terms are called *1-subsections*.
- Subscale 2 is the scale of each term of  $\prod_{i=0}^{q_n-1} \prod_{j=0}^{k_n-1} (b^{q_n-j_i} w_j^{l_n-1} e^{j_i})$  that has the form  $\prod_{j=0}^{k_n-1} (b^{q_n-j_i} w_j^{l_n-1} e^{j_i})$  and these terms are called *2-subsections*.

**10.3. Categories  $\mathcal{OB}$  and  $\mathcal{CB}$  and the functor  $\mathcal{F} : \mathcal{OB} \rightarrow \mathcal{CB}$ .** For a fixed circular coefficient sequence  $(k_n, l_n)_{n \in \mathbb{N}}$  we consider two categories  $\mathcal{OB}$  and  $\mathcal{CB}$  whose objects are odometer-based and circular systems, respectively. The morphisms in these categories are (synchronous and anti-synchronous) graph joinings.

**Definition 79.** If  $\mathbb{K}$  is an odometer-based system, we denote its odometer base by  $\mathbb{K}^\pi$  and let  $\pi : \mathbb{K} \rightarrow \mathbb{K}^\pi$  be the canonical factor map. Similarly, if  $\mathbb{K}^c$  is a circular system, we let  $(\mathbb{K}^c)^\pi$  be the circle factor  $\mathcal{K}$  and let  $\pi : \mathbb{K}^c \rightarrow \mathcal{K}$  be the canonical factor map.

- (1) Let  $\mathbb{K}$  and  $\mathbb{L}$  be odometer based systems with the same coefficient sequence and let  $\rho$  be a joining between  $\mathbb{K}$  and  $\mathbb{L}^{\pm 1}$ . Then  $\rho$  is called *synchronous*

if  $\rho$  joins  $\mathbb{K}$  and  $\mathbb{L}$  and the projection of  $\rho$  to a joining on  $\mathbb{K}^\pi \times \mathbb{L}^\pi$  is the graph joining determined by the identity map. The joining  $\rho$  is called *anti-synchronous* if  $\rho$  joins  $\mathbb{K}$  and  $\mathbb{L}^{-1}$  and the projection of  $\rho$  to a joining on  $\mathbb{K}^\pi \times (\mathbb{L}^{-1})^\pi$  is the graph joining determined by the map  $x \mapsto -x$ .

- (2) Let  $\mathbb{K}^c$  and  $\mathbb{L}^c$  be circular systems with the same coefficient sequence and let  $\rho$  be a joining between  $\mathbb{K}^c$  and  $(\mathbb{L}^c)^{\pm 1}$ . Then  $\rho$  is called *synchronous* if  $\rho$  joins  $\mathbb{K}^c$  and  $\mathbb{L}^c$  and the projection of  $\rho$  to a joining on  $\mathcal{K} \times \mathcal{L}$  is the graph joining determined by the identity map. The joining  $\rho$  is called *anti-synchronous* if  $\rho$  joins  $\mathbb{K}^c$  and  $(\mathbb{L}^c)^{-1}$  and the projection of  $\rho$  to a joining on  $\mathcal{K} \times \mathcal{L}^{-1}$  is the graph joining determined by the map  $rev(\cdot) \circ \natural$ .

In [FW19b] Foreman and Weiss define a functor taking odometer-based systems to circular system that preserves the factor and conjugacy structure. To review the definition of the functor we fix a circular coefficient sequence  $(k_n, l_n)_{n \in \mathbb{N}}$ . Let  $\Sigma$  be a finite alphabet and  $(\mathcal{W}_n)_{n \in \mathbb{N}}$  be a construction sequence for an odometer-based system with coefficients  $(k_n)_{n \in \mathbb{N}}$ . Then we define a circular construction sequence  $(\mathcal{W}_n^c)_{n \in \mathbb{N}}$  and bijections  $c_n : \mathcal{W}_n \rightarrow \mathcal{W}_n^c$  by induction:

- Let  $\mathcal{W}_0^c = \Sigma$  and  $c_0$  be the identity map.
- Suppose that  $\mathcal{W}_n, \mathcal{W}_n^c$  and  $c_n$  have already been defined. Then we define

$$\mathcal{W}_{n+1}^c = \{ \mathcal{C}_n(c_n(w_0), c_n(w_1), \dots, c_n(w_{k_n-1})) : w_0 w_1 \dots w_{k_n-1} \in \mathcal{W}_{n+1} \}$$

and the map  $c_{n+1}$  by setting

$$c_{n+1}(w_0 w_1 \dots w_{k_n-1}) = \mathcal{C}_n(c_n(w_0), c_n(w_1), \dots, c_n(w_{k_n-1})).$$

In particular, the prewords are

$$P_{n+1} = \{ c_n(w_0) c_n(w_1) \dots c_n(w_{k_n-1}) : w_0 w_1 \dots w_{k_n-1} \in \mathcal{W}_{n+1} \}.$$

**Definition 80.** Suppose that  $\mathbb{K}$  is built from a construction sequence  $(\mathcal{W}_n)_{n \in \mathbb{N}}$  and  $\mathbb{K}^c$  has the circular construction sequence  $(\mathcal{W}_n^c)_{n \in \mathbb{N}}$  as constructed above. Then we define a map  $\mathcal{F}$  from the set of odometer-based systems (viewed as subshifts) to circular systems (viewed as subshifts) by

$$\mathcal{F}(\mathbb{K}) = \mathbb{K}^c.$$

*Remark 81.* The map  $\mathcal{F}$  is a bijection between odometer-based symbolic systems with coefficients  $(k_n)_{n \in \mathbb{N}}$  and circular symbolic systems with coefficients  $(k_n, l_n)_{n \in \mathbb{N}}$  that preserves uniformity. Since the construction sequences for our odometer-based systems are uniquely readable, the corresponding circular construction sequences automatically satisfy the strong readability assumption.

In [FW19b] it is shown that  $\mathcal{F}$  gives an isomorphism between the categories  $\mathcal{OB}$  and  $\mathcal{CB}$ . We state the following fact which is part of the main result in [FW19b].

**Lemma 82.** *For a fixed circular coefficient sequence  $(k_n, l_n)_{n \in \mathbb{N}}$  the categories  $\mathcal{OB}$  and  $\mathcal{CB}$  are isomorphic by the functor  $\mathcal{F}$  that takes synchronous isomorphisms to synchronous isomorphisms and anti-synchronous isomorphisms to anti-synchronous isomorphisms.*

*Remark 83.* Using the operator  $\mathcal{C}_n^r$  from equation (10.1) we can give a construction sequence  $\{ rev(\mathcal{W}_n^c) \}_{n \in \mathbb{N}}$  for  $\mathcal{F}(\mathbb{K})^{-1} = (\mathbb{K}^c)^{-1}$  via

$$rev(\mathcal{W}_{n+1}^c) = \{ \mathcal{C}_n^r(rev(c_n(w_0)), rev(c_n(w_1)), \dots, rev(c_n(w_{k_n-1}))) : w_0 w_1 \dots w_{k_n-1} \in \mathcal{W}_{n+1} \}.$$

We recall that the length of a  $n$ -block  $w$  in the odometer-based system is  $h_n = \prod_{i=0}^{n-1} k_i$ , if  $n > 0$ , and  $h_0 = 1$ , while the length of a  $n$ -block in the circular system is  $q_n$ , that is,  $|c_n(w)| = q_n$ . Moreover, we will use the following map from substrings of the underlying odometer-based system to the circular system:

$$C_{n,i}(w_s w_{s+1} \dots w_t) = \prod_{j=s}^t \left( b^{q_n - j_i} (c_n(w_j))^{l_n - 1} e^{j_i} \right)$$

for any  $0 \leq i \leq q_n - 1$  and  $0 \leq s \leq t \leq k_n - 1$ .

*Remark 84.* As in [FW3, section 4.9] we also propagate our equivalence relations and group actions to the circular system. For all  $n \in \mathbb{N}$ ,  $w_1, w_2 \in \mathcal{W}_n$  we let  $(c_n(w_1), c_n(w_2))$  be in the relation  $(\mathcal{Q}_s^n)^c$  if and only if  $(w_1, w_2) \in \mathcal{Q}_s^n$ . Moreover, given  $g \in G_s^n$ , we let  $g[c_n(w_1)]_s = [c_n(w_2)]_s$  if and only if  $g[w_1]_s = [w_2]_s$ .

**10.4.  $\bar{f}$ -estimates in the circular system.** We recall that we enumerate construction sequences of our odometer-based systems as  $\{\mathcal{W}_n\}_{n \in \mathbb{N}}$  in this section, that is,  $(n+1)$ -words are built by concatenating  $n$ -words. Before we analyze the circular system we foreshadow that we will choose the circular coefficients  $(l_n)_{n \in \mathbb{N}}$  as the last sequence of parameters in Subsection 10.6 to grow fast enough to allow the realization as diffeomorphisms. We stress that none of the other parameters depends on  $(l_n)_{n \in \mathbb{N}}$ . In particular, we can choose  $(l_n)_{n \in \mathbb{N}}$  to satisfy

$$(10.2) \quad l_n \geq \max(4R_{n+2}^2, 9l_{n-1}^2).$$

For the  $\bar{f}$ -estimates in the circular system we make use of the  $\bar{f}$ -estimates in the underlying odometer-based system. We will also use a sequence  $(R_n^c)_{n=1}^\infty$ , where  $R_1^c = R_1$  (with  $R_1$  from the sequence  $(R_n)_{n \in \mathbb{N}}$  in the construction of the odometer-based system in Section 8) and  $R_n^c = \lfloor \sqrt{l_{n-2}} \cdot k_{n-2} \cdot q_{n-2}^2 \rfloor$  for  $n \geq 2$ . We note that for  $n \geq 2$ ,

$$(10.3) \quad \frac{q_n}{R_n^c} = \frac{k_{n-1} \cdot l_{n-1} \cdot (k_{n-2} \cdot l_{n-2} \cdot q_{n-2}^2) \cdot q_{n-1}}{\lfloor \sqrt{l_{n-2}} \cdot k_{n-2} \cdot q_{n-2}^2 \rfloor} \geq \sqrt{l_{n-2}} \cdot k_{n-1} \cdot l_{n-1} \cdot q_{n-1}.$$

Hence, for  $n \geq 2$  a substring of at least  $q_n/R_n^c$  consecutive symbols in a circular  $n$ -block contains at least  $\sqrt{l_{n-2}} - 1$  complete 2-subsections, which have length  $k_{n-1}l_{n-1}q_{n-1}$  (recall the notion of a 2-subsection from the end of Subsection 10.2). When conducting estimates in Lemma 88 this will allow us to ignore incomplete 2-subsections at the ends of the substring.

Now we show  $\bar{f}$ -estimates in the circular system by induction. We start with the base case  $n = 1$ .

**Lemma 85.** *Let  $w, \bar{w} \in \mathcal{W}_1$  and  $\mathcal{A}, \bar{\mathcal{A}}$  be any substrings of at least  $q_1/R_1^c$  consecutive symbols in  $c_1(w)$  and  $c_1(\bar{w})$ , respectively.*

(1) *If  $[w]_1 \neq [\bar{w}]_1$ , then we have*

$$(10.4) \quad \bar{f}(\mathcal{A}, \bar{\mathcal{A}}) > \frac{1}{2} - \frac{3}{R_1} - \frac{2}{l_0}.$$

(2) *If  $w \neq \bar{w}$ , then we have*

$$(10.5) \quad \bar{f}(\mathcal{A}, \bar{\mathcal{A}}) > \frac{1}{p_1} \cdot \left( \frac{1}{2} - \frac{1}{R_1} \right) - \frac{3}{R_1} - \frac{2}{l_0}.$$

*Proof.* Since  $q_0 = 1$ , we have

$$\frac{q_1}{R_1^c} = \frac{k_0 \cdot l_0}{R_1}.$$

By adding fewer than  $2l_0$  symbols to each of  $\mathcal{A}$  and  $\overline{\mathcal{A}}$  we can complete partial strings  $bw_j^{l_0-1}$ , where  $w_j \in \mathcal{W}_0$ , at the beginning and end. The spacer  $b$  occupies a proportion of  $\frac{1}{l_0}$  in the augmented strings of  $\mathcal{A}$  and  $\overline{\mathcal{A}}$ , respectively. We ignore the spacers in the following consideration and denote the modified strings by  $\mathcal{A}'$  and  $\overline{\mathcal{A}'}$ . By Fact 9 we have

$$\overline{f}(\mathcal{A}, \overline{\mathcal{A}}) > \overline{f}(\mathcal{A}', \overline{\mathcal{A}'}) - \frac{2R_1}{k_0} - \frac{2}{l_0} > \overline{f}(\mathcal{A}', \overline{\mathcal{A}'}) - \frac{1}{R_1} - \frac{2}{l_0}.$$

Since 0-words are symbols, we obtain from Remark 54 that

$$\overline{f}(\mathcal{A}', \overline{\mathcal{A}'}) > \begin{cases} \frac{1}{2} - \frac{2}{R_1} - \frac{2}{p_1}, & \text{if } [w]_1 \neq [\overline{w}]_1, \\ \frac{1}{p_1} \cdot \left(\frac{1}{2} - \frac{1}{R_1}\right) - \frac{2}{R_1}, & \text{if } w \neq \overline{w}. \end{cases}$$

□

This motivates the definitions

$$\alpha_{1,1}^c = \min\left(\frac{1}{2} - \frac{3}{R_1} - \frac{2}{p_1} - \frac{2}{l_0}, \frac{1}{9}\right), \quad \beta_1^c = \min\left(\frac{1}{p_1} \cdot \left(\frac{1}{2} - \frac{1}{R_1}\right) - \frac{3}{R_1} - \frac{2}{l_0}, \alpha_{1,1}^c\right).$$

We note that by  $R_1 \geq 40p_1$  from condition (8.6), we have  $\beta_1^c > \frac{1}{4p_1} \geq \frac{10}{R_1}$ . Hence,  $R_1 > \frac{10}{\beta_1^c}$ .

We assume for  $n \geq 1$  that there are  $\frac{1}{8} > \alpha_{1,n}^c > \dots > \alpha_{s(n),n}^c > \beta_n^c$  such that in addition to (8.2) we have

$$(10.6) \quad R_n \geq \frac{7}{\beta_n^c},$$

and the following estimates on  $\overline{f}$  distances hold:

- For every  $s \in \{1, \dots, s(n)\}$  we assume that if  $w, \overline{w} \in \mathcal{W}_n$  with  $[w]_s \neq [\overline{w}]_s$  for some  $s \leq s(n)$ , then

$$(10.7) \quad \overline{f}(\mathcal{A}, \overline{\mathcal{A}}) > \alpha_{s,n}^c$$

on any substrings  $\mathcal{A}, \overline{\mathcal{A}}$  of at least  $q_n/R_n^c$  consecutive symbols in  $c_n(w)$  and  $c_n(\overline{w})$ , respectively.

- For  $w, \overline{w} \in \mathcal{W}_n$  with  $w \neq \overline{w}$ , we have

$$(10.8) \quad \overline{f}(\mathcal{A}, \overline{\mathcal{A}}) > \beta_n^c$$

on any substrings  $\mathcal{A}, \overline{\mathcal{A}}$  of at least  $q_n/R_n^c$  consecutive symbols in  $c_n(w)$  and  $c_n(\overline{w})$ , respectively.

We also recall the assumption  $p_{n+1} > \max\left(4R_n, 2^{n+1}, \frac{1}{\epsilon_{n+1}}\right)$  from (8.5).

At stage  $n+1$  of the construction we choose  $R_{n+1}$  such that besides (8.6) it also satisfies

$$(10.9) \quad R_{n+1} \geq \frac{40p_{n+1}}{\beta_n^c}.$$

Our assumptions (10.7) and (10.8) immediately imply  $\overline{f}$ -estimates for 1-subsections.



**Lemma 86.** Let  $w, \bar{w} \in \mathcal{W}_n$ ,  $0 \leq i_1, i_2 < q_n$ , and  $\mathcal{B}, \bar{\mathcal{B}}$  be any substrings of at least  $\sqrt{l_n}q_n$  consecutive symbols in  $\mathcal{C}_{n,i_1}(w)$  and  $\mathcal{C}_{n,i_2}(\bar{w})$ , respectively.

(1) If  $[w]_s \neq [\bar{w}]_s$  for any  $s \leq s(n)$ , then we have

$$(10.10) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \alpha_{s,n}^c - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_n}}.$$

(2) If  $w \neq \bar{w}$ , then we have

$$(10.11) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \beta_n^c - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_n}}.$$

*Proof.* The newly introduced spacers  $b$  and  $e$  occupy a proportion of at most  $\frac{q_n}{\sqrt{l_n}q_n} = \frac{1}{\sqrt{l_n}}$  in each  $\mathcal{B}$  and  $\bar{\mathcal{B}}$ . We ignore them in the following consideration, which might increase the  $\bar{f}$ -distance by at most  $\frac{2}{\sqrt{l_n}}$  according to Fact 9. Then we apply Lemma 13 with  $\bar{f} = 1$  and our assumptions (10.7) and (10.8), respectively.  $\square$

Hereby, we can prove  $\bar{f}$ -estimates for substantial substrings of 2-subsections.

**Lemma 87.** Let  $w, \bar{w} \in \mathcal{W}_{n+1}$ ,  $0 \leq i_1, i_2 < q_n$ , and  $\mathcal{B}, \bar{\mathcal{B}}$  be any substrings of at least  $\frac{k_n l_n q_n}{R_{n+1}}$  consecutive symbols in  $\mathcal{C}_{n,i_1}(w)$  and  $\mathcal{C}_{n,i_2}(\bar{w})$ , respectively.

(1) If  $s(n+1) = s(n)$ , then we have

$$(10.12) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \begin{cases} \alpha_{s,n}^c - \frac{1}{R_n} - \frac{2}{R_{n+1}} - \frac{1}{R_n^c} - \frac{3}{\sqrt{l_n}}, & \text{if } [w]_s \neq [\bar{w}]_s, \\ \beta_n^c - \frac{1}{R_n} - \frac{2}{R_{n+1}} - \frac{1}{R_n^c} - \frac{3}{\sqrt{l_n}}, & \text{if } w \neq \bar{w}. \end{cases}$$

(2) If  $s(n+1) = s(n) + 1$ , then we have

$$(10.13) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \begin{cases} \alpha_{s,n}^c - \frac{1}{R_n} - \frac{3}{R_{n+1}} - \frac{1}{R_n^c} - \frac{3}{\sqrt{l_n}}, & \text{if } [w]_s \neq [\bar{w}]_s, \\ \beta_n^c - \frac{1}{R_n} - \frac{3}{R_{n+1}} - \frac{1}{R_n^c} - \frac{3}{\sqrt{l_n}}, & \text{if } [w]_{s(n)+1} \neq [\bar{w}]_{s(n)+1}, \\ \frac{1}{2p_{n+1}} \cdot \left( \beta_n^c - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_n}} \right) - \frac{1}{R_{n+1}} - \frac{1}{\sqrt{l_n}}, & \text{if } w \neq \bar{w}. \end{cases}$$

*Proof.* By adding fewer than  $2l_n q_n$  symbols to each of  $\mathcal{B}$  and  $\bar{\mathcal{B}}$  we can complete any partial  $\mathcal{C}_{n,i_1}(w_j)$  at the beginning and end of  $\mathcal{B}$  and any partial  $\mathcal{C}_{n,i_2}(\bar{w}_j)$  at the beginning and end of  $\bar{\mathcal{B}}$ , where  $w_j, \bar{w}_j \in \mathcal{W}_n$ . Let  $\mathcal{B}_{aug}$  and  $\bar{\mathcal{B}}_{aug}$  be the augmented  $\mathcal{B}$  and  $\bar{\mathcal{B}}$  strings obtained in this way. By Fact 9 we have

$$(10.14) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) \geq \bar{f}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) - \frac{2R_{n+1}}{k_n} > \bar{f}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) - \frac{1}{R_{n+1}}.$$

We consider the corresponding strings  $\mathcal{B}_{aug}$  and  $\bar{\mathcal{B}}_{aug}$  in the words  $w, \bar{w} \in \mathcal{W}_{n+1}$  expressed in the alphabets  $\Sigma_{n,s} = (\mathcal{W}_n/\mathcal{Q}_s^n)^*$  or  $\Sigma_n = (\mathcal{W}_n)^*$ . They have length at least  $\frac{k_n}{R_{n+1}}$ . Hence, we can apply Remarks 47 or 53 depending on whether we are in case 1 ( $s(n+1) = s(n)$ ) or case 2 ( $s(n+1) = s(n) + 1$ ) of the construction.

We examine case 2. If  $[w]_s \neq [\bar{w}]_s$  for  $s \leq s(n)$ , then we have  $\bar{f}_{\Sigma_{n,s}}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) \geq 1 - \frac{1}{R_n} - \frac{2}{R_{n+1}}$ . We apply Lemma 13 with  $\bar{f} \geq 1 - \frac{1}{R_n} - \frac{2}{R_{n+1}}$  and the estimate

from (10.10) to obtain

$$\begin{aligned}\bar{f}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) &\geq \left(1 - \frac{1}{R_n} - \frac{2}{R_{n+1}}\right) \cdot \left(\alpha_{s,n}^c - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_n}}\right) - \frac{1}{\sqrt{l_n}} \\ &\geq \alpha_{s,n}^c - \frac{1}{R_n} - \frac{2}{R_{n+1}} - \frac{1}{R_n^c} - \frac{3}{\sqrt{l_n}}.\end{aligned}$$

Together with (10.14) we deduce the first inequality in (10.13). In an analogous manner, we obtain the second inequality in (10.13) from (10.11) and  $\bar{f}_{\Sigma_n}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) \geq 1 - \frac{1}{R_n} - \frac{2}{R_{n+1}}$  following from (8.24). To see the third inequality, we note that  $\bar{f}_{\Sigma_n}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) \geq \frac{1}{2p_{n+1}}$  by (8.23) in Remark 53. Thus, we apply Lemma 13 and the estimate from (10.11) to obtain

$$\bar{f}(\mathcal{B}_{aug}, \bar{\mathcal{B}}_{aug}) \geq \frac{1}{2p_{n+1}} \cdot \left(\beta_n^c - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_n}}\right) - \frac{1}{\sqrt{l_n}},$$

which yields the claim with the aid of (10.14).

By the same method we can also examine case 1 and conclude (10.12).  $\square$

As an immediate consequence from the previous lemma we obtain  $\bar{f}$ -estimates for  $(n+1)$ -words in the circular system.

**Lemma 88.** *Let  $w, \bar{w} \in \mathcal{W}_{n+1}$  and  $\mathcal{B}, \bar{\mathcal{B}}$  be any substrings of at least  $q_{n+1}/R_{n+1}^c$  consecutive symbols in  $c_{n+1}(w)$  and  $c_{n+1}(\bar{w})$ , respectively.*

(1) *If  $s(n+1) = s(n)$ , then we have*

$$(10.15) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \begin{cases} \alpha_{s,n}^c - \frac{2}{R_n} - \frac{3}{R_{n+1}}, & \text{if } [w]_s \neq [\bar{w}]_s, \\ \beta_n^c - \frac{2}{R_n} - \frac{3}{R_{n+1}}, & \text{if } w \neq \bar{w}. \end{cases}$$

(2) *If  $s(n+1) = s(n) + 1$ , then we have*

$$(10.16) \quad \bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \begin{cases} \alpha_{s,n}^c - \frac{2}{R_n} - \frac{4}{R_{n+1}}, & \text{if } [w]_s \neq [\bar{w}]_s, \\ \beta_n^c - \frac{2}{R_n} - \frac{4}{R_{n+1}}, & \text{if } [w]_{s(n)+1} \neq [\bar{w}]_{s(n)+1}, \\ \frac{1}{2p_{n+1}} \cdot \left(\beta_n^c - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_n}}\right) - \frac{3}{R_{n+1}}, & \text{if } w \neq \bar{w}. \end{cases}$$

*Proof.* In case of  $n > 0$  we note that  $\mathcal{B}$  and  $\bar{\mathcal{B}}$  have at least the length of  $\sqrt{l_{n-1}}$  complete 2-subsections by equation (10.3). By adding fewer than  $2l_n k_n q_n$  symbols to each of  $\mathcal{B}$  and  $\bar{\mathcal{B}}$ , we can complete any partial 2-subsections at the beginning and end of  $\mathcal{B}$  and  $\bar{\mathcal{B}}$ . This change will increase the  $\bar{f}$  distance between  $\mathcal{B}$  and  $\bar{\mathcal{B}}$  by less than  $2/\sqrt{l_{n-1}}$  due to Fact 9. Then we apply Lemma 13 with  $\bar{f} = 1$  and the corresponding estimates from Lemma 87. In case of  $s(n+1) = s(n)$  this yields

$$\bar{f}(\mathcal{B}, \bar{\mathcal{B}}) > \begin{cases} \alpha_{s,n}^c - \frac{1}{R_n} - \frac{3}{R_{n+1}} - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_{n-1}}} - \frac{3}{\sqrt{l_n}}, & \text{if } [w]_s \neq [\bar{w}]_s, \\ \beta_n^c - \frac{1}{R_n} - \frac{3}{R_{n+1}} - \frac{1}{R_n^c} - \frac{2}{\sqrt{l_{n-1}}} - \frac{3}{\sqrt{l_n}}, & \text{if } w \neq \bar{w}. \end{cases}$$

We use the conditions on the sequence  $(l_n)_{n \in \mathbb{N}}$  from (10.2) to estimate

$$\frac{1}{R_n^c} + \frac{2}{\sqrt{l_{n-1}}} + \frac{3}{\sqrt{l_n}} \leq \frac{1}{\sqrt{l_{n-2}}} + \frac{3}{\sqrt{l_{n-1}}} \leq \frac{2}{\sqrt{l_{n-2}}} \leq \frac{1}{R_n},$$

which implies (10.15). Similarly, we obtain (10.16).  $\square$

This accomplishes the induction step. In particular, we note with the aid of (10.9) that

$$\beta_{n+1}^c > \frac{\beta_n^c}{4p_{n+1}} - \frac{3}{R_{n+1}} \geq \frac{7\beta_n^c}{40p_{n+1}} \geq \frac{7}{R_{n+1}},$$

that is,  $R_{n+1} > \frac{7}{\beta_{n+1}^c}$  and the induction assumption (10.6) for the next step is satisfied. Since we have

$$\alpha_{s,M(s)}^c = \beta_{M(s)-1}^c - \frac{2}{R_{M(s)-1}} - \frac{4}{R_{M(s)}} \geq \beta_{M(s)-1}^c - \frac{2}{7}\beta_{M(s)-1}^c - \frac{\beta_{M(s)-1}^c}{10p_{M(s)}} > \frac{\beta_{M(s)-1}^c}{2}$$

by (10.6) and (10.9), we also obtain

$$\alpha_{s,n}^c \geq \alpha_{s,M(s)}^c - \frac{2}{R_{M(s)}} - \sum_{j>M(s)} \frac{6}{R_j} \geq \alpha_{s,M(s)}^c - \frac{\beta_{M(s)-1}^c}{20p_{M(s)}} - \frac{6\beta_{M(s)-1}^c}{40} > \frac{\alpha_{s,M(s)}^c}{2}$$

for any  $n > M(s)$  by conditions (10.6) and (10.9). Hereby, we see that for every  $s \in \mathbb{N}$  the decreasing sequence  $(\alpha_{s,n}^c)_{n \geq M(s)}$  is bounded from below by  $\frac{\alpha_{s,M(s)}^c}{2}$ , and we denote its limit by  $\alpha_s^c$ . Altogether, this proves the following proposition.

**Proposition 89.** *For every  $n \geq M(s)$ , we have*

$$(10.17) \quad \bar{f}(\mathcal{A}, \bar{\mathcal{A}}) > \alpha_s^c$$

on any substrings  $\mathcal{A}, \bar{\mathcal{A}}$  of at least  $q_n/R_n^c$  consecutive symbols in  $c_n(w)$  and  $c_n(\bar{w})$ , respectively, for  $w, \bar{w} \in \mathcal{W}_n$  with  $[w]_s \neq [\bar{w}]_s$ .

**10.5. Proof of Non-Kakutani Equivalence.** We follow the strategy in Section 9 to show that for a tree  $\mathcal{T} \in \text{Trees}$  without an infinite branch  $T_c = \mathcal{F}(\Psi(\mathcal{T}))$  and  $T_c^{-1} = \mathcal{F}(\Psi(\mathcal{T}))^{-1}$  are not Kakutani equivalent.

**Lemma 90.** *Let  $\mathcal{T} \in \text{Trees}$ ,  $T_c = \mathcal{F}(\Psi(\mathcal{T}))$ , and  $f : X \rightarrow \mathbb{Z}^+$  be integrable. If  $T_c^f \cong T_c$ , then  $\int f d\mu = 1$ .*

*Proof.* The proof follows along the lines of Proposition 56. This time, one decomposes the name  $a(T_c, P_c, x)$  in the circular system into strings of the form  $\mathcal{C}_{n,j}(F_{n,i})$ , where  $0 \leq j < q_n$  and  $F_{n,i}$  is a Feldman pattern of 1-equivalence classes of  $n$ -words in the odometer-based system (as considered in the proof of Proposition 56).  $\square$

Hereby, we can show as in Proposition 67 that if  $\mathbb{K}^c = \mathcal{F}(\Psi(\mathcal{T}))$  and  $(\mathbb{K}^c)^{-1} = \mathcal{F}(\Psi(\mathcal{T}))^{-1}$  are Kakutani equivalent, then it can only be by an even equivalence. We show that  $\mathbb{K}^c$  and  $(\mathbb{K}^c)^{-1}$  are not evenly equivalent if  $\mathcal{T} \in \text{Trees}$  does not have an infinite branch. For this purpose, we prove the following analogues of Lemmas 71 and 72.

**Lemma 91.** *Let  $s \in \mathbb{N}$  and  $\phi$  be a finite code from  $\mathbb{K}^c$  to  $\text{rev}(\mathbb{K}^c)$ . Then for sufficiently large  $n \in \mathbb{N}$ ,  $0 \leq i_1, i_2 < q_n$ , and any pair of  $s$ -Feldman patterns  $P_{n,\ell}$  in  $\mathbb{K}$  and  $\bar{P}_{n,\bar{\ell}}$  in  $\text{rev}(\mathbb{K})$  of different pattern type we have*

$$\bar{f}(\phi(\mathcal{G}), \bar{\mathcal{G}}) > \frac{\alpha_s^c}{10}$$

for any substrings  $\mathcal{G}, \bar{\mathcal{G}}$  of at least  $\frac{|P_{n,\ell}| \cdot l_n q_n}{2^{2e(n)}} = \frac{|\bar{P}_{n,\bar{\ell}}| \cdot l_n q_n}{2^{2e(n)}}$  consecutive symbols in  $\mathcal{C}_{n,i_1}(P_{n,\ell})$  and  $\mathcal{C}_{n,i_2}^r(\bar{P}_{n,\bar{\ell}})$ , respectively.

*Proof.* The proof follows along the lines of Lemma 71.  $\square$

**Lemma 92.** *Let  $s \in \mathbb{N}$  and  $\phi$  be a finite code from  $\mathbb{K}^c$  to  $\text{rev}(\mathbb{K}^c)$ . Moreover, let  $n \in \mathbb{N}$  be sufficiently large,  $0 \leq i_1, i_2 < q_n$ , and  $w, \bar{w} \in \mathcal{W}_{n+1}$  be any pair such that  $w$  and  $\text{rev}(\bar{w})$  are of different  $s$ -pattern type. Then we have*

$$\bar{f}(\phi(\mathcal{A}), \bar{\mathcal{A}}) > \frac{\alpha_s^c}{16}$$

on any substrings  $\mathcal{A}, \bar{\mathcal{A}}$  of at least  $\frac{k_n l_n q_n}{2^{\varepsilon(n)}}$  consecutive symbols in  $\mathcal{C}_{n, i_1}(w)$  and  $\mathcal{C}_{n, i_2}^r(\bar{w})$ , respectively.

*Proof.* Analogously to the proof of Lemma 72, we divide  $\mathcal{A}$  and  $\bar{\mathcal{A}}$  into circular images  $\mathcal{P}_{n, j} = \mathcal{C}_{n, i_1}(P_{n, j})$  and  $\bar{\mathcal{P}}_{n, j} = \mathcal{C}_{n, i_2}^r(\bar{P}_{n, j})$ , where  $P_{n, j}$  and  $\bar{P}_{n, j}$  are  $s$ -Feldman patterns in  $w$  and  $\bar{w}$ , respectively. Then the proof follows along the lines of Lemma 72.  $\square$

We can use Lemmas 91 and 92 to identify a well-approximating finite code with an element of the group action in a similar way that we used Lemmas 71 and 72 in the proof of Lemma 73.

**Lemma 93.** *Let  $s \in \mathbb{N}$ ,  $0 < \delta < \frac{1}{4}$ ,  $0 < \varepsilon < \frac{\alpha_s^c}{200}\delta$ , and  $\phi$  be a finite code from  $\mathbb{K}^c$  to  $\text{rev}(\mathbb{K}^c)$ . Then for  $n$  sufficiently large, any  $0 \leq i_1 < q_n$ , any  $w \in \mathcal{W}_{n+1}$ , and any string  $\bar{\mathcal{A}}$  in  $\text{rev}(\mathbb{K}^c)$  with*

$$\bar{f}(\phi(\mathcal{C}_{n, i_1}(w)), \bar{\mathcal{A}}) < \varepsilon,$$

there must be exactly one string of the form  $\mathcal{C}_{n, i_2}^r(\bar{w})$  with  $|\bar{\mathcal{A}} \cap \mathcal{C}_{n, i_2}^r(\bar{w})| \geq (1 - \delta)k_n l_n q_n$  for some  $0 \leq i_2 < q_n$ ,  $\bar{w} \in \mathcal{W}_{n+1}$ . Moreover,  $\text{rev}(\bar{w})$  must be of the form  $[\text{rev}(\bar{w})]_s = \eta_g [w]_s$  for a unique  $g \in G_s$ .

*Proof.* As in the proof of Lemma 73 we neglect end effects and conclude that there can be at most one string of the form  $\mathcal{C}_{n, i_2}^r(\bar{w})$  with  $|\bar{\mathcal{A}} \cap \mathcal{C}_{n, i_2}^r(\bar{w})| \geq (1 - \delta)k_n l_n q_n$ . Suppose that there is no such  $\mathcal{C}_{n, i_2}^r(\bar{w})$ , but two strings  $\mathcal{C}_{n, i_{2,1}}^r(\bar{w}_1)$ ,  $\mathcal{C}_{n, i_{2,2}}^r(\bar{w}_2)$  with  $\bar{w}_1, \bar{w}_2 \in \mathcal{W}_n$  and  $0 \leq i_{2,1}, i_{2,2} < q_n$  such that  $\bar{\mathcal{A}}$  is a substring of  $\mathcal{C}_{n, i_{2,1}}^r(\bar{w}_1)\mathcal{C}_{n, i_{2,2}}^r(\bar{w}_2)$  and

$$|\bar{\mathcal{A}} \cap \mathcal{C}_{n, i_{2,j}}^r(\bar{w}_j)| \geq (\delta - 3\varepsilon) \cdot k_n l_n q_n > \frac{180}{\alpha_s^c} \varepsilon k_n l_n q_n \text{ for } j = 1, 2.$$

As before there are two possible cases: Either at least one of  $\text{rev}(\bar{w}_1), \text{rev}(\bar{w}_2)$  has a different  $s$ -pattern type from that of  $w$  or  $\text{rev}(\bar{w}_1), \text{rev}(\bar{w}_2)$  have the same  $s$ -pattern type as  $w$ . We recall that the  $s$ -pattern structure of the beginning and end of  $w$  are different from each other by construction. To investigate both cases we denote  $\bar{\mathcal{A}}_j := \bar{\mathcal{A}} \cap \mathcal{C}_{n, i_{2,j}}^r(\bar{w}_j)$ . Let  $\pi$  be a best possible match in  $\bar{f}$  between  $\phi(\mathcal{C}_{n, i_1}(w))$  and  $\bar{\mathcal{A}}$ . Then we denote the parts in  $\phi(\mathcal{C}_{n, i_1}(w))$  corresponding to  $\bar{\mathcal{A}}_j$  under  $\pi$  by  $\mathcal{A}_j$ . If  $\bar{f}(\mathcal{A}_j, \bar{\mathcal{A}}_j) > \alpha_s^c$ , we stop. Otherwise, we have  $\frac{8}{9} \leq \frac{|\mathcal{A}_j|}{|\bar{\mathcal{A}}_j|} \leq \frac{9}{8}$ , that is,  $|\mathcal{A}_j| \geq \frac{8}{9}|\bar{\mathcal{A}}_j| \geq \frac{160}{\alpha_s^c} \varepsilon k_n l_n q_n$ . Hence, for sufficiently large  $n$  the strings are long enough such that we can apply Lemmas 91 and 92. In both aforementioned cases this yields with the aid of Fact 10 that

$$\bar{f}(\phi(w), \bar{\mathcal{A}}) \geq \frac{80}{\alpha_s^c} \varepsilon \cdot \frac{\alpha_s^c}{16} = 5\varepsilon,$$

which contradicts the assumption  $\bar{f}(\phi(\mathcal{C}_{n, i_1}(w)), \bar{\mathcal{A}}) < \varepsilon$ . Hence, there must be exactly one  $\mathcal{C}_{n, i_2}^r(\bar{w})$  with  $|\bar{\mathcal{A}} \cap \mathcal{C}_{n, i_2}^r(\bar{w})| \geq (1 - \delta)k_n l_n q_n$ .

Lemma 92 also implies that  $w$  and  $\text{rev}(\bar{w})$  need to have the same  $s$ -pattern structure. As before this only happens if  $[\text{rev}(\bar{w})]_s = \eta_g[w]_s$  for  $g \in G_s$ .  $\square$

Assume  $\mathbb{K}^c$  and  $(\mathbb{K}^c)^{-1}$  are evenly equivalent. Let  $(\varepsilon_\ell)_{\ell \in \mathbb{N}}$  be a sequence of positive reals satisfying  $\sum_{i=\ell+1}^{\infty} \varepsilon_i < \varepsilon_\ell$ . Then we take a sequence of  $(\varepsilon_\ell, K_\ell)$ -finite  $\bar{f}$  codes  $\phi_\ell^c$  from  $\mathbb{K}^c$  to  $(\mathbb{K}^c)^{-1}$  as produced by Lemma 68. In the following we use  $\gamma_s^c = \frac{(\alpha_s^c)^4}{5 \cdot 10^9}$  for each  $s \in \mathbb{N}$ . We apply Corollary 70 with  $\gamma_s^c$  on our sequence of codes and obtain  $N(s) \in \mathbb{Z}^+$  such that for every  $N \geq N(s)$  and  $k \in \mathbb{Z}^+$  there is  $n(s, N, k) \in \mathbb{Z}^+$  sufficiently large such that for all  $n \geq n(s, N, k)$  Lemma 93 with  $\delta_s^c = \frac{(\alpha_s^c)^3}{2 \cdot 10^7}$  and  $\phi_N$  holds and there exists a collection of words  $\mathcal{W}'_n \subset \mathcal{W}_n^c$  of circular  $n$ -words satisfying the analogues of (C1) and (C2).

**Lemma 94.** *Suppose  $\mathbb{K}^c$  and  $(\mathbb{K}^c)^{-1}$  are evenly equivalent and let  $s \in \mathbb{N}$ . There is a unique  $g_s \in G_s$  such that for every  $N \geq N(s)$  and sufficiently large  $n \in \mathbb{N}$  we have for every  $w^c \in \mathcal{W}'_n$  that there is  $\bar{w}^c \in \text{rev}(\mathcal{W}_n^c)$  with  $[\bar{w}^c]_s = \eta_{g_s}[w^c]_s$  such that*

$$\bar{f}(\phi_N^c(w^c), \bar{w}^c) < \frac{\alpha_s^c}{4},$$

where  $(\phi_\ell^c)_\ell$  is a sequence of codes as described above. Moreover,  $g_s$  is of odd parity and the sequence  $(g_s)_{s \in \mathbb{N}}$  satisfies  $g_s = \rho_{s+1, s}(g_{s+1})$  for all  $s \in \mathbb{N}$ .

*Proof.* Let  $w^c \in \mathcal{W}'_{n+1} \subset \mathcal{W}_{n+1}^c$  and we recall that  $w^c = \prod_{i=0}^{q_n-1} \mathcal{C}_{n,i}(w)$  with  $w \in \mathcal{W}_{n+1}$ . As in the proof of Lemma 74 we use property (C1) to choose  $n$  sufficiently large such that there exists  $z \in \text{rev}(\mathbb{K}^c)$  with

$$\bar{f}(\phi_N^c(\mathcal{C}_{n,i}(w)), z \upharpoonright [0, k_n l_n q_n - 1]) < \gamma_s^c.$$

We denote  $\bar{\mathcal{A}}_i := z \upharpoonright [0, k_n l_n q_n - 1]$  in  $\text{rev}(\mathbb{K}^c)$ . By Lemma 93 there is a string of the form  $\mathcal{C}_{n,j_i}^r(\bar{w})$  with  $|\bar{\mathcal{A}}_i \cap \mathcal{C}_{n,j_i}^r(\bar{w})| \geq (1 - \delta_s^c) k_n l_n q_n$  for some  $0 \leq j_i < q_n$ ,  $\bar{w} \in \mathcal{W}_{n+1}$ , and  $\text{rev}(\bar{w})$  must be of the form  $[\text{rev}(\bar{w})]_s = \eta_g[w]_s$  for a unique  $g \in G_s$ . By Fact 9 we conclude  $\bar{f}(\phi_N^c(\mathcal{C}_{n,i}(w)), \mathcal{C}_{n,j_i}^r(\bar{w})) < \gamma_s^c + \delta_s^c$ . Since this holds true for every  $i \in \{0, \dots, q_n - 1\}$  and the newly introduced spacers occupy a proportion of at most  $\frac{1}{l_n}$ , we obtain for  $n$  sufficiently large that  $\bar{f}(\phi_N^c(w^c), \mathcal{C}_n^r(\bar{w})) < \gamma_s^c + \delta_s^c + \frac{2}{l_n} < 10^{-7} (\alpha_s^c)^3$ . Hence, we choose  $\bar{w}^c = \mathcal{C}_n^r(\bar{w}) \in \text{rev}(\mathcal{W}_{n+1}^c)$  which satisfies  $[\bar{w}^c]_s = \eta_g[w^c]_s$  by  $[\text{rev}(\bar{w})]_s = \eta_g[w]_s$  and Remark 84. To see that the same  $g \in G_s$  is supposed to work for all  $w \in \mathcal{W}_{n+1}$  we repeat the argument as in Lemma 74 for a word in  $\mathcal{W}_{n+2}^c$  which is a concatenation of  $(n+1)$ -words by construction. By the same arguments as in Lemma 74 one can also show that  $g_s = \rho_{s+1, s}(g_{s+1})$  for all  $s \in \mathbb{N}$ .  $\square$

As in Subsection 9.3 we conclude that  $\mathbb{K}^c$  and  $(\mathbb{K}^c)^{-1}$  cannot be Kakutani equivalent.

**10.6. Proofs of Theorems 24 and 25.** Now we can complete the transfer to the setting of diffeomorphisms.

*Proof of Theorem 24.* In the proof of Theorem 21 we constructed a continuous reduction  $\Psi : \text{Trees} \rightarrow \mathcal{E}$  that took values in the strongly uniform odometer-based transformations. Under the functor  $\mathcal{F}$  from Subsection 10.3 these are transferred to strongly uniform circular systems. During the course of the construction several parameters with numerical conditions about growth and decay rates appear. In particular, at stage  $n$  the parameter  $l_n$  is chosen last. Hence, we can choose

it sufficiently large to guarantee condition (10.2), as well as the convergence to a  $C^\infty$  diffeomorphism that is measure-theoretically isomorphic to the circular system asserted in Lemma 77. We denote this realization map with our choice of the sequence  $(l_n)_{n \in \mathbb{N}}$  by  $R$ . Hereby, we define the map  $F^s : \mathcal{Trees} \rightarrow \text{Diff}^\infty(M, \lambda)$  by  $F^s = R \circ \mathcal{F} \circ \Psi$ . We want to show that  $F^s$  is a continuous reduction.

**Lemma.**  $F^s : \mathcal{Trees} \rightarrow \text{Diff}^\infty(M, \lambda)$  is continuous.

*Proof.* Let  $T = F^s(\mathcal{T})$  for  $\mathcal{T} \in \mathcal{Trees}$  and  $U$  be an open neighborhood of  $T$  in  $\text{Diff}^\infty(M, \lambda)$ . Let  $T^c = \mathcal{F} \circ \Psi(\mathcal{T})$  be the circular system such that  $R(T^c) = T$ . By [FW19a, Proposition 61] there is  $M \in \mathbb{N}$  sufficiently large such that for all  $S \in \text{Diff}^\infty(M, \lambda)$  in the range of  $R$  we have the following property: If  $(\mathcal{W}_n^c(T^c))_{n \leq M} = (\mathcal{W}_n^c(S^c))_{n \leq M}$ , then  $S \in U$ . Here,  $S^c$  denotes the circular system such that  $S = R(S^c)$ . Moreover,  $(\mathcal{W}_n(T^c))_{n \in \mathbb{N}}$  and  $(\mathcal{W}_n(S^c))_{n \in \mathbb{N}}$  denote the construction sequences of  $T^c$  and  $S^c$ , respectively. We recall from Subsection 10.3 that  $(\mathcal{W}_n(T^c))_{n \leq M}$  is determined by the first  $M + 1$  members in the construction sequence of the odometer based system  $F(\mathcal{T})$ , that is, it is determined by  $(\mathcal{W}_n(\mathcal{T}))_{n \leq M}$ . By (8.1) there is a basic open set  $V \subseteq \mathcal{Trees}$  containing  $\mathcal{T}$  such that for all  $\mathcal{S} \in V$  the first  $M + 1$  members of the construction sequences of  $\Psi(\mathcal{T})$  and  $\Psi(\mathcal{S})$  are the same, that is,  $(\mathcal{W}_n(\mathcal{T}))_{n \leq M} = (\mathcal{W}_n(\mathcal{S}))_{n \leq M}$ . Then it follows that  $F^s(\mathcal{S}) \in U$  for all  $\mathcal{S} \in V$  which yields the continuity of  $F^s$  by Fact 17.  $\square$

**Lemma.**  $F^s : \mathcal{Trees} \rightarrow \text{Diff}^\infty(M, \lambda)$  reduces the collection of ill-founded trees to the collection of ergodic  $C^\infty$  diffeomorphisms  $T$  such that  $T$  and  $T^{-1}$  are Kakutani equivalent.

*Proof.* Since the realization map  $R$  maps the circular system to a measure-theoretically isomorphic diffeomorphism, it preserves Kakutani equivalence. Hence, to see that  $F^s$  is a reduction, it suffices to check that  $\mathcal{F} \circ \Psi$  is a reduction. Here, we let  $\mathbb{K} = \Psi(\mathcal{T})$  and  $\mathbb{K}^c = \mathcal{F} \circ \Psi(\mathcal{T})$  for  $\mathcal{T} \in \mathcal{Trees}$ .

Suppose  $\mathcal{T} \in \mathcal{Trees}$  has an infinite branch. By the proof of part (1) of Proposition 23 in Section 5 there is an anti-synchronous isomorphism  $\phi : \mathbb{K} \rightarrow \mathbb{K}^{-1}$ . Then we can apply Lemma 82 on the functor  $\mathcal{F}$  and obtain the existence of an anti-synchronous isomorphism  $\phi^c : \mathbb{K}^c \rightarrow (\mathbb{K}^c)^{-1}$ . The converse direction follows from Subsection 10.5, where we have shown that if  $\mathcal{T} \in \mathcal{Trees}$  does not have an infinite branch, then  $\mathbb{K}^c$  and  $(\mathbb{K}^c)^{-1}$  are not Kakutani equivalent.  $\square$

Both statements together yield that  $F^s$  is a continuous reduction and we conclude the statement as in the proof of Theorem 20.  $\square$

In an analogous manner we can also prove Theorem 25.

*Proof of Theorem 25.* This time we let  $R$  be the realization map from circular systems with fast growing coefficients  $(l_n)_{n \in \mathbb{N}}$  to  $\text{Diff}_\rho^\omega(\mathbb{T}^2, \lambda)$  obtained by Lemma 78. Then the proof follows along the lines of the previous one, using [BKpp, Proposition 7.10] instead of [FW19a, Proposition 61].  $\square$

**Acknowledgements.** We thank M. Foreman and B. Weiss for helpful discussions. We especially thank Foreman for his encouragement and his detailed answers to our questions regarding his work with D. Rudolph and B. Weiss.

## REFERENCES

- [AK70] D. Anosov and A. Katok: New examples in smooth ergodic theory. Ergodic diffeomorphisms. Trudy Moskov. Mat. Obsc., 23: 3-36, 1970.
- [Ba17] S. Banerjee: Real-analytic AbC method on the torus. PhD thesis, available at <https://etda.libraries.psu.edu/catalog/14089sub232>
- [BKpp] S. Banerjee and P. Kunde: Real-analytic realization of uniform circular systems and some applications. Preprint, arXiv:1705.05079, to appear in J. Anal. Math.
- [BF96] F. Beleznyay and M. Foreman: The complexity of the collection of measure-distal transformations. Ergodic Theory Dynam. Systems, 16(5): 929-962, 1996.
- [Be15] M. Benhenda: An uncountable family of pairwise non-Kakutani equivalent smooth diffeomorphisms. J. Anal. Math. 127: 129-178, 2015.
- [Fe74] J. Feldman: Borel structures and invariants for measurable transformations. Proc. Amer. Math. Soc. 46: 383-394, 1974.
- [Fe76] J. Feldman: New K-automorphisms and a problem of Kakutani. Israel J. Math., 24 (1): 16-38, 1976.
- [FJR94] A. Fieldsteel, A. del Junco and D. Rudolph:  $\alpha$ -equivalence: a refinement of Kakutani equivalence. Ergodic Theory Dynam. Systems 14 (1): 69-102, 1994.
- [FR92] A. Fieldsteel and D. Rudolph: An ergodic transformation with trivial centralizer. Ergodic Theory Dynam. Systems 12: 459-478, 1992.
- [Fo10] M. Foreman: Models for measure preserving transformations, Topology and its Applications 157: 1404-1414, 2010.
- [Fo18] M. Foreman: What is a Borel reduction? Notices Amer. Math. Soc. 65 (10): 1263-1268, 2018.
- [FRW06] M. Foreman, D. Rudolph, and B. Weiss: On the conjugacy relation in ergodic theory. C. R. Math. Acad. Sci. Paris 343 (10): 653-656, 2006.
- [FRW11] M. Foreman, D. Rudolph, and B. Weiss: The conjugacy problem in ergodic theory. Ann. of Math. (2), 173(3): 1529-1586, 2011.
- [FW04] M. Foreman and B. Weiss: An anti-classification theorem for ergodic measure-preserving transformations. J. Eur. Math. Soc. 6, no. 3: 277-292, 2004.
- [FW19a] M. Foreman and B. Weiss: A symbolic representation for Anosov-Katok systems. J. Anal. Math., 137(2): 603-661, 2019.
- [FW19b] M. Foreman and B. Weiss: From odometers to circular Systems. J. Mod. Dyn., 15: 345-423, 2019.
- [FW3] M. Foreman and B. Weiss: Measure preserving diffeomorphisms of the torus are unclassifiable. Preprint, arXiv:1705.04414, to appear in J. Eur. Math. Soc.
- [FW4] M. Foreman and B. Weiss: Odometer based systems. Preprint, arXiv:2009.10162.
- [GKpp] M. Gerber and P. Kunde: Loosely Bernoulli odometer-based systems whose corresponding circular systems are not loosely Bernoulli. Preprint, arXiv:1912.12684, to appear in Ergodic Theory Dynam. Systems.
- [GLpp] E. Gardella and M. Lupini: The complexity of conjugacy, orbit equivalence, and von Neumann equivalence of actions of nonamenable groups. Preprint, arXiv:1708.01327.
- [HN42] P. Halmos and J. von Neumann: Operator methods in classical mechanics. II. Ann. of Math. (2), 43: 332-350, 1942.
- [Hj01] G. Hjorth: On invariants for measure preserving transformations. Fund. Math., 169(1): 51-84, 2001.
- [KVW21] A. Kanigowski, K. Vinhage and D. Wei: Kakutani equivalence of unipotent flows. Duke Math. J. 170 (7): 1517-1583, 2021.
- [Ka77] A. Katok: Monotone equivalence in ergodic theory. Mathematics of the USSR-Izvestiya 11 (1): 99-146, 1977.
- [Ke95] A. Kechris: Classical descriptive set theory, Graduate Texts in Mathematics, 156. Springer-Verlag, New York, 1995.
- [Ki86] J. King: The commutant is the weak closure of the powers, for rank-1 transformations. Ergodic Theory Dynam. Systems 6 (3): 363-384, 1986.
- [Ku65] A. Kushnirenko: An upper bound for the entropy of a classical dynamical system. Sov. Math. Dokl 6, 360-362, 1965.
- [Ne32] J. von Neumann: Zur Operatorenmethode in der klassischen Mechanik. Ann. of Math. (2), 33(3): 587 - 642, 1932.

- [Or70] D. Ornstein: Bernoulli shifts with the same entropy are isomorphic. *Advances in Math.* 4: 337 - 352, 1970.
- [ORW82] D. Ornstein, D. Rudolph and B. Weiss: Equivalence of measure preserving transformations. *Mem. Amer. Math. Soc.* 37, no. 262, 1982.
- [OS73] D. Ornstein and P. Shields: An uncountable family of K-automorphisms. *Advances in Math.* 10, 63-88, 1973.
- [Ra81] M. Ratner: Some invariants of Kakutani equivalence. *Israel J. Math.* 38 (3): 231-240, 1981.
- [VO16] M. Viana and K. Oliveira: *Foundations of ergodic theory*. Cambridge Studies in Advanced Mathematics, 151. Cambridge University Press, Cambridge, 2016.
- [We14] B. Weiss: On the classification of measure preserving diffeomorphisms of the torus. Talk at “Recent Progress in Dynamical Systems and Related Topics”, 2014, available at <http://www.birs.ca/events/2014/5-day-workshops/14w5058/videos/watch/201408130935-Weiss.html>