

SATO-TATE DISTRIBUTIONS OF CATALAN CURVES

HEIDI GOODSON

ABSTRACT. For distinct odd primes p and q , we define the Catalan curve $C_{p,q}$ by the affine equation $y^q = x^p - 1$. In this article we construct the Sato-Tate groups of the Jacobians in order to study the limiting distributions of coefficients of their normalized L-polynomials. Catalan Jacobians are nondegenerate and simple with noncyclic Galois groups (of the endomorphism fields over \mathbb{Q}), thus making them interesting varieties to study in the context of Sato-Tate groups. We compute both statistical and numerical moments for the limiting distributions. Lastly, we determine the Galois endomorphism types of the Jacobians using both old and new techniques.

1. INTRODUCTION

Let p and q be distinct odd primes. The nonsingular curve $C_{p,q}$ defined by the affine equation

$$y^q = x^p - 1$$

is called *Catalan curve*, likely named after the famous Catalan conjecture¹ regarding consecutive integers that are perfect powers. The primary goals of this article are to study the limiting distributions of coefficients of normalized L-polynomials of the Jacobians of Catalan curves and to study their real endomorphism algebras. These types of distributions are called *Sato-Tate distributions*, named for Mikio Sato and John Tate who independently made conjectures for those associated to elliptic curves in the 1960s. Their conjectures are known to be true for elliptic curves without complex multiplication defined over totally real fields and for all elliptic curves with complex multiplication.

Recent work has expanded this field to the study of distributions for abelian varieties of dimension at least 2. The generalized Sato-Tate conjecture predicts that the distributions converge to the distributions of traces in a compact Lie group referred to as the Sato-Tate group. The Sato-Tate group is related to the Mumford-Tate group, Hodge group, motivic Galois group, and l -adic monodromy group, thus placing this area of research at the intersection of many important fields in number theory, group theory, and algebraic geometry.

Before describing the work in higher genus, we first recall the original Sato-Tate conjecture for elliptic curves. Let F be a number field, E/F be an elliptic curve without complex multiplication (CM), and v be a finite prime of F such that E has good reduction at v . By a theorem of Hasse, the number of F_{q_v} points of E is $q_v + 1 - a_v$, where F_{q_v} denotes the residue field of v and a_v is an integer (called the trace of Frobenius) satisfying $|a_v| \leq 2q_v^{1/2}$. The Sato-Tate conjecture predicts that, as v varies through the primes of good reduction for E , the normalized Frobenius

2010 *Mathematics Subject Classification*. 11M50, 11G10, 11G20, 14G10.

Key words and phrases. Sato-Tate groups, Sato-Tate distributions, Jacobian varieties, endomorphism algebras.

¹The Catalan conjecture was proved by Mihăilescu [24] more than 150 years after Catalan published his conjecture.

traces $a_v/q_v^{1/2}$ are distributed in the interval $[-2, 2]$ with respect to the image of the Haar measure on the special unitary group $SU(2)$. This conjecture has recently been proved in 2008 for non-CM elliptic curves defined over totally real fields (see [6, 7, 17, 32]).

The distributions of the normalized Frobenius traces were known much earlier for CM elliptic curves over all fields: they are distributed with respect to the image of the Haar measure on either the unitary group $U(1)$ or the normalizer of $U(1)$ in $SU(2)$, depending on whether or not the field of definition contains the field of complex multiplication (see the exposition in [5]).

The generalized Sato-Tate conjecture for an abelian variety predicts the existence of a compact Lie group that determines the limiting distribution of normalized local Euler factors (see, for example, [31]). In our work, the abelian varieties will be the Jacobians of curves, and so we now state the conjecture specifically for Jacobian varieties.

Let C be a smooth, projective, genus g curve defined over a number field F . The Sato-Tate group of the Jacobian of C , $ST(\text{Jac}(C)) \subseteq \text{USp}(2g)$, is a compact Lie group satisfying the following property: for each prime \mathfrak{p} at which C has good reduction, there exists a conjugacy class of $ST(\text{Jac}(C))$ whose characteristic polynomial equals the normalized L-polynomial

$$\overline{L}_{\mathfrak{p}}(C, T) = T^{2g} + a_1 T^{2g-1} + a_2 T^{2g-2} + \cdots + a_{g-1} T^2 + a_g T + 1. \quad (1)$$

Conjecture 1.1. (*Generalized Sato-Tate Conjecture*) *Let $(x_{\mathfrak{p}})$ be the sequence of conjugacy classes of normalized images of Frobenius elements in $ST(\text{Jac}(C))$ at primes \mathfrak{p} of good reduction for $\text{Jac}(C)$, ordered by norm. Then the sequence $(x_{\mathfrak{p}})$ is equidistributed with respect to the pushforward of the Haar measure of $ST(\text{Jac}(C))$ to its space of conjugacy classes.*

The generalized Sato-Tate conjecture was proved for CM abelian varieties in [22]. Beyond proving the generalized Sato-Tate conjecture, it is also important to determine the explicit embedding of the Sato-Tate group inside $\text{USp}(2g)$ in order to provide an explicit description of the limiting distributions of the normalized L-polynomial. The distributions can also give us information about certain arithmetic invariants, such as the real endomorphism algebra and the Néron-Severi group of the abelian variety.

Determining these Sato-Tate groups is the source of ongoing interest and work. The current literature contains many articles for abelian varieties of fixed, small genus. For example, [11] and [12, 16] determine all possible Sato-Tate groups in dimension 2 and 3, respectively, by determining which subgroups of the unitary symplectic group satisfy certain axioms (see Section 2.4). Other articles determine Sato-Tate groups for certain families of genus 2 and 3 curves (see [15, 23]) or for twists of curves (see [3, 13, 14]).

As noted in [16], it is not expected, in dimension greater than 3, that every group satisfying the Sato-Tate axioms (see Section 2.4) can be realized using Jacobians of curves. There are currently two articles that study families of Jacobian varieties of arbitrarily high genus (see [9, 10]). In this article we provide a new example of an infinite family of Sato-Tate groups that can be realized by higher dimensional Jacobian varieties. Unlike the earlier articles, the component groups of these Sato-Tate groups are products of cyclic groups. Furthermore, we go a step further by computing the Galois endomorphism types of the Jacobians.

Organization of the paper. In Section 2 we provide an overview of this area of research, as well as some results that will be applied when computing the Sato-Tate groups of Jacobians of Catalan curves (Catalan Jacobians). We begin the section with a precise definition of the algebraic Sato-Tate group, which is needed in order to define the Sato-Tate group of an abelian variety. We then discuss nondegeneracy and its relationship to the Hodge conjecture in Section 2.2. In Section 2.3 we remind readers of the definition of the twisted Lefschetz group. We prove in Proposition 2.3 that, for Catalan Jacobians, this group is equal to the algebraic Sato-Tate group. We end the background section with a discussion of the Sato-Tate axioms in Section 2.4.

In Section 3 we define a generator for the group of endomorphisms of the Catalan Jacobian. Catalan curves have CM by the field $\mathbb{Q}(\zeta_{pq})$, and we study the action of $\text{Gal}(\mathbb{Q}(\zeta_{pq})/\mathbb{Q})$ generators on the endomorphism. We give complete descriptions of the actions in Propositions 3.4 and 3.5. These descriptions are needed in order to determine the twisted Lefschetz groups of Catalan Jacobians.

The main goal of Section 4 is to compute the Sato-Tate groups of Catalan Jacobians. Since the algebraic Sato-Tate conjecture for Catalan Jacobians is explained by endomorphisms, the component group of the Sato-Tate group is isomorphic to the Galois group $\text{Gal}(\mathbb{Q}(\zeta_{pq})/\mathbb{Q})$. This immediately tells us that the component group is a product of cyclic groups of orders $p - 1$ and $q - 1$. Still, we give explicit generators for the component group in Theorem 4.6 in order to obtain the limiting distribution of normalized local Euler factors.

There is an additional interesting result in Section 4. Lemma 4.2, which we use to determine component group generators, can be applied more broadly than the family of Jacobian varieties that we study in this paper. For any nondegenerate abelian variety with endomorphism given by a diagonal matrix and with CM by a cyclotomic field, this lemma can be used to determine the component group generators. This may seem like a very specific scenario, but we note that this describes the Jacobians considered in, for example, [9, 10, 23] and [15] (for certain values of c).

In Section 5, we compute moment statistics associated to the Sato-Tate groups of Catalan Jacobians. These moment statistics can be used to verify the equidistribution statement of the generalized Sato-Tate conjecture by comparing them to moment statistics obtained for the traces a_i in the normalized L -polynomial. The numerical moment statistics are an approximation since one can only ever compute them up to some prime. We list moments for some Catalan Jacobians in Table 5 of Section 5.

In Section 6, we study the Galois endomorphism types of Catalan Jacobians. There is a nice correspondence between the Sato-Tate group and the real endomorphism algebra for abelian varieties of dimension $g \leq 3$ (see Theorem 1.4 of [11]). This may not always extend to higher dimension, but it does for Catalan Jacobians and other nondegenerate abelian varieties. We approach this problem in two ways: through data obtained from moment statistics and by working with Rosati forms. The latter is the more traditional method of determining real endomorphism algebras, whereas the former is a new technique that uses recent results of Costa, Fité, and Sutherland [8]. We explore the limits of the four results of [8] and determine when they can be used to uniquely determine the endomorphism algebra from moment statistics.

Notation and conventions. We begin by fixing notation used in later sections. Let A be an abelian variety defined over a number field F . The ring of endomorphisms of A/F is denoted by $\text{End}(A_F)$, or simply $\text{End}(A)$ if the field of definition is clear from the context.

The curve $y^q = x^p - 1$ is denoted by $C_{p,q}$, and we assume throughout the paper that p and q are distinct odd primes. We will write ζ_m for a primitive m^{th} root of unity. For any rational number x whose denominator is coprime to an integer r , $\langle x \rangle_r$ denotes the unique representative of x modulo r between 0 and $r - 1$.

Let I denote the 2×2 identity matrix and define the matrices

$$J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (2)$$

and

$$Z = Z_{pq} = \text{diag}(\zeta_{p,q}, \overline{\zeta_{p,q}}) \quad (3)$$

The symplectic form considered throughout the paper is given by

$$H = \text{diag}(\underbrace{J, \dots, J}_g). \quad (4)$$

Lastly, for any positive integer n , we define the following subgroup of the unitary symplectic group $\text{USp}(2n)$

$$\text{U}(1)^n := \langle \text{diag}(u_1, \overline{u_1}, \dots, u_n, \overline{u_n}) : u_i \in \mathbb{C}^\times, |u_i| = 1 \rangle.$$

2. BACKGROUND

2.1. An ℓ -adic construction of the Sato-Tate group. We begin by defining both the algebraic Sato-Tate group and the Sato-Tate group. We follow the exposition of [9] and [31, Section 3.2]. See also [29, Chapter 8].

Let A/F be an algebraic variety of dimension g defined over the number field F . We define the Tate module $T_\ell := \varprojlim_n A[\ell^n]$, where ℓ is prime, to be a free \mathbb{Z}_ℓ -module of rank $2g$, and the rational Tate module $V_\ell := T_\ell \otimes_{\mathbb{Z}} \mathbb{Q}$ to be a \mathbb{Q}_ℓ -vector space of dimension $2g$. The Galois action on the Tate module is given by an ℓ -adic representation

$$\rho_{A,\ell} : \text{Gal}(\overline{F}/F) \rightarrow \text{Aut}(V_\ell) \cong \text{GL}_{2g}(\mathbb{Q}_\ell). \quad (5)$$

The ℓ -adic monodromy group of A , denoted $G_{A,\ell}$, is the Zariski closure of the image of this map in $\text{GL}_{2g}(\mathbb{Q}_\ell)$, and we define $G_{A,\ell}^1 := G_{A,\ell} \cap \text{Sp}_{2g}(\mathbb{Q}_\ell)$. Banaszak and Kedlaya proposed in [5, Conjecture 2.1] the following conjecture that was partly intended to be a refinement of the Mumford-Tate conjecture.

Conjecture 2.1 (Algebraic Sato-Tate Conjecture). *There is an algebraic subgroup $\text{AST}(A)$ of GSp_{2g} over \mathbb{Q} , called the **algebraic Sato-Tate group of A** , such that $\text{AST}^0(A)$ is reductive and, for each prime ℓ , $G_{A,\ell}^1 = \text{AST}(A) \otimes_{\mathbb{Q}} \mathbb{Q}_\ell$.*

With this set up, we define the **Sato-Tate group** of A , denoted $\text{ST}(A)$, to be a maximal compact Lie subgroup of $G_{A,\ell}^1 \otimes_{\mathbb{Q}_\ell} \mathbb{C} = \text{AST}(A) \otimes_{\mathbb{Q}} \mathbb{C}$ contained in $\text{USp}(2g)$. It is conjectured that $\text{ST}(A)$ is, up to conjugacy in $\text{USp}(2g)$, independent of the choice of the prime ℓ and of the embedding of \mathbb{Q}_ℓ in \mathbb{C} and so we will refer to $\text{ST}(A)$ as *the Sato-Tate group of A* (see, for example, [31]). While the Sato-Tate group is a compact Lie group, it may not be connected [15]. We denote the connected component of the identity (also called the identity component) of $\text{ST}(A)$ by $\text{ST}^0(A)$.

2.2. Nondegeneracy. Let A be a nonsingular projective variety over \mathbb{C} . We denote the (complexified) Hodge ring of A by

$$\mathcal{B}^*(A) := \sum_{d=0}^{\dim(A)} \mathcal{B}^d(A),$$

where $\mathcal{B}^d(A) = (H^{2d}(A, \mathbb{Q}) \cap H^{d,d}(A)) \otimes \mathbb{C}$ is the \mathbb{C} -span of Hodge cycles of codimension d on A . Furthermore, we define the ring

$$\mathcal{D}^*(A) := \sum_{d=0}^{\dim(A)} \mathcal{D}^d(A)$$

where $\mathcal{D}^d(A)$ is the \mathbb{C} -span of classes of intersection of d -divisors. This is the subring of $\mathcal{B}^*(A)$ generated by the divisor classes, i.e. generated by $\mathcal{B}^1(A)$. In general, it is known that we have containment $\mathcal{D}^*(A) \subseteq \mathcal{B}^*(A)$, and we say that an abelian variety A is **nondegenerate** if we have equality, i.e. $\mathcal{D}^*(A) = \mathcal{B}^*(A)$ (see [1]). Furthermore, an abelian variety A is said to be **stably nondegenerate** if, for any integer $k \geq 1$, $\mathcal{D}^*(A^k) = \mathcal{B}^*(A^k)$ [1, 20]. If $\mathcal{B}^*(A)$ is not generated by the divisor classes $\mathcal{B}^1(A)$, i.e. $\mathcal{D}^*(A) \neq \mathcal{B}^*(A)$, then A is said to be **degenerate**. The additional Hodge cycles that are not generated by divisor classes are referred to as **exceptional cycles** or **exceptional classes** (see, for example, [25]).

While nondegeneracy extends to powers of stably nondegenerate abelian varieties, it does not necessarily apply to products of nonisogenous varieties. For example, Shioda demonstrates in [30] that the Jacobian of the genus 4 curve $y^2 = x^9 - 1$ is degenerate, though it is the product of two simple, nondegenerate varieties. Aoki proves in [2] that generalized Catalan curves $C_{p^\mu, q^\nu} : y^{q^\nu} = x^{p^\mu} - 1$, where $(\mu, \nu) \neq (1, 1)$, also exhibit this phenomenon: each of the factors are nondegenerate but C_{p^μ, q^ν} is itself degenerate. Hazama explains in [20] that this can occur when one of the simple factors is of type-IV in the Albert's classification.

Hazama proves in Theorem 1.2 of [20] that A is stably nondegenerate if and only if the dimension of its Hodge group is maximal². When A is an abelian variety with complex CM and is absolutely simple, this is equivalent to saying that the CM type is nondegenerate (see, for example, [1, 10]). The following result is proved in [21].

Proposition 2.2. *The Jacobian variety $\text{Jac}(C_{p,q})$ of the Catalan curve is absolutely simple and nondegenerate.*

Hazama proves the nondegeneracy by proving that the CM-type is nondegenerate. This is done by verifying the non-vanishing of certain character sums attached to the Jacobian.

Nondegeneracy is related to the Hodge conjecture. Let $\mathcal{C}^d(A)$ be the subspace of $\mathcal{B}^d(A)$ generated by the classes of algebraic cycles on A of codimension d . Then

$$\mathcal{D}^d(A) \subseteq \mathcal{C}^d(A) \subseteq \mathcal{B}^d(A)$$

and the **Hodge conjecture** for A asserts that $\mathcal{C}^d(A) = \mathcal{B}^d(A)$ for all d [1, 30]. It is clear from the definition that if A is nondegenerate then the Hodge conjecture holds, and if the Hodge conjecture does not hold then A must be degenerate. However, there are many cases where the Hodge conjecture holds for degenerate abelian varieties (see, for example, [26, 30]).

²In [10], the authors use the word *nondegenerate* to describe abelian varieties with this property.

2.3. The Twisted Lefschetz group. Banaszak and Kedlaya introduced in [5] the **twisted Lefschetz group**, denoted $\mathrm{TL}(A)$, which is a closed algebraic subgroup of Sp_{2g} defined by

$$\mathrm{TL}(A) := \bigcup_{\tau \in \mathrm{Gal}(\overline{F}/F)} \mathrm{L}(A)(\tau), \quad (6)$$

where $\mathrm{L}(A)(\tau) := \{\gamma \in \mathrm{Sp}_{2g} \mid \gamma\alpha\gamma^{-1} = \tau(\alpha) \text{ for all } \alpha \in \mathrm{End}(A_{\overline{F}})_{\mathbb{Q}}\}$. When τ is the identity automorphism, $\mathrm{L}(A)(\tau)$ forms a group, called the Lefschetz group, which is denoted in the literature as simply $\mathrm{L}(A)$.

Banaszak and Kedlaya prove in [5, Theorem 6.1] that if the Mumford-Tate conjecture is explained by endomorphisms and the twisted Lefschetz group over \overline{F} is connected, then the algebraic Sato-Tate conjecture holds and $\mathrm{AST}(A) = \mathrm{TL}(A)$. In this case, we say that the algebraic Sato-Tate conjecture is also explained by endomorphisms. It follows from [5, Theorem 6.1] and work of Serre in [29, Section 8.3.4] that when the algebraic Sato-Tate conjecture is explained by endomorphisms the component group $\mathrm{ST}(A)/\mathrm{ST}^0(A)$ is isomorphic to the Galois group $\mathrm{Gal}(K/F)$, where K is the **endomorphism field** of A , i.e. the minimal extension over which all the endomorphisms of the abelian variety A are defined (see, for example, [31, Theorem 3.12] and [11, Proposition 2.17]). We can use the twisted Lefschetz group to write down explicit generators of the component group of the Sato-Tate group.

The Jacobian varieties we study in this paper satisfy these conditions, and so we have the following result.

Proposition 2.3. *The algebraic Sato-Tate Conjecture holds for the Catalan curve $\mathrm{Jac}(C_{p,q})$ with $\mathrm{AST}(\mathrm{Jac}(C_{p,q})) = \mathrm{TL}(\mathrm{Jac}(C_{p,q}))$.*

Proof. In Proposition 2.2 we saw that $\mathrm{Jac}(C_{p,q})$ is a nondegenerate CM abelian variety. The result then follows from Theorems 6.1 and 6.6 in [5]. \square

2.4. The Sato-Tate axioms. We conclude the background section by recalling some necessary conditions for a subgroup of $\mathrm{USp}(2g)$ to occur as a Sato-Tate group. These axioms will not be used in our work, but they are useful when attempting to classify all possible Sato-Tate groups for a given dimension g (see, for example, [11, 12, 16]). We state the axioms as they are laid out in [16]. For a group G with identity component G^0 , the Sato-Tate group axioms in dimension g are as follows.

- (ST1) The group G is a closed subgroup of $\mathrm{USp}(2g)$.
- (ST2) (Hodge condition) There exists a homomorphism $\theta: \mathrm{U}(1) \rightarrow G^0$ such that $\theta(u)$ has eigenvalues u, u^{-1} each with multiplicity g . The image of such a θ is called a *Hodge circle*, and the set of all Hodge circles generates a dense subgroup of G^0 .
- (ST3) (Rationality condition) For each component H of G and irreducible character χ of $\mathrm{GL}(\mathbb{C}^{2g})$, the expected value (under the Haar measure) of $\chi(\gamma)$ over $\gamma \in H$ is an integer.
- (ST4) (Lefschetz condition) The subgroup of $\mathrm{USp}(2g)$ fixing $\mathrm{End}(\mathbb{C}^{2g})^{G^0}$ is equal to G^0 .

Proposition 3.2 of [11] proves that, for any abelian variety satisfying both the Mumford-Tate conjecture and the algebraic Sato-Tate conjecture, the Sato-Tate group $G = \mathrm{ST}(A)$ satisfies the first three axioms. The last axiom is satisfied if, in addition, the Hodge group of A equals the Lefschetz group of A (see Proposition 3.3 of [16]). Catalan Jacobians satisfy these properties, and so their Sato-Tate groups satisfy the four axioms.

3. ENDOMORPHISMS OF CATALAN JACOBIANS

Let $p \neq q$ be odd primes. The Jacobian of $C_{p,q}$ has complex multiplication by the field $K = \mathbb{Q}_{pq}$, and so the Galois group $\text{Gal}(K/\mathbb{Q})$ is isomorphic to a product of cyclic groups $\mathbb{Z}/p\mathbb{Z}^\times \times \mathbb{Z}/q\mathbb{Z}^\times$. The main goals of this section are to give an explicit model of a generator for the group of endomorphisms of $\text{Jac}(C_{p,q})$ and to describe how the Galois group acts on the endomorphism. We begin by defining some necessary notation that will be used when defining both the endomorphism and the Sato-Tate group of $\text{Jac}(C_{p,q})$.

3.1. Galois elements and some notation. Let c be a generator for $\mathbb{Z}/p\mathbb{Z}^\times$ and d be a generator for $\mathbb{Z}/q\mathbb{Z}^\times$. These elements yield the following generators of $\text{Gal}(K/\mathbb{Q})$:

$$\sigma_p: \begin{cases} \zeta_p \mapsto \zeta_p^c \\ \zeta_q \mapsto \zeta_q \end{cases} \qquad \sigma_q: \begin{cases} \zeta_p \mapsto \zeta_p \\ \zeta_q \mapsto \zeta_q^d. \end{cases}$$

In this paper, we will be working with powers of ζ_{pq} of the form $\zeta_{pq}^{q(a+1)-pb}$. We compute the actions of the Galois elements on these powers of ζ_{pq} to be

$$\sigma_p \left(\zeta_{pq}^{q(a+1)-pb} \right) = \zeta_{pq}^{qc(a+1)-pb} \quad \text{and} \quad \sigma_q \left(\zeta_{pq}^{q(a+1)-pb} \right) = \zeta_{pq}^{q(a+1)-pdb}. \quad (7)$$

We now define some additional notation. Let $k_0 = 0$ and for any b satisfying $1 \leq b \leq q-1$, define

$$k_b = \left\lfloor \frac{pb - q - 1}{q} \right\rfloor. \quad (8)$$

The k_b -values are increasing with respect to b : $k_{b_1} \leq k_{b_2}$ whenever $b_1 \leq b_2$. When $p < q$, the value k_b may be negative for some values of b , and we let κ_t be the sum

$$\kappa_t = \sum_{\substack{b=0 \\ k_b \geq 0}}^t (k_b + 1), \quad (9)$$

where we restrict to adding only the nonnegative values k_b .

Remark 1. The quantities in Equations (8) and (9) are relevant to our work in the following way. The block entries of the endomorphism of $\text{Jac}(C_{p,q})$ will be defined in terms of combinations of values of b and a (defined below in Inequalities (10) and (11), respectively). The number of a -values paired with a particular b -value will be $k_b + 1$ whenever $k_b \geq 0$. The value κ_b will help us keep track of the columns associated to particular (a, b) -pairings. Furthermore, κ_{q-1} gives the total number of (a, b) -pairings, and so it equals the genus of $C_{p,q}$.

3.2. Defining the endomorphism. Let $\alpha: C_{p,q} \rightarrow C_{p,q}$ be the curve automorphism defined by $\alpha(x, y) = (\zeta_p x, \zeta_q y) = (\zeta_{pq}^q x, \zeta_{pq}^p y)$. This automorphism has order pq , and so it is a generator of the automorphism group of $C_{p,q}$.

The genus of $C_{p,q}$ is $g = (p-1)(q-1)/2$ and we take as a basis for the space of regular 1-forms $\Omega^1(C_{p,q})$ the set $\{\omega_{a,b} = \frac{x^a dx}{y^b}\}_{a,b}$ where a and b satisfy

$$1 \leq b \leq q-1 \quad (10)$$

$$0 \leq a < p \quad \text{and} \quad a \leq k_b. \quad (11)$$

We compute pullbacks of the differentials with respect to α in order to determine the associated endomorphism α of $\text{Jac}(C_{p,q})$:

$$\alpha^*(\omega_{a,b}) = \zeta_{pq}^{q(a+1)-pb} \omega_{a,b}.$$

In order to determine explicit generators for the Sato-Tate group, we must write an explicit expression for the endomorphism α of $\text{Jac}(C_{p,q})$. We start by choosing an ordering of the basis elements. Start with $b = 1$ and list all $\omega_{a,b}$, where the a are increasing and satisfy the inequalities in (11). Increment b and repeat this process while Equation (10) is still satisfied. Any choice that is made for the ordering of the basis elements of $\Omega^1(C_{p,q})$ will lead to an isomorphic Sato-Tate group of $\text{Jac}(C_{p,q})$. We can now define the induced endomorphism α of $H_1(\text{Jac}(C_{p,q})_{\mathbb{C}}, \mathbb{Q})$.

Definition 3.1. *By taking the symplectic basis of $H_1(\text{Jac}(C_{p,q})_{\mathbb{C}}, \mathbb{C})$ (with respect to the matrix $H = \text{diag}(J, \dots, J)$) given above, α is the diagonal matrix whose i^{th} diagonal 2×2 block is*

$$\alpha[i] = Z^{q(a_i+1)-pb_i},$$

where $a_i = i - \kappa_{b_i-1} - 1$ and b_i is the largest integer satisfying (10) such that $a_i > 0$.

Example 3.2. *Consider the genus 6 curve $C_{7,3}: y^3 = x^7 - 1$. Using the strategy described above, we write the basis of $\Omega^1(C_{p,q})$ as $\{\omega_{0,1}, \omega_{1,1}, \omega_{0,2}, \omega_{1,2}, \omega_{2,2}, \omega_{3,2}\}$ and the endomorphism α as*

$$\alpha = \text{diag}(Z^{q-p}, Z^{2q-p}, Z^{q-2p}, Z^{2q-2p}, Z^{3q-2p}, Z^{4q-2p}).$$

3.3. The actions of the Galois elements on α . The main results of this subsection are Propositions 3.4 and 3.5, where we describe the action of the Galois elements on the endomorphism α . The following lemma will be referenced in the proofs of these propositions.

Lemma 3.3. *Let β be a positive integer satisfying Inequality (10), and define two quantities $y_1 = (p(q - \beta) - q + 1)/q$ and $y_2 = (p(q - \beta) - q - 1)/q$. Then any integer λ satisfying $\lambda < y_1$ also satisfies $\lambda \leq \lfloor y_2 \rfloor$.*

Proof. The quantities y_1 and y_2 differ by $2/q$: $y_2 = y_1 - 2/q$. Let λ be any integer less than y_1 . We split the proof of the lemma into two cases: $y_1 \in \mathbb{Z}$ and $y_1 \notin \mathbb{Z}$.

First suppose that $y_1 \in \mathbb{Z}$ so that $\lambda \leq y_1 - 1$. Since $0 < 2/q < 1$, the inequalities $y_1 - 1 < y_1 - 2/q < y_1$ hold. Hence, $y_1 - 1 = \lfloor y_1 - 2/q \rfloor = \lfloor y_2 \rfloor$, and so $\lambda \leq \lfloor y_2 \rfloor$.

Suppose instead that $y_1 \notin \mathbb{Z}$. We claim that $y_1 - 1/q$ is also not an integer. Observe that

$$y_1 - \frac{1}{q} = \frac{p(q - \beta) - q}{q} = p - 1 - \frac{p\beta}{q}.$$

Since both p and β are relatively prime to q , the fractional term $\frac{p\beta}{q}$ and, hence, $y_1 - 1/q$ are not integers. This implies that $y_1 > \lfloor y_1 - 1/q \rfloor = \lfloor y_1 - 2/q \rfloor = \lfloor y_2 \rfloor$. Combining this with the requirement that λ is an integer less than y_1 yields the desired result of $\lambda \leq \lfloor y_2 \rfloor$. \square

We will now give a complete description of the action of σ_q on the endomorphism α . We will do this by studying the action on an arbitrary diagonal block entry of the matrix. Using the action described in Equation (7), we see that

$$\sigma_q \alpha[i] = Z^{q(a_i+1)-p\beta b_i}.$$

A particularly nice property of this action is that $\sigma_q \alpha[i]$ is either equal to or is the conjugate of some block diagonal entry of the matrix α (we denote the conjugate of $\alpha[j]$ by $\overline{\alpha[j]}$).

Proposition 3.4. *Let d be a generator of the cyclic group $\mathbb{Z}/q\mathbb{Z}^\times$ and let $t_i = \langle db_i \rangle_q$. Define the quantities $\iota_1 = \kappa_{t_i-1} + a_i + 1$ and $\iota_2 = \kappa_{t'_i-1} + p - (a_i + 1)$, where $t'_i = q - t_i$. Then $\sigma^q \alpha$ is the diagonal matrix whose i^{th} diagonal 2×2 block is*

$$\sigma^q \alpha[i] = \begin{cases} \alpha[\iota_1] & \text{if } 0 \leq a_i \leq k_{t_i}, \\ \alpha[\iota_2] & \text{if } a_i > k_{t_i}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. First suppose that a_i satisfies $0 \leq a_i \leq k_{t_i}$. Then t_i and a_i together satisfy the bounds of Equations (10) and (11). Hence, $\sigma^q \alpha[i] = Z^{q(a_i+1)-pt_i} = \alpha[\iota]$, for some ι . Comparing this to the definition of κ and doing a bit of arithmetic shows that $\iota = \iota_1$.

Suppose instead that a_i satisfies

$$a_i \geq k_{t_i} + 1 > \frac{pt_i - q - 1}{q}.$$

Note that this includes the case where $k_{t_i} < 0$. Then t_i and a_i do not satisfy the bounds of Equations (10) and (11). Let $a'_i = p - a_i - 2$ and $t'_i = q - t_i$. Then

$$Z^{q(a_i+1)-pt_i} = \overline{Z^{q(a'_i+1)-pt'_i}}.$$

The bounds on t_i imply that t'_i satisfies $1 \leq t'_i \leq q - 1$ and the bounds on a_i imply

$$\begin{aligned} a'_i &< p - \frac{pt_i - q - 1}{q} - 2 \\ &= \frac{p(q - t_i) - q + 1}{q} \\ &= \frac{pt'_i - q + 1}{q}. \end{aligned}$$

By Lemma 3.3, we can conclude that $a'_i \leq \lfloor (pt'_i - q - 1)/q \rfloor = k_{t'_i}$. Thus, a'_i and t'_i together satisfy the bounds of Inequalities (10) and (11). Hence, $Z^{q(a'_i+1)-pt'_i}$ appears as a block entry of α and

$$\sigma^q \alpha[i] = \overline{Z^{q(a'_i+1)-pt'_i}} = \overline{\alpha[\iota]}$$

for some ι . Comparing this to the definition of κ and working out a bit of arithmetic shows that $\iota = \iota_2$. □

We can use a similar strategy to describe of the action of σ_p on the endomorphism α . Using the action described in Equation (7), we see that

$$\sigma_p \alpha[i] = Z^{qc(a_i+1)-pb_i}.$$

This leads to the following proposition.

Proposition 3.5. *Let c be a generator of the cyclic group $\mathbb{Z}/p\mathbb{Z}^\times$ and let $s_i = \langle c(a_i + 1) \rangle_p - 1$. Let $\iota_1 = \kappa_{b_i-1} + s_i + 1$ and $\iota_2 = \kappa_{b'_i-1} + p - (s_i + 1)$, where $b'_i = q - b_i$. Then $\sigma^q \alpha$ is the diagonal matrix whose i^{th} diagonal 2×2 block is*

$$\sigma^p \alpha[i] = \begin{cases} \alpha[\iota_1] & \text{if } 0 \leq s_i \leq k_{b_i}, \\ \alpha[\iota_2] & \text{if } s_i > k_{b_i}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The bounds for s_i are $0 \leq s_i < p - 1$, and $k_{b_i} \geq 0$ since it is associated to a pair a_i, b_i that appear in an entry of the endomorphism α . We follow the method of proof for Proposition 3.4 and split into two cases.

First suppose that s_i satisfies $0 \leq s_i \leq k_{b_i}$. Then b_i and s_i together satisfy the bounds of Equations (10) and (11). Hence, $\sigma_p \alpha[i] = Z^{q(s_i+1)-pb_i} = \alpha[\iota]$, for $\iota = \iota_1$.

Suppose instead that s_i satisfies

$$s_i \geq k_{b_i} + 1 > \frac{pb_i - q - 1}{q}.$$

Let $s'_i = p - s_i - 2$ and $b'_i = q - b_i$. Then $Z^{q(s_i+1)-pb_i} = \overline{Z^{q(s'_i+1)-pb'_i}}$. The bounds on s_i imply that

$$\begin{aligned} s'_i &< p - \frac{pb_i - q - 1}{q} - 2 \\ &= \frac{p(q - b_i) - q + 1}{q} \\ &= \frac{pb'_i - q + 1}{q}. \end{aligned}$$

By Lemma 3.3, this proves that $s'_i \leq [(pb'_i - q - 1)/q] = k_{b'_i}$. Thus, b'_i and s'_i together satisfy the bounds of Equations (10) and (11). Hence, $Z^{q(s'_i+1)-pb'_i}$ appears as a block entry of α and

$$\sigma_q \alpha[i] = \overline{Z^{q(s'_i+1)-pb'_i}}$$

which is the conjugate of the block $\alpha[\iota]$, for $\iota = \kappa_{b'_i-1} + s'_i + 1 = \kappa_{b'_i-1} + p - s_i - 1 = \iota_2$. \square

Example 3.6. We return to the curve $C_{7,3}$ from Example 3.2. The action of σ_q (using $d = 2$) on α yields the diagonal matrix

$$\sigma_q \alpha = \text{diag}(Z^{q-2p}, Z^{2q-2p}, Z^{q-p}, Z^{2q-p}, \overline{Z^{4q-2p}}, \overline{Z^{3q-2p}}).$$

The action of σ_p (using $c = 3$) on α yields

$$\sigma_p \alpha = \text{diag}(\overline{Z^{4q-2p}}, \overline{Z^{q-2p}}, Z^{3q-2p}, \overline{Z^{q-p}}, Z^{2q-2p}, \overline{Z^{2q-p}}).$$

4. THE SATO-TATE GROUP OF $\text{Jac}(C_{p,q})$

We begin with a definition and a lemma that we need in order to define the component group and to prove Theorem 4.6.

4.1. Preliminaries.

Definition 4.1. A *signed permutation matrix* is square matrix that has exactly one entry of 1 or -1 in each row and each column and 0s everywhere else. We will define a **block signed permutation matrix** to be an even dimension signed permutation matrix that is partitioned into 2×2 blocks that are either I , J , or the zero matrix.

For any matrix partitioned X into 2×2 blocks, we will let $X[i, j]$ be the block in the i^{th} row partition and j^{th} column partition.

The general strategy for writing down a γ associated to a Galois element σ is described in the following lemma.

Lemma 4.2. *Let A be a diagonal matrix partitioned into 2×2 blocks and let B be a block signed permutation matrix. Then BAB^{-1} is a diagonal matrix. Furthermore, if $B[i, j] = I$ then $BAB^{-1}[i, i] = A[j, j]$ and if $B[i, j] = J$ then $BAB^{-1}[i, i] = -JA[j, j]J$.*

Proof. The proof requires little more than elementary linear algebra. The first statement, that BAB^{-1} is a diagonal matrix, comes from the fact that A is diagonal and B is a block signed permutation matrix.

Now note that the block entry $B^{-1}[j, i]$ is the inverse of the block $B[i, j]$. The i^{th} diagonal block entry of BAB^{-1} is the product

$$BAB^{-1}[i, i] = B[i, j] \cdot A[j, j] \cdot B^{-1}[j, i].$$

Thus, if $B[i, j] = I$ then the i^{th} entry of BAB^{-1} is $A[j, j]$. If, instead, $B[i, j] = J$ then $BAB^{-1}[i, i] = JA[j, j]J^{-1} = -JA[j, j]J$. \square

Remark 2. As noted in the Introduction, this lemma can be applied to more varieties than those we study in this paper. For example, this lemma could be used for the varieties studied in [9, 10, 23] and [15] (for certain values of c).

4.2. The identity component and the component group. We begin this subsection with a result about the identity component of the Sato-Tate group.

Proposition 4.3. *The identity component of the Sato-Group of $\text{Jac}(C_{p,q})$ is*

$$\text{ST}^0(\text{Jac}(C_{p,q})) \simeq \text{U}(1)^g,$$

where $g = (p-1)(q-1)/2$ is the genus of $C_{p,q}$.

Proof. Recall the definition of the algebraic Sato-Tate group $\text{AST}(\text{Jac}(C_{p,q}))$ in Conjecture 2.1. If we consider the same construction but restrict the domain of the ℓ -adic representation $\rho_{A,\ell}$ to $\text{Gal}(\overline{\mathbb{Q}}/K)$, where K is the CM field of $\text{Jac}(C_{p,q})$, we obtain the identity component of $\text{AST}(\text{Jac}(C_{p,q}))$. The identity component of the Sato-Tate group of $\text{Jac}(C_{p,q})$ is a maximal compact subgroup of this.

This restricted map is exactly the map that appears in Theorem A of [4], and so we can apply that result to our situation. The theorem states that the image of the restricted map is contained in the group of diagonal matrices of the form

$$\{\text{diag}(x_1, y_1, \dots, x_g, y_g) \mid x_i, y_i \in \mathbb{Q}_\ell^\times, x_1 y_1 = \dots = x_g y_g = 1\}.$$

The Jacobian of $C_{p,q}$ is absolutely simple and nondegenerate (see Proposition 2.2), and so, as in [9, 10], we can go one step further and conclude that the containment is actually an equality. It follows that we can choose $\text{ST}^0(\text{Jac}(C_{p,q}))$ to be the maximal compact subgroup $\text{U}(1)^g$. \square

We now define two matrices. We will then prove that they form a generating set for the component group of the Sato-Tate group. For the following definitions, let g be the genus of $C_{p,q}$.

Definition 4.4. *Let d be a generator of the cyclic group $\mathbb{Z}/q\mathbb{Z}^\times$. For any integer i satisfying $1 \leq i \leq g$, let $t_i = \langle db_i \rangle_q$ and $r_i = i - \kappa_{b_i-1} - 1$, where b_i is the largest positive integer such that r_i is nonnegative. Define γ_q to be the $2g \times 2g$ block signed permutation matrix whose ij^{th} block is*

$$\gamma_q[i, j] = \begin{cases} I & \text{if } j = \kappa_{t_i-1} + r_i + 1 \text{ and } 0 \leq r_i \leq k_{t_i}, \\ J & \text{if } j = \kappa_{t_i-1} + p - (r_i + 1), \text{ and } r_i > k_{t_i}, \\ 0 & \text{otherwise,} \end{cases}$$

where $t'_i = q - t_i$.

Definition 4.5. Let c be a generator of the cyclic group $\mathbb{Z}/p\mathbb{Z}^\times$, r_i and b_i be as defined in Definition 4.4, and $s_i = \langle cr_i \rangle_p - 1$. Define γ_p to be the $2g \times 2g$ block signed permutation matrix whose ij^{th} block is

$$\gamma_p[i, j] = \begin{cases} I & \text{if } j = \kappa_{b_{i-1}} + s_i + 1 \text{ and } 0 \leq s_i \leq \kappa_{b_i}, \\ J & \text{if } j = \kappa_{b'_{i-1}} + p - (s_i + 1) \text{ and } s_i > \kappa_{b_i}, \\ 0 & \text{otherwise,} \end{cases}$$

where $b'_i = q - b_i$.

Theorem 4.6. Let $q \neq p$ be odd primes and γ_q, γ_p be as defined in Definitions 4.4 and 4.5. The Sato-Tate group of the Jacobian of the genus g curve $C_{p,q}$ is

$$\text{ST}(\text{Jac}(C_{p,q})) \simeq \langle U(1)^g, \gamma_q, \gamma_p \rangle.$$

Proof. We proved in Proposition 4.3 that the identity component $\text{ST}^0(\text{Jac}(C_{p,q}))$ is $U(1)^g$. We will now prove the claim regarding the component group.

It follows from Proposition 2.3 that we can use the twisted Lefschetz group of $\text{Jac}(C_{p,q})$ to write down explicit generators of the component group. Choose generators c and d of $\mathbb{Z}/p\mathbb{Z}^\times$ and $\mathbb{Z}/q\mathbb{Z}^\times$, respectively. Propositions 3.4 and 3.5 show that the actions of the Galois elements simply rearrange and sometimes conjugate the diagonal entries of the endomorphism α . Since $-JZJ = \overline{Z}$, where Z is defined in Equation (3), we can apply Lemma 4.2 to determine the component group generators.

From here, we simply compare Definition 4.4 to Proposition 3.4 and Definition 4.5 to Proposition 3.5. For the former, $r_i = a_i$ and so the definition of γ_q yields the correct rearranging and conjugating as determined by the action of σ_q . For the latter, we used the same notation and so it is clear that the definition of γ_p yields the rearranging and conjugating determined by the action of σ_p . Thus, γ_q and γ_p are generators of $\text{TL}(\text{Jac}(C_{p,q}))$. \square

The description of γ_q simplifies rather nicely in the case where $q = 3$, and so we present the following corollary.

Corollary 4.7. Let $q = 3$, $p > 3$ be prime, and $m = \lfloor \frac{p-4}{3} \rfloor + 1$. Then the component group generator γ_q is

$$\gamma_q = \left(\begin{array}{c|c|c} & I_{2m} & \\ \hline I_{2m} & & \\ \hline & & J_{g-2m} \end{array} \right),$$

where I_{2m} is the $2m \times 2m$ identity matrix and $J_{g-2m} = \underbrace{\text{antidiag}(J, J, \dots, J)}_{g-2m}$.

Proof. If $q = 3$, then the genus is $g = p - 1$ and the values of b satisfying Equation (10) are $b_1 = 1$ and $b_2 = 2$. We choose $d = 2$ as the generator of $\mathbb{Z}/q\mathbb{Z}^\times$, which leads to $t_1 = 2$ and $t_2 = 1$. Furthermore, $k_1 = m - 1$, $\kappa_1 = m$, and $k_2 = \lfloor (2p - 4)/3 \rfloor$.

The values of i associated to b_1 are $1 \leq i \leq m$ (so that $r_i \geq 0$). For these values of i , $0 \leq r_i \leq$ is always satisfied. Hence, $\gamma_q[i, j] = I$ when

$$j = \kappa_1 + r_i + 1 = m + i.$$

This verifies that the I_{2m} in the top rows of γ_q in the statement of the corollary are correct.

The values of i associated to b_2 are then $m + 1 \leq i \leq g = p - 1$. The associated values are r_i are split into two intervals: $0 \leq r_i \leq m - 1$ and $r_i \geq m$. In the first interval we have $m + 1 \leq i \leq 2m$, which leads to $\gamma_q[i, j] = I$ when $j = i - m$. This verifies that the I_{2m} in the middle rows of γ_q in the statement of the corollary are correct.

In the second interval, $r_i \geq m$ and we need to determine which values of j will lead to $\gamma_q[i, j] = J$. Let $i = 2m + h$, where $1 \leq h \leq g - 2m$, so that $r_i = m + h - 1$. Then the value of j we need is

$$j = m + p - (m + h) = p - h = g - (h - 1).$$

Thus, the J blocks will start in row $i = 2m + 1$, column $j = g$, and will then cascade down and to the left. This verifies that the J_{g-2m} in the bottom rows of γ_q in the statement of the corollary are correct. \square

Remark 3. See the Appendix for examples of the component group generators.

5. MOMENT STATISTICS

In this section, we describe the distributions of the coefficients of the characteristic polynomial of random conjugacy classes in the Sato-Tate groups in Theorem 4.6. For comparison, we compute the numerical a_1 -moments of the normalized L-polynomial for some small genus Catalan curves.

5.1. Preliminaries. The following background information has been adapted from [9, Section 6].

We start by recalling some basic properties of moment statistics. We define the n^{th} moment (centered at 0) of a probability density function to be the expected value of the n^{th} power of the values, i.e. $M_n[X] = E[X^n]$. Recall that for independent variables X and Y we have $E[X + Y] = E[X] + E[Y]$ and $E[XY] = E[X]E[Y]$ (see, for example, [23]). This yields the following identities for moment statistics: $M_n[XY] = M_n[X]M_n[Y]$, $M_a[X]M_b[X] = M_{a+b}[X]$, and

$$M_n[X_1 + \cdots + X_m] = \sum_{\beta_1 + \cdots + \beta_m = n} \binom{n}{\beta_1, \dots, \beta_m} M_{\beta_1}[X_1] \cdots M_{\beta_m}[X_m].$$

Furthermore, for any constant b , we have $M_n[b] = b^n$.

We start with the unitary group $U(1)$ and consider the trace map on $U \in U(1)$ defined by $z := \text{tr}(U) = u + \bar{u} = 2 \cos(\theta)$, where $u = e^{i\theta}$. This trace map takes values in $[-2, 2]$. From here we see that $dz = 2 \sin(\theta) d\theta$ and

$$\mu_{U(1)} = \frac{1}{\pi} \frac{dz}{\sqrt{4 - z^2}} = \frac{1}{\pi} d\theta$$

gives a uniform measure of $U(1)$ on $\theta \in [-\pi, \pi]$ (see [31, Section 2]). We can deduce the following pushforward measure

$$\mu_{U(1)^n} = \prod_{i=1}^n \frac{1}{\pi} \frac{dz_i}{\sqrt{4 - z_i^2}} = \prod_{i=1}^n \frac{1}{\pi} d\theta_i.$$

We can now define the moment sequence $M[\mu]$, where μ is a positive measure on some interval $I = [-d, d]$. The n^{th} moment $M_n[\mu]$ is, by definition, the expected value of $\mu(\phi_n)$, where ϕ_n is the function $z \mapsto z^n$. It is therefore given by

$$M_n[\mu] = \int_I z^n \mu(z).$$

For $U(1)$ we have $M_n[\mu_{U(1)}] = \binom{n}{n/2}$, where $\binom{n}{n/2} = 0$ if n is odd. Hence, $M[\mu_{U(1)}] = (1, 0, 2, 0, 6, 0, 20, 0, \dots)$. From here, we take binomial convolutions to obtain

$$M_n[\mu_{U(1)^g}] = \sum_{\beta_1 + \dots + \beta_g = n} \binom{n}{\beta_1, \dots, \beta_g} M_{\beta_1}[\mu_{U(1)}] \cdots M_{\beta_g}[\mu_{U(1)}]. \quad (12)$$

In what follows, for each $i \in \{1, 2, \dots, g\}$, denote by μ_i the projection of the Haar measure onto the interval $[-\binom{2g}{i}, \binom{2g}{i}]$. We can compute $M_n[\mu_i]$ by averaging over the components of the Sato-Tate group.

5.2. Characteristic polynomials. In this subsection, we give results for the characteristic polynomials in each component of the Sato-Tate group. Let U be a random matrix in the identity component of $\text{ST}(\text{Jac}(C_{p,q}))$ and let $A_{m,n} = \gamma_p^m \gamma_q^n$ be a matrix in the component group $\text{ST}(\text{Jac}(C_{p,q}))/\text{ST}^0(\text{Jac}(C_{p,q}))$. The characteristic polynomial of $U \cdot A_{m,n}$ is a degree $2g$ palindromic polynomial of the form

$$P_{m,n}(T) = T^{2g} + b_1 T^{2g-1} + b_2 T^{2g-2} + \dots + b_2 T^2 + b_1 T + 1.$$

We now present two propositions regarding the coefficients of the characteristic polynomials.

Proposition 5.1. *Only the identity component $U(1)^g$ contributes to the b_1 -coefficient of the characteristic polynomial of any matrix in $\text{ST}(\text{Jac}(C_{p,q}))$.*

Proof. The b_1 -coefficient of the characteristic polynomial of $P_{m,n}$ is the trace of the matrix product $U \cdot A_{m,n}$. We will show that this trace is nonzero if and only if $m \equiv 0 \pmod{p-1}$ and $n \equiv 0 \pmod{q-1}$.

To do this it is sufficient to show that $A_{m,n}$ has no nonzero diagonal entries. If $A_{m,n}$ has a nonzero diagonal entry then, for some j , the j^{th} diagonal block entry of $A_{m,n}$ is I and α satisfies

$$(\sigma_p)^m \circ (\sigma_q)^n \alpha[j] = \alpha[j]. \quad (13)$$

To determine if and when this is possible, we consider the action of the corresponding Galois element $(\sigma_p)^m \circ (\sigma_q)^n$. Let $\alpha[j] = Z^{q(a+1)-pb}$. Equation (13) holds if and only if the following is true:

$$\zeta_{pq}^{q(a+1)-pb} = (\sigma_p)^m \circ (\sigma_q)^n \left(\zeta_{pq}^{q(a+1)-pb} \right) = \zeta_{pq}^{qc^m(a+1)-pd^nb}.$$

This equality implies that

$$\zeta_p^{(a+1)} \zeta_q^{-b} = \zeta_p^{c^m(a+1)} \zeta_q^{-d^nb}.$$

Since p and q are relatively prime, we can conclude from here that

$$\zeta_p^{(a+1)} = \zeta_p^{c^m(a+1)} = (\sigma_p)^m \zeta_p^{(a+1)} \quad \text{and} \quad \zeta_q^{-b} = \zeta_q^{-d^nb} = (\sigma_q)^n \zeta_q^{-b}.$$

Since σ_p and σ_q are generators of $\text{Gal}(K/\mathbb{Q})$ and have orders $p-1$ and $q-1$, respectively, the above equalities hold if and only if $m \equiv 0 \pmod{p-1}$ and $n \equiv 0 \pmod{q-1}$. \square

One benefit of this result is that, in order to compute the b_1 -moment statistics, we need only compute moments for a random matrix U in the identity component of $\text{ST}(\text{Jac}(C_{p,q}))$. To average over the components we simply divide this value by the size of the Galois group.

Proposition 5.2. *Let $m = (p-1)/2$ and $n = (q-1)/2$. Then*

$$P_{m,n}(T) = (T^2 + 1)^g.$$

Proof. We will show that the matrix $A_{m,n}$ for these values of m and n has J or $-J$ as each of its g diagonal blocks. Since the characteristic polynomial of $\pm J$ is $T^2 + 1$, this will yield the desired result.

The actions and orders of the Galois elements σ_p and σ_q tell us that $\sigma_p^m(\zeta_p) = \bar{\zeta}_p$ and $\sigma_q^n(\zeta_q) = \bar{\zeta}_q$. Thus,

$$\begin{aligned} \sigma_p^m \circ \sigma_q^n(\zeta_{pq}^{q(a+1)-pb}) &= \sigma_p^m \circ \sigma_q^n(\zeta_p^{(a+1)} \bar{\zeta}_q^b) \\ &= \bar{\zeta}_p^{(a+1)} \zeta_q^b \\ &= \overline{\zeta_{pq}^{q(a+1)-pb}}. \end{aligned}$$

Thus, every diagonal entry of α satisfies

$$\sigma_p^m \circ \sigma_q^n \alpha[j] = \overline{\alpha[j]},$$

which means that the block diagonal entries of the corresponding matrix are all either J or $-J$. \square

Example 5.3. We determine moment statistics for the genus 4 curve $C_{5,3} : y^3 = x^5 - 1$ over each subfield of the CM field $K = \mathbb{Q}(\zeta_{15})$. Using characteristic polynomials, we can compute the n th moments for each μ_i , $1 \leq i \leq 4$. There is a table corresponding to each of the four subfields: \mathbb{Q} (Table 1), $\mathbb{Q}(\zeta_3)$ (Table 2), $\mathbb{Q}(\zeta_5)$ (Table 3), and $\mathbb{Q}(\zeta_{15})$ (Table 4). These moments were computed using Sage [27].

$M[\mu_1]$	(1, 0, 1, 0, 21, 0, 640, 0, 23765, ...)
$M[\mu_2]$	(1, 1, 8, 76, 1168, 20956, 414284, 8643328, 187416464, ...)
$M[\mu_3]$	(1, 0, 13, 0, 11745, 0, 17177080, 0, 31036079585, ...)
$M[\mu_4]$	(1, 2, 27, 476, 18391, 689812, 34599990, 1677458008, 91894386279, ...)

TABLE 1. Moment Statistics for $y^3 = x^5 - 1$ over \mathbb{Q} .

$M[\mu_1]$	(1, 0, 2, 0, 42, 0, 1280, 0, 47530, ...)
$M[\mu_2]$	(1, 1, 11, 136, 2263, 41656, 827444, 17282560, 374815319, ...)
$M[\mu_3]$	(1, 0, 26, 0, 23490, 0, 34354160, 0, 62072159170, ...)
$M[\mu_4]$	(1, 2, 42, 890, 36418, 1377502, 69187410, 3354841408, 183788328258, ...)

TABLE 2. Moment Statistics for $y^3 = x^5 - 1$ over $\mathbb{Q}(\zeta_3)$.

$M[\mu_1]$	(1, 0, 4, 0, 84, 0, 2560, 0, 95060, ...)
$M[\mu_2]$	(1, 2, 22, 272, 4526, 83312, 1654888, 34565120, 749630638, ...)
$M[\mu_3]$	(1, 0, 52, 0, 46980, 0, 68708320, 0, 124144318340, ...)
$M[\mu_4]$	(1, 4, 82, 1780, 72830, 2755004, 138374800, 6709682816, 367576656446, ...)

TABLE 3. Moment Statistics for $y^3 = x^5 - 1$ over $\mathbb{Q}(\zeta_5)$.

$M[\mu_1]$	(1, 0, 8, 0, 168, 0, 5120, 0, 190120, ...)
$M[\mu_2]$	(1, 4, 40, 544, 9016, 166624, 3309376, 69130240, 1499256376, ...)
$M[\mu_3]$	(1, 0, 104, 0, 93960, 0, 137416640, 0, 248288636680, ...)
$M[\mu_4]$	(1, 6, 156, 3528, 145512, 5509296, 276746016, 13419347136, 735153215448, ...)

TABLE 4. Moment Statistics for $y^3 = x^5 - 1$ over $\mathbb{Q}(\zeta_{15})$.

5.3. Tables of μ_1 - and a_1 -moment statistics. Table 5 gives both $M[\mu_1]$ and the numerical moments of the normalized L -polynomial of some Catalan Jacobians (over \mathbb{Q}). The numerical moments were computed for primes up to 2^N , where N is given in the last column of the table, using an algorithm described in [18] and [19]. As the genus grows, the computations require more processing power, and so a smaller bound N was used. This leads to less accurate estimates for the numerical moments in higher genus, which make it more difficult to compare them to the theoretical moments obtained from the Sato-Tate groups.

In Proposition 5.1 we proved that only the identity component contributes to the b_1 -coefficient of the characteristic polynomial. Let $M_n[\mu_1]$ denote the n^{th} moment of the identity component. As noted in the proof of Proposition 5.1, the b_1 -coefficient of the characteristic polynomial is the trace of a random matrix $U \in \text{U}(1)^g$. Thus, $M_n[\mu_1] = M_n[\mu_{\text{U}(1)^g}]$ and we can use the formula in Equation (12). To compute $M_n[\mu_1]$, we average over the components by dividing $M_n[\mu_{\text{U}(1)^g}]$ by the size of the Galois group.

(p, q)	g		M_2	M_4	M_6	M_8	N
(5,3)	4	μ_1	1	21	640	23765	23
		a_1	0.994	20.726	625.606	22961.668	
(7,3)	6	μ_1	1	33	1660	106785	23
		a_1	0.998	32.718	1628.656	103534.931	
(11,3)	10	μ_1	1	57	5140	615545	19
		a_1	0.972	55.306	4836.457	532489.325	
(13,3)	12	μ_1	1	69	7600	1121925	19
		a_1	0.998	68.164	7570.201	1178539.197	
(7,5)	12	μ_1	1	69	7600	1121925	19
		a_1	0.998	68.164	7570.201	1178539.197	
(17,3)	16	μ_1	1	93	13960	2840285	18
		a_1	0.954	91.335	15348.411	3864179.550	

TABLE 5. Table of some μ_1 - and a_1 -moments $\text{Jac}(C_{p,q})$ over \mathbb{Q} .

Note that $M_2[\mu_1]$ equals 1 for each example in Table 5. We prove that this is true for every Catalan Jacobian in Proposition 5.4.

Remark 4. In many cases, the μ_1 -moments of Catalan Jacobians match those of the Fermat quotients considered in [10]. This occurs when the genera are the same and the identity components are the same (some of the identity components in [10] are of the form $(\text{U}(1)_{g/3})^3$). These Jacobian varieties have a similar property to ours: only the identity components of the Sato-Tate groups contribute to the b_1 -coefficients of the characteristic polynomials. However we expect to see a difference in the moments of the higher traces.

For example, consider the two genus 6 curves $y^3 = x^7 - 1$ (Catalan curve) and $v^{13} = u(u+1)^{10}$ (a curve from [10]) over \mathbb{Q} . Both of their associated Sato-Tate groups have identity component $U(1)^6$, but their component groups are different. We can see this difference in the μ_2 -moments:

$$\begin{aligned} y^3 = x^7 - 1: \quad M[\mu_2] &= (1, 1, 12, 206, 5796, \dots), \\ v^{13} = u(u+1)^{10}: \quad M[\mu_2] &= (1, 1, 11, 206, 5781, \dots). \end{aligned}$$

5.4. General results for moment statistics.

Proposition 5.4. *Over \mathbb{Q} , the second moment $M_2[\mu_1]$ equals 1 for every Catalan Jacobian. Over $\mathbb{Q}(\zeta_p)$, $\mathbb{Q}(\zeta_q)$, and $\mathbb{Q}(\zeta_{pq})$, the moment $M_2[\mu_1]$ is $p-1$, $q-1$, and $(p-1)(q-1)$, respectively.*

Proof. In Proposition 5.1 we proved that only the identity component contributes to the b_1 -coefficient of the characteristic polynomial. We will demonstrate that the value obtained for the second moment $M_2[{}^0\mu_1]$ is $(p-1)(q-1)$, so that when averaging over all components we obtain the desired results.

This amounts to computing the sum in Equation (12) when $n = 2$. In this situation, we are summing over β_i values that add to 2. There are two types of summands to consider: ones where there are only two nonzero values $\beta_i = \beta_j = 1$ for some $i \neq j$ and ones where there is only one nonzero value $\beta_i = 2$ for some i .

The first type does not contribute to the sum because $M_1[\mu_{U(1)}] = 0$ and, hence the entire term equals 0. On the other hand, each term of the second type will simplify to

$$\binom{2}{2, 0, \dots, 0} M_2[\mu_{U(1)}] M_0[\mu_{U(1)}] \cdots M_0[\mu_{U(1)}] = 2$$

since $M_0[\mu_{U(1)}] = 1$ and $M_2[\mu_{U(1)}] = 2$. There are exactly g terms of this form since this is essentially counting the number of g -tuples with a single nonzero entry. Hence, $M_n[{}^0\mu_1] = 2g = (p-1)(q-1)$.

We now consider the size of the component group for each of the subfields of $\mathbb{Q}(\zeta_{pq})$. Over \mathbb{Q} , the component group has size $(p-1)(q-1)$. Over $\mathbb{Q}(\zeta_p)$, $\mathbb{Q}(\zeta_q)$, $\mathbb{Q}(\zeta_{pq})$, the sizes are $q-1$, $p-1$, and 1, respectively. Averaging $M_n[{}^0\mu_1]$ over these values yields the desired results. \square

6. GALOIS ENDOMORPHISM TYPES AND RELATED RESULTS

We begin this section with some notation and terminology. Let A/\mathbb{Q} be an abelian variety and K be the endomorphism field of A . We will denote the real endomorphism algebra of A/\mathbb{Q} by $\text{End}(A_K)_{\mathbb{R}} := \text{End}(A_K) \otimes_{\mathbb{Z}} \mathbb{R}$. Let \mathcal{C} be the category of pairs (G, E) , where G is a finite group and E is an \mathbb{R} -algebra equipped with an \mathbb{R} -linear action of G . The **Galois endomorphism type** of A/\mathbb{Q} is the isomorphism class in \mathcal{C} of the pair $[\text{Gal}(K/\mathbb{Q}), \text{End}(A_{\mathbb{Q}})_{\mathbb{R}}]$ [11, Definition 1.3]. If L/\mathbb{Q} is a subextension of K/k corresponding to a subgroup $N \subseteq \text{Gal}(K/\mathbb{Q})$, then $\text{End}(A_L)_{\mathbb{R}} \simeq (\text{End}(A_K)_{\mathbb{R}})^N$ (see, for example, [15, Section 6]).

Let \mathbb{H} denote the quaternions and let $M_n(R)$ denote the $n \times n$ matrix ring over a ring R . Wedderburn's structure theorem tells us that the \mathbb{R} -algebra $\text{End}(A_{\mathbb{Q}})_{\mathbb{R}}$ of an abelian variety A/k satisfies

$$\text{End}(A_K)_{\mathbb{R}} \simeq \prod_i M_{t_i}(\mathbb{R}) \times \prod_i M_{n_i}(\mathbb{H}) \times \prod_i M_{p_i}(\mathbb{C}), \quad (14)$$

for some nonnegative integers t_i, n_i, p_i .

In the subsections that follow, we will prove the following result for the Galois endomorphism types of Catalan Jacobians.

Theorem 6.1. *Let $K = \mathbb{Q}(\zeta_{pq})$. Then the Galois endomorphism types associated to A for each subextension L/\mathbb{Q} of K/\mathbb{Q} are as given in Table 6.*

Field L	$N \subseteq \text{Gal}(K/\mathbb{Q})$	$\text{End}(\text{Jac}(C_{p,q})_L)_{\mathbb{R}}$
\mathbb{Q}	1	\mathbb{R}
$\mathbb{Q}(\zeta_p)$	C_{q-1}	$\mathbb{C}^{(p-1)/2}$
$\mathbb{Q}(\zeta_q)$	C_{p-1}	$\mathbb{C}^{(q-1)/2}$
$\mathbb{Q}(\zeta_{pq})$	$C_{p-1} \times C_{q-1}$	\mathbb{C}^g

TABLE 6. Galois endomorphism types of $\text{Jac}(C_{p,q})_L$.

We will prove this result using two methods: using data obtained from moment statistics and working with Rosati forms. The latter is the more traditional method of determining real endomorphism algebras, whereas the former is a new technique using recent results of Costa, Fité, and Sutherland [8].

The results of Proposition 6.3, Proposition 6.11, and Corollary 6.12 together prove Theorem 6.1.

6.1. Data obtained from moment statistics. In this section we use the results of [8] to obtain information about the real endomorphism algebras of Catalan Jacobians. There are four main results in that paper, and we apply them all. We start by using the following proposition.

Proposition 6.2. [8, Proposition 1] *Let A be an abelian variety defined over k .*

$$M_2[a_1] = \text{rk}_{\mathbb{Z}}(\text{End}(A_k)).$$

Combining this with the results of Proposition 5.4, yields the values in Table 7. From here we can immediately conclude the result in Proposition 6.3.

Field L	$\text{rk}_{\mathbb{Z}}(\text{End}(\text{Jac}(C_{p,q})_L))$
\mathbb{Q}	1
$\mathbb{Q}(\zeta_p)$	$p - 1$
$\mathbb{Q}(\zeta_q)$	$q - 1$
$\mathbb{Q}(\zeta_{pq})$	$(p - 1)(q - 1)$

TABLE 7. Rank of the endomorphism ring of $\text{Jac}(C_{p,q})_L$.

Proposition 6.3. *The real endomorphism algebra $\text{End}(\text{Jac}(C_{p,q})_{\mathbb{Q}})_{\mathbb{R}}$ for any Catalan Jacobian is \mathbb{R} .*

Proof. Since $\text{rk}_{\mathbb{Z}}(\text{End}(\text{Jac}(C_{p,q})_{\mathbb{Q}})) = 1$, the only possibility for $\text{End}(\text{Jac}(C_{p,q})_{\mathbb{Q}})_{\mathbb{R}}$ is \mathbb{R} . \square

Let $\text{NS}(A_F)$ denote the Néron-Severi group of a variety A defined over F .

Corollary 6.4. *The rank of the Néron-Severi group of $\text{Jac}(C_{p,q})_{\mathbb{Q}}$ is 1.*

Proof. This follows from Proposition 6.3 and the fact that the Néron-Severi group of a principally polarized abelian variety A embeds into $\text{End}(A) \otimes_{\mathbb{Z}} \mathbb{Q}$.

Alternatively, we can use Lemma 3 of [8, Proposition] to prove the result. Applying this lemma to our situation yields the following equation:

$$1 - 2 \cdot \text{rk}_{\mathbb{Z}}(\text{NS}(\text{Jac}(C_{p,q})_{\mathbb{Q}})) = -1.$$

Hence, $\text{rk}_{\mathbb{Z}}(\text{NS}(\text{Jac}(C_{p,q})_{\mathbb{Q}})) = 1$. \square

Proposition 2 of [8] gives an additional identity for the the rank of $\text{NS}(A_F)$:

$$M_1[a_2] = \text{rk}_{\mathbb{Z}}(\text{NS}(A_F)). \quad (15)$$

Thus, we obtain the following corollary.

Corollary 6.5. *For the Catalan Jacobian $\text{Jac}(C_{p,q})_{\mathbb{Q}}$, we have*

$$M_1[a_2] = 1.$$

We now consider the moment $M_1[s_2] = M_2[a_1] - 2M_1[a_2]$. This moment can be interpreted as a Frobenius-Schur indicator of the standard representation of $\text{ST}(A)$ (see the exposition in Remark 5 of [8] for more details). The value of $M_1[s_2]$, therefore, can only be $-1, 0$, or 1 (see Proposition 39 of [28]). We now prove that, for Catalan Jacobians, the moment $M_1[s_2]$ is always either -1 or 0 .

Proposition 6.6. *Over \mathbb{Q} , the moment $M_1[s_2]$ equals -1 for every Catalan Jacobian. Over $\mathbb{Q}(\zeta_p), \mathbb{Q}(\zeta_q)$, and $\mathbb{Q}(\zeta_{pq})$, the moment is $M_1[s_2] = 0$.*

Proof. To prove this result, we use the results of Proposition 5.4, the identity $M_1[s_2] = M_2[a_1] - 2M_1[a_2]$, and the fact that $M_1[s_2]$ can only equal $-1, 0$, or 1 by Proposition 39 of [28].

Over \mathbb{Q} , we have already proved that $M_2[a_1] = 1$ and $M_1[a_2] = 1$ in Proposition 5.4 and Corollary 6.5. Thus, $M_1[s_2] = 1 - 2 = -1$ as desired.

Now consider the Jacobian over the field $\mathbb{Q}(\zeta_p)$. We have already shown in Proposition 5.4 that $M_2[a_1] = p - 1$. Thus,

$$M_1[s_2] = p - 1 - 2M_1[a_2].$$

If $M_1[s_2]$ equals 1 or -1 , then $M_1[a_2] = \frac{p-1 \pm 1}{2}$. However neither of these values are integers, and the moment $M_1[a_2]$ must be an integer. Thus, we must have $M_1[s_2] = 0$.

A similar argument proves that $M_1[s_2] = 0$ when working over the fields $\mathbb{Q}(\zeta_q)$ and $\mathbb{Q}(\zeta_{pq})$. \square

Corollary 6.7. *Over $\mathbb{Q}(\zeta_p), \mathbb{Q}(\zeta_q)$, and $\mathbb{Q}(\zeta_{pq})$, the moment $M_1[a_2]$ equals $\frac{p-1}{2}, \frac{q-1}{2}$, and $\frac{(p-1)(q-1)}{2}$, respectively.*

Proof. This follows from work in the proof of Proposition 6.6. \square

Corollary 6.8. *Over $\mathbb{Q}(\zeta_p), \mathbb{Q}(\zeta_q)$, and $\mathbb{Q}(\zeta_{pq})$, the the ranks of the Néron-Severi groups are $\frac{p-1}{2}, \frac{q-1}{2}$, and $\frac{(p-1)(q-1)}{2}$, respectively.*

Proof. This follows from Corollary 6.7 and Equation (15). \square

The value that we obtain for the moment $M_1[s_2]$ can give us some information about the real endomorphism algebra of $\text{Jac}(C_{p,q})_L$ for each subextension L/\mathbb{Q} of K/\mathbb{Q} . Corollary 4 of [8] shows that

$$M_1[s_2] = 2 \sum_i n_i - \sum_i t_i. \quad (16)$$

In some cases, this identity combined with other moment results is enough to determine the real endomorphism algebra.

Proposition 6.9. *Let one of p or q be equal to 3. Then the real endomorphism algebra $\text{End}(\text{Jac}(C_{p,q})_{\mathbb{Q}(\zeta_3)})_{\mathbb{R}}$ is \mathbb{C} .*

Proof. For ease of notation, let $L = \mathbb{Q}(\zeta_3)$. First note that, by Propositions 5.4 and 6.2, we have $\text{rk}_{\mathbb{Z}}(\text{End}(\text{Jac}(C_{p,q})_L)) = 2$. This immediately tells us that the real endomorphism algebra is either $\mathbb{R} \times \mathbb{R}$ or \mathbb{C} . We now work to show that the algebra must, in fact, be \mathbb{C} .

We proved in Proposition 6.6 that, over the field L , $M_1[s_2] = 0$. Combining this with the result in Corollary 4 of [8] yields

$$2 \sum_i n_i = \sum_i t_i.$$

This implies that if the real endomorphism algebra has a factor of $M_{t_i}(\mathbb{R})$, for some nonzero t_i , then it must also have a factor of $M_{n_i}(\mathbb{H})$ for a nonzero n_i . Thus, the real endomorphism algebra cannot be $\mathbb{R} \times \mathbb{R}$, and so it must be \mathbb{C} . \square

Proposition 6.10. *Let one of p or q be equal to 5. Then the real endomorphism algebra $\text{End}(\text{Jac}(C_{p,q})_{\mathbb{Q}(\zeta_5)})_{\mathbb{R}}$ is $\mathbb{C} \times \mathbb{C}$.*

Proof. For ease of notation, let $L = \mathbb{Q}(\zeta_5)$. First note that, by Propositions 5.4 and 6.2, we have $\text{rk}_{\mathbb{Z}}(\text{End}(\text{Jac}(C_{p,q})_L)) = 4$. This immediately tells us that the real endomorphism algebra is one of the following:

$$\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}, M_2(\mathbb{R}), \mathbb{H}, \mathbb{C} \times \mathbb{R} \times \mathbb{R}, \text{ or } \mathbb{C} \times \mathbb{C}.$$

By a similar argument to that in the proof of Proposition 6.9, we determine that there are no $M_{t_i}(\mathbb{R})$ or $M_{n_i}(\mathbb{H})$ factors. The only remaining possibility is $\mathbb{C} \times \mathbb{C}$. \square

It does not appear to be possible to use these techniques to uniquely determine the real endomorphism algebra for larger values of p and q or for pq with the primes 3 and 5. For example, suppose one of the primes equals 7. Then the rank of the real endomorphism algebra is 6, and $\mathbb{R} \times \mathbb{R} \times \mathbb{H}$ and $\mathbb{C} \times \mathbb{C} \times \mathbb{C}$ both satisfy all of the results we have used thus far in our discussion. In order to further classify real endomorphism algebras associated to Catalan Jacobians, we will use Rosati forms.

6.2. Further data from Rosati forms. In this section we use the fact that, for any nondegenerate abelian variety A/F , the Galois endomorphism type is uniquely determined by its Sato-Tate group (see Propositions 2.16 and 2.19 of [11]).

Let $K = \mathbb{Q}(\zeta_{pq})$ and $G = \text{ST}(\text{Jac}(C_{p,q})_K)$. Then $\text{End}(\text{Jac}(C_{p,q})_K) \otimes_{\mathbb{Z}} \mathbb{C}$ is the subspace of $M_{2g}(\mathbb{C})$ fixed by the action of G^0 , and the real endomorphism algebra $\text{End}(\text{Jac}(C_{p,q})_K)_{\mathbb{R}}$ is the subspace of half the dimension for which the Rosati form is positive definite (see Definition 2.18 in [11]).

The Sato-Tate group $\text{ST}(\text{Jac}(C_{p,q})_K)$ is connected: it is equal to the identity component of $\text{ST}(\text{Jac}(C_{p,q})_{\mathbb{Q}})$, which we proved in Proposition 4.3 is $\text{U}(1)^g$. This allows us to prove the following statement.

Proposition 6.11. *The real endomorphism algebra $\text{End}(\text{Jac}(C_{p,q})_K)_\mathbb{R}$ is \mathbb{C}^g .*

Proof. We will prove the result by determining the subspace of $M_{2g}(\mathbb{C})$ that is positive definite for the real part of the Rosati form. This amounts to determining the matrices Φ for which $\text{Trace}(\Phi H^t \Phi^t H) \geq 0$, where H is the symplectic matrix in Equation (4).

The matrices in $M_{2g}(\mathbb{C})$ that commute with the identity component $G^0 = \text{U}(1)^g$ are

$$\{\Phi = \text{diag}(z_1, z_2, \dots, z_{2g}) \mid z_j \in \mathbb{C}\}.$$

We compute the matrix $\Phi H^t \Phi^t H$ to be

$$\text{diag}(z_1 z_2, z_1 z_2, \dots, z_{2g-1} z_{2g}, z_{2g-1} z_{2g}).$$

The trace of this matrix is $2z_1 z_2 + 2z_3 z_4 + \dots + 2z_{2g-1} z_{2g}$, and each term in this sum can be expressed as

$$\frac{(z_j + z_{j+1})^2 - (z_j - z_{j+1})^2}{2}.$$

The positivity of the Rosati form requires that $z_j + z_{j+1} \in \mathbb{R}$ and $z_j - z_{j+1} \in i\mathbb{R}$. Thus, $z_j = x_k + iy_k$ and $z_{j+1} = x_k - iy_k$ for some real numbers x_k, y_k . From this we can conclude that each 2×2 block of Φ is of the form $\text{diag}(x_k + iy_k, x_k - iy_k)$. There are g such blocks, so $\text{End}(\text{Jac}(C_{p,q})_K)_\mathbb{R} \simeq \mathbb{C}^g$. \square

Having established the real endomorphism algebra corresponding to the endomorphism field of $\text{Jac}(C_{p,q})$, we can now determine the real endomorphism algebra $\text{End}(\text{Jac}(C_{p,q})_L)$ for each subextension L/\mathbb{Q} of K/\mathbb{Q} .

Corollary 6.12. *The real endomorphism algebras over the fields $\mathbb{Q}(\zeta_p)$ and $\mathbb{Q}(\zeta_q)$ are $\mathbb{C}^{(p-1)/2}$ and $\mathbb{C}^{(q-1)/2}$, respectively.*

Proof. We proved in Proposition 6.11 that $\text{End}(\text{Jac}(C_{p,q})_K)_\mathbb{R} \simeq \mathbb{C}^g$. Thus, for any subextension L/\mathbb{Q} , $\text{End}(\text{Jac}(C_{p,q})_L)_\mathbb{R}$ does not contain any $M_{n_i}(\mathbb{H})$ factors. We proved in Proposition 6.6 that $M_1[s_2] = 0$ for both $\mathbb{Q}(\zeta_p)$ and $\mathbb{Q}(\zeta_q)$. Using Equation (16) and the notation in Equation 14, we can conclude that there are no factors of the form $M_{t_i}(\mathbb{R})$ in the real endomorphism algebra.

Thus, the real endomorphism algebras over both fields are powers of \mathbb{C} . The ranks of the real endomorphism algebras in Table 7 confirm that the powers are as given in the statement of the corollary. \square

ACKNOWLEDGEMENTS

Many thanks to Francesc Fité, Edgar Costa, and Drew Sutherland while working on Section 6.1. Thanks also to McKenzie West for patiently answering my Sage and GitHub questions while I was working on Table 5. I would like to thank Drew Sutherland for sharing access to a server at MIT while I was computing the numerical moments given in Table 5.

I am grateful for the support and encouragement of Matt Montesano, Alanna Hoyer-Letizel, and the Rethinking Number Theory Workshop co-organizers, project leaders, and participants. They provided a sense of community during a difficult time, and it would not have been possible to write this article during a pandemic without them.

My research is supported by a PSC-CUNY Award, jointly funded by The Professional Staff Congress and The City University of New York.

A. EXAMPLES OF THE COMPONENT GROUP GENERATORS

In Table 8 we give examples of the matrices γ_q and γ_p from Definitions 4.4 and 4.5. These were computed in Sage [27] using Sage's chosen generators for $(\mathbb{Z}/q\mathbb{Z})^\times$ and $(\mathbb{Z}/p\mathbb{Z})^\times$.

(p, q)	γ_p	γ_q
$(5, 3)$	$\begin{pmatrix} 0 & 0 & 0 & J \\ 0 & 0 & I & 0 \\ J & 0 & 0 & 0 \\ 0 & I & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & I & 0 & 0 \\ I & 0 & 0 & 0 \\ 0 & 0 & 0 & J \\ 0 & 0 & J & 0 \end{pmatrix}$
$(7, 3)$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & J \\ 0 & 0 & J & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & I & 0 \\ J & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I & 0 & 0 \\ 0 & J & 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & I & 0 & 0 & 0 \\ 0 & 0 & 0 & I & 0 & 0 \\ I & 0 & 0 & 0 & 0 & 0 \\ 0 & I & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & J \\ 0 & 0 & 0 & 0 & J & 0 \end{pmatrix}$

TABLE 8. Examples of γ_p and γ_q matrices for $y^q = x^p - 1$.

REFERENCES

- [1] Noboru Aoki. Hodge cycles on CM abelian varieties of Fermat type. *Comment. Math. Univ. St. Paul.*, 51(1):99–130, 2002.
- [2] Noboru Aoki. The Hodge conjecture for the Jacobian varieties of generalized Catalan curves. *Tokyo J. Math.*, 27(2):313–335, 2004.
- [3] Sonny Arora, Victoria Cantoral-Farfán, Aaron Landesman, Davide Lombardo, and Jackson S. Morrow. The twisting Sato-Tate group of the curve $y^2 = x^8 - 14x^4 + 1$. *Math. Z.*, 290(3-4):991–1022, 2018.
- [4] G. Banaszak, W. Gajda, and P. Krasoń. On Galois representations for abelian varieties with complex and real multiplications. *J. Number Theory*, 100(1):117–132, 2003.
- [5] Grzegorz Banaszak and Kiran S. Kedlaya. An algebraic Sato-Tate group and Sato-Tate conjecture. *Indiana Univ. Math. J.*, 64(1):245–274, 2015.
- [6] Tom Barnet-Lamb, David Geraghty, Michael Harris, and Richard Taylor. A family of Calabi-Yau varieties and potential automorphy II. *Publ. Res. Inst. Math. Sci.*, 47(1):29–98, 2011.
- [7] Laurent Clozel, Michael Harris, and Richard Taylor. Automorphy for some l -adic lifts of automorphic mod l Galois representations. *Publ. Math. Inst. Hautes Études Sci.*, (108):1–181, 2008. With Appendix A, summarizing unpublished work of Russ Mann, and Appendix B by Marie-France Vignéras.
- [8] Edgar Costa, Francesc Fité, and Andrew V. Sutherland. Arithmetic invariants from Sato-Tate moments. *C. R. Math. Acad. Sci. Paris*, 357(11-12):823–826, 2019.
- [9] Melissa Emory and Heidi Goodson. Sato-Tate Distributions of $y^2 = x^p - 1$ and $y^2 = x^{2p} - 1$. *arXiv e-prints*, page arXiv:2004.10583, April 2020.
- [10] Francesc Fité, Josep González, and Joan-Carles Lario. Frobenius distribution for quotients of Fermat curves of prime exponent. *Canad. J. Math.*, 68(2):361–394, 2016.
- [11] Francesc Fité, Kiran S. Kedlaya, Víctor Rotger, and Andrew V. Sutherland. Sato-Tate distributions and Galois endomorphism modules in genus 2. *Compos. Math.*, 148(5):1390–1442, 2012.
- [12] Francesc Fité, Kiran S. Kedlaya, and Andrew V. Sutherland. Sato-Tate groups of abelian threefolds: a preview of the classification. In *Arithmetic, geometry, cryptography and coding theory*, volume 770 of *Contemp. Math.*, pages 103–129. Amer. Math. Soc., [Providence], RI, [2021].
- [13] Francesc Fité, Elisa Lorenzo García, and Andrew V. Sutherland. Sato-Tate distributions of twists of the Fermat and the Klein quartics. *Res. Math. Sci.*, 5(4):Paper No. 41, 40, 2018.
- [14] Francesc Fité and Andrew V. Sutherland. Sato-Tate distributions of twists of $y^2 = x^5 - x$ and $y^2 = x^6 + 1$. *Algebra Number Theory*, 8(3):543–585, 2014.

- [15] Francesc Fité and Andrew V. Sutherland. Sato-Tate groups of $y^2 = x^8 + c$ and $y^2 = x^7 - cx$. In *Frobenius distributions: Lang-Trotter and Sato-Tate conjectures*, volume 663 of *Contemp. Math.*, pages 103–126. Amer. Math. Soc., Providence, RI, 2016.
- [16] Francesc Fité, Kiran S. Kedlaya, and Andrew V. Sutherland. Sato-tate groups of abelian threefolds. *arXiv e-prints*, page arxiv:2106.13759, 2021.
- [17] Michael Harris, Nick Shepherd-Barron, and Richard Taylor. A family of Calabi-Yau varieties and potential automorphy. *Ann. of Math. (2)*, 171(2):779–813, 2010.
- [18] David Harvey and Andrew V. Sutherland. Computing Hasse-Witt matrices of hyperelliptic curves in average polynomial time. *LMS J. Comput. Math.*, 17(suppl. A):257–273, 2014.
- [19] David Harvey and Andrew V. Sutherland. Computing Hasse-Witt matrices of hyperelliptic curves in average polynomial time, II. In *Frobenius distributions: Lang-Trotter and Sato-Tate conjectures*, volume 663 of *Contemp. Math.*, pages 127–147. Amer. Math. Soc., Providence, RI, 2016.
- [20] Fumio Hazama. Algebraic cycles on nonsimple abelian varieties. *Duke Math. J.*, 58(1):31–37, 1989.
- [21] Fumio Hazama. Hodge cycles on the Jacobian variety of the Catalan curve. *Compositio Math.*, 107(3):339–353, 1997.
- [22] Christian Johansson. On the Sato-Tate conjecture for non-generic abelian surfaces. *Trans. Amer. Math. Soc.*, 369(9):6303–6325, 2017. With an appendix by Francesc Fité.
- [23] Joan-Carles Lario and Anna Somoza. The Sato-Tate conjecture for a Picard curve with complex multiplication (with an appendix by Francesc Fité). In *Number theory related to modular curves—Momose memorial volume*, volume 701 of *Contemp. Math.*, pages 151–165. Amer. Math. Soc., Providence, RI, 2018.
- [24] Preda Mihailescu. Primary cyclotomic units and a proof of Catalan’s conjecture. *J. Reine Angew. Math.*, 572:167–195, 2004.
- [25] V. Kumar Murty. Exceptional Hodge classes on certain abelian varieties. *Math. Ann.*, 268(2):197–206, 1984.
- [26] Henry Pohlmann. Algebraic cycles on abelian varieties of complex multiplication type. *Ann. of Math. (2)*, 88:161–180, 1968.
- [27] Inc. SageMath. *CoCalc Collaborative Computation Online*, 2020. <https://cocalc.com/>.
- [28] Jean-Pierre Serre. *Linear representations of finite groups*. Graduate Texts in Mathematics, Vol. 42. Springer-Verlag, New York-Heidelberg, 1977. Translated from the second French edition by Leonard L. Scott.
- [29] Jean-Pierre Serre. *Lectures on $N_X(p)$* , volume 11 of *Chapman & Hall/CRC Research Notes in Mathematics*. CRC Press, Boca Raton, FL, 2012.
- [30] Tetsuji Shioda. Algebraic cycles on abelian varieties of Fermat type. *Math. Ann.*, 258(1):65–80, 1981/82.
- [31] Andrew V. Sutherland. Sato-Tate distributions. In *Analytic methods in arithmetic geometry*, volume 740 of *Contemp. Math.*, pages 197–248. Amer. Math. Soc., Providence, RI, 2019.
- [32] Richard Taylor. Automorphy for some l -adic lifts of automorphic mod l Galois representations. II. *Publ. Math. Inst. Hautes Études Sci.*, (108):183–239, 2008.

DEPARTMENT OF MATHEMATICS, BROOKLYN COLLEGE, CITY UNIVERSITY OF NEW YORK; 2900
 BEDFORD AVENUE, BROOKLYN, NY 11210 USA
 Email address: heidi.goodson@brooklyn.cuny.edu