

Analytic AC conductivities from holography

Jie Ren and Wenni Zheng

School of Physics, Sun Yat-sen University, Guangzhou, 510275, China

(Dated: September 17, 2021)

We find exact, analytic solutions of the holographic AC conductivity at arbitrary frequency ω and temperature T , in contrast to previous works where the AC conductivity was analytically obtained usually at small ω and T . These solutions enable us to study the analyticity properties of the current-current correlator $G(\omega)$ in detail. The first system we study is the AdS₅ planar black hole with momentum dissipation, whose extremal limit has an AdS₂ factor. Then we study an AdS₄ Einstein-dilaton system whose special cases are maximal gauged supergravities. The solutions show how the poles move and how branch cuts emerge as the temperature varies. As a byproduct, we obtain an R -current correlator in $\mathcal{N} = 4$ super-Yang-Mills theory on a sphere at finite temperature in the large N and strong coupling limit.

INTRODUCTION

The AdS/CFT correspondence [1–3] provides a powerful tool to study strongly interacting quantum systems, where the perturbative method does not work. An example is condensed matter systems without quasiparticles [4–6]. A key observable to calculate is the electric conductivity, which is related to the current-current correlation function by the Kubo formula. Analyticity properties of correlation functions are of great interest to theoretical physicists. There have been a large number of works on calculating the conductivity. Specifically, the DC conductivity can often be calculated analytically.

However, analytic solutions of AC conductivities are rare. Known solutions are usually from constant curvature spacetimes, such as pure AdS or the BTZ black hole [7, 8]. Another known solution is from the planar AdS₅ black hole, corresponding to the R -current correlator of the $\mathcal{N} = 4$ super-Yang-Mills (SYM) theory [9, 10]. However, in this system, all finite temperatures are equivalent. It is desirable to obtain analytic solutions of the AC conductivity that has a nontrivial temperature dependence. The analyticity properties of the conductivity can be studied as the temperature varies and the extremal limit is approached.

Momentum dissipation exists in real-world materials. Without any momentum dissipation, the DC conductivity for finite density systems is divergent. A simple way to introduce momentum dissipation is by adding axions (massless scalar fields) in the system [11–13]. We consider zero density systems for simplicity. Since the momentum dissipation introduces another scale, the system has a nontrivial temperature dependence for the planar black hole. In the extremal limit, the IR geometry may have an AdS₂ factor or be a hyperscaling-violating geometry.

In this manuscript, we add momentum dissipation in zero density systems that are closely related to maximal gauged supergravities. We obtain a number of analytic solutions of the AC conductivity in an AdS₅ system and a one-parameter family of AdS₄ systems. Analyticity prop-

erties of the retarded current-current correlator $G_R(\omega)$ are studied. At finite temperature, there are poles in the lower-half complex ω -plane. When the extremal limit is taken, the poles will become branch cuts via different moving patterns. Finally, as a byproduct, an R -current correlator in SYM theory on a sphere at finite temperature is analytically calculated.

EINSTEIN-DILATON-AXION SYSTEMS

We study zero density systems dual to AdS _{$d+1$} with momentum dissipation. The background geometry is determined by Einstein gravity, a neutral scalar ϕ , and axions χ_i ($i = 1, \dots, d-1$). The holographic conductivity is calculated by perturbing the system with a Maxwell field A_μ . The Lagrangian density is [14, 15]

$$\mathcal{L} = R - \frac{1}{4}Z(\phi)F^2 - \frac{1}{2}(\partial\phi)^2 - V(\phi) - \frac{1}{2}\sum_{i=1}^{d-1}(\partial\chi_i)^2, \quad (1)$$

where $\chi_i = ax_i$ satisfies the equations of motion and breaks the translation symmetry; a is a parameter describing the strength of momentum dissipation. We consider solutions of neutral planar AdS black holes with the metric

$$ds^2 = -f(r)dt^2 + f(r)^{-1}dr^2 + U(r)d\vec{x}^2. \quad (2)$$

The conductivity as a function of frequency in strongly coupled systems dual to the AdS gravity can be calculated from the retarded current-current correlator by the AdS/CFT prescription. To calculate the AC conductivity, we perturb the system by an alternating electric field along the $x \equiv x_1$ direction, which is achieved by adding a vector potential $\delta A_x(r, t) = e^{-i\omega t}A_x(r)$. The Maxwell equation for the zero density system is given by

$$\frac{1}{f(r)Z(\phi)}(f(r)Z(\phi)A_x')' + \frac{\omega^2}{f(r)^2}A_x(r) = 0, \quad (3)$$

where the prime denotes derivative with respect to the radial coordinate r . To obtain the retarded Green's function, we impose the infalling boundary condition at the

horizon $r = r_h$: $A_x \propto (r - r_h)^{-i\omega/|f'(r_h)|}$. At the AdS boundary, the leading and subleading terms of $A_x(r)$ are

$$A_x(r) = \mathcal{A} + \frac{\mathcal{B}}{r^{d-2}} + \frac{\tilde{\mathcal{B}}}{r^{d-2}} \log(\Lambda r), \quad (4)$$

where Λ is a cutoff scale taken to be 1. In AdS₅, $\tilde{\mathcal{B}} = \mathcal{A}\omega^2 L^4/2$, while in AdS₄, $\tilde{\mathcal{B}} = 0$. According to the AdS/CFT prescription [2, 3, 16], \mathcal{A} and \mathcal{B} are the dual source of the current \mathcal{O} and its expectation value $\langle \mathcal{O} \rangle$ in the boundary field theory, respectively. Then the retarded Green's function corresponding to the current-current correlator $\langle J^x J^x \rangle$ is [17]

$$G(\omega) = \begin{cases} \mathcal{B}/\mathcal{A} & \text{for AdS}_4, \\ 2\mathcal{B}/\mathcal{A} - \omega^2 L^4/2 & \text{for AdS}_5. \end{cases} \quad (5)$$

By the Kubo formula, the conductivity is $\sigma(\omega) = G/i\omega$.

PLANAR BLACK HOLE IN ADS₅

We start with the planar Schwarzschild-AdS₅ black hole with momentum dissipation. We take $V = -12/L^2$, $Z = 1$ and $\phi = 0$ in Eq. (1). The blackening factor of the metric with $U(r) = r^2$ is

$$f(r) = \frac{1}{L^2} \left(r^2 + r_h^2 - \frac{a^2 L^2}{4} \right) \left(1 - \frac{r_h^2}{r^2} \right), \quad (6)$$

where r_h is the horizon size. The Hawking temperature of this black hole is

$$T = \frac{8r_h^2 - a^2 L^2}{8\pi r_h L^2}. \quad (7)$$

The extremal limit is at $a = 2\sqrt{2}r_h/L$, at which the IR geometry is AdS₂ \times \mathbb{R}^3 . As a comparison, the geometry at $T = 0$ without axions is the pure AdS. Without loss of generality, we set $L = 1$ and $r_h = 1$.

The solution of the perturbation equation (3) with the infalling boundary condition at the horizon is

$$A_x = \left(\frac{r^2 - 1}{r^2 - 1 + 2\pi T} \right)^{-\frac{i\omega}{4\pi T}} {}_2F_1 \left(-\frac{i\omega(1 - \sqrt{1 - 2\pi T})}{4\pi T}, -\frac{i\omega(1 + \sqrt{1 - 2\pi T})}{4\pi T}; 1 - \frac{i\omega}{2\pi T}; \frac{r^2 - 1}{r^2 - 1 + 2\pi T} \right), \quad (8)$$

where the parameter a in the metric has been replaced by the temperature T . To obtain the Green's function, we expand $A_x(r)$ at asymptotic infinity as Eq. (4). By Eq. (5), the Green's function is

$$G(\omega) = i\omega - \frac{1}{2}\omega^2 \left[\psi \left(1 - \frac{i\omega(1 - \sqrt{1 - 2\pi T})}{4\pi T} \right) + \psi \left(1 - \frac{i\omega(1 + \sqrt{1 - 2\pi T})}{4\pi T} \right) + \log(2\pi T) + 2\gamma \right], \quad (9)$$

where ψ is the digamma function $\psi(z) = \Gamma'(z)/\Gamma(z)$ and γ is Euler's constant. The poles of the Green's function are at

$$\omega_n = -2in(1 \pm \sqrt{1 - 2\pi T}), \quad n = 1, 2, 3, \dots \quad (10)$$

Interestingly, the real part of the conductivity can be expressed by an elementary function as

$$\text{Re}[\sigma(\omega)] = \frac{\pi\omega}{4} \left(\coth \frac{(1 - \sqrt{1 - 2\pi T})\omega}{4T} + \coth \frac{(1 + \sqrt{1 - 2\pi T})\omega}{4T} \right). \quad (11)$$

From the indefinite integral over the real part of the conductivity, we obtain the following identity

$$\int_0^\infty \left(\text{Re}[\sigma(\omega)] - \frac{\pi\omega}{2} \right) d\omega = \frac{\pi}{3}(1 - \pi T) = \frac{\pi a^2}{24}. \quad (12)$$

When $a = 0$, i.e., without the momentum dissipation, the right-hand side is zero, which matches the sum rule in [18]. This implies that the sum rule depends on the conservation of momentum.

When the axions vanish ($a = 0$), the poles of the retarded Green's function are distributed below the real axis in the complex ω -plane, where the solution can be found in Ref. [9]. As the temperature decreases, i.e., a increases, the poles move towards the negative imaginary axis. Once a equals a critical value $a_0 = 2r_h/L$, the poles become constrained on the negative imaginary axis. As the temperature is lowered, the poles become denser. More precisely, the poles which were on the right of the imaginary axis become sparser, and the poles which were on the left become denser. The quasinormal frequencies as poles of the Green's function are shown in Fig. 1.

When the temperature is zero, $A_x(r)$ with the infalling boundary condition at the horizon can be solved by the Whittaker functions $W_{\kappa, \mu}(z)$:

$$A_x = W_{\frac{i\omega}{4}, \frac{1}{2}} \left(-\frac{i\omega}{r^2 - 1} \right). \quad (13)$$

The Green's function is

$$G(\omega) = i\omega - \frac{1}{2}\omega^2 \left[\psi \left(1 - \frac{i\omega}{4} \right) + \log(-i\omega) + 2\gamma \right]. \quad (14)$$

There are poles at $\omega_n = -4in$ ($n = 1, 2, 3, \dots$) and a branch cut due to the logarithmic term on the negative imaginary axis of the complex ω -plane, which matches perfectly with the extremal limit of the finite temperature result.

CONDUCTIVITIES IN ADS₄

If we change the dimension to $d = 3$ and perform the same calculation, we obtain that the conductivity




FIG. 1. Quasinormal modes in the complex ω -plane for the Schwarzschild black hole in AdS₅ with momentum dissipation as the temperature decreases.

is a constant. Without momentum dissipation, it is well known that the conductivity is a constant due to an electromagnetic self-duality [19]. However, nonconstant conductivity in AdS₄ can be obtained in other systems. We take the potential of the scalar field ϕ given by [20]

$$V(\phi) = -\frac{2}{(1+\alpha^2)^2 L^2} \left[\alpha^2 (3\alpha^2 - 1) e^{-\phi/\alpha} + 8\alpha^2 e^{(\alpha-1/\alpha)\phi/2} + (3-\alpha^2) e^{\alpha\phi} \right], \quad (15)$$

and $Z(\phi) = e^{-\alpha\phi}$ [21], where α is a parameter, and the values of $\alpha = 1/\sqrt{3}$, 1, and $\sqrt{3}$ correspond to special cases of STU supergravity. We call them 3-charge, 2-charge, and 1-charge black holes in AdS₄, respectively [22].

There is a neutral planar black hole solution for the system (1) with the potential (15) [23]. The factors $f(r)$ and $U(r)$ in the metric (2) and the scalar field ϕ are

$$f(r) = -\frac{a^2}{2} \left(1 - \frac{b}{r}\right)^{\frac{1-\alpha^2}{1+\alpha^2}} + \frac{r^2}{L^2} \left(1 - \frac{b}{r}\right)^{\frac{2\alpha^2}{1+\alpha^2}},$$

$$U(r) = r^2 \left(1 - \frac{b}{r}\right)^{\frac{2\alpha^2}{1+\alpha^2}}, \quad e^{\alpha\phi} = \left(1 - \frac{b}{r}\right)^{\frac{2\alpha^2}{1+\alpha^2}}. \quad (16)$$

For $\alpha \neq 0$, there are curvature singularities at $r = b$ and $r = 0$. When $a \neq 0$, the curvature singularities are always enclosed by a horizon at $r = r_h$ at finite temperature. Either a or b will be replaced by the black hole horizon size r_h as needed.

First, we consider the case without momentum dissipation. When $a = 0$, Eq. (16) describes a spacetime without a regular horizon. Since the AdS boundary is at $r \rightarrow \infty$, the IR is at $r = b$ when $b > 0$, and at $r = 0$ when $b < 0$.

This geometry can be regarded as the extremal limit of a regular black hole, if the Gubser criterion is satisfied [24]. For the above background with $a = 0$, the perturbation equation (3) can be solved by Bessel functions as

$$A_x = C_1 \sqrt{1 - \frac{b}{r}} J_\nu \left(\frac{2\nu\omega L^2}{b} \left(1 - \frac{b}{r}\right)^{\frac{1}{2\nu}} \right) + C_2 \sqrt{1 - \frac{b}{r}} K_\nu \left(-\frac{2i\nu\omega L^2}{b} \left(1 - \frac{b}{r}\right)^{\frac{1}{2\nu}} \right), \quad (17)$$

where

$$\nu = \frac{1 + \alpha^2}{2(1 - \alpha^2)}. \quad (18)$$

It is crucial to examine the boundary conditions in the IR [25]. We assume $b > 0$ first, in which case the IR is at $r = b$. If $0 < \alpha < 1$, there is no infalling wave, and we require that the solution is normalizable at $r = b$. Thus the solution is Eq. (17) with $C_2 = 0$ [26]. When $b > 0$ and $0 < \alpha < 1$, the Green's function is

$$G(\omega) = -\omega L^2 \frac{J_{\nu-1}(2\nu\omega L^2/b)}{J_\nu(2\nu\omega L^2/b)}. \quad (19)$$

The Green's function has poles on the real axis of the complex ω -plane. If $\alpha > 1$, we can impose infalling boundary condition in the IR, and it is more convenient to choose the modified Bessel function K_ν (or Hankel function $H_\nu^{(1)}$). When $b > 0$ and $\alpha > 1$, the Green's function is

$$G(\omega) = i\omega L^2 \frac{K_{\nu-1}(2i\nu\omega L^2/b)}{K_\nu(2i\nu\omega L^2/b)}. \quad (20)$$

The Green's function has a branch cut on the negative imaginary axis of the complex ω -plane. When $\alpha = 1$, the result is Eq. (31) below. For all α , the real part of the conductivity has a delta function at $\omega = 0$, despite this is a zero density system.

For the case $b < 0$, the IR geometry is at $r = 0$. When $0 < \alpha < 1$, there is infalling wave in the IR, and the Green's function is Eq. (20). When $\alpha > 1$, the boundary condition in the IR is normalizability, and the Green's function is Eq. (19). The conductivity no longer has a delta function at $\omega = 0$. Depending on the parameters b and α , the above dilatonic model has rich features analogous to strongly correlated electronic systems, such as the twisted bilayer graphene [27].

When the momentum dissipation is nonzero, the solution (16) describes finite temperature black holes. In the following, we obtain analytic solutions of the holographic AC conductivity from the 3-charge and 2-charge black holes in AdS₄ with momentum dissipation.

3-CHARGE BLACK HOLE IN ADS₄

The solution for the 3-charge black hole in AdS₄ is given by Eq. (16) with $\alpha = 1/\sqrt{3}$. The temperature is

[28]

$$T = \frac{\sqrt{r_h(r_h - b)}}{2\pi L^2}, \quad (21)$$

where $r_h = aL/\sqrt{2}$. The solution of the perturbation equation (3) with the infalling boundary condition at the horizon is

$$A_x = \frac{r-b}{r+r_h} \left(\frac{r-r_h}{r+r_h} \right)^{-\frac{i\omega}{4\pi T}} {}_2F_1 \left(\tilde{a}, \tilde{b}; \tilde{c}; \tilde{x} \frac{r-r_h}{r+r_h} \right), \quad (22)$$

where ${}_2F_1(\tilde{a}, \tilde{b}; \tilde{c}; \tilde{x})$ is the hypergeometric function and

$$\begin{aligned} \tilde{a} &= 1 - \frac{i\omega L^2}{2} \left(\frac{1}{\sqrt{r_h(r_h - b)}} - \frac{1}{\sqrt{r_h(r_h + b)}} \right), \\ \tilde{b} &= 1 - \frac{i\omega L^2}{2} \left(\frac{1}{\sqrt{r_h(r_h - b)}} + \frac{1}{\sqrt{r_h(r_h + b)}} \right), \\ \tilde{c} &= 1 - \frac{i\omega L^2}{\sqrt{r_h(r_h - b)}}, \quad \tilde{x} = \frac{b+r_h}{b-r_h}. \end{aligned} \quad (23)$$

By the AdS/CFT prescription, the Green's function is given by

$$G(\omega) = -b - r_h \left(\tilde{c} + \frac{2\tilde{a}\tilde{b}\tilde{x}}{\tilde{c}} \cdot \frac{{}_2F_1(1 + \tilde{a}, 1 + \tilde{b}; 1 + \tilde{c}; \tilde{x})}{{}_2F_1(\tilde{a}, \tilde{b}; \tilde{c}; \tilde{x})} \right). \quad (24)$$

We need to treat the zero temperature limit more carefully, since the $b < 0$ and $b > 0$ cases have different extremal geometries. When $b < 0$, the extremal limit is at $r_h = 0$, in which case the axions vanish ($a = 0$). The conductivity is given by Eq. (19). When $b > 0$, the extremal limit is at $r_h = b$, in which case the axions do not vanish. In this extremal geometry, we solve the perturbation equation (3) and obtain the Green's function at $T = 0$. After imposing the infalling boundary condition in the IR, the solution of $A_x(r)$ is

$$A_x = K_{\tilde{\nu}} \left(-\tilde{\nu} \sqrt{\frac{r+b}{r-b}} \right), \quad \tilde{\nu} = \frac{i\omega L^2}{\sqrt{2b}}. \quad (25)$$

The Green's function is

$$G(\omega) = \frac{i\omega L^2}{\sqrt{2}} \left(\frac{K_{\tilde{\nu}+1}(-\tilde{\nu})}{K_{\tilde{\nu}}(-\tilde{\nu})} + 1 \right). \quad (26)$$

The Green's function has a branch cut on the negative imaginary axis of the complex ω -plane.

2-CHARGE BLACK HOLE IN ADS₄

The solution for the 2-charge black hole in AdS₄ is given by Eq. (16) with $\alpha = 1$. The temperature is

$$T = \frac{2r_h - b}{4\pi L^2} = \frac{2r_h^2 + a^2 L^2}{8\pi r_h L^2}, \quad (27)$$

where $a^2 L^2 = 2r_h(r_h - b)$. There is a minimum temperature. The extremal limit is always at $a = 0$, in contrast to the 3-charge black hole in AdS₄. The extremal geometry apparently has a finite temperature despite there is no horizon enclosing the spacetime singularity. However, the dimensionless temperature $\tilde{T} = T/a$ is infinite in the extremal limit. The solution of the perturbation equation (3) with the infalling boundary condition at the horizon is

$$A_x = \left(\frac{r-r_h}{r+r_h-b} \right)^{-\frac{i\omega}{4\pi T}} H\ell \left(\frac{(2r_h-b)b}{r_h^2}, -\frac{i\omega L^2 b}{r_h^2}; 0, 2, 1 - \frac{2i\omega L^2}{2r_h-b}, 1; \frac{b(r-r_h)}{r_h(r-b)} \right), \quad (28)$$

where $H\ell(a, q; \alpha, \beta, \gamma, \delta; z)$ is the (local) Heun function. The Green's function is given by

$$G(\omega) = iL^2\omega - \frac{(r_h-b)b}{r_h} \times \frac{H\ell' \left(\frac{(2r_h-b)b}{r_h^2}, -\frac{i\omega L^2 b}{r_h^2}; 0, 2, 1 - \frac{2i\omega L^2}{2r_h-b}, 1; \frac{b}{r_h} \right)}{H\ell \left(\frac{(2r_h-b)b}{r_h^2}, -\frac{i\omega L^2 b}{r_h^2}; 0, 2, 1 - \frac{2i\omega L^2}{2r_h-b}, 1; \frac{b}{r_h} \right)}. \quad (29)$$

where $H\ell'(a, q; \alpha, \beta, \gamma, \delta; z)$ is the derivative of the Heun function.

The conductivity without the momentum dissipation has been obtained in Ref. [15] for $b > 0$, and an AdS₅ counterpart was obtained in Ref. [29]. The following result is for general b . When the axions vanish ($a = 0$), the solution of $A_x(r)$ is

$$A_x = C_1 \left(\frac{r-b}{r} \right)^{\frac{b+\sqrt{b^2-4\omega^2 L^4}}{2b}} + C_2 \left(\frac{r-b}{r} \right)^{\frac{b-\sqrt{b^2-4\omega^2 L^4}}{2b}}. \quad (30)$$

When $2\omega L^2 > |b|$, the first term with the $\omega \rightarrow \omega + i\epsilon$ prescription describes infalling wave in the IR for either $b > 0$ or $b < 0$. Thus the infalling boundary condition requires $C_2 = 0$. The Green's function is given by

$$G(\omega) = -\frac{1}{2} \left(b + \sqrt{b^2 - 4\omega^2 L^4} \right). \quad (31)$$

There are branch cuts $4\omega^2 L^4 > b^2$ on the real axis of the complex ω -plane. For small a , the branch cuts become dense poles in the lower-half plane. As a increases, the poles become increasingly sparser and closer to the imaginary axis of the complex ω -plane. The quasinormal frequencies as poles of the Green's function are shown in Fig. 2.

The conductivity is obtained by $\sigma = G/i\omega|_{\omega \rightarrow \omega + i\epsilon}$, and the real part is

$$\text{Re}[\sigma(\omega)] = \pi b \theta(b) \delta(\omega) + \theta(4\omega^2 L^4 - b^2) \frac{\sqrt{4\omega^2 L^4 - b^2}}{2\omega}, \quad (32)$$




FIG. 2. Quasinormal modes in the complex ω -plane for the 2-charge black hole in AdS₄ with momentum dissipation as the extremal limit is approached.




FIG. 3. Real part of the conductivity as a function of frequency plotted for the 2-charge black hole in AdS₄. We take $b = 1$ and $L = 1$. The red, green, blue, and purple curves are for $r_h = 1.5, 1.2, 1.01, \text{ and } 1.001$, respectively.

where $\theta(x)$ is the Heaviside step function. There is a delta function at $\omega = 0$ when $b > 0$ and there is a gap for $0 < 2\omega L^2 < |b|$. When a is close to zero, we expect that the delta function becomes a Drude-like peak. Note that the system is at zero density. It was pointed out that this type of delta function at $\omega = 0$ together with the hard gap is closely related to superconductivity [29], and an example was given in Ref. [30]. Figure 3 shows the real part of the conductivity calculated from Eq. (29) as the extremal limit is approached.

R-CURRENT CORRELATOR OF $\mathcal{N} = 4$ SYM ON A SPHERE

Although the planar black holes with momentum dissipation are from the bottom-up approach, they are closely related to spherical black holes in the top-down approach. The blackening factor for a hyperbolic black hole is the same as a planar black hole with momentum dissipation [14, 31, 32]. By analytic continuation of the solution for a hyperbolic black hole, we can obtain the solution for a spherical black hole.

An R -current correlator of $\mathcal{N} = 4$ SYM on a sphere

at finite temperature in the large N and strong coupling limit is obtained as follows, as a generalization of the previous result in which the CFT lives on flat space [9, 10]. The metric for the spherical Schwarzschild-AdS₅ black hole is

$$ds^2 = -f(r)dt^2 + f(r)^{-1}dr^2 + r^2 d\Omega_3^2, \quad (33)$$

where $d\Omega_3^2 = d\rho^2 + \sin^2 \rho (d\theta^2 + \sin^2 \theta d\varphi^2)$ is the metric for a 3-dimensional sphere with unit radius. The solution of $f(r)$ is Eq. (6) with $a^2 = -4$ [33]. The boundary CFT lives on a sphere with radius L . The temperature is given by

$$T = \frac{2r_h^2 + L^2}{2\pi r_h L^2}. \quad (34)$$

The black hole has a minimum temperature at $r_h = L/\sqrt{2}$. There is a Hawking-Page phase transition between the large black hole and thermal AdS [34, 35].

We perturb the system by [36]

$$\delta A_\varphi = e^{-i\omega t} A_\varphi(r). \quad (35)$$

The Maxwell equation is exactly the same as Eq. (3), provided that we use the new $f(r)$ for the spherical geometry. The solution with the infalling boundary condition at the horizon is

$$A_\varphi = \left(\frac{r^2 - r_h^2}{r^2 + r_h^2 + L^2} \right)^{-\frac{i\omega}{4\pi T}} {}_2F_1 \left(\frac{\omega L^2 (\sqrt{r_h^2 + L^2} - ir_h)}{2(2r_h^2 + L^2)}, \right. \\ \left. - \frac{\omega L^2 (\sqrt{r_h^2 + L^2} + ir_h)}{2(2r_h^2 + L^2)}; 1 - \frac{i\omega L^2 r_h}{2r_h^2 + L^2}; \frac{r^2 - r_h^2}{r^2 + r_h^2 + L^2} \right). \quad (36)$$

By the AdS/CFT prescription, the Green's function is obtained as

$$G(\omega) = -i\omega L^2 r_h - \frac{1}{2}\omega^2 L^4 \left[\psi \left(\frac{\omega L^2 (\sqrt{r_h^2 + L^2} - ir_h)}{2(2r_h^2 + L^2)} \right) \right. \\ \left. + \psi \left(-\frac{\omega L^2 (\sqrt{r_h^2 + L^2} + ir_h)}{2(2r_h^2 + L^2)} \right) + 2\gamma + \log(2r_h^2 + L^2) \right]. \quad (37)$$

The poles of the Green's function are at

$$\omega_n L^2 = 2n \left(\pm \sqrt{r_h^2 + L^2} - ir_h \right), \quad (38)$$

where $n = 1, 2, 3, \dots$.

DISCUSSION

We have obtained analytic solutions of the AC conductivity at arbitrary frequency and temperature from the planar Schwarzschild-AdS₅ black hole and a dilatonic system related to STU supergravity in AdS₄ with momentum dissipation. With the analytic solutions at hand, we

study the analyticity properties of the Green's function, especially how the poles move as the temperature varies. Typically, branch cuts emerge when the extremal limit is approached. The real part of the conductivity can have a delta function at $\omega = 0$ and a hard gap, depending on the parameters. The system may describe insulators, conductors, or superconductors.

A number of aspects need more study in the future. (i) We have not explored the full parameter space for the dilatonic model in AdS_4 . It may shed some light on understanding strongly correlated materials. (ii) The 1-charge black hole in AdS_4 can also give a solution for conductivity in terms of the Heun function, but the result is more involved. (iii) It is unclear that how the strongly coupled result relates to the weakly coupled result as we tune the coupling constant [37]. It would be interesting to make a comparison between the correlators calculated by AdS/CFT and field theory techniques.

JR thanks Christopher Herzog, Elias Kiritsis, and Li Li for helpful discussions. This work was supported in part by the NSF of China under Grant No. 41030001.

-
- [1] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2**, 231 (1998) [hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, Phys. Lett. B **428**, 105 (1998) [hep-th/9802109].
- [3] E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. **2**, 253 (1998) [hep-th/9802150].
- [4] S. A. Hartnoll, A. Lucas and S. Sachdev, *Holographic quantum matter*, arXiv:1612.07324.
- [5] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, *Building a holographic superconductor*, Phys. Rev. Lett. **101**, 031601 (2008) [arXiv:0803.3295].
- [6] T. Faulkner, H. Liu, J. McGreevy and D. Vegh, *Emergent quantum criticality, Fermi surfaces, and AdS_2* , Phys. Rev. D **83**, 125002 (2011) [arXiv:0907.2694].
- [7] M. Banados, C. Teitelboim and J. Zanelli, *The black hole in three-dimensional space-time*, Phys. Rev. Lett. **69**, 1849 (1992) [hep-th/9204099].
- [8] J. Ren, *One-dimensional holographic superconductor from $\text{AdS}_3/\text{CFT}_2$ correspondence*, JHEP **11**, 055 (2010) [arXiv:1008.3904].
- [9] P. K. Kovtun and A. O. Starinets, *Quasinormal modes and holography*, Phys. Rev. D **72**, 086009 (2005) [hep-th/0506184].
- [10] R. C. Myers, A. O. Starinets and R. M. Thomson, *Holographic spectral functions and diffusion constants for fundamental matter*, JHEP **11**, 091 (2007) [arXiv:0706.0162].
- [11] T. Andrade and B. Withers, *A simple holographic model of momentum relaxation*, JHEP **05**, 101 (2014) [arXiv:1311.5157].
- [12] R. A. Davison and B. Gouteraux, *Momentum dissipation and effective theories of coherent and incoherent transport*, JHEP **01**, 039 (2015) [arXiv:1411.1062].
- [13] M. Baggioli, K. Y. Kim, L. Li and W. J. Li, *Holographic axion model: a simple gravitational tool for quantum matter*, Sci. China Phys. Mech. Astron. **64**, 270001 (2021) [arXiv:2101.01892].
- [14] B. Gouteraux, *Charge transport in holography with momentum dissipation*, JHEP **04**, 181 (2014) [arXiv:1401.5436].
- [15] E. Kiritsis and J. Ren, *On Holographic insulators and supersolids*, JHEP **09**, 168 (2015) [arXiv:1503.03481].
- [16] D. T. Son and A. O. Starinets, *Minkowski space correlators in AdS/CFT correspondence: recipe and applications*, JHEP **09**, 042 (2002) [hep-th/0205051].
- [17] G. T. Horowitz and M. M. Roberts, *Holographic superconductors with various condensates*, Phys. Rev. D **78**, 126008 (2008) [arXiv:0810.1077].
- [18] D. R. Gulotta, C. P. Herzog and M. Kaminski, *Sum rules from an extra dimension*, JHEP **01**, 148 (2011) [arXiv:1010.4806].
- [19] C. P. Herzog, P. Kovtun, S. Sachdev and D. T. Son, *Quantum critical transport, duality, and M-theory*, Phys. Rev. D **75**, 085020 (2007) [hep-th/0701036].
- [20] C. J. Gao and S. N. Zhang, *Dilaton black holes in de Sitter or anti-de Sitter universe*, Phys. Rev. D **70** (2004) 124019 [hep-th/0411104].
- [21] When $Z = 1$, the conductivity is a constant for arbitrary α and a .
- [22] There are $U(1)^4$ gauge fields in STU supergravity in AdS_4 with charges Q_i . We call the system 3-charge black hole if $Q_1 = Q_2 = Q_3 = Q$ and $Q_4 = 0$; 2-charge black hole if $Q_1 = Q_2 = Q$ and $Q_3 = Q_4 = 0$; 1-charge black hole if $Q_1 = Q$ and $Q_2 = Q_3 = Q_4 = 0$.
- [23] M. M. Caldarelli, A. Christodoulou, I. Papadimitriou and K. Skenderis, *Phases of planar AdS black holes with axionic charge*, JHEP **04**, 001 (2017) [arXiv:1612.07214].
- [24] S. S. Gubser, *Curvature singularities: the good, the bad, and the naked*, Adv. Theor. Math. Phys. **4**, 679 (2000) [hep-th/0002160].
- [25] C. Charmousis, B. Gouteraux, B. S. Kim, E. Kiritsis and R. Meyer, *Effective holographic theories for low-temperature condensed matter systems*, JHEP **11**, 151 (2010) [arXiv:1005.4690].
- [26] For certain values of α , there are ambiguities for the boundary condition in the IR. See [15, 25].
- [27] Y. Cao *et al.*, *Unconventional superconductivity in magic-angle graphene superlattices*, Nature **556**, 43 (2018) [arXiv:1803.02342].
- [28] For the 1-charge black hole in AdS_4 , the temperature is Eq. (21) with $r_h = aL/\sqrt{2} - b$.
- [29] O. DeWolfe, S. S. Gubser and C. Rosen, *Fermi surfaces in $\mathcal{N} = 4$ super-Yang-Mills theory*, Phys. Rev. D **86**, 106002 (2012) [arXiv:1207.3352].
- [30] F. Aprile, *Holographic superconductors in a cohesive phase*, JHEP **10**, 009 (2012) [arXiv:1206.5827].
- [31] Y. Bardoux, M. M. Caldarelli and C. Charmousis, *Shaping black holes with free fields*, JHEP **05**, 054 (2012) [arXiv:1202.4458].
- [32] J. Ren, *Phase transitions of hyperbolic black holes in anti-de Sitter space*, arXiv:1910.06344.
- [33] By setting $a^2 = 4$, we can obtain a hyperbolic black hole; the boundary theory lives in a hyperbolic space with radius L .
- [34] S. W. Hawking and D. N. Page, *Thermodynamics of black holes in anti-de Sitter space*, Commun. Math. Phys. **87**,

- 577 (1983).
- [35] E. Witten, *Anti-de Sitter space, thermal phase transition, and confinement in gauge theories*, Adv. Theor. Math. Phys. **2**, 505 (1998) [hep-th/9803131].
- [36] The same perturbation equation can be obtained by $\delta A_\rho = e^{-i\omega t} A_\rho(r) \csc^2 \rho$ or $\delta A_\theta = e^{-i\omega t} A_\theta(r) \csc \theta$.
- [37] S. A. Hartnoll and S. P. Kumar, *AdS black holes and thermal Yang-Mills correlators*, JHEP **12**, 036 (2005) [hep-th/0508092].