

LOCAL PARAMETERS OF SUPERCUSPIDAL REPRESENTATIONS

WEE TECK GAN, MICHAEL HARRIS, AND WILL SAWIN

ABSTRACT. Let G be a connected reductive group over the non-archimedean local field F and let π be a supercuspidal representation of $G(F)$. The local Langlands conjecture posits that to such a π can be attached a parameter $L(\pi)$, which is an equivalence class of homomorphisms from the Weil-Deligne group with values in the Langlands L -group ${}^L G$ over an appropriate algebraically closed field C of characteristic 0. When F is of positive characteristic p then Genestier and Lafforgue have defined a parameter, $L^{ss}(\pi)$, which is a homomorphism $W_F \rightarrow {}^L G(C)$ that is *semisimple* in the sense that, if the image of $L^{ss}(\pi)$, intersected with the Langlands dual group $\hat{G}(C)$, is contained in a parabolic subgroup $P \subset \hat{G}(C)$, then it is contained in a Levi subgroup of P . If the Frobenius eigenvalues of $L^{ss}(\pi)$ are pure in an appropriate sense, then the local Langlands conjecture asserts that the image of $L^{ss}(\pi)$ is in fact *irreducible* – its image is contained in no proper parabolic P . In particular, unless $G = GL(1)$, $L^{ss}(\pi)$ is ramified: it is non-trivial on the inertia subgroup $I_F \subset W_F$. In this paper we prove, at least when G is split and semisimple, that this is the case provided π can be obtained as the induction of a representation of a compact open subgroup $U \subset G(F)$, and provided the constant field of F is of order greater than 3. Conjecturally every π is compactly induced in this sense, and the property was recently proved by Fintzen to be true as long as p does not divide the order of the Weyl group of G . The proof is an adaptation of the globalization method of [GLo] when the base curve is \mathbb{P}^1 , and a simple application of Deligne’s Weil II.

CONTENTS

1. Introduction
- Acknowledgments
2. Local systems on open subsets of \mathbb{P}^1
 - 2.1. Purity
 - 2.2. Kummer theory

Date: October 1, 2021.

W.T. Gan was partially supported by a Singapore government MOE Tier 1 grant R-146-000-320-114.

M. Harris was partially supported by NSF Grant DMS-2001369, and by a Simons Foundation Fellowship, Award number 663678 .

W. Sawin was partially supported by NSF grant DMS-2101491 and served as a Clay Research Fellow.

3. Global and local Langlands-Lafforgue parameters
 - 3.1. Review of the results of Lafforgue and Genestier-Lafforgue
 - 3.2. Ramification and pure semisimple representations
4. Compact open subgroups of $G(\mathbf{A}_{k(t)})$.
5. Poincaré series and the Main Theorem: split case
6. Wild ramification
7. Pure Weil-Deligne parameters
 - 7.1. Tempered parameters and pure Weil-Deligne parameters
 - 7.2. An application of the purity of the monodromy weight filtration
 - 7.3. Existence of pure completions of local parameters of supercuspidal representations
 - 7.4. Generalization to discrete series representations
8. Base change and incorrigible representations
 - 8.1. Existence of tempered base change
 - 8.2. The case of $GL(n)$
9. Applications to groups over p -adic fields
- References

1. INTRODUCTION

Let G be a connected reductive group over the non-archimedean local field F and let π be a supercuspidal representation of $G(F)$. The local Langlands conjecture posits that to such a π can be attached a parameter $\mathcal{L}(\pi)$, which is a homomorphism from the Weil-Deligne group with values in the Langlands L -group ${}^L G$ over an appropriate algebraically closed field C of characteristic 0. When F is of positive characteristic p then Genestier and Lafforgue have defined a parameter, $\mathcal{L}^{ss}(\pi)$, which is a homomorphism $W_F \rightarrow {}^L G(C)$ that is *semisimple* in the sense that, if the image of $\mathcal{L}^{ss}(\pi)$, intersected with the Langlands dual group $\hat{G}(C)$, is contained in a parabolic subgroup $P \subset \hat{G}(C)$, then it is contained in a Levi subgroup of P [GLa]. If the Frobenius eigenvalues of $\mathcal{L}^{ss}(\pi)$ are pure in an appropriate sense, then the local Langlands conjecture asserts that the image of $\mathcal{L}^{ss}(\pi)$ is in fact *irreducible* – its image is contained in no proper parabolic P . In particular, unless $G = GL(1)$, $\mathcal{L}^{ss}(\pi)$ is ramified: it is non-trivial on the inertia subgroup $I_F \subset W_F$.

In what follows we take $C = \overline{\mathbb{Q}_\ell}$ as the coefficient field for the Genestier-Lafforgue parameter. In this paper we prove the following theorem.

Theorem 1.1. *Suppose F is the local non-archimedean field $k((t))$, where $k = \mathbb{F}_q$ is a finite field of order $q = p^m$ for some prime p . Let G be a split semisimple group over F . Assume $q > 5$ and π is a supercuspidal (irreducible) representation of $G(F)$ that can be obtained as the induction of a representation of a compact open subgroup $U \subset G(F)$:*

$$\pi \xrightarrow{\sim} \mathfrak{c} - \text{Ind}_U^G(F)\tau$$

for some (irreducible) smooth representation τ of U with coefficients in $\overline{\mathbb{Q}_\ell}$. Suppose finally that $\mathcal{L}^{ss}(\pi)$ is pure: if $\text{Frob}_F \in W_F$ is any Frobenius element, then the eigenvalues of $\mathcal{L}^{ss}(\pi)(\text{Frob}_F)$ are all Weil q -numbers of the same weight. (In this case we say that π is a pure supercuspidal representation.)

Then $\mathcal{L}^{ss}(\pi) : W_F \rightarrow {}^L G(\overline{\mathbb{Q}_\ell})$ is a ramified parameter: the image of the inertia group I_F under $\mathcal{L}^{ss}(\pi)$ is non-trivial.

Conjecturally every π is compactly induced in this sense, and the property was recently proved by Fintzen to be true as long as p does not divide the order of the Weyl group of G . When $G = \text{GL}(n)$ the result is part of the local Langlands correspondence proved by Laumon, Rapoport, and Stuhler [LRS93]. When G is a split classical group the result was proved in most cases by Ganapathy and Varma by comparison with Arthur’s proof of the local Langlands correspondence over p -adic fields, by the method of close local fields [GV, A]. Thus the novelty of this paper is that, like the Genestier-Lafforgue result, Theorem 1.1 is valid uniformly for all split semisimple groups, including exceptional groups. (In a future draft we plan to extend the result to quasi-split reductive groups.)

Of course, Theorem 1.1 is a much weaker statement than the expected local Langlands correspondence, which not only asserts that $\mathcal{L}^{ss}(\pi)$ is irreducible when π is pure but gives a complete description of the L -packet of representations with the given local parameter. However, all proofs of the local Langlands correspondence for $\text{GL}(n)$ begin with the observation that a supercuspidal representation π of $\text{GL}(n, F)$ is not *incorrigible* [H19] – that it becomes a constituent of a principal series representation after a finite series of base changes corresponding to cyclic extensions F_{i+1}/F_i , $i = 0, \dots, n$, where $F_0 = F$. Since one can choose the F_i so that the Galois parameter restricted to $W_{F_{n+1}}$ is unramified, this is obvious if we know that

- (1) no supercuspidal representation of any Levi factor of $\text{GL}(n, F)$ other than $\text{GL}(1)^n$ has an unramified parameter, and
- (2) the parametrization is compatible with cyclic stable base change.

In the proofs of the local Langlands correspondence in [LRS93, HT01, He00], point (1) is proved by reference to Henniart’s numerical local correspondence [He88], whereas point (2) is proved by a global method. In Scholze’s proof of the local correspondence for $\text{GL}(n)$ [Sch13], condition (1) is proved by a geometric argument using a study of nearby cycles in an integral model of the Lubin-Tate local moduli space.

Starting with point (1), the local correspondence is deduced by a study of the fibers of (stable) base change for cyclic extensions of p -adic fields. This is ultimately a consequence of deep properties – the “advanced theory” – of the Arthur-Selberg trace formula, that are established by Arthur and Clozel in [AC]. For $\text{GL}(n)$ over a number field, the trace formula and its twisted analogue are automatically stable, but there is as yet no stable trace formula for general groups over function fields. If and when a stable trace formula is developed in this generality, it is likely that Theorem 1.1 will provide the

starting point for an inductive proof of the local correspondence, at least for supercuspidal representations with pure parameters.

The proof is an adaptation of the globalization method of [GLo] when the base curve is $Y = \mathbb{P}^1$, and a simple application of Deligne's work [De80]. Let K denote the global function field $k(Y) = k(t)$. By the method of Poincaré series used in [GLo], we construct a cuspidal automorphic representation Π of $G(\mathbf{A}_K)$ that is unramified outside the set of places of K corresponding to the set $Y(k)$ of k -rational points of Y . At the points $v = 0$ and $v = \infty$ we arrange for Π to be tamely ramified, with $\Pi_\infty^{I_-(p)} \neq 0$ and $\Pi_0^{I_+(p)} \neq 0$; here I_- and I_+ denote opposite minimal parahoric subgroups of $G(k(\frac{1}{t}))$ and $G(k(t))$, respectively; $I_-(p) \subset I_-$ and $I_+(p) \subset I_+$ are the corresponding maximal pro- p subgroups. At the other rational points v we assume Π_v to be isomorphic to the fixed supercuspidal π . For the sake of this sketch, we assume π to be compactly induced from an open subgroup of a hyperspecial maximal compact. Moreover, we assume Π_0 contains a sufficiently regular character α of the Iwahori subgroup I^+ . V. Lafforgue attaches to such a Π a local system $L(\pi)$ over $Y \setminus \mathbb{P}^1(k)$. Suppose $L^{ss}(\pi)$ is pure and unramified. By Deligne [De80], we then know that $L(\pi)$ extends to a local system over \mathbb{G}_m . Provided $q > 3$, we can choose α to obtain a contradiction.

The proof of Theorem 1.1 is contained in the first five sections. The remaining five sections provide refinements and extensions of the methods used in its proof. In §6 we prove that supercuspidal representations compactly induced from sufficiently small compact mod center subgroups have wildly ramified Galois parameters, provided they are pure. In §7 we consider supercuspidal representations that are not necessarily pure and prove, in nearly all cases, that their semisimple Galois parameters can be extended to Weil-Deligne parameters that satisfy purity of the monodromy weight filtration, in an appropriate sense. The result does not depend on realizing the representation by compact induction. In this way we can attach full Weil-Deligne parameters to arbitrary supercuspidal representations, and the method extends to general discrete series representations if we admit a version of the stable trace formula over function fields. In §8 we explain the implications of our results for incorrigible representations and prove a weak version of the local Langlands correspondence for $GL(n)$: that all supercuspidal representations of $GL(n, F)$ are either pure with ramified Galois parameters or have wildly ramified parameters. Finally, in §9 we propose the obvious conjecture linking the Fargues-Scholze parametrization of representations of groups over p -adic fields to the Genestier-Lafforgue parametrization, by means of the Deligne-Kazhdan method of close local fields.

ACKNOWLEDGMENTS

The authors thank Hélène Esnault, Tony Feng, Jessica Fintzen, Dennis Gaitsgory, Tasho Kaletha, Jean-Pierre Labesse, Bertrand Lemaire, and Zhiwei Yun for helpful discussions of various aspects of the paper. We benefited

from exchanges with Laurent Fargues and Peter Scholze regarding the material in the final section. Finally, we are especially grateful to Luis Lomelí for his role, in collaboration with one of us, in developing the main techniques on which our results are based, and for his comments in the early stages of this project.

2. LOCAL SYSTEMS ON OPEN SUBSETS OF \mathbb{P}^1

2.1. Purity. Let $k = \mathbb{F}_q$ be a finite field of characteristic p . Let $Z, X \subset \mathbb{P}^1(k)$ be disjoint finite subsets, $|Z|$ and $|X|$ the corresponding reduced subschemes of $\mathbb{P}^1(k)$, and let $Y = \mathbb{P}_k^1 \setminus |X| \cup |Z|$, viewed as a scheme over $\text{Spec}(k)$. Let L be a finite extension of \mathbb{Q}_ℓ and consider a local system \mathcal{V} over Y with coefficients in L (a lisse ℓ -adic sheaf). Picking a point y of Y (the generic point, for example), and letting V denote the finite-dimensional vector space \mathcal{V}_y over L , we can identify \mathcal{V} with a representation

$$\rho : \pi_1(Y, y) \rightarrow \text{Aut}(V).$$

Let $F_z = k(Y)_z$ be the completion of $k(Y)$ at the point $z \in Z$. The monodromy of \mathcal{V} at z is the restriction of ρ (up to conjugation) to a homomorphism

$$(2.1) \quad \rho_z : \text{Gal}(F_z^{sep}/F_z) \rightarrow \text{Aut}(V).$$

Let $I_z \subset \text{Gal}(F_z^{sep}/F_z)$ denote the inertia group and let $\text{Frob}_z \in \text{Gal}(F_z^{sep}/F_z)$ denote any geometric Frobenius element; i.e., any lift to an automorphism of F_z^{sep} of the automorphism $x \mapsto x^{\frac{1}{q}}$ of \bar{k} .

If Γ is a topological group and $\rho : \Gamma \rightarrow \text{Aut}(V)$ is a continuous homomorphism, we let ρ^{ss} denote its semisimplification – the sum of its Jordan-Hölder constituents.

Theorem 2.2 ([De80], Corollary 1.7.6). *Suppose that Z is not empty and, for all $z \in Z$, the following conditions are satisfied:*

- ρ_z^{ss} is unramified;
- the Frobenius eigenvalues $\rho_z(\text{Frob}_z)$ are all Weil q -numbers of the same weight.

Then \mathcal{V} extends to a local system over $\mathbb{P}_k^1 \setminus |X|$.

Proof. In fact, Deligne’s result only requires the Frobenius eigenvalues to have the same weight for a single complex embedding; under this condition, $\rho_z(I_z)$ is a finite group. But we have assumed that ρ_z^{ss} is unramified, thus $\rho_z(I_z)$ is unipotent. It follows that $\rho_z(I_z)$ must be trivial. \square

Corollary 2.3. *Under the hypotheses of Theorem 2.2, suppose $X = \{0, \infty\}$ and \mathcal{V} is semisimple. Suppose moreover that the monodromy of \mathcal{V} at 0 and ∞ is tame. Let*

$$\rho : \pi_1(\mathbb{P}^1 \setminus |X|, y) \rightarrow \text{Aut}(V)$$

be the corresponding monodromy representation. Suppose the semisimplified local monodromy representation at 0 or ∞ is abelian and Frobenius-semisimple.

Then, possibly after replacing L by a finite extension L' , \mathcal{V} breaks up as the sum of 1-dimensional local systems.

Proof. Theorem 2.2 implies that \mathcal{V} extends to a tame local system, also called \mathcal{V} , over \mathbb{G}_m . By [Laf02], Corollary VII.8, \mathcal{V} can be written as a (finite) direct sum

$$\mathcal{V} = \bigoplus \mathcal{V}_\chi \otimes \chi$$

where each \mathcal{V}_χ is a mixed local system and χ is the pullback to $\mathbb{P}^1 \setminus |X|$ of a rank 1 ℓ -adic local system over $\text{Spec}(k)$, in other words a continuous ℓ -adic character of $\text{Gal}(\bar{k}/k)$. We thus reduce to the case where \mathcal{V} is a mixed ℓ -adic local system, and since \mathcal{V} is assumed semisimple we may assume \mathcal{V} to be irreducible. Then the purity hypothesis at the points z in the non-empty set Z implies that \mathcal{V} is itself pure. But the tame fundamental group of $\mathbb{G}_{m,\bar{k}}$ is just $\prod_{p' \neq p} \mathbb{Z}_{p'}(1)$. It follows from the purity that the image J under ρ of the geometric fundamental group of \mathbb{G}_m is finite and necessarily abelian. The hypothesis on the local monodromy at 0 or ∞ implies that the conjugation action of $\text{Gal}(\bar{k}/k)$ on J is trivial, and thus the image under ρ of the arithmetic fundamental group $\pi_1(\mathbb{P}^1 \setminus |X|, y)$ is also abelian; moreover, the image of $\text{Gal}(\bar{k}/k)$ is semisimple by hypothesis. This completes the proof. \square

2.2. Kummer theory. Let $K = k(\mathbb{P}^1) = k(t)$ be the global function field of \mathbb{P}^1 , and let K' denote the union of all abelian extensions of K of degree dividing $q - 1$, which is the order of the group of roots of unity in K . Homomorphisms

$$\alpha : \text{Gal}(K'/K) \rightarrow \overline{\mathbb{Q}}_\ell^\times$$

are classified by Kummer theory:

$$(2.4) \quad \psi : K^\times / (K^\times)^{q-1} \xrightarrow{\sim} X(K) := \text{Hom}(\text{Gal}(K'/K), \overline{\mathbb{Q}}_\ell^\times).$$

Let $C_T \subset K^\times / (K^\times)^{q-1}$ be the subgroup generated by k^\times and the parameter T , and let $X_T = \psi(C_T)$. We also consider Kummer theory for the constant field k :

$$(2.5) \quad \psi_k : k^\times \xrightarrow{\sim} X(k) := \text{Hom}(\text{Gal}(k'/k), \overline{\mathbb{Q}}_\ell^\times),$$

where k' is the cyclic extension of k of order q^{q-1} . For $b \in k^\times$, $\beta \in \text{Gal}(k'/k)$, let $\beta^b = \psi_k(b)$.

Any $z \in k^\times$ defines the geometric point $z \in \mathbb{G}_{m,k}$; let $\Gamma_z \subset \text{Gal}(K'/K)$ denote its decomposition group, I_z its inertia group. An element $\alpha \in X_T$ is unramified at such a point z , and thus defines a homomorphism

$$\alpha^{(z)} : \Gamma_z / I_z \xrightarrow{\sim} \text{Gal}(k'/k) \rightarrow \overline{\mathbb{Q}}_\ell^\times.$$

For $a = 1, \dots, q - 1$ we let $\alpha_a = \psi(T^a)$. Then we have the following reciprocity law:

Lemma 2.6. *For $z \in \mathbb{G}_m(k)$ and $a = 1, \dots, q - 1$ we have*

$$(\alpha_a)^{(z)} = \beta^{az}.$$

Proof. Let K_z denote the completion of K at z , \mathcal{O}_z its integer ring, k_z its residue field, which the inclusion of k in K canonically identifies with k . The image of the element $T \in \mathcal{O}_z$ in k_z is identified with the element $z \in k^\times$, so the residue field of the unramified extension $\mathcal{O}_z[T^{1/q-1}]$ is just $k(z^{1/q-1})$. The identity is then obvious. \square

Lemma 2.7. *Let \mathcal{L} be a rank one local system on $\mathbb{G}_{m,k}$. Then*

$$\mathrm{tr}(\mathrm{Frob}_z, \mathcal{L}) = \mathrm{tr}(\mathrm{Frob}_1, \mathcal{L}) \mathrm{tr}(m_z^{-1}, \mathcal{L})$$

where m_z is the element of the inertia group I_0 which acts by multiplication by a on the $(q-1)$ st root of T .

Proof. (Class field theory) It suffices to show that $\mathrm{Frob}_z = \mathrm{Frob}_1 \cdot m_z^{-1}$ in the abelianization of the tame fundamental group of \mathbb{G}_m . Class field theory gives an isomorphism between this abelianization and the profinite completion of

$$K^\times \backslash \mathbb{A}_K^\times / \left(\prod_{\substack{y \in |\mathbb{P}_k^1| \\ y \notin \{0, \infty\}}} \mathcal{O}_{K_y}^\times \times \prod_{y \in \{0, \infty\}} U_{K_y, 1} \right).$$

Moreover, Frob_z corresponds to the idèle which is a uniformizer at z and the identity at all other places, Frob_1 corresponds to the idèle which is a uniformizer at 1 and the identity at all other places, and m_z corresponds to the idèle which is z at 0 and the identity at all other places. So it suffices to check that the first idèle class is the product of the second two idèle classes. This can be checked by multiplying by $\frac{T-z}{T-1} \in K^\times$, which is a uniformizer at z , the inverse of a uniformizer at 1, and z at 0.

(Kummer theory) We can construct a one-dimensional representation of $\mathrm{Gal}(\bar{k}/k)$ on which Frob_q acts by multiplication by $\mathrm{tr}(\mathrm{Frob}_1, \mathcal{L})$. On tensoring \mathcal{L} with the inverse representation, we reduce to the case where Frob_1 acts trivially on \mathcal{L} . Since the tame fundamental group of \mathbb{G}_m is generated by the geometric tame fundamental group and Frobenius, the image of the Galois group acting on \mathcal{L} must equal the image of the geometric tame fundamental group. The action of the geometric tame fundamental group on \mathcal{L} must be by a Frob_q -invariant character, which means it factors through the Frob_q -coinvariants of the abelianization of the tame geometric fundamental group, which are $\mathbb{Z}/(q-1)$. So the action of the geometric fundamental group is by a character of order $q-1$. This now follows from Lemma 2.6. \square

3. GLOBAL AND LOCAL LANGLANDS-LAFFORGUE PARAMETERS

3.1. Review of the results of Lafforgue and Genestier-Lafforgue.

In this section Y is a smooth projective curve over k and $K = k(Y)$ is the global function field of Y . Let G be a connected reductive algebraic group over K . Let \hat{G} be the Langlands dual group of G with coefficients in $\overline{\mathbb{Q}_\ell}$, and

let ${}^L G = \hat{G} \rtimes \text{Gal}(K^{\text{sep}}/K)$ be the Langlands L -group of G (in the Galois form).

Let $\mathcal{A}_0(G) = \mathcal{A}_0(G, Y)$ denote the set of cuspidal automorphic representations of G with central character of finite order. We let $\mathcal{G}^{\text{ss}}(G)$ denote the set of equivalence classes of compatible families of semisimple ℓ -adic homomorphisms, for $\ell \neq p$:

$$\rho_\ell : \text{Gal}(K^{\text{sep}}/K) \rightarrow {}^L G.$$

The term *semisimple* is understood to mean that if $\rho_\ell(\text{Gal}(K^{\text{sep}}/K)) \cap \hat{G}$ is contained in a parabolic subgroup $P \subset \hat{G}$, then it is contained in a Levi subgroup of P .

If $\nu : G \rightarrow \mathbb{G}_m$ is an algebraic character, with \mathbb{G}_m here designating the split 1-dimensional torus over K , the theory of L -groups provides a dual character $\hat{\nu} : \mathbb{G}_m \rightarrow \hat{G} \subset {}^L G$. If Z is the center of G , and if $c : \mathbb{G}_m \rightarrow Z \subset G$ is a homomorphism, then the theory of L -groups provides an algebraic character ${}^L c : {}^L G \rightarrow \mathbb{G}_m$.

We state Lafforgue's theorem for general reductive groups over general global function fields, but we will mainly be applying it to $K = k(\mathbb{P}^1)$. In what follows, a representation of G over a local field F is “unramified” if it contains a vector invariant under a hyperspecial maximal compact subgroup; in particular, the group itself is assumed to be unramified over F .

Theorem 3.1. *Let K be any global function field and G a connected reductive algebraic group over K . (i) [Laf18][Théorème 0.1] There is a map*

$$\mathcal{L} : \mathcal{A}_0(G) \rightarrow \mathcal{G}^{\text{ss}}(G)$$

with the following property: if v is a place of K and $\Pi \in \mathcal{A}_0(G)$ is a cuspidal automorphic representation such that Π_v is unramified, then $\mathcal{L}(\Pi)$ is unramified at v , and $\mathcal{L}^{\text{ss}}(\Pi)|_{W_{K_v}}$ is the Satake parameter of Π_v .¹

(ii) Suppose $\nu : G \rightarrow \mathbb{G}_m$ is an algebraic character. Suppose

$$\chi : \mathbf{A}_K^\times / K^\times \rightarrow \overline{\mathbb{Q}}_\ell^\times = GL(1, \overline{\mathbb{Q}}_\ell)$$

is any continuous character of finite order. For any $\Pi \in \mathcal{G}_0(G)$, let $\Pi \otimes \chi \circ \nu$ denote the twist of Π by the character $\chi \circ \nu$ of $G(\mathbf{A})$. Then

$$\mathcal{L}(\Pi \otimes \chi \circ \nu) = \mathcal{L}(\Pi) \cdot \hat{\nu} \hat{\chi}$$

where $\hat{\nu}$ is as above and where $\hat{\chi} : \text{Gal}(K^{\text{sep}}/K)^{\text{ab}} \rightarrow GL(1, \overline{\mathbb{Q}}_\ell)$ is the character corresponding to χ by local class field theory.

(iii) [GLa][Théorème 0.1] Let v be a place of K and let $F = K_v$. Then the semisimplification of the restriction of $\mathcal{L}(\Pi)$ to W_{K_v} depends only on F

¹Here and below we will mainly refer to the restriction of a global Galois parameter to the local Weil group, rather than to the local Galois group, because the unramified Langlands correspondence relates spherical representations to unramified homomorphisms of the local Weil group to the L -group. But the difference is inessential.

and the local component Π_v of Π , which is an irreducible admissible representation of $G(F)$, and not on the rest of the automorphic representation Π , nor on the global field K . We denote this parameter $\mathcal{L}^{ss}(\Pi_v)$.

Let F be a non-archimedean local field of characteristic p ; $F \xrightarrow{\sim} k'((t))$ for some finite extension k' of \mathbb{F}_p . Let $\mathcal{A}(G, F)$ denote the set of equivalence classes of admissible irreducible representations of $G(F)$, and let $\mathcal{G}^{ss}(G, F)$ denote the set of equivalence classes of semisimple ℓ -adic homomorphisms

$$\rho : W_F \rightarrow {}^L G.$$

By (iii) above, we thus obtain a (semisimple) parametrization of $\mathcal{A}(G, F)$:

$$(3.2) \quad \mathcal{L}^{ss} : \mathcal{A}(G, F) \rightarrow \mathcal{G}^{ss}(G, F).$$

Of course, the theorem quoted above only constructs $\mathcal{L}^{ss}(\pi)$ when π can be realized as the local component Π_v of a global cuspidal automorphic representation Π , but the statement of Théorème 0.1 of [GLa] includes the extension to a (semisimple) parametrization of all members of $\mathcal{A}(G, F)$. Theorem 3.1 continues:

Theorem 3.3 ([GLa], Théorème 0.1). *(iv) The local parametrization \mathcal{L}^{ss} is compatible with parabolic induction in the following sense: Let $P \subset G$ be an F -rational parabolic subgroup with Levi factor M and let $\sigma \in \mathcal{A}(M, F)$. If $\pi \in \mathcal{A}(G, F)$ is an irreducible constituent of $\text{Ind}_{P(F)}^{G(F)} \sigma$ (normalized induction) and $i_M : {}^L M \rightarrow {}^L G$ is the inclusion of the L -group of M as a Levi factor of a parabolic subgroup in the L -group of G , then*

$$\mathcal{L}^{ss}(\pi) = i_M \circ \mathcal{L}^{ss}(\sigma).$$

(v) In particular, suppose G is quasi-split over F with Borel subgroup B , $T \subset B$ is a maximal torus, $I \subset G(F)$ is an Iwahori subgroup, and $I(p) \subset I$ is its maximal pro- p subgroup. Suppose $\pi^{I(p)} \neq 0$. Then $\mathcal{L}^{ss}(\pi)$ takes values in the L -group of T .

(vi) The local and global parametrizations \mathcal{L} and \mathcal{L}^{ss} are compatible with homomorphisms

$$\Upsilon : G_1 \rightarrow G_2$$

of reductive groups where G_2 is split reductive and the image of Υ is a normal subgroup, in the following sense. Let ${}^L \Upsilon : \hat{G}_2 \rightarrow \hat{G}_1$ be the homomorphism of dual groups corresponding to Υ . Then for any cuspidal automorphic (resp. admissible irreducible) representation π of G_2 over K (resp. over F),

$$\mathcal{L}(\pi \circ \Upsilon) = {}^L \Upsilon \circ \mathcal{L}(\pi)$$

(resp.

$$\mathcal{L}^{ss}(\pi \circ \Upsilon) = {}^L \Upsilon \circ \mathcal{L}^{ss}(\pi).$$

This applies in particular if Υ is an isogeny of split semisimple groups.

Proof. In the situation of (v), the condition $\pi^{I(p)} \neq 0$ implies that π contains an eigenvector for I , and thus is a constituent of a principal series representation. The assertion (v) then follows from (iv). Assertion (vi) in the local setting is the last assertion of [GLa, Théorème 0.1]; in the global setting it is [Laf18, Proposition 12.5]. \square

Corollary 3.4. *Let $\Pi \in \mathcal{A}_0(G)$, and suppose $|X| \subset Y$ is the largest closed subset such that Π_v is unramified for $v \notin |X|$.*

(i) *The parametrization of Theorem 3.1 defines a symmetric monoidal functor from the category $\text{Rep}({}^L G)$ of finite-dimensional algebraic representations of ${}^L G$ to the category of completely reducible local systems on $Y \setminus |X|$: to any $\tau : {}^L G \rightarrow GL(N)$ one attaches the N -dimensional local system attached to the representation*

$$\mathcal{V}(\Pi, \tau) := \tau \circ \mathcal{L}(\Pi)$$

of $\pi_1(Y \setminus |X|)$ (any base point).

(ii) *Similarly, let $\sigma \in \mathcal{A}(G, F)$. The local parametrization \mathcal{L}^{ss} of (3.2) defines a symmetric monoidal functor from the category $\text{Rep}({}^L G)$ of finite-dimensional algebraic representations of ${}^L G$ to the category of completely reducible representations of W_F :*

$$\tau \mapsto \mathcal{L}^{ss}(\sigma, \tau) := \tau \circ \mathcal{L}^{ss}(\sigma).$$

(iii) *Suppose Π has the property that $\mathcal{V}(\Pi, \tau)$ is abelian and semisimple for some faithful representation τ . Then the image of $\mathcal{L}(\Pi)$, intersected with \hat{G} , lies in a maximal torus of \hat{G} .*

Proof. The first assertion is the standard characterization of \hat{G} -local systems as symmetric monoidal functors; the second assertion is obvious. \square

Definition 3.5. *With notation as in Theorem 3.3, let $\sigma \in \mathcal{A}(G, F)$. We say σ is **pure** if, for some (equivalently, for all) faithful representations $\tau : {}^L G \rightarrow GL(N)$, $\mathcal{L}^{ss}(\sigma, \tau)$ has the property that, for any Frobenius element $\text{Frob}_F \in W_F$, the eigenvalues of $\mathcal{L}^{ss}(\sigma, \tau)(\text{Frob})$ are all Weil q -numbers of the same weight.*

3.2. Ramification and pure semisimple representations. We now return to our assumption that $K = k(\mathbb{P}^1)$.

In the statement of the following theorem, F is the local field $k((t))$. We assume henceforward that G is a split connected reductive group over $K = k(\mathbb{P}^1) = k(t)$ that is unramified at all places of K ; at every place y of K we let $B_y = T_y \cdot N_y \subset G(K_y)$ denote a fixed Borel subgroup, with T_y a maximal torus and N_y the unipotent radical of B_y . Let I and $I(p)$ be as in Theorem 3.3 (v).

Theorem 3.6. *Let $F = k((t))$ and let σ be a supercuspidal representation of $G(F)$. Let $K = k(\mathbb{P}^1)$ and suppose there exists a cuspidal automorphic representation Π of $G(\mathbf{A}_K)$, with the following properties:*

- (a) *At every $z \in \mathbb{G}_m(k) \subset \mathbb{P}^1(k)$, $\Pi_z \xrightarrow{\sim} \sigma$.*

- (b) The representation Π_∞ has a vector invariant under $I(p)$.
- (c) The representation Π_0 is a constituent of a principal series $\text{Ind}_{B_0(k((t)))}^{G(k((t)))} \chi$, where χ is a character of $T_0(k((t)))$.
- (d) The local components Π_y of Π for all $y \notin \mathbb{P}^1(k)$ are unramified.

Suppose in addition

- (e) The semisimple local parameter $\mathcal{L}^{ss}(\sigma)$ is pure.

Suppose $q > 3$. Let χ_k denote the restriction of χ to the subgroup $T_y(k) \subset T_y(k((t)))$. Then if

- (f) χ_k is the composition of a homomorphism $T_y(k) \rightarrow \mathbb{G}_m(k)$ induced by a character $T_y(k) \rightarrow \mathbb{G}_m(k)$ which corresponds to a root α of G with a faithful character $\mathbb{G}_m(k) \rightarrow \overline{\mathbb{Q}}_\ell^\times$

then the semisimple local parameter $\mathcal{L}^{ss}(\sigma)$ is ramified – its restriction to the inertia subgroup of $\text{Gal}(F^{sep}/F)$ is non-trivial.

Proof. Assume $\mathcal{L}^{ss}(\sigma)$ is unramified.

Let β be the root of \hat{G} dual to α . Let $\tau : \hat{G} \rightarrow GL(N)$ be the representation with highest weight β . Consider the (semisimple) local system $\mathcal{V}(\Pi, \tau)$ on $\mathbb{P}^1 \setminus |\mathbb{P}^1(k)|$. Let $z \in \mathbb{G}_m(k)$, $\Gamma_z \subset \pi_1(\mathbb{P}^1 \setminus |\mathbb{P}^1(k)|)$ its decomposition group, and let

$$\mathcal{L}(\Pi, \tau)_z : \Gamma_z \rightarrow GL(N, \overline{\mathbb{Q}}_\ell)$$

denote the local monodromy representation at z corresponding to $\mathcal{V}(\Pi, \tau)$, $\mathcal{L}(\Pi, \tau)_z^{ss}$ its semisimplification. By property (a) and Theorem 3.1, (iii),

$$(3.7) \quad \mathcal{L}(\Pi, \tau)_z^{ss} \xrightarrow{\sim} \mathcal{L}^{ss}(\sigma, \tau);$$

in other words, they are equivalent in the \hat{G} -conjugacy class of homomorphisms from W_F to ${}^L G$. By our assumption, $\mathcal{L}(\Pi, \tau)_z^{ss}$ is therefore unramified. Hypothesis (e) and Theorem 2.2 then imply that $\mathcal{V}(\Pi, \tau)$ extends to a local system on \mathbb{G}_m . Properties (b) and (c), and Theorem 3.3 (iv), imply that the monodromy representations at 0 and ∞ are tamely ramified and satisfy the hypotheses of Corollary 2.3. It follows that $\mathcal{V}(\Pi, \tau)$ breaks up as the sum of 1-dimensional local systems $\mathcal{L}_1, \dots, \mathcal{L}_n$. It now follows from (3.7) that the local parameters $\mathcal{L}(\Pi, \tau)_z^{ss}$, for $z \in \mathbb{G}_m(k)$, are all conjugate under the action of the Weyl group of \hat{G} . In particular, using Lemma 2.7,

$$(3.8) \quad \text{tr}(\text{Frob}_z, \mathcal{V}(\Pi, \tau)) = \sum_{i=1}^N \text{tr}(\text{Frob}_z, \mathcal{L}_i) = \sum_{i=1}^n \text{tr}(\text{Frob}_1, \mathcal{L}_i) \text{tr}(m_z, \mathcal{L}_i)$$

is a constant function of z .

Let us now evaluate $\text{tr}(m_z, \mathcal{L}_i)$. The restriction of $\mathcal{V}(\Pi, \tau)$ to the Galois group of $k((t))$ is the image under τ of the Genestier-Lafforgue parameter of Π_0 , which by assumption (c) and Theorem 3.1 is the character $\hat{\chi} : \text{Gal}(k((t))^{sep}/k((t))) \rightarrow \hat{T}(\overline{\mathbb{Q}}_\ell)$ corresponding to χ under class field theory. Hence the restriction $\hat{\chi}_k$ of $\hat{\chi}$ to the tame inertia k^\times in the abelianization of $\text{Gal}(k((t))^{sep}/k((t)))$ corresponds to $\chi_k : T(\mathbb{F}_q) \rightarrow \overline{\mathbb{Q}}_\ell^\times$. In particular, because

χ_k is the character of T corresponding to α composed with a faithful character $\chi_0 : \mathbb{F}_q^\times \rightarrow \overline{\mathbb{Q}_\ell}^\times$, $\hat{\chi}_k$ is the cocharacter of \hat{T} corresponding to α composed with χ_0 .

Thus the restriction of $\mathcal{V}(\Pi, \tau)$ to the tame inertia group of $k((t))$ is the map $\chi_0 : \mathbb{F}_q^\times \rightarrow \overline{\mathbb{Q}_\ell}^\times$ composed with the composition

$$\overline{\mathbb{Q}_\ell}^\times \rightarrow \hat{T}(\overline{\mathbb{Q}_\ell}^\times) \rightarrow \hat{G}(\overline{\mathbb{Q}_\ell}^\times) \rightarrow GL_N(\overline{\mathbb{Q}_\ell}^\times)$$

where the first map is associated to the coroot α of \hat{G} , the second the inclusion of the maximal torus, and the third the representation associated to β . This composition defines a representation

$$\overline{\mathbb{Q}_\ell}^\times \rightarrow GL_N(\overline{\mathbb{Q}_\ell}^\times)$$

that can be equivalently viewed as the sum over coweights β' of τ of the one-dimensional representation $x \mapsto x^{\langle \alpha, \beta' \rangle}$ of $\overline{\mathbb{Q}_\ell}^\times$.

Since any two decompositions of a representation into one-dimensional summands have isomorphic one-dimensional summands, we can choose for each i from 1 to n a weight β' of τ such that

$$\mathrm{tr}(m_z, \mathcal{L}_i) = \chi_0(z)^{\langle \alpha, \beta_i \rangle}$$

and each weight β' occurs a number of times equal to its multiplicity in τ . Because χ_0 is a character of order $q - 1$, and $q - 1 > 2$, $z \mapsto \chi_0(z)^2$ is a nontrivial character of \mathbb{F}_q^\times . Thus by orthogonality of characters and (3.8)

(3.9)

$$\begin{aligned} 0 &= \sum_{z \in \mathbb{F}_q^\times} \chi_0(z)^{-2} \mathrm{tr}(\mathrm{Frob}_z, \mathcal{V}(\Pi, \tau)) = \sum_{z \in \mathbb{F}_q^\times} \chi_0(z)^{-2} \sum_{i=1}^n \mathrm{tr}(\mathrm{Frob}_1, \mathcal{L}_i) \chi_0(z)^{\langle \alpha, \beta_i \rangle} \\ &= \sum_{i=1}^n \sum_{z \in \mathbb{F}_q^\times} \chi_0(z)^{-2} \mathrm{tr}(\mathrm{Frob}_1, \mathcal{L}_i) \chi_0(z)^{\langle \alpha, \beta_i \rangle} = \sum_{\substack{i \in \{1, \dots, n\} \\ \langle \alpha, \beta_i \rangle \equiv 2 \pmod{q-1}}} \mathrm{tr}(\mathrm{Frob}_1, \mathcal{L}_i) (q-1). \end{aligned}$$

Because α is the coroot dual to β , we have $\langle \alpha, \beta \rangle = 2$. Because β is the highest weight of τ , for any root β' not equal to β , we have $-2 \leq \langle \alpha, \beta' \rangle < 2$, so, because $\langle \alpha, \beta' \rangle \in \mathbb{Z}$, we have $\langle \alpha, \beta' \rangle \in \{-2, -1, 0, 1\}$. Hence because $q > 1$, $\langle \alpha, \beta' \rangle$ is not congruent to 2 mod $q - 1$ unless $\beta' = \beta$. Furthermore, because β is the highest weight, it occurs with multiplicity 1. So in fact the sum (3.9) has length 1, and so its unique term $\mathrm{tr}(\mathrm{Frob}_1, \mathcal{L}_i)(q - 1)$ must vanish, but this is impossible as \mathcal{L}_i is a one-dimensional representation, giving the desired contradiction. \square

4. COMPACT OPEN SUBGROUPS OF $G(\mathbf{A}_{k(t)})$.

Let us first prove a technical lemma on the matrix entries of elements of parahoric subgroups that will be useful in this section and later.

Lemma 4.1. *Let G be an unramified semisimple group over a local field $K = \kappa((t))$. Let T be a maximal split torus of G . Let c be a point in the apartment of T in the Bruhat-Tits building and let G_c be the associated parahoric subgroup.*

Let $\rho: G \rightarrow \text{Aut}(V)$ be a faithful representation of G , and let e_1, \dots, e_n be a basis of V on which T acts diagonally. Assume V and e_1, \dots, e_n to be defined over κ . (This is always possible because G is unramified and thus defined over κ .) Let $\lambda_i \in X^(T)$ be the cocharacter by which T acts on e_i .*

Then, for any $g \in G_c \subset G(K)$, we have

$$(4.2) \quad v(\rho_{ij}(g)) \geq c \cdot (\lambda_i - \lambda_j).$$

Furthermore, if g lies in the Moy-Prasad subgroup $G_{c,r+}$ for some $r \geq 0$, then we have

$$(4.3) \quad v(\rho_{ij}(g) - \delta_{ij}) > c \cdot (\lambda_i - \lambda_j) + r.$$

Proof. We give the proof in the G_c case, and then describe the modifications necessary for $G_{c,r+}$.

If (4.2) holds for g_1 and g_2 , then it holds for $g_1 g_2$, so it suffices to verify (4.2) for a set of generators of the parahoric G_c . In the description given by Moy and Prasad, the parahoric G_c is generated by the subgroups U_ψ for affine functions ψ on the apartment of T such that $\psi(c) \geq 0$. Here ψ is an affine function of the form $\psi(w) = b \cdot w - s$ for a root b and a real number s , $U_b \subset G$ is a root group carrying a natural isomorphism to \mathbb{G}_a , and U_ψ consists of elements sent to elements with valuation $\geq s$ under that isomorphism.

To obtain (4.2), it suffices to check that for $g \in U_b$, if $\rho_{ij}(g) \neq 0$ then $\lambda_i - \lambda_j = nb$ for some $n \geq 0$, and that $\rho_{ij}(g)$ is a homogeneous polynomial function on U_b defined over k of degree n , as then

$$v(\rho_{ij}(g)) \geq ns \geq ns + n\psi(c) = nb \cdot c = (\lambda_i - \lambda_j) \cdot c.$$

The group G has a lift from the field κ to the Witt vectors $W(\kappa)$, unique up to not-necessarily-unique isomorphism. Since irreducible representations are classified by Galois orbits of highest-weights, and κ and $W(\kappa)$ have the same Galois action, we may lift the representation V from the field κ to its Witt vectors $W(\kappa)$. Both these vanishing and homogeneous polynomiality properties are preserved by reduction mod p , and thus may be checked over the fraction field of $W(\kappa)$, where U_b is obtained by exponentiating the root b , and where they are automatic.

For $G_{c,r+}$, note that (4.3) is also stable under matrix multiplication since $r \geq 0$. Because the subgroups $G_{c,r+}$ are generated by U_ψ for those ψ with $\psi(c) > r$ and the subgroups $Z_{n\delta}^a$ for a maximal torus Z containing T , for $n > r$.

That these U_ψ satisfy (4.3) by the same facts that $\lambda_i - \lambda_j = nb$ for $n \geq 0$ if ρ_{ij} is nonzero on U_b , that $\rho_{ij}(g)$ is a homogeneous polynomial of degree n , and the addition fact that ρ_{ij} is the constant function 1 on U_b if $n = 0$,

as then

$$v(\rho_{ij}(g)) \geq ns \geq ns + n\psi(c) + nr = nb \cdot c + nr > (\lambda_i - \lambda_j) \cdot c + r$$

if $n > 0$ and $v(\rho_{ij}(g) - 1) = \infty$ if $n = 0$.

So it remains to show that elements $g \in Z_{n\delta}^a$ satisfy (4.3). Because Z commutes with T , if $g \in Z_{n\delta}^a \subseteq Z$ then $\rho_{ij}(g) = 0$ unless $\lambda_i = \lambda_j$, so it suffices to show $v(\rho_{ij}(g) - 1) \geq n > r$. Here a is a simple root of Z in G , and $Z_{n\delta}^a$ is defined as a subset of the maximal torus Z_a of the copy of SL_2 arising from the root a consisting of elements that, in the natural identification of the maximal torus of SL_2 with \mathbb{G}_m , differ from 1 by an element of valuation at least n . These roots and this copy of SL_2 may be defined over \bar{k} , so the action of the torus must be, and therefore the matrix entries are given by polynomial functions on \mathbb{G}_m over \bar{k} , which are equal to δ_{ij} at 1, and thus the difference between $\rho_{ij}(g)$ and 1 has valuation at least n , as desired. \square

In this section G is a split semisimple group over $K = k(t)$, with Borel subgroup B containing maximal split torus T as above. Let $U^* \subset G(k((t)))$ be a fixed maximal compact subgroup. We choose an open compact subgroup of $G(\mathbf{A}_{k(t)})$ of the form $U = \prod_x U_x$, where x runs over the places of K , with the following properties.

- (a) At every $z \in \mathbb{G}_m(k) \subset \mathbb{P}^1(k)$, U_z equals the fixed U^* , with respect to a chosen isomorphism

$$G(K_z) \xrightarrow{\sim} G(k((t)));$$

- (b) For $x \notin |\mathbb{P}^1(k)|$, U_x is the hyperspecial maximal compact subgroup $G(\mathcal{O}_x)$;
 (c) The subgroups U_0 and U_∞ are parahoric subgroups that remain to be chosen.

Proposition 4.4. *The subgroups U_0 and U_∞ can be chosen in such a way that*

$$G(K) \cap U = T(k).$$

Remark 4.5. *Suppose $U_z = G(\mathcal{O}_z)$ for $z \in \mathbb{G}_m(k)$. Then we can take $U_0 = I^+$ and $U_\infty = I^-$ to be respectively the upper and lower Iwahori subgroups; by this we mean that U_0 (resp. U_∞) is the Iwahori subgroup corresponding to the positive (resp. negative) root system corresponding to our chosen Borel subgroup B . Then*

$$G(K) \cap U \subset G(K) \cap \prod_x G(\mathcal{O}_x).$$

The right-hand side is the group of global maps from \mathbb{P}_k^1 to the affine group G_k . Any such map must be constant, because \mathbb{P}^1 is projective, and thus must belong to $G(k)$ because it is defined over k . Thus

$$G(K) \cap U = (U_0 \times U_\infty) \cap G(k) = T(k).$$

Proof. Let Λ be the cocharacter lattice of T , and view $\Lambda \otimes \mathbb{R}$ as an apartment in the Bruhat-Tits building of G ; let $y \in \Lambda \otimes \mathbb{R}$ be the fixed point of U^* . Choose $a, b \in \Lambda \otimes \mathbb{R}$ such that

$$a + b + (q - 1)y = 0$$

and such that a is sufficiently generic.

Define $f(v)$ for a valuation v of K by $f(v) = a$, $f(\infty) = b$, $f(z) = y$ for any $z \in \mathbb{G}_m(k)$, and $f(x) = 0$ if $x \notin |\mathbb{P}^1(k)|$. Under this definition, U_v is the parhoric $G_{f(v)}$ associated to $f(v)$, so U consists of elements which at each place v lie in $G_{f(v)}$.

We are now ready to prove that $G(K) \cap U = T(k)$. To do this, we choose a representation V and a basis e_1, \dots, e_n , where for all i , e_i is an eigenvalue of T with eigenvector a cocharacter λ_i , as in Lemma 4.1

Now if $g \in U \cap G(K)$ and $\rho_{ij}(g) \neq 0$ for some i, j , then we have

$$0 = \sum_v v(\rho_{ij}(g)) \geq \sum_v f(v) \cdot (\lambda_i - \lambda_j) = 0 \cdot (\lambda_i - \lambda_j) = 0.$$

Because both sides must be 0, the inequality is sharp, so

$$v(\rho_{ij}(g)) = f(v) \cdot (\lambda_i - \lambda_j)$$

for all v . This implies that $f(v) \cdot (\lambda_i - \lambda_j)$ is an integer, and in particular that $a \cdot (\lambda_i - \lambda_j)$ is an integer. Since, by assumption, a is generic, we can assume that $a \cdot (\lambda_i - \lambda_j)$ is not an integer for any i, j from 1 to n , except if $\lambda_i = \lambda_j$.

We conclude that, for $g \in U$, $\rho_{ij}(g) = 0$ unless $\lambda_i = \lambda_j$. In other words, $\rho(g)$ commutes with $\rho(T)$, and thus $g \in T(K)$. Furthermore, for i, j with $\rho_{ij}(g) \neq 0$, we have

$$v(\rho_{ij}(g)) = f(v) \cdot (\lambda_i - \lambda_j) = f(v) \cdot 0 = 0$$

for all v , so $\rho_{ij}(g) \in k$, and thus $g \in T(k)$, so $U = T(k)$, as desired. \square

The following is then obvious.

Corollary 4.6. *Under the hypotheses of Proposition 4.4, one can choose Iwahori subgroups $I_0 \subset U_0$ and $I_\infty \subset U_\infty$ such that, if $I_\infty(p) \subset I_\infty$ is the maximal pro- p subgroup, then*

$$G(K) \cap \prod_{x \neq 0, \infty} U_x \times I_\infty(p) \times I_0 = \{1\}.$$

5. POINCARÉ SERIES AND THE MAIN THEOREM: SPLIT CASE

We continue to assume G split semisimple. Let $F = k((t))$ and let σ be a supercuspidal representation of $G(F)$. We say σ contains an \mathfrak{s} -type, if there is an open compact subgroup $U^* \subset G(F)$ and an irreducible representation ρ of U^* such that

$$(5.1) \quad \sigma \xrightarrow{\sim} c - \text{Ind}_{U^*}^{G(F)} \rho.$$

Theorem 5.2. *Let $K = k(T) = k(\mathbb{P}^1)$. Let $F = k((t))$ and let σ be a supercuspidal representation of $G(F)$. Suppose*

$$(5.3) \quad \sigma \text{ contains an } \mathfrak{s} - \text{type}(U^*, \rho).$$

Then there exists a cuspidal automorphic representation Π of $G(K)$ satisfying the conditions (a)-(d) and (f) of Theorem 3.6.

Proof. The construction follows the method of [GLo]. Let $U = \prod_x U_x \subset G(\mathbf{A}_K)$ be an open compact subgroup as in Proposition 4.4. Hypothesis (5.3) implies that σ has a matrix coefficient φ^* with support in U^* , with the property that $\varphi^*(1) = 1$. Define a function $\varphi = \prod_x \varphi_x : G(\mathbf{A}_K) \rightarrow \mathbb{C}$ as follows.

- (a) At every $z \in \mathbb{G}_m(k) \subset \mathbb{P}^1(k)$, $\varphi_z = \varphi^*$;
- (b) For $x \notin |\mathbb{P}^1(k)|$, φ_x is the characteristic function of the hyperspecial maximal compact subgroup $G(\mathcal{O}_x)$;
- (c) φ_∞ is the characteristic function of the group $I_\infty(p)$ of Corollary 4.6;
- (d) φ_0 is the character $\chi_k : I_0/I_0(p) \rightarrow \mathbb{C}^\times$, where χ_k is as in the proof of Theorem 3.6.

Define the Poincaré series $P_\varphi : G(K) \backslash G(\mathbf{A}_K) \rightarrow \mathbb{C}$:

$$P_\varphi(g) = \sum_{\gamma \in G(K)} \varphi(\gamma \cdot g).$$

The sum converges absolutely, as in [GLo]. It follows from Corollary 4.6 that

$$P_\varphi(1) = \prod_{z \in \mathbb{G}_m(k)} \varphi^*(1) = 1.$$

In particular, $P_\varphi \neq 0$. □

Corollary 5.4. *Let G be split semisimple over a non-archimedean local field F of characteristic p , with constant field k of order $q > 3$. Suppose p does not divide the order of the Weyl group of G . Let $K = k(T) = k(\mathbb{P}^1)$ and let σ be a supercuspidal representation of $G(F)$. Then there exists a cuspidal automorphic representation Π of $G(K)$ satisfying the conditions (a)-(d) and (f) of Theorem 3.6.*

In particular, if σ is pure and $q > 3$ then $\mathcal{L}^{ss}(\sigma)$ is not unramified.

Proof. Thanks to the results of [Fi], the supercuspidal representation σ of $G(F)$ contains an \mathfrak{s} -type by our hypothesis on p . The claim now follows immediately from Theorems 3.6 and 5.2. □

Corollary 5.5. *Let $F = k((t))$ and let σ be an irreducible representation of $G(F)$. Suppose the Genestier-Lafforgue parameter $\mathcal{L}^B(\sigma)$ is pure and unramified. Suppose every supercuspidal representation of every Levi factor of $G(F)$ contains an \mathfrak{s} -type; this is the case in particular if p does not divide*

the order of the Weyl group of G . Then σ is an irreducible constituent of an unramified principal series representation of $G(F)$.

Proof. In any case σ is an irreducible constituent of a representation of the form $\text{Ind}_{P(F)}^{G(F)} \tau$ for some parabolic subgroup $P(F) = L(F) \cdot N(F) \subset G(F)$, where $L(F)$ is a Levi subgroup of $P(F)$, $N(F)$ its unipotent radical, and τ is a supercuspidal representation of $L(F)$. Since the Genestier-Lafforgue parametrization is compatible with parabolic induction by Theorem 3.3, it follows that $\mathcal{L}^{ss}(\tau)$ is unramified. Since τ is supercuspidal, it follows from Theorem 5.2 3.6 that $L(F)$ must be a torus and τ an unramified character. \square

6. WILD RAMIFICATION

Let G be a reductive group over a finite field \mathbb{F}_q , and let H be a finite subgroup of $G(\mathbb{F}_q)$.

We say that H is *sufficiently small* if there is a Borel subgroup B of G such that $B(\mathbb{F}_q) \cap H$ is trivial.

One example of a sufficiently small subgroup H is the maximal unipotent subgroup of a Borel (by the Bruhat decomposition).

Remark 6.1. *This condition could likely be weakened and the below argument would still work. In principle as long as $B(\mathbb{F}_q) \cap H$ is contained in a proper algebraic subgroup of the maximal torus of B , this should be fine for q sufficiently large.*

For G a reductive group over a local field $\mathbb{F}_q((t))$ of equal characteristic p , and G_x a parahoric subgroup, we say $J \subseteq G_x$ is *sufficiently small* if its projection to the reductive group $G_x/G_{x,0+}$ is sufficiently small. For example, $G_{x,0+}$ is sufficiently small.

Proposition 6.2. *Let G be an unramified reductive group over a local field $\mathbb{F}_q((t))$ of equal characteristic p , G_x a parahoric subgroup, and J an open subgroup of G_x . Let ρ be a representation of J such that $\pi = \text{c-Ind}_J^{G(\mathbb{F}_q((t)))} \rho$ is irreducible.*

If J is sufficiently small and $q > 2$, then the Genestier-Lafforgue parameter of π is wildly ramified.

Proof. We can assume G is defined over \mathbb{F}_q .

We will prove this by constructing two different automorphic forms on $G(\mathbb{F}_q(t))$ whose local representation at 0 is ρ , with different local representations at ∞ examining their global Lafforgue parameters, deducing if the Genestier-Lafforgue parameter of π is tame that they have the same local parameter at ∞ , and deriving a contradiction. We construct the automorphic forms via Poincaré series.

Let x be the point in the Bruhat-Tits building of $G(\mathbb{F}_q((t)))$ whose associated parahoric subgroup is G_x . By conjugation, we may assume x is contained in the apartment of the standard maximal torus. Let y be a small

perturbation of the point $-x$ in the Bruhat-Tits building of $G(\mathbb{F}_q((u))) = G(\mathbb{F}_q((t^{-1})))$. Specifically, we choose $y - x$ to be a small rational multiple of a cocharacter which lies in the interior of a Weyl chamber of G . Because $y - x$ does not lie on the walls of the Weyl chamber, y does not lie on any wall of the apartment that also contains x , and because $y - x$ is small, y does not lie on any wall of the apartment that doesn't contain x , so y lies in the interior of a Weyl alcove and thus G_y is a parahoric subgroup.

Let K_0 be the compact open subgroup of $G(\mathbb{A}_{\mathbb{F}_q(t)})$ defined as the product of G_x at the place 0, G_y at the place ∞ , and the standard hyperspecial maximal compact G_0 at each other place. I claim the natural map

$$K_0 \cap G(F) \rightarrow G_x \rightarrow G_x/G_{x,0+}$$

is injective with image a Borel subgroup of $G_x/G_{x,0+}$.

To check this, we apply Lemma 4.1. First, we choose a faithful representation $\rho: G \rightarrow \text{Aut}(V)$ of G on a vector space V , and a basis e_1, \dots, e_n of V , where e_i is an eigenvector of T with eigenvalue a cocharacter λ_i , all defined over \mathbb{F}_q .

We now observe using the product formula and Lemma 4.1 that for $g \in K_0 \cap G(F)$, if some coordinate $\rho_{ij}(g) \neq 0$ then

$$(6.3) \quad 0 = \sum_v v(\rho_{ij}(g)) \geq \sum_v [f(v) \cdot (\lambda_i - \lambda_j)] = [x \cdot (\lambda_i - \lambda_j)] + [y \cdot (\lambda_i - \lambda_j)].$$

Since y is a small perturbation of $-x$, (6.3) is only satisfied if $x \cdot (\lambda_i - \lambda_j)$ is an integer and $y \cdot (\lambda_i - \lambda_j) \leq -x \cdot (\lambda_i - \lambda_j)$.

Let $a: \mathbb{G}_m \rightarrow T$ be a cocharacter which is a positive multiple of $x + y$. Then for any i, j with $\rho_{ij}(g) \neq 0$, we have $(x + y) \cdot (\lambda_i - \lambda_j) \leq 0$ and thus $a \cdot (\lambda_i - \lambda_j) \leq 0$. It follows that $a(\alpha)ga(\alpha)^{-1}$ is bounded as $\alpha \rightarrow 0$, since all the nonvanishing coordinates of ρ_{ij} of g are eigenvectors of conjugation by $a(\alpha)$ with eigenvalue a positive power of α .

The image of a in $G_{0,x}/G_{0,x+}$ defines a cocharacter of the maximal torus of $G_{0,x}/G_{0,x+}$, and the image of g in $G_{0,x}/G_{0,x+}$ remains bounded under conjugation by that character. Hence the image of g in $G_{0,x}/G_{0,x+}$ lies in the parabolic P_a associated to that cocharacter (under the ‘‘dynamical’’ definition of parabolic subgroups. Because a lies in the interior of a Weyl chamber, this is a Borel subgroup, so the image of $K_0 \cap G(F)$ in $G_{0,x}/G_{0,x+}$ lies in a Borel subgroup, as desired.

Furthermore, if $\rho_{ij}(g) \neq 0$, then because $x \cdot (\lambda_i - \lambda_j)$ is an integer and $y \cdot (\lambda_i - \lambda_j) \leq -x \cdot (\lambda_i - \lambda_j)$, the right side of (6.3) is 0, so every inequality is sharp and thus the valuation of $\rho_{ij}(g)$ at 0 is equal to $x \cdot (\lambda_i - \lambda_j)$.

On the other hand, by the second part of Lemma 4.1, if $g \in G_{0,x+}$, then the valuation of $\rho_{ij}(g)$ is strictly greater than $x \cdot (\lambda_i - \lambda_j)$ if $(\lambda_i - \lambda_j) \neq 0$, and the valuation of $(\rho_{ij}(g) - \delta_{ij})$ is > 0 if $(\lambda_i - \lambda_j) = 0$. Thus, if $i \neq j$, then we obtain a contradiction from the assumption $\rho_{ij}(g) \neq 0$, so $\rho_{ij}(g) = 0$, and if $i = j$, then $\rho_{ij}(g)$ has valuation 0 at every place, and thus lies in \mathbb{F}_q , but its difference from 1 has positive valuation at 0, so it is equal to 1.

Thus, if $g \in G_{0,x+}$, then $\rho(g)$ is the identity, so because ρ is faithful, g is the identity.

Let B be the Borel subgroup of $G_x/G_{x,0+}$ defined as the image of K_0 . Let H be the image of J in $G_x/G_{x,0+}$. If J is sufficiently small, then by definition there exists g in $G_x/G_{x,0+}$ with $J \cap gBg^{-1}$ trivial. By conjugating J by a lift of g , we may assume $g = 1$.

Let K_1 be the compact open subgroup of $G(\mathbb{A}_{\mathbb{F}_q(t)})$ defined as the product of J at the place 0, G_y at the place ∞ , and the standard hyperspecial maximal compact G_0 at each other place. Then the natural map $K_1 \cap G(\mathbb{F}_q(t)) \rightarrow G_x/G_{x,0+}$ is injective with image contained in $H \cap B$, hence with trivial image, so $K_1 \cap G(\mathbb{F}_q(t))$ is trivial.

The quotient $G_y/G_{y,0+}$ is a torus of rank r , hence has at least $(q-1)^r$ points, and thus has a non-trivial character χ because $q > 2$. We can view ρ and $\rho \otimes \chi$ as representations of K_1 by projection.

Let f_1 be a function on $G(\mathbb{F}_q(t)) \backslash G(\mathbb{A}_{\mathbb{F}_q(t)})$ created by applying the Poincaré series construction to a matrix coefficient of ρ , and let f_2 be a function on $G(\mathbb{F}_q(t))$ created by applying the Poincaré series construction to $\rho \otimes \chi$.

Both f_1 and f_2 must be a sum of generalized eigenforms of the excursion operators. Let \mathcal{L}_1 and \mathcal{L}_2 be the respective Lafforgue-Langlands parameters of these eigenforms, each a representation $\text{Gal}(\mathbb{F}_q(t)) \rightarrow G(\overline{\mathbb{Q}_\ell})$.

Because f_1 and f_2 are invariant under the hyperspecial maximal compact subgroup at each place other than 0 and ∞ , the associated parameters \mathcal{L}_1 and \mathcal{L}_2 are unramified away from 0 and ∞ . Because f_1 and f_2 transform under the parahoric subgroup G_y at ∞ by the trivial character and the character χ respectively, each is a sum of automorphic forms that lie in local representations that contain a G_y -equivariant vector. Such representations are subquotients of parabolic inductions from the Borel of characters of the maximal torus $T(\mathbb{F}_q((t)))$ whose restrictions to $T(\mathbb{F}_q)$ are 1 and χ respectively. Thus their associated Genestier-Lafforgue parameters are tame, and their restrictions to the tame inertia group at ∞ correspond under class field theory to 1 and χ , up to semisimplification. In particular, the restriction of \mathcal{L}_1 to the tame inertia group at ∞ is trivial (up to semisimplification), while the restriction of \mathcal{L}_2 to the tame inertia group at infinity is nontrivial.

Finally, because f_1 and f_2 transform under the representation π at 0, the parameters \mathcal{L}_1 and \mathcal{L}_2 at 0 are isomorphic to the Genestier-Lafforgue parameter of ρ , up to semisimplification. Now suppose the Genestier-Lafforgue parameter of ρ is tame. Since wildness is preserved by semisimplification, it follows that \mathcal{L}_1 and \mathcal{L}_2 are unramified away from 0 and ∞ and tame at ∞ . Thus \mathcal{L}_1 and \mathcal{L}_2 factor through the tame fundamental group of $\mathbb{G}_{m,\mathbb{F}_q}$. In the tame fundamental group of $\mathbb{G}_{m,\mathbb{F}_q}$, the inertia subgroups at 0 and ∞ are the same subgroup, both being equal to the tame geometric fundamental group. Since \mathcal{L}_1 and \mathcal{L}_2 are isomorphic when restricted to the tame inertia group at 0 and semisimplified, they are also isomorphic when restricted to the tame inertia group at ∞ and semisimplified. This contradicts our earlier claim that \mathcal{L}_1 is trivial while \mathcal{L}_2 is nontrivial when restricted to the tame

inertia group at ∞ and then semisimplified. This gives a contradiction, so indeed the Genestier-Lafforgue parameter of π is wildly ramified. \square

When the group J is contained in the principal congruence subgroup of a hyperspecial maximal compact, there is a shorter argument.

Proposition 6.4. *Under the hypotheses of Proposition 6.2, suppose G_x is a hyperspecial maximal compact and J is contained its principal congruence subgroup $G_{x,+}$. Let π be a pure supercuspidal representation of $G(\mathbb{F}_q)$ compactly induced from an irreducible representation ρ of J . Then the Genestier-Lafforgue parameter of π is wildly ramified.*

Proof. We construct a Poincaré series as in Theorem 5.2 using a function φ which at ∞ is a matrix coefficient of ρ and is a characteristic function of a hyperspecial maximal compact U_v at all other places v . Now the support of φ_∞ is contained in J , and $J \times \otimes_{v \neq \infty} U_v \cap G(K) = \{1\}$. Thus the Poincaré series P_φ does not vanish at 1, and the space of cusp forms it generates contains a cuspidal automorphic representation Π that is unramified except at ∞ . So $\mathcal{L}(\Pi)$ is a parameter with values in \hat{G} that is unramified away from ∞ . On the other hand, by Theorem 3.6 $\mathcal{L}^{ss}(\pi)$ is ramified. Since a tamely ramified local system on \mathbb{A}^1 is unramified, it follows that the ramification at ∞ must be wild. \square

7. PURE WEIL-DELIGNE PARAMETERS

7.1. Tempered parameters and pure Weil-Deligne parameters. Let F be a non-archimedean local field. A Weil-Deligne parameter for F with values in ${}^L G(C)$ is a pair (ρ, N) where $\rho : W_F \rightarrow {}^L G(C)$ is a continuous homomorphism (for the discrete topology on C) and $N \in \text{Lie}(G)(C)$ is a unipotent element such that $(ad \circ \rho, N)$ is a homomorphism from the Weil-Deligne group of F to $\text{Aut}(\text{Lie}(G))(C)$.

Without loss of generality we may assume $C = \mathbb{C}$. There is a simple recipe for converting a Weil-Deligne parameter (ρ, N) for F with values in ${}^L G(\mathbb{C})$ to a homomorphism

$$\varphi = \varphi_{\rho, N} : W_F \times SL(2, \mathbb{C}) \rightarrow {}^L G(\mathbb{C}),$$

which is algebraic (over \mathbb{R}) in the second factor, and vice versa. Suppose $G = GL(d, F)$, so ${}^L G(\mathbb{C}) = GL(d, \mathbb{C}) \times W_F$ and $\rho : W_F \rightarrow GL(V)$ is a representation of W_F on a d -dimensional complex vector space V . The map from $W_F \times SL(2, \mathbb{C})$ to the second factor W_F of ${}^L G(\mathbb{C})$ is just the projection to W_F and we ignore it in what follows. Write $V = \oplus_i V_i$ as a sum of Jordan blocks for N , with $N_i = N|_{V_i}$, so that each N_i is of order $d_i = \dim V_i - 1$. The restriction of φ to $\{1\} \times SL(2, \mathbb{C})$ has image in $\prod_i GL(V_i)$ and induces the irreducible d_i -dimensional representation ψ_i of $SL(2, \mathbb{C})$ on V_i , in such a way that $u_i = \exp(N_i)$ is identified with $\psi_i\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right)$ on V_i . Since $\rho(I_F) \subset \rho(W_F)$

commutes with N it stabilizes each V_i and $\varphi|_{I_F} = \rho|_{I_F}$. Finally, choose a Frobenius element $Frob_F \in W_F$ so that $\rho(Frob_F) = s$ satisfies $su_i s^{-1} = u_i^q$, and define

$$\varphi(Frob_F \times 1) := \rho(Frob_F) \times \psi_i \left(\begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right).$$

The equivalence class of φ is then well-defined.

For general G , we choose a faithful representation $\sigma : {}^L G(\mathbb{C}) \hookrightarrow GL(d, \mathbb{C})$ for some d . Then $\varphi_{\sigma \circ \rho, \sigma \circ N}$ factors (uniquely) through a homomorphism $\varphi_{\rho, N} : W_F \times SL(2, \mathbb{C}) \rightarrow {}^L G(\mathbb{C})$.

Definition 7.1. *The Weil-Deligne parameter (ρ, N) is tempered if the restriction of $\varphi_{\rho, N}$ to $W_F \times SU(2)$ has bounded image after projection to $\hat{G}(\mathbb{C})$. The parameter (ρ, N) is essentially tempered if its image in \hat{G}^{ad} is bounded.*

Definition 7.2. (a) *The Weil-Deligne representation (ρ, N) with values in $GL(V)$ is pure if there is a complex number t such that*

- (i) *The eigenvalues of $\rho_t(Frob_F) := \rho(Frob_F)q^t$ are all q -numbers of integer weight.*
- (ii) *The subspace $W_a V \subset V$ of eigenvectors for $\rho_t(Frob_F)$ with eigenvalues of weight $\leq a$ is invariant under (ρ, N) ;*
- (iii) *Letting $gr_a V = W_a V / W_{a-1} V$, there is an integer w such that, for all $i \geq 0$, the map*

$$N : gr_{w-i} V \rightarrow gr_{w+i} V$$

is an isomorphism.

The integer w in (iii) is then called the weight of (ρ, N) (twisted by t).

(b) *Let G be a connected reductive group over F . The Weil-Deligne parameter (ρ, N) with values in ${}^L G(\mathbb{C})$ is pure if $(\sigma \circ \rho, N)$ is pure of some weight for some (equivalently every) faithful representation σ of ${}^L G$.*

In particular, if $N = 0$ then the Weil-Deligne parameter (ρ, N) is pure if and only if the Weil group parameter ρ is pure (in the usual sense) of some weight, (up to twist by a power of the norm). We distinguish the two notions of purity by referring to “pure Weil-Deligne parameters” and “pure Weil group parameters”, or “pure semi-simple parameters,” respectively. Purity in previous sections referred exclusively to the latter notion. We have tried to determine whether there is a convention in the literature to designate the Weil-Deligne parameters (ρ, N) that satisfy the conditions of Definition 7.2 when ρ is not itself pure. The article [TY] calls such parameters pure but then uses the expression *strictly pure*, which does not seem to have caught on, for what we are calling pure Weil group parameters.

The following facts are well-known and in any case are easy to verify.

Lemma 7.3. (a) *The Weil-Deligne parameter (ρ, N) is pure if and only if it is essentially tempered.*

(b) Let $\rho : W_F \rightarrow {}^L G(C)$ be a homomorphism. There is at most one way to complete ρ to a pure Weil-Deligne parameter (ρ, N) . If it exists, we call it the pure completion of ρ .

7.2. An application of the purity of the monodromy weight filtration. The parametrization of Genestier-Lafforgue only attaches a semisimple parameter $\mathcal{L}^{ss}(\pi)$ to a local representation π . But when π is realized as a local component of a cuspidal automorphic representation Π of G over some global function field K – say $\pi = \Pi_x$ for some place x of K – then the restriction $\mathcal{L}^{ss}(\Pi)_x$ to the decomposition group Γ_x at x need not be semisimple. We let $\mathcal{L}_\Pi(\pi) = (\mathcal{L}^{ss}(\pi), N_\Pi)$ denote the Weil-Deligne parameter associated to the Galois parameter $\mathcal{L}^{ss}(\Pi)_x$.

The following result of Sawin and Templier generalizes Deligne’s theorem on the purity of the monodromy weight filtration to G -local systems:

Lemma 7.4. [ST, Lemma 11.4] *Let X be a smooth projective curve over a finite field k , with function field $K = k(X)$, and let $\rho : \text{Gal}(K^{sep}/K) \rightarrow {}^L G(C)$ be an irreducible homomorphism (with image not contained in a proper parabolic subgroup). Let z be any place of K , and let (ρ_z^{ss}, N_z) be the Weil-Deligne parameter associated to the restriction ρ_z of ρ to a decomposition group at z . Then (ρ_z^{ss}, N_z) is a pure Weil-Deligne parameter.*

Remark 7.5. *The lemma is stated in [ST] when G is split semisimple but the proof is valid for any reductive G , with the appropriate extension of the property of being contained in a proper parabolic subgroup to disconnected ${}^L G(C)$. We leave it to the reader to formulate the definition.*

Corollary 7.6. *Let π be an irreducible admissible representation of $G(F)$. Suppose X is a smooth projective curve over a finite field k and z is a place of the function field $K = k(X)$ such that $K_z \xrightarrow{\sim} F$. Suppose there is an automorphic representation Π of G_K , with $\Pi_z \xrightarrow{\sim} \pi$, such that the global parameter $\mathcal{L}^{ss}(\Pi)$ is irreducible.*

Then the Weil-Deligne parameter $\mathcal{L}_\Pi(\pi)$ is pure. Moreover, it is the pure completion of the semisimple parameter $\mathcal{L}^{ss}(\pi)$.

7.3. Existence of pure completions of local parameters of supercuspidal representations. Not every semisimple parameter admits a completion to a pure Weil-Deligne parameter. Corollary 7.6 asserts that $\mathcal{L}^{ss}(\pi)$ does admit a pure completion provided π can be realized as a local component of an automorphic representation whose global parameter is irreducible in the indicated sense. In this section we construct such realizations when π is supercuspidal. This is consistent with the expectation that the full Weil-Deligne parameter that the local Langlands correspondence attaches to any supercuspidal is pure, and that this holds as well for unipotent supercuspidals.

Proposition 7.7. *Let G be a reductive group over F . If G is of type B_n or C_n assume $p > 2$; if G is of type G_2 or F_4 assume $p > 3$ (in other words,*

we assume p is a good prime for these groups). Let π be a supercuspidal representation of $G(F)$. Let X be a smooth projective curve over a finite field k and let z be a place of the function field $K = k(X)$ such that $K_z \xrightarrow{\sim} F$. Then there exist infinitely many cuspidal automorphic representations Π of G_K , with $\Pi_z \xrightarrow{\sim} \pi$, such that the global parameter $\mathcal{L}^{ss}(\Pi)$ is irreducible.

In particular, the semisimple parameter $\mathcal{L}^{ss}(\pi)$ admits a completion to a pure Weil-Deligne parameter that is realized as the specialization at x of the global parameter $\mathcal{L}^{ss}(\Pi)$, for any such Π .

Proof. Let $x \neq z$ be any other place of K . Since there are infinitely many such x , it suffices to show

- (i) There is a supercuspidal representation π' of $G(K_x)$ such that $\mathcal{L}^{ss}(\pi')$ is irreducible, and
- (ii) There is an automorphic representation Π of G_K with $\Pi_x \xrightarrow{\sim} \pi'$ and $\Pi_z \xrightarrow{\sim} \pi$, and with Π_y tamely ramified for all $y \neq x, z$.

The existence of Π as in (ii) is proved in [GLo]. It remains to show (i). We let $k' = k(x)$ denote the residue field at x and construct the Kloosterman sheaf $\mathrm{Kl}_{\hat{G}}(\phi, \chi)$ of [HNY] over $\mathbb{P}_{k'}^1$. Let π' be the local component at ∞ of the corresponding automorphic representation $\pi(\phi, \chi)$. Under the hypotheses on G and p , [HNY, Theorem 2] implies that the restriction to the inertia group at ∞ of the local monodromy representation of $\mathrm{Kl}_{\hat{G}}(\phi, \chi)$ is irreducible. On the other hand, it follows from construction that $\mathrm{Kl}_{\hat{G}}(\phi, \chi)$ is the global parameter attached by [Laf18] to $\pi(\phi, \chi)$. More precisely, [HNY] computes the local monodromy of $\mathrm{Kl}_{\hat{G}}(\phi, \chi)$ at unramified places and identifies it with that obtained from the Satake parameters of $\pi(\phi, \chi)$. (The construction of $\pi(\phi, \chi)$ is made explicit in [Y16] but its existence is implicit in [HNY].) The claim for the local parameter at ∞ then follows by Chebotarev density. \square

The idea behind Corollary 7.6 and Proposition 7.7 was already applied in the case of classical groups in the proof of [GLo, Proposition 7.3]. Instead of using the Kloosterman representations of [HNY] to establish irreducibility, [GLo] used depth zero supercuspidals, but introduced a hypothesis because the irreducibility of the corresponding parameter has not (yet) been established for local fields of positive characteristic.

7.4. Generalization to discrete series representations. Since discrete series representations are essentially tempered, Proposition 7.7 is expected to hold when π is any discrete series representation, not necessarily supercuspidal. To establish this we can replace the compactly supported matrix coefficient that we used for supercuspidal π by a pseudocoefficient f_π of the discrete series π , whose properties are recalled in [GLo, §8.1]. This has the obvious disadvantage that the Poincaré series can no longer be shown to converge absolutely. If we assume that the stable Arthur trace formula is valid over function fields, we can still prove the analogue of Proposition 7.7 for discrete series representations, though the proof is less direct. Gan and

Lomelí formulate a version of the trace formula that is adequate for these purposes as Working Hypothesis II in §8.2 of their paper. We then have the following generalization of Proposition 7.7.

Proposition 7.8. *Assume $G(F)$ is split semisimple. If G is of type B_n or C_n assume $p > 2$; if G is of type G_2 or F_4 assume $p > 3$ (in other words, we assume p is a good prime for these groups). Assume Working Hypothesis II of [GLo, §8.2] Let π be a discrete series representation of $G(F)$. Let X be a smooth projective curve over a finite field k and let z be a place of the function field $K = k(X)$ such that $K_z \xrightarrow{\sim} F$. Then there exist infinitely many cuspidal automorphic representations Π of G_K , with $\Pi_z \xrightarrow{\sim} \pi$, such that the global parameter $\mathcal{L}^{ss}(\Pi)$ is irreducible.*

In particular, the semisimple parameter $\mathcal{L}^{ss}(\pi)$ admits a pure completion that is realized as the specialization at x of the global parameter $\mathcal{L}^{ss}(\Pi)$, for any such Π .

Proof. As in the proof of Proposition 7.7, it would be enough to show the existence of Π with $\Pi_x \xrightarrow{\sim} \pi'$ and $\Pi_z \xrightarrow{\sim} \pi$, and with Π_y tamely ramified for all $y \neq x, z$. Using pseudocoefficients we cannot prove this, however. What we can show, as in the proof of [GLo, Proposition 8.2], is the existence of a Π with $\Pi_x \xrightarrow{\sim} \pi'$ and $Tr(\Pi_z)(f_\pi) \neq 0$, where f_π is a pseudocoefficient of π , as above. This implies (by [GLo, Lemma 8.1]) that Π_z has the same cuspidal support as π , and therefore that

$$\mathcal{L}^{ss}(\Pi_z) = \mathcal{L}^{ss}(\pi)$$

since the semisimple Genestier-Lafforgue parameter is compatible with parabolic induction. On the other hand, the proof of Proposition 7.7 shows that $\mathcal{L}^{ss}(\Pi_z)$ admits a pure completion, and this is enough to complete the proof. \square

8. BASE CHANGE AND INCORRIGIBLE REPRESENTATIONS

Let $E = k((t))$ for a finite field k and let G be a connected reductive group over E . Let $\mathcal{T}(G, F) \subset \mathcal{A}(G, F)$ denote the set of irreducible tempered (admissible) representations of $G(F)$ with coefficients in C , and let $P(\mathcal{T}(G, F))$ be the set of non-empty subsets of $\mathcal{T}(G, F)$. We write \mathcal{L}_F^{ss} for the semisimple Langlands parametrization of (3.2) for $\mathcal{A}(G, F)$, with a subscript to indicate the base field. Say G admits tempered base change if, for every pair $F \subset F'$ of finite extensions of E , with F'/F a cyclic extension of prime order, there is a map

$$BC_{F'/F} : \mathcal{T}(G, F) \rightarrow P(\mathcal{T}(G, F'))$$

so that for any $\pi \in \mathcal{T}(G, F)$ and any $\pi' \in BC_{F'/F}(\pi)$,

$$(8.1) \quad \mathcal{L}_{F'}^{ss}(\pi') = \mathcal{L}_F^{ss}(\pi) |_{W_{F'}}.$$

Of course it is expected that every G admits tempered base change and that the set $BC(\pi)$ is characterized intrinsically inside $\mathcal{A}(G, F)$ in terms of π by relations of distribution characters. It is also assumed that $BC(\pi)$ is always

a finite set, but we do not make that assumption. There is a brief discussion of the existence of tempered base change at the end of this section.

Definition 8.2. *Suppose G admits tempered base change. The supercuspidal representation π of $G(F)$ is **in corrigible** if for any sequence*

$$(8.3) \quad F = F_0 \subset F_1 \subset \cdots \subset F_r$$

of extensions, where F_i/F_{i-1} is cyclic of prime order for all $i \geq 1$, the set

$$BC_{F_r/F}(\pi) := BC_{F_r/F_{r-1}}(BC_{F_{r-1}/F_{r-2}} \cdots (BC_{F_1/F}(\pi) \cdots))$$

contains a supercuspidal member.

Proposition 8.4. *Assume G admits tempered base change and $p \nmid |W|$. Then for any extension F/E , no pure supercuspidal representation of $G(F)$ is incorrigible. More generally, suppose G has the property that, for every Levi subgroup $L(F)$ of $G(F)$, every supercuspidal representation of $L(F)$ contains an \mathfrak{s} -type. Then no pure supercuspidal representation of $G(F)$ is incorrigible.*

Proof. Let π be a pure supercuspidal representation of $G(F)$. The image of the inertia group under $\mathcal{L}_F^{ss}(\pi)$ is finite, so there is a sequence of cyclic extensions of prime order, as in (8.3), such that, if π_r is any member of $BC_{F_r/F}(\pi)$, we have

$$\mathcal{L}_{F_r}^{ss}(\pi_r) = \mathcal{L}_F^{ss}(\pi) |_{W_{F_r}}$$

is unramified. It then follows from Corollary 5.5 that π_r is a constituent of an unramified principal series representation. In particular, no member of $BC_{F_r/F}(\pi)$ can be supercuspidal. \square

8.1. Existence of tempered base change. Suppose for the moment that F is a p -adic field. A procedure for defining cyclic base change can be constructed using the methods of [Lab99]. Suppose first that π is supercuspidal with central character of finite order. Then there is

- (i) A totally real number field K with a local place w such that $K_w \xrightarrow{\sim} F$;
- (ii) A totally real cyclic extension K'/K with $K'_w = K' \otimes_K F \xrightarrow{\sim} F'$,
- (iii) A connected reductive group \mathcal{G} over K , with $\mathcal{G}(K_w) \xrightarrow{\sim} G(F)$ and
- (iv) $\mathcal{G}(K_\sigma)$ compact modulo center for all archimedean places σ of K ,
- (v) And an automorphic representation Π such that $\Pi_w \xrightarrow{\sim} \pi$ and
- (vi) Π_{v_i} isomorphic to a Steinberg representation at any chosen finite set of places v_i .

Then it is proved in [Lab99], using the stable twisted trace formula, that there is a non-empty collection $\{\Pi'\}$ of automorphic representations of $\mathcal{G}_{K'}$ such that, at every place u at which Π and K'/K are unramified, Π'_u is the unramified base change of Π_u . We can then define $BC(\pi)$ to be the collection of Π'_w for all such K, K', \mathcal{G}, Π . The stable twisted characters of such collections of Π'_w are related to the stable character of π but in the absence of a canonical notion of L -packet we omit the precise statement.

When Labesse defined his construction in [Lab99], the stabilization of the trace formula had not yet been established in either the twisted or the untwisted setting. Arthur reduced the stabilization to the fundamental lemma in a series of papers shortly thereafter, and a few years later the main step in the proof of the fundamental lemma was completed by Ngô. The stabilization of the twisted trace formula is contained in [MW]; see also [1] for references for the untwisted case. Labesse needed hypothesis (vi) in the above list in order to work with a simplified version of the trace formula. In principle (vi) is no longer necessary, but the necessary consequences of the full stabilized trace formula have not yet been established (see [1, §CHL.IV.B] for an example of the kind of work required). Thus for practical purposes, the construction in [Lab99] still provides the most complete definition of local supercuspidal base change in general.

Since we are only looking for semisimple Langlands parameters, we can reduce a general tempered π to the supercuspidal case by means of parabolic induction. Unfortunately, although Fargues and Scholze have defined a semisimple parametrization of all irreducible admissible representations of $G(F)$, when F is a p -adic field, there is no way to relate the parameter attached to π to that attached to the elements of the set $BC(\pi)$ defined by Labesse's construction, unless we know how to attach Galois parameters to the globalizations Π as in (v). If we could do that, then the relation between the parameters of Π_u and Π'_u at unramified places would suffice, by Chebotarev density, to establish the relation (8.1) at the place of interest. This reasoning has in fact been applied for most classical groups, and for G_2 , but it is not available in general.

Now suppose $F = k((t))$ as above. We can certainly globalize π as in the number field case, and then [Laf18] supplies the necessary global parametrization that is missing for number fields. However, the stable twisted trace formula is lacking for function fields, so for most G tempered base change is not available. It should nevertheless be (relatively) straightforward, although time-consuming, for specialists to prove the necessary statements, starting with the construction of the non-invariant trace formula by Labesse and Lemaire [LL], when p is large relative to the group G .

8.2. The case of $GL(n)$. We again assume F is a local field of positive characteristic. Let $G = GL(n)$. It has been proved by Henniart and Lemaire in [HeLe] that G admits tempered base change; they even prove that $BC_{F'/F}(\pi)$ is a singleton for any π . Moreover, every supercuspidal representation of every Levi subgroup of $G(F)$ contains an \mathfrak{s} -type [BK]. Thus the above discussion applies. Before drawing the relevant conclusion, we make the following observation.

The following lemma may be derived from the work of Bushnell and Kutzko.

Lemma 8.5. *Let π be a supercuspidal representation of $GL(n, F)$ for F a local field. One of the following is true:*

(a) *There exists a principal hereditary order $\mathfrak{A} \subset M_n(F)$, a field $E \subset M_n(F)$ which is an extension of F of degree > 1 , such that E^\times normalizes \mathfrak{A} , and, writing \mathfrak{B} for the centralizer of E in \mathfrak{A} and \mathfrak{P} for the maximal ideal of \mathfrak{A} , a representation Λ of $E^\times \cdot (\mathfrak{A}^\times \cap (\mathfrak{B} + \mathfrak{P}))$ such that π is the induced representation of Λ .*

(b) *There exists a representation Λ of $F^\times GL_n(\mathcal{O}_F)$ such that π is the induced representation of Λ .*

Proof. By [BK, Theorem 8.4.1], there exists a simple type (J, λ) in G such that $\pi \upharpoonright J$ contains λ . The definition [BK, Definition 5.5.10] of simple type splits into two cases, (a) and (b).

The case (b) will be easier to handle – so much so that it is not even necessary to give the definition of this case. Instead, we note that in this case, by [BK, Theorem 8.4.1(iii)], there exists a representation Λ of $F^\times \mathbf{U}(\mathfrak{A})$, where $\mathbf{U}(\mathfrak{A})$ is the unit group of a hereditary order \mathfrak{A} , such that $\pi = \text{Ind}(\Lambda)$. Because \mathfrak{A} is a hereditary order, $\mathbf{U}(\mathfrak{A})$ is a compact open subgroup of $GL(n, F)$ and thus, up to conjugation, is contained in the maximal compact $GL_n(\mathcal{O}_F)$. By first inducing Λ from $F^\times \mathbf{U}(\mathfrak{A})$ to $F^\times GL_n(\mathcal{O}_F)$, we obtain case (b) above.

Now we turn to case (a). Let A be the algebra of $n \times n$ matrices over F . In this case, \mathfrak{A} is a principal \mathcal{O}_F -order in A and $[\mathfrak{A}, n, 0, \beta]$ is a simple stratum. Let us unpack this definition. The order \mathfrak{A} is a compact open \mathcal{O}_F -subalgebra of A . We write \mathfrak{P} for the Jacobson radical of \mathfrak{A} . We let β be an element of A and take n to be a positive integer. We define $E = F[\beta]$, \mathfrak{B} the centralizer of E in \mathfrak{A} , and \mathfrak{P} the maximal ideal of \mathfrak{A} . We set [BK, Definitions 1.4.3 and 1.4.5]

$$k_0(\beta, \mathfrak{A}) = \max\{k \in \mathbb{Z} \mid \{x \in \mathfrak{A} \mid \beta x - x\beta \in \mathfrak{P}^k\} \not\subseteq \mathfrak{B} + \mathfrak{P}\}.$$

Then the assumption that $[\mathfrak{A}, n, 0, \beta]$ is a simple stratum means that \mathfrak{A} is hereditary, E^\times normalizes \mathfrak{A} , $v_{\mathfrak{A}}(\beta) = -n$, and $0 < -k_0(\beta, \mathfrak{A})$ [BK, Definition 1.5.5].

In this context, [BK, Definition 3.1.8] defines an algebra $\mathfrak{J}(\beta)$ by an inductive procedure. For us, the only relevant feature of this definition is that it is contained in $\mathfrak{B} + \beta^{\lfloor \frac{r+1}{2} \rfloor}$ where $r = -k_0(\beta, \mathfrak{A}) > 0$ ([BK, 3.1.6]) and thus $\mathfrak{J} \subseteq \mathfrak{B} + \mathfrak{P}$. We can next define [BK, (3.1.14)] a group $J(\beta, \mathfrak{A})$ as the intersection of \mathfrak{J} with the group of units $\mathbf{U}^0(\mathfrak{A}) = \mathfrak{A}^\times$.

We can finally state the conclusion of [BK, Theorem 8.4.1(ii)] in this case, which is that there is a representation Λ of $E^\times J(\beta, \mathfrak{A})$ such that $\pi = \text{Ind}(\Lambda)$. We will use only this structure, and will not concern ourselves with the exact construction of Λ . If $E = F$ then, because $J(\beta, \mathfrak{A})$ is contained in a maximal compact subgroup, we are again in case (b), so we may assume $E \neq F$.

Since $J(\beta, \mathfrak{A})$ is contained in $\mathfrak{A}^\times \cap (\mathfrak{B} + \mathfrak{P})$, we can induce Λ from $E^\times J(\beta, \mathfrak{A})$ to $E^\times \cdot (\mathfrak{A}^\times \cap (\mathfrak{B} + \mathfrak{P}))$, and we are in case (a) above. \square

Lemma 8.6. *If π satisfies case (b) of Lemma 8.5, then π is pure and ramified.*

Proof. The determinants of elements of $GL_n(\mathcal{O}_F)$ must have zero valuation, and because F^\times consists of scalars, the determinants of its elements must have valuation divisible by n . It follows that, if α is the character of $GL(n, F)$ defined by $\alpha(g) = e^{\frac{2\pi i}{n}v(\det(g))}$ then

$$\pi \otimes \alpha = \text{Ind}(\Lambda) \otimes \alpha = \text{Ind}(\Lambda \otimes \alpha) = \text{Ind}(\Lambda) = \pi.$$

By [GLa, Remarque 0.2], it follows that the Genestier-Lafforgue parameter σ_π is stable under tensor-product with the one-dimensional unramified representation of W_F sending Frobenius to $e^{2\pi i/n}$. Let a be a matrix representing that isomorphism, then $\sigma_\pi(\text{Frob}_q) \circ a = e^{2\pi i/n} a \circ \sigma_\pi(\text{Frob}_q)$ so a is conjugate to $e^{2\pi i/n} a$. Thus the set of eigenvalues of a is stable under multiplication by $e^{2\pi i/n}$, so a has n distinct eigenvalues, and thus a has a one-dimensional eigenspace. This one-dimensional eigenspace is a one-dimensional representation χ of the subgroup of W_F corresponding to the degree n unramified extension F_n of F , which, because Frobenius permutes the eigenspaces, gives an isomorphism $\text{Ind}_{W_{F_n}}^{W_F} \chi \cong \sigma_\pi$. Since χ is a one-dimensional representation, it is automatically pure of some weight (possibly non-integral), so σ_π is pure. By our previous result, σ_π is ramified. \square

We now assume given π in case (a) of Lemma 8.5, and prepare for the construction of a Poincaré series by choosing compact subgroups at each place.

It follows from an elementary calculation or [BK, p. 76 and p. 183] that a principal hereditary order \mathfrak{A} , after choosing a basis, can be put into the following form: We choose a divisor e of n and then we take all $n \times n$ matrices decomposed into $n/e \times n/e$ blocks of size $e \times e$, where all the blocks on the diagonal and above have entries in the ring of integers \mathcal{O}_F , and all the blocks below the diagonal have entries in the maximal ideal of \mathcal{O}_F . We fix for the remainder of this argument a basis where \mathfrak{A} has this form.

We can choose a finite field \mathbb{F}_q such that $F \cong \mathbb{F}_q((t))$ and fix such an isomorphism. This identifies F with the local field $\mathbb{F}_q(t)_0$ of $\mathbb{P}_{\mathbb{F}_q}^1$ at 0. We now define subgroups U_x of $GL_n(\mathbb{F}_q(t)_x)$ for each place x of $\mathbb{F}_q(t)$. For $x \neq 0, \infty$, we take U_x to be the standard maximal compact subgroup. We take U_∞ to consist of block-diagonal matrices in $GL_n(\mathbb{F}_q((t^{-1})))$ where all the blocks on the diagonal and below are in $M_e(\mathbb{F}_q[[t^{-1}]])$, all the blocks above the diagonal are in $M_e(t^{-1}\mathbb{F}_q[[t^{-1}]])$, and all the diagonal blocks, modulo t^{-1} are in n/e fixed Borel subgroups of GL_n , to be chosen in Lemma 8.7. We take U_0 to be $E^\times(\beta, \mathfrak{A})$.

Let $S = GL_n(\mathbb{F}_q(t)) \cap \prod_x U_x$.

Lemma 8.7. *For a suitable choice of Borel subgroups, the group S is contained in a maximal torus T of $GL_n(\mathbb{F}_q)$, and furthermore, that the n -dimensional vector space, when viewed as a representation of S , is a sum of n/k characters each repeated k times for some $k > 1$ a divisor of n .*

Proof. First observe that the valuation of the determinant of any element of U_x for any $x \neq 0$ is 0. It follows that for $g \in S$, $\det g$ has valuation 0 at every place away from 0, thus by the product formula has valuation 0 at 0 as well. The valuation of the determinant of every element of \mathfrak{A}^\times is 0, so g must be the product of an element of $\mathfrak{A}^\times \cap (\mathfrak{B} + \mathfrak{P})$ with a unit $u \in E^\times$. Now every unit of E^\times is contained in \mathfrak{A}^\times by [BK, Proposition 1.2.1(i)]. Because $u \in E^\times$ and thus u commutes with β , we have $u \in \mathfrak{B}$. Thus

$$g \in u \cdot (\mathfrak{A}^\times \cap (\mathfrak{B} + \mathfrak{P})) = \mathfrak{A}^\times \cap (\mathfrak{B} + \mathfrak{P}).$$

In particular, g , at 0, is contained in \mathfrak{A} . Together with our assumptions at other places, we conclude that every matrix entry of g is integral at each place of $\mathbb{F}_q(t)$, hence in fact lies in \mathbb{F}_q , so $g \in GL_n(\mathbb{F}_q)$. Furthermore, by the integrality conditions at 0, we see that every $e \times e$ block of g below the diagonal vanishes, and by the integrality conditions at ∞ , every $e \times e$ block of g above the diagonal vanishes. So g is a block-diagonal matrix, i.e. $g \in (GL_e(\mathbb{F}_q))^{n/e}$. Furthermore, each of the diagonal matrices must be contained in some fixed Borel, to be chosen later.

We now use the fact that E/F is a nontrivial extension. Thus it either ramifies or contains an extension of the residue field.

We handle the case when E is ramified first. In this case, E contains an element x whose determinant does not have valuation divisible by n , and thus the action of x on \mathfrak{A} must permute the n/e diagonal blocks. The action on the blocks divides them into $n/(ek)$ orbits of size k for some $k > 1$ dividing n/e . Because elements of \mathfrak{B} commute with E , and thus in particular with x , their value mod \mathfrak{p}_F on one block in the orbit determines their value on every other block in the orbit. So, mod \mathfrak{P} , the elements of \mathfrak{B} lie in $M_e(\mathbb{F}_q)^{n/(ek)}$, and the units of \mathfrak{B} lie in $GL_e(\mathbb{F}_q)^{n/(ek)}$. To define U_∞ , we choose n/e Borel subgroups of $GL_e(\mathbb{F}_q)$ such that, in each orbit, at least one of the Borel subgroups is sent by x to a Borel in general position with respect to the next Borel in the orbit. It follows that each element of $S \subset GL_e(\mathbb{F}_q)^{n/(ek)}$ lies in the intersection of two Borels in general position, and thus lies in the maximal torus, which is contained in a maximal torus of GL_n . Furthermore, the n -dimensional representation of S is the restriction of the n -dimensional representation of $GL_e(\mathbb{F}_q)^{n/(ek)}$, which is the sum of $n/(ek)$ standard representations, each repeated k times, so as a representation of S it is the sum of n/k characters, each repeated k times, as desired.

We handle the case when E contains a residue field extension \mathbb{F}_{q^k} next. In this case, E contains an element x which generates the residue field extension. Because x is a unit, x lies in \mathfrak{A} . Restricting to each of the n/e diagonal blocks and modding out by \mathfrak{p}_F , the element x satisfies the characteristic polynomial of a generator of \mathbb{F}_{q^k} and thus generates a subfield

isomorphic to \mathbb{F}_{q^k} . Thus, the centralizer of x in each block is isomorphic to $GL_{e/k}(\mathbb{F}_{q^k})$, and every unit in $\mathfrak{B} + \mathfrak{P}$, restricted to each diagonal block and reduced modulo \mathfrak{p}_F , must lie in this $GL_{e/k}(\mathbb{F}_{q^k})$.

We choose a Borel as follows: Fix a \mathbb{F}_{q^k} -basis of the e -dimensional vector space $v_1, \dots, v_{e/k}$, and then extend the sequence

$$v_1, \dots, v_{e/k}, xv_{e/k}, xv_{e/k-1}, \dots, xv_1$$

arbitrarily to an \mathbb{F}_q -basis of this space. For each block, choose a Borel consisting of linear transformations which are upper-triangular with respect to this basis, and use these to define U_∞ . Then every $g \in S$, restricted to this block, will lie in the intersection of $GL_{e/k}(\mathbb{F}_{q^k})$ with this Borel. Hence g will be upper-triangular with respect to the \mathbb{F}_{q^k} -basis v_1, \dots, v_k and upper-triangular with respect to the \mathbb{F}_{q^k} -basis $xv_{e/k}, xv_{e/k-1}, \dots, xv_1$, thus \mathbb{F}_{q^k} -diagonal with respect to the basis v_1, \dots, v_k . Furthermore, g must act on v_1, \dots, v_k by multiplication by elements of \mathbb{F}_q , since otherwise it would not preserve the \mathbb{F}_q -subspace generated by v_1, \dots, v_k , so in fact g is contained in the diagonal of $GL_{e/k}(\mathbb{F}_q)$ inside $GL_{e/k}(\mathbb{F}_{q^k})$ inside $GL_e(\mathbb{F}_q)$. Passing from $GL_{e/k}(\mathbb{F}_q)$ to $GL_e(\mathbb{F}_q)$ in this way sends the maximal torus to a subset of the maximal torus, as desired, and expresses the e -dimensional standard representation of GL_e as a sum of k copies of the e/k -dimensional standard representation of $GL_{e/k}$, hence the sum of k copies of e/k characters of S , as desired. \square

Note that U_∞ is an Iwahori subgroup of $GL_n(\mathbb{F}_q((t^{-1})))$ and thus surjects onto $((\mathbb{F}_q)^\times)^n$.

Lemma 8.8. *If $q > 2$, any (one-dimensional) character of S may be extended to a nontrivial character of U_∞ .*

Furthermore, if $q > 3$, any character of S may be extended to two characters, χ_a, χ_b , both pulled back from $((\mathbb{F}_q)^\times)^n$, where χ_a is trivial on at least one copy of \mathbb{F}_q^\times and χ_b is trivial on no copies of \mathbb{F}_q^\times .

Proof. For each case, note that the subgroup U_∞ is an Iwahori subgroup of $GL_n(\mathbb{F}_q((t^{-1})))$, so its quotient by its maximal pro- p subgroup is $((\mathbb{F}_q)^\times)^n$. Since S is a subgroup of the maximal torus of $GL_n(\mathbb{F}_q)$ by Lemma 8.7, its order is prime to p , and so the quotient of U_∞ by its maximal pro- p subgroup is faithful. Thus χ_S is a character of the image of S inside $((\mathbb{F}_q)^\times)^n$.

For the first case, it suffices to extend χ_S^{-1} to a nontrivial character of $(\mathbb{F}_q^\times)^n$ and inflate to U_∞ . We can do this unless χ_S is trivial and $S = ((\mathbb{F}_q)^\times)^n$, which contradicts our claim that the characters of S are repeated unless \mathbb{F}_q^\times is trivial, which implies $q = 2$.

For the second case, if $q > 3$, note that since the standard representation of S is the sum of n/k characters repeated k times, the image of S inside $(\mathbb{F}_q^\times)^n$ is contained in $(\mathbb{F}_q^\times)^{n/k}$, repeated k times. It follows that when we extend χ_S^{-1} to a character χ of $(\mathbb{F}_q^\times)^n$, which can be viewed as a tuple of n characters χ_1, \dots, χ_n of \mathbb{F}_q , we can choose any tuple of characters as long

as the product of k characters in each of n/k orbits takes some fixed value depending on the orbit. So we can certainly choose $k - 1$ of the characters in each orbit to be trivial, and the last one to take the fixed value, to produce χ_a . To produce χ_b , we choose $k - 2$ of the characters in each orbit to be an arbitrary nontrivial character. The product of the two remaining characters is then determined. Since $q > 3$, there are more than two characters, and so we can choose the next character to be a character which is neither trivial nor the determined product. It follows that the last character is nontrivial, so χ_b indeed contains no trivial characters. \square

We are now ready to construct our Poincaré series. Since S is a finite abelian group, λ restricted to $S \subset E^\times J(\beta, \mathfrak{A})$ is a sum of one-dimensional characters. Let χ_S be one of these.

Proposition 8.9. *Assume the residue field of F has more than 2 elements. Then every supercuspidal representation of $GL(n, F)$ is ramified. Moreover, if the residue field of F has more than 3 elements, then every supercuspidal representation of $GL(n, F)$ is either ramified and pure or wildly ramified.*

Proof. Let $F = \mathbb{F}_q((t))$. Fix a supercuspidal representation π . We apply Lemma 8.5. In case (b), π is pure and ramified by Lemma 8.6, so we may reduce to case (a), where we will prove that π is ramified if $q > 2$ and wildly ramified if $q > 3$.

To do this, we construct a Poincaré series, using all the notation above. Since S is a finite abelian group, λ restricted to $S \subset E^\times J(\beta, \mathfrak{A})$ is a sum of one-dimensional characters. Let χ_S be one of these. We extend χ_S^{-1} to a nontrivial character χ of U_∞ using Lemma 8.8. We take the representation $\lambda \otimes \chi$ of $\prod_x U_x$, which is trivial by construction on $S = GL_n(F) \cap \prod_x U_x$. Using a matrix coefficient of this representation, we can construct a Poincaré series. After decomposing into Hecke eigenforms, this series generates an automorphic representation Π which is unramified outside of 0 and ∞ , of type $\text{Ind}(\lambda) = \pi$ at 0, and whose local factor at ∞ contains a vector that transforms according to χ under U_∞ . By Lafforgue, associated to this representation is a global parameter σ_Π which matches σ_π at 0, and is unramified away from 0 and ∞ .

Furthermore, at ∞ , the local factor of Π is contained in a parabolic induction from the Borel of a tamely ramified character of the maximal torus, which, restricted to the \mathbb{F}_q -points of the maximal torus, is χ . Thus, by the compatibility of Genestier-Lafforgue with parabolic induction and the local Langlands correspondence for GL_1 , the Genestier-Lafforgue parameter at ∞ is a sum of n tamely ramified characters whose restriction to the inertia group is trivial if and only if the restriction of χ to the corresponding copy of \mathbb{F}_q^\times is trivial. Thus the restriction of the global parameter σ_Π to ∞ is tamely ramified, and is unipotent if and only if χ is trivial.

If π is unramified, then σ_Π is unramified away from 0 and ∞ and has at worst unipotent ramification at 0. Because σ_Π is unramified away from 0

and ∞ and tamely ramified at 0 and ∞ , the restrictions of σ_Π to the inertia groups at 0 and ∞ are isomorphic, which contradicts the fact that σ_Π is unipotent at 0 and σ_Π is non-unipotent at ∞ , it is non-unipotent at 0. So in fact σ_π is ramified.

If $q > 3$ and π is tamely ramified, then we apply the second case of Lemma 8.8 to produce two characters, χ_a and χ_b , and follow the same procedure above to produce two globalizations Π_a, Π_b . Again $\sigma_{\Pi_a}, \sigma_{\Pi_b}$ are tame at 0 and ∞ , so the restrictions of σ_{Π_a} to the inertia subgroup at ∞ is isomorphic to its restriction to the inertia subgroup at 0 and thus isomorphic, up to semisimplification, to σ_π . The same is true for σ_{Π_b} , so σ_{Π_a} and σ_{Π_b} are isomorphic up to semisimplification. However, the semisimplification of σ_{Π_a} is a sum of one-dimensional characters, at least one trivial, whereas the semisimplification of σ_{Π_b} is a sum of one-dimensional characters, all nontrivial, so they cannot be isomorphic. This is a contradiction, hence π is wildly ramified in this case. \square

Corollary 8.10. *Let π be a supercuspidal representation of $GL(n, F)$. Then there is a sequence of cyclic extensions as in (8.3) such that $BC_{F_r/F}(\pi)$ is an irreducible constituent of an unramified principal series representation.*

Proof. By 5.5 and induction we know that we can find a sequence of cyclic extensions as in (8.3) such that $BC_{F_r/F}(\pi)$ is a irreducible constituent of an principal series representation whose parameter is unramified. By Proposition 8.9, such a principal series must be a parabolic induction from the Borel, as a supercuspidal on any Levi except the maximal torus would have a ramified Langlands parameter. By the local Langlands conjecture for GL_1 , it must be an induction of an unramified representation of the maximal torus, i.e. an unramified principal series. \square

Corollary 8.10 is the key point in (almost) every proof of the local Langlands correspondence for $GL(n)$. It was already mentioned in the introduction that in [LRS93, HT01, He00], this result is obtained as a consequence of Henniart's numerical correspondence; whereas in [Sch13] it is proved by a geometric argument involving nearby cycles in the local model. Starting with Corollary 8.10 one obtains the full local correspondence by an inductive study of the fibers of local base change.

Here we obtain Corollary 8.10 by a completely different argument, based on the geometric considerations of [Laf18], the global base change of [HeLe], and the existence of types for supercuspidal representations [BK]. It is questionable whether this argument is genuinely new, because the existence of the global parametrization in [Laf18], combined with the converse theorem as in [Laf02], already suffices to prove the local as well as the global correspondence. However, the reasoning used here can be applied to any G for which global base change and the existence of types is known.

9. APPLICATIONS TO GROUPS OVER p -ADIC FIELDS

Let $F = k((t))$ be a local field of characteristic p as before, and let F^\sharp be a p -adic field. We write $\mathcal{O}_F, \mathcal{O}_{F^\sharp}$ for their integer rings, and m_F, m_{F^\sharp} for the corresponding maximal ideals.

Definition 9.1. *Let n be a positive integer. We say that F and F^\sharp are n -close if*

$$\mathcal{O}_F/m_F^n \xrightarrow{\sim} \mathcal{O}_{F^\sharp}/m_{F^\sharp}^n$$

as rings.

We write $F^\? for either F or F^\sharp . Let G be a connected reductive group over $F^\? The depth of a parameter $\phi \in \mathcal{G}^{ss}(G, F)$ is defined to be the maximum n such that ϕ is trivial on the subgroup $I_{F^\?}^n$ of the inertia group $I_{F^\?}$, where we are using the upper numbering. Let $\mathcal{G}^{ss,n}(G, F) \subset \mathcal{G}^{ss}(G, F)$ denote the subset of parameters of depth at most n .$$

Theorem 9.2 (Deligne). *If F and F^\sharp are n -close, then there is a natural bijection*

$$(9.3) \quad \mathcal{G}^{ss,n}(G, F) \simeq \mathcal{G}^{ss,n}(G, F^\sharp).$$

Assume now that G is split, and let $I_n(F^\?) \subset G(\mathcal{O}_{F^\?})$ denote the n -th Iwahori filtration subgroup, as defined in [G, §3]; thus $I = I_0$ is the usual Iwahori subgroup, and I_n is the kernel of the reduction map $\mathbf{I}(\mathcal{O}_{F^\?}) \rightarrow \mathbf{I}(\mathcal{O}_{F^\?}/m_{F^\?}^n)$. Let $H(G(F^\?), I_n)$ denote the Hecke algebra of $I_n(F^\?)$ -biinvariant functions on $G(F^\?)$ with coefficients in the algebraically closed field C . Let $\mathcal{A}^n(G, F^\?) \subset \mathcal{A}(G, F)$ denote the subset of equivalence classes of irreducible admissible representations generated by vectors fixed under $I_n(F^\?)$. Any $\pi \in \mathcal{A}^n(G, F^\?)$ is then determined up to isomorphism by the representation of $H(G(F^\?), I_n)$ on its invariant subspace $\pi^{I_n(F^\?)}$. The following is Ganapathy's refinement of a theorem of Kazhdan:

Theorem 9.4 (Ganapathy-Kazhdan, [G]). *If F^\sharp and F are n -close, then there is a natural isomorphism*

$$(9.5) \quad H(G(F), I_n) \xrightarrow{\sim} H(G(F^\sharp), I_n)$$

of finitely-generated C -algebras. Moreover, there is a bijection

$$(9.6) \quad \mathcal{A}^n(G, F) \simeq \mathcal{A}^n(G, F^\sharp)$$

with the property that, if $\pi \in \mathcal{A}^n(G, F)$ corresponds to $\pi^\sharp \in \mathcal{A}^n(G, F^\sharp)$, then the invariant subspaces $\pi^{I_n(F)}$ and $\pi^{\sharp, I_n(F^\sharp)}$ are isomorphic as modules with respect to the isomorphism (9.5).

In view of these results, it would be unnatural not to make the following conjecture:

Conjecture 9.7. *For any positive integer n the following diagram commutes:*

$$\begin{array}{ccc}
 \mathcal{A}^n(G/F) & \xrightarrow{\text{Genestier-Lafforgue}} & \mathcal{G}^{ss,n}(G/F) \\
 (9.6) \downarrow & & \downarrow (9.3) \\
 \mathcal{A}^n(G/F^\sharp) & \xrightarrow{\text{Fargues-Scholze}} & \mathcal{G}^{ss,n}(G/F^\sharp)
 \end{array}$$

Here the top line is the parametrization of Genestier and Lafforgue (Theorem 3.1 (iii)), while the bottom line is the parametrization defined by Fargues and Scholze in [2] for groups over p -adic fields.

The following corollary is an immediate consequence of the Conjecture and the results of the previous sections.

Corollary 9.8. *Let F^\sharp be a finite extension of \mathbb{Q}_p with residue field k of characteristic p . Let $n \geq 0$ and assume there is a local field F of characteristic p such that F^\sharp and F are n -close. Let G be a split semisimple group over F . Suppose $|k| > 3$ and p does not divide the order of the Weyl group of G . Finally, suppose Conjecture 9.7 holds for the p -adic field F^\sharp .*

Let σ be a supercuspidal representation of $G(F^\sharp)$. Suppose the semisimple Fargues-Scholze parameter $\mathcal{L}^{ss}(\sigma)$ attached to σ is pure. Then $\mathcal{L}^{ss}(\sigma)$ is ramified.

More generally, let σ be any pure irreducible representation of $G(F^\sharp)$ with unramified Fargues-Scholze parameter. Then σ is an irreducible constituent of an unramified principal series representation.

Since every p -adic field has a local field F of characteristic p that is 0-close to F^\sharp , we have the following special case of Corollary 9.8

Corollary 9.9. *Let F^\sharp be a finite extension of \mathbb{Q}_p with residue field k of characteristic p . Let G be a split semisimple group over F . Suppose $|k| > 3$ and p does not divide the order of the Weyl group of G . Finally, suppose Conjecture 9.7 holds for the p -adic field F^\sharp . Then any pure depth zero supercuspidal representation of $G(F^\sharp)$ has ramified Fargues-Scholze parameter.*

One can hope to derive consequences of Conjecture 9.7 for general supercuspidal representations of a general p -adic field F^\sharp , one can hope to replace F^\sharp by a Galois extension F' with an n -close F for any n . This is the strategy followed by Henniart in his proof of the numerical local correspondence for p -adic fields in [He88]; the Galois extension F' in his construction was obtained by applying the results of [AC] on cyclic base change. For general groups over p -adic fields one can define a version of local cyclic base change using the global methods of [Lab99], as explained in §8.1. In order to apply this to prove results about purity, we need to know the answers to the following questions:

Question 9.10. *Does the Fargues-Scholze parametrization commute with cyclic base change, as described in §8.1?*

Question 9.11. *Let F'/F^\sharp be a cyclic extension of prime order. Suppose $\pi \in \mathcal{A}^n(G/F^\sharp)$, and π' belongs to the base change of π to $G(F')$. Is $\pi' \in \mathcal{A}^n(G/F')$?*

REFERENCES

- [A] J. Arthur, *The endoscopic classification of representations—orthogonal and symplectic groups*, *American Mathematical Society Colloquium Publications*, **61** (2013).
- [AC] J. Arthur and L. Clozel, *Simple algebras, base change, and the advanced theory of the trace formula*, *Annals of Math. Studies*, **120**, Princeton: Princeton University Press (1989).
- [BHKT] G. Böckle, M. Harris, C. Khare, J. Thorne, \hat{G} -local systems on smooth projective curves are potentially automorphic, *Acta Math.*, **223** (2019) 1–111.
- [BK] C. J. Bushnell, P. Kutzko, *The admissible dual of $GL(N)$ via compact open subgroups*, *Annals of Mathematics Studies*, **129**. Princeton University Press, Princeton, NJ, (1993).
- [De80] P. Deligne, La conjecture de Weil. II. *Publ. Math. IHES*, **52** (1980) 137–252.
- [1] L. Clozel, M. Harris, J.-P. Labesse, B.C. Ngô, *The stable trace formula, Shimura varieties, and arithmetic applications, Book 1: On the stabilization of the trace formula*, Somerville, MA: International Press (2011).
- [2] L. Fargues, P. Scholze, Geometrization of the local Langlands correspondence,
- [Fi] J. Fintzen, *Types for tame p -adic groups*, *Ann. of Mathematics*, **193** (2021) 303–346.
- [GLo] W.-T. Gan and L. Lomelí, Globalization of supercuspidal representations over function fields and applications, *J. Eur. Math. Soc.*, **20** (2018) 2813–2858.
- [G] R. Ganapathy, The local Langlands correspondence for GSp_4 over local function fields, *Am. J. Mathematics*, **137** (2015) 1441–1534.
- [GV] R. Ganapathy, S. Varma, On the local Langlands correspondence for split classical groups over local function fields. *J. Math. Inst. Jussieu*, **16** (2017) 987–1074.
- [GLa] A. Genestier and V. Lafforgue, Chtoucas restreints pour les groupes réductifs et paramétrisation de Langlands locale.
- [HT01] M. Harris, R. Taylor, *The geometry and cohomology of some simple Shimura varieties*. With an appendix by Vladimir G. Berkovich. *Annals of Mathematics Studies*, **151** Princeton University Press, Princeton, NJ (2001).
- [H19] M. Harris, Incorrigible representations, arXiv
- [HNY] J. Heinloth, B.-C. Ngô, Z. Yun, Kloosterman sheaves for reductive groups, *Annals of Math.*, **177** (2013) 241–310.
- [He88] G. Henniart, La conjecture de Langlands locale numérique pour $GL(n)$. *Ann. Sci. E.N.S.* **21** (1988) 497–544.
- [He00] G. Henniart, Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique. *Invent. Math.* **139** (2000) 439–455.
- [HeLe] G. Henniart and B. Lemaire, Changement de base et induction automorphe pour GL_n en caractéristique non nulle, *Mém. Soc. Math. Fr.*, **124** (2011).
- [HeLo] G. Henniart and L. Lomelí, *Uniqueness of Rankin-Selberg factors*, *Journal of Number Theory* **133** (2013), 4024–4035.
- [Lab99] J.-P. Labesse, Cohomologie, stabilisation, et changement de base, *Astérisque*, **257** (1999).
- [LL] J.-P. Labesse, B. Lemaire, La formule des traces tordue pour les corps de fonctions, <https://arxiv.org/pdf/2102.02517v1.pdf>.
- [Laf02] L. Lafforgue, Chtoucas de Drinfeld et correspondance de Langlands, *Invent. Math.*, **147** (2002) 1–241.
- [Laf18] V. Lafforgue, Chtoucas pour les groupes réductifs et paramétrisation de Langlands globale, *JAMS*, **31** (2018) 719–891.

- [LRS93] G. Laumon, M. Rapoport, and U. Stuhler, D-elliptic sheaves and the Langlands correspondence, *Invent. Math.*, **113** (1993) 217–338.
- [Lo19] L. Lomelí, Rationality and holomorphy of Langlands-Shahidi L -functions over function fields, *Math. Z.* **291** (2019) 711–739.
- [MW] C. Mœglin and J.-L. Waldspurger, Stabilisation de la formule des traces tordue. Vol. 1, 2. *Progress in Mathematics*, 316-317. Birkhäuser/Springer, Cham, (2016).
- [ST] W. Sawin and N. Templier, On the Ramanujan conjecture for automorphic forms over function fields, I. Geometry, <https://arxiv.org/pdf/1805.12231.pdf>.
- [Sch13] P. Scholze, The local Langlands correspondence for GL_n over p -adic fields. *Invent. Math.*, **192** (2013) 663–715.
- [TY] R. Taylor and T. Yoshida, Compatibility of local and global Langlands correspondences. *J. Amer. Math. Soc.*, **20** (2007) 467–493.
- [Y16] Z. Yun, Epipelagic representations and rigid local systems, *Sel. Math. New Ser.* , **22** (2016) 1195–1243.

WEE TECK GAN, MATHEMATICS DEPARTMENT, NATIONAL UNIVERSITY OF SINGAPORE, BLOCK S17, 10 LOWER KENT RIDGE ROAD SINGAPORE 119076

Email address: `matgwt@nus.edu.sg`

MICHAEL HARRIS, DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, USA

Email address: `harris@math.columbia.edu`

WILL SAWIN, DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, USA

Email address: `sawin@math.columbia.edu`