

Lattices, injective metrics and the $K(\pi, 1)$ conjecture

Thomas Haettel

September 17, 2021

ABSTRACT. Starting with a lattice with an action of \mathbb{Z} or \mathbb{R} , we build a Helly graph or an injective metric space. We deduce that the ℓ^∞ orthoscheme complex of any bounded graded lattice is injective. We also prove a Cartan-Hadamard result for locally injective metric spaces. We apply this to show that any Garside group acts on an injective metric space and on a Helly graph. We also deduce that the natural piecewise ℓ^∞ metric on any Euclidean building of type \widetilde{A}_n extended, \widetilde{B}_n , \widetilde{C}_n or \widetilde{D}_n is injective, and its thickening is a Helly graph.

Concerning Artin groups of Euclidean types \widetilde{A}_n and \widetilde{C}_n , we show that the natural piecewise ℓ^∞ metric on the Deligne complex is injective, the thickening is a Helly graph, and it admits a convex bicombing. This gives a metric proof of the $K(\pi, 1)$ conjecture, as well as several other consequences usually known when the Deligne complex has a CAT(0) metric.

Contents

1	Review of posets, Artin groups and injective metric spaces	6
1.1	Posets and lattices	6
1.2	Coxeter groups, Artin groups and Deligne complexes	7
1.3	Injective metrics and Helly graphs	9
2	The thickening of a lattice	12
3	The affine version of a lattice	15
4	Application to Euclidean buildings and the Deligne complex of type \widetilde{A}_n	23
5	The lattice of maximal parabolic subgroups of braid groups	27
5.1	The lattice of cut-curves	27
5.2	Cosets in braid groups	29
6	The thickening of a semilattice	31
7	Application to Euclidean buildings and the Deligne complex of other Euclidean types	34
8	Bicombings on Deligne complexes in types \widetilde{A}_n and \widetilde{C}_n	36

Keywords : Injective metrics, Helly graphs, lattices, Artin groups, Euclidean buildings, Cartan-Hadamard. **AMS codes** : 52A35, 20E42, 05B35, 06A12

Introduction

Injective metric spaces are geodesic metric spaces where every family of pairwise intersecting balls has a non-empty global intersection. Their discrete counterpart are called Helly graphs. Their use in geometric group theory is recent and growing, see notably [Lan13], [HO19], [CCG⁺20], [HHP20], [OV20], [Hae21]. Roughly speaking, CAT(0) spaces are typically locally Euclidean spaces, whereas injective metric spaces are typically locally ℓ^∞ metric spaces. Injective metric spaces display many nonpositive curvature features observed in CAT(0) spaces. We believe that Helly graphs and injective metric spaces may also reveal more powerful than CAT(0) spaces for some purposes.

Firstly, it appears that many nonpositively curved groups (notably most CAT(0) groups) have a nice action on an injective metric space, so that the theory encompasses a vast class of groups: CAT(0) cubical groups, hyperbolic groups, relatively hyperbolic groups, uniform lattices in semisimple Lie groups over local fields, braid groups and more generally Garside groups, Artin groups of type FC, mapping class groups and more generally hierarchically hyperbolic groups. For instance, Helly graphs admit a simple combinatorial local characterization (see Theorem 1.11), which makes them potentially as powerful as CAT(0) cube complexes, whereas piecewise Euclidean CAT(0) complexes desperately lack a combinatorial characterization.

Secondly, we can still deduce many of the results that hold true for CAT(0) spaces and CAT(0) groups in the setting of injective spaces and injective groups (groups acting geometrically on injective spaces). For instance, injective metric spaces have a conical geodesic bicombing and are thus contractible, and every isometric bounded group action has a fixed point. Moreover, every injective group is semi-hyperbolic (see [AB95]) and satisfies the Farrell-Jones conjecture (see [KR17]). Sometimes we can even deduce stronger results than in the CAT(0) setting: as a sample result, note that any group acting properly and cocompactly on a Helly graph is biautomatic (see [CCG⁺20]), whereas not all CAT(0) groups are biautomatic (see [LM21]).

In this article, we will pursue this philosophy. On one hand, we develop results useful to prove that some metric spaces are injective. On the other hand, we apply these results to Euclidean buildings and Deligne complexes of Euclidean Artin groups. We believe, however, that the scope of our results will not be limited to Artin groups of Euclidean type, and could concern much larger classes of groups.

We now present a very simple criterion (already appearing in a restricted form in [Hae21]) showing how to build a Helly graph or an injective metric space, starting with a lattice and an action of \mathbb{Z} or \mathbb{R} .

Theorem A. (*Theorem 2.1*) *Assume that L is a lattice such that each upperly bounded subset of L has a join. Assume that there is an order-preserving increasing continuous action $(f_t)_{t \in H}$ of $H = \mathbb{Z}$ or \mathbb{R} on L , such that*

$$\forall x, y \in L, \exists t \in H_+, f_{-t}(x) \leq y \leq f_t(x).$$

Let us define the following metric d on L :

$$\forall x, y \in L, d(x, y) = \inf\{t \in H_+ \mid f_{-t}(x) \leq y \leq f_t(x)\}.$$

- *If $H = \mathbb{Z}$, the metric space (L, d) is the vertex set of a Helly graph.*
- *If $H = \mathbb{R}$, the metric space (L, d) is injective.*

We applied this criterion in [Hae21] to prove that the thickening of a Bruhat-Tits Euclidean building of extended type \widetilde{A}_n is a Helly graph. It turns out that the Bruhat-Tits restriction is unnecessary.

Theorem B. (Theorems 4.4 and 7.3) *The natural ℓ^∞ metric on any Euclidean building of type \widetilde{A}_n extended, \widetilde{B}_n , \widetilde{C}_n or \widetilde{D}_n is injective. Furthermore, the thickening of its vertex set is a Helly graph.*

Since Euclidean buildings are also endowed with CAT(0) metrics, we may wonder about the importance of this result. For once, it gives a lot of new examples to the theory of injective metric spaces and Helly graphs. Another consequence is another approach to Swiatkowski's result (see [S06]) that cocompact lattices in Euclidean buildings are biautomorphic, in types \widetilde{A}_n , \widetilde{B}_n , \widetilde{C}_n or \widetilde{D}_n . The proof for types \widetilde{B}_n , \widetilde{C}_n and \widetilde{D}_n rely on a generalization for graded semilattices, see Section 6.

Another immediate consequence of Theorem A is the following.

Theorem C. (Corollary 2.2) *The thickening of the Cayley graph of any Garside group with respect to its simple elements is a Helly graph.*

Note that, in the case of a finite type Garside group, this is due to Huang and Osajda (see [HO19]). However, our proof is different, and does not rely on the deep local-to-global result for Helly graphs (see Theorem 1.11). Besides, it also works in the case of a Garside group with infinite set of simple elements.

One application is the study of the orthoscheme complex of a bounded, graded lattice L . Note that simplices of the geometric realization $|L|$ of L correspond to chains in L , so that one can endow each simplex with metric of the standard ℓ^∞ orthosimplex associated to this order on vertices (see Section 1 for details). We endow the geometric realization $|L|$ of L with a lattice structure, and use Theorem A to deduce the following.

Theorem D. (Theorem 3.9) *Let L denote a bounded, graded lattice. Then the orthoscheme complex $|L|$ of L , with the piecewise ℓ^∞ metric, is injective.*

When L is a bounded, graded lattice, one may endow its orthoscheme complex with the piecewise Euclidean metric. Deciding whether it is a CAT(0) metric space is a very difficult question. It turns out that for orthoscheme complexes of posets, the CAT(0) property is more restrictive than the injective property, see Theorem 3.10.

One famous conjecturally CAT(0) example is the dual braid complex, defined by Brady and McCammond in [BM10]. The n -strand braid group B_n has a standard Garside structure, associated to the "half-turn" as Garside element. The braid group also enjoys a dual Garside structure defined by Bessis (see [Bes03]), associated to the rotation of a n th of a turn as Garside element. Associated to this dual Garside structure, one may consider the geometric realization and endow it with the piecewise Euclidean orthoscheme metric (see Section 1). Brady and McCammond conjecture that this so-called dual braid complex is CAT(0) for all braid groups, but it has only been proved for $n \leq 7$ (see [BM10], [HKS16], [Jeo20]). However, an immediate consequence of Theorem D is the following.

Theorem E. (Corollary 3.11) *The Garside complex of any Garside group, endowed with the piecewise ℓ^∞ orthoscheme metric, is injective.*

As a particular case, the dual braid complex, endowed with the piecewise ℓ^∞ orthoscheme metric, is injective for all braid groups. This also holds, more generally, for every spherical type Artin group with some Garside structure. Note that Theorem D proves local injectivity, and one also needs a Cartan-Hadamard result for injective metric spaces in order to conclude. We therefore rely on the local-to-global result for Helly graphs to prove the following generalization of [Mie17] in the non-proper setting (see Section 1 for the definition of semi-uniformly locally injective). This result is clearly of independent interest.

Theorem F (Cartan-Hadamard for injective metric spaces). (*Theorem 1.13*) *Let X denote a complete, simply connected, semi-uniformly locally injective metric space. Then X is injective.*

Another very promising family of examples are Deligne complexes of Artin groups (see Section 1 for definitions). Note that buildings and Deligne complexes of Artin groups are closely related: in fact in his original article (see [Del72]), Deligne called "buildings for generalized braid groups" the complexes later called Deligne complexes. However, to the best of our knowledge, the close relationship between Euclidean buildings and Deligne complexes of Euclidean type Artin groups has not yet been exploited in the literature. Notably, the automorphism groups of Euclidean buildings do not possess a Garside structure, and the Deligne complexes do not have an apartment system as rich as in the building case. However, one common feature is that they locally look like a lattice, which is the key combinatorial property we are using in this article.

Associated to every Coxeter graph Γ with vertex set S , we may define the Coxeter group $W(\Gamma)$, the Artin group $A(\Gamma)$ and the hyperplane complement $M(\Gamma)$ (see Section 1). The Coxeter group $W(\Gamma)$ acts naturally on $M(\Gamma)$, and $A(\Gamma)$ is the fundamental group of the quotient $W(\Gamma)\backslash M(\Gamma)$. One very natural question is to decide whether it is a classifying space. This is the statement of the following conjecture.

Conjecture G (The $K(\pi, 1)$ conjecture). *The hyperplane complement $M(\Gamma)$ is aspherical.*

The $K(\pi, 1)$ conjecture has been proved for spherical type Artin groups by Deligne in [Del72], and for Euclidean type Artin groups by Paolini and Salvetti in [PS21] very recently, even for the type \widetilde{D}_n . Another approach, more closely related to the metric approach of this article, has been used by Charney and Davis in [CD95b] to prove the $K(\pi, 1)$ conjecture for Artin groups of type FC or of 2-dimensional type. Their proof relies on the use of a simplicial complex, called the Deligne complex $\Delta(\Gamma)$, which is the geometric realization of the poset of cosets of spherical type parabolic subgroups (see Section 1). This complex (in this form) has been defined by Charney and Davis in [CD95b], where they proved that $\Delta(\Gamma)$ is homotopy equivalent to the universal cover of $W(\Gamma)\backslash M(\Gamma)$. In other words, the $K(\pi, 1)$ conjecture is equivalent to the contractibility of the Deligne complex $\Delta(\Gamma)$.

Charney and Davis's method for proving that the Deligne complex is contractible is to endow it with a CAT(0) metric. This works for Artin groups of type FC or of 2-dimensional type. However, in the general case, the key question is to decide whether the Deligne complex for the braid group is CAT(0) with Moussong's metric. It is only known for the braid group up to 4 strands only (see [Cha04]). However, we will see that the natural piecewise ℓ^∞ metric is injective for all braid groups, up to taking the product with \mathbb{R} .

Theorem H. (*Theorem 4.4*) *Let Δ denote the Deligne complex of the Artin group of Euclidean type \widetilde{A}_n . Then the natural piecewise ℓ^∞ length metric on $\Delta \times \mathbb{R}$ is injective. Moreover, the thickening of $\Delta \times \mathbb{R}$ is a Helly graph.*

The proof consists in applying Theorem D to prove that the Deligne complex is locally injective. One key combinatorial property is that the Deligne complex in type A_n is essentially a lattice, see Section 5. The proof of the lattice property, through the cut-curve lattice, is due to Crisp and McCammond, copied here with their permission.

Note that one may wonder whether it is necessary to consider the direct product with \mathbb{R} . In fact, Hoda proved (see [Hod20]) that the Euclidean Coxeter group of type \widetilde{A}_n is not Helly for $n \geq 2$, even though its direct product with \mathbb{Z} is. We made a similar distinction for automorphism groups of Euclidean buildings of type \widetilde{A}_n in [Hae21]. We therefore strongly

believe that there is no injective metric on the Deligne complex of type \widetilde{A}_n itself which is invariant under the Artin group. However, we will see in Theorem L below that there is a convex bicombing on the Deligne complex itself.

In order to deal with the Euclidean type \widetilde{C}_n , we first prove a generalization of Theorem D for graded semilattices, see Section 6.

Theorem I. *(Theorem 7.3) Let Δ denote the Deligne complex of the Artin group of Euclidean type \widetilde{C}_n . Then the natural piecewise ℓ^∞ length metric on Δ is injective. Moreover, the thickening of Δ is a Helly graph.*

An immediate consequence is another proof of the $K(\pi, 1)$ conjecture in Euclidean types \widetilde{A}_n and \widetilde{C}_n , originally due to Okonek (see [Oko79]). The novelty is that it is the first metric proof. Moreover, in Charney and Davis's approach to the $K(\pi, 1)$ conjecture by showing that Mousson's metric is CAT(0), the main difficulty is to prove that it is locally CAT(0) for the braid groups. It is precisely this statement that we are able to prove in the injective setting.

Corollary J (Okonek). *The Deligne complex Δ of Euclidean type \widetilde{A}_n or \widetilde{C}_n is contractible. In particular, the $K(\pi, 1)$ conjecture holds in these cases.*

Moreover, several results have been proved, relying on the assumption that one may endow the Deligne complex with a piecewise Euclidean CAT(0) metric. Crisp studied the fixed-point subgroup under a symmetry group of the Artin system (see [Cri00]). Godelle studied the centralizer and normalizer of standard parabolic subgroups (see [God07]). Morris-Wright studied the intersections of parabolic subgroups (see [MW19]). It turns out almost all the arguments merely used the existence of an equivariant geodesic bicombing on the Deligne complex (see the end of Section 3 for a definition of a bicombing). Moreover, work of Descombes and Lang notably (see [DL16] and [DL15]) justify the importance of convex geodesic bicomblings for themselves. We therefore state the following conjecture, which may be seen as a metric strategy for the proof of the $K(\pi, 1)$ conjecture.

Conjecture K. *The Deligne complex of any Artin group A has an A -invariant metric that admits a convex, consistent, reversible geodesic bicombing.*

We are able to prove this conjecture in spherical types A_n , B_n and Euclidean types \widetilde{A}_n , \widetilde{C}_n , and we believe that our result represents a major step towards the general case.

Theorem L. *(Theorem 8.1) Let Δ denote the Deligne complex of the Artin group of spherical type A_n or B_n or Euclidean type \widetilde{A}_n or \widetilde{C}_n . There exists a metric on Δ , invariant under the Artin group, which admits a convex, consistent, reversible geodesic bicombing.*

If an Artin group satisfies Conjecture K, then the $K(\pi, 1)$ conjecture follows. Moreover, we may also list consequences of work of Crisp (see [Cri00]), Godelle (see [God07]) and Morris-Wright (see [MW19] and [CGGMW19]) that rely on the assumption that the Deligne complex has a CAT(0) metric. However, most of the arguments only use the geodesic bicombing. The following results were only known for Artin groups of type FC or of 2-dimensional type.

Corollary M. *(Corollaries 8.3, 8.4 and 8.5) Let A denote the Artin group of Euclidean type \widetilde{A}_n or \widetilde{C}_n .*

- *The intersection of any parabolic subgroups of A is a parabolic subgroup.*
- *A satisfies Properties (\star) , $(\star\star)$ and $(\star\star\star)$ from [God07], notably:*

★ For any subset $X \subset S$, we have

$$\text{Com}_A(A_X) = N_A(A_X) = A_X \cdot QZ_A(X),$$

where the quasi-centralizer of X is $QZ_A(X) = \{g \in A \mid g \cdot X = X\}$.

★ For any subsets $X, Y \subset S$ and any $g \in A$, we have

$$gA_Xg^{-1} \subset A_Y \implies \exists h \in A_Y, \exists Z \subset X, gA_Xg^{-1} = hA_Zh^{-1}.$$

- For any group G of symmetries of the Artin system, the fixed point subgroup A^G is isomorphic to an Artin group.

In view of Conjecture K, looking for other consequences of the existence of a convex bicombing on the Deligne complex may reveal fruitful.

Structure of the article

In Section 1, we review basic definitions of posets, lattices, Artin groups, Deligne complexes, injective metric spaces and Helly graphs. We also prove the Cartan-Hadamard Theorem for injective metric spaces. In Section 2, we prove the central simple criterion showing how to produce an injective metric space or a Helly graph starting from a lattice with an action of \mathbb{Z} or \mathbb{R} . In Section 3, we apply this criterion to prove that the orthoscheme complex of a bounded, graded lattice is injective. In Section 4, we use this to prove that, for Euclidean buildings and the Deligne complex in Euclidean type \widetilde{A}_n , the natural piecewise ℓ^∞ metric is injective. In Section 6, we show how to adapt the criterion to a mere semilattice with some extra property. We then apply it in Section 7 to prove that, for Euclidean buildings and the Deligne complex in Euclidean type \widetilde{C}_n , the natural piecewise ℓ^∞ metric is injective. In the last Section 8, we use the convex bicombing on the Deligne complexes to deduce many corollaries about parabolic subgroups of Artin groups.

Acknowledgments: We would like to thank very warmly Jon McCammond, Jingyin Huang, Damian Osajda and Luis Paris for very interesting discussions. We are indebted to Jon McCammond and John Crisp for their authorization to copy their work about the lattice of cut-curves. We also would like to thank the organizers of the ICMS Edinburgh 2021 meeting "Perspectives on Artin groups", where we had fruitful discussions. We also would like to thank Jérémie Chalopin, Victor Chepoi, Anthony Genevois, Hiroshi Hirai, Alexandre Martin and Bert Wiest for various discussions.

We acknowledge support from French project ANR-16-CE40-0022-01 AGIRA.

1 Review of posets, Artin groups and injective metric spaces

1.1 Posets and lattices

We recall briefly basic definitions related to posets and lattices.

Definition 1.1 (Poset).

Let L denote a poset. A *chain* of L is a totally ordered non-empty subset of L . A *maximal chain* is a chain that is maximal with respect to inclusion. A finite chain of $n + 1$ elements $x_0 < x_1 < \dots < x_n$ is called of *length* n . The poset L is called *bounded below* (resp. *above*) if it has a global minimum denoted 0 (resp. global maximum denoted 1). The poset L is called *bounded* if it is both bounded above and below. The poset L has *rank* n if it is bounded and all maximal chains have length n .

Definition 1.2 (Interval).

Given two elements $x \leq y$ in a poset L , we denote the *interval*

$$I(x, y) = \{z \in L \mid x \leq z \leq y\}.$$

In case of possible confusion, we may also denote the interval $I_L(x, y)$ to emphasize that we are considering the interval in the poset L . The poset L is called *graded* if every interval of L has a rank. Let x denote an element in a graded poset L that is bounded below, the *rank* of x is the rank of the interval $I(0, x)$.

Definition 1.3 (Lattice).

Given elements x, y in a poset L , if there exists a unique maximal lower bound to $\{x, y\}$, it is called the *meet* of x and y and denoted $x \wedge y$. Similarly, if there exists a unique minimal upper bound to $\{x, y\}$, it is called the *join* of x and y and denoted $x \vee y$. If any two elements of L have a meet (resp. a join), the poset L is called a *meet-semilattice* (resp. *join-semilattice*). If any two elements of L have a meet and a join, the poset L is called a *lattice*.

We will now describe a very simple criterion due to Brady and McCammond (see [BM10]) to decide when a bounded graded poset is a lattice.

Definition 1.4 (Bowtie).

In a poset L , a *bowtie* consists of 4 distinct elements a, b, c, d such that a, c are minimal upper bounds of b, d , and b, d are maximal lower bounds for a, c .

Proposition 1.5. [BM10, Proposition 1.5] *Let L denote a bounded graded poset. Then L is a lattice if and only if L does not contain a bowtie.*

1.2 Coxeter groups, Artin groups and Deligne complexes

We recall the definitions of Coxeter groups, Artin groups, and their associated Deligne complex.

For every finite simple graph Γ with vertex set S and with edges labeled by some integer in $\{2, 3, \dots\}$, one associates the Coxeter group $W(\Gamma)$ with the following presentation:

$$W(\Gamma) = \langle S \mid \forall \{s, t\} \in \Gamma^{(1)}, \forall s \in S, s^2 = 1, \forall s, t \in S, [s, t]_m = [t, s]_m \text{ if the edge } \{s, t\} \text{ is labeled } m \rangle,$$

where $[s, t]_m$ denotes the word $ststs\dots$ of length m .

The associated Artin group $A(\Gamma)$ is defined by a similar presentation:

$$A(\Gamma) = \langle S \mid \forall \{s, t\} \in \Gamma^{(1)}, \forall s, t \in S, [s, t]_m = [t, s]_m \text{ if the edge } \{s, t\} \text{ is labeled } m \rangle.$$

The groups $A(\Gamma)$ are also called Artin-Tits groups, since they have been defined by Tits in [Tit66].

Note that only the relations $s^2 = 1$ have been removed, so that there is a natural surjective morphism from $A(\Gamma)$ to $W(\Gamma)$. Also note that when $m = 2$, then s and t commute, and when $m = 3$, then s and t satisfy the classical braid relation $sts = tst$.

The knowledge of general Artin groups is extremely limited (see notably [McC17], [Cha], [GP12]). In particular, we do not know whether the word problem is solvable, nor whether they are torsion-free.

Most results about Artin-Tits groups concern particular classes. The Artin group $A(\Gamma)$ is called:

- of *spherical type* if its associated Coxeter group $W(\Gamma)$ is finite, i.e. may be realized as a reflection group of a sphere.





	Spherical type		Euclidean type
$A_n, n \geq 2$	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \bullet$	\widetilde{A}_n	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \bullet$ 
$B_n, n \geq 2$	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \underline{4} \bullet$	\widetilde{B}_n	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \underline{4} \bullet$ 
$C_n = B_n, n \geq 2$	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \underline{4} \bullet$	\widetilde{C}_n	$\bullet \text{---} \underline{4} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \underline{4} \bullet$ 
$D_n, n \geq 4$	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \begin{matrix} \bullet \\ \bullet \end{matrix}$	\widetilde{D}_n	$\bullet \text{---} \bullet \text{---} \bullet \quad \dots \quad \bullet \text{---} \bullet \text{---} \begin{matrix} \bullet \\ \bullet \end{matrix}$ 

Table 1: Spherical and Euclidean irreducible diagrams of type A , B , C and D

- of *Euclidean type* if its associated Coxeter group $W(\Gamma)$ may be realized as a reflection group of a Euclidean space.

We recall in Table 1 the classification of the four infinite families of spherical and Euclidean irreducible diagrams, see [Bou02] for the full classification. We only present those because we will only consider these types in this article. Note that we use in this table the convention of Dynkin diagrams: vertices that are not joined by an edge commute, and we drop the label 3 from edges.

Artin groups are closely related to hyperplane complements, which can be presented in a simple way in spherical and Euclidean types. Fix a Coxeter group $W = W(\Gamma)$ of spherical type or Euclidean type acting by reflections on a sphere \mathbb{S}^{n-1} or a Euclidean space \mathbb{R}^n . In case W is of spherical type, consider the associated linear action on \mathbb{R}^n . A conjugate of an element of the standard generating set S is called a *reflection*. Let \mathcal{R} denote the set of reflections in W . Consider the family of affine hyperplanes of \mathbb{R}^n

$$\mathcal{H} = \{H_r \mid r \in \mathcal{R}\},$$

where $H_r \subset \mathbb{R}^n$ denotes the fixed point set of the reflection r .

The complexified hyperplane arrangement is

$$M(\Gamma) = \mathbb{C}^n \setminus \bigcup_{r \in \mathcal{R}} (\mathbb{C} \otimes H_r).$$

Note that W acts naturally on M , and we have the following (see [VdL83]):

$$\pi_1(W(\Gamma) \backslash M(\Gamma)) \simeq A(\Gamma).$$

So the Artin group $A(\Gamma)$ appears as the fundamental group of (a quotient of) a complexified hyperplane arrangement. One very natural question is to decide whether it is a classifying space. This is the statement of the following conjecture.

Conjecture ($K(\pi, 1)$ conjecture). *The hyperplane arrangement $M(\Gamma)$ is aspherical.*

This conjecture has been proved for spherical type Artin groups by Deligne in [Del72], and for Euclidean type Artin groups by Paolini and Salvetti in [PS21] very recently, even for the type \widetilde{D}_n . Another approach, more closely related to the content of this article, has been used by Charney and Davis in [CD95a] to prove the $K(\pi, 1)$ conjecture for Artin groups of type FC or of 2-dimensional type. Their proof relies on the use of a simplicial complex, called the Deligne complex, and endow it with a particular metric to show that it is contractible.

We will now recall the definition of the Deligne complex of an Artin group $A = A(\Gamma)$. A *standard parabolic* subgroup of A is the subgroup $A_T = \langle T \rangle$ generated by a subset T

of S , the standard generating set of A . A *parabolic* subgroup denotes any conjugate of a standard parabolic subgroup. Let us denote

$$\mathcal{S}_f = \{T \subset S \mid W_T \text{ is finite}\}.$$

The *Deligne complex* $\Delta = \Delta(\Gamma)$ is the order complex of the set of cosets of parabolic subgroups

$$\{gA_T \mid g \in A, T \in \mathcal{S}_f\},$$

where the partial order is given by the inclusion $gA_T \subset g'A_{T'}$ of cosets.

One key property of the Deligne complex is that it has the same homotopy type as the universal cover of the hyperplane complement:

Theorem 1.6. [CD95b, Theorem 1.5.1] *The Deligne complex $\Delta(\Gamma)$ is homotopy equivalent to the universal cover of the quotient of the hyperplane complement $W(\Gamma) \backslash M(\Gamma)$.*

In particular, the $K(\pi, 1)$ conjecture amounts to proving that the Deligne complex is contractible.

1.3 Injective metrics and Helly graphs

We briefly recall basic definitions of injective metric spaces and Helly graphs. We will also state the local-to-global result for Helly graphs and deduce the analogous Cartan-Hadamard result for injective metric spaces.

A geodesic metric space is called *injective* (one may also say *hyperconvex*, or *absolute 1-Lipschitz retract*) if any family of pairwise intersecting closed balls has a non-empty global intersection, the so-called *Helly property*. We refer the reader to [Lan13] for a presentation of injective metric spaces.

A connected graph X is called *Helly* if any family of pairwise intersecting combinatorial balls of X has a non-empty global intersection, in other words the combinatorial balls satisfy the Helly property. We refer the reader to [CCG⁺20] for a presentation of Helly graphs and Helly groups.

In order to endow a simplicial complex with a potentially injective (or CAT(0)) metric, it is natural to ask for metric simplices which may tile the Euclidean space \mathbb{R}^n . One choice is to consider the barycentric subdivision of the standard cubical tiling of \mathbb{R}^n , whose simplices are orthosimplices, which we now formally define.

Definition 1.7 (Orthosimplex). The *standard orthosimplex* of dimension n is the simplex of \mathbb{R}^n with vertices $(0, \dots, 0), (1, 0, \dots, 0), \dots, (1, 1, \dots, 1)$ (see Figure 1). One may endow the simplex with the standard ℓ^p metric on \mathbb{R}^n , for any $p \in [1, \infty]$. Throughout this article (unless in Theorem 3.10), we will only consider the ℓ^∞ metric, called the ℓ^∞ orthosimplex. Note that any n -simplex with a total order on its vertices $v_0 < v_1 < \dots < v_n$ may be identified uniquely with the ℓ^∞ orthosimplex of dimension n , where each v_i is identified with the vertex $(1, \dots, 1, 0, \dots, 0)$ with i ones and $n - i$ zeros. Also note that reversing the total order on the vertices gives rise to an isometry of the orthosimplex.

Definition 1.8 (Simplicial complex with ordered simplices and maximal edges). A simplicial complex X is called *with ordered simplices* if each simplex of X has a total order on its vertex set, which is consistent with respect to inclusions of simplices of X . Moreover, we say that X has *maximal edges* if, given any simplex σ of X , there exist adjacent vertices a, b in X such that $\sigma \cup \{a, b\}$ is a simplex of X , and such that any vertex c of X adjacent to both a and b satisfies $a \leq c \leq b$. Such an edge $\{a, b\}$ is then called a maximal edge of X .




Figure 1: The standard 3-dimensional orthosimplex.

Definition 1.9 (Orthoscheme complex). Let X denote a finite-dimensional simplicial complex with ordered simplices. Then one may endow each simplex of X with the associated ℓ^∞ orthoscheme metric. Then the geometric realization of $|X|$, endowed with the length metric associated with the ℓ^∞ orthoscheme metric on each simplex, is called the ℓ^∞ orthoscheme complex of X .

As a particular case, if L is a graded poset with bounded rank, then the geometric realization of $|L|$ satisfies the assumptions, so we may talk about the ℓ^∞ orthoscheme complex of L . The following is an immediate adaptation of [BH99, Theorem 7.13].

Theorem 1.10. *Let X denote a finite-dimensional simplicial complex with ordered simplices. Then its ℓ^∞ orthoscheme complex is a complete length space.*

Proof. Note that since X has dimension n , the ℓ^∞ orthoscheme complex of X has finitely many isometry types of cells: the standard orthoscheme k -simplices, where $k \leq n$. The proof of [BH99, Theorem 7.13] adapt without change to this situation. \square

One key property in the study of Helly graphs is the following local-to-global statement (see [CCHO21]). Recall that a *clique* of a graph is a complete subgraph. A graph is called *clique-Helly* if its family of maximal cliques satisfies the Helly property. The triangle complex of a simplicial graph X is the simplicial 2-complex whose 1-skeleton is X , and whose 2-simplices correspond to triangles in X .

Theorem 1.11. [CCHO21, Theorem 3.5] *Let X denote a graph. Then X is Helly if and only if X is clique-Helly and its triangle complex is simply connected.*

In order to transfer this local-to-global property to injective metric spaces, we will need the following technical lemma. Say that a metric space is ε -*coarsely injective* (for some $\varepsilon \geq 0$) if, for any families $(x_i)_{i \in I}$ in X and $(r_i)_{i \in I}$ in \mathbb{R}_+ such that $\forall i, j \in I, d(x_i, x_j) \leq r_i + r_j$, we have $\bigcap_{i \in I} B_X(x_i, r_i + \varepsilon) \neq \emptyset$. Note that when $\varepsilon = 0$, we recover the definition of an injective metric space.

Lemma 1.12. *Let X denote a complete metric space that is ε -coarsely injective for every $\varepsilon > 0$. Then X is injective.*

Proof. Fix a family $(B_X(x_i, r_i))_{i \in I}$ such that $\forall i, j \in I, d(x_i, x_j) \leq r_i + r_j$. For any $\varepsilon > 0$, let us denote $A_\varepsilon = \bigcap_{i \in I} B_X(x_i, r_i + \varepsilon)$, which is non-empty by assumption.

Fix $0 < \varepsilon \leq \varepsilon'$, we will prove that the Hausdorff distance between A_ε and $A_{\varepsilon'}$ is at most $\varepsilon + \varepsilon'$. Note that $A_\varepsilon \subset A_{\varepsilon'}$. Fix $x \in A_{\varepsilon'}$: we know that the ball $B_X(x, \varepsilon')$

intersects each ball $B(x_i, r_i)$, for $i \in I$. So we deduce that the ball $B_X(x, \varepsilon' + \varepsilon)$ intersects $A_\varepsilon = \bigcap_{i \in I} B_X(x_i, r_i + \varepsilon)$. So we have proved that the Hausdorff distance between A_ε and $A_{\varepsilon'}$ is at most $\varepsilon + \varepsilon'$.

For each $n \in \mathbb{N}$, consider by induction $x_n \in A_{2^{-n}}$ such that $\forall n \geq 0, d_X(x_{n+1}, x_n) \leq 2^{-n} + 2^{-(n+1)} \leq 2^{-n+1}$. For each $0 \leq n \leq m$, we have $d_X(x_n, x_m) \leq 2^{-n+2}$, hence the sequence $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in X . Since X is complete, it has a limit $y \in X$. For each $n \in \mathbb{N}$, we have $y \in A_{2^{-n}}$, so for each $i \in I$ we have $d_X(y, x_i) \leq r_i + 2^{-n}$. We deduce that, for each $i \in I$, we have $d_X(y, x_i) \leq r_i$. In other words, y belongs to the intersection $\bigcap_{i \in I} B(x_i, r_i)$: we have proved that X is injective. \square

Say that a metric space is *uniformly locally injective* if there exists $\varepsilon > 0$ such that each ball of radius ε is injective. Say that a metric space is *semi-uniformly locally injective* if there exists $\varepsilon > 0$ such that each ball of radius ε is uniformly locally injective. For instance, any locally compact, locally injective metric space is semi-uniformly locally injective.

As a concrete example, if X denotes the injective hull of the hyperbolic plane $\mathbb{H}_{\mathbb{R}}^2$ and Γ is a non-uniform lattice in $\mathrm{PGL}(2, \mathbb{R}) = \mathrm{Isom}(\mathbb{H}_{\mathbb{R}}^2)$, then the quotient $\Gamma \backslash X$ is semi-uniformly locally injective, but not uniformly locally injective. There are similar examples in higher rank, see [Hae21].

There are also non-locally compact examples: let X denote a metric simplicial graph, such that the systole at each vertex of X is bounded below. Then X is locally uniformly locally injective. However, if the systole of X is 0, then X is not uniformly locally injective.

We now prove a Cartan-Hadamard theorem for such injective metric spaces, relying on the local-to-global property for Helly graphs, Theorem 1.11. Note that this statement generalizes [Mie17, Theorem 1.2] without the local compactness assumption.

Theorem 1.13 (Cartan-Hadamard for injective metric spaces). *Let X denote a complete, simply connected, semi-uniformly locally injective metric space. Then X is injective.*

Proof. We will first prove the statement when X is uniformly locally injective. Fix $\varepsilon > 0$ small enough such that balls in X of radius at most 2ε are injective. Consider the graph Γ_ε with vertex set X , and with an edge between $x, y \in X$ if $d(x, y) \leq \varepsilon$. Since X is geodesic, Γ_ε is a connected graph. Also note that, for any $x \in X$ and $n \in \mathbb{N}$, we have

$$B_{\Gamma_\varepsilon}(x, n) = B_X(x, n\varepsilon).$$

We will prove that the family of combinatorial 1-balls in Γ_ε satisfies the Helly property. Fix a family of vertices $(x_i)_{i \in I}$ of vertices of Γ_ε such that $\forall i, j \in I, d_{\Gamma_\varepsilon}(x_i, x_j) \leq 2$. Then the family of metric balls $(B_X(x_i, \varepsilon))_{i \in I}$ in X pairwise intersects: since such balls have the Helly property by assumption, so we deduce that there exists $y \in X$ such that $\forall i \in I, d_X(x_i, y) \leq \varepsilon$. In other words, the vertex $y \in \Gamma_\varepsilon$ lies in the intersection of all combinatorial 1-balls $(B_{\Gamma_\varepsilon}(x_i, 1))_{i \in I}$.

We now deduce that Γ_ε is clique-Helly: fix a family of pairwise intersecting maximal cliques $(\sigma_i)_{i \in I}$ of Γ_ε . Then the family of combinatorial 1-balls centered at each vertex of each clique σ_i , for $i \in I$, pairwise intersects: according to the previous paragraph, we deduce that there exists a vertex $y \in \Gamma_\varepsilon$ adjacent to each vertex of each clique σ_i , for $i \in I$. Since each clique σ_i is maximal, we deduce that y belongs to the intersection of all σ_i , for $i \in I$. The graph Γ_ε is clique-Helly.

We now prove that the triangle complex of Γ_ε is simply connected. Fix a combinatorial loop ℓ in Γ_ε . Since X is simply connected, there exists a disk D in X bounding ℓ . Consider a triangulation T of D such that triangles have diameter for d_X at most ε . Then the vertex set of each triangle of T is a clique in Γ_ε , therefore ℓ is null-homotopic in the triangle complex of Γ_ε . So the triangle complex of Γ_ε is simply connected.

The graph Γ_ε is clique-Helly and has a simply connected triangle complex, so according to Theorem 1.11, we deduce that Γ_ε is a Helly graph.

Note that, for any $\varepsilon > 0$, we have $d_X \leq \varepsilon d_{\Gamma_\varepsilon} \leq d_X + \varepsilon$, and balls for the metric d_{Γ_ε} with integral radius satisfy the Helly property. We deduce that, for any $\varepsilon > 0$, the metric space X is ε -coarsely injective. According to Lemma 1.12, we conclude that X is injective.

We now turn to the general case, when X is only semi-uniformly locally injective: there exists $\varepsilon > 0$ such that each ball of radius ε in X is uniformly locally injective. According to the uniformly locally injective case, we deduce that each ball of radius ε is injective: this means that X is uniformly locally injective. According to the uniformly locally injective case again, we deduce that X is injective. \square

We now see that, under a mild assumption on a simplicial complex with ordered simplices, saying that the ℓ^∞ orthoscheme realization is injective is equivalent to saying that the thickening of the 1-skeleton is Helly.

Theorem 1.14. *Let X denote a finite-dimensional simplicial complex with ordered simplices, and with maximal edges. Let Γ denote the graph with vertex set $X^{(0)}$, and with an edge between $x, y \in X^{(0)}$ if there exist $a, b \in X^{(0)}$ and ordered triangles $a \leq x \leq b$ and $a \leq y \leq b$ in X . If the ℓ^∞ orthoscheme complex of X is injective, then the thickening Γ of X is a Helly graph.*

Proof. We see that maximal cliques in Γ correspond to intervals $I_{ab} = \{x \in X^{(0)} \mid a \leq x \leq b \text{ is an ordered triangle in } X\}$, for any maximal edge $a \leq b$ in X . Let $m_{ab} \in |X|$ denote the midpoint of the maximal edge $a \leq b$. By assumption, any simplex of X containing a and b has for vertex set a chain from a to b . Then $B_{|X|}(m_{ab}, \frac{1}{2})$ is a subcomplex of $|X|$ with vertex set $B_{|X|}(m_{ab}, \frac{1}{2}) \cap X^{(0)} = I_{ab}$.

We will prove that Γ is clique-Helly: let $(I_{a_i b_i})_{i \in I}$ denote a family of pairwise intersecting maximal cliques in Γ . Since $|X|$ is injective, there exists $z \in \bigcap_{i \in I} B_{|X|}(m_{a_i b_i}, \frac{1}{2})$. Since each such ball is a subcomplex of $|X|$, we may assume that z is a vertex of X . We deduce that z belongs to each clique $I_{a_i b_i}$, for $i \in I$. So Γ is clique-Helly.

We will now prove that the triangle complex of Γ is simply connected. Let ℓ denote a combinatorial loop in the 1-skeleton of Γ . Up to homotopy in the triangle complex of Γ , we may assume that ℓ lies in the 1-skeleton of X . Since $|X|$ is injective, it is contractible, so the 2-skeleton of X is simply connected. As the 2-skeleton of X is contained in the triangle complex of Γ , we conclude that ℓ is null-homotopic in the triangle complex of Γ .

According to Theorem 1.11, we deduce that Γ is Helly. \square

2 The thickening of a lattice

We will explain a very simple construction of Helly graphs and injective metric spaces, starting with a lattice endowed with an action of the group \mathbb{Z} or \mathbb{R} .

Assume that L is a lattice, such that each upperly bounded subset of L has a join. Assume that there is an order-preserving, increasing, continuous (with respect to the order topology on L), action $(f_t)_{t \in H}$ of $H = \mathbb{Z}$ or \mathbb{R} on L , such that

$$\forall x, y \in L, \exists t \in H_+, f_{-t}(x) \leq y \leq f_t(x).$$

Let us define the following metric d on L :

$$\forall x, y \in L, d(x, y) = \inf\{t \in H_+ \mid f_{-t}(x) \leq y \leq f_t(x)\}.$$

Theorem 2.1. *If $H = \mathbb{Z}$, then (L, d) is a Helly graph with the combinatorial distance.*

If $H = \mathbb{R}$, then (L, d) is injective.

Proof. We start by proving that d is indeed a metric on L . If $x, y \in L$ and $t \in H_+$ are such that $f_{-t}(x) \leq y \leq f_t(x)$, then by applying f_t and f_{-t} we deduce that $f_{-t}(y) \leq x \leq f_t(y)$: the metric d is symmetric.

Let $x, y, z \in L$, and assume that $f_{-t}(x) \leq y \leq f_t(x)$ and $f_{-s}(y) \leq z \leq f_s(y)$ for some $t, s \in H_+$. Then $f_{-t-s}(x) \leq f_{-s}(y) \leq z \leq f_s(y) \leq f_{s+t}(x)$, so we deduce that $d(x, z) \leq d(x, y) + d(y, z)$.

Assume that $x, y \in L$ are distinct.

- Assume first that x, y are comparable, for instance $x < y$. Since $\{z \in L \mid z < y\}$ is open and contains x , by continuity of the action there exists $\varepsilon > 0$ such that, for any $|t| \leq \varepsilon$, we have $f_t(x) < y$. In particular $d(x, y) \geq \varepsilon$.
- Assume now that x, y are not comparable. Since $U = \{z \in L \mid x \wedge y < z < x \vee y\}$ is open and contains x , by continuity of the action there exists $\varepsilon > 0$ such that, for any $|t| \leq \varepsilon$, we have $f_t(x) \in U$. For any $0 \leq t < \varepsilon$, since $x \leq f_t(x) < x \vee y$, we have $f_t(x) \not\leq y$. In particular, $d(x, y) \geq \varepsilon$.

So we have proved that d is a metric on L .

We will first consider the case $H = \mathbb{Z}$. Let us define a graph \hat{L} with vertex set L , and with an edge between $x, y \in L$ if $f_{-1}(x) \leq y \leq f_1(x)$. We will first prove, by induction on $k \geq 0$, that for any $x \in L$, the ball $B_{\hat{L}}(x, k)$ in the graph \hat{L} coincides with the interval $I(f_{-k}(x), f_k(x)) = \{y \in L \mid f_{-k}(x) \leq y \leq f_k(x)\}$.

For $k \leq 1$ it is the definition of the edges of \hat{L} , so fix $k \geq 2$ and assume that the statement holds for $k-1$. Fix $y \in I(f_{-k}(x), f_k(x))$, we will prove that $y \in B_{\hat{L}}(x, k)$. Since $y \geq f_{-k}(x)$, we deduce that $f_1(y) \geq f_{-k+1}(x)$, and also since $y \leq f_k(x)$ we deduce that $f_{-1}(y) \leq f_{k-1}(x)$. So we have that both $f_1(y)$ and $f_{k-1}(x)$ are superior to both $f_{-1}(y)$ and $f_{-k+1}(x)$: since M is a lattice, there exists some element $z \in L$ in the intersection $I(f_{-1}(y), f_1(y)) \cap I(f_{-k+1}(x), f_{k-1}(x))$. In particular, y and z are adjacent in \hat{L} , and by induction we know that $d_{\hat{L}}(y, x) \leq k-1$, so $d_{\hat{L}}(x, z) \leq k$. Conversely, it is clear that the ball $B_{\hat{L}}(x, k)$ is included in the interval $I(f_{-k}(x), f_k(x))$. So we have $B_{\hat{L}}(x, k) = I(f_{-k}(x), f_k(x))$.

By assumption on the lattice L , we deduce that the graph \hat{L} is connected, and furthermore that $d_{\hat{L}} = d$.

Remark now that intervals in the lattice L satisfy the Helly property. Fix any collection $(I(x_i, y_i))_{i \in I}$ of pairwise intersecting intervals of L . Fix $j_0 \in I$. For any $i \in I$, we have $x_i \leq y_{j_0}$, so the set $\{x_i, i \in I\}$ is upperly bounded. By assumption, the set $\{x_i, i \in I\}$ has a join $z \in L$, such that $z \leq y_{j_0}$. This holds for any $j_0 \in I$, so z belongs to the intersection of all intervals $(I(x_i, y_i))_{i \in I}$.

Hence the graph \hat{L} is connected, and its balls satisfy the Helly property: it is a Helly graph.

We now turn to the case $H = \mathbb{R}$. First note that balls in (L, d) are intervals in L , so according to the previous argument, we know that balls in (L, d) satisfy the Helly property. We only have to prove that (L, d) is a geodesic metric space. It is enough to prove that (M, d) is weakly geodesic, i.e. for any $x, y \in L$ and any $0 \leq r \leq d(x, y)$, there exists $z \in B(x, r) \cap B(y, d(x, y) - r)$.

For each $k \in \mathbb{N} \setminus \{0\}$, let us consider the action of $\frac{1}{k}\mathbb{Z} \subset \mathbb{R}$ on L , and the associated Helly graph distance

$$\forall x, y \in L, d_k(x, y) = \inf \left\{ t \in \frac{1}{k}\mathbb{N} \mid f_{-t}(x) \leq y \leq f_t(x) \right\}.$$

Fix $x, y \in L$, and $0 \leq r \leq d(x, y)$. For each $k \in \mathbb{N} \setminus \{0\}$, there exists $z_k \in L$ such that $d_k(x, z_k) \leq r + \frac{1}{k}$ and $d_k(z_k, y) \leq d_k(x, y) - r + \frac{1}{k} \leq d(x, y) - r + \frac{2}{k}$. So the intervals $I_k = I(f_{-r-\frac{1}{k}}(x), f_{r+\frac{1}{k}}(x))$ and $J_k = I(f_{-d(x,y)+r-\frac{2}{k}}(y), f_{d(x,y)-r+\frac{2}{k}}(y))$ in L intersect. If $k \leq k'$, we know that $I_{k'} \subset I_k$ and $J_{k'} \subset J_k$. We deduce that the family of intervals $\{I_k\}_{k \in \mathbb{N} \setminus \{0\}} \cup \{J_k\}_{k \in \mathbb{N} \setminus \{0\}}$ pairwise intersects. By the Helly property for intervals, the global intersection is non-empty: let us denote z some element in the intersection.

We know that $\lim_{k \rightarrow +\infty} d_k(x, z) = d(x, z) \leq r$ and $\lim_{k \rightarrow +\infty} d_k(y, z) = d(y, z) \leq d(x, y) - r$, so we have proved that (L, d) is a weakly geodesic metric space. Since balls in (L, d) satisfy the Helly property, we conclude that (L, d) is injective. \square

An immediate consequence concerns Garside groups. Recall that a group G is called *Garside* if there exists a subset $S \subset G$ and an element $\delta \in G$ such that the following hold:

- S spans the group G .
- For each element in the semigroup $\langle S \rangle^+$, there is a bound on the length of factorizations over S .
- The element δ belongs to the semigroup $\langle S \rangle^+$, and δ is *balanced*: the set of prefixes of δ coincide with the set of suffixes of δ .
- The poset of prefixes of δ in $\langle S \rangle^+$ is a lattice.

Note that G is endowed with two natural orders, the left order \leq_L and the right order \leq_R :

- $g \leq_L h$ if and only if $g^{-1}h \in \langle S \rangle^+$.
- $g \leq_R h$ if and only if $hg^{-1} \in \langle S \rangle^+$.

Authors sometimes add the requirement that S is finite, in which case G may also be called a Garside group of finite type.

Fix a Garside group (G, S, δ) . Let X denote the graph with vertex set G , with an edge between $g, h \in G$ if $g\delta^{-1} \leq_L h \leq_L g\delta$. The graph X is called the *thickening* of G .

Corollary 2.2. *The thickening of any Garside group is a Helly graph.*

Proof. The left order on G is a lattice order (see [Deh15]). Furthermore, consider the action of \mathbb{Z} on G by right multiplication by δ . For any $g \in G$, there exists $n \in \mathbb{N}$ such that $\delta^{-n} \leq_L g \leq_L \delta^n$. So this action satisfies the assumptions of Theorem 2.1: we deduce that the graph X is Helly. \square

Note that this applies, in particular, to Garside groups of infinite type, such as crystallographic Garside groups (see [MS17]). However, we do not have yet an application of this simply transitive action of a Garside group on a locally infinite Helly graph.

In the case of Garside groups of finite type, we recover a particularly simple proof of the following result by Huang and Osajda (see [HO19]). In particular, our proof does not rely on the deep result that local-to-global result for Helly graphs (Theorem 1.11, see [CCHO21]).

Corollary 2.3 (Huang-Osajda). *Any Garside group of finite type is a Helly group.*

In particular, this leads to a particularly simple proof that braids groups are Helly, relying only on some Garside structure.

3 The affine version of a lattice

In this section, we will prove that the orthoscheme complex of a bounded graded lattice, endowed with the orthoscheme ℓ^∞ metric, is injective. In order to do so, we will apply results from the previous section, and endow the geometric realization of a lattice with a partial order, which is a lattice.

Assume that L is a bounded, graded lattice of rank n . Let \leq_L denote the order on L . Let H denote either a cyclic subgroup of $(\mathbb{R}, +)$, or $H = \mathbb{R}$. We will define a new poset M_H , which will be called the *affine version* of L over H . If there is no ambiguity about H , we will simply denote $M = M_H$. Let $C(L)$ denote the set of maximal chains $c_{0,1} = 0 <_L c_{1,2} <_L \dots <_L c_{n-1,n} <_L c_{n,n+1} = 1$ in L . We will use the convention that the element denoted $c_{i,i+1}$ has rank i .

Let us consider the subspace

$$\sigma = \{u \in H^n \mid u_1 \leq u_2 \leq \dots \leq u_n\}$$

of H^n .

For each maximal chain $c \in C(L)$, let σ_c denote a copy of σ .

Let us consider the space

$$M = \bigcup_{c \in C(L)} \sigma_c / \sim,$$

where for each $c, c' \in C(L)$, if we denote $I = \{1 \leq i \leq n-1 \mid c_{i,i+1} \neq c'_{i,i+1}\}$, we identify σ_c and $\sigma_{c'}$ along the subspaces

$$\{u \in \sigma_c \mid \forall i \in I, u_i = u_{i+1}\} \simeq \{u \in \sigma_{c'} \mid \forall i \in I, u_i = u_{i+1}\}.$$

We can describe elements of M as a quotient of the space $M_0 = C(L) \times \sigma$.

Example. One illustrating example is the following: consider the boolean lattice L of rank n , i.e. the lattice of subsets of the finite set $\{1, \dots, n\}$, with the inclusion order. Maximal chains in L correspond to permutations of $\{1, \dots, n\}$. The space M_H may be identified with H^n , where for each permutation w of $\{1, \dots, n\}$, the subspace σ_w is

$$\sigma_w = \{x \in H^n \mid x_{w(1)} \leq x_{w(2)} \leq \dots \leq x_{w(n)}\}.$$

If $c \in C(L)$ and $u \in \sigma$, let us denote by $[c, u]$ the equivalence class of $(c, u) \in M_0$ in M .

For each $c \in C(L)$, let us endow σ_c with the partial order from $H^n \subset \mathbb{R}^n$: $u \leq v$ if $\forall 1 \leq i \leq n, u_i \leq v_i$. Let us endow M with the induced partial order: we have $\alpha \leq \beta$ in M if there exists a sequence $\alpha_0 = \alpha, \alpha_1, \dots, \alpha_m = \beta$ in M such that, for each $0 \leq i \leq m-1$, there exists $c \in C(L)$ and $x \leq y$ in σ_c such that $\alpha_i = [c, x]$ and $\alpha_{i+1} = [c, y]$.

Let \leq_M denote the order on M . We will prove the following.

Theorem 3.1. *If H is a discrete subgroup of \mathbb{R} , the poset M_H is a lattice.*

Before proving Theorem 3.1, we will gather some preliminary results. Without loss of generality, assume that $H = \mathbb{Z}$. To simplify notations, we will let $M = M_H$.

First notice that if $(a, u) \sim (b, v)$, then $u = v$. Therefore the second projection $M_0 \rightarrow \sigma$ defines a projection $\pi : M \rightarrow \sigma$. Fix $\beta = [b, v], \gamma = [c, w] \in M$, and fix $\alpha \leq_M \beta, \gamma$ in M . We will prove that β, γ have a join, and α will play an auxiliary role.

We say that β is *elementarily superior* to α if there are representatives $\alpha = [a, u]$ and $\beta = [b, v]$ such that $a = b$ and there exist $1 \leq i \leq j \leq n$ such that:

- $u_i = u_{i+1} = \dots = u_j$,
- $v_i = v_{i+1} = \dots = v_j = u_j + 1$, and
- $\forall k \notin [i, j], u_k = v_k$.

Lemma 3.2. Fix $\alpha = [a, u] \in M$. Any element $\beta = [b, v]$ of M elementarily superior to α is uniquely determined by:

- Integers $1 \leq i \leq j \leq n$, such that $u_j < u_{j+1}$ if $j < n$.
- Some element $b_{i-1, i}$ of rank $i - 1$ in the interval $I(a_{i_0-1, i_0}, a_{j, j+1})$, where $i_0 \in [1, i]$ is minimal such that $u_{i_0} = u_i$.

Let us denote $\beta = \alpha[i, j, b_{i-1, i}]$.

Proof. Let us define the element $\beta = [b, v]$ in M by:

$$\begin{aligned} \forall k \notin [i, j], v_k &= u_k \\ \forall k \in [i, j], v_k &= u_j + 1 \\ \forall k \leq i_0 - 1, b_{k, k+1} &= a_{k, k+1} \\ \forall k \geq j, b_{k, k+1} &= a_{k, k+1}. \end{aligned}$$

Note that v is non-decreasing, hence $v \in \sigma$. Furthermore, since $v_{i_0} = v_{i_0+1} = \dots = v_{i-1}$ and $v_i = v_{i+1} = \dots = u_j + 1 \leq v_{j+1}$, it is enough to define $b_{k, k+1}$ for $k \leq i_0 - 1$, $k = i - 1$ and $k \geq j$. Hence β is well-defined, and it is elementarily superior to α . It is clear that β is the only such element in M . \square

Lemma 3.3. Given $\alpha \leq_M \beta$ in M , there exist $m \geq 0$ and a sequence $\beta_0 = \alpha, \beta_1, \dots, \beta_m = \beta$ for which, for each $0 \leq i \leq m - 1$, β_{i+1} is elementarily superior to β_i .

Proof. According to the definition of the order on M , it is sufficient to prove the statement for $\alpha = [a, u]$ and $\beta = [b, v]$ such that $a = b$. We have $u \leq v$ in $\mathbb{Z}^n \subset \mathbb{R}^n$. Then consider a sequence $u^0 = u \leq u^1 \leq \dots \leq u^m = v$ such that, for each $0 \leq k \leq m - 1$, there exist $1 \leq i_k \leq j_k \leq n$ such that:

- $u_{i_k}^k = u_{i_k+1}^k = \dots = u_{j_k}^k$,
- $u_{i_k}^{k+1} = u_{i_k+1}^{k+1} = \dots = u_{j_k}^{k+1} = u_{j_k}^k + 1$, and
- $\forall \ell \notin [i_k, j_k], u_\ell^k = u_\ell^{k+1}$.

For each $0 \leq k \leq m - 1$, the element $\beta_{k+1} = [a, u^{k+1}]$ is elementarily superior to $\beta_k = [a, u^k]$. \square

Lemma 3.4. We have $\alpha = [a, u] \leq \beta = [b, v]$ in M if and only if $u \leq v$ and for every $0 \leq j \leq n - 1$ such that $u_j < u_{j+1}$, if we denote by $i \in \{0, \dots, j\}$ the minimal element such that $v_{i+1} \geq u_{j+1}$, we have $b_{i, i+1} \leq_L a_{j, j+1}$.

Proof. Let us denote by \prec this relation. We first show that \prec is transitive and antisymmetric.

Assume that $\alpha = [a, u] \preceq \beta = [b, v]$ and $u = v$, we will prove that $\alpha = \beta$. Fix $0 \leq j \leq n - 1$, and assume that $a_{j, j+1} \neq b_{j, j+1}$. Since these two elements of L have the same rank j , we deduce that they are not comparable: $b_{j, j+1} \not\leq_L a_{j, j+1}$ and $a_{j, j+1} \not\leq_L b_{j, j+1}$. We will prove that $u_j = u_{j+1}$: by contradiction, if $u_j < u_{j+1}$, then the minimal $i \in \{0, \dots, j\}$ such that $v_{i+1} \geq u_{j+1}$ is equal to j , so $b_{j, j+1} \leq_L a_{i, i+1}$. So $u_j = u_{j+1}$. This proves that $\alpha = \beta$.

This implies in particular that \prec is reflexive: indeed assume that $\alpha \preceq \beta$ and $\beta \preceq \alpha$. Then $u \leq v$ and $v \leq u$, so $u = v$, hence $\alpha = \beta$.

We now prove that \prec is transitive: assume that $\alpha = [a, u] \preceq \beta = [b, v] \preceq \gamma = [c, w]$. Then $u \leq v \leq w$, so $u \leq w$. Let $0 \leq k \leq n-1$ such that $u_k < u_{k+1}$, let $i \in \{0, \dots, k\}$ be minimal such that $w_{i+1} \geq u_{k+1}$: we want to prove that $c_{i,i+1} \leq_L a_{k,k+1}$. Let $j \in \{0, \dots, k\}$ be minimal such that $v_{j+1} \geq u_{k+1}$: we know that $b_{j,j+1} \leq_L a_{k,k+1}$. Since j is minimal, we know that either $j = 0$ or $v_j < u_{k+1}$.

If $j = 0$, then since $i \leq j$ we know that $i = 0$ and so $c_{i,i+1} = 0 = b_{j,j+1} \leq_L a_{k,k+1}$.

If $v_j < u_{k+1}$ then $v_j < v_{j+1}$, and let $i' \in \{0, \dots, j\}$ be minimal such that $w_{i'+1} \geq v_{j+1}$. We know that $c_{i',i'+1} \leq_L b_{j,j+1}$. But since $w_i < u_{k+1} \leq v_{j+1}$, we have $i \leq i'$, so $c_{i,i+1} \leq_L c_{i',i'+1} \leq_L b_{j,j+1} \leq_L a_{k,k+1}$.

Hence $\alpha \preceq \gamma$.

We will now prove that if $\beta = [b, v]$ is elementarily superior to $\alpha = [a, u]$, we have $\alpha \preceq \beta$. Let us denote $\beta = \alpha[i, j, b_{i-1,i}]$. First note that $u \leq v$. If $0 \leq k \leq n-1$ is such that $u_k < u_{k+1}$, then either $k \leq i-1$ or $k \geq j$. Let $\ell \in \{0, \dots, k\}$ denote the minimal element such that $v_{\ell+1} \geq u_{k+1}$, we will prove that $b_{\ell,\ell+1} \leq_L a_{k,k+1}$.

- If $k \leq i-1$, then $\ell = k$ and $b_{\ell,\ell+1} = a_{k,k+1}$.
- If $k \geq j$ and $v_{k+1} = v_j$, then $\ell = i-1$ and $b_{\ell,\ell+1} = b_{i-1,i} \leq_L a_{j,j+1} \leq_L a_{k,k+1}$.
- If $k \geq j$ and $v_{k+1} > v_j$, then $\ell \geq j$ and $b_{\ell,\ell+1} = a_{\ell,\ell+1} \leq_L a_{k,k+1}$.

According to Lemma 3.3, we deduce that if $\alpha \leq_M \beta$, then $\alpha \preceq \beta$.

We will now prove that if $\alpha = [a, u] \preceq \beta = [b, v]$ with $\alpha \neq \beta$, there exists γ elementarily superior to α such that $\gamma \preceq \beta$. If $u = v$, we have seen that $\alpha = \beta$, so let us assume that $u < v$. Let $1 \leq i \leq n$ be minimal such that $u_i < v_i$. Let $j \in \{i, \dots, n\}$ be maximal such that $u_i = u_j$.

Let $i_0 \in \{1, \dots, i\}$ be minimal such that $u_{i_0} = u_i$. Since for all $k \leq i-1$ we have $u_k = v_k$, there exist representatives $\alpha = [a, u]$ and $\beta = [b, v]$ such that, for any $1 \leq k \leq i-1$, we have $a_{k-1,k} = b_{k-1,k}$.

We will also prove that we may assume that $a_{i-1,i} = b_{i-1,i}$.

Assume first that $u_{i-1} < u_i$. Let $p \in \{0, \dots, i-1\}$ be minimal such that $v_{p+1} \geq u_i$. Since $v_{i-1} = u_{i-1} < u_i$, we deduce that $p = i-1$. Since $\alpha \preceq \beta$, we have $b_{i-1,i} \leq_L a_{i-1,i}$. As both elements have the same rank $i-1$ in L , we have $a_{i-1,i} = b_{i-1,i}$.

Assume now that $u_{i-1} = u_i$. Then $u_j < u_{j+1}$: let $p \in \{0, \dots, j\}$ be minimal such that $v_{p+1} \geq u_{j+1}$. Since $v_{i-1} = u_{i-1} < u_{j+1}$, we deduce that $p \geq i-1$. Since $\alpha \preceq \beta$, we have $b_{i-1,i} \leq_L b_{p,p+1} \leq_L a_{j,j+1}$. Since $u_{i-1} = u_i = \dots = u_j$, we may choose a representative of α such that $a_{i-1,i} = b_{i-1,i}$.

In particular $b_{i-1,i} \geq_L b_{i_0-1,i_0} = a_{i_0-1,i_0}$. Since $u_j < u_{j+1}$ and $\alpha \preceq \beta$, if we denote by $i' \in \{0, \dots, j\}$ the minimal element such that $v_{i'+1} \geq u_{j+1}$, we have $b_{i',i'+1} \leq_L a_{j,j+1}$. Since $v_{i-1} = u_{i-1} < u_{j+1}$, we have $i' > i-2$, so $b_{i-1,i} \leq_L b_{i',i'+1} \leq_L a_{j,j+1}$.

So we know that $b_{i-1,i} \in I(a_{i_0-1,i_0}, a_{j,j+1})$: we can define $\gamma = \alpha[i, j, b_{i-1,i}] = [w, c]$. We will now prove that $\gamma \preceq \beta$. Fix $0 \leq k \leq n-1$ such that $w_k < w_{k+1}$, and denote by $\ell \in \{0, \dots, k\}$ the minimal element such that $v_{\ell+1} \geq w_{k+1}$, we will prove that $b_{\ell,\ell+1} \leq_L c_{k,k+1}$. Recall that either $k \leq i-1$ or $k \geq j$.

- If $k \leq i-2$, then $w_{k+1} = u_{k+1}$. Since $\alpha \preceq \beta$, we know that $b_{\ell,\ell+1} \leq_L a_{k,k+1} = c_{k,k+1}$.
- If $k = i-1$, then $w_{k+1} = w_i = u_i + 1 > u_i$. Then $v_{i-1} = u_{i-1} \leq u_i < w_{k+1}$, so $\ell = i-1 = k$. Hence $b_{\ell,\ell+1} = b_{i-1,i} = a_{i-1,i} = c_{i-1,i} = c_{k,k+1}$.

- If $k \geq j$, then $w_{k+1} = u_{k+1}$. Since $\alpha \preceq \beta$, we know that $b_{\ell, \ell+1} \leq a_{k, k+1}$. We also have $a_{k, k+1} = c_{k, k+1}$, so $b_{\ell, \ell+1} \leq c_{k, k+1}$.

So we conclude that $\gamma \preceq \beta$.

Now fix any $\alpha = [a, u] \preceq \beta = [b, v]$. By induction on $\|v - u\|_1$, we see that there is a bound on sequences of elementarily superior elements starting from α which all are $\preceq \beta$. Therefore we conclude that $\alpha \leq_M \beta$.

In conclusion, the two orders \leq_M and \prec coincide. \square

Given $\alpha \leq_M \beta$ in M , let us denote by $D(\alpha, \beta)$ the minimal number $m \geq 0$ such that there exists a sequence $\beta_0 = \alpha, \beta_1, \dots, \beta_m = \beta$ for which, for each $0 \leq i \leq m - 1$, β_{i+1} is elementarily superior to β_i .

We will prove, by induction on $D(\alpha, \beta) + D(\alpha, \gamma)$, that β and γ have a join.

Lemma 3.5. *Assume that $\alpha, \beta, \gamma \in M$ are such that $\alpha \leq_M \beta$, $\alpha \leq_M \gamma$ and $D(\alpha, \beta) = D(\alpha, \gamma) = 1$. Then β, γ have a join δ such that $D(\beta, \delta) \leq 1$ and $D(\gamma, \delta) \leq 1$.*

Proof. Consider representatives $\alpha = [a, u]$, $\beta = [b, v]$ and $\gamma = [c, w]$ of α , β and γ respectively. According to Lemma???, there exist $1 \leq i \leq j \leq n$ and $b_{i, i+1}$ such that $\beta = \alpha[i, j, b_{i, i+1}]$, and there exist $1 \leq i' \leq j' \leq n$ and $c_{i', i'+1}$ such that $\gamma = \alpha[i', j', c_{i', i'+1}]$.

First case: assume that the intervals $[i, j]$ and $[i', j']$ are disjoint, for instance $j < i'$. Let us define $\delta = \beta[i', j', c_{i', i'+1}]$: we will see that δ is well-defined and that $\delta = \gamma[i, j, b_{i, i+1}]$.

First note that $v_{j'} = u_{j'} < u_{j'+1} = v_{j'+1}$. Furthermore, let $i'_0 \in [1, i']$ be minimal such that $u_{i'_0} = u_{i'}$. Since $u_j < u_{j+1}$, we know that $j + 1 \leq i'_0$. So we have $v_{i'_0} = u_{i'_0} = u_{i'} = v_{i'}$. So, if we denote $i''_0 \in [1, i']$ the minimal integer such that $v_{i''_0} = v_{i'}$, we have $i''_0 \leq i'_0$.

By definition we have $c_{i', i'+1} \in I(a_{i'_0-1, i'_0}, a_{j', j'+1})$. Since $j < j'$, we have $b_{j', j'+1} = a_{j', j'+1}$, so $c_{i', i'+1} \leq_L b_{j', j'+1}$. And as $b_{i''_0-1, i''_0} \leq_L b_{i'_0-1, i'_0} = a_{i'_0-1, i'_0}$, we deduce that $c_{i', i'+1} \geq a_{i'_0-1, i'_0} \geq b_{i''_0-1, i''_0}$. So we have $c_{i', i'+1} \in I(b_{i''_0-1, i''_0}, b_{j', j'+1})$. Hence $\delta = \beta[i', j', c_{i', i'+1}]$ is well-defined, and it is elementarily superior to β .

Following the same argument, the element $\delta' = \gamma[i, j, b_{i, i+1}]$ is well-defined. We will prove that $\delta = \delta'$. Let $i_0 \in [1, i]$ be minimal such that $u_{i_0} = u_i$. According to the proof of Lemma 3.2, we can see that $\delta = \delta' = [d, x]$ are explicitly equal to the following:

$$\begin{aligned}
\forall k \notin [i, j] \cup [i', j'], x_k &= u_k \\
\forall k \in [i, j], x_k &= u_j + 1 \\
\forall k \in [i', j'], x_k &= u_{j'} + 1 \\
\forall k \leq i_0 - 1, d_{k, k+1} &= a_{k, k+1} \\
d_{i, i+1} &= b_{i, i+1} \\
\forall k \in [j, i'_0 - 1], b_{k, k+1} &= a_{k, k+1} \\
d_{i', i'+1} &= c_{i', i'+1} \\
\forall k \geq j', b_{k, k+1} &= a_{k, k+1}.
\end{aligned}$$

So the element $\delta = \delta'$ is elementarily superior to both β and γ .

We will now prove that δ is the minimal element of M superior to both β and γ . Fix $\theta = [e, y] \in M$ any element superior to both β and γ , we will prove that $\delta \leq \theta$. Since $y \geq v$ and $y \geq w$, we deduce that $y \geq x$. Fix any $0 \leq k \leq n - 1$ such that $x_k < x_{k+1}$, and let $\ell \in \{0, \dots, k\}$ denote the minimal element such that $y_{\ell+1} \geq x_{k+1}$. We will prove that $e_{\ell, \ell+1} \leq_L d_{k, k+1}$.

- Assume that $k \leq j-1$. Then $x_{k+1} = v_{k+1}$, and since $\beta \leq \theta$, we deduce by Lemma 3.4 that $e_{\ell, \ell+1} \leq_L b_{k, k+1} = d_{k, k+1}$.
- Assume that $k \geq j$. Then $x_{k+1} = w_{k+1}$, and since $\gamma \leq \theta$, we deduce by Lemma 3.4 that $e_{\ell, \ell+1} \leq_L c_{k, k+1} = d_{k, k+1}$.

According to Lemma 3.4, we deduce that $\delta \leq_M \theta$: δ is the minimal element of M superior to both β and γ . Hence $\delta = \beta \vee_M \gamma$. Furthermore, we have noticed that δ is elementarily superior to both β and γ , so $D(\beta, \delta) = D(\gamma, \delta) = 1$.

Second case: assume now that the intervals $[i, j]$ and $[i', j']$ intersect. Without loss of generality, assume that $i \leq i'$. Since $u_j < u_{j+1}$, we deduce that $j = j'$. Let $1 \leq i_0 \leq n$ be minimal such that $u_{i_0} = u_i$. The elements $b_{i-1, i}$ and $c_{i'-1, i'}$ both belong to the interval $I(a_{i_0-1, i_0}, a_{j, j+1})$.

If $b_{i-1, i} = c_{i'-1, i'}$, then $\beta = \gamma$ and they have an trivial join $\delta = \beta = \gamma$. So we may assume that $b_{i-1, i} \neq c_{i'-1, i'}$.

Consider the meet $g = b_{i-1, i} \wedge_L c_{i'-1, i'} \in L$: its rank $r-1$ is such that $i_0 \leq r < i, i'$. Let us define $\delta = \alpha[r, j, g] \in M$. We see that $\delta = \beta[r, i-1, g] = \gamma[r, i'-1, g]$, so δ is elementarily superior to β and γ .

We will now prove that $\delta = [d, x]$ is the minimal element of M superior to both β and γ . Fix $\theta = [e, y] \in M$ any element superior to both β and γ , we will prove that $\delta \leq \theta$.

We will first prove that $x \leq y$. Since $\beta, \gamma \leq \theta$, we deduce that $v, w \leq y$. In particular, for any $m < r$ or $m \geq i$, we have $y_m \geq b_m = x_m$. And for $r \leq m \leq i-1$, we have $y_m \geq a_m = x_m - 1$. Assume by contradiction that there exists $m \in \{r, \dots, i-1\}$ such that $y_m = x_m - 1$, and choose such m maximal. Since $x_r = x_{r+1} = \dots = x_j$, we have $y_{m+1} \geq x_{m+1} = x_i = v_i = x_{i'} = w_{i'}$. Since $\beta \leq_L \theta$ and $\gamma \leq_L \theta$, according to Lemma 3.4, we know that $e_{m, m+1} \leq_L b_{i-1, i}$ and $e_{m, m+1} \leq_L c_{i'-1, i'}$. In particular, we deduce that $e_{m, m+1} \leq_L b_{i-1, i} \wedge_L c_{i'-1, i'} = g$. Note that the rank of $e_{m, m+1}$ is m , whereas the rank of g is $r-1$. Since $m > r-1$, this is a contradiction. Hence $x \leq y$.

Fix any $0 \leq k \leq n-1$ such that $x_k < x_{k+1}$, and let $\ell \in \{0, \dots, k\}$ denote the minimal element such that $y_{\ell+1} \geq x_{k+1}$. We will prove that $e_{\ell, \ell+1} \leq_L d_{k, k+1}$.

- Assume that $k \leq r-2$. Then $x_{k+1} = v_{k+1}$, and since $\beta \leq \theta$ we deduce by Lemma 3.4 that $e_{\ell, \ell+1} \leq_L b_{k, k+1} = d_{k, k+1}$.
- Assume that $r-1 \leq k \leq j-1$. Then $x_{k+1} = v_i = w_{i'}$. Since $\beta \leq \theta$, we deduce by Lemma 3.4 that $e_{\ell, \ell+1} \leq_L b_{i-1, i}$. And since $\gamma \leq \theta$, we also deduce that $e_{\ell, \ell+1} \leq_L c_{i'-1, i'}$. Hence $e_{\ell, \ell+1} \leq_L b_{i-1, i} \wedge_L c_{i'-1, i'} = g = d_{r-1, r} \leq_L d_{k, k+1}$.
- Assume that $k \geq j$. Then $x_{k+1} = v_{k+1}$, and since $\beta \leq \theta$ we deduce by Lemma 3.4 that $e_{\ell, \ell+1} \leq_L b_{k, k+1} = d_{k, k+1}$.

According to Lemma 3.4, we deduce that $\delta \leq_M \theta$: δ is the minimal element of M superior to both β and γ . Hence $\delta = \beta \vee_M \gamma$. Furthermore, we have noticed that δ is elementarily superior to both β and γ , so $D(\beta, \delta) = D(\gamma, \delta) = 1$. \square

Lemma 3.6. *Assume that $\alpha, \beta, \gamma \in M$ are such that $\alpha \leq_M \beta$, $\alpha \leq_M \gamma$, $D(\alpha, \beta) = m$ and $D(\alpha, \gamma) = m'$ for some $m, m' \in \mathbb{N}$. Then β, γ have a meet δ such that $D(\beta, \delta) \leq m'$ and $D(\gamma, \delta) \leq m$.*

Proof. We proceed by induction on $m + m'$: when $m + m' \leq 2$, the statement holds by Lemma 3.5. Now fix $k \geq 3$, and assume that the statement holds when $m + m' < k$. Fix m, m' such that $m + m' = k$, and without loss of generality assume that $m \geq 2$. Choose

$\beta_0 = \alpha, \beta_1, \dots, \beta_m = \beta$ an elementary sequence from α to β , with $m = D(\alpha, \beta)$. We have $D(\alpha, \beta_1) + D(\alpha, \gamma) = 1 + m' < k$, so by induction there exists $\delta' = \beta_1 \vee_M \gamma$ with $D(\beta_1, \delta') \leq m'$ and $D(\gamma, \delta') \leq 1$. Since $D(\beta_1, \beta) + D(\beta_1, \delta') = m - 1 + m' < k$, by induction there exists $\delta = \beta \vee_M \delta'$ such that $D(\beta, \delta) \leq m'$ and $D(\delta', \delta) \leq m - 1$. So we deduce that $D(\gamma, \delta) \leq m$.

We will now prove that δ is the meet of β and γ . We have $\delta \geq_M \beta$ and $\delta \geq_M \delta' \geq_M \gamma$. Furthermore, consider any $\theta \in M$ such that $\theta \geq_M \beta$ and $\theta \geq_M \gamma$. As $\beta \geq_M \beta_1$, we deduce that $\theta \geq_M \beta_1 \vee_M \gamma = \delta'$. And we deduce that $\theta \geq_M \beta \vee_M \delta' = \delta$. So we have proved that $\delta = \beta \vee_M \gamma$. \square

Proof. [of Theorem 3.1] Fix any $\beta = [b, v], \gamma = [c, w] \in M$. Let $k \geq 0$ such that $v_n - k < w_1$. Let $u = (v_1, k, v_2 - k, \dots, v_n - k)$. Then $\alpha = [b, u] \in M$ is inferior to both β and γ , so we can apply Lemma 3.6 to deduce that β and γ have a meet in M . By symmetry of the construction, β and γ also have a join in M . So M is a lattice. \square

If $H = \mathbb{R}$, the affine version $M_{\mathbb{R}}$ of L over \mathbb{R} is a gluing of subspaces $\sigma \subset \mathbb{R}^n$. We may therefore endow $M_{\mathbb{R}}$ with the piecewise length metric $d_{\mathbb{R}}$ induced by the standard ℓ^∞ metric on each $\sigma \subset \mathbb{R}^n$.

Theorem 3.7. *If $H = \mathbb{R}$, the poset $M_{\mathbb{R}}$ is a lattice, and the metric space $(M_{\mathbb{R}}, d_{\mathbb{R}})$ is injective.*

Proof. Remark that, for each $\theta > 0$, the space $M_{\theta\mathbb{Z}}$ may be realized naturally as a closed subspace of $M_{\mathbb{R}}$. Furthermore, the sequence of closed subsets $M_{\theta\mathbb{Z}}$ of $M_{\mathbb{R}}$ converges to $M_{\mathbb{R}}$ as $\theta \rightarrow 0$. We will use this convergence to prove that $M_{\mathbb{R}}$ is a lattice.

Fix $\alpha, \beta \in M_{\mathbb{R}}$, and consider maximal chains a, b in L such that $\alpha \in \sigma_a$ and $\beta \in \sigma_b$. For any $\theta > 0$, note that for any $x \in \mathbb{R}$, there exists $x_\theta \in \theta\mathbb{Z}$ such that $x_\theta - \theta \leq x \leq x_\theta + \theta$. So we may consider $\alpha_\theta \in M_{\theta\mathbb{Z}} \cap \sigma_a$ and $\beta_\theta \in M_{\theta\mathbb{Z}} \cap \sigma_b$ such that $(-\theta) \cdot \alpha_\theta \leq_M \alpha \leq_M \theta \cdot \alpha_\theta$, and similarly $(-\theta) \cdot \beta_\theta \leq_M \beta \leq_M \theta \cdot \beta_\theta$.

According to Theorem 3.1, the poset $M_{\theta\mathbb{Z}}$ is a lattice: consider $\gamma_\theta = \alpha_\theta \vee_{M_{\theta\mathbb{Z}}} \beta_\theta$. Let $C_{a,b} \subset L$ denote the smallest subset of L containing a, b , and which is stable under meet. Since L is bounded and graded, $C_{a,b}$ is finite. According to the proof of Theorem 3.1, we see that, for every $\theta > 0$, there exists a maximal chain $c_\theta \subset C_{a,b}$ such that $\gamma_\theta \in \sigma_{c_\theta}$. Since $C_{a,b}$ is finite, σ is locally compact and $(\gamma_\theta)_{\theta > 0}$ is bounded, there exists a sequence $\theta_k \xrightarrow[k \rightarrow +\infty]{} 0$ such that the sequence $(\gamma_{\theta_k})_{k \in \mathbb{N}}$ converges to some $\gamma \in M_{\mathbb{R}}$. Since $\gamma_\theta \geq_M \alpha_\theta \geq_M (\theta) \cdot \alpha$, we deduce that $\gamma \geq_M \alpha$. Similarly $\gamma \geq_M \beta$.

Assume that there exists $\delta \in M_{\mathbb{R}}$ such that $\alpha, \beta \leq_M \delta$. We will prove that $\gamma \leq_M \delta$. For any $\theta > 0$, fix $\delta_\theta \in M_{\theta\mathbb{Z}}$ such that $(-\theta) \cdot \delta_\theta \leq_M \delta \leq_M \theta \cdot \delta_\theta$. In particular $(2\theta) \cdot \delta_\theta \geq_M \theta \cdot \delta \geq_M \theta \cdot \alpha \geq_M \alpha_\theta$, and similarly $(2\theta) \cdot \delta_\theta \geq_M \beta_\theta$. We deduce that $(2\theta) \cdot \delta_\theta \geq_M \alpha_\theta \vee_{M_{\theta\mathbb{Z}}} \beta_\theta = \gamma_\theta$. Considering the limit along $(\theta_k)_{k \in \mathbb{N}}$ as $k \rightarrow +\infty$, we deduce that $\delta \geq_M \gamma$.

So we have proved that α and β have a meet in $M_{\mathbb{R}}$. By symmetry of the construction, they also have a join, so $M_{\mathbb{R}}$ is a lattice.

According to Theorem 2.1, we know that the metric space $(M_{\mathbb{R}}, d)$ is injective, with respect to the metric

$$\forall x, y \in M_{\mathbb{R}}, d(x, y) = \inf\{t \geq 0 \mid (-t) \cdot x \leq y \leq t \cdot x\}.$$

Note that the metric d is geodesic, and it restricts on each $\sigma_c \subset M_{\mathbb{R}}$, for $c \in C(L)$, to the natural ℓ^∞ metric on $\sigma_c \subset \mathbb{R}^n$. Therefore d coincides with the length metric $d_{\mathbb{R}}$.

So we conclude that $(M_{\mathbb{R}}, d_{\mathbb{R}})$ is injective. \square

We have seen in the introduction that the existence of a bicombing may be extremely useful, notably in the case of Deligne complexes of Artin groups. Let us recall that a *geodesic bicombing* on a metric space X is a map $\sigma : X \times X \times [0, 1] \rightarrow X$ such that, for all $x, y \in X$, the map $t \in [0, 1] \mapsto \sigma(x, y, t)$ is a constant speed geodesic from x to y .

The bicombing σ is called:

- *reversible* if $\forall x, y \in X, \forall t \in [0, 1], \sigma(x, y, t) = \sigma(y, x, 1 - t)$,
- *consistent* if $\forall x, y \in X, \forall s, t \in [0, 1], \sigma(x, \sigma(x, y, t), s) = \sigma(x, y, st)$ and
- *convex* if $\forall x, x', y, y' \in X$, the map $t \in [0, 1] \mapsto d(\gamma(x, y, t), \gamma(x', y', t))$ is convex.

Theorem 3.8. *The metric space $(M_{\mathbb{R}}, d_{\mathbb{R}})$ has a convex reversible consistent geodesic bicombing.*

Proof. Fix $x, y \in M_{\mathbb{R}}$ and $t \in [0, 1]$, and denote $D = d(x, y)$. Let us define

$$\sigma(x, y, t) = (tD) \cdot x \wedge ((1 - t)D) \cdot y.$$

We will first prove that σ is a geodesic bicombing, i.e. that $d(x, \sigma(x, y, t)) = tD$. First note that $\sigma(x, y, t) \leq (tD) \cdot x$. Furthermore, since $D \cdot y \geq x$, we have $((1 - t)D) \cdot y \geq (-tD) \cdot x$, hence $\sigma(x, y, t) \geq (-tD) \cdot x$. We conclude that $d(x, \sigma(x, y, t)) \leq tD$. By symmetry, we also have $d(y, \sigma(x, y, t)) \leq (1 - t)D$. According to the triangular inequality, we have $d(x, \sigma(x, y, t)) = tD$. Hence σ is a reversible geodesic bicombing on $M_{\mathbb{R}}$.

We will now prove that σ is consistent: fix $x, y \in M_{\mathbb{R}}$ and denote $D = d(x, y)$. Fix $t \in (0, 1)$, and let $z = \sigma(x, y, t) = (tD) \cdot x \wedge ((1 - t)D) \cdot y$. Fix $s \in [0, 1]$, we will prove that $\sigma(x, z, s) = \sigma(x, y, st)$. Since $d(x, z) = tD$, by definition of σ we have

$$\begin{aligned} \sigma(x, z, s) &= (stD) \cdot x \wedge ((1 - s)tD) \cdot z \\ &= (stD) \cdot x \wedge ((1 - s)tD) \cdot ((tD) \cdot x \wedge ((1 - t)D) \cdot y) \\ &= (stD) \cdot x \wedge ((1 - s)tD + tD) \cdot x \wedge ((1 - st)D) \cdot y \\ &= (stD) \cdot x \wedge ((1 - st)D) \cdot y \\ &= \sigma(x, y, st). \end{aligned}$$

So we have proved that σ is a consistent bicombing.

We will now prove that, for every $x, y \in M_{\mathbb{R}}$ with $d(x, y) = D$ and every $t \in [0, 1]$, we have a symmetrical formula for σ :

$$\sigma(x, y, t) = (tD) \cdot x \wedge ((1 - t)D) \cdot y = (-tD) \cdot x \vee (-(1 - t)D) \cdot y. \quad (1)$$

We start by proving the formula for $t = \frac{1}{2}$: since both x and y are superior to both $(-D) \cdot x$ and $(-D) \cdot y$, we deduce that $x \wedge y \geq (-D) \cdot x \vee (-D) \cdot y$, so $(\frac{1}{2}D) \cdot x \wedge (\frac{1}{2}D) \cdot y \geq (-\frac{1}{2}D) \cdot x \vee (-\frac{1}{2}D) \cdot y$. Similarly, since x and y are inferior to both $(D) \cdot x$ and $(D) \cdot y$, we deduce that $(-\frac{1}{2}D) \cdot x \vee (-\frac{1}{2}D) \cdot y \leq (\frac{1}{2}D) \cdot x \wedge (\frac{1}{2}D) \cdot y$. We conclude that

$$(-\frac{1}{2}D) \cdot x \vee (-\frac{1}{2}D) \cdot y = (\frac{1}{2}D) \cdot x \wedge (\frac{1}{2}D) \cdot y = \gamma(x, y, \frac{1}{2}).$$

Since σ is consistent and continuous, we deduce that Formula 1 holds for any $t \in [0, 1]$.

We will now prove that, for any $x, y, y' \in M_{\mathbb{R}}$, we have

$$d\left(\sigma\left(x, y, \frac{1}{2}\right), \sigma\left(x, y', \frac{1}{2}\right)\right) \leq \frac{d(y, y')}{2}.$$

Fix $x, y, y' \in M_{\mathbb{R}}$, and let us denote $D = \frac{d(x, y)}{2}$ and $D' = \frac{d(x, y')}{2}$. Let us write $m = \sigma(x, y, \frac{1}{2}) = (-D) \cdot x \vee (-D) \cdot y$ and $m' = \sigma(x, y', \frac{1}{2}) = (D') \cdot x \wedge (D') \cdot y'$. Let $A = \frac{d(y, y')}{2}$. We have $(A) \cdot m' = (D' + A) \cdot x \wedge (D' + A) \cdot y'$, we will prove that $m \leq (A) \cdot m'$. We have the following:

- $(-D) \cdot x \leq (D' + A) \cdot x$ because $D + D' + A \geq 0$,
- $(-D) \cdot x \leq (D' + A) \cdot y'$ because $D + D' + A = \frac{1}{2}(d(x, y) + d(x, y') + d(y, y')) \geq d(x, y') = 2D'$,
- $(-D) \cdot y \leq (D' + A) \cdot x$ because $D + D' + A = \frac{1}{2}(d(x, y) + d(x, y') + d(y, y')) \geq d(x, y) = 2D$ and
- $(-D) \cdot y \leq (D' + A) \cdot y'$ because $D + D' + A = d(x, y) + d(x, y') \geq d(y, y')$.

As a consequence, we have $m \leq (A) \cdot m'$. By the symmetry of the roles of y and y' , we also have $m' \leq (A) \cdot m$, hence $d(m, m') \leq A = \frac{d(y, y')}{2}$.

We deduce that, for any $x, x', y, y' \in M_{\mathbb{R}}$, we have

$$d\left(\sigma\left(x, y, \frac{1}{2}\right), \sigma\left(x', y', \frac{1}{2}\right)\right) \leq \frac{d(x, x') + d(y, y')}{2}.$$

Since σ is a consistent bicombing, we deduce that σ is convex. \square

Theorem 3.9. *Let L denote a bounded, graded lattice. The orthoscheme realization $|L|$ of L , endowed with the piecewise ℓ^∞ metric, is injective. Moreover, $|L|$ has a unique convex reversible consistent geodesic bicombing.*

Proof. Consider the affine version $M = M_{\mathbb{R}}$ of L over \mathbb{R} . For some maximal chain $c \in C(L)$, consider the elements $0_M = [(0, \dots, 0), c] \in M$, $\mu_M = [(\frac{1}{2}, \dots, \frac{1}{2}), c] \in M$ and $1_M = [(1, \dots, 1), c] \in M$: note that 0_M , μ_M and 1_M do not depend on c .

Note that the interval $I(0_M, 1_M)$ coincides with the ball $B(\mu_M, \frac{1}{2})$ in M for the metric $d_{\mathbb{R}}$. According to Theorem 3.7, $(M, d_{\mathbb{R}})$ is injective. So the ball $B(\mu_M, \frac{1}{2}) = I(0_M, 1_M)$ is injective.

We remark that the interval $I(0_M, 1_M)$ of M , endowed with the metric $d_{\mathbb{R}}$, is isometric to the orthoscheme realization of L , endowed with the piecewise ℓ^∞ metric. Indeed, for each maximal chain $c \in C(L)$, notice that $\sigma_C \cap I(0_M, 1_M)$ identifies with the standard orthoscheme $\{x \in \mathbb{R}^n \mid 0 \leq x_1 \leq \dots \leq x_n \leq 1\}$, where n denotes the rank of L . We conclude that the orthoscheme realization of L is injective.

According to Theorem 3.8, we know that $|L|$ has a convex reversible consistent geodesic bicombing σ . Note that $|L|$ has combinatorial dimension at most n . According to [DL15, Theorem 1.2], we deduce that σ is the only convex bicombing on $|L|$. \square

Note that knowing when the orthoscheme complex of a lattice, endowed with the piecewise Euclidean metric, is CAT(0) is a very subtle question (see the introduction). We can prove the following.

Theorem 3.10. *Let L denote a bounded, graded poset, let $|L|$ denote the geometric realization of L , and let d_p denote the ℓ^p orthoscheme metric on $|L|$, for $p \in \{2, \infty\}$. If $(|L|, d_2)$ is CAT(0), then $(|L|, d_\infty)$ is injective. The converse is false.*

Proof. Assume that L is a bounded, graded poset such that $(|L|, d_\infty)$ is not injective. According to Theorem 3.9, we deduce that L is not a lattice. According to Proposition 1.5, there exists a bowtie in L : consider $x_1, x_2, x_3, x_4 \in L$ such that x_1 and x_3 are maximal elements inferior to both x_2 and x_4 , and such that x_2 and x_4 are minimal elements superior to both x_1 and x_3 . In the link $L_{[0,1]}$ of the diagonal edge $[0, 1]$ in $(|L|, d_2)$, consider the piecewise geodesic loop ℓ going through x_1, x_2, x_3, x_4, x_1 .

According to [BM10, Proposition 4.8], each geodesic segment $[x_i, x_{i+1}]$ has length smaller than $\frac{\pi}{2}$, hence ℓ has length smaller than 2π . On the other hand, according to [BM10,

Lemma 7.2] the loop ℓ is locally geodesic in $L_{[0,1]}$. Hence $L_{[0,1]}$ is not CAT(1), and $(|L|, d_2)$ is not CAT(0).

We will now prove that the converse does not hold. Consider the Coxeter symmetric group $W = \mathfrak{S}_4$, with standard generators $S = \{s_1, s_2, s_3\}$ and standard Garside longest element $\Delta = s_1 s_2 s_3 s_1 s_2 s_1$. Let L denote the poset W , with order relation "being a left prefix for a shortest representative in S ". Then L is a lattice, and so $(|L|, d_\infty)$ is injective according to Theorem 3.9.

On the other hand, consider the piecewise geodesic loop ℓ in the link $L_{[0,1]}$ of the diagonal edge $[0, 1]$ in $(|L|, d_2)$ going through the vertices $s_1, s_1 s_2 s_1, s_2, s_2 s_3 s_2, s_3, s_3 s_1, s_1$. According to [BM10, Proposition 4.8], its length is

$$2 \arccos \left(\sqrt{\frac{14}{25}} \right) + 4 \arccos \left(\sqrt{\frac{13}{35}} \right) \simeq 0.987(2\pi) < 2\pi.$$

Since $L_{[0,1]}$ has the homotopy type of a circle, ℓ is not nullhomotopic in $L_{[0,1]}$. So $L_{[0,1]}$ is not CAT(1), and $(|L|, d_2)$ is not CAT(0). \square

We also deduce an immediate consequence concerning the Garside complex of a Garside group. Fix a Garside group (G, S, δ) . Let X denote the *Garside complex* of G , i.e. the simplicial complex with vertex set G , and with simplices corresponding to chains $g_1 <_L g_2 <_L \cdots <_L g_n$ such that $g_n \leq_L g_1 \delta$. Since simplices of X have a total order on their vertices, X is a simplicial complex with ordered simplices, so we may endow X with the piecewise ℓ^∞ orthoscheme metric.

Corollary 3.11. *The Garside complex of any Garside group, endowed with the piecewise ℓ^∞ orthoscheme metric, is injective.*

This applies, in particular, to the dual braid complex studied by Brady and McCammond in [BM10]. This complex, endowed with the piecewise orthoscheme Euclidean metric, is conjectured to be CAT(0), but it is only known for a very small number of strands (see [BM10], [HKS16], [Jeo20]). On the other hand, if we endow the dual braid complex with the piecewise orthoscheme ℓ^∞ metric, we see that it is injective.

4 Application to Euclidean buildings and the Deligne complex of type \widetilde{A}_n

Let us consider the Euclidean Coxeter group $W \simeq \mathfrak{S}_n \times \mathbb{Z}^{n-1}$ of type \widetilde{A}_{n-1} . Its Coxeter complex may be identified with

$$\Sigma = \{x \in \mathbb{R}^n \mid x_1 + \cdots + x_n = 0\}.$$

Up to homothety, we may choose the following affine hyperplanes to define Σ :

$$\{x_i - x_j = k \mid 1 \leq i \neq j \leq n, k \in \mathbb{Z}\},$$

so that maximal simplices of Σ identify with the W -orbit of

$$K = \{x \in \Sigma \mid x_1 \leq x_2 \leq \cdots \leq x_n \leq x_1 + 1\}.$$

The vertex set of Σ identifies with $(\frac{1}{n}\mathbb{Z})^n \cap \Sigma$. Since the simplex K is a strict fundamental domain for the action of W on Σ , one may define a W -invariant type function τ on the vertex set of Σ :

$$\begin{aligned} \tau : \Sigma^{(0)} &\rightarrow \mathbb{Z}/n\mathbb{Z} \\ x \in W \cdot v_i &\mapsto i, \end{aligned}$$

where $v_i = (\frac{n-i}{n}, \dots, \frac{n-i}{n}, \frac{-i}{n}, \dots, \frac{-i}{n})$ is the vertex of K whose first i coordinates equal $\frac{n-i}{n}$ and the $n-i$ last coordinates equal $\frac{-i}{n}$.

This type function is such that adjacent vertices have distinct types.

Let us define the *extended* Coxeter complex

$$\hat{\Sigma} = \Sigma \times \mathbb{R} = \mathbb{R}^n,$$

where the action of the standard generators w_1, \dots, w_n of W on $\hat{\Sigma}$ is given by

$$\begin{aligned} \forall 1 \leq i \leq n-1, w_i \cdot (x_1, \dots, x_i, x_{i+1}, \dots, x_n) &= (x_1, \dots, x_{i+1}, x_i, \dots, x_n) \\ \text{and } w_n \cdot (x_1, \dots, x_n) &= (x_n - 1, x_2, \dots, x_{n-1}, x_1 + 1). \end{aligned}$$

Also note that $\hat{\Sigma}$ has a natural simplicial complex structure, with vertex set \mathbb{Z}^n , with maximal simplices corresponding to the W -orbits of the simplices

$$\{k \leq x_i \leq x_{i+1} \leq x_{i+2} \leq \dots \leq x_{i+n-1} \leq k+1 \mid k \in \mathbb{Z}, i \in \mathbb{Z}/n\mathbb{Z}\},$$

where indices in \mathbb{R}^n are considered modulo n . Note that the action of the Coxeter group W preserves τ .

A fundamental domain for the action of W on Σ is the simplex

$$K = \{x \in \Sigma \mid x_1 \leq x_2 \leq \dots \leq x_n \leq x_1 + 1\},$$

and a fundamental domain for the action of W on $\hat{\Sigma}$ is the column

$$\hat{K} = \{x \in \hat{\Sigma} \mid x_1 \leq x_2 \leq \dots \leq x_n \leq x_1 + 1\}.$$

Note that such columns have been studied by Brady-McCammond in [BM10], and by Dougherty, McCammond and Witzel in [DMW].

We will endow $\hat{\Sigma}$ with the standard ℓ^∞ metric from \mathbb{R}^n .

Let us consider a simplicial complex X such that:

- either X is a Euclidean building of type \widetilde{A}_{n-1}
- or X is the Deligne complex of the Euclidean Artin group A of type \widetilde{A}_{n-1} , with a coarser simplicial structure.

We will define the *extended* version of X , denoted \hat{X} . It is a simplicial complex whose geometric realization is homeomorphic to $X \times \mathbb{R}$.

Before giving the precise definition of \hat{X} , let us first recall briefly the definition of the Deligne complex Δ of the Euclidean Artin group A of type \widetilde{A}_{n-1} (see also Section 1.2).. Let $S \simeq \mathbb{Z}/n\mathbb{Z}$ denote the standard generating set of A . Consider the set

$$\{gA_T \mid g \in A, T \subsetneq S\},$$

endowed with the following partial order: $gA_T \leq g'A_{T'}$ if $gA_T \subset g'A_{T'}$. Then the Deligne complex Δ of A is the geometric realization of this poset. We will define a coarser simplicial structure X on Δ . Note that, for any minimal vertex $gA_\emptyset \in \Delta$ (where $g \in A$), the 1-neighbourhood of gA_\emptyset in Δ is

$$\{gA_T \mid T \subsetneq S\},$$

which is precisely the barycentric subdivision of the simplex with vertex set

$$\{gA_T \mid T \subsetneq S, |S \setminus T| = 1\}$$

consisting of (the g -translates of) all maximal proper standard parabolic subgroups of A .

Therefore we may define the simplicial complex X with vertex set $\{gA_T \mid g \in A, T \subsetneq S, |S \setminus T| = 1\}$, and such that $g_1A_{T_1}, \dots, g_kA_{T_k}$ span a simplex in X if and only if $\bigcap_{i=1}^k g_iA_{T_i} \neq \emptyset$. Then Δ identifies with the barycentric subdivision of X , and also the geometric realizations of Δ and X are homeomorphic.

Note that, in both cases (Euclidean building or Deligne complex), there is a well-defined type function $\tau : X^{(0)} \rightarrow \mathbb{Z}/n\mathbb{Z}$ such that adjacent vertices have different types.

- In the case X is a Euclidean building of type \widetilde{A}_{n-1} , each apartment is identified with the Coxeter complex Σ , we may define the type of a vertex v of X to be its type in any apartment containing v . Since the Coxeter group W preserves the type, this definition does not depend on the choice of apartment.
- In the case X is the Deligne complex of type \widetilde{A}_{n-1} , one may either use the projection onto the Coxeter complex, or use a direct definition: if gA_T is a vertex of X , with $g \in A$ and $T = S \setminus \{i\}$, its type is i .

The complex X may also be defined as a particular gluing of copies of the Coxeter complex Σ . Roughly speaking, \hat{X} will be the same gluing of copies of $\hat{\Sigma}$.

More precisely, let $\hat{X} = X \times \mathbb{R}$, and we will define a simplicial complex structure on \hat{X} . The vertex set of \hat{X} is $\{(x, i) \in X^{(0)} \times \mathbb{Z} \mid \tau(x) = i + n\mathbb{Z}\}$, where $X^{(0)}$ denotes the vertex set of X . Using the type function τ , remark that vertices of maximal simplices of X have a well-defined cyclic ordering in $\mathbb{Z}/n\mathbb{Z}$. The maximal simplices of \hat{X} are

$$\{(x_i, kn+i), (x_{i+1}, kn+i+1), \dots, (x_n, kn+n), (x_1, kn+n+1), (x_2, kn+n+2), \dots, (x_i, kn+n+i)\},$$

for any $k \in \mathbb{Z}$, any $1 \leq i \leq n$ and any maximal simplex (x_1, x_2, \dots, x_n) of X with $\forall 1 \leq i \leq n, \tau(x_i) = i$. We endow each such simplex with the ℓ^∞ orthoscheme metric for the given ordering. Endow \hat{X} with the associated length metric.

Note that the translation action on the \mathbb{R} factor of $\hat{X} = X \times \mathbb{R}$ defines an isometric action denoted θ . Also note that θ does not preserve the simplicial structure of \hat{X} , but only its restriction to the subgroup $n\mathbb{Z}$ of \mathbb{R} .

If X is a Euclidean building of type \widetilde{A}_{n-1} , then \hat{X} is called a Euclidean building of extended type \widetilde{A}_{n-1} (see [BT84], and also [Hae21]).

If X is the Deligne complex of the Euclidean Artin group A of type \widetilde{A}_{n-1} , we will give another description of \hat{X} . Recall that the classical Deligne complex X may be defined as

$$X = (A \times K) / \sim,$$

where $(g, x) \sim (g', x')$ if $x = x'$ and, if the stabilizer of x in Σ equals W_T for some $T \subsetneq S$, then $g^{-1}g' \in A_T$. Then the extended Deligne complex may also be defined similarly as

$$\hat{X} = (A \times \hat{K}) / \sim,$$

where $(g, x) \sim (g', x')$ if $x = x'$ and, if the stabilizer of x in $\hat{\Sigma}$ equals W_T for some $T \subsetneq S$, then $g^{-1}g' \in A_T$.

Fix any vertex $x \in X$, and let $L_{x,0}$ denote the set of vertices of X adjacent to x . Without loss of generality, we may assume that x has type $\tau(x) = 0$. Note that vertices in $L_{x,0}$ have a type in $\mathbb{Z}/n\mathbb{Z} \setminus \{\tau(x)\} = \{1, \dots, n\}$, which is an interval. Since maximal simplices of X have a natural cyclic ordering in $\mathbb{Z}/n\mathbb{Z}$, there is a natural induced order on $L_{x,0}$ that is consistent with the type function $\tau : L_{x,0} \rightarrow \{1, \dots, n\}$. Consider the poset $L_x = L_{x,0} \cup \{0, 1\}$, where 0 and 1 are defined to be the minimum and the maximum of L_x respectively.

Proposition 4.1. *Consider any $p \in \hat{X}$ whose projection p_X onto X is contained in the open star of the vertex x . Then \hat{X} is locally isometric at p to a neighbourhood of a point in the ℓ^∞ orthoscheme realization of the poset L_x .*

Proof. Remember that there is a diagonal, isometric action θ of \mathbb{R} on \hat{X} , whose quotient is X . So, up to isometry, we may assume that p lies in the open star of the diagonal edge joining the vertices $(x, 0)$ and (x, n) of \hat{X} . Note that each simplex of \hat{X} containing p corresponds to a chain in L_x containing $(x, 0)$ (identified with $0 \in L_x$) and (x, n) (identified with $1 \in L_x$). As a consequence, a neighbourhood of p in \hat{X} is isometric to a neighbourhood of a point in the orthoscheme realization of L_x . \square

We now prove that, in the case of a Euclidean building, the poset L_x is a lattice.

Proposition 4.2. *Let L_0 denote the vertex set of a (possibly non-thick) spherical building of type A_{n-1} , for some $n \geq 1$, and let $L = L_0 \cup \{0, 1\}$. In other words, L is the poset of linear subspaces of a projective space of dimension $n - 1$. Then L is a bounded, graded lattice of rank $n - 1$.*

Proof. The lattice property is obvious: if S, S' are linear subspaces of a projective space X , then $S \wedge S' = S \cap S'$ and $S \vee S'$ is the linear subspace spanned by $S \cup S'$. \square

As a consequence, if X is a Euclidean building of type \widetilde{A}_{n-1} , for any vertex $x \in X$, the poset L_x is a bounded graded lattice.

We now turn to the case of the Deligne complex of type \widetilde{A}_{n-1} . We defer the proof of the following result to Section 5.

Theorem 4.3 (Crisp-McCammond).

Let X denote the Deligne complex of type \widetilde{A}_{n-1} . For any vertex $x \in X$, the poset L_x is a bounded, graded lattice.

We may now deduce the main result.

Theorem 4.4. *Any Euclidean building of extended type \widetilde{A}_{n-1} , or the extended Deligne complex of type \widetilde{A}_{n-1} , endowed with the piecewise ℓ^∞ metric, is injective.*

Proof. According to Proposition 4.1 and Theorem 4.3, we know that \hat{X} is locally isometric to the ℓ^∞ orthoscheme complex of a bounded graded lattice. According to Theorem 3.9, we deduce that \hat{X} is uniformly locally injective.

If X is a Euclidean building, X and \hat{X} are contractible, and in particular simply connected. If X is a Deligne complex, according to Theorem 1.6, \hat{X} is simply connected.

According to Theorem 1.10, \hat{X} is complete. We deduce with Theorem 1.13 that \hat{X} is injective. \square

We can also deduce that the thickening is a Helly graph.

Corollary 4.5. *The thickening of the vertex set of any Euclidean building of extended type \widetilde{A}_{n-1} , or the extended Deligne complex of type \widetilde{A}_{n-1} , is a Helly graph.*

Proof. Let X denote either a Euclidean building of extended type \widetilde{A}_{n-1} or the extended Deligne complex of type \widetilde{A}_{n-1} . We will see that X satisfies the assumptions of Theorem 1.14.

Since simplices of X have a well-defined order, we see that X is a simplicial complex with ordered simplices. For each maximal simplex σ of X , the minimal and maximal vertices of σ form a maximal edge in X . Therefore, X has maximal edges as in Definition 1.8. According to Theorem 1.14, we deduce that the thickening of X is a Helly graph. \square

We will now deduce a bicombing on X , considered as the quotient of $(\hat{X}, d_{\hat{X}})$ by the diagonal action θ of \mathbb{R} on \hat{X} . Let us define the quotient metric d_X on X :

$$\forall x, y \in X, d_X(x, y) = \inf_{t \in \mathbb{R}} d_{\hat{X}}(x, \theta(t) \cdot y).$$

Theorem 4.6. *Any Euclidean building X of type \widetilde{A}_{n-1} , or the Deligne complex X of type \widetilde{A}_{n-1} , has a metric d_X that admits a unique convex, consistent, reversible geodesic bicombing σ . Moreover, d_X and σ are invariant under the group of type-preserving automorphisms of X .*

Proof. We will prove it locally, i.e. for the star Y of a vertex v of X . According to Theorem 3.9, there exists a unique convex, consistent, reversible bicombing $\hat{\sigma}$ on \hat{Y} . For each $x, y \in Y$, choose lifts $\hat{x}, \hat{y} \in \hat{Y}$ such that $d_{\hat{X}}(\hat{x}, \hat{y}) = d_X(x, y)$. For each $t \in [0, 1]$, let us define $\sigma(x, y, t) = \theta(\mathbb{R}) \cdot \hat{\sigma}(\hat{x}, \hat{y}, t) \in Y$.

We will first see that σ is well-defined: indeed if there exists $t \neq 0$ such that $d_{\hat{X}}(\hat{x}, \theta(t) \cdot \hat{y}) = D$, then $d_{\hat{X}}(\hat{x}, \theta(\frac{t}{2}) \cdot \hat{y}) < D$, which is a contradiction. Hence σ is well-defined. Moreover, it is easy to see that σ is a reversible, consistent, geodesic bicombing.

We will now see that σ is convex. Consider $x, x', y, y' \in Y$, and choose lifts $\hat{x}, \hat{x}', \hat{y}, \hat{y}' \in \hat{Y}$ such that $d_{\hat{X}}(\hat{x}, \hat{y}) = d_X(x, y)$ and $d_{\hat{X}}(\hat{x}', \hat{y}') = d_X(x', y')$. Fix any $t \in [0, 1]$ and $s \in \mathbb{R}$. We have

$$\begin{aligned} d_X(\sigma(x, y, t), \sigma(x', y', t)) &= \inf_{s \in \mathbb{R}} d_{\hat{X}}(\hat{\sigma}(\hat{x}, \hat{y}, t), \theta(s) \cdot \hat{\sigma}(\hat{x}', \hat{y}', t)) \\ &\leq d_{\hat{X}}(\hat{\sigma}(\hat{x}, \hat{y}, t), \hat{\sigma}(\hat{x}', \hat{y}', t)) \\ &\leq (1-t)d_{\hat{X}}(\hat{x}, \hat{x}') + td_{\hat{X}}(\hat{y}, \hat{y}') \\ &\leq (1-t)d_X(x, x') + td_X(y, y'), \end{aligned}$$

so σ is a convex bicombing.

We have seen that (X, d_X) is locally convexly reversibly consistently bicomvable. According to [Mie17, Theorem 1.1], we deduce that (X, d_X) has a unique global convex reversible geodesic consistent bicombing σ that is consistent with local bicombing. Since the local bicombing only depends on the local combinatorics of X and the type, we deduce that σ is invariant under type-preserving automorphisms of X . \square

5 The lattice of maximal parabolic subgroups of braid groups

This whole section is unpublished work of John Crisp and Jon McCammond, copied here with their permission. The results concerning the lattice of cut-curves are contained in Bessis's work (see [Bes06]) about Garside structures on free groups. For consistency, we choose to follow Crisp and McCammond's presentation instead.

5.1 The lattice of cut-curves

Let \mathbb{D}^2 denote the unit disk in \mathbb{R}^2 , and let $\{p_1, \dots, p_n\}$ denote a set of n distinct points in \mathbb{D}^2 , as shown in Figure 2. We write $\mathbb{D}_* = \mathbb{D}^2 \setminus \{p_1, \dots, p_n\}$. Let $a = (0, 1)$ and $b = (0, -1)$ denote the top and bottom points of the boundary $\partial\mathbb{D}^2$.

Definition 5.1. By a *cut-curve* or *curve* on \mathbb{D}_* we shall mean a smoothly embedded curve c in \mathbb{D}_* which meets $\partial\mathbb{D} = \partial\mathbb{D}^2$ precisely at its endpoints, and which separates the boundary points a and b .




Figure 2: An example of cut-curve c in the punctured disk \mathbb{D}_* .

We shall say that two cut-curves are isotopic if they are isotopic in \mathbb{D}_* relative to $\{a, b\}$. We denote by $[c]$ the isotopy class of a curve c , and write \mathcal{C} for the set of all isotopy classes of cut-curves in \mathbb{D}_* .

Observe that any cut-curve c separates \mathbb{D}_* into two regions, an upper region containing a and a lower region containing b , and induces, in particular, a partition of the points $\{p_1, \dots, p_n\}$ into two sets. In general we shall say that the contents of the region containing a lie *above* c and the contents of the region containing b lie *below* c . For each curve c , we write $\deg(c)$ for the number of points p_i which lie below c . Clearly this number is invariant under isotopy, and so defines a degree function on \mathcal{C} by $\deg([c]) = \deg(c)$.

Let c_1, c_2 be two curves. We say that c_1 and c_2 are *in minimal position* with respect to one another if they do not cobound any disk regions ("bigons") in \mathbb{D}_* . This includes triangular regions against the boundary. Any such disk regions can always be removed by modifying just one of the two curves in its isotopy class without changing the other. That is to say that, for any two curves c_1, c_2 we can always find c'_1 such that $[c'_1] = [c_1]$ and c'_1 is in minimal position with respect to c_2 .

Here is another argument about minimal position using hyperbolic geometry. Such an argument has been used in [Bes06]. If $n \geq 2$, one may also endow \mathbb{D}^* with a fixed complete hyperbolic metric. Fix $p^+ = (1, 0), p^- = (-1, 0) \in \partial\mathbb{D}$ points on each connected component of $\partial\mathbb{D} \setminus \{a, b\}$. Then, for each isotopy class $[c]$, we may consider the unique geodesic line in \mathbb{D}^* in the isotopy class $[c]$ with endpoints $p^+, p^- \in \partial\mathbb{D}$. Then, for any $[c_1], [c_2] \in \mathcal{C}$, the geodesic representatives are in minimal position.

Definition 5.2. Let c_1, c_2 be two curves. We say that $[c_1] \leq [c_2]$ if c_1 is isotopic to a curve which lies below c_2 . It is easily checked that this defines a partial order on the set \mathcal{C} of cut-curves classes.

Note that if c_1 and c_2 are in minimal position with respect to one another then $[c_1] \leq [c_2]$ if and only if c_1 lies below c_2 (in particular they are disjoint). Note also that the function $\deg : \mathcal{C} \rightarrow \{1, \dots, n\}$ is a strict order preserving map, a grading on the poset (\mathcal{C}, \leq) .

Theorem 5.3. [Bes06, Theorem 2.6] *The graded poset (\mathcal{C}, \leq) is a lattice.*

Proof. Let $x = [c_1], y = [c_2]$ be arbitrary elements of \mathcal{C} . Suppose that the curves c_1 and c_2 are in minimal position with respect to one another. Now consider how the union of c_1

and c_2 cut \mathbb{D}_* up into connected regions. There is a unique lowermost such region R which lies below both c_1 and c_2 (and contains the point b), and another uppermost such region R' which lies above both c_1 and c_2 (and contains the point a). Let c , resp. c' , denote the curves which skirt along the part of the boundary of R , resp. R' , lying in the interior of \mathbb{D}_* (i.e. not along $\partial\mathbb{D}$). Then we claim that $x \wedge y = [c]$ and $x \vee y = [c']$.

We check just the first of these two claims. Suppose that c_0 represents a common lower bound for x and y . Then, by a sequence of bigon removing isotopies (or by considering the geodesic representative), we may choose the representative c_0 to be in minimal position with respect to both c_1 and c_2 . Since c_0 represents a common lower bound, c_0 lies below both c_1 and c_2 , hence lies below the curve c . \square

Recall that the n -strand braid group B_n is isomorphic to the mapping class group $\text{MCG}(\mathbb{D}_*, \partial\mathbb{D})$. As a consequence, B_n acts naturally on the set \mathcal{C} of isotopy classes of cut-curves.

Lemma 5.4. *The action of B_n on \mathcal{C} preserves the order and the degree.*

Proof. Let $[c] \in \mathcal{C}$ be a curve of degree $1 \leq k \leq n-1$ and $g \in B_n$. Then k points among $\{p_1, \dots, p_n\}$ lie below c . Since the action B_n fixes the boundary of \mathbb{D}_* pointwise, we know that k points among $\{p_1, \dots, p_n\}$ lie below $g(c)$. Hence $\deg g([c]) = \deg [c]$.

Similarly, consider two curves $[c_1], [c_2] \in \mathcal{C}$ such that $[c_1] \leq [c_2]$. Consider c_1, c_2 in minimal position, so that c_1 is below c_2 . For any $g \in B_n$, we have that $g(c_1)$ is below $g(c_2)$, hence $g([c_1]) \leq g([c_2])$. We conclude that the action of B_n preserves the order on \mathcal{C} . \square

Note that B_n acts transitively on the set of cut-curves with fixed degree.

5.2 Cosets in braid groups

We will show how the cut-curve lattice (\mathcal{C}, \leq) can be reinterpreted in purely algebraic terms.

Definition 5.5. Recall that the braid group B_n is generated by the set of standard generators $S = \{\sigma_1, \sigma_2, \dots, \sigma_{n-1}\}$. For each $1 \leq k \leq n-1$, let P_k denote the maximal parabolic subgroup of B_n generated by $S \setminus \{\sigma_k\}$. Thus P_k is isomorphic to a product $B_k \times B_{n-k}$. Note also that these subgroups are distinct for distinct values of k . We define the following augmented collection of cosets in B_n :

$$\mathcal{B} = \{gP_k \mid g \in B_n, 1 \leq k \leq n-1\} \cup \{0, 1\}.$$

For $b \in \mathcal{B}$, we write $\deg(b) = k$ if $b = gP_k$ for $1 \leq k \leq n-1$, and $\deg(0) = 0$, $\deg(1) = n$. We define an order relation $\leq_{\mathcal{B}}$ on \mathcal{B} as follows. First, define $0 \leq b \leq 1$ for all $b \in \mathcal{B}$. Otherwise, for $g_1, g_2 \in B_n$ and $1 \leq k_1, k_2 \leq n-1$, we write $g_1P_{k_1} \leq_{\mathcal{B}} g_2P_{k_2}$ if $k_1 \leq k_2$ and $g_1P_{k_1} \cap g_2P_{k_2} \neq \emptyset$.

For simplicity, we shall henceforth identify B_n with the mapping class group of \mathbb{D}_* (relative to $\partial\mathbb{D}$). Let us consider the base maximal chain $\alpha = (c_1, c_2, \dots, c_{n-1})$ of \mathcal{C} depicted in Figure 3.

Definition 5.6. Define a map $\Phi : \mathcal{C} \rightarrow \mathcal{B}$ by setting $\Phi(c) = 0$ if $\deg(c) = 0$, $\Phi(c) = 1$ if $\deg(c) = n$, and otherwise

$$\Phi(c) = \{g \in B_n \mid g(\alpha) \text{ contains } c\}.$$

Lemma 5.7. *If $c \in \alpha$ and $1 \leq \deg(c) = k \leq n-1$, then $\Phi(c) = P_k$.*




Figure 3: The base maximal chain $\alpha = (c_1, c_2, \dots, c_{n-1})$ of \mathcal{C}

Proof. The curve c is a base curve such that p_1, \dots, p_k lie below c and p_{k+1}, \dots, p_n lie above c . Let R^- denote the connected component of $\mathbb{D}^* \setminus c$ below c , and R^+ denote the connected component of $\mathbb{D}^* \setminus c$ above c . The stabilizer of c in the mapping class group $B_n = \text{MCG}(\mathbb{D}_*, \partial\mathbb{D})$ is the direct product of $\text{MCG}(R^-, \partial R^-) = \langle \sigma_1, \dots, \sigma_{k-1} \rangle$ and $\text{MCG}(R^+, \partial R^+) = \langle \sigma_{k+1}, \dots, \sigma_{n-1} \rangle$. Hence the stabilizer of c in B_n equals P_k . \square

Lemma 5.8. *The map $\Phi : \mathcal{C} \rightarrow \mathcal{B}$ is well-defined, surjective, and respects degrees.*

Proof. Fix $c \in \mathcal{C}$ with $1 \leq \deg(c) = k \leq n-1$. Fix some $g \in B_n$ such that the curve c lies in $g(\alpha)$. Let $c_0 = g^{-1}(c)$, the degree k curve in α . For any $h \in B_n$, we have

$$h \in \Phi(c) \Leftrightarrow c \in h(\alpha_0) \Leftrightarrow g(c_0) = h(c_0).$$

According to Lemma 5.7, this is equivalent to $gP_k = hP_k$. Hence $\Phi(c) = gP_k$ is well-defined. Moreover it is clear that Φ is surjective, and that for all $c \in \mathcal{C}$ we have $\deg(\Phi(c)) = \deg(c)$. \square

Lemma 5.9. *The map Φ is injective, so it is a bijection between \mathcal{C} and \mathcal{B} .*

Proof. Let $c_1, c_2 \in \mathcal{C}$ such that $\Phi(c_1) = \Phi(c_2)$. We may suppose that c_1 lies in $g_1(\alpha)$ and c_2 in $g_2(\alpha)$, for some $g_1, g_2 \in B_n$. According to Lemma 5.8, we deduce that c_1 and c_2 have the same degree $0 \leq k \leq n$. Since \mathcal{C} and \mathcal{B} have unique elements of degree 0 and n , we may restrict to the case $1 \leq k \leq n$.

Since $\Phi(c_1) = \Phi(c_2)$, we deduce that $g_1P_k = g_2P_k$, so $g_1^{-1}g_2 \in P_k$. Let $h \in P_k$ such that $g_2 = g_1h$. Let c_0 denote the degree k curve of α . Then $c_1 = g_1(c_0)$ and $c_2 = g_2(c_0)$. Since $h \in P_k$, h fixes c_0 so $c_1 = c_2$. \square

Theorem 5.10 (Crisp-McCammond). *The bijection $\Phi : \mathcal{C} \rightarrow \mathcal{B}$ is an order isomorphism between (\mathcal{C}, \leq) and $(\mathcal{B}, \leq_{\mathcal{B}})$. As a consequence, $(\mathcal{B}, \leq_{\mathcal{B}})$ is a lattice.*

Proof. To prove the theorem, we need to show that for $c_1, c_2 \in \mathcal{C}$, we have $c_1 \leq c_2$ if and only if $\Phi(c_1) \leq_{\mathcal{B}} \Phi(c_2)$. Write $k_i = \deg(c_i)$ for $i = 1, 2$, and suppose, without loss, that $0 \leq k_1 \leq k_2 \leq n$. Then $\Phi(c_1) \leq_{\mathcal{B}} \Phi(c_2)$ if and only if $\Phi(c_1) \cap \Phi(c_2) \neq \emptyset$ if and only if there exists a maximal chain $\alpha' \subset \mathcal{C}$ which contains both c_1 and c_2 , if and only if $c_1 \leq c_2$ (since we already know that $\deg(c_1) \leq \deg(c_2)$). \square

We deduce the proof of Theorem 4.3 that each local poset L_x in the Deligne complex of type \tilde{A}_{n-1} , is a lattice.

Proof. [of Theorem 4.3] Let x denote a vertex of the Deligne complex X of type \tilde{A}_{n-1} . Without loss of generality, we may assume that x correspond to the maximal proper parabolic subgroup $A(A_{n-1}) = \langle \sigma_1, \dots, \sigma_{n-1} \rangle$, which is isomorphic to the Artin group of type A_{n-1} , i.e. the n -strand braid group. Since $A(\tilde{A}_{n-1})$ is a maximal parabolic subgroup of spherical type of the Euclidean Artin group $A(\tilde{A}_{n-1})$, the star of x in X identifies with the Deligne complex of the Artin group $A(A_{n-1})$. In particular, vertices in X adjacent to x may be identified with cosets of proper maximal parabolic subgroups of the braid group $A(A_{n-1})$. Furthermore, given two such cosets gP_i and hP_j , for $g, h \in A(A_{n-1})$ and $1 \leq i, j \leq n-1$, the corresponding vertices of X are adjacent if and only if $gP_i \cap hP_j \neq \emptyset$. As a consequence, the poset $L_{x,0}$ is isomorphic to the poset $\mathcal{B} \setminus \{0, 1\}$, and L_x is isomorphic to the poset \mathcal{B} . According to Theorem 5.10, we deduce that L_x is a lattice. \square

6 The thickening of a semilattice

We will now consider a generalization of Theorem 3.9 to the case of a semilattice. This will be useful to consider Euclidean buildings and Deligne complexes in type \tilde{C}_n .

Theorem 6.1. *Let L denote a graded poset with minimum 0 and with bounded rank such that:*

- L is a meet-semilattice.
- If $x, y, z \in L$ are pairwise upperly bounded, then $\{x, y, z\}$ has a join.

Then the ℓ^∞ orthoscheme realization of $|L|$ is injective.

Proof. Let us consider $\bar{L} = L \cup \{1\}$: it is a bounded graded lattice. Let us denote by $\bar{M} = |\bar{L}|$ the geometric realization of \bar{L} , and $M = |L| \subset \bar{M}$ the geometric realization of L . Note that we consider \bar{L} as a subset of \bar{M} , its vertex set. The orthoscheme realization of M is the induced length metric on the subspace of the orthoscheme realization of \bar{M} .

According to Theorem 3.9, the orthoscheme realization of \bar{M} is injective. Let us denote by d the length metric of M , and by \bar{d} the injective metric on \bar{M} .

Fix $\varepsilon > 0$, and consider the graph Γ_ε with vertex set M , and with an edge between $x, y \in M$ if $\bar{d}(x, y) \leq \varepsilon$. We will prove that Γ_ε is Helly, by proving that it is clique-Helly and that its triangle complex is simply connected.

Let $\sigma \subset \Gamma_\varepsilon$ denote a maximal clique: we claim that there exists $m \in \bar{L}$ such that σ is the intersection of an interval $I_{\bar{M}}(a, b)$ in \bar{M} with M . Since maximal sets of diameter ε in \bar{M} are intervals, we deduce that there exists $a \leq b$ in \bar{M} such that $\sigma \subset I_{\bar{M}}(a, b)$ and $\bar{d}(a, b) \leq \varepsilon$. Since σ is a maximal clique, we deduce that $\sigma = I_{\bar{M}}(a, b) \cap M$.

We will prove that Γ_ε is clique-Helly. Consider a family of pairwise intersecting maximal cliques $(\sigma_i)_{i \in I}$ in Γ_ε . For each $i \in I$, according to the previous paragraph, there exists $a_i \leq b_i$ in \bar{M} such that $\sigma_i = I_{\bar{M}}(a_i, b_i) \cap M$. For each $i \in I$, let $m_i \in L$ minimal such that $a_i \leq m_i$. For each $i \neq j$ in I , since σ_i and σ_j intersect, the intersection $I_{\bar{M}}(a_i, b_i) \cap I_{\bar{M}}(a_j, b_j) \cap M$ is non-empty. In particular, the elements m_i and m_j have a common upper bound in L .

The family $(m_i)_{i \in I}$ is pairwise upperly bounded: by assumption, and since L is a graded lattice with bounded rank, we deduce that this family has a join $m = \bigvee_{i \in I} m_i$ in L . In particular, the family $(\sigma_i \cap I_{\bar{M}}(0, m))_{i \in I}$ of intervals in the lattice $I_{\bar{M}}(0, m) = I_M(0, m)$ has a non-empty intersection. So the graph Γ_ε is clique-Helly.



Figure 4: The Dynkin diagram of type B_n

We will now prove that the triangle complex of Γ_ε is simply connected. For each $t \in [0, 1]$, consider the map $\pi_t : M \rightarrow M$ sending each $x \in I(0, m)$ to the point on the affine segment joining 0 to x at distance $td(0, x)$ from 0. Note that π_t is 1-Lipschitz with respect to the distance d . Furthermore, if $t, t' \in [0, 1]$ are such that $|t - t'| \leq \varepsilon$, then for each $x \in M$ we have $d(\pi_t(x), \pi_{t'}(x)) \leq \varepsilon$. As a consequence, any combinatorial loop γ in Γ_ε may be homotoped in the triangle complex of Γ_ε to the loop $\pi_0(\gamma)$, which is the constant loop at 0. So we have proved that the triangle complex of Γ_ε is simply connected.

According to Theorem 1.11, we deduce that the graph Γ_ε is Helly. In particular, for each $\varepsilon > 0$, the metric space M is ε -coarsely injective. Since M is complete according to Theorem 1.10, we deduce by Lemma 1.12 that M is injective. \square

We will now give natural examples of such semilattices, which will be used in the sequel for Euclidean buildings and Deligne complexes of Euclidean type different from \widetilde{A}_n .

Proposition 6.2. *Let L_0 denote the vertex set of a (possibly non-thick) spherical building of type B_n , for some $n \geq 1$, and let $L = L_0 \cup \{0\}$. In other words, L is the poset of subspaces of a polar space of projective dimension $n - 1$. Then L is a graded semi-lattice of rank n with minimum 0, such that any pairwise upperly bounded subset of L has a join.*

Proof. The semi-lattice property is obvious: if S, S' are subspaces of a polar space X , then $S \wedge S' = S \cap S'$. If $A \subset L$ are pairwise upperly bounded in L , let $A_X = \bigcup_{S \in A} S \subset X$. For any $x, y \in A_X$, there exists a subspace of X containing x and y . According to [Tit74, 7.2.1], the subset A_X is contained in a subspace S_0 of X . Hence, for any $S \in A$, we have $S \subset S_0$. So the intersection of all subspaces of X containing $\bigcup_{S \in A} S$ is the join of A . \square

Another application concerns the Artin group $A(B_n)$ of spherical type B_n , with the following Dynkin diagram (see Figure 4).

Let s_1, \dots, s_n denote the standard generators of $A(B_n)$, with $s_{n-1}s_n s_{n-1}s_n = s_n s_{n-1} s_n s_{n-1}$. For each $1 \leq i \leq n$, let P_i denote the maximal proper standard parabolic subgroup of $A(B_n)$

$$P_i = \langle s_1, \dots, s_{i-1} \rangle \times \langle s_{i+1}, \dots, s_n \rangle.$$

Let $L = \{0\} \cup \{gP_i \mid g \in A(B_n), 1 \leq i \leq n\}$. We define an order relation on L as follows. First, define $0 \leq gP_i$ for each $gP_i \in L$. Otherwise, for $g_1, g_2 \in A(B_n)$ and $1 \leq k_1, k_2 \leq n$, define $g_1P_{k_1} \leq g_2P_{k_2}$ if $k_1 \leq k_2$ and $g_1P_{k_1} \cap g_2P_{k_2} \neq \emptyset$.

Note that we are grateful to Luis Paris for his help in the following proof, notably the use of normal forms.

Proposition 6.3. *L is a graded semi-lattice of rank n with minimum 0, such that any pairwise upperly bounded subset of L has a join.*

Proof. Let t_1, \dots, t_{2n-1} the standard generators of the braid group $A(A_{2n-1})$. Consider the following morphism

$$\begin{aligned} \phi : A(B_n) &\rightarrow A(A_{2n-1}) \\ \forall 1 \leq i \leq n-1, s_i &\mapsto t_i t_{2n-i} \\ s_n &\mapsto t_n. \end{aligned}$$

This morphism ϕ is injective, see for instance [DP99], [Mic99] and [Cri00]. Let σ denote the involution of $A(A_{2n-1})$ defined by $\forall 1 \leq i \leq 2n-1, \sigma(t_i) = t_{2n-i}$. According to [Cri00, Theorem 4], we know furthermore that the image of ϕ coincides with the fixed point set of σ .

For each $1 \leq i \leq 2n-1$, consider the standard proper maximal parabolic subgroup $Q_i = \langle t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_{2n-1} \rangle$ of $A(A_{2n-1})$. Let M_0 denote the poset of cosets of maximal proper parabolic subgroups of $A(A_{2n-1})$, and $M = M_0 \cup \{0, 1\}$. We will define a poset map $\psi : L_0 \rightarrow M_0$.

For each $g \in A(B_n)$ and each $1 \leq i \leq n$, let us define $\psi(gP_i) = \phi(g)Q_i$. Since $\phi(P_i) \subset Q_i$, the map $\psi : L_0 \rightarrow M_0$ is well-defined. Assume that $gP_i \leq hP_j$, i.e. $i \leq j$ and $h^{-1}g \in P_jP_i$. Then $\phi(h)^{-1}\phi(g) \in \phi(P_j)\phi(P_i) \subset Q_jQ_j$, so $\phi(g)Q_i \leq \phi(h)Q_j$. As a consequence, ψ is a rank-preserving injective poset map.

Note that the involution σ extends naturally to an order-reversing involution on M , by letting $\sigma(gQ_i) = \sigma(g)Q_{2n-i}$. And for each $g \in A(B_n)$ and each $1 \leq i \leq n$, we have $\sigma(\psi(gP_i)) = \phi(g)Q_{2n-i}$.

We will now prove that L is a meet-semilattice. Fix $a, b \in A(B_n)$ and $1 \leq i, j \leq n$, we will prove that aP_i and bP_j have a meet in L . Consider the elements $\phi(a)Q_i$ and $\phi(b)Q_j$ in M : they have a meet γQ_k , for some $\gamma \in A(A_{2n-1})$ and $0 \leq k \leq i, j$. Then as $\phi(a)Q_i \leq \phi(a)Q_{2n-i}$ and $\phi(b)Q_j \leq \phi(b)Q_{2n-j}$, we deduce that $\gamma Q_k \leq \sigma(\gamma)Q_{2n-k}$.

So, up to the choice of $\gamma \in \gamma Q_k$, we may assume that $\sigma(\gamma) \in \gamma Q_{2n-k}$: let $q \in Q_{2n-k}$ such that $\sigma(\gamma) = \gamma q$. Since σ is an involution, we have $\gamma = \gamma q \sigma(q)$, so $q = \sigma(q)^{-1} \in Q_k \cap Q_{2n-k}$.

We claim that if $g \in A(A_{2n-1})$ is such that $\sigma(g) = g^{-1}$, there exists $h \in A(A_{2n-1})^+$ such that $g = h\sigma(h)^{-1}$. According to [Cha95], there exist unique $h, h' \in A(A_{2n-1})^+$ such that $g = hh'^{-1}$ and the right greatest common divisor of h and h' in the Garside monoid $A(A_{2n-1})^+$ is 1. Note that σ preserves the monoid $A(A_{2n-1})^+$, hence we have $g^{-1} = h'h^{-1}$ on one side, and $g^{-1} = \sigma(g) = \sigma(h)\sigma(h')^{-1}$ on the other side. By uniqueness of h, h' , we deduce that $h' = \sigma(h)$. Hence $g = h\sigma(h)^{-1}$. Furthermore, note that if $g \in Q_k \cap Q_{2n-k}$, then $h \in Q_k \cap Q_{2n-k}$.

According to the claim, there exists $q' \in Q \cap Q_{2n-k}$ such that $q = q'\sigma(q')^{-1}$. So, up to replacing γ with $\gamma' = \gamma q' \in h_i Q_i$, we have

$$\sigma(\gamma') = \sigma(\gamma)\sigma(q') = \gamma q \sigma(q') = \gamma q' = \gamma'.$$

So we may assume that γ' is fixed by σ : according to [Cri00, Theorem 4], we may consider $c \in A(B_n)$ such that $\phi(c) = \gamma'$. So we deduce that $cP_k \leq aP_i, bP_j$. Conversely, for any $c' \in A(B_n)$ and $k' \leq i, j$ such that $c'P_{k'} \leq aP_i, bP_j$, we have $\phi(c')Q_{k'} \leq \phi(a)Q_i \wedge \phi(b)Q_j = \phi(c)Q_k$, so $c'P_{k'} \leq cP_k$. We conclude that cP_k is the meet of aP_i and bP_j in L .

We will now prove that any three pairwise upperly bounded elements of L have an upper bound. Fix $g_1, g_2, g_3 \in A(B_n)$ and $1 \leq k_1, k_2, k_3 \leq n$ such that, for each $i \neq j$, the elements $g_i P_{k_i}$ and $g_j P_{k_j}$ have a join $g_{ij} P_{k_{ij}}$ in L . Consider the elements $\phi(g_{ij})Q_{k_{ij}}$ of M : they have a join γQ_k in M . Since, for each $1 \leq i, j \leq 3$, we have $\phi(g_i)Q_i \leq \phi(g_{ij})Q_{k_{ij}} \leq \phi(g_{ij})Q_{2n-k_{ij}} \leq \phi(g_j)Q_{2n-k_j}$, we deduce that γQ_k is inferior to the meet $\sigma(\gamma)Q_{2n-k}$ of the elements $\phi(g_{ij})Q_{2n-k_{ij}}$. We deduce that $k \leq n$. Also, as in the previous paragraph, we deduce that we can choose $\gamma \in A(A_{2n-1})$ such that $\gamma = \phi(c)$, for some $c \in A(B_n)$. Hence cP_k is a common upper bound for the elements $(g_i P_{k_i})_{1 \leq i \leq 3}$.

Since L is graded without finite rank, we deduce that any family of pairwise upperly bounded elements of L have a join. \square

We believe that a similar statement holds for the Artin group of spherical type D_n , but we have no embedding into an Artin group of type A in order to use a similar proof.

7 Application to Euclidean buildings and the Deligne complex of other Euclidean types

In this section, we will prove that we can deduce from Theorem 4.4 an injective metric on Euclidean buildings and Deligne complexes of Euclidean types other than \widetilde{A}_n .

More precisely, let us consider a simplicial complex X that is:

- either a Euclidean building of type \widetilde{B}_n , \widetilde{C}_n or \widetilde{D}_n ,
- or the Deligne complex of Euclidean type \widetilde{C}_n .

Note that the Coxeter groups of types \widetilde{B}_n and \widetilde{D}_n may be considered as subgroups of the Coxeter group of Euclidean type \widetilde{C}_n , spanned by reflections with respect to subarrangements of hyperplanes, see below for more details. As a consequence, if X is a Euclidean building of type \widetilde{B}_n or \widetilde{D}_n , we will consider it as a (possibly non-thick) Euclidean building of type \widetilde{C}_n .

Let $W(\widetilde{C}_n)$ denote the Euclidean Coxeter group of type \widetilde{C}_n . Its Coxeter complex $\Sigma(\widetilde{C}_n)$ identifies with \mathbb{R}^n , and reflections of $W(\widetilde{C}_n)$ correspond to reflections with respect to hyperplanes

$$\{x_i = k \mid 1 \leq i \leq n, k \in \mathbb{Z}\} \text{ and } \{x_i \pm x_j = 2k \mid 1 \leq i \neq j \leq n, k \in \mathbb{Z}\}.$$

Note that we can see the following subarrangement has type \widetilde{B}_n :

$$\{x_i = 2k \mid 1 \leq i \leq n, k \in \mathbb{Z}\} \text{ and } \{x_i \pm x_j = 2k \mid 1 \leq i \neq j \leq n, k \in \mathbb{Z}\}.$$

Also the following subarrangement has type \widetilde{D}_n :

$$\{x_i \pm x_j = 2k \mid 1 \leq i \neq j \leq n, k \in \mathbb{Z}\}.$$

This justifies that, in the case of Euclidean buildings, we may assume that X is a (possibly non-thick) Euclidean building of type \widetilde{C}_n .

The vertex set of $\Sigma(\widetilde{C}_n)$ identifies with \mathbb{Z}^n , and a strict fundamental domain of the action of $W(\widetilde{C}_n)$ on $\Sigma(\widetilde{C}_n)$ is given by the standard orthoscheme simplex σ with vertices

$$v_0 = (0, \dots, 0), v_1 = (1, 0, \dots, 0), \dots, v_n = (1, 1, \dots, 1).$$

Hence v_0, v_1, \dots, v_n are representatives of the $n + 1$ orbits of vertices of $\Sigma(\widetilde{C}_n)$ under the action of $W(\widetilde{C}_n)$. This enables to define a type function τ on vertices of $\Sigma(\widetilde{C}_n)$:

$$\begin{aligned} \tau : \Sigma(\widetilde{C}_n)^{(0)} &\rightarrow \{0, 1, \dots, n\} \\ g \cdot v_i &\mapsto i. \end{aligned}$$

More generally, we can define a partial order on vertices of $\Sigma(\widetilde{C}_n)$: say that $v < v'$ if v, v' are adjacent vertices and $\tau(v) < \tau(v')$. We can therefore also view $\Sigma(\widetilde{C}_n)$ as the geometric realization of the poset of its vertices.

We will now see how to extend the type function and the partial order on vertices of X .

Proposition 7.1. *Assume that X is a Euclidean building of type \widetilde{C}_n or the Deligne complex of Euclidean type \widetilde{C}_n . There exists a type function $\tau : X^{(0)} \rightarrow \{0, 1, \dots, n\}$ such that adjacent vertices of X have different types. Moreover, let us define a partial order on vertices of X by setting $v < v'$ if v and v' are adjacent in X and $\tau(v) < \tau(v')$. Then X is the geometric realization of the poset of its vertices.*

Proof. Let us first consider the case where X is a Euclidean building of type \widetilde{C}_n . Given any vertex v of X , consider any apartment $A \subset X$ containing v , and let us define $\tau(v)$ as defined with respect to the apartment $A \simeq \Sigma(\widetilde{C}_n)$. Since two apartments containing v differ by an element of the Weyl group $W(\widetilde{C}_n)$, which preserves the type, we deduce that τ is well-defined on $X^{(0)}$. Similarly, given any two adjacent vertices v, v' in X , say that $v < v'$ if $\tau(v) < \tau(v')$.

Let us now consider the case where X is the Deligne complex of Euclidean type \widetilde{C}_n . Let $s_0, s_1, \dots, s_{n-1}, s_n$ denote the standard generators of the Artin group $A(\widetilde{C}_n)$. For each $0 \leq i \leq n$, consider the maximal spherical type standard parabolic subgroup

$$P_i = \langle s_0, s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_n \rangle \simeq A(B_i) \times A(B_{n-i}).$$

For each vertex gP_i in X , where $g \in A(\widetilde{C}_n)$ and $0 \leq i \leq n$, let us define $\tau(gP_i) = i$. Given any two vertices gP_i, hP_j of X , say that $gP_i < hP_j$ if they are adjacent in X (i.e. $gP_i \cap hP_j \neq \emptyset$) and $i < j$. \square

We will endow X with the piecewise ℓ^∞ orthoscheme metric d given by the geometric realization of its poset of vertices. We will now describe the local structure of X at any vertex.

Proposition 7.2. *Assume that X is a Euclidean building of type \widetilde{C}_n or the Deligne complex of Euclidean type \widetilde{C}_n . Fix a vertex $v \in X$ of type $\tau(v) = i \in \{0, 1, \dots, n\}$.*

There exist posets $L_{1,i}$ (resp. $L'_{0,n-i}$), which are either the poset of vertices of a spherical building of type B_i (resp. B_{n-i}) or the poset of maximal proper parabolic subgroups of the Artin group of spherical type B_i (resp. B_{n-i}). Let us define the posets $L_i = \{1\} \cup L_{1,i}$ and $L'_{n-i} = L'_{0,n-i} \cup \{0\}$, and let us consider the geometric realizations $|L_i|$ and $|L'_{n-i}|$, endowed with the piecewise ℓ^∞ orthoscheme metrics.

Then X is locally isometric at v to the ℓ^∞ direct product $|L_i| \times |L'_{n-i}|$.

Proof. The space X is locally isometric at v to the ℓ^∞ orthoscheme realization of the poset L_v of vertices of X comparable to v . The poset L_v is the disjoint union of $L_{1,i} = \{w \in L_v \mid w < v\} \sqcup \{v\} \sqcup L'_{0,n-i} = \{w \in L_v \mid w > v\}$, such that $L_{1,i} < \{v\} < L'_{0,n-i}$. We may identify the poset $L_i = L_{1,i} \cup \{1\}$ with the interval $L_{1,i} \cup \{v\}$ in L_v . Similarly, we may identify the poset $L'_{n-i} = L'_{0,n-i} \cup \{0\}$ with the interval $L'_{0,n-i} \cup \{v\}$ in L_v . Then the orthoscheme realization $|L_v|$ of L_v is locally isometric at v to the direct ℓ^∞ product $|L_i| \times |L'_{n-i}|$ of the orthoscheme realizations of L_i and L'_{n-i} .

In case X is a Euclidean building of type \widetilde{C}_n , note that $L_{1,i}$ is the vertex set of a spherical building of type B_i (with reversed order), and that $L'_{0,n-i}$ is the vertex set of a spherical building of type B_{n-i} .

In case X is the Deligne complex of type \widetilde{C}_n , note that $L_{1,i}$ is the poset of proper parabolic subgroups of the Artin group of type B_i (with reversed order), and that $L'_{0,n-i}$ is the poset of proper parabolic subgroups of the Artin group of type B_{n-i} . \square

We can now prove the following.

Theorem 7.3. *Any Euclidean building of type $\widetilde{B}_n, \widetilde{C}_n$ or \widetilde{D}_n , or the Deligne complex of type \widetilde{C}_n , endowed with the piecewise ℓ^∞ metric d , is injective.*

Proof. According to Proposition 7.2, the space X is locally isometric to a product of orthoscheme complexes of the type $|L|$, where $L = L_0 \cup \{0\}$ is a poset, and L_0 is either the poset of vertices of a spherical building of type B_i or the poset of maximal proper parabolic subgroups of the Artin group of spherical type B_i , for some $0 \leq i \leq n$.

According to Propositions 6.2 and 6.3, we know that in each case, L is a graded meet-semilattice with minimum 0 of rank i , such that any upperly bounded subset has a join. According to Theorem 6.1, we deduce that $|L|$ is injective.

We know that X is uniformly locally injective. According to Theorem 1.13, we deduce that X is injective. \square

We can also deduce that the thickening is a Helly graph.

Corollary 7.4. *The thickening of the vertex set of any Euclidean building of type \widetilde{B}_n , \widetilde{C}_n or \widetilde{D}_n , or the extended Deligne complex of type \widetilde{C}_n , is a Helly graph.*

Proof. Let X denote either a Euclidean building of type \widetilde{B}_n , \widetilde{C}_n or \widetilde{D}_n , or the extended Deligne complex of type \widetilde{C}_n . We will see that X satisfies the assumptions of Theorem 1.14.

Since simplices of X have a well-defined order, we see that X is a simplicial complex with ordered simplices. For each maximal simplex σ of X , the minimal and maximal vertices of σ form a maximal edge in X . Therefore, X has maximal edges as in Definition 1.8. According to Theorem 1.14, we deduce that the thickening of X is a Helly graph. \square

8 Bicomblings on Deligne complexes in types \widetilde{A}_n and \widetilde{C}_n

We now see that, in the Deligne complex of spherical types A_n , B_n and Euclidean types \widetilde{A}_n , \widetilde{C}_n , we may find a convex bicombling.

Theorem 8.1. *Let X denote the Deligne complex of the Artin group A of spherical type A_n , B_n or Euclidean type \widetilde{A}_n , \widetilde{C}_n . There is a metric d_X on X that admits a convex, consistent, reversible geodesic bicombling σ . Moreover, d_X and σ are invariant under A .*

Proof. The statement for types A_n and \widetilde{A}_n is Theorem 4.6.

Assume that X is the Deligne complex of type B_n . We have seen in the proof of Proposition 6.3 that X may be realized as the fixed point subspace of the Deligne complex Y of type A_{2n-1} for an involution s . Note that, since s is order-reversing, s induces an isometry of Y . According to Theorem 4.6, there exists a unique reversible, convex, consistent, geodesic bicombling σ_Y on Y for a metric d_Y . Since σ_Y is unique, we deduce that s preserves σ_Y , and that the fixed point subspace X is σ -stable. Let d_X denote the restriction of d_Y to X , and let σ_X denote the restriction of σ_Y to X . We deduce that σ_X is a convex, consistent, reversible geodesic bicombling on (X, d_X) . Moreover, d_X and σ_X are invariant under A .

Assume that X is the Deligne complex of type \widetilde{C}_n . According to [Dig12, Theorem 5.2] (it is also a consequence of Corollary 8.5), we see that X may be realized as the fixed point subspace of the Deligne complex Y of type \widetilde{A}_{2n-1} for an involution s . Following the same arguments as in the type B_n , we conclude that there exists a metric d_X on X and a convex, consistent, reversible geodesic bicombling σ_X on (X, d_X) that are both invariant under A . \square

We will describe several consequences we can derive from the fact that the Deligne complex has a consistent convex bicombling, that were usually known when the Deligne complex had a CAT(0) metric.

Corollary 8.2 (Okonek). *The Deligne complex X of Euclidean type \widetilde{A}_n or \widetilde{C}_n is contractible. In particular, the $K(\pi, 1)$ conjecture holds in these cases.*

Proof. Metric spaces with a convex bicombling are contractible. \square

The proof of Morris-Wright (see [MW19]) for the intersection of parabolic subgroups in FC type adapts directly to our situation. It relies mainly of the result by Cumplido et al. (see [CGGMW19]) that in a spherical type Artin group, the intersection of parabolic subgroups is a parabolic subgroup.

Corollary 8.3. *Let A denote the Artin group of Euclidean type \widetilde{A}_n or \widetilde{C}_n . The intersection of any family of parabolic subgroups of A is a parabolic subgroup.*

Proof. The proof of [MW19, Theorem 3.1] uses only the existence of an A -equivariant consistent geodesic bicombing on X . \square

The results by Godelle (see [God07]) describing centralizers and normalizers of parabolic subgroups also adapt to our case.

Corollary 8.4. *Let A denote the Artin group of Euclidean type \widetilde{A}_n or \widetilde{C}_n . Then A satisfies Properties (\star) , $(\star\star)$ and $(\star\star\star)$ from [God07], notably:*

- For any subset $X \subset S$, we have

$$\text{Com}_A(A_X) = N_A(A_X) = A_X \cdot QZ_A(X),$$

where the quasi-centralizer of X is $QZ_A(X) = \{g \in A \mid g \cdot X = X\}$.

- For any subsets $X, Y \subset S$ and any $g \in A$, we have

$$gA_Xg^{-1} \subset A_Y \implies \exists h \in A_Y, \exists Z \subset X, gA_Xg^{-1} = hA_Zh^{-1}.$$

Proof. The proof of [God07, Theorem 3.1] uses only the following properties of the Deligne complex, which hold for the Deligne complex X :

- There exists an A -equivariant consistent geodesic bicombing σ_X on X .
- Closed cells of X are σ_X -stable.

These assumptions are satisfied. \square

The results by Crisp about symmetrical subgroups of Artin groups also extend to our case (see [Cri00]).

Corollary 8.5. *Let A denote the Artin group of Euclidean type \widetilde{A}_n or \widetilde{C}_n . For any group G of symmetries of the Artin system, the fixed point subgroup A^G is isomorphic to an Artin group.*

For the explicit description of the Artin group A^G , we refer the reader to [Cri00].

Proof. The proof of [Cri00, Theorem 23] uses only the following properties of the Deligne complex X :

- There exists an A -equivariant consistent geodesic bicombing σ_X on X .
- Subsets of X which are σ_X -stable are contractible.

Let us check the last condition: if $C \subset X$ is a σ_X -stable subset, then σ_X restricts to a convex bicombing on C , which implies that C is contractible. \square

One particular case of interest is when A is of Euclidean type \widetilde{A}_n , and the group G of symmetries of the Artin system (the n -cycle) is generated by a symmetry of the cycle defining A . We recover the result from Digne (see [Dig12]) that the fixed subgroup is isomorphic to the Artin group of type $\widetilde{C}_{\lfloor n/2 \rfloor}$.

For any divisor $1 < k \leq n$ of n , one may also consider the group G of symmetries of the Artin system of type \widetilde{A}_n (the n -cycle) generated by a rotation of k . The fixed point subgroup is then isomorphic to the Artin group of Euclidean type \widetilde{A}_k . When we see $A(\widetilde{A}_n)$ as the group of braids with n strands on the annulus, we may think of the fixed subgroup as the subgroup of braids which are invariant by a rotation of the annulus of angle $\frac{2\pi k}{n}$.

Thomas Haettel
IMAG, Univ Montpellier, CNRS, France
thomas.haettel@umontpellier.fr

References

- [AB95] J. M. ALONSO & M. R. BRIDSON – “Semihyperbolic groups”, *Proc. London Math. Soc. (3)* **70** (1995), no. 1, p. 56–114.
- [Bes03] D. BESSIS – “The dual braid monoid”, *Ann. Sci. École Norm. Sup. (4)* **36** (2003), no. 5, p. 647–683.
- [Bes06] —, “A dual braid monoid for the free group”, *J. Algebra* **302** (2006), no. 1, p. 55–69.
- [BH99] M. R. BRIDSON & A. HAEFLIGER – Grund. math. Wiss., *Metric spaces of non-positive curvature* **319**, Springer, 1999.
- [BM10] T. BRADY & J. MCCAMMOND – “Braids, posets and orthoschemes”, *Algebr. Geom. Topol.* **10** (2010), no. 4, p. 2277–2314.
- [Bou02] N. BOURBAKI – *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley.
- [BT84] F. BRUHAT & J. TITS – “Schémas en groupes et immeubles des groupes classiques sur un corps local”, *Bull. Soc. Math. France* **112** (1984), p. 259–301.
- [CCG⁺20] J. CHALOPIN, V. CHEPOI, A. GENEVOIS, H. HIRAI & D. OSAJDA – “Helly groups”, *arXiv preprint arXiv:2002.06895* (2020).
- [CCHO21] J. CHALOPIN, V. CHEPOI, H. HIRAI & D. OSAJDA – “Weakly modular graphs and nonpositive curvature”, *Mem. Amer. Math. Soc.* (2021).
- [CD95a] R. CHARNEY & M. W. DAVIS – “Finite $K(\pi, 1)$ s for Artin groups”, *Ann. of Math. Stud.*, in *Prospects in topology (Princeton, NJ, 1994)* **138**, Princeton Univ. Press, 1995, p. 110–124.
- [CD95b] —, “The $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups”, *J. Amer. Math. Soc.* **8** (1995), no. 3, p. 597–627.
- [CGGMW19] M. CUMPLIDO, V. GEBHARDT, J. GONZÁLEZ-MENESES & B. WIEST – “On parabolic subgroups of Artin-Tits groups of spherical type”, *Adv. Math.* **352** (2019), p. 572–610.

- [Cha] R. CHARNEY – “Problems related to Artin groups”, American Institute of Mathematics, http://people.brandeis.edu/~charney/papers/_probs.pdf.
- [Cha95] —, “Geodesic automation and growth functions for Artin groups of finite type”, *Math. Ann.* **301** (1995), no. 2, p. 307–324.
- [Cha04] —, “The Deligne complex for the four-strand braid group”, *Trans. Amer. Math. Soc.* **356** (2004), no. 10, p. 3881–3897.
- [Cri00] J. CRISP – “Symmetrical subgroups of Artin groups”, *Adv. Math.* **152** (2000), no. 1, p. 159–177.
- [Deh15] P. DEHORNOY – EMS Tracts in Mathematics, *Foundations of Garside theory* **22**, European Mathematical Society (EMS), Zürich, 2015, With François Digne, Eddy Godelle, Daan Krammer and Jean Michel, Contributor name on title page: Daan Kramer.
- [Del72] P. DELIGNE – “Les immeubles des groupes de tresses généralisés”, *Invent. Math.* **17** (1972), p. 273–302.
- [Dig12] F. DIGNE – “A Garside presentation for Artin-Tits groups of type \widetilde{C}_n ”, *Ann. Inst. Fourier (Grenoble)* **62** (2012), no. 2, p. 641–666.
- [DL15] D. DESCOMBES & U. LANG – “Convex geodesic bicomings and hyperbolicity”, *Geom. Dedicata* **177** (2015), p. 367–384.
- [DL16] —, “Flats in spaces with convex geodesic bicomings”, *Anal. Geom. Metr. Spaces* **4** (2016), no. 1, p. 68–84.
- [DMW] M. DOUGHERTY, J. MCCAMMOND & S. WITZEL – “Boundary braids, journal = arXiv:1811.05603, year = 2018,”.
- [DP99] P. DEHORNOY & L. PARIS – “Gaussian groups and Garside groups, two generalisations of Artin groups”, *Proc. London Math. Soc. (3)* **79** (1999), no. 3, p. 569–604.
- [God07] E. GODELLE – “Artin-Tits groups with CAT(0) Deligne complex”, *J. Pure Appl. Algebra* **208** (2007), no. 1, p. 39–52.
- [GP12] E. GODELLE & L. PARIS – “Basic questions on Artin-Tits groups”, CRM Series, in *Configuration spaces* **14**, Ed. Norm., Pisa, 2012, p. 299–311.
- [Hae21] T. HAETTEL – “Injective metrics on buildings and symmetric spaces”, arXiv:2101.09367, 2021.
- [HHP20] T. HAETTEL, N. HODA & H. PETYT – “The coarse Helly property, hierarchical hyperbolicity, and semihyperbolicity”, *arXiv preprint arXiv:2009.14053* (2020).
- [HKS16] T. HAETTEL, D. KIELAK & P. SCHWER – “The 6-strand braid group is CAT(0)”, *Geom. Dedicata* **182** (2016), p. 263–286.
- [HO19] J. HUANG & D. OSAJDA – “Helly meets Garside and Artin”, *arXiv:1904.09060* (2019).
- [Hod20] N. HODA – “Crystallographic Helly groups”, *arXiv preprint arXiv:2010.07407* (2020).

- [Jeo20] S. G. JEONG – “The seven-strand braid group is CAT(0)”, arXiv:2009.09350, 2020.
- [KR17] D. KASPROWSKI & H. RÜPING – “The Farrell-Jones conjecture for hyperbolic and CAT(0)-groups revisited”, *J. Topol. Anal.* **9** (2017), no. 4, p. 551–569.
- [Lan13] U. LANG – “Injective hulls of certain discrete metric spaces and groups”, *J. Topol. Anal.* **5** (2013), no. 3, p. 297–331.
- [LM21] I. LEARY & A. MINASYAN – “Commensurating HNN-extensions: non-positive curvature and biautomaticity”, *Geom. Topol.* (2021).
- [McC17] J. MCCAMMOND – “The mysterious geometry of Artin groups”, (2017), <http://web.math.ucsb.edu/~jon.mccammond/papers/mysterious-geometry.pdf>.
- [Mic99] J. MICHEL – “A note on words in braid monoids”, *J. Algebra* **215** (1999), no. 1, p. 366–377.
- [Mie17] B. MIESCH – “The Cartan-Hadamard theorem for metric spaces with local geodesic bicomings”, *Enseign. Math.* **63** (2017), no. 1-2, p. 233–247.
- [MS17] J. MCCAMMOND & R. SULWAY – “Artin groups of Euclidean type”, *Invent. Math.* **210** (2017), no. 1, p. 231–282.
- [MW19] R. MORRIS-WRIGHT – “Parabolic subgroups of Artin groups of FC type”, arXiv:1906.07058, 2019.
- [Oko79] C. OKONEK – “Das $K(\pi, 1)$ -Problem für die affinen Wurzelsysteme vom Typ A_n, C_n ”, *Math. Z.* **168** (1979), no. 2, p. 143–148.
- [OV20] D. OSAJDA & M. VALIUNAS – “Helly groups, coarse Helly groups, and relative hyperbolicity”, arXiv:2012.03246, 2020.
- [PS21] G. PAOLINI & M. SALVETTI – “Proof of the $K(\pi, 1)$ conjecture for affine Artin groups”, *Invent. Math.* **224** (2021), no. 2, p. 487–572.
- [Ś06] J. ŚWIĄTKOWSKI – “Regular path systems and (bi)automatic groups”, *Geom. Dedicata* **118** (2006), p. 23–48.
- [Tit66] J. TITS – “Normalisateurs de tores. I. Groupes de Coxeter étendus”, *J. Algebra* **4** (1966), p. 96–116.
- [Tit74] J. TITS – *Buildings of spherical type and finite BN-pairs*, Lect. Notes in Math. **386**, Springer, 1974.
- [VdL83] H. VAN DER LEK – “The homotopy type of complex hyperplane complements”, (1983), PhD thesis, Radboud University of Nijmegen.