

G -TORSORS AND UNIVERSAL TORSORS OVER NONSPLIT DEL PEZZO SURFACES

ULRICH DERENTHAL AND NORBERT HOFFMANN

ABSTRACT. Let S be a smooth del Pezzo surface that is defined over a field K and splits over a Galois extension L . Let G be either the split reductive group given by the root system of S_L in $\text{Pic } S_L$, or a form of it containing the Néron–Severi torus. Let \mathcal{G} be the G -torsor over S_L obtained by extension of structure group from a universal torsor \mathcal{T} over S_L . We prove that \mathcal{G} does not descend to S unless \mathcal{T} does. This is in contrast to a result of Friedman and Morgan that such \mathcal{G} always descend to singular del Pezzo surfaces over \mathbb{C} from their desingularizations.

1. INTRODUCTION

One of the most famous theorems of 19th century algebraic geometry states that cubic surfaces over \mathbb{C} contain precisely 27 lines. The symmetry group of their configuration is the Weyl group W of the root system \mathbf{E}_6 .

Cubic surfaces over fields K that are not algebraically closed exhibit a substantially wider range of phenomena. Given such a cubic surface S over K , there is a finite Galois extension L/K over which S splits (i.e., over which all 27 lines are defined). Its Galois group Γ acts on the configuration of lines via W . This action plays a major role in the arithmetic of S .

An important tool in the study of such cubic surfaces S are the universal torsors introduced and studied by Colliot-Thélène and Sansuc [2]. These are certain torsors over S under the Néron–Severi torus. However, such universal torsors do not always exist in the nonsplit case, and they do not descend to singular cubic surfaces from their desingularizations. In particular, this does not provide degenerations of universal torsors to singular cubic surfaces.

Friedman and Morgan suggested to replace the Néron–Severi torus here by a reductive algebraic group G of type \mathbf{E}_6 . They construct natural G -torsors over singular cubic surfaces over \mathbb{C} [5]. These G -torsors are compatible with degenerations of split cubic surfaces [4].

In this paper, we deal with the existence of natural G -torsors over nonsplit smooth cubic surfaces. Here, the natural candidates for G are the split group of type \mathbf{E}_6 , and forms of it that contain the Néron–Severi torus, because both reflect the combinatorial information coming from S . Motivated by the result of Friedman and Morgan, there has been a hope that such G -torsors could exist also in situations where universal torsors do not exist. However, the main result of the present paper is that such G -torsors do not exist unless universal torsors exist.

Date: September 16, 2021.

2020 Mathematics Subject Classification. 14J26 (14L30, 11E57, 14G27).

Key words and phrases. Universal torsor, del Pezzo surface, reductive group.

More generally, let S be a smooth del Pezzo surface over a field K . Let L/K be a finite Galois extension such that the base change S_L is split. Then the Galois group $\Gamma = \text{Gal}(L/K)$ acts on $\Lambda_S := \text{Pic } S_L$.

Manin [7] discovered that the Picard group Λ_S of the split del Pezzo surface S_L together with the intersection form $(-, -)$ provide us with a reduced root datum $(\Phi_S \subset \Lambda_S, \Phi_S^\vee \subset \Lambda_S^\vee)$ given as follows: The set of roots is

$$(1) \quad \Phi_S := \{\alpha \in \Lambda_S \mid (\alpha, \alpha) = -2, (\alpha, -K_{S_L}) = 0\}.$$

With $\alpha^\vee \in \Lambda_S^\vee$ given by $\langle \alpha^\vee, \lambda \rangle := (-\alpha, \lambda)$ for $\lambda \in \Lambda_S$, the set of coroots is

$$\Phi_S^\vee := \{\alpha^\vee \in \Lambda_S^\vee \mid \alpha \in \Phi_S\}.$$

Its Weyl group $W_S \subset \text{Aut } \Lambda_S$ is the symmetry group of the configuration of (-1) -curves on S_L .

The Néron–Severi torus for S_L is the split L -torus T with character lattice $\text{Hom}(T, \mathbb{G}_{m,L}) = \Lambda_S$ as abelian groups. A universal torsor for S_L is a T -torsor \mathcal{T} over S_L whose type $\text{Hom}(T, \mathbb{G}_m) \rightarrow \text{Pic } S_L$ (in the sense of [2], see Section 3) is the identity on Λ_S . Such a \mathcal{T} exists and is unique up to isomorphism.

The Néron–Severi torus for S is the K -torus T that splits over L with character lattice $\text{Hom}(T_L, \mathbb{G}_{m,L}) = \Lambda_S$ as Γ -modules. A universal torsor for S is a T -torsor \mathcal{T} over S whose base change \mathcal{T}_L is a universal torsor for S_L . Such a \mathcal{T} may or may not exist, and may or may not be unique [2].

In this situation, our main result on the existence of G -torsors is:

Theorem. *Let T be*

- (i) *the split torus over K with character lattice $\text{Hom}(T, \mathbb{G}_{m,K}) = \Lambda_S$, or*
- (ii) *the Néron–Severi torus for S over K .*

Let G be a reductive group over K with maximal torus $\iota : T \hookrightarrow G$ and resulting reduced root datum $(\Phi_S \subset \Lambda_S, \Phi_S^\vee \subset \Lambda_S^\vee)$.

Let \mathcal{T} be a universal T_L -torsor over S_L , and let $\mathcal{G} := (\iota_L)_ \mathcal{T}$ be the G_L -torsor over S_L obtained by extension of structure group. Then \mathcal{G} descends to a G -torsor over S if and only if universal torsors for S exist.*

We prove this as an equivalence of categories. Part (ii) is contained in Theorem 4.2 below, and part (i) is contained in its Corollary 4.3.

The proof of Theorem 4.2 is based on descent from the split situation. It turns out that the relevant T -torsors and G -torsors have the same descent data because they have the same automorphism group. Corollary 4.3 then follows using that the split group G over K contains a Néron–Severi torus, a result due independently to Gille [6] and Raghunathan [8].

The structure of our paper is as follows. In Section 2, we collect a few basic facts about G -torsors, with a view towards their descent. We circumvent the difficulty of two Galois actions (on S_L and G_L) as follows: Instead of working with G_L -torsors over S_L , we view them as G -torsors over S_L regarded as a K -scheme. In Section 3, we compare automorphisms of T -torsors and G -torsors in the split case. Finally, in Section 4, we apply our descent setup to prove the main results.

Acknowledgments. We learned about this question from Yuri Tschinkel. The first author was partly supported by grant DE 1646/4-2 of the Deutsche Forschungsgemeinschaft. Some of this work was done while the second author was on sabbatical leave at the Riemann Center for Geometry and Physics of Leibniz Universität Hannover, supported by Mary Immaculate College Limerick through the PLOA program.

2. TORSORS

Let X be a smooth projective scheme over K . We assume that X is connected (but not necessarily geometrically connected over K). Let G be a linear algebraic group over K .

Definition 2.1. A G -torsor over X is a faithfully flat morphism $\mathcal{G} \rightarrow X$ of finite type together with a right action $\mathcal{G} \times_K G \rightarrow \mathcal{G}$ over X such that the map $\mathcal{G} \times_K G \rightarrow \mathcal{G} \times_X \mathcal{G}$ given by $(u, g) \mapsto (u, ug)$ is an isomorphism.

Definition 2.2. Let $\phi : G \rightarrow H$ be a homomorphism of linear algebraic groups over K . The *extension of structure group of \mathcal{G} along ϕ* is the H -torsor

$$\phi_*\mathcal{G} := \mathcal{G} \times^G H = (\mathcal{G} \times_K H)/G \rightarrow X,$$

where G acts on \mathcal{G} from the right, and on H via ϕ from the left.

Let $f : Y \rightarrow X$ be a morphism of connected smooth projective K -schemes.

Definition 2.3. The *pullback of \mathcal{G} along f* is the G -torsor

$$f^*\mathcal{G} := \mathcal{G} \times_X Y \rightarrow Y.$$

Remark 2.4. Generalizing Definition 2.2, if ϕ is a homomorphism of group schemes over X from $G_X := G \times_K X$ to $H_X := H \times_K X$, then

$$\phi_*\mathcal{G} := (\mathcal{G} \times_X H_X)/G_X$$

is still an H -torsor over X . Its pullback along f satisfies

$$(2) \quad f^*(\phi_*\mathcal{G}) \cong (f^*\phi)_*(f^*\mathcal{G}),$$

where $f^*\phi : G_Y \rightarrow H_Y$ is the pullback of ϕ .

Definition 2.5. A morphism $f : Y \rightarrow X$ between connected smooth projective schemes over K is a *Galois covering* if f is finite, surjective and étale such that the natural map

$$\Sigma \times Y \rightarrow Y \times_X Y, \quad (\sigma, y) \mapsto (\sigma y, y)$$

is an isomorphism for the discrete group $\Sigma := \text{Aut}_X Y$ [1, §6.2B].

Here, each $\sigma \in \Sigma$ is a K -morphism $\sigma : Y \rightarrow Y$. Hence any G -torsor \mathcal{G} over Y can be pulled back along σ , resulting in another G -torsor $\sigma^*\mathcal{G}$ over Y .

Suppose that $\mathcal{G} = f^*\mathcal{G}^\circ$ for some G -torsor \mathcal{G}° over X . For each $\sigma \in \Sigma$, let $c_\sigma : \sigma^*\mathcal{G} \rightarrow \mathcal{G}$ be the canonical isomorphism $\sigma^*\mathcal{G} = \sigma^*f^*\mathcal{G}^\circ \rightarrow f^*\mathcal{G}^\circ = \mathcal{G}$ of G -torsors over Y coming from the fact that $f \circ \sigma = f : Y \rightarrow X$. By construction, they satisfy the cocycle condition

$$(3) \quad c_{\rho\sigma} = c_\sigma \circ \sigma^*(c_\rho)$$

as isomorphisms $(\varrho\sigma)^*\mathcal{G} = \sigma^*\varrho^*\mathcal{G} \rightarrow \sigma^*\mathcal{G} \rightarrow \mathcal{G}$ of G -torsors over Y for every $\varrho, \sigma \in \Sigma$. Conversely, we have the following descent result.

Lemma 2.6. *Given a G -torsor \mathcal{G} over Y , and isomorphisms $c_\sigma : \sigma^*\mathcal{G} \rightarrow \mathcal{G}$ for $\sigma \in \Sigma$ satisfying (3) for all $\varrho, \sigma \in \Sigma$, there is a unique G -torsor \mathcal{G}° over X such that $f^*\mathcal{G}^\circ$ with the induced cocycle is isomorphic to \mathcal{G} with $(c_\sigma)_{\sigma \in \Sigma}$.*

Proof. The relatively affine scheme \mathcal{G} over Y is the relative spectrum of a quasicohherent \mathcal{O}_Y -algebra \mathcal{A} , which descends to a unique quasicohherent \mathcal{O}_X -algebra \mathcal{A}° according to [9, Lemma 0D1V]. We define \mathcal{G}° as the relative spectrum of \mathcal{A}° over X .

The group action $\mathcal{G} \times_K G \rightarrow \mathcal{G}$ corresponds to a morphism $\mathcal{A} \rightarrow \mathcal{A} \otimes_K K[G]$ of quasi-coherent \mathcal{O}_Y -algebras, where $K[G] := \Gamma(G, \mathcal{O}_G)$ is the coordinate ring of G . Since σ is a K -morphism, we have a canonical isomorphism

$$\sigma^*(\mathcal{A} \otimes_K K[G]) = (\sigma^*\mathcal{A}) \otimes_K K[G],$$

which allows us to use [9, Lemma 0D1V] again to descend the given group action to a unique group action $\mathcal{G}^\circ \times_K G \rightarrow \mathcal{G}^\circ$. This turns \mathcal{G}° into a G -torsor over X , as required. \square

3. AUTOMORPHISMS OF T -TORSORS AND G -TORSORS

In this section, we assume that S is a *split* del Pezzo surface over K . Let T be a split torus over K with character lattice Λ_T , and let \mathcal{T} be a T -torsor over S .

Recall from [2] that the *type* of \mathcal{T} is the homomorphism $\tau : \Lambda_T \rightarrow \Lambda_S$ that sends a character $\chi : T \rightarrow \mathbb{G}_m$ to the class $[\chi_*\mathcal{T}]$ of the line bundle given by the \mathbb{G}_m -torsor $\chi_*\mathcal{T}$ over S . For a homomorphism $\phi : T \rightarrow T'$ of split K -tori, the extension of structure group $\phi_*\mathcal{T}$ has type $\tau \circ \phi^* : \Lambda_{T'} \rightarrow \Lambda_T \rightarrow \Lambda_S$.

Let G be a split reductive group over K with maximal torus $\iota : T \hookrightarrow G$ and resulting reduced root datum $(\Phi_G \subset \Lambda_T, \Phi_G^\vee \subset \Lambda_T^\vee)$. We assume that the type $\tau : \Lambda_T \rightarrow \Lambda_S$ of \mathcal{T} is an isomorphism from this root datum to $(\Phi_S \subset \Lambda_S, \Phi_S^\vee \subset \Lambda_S^\vee)$, i.e., τ is bijective, $\tau(\Phi_G) = \Phi_S$, and $\tau^*(\Phi_S^\vee) = \Phi_G^\vee$. This determines G up to isomorphism [3, Exposé XXV, Corollaire 1.2].

Remark 3.1. This group G can be described more explicitly as follows. The anticanonical class $-K_S \in \Lambda_S$ vanishes on all coroots and hence defines a character

$$\chi := \tau^{-1}(-K_S) : G \rightarrow \mathbb{G}_m.$$

If the degree of S is at most 6, then the coroots $\alpha^\vee \in \Lambda_S^\vee$ generate the subgroup $\{\phi \in \Lambda_S^\vee \mid \phi(-K_S) = 0\}$, so we obtain a short exact sequence

$$1 \rightarrow [G, G] \rightarrow G \xrightarrow{\chi} \mathbb{G}_m \rightarrow 1,$$

in which $\ker(\chi)$ is simply connected. Therefore, the commutator subgroup $[G, G]$ is the semisimple and simply connected algebraic group of type \mathbf{E}_8 , \mathbf{E}_7 , \mathbf{E}_6 , \mathbf{D}_5 , \mathbf{A}_4 , $\mathbf{A}_2 + \mathbf{A}_1$ if the degree of S is 1, 2, 3, 4, 5, 6, respectively.

In degree 7, we have $G \cong \mathbb{G}_m \times \mathrm{GL}_2$. In degree 8, we have $G \cong \mathrm{GL}_2$ if $S \cong \mathbb{P}^1 \times \mathbb{P}^1$, and $G \cong \mathbb{G}_m^2$ if S is the blow-up of \mathbb{P}^2 in a point. Finally, in degree 9, we have $G \cong \mathbb{G}_m$.

Let $\mathcal{G} := \iota_*\mathcal{T}$ denote the G -torsor over S obtained from \mathcal{T} by extension of structure group. We can view the total space \mathcal{T} as contained in \mathcal{G} .

Proposition 3.2. *Let \mathcal{T}_1 be another T -torsor over S of the same type τ as \mathcal{T} , and let $\mathcal{G}_1 = \iota_* \mathcal{T}_1$. Every isomorphism $\mathcal{G}_1 \rightarrow \mathcal{G}$ of G -torsors over S restricts to an isomorphism $\mathcal{T}_1 \rightarrow \mathcal{T}$ of T -torsors over S .*

Proof. We have $\mathcal{T} \cong \mathcal{T}_1$ since both are torsors of the same type τ under the split torus T . Choosing one such isomorphism, it suffices to show that the natural map

$$T = \text{Aut } \mathcal{T} \rightarrow \text{Aut } \mathcal{G}$$

is an isomorphism. Here the automorphism group $\text{Aut } \mathcal{G}$ of the G -torsor \mathcal{G} over S is a linear algebraic group over K . To see this, choose a closed embedding $G \hookrightarrow \text{GL}_n$, and let $\mathcal{V} := \mathcal{G} \times^G \mathbb{A}^n$ denote the associated vector bundle over S . Then $\text{Aut } \mathcal{G}$ is a closed subgroup of $\text{Aut } \mathcal{V}$, which is the unit group of the finite-dimensional K -algebra $\text{End } \mathcal{V}$. Hence $\text{Aut } \mathcal{G}$ is indeed a linear algebraic group over K .

The Lie algebra of $\text{Aut } \mathcal{G}$ is

$$H^0(S, \mathcal{G} \times^G \mathfrak{g}) = H^0(S, \mathcal{T} \times^T \mathfrak{g}),$$

where \mathfrak{g} denotes the Lie algebra of G . Let

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \bigoplus_{\alpha \in \Phi_G} \mathfrak{g}_\alpha$$

be the root space decomposition of \mathfrak{g} into its eigenspaces under T . For any root $\alpha \in \Phi_G$, the line bundle

$$L_\alpha := \mathcal{T} \times^T \mathfrak{g}_\alpha$$

over S has as isomorphism class $\tau(\alpha)$ since \mathcal{T} has type τ . In particular, the anticanonical degree of L_α is $(-K_S, \tau(\alpha)) = 0$, and $L_\alpha \not\cong \mathcal{O}_S$. Therefore, $H^0(S, L_\alpha) = 0$ for each root α , and hence

$$H^0(S, \mathcal{T} \times^T \mathfrak{g}) = H^0(S, \mathcal{T} \times^T \mathfrak{g}_0) = H^0(S, \mathcal{O}_S \otimes \mathfrak{g}_0) = \mathfrak{g}_0 = \mathfrak{t},$$

the Lie algebra of T . This shows that the natural embedding $T \hookrightarrow \text{Aut } \mathcal{G}$ induces an isomorphism on tangent spaces.

Since T is smooth, it follows that T is an open subgroup of $\text{Aut } \mathcal{G}$. Being also connected, T is normal in $\text{Aut } \mathcal{G}$. As a global section of the automorphism group scheme $\mathcal{G} \times^G G$ over S , where G acts on itself by conjugation, each element in $\text{Aut } \mathcal{G}$ therefore lies already in

$$\mathcal{G} \times^G N = \mathcal{T} \times^T N$$

for the normalizer N of T in G . Since T acts trivially on the Weyl group $W_G := N/T$, we have an exact sequence

$$1 \rightarrow \mathcal{T} \times^T T \rightarrow \mathcal{T} \times^T N \rightarrow W_G \rightarrow 1.$$

Suppose that $w \in W_G$ is in the image of $\text{Aut } \mathcal{G}$. Then \mathcal{T} is isomorphic to its extension of structure group $w_* \mathcal{T}$ along the automorphism $w_* : T \rightarrow T$. As \mathcal{T} has type τ , the type of $w_* \mathcal{T}$ is $\tau \circ w^* : \Lambda_T \rightarrow \Lambda_T \rightarrow \Lambda_S$. Since isomorphic torsors must have the same type, we conclude that $w = 1$.

Hence every automorphism of \mathcal{G} is a global section of $\mathcal{T} \times^T T = S \times T$, and thus an element of T . \square

4. PROOF OF THE MAIN RESULTS

We use the notation of Section 1. In particular, S and S_L are both connected smooth projective schemes over K (even though S_L is not geometrically connected over K unless $L = K$), and the projection $\pi : S_L \rightarrow S$ is a K -morphism. More precisely, π is a Galois covering with group

$$\Sigma = \{\text{id}_S \times \gamma^* \mid \gamma \in \Gamma\} \cong \Gamma^{\text{op}}$$

in the sense of Definition 2.5.

Let \mathcal{G} be a G -torsor over the K -scheme S_L , as in Definition 2.1. Then the same total space \mathcal{G} can also be viewed as a G_L -torsor over the L -scheme S_L . Indeed, the required action $\mathcal{G} \times_L G_L \rightarrow \mathcal{G}$ of G_L comes from the given action $\mathcal{G} \times_K G \rightarrow \mathcal{G}$ of G , since the two fiber products are canonically isomorphic and the axioms are clearly equivalent. Conversely, every G_L -torsor over the L -scheme S_L can be viewed as a G -torsor over the K -scheme S_L .

Let T be a K -torus that splits over L with character lattice Λ_T , and let \mathcal{T} be a T -torsor over S_L . Then we can in particular view \mathcal{T} also as a T_L -torsor over S_L , which has a type $\tau : \Lambda_T \rightarrow \Lambda_S$ in the sense of Section 3.

Lemma 4.1. *For $\gamma \in \Gamma$, put $\sigma = \text{id}_S \times \gamma^* \in \Sigma$. Then the T -torsor $\sigma^*\mathcal{T}$ over S_L has type*

$$\sigma^* \circ \tau \circ (\gamma_*^T)^{-1} : \Lambda_T \rightarrow \Lambda_S,$$

where $\gamma_*^T : \Lambda_T \rightarrow \Lambda_T$ is induced by γ .

Proof. Every element of Λ_T has the form $\gamma_*^T \chi$ for a unique $\chi \in \Lambda_T$. The type of $\sigma^*\mathcal{T}$ sends $\gamma_*^T \chi$ to the class of the line bundle

$$(\gamma_*^T \chi)_*(\sigma^*\mathcal{T}) \cong \sigma^*(\chi_*\mathcal{T})$$

over S_L , where the isomorphism is a special case of (2). This class is $(\sigma^* \circ \tau)(\chi)$, as required. \square

Theorem 4.2. *Let $\iota : T \hookrightarrow G$ be a maximal torus in a reductive group over K such that T splits over L , with resulting root datum $(\Phi_G \subset \Lambda_T, \Phi_G^\vee \subset \Lambda_T^\vee)$. Let \mathcal{T} be a T -torsor over S_L whose type $\tau : \Lambda_T \rightarrow \Lambda_S$ is a Γ -equivariant isomorphism from this root datum to $(\Phi_S \subset \Lambda_S, \Phi_S^\vee \subset \Lambda_S^\vee)$. We consider the G -torsor $\mathcal{G} := \iota_*\mathcal{T}$ over S_L obtained by extension of structure group.*

Then the functor ι_ from the groupoid of T -torsors \mathcal{T}° over S such that $\mathcal{T}_L^\circ \cong \mathcal{T}$ to the groupoid of G -torsors \mathcal{G}° over S such that $\mathcal{G}_L^\circ \cong \mathcal{G}$ is an equivalence of categories.*

Proof. Let \mathcal{T}° be a T -torsor over S such that $\mathcal{T}_L^\circ \cong \mathcal{T}$. Let $\mathcal{G}^\circ := \iota_*\mathcal{T}^\circ$. Then the base change of \mathcal{G}° to L is

$$\mathcal{G}_L^\circ = (\iota_*\mathcal{T}^\circ)_L \cong \iota_*(\mathcal{T}_L^\circ) \cong \iota_*\mathcal{T} = \mathcal{G}.$$

Clearly any isomorphism ϕ between such T -torsors over S induces an isomorphism $\iota_*\phi$ between G -torsors over S . Hence ι_* is indeed a functor.

Let \mathcal{T}^\bullet be another T -torsor over S such that $\mathcal{T}_L^\bullet \cong \mathcal{T}$, and $\mathcal{G}^\bullet := \iota_*\mathcal{T}^\bullet$. Let $\psi : \mathcal{G}^\bullet \rightarrow \mathcal{G}^\circ$ be an isomorphism of G -torsors over S . Then Proposition 3.2 applies to the base change

$$\psi_L : \iota_*(\mathcal{T}_L^\bullet) = \mathcal{G}_L^\bullet \rightarrow \mathcal{G}_L^\circ = \iota_*(\mathcal{T}_L^\circ)$$

and shows that ψ_L restricts to an isomorphism $\mathcal{T}_L^\bullet \rightarrow \mathcal{T}_L^\circ$. In particular, $\psi_L(\mathcal{T}_L^\bullet) = \mathcal{T}_L^\circ$ as closed subschemes of \mathcal{G}_L° , and hence $\psi(\mathcal{T}^\bullet) = \mathcal{T}^\circ$ as closed subschemes of \mathcal{G}° . The restriction $\mathcal{T}^\bullet \rightarrow \mathcal{T}^\circ$ of ψ becomes an isomorphism of T -torsors over S_L after base change, and therefore is an isomorphism of T -torsors over S . This proves that the functor ι_* is fully faithful.

Now assume that the G -torsor $\mathcal{G} = \iota_*\mathcal{T}$ descends to a G -torsor \mathcal{G}° over S . As explained in Section 2, we have an isomorphism

$$c_\sigma : \sigma^*\mathcal{G} \rightarrow \mathcal{G}$$

of G -torsors over S_L for each $\sigma \in \Sigma$, satisfying the cocycle conditions (3). Here, $\sigma^*\mathcal{G} = \iota_*\sigma^*\mathcal{T}$. Since τ is Γ -equivariant, $\sigma^*\mathcal{T}$ has the same type τ as \mathcal{T} due to Lemma 4.1. Therefore, Proposition 3.2 shows that c_σ restricts to an isomorphism

$$c'_\sigma : \sigma^*\mathcal{T} \rightarrow \mathcal{T}.$$

The c'_σ satisfy the cocycle condition because the c_σ do. Therefore, \mathcal{T} descends to a T -torsor \mathcal{T}° over S . By construction, $\iota_*\mathcal{T}^\circ \cong \mathcal{G}^\circ$. This proves that the functor ι_* is essentially surjective. \square

Corollary 4.3. *Let $\iota : T \hookrightarrow G$ be a split maximal torus in a split reductive group over K , with resulting root datum $(\Phi_G \subset \Lambda_T, \Phi_G^\vee \subset \Lambda_T^\vee)$. Let \mathcal{T} be a T -torsor over S_L whose type $\tau : \Lambda_T \rightarrow \Lambda_S$ is a \mathbb{Z} -linear isomorphism from this root datum to $(\Phi_S \subset \Lambda_S, \Phi_S^\vee \subset \Lambda_S^\vee)$. We consider the G -torsor $\mathcal{G} := \iota_*\mathcal{T}$ over S_L obtained by extension of structure group.*

Then the groupoid of G -torsors \mathcal{G}° over S such that $\mathcal{G}_L^\circ \cong \mathcal{G}$ is equivalent to the groupoid of universal torsors over S .

Proof. Let $N \subset G$ denote the normalizer of T in G . The exact sequence

$$1 \rightarrow T \rightarrow N \rightarrow W_G \rightarrow 1$$

defines the Weyl group W_G of G . For $g \in G(L)$, we denote by

$$\text{conj}_g := g \cdot - \cdot g^{-1} : G_L \rightarrow G_L$$

the conjugation. For $n \in N(L)$, conj_n restricts to an automorphism of T_L , which depends only on the image of n in W_G . This defines a left action of W_G on T_L , and consequently a right action of W_G on Λ_T , where $w \in W_G$ acts via the pullback of characters $w^* : \Lambda_T \rightarrow \Lambda_T$. Sending $w \in W_G$ to $(w^{-1})^* : \Lambda_T \rightarrow \Lambda_T$, we identify W_G with the Weyl group $W(\Phi_G) \subset \text{Aut } \Lambda_T$ of our root datum of G .

Since τ is an isomorphism of root data, it induces an isomorphism

$$\tau_W : W_G \rightarrow W_S,$$

where $W_S := W(\Phi_S) \subset \text{Aut } \Lambda_S$ is the Weyl group of our root datum of S .

The Galois action of Γ on Λ_S factors through the Weyl group action of W_S on Λ_S via a natural map

$$(4) \quad \rho_S : \Gamma \rightarrow W_S.$$

Applying [6, Théorème 5.1(b)] or [8, Theorem 1.1] to the semisimple commutator subgroup $G^c := [G, G] \subset G$, there is a $g \in G^c(L) \subset G(L)$ such that, for each $\gamma \in \Gamma$, the element $g^{-1}\gamma(g) \in G(L)$ is in $N(L)$, and τ_W maps its image in W_G to $\rho_S(\gamma)$. The former implies that there is a unique maximal

torus $\iota' : T' \hookrightarrow G$ over K such that $T'_L = gT_Lg^{-1}$, and the latter says that the square on the right in the following diagram commutes:

$$(5) \quad \begin{array}{ccccc} \Lambda_{T'} & \xrightarrow{\text{conj}_g^*} & \Lambda_T & \xrightarrow{\tau} & \Lambda_S \\ \downarrow \gamma_*^{T'} & & \downarrow \text{conj}_{\gamma(g^{-1})}^* & & \downarrow \rho_S(\gamma) \\ \Lambda_{T'} & \xrightarrow{\text{conj}_g^*} & \Lambda_T & \xrightarrow{\tau} & \Lambda_S \end{array}$$

In order to determine the Galois action $\gamma_*^{T'}$ coming from the fact that T' is defined over K , we note that the commutative diagram

$$\begin{array}{ccccc} G_L & \xrightarrow{\gamma(g)\cdot-} & G_L & \xrightarrow{-\gamma(g^{-1})} & G_L \\ \downarrow \gamma^* & & \downarrow \gamma^* & & \downarrow \gamma^* \\ G_L & \xrightarrow{g\cdot-} & G_L & \xrightarrow{-g^{-1}} & G_L \end{array}$$

restricts to a commutative diagram

$$\begin{array}{ccc} T_L & \xrightarrow{\text{conj}_{\gamma(g)}} & T'_L \\ \downarrow \gamma^* & & \downarrow \gamma^* \\ T_L & \xrightarrow{\text{conj}_g} & T'_L, \end{array}$$

which in turn induces the commutative diagram of character lattices

$$\begin{array}{ccc} \Lambda_T & \xleftarrow{\text{conj}_{\gamma(g)}^*} & \Lambda_{T'} \\ \uparrow \gamma_*^T & & \uparrow \gamma_*^{T'} \\ \Lambda_T & \xleftarrow{\text{conj}_g^*} & \Lambda_{T'}. \end{array}$$

Since T is split over K , the map γ_*^T here is the identity, and hence

$$\gamma_*^{T'} = \text{conj}_{g\gamma(g^{-1})}^*.$$

This shows that the square on the left in (5) commutes as well. Therefore, the composition

$$(6) \quad \Lambda_{T'} \xrightarrow{\text{conj}_g^*} \Lambda_T \xrightarrow{\tau} \Lambda_S$$

is Galois-equivariant. In particular, T' is a Néron–Severi torus.

Now Theorem 4.2 applies to $\iota' : T' \hookrightarrow G$, the T' -torsor $\mathcal{T}' := (\text{conj}_g)_* \mathcal{T}$ and the resulting G -torsor $\mathcal{G}' := \iota'_* \mathcal{T}'$ over S_L . Indeed, the type of \mathcal{T}' is by construction the composition (6). This composition is Galois-equivariant, as we have just seen. It is also an isomorphism of root data since conj_g^* preserves the root datum of G and τ respects the relevant root data by assumption. So the hypotheses of Theorem 4.2 are satisfied.

Therefore, the functor ι'_* is an equivalence from the groupoid of T' -torsors \mathcal{T}° over S such that $\mathcal{T}_L^\circ \cong \mathcal{T}'$ to the groupoid of G -torsors \mathcal{G}° over S such that $\mathcal{G}_L^\circ \cong \mathcal{G}'$. The former groupoid is equivalent to the groupoid of universal torsors over S because T' is a Néron–Severi torus and the type of \mathcal{T}' is a Galois-equivariant isomorphism. The latter groupoid is the groupoid in the claim because $\mathcal{G}' \cong \mathcal{G}$. Indeed, $\mathcal{G}' = (\text{conj}_g)_* \mathcal{G}$ because $\iota'_L \circ \text{conj}_g = \text{conj}_g \circ \iota_L$, and conj_g is an inner automorphism of G_L . \square

REFERENCES

- [1] S. Bosch, W. Lütkebohmert, and M. Raynaud. *Néron models*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1990.
- [2] J.-L. Colliot-Thélène and J.-J. Sansuc. La descente sur les variétés rationnelles. II. *Duke Math. J.*, 54(2):375–492, 1987.
- [3] M. Demazure and A. Grothendieck. *Schémas en groupes (SGA 3). Tome III. Structure des schémas en groupes réductifs*, volume 153 of *Lecture Notes in Mathematics*. Springer-Verlag, Heidelberg, 1970.
- [4] U. Derenthal and N. Hoffmann. Degeneration of torsors over families of del Pezzo surfaces. *Math. Nachr.*, 293(12):2306–2334, 2020.
- [5] R. Friedman and J. W. Morgan. Exceptional groups and del Pezzo surfaces. In *Symposium in Honor of C. H. Clemens (Salt Lake City, UT, 2000)*, volume 312 of *Contemp. Math.*, pages 101–116. Amer. Math. Soc., Providence, RI, 2002.
- [6] P. Gille. Type des tores maximaux des groupes semi-simples. *J. Ramanujan Math. Soc.*, 19(3):213–230, 2004.
- [7] Yu. I. Manin. *Cubic forms*, volume 4 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, second edition, 1986.
- [8] M. S. Raghunathan. Tori in quasi-split-groups. *J. Ramanujan Math. Soc.*, 19(4):281–287, 2004.
- [9] The Stacks project authors. The Stacks project. <https://stacks.math.columbia.edu>, 2021.

INSTITUT FÜR ALGEBRA, ZAHLENTHEORIE UND DISKRETE MATHEMATIK, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER, GERMANY
Email address: `derenthal@math.uni-hannover.de`

DEPARTMENT OF MATHEMATICS AND COMPUTER STUDIES, MARY IMMACULATE COLLEGE, SOUTH CIRCULAR ROAD, LIMERICK, IRELAND
Email address: `norbert.hoffmann@mic.ul.ie`