

# Families of polytopes with rational linear precision in higher dimensions

Isobel Davies, Eliana Duarte, Irem Portakal, Miruna-Ştefana Sorea

September 20, 2021

## Abstract

In this article we introduce a new family of lattice polytopes with rational linear precision. For this purpose, we define a new class of discrete statistical models that we call multinomial staged tree models. We prove that these models have rational maximum likelihood estimators (MLE) and give a criterion for these models to be log-linear. Our main result is then obtained by applying Garcia-Puente and Sottile's theorem that establishes a correspondence between polytopes with rational linear precision and log-linear models with rational MLE. Throughout this article we also study the interplay between the primitive collections of the normal fan of a polytope with rational linear precision and the shape of the Horn matrix of its corresponding statistical model. Finally, we investigate lattice polytopes arising from toric multinomial staged tree models, in terms of the combinatorics of their tree representations.

## 1 Introduction

In Geometric Modeling, pieces of parametrised curves and surfaces are used as building blocks to describe geometric shapes in 2D and 3D. Some of the most widely used parametric units for this purpose are Bézier curves, triangular Bézier surfaces and tensor product surfaces. These pieces of curves and surfaces are constructed using a set of polynomial blending functions defined on the convex hull of a set of points  $\mathcal{A}$ , together with a set of control points. Taking as inspiration the theory of toric varieties and the form of the blending functions for the previous examples, Krasauskas introduced the more general notion of a *toric patch* whose domain is a lattice polytope  $P \subseteq \mathbb{R}^d$  [14]. The blending functions,  $\{\beta_{w,m} : P \rightarrow \mathbb{R}\}_{m \in \mathcal{A}}$ , of a toric patch are constructed from the set of lattice points  $\mathcal{A} := P \cap \mathbb{Z}^d$  and a vector of positive weights  $w$  associated to each point in  $\mathcal{A}$ .

A significant difference between an arbitrary toric patch and one of the triangular or tensor product patches is that its blending functions do not necessarily satisfy the property

---

*MSC2020 Subject Classification:* Primary: 52B20; Secondary: 14M25, 62R01, 65D17.

*Key words and phrases:* lattice polytope, Horn parametrisation, rational linear precision, primitive collection, log-linear model, maximum likelihood estimator, toric variety, staged tree.

of *linear precision*. The collection of blending functions  $\{\beta_{w,m} : P \rightarrow \mathbb{R}\}_{m \in \mathcal{A}}$  has linear precision if for any affine function  $\Lambda : \mathbb{R}^d \rightarrow \mathbb{R}$ ,

$$\Lambda(u) = \sum_{m \in \mathcal{A}} \Lambda(m) \beta_{w,m}(u), \text{ for all } u \in P.$$

Thus, linear precision is the ability of the blending functions to replicate affine functions and it is desirable from the practical standpoint [10]. If a pair  $(P, w)$  has *rational linear precision* then its blending functions  $\{\beta_{w,m} : P \rightarrow \mathbb{R}\}_{m \in \mathcal{A}}$  admit a rational reparametrisation  $\{\hat{\beta}_{w,m} : P \rightarrow \mathbb{R}\}_{m \in \mathcal{A}}$  with linear precision. It is an open problem, motivated by Geometric Modeling, to characterise all pairs  $(P, w)$  that have rational linear precision in dimension  $d \geq 3$  [4, 14]. The classification of all such pairs in dimension  $d = 2$  is given in [11].

Garcia-Puente and Sottile studied the property of rational linear precision for toric patches by associating a scaled projective toric variety  $X_{\mathcal{A},w}$  to the pair  $(P, w)$  [10]. The variety  $X_{\mathcal{A},w}$  is the image of the map  $[w\chi]_{\mathcal{A}} : (\mathbb{C}^*)^d \rightarrow \mathbb{P}^{s-1}$  defined by  $t \mapsto [w_1 t^{m_1} : w_2 t^{m_2} : \dots : w_s t^{m_s}]$  where  $\mathcal{A} = \{m_1, \dots, m_s\}$ . One of their main results states that a pair  $(P, w)$  has rational linear precision if and only if the variety  $X_{\mathcal{A},w}$ , seen as a discrete statistical model, has rational maximum likelihood estimator (MLE). This result establishes a communication channel between Geometric Modeling and Algebraic Statistics. Thus, it is natural to use ideas from Algebraic Statistics to study the property of rational linear precision.

Models with rational MLE are algebraic varieties that admit a special parametrisation known as Horn uniformisation [9, 13]. This parametrisation depends on a *Horn matrix*  $H$  and a coefficient for each column of  $H$ . In their recent study of moment maps of toric varieties [4], Clarke and Cox go one step further in strengthening the relationship between pairs  $(P, w)$  with rational linear precision and models  $X_{\mathcal{A},w}$  with rational MLE by using Horn matrices to characterise all pairs  $(P, w)$  that satisfy the more restrictive property of *strict linear precision*. They propose the use of Horn matrices to study polytopes with rational linear precision and state several questions and conjectures about the relationship between the Horn matrix of  $X_{\mathcal{A},w}$  and the primitive collections of the normal fan of  $P$ .

In this article we study the property of rational linear precision of pairs  $(P, w)$  from the point of view of Algebraic Statistics. Our main contribution is Theorem 4.1, which introduces a new family of polytopes (with associated weights) that has rational linear precision. We construct this family from a subclass of discrete statistical models introduced in Section 4 that we call *multinomial staged trees*. Looking at specific members of this family in 3D, we settle some of the questions raised in [4] related to Horn matrices and primitive collections.

This paper is structured as follows: In Sections 2.1-2.4 we provide background material on rational linear precision, discrete statistical models with rational MLE and Horn matrices. In Section 2.5 we state Questions 1 and 2 which guided our investigations related to Horn matrices and primitive collections. These questions are followed by a quick outline referring to the places in this article where they are addressed. In Section 3, we characterise the shape of the Horn matrix for pairs  $(P, w)$  in 2D. We also present a family of pairs  $(P, w)$  in 3D that has rational linear precision and explain several aspects of this family that relate to Questions 1 and 2. In Section 4.1 we define multinomial staged tree models, we prove they have rational MLE in Section 4.6 and we characterise the subclass of these models that are toric varieties in Section 4.5. These results lead to our main theorem, Theorem 4.1. Finally, in Section 5, we show that the examples from Section 3 are all multinomial staged trees

and prove our conjectures about the relationship between the combinatorics of the trees and primitive collections.

## 2 Preliminaries

We assume the reader is familiar with introductory material on computational algebraic geometry and toric geometry at the level of [6] and [7].

### 2.1 Notation and conventions

We consider pairs  $(P, w)$  where  $P$  is a  $d$ -dimensional lattice polytope in  $\mathbb{R}^d$ ,  $\mathbb{Z}^d$  is the fixed lattice,  $\mathcal{A} = P \cap \mathbb{Z}^d = \{m_1, \dots, m_s\}$  and  $w$  is a vector of positive weights indexed by  $\mathcal{A}$ . Fix  $n_1, \dots, n_r$  to be the inward facing primitive normal vectors of  $P$  corresponding to the facets  $F_1, \dots, F_r$  of  $P$  and let  $a_1, \dots, a_r$  be the corresponding integer translates in the facet presentation of  $P$  given by  $P = \{p \in \mathbb{R}^d : \langle p, n_i \rangle \geq -a_i, \forall i \in \{1, \dots, r\}\}$ . The lattice distance from  $p \in \mathbb{R}^d$  to  $F_i$  is

$$h_i(p) = \langle p, n_i \rangle + a_i, \quad i = 1, \dots, r.$$

The value  $h_i(m_j)$  is the lattice distance from the  $j$ -th lattice point to the  $i$ -th facet. The matrix with  $ij$  entry equal to  $h_i(m_j)$  is the *lattice distance matrix* of  $\mathcal{A}$ . We will often consider products of linear forms or variables whose exponents are given by vectors. For vectors  $v = (v_1, \dots, v_n), h = (h_1, \dots, h_n)$  we use  $v^h$  to denote the product  $\prod_{i=1}^n v_i^{h_i}$  and use the convention that  $0^0 = 1$ . If  $P$  is a polytope and  $a \geq 1$  is an integer,  $aP$  denotes its dilation.

### 2.2 Rational linear precision

In this section we follow closely the exposition in [4]. A more elementary introduction to this topic is available in [5, Chapter 3].

**Definition 2.1.** Let  $P \subseteq \mathbb{R}^d$  be a full dimensional polytope and let  $w = (w_1, \dots, w_s)$  be a vector of positive weights.

1. For  $1 \leq j \leq s$  and  $p \in P$ ,  $\beta_j(p) := \prod_{i=1}^r h_i(p)^{h_i(m_j)}$ .
2. The functions  $\beta_{w,j} := w_j \beta_j / \beta_w$  are the *toric blending functions* of  $(P, w)$ , where  $\beta_w(p) := \sum_{j=1}^s w_j \beta_j(p)$ .
3. Given *control points*  $\{Q_j\}_{1 \leq j \leq s} \in \mathbb{R}^n$ , the *toric patch*  $F : P \rightarrow \mathbb{R}^n$  is defined by

$$p \mapsto \frac{1}{\beta_w(p)} \sum_{j=1}^s w_j \beta_j(p) Q_j. \quad (1)$$

In part (3) of the previous definition, it is natural to choose the set control points to be  $\mathcal{A}$ .

**Definition 2.2.** Let  $(P, w)$  be as in Definition 2.1.

1. The *tautological patch*  $K_w : P \rightarrow P$  is the toric patch (1) where  $\{Q_j = m_j\}_{1 \leq j \leq s}$ .

2. The pair  $(P, w)$  has *strict linear precision* if  $K_w$  is the identity on  $P$ , that is

$$p = \frac{1}{\beta_w(p)} \sum_{j=1}^s w_j \beta_j(p) m_j, \text{ for all } p \in P.$$

3. The pair  $(P, w)$  has *rational linear precision* if  $K_w$  is a homeomorphism with a rational inverse.

*Remark 2.1.* We are interested in the property that the toric blending functions of a pair  $(P, w)$  have linear precision. By [10, Proposition 2.6], this happens if and only if the pair  $(P, w)$  has strict linear precision. In the case that  $(P, w)$  has rational linear precision, then the inverse of  $K_w$  can be used to reparametrise the toric blending functions  $\{\beta_{w,j} : P \rightarrow \mathbb{R}\}$  to obtain a set of blending functions  $\{\hat{\beta}_j = \beta_{w,j} \circ K_w^{-1} : P \rightarrow \mathbb{R}\}$  with linear precision.

*Remark 2.2.* An alternative way to specify a pair  $(P, w)$  is by using a homogeneous polynomial  $F_{\mathcal{A},w}$  whose dehomogenisation  $f_{\mathcal{A},w} = \sum_{j=1}^s w_j t^{m_j}$  encodes the weights in the coefficients and the lattice points in  $\mathcal{A}$  as exponents. We use this notation in Section 3 to describe toric patches in 2D and 3D.

*Remark 2.3.* If  $(P, w)$  has rational linear precision then  $(aP, \tilde{w})$ ,  $a \geq 1$ , also has this property where  $\tilde{w}$  is the vector of coefficients of  $(f_{\mathcal{A},w})^a$ . This follows by [11, Lemma 2.2].

**Example 2.1.** Consider the trapezoid  $P = \text{conv}((0,0), (3,0), (1,2), (0,2))$ , with ordered set of lattice points  $\mathcal{A} = \{(0,2), (1,2), (0,1), (1,1), (2,1), (0,0), (1,0), (2,0), (3,0)\}$  and vector of weights  $w = (1, 1, 2, 4, 2, 1, 3, 3, 1)$ . The polynomial  $f_{\mathcal{A},w}(s, t) = (1+s)(1+s+t)^2$  encodes  $(P, w)$ . The lattice distance functions for the facets of  $P$  are:

$$h_1(s, t) = s, \quad h_2(s, t) = t, \quad h_3(s, t) = 3 - t - s, \quad h_4(s, t) = 2 - t.$$

The toric blending functions for  $(P, w)$  are

$$\beta_{w,(i,j)}(s, t) = \frac{\binom{2}{j} \binom{3-j}{i} s^i t^j (3-s-t)^{3-i-j} (2-t)^{2-j}}{(1+s)(1+s+t)^2}, \text{ where } (i, j) \in \mathcal{A}.$$

The pair  $(P, w)$  does not have strict linear precision, but rational linear precision. The reparametrised toric blending functions with linear precision are:

$$\hat{\beta}_{w,(i,j)}(s, t) = \frac{\binom{2}{j} \binom{3-j}{i} s^i t^j (3-s-t)^{3-i-j} (2-t)^{2-j}}{4(3-t)^{3-j}}, \text{ where } (i, j) \in \mathcal{A}.$$

**Example 2.2.** Let  $\Delta_d = \{x \in \mathbb{R}^d : x_1 + \dots + x_d \leq 1, x_i \geq 0\}$  be the standard simplex in  $\mathbb{R}^d$  and  $k\Delta_d$  be its dilation by the integer  $k \geq 1$ . To a point  $m = (a_1, \dots, a_d) \in \mathcal{A} = k\Delta_d \cap \mathbb{Z}^d$  we associate the weight

$$w_m = \binom{k}{m} = \binom{k}{k - |m|, a_1, \dots, a_d}, \text{ where } |m| = a_1 + \dots + a_d.$$

The pair  $(k\Delta_d, w)$  has strict linear precision, see [10, Example 4.7]. By [4, Example 4.6], the product of two pairs,  $(P, w)$  and  $(Q, \tilde{w})$ , with strict linear precision also has strict linear precision. Hence, *the Bézier simploids*, which are polytopes of the form  $k_1\Delta_{d_1} \times \dots \times k_r\Delta_{d_r}$  for positive integers  $k_1, \dots, k_r, n_1, \dots, n_r$ , have strict linear precision. Conjecture 4.8 in [4] states that these are the only polytopes with strict linear precision.

## 2.3 Discrete statistical models with rational MLE

A probability distribution of a discrete random variable  $X$  with state space  $\{0, \dots, n\}$ , is a vector  $(p_0, \dots, p_n) \in \mathbb{R}^{n+1}$  where  $p_i = P(X = i)$ ,  $i \in \{0, \dots, n\}$ ,  $p_i \geq 0$  and  $\sum_{i=0}^n p_i = 1$ . The open probability simplex

$$\Delta_n^\circ = \{(p_0, \dots, p_n) \in \mathbb{R}^{n+1} \mid p_i > 0, p_0 + \dots + p_n = 1\}$$

consists of all strictly positive probability distributions for a discrete random variable with state space  $\{0, \dots, n\}$ . A discrete statistical model  $\mathcal{M}$  is a subset of  $\Delta_n^\circ$ .

Given a set of independent and identically distributed observations  $\mathcal{D} = \{X_1, \dots, X_N\}$  of  $X$ , we let  $u = (u_0, \dots, u_n)$  be the vector where  $u_i$  is the number of times the outcome  $i$  appears in  $\mathcal{D}$ . The likelihood function  $L(p, u) : \mathcal{M} \rightarrow \mathbb{R}_{\geq 0}$  defined by  $(p_0, \dots, p_n) \mapsto \prod p_i^{u_i}$  records the probability of observing the set  $\mathcal{D}$ . The *maximum likelihood estimator* (MLE) of the model  $\mathcal{M}$  is the function  $\Phi : \mathbb{R}^{n+1} \rightarrow \mathcal{M}$  that sends each vector  $(u_0, \dots, u_n)$  to the maximiser of  $L(p, u)$ , i.e.

$$\Phi(u) := \arg \max L(p, u).$$

For an arbitrary model  $\mathcal{M}$ , the problem of estimating  $\arg \max L(p, u)$  is a difficult one. However, for special families, such as discrete exponential families, there are theorems that guarantee the existence and uniqueness of  $\arg \max L(p, u)$  when  $u$  has nonzero entries. We are interested in the case where  $\Phi$  is a rational function of  $u$ .

**Definition 2.3.** Let  $\mathcal{M}$  be a discrete statistical model with MLE  $\Phi : \mathbb{R}^{n+1} \rightarrow \mathcal{M}$ ,  $u \mapsto \hat{p}$ . The model  $\mathcal{M}$  has *rational MLE* if the coordinate functions of  $\Phi$  are rational functions in  $u$ .

**Example 2.3.** Consider the model  $\mathcal{M}$  of two independent binary random variables  $X, Y$ , with outcome set  $\{0, 1\}$  and  $p_{ij} = P(X = i, Y = j)$ . This model is the set of all points  $(p_{00}, p_{01}, p_{10}, p_{11})$  in  $\Delta_3^\circ$  that satisfy the equation  $p_{00}p_{11} - p_{10}p_{01} = 0$ . The model has rational MLE  $\Phi : \mathbb{R}^4 \rightarrow \mathcal{M}$  where

$$(u_{00}, u_{01}, u_{10}, u_{11}) \mapsto \left( \frac{u_{0+}u_{+0}}{u_{++}^2}, \frac{u_{0+}u_{+1}}{u_{++}^2}, \frac{u_{1+}u_{+0}}{u_{++}^2}, \frac{u_{1+}u_{+1}}{u_{++}^2} \right)$$

and  $u_{i+} = u_{i0} + u_{i1}$ ,  $u_{+j} = u_{0j} + u_{1j}$ ,  $u_{++} = \sum_{i,j \in \{0,1\}} u_{ij}$ .

**Definition 2.4.** A *Horn matrix* is an integer matrix whose column sums are equal to zero. Given an Horn matrix  $H$ , with columns  $h_1, \dots, h_r$ , and a vector  $\lambda \in \mathbb{R}^r$ , the Horn parametrisation  $\varphi_{(H,\lambda)} : \mathbb{R}^r \rightarrow \mathbb{R}^r$  is the rational map given by

$$u \mapsto (\lambda_1(Hu)^{h_1}, \lambda_2(Hu)^{h_2}, \dots, \lambda_r(Hu)^{h_r}).$$

**Example 2.4.** The MLE  $\Phi$  in Example 2.3 is given by a Horn parametrisation  $\varphi_{(H,\lambda)}$ , where

$$u = \begin{pmatrix} u_{00} \\ u_{01} \\ u_{10} \\ u_{11} \end{pmatrix}, \quad H = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ -2 & -2 & -2 & -2 \end{pmatrix}, \quad Hu = \begin{pmatrix} u_{0+} \\ u_{1+} \\ u_{+0} \\ u_{+1} \\ -2u_{++} \end{pmatrix}, \quad \text{and } \lambda = (4, 4, 4, 4).$$

**Definition 2.5.** We say that  $(H, \lambda)$  is a Horn pair if: (1) the sum of the coordinates of  $\varphi_{(H, \lambda)}$  as rational functions in  $u$  is equal to 1 and (2) the map  $\varphi_{(H, \lambda)}$  is defined for all positive vectors and it sends these to positive vectors in  $\mathbb{R}^r$ .

**Theorem 2.1** ([9, Theorem 1]). *A discrete statistical model  $\mathcal{M}$  has rational MLE  $\Phi$  if and only if there exists a Horn pair  $(H, \lambda)$  such that  $\mathcal{M}$  is the image of the Horn parametrisation  $\varphi_{(H, \lambda)}$  restricted to the open orthant  $\mathbb{R}_{>0}^r$  and  $\Phi = \varphi_{(H, \lambda)}$  on  $\mathbb{R}_{>0}^r$ .*

It is possible that two Horn parametrisations  $\varphi_{(H, \lambda)}$  and  $\varphi_{(\tilde{H}, \tilde{\lambda})}$  are equal even if  $H \neq \tilde{H}$  and  $\lambda \neq \tilde{\lambda}$ . A Horn matrix  $H$  is *minimal* if it has no zero rows and no two rows are linearly dependent. By [4, Proposition 6.11] there exists a unique, up to permutation of the rows, minimal Horn matrix that defines  $\varphi_{(H, \lambda)}$ . Any other pair  $(H, \lambda)$  that defines the same Horn parametrisation may be transformed into one where  $H$  is a minimal Horn matrix; this is done by adding collinear rows, deleting zero rows and adjusting the vector  $\lambda$  accordingly, see [9, Lemma 3]. We end this section by noting that [9, Proposition 23] states that if  $(H, \lambda)$  is a minimal Horn pair, then every row of  $H$  has either all entries greater than or equal zero or all entries less than or equal to zero. We call the submatrix of  $H$  that consists of all rows with nonnegative entries, the positive part of  $H$ , and its complement the negative part of  $H$ .

## 2.4 The two properties combined

Recall that if a pair  $(P, w)$  has rational linear precision then the reparametrisations of its toric blending functions by the inverse of the tautological patch  $K_w$  have linear precision. To understand when such a reparameterisation is possible it is useful to consider all possible reparameterisations of blending functions at once. Garcia-Puente and Sottile [10] achieve this by associating a toric variety  $X_{\mathcal{A}, w}$  to the pair  $(P, w)$ .

Given a pair  $(P, w)$ , the *scaled projective toric variety*  $X_{\mathcal{A}, w}$  is the image of the map  $[w\chi]_{\mathcal{A}} : (\mathbb{C}^*)^d \rightarrow \mathbb{P}^{s-1}$  defined by  $t \mapsto [w_1 t^{m_1} : w_2 t^{m_2} : \dots : w_s t^{m_s}]$ . To consider the maximum likelihood estimation problem in the realm of complex algebraic geometry we consider the variety  $W = V(x_1 \dots x_s (x_1 + \dots + x_s)) \subset \mathbb{P}^{s-1}$  and the map

$$X_{\mathcal{A}, w} \setminus W \rightarrow (\mathbb{C}^*)^s, [x_1 : \dots, x_s] \mapsto \frac{1}{x_1 + \dots + x_s} (x_1, \dots, x_s).$$

The image of this map is closed and denoted by  $Y_{\mathcal{A}, w}$ . We call  $Y_{\mathcal{A}, w}$  a *scaled very affine toric variety*. The set  $\mathcal{M}_{\mathcal{A}, w} = Y_{\mathcal{A}, w} \cap \mathbb{R}_{>0}^s$  is a subset of the open simplex  $\Delta_{s-1}^\circ$  and as such it is a statistical model. This class of models, of the form  $\mathcal{M}_{\mathcal{A}, w}$ , are known as *log-linear models*.

*Remark 2.4.* The variety  $Y_{\mathcal{A}, w}$  admits two parameterisations, one by monomials and another by toric blending functions [4, Proposition 5.2]. These are

$$\begin{aligned} \overline{w_{\chi_{\mathcal{A}}}} : \mathbb{C}^d &\rightarrow Y_{\mathcal{A}, w}, & t &\mapsto \left( \frac{w_1 t^{m_1}}{\sum_{j=1}^s w_j t^{m_j}}, \dots, \frac{w_s t^{m_s}}{\sum_{j=1}^s w_j t^{m_j}} \right), \\ \overline{w_{\beta_{\mathcal{A}}}} : \mathbb{C}^d &\rightarrow Y_{\mathcal{A}, w}, & t &\mapsto \left( \frac{w_1 \beta_1(t)}{\beta_w(t)}, \dots, \frac{w_s \beta_s(t)}{\beta_w(t)} \right). \end{aligned} \tag{2}$$

We now consider the maximum likelihood estimation problem for log-linear models. Given a vector of counts  $u$ , we let  $\bar{u} := u/|u| \in \Delta_{s-1}^\circ$  be the empirical distribution, where  $|u| = \sum u_j$ . We define the tautological map  $\tau_{\mathcal{A}}$  following the convention in [4],

$$\tau_{\mathcal{A}} : \Delta_{s-1}^\circ \rightarrow P^\circ, \quad (\bar{u}_1, \dots, \bar{u}_s) \mapsto \sum_{j=1}^s \bar{u}_j m_j. \quad (3)$$

The maximum likelihood estimate of  $\bar{u}$  for the model  $\mathcal{M}_{\mathcal{A},w}$  exists and it is unique whenever all entries of  $\bar{u}$  are positive.

**Theorem 2.2.** [16, Corollary 7.3.9] *The maximum likelihood estimate in  $\mathcal{M}_{\mathcal{A},w}$  for the empirical distribution  $\bar{u} \in \Delta_{s-1}^\circ$  is the unique point  $\hat{p} \in \mathcal{M}_{\mathcal{A},w}$  that satisfies  $\tau_{\mathcal{A}}(\hat{p}) = \tau_{\mathcal{A}}(\bar{u})$ .*

In the Algebraic Statistics literature, models with rational MLE are also known as models with *maximum likelihood degree* equal to 1. Even though the previous theorem guarantees the existence of the MLE, it is not true that every log-linear model has rational MLE. We refer the reader to [1] for several examples of log-linear models that do not have rational MLE, or equivalently for examples of models with maximum likelihood degree greater than 1. We end this section by recalling two theorems that connect models with rational MLE and pairs with rational linear precision.

**Theorem 2.3.** [10, Proposition 5.1] *The pair  $(P, w)$  has rational linear precision if and only if the model  $\mathcal{M}_{\mathcal{A},w}$  has rational MLE.*

**Theorem 2.4** ([4]). *Set  $a_P := \sum_{i=1}^r a_i$  and  $n_P := \sum_{i=1}^r n_i$ . The following are equivalent:*

1. *The pair  $(P, w)$  has strict linear precision.*
2.  *$n_P = 0$  and  $\beta_w(p) = \sum_{j=1}^s w_j \beta_j(p) = \sum_{j=1}^s w_j \prod_{i=1}^r h_i(p)^{h_i(m_j)}$  is a nonzero constant  $c$ .*
3.  *$\mathcal{M}_{\mathcal{A},w}$  has rational MLE with minimal Horn pair  $(H, \lambda)$  given by*

$$H = \begin{pmatrix} h_1(m_1) & h_1(m_2) & \dots & h_1(m_s) \\ h_2(m_1) & h_2(m_2) & \dots & h_2(m_s) \\ \vdots & \vdots & & \vdots \\ h_r(m_1) & h_r(m_2) & \dots & h_r(m_s) \\ -a_P & -a_P & \dots & -a_P \end{pmatrix}, \quad \lambda_j = \frac{w_j}{c} (-a_P)^{a_P}$$

## 2.5 Primitive collections and Horn pairs

The notion of primitive collections was first introduced by Batyrev in [3] for a smooth and projective toric variety  $X_{\Sigma_P}$  of the polytope  $P$ . It provides an elegant description of the nef cone for  $X_{\Sigma_P}$ . This result has been generalized to the simplicial case and the definition of primitive collections for the non-simplicial case has been introduced in [8].

**Definition 2.6.** Let  $\Sigma_P$  be a normal fan. A subset  $C \subseteq \Sigma_P(1)$  of 1-faces of the normal fan is called a *primitive collection* if

1.  $C \not\subseteq \sigma(1)$  for all  $\sigma \in \Sigma_P$ .

2. For every proper subset  $C' \subsetneq C$ , there exists  $\sigma \in \Sigma_P$  such that  $C' \subseteq \sigma(1)$ .

In particular, if  $\Sigma_P$  is simplicial,  $C$  is a primitive collection if  $C$  does not generate a cone of  $\Sigma_P$  but every proper subset does.

For strict linear precision, Theorem 2.4 gives the minimal Horn pair based only on the lattice distance functions of the facets of the polytope. The authors in [4] raise the question whether it is possible to obtain a similar description of minimal Horn pairs of polytopes with rational linear precision.

**Question 1.** Is the positive part of the minimal Horn matrix of a pair  $(P, w)$  with rational linear precision always equal to the lattice distance matrix of  $\mathcal{A}$ ?

For pairs  $(P, w)$  in 2D with rational linear precision, and the family of prisms in Section 3.2, the answer to Question 1 is affirmative, see Theorem 3.2, Proposition 3.3, and Appendix A. In [4] there are two examples, one of a trapezoid [4, Section 8.1] and one of a decomposable graphical model [4, Section 8.3], where the positive part of the Horn matrix is the lattice distance matrix of  $\mathcal{A}$  and the negative rows are obtained via the primitive collections of the normal fan of  $P$ . These examples motivate the next definition and Question 2:

**Definition 2.7.** To a pair  $(P, w)$  we associate the matrix  $M_{\mathcal{A}, \Sigma_P}$  which consists of the lattice distance matrix of  $\mathcal{A}$ , with  $ij$ -th entry  $h_i(m_j)$ , together with negative rows given by summing the rows of the lattice distance functions  $-h_i$ , for which the facet normals  $n_i$  belong to the same primitive collection of  $\Sigma_P$ .

**Question 2.** For a pair  $(P, w)$  with rational linear precision is there a Horn pair  $(H, \lambda)$  for which  $H = M_{\mathcal{A}, \Sigma_P}$ ?

For pairs  $(P, w)$  in 2D with rational linear precision, the answer to Question 2 is affirmative, see Theorem 3.2. For the family of prisms in Section 3.2, Question 2 is affirmative only for certain subclasses, see Theorem 3.5. For an arbitrary pair  $(P, w)$  with rational linear precision, the matrix  $M_{\mathcal{A}, \Sigma_P}$  is not necessarily a Horn matrix, see Section 3.3.1. Even in the case that  $M_{\mathcal{A}, \Sigma_P}$  is a Horn matrix, it does not necessarily give rise to a Horn pair for  $(P, w)$ , see Section 3.3.2. In Section 3 we see a number of special cases for which the answer to Question 2 is affirmative. In Section 5, we give a condition on  $(P, w)$  which guarantees the existence of a Horn pair  $(H, \lambda)$  with  $H = M_{\mathcal{A}, \Sigma_P}$ . We also provide an explanation for the negative rows of the Horn matrix in the language of multinomial staged tree models - introduced in Section 4.

### 3 Examples of Horn pairs in 2D and 3D

In this section we present families of 2D and 3D pairs  $(P, w)$  with rational linear precision and explore the connection between the geometry of the polytope and the shape of its corresponding Horn pair.



### 3.1 Toric surface patches and Horn pairs in 2D

By [11], the only 2D toric patches with rational linear precision are the Bézier triangles, tensor product patches and trapezoidal patches, seen in Figure 1. This family of polygons, that we denote by  $\mathcal{F}$ , consists of all the Newton polytopes of the polynomials

$$f_{a,b,d}(s,t) := (1+s)^a((1+s)^d+t)^b \text{ for } a,b,d \geq 0.$$

For general  $a,b,d$ ,  $T_{a,b,d}$  will be a trapezoidal patch, in the special cases  $T_{0,b,1} = b\Delta_2$  and  $T_{a,b,0} = a\Delta_1 \times b\Delta_1$ , we will have the more familiar Bézier triangles and tensor product patches. The lattice points in  $T_{a,b,d} \cap \mathbb{Z}^2$  are  $\mathcal{A} = \{(i,j) : 0 \leq j \leq b, 0 \leq i \leq a+d(b-j)\}$ . By Theorems 2.1 and 2.3 we know that the statistical model associated to a pair in  $\mathcal{F}$  admits a Horn pair.




Figure 1: Left: Bézier triangles. Middle: Tensor product patches. Right: Trapezoids. The normal fan of each polygon is displayed in the bottom row; two rays with the same color are in the same primitive collection.

**Proposition 3.1.** *A Horn pair  $(H, \lambda)$  of a polygon in the family  $\mathcal{F}$  is given by:*

$$H = \begin{pmatrix} h_1(m_1) & \dots & h_1(i,j) & \dots & h_1(m_s) \\ h_2(m_1) & \dots & h_2(i,j) & \dots & h_2(m_s) \\ h_3(m_1) & \dots & h_3(i,j) & \dots & h_3(m_s) \\ h_4(m_1) & \dots & h_4(i,j) & \dots & h_4(m_s) \\ -(h_1+h_3)(m_1) & \dots & -(h_1+h_3)(i,j) & \dots & -(h_1+h_3)(m_s) \\ -(h_2+h_4)(m_1) & \dots & -(h_2+h_4)(i,j) & \dots & -(h_2+h_4)(m_s) \end{pmatrix},$$

$$\lambda_{ij} = (-1)^{a+d(b-j)+b} \binom{b}{j} \binom{a+d(b-j)}{i},$$

where  $(i,j) \in \mathcal{A}$  and  $h_1(s,t) = s$ ,  $h_2(s,t) = t$ ,  $h_3(s,t) = a+db-s-dt$ ,  $h_4(s,t) = b-t$ .

*Proof.* We use [4, Proposition 8.4]. The terms of the polynomial  $f_{a,b,d}(s,t)$  specify weights and lattice points in  $T_{a,b,d} \cap \mathbb{Z}^2$ . The monomial parametrisation (2) of  $Y_{\mathcal{A},w}$  is

$$\overline{w\chi_{\mathcal{A}}}(s,t) = \frac{1}{f_{a,b,d}(s,t)} (S_{a+db,0}, \dots, S_{1,0}, S_{0,1}, S_{0,0}),$$

where  $S_{i,j} = w_{ij}s^i t^j$  and  $w_{ij} = \binom{b}{j} \binom{a+db-dj}{i}$ . Composing the monomial parametrisation with the tautological map (3) gives the following birational map:

$$(\tau_{\mathcal{A}} \circ \overline{w\chi_{\mathcal{A}}})(s, t) = \left( \frac{((a+db)(1+s)^d + at)s}{((1+s)^d + t)(1+s)}, \frac{bt}{((1+s)^d + t)} \right)$$

with the following inverse:

$$\varphi(s, t) = \left( \frac{s}{a+db-s-dt}, \frac{(a+db-dt)^d t}{(a+db-s-dt)^d (b-t)} \right)$$

The  $i, j$ -th component of the monomial parametrisation composed with  $\varphi(s, t)$  is given by

$$\begin{aligned} ((\overline{w\chi_{\mathcal{A}}} \circ \varphi)(s, t))_{i,j} &= \frac{S_{ij}(\varphi(s, t))}{f_{a,b,d}(\varphi(s, t))} \\ &= \binom{b}{j} \binom{a+db-dj}{i} \frac{s^i t^j (a+db-s-dt)^{a+db-i-dj} (b-t)^{b-j}}{(a+db-dt)^{a+db-dj} b^b} \\ &= w_{i,j} \frac{h_1(s, t)^{h_1(i,j)} h_2(s, t)^{h_2(i,j)} h_3(s, t)^{h_3(i,j)} h_4(s, t)^{h_4(i,j)}}{h_5(s, t)^{h_5(i,j)} b^b}, \end{aligned} \tag{4}$$

where  $h_1, h_2, h_3, h_4$  are as in the statement of the proposition and  $h_5 = h_1 + h_3$ . It follows from [4, Proposition 8.4] that the Horn parametrisation is  $(\overline{w\chi_{\mathcal{A}}} \circ \varphi)(p)$  where

$$p = \sum_{(i,j) \in \mathcal{A}} \frac{u_{i,j}}{u_+} (i, j), \quad u_+ = \sum_{(i,j) \in \mathcal{A}} u_{i,j}.$$

Therefore, the columns of the Horn matrix are the exponents of

$$\frac{h_1(p)^{h_1(i,j)} h_2(p)^{h_2(i,j)} h_3(p)^{h_3(i,j)} h_4(p)^{h_4(i,j)}}{h_5(p)^{h_5(i,j)} b^b},$$

taking into account the negative sign of the exponents in the denominator. From the sum-to-one condition  $\sum_{i,j} ((\overline{w\chi_{\mathcal{A}}} \circ \varphi)(p))_{i,j} = 1$  it follows that  $\lambda_{i,j} = (-1)^{a+d(b-j)+b} w_{i,j}$ .  $\square$

*Remark 3.1.* The reparametrised blending functions for each pair  $(P, w)$  in  $\mathcal{F}$  are given in equation (4) in the previous proof. For the case  $a = d = 1$  and  $b = 2$ , these are in Example 2.1.

*Remark 3.2.* For general  $a, b, d$ , Proposition 3.1 gives the minimal Horn pair for  $T_{a,b,d}$ ; this is not the case for  $T_{0,b,1}$  and  $T_{a,b,0}$ . For the last two cases, the minimal Horn pair is obtained after row reduction operations or alternatively from Theorem 2.4.

Using Proposition 3.1 and Theorem 2.4 we obtain an affirmative answer to Question 1 for pairs  $(P, w)$  in 2D. A closer look at the primitive collections in Figure 1 also reveals an affirmative answer to Question 2. This is contained in the next theorem.




Figure 2: The general representative of a prismatoid in  $\mathcal{P}$  is the convex hull of two trapezoids,  $\text{conv}(T_{a,b,d} \times \{0\}, T_{a',b',d} \times \{1\})$ , dilated by  $l$ .

**Theorem 3.2.** *Every pair  $(P, w)$  in 2D with rational linear precision has a Horn pair  $(H, \lambda)$  with  $H = M_{\mathcal{A}, \Sigma_P}$ .*

*Proof.* The normal fans of the polygons in  $\mathcal{F}$  are depicted in Figure 1, in each subcase the shape of the normal fan and its primitive collections are independent of the values of  $a, b, d$ . The minimal Horn pair  $(H, \lambda)$  for the 2D simplex,  $T_{0,b,1} = b\Delta_2$ , given in Theorem 2.4 satisfies  $H = M_{\mathcal{A}, \Sigma_P}$ . This follows because  $b\Delta_2$  has one primitive collection,  $\{n_1, n_2, n_3\}$  and hence  $M_{\mathcal{A}, \Sigma_P}$  has a single negative row. For the tensor product patch  $T_{a,b,0} = a\Delta_1 \times b\Delta_1$  and the general trapezoid  $T_{a,b,d}$ , the primitive collections are  $\{n_1, n_3\}$  and  $\{n_2, n_4\}$ . In these cases, the Horn pair  $(H, \lambda)$  in Proposition 3.1 satisfies  $H = M_{\mathcal{A}, \Sigma_P}$ .  $\square$

### 3.2 A family of prismatoids with rational linear precision

Unlike the 2D case, there is no classification for 3D lattice polytopes with rational linear precision. In this section we consider the family of prismatoids

$$\begin{aligned} \mathcal{P} := \{ & (P, w) : P \text{ is the Newton polytope of } f_{\mathcal{A}, w}(s, t, v) = (f_{a,b,d}(s, t) + v f_{a',b',d}(s, t))^l, \\ & w \text{ is the vector of coefficients of } f_{\mathcal{A}, w}(s, t, v), \\ & a, a', b, b', d, l \in \mathbb{Z}_{\geq 0} \text{ with } a' \leq a, b' \leq b \}. \end{aligned}$$

A general element of  $\mathcal{P}$  is depicted in Figure 2, prismatoids for different specialisations of  $a, a', b, b', d$  are displayed in Table 1. Note that some 3D Bézier simploids are also obtained by specialisation. Even though Remark 2.3 says it suffices to show that  $P$  has rational linear precision for  $l = 1$ , we do not use this extra assumption.

| <b>(A) Primatoids with trapezoidal base <math>a &gt; 0, b &gt; 0, d &gt; 0, l &gt; 0</math></b>                 |  |  |  |
|---|--|--|--|
| Trapezoidal frusta<br>$a' > 0, b' > 0$  |  | Triangle top<br>(simplex if $d = 1$ )<br>$a' = 0, b' > 0$                                    |  |
| Trapezoidal wedges<br>$a' > 0, b' = 0$  |  | Trapezoidal Pyramids<br>$a' = 0, b' = 0$   |  |
| <b>(B) Primatoids with tensor product base <math>a &gt; 0, b &gt; 0, d = 0, l &gt; 0</math></b>                 |  |  |  |
| Tensor product frusta<br>$a' > 0, b' > 0$   |  | Tensor product wedges<br>$a' = 0, b' > 0$  |  |
| Tensor product pyramids<br>$a' = 0, b' = 0$   |  | Tensor product wedges<br>$a' > 0, b' = 0$  |  |
| <b>(C) Primatoids with triangular base <math>a = 0, b &gt; 0, d &gt; 0, l &gt; 0</math></b>                     |  |  |  |
| Triangular frusta<br>(simplices if $d = 1$ )<br>$a' = 0, b' > 0$  |  | Pyramid (simplex-based if $d = 1$ )<br>$a' = 0, b' = 0$                                      |  |
| <b>(D) 3D Bézier simploids <math>l &gt; 0</math></b>  |  |  |  |
| 3D Tensor Product<br>$a' = a > 0$<br>$b' = b > 0$<br>$d = 0$<br>$la\Delta_1 \times lb\Delta_1 \times l\Delta_1$ |  | Triangular Prism<br>$a' = a = 0$<br>$b' = b > 0$<br>$d = 1$<br>$lb\Delta_2 \times l\Delta_1$ |  |
| 3D simplex<br>$a' = a = 0$<br>$b' = 0, b = 1$<br>$d = 1$<br>$l\Delta_3$   |  |  |  |

Table 1: Representatives of members of  $\mathcal{P}$  for different choices of parameters  $a, a', b, b', d$

**Proposition 3.3.** *The pairs in  $\mathcal{P}$  have rational linear precision with a Horn pair  $(H, \lambda)$ :*

$$H = \begin{pmatrix} h_1(m_1) & \dots & h_1(i, j, k) & \dots & h_1(m_s) \\ h_2(m_1) & \dots & h_2(i, j, k) & \dots & h_2(m_s) \\ h_3(m_1) & \dots & h_3(i, j, k) & \dots & h_3(m_s) \\ h_4(m_1) & \dots & h_4(i, j, k) & \dots & h_4(m_s) \\ h_5(m_1) & \dots & h_5(i, j, k) & \dots & h_5(m_s) \\ h_6(m_1) & \dots & h_6(i, j, k) & \dots & h_6(m_s) \\ -(h_1 + h_4)(m_1) & \dots & -(h_1 + h_4)(i, j, k) & \dots & -(h_1 + h_4)(m_s) \\ -(h_2 + h_5)(m_1) & \dots & -(h_2 + h_5)(i, j, k) & \dots & -(h_2 + h_5)(m_s) \\ -(h_3 + h_6)(m_1) & \dots & -(h_3 + h_6)(i, j, k) & \dots & -(h_3 + h_6)(m_s) \end{pmatrix},$$

$$\lambda_{i,j,k} = (-1)^{(\sum_{i=1}^6 h_i)(i,j,k)} \binom{l}{k} \binom{bl - (b-b')k}{j} \binom{(a+db)l - ((a+db) - (a'+db'))k - dj}{i},$$

where  $m_1, \dots, m_n$  is an ordered list of lattice points  $(i, j, k)$  in  $\mathcal{A}$  and  $h_1, \dots, h_6$  are

$$\begin{aligned} h_1(s, t, v) &= s, & h_4(s, t, v) &= (a+db)l - s - dt - ((a+db) - (a'+db'))v, \\ h_2(s, t, v) &= t, & h_5(s, t, v) &= bl - t - (b-b')v, \\ h_3(s, t, v) &= v, & h_6(s, t, v) &= l - v. \end{aligned}$$

*Proof.* The polynomial  $f_{\mathcal{A},w}(s, t, v)$  in the definition of  $\mathcal{P}$ , can be expressed as a sum

$$f_{\mathcal{A},w}(s, t, v) = \sum_{k=0}^l \sum_{j=0}^{bl - (b-b')k} \sum_{i=0}^{(a+db)l - ((a+db) - (a'+db'))k - dj} S_{i,j,k}$$

where

$$S_{i,j,k}(s, t, v) = \binom{l}{k} \binom{bl - (b-b')k}{j} \binom{(a+db)l - ((a+db) - (a'+db'))k - dj}{i} s^i t^j v^k.$$

We let

$$(\overline{w\chi_{\mathcal{A}}}(s, t, v))_{i,j,k} = \frac{S_{i,j,k}(s, t, v)}{S(s, t, v)},$$

then, the vector of all  $(\overline{w\chi_{\mathcal{A}}}(s, t, v))_{i,j,k}$  gives the monomial parametrisation (2) of  $Y_{\mathcal{A},w}$  with weights

$$w_{i,j,k} = \binom{l}{k} \binom{bl - (b-b')k}{j} \binom{(a+db)l - ((a+db) - (a'+db'))k - dj}{i}.$$

Composing the monomial parametrisation with the tautological map (3) gives the following

birational map:

$$\begin{aligned}
(\tau_{\mathcal{A}} \circ \overline{w\chi_{\mathcal{A}}})(s, t, v) &= \frac{1}{S(s, t, v)} \left( \sum_{k=0}^l \sum_{j=0}^{bl-(b-b')k} \sum_{i=0}^{(a+db)l-((a+db)-(a'+db'))k-dj} S_{i,j,k} \right) (i, j, k) \\
&= \left( \frac{ls \left( ((a+db)(1+s)^d + at)(1+s)^a ((1+s)^d + t)^{b'} + v(a'+db')(1+s)^d + a't \right) (1+s)^{a'} ((1+s)^d + t)^{b'}}{(1+s)((1+s)^d + t)((1+s)^a ((1+s)^d + t)^{b+v} (1+s)^{a'} ((1+s)^d + t)^{b'}} \right. \\
&\quad \frac{lt(b'v(1+s)^{a'} ((1+s)^d + t)^{b'} + b(1+s)^a ((1+s)^d + t)^b)}{((1+s)^d + t)((1+s)^a ((1+s)^d + t)^{b+v} (1+s)^{a'} ((1+s)^d + t)^{b'}} \\
&\quad \left. \frac{lv(1+s)^{a'} ((1+s)^d + t)^{b'}}{(1+s)^a ((1+s)^d + t)^{b+v} (1+s)^{a'} ((1+s)^d + t)^{b'}} \right)
\end{aligned}$$

with the following inverse:

$$\begin{aligned}
\varphi(s, t, v) &= \left( \frac{s}{(a+db)l - s - dt - ((a+db) - (a'+db'))v}, \right. \\
&\quad \frac{t((a+db)l - dt - ((a+db) - (a'+db'))v)^d}{((a+db)l - s - dt - ((a+db) - (a'+db'))v)^d (bl - t - (b-b')v)}, \\
&\quad \left. \frac{v((a+db)l - dt - ((a+db) - (a'+db'))v)^{(a+db)-(a'+db')}(bl - (b-b')v)^{b-b'}}{((a+db)l - s - dt - ((a+db) - (a'+db'))v)^{(a+db)-(a'+db')}(bl - t - (b-b')v)^{b-b'}(l-v)} \right)
\end{aligned}$$

Composing  $\varphi(s, t, v)$  with the monomial parametrisation gives

$$\frac{S_{i,j,k}(\varphi(s, t, v))}{S(\varphi(s, t, v))} = w_{i,j,k} \frac{s^i t^j v^k ((a+db)l - s - dt - ((a+db) - (a'+db'))v)^{(a+db)l - i - dj - ((a+db) - (a'+db'))k} (bl - (b-b')v - t)^{bl - (b-b')k - j} (l-v)^{l-k}}{l^l ((a+db)l - dt - ((a+db) - (a'+db'))v)^{(a+db)l - dj - ((a+db) - (a'+db'))k} (bl - (b-b')v)^{bl - (b-b')k}} \quad (5)$$

$$= w_{i,j,k} \frac{h_1(s, t, v)^{h_1(i,j,k)} h_2(s, t, v)^{h_2(i,j,k)} h_3(s, t, v)^{h_3(i,j,k)} h_4(s, t, v)^{h_4(i,j,k)} h_5(s, t, v)^{h_5(i,j,k)} h_6(s, t, v)^{h_6(i,j,k)}}{h_7(s, t, v)^{h_7(i,j,k)} h_8(s, t, v)^{h_8(i,j,k)} h_9(s, t, v)^{h_9(i,j,k)}} \quad (6)$$

where  $h_1, \dots, h_6$  are as in the statement of the theorem and

$$h_7 = h_1 + h_4 \quad h_8 = h_2 + h_5 \quad h_9 = h_3 + h_6.$$

According to [4, Proposition 8.4], the polytope has rational linear precision with weights  $w_{i,j,k}$  as defined above and the Horn parametrisation of  $Y_{\mathcal{A},w}$  is given by:

$$\frac{S_{i,j,k}(\varphi(p))}{S(\varphi(p))} = w_{i,j,k} \frac{h_1(p)^{h_1(i,j,k)} h_2(p)^{h_2(i,j,k)} h_3(p)^{h_3(i,j,k)} h_4(p)^{h_4(i,j,k)} h_5(p)^{h_5(i,j,k)} h_6(p)^{h_6(i,j,k)}}{h_7(p)^{h_7(i,j,k)} h_8(p)^{h_8(i,j,k)} h_9(p)^{h_9(i,j,k)}}$$

where

$$p = \sum_{(i,j,k) \in \mathcal{A}} \frac{u_{i,j,k}}{u_+} (i, j, k), \quad u_+ = \sum_{(i,j,k) \in \mathcal{A}} u_{i,j,k}.$$

Since the Horn parametrisation is, by definition, a product of linear forms whose exponents match their coefficients, we know that the columns of  $H$  are the exponents  $(h_1, h_2, h_3, h_4, h_5, h_6, -h_7, -h_8, -h_9)$  all evaluated at a lattice point  $(i, j, k)$ . It follows from  $\sum_{i,j,k} ((\overline{w\chi_{\mathcal{A}}} \circ \varphi)(p))_{i,j,k} = 1$  that  $\lambda_{i,j,k} = (-1)^{(\sum_{i=1}^6 h_i)(i,j,k)} w_{i,j,k}$ .  $\square$

### 3.3 Minimal Horn pairs for prisms in $\mathcal{P}$

We now study Questions 1 and 2 for elements in  $\mathcal{P}$ . Proposition 3.3 gives a Horn pair  $(H, \lambda)$  for each  $(P, w) \in \mathcal{P}$  in Table 1, however  $H$  need not be the minimal Horn matrix in each case. By [9, Lemma 9], we can find *the minimal Horn matrix associated to  $(P, w)$*  using row reduction operations on  $H$ .

**Notation 3.4.** *We will denote the facets of a general element in  $\mathcal{P}$  as follows:*

$$\begin{array}{lll} F_1 = \text{left facet}, & F_2 = \text{front facet}, & F_3 = \text{bottom facet}, \\ F_4 = \text{right facet}, & F_5 = \text{back facet}, & F_6 = \text{upper facet}. \end{array}$$

*This labelling is used in Figure 2. The normal vectors corresponding to each facet are:*

$$\begin{array}{lll} n_1 = (1, 0, 0), & n_2 = (0, 1, 0), & n_3 = (0, 0, 1), \\ n_4 = (-1, -d, -((a + db) - (a' + db'))), & n_5 = (0, -1, -(b - b')), & n_6 = (0, 0, -1). \end{array}$$

#### 3.3.1 The non-simple prisms

The trapezoidal pyramids, tensor product pyramids and prisms with triangle on top, depicted in Table 1 (A) and (B), are all examples of non-simple polytopes in  $\mathcal{P}$ . Their primitive collections are:

$$\begin{array}{ll} \text{Prisms with triangle on top} & \{n_1, n_3, n_4\}, \{n_1, n_2, n_4\}, \{n_2, n_5\}, \{n_3, n_6\} \\ \text{Trapezoidal pyramids} & \{n_1, n_3, n_4\}, \{n_2, n_3, n_5\} \\ \text{Tensor product pyramids} & \{n_1, n_3, n_4\}, \{n_2, n_3, n_5\}. \end{array}$$

There is no  $n_6$  for the two pyramids since the facet  $F_6$  has collapsed to a point.

For a pair  $(P, w)$  in the subfamily of non-simple prisms in  $\mathcal{P}$ , the matrix  $M_{\mathcal{A}, \Sigma_P}$  cannot be a Horn matrix since the primitive collections are not a partition of the 1-dimensional rays of the normal fan and therefore the columns cannot add to zero.

**Example 3.1.** It follows from Proposition 3.3 that the minimal Horn matrix associated to the tensor product pyramid in Table 1 (B) is:

$$H = \begin{pmatrix} h_1(m_1) & \dots & h_1(m_s) \\ h_2(m_1) & \dots & h_2(m_s) \\ h_3(m_1) & \dots & h_3(m_s) \\ h_4(m_1) & \dots & h_4(m_s) \\ h_5(m_1) & \dots & h_5(m_s) \\ -(h_1 + h_2 + h_4 + h_5 - h_6)(m_1) & \dots & -(h_1 + h_2 + h_4 + h_5 - h_6)(m_s) \\ -(h_3 + h_6)(m_1) & \dots & -(h_3 + h_6)(m_s) \end{pmatrix}$$

where  $m_1, \dots, m_s$  are the points in  $\mathcal{A}$  and  $h_1, \dots, h_6$  are defined to be

$$\begin{array}{lll} h_1(s, t, v) = s, & h_2(s, t, v) = t, & h_3(s, t, v) = v, \\ h_4(s, t, v) = al - s - av & h_5(s, t, v) = bl - t - bv, & h_6(s, t, v) = l - v. \end{array}$$

We were able to add  $h_6$  to the negative rows  $-(h_1 + h_4)$  and  $-(h_2 + h_5)$  since all three rows are colinear in this case. As a result, the positive part of the minimal Horn matrix coincides with the lattice distance matrix of  $\mathcal{A}$ .

### 3.3.2 The simple prisms with fewer facets

The trapezoidal wedges (A), tensor product wedges (B), triangular frusta (C) and triangular based pyramids (C) from Table 1 are simple prisms with less than 6 facets. The primitive collections in each case are:

|                                |                                   |
|--------------------------------|-----------------------------------|
| Trapezoidal wedges             | $\{n_1, n_4\}, \{n_2, n_3, n_5\}$ |
| Tensor product wedges $a' = 0$ | $\{n_1, n_3, n_4\}, \{n_2, n_5\}$ |
| Tensor product wedges $b' = 0$ | $\{n_1, n_4\}, \{n_2, n_3, n_5\}$ |
| Triangular based pyramid       | $\{n_1, n_2, n_3, n_4\}$          |
| Triangular frusta              | $\{n_1, n_2, n_4\}, \{n_3, n_6\}$ |

None of the polytopes above, except the triangular frusta, have an upper facet  $F_6$  and hence their normal fans and primitive collections do not include  $n_6$ . Also, the triangular based pyramid and triangular frusta have no back facet  $F_5$  and hence their normal fans and primitive collections do not include  $n_5$ . In each case, the primitive collections give a partition of the rays in the normal fan, hence the matrix  $M_{\mathcal{A}, \Sigma_P}$  associated to  $(P, w)$  is a Horn matrix for these cases. The question is whether this Horn matrix belongs to a Horn pair for  $(P, w)$ .

**Example 3.2.** Proposition 3.3 gives a Horn pair for the trapezoidal wedge in Table 1 (A), which can be reduced to a Horn pair  $(H, \lambda)$ , with:

$$H = \begin{pmatrix} h_1(m_1) & \dots & h_1(m_s) \\ h_2(m_1) & \dots & h_2(m_s) \\ h_3(m_1) & \dots & h_3(m_s) \\ h_4(m_1) & \dots & h_4(m_s) \\ h_5(m_1) & \dots & h_5(m_s) \\ -(h_1 + h_4)(m_1) & \dots & -(h_1 + h_4)(m_s) \\ -(h_2 + h_5 - h_6)(m_1) & \dots & -(h_2 + h_5 - h_6)(m_s) \\ -(h_3 + h_6)(m_1) & \dots & -(h_3 + h_6)(m_s) \end{pmatrix}$$

where  $m_1, \dots, m_s$  are the points in  $\mathcal{A}$  and  $h_1, \dots, h_9$  are defined to be

$$\begin{aligned} h_1(s, t, v) &= s, & h_2(s, t, v) &= t, & h_3(s, t, v) &= v \\ h_4(s, t, v) &= (a + db)l - s - dt - (a - a' + db)v, & h_5(s, t, v) &= bl - bv - t, & h_6(s, t, v) &= l - v. \end{aligned}$$

Let us compare  $H$  with the matrix  $M_{\mathcal{A}, \Sigma_P}$

$$M_{\mathcal{A}, \Sigma_P} = \begin{pmatrix} h_1(m_1) & \dots & h_1(m_s) \\ h_2(m_1) & \dots & h_2(m_s) \\ h_3(m_1) & \dots & h_3(m_s) \\ h_4(m_1) & \dots & h_4(m_s) \\ h_5(m_1) & \dots & h_5(m_s) \\ -(h_1 + h_4)(m_1) & \dots & -(h_1 + h_4)(m_s) \\ -(h_2 + h_3 + h_5)(m_1) & \dots & -(h_2 + h_3 + h_5)(m_s) \end{pmatrix}$$



where  $m_1, \dots, m_s$  and  $h_1, \dots, h_5$  are as in  $H$ . For  $b = 1$ , we see that  $h_2 + h_5 - h_6 = 0$ ,  $h_3 + h_6 = l$ , and  $h_2 + h_3 + h_5 = l$ , thus  $H = M_{\mathcal{A}, \Sigma_P}$ . For  $b > 1$ ,  $H$  and  $M_{\mathcal{A}, \Sigma_P}$  are minimal Horn matrices, hence, by uniqueness,  $M_{\mathcal{A}, \Sigma_P}$  cannot give rise to a Horn pair for  $(P, w)$ .

For all other examples of simple prisms with fewer facets, we noticed a similar phenomenon. Firstly, if  $n_i$  is not in the normal fan, then the positive row  $h_i$  is collinear with a negative row. In particular, for all these examples the positive part of the minimal Horn matrix coincides with the lattice distance matrix of  $\mathcal{A}$ . Below we summarise for which parameters the matrix  $M_{\mathcal{A}, \Sigma_P}$  gives rise to a Horn pair for  $(P, w)$ , this is not true in general for these families of ‘simple prisms with fewer facets’.

|                                |             |
|--------------------------------|-------------|
| Trapezoidal wedges             | $b = 1$     |
| Tensor product wedges $a' = 0$ | $a = 1$     |
| Tensor product wedges $b' = 0$ | $b = 1$     |
| Triangular based pyramid       | $b = d = 1$ |
| Triangular frusta              | $d = 1$     |

These seemingly arbitrary constraints have a nice geometrical interpretation. The constraint  $b = 1$  forces the triangular facet  $F_1$  in the trapezoidal wedges and tensor product wedges ( $b' = 0$ ) to be a simplex. The constraint  $a = 1$  forces the triangular facet  $F_2$  in the tensor product wedges ( $a' = 0$ ) to be a simplex. The constraint  $b = d = 1$  on the triangular based pyramid, means it is a 3D simplex and the constraint  $d = 1$  on the triangular frusta forces the two triangular facets  $F_3$  and  $F_6$  to be simplices. All the prisms considered in this section, except the ones just described, are examples of polytopes with simplicial normal fans for which the answer to Question 2 is negative.

### 3.3.3 The trapezoidal and tensor product frusta

The primitive collections for the trapezoidal frusta and the tensor product frusta are:

$$\{n_1, n_4\} \quad \{n_2, n_5\} \quad \{n_3, n_6\}$$

It follows easily that the Horn matrix given by Proposition 3.3 is  $M_{\mathcal{A}, \Sigma_P}$ . This matrix is also the minimal Horn matrix for all trapezoidal frusta and for general tensor product frusta. However, there are cases of tensor product frusta, where two or more rows of this matrix are collinear and hence the minimal Horn matrix is not exactly  $M_{\mathcal{A}, \Sigma_P}$ . For an overview of all minimal Horn matrices for the family  $\mathcal{P}$  of prisms, see Table 2 in Appendix A.

**Example 3.3.** A Horn matrix associated to the tensor product frusta in Table 1 (B) is

$$M_{\mathcal{A}, \Sigma_P} = \begin{pmatrix} h_1(m_1) & \dots & h_1(m_n) \\ h_2(m_1) & \dots & h_2(m_n) \\ h_3(m_1) & \dots & h_3(m_n) \\ h_4(m_1) & \dots & h_4(m_n) \\ h_5(m_1) & \dots & h_5(m_n) \\ h_6(m_1) & \dots & h_6(m_n) \\ -(h_1 + h_4)(m_1) & \dots & -(h_1 + h_4)(m_n) \\ -(h_2 + h_5)(m_1) & \dots & -(h_2 + h_5)(m_n) \\ -(h_3 + h_6)(m_1) & \dots & -(h_3 + h_6)(m_n) \end{pmatrix}$$

where  $m_1, \dots, m_n$  are the points in  $\mathcal{A}$  and  $h_1, \dots, h_6$  are defined to be

$$\begin{aligned} h_1(s, t, v) &= s, & h_2(s, t, v) &= t, & h_3(s, t, v) &= v, \\ h_4(s, t, v) &= al - s - (a - a')v & h_5(s, t, v) &= bl - t - (b - b')v, & h_6(s, t, v) &= l - v. \end{aligned}$$

If we consider the subfamily of tensor product frusta such that  $a = \lambda b$ ,  $a' = \lambda b'$  for  $\lambda \geq 1$  or  $\lambda = \frac{1}{\mu}$  with  $\mu \geq 1$ , then the minimal Horn matrix is

$$H = \begin{pmatrix} h_1(m_1) & \dots & h_1(m_n) \\ h_2(m_1) & \dots & h_2(m_n) \\ h_3(m_1) & \dots & h_3(m_n) \\ h_4(m_1) & \dots & h_4(m_n) \\ h_5(m_1) & \dots & h_5(m_n) \\ h_6(m_1) & \dots & h_6(m_n) \\ -(h_1 + h_2 + h_4 + h_5)(m_1) & \dots & -(h_1 + h_2 + h_4 + h_5)(m_n) \\ -(h_3 + h_6)(m_1) & \dots & -(h_3 + h_6)(m_n) \end{pmatrix}$$

where  $m_1, \dots, m_n$  are the points in  $\mathcal{A}$  and  $h_1, \dots, h_6$ , are as above.

**Theorem 3.5.** *For all pairs in  $\mathcal{P}$ , the positive part of the minimal Horn matrix is the lattice distance matrix of  $\mathcal{A}$ . For the following subfamilies of  $\mathcal{P}$ , the matrix  $M_{\mathcal{A}, \Sigma_P}$  gives rise to a Horn pair for  $(P, w)$ .*

| Name of subfamily                  | Constraints on $a' \leq a, b' \leq b, d$ |
|------------------------------------|--|
| Trapezoidal wedges                 | $a' > 0, b' = 0, b = 1, d > 0$           |
| Tensor product wedges ( $a' = 0$ ) | $a' = 0, a = 1, b' > 0, d = 0$           |
| Tensor product wedges ( $b' = 0$ ) | $a' > 0, b' = 0, b = 1, d = 0$           |
| 3D simplex                         | $a' = a = b' = 0, b = 1, d = 1$          |
| Triangular frusta                  | $a' = a = 0, b > 0, d = 1$               |
| Tensor product frusta              | $a' > 0, b' > 0, d = 0$                  |
| Trapezoidal frusta                 | $a' > 0, b' > 0, d > 0$                  |

*Proof.* Let  $(P, w) \in \mathcal{P}$ , if  $M_{\mathcal{A}, \Sigma_P}$  is a Horn matrix, then after row reduction operations, we get a minimal Horn matrix. Comparing this matrix with the minimal Horn matrix associated to  $(P, w)$  in Table 2 (Appendix A) and by uniqueness of minimal Horn matrices, one can verify both statements on the theorem.  $\square$

## 4 Multinomial staged tree models

In this section we define multinomial staged tree models, we prove that every such model has rational MLE, Theorem 4.11, and we give a criterion to determine when such models are toric varieties for binary multinomial staged trees, Theorem 4.10. To each toric binary multinomial staged tree one can associate a polytope, by Theorem 2.3 such polytope has rational linear precision. These results imply our main theorem:

**Theorem 4.1.** *Polytopes of toric binary multinomial staged trees have rational linear precision.*

Our motivation to introduce this model class arose from the observation that the Horn pair of all 2D and 3D polytopes in Section 3 could be interpreted as a statistical model defined by an event tree with a specific choice of parametrisation. Multinomial staged tree models improve the understanding of polytopes with rational linear precision in 2D and 3D and give their generalization for higher dimensions.

### 4.1 Definition of multinomial staged tree

Let  $\mathcal{T} = (V, E)$  denote a directed rooted tree with vertex set  $V$  and edge set  $E$  such that all edges are directed away from the root. For  $v, w \in V$ , the directed edge from  $v$  to  $w$  is denoted  $v \rightarrow w$  and the set of edges in the path from  $v$  to  $w$  is denoted  $\lambda_{v,w}$ . The set of children of  $v$  is  $\text{ch}(v) := \{u : v \rightarrow u \in E\}$  and the set of outgoing edges from  $v$  is  $E(v) := \{v \rightarrow u : u \in \text{ch}(v)\}$ . An element  $v \in V$  is a leaf if  $\text{ch}(v) = \emptyset$  and  $\tilde{V}$  denotes the set of non-leaf vertices of  $\mathcal{T}$ .

Let  $S = \{s_i : i \in I\}$  be a set of symbols indexed by the set  $I$ , and  $I_1, \dots, I_m$  a partition of  $I$  which induces a partition  $S_1, \dots, S_m$  of  $S$ . For  $a \in \mathbb{Z}_{\geq 1}$  and  $\ell \in [m]$ , a *floret of degree  $a$* , on  $S_\ell$ , is the set of terms in the expansion of the polynomial expression  $(\sum_{i \in I_\ell} s_i)^a$ , we denote this set by  $f_{\ell,a}$ . For instance if  $S_\ell = \{s_0, s_1\}$  and  $a = 3$ ,  $f_{\ell,3} = \{s_0^3, 3s_0^2s_1, 3s_0s_1^2, s_1^3\}$ . Note that  $f_{\ell,1} = S_\ell$ . A function  $\mathcal{L} : E \rightarrow \bigcup_{\ell \in [m], a \in \mathbb{Z}_{\geq 1}} f_{\ell,a}$  is a *labeling* of  $\mathcal{T}$  if for every  $v \in \tilde{V}$ ,  $\mathcal{L}(E(v)) = f_{\ell,a}$  for some  $\ell \in [m]$ ,  $a \in \mathbb{Z}_{\geq 1}$ , and the restriction  $\mathcal{L}_v : E(v) \rightarrow f_{\ell,a}$  is a bijection. Thus, for each  $v \in \tilde{V}$  and  $e \in E(v)$ ,  $\mathcal{L}_v(e)$  is a term in the multinomial expansion of  $(\sum_{i \in I_\ell} s_i)^a$ . When  $\text{im}(\mathcal{L}_v) = f_{\ell,a}$ , we fix the convention to index the children of  $v$  by  $v(K)$  where  $i_1, \dots, i_{|I_\ell|}$  is a fixed ordering of the elements in  $I_\ell$  and  $K = (k_{i_1}, \dots, k_{i_{|I_\ell|}}) \in \mathbb{N}^{|I_\ell|}$  is a tuple of nonnegative integers that add to  $a$ . For any  $K \in \mathbb{N}^{|I_\ell|}$ ,  $|K| := \sum_{q=1}^{|I_\ell|} k_{i_q}$ . Hence, for  $\text{im}(\mathcal{L}_v) = f_{\ell,a}$ ,  $E(v) = \{v \rightarrow v(K) : K \in \mathbb{N}^{|I_\ell|}, |K| = a\}$ . To illustrate, if  $v$  is such that  $\text{im}(\mathcal{L}_v) = f_{\ell,3}$  and  $S_\ell = \{s_0, s_1\}$  with this convention,  $\text{ch}(v) = \{v(3,0), v(2,1), v(1,2), v(0,3)\}$  and  $E(v) = \{v \rightarrow v(3,0), v \rightarrow v(2,1), v \rightarrow v(1,2), v \rightarrow v(0,3)\}$ . We further assume that the indexing of the children is compatible with the labeling  $\mathcal{L}$ , namely for all multinomial stages trees,  $\mathcal{L}_v(v \rightarrow v(K)) = \binom{a}{k_{i_1}, \dots, k_{i_{|I_\ell|}}} \prod_{q=1}^{|I_\ell|} s_{i_q}^{k_{i_q}}$ . To simplify notation we also write  $\binom{a}{K}$  for the previous multinomial coefficient.

**Definition 4.1.** Let  $\mathcal{T}$  be a tree and  $\mathcal{L}$  a fixed labelling of  $\mathcal{T}$ . A *multinomial staged tree* is a pair  $(\mathcal{T}, \mathcal{L})$ . The sets  $S_1, \dots, S_m$  in the partition of  $S$  are called the *stages* of  $\mathcal{T}$ . If  $\mathcal{T}$  is such that  $|S_\ell| = 2$  for all  $\ell \in [m]$ , then  $(\mathcal{T}, \mathcal{L})$  is a *binary multinomial staged tree*. We write

$v \sim w$  whenever  $E(v)$  and  $E(w)$  are labeled by florets on the same set of symbols  $S_\ell$ ,  $\ell \in [m]$ . Fix  $J$  to denote the set of root-to-leaf paths in  $\mathcal{T}$ . For  $j \in J$ , define  $p_j$  to be the product of all edge labels in the path  $j$ . Let  $c_j$  be the coefficient of  $p_j$  and  $a_j$  the exponent vector of the symbols  $(s_i)_{i \in I}$  in  $p_j$ . With this notation,  $p_j = c_j \prod_{i \in I} s_i^{a_{ij}}$ , where  $a_{ij}$  are the entries of  $a_j$ .

*Remark 4.1.* If all of the florets in a multinomial staged tree have degree one, then it is called a staged tree. Multinomial staged tree models are a generalization of staged tree models which were introduced in [15].

*Remark 4.2.* We will always consider a multinomial staged tree  $(\mathcal{T}, \mathcal{L})$  as an embedded tree in the plane. This means the tree has a fixed ordering of its edges and vertices. We also assume that each root-to-leaf path contains at most one monomial term from each floret. Several figures in Section 5 contain multinomial staged trees, in these pictures, for simplicity, we omit the coefficients of the monomial edge labels.

**Definition 4.2.** Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree, with index set  $I = \sqcup_{\ell \in [m]} I_\ell$ . Fix  $J$  to be the set of root-to-leaf paths in  $\mathcal{T}$ , with  $|J| = n + 1$ . Define the parameter space

$$\Theta_{\mathcal{T}} := \{(\theta_i)_{i \in I} \in (0, 1)^{|I|} : \sum_{i \in I_\ell} \theta_i = 1 \text{ for all } \ell \in [m]\}$$

The *multinomial staged tree model*  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  is the image of the parameterisation

$$\phi_{\mathcal{T}} : \Theta_{\mathcal{T}} \longrightarrow \Delta_n^\circ, (\theta_i)_{i \in I} \mapsto (c_j \prod_{i \in I} \theta_i^{a_{ij}})_{j \in J}.$$

*Remark 4.3.* The sum-to-one conditions on the parameter space  $\Theta_{\mathcal{T}}$  imply that the image of  $\phi_{\mathcal{T}}$  is contained in  $\Delta_n^\circ$ . The multinomial coefficients on the labels of  $\mathcal{T}$  are necessary for this condition to hold. The model  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  is an algebraic variety inside  $\Delta_n^\circ$  with an explicit parameterisation given by  $\phi_{\mathcal{T}}$ . For  $\theta \in \Theta_{\mathcal{T}}$ ,  $\text{eval}_\theta$  is the evaluation map  $s_i \mapsto \theta_i$ . The  $j$ -th coordinate of  $\phi_{\mathcal{T}}$  is  $\text{eval}_\theta(p_j)$ , where  $p_j = c_j \prod_{i \in I} s_i^{a_{ij}}$  (Definition 4.1). For this reason we also use  $p_j$  to denote the  $j$ -th coordinate in the probability simplex  $\Delta_n^\circ$ .

**Example 4.1.** The multinomial model which encodes the experiment of rolling a  $k$ -sided die  $N$  times and recording all possible outcomes is a multinomial staged tree model. In this case there is a single stage  $\{s_1, \dots, s_k\}$ , each symbol in the stage represents the probability of the die showing face  $i$  at the top. The tree consists of single floret of edges whose labels are the terms in the expansion of  $(s_1 + \dots + s_k)^N$ .

**Example 4.2.** Consider the following experiment with two independent coins: Toss the first coin  $b$  times and record the number of tails, say this number is  $j$ . Then toss the second coin  $a + d(b - j)$  times, record the number of heads, say  $i$ . An outcome of this experiment is a pair  $(i, j)$  where  $i$  is the number of tails in the second sequence of coin tosses and  $j$  is the number of tails in the first. Let  $\mathcal{M}_{a,b,d} \subset \Delta_n$  be the statistical model consisting of all probability distributions that follow the experiment just described and let  $p_{ij}$  denote the probability of the outcome  $(i, j)$ . The model  $\mathcal{M}_{a,b,d}$  is parameterized by the map  $\phi : \Delta_1^\circ \times \Delta_1^\circ \rightarrow \mathcal{M}_{a,b,d}$ ,

$$(\theta_0, \theta_1) \times (\theta_2, \theta_3) \mapsto (p_{ij})_{\substack{0 \leq j \leq b \\ 0 \leq i \leq a + d(b-j)}} \text{ where } p_{ij} = \binom{b}{j} \binom{a + d(b-j)}{i} \theta_0^j \theta_1^{b-j} \theta_2^i \theta_3^{a+d(b-j)-i}.$$

This model depends on two independent parameters, thus it has dimension two. The model  $\mathcal{M}_{a,b,d}$  is a binary multinomial staged tree model, it is represented by  $\mathcal{T}_{a,b,d}$  in Figure 3.

**Definition 4.3.** Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree. Fix the polynomial rings  $\mathbb{R}[P_j : j \in J]$ ,  $\mathbb{R}[s_i : i \in I]$  and  $\mathbb{R}[s_i : i \in I]/\mathfrak{q}$  where  $\mathfrak{q} = \langle 1 - \sum_{i \in I_\ell} s_i : \ell \in [m] \rangle$ . We define

$$\Psi_{\mathcal{T}}^{\text{toric}} : \mathbb{R}[P_j : j \in J] \rightarrow \mathbb{R}[s_i : i \in I] \text{ by } P_j \mapsto c_j \prod_{i \in I} s_i^{a_{ij}}, \text{ and}$$

$$\Psi_{\mathcal{T}} : \mathbb{R}[P_j : j \in J] \rightarrow \mathbb{R}[s_i : i \in I]/\mathfrak{q} \text{ by } \Psi_{\mathcal{T}} = \pi \circ \Psi_{\mathcal{T}}^{\text{toric}}$$

where  $\pi : \mathbb{R}[s_i : i \in I] \rightarrow \mathbb{R}[s_i : i \in I]/\mathfrak{q}$  is the canonical projection to the quotient ring. The ideal  $\ker(\Psi_{\mathcal{T}}^{\text{toric}})$  is the toric ideal associated to  $(\mathcal{T}, \mathcal{L})$  and  $\ker(\Psi_{\mathcal{T}})$  is the model ideal associated to  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$ . Whenever  $\ker(\Psi_{\mathcal{T}}) = \ker(\Psi_{\mathcal{T}}^{\text{toric}})$ , we call  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  a toric model.

*Remark 4.4.* The ideal  $\ker(\Psi_{\mathcal{T}})$  defines the model  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  implicitly, namely  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})} = V(\ker(\Psi_{\mathcal{T}})) \cap \Delta_n^\circ$ . Because of the containment  $\ker(\Psi_{\mathcal{T}}^{\text{toric}}) \subset \ker(\Psi_{\mathcal{T}})$ ,  $V(\ker(\Psi_{\mathcal{T}}^{\text{toric}}))$  is a toric variety that contains  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$ . The polynomial  $1 - \sum_{j \in J} P_j$  is always an element in  $\ker(\Psi_{\mathcal{T}})$ , hence using this polynomial as a homogenising element, we shall always consider  $\ker(\Psi_{\mathcal{T}})$  as a homogeneous ideal in  $\mathbb{R}[P_j : j \in J]$ .

## 4.2 The ideal of model invariants for $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$

As is common in algebraic geometry, finding the explicit equations of the prime ideal  $\ker(\Psi_{\mathcal{T}})$  is hard. Luckily, the statistical insight of the problem allows us to find a nonprime ideal, usually referred to as the ideal of model invariants, that defines the model inside the probability simplex. In this section we define such an ideal and postpone the proof that it has the aforementioned property to Section 4.4.

For a vertex  $v \in V$ , define  $[v] := \{j \in J : \text{the path } j \text{ goes through the vertex } v\}$  and for  $(p_0, \dots, p_n) \in \Delta_n^\circ$  set  $p_{[v]} := \sum_{j \in [v]} p_j$ . The next lemma gives several equations that hold between a point in  $\Theta_{\mathcal{T}}$  and its image under  $\phi_{\mathcal{T}}$ .

**Lemma 4.2.** *Let  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  be a multinomial staged tree model where  $\mathcal{T} = (V, E)$ . Fix  $v \in \tilde{V}$  and suppose  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$  for  $\ell \in [m]$ ,  $a \in \mathbb{Z}_{\geq 1}$ . Set  $i_1, \dots, i_{|I_\ell|}$  to be a fixed ordering of the elements in  $I_\ell$ . Let  $(p_0, \dots, p_n) \in \mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  and  $\theta = (\theta_i)_{i \in I} \in \Theta_{\mathcal{T}}$  be such that  $\phi_{\mathcal{T}}(\theta) = (p_0, \dots, p_n)$ .*

(1) *For each  $K \in \mathbb{N}^{|I_\ell|}$  with  $|K| = a$ ,*

$$\frac{p_{[v(K)]}}{p_{[v]}} = \binom{a}{k_{i_1}, \dots, k_{i_{|I_\ell|}}} \prod_{q=1}^{|I_\ell|} \theta_{i_q}^{k_{i_q}}.$$

(2) *Let  $K^1, K^2, K^3, K^4 \in \mathbb{N}^{|I_\ell|}$  with  $|K^1| = |K^2| = |K^3| = |K^4| = a$ , be such that  $K^1 + K^2 = K^3 + K^4$ . Define  $C_{(K^1, K^2)} := \binom{a}{K^1} \binom{a}{K^2}$  and similarly for  $C_{(K^3, K^4)}$ . Then*

$$C_{(K^3, K^4)} p_{[v(K^1)]} p_{[v(K^2)]} - C_{(K^1, K^2)} p_{[v(K^3)]} p_{[v(K^4)]} = 0$$

(3) *For each  $i_q \in I_\ell$ ,  $1 \leq q \leq |I_\ell|$*

$$\theta_{i_q} = \frac{\sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} p_{[v(K)]}}{a p_{[v]}}.$$

(4) Let  $w \in V$  and  $\text{im}(\mathcal{L}_w) = f_{\ell,b}$ . For all  $i_q \in I_\ell$ :

$$bp_{[w]} \left( \sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} p_{[v(K)]} \right) - ap_{[v]} \left( \sum_{|K'|=b, k'_{i_q} \geq 1} k'_{i_q} p_{[w(K')]} \right) = 0$$

*Proof.* (1) Since  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  is a probability tree, the transition probability from  $v$  to  $v(K)$  is the probability of arriving at  $v(K)$  divided by the probability of arriving at  $v$ , namely  $p_{[v(K)]}/p_{[v]}$ . By definition of  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$ , and since  $\mathcal{L}_v(v \rightarrow v(K)) = \binom{a}{k_{i_1}, \dots, k_{i_{|I_\ell|}}} \prod_{\alpha=1}^{|I_\ell|} s_{i_\alpha}^{k_{i_\alpha}}$ , this probability is exactly  $\binom{a}{k_{i_1}, \dots, k_{i_{|I_\ell|}}} \prod_{\alpha=1}^{|I_\ell|} \theta_{i_\alpha}^{k_{i_\alpha}}$ .

(2) This equality follows by direct substitution for the values from (1) and by noting that the coefficients  $C_{(K^1, K^2)}, C_{(K^3, K^4)}$  are needed to achieve cancellation.

(3) We start from the right-hand side, use (1), the fact that  $\sum_{i \in I_\ell} \theta_i = 1$ , and simplification with multinomial coefficients to arrive at  $\theta_{i_q}$ :

$$\begin{aligned} \frac{\sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} p_{[v(K)]}}{ap_{[v]}} &= \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} \frac{k_{i_q}}{a} \frac{p_{[v(K)]}}{p_{[v]}} = \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} \frac{k_{i_q}}{a} \binom{a}{k_{i_1}, \dots, k_{i_{|I_\ell|}}} \prod_{\alpha=1}^{|I_\ell|} \theta_{i_\alpha}^{k_{i_\alpha}} \\ &= \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} \frac{k_{i_q}}{k_{i_q}} \frac{a}{a} \binom{a-1}{k_{i_1}, \dots, k_{i_q}-1, \dots, k_{i_{|I_\ell|}}} \theta_{i_q} \theta_{i_q}^{k_{i_q}-1} \prod_{\substack{\alpha=1 \\ \alpha \neq q}}^{|I_\ell|} \theta_{i_\alpha}^{k_{i_\alpha}} \\ &= \theta_{i_q} \sum_{|K|=a-1} \binom{a-1}{k_{i_1}, \dots, k_{i_q}, \dots, k_{i_{|I_\ell|}}} \prod_{\alpha=1}^{|I_\ell|} \theta_{i_\alpha}^{k_{i_\alpha}} \\ &= \theta_{i_q} \left( \sum_{\alpha=1}^{|I_\ell|} \theta_{i_\alpha} \right)^{a-1} = \theta_{i_q}. \end{aligned}$$

(4) Applying part (3) to  $i_q$  for  $v$  and  $w$  separately yields

$$\theta_{i_q} = \frac{\sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} p_{[v(K)]}}{ap_{[v]}} = \frac{\sum_{|K'|=b, k'_{i_q} \geq 1} k'_{i_q} p_{[w(K')]}}{bp_{[w]}}.$$

After cross multiplication we get the desired equation in (4).  $\square$

From Lemma 4.2 parts (2) and (4), we obtain an idea to define the ideal of model invariants for a multinomial staged tree model. Namely, there are equations that must hold for every vertex, these are described in (2), and there are equations that must hold for every pair of vertices with the same associated set of symbols, these are described in (4).

**Definition 4.4.** Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree. We define

$$\begin{aligned} I_{\text{stages}} &:= \left\langle bp_{[w]} \left( \sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]} \right) - ap_{[v]} \left( \sum_{|K'|=b, k'_{i_q} \geq 1} k'_{i_q} P_{[w(K')]} \right) \right\rangle : \\ &v \sim w, \text{im}(\mathcal{L}_v) = f_{\ell,a}, \text{im}(\mathcal{L}_w) = f_{\ell,b}, \ell \in [m], 1 \leq q \leq |I_\ell|, \text{ and} \end{aligned}$$

$$I_{\text{vertices}} := \langle C_{(K^1, K^2)} P_{[v(K^3)]} P_{[v(K^4)]} - C_{(K^1, K^2)} P_{[v(K^3)]} P_{[v(K^4)]} : v \in \tilde{V}, \text{im}(\mathcal{L}_v) = f_{\ell, a}, \\ K^1, K^2, K^3, K^4 \in \mathbb{N}^{|\ell|}, |K^1| = |K^2| = |K^3| = |K^4| = a, K^1 + K^2 = K^3 + K^4 \rangle.$$

The *ideal of model invariants* of  $(\mathcal{T}, \mathcal{L})$  is  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} := I_{\text{stages}} + I_{\text{vertices}} + \langle 1 - \sum_{j \in J} P_j \rangle$ .

*Remark 4.5.* The generators of  $I_{\text{vertices}}$  for each fixed vertex  $v$  are similar to the Veronese relations of the embedding  $\nu_a : \mathbb{P}^{|\ell|-1} \rightarrow \mathbb{P}^M$  by monomials of total degree  $a$ . The only difference is the coefficients, defined in Lemma 4.2 part (2), that are needed for cancellation.

*Remark 4.6.* By definition,  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$  always contains the sum to one condition  $1 - \sum_{j \in J} P_j$ , thus in a similar way as for  $\ker(\Psi_{\mathcal{T}})$  in Remark 4.4, we always consider  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$  as a homogeneous ideal generated by  $I_{\text{stages}}$  and  $I_{\text{vertices}}$ .

### 4.3 Algebraic lemmas for multinomial staged trees

To understand the defining equations of  $\ker(\Psi_{\mathcal{T}})$  and the case when this ideal is toric, it is important to establish several lemmas that describe algebraic relations that hold in  $\mathbb{R}[P_j : j \in J]$ ,  $\mathbb{R}[s_i : i \in I]$ ,  $\mathbb{R}[P_j : j \in J]/\ker(\Psi_{\mathcal{T}})$  and  $\mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$ . The reader may decide to skip this section and only get back to it when the lemmas are used in the proofs of Theorem 4.8 and Theorem 4.10.

**Definition 4.5.** Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree with  $\mathcal{T} = (V, E)$ . For  $v \in V$ , let  $\Lambda_v$  denote the set of all  $v$ -to-leaf paths in  $\mathcal{T}$ . A path  $\lambda \in \Lambda_v$  is a sequence of edges  $v \rightarrow v_1 \rightarrow \dots \rightarrow v_\alpha$  where  $v_\alpha$  is a leaf of  $\mathcal{T}$ . For each  $v \in V$  we define the *interpolating polynomial* of  $v$ ,  $t(v) \in \mathbb{R}[s_i : i \in I]$ , by

$$t(v) := \sum_{\lambda \in \Lambda_v} \prod_{e \in \lambda} \mathcal{L}(e).$$

If  $v$  is a leaf,  $t(v) := 1$ . We denote by  $\overline{t(v)}$ , the image of  $t(v)$  under the canonical projection to  $\mathbb{R}[s_i : i \in I]/\mathfrak{q}$ . Note that for all  $v \in V$ ,  $\overline{t(v)} = 1$ .

**Lemma 4.3.** *Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree where  $\mathcal{T} = (V, E)$  and let  $v \in \tilde{V}$  be such that  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$ .*

(1) *The polynomial  $t(v)$  satisfies*

$$t(v) = \sum_{|K|=a} \binom{a}{K} \prod_{i \in I_\ell} s_i^{k_i} \cdot t(v(K));$$

(2) *The image of  $P_{[v]}$  under  $\Psi_{\mathcal{T}}^{\text{toric}}$  is  $(\prod_{e \in \lambda_{r,v}} \mathcal{L}(e)) \cdot \overline{t(v)}$ , where  $\lambda_{r,v}$  is the set of edges in the root-to- $v$  path in  $\mathcal{T}$ . Moreover  $\Psi_{\mathcal{T}}(P_{[v]}) = \prod_{e \in \lambda_{r,v}} \mathcal{L}(e)$ .*

*Proof.* (1) Any path in  $\Lambda_v$  goes through a child  $v(K)$  of  $v$ . The sum of all the edge products corresponding to the paths that go through child  $v(K)$  is equal to the sum of all the edge products corresponding to the paths starting at  $v(K)$  ( $t(v(K))$ ) multiplied by the label of the edge from  $v$  to  $v(K)$  ( $\binom{a}{K} \prod_{i \in I_\ell} s_i^{k_i}$ ). Taking the sum of this expression over all children of  $v$  gives the desired result.

(2) Let  $j$  be a root-to-leaf path that goes through  $v$ . Then  $j$  is the concatenation of a path from the root to  $v$ , denoted by  $\lambda_{r,v}$  and a path from  $v$  to the leaf denoted by  $\lambda_{v,j}$ . Then

$$\Psi_{\mathcal{T}}^{\text{toric}}(P_{[v]}) = \sum_{j \in [v]} \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \prod_{e \in \lambda_{v,j}} \mathcal{L}(e) = \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) \left( \sum_{j \in [v]} \prod_{e \in \lambda_{v,j}} \mathcal{L}(e) \right) = \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) t(v).$$

The second statement follows by noting that  $\overline{t(v)} = 1$ .  $\square$

**Lemma 4.4.** *Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree, there is a containment of ideals  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} \subset \ker(\Psi_{\mathcal{T}})$  in  $\mathbb{R}[P_j : j \in J]$ .*

*Proof.* To show that  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} \subset \ker(\Psi_{\mathcal{T}})$  it suffices to show that the generators of  $I_{\text{stages}}$  and  $I_{\text{vertices}}$  are zero after applying  $\Psi_{\mathcal{T}}$ . We present the proof for the generators of  $I_{\text{stages}}$ , the proof for  $I_{\text{vertices}}$  is similar and also uses Lemma 4.3. A generator of  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$  is of the form  $bP_{[w]} \left( \sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]} \right) - aP_{[v]} \left( \sum_{|K'|=b, k'_{i_q} \geq 1} k'_{i_q} P_{[w(K')]} \right)$  where  $v, w \in \tilde{V}$ ,  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$ ,  $\text{im}(\mathcal{L}_w) = f_{\ell, b}$  for some  $\ell \in [m]$  and a fixed  $q$ ,  $1 \leq q \leq |I_{\ell}|$ .

Claim:  $\Psi_{\mathcal{T}} \left( \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} k_{i_q} P_{[v(K)]} \right) = a s_{i_q} \prod_{e \in \lambda_{r,v}} \mathcal{L}(e)$ .

$$\begin{aligned} \Psi_{\mathcal{T}} \left( \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} k_{i_q} P_{[v(K)]} \right) &= \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} k_{i_q} \Psi_{\mathcal{T}}(P_{[v(K)]}) \stackrel{\text{Lemma 4.3}}{=} \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} k_{i_q} \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) \binom{a}{K} \prod_{\alpha=1}^{|I_{\ell}|} s_i^{k_{i\alpha}} \\ &= \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) \left( \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} k_{i_q} \binom{a}{K} \prod_{\alpha=1}^{|I_{\ell}|} s_i^{k_{i\alpha}} \right) \\ &= \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) \left( \sum_{\substack{|K|=a \\ k_{i_q} \geq 1}} k_{i_q} \frac{a(a-1)!}{k_{i_1}! \cdots k_{i_{|I_{\ell}|}}!} s_{i_q} s_{i_q}^{k_q-1} \prod_{\substack{\alpha=1 \\ \alpha \neq q}}^{|I_{\ell}|} s_i^{k_{i\alpha}} \right) \\ &= \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) \left( \sum_{|K|=a-1} a s_{i_q} \binom{a-1}{K} \prod_{\alpha=1}^{|I_{\ell}|} s_i^{k_{i\alpha}} \right) \\ &= \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) a s_{i_q} \left( \sum_{\alpha=1}^{|I_{\ell}|} s_{i\alpha} \right)^{a-1} = \left( \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \right) a s_{i_q}. \end{aligned}$$

The last equality follows from the fact that  $\sum_{\alpha=1}^{|I_{\ell}|} s_{i\alpha} = 1$  in  $\mathbb{R}[s_i : i \in I]/\mathfrak{q}$ . The claim applied to  $w \in \tilde{V}$ , implies  $\Psi_{\mathcal{T}} \left( \sum_{\substack{|K|=b \\ k_{i_q} \geq 1}} k_{i_q} P_{[w(K)]} \right) = \left( \prod_{e \in \lambda_{r,w}} \mathcal{L}(e) \right) b s_{i_q}$ . Thus, by Lemma 4.3,

$$\begin{aligned} \Psi_{\mathcal{T}} \left( bP_{[w]} \left( \sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]} \right) - aP_{[v]} \left( \sum_{|K'|=b, k'_{i_q} \geq 1} k'_{i_q} P_{[w(K')]} \right) \right) &= \\ b \prod_{e \in \lambda_{r,w}} \mathcal{L}(e) \cdot a s_{i_q} \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) - a \prod_{e \in \lambda_{r,v}} \mathcal{L}(e) \cdot b s_{i_q} \prod_{e \in \lambda_{r,w}} \mathcal{L}(e) &= 0. \end{aligned}$$

$\square$



### 4.3.1 Binary multinomial staged trees

From this point on we assume in  $(\mathcal{T}, \mathcal{L})$  is a binary multinomial staged tree, and modify our notation according to this assumption.

**Lemma 4.5.** *Let  $(\mathcal{T}, \mathcal{L})$  be a binary multinomial staged tree where  $\mathcal{T} = (V, E)$  and let  $v \in \tilde{V}$  be such that  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$ .*

*Fix  $k_0 \in \{0, \dots, a\}$ . The following equalities hold in  $\mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$ :*

$$\binom{a}{k_0 + k} P_{[v(k_0, a-k_0)]} l_1^{k_0+k} l_2^{a-(k_0+k)} = \binom{a}{k_0} P_{[v(k_0+k, a-(k_0+k))]} l_1^{k_0} l_2^{a-k_0}, \quad 1 \leq k \leq a - k_0, \quad (7)$$

$$\binom{a}{k_0 - k} P_{[v(k_0, a-k_0)]} l_1^{k_0-k} l_2^{a-(k_0-k)} = \binom{a}{k_0} P_{[v(k_0-k, a-(k_0-k))]} l_1^{k_0} l_2^{a-k_0}, \quad 1 \leq k \leq k_0, \quad (8)$$

where  $l_1 := \sum_{k=1}^a k P_{[v(k, a-k)]}$  and  $l_2 := \sum_{k=1}^a k P_{[v(a-k, k)]}$ .

For the proof of Lemma 4.5, we use Lemma 4.6.

**Lemma 4.6.** *Under the hypotheses from Lemma 4.5, the following equality holds in  $\mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$ :*

$$(a - k_0) P_{[v(k_0, a-k_0)]} l_1 = (k_0 + 1) P_{[v(k_0+1, a-(k_0+1))]} l_2. \quad (9)$$

*Proof.* By Definition 4.4 of the ideal of model invariants, we have

$$\binom{a}{k_0 + 1} \binom{a}{k - 1} P_{[v(k_0, a-k_0)]} P_{[v(k, a-k)]} - \binom{a}{k_0} \binom{a}{k} P_{[v(k_0+1, a-(k_0+1))]} P_{[v(k-1, a-(k-1))]} = 0$$

in  $\mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$ , or equivalently,

$$\frac{a! \left( (a - k_0) k P_{[v(k_0, a-k_0)]} P_{[v(k, a-k)]} - (k_0 + 1) (a - (k - 1)) P_{[v(k_0+1, a-(k_0+1))]} P_{[v(k-1, a-(k-1))]} \right)}{(k_0 + 1)! (a - k_0)! k! (a - (k - 1))!} = 0.$$

Thus we obtain  $(a - k_0) k P_{[v(k_0, a-k_0)]} P_{[v(k, a-k)]} = (k_0 + 1) (a - (k - 1)) P_{[v(k_0+1, a-(k_0+1))]} P_{[v(k-1, a-(k-1))]}$ . Using this identity and the definition of  $l_1$  in Lemma 4.5, it follows that

$$\begin{aligned} (a - k_0) P_{[v(k_0, a-k_0)]} l_1 &= (a - k_0) P_{[v(k_0, a-k_0)]} \sum_{k=1}^a k P_{[v(k, a-k)]} = \sum_{k=1}^a (a - k_0) k P_{[v(k_0, a-k_0)]} P_{[v(k, a-k)]} \\ &= \sum_{k=1}^a (k_0 + 1) (a - (k - 1)) P_{[v(k_0+1, a-(k_0+1))]} P_{[v(k-1, a-(k-1))]} \\ &= (k_0 + 1) P_{[v(k_0+1, a-(k_0+1))]} l_2. \end{aligned}$$

□

We are now ready to prove Lemma 4.5.

*Proof.* Let us first prove equality (7). We will do this by mathematical induction on  $k$ . First we show that (7) holds for  $k = 1$ :

$$\begin{aligned} \binom{a}{k_0+1} P_{[v(k_0, a-k_0)]} l_1^{k_0+1} l_2^{a-(k_0+1)} &\stackrel{(9)}{=} \binom{a}{k_0+1} \frac{k_0+1}{a-k_0} P_{[v(k_0+1, a-(k_0+1))]} l_1^{k_0} l_2^{a-k_0} \\ &= \binom{a}{k_0} P_{[v(k_0+1, a-(k_0+1))]} l_1^{k_0} l_2^{a-k_0}. \end{aligned}$$

Let us now suppose that (7) holds for  $k$ , and prove it for  $k+1$ .

$$\begin{aligned} \binom{a}{k_0+k+1} P_{[v(k_0, a-k_0)]} l_1^{k_0+k+1} l_2^{a-(k_0+k+1)} \\ &= \frac{a!(a-(k_0+k))}{(k_0+k+1)(k_0+k)!(a-(k_0+k))!} P_{[v(k_0, a-k_0)]} l_1^{k_0+k+1} l_2^{a-(k_0+k+1)} \\ &= \binom{a}{k_0+k} \frac{a-(k_0+k)}{k_0+k+1} P_{[v(k_0, a-k_0)]} l_1^{k_0+k} l_2^{a-(k_0+k+1)} l_2^{-1} \end{aligned}$$

Using (7) for  $k$ , we further simplify to

$$\begin{aligned} &\stackrel{hyp.(7)}{=} \binom{a}{k_0} P_{[v(k_0+k, a-(k_0+k))]} l_1^{k_0} l_2^{a-k_0} \frac{a-(k_0+k)}{k_0+k+1} l_1 l_2^{-1} \\ &\stackrel{(9)}{=} \binom{a}{k_0} \frac{a-(k_0+k)}{k_0+k+1} \frac{k_0+k+1}{a-(k_0+k)} P_{[v(k_0+k+1, a-(k_0+k+1))]} l_1^{k_0} l_2^{a-k_0} l_2^{-1} \\ &= \binom{a}{k_0} P_{[v(k_0+k+1, a-(k_0+k+1))]} l_1^{k_0} l_2^{a-k_0}. \end{aligned}$$

Let us now prove equality (8). By (9), we have

$$(a-k_0-1) P_{[v(k_0-1, a-(k_0-1))]} l_1 = k_0 P_{[v(k_0, a-k_0)]} l_2.$$

We will again use mathematical induction on  $k$ . First we show that (8) holds for  $k = 1$ :

$$\begin{aligned} \binom{a}{k_0-1} P_{[v(k_0, a-k_0)]} l_1^{k_0-1} l_2^{a-(k_0-1)} &= \binom{a}{k_0-1} \frac{a-(k_0-1)}{k_0} P_{[v(k_0-1, a-(k_0-1))]} l_1 l_2^{-1} l_1^{k_0-1} l_2^{a-(k_0-1)} \\ &= \binom{a}{k_0} P_{[v(k_0-1, a-(k_0-1))]} l_1^{k_0} l_2^{a-k_0}. \end{aligned}$$

Let us now suppose that (8) holds for  $k$ , and prove it for  $k+1$ .

$$\begin{aligned} \binom{a}{k_0-(k+1)} P_{[v(k_0, a-k_0)]} l_1^{k_0-(k+1)} l_2^{a-(k_0-(k+1))} \\ &= \binom{a}{k_0-k} \frac{k_0-k}{a-(k_0-(k+1))} P_{[v(k_0, a-k_0)]} l_1^{k_0-k} l_1^{-1} l_2^{a-(k_0-k)} l_2 \\ &\stackrel{hyp.(8)}{=} \binom{a}{k_0} P_{[v(k_0-k, a-(k_0-k))]} l_1^{k_0} l_2^{a-k_0} \frac{k_0-k}{a-(k_0-(k+1))} l_1^{-1} l_2 \\ &\stackrel{(9)}{=} \binom{a}{k_0} \frac{k_0-k}{a-(k_0-(k+1))} \frac{a-(k_0-(k+1))}{k_0-k} l_1 P_{[v(k_0-k-1, a-(k_0-k-1))]} l_1^{k_0} l_2^{a-k_0} l_1^{-1} \\ &= \binom{a}{k_0} P_{[v(k_0-(k+1), a-(k_0-(k+1)))]} l_1^{k_0} l_2^{a-k_0}. \end{aligned}$$

□

**Lemma 4.7.** *Under the same hypotheses as in Lemma 4.5, the following equalities hold in  $\mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$ :*

$$\binom{a}{k_0} \left( \frac{l_1}{aP[v]} \right)^{k_0} \left( \frac{l_2}{aP[v]} \right)^{a-k_0} = \frac{P_{[v(k_0, a-k_0)]}}{P_{[v]}}. \quad (10)$$

*Proof.* First, note that  $aP_{[v]} = (l_1 + l_2)$ . Indeed

$$\begin{aligned} l_1 + l_2 &= (P_{[v(1, a-1)]} + 2P_{[v(2, a-2)]} + \dots + (a-2)P_{[v(a-2, 2)]} + (a-1)P_{[v(a-1, 1)]} + aP_{[v(a, 0)]}) \\ &\quad + (aP_{[v(0, a)]} + (a-1)P_{[v(1, a-1)]} + (a-2)P_{[v(2, a-2)]} + \dots + 2P_{[v(a-2, 2)]} + P_{[v(a-1, 1)]}) \\ &= a(P_{[v(a, 0)]} + P_{[v(1, a-1)]} + P_{[v(2, a-2)]} + \dots + P_{[v(1, a-1)]} + P_{[v(0, a)]}) = aP_{[v]}. \end{aligned}$$

Therefore we have  $P_{[v(k_0, a-k_0)]}(aP_{[v]})^a = P_{[v(k_0, a-k_0)]}(l_1 + l_2)^a$ . Now, by Lemma 4.5, we obtain:

$$\begin{aligned} P_{[v(k_0, a-k_0)]}(l_1 + l_2)^a &= P_{[v(k_0, a-k_0)]} \sum_{k=0}^a \binom{a}{k} l_1^k l_2^{a-k} \\ &= \sum_{k=0}^a \binom{a}{k} P_{[v(k_0, a-k_0)]} l_1^k l_2^{a-k} \\ &= \sum_{k=1}^{k_0} \binom{a}{k_0 - k} P_{[v(k_0, a-k_0)]} l_1^{k_0 - k} l_2^{a - (k_0 - k)} + \binom{a}{k_0} P_{[v(k_0, a-k_0)]} l_1^{k_0} l_2^{a - k_0} \\ &\quad + \sum_{k=1}^{a-k_0} \binom{a}{k_0 + k} P_{[v(k_0, a-k_0)]} l_1^{k_0 + k} l_2^{a - (k_0 + k)} \\ &= \binom{a}{k_0} l_1^{k_0} l_2^{a - k_0} \left( \sum_{k=1}^{k_0} P_{[v(k_0 - k, a - (k_0 - k))]} + P_{[v(k_0, a - k_0)]} + \sum_{k=1}^{a - k_0} P_{[v(k_0 + k, a - (k_0 + k))]} \right) \\ &= \binom{a}{k_0} l_1^{k_0} l_2^{a - k_0} P_{[v]}. \end{aligned}$$

□

## 4.4 Defining equations of binary multinomial staged trees

In this section and the next, we prove Theorem 4.8 and Theorem 4.10 for binary multinomial staged trees; despite being unable to provide a proof, we believe these statements also hold for non-binary multinomial staged trees. First we show that the ring homomorphism  $\Psi_{\mathcal{T}}$  admits an inverse when localised at a suitable element. From this it follows as a corollary that the ideal of model invariants defines  $\mathcal{M}(\mathcal{T}, \mathcal{L})$  inside the probability simplex.

**Theorem 4.8.** *Let  $(\mathcal{T}, \mathcal{L})$  be a binary multinomial staged tree and set  $\mathbf{P} := \prod_{v \in V} P_{[v]}$ . Then, the localised map*

$$(\Psi_{\mathcal{T}})_{\mathbf{P}} : \left( \mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} \right)_{\mathbf{P}} \rightarrow \left( \mathbb{R}[s_i : i \in I]/\mathfrak{q} \right)_{\Psi_{\mathcal{T}}(\mathbf{P})},$$

*is an isomorphism of  $\mathbb{R}$ -algebras. Therefore  $(I_{\mathcal{M}(\mathcal{T}, \mathcal{L})})_{\mathbf{P}} = (\ker(\Psi_{\mathcal{T}}))_{\mathbf{P}}$  and  $(I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} : \mathbf{P}^{\infty}) = \ker(\Psi_{\mathcal{T}})$ .*

*Proof.* We define a ring homomorphism

$$\varphi : (\mathbb{R}[s_i : i \in I]/\mathfrak{q})_{\Psi_{\mathcal{T}}(\mathbf{P})} \rightarrow (\mathbb{R}[P_j : j \in J]/I_{\mathcal{M}(\mathcal{T}, \mathcal{L})})_{\mathbf{P}}$$

and show that it is a two sided inverse for  $(\Psi_{\mathcal{T}})_{\mathbf{P}}$ . For  $\ell \in [m]$  and  $1 \leq q \leq |I_\ell|$ , let  $v$  be a vertex with  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$  and define

$$\varphi(s_{i_q}) = \frac{\sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]}}{a P_{[v]}}.$$

Note that  $\varphi$  is well defined: If  $w$  is another vertex with  $\text{im}(\mathcal{L}_w) = f_{\ell, b}$ , then

$$\varphi(s_{i_q}) = \frac{\sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]}}{a P_{[v]}} = \frac{\sum_{|K'|=b, k'_{i_q} \geq 1} k_{i_q} P_{[w(K')]} }{b P_{[w]}}$$

because the polynomial  $b P_{[w]} \left( \sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]} \right) - a P_{[v]} \left( \sum_{|K'|=b, k'_{i_q} \geq 1} k'_{i_q} P_{[w(K')]} \right)$  is in  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$ . First, we check that  $(\Psi_{\mathcal{T}})_{\mathbf{P}} \circ \varphi = \text{Id}$ ,

$$\begin{aligned} (\Psi_{\mathcal{T}})_{\mathbf{P}}(\varphi(s_{i_q})) &= (\Psi_{\mathcal{T}})_{\mathbf{P}} \left( \frac{\sum_{|K|=a, k_{i_q} \geq 1} k_{i_q} P_{[v(K)]}}{a P_{[v]}} \right) = \sum_{|K|=a, k_{i_q} \geq 1} \frac{k_{i_q}}{a} (\Psi_{\mathcal{T}})_{\mathbf{P}} \left( \frac{P_{[v(K)]}}{P_{[v]}} \right) \\ &= \sum_{|K|=a, k_{i_q} \geq 1} \frac{k_{i_q}}{a} \frac{\Psi_{\mathcal{T}}(P_{[v(K)]})}{\Psi_{\mathcal{T}}(P_{[v]})} = \sum_{|K|=a, k_{i_q} \geq 1} \frac{k_{i_q}}{a} \binom{a}{K} \prod_{\alpha=1}^{|I_\ell|} s_{i_\alpha}^{k_{i_\alpha}} = s_{i_q}. \end{aligned}$$

The second to last equality follows by using the expression for  $\Psi_{\mathcal{T}}(P_{[v]})$  presented in Lemma 4.3 part (2), a variation of the same result to compute  $\Psi_{\mathcal{T}}(P_{[v(K)]})$ , and finally noting that their quotient  $\Psi_{\mathcal{T}}(P_{[v(K)]})/\Psi_{\mathcal{T}}(P_{[v]})$  is exactly  $\binom{a}{K} \prod_{\alpha=1}^{|I_\ell|} s_{i_\alpha}^{k_{i_\alpha}}$ . The last equality is obtained by using the same argument as in Lemma 4.2 part (3).

Next, we verify  $\varphi \circ (\Psi_{\mathcal{T}})_{\mathbf{P}} = \text{Id}$ , which amounts to proving that  $(\varphi \circ \Psi_{\mathcal{T}, \mathbf{P}})(P_j) = P_j$  for each  $j \in J$ . From this point on we further assume that  $|I_\ell| = 2$  for all  $\ell \in [m]$ . Fix  $j \in J$  and let  $v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_\alpha$  be the root-to-leaf path  $j$ . By Definition 4.3

$$(\Psi_{\mathcal{T}})_{\mathbf{P}}(P_j) = c_j \prod_{i \in I} s_i^{a_{ij}} = \mathcal{L}_{v_1}(v_1 \rightarrow v_2) \cdots \mathcal{L}_{v_{\alpha-1}}(v_{\alpha-1} \rightarrow v_\alpha). \quad (11)$$

Where for each  $\gamma \in [\alpha - 1]$ ,  $\text{im}(\mathcal{L}_{v_\gamma}) = f_{\ell_\gamma, a_\gamma}$  for some  $\ell_\gamma \in [m]$  and  $a_\gamma \in \mathbb{Z}_{\geq 1}$ . By Remark 4.2, none of the florets  $f_{\ell_\gamma, a_\gamma}$  share the same set of symbols. Moreover, for each  $\gamma \in [\alpha - 1]$ ,

$$\mathcal{L}_{v_\gamma}(v_\gamma \rightarrow v_{\gamma+1}) = \binom{a_\gamma}{k_\gamma} s_{\gamma, i_1}^{k_\gamma} s_{\gamma, i_2}^{a_\gamma - k_\gamma} \text{ where } S_{\ell_\gamma} = \{s_{\gamma, i_1}, s_{\gamma, i_2}\}, 0 \leq k_\gamma \leq a_\gamma. \quad (12)$$

With this notation, we also deduce that  $v_{\gamma+1} = v_\gamma(k_\gamma, a - k_\gamma)$ . Now we apply  $\varphi$  to (11), use that  $\varphi$  is a ring homomorphism and use equation (12) to obtain

$$\begin{aligned} \varphi((\Psi_{\mathcal{T}})_{\mathbf{P}}(P_j)) &= \varphi(\mathcal{L}_{v_1}(v_1 \rightarrow v_2)) \cdots \varphi(\mathcal{L}_{v_{\alpha-1}}(v_{\alpha-1} \rightarrow v_\alpha)) \\ &= \prod_{\gamma=1}^{\alpha-1} \binom{a_\gamma}{k_\gamma} \varphi(s_{\gamma, i_1})^{k_\gamma} \varphi(s_{\gamma, i_2})^{a_\gamma - k_\gamma}. \end{aligned} \quad (13)$$

Using the definition of  $\varphi$ , for each  $\gamma \in [\alpha - 1]$ ,

$$\varphi(s_{\gamma, i_1}) = \frac{\sum_{k=1}^{a_\gamma} k P_{[v_\gamma(k, a_\gamma - k)]}}{a_\gamma P_{[v_\gamma]}} \quad \text{and} \quad \varphi(s_{\gamma, i_2}) = \frac{\sum_{k=1}^{a_\gamma} k P_{[v_\gamma(a_\gamma - k, k)]}}{a_\gamma P_{[v_\gamma]}}.$$

By Lemma 4.7, with  $a = a_\gamma$ ,  $l_1 = \sum_{k=1}^{a_\gamma} k P_{[v_\gamma(k, a_\gamma - k)]}$ ,  $l_2 = \sum_{k=1}^{a_\gamma} k P_{[v_\gamma(a_\gamma - k, k)]}$  and  $k_0 = k_\gamma$ , we conclude that

$$\binom{a_\gamma}{k_\gamma} \varphi(s_{\gamma, i_1})^{k_\gamma} \varphi(s_{\gamma, i_2})^{a - k_\gamma} = \frac{P_{[v_\gamma(k_\gamma, a - k_\gamma)]}}{P_{[v_\gamma]}}.$$

Thus, continuing from (13) and using that for each  $\gamma \in [\alpha - 1]$ ,  $v_{\gamma+1} = v_\gamma(k_\gamma, a - k_\gamma)$ , we have

$$\varphi((\Psi_{\mathcal{T}})_{\mathbf{P}}(P_j)) = \prod_{\gamma=1}^{\alpha-1} \frac{P_{[v_\gamma(k_\gamma, a - k_\gamma)]}}{P_{[v_\gamma]}} = \frac{P_{[v_1(k_1, a_1 - k_1)]}}{P_{[v_1]}} \frac{P_{[v_2(k_2, a_2 - k_2)]}}{P_{[v_2]}} \dots \frac{P_{[v_{\alpha-1}(k_{\alpha-1}, a_{\alpha-1} - k_{\alpha-1})]}}{P_{[v_{\alpha-1}]}} \quad (14)$$

$$= \frac{P_{[v_1(k_1, a_1 - k_1)]}}{P_{[v_1]}} \frac{P_{[v_2(k_2, a_2 - k_2)]}}{P_{[v_1(k_1, a_1 - k_1)]}} \dots \frac{P_{[v_{\alpha-1}(k_{\alpha-1}, a_{\alpha-1} - k_{\alpha-1})]}}{P_{[v_{\alpha-2}(k_{\alpha-2}, a_{\alpha-2} - k_{\alpha-2})]}} \frac{P_{[v_\alpha]}}{P_{[v_{\alpha-1}(k_{\alpha-1}, a_{\alpha-1} - k_{\alpha-1})]}}. \quad (15)$$

Note that  $P_{[v_1]} = 1$  by definition of  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$  and  $P_{[v_\alpha]} = P_j$  because  $v_\alpha$  is the last vertex in the path  $j$ . Thus  $\varphi((\Psi_{\mathcal{T}})_{\mathbf{P}}(P_j)) = P_j$ . The second statement of the theorem follows from the fact that  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} \subset \ker(\Psi_{\mathcal{T}})$  and that the localisation  $(\Psi_{\mathcal{T}})_{\mathbf{P}}$  is an isomorphism.  $\square$

**Corollary 4.9.** *The ideal of model invariants defines the binary multinomial staged tree model inside the probability simplex, i.e.  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})} = V(I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}) \cap \Delta_n^\circ$ .*

*Proof.* The variety  $V(I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} : \mathbf{P}^\infty)$  exactly describes the points in  $V(I_{\mathcal{M}(\mathcal{T}, \mathcal{L})})$  that are not in  $V(\mathbf{P})$ . The latter variety contains the boundary of the simplex, hence restricting to positive points that add to one, yields  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})} = V(I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}) \cap \Delta_n^\circ$ .  $\square$

## 4.5 Toric binary multinomial staged tree models

It is not true in general the the ideal  $\ker(\Psi_{\mathcal{T}})$  of a multinomial staged tree is toric. For the case of staged trees, a characterization of when  $\ker(\Psi_{\mathcal{T}})$  is equal to a subideal generated by binomials is available in [9]. The goal of this section is to establish a similar criterion, based on interpolating polynomials from Definition 4.5, for multinomial staged trees. This criterion will allow us to study the polyhedral geometry of these models in Section 5.

**Definition 4.6.** Let  $(\mathcal{T}, \mathcal{L})$  be a multinomial staged tree and let  $v, w$  be two vertices in the same stage with  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$  and  $\text{im}(\mathcal{L}_w) = f_{\ell, b}$  for some  $\ell \in [m]$ .

- (1) The vertex  $v$  is balanced if for all  $K^1, K^2, K^3, K^4 \in \mathbb{N}^{|\ell|}$  with  $|K^1| = |K^2| = |K^3| = |K^4| = a$  and  $K^1 + K^2 = K^3 + K^4$ , the next identity holds in  $\mathbb{R}[s_i : i \in I]$

$$t(v(K^1))t(v(K^2)) = t(v(K^3))t(v(K^4)).$$

- (2) The pair of vertices  $v, w$  is balanced if for all tuples  $K, K', Q, Q' \in \mathbb{N}^{|\ell|}$  with  $|K| = |K'| = a$  and  $|Q| = |Q'| = b$  with  $K + Q = K' + Q'$  the following identity holds in  $\mathbb{R}[s_i : i \in I]$

$$t(v(K)) \cdot t(w(Q)) = t(v(K')) \cdot t(w(Q)).$$

The multinomial staged tree  $(\mathcal{T}, \mathcal{L})$  is *balanced* if every vertex is balanced and every pair of vertices in the same stage is balanced.

*Remark 4.7.* Condition (1) in Definition 4.6 is an empty condition for florets of degree one. For staged trees, condition (2) specializes to the definition of balanced stated in [2].

*Remark 4.8.* If all root-to-leaf paths in  $(\mathcal{T}, \mathcal{L})$  have length 1, then  $(\mathcal{T}, \mathcal{L})$  is vacuously balanced. If  $(\mathcal{T}, \mathcal{L})$  has all root-to-leaf paths of length 2, such as  $\mathcal{T}_{a,b,d}$  in Figure 3, it suffices to check that the root is balanced. For the other vertices, the conditions in Definition 4.6 reduce to the trivial equality  $1 \cdot 1 = 1 \cdot 1$ .

**Theorem 4.10.** *Let  $(\mathcal{T}, \mathcal{L})$  be a binary multinomial staged tree. The model  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  is toric if and only if  $(\mathcal{T}, \mathcal{L})$  is balanced.*

*Proof.* We prove that  $\ker(\Psi_{\mathcal{T}}) = \ker(\Psi_{\mathcal{T}}^{\text{toric}})$  if and only if  $(\mathcal{T}, \mathcal{L})$  is balanced. Define the ideal  $J$  to be generated by all polynomials of the form

$$C_{(K',Q)}P_{[v(K)]}P_{[w(Q')]} - C_{(K,Q')}P_{[v(K')] }P_{[w(Q)]}, \quad C_{(K^3,K^4)}P_{[v(K^1)]}P_{[v(K^2)]} - C_{(K^1,K^2)}P_{[v(K^3)]}P_{[v(K^4)]}$$

where  $v, w \in V$  are in the same stage and  $K, K', Q, Q'$  obey the condition (2) and  $K^1, K^2, K^3, K^4$  obey the condition (1) in Definition 4.6.

Claim 1:  $J \subset \ker(\Psi_{\mathcal{T}}^{\text{toric}})$  if and only if  $(\mathcal{T}, \mathcal{L})$  is balanced.

By Lemma 4.3,

$$\Psi_{\mathcal{T}}^{\text{toric}}(P_{[v(K)]}P_{[w(Q')]})) = \left( \prod_{e \in \lambda_{r,v(K)}} \mathcal{L}(e) \right) t(v(K)) \left( \prod_{e \in \lambda_{r,w(Q')}} \mathcal{L}(e) \right) t(w(Q')) \text{ and} \quad (16)$$

$$\Psi_{\mathcal{T}}^{\text{toric}}(P_{[v(K')] }P_{[w(Q)]}) = \left( \prod_{e \in \lambda_{r,v(K')}} \mathcal{L}(e) \right) t(v(K')) \left( \prod_{e \in \lambda_{r,w(Q)}} \mathcal{L}(e) \right) t(w(Q)). \quad (17)$$

Note that the right hand side of the two equations above share the common factor  $\prod_{e \in \lambda_{v,w}} \mathcal{L}(e)$  where  $\lambda_{v,w}$  is the set of edges in the path from  $v$  to  $w$ . Thus we extract this factor from the two previous equations and multiply times the labels of the edges  $v \rightarrow v(K), w \rightarrow w(Q)$  and  $v \rightarrow v(K'), w \rightarrow w(Q')$ , respectively, to further simplify the two expressions into

$$\Psi_{\mathcal{T}}^{\text{toric}}(P_{[v(K)]}P_{[w(Q')]})) = \prod_{e \in \lambda_{v,w}} \mathcal{L}(e) \left( \binom{a}{K} \binom{b}{Q'} \prod_{i \in I_{\ell}} s_i^{k_i+q'_i} \right) t(v(K))t(w(Q')) \text{ and} \quad (18)$$

$$\Psi_{\mathcal{T}}^{\text{toric}}(P_{[v(K')] }P_{[w(Q)]}) = \prod_{e \in \lambda_{v,w}} \mathcal{L}(e) \left( \binom{a}{K'} \binom{b}{Q} \prod_{i \in I_{\ell}} s_i^{k'_i+q_i} \right) t(v(K'))t(w(Q)) \quad (19)$$

Finally, since  $K + Q' = K' + Q$  and  $(\mathcal{T}, \mathcal{L})$  is balanced, we obtain

$$\begin{aligned} & \Psi_{\mathcal{T}}^{\text{toric}}(C_{(K',Q)}P_{[v(K)]}P_{[w(Q')]} - C_{(K,Q')}P_{[v(K')] }P_{[w(Q)]}) = \\ & C_{(K,Q')}C_{(K',Q)} \prod_{e \in \lambda_{v,w}} \mathcal{L}(e) \prod_{i \in I_{\ell}} s_i^{k_i+q'_i} (t(v(K))t(w(Q)) - t(v(K'))t(w(Q))) = 0 \end{aligned}$$

A similar calculation shows that  $\Psi_{\mathcal{T}}^{\text{toric}}(C_{(K^3,K^4)}P_{[v(K^1)]}P_{[v(K^2)]} - C_{(K^1,K^2)}P_{[v(K^3)]}P_{[v(K^4)]}) = 0$  if  $\mathcal{T}$  is balanced. Conversely, note that if  $J \subset \ker(\Psi_{\mathcal{T}}^{\text{toric}})$ , the same equations as before show

that  $\mathcal{T}$  must be balanced.

Claim 2:  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})} \subset J$ . The ideal  $I_{\mathcal{M}(\mathcal{T}, \mathcal{L})}$  is the sum of  $I_{\text{stages}}$  and  $I_{\text{vertices}}$ . By definition, the generators of  $I_{\text{vertices}}$  are also generators of  $J$ . Hence it suffices to show that the generators of  $I_{\text{stages}}$  are polynomial combinations of the generators of  $J$ . From this point on we further assume that  $(\mathcal{T}, \mathcal{L})$  is binary. Suppose  $v, w$  are in the same stage, where  $\text{im}(\mathcal{L}_v) = f_{\ell, a}$ ,  $\text{im}(\mathcal{L}_w) = f_{\ell, b}$  and  $|I_\ell| = 2$ . There are two equations that hold for this stage, one for each element in  $I_\ell$ . We will show that the equation

$$bP_{[w]} \left( \sum_{k_1=1}^a k_1 P_{[v(k_1, a-k_1)]} \right) - aP_{[v]} \left( \sum_{k_2=1}^b k_2 P_{[w(k_2, b-k_2)]} \right), \quad (20)$$

which is the equation for the first element in  $I_\ell$ , is a combination the generators of  $J$ , defined at the beginning. The one for the second element in  $I_\ell$  follows an analogous argument. We use the following two identities:

$$bP_{[w]} = \sum_{k_2=1}^b k_2 P_{[w(k_2, b-k_2)]} + \sum_{k_2=1}^b k_2 P_{w[(b-k_2, k_2)]} \quad \text{and} \quad aP_{[v]} = \sum_{k_1=1}^a k_1 P_{[v(k_1, a-k_1)]} + \sum_{k_1=1}^a k_1 P_{[v(a-k_1, k_1)]}.$$

Working from equation (20), using the identities, we have

$$\begin{aligned} & \left( \sum_{k_2=1}^b k_2 P_{[w(k_2, b-k_2)]} + \sum_{k_2=1}^b k_2 P_{[w(b-k_2, k_2)]} \right) \left( \sum_{k_1=1}^a k_1 P_{[v(k_1, a-k_1)]} \right) - \\ & \left( \sum_{k_1=1}^a k_1 P_{[v(k_1, a-k_1)]} + \sum_{k_1=1}^a k_1 P_{[v(a-k_1, k_1)]} \right) \left( \sum_{k_2=1}^b k_2 P_{[w(k_2, b-k_2)]} \right) = \\ & \left( \sum_{k_2=1}^b k_2 P_{[w(b-k_2, k_2)]} \right) \left( \sum_{k_1=1}^a k_1 P_{[v(k_1, a-k_1)]} \right) - \left( \sum_{k_1=1}^a k_1 P_{[v(a-k_1, k_1)]} \right) \left( \sum_{k_2=1}^b k_2 P_{[w(k_2, b-k_2)]} \right) = \\ & \left( \sum_{k_2=1}^b \sum_{k_1=1}^a k_2 k_1 P_{[w(b-k_2, k_2)]} P_{[v(k_1, a-k_1)]} \right) - \left( \sum_{k_2=1}^b \sum_{k_1=1}^a k_1 k_2 P_{[w(k_2, b-k_2)]} P_{[v(a-k_1, k_1)]} \right) = \end{aligned} \quad (21)$$

After rearranging the second summation in equation (21) and combining into a single double summation in (22) we get

$$\sum_{k_2=1}^b \sum_{k_1=1}^a k_2 k_1 P_{[w(b-k_2, k_2)]} P_{[v(k_1, a-k_1)]} - (b - (k_2 - 1))(a - (k_1 - 1)) P_{[w(b-(k_2-1), k_2-1)]} P_{[v(k_1-1, a-(k_1-1))]} \quad (22)$$

Finally, the  $(k_1, k_2)$  summand in (22) is a multiple of the generator of  $J$ , where  $Q' = (b - k_2, k_2)$ ,  $K = (k_1, a - k_1)$ ,  $K' = (k_1 - 1, a - (k_1 - 1))$ ,  $Q = (b - (k_2 - 1), k_2 - 1)$ . The generator of  $J$  corresponding to this choice of  $K, K', Q, Q'$  is

$$\binom{b}{k_2 - 1} \binom{a}{k_1 - a} P_{[w(b-k_2, k_2)]} P_{[v(k_1, a-k_1)]} - \binom{b}{k_2} \binom{a}{k_1} P_{[w(b-(k_2-1), k_2-1)]} P_{[v(k_1, a-(k_1-1))]} \quad (23)$$

Note that  $(b - k_2, k_2) + (k_1, a - k_1) = (b - (k_2 - 1), k_2 - 1) + (k_1 - 1, a - (k_1 - 1))$ , thus  $K + Q' = K' + Q$ . Multiplying equation (23) times  $\frac{(b-(k_2-1))!(a-(k_1-1))!k_1!k_2!}{a!b!}$  gives the  $(k_1, k_2)$  summand in (22). This implies that (22) is a sum of multiples of the generators in  $J$ , hence  $I_{\text{stages}} \subset J$ .  $\square$

## 4.6 Multinomial staged tree models have rational MLE

In this last section on multinomial staged trees we prove that they have rational MLE. This fact together with Theorem 4.10 establishes Theorem 4.1 and thus provides a new class of polytopes that have rational linear precision.

**Theorem 4.11.** *The multinomial staged tree model  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  has rational MLE  $\Phi$ . The  $j$ -th coordinate of  $\Phi$  is*

$$\Phi_j(u_0, \dots, u_n) = c_j \prod_{i \in I} \hat{\theta}_i^{a_{ij}}, \quad \text{where for } i \in I_\ell, \hat{\theta}_i = \frac{\sum_{j \in J} u_j a_{ij}}{\sum_{i \in I_\ell} (\sum_{j \in J} u_j a_{ij})}.$$

*Proof.* Let  $u = (u_0, \dots, u_n)$  be a vector of counts. The likelihood function for  $\mathcal{M}_{\mathcal{T}, \mathcal{L}}$  is

$$\begin{aligned} L(p|u) &= \prod_{j \in J} p_j^{u_j} = \prod_{j \in J} \left( \prod_{i \in I} c_j \theta_i^{a_{ij}} \right)^{u_j} \\ &= \prod_{j \in J} \left( \prod_{i \in I} c_j^{u_j} \theta_i^{a_{ij} u_j} \right) = \left( \prod_{i \in I} c_0^{u_0} \theta_i^{a_{i0} u_0} \right) \dots \left( \prod_{i \in I} c_n^{u_n} \theta_i^{a_{in} u_n} \right) \\ &= \left( \prod_{j \in J} c_j^{u_j} \right) \left( \prod_{i \in I} \theta_i^{\sum_{j \in J} u_j a_{ij}} \right), \quad \text{let } C = \prod_{j \in J} c_j^{u_j} \\ &= C \left( \prod_{i \in I_1} \theta_i^{\sum_{j \in J} u_j a_{ij}} \right) \dots \left( \prod_{i \in I_k} \theta_i^{\sum_{j \in J} u_j a_{ij}} \right) = C L_1 \dots L_k, \end{aligned}$$

where  $L_1, \dots, L_m$  denote the factors before the last equality in the previous line. The function  $L(p|u)$  is maximized when each factor is maximized. This is because the parameters are partitioned by  $I_1, \dots, I_m$  and hence each factor is independent. Thus we find the maximizers of each factor. The function  $L_\ell$ ,  $\ell \in [m]$ , is the likelihood function of the saturated model  $\Delta_{|I_\ell|-1}$  with parameters  $(\theta_i)_{i \in I_\ell}$  and vector of counts  $(\sum_{j \in J} u_j a_{ij})_{i \in I_\ell}$ . Therefore  $\hat{\theta}_i = \frac{\sum_{j \in J} u_j a_{ij}}{\sum_{i \in I_\ell} \sum_{j \in J} u_j a_{ij}}$ ,  $i \in I_\ell$ .  $\square$

**Corollary 4.12.** *Let  $H_{(\mathcal{T}, \mathcal{L})}$  be the  $(|I| + m) \times (|J|)$  matrix with entries*

$$h_{ij} = a_{ij}, \quad i \in I, \quad h_{\ell j} = - \sum_{i \in I_\ell} a_{ij}, \quad \ell \in [m], \quad \text{and } \lambda_j := (-1)^{\sum_{i \in I} a_{ij}} c_j.$$

*Then  $(H_{(\mathcal{T}, \mathcal{L})}, \lambda)$  is a Horn pair for  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$ .*

*Proof.* It suffices to check that the  $j$ -th coordinate of  $\varphi_{(H, \lambda)}$  is equal to  $\Phi_j$  in Theorem 4.11. Let  $u = (u_0, \dots, u_n)$ , then

$$(H_{(\mathcal{T}, \mathcal{L})} u)^T = \left( \sum_{j \in J} a_{1j} u_j, \dots, \sum_{j \in J} a_{|I|j} u_j, - \sum_{j \in J} \left( \sum_{i \in I_1} a_{ij} \right) u_j, \dots, - \sum_{j \in J} \left( \sum_{i \in I_m} a_{ij} \right) u_j \right).$$



The  $j$ -th coordinate of  $\varphi_{(H(\mathcal{T}, \mathcal{L}), \lambda)}$  is

$$\begin{aligned}
\lambda_j(H(\mathcal{T}, \mathcal{L})u)^{h_j} &= (-1)^{\sum_{i \in I} a_{ij}} c_j \cdot \frac{(\sum_{j \in J} a_{1j} u_j)^{a_{1j}} \cdots (\sum_{j \in J} a_{|I|j} u_j)^{a_{|I|j}}}{(-\sum_{j \in J} (\sum_{i \in I_1} a_{ij}) u_j)^{\sum_{i \in I_1} a_{ij}} \cdots (-\sum_{j \in J} (\sum_{i \in I_m} a_{ij}))^{\sum_{i \in I_m} a_{ij}}} \\
&= c_j \cdot \frac{(\sum_{j \in J} a_{1j} u_j)^{a_{1j}} \cdots (\sum_{j \in J} a_{|I|j} u_j)^{a_{|I|j}}}{(\sum_{j \in J} (\sum_{i \in I_1} a_{ij}) u_j)^{\sum_{i \in I_1} a_{ij}} \cdots (\sum_{j \in J} (\sum_{i \in I_m} a_{ij}))^{\sum_{i \in I_m} a_{ij}}} \\
&= c_j \cdot \left( \frac{\sum_{j \in J} a_{1j} u_j}{\sum_{j \in J} (\sum_{i \in I_1} a_{ij}) u_j} \right)^{a_{1j}} \cdots \left( \frac{\sum_{j \in J} a_{|I|j} u_j}{\sum_{j \in J} (\sum_{i \in I_m} a_{ij}) u_j} \right)^{a_{|I|j}} = c_j \prod_{i \in I} \hat{\theta}_i^{a_{ij}} = \Phi_j.
\end{aligned}$$

□

## 5 Polytopes arising from toric multinomial staged trees

The aim of this section is to bring together the examples of 2D and 3D polytopes with rational linear precision (Section 3) and multinomial staged trees (Section 4). To this end, we investigate certain properties of the lattice polytopes arising from toric multinomial staged trees. This leads to an understanding for the negative part of the Horn matrix from a wider perspective than the one provided by using primitive collections. Recall that  $J$  denotes the set of root-to-leaf paths in  $\mathcal{T}$ . For  $j \in J$ ,  $p_j$  is defined to be the product of all edge labels in the path  $j$ . We denote the stages of  $(\mathcal{T}, \mathcal{L})$  by  $S_1, \dots, S_m$ .

**Definition 5.1.** The lattice polytope  $P_{\mathcal{T}}$  of a balanced multinomial staged tree  $(\mathcal{T}, \mathcal{L})$  is the convex hull of exponent vectors  $a_j$  of  $p_j$  for every root-to-leaf path  $j$  in  $\mathcal{T}$ .

Note that  $P_{\mathcal{T}} \subset \mathbb{R}^d$  is not a full-dimensional polytope for  $d = |S_1| + \dots + |S_m|$ . This can be observed e.g. in Figure 3 (left) for  $P_{\mathcal{T}_{b\Delta_2}} \cong b\Delta_2$  (unimodularly equivalent). We call  $(\mathcal{T}, \mathcal{L})$  a *multinomial staged tree representation* of a full-dimensional polytope  $P \cong P_{\mathcal{T}}$ .

### 5.1 Two dimensional multinomial staged tree models

The polytopes in 2D from Section 3.1 admit a multinomial staged tree representation.

**Proposition 5.1.** *All statistical models associated to pairs  $(P, w)$  in 2D with rational linear precision are toric multinomial staged tree models. The multinomial staged tree representations for each family in 2D are described in Figure 3.*

*Proof.* For the model  $b\Delta_2 = T_{0,b,1}$ , it suffices to note that the polytope  $b\Delta_n$  with weights given by multinomial coefficients has a Horn pair given by Theorem 2.4 which is equal to that one described in [9, Example 20] for multinomial models with  $b$  trials and  $n+1$  outcomes. The statistical model for  $T_{a,b,d}$  is the binary multinomial staged tree  $\mathcal{M}_{a,b,d}$  in Example 4.2, denoted by  $\mathcal{T}_{a,b,d}$  in Figure 3. The Horn matrix in Proposition 3.1, associated to the model for  $T_{a,b,d}$ , is equal to the Horn matrix of the model  $\mathcal{M}_{a,b,d}$ . Firstly, in both cases the columns are indexed by pairs  $(i, j)$  such that  $0 \leq j \leq b, 0 \leq i \leq a + d(b - j)$  so these matrices have the same number of columns. Using Corollary 4.12, we see that the column corresponding to

the outcome  $(i, j)$  in  $\mathcal{M}_{a,b,d}$  is  $(i, j, a + d(b - j) - i, b - j, -(a + d(b - j)), -b)$ , which equals the column associated to the lattice point  $(i, j)$  in Proposition 3.1. Uniqueness of the minimal Horn matrix, implies that the model associated to  $T_{a,b,d}$  is  $\mathcal{M}_{a,b,d}$ .

It remains to show that  $\mathcal{T}_{b\Delta_2}$  and  $\mathcal{T}_{a,b,d}$  are balanced. By Remark 4.8,  $\mathcal{T}_{b\Delta_2}$  is balanced because all root-to-leaf paths have length 1. For  $\mathcal{T}_{a,b,d}$ , it suffices to prove that the root  $r$  is balanced. Following the notation in Definition 4.6, let  $K^1 = (j_1, b - j_1)$ ,  $K^2 = (j_2, b - j_2)$ ,  $K^3 = (j_3, b - j_3)$ , and  $K_4 = (j_4, b - j_4)$  be such that  $K^1 + K^2 = K^3 + K^4$ . Then

$$\begin{aligned} t(r(K^1))t(r(K^2)) &= (s_2 + s_3)^{a+d(b-j_1)}(s_2 + s_3)^{a+d(b-j_2)} = (s_2 + s_3)^{2a+2db-d(j_1+j_2)} \\ &= (s_2 + s_3)^{2a+2db-d(j_3+j_4)} = (s_2 + s_3)^{a+d(b-j_3)}(s_2 + s_3)^{a+d(b-j_4)} \\ &= t(r(K^3))t(r(K^4)) \end{aligned}$$

□

Note that we obtain  $P_{\mathcal{T}_{0,b,1}} \cong b\Delta_2$  i.e. we have two different tree representations of  $b\Delta_2$ :  $\mathcal{T}_{b\Delta_2}$  and  $\mathcal{T}_{0,b,1}$ . For the investigation of the shape of a Horn matrix, we will be interested in those trees  $\mathcal{T}$  where the positive part of the Horn matrix  $H_{(\mathcal{T}, \mathcal{L})}$  from Corollary 4.12 is the lattice distance matrix of  $P_{\mathcal{T}}$  (Definition 5.3)(1). For simple polytopes  $P_{\mathcal{T}}$ , these trees with an additional property provide us an explanation for the negative part of  $H_{(\mathcal{T}, \mathcal{L})}$  in terms of primitive collections in Theorem 5.5.

## 5.2 Three dimensional binary multinomial staged tree models

Before we examine the multinomial staged tree representations more generally, we present the multinomial staged trees for the family of prisms in Table 1.

**Proposition 5.2.** *All statistical models associated to pairs in  $\mathcal{P}$  are toric binary multinomial staged trees.*

*Proof.* We first show that the Horn matrix of the statistical model associated to a general element in  $\mathcal{P}$  is equal to the Horn matrix of a binary multinomial staged tree. The general element in  $\mathcal{P}$  is a frustum with parameters  $a, a', b, b', d, l > 0$ . Let  $S = \{\{s_0, s_1\}, \{s_2, s_3\}, \{s_4, s_5\}\}$  be a set of symbols. We define the labelled tree  $(\mathcal{T}, \mathcal{L})$  by specifying its set of leaves, its set of root-to-leaf paths, and the labelling for the edges in each path. The set of leaves in  $\mathcal{T}$  is

$$J = \{(i, j, k) : 0 \leq k \leq l, \ 0 \leq j \leq bl - (b - b')k, \ 0 \leq i \leq (a + db)l - ((a + db) - (a' + db')k - dj)\}.$$

the labelled root-to-leaf path that ends at leaf  $(i, j, k)$  is

$$r \xrightarrow{\binom{l}{k} s_0^k s_1^{l-k}} v \xrightarrow{\binom{b-(b-b')k}{j} s_2^j s_3^{bl-(b-b')k-j}} w \xrightarrow{\binom{(a+db)l-((a+db)-(a+db')k-dj)}{i} s_4^i s_5^{(a+db)l-((a+db)-(a+db')k-dj)-i}} (i, j, k).$$

Thus  $p_{(i,j,k)}$  is the product of the labels in this path. A picture of this tree when  $l = 1$  is contained in Figure 3 as  $\mathcal{T}_{(A)_1}$ . Using Corollary 4.12, applied to  $(\mathcal{T}, \mathcal{L})$  just defined, we see that the column corresponding to  $(i, j, k)$  in the Horn matrix for  $\mathcal{M}_{(\mathcal{T}, \mathcal{L})}$  is equal to the column of the matrix in Proposition 3.3 evaluated at  $(i, j, k)$ .

It remains to prove that  $(\mathcal{T}, \mathcal{L})$  is balanced. Let us first prove that the root  $r$  of  $(\mathcal{T}, \mathcal{L})$




Figure 3: Multinomial staged trees for pairs in 2D and 3D (Table 1(A)) with rational linear precision.

is balanced. The exponents of the outgoing edges of  $r$  can be written as pairs of natural numbers that sum to the degree  $\ell$  of the floret. Thus they are pairs of the form  $(k, \ell - k)$ . Let us consider four such pairs, denoted by  $Q_1 := (k_1, \ell - k_1)$ ,  $Q_2 := (k_2, \ell - k_2)$ ,  $Q_3 := (k_3, \ell - k_3)$ ,  $Q_4 := (k_4, \ell - k_4)$ . Suppose that  $Q_1 + Q_2 = Q_3 + Q_4$ . Then we have  $k_1 + k_2 = k_3 + k_4$ .

We further need to check the following equality:

$$t(r(Q_1))t(r(Q_2)) = t(r(Q_3))t(r(Q_4)).$$

We have:

$$\begin{aligned} t(r(Q_1)) &= \sum_{j=0}^{bl-(b-b')k_1} \left[ \binom{bl-(b-b')k_1}{j} s_2^j s_3^{bl-(b-b')k_1-j} t(v(j, bl-(b-b')k_1-j)) \right] \\ &= \sum_{j=0}^{bl-(b-b')k_1} \left[ \binom{bl-(b-b')k_1}{j} s_2^j s_3^{bl-(b-b')k_1-j} (s_4 + s_5)^{(a+db)\ell - ((a+db)-(a'+db'))k_1 - dj} \right] \\ &= (s_4 + s_5)^{(a+db)\ell - ((a+db)-(a'+db'))k_1} \sum_{j=0}^{bl-(b-b')k_1} \left[ \binom{bl-(b-b')k_1}{j} \left( \frac{s_2}{(s_4 + s_5)^d} \right)^j s_3^{bl-(b-b')k_1-j} \right] \\ &= (s_4 + s_5)^{(a+db)\ell - ((a+db)-(a'+db'))k_1} \left( \frac{s_2}{(s_4 + s_5)^d} + s_3 \right)^{bl-(b-b')k_1}. \end{aligned}$$

We obtain similar formulae for  $t(r(Q_2))$ ,  $t(r(Q_3))$  and  $t(r(Q_4))$ . It follows that

$$\begin{aligned} t(r(Q_1))t(r(Q_2)) &= (s_4 + s_5)^{2(a+db)\ell - ((a+db)-(a'+db'))(k_1+k_2)} \left( \frac{s_2}{(s_4 + s_5)^d} + s_3 \right)^{2bl-(b-b')(k_1+k_2)} \\ &= (s_4 + s_5)^{2(a+db)\ell - ((a+db)-(a'+db'))(k_3+k_4)} \left( \frac{s_2}{(s_4 + s_5)^d} + s_3 \right)^{2bl-(b-b')(k_3+k_4)} \\ &= t(r(Q_3))t(r(Q_4)). \end{aligned}$$

A similar reasoning can be applied in to prove that the children of the root are balanced vertices. Next, let us denote by  $v$  such a vertex, whose parent is  $r$ . By Remark 4.8, any child of  $v$  is trivially balanced. Finally, we prove that all pairs of vertices in the same stage are balanced. There are three stages  $S_1 = \{s_0, s_1\}$ ,  $S_2 = \{s_2, s_3\}$  and  $S_3 = \{s_4, s_5\}$ . Denote by  $v$  and  $v'$  two children of the root  $r$ . The exponents of the outgoing edges of  $v$  can be written as pairs of natural numbers that sum to the degree  $bl - (b - b')k_1$  of the floret. Thus they are pairs of the form  $(j, bl - (b - b')k_1 - j)$ . Let us consider two such pairs, denoted by  $Q_1 := (j_1, bl - (b - b')k_1 - j_1)$ ,  $Q_2 := (j_2, bl - (b - b')k_1 - j_2)$ . Similarly, we consider two children of  $v'$  and we denote by  $Q_3 := (j_3, bl - (b - b')k_2 - j_3)$ ,  $Q_4 := (j_4, bl - (b - b')k_2 - j_4)$ . Suppose that  $Q_1 + Q_4 = Q_2 + Q_3$ . It follows that  $j_1 + j_4 = j_2 + j_3$ . We want to prove that

$$t(v(Q_1))t(v(Q_4)) = t(v(Q_2))t(v(Q_3)).$$

We obtain the following equality:

$$t(v(Q_1)) = t(v(j_1, bl - (b - b')k_1 - j_1)) = (s_4 + s_5)^{(a+db)\ell - ((a+db)-(a'+db'))k_1 - dj_1},$$

and its analogues for  $t(v(Q_2))$ ,  $t(v(Q_3))$  and  $t(v(Q_4))$ . Then

$$\begin{aligned} t(v(Q_1))t(v(Q_4)) &= (s_4 + s_5)^{2(a+db)\ell - ((a+db) - (a'+db'))(k_1+k_2) - d(j_1+j_4)} \\ &= (s_4 + s_5)^{2(a+db)\ell - ((a+db) - (a'+db'))(k_1+k_2) - d(j_2+j_3)} \\ &= t(v(Q_2))t(v(Q_3)). \end{aligned}$$

Therefore the pairs of vertices we considered are balanced.  $\square$

To obtain the multinomial staged tree representations for the models of the polytopes in Table 1, we use the tree in the proof of Proposition 5.2 and specialize the values of the parameters  $a, a', b, b', d, l$  accordingly. The trees for the family of prismatoids with trapezoidal base in Table 1 (A), with  $l = 1$ , are depicted in Figure 3. The trapezoidal frusta is represented by  $\mathcal{T}_{(A)_1}$ , the upper branch is the model for  $T_{a,b,d}$  and the lower branch is the the model for  $T_{a',b',d}$ . The other trees,  $\mathcal{T}_{(A)_2}$ ,  $\mathcal{T}_{(A)_3}$  and  $\mathcal{T}_{(A)_4}$ , have the same upper branch as  $\mathcal{T}_{(A)_1}$ . For the prismatoid with simplex on top, the substitution  $a' = 0$  has the effect of chopping a floret from  $\mathcal{T}_{(A)_1}$ , this gives  $\mathcal{T}_{(A)_2}$ . For the trapezoidal wedge,  $b' = 0$ , the edges in  $\mathcal{T}_{(A)_1}$  that contain  $b'$  contract to a single vertex, yielding  $\mathcal{T}_{(A)_3}$ . For the trapezoidal pyramid,  $a' = b' = 0$ , we chop off the lower part of the tree after the edge labelled by  $s_1$ . The trees for the remaining part of the Table 1, (B), (C) and (D) are obtained similarly.

### 5.3 Properties of the polytope $P_{\mathcal{T}}$

In this section we study certain properties of  $P_{\mathcal{T}}$  that can be formulated in terms of the combinatorics of its tree  $\mathcal{T}$ . We start by looking at root-to-leaf paths in  $\mathcal{T}$  that represent vertices of  $P_{\mathcal{T}}$ , this allows us to work with the normal fan  $\Sigma_{P_{\mathcal{T}}}$ . For simplicity we assume that  $(\mathcal{T}, \mathcal{L})$  has a root-to-leaf path of length  $m$  where  $S_1, \dots, S_m$  are the stages of  $\mathcal{T}$ .

**Definition 5.2.** A root-to-leaf path  $j$  is *vertex representing* if the exponent vector  $a_j$  of  $p_j$  is a vertex of  $P_{\mathcal{T}}$ .

**Lemma 5.3.** Let  $P_{\mathcal{T}} \subset \mathbb{R}^d$  be a polytope where  $(\mathcal{T}, \mathcal{L})$  is a multinomial staged tree from Proposition 5.1 or Proposition 5.2. Then the vertex representing paths in  $P_{\mathcal{T}}$  are those for which  $p_j$  is divisible by at most one symbol from each stage.

*Proof.* The upper and lower branches of the tree  $\mathcal{T}_{(A)_1}$  are the same up to a choice of parameters, thus we prove it only for the upper branch. Consider a root-to-leaf path  $\mathbf{j}$  in  $\mathcal{T}_{(A)_1}$  such that  $a_{\mathbf{j}} = (1, 0, j, b-j, k, a+d(b-j)-k)$  for  $0 < j < b$  and  $0 < k < a+d(b-j)$ . Let  $\mathbf{j}_1$  and  $\mathbf{j}_2$  be two root-to-leaf paths such that  $a_{\mathbf{j}_1} = (1, 0, j, b-j, a+d(b-j), 0)$  and  $a_{\mathbf{j}_2} = (1, 0, j, b-j, 0, a+d(b-j))$ . Then we obtain the equality  $a_{\mathbf{j}} = \frac{k}{a+d(b-j)}a_{\mathbf{j}_1} + \frac{a+d(b-j)-k}{a+d(b-j)}a_{\mathbf{j}_2}$  and hence  $a_{\mathbf{j}}$  cannot be a vertex of  $P_{\mathcal{T}_{(A)_1}}$ . It remains to show that  $a_{\mathbf{j}} = (1, 0, j, b-j, a+d(b-j), 0)$  is not a vertex. Let now  $\mathbf{j}'_1$  and  $\mathbf{j}'_2$  be two root-to-leaf paths such that  $p_{\mathbf{j}'_1}$  and  $p_{\mathbf{j}'_2}$  are divisible by one symbol from each stage and  $a_{\mathbf{j}'_1} = (1, 0, b, 0, a, 0)$ ,  $a_{\mathbf{j}'_2} = (1, 0, 0, b, a+db, 0)$ . Hence  $a_{\mathbf{j}} = \frac{j}{b}a_{\mathbf{j}'_1} + \frac{b-j}{b}a_{\mathbf{j}'_2}$  and  $a_{\mathbf{j}}$  is not a vertex of  $P_{\mathcal{T}_{(A)_1}}$ . The proofs for the remaining trees follow similarly.  $\square$

Next, we investigate the relation between primitive collections, Horn matrices, and stages of multinomial staged trees. The following definition was motivated by the observations on the trees from Section 5.1, 5.2 and by an attempt for answering Question 2.

**Definition 5.3.** Let  $(\mathcal{T}, \mathcal{L})$  be a balanced multinomial staged tree representation of  $P \cong P_{\mathcal{T}}$ . We say that the polytope  $P_{\mathcal{T}}$  has property  $(\star)$  if,

1. The positive part of the Horn matrix  $H_{(\mathcal{T}, \mathcal{L})}$  from Corollary 4.12 is the lattice distance matrix of  $P_{\mathcal{T}}$ .
2. The vertices of  $P_{\mathcal{T}}$  satisfy the conclusion of Lemma 5.3.

Note that isomorphic polytopes have the same lattice distance matrices.

**Lemma 5.4.** *Let  $P_{\mathcal{T}} \subset \mathbb{R}^d$  be a polytope with property  $(\star)$ . Then  $P_{\mathcal{T}}$  is simple if and only if all root-to-leaf paths have the same length  $m$  and  $|S_1| + \dots + |S_m| - m = \dim(P_{\mathcal{T}})$ .*

*Proof.* Recall that we assumed that there exists a root-to-leaf path of length  $m$ . All root-to-leaf paths have the same length  $m$  if and only if all vertex representing root-to-leaf paths have the same length  $m$ . Recall also that the vertices of  $P_{\mathcal{T}}$  are in one-to-one correspondence with the maximal cones of the normal fan  $\Sigma_{P_{\mathcal{T}}}$ . First suppose that  $P_{\mathcal{T}}$  is simple and there exists a vertex representing root-to-leaf path  $j'$  of length  $m' < m$ . Since the positive part of  $H_{(\mathcal{T}, \mathcal{L})}$  is the lattice distance matrix of  $P_{\mathcal{T}}$ , the symbols which do not divide  $p_{j'}$  represent the facets of  $P_{\mathcal{T}}$  which are lattice distance 0 to  $a_{j'}$ . Since  $P_{\mathcal{T}}$  satisfies the conclusion of Lemma 5.3, the maximal cone associated to the vertex  $a_{j'}$  in  $\Sigma_{P_{\mathcal{T}}}$  has more 1-face (ray) generators than the one associated to  $a_j$  where  $j$  has length  $m$ . Thus  $\Sigma_{P_{\mathcal{T}}}$  is not simplicial, contradiction. Moreover we obtain that the maximal cone associated to a vertex  $a_j$  is generated by the normal vectors associated to  $\cup_{l=1}^m S_l \setminus s_{i_l}$  for some  $s_{i_l} \in S_l$  and where  $\prod_{l=1}^m s_{i_l}$  divides  $p_j$ . This implies that  $\dim(P_{\mathcal{T}}) = |S_1| + \dots + |S_m| - m$ . Now suppose that all vertex representing root-to-leaf paths have the same length  $m$ . Then the number of symbols which do not divide  $p_j$  is  $|S_1| + \dots + |S_m| - m$  where  $j$  is a vertex representing root-to-leaf path. If this number is equal to  $\dim(P_{\mathcal{T}})$ , then  $P_{\mathcal{T}}$  is simple.  $\square$

Remark that the equality  $|S_1| + \dots + |S_m| - m = \dim(P_{\mathcal{T}})$  holds for all models from Proposition 5.1 and Proposition 5.2.

**Example 5.1.** The multinomial staged tree  $\mathcal{T}_{(A)_4}$  in Figure 3 for the trapezoidal pyramid, does not satisfy Definition 5.3(1). However when  $b = 1$ , we can find such a balanced multinomial staged tree representation for this polytope, it is shown in Figure 4 (left). This tree  $\mathcal{T}$  and  $\mathcal{T}_{(A)_4}$  represent the same model because their minimal Horn matrices are equal. When  $a = b = d = 1$ , the tree and its polytope are in Figure 4 (center) and (right). There are five vertex representing root-to-leaf paths namely 1, 3, 4, 5, and 6 and thus  $a_1, a_3, a_4, a_5$  and  $a_6$  are the vertices of the trapezoidal pyramid. In particular  $a_2 = \frac{1}{2}a_1 + \frac{1}{2}a_3$ . Hence  $P_{\mathcal{T}}$  has property  $(\star)$ . Moreover,  $P_{\mathcal{T}}$  is not simple by Lemma 5.4, since not all root-to-leaf paths have the length 2.




Figure 4: Multinomial staged tree representation of the non-simple trapezoidal pyramid,  $b = 1$ .

Even more, the minimal Horn matrix for this example with  $b = 1$  (below left) coincides with  $H_{(\mathcal{T}, \mathcal{L})}$ . As mentioned also in Section 3.3.1, the primitive collections  $\{n_1, n_3, n_4\}, \{n_2, n_3, n_5\}$  do not offer an explanation for the negative part of the minimal Horn matrix, however the stages  $\{s_0, s_1, s_2\}, \{s_3, s_4\}$  do.

|                      | $m_1$  | $m_2$               | $m_3$  | $m_4$       | $m_5$ | $m_6$ |  | $m_1$ | $m_2$ | $m_3$ | $m_4$ | $m_5$ | $m_6$ | $m_7$ | $m_8$ | $m_9$ | $m_{10}$ |
|----------------------|--|---------------------|--|-------------|-------|-------|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|----------|
| $s_0 = h_5$          | $\begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 2 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ -1 & -1 & -1 & -1 & -1 & -1 \\ -2 & -2 & -2 & -1 & -1 & 0 \end{bmatrix}$ | $s_1 = h_3$         | $\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 2 & 1 & 1 & 1 & 0 \\ 2 & 2 & 2 & 2 & 0 & 0 & 1 & 1 & 1 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & 1 & 2 & 3 & 0 & 1 & 0 & 1 & 2 & 0 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & 0 \\ -3 & -3 & -3 & -3 & -1 & -1 & -3 & -2 & -2 & 0 \end{bmatrix}$ | $s_2 = h_2$ |       |       |  |       |       |       |       |       |       |       |       |       |          |
| $s_1 = h_2$          |  | $s_3 = h_5$         |  |             |       |       |  |       |       |       |       |       |       |       |       |       |          |
| $s_2 = h_3$          |  | $s_4 = h_4$         |  |             |       |       |  |       |       |       |       |       |       |       |       |       |          |
| $s_3 = h_4$          |  | $s_5 = h_1$         |  |             |       |       |  |       |       |       |       |       |       |       |       |       |          |
| $s_4 = h_1$          |  | $-(s_0 + s_1)$      |  |             |       |       |  |       |       |       |       |       |       |       |       |       |          |
| $-(s_0 + s_1 + s_2)$ |  | $s_0 - (s_2 + s_3)$ |  |             |       |       |  |       |       |       |       |       |       |       |       |       |          |
| $-(s_3 + s_4)$       | $(s_4 + s_5)$  |                     |  |             |       |       |  |       |       |       |       |       |       |       |       |       |          |

On the other hand we observe that there exists no multinomial staged tree representation for  $b = 2$ ,  $a = d = 1$  fitting Definition 5.3 by looking at the lattice distance matrix seen in the positive part of its minimal Horn matrix (above right). This matrix can also be obtained by applying Corollary 4.12 to  $\mathcal{T}_{(A)_4}$  and performing the row operations explained in [9, Lemma 3] eliminating the row  $s_0 = h_6$ . This demonstrates how multinomial staged trees provide a wider understanding for the negative part of the Horn matrix.

For simple polytopes  $P_{\mathcal{T}}$  with property  $(\star)$  we show that the stages coincide with the primitive collections of  $\Sigma_{P_{\mathcal{T}}}$ .

**Theorem 5.5.** *Let  $P_{\mathcal{T}} \subset \mathbb{R}^d$  be a simple polytope with property  $(\star)$ . Then the primitive collections of the simplicial normal fan  $\Sigma_{P_{\mathcal{T}}}$  are represented by the stages  $S_1, \dots, S_m$ .*

*Proof.* By Definition 5.3(1), the symbols of the stages represent the facets of  $P_{\mathcal{T}}$ . Let now  $j$  be a vertex representing root-to-leaf path. Recall by the proof of Lemma 5.4, the maximal cone associated to  $a_j$  is generated by the normal vectors (1-faces) associated to  $\bigcup_{l=1}^m S_l \setminus s_{i_l}$  for some  $s_{i_l} \in S_l$  and where  $\prod_{l=1}^m s_{i_l}$  divides  $p_j$ . Since any intersection of two cones in  $\Sigma_{P_{\mathcal{T}}}$  is also a cone in  $\Sigma_{P_{\mathcal{T}}}$ , we obtain that  $\bigcup_{l=1}^m S_l \setminus S'_l$  for all  $S'_l \subseteq S_l$  with  $|S'_l| \geq 1$  is a cone of  $\Sigma_{P_{\mathcal{T}}}$ .

By Definition 2.6, since  $\Sigma_{P_{\mathcal{T}}}$  is simplicial, a primitive collection is a set of 1-faces which does not generate a cone itself but any proper subset does. This concludes that the partition  $S_1, \dots, S_m$  are the primitive collections of  $\Sigma_{P_{\mathcal{T}}}$ .  $\square$

The following corollary gives an affirmative answer to Question 2.

**Corollary 5.6.** *Let  $P_{\mathcal{T}} \subset \mathbb{R}^d$  be a simple polytope with property  $(\star)$ . Then we have  $H_{(\mathcal{T}, \mathcal{L})} = M_{\mathcal{A}, \Sigma_{P_{\mathcal{T}}}}$ , i.e. the negative rows are given by the primitive collections of  $\Sigma_{P_{\mathcal{T}}}$ .*

*Proof.* It follows from Corollary 4.12 and Theorem 5.5.  $\square$

**Example 5.2.** The multinomial staged trees  $\mathcal{T}_{b\Delta_2}$  and  $\mathcal{T}_{a,b,d}$  satisfy Definition 5.3(1) for the simplex and trapezoid ( $a, b, d > 0$ ) respectively. That means the facets of the polytopes are in one-to-one correspondence with the symbols in the stages. Moreover  $P_{\mathcal{T}_{b\Delta_2}}$  and  $P_{\mathcal{T}_{a,b,d}}$  are simple polytopes. Hence by Theorem 5.5 we obtain that the primitive collections are given by the partition of the stages. For the simplex  $P_{\mathcal{T}_{b\Delta_2}} \cong a\Delta_2$  we have only one primitive collection  $\{s_0, s_1, s_2\}$ . Similarly, for  $P_{\mathcal{T}_{a,b,d}} \cong T_{a,b,d}$  we have the partition of the stages as  $\{s_0, s_1\}$  and  $\{s_2, s_3\}$ , which correspond exactly to the primitive collections obtained in Theorem 3.2.




Figure 5: Multinomial staged tree representation of the simple trapezoidal wedge with  $b = 1$  considered in Example 5.3.

**Example 5.3.** Let us consider the balanced multinomial staged tree representation, satisfying Definition 5.3, of the trapezoidal wedge from Table 1 (A) with  $b = 1$  seen in Figure 5. This tree representation encodes the same model as the tree  $\mathcal{T}_{(A)_3}$  in Figure 3, because they have the same minimal Horn matrix. In particular we observe by Lemma 5.4 that  $P_{\mathcal{T}}$  is simple. By Theorem 5.5, the primitive collections are represented by the partition of the stages:  $\{n_2, n_3, n_5\} = \{s_0, s_1, s_2\}$  and  $\{n_1, n_4\} = \{s_3, s_4\}$ . By Corollary 5.6 the negative part of the minimal Horn matrix  $a = b = d = a' = 1$  (left), is explained by primitive collections. In Section 3.3.2 we saw that even for simple polytopes, the negative part of the Horn matrix is not always explained by the primitive collections. From the perspective of staged trees, for  $a' = a = d = 1, b = 2$ , this polytope can be represented by  $\mathcal{T}_{(A)_3}$ . After minimising the Horn matrix  $H_{(\mathcal{T}_{(A)_3}, \mathcal{L}_{(A)_3})}$ , constructed by Corollary 4.12, we obtain the matrix (right) whose positive part is the lattice distance matrix of  $P_{\mathcal{T}_{(A)_3}}$  (see Table 2). However a simple



computation shows that there exists no tree  $(\mathcal{T}, \mathcal{L})$  such that the positive part of  $H_{(\mathcal{T}, \mathcal{L})}$  is the lattice distance matrix of  $P_{\mathcal{T}_{(A)_3}}$  i.e. satisfying Definition 5.3(1).

$$\begin{array}{l}
 h_5=s_0 \\
 h_2=s_1 \\
 h_3=s_2 \\
 h_4=s_3 \\
 h_1=s_4 \\
 -(s_0+s_1+s_2) \\
 -(s_3+s_4)
 \end{array}
 \begin{bmatrix}
 m_1 & m_2 & m_3 & m_4 & m_5 & m_6 & m_7 \\
 1 & 1 & 0 & 0 & 0 & 0 & 1 \\
 0 & 0 & 1 & 1 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 1 & 1 & 0 \\
 2 & 0 & 1 & 0 & 1 & 0 & 1 \\
 0 & 2 & 0 & 1 & 0 & 1 & 1 \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 -2 & -2 & -1 & -1 & -1 & -1 & -2
 \end{bmatrix}
 \begin{array}{l}
 h_3=s_1 \\
 h_5=s_2 \\
 h_2=s_3 \\
 h_4=s_4 \\
 h_1=s_5 \\
 -s_0-s_1 \\
 s_0-s_2-s_3 \\
 -s_4-s_5
 \end{array}
 \begin{bmatrix}
 m_1 & m_2 & m_3 & m_4 & m_5 & m_6 & m_7 & m_8 & m_9 & m_{10} & m_{11} \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\
 2 & 2 & 2 & 2 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 0 & 0 \\
 3 & 2 & 1 & 0 & 2 & 1 & 0 & 1 & 0 & 1 & 0 \\
 0 & 1 & 2 & 3 & 0 & 1 & 2 & 0 & 1 & 0 & 1 \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & 0 & 0 \\
 -3 & -3 & -3 & -3 & -2 & -2 & -2 & -1 & -1 & -1 & -1
 \end{bmatrix}$$

## Acknowledgments

Computations using the software system Sagemath [17] and Macaulay2 [12] were crucial for the development of this paper. Eliana Duarte was supported by the Deutsche Forschungsgemeinschaft DFG under grant 314838170, GRK 2297 MathCoRe. We thank Bernd Sturmfels for introducing us to each other and for encouraging us to work on this project.

## References

- [1] Carlos Améndola, Nathan Bliss, Isaac Burke, Courtney R. Gibbons, Martin Helmer, Serkan Hoşten, Evan D. Nash, Jose Israel Rodriguez, and Daniel Smolkin, *The maximum likelihood degree of toric varieties*, J. Symbolic Comput. **92** (2019), 222–242. MR 3907355
- [2] Lamprini Ananiadi and Eliana Duarte, *Gröbner bases for staged trees*, Algebr. Stat. **12** (2021), no. 1, 1–20. MR 4251251
- [3] Victor V. Batyrev, *On the classification of smooth projective toric varieties*, Tohoku Math. J. (2) **43** (1991), no. 4, 569–585. MR 1133869
- [4] Patrick Clarke and David A. Cox, *Moment maps, strict linear precision, and maximum likelihood degree one*, Adv. Math. **370** (2020), 107233, 51. MR 4103774
- [5] David A. Cox, *Applications of polynomial systems*, vol. 134, American Mathematical Soc., 2020.
- [6] David A. Cox, John Little, and Donal O’Shea, *Ideals, varieties, and algorithms*, fourth ed., Undergraduate Texts in Mathematics, Springer, Cham, 2015, An introduction to computational algebraic geometry and commutative algebra. MR 3330490
- [7] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR 2810322

- [8] David A. Cox and Christine von Renesse, *Primitive collections and toric varieties*, Tohoku Math. J. (2) **61** (2009), no. 3, 309–332. MR 2568257
- [9] Eliana Duarte, Orlando Marigliano, and Bernd Sturmfels, *Discrete statistical models with rational maximum likelihood estimator*, Bernoulli **27** (2021), no. 1, 135–154. MR 4177364
- [10] Luis David Garcia-Puente and Frank Sottile, *Linear precision for parametric patches*, Adv. Comput. Math. **33** (2010), no. 2, 191–214. MR 2659586
- [11] Hans-Christian Graf von Bothmer, Kristian Ranestad, and Frank Sottile, *Linear precision for toric surface patches*, Found. Comput. Math. **10** (2010), no. 1, 37–66. MR 2591838
- [12] Daniel R. Grayson and Michael E. Stillman, *Macaulay2, a software system for research in algebraic geometry*, Available at <https://math.uiuc.edu/Macaulay2/>.
- [13] June Huh, *Varieties with maximum likelihood degree one*, J. Algebr. Stat. **5** (2014), no. 1, 1–17. MR 3279951
- [14] Rimvydas Krasauskas, *Toric surface patches*, Advances in Computational Mathematics **17** (2002), no. 1, 89–113. MR 1902537
- [15] Jim Q. Smith and Paul E. Anderson, *Conditional independence and chain event graphs*, Artificial Intelligence **172** (2008), no. 1, 42–68. MR 2388535
- [16] Seth Sullivant, *Algebraic statistics*, Graduate Studies in Mathematics, vol. 194, American Mathematical Society, Providence, RI, 2018. MR 3838364
- [17] The Sage Developers, *Sagemath, the Sage Mathematics Software System (Version 9.1)*, 2020, <https://www.sagemath.org>.

**Authors' addresses:**

Isobel Davies

Otto-von-Guericke Universität Magdeburg, Universitätsplatz 2, 39106 Magdeburg, Germany.

[isobel.davies@ovgu.de](mailto:isobel.davies@ovgu.de)

Eliana Duarte

Otto-von-Guericke Universität Magdeburg, Universitätsplatz 2, 39106 Magdeburg, Germany.

[eliana.duarte@ovgu.de](mailto:eliana.duarte@ovgu.de)

Irem Portakal

Max-Planck-Institut für Mathematik in den Naturwissenschaften, Inselstraße 22, 04103, Leipzig, Germany.

[mail@irem-portakal.de](mailto:mail@irem-portakal.de)

Miruna-Ştefana Sorea, SISSA, Trieste, Italy and RCMA Lucian Blaga University Sibiu, Romania.

[msorea@sissa.it](mailto:msorea@sissa.it)

# A Appendix A

| Name of subfamily   | Columns of minimal Horn matrix   |
|---|--|
| <b>(A) Prisms with trapezoidal base</b> $a > 0, b > 0, d > 0, l > 0$                                      |  |
| Trapezoidal frusta  | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_1 + h_4), -(h_2 + h_5), -(h_3 + h_6))$   |
| Triangle top  | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_1 + h_4), -(h_2 + h_5), -(h_3 + h_6))$   |
| Trapezoidal wedges with $b \neq 1$  | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4), -(h_2 + h_5 - h_6), -(h_3 + h_6))$  |
| Trapezoidal wedges with $b = 1$   | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4), -(h_2 + h_3 + h_5))$                |
| Trapezoidal pyramids with $b \neq 1$  | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4), -(h_2 + h_5 - h_6), -(h_3 + h_6))$  |
| Trapezoidal pyramids with $b = 1$   | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4), -(h_2 + h_3 + h_5))$                |
| <b>(B) Prisms with tensor product base</b> $a > 0, b > 0, d = 0, l > 0$                                   |  |
| General tensor product frusta   | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_1 + h_4), -(h_2 + h_5), -(h_3 + h_6))$   |
| Tensor product frusta with $a' = a$   | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_2 + h_5), -(h_1 + h_3 + h_4 + h_6))$     |
| Tensor product frusta with $b' = b$   | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_1 + h_4), -(h_2 + h_3 + h_5 + h_6))$     |
| 3D Tensor Product   | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_1 + h_2 + h_3 + h_4 + h_5 + h_6))$       |
| Tensor product frusta with $(a', a) = \lambda(b', b)$ or $\mu(a', a) = (b', b)$ ( $\lambda, \mu \geq 1$ ) | $(h_1, h_2, h_3, h_4, h_5, h_6, -(h_1 + h_2 + h_4 + h_5), -(h_3 + h_6))$     |
| Tensor product wedges ( $a' = 0$ ) with $a \neq 1$  | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4 - h_6), -(h_2 + h_5), -(h_3 + h_6))$  |
| Tensor product wedges ( $a' = 0$ ) with $a = 1$   | $(h_1, h_2, h_3, h_4, h_5, -(h_2 + h_5), -(h_1 + h_3 + h_4))$                |
| Tensor product wedges ( $b' = 0$ ) with $b \neq 1$  | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4), -(h_2 + h_5 - h_6), -(h_3 + h_6))$  |
| Tensor product wedges ( $b' = 0$ ) with $b = 1$   | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_4), -(h_2 + h_3 + h_5))$                |
| Tensor product pyramids   | $(h_1, h_2, h_3, h_4, h_5, -(h_1 + h_2 + h_4 + h_5 - h_6), -(h_3 + h_6))$    |
| <b>(C) Prisms with triangular base</b> $a = 0, b > 0, d > 0, l > 0$                                       |  |
| Triangular frusta ( $b' \neq b$ ) with $d \neq 1$   | $(h_1, h_2, h_3, h_4, h_6, -(h_1 + h_4 - h_5), -(h_2 + h_5), -(h_3 + h_6))$  |
| Triangular prism ( $b' = b$ ) with $d \neq 1$   | $(h_1, h_2, h_3, h_4, h_6, -(h_1 + h_4 - h_5), -(h_2 + h_3 + h_5 + h_6))$    |
| Triangular frusta ( $b' \neq b$ ) with $d = 1$  | $(h_1, h_2, h_3, h_4, h_6, -(h_1 + h_2 + h_4), -(h_3 + h_6))$                |
| Triangular prism ( $b' = b$ ) with $d = 1$  | $(h_1, h_2, h_3, h_4, h_6, -(h_1 + h_2 + h_3 + h_4 + h_6))$                  |
| Triangular based pyramid with $b \neq 1$ and $d \neq 1$   | $(h_1, h_2, h_3, h_4, -(h_1 + h_4 - h_5), -(h_2 + h_5 - h_6), -(h_3 + h_6))$ |
| Triangular based pyramid with $b = 1$ and $d \neq 1$  | $(h_1, h_2, h_3, h_4, -(h_1 + h_4 - h_5), -(h_2 + h_3 + h_5))$               |
| Triangular based pyramid with $b \neq 1$ and $d = 1$  | $(h_1, h_2, h_3, h_4, -(h_1 + h_2 + h_4 - h_6), -(h_3 + h_6))$               |
| 3D simplex  | $(h_1, h_2, h_3, h_4, -(h_1 + h_2 + h_3 + h_4))$                             |

Table 2: Minimal Horn matrices for pairs in  $\mathcal{P}$ . The right column in the table contains the columns of the Horn matrix in terms of the lattice distance functions of each polytope.