

Extensions of characters in type D and the inductive McKay condition, I

Britta Späth*

September 20, 2021

Abstract

This is a contribution to the study of $\text{Irr}(G)$ as an $\text{Aut}(G)$ -set for G a finite quasi-simple group. Focusing on the last open case of groups of Lie type D and ${}^2\text{D}$, a crucial property is the so-called condition $A'(\infty)$ expressing that diagonal automorphisms and graph-field automorphisms of G have transversal orbits in $\text{Irr}(G)$. This is part of the stronger $A(\infty)$ condition introduced in the context of the reduction of the McKay conjecture to a question on quasi-simple groups. Our main theorem is that a minimal counter-example to condition $A(\infty)$ for groups of type D would still satisfy $A'(\infty)$. This will be used in a second paper to fully establish $A(\infty)$ for any type and rank. The present paper uses Harish-Chandra induction as a parametrization tool. We give a new, more effective proof of the theorem of Geck and Lusztig ensuring that cuspidal characters of arbitrary standard Levi subgroups of $G = \text{D}_{l,\text{sc}}(q)$ extend to their stabilizers in the normalizer of that Levi subgroup. This allows to control the action of automorphisms on these extensions. From there Harish-Chandra theory leads naturally to a detailed study of associated relative Weyl groups and other extendibility problems in that context.

Contents

1	Basic Considerations	5
1.A	Notation and Condition $A(\infty)$	5
1.B	Action of $\text{Aut}(G)$ on Harish-Chandra induced characters	6
1.C	Action on characters of normalizers of Levi subgroups	9
1.D	Reminder on cuspidal characters	10

*This material is partially based upon work supported by the National Science Foundation under Grant No. DMS-1440140 while the author was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Spring 2018 semester. Some research was conducted in the framework of the research training group *GRK 2240: Algebro-Geometric Methods in Algebra, Arithmetic and Topology*, funded by the DFG.

School of Mathematics and Natural Sciences University of Wuppertal, Gaußstr. 20, 42119 Wuppertal, Germany, e-mail: bspaeth@uni-wuppertal.de

Mathematics Subject Classification (2010): 20C20 (20C33 20C34)

2	The Levi subgroup and its normalizer	12
2.A	Subgroups of the Levi subgroup L	13
2.B	The structure of N/L	16
2.C	A supplement of L in \overline{N}	18
3	Extending cuspidal characters of Levi subgroups	20
3.A	The inclusion $H_1 \triangleleft \overline{V}_1$	20
3.B	The inclusion $K_d \triangleleft K_d \overline{V}_d$ for $d \geq 2$	21
3.C	Consideration on $K_{-1} \triangleleft K_{-1} \overline{V}_{-1}$	24
3.D	Proof of Theorem B	25
4	More on cuspidal characters	28
5	Character theory for the relative inertia groups $W(\lambda)$	31
5.A	Understanding $\text{Irr}_{\text{cusp}}(N)$ via characters of subgroups of W	33
5.B	Cuspidal characters of L_I	36
5.C	Clifford theory for $W(\hat{\lambda}) \triangleleft W(\lambda)$ in the case of $\tilde{L}\tilde{L}_\lambda = \tilde{L}$	40
5.D	Clifford theory for $W(\hat{\lambda}) \triangleleft W(\lambda)$ in the case of $\tilde{L}_\lambda = \tilde{L}$	42
6	Proof of Theorem A	47
	References	51
	Index	53

Introduction

After the classification of finite simple groups and with the knowledge of their representations having also greatly expanded in the last decades it seems overdue to determine for each quasi-simple group G the action of its outer-automorphism group $\text{Out}(G)$ on its set of irreducible (complex) characters $\text{Irr}(G)$. This is important in order to use our results on representations of simple groups to get theorems about arbitrary finite groups. A crucial example is the McKay conjecture asserting

$$|\text{Irr}_{p'}(X)| = |\text{Irr}_{p'}(N_X(P))|$$

for p a prime, X a finite group, P one of its Sylow p -subgroups and $\text{Irr}_{p'}(X)$ the set of irreducible characters of degree prime to p . It is fairly clear that once this is solved for a normal subgroup Y of X the next step to deduce something for X is to determine the action of X on at least $\text{Irr}_{p'}(Y)$. The McKay conjecture has been reduced to a so-called inductive McKay condition about quasi-simple groups by Isaacs-Malle-Navarro [IMN07] and the first requirement is an $\text{Out}(X)$ -equivariant bijection realizing McKay's equality. Determining the action of $\text{Out}(G)$ on $\text{Irr}(G)$ for all quasi-simple groups G would also have applications to other conjectures about characters with similar reductions such as the Alperin-McKay conjecture or the Dade conjecture (see [S13], [S17]) or even conjectures about modular characters (see [NT11]) through the unitriangularity of decomposition matrices (see [BDT20]).

For alternating and sporadic groups the action of $\text{Out}(G)$ on $\text{Irr}(G)$ is easy to deduce from the available description of $\text{Irr}(G)$. When G is the universal covering group of a finite simple group of Lie type, this is Question 5.9 in [GM]. Previous research on the subject has left open only the case of groups of type D , see [CS19, 2.5]. The present paper is the

first part of a solution to that problem. A second part [S21a] will finish the determination. The splitting is due to the quite different methods used here and in [S21a]. A third part will focus on applications to the McKay conjecture [S21b].

In order to be more specific about intermediary goals and results, let us introduce some notation. Let $G = \mathbf{G}^F$ for $F: \mathbf{G} \rightarrow \mathbf{G}$ a Frobenius endomorphism of a simply-connected simple algebraic group \mathbf{G} . Upon choosing an F -stable maximal torus and a Borel subgroup containing it, one can define a group E of so-called field and graph automorphisms of G . One can also define a reductive group $\tilde{\mathbf{G}}$ realizing a regular embedding for \mathbf{G} , that is $\mathbf{G} = [\tilde{\mathbf{G}}, \tilde{\mathbf{G}}]$ with connected $Z(\tilde{\mathbf{G}})$. We can also assume that F extends to $\tilde{\mathbf{G}}$ into a Frobenius endomorphism and that E also acts on $\tilde{G} := \tilde{\mathbf{G}}^F$. Then $\text{Aut}(G)$ is induced by the semi-direct product $\tilde{G} \rtimes E$ (see for instance [GLS, 2.5.12]).

The determination of the action of $\tilde{G} \rtimes E$ on $\text{Irr}(G)$ essentially relies on establishing that \tilde{G} -orbits and E -orbits are somehow transversal. More precisely, one shows the following property

$A'(\infty)$: *There exists an E -stable \tilde{G} -transversal in $\text{Irr}(G)$.*

This, combined with the present knowledge of $\text{Irr}(\tilde{G})$, is enough to determine $\text{Irr}(G)$ as an $\text{Out}(G)$ -set (see [CS19, 2.5]). However, in order to deduce any valuable statement about representation of almost-simple groups, it is also important to answer extendibility questions. For instance a difficult theorem of Lusztig essentially focusing on the case of type D shows that any element of $\text{Irr}(G)$ extends to its stabilizer in \tilde{G} (see [L88], [L08]). This notably leads to the determination of the action of E on the set of \tilde{G} -orbits in $\text{Irr}(G)$.

The following strengthening of $A'(\infty)$ was introduced in [S12] in order to check the inductive McKay condition for the defining characteristic.

$A(\infty)$: *There exists an E -stable \tilde{G} -transversal \mathbb{T} in $\text{Irr}(G)$ and any $\chi \in \mathbb{T}$ extends as an irreducible character of its stabilizer $G \rtimes E_\chi$.*

The aim of the present paper and its sequel [S21a] is to prove $A(\infty)$ for G of type D and ${}^2\text{D}$. In the present paper G will be indeed some $\text{D}_{l,sc}(q)$ ($l \geq 4$, q a power of an odd prime); the case of twisted types ${}^2\text{D}$ will also be deduced in [S21a]. Condition $A(\infty)$ in the case of $\text{D}_{l,sc}(2^m)$ and ${}^2\text{D}_{l,sc}(2^m)$, essentially a corollary of [CS19, 3.9], is also established in [S21a].

Our main theorem here can be seen as showing that a counterexample to $A(\infty)$ with minimal l still satisfies $A'(\infty)$.

Theorem A *Let $G = \text{D}_{l,sc}(q)$ ($l \geq 4$, q a power of an odd prime), let \tilde{G} and E as above (see also 1.2). If any $\text{D}_{l',sc}(q)$ for $4 \leq l' < l$ satisfies $A(\infty)$, then G satisfies $A'(\infty)$.*

More precisely we assume Hypothesis 1.14, i.e. that condition $A(\infty)$ holds for the cuspidal characters of $G' = \text{D}_{l',sc}(q)$ with $4 \leq l' < l$.

Our proof uses as a starting point a theorem of Malle [M17] showing the existence part of the above statement for cuspidal characters. Our strategy thereof is through the parametrization of $\text{Irr}(G)$ given by Harish Chandra theory. In particular we take the standard Levi supplement \mathbf{L} of a parabolic subgroup \mathbf{P} containing our chosen Borel subgroup and consider parabolic induction $\text{R}_{\mathbf{L}}^{\mathbf{G}} \lambda$ of cuspidal characters $\lambda \in \text{Irr}_{\text{cusp}}(\mathbf{L}^F)$.

An essential ingredient of that parametrization is the deep theorem by Lusztig and Geck ([L, 8.6] and [G93, Cor. 2]) that any $\lambda \in \text{Irr}_{\text{cusp}}(\mathbf{L}^F)$ extends to its stabilizer in $N := \text{N}_{\mathbf{G}}(\mathbf{L})^F$. In order to put that parametrization to use for our purpose of tracking automorphism actions it is important to find an equivariant version of that statement. This does not seem possible from the available proofs, so we devise a new one in this paper, showing namely with the same notations for $G = \text{D}_{l,sc}(q)$, \tilde{G} , \mathbf{L} , N , E

Theorem B *Let $\lambda \in \text{Irr}_{\text{cusp}}(\mathbf{L}^F)$. Assume Hypothesis 1.14 holds for $D_{l',sc}(q)$ ($4 \leq l' < l$). Then some $(Z(\tilde{\mathbf{G}})\mathbf{L})^F$ -conjugate λ_0 of λ has an $(NN_E(\mathbf{L}))_{\lambda_0}$ -stable extension to N_{λ_0} .*

Studying the group structure of N , our proof uses essentially the Steinberg relations for the structure of G , not its realization as spin group, making probably more uniform a case-by-case but effective proof of Geck-Lusztig's theorem for other quasi-simple groups of Lie type (see [BS20, 4.3] and [CSS21, 4.13] for types A and C).

We should point out that the above extendibility property is part of the following wider problem where (\mathbf{H}, F) is a reductive group defined over a finite field and F is its associated Frobenius endomorphism.

(P) *Let \mathbf{S} be an F -stable torus of \mathbf{G} . Does every $\psi \in \text{Irr}(\mathbf{C}_{\mathbf{H}}(\mathbf{S})^F)$ extend to its stabilizer in $\mathbf{N}_{\mathbf{H}}(\mathbf{S})^F$?*

This was solved in the affirmative in the case where \mathbf{S} is a Sylow ϕ -torus (ϕ any cyclotomic polynomial), see [S09, S10a, S10b]. Lusztig's theorem on the case where \mathbf{S} is split and ψ is cuspidal was essential to turn Deligne-Lusztig theory into a parametrization of $\text{Irr}(\mathbf{H}^F)$ when \mathbf{H} has connected center [L]. It seems that even partial answers to (P) have quite interesting applications (see also [B06, §15], [M14, 2.9]).

Let for now $\text{Irr}_{\text{cusp}}(N)$ be the set of characters of N whose restriction to L is a sum of cuspidal characters. Theorem B then can be seen as the starting point of a specific parametrization of $\text{Irr}_{\text{cusp}}(N)$ bearing similarities with the parametrizations of characters of normalizers of Sylow ϕ -tori given in the author's work just mentioned but with a special emphasis on outer automorphism actions.

Through preparations recalled at the start of the paper and similar to a method developed in [MS16] where \mathbf{L} was a torus, our main goal Theorem A reduces to a weak analogue of it for $\text{Irr}_{\text{cusp}}(N)$. This is Theorem 5.1. It is checked through a strategy prescribed by Clifford theory. In particular this entails a quite detailed analysis of the relative Weyl groups

$$W(\lambda) := N_{\lambda}/\mathbf{L}^F$$

and their various embeddings related to $\tilde{\mathbf{G}}$ and E .

Structure of the paper. In Section 1 we recall notation on quasi-simple groups of Lie type, their automorphisms and the conditions $A(\infty)$, $A'(\infty)$. Then we collect the basic facts about cuspidal characters and Harish Chandra theory for finite groups. This leads to Theorem 1.8 which sums up the methods from [MS16] to establish condition $A(\infty)$ through Harish Chandra theory. This is roughly the road map for the rest of the paper, in particular splitting the task into two halves that will be addressed in Sections 2-3 and Sections 5-6.

The rest of the paper is specific to type D (untwisted) in odd characteristic. After recalling a method from [CSS21] for constructing extensions, the main objective of Section 2-3 is Theorem B. Section 2 is a description of certain group theoretical aspects of the groups $L := \mathbf{L}^F$ and N , using also the classic embedding $\mathbf{G} \leq \overline{\mathbf{G}}$ of type D_l into type B_l . The root system Φ' of \mathbf{L} is the direct product of irreducible root systems of type A and type D. Roughly speaking, the factors of type A_{d-1} form a root system Φ_d and the factor of type D gives Φ_{-1} . Along the way we introduce a set \mathbb{D} determining the types occurring as factors of Φ' . This description will be used in the whole paper. For each $d \in \mathbb{D}$ we describe a normal inclusion $H_d \triangleleft V_d \leq \overline{N} := N_{\overline{\mathbf{G}}}(\mathbf{L})^F$ where $H_d = L \cap V_d$ is an elementary abelian 2-group and $L \langle V_d \mid d \in \mathbb{D} \rangle = \overline{N}$. This normal inclusion $H_d \triangleleft V_d$ concentrates the equivariant extendibility problem we have to solve.

In Section 3 we draw the consequences of the structure of N in terms of characters. One has to take care of all the factors involved and deal with the inclusion in type B

which provides the graph automorphism specific to type D. Concerning the diagonal automorphisms we avoid choosing a regular embedding \tilde{G} and instead consider inclusions $L \triangleleft \mathcal{L}^{-1}(Z) \cap \mathbf{L}$ where $Z \leq Z(\mathbf{G})$ and \mathcal{L} is the Lang map $x \mapsto x^{-1}F(x)$.

Theorem B being proved, we study in Section 4 how automorphisms act on cuspidal characters in types A and D, making use in the latter case of Malle's theorem [M17] mentioned above and some results about semi-simple characters already used in the study of the McKay conjecture for the defining characteristic (see [Ma10, §8]).

In Section 5, the most technical of the paper, the objective is to prove Theorem 5.1, showing that $\text{Irr}_{\text{cusp}}(N)$ satisfies a version of $A(\infty)$. As already shown in Section 1, this translates into requirements on $\text{Irr}(N_\lambda/L)$, the characters of the relative Weyl group $W(\lambda)$ associated to a cuspidal character λ of L . The comparison of the action of diagonal versus graph-field automorphisms on $\text{Irr}_{\text{cusp}}(N)$ relates with the induced action of related characters of relative Weyl groups. The proof splits naturally into the various cases for the stabilizer of λ in $L \cap \mathcal{L}^{-1}(Z(\mathbf{G}))/L$. This leads to Proposition 5.28 and Proposition 5.35 describing the situation in the two main cases. In the proofs graph-field automorphisms are taken care by embedding the relative Weyl group $W(\lambda)$ into overgroups $K(\lambda)$ and $\widehat{K}(\lambda)$ (see Notation 5.4) for field automorphisms and the embedding into type B for the graph automorphism of order 2.

In Section 6 we essentially put together all the material of the preceding section to establish Theorem 5.1 and with some extra effort Theorem A.

Acknowledgements. This material is partly based upon work supported by the NSF under Grant DMS-1440140 while the author was in residence at the MSRI, Berkeley CA. Some research was conducted in the framework of the research training group *GRK 2240: Algebro-Geometric Methods in Algebra, Arithmetic and Topology*, funded by the DFG. I thank Julian Brough and Lucas Ruhstorfer for their remarks on preliminary versions of the paper and Gunter Malle for reading an earlier version.

1 Basic Considerations

We first recall here some notation around characters, recall Condition $A(\infty)$ and give a rephrasing that provides alternative approaches for the proof of Theorem A. In 1.B we recall relevant results from Harish-Chandra theory. We conclude with general considerations on cuspidal characters in 1.C.

1.A Notation and Condition $A(\infty)$

In general we follow the notation about characters as introduced in [I]. Additionally we use some terminology from [S09, S10a, S10b] that is recalled in the following paragraph.

Notation 1.1 Let $X \triangleleft Y$ be finite groups and $\mathbb{T} \subseteq \text{Irr}(X)$. An *extension map with respect to $X \triangleleft Y$* for \mathbb{T} is a map $\Lambda : \mathbb{T} \rightarrow \coprod_{X \leq I \leq Y} \text{Irr}(I)$ such that every $\lambda \in \mathbb{T}$ is mapped to an extension of λ to Y_λ the inertia subgroup of λ in Y . We say that *maximal extendibility holds with respect to $X \triangleleft Y$* for \mathbb{T} if such an extension map exists, see also [CS17a, Def. 5.7]. In such a case the map can be chosen Y -equivariant, provided \mathbb{T} is Y -stable, see [CS17a, Thm. 4.1]. Whenever $\mathbb{T} = \text{Irr}(X)$ we omit to mention \mathbb{T} . For $\lambda \in \text{Irr}(X)$ and $\psi \in \text{Irr}(Y)$ we write λ^Y for the character induced to Y and $\psi|_X$ for the restricted character. For any generalized character κ we denote by $\text{Irr}(\kappa)$ the set of (irreducible) constituents of κ . If $\sigma \in \text{Aut}(X)$ and $\lambda \in \text{Irr}(X)$ we write $\lambda^\sigma = \sigma^{-1}\lambda$

for the character with $\sigma^{-1}\lambda(x) = \lambda^\sigma(x) = \lambda(\sigma(x))$ for $x \in X$.

If two subgroups $H_1, H_2 \leq Y$ satisfy $[H_1, H_2] = 1$, we write $H_1.H_2$ for $\langle H_1, H_2 \rangle \leq Y$. If $\lambda_i \in \text{Irr}(H_i)$ for $i \in \underline{2}$ and $\text{Irr}(\lambda_1|_{H_1 \cap H_2}) = \text{Irr}(\lambda_2|_{H_1 \cap H_2})$, there exists a unique character $\phi \in \text{Irr}(\langle H_1, H_2 \rangle)$ with $\text{Irr}(\phi|_{H_i}) = \{\lambda_i\}$ according to [IMN07, §5] and we write $\lambda_1 \cdot \lambda_2$ for this character. Let \mathbb{I} be a finite set, Z and H_i ($i \in \mathbb{I}$) finite groups with $Z \leq H_i$. If $[H_i, H_{i'}] = 1$ for every $i, i' \in \mathbb{I}$ with $i \neq i'$ and $H_i \cap \langle H_j \mid j \in \mathbb{I} \setminus \{i\} \rangle = Z$, we write $\otimes_{i \in \mathbb{I}} H_i$ for the central product of the groups H_i . Given $\nu \in \text{Irr}(Z)$ and $\lambda_i \in \text{Irr}(H_i \mid \nu)$ we denote by $\odot_{i \in \mathbb{I}} \lambda_i \in \text{Irr}(H_i)$ the character $\phi \in \text{Irr}(\otimes_{i \in \mathbb{I}} H_i)$ with $\text{Irr}(\phi|_{H_i}) = \{\lambda_i\}$ for every $i \in \mathbb{I}$, see also [IMN07, §5].

Next we introduce the groups and automorphisms considered in the following.

Notation 1.2 — Simple Groups of Lie type. Let \mathbf{G} be a simple linear algebraic group of simply-connected type over an algebraic closure \mathbb{F} of \mathbb{F}_p for p a prime. Additionally let $F : \mathbf{G} \rightarrow \mathbf{G}$ be a Frobenius endomorphism defining an \mathbb{F}_q -structure on \mathbf{G} . Let \mathbf{T}_0 be an F -stable maximally split torus and \mathbf{B} be an F -stable Borel subgroup of \mathbf{G} with $\mathbf{T}_0 \subseteq \mathbf{B}$ and $\mathbf{N}_0 := \mathbf{N}_{\mathbf{G}}(\mathbf{T}_0)$. According to [MT, Thm. 24.11] the group $G := \mathbf{G}^F$ has a split BN -pair with respect to $B := \mathbf{B}^F$, $T_0 := \mathbf{T}_0^F$ and $N_0 := \mathbf{N}_0^F$. Let $E(\mathbf{G}^F)$, often just E , be the subgroup of $\text{Aut}(\mathbf{G}^F)_{\mathbf{B}, \mathbf{T}_0}$ generated by graph automorphisms and some Frobenius endomorphism F_0 as in [GLS, Thm. 2.5.1] and [CS19, §2.A].

Let $\iota : \mathbf{G} \rightarrow \tilde{\mathbf{G}}$ be a regular embedding, i.e., a closed inclusion of algebraic groups with $\tilde{\mathbf{G}} = \mathbf{Z}(\tilde{\mathbf{G}})\mathbf{G}$ and connected $\mathbf{Z}(\tilde{\mathbf{G}})$. Then $\tilde{\mathbf{T}}_0 := \mathbf{Z}(\tilde{\mathbf{G}})\mathbf{T}_0$ is a maximal torus of $\tilde{\mathbf{G}}$. Let $\tilde{T}_0 := \tilde{\mathbf{T}}_0^F$. Assume that $F : \tilde{\mathbf{G}} \rightarrow \tilde{\mathbf{G}}$ is a Frobenius endomorphism extending the one of \mathbf{G} , see also [MS16, §2]. Then $\tilde{G} := \tilde{\mathbf{G}}^F$ has again a split BN -pair with respect to the groups $\tilde{B} := \tilde{T}_0 B$ and $\tilde{N}'_0 := \tilde{T}_0 N_0$, see [MT, Thm. 24.11]. Often the action of \tilde{N}'_0 on \mathbf{G} will be studied via the group $\tilde{N}_0 := \{x \in \mathbf{N}_{\mathbf{G}}(\mathbf{T}_0) \mid x^{-1}F(x) \in \mathbf{Z}(\mathbf{G})\}$, which will be shown to induce the same automorphisms on \mathbf{G} , see 1.16.

Via the convention given in [MS16, §2], $E(\mathbf{G}^F)$ also acts on \tilde{G} .

We recall the abbreviating notation from [CS19, Def. 2.2]:

Condition 1.3 (On stabilizers of irreducible characters of \mathbf{G}^F).

$A(\infty)$: There exists some E -stable $\tilde{\mathbf{G}}^F$ -transversal \mathbb{T} in $\text{Irr}(\mathbf{G}^F)$, such that every $\chi \in \mathbb{T}$ extends to $\mathbf{G}^F E_\chi$.

$A'(\infty)$: There exists some E -stable $\tilde{\mathbf{G}}^F$ -transversal \mathbb{T} in $\text{Irr}(\mathbf{G}^F)$.

Condition $A'(\infty)$ implies a weak version of Assumption 2.12(v) of [S12].

Lemma 1.4 Let \hat{Y} and \tilde{X} be two subgroups of a group Z with $\tilde{X} \triangleleft Z$ and $Z = \hat{Y}\tilde{X}$.

For $X := \tilde{X} \cap \hat{Y}$ let $\mathcal{M} \subseteq \text{Irr}(X)$ be Z -stable. Then the following are equivalent:

- (i) there is a \hat{Y} -stable \tilde{X} -transversal \mathcal{M}_0 in \mathcal{M} ;
- (ii) every $\zeta' \in \mathcal{M}$ is \tilde{X} -conjugate to some ζ such that $(\tilde{X}\hat{Y})_\zeta = \tilde{X}_\zeta \hat{Y}_\zeta$;
- (iii) every $\zeta' \in \mathcal{M}$ satisfies $(\tilde{X}\hat{Y})_{\zeta'} = (\hat{Y}^x)_\zeta \tilde{X}_\zeta$ for some $x \in \tilde{X}$.

Proof. This follows from Remark 3.3 of [CSS21]. □

1.B Action of $\text{Aut}(G)$ on Harish-Chandra induced characters

Using detailed analysis of Harish-Chandra induction, the results of [MS16] describe the action of $\text{Aut}(\mathbf{G}^F)$ in terms of cuspidal characters and their relative Weyl groups. The action is expressed in terms of the labels given by Howlett-Lehrer theory.

Notation 1.5 Let L be a standard Levi subgroup of G with respect to B and T_0 , i.e. $L = \mathbf{L}^F$ for some standard Levi subgroup \mathbf{L} of \mathbf{G} such that $\mathbf{T}_0 \leq \mathbf{L}$ and $\mathbf{L}\mathbf{B}$ is a parabolic subgroup. We set $N := N_{\mathbf{G}^F}(\mathbf{L})$, $W := N/L$ and $E_L = \text{Stab}_{E(\mathbf{G}^F)}(L)$. We write $\text{Irr}_{\text{cusp}}(L)$ for the set of cuspidal characters of L as defined in [C, 9.1] and $\text{Irr}_{\text{cusp}}(N) := \bigcup_{\lambda \in \text{Irr}_{\text{cusp}}(L)} \text{Irr}(\lambda^N)$. Additionally let $\text{Irr}(G \mid (L, \lambda)) := \text{Irr}(\mathbf{R}_L^G(\lambda))$, in other places also denoted as $\mathcal{E}(G, (L, \lambda))$ and $\text{Irr}(G \mid (L, \mathbb{T})) := \bigcup_{\lambda \in \mathbb{T}} \text{Irr}(G \mid (L, \lambda))$ for $\mathbb{T} \subseteq \text{Irr}_{\text{cusp}}(L)$, where \mathbf{R}_L^G denotes here the Harish-Chandra induction from L to G .

1.6. Let $\text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$ be the subgroup of $\text{Aut}(\mathbf{G}^F)$ generated by the automorphisms of \mathbf{G}^F induced by N and $\text{Aut}(\mathbf{G}^F)_{BL, L}$. Note $E_L \leq \text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$. According to Howlett-Lehrer theory (see [C, §10]), fixing an extension $\tilde{\lambda} \in \text{Irr}(N_\lambda)$ of $\lambda \in \text{Irr}_{\text{cusp}}(L)$ defines a unique labelling of $\text{Irr}(G \mid (L, \lambda))$ by $\text{Irr}(W(\lambda))$ where $W(\lambda) := N_\lambda/L$. We write $\mathbf{R}_L^G(\lambda)_\eta$ for the character of $\text{Irr}(G \mid (L, \lambda))$ associated to $\eta \in \text{Irr}(W(\lambda))$ via the extension $\tilde{\lambda}$.

Accordingly the parametrization of $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$ depends on an extension map Λ_L with respect to $L \triangleleft N$ for $\text{Irr}_{\text{cusp}}(L)$. For $\lambda \in \text{Irr}_{\text{cusp}}(L)$ let $R(\lambda) \triangleleft W(\lambda)$ be defined as in [C, Prop. 10.6.3]. If $\lambda \in \text{Irr}_{\text{cusp}}(L)$ and $\sigma \in \text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$ let $\delta_{\lambda, \sigma}$ be the unique linear character of $W(\sigma\lambda)$ satisfying

$${}^\sigma \Lambda_L(\lambda) = \Lambda_L(\sigma\lambda)\delta_{\lambda, \sigma}. \quad (1.1)$$

We only use the formula with some simplifying assumptions on $R(\lambda)$ and $\delta_{\lambda, \sigma}$.

Theorem 1.7 — Malle-Sp ath [MS16, Thm. 4.6 and 4.7]. *Let $\sigma \in \text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$ and Λ_L be an N -equivariant extension map with respect to $L \triangleleft N$ for $\text{Irr}_{\text{cusp}}(L)$. Assume that $\mathbf{R}_L^G(\lambda)_\eta$ ($\lambda \in \text{Irr}_{\text{cusp}}(L)$, $\eta \in \text{Irr}(W(\lambda))$) is defined using Λ_L and*

$$R(\sigma\lambda) \leq \ker(\delta_{\lambda, \sigma}) \quad \text{for every } \lambda \in \text{Irr}_{\text{cusp}}(L). \quad (1.2)$$

Then ${}^\sigma(\mathbf{R}_L^G(\lambda)_\eta) = \mathbf{R}_L^G(\sigma\lambda)_{\sigma\eta\delta_{\lambda, \sigma}^{-1}}$ for every $\lambda \in \mathbb{T}$ and $\eta \in \text{Irr}(W(\lambda))$.

In Section 5 of [MS16], the analogue of Theorem A was proven for characters in $\text{Irr}(G \mid (T_0, \text{Irr}_{\text{cusp}}(T_0)))$ by studying $\text{Irr}_{\text{cusp}}(N_{\mathbf{G}}(\mathbf{T}_0)^F)$. For other standard Levi subgroups the strategy from [MS16] leads naturally to the following statement where we focus on a single L and its stabilizer in E . Sections 2-5 will ensure the assumptions for the groups from 1.2 whenever $\mathbf{G}^F = D_{l, \text{sc}}(q)$.

Theorem 1.8 *Let $\tilde{L}' := \tilde{T}_0 L$, $\tilde{N}' := \tilde{T}_0 N$ and $\hat{N} := NE_L$. Assume there exist*

- (i) *an \hat{N} -stable \tilde{L}' -transversal \mathbb{T} in $\text{Irr}_{\text{cusp}}(L)$, an N -equivariant extension map $\Lambda_{L, \mathbb{T}}$ with respect to $L \triangleleft N$ for \mathbb{T} such that any $\lambda \in \mathbb{T}$ satisfies Equation (1.2); and*
- (ii) *some E_L -stable \tilde{N}' -transversal in $\text{Irr}_{\text{cusp}}(N)$.*

Then there exists an E_L -stable \tilde{G} -transversal in $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$.

For the proof of Theorem 1.8 we parametrize $\text{Irr}_{\text{cusp}}(N)$ via a set $\mathcal{P}(L)$ using an extension map Λ_L with respect to $L \triangleleft N$ for $\text{Irr}_{\text{cusp}}(L)$ deduced from $\Lambda_{L, \mathbb{T}}$.

Notation 1.9 By the assumption \mathbb{T} is an \hat{N} -stable \tilde{L}' -transversal in $\text{Irr}_{\text{cusp}}(L)$. For each $\lambda \in \mathbb{T}$ we denote by \mathcal{O}_λ its N -orbit in $\text{Irr}_{\text{cusp}}(L)$. Note $\mathcal{O}_\lambda \subseteq \text{Irr}_{\text{cusp}}(L)$. Let $M(\lambda) \subseteq \tilde{L}'$ be a set of representatives of the \tilde{L}'_λ -cosets in \tilde{L}' . We define an extension map Λ_L on \mathcal{O}_λ by

$$\Lambda_L(\lambda^m) = \Lambda_{L, \mathbb{T}}(\lambda')^m \quad \text{for every } \lambda' \in \mathcal{O}_\lambda \text{ and } m \in M(\lambda).$$

Hence, Λ_L is defined, but depends on the choice of $M(\lambda)$. The map $\Lambda'_L : \text{Irr}_{\text{cusp}}(L) \longrightarrow \prod_{L \leq I \leq N} \text{Irr}(I)$ with $\Lambda'_L(\mu) := \Lambda_L(\mu) \upharpoonright_{N_{\tilde{\mu}}}$ for every $\mu \in \text{Irr}_{\text{cusp}}(L)$ is well-defined, where $\tilde{\mu} \in \text{Irr}(\tilde{L}'_{\mu})$ is an extension of μ . In contrast to Λ_L we see that Λ'_L is independent of the choice of $M(\lambda)$. Observe $[N/L, \tilde{L}'/L] = 1$ via conjugation. The map Λ'_L is even $\widehat{N}\tilde{L}'$ -equivariant since Λ_L is N -equivariant and $\Lambda_{L, \mathbb{T}}$ is \widehat{N} -equivariant.

We write $\mathcal{P}^\circ(L)$ for the set of pairs (λ, η) with $\lambda \in \text{Irr}_{\text{cusp}}(L)$ and $\eta \in \text{Irr}(W(\lambda))$. The groups N and W act naturally via conjugation on $\mathcal{P}^\circ(L)$. We denote by $\mathcal{P}(L)$ the set of N -orbits in $\mathcal{P}^\circ(L)$ and by $\overline{(\lambda, \eta)}$ the N -orbit containing (λ, η) . Since L is mostly clear from the context, we omit it, writing \mathcal{P}° and \mathcal{P} .

The parametrization of $\text{Irr}_{\text{cusp}}(N)$ is given by the following.

Proposition 1.10 *Let Λ_L , \mathcal{P}° and \mathcal{P} be as in 1.8 and 1.9.*

(a) *Then the map*

$$\Upsilon : \mathcal{P} \longrightarrow \text{Irr}_{\text{cusp}}(N) \text{ with } \overline{(\lambda, \eta)} \longmapsto (\Lambda_L(\lambda)\eta)^N$$

is a well-defined bijection.

(b) *$\sigma \Upsilon(\overline{(\lambda, \eta)}) = \Upsilon(\overline{(\sigma\lambda, \sigma\eta\delta_{\lambda, \sigma})})$ for every $\sigma \in \text{Aut}(G)_{L, \text{HC}}$ and $\overline{(\lambda, \eta)} \in \mathcal{P}$, where $\delta_{\lambda, \sigma} \in \text{Irr}(W(\sigma\lambda))$ is given as in 1.6.*

Proof. Clifford theory together with Gallagher's Lemma [I, 6.17] proves part (a). The definition of $\delta_{\lambda, \sigma}$ in Equation (1.1) from 1.6 leads to part (b). \square

In combination with Theorem 1.7 we obtain a proof of Theorem 1.8.

Proof of Theorem 1.8. For the application of Theorem 1.7 we have to ensure that by the assumptions the Equation (1.2) holds for characters $\lambda \in \mathbb{T}$ and $\sigma \in \text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$. For every $\lambda \in \text{Irr}_{\text{cusp}}(L)$, the character $\Lambda_L(\lambda)$ is an extension of $\Lambda'_L(\lambda)$. Accordingly $\delta_{\lambda, \sigma}$ defined as the unique linear character of $W(\sigma\lambda)$ such that $\sigma\Lambda_L(\lambda) = \Lambda_L(\sigma\lambda)\delta_{\lambda, \sigma}$ satisfies as well $\sigma\Lambda'_L(\lambda) \upharpoonright_{N_{\tilde{\lambda}\sigma}} = \Lambda'_L(\sigma\lambda) \delta_{\lambda, \sigma} \upharpoonright_{N_{\tilde{\lambda}\sigma}}$. Since $\Lambda'_L(\lambda)$ is $\tilde{N}'E_L$ -equivariant, we see that $\delta_{\lambda, \sigma} \upharpoonright_{N_{\tilde{\lambda}\sigma}}$ is trivial. Accordingly $\ker(\delta_{\lambda, \sigma}) \geq N_{\sigma\tilde{\lambda}}/L$ for every $\lambda \in \text{Irr}_{\text{cusp}}(L)$ and $\sigma \in \text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$ where $\tilde{\lambda}$ denotes an extension of λ to \tilde{L}'_{λ} . Recall $W(\sigma\tilde{\lambda}) = N_{\sigma\tilde{\lambda}}/L$. In combination with the inclusion $R(\sigma\lambda) \leq W(\sigma\tilde{\lambda})$ from [CSS21, Lem. 4.14] we obtain the required containment (1.2).

Via Harish-Chandra induction the map

$$\Upsilon' : \mathcal{P} \longrightarrow \text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L))) \text{ with } \overline{(\lambda, \eta)} \longmapsto \mathbf{R}_L^G(\lambda)_\eta$$

is well-defined according to [MS16, Thm. 4.7] and bijective. Hence $\Upsilon' \circ \Upsilon^{-1}$ is a bijection between $\text{Irr}_{\text{cusp}}(N)$ and $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$. Via Υ and Υ' the groups $\text{Aut}(\mathbf{G}^F)_{L, \text{HC}}$ and hence $\tilde{N}'E_L$ act on \mathcal{P} . By the description of this action given in 1.7 and 1.10 these actions coincide. Hence $\Upsilon' \circ \Upsilon^{-1}$ is $\tilde{N}'E_L$ -equivariant. By Assumption 1.8(ii) every $\psi_0 \in \text{Irr}(N \mid \text{Irr}_{\text{cusp}}(L))$ has a \tilde{L}' -conjugate ψ such that $(\tilde{N}'E_L)_\psi = \tilde{N}'_\psi E_{L, \psi}$. Hence every $\chi_0 \in \text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$ has an \tilde{N}' -conjugate χ with $(\tilde{G}E_L)_\chi = G(\tilde{N}'E_L)_\chi = G(\tilde{N}'_\chi)E_{L, \chi} = \tilde{G}_\chi E_{L, \chi}$. This implies the statement, see Lemma 1.4. \square

In the following sections we verify the assumptions of Theorem 1.8: we prove Assumption 1.8(ii), i.e., that every $\psi \in \text{Irr}_{\text{cusp}}(N)$ is \tilde{L}' -conjugate to some ψ_0 with $(\tilde{N}'E_L)_{\psi_0} = \tilde{N}'_{\psi_0} E_{L, \psi_0}$, and prove the existence of an extension map as required in 1.8(i). Note that

by Lusztig [L] and Geck [G93] an extension map exists. Their proofs are indirect and we don't see how the required properties can be deduced from their proofs. In later sections we give an independent explicit construction of the required extension map.

1.C Action on characters of normalizers of Levi subgroups

In the following we discuss some basic considerations that will be applied to ensure Assumption 1.8(ii). In the case where $L = T_0$ the assumption 1.8(ii) holds, whenever the underlying group \mathbf{G}^F is of simply-connected type, see [MS16, Proof of Cor. 5.3]. The assumption on the characters $\text{Irr}_{\text{cusp}}(N)$ is very similar to the results [CS17a, Prop. 5.13], [CS17b, Thm. 5.1] and [CS19, 5.E] on $\text{Irr}(\mathbf{N}_{\mathbf{H}}(\mathbf{S})^F)$ for Sylow Φ_d -tori \mathbf{S} of (\mathbf{H}, F) , where \mathbf{H} is a simply-connected group of type different from D_l and d is a positive integer. The proof there relies on Theorem 4.3 of [CS17b] and we use here a similar strategy. The following proposition gives the road map for the verification of Assumption 1.8(ii).

We set $W(\phi) = N_\phi/L$ for every $L \leq M \leq \tilde{L}$ and $\phi \in \text{Irr}(M)$.

Proposition 1.11 *Let \hat{N} , \tilde{L}' be as in 1.8, \mathbb{T} and $\Lambda_{L, \mathbb{T}}$ as in Assumption 1.8(i) and Υ from Proposition 1.10. Let $\lambda \in \mathbb{T}$, $\tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda | \lambda)$, $\eta \in \text{Irr}(W(\lambda))$ and $\eta_0 \in \text{Irr}(\eta|_{W(\tilde{\lambda})})$. We set $\widehat{W} := \hat{N}/L = NE_L/L$ and $\widehat{K}(\lambda) := \widehat{W}_\lambda$. If η is $\widehat{K}(\lambda)_{\eta_0}$ -stable, then*

$$(\hat{N}\tilde{L}')_{\Upsilon(\overline{(\lambda, \eta)})} = \hat{N}_{\Upsilon(\overline{(\lambda, \eta)})} \tilde{L}'_{\Upsilon(\overline{(\lambda, \eta)})}.$$

We adapt the arguments from the proof of [CS17b, Thm. 4.3], where η is assumed to be $N_{W \rtimes E_L}(W(\tilde{\lambda}))_{\eta_0}$ -stable. Note that $\widehat{K}(\lambda)$ normalizes $W(\lambda)$ but this group is in general different from $N_{W \rtimes E_L}(W(\tilde{\lambda}))$.

Proof. Recall $\psi = \Upsilon(\overline{(\lambda, \eta)}) = (\Lambda_{L, \mathbb{T}}(\lambda)\eta)^N$. By the assumptions on \mathbb{T} , $(\hat{N}\tilde{L}')_\lambda = \hat{N}_\lambda \tilde{L}'_\lambda$ for every $\lambda \in \mathbb{T}$.

Let $\tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda | \lambda)$ and $\eta_0 \in \text{Irr}(\eta|_{W(\tilde{\lambda})})$. According to [CE, 15.11], $\tilde{\lambda}$ is an extension of λ . The group $\tilde{L}'_\lambda/(LZ(\tilde{G}))$ acts by multiplication with linear characters of $W(\lambda)/W(\tilde{\lambda})$ on $\text{Irr}(W(\lambda) | \eta_0)$. Computing the action of $W(\lambda)/W(\tilde{\lambda})$ on $\text{Irr}(\tilde{L}'_\lambda | \lambda)$ show that the action of \tilde{L}'_λ/L on $\text{Irr}(W(\lambda) | \eta_0)$ is transitive. Hence the characters $\{(\Lambda_L(\lambda)\eta')^N | \eta' \in \text{Irr}(W(\lambda) | \eta_0)\}$ are the \tilde{L}'_λ -conjugates of ψ .

Let $l \in \tilde{L}'$ and $\hat{n} \in \hat{N}$ with $\psi^l = (\psi)^{\hat{n}}$. Note that $\psi^{\hat{n}} \in \text{Irr}(N | \mathbb{T})$ since \mathbb{T} is \hat{N} -stable. Then $\text{Irr}(\psi^l|_L)$ is the N -orbit of λ^l . Recall \mathbb{T} is an \tilde{L} -transversal. If $l \notin \tilde{L}'_\lambda$ then $\lambda^l \neq \lambda$ and $\lambda^l \notin \mathbb{T}$, in particular $\psi^l \notin \text{Irr}(N | \mathbb{T})$. This implies $l \in \tilde{L}'_\lambda$ and $\psi^l = (\Lambda_L(\lambda)\eta\nu)^N$ for some linear character ν of $\text{Irr}(W(\lambda)/W(\tilde{\lambda}))$. Accordingly $(\psi)^{\hat{n}} \in \text{Irr}(N | \lambda)$ and hence $(\psi)^{\hat{n}} = (\psi)^{\hat{n}'}$ for some $\hat{n}' \in \hat{N}_\lambda$. Note that

$$(\psi)^{\hat{n}'} = ((\Lambda_L(\lambda)\eta)^{\hat{n}'})^N = (\Lambda_L(\lambda)\eta^{\hat{n}'})^N.$$

The equality $\psi^l = (\psi)^{\hat{n}'}$ implies $\eta^{\hat{n}'} = \eta\nu$ and hence $\hat{n}'L \in W(\lambda)\widehat{K}(\lambda)_{\eta_0}$. As η is $\widehat{K}(\lambda)_{\eta_0}$ -stable, $\eta^{\hat{n}'} = \eta$ and hence $\psi^{\hat{n}'} = \psi$. This shows $(\hat{N}\tilde{L}')_\psi = \hat{N}_\psi \tilde{L}'_\psi$. \square

The above proposition allows the following result showing how to construct an \hat{N} -stable \tilde{L}' -transversal in $\text{Irr}_{\text{cusp}}(N)$.

Proposition 1.12 *In the situation of 1.8 assume*

- (i) $(\widehat{N}\widehat{L}')_\lambda = \widehat{N}_\lambda\widehat{L}'_\lambda$ for every $\lambda \in \mathbb{T}$,
- (ii) there exists an \widehat{N} -equivariant extension map $\Lambda_{L,\mathbb{T}}$ with respect to $L \triangleleft N$ for \mathbb{T} , and
- (iii) for every $\lambda \in \mathbb{T}$, $\tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda \mid \lambda)$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there exists some $\widehat{K}(\lambda)_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$.

Then there exists some \widehat{N} -stable \tilde{L}' -transversal in $\text{Irr}(N \mid \overline{\mathbb{T}})$, where $\overline{\mathbb{T}} \subseteq \text{Irr}_{\text{cusp}}(L)$ is the set of characters that are \tilde{L}' -conjugate to one in \mathbb{T} .

Proof. By the assumptions there exist $\mathcal{P}_1 \subseteq \mathcal{P}(L)$ such that

- if $(\lambda, \eta) \in \mathcal{P}_1$, then $\lambda \in \mathbb{T}$ and η is $\widehat{K}(\lambda)_{\eta_0}$ -stable for some $\tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda \mid \lambda)$ and $\eta_0 \in \text{Irr}(\eta \upharpoonright_{W(\tilde{\lambda})})$; and
- for each $\lambda \in \mathbb{T}$, $\tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda \mid \lambda)$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there exists some $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$ with $(\lambda, \eta) \in \mathcal{P}_1$.

According to Proposition 1.11 the characters $\Upsilon(\mathcal{P}_1)$ can form part of an E_L -stable \tilde{L}' -transversal.

According to Proposition 1.10, for every $\lambda \in \mathbb{T}$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$, the group \tilde{L}'_λ acts transitively on the set $\text{Irr}(W(\lambda) \mid \eta_0)$. Since for each $\lambda \in \mathbb{T}$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there exists some $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$ such that $(\lambda, \eta) \in \mathcal{P}_1$, each \tilde{L}' -orbit has a non-empty intersection with $\Upsilon(\mathcal{P}_1)$. This implies that every character in $\Upsilon(\mathcal{P}_1)$ has the property required, see Lemma 1.4. \square

1.D Reminder on cuspidal characters

The considerations of Section 1.B explain how the action of automorphisms on non-cuspidal characters depends on the underlying cuspidal character and a character of the relative Weyl group associated to a cuspidal pair. For the proof of Theorem A we require some general results on cuspidal characters that we collect here. By a theorem of Malle, stabilizers of cuspidal characters coincide with those of semi-simple characters (see [B06, 15.A] for a definition of semi-simple characters).

Theorem 1.13 *Let \mathbf{H} be a simply-connected simple linear algebraic group with an \mathbb{F}_q -structure given by a Frobenius map $F : \mathbf{H} \rightarrow \mathbf{H}$. Let $\mathbf{H} \rightarrow \tilde{\mathbf{H}}$ be a regular embedding and $E(\mathbf{H}^F)$ be a group of automorphisms of \mathbf{H}^F generated by graph and field automorphisms as in 1.2. Then there exists some $E(\mathbf{H}^F)$ -stable $\tilde{\mathbf{H}}^F$ -transversal in $\text{Irr}_{\text{cusp}}(\mathbf{H}^F)$.*

Proof. We abbreviate $E(\mathbf{H}^F)$ as E . Let $\chi \in \text{Irr}_{\text{cusp}}(\mathbf{H}^F)$. According to Lemma 1.4 it is sufficient to prove that χ has some $(\tilde{\mathbf{H}})^F$ -conjugate χ_0 with $(\tilde{\mathbf{H}}^F E)_{\chi_0} = \tilde{\mathbf{H}}_{\chi_0}^F E_{\chi_0}$. By [M17, Thm. 1] there exists a semi-simple character ρ of \mathbf{H}^F , such that ρ and χ have the same stabilizer. By [S12, proof of 3.4(c)] the semi-simple character ρ has some $(\tilde{\mathbf{H}})^F$ -conjugate ρ_0 with $(\tilde{\mathbf{H}}^F E)_{\rho_0} = \tilde{\mathbf{H}}_{\rho_0}^F E_{\rho_0}$. \square

In our considerations on $D_{l,sc}(q)$ we assume the following for all $l' < l$, which amounts to $A(\infty)$ for cuspidal characters in rank $< l$. This was called A_{cusp} in our Introduction. In [S21a] we will see that it is actually always satisfied.

Hypothesis 1.14 — Extension of cuspidal characters of $D_{l',sc}(q)$. Let \mathbf{H} be a simply-connected simple group of type $D_{l'}$ ($l' \geq 4$) and $F : \mathbf{H} \rightarrow \mathbf{H}$ a standard Frobenius endomorphism. Then there exists some $E(\mathbf{H}^F)$ -stable $\tilde{\mathbf{H}}^F$ -transversal \mathbb{T} in $\text{Irr}_{\text{cusp}}(\mathbf{H}^F)$

such that every $\chi \in \mathbb{T}$ extends to $\mathbf{H}^F E(\mathbf{H}^F)_\chi$.

The following facts are well-known (see also [B06, 12.1]).

Lemma 1.15 *Let \mathbf{G} be a simply-connected simple group with Frobenius endomorphism $F : \mathbf{G} \rightarrow \mathbf{G}$, \mathbf{L} an F -stable Levi subgroup of \mathbf{G} , $L := \mathbf{L}^F$, $L_0 := [\mathbf{L}, \mathbf{L}]^F$, and $\lambda \in \text{Irr}_{\text{cusp}}(L)$.*

- (a) *Then $\text{Irr}(\lambda|_{L_0}) \subseteq \text{Irr}_{\text{cusp}}(L_0)$*
- (b) *If $\mathbf{L}_0 := [\mathbf{L}, \mathbf{L}]$ is a central product of F -stable reductive groups \mathbf{H}_1 and \mathbf{H}_2 , then $\text{Irr}(\lambda|_{\mathbf{H}_1^F}) \subseteq \text{Irr}_{\text{cusp}}(\mathbf{H}_1^F)$.*
- (c) *Let $\tilde{\mathbf{G}}$ be a reductive group with \mathbb{F}_q -structure given by $F : \tilde{\mathbf{G}} \rightarrow \tilde{\mathbf{G}}$ such that $\tilde{\mathbf{G}} = \mathbf{G}\tilde{\mathbf{T}}$ for some maximally split torus $\tilde{\mathbf{T}}$ of $\tilde{\mathbf{G}}$. If $\tilde{\mathbf{T}} \cap \mathbf{L}$ is a maximal torus of \mathbf{G} , every $\tilde{\lambda} \in \text{Irr}((\tilde{\mathbf{T}}\mathbf{L})^F | \lambda)$ is cuspidal.*

Proof. For a finite group H with a split BN-pair of characteristic p , a given $\chi \in \text{Irr}(H)$ is cuspidal if and only if the corresponding representation space has no non-zero fixed point under any $O_p(P)$ for any proper parabolic subgroup P of H . It is then clear that for any $H' \triangleleft H$ with p' -index, one has $\chi \in \text{Irr}_{\text{cusp}}(H)$ if and only if $\chi|_{H'}$ has a cuspidal irreducible component (and then all are). This gives (a), and (c). For (b), note that $\mathbf{H}_1 \cap \mathbf{H}_2$ is a group of semi simple elements, so that the $O_p(P)$'s of above for $H := \mathbf{L}_0^F$ are direct products of corresponding subgroups of \mathbf{H}_1^F and \mathbf{H}_2^F . \square

Remark 1.16 (a) Let \mathbf{G} be a simply-connected simple group and $\tilde{\mathbf{G}}$ a connected algebraic group with $\tilde{\mathbf{G}} = \mathbf{G}Z(\tilde{\mathbf{G}})$. Let $F : \tilde{\mathbf{G}} \rightarrow \tilde{\mathbf{G}}$ be a Frobenius endomorphism stabilizing \mathbf{G} . Then $x \in \tilde{\mathbf{G}}^F$ can be written as $x = gz$ with $g \in \mathbf{G}$ and $z \in Z(\tilde{\mathbf{G}})$, such that $g^{-1}F(g) = zF(z^{-1})$. If $\mathcal{L} : \mathbf{G} \rightarrow \mathbf{G}$ is defined by $g \mapsto g^{-1}F(g)$ and $\hat{\mathbf{G}} := \mathcal{L}^{-1}(Z(\mathbf{G}))$, we see

$$\tilde{\mathbf{G}}^F \leq \hat{\mathbf{G}}.\hat{Z},$$

where $\tilde{\mathcal{L}} : \tilde{\mathbf{G}} \rightarrow \tilde{\mathbf{G}}$ is given by $g \mapsto g^{-1}F(g)$ and $\hat{Z} := \tilde{\mathcal{L}}^{-1}(Z(\mathbf{G})) \cap Z(\tilde{\mathbf{G}})$. Note that $\hat{\mathbf{G}}$ by its construction is independent of the choice of $\tilde{\mathbf{G}}$.

- (b) If additionally $Z(\tilde{\mathbf{G}})$ is connected the (outer) automorphisms of \mathbf{G}^F induced by conjugation with some element $g \in \tilde{\mathbf{G}}^F$ are called *diagonal automorphisms* and they are parametrized by $\mathcal{L}(g)[Z(\mathbf{G}), F]$, see also [GM, 1.5.12].

This allows the following conclusion for the group $\hat{\mathbf{G}}$ from Remark 1.16.

Lemma 1.17 *Maximal extendibility holds with respect to $\mathbf{G}^F \triangleleft \hat{\mathbf{G}}$.*

Proof. Let $\tilde{\mathbf{G}}$ be a group with connected centre, such that there exists a regular embedding $\mathbf{G} \rightarrow \tilde{\mathbf{G}}$ that is also an \mathbb{F}_q -morphism as in 1.2. Then according to a theorem of Lusztig (see [CE, 15.11]) maximal extendibility holds with respect to $\mathbf{G}^F \triangleleft \tilde{\mathbf{G}}^F$, and χ has an extension $\tilde{\chi}$ to $\tilde{\mathbf{G}}_\chi^F$. According to the above, $\tilde{\mathbf{G}}^F \leq \hat{\mathbf{G}}.\hat{Z}$. Clearly $\tilde{\chi}$ extends to $\tilde{\mathbf{G}}_\chi^F \hat{Z}$ since \hat{Z} is abelian and $[\hat{Z}, \tilde{\mathbf{G}}^F] = 1$. Now we see that $\tilde{\mathbf{G}}_\chi^F \hat{Z} = \hat{\mathbf{G}}_\chi \hat{Z}$ and hence χ extends to $\hat{\mathbf{G}}_\chi$ as well. \square

Lemma 1.18 *In the situation of Remark 1.16, let $\mathbf{K} \leq \mathbf{G}$ an F -stable reductive subgroup with $\mathbf{T}_0 \leq \mathbf{K}$. Let $\tilde{\mathbf{K}} := \mathbf{K}Z(\tilde{\mathbf{G}})$ and $\hat{\mathbf{K}} := \mathcal{L}^{-1}(Z(\mathbf{G})) \cap \mathbf{K}$. Let $\chi \in \text{Irr}(\mathbf{K}^F)$, $\tilde{\chi} \in \text{Irr}(\tilde{\mathbf{K}}^F | \chi)$ and $\nu \in \text{Irr}(\tilde{\chi}|_{Z(\tilde{\mathbf{G}}^F)})$. Assume that χ extends to $\tilde{\mathbf{K}}_\chi^F$. Let $\gamma \in E(\mathbf{G}^F)_{\chi, \mathbf{K}}$ and $\mu \in \text{Irr}(\tilde{\mathbf{K}}^F/\mathbf{K}^F)$ with $\tilde{\chi}^\gamma = \tilde{\chi}\mu$. Then the following are equivalent:*

- (i) χ has a γ -stable extension to $\hat{\mathbf{K}}_\chi$.

- (ii) for $\widehat{Z}' := \mathcal{L}(\widehat{K}_\chi)$ there exists some extension $\tilde{\nu} \in \text{Irr}(\widehat{Z}')$ of ν such that $\mu(tz) = \tilde{\nu}(z)(\tilde{\nu}(z)^\gamma)^{-1}$ for every $t \in \widehat{K}_\chi$ and $z \in \widehat{Z}'$ with $tz \in \widetilde{\mathbf{K}}_\chi^F$.

Proof. We prove the statement only in the case where $\mathbf{K} = \mathbf{G}$. The results transfer to a general \mathbf{K} as only the quotient groups are relevant to our considerations. Let $\widehat{\chi}$ be a γ -stable extension to \widehat{G}_χ , then there exists an extension $\tilde{\nu} \in \text{Irr}(\widehat{Z}')$ of ν such that $\tilde{\chi} := (\widehat{\chi} \cdot \tilde{\nu})|_{\widetilde{\mathbf{G}}_\chi^F}$. We observe $(\widehat{\chi} \cdot \tilde{\nu})^\gamma = \widehat{\chi} \cdot \tilde{\nu}^\gamma$. This leads to the given formula for μ in (ii).

For the other direction let χ_0 be the extension of χ to $\widetilde{\mathbf{G}}_\chi^F$ such that $\tilde{\chi} = \chi_0^{\widetilde{\mathbf{G}}_\chi^F}$. Then $\chi_0^\gamma = \chi_0 \mu|_{\widetilde{\mathbf{G}}_\chi^F}$ and $\chi_0 \cdot \tilde{\nu}$ is an extension of χ to $\widetilde{\mathbf{G}}_\chi^F \widehat{Z}' = \widehat{G}_\chi \widehat{Z}'$. The character $\widehat{\chi} := (\chi_0 \cdot \tilde{\nu})|_{\widehat{G}_\chi}$ satisfies

$$\widehat{\chi}^\gamma \cdot \tilde{\nu}^\gamma = (\widehat{\chi} \cdot \tilde{\nu})^\gamma = (\chi_0 \cdot \tilde{\nu})^\gamma = \chi_0^\gamma \cdot \tilde{\nu}^\gamma = \chi_0 \mu|_{\widetilde{\mathbf{G}}_\chi^F} \cdot \tilde{\nu}^\gamma.$$

There is some $\kappa \in \text{Irr}(\widehat{G}_\chi/\mathbf{G}^F)$ with $\widehat{\chi}^\gamma = \widehat{\chi}\kappa$. According [I, (6.17)] the above equality of characters implies $\kappa(t)\tilde{\nu}^\gamma(z)(\tilde{\nu}(z))^{-1} = \mu(tz)$, whenever $t \in \widehat{G}_\chi$ and $z \in \widehat{Z}'$ with $tz \in \widetilde{\mathbf{G}}_\chi^F$. By the assumption on μ and $\tilde{\nu}$ this leads to $\kappa = 1$. Then χ has a γ -stable extension to \widehat{G}_χ . \square

For later we restate $A(\infty)$ for type A (see [CS17a]).

Proposition 1.19 *Let $G = \text{SL}_n(q)$, $\widetilde{G} := \text{GL}_n(q)$ and write $E(\text{SL}_n(q))$ for the group of field and graph automorphisms of G and \widetilde{G} with regard to the usual BN-pair.*

- (a) *Then there exists an $E(\text{SL}_n(q))$ -stable $\text{GL}_n(q)$ -transversal \mathbb{T} in $\text{Irr}(\text{SL}_n(q))$, such that every $\chi \in \mathbb{T}$ extends to $\text{SL}_n(q)E(\text{SL}_n(q))_\chi$.*
- (b) *Let γ' be the automorphism of $\text{SL}_n(q)$ given by transpose-inverse and $E'(\text{SL}_n(q)) \leq \text{Aut}(\text{SL}_n(q))$ be the subgroup generated by γ' and the field automorphisms described above. Then $E'(\text{SL}_n(q))$ is abelian and there exists an $E'(\text{SL}_n(q))$ -stable $\text{GL}_n(q)$ -transversal \mathbb{T} in $\text{Irr}(\text{SL}_n(q))$, such that every $\chi \in \mathbb{T}$ extends to $\text{SL}_n(q)E'(\text{SL}_n(q))_\chi$.*

Proof. Part (a) follows from [CS17a, Thm. 4.1] using Lemma 1.4.

Let $\gamma \in E(\text{SL}_n(q))$ be the graph automorphism. Following the considerations in [CS17a, 3.2] we see that γ' and $v_0\gamma$ induce the same automorphism of $\text{SL}_n(q)$, where $v_0 \in \text{SL}_n(p)$ is defined as in [CS17a, 3.2] and p is the prime dividing q . This proves that \mathbb{T} is also $E'(\text{SL}_n(q))$ -stable. For part (b) we have to prove that every $\chi \in \mathbb{T}$ extends to its inertia group in $\text{SL}_n(q)E'(\text{SL}_n(q))$. This statement is clear whenever $E'(\text{SL}_n(q))_\chi$ is cyclic, see [I, (9.12)]. If for $\chi \in \mathbb{T}$ the group $E'(\text{SL}_n(q))_\chi$ is non-cyclic, we see $\gamma' \in E'(\text{SL}_n(q))_\chi$. Let $F_{q'} \in E'(\text{SL}_n(q))$ be a field automorphism such that $E'(\text{SL}_n(q))_\chi = \langle F_{q'}, \gamma' \rangle$. By (a) there exists some γ -stable extension of χ to $G \langle F_{q'} \rangle$. This extension is then also γ' and hence γv_0 -stable as $[v_0, F_{q'}] = 1$. From this we deduce that χ extends to $\text{SL}_n(q)E'(\text{SL}_n(q))_\chi$. \square

2 The Levi subgroup and its normalizer

In this and the following section we reprove with quite different methods that for every standard Levi subgroup L of $D_{l,sc}(q)$, every $\lambda \in \text{Irr}_{\text{cusp}}(L)$ extends to its stabilizer inside $N_{\mathbf{G}^F}(\mathbf{L})$, which follows from the mentioned results by Geck and Lusztig. For $E(\mathbf{G}^F) \leq \text{Aut}(\mathbf{G}^F)$ from Section 1.B, we construct a \widetilde{T} -transversal \mathbb{T} of $\text{Irr}_{\text{cusp}}(L)$ and an $N \text{Stab}_{E(\mathbf{G}^F)}(L)$ -equivariant extension map with respect to $L \triangleleft N$ for \mathbb{T} .

Theorem 2.1 *Let L be a standard Levi subgroup of $\mathbf{G}^F = D_{l,sc}(q)$. Let $E_L := \text{Stab}_{E(\mathbf{G}^F)}(L)$, N , $\widehat{N} := NE_L$ and $\widetilde{L}' := \widetilde{T}_0 L$ be associated to L as in 1.8. If $D_{l',sc}(q)$ is a direct factor of $[L, L]$, then assume Hypothesis 1.14 holds for $D_{l',sc}(q)$. Then:*

- (a) *There exists an \widehat{N} -stable \widetilde{L}' -transversal $\mathbb{T} \subseteq \text{Irr}_{\text{cusp}}(L)$.*
- (b) *There exists an \widehat{N} -equivariant extension map $\Lambda_{L \triangleleft N}$ with respect to $L \triangleleft N$ for \mathbb{T} .*

This implies Theorem B and ensures Assumptions (i) and (ii) of Proposition 1.12. In [BS20, Thm. 4.3] and [CSS21, Prop. 4.13] the analogous result was shown in the case where \mathbf{G} is of type A_l or C_l . The interested reader may notice that without assuming Hypothesis 1.14 for smaller ranks, the proof we give implies a version of the theorem without the equivariance statement.

Like in the proofs given in [BS20] and [CSS21], we essentially apply the following statement providing an extension map for non-linear characters.

Proposition 2.2 — [CSS21, Prop. 4.1]. *Let $K \triangleleft M$ be finite groups, let the group D act on M , stabilizing K and let $\mathbb{K} \subseteq \text{Irr}(K)$ be MD -stable. Assume there exist D -stable subgroups K_0 and V of M such that*

- (i) *the groups satisfy:*
 - (i.1) $K = K_0(K \cap V)$ and $H := K \cap V \leq Z(K)$,
 - (i.2) $M = KV$;
- (ii) *for $\mathbb{K}_0 := \bigcup_{\lambda \in \mathbb{K}} \text{Irr}(\lambda|_{K_0})$ there exist*
 - (ii.1) *a VD -equivariant extension map Λ_0 with respect to $H \triangleleft V$; and*
 - (ii.2) *an $\epsilon(V)D$ -equivariant extension map Λ_ϵ with respect to $K_0 \triangleleft K_0 \rtimes \epsilon(V)$ for \mathbb{K}_0 , where $\epsilon: V \rightarrow V/H$ denotes the canonical epimorphism.*

Then there exists an MD -equivariant extension map with respect to $K \triangleleft M$ for \mathbb{K} .

In this section we construct the set \mathbb{T} for part (a) of Theorem 2.1 and introduce groups H , K , K_0 (see Lemma 2.11), M , D and V (in Corollary 2.23) for a later application of Proposition 2.2 in the proof of Theorem 2.1(b). Here we show that the groups introduced satisfy the group-theoretic assumptions made in 2.2(i). Afterwards, in Section 3 we ensure the character-theoretic assumptions, namely 2.2(ii) in order to prove Theorem 2.1(b).

2.A Subgroups of the Levi subgroup L

As a first step we dissect the root system of \mathbf{L} and introduce subgroups of L with those new root systems. For a non-negative integer i let $\underline{i} := \{1, \dots, i\}$. For computations with elements of \mathbf{G} we use the Steinberg generators satisfying the Chevalley relations together with an explicit embedding of $D_{l,sc}(\mathbb{F})$ into $B_{l,sc}(\mathbb{F})$.

Notation 2.3 — The groups \mathbf{G} and $\overline{\mathbf{G}}$, their root systems and their elements. In this and the following section we assume that the simply-connected simple group \mathbf{G} from 1.2 is of type D_l ($l \geq 4$). Hence $\mathbf{G} \cong D_{l,sc}(\mathbb{F})$. Moreover let $\Phi := \{\pm e_i \pm e_j \mid i, j \in \underline{l}, i \neq j\}$ be its root system with simple roots $\alpha_2 := e_2 + e_1$, $\alpha_1 = e_2 - e_1$ and $\alpha_i := e_i - e_{i-1}$ ($i \geq 3$),

$$\Delta := \{\alpha_i \mid i \in \underline{l}\},$$

see [GLS, Rem. 1.8.8], where the set $\{e_i\}$ is an orthonormal basis of \mathbb{R}^l . The Chevalley generators $\mathbf{x}_\alpha(t)$, $\mathbf{n}_\alpha(t')$ and $\mathbf{h}_\alpha(t')$ ($\alpha \in \Phi$, $t, t' \in \mathbb{F}$ with $t' \neq 0$) together with the Chevalley relations describe the group structure of \mathbf{G} , see [GLS, Thm. 1.12.1].

Let $\overline{\Phi} := \{\pm e_i, \pm e_i \pm e_j \mid i, j \in \underline{l}, i \neq j\}$, $\overline{\mathbf{G}} := B_{l,sc}(\mathbb{F})$ with Chevalley generators

$\bar{\mathbf{x}}_\alpha(t)$, $\bar{\mathbf{n}}_\alpha(t')$ and $\bar{\mathbf{h}}_\alpha(t')$ ($\alpha \in \bar{\Phi}$, $t, t' \in \mathbb{F}$ with $t' \neq 0$). Assume that the structure constants of \mathbf{G} and $\bar{\mathbf{G}}$ are chosen such that $\mathbf{x}_\alpha(t) \mapsto \bar{\mathbf{x}}_\alpha(t)$ ($\alpha \in \Phi$, $t \in \mathbb{F}$) defines an embedding $\iota_D : \mathbf{G} \rightarrow \bar{\mathbf{G}}$. For simplicity of notation we write $\mathbf{x}_\alpha(t)$, $\mathbf{n}_\alpha(t')$ and $\mathbf{h}_\alpha(t')$ for the generators of $\bar{\mathbf{G}}$ and thus identify \mathbf{G} with the corresponding subgroup of $\bar{\mathbf{G}}$. This is possible according to [S10a, 10.1], see also [MS16, 2.C]. Let $\mathbf{X}_\alpha := \langle \mathbf{x}_\alpha(t) \mid t \in \mathbb{F} \rangle$ for $\alpha \in \bar{\Phi}$, $\mathbf{T} := \langle \mathbf{h}_\alpha(t') \mid \alpha \in \bar{\Phi}, t' \in \mathbb{F}^\times \rangle$ and $\bar{\mathbf{T}} := \langle \bar{\mathbf{h}}_\alpha(t') \mid \alpha \in \bar{\Phi}, t' \in \mathbb{F}^\times \rangle$. Note $\mathbf{T} = \bar{\mathbf{T}}$ following straight-forward calculations, see also [S10a, 10.1]. The group \mathbf{T} can be chosen as the group \mathbf{T}_0 from 1.2 and $\mathbf{T} \langle \mathbf{X}_\alpha \mid \alpha \in \Delta \rangle$ as the group \mathbf{B} .

Let p be the characteristic of \mathbb{F} and for every positive integer i let $F_{p^i} : \bar{\mathbf{G}} \rightarrow \bar{\mathbf{G}}$ be the Frobenius endomorphism given by $\mathbf{x}_\alpha(t) \mapsto \mathbf{x}_\alpha(t^{p^i})$ for $t \in \mathbb{F}$ and $\alpha \in \bar{\Phi}$. We write γ for the graph automorphism of \mathbf{G} given by $\mathbf{x}_{\epsilon\alpha}(t) \mapsto \mathbf{x}_{\epsilon\gamma_0(\alpha)}(t)$ for $t \in \mathbb{F}$, $\epsilon \in \{\pm 1\}$ and $\alpha \in \Delta$, where γ_0 denotes the symmetry of the Dynkin diagram of Δ of order 2 with $\alpha_2 \mapsto \alpha_1$. If $l = 4$ we denote by γ_3 the graph automorphism of \mathbf{G} induced by the symmetry of the Dynkin diagram of Δ with order 3 sending $\alpha_2 \mapsto \alpha_1$ and $\alpha_1 \mapsto \alpha_4$. We assume that $F = F_q$ for $q := p^f$, where f is a positive integer. Note that the group $E(\mathbf{G}^F)$ from 1.2 satisfies accordingly $E(\mathbf{G}^F) = \langle F_p, \gamma \rangle$ whenever $l \geq 5$, otherwise $l = 4$ and $E(\mathbf{G}^F) = \langle F_p, \gamma, \gamma_3 \rangle$.

We recall that the graph automorphism γ of \mathbf{G} is induced by an element of $\bar{\mathbf{G}}$ (see [GLS, 2.7] for the corresponding statement for classical groups). We denote the greatest common divisor of two integers i_1 and i_2 by (i_1, i_2) or $\gcd(i_1, i_2)$.

Remark 2.4 Let $\varpi \in \mathbb{F}^\times$ be a 4th root of unity. By [S10a, 10.1], see also [MS16, 2.C], the automorphism γ of \mathbf{G} is induced by conjugating with $\mathbf{n}_{e_1}(\varpi) \in \bar{\mathbf{G}}$.

Notation 2.5 Let \mathbf{L} be a Levi subgroup of \mathbf{G} such that \mathbf{BL} is a parabolic subgroup of \mathbf{G} and $\mathbf{T} \subseteq \mathbf{L}$. Let $L := \mathbf{L}^F$ and Φ' be the root system of \mathbf{L} , i.e., $\mathbf{L} = \mathbf{T} \langle \mathbf{X}_\alpha \mid \alpha \in \Phi' \rangle$. As Φ' is a parabolic root subsystem of Φ there exists a basis $\Delta' \subseteq \Delta$ of Φ' . We assume that one of the following holds:

- (i) $\Delta' \subseteq \{e_2 - e_1, e_3 - e_2, \dots, e_l - e_{l-1}\}$, or
- (ii) $\{e_1 - e_2, e_1 + e_2\} \subseteq \Delta'$.

The assumption on Δ' is justified by the following statement. Recall that a split Levi subgroup of \mathbf{G} containing \mathbf{T} is called standard if it is generated by \mathbf{T} and the \mathbf{X}_α 's such that $\alpha \in \pm\Delta'$ for some subset $\Delta' \subseteq \Delta$.

Lemma 2.6 Every standard Levi subgroup of \mathbf{G} containing \mathbf{T} is $\langle \gamma \rangle$ -conjugate to a standard Levi subgroup whose root system has a basis $\Delta' \subseteq \Delta$ satisfying 2.5(i) or 2.5(ii).

Proof. If Δ'_0 is the basis of our standard Levi subgroup, then one of the following holds:

- (i) $e_2 + e_1 \notin \Delta'_0$, and equivalently $\Delta'_0 \subseteq \{e_2 - e_1, e_3 - e_2, \dots, e_l - e_{l-1}\}$,
- (ii) $\{e_1 - e_2, e_1 + e_2\} \subseteq \Delta'_0$, or
- (iii) $e_2 - e_1 \notin \Delta'_0$, and equivalently $\Delta'_0 \subseteq \{e_2 + e_1, e_3 - e_2, \dots, e_l - e_{l-1}\}$.

In the last case $\gamma(\Delta'_0)$ satisfies $e_2 + e_1 \notin \gamma(\Delta'_0)$ and then $\gamma(\Delta'_0)$ satisfies 2.5(i). \square

In the following we decompose Φ' into smaller root systems, which are the disjoint union of irreducible root systems of the same type. By $\text{type}(\Psi)$ we denote the type of the root system Ψ .

2.7 (Decomposing Φ'). Since Φ' is a parabolic root subsystem of Φ , Φ' decomposes as a disjoint union of indecomposable root systems of type D and type A, that are called components of Φ' .

If Δ' satisfies Assumption 2.5(i), let Φ_d be the union of components of Φ' of type A_{d-1} ($d \geq 2$). If Δ' satisfies Assumption 2.5(ii), let Φ_{-1} be the union of components of Φ' that have a non-trivial intersection with $\{e_1 - e_2, e_1 + e_2\}$ and let Φ_d be the union of components of $\Phi' \setminus \Phi_{-1}$ of type A_{d-1} ($d \geq 2$). If Δ' satisfies Assumption 2.5(ii), $\text{type}(\Phi_{-1}) \in \{A_3, A_1 \times A_1, D_m \mid m \geq 4\}$.

Let \mathbb{D}' be the set of integers d such that Φ_d is defined and non-empty, that is $\text{SL}_d(\mathbb{F})$ is a summand of $[\mathbf{L}, \mathbf{L}]$. Then $\Phi' = \bigsqcup_{d \in \mathbb{D}'} \Phi_d$, a disjoint union.

Recall that $W_{\overline{\Phi}}$, the group generated by the reflections along the roots of $\overline{\Phi}$ coincides with $W_0 := \overline{N}_0/T_0$, can be identified with $\mathcal{S}_{\pm l}$ and hence acts on \underline{l} , see [GLS, Rem. 1.8.8].

2.8 (Orbits of $W_{\Phi'}$ on \underline{l}). Let \mathcal{O} be the set of $W_{\Phi'}$ -orbits on \underline{l} , $\mathcal{O}_1 \subseteq \mathcal{O}$ the set of singletons in \mathcal{O} and \mathcal{O}_d be the set of W_{Φ_d} -orbits on \underline{l} contained in $\mathcal{O} \setminus \mathcal{O}_1$, whenever $d \in \mathbb{D}'$. We define

$$\mathbb{D}(\underline{l}) = \mathbb{D} = \begin{cases} \mathbb{D}' \cup \{1\} & , \text{ if } \mathcal{O}_1 \neq \emptyset, \\ \mathbb{D}' & , \text{ otherwise.} \end{cases}$$

For $d \in \mathbb{D} \setminus \{-1\}$ let $a_d := |\mathcal{O}_d|$.

For $I \subseteq \underline{l}$ let $\Phi_I := \Phi' \cap \langle e_k \mid k \in I \rangle$ and $\overline{\Phi}_I := \overline{\Phi} \cap \langle e_k \mid k \in I \rangle$. For $d \in \mathbb{D}$ let $J_d := \bigcup_{o \in \mathcal{O}_d} o$, and $\overline{\Phi}_{J_d} := \overline{\Phi}_{J_d}$.

Next we introduce groups K, K_0 and H that will later be proven to satisfy Assumption 2.2(i) with a group M .

Notation 2.9 — Subgroups of \mathbf{L} and L . Let $\varpi \in \mathbb{F}^\times$ as in 2.4, $h_0 := \mathbf{h}_{e_1}(-1)$ and $\mathbf{h}_I(t) := \prod_{i \in I} \mathbf{h}_{e_i}(t)$ for $I \subseteq \underline{l}$ and $t \in \mathbb{F}^\times$. For $I \in \mathcal{O}$ let $\mathbf{G}_I = \langle \mathbf{X}_\alpha \mid \alpha \in \Phi_I \rangle$ and $G_I := \mathbf{G}_I^F$. Note that for $I \in \mathcal{O}_1$ the group \mathbf{G}_I is trivial. Let $H_0 := \langle h_0, \mathbf{h}_{e_i}(\varpi) \mathbf{h}_{e_{i'}}(-\varpi) \mid i, i' \in \underline{l} \rangle$. For $d \in \mathbb{D}$ let $\tilde{H}_d := \langle h_0, \mathbf{h}_I(\varpi) \mid I \in \mathcal{O}_d \rangle$, $H_d := \langle h_0, \mathbf{h}_I(\varpi) \mathbf{h}_{I'}(-\varpi) \mid I, I' \in \mathcal{O}_d \rangle$ and

$$H := \langle \tilde{H}_d \mid d \in \mathbb{D} \rangle \cap H_0.$$

Lemma 2.10 Let $\mathbb{D}_{\text{even}} := \mathbb{D} \cap 2\mathbb{Z}$ and $\mathbb{D}_{\text{odd}} := \mathbb{D} \setminus \mathbb{D}_{\text{even}}$. If $H_\epsilon := \langle \tilde{H}_d \mid d \in \mathbb{D}_\epsilon \rangle \cap H_0$ for $\epsilon \in \{\text{odd}, \text{even}\}$, then $H = H_{\text{even}} \times H_{\text{odd}}$.

Proof. An element $t \in \mathbf{T}$ can be written as $\prod_{i=1}^l \mathbf{h}_{e_i}(t_i)$ ($t_i \in \mathbb{F}^\times$). We have $t \in H_0$ if $t_i \in \langle \varpi \rangle$ and $\prod_{i=1}^l t_i^2 = 1$. In particular $\mathbf{h}_I(\varpi) \in H_0$ if and only if $|I|$ even. This implies $H_d \leq H_0$ whenever $2 \mid d$. On the other hand $\tilde{H}_d \not\leq H_0$ for every $d \in \mathbb{D}_{\text{odd}}$. \square

With this notation $Z(\mathbf{G}) = \langle h_0, \mathbf{h}_l(\varpi) \rangle$, see [GLS, Table 1.12.6].

Lemma 2.11 $H \leq Z(\mathbf{L})$.

The groups $K_0 := \langle G_I \mid I \in \mathcal{O} \rangle$ and $K := K_0 H$ then satisfy Assumption 2.2(i.1) for H .

Proof. We see that $[\mathbf{h}_I(\varpi), \mathbf{G}_I] = 1$ by the Chevalley relations and this implies the statement by the definition of H . \square

For understanding later the action of $N_{\mathbf{G}^F}(\mathbf{L})$ on $\text{Irr}(K)$ we analyse the structure of L by introducing several subgroups.

2.12 (Structure of \mathbf{L}). We observe that the Levi subgroup \mathbf{L} satisfies $\mathbf{L} = \mathbf{T} \langle \mathbf{G}_I \mid I \in \mathcal{O} \rangle$. Let $\mathbf{T}_I := \langle \mathbf{h}_{e_i}(t) \mid i \in I, t \in \mathbb{F}^\times \rangle$ for $I \in \mathcal{O}$. For $I, I' \in \mathcal{O}$ with $I \neq I'$ we see that no non-trivial linear combination of a root of Φ_I and $\Phi_{I'}$ is a root in Φ as well and hence $[\mathbf{G}_I, \mathbf{G}_{I'}] = 1$ according to Chevalley's commutator formula. By the Steinberg relations we see $[\mathbf{G}_I, \mathbf{T}_{I'}] = 1$. The group \mathbf{G}_I is either trivial or a simply-connected simple group unless $I = \mathcal{O}_{-1}$ and $\text{type}(\Phi_{-1}) = A_1 \times A_1$. Accordingly $[\mathbf{L}, \mathbf{L}] = \langle \mathbf{G}_I \mid I \in \mathcal{O} \rangle$.

We observe that $\mathbf{G}_I \cap \mathbf{T} \leq \mathbf{T}_I$ and computations with the coroot lattices prove that \mathbf{T} is the central product of the groups \mathbf{T}_I ($I \in \mathcal{O}$) over $\langle h_0 \rangle$. This implies that \mathbf{L} is the central product of the groups \mathbf{L}_I ($I \in \mathcal{O}$) where $\mathbf{L}_I := \mathbf{T}_I \mathbf{G}_I$.

Analogously we see that \mathbf{L} is the central product of the groups \mathbf{L}_d ($d \in \mathbb{D}$) over the group $\langle h_0 \rangle$, where $\mathbf{L}_d := \langle \mathbf{L}_I \mid I \in \mathcal{O}_d \rangle$.

The structure of \mathbf{L} studied above implies the following results on L . Recall $K_0 := \langle G_I \mid I \in \mathcal{O} \rangle$ from Lemma 2.11.

Lemma 2.13 Recall $\mathcal{L} : \mathbf{G} \rightarrow \mathbf{G}$ the Lang map defined by $g \mapsto g^{-1}F(g)$, let $\widehat{L} := \mathbf{L} \cap \mathcal{L}^{-1}(\langle h_0 \rangle)$ and $\widetilde{L} := \mathbf{L} \cap \mathcal{L}^{-1}(Z(\mathbf{G}))$.

- (a) If $L_I := \mathbf{L}_I^F$ for every $I \in \mathcal{O}$ and $L_0 := \langle L_I \mid I \in \mathcal{O} \rangle$, then $L_0 \triangleleft L$.
- (b) Let $I \in \mathcal{O}$ and $\widehat{L}_I := \mathbf{L}_I \cap \widehat{L}$. Then $\widehat{L}_I = \langle L_I, t_I \rangle$ for every $t_I \in \mathbf{T}_I \cap \mathcal{L}^{-1}(h_0)$. The group \widehat{L} is the central product of \widehat{L}_I ($I \in \mathcal{O}$) and for $d \in \mathbb{D}$, $\widehat{L}_d := \widehat{L} \cap \mathbf{L}_d$ is the central product of \widehat{L}_I ($I \in \mathcal{O}_d$).
- (c) $L = \langle \mathbf{L}_I^F, t_I t_{I'} \mid I, I' \in \mathcal{O} \rangle$.
- (d) K_0 is the direct product of all G_I , $K_0 \triangleleft \widetilde{L}$ and \widetilde{L}/K_0 is abelian.
- (e) If $\zeta \in \mathbb{F}^\times$ with $\zeta^{(q-1)^2} = \varpi$ and $t_{Q,2} := \mathbf{h}_Q(\zeta)$ for every $Q \subseteq \underline{l}$, then $\widetilde{L} = \widehat{L} \langle t_{\underline{l},2} \rangle$.

The arguments of Remark 1.16 show that \widetilde{L}' from 1.8 and \widetilde{L} induce the same automorphisms on \mathbf{G} .

Proof. Recall that \mathbf{L} is the central product of the groups \mathbf{L}_I , where each \mathbf{L}_I is F -stable. Every $x \in \mathbf{L}$ can be written as $\prod_{I \in \mathcal{O}} x_I$ with $x_I \in \mathbf{L}_I$. Clearly $x \in L$ if and only if $\mathcal{L}(x) = 1$. We see that $\mathcal{L}(x) = \prod_{I \in \mathcal{O}} \mathcal{L}(x_I)$ by the structure of \mathcal{L} and hence $x \in L$ implies $\mathcal{L}(x_I) \in \langle h_0 \rangle$. The group L_0 is the group of elements $\prod_{I \in \mathcal{O}} x_I$ with $x_I \in L_I := \mathbf{L}_I^F$. The group $\widehat{L} := \mathcal{L}(\langle h_0 \rangle) \cap \mathbf{L}$ is the group of elements $\prod x_I$ with $x_I \in \mathbf{L}_I$ and $\mathcal{L}(x_I) \in \langle h_0 \rangle$. Hence \widehat{L} is the central product of \widehat{L}_I ($I \in \mathcal{O}$) over $\langle h_0 \rangle$. Clearly $L_0 \triangleleft \widehat{L}$, $\widehat{L}_I = \langle L_I, t_I \rangle$ and $L = L_0 \langle t_I t_{I'} \mid I, I' \in \mathcal{O} \rangle$. This ensures the parts (a), (b) and (c).

Part (d) follows from the fact that $\mathbf{L}/\langle \mathbf{G}_I \mid I \in \mathcal{O} \rangle$ is isomorphic to a quotient of \mathbf{T} and hence abelian. For part (e) we observe $\mathcal{L}(\mathbf{h}_Q(\zeta)) = \mathbf{h}_Q(\varpi)$ for every $Q \subseteq \underline{l}$ and recall $Z(\mathbf{G}) = \langle h_0, \mathbf{h}_{\underline{l}}(\varpi) \rangle$. \square

2.B The structure of N/L

We analyse $N := N_{\mathbf{G}^F}(\mathbf{L})$ and $\overline{N} := N_{\overline{\mathbf{G}}^F}(\mathbf{L})$. In the following we identify $W_{\overline{\mathfrak{F}}}$ with $\mathcal{S}_{\pm \underline{l}}$ via the action on $\{\pm e_i \mid i \in \underline{l}\}$ and $W_{\mathfrak{F}}$ with $\mathcal{S}_{\pm \underline{l}}^D$. We generalize the notation of those permutation groups in order to describe N/L .

Notation 2.14 — Young-like subgroups, \mathcal{S}_M and \mathcal{Y}_J . Let M be a set. Given a map $\|\cdot\| : M \rightarrow \mathbb{Z}$ with $m \mapsto \|m\|$ we define \mathcal{S}_M to be the group of bijections $\pi : M \rightarrow M$ with $\|\pi(m)\| = \|m\|$ for every $m \in M$ and we write $\mathcal{S}_{\pm M}$ for the bijections $\pi : \{\pm 1\} \times M \rightarrow \{\pm 1\} \times M$ satisfying $\pi(-1, m) = (-\epsilon, m')$ and $\|m\| = \|m'\|$, whenever $m, m' \in M$ with $\pi(1, m) = (\epsilon, m')$.

For denoting the elements of \mathcal{S}_M and $\mathcal{S}_{\pm M}$ we fix a bijection $f : M \rightarrow \{1, \dots, |M|\}$.

This induces a canonical embedding $\iota : \mathcal{S}_M \rightarrow \mathcal{S}_{|M|}$ and an embedding $\iota_{\pm} : \mathcal{S}_{\pm M} \rightarrow \mathcal{S}_{\pm|M|}$. For r pairwise distinct elements $m_1, m_2, \dots, m_r \in M$ we write $(m_1, m_2, \dots, m_r) \in \mathcal{S}_M$ for the element $\iota^{-1}(f(m_1), f(m_2), \dots, f(m_r))$. Via ι_{\pm} we obtain also cycle notations for elements of $\mathcal{S}_{\pm M}$.

If J is a partition of M we write $J \vdash M$ for short. For $J \vdash M$ we set

$$\begin{aligned} \mathcal{Y}_J &:= \{\pi \in \mathcal{S}_M \mid \pi(J') = J' \text{ for every } J' \in J\}, \text{ and} \\ \mathcal{Y}_{\pm J} &:= \{\pi \in \mathcal{S}_{\pm M} \mid \pi(\{\pm 1\} \times J') = \{\pm 1\} \times J' \text{ for every } J' \in J\}. \end{aligned}$$

Let $M_{\text{odd}} := \{m \in M \mid \|m\| \text{ odd}\}$ and

$$\mathcal{S}_{\pm M}^{\text{D}} = \{\pi \in \mathcal{S}_{\pm M} \mid |(\{1\} \times M_{\text{odd}}) \cap \pi^{-1}(\{-1\} \times M_{\text{odd}})| \text{ is even}\}.$$

We use the above notation for permutation groups on the set \mathcal{O} from 2.8.

Definition 2.15 Let $\|\cdot\| : \mathcal{O} \rightarrow \mathbb{Z}$ be given by $\|I\| = d$ for every $I \in \mathcal{O}_d$ and let $\mathcal{S}_{\pm \mathcal{O}}$, $\mathcal{S}_{\pm \mathcal{O}}^{\text{D}}$ and $\mathcal{S}_{\mathcal{O}}$ be the permutation groups on \mathcal{O} defined as in 2.14 with respect to $\|\cdot\|$.

Recall that we denote by \mathbf{T} a maximal torus of \mathbf{G} and that \mathbf{L} is a standard Levi subgroup of \mathbf{G} with $\mathbf{T} \subseteq \mathbf{L}$, see 2.3 and 2.5. For $\overline{\mathbf{N}}_0 := \mathbf{N}_{\overline{\mathbf{G}}}(\mathbf{T})$ we identify the Weyl group $\overline{\mathbf{N}}_0/\mathbf{T}$ with $\mathcal{S}_{\pm l}$, the epimorphism $\rho_{\mathbf{T}} : \overline{\mathbf{N}}_0 \rightarrow \mathcal{S}_{\pm l}$ is given by

$$\rho_{\mathbf{T}}(\mathbf{n}_{e_i}(-1)) = (i, -i) \text{ and } \rho_{\mathbf{T}}(\mathbf{n}_{e_i - e_j}(-1)) = (i, j)(-i, -j).$$

With this notation we can compute the relative Weyl group of L in G . Recall $N := \mathbf{N}_{\mathbf{G}^F}(\mathbf{L})$,

Proposition 2.16 Let $N_0 := \mathbf{N}_{\mathbf{G}^F}(\mathbf{T})$, $\overline{N}_0 := \mathbf{N}_{\overline{\mathbf{G}}^F}(\mathbf{T})$, and $\overline{N} := \mathbf{N}_{\overline{\mathbf{G}}^F}(\mathbf{L})$. Then

$$\rho_{\mathbf{T}}(\overline{N} \cap N_0) / \rho_{\mathbf{T}}(L \cap N_0) \cong \mathcal{S}_{\pm \mathcal{O}} \text{ and } \rho_{\mathbf{T}}(N \cap N_0) / \rho_{\mathbf{T}}(L \cap N_0) \cong \mathcal{S}_{\pm \mathcal{O}}^{\text{D}}.$$

Proof. According to the considerations in [C, 9.2], $\rho_{\mathbf{T}}(N \cap N_0) / \rho_{\mathbf{T}}(L \cap N_0) \cong \mathbf{N}_{\overline{W}_0}(W_{\Phi'}) / W_{\Phi'}$, where $\overline{W}_0 := \overline{N}_0 / \overline{T}_0$. Note that $\mathbf{N}_{\overline{W}_0}(W_{\Phi'}) = \text{Stab}_{\overline{W}_0}(\Phi') = W_{\Phi'} \text{Stab}_{\overline{W}_0}(\Delta')$.

From the definition of Φ_{-1} one can check that $\text{Stab}_{\overline{W}_0}(\Delta')$ stabilizes $\Phi_{-1} \cap \Delta'$. This implies that $\text{Stab}_{\overline{W}_0}(\Delta')$ stabilizes $\Phi_d \cap \Delta'$ for every $d \in \mathbb{D}$, and

$$\text{Stab}_{W_{\overline{\Phi}_d}}(\Phi_d \cap \Delta') = \mathcal{S}_{\pm \mathcal{O}_d}.$$

We have $\mathbf{N}_{\overline{W}_0}(W_{\Phi'}) = W_{\overline{\Phi}_1} \times \prod_{d \in \mathbb{D}} \text{Stab}_{W_{\overline{\Phi}_d}}(\Phi_d) = \text{Stab}_{W_{\overline{\Phi}_{-1}}}(\Phi_{-1}) \times W_{\overline{\Phi}_1} \times \prod_{\substack{d \in \mathbb{D} \\ d > 1}} \text{Stab}_{W_{\overline{\Phi}_d}}(\Phi_d)$ with

$$\text{Stab}_{W_{\overline{\Phi}_{-1}}}(\Phi_{-1}) = W_{\overline{\Phi}_{-1}} \langle (1, -1) \rangle = W_{\overline{\Phi}_{-1}},$$

and

$$\text{Stab}_{W_{\overline{\Phi}_d}}(\Phi_d) = W_{\overline{\Phi}_d} \rtimes \mathcal{S}_{\pm \mathcal{O}_d}$$

for $d \in \mathbb{D}$ with $d > 1$. Hence $\rho_{\mathbf{T}}(N \cap N_0) / \rho_{\mathbf{T}}(L \cap N_0) \cong \mathcal{S}_{\pm \mathcal{O}}^{\text{D}}$. \square

By the proof we see that $\mathcal{S}_{\pm \mathcal{O}}$ corresponds to $\text{Stab}_{\overline{W}_0}(\Phi' \cap \Delta)$ and hence there exists some embedding of $\mathcal{S}_{\pm \mathcal{O}}$ into $\mathcal{S}_{\pm l}$. We fix some more notation to describe explicitly the permutations in $\mathcal{S}_{\pm l}$ corresponding to $\text{Stab}_{\overline{W}_0}(\Delta')$.

Notation 2.17 For $d \in \mathbb{D} \setminus \{-1\}$ we fix orderings on \mathcal{O}_d and the sets $I \in \mathcal{O}_d$: We write $I_{d,j}$ ($j \in \underline{a}_d$) for the sets in \mathcal{O}_d and $I_{d,j}(k) \in I_{d,j}$ ($j \in \underline{a}_d, k \in \underline{d}$) for the elements of $I_{d,j}$.

For each $k \in \underline{d}$ let $f_k^{\circ,(d)} : \underline{a}_d \rightarrow J_d$ be given by $j \mapsto I_{d,j}(k)$. Let $f_k^{(d)} : \underline{l} \rightarrow \underline{l}$ be a bijection such that $f_k^{(d)}(j) = f_k^{\circ,(d)}(j)$ for every $j \in \underline{a}_d$ and $f_k^{(d)}$ has the maximal number of fixed points. Then $f_k^{(d)}$ defines an element of $\mathcal{S}_{\pm \underline{l}}$ without sign changes, that we by abuse of notation also denote by $f_k^{(d)}$.

In the following we use that for every $Q \subseteq \underline{l}$, $\mathcal{S}_{\pm Q}$ can be seen naturally as a subgroup of $\mathcal{S}_{\pm \underline{l}}$.

Lemma 2.18 (a) Let $d \in \mathbb{D} \setminus \{-1\}$ and $\bar{\kappa}_d : \mathcal{S}_{\pm a_d} \rightarrow \mathcal{S}_{\pm J_d}$ be given by $\pi \mapsto \prod_{k \in \underline{d}} \pi^{f_k^{(d)}}$. Then $\bar{\kappa}_d$ is injective and $\text{Stab}_{\mathcal{S}_{\pm J_d}}(\Phi_d) = W_{\Phi_d} \rtimes \bar{\kappa}_d(\mathcal{S}_{\pm a_d})$
 (b) If $-1 \in \mathbb{D}$ let $\bar{\kappa}_{-1} : \mathcal{S}_{\pm 1} \rightarrow \mathcal{S}_{\pm J_{-1}}$ be the morphism with $\bar{\kappa}_{-1}(\mathcal{S}_{\pm 1}) = \langle (1, -1) \rangle$. Let $\bar{W}_d := \bar{\kappa}_d(\mathcal{S}_{\pm a_d})$ and $W^\circ(L) := \prod_{d \in \mathbb{D}} \bar{W}_d$. Then $\text{Stab}_{\bar{W}_0}(\Phi') = W_{\Phi'} W^\circ(L)$.

Proof. For (a) we observe that the sets $\bigcup_{j \in \underline{a}_d} I_{d,j}(k)$ ($k \in \underline{d}$) form a partition of J_d . This implies that the groups $\mathcal{S}_{\pm a_d}^{f_k^{(d)}}$ and $\mathcal{S}_{\pm a_d}^{f_{k'}^{(d)}}$ commute and are disjoint. We see that $\bar{\kappa}_d(\mathcal{S}_{\pm a_d})$ stabilizes \mathcal{O}_d . This proves (a). Part (b) is clear from the definitions. \square

We can choose $I_{d,j}(k)$ ($d \in \mathbb{D} \setminus \{\pm 1\}$, $j \in \underline{a}_d$ and $k \in \underline{d}$) such that $e_{I_{d,j}(k+1)} - e_{I_{d,j}(k)} \in \Delta'$ for every $j \in \underline{a}_d$ and $k \in \underline{d}-1$. With this choice, $\bar{\kappa}_d(\mathcal{S}_{\pm a_d})$ stabilizes Δ' and hence coincides $\text{Stab}_{\bar{W}_{\Phi_d}}(\Delta')$.

2.C A supplement of L in \bar{N}

In the following we determine a supplement $\bar{V} \leq \bar{N}_0$ with $\bar{N} = L\bar{V}$ and $\rho_{\mathbf{T}}(\bar{V}) = W^\circ(L)$ where $W^\circ(L)$ is the group from Lemma 2.18. We construct the group \bar{V} using extended Weyl groups \bar{V}'_{Weyl} , see 2.19. Extended Weyl groups are known as supplements of T_0 in \bar{N}_0 .

In a first step we define for every $d \in \mathbb{D}$ a subgroup $\bar{V}_d \leq \bar{N}_0$ with $\rho_{\mathbf{T}}(\bar{V}_d) = \bar{\kappa}_d(\mathcal{S}_{\pm a_d})$. We construct $\bar{\kappa}_d$, a lifting of $\bar{\kappa}_d$ via $\rho_{\mathbf{T}}$. This construction will later simplify some arguments by providing a tool to transfer results from [MS16].

By definition the group \bar{N}_0 is an extension of \bar{W}_0 by \mathbf{T} . It has proven to be more convenient to work with an extension of \bar{W}_0 by an elementary abelian 2-group, the extended Weyl group (introduced first by Tits), here denoted by \bar{V}'_{Weyl} . (Note that if $2 \mid q$ the group \bar{N}_0 is the semi-direct product of \mathbf{T} and a group isomorphic to the Weyl group.) In consideration of Remark 2.4 we work here with the group \bar{V}_0 , a \mathbf{T} -conjugate of \bar{V}'_{Weyl} . Then the graph automorphism of G is induced by an element of \bar{V}_0 (see Remark 2.4).

Notation 2.19 — The groups \bar{V}_0 , \bar{V}_I and V_I . The group $\bar{V}'_{Weyl} := \langle \bar{\mathbf{n}}'_i \mid i \in \underline{l} \rangle$ with $\bar{\mathbf{n}}'_1 := \mathbf{n}_{e_1}(1)$ and $\bar{\mathbf{n}}'_i := \mathbf{n}_{\alpha_i}(-1)$, where $\alpha_i := e_i - e_{i-1}$ ($2 \leq i \leq l$) is known as the extended Weyl group of type B_l .

Let $\zeta_8 \in \mathbb{F}$ with $\zeta_8^2 = \varpi$. The group $\bar{V}_0 := (V'_{Weyl})^{\mathbf{h}_L(\zeta_8)}$ is accordingly generated by $\bar{\mathbf{n}}_1 := (\bar{\mathbf{n}}'_1)^{\mathbf{h}_L(\zeta_8)} = \mathbf{n}_{e_1}(\varpi)$ and $\bar{\mathbf{n}}_i := (\bar{\mathbf{n}}'_i)^{\mathbf{h}_L(\zeta_8)} = \mathbf{n}_{\alpha_i}(-1)$. The group \bar{V}_0 satisfies $\bar{V}_0 \cap \bar{T}_0 = H_0$ where H_0 is defined as $\langle \mathbf{h}_\alpha(-1) \mid \alpha \in \Phi \rangle$ in 2.19. According to 2.4, $\bar{\mathbf{n}}_1 \in \bar{V}_0$ and γ induce the same automorphism of \mathbf{G} .

For $I \subseteq \underline{l}$ we set

$$V_I := \langle \mathbf{h}_0, \mathbf{n}_{\pm e_i \pm e_{i'}}(1) \mid i, i' \in I \text{ with } i \neq i' \rangle \text{ and } \bar{V}_I := V_I \langle \mathbf{n}_{e_i}(\varpi) \mid i \in I \rangle. \quad (2.1)$$

Let $\tilde{H}_I := \langle \mathbf{h}_{e_i}(\varpi) \mid i \in I \rangle$ and $\tilde{H}_0 := \tilde{H}_{\underline{l}}$.

2.20 (Facts around $H_I \triangleleft \overline{V}_I$). Maximal extendibility holds with respect to $H_I \triangleleft \overline{V}_I$, since those groups are conjugate to those considered in [MS16, Prop. 3.8] for the case where the underlying root system is of type $B_{|I|}$. For $H_I := \langle h_0, \mathbf{h}_{\pm e_i \pm e_{i'}}(-1) \mid i, i' \in I \rangle$ we obtain $\overline{V}_I \cap \mathbf{T} = H_I$.

For disjoint sets $I, I' \subseteq \underline{l}$ the Steinberg relations imply

$$[\overline{V}_I, V_{I'}] = 1 \text{ and } [\overline{V}_I, \overline{V}_{I'}] = \langle h_0 \rangle. \quad (2.2)$$

We introduce maps $\kappa_d : \tilde{H}_{\underline{a}_d} V_{\underline{a}_d} \longrightarrow \tilde{H}_0 \overline{V}_0$ with $\rho_{\mathbf{T}} \circ \kappa_d = \overline{\kappa}_d \circ \rho_{\underline{a}_d}$ for the canonical epimorphism $\rho_{\underline{a}_d} : \overline{V}_{\underline{a}_d} \longrightarrow \mathcal{S}_{\pm \underline{a}_d}$.

The following defines a lift of $\overline{W}_d := \overline{\kappa}_d(\mathcal{S}_{\pm \underline{a}_d})$ that is a subgroup of \overline{V}_0 . In 2.17 we introduced the elements $f_k^{(d)} \in \mathcal{S}_{\pm \underline{l}}$ ($d \in \mathbb{D} \setminus \{-1\}$, $k \in \underline{a}_d$) without sign changes.

Lemma 2.21 Let $d \in \mathbb{D} \setminus \{-1\}$, $m_k^{(d)} \in \overline{V} \cap \rho_{\mathbf{T}}^{-1}(f_k^{(d)})$ ($k \in \underline{a}_d$) and

$$\kappa_d : \tilde{H}_{\underline{a}_d} \overline{V}_{\underline{a}_d} \longrightarrow \tilde{H}_0 \overline{V}_0 \text{ with } x \mapsto \prod_{k=1}^d x^{m_k^{(d)}}$$

for a fixed order in \underline{a}_d . Set $\overline{V}_d := \langle \kappa_d(\overline{V}_{\underline{a}_d}) \rangle$. Then:

- (a) $\kappa_d|_{V_{\underline{a}_d}}$ is a morphism of groups;
- (b) $\kappa_d(v^x) = \kappa_d(v)^{\kappa_d(x)}$ for every $x \in \overline{V}_{\underline{a}_d}$ and $v \in V_{\underline{a}_d}$;
- (c) $\kappa_d(H_{\underline{a}_d}) = \langle h_0^d, \mathbf{h}_I(\varpi) \mathbf{h}_{I'}(-\varpi) \mid I, I' \in \mathcal{O}_d \rangle \leq H_d$;
- (d) $\kappa_d(\tilde{H}_{\underline{a}_d}) = \langle h_0^d, \mathbf{h}_I(\varpi) \mid I \in \mathcal{O}_d \rangle$,
- (e) $\rho_{\mathbf{T}} \circ \kappa_d = \overline{\kappa}_d \circ \rho_{\underline{a}_d}$ for the canonical epimorphism $\rho_{\underline{a}_d} : \overline{V}_{\underline{a}_d} \longrightarrow \mathcal{S}_{\pm \underline{a}_d}$, in particular $\rho_{\mathbf{T}}(\overline{V}_d) = \overline{W}_d = \kappa_d(\mathcal{S}_{\pm \underline{a}_d})$.

Proof. The sets $J_d(k) := f_k^{(d)}(\underline{a}_d)$ form a partition of J_d . For $x \in V_{\underline{a}_d}$ we see $x^{m_k^{(d)}} \in V_{J_d(k)}$ and hence $\kappa_d|_{V_{\underline{a}_d}}$ is independent of the order chosen in \underline{a}_d . Then $\kappa_d|_{V_{\underline{a}_d}}$ is a diagonal embedding of $V_{\underline{a}_d}$ into the central product of the groups $V_{J_d(k)}$ ($k \in \underline{a}_d$) over $\langle h_0 \rangle$. This implies that $\kappa_d|_{V_{\underline{a}_d}}$ is a morphism of groups. This proves (a).

By part (a) it is enough to prove part (b) for $x = \mathbf{n}_1$ and $v \in \{\overline{\mathbf{n}}_2^{\overline{\mathbf{n}}_1}, \overline{\mathbf{n}}_2, \overline{\mathbf{n}}_3, \dots, \overline{\mathbf{n}}_{\underline{a}_d}\}$, since $V_{\underline{a}_d}$ is generated by $\{\overline{\mathbf{n}}_2^{\overline{\mathbf{n}}_1}, \overline{\mathbf{n}}_2, \overline{\mathbf{n}}_3, \dots, \overline{\mathbf{n}}_{\underline{a}_d}\}$. The equation $\kappa_d(\overline{\mathbf{n}}_i^{\overline{\mathbf{n}}_1}) = \kappa_d(\overline{\mathbf{n}}_i)^{\kappa_d(\overline{\mathbf{n}}_1)}$ for $i \geq 3$ is clear since no non-trivial linear combination of those roots is a root. Computations show $\kappa_d(\overline{\mathbf{n}}_2^{\overline{\mathbf{n}}_1}) = \kappa_d(\overline{\mathbf{n}}_2)^{\kappa_d(\overline{\mathbf{n}}_1)}$ and hence part (b).

For part (c) we observe that $\ker(\kappa_d|_{H_{\underline{a}_d}}) = \langle h_0^{d-1} \rangle$. The equation $\rho_{\mathbf{T}} \circ \kappa_d = \overline{\kappa}_d \circ \rho_{\underline{a}_d}$ in (e) follows from $\rho_{\mathbf{T}}(m_k^{(d)}) = f_k^{(d)}$. \square

Recall that the group H from Notation 2.9 is a subgroup associated to \mathbf{L} . For understanding the above construction we consider the following statement.

Theorem 2.22 If $-1 \in \mathbb{D}$, set $\overline{V}_{-1} := \langle H_{-1}, \overline{\mathbf{n}}_1 \rangle$. Let $\overline{V} := H \langle \overline{V}_d \mid d \in \mathbb{D} \rangle$ and $V_{\mathbb{D}} := \overline{V} \cap \mathbf{G}$.

- (a) $N = LV_{\mathbb{D}}$.

(b) If $\gamma \in E_L$, then $\bar{\mathbf{n}}_1 \in \bar{V}$.

Proof. Because of $\rho_{\mathbf{T}}(\bar{V}_d) = \bar{W}_d$ we see $\rho_{\mathbf{T}}(\bar{V}) = W^\circ(L)$. Clearly $V_D \leq N$. Additionally we see that \bar{V} normalizes L and \mathbf{L} by definition. If \mathbf{L} is γ -stable, then $\bar{\mathbf{n}}_1 \in \bar{V}$. According to Remark 2.4, $\bar{\mathbf{n}}_1$ and γ induce the same automorphism of \mathbf{G} . \square

Corollary 2.23 *The groups K , K_0 and H from 2.9 and Lemma 2.11 together with $V := V_D$, $M := KV$ and $D := E_L$ satisfy the Assumption 2.2(i).*

Proof. According to Lemma 2.11, $K = HK_0$ and $H \leq Z(K)$. This is Assumption 2.2.(i.1). The equality $M = KV$ follows from the definition of M . For proving $H = V \cap K$ we show $\bar{V}_d \cap \mathbf{L} \leq H_d$ for every $d \in \mathbb{D}$. Since $\bar{V}_d \leq \bar{V}_0$ by construction and $\bar{V}_d := \langle \kappa_d(\bar{V}_{a_d}) \rangle = \bar{W}_d$, we observe that $\kappa_d(H_{a_d}) \leq H_d$ according to Lemma 2.21(c).

By the construction V_D is $\bar{V} \langle F_p \rangle$ -stable and hence E_L -stable. By the construction we also see that K_0 and V_D are D -stable. \square

3 Extending cuspidal characters of Levi subgroups

This section now focuses on the character theory of our groups. We ensure the character-theoretic Assumptions 2.2(ii) and apply Proposition 2.2 in the proof of Theorem 2.1(b). We analyse the action of \bar{V} on K_0 and consider subgroups of \bar{N} and L associated to each $d \in \mathbb{D}$. For every $d \in \mathbb{D}$ we define subgroups $K_{0,d}$ and K_d and study them separately for $d = 1$, $d \geq 2$ and $d = -1$.

3.A The inclusion $H_1 \triangleleft \bar{V}_1$

We recall here some results on the extended Weyl groups. If $1 \in \mathbb{D}$, then $H_1 \cong H_{J_1}$ and $\bar{V}_1 \cong \bar{V}_{J_1}$ for the group \bar{V}_{J_1} from 2.19. (Recall $J_d = \bigcup_{I \in \mathcal{O}_d} I$ for $d \in \mathbb{D}$, see 2.9.) We set $K_{0,1} := 1$ and $K_1 := H_1$. In order to apply 2.2 we investigate the Clifford theory for $H_1 \triangleleft \bar{V}_1$. The results are also relevant for studying $H_d \triangleleft \bar{V}_d$ for $d \geq 1$.

Proposition 3.1 *Let $l' \leq l$ be some positive integer, $\tilde{H}' := \tilde{H}_{\underline{l}'}$, $H' := \tilde{H}' \cap H_0$, $\bar{V}' := \bar{V}_{\underline{l}'}$ and $\rho' : \bar{V}' \rightarrow \mathcal{S}_{\pm l'}$ the canonical epimorphism.*

- (a) *Maximal extendibility holds with respect to $H' \triangleleft \bar{V}'$.*
- (b) *Let $\lambda \in \text{Irr}(H')$ with $h_0 \notin \ker(\lambda)$. Then some \bar{V}' -conjugate λ' of λ has some extension $\tilde{\lambda}'$ to \tilde{H}' such that $\rho'(\bar{V}'_{\tilde{\lambda}'}) = \mathcal{S}_{l'}$ and $\bar{V}'_{\lambda'} = \bar{V}'_{\tilde{\lambda}'} \langle c \rangle$ for some $c \in \bar{V}'$ with $\rho_{\mathbf{T}}(c) = \prod_{i \in \underline{l}'} (i, -i)$.*

Proof. By [MS16, Prop. 3.10] maximal extendibility holds with respect to $H' \triangleleft \bar{V}'$. Note that \bar{V}' coincides with the group \bar{V}^t considered in [MS16]. This proves part (a).

Let $\lambda \in \text{Irr}(H')$ with $\lambda(h_0) = -1$ and $\tilde{\lambda} \in \text{Irr}(\tilde{H}' \mid \lambda)$. Note that \tilde{H}' is the l' -fold central product of the cyclic groups $\langle \mathbf{h}_{e_i}(\varpi) \rangle$ ($i \in \underline{l}'$) over $\langle h_0 \rangle$. The group \bar{V}' acts by permutation and inversion on the factors. It is then easy to see that some \bar{V}' -conjugate λ' of λ has an extension $\tilde{\lambda}' \in \text{Irr}(\tilde{H}')$ such that

$$\tilde{\lambda}'(\mathbf{h}_{e_i}(\varpi)) = \tilde{\lambda}'(\mathbf{h}_{e_{i'}}(\varpi)) \text{ for every } i, i' \in \underline{l}'.$$

The other extension of λ' to \tilde{H} is $(\tilde{\lambda}')^{-1}$. Observe that $(\tilde{\lambda}')^2$ is the character with kernel H' . (Recall $\tilde{H}'/H' \cong C_2$ and hence there is exactly one character with this property.) The

element $c \in \overline{V}'$ with $\rho'(c) = \prod_{i=1}^{l'}(i, -i)$ satisfies $(\tilde{\lambda}')^c = (\tilde{\lambda}')^{-1}$ and hence $\overline{V}'_{\tilde{\lambda}'} = \overline{V}'_{\tilde{\lambda}'} \langle c \rangle$. According to (a) there exists some extension ϕ_0 of λ' to $\overline{V}'_{\tilde{\lambda}'}$. Then $\phi_0|_{\overline{V}'_{\tilde{\lambda}'}}$ and $\tilde{\lambda}'$ determine a common extension ϕ to $\tilde{H}'\overline{V}'_{\tilde{\lambda}'}$, see [S10b, 4.1(a)]. By this construction $\phi|_{\overline{V}'_{\tilde{\lambda}'}}$ is c -stable. \square

3.B The inclusion $K_d \triangleleft K_d \overline{V}_d$ for $d \geq 2$

In the following we investigate the groups $K_d := H_d \langle G_I \mid I \in \mathcal{O}_d \rangle$ and $K_d \overline{V}_d$ for $d \in \mathbb{D} \setminus \{\pm 1\}$, where $\mathbf{G}_I = \langle \mathbf{X}_\alpha \mid \alpha \in \Phi_I \rangle$ and $G_I = \mathbf{G}_I^F$, see Notation 2.9 and Lemma 2.13. Again $\gcd(i_1, i_2)$ is the greatest common divisor of two integers i_1 and i_2 .

Lemma 3.2 *Let $I \in \mathcal{O} \setminus (\{J_{-1}\} \cup \mathcal{O}_1)$ and $\mathbf{Z}_I := \mathbf{h}_I(\mathbb{F}^\times)$. Then:*

- (a) $\mathbf{G}_I \cong \mathrm{SL}_{|I|}(\mathbb{F})$ and $G_I \cong \mathrm{SL}_{|I|}(q)$;
- (b) $\mathbf{L}_I = \mathbf{G}_I \mathbf{Z}_I$, $\mathbf{L}_I / \langle h_0 \rangle \cong \mathrm{GL}_{|I|}(\mathbb{F})$ and $|\mathbf{G}_I \cap \mathbf{Z}_I| = \frac{|I|_{p'}}{\gcd(2, |I|)}$; and
- (c) $L_I \cong \mathrm{GL}_{|I|}(q)$ if $2 \nmid |I|$.

Proof. By the assumptions $d := |I| \geq 1$ and Φ_I is a root system of type A_d . The isomorphism in (a) follows from the Chevalley relations. By definition the groups \mathbf{G}_I are simply-connected, since Φ_I is a parabolic subsystem of Φ' of type $A_{|I|-1}$. Note $I \neq J_{-1}$.

Any element of \mathbf{T}_I can be written as $\prod_{i \in I} \mathbf{h}_{e_i}(t_i)$ for some $t_i \in \mathbb{F}^\times$. Let $\kappa \in \mathbb{F}$ with $\kappa^{|I|} = \prod_{i \in I} t_i$ and fix $j \in I$. Then by the Chevalley relations

$$\begin{aligned} \prod_{i \in I} \mathbf{h}_{e_i}(t_i) &= \mathbf{h}_{e_j}(t_j \kappa^{-1}) \mathbf{h}_{e_j}(\kappa) \prod_{\substack{i \in I \\ i \neq j}} (\mathbf{h}_{e_j}(t_i^{-1} \kappa)^{-1} \mathbf{h}_{e_j}(t_i^{-1} \kappa) \mathbf{h}_{e_i}(t_i \kappa^{-1}) \mathbf{h}_{e_i}(\kappa)) = \\ &= \left(\mathbf{h}_{e_j}(\kappa^{-|I|} \prod_{i \in I} t_i) \right) \mathbf{h}_{e_j}(\kappa) \prod_{\substack{i \in I \\ i \neq j}} (\mathbf{h}_{e_i - e_j}((t_i \kappa^{-1})^2) \mathbf{h}_{e_i}(\kappa)) = \\ &= \left(\prod_{\substack{i \in I \\ i \neq j}} \mathbf{h}_{e_i - e_j}((t_i \kappa^{-1})^2) \right) \mathbf{h}_I(\kappa). \end{aligned}$$

Accordingly $\mathbf{T}_I = (\mathbf{T}_I \cap \mathbf{G}_I) \mathbf{Z}_I$. We note that $\mathbf{G}_I \cong \mathrm{SL}_{|I|}(\mathbb{F})$ and $G_I \cong \mathrm{SL}_{|I|}(q)$ as F acts on \mathbf{G}_I as standard Frobenius endomorphism. By the Chevalley relations $\mathbf{Z}_I \leq \mathrm{C}_L(\mathbf{G}_I)$ and $\mathbf{L}_I = \mathbf{T}_I \mathbf{G}_I = \mathbf{Z}_I \mathbf{G}_I$.

The calculations above show that an element of $\mathbf{Z}_I \cap \mathbf{G}_I$ can be written as $\prod_{i \in I} \mathbf{h}_{e_i}(t_i)$ with $t_j = t_i$ and $t_i^{|I|} = 1$ for every $i \in I$. For $d \in \mathbb{D}_{\text{even}}$, the element $\prod_{i=1}^d \mathbf{h}_{e_i}(-1)$ is trivial and hence $|\mathbf{Z}_I \cap \mathbf{G}_I| = \frac{|I|_{p'}}{\gcd(2, |I|_{p'})}$.

If $2 \mid d$, then with similar considerations as above we see

$$h_0 = \mathbf{h}_I(\zeta) \prod_{\substack{i \in I \\ i \neq j}} \mathbf{h}_{e_i - e_j}(\zeta^{-2}), \quad (3.1)$$

where $\zeta \in \mathbb{F}^\times$ is a root of 1 of order $2|I|_2$. This implies that $\mathbf{L}_I / \langle h_0 \rangle$ is the central product of the one-dimensional torus $\mathbf{Z}_I / \langle h_0 \rangle$ and $\mathbf{G}_I / \langle h_0 \rangle$ over $Z(\mathbf{G}_I)$. Accordingly $\mathbf{L}_I / \langle h_0 \rangle \cong \mathrm{GL}_{|I|}(\mathbb{F})$.

For odd d this implies analogously $\mathbf{L}_I \cong \mathrm{GL}_d(\mathbb{F})$ and $L_I \cong \mathrm{GL}_d(q)$. This is the statement in (c). \square

Next we study how \tilde{L} acts on $K_{0,d}$, which includes the action induced by t_I ($I \in \mathcal{O}$) and $t_{\underline{L},2}$ from Lemma 2.13. Recall that $\hat{L}_I := \mathbf{L}_I \cap \mathcal{L}^{-1}(\langle h_0 \rangle)$ satisfies $\hat{L}_I = \langle L_I, t_I \rangle$ for some $t_I \in \mathbf{T}_I \cap \mathcal{L}^{-1}(h_0)$, and $\mathcal{L}^{-1}(\mathbf{h}_{\underline{L}}(\varpi)) \cap \mathbf{L} = \langle \hat{L}, t_{\underline{L},2} \rangle$ with $t_{\underline{L},2} = \mathbf{h}_{\underline{L}}(\zeta)$. According to Remark 1.16(b), diagonal automorphisms of G_I are parametrized by $Z(\mathbf{G}_I)/[Z(\mathbf{G}_I), F]$.

Lemma 3.3 *Let $I \in \mathcal{O} \setminus (\mathcal{O}_1 \cup \{J_{-1}\})$.*

- (a) *If $2 \nmid |I|$, then $\hat{L}_I = L_I \mathbf{C}_{\mathbf{T}_I}(L_I)$, in particular t_I from Lemma 2.13(b) can be chosen such that $t_I \in \mathbf{C}_{\mathbf{T}_I}(L_I)$.*
- (b) *For $2 \mid |I|$, the element t_I induces on G_I a diagonal automorphism corresponding to $g[Z(\mathbf{G}_I), F]$ with $g \in Z(\mathbf{G}_I)$ of order $|Z(\mathbf{G}_I)|_2$.*
- (c) *$t_{\underline{L},2} \in \mathbf{C}_{\hat{L}}(L_I)$.*

Proof. Keep $d := |I|$. According to the theorem of Lang we can choose $t_I \in \mathbf{T}_I$ such that $t_I^{-1}F(t_I) = h_0$ as \mathbf{T}_I is connected.

If $2 \nmid |I|$ we see that $h_0 = \mathbf{h}_I(-1)$ and hence $h_0 \in \mathbf{Z}_I$. Since \mathbf{Z}_I is again connected t_I can be chosen in $Z(\mathbf{L}_I)$, whence (a).

Following (3.1), $h_0 = z_1 z_2$ for some $z_1 \in \mathbf{Z}_I$ and $z_2 \in Z(\mathbf{G}_I)$. Here z_2 is an element of order $|I|_2 = d_2$. Then the element t_I can be analogously written as $z g$ with $z \in Z(\mathbf{G}_I)$ and $g \in \mathbf{G}$ such that $\mathcal{L}(z) = z_1$ and $\mathcal{L}(g) = z_2$. As g induces on \mathbf{G} a diagonal automorphism associated to $z_2[Z(\mathbf{G}_I), F]$, the element $t_I \in \hat{\mathbf{G}}_I$ with $x^{-1}F(x) = h_0$ induces the same diagonal automorphism. This gives (b).

The element $t_{\underline{L},2} = \mathbf{h}_{\underline{L}}(\zeta)$ from Lemma 2.13(e) centralizes \mathbf{G}_I since the Weyl group of \mathbf{G}_I centralizes $t_{\underline{L},2}$. \square

Recall the groups $\tilde{H}_d = \langle h_0, \mathbf{h}_I(\varpi) \mid I \in \mathcal{O}_d \rangle$, $H_0 = \langle \mathbf{h}_\alpha(-1) \mid \alpha \in \Phi \rangle$ and $H_d = \tilde{H}_d \cap H_0$ defined in 2.9 for every $d \in \mathbb{D}$. Using the groups G_I from Lemma 2.13 let $\tilde{K}_{0,d} := \langle G_I \mid I \in \mathcal{O}_d \rangle$ and $K_d := H_d \tilde{K}_{0,d}$. If $\mathbb{D} = \{d\}$, then $K_0 = K_{0,d}$ and $K = K_d$. As $\bar{V}_d \cap K_d \leq \mathbf{C}_{\mathbf{L}}(\mathbf{G}_I)$ as a consequence of Lemma 2.11, there is a well-defined action of \bar{V}_d/H_d on K_d .

Lemma 3.4 — The action of V_d on $K_{0,d}$. *Let $d \in \mathbb{D}$. Let $\epsilon_d : \bar{V}_d \rightarrow \bar{V}_d/H_d$ be the canonical morphism and $\bar{\mathbf{n}}_1^{(d)} := \kappa_d(\bar{\mathbf{n}}_1)$. Then:*

- (a) *$K_{0,d} \rtimes \epsilon(\bar{V}_d) \cong (G_{I_{d,1}} \rtimes \langle \epsilon(\bar{\mathbf{n}}_1^{(d)}) \rangle) \wr \mathcal{S}_{a_d}$.*
- (b) *Then $\bar{\mathbf{n}}_1^{(d)}$ induces a non-trivial graph automorphism on $G_{I_{d,1}}$.*
- (c) *If $2 \nmid d$, $\bar{\mathbf{n}}_1^{(d)}$ induces on $L_{I_{d,1}}$ a combination of a non-trivial graph and an inner automorphism via the isomorphism of $L_{I_{d,1}} \cong \mathrm{GL}_d(q)$ from Lemma 3.2(c).*

Proof. Part (a) follows from the Steinberg presentation.

For part (b) we see that $G_{I_{d,1}}^{\bar{\mathbf{n}}_1^{(d)}} = G_{I_{d,1}}$ and $G_{I_{d,1}} \cap \langle h_0 \rangle = \{1\}$ from the Chevalley relations. We compute the action of $\bar{\mathbf{n}}_1^{(d)}$ on $G_{I_{d,1}}$ in the quotient $G/\langle h_0 \rangle$ or $G_{I_{d,1}} \times \langle h_0 \rangle / \langle h_0 \rangle$, respectively. In [GLS, 2.7] the group $\mathbf{G}/\langle h_0 \rangle$ and its Steinberg generators are given explicitly as subgroup and elements of $\mathrm{SO}_{2l}(q)$. The element $\bar{\mathbf{n}}_1^{(d)}$ acts on $\mathbf{G}_{I_{d,1}} \langle h_0 \rangle / \langle h_0 \rangle$ by transpose-inverse via $G_{I_{d,1}} \cong \mathrm{SL}_d(q)$. Computations in that group show part (b).

The element $\bar{\mathbf{n}}_1^{(d)}$ acts by inversion on $\mathbf{Z}_{I_{d,1}}$ and hence $\bar{\mathbf{n}}_1^{(d)}$ satisfies the statement in part (c) as $\mathbf{L}_{I_{d,1}} = \mathbf{G}_{I_{d,1}} \mathbf{Z}_{I_{d,1}}$. \square

Next we study an analogue of \tilde{L} from Lemma 2.13 associated to $d \in \mathbb{D}$, that is defined using the Lang map \mathcal{L} from there. Note that $\mathbf{h}_{\underline{L}}(\varpi) \notin \mathbf{T}_{J_d}$ whenever $\mathbb{D} \neq \{d\}$, but $\mathbf{h}_{\underline{L}}(\varpi) = \prod_{d \in \mathbb{D}} \mathbf{h}_{J_d}(\varpi)$.

Proposition 3.5 *Let $d \in \mathbb{D} \setminus \{\pm 1\}$, $\epsilon_d : \overline{V}_d \rightarrow \overline{V}_d/H_d$ be as in Lemma 3.4, $\tilde{T}_d := \mathbf{T}_{J_d} \cap \mathcal{L}^{-1}(\langle h_0, \mathbf{h}_{J_d}(\varpi) \rangle)$ and $\tilde{L}_d := \tilde{T}_d K_{0,d}$. Then:*

- (a) *There exists some $\overline{V}_d \langle F_p \rangle$ -stable \tilde{L}_d -transversal \mathbb{T}_d° in $\text{Irr}_{\text{cusp}}(K_{0,d})$.*
- (b) *There exists an $\epsilon_d(\overline{V}_d) \times \langle F_p \rangle$ -equivariant extension map Λ_{ϵ_d} with respect to $K_{0,d} \triangleleft K_{0,d} \rtimes \epsilon_d(V_d)$ for \mathbb{T}_d° .*
- (c) *Maximal extendibility holds with respect to $K_{0,d} \triangleleft \tilde{L}_d$ and $K_d \triangleleft \tilde{L}_d$.*

For the proof of part (b) we require a strengthening of a result on wreath products that can for example be found in [K20, Thm. 2.10].

Lemma 3.6 *Let $X \rtimes Y$ be a finite group and A be a group of automorphisms of $X \rtimes Y$, stabilizing X, Y and some $\mathbb{K} \subseteq \text{Irr}(X)$. Let a be a positive integer. Note that A acts on $X^a \triangleleft (X \rtimes Y) \wr \mathcal{S}_a$ by simultaneously acting on $(X \rtimes Y)^a$ and trivially on \mathcal{S}_a . In this context we write then ΔA for that group. If there exists an $(X \rtimes Y) \rtimes A$ -equivariant extension map $X \triangleleft X \rtimes Y$ for \mathbb{K} there exists an $((X \rtimes Y) \wr \mathcal{S}_a) \rtimes \Delta A$ -equivariant extension map with respect to $X^a \triangleleft (X \rtimes Y) \wr \mathcal{S}_a$ for $\mathbb{K}^a := \{\chi_1 \otimes \cdots \otimes \chi_a \mid \chi_i \in \mathbb{K}\}$.*

Proof. This follows by the considerations in the proof of [K20, Thm. 2.10] using the construction of representation of wreath products given in [N, 10.1]. \square

Proof of Proposition 3.5. Let $I_1 := I_{d,1}$. Via the isomorphism $G_{I_1} \cong \text{SL}_d(q)$ from Lemma 3.2 the $E(\text{SL}_d(q))$ -stable $\text{GL}_d(q)$ -transversal in $\text{Irr}(\text{SL}_d(q))$ from Proposition 1.19 determines a subset $\mathbb{T}_{I_1} \subseteq \text{Irr}_{\text{cusp}}(G_{I_1})$. According to Lemma 3.4(b) this set is $\overline{\mathbf{n}}_1^{(d)}$ -stable. The $E(\text{SL}_d(q))$ -stable $\text{GL}_d(q)$ -transversal in $\text{Irr}(\text{SL}_d(q))$ can even be chosen such that each character extends to its inertia group in $\text{SL}_d(q) \times E(\text{SL}_d(q))$. Accordingly maximal extendibility holds with respect to $G_{I_1} \triangleleft G_{I_1} \rtimes \langle F_p, \epsilon_d(\overline{\mathbf{n}}_1^{(d)}) \rangle$ for \mathbb{T}_{I_1} .

Note that \mathbb{T}_{I_1} is $N_{\overline{V}_d}(G_{I_1})$ -stable, as $N_{\overline{V}_d}(G_{I_1})$ acts as $\langle \overline{\mathbf{n}}_1^{(d)} \rangle$. Accordingly via conjugation with elements of \overline{V}_d the set \mathbb{T}_{I_1} determines characters $\mathbb{T}_I \subseteq \text{Irr}_{\text{cusp}}(G_I)$ for every $I \in \mathcal{O}_d$. Recall that by Lemma 2.13(d) the group K_0 is the direct product of the groups G_I ($I \in \mathcal{O}$). Analogously $K_{0,d}$ is the direct product of the groups G_I ($I \in \mathcal{O}_d$).

The product \mathbb{T}_d° of these characters $\prod_{I \in \mathcal{O}_d} \mathbb{T}_I$ defines a $\overline{V}_d \langle F_p \rangle$ -stable set in $\text{Irr}_{\text{cusp}}(K_{0,d})$. By this construction \mathbb{T}_d° is $\overline{V}_d \langle F_p \rangle$ -stable. Following the description of the action of \tilde{L} on $K_{0,d}$ given in Lemma 3.3 we see that \mathbb{T}_d° is an \tilde{L}_d -transversal in $\text{Irr}_{\text{cusp}}(K_{0,d})$. This proves part (a).

Recall $K_{0,d} \rtimes \epsilon_d(\overline{V}_d) \cong (G_{I_1} \rtimes \langle \epsilon(\overline{\mathbf{n}}_1^{(d)}) \rangle) \wr \mathcal{S}_{a_d}$ from Lemma 3.4. As stated above, maximal extendibility holds with respect to $G_{I_1} \triangleleft G_{I_1} \rtimes \langle F_p, \epsilon(\overline{\mathbf{n}}_1^{(d)}) \rangle$ for \mathbb{T}_{I_1} . According to Lemma 3.6 this implies by the choice of \mathbb{T}_d° that there is an $\epsilon_d(\overline{V}_d) \langle F_p \rangle$ -equivariant extension map Λ_{ϵ_d} with respect to $K_{0,d} \triangleleft K_{0,d} \rtimes \epsilon_d(V_d)$ for \mathbb{T}_d° .

According to Lemma 1.17 maximal extendibility holds with respect to $G_I \triangleleft \tilde{G}_I$ where $\tilde{G}_I := \mathbf{G}_I \cap \mathcal{L}^{-1}(Z(\mathbf{G}_I))$. Additionally $[\mathbf{G}_I, \mathbf{Z}_I] = 1$ for $\mathbf{Z}_I := \mathbf{h}_I(\mathbb{F}^\times)$ from Lemma 3.2. We observe that $\tilde{L}_d \leq \langle \tilde{G}_I \mid I \in \mathcal{O}_d \rangle \langle \mathbf{Z}_I \mid I \in \mathcal{O}_d \rangle$, even more precisely

$$\tilde{L}_d \leq \langle \tilde{G}_I \mid I \in \mathcal{O}_d \rangle \langle \hat{\mathbf{Z}}_I \mid I \in \mathcal{O}_d \rangle,$$

where $\widehat{Z}_I := \mathcal{L}^{-1}(\mathbf{Z}(\mathbf{G}_I) \cap \mathbf{Z}_I) \cap \mathbf{Z}_I$. We see that maximal extendibility holds with respect to $K_{0,d} \triangleleft \langle \widetilde{G}_I \mid I \in \mathcal{O}_d \rangle \triangleleft \langle \widehat{Z}_I \mid I \in \mathcal{O}_d \rangle$. Hence maximal extendibility holds with respect to $K_{0,d} \triangleleft \widetilde{L}_d$ and $K_d \triangleleft \widetilde{L}_d$, as $\widetilde{L}_d/K_{0,d}$ is abelian. \square

Lemma 3.7 *Let $d \in \mathbb{D} \setminus \{\pm 1\}$.*

- (a) *Maximal extendibility holds with respect to $H_d \triangleleft \overline{V}_d$.*
- (b) *If $2 \nmid d$, $\lambda \in \text{Irr}(H_d)$ with $\lambda(h_0) = -1$ and $\tilde{\lambda} \in \text{Irr}(\widetilde{H}_d \mid \lambda)$, then $(\overline{V}_d)_{\tilde{\lambda}} \leq V_D$ and $(\overline{V}_d)_\lambda = (\overline{V}_d)_{\tilde{\lambda}} \langle c_d \rangle$ for some $c_d \in \overline{V}_d$.*

Proof. Recall that by [MS16, Prop. 3.8] maximal extendibility holds with respect to $H_{a_d} \triangleleft \overline{V}_{a_d}$. Via the map $\kappa_d : \overline{V}_{a_d} \rightarrow \overline{V}_d$ from Lemma 2.21 the maximal extendibility with respect to $H_{a_d} \triangleleft \overline{V}_{a_d}$ gives a \overline{V}_d -equivariant extension map for $\kappa_d(H_{a_d}) \triangleleft \kappa_d(V_{a_d})$. This implies part (a) according to [S10a, 4.1(a)].

In part (b) we assume $2 \nmid d$ and hence $\kappa_d(H_{a_d}) = H_d$. The character $\lambda \in \text{Irr}(H_d)$ with $\lambda(h_0) = -1$ corresponds via κ_d , λ to some $\lambda_0 \in \text{Irr}(H_{a_d})$ with $\lambda_0(h_0) = -1$. Proposition 3.1(b) implies that via κ_d there is some \overline{V}_d -conjugate λ' of λ with $\rho_{\mathbf{T}}(\overline{V}_{\lambda'}) = \mathcal{S}_{\mathcal{O}_d}$ for any $\tilde{\lambda}' \in \text{Irr}(\widetilde{H}_d \mid \lambda')$ and $\overline{V}_{d,\lambda'} = \overline{V}_{\tilde{\lambda}'} \langle c'_d \rangle$ for some $c'_d \in \text{Irr}(\overline{V}_d)$ with $\rho_{\mathbf{T}}(c'_d) = \prod_{i \in J_d} (i, -i)$. We observe that $\overline{V}_{d,\tilde{\lambda}'} \leq V_D$. Because of $V_D \triangleleft \overline{V}$ this implies $(\overline{V}_d)_{\tilde{\lambda}} \leq V_D$ and $(\overline{V}_d)_\lambda = (\overline{V}_d)_{\tilde{\lambda}} \langle c_d \rangle$ for some $c_d \in \overline{V}_d$. This proves part (b). \square

3.C Consideration on $K_{-1} \triangleleft K_{-1} \overline{V}_{-1}$

The group structure of $G_{J_{-1}}$ depends on $\text{type}(\Phi_{-1})$. By its definition, $\text{type}(\Phi_{-1}) \in \{\mathbf{A}_1 \times \mathbf{A}_1, \mathbf{A}_3, \mathbf{D}_{|J_{-1}|}\}$. For the application of Proposition 2.2 we prove the following statement. Recall $\overline{V}_{-1} = \langle \mathbf{n}_{e_1}(\varpi), h_0 \rangle$, $\widetilde{H}_{-1} = \langle \mathbf{h}_{J_{-1}}(\varpi), h_0 \rangle$, $H_{-1} = \widetilde{H}_{-1} \cap H_0$ and $G_{J_{-1}} = \langle \mathbf{X}_\alpha \mid \alpha \in \Phi_{-1} \rangle^F$. As before we set $K_{0,-1} := G_{J_{-1}}$ and $K_{-1} := H_{-1} G_{J_{-1}}$.

Proposition 3.8 *Assume Hypothesis 1.14 holds for $\mathbf{G}_{J_{-1}}^F$ if Φ_{-1} is of type D. Let $\epsilon_{-1} : \overline{V}_{-1} \rightarrow \overline{V}_{-1}/H_{-1}$ be the canonical epimorphism and $\widetilde{L}_{-1} := (\mathbf{G}_{J_{-1}} \mathbf{T}_{J_{-1}}) \cap \mathcal{L}^{-1}(\langle h_0, \mathbf{h}_{J_{-1}}(\varpi) \rangle)$. Then:*

- (a) *There exists some $\overline{V}_{-1} \langle F_p \rangle$ -stable \widetilde{L}_{-1} -transversal \mathbb{T}_{-1}° in $\text{Irr}_{\text{cusp}}(K_{0,-1})$.*
- (b) *There exists an $\epsilon_{-1}(\overline{V}_{-1}) \langle F_p \rangle$ -equivariant extension map $\Lambda_{\epsilon_{-1}}$ with respect to $K_{0,-1} \triangleleft K_{0,-1} \rtimes \epsilon_{-1}(\overline{V}_{-1})$ for \mathbb{T}_{-1}° .*
- (c) *Maximal extendibility holds with respect to $K_{0,-1} \triangleleft \widetilde{L}_{-1}$ and $K_{-1} \triangleleft \widetilde{L}_{-1}$.*

Proof. As in the proof of Lemma 2.13 we see that $\widetilde{L}_{-1} = T_{-1} G_{J_{-1}} \langle t_{J_{-1}}, t_{J_{-1},2} \rangle$, where $T_{-1} := \mathbf{T}_{J_{-1}}^F$, $\zeta \in \mathbb{F}^\times$ with $\zeta^{(q-1)^2} = \varpi$, $t_{J_{-1}} := \mathbf{h}_{e_1}(\zeta^2)$ and $t_{J_{-1},2} := \mathbf{h}_{J_{-1}}(\zeta)$. Note that the action of \widetilde{L} on $G_{J_{-1}}$ coincides with the one of $\langle T_{-1}, t_{J_{-1}}, t_{J_{-1},2} \rangle$ up to inner automorphisms. By the definition of $G_{J_{-1}}$ we see

$$G_{J_{-1}} \cong \begin{cases} \mathbf{D}_{|J_{-1}|,sc}(q) & \text{if } \text{type}(\Phi_{-1}) = \mathbf{D}_{|J_{-1}|}, \\ \mathbf{SL}_4(q) & \text{if } \text{type}(\Phi_{-1}) = \mathbf{A}_3, \\ \mathbf{SL}_2(q) \times \mathbf{SL}_2(q) & \text{if } \text{type}(\Phi_{-1}) = \mathbf{A}_1 \times \mathbf{A}_1. \end{cases}$$

Assume $\text{type}(\Phi_{-1}) = \mathbf{D}_{l_{-1}}$ with $l_{-1} := |J_{-1}|$. Then $T_{-1} \leq G_{J_{-1}}$. The elements $t_{J_{-1}}$ and $t_{J_{-1},2}$ act as diagonal automorphisms on $G_{J_{-1}}$. Part (a) follows from Theorem 1.13. By Hypothesis 1.14 we can choose a $\overline{V}_{-1} \langle F_p \rangle$ -stable \widetilde{L}_{-1} -transversal \mathbb{T}_{-1}° in $\text{Irr}_{\text{cusp}}(K_{0,-1})$

such that maximal extendibility holds with respect to $K_{0,-1} \triangleleft K_{0,-1} \rtimes \langle \gamma, F_p \rangle$ for \mathbb{T}_{-1} . Note that $K_{0,-1} \rtimes \langle \gamma, F_p \rangle = K_{0,-1} \rtimes (\epsilon_{-1}(\overline{V}_{-1}) \times \langle F_p \rangle)$. By this choice we see that an extension map Λ_{e_1} as required in part (b) exists. Note that the actions on $G_{J_{-1}}$ induced by γ and $\mathbf{n}_{e_1}(\varpi)$ coincide by 2.3. According to Lemma 1.17 maximal extendibility holds with respect to $G_{J_{-1}} \triangleleft \widetilde{L}_{-1}$. This proves part (c) in the case where $\text{type}(\Phi_{-1}) = D_{l_{-1}}$.

Assume $\text{type}(\Phi_{-1}) = A_1 \times A_1$, then $t_{J_{-1}}$ induces on both factors a non-inner diagonal automorphism, while $t_{J_{-1},2}$ induces a non-inner diagonal automorphism only on one factor, since $h_0 = \mathbf{h}_{e_1-e_2}(-1)\mathbf{h}_{e_1+e_2}(-1)$ and $\mathbf{h}_{e_1}(\varpi)\mathbf{h}_{e_2}(\varpi) = \mathbf{h}_{e_1+e_2}(-1)$. Clearly \overline{V}_{-1} acts by permutation of the two factors. Let $\mathbb{T}(\text{SL}_2(q))$ be an $\langle F_p \rangle$ -stable $\text{GL}_2(q)$ -transversal in $\text{Irr}_{\text{cusp}}(\text{SL}_2(q))$, see Proposition 1.19. Then $\mathbb{T}_{-1}^\circ := \mathbb{T}(\text{SL}_2(q)) \times \mathbb{T}(\text{SL}_2(q))$ is a \overline{V}_{-1} - $\langle F_p \rangle$ -stable \widetilde{L}_{-1} -transversal in $\text{Irr}_{\text{cusp}}(G_{J_{-1}})$. This proves part (a) in that case. Part (b) follows from the fact that $K_{0,-1} \rtimes \epsilon_{-1}(\overline{V}_{-1}) \cong \text{SL}_2(q) \wr C_2$, see also Lemma 3.6. Part (c) follows again from the fact that \widetilde{L}_{-1} is $(\widehat{\text{SL}}_2(q))^2$, where

$$\widehat{\text{SL}}_2(q) := \{x \in \text{SL}_2(\mathbb{F}) \mid F_q(x) = \pm x\}.$$

Assume $\text{type}(\Phi_{-1}) = A_3$. Recall $\alpha_2 = e_2 + e_1$, $\alpha_1 = e_2 - e_1$ and $\alpha_i := e_i - e_{i-1}$ ($i \geq 3$) for the simple roots in Δ . In this case $G_{J_{-1}} \cong \text{SL}_4(q)$ and $\mathbf{n}_{e_1}(\varpi)$ acts on $G_{J_{-1}}$ as a non-trivial graph automorphism. In order to see the automorphisms induced by $t_{J_{-1}}$ and $t_{J_{-1},2}$ we use again the equation $h_0 = \mathbf{h}_{\alpha_1}(-1)\mathbf{h}_{\alpha_2}(-1)$ and additionally the equation

$$\mathbf{h}_3(\varpi) = \mathbf{h}_{\alpha_1}(-\varpi)\mathbf{h}_{\alpha_2}(-\varpi)\mathbf{h}_{\alpha_3}(-1).$$

This implies that $t_{J_{-1}}$ induces on $G_{J_{-1}}$ some non-inner diagonal automorphism of $\text{SL}_4(q)$ corresponding to the central involution, while $t_{J_{-1},2}$ induces a diagonal automorphism of $\text{SL}_4(q)$ associated to a generator of the centre. Let $E(\text{SL}_4(q))$ be the subgroup of $\text{Aut}(\text{SL}_4(q))$ from 1.19. According to Proposition 1.19 there exists a $\text{GL}_4(q)$ -transversal $\mathbb{T}(\text{SL}_4(q))$ in $\text{Irr}(\text{SL}_4(q))$, that is stable under the group $E(\text{SL}_4(q))$ generated by graph and field automorphisms of $\text{SL}_4(q)$ and such that maximal extendibility holds with respect to $\text{SL}_4(q) \triangleleft \text{SL}_4(q) \rtimes E(\text{SL}_4(q))$ for $\mathbb{T}(\text{SL}_4(q))$. This choice guarantees part (b). As $\widetilde{L}_{-1}/G_{J_{-1}}$ is cyclic, part (c) holds in that case, as well. \square

Recall $\widetilde{H}_1 := \langle h_0, \mathbf{h}_{J_{-1}}(\varpi) \rangle$.

Lemma 3.9 (a) *There exists some \overline{V}_{-1} -equivariant extension map $\Lambda_{0,-1}$ with respect to $H_{-1} \triangleleft \overline{V}_{-1}$.*

(b) *If $\lambda \in \text{Irr}(H_{-1})$ with $\lambda(h_0) = -1$ and $\widetilde{\lambda} \in \text{Irr}(\widetilde{H}_{-1} \mid \lambda)$, then $(\overline{V}_{-1})_{\widetilde{\lambda}} = H_{-1}$.*

Proof. As \overline{V}_{-1}/H_{-1} is cyclic there exists an extension map as required in (a). For the proof of (b) note that the equality $[\overline{\mathbf{n}}_1, \mathbf{h}_{J_{-1}}(\varpi)] = h_0$ implies $\widetilde{\lambda}^{\overline{\mathbf{n}}_1} \neq \widetilde{\lambda}$ as $\widetilde{\lambda}(h_0) = -1$. \square

3.D Proof of Theorem B

We now finish the proof of Theorem 2.1 and therefore Theorem B. The above allows us now to verify the character-theoretic assumptions from Proposition 2.2 for the groups K , K_0 , $K_{0,d}$ and V_D , introduced in Lemma 2.11 and Theorem 2.22. From the definitions of $K_{0,d}$ before 3.4 we see $K_0 = \langle K_{0,d} \mid d \in \mathbb{D} \rangle$, even more K_0 is the central product of the groups $K_{0,d}$ ($d \in \mathbb{D}$).

By abuse of notation we write $\text{Irr}_{\text{cusp}}(K)$ for $\bigcup_{\chi \in \text{Irr}_{\text{cusp}}(L)} \text{Irr}(\chi|_K) \subseteq \text{Irr}(K)$.

Proposition 3.10 *There exists a $\overline{V} \langle F_p \rangle$ -stable \tilde{L} -transversal \mathbb{T}° in $\text{Irr}_{\text{cusp}}(K_0)$. Moreover $\mathbb{K} := \text{Irr}(K \mid \mathbb{K}_0)$ and $\mathbb{T} = \text{Irr}(L \mid \mathbb{K})$ are NE_L -stable \tilde{L} -transversals in $\text{Irr}_{\text{cusp}}(K)$ and $\text{Irr}_{\text{cusp}}(L)$, respectively.*

Note that this implies Theorem 2.1(a).

Proof. For $d \in \mathbb{D} \setminus \{1\}$ let \mathbb{T}_d° be the $\overline{V}_d \langle F_p \rangle$ -stable \tilde{L}_d -transversal in $\text{Irr}_{\text{cusp}}(K_{0,d})$ from Proposition 3.5 and Proposition 3.8. Note $K_{0,1} = 1$. The group K_0 is a central product of the groups $K_{0,d}$ ($d \in \mathbb{D}$) according to Lemma 2.13. Hence the irreducible characters of K_0 are obtained as the products of the irreducible characters of $K_{0,d}$. The central product of the characters in \mathbb{T}_d° form a subset $\mathbb{K}_0 \subseteq \text{Irr}(K_0)$. We see that \mathbb{K}_0 is $V_{\mathbb{D}}E_L$ -stable since $V_{\mathbb{D}}E_L$ and $\overline{V} \langle F_p \rangle$ act on each factor $K_{0,d}$ as $\overline{V}_d \langle F_p \rangle$. Let $\widehat{\mathbb{T}} := \mathbb{T} \cap \mathcal{L}^{-1}(\langle h_0 \rangle)$. The automorphisms of \tilde{L} induced on K_0 are induced by K_0 , $\widehat{\mathbb{T}} = \prod_{d \in \mathbb{D}} \widehat{\mathbb{T}}_d$ and $t_{L,2} = \prod_{d \in \mathbb{D}} t_{J_d,2}$, see Lemma 3.2. According to Lemma 3.3 the element $t_{J_d,2}$ acts trivially on G_d , whenever $d \geq 1$. Hence \mathbb{K}_0 is an \tilde{L} -transversal of $\text{Irr}_{\text{cusp}}(K_0)$ as well. According to Proposition 3.5 and Proposition 3.8 maximal extendibility holds with respect to $K_{0,d} \triangleleft \tilde{L}_d$. Since $[\tilde{L}_d, \tilde{L}_{d'}] = 1$ for every $d, d' \in \mathbb{D}$ with $d \neq d'$, this implies that maximal extendibility holds also with respect to $K_0 \triangleleft \tilde{L}$ as $\tilde{L} \leq \langle \tilde{L}_d \mid d \in \mathbb{D} \rangle$. Since \tilde{L}/K_0 is abelian by Lemma 2.13, \mathbb{K} and \mathbb{T} are again NE_L -stable \tilde{L} -transversals in $\text{Irr}_{\text{cusp}}(K)$ and $\text{Irr}_{\text{cusp}}(L)$, respectively. \square

We apply the following statement in order to construct some extension map with respect to $L \triangleleft N$ for $\text{Irr}_{\text{cusp}}(L)$ satisfying Equation (1.2) from Theorem 1.7.

Proposition 3.11 *There exists a $V_{\mathbb{D}} \rtimes E_L$ -equivariant extension map Λ_0 with respect to $H \triangleleft V_{\mathbb{D}}$.*

This ensures Assumption 2.2(ii.1) with the choice made in Lemma 2.11.

Proof. Recall $\overline{V} := H \langle \overline{V}_d \mid d \in \mathbb{D} \rangle$ and $V_{\mathbb{D}} := \overline{V} \cap \mathbf{G}$, see Theorem 2.22. Let $\tilde{H}_\epsilon := \langle \tilde{H}_d \mid d \in \mathbb{D}_\epsilon \rangle$. We apply the extension maps from Proposition 3.1, Lemma 3.7 and Proposition 3.8(c) for constructing a \overline{V} -equivariant extension map for $H \triangleleft V_{\mathbb{D}}$. Note that by the definition of \overline{V} , $\overline{\mathbf{n}}_1 \in \overline{V} \setminus V_{\mathbb{D}}$ whenever $\gamma \in E_L$, and then $\overline{\mathbf{n}}_1$ and γ induce the same automorphisms on \mathbf{G} according to Remark 2.4. By Theorem 2.22, $[F_p, V_{\mathbb{D}}] = 1$. Altogether it is sufficient to prove that maximal extendibility holds with respect to $H \triangleleft \overline{V}$.

Let $\lambda \in \text{Irr}(H)$ and $\tilde{\lambda} \in \text{Irr}(\tilde{H} \mid \lambda)$. The character $\tilde{\lambda}$ writes as $\odot_{d \in \mathbb{D}} \tilde{\lambda}_d$ with $\tilde{\lambda}_d \in \text{Irr}(\tilde{H}_d)$ ($d \in \mathbb{D}$). Let ψ_d be the extension of $\lambda_d := \lambda|_{H_d}$ to $(V_d)_{\lambda_d}$ given by Proposition 3.1, Lemma 3.7 and Proposition 3.8(c).

Assume $\lambda(h_0) = 1$. Let $\overline{\lambda} \in \text{Irr}(H/\langle h_0 \rangle)$ be associated with λ . It is sufficient to show that $\overline{\lambda}$ extends to $\overline{V}_{\overline{\lambda}}/\langle h_0 \rangle$. Since $[\overline{V}_d/\langle h_0 \rangle, \overline{V}_{d'}/\langle h_0 \rangle] = 1$ according to (2.2), the group $\overline{V}/\langle h_0 \rangle$ is the central product of the groups $\overline{V}_d/\langle h_0 \rangle$. The characters ψ_d ($d \in \mathbb{D}$) define extensions $\overline{\psi}_d$ of $\tilde{\lambda}_d$ to $(\overline{V}_d)_{\lambda_d}/\langle h_0 \rangle$ and $\overline{\psi} := \odot_{d \in \mathbb{D}} \overline{\psi}_d$ lifts to a character ψ° of $\langle (\overline{V}_d)_{\lambda_d} \mid d \in \mathbb{D} \rangle$. Recall $H \geq \langle H_d \mid d \in \mathbb{D} \rangle$. According to [S10b, 4.1(a)] we see that λ has an extension ψ to \overline{V}_λ such that $\psi|_{\langle V_{d,\lambda_d} \mid d \in \mathbb{D} \rangle_\lambda} = \psi^\circ|_{\langle V_{d,\lambda_d} \mid d \in \mathbb{D} \rangle_\lambda}$. The extension map with respect to $H \triangleleft V_{\mathbb{D}}$ for $\text{Irr}(H \mid 1_{\langle h_0 \rangle})$ obtained this way is then automatically $\overline{V} \rtimes \langle F_p \rangle$ -equivariant.

Assume otherwise $\lambda(h_0) = -1$. As in 2.10 let $\mathbb{D}_{\text{odd}} := \{i \in \mathbb{D} \mid i \text{ odd}\}$ and $\mathbb{D}_{\text{even}} := \{i \in \mathbb{D} \mid i \text{ even}\}$. For $\epsilon \in \{\text{odd}, \text{even}\}$ recall

$$\tilde{H}_\epsilon := \langle \tilde{H}_d \mid d \in \mathbb{D}_\epsilon \rangle, \quad H_\epsilon := \tilde{H}_\epsilon \cap H_0,$$

and $H = H_{\text{even}} \times H_{\text{odd}}$, see Lemma 2.10. Analogously we define

$$\overline{V}_\epsilon := H_\epsilon \langle \overline{V}_d \mid d \in \mathbb{D}_\epsilon \rangle \quad \text{and} \quad \tilde{V}_\epsilon := \tilde{H}_\epsilon V_\epsilon.$$

Notice that by this definition $V_{\text{even}} \leq V_D$ and hence $V_D = H(V_{\text{even}} \cdot (V_{\text{odd}} \cap V_D))$.

Let $\tilde{\lambda}_\epsilon := \tilde{\lambda} \upharpoonright_{\tilde{H}_\epsilon}$ and $\lambda_\epsilon := \lambda \upharpoonright_{H_\epsilon}$. Since $[\overline{V}_d, \overline{V}_{d'}] = 1$ for every $d \in \mathbb{D}_{\text{even}}$ and $d' \in \mathbb{D}$ by (2.2), the extensions ψ_d ($d \in \mathbb{D}_{\text{even}}$) allow us to define an extension of λ_{even} to $\overline{V}_{\text{even}, \lambda_{\text{even}}}$.

For $\epsilon = \text{even}$, H_{even} is the central product of the groups H_d ($d \in \mathbb{D}_{\text{even}}$) and $V_{\text{even}, \lambda_{\text{even}}}$ is analogously the central product of the groups $\overline{V}_{d, \lambda_d}$. Hence λ_ϵ extends to some $\tilde{\lambda}_\epsilon \in \text{Irr}(V_\epsilon, \lambda_\epsilon)$: The product of the characters ψ_d ($d \in \mathbb{D}_{\text{even}}$) defines an extension $\hat{\lambda}_{\text{even}}$ of λ_{even} to $V_{\text{even}, \lambda_{\text{even}}}$.

For extending λ_{odd} to $V_{\text{odd}, \lambda_{\text{odd}}}$ we first extend $\tilde{\lambda}_{\text{odd}} := \tilde{\lambda} \upharpoonright_{\tilde{H}_{\text{odd}}}$. Again $\tilde{\lambda}_{\text{odd}}$ is the central product of characters $\tilde{\lambda}_d$ ($d \in \mathbb{D}_{\text{odd}}$). According to Proposition 3.1(b) and Lemma 3.7(b) we have $\overline{V}_{d, \tilde{\lambda}_d} \leq V_D$. The same holds also for $d = -1$ by straight-forward calculations.

Let $\nu \in \text{Irr}(\tilde{H}_{\text{odd}})$ with $\ker(\nu) = H_{\text{odd}}$. According to Lemma 3.7(b) there exists some element $c_d \in \overline{V}_d$ such that $\overline{V}_{d, \lambda_d} = \langle \overline{V}_{d, \tilde{\lambda}_d}, c_d \rangle$, which satisfies $\tilde{\lambda}_d^{c_d} = \tilde{\lambda}_d \nu \upharpoonright_{\tilde{H}_d}$. The extensions $\psi_d \upharpoonright_{\overline{V}_{d, \tilde{\lambda}_d}}$ define also extensions ψ'_d of $\tilde{\lambda}_d$ on $\tilde{H}_d \overline{V}_{d, \tilde{\lambda}_d}$. The restriction $\psi'_d \upharpoonright_{V_{d, \tilde{\lambda}_d}}$ is then c_d -stable. Since $\overline{V}_{d, \tilde{\lambda}_d}$ is contained in V_D the group $\overline{V}_{\text{odd}, \tilde{\lambda}_{\text{odd}}}$ is the central product of the groups $\tilde{H}_d \overline{V}_{d, \tilde{\lambda}_d}$ ($d \in \mathbb{D}_{\text{odd}}$). The product $\psi' := \prod_{d \in \mathbb{D}_{\text{odd}}} \psi'_d$ determines uniquely an extension ψ'' of $\tilde{\lambda}_{\text{odd}}$ to $\tilde{H}_{\text{odd}} \overline{V}_{\text{odd}, \tilde{\lambda}_{\text{odd}}}$. Routine calculations show that $\overline{V}_{\text{odd}, \lambda_{\text{odd}}} = \overline{V}_{\text{odd}, \tilde{\lambda}_{\text{odd}}} \langle c_{\text{odd}} \rangle$ where $c_{\text{odd}} = \prod_{d \in \mathbb{D}_{\text{odd}}} c_d$. The character $\psi'' \upharpoonright_{H_{\text{odd}} \overline{V}_{\text{odd}, \tilde{\lambda}_{\text{odd}}}}$ is then c_{odd} -stable and extends to $\overline{V}_{\text{odd}, \lambda_{\text{odd}}}$. This way we obtain an extension $\hat{\lambda}_{\text{odd}}$ of λ_{odd} to $\overline{V}_{\text{odd}, \lambda_{\text{odd}}}$.

Recall $[\overline{V}_{\text{odd}}, \overline{V}_{\text{even}}] = 1$. Hence the extensions $\hat{\lambda}_{\text{odd}}$ and $\hat{\lambda}_{\text{even}}$ determine an extension of λ to \overline{V}_λ , see [S09, Lem. 4.2]. \square

In the next step we show that there exists an extension map with respect to $K_0 \triangleleft K_0 \rtimes \epsilon(V_D)$ for the set \mathbb{K}_0 from Proposition 3.10 as required in Proposition 2.2.

Proposition 3.12 *There exists a $V_D E_L$ -equivariant extension map Λ_ϵ with respect to $K_0 \triangleleft K_0 \rtimes \epsilon(V_D)$ for \mathbb{K}_0 , where $\epsilon : V_D E_L \rightarrow V_D E_L / H$ is the canonical morphism.*

Proof. By Proposition 3.5 and Proposition 3.8 there exist $\overline{V}_d \langle F_p \rangle$ -equivariant extension maps Λ_{ϵ_d} with respect to $K_{0,d} \triangleleft K_{0,d} \rtimes \epsilon(\overline{V}_d)$ for \mathbb{T}_d° , whenever $d \in \mathbb{D}$ with $d \neq 1$. Note $K_1^\circ = 1$ and hence the trivial map gives a map as required for $d = 1$. The group $K_0 \rtimes \epsilon(\overline{V})$ is the direct product of the groups $K_{0,d} \rtimes \epsilon_d(\overline{V}_d)$. Using the maps Λ_{ϵ_d} ($d \in \mathbb{D}$) we obtain an extension map Λ_ϵ as required. \square

This leads to the following statement. We use the set

$$\mathbb{K} := \text{Irr}(K \mid \mathbb{K}_0)$$

with \mathbb{K}_0 from Proposition 3.10. For the application of Proposition 2.2 we use the group $M = KV_D$, see also Corollary 2.23.

Proposition 3.13 *There exists a VE_L -equivariant extension map $\Lambda_{K \triangleleft M}$ with respect to $K \triangleleft M$ for \mathbb{K} .*

Proof. By the above all the assumptions of Proposition 2.2 are satisfied. The groups satisfy the required assumptions in 2.2(i) according to 2.23. Using the set \mathbb{K} , given as $\text{Irr}(K \mid \mathbb{K}_0)$ from Proposition 3.10 the set \mathbb{K}_0 coincides with $\bigcup_{\lambda \in \mathbb{K}} \text{Irr}(\lambda|_{K_0})$. With the $V_D E_L$ -equivariant extension map Λ_0 for $H \triangleleft V_D$ from Proposition 3.11 and the extension map Λ_ϵ for $K_0 \triangleleft K_0 \rtimes \epsilon(\overline{V})$ from Proposition 3.12 the Assumption 2.2(ii) is satisfied. The application of this statement implies the result. \square

For the set \mathbb{T} defined as $\text{Irr}(L \mid \mathbb{K})$ in Proposition 3.10 we verify that there exists an NE_L -equivariant extension map with respect to $L \triangleleft N$ for \mathbb{T} .

Proof of Theorem 2.1(b) and Theorem B. For the proof it is sufficient to construct for every $\lambda \in \mathbb{T} = \text{Irr}(L \mid \mathbb{K})$ some NE_L -stable extension of λ to N_λ . A character $\lambda \in \mathbb{T}$ lies above a unique $\lambda_0 \in \mathbb{K} = \text{Irr}(K \mid \mathbb{K}_0)$. Moreover some extension $\tilde{\lambda}_0 \in \text{Irr}(L_{\lambda_0})$ to L_{λ_0} satisfies $(\tilde{\lambda}_0)^L = \lambda$. By the properties of \mathbb{K} we see $N_{\lambda_0} = L_{\lambda_0} M_{\lambda_0}$, that is $(NE_L)_{\tilde{\lambda}_0}$ -stable. By Proposition 3.13 the character λ_0 has a $(\overline{V} \langle F_p \rangle)_{\lambda_0}$ -stable extension to M_{λ_0} . According to [S10b, 4.1] this defines an extension ϕ of $\tilde{\lambda}_0$ to $N_{\tilde{\lambda}_0}$ since $N_{\tilde{\lambda}_0} \leq L_{\lambda_0} M_{\lambda_0}$. By the construction we see that ϕ^{N_λ} is an extension of λ .

As \mathbb{T} is an M -stable \tilde{L} -transversal $(\tilde{N})_{\lambda_0} = \tilde{L}_{\lambda_0} M_{\lambda_0}$ and $(\tilde{NE}_L)_{\lambda_0} = \tilde{L}_{\lambda_0} (\widehat{M})_{\lambda_0}$. Hence this extension of λ_0 defines an extension of λ as required. \square

Later this ensures Assumption 1.12(ii).

Remark 3.14 While Theorem 2.1(b) assumes q to be odd, the proof would give a similar conclusion in the other case. For even q every $\chi \in \text{Irr}(G)$ satisfies $(\tilde{G}E)_\chi = \tilde{G}_\chi E_\chi$ since $\tilde{G} = G$ in the notation of 1.2. Nevertheless the conclusion of Theorem 2.1(b) holds as well. We observe that the arguments from before prove that there exists some NE_L -equivariant extension map $K_0 \triangleleft K_0 V_D$ for $\text{Irr}_{\text{cusp}}(L)$, where V_D is isomorphic to N/L and is defined as before with $1 = \pm \varpi$ in the argument of the Chevalley generators.

4 More on cuspidal characters

In order to prove our main theorem we need more specific results on cuspidal characters, especially with regard to automorphisms. We keep q a power of an odd prime.

Proposition 4.1 *Let $n \geq 3$, $\chi \in \text{Irr}_{\text{cusp}}(\text{GL}_n(q))$ and $\gamma \in \text{Aut}(\text{GL}_n(q))$ given by transpose-inverse up to some inner automorphism.*

- (a) *If $\chi^\gamma = \chi$, then $2 \mid n$ and $Z(\text{GL}_n(q)) \leq \ker(\chi)$.*
- (b) *If $\chi^\gamma = \chi\delta$ for $\delta \in \text{Irr}(\text{GL}_n(q))$ a linear character of multiplicative order 2, then $2 \mid n$.*

Proof. Let us recall the form of elements of $\text{Irr}_{\text{cusp}}(\text{GL}_n(q))$. We write $\mathbf{K} := \text{GL}_n(\mathbb{F})$ and $\mathbf{K}^* := \text{GL}_n(\mathbb{F})$ as the dual with \mathbb{F}_q -structures given by F . Let $s \in (\mathbf{K}^*)^F = \text{GL}_n(q)$ be such that the Lusztig series $\mathcal{E}(\mathbf{K}^F, (s))$ associated to s contains a cuspidal character. Combining [GM, 3.2.22] and the fact that the group $\mathbf{C}_{\mathbf{K}^*}(s)^F$ of type A can have cuspidal unipotent characters only when it is a torus (see for instance [GM, Ex. 2.4.20]), we get that s is regular and $\mathbf{C}_{\mathbf{K}^*}(s)$ is a Coxeter torus. This can be summed up in the fact that the spectrum of s is a single orbit of order n under F , or equivalently $\mathbb{F}_q[\zeta] = \mathbb{F}_{q^n}$ for any eigenvalue ζ of s . Concerning the action of γ , note that an element of $\mathcal{E}(\mathbf{K}^F, (s))$ is sent to an element of $\mathcal{E}(\mathbf{K}^F, (s^{-1}))$ (apply [CS13, 3.1]).

For the proof of (a) let $\chi \in \mathcal{E}(\mathbf{K}^F, (s))$ be invariant under γ . Then s and s^{-1} have the same spectrum. If 1 or -1 is an eigenvalue of s , then $s \in \{1, -1\}$ and $n = 1$ since the eigenvalues of s form a single F -orbit. This is impossible so inversion is without fixed point on the spectrum of s . This implies that n is even and that the product of the eigenvalues of s is 1. So $s \in [\mathbf{K}^*, \mathbf{K}^*]^F$ and this implies that all characters of $\mathcal{E}(\mathbf{K}^F, (s))$ have $Z(\mathrm{GL}_n(q))$ in their kernel (see [CE, 15.3]).

For the proof of part (b) note that by the assumptions q is odd and $\mathrm{SL}_n(q)$ is perfect, see [MT, 24.17]. By the correspondence induced by duality between (linear) characters of $\mathbf{K}^F/[\mathbf{K}, \mathbf{K}]^F$ and elements of $Z(\mathbf{K}^*)^F$ (see for instance [DM20, 11.4.6]) we have $\delta\mathcal{E}(\mathbf{K}^F, (s)) = \mathcal{E}(\mathbf{K}^F, (-s))$. Assuming $\chi^\gamma = \chi\delta$, the same considerations as above show that s and $-s^{-1}$ have the same eigenvalues. The spectrum of s is of the form $\{F(\zeta), F^2(\zeta), \dots, F^n(\zeta) = \zeta\}$ with $\mathbb{F}_q[\zeta] = \mathbb{F}_{q^n}$. Since s and $-s^{-1}$ have the same eigenvalues, then $-\zeta^{-1} = F^a(\zeta)$ for some $1 \leq a \leq n$. We have $F^{2a}(\zeta) = -F^a(\zeta)^{-1} = \zeta$ and therefore $\mathbb{F}_{q^{2a}} \supseteq \mathbb{F}_q[\zeta] = \mathbb{F}_{q^n}$. Then n divides $2a$. Assume now that n is odd, this implies that n divides $a \leq n$. So $a = n$ and $-\zeta^{-1} = F^n(\zeta) = \zeta$. But then $\zeta^2 = -1$ and $\mathbb{F}_q[\zeta] \subseteq \mathbb{F}_{q^2}$ which contradicts $n \geq 3$. So we get our claim that $2 \mid n$. \square

The following statement is used later for computing the relative Weyl groups associated to cuspidal characters of a Levi subgroup of $\mathrm{D}_{l,\mathrm{sc}}(q)$.

Proposition 4.2 *Let $n \geq 2$, $\psi \in \mathrm{Irr}_{\mathrm{cusp}}(\mathrm{SL}_n(q))$ and $\gamma \in \mathrm{Aut}(\mathrm{GL}_n(q))$ given by transpose-inverse up to some inner automorphism. If $|\mathrm{GL}_n(q) : \mathrm{GL}_n(q)_\psi|$ is even and $\psi^\gamma = \psi$, then $n = 2$ and ψ is one of the two characters $\mathrm{R}'_\sigma(\theta_0)$ ($\sigma \in \{\pm 1\}$) of degree $\frac{q-1}{2}$ from Table 5.4 of [B].*

Proof. According to Table 5.4 of [B] the two characters $\mathrm{R}'_\sigma(\theta_0)$ ($\sigma \in \{\pm 1\}$) are the only characters of $\mathrm{SL}_2(q)$ that are cuspidal and not $\mathrm{GL}_2(q)$ -stable. The characters $\mathrm{R}(\theta)$ given there are $\mathrm{GL}_2(q)$ -stable and the other characters $\mathrm{R}_\sigma(\alpha_0)$ ($\sigma \in \{\pm 1\}$) are in the principal Harish-Chandra series. Note that γ then restricts to an interior automorphism of $\mathrm{SL}_2(q)$.

Now consider $n \geq 4$. Let ψ be as in the proposition, let $\chi \in \mathrm{Irr}(\mathrm{GL}_n(q) \mid \psi)$, so that χ is cuspidal thanks to Lemma 1.15(c). We keep the notation of the proof of Proposition 4.1 with $\chi \in \mathcal{E}(\mathrm{GL}_n(q), (s))$ and ζ some eigenvalue of s .

By Clifford theory χ is induced from a character of $\mathrm{GL}_n(q)_\psi$. Then the assumption $2 \mid |\mathrm{GL}_n(q) : \mathrm{GL}_n(q)_\psi|$ implies $\chi = \nu_1\chi$ for $\nu_1 \in \mathrm{Irr}(\mathrm{GL}_n(q))$ the linear character of order 2 with kernel containing $\mathrm{SL}_n(q)$. Hence s is $\mathrm{GL}_n(q)$ -conjugate to $-s$. Then $-\zeta \in \{F(\zeta), F^2(\zeta), \dots, F^n(\zeta) = \zeta\}$ since this is the spectrum of s .

Clifford theory also tells us that the assumption $\psi^\gamma = \psi$ implies $\chi^\gamma = \nu_2\chi$ for some linear character ν_2 of $\mathrm{GL}_n(q)$ with $\mathrm{SL}_n(q)$ in its kernel. Then s^{-1} is conjugate to λs for some $\lambda \in \mathbb{F}_q^\times$. As before we obtain $\zeta^{-1} \in \{\lambda F(\zeta), \lambda F^2(\zeta), \dots, \lambda F^n(\zeta) = \lambda\zeta\}$.

We can now write $-\zeta = F^a(\zeta)$ and $\lambda\zeta^{-1} = F^b(\zeta)$ for $1 \leq a, b \leq n$. The first equality gives $F^{2a}(\zeta) = -F^a(\zeta) = \zeta$ and the second $F^{2b}(\zeta) = \lambda F^b(\zeta)^{-1} = \zeta$ since $\lambda \in \mathbb{F}_q$. So $\zeta \in \mathbb{F}_{q^{2a}} \cap \mathbb{F}_{q^{2b}}$, but since $\mathbb{F}_q[\zeta] = \mathbb{F}_{q^n}$ we get that n divides both $2a$ and $2b$. The latter are at most $2n$, so $2a, 2b \in \{n, 2n\}$. Having $a = n$ would imply $-\zeta = F^n(\zeta) = \zeta$ which is impossible because q is odd. So n is even and $a = \frac{n}{2}$. On the other hand, if $b = n$ then $\zeta = F^n(\zeta) = \lambda\zeta^{-1}$ and therefore $\zeta^2 \in \mathbb{F}_q$. Then $\mathbb{F}_q[\zeta] \subseteq \mathbb{F}_{q^2}$ and this gives our claim that $n = 2$.

There remains the case when $b = \frac{n}{2} = a$. Then $\lambda\zeta^{-1} = F^a(\zeta) = -\zeta$ and again $\zeta^2 \in \mathbb{F}_q$. This yields $n = 2$ as seen before. \square

We complement the above by a result on cuspidal characters of $\mathrm{D}_{l,\mathrm{sc}}(q)$, which follows from a combination of results of Malle and Maslowski. We use \mathbf{G} , F , γ from

Notation 2.3 and h_0 from Notation 2.9. Recall the Lang map \mathcal{L} defined on \mathbf{G} by $\mathcal{L}(g) = g^{-1}F(g)$.

Proposition 4.3 *Let $\tilde{G} := \mathcal{L}^{-1}(Z(\mathbf{G}))$. If $\lambda \in \text{Irr}_{\text{cusp}}(\mathbf{G}^F | 1_{\langle h_0 \rangle})$ with $\tilde{G}_\lambda \leq \mathcal{L}^{-1}(\langle h_0 \rangle)$, then γ acts trivially on λ and $\text{Irr}(\mathcal{L}^{-1}(\langle h_0 \rangle) | \lambda)$.*

Proof. Recall that a character of $\tilde{\mathbf{G}}^F$ is called semi-simple when it corresponds to a trivial unipotent character through the Jordan decomposition of characters. The components of their restrictions to \mathbf{G}^F are also called the semi-simple characters of \mathbf{G}^F . In particular both are of degree prime to p (see [GM, 2.6.11]).

According to [M17, Thm. 1] there exists a semi-simple character $\rho \in \text{Irr}(\mathbf{G}^F)$ with $(\tilde{G}E)_\rho = (\tilde{G}E)_\lambda$, where ρ and λ lie in the same rational Lusztig series. We use now results from [S12] to investigate ρ further. In a first step we prove that γ acts trivially on ρ and $\text{Irr}(\mathcal{L}^{-1}(\langle h_0 \rangle) | \rho)$.

We assume that \mathbf{G} , \mathbf{T} and Δ are given as in 2.3 and let $\mathbf{U} := \langle \mathbf{X}_\alpha | \alpha \in \Delta \rangle$ and $\mathbf{B} := \mathbf{T}\mathbf{U}$. As group $\tilde{\mathbf{G}}$ introduced in 1.2 we use the particular choice from [S12, 3.1]. Then $\tilde{\mathbf{G}}^F$ and \tilde{G} induce the same automorphisms on $G := \mathbf{G}^F$. Let $\tilde{\mathbf{B}} := \mathbf{B}Z(\tilde{\mathbf{G}})$. Let $\tilde{\Omega} : \text{Irr}_{p'}(\tilde{\mathbf{G}}^F) \rightarrow \text{Irr}_{p'}(\tilde{\mathbf{B}}^F)$ be the $\text{Irr}(\tilde{\mathbf{G}}^F/\mathbf{G}^F) \times E(\mathbf{G}^F)$ -equivariant bijection with $\text{Irr}(\tilde{\psi}|_{Z(\tilde{\mathbf{G}}^F)}) = \text{Irr}(\tilde{\Omega}(\tilde{\psi})|_{Z(\tilde{\mathbf{B}}^F)})$ for every $\tilde{\psi} \in \text{Irr}_{p'}(\tilde{\mathbf{G}}^F)$ from [S12, 3.3(a)].

Let $\tilde{\rho} \in \text{Irr}(\tilde{\mathbf{B}}^F | \rho)$, $\tilde{\phi} := \tilde{\Omega}(\tilde{\rho})$ and $\phi \in \text{Irr}(\tilde{\phi}|_{\mathbf{B}^F})$. Let Δ be the basis of the root system of \mathbf{G} given by \mathbf{B} , C the Cartan matrix associated to Δ and $C^{-1} = (c'_{\alpha\beta})$ its inverse. Let $\zeta \in \mathbb{F}^\times$ a root of unity of order $\det(C)(q-1)$. For $\alpha \in \Delta$ we set

$$t_\alpha^{(0)} := \prod_{\beta \in \Delta} \mathbf{h}_\beta(\zeta^{\det(C)c'_{\alpha\beta}}),$$

see also [Ma10, 8.1]. Then we can choose elements $t_\alpha \in t_\alpha^{(0)} Z(\tilde{\mathbf{G}}) \cap \tilde{\mathbf{T}}^F$ such that $\tilde{\mathbf{T}}^F = Z(\tilde{\mathbf{G}}^F) \langle t_\alpha | \alpha \in \Delta \rangle$, see [Ma10, §8]. Assume that Δ is given as in 2.3 and let $\alpha \in \{e_2 \pm e_1\}$. The entries of C^{-1} can be found in [OV, p. 296]. We see $\mathcal{L}(t_\alpha^{(0)}) = (t_\alpha^{(0)})^{q-1} \in \mathbf{h}_\alpha(\varpi) \langle h_0 \rangle$. Hence t_α induces a diagonal automorphism of \mathbf{G}^F associated to some element in $\mathbf{h}_\alpha(\varpi) \langle h_0 \rangle$ in the notation of Remark 1.16(b).

We abbreviate $\tilde{G} := \mathcal{L}^{-1}(\langle h_0 \rangle)$. (Note that \tilde{G} then differs from the group introduced in 1.16.) The assumption $\tilde{G}_\lambda \leq \tilde{G}$ implies $\lambda^{t_\alpha} \neq \lambda$. Via the construction we have $(\tilde{G}E)_\rho = (\tilde{G}E)_\lambda$ and hence $\rho^{t_\alpha} \neq \rho$. By Clifford theory the character $\tilde{\rho}$ satisfies $\tilde{\rho}(t_\alpha \mathbf{G}^F) = 0$ and is stable under multiplying with linear characters with kernel

$$\{g \in \tilde{\mathbf{G}}^F | g \text{ induces diagonal autom. of } \mathbf{G}^F \text{ associated to } \langle h_0 \rangle\}.$$

As $\tilde{\Omega}$ is an $\text{Irr}(\tilde{\mathbf{G}}^F/\mathbf{G}^F)$ -equivariant bijection, the character $\tilde{\phi}$ has to satisfy $\tilde{\phi}(t_\alpha) = 0$ as well. As in Remark 1.16 the character $\tilde{\phi}$ can be extended to some κ on $\tilde{\mathbf{B}}^F \tilde{Z} = \tilde{B}_0 \tilde{Z}$, where $\tilde{\mathcal{L}}$ is the Lang map on $\tilde{\mathbf{G}}$, $\tilde{Z} := \tilde{\mathcal{L}}^{-1}(Z(\mathbf{G})) \cap Z(\tilde{\mathbf{G}})$ and $\tilde{B}_0 := \mathcal{L}^{-1}(Z(\mathbf{G})) \cap \mathbf{B}$. Note that $\tilde{B}_0 = Z(\mathbf{G}^F) \langle t_\beta^{(0)} | \beta \in \Delta \rangle \mathbf{U}^F$. Then $\kappa(t_\alpha^{(0)}) = 0$. The character $\kappa|_{\tilde{B}_0}$ is γ -stable, since $\text{Irr}(\kappa|_{Z(\mathbf{G}^F)}) = \text{Irr}(\phi|_{Z(\mathbf{G}^F)})$ is γ -stable because of $h_0 \in \ker(\phi)$ and $t_\beta^{(0)}$ is γ -fixed for every $\beta \in \Delta \setminus \{e_2 \pm e_1\}$ according to the explicit value of C^{-1} .

As κ is γ -stable, some $\phi \in \text{Irr}(\kappa|_{\mathbf{B}^F}) = \text{Irr}(\tilde{\phi}|_{\mathbf{B}^F})$ is γ -stable, see [S12, 3.6(a)]. By Clifford theory $\kappa|_{\tilde{B}_0}$ is of the form $\hat{\phi}^{\tilde{B}_0}$ for a unique $\hat{\phi} \in \text{Irr}(\tilde{B}_0, \phi)$. As ϕ extends to $\tilde{\mathbf{B}}_\phi^F$ according to [S12, Thm. 3.5(a)], the character $\hat{\phi}$ is an extension of ϕ . As κ and ϕ are γ -stable, $\hat{\phi}$ is γ -stable. Note that $\tilde{B}_0, \phi \leq \tilde{B} := \mathcal{L}^{-1}(\langle h_0 \rangle) \cap \mathbf{B}$.

Via the statement given in Lemma 1.18 some \tilde{G} -conjugate ρ' of ρ is γ -stable and has also a γ -stable extension to \hat{G} . If $\rho' \neq \rho$ we observe that $(\hat{G}\langle\gamma\rangle)^{t'} = \hat{G}\langle\gamma\rangle$ for every $t' \in \mathcal{L}^{-1}(\mathbf{h}_2(\varpi))$. Hence the character ρ extends to $\hat{G}\langle\gamma\rangle$, as well.

We deduce from this result on ρ the analogous property of λ . Recall that by definition λ and ρ are in the same rational Lusztig series and $(\tilde{\mathbf{G}}^F E)_\rho = (\tilde{\mathbf{G}}^F E)_\lambda$, in particular $\tilde{\mathbf{G}}_\rho^F = \tilde{\mathbf{G}}_\lambda^F$. Recall $\tilde{\rho} \in \text{Irr}(\tilde{\mathbf{G}}^F | \rho)$ and $\text{Irr}(\tilde{\mathbf{G}}^F / \mathbf{G}^F)$ acts on $\text{Irr}(\tilde{\mathbf{G}}^F)$ acts by multiplication with linear characters. As $\tilde{\mathbf{G}}^F / \mathbf{G}^F$ is abelian and maximal extendibility holds with respect to $\mathbf{G}^F \triangleleft \tilde{\mathbf{G}}^F$ we see

$$\text{Irr}(\tilde{\mathbf{G}}^F / G)_{\tilde{\lambda}} = \text{Irr}(\tilde{\mathbf{G}}^F / \tilde{\mathbf{G}}_\lambda^F) = \text{Irr}(\tilde{\mathbf{G}}^F / \tilde{\mathbf{G}}_\rho^F) = \text{Irr}(\tilde{\mathbf{G}}^F / G)_{\tilde{\rho}}.$$

Let $\mathcal{E}(\tilde{\mathbf{G}}^F, s)$ be the rational Lusztig series containing $\tilde{\rho}$. The character $\tilde{\rho}$ is semi-simple. The series $\mathcal{E}(\tilde{\mathbf{G}}^F, s)$ contains exactly one regular character, see [DM20, 12.4.10]. By the definition of semi-simple and regular in [DM20, 12.4.1] we see that there exists also a unique regular character in that series. Let $\tilde{\rho}' \in \text{Irr}(\tilde{\mathbf{G}}^F)$ be the Alvis-Curtis dual of $\tilde{\rho}$ up to a sign, see [DM20, 7.2]. Then $\{\tilde{\rho}'\} = \text{Irr}(\Gamma(\tilde{\mathbf{G}}^F)) \cap \mathcal{E}(\tilde{\mathbf{G}}^F, s)$, where $\Gamma(\tilde{\mathbf{G}}^F)$ denotes the Gelfand-Graev character of $\tilde{\mathbf{G}}^F$. Since it vanishes outside unipotent elements, the Gelfand-Graev character is stable under $\text{Irr}(\tilde{\mathbf{G}}^F / \mathbf{G}^F)$. Hence $\text{Irr}(\tilde{\mathbf{G}}^F / \mathbf{G}^F)_{\tilde{\rho}'}$ coincides with the stabiliser of $\mathcal{E}(\tilde{\mathbf{G}}^F, s)$ in $\text{Irr}(\tilde{\mathbf{G}}^F / \mathbf{G}^F)$. This group is called $B(s)$ in [CE, 15.13]. By the construction of Alvis-Curtis duality this implies $\text{Irr}(\tilde{\mathbf{G}}^F / G)_{\tilde{\rho}} = B(s)$. The characters $\tilde{\rho}^\gamma$ and $\tilde{\lambda}^\gamma$ belong to $\mathcal{E}(\tilde{\mathbf{G}}^F, \gamma^{-1}(s))$. As λ and ρ are γ -stable, $\tilde{\rho}^\gamma = \tilde{\rho}\mu$ and $\tilde{\lambda}^\gamma = \tilde{\lambda}\mu'$ for linear characters $\mu, \mu' \in \text{Irr}(\tilde{\mathbf{G}}^F)$. Since $\tilde{\rho}^\gamma$ and $\tilde{\lambda}^\gamma$ are in the same rational series, $\mu \in \mu' B(s)$ or equivalently $\mu \upharpoonright_{\tilde{\mathbf{G}}_\rho^F} = \mu' \upharpoonright_{\tilde{\mathbf{G}}_\lambda^F}$.

Because of $\text{Irr}(\tilde{\rho} \upharpoonright_{Z(\tilde{\mathbf{G}}^F)}) = \text{Irr}(\tilde{\lambda} \upharpoonright_{Z(\tilde{\mathbf{G}}^F)})$, Lemma 1.18 allows to conclude that λ has a γ -stable extension to \hat{G} as ρ has such an extension. \square

5 Character theory for the relative inertia groups $W(\lambda)$

The aim of this section is to ensure Assumption 1.8(ii), namely to prove (a main step towards) the following statement.

Theorem 5.1 *Let $l \geq 4$. Let $G = \mathbf{G}^F = D_{l,sc}(q)$ with odd q and $L = \mathbf{L}^F$ a standard Levi subgroup of G (see Notation 2.3). Let $N, \tilde{N}' := \tilde{T}_0 N$ and $E_L := \text{Stab}_{E(\mathbf{G}^F)}(\mathbf{L}^F)$ be associated to L as in Theorem 1.8. If Hypothesis 1.14 holds for every l' with $4 \leq l' < l$, then there exists some E_L -stable \tilde{N}' -transversal in $\text{Irr}_{\text{cusp}}(N)$.*

Some technical difficulties (mainly in the case where $\mathbf{G} = D_{4,sc}(\mathbb{F})$) delay the complete proof until Section 6. We construct the E_L -stable \tilde{N}' -transversal as subset of $\text{Irr}(N | \mathbb{T})$, where \mathbb{T} is the \hat{N} -stable \tilde{N}' -transversal from Theorem 2.1(a). In Lemma 5.3 we find some E_L -stable \tilde{N}' -transversal of $\text{Irr}(N | \{\lambda \in \text{Irr}_{\text{cusp}}(L) \mid \tilde{L}'_\lambda = L\})$.

In order to find the transversal of $\text{Irr}(N | \{\lambda \in \text{Irr}_{\text{cusp}}(L) \mid \tilde{L}'_\lambda \neq L\})$ with the required properties we apply the strategy mapped out by Proposition 1.12, itself based on the parametrization of Proposition 1.10. Thanks to Theorem 2.1 the two first assumptions of Proposition 1.12 can be assumed, in particular there exist some \hat{N} -equivariant extension map $\Lambda_{L, \mathbb{T}}$ with respect to $L \triangleleft N$ for \mathbb{T} , where $\hat{N} = N E_L$. We have to ensure the remaining Assumption 1.12(iii) and study the characters of the relative Weyl groups and their Clifford theory.

As already discussed in 1.C characters in such a transversal have a stabilizer in $\tilde{N}'E_L$ with a specific structure, namely such a $\psi \in \text{Irr}(N)$ satisfies

$$(\widehat{N\tilde{L}'})_\psi = \widehat{N}_\psi \tilde{L}'_\psi, \quad (5.1)$$

see also Lemma 1.4. For studying a character $\psi \in \text{Irr}(N | \mathbb{T})$ we apply the parametrization Υ from Proposition 1.10(a) and the extension map $\Lambda_{L,\mathbb{T}}$ with respect to $L \triangleleft N$ for \mathbb{T} . Then $\psi = \Upsilon(\overline{(\lambda, \eta)}) = (\Lambda_{L,\mathbb{T}}(\lambda)\eta)^N$ with $\lambda \in \mathbb{T}$ and $\eta \in \text{Irr}(W(\lambda))$. According to Proposition 1.11 the character $\psi = \Upsilon(\overline{(\lambda, \eta)})$ satisfies Equation (5.1) if

$$\eta \text{ is } \widehat{K}(\lambda)_{\eta_0}\text{-stable, where } \tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda | \lambda) \text{ and } \eta_0 \in \text{Irr}(W(\tilde{\lambda})),$$

where $\widehat{W} = NE_L/L$ and $\widehat{K}(\lambda) = \widehat{W}_\lambda$. The aim of this section is Corollary 5.36, namely to prove that for every $\lambda \in \mathbb{T}$, $\tilde{\lambda} \in \text{Irr}(\tilde{L}'_\lambda | \lambda)$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$

$$\text{there exists some } K(\lambda)_{\eta_0}\text{-stable } \eta \in \text{Irr}(W(\lambda) | \eta_0),$$

where $K(\lambda)$ is the group from 5.4. According to Lemma 5.5 such a character η is also $\widehat{K}(\lambda)_{\eta_0}$ -stable, whenever \mathbf{G} is not of type D_4 .

In the proof some arguments depend on the group \tilde{L}'_λ . As in Remark 1.16 we relate the group \tilde{L}' to subgroups of \mathbf{G} .

Notation 5.2 Recall the definitions $\tilde{L} := \mathcal{L}^{-1}(Z(\mathbf{G})) \cap \mathbf{L}$ and $\widehat{L} := \mathcal{L}^{-1}(\langle h_0 \rangle) \cap \mathbf{L}$ from Lemma 2.13, where $\mathcal{L} : \mathbf{G} \rightarrow \mathbf{G}$ is given by $x \mapsto x^{-1}F(x)$. Recall $\tilde{N}' := \tilde{T}_0N$ and set analogously $\tilde{N} := \tilde{L}N$. Then \tilde{L}' and \tilde{N}' induce on \mathbf{G} the same automorphisms as \tilde{L} and \tilde{N} , respectively.

Note that by an application of Lang's theorem $\mathcal{L}(\mathbf{L}) = \mathbf{L} \supseteq Z(\mathbf{G})$ so that $L \leq \widehat{L} \leq \tilde{L}$.

For $\lambda \in \mathbb{T}$ the characters of $W(\lambda)$ and $W(\tilde{\lambda})$ defined as above will be investigated depending on the value of \tilde{L}_λ . The set $\text{Irr}_{\text{cusp}}(L)$ can be partitioned in the following way:

$$\text{Irr}_{\text{cusp}}(L) = \mathbb{M}^{(L)} \sqcup \mathbb{M}^{(\widehat{L})} \sqcup \mathbb{M}^{(\tilde{L})} \sqcup \mathbb{M}_0,$$

where $\mathbb{M}^{(X)} := \{\lambda \in \text{Irr}_{\text{cusp}}(L) \mid \tilde{L}_\lambda = X\}$ for any subgroup $L \leq X \leq \tilde{L}$ and $\mathbb{M}_0 := \text{Irr}_{\text{cusp}}(L) \setminus (\mathbb{M}^{(L)} \cup \mathbb{M}^{(\widehat{L})} \cup \mathbb{M}^{(\tilde{L})})$. (In case of $|Z(\mathbf{G})| = 2$ one has $\mathbb{M}^{(L)} = \emptyset$.) Note that the sets are by definition $\tilde{N}E_L$ -stable as L , \widehat{L} and \tilde{L} are $\tilde{N}E_L$ -stable. In the following we construct an E_L -stable \tilde{N} -transversal in $\text{Irr}_{\text{cusp}}(N | \mathbb{M}')$ for those given NE_L -stable subsets $\mathbb{M}' \subseteq \text{Irr}_{\text{cusp}}(L)$.

Lemma 5.3 Let $\mathbb{T}^{(L)} := \mathbb{T} \cap \mathbb{M}^{(L)}$. Then $\text{Irr}(N | \mathbb{T}^{(L)})$ is an E_L -stable \tilde{N}' -transversal in $\text{Irr}(N | \mathbb{M}^{(L)})$.

Proof. The set \mathbb{T} is by construction NE_L -stable and no two elements are \tilde{L} -conjugate. Hence for $\lambda \in \mathbb{T}^{(L)}$ we have $(NE_L\tilde{L})_\lambda = (NE_L)_\lambda$ by Lemma 1.4. By Clifford theory $\text{Irr}(N | \mathbb{T}^{(L)})$ is an \tilde{N} -transversal in $\text{Irr}(L | \mathbb{M}^{(L)})$ and is NE_L -stable. \square

Determining an NE_L -stable \tilde{L} -transversal in $\text{Irr}(N | \mathbb{M}')$ for the other sets \mathbb{M}' is more involved. We start by some general descriptions of $W(\lambda)$ and related groups for $\lambda \in \text{Irr}_{\text{cusp}}(L)$, see Proposition 5.28. Afterwards we collect some particular results on cuspidal characters. In the following two subsections we verify for characters of $W(\lambda)$ the above condition under the assumption that $\lambda \in \mathbb{M}^{(\tilde{L})} \cup \mathbb{M}_0$ or $\lambda \in \mathbb{M}^{(\widehat{L})}$.

In 5.D we ensure a closely related condition on characters of $W(\lambda)$ for $\lambda \in \text{Irr}_{\text{cusp}}(L)$ with $\tilde{L}_\lambda = \widehat{L}$. In Section 6 we show how these considerations prove Theorem 5.1 and how this implies Theorem A.

5.A Understanding $\text{Irr}_{\text{cusp}}(N)$ via characters of subgroups of W

We start by recalling some basic notation and introducing subgroups of $\widehat{W} := \widehat{N}/L = NE_L/L$ as in 1.8. Additionally let $\overline{N} := N_{\mathbf{G}^F}(\mathbf{L})$ and $\overline{W} = \overline{N}/L$, see also Proposition 2.16.

Notation 5.4 Let \mathbf{G} and $F : \mathbf{G} \rightarrow \mathbf{G}$ be as in 2.3 with odd q . Let \mathbf{L} be the standard Levi subgroup of (\mathbf{G}, F) such that $L := \mathbf{L}^F$. For any J with $L \leq J \leq \widetilde{L}$ and $\lambda \in \mathbb{Z}\text{Irr}(J)$ we set $W(\lambda) := N_\lambda/L$. If additionally J is E_L -stable, \overline{W} acts on $\text{Char}(J)$, hence we can define $\overline{W}(\lambda) := \overline{W}_\lambda$ and $K(\lambda) := \overline{W}_{\lambda^{J\langle F_p \rangle}} \Big|_J$.

The groups $K(\lambda)$ and $\widehat{K}(\lambda)$ are strongly related, in particular by the following result it is sufficient to consider $K(\lambda)$ instead of $\widehat{K}(\lambda)$ if \mathbf{G} is not of type D_4 . Recall that γ is the graph automorphism of \mathbf{G} of order 2 swapping α_1 and α_2 .

Lemma 5.5 Let $E^\circ := \langle F_p, \gamma \rangle$, $E_L^\circ := E^\circ \cap E_L$, $\lambda \in \mathbb{T}$, $\widetilde{\lambda}$ defined as above, $\eta_0 \in \text{Irr}(W(\widetilde{\lambda}))$ and $\eta \in \text{Irr}(W(\lambda))$. Then η is $\widehat{K}(\lambda)_{\eta_0} \cap (W \rtimes E_L^\circ)$ -stable if and only if it is $K(\lambda)_{\eta_0}$ -stable.

Proof. Note $F_p \in Z(\widehat{W})$, hence η and η_0 are F_p -stable. Recall \overline{W} can be identified with the quotient $(\widehat{W} \cap (W \rtimes E_L^\circ)) / \langle F_p \rangle$. The group $\widehat{K}(\lambda) \cap (W \rtimes E_L^\circ)$ then projects to $K(\lambda)$, i.e. for every $w \in \overline{W}$ and $e \in \langle F_p \rangle$ with $\lambda^{we} = \lambda$ we see $(\lambda^{LE_L^\circ} \Big|_L)^w = \lambda$. This implies $\widehat{K}(\lambda) \langle F_p \rangle \cap \overline{W} = K(\lambda)$ and $\widehat{K}(\lambda)_{\eta_0} \langle F_p \rangle \cap \overline{W} = K(\lambda)_{\eta_0}$. As F_p stabilizes η this implies the statement. \square

For $\lambda \in \mathbb{T}$ and $\widetilde{\lambda} \in \text{Irr}(\widetilde{L}_\lambda \mid \lambda)$ the group $W(\widetilde{\lambda})$ is determined by $\widetilde{\lambda} \Big|_L$, as W acts trivially on the characters of \widetilde{L}_λ/L . Note that $W(\widetilde{\lambda}) \neq W(\widetilde{\lambda} \Big|_L)$ in general. We can work with the group \widetilde{L} instead of \widetilde{L}' because of the following observation.

Lemma 5.6 Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$, $\widehat{\lambda} \in \text{Irr}(\widehat{L} \mid \lambda)$ and $\widetilde{\lambda} \in \text{Irr}(\widetilde{L} \mid \widehat{\lambda})$.

- (a) Then $\overline{W}(\widetilde{\lambda}) = \overline{W}(\widehat{\lambda})$ and $W(\widetilde{\lambda}) = W(\widehat{\lambda})$ for every $\widetilde{\lambda}' \in \text{Irr}(\widetilde{L}' \mid \lambda)$.
- (b) Then $\overline{W}(\widetilde{\lambda}) \leq \overline{W}(\widehat{\lambda}) \leq \overline{W}(\lambda)$ and $W(\widetilde{\lambda}) \leq W(\widehat{\lambda}) \leq W(\lambda)$.

Proof. By the construction of \mathbb{T} , the character $\lambda \in \mathbb{T}$ satisfies $(\widehat{N}\widehat{L})_\lambda = \widehat{N}_\lambda \widehat{L}_\lambda$. Because of $\widetilde{\mathbf{G}} = Z(\mathbf{G})\mathbf{G}$ the group \widetilde{L}' is a subgroup of $\widetilde{L}Z(\widetilde{\mathbf{G}})$ and \widetilde{L} is its projection of \widetilde{L}' on \mathbf{L} . This implies $(NE_L)_{\widetilde{\lambda}} = (NE_L)_{\widetilde{\lambda}'}$, and hence part (a), see Remark 1.16 for a similar argument.

As h_0 is centralized by \widehat{N} , the group \widehat{L} is normalized by N and \overline{N} . The equalities from part (b) follow from this by straightforward considerations. \square

For $\lambda \in \mathbb{T}$ and $\widetilde{\lambda} \in \text{Irr}(\widetilde{L} \mid \lambda)$ we compute $\overline{W}(\widetilde{\lambda})$ as an approximation of $W(\widetilde{\lambda})$. For $I \in \mathcal{O}$ and $d \in \mathbb{D}$ we use the groups \mathbf{G}_I , L_I , \widehat{L}_I and \mathbf{L}_d from Notation 2.9, Lemma 2.12 and Lemma 2.13. In Lemma 2.13 the structure of \mathbf{L} and some of its subgroups was already studied. Additionally we use the following properties of \widehat{L} .

Proposition 5.7 — The structure of \widehat{L} . For $d \in \mathbb{D}$ and $I \in \mathcal{O}$ let $\widehat{L}_d := \widehat{L} \cap \mathbf{L}_d$.

- (a) L_1 is a split torus of rank $|J_1|$.
- (b) \widehat{L} is the central product of \widehat{L}_d ($d \in \mathbb{D}$) over $\langle h_0 \rangle$.
- (c) \widehat{L}_d is the central product of \widehat{L}_I ($I \in \mathcal{O}_d$) over $\langle h_0 \rangle$.
- (d) $[\widehat{L}_I, \widehat{L}_{I'}] = 1$ for all $I, I' \in \mathcal{O}$ with $I \neq I'$.

Proof. The first three parts follow from Lemma 2.13(c).

Part (d) is clear if $I \in \mathcal{O}_1$ or $I' \in \mathcal{O}_1$. Note that the groups \widehat{L}_I and $\widehat{L}_{I'}$ contain the root subgroups for Φ_I and $\Phi_{I'}$, that are orthogonal to each other. At least one of them is of type A_I . Hence no non-trivial linear combination of roots from Φ_I and $\Phi_{I'}$ is a root itself. Hence by Chevalley's commutator formula we see that the commutator of the groups is trivial. \square

We continue using the groups \overline{V}_d from 2.21 for the description of $\overline{W}(\widehat{\lambda})$. We write $\text{Irr}_{\text{cusp}}(\widehat{L})$ for $\text{Irr}(\widehat{L} \mid \text{Irr}_{\text{cusp}}(L))$.

Lemma 5.8 — Characters of \widehat{L} . *Let $\widehat{\lambda} \in \text{Irr}_{\text{cusp}}(\widehat{L})$, $\widehat{\lambda}_d \in \text{Irr}(\widehat{\lambda} \upharpoonright_{\widehat{L}_d})$ for every $d \in \mathbb{D}$ and $\widehat{\lambda}_I \in \text{Irr}(\widehat{\lambda} \upharpoonright_{\widehat{L}_I})$ for every $I \in \mathcal{O}$. Then:*

- (a) $\widehat{\lambda} = \odot_{d \in \mathbb{D}} \widehat{\lambda}_d$ and $\widehat{\lambda}_d = \odot_{I \in \mathcal{O}_d} \widehat{\lambda}_I$ for every $d \in \mathbb{D}$,
- (b) $\widehat{\lambda}_d \in \text{Irr}_{\text{cusp}}(\widehat{L}_d)$ and $\widehat{\lambda}_I \in \text{Irr}_{\text{cusp}}(\widehat{L}_I)$,
- (c) $\overline{W}(\widehat{\lambda})$ is the direct product of the groups $\overline{W}_d(\widehat{\lambda}) := \overline{V}_{d, \widehat{\lambda}}/H_d$ ($d \in \mathbb{D}$), and
- (d) $\overline{V}_{d, \widehat{\lambda}}/H_d = \overline{V}_{d, \widehat{\lambda}_d}/H_d$.

Proof. The description of $\widehat{\lambda}$ and $\widehat{\lambda}_d$ in (a) follows from the structure of \widehat{L} and \widehat{L}_d given in Proposition 5.7. The characters $\widehat{\lambda}_d$ and $\widehat{\lambda}_I$ cover a cuspidal character of L_d and L_I , respectively (see Lemma 1.15). Considering the roots underlying V_d and $\mathbf{L}_{d'}$ we see that the Chevalley relations imply $[V_d, \mathbf{L}_{d'}] = 1$ for $d, d' \in \mathbb{D}$ with $d \neq d'$. This implies the parts (c) and (d). \square

For a more explicit description of the groups $\overline{W}(\widehat{\lambda})$ we introduce some elements of \overline{V} using the maps κ_d ($d \in \mathbb{D}$) from Lemma 2.21. For $d \in \mathbb{D} \setminus \{-1\}$, recall $\mathcal{O}_d = \{I_{d,1}, \dots, I_{d,a_d}\}$ from Notation 2.17.

Notation 5.9 Let $d \in \mathbb{D} \setminus \{-1\}$ and $c_{I_{d,j}} := \kappa_d(\mathbf{n}_{e_j}(\varpi)) \in \overline{V}_d$ for every $j \in \underline{a_d}$. Note that for every $I \in \mathcal{O}_d$, c_I is some \overline{V}_d -conjugate of $\overline{\mathbf{n}}_1^{(d)}$ and $\rho_{\mathbf{T}}(c_I) = \prod_{i \in I} (i, -i)$, where $\rho_{\mathbf{T}} : \overline{N}_0 \rightarrow \mathcal{S}_{\pm 1}$ is the natural epimorphism, see before Proposition 2.16. If $2 \nmid |I|$ and $I \notin \mathcal{O}_{-1} \cup \mathcal{O}_1$, then by considerations as in the proof of 3.4(b), c_I acts as by transpose-inverting on L_I via the identification of L_I with $\text{GL}_{|I|}(q)$.

We define additionally the subgroups

$$\overline{V}_{d,S} := H_d \langle \kappa_d(\mathbf{n}_{e_i - e_{i+1}}(-1)) \mid i \in \underline{a_d - 1} \rangle \quad (5.2)$$

and $V_S := \langle V_{d,S} \mid d \in \mathbb{D} \setminus \{-1\} \rangle$. Then $\rho_{\mathbf{T}}(V_S(L \cap N_0))/\rho_{\mathbf{T}}(L \cap N_0) = \mathcal{S}_{\mathcal{O}} \leq \mathcal{S}_{\pm \mathcal{O}}$.

If $-1 \in \mathbb{D}$, then we set $c_{J_{-1}} := \overline{\mathbf{n}}_1$ from 2.19.

Using the notion of permutation groups given in 2.14 we identify the group $\overline{W} = \overline{N}/L$ with $\mathcal{S}_{\pm \mathcal{O}}$. Computations in \overline{W} show that $\overline{V} = (H \langle c_I \mid I \in \mathcal{O} \rangle) V_S$.

Definition 5.10 Let $\widehat{\lambda} \in \text{Irr}(\widehat{L})$. We call $\widehat{\lambda}$ *standardized* if for every $I, I' \in \mathcal{O}$ the characters $\widehat{\lambda}_I$ and $\widehat{\lambda}_{I'}$ are either V_S -conjugate or not \overline{V} -conjugate. We call the characters in $\text{Irr}(\widehat{\lambda} \upharpoonright_L)$ also *standardized*.

Computations show that every standardized character $\widehat{\lambda}$ satisfies $\overline{V}_{\widehat{\lambda}} = H \langle c_I \mid I \in \mathcal{O} \rangle_{\widehat{\lambda}} V_{S, \widehat{\lambda}}$ and every \overline{N} -orbit in $\text{Irr}(\widehat{L})$ contains a standardized character. For a more explicit description of $\overline{W}_d(\widehat{\lambda})$ we use the following notation.

Notation 5.11 Let E be a set and M a subset of 2^E , the set of all subsets of E . For $m' \subseteq E$ we write $m' \subset\subset M$ if $m' \subseteq m$ for some $m \in M$.

Using the notion of permutation groups given in 2.14 we identify the group $\overline{W} = \overline{N}/L$ with $\mathcal{S}_{\pm\mathcal{O}}$. In the following we describe $\overline{W}_d(\widehat{\lambda}_d)$ as subgroup of $\mathcal{S}_{\pm\mathcal{O}_d}$. We use the Young-like subgroups of $\mathcal{S}_{\pm\mathcal{O}_d}$ from Notation 2.14 that are associated to a partition of \mathcal{O}_d .

Lemma 5.12 Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$ be standardized. We set

$$\mathcal{O}_c(\widehat{\lambda}) := \{I \in \mathcal{O} \mid (\widehat{\lambda}_I)^{c_I} = \widehat{\lambda}_I\}.$$

Let $Y(\widehat{\lambda}) \vdash \mathcal{O}_c(\widehat{\lambda})$ and $Y'(\widehat{\lambda}) \vdash (\mathcal{O}(\widehat{\lambda}) \setminus \mathcal{O}_c(\widehat{\lambda}))$ be the partitions such that $\{I, I'\} \subset\subset Y(\widehat{\lambda})$ or $\{I, I'\} \subset\subset Y'(\widehat{\lambda})$ if and only if $\widehat{\lambda}_I$ and $\widehat{\lambda}_{I'}$ are V_S -conjugate. Then

$$\overline{W}(\widehat{\lambda}) = \mathcal{Y}_{\pm Y(\widehat{\lambda})} \times \mathcal{Y}_{Y'(\widehat{\lambda})},$$

where $\mathcal{Y}_{\pm Y(\widehat{\lambda})}$ and $\mathcal{Y}_{Y'(\widehat{\lambda})}$ are defined as in 2.14.

Proof. Note $Y'(\widehat{\lambda}) \cup Y(\widehat{\lambda}) \vdash \mathcal{O}$. As $\widehat{\lambda}$ is standardized

$$\overline{W}_d(\widehat{\lambda}) = \langle (I, -I) \mid I \in \mathcal{O}_d \rangle_{\widehat{\lambda}} \rtimes \langle (I, I')(-I, -I') \mid I, I' \in \mathcal{O}_d \rangle_{\widehat{\lambda}} \text{ for every } d \in \mathbb{D}.$$

This gives our claim. \square

Let $\zeta \in \mathbb{F}^\times$ with $\varpi = \zeta^{(q-1)^2}$ and $t_{I,2} := \mathbf{h}_I(\zeta)$ for every $I \subseteq \underline{l}$ as in Lemma 2.13. For $I \in \mathcal{O} \setminus \{J_{-1}\}$ the element $t_{I,2}$ satisfies $[\mathbf{L}_I, t_{I,2}] = 1$.

Lemma 5.13 — Structure of $\overline{W}(\widehat{\lambda})$. Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$, $\widehat{\lambda} \in \text{Irr}(\widehat{L} \mid \lambda)$, $\widetilde{\lambda} \in \text{Irr}(\widetilde{L} \mid \widehat{\lambda})$, $\widehat{\lambda}_I \in \text{Irr}(\widehat{\lambda} \upharpoonright_{\widehat{L}_I})$ and $\widetilde{\lambda}_I \in \text{Irr}(\widetilde{L}_I \mid \widehat{\lambda}_I)$ ($I \in \mathcal{O}_c(\widehat{\lambda})$). Assume that $\widehat{\lambda}$ is standardized and $\widetilde{L}_{\widehat{\lambda}} = \widetilde{L}$. We set

$$\mathcal{O}_{c,1}(\widehat{\lambda}) := \{I \in \mathcal{O}_c(\widehat{\lambda}) \mid (\widetilde{\lambda}_I)^{c_I} = \widetilde{\lambda}_I\} \text{ and } \mathcal{O}_{c,-1}(\widehat{\lambda}) := \{I \in \mathcal{O}_c(\widehat{\lambda}) \mid (\widetilde{\lambda}_I)^{c_I} \neq \widetilde{\lambda}_I\}.$$

(a) For $I \in \mathcal{O}_c(\widehat{\lambda}) \setminus \{J_{-1}\}$ and $\epsilon = \pm 1$: $I \in \mathcal{O}_{c,\epsilon}(\widehat{\lambda}) \Leftrightarrow \widehat{\lambda}_I(t_{I,2}^2) = \epsilon \widehat{\lambda}_I(1)$.

(b) $\overline{W}(\widetilde{\lambda}) \leq \mathcal{S}_{\pm\mathcal{O}_{c,1}(\widehat{\lambda})} \times \mathcal{S}_{\pm\mathcal{O}_{c,-1}(\widehat{\lambda})}^{\text{D}}$, more precisely

$$\overline{W}(\widetilde{\lambda}) = \left(\langle (I, -I) \mid I \in \mathcal{O}_{c,1}(\widehat{\lambda}) \rangle \langle (I, -I)(I', -I') \mid I, I' \in \mathcal{O}_{c,-1}(\widehat{\lambda}) \rangle \right) \rtimes \mathcal{Y}_{Y(\widehat{\lambda}) \cup Y'(\widehat{\lambda})}.$$

If from the context the character $\widehat{\lambda}$ is clear we write $\mathcal{O}_{c,\epsilon}$ instead of $\mathcal{O}_{c,\epsilon}(\widehat{\lambda})$.

Proof. Note that the description of \widetilde{L} given in Lemma 2.13(e) shows that $\widetilde{\lambda}$ extends to \widetilde{L} if and only if $\widehat{\lambda}_I$ extends to $\widetilde{L}_I := \widehat{L}_I \langle t_{I,2} \rangle$ for every $I \in \mathcal{O}$.

We have $t_{\underline{l},2} := \prod_{I \in \mathcal{O}} t_{I,2}$, $\mathcal{L}(t_{\underline{l},2}) = \mathbf{h}_{\underline{l}}(\varpi)$ and $\widetilde{L} = \langle \widehat{L}, t_{\underline{l},2} \rangle$, see Lemma 2.13. This implies $\widetilde{L} \leq \langle \widetilde{L}_I \mid I \in \mathcal{O} \rangle$. By the Chevalley relations we see $[V_S, t_{\underline{l},2}] = 1$. Let $I, I' \in \mathcal{O}$ such that $\widehat{\lambda}_I$ and $\widehat{\lambda}_{I'}$ are V_S -conjugate. Then we can choose their extensions $\widetilde{\lambda}_I$ and $\widetilde{\lambda}_{I'}$ to \widetilde{L}_I and $\widetilde{L}_{I'}$ such that they are V_S -conjugate, as $N_{V_S}(\widehat{L}_I) = HC_{V_S}(\widehat{L}_I)$, and therefore $\widetilde{\lambda}_{I'}$ is uniquely determined by $\widetilde{\lambda}_I$.

Let $\phi \in \text{Irr}(\langle \tilde{L}_I \mid I \in \mathcal{O} \rangle)$ with $\phi|_{\tilde{L}_I} = \tilde{\lambda}_I$ for every $I \in \mathcal{O}$. Without loss of generality we may assume $\phi|_{\tilde{L}} = \tilde{\lambda}$. By the above construction we have $V_{S,\phi} = V_{S,\tilde{\lambda}}$. Because of $\overline{V}_{\tilde{\lambda}} = H \langle c_I \mid I \in \mathcal{O} \rangle_{\tilde{\lambda}} V_{S,\tilde{\lambda}}$, it is sufficient to determine $\langle c_I \mid I \in \mathcal{O} \rangle_{\tilde{\lambda}}$ for computing $\overline{V}_{\tilde{\lambda}}$.

Let $\mu_I \in \text{Irr}(\tilde{L}_I)$ be the linear character with $\ker(\mu_I) = \hat{L}_I$. For any $Q \subseteq \mathcal{O}$ let $\mu_Q \in \text{Irr}(\langle \tilde{L}_I \mid I \in \mathcal{O} \rangle)$ be the linear character with $\langle L_I \mid I \in \mathcal{O} \rangle \leq \ker(\mu_Q)$ such that for every $I \in \mathcal{O}$ the inclusion $\tilde{L}_I \leq \ker(\mu_Q)$ holds if and only if $I \notin Q$. Note that $\mu_Q(t_{I,2}) = 1$ if and only if $|Q|$ is even.

For $Q \subseteq \mathcal{O}$ let $c_Q := \prod_{I \in Q} c_I \in \overline{V}$. If $Q' \subseteq \mathcal{O}_c(\hat{\lambda})$, then $c_{Q'} \in \overline{V}_{\tilde{\lambda}}$ and we see that $\phi^{c_{Q'}} = \phi \mu_{Q' \cap \mathcal{O}_{c,-1}}$. As $\mu_{Q' \cap \mathcal{O}_{c,-1}}(t_{I,2}) = (-1)^{|Q' \cap \mathcal{O}_{c,-1}|}$ this leads to a proof of part (b), in particular

$$\overline{W}(\tilde{\lambda}) = (\langle (I, -I) \mid I \in \mathcal{O}_{c,1} \rangle \langle (I, -I)(I', -I') \mid I, I' \in \mathcal{O}_{c,-1} \rangle) \rtimes \mathcal{Y}_{Y(\hat{\lambda}) \cup Y'(\hat{\lambda})}.$$

Let $I \in \mathcal{O}_c(\hat{\lambda}) \setminus \{J_{-1}\}$. Then c_I acts by inverting on \mathbf{T}_I , in particular $t_{I,2}^{c_I} = t_{I,2}^{-1}$ and $[c_I, t_{I,2}] = t_{I,2}^{-2}$. Because of $t_{I,2} \in Z(\mathbf{L}_I)$ we see that $[t_I, \overline{V}_d] \subseteq Z(\mathbf{L})$. Any extension $\tilde{\lambda}_I$ of $\hat{\lambda}_I$ to $\langle \hat{L}_I, t_{I,2} \rangle$ satisfies $\tilde{\lambda}_I(t_{I,2}) \neq 0$ since $t_{I,2} \in Z(\tilde{L}_I)$.

Note $\nu_I \in \text{Irr}(\tilde{\lambda}|_{Z(\tilde{L}_I)})$ is linear. As $\hat{\lambda}_I$ is c_I -stable ν_I has multiplicative order 1 or 2. We observe that $t_{I,2}^{c_I} = t_{I,2}^{-1} \in Z(\tilde{L}_I)$ and hence

$$\tilde{\lambda}_I(t_{I,2}^{c_I}) = \lambda_I(1)\nu_I(t_{I,2}^{c_I}) = \tilde{\lambda}(t_{I,2})\nu_I([t_{I,2}, c_I]) = \tilde{\lambda}(t_{I,2})\nu_I(t_{I,2}^{-2}).$$

Accordingly $\tilde{\lambda}_I$ is c_I -invariant if and only if $[c_I, t_{I,2}] \in \ker(\nu_I) = \ker(\hat{\lambda}) \cap Z(\hat{L}_I)$. This proves (a). \square

The group $W(\lambda)$ is then generated by $W(\hat{\lambda})$ and an element that is described below.

Lemma 5.14 *Let $\mu \in \text{Irr}(\hat{L})$ with $\ker(\mu) = L$ and $\hat{\lambda} \in \text{Irr}(\hat{L})$. Additionally for every $I \in \mathcal{O}$ let $\mu_I \in \text{Irr}(\hat{L}_I)$ with $\ker(\mu_I) = L_I$, $\hat{\lambda}_I \in \text{Irr}(\hat{\lambda}|_{\hat{L}_I})$ and $\lambda_I \in \text{Irr}(\hat{\lambda}_I|_{L_I})$.*

- (a) *Let $x \in \overline{W} \setminus \overline{W}(\hat{\lambda})$ and $\lambda \in \text{Irr}(\hat{\lambda}|_L)$. Then $x \in \overline{W}(\lambda)$ if and only if for every $I \in \mathcal{O}$ the equality $(\hat{\lambda}_I)^x = \hat{\lambda}_{I'}\mu_{I'}$ holds, where $I' \in \mathcal{O}$ with $(\hat{L}_I)^x = \hat{L}_{I'}$.*
- (b) *We set $\mathcal{O}_{\text{ext}} := \{I \in \mathcal{O} \mid \hat{\lambda}_I|_{L_I} = \lambda_I\}$ and $\mathcal{O}_{\text{ind}} := \mathcal{O} \setminus \mathcal{O}_{\text{ext}}$. Then $W(\lambda)$ stabilizes \mathcal{O}_{ext} and \mathcal{O}_{ind} .*

Proof. Since \hat{L}/L has order 2, we see that $\mu_{\mathcal{O}}$, the product of the characters $\hat{\mu}_I$ ($I \in \mathcal{O}$) defined as in the proof of Lemma 5.13, is an extension of μ . This implies part (a).

For part (b) we observe that for $I \in \mathcal{O}$, $\sigma \in W(\lambda)$ and $I' := \sigma^{-1}(I)$ the characters $\hat{\lambda}_I|_{L_I}$ and $(\hat{\lambda}_I)^\sigma|_{L_{I'}} = \hat{\lambda}_{I'}\mu_{I'}|_{L_{I'}}$ have the same number of constituents. This proves part (b) since $I \in \mathcal{O}_{\text{ind}}$ if and only if $\hat{\lambda}_I|_{L_I}$ is reducible. \square

5.B Cuspidal characters of L_I

The aim here is to describe the structure of $\overline{W}(\tilde{\lambda})$ by analysing $\mathcal{O}_{c,-1}(\hat{\lambda})$ (see Lemma 5.13). We show in this section that for some $I \in \mathcal{O}$ there exists no or only few c_I -stable cuspidal characters of L_I and study the kernel of those characters, see Corollary 5.22.

For $I \in \mathcal{O}$, let $\text{Irr}_{\text{cusp}}(\hat{L}_I) := \text{Irr}(\hat{L}_I \mid \text{Irr}_{\text{cusp}}(L_I))$ and call those characters *cuspidal* as well.

Lemma 5.15 *Let $I \in \mathcal{O}_d$ for some $d \in \mathbb{D}_{\text{odd}} \setminus \{\pm 1\}$. There exists no c_I -stable character in $\text{Irr}_{\text{cusp}}(\widehat{L}_I)$.*

Proof. According to Lemma 3.2(c), $L_I \cong \text{GL}_d(q)$ and the element c_I defined in 5.9 induces on G_I a combination of an inner automorphism and the non-trivial graph automorphism according to Lemma 3.4(b). The element c_I acts on the torus $\mathbf{Z}_I := \mathbf{h}_I(\mathbb{F}^\times)$ from Lemma 3.2 by inverting. Hence via the isomorphism $L_I \cong \text{GL}_d(q)$ the element c_I induces on L_I a combination of an inner automorphism and the non-trivial graph automorphism.

According to Proposition 4.1(a) there is no cuspidal character of $\text{GL}_d(q)$ that is invariant under transpose-inverse. So no character in $\text{Irr}_{\text{cusp}}(L)$ is c_I -stable. Now the element t_I from Lemma 2.13 can be chosen such that $[t_I, \mathbf{L}_I] = 1$, see Lemma 3.3. This implies that every character of \widehat{L}_I is an extension of a cuspidal character of L_I . This proves that there is no c_I -stable character in $\text{Irr}_{\text{cusp}}(\widehat{L}_I)$. \square

With the following statement the above shows that $\mathcal{O}_{c,1}(\widehat{\lambda}) \cap \mathcal{O}_d = \emptyset$ for every $d \in \mathbb{D} \setminus \{\pm 1\}$ and $\widehat{\lambda} \in \text{Irr}_{\text{cusp}}(\widehat{L})$ with $h_0 \in \ker(\widehat{\lambda})$.

Proposition 5.16 *Let $I \in \mathcal{O}_d$ for some $d \in \mathbb{D} \setminus \{\pm 1\}$. Then every $\psi \in \text{Irr}_{\text{cusp}}(\widehat{L}_I | 1_{\langle h_0 \rangle})$ with $\psi^{c_I} = \psi$ satisfies $\mathbf{Z}_I^F \leq \ker(\psi)$, where $\mathbf{Z}_I := \mathbf{h}_I(\mathbb{F}^\times)$ as in Lemma 3.2.*

Proof. Under the isomorphism $\mathbf{L}_I / \langle h_0 \rangle \cong \text{GL}_d(\mathbb{F})$ from Lemma 3.2 we obtain $\widehat{L}_I / \langle h_0 \rangle \cong \text{GL}_d(q)$. Via this isomorphism \mathbf{Z}_I^F is mapped to $Z(\text{GL}_d(q))$. Let $\psi \in \text{Irr}_{\text{cusp}}(\widehat{L}_I | 1_{\langle h_0 \rangle})$. If ψ is c_I -invariant, then it corresponds to a cuspidal character of $\text{GL}_d(q)$ that is invariant under transpose-inverse, see Lemma 3.4(b). According to Proposition 4.1(a) such a character is trivial on the centre. This implies $\mathbf{Z}_I^F \leq \ker(\psi)$. \square

Theorem 5.17 *Let $\nu \in \text{Irr}(\langle h_0 \rangle)$ be non-trivial, $d \in \mathbb{D}_{\text{even}}$, $I \in \mathcal{O}_d$ and let $t_{I,2}$ be as defined before Lemma 5.13.*

- (a) *If $d \geq 4$, every $\psi \in \text{Irr}_{\text{cusp}}(\widehat{L}_I | \nu)$ with $\psi^{c_I} = \psi$ satisfies $t_{I,2}^2 \in \ker(\psi)$.*
- (b) *If $d = 2$ and $4 \mid (q - 1)$, there is a unique $\psi \in \text{Irr}_{\text{cusp}}(\widehat{L}_I | \nu)$ with $\psi^{c_I} = \psi$ and $t_{I,2}^2 \notin \ker(\psi)$.*

The proof goes through the next three lemmas. We keep ν the non-trivial irreducible character of $\langle h_0 \rangle$. As a first step towards a proof of the above we determine the inertia group in \widehat{L}_I of cuspidal c_I -stable characters of L_I .

Lemma 5.18 *Let $d \in \mathbb{D}_{\text{even}}$, $I \in \mathcal{O}_d$, $\psi \in \text{Irr}_{\text{cusp}}(L_I | \nu)$ with $\psi^{c_I} = \psi$ and $t_{I,2}^2 \notin \ker(\psi)$. Then $\widehat{L}_{I,\psi} = L_I$.*

Proof. For the proof it is sufficient to show that a character ψ with the above properties and $\widehat{L}_{I,\psi} = \widehat{L}_I$ cannot exist. Recall $t_{I,2}^2 = \mathbf{h}_I(\zeta')$, where $\zeta' \in \mathbb{F}^\times$ is a root of unity of order $2(q - 1)_2$.

Let $\mathbf{G}' := \text{D}_{2d,sc}(\mathbb{F})$ with an \mathbb{F}_q -structure given by a standard Frobenius endomorphism $F_1 : \mathbf{G}' \rightarrow \mathbf{G}'$. Let \mathbf{L}' be the Levi subgroup of \mathbf{G}' of type $\text{A}_{d-1} \times \text{A}_{d-1}$ such that $\mathcal{O}(\mathbf{L}') = \mathcal{O}_d(\mathbf{L}') = \{I_1, I_2\}$ be defined by \mathbf{L}' as in 2.8. Then ψ defines cuspidal characters $\lambda_{I_1} \in \text{Irr}_{\text{cusp}}(L_{I_1})$ and $\lambda_{I_2} \in \text{Irr}_{\text{cusp}}(L_{I_2})$ that have extensions to \widehat{L}_{I_1} and \widehat{L}_{I_2} and are V'_S -stable, where V'_S is associated to \mathbf{G}' and \mathbf{L}' as in Definition 5.10. We can choose $\widehat{\lambda}_{I_j} \in \text{Irr}(\widehat{L}_{I_j} | \lambda_{I_j})$ ($j = 1, 2$) such that they are not V'_S -conjugate. The group $\widehat{L}' := \widehat{L}_{I_1} \cdot \widehat{L}_{I_2}$ is a central product of the groups \widehat{L}_{I_j} ($j \in \underline{2}$) over $\langle h_0 \rangle$. Let $\widehat{\lambda}' := \widehat{\lambda}_{I_1} \cdot \widehat{\lambda}_{I_2} \in \text{Irr}(\widehat{L}')$, $\lambda' = \widehat{\lambda}' \Big|_{L'}$

and $\tilde{\lambda}' \in \text{Irr}(\tilde{L}' | \hat{\lambda}')$ where $L' := (\mathbf{L}')^F$, $\hat{L}' := \mathcal{L}'^{-1}(\langle h_0 \rangle) \cap \mathbf{L}'$ and $\tilde{L}' := \mathcal{L}'^{-1}(\mathbf{Z}(\mathbf{G}')) \cap \mathbf{L}'$ for the Lang map $\mathcal{L}': x \mapsto x^{-1}F'(x)$ of \mathbf{G}' .

Defining W, \bar{W} from the above for \mathbf{G}' and \mathbf{L}' , note that $W(\hat{\lambda}') = \bar{W}(\hat{\lambda}') = \langle (I_1, -I_1), (I_2, -I_2) \rangle$ and $W(\lambda') = \bar{W}(\lambda') = \mathcal{S}_{\pm \mathcal{O}(\mathbf{L}')}.$ Note also that $W(\tilde{\lambda}') = \langle (I_1, -I_1)(I_2, -I_2) \rangle \leq \mathbf{Z}(W(\lambda')) = [W(\lambda'), W(\lambda')].$

Now observe that the non-trivial character of $W(\tilde{\lambda}')$ is $W(\lambda')$ -stable but does not extend to $W(\lambda')$ as the kernel of any linear character of $W(\lambda')$ contains $\mathbf{Z}(W(\lambda')) = [W(\lambda'), W(\lambda')].$ This also implies that for some character $\eta \in \text{Irr}(W(\lambda'))$ the constituent η_0 of $\eta|_{W(\tilde{\lambda}')}$ has multiplicity 2 in $\eta|_{W(\tilde{\lambda}')}.$ The character $\mathbf{R}_{\tilde{L}'}^{\tilde{G}'}(\tilde{\lambda}')_{\eta_0}$ restricts to $(\mathbf{G}')^F$ and has only constituents with multiplicity 1 according to [CE, 15.11].

Like in other places these results are considering first the situation of Harish-Chandra induction for a group $(\tilde{\mathbf{G}}')^{F_1}$ that comes from a regular embedding of \mathbf{G}' into a group with connected centre. These results can be then applied to the groups $\tilde{G}' := \mathcal{L}^{-1}(\mathbf{Z}(\mathbf{G}'))$ and the subgroup $\tilde{L}'.$

On the other hand according to [B06, 13.9(b)], the character $\mathbf{R}_{L'}^{G'}(\lambda')_{\eta}$ has multiplicity 2 in $\mathbf{R}_{\tilde{L}'}^{\tilde{G}'}(\tilde{\lambda}')_{\eta_0}.$ This is a contradiction. This implies that a character ψ with the above properties cannot exist and proves the statement. \square

In the next step we continue to consider the case where $I \in \mathcal{O}_d$ with $2 \mid d.$

Lemma 5.19 *Let $\nu \in \text{Irr}(\langle h_0 \rangle)$ be non-trivial and $I \in \mathcal{O}_d$ for some $d \in \mathbb{D}_{\text{even}}$ with $d > 2.$ Then every $\psi \in \text{Irr}_{\text{cusp}}(L_I | \nu)$ with $\psi^{c_I} = \psi$ satisfies $t_{I,2}^2 \in \ker(\psi).$*

Proof. Let $z := t_{I,2}^2 = \mathbf{h}_I(\zeta')$ for some $\zeta' \in \mathbb{F}^\times$ a root of unity of order $2(q-1)_2,$ and $\psi \in \text{Irr}_{\text{cusp}}(L_I | \nu)$ with $\psi^{c_I} = \psi$ and $z \notin \ker(\psi).$ According to Lemma 5.18, $\psi^{\hat{L}_I}$ is irreducible. Note $z \in \mathbf{Z}(\mathbf{L}_I).$ Since $d \geq 4$ it is sufficient to show the statement in the case where $I = \underline{l}$ and hence $L_I = L.$

The group $L_0 := [\mathbf{L}, \mathbf{L}]^F$ satisfies $L_0 \cong \text{SL}_l(q),$ see Lemma 3.2. Let $\psi_0 \in \text{Irr}(\psi|_{L_0}).$ According to Lemma 1.15, ψ_0 is cuspidal. Following Lemma 3.3 the automorphisms of L_0 induced by \hat{L} are diagonal automorphisms of $L_0.$ Since $C_{\hat{L}}(L_0) \leq \mathbf{Z}(\mathbf{L})$ and $\hat{L}/(C_{\hat{L}}(L_0)L_0)$ is cyclic, we can see that maximal extendibility holds with respect to $L_0 \triangleleft L.$ As $\psi^{\hat{L}}$ is irreducible, $\hat{L}\psi = L.$ As \hat{L}/L_0 is abelian this implies $\hat{L}\psi_0 \leq L.$

We now use the fact $L_0 \cong \text{SL}_l(q).$ Let $\mathbf{H} := \text{GL}_l(\mathbb{F})$ and let $F' : \mathbf{H} \rightarrow \mathbf{H}$ be a Frobenius endomorphism giving an \mathbb{F}_q -structure such that $\mathbf{H}^{F'} \cong \text{GL}_l(q).$ Via $[\mathbf{H}, \mathbf{H}] \cong [\mathbf{L}, \mathbf{L}]$ we identify $[\mathbf{H}, \mathbf{H}]$ with $\mathbf{L}_0.$ Hence ψ_0 defines $\psi'_0 \in \text{Irr}_{\text{cusp}}([\mathbf{H}, \mathbf{H}]^{F'}).$ By the above this implies

$$2 \mid |\mathbf{H}^F : \mathbf{H}_{\psi'_0}^F|.$$

The character $\psi^{\hat{L}}$ is c_I -stable. Hence $\psi|_{L_0}$ is c_I -stable. Following Notation 5.9, c_I acts on L_0 by a graph automorphism and \hat{L} acts on L_0 as diagonal automorphisms.

As $\psi|_{L_0}$ is c_I -stable, we can choose ψ' to be stable under the graph automorphism of $\text{SL}_l(q)$ and it is cuspidal according to Lemma 1.15. In this situation ψ_0 only exists if $l = 2,$ see Proposition 4.2. By the assumption $d > 2,$ so we get a contradiction. This implies our claim that any c_I -stable character ψ satisfies $t_{I,2}^2 \in \ker(\psi).$ \square

Lemma 5.20 *Let $I \in \mathcal{O}_2$ and ν as in Lemma 5.19. There are exactly two characters $\psi \in \text{Irr}_{\text{cusp}}(L_I | \nu)$ with $\psi^{c_I} = \psi$ and $t_{I,2}^2 \notin \ker(\psi).$ Those characters are \hat{L} -conjugate.*

Proof. From the proof of Lemma 5.19 and Proposition 4.2 we see that there are two $\mathrm{GL}_2(q)$ -conjugate characters $\psi_0 \in \mathrm{Irr}(\psi|_{[\mathbf{L}_I, \mathbf{L}_I]^F})$, that are the only possible constituents of ψ . If $I = \{i, i'\}$, then $\psi_0(\mathbf{h}_{e_i - e_{i'}}(-1)) = (-1)^{\frac{q+1}{2}} \psi_0(1)$ according to [B, Tab. 5.4]. Then $\mathbf{L}^F \cong \mathrm{SL}_2(q) \times \mathbf{Z}_I^F$ by Lemma 3.2, in particular $h_0 = \mathbf{h}_I(\varpi) \mathbf{h}_{e_i - e_{i'}}(-1)$. Because of $\psi(h_0) = -\psi(1)$ this implies $\psi(\mathbf{h}_I(\varpi)) = -(-1)^{\frac{q+1}{2}} \psi(1) = (-1)^{\frac{q-1}{2}} \psi(1)$.

Let $\kappa \in \mathrm{Irr}(\mathbf{Z}_I^F)$ such that $\psi = \psi_0 \times \kappa$. As c_I acts by inverting on \mathbf{Z}_I , κ has multiplicative order 1 or 2. The assumption $t_{I,2}^2 \notin \ker(\psi)$ implies that κ has order 2. This proves that given ψ_0 , the character κ is uniquely determined by the fact that ψ is c_I -stable and $t_{I,2}^2 \notin \ker(\psi)$. Hence the only characters with the given properties are \widehat{L} -conjugate. \square

By the above three statements we can show Theorem 5.17.

Proof of Theorem 5.17. Let $\nu \in \mathrm{Irr}(\langle h_0 \rangle)$ be non-trivial, $d \in \mathbb{D}_{\mathrm{even}}$, $I \in \mathcal{O}_d$ and $\psi \in \mathrm{Irr}_{\mathrm{cusp}}(\widehat{L}_I | \nu)$ with $\psi^{c_I} = \psi$. If $d > 2$, then $t_{I,2}^2 \in \ker(\psi)$ according to Lemma 5.19. This shows part (a).

Assume $d = 2$ and $t_{I,2}^2 \notin \ker(\psi)$. The set $\mathrm{Irr}(\psi|_{L_I})$ contains two characters according to Lemma 5.18. Following Lemma 3.4(b) together with Proposition 1.19 the character $\psi' \in \mathrm{Irr}(\psi|_{L_I})$ is cuspidal, satisfies $(\psi')^{c_I} = \psi'$ and $t_{I,2}^2 \notin \ker(\psi')$. Then there are exactly two \widehat{L}_I -conjugate characters in $\mathrm{Irr}_{\mathrm{cusp}}(L_I | \nu)$ with those properties, see Lemma 5.20. Since $|\widehat{L}_I : L_I| = 2$ this implies that there is only one character ψ with the given properties. This proves (b). \square

Lemma 5.21 *If $\nu \in \mathrm{Irr}(\langle h_0 \rangle)$ is non-trivial, then every $\psi \in \mathrm{Irr}_{\mathrm{cusp}}(\widehat{L}_{J-1} | \nu)$ satisfies $\psi^{c_{J-1}} \neq \psi$.*

Proof. Note $\mathbf{h}_{J-1}(\varpi) \in \mathbf{Z}(\mathbf{L}_{-1})$ and $[c_{J-1}, \mathbf{h}_{J-1}(\varpi)] = h_0$. No extension $\widehat{\nu} \in \mathrm{Irr}(\mathbf{Z}(\mathbf{L}_{-1})^F | \nu)$ is c_{J-1} -stable. This implies that ψ_{-1} is not c_{J-1} -stable. \square

The above leads to the following statement on the sets $\mathcal{O}_c(\widehat{\lambda})$, $\mathcal{O}_{c,-1}(\widehat{\lambda})$ and $Y(\widehat{\lambda})$ introduced earlier in 5.12 and 5.13. We use the notation $o(\mu)$ to denote the multiplicative order of a linear character μ of a finite group.

Corollary 5.22 *Let $\lambda \in \mathrm{Irr}_{\mathrm{cusp}}(L)$, $\widehat{\lambda}$ and $\widehat{\lambda}_I$ associated to λ as in Lemma 5.13. If $\widehat{\lambda}$ is standardized, then:*

- (a) $\mathcal{O}_c(\widehat{\lambda}) \subseteq \bigcup_{d \in \mathbb{D}_{\mathrm{even}} \cup \{1, -1\}} \mathcal{O}_d$.
- (b) If $h_0 \in \ker(\lambda)$, then $\mathcal{O}_{c,-1}(\widehat{\lambda}) \subseteq J_{-1} \cup \{I \in \mathcal{O}_1 \mid o(\widehat{\lambda}_I) \mid 2\}$.
- (c) If $h_0 \notin \ker(\lambda)$, then $\mathcal{O}_{c,-1}(\widehat{\lambda}) \subseteq \mathcal{O}_2$ and all $\{\widehat{\lambda}_I \mid I \in \mathcal{O}_{c,-1}(\widehat{\lambda})\}$ are V_S -conjugate, i.e. $\mathcal{O}_{c,-1}(\widehat{\lambda}) \in Y(\widehat{\lambda})$.

Proof. Lemma 5.15 implies that $\mathcal{O}_{c,-1}(\widehat{\lambda}) \cap \mathcal{O}_d = \emptyset$ for every $d \in \mathbb{D}_{\mathrm{odd}} \setminus \{\pm 1\}$. This gives (a).

For the proof of (b) assume $h_0 \in \ker(\lambda)$. Then Proposition 5.16 implies $\mathcal{O}_{c,-1}(\widehat{\lambda}) \subseteq \mathcal{O}_{-1} \cup \mathcal{O}_1$. For $I \in \mathcal{O}_1$, the character $\widehat{\lambda}_I$ is c_I -stable if and only if $o(\widehat{\lambda}_I) \mid 2$.

For the proof of (c) assume $h_0 \notin \ker(\lambda)$. Then $\mathcal{O}_{c,-1}(\widehat{\lambda}) \cap \mathcal{O}_1 = \emptyset$ and $\mathcal{O}_{c,-1}(\widehat{\lambda}) \subseteq \mathcal{O}_2$ according to Theorem 5.17. Lemma 5.20 proves that $\{\widehat{\lambda}_I \mid I \in \mathcal{O}_{c,-1}(\widehat{\lambda})\}$ are V_S -conjugate. Hence the partition $Y(\widehat{\lambda})$ from Lemma 5.12 contains $\mathcal{O}_{c,-1}(\widehat{\lambda})$. \square

Recall $K(\lambda) := \overline{W}_{\lambda^{L\langle F_p \rangle}}|_L$. For any \overline{W} -stable $L \leq J \leq \widetilde{L}$, $\kappa \in \mathrm{Irr}(J)$ and $Q \subseteq \mathcal{O}$ let $\overline{W}^Q(\kappa) := \overline{W}(\kappa) \cap \mathcal{S}_{\pm Q}$ and $W^Q(\kappa) := \overline{W}^Q(\kappa) \cap W$.

Proposition 5.23 Let $\lambda \in \mathbb{T}$, $\widehat{\lambda} \in \text{Irr}_{\text{cusp}}(\widehat{L} \mid \lambda)$, $\widehat{\lambda}_I \in \text{Irr}(\widehat{\lambda} \upharpoonright_{\widehat{L}_I})$ ($I \in \mathcal{O}$) and $\widetilde{\lambda} \in \text{Irr}(\widetilde{L} \mid \widehat{\lambda})$. Assume that $\widehat{\lambda}$ is standardized in the sense of 5.10. We set

$$Q^1(\widehat{\lambda}) := \begin{cases} \{I \in \mathcal{O}_1 \mid o(\widehat{\lambda}_I) \mid 2\} \cup \mathcal{O}_{-1} & , \text{ if } h_0 \in \ker(\lambda), \\ \mathcal{O}_{c,-1}(\widehat{\lambda}) & \text{ otherwise.} \end{cases}$$

Then:

- (a) $Q^1(\widehat{\lambda})$ is $K(\lambda)$ -stable, and
- (b) $W(\widetilde{\lambda}) = W^1(\widetilde{\lambda}) \times W^2(\widetilde{\lambda})$, where $Q^2(\widehat{\lambda}) := \mathcal{O} \setminus Q^1(\widehat{\lambda})$, and $W^j(\widetilde{\lambda}) := W^{Q^j(\widehat{\lambda})}(\widetilde{\lambda})$.

Proof. Let $e \in \langle F_0 \rangle$ such that λ and λ^e are \overline{N} -conjugate. As $\widehat{\lambda}$ is standardized, then $\widehat{\lambda}^e$ is also standardized. As the orders of $\widehat{\lambda}_I$ and $(\widehat{\lambda}_I)^e$ coincide for every $I \in \mathcal{O}$, we see that $Q^1(\widehat{\lambda}) = Q^1(\widehat{\lambda}^e)$ from the definition, whenever $h_0 \in \ker(\lambda) = \ker(\widehat{\lambda}^e)$ and hence $h_0 \in \ker(\widehat{\lambda}^e)$.

Assume $h_0 \notin \ker(\lambda)$. Then $h_0 \notin \ker(\lambda^e)$. Let $I \in \mathcal{O} \setminus \{J_{-1}\}$. Because of $c_I^e \in c_I \langle h_0 \rangle$ we see

$$\widehat{\lambda}_I^{c_I} = \widehat{\lambda}_I \Leftrightarrow \text{Irr}(\widehat{\lambda}^e \upharpoonright_{\widehat{L}_I})^{c_I} = \text{Irr}(\widehat{\lambda}^e \upharpoonright_{\widehat{L}_I}).$$

In the case of $\widehat{\lambda}_I^{c_I} = \widehat{\lambda}_I$ the character $\widehat{\lambda}_I$ has some c_I -stable extension to \widetilde{L}_I if and only if the unique character in $\text{Irr}(\widehat{\lambda}^e \upharpoonright_{\widehat{L}_I})$ has some c_I -stable extension to \widetilde{L}_I . (The set $\text{Irr}(\phi \upharpoonright_{\widehat{L}_I})$ is a singleton for every $\phi \in \text{Irr}_{\text{cusp}}(\widehat{L})$, since \widehat{L} is the central product of the groups \widehat{L}_I over $\langle h_0 \rangle$.) This shows $\mathcal{O}_{c,-1}(\widehat{\lambda}) = \mathcal{O}_{c,-1}(\widehat{\lambda}^e)$ and $Q^1(\widehat{\lambda}) = Q^1(\widehat{\lambda}^e)$ by the definition of those sets. Let $w \in \overline{W}$ and $e \in \langle F_p \rangle$ with $we \in \widehat{K}(\lambda)$. Then $\widehat{\lambda}^w$ is standardized and $Q^1(\widehat{\lambda})^w = Q^1(\widehat{\lambda}^w)$. Accordingly $w \in K(\lambda)$ stabilizes $Q^1(\widehat{\lambda})$. This implies part (a).

For part (b) recall the description of $\overline{W}(\widetilde{\lambda})$ from Lemma 5.6:

$$\overline{W}(\widetilde{\lambda}) = \left(\left\langle (I, -I) \mid I \in \mathcal{O}_{c,1}(\widehat{\lambda}) \right\rangle \left\langle (I, -I)(I', -I') \mid I, I' \in \mathcal{O}_{c,-1}(\widehat{\lambda}) \right\rangle \right) \rtimes \mathcal{Y}_{Y(\widehat{\lambda}) \cup Y'(\widehat{\lambda})}.$$

First assume $h_0 \in \ker(\lambda)$. By construction $\mathcal{O}_c(\widehat{\lambda}) \subseteq Q^1(\widehat{\lambda}) \cup \bigcup_{d \in \mathbb{D}_{\text{even}}} \mathcal{O}_d$ and hence $\overline{W}(\widetilde{\lambda}) = \overline{W}^1(\widetilde{\lambda}) \times \overline{W}^2(\widetilde{\lambda})$. According to Corollary 5.22(a) we observe $(I, -I) \in W$ for every $I \in \mathcal{O}_c(\widehat{\lambda}) \setminus Q^1(\widehat{\lambda})$. This implies $\overline{W}^2(\widetilde{\lambda}) \leq W$ and $W(\widetilde{\lambda}) = W^1(\widetilde{\lambda}) \times W^2(\widetilde{\lambda})$ by the definition of W .

It remains to consider the case where $h_0 \notin \ker(\lambda)$. Then $Q^1(\widehat{\lambda}) = \mathcal{O}_{c,-1}(\widehat{\lambda}) \subseteq \mathcal{O}_2$ by Corollary 5.22(c) and hence $\overline{W}^1(\widetilde{\lambda}) \leq W$. By the structure of $\overline{W}(\widetilde{\lambda})$ described in Lemma 5.6 we see $W(\widetilde{\lambda}) = W^1(\widetilde{\lambda}) \times W^2(\widetilde{\lambda})$. \square

5.C Clifford theory for $W(\widetilde{\lambda}) \triangleleft W(\lambda)$ in the case of $\widehat{L}\widetilde{L}_\lambda = \widetilde{L}$

In this section we study the characters of $W(\widetilde{\lambda})$, in particular their Clifford theory with respect to $K(\lambda)$. Assuming $\widehat{L}\widetilde{L}_\lambda = \widetilde{L}$ we prove maximal extendibility with respect to $W(\widetilde{\lambda}) \triangleleft K(\lambda)$. This result is required for a later application of Proposition 1.11. We consider the following situation.

Notation 5.24 Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$, $\widehat{\lambda} \in \text{Irr}(\widehat{L} \mid \lambda)$ and $\widetilde{\lambda} \in \text{Irr}(\widetilde{L} \mid \widehat{\lambda})$ such that $\widehat{\lambda}$ is standardized and $\widehat{L}\widetilde{L}_\lambda = \widetilde{L}$ (or equivalently $\widehat{\lambda} \upharpoonright_{\widehat{L}} = \widetilde{\lambda}$).

For further computations we use the groups $K^j(\lambda)$ associated to the subsets $Q^j(\lambda) \subseteq \mathcal{O}$ from Proposition 5.23, where $K^j(\lambda) := (K(\lambda)\mathcal{S}_{\pm(\mathcal{O} \setminus Q^j(\lambda))}) \cap \mathcal{S}_{\pm Q^j(\lambda)}$ for $j \in \underline{2}$.

Lemma 5.25 *If maximal extendibility holds with respect to $W^j(\tilde{\lambda}) \triangleleft K^j(\lambda)$ for every $j \in \underline{2}$, then maximal extendibility holds with respect to $W(\tilde{\lambda}) \triangleleft K(\lambda)$, in particular for every $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there exists some $K(\lambda)_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$.*

In that situation the above statement will ensure Assumption 1.12(iii) for $(\overline{\lambda}, \eta) \in \mathcal{P}(L)$ via Lemma 5.5.

Proof. Recall $K(\lambda) := \overline{W}_{\lambda^{L(F_p)}} \Big|_L$ by the definition in 5.4. As $K(\lambda)$ stabilizes $Q^1(\widehat{\lambda})$ and $Q^2(\widehat{\lambda})$ by Proposition 5.23, $K(\lambda) \leq \mathcal{S}_{\pm Q^1(\widehat{\lambda})} \times \mathcal{S}_{\pm Q^2(\widehat{\lambda})}$. This rewrites as $K(\lambda) \leq K^1(\lambda) \times K^2(\lambda)$. Recall that $\mathcal{O}_{c,-1}(\widehat{\lambda}) \subseteq Q^1(\widehat{\lambda})$.

Since maximal extendibility holds with respect to $W^j(\tilde{\lambda}) \triangleleft K^j(\lambda)$ for $j \in \underline{2}$ by assumption, maximal extendibility holds with respect to

$$W(\tilde{\lambda}) = W^1(\tilde{\lambda}) \times W^2(\tilde{\lambda}) \triangleleft K^1(\lambda) \times K^2(\lambda).$$

This implies the statement as $K(\lambda) \leq K^1(\lambda) \times K^2(\lambda)$. \square

For $\lambda \in \mathbb{T}$ with $\tilde{L}_\lambda = \tilde{L}$ we study first the Clifford theory of $W^2(\tilde{\lambda}) \triangleleft K^2(\lambda)$ for the groups from Lemma 5.25.

Lemma 5.26 *Let $W^1(\tilde{\lambda})$, $W^2(\tilde{\lambda})$, $K^1(\lambda)$ and $K^2(\lambda)$ be the groups from Lemma 5.25.*

Then:

- (a) *maximal extendibility holds with respect to $W^2(\tilde{\lambda}) \triangleleft K^2(\lambda)$, and*
- (b) *maximal extendibility holds with respect to $W^1(\tilde{\lambda}) \triangleleft K^1(\lambda)$, if $h_0 \notin \ker(\lambda)$.*

Proof. Let $Y(\widehat{\lambda}) \vdash \mathcal{O}_c(\widehat{\lambda})$ and $Y'(\widehat{\lambda}) \vdash \mathcal{O} \setminus \mathcal{O}_c(\widehat{\lambda})$ be the partitions from Lemma 5.12. For proving part (a) we can assume $Q^2(\widehat{\lambda}) = \mathcal{O}$ without loss of generality. We have $\overline{W}(\tilde{\lambda}) = \overline{W}^2(\lambda) = \mathcal{Y}_{\pm Y(\widehat{\lambda})} \times \mathcal{Y}_{Y'(\widehat{\lambda})}$, see Lemma 5.13.

If $h_0 \in \ker(\lambda)$ then $\mathcal{O}_c(\widehat{\lambda}) \cap (\mathcal{O}_1 \cup \mathcal{O}_{-1}) = \emptyset$ by the choice of $Q^1(\widehat{\lambda})$ according to Corollary 5.22. If $h_0 \notin \ker(\lambda)$, Lemma 5.21 implies $\mathcal{O}_c(\widehat{\lambda}) \cap \mathcal{O}_{-1} = \emptyset$ and analogously we see $\mathcal{O}_c(\widehat{\lambda}) \cap \mathcal{O}_1 = \emptyset$.

This implies $\mathcal{O}_c(\widehat{\lambda}) \subseteq \bigcup_{d \in \mathbb{D}_{\text{even}}} \mathcal{O}_d$. Accordingly $W(\tilde{\lambda})$ is the direct product of groups $W_d(\tilde{\lambda})$ for $d \in \mathbb{D}$. It suffices to consider the case where $\mathcal{O} = \mathcal{O}_d = Q^2(\widehat{\lambda})$ for some $d \in \mathbb{D}$ and $\mathcal{O}_c(\widehat{\lambda}) \in \{\mathcal{O}, \emptyset\}$. Additionally we can assume that $Y(\widehat{\lambda})$ and $Y'(\widehat{\lambda})$ are partitions whose elements have all the same cardinality. If $\mathcal{O}_c(\widehat{\lambda}) = \mathcal{O}$, then $W^2(\lambda) \cong (C_2 \wr \mathcal{S}_k)^a$ for some positive integers k and a . Then $K^2(\lambda) \cong (C_2 \wr \mathcal{S}_k) \wr \mathcal{S}_a$, and hence maximal extendibility holds with respect to $W^2(\tilde{\lambda}) \triangleleft K^2(\lambda)$.

If $\mathcal{O}_c(\widehat{\lambda}) = \emptyset$, then $W^2(\lambda) = \mathcal{Y}_{Y'}$ and hence it is isomorphic to a direct product of symmetric groups. The group $K^2(\lambda) \leq \text{N}_{\mathcal{S}_{\pm \mathcal{O}_d}}(\mathcal{Y}_Y)$ is isomorphic to $(C\mathcal{Y}_Y) \rtimes \mathcal{S}_Y$, where $C := \langle \prod_{k \in y} (k, -k) \mid y \in Y \rangle \leq \mathcal{S}_{\pm \mathcal{O}_d}$. By Lemma 3.6 maximal extendibility holds with respect to $W_2(\tilde{\lambda}) \triangleleft K_2(\lambda)$. This proves part (a).

For part (b) we assume $\mathcal{O}_{c,-1}(\lambda) = \mathcal{O}$, $h_0 \notin \ker(\lambda)$, and as before $Q^1(\widehat{\lambda}) = \mathcal{O}$. By Corollary 5.22(c) we have $K(\lambda) = W$ and $|W : W(\tilde{\lambda})| = 2$, see Proposition 5.16(c). As $W(\lambda)/W(\tilde{\lambda})$ is cyclic, maximal extendibility holds with respect to $W(\tilde{\lambda}) \triangleleft K(\lambda)$. \square

It remains to prove the following.

Proposition 5.27 *Maximal extendibility holds with respect to $W^1(\tilde{\lambda}) \triangleleft K^1(\lambda)$, if $h_0 \in \ker(\lambda)$.*

Proof. Let $O_{1,i} = \{I \in \mathcal{O}_1 \mid o(\widehat{\lambda}_I) = i\}$ for $i \in \underline{4}$ and $l_i := |O_{1,i}|$. By Lemma 5.12 and Lemma 5.13 we have $\overline{W}_{Q^1(\widehat{\lambda})}(\tilde{\lambda}) \leq C \times \mathcal{S}_{\pm O_{1,1}} \times \mathcal{S}_{\pm O_{1,2}}$, where $C \leq W_{\mathcal{O}_{-1}}(\widehat{\lambda})$ and C is then either trivial or a cyclic group of order 2. The group structures depend on J_{-1} and those groups are described in Table 5.1, where $W(B_j)$ and $W(D_j)$ are Coxeter groups of type B_j and D_j , respectively.

	$W^1(\tilde{\lambda})$	$K^1(\lambda)$
$J_{-1} \notin \mathcal{O}_c(\widehat{\lambda})$	$W(D_{l_1}) \times W(D_{l_2})$	$C_2 \times (W(B_{l_1}) \wr C_2)$, if $l_1 = l_2$ $C_2 \times W(B_{l_1}) \times W(B_{l_2})$, if $l_1 \neq l_2$
$J_{-1} \in \mathcal{O}_{c,1}$	$W(B_{l_1}) \times W(D_{l_2})$	$C_2 \times W(B_{l_1}) \times W(B_{l_2})$
$J_{-1} \in \mathcal{O}_{c,-1}$	$W(D_{l_1}) \times W(B_{l_2})$	$C_2 \times W(B_{l_1}) \times W(B_{l_2})$

Table 5.1: Isomorphism types of $W^1(\tilde{\lambda})$ and $K^1(\lambda)$

Note that in all cases $\overline{W}_{O_{1,1}}(\tilde{\lambda})$ is a Coxeter group of type B_{l_1} . Considering the structure we observe that in all cases the statement holds according to Lemma 3.6. \square

Recall that $\mathbb{M}^{(X)} = \{\lambda \in \text{Irr}_{\text{cusp}}(L) \mid \tilde{L}_\lambda = X\}$ and $\mathbb{M}_0 := \text{Irr}_{\text{cusp}}(L) \setminus (\mathbb{M}^{(L)} \cup \mathbb{M}^{(\tilde{L})})$ for X with $L \leq X \leq \tilde{L}$. For characters in $\mathbb{M}^{(\tilde{L})} \cup \mathbb{M}_0$ the above proves the following:

Proposition 5.28 *Let $\lambda \in \mathbb{M}^{(\tilde{L})} \cup \mathbb{M}_0$, i.e. $\tilde{L}\tilde{L}_\lambda = \tilde{L}$. For every $\tilde{\lambda} \in \text{Irr}(\tilde{L} \mid \lambda)$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there exists some $K(\lambda)_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$.*

Proof. According to Lemma 5.25, Proposition 5.27 and Lemma 5.26 imply the statement. \square

5.D Clifford theory for $W(\widehat{\lambda}) \triangleleft W(\lambda)$ in the case of $\tilde{L}_\lambda = \widehat{L}$

We now study $W(\lambda)$ and $W(\widehat{\lambda})$ for characters $\lambda \in \mathbb{M}^{(\widehat{L})}$, where $\widehat{\lambda} \in \text{Irr}(\widehat{L} \mid \lambda)$ is standardized. We prove statements on the characters of $W(\widehat{\lambda})$ and their possible extensions to $W(\lambda)$. The results later imply that there exists some E_L -stable \tilde{N} -transversal in $\text{Irr}(N \mid \mathbb{M}^{(\widehat{L})})$.

In the following we study the Clifford theory of $W(\widehat{\lambda}) \triangleleft K(\lambda)$ for $\lambda \in \mathbb{M}^{(\widehat{L})}$, where $K(\lambda) = \overline{W}_{\lambda^{L\langle F_p \rangle}} \Big|_L$.

Lemma 5.29 *Assume $|Z(G)| = 2$ or equivalently $q \equiv 3(4)$ and $2 \nmid l$. Every $\lambda \in \text{Irr}_{\text{cusp}}(L)$ satisfies $\widehat{L} \leq \tilde{L}_\lambda$. Then $\text{Irr}_{\text{cusp}}(N \mid \mathbb{T} \cap \mathbb{M}^{(\widehat{L})})$ is an E_L -stable \tilde{L} -transversal of $\text{Irr}_{\text{cusp}}(N \mid \mathbb{M}^{(\widehat{L})})$.*

Proof. The arguments given in Lemma 5.3 show the statement. \square

According to Lemma 5.29 we can now assume $Z(G) = Z(\mathbf{G})$. We do that until the end of the section.

Lemma 5.30 *If $|Z(G)| = 4$ and $\lambda \in \text{Irr}_{\text{cusp}}(L)$ with $\tilde{L}_\lambda = \hat{L}$, then $-1 \in \mathbb{D}$ and $\hat{\lambda}_{-1}^{t_{L,2}} \neq \hat{\lambda}_{-1}$. Moreover $\hat{\lambda}_{-1}^{\mathbf{n}_{e_1}(\varpi)} = \hat{\lambda}_{-1}$, if $h_0 \in \ker(\lambda)$ or $\text{type}(\Phi_{-1})$ is not of type $D_{|J_{-1}|}$.*

Proof. Recall that maximal extendibility holds with respect to $L \triangleleft \tilde{L}$, see Lemma 1.17. Accordingly $\tilde{L}_\lambda = \hat{L}$ implies that λ is not $t_{L,2}$ -stable for the element $t_{L,2} \in \mathbf{T}$ from Lemma 2.13. If $\zeta \in \mathbb{F}^\times$ with $\zeta^{(q-1)^2} = \varpi$ and $t_{I,2} := \mathbf{h}_I(\zeta)$ as in Lemma 5.13, then $t_{L,2} = \prod_{I \in \mathcal{O}} t_{I,2}$. Recall that $\tilde{L} = \hat{L} \langle t_{L,2} \rangle$. The character $\hat{\lambda}$ is $t_{L,2}$ -stable, if $\hat{\lambda}_I$ is $t_{I,2}$ -stable for every $I \in \mathcal{O}$. For $I \in \mathcal{O} \setminus \{J_{-1}\}$ we see $t_{I,2} \in C_{L_I}(\hat{L}_I)$ and hence $\hat{\lambda}_I$ is $t_{I,2}$ -stable. As $\hat{\lambda}$ is not $t_{L,2}$ -stable, $-1 \in \mathbb{D}$ and $\hat{\lambda}_{-1}$ is not $t_{L,2}$ -stable.

In the next step we prove $\hat{\lambda}_{-1}^{\mathbf{n}_{e_1}(\varpi)} = \hat{\lambda}_{-1}$. Since $\lambda_{-1}^{t_{L,2}} \neq \lambda_{-1}$, Proposition 4.3 implies that $\hat{\lambda}_{-1}$ is γ -stable, if $\text{type}(\Phi_{-1}) = D_{|J_{-1}|}$. We consider the other possible values of $\text{type}(\Phi_{-1})$. We first assume $\text{type}(\Phi_{-1}) = A_1 \times A_1$. Then $L_{-1} = \text{SL}_2(q) \times \text{SL}_2(q)$. Let $\lambda_{-1,1}, \lambda_{-1,2} \in \text{Irr}(\text{SL}_2(q))$ such that $\lambda|_{L_{-1}} = \lambda_{-1,1} \times \lambda_{-1,2}$. By the proof of Proposition 3.8, $\tilde{L}_\lambda \leq \hat{L}$ implies that both characters $\lambda_{-1,1}$ and $\lambda_{-1,2}$ are not $\text{GL}_2(q)$ -stable. Additionally they are cuspidal. Following [B, Tab. 5.4], the characters $\lambda_{-1,1}$ and $\lambda_{-1,2}$ are uniquely determined up to $\text{GL}_2(q)$ -conjugation. After applying some \tilde{L} -conjugation we obtain that $\lambda_{-1,1}$ and $\lambda_{-1,2}$ are $\mathbf{n}_{e_1}(\varpi)$ -conjugate. As \tilde{L} induces on the $\text{SL}_2(q)$ -factors of L_{-1} simultaneously (non-inner) diagonal automorphisms, the set $\text{Irr}(\hat{L}_{-1} | \lambda_{-1})$ contains only one character hence $\hat{\lambda}_{-1}$ is again $\mathbf{n}_{e_1}(\varpi)$ -stable.

It remains to consider the case where $\text{type}(\Phi_{-1}) = A_3$. Again the character λ_{-1} is not \tilde{L} -stable. Via the isomorphism $L_{-1} \cong \text{SL}_4(q)$ we see that $t_{L,2}$ induces on $\text{SL}_4(q)$ a diagonal automorphism corresponding to a generator of $Z(\text{SL}_4(q))$ in the sense of 1.16(b), see also the proof of Proposition 3.8. We take any $\chi \in \text{Irr}(\text{GL}_4(q) | \lambda_{-1})$. Then χ is cuspidal, see Lemma 1.15(c). Using the description of cuspidal characters of general linear groups recalled in the proof of Proposition 4.1, we let $s \in \text{GL}_4(q)$ and $\zeta \in \mathbb{F}^\times$ such that χ belongs to the rational Lusztig series of s and $\zeta \in \mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}$ an eigenvalue of s . Let $\det : \text{GL}_4(\mathbb{F}) \rightarrow \mathbb{F}^\times$ denote the determinant and by \det^* the associated linear character of $\text{GL}_4(\mathbb{F})$ with kernel $\text{SL}_4(\mathbb{F})$. By the assumptions on χ we see that $\chi = \chi(\det^*)^{\frac{q-1}{2}}$ and hence s and $-s$ are conjugate. Then $-\zeta \in \{\zeta, \zeta^q, \zeta^{q^2}, \zeta^{q^3}\}$. Hence, using again o to denote multiplicative order, $o(\zeta)_2 = 2(q^2 - 1)_2$, as $-\zeta \in \{\zeta, \zeta^q, \zeta^{q^3}\}$ would imply that $\zeta \in \mathbb{F}_{q^2}$ or $\zeta \in \mathbb{F}_{q^6}$, contradicting $\zeta \in \mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}$. For computing $\ker(\chi|_{Z(\text{SL}_4(q))})$ we see that $\det s = \zeta^{\frac{q^4-1}{q-1}}$ is not a square in \mathbb{F}_q^\times since $o(\zeta)_2 = 2(q^2 - 1)_2$. This contradicts $h_0 \in \ker(\lambda)$, as h_0 corresponds to the central involution of $\text{SL}_4(q)$. Hence there exists no cuspidal character λ_{-1} of L_{-1} that satisfies $h_0 \in \ker(\lambda)$ and $\lambda_{-1}^{t_{L,2}} \neq \lambda_{-1}$. This shows that $\text{type}(\Phi_{-1}) = A_3$ is not possible. This finishes our proof. \square

Lemma 5.31 *Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$ and $\hat{\lambda} \in \text{Irr}(\hat{L} | \lambda)$ such that $h_0 \notin \ker(\lambda)$ and $W(\hat{\lambda}) \neq W(\lambda)$. Then maximal extendibility holds with respect to $W(\hat{\lambda}) \triangleleft K(\lambda)$.*

Proof. We first determine $W(\hat{\lambda})$. Denote $c_{-1} := c_{J_{-1}}$. As the character $\text{Irr}(\lambda|_{Z(L_{-1})})$ is not c_{-1} -stable, λ_{-1} is not c_{-1} -stable. This implies $J_{-1} \notin \mathcal{O}_c(\hat{\lambda})$.

If $I \in \mathcal{O}_1$, $\hat{\lambda}_I$ is a linear character. Since $h_0 \notin \ker(\lambda)$ and hence $h_0 \notin \ker(\hat{\lambda}_I)$, the order of $\hat{\lambda}_I$ is divisible by $2(q-1)_2 \geq 4$. Hence $\mathcal{O}_c(\hat{\lambda}) \cap \mathcal{O}_1 = \emptyset$.

Together with Lemma 5.14 and Corollary 5.22(a) this leads to $\mathcal{O}_c(\hat{\lambda}) \subseteq \bigcup_{d \in \mathbb{D}_{\text{even}}} \mathcal{O}_d$. The structure of $W(\hat{\lambda})$ is given by Lemma 5.13 and we observe $\overline{W}(\hat{\lambda}) = W(\hat{\lambda})$. As in the proof of Lemma 5.26 we can apply Lemma 3.6, and we see that maximal extendibility

holds with respect to $W(\widehat{\lambda}) \triangleleft N_{\overline{W}}(W(\widehat{\lambda}))$. Because of $K(\lambda) \leq N_{\overline{W}}(W(\widehat{\lambda}))$, this proves maximal extendibility with respect to $W(\widehat{\lambda}) \triangleleft K(\lambda)$. \square

As in Lemma 5.25 we associate to λ subsets $Q^1(\widehat{\lambda})$ and $Q^2(\widehat{\lambda})$ of \mathcal{O} . Recall $K(\lambda) = \overline{W}_{\lambda \langle F_p \rangle} \Big|_L$, whenever \mathbf{G} is not of type D_4 .

Lemma 5.32 *Let $\lambda \in \mathbb{M}(\widehat{L}) \cap \text{Irr}(L \mid 1_{\langle h_0 \rangle})$ and $\widehat{\lambda} \in \text{Irr}(\widehat{L} \mid \lambda)$ with $W(\lambda) \neq W(\widehat{\lambda})$. Let*

$$Q^1(\widehat{\lambda}) := \{I \in \mathcal{O}_1 \mid o(\widehat{\lambda}_I) \in \{1, 2, 4\}\} \cup \mathcal{O}_{-1} \text{ and } Q^2(\widehat{\lambda}) := \mathcal{O} \setminus Q^1(\widehat{\lambda}).$$

Let $\overline{W}^j(\widehat{\lambda}) := \overline{W}(\widehat{\lambda}) \cap \mathcal{S}_{\pm Q^j(\widehat{\lambda})}$, $W^j(\widehat{\lambda}) := \overline{W}^j(\widehat{\lambda}) \cap W$, $\widehat{L}^{(i)} := \langle \widehat{L}_I \mid I \in Q^i(\widehat{\lambda}) \rangle$ and $L^{(i)} := L \cap \widehat{L}^{(i)}$.

(a) Then $K(\lambda)$ stabilizes $Q^1(\widehat{\lambda})$ and

$$W(\widehat{\lambda}) = W^1(\widehat{\lambda}^{(1)}) \times W^2(\widehat{\lambda}^{(2)}),$$

where $\widehat{\lambda}^{(i)} \in \text{Irr}(\widehat{\lambda} \Big|_{\widehat{L}^{(i)}})$.

(b) If $x \in W(\lambda) \setminus W(\widehat{\lambda})$, then $x = x_1 x_2$ for some $x_i \in W^i(\lambda^{(i)})$ ($i \in \underline{2}$), where $\lambda^{(i)} \in \text{Irr}(\lambda \Big|_{L^{(i)}})$.

(c) $|\mathcal{O}_d \cap Q^2(\widehat{\lambda})|$ is even for every $d \in \mathbb{D}_{\text{odd}} \setminus \{-1\}$.

(d) Let $\overline{W}^i := \overline{W} \cap \mathcal{S}_{\pm Q^i(\widehat{\lambda})}$ and $K^i(\lambda) := (\overline{W}^i)_{(\lambda^{(i)})L^{(i)}\langle F_p \rangle} \Big|_{L^{(i)}}$. If $\mathcal{O}_1 \cap Q^1(\widehat{\lambda}) \neq \emptyset$,

then $K(\lambda) \leq K^1(\lambda) \times N_{\overline{W}^2}(W^2(\widehat{\lambda}))$.

Proof. First note that \widehat{N} normalizes the groups \mathbf{L}_I ($I \in \mathcal{O}$) and hence there is a well-defined action of \widehat{W} on \mathcal{O} . Now $Q^1(\widehat{\lambda})$ is defined using $\widehat{\lambda}$ (and is independent of the choice of $\widehat{\lambda} \in \text{Irr}(\widehat{L} \mid \lambda)$). Note that by this definition any element in \widehat{N}_λ stabilizes $Q^1(\widehat{\lambda})$. Without loss of generality we can assume that $\widehat{\lambda}$ is standardized and hence $\overline{W}(\lambda)$ is given in Lemma 5.13.

Accordingly $\overline{W}(\widehat{\lambda}) = \overline{W}^1(\widehat{\lambda}) \times \overline{W}^2(\widehat{\lambda})$. As $\lambda \in \text{Irr}(L \mid 1_{\langle h_0 \rangle})$, Corollary 5.22 implies $\mathcal{O}_c(\widehat{\lambda}) \subseteq \bigcup_{d \in \mathbb{D}_{\text{even}}} \mathcal{O}_d$. By the definition of $Q^2(\widehat{\lambda})$ we observe $\overline{W}^2(\widehat{\lambda}) = W^2(\widehat{\lambda})$.

According to Lemma 5.8(d), $\overline{W}(\widehat{\lambda})$ is the direct product of the groups $\overline{W}_d(\widehat{\lambda})$, where $\overline{W}_d(\widehat{\lambda}) := \overline{V}_{d, \widehat{\lambda}}/H_d$. For $\overline{W}^1(\widehat{\lambda})$ we note that

$$\overline{W}^1(\widehat{\lambda}) = \overline{W}^{1,1}(\widehat{\lambda}) \times \overline{W}^{1,2}(\widehat{\lambda}) \times \overline{W}^{1,4}(\widehat{\lambda}),$$

where $Q^{1,j}(\lambda) := \{I \in \mathcal{O}_1 \mid o(\widehat{\lambda}_I) = j\}$ and $\overline{W}^{1,j}(\widehat{\lambda}) := \overline{W}^1(\widehat{\lambda}) \cap \mathcal{S}_{\pm Q^{1,j}(\lambda)}$. This proves that $\overline{W}(\widehat{\lambda}) = \overline{W}^1(\widehat{\lambda}) \times \overline{W}^2(\widehat{\lambda})$. By the above $W^2(\widehat{\lambda}) = \overline{W}^2(\widehat{\lambda})$ and hence $W(\widehat{\lambda}) = \overline{W}^1(\widehat{\lambda}) \times W^2(\widehat{\lambda}) = W^1(\widehat{\lambda}) \times W^2(\widehat{\lambda})$. Since $\widehat{\lambda} = \widehat{\lambda}^{(1)} \times \widehat{\lambda}^{(2)}$ we observe $W(\widehat{\lambda}^{(1)}) = W^1(\widehat{\lambda}^{(1)})$ and $W^2(\widehat{\lambda}) = W^2(\widehat{\lambda}^{(2)})$.

As $x \in K(\lambda)$ stabilizes $Q^1(\widehat{\lambda})$ by (a), it can be written as product $x_1 x_2$ where $x_i \in \overline{W}^i(\lambda^{(i)})$. Since $\widehat{\lambda}^x = \widehat{\lambda} \mu$ for the faithful character μ of \widehat{L}/L , it satisfies $(\widehat{\lambda}^{(i)})^x = \widehat{\lambda}^{(i)} \mu^{(i)}$ where $\mu^{(i)} = \mu \Big|_{\widehat{L}^{(i)}}$ with $\widehat{L}^{(i)} := \langle \widehat{L}_I \mid I \in Q^i(\lambda) \rangle$. Hence $(\widehat{\lambda}^{(i)})^{x_i} = \widehat{\lambda}^{(i)} \mu^{(i)}$.

In the following we show that any element $x_2 \in \overline{W}^2$ with $(\widehat{\lambda}^{(2)})^{x_i} = \widehat{\lambda}^{(2)} \mu^{(i)}$ also satisfies $x_2 \in W$. This then implies the statement in part (b). Recall that $\overline{W}_d \leq W$ for $d \in \mathbb{D}_{\text{even}}$. Hence, without loss of generality we can assume that $Q^2(\widehat{\lambda}) \subseteq \mathcal{O}_d$ for some $d \in \mathbb{D}_{\text{odd}} \setminus \{-1\}$.

For $I_0 \in \mathcal{O}$ and $\kappa \in \text{Irr}_{\text{cusp}}(\tilde{L}_{I_0})$ we set

$$\mathcal{O}_\kappa(\hat{\lambda}) := \{I \in \mathcal{O} \mid \hat{\lambda}_I \text{ is } V_S\text{-conjugate to } \kappa \text{ or } \kappa^{c_{I_0}}\}.$$

Let μ_I be defined as in Lemma 5.14. Then $x_2(\mathcal{O}_\kappa(\hat{\lambda})) = \mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})$, see 5.14(a). With $\overline{\mathcal{O}}_\kappa(\hat{\lambda}) := \mathcal{O}_\kappa(\hat{\lambda}) \cup \mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})$ the element x can be written as product of $x_{\overline{\mathcal{O}}_\kappa(\lambda)} \in \mathcal{S}_{\pm\overline{\mathcal{O}}_\kappa(\lambda)}$ where $I \in \mathcal{O}$ and $\kappa \in \text{Irr}_{\text{cusp}}(\tilde{L}_I)$ runs over the $\langle \mu_I \rangle \times \langle c_I \rangle$ -orbits in $\text{Irr}_{\text{cusp}}(\tilde{L}_I)$. For proving $x_2 \in W$ it is sufficient to prove $x_{\overline{\mathcal{O}}_\kappa(\lambda)} \in W$. Hence we assume $Q^2(\hat{\lambda}) = \mathcal{O}_\kappa(\hat{\lambda}) \cup \mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})$ for some $\kappa \in \text{Irr}(\tilde{L}_{I_0})$.

If $I_0 \in \mathcal{O}_1$, we observe that $o(\kappa) \notin \{1, 2, 4\}$ by the definition of $Q^2(\hat{\lambda})$. This implies $\kappa\mu_{I_0} \notin \{\kappa, \kappa^{c_{I_0}}\}$ and hence $\mathcal{O}_\kappa(\hat{\lambda}) \cap \mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda}) = \emptyset$. Note that $\overline{W}^2(\hat{\lambda}) \leq W$. The element x_2 satisfies $x_2(\mathcal{O}_\kappa(\hat{\lambda})) = \mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})$ as element of $\mathcal{S}_{Q^2(\hat{\lambda})}$. Recall $\hat{\lambda}$ is standardized. Let $I \in \mathcal{O}_\kappa(\hat{\lambda})$ and $I' \in \mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})$. If $\hat{\lambda}_I$ and $\hat{\lambda}_{I'}$ are V_S -conjugate,

$$x_2(\epsilon\mathcal{O}_\kappa(\hat{\lambda})) = \epsilon\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda}) \text{ and } x_2(\epsilon\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})) = \epsilon\mathcal{O}_\kappa(\hat{\lambda}) \quad (5.3)$$

for every $\epsilon \in \{\pm 1\}$ as element of $\mathcal{S}_{\pm Q^2(\hat{\lambda})}$. Otherwise

$$x_2(\epsilon\mathcal{O}_\kappa(\hat{\lambda})) = -\epsilon\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda}) \text{ and } x_2(\epsilon\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})) = -\epsilon\mathcal{O}_\kappa(\hat{\lambda}) \quad (5.4)$$

for every $\epsilon \in \{\pm 1\}$ as element of $\mathcal{S}_{\pm Q^2(\hat{\lambda})}$. In both cases we see $x_2 \in W$.

Assume $I_0 \in \mathcal{O}_d$ for $d \in \mathbb{D}_{\text{odd}} \setminus \{\pm 1\}$. Hence $L_{I_0} \cong \text{GL}_d(q)$ by Lemma 3.2 and c_{I_0} acts on L_{I_0} as a graph automorphism by Lemma 3.4(c). According to Lemma 5.15 we have $\kappa \neq \kappa^{c_{I_0}}$ and $\overline{W}^2(\hat{\lambda}) \leq W$. Proposition 4.1(b) leads to $\kappa\mu_{I_0} \notin \{\kappa, \kappa^{c_{I_0}}\}$. We see again that $\mathcal{O}_\kappa(\hat{\lambda})$ and $\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})$ are disjoint. This implies again that there exists some $\epsilon' \in \{\pm 1\}$ such that

$$x_2(\epsilon\mathcal{O}_\kappa(\hat{\lambda})) = \epsilon'\epsilon\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda}) \text{ and } x_2(\epsilon\mathcal{O}_{\kappa\mu_{I_0}}(\hat{\lambda})) = \epsilon'\epsilon\mathcal{O}_\kappa(\hat{\lambda}) \quad (5.5)$$

for every $\epsilon \in \{\pm 1\}$ as element of $\mathcal{S}_{\pm Q^2(\hat{\lambda})}$. Again $x_2 \in W$. Altogether this proves part (b).

The considerations above imply $|\mathcal{O}_\kappa(\hat{\lambda})| = |\mathcal{O}_{\kappa\mu_I}(\hat{\lambda})|$ for every $I \in \mathcal{O}$ and $\kappa \in \text{Irr}_{\text{cusp}}(\tilde{L}_I)$. If $\mathcal{O}_\kappa(\hat{\lambda}) \subseteq Q^2(\hat{\lambda})$, the sets are disjoint such that $2 \mid |\overline{\mathcal{O}}_\kappa(\hat{\lambda})|$. This also applies if $I \in \mathcal{O}_1$ and $\kappa \in \text{Irr}_{\text{cusp}}(\tilde{L}_I)$ with $o(\kappa) \mid 2$. This implies part (c). \square

We study first the Clifford theory $W(\hat{\lambda}) \triangleleft W(\lambda)$ via considering subgroups associated to $Q^1(\hat{\lambda})$ and $Q^2(\hat{\lambda})$.

Lemma 5.33 *In the situation of Lemma 5.32 maximal extendibility holds with respect to $W^2(\hat{\lambda}^{(2)}) \triangleleft N_{\overline{W}^2}(W(\hat{\lambda}^{(2)}))$.*

Proof. Without loss of generality we can assume that $\hat{\lambda}$ is standardized. The structure of $\overline{W}(\hat{\lambda})$ is then given by Lemma 5.12. As in the proof of Lemma 5.26(a) the groups $W^2(\hat{\lambda}^{(2)})$ and $N_{\overline{W}^2}(W(\hat{\lambda}^{(2)}))$ satisfy Lemma 3.6 and accordingly maximal extendibility holds. \square

Proposition 5.34 *Let $\lambda \in \text{Irr}_{\text{cusp}}(L \mid 1_{(h_0)})$, $\hat{\lambda} \in \text{Irr}(\hat{L} \mid \lambda)$ and $\eta_0 \in \text{Irr}(W(\hat{\lambda}))$ with $\tilde{L}_\lambda = \hat{L}$.*

- (a) *If $Q^1(\hat{\lambda}) = \mathcal{O}$, then every $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$ is $K(\lambda)_{\eta_0}$ -stable.*
- (b) *Maximal extendibility holds with respect to $W^1(\hat{\lambda}^{(1)}) \triangleleft K^1(\lambda^{(1)})$.*

Proof. The statement in (a) is trivial if $W(\widehat{\lambda}) = W(\lambda)$. Hence we assume in the following $W(\widehat{\lambda}) \neq W(\lambda)$ and $Q^1(\widehat{\lambda}) = \mathcal{O}$. According to Lemma 5.30, $-1 \in \mathbb{D}$ and $\widehat{\lambda}_{-1}$ is $c_{J_{-1}}$ -stable, i.e.,

$$|\overline{W}_{-1}(\widehat{\lambda})| = 2.$$

For studying those groups we introduce more notation: For $j \in \{1, 2, 4\}$ let $Q^{1,j} = \{I \in \mathcal{O}_1 \mid o(\widehat{\lambda}_I) = j\}$ and $l_j := |Q^{1,j}|$. Then

$$\overline{W}^{Q^{1,j}}(\widehat{\lambda}^{(1)}) = \begin{cases} \mathcal{S}_{\pm Q^{1,j}} & \text{if } j \in \{1, 2\}, \\ \mathcal{S}_{Q^{1,4}} & \text{if } j = 4. \end{cases}$$

Accordingly $\overline{W}^1(\widehat{\lambda}) = \overline{W}_{-1}(\widehat{\lambda}) \times \mathcal{S}_{\pm Q^{1,1}} \times \mathcal{S}_{\pm Q^{1,2}} \times \mathcal{S}_{Q^{1,4}}$. Additionally $\overline{W}_{\widehat{\lambda}^{E_L}]_{\widehat{L}}}$ stabilizes the sets J_{-1} and $Q^{1,j}$ ($j \in \{1, 2, 4\}$). If $W(\lambda) \neq W(\widehat{\lambda})$, every $x \in W(\lambda) \setminus W(\widehat{\lambda})$ satisfies $x(Q^{1,1}) = Q^{1,2}$ as element of $\mathcal{S}_{Q^1(\widehat{\lambda})}$, see the proof of Lemma 5.32. Hence in that case $l_1 = l_2$.

Following the arguments given in the proof of Lemma 5.32 the element $x_1 \in W^1(\lambda) \setminus W^1(\widehat{\lambda})$ can be written as $x_{-1}x_{\{1,2\}}x_4$, where $x_{-1} \in \langle (J_{-1}, -J_{-1}) \rangle$, $x_{\{1,2\}} \in \mathcal{S}_{\pm(Q^{1,1} \cup Q^{1,2})}$ with $x_{\{1,2\}}(Q^{1,1}) = Q^{1,2}$ as element of $\mathcal{S}_{(Q^{1,1} \cup Q^{1,2})}$ and $x_4 \in \langle x_4^\circ \rangle$ with

$$x_4^\circ := \prod_{i \in Q^{1,4}} (i, -i) \in \mathcal{S}_{\pm Q^{1,4}}.$$

Note that $-1 \in \mathbb{D}$ according to Lemma 5.30 and hence $\gamma \in E_L$. In this notation we have

$$K(\lambda) \leq K^1 := \langle (J_{-1}, -J_{-1}) \rangle \times ((\mathcal{S}_{\pm Q^{1,1}} \times \mathcal{S}_{\pm Q^{1,2}}) \rtimes \langle x_{\{1,2\}} \rangle) \times (\mathcal{S}_{Q^{1,4}} \rtimes \langle x_4^\circ \rangle).$$

We see $W(\widehat{\lambda}) \cong \mathcal{S}_{\pm Q^{1,1}} \times \mathcal{S}_{\pm Q^{1,2}} \times \mathcal{S}_{Q^{1,4}}$. Since $K(\lambda) = \overline{W}_{\lambda^{L(\mathbb{F}_p)}]_L$, we have

$$K(\lambda) \leq \langle (J_{-1}, -J_{-1}) \rangle \times ((\mathcal{S}_{\pm Q^{1,1}} \times \mathcal{S}_{\pm Q^{1,2}}) \rtimes \langle x_{\{1,2\}} \rangle) \times \langle x_4 \rangle \mathcal{S}_{Q^{1,4}} \leq W(\widehat{\lambda}) \langle x, c_{-1}, x_4^\circ \rangle$$

for the element x from above and with $c_{-1} := \rho_{\mathbf{T}}(c_{J_{-1}})$. Note $W(\widehat{\lambda}) \langle x, c_{-1}, x_4 \rangle \leq K^1 = W(\lambda) \langle c_{-1}, x_4^\circ \rangle$. We observe $c_{-1}, x_4^\circ \in \mathbf{Z}(W(\lambda) \langle c_{-1}, x_4^\circ \rangle)$. This implies that every character $\eta \in \text{Irr}(W^1(\lambda))$ is K^1 -stable. This proves part (a), even that every character of $\text{Irr}(W(\lambda))$ extends to K^1 .

Now by the definition of $W^1(\widehat{\lambda}^{(1)})$ and $K^1(\lambda^{(1)})$ we see that in the general case the groups obtained as $K^1(\lambda)$ coincide with $K(\lambda)$ for a group of smaller rank where for the character $\widehat{\lambda}^{(1)}$ part (a) can be applied. This then proves part (b). \square

We consider the general case.

Proposition 5.35 *Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$ with $\widetilde{L}_\lambda = \widehat{L}$ and $\eta_0 \in \text{Irr}(W(\widehat{\lambda}))$. Then every character in $\text{Irr}(W(\lambda) \mid \eta_0)$ is $K(\lambda)_{\eta_0}$ -stable.*

Proof. Note that because of $|\widehat{L}_\lambda : L| = 2$ it is sufficient to prove that some character in $\text{Irr}(W(\lambda) \mid \eta_0)$ is $K(\lambda)_{\eta_0}$ -stable. According to Lemma 5.33 and Proposition 5.34 we can assume $Q^1(\widehat{\lambda}) \neq \mathcal{O} \neq Q^2(\widehat{\lambda})$ for the sets $Q^1(\widehat{\lambda})$ and $Q^2(\widehat{\lambda})$ from Lemma 5.32.

By Lemma 5.31 we can assume $h_0 \in \ker(\lambda)$. The groups $W^i(\widehat{\lambda})$ ($i \in \underline{2}$) satisfy $W(\widehat{\lambda}) = W^1(\widehat{\lambda}) \times W^2(\widehat{\lambda})$, see Lemma 5.32.

If $\mathcal{O}_1 \cap Q^1(\widehat{\lambda}) \neq \emptyset$, then $K(\lambda) \leq K^1(\lambda) \times N_{\overline{W}^2}(W^2(\widehat{\lambda}^{(2)}))$, see Lemma 5.32(d). Let $\eta_i \in \text{Irr}(W^i(\widehat{\lambda}))$ such that $\eta_0 = \eta_1 \times \eta_2$. According to Proposition 5.34, η_1 has a $K^1(\lambda)_{\eta_1}$ -stable extension to $W^1(\lambda)_{\eta_1}$ and maximal extendibility holds with respect to $W^2(\widehat{\lambda}) \triangleleft N_{\overline{W}^2}(W^2(\widehat{\lambda}^{(2)}))$ according to Lemma 5.33. This proves the statement in that case.

If $\mathcal{O}_1 \cap Q^1(\widehat{\lambda}) = \emptyset$, then $Q^1(\widehat{\lambda}) = \{J_{-1}\}$. Then $|W^1(\widehat{\lambda})| = 1$ and hence the character of $W(\lambda)$ has the required property. \square

Together with Proposition 5.28 this leads to the following statement.

Corollary 5.36 *Let $\lambda \in \mathbb{T}$ with $\widetilde{L}_\lambda \neq L$, $\widetilde{\lambda} \in \text{Irr}(\widetilde{L}_\lambda \mid \lambda)$ and $\eta_0 \in \text{Irr}(W(\widetilde{\lambda}))$ there exists some $K(\lambda)_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$.*

Proof. For $\lambda \in \mathbb{M}(\widetilde{L}) \cup \mathbb{M}_0$ this is Proposition 5.28. For $\lambda \in \text{Irr}_{\text{cusp}}(L)$ with $\widetilde{L}_\lambda = \widehat{L}$ the statement follows from Proposition 5.35. \square

6 Proof of Theorem A

In the following we explain how Corollary 5.36 about the action of $K(\lambda)$ on $\text{Irr}(W(\lambda))$ proves Theorem 5.1. As already sketched in the beginning of Section 5 based on Proposition 1.11, knowing the action of $\widehat{K}(\lambda)$ on $\text{Irr}(W(\lambda))$ is crucial to verify Theorem 5.1. Unless \mathbf{G} is of type D_4 the action $\widehat{K}(\lambda)$ on $\text{Irr}(W(\lambda))$ is given by the action of $K(\lambda)$, see Lemma 5.5.

Via Harish-Chandra induction we transfer this result on characters of N to a weak version of Theorem A. Special considerations are needed to determine the stabilisers in $\widetilde{G}E$ of characters $\chi \in \text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$, whenever L is not $E(\mathbf{G}^F)$ -stable.

Lemma 6.1 *Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$, $\widetilde{\lambda} \in \text{Irr}(\widetilde{L}_\lambda \mid \lambda)$ and $\eta_0 \in \text{Irr}(W(\lambda))$. Then some character in $\text{Irr}(W(\lambda) \mid \eta_0)$ is $\widehat{K}(\lambda)_{\eta_0}$ -stable.*

Proof. According to Corollary 5.36 there exists some $K(\lambda)_{\eta_0}$ -stable η in $\text{Irr}(W(\lambda) \mid \eta_0)$. According to Lemma 5.5 the character η is $\widehat{K}(\lambda)_{\eta_0} \cap (W \rtimes E_L^\circ)$ -stable, where $E^\circ := \langle F_p, \gamma_0 \rangle$ and $E_L^\circ := E^\circ \cap E_L$. If \mathbf{G} is not of type D_4 or $E_L \leq \langle F_p, \gamma_0 \rangle$ this is the above statement.

Accordingly we can assume in the following that \mathbf{G} is of type D_4 and L is γ_3 -stable for the graph automorphism γ_3 of $D_4(\mathbb{F})$ from 2.3. If $L = T_0$ the statement follows from Theorem 3.7 of [MS16]. Otherwise L is one of the two possible γ_3 -stable Levi subgroups. In both cases easy calculations show that $W(L)$ is a 2-group. According to our considerations above we know that there is some $K(\lambda)_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$. As $[K(\lambda) : W(\lambda)] \in \{1, 2\}$ the character η_0 extends to its inertia group $K(\lambda)_{\eta_0}$. This shows that maximal extendibility holds with respect to $W(\widetilde{\lambda}) \triangleleft K(\lambda)$. Let $K'(\lambda) = (\widehat{K}(\lambda) \langle F_p \rangle) \cap (W \rtimes \langle \gamma, \gamma_3 \rangle)$. When we identify \overline{W} with $W \rtimes \langle \gamma \rangle$, we can see $K(\lambda)$ as a subgroup of $K'(\lambda)$ with index 1 or 3. Hence $K(\lambda)_{\eta_0}$ has index 1 or 3 in $K'(\lambda)_{\eta_0}$. The character η_0 extends to $K(\lambda)_{\eta_0}$ by the above.

Assume that $K(\lambda)_{\eta_0}/W(\widetilde{\lambda})$ is a Sylow 2-subgroup of $K'(\lambda)/W(\widetilde{\lambda})$. Let K_3 be a subgroup of $K'(\lambda)_{\eta_0}$ with $W(\lambda)_{\eta_0} \leq K_3$ such that $K_3/W(\lambda)_{\eta_0}$ is a Sylow 3-subgroup $K'(\lambda)_{\eta_0}/W(\widetilde{\lambda})$. The character η_0 extends to K_3 as $|W(\widetilde{\lambda})|$ is coprime to 3 according to [I, (11.32)]. This implies that η_0 extends to $K(\lambda)_{\eta_0}$. maximal extendibility holds with respect to $W(\widetilde{\lambda}) \triangleleft K'(\lambda)$ as well. (This can be seen via an application of [I, (11.31)].)

If $K(\lambda)_{\eta_0}/W(\widetilde{\lambda})$ is not a Sylow 2-subgroup of $K'(\lambda)_{\eta_0}/W(\widetilde{\lambda})$, the group $K(\lambda^{\gamma_3})_{\eta_0^{\gamma_3}} = (K(\lambda)_{\eta_0})^{\gamma_3}$ contains a Sylow 2-subgroup of $K'(\lambda^{\gamma_3})_{\eta_0^{\gamma_3}}$ and hence by the above $(\eta_0)^{\gamma_3}$ extends to $K'(\lambda^{\gamma_3})_{\eta_0^{\gamma_3}}$. Via conjugation this implies that η_0 extends to $K'(\lambda)_{\eta_0}$. \square

We can now show Theorem 5.1.

Proof of Theorem 5.1. Recall $\mathbb{M}^{(X)} := \{\lambda \in \text{Irr}_{\text{cusp}}(L) \mid \tilde{L} = X\}$ for the subgroups $L \leq X \leq \tilde{L}$ and $\mathbb{M}_0 := \text{Irr}_{\text{cusp}}(L) \setminus (\mathbb{M}^{(L)} \cup \mathbb{M}^{(\tilde{L})} \cup \mathbb{M}^{(\tilde{L})})$, see before Lemma 5.3. As the sets $\mathbb{M}^{(L)}$, $\mathbb{M}^{(\tilde{L})}$, $\mathbb{M}^{(\tilde{L})}$ and \mathbb{M}_0 are E_L -stable, it is sufficient to construct an \tilde{N} -stable \tilde{L} -transversal in $\text{Irr}(N \mid \mathbb{M}')$ for $\mathbb{M}' \in \{\mathbb{M}^{(L)}, \mathbb{M}^{(\tilde{L})}, \mathbb{M}^{(\tilde{L})}\}$. Note that since every character of N is N -stable, one can equivalently also construct E_L -stable \tilde{N}' -transversals. Lemma 5.3 provides an \tilde{N} -stable \tilde{L} -transversal in $\text{Irr}(N \mid \mathbb{M}^{(L)})$.

Lemma 6.1 shows that for every $\lambda \in \mathbb{M}_0 \cup \mathbb{M}^{(\tilde{L})} \cup \mathbb{M}^{(\tilde{L})}$ and every character $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there is some $K(\tilde{\lambda})_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda) \mid \eta_0)$, where $\tilde{\lambda} \in \text{Irr}(\tilde{L}_\lambda \mid \lambda)$.

Assumptions (i) and (ii) of Proposition 1.11 are satisfied with $\mathbb{T}' := \mathbb{T} \cap (\mathbb{M}_0 \cup \mathbb{M}^{(\tilde{L})} \cup \mathbb{M}^{(\tilde{L})})$ from Proposition 3.10 and the extension map Λ from Theorem 2.1. For every $\lambda \in \mathbb{T}'$ and $\eta_0 \in \text{Irr}(W(\tilde{\lambda}))$ there exists some $K(\lambda)_{\eta_0}$ -stable $\eta \in \text{Irr}(W(\lambda))$. This allows to apply Proposition 1.12 and hence some \tilde{N} -stable \tilde{N} -transversal in $\text{Irr}_{\text{cusp}}(N \mid \mathbb{M}_0 \cup \mathbb{M}^{(\tilde{L})} \cup \mathbb{M}^{(\tilde{L})})$ exists. \square

Theorem 5.1 implies according to Theorem 1.8 that the equation $(\tilde{G}E_L)_\chi = \tilde{G}_\chi(E_L)_\chi$ holds for every character χ of a \tilde{G} -transversal in $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$. Accordingly we have constructed an E_L -stable \tilde{G} -transversal of $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$, see Lemma 1.4.

Corollary 6.2 *Let $\mathbf{G} = \text{D}_{l, \text{sc}}(\mathbb{F})$, $F : \mathbf{G} \rightarrow \mathbf{G}$ a standard Frobenius endomorphism and E be defined as in 2.3 and $\tilde{G} := \mathcal{L}^{-1}(Z(\mathbf{G}))$ for the Lang map defined by F . Let L be a standard Levi subgroup of \mathbf{G}^F and E_L its stabilizer in $E(\mathbf{G}^F)$. If Hypothesis 1.14 holds for every $l' < l$, then there exists an E_L -stable \tilde{G} -transversal in $\text{Irr}(\mathbf{G}^F \mid (L, \text{Irr}_{\text{cusp}}(L)))$.*

Proof. For a given fixed Levi subgroup L we apply Theorem 1.8 whose assumptions follow from Theorem 2.1 and Theorem 5.1. \square

Condition $A'(\infty)$ from 1.3 and equivalently Theorem A require to replace in the above statement E_L by E and study $(\tilde{G}E)_\chi$. Hence we study the stabilizers of characters in $\text{Irr}(\mathbf{G}^F \mid (L, \text{Irr}_{\text{cusp}}(L)))$ in the case where L is a standard Levi subgroup that is not E -stable.

Proposition 6.3 *We keep $\mathbf{G} = \text{D}_{l, \text{sc}}(\mathbb{F})$ and assume Hypothesis 1.14 holds for every $l' < l$. Let \mathbf{T} and \mathbf{L} as in Notation 1.5. Let $E^\circ := \langle F_p, \gamma \rangle \leq E$ in the notation of 2.3. Assume that no $N_{\mathbf{G}^F}(\mathbf{T})$ -conjugate of \mathbf{L} is E -stable. Let $\chi \in \text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$. Then $\tilde{G}_\chi = \tilde{G}$ or $(\tilde{G}E^\circ)_\chi \leq \tilde{G}_\chi(E^\circ \cap E_L)$.*

Proof. Let $N_0 := N_{\mathbf{G}^F}(\mathbf{T})$. We consider first the possible structure of L , in particular the values of $\mathbb{D}(L)$. Then we give the possible values of \tilde{L}_χ via describing $W(\lambda)$.

We see that L is F_p -stable. If L is γ -stable, then $E(\mathbf{G}^F) = E_L$. By our assumption $E_L \neq E$ we have $\gamma \notin E_L$. We observe that then $-1 \notin \mathbb{D}(L)$, as otherwise the system of simple roots Δ' , associated to L as in 2.5 is γ -stable, which then implies $\gamma \in E_L$.

If $1 \in \mathbb{D}(L)$, then some N_0 -conjugate of L is γ -stable: The conjugation is given by some element $v \in N_0 := N_{\mathbf{G}^F}(\mathbf{T}_0)$ that corresponds to some $\sigma \in \mathcal{S}_{\pm l}$ with the following properties: $\sigma(l) = \underline{l}$, $1 \in \sigma(J_1)$ and $\sigma(\Delta') \subseteq \Delta$. The Levi subgroup l satisfies accordingly $1 \notin \mathbb{D}(L)$.

Let $W_0 = N_0/\mathbf{T}^F$. For the proof of the statement we consider $\chi \in \text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$ with $(\tilde{G}E)_\chi \leq \tilde{G}E_L$. Then χ^γ and χ are $\tilde{G}E_L$ -conjugate. For the statement we have to

show that $\tilde{G}_\chi = \tilde{G}$. We assume that χ^γ and χ are $\tilde{G}E_L$ -conjugate. Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$ with $\chi \in \text{Irr}(G \mid (L, \lambda))$. Then (L, λ) and $(\gamma(L), \lambda^\gamma)$ are $\tilde{G}E_L$ -conjugate, in particular $\gamma(L)$ and L are N_0 -conjugate. This shows $\overline{W}(L) \neq W(L)$, or equivalently $\mathbb{D}_{\text{odd}} \neq \emptyset$. Let $\mathcal{O}_{\text{odd}} = \bigcup_{d \in \mathbb{D}_{\text{odd}}} \mathcal{O}_d$, $\mathcal{O}_{\text{even}} = \bigcup_{d \in \mathbb{D}_{\text{even}}} \mathcal{O}_d$ and $I_0 \in \mathcal{O}_{\text{odd}}$. Without loss of generality we assume $1 \in I_0$. Otherwise we replace L by some N_0 -conjugate. Let $w_0 := \prod_{i \in I_0} (i, -i) \in W_0$ and $n_0 \in \overline{N} = N_{\overline{\mathbf{G}}^F}(\mathbf{T})$ the corresponding element. Hence $w_0 \in \gamma N_0$. We observe that N induces on L the outer automorphisms W , while any $w' \in \overline{W} \setminus W$ is induced by elements of $N_{\mathbf{G}^F \langle \gamma \rangle}(L) \setminus N$. This proves that in this case L and $\gamma(L)$ are actually N_0 -conjugate. Hence the Harish-Chandra series satisfy $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L))) = \text{Irr}(G \mid (\gamma(L), \text{Irr}_{\text{cusp}}(\gamma(L))))$.

Let $\lambda \in \text{Irr}_{\text{cusp}}(L)$. Assume that (L, λ) and $(L, \lambda)^{e\gamma}$ are G -conjugate for some $e \in \langle F_p \rangle$. This implies that $(L, \lambda) = (L, \lambda)^{e\overline{n}}$ for some $\overline{n} \in \overline{N} \setminus N$. Note that $W(\lambda)^{\overline{n}} = W(\lambda)$. Because of $-1 \notin \mathbb{D}(L)$ and $\mathbb{D}_{\text{odd}} \neq \emptyset$ we observe $\tilde{L}_\lambda = \tilde{L}$, as $\langle \hat{L}_{I_0}, t_{l,2} \rangle \leq C_{\mathbf{G}}(L)L$.

Assume $h_0 \in \ker(\lambda)$ and that some $\chi \in \text{Irr}(\mathbf{G}^F \mid (L, \lambda))$ satisfies $\tilde{G}_\chi \neq \tilde{G}$. According to Corollary 5.22 the equation $W(\hat{\lambda}) = W(\tilde{\lambda})$ holds, as $\{\pm 1\} \cap \mathbb{D}(L) = \emptyset$. If $W(\lambda) = W(\hat{\lambda})$, then $\tilde{G}_\chi = \tilde{G}$. Hence we assume $W(\lambda) \neq W(\hat{\lambda})$ in the following. Let $W^{\text{odd}}(\lambda) := (W(\lambda)\mathcal{S}_{\pm\mathcal{O}_{\text{even}}}) \cap \mathcal{S}_{\pm\mathcal{O}_{\text{odd}}}$. Without loss of generality we can assume λ to be standardized, as in every \overline{N} -orbit in $\text{Irr}_{\text{cusp}}(L)$ is at least one standardized character, see after 5.10. As $W(\lambda) \neq W(\hat{\lambda})$ and λ is standardized, $W^{\text{odd}}(\hat{\lambda}) \leq \mathcal{S}_{\mathcal{O}_{\text{odd}}}$ and $x \in W(\lambda) \setminus W(\hat{\lambda})$ can be chosen as an involution with no fixpoint in \mathcal{O}_{odd} . We observe $N_{\mathcal{S}_{\pm\mathcal{O}_{\text{odd}}}}(W^{\text{odd}}(\lambda)) \leq W$. This implies $N_{\overline{W}}(W(\lambda)) \leq W$ and hence (L, λ) and $(L, \lambda)^{e\gamma}$ are not G -conjugate for any $e \in \langle F_p \rangle$. This proves that $(L, \lambda)^\gamma$ is not G -conjugate to any element of the E_L -orbit of (L, λ) , when $h_0 \in \ker(\lambda)$.

Assume $h_0 \notin \ker(\lambda)$ and $\nu \in \text{Irr}(\lambda|_{Z(G)})$. In the following we assume $|Z(G)| = 4$. We observe that then $E_\nu^\circ = \langle F_p \rangle = C_E(Z(G))$ and hence $(\tilde{G}E)_\chi \leq \tilde{G}E_\nu^\circ$ for every $\chi \in \text{Irr}(G \mid (L, \lambda))$. If $C_E(Z(G)) = E_L$, this implies $(\tilde{G}E)_\chi \leq \tilde{G}E_L$ as required. Note that if $2 \mid l$, then $C_E(Z(G)) = E_L$. In the following we prove $2 \mid |\mathcal{O}_{\text{odd}}|$ as this implies $2 \mid l$.

For $I \in \mathcal{O}$ let Z_I be defined as in 3.2, $Z_I := Z_I^F$ and $\delta_I \in \text{Irr}(\lambda|_{Z_I})$. Fix $d \in \mathbb{D}_{\text{odd}}$ and $I_d \in \mathcal{O}_d$. For $\kappa \in \text{Irr}(Z_{I_d})$ we define

$$a_\kappa(\lambda) := |\{I \in \mathcal{O}_d \mid \kappa \text{ and } \delta_I \text{ are } \overline{V}_d\text{-conjugate}\}|.$$

Recall $I_d \in \mathcal{O}_{\text{odd}}$ and $h_0 \notin \ker(\lambda)$. Hence $Z_{I_d} \cong C_{q-1}$ and $o(\delta_{I_d})_2 = (q-1)_2$. On the other hand we see that

$$\sum_{\kappa} a_\kappa(\lambda) = |\mathcal{O}_d|,$$

where κ runs over the $\langle c_{I_d} \rangle$ -orbits in $\text{Irr}(Z_{I_d})$. By the above $a_\kappa(\lambda) = 0$ for every $\kappa \in \text{Irr}(Z_{I_d})$ with $o(\kappa)_2 \neq (q-1)_2$. If $\lambda' \in \text{Irr}_{\text{cusp}}(L)$ is \overline{N} -conjugate to λ then $a_\kappa(\lambda) = a_\kappa(\lambda')$ for every $\kappa \in \text{Irr}(Z_{I_d})$ according to the action of \overline{V}_d on the groups Z_I ($I \in \mathcal{O}_d$). Note that $a_\kappa(\lambda) = 0$ as $o(\delta_I)_2 = (q-1)_2$ for every $I \in \mathcal{O}_d$.

Recall that we assume (L, λ) and $(\gamma(L), \lambda^{\gamma e})$ are G -conjugate. As the order of γe is even, we can choose some $F_0 \in \langle F_p \rangle$ such that $\langle F_0 \rangle$ is a Sylow 2-subgroup of $\langle e \rangle$. Then the \overline{N} -orbit of λ is F_0 -stable. Then F_0 acts on the characters of $\text{Irr}(Z_{I_d})$, inducing an action on the set of $\langle c_{I_d} \rangle$ -orbits in $\text{Irr}(Z_{I_d})$. We denote this set by $\text{Irr}(Z_{I_d})/\langle c_{I_d} \rangle$. If $\kappa \in \text{Irr}(Z_{I_d})$ with $o(\kappa)_2 = (q-1)_2$, the $\langle c_{I_d} \rangle$ -orbit of κ is not F_0 -stable. Hence the F_0 -orbit in $\text{Irr}(Z_{I_d})/\langle c_{I_d} \rangle$ containing κ has an even length. Since the \overline{N} -orbit of λ is F_0 -stable, we see that $a_\kappa(\lambda^{F_0}) = a_{\kappa^{F_0}}(\lambda)$. Accordingly

$$2 \mid \sum_{\kappa} a_\kappa(\lambda),$$

whenever κ runs over a $\langle c_{I_d} \rangle$ -transversal in $\{\kappa' \in \text{Irr}(Z_{I_d}) \mid o(\kappa')_2 = (q-1)_2\}$. By the above $a_\kappa(\lambda) = 0$ for every $\kappa \in \text{Irr}(Z_{I_d})$ with $o(\kappa)_2 \neq (q-1)_2$. Altogether this implies $2 \mid \sum_\kappa a_\kappa(\lambda) = |\mathcal{O}_d|$, where $\kappa \in \text{Irr}(Z_{I_d})$ runs over a $\langle c_{I_d} \rangle$ -transversal.

As $l = \sum_{d \in \mathbb{D}} d|\mathcal{O}_d|$ and hence $l \equiv |\mathcal{O}_{\text{odd}}| \pmod{2}$, the rank l is even and $\text{C}_E(\text{Z}(G)) = \langle F_p \rangle = E_L$. As explained above this leads to $(\tilde{G}E)_\chi \leq \tilde{G}E_{\chi|_{\text{Z}(G)}} = \tilde{G}E_L$ and hence a contradiction to the assumption on χ .

It remains to study the case of $|\text{Z}(G)| = 2$. Then $2 \nmid l$ and $4 \nmid (q-1)$. Note that $4 \nmid (q-1)$ implies $2 \nmid |E_L|$ and hence $|E_L|$ and $o(\gamma) = 2$ are coprime. If (L, λ) and $(\gamma(L), \lambda^\gamma)$ are GE_L -conjugate, then the pairs are already G -conjugate. In the following we see that the G -orbit of (L, λ) can not be γ -stable. By the above we have $\mathbb{D}_{\text{odd}} \neq \emptyset$ and $-1 \notin \mathbb{D}(L)$. According to 3.3, λ is \hat{L} -stable, as each λ_I is \hat{L}_I -stable for $I \in \mathcal{O}_{\text{odd}}$. Hence λ is \tilde{L} -stable, see also Lemma 3.3. According to Lemma 5.14 the assumption $2 \nmid l$ implies $W(\lambda) = W(\hat{\lambda})$, even more $\overline{W}(\lambda) = \overline{W}(\hat{\lambda}) \leq W$. But this implies that (L, λ) and $(L, \lambda)^\gamma$ are not G -conjugate. Hence $\gamma \notin (\tilde{G}E)_\chi$ and hence $(\tilde{G}E)_\chi \leq \tilde{G}E_{L, \chi}$. \square

A last obstacle is formed by the groups $D_{4,sc}(q)$. We keep the same notation.

Proposition 6.4 *If $G = D_{4,sc}(q)$, every \tilde{G} -orbit of $\text{Irr}(G)$ contains some χ with $(\tilde{G}E)_\chi = \tilde{G}_\chi E_\chi$.*

Proof. Let $\chi_0 \in \text{Irr}(G)$ and $E^\circ := \langle \gamma, F_p \rangle$. Then some Sylow 2-subgroup of E is contained in E° . We can assume that $\tilde{G}E^\circ/G$ contains a Sylow 2-subgroup of $(\tilde{G}E^\circ)_\chi/G$. (Otherwise we can replace χ_0 by some of its E -conjugates.) Some \tilde{G} -conjugate χ of χ_0 satisfies $(\tilde{G}E^\circ)_\chi = \tilde{G}_\chi E_\chi^\circ$ according to Proposition 6.3. This proves the statement if $(\tilde{G}E)_\chi \leq \tilde{G}E^\circ$. Additionally, $(\tilde{G}E)_\chi = \tilde{G}_\chi E_\chi$ holds if $\tilde{G}_\chi = \tilde{G}$.

Accordingly there is some $f \in \langle F_p \rangle$ and $t \in \tilde{G}$ such that χ is $\gamma_3 f t$ -stable and $\gamma_3 f t$ has 3-power order in $\tilde{G}E/G$. If $t \in G$ the equation $(\tilde{G}E)_\chi = \tilde{G}_\chi E_\chi$ holds. Clearly $\tilde{G}_\chi \triangleleft (\tilde{G}E)_\chi$. Hence \tilde{G}_χ is normalized by $\gamma_3 f t$. But via the isomorphism $\tilde{G}/G = \text{Z}(G)$ we see that $\tilde{G}_\chi = G$, as there is no $\gamma_3 f$ -stable subgroups of $\text{Z}(G)$ apart from $\{1\}$ and $\text{Z}(G)$.

The element $\gamma_3 f$ acts on $\text{Z}(G)$ such that only the trivial element is fixed by $\gamma_3 f$ and $[\gamma_3 f, \text{Z}(G)] = \text{Z}(G) \setminus \{1\}$.

Hence some \tilde{G} -conjugate χ' of χ satisfies $\gamma_3 f \in (\tilde{G}E)_{\chi'}$. We observe that $\gamma_3 f$ is a 3-element and hence $o(f)$ is a power of 3. Note that f acts trivially on $\text{Z}(G)$. Since χ satisfies $(\tilde{G}E^\circ)_\chi = GE_\chi^\circ$ and $[\text{Z}(G), F_p] = 1$, this leads to $(\tilde{G}E^\circ)_{\chi'} \in \{GE_\chi^\circ, G \langle F_p, \gamma \hat{t} \rangle_{\chi'}\}$ for some $\hat{t} \in \tilde{G}$ with $\mathcal{L}(\hat{t}) = h_0$. Let $\hat{G} := \mathcal{L}^{-1}(\langle h_0 \rangle)$. In the latter case

$$\hat{t}^{-1} \gamma_3(\hat{t}) f^2 = \gamma_3 f (\gamma_3 f)^{\gamma \hat{t}} \in (\hat{G}E)_{\chi'}.$$

Recalling that the orders of $\hat{G}/(\text{Z}(\hat{G})G)$ and f are coprime, we observe $\hat{t}^{-1} \gamma_3(\hat{t}) \in \langle \hat{t}^{-1} \gamma_3(\hat{t}) f^2 \rangle$, but $\hat{t}^{-1} \gamma_3(\hat{t}) \in \hat{G}_\chi$ and $\hat{t}^{-1} \gamma_3(\hat{t}) \notin G$. This leads to a contradiction and we see that $(\tilde{G}E^\circ)_{\chi'} = GE_\chi^\circ$ and hence $(\tilde{G}E)_{\chi'} = G \langle E_{\chi'}^\circ, \gamma_3 f \rangle$. \square

We can now deduce Theorem A from Theorem 1.8.

Proof of Theorem A. For a given fixed Levi subgroup L of G we apply Theorem 1.8 whose assumptions follow from Theorem 2.1 and Theorem 5.1. In this way we obtain an E_L -stable \tilde{G} -transversal in $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusps}}(L)))$, see Corollary 6.2. If L is E -stable, $E_L = E$ and this gives the required statement. If L has an E -stable G -conjugate L' , then we

observe that $\text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L))) = \text{Irr}(G \mid (L', \text{Irr}_{\text{cusp}}(L')))$ and there exists an E -stable \tilde{G} -transversal in $\text{Irr}(G \mid (L', \text{Irr}_{\text{cusp}}(L')))$.

It remains to consider the case where $E_L \neq E$ and no G -conjugate of L is E -stable. Then according to Propositions 6.3 and 6.4 every $\chi' \in \text{Irr}(G \mid (L, \text{Irr}_{\text{cusp}}(L)))$ has some \tilde{G} -conjugate χ with $(\tilde{G}E)_\chi = \tilde{G}_\chi E_\chi$. \square

References

- [B] C. BONNAFÉ, *Representations of $\text{SL}_2(\mathbb{F}_q)$* . Springer, London, 2011.
- [B06] C. BONNAFÉ, *Sur les caractères des groupes réductifs finis à centre non connexe: applications aux groupes spéciaux linéaires et unitaires*. Astérisque **306** (2006).
- [BS20] J. BROUGH AND B. SPÄTH, On the Alperin-McKay conjecture for simple groups of type A. *J. Algebra* **558**, 221–259 (2020).
- [BDT20] O. BRUNAT, O. DUDAS AND J. TAYLOR, Unitriangular shape of decomposition matrices of unipotent blocks. *Ann. of Math. (2)* **192**, 583–663 (2020).
- [C] R.W. CARTER, *Finite Groups of Lie Type*. John Wiley & Sons Inc., New York, 1985.
- [CE] M. CABANES AND M. ENGUEHARD, *Representation Theory of Finite Reductive Groups*. Cambridge University Press, Cambridge, 2004.
- [CS13] M. CABANES AND B. SPÄTH, Equivariance and extendibility in finite reductive groups with connected center. *Math. Z.* **275**, 689–713 (2013).
- [CS17a] M. CABANES AND B. SPÄTH, Equivariant character correspondences and inductive McKay condition for type A. *J. Reine Angew. Math.* **728**, 153–194 (2017).
- [CS17b] M. CABANES AND B. SPÄTH, The inductive McKay condition for type C. *Represent. Theory* **21**, 61–81 (2017).
- [CS19] M. CABANES AND B. SPÄTH, Descent equalities and the inductive McKay condition for types B and E. *Adv. Math.* **356**, 106820 (2019).
- [CSS21] M. CABANES, A. SCHAEFFER FRY AND B. SPÄTH, On the inductive Alperin-McKay conditions in the maximally split case. to appear in *Math. Z.*, (2021).
- [DM20] F. DIGNE AND J. MICHEL, *Representations of Finite Groups of Lie Type*. 2nd Edition. Cambridge University Press, Cambridge, 2020.
- [G93] M. GECK, A note on Harish-Chandra induction. *Manuscripta Math.* **80**, 393–401 (1993).
- [GM] M. GECK AND G. MALLE *The Character Theory of Finite Groups of Lie Type: A Guided Tour*. Cambridge University Press, Cambridge, 2020.
- [GLS] D. GORENSTEIN, R. LYONS, AND R. SOLOMON, *The Classification of the Finite Simple Groups. Number 3*. American Mathematical Society, Providence, RI, 1998.
- [I] I.M. ISAACS, *Character Theory of Finite Groups*. Academic Press, New York, 1976.
- [IMN07] I.M. ISAACS, G. MALLE, AND G. NAVARRO, A reduction theorem for the McKay conjecture. *Invent. Math.* **170**, 33–101 (2007).

- [K20] M. KLUPSCH, On Extending Unipotent Representations to their Stabilizers. *Preprint* arXiv:2012.02580, (2020).
- [L] G. LUSZTIG, *Characters of Reductive Groups over a Finite Field*. Princeton University Press, 1984.
- [L88] G. LUSZTIG, On the representations of reductive groups with disconnected centre, *in: Orbites unipotentes et représentations, I*. Astérisque **168**, 157–166 (1988).
- [L08] G. LUSZTIG, Irreducible representations of finite spin groups. *Represent. Theory* **12**, 1–36 (2008).
- [Ma10] J. MASLOWSKI, Equivariant character bijections in groups of Lie type. Dissertation, <http://kluedo.ub.uni-kl.de/volltexte/2010/2537/>, Technische Universität Kaiserslautern, 2010.
- [M14] G. MALLE, On the inductive Alperin-McKay and Alperin weight conjecture for groups with abelian Sylow subgroups. *J. Algebra* **397**, 190–208 (2014).
- [M17] G. MALLE, Cuspidal characters and automorphisms. *Adv. Math.* **320**, 887–903 (2017).
- [MS16] G. MALLE AND B. SPÄTH, Characters of odd degree. *Ann. Math.* **184**, 869–908 (2016).
- [MT] G. MALLE AND D. TESTERMAN, *Linear Algebraic Groups and Finite Groups of Lie Type*. Cambridge University Press, Cambridge, 2011.
- [N] G. NAVARRO, *Character theory and the McKay Conjecture*. Cambridge University Press, Cambridge, 2018.
- [NT11] G. NAVARRO AND P. TIEP, A reduction theorem for the Alperin weight conjecture, *Invent. Math.* **184**, 529–565, (2011).
- [OV] A. ONISHCHIK AND È. VINBERG, *Lie groups and algebraic groups*. Springer, Berlin, 1990.
- [S09] B. SPÄTH, The McKay conjecture for exceptional groups and odd primes. *Math. Z.* **261**, 571–595 (2009).
- [S10a] B. SPÄTH, Sylow d -tori of classical groups and the McKay conjecture. I. *J. Algebra* **323**, 2469–2493 (2010).
- [S10b] B. SPÄTH, Sylow d -tori of classical groups and the McKay conjecture. II. *J. Algebra* **323**, 2494–2509 (2010).
- [S12] B. SPÄTH, Inductive McKay condition in defining characteristic. *Bull. Lond. Math. Soc.* **44**, 426–438 (2012).
- [S13] B. SPÄTH, A reduction theorem for the Alperin-McKay conjecture. *J. Reine Angew. Math.* **680**, 153–189 (2013).
- [S17] B. SPÄTH, A reduction theorem for Dade’s projective conjecture. *J. Eur. Math. Soc. (JEMS)* **19** no. 4, 1071–1126 (2017).
- [S21a] B. SPÄTH, Extensions of characters in type D and the inductive McKay condition, II. Preprint, 2021.
- [S21b] B. SPÄTH, Extensions of characters in type D and the inductive McKay condition, III. In preparation, 2021.

Index

- $A'(\infty)$, 6
 $A(\infty)$, 6
 γ , 14
 γ_0 , 14
 γ_3 , 14
 $\lambda_1 \cdot \lambda_2$, 6
 $\otimes_{i \in \mathbb{I}} H_i$, 6
 $\overline{W}^j(\widehat{\lambda})$, 44
 α_1 , 13
 α_2 , 13
 α_i , 13

 $\|\cdot\|$, 17
 a_d , 15
 $\text{Aut}(\mathbf{G}^F)_{L,\text{HC}}$, 7
 automorphism
 diagonal, 11

 B , 6
 \mathbf{B} , 6
 \widetilde{B} , 6

 character
 cuspidal of \widehat{L}_I , 36
 restricted, 5
 standardized, 34
 $c_{I_d,j}$, 34
 $\text{Irr}_{\text{cusp}}(\widehat{L}_I)$, 36
 cuspidal character of \widehat{L}_I ,
 36

 D , 20
 \mathbb{D} , 15
 \mathbb{D}' , 15
 $\mathbb{D}(L)$, 15
 Δ , 13
 Δ' , 14
 $\delta_{\lambda,\sigma}$, 7
 \mathbb{D}_{even} , 15, 26
 diagonal automorphism,
 11
 \mathbb{D}_{odd} , 15, 26

 E , 6
 $E(\mathbf{G}^F)$, 14
 E_L° , 33
 $E(\mathbf{G}^F)$, 6
 $\mathcal{E}(G, (L, \lambda))$, 7

 e_i , 13
 E_L , 7
 ϵ_d , 22
 ϵ_{-1} , 24
 $E(\text{SL}_n(q))$, 12
 extendibility
 maximal, 5
 extension map, 5

 F , 6
 f , 14
 F_0 , 6
 F_{p^i} , 14
 \mathbb{F} , 6
 $f_k^{(d)}$, 18
 $f_k^{\circ,(d)}$, 18

 G , 6
 \mathbf{G} , 6, 13
 $\overline{\mathbf{G}}$, 13
 $\text{gcd}(i_1, i_2)$, 14, 21
 \widehat{G} , 11, 30
 G_I , 15
 \mathbf{G}_I , 15
 \widetilde{G} , 6
 $\widetilde{\mathbf{G}}$, 6

 H , 15
 H_0 , 15, 22
 h_0 , 15
 \widetilde{H}_0 , 19
 $\mathbf{h}_\alpha(t')$, 13
 Harish-Chandra
 induction, 7

 H_d , 15
 \widehat{H}_d , 22
 \widetilde{H}_d , 22
 \widetilde{H}_d , 15
 H_ϵ , 15
 H_ϵ , 26
 H_I , 19
 $\mathbf{h}_I(t)$, 15
 \widetilde{H}_I , 19
 \widetilde{H}_1 , 25
 \widetilde{H}_ϵ , 26
 \widetilde{H}_ϵ , 26

 (i_1, i_2) , 14

 $I_{d,j}$, 18
 $I_{d,j}(k)$, 18
 induced character, 5
 $\iota : \mathbf{G} \rightarrow \widetilde{\mathbf{G}}$, 6
 $\text{Irr}_{\text{cusp}}(L)$, 7
 $\text{Irr}_{\text{cusp}}(\widehat{L})$, 34
 $\text{Irr}_{\text{cusp}}(N)$, 7
 $\text{Irr}(G \mid (L, \lambda))$, 7
 $\text{Irr}(G \mid (L, \mathbb{T}))$, 7
 $\text{Irr}(\kappa)$, 5

 J_d , 15

 K , 15
 \mathbb{K} , 26
 K_{-1} , 24
 K_0 , 15
 \overline{K}_{-1} , 18
 κ_d , 19
 \overline{K}_d , 18
 $K_{0,-1}$, 24
 K_d , 21, 22
 $K_{0,d}$, 22
 $\widehat{K}(\lambda)$, 9
 $K^i(\lambda)$, 44
 $K^j(\lambda)$, 40
 $K(\lambda)$, 33

 L , 7, 14
 Λ_0 , 26
 Λ_ϵ , 27
 $\Lambda_{\epsilon_{-1}}$, 24
 Λ_{ϵ_d} , 23
 $\Lambda_{K \triangleleft M}$, 27
 Λ_L , 7
 $\Lambda_{L \triangleleft N}$, 13
 λ^σ , 5
 $\overline{\lambda^Y}$, 5
 (λ, η) , 8
 \mathbf{L} , 7
 \mathbf{L}_d , 16
 \mathcal{L} , 11
 $\widetilde{\mathcal{L}}$, 16
 $\widetilde{\mathcal{L}}$, 11
 L_0 , 16
 \widehat{L} , 16
 \widehat{L}_d , 16, 33
 $\widehat{L}^{(i)}$, 44

L_I , 16	\mathcal{P}° , 8	\tilde{T}_d , 23
$L^{(i)}$, 44	$\mathcal{P}^\circ(L)$, 8	$t_{I,2}$, 22
\mathbf{L}_I , 16	Φ , 13	$\text{type}(\Psi)$, 14
\widehat{L}_I , 16	Φ' , 14	
\widetilde{L} , 16	Φ_{-1} , 15	\dot{i} , 13
\widetilde{L}' , 7	Φ_d , 15	Υ , 8
	Φ_I , 15	Υ' , 8
M , 20	$\overline{\Phi}$, 13	
$m' \subset\subset M$, 35	$\overline{\Phi}_d$, 15	V , 20
\mathbb{M}_0 , 32	$\overline{\Phi}_I$, 15	\overline{V}_{-1} , 19
map	$\mathcal{P}(L)$, 8	ϖ , 14
extension, 5	$\psi \upharpoonright_X$, 5	V_D , 19
maximal extendibility, 5		V_I , 18
$\mathbb{M}^{(X)}$, 32	q , 14	\overline{V}_I , 18
	$Q^1(\widehat{\lambda})$, 40	\overline{V} , 19
N , 7	$Q^{1,j}(\lambda)$, 44	\overline{V}_0 , 18
\overline{N} , 17	$Q^1(\lambda)$, 44	\overline{V}_d , 19
N_0 , 6	$Q^2(\widehat{\lambda})$, 44	$\overline{V}_{d,S}$, 34
\overline{N}_0 , 17		\overline{V}_ϵ , 27
\widetilde{N}_0 , 17	$R(\lambda)$, 7	V_S , 34
\widetilde{N}'_0 , 6	restricted character, 5	\widetilde{V}_ϵ , 27
$\overline{\mathbf{n}}_1^{(d)}$, 22	$\rho_{a_d} : \overline{V}_{a_d} \rightarrow \mathcal{S}_{\pm a_d}$, 19	V'_{Weyl} , 18
$\overline{\mathbf{n}}_1$, 18	$\rho_{\mathbf{T}}$, 17	
N_0 , 17	R_L^G , 7	W , 7
$\mathbf{n}_\alpha(t')$, 13	$R_L^G(\lambda)_\eta$, 7	$W(\phi)$, 9
\mathbf{N}_0 , 6		$W^\circ(L)$, 18
\widehat{N} , 7	standardized character,	\widehat{W} , 9
$\overline{\mathbf{n}}_i$, 18	34	$W^j(\widehat{\lambda})$, 44
\widetilde{N}' , 7	$\mathcal{S}_{\pm M}$, 16	$W^j(\lambda)$, 40
\widetilde{N} , 32	$\mathcal{S}_{\pm M}^D$, 17	$W(\lambda)$, 7
\widetilde{N}_0 , 6		\overline{W} , 33
	\mathbb{T} , 26	\overline{W}_0 , 17
\mathcal{O}_1 , 15	\mathbf{T} , 14	\overline{W}_d , 18
$\mathcal{O}_{c,-1}(\widehat{\lambda})$, 35	$\overline{\mathbf{T}}$, 14	$\overline{W}_d(\widehat{\lambda})$, 34
$\mathcal{O}_{c,1}(\widehat{\lambda})$, 35	\mathbb{T}° , 24	$\overline{W}^Q(\kappa)$, 39
\mathcal{O} , 15	T_0 , 6	$W^Q(\kappa)$, 39
\mathcal{O}_d , 15	\mathbf{T}_0 , 6	
$\mathcal{O}_{c,\epsilon}$, 35	\widetilde{T}_0 , 6	
$\mathcal{O}_c(\widehat{\lambda})$, 35	$\widehat{\mathbf{T}}_0$, 6	
$\mathcal{O}_{\text{even}}$, 49	\mathbb{T}° , 26	
$\odot_{i \in \mathbb{I}} \lambda_i$, 6	\mathbb{T}°_d , 23	
$\mathcal{O}_\kappa(\widehat{\lambda})$, 45	\widehat{T} , 26	
\mathcal{O}_{odd} , 49	\mathbf{T}_I , 16	
$\overline{\mathcal{O}}_\kappa(\widehat{\lambda})$, 45	t_I , 22	
$o(\mu)$, 39	$t_{I,2}$, 35	
	t_{J-1} , 24	
\mathcal{P} , 8	$t_{J-1,2}$, 24	
p , 14	$\mathbb{T}^{(L)}$, 32	
		ζ , 16
		\mathbf{Z}_I , 21