

A CHARACTERIZATION OF STRONGLY MONOTYPIC POLYTOPES

VUONG BUI

ABSTRACT. We characterize all the strongly monotypic polytopes. Hadwiger's conjecture for this class of polytopes is deduced from the characterization.

1. INTRODUCTION

Monotypic polytopes and strongly monotypic polytopes were first introduced in [1]. Monotypic polytopes can be seen as a subclass of simple polytopes: A polytope P is monotypic if every polytope having the same set of normals as P is combinatorially equivalent to P .

On the other hand, strongly monotypic polytopes is a subclass of monotypic polytopes: A polytope P is strongly monotypic if every polytope Q having the same set of normals as P satisfies $\mathcal{A}(P)$ and $\mathcal{A}(Q)$ being combinatorial equivalent, where $\mathcal{A}(P)$ is the arrangement of hyperplanes containing the facets of P . An interesting property of such a polytope P is that: The intersection of any two translates of P is always a Minkowski summand of P . In other words, a strongly monotypic polytope is always a generating set. The other direction whether a polytope with the generating property is always a strongly monotypic polytope is still open. However, it is shown to be the case for \mathbb{R}^3 in [1].

Note that the property with a convex set in the place of a polytope and any multiple number of sets instead of 2 is actually the definition of generating sets. It is shown in [2] that to check if a set is generating, one just needs to check for every pair of its translates. That is we can replace every pair of translates by any collection of translates in the property of strongly monotypic polytopes.

The paper [1] gives a partial characterization of the strongly monotypic polytopes in \mathbb{R}^3 , which was later extended to a full characterization in [3]. In this paper, a treatment for higher dimensions will be given.

Monotypic polytopes and strongly monotypic polytopes P can be recognized by the set of normals $N(P)$. In [1], several equivalent versions of the necessary and sufficient conditions are given. In particular, Condition $M3'$ of a monotypic polytope P says that: If V_1 and V_2 are disjoint primitive subsets of $N(P)$ then $\text{pos } V_1 \cap \text{pos } V_2 = \{0\}$. (V is a primitive subset of $N(P)$ if V is linearly independent and $\text{pos } V \cap N(P) = V$.) Also, the condition $S4'$ for a strongly monotypic polytope P says that: If Q is any polytope with $N(Q) \subseteq N(P)$ then Q is monotypic.

In order to characterize the set of normals of strongly monotypic polytopes more conveniently, we give another equivalent condition.

Theorem 1. *An n -dimensional polytope P is monotypic if and only if every $n+1$ normals of P is either (i) not separated from 0 or (ii) if they are in a conical position, then their positive hull contains another normal of P .*

The author is supported by the Deutsche Forschungsgemeinschaft (DFG) Graduiertenkolleg "Facets of Complexity" (GRK 2434). The work was initiated during the time the author studied at Moscow Institute of Physics and Technology.

In this text, a set of points is said to be separated from 0 if there is a hyperplane strictly separating the set from 0. Also, a set of points is said to be in *conical position* if they are separated from 0 and there is no point in the positive hull of the rest.

Theorem 2. *An n -dimensional polytope P is strongly monotypic if and only if every $n + 1$ normals of P is either (i) not separated from 0 or (ii) one of them is in the positive hull of the others.*

These equivalences are verified in Section 2, by using Conditions $M3'$ and $S4'$.

For convenience, we give the following term with respect to Theorem 2.

Definition 1. *A set of points in \mathbb{R}^n is said to be in a good position if either their convex hull contains 0 or one of them is in the positive hull of the rest. Otherwise they are said to be in a bad position.*

Since two strongly monotypic polytopes of the same set of normals are combinatorially equivalent, we characterize their sets of normals instead of the polytopes themselves. The problem of characterizing the set of normals of strongly monotypic polytopes can be reduced to the following problem, due to Theorem 2.

Problem 1. *Characterise all the finite sets $X \subset \mathbb{R}^n \setminus \{0\}$ such that (i) 0 is in the interior of the convex hull of X , none $x \in X$ is a positive scale of another $y \in X$ and (ii) every $n + 1$ points of X are in a good position.*

The requirement (i) is actually to ensure that the set X is a set of normal vectors of a polytope. From now on, all the sets are understood as the set of normals, that is no element is a positive scale of another element. They may be normalized under some convention, for which the set of normals of a polytope is unique (e.g. the norm of each normal is 1, or Convention 1, which we actually use and describe later).

Throughout the text we only treat the problem for $n \geq 3$, partly because the problem is trivial on the plane and partly because some statements are not true on the plane. (Note that all polygons are strongly monotypic.)

We characterize X by showing that X must contain a “skeleton” as defined in the following theorem ¹.

Theorem 3. *Every X satisfying Problem 1 for $n \geq 3$ contains some k disjoint subsets X_1, \dots, X_k such that (i) each X_i is the vertices of a simplex whose relative interior contains 0, (ii) the linear spaces spanned by each X_i are linearly independent and (iii) these linear spaces directly sum up to \mathbb{R}^n . Such a collection of X_1, \dots, X_k is said to be a skeleton of X .*

We can assume that the linear spaces spanned by the sets X_i are orthogonal (otherwise applying an appropriate linear transform to obtain it). We also assume that the sum of the elements in each X_i is the origin. Theorem 3 is proved in Section 6.

The skeleton gives a bound for the number of translates in Hadwiger’s conjecture ².

Theorem 4. *If the set of normals X of a strongly monotypic polytope $P \subset \mathbb{R}^n$ ($n \geq 3$) has a skeleton X_1, \dots, X_k then P can be covered by at most $|X_1| \dots |X_k|$ translates of $(1 - \epsilon)P$ for a small enough $\epsilon > 0$.*

The proof of Theorem 4 does not use all the characterization of X but only a handful of propositions. It is perhaps not so easy to deduce Theorem 4 directly from Theorem

¹It does not hold in \mathbb{R}^2 .

²We are addressing the combinatorial geometry one, not the graph theoretical one.

2 since the later only provides a local picture of the normals while the former asks for a more global picture.

Hadwiger's conjecture for this class of polytopes follows as $|X_1| \dots |X_k| \leq 2^n$ for $P \subset \mathbb{R}^n$. Note that the case of convex bodies in \mathbb{R}^2 was confirmed in [4] by Levi.

Once we have the skeleton, we may have to add more points to obtain X . The following problem is the instance $k = 1$. Instances of higher k can be reduced to this problem as in Section 4.

Problem 2. *Given the vertices $E = \{e_0, \dots, e_n\}$ of a simplex in \mathbb{R}^n for $n \geq 3$ such that $e_0 + \dots + e_n = 0$ (i.e. 0 is in the interior), characterize all the set $X \supseteq E$ such that every $n + 1$ points of X are in a good position.*

The solution for Problem 2 itself is given in Section 5. After characterizing the set of normals, we have enough tools to prove Theorem 4, which is done in Section 6.

2. PROOF FOR THE EQUIVALENCE OF THE CHARACTERIZATIONS

Let D denote the condition in Theorem 1. In order to prove Theorem 1, we will show that the two definitions $M3'$ and D are equivalent by verifying both directions.

Proposition 2.1. *$M3'$ implies D .*

Proof. Suppose we do not have D , which means there exist $n + 1$ normals in conical position with the positive hull not containing any other normal. Let H be a hyperplane not through 0 such that for each a of the $n + 1$ normals, ray $\overrightarrow{0a}$ cuts the hyperplane at only a point. Replace every normal a by the intersection of ray $\overrightarrow{0a}$ and the hyperplane, apply Radon's theorem to this set on the affine space, there will be two disjoint subsets whose intersection is nonempty. It means in the linear space, we have V_1 and V_2 contradicting $M3'$ □

To show that D implies $M3'$, we need the following lemmas.

Lemma 1. *Given two disjoint primitive subsets V_1, V_2 of a set of normals. If $\text{pos } V_1$ and $\text{pos } V_2$ intersect at a point other than 0, then there are $V'_1 \subseteq V_1, V'_2 \subseteq V_2$ such that $(\text{pos } V'_1 \cap \text{pos } V'_2) \setminus \{0\}$ is precisely a ray in the relative interior of both the positive hulls. Moreover, the union $V'_1 \cup V'_2$ is separated from 0.*

Proof. Let $U_1 \subseteq V_1$ and $U_2 \subseteq V_2$ be minimal subsets so that $\text{pos } U_1 \cap \text{pos } U_2 \neq \{0\}$.

We show that $(\text{pos } U_1 \cap \text{pos } U_2) \setminus \{0\}$ is a ray in the relative interior of both the positive hulls. Suppose there are two rays $\overrightarrow{0p}, \overrightarrow{0p'}$ in the intersection. Rotating $0p$ to $0p'$ and going further, we will pass the relative boundary of one of the two positive hulls for the first time, say $\text{pos } U_1$, at sometime before getting out of the positive hull. Let $0q$ be the ray on the relative boundary at that time. We have a contradiction as we can set U_1 to be a smaller subset U_1 so that $0q$ is still in the positive hull of the subset. In fact, the only ray excluding 0 is in the relative interior of both $\text{pos } U_1$ and $\text{pos } U_2$, as otherwise U_1, U_2 can be set to be smaller subsets.

We continue to prove that is $U_1 \cup U_2$ separated from 0. Suppose $0 \in \text{conv}(U_1 \cup U_2)$, that is

$$-\sum_{x_i \in U_1} \lambda_i x_i = \sum_{x_i \in U_2} \lambda_i x_i,$$

for some negative coefficients λ_i .

Since $\text{pos } U_1 \cap \text{pos } U_2$ excluding 0 is a ray in the relative interior of both positive hulls,

$$\sum_{x_i \in U_1} \theta_i x_i = \sum_{x_i \in U_2} \theta_i x_i,$$

where all the coefficients θ_i are positive.

Let α be so that $\min\{-\lambda_i + \alpha\theta_i : x_i \in U_1\} = 0$ (that is $\alpha = \max\{\lambda_i/\theta_i : x_i \in U_1\}$), we have

$$\sum_{x_i \in U_1} (-\lambda_i + \alpha\theta_i)x_i = \sum_{x_i \in U_2} (\lambda_i + \alpha\theta_i)x_i,$$

with at least one coefficient on the left is 0. It means that $\text{pos } U_1$ also intersects $\text{pos } U_2$ at a point other than 0 and not in $\text{rel int pos } U_1$, contradiction.

The sets U_1, U_2 confirm the conclusion. \square

Lemma 2. *The followings are equivalent for a point set X that spans \mathbb{R}^n :*

(i) *There exist $n + 1$ points of X in conical position with the positive hull empty of other points of X .*

(ii) *For some $d \leq n$, there exist $d + 1$ points of X that are d -dimensional and in conical position with the positive hull empty of other points of X .*

Proof. The direction (i) \implies (ii) is trivial, just take any $(\dim \text{span } X) + 1$ points among them.

We show the other direction, that is (ii) \implies (i).

At first, we start with the given $d + 1$ points X' of dimension d , and increase their dimension one by one, by adding one point in each step, until we have $n + 1$ points. If the current number of points is less than $n + 1$, the dimension is therefore less than n , which means the existence of another point p not in the span of X' . If the positive hull of $X' \cup \{p\}$ is not empty of other points of X , replace p by any point of X in the positive hull other than those in X' , and recursively repeat it until the positive hull is empty of other points of X . The process will eventually terminate as the positive hull contains fewer points after each step. So, for any X' , we can increase the dimension by one, by adding one point but still keeping the positive hull empty of other points. Therefore, we can come up with $n + 1$ points satisfying the hypothesis, thus achieving (i). \square

Remark 1. *If we have (ii), we also have the same (ii) for every higher d , including n , which is itself a stronger statement than (i).*

Now comes the verification of D implies $M3'$:

Proposition 2.2. *D implies $M3'$.*

Proof. Suppose we do not have $M3'$, which means there are two disjoint primitive subsets V_1, V_2 of $N(P)$ such that $(\text{pos } V_1 \cap \text{pos } V_2) \setminus \{0\}$ is nonempty. Over all such pairs V_1, V_2 , we consider a pair so that $\text{pos}(V_1 \cup V_2)$ is minimal (up to inclusion). Due to Lemma 1, we have $(\text{pos } V_1 \cap \text{pos } V_2) \setminus \{0\}$ is a precisely a ray in the relative interior of both $\text{pos } V_1, \text{pos } V_2$. Moreover, $V_1 \cup V_2$ are separated from 0.

Consider a hyperplane that cuts every ray $0p$ for each $p \in V_1 \cup V_2$ at exactly a point p' . Let U_1, U_2 be the corresponding sets of those points p' . Although U_1, U_2 are contained in an affine space, it is more convenient to see the hyperplane as a linear space by taking $\text{conv } U_1 \cap \text{conv } U_2$ as the origin. Moreover, w.l.o.g. we assume the spaces spanned by U_1 and U_2 are orthogonal (otherwise, we transform the space). Let U be the intersection of all the normals in $\text{pos}(V_1 \cup V_2)$ with the hyperplane. Note that $\text{conv } U_1$ and $\text{conv } U_2$ are empty of other points in U (as V_1, V_2 are primitive).

Suppose we have D , that is $U \setminus (U_1 \cup U_2)$ is nonempty.

Let p be a point in $U \setminus (U_1 \cup U_2)$ that has the smallest distance to $\text{conv } U_1$. It follows that $\text{conv}(\{p\} \cup U_1)$ is empty of other points in U since otherwise another point would be closer to $\text{conv } U_1$ than p . Also note that the points $\{p\} \cup U_1$ are in convex position since p is not in the affine space spanned by U_1 .

Let p' be the projection of p onto $\text{conv } U_1$. Since U_1 is the vertices of a simplex containing 0, there is a proper subset $U'_1 \subset U_1$ so that $0 \in \text{conv}(\{p\} \cup U'_1)$. We then have $\text{conv}(\{p\} \cup U'_1)$ and $\text{conv } U_2$ intersecting. Their corresponding normals are the two sets satisfying the condition of V_1, V_2 but having a smaller positive hull of the union, contradiction. It follows that we do not have D . \square

We now have verified Theorem 1.

Proof of Theorem 1. Theorem 1 follows from Propositions 2.1 and 2.2. \square

Let DD denote the condition in Theorem 2, we show the equivalence of Conditions DD and $S4'$ in order to prove Theorem 2.

Proof of Theorem 2. In one direction, if a monotypic polytope P is also strongly monotypic, it should not have $n + 1$ normals in conical position (i.e. Condition DD). Indeed, suppose V is the set of such $n + 1$ normals, we consider the subset after removing every other normal in the positive hull (except the $n + 1$ normals), which is $N(P) \setminus ((\text{pos } V) \setminus V)$. Any polytope taking this subset as the set of normals is not monotypic, due to the existence of a positive hull of $n + 1$ normals of V in conical position but not containing any other normal.

In the other direction, if a polytope P has Condition DD as above, then every polytope Q whose set of normals is a subset of the normal set of P should be monotypic, too. It is rather clear from definitions D and DD , as the condition (ii) of DD means the hypothesis in the condition (ii) of D doesn't happen. \square

3. PROOF OF THEOREM 3

Among the elements of X take a set of points B such that B is in conical position and their positive hull is maximal in the sense that no other such points have the positive hull being a superset of the positive hull of B . Since every $n + 1$ points of X is in a good position, it follows that B is a set of n linearly independent points b_1, \dots, b_n .

Consider any other point $x = \sum_{i=1}^n \lambda_i b_i$ in X . The $n + 1$ points $\{x, b_1, \dots, b_n\}$ either (i) have the convex hull containing 0, for which all λ_i are nonpositive or (ii) have one of them in the positive hull of the rest, for which all λ_i are nonnegative. The other (impossible) possibility is the coefficients λ_i are of mixed signs, we then have precisely one positive coefficient, since if there are at least two positive coefficients, together with at least one negative coefficients, the signs of coefficients indicates that $\{x, b_1, \dots, b_n\}$ are in conical position, contradiction. As there are now only one positive coefficient, say λ_i , and other nonpositive coefficients, we have $b_i \in \text{pos}\{x, b_1, \dots, b_n\} \setminus \{b_i\}$ for some i ; a contradiction since the n points $\{x, b_1, \dots, b_n\} \setminus \{b_i\}$ has a bigger positive hull than the positive hull of $\{b_1, \dots, b_n\}$ (the former contains x while the latter does not). So, it is always either (i) or (ii).

Let the *Cartesian support*³ of a point x be the set $\{b_i : \lambda_i \neq 0\}$ for $x = \sum_{i=1}^n \lambda_i b_i$. We show that the Cartesian support of two points both in $\text{pos}\{b_1, \dots, b_n\}$ or both in $\text{pos}\{-b_1, \dots, -b_n\}$ are either disjoint or one is a subset of the other. Let the two points be x, y with $x = \sum_{i=1}^n \lambda_i b_i$ and $y = \sum_{i=1}^n \theta_i b_i$. Suppose the Cartesian support of x be

³The leading "Cartesian" is to distinguish with the other definition of support as in Definition 2, which dominates the rest of the text.

$\{b_1, \dots, b_{k_2}\}$ and the Cartesian support of y be $\{b_{k_1}, \dots, b_{k_3}\}$ with $1 < k_1 \leq k_2 < k_3$ (we need the assumption of \mathbb{R}^3 here). The unique linear relation (upto scaling)

$$x = \sum_{i=1}^{k_1-1} \lambda_i b_i + \frac{\lambda_{k_1}}{\theta_{k_1}} y + \sum_{i=k_1+1}^{k_2} (\lambda_i - \frac{\lambda_{k_1}}{\theta_{k_1}} \theta_i) b_i - \sum_{i=k_2+1}^{k_3} \frac{\lambda_{k_1}}{\theta_{k_1}} \theta_i b_i$$

raises a contradiction as the $n+1$ points $\{x, y\} \cup \{b_1, \dots, b_n\} \setminus \{b_{k_1}\}$ are in a bad position (one positive coefficient on the left hand side, two positive and one negative coefficients on the right hand side).

Since $0 \in \text{int conv } X$, there must be some k points x_1, \dots, x_k all in $\text{pos}\{-b_1, \dots, -b_n\}$ and their Cartesian supports are disjoint while the union of all Cartesian supports is $\{b_1, \dots, b_n\}$. Let X_i for each $i = 1, \dots, k$ be the union of $\{x_i\}$ and the Cartesian support of x_i , we obtain the desired X_1, \dots, X_k , which completes the proof.

4. REDUCTION OF PROBLEM 1 TO PROBLEM 2

The approach to Problem 1: We first start with a skeleton X_1, \dots, X_k of X and then try to constrain all the additional points in $X \setminus (X_1 \cup \dots \cup X_k)$. The following propositions give the necessary condition for the additional points. Their proofs are rather not so hard and will be given later in Section 7. Theorem 5 reduces the Problem 1 to Problem 2 when the remaining work is to characterize the points in certain spaces span X_i .

We begin with a definition.

Definition 2. *Given a skeleton X_1, \dots, X_k . If p_1, \dots, p_k is the projection of p onto the linear spaces spanned by X_1, \dots, X_k and $X'_i \subset X_i$ is the minimal subset of X_i such that p_i is in the positive hull of X'_i , then $S = X'_1 \cup \dots \cup X'_k$ is called the support of p .*

One can see that $|S_p|$ is at least 2 except for $p \in X_1 \cup \dots \cup X_k$, and $|S_p|$ cannot exceed n .

The following propositions are the necessary conditions for additional points in X .

Proposition 4.1. *If $x \in X$ is not in the space spanned by any X_i then $x \in \text{pos}\{x_i, x_j\}$ for some $x_i \in X_i, x_j \in X_j$.*

Proposition 4.2. *If each of $x, y \in X$ is not in the space spanned by any X_i with $x \in \text{pos}\{x_i, x_j\}$, $y \in \text{pos}\{y_{i'}, y_{j'}\}$, $x_i \in X_i, x_j \in X_j, y_{i'} \in X_{i'}, y_{j'} \in X_{j'}$, then the sets $\{i, j\}$ and $\{i', j'\}$ are either equal or disjoint.*

Proposition 4.3. *Given two points $x, y \in X$ not in the space spanned by any X_i with $x \in \text{pos}\{x_i, x_j\}$, $y \in \text{pos}\{y_i, y_j\}$, $x_i, y_i \in X_i, x_j, y_j \in X_j$. If $|X_i| > 2$ then $x_i = y_i$.*

Proposition 4.4. *Suppose $x \in X$ is not in the space spanned by any X_i , $x \in \text{pos}\{x_i, x_j\}$ for some $x_i \in X_i, x_j \in X_j$. If there is any other point $y \in X$ in the space spanned by X_i than those points in X_i , then the support of y cannot contain x_i .*

Proposition 4.5. *If $x \in X$ is not in the space spanned by any X_i , $x \in \text{pos}\{x_i, x_j\}$ for some $x_i \in X_i, x_j \in X_j$, then every point $y \in X$ in the space spanned by X_i is a positive scale of $\sum_{e_t \in S_y} e_t$ (we will use the convention that $y = \sum_{e_t \in S_y} e_t$). Further more, if p, q are in the space spanned by X_i then either their supports S_p, S_q are disjoint, or one is a subset of the other.*

An X_i such that there is a point $x \in X$ contained in the relative interior of $\text{pos}\{x_i, x_j\}$ for some $x_i \in X_i, x_j \in X_j$ is said to be an *involved* X_i , otherwise it is called an *uninvolved* X_i . If every $n+1$ points of X in the space spanned by such an X_i are always in a good position, then every $n+1$ points in X are always in a good position as in Theorem 5

below. That being said we have reduced Problem 1 to Problem 2. Note that our setting for the points in the space spanned by an involved X_i also satisfy the solution of Problem 2 (as the readers can compare with Section 5).

Theorem 5. *If the requirements in all the above propositions are satisfied and the points in the space spanned by an uninvolved X_i satisfy Problem 2, then the points of X satisfies Problem 1.*

Proof. Consider a set C of any $n + 1$ points of X . If the space of any uninvolved X_i has more than $\dim X_i$ points then the points are in a good position.

In the remaining case, there must be a pair X_i, X_j such that there are more points of C in the space spanned by $X_i \cup X_j$ than the dimension of $X_i \cup X_j$ (by Proposition 4.2).

Let us first start with a lemma.

Lemma 3. *Let X_i be uninvolved by a point in $\text{pos}\{x_i, x_j\}$ for $x_i \in X_i$. Consider some $|X_i| - 1$ points in the space spanned by an involved X_i , then either those points are in a good position, or we have a linear combination of all nonnegative coefficients with value $-x_i$, or x_i can be found among the considered points. Moreover, if $|X_i|$ points are considered, then those points are in a good position.*

Proof. (i) $|X_i| - 1$ points are considered.

If x_i is among the considered points, then the conclusion holds.

We assume otherwise. That is all the $\dim X_i$ points take only the subsets of $X_i \setminus \{x_i\}$ as their supports (by Proposition 4.4). Consider the forest structure defined by these supports where a support is a father of another support if the former is superset of the latter (by Proposition 4.5).

If the union of all the roots is $X_i \setminus \{x_i\}$, the conclusion holds as $-x_i$ is in the positive hull of the considered points. Indeed, let the points whose supports are the roots be p_1, \dots, p_t , then $\sum_j p_j = \sum_{e \in X_i \setminus \{x_i\}} e = -e_i$ (note that each p_j is the sum of elements in the support of p_j , by Proposition 4.5).

If the union of all the roots is a proper subset of $X_i \setminus \{x_i\}$, then more point are considered than the cardinality of the union of the roots. It follows that there is a vertex being the union of its children, which indicates a good position since the point taking the father as the support is the sum of the points taking the children as the supports (also by Proposition 4.5).

(ii) $|X_i|$ points are considered.

If the union of the roots is X_i , then the sum of of the points taking the roots as the supports is $\sum_{e \in X_i} e = 0$, which indicates a good position.

If the union of the root is a proper subset of X_i , then there are more points considered than the cardinality of the union of the root. We also have a good position in this case (as already reasoned in (i)). \square

We assume the dimension of $X_i \cup X_j$ is greater than 2, otherwise, any three points from the plane are in a good position.

In case one of X_i, X_j is 1-dimensional, we always assume it is X_j .

Case 0: If there is 0 point of C in $\text{pos}(X_i \cup X_j) \setminus \text{pos} X_i \cup \text{pos} X_j$, then the space spanned by either X_i or X_j has more points of C than its dimension, hence, they are in a good position.

Case 1: There is precisely 1 point of C in $\text{pos}(X_i \cup X_j) \setminus (\text{pos} X_i \cup \text{pos} X_j)$, then the point is in $\text{pos}\{x_i, x_j\}$ for some $x_i \in X_i, x_j \in X_j$.

We consider the case when $\text{span } X_i$ has $\dim X_i$ points of C and $\text{span } X_j$ has $\dim X_j$ points of C , as otherwise one of them has more points than the dimension, which indicates a good position.

Let the considered point be $y = \alpha x_i + \beta x_j$.

By Lemma 3 either $x_i \in C$ or $-x_i$ can be presented by a linear combination of the points in $C \cap \text{span } X_i$ with all nonnegative coefficients. We have the same for x_j . In either case, we always have a good position.

Case 2: There are precisely 2 points y, z of C in $\text{pos}(X_i \cup X_j) \setminus \text{pos } X_i \cup \text{pos } X_j$. Let one point is in $\text{pos}\{x_i, x_j\}$ for some $x_i \in X_i, x_j \in X_j$. The other point must be in the same positive hull, or in $\text{pos}\{x_i, -x_j\}$ in case $\text{pos } X_j$ is 1-dimensional (by Proposition 4.3). To cover all the cases, we say the two points are in $\text{pos}\{x_i, x_j\} \cup \text{pos}\{x_i, -x_j\}$.

The space spanned by either X_i or X_j must have at least as many points as its dimension.

(i) Suppose it is X_i . By Lemma 3 either $x_i \in C$ or $-x_i$ can be presented by a linear combination of the points in $C \cap \text{pos } X_i$ with all nonnegative coefficients. No matter x_i is in $\text{pos}\{y, z\}$ or $y \in \text{pos}\{x_i, z\}$ (or $z \in \text{pos}\{x_i, y\}$), with the representation of x_i by the points in $C \cap \text{pos } X_i$, we always have a good position.

(ii) If it is the case for X_j , then either x_j or $-x_j$ is in C . No matter $y \in \text{pos}\{x_j, z\}$ ($z \in \text{pos}\{x_j, y\}$) or $y \in \text{pos}\{-x_j, z\}$ ($z \in \text{pos}\{-x_j, y\}$), we always have a good position of the three points y, z, x_j or $y, z, -x_j$.

Case 3: If there are at least 3 points of C in $(\text{pos}\{x_i, x_j\} \cup \text{pos}\{x_i, -x_j\}) \setminus (\text{pos } X_i \cup \text{pos } X_j)$ for some $x_i \in X_i, x_j \in X_j$, then the three points are in a good position.

All the cases were covered, and the conclusion follows. \square

5. SOLUTION OF PROBLEM 2

We first give all the necessary conditions for additional points of $X \setminus E$ in the following propositions. Their proofs are not so hard and will be given in Section 7. Finally, Theorem 6 claims that these necessary conditions are also sufficient.

As two vectors with one being a positive scale of the other present the same normal, we first give a convention for convenience later.

Convention 1. A point $p = \sum_{e_i \in S_p} \lambda_i e_i$ (⁴) in X will be normalized so that $\min_{e_i \in S_p} \lambda_i = 1$.

Proposition 5.1. A point $p = \sum_{e_i \in S_p} \lambda_i e_i$ in X has all $\lambda_i = 1$ except at most one $\lambda_j > 1$.

Definition 3. Given $p = \sum_{e_i \in S_p} \lambda_i e_i$ in X , the point $e_j \in S_p$ such that $\lambda_j > 1$ if exists is said to be the strange index of p . If there is such p , we also say S_p has a strange index ⁵.

Proposition 5.2. Given two points p, q in X , then either (i) $S_p \cap S_q = \emptyset$, or (ii) one of S_p, S_q is a subset of the other, or (iii) $S_p \cup S_q = E$ and $|S_p \cap S_q| = 1$, or (iv) $S_p \cup S_q = E$ and $|S_p \cap S_q| = n - 1$.

Proposition 5.3. Given two points p, q in X having the same support $S_p = S_q$ of more than two points, then if both of them have strange indices, these indices are the same.

Proposition 5.4. Given two points p, q in X with $S_p \cap S_q = \emptyset$ then at most one of p, q can have a strange index.

⁴In this section, we use the same definition of support as in Definition 2 (for $k = 1$), and denote by S_p the support of p .

⁵For convenience, we use the point e_i for the strange index instead of the index i as the name may suggest.

Proposition 5.5. *Given two points p, q in X with $S_p \subset S_q, |S_p| \geq 2, p = \sum_{e_i \in S_p} \lambda_i e_i, q = \sum_{e_i \in S_q} \theta_i e_i$. We then have $\lambda_i = \theta_i$ for every $e_i \in S_p$.*

Proposition 5.6. *Given two points p, q in X with $S_p \cup S_q = E$ and $S_p \cap S_q = \{e_k\}$. If each of S_p, S_q has at least 3 points, then the strange index of any point of p, q , if any, is e_k . If $|S_p| = 2$ then the strange index of S_q , if any, is e_k .*

Proposition 5.7. *Given two points p, q in X with $S_p \cup S_q = E$ and $|S_p \cap S_q| = n - 1$ then $\lambda_a > 1, \theta_b > 1$ and $\lambda_a \theta_b \geq \lambda_a + \theta_b$ where $\{e_a\} = S_p \setminus S_q, \{e_b\} = S_q \setminus S_p$ and $p = \sum_{e_i \in S_p} \lambda_i e_i, q = \sum_{e_i \in S_q} \theta_i e_i$.*

Proposition 5.8. *Given two points p, q in X with $S_p \cup S_q = E, |S_p \cap S_q| = 1$ and $|S_p| \geq 3, |S_q| \geq 3$, then for every point $r \in X$ with $|S_r| \geq 2$, the support S_r is a subset of either S_p or S_q .*

Proposition 5.9. *Given two points p, q in X with $S_p \cup S_q = E, |S_p \cap S_q| = n - 1$ and a point $r \in X$ with $S_r \neq S_p, S_r \neq S_q$ and $|S_r| \geq 2$, then S_r is either a subset of $S_p \cap S_q$ or $S_r = (S_p \setminus S_q) \cup (S_q \setminus S_p)$.*

Proposition 5.10. *Given two points p, q in X with $|S_p \cap S_q| = 1, |S_p| = 2, |S_q| = n$ and a point $r \in X$ with $S_r \neq S_p, S_r \neq S_q$ and $|S_r| \geq 2$, then S_r is either a subset of $S_q \setminus S_p$ or $S_r = (S_p \setminus S_q) \cup (S_q \setminus S_p)$.*

We show that the above necessary requirements are actually sufficient.

Theorem 6. *If a set X satisfies the conditions in all above propositions, then X also satisfies Problems 2.*

Proof. Take any $n + 1$ points of X , consider the graph whose vertices are the supports of the points and two supports are connected by an edge if one is a subset of the other. This graph is actually a partial order, but for convenience, we remove some edges if necessary to obtain a forest in the way that: If a vertex v is a subset of at least one vertex, then v is connected to precisely one of them. (Note that one support may correspond to more than one point.) From now on in this proof, we only treat these $n + 1$ points instead the whole X .

Lemma 4. *If a tree supports at least as many points as the cardinality of the root then either (i) one can obtain the sum of the elements in S in the root by a linear combination of the points with at most one negative coefficient (and zero negative coefficients when S has no strange index), or (ii) one of the points is in the positive hull of the rest. If the number of points is strictly greater than the number of vertices, then we always have (ii).*

Proof. Let S be the root.

The statement is obviously true when S contains only one point. Also, it is true when $|S| = 2$. Indeed, if there are two points x, y in $\text{pos}\{e_0, e_1\}$. The desired sum $e_0 + e_1$ is in the middle of the positive hull. Therefore, regardless of the position of x, y , the sum is always obtained by a linear combination of the two linearly independent vector x, y . Note that the coefficients are not all negative, since otherwise, $e_0 + e_1$ must be in the other quadrant $-\text{pos}\{e_0, e_1\}$. Also, when there is no strange index, x, y are actually e_0, e_1 , hence, there is no negative coefficient.

If there are 3 points then one of them is in the positive hull of the other two points since one vector is in the acute angle formed by the other two vectors.

We now prove the lemma for any $|S| \geq 3$ with the assumption that the lemma holds for S of lower cardinalities.

(a) Consider the case the tree supports exactly as many points as the cardinality of S .

When there is no strange index, we obtain (i) with the linear combination of only one summand which is the point supported by S . We only consider the case we have a strange index hereafter. Note that the strange index must be unique due to Proposition 5.5.

If there are 2 points whose support is S , then the strange indices, if both have, are the same by Proposition 5.3. Let the strange index be s_k , and the two points be $p = \sum_{e_i \in S} \lambda_i e_i$, $q = \sum_{e_i \in S} \theta_i e_i$ with $\lambda_k > \theta_k$, we then have

$$\frac{\lambda_k - 1}{\lambda_k - \theta_k} q - \frac{\theta_k - 1}{\lambda_k - \theta_k} p = \sum_{e_i \in S} e_i.$$

We treat the case when only one point $p = \sum_{e_i \in S} \lambda_i e_i$ is supported by S .

We have to consider only the case when no subtree supports more points than the cardinality of the root of that subtree, otherwise we have (ii) already by the induction hypothesis. Such a situation implies that every subtree supports precisely as many points as the cardinality of the root of that subtree. Also, the union of the children of a vertex S' cannot be S' but $S' \setminus \{e_j\}$ for some $e_j \in S'$.

Case 1: the point e_j is the strange index. Let S_1, S_2, \dots, S_m be the children of S . By induction hypothesis, we have a linear combination L_i of nonnegative coefficients whose value is the sum of elements in each S_i (note that no strange index is in any S_i). It follows that

$$\sum_{j=1}^m L_i = \sum_{e_i \in S \setminus \{e_j\}} e_i.$$

Now we have the representation of the desired sum

$$\left(1 - \frac{1}{\lambda_j}\right) \left(\sum_{t=1}^m L_t\right) + \frac{1}{\lambda_j} p = \sum_{e_i \in S} e_i.$$

Case 2: some e_k other than e_j is the strange index. Consider the smallest subtree whose root contains e_k . If that subtree is a leaf e_k itself, i.e. we have point e_k among the considered points, we have

$$p - (\lambda_k - 1)e_k = \sum_{e_i \in S} e_i.$$

If that subtree is rooted by T with some children (none of them has e_k), then the union of the children, say $T_1, \dots, T_{m'}$, is $T \setminus \{e_k\}$. By induction hypothesis, for each T_i there is a linear combination L_i of nonnegative coefficients whose result is the sum of elements in T_i . Let $q = \sum_{e_i \in T} \theta_i e_i$ be the point supported by T , we have

$$p + \frac{\lambda_k - 1}{\theta_k - 1} \left(\sum_{j=1}^{m'} L_i - q\right) = \sum_{e_i \in S} e_i.$$

(b) Consider the case the tree supports more points than the cardinality of S .

If there are 3 points whose support is S then one of them is in the positive hull of the other two.

Consider the case when there are only two points whose support is S .

If the strange index e_j is among the $n + 1$ points, one of the 2 points is in the positive hull of the other and e_j .

In the other case, dropping the root S and the two points, the cardinality of the union of the roots of the remaining trees is at most $|S| - 1$, while the number of points is reduced by 2. That is we have as many points as the cardinality of the union of the roots of the

remaining forest. We are done if some tree satisfies Case (ii) of the conclusion. If all the remaining subtrees satisfy Case (i), then one of the two points is in the positive hull of the other point and the remaining points.

If there is only 1 points supported by S , then one of the subtrees satisfies the requirement (b) of S . By induction hypothesis, we have Case (ii) of the conclusion. \square

Corollary 1. *If a forest with disjoint roots supports at least as many points as the cardinality of the union of the roots then either (i) one can obtain the sum of all e_i in the roots by a linear combination of the points with at most one negative coefficient (and zero negative coefficients when no root has a strange index), or (ii) one of the points is in the positive hull of the rest. If the number of points is strictly greater, then we always have (ii).*

If a tree of the forest supports more points than the cardinality of the root then we are done, by Conclusion (ii) of Lemma 4.

Consider the remaining case that each tree supports at most as many points as the cardinality of the root. The following cases are exclusive due to Propositions 5.2, 5.9 and 5.10.

Case 1: The roots are disjoint (and the union is E), thus each root supports precisely as many points as the cardinality of the root. We are done if Conclusion (ii) applies for any tree. If Conclusion (i) applies to all trees, then a linear combination of all points with at most one negative coefficient is $\sum_{e_i \in E} e_i = 0$, hence giving a good position. (Note that there is at most one support among the disjoint supports that has a strange index, by Proposition 5.4.)

Case 2: There are precisely two roots S_1, S_2 intersecting at only some e_k and the union is E .

Subcase (a): $|S_1|, |S_2| \geq 3$.

If e_k is among the $n + 1$ points, then e_k is in the positive hull of two points, each from one root since

$$p + q - (\lambda_k + \theta_k - 1)e_k = \sum_{e_i \in E} e_i = 0.$$

(Note that the strange index of any point of p, q , if any, is e_k , by Proposition 5.6.)

Consider the case there is no e_k among the $n + 1$ points, hence, no support containing e_k , except the two roots (by Proposition 5.8).

One of the two trees supports as many points as the cardinality of the root, say S_1 , since otherwise the total number of points is less than $n + 1$.

If there are two points supported by this root: Suppose the two points be $p = \sum_{i=0}^k \lambda_i e_i, p' = \sum_{i=0}^k \lambda'_i e_i$ with $\lambda_k < \lambda'_k$. Let a point supported by S_2 be $\sum_{i=k}^n \theta_i e_i$.

The linear relation

$$up + u'p' + q = \sum_{e_i \in E} e_i = 0,$$

where u, u' satisfy $u + u' = 1$ and $\lambda_k u + \lambda_k u' + \theta_k = 1$, shows that the three points p, p', q are in a good position (any three points on the same plane through 0 are in a good position).

If there is only one point p supported by the root S_1 , excluding p and the support itself, there remains a forest with at least as many points as the cardinality of the union of the roots. The union of the roots of the remaining forest is a subset of $S_1 \setminus \{e_k\}$. We are done if the union is a proper subset, since we have Conclusion (ii) of Corollary 1 then. It remains to consider the case the union is precisely $S_1 \setminus \{e_k\}$. The point p is then in the

positive hull of the other points of the tree of S_1 and q where $q = \sum_{e_i \in S_2} \theta_i e_i$ is a point supported by S_2 since

$$q - \frac{\theta_k - 1}{\lambda_k} p + (1 + \frac{\theta_k - 1}{\lambda_k}) L = \sum_{e_i \in E} e_i = 0,$$

where L is the linear combination of nonnegative coefficient with value being the sum of the elements in $S_1 \setminus \{e_k\}$ (by Corollary 1). (Note that there is no other strange index in $S_1 \setminus \{e_k\}$ by Proposition 5.5.)

Subcase (b): One of $|S_1|, |S_2|$ is 2, say $|S_1| = 2$.

Suppose $S_1 = \{e_0, e_1\}$, $S_2 = \{e_1, \dots, e_n\}$.

Consider the case S_1 supports two points $p = \sum_{e_i \in S_1} \lambda_i e_i$, $p' = \sum_{e_i \in S_1} \lambda'_i e_i$. The linear relation

$$up + u'p' + q = \sum_{e_i \in E} e_i = 0,$$

where u, u' satisfy $u\lambda_0 + u'\lambda'_0 = 1, u\lambda_1 + u'\lambda'_1 + \theta_1 = 1$, shows a good position of the considered points.

Consider the case S_2 supports two points $q = \sum_{i=1}^n \theta_i e_i$, $q' = \sum_{i=1}^n \theta'_i e_i$, the linear relation

$$uq + u'q' + \frac{1}{\lambda_0} p = \sum_{e_i \in E} e_i = 0,$$

where u, u' satisfy $u + u' = 1, u\theta_1 + u'\theta'_1 + \frac{\lambda_1}{\lambda_0} = 1$, shows a good position of the considered points. (Note that $\theta_1 \neq \theta'_1$, by Proposition 5.6.)

It remains to consider the case there are no two points both supported by either S_1 or S_2 .

If e_0 is among the considered points then the linear relation

$$up + ve_0 + q = \sum_{e_i \in E} e_i = 0,$$

where u, v satisfy $\lambda_0 u + v = 1$ and $\lambda_1 u + \theta_1 = 1$, shows a good position of the considered points.

If e_1 is among the points then the linear relation

$$up + ve_1 + q = \sum_{e_i \in E} e_i = 0,$$

where $\lambda_0 u = 1$ and $\lambda_1 u + v + \theta_1 = 1$, shows a good position of the considered points.

If neither e_0 nor e_1 is among the points then all other points than p, q are supported by subsets of $\{e_2, \dots, e_n\}$. Applying Corollary 1 to these $n - 1$ remaining points, one obtains a good position already (Conclusion (ii)) or otherwise a linear combination L whose result is the sum of e_2, \dots, e_n with at most one negative coefficient. The linear relation

$$\frac{1}{\lambda_0} p + \frac{1 - \frac{\lambda_1}{\lambda_0}}{\theta_1} q + (1 - \frac{1 - \frac{\lambda_1}{\lambda_0}}{\theta_1}) L = \sum_{e_i \in E} e_i = 0$$

shows a good position of the considered points.

Case 3: There are precisely two roots S_1, S_2 intersecting at precisely $n - 1$ points and their union is E . Suppose $S_1 = \{e_0, \dots, e_{n-1}\}$ and $S_2 = \{e_1, \dots, e_n\}$.

Consider the case there are two points $p = \sum_{i=0}^{n-1} \lambda_i e_i$, $p' = \sum_{i=0}^{n-1} \lambda'_i e_i$ of the same support S_1 . Let the point supported by S_2 be $q = \sum_{i=1}^n \theta_i e_i$. By Proposition 5.7, we have $\lambda_i = \lambda'_i = \theta_i = 1$ for $i = 1, \dots, n - 1$.

The linear relation

$$up + u'p' + vq = \sum_{e_i \in E} e_i = 0,$$

where u, u', v satisfy $u\lambda_0 + u'\lambda'_0 = u + u' + v = v\theta_n = 1$ indicates that p, p', q are in a good position. (Note that $\lambda_0 \neq \lambda'_0$).

The case S_2 supports two points is treated likewise.

It remains to check the case each S_i supports only one point.

As there is no other root, the support of every other point of the remaining $n - 1$ points is a subset of $\{e_1, \dots, e_{n-1}\}$ (by Proposition 5.9). By Corollary 1, we have either Case (i) or Case (ii) of the conclusion. We are done if we already have Case (ii), that is one point is in the positive hull of the rest. Otherwise, we can represent the sum of e_1, \dots, e_{n-1} as a linear combination L with at most one negative coefficient by Case (i). It follows that

$$(\lambda_0\theta_n - \lambda_0 - \theta_n)L + \theta_np + \lambda_0q = \sum_{e_i \in E} e_i = 0,$$

where the coefficient $\lambda_0\theta_n - \lambda_0 - \theta_n$ of L is nonnegative by Proposition 5.7.

Case 4: There are two roots S_1, S_2 as in Case 3 and additionally a root $S_0 = \{e_0, e_n\}$. Let $r = \phi_0e_0 + \phi_ne_n$ be the point supported by S_0 , the relation

$$up + vq + tr = \sum_{e_i \in E} e_i = 0,$$

where u, v, t satisfy $u\lambda_0 + t\phi_0 = 1, u + v = 1, v\theta_n + t\phi_n = 1$, indicates a good position of the considered points. (The readers can check that the determinant of the matrix corresponding to the system is $\phi_0\theta_n + \lambda_0\phi_n > 0$.) \square

6. PROOF OF THEOREM 4

The proof uses the characterization in Sections 4 and 5. The readers just need to check the propositions there.

At first, we revise an alternate formulation with illumination of Hadwiger's conjecture whose exposition can be found in [5, Chapter VI].

Proposition 6.1 (Boltyanskii). *A convex body P can be covered by h translates of $(1-\epsilon)P$ (for ϵ small enough) if and only if there exists a set H of h vectors such that for each point x on the boundary of P , at least one vector $v \in H$ gives $x + v \in \text{int } P$.*

A point x on the boundary of a polytope is contained in the intersection of some facets with the hyperplanes $\{x : \langle a_i, x \rangle = c_i\}$. The set of vectors v allows $x + v \in \text{int } P$ is

$$\bigcap_i \{v : \langle a_i, v \rangle < 0\}.$$

We can see that it is sufficient to find a set of vectors H for the vertices of P only, instead of all the points on the boundary.

We give a trivial observation.

Observation 1. *Given a set of normals $\{a_i\}_i$ and $v \in \bigcap_i \{v : \langle a_i, v \rangle < 0\}$, if $a \in \text{pos}\{a_i\}$ then $\langle a, v \rangle < 0$.*

The above observation easily implies the following result.

Lemma 5. *Given a set of normal vectors a'_1, \dots, a'_m contained in the positive hull of normals a_1, \dots, a_n , then*

$$\bigcap_i \{v : \langle a'_i, v \rangle < 0\} \supseteq \bigcap_i \{v : \langle a_i, v \rangle < 0\}.$$

By the property of the skeleton, the union of the positive hulls of $X'_1 \cup \dots \cup X'_k$ over all X'_i being a subset of X_i obtained by excluding an element $x_i \in X_i$ is \mathbb{R}^n , that is

$$\bigcup_{x_1 \in X_1, \dots, x_n \in X_n} \text{pos}\left(\bigcup_i X_i \setminus \{x_i\}\right) = \mathbb{R}^n.$$

From each of $|X_1| \dots |X_k|$ such sets of n linearly independent normals a_i , we pick any $v \in \bigcap_i \{v : \langle a_i, v \rangle < 0\}$ to add to the desired set H . Note that a vertex is the intersection of the facets of n primitive normals and no other facets as otherwise there is $n+1$ normals in conical position with the positive hull empty of other normals. Using Lemma 5, we can see that it suffices to show that every n primitive normals are contained in one of the $|X_1| \dots |X_k|$ positive hulls.

Assume otherwise, among the n normals there are two normals a, b not in the same positive hull. We investigate the possible cases and show that $\text{pos}\{a, b\}$ contains some normal of $X_1 \cup \dots \cup X_k$ or $0 \in \text{conv}\{a, b\}$ (which actually contradicts with the primitivity of the n normals).

Case 1: $a \in \text{pos } X_i, b \in \text{pos } X_j$ for different i, j .

The positive hull of $X'_1 \cup \dots \cup X'_k$ for $a \in X'_i, b \in X'_j$ contains both a, b .

Case 2: a, b both in $\text{pos } X_i$ for some i .

Since a, b are not in the positive hull of any $X'_i \subset X_i$, the union of supports is $S_a \cup S_b = X_i$. Let $X_i = \{e_0, \dots, e_m\}$.

By Proposition 5.2, we just need to treat the following situations.

Suppose $|S_a| = |S_b| = |X_i| - 1$, say $S_a = \{e_0, \dots, e_{m-1}\}, S_b = \{e_1, \dots, e_m\}, a = \sum_{i=0}^{m-1} \lambda_i e_i, b = \sum_{i=1}^m \theta_i e_i$, with $\lambda_0 > \lambda_1 = \dots = \lambda_{m-1} = 1, 1 = \theta_1 = \dots = \theta_{m-1} < \theta_m$ and $\lambda_0 \theta_m \geq \lambda_0 + \theta_m$ (by Proposition 5.7). One obtains either $0 \in \text{conv}\{a, b\}$ or $e_m \in \text{pos}\{a, b\}$ since

$$a + (\lambda_0 - 1)b = \left(\lambda_0 \sum_{i=0}^{m-1} e_i\right) + (\lambda_0 - 1)\theta_m e_m = ((\lambda_0 - 1)\theta_m - \lambda_0)e_m$$

with $(\lambda_0 - 1)\theta_m - \lambda_0 \geq 0$ as $\lambda_0 \theta_m \geq \lambda_0 + \theta_m$ (we have $0 \in \text{conv}\{a, b\}$ for the equality).

Suppose $|S_a \cap S_b| = 1$ with $|S_a| \leq |S_b|$: say $S_a = \{e_0, \dots, e_k\}, S_b = \{e_k, \dots, e_m\}, a = \sum_{i=0}^k \lambda_i e_i, b = \sum_{j=k}^m \theta_j e_j$ with $\theta_k \geq \theta_{k+1} = \dots = \theta_m$ (by Proposition 5.6).

If $1 = \lambda_0 = \dots = \lambda_{k-1} \leq \lambda_k$, then $e_k \in \text{pos}\{a, b\}$ as

$$a + b = (\lambda_k + \theta_k)e_k.$$

If $k = 1$ and $\lambda_0 > \lambda_1 = 1$, then $e_1 \in \text{pos}\{a, b\}$ as

$$\frac{1}{\lambda_0}a + b = \left(\frac{1}{\lambda_0} + \theta_1\right)e_1.$$

It remains to consider the case $S_a \cap S_b = \emptyset$. If neither a or b has a strange index, we have $a + b = 0$. Otherwise, only one of them has a strange index (by Proposition 5.4), say a with $\lambda_0 > 1$ for $a = \sum_i \lambda_i e_i, b = \sum_i \theta_i e_i$. We then have $a + b = (\lambda_0 - 1)e_0$, that is $e_0 \in \text{pos}\{a, b\}$, contradiction.

Case 3: $a \in \text{pos } X_i$ for some i but b is not in the span of any X_j .

If $b \in \text{pos}\{x_j, x_{j'}\}$ for $x_j \in X_j, x_{j'} \in X_{j'}$ (by Proposition 4.1) where none of j, j' is i , then the positive hull of $X'_1 \cup \dots \cup X'_k$ for $S_a \subseteq X'_i, x_j \in X'_j, x_{j'} \in X'_{j'}$ contains both a, b .

Consider the case $b \in \text{pos}\{x_i, x_j\}$ for $x_i \in X_i, x_j \in X_j$.

If $|S_a| < |X_i| - 1$, or $|S_a| = |X_i| - 1$ but $x_i \in S_a$, then take any $X'_i \subset X_i$ of $|X_i| - 1$ points such that $S_a \cup \{x_i\} \subseteq X'_i$, the positive hull of $X'_1 \cup \dots \cup X'_k$ with the chosen X'_i and $x_j \in X'_j$ contains both a, b .

If $|S_a| \sqcup \{x_i\} = X_i$, by Proposition 4.5, we have $a = \sum_{e \in S_a} e$, hence $x_j \in \text{pos}\{a, b\}$ since

$$a + \frac{1}{\alpha}b = \sum_{e \in S_a} e + x_i + \frac{\beta}{\alpha}x_j = \frac{\beta}{\alpha}x_j$$

for $b = \alpha x_i + \beta x_j$.

Case 4: both a, b are not in the span of any X_i . Let $a \in \text{pos}\{x_i, x_j\}, b \in \text{pos}\{y_{i'}, y_{j'}\}$ for $x_i \in X_i, x_j \in X_j, y_{i'} \in X_{i'}, y_{j'} \in X_{j'}$.

By Proposition 4.2, we just need to consider the following situations.

First, suppose $i = i', j = j'$.

If $|X_i| = |X_j| = 2$, then we have the picture of the plane with four vectors $(0, 1), (0, -1), (1, 0), (-1, 0)$. Any two additional normals a, b not in the same quadrant would either have convex hull containing 0 or their positive hull contains one of the four basis vectors.

If both X_i and X_j have at least 3 elements, the positive hull of $X'_1 \cup \dots \cup X'_k$ for $x_i, y_i \in X'_i$ and $x_j, y_j \in X'_j$ contains both a, b .

If only one of them has at least 3 elements, say $|X_i| \geq 3$, then $x_i = y_i$ due to Proposition 4.3. We have the picture of the upper half-plane with 3 vectors $(0, 1), (-1, 0), (1, 0)$. Any two additional normals not in the same quadrant would have positive hull containing $(0, 1)$.

Second, suppose $\{i, j\}$ and $\{i', j'\}$ are disjoint.

The positive hull of $X'_1 \cup \dots \cup X'_k$ for $x_i \in X'_i, x_j \in X'_j, y_{i'} \in X'_{i'}, y_{j'} \in X'_{j'}$ contains both a, b .

We have shown contradictions in all the cases, hence no two primitive normals could be in two different positive hulls of the $|X_1| \dots |X_k|$ positive hulls. Therefore, every n primitive normals are contained in the same positive hull, which completes the theorem.

7. PROOFS OF PROPOSITIONS

At first, we present the approach that will be used throughout the section. In each proof, we will show a set of $n + 1$ points p_0, \dots, p_n so that some n points of them are linearly independent but all $n + 1$ points are linearly dependent by a unique (up to scaling) relation $\sum_i \lambda_i p_i = 0$ with 2 positive coefficients and 2 negative coefficients. This set of points clearly is in a bad position. Usually, some n points of them are shown to be linearly independent by Lemma 6.

For convenience, λ_i and θ_i will represent the points p and q in the way that: $p = \sum_i \lambda_i e_i$ and $q = \sum_i \theta_i e_i$. These representations will be used throughout the section without reintroducing λ_i and θ_i .

At first, we will present the following useful lemma to prove linear independency.

Lemma 6. *Given $E = \{e_0, \dots, e_n\}$ be the vertices of a simplex whose interior contains 0, then every p is not in the linear span of $E \setminus \{e_a, e_b\}$ for any $e_a \in S_p$ and any $e_b \in E \setminus S_p$, where S_p is the support⁶ of p .*

Proof. Let $p = \sum_{e_i \in S_p} \lambda_i e_i$.

Suppose p is in the positive hull of $E \setminus \{e_a, e_b\}$, we denote $p = \sum_{e_i \in E \setminus \{e_a, e_b\}} \alpha_i e_i$.

Taking the difference between the two representations,

$$\left(\sum_{e_i \in S_p \setminus \{e_a\}} (\lambda_i - \alpha_i) e_i \right) + \lambda_a e_a - \sum_{e_i \in (E \setminus S_p) \setminus \{e_b\}} \alpha_i e_i = 0.$$

⁶We use the same definition as Definition 2 but for $k = 1$.

It follows that 0 is a linear combination of $|E \setminus \{e_b\}| = n$ points in E with at least one nonzero coefficient λ_a . This is a contradiction as the n points are linearly independent. The conclusion follows. \square

Lemma 6 will be implicitly used to show that some n points are linearly independent in the way that: One of them is not in the span of the remaining points, which are themselves linearly independent.

Proof of Proposition 4.1. As x is not in the space spanned by any X_i , there is a partition of $1, \dots, k$ into two nonempty parts, say $L \sqcup M = \{1, \dots, k\}$ such that x is neither in the linear space \mathcal{L} spanned by $X_L = \bigcup_{i \in L} X_i$ nor the linear space \mathcal{M} spanned by $X_M = \bigcup_{i \in M} X_i$.

Let the projection of x onto the spaces \mathcal{L} and \mathcal{M} be $x_{\mathcal{L}}$ and $x_{\mathcal{M}}$.

We show that $x_{\mathcal{L}}$ is in X_L . Suppose otherwise, choose $\dim \mathcal{L}$ linearly independent points $\{y_i\}_i$ of X_L whose positive hull contains $x_{\mathcal{L}}$. If $x_{\mathcal{L}} = \sum_i \lambda_i y_i$, then at least two λ_i are positive. Choose $\dim \mathcal{M}$ linearly independent points $\{z_j\}_j$ of X_M but their positive hull does not contain $x_{\mathcal{M}}$. If $x_{\mathcal{M}} = \sum_j \theta_j z_j$, then at least one θ_j is negative.

In total, they are n linearly independent points, and x can be uniquely represented as

$$x = x_{\mathcal{L}} + x_{\mathcal{M}} = \sum_i \lambda_i y_i + \sum_j \theta_j z_j.$$

Since at least two λ_i are positive and at least one θ_j is negative, we obtain a bad position of x and the n points, a contradiction.

Therefore, $x_{\mathcal{L}}$ is just a positive scale of a point in X_L . Exchanging the role of \mathcal{L} and \mathcal{M} , we obtain the conclusion that x is in the positive hull of two points $x_i \in X_i, x_j \in X_j$ for different X_i, X_j . \square

Proof of Proposition 4.2. Suppose there exist $x \in \text{pos}\{x_1, x_2\}$ for $x_1 \in X_1, x_2 \in X_2$ and $y \in \text{pos}\{y_2, y_3\}$ for $y_2 \in X_2, y_3 \in X_3$.

Let $x = \alpha_1 x_1 + \alpha_2 x_2$ and $y = \beta_2 y_2 + \beta_3 y_3$.

Consider the case $x_2 \neq y_2$, the linear relation

$$\frac{1}{\alpha_2} x + \frac{1}{\beta_2} y = \frac{\alpha_1}{\alpha_2} x_1 + \frac{\beta_3}{\beta_2} y_3 - \sum_{p \in X_j \setminus \{x_2, y_2\}} p$$

shows that the $n + 1$ points $x, y, X'_1, X_2 \setminus \{x_2, y_2\}, X'_3, \dots, X'_k$ for any $X'_i \subset X_i$ satisfying $|X'_i| = |X_i| - 1, x_1 \in X'_1, y_3 \in X'_3$ are in a bad position.

Consider the case $x_2 = y_2$, the linear relation

$$\frac{1}{\alpha_2} x - \frac{1}{\beta_2} y = \frac{\alpha_1}{\alpha_2} x_1 - \frac{\beta_3}{\beta_2} y_3$$

shows that the $n + 1$ points $x, y, X'_1, X_2'', X'_3, \dots, X'_k$ for any $X'_i \subset X_i$ satisfying $|X'_i| = |X_i| - 1, x_1 \in X'_1, y_3 \in X'_3, X_2'' \subset X_2, |X_2''| = |X_2| - 2, x_2 \notin X_2''$ are in a bad position.

The conclusions follows from the bad positions in the cases covered. \square

Proof of Proposition 4.3. This proof is almost identical to the proof of Proposition 4.2.

Suppose $i = 1, j = 2$, let $x = \alpha_1 x_1 + \alpha_2 x_2, y = \beta_1 y_1 + \beta_2 y_2$.

Assume the conclusion does not hold, that is $x_1 \neq y_1$ while $|X_1| > 2$.

Consider the case $x_2 \neq y_2$, the linear relation

$$\frac{1}{\alpha_2} x + \frac{1}{\beta_2} y = \frac{\alpha_1}{\alpha_2} x_1 + \frac{\beta_1}{\beta_2} y_1 - \sum_{p \in X_j \setminus \{x_2, y_2\}} p$$

shows that the $n + 1$ points $x, y, X'_1, X_2 \setminus \{x_2, y_2\}, X'_3, \dots, X'_k$ for any $X'_i \subset X_i$ satisfying $|X'_i| = |X_i| - 1$, $x_1, y_1 \in X'_1$ are in a bad position.

Consider the case $x_2 = y_2$, the linear relation

$$\frac{1}{\alpha_2}x - \frac{1}{\beta_2}y = \frac{\alpha_1}{\alpha_2}x_1 - \frac{\beta_1}{\beta_2}y_1$$

shows that the $n + 1$ points $x, y, X'_1, X''_2, X'_3, \dots, X'_k$ for any $X'_i \subset X_i$ satisfying $|X'_i| = |X_i| - 1$, $x_1, y_1 \in X'_1$, $X''_2 \subset X_2$, $|X''_2| = |X_2| - 2$, $x_2 \notin X''_2$ are in a bad position.

The bad positions imply that $x_1 = y_1$. \square

Proof of Proposition 4.4. Consider a point $y \notin X_i$ in the space spanned by X_i , then there exists a subset $X'_i \subset X_i$ of $|X_i| - 1$ points whose positive hull contains y . Let $y = \sum_{e_t \in X'_i} \lambda_t e_t$ such that all λ_t are nonnegative with at least two of them positive.

Assume $x_i \in S_y$, we have

$$x_i = \frac{1}{\lambda_i}y - \sum_{e_t \in S_y \setminus \{x_i\}} \frac{\lambda_t}{\lambda_i}e_t.$$

Choose the k subsets $X'_t \subset X_t$ for $t = 1, \dots, k$ so that $|X'_t| = |X_t| - 1$, $S_y \subseteq X'_i$ and $x_j \in X'_j$. The points in $X'_i \cup X'_j \cup \{y\} \setminus \{x_i\}$ are linearly independent, thus, the following unique relation

$$x = \alpha \left(\frac{1}{\lambda_i}y - \sum_{e_t \in S_y \setminus \{x_i\}} \frac{\lambda_t}{\lambda_i}e_t \right) + \beta x_j,$$

for $x = \alpha x_i + \beta x_j$, indicates a bad position for the $n + 1$ points $\{x, y\} \cup (X'_1 \cup \dots \cup X'_k) \setminus \{x_i\}$, contradiction.

Therefore, $x_i \notin S_y$. \square

Proof of Proposition 4.5. Let $x = \alpha x_i + \beta x_j$.

Assume there exists a point y in the span of X_i but not a positive scale of $\sum_{e_t \in S_y} e_t$. Let the support of y be $S_y = \{e_0, \dots, e_m\} \subset X_i$, by Proposition 5.1, with the strange index supposed to be e_0 , we have $y = \theta_0 e_0 + \sum_{t=1}^m e_t$ for $\theta_0 > 1$.

By Proposition 4.4, $x_i \notin S_y$. The linear relation

$$\begin{aligned} y &= (\theta_0 - 1)e_0 + \sum_{t=0}^m e_t \\ &= (\theta_0 - 1)e_0 - \sum_{e_t \in X_i \setminus S_y} e_t \\ &= (\theta_0 - 1)e_0 - \left(\sum_{e_t \in (X_i \setminus S_y) \setminus \{x_i\}} e_t \right) - \frac{1}{\alpha}x + \frac{\beta}{\alpha}x_j \end{aligned}$$

shows a bad position for the points $\{x, y\} \cup X'_1 \cup \dots \cup X'_{i-1} \cup X_i \setminus \{x_i, x_{i'}\} \cup X'_{i+1} \cup \dots \cup X'_k$, where $X'_t \subset X_t$ is any set with $|X'_t| = |X_t| - 1$ and $x_{i'} \in S_y \setminus \{e_0\}$, $x_j \in X'_j$.

The conclusion on the relation between S_p and S_q can be deduced from Proposition 5.2 with the restriction from Proposition 4.4. We however give a self-contained argument below.

Assume there exist p, q in the space spanned by X_i such that $S_p \cap S_q \neq \emptyset$, $S_p \setminus S_q \neq \emptyset$, $S_q \setminus S_p \neq \emptyset$. By Proposition 4.4, neither S_p or S_q contains x_i , hence $q, S_p \setminus S_q, S_q \setminus S_p$ are

linearly independent. Therefore, let $p = \sum_{e \in S_p} e$, $q = \sum_{e \in S_q} e$, the linear relation

$$p - q = \sum_{e \in S_p \setminus S_q} e - \sum_{e \in S_q \setminus S_p} e$$

shows a bad position for the $n+1$ points $\{p, q\} \cup X'_1 \cdots \cup X'_{i-1} \cup X_i \setminus \{x_i, x_{i'}\} \cup X'_{i+1} \cup \cdots \cup X'_k$ for any $x_{i'} \in S_p \cap S_q$ and any $X'_t \subset X_t$ with $|X'_t| = |X_t| - 1$. \square

Proof of Proposition 5.1. Assume otherwise, there are three indices of S_p , namely 0, 1, 2, such that $1 = \lambda_0$, $1 < \lambda_1$, $1 < \lambda_2$ for $p = \sum_{i=0}^k \lambda_i e_i$ (that is $S_p = \{e_0, \dots, e_k\}$).

Since $e_0 = -\sum_{i=1}^n e_i$, the unique relation

$$p = \sum_{i=0}^k \lambda_i e_i = \sum_{i=1}^k \lambda_i e_i - \sum_{i=1}^n e_i = (\lambda_1 - 1)e_1 + (\lambda_2 - 1)e_2 + \sum_{i=3}^k (\lambda_i - 1)e_i - \sum_{i=k+1}^n e_i$$

shows that the $n+1$ points p, e_1, \dots, e_n are in a bad position, contradiction. \square

Proof of Proposition 5.2. Suppose there are p, q with $S_p \cap S_q \neq \emptyset$, $S_p \setminus S_q \neq \emptyset$, $S_q \setminus S_p \neq \emptyset$.

We first show that $S_p \cup S_q = E$. Assume otherwise, consider any $e_j \in S_p \cap S_q$,

$$\theta_j p - \lambda_j q = \sum_{e_i \in S_p \setminus S_q} \theta_j \lambda_i e_i - \sum_{e_i \in S_q \setminus S_p} \lambda_j \theta_i e_i + \sum_{e_i \in (S_p \cap S_q) \setminus \{e_j\}} (\theta_j \lambda_i - \lambda_j \theta_i) e_i.$$

Since $S_p \cup S_q \neq E$, p is not in the linear span of $S_p \cup S_q \setminus \{e_j\}$, by Lemma 6. The above linear relation shows that the $n+1$ points $\{p, q\} \cup S_p \cup S_q \setminus \{e_j\}$ are in a bad position, contradiction.

Assume $|S_p \cap S_q| \geq 2$, we show that $|S_p \cap S_q| = n - 1$.

Take any a, b such that $\lambda_a = \max_i \lambda_i$ and $\theta_b = \max_i \theta_i$.

If $e_a, e_b \in S_p \cap S_q$, then the linear relation

$$\begin{aligned} p + q &= p + q - \sum_{e_i \in E} e_i = \sum_{e_i \in S_p \cup S_q} (\lambda_i + \theta_i - 1) e_i - \sum_{e_i \in E \setminus (S_p \cup S_q)} e_i \\ &= \sum_{e_i \in S_p \cap S_q} (\lambda_i + \theta_i - 1) e_i - \sum_{e_i \in E \setminus (S_p \cup S_q)} e_i \end{aligned}$$

shows that the $n+1$ points $\{p, q\} \cup E \setminus \{e_i, e_j\}$ for any $e_i \in S_p \setminus S_q$ and any $e_j \in S_q \setminus S_p$ are in a bad position, contradiction.

If $e_a \in S_p \cap S_q, e_b \in S_q \setminus S_p$, then the relation

$$\begin{aligned} \theta_b p + q &= \theta_b p + q - \theta_b \sum_{e_i \in E} e_i \\ &= \sum_{e_i \in S_p \setminus S_q} \theta_b e_i + \sum_{e_i \in S_p \cap S_q} (\theta_b \lambda_i + 1) e_i + \left(\sum_{e_i \in (S_q \setminus S_p) \setminus \{e_b\}} e_i \right) + \theta_b e_b \\ &= \sum_{e_i \in S_p \cap S_q} (\theta_b \lambda_i + 1) e_i + \sum_{e_i \in (S_q \setminus S_p) \setminus \{e_b\}} e_i - \theta_b \left(\sum_{e_i \notin (S_p \setminus S_q) \cup \{e_b\}} e_i \right) \\ &= \sum_{e_i \in S_p \cap S_q} (\theta_b \lambda_i + 1 - \theta_b) e_i + \sum_{e_i \in (S_q \setminus S_p) \setminus \{e_b\}} (1 - \theta_b) e_i \end{aligned}$$

shows that the $n+1$ points $\{p, q\} \cup E \setminus \{e_i, e_b\}$ for some $e_i \in S_p \setminus S_q$ are in a bad position, contradiction.

If $e_a \in S_p \setminus S_q, e_b \in S_p \cap S_q$, we have a similar argument due to symmetry.

If $e_a \in S_p \setminus S_q, e_b \in S_q \setminus S_p$, and $|S_p \setminus S_q| \geq 2$, then the relation

$$\begin{aligned} \theta_b p + q &= \theta_b \lambda_a e_a + \sum_{e_i \in (S_p \setminus S_q) \setminus \{e_a\}} \theta_b e_i + \sum_{e_i \in S_p \cap S_q} (\theta_b + 1) e_i \\ &\quad + \left(\sum_{e_i \in (S_q \setminus S_p) \setminus \{e_b\}} e_i \right) + \theta_b e_b - \theta_b \sum_{e_i \in E} e_i \\ &= (\theta_b \lambda_a - \theta_b) e_a + \sum_{e_i \in S_p \cap S_q} e_i + \sum_{e_i \in (S_q \setminus S_p) \setminus \{e_b\}} (1 - \theta_b) e_i \end{aligned}$$

shows that the $n + 1$ points $\{p, q\} \cup E \setminus \{e_i, e_b\}$ for some $e_i \in (S_p \setminus S_q) \setminus \{e_a\}$ are in a bad position, contradiction.

A similar argument applies for the case $e_a \in S_p \setminus S_q, e_b \in S_q \setminus S_p$, and $|S_q \setminus S_p| \geq 2$.

The only remaining case is $|S_p \setminus S_q| = |S_q \setminus S_p| = 1$, that is $|S_p \cap S_q| = n - 1$. The conclusion follows.

(Note that in Case (iv), if p, q have strange indices, they can be the points in $S_p \setminus S_q$ and $S_q \setminus S_p$ only. This fact will be used in the proof of Proposition 5.7.) \square

Proof of Proposition 5.3. Assume distinct e_a, e_b are the strange indices of p, q respectively. The linear relation

$$p - q = (\lambda_a - 1)e_a - (\theta_b - 1)e_b$$

shows that the points $p, q, E \setminus \{e_i, e_j\}$ for any $e_i \in S_p \setminus \{e_a, e_b\}, e_j \in E \setminus S_p$ are in a bad position, contradiction. \square

Proof of Proposition 5.4. Assume e_a, e_b are the strange indices of p, q respectively, then the linear relation

$$p + q - (\lambda_a - 1)e_a - (\theta_b - 1)e_b + \sum_{e_i \in E \setminus (S_p \cup S_q)} e_i = 0$$

shows that the $n + 1$ points $p, q, E \setminus \{e_i, e_j\}$ for any $e_i \in S_p \setminus \{e_a\}, e_j \in S_q \setminus \{e_b\}$ are in a bad position, contradiction. \square

Proof of Proposition 5.5. Suppose $S_p = \{e_0, \dots, e_k\}, S_q = \{e_0, \dots, e_\ell\}$ with $0 < k < \ell < n$.

Since there is always some $\lambda_i = 1$, we assume that $\lambda_0 = 1$. Consider the relation

$$\lambda_0 q - \theta_0 p = \sum_{i=1}^k (\lambda_0 \theta_i - \theta_0 \lambda_i) e_i + \sum_{i=k+1}^{\ell} \lambda_0 \theta_i e_i.$$

We need $\lambda_0 \theta_i - \theta_0 \lambda_i \geq 0$ for each $i = 1, \dots, k$, otherwise the $n + 1$ points $\{p, q\} \cup E \setminus \{e_0, e_n\}$ would be in a bad position.

If $\theta_i = 1$ for some $1 \leq i \leq k$, then $\lambda_0 \theta_i - \theta_0 \lambda_i \geq 0$ implies $\theta_0 = 1$ and $\lambda_i = 1$.

If $\theta_i > 1$ for some $1 \leq i \leq k$, then $\lambda_0 \theta_i - \theta_0 \lambda_i \geq 0$ implies $\theta_i \geq \lambda_i$. (Note that we still have $\theta_0 = 1$ in this case as $\theta_i > 1$.)

Suppose $\theta_i > \lambda_i$ for some i , say $\theta_k > \lambda_k$, we have

$$\lambda_k q - \theta_k p = \sum_{i=0}^{k-1} (\lambda_k \theta_i - \theta_k \lambda_i) e_i + \sum_{i=k+1}^{\ell} \lambda_k \theta_i e_i.$$

Since $\theta_0 = \lambda_0 = 1$, it follows that $\lambda_k \theta_0 - \theta_k \lambda_0 < 0$. It means the $n + 1$ points $\{p, q\} \cup E \setminus \{e_k, e_n\}$ are in a bad position, contradiction.

Therefore, we obtain the conclusion that $\lambda_i = \theta_i$ for $i = 0, \dots, k$. \square

Proof of Proposition 5.6. Let $e_a \in S_p$, $e_b \in S_q$ so that $\lambda_a = \max_{e_i \in S_p} \lambda_i$ and $\theta_b = \max_{e_i \in S_q} \theta_i$. If $\lambda_a = \lambda_k$, we set $a = k$. Likewise, if $\theta_b = \theta_k$, we set $b = k$.

Consider the case $|S_p| \geq 3$, $|S_q| \geq 3$.

If $a \neq k$ and $b \neq k$, then the relation

$$p + q - (\lambda_a - 1)e_a - (\theta_b - 1)e_b - (\lambda_k + \theta_k - 1)e_k = 0$$

shows that the $n+1$ points $\{p, q\} \cup E \setminus \{e_i, e_j\}$ for any $e_i \in S_p \setminus \{e_a, e_k\}$ and $e_j \in S_q \setminus \{e_b, e_k\}$ are in a bad position, contradiction.

If $a = k$ and $b \neq k$, then the relation

$$p + q - (\lambda_k + \theta_k - 1)e_k - (\theta_b - 1)e_b = 0$$

shows that the $n+1$ points $\{p, q\} \cup E \setminus \{e_i, e_j\}$ for any $e_i \in S_p \setminus \{e_k\}$ and $e_j \in S_q \setminus \{e_b, e_k\}$ are in a bad position, contradiction. (Note that this also works for $|S_p| \geq 2$.)

A similar argument works for the case $a \neq k$ and $b = k$.

For the case $|S_p| = 2$ (hence $|S_q| \geq 3$), we suppose $S_p = \{e_0, e_1\}$, $S_q = \{e_1, \dots, e_n\}$ (then $k = 1$).

If $a = 1$, then the above linear relation shows that $b = 1$, otherwise we have same set of points in a bad position, contradiction.

If $a = 0$, suppose $b \neq 1$ then the linear relation

$$\frac{1}{\lambda_0}p + q - \frac{\lambda_1}{\lambda_0}e_1 - (\theta_b - 1)e_b = 0$$

shows that the points $p, q, E \setminus \{e_0, e_b\}$ are in a bad position, contradiction.

All the contradictions confirm the conclusion. \square

Proof of Proposition 5.7. As in the proof of Proposition 5.2, we have shown that for the condition given in Proposition 5.7, the strange indices of p, q , if any, are e_a, e_b , respectively. It means $\lambda_i = \theta_i = 1$ for $i \in S_p \cap S_q$.

We now prove that $\lambda_a > 1$, $\theta_b > 1$ and $\lambda_a \theta_b \geq \lambda_a + \theta_b$. Assume otherwise, the linear relation

$$\begin{aligned} \theta_b p + \lambda_a q &= \theta_b \lambda_a e_a + \lambda_a \theta_b e_b + \sum_{e_i \in E \setminus \{e_a, e_b\}} (\lambda_a + \theta_b) e_i - \lambda_a \theta_b \sum_{e_i \in E} e_i \\ &= \sum_{e_i \in E \setminus \{e_a, e_b\}} (\lambda_a + \theta_b - \lambda_a \theta_b) e_i \end{aligned}$$

has the positive coefficient $\lambda_a + \theta_b - \lambda_a \theta_b$ of e_i for $e_i \in E \setminus \{e_a, e_b\}$ (note that $|E \setminus \{e_a, e_b\}| \geq 2$), a contradiction by the point set $\{p, q\} \cup E \setminus \{e_a, e_b\}$. \square

Proof of Proposition 5.8. Assume otherwise, there exist $e_i \in S_r \setminus S_q$ and $e_j \in S_r \setminus S_p$. In fact, $e_i \in S_p \setminus S_q$ and $e_j \in S_q \setminus S_p$.

Either S_p or S_q is not a subset of S_r since otherwise $S_r = E$. Suppose S_q is not a subset of S_r . It follows that $|S_r \cap S_q|$ is either 1 or $n-1$ (by Proposition 5.2).

Suppose $|S_r \cap S_q| = 1$, that is $S_r \cap S_q = \{e_j\}$. It follows that $S_r = E \setminus S_1 \cup \{e_j\}$ by Proposition 5.2. However, we then have $|S_p \cap S_r| = |S_p| - 1$, which is neither 1 nor $n-1$, contradicting Proposition 5.2.

It follows that $|S_r \cap S_q| = n-1$. However, obviously $|S_r \cap S_q| \leq |S_q| < n-1$ since $|S_p| \geq 3$, contradiction. \square

Proof of Proposition 5.9. Suppose $S_p = \{e_0, \dots, e_{n-1}\}$, $S_q = \{e_1, \dots, e_n\}$. Assume S_r is not a subset of $S_p \cap S_q$, we show that $S_r = \{e_0, e_n\}$.

Firstly, $|S_r| < n$. Assume otherwise, let $S_r = E \setminus \{e_k\}$ ($k \neq 0, n$). Because $S_r \cup S_p = E$, $|S_r \cap S_p| = n - 1$ and $S_p \setminus S_r = \{e_k\}$, one obtains $\lambda_k > 1$ but we already have $\lambda_0 > 1$ (by Proposition 5.7), contradiction.

We continue by showing that $|S_r|$ cannot exceed 2. Assume otherwise $|S_r| > 2$. As S_r is not a subset of $S_p \cap S_q$, either e_0 or e_n is in S_r , say $e_0 \in S_r$. Hence, $|S_r \cap S_q| = |S_r| - 1$. It follows that $1 < |S_r \cap S_q| < n - 1$, contradiction.

As $|S_r| = 2$, if S_r is not a subset of $S_p \cap S_q$, then either $S_r = \{e_0, e_k\}$, or $S_r = \{e_k, e_n\}$, or $S_r = \{e_0, e_n\}$ for some $k \neq 0, n$. Assume $S_r = \{e_0, e_k\}$, then $S_r \cap S_p = \{e_k\}$, by Proposition 5.6 we obtain $\lambda_k > 1$ as the strange index of p , but $\lambda_0 > 1$, a contradiction. A similar contradiction is raised by the case $S_r = \{e_k, e_n\}$. Therefore, $S_r = \{e_0, e_n\}$. \square

Proof of Proposition 5.10. Suppose $S_p = \{e_0, e_1\}$, $S_q = \{e_1, \dots, e_n\}$. Assume S_r is not a subset of $S_q \setminus S_p$, that is S_r contains either e_0 or e_1 .

If $|S_r| = 2$ then S_r is either $\{e_0, e_k\}$ or $\{e_1, e_k\}$ for $k \neq 0, 1$. The intersection $S_r \cap S_p$ is only a point but their union is not E , contradicting Proposition 5.2.

Assume $2 < |S_r| < n$, one obtains $e_0 \notin S_r$, as otherwise $1 < |S_r \cap S_q| < n - 1$. Also, $e_1 \notin S_r$, as otherwise $S_p \cup S_r$ is not E , contradiction. It follows that $|S_r| = n$.

Suppose $S_r = E \setminus \{e_k\}$ for $k \neq 1$. As $|S_r \cap S_q| = n - 1$, it follows that $\theta_k > 1$ by Proposition 5.7. However, we also have $\theta_1 > 1$ by Proposition 5.6, contradiction. Therefore, $S_r = E \setminus \{e_1\} = (S_p \setminus S_q) \cup (S_q \setminus S_p)$. \square

ACKNOWLEDGEMENT

The author would like to thank Roman Karasev for his encouragement in characterizing this class of polytopes and his patient reading and commenting various pieces of this paper.

REFERENCES

- [1] P McMullen, R Schneider, and GC Shephard. Monotypic polytopes and their intersection properties. *Geometriae Dedicata*, 3(1):99–129, 1974.
- [2] RN Karasëv. On the characterization of generating sets. *Modelir. Anal. Inf. Sist*, 8(2):3–9, 2001.
- [3] D Borowska and J Grzybowski. Strongly monotypic polytopes in R^3 . *Demonstratio Mathematica*, 41(1):165–170, 2008.
- [4] Friedrich Wilhelm Levi. Überdeckung eines eibereiches durch parallelverschiebung seines offenen kerns. *Archiv der Mathematik*, 6(5):369–370, 1955.
- [5] Vladimir Boltyanski, Horst Martini, and Petru S Soltan. *Excursions into combinatorial geometry*. Springer Science & Business Media, 2012.

VUONG BUI, INSTITUT FÜR INFORMATIK, FREIE UNIVERSITÄT BERLIN, TAKUSTRASSE 9, 14195 BERLIN, GERMANY

Email address: bui.vuong@fu-berlin.de