

GEOMETRY OF LINES ON A CUBIC FOURFOLD

FRANK GOUNELAS AND ALEXIS KOUVIDAKIS

ABSTRACT. For a general cubic fourfold $X \subset \mathbb{P}^5$ with Fano scheme of lines F , we prove a number of properties of the universal family of lines $I \rightarrow F$ and various subloci. We first describe the moduli and ramification theory of the genus four fibration $p : I \rightarrow X$ and explore its relation to a birational model of F in I . The main part of the paper is devoted to describing the locus $V \subset F$ of triple lines, i.e., the fixed locus of the Voisin map $\phi : F \dashrightarrow F$, in particular proving it is an irreducible projective singular surface of class $21c_2(\mathcal{U}_F)$ and detailing its intersection with the locus S of second type lines. A consequence of the analysis of the singularities of V is a geometric proof of the fact that if X is very general, then the number of singular (necessarily 1-nodal) rational curves in F of primitive class is 3780.

CONTENTS

1. Introduction	1
2. Background and notation	3
3. The curve of lines through a point	8
4. Universal families and the surface V	15
5. Geometry of the ramification locus	31
Appendix A. Admissible covers	35
References	37

1. INTRODUCTION

Let $X \subset \mathbb{P}_{\mathbb{C}}^5$ be a smooth cubic fourfold and $F \subset G(2, 6)$ its Fano scheme of lines. The geometry of these varieties has received a lot of attention since the beginning of the 20th century and in particular since the foundational papers [CG72, AK77]. Consider the universal family $I = \mathbb{P}(\mathcal{U}_F)$ of lines, for \mathcal{U}_F the universal subbundle of the Grassmannian $G(2, 6)$ restricted to F , with the two projections

$$\begin{array}{ccc} I & \xrightarrow{p} & X \\ \downarrow q & & \\ F & & \end{array}$$

It turns out that the morphism p is a fibration of $(2, 3)$ -complete intersections in \mathbb{P}^3 associated to the geometry of X , and all but finitely many fibres are genus 4 curves. As such, each fibre $C_x = p^{-1}(x) \subset I$, which parametrises

Date: September 20, 2021.

2020 Mathematics Subject Classification. 14J70, 14J35, 14H10, 14M15.

lines in X through x , comes equipped with two g_3^1 linear systems by restricting the rulings of the corresponding quadric. The aim of this paper is to study the fibrations p, q and the geometry and intersection theory of various geometric loci in each variety.

Recall that there are two types of points $[\ell] \in F$, i.e., lines $\ell \subset X$, depending on the decomposition of the normal bundle $N_{\ell/X}$. The generic line is called of *first type*, whereas there is a surface $S \subset F$ parametrising those of *second type* (see Section 2 for formal definitions).

Our first goal is to further expand in Section 3 on the geometry of the curves C_x , which we achieve by studying properties of the fibration p on I but also the induced restricted map over $q^{-1}(S) \subset I$. To name one example, we obtain the following as a combination of Proposition 3.5, Corollary 3.7.

Theorem A. *For $x \in X \subset \mathbb{P}^5$ a general point of a general cubic fourfold, C_x is Brill–Noether general and the morphism*

$$p|_{q^{-1}(S)} : q^{-1}(S) \rightarrow W \subset X$$

is birational onto its image W , an irreducible divisor in X . If $x \in W$ is general, then C_x is an irreducible curve with one node whose normalisation is Brill–Noether general.

Moving now to geometric constructions on F , Voisin [Voi04] defined a map

$$\phi : F \dashrightarrow F$$

taking a point $[\ell]$ corresponding to a line $\ell \subset X$ to the *residual line* $[\ell']$, in the sense that, if ℓ is general, there is a unique $\Pi_\ell = \mathbb{P}^2$ so that $X \cap \Pi_\ell = 2\ell + \ell'$. This map has been studied in detail in [Ame11] and if X does not contain any planes (the generic case, which we are in), is resolved by a single blowup of the locus S . This gives impetus to the construction of the following map

$$\begin{aligned} \psi : F &\dashrightarrow X \\ [\ell] &\mapsto \ell \cap \ell'. \end{aligned}$$

The locus of indeterminacy of ψ also includes the locus $V \subset F$ of *triple lines*, i.e., lines for which there is a $\Pi_\ell = \mathbb{P}^2$ so that $X \cap \Pi_\ell = 3\ell$. To the best of our knowledge, this locus has so far eluded study. Note that it can be viewed as the closure of the fixed locus

$$\text{Fix}(\phi) \subset F$$

of the Voisin map ϕ . We will prove that it consists of both first and second type lines. Using results of Section 3, we may interpret a general point $[\ell] \in V$ as a triple ramification point of one of the g_3^1 's on C_x for $x \in \ell$.

The main part of this paper is devoted, in Section 4, to the study of various universal families related to V, S and F . We conclude the following, where \mathcal{U}_F is the restriction of the universal bundle from the Grassmannian and $H_F = c_1(\mathcal{U}_F^\vee)$ the restriction of the Plücker polarisation.

Theorem B. *If $X \subset \mathbb{P}^5$ is a general cubic, then the locus $V \subset F$ of triple lines is a projective irreducible singular surface, whereas the locus $C = S \cap V$ is pure 1-dimensional and nodal. Their classes are*

$$[V] = 21c_2(\mathcal{U}_F) \in \text{CH}^2(F),$$

$$[C] = \frac{35}{2}H_F^3 \in H_2(F, \mathbb{Q}).$$

In particular, the class of V is a multiple of the generator (cf. [Voi10]) of the known extremal ray of the effective cone $\text{Pseff}^2(F)$. As a corollary to the analysis of the singularities of V , we obtain the following, which to the best of our knowledge was not previously known (despite similar results, e.g., [NO21, Theorem 1.3]).

Theorem C. *If $X \subset \mathbb{P}_{\mathbb{C}}^5$ is a very general cubic, then the number of*

- (1) *nodal rational curves in F of primitive class $\beta \in H_2(F, \mathbf{Z})$,*
- (2) *singularities of V , all of which are isolated non-normal with degree two normalisation,*
- (3) *nodes of $C = S \cap V$,*

are all equal to 3780.

As a further application of the analysis above, in Section 5 we study the loci

$$R, R', N \subset I$$

which are the closure of the locus of ramification, residual to all ramification and triple ramification points respectively, of the two g_3^1 's on the smooth fibres C_x . The projection $R' \rightarrow F$ is generically finite of degree 16, the degree of the Voisin map ϕ . On the other hand, we compute that the map ψ from above has degree 24, and in Corollary 5.5 we prove that it is resolved by blowing up S and then the strict transform of V .

Finally, putting everything together, we prove the following in Theorem 5.3 and compute in Proposition 5.7 the classes of the above loci in the Chow group of I in terms of the tautological line bundle $l = \mathcal{O}_I(1) = p^*\mathcal{O}_X(1)$ and the pullback q^*H_F of the class of the Plücker polarisation (or the relative canonical divisor ω_p of the fibration $p : I \rightarrow X$).

Theorem D. *If $X \subset \mathbb{P}^5$ is a general cubic fourfold, then the locus of ramification points R is irreducible and smooth outside $q^{-1}(S) \subset R$ and has normalisation the blowup of F first at S and then the strict transform of V . The classes of R, R' in $\text{CH}^1(I)$ and N in $\text{CH}^2(I)$ are given as follows*

$$\begin{aligned} [R] &= 4q^*H_F + l = 4c_1(\omega_p) - 3l, \\ [R'] &= 4q^*H_F + 16l = 4c_1(\omega_p) + 12l, \\ [N] &= 21(l^2 - lq^*H_F) = 21(2l^2 - lc_1(\omega_p)). \end{aligned}$$

In the appendix, we reprove the first two of the above class computations using the genus four fibration $I \rightarrow X$ and some intersection theory on the moduli space of admissible covers.

Acknowledgements We would like to thank Olof Bergvall and Daniel Huybrechts for helpful correspondence. The first author was supported by the ERC Consolidator Grant 681838 “K3CRYSTAL”.

2. BACKGROUND AND NOTATION

As the notation surrounding cubic fourfolds is substantial, we devote this section to fixing that used in the paper and recalling some basic properties, so that it acts as a reference for later sections.

For a vector bundle E we denote by $\mathbb{P}(E) = \text{Proj}(\text{Sym}(E^\vee))$, so that projective space parametrises one dimensional subspaces. We denote by $G(k, n)$ the space of k -dimensional subspaces of \mathbb{C}^n , with universal bundle \mathcal{U} of rank k and universal quotient bundle \mathcal{Q} of rank $n - k$. When multiple such bundles are in question, we will denote them by \mathcal{U}_k . Finally, we denote by σ_I the standard Schubert cycles for an index I so that, e.g., $\sigma_i = c_i(\mathcal{Q})$ for $i \geq 1$.

Throughout, $X \subset \mathbb{P}^5$ will be a smooth cubic fourfold with $H_X = \mathcal{O}_X(1)$ and $F \subset G(2, 6)$ the Fano scheme of lines contained in X which is a hyperkähler fourfold [BD85]. To unburden notation, we will often be sloppy in distinguishing a line $\ell \subset X$ and the point $[\ell] \in F$ that it defines. We denote by $\mathcal{U}_F, \mathcal{Q}_F$ the restrictions of \mathcal{U}, \mathcal{Q} to F . The universal family of lines sits in a diagram

$$\begin{array}{ccc} \mathbb{P}(\mathcal{U}) & \xrightarrow{p} & \mathbb{P}^5 \\ \downarrow q & & \\ G(2, 6) & & \end{array}$$

and note that we have

$$p^* \mathcal{O}_{\mathbb{P}^5}(1) \cong \mathcal{O}_{\mathbb{P}(\mathcal{U})}(1).$$

We use the same notation p, q for the induced maps on

$$I := \mathbb{P}(\mathcal{U}_F)$$

the universal family of lines on F .

The subvariety $F \subset G(2, 6)$ is given by a section of the rank four bundle $\text{Sym}^3 \mathcal{U}^* \cong q_* p^* \mathcal{O}_{\mathbb{P}^5}(3)$ - in fact it is the section induced, under this isomorphism, by $f \in k[x_0, \dots, x_5]_3$ whose vanishing is X (see [EH16, Proposition 6.4]) so its cohomology class in the Grassmannian is given by $c_4(\text{Sym}^3 \mathcal{U}^*)$ which can be computed as follows (see [Ful98, Example 14.7.13])

$$\begin{aligned} [F] &= 18c_1(\mathcal{U}^*)^2 c_2(\mathcal{U}^*) + 9c_2(\mathcal{U}^*)^2 \\ (1) \quad &= 18\sigma_1^2 \sigma_{1,1} + 9\sigma_{1,1}^2 \\ &= 27\sigma_2^2 - 9\sigma_1 \sigma_3 - 18\sigma_4. \end{aligned}$$

Consider now the morphism $p : I \rightarrow X$. We denote by

$$C_x := p^{-1}(x)$$

the fibre over x , which parametrises the lines in X containing x . If X is general, then C_x is 1-dimensional for all x (see [LSV17, Lemma 2.5]), whereas for arbitrary X there are only finitely many points in X where the fibre can be two dimensional (see [CS09, Corollary 2.2]). We defer to Proposition 3.5 for more properties of C_x . In any case, C_x embeds in F via q and can also be realised as the $(2, 3)$ -complete intersection in \mathbb{P}^3 formed by the intersection points of the lines through x with $T_x X \cap A$, where A a hyperplane not containing x : for $R_x X = \mathbf{V}(\sum x_i \partial_i f)$ the polar quadric, C_x is the intersection $R_x X \cap T_x X \cap X \cap A$. As such, if it is 1-dimensional, it is of arithmetic genus 4 and has two g_3^1 's, counted with multiplicities [EH16, Corollary D.11], namely the restrictions of the rulings of the quadric $R_x X \cap T_x X \cap A$.

If X is very general, $\text{Pic}(F) = \mathbf{Z}$ and we denote by β the generator of $H_2(F, \mathbf{Z})^{\text{alg}}$. In [Ame11] it is proven that

$$[C_x] = 2\beta \in H_2(F, \mathbf{Z}).$$

Following [CG72], there are two types of lines $\ell \in F$, depending on the decomposition of the normal bundle $N_{\ell/X}$.

Definition 2.1. We say that a line $\ell \subset X$ is

- (1) of *first type* if $N_{\ell/X} \cong \mathcal{O}(1) \oplus \mathcal{O}^2$,
- (2) of *second type* if $N_{\ell/X} \cong \mathcal{O}(1)^2 \oplus \mathcal{O}(-1)$.

An equivalent geometric description is as follows: ℓ is of

- (1) first type if there is a unique $\Pi_\ell = \mathbb{P}^2$ tangent to X along ℓ .
- (2) second type if there is a unique $H_\ell = \mathbb{P}^3$ tangent to X along ℓ (equivalently, a family $\Pi_{\ell,t} = \mathbb{P}^2$, $t \in \mathbb{P}^1$, of 2-planes tangent to X along ℓ).

Denote by

$$\begin{aligned} S &:= \{\ell : \ell \text{ is of second type}\} \subset F, \\ W &:= p(\mathbb{P}(\mathcal{U}_S)) = p(q^{-1}(S)) \subset X, \end{aligned}$$

the *locus of second type lines* and the locus of points through which there passes a second type line respectively. The universal family of second type lines $q^{-1}(S) = \mathbb{P}(\mathcal{U}_S)$ is the singular locus $\text{Sing}(p)$ of the morphism $p : I \rightarrow X$ from Lemma 3.1.

Lemma 2.2. ([Huy20, Corollary 2.2.14]) *Let $X \subset \mathbb{P}^5$ be a smooth cubic fourfold. Then W is 3-dimensional, and if X is general it is irreducible.*

Denote by $H_F = c_1(\mathcal{U}_F^\vee)$ the Plücker ample line bundle on F and by H_S the restriction on S . The following is a combination of [Ame11, Lemma 1], [OSY19, §3], [Huy20, Proposition 5.4.7].

Theorem 2.3. *If $X \subset \mathbb{P}^5$ is a cubic fourfold then S is 2-dimensional and is the degeneracy locus of the Gauss map, i.e., the following morphism of vector bundles*

$$\text{Sym}^2 \mathcal{U}_F \rightarrow \mathcal{Q}_F^\vee.$$

In particular $c_1(K_S) = 3H_S$ in $H^2(S, \mathbb{Q})$ and the class of S in $\text{CH}^2(F)$ is given by

$$[S] = 5(c_1(\mathcal{U}_F^\vee)^2 - c_2(\mathcal{U}_F^\vee)) = 5c_2(\mathcal{Q}_F) = 5\sigma_2|_F.$$

If X is general, S is a smooth projective irreducible surface.

Let

$$(2) \quad \begin{aligned} \phi : F &\dashrightarrow F \\ \ell &\mapsto \ell' \end{aligned}$$

be the *Voisin map* of [Voi04], taking a general line ℓ and giving the residual line ℓ' in the tangent 2-plane Π_ℓ to ℓ , i.e., $\Pi_\ell \cap X = 2\ell + \ell'$. Note that this is not defined on S nor on any lines contained in a plane contained inside X . Containing a plane is a divisorial condition in the moduli space, so for X outside this locus, we can resolve this map with one blowup $\text{Bl}_S F$ along the

surface S . We denote by E_S the exceptional divisor which is a \mathbb{P}^1 fibration over S

$$(3) \quad \begin{array}{ccccc} & & E_S & \hookrightarrow & \text{Bl}_S F \\ & \swarrow & & \searrow & \tilde{\phi} \\ & & \pi & & \\ S & \xrightarrow{\quad} & F & \xrightarrow{\quad} & F \\ & & \phi & & \end{array}$$

As it will be relevant for the count of nodal rational curves later, we record the following.

Theorem 2.4. ([OSY19, Theorem 0.2]) *If X is a very general cubic fourfold, then for every rational curve $C \subset F$ of class β there exists a unique $s \in S$ so that $C = \phi(q^{-1}(s))$.*

One can also define the following rational map

$$(4) \quad \begin{array}{l} \psi : F \dashrightarrow X \\ [\ell] \mapsto \ell \cap \ell' \end{array}$$

taking a general line to the intersection point with its residual in the unique tangent 2-plane. This is not defined on S but also on the following locus.

Definition 2.5. Denote by $V \subset F$ the locus of points $[\ell]$ so that there is a 2-plane Π_ℓ so that $\Pi_\ell \cap X = 3\ell$.

This locus appears briefly in the literature ([SV16, Definition 20.4] and also [CG72, Lemma 10.15] where in the case of a cubic threefold it is proven to be a finite set) and can also be viewed as the set of lines fixed by the Voisin map ϕ , that is those $[\ell] \in F$ with $\phi([\ell]) = [\ell]$.

We will need the follow results from intersection theory, which we put together for later use. In the following we denote by $l := \mathcal{O}_{I/F}(1)$ the tautological bundle, $\omega_p := \det(\Omega_p^1)$ the determinant of the relative cotangent bundle, and by $\lambda := \det(p_*\omega_p)$ the determinant of the Hodge bundle of the fibration p .

Lemma 2.6. *Let $X \subset \mathbb{P}^5$ be a smooth cubic fourfold. We have the following equalities in the Chow ring (or in cohomology)*

- (1) $H_F^4 = 108$, $H_F^2 c_2(\mathcal{U}_F) = 45$, $c_2^2(\mathcal{U}_F) = 27$,
- (2) $H_F^2 \sigma_2 = 63$, $\sigma_2^2|_F = 45$,
- (3) $\beta H_F = 3$, $\beta = \frac{1}{36} H_F^3$ and so $q_*[C_x] = \frac{1}{18} H_F^3$,
- (4) $q_* p^* H_X^2 = H_F$, $q_* p^* H_X^3 = \sigma_2|_F$, $q_* p^* H_X^4 = q_* p^* 3[x] = \frac{1}{6} H_F^3$,
- (5) $c_1(\omega_p) = q^* H_F + l$,
- (6) $q^* c_2(\mathcal{U}_F^\vee) = l^2 - l q^* H_F$,
- (7) $p_* q^* H_F^2 = 21 H_X$, $p_*(l^2) = 0$, $p_*(l q^* H_F) = 6 H_X$,
- (8) $[W] = 75 H_X$,
- (9) $p_*(c_1(\omega_p) q^*[S]) = 180 H_X^2$,
- (10) $\lambda = 9 H_X$,
- (11) $c_1(N_{S/F}) = 3 H_S \in H^2(S, \mathbf{Q})$, $c_2(N_{S/F}) = 1125$.

Proof. In the following we will repeatedly use the fact that if x a cycle in X and y a cycle in F with $\text{codim}(x) + \text{codim}(y) = 5$, then $q_* p^*(x) y = x p_* q^* y \in \mathbb{Z}$, from the projection formula.

(1) and (3) are in [Ame11, Lemma 4] (see also [OSY19]), whereas (2), (4) are standard intersection theory of Schubert classes combined with formula of the class of F given in (1).

(5) follows as we have $\omega_p = \omega_I - p^*\omega_X$, so by combining the facts that $\omega_X = -3H_X$, $p^*H_X = l$ and $\omega_I = \omega_q = q^*H_F - 2l$ we get the result. (6) is the Grothendieck relation.

For (7), the first follows as if we let $p_*q^*H_F^2 = aH_X$, then $p_*q^*H_F^2H_X^3 = 3a$ so, by Theorem 2.3 and (4), we have

$$\begin{aligned} p_*q^*H_F^2H_X^3 &= H_F^2q_*p^*(H_X^3) \\ &= H_F^2\sigma_2 \\ &= 63. \end{aligned}$$

For the second, $l^2 = p^*H_X^2$ implies $p_*l^2 = 0$. For the third,

$$\begin{aligned} p_*(lq^*H_F) &= p_*(p^*H_Xq^*H_F) = H_Xp_*q^*H_F \\ &= 6H_X, \end{aligned}$$

where for the final equality, if $p_*q^*H_F = m[1_X]$ with $m = p_*q^*H_F[x]$ for $x \in X$, then $m = H_Fq_*p^*[x] = H_F[C_x] = 6$.

For (8) let $[W] = aH_X$. Then $[W]H_X^3 = aH_X^4 = 3a$. We will show later in Proposition 3.7 that for X general, $p : q^{-1}(S) \rightarrow W$ is birational. Assuming this for the moment, we have $[W] = p_*q^*[S]$ and therefore

$$\begin{aligned} [W]H_X^3 &= p_*q^*[S]H_X^3 = [S]q_*p^*H_X^3 \\ &= 5\sigma_2^2|_F \\ &= 225, \end{aligned}$$

giving (8) if X is general (see below for the case of arbitrary smooth X).

For (9), recall from (5) that $c_1(\omega_p) = p^*H_X + q^*H_F$ and, for X general, that

$$\begin{aligned} p_*(p^*H_Xq^*[S]) &= H_Xp_*q^*[S] = H_X[W] \\ &= 75H_X^2. \end{aligned}$$

Assuming that X is very general, we have that $H^{2,2}(X, \mathbf{Z})_{\text{pr}} = 0$ from Deligne's Invariant Cycle Theorem (see [Huy20, Corollary 1.2.12]). Hence $p_*(q^*H_Fq^*[S]) = mH_X^2$ for some integer m , giving

$$\begin{aligned} 3m &= p_*(q^*H_Fq^*[S])H_X^2 = p_*q^*(H_F[S])H_X^2 \\ &= H_F[S]q_*p^*H_X^2 \\ &= 5H_F^2\sigma_2 = 5 \cdot 63. \end{aligned}$$

Therefore $p_*(q^*H_Fq^*[S]) = 105H_X^2$ giving the result if X is very general.

To extend (8) and (9) to every smooth cubic, note that all classes in question are defined globally on the universal family of smooth cubic fourfolds $f : \mathcal{X} \rightarrow |\mathcal{O}_{\mathbb{P}^5}(3)|_{\text{sm}}$. Hence, e.g., in (7), if we consider the class $\alpha = [\mathcal{W}] - 75H_{\mathcal{X}} \in H^2(\mathcal{X}, \mathbf{Z})$, then as $R^2f_*\mathbf{Z}$ is a local system and $\alpha_t = 0$ for t in the complement of a countable union of closed subsets, we obtain that $\alpha_t = 0$ for all $t \in U$. The analogous result for Chow groups is [Voi14, Lemma 3.2].

For (10), from the standard formula for λ (e.g., from [HM98, 3.110]) and (5) we have

$$\begin{aligned}\lambda &= \frac{1}{12}(p_*\omega_p^2 + [W]) \\ &= \frac{1}{12}(p_*(l + q^*H_F)^2 + 75H_X) \\ &= 9H_X. \quad \square\end{aligned}$$

For (11), note that $K_F = 0$ and Theorem 2.3 imply the first equality, whereas the second also follows from the description of S as a degeneracy locus and some computations involving the Harris–Tu formula, see [GK20, Proposition 4.1].

3. THE CURVE OF LINES THROUGH A POINT

In this section we collect and extend some facts about the $(2, 3)$ -intersection C_x parametrising lines through a point $x \in X \subset \mathbb{P}^5$ of a smooth cubic fourfold. As mentioned in the introduction, if X is general then C_x is always a curve, whereas for arbitrary X there can be only finitely many points where it is a surface. The following is well known to experts but we include the argument so as to fix notation and extract a bit further out of its proof.

Lemma 3.1. *Let $X \subset \mathbb{P}^5$ be a smooth cubic fourfold and $x \in X$ such that $C_x = p^{-1}(x)$ is 1-dimensional.*

- (1) *If ℓ is a line of first type through x , then C_x is smooth at $[\ell]$.*
- (2) *If ℓ is a line of second type through x , then C_x is singular at the point $[\ell]$.*

Proof. Let ℓ be a line of first type in X . We may assume that it is given by the equations $\mathbf{V}(x_2, x_3, x_4, x_5)$. Then the equation of X may take the form [CG72, 6.9].

$$(5) \quad \begin{aligned}F &= x_4x_0^2 + x_5x_0x_1 + x_3x_1^2 + x_0Q_0(x_2, x_3, x_4, x_5) + \\ &\quad x_1Q_1(x_2, x_3, x_4, x_5) + P(x_2, x_3, x_4, x_5) = 0.\end{aligned}$$

Since $x \in \ell$ we may assume that $x = [1, a, 0, 0, 0] \in \ell$ (the case $x = [0, 1, 0, 0, 0]$ is treated similarly). To find C_x , we write the equation as

$$\begin{aligned}x_4x_0^2 + x_5x_0(x_1 - ax_0) + ax_5x_0^2 + x_3(x_1 - ax_0)^2 + 2ax_0x_3(x_1 - ax_0) \\ + a^2x_0^2x_3 + x_0Q_0 + (x_1 - ax_0)Q_1 + ax_0Q_1 + P = 0.\end{aligned}$$

Putting $x_0 = 1$ and $x'_1 = x_1 - a$ we get

$$[a^2x_3 + x_4 + ax_5] + [x_5x'_1 + 2ax_3x'_1 + Q_0 + aQ_1] + [x_3x'^2_1 + x'_1Q_1 + P] = 0.$$

Note in this case that $\Pi_\ell = \mathbf{V}(x_3, x_4, x_5)$ is the tangent 2-plane to X along ℓ . The lines through $x = (0, 0, 0, 0, 0) \in \mathbb{A}^5$ (in the x'_1, x_2, x_3, x_4, x_5 coordinates) are determined by their slopes $[x'_1 : x_2 : x_3 : x_4 : x_5] \in \mathbb{P}^4$, with the slope of the line ℓ being $[1 : 0 : 0 : 0 : 0]$. They are parametrised by the following curve which is given by the system

$$\begin{aligned}a^2x_3 + x_4 + ax_5 &= 0, \\ T_2 = x_5x'_1 + 2ax_3x'_1 + Q_0(x_2, x_3, x_4, x_5) + aQ_1(x_2, x_3, x_4, x_5) &= 0,\end{aligned}$$

$$T_3 = x_3x_1'^2 + x_1'Q_1(x_2, x_3, x_4, x_5) + P(x_2, x_3, x_4, x_5) = 0.$$

Substituting x_4 from the 1st equation, $T_2 = 0, T_3 = 0$ become equations in the variables x_1', x_2, x_3, x_5 in \mathbb{P}^3 . The point in C_x which corresponds to the line ℓ is then $[1 : 0 : 0 : 0]$ (in the x_1', x_2, x_3, x_5 -coordinates). At this point the gradients of the surfaces T_2, T_3 are $\langle 0, 0, 2a, 1 \rangle$ and $\langle 0, 0, 1, 0 \rangle$ respectively. Hence the intersection is transversal and the curve C_x is smooth at the point $[\ell]$ with the tangent line given by $x_3 = x_5 = 0$ in \mathbb{P}^3 . Note then that the plane spanned by the line ℓ and the above tangent line is the tangent plane Π_ℓ to ℓ in X .

Let now ℓ be a line of second type, given again by $\mathbf{V}(x_2, x_3, x_4, x_5)$. Then the equation of X may take the form [CG72, 6.10]

$$(6) \quad \begin{aligned} F = & x_4x_0^2 + x_5x_1'^2 + x_0Q_0(x_2, x_3, x_4, x_5) + x_1'Q_1(x_2, x_3, x_4, x_5) \\ & + P(x_2, x_3, x_4, x_5) = 0. \end{aligned}$$

A similar calculation shows that with centre the point $x = [1 : a : 0 : 0 : 0 : 0] \in \ell$ (the case $x = [0 : 1 : 0 : 0 : 0 : 0]$ is treated similarly) the equation of X becomes (by putting $x_0 = 1$ and $x_1 - a = x_1'$)

$$[x_4 + a^2x_5] + [2ax_5x_1' + Q_0 + aQ_1] + [x_5x_1'^2 + x_1'Q_1 + P] = 0.$$

Note in this case that $H_\ell = \mathbf{V}(x_4, x_5)$ is the tangent 3-space to X along ℓ . Then C_x is the intersection of two surfaces in a 3-dimensional projective space in variables x_1', x_2, x_3, x_5 with gradients at $\langle 0, 0, 0, 2a \rangle$ and $\langle 0, 0, 0, 1 \rangle$ respectively at the point $[1 : 0 : 0 : 0]$ corresponding to ℓ . Hence the intersection is not transversal at the point $[\ell]$, C_x is singular there with the tangent cone given by the plane $x_5 = 0$ in \mathbb{P}^3 . Note then that the space spanned by the line ℓ and the above tangent cone is the tangent space H_ℓ to ℓ in X . \square

To summarise the geometry, for $x \in \ell \subset X$, the tangent space to the line $\ell \subset X$ is spanned by ℓ and the tangent cone to C_x at the point $[\ell]$. This space has dimension 2 (i.e., $[\ell]$ is a line of first type) if and only if C_x is smooth at $[\ell]$ and has dimension 3 (i.e., $[\ell]$ is a line of second type) if and only if C_x has a singularity at $[\ell]$ (with tangent cone of dimension two). When we consider C_x as a curve in F (i.e., as $q(p^{-1}(x))$), the singular points correspond to the intersection points with the surface S . When $x \notin W$ this intersection is empty and the curve C_x is smooth.

Remark 3.2. In [CG72, Definition 6.6], lines of first or second type are defined as those for which the image of the dual mapping is a smooth plane conic or a two-to-one covering of the projective line respectively.

In fact, when the above line ℓ is of second type, $\nabla F([s : t : 0 : 0 : 0 : 0]) = \langle 0, 0, 0, 0, s^2, t^2 \rangle$ and the fibres of the two-to-one covering are formed by the points $[s : \pm t : 0 : 0 : 0 : 0]$, with ramification points the two points at infinity $x_0 = [1 : 0 : 0 : 0 : 0 : 0]$, $x_1 = [0 : 1 : 0 : 0 : 0 : 0]$. With $x = [1 : a : 0 : 0 : 0 : 0] \in \ell$, $a \neq 0$ and $Y_x = T_x X \cap X$, one sees that Y_x contains ℓ and has two singular points on ℓ , namely x and $x' = [1 : -a : 0 : 0 : 0 : 0]$, the conjugate of x under the dual mapping. For $a = 0$, Y_{x_0} has a non-ordinary singularity at x_0 and the same for Y_{x_1} . Finally note that for a point $x \in \ell$, with $[\ell] \in S$, the singularity of C_x corresponding

to ℓ has the same type as the section $Y_x = Y_{x'}$ has at the conjugate point x' (see [W05, Section 2, pg.6, case (i)], [CML09, Section 3.1]). Hence if x' is not in the Hessian this singularity is nodal. If there are several second type lines containing x , each one induces a singularity as above.

On the other hand, if $[\ell] \in F \setminus S$ and $x \in \ell$ then Y_x does not have a singularity on ℓ other than x . In particular, if $x \notin W$ then the only singularity of Y_x is at x .

Lemma 3.3. *We have the following two correspondences*

- (1) *between pairs (Y, x) where $Y \subset \mathbb{P}^4$ is a cubic threefold with a singular ordinary double point x (resp. A_2 singularity at the point x), and smooth non-hyperelliptic curves of genus 4 with canonical image which lies on a smooth quadric (resp. quadric cone).*
- (2) *between pairs (Y, x) where $Y \subset \mathbb{P}^4$ is a cubic threefold with two collinear ordinary double points so that x is one of them, and smooth non-hyperelliptic genus 3 curves with two marked points and with nodal canonical image lying on a smooth quadric.*

In both cases the canonical image of the corresponding curve parametrises the lines in Y through x .

Remark 3.4. The cases of (1) above are exactly the smooth (2, 3) complete intersections in \mathbb{P}^3 . Note that in the case of a quadric cone, as the curve is smooth, it necessarily does not pass through the vertex of the cone.

Proof. The first has appeared many times in the literature, see, e.g., [CG72, p.306-307] and [CML09, §3.1]. The second is a minor modification of this so we briefly sketch the construction. For $(C', p_1, p_2) \in \mathcal{M}_{3,2}^{nhyp}$, the linear system $|K_{C'} + p_1 + p_2|$ gives a morphism $C' \rightarrow \mathbb{P}^3$ with image a sextic curve C with one node p . Just as in the classical construction of the canonical embedding of a non-hyperelliptic genus 4 curves, $H^0(\mathbb{P}^3, I_C(2)) = \mathbf{C}$ so there is a unique quadric Q containing C , which we may assume is smooth by genericity. Also, the linear system $|I_C(3)|$ induces a birational map $h : \mathbb{P}^3 \dashrightarrow \mathbb{P}^4$ with image a singular cubic threefold Y . This can also be realised by blowing up \mathbb{P}^3 at the curve C to obtain a variety \tilde{Y} with one node (C and \tilde{Y} have the same singularity count and type) lying above the node of C , and blowing down the strict transform of the quadric Q to obtain Y . The threefold Y will be singular both at the image x of the quadric but also the image of the singular point of \tilde{Y} and these two must lie on the same line. Projecting from x gives the inverse map to h . \square

Note that from Lemma 2.2, if X is general, the locus $W \subset X$ spanned by lines of second type is an irreducible divisor.

Proposition 3.5. *Let $X \subset \mathbb{P}^5$ be a general cubic fourfold and $W \subset X$ the locus spanned by lines of second type. For any $x \in X$ the curve $C_x \subset F$ parametrising lines passing through x is reduced and connected. Moreover, the following hold.*

- (1) *For $x \in X \setminus W$ the curve C_x is irreducible and smooth of genus 4 and if x is general then it is Brill–Noether general.*

- (2) For $x \in W$ general, C_x is irreducible and has only one node and its normalisation is a Brill–Noether general curve of genus 3 with the two preimages of the node being general points.
- (3) For any $x \in W$ the total Tjurina number of C_x is ≤ 6 , and if reducible, C_x is the union of two irreducible, at worst nodal, plane cubics meeting at 3 distinct points. Hence, there can be at most 4 lines of second type through x .

Proof. From [LSV17, Lemma 2.7] we know that C_x is reduced for all $x \in X$, and that it is connected follows from [BVdV79]. For any x not in W , from Lemma 3.1 it follows that C_x is also smooth and hence irreducible, proving the first part of (1). We now prove the remainder of (1) and (2).

Let C be a general curve of genus 4 (resp. a general arithmetic genus 4 curve with one node). From Lemma 3.3 we obtain a cubic threefold Y with one (resp. two) nodes, in such a way that C parametrises lines contained in Y passing through the marked node p_0 . Write the equation determining Y as $F(x_0, \dots, x_4) = 0$ with singularity at the marked node $p_0 = [1 : 0 : \dots : 0]$ (in such a way that the other node is not on the $x_0 = 0$ hyperplane), and consider the cubic equation

$$G(x_0, \dots, x_5) = F(x_0, \dots, x_4) + x_0^2 x_5.$$

It is easy to check that $X_0 = \mathbf{V}(G)$ defines a smooth cubic fourfold with Y as the tangent hyperplane section at the point $[1 : 0 : \dots : 0]$, so that we have $p_0 \in X_0$ and that $C = C_{p_0}$ is the curve of lines through p_0 . Note that in the second case, Y will be a hyperplane section of X_0 with two nodes, and the line between these nodes is of second type (cf., Remark 3.2 for the converse statement that the tangent hyperplane at a general point of a general line of second type has two singularities).

Let now

$$\mathcal{X} \rightarrow |\mathcal{O}_{\mathbb{P}^5}(3)|_{\text{sm}}$$

the universal family of smooth cubic fourfolds, $p : \mathcal{I} \rightarrow \mathcal{X}$ the projection from the universal family of lines on all of them, and $q : \mathcal{I} \rightarrow \mathcal{F}$ the morphism to the universal Fano variety of lines (we discuss this and some of the universal families mentioned below in the beginning of the following section). From semicontinuity of fibre dimension, the generic fibre of the morphism $p : \mathcal{I} \rightarrow \mathcal{X}$ is a smooth curve of genus 4. In fact, [CS09, Lemma 2.1, Corollary 2.2] prove that for any smooth cubic fourfold X , there are only finitely many points in X over which the fibre of $p : I \rightarrow X$ is not 1-dimensional, so this locus is codimension 4 in \mathcal{X} . The universal locus $\mathcal{S} \subset \mathcal{F}$ of second type lines admits a projection to the Grassmannian $G(2, 6)$ with isomorphic, irreducible fibres (see [Huy20, Proposition 2.2.13] and its proof). Hence \mathcal{S} is irreducible and of codimension 2 in \mathcal{F} so that the total space $q^{-1}(\mathcal{S})$ of second type lines in \mathcal{I} is irreducible and likewise its image $\mathcal{W} \subset \mathcal{X}$, the locus of points on lines of second type, which is a divisor in \mathcal{X} since $q^{-1}(\mathcal{S}) \rightarrow \mathcal{W}$ is generically finite from Lemma 2.2. We conclude that the non-empty open subset of \mathcal{W} parametrising points where the fibres of p are curves is irreducible.

Since a general X degenerates to X_0 , so will the corresponding family of curves C_x degenerate to C , proving (1), since being Brill–Noether general is

an open condition. Similarly, the curve C_x for a general point $x \in W \subset X$ must be Brill–Noether general and irreducible with one node, since being nodal is an open condition in a flat family of curves and W is irreducible. This completes (2).

We now prove (3). As mentioned in Section 2, the class of C_x in cohomology is 2β , and since for the very general X the class β generates the algebraic part of cohomology, it must be the case that for general X the class β is irreducible, so that every curve representing it must be irreducible, implying C_x has at most two irreducible components. From [NO21, Lemma 3.8] it follows that each such irreducible component has arithmetic genus 1 and at worst nodes, implying that the two components must meet at three distinct points as the total arithmetic genus is 4. The statement about singularities follows from the correspondence between singularities of the blow up \tilde{Y}_x at x of the tangent hyperplane section $Y_x = T_x X \cap X$ and those of C_x of [CML09, 3.1-3.3] and from the fact that for a general X , every hyperplane has isolated singularities of Tjurina number at most 6 from [LSV17, Corollary 3.7]. As the arithmetic genus of C_x is four and from Lemma 3.1 every singularity corresponds to a second type line through x , the above discussion implies that there can be at most 4 lines of second type through x . \square

Remark 3.6. In particular, from Lemma 3.1 the curve C_x can have at most four singular points for x an arbitrary point of a general cubic fourfold. It would be desirable to have a complete list of possible singularities which occur for C_x and a description of the corresponding loci in X .

Corollary 3.7. *For $X \subset \mathbb{P}^5$ a general cubic fourfold, the morphism $p : \mathbb{P}(\mathcal{U}_S) \rightarrow W$ is birational and the degree of W in \mathbb{P}^5 is 225.*

Proof. That through a general point $x \in W$ there passes precisely one second type line follows from the fact that each second type line through x forces an extra singularity of C_x by Lemma 3.1, so we conclude from Proposition 3.5, as the generic $x \in W$ has C_x irreducible with one node. That the degree of W is 225 was already computed in Lemma 2.6, under the assumption that the degree of the above map is one. \square

Remark 3.8. More generally, let $n \geq 3$ and let $X \subset \mathbb{P}^{n+1}$ be a general cubic n -fold, F its Fano scheme of lines and $F_2 \subset F$ the locus of lines of second type, which is smooth of dimension $n - 2 = \frac{1}{2} \dim F$ from [Huy20, Proposition 2.2.13]. Consider the induced morphism from the family of lines

$$\begin{array}{ccc} \mathbb{P}(\mathcal{U}_{F_2}) & \xrightarrow{p} & W \subset X \\ \downarrow q & & \\ F_2 & & \end{array}$$

for \mathcal{U} the universal bundle on the Grassmannian $G(2, n + 2)$. The same argument as in Proposition 3.5 proves that p is birational onto its image, a divisor in X , and we summarise this in the following two paragraphs.

First, there is a correspondence between pairs (Y, x) where $Y \subset \mathbb{P}^n$ is a cubic $(n - 1)$ -fold with two collinear ordinary double points, one of which is x , and $(2, 3)$ -complete intersections in \mathbb{P}^{n-1} with one ordinary double point.

For example (see [DvGK05]), if $n = 3$, this will be between cubic surfaces with two collinear nodes, and 0-cycles in \mathbb{P}^2 of length 6 with support on 5 distinct points, one of which is non-reduced. To explain the correspondence, call Q and C the quadric and cubic respectively, and M their intersection. The correspondence follows from the diagram

$$\begin{array}{ccc} & \text{Bl}_x(Y) & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ Y & \overset{h}{\dashrightarrow} & \mathbb{P}^{n-1} \end{array}$$

where π_1 is the blowup map, π_2 contracts the family of lines through x and h is the projection from the marked node. The image of the exceptional divisor $\pi_1^{-1}(x)$ under π_2 is the quadric Q , whereas π_2 is the blowup of the intersection M of Q with a cubic C . The sheaf $I_M(3)$ has $n + 1$ sections, and this linear system induces the inverse rational map to h with image Y . Starting with M , the blowup $\text{Bl}_M(\mathbb{P}^{n-1})$ is singular at a point y , and the strict transform of Q can be contracted to a point x , collinear to y , in a cubic $(n - 1)$ -fold Y with ordinary double points at x, y .

Now for $x \in X$ a general point on a general line ℓ of second type, the tangent hyperplane section $Y_x = T_x X \cap X$ will have a node at x and at precisely one other point on ℓ from Remark 3.2. The corresponding $(2, 3)$ -complete intersection parametrises lines through x and will have precisely one ordinary double point. As in the fourfold case, this implies the map p is birational, or that the ramification of the map $p : I \rightarrow X$ is generically as simple as possible along W .

We give now a geometric interpretation of the fact that ℓ and its residual ℓ' (i.e., the image under the Voisin map) meet at a point $x \in \ell \cap \ell' \subset X$ in terms of ramification points of the g_3^1 's on C_x . In particular, we will be interested in the degenerate case $\ell = \ell'$.

Lemma 3.9. *Let (Y, x) be a pair of a cubic threefold $Y \subset \mathbb{P}^4$ and a point $x \in Y$ as in either of cases (1) or (2) of Lemma 3.3 and let C_x be the sextic curve parametrising lines through x . For $t \in C_x$, we denote the corresponding line by ℓ_t . Then the following are true.*

- (1) *For $t_1, t_2 \in C_x$ and $\langle \ell_{t_1}, \ell_{t_2} \rangle \cap Y = \ell_{t_1} + \ell_{t_2} + \ell$, we have that $[\ell] \in C_x$ if and only if t_1, t_2 are in the same ruling of the quadric containing C_x . In this case $\ell = \ell_{t_{12}}$ for $t_{12} \in C_x$ the third point of intersection of C_x with that ruling.*
- (2) *The residual line $\ell_{t'}$ in the intersection of Y with a 2-plane tangent to the cone of lines through x at the line ℓ_t also passes through x if and only if t is a ramification point of one of the possibly two g_3^1 's that C_x possesses. In particular, $\ell_t = \ell_{t'}$ if t is a triple ramification point of one of the possibly two g_3^1 's that C_x possesses.*

Proof. The first part is basically [CG72, Lemma 8.6], whereas the second follows from the first. \square

Recall that for a cubic fourfold $X \subset \mathbb{P}^5$, the divisor $W \subset X$ is the locus of points x whose curve C_x is singular. Proposition 3.5 implies that if $W' \subset W$

is the closure of the set

$$(7) \quad \{x \in X : C_x \text{ has worse than irreducible 1-nodal singularities}\}$$

then W' has codimension at least 2 in X .

Proposition 3.10. *Let $X \subset \mathbb{P}^5$ be a general cubic fourfold and $x \in X \setminus W'$. For ξ a 0-dimensional subscheme of length 2 on C_x , i.e., $\xi = t_1 + t_2$ for two possibly non-distinct points in C_x , the 2-plane spanned by ℓ_{t_1}, ℓ_{t_2} (and in the case $t_1 = t_2$, a 2-plane tangent to ℓ_{t_1}) meets X at a line ℓ_{t_3} which passes through x if and only if t_1, t_2, t_3 are a fibre of one of the possibly two g_3^1 's on C_x .*

Proof. Apply the previous lemma to the nodal cubic threefold $Y = T_x X \cap X$, noting that such lines ℓ_{t_i} are contained in Y - recall, also, the summary of the geometry following Lemma 3.1. \square

Suppose we are given a pair (ℓ, Π) , with ℓ a line in X and Π a tangent 2-plane to X along ℓ , and a point $x \in \ell \setminus W'$. Let $\ell = \ell_t$ and let $\ell' = \ell_{t'}$ the residual line in the intersection of X with Π , with $t, t' \in C_x$. Then by the above proposition we have that $x \in \ell_t \cap \ell_{t'}$ if and only if t is a ramification point of one of the (possibly) two g_3^1 's on C_x and t' the residual to that ramification point in the corresponding g_3^1 . In particular we have $\ell' = \ell = \ell_t$ if and only if t is a triple ramification point. In Section 5 we will study the locus in the universal family of lines I of ramification points (and their residuals) of the g_3^1 's of the curves C_x .

From Proposition 3.5 there is a rational map

$$j : X \dashrightarrow \overline{\mathcal{M}}_4$$

not defined on a codimension two locus, and whose image hits the boundary in the component Δ_0 of 1-nodal curves. In the following, denote by $D_3 \subset \overline{\mathcal{M}}_4$ the divisor which is the closure of the divisor parametrising smooth genus four curves which have a g_3^1 which admits a triple ramification point.

Lemma 3.11. *Let X be a general cubic fourfold and consider $j : X \dashrightarrow \overline{\mathcal{M}}_4$ the induced moduli map, taking a point x to $[C_x]$. Then*

$$j^*[D_3] = 126H_X.$$

Proof. From [Har84], the class of D_3 on the moduli space can be computed as follow

$$[D_3] = 2(132\lambda - 15\delta_0 + \text{other boundary components}).$$

By Lemma 2.6 we get that

$$j^*[D_3] = 2(132 \cdot 9 - 15 \cdot 75)H_X = 126H_X. \quad \square$$

Corollary 3.12. *Let $X \subset \mathbb{P}^5$ be a general cubic fourfold. The locus $V \subset F$ of triple lines is non-empty.*

Proof. From Lemma 3.11 we obtain that the locus $x \in X$ so that C_x has a triple ramification point is non-empty and divisorial. In particular, from Lemma 3.9 and Proposition 3.10 we see that V must be non-empty as required. \square

Remark 3.13. It is also possible to give an equivalent to Corollary 3.7 for the locus V of triple lines. We will later in Theorem 4.7 compute that $[V] = 21c_2(\mathcal{U}_F) \in \text{CH}^2(F)$, so as from Lemma 2.6, $q_*p^*H_X^3 = \sigma_2|_F$, we get $21c_2(\mathcal{U}_F)\sigma_2 = 21(c_2(\mathcal{U}_F)H_F^2 - c_2^2(\mathcal{U}_F)) = 3 \cdot 126$ and as $H_X^4 = 3$, we conclude that

$$p_*q^*[V] = 126H_X.$$

As this agrees with j^*D_3 above, we obtain that the morphism

$$q^{-1}(V) \rightarrow p(q^{-1}(V))$$

is birational. In particular, the generic point $x \in X$ which has a triple line through it has exactly one triple line through it.

4. UNIVERSAL FAMILIES AND THE SURFACE V

Throughout this section $X \subset \mathbb{P}^5$ will be a general cubic fourfold, F its Fano variety of lines and we will denote by V, S etc. as in Section 2. We will often consider the following special linear subspaces in \mathbb{P}^5

$$(8) \quad \ell_0 = \mathbf{V}(x_2, x_3, x_4, x_5), \quad \Pi_0 = \mathbf{V}(x_3, x_4, x_5), \quad H_0 = \mathbf{V}(x_4, x_5).$$

Recall first that the Voisin map $\phi : F \dashrightarrow F$ is resolved by blowing up the locus S to give Diagram (3). Consider now the incidence variety

$$\tilde{F} = \{(\ell, \Pi) : \Pi \text{ is tangent to } X \text{ along } \ell\} \subset \text{G}(2, 6) \times \text{G}(3, 6).$$

Lemma 4.1. ([Huy20, Remark 2.2.18]) *The blowup $\text{Bl}_S(F)$ is isomorphic to the incidence variety \tilde{F} , so in particular the fibre $E_{S, [\ell]} \cong \mathbb{P}^1$ of $\pi_{E_S} : E_S \rightarrow S$ over $[\ell]$ parametrises the 2-planes tangent along ℓ .*

The map ψ of (4) lifts to a rational map $\psi : \tilde{F} \dashrightarrow X$, not defined at the locus of pairs (ℓ, Π) as above for which $\Pi \cap X = 3\ell$. This is the strict transform \tilde{V} in \tilde{F} of the locus V of Definition 2.5. We will see later in Corollary 5.5 that the map ψ is resolved by blowing up \tilde{V} .

In this section we will need to construct the universal families of the varieties mentioned above and derive various of their properties from these constructions. First, we note that we may construct the universal cubic hypersurface $\mathcal{X} \subset \mathbb{P}^5 \times |\mathcal{O}_{\mathbb{P}^5}(3)|$ as the vanishing locus of the universal section $s_{\mathcal{X}} \in \text{H}^0(\mathbb{P}^5 \times |\mathcal{O}_{\mathbb{P}^5}(3)|, \mathcal{O}(3, 1))$, and it is not hard to also construct the universal family $\mathcal{F} \subset |\mathcal{O}_{\mathbb{P}^5}(3)| \times \text{G}(2, 6)$, whose fibre under the first projection over a cubic $X = \mathbf{V}(G)$ is the Fano scheme of lines F of X (see [Huy20, §2]). The second projection realises \mathcal{F} as a projective bundle $\mathbb{P}(K)$ where K is a rank 52 vector bundle on $\text{G}(2, 6)$ sitting in the following sequence

$$(9) \quad 0 \rightarrow K \rightarrow S^3(\mathbf{C}^6)^\vee \otimes \mathcal{O}_{\text{G}(2,6)} \rightarrow S^3\mathcal{U}^\vee \rightarrow 0$$

whose fibre over a line $[\ell] \in \text{G}(3, 6)$ is $\mathbb{P}(\text{H}^0(\mathbb{P}^5, I_\ell \otimes \mathcal{O}(3)))$. Finally, it was proven in [Ame11, Lemma 1] (see also [Huy20, §3]) that the universal family $\mathcal{S} \subset \mathcal{F}$ of surfaces of second type lines, with fibre over a smooth $X = \mathbf{V}(G)$ the surface S of type two lines contained in X , is smooth when restricted to the locus of smooth cubics, and has non-empty fibres there.

We continue now by exploring universal constructions for \tilde{F} and \tilde{V} . Define first the two spaces

$$(10) \quad \begin{aligned} \mathcal{G} &= \{(\ell, \ell', \Pi) : \ell, \ell' \subset \Pi\} \subset \mathbf{G}(2, 6) \times \mathbf{G}(2, 6) \times \mathbf{G}(3, 6), \\ \mathcal{G}_\Delta &= \{(\ell, \ell', \Pi) \in \mathcal{G} : \ell = \ell'\}. \end{aligned}$$

Note that \mathcal{G} is smooth, irreducible and of dimension 13 since it is the self-product of the universal family of lines contained in a 2-plane in \mathbb{P}^5 , i.e., is a $\mathbb{P}^2 \times \mathbb{P}^2$ -fibration over the 9-dimensional $\mathbf{G}(3, 6)$. Similarly, \mathcal{G}_Δ is smooth, irreducible and of dimension 11 and in this case we refer to its points simply as tuples (ℓ, Π) . As it will be needed in various statements in the rest of this section, we detail the local structure of \mathcal{G} .

Our aim is to describe an affine open set U of \mathcal{G} . A choice of local coordinates $x = (x_{03}, \dots, x_{25})$ of an affine open \mathbf{C}^9 in $\mathbf{G}(3, 6)$ determines a 2-plane $\Pi(x)$ given by the row space of the matrix

$$A(x) = \begin{pmatrix} 1 & 0 & 0 & x_{03} & x_{04} & x_{05} \\ 0 & 1 & 0 & x_{13} & x_{14} & x_{15} \\ 0 & 0 & 1 & x_{23} & x_{24} & x_{25} \end{pmatrix}.$$

Then, for $y = (y_1, y_2)$ affine coordinates of \mathbf{C}^2 , a line $\ell(x, y) \subset \Pi(x)$ is the row space of the matrix

$$B(x, y) = \begin{pmatrix} 1 & 0 & y_1 & x_{03} + x_{23}y_1 & x_{04} + x_{24}y_1 & x_{05} + x_{25}y_1 \\ 0 & 1 & y_2 & x_{13} + x_{23}y_2 & x_{14} + x_{24}y_2 & x_{15} + x_{25}y_2 \end{pmatrix}.$$

In parametric form, if $[s_0 : s_1 : s_2]$ are the coordinates of \mathbb{P}^2 , then the 2-plane $\Pi(x)$ in \mathbb{P}^5 is given by $[(s_0, s_1, s_2)A(x)]$, whereas for $[\lambda : \mu]$ coordinates of \mathbb{P}^1 , the line $\ell(x, y)$ is given by $[(\lambda, \mu)B(x)]$, and its Cartesian equation in $\mathbb{P}^2 \cong \Pi(x)$ is $s_0y_1 + s_1y_2 - s_2 = 0$. We then define

$$(11) \quad U = \{(\ell(x, y), \ell(x, y'), \Pi(x)) : (x, y, y') \in \mathbf{C}^9 \times \mathbf{C}^2 \times \mathbf{C}^2\}.$$

Geometrically, the points of U correspond to projective planes $\Pi(x)$ in \mathbb{P}^5 which do not intersect the projective plane $\Pi(0) = \Pi_0 = \mathbf{V}(x_0, x_1, x_2)$ and lines ℓ, ℓ' in $\Pi(x)$ which do not contain the point $[0, 0, 1]$ of $\mathbb{P}^2 \cong \Pi(x)$. Note that when $y = y'$ then we are in the subspace \mathcal{G}_Δ , the intersection is non-empty and for $x = y = y' = 0$ we are centered at the point $(\ell_0, \Pi_0) \in \mathcal{G}_\Delta$.

For $X = \mathbf{V}(G)$ a cubic fourfold, denote by

$$(12) \quad \begin{aligned} \tilde{\mathcal{F}} &= \{(X, (\ell, \ell', \Pi)) : X \cap \Pi = 2\ell + \ell'\} \subset |\mathcal{O}_{\mathbb{P}^5}(3)| \times \mathcal{G}, \\ \tilde{\mathcal{V}} &= \{(X, (\ell, \ell', \Pi)) : \ell = \ell'\} \subset \tilde{\mathcal{F}}. \end{aligned}$$

We have a diagram

$$\begin{array}{ccc} \tilde{\mathcal{F}} & \xrightarrow{\text{pr}_1} & |\mathcal{O}_{\mathbb{P}^5}(3)| \\ & \downarrow \text{pr}_2 & \\ & \mathcal{G} & \end{array}$$

and observe that $\tilde{\mathcal{V}} = \text{pr}_2^{-1}(\mathcal{G}_\Delta)$. The fibre of $\tilde{\mathcal{F}}$ over a general point $X \in |\mathcal{O}(3)|$ is the graph of the Voisin map, i.e., isomorphic to the blowup \tilde{F} of F at S under the map $(X, (\ell, \ell', \Pi)) \mapsto (X, (\ell, \Pi))$ (see Lemma 4.1). The

fibre of $\tilde{\mathcal{V}}$ is the strict transform \tilde{V} of the locus V of triple lines in X in the blowup \tilde{F} , which as we saw in Corollary 3.12 is non-empty.

Proposition 4.2. *The varieties $\tilde{\mathcal{F}}, \tilde{\mathcal{V}}$ are \mathbb{P}^{46} -fibrations over \mathcal{G} , respectively \mathcal{G}_Δ . In particular they are both smooth.*

Proof. The action of $\mathrm{PGL}(5)$ on \mathbb{P}^5 induces a transitive action on both $\tilde{\mathcal{F}} \setminus \tilde{\mathcal{V}}$ and $\tilde{\mathcal{V}}$ which commutes with the projections to $\mathcal{G} \setminus \mathcal{G}_\Delta$ and \mathcal{G}_Δ respectively and sends fibres isomorphically to fibres. The $\mathrm{PGL}(5)$ -orbit of U in \mathcal{G} gives a covering of \mathcal{G} by affine open sets isomorphic to U over which the fibrations $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{V}}$ are isomorphic to the corresponding ones over U . To show the above claim, it suffices then to show that the above universal families are, over U , projectivisations of affine bundles. Taking this for granted, we have a diagram

$$(13) \quad \begin{array}{ccc} \tilde{\mathcal{F}}_U^a & \xrightarrow{\mathrm{pr}_1} & M = \mathrm{H}^0(\mathbb{P}^5, \mathcal{O}_{\mathbb{P}^5}(3)) \\ \uparrow & \nearrow \mathrm{pr}_1 & \\ \tilde{\mathcal{V}}_U^a & & \end{array}$$

namely the affine bundle version of that in (12).

We now show the last claim. Let $w = (w_{k_0 \dots k_5} : \sum k_i = 3)$ be coordinates for $M \cong \mathbf{C}^{56}$, and for one such w , denote by G_w the corresponding cubic hypersurface. From the above local analysis, $\tilde{\mathcal{F}}_U^a \subset M \times U$ consists of those $(G_w, (\ell(x, y), \ell(x, y'), \Pi(x)))$ so that

$$(14) \quad \begin{aligned} G_w(s_0, s_1, s_2, s_0x_{03} + s_1x_{13} + s_2x_{23}, s_0x_{04} + s_1x_{14} + s_2x_{24}, \\ s_0x_{05} + s_1x_{15} + s_2x_{25}) = c(s_0y_1 + s_1y_2 - s_2)^2(s_0y'_1 + s_1y'_2 - s_2) \end{aligned}$$

for some constant c . Letting z be a coordinate for \mathbf{C} parametrising c , we have that $\tilde{\mathcal{F}}_U^a$ is the image of the locus $\mathcal{A} \subset M \times U \times \mathbf{C}$ cut out by the above equation (14) (with c replaced by z). This is a cubic polynomial equation in the variables s_0, s_1, s_2 and therefore corresponds to a system of 10 polynomial equations $E_{t_0 t_1 t_2}$ indexed by $s_0^{t_0} s_1^{t_1} s_2^{t_2}$ so that $t_0 + t_1 + t_2 = 3$. They have the form

$$E_{t_0 t_1 t_2} : w_{t_0 t_1 t_2 000} + (\text{terms not containing } w_{t_0 t_1 t_2 000}) = 0,$$

so that the Jacobian matrix of the system, which is a 10×70 matrix, is such that the first 10×10 minor is the identity. Hence, the above system defines a smooth complete intersection of dimension 60. Note also that $c = -G_w(0, 0, 1, x_{23}, x_{24}, x_{25})$, so that \mathcal{A} is bijective onto its image $\tilde{\mathcal{F}}_U^a \subset M \times U$. Moreover when we fix a point $(x, y, y') \in U$, the fibre of \mathcal{A} over that point (resp. its image in $M \times U$) is a linear subspace of $M \times \mathbf{C} \cong \mathbf{C}^{57}$ (resp. of $M \cong \mathbf{C}^{56}$). In fact, when we fix the plane $\Pi(0)$ (as we may do up to a projective transformation) the fibre over the point $(x = 0, y, y') \in U$ is given by the polynomials

$$(15) \quad \begin{aligned} G = & x_5 Q_5(x_0, \dots, x_5) + x_4 Q_4(x_0, \dots, x_4) + x_3 Q_3(x_0, \dots, x_3) + \\ & c(x_0 y_1 + x_1 y_2 - x_2)^2 (x_0 y'_1 + x_1 y'_2 - x_2). \end{aligned}$$

Observe that there are $47 = \binom{7}{2} + \binom{6}{2} + \binom{5}{2} + 1$ coefficients involved. All in all, the above equations define an affine bundle over the thirteen dimensional base U of rank $47 = 60 - 13$. Finally, when $y = y'$ the above gives a local description of $\tilde{\mathcal{V}}^a$ over an open of \mathcal{G}_Δ , isomorphic to $\mathbf{C}^9 \times \mathbf{C}^2$. The varieties $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{V}}$ locally over U are the projectivisations of the affine bundles $\tilde{\mathcal{F}}_U^a$ and $\tilde{\mathcal{V}}_U^a$ respectively. Under the action of $\mathrm{PGL}(5)$, we get that they are locally projectivisations of affine bundles. \square

With notation and local study of the universal families established, we may now proceed to studying the main object of this section, namely the surface V .

Lemma 4.3. *The locus $\tilde{V} \subset \tilde{F}$ is closed, smooth and of pure dimension 2. Hence, also its image $V \subset F$ is closed and 2-dimensional.*

Proof. From Proposition 4.2 $\tilde{\mathcal{V}}$ is of dimension 57 and smooth. As from Corollary 3.12 the first projection $\mathrm{pr}_1 : \tilde{\mathcal{V}} \rightarrow |\mathcal{O}_{\mathbb{P}^5}(3)|$ is onto its image (a space of dimension 55), we obtain from generic smoothness that the generic fibre is closed, smooth and of pure dimension 2. \square

Theorem 4.4. *The surface \tilde{V} is irreducible, hence also its image V in F .*

Proof. Note first that irreducibility will follow from Lemma 4.3 if we can show connectedness. We follow the strategy of [BVdV79, Theorem 6], which we shall summarise before delving into the technical details. We continue using the notation established in various proofs above.

Consider now again the projection

$$\mathrm{pr}_1 : \tilde{\mathcal{V}} \rightarrow M = H^0(\mathbb{P}^5, \mathcal{O}_{\mathbb{P}^5}(3)),$$

which for a smooth cubic equation $G \in M$ so that $X = \mathbf{V}(G)$, has fibre \tilde{V} the strict transform in \tilde{F} of the surface V of triple lines in X . The Stein factorisation of pr_1 is as follows

$$\begin{array}{ccc} \tilde{\mathcal{V}} & & \\ \downarrow \mathrm{pr}_1 & \searrow \rho & \\ & & W \\ & \swarrow \sigma & \\ M & & \end{array}$$

where ρ has connected fibres, σ is finite and W is a normal variety. As the base M is simply connected, either σ is a bijection in which case the theorem is proven, or there is a ramification divisor $D \subset M$ and a branch divisor $D' \subset W$. As explained above, \mathcal{G}_Δ is covered by opens isomorphic to $U = \{(\ell(x, y), \Pi(x)) : (x, y) \in \mathbf{C}^9 \times \mathbf{C}^2\}$, (i.e., (11) restricted to \mathcal{G}_Δ where we have taken $y = y'$) over which the fibration $\tilde{\mathcal{V}}$ is isomorphic to the one over U , that is isomorphic to the projectivisation of the affine bundle $\tilde{\mathcal{V}}_U^a$. We now restrict to the latter and consider the analogue of the above diagram there.

If we denote by $\tilde{D} \subset \tilde{\mathcal{V}}_U^a$ an irreducible component of the inverse image of D' , then the differential

$$dpr_1 : T_{\tilde{\mathcal{V}}_U^a} \rightarrow T_M$$

cannot be surjective along points of \tilde{D} . We will now arrive at a contradiction by arguing that the locus of points where the differential is not surjective lies in codimension at least 2, thus ruling out this eventuality. We will prove this by arguing that the non-surjective locus is of codimension at least two in every fibre of $pr_2 : \tilde{\mathcal{V}}_U^a \rightarrow U \subset \mathcal{G}_\Delta$, with notation as in (12), (10). As all fibres are isomorphic, it suffices to prove this over the point (ℓ_0, Π_0) as in (8).

Recall that $\mathcal{A} \cong \tilde{\mathcal{V}}_U^a$ and let $\beta : \mathcal{A} \rightarrow M$ the composition of pr_1 with this isomorphism. The partial derivatives $\partial_{w_{k_0 \dots k_5}}, \partial_{x_{ij}}, \partial_{y_t}, \partial_z$ for $\sum k_\ell = 3, i \in \{0, 1, 2\}, j \in \{3, 4, 5\}$ and $t \in \{1, 2\}$ generate the tangent space $T_{M \times U \times \mathbf{C}}$, and an explicit computation shows that a vector

$$(\{W_{k_0 \dots k_5}\}, \{X_{ij}\}, \{Y_t\}, Z) \in T_{M \times U \times \mathbf{C}}$$

is in the subspace $T_{\mathcal{A}|_{pr_2^{-1}(\ell_0, \Pi_0)}}$ at the point (G_b, c) if and only if

$$(16) \quad \sum_{j=3,4,5} \frac{\partial G_b}{\partial x_j}(s_0, s_1, s_2, 0, 0, 0)(s_0 X_{0j} + s_1 X_{1j} + s_2 X_{2j}) \\ + \sum_{k_0+k_1+k_2=3} W_{k_0 k_1 k_2 000} s_0^{k_0} s_1^{k_1} s_2^{k_2} \\ = s_2^2(3c(Y_1 s_0 + Y_2 s_1) - Z s_2).$$

The map $d\beta$ sends $(\{W_{k_0 \dots k_5}\}, \{X_{ij}\}, \{Y_t\}, Z) \mapsto (\{W_{k_0 \dots k_5}\}) \in T_{G_b} M$.

As Equation (16) imposes no conditions on the $W_{k_0 \dots k_5}$ when $k_0 + k_1 + k_2 \leq 2$, the map $d\beta$ is then onto at the point $(G_b, (\ell_0, \Pi_0), c)$ if and only if for any tuple $(\{W_{k_0 k_1 k_2 000}\} : k_0 + k_1 + k_2 = 3)$, there exists $(\{X_{ij}\}, \{Y_t\}, Z)$ such that Equation (16) is satisfied. This translates into the surjectivity of the following linear map

$$\phi : \left(\bigoplus_{j=3,4,5} (\mathbf{C}^3)^* \right) \bigoplus (\mathbf{C}^2)^* \bigoplus \mathbf{C}^* \rightarrow \mathbf{H}^0(\mathbb{P}^2, \mathcal{O}(3))$$

defined by

$$(l_3, l_4, l_5, l, l') \xrightarrow{\phi} \sum_{j=3,4,5} \frac{\partial G_b}{\partial x_j}(s_0, s_1, s_2, 0, 0, 0) l_j + s_2^2(c l + l'),$$

with coordinates s_0, s_1, s_2 for $(\mathbf{C}^3)^*$, s_0, s_1 for $(\mathbf{C}^2)^*$ and s_2 for \mathbf{C}^* .

Choosing the natural basis for the 12-dimensional space on the left and 10-dimensional space on the right respectively, an easy but lengthy computation which we omit allows one to write down the 10×12 matrix corresponding to ϕ explicitly. We note that out of the 46 coefficients involved in defining a G_b which has the pair (ℓ_0, Π_0) as a triple line (i.e., of the form (15)), only 19 of them arise in the matrix corresponding to the map ϕ , i.e., ϕ is a 10×12 matrix M with 19 variables involved in its entries. Namely for A_{ijk} six 1×3

vectors of coefficients of G_b (cf. [BVdV79, p.100]) and c , the matrix has the form

$$(17) \quad M = \left(\begin{array}{cccccc|cccc} A_{200} & & & & & & s_0^3 & & & & & \\ A_{110} & A_{200} & & & & & s_0^2 s_1 & & & & & \\ A_{101} & & A_{200} & & & & s_0^2 s_2 & & & & & \\ A_{020} & A_{110} & & & & & s_0 s_1^2 & & & & & \\ A_{011} & A_{101} & A_{110} & & & & s_0 s_1 s_2 & & & & & \\ A_{002} & & A_{101} & c & & & s_0 s_2^2 & & & & & \\ & & A_{020} & & & & s_1^3 & & & & & \\ & & A_{011} & A_{020} & & & s_1^2 s_2 & & & & & \\ & & A_{002} & A_{011} & c & & s_1 s_2^2 & & & & & \\ & & & A_{002} & & 1 & s_2^3 & & & & & \end{array} \right)$$

where the final column corresponds to the basis of $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(3))$.

As being of maximal rank is invariant under multiplying by a scalar, to show that the locus where ϕ is not surjective is in codimension at least 2, it suffices to construct a $\mathbb{P}^1 \subset \mathbb{P}H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(3))$, over every point of which ϕ is surjective - as if it were in codimension ≤ 1 , then this \mathbb{P}^1 would necessarily meet this locus of non-surjectivity. In other words, to conclude we need to choose two linearly independent vectors in \mathbf{C}^{19} , i.e., entries for the A_{ijk} and c above, so that the plane they span corresponds to a 2-dimensional subspace of the space of coefficients of cubic polynomials of the form (15), at all of which the matrix defined by ϕ has rank 10. To this aim, we let κ, λ be coordinates of such a \mathbb{P}^1 and consider the following matrix (chosen essentially arbitrarily to be in the form (17))

$$M(\kappa, \lambda) = \left(\begin{array}{cccccccccccc} \kappa & 0 & \lambda & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \kappa & \kappa & 0 & \lambda & 0 & 0 & 0 & 0 & 0 & 0 \\ \lambda & \lambda & \kappa & 0 & 0 & 0 & \kappa & 0 & \lambda & 0 & 0 & 0 \\ 0 & \lambda & \kappa & 0 & 0 & \kappa & 0 & 0 & 0 & 0 & 0 & 0 \\ \lambda & \kappa & 0 & \lambda & \lambda & \kappa & 0 & 0 & \kappa & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \lambda & \lambda & \kappa & \kappa & 0 & 0 \\ 0 & 0 & 0 & 0 & \lambda & \kappa & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \lambda & \kappa & 0 & 0 & \lambda & \kappa & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \lambda & \kappa & 0 & 0 & \kappa & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

It is now not hard to check by a Gröbner basis computation using Macaulay2 that the 10×10 minors of the above matrix have no common solution for $(\kappa, \lambda) \neq (0, 0)$ which concludes the proof. \square

Remark 4.5. In fact the proof of Theorem 4.4 gives that the locus of triple lines in any cubic fourfold X (i.e., not just smooth) is connected. If X is general, then V is irreducible even.

We recall that \mathcal{U}_k denotes the universal subbundle on $G(k, 6)$.

Lemma 4.6. *The normal bundle of the inclusion $\tilde{V} \subset \tilde{F}$ is given by*

$$N_{\tilde{V}/\tilde{F}} \cong (\mathcal{U}_2^\vee \boxtimes \bigwedge^3 \mathcal{U}_3 \boxtimes \bigwedge^2 \mathcal{U}_2^\vee)|_{\tilde{V}}.$$

In particular, for $H = \pi^* H_F$ the pullback of the Plücker line bundle to \tilde{F}

$$\begin{aligned} \bigwedge^2 N_{\tilde{V}/\tilde{F}} &= (-3H + 2E_S)_{\tilde{V}}, \\ K_{\tilde{V}} &= 3(E_S - H)_{\tilde{V}}. \end{aligned}$$

Proof. Recalling the definitions of \mathcal{G} , $\tilde{\mathcal{F}}$, $\tilde{\mathcal{V}}$ from (10), (12), we note that \mathcal{G}_Δ is isomorphic to the locus $\mathcal{D} = \mathbf{G}(2, \mathcal{U}_3) = \{(\ell, \Pi) : \ell \subset \Pi\} \subset \mathbf{G}(2, 6) \times \mathbf{G}(3, 6)$, so the inclusion $\mathcal{G}_\Delta \hookrightarrow \mathcal{G} \subset \mathbf{G}(2, 6) \times \mathbf{G}(2, 6) \times \mathbf{G}(3, 6)$ is a section δ to the projection $\pi_{2,3} : \mathcal{G} \rightarrow \mathcal{D}$. Let $\gamma : \mathcal{D} \rightarrow \mathbf{G}(2, 6)$ and $\beta : \mathcal{D} \rightarrow \mathbf{G}(3, 6)$ the projections. The variety \mathcal{G} is the Grassmann bundle $h : \mathbf{G}(2, \beta^* \mathcal{U}_3) \rightarrow \mathcal{D}$ and this section is determined (see [Ful98, B.5.7]) by a rank two subbundle of $\beta^* \mathcal{U}_3$ on \mathcal{D} . We claim now that this subbundle is $\gamma^* \mathcal{U}_2$. Note that $\gamma^* \mathcal{U}_2$ is the universal bundle over $\mathbf{G}(2, \mathcal{U}_3)$ so that we have a sequence:

$$(18) \quad 0 \rightarrow \gamma^* \mathcal{U}_2 \rightarrow \beta^* \mathcal{U}_3 \rightarrow Q \rightarrow 0,$$

with Q a line bundle. Indeed if $\mathcal{U}_\mathcal{G}$ the universal subbundle on \mathcal{G} , then at the level of fibres, we have

$$\delta^* \mathcal{U}_\mathcal{G}|_{(\ell, \Pi)} = \gamma^* \mathcal{U}_2|_{(\ell, \Pi)} = \ell \subset \Pi = \beta^* \mathcal{U}_3|_{(\ell, \Pi)}.$$

This in particular implies that the normal bundle of the inclusion of the section is

$$N_{\mathcal{G}_\Delta/\mathcal{G}} \cong \gamma^* \mathcal{U}_2^\vee \otimes Q \cong \gamma^* \mathcal{U}_2^\vee \otimes \bigwedge^3 \beta^* \mathcal{U}_3 \otimes \bigwedge^2 \gamma^* \mathcal{U}_2^\vee.$$

and from [Sta21, Tag 0473] that $N_{\tilde{V}/\tilde{\mathcal{F}}} = \text{pr}_2^* N_{\mathcal{G}_\Delta/\mathcal{G}}$. Restricting to a fibre $\tilde{F} \subset \tilde{\mathcal{F}}$ over a general smooth $X \in |\mathcal{O}(3)|$, we have a surjection $N_{\tilde{V}/\tilde{\mathcal{F}}}^\vee|_{\tilde{V}} \rightarrow N_{\tilde{V}/\tilde{F}}^\vee$, and as both are locally free of rank two (the latter from Lemma 4.3 and Theorem 4.4), they must be isomorphic so we compute $N_{\tilde{V}/\tilde{F}}$ as in the statement of the lemma. As the pullback (under the isomorphism of Lemma 4.1) of $\det \mathcal{U}_3$ to \tilde{F} is $-3H + E_S$ (see Lemma 4.8) and the pullback of $\det \mathcal{U}_2^\vee$ is H , we obtain the second equality, whereas the third follows from the first and the fact that $K_{\tilde{F}} = E_S$ as \tilde{F} is the blowup of F at S . \square

Theorem 4.7. *The class of V in $\text{CH}^2(F)$ is given by*

$$[V] = 21c_2(\mathcal{U}_F).$$

Proof. Keeping the notation established in the proof of the above Lemma 4.6, consider

$$\mathbb{P}(\mathcal{U}_3^\vee) \cong \mathbf{G}(2, \mathcal{U}_3) = \mathcal{D} = \{(\ell, \Pi), \ell \subset \Pi\} \subset \mathbf{G}(2, 6) \times \mathbf{G}(3, 6).$$

We have the fibre diagram

$$\begin{array}{ccc} \mathcal{G} & \xrightarrow{\alpha_1} & \mathbb{P}(\mathcal{U}_3^\vee) \\ \downarrow \alpha_2 & & \downarrow \beta \\ \mathbb{P}(\mathcal{U}_3^\vee) & \xrightarrow{\beta} & \mathbf{G}(3, 6). \end{array}$$

Let $\alpha = \alpha_i \beta$ and $\delta : \mathbb{P}(\mathcal{U}_3^\vee) \rightarrow \mathcal{G}$ the diagonal embedding with image \mathcal{G}_Δ . We have $\alpha_i \delta = 1$ and hence $\delta^* \alpha_i^* = 1 = \alpha_{i*} \delta_*$. Let $h = \mathcal{O}_{\mathbb{P}(\mathcal{U}_3^\vee)}(1)$ the tautological bundle and $h_i = \alpha_i^* h$ on \mathcal{G} . The intersection ring of \mathcal{G} is generated

by $\langle h_1, h_2, \alpha^* A(G(3, 6)) \rangle$. Aiming to compute the class of \mathcal{G}_Δ in this ring, we may write

$$[\mathcal{G}_\Delta] = s_{11}h_1^2 + s_{12}h_1h_2 + s_{22}h_2^2 + h_1\alpha^*c_1 + h_2\alpha^*c_2 + \alpha^*c$$

for some s_{ij} integers, c_1, c_2 multiples of the Schubert cycle $\sigma_1(3)$ on $G(3, 6)$ and c a linear combination of $\sigma_2(3), \sigma_{1,1}(3)$ which we will now determine. Note that we have the relations

$$\begin{aligned} \alpha_{1*}\alpha_2^* &= \beta^*\beta_* = \alpha_{2*}\alpha_1^*, \\ \alpha_i^*a \cdot [\mathcal{G}_\Delta] &= \delta_*a, \\ (\alpha_1^*a \cdot \alpha_2^*a') \cdot [\mathcal{G}_\Delta] &= \delta_*(aa'), \\ \beta_*(h^{i+2}) &= s_i(\mathcal{U}_3^\vee). \end{aligned}$$

In particular we have $\beta_*(h^2) = 1$, $\beta_*(h^3) = s_1(\mathcal{U}_3^\vee) = c_1(\mathcal{U}_3)$, $\beta_*(h^4) = s_2(\mathcal{U}_3^\vee) = c_1^2(\mathcal{U}_3) - c_2(\mathcal{U}_3)$. We now intersect $[\mathcal{G}_\Delta]$ with h_1 and use the second relation above to obtain $s_{22} = 1$, by pushing forward by α_1 , whereas the same analogous intersection with h_2 gives $s_{12} = 1$ and $c_1 = -c_1(\mathcal{U}_3)$. By symmetry we also obtain $s_{11} = 1$ and $c_2 = -c_1(\mathcal{U}_3)$. Intersecting instead with h_1^2 and pushing forward via α_2 we get $c = c_2(\mathcal{U}_3)$, giving overall

$$(19) \quad [\mathcal{G}_\Delta] = h_1^2 + h_1h_2 + h_2^2 - h_1\alpha^*c_1(\mathcal{U}_3) - h_2\alpha^*c_1(\mathcal{U}_3) + \alpha^*c_2(\mathcal{U}_3).$$

Now [Ful98, Example 14.6.5] gives that if $\phi : \mathbb{P}(\mathcal{U}_3^\vee) = G(1, \mathcal{U}_3^\vee) \rightarrow G(2, \mathcal{U}_3)$ the natural isomorphism, then $\phi^*(Q^{-1}) = \mathcal{O}_{\mathbb{P}(\mathcal{U}_3^\vee)}(-1)$, with Q as in Sequence (18), so that the class of $\phi^*(Q)$ is h . Sequence (18) then gives the following two formulas.

$$(20) \quad h = \beta^*c_1(\mathcal{U}_3) - \gamma^*c_1(\mathcal{U}_2),$$

$$(21) \quad \beta^*c_2(\mathcal{U}_3) = h\gamma^*c_1(\mathcal{U}_2) + \gamma^*c_2(\mathcal{U}_2).$$

For $\tilde{F} = \text{Bl}_S(F)$ and Γ_ϕ the graph of the Voisin map, we have observed already that we have an isomorphism $\pi' : \Gamma_\phi \rightarrow \tilde{F}$ sending (ℓ, ℓ', Π) to (ℓ, Π) (cf. Lemma 4.1). We now consider the one-to-one morphism $\iota_\phi : \Gamma_\phi \rightarrow \mathcal{G}$ and let $\tilde{\iota} : \tilde{F} \hookrightarrow \mathbb{P}(\mathcal{U}_3^\vee)$ the identity inclusion $\tilde{\iota}((\ell, \Pi)) = (\ell, \Pi)$. Then $\alpha_1\iota_\phi$ factors through the isomorphism $\pi' : \Gamma_\phi \rightarrow \tilde{F}$. All in all, we have a diagram

$$\begin{array}{ccccc} \mathcal{G} & \xleftarrow{\iota_\phi} & \Gamma_\phi & \xrightarrow{\iota_\phi} & \mathcal{G} \\ \alpha_2 \downarrow & & \downarrow \pi' & & \downarrow \alpha_1 \\ G(3, 6) & \xleftarrow{\beta} & \mathbb{P}(\mathcal{U}_3^\vee) & \xrightarrow{\tilde{\iota}} & \mathbb{P}(\mathcal{U}_3^\vee) & \xrightarrow{\beta} & G(3, 6) \\ \gamma \downarrow & & \downarrow \pi & & \downarrow \gamma \\ G(2, 6) & & F & \xrightarrow{\iota} & G(2, 6) \end{array}$$

The map $\tilde{\phi} = \gamma\alpha_2\iota_\phi\pi'^{-1} : \tilde{F} \rightarrow G(2, 6)$ is the Voisin map $\phi : \tilde{F} \rightarrow F$ combined with the embedding of F into $G(2, 6)$.

For the duration of this proof, denote for simplicity the class of E_S by E and the following cycles $H = \pi^*H_F$ and $c_2 = \pi^*c_2(\mathcal{U}_F)$ on \tilde{F} . We have that $c_1(\pi^*\iota^*\mathcal{U}_2) = -H$.

Lemma 4.8. ([Huy20, Remark 2.2.19], [Ame11, Proposition 6]) *On \tilde{F} we have*

$$\begin{aligned}\tilde{\iota}^* \beta^* c_1(\mathcal{U}_3) &= -3H + E, \\ \tilde{\phi}^* c_1(\mathcal{U}_2) &= -7H + 3E.\end{aligned}$$

Relation (20) combined with the above lemma gives on $\tilde{F} \cong \Gamma_\phi$ that

$$\begin{aligned}\pi'_* \iota_\phi^* h_1 &= -(2H - E), \\ \pi'_* \iota_\phi^* h_2 &= 2(2H - E).\end{aligned}$$

Indeed, for the first recall that $h_i = \alpha_i^* h$ and observe that $\iota_\phi^* \alpha_1^* = \pi'^* \tilde{\iota}^*$ and for the second, observe that $\beta \alpha_2 = \alpha = \beta \alpha_1$ and recall the above mentioned formula for $\tilde{\phi}$.

Observe also that (20) and (21) give that $\beta^* c_2(\mathcal{U}_3) = \beta^* c_1(\mathcal{U}_3) \gamma^* c_1(\mathcal{U}_2) - \gamma^* c_1^2(\mathcal{U}_2) + \gamma^* c_2(\mathcal{U}_2)$ and combining with the above lemma we get

$$\pi'_* \iota_\phi^* \alpha^* c_2(\mathcal{U}_3) = (3H - E)H - H^2 + c_2.$$

Relation (19) leads now to the computation of the class of \tilde{V} as follows

$$\begin{aligned}(22) \quad [\tilde{V}] &= \pi'_* \iota_\phi^*(\mathcal{G}_\Delta) \\ &= 20H^2 - 18HE + 4E^2 + c_2,\end{aligned}$$

from which we may now conclude, by pushing forward via π and using Theorem 2.3

$$\begin{aligned}[V] &= \pi_*[\tilde{V}] \\ &= 20H_F^2 - 4[S] + c_2(\mathcal{U}_F) \\ &= 21c_2(\mathcal{U}_F).\end{aligned} \quad \square$$

To summarise the above, we conclude the following.

Corollary 4.9. *The locus $V \subset F$ of triple lines is an irreducible projective surface of class $21c_2(\mathcal{U}_F)$.*

In order to analyse singularities of V we will now need to study the intersections

$$\begin{aligned}C &= S \cap V \subset F, \\ \tilde{C} &= E_S \cap \tilde{V} \subset \tilde{F}.\end{aligned}$$

Corollary 4.10. *We have $C \neq \emptyset$.*

Proof. Theorems 4.7, 2.3 and Lemma 2.6 give

$$[S][V] = 5(H_F^2 - c_2(\mathcal{U}_F^\vee)) 21c_2(\mathcal{U}_F) = 1890,$$

so that in particular $S \cap V$ cannot be empty. \square

Proposition 4.11. *The variety \tilde{V} intersects E_S transversely at a smooth pure 1-dimensional variety \tilde{C} .*

Proof. Denote by $\mathcal{E} \subset \tilde{\mathcal{F}}$ the inverse image of the universal type two locus $\mathcal{S} \subset |\mathcal{O}_{\mathbb{P}^5}(3)| \times \mathbf{G}(2,6)$ defined at the beginning of this section. The fibre $\mathrm{pr}_1^{-1}(X)$ restricted to \mathcal{E} is E_{S_X} the exceptional divisor of the blow up, whose fibres over the type two locus S_X of X parametrise 2-planes tangent along a line in S_X . We show now that all fibres of $\mathrm{pr}_2 : \mathcal{E} \rightarrow \mathcal{G}$ are isomorphic. Indeed, a projective transformation takes a second type line of a smooth cubic to a second type line since for a smooth cubic $X = \mathbf{V}(G)$, ℓ being of second type is equivalent to

$$\langle \partial_i G|_{\ell} \rangle_{i=0,\dots,5} \subset \mathbf{H}^0(\ell, \mathcal{O}(2))$$

being 2-dimensional (instead of 3-dimensional which is the generic case). Keeping the notation established in the beginning of this sections and restricting at the fibre over $(x=0, y=0, y')$, i.e. $\Pi = \Pi_0$, $\ell = \ell_0$ as in (8), then using the equation of G from (15), we see that

$$\begin{aligned} \langle \partial_i G|_{\ell_0} \rangle_{i=0,\dots,5} &= \langle 0, 0, 0, Q_3(x_0, x_1, 0, 0), \\ &\quad Q_4(x_0, x_1, 0, 0, 0), Q_5(x_0, x_1, 0, 0, 0) \rangle. \end{aligned}$$

Since three quadrics in two variables are linearly dependent if and only if the determinant of the corresponding 3×3 matrix vanishes, we see that this imposes one extra condition. We also conclude that all the fibres of the map $\mathrm{pr}_2 : \mathcal{E} \rightarrow \mathcal{G}$ are isomorphic to a determinant cubic hypersurface (inside the fibre \mathbb{P}^{46} of the second projection of $\tilde{\mathcal{F}}$), which is smooth outside the codimension 4 locus of the generic determinantal variety $M_1(3,3)$. This singular locus maps under the first projection to the locus of singular cubics since in this case the cubic is singular at a point of the line. The degeneracy locus $M_1(3,3)$ can be constructed universally and as it maps into the locus of singular cubics in $|\mathcal{O}_{\mathbb{P}^5}(3)|$ (cf. [Huy20, Proposition 2.2.13]), we obtain also that $\mathcal{E}_{\mathrm{sm}} = \mathrm{pr}_1^{-1}(|\mathcal{O}_{\mathbb{P}^5}(3)|_{\mathrm{sm}}) \subset \mathcal{E}$ is smooth.

The intersection $\tilde{\mathcal{V}} \cap \mathcal{E}$ is simply the restriction of \mathcal{E} to the locus \mathcal{G}_{Δ} under pr_2 . Restricting to the locus of smooth cubics, we have a diagram

$$\begin{array}{ccc} \tilde{\mathcal{V}} \cap \mathcal{E}_{\mathrm{sm}} & \xrightarrow{\mathrm{pr}_1} & |\mathcal{O}_{\mathbb{P}^5}(3)|_{\mathrm{sm}} \\ \downarrow \mathrm{pr}_2 & & \\ \mathcal{G}_{\Delta} & & \end{array}$$

where any fibre over \mathcal{G}_{Δ} is the fixed determinantal cubic hypersurface in \mathbb{P}^{46} from above. The restricted map pr_1 is onto from Lemma 4.10. In particular, as the base \mathcal{G}_{Δ} is smooth and the intersection above is just the restriction to \mathcal{G}_{Δ} , so also the total space $\tilde{\mathcal{V}} \cap \mathcal{E}_{\mathrm{sm}}$ is smooth. By generic smoothness, also $\mathrm{pr}_1^{-1}(\mathbf{V}(G))$ is smooth for generic $\mathbf{V}(G) \in |\mathcal{O}_{\mathbb{P}^5}(3)|_{\mathrm{sm}}$ and must have pure 1-dimensional fibres. This implies that $\tilde{\mathcal{V}}$ meets E_S transversely for a fixed general cubic $\mathbf{V}(G)$. \square

We note that the above proof does not conclude that \tilde{C} is irreducible, and as a consequence, we also do not conclude the irreducibility of C . Even though we expect this to be true, it will not be necessary for what follows so we do not pursue it.

We consider now the 12-dimensional family

$$(23) \quad \mathcal{H} = \{(\ell, H) \text{ with } \ell \subset H\} \subset \mathbf{G}(2, 6) \times \mathbf{G}(4, 6),$$

which is a $\mathbf{G}(2, 4)$ -fibration over $\mathbf{G}(2, 6)$. Since any two such pairs differ by a projective transformation, the family of smooth cubic fourfolds which contain ℓ as a line of second type with corresponding 3-plane H are all isomorphic as we vary ℓ and H . We may then assume ℓ_0, H_0 are as in (8). The smooth cubics in \mathbb{P}^5 which contain ℓ_0 as a line of second type and whose corresponding tangent 3-plane is H_0 form an open set inside the family of cubics given by the following equation

$$(24) \quad F = (c_0x_4 + d_0x_5)x_0^2 + (c_{01}x_4 + d_{01}x_5)x_0x_1 + (c_1x_4 + d_1x_5)x_1^2 + x_0Q_0(x_2, x_3, x_4, x_5) + x_1Q_1(x_2, x_3, x_4, x_5) + P(x_2, x_3, x_4, x_5),$$

with Q_0, Q_1 quadrics and P a cubic (cf. Equation (6) where one member of the above family was chosen). The above family has dimension 45 and thus the variety

$$\{(\mathbf{V}(G), (\ell, H)) : \ell \subset \mathbf{V}(G) \text{ has tangent 3-plane } H\} \subset |\mathcal{O}_{\mathbb{P}^5}(3)|_{\text{sm}} \times \mathcal{H}$$

is irreducible of dimension 57 and smooth - note here that this variety is isomorphic to \mathcal{S} introduced above since every type two line has a unique triple tangent hyperplane.

The pencil of 2-planes $\Pi_{\lambda, \mu}$ inside $H_0 \cong \mathbb{P}^3$ (with coordinates x_0, x_1, x_2, x_3) which contain ℓ_0 is given by $\mu x_2 - \lambda x_3 = 0$. The cubic surface $\mathbf{V}(G) \cap H_0$ is given by the equation

$$x_0Q_0(x_2, x_3) + x_1Q_1(x_2, x_3) + P(x_2, x_3) = 0,$$

where $Q_0(x_2, x_3) = Q_0(x_2, x_3, 0, 0)$ etc. The intersection of $\Pi_{\lambda, \mu}$ with $\mathbf{V}(G)$ is then given by the equation (without loss of generality we assume that $\lambda \neq 0$ in the following)

$$\begin{aligned} x_0x_2^2Q_0(\lambda, \mu) + x_1x_2^2Q_1(\lambda, \mu) + x_2^3\lambda^{-1}P(\lambda, \mu) \\ = x_2^2[x_0Q_0(\lambda, \mu) + x_1Q_1(\lambda, \mu) + x_2\lambda^{-1}P(\lambda, \mu)] = 0, \end{aligned}$$

and the solution $x_2^2 = 0$ is the double line ℓ_0 whereas $x_0Q_0(\lambda, \mu) + x_1Q_1(\lambda, \mu) + x_2\lambda^{-1}P(\lambda, \mu) = 0$ is the residual line $\ell_{\lambda, \mu}$ (the equations take place in the plane $\Pi_{\lambda, \mu}$). Recall that when $X = \mathbf{V}(G)$ is general then it does not contain a 2-plane. Hence, in this case, for any $[\lambda, \mu]$ the last equation is not the null equation, otherwise $\Pi_{\lambda, \mu} \subset X$. The intersection point of the residual line with ℓ_0 is given by

$$(25) \quad x_0Q_0(\lambda, \mu) + x_1Q_1(\lambda, \mu) = 0.$$

Finally, Q_0, Q_1 have a common root $[\lambda_0, \mu_0]$ if and only if there exists a 2-plane, namely Π_{λ_0, μ_0} , which intersects X at x_2^3 , in which case $\ell_0 \in S \cap V$. By a dimension count, none of them is identically zero and they cannot have a common double root as X is general.

We note the following corollary of the above discussion, which will be used in the next section.

Corollary 4.12. *For $s \in S \setminus (S \cap V)$, the restriction of the map*

$$\psi_{E_S} : E_S \dashrightarrow W$$

to $E_{S,s} \cong \mathbb{P}^1$ is a two-to-one morphism.

Proof. The map ψ_{E_S} is not defined on the fibres of E_S over $V \cap S$ and from the discussion above this corresponds to the case in which Q_0, Q_1 have a common root. If this is not the case, then given a point $[x_0 : x_1] \in \ell_0$ there are two pairs $[\lambda, \mu]$ (counted with multiplicity) which satisfy Equation (25) which proves the claim. \square

Viewed geometrically, given $p \in \ell$, $[\ell] \in S$ general, the line ℓ corresponds to the node of C_p , the nodal genus four curve which parametrises lines through $p \in \ell$ (see Lemma 3.1). As a generic nodal genus four curve possesses two g_3^1 's and the fibre through the node of each one of these defines a conjugate point, the above proposition (and Proposition 3.10) give that we have two conjugate lines intersecting ℓ at the point p .

Lemma 4.13. *The smooth irreducible surface \tilde{V} does not contain any fibres of $E_S \rightarrow S$, and thus C is a pure 1-dimensional variety whose normalisation is \tilde{C} . The locus of $s \in S$ over which $E_{S,s}$ meets \tilde{V} in two points is a finite subset of $C = S \cap V$.*

Proof. From Proposition 4.11, $\tilde{C} = E_S \cap \tilde{V}$ is smooth 1-dimensional. Note now that the above analysis gives further information on \tilde{V} . First, $\tilde{V} \subset \tilde{F}$ cannot contain an entire fibre of $E_S \rightarrow S$ since then $Q_0(\lambda, \mu), Q_1(\lambda, \mu)$ of Equation (25) would both be identically zero, which cannot happen for a general fourfold. In particular, Q_0 and Q_1 can have at most two common roots, and these correspond to the number of intersection points of \tilde{V} with a fibre of E_S . In other words $C = S \cap V = \pi(\tilde{C})$ is pure 1-dimensional and $\pi : \tilde{C} \rightarrow C$ is the normalisation morphism. As from Corollary 4.10 $V \cap S \neq \emptyset$ for a general cubic, having one common root imposes one condition in the fibre S of \mathcal{S} over $X = \mathbf{V}(G) \in |\mathcal{O}_{\mathbb{P}^5}(3)|$, whereas having two common roots is a codimension two condition in S , i.e., a finite number of points on C . \square

Lemma 4.14. *The class of C is given by*

$$[C] = \frac{35}{2} H_F^3 \in H_2(F, \mathbf{Q})$$

(or $6H_S \in H^2(S, \mathbf{Q})$). We have $p_a(\tilde{C}) = 4726$, $p_a(C) = 8506$.

Proof. From (22), we have

$$[\tilde{C}] = [\tilde{V}] \cdot E_S = 20H^2 E_S - 18H E_S^2 + 4E_S^3 + E_S c_2.$$

To compute the terms, we use the standard diagram

$$\begin{array}{ccc} E_S = \mathbb{P}(N_{S/F}) & \xhookrightarrow{j} & \tilde{F} \\ \downarrow \pi' & & \downarrow \pi \\ S & \xhookrightarrow{i} & F. \end{array}$$

Let ξ be the tautological bundle of $E_S = \mathbb{P}(N_{S/F})$. Then $j^* E_S = \xi$, which implies that $\pi_* E_S^k = i_* \pi'_* \xi^{k-1}$, and let $\xi^2 - c_1(\pi'^* N_{S/F}) \xi + c_2(\pi'^* N_{S/F}) = 0$

the standard relation. We then have $\pi_*(H^2E_S) = \pi_*(c_2E_S) = 0$, $\pi_*HE_S^2 = -H_S$ and $\pi_*E_S^3 = -c_1(N_{S/F})$. As $K_F = 0$, we obtain $c_1(N_{S/F}) = K_S = 3H_S$ from Theorem 2.3. Then $[C] = \pi_*[\tilde{C}] = 6H_S$. Theorem 2.3 and Lemma 2.6 implies the result.

To compute the genus of $\tilde{C} \subset \tilde{V}$ we use the adjunction formula and the computations of Lemma 4.6 as follows. From the above computation of $[\tilde{C}]$ we see that

$$\tilde{C}(\tilde{C} + K_{\tilde{V}}) = 16E_S^4 - 84HE_S^3 + 134H^2E_S^2 + 4E_S^2c_2.$$

As above, by using Theorem 2.3 and Lemma 2.6 we get $E_S^4 = -c_1^2(N_{S/F}) + c_2(N_{S/F}) = -1710$, $HE_S^3 = -c_1(N_{S/F})H_S = -945$, $H^2E_S^2 = -[S]H_F^2 = -315$, and $E_S^2c_2 = -[S]c_2(\mathcal{U}_F) = -90$ which compute the above expression to be 9450, implying the first genus computation. Similarly, computing now via adjunction for $C \subset S$ we have from the above that $C(C + K_S) = 54H_S^2 = 17010$ so that $p_a(C) = 8506$. \square

In the following, we say that a point x in a variety X has *normal crossing singularities* if

$$\hat{\mathcal{O}}_{X,x} \cong \mathbf{C}[[x_1, \dots, x_n]]/(x_1 \cdots x_r).$$

Proposition 4.15. *The universal intersection curve $\mathcal{C} = \mathcal{S} \cap \mathcal{V} \subset \mathcal{F}$ is irreducible and there is an open subset $U \subset |\mathcal{O}(3)|$ so that $\mathcal{C}|_U$, the restriction to the inverse image of U , has normal crossings singularities and its normalisation $\mathcal{C}|_U^\nu \rightarrow \mathcal{C}|_U$ identifies pairs of points of a divisor.*

Proof. From the analysis around Equation (23), the second projection of the universal type two locus $\mathcal{S} \subset |\mathcal{O}(3)|_{\text{sm}} \times \mathcal{H} \rightarrow \mathcal{H}$ is a \mathbb{P}^{45} -fibration over the locus \mathcal{H} of pairs (ℓ, H) where $\ell \subset H \cong \mathbb{P}^3 \subset \mathbb{P}^5$ (see (23)), and the line ℓ will be a type two triple line (i.e., in $S \cap V$) if and only if for G of type (24), the plane quadrics Q_0, Q_1 have a common root (see Lemma 4.13). In particular the fibre of \mathcal{C} over \mathcal{H} in \mathbb{P}^{45} - i.e., the locus parametrising cubics G for which a fixed pair $(\ell_0, H_0) \in \mathcal{H}$ is a triple type two line - is a cone over the corresponding resultant R in 6 variables, namely the coefficients of the Q_i . Note also that the induced PGL_5 action on points of \mathcal{H} is transitive, as the property of being a triple type two line is preserved under this action.

Denoting these coefficients x_1, x_2, x_3 (resp. y_i) of Q_1 (resp. Q_2), the resultant is given by

$$R = \mathbf{V}((x_1y_3 - x_3y_1)^2 - (x_1y_2 - x_2y_1)(x_2y_3 - x_3y_2)).$$

Since R is irreducible, the universal locus \mathcal{C} must also be irreducible.

Observe that the family of vertices of the cones over the resultant in each fibre \mathbb{P}^{45} are in high codimension when mapped to $|\mathcal{O}(3)|$, so as our cubics are assumed general, we may ignore this locus, and we denote by $\hat{R} \subset \mathbb{P}^{45}$ its complement in the cone, which is of dimension 44.

Lemma 4.16. *The resultant R generically has normal crossing singularities, resolved by a degree 2 normalisation.*

Proof. The normalisation R^ν of the resultant R is given by the vanishing locus of the polynomials

$$\begin{aligned} wx_3 - w'y_3, \\ wx_2 - w'y_2 + x_3y_1 - x_1y_3, \\ wx_1 - w'y_1 + x_2y_1 - x_1y_2, \\ w'^2 - w'x_2 + x_1x_3, \\ ww' - w'y_2 + x_3y_1, \\ w^2 - wy_2 + y_1y_3 \end{aligned}$$

in $\mathbf{C}[w, w', x_1, \dots, y_3]$. The singularities of the normalisation are in codimension 4 so we may ignore them as we are working generically. The singular locus of the resultant is the smooth rank ≤ 1 determinantal locus given by the matrix

$$\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}.$$

It consists of points of the form

$$[x_1 : x_2 : x_3 : tx_1 : tx_2 : tx_3] \text{ or } [ty_1 : ty_2 : ty_3 : y_1 : y_2 : y_3].$$

and over these points the normalisation has, generically, two points, e.g., over the first type of point the two points are

$$[tw'_i : w'_i : x_1 : x_2 : x_3 : tx_1 : tx_2 : tx_3], \quad i = 1, 2,$$

where w'_i are the roots of the equation $w'^2 - w'x_2 + x_1x_3$. It has one point over the locus $x_2^2 = 4x_1x_3$ (where the quadric has a double root, i.e., $w' = x_2/2$).

A calculation shows that, after performing row transformations, the Jacobian matrix at the the point $[tw' : w' : x_1 : x_2 : x_3 : tx_1 : tx_2 : tx_3]$ is in all cases equivalent to the matrix (we omit zero rows)

$$J = \begin{pmatrix} 0 & 0 & tx_3^2 & -tx_3w' & tw'^2 & -x_3^2 & x_3w' & -w'^2 \\ 0 & 2w' - x_2 & x_3 & -w' & x_1 & 0 & 0 & 0 \\ 2w' - x_2 & 0 & 0 & 0 & 0 & x_3 & -w' & x_1 \end{pmatrix},$$

except when $(w', x_3) = (0, 0)$, in which case we get the matrix

$$J = \begin{pmatrix} 0 & 0 & -tx_2^2 & tx_1x_2 & -tx_1^2 & x_2^2 & -x_1x_2 & x_1^2 \\ 0 & -x_2 & 0 & 0 & x_1 & 0 & 0 & 0 \\ -x_2 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 \end{pmatrix}.$$

The above matrices have, outside the locus $x_2^2 = 4x_1x_3$, the first 3×2 minor of rank 2. Hence, outside the above locus, the differential of the map $p : R^\nu \rightarrow R$ (which is the restriction of the projection $p : \mathbb{P}^7 \dashrightarrow \mathbb{P}^5$ to the last six coordinates) at the above point is injective: the tangent space is the linear space of vectors v with $Jv = 0$. The kernel of the differential consists of those vectors with the last six coordinates zero, but in this case the first two coordinates are also zero.

Moreover, locally the images of the two sheets of the normalisation, where the points $[tw'_i : w'_i : x_1 : x_2 : x_3 : tx_1 : tx_2 : tx_3]$, $i = 1, 2$, belong, are smooth hypersurfaces in \mathbb{P}^5 . The corresponding perp vectors of those images at the

points $[x_1 : x_2 : x_3 : tx_1 : tx_2 : tx_3]$, $i = 1, 2$, when $(w', x_3) \neq (0, 0)$ are given by

$$\langle tx_3^2, -tx_3w', tw'^2, -x_3^2, x_3w', -w'^2 \rangle$$

with $w' = w'_i$ and by

$$\langle tx_2^2, tx_1x_2, -tx_1^2, x_2^2, -x_1x_2, x_1^2 \rangle$$

when $(w', x_3) = (0, 0)$. These are not parallel in all cases and hence we have two branches with normal crossing there. \square

As $\mathcal{C} \rightarrow \mathcal{H}$ is generically isotrivial, a standard argument with the Isom-scheme (see, e.g., [Bui94, p.3]) shows that there exists an open neighbourhood U , and a finite étale cover $U' \rightarrow U$ so that the pullback of \mathcal{C} (minus the vertices of the cones as above) is isomorphic to

$$U' \times \widehat{R}.$$

The transitivity of the PGL_5 -action on \mathcal{H} implies that any point in \mathcal{H} has a finite étale neighbourhood which trivialises \mathcal{C} . Clearly $U' \times \widehat{R}$ has normal crossing singularities outside a codimension 2 locus from Lemma 4.16. As being normal crossings can be checked étale locally, we deduce that \mathcal{C} has normal crossing singularities outside a codimension 2 closed subset too. By a dimension count, the non-normal-crossing locus cannot surject onto $|\mathcal{O}(3)|_{\mathrm{sm}}$, so after removing its image, we may assume there is an open in $|\mathcal{O}(3)|$ over which the universal family \mathcal{C} has normal crossings. \square

Theorem 4.17. *The curve C has 3780 nodes and no other singular points.*

Proof. From Corollary 4.14, as the difference between the genus of C and its normalisation \widetilde{C} is exactly 3780 (in particular positive), we obtain that C must be singular for a generic cubic fourfold. To conclude the result, it suffices thus to show that C is nodal, by which we mean it is a pure 1-dimensional, possibly disconnected, variety with at worst nodes as singularities.

To this aim, we note that Proposition 4.15 proves that \mathcal{C} has normal crossings over an open $U \subset |\mathcal{O}(3)|$. If we denote by $\mathcal{C}_{\mathrm{fs}}$ the locus of singular points in the fibres over U , then as pointed out above, $\mathcal{C}_{\mathrm{fs}} \rightarrow U$ is surjective. Let

$$\begin{array}{ccc} \mathcal{C}^\nu & \xrightarrow{\nu} & \mathcal{C} \\ & \searrow & \swarrow \\ & U & \end{array}$$

be the normalisation of \mathcal{C} . As a consequence of the proof of Lemma 4.16, we may shrink U further and assume that \mathcal{C}^ν is in fact smooth. The generic fibre of $\mathcal{C}^\nu \rightarrow U$ is also smooth, so it can be checked from Zariski's Main Theorem that after shrinking U further, $\nu_u : \mathcal{C}_u^\nu \rightarrow \mathcal{C}_u$ is the normalisation of every fibre over $u \in U$. Since $\nu^{-1}(\mathcal{C}_{\mathrm{fs}}) \rightarrow U$ has finite fibres, by purity of the branch locus we may shrink U further and assume that every fibre of \mathcal{C}^ν meets $\nu^{-1}(\mathcal{C}_{\mathrm{fs}})$ transversely (at an even number of points). The map $\nu_u : \mathcal{C}_u^\nu \rightarrow \mathcal{C}_u$ identifies in pairs the above points forming nodes in the image because of the above mentioned transversality condition. Hence, as C is

nodal and its arithmetic genus differs by 3780 from its normalisation, it must have precisely 3780 nodes. \square

Corollary 4.18. *The surface $V \subset F$ is irreducible and projective and has precisely 3780 isolated non-normal singularities, whose normalisation $\tilde{V} \rightarrow V$ is two-to-one over this non-normal locus.*

Proof. The first claims are from Corollary 4.9. Using notation around Equation (25) and Lemma 4.13, \tilde{V} meets a fibre $E_{S, [\ell]} \cong \mathbb{P}^1$ in either one or two distinct points, namely the number of common roots of Q_0, Q_1 (noting that the case of a double root is ruled out by generality). From Lemma 4.13 and Theorem 4.17, there are 3780 triple lines of second type $\ell \in S \cap V$ for which there are precisely two distinct 2-planes tangent to ℓ which have the same conjugate line, namely ℓ . A local computation shows that V cannot be singular along smooth points of C : over a smooth point of C , \tilde{V} meets the fibre of the exceptional divisor E_S at one point with multiplicity one and hence cannot have singular image. Hence the map $\tilde{V} \rightarrow V$ identifies these 3780 pairs of points giving as many non-normal isolated singularities as such pairs, and is an isomorphism otherwise. \square

Theorem 4.19. *If X is very general, every singular rational curve in F of primitive class β has precisely one node and there are 3780 such curves.*

Proof. From [Ame11, Proposition 6], the rational curves $\phi(E_{S, [\ell]})$ are of class β in $H_2(F, \mathbf{Z})$ and Theorem 2.4 proves that if X is very general, every rational curve in F of class β is of this form. Next, [NO21, Proposition 1.4, Lemma 3.8] proves that every curve in F of class β is as worst nodal. If a fibre $E_{S, [\ell]}$ of E_S over $[\ell]$ maps to a nodal curve under ϕ , then the preimages of the node are two distinct points of $E_{S, [\ell]}$. These correspond to two distinct planes, tangent along ℓ and having the same residual line ℓ' . But the only case in which this can happen is if $\ell = \ell'$ and these planes are triply tangent to the line $[\ell]$. This implies that $[\ell] \in C = S \cap V$ and the two planes correspond to the two distinct common roots of the quadrics Q_0, Q_1 of Equation (25). In this case, the image $\phi(E_{S, [\ell]})$ has exactly one node, at the point $[\ell]$. Alternatively, one could also argue that the images of these curves have at most one node as in [NO21, §1.4], which implies that every curve of class β is either a smooth rational curve or a plane cubic, hence is of arithmetic genus 0 or 1-nodal of arithmetic genus 1.

In other words, singular curves in F of class β are all 1-nodal and rational, and are images of fibres of E_S over the points of $C = S \cap V$ where the Q_0, Q_1 have two distinct common roots. As discussed above, these correspond exactly to the nodes of C and so Theorem 4.17 gives that there are 3780 of them. \square

To make the geometry of the above count more explicit: each nodal rational curve of class $\beta \in H^2(F, \mathbf{Z})$ corresponds to a singularity of the curve $C = S \cap V \subset F$, in the sense that it is the image of $\mathbb{P}^1 \cong E_s \subset \tilde{F}$ for $s \in C_{\text{sing}}$ under the Voisin map $\phi : \tilde{F} \rightarrow F$. In turn, each such singular point of C corresponds to a triple type two line which has two distinct triple tangent 2-planes and these can be counted intersection-theoretically.

Remark 4.20. For general X , [NO21, Lemmas 3.7-3.9] prove that for $\Sigma \subset \text{CH}_\beta(F)$ the locus of irreducible curves of arithmetic genus 1, every singular member is 1-nodal and that the morphism $\sigma : \Sigma \rightarrow \mathbb{P}^1$ taking a curve to its j -invariant is of degree 3780. In particular, the fibre over the boundary point at infinity corresponds to the number of 1-nodal rational curves in F of class β , which as discussed in the proof above is the image of a line $E_{S, [\ell]}$ for $[\ell] \in S \cap V$. Theorem 4.19 thus proves that σ is unramified at infinity.

5. GEOMETRY OF THE RAMIFICATION LOCUS

As above, we work with a general cubic fourfold $X \subset \mathbb{P}^5$. We begin with some basic facts stemming from the geometric description of the ramification points of the g_3^1 's on the curves C_x given in Proposition 3.10 and the discussion following that. As in the introduction, we define the following loci, recalling that $W' \subset X$ was defined in (7) as the locus of points x whose corresponding curve C_x has worse than 1-nodal singularities.

Definition 5.1. Denote by

$$R, R' \subset I$$

the closures of the locus of all ramification points and the residual to all ramification points respectively, of one of the possibly two g_3^1 of the family of curves $p : I \rightarrow X$ restricted to $X \setminus W'$. Denote also by N the (closure of the) locus of triple ramification points.

Note that the locus R contains the locus $q^{-1}(S)$ as can be seen from the following. On a family of smooth curves with a family of g_3^1 's, if we degenerate to a nodal curve, two of the ramification points “converge” to the node (as for example can be seen by the theory of admissible covers). A point $t \in R$ lying on the smooth part of the curve C_x , with $x \in X \setminus W'$, is a ramification point of one of the (possibly) two g_3^1 that C_x possesses. As explained at the end of Section 3, if $t' \in C_x$ is the residual to t in the above g_3^1 , then $x \in \ell_t \cap \ell_{t'}$, the 2-plane Π spanned by ℓ_t and $\ell_{t'}$ is tangent to X along ℓ and then $\ell_{t'}$ is the line which is residual to ℓ_t in the intersection of Π with X . In particular, a point of $t \in N$ lying on a curve C_x as above corresponds to a triple tangent plane Π to X along ℓ_t . Note also that by the discussion following Corollary 4.12, a node t of C_x corresponds to a line ℓ_t of second type and (in general) a pair of tangent planes to X along ℓ_t so that the two residual lines to ℓ_t in the intersection of those planes with X intersect ℓ_t at the point x .

Lemma 5.2. For $q : I \rightarrow F$ the projection and $V \subset F$ the locus of triple lines, we have

$$q^{-1}(V) = N \subset I.$$

Proof. As the intersection of $q^{-1}(V)$ with the family of curves $p : I \rightarrow X$ restricted to $X \setminus W'$ is the triple ramification locus from Proposition 3.10, we certainly have that $N \subseteq q^{-1}(V)$. We now prove the reverse inclusion $N \supseteq q^{-1}(V)$. From Theorem 4.4, V is irreducible. Hence, $q^{-1}(V)$ is not contained in N only if its intersection with $p^{-1}(W')$ contains an open set of $q^{-1}(V)$. But this cannot happen because $V \neq S$ so the generic point $[\ell] \in V$ represents a line of first type, which in turn implies that $p(q^{-1}[\ell])$ intersects W' at most at finitely many points. \square

We recall that $I = \{(p, \ell), p \in \ell\} \subset X \times F$ and that

$$\tilde{F} = \{(\ell, \Pi), \Pi \text{ tangent to } X \text{ along } \ell\} \subset F \times G(3, 6).$$

The pullback of the family I from F to \tilde{F} , under the blow up $\pi : \tilde{F} \rightarrow F$ of S , is given by $\tilde{I} = \{(p, (\ell, \Pi)), p \in \ell\} \subset X \times \tilde{F}$. Let $\tilde{\pi} : \tilde{I} \rightarrow I$, with $(p, (\ell, \Pi)) \rightarrow (p, \ell)$ be the natural morphism. Note that by identifying \tilde{F} with the graph of the Voisin map, we may represent the points of \tilde{I} by tuples $(p, (\ell, \ell', \Pi))$, with $(\pi, (\ell, \Pi))$ as above and ℓ' the residual line to ℓ in the intersection of Π with X . Define now

$$\tilde{R} = \{(p, (\ell, \ell', \Pi)) \in \tilde{I}, p \in \ell \cap \ell'\}$$

and consider the diagram

$$(26) \quad \begin{array}{ccccc} \tilde{R} & \hookrightarrow & \tilde{I} & \xrightarrow{\tilde{q}} & \tilde{F} \\ & & \downarrow \tilde{\pi} & & \downarrow \pi \\ R & \hookrightarrow & I & \xrightarrow{q} & F. \end{array}$$

Theorem 5.3. *We have that*

- (1) $\text{Bl}_{\tilde{V}} \tilde{F} \cong \tilde{R}$,
- (2) $\tilde{\pi}(\tilde{R}) \subset R$ and
- (3) *the induced map $\tilde{\pi} : \tilde{R} \rightarrow R$ is generically two-to-one over $q^{-1}(S)$ and is an isomorphism otherwise.*

As a consequence, R is irreducible with normalisation \tilde{R} , and is smooth outside $q^{-1}(S)$ over which the normalisation is generically two-to-one.

Proof. Continuing with the notation established in the previous section, denote by $\text{pr}_1 : \mathcal{G} \rightarrow G(2, 6)$ the projection $(\ell, \ell', \Pi) \rightarrow \ell$. Define also

$$\tilde{\mathcal{I}}_{\mathcal{G}} = \mathbb{P}(\text{pr}_1^*(\mathcal{U}_2)) \subset \mathbb{P}^5 \times \mathcal{G}$$

and its subset

$$\tilde{\mathcal{R}}_{\mathcal{G}} = \{(p, (\ell, \ell', \Pi)), p \in \ell \cap \ell'\} \subset \tilde{\mathcal{I}}_{\mathcal{G}}.$$

Any two fibres of $\tilde{\mathcal{R}}_{\mathcal{G}} \rightarrow \mathbb{P}^5$ are isomorphic, and over a point $p \in \mathbb{P}^5$, this is $\{(\ell, \ell', \Pi), p \in \ell \cap \ell'\} \subset \mathcal{G}$ and this fibre is itself a \mathbb{P}^1 -fibration over $\{(\ell, \Pi), p \in \ell \subset \Pi\}$, parametrising lines in a plane passing through a point. Finally $\{(\ell, \Pi), p \in \ell \subset \Pi\}$ is again a \mathbb{P}^1 -fibration over the Schubert cycle $\Sigma_3^p \subset G(3, 6)$ parametrising 2-planes through the point $p \in \mathbb{P}^5$. Note now that if we fix a hyperplane $H = \mathbb{P}^4 \subset \mathbb{P}^5$ not containing p , we obtain an isomorphism $\Sigma_3^p \cong G(2, H) = G(2, 5)$ by taking a 2-plane Π containing p and giving the line $\Pi \cap H$. Hence the image in $G(3, 6)$ is smooth as it is isomorphic to a Grassmannian itself. All in all, we obtain that $\tilde{\mathcal{R}}_{\mathcal{G}}$ is smooth and irreducible.

The projection $\tilde{\mathcal{R}}_{\mathcal{G}} \rightarrow \mathcal{G}$ is a birational morphism with exceptional locus a \mathbb{P}^1 -fibration over \mathcal{G}_{Δ} . Since $\tilde{\mathcal{R}}_{\mathcal{G}}$ is smooth, it is necessarily isomorphic to the blow up of \mathcal{G} at \mathcal{G}_{Δ} (see [GH78, p.604, item 4]).

We now pull everything back to $\mathbb{P}^5 \times \tilde{\mathcal{F}}$ via the projection $\text{pr} = (\text{id}, \text{pr}_2)$ for $\text{pr}_2 : \tilde{\mathcal{F}} \rightarrow \mathcal{G}$. In other words, denote by

$$\tilde{\mathcal{I}} = \text{pr}^{-1}(\tilde{\mathcal{I}}_{\mathcal{G}}) \subset \mathbb{P}^5 \times \tilde{\mathcal{F}}$$

and $\tilde{\mathcal{R}} \subset \tilde{\mathcal{I}}$ the inverse image of $\tilde{\mathcal{R}}_{\mathcal{G}}$. The locus $\tilde{\mathcal{R}}$ is smooth and irreducible as it admits a smooth morphism over $\tilde{\mathcal{R}}_{\mathcal{G}}$ which is smooth and irreducible as we showed above. Note that for $X = \mathbf{V}(G) \in |\mathcal{O}_{\mathbb{P}^5}(3)|$ we have that the restrictions over X are

$$\begin{aligned}\tilde{\mathcal{I}}|_X &= \tilde{I}_X, \\ \tilde{\mathcal{R}}|_X &= \tilde{R}_X,\end{aligned}$$

where \tilde{I}_X is as in Diagram (26), i.e., is the blowup of the universal family of lines I_X over X at the universal locus of type two lines $q^{-1}(S_X)$, and \tilde{R}_X a subvariety of it, both denoted with the subscript X to distinguish between the fibres.

As $\tilde{\mathcal{R}}$ is smooth irreducible, the fibre $\tilde{R}_X = \tilde{\mathcal{R}}|_X$ over a general point $X \in |\mathcal{O}(3)|$ is smooth. It is also birational to \tilde{F}_X , and over the locus \tilde{V}_X it is codimension one since it is a \mathbb{P}^1 -fibration over it. Now the irreducibility of $\tilde{\mathcal{R}}$ implies that a general fibre \tilde{R}_X is equidimensional, so from the aforementioned birationality it must be irreducible. As the fibre of \tilde{R}_X over \tilde{V}_X is a line \mathbb{P}^1 , and it is otherwise isomorphic to \tilde{F}_X , we obtain from [GH78, p.604, item 4] that

$$(27) \quad \tilde{R}_X \cong \mathrm{Bl}_{\tilde{V}}(\tilde{F}).$$

By construction

$$\tilde{R}_X = \{(p, (\ell, \ell', \Pi)), p \in \ell \cap \ell'\} \subset X \times \tilde{F}.$$

By what we discussed after Proposition 3.10 and Definition 5.1, when $p \notin W'$ (i.e., the generic case), then $(p, (\ell, \ell', \Pi))$ maps to a general point of R . Hence from the irreducibility of \tilde{R}_X , we get a surjection $\tilde{\pi} : \tilde{R}_X \rightarrow R_X$. Over the complement of $q^{-1}(S)$ it is an isomorphism since there the map $\tilde{I}_X \rightarrow I_X$ is an isomorphism. Over $q^{-1}(S)$ it is generically two to one from Corollary 4.12. \square

Remark 5.4. In fact it is not hard to see, e.g., using [EH00, Proposition IV-21] that the isomorphism of Equation (27) holds globally, i.e., that $\tilde{\mathcal{R}} \cong \mathrm{Bl}_{\tilde{V}}(\tilde{\mathcal{F}})$.

By composing the blowup map with the projection to X , we obtain

$$\mathrm{Bl}_{\tilde{V}} \tilde{F} \longrightarrow \tilde{I} \longrightarrow I \longrightarrow X.$$

Outside the exceptional divisor, the first morphism sends a point (ℓ, ℓ', Π) , so that $X \cap \Pi = 2\ell + \ell'$, to the point in \tilde{I} in the fibre over $[\ell]$ which is the point of intersection with ℓ' . On the other hand, it identifies the points in the exceptional divisor in the fibre over (ℓ, ℓ, Π) with the points of ℓ .

Corollary 5.5. *The map $\psi : \tilde{F} \dashrightarrow X$ is resolved by blowing up \tilde{V} , and the resolution morphism is given by the above composition.*

From the proposition above, the projection $q : R \rightarrow F$ is birational, thus R is a birational model of F inside I .

Lemma 5.6. *Let $X \subset \mathbb{P}^5$ be a general cubic fourfold. The map $\psi : F \dashrightarrow X$ defined in Diagram (4) is a dominant rational map of degree 24.*

Proof. Let $x \in X$ be a general point. We want to count the number of pairs (ℓ, ℓ') of distinct lines through x such that the 2-plane spanned by ℓ, ℓ' is tangent to ℓ . From Proposition 3.10 the number of such pairs is exactly the number of ramification points of the two g_3^1 's on C_x , which is 24. Each contributes a distinct point as the ramification is simple at all points since the curve C_x is Brill–Noether general from Proposition 3.5. \square

We now compute the classes of R, R' and N in the Chow group of I . Given Proposition 3.5, perhaps the most natural way to do this is via admissible covers on the moduli space of curves, which we include as Appendix A and is independent of the proof below (one uses Lemma 2.6 to write these classes in terms of q^*H_F, l), but does not recover the class of N .

Proposition 5.7. *The classes of R, R' and N in I are as follows, for $l = c_1(\mathcal{O}_I(1))$ the tautological class*

$$\begin{aligned} [R] &= 4q^*H_F + l, \\ [R'] &= 4q^*H_F + 16l, \\ [N] &= 4l^2 - 4lq^*H_F + 25q^*c_2(\mathcal{U}_F). \end{aligned}$$

Remark 5.8. We note that using the computation of the class $[R']$ as a consequence of Appendix A one recovers that the degree of the Voisin map satisfies

$$\deg \phi = [R']q^*[\ell] = 16.$$

In an earlier version of this paper we computed the classes of R, R' and N (and as a consequence the class of V) by avoiding the techniques of Section 4 and instead doing intersection theory on a partial resolution of singularities J of $I \times_X I$, and in particular interpreting R as the image of a natural degeneracy locus in J - i.e., this method constructed a natural geometric correspondence over (F, X) inside $I \times_X I$. This involved various technicalities to step around the locus $W' \subset X$ parametrising the worse than 1-nodal fibres of $p : I \rightarrow X$, so in this version of the paper we chose instead to construct \tilde{R} via the universal family in Proposition 5.3, and then to resort to computations of [Ame11, SV16, OSY19] for the following proof of Proposition 5.7.

Proof of 5.7. The computation of the class of N follows from Theorem 4.7 and Lemmas 5.2, 2.6. For the other two classes, we follow closely the notation of Theorem 4.7 and Proposition 5.3.

The inclusion $\gamma^*\mathcal{U}_2 \hookrightarrow \beta^*\mathcal{U}_3$ in the exact sequence (18) yields, for $i = 1, 2$, inclusions $\iota_\phi^*\alpha_i^*\gamma^*\mathcal{U}_2 \hookrightarrow \iota_\phi^*\alpha^*\mathcal{U}_3$ and thus an embedding

$$j_i : P_i := \mathbb{P}(\iota_\phi^*\alpha_i^*\gamma^*\mathcal{U}_2) \hookrightarrow P := \mathbb{P}(\iota_\phi^*\alpha^*\mathcal{U}_3)$$

of bundles over Γ_ϕ . Let $u : P \rightarrow \Gamma_\phi$ and $u_i = u|_{P_i} : P_i \rightarrow \Gamma_\phi$ the natural maps. For $i, k \in \{1, 2\}, i \neq k$, [EH16, Proposition 9.13] gives

$$(28) \quad j_i^*j_{k*}[P_k] = c_1(\mathcal{O}_{P_i}(1)) + u_i^*(\iota_\phi^*\alpha^*c_1(\mathcal{U}_3)) - \iota_\phi^*\alpha_k^*\gamma^*c_1(\mathcal{U}_2)$$

on P_i .

We now identify Γ_ϕ with \tilde{F} (via the isomorphism π'). Under this identification, $P_1 = \tilde{I} := \pi^*I$ and $P_2 = \tilde{I}' := \phi^*I$. We denote by $[\tilde{R}] = j_1^*j_{2*}[P_2]$ in

\tilde{I} and by $[\tilde{R}'] = j_2^* j_{1*} [P_1]$ in \tilde{I}' . The cycle $[\tilde{R}]$ in \tilde{I} is geometrically the cycle representing the locus which on a fibre ℓ of \tilde{I} over a general point (ℓ, Π) of \tilde{F} is the intersection point of ℓ with the residual line ℓ' in the plane Π . Thus, $\tilde{\pi}_*[\tilde{R}] = [R]$, with $\tilde{\pi} : \tilde{I} \rightarrow I$ the natural map corresponding to the blow up map $\pi : \tilde{F} \rightarrow F$. Similarly, $\tilde{\phi}_*[\tilde{R}'] = [R']$, with $\tilde{\phi} : \tilde{I}' \rightarrow I$ the natural map corresponding to the Voisin map $\phi : \tilde{F} \rightarrow F$.

We then have from (28) and Lemma 4.8

$$\begin{aligned} [\tilde{R}] &= c_1(\mathcal{O}_{\tilde{I}}(1)) + u_1^*((-3H_F + E) + (7H_F - 3E)) \\ &= c_1(\mathcal{O}_{\tilde{I}}(1)) + u_1^*(4H_F - 2E). \end{aligned}$$

By pushing forward to I by the map $\tilde{\pi}$ we get $[R] = 4q^*H_F + l$ as claimed.

The computation for $[\tilde{R}']$ is similar. We have first from (28)

$$\begin{aligned} [\tilde{R}'] &= c_1(\mathcal{O}_{\tilde{I}'}(1)) + u_2^*((-3H + E) + H) \\ &= c_1(\mathcal{O}_{\tilde{I}'}(1)) + u_2^*(-2H + E). \end{aligned}$$

By pushing forward to I by the map $\tilde{\phi}$, we get $\tilde{\phi}_*c_1(\mathcal{O}_{\tilde{I}'}(1)) = 16l$ as the degree of v is 16. Finally, $\tilde{\phi}_*u_2^*H = 28q^*H_F$ (see [SV16, Proposition 21.4]) and $\tilde{\phi}_*u_2^*E = 60q^*H_F$ from [OSY19, Relation (18)]. We conclude $[R'] = 4q^*H_F + 16l$. \square

APPENDIX A. ADMISSIBLE COVERS

In this section we reinterpret the divisors $R, R' \subset I$ of Section 5 using Hurwitz spaces, and in particular obtain again some of the results of Proposition 5.7.

Proposition A.1. *Let λ be the determinant of the Hodge bundle of the curve fibration $p : I \rightarrow X$. Then*

- $[R] = 4\omega_p + 8p^*\lambda - p^*[W]$,
- $[R'] = 4\omega_p + 68p^*\lambda - 8p^*[W]$.

Proof. We denote by $\tilde{\mathcal{H}}_{4,3}$ the part of the space of admissible covers of genus 4 and degree 3, with $b = 12$ labelled branched points, consisting of the union of the locus of smooth admissible covers and the boundary divisors E_0 and E_3 (see [vdGK12] for the notation) where a point of E_0 corresponds to a map $X \rightarrow P$, with $X = (C_1 \cup R_1) \cup C_2$ and $P = \mathbb{P}^1 \cup \mathbb{P}^1$, with C_2 a smooth genus 3 curve and all other curves rational, as in [vdGK12, Proposition 3.1, ii] for $k = 2$. A point of E_3 corresponds to a map $X \rightarrow P$, with $X = S \cup C$, where C is a smooth genus 4 curve and all other curves rational, as in [vdGK12, Proposition 4.1, ii] for $k = 2$. We have the diagram of universal curves

$$\begin{array}{ccccc} \overline{\mathcal{M}}_{0,b+1} & \xleftarrow{f} & \mathcal{C}_{\tilde{\mathcal{H}}_{4,3}} & \xrightarrow{\phi} & \overline{\mathcal{M}}_{4,1} \\ \pi_b \downarrow & & \downarrow t & & \downarrow \pi \\ \overline{\mathcal{M}}_{0,b} & \xleftarrow{j} & \tilde{\mathcal{H}}_{4,3} & \xrightarrow{h} & \overline{\mathcal{M}}_4. \end{array}$$

The map h is generically finite of degree $N_0 = 2b!$ which is the number of g_3^1 's times the possible permutations of the branch points. The map t is

equipped with b disjoint sections $\tau_i : \tilde{\mathcal{H}}_{4,3} \rightarrow \mathcal{C}_{\tilde{\mathcal{H}}_{4,3}}$ corresponding to the b labelled ramification points sitting over the labelled branched points. These commute with the natural sections $s_i : \overline{\mathcal{M}}_{0,b} \rightarrow \overline{\mathcal{M}}_{0,b+1}$, i.e., $f\tau_i = s_i j$. Let T_i be the image of τ_i . Note that each ramification point of a 3-to-1 map $F : X \rightarrow P$ admits all the labels - in other words, if $i \in \{1, \dots, b\}$ and p is a ramification point of F then there exists an $h \in \tilde{\mathcal{H}}_{4,3}$ so that the fibre over h represents the map F and $\tau_i(h) = p$. Therefore the images of the T_i 's under the map ϕ coincide and are equal to the union of all ramification points of all the g_3^1 's.

The map h is simply branched at E_0 and E_3 . The map ϕ factors via the fibre product $\mathcal{X} = \overline{\mathcal{M}}_{4,1} \times_{\overline{\mathcal{M}}_4} \tilde{\mathcal{H}}_{4,3}$. Let

$$\begin{array}{ccccc} \mathcal{C}_{\tilde{\mathcal{H}}_{4,3}} & \xrightarrow{\rho} & \mathcal{X} & \xrightarrow{\tilde{h}} & \overline{\mathcal{M}}_{4,1} \\ & \searrow t & \downarrow \tilde{\pi} & & \downarrow \pi \\ & & \tilde{\mathcal{H}}_{4,3} & \xrightarrow{h} & \overline{\mathcal{M}}_4 \end{array}$$

the induced diagram, with $\phi = \tilde{h}\rho$. We have $\omega_{\tilde{\pi}} = \tilde{h}^*\omega_{\pi}$. The map h is ramified at the points $x \in E_0$ and therefore the fibre of $\tilde{\pi}$ over x has an A_1 singularity corresponding to the node over the fibre of π . The map ρ resolves the singularity and the corresponding exceptional divisor \mathcal{C}_1 (the union of the C_1 from above) consists of (-2) -curves. In addition, on the fibres over E_0 the map ρ contracts an exceptional divisor \mathcal{R}_1 which consists of (-1) -curves - blow-up along a smooth locus (the union of the tails R_1 as above). The map h is also ramified at the points $x \in E_3$ but since the fibre over x maps to a smooth curve, the fibre of $\tilde{\pi}$ over x is a smooth curve. The map ρ is the blow-up at the triple ramification point locus of the fibres of $\tilde{\pi}$. The exceptional divisor \mathcal{S} consists of (-1) -curves (the curves S as above). Since the restriction of ω_t on (-2) -curves is trivial as C_1 gets contracted via ϕ , we get that $\omega_t = \phi^*\omega_{\pi} + B$ with $B = \mathcal{R}_1 + \mathcal{S}$.

Let α be a divisor class on $\mathcal{C}_{\tilde{\mathcal{H}}_{4,3}}$. We then have

$$\begin{aligned} \pi_*(\omega_{\pi}\phi_*\alpha) &= \pi_*\phi_*(\phi^*\omega_{\pi}\alpha) \\ &= h_*t_*(\phi^*\omega_{\pi}\alpha) \\ &= h_*t_*(\omega_t\alpha - B\alpha) \\ &= h_*t_*(\omega_t\alpha) - h_*t_*(B\alpha). \end{aligned}$$

We have $\omega_t = f^*\omega_{\pi_b} + R_f$, with R_f the ramification divisor of the map f , that is $R_f = \sum_{i=1}^b T_i$. Then

$$\begin{aligned} h_*t_*(\omega_t\alpha) &= h_*t_*(f^*\omega_{\pi_b}\alpha) + h_*t_*(R_f\alpha) \\ &= h_*t_*(f^*\omega_{\pi_b}\alpha) + \sum_{i=1}^b h_*t_*(\tau_i^*\alpha) \\ &= h_*t_*(f^*\omega_{\pi_b}\alpha) + \sum_{i=1}^b h_*(\tau_i^*\alpha). \end{aligned}$$

Suppose now that $\alpha = R_f = \sum_{i=1}^b T_i$. Note that the degree of R over $\overline{\mathcal{M}}_4$ is $2b$ so we have

$$R = \frac{2}{N_0} \phi_* R_f.$$

By [vdGK11, Lemma 4.2] we have

$$\begin{aligned} \sum_{i=1}^b \tau_i^* R_f &= \sum_{i=1}^b \tau_i^* T_i = \sum_{i=1}^b t_* \tau_{i*} \tau_i^* T_i \\ &= \sum_{i=1}^b t_* T_i^2 = t_*(R_f^2) = -\frac{1}{2} j^* \psi \end{aligned}$$

and $t_*(f^* \omega_{\pi_b} R_f) = j^* \psi$. Therefore $h_* t_*(\omega_t R_f) = \frac{1}{2} h_* j^* \psi$. Also, $t_*(BR_f) = 2E_3$. This is because \mathcal{S} has two ramification points and \mathcal{R}_1 has none. Therefore

$$\pi_*(\omega_\pi \phi_* R_f) = \frac{1}{2} h_* j^* \psi - 2h_* E_3.$$

Following formulas in [vdGK12] we have:

$$\begin{aligned} h_* j^* \psi &= 40N_0(9\lambda - \delta_0) \\ 2h_* E_3 &= N_0(132\lambda - 15\delta_0) \end{aligned}$$

and so,

$$\begin{aligned} \pi_*(\omega_\pi R) &= \frac{2}{N_0} \pi_*(\omega_\pi \phi_* R_f) = 40(9\lambda - \delta_0) - 2(132\lambda - 15\delta_0) \\ &= 96\lambda - 10\delta_0. \end{aligned}$$

Writing $R = a\omega_\pi + b\pi^*\lambda + c\pi^*\delta_0$, we get $a = 4$ (because the fibre has 24 points and $24 = 4 \cdot 6$, with $6 = 2g - 2$). This implies

$$\pi_*(\omega_\pi R) = 4(12\lambda - \delta_0) + 6b\lambda + 6c\delta_0 = (48 + 6b)\lambda + (-4 + 6c)\delta_0.$$

Hence $48 + 6b = 96$ which implies $b = 8$ and $-4 + 6c = -10$ giving $c = -1$.

Suppose now that $\alpha = A$ is the sum of the residual sections to the T_i 's. Then $R' = \frac{2}{N_0} \phi_* A$. We have $h_* t_*(\omega_t A) = h_* t_*(f^* \omega_{\pi_b} A) = h_* t_*(f^* \omega_{\pi_b} A) = j^* \psi$ and $B = 2E_0 + 2E_3$. This is because both \mathcal{R}_1 and \mathcal{S} contain two points which are residual to the ramification points. Then

$$\pi_*(\omega_\pi \phi_* A) = h_* j^* \psi - 2h_* E_0 - 2h_* E_3.$$

We have $h_* E_0 = \frac{N_0}{2} \delta_0$. Therefore $\pi_*(\omega_\pi R_1) = \frac{2}{N_0} \pi_*(\omega_\pi \phi_* A) = 80(9\lambda - \delta_0) - 2(132\lambda - 15\delta_0) - 2\delta_0 = 456\lambda - 52\delta_0$. Writing now $R' = 4\omega_\pi + b\pi^*\lambda + c\pi^*\delta_0$ we have as above that $48 + 6b = 456$ which implies $b = 68$ and $-4 + 6c = -52$ giving $c = -8$. \square

REFERENCES

- [AK77] Allen B. Altman and Steven L. Kleiman. Foundations of the theory of Fano schemes. *Compositio Math.*, 34(1):3–47, 1977.
- [Ame11] Ekaterina Amerik. A computation of invariants of a rational self-map. *Annales de la faculté des sciences de Toulouse Mathématiques*, 18(3):481–493, 2011.
- [BVdV79] W. Barth and A. Van de Ven. Fano varieties of lines on hypersurfaces. *Arch. Math. (Basel)*, 31(1):96–104, 1978/79.

- [BD85] Arnaud Beauville and Ron Donagi. La variété des droites d’une hypersurface cubique de dimension 4. *C. R. Acad. Sci. Paris Sér. I Math.*, 301(14):703–706, 1985.
- [Bui94] Alexandru Buium. *Differential algebra and Diophantine geometry*. Actualités Mathématiques. [Current Mathematical Topics]. Hermann, Paris, 1994.
- [CML09] Sebastian Casalaina-Martin and Radu Laza. The moduli space of cubic threefolds via degenerations of the intermediate Jacobian. *J. Reine Angew. Math.*, 633:29–65, 2009.
- [CG72] C. Herbert Clemens and Phillip A. Griffiths. The intermediate Jacobian of the cubic threefold. *Ann. of Math. (2)*, 95:281–356, 1972.
- [CTS21] Jean-Louis Colliot-Thélène and Alexei N. Skorobogatov. *The Brauer-Grothendieck group*, volume 71 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer, Cham, [2021] ©2021.
- [CS09] Izzet Coskun and Jason Starr. Rational curves on smooth cubic hypersurfaces. *Int. Math. Res. Not. IMRN*, (24):4626–4641, 2009.
- [DvGK05] I. Dolgachev, B. van Geemen, and S. Kondō. A complex ball uniformization of the moduli space of cubic surfaces via periods of $K3$ surfaces. *J. Reine Angew. Math.*, 588:99–148, 2005.
- [Ele85] Georges Elençwajg. Brauer group of fibrations and symmetric products of curves. *Proc. Amer. Math. Soc.*, 94(4):597–602, 1985.
- [EH87] David Eisenbud and Joe Harris. The Kodaira dimension of the moduli space of curves of genus ≥ 23 . *Invent. Math.*, 90(2):359–387, 1987.
- [EH00] David Eisenbud and Joe Harris. *The geometry of schemes*, volume 197 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000.
- [EH16] David Eisenbud and Joe Harris. *3264 and all that—a second course in algebraic geometry*. Cambridge University Press, Cambridge, 2016.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.
- [GK20] Frank Gounelas and Alexis Kouvidakis. On some invariants of cubic fourfolds. *arXiv e-prints*, page [arXiv:2008.05162](https://arxiv.org/abs/2008.05162), August 2020.
- [GH78] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics.
- [Har84] J. Harris. On the Kodaira dimension of the moduli space of curves. II. The even-genus case. *Invent. Math.*, 75(3):437–466, 1984.
- [HM98] Joe Harris and Ian Morrison. *Moduli of curves*, volume 187 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1998.
- [Huy20] Daniel Huybrechts. The geometry of cubic hypersurfaces. Lecture notes. <http://www.math.uni-bonn.de/people/huybrech/>, 2021.
- [LSV17] Radu Laza, Giulia Saccà, and Claire Voisin. A hyper-kähler compactification of the intermediate jacobian fibration associated with a cubic 4-fold. *Acta Mathematica*, 218(1):55–135, 2017.
- [NO21] Denis Nesterov and Georg Oberdieck. Elliptic curves in hyper-Kähler varieties. *Int. Math. Res. Not. IMRN*, (4):2962–2990, 2021.
- [OSY19] Georg Oberdieck, Junliang Shen, and Qizheng Yin. Rational curves in holomorphic symplectic varieties and Gromov-Witten invariants. *Adv. Math.*, 357:106829, 8, 2019.
- [SV16] Mingmin Shen and Charles Vial. The Fourier transform for certain hyperkähler fourfolds. *Mem. Amer. Math. Soc.*, 240(1139):vii+163, 2016.
- [Sta21] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2021.

- [vdGK11] Gerard van der Geer and Alexis Kouvidakis. The Hodge bundle on Hurwitz spaces. *Pure Appl. Math. Q.*, 7(4, Special Issue: In memory of Eckart Viehweg):1297–1307, 2011.
- [vdGK12] Gerard van der Geer and Alexis Kouvidakis. The class of a Hurwitz divisor on the moduli of curves of even genus. *Asian J. Math.*, 16(4):787–805, 2012.
- [Voi04] Claire Voisin. Intrinsic pseudo-volume forms and K -correspondences. In *The Fano Conference*, pages 761–792. Univ. Torino, Turin, 2004.
- [Voi10] Claire Voisin. Coniveau 2 complete intersections and effective cones. *Geom. Funct. Anal.*, 19(5):1494–1513, 2010.
- [Voi14] Claire Voisin. *Chow rings, decomposition of the diagonal, and the topology of families*, volume 187 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2014.
- [W05] C.T.C. Wall. Sextic curves and quartic surfaces with higher singularities. Preprint, 2005.

GEORG-AUGUST-UNIVERSITÄT GÖTTINGEN, FAKULTÄT FÜR MATHEMATIK UND INFORMATIK, BUNSENSTR. 3-5, 37073 GÖTTINGEN, GERMANY

Email address: `gounelas@mathematik.uni-goettingen.de`

DEPT. OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF CRETE, 70013 HERAKLION, GREECE.

Email address: `kouvid@uoc.gr`