

MAHLER MEASURE NUMEROLOGY

WADIM ZUDILIN

*To David Boyd, an outstanding conductor of the Mahler measures,
on the occasion of his 80th birthday*

ABSTRACT. We discuss some (conjectural) evaluations of L -values attached to elliptic curves of conductors 15, 21, 24 and 32 as ‘hypergeometric periods’. These numerical observations are motivated by the Mahler measures of three-variable polynomials.

The (logarithmic) Mahler measure

$$m(P(x_1, \dots, x_k)) = \frac{1}{(2\pi i)^k} \int \cdots \int_{|x_1|=\dots=|x_k|=1} \log |P(x_1, \dots, x_k)| \frac{dx_1}{x_1} \cdots \frac{dx_k}{x_k}$$

of an k -variable (Laurent) polynomial $P(x_1, \dots, x_k) \in \mathbb{C}[x_1, \dots, x_k]$ is a quite unique attractor of numerous problems in mathematics [3]. One of big problems (well open even in the case $k = 1$!) is the range of the Mahler measures attached to polynomials with *integer* coefficients [1]. A particular aspect of this problem is a remarkable connection of such Mahler measures to the L -values of algebraic varieties, usually related to the zero loci of the underlying polynomials $P(x_1, \dots, x_k)$. Research in this direction originated in the 1990s: the works of Deninger [4], Boyd [2] and Rodriguez Villegas [8] set up in depth the story for the case of two-variable polynomials $P(x, y)$ such that $P(x, y) = 0$ defines an elliptic curve. The papers all feature the family of two-variable Mahler measures

$$\mu(k) = m(k + (x + 1/x)(y + 1/y)),$$

which is indeed quite special. Illustrative examples for the range $0 < k < 4$ include

$$\begin{aligned} \mu(1) &= L'(E_{15a8}, 0), & \mu(\sqrt{2}) &= 4L'(E_{56a1}, 0), & \mu(2) &= L'(E_{24a4}, 0), \\ \mu(2\sqrt{2}) &= L'(E_{32a1}, 0), & \mu(3) &= \frac{1}{2}L'(E_{21a4}, 0), & \mu(2\sqrt{3}) &\stackrel{?}{=} 2L'(E_{72a1}, 0), \end{aligned}$$

where E_{NxM} refers to the elliptic curve labeled in accordance with [5] (the label corresponds to a Weierstrass form of the curve $k + (x + 1/x)(y + 1/y) = 0$ which is defined over \mathbb{Q} whenever $k^2 \in \mathbb{Q} \setminus \{0, 16\}$). Notice that for this range

$$\mu(k) = \frac{k}{4} \cdot {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2} \mid \frac{k^2}{16}\right) = \int_0^{k/4} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \mid x^2\right) dx, \quad (1)$$

so that the Mahler measure evaluations can be translated into equalities of the L -values and the values of hypergeometric functions.

Date: 17 September 2021.

2020 Mathematics Subject Classification. Primary 11R06; Secondary 11G05, 14G10, 33C20, 33C75.

The L -values of conductor 15, 21 and 24 elliptic curves also show up, conjecturally(!), in the three-variable Mahler measures¹

$$\begin{aligned} m(1+x+y+xy+z) &\stackrel{?}{=} -2L'(E_{15}, -1) = \frac{15^2}{4\pi^4}L(E_{15}, 3) = 0.4839979734\dots, \\ m((1+x)^2+y+z) &\stackrel{?}{=} -L'(E_{24}, -1) = \frac{7^2}{\pi^4}L(E_{24}, 3) = 0.7025655062\dots, \\ m(1+x+y-xy+z) &\stackrel{?}{=} -\frac{5}{4}L'(E_{21}, -1) = \frac{5 \cdot 21^2}{32\pi^4}L(E_{21}, 3) = 0.6242499823\dots \end{aligned}$$

The Mahler measures in all these cases are reduced to hypergeometric integrals [3, Section 6.3] (see also [9]), and there are also hypergeometric expressions available for $L'(E_{32}, -1)$ [10]. Here we highlight the corresponding formulae for the L -values, dropping off the intermediate Mahler measures, and give some additional formulae that are not linked to known conjectures on the Mahler measures but reflect similarities with the two-variable expression (1).

Conductor 15 ($k = 1$): The expectation in this case is

$$\begin{aligned} -L'(E_{15}, -1) &\stackrel{?}{=} \frac{1}{\pi} \int_{1/4}^1 {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-x^2\right) \log(4x) \, dx \\ &= -\frac{1}{\pi} \int_0^{1/4} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-x^2\right) \log(4x) \, dx. \end{aligned} \quad (2)$$

Notice the induced evaluation

$$\frac{1}{\pi} \int_0^1 {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-x^2\right) \log(4x) \, dx = 0 \quad (3)$$

established in [9]. There is a complimentary observation of H. Cohen (2018) in this case, not related to a Mahler measure:

$$L'(E_{15}, -2) \stackrel{?}{=} \frac{12}{\pi^2} \int_0^{1/4} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| x^2\right) \log^2(4x) \, dx. \quad (4)$$

Conductor 21 ($k = 3$): Here we expect

$$-5L'(E_{21}, -1) \stackrel{?}{=} \frac{2}{\pi} \int_{1/4}^2 {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-(1-x)^2\right) \log(4x) \, dx. \quad (5)$$

The argument $2x - x^2$ of the hypergeometric integrand in (5) ranges between $1 - k^2/16$ and 1 when $\frac{1}{4} < x < 1$ and then between 1 and 0 when $1 < x < 2$. Thus, taking into account the evaluation²

$$\frac{7\zeta(3)}{\pi^2} = -28\zeta'(-2) = \frac{1}{\pi} \int_0^1 {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-x^2\right) \log(4(1+x)) \, dx \quad (6)$$

¹The notation ' $\stackrel{?}{=}$ ' is used for equalities numerically observed.

²This is kindly reported to us by Ringeling [7] based on the findings in his work [6].

for the second part (see also (3)), our main expectation is

$$-\frac{5}{2}L'(E_{21}, -1) + 28\zeta'(-2) \stackrel{?}{=} \frac{1}{\pi} \int_0^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-x^2\right) \log(4(1-x)) dx. \quad (7)$$

Conductor 24 ($k = 2$): We know in this case [9] that

$$\begin{aligned} -L'(E_{24}, -1) &\stackrel{?}{=} \frac{2\sqrt{2}}{\pi\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n \Gamma(\frac{n}{2} + \frac{3}{4})}{n! \Gamma(\frac{n}{2} + \frac{5}{4}) (2n+1)^2} \left(\frac{1}{2}\right)^n \\ &= \frac{8\Gamma(\frac{3}{4})^2}{\pi^{5/2}} {}_5F_4\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4} \middle| \frac{1}{4}\right) + \frac{\Gamma(\frac{1}{4})^2}{54\pi^{5/2}} {}_5F_4\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{5}{4}, \frac{5}{4} \middle| \frac{1}{4}\right). \end{aligned} \quad (8)$$

However there is an identification of $L'(E_{24}, -1)$ similar to (2) and (7) but not related to a Mahler measure:

$$-2L'(E_{24}, -1) + 28\zeta'(-2) \stackrel{?}{=} \frac{1}{\pi} \int_0^{2/4} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| 1-x^2\right) \log\left(\frac{1-x}{x}\right) dx. \quad (9)$$

In a different direction, here is a striking connection of (8) with other L -values of the same elliptic curve³:

$$L'(E_{24}, 0) \stackrel{?}{=} \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n \Gamma(\frac{n}{2} + \frac{3}{4})}{n! \Gamma(\frac{n}{2} + \frac{5}{4}) (2n+1)} \left(-\frac{1}{2}\right)^n, \quad (10)$$

$$L(E_{24}, 1) = \frac{\sqrt{\pi}}{4\sqrt{2}} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n \Gamma(\frac{n}{2} + \frac{3}{4})}{n! \Gamma(\frac{n}{2} + \frac{5}{4})} \left(\frac{1}{2}\right)^n. \quad (11)$$

These two evaluations can be compared with

$$L'(E_{24}, 0) = \frac{1}{2} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^2}{n!^2 (2n+1)} \left(\frac{1}{4}\right)^n, \quad (12)$$

$$L(E_{24}, 1) = \frac{\pi}{8} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^2}{n!^2} \left(1 - \frac{1}{4}\right)^n, \quad (13)$$

where the first one comes out of the formula for $\mu(2)$ interpreted through (1).

³The proof of (11) we are aware of is somewhat tricky. One starts with

$$\begin{aligned} \frac{\sqrt{\pi}}{2} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n \Gamma(\frac{n}{2} + \frac{3}{4})}{n! \Gamma(\frac{n}{2} + \frac{5}{4})} z^n &= \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n}{n!} z^n \int_0^1 t^{n+1/2} (1-t^2)^{-1/2} dt = \int_0^1 \frac{\sqrt{t}}{\sqrt{(1-t^2)(1-zt)}} dt \\ &= \sqrt{2} \left(\int_0^1 \frac{du}{(1-\frac{1}{2}u^2)\sqrt{(1-u^2)(1-\frac{1}{2}(1+z)u^2)}} - \int_0^1 \frac{du}{\sqrt{(1-u^2)(1-\frac{1}{2}(1+z)u^2)}} \right) \end{aligned}$$

to arrive, when $z = \frac{1}{2}$, at

$$\int_0^1 \frac{du}{(1-\frac{1}{2}u^2)\sqrt{(1-u^2)(1-\frac{3}{4}u^2)}} = \frac{3}{2} \int_0^1 \frac{du}{\sqrt{(1-u^2)(1-\frac{3}{4}u^2)}},$$

and this allows one to conclude with (11) from (13). There seems to be no simple relation like this for $z = -\frac{1}{2}$ (and for other values of z from the real interval $|z| < 1$). This fact makes it difficult to lift the finding to (10).

Conductor 32 ($k = 2\sqrt{2}$): Here we have the following [10]:

$$\begin{aligned}
 -L'(E_{32}, -1) &= \frac{\Gamma(\frac{1}{4})^2}{6\sqrt{2}\pi^{5/2}} {}_4F_3\left(\begin{matrix} 1, 1, 1, \frac{1}{2} \\ \frac{7}{4}, \frac{3}{2}, \frac{3}{2} \end{matrix} \middle| 1\right) + \frac{4\Gamma(\frac{3}{4})^2}{\sqrt{2}\pi^{5/2}} {}_4F_3\left(\begin{matrix} 1, 1, 1, \frac{1}{2} \\ \frac{5}{4}, \frac{3}{2}, \frac{3}{2} \end{matrix} \middle| 1\right) \\
 &\quad + \frac{\Gamma(\frac{1}{4})^2}{2\sqrt{2}\pi^{5/2}} {}_4F_3\left(\begin{matrix} 1, 1, 1, \frac{1}{2} \\ \frac{3}{4}, \frac{3}{2}, \frac{3}{2} \end{matrix} \middle| 1\right). \tag{14}
 \end{aligned}$$

Conjecturally we find out the following representation resembling (2), (7) and (9):

$$-L'(E_{32}, -1) \stackrel{?}{=} \frac{1}{\pi} \int_0^{2\sqrt{2}/4} {}_2F_1\left(\begin{matrix} \frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix} \middle| 1-x^2\right) \log\left(\frac{1-x^2}{x^2}\right) dx. \tag{15}$$

Acknowledgements. I thank François Brunault and Berend Ringeling for inspiring conversations and assistance.

REFERENCES

- [1] D. W. BOYD, Speculations concerning the range of Mahler’s measure, *Canad. Math. Bull.* **24** (1981), 453–469.
- [2] D. W. BOYD, Mahler’s measure and special values of L -functions, *Experiment. Math.* **7** (1998), 37–82.
- [3] F. BRUNAULT and W. ZUDILIN, *Many Variations of Mahler Measures. A Lasting Symphony*, Austral. Math. Soc. Lecture Ser. **28** (Cambridge University Press, Cambridge, 2020).
- [4] C. DENINGER, Deligne periods of mixed motives, K -theory and the entropy of certain \mathbb{Z}^n -actions, *J. Amer. Math. Soc.* **10** (1997), 259–281.
- [5] THE LMFDB COLLABORATION, The L -functions and modular forms database, www.lmfdb.org (2013–2021).
- [6] B. RINGELING, Special zeta Mahler functions, *Preprint* arXiv:2012.09506 [math.NT] (2020).
- [7] B. RINGELING, An identity, *Private communication* (16 December 2020).
- [8] F. RODRIGUEZ VILLEGAS, Modular Mahler measures. I, in *Topics in number theory* (University Park, PA, 1997), Math. Appl. **467** (Kluwer Acad. Publ., Dordrecht, 1999), 17–48.
- [9] A. STRAUB and W. ZUDILIN, Short walk adventures, in *From analysis to visualization: A celebration of the life and legacy of Jonathan M. Borwein* (Callaghan, Australia, September 2017), B. Sims et al. (eds.), Springer Proceedings in Math. Stat. **313** (Springer, New York, 2020), 423–439.
- [10] W. ZUDILIN, Period(d)ness of L -values, in *Number Theory and Related Fields, In memory of Alf van der Poorten*, J.M. Borwein et al. (eds.), *Springer Proceedings in Math. Stat.* **43** (Springer, New York, 2013), 381–395.

DEPARTMENT OF MATHEMATICS, IMAPP, RADBOD UNIVERSITY, PO Box 9010, 6500 GL NIJMEGEN, NETHERLANDS

Email address: w.zudilin@math.ru.nl