

NONTRIVIAL EXAMPLES OF JN_p AND VJN_p FUNCTIONS

TIMO TAKALA

ABSTRACT. We study the John-Nirenberg space JN_p , which is a generalization of the space of bounded mean oscillation. In this paper we construct new JN_p functions, that increase the understanding of this function space. It is already known that $L^p(Q_0) \subsetneq JN_p(Q_0) \subsetneq L^{p,\infty}(Q_0)$. We construct a nonnegative function f such that $f^{1/p} \notin JN_p(Q_0)$ even though $f^{1/q} \in JN_q(Q_0)$, for every $q \in (p, \infty)$. We also show that if $|f|^{1/p} \in JN_p(Q_0)$, then $|f|^{1/q} \in JN_q(Q_0)$, where $q \geq p$. We present functions $g \in JN_p(Q_0) \setminus VJN_p(Q_0)$ and $g_0 \in VJN_p(Q_0) \setminus L^p(Q_0)$, proving the nontriviality of the vanishing subspace VJN_p . Finally we extend some results from JN_p type functions defined in a bounded cube to functions that are defined in the whole space \mathbb{R}^n .

1. INTRODUCTION

In their seminal paper [7] John and Nirenberg studied the famous space of bounded mean oscillation BMO and proved the profound John-Nirenberg inequality for BMO functions. In the same paper they also defined a generalization of BMO which has since become known as the John-Nirenberg space, or JN_p , for some parameter $p \in (1, \infty)$. The space JN_p is a generalization of BMO in the sense that the BMO norm of a function is the limit of its JN_p norm when p tends to infinity.

In this paper we define JN_p as in Definition 2.1 below: for a bounded cube $Q_0 \subset \mathbb{R}^n$ and a number $p \in (1, \infty)$, a function f is in $JN_p(Q_0)$ if $f \in L^1(Q_0)$ and

$$\|f\|_{JN_p(Q_0)}^p := \sup \sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p < \infty,$$

where the supremum is taken over all countable collections of pairwise disjoint cubes $(Q_i)_{i=1}^{\infty}$ that are contained in Q_0 . Here f_{Q_i} denotes the integral average of f over Q_i :

$$f_{Q_i} := \int_{Q_i} f := \frac{1}{|Q_i|} \int_{Q_i} f.$$

There are many other definitions for JN_p , some of which are not equivalent with this definition. This is because the space depends on how much we let the sets Q_i overlap. For example many of the definitions in the more general metric measure space, such as in [1, 6, 9, 11], use balls B_i instead of cubes, and these balls may overlap with each other in some definitions. This results in different spaces.

Date: September 20, 2021.

2020 Mathematics Subject Classification. 42B35, 46E30, 46E35.

Key words and phrases. John–Nirenberg space, vanishing subspace, Euclidian space, cube, bounded mean oscillation, John-Nirenberg inequality.

Other related function spaces include the dyadic JN_p [8], the John-Nirenberg-Campanato spaces [15], their localized versions [14] and the sparse JN_p [5]. For an extensive survey of John-Nirenberg type spaces see [17].

It is well-known that $L^p(Q_0) \subset JN_p(Q_0)$. The space JN_p is also embedded in the weak L^p space, $JN_p(Q_0) \subset L^{p,\infty}(Q_0)$. Further both of these inclusions are strict. An example of a function in $JN_p \setminus L^p$ was constructed in a recent paper [4]. Thus the space JN_p is a nontrivial space between L^p and $L^{p,\infty}$. Other than that, the behaviour of JN_p functions is still very much unknown.

One of the properties with L^p spaces is that $|f|^{1/q} \in L^q$ if and only if $|f|^{1/p} \in L^p$ whenever $p, q \in [1, \infty)$. A similar equivalence holds for weak L^p spaces. Since JN_p is a space between these spaces, this raises the question: does the equivalence hold for JN_p spaces? In Section 3 we give a negative answer to this question by constructing an example of a nonnegative function f such that $f^{1/p} \notin JN_p(Q_0)$ even though $f^{1/q} \in JN_q(Q_0)$ for every $q \in (p, \infty)$. This is our main result. The previous existing example of a JN_p function does not have this property of distinguishing JN_p spaces with different values of p . We also show a proof that if $|f|^{1/p} \in JN_p$, then $|f|^{1/q} \in JN_q$ for every $q \geq p$.

Additionally we study the vanishing subspace of JN_p , which is denoted as VJN_p . This subspace has been studied by Brudnyi [3] and by Tao et al. [16]. The space VJN_p is defined as a John-Nirenberg space counterpart to the famous space of vanishing mean oscillation, VMO , which is a subspace of BMO , and was first studied by Sarason [13]. It is already well-known that $L^p(Q_0) \subset VJN_p(Q_0) \subset JN_p(Q_0)$. Tao et al. proved that $L^p(Q_0) \neq VJN_p(Q_0)$ by showing that the dual of the dual of VJN_p is JN_p [16]. In Section 4 we present examples of functions in $JN_p \setminus VJN_p$ and in $VJN_p \setminus L^p$, thereby proving that $VJN_p(Q_0) \neq JN_p(Q_0)$. One of these functions is the same as the example of a function in $JN_p \setminus L^p$ presented by Dafni et al. [4]. The other is a function with a similar fractal construction.

Finally in Section 5 we study JN_p and VJN_p when the functions have an unbounded domain. More specifically we define these John-Nirenberg type spaces in \mathbb{R}^n . The embedding $JN_p \subset L^{p,\infty}$ has been proved in many different ways for a bounded domain - originally in [7] and further discussion can be found in [1, 2, 6, 9, 10, 15]. However, as far as we know, the question of whether this holds for unbounded domains has not been addressed in the literature. For the sake of completeness we prove that this embedding indeed holds in the case of the whole space \mathbb{R}^n , if we replace $L^{p,\infty}$ with $L^{p,\infty}/\mathbb{R}$ - the space of functions in $L^{p,\infty}$ modulo constant.

We also consider the space CJN_p : a subspace of JN_p , that has been studied by Tao et al. [16]. The space CJN_p is defined analogously to the space of continuous mean oscillation, CMO , which is a subspace of BMO , and was first announced by Neri [12]. We study CJN_p only on \mathbb{R}^n as on bounded domains it coincides with VJN_p . It is clear from the definitions that $L^p/\mathbb{R} \subset CJN_p \subset VJN_p \subset JN_p$. By extending the functions in Section 4 into the whole space \mathbb{R}^n we get functions that are in $JN_p(\mathbb{R}^n) \setminus VJN_p(\mathbb{R}^n)$ and in $CJN_p(\mathbb{R}^n) \setminus L^p(\mathbb{R}^n)/\mathbb{R}$, proving that the respective inclusions are strict. This answers [16, Question 5.8] and it partially answers [16, Question 5.6] and [17, Question 5.27].

Throughout this paper we assume that a function has its domain in the Euclidian space \mathbb{R}^n or in a bounded cube $Q_0 \subset \mathbb{R}^n$. Every cube in \mathbb{R}^n is assumed to have edges parallel to the coordinate axes.

2. PRELIMINARIES

2.1. Definition (JN_p). Let $Q_0 \subset \mathbb{R}^n$ be a bounded cube and let $1 < p < \infty$. A function f is in $JN_p(Q_0)$ if $f \in L^1(Q_0)$ and there is a constant $K < \infty$ such that

$$\sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p \leq K^p$$

for all countable collections of pairwise disjoint cubes $(Q_i)_{i=1}^{\infty}$ in Q_0 . We denote the smallest such number K by $\|f\|_{JN_p}$.

2.2. Remark. JN_p can also be defined with infimums instead of integral averages. This definition is equivalent with Definition 2.1, because

$$\inf_{c_i} \int_{Q_i} |f - c_i| \leq \int_{Q_i} |f - f_{Q_i}| \leq 2 \inf_{c_i} \int_{Q_i} |f - c_i|.$$

The infimum approach makes calculations sometimes much more simple. We denote the JN_p norm obtained with infimums as $\|f\|_{JN_p, \text{inf}}$.

One can also define JN_p by using medians which allows us to not use integrals at all. This approach is studied by Myyryläinen [11]. Even though we don't use the median type approach in the proofs, we would still like to acknowledge it here, because it was useful in constructing the function in Section 3. Notice that there is no point in defining JN_p as in Definition 2.1 when $p = 1$. This is because then we would simply have $JN_1(Q_0) = L^1(Q_0)$.

It is clear that if a function f is in $L^p(Q_0)$, then it is also in $JN_p(Q_0)$. This follows from Hölder's inequality:

$$\sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p \leq \sum_{i=1}^{\infty} |Q_i| \int_{Q_i} |f - f_{Q_i}|^p \leq \sum_{i=1}^{\infty} 2^p \int_{Q_i} |f|^p \quad (2.3)$$

for every partition $(Q_i)_{i=1}^{\infty}$ of Q_0 into disjoint cubes. The inclusion $L^p \subset JN_p$ is strict, but functions in $JN_p \setminus L^p$ are very complicated. An example of a function in $JN_p \setminus L^p$ was constructed in a recent paper [4].

If a function is in $JN_p(Q_0)$, then it is also in the weak L^p -space $L^{p,\infty}(Q_0)$. This result is a JN_p space counterpart of the famous John-Nirenberg lemma for BMO functions.

2.4. Theorem. Let $1 < p < \infty$, $Q_0 \subset \mathbb{R}^n$ a bounded cube and $f \in JN_p(Q_0)$. Then $f \in L^{p,\infty}(Q_0)$ and

$$\|f - f_{Q_0}\|_{L^{p,\infty}(Q_0)} \leq c \|f\|_{JN_p(Q_0)}$$

with some constant $c = c(n, p)$.

Conveniently the constant c does not depend on Q_0 even though the proof does rely on the boundedness of Q_0 . This embedding was originally proved in [7, Lemma 3]. In [6] this was generalized from cubes into John domains. In [1] and [9] the result was proven in a more general metric space with a doubling measure. In [15] a similar result was proven for John-Nirenberg-Campanato spaces, which are a generalization of John-Nirenberg spaces. Milman gave a new characterization of the weak L^p space as the Garsia-Rodemich space and proved the embedding of $JN_p(Q_0)$ in this space [10]. Berkovits et al. proved a good- λ inequality and used this to prove multiple embedding theorems including the embedding of JN_p to the weak L^p -space [2].

The inclusion $JN_p(Q_0) \subset L^{p,\infty}(Q_0)$ is strict as well. Consider the counterexample $f(x) = x^{-1/p}$ in the one-dimensional case $n = 1$ with $Q_0 =]0, 1[$. In this case f is in the weak L^p space. To prove that f is not in the JN_p space, we divide Q_0 into subintervals $Q_i = [2^{-i}, 2^{-i+1}[$ for every $i \in \mathbb{Z}_+$. Then it turns out that

$$|Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p = |Q_1| \left(\int_{Q_1} |f - f_{Q_1}| \right)^p > 0$$

for all $i \geq 1$. Hence the JN_p norm of f is infinite, and therefore $f \notin JN_p(Q_0)$. This counterexample can also be easily extended into the multidimensional case $Q_0 \subset \mathbb{R}^n$ by using the following proposition.

2.5. Proposition. *Let $Q_0 \subset \mathbb{R}^n$ be a cube, $f \in L^1(Q_0)$, and $\tilde{f}(x, t) := f(x)$ its trivial extension to $(x, t) \in \tilde{Q}_0 := Q_0 \times [0, l(Q_0)) \subset \mathbb{R}^{n+1}$. Then $f \in JN_p(Q_0)$ if and only if $\tilde{f} \in JN_p(\tilde{Q}_0)$, and*

$$2^{-1/p} \|f\|_{JN_p(Q_0)} l(Q_0)^{1/p} \leq \|\tilde{f}\|_{JN_p(\tilde{Q}_0)} \leq \|f\|_{JN_p(Q_0)} l(Q_0)^{1/p}.$$

For the proof we refer to [4, Proposition 4.1]. In conclusion we know that

$$L^p(Q_0) \subsetneq JN_p(Q_0) \subsetneq L^{p,\infty}(Q_0).$$

This raises many questions about the properties of JN_p functions. For example it is clear that for nonnegative L^p functions the equivalence

$$f^{1/p} \in L^p(Q_0) \iff f^{1/q} \in L^q(Q_0) \quad (2.6)$$

holds for all $p, q \in [1, \infty[$. This also holds for weak L^p spaces.

$$f^{1/p} \in L^{p,\infty}(Q_0) \iff f^{1/q} \in L^{q,\infty}(Q_0).$$

Our example in Section 3 proves that JN_p spaces do not have the same property. However there is a weaker result.

2.7. Proposition. *If $f : Q_0 \rightarrow \mathbb{R}$ is a nonnegative function, then*

$$f^{1/p} \in JN_p(Q_0) \implies f^{1/q} \in JN_q(Q_0)$$

whenever $1 < p \leq q < \infty$.

Proof. We shall prove this result by using the $JN_{p,inf}$ -norm. Since it is equivalent with JN_p , the theorem follows. Firstly we notice that $f^{1/q} \in L^1(Q_0)$, because $f^{1/q} \leq f^{1/p}$ whenever $f \geq 1$, $f^{1/p} \in L^1(Q_0)$ by assumption, and Q_0 is a bounded cube so it has finite measure. We notice that for any $c_i \in \mathbb{R}$

$$\begin{aligned} \left(\inf_{c'_i \in \mathbb{R}} \int_{Q_i} |f^{1/q} - c'_i| \right)^q &\leq \left(\int_{Q_i} |f^{1/q} - |c_i|^{p/q}| \right)^q \\ &\leq \left(\int_{Q_i} |f^{1/q} - |c_i|^{p/q}|^{q/p} \right)^p \\ &\leq \left(\int_{Q_i} |f^{1/p} - c_i| \right)^p. \end{aligned}$$

Here we first used Hölder's inequality and then the well known inequality $a^r + b^r \leq (a + b)^r$, when $a, b \geq 0$ and $r \geq 1$. Also $|f^{1/p} - |c_i|| \leq |f^{1/p} - c_i|$. By taking the infimum over c_i , this implies that

$$|Q_i| \left(\inf_{c'_i \in \mathbb{R}} \int_{Q_i} |f^{1/q} - c'_i| \right)^q \leq |Q_i| \left(\inf_{c_i \in \mathbb{R}} \int_{Q_i} |f^{1/p} - c_i| \right)^p$$

$$\sup_{Q_i \subset Q_0} \sum_{i=1}^{\infty} |Q_i| \left(\inf_{c'_i \in \mathbb{R}} \int_{Q_i} |f^{1/q} - c'_i| \right)^q \leq \sup_{Q_i \subset Q_0} \sum_{i=1}^{\infty} |Q_i| \left(\inf_{c_i \in \mathbb{R}} \int_{Q_i} |f^{1/p} - c_i| \right)^p$$

$$\|f^{1/q}\|_{JN_{q,inf}}^q \leq \|f^{1/p}\|_{JN_{p,inf}}^p$$

This completes the proof. \square

The implication in Proposition 2.7 does not hold in the other direction. In general

$$f^{1/q} \in JN_q(Q_0) \not\Rightarrow f^{1/p} \in JN_p(Q_0) \quad (2.8)$$

when $1 < p < q < \infty$. The proof is in the next section.

3. AN EXAMPLE THAT DISTINGUISHES JN_p SPACES WITH DIFFERENT PARAMETERS p

In this section we construct a function f that distinguishes JN_p spaces with different p in the sense that $f^{1/p} \notin JN_p$ but $f^{1/q} \in JN_q$, where $q > p$. Clearly we need to have $f^{1/q} \in JN_q \setminus L^q$, because if $f^{1/q} \in L^q$, then it follows from (2.6) that $f^{1/p} \in L^p \subset JN_p$. This is why the example is influenced by the function $g \in JN_p \setminus L^p$ given in [4]. Since the spaces L^p and JN_p coincide for monotone functions [4, Theorem 2.1], all examples in $JN_p \setminus L^p$ must have a similar highly oscillatory structure.

In the case of g , we have $g \in JN_p$ and also $g^{p/q} \in JN_q$ for any $q > 1$. More specifically this follows because g^p does not depend on p . Therefore it is impossible for that function to separate JN_p spaces with different parameters p . The function had to be modified substantially to get this example which does prove the result (2.8). It is enough to construct this function in the interval $[0, 1]$. Then we can make a change of variable to get a similar function in an arbitrary interval.

3.1. Theorem. *Let $Q_0 = [0, 1] \subset \mathbb{R}$ and let $1 < p < \infty$. Then there exists a nonnegative function f such that $f^{1/q} \notin JN_q(Q_0)$ whenever $1 < q \leq p$, and $f^{1/q} \in JN_q(Q_0)$ for every $q > p$.*

First we define a nonnegative function $u = f^{1/p}$. Then we show that $u \notin JN_p(Q_0)$, which together with Proposition 2.7 implies that $f^{1/q} \notin JN_q(Q_0)$ whenever $1 < q \leq p$. Finally we show that $u^{p/q} = f^{1/q} \in JN_q(Q_0)$ for all $q > p$.

The function u consists of "towers" of width l_i where $i \in \mathbb{Z}_+$. Each tower has an inclined roof with its left side located at height a_i and the right side at height b_i . The roof of the tower is linear. On both sides of the tower, at distance d_i , there are two similar narrower towers of width l_{i+1} and height ranging from a_{i+1} to b_{i+1} . This construction continues indefinitely, with every tower at a given level i having two towers of the next level $i + 1$ on both side at distance d_i . So there is one tower on level 1, two towers on level 2, 4 towers on level 3 etc. See Figure 1 to get a better understanding of the function.

To give the exact definition let us use the following dyadic notation. For any dyadic interval $I \subset [0, \frac{1}{2})$ of length 2^{-i} , there is a corresponding interval $\hat{I} \subset Q_0$ of length l_i . We start the indexing from $[0, \frac{1}{2})$ instead of $[0, 1)$ for technical reasons. The locations of \hat{I} in Q_0 are defined recursively. For $I^1 := [0, \frac{1}{2})$, let \hat{I}^1 be the interval of length l_1 located in the center of Q_0 . If \hat{I} is already defined and I' is

the left (right) half of I , let \hat{I}' be the interval of length l_{i+1} positioned on the left (right) side of \hat{I} in such a way that $\text{dist}(\hat{I}', \hat{I}) = d_i$.

We denote the intervals as $\hat{I} = [t_I, t_I + l_i]$. For every interval \hat{I} we define the function

$$u^I(x) := \frac{b_i - a_i}{l_i}(x - t_I) + a_i$$

for every $x \in \hat{I}$. The function u^I is identically zero elsewhere. Note that u^I is supported in \hat{I} , not in I . Finally we define the function u as the sum of all these functions u^I :

$$u(x) := \sum_{\substack{I \subset [0, \frac{1}{2}] \\ I \text{ dyadic}}} u^I(x)$$

for every $x \in Q_0$.




FIGURE 1. First three generations ($i = 1, 2, 3$) in the construction of u (with $p = 2$). δ_1 (resp. D_1) is the distance from $[0, \frac{1}{2}]$ to the nearest (resp. farthest) \hat{I} after infinitely many iterations of the construction. Notice that $a_1 = 0$ by definition.

The heights, widths and distances are defined as

$$\begin{aligned} l_i &= 2^{-i^2-i}, \\ d_i &= 2^{-i^2-2i}, \\ a_i &= (1 - i^{-1/p})2^{i^2/p}, \text{ and} \\ b_i &= (1 + i^{-1/p})2^{i^2/p} \end{aligned} \tag{3.2}$$

for every $i \geq 1$. From these definitions it follows that the intervals \hat{I} are disjoint. Indeed given an interval of length l_i , its distance to any other interval \hat{I}' is at least

$$\begin{aligned}
\delta_i &:= d_i - d_{i+1} - l_{i+2} - d_{i+2} - l_{i+3} - \dots = d_i - \sum_{j=1}^{\infty} d_{i+j} - \sum_{j=2}^{\infty} l_{i+j} \\
&= 2^{-i^2-2i} - \sum_{j=1}^{\infty} 2^{-(i+j)^2-2(i+j)} - \sum_{j=2}^{\infty} 2^{-(i+j)^2-i-j} \\
&\geq 2^{-i^2-2i} - 2^{-i^2-2i} \sum_{j=1}^{\infty} 2^{-j^2-2j} - 2^{-i^2-2i} \sum_{j=2}^{\infty} 2^{-j^2-j} \\
&\geq 2^{-i^2-2i} \left(1 - \sum_{j=1}^{\infty} 2^{-j-2} - \sum_{j=2}^{\infty} 2^{-j-1} \right) \\
&= \left(1 - \frac{1}{4} - \frac{1}{4} \right) 2^{-i^2-2i} = \frac{1}{2} d_i.
\end{aligned}$$

This means that the functions u^I have disjoint supports. Similarly we can compute a bound for D_i , the largest distance from an interval \hat{I} at level i to any of its descendants:

$$\begin{aligned}
D_i &:= d_i + l_{i+1} + d_{i+1} + l_{i+2} + d_{i+2} + l_{i+3} + \dots = d_i + \sum_{j=1}^{\infty} d_{i+j} + \sum_{j=1}^{\infty} l_{i+j} \\
&= 2^{-i^2-2i} + \sum_{j=1}^{\infty} 2^{-(i+j)^2-2(i+j)} + \sum_{j=1}^{\infty} 2^{-(i+j)^2-i-j} \\
&\leq 2^{-i^2-2i} + 2^{-i^2-2i} \sum_{j=1}^{\infty} 2^{-j^2-2j} + 2^{-i^2-2i} \sum_{j=1}^{\infty} 2^{-j^2-j} \\
&\leq 2^{-i^2-2i} \left(1 + \sum_{j=1}^{\infty} 2^{-j-2} + \sum_{j=1}^{\infty} 2^{-j-1} \right) \\
&= \left(1 + \frac{1}{4} + \frac{1}{2} \right) 2^{-i^2-2i} \leq 2d_i.
\end{aligned}$$

From this it follows that the support of u is contained within an interval of length $l_1 + 2D_1 \leq \frac{1}{4} + 4 \cdot \frac{1}{8} \leq 1$. This means that the function is indeed supported in $[0, 1]$. It is also useful to notice that $b_i = (1 + i^{-1/p})2^{i^2/p} \leq 2 \cdot 2^{i^2/p}$ for every $i \geq 1$. We summarize the results so far into the following lemma.

3.3. Lemma. *For every $i \geq 1$ we have*

$$\begin{aligned}
\frac{1}{2} d_i &\leq \delta_i \leq d_i \leq D_i \leq 2d_i, \text{ and} \\
b_i^p &\leq 2^p \cdot 2^{i^2}.
\end{aligned}$$

3.4. Proposition. $u = f^{1/p} \notin JN_p(Q_0)$.

Proof. Because u is linear in the intervals \hat{I} , it has a constant slope within \hat{I} , which we denote by k_I . Then it is easy to see that

$$\int_{\hat{I}} |u - u_{\hat{I}}| = \frac{|\hat{I}|}{2} \cdot k_I \cdot \frac{|\hat{I}|}{2} = \frac{l_i^2}{4} \cdot \frac{b_i - a_i}{l_i}.$$

By plugging in the values of a_i and b_i we get

$$\int_{\hat{I}} |u - u_{\hat{I}}| = \frac{1}{4} \cdot 2 \cdot i^{-1/p} \cdot 2^{i^2/p}.$$

This finally implies that

$$\begin{aligned} \|u\|_{JN_p(Q_0)}^p &\geq \sum_{I \text{ dyadic}} |\hat{I}| \left(\int_{\hat{I}} |u - u_{\hat{I}}| \right)^p \\ &= \sum_{i=1}^{\infty} 2^{i-1} \cdot 2^{-i^2-i} \left(\frac{1}{2} \cdot i^{-1/p} \cdot 2^{i^2/p} \right)^p \\ &= \sum_{i=1}^{\infty} 2^{-1-p} \cdot \frac{1}{i} = \infty, \end{aligned}$$

since the harmonic series diverges. \square

From now on we fix a number $q \in (p, \infty)$. Our goal is to prove that $u^{p/q} \in JN_q(Q_0)$. First we need to prove the following lemmas.

3.5. Lemma. *Let I be a dyadic interval of length 2^{-i} . Then*

$$\int_I u^{p/q} \leq c 2^{-i^2/q' - i},$$

where $q' = \frac{q}{q-1}$ is the Hölder conjugate of q . Also

$$\sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{I'} u^{p/q} \leq c 2^{-(i+1)^2/q' - i}.$$

In particular $u^{p/q} \in L^1(Q_0)$. Here the constants c depend only on p and q .

Proof. Because $u \leq b_i$ in \hat{I} , we have

$$\int_{\hat{I}} u^{p/q} \leq |\hat{I}| \cdot b_i^{p/q} \leq l_i \left(c 2^{i^2} \right)^{1/q} = 2^{-i^2-i} \cdot c \cdot 2^{i^2/q} = c 2^{-i^2/q' - i}.$$

Additionally

$$\begin{aligned} \sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{I'} u^{p/q} &\leq \sum_{j=1}^{\infty} 2^j \cdot c 2^{-(i+j)^2/q' - (i+j)} = c \sum_{j=1}^{\infty} 2^{-(i+j)^2/q' - i} \\ &\leq c 2^{(-i^2-2i)/q' - i} \sum_{j=1}^{\infty} 2^{-j^2/q'} \\ &\leq c 2^{-(i+1)^2/q' - i}. \end{aligned}$$

The fact that $u^{p/q} \in L^1$ follows from the previous result by having $I = [0, \frac{1}{2})$, because then

$$\|u^{p/q}\|_{L^1(Q_0)} = \int_{\hat{I}} u^{p/q} + \sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{I'} u^{p/q} \leq c2^{-1/q'-1} + c2^{-2^2/q'-2} < \infty.$$

□

The following lemma is easy to prove by finding the local minima of the function $h(x) = 2x - (1+x)^{p/q} + (1-x)^{p/q}$, where $0 \leq x \leq 1$, or by using the Taylor approximation $(1+x)^r \approx 1+rx$, when $|x|$ is small.

3.6. Lemma. *Let $1 < p < q < \infty$ and $i \in \mathbb{Z}_+$. Then*

$$\left(1 + i^{-1/p}\right)^{p/q} - \left(1 - i^{-1/p}\right)^{p/q} \leq 2i^{-1/p}.$$

To prove that $u^{p/q} \in JN_q(Q_0)$, we let \mathcal{J} be a countable collection of disjoint subintervals of Q_0 . For $J \in \mathcal{J}$ we define

$$F(J) := |J| \left(\int_J \left| u^{p/q} - (u^{p/q})_J \right| \right)^q.$$

It is clear that if J does not intersect any of the intervals \hat{I} , then $F(J) = 0$. Also if J does intersect some of the intervals, then there is a unique widest interval I_J , such that J intersects \hat{I}_J . We look into four separate cases how J relates to I_J .

3.7. Definition.

- 1) J is contained, if $|J \setminus \hat{I}_J| = 0$ i.e. $J \subset \hat{I}_J$
- 2) J is short, if $0 < |J \setminus \hat{I}_J| \leq \delta_I$
- 3) J is medium, if $\delta_I < |J \setminus \hat{I}_J| \leq 2D_I$
- 4) J is long, if $|J \setminus \hat{I}_J| > 2D_I$.

3.8. Lemma. *Assume that J intersects some of the intervals \hat{I} and $I = I_J$. Then*

$$F(J) \leq c \left[|J|^{1-q} \left(\int_{J \cap \hat{I}} u^{p/q} \right)^q + \min(|J \setminus \hat{I}|, |J \cap \hat{I}|) b_I^p + |J \cap \hat{I}| i^{-q/p} \cdot 2^{i^2} \right],$$

where $c = c(q)$ and $|I| = 2^{-i}$.

Proof. It is clear that

$$\left(u^{p/q}\right)_J = \frac{1}{|J|} \int_{J \cap \hat{I}} u^{p/q} + \frac{1}{|J|} \int_{J \setminus \hat{I}} u^{p/q}$$

and so we get from the triangle inequality

$$\begin{aligned} F(J) &= |J|^{1-q} \left(\int_{J \cap \hat{I}} \left| u^{p/q} - \left(u^{p/q}\right)_J \right| + \int_{J \setminus \hat{I}} \left| u^{p/q} - \left(u^{p/q}\right)_J \right| \right)^q \\ &\leq |J|^{1-q} \left(\int_{J \cap \hat{I}} \left| u^{p/q} - \frac{1}{|J|} \int_{J \cap \hat{I}} u^{p/q} \right| + \frac{|J \cap \hat{I}|}{|J|} \int_{J \setminus \hat{I}} u^{p/q} \right. \\ &\quad \left. + \int_{J \setminus \hat{I}} u^{p/q} + \frac{|J \setminus \hat{I}|}{|J|} \int_{J \cap \hat{I}} u^{p/q} + \frac{|J \setminus \hat{I}|}{|J|} \int_{J \setminus \hat{I}} u^{p/q} \right)^q. \end{aligned}$$

Define $\alpha := \frac{1}{|J|} \int_{J \cap \hat{I}} u^{p/q}$. If $\alpha \leq (u(x))^{p/q}$ for all $x \in J \cap \hat{I}$, then $F(J)$ is bounded by

$$\begin{aligned}
& |J|^{1-q} \left(\int_{J \cap \hat{I}} u^{p/q} - \frac{|J \cap \hat{I}|}{|J|} \int_{J \cap \hat{I}} u^{p/q} + 2 \int_{J \setminus \hat{I}} u^{p/q} + \frac{|J \setminus \hat{I}|}{|J|} \int_{J \cap \hat{I}} u^{p/q} \right)^q \\
&= 2^q |J|^{1-q} \left(\int_{J \setminus \hat{I}} u^{p/q} + \frac{|J \setminus \hat{I}|}{|J|} \int_{J \cap \hat{I}} u^{p/q} \right)^q \\
&\leq c \left[|J|^{1-q} \left(\int_{J \setminus \hat{I}} u^{p/q} \right)^q + |J|^{1-q} \frac{|J \setminus \hat{I}|^q}{|J|^q} |J \cap \hat{I}|^q \left(b_I^{p/q} \right)^q \right] \\
&\leq c \left[|J|^{1-q} \left(\int_{J \setminus \hat{I}} u^{p/q} \right)^q + \min(|J \setminus \hat{I}|, |J \cap \hat{I}|) b_I^p \right].
\end{aligned}$$

On the other hand if $\alpha \geq \inf\{(u(x))^{p/q} : x \in J \cap \hat{I}\}$, then we know that $a_i^{p/q} \leq \alpha \leq b_i^{p/q}$ and $a_i^{p/q} \leq u^{p/q} \leq b_i^{p/q}$ in the interval $J \cap \hat{I}$ and so

$$\begin{aligned}
\int_{J \cap \hat{I}} |u^{p/q} - \alpha| &\leq \int_{J \cap \hat{I}} b_I^{p/q} - a_I^{p/q} = |J \cap \hat{I}| \left(b_I^{p/q} - a_I^{p/q} \right) \\
&= |J \cap \hat{I}| \left(\left(1 + i^{-1/p} \right)^{p/q} - \left(1 - i^{-1/p} \right)^{p/q} \right) \left(2^{i^2/p} \right)^{p/q} \\
&\leq |J \cap \hat{I}| \cdot 2 \cdot i^{-1/p} \cdot 2^{i^2/q}.
\end{aligned}$$

Here we used Lemma 3.6 to get the final inequality. Then we get

$$\begin{aligned}
F(J) &\leq |J|^{1-q} \left(|J \cap \hat{I}| \cdot 2 \cdot i^{-1/p} \cdot 2^{i^2/q} + 2 \int_{J \setminus \hat{I}} u^{p/q} + \frac{|J \setminus \hat{I}|}{|J|} \int_{J \cap \hat{I}} u^{p/q} \right)^q \\
&\leq c |J|^{1-q} |J \cap \hat{I}|^q \cdot i^{-q/p} \cdot 2^{i^2} + c |J|^{1-q} \left(\int_{J \setminus \hat{I}} u^{p/q} \right)^q \\
&\quad + c |J|^{1-q} \frac{|J \setminus \hat{I}|^q}{|J|^q} |J \cap \hat{I}|^q \left(b_I^{p/q} \right)^q \\
&\leq c \left[|J|^{1-q} \left(\int_{J \setminus \hat{I}} u^{p/q} \right)^q + \min(|J \setminus \hat{I}|, |J \cap \hat{I}|) b_I^p + |J \cap \hat{I}| i^{-q/p} \cdot 2^{i^2} \right].
\end{aligned}$$

Hence in all cases the lemma is true. \square

3.9. Proposition. *If J is contained, then*

$$F(J) \leq c(q) |J| \cdot i^{-q/p} \cdot 2^{i^2}$$

and consequently

$$\sum_{\substack{J \in \mathcal{J} \\ J \text{ contained}}} F(J) \leq c_1(p, q) < \infty.$$

Proof. In this case the first two terms in Lemma 3.8 are just zero, and so we get the bound for $F(J)$. Then it follows that

$$\begin{aligned}
\sum_{\substack{J \in \mathcal{J} \\ J \text{ contained}}} F(J) &= \sum_{I \text{ dyadic}} \sum_{\substack{J \in \mathcal{J} \\ J \subset \hat{I}}} F(J) \leq \sum_{i=1}^{\infty} \sum_{\substack{I \text{ dyadic} \\ |I|=2^{-i}}} \sum_{\substack{J \in \mathcal{J} \\ J \subset \hat{I}}} c|J| \cdot i^{-q/p} \cdot 2^{i^2} \\
&\leq \sum_{i=1}^{\infty} \sum_{\substack{I \text{ dyadic} \\ |I|=2^{-i}}} c \cdot l_i \cdot i^{-q/p} \cdot 2^{i^2} \\
&= c \sum_{i=1}^{\infty} 2^{i-1} \cdot 2^{-i^2-i} \cdot i^{-q/p} \cdot 2^{i^2} \\
&= c \sum_{i=1}^{\infty} i^{-q/p} = c_1(p, q) < \infty.
\end{aligned}$$

This is an over-harmonic series, so it converges. \square

We move on to the other cases where $|J \cap \hat{I}_J| > 0$. In these cases J must intersect the boundary of the interval \hat{I}_J and so for each I there are at most two intervals J such that $I = I_J$ and $|J \cap \hat{I}| > 0$.

3.10. Proposition. *If J is short, then*

$$F(J) \leq c(p, q) \left(2^{-2i} + 2^{-i} \cdot i^{-q/p} \right)$$

and consequently

$$\sum_{\substack{J \in \mathcal{J} \\ J \text{ short}}} F(J) \leq c_2(p, q) < \infty.$$

Proof. In this case u is zero in $J \setminus \hat{I}$ and so the first term in Lemma 3.8 is just zero. Then we get

$$\begin{aligned}
F(J) &\leq c \left[\min(|J \setminus \hat{I}|, |J \cap \hat{I}|) b_I^p + |J \cap \hat{I}| \cdot i^{-q/p} \cdot 2^{i^2} \right] \\
&\leq c \left[\delta_I b_I^p + l_I \cdot i^{-q/p} \cdot 2^{i^2} \right] \\
&\leq c \left(2^{-i^2-2i} \cdot 2^p \cdot 2^{i^2} + 2^{-i^2-i} \cdot i^{-q/p} \cdot 2^{i^2} \right) \\
&\leq c(p, q) \left(2^{-2i} + 2^{-i} \cdot i^{-q/p} \right).
\end{aligned}$$

Using this bound we get

$$\begin{aligned}
\sum_{\substack{J \in \mathcal{J} \\ J \text{ short}}} F(J) &\leq \sum_{i=1}^{\infty} 2^{i-1} \cdot 2 \cdot c \left(2^{-2i} + 2^{-i} \cdot i^{-q/p} \right) \\
&= \sum_{i=1}^{\infty} c 2^{-i} + \sum_{i=1}^{\infty} c i^{-q/p} = c_2(p, q) < \infty.
\end{aligned}$$

\square

3.11. Proposition. *If J is medium, then*

$$F(J) \leq c(p, q) \left(2^{-iq} + 2^{-2i} + 2^{-i} \cdot i^{-q/p} \right)$$

and consequently

$$\sum_{\substack{J \in \mathcal{J} \\ J \text{ medium}}} F(J) \leq c_3(p, q) < \infty.$$

Proof. In this case the bounds for the second and third term in Lemma 3.8 are essentially the same as in Proposition 3.10 because $|J \setminus \hat{I}| \leq 2D_I \leq 4d_I$. For the first term we use Lemma 3.5 to get

$$\int_{J \setminus \hat{I}} u^{p/q} \leq \sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{\hat{I}'} u^{p/q} \leq c 2^{-(i+1)^2/q' - i}.$$

Then we use the fact that $|J| \geq |J \setminus \hat{I}| > \delta_I$ to get

$$\begin{aligned} |J|^{1-q} \left(\int_{J \setminus \hat{I}} u^{p/q} \right)^q &\leq \delta_I^{1-q} \left(c 2^{-(i+1)^2/q' - i} \right)^q \\ &\leq c \left(2^{-i^2 - 2i} \right)^{1-q} \cdot 2^{-(i+1)^2(q-1)} \cdot 2^{-iq} \\ &\leq c 2^{-iq} \end{aligned}$$

and so we have the bound for $F(J)$. This implies that

$$\begin{aligned} \sum_{\substack{J \in \mathcal{J} \\ J \text{ medium}}} F(J) &\leq \sum_{i=1}^{\infty} 2^{i-1} \cdot 2c \left(2^{-iq} + 2^{-2i} + 2^{-i} \cdot i^{-q/p} \right) \\ &= \sum_{i=1}^{\infty} c 2^{(1-q)i} + \sum_{i=1}^{\infty} c 2^{-i} + \sum_{i=1}^{\infty} c i^{-q/p} \\ &= c_3(p, q) < \infty. \end{aligned}$$

□

3.12. Proposition. *If J is long, then*

$$F(J) \leq c(p, q) |I_J|$$

and the corresponding intervals I form a Carleson family, in the sense that

$$\sum_{\substack{I \text{ dyadic} \\ \exists J \text{ long s.t. } I=I_J}} |I| \leq 1.$$

Therefore

$$\sum_{\substack{J \in \mathcal{J} \\ J \text{ long}}} F(J) \leq c_4(p, q) < \infty.$$

Proof. The proof of the corresponding intervals forming a Carleson family is essentially the same as the proof of [4, Lemma 3.9], which presents a function with a similar construction.

To get the bound for $F(J)$ we use Lemma 3.8. First we notice that the bound for the first term is essentially the same as in Proposition 3.11, because $|J| \geq |J \setminus \hat{I}| > 2D_I \geq 2d_I$. For the second and third term we use the fact that $|J \cap \hat{I}| \leq |\hat{I}| = l_I$. Then we get

$$F(J) \leq c \left[c 2^{-iq} + l_I b_I^p + l_I \cdot i^{-q/p} \cdot 2^{i^2} \right] \leq c \left[2^{-iq} + 2^{-i^2 - i} \cdot c 2^{i^2} + 2^{-i^2 - i} \cdot 2^{i^2} \right]$$

$$\leq c [2^{-iq} + 2^{-i}] \leq c(p, q)|I|.$$

Let us say that a dyadic interval I is long, if it is of the form I_J for some long J . Then we conclude that

$$\sum_{\substack{J \in \mathcal{J} \\ J \text{ long}}} F(J) = \sum_{I \text{ long}} \sum_{\substack{J \text{ long} \\ I=I_J}} F(J) \leq \sum_{I \text{ long}} 2c|I| \leq 2c = c_4(p, q) < \infty. \quad \square$$

Now we are ready to prove the following proposition, which also concludes the proof of Theorem 3.1.

3.13. Proposition. $u^{p/q} = f^{1/q} \in JN_q(Q_0)$.

Proof. This follows from Propositions 3.9, 3.10, 3.11 and 3.12. If we take any partition \mathcal{J} of Q_0 into disjoint subintervals $J \in \mathcal{J}$, then

$$\begin{aligned} \sum_{J \in \mathcal{J}} |J| \left(\int_J |u^{p/q} - (u^{p/q})_J| \right)^q &= \sum_{\substack{J \in \mathcal{J} \\ J \text{ contained}}} F(J) + \sum_{\substack{J \in \mathcal{J} \\ J \text{ short}}} F(J) \\ &+ \sum_{\substack{J \in \mathcal{J} \\ J \text{ medium}}} F(J) + \sum_{\substack{J \in \mathcal{J} \\ J \text{ long}}} F(J) \\ &\leq c_1 + c_2 + c_3 + c_4 < \infty. \end{aligned}$$

As the estimate does not depend on \mathcal{J} , we can conclude that $\|u^{p/q}\|_{JN_q(Q_0)}^q < \infty$ and therefore $u^{p/q} \in JN_q(Q_0)$. \square

The function u can be extended from the one-dimensional case $Q_0 \subset \mathbb{R}$ into the multidimensional case $Q_0 \subset \mathbb{R}^n$ with an arbitrary n , by repeatedly using Proposition 2.5.

3.14. Theorem. *Let $Q_0 \subset \mathbb{R}^n$ be a bounded cube and let $1 < p < \infty$. Then there exists a nonnegative function f such that $f^{1/q} \notin JN_q(Q_0)$ whenever $1 < q \leq p$, and $f^{1/q} \in JN_q(Q_0)$ for every $q > p$.*

4. THE VANISHING SUBSPACE VJN_p

The vanishing subspace VJN_p of JN_p is defined analogously to the space of vanishing mean oscillation VMO which is a subspace of BMO .

4.1. Definition (VJN_p). Let $1 < p < \infty$ and $Q_0 \subset \mathbb{R}^n$ a bounded cube. Then $f \in VJN_p(Q_0)$ if $f \in JN_p(Q_0)$ and

$$\lim_{a \rightarrow 0} \sup_{\substack{Q_i \subset Q_0 \\ l(Q_i) \leq a}} \sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p = 0$$

where the supremum is taken over all collections of pairwise disjoint subcubes $Q_i \subset Q_0$ such that the side length of each Q_i is at most a .

The following theorem is a characterization of $VJN_p(Q_0)$.

4.2. Theorem.

$$VJN_p(Q_0) = \overline{C^\infty(Q_0)}^{JN_p(Q_0)}$$

where $C^\infty(Q_0)$ is the set of smooth functions in \mathbb{R}^n that have been restricted to Q_0 .

For the proof we refer to [16, Theorem 5.3]. To be precise, that theorem applies to the so-called John-Nirenberg-Campanato spaces $JN_{(p,q,s)\alpha}(X)$ that were studied by Tao et al. [15]. Here $p \in (1, \infty)$, $q \in [1, \infty)$, $\alpha \in [0, \infty)$, s is a nonnegative integer and X is either \mathbb{R}^n or a bounded cube $Q_0 \subset \mathbb{R}^n$. Brudnyi studied the vanishing subspace of this John-Nirenberg-Campanato space, denoted as $VJN_{(p,q,s)\alpha}(X)$ [3].

By choosing $q = 1$, $s = \alpha = 0$ and $X = Q_0$, the John-Nirenberg-Campanato space $JN_{(p,q,s)\alpha}(X)$ becomes the John-Nirenberg space, $JN_{(p,1,0)_0}(Q_0) = JN_p(Q_0)$. The same holds for the vanishing subspaces, i.e. $VJN_p(Q_0) = VJN_{(p,1,0)_0}(Q_0)$. So by choosing the parameters appropriately, [16, Theorem 5.3] simplifies into the form in Theorem 4.2.

It follows directly from Theorem 4.2 that $L^p(Q_0) \subset VJN_p(Q_0)$, by density of C^∞ in L^p and by Equation (2.3). Thus it is immediate that

$$L^p(Q_0) \subseteq VJN_p(Q_0) \subseteq JN_p(Q_0).$$

The obvious question is now if these inclusions are strict. We answer this question by examining two functions. The counterexample presented in [4] is in $JN_p \setminus VJN_p$. By modifying the function, we get another function which is in $VJN_p \setminus L^p$.

4.3. Proposition. *Let $1 < p < \infty$ and $Q_0 \subset \mathbb{R}$ a bounded interval. Then there is a function $g : Q_0 \rightarrow \mathbb{R}$ such that $g \in JN_p(Q_0) \setminus VJN_p(Q_0)$.*

Proof. Let g be the function defined in [4, Section 3]. In the paper it is already shown that $g \in JN_p(Q_0)$. The function is similar to the function in Section 3 of this paper. The key difference is that in this case the function is constant in every interval \hat{I} . In other words for this function we have $a_i = b_i = 2^{i^2/p}$ in (3.2). To show that $g \notin VJN_p(Q_0)$, let $a > 0$ be arbitrarily small. Then there is a positive integer m such that $2l_m \leq a$. For any integer $k \geq m$ we notice that $d_{k-1} \geq l_k \geq D_k$. This means that for each interval \hat{I} of length l_k we can choose an interval J of length $2l_k$ such that it covers \hat{I} and all of its descendants on one side and the interval J doesn't intersect any other interval \hat{I}' . Then we get

$$|J| \left(\int_J |g - g_J| \right)^p \geq |J|^{1-p} \left(\int_{\hat{I}} |g - g_J| \right)^p = (2l_k)^{1-p} (l_k(h_k - g_J))^p.$$

On the other hand we know that

$$\begin{aligned} g_J &= \int_J g = \frac{1}{2l_k} \left(h_k l_k + \sum_{i=1}^{\infty} 2^{i-1} h_{k+i} l_{k+i} \right) \\ &= h_k \left(\frac{1}{2} + \frac{1}{4} \sum_{i=1}^{\infty} 2^{(\frac{1}{p}-1)(i^2+2ki)} \right) \\ &\leq h_k \left(\frac{1}{2} + \frac{1}{4} 2^{(\frac{1}{p}-1)2k} \sum_{i=1}^{\infty} 2^{(\frac{1}{p}-1)i^2} \right) \\ &\leq \frac{3}{4} h_k \end{aligned}$$

at least when k is large enough. Thus for $k \geq m$ large enough, we have

$$\begin{aligned} |J| \left(\int_J |g - g_J| \right)^p &\geq (2l_k)^{1-p} (l_k(h_k - g_J))^p \\ &\geq 2^{1-p} l_k \left(\frac{h_k}{4} \right)^p \end{aligned}$$

$$\begin{aligned}
&= 2^{1-3p} \cdot 2^{-k^2-k-1/4} \cdot 2^{k^2} \\
&= 2^{\frac{3}{4}-3p} \cdot 2^{-k}.
\end{aligned}$$

Since there are 2^k many intervals \hat{I} of length l_k , this means that

$$\sup_{\substack{Q_i \subset Q_0 \\ l(Q_i) \leq a}} \sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |g - g_{Q_i}| \right)^p \geq 2^k \cdot 2^{\frac{3}{4}-3p} \cdot 2^{-k} = 2^{\frac{3}{4}-3p}$$

and this is true for all $a > 0$, so especially the limit as $a \rightarrow 0$ is positive. So by definition $g \notin VJN_p(Q_0)$. \square

4.4. Proposition. *Let $1 < p < \infty$ and $Q_0 \subset \mathbb{R}$ a bounded interval. Then there is a function $g_0 : Q_0 \rightarrow \mathbb{R}$ such that $g_0 \in VJN_p(Q_0) \setminus L^p(Q_0)$.*

Proof. Let us construct g_0 by modifying the function g in the proof of Proposition 4.3. For g_0 we set the height of every tower, with $i \geq 1$, as $h_i = \left(\frac{2^{i^2}}{i}\right)^{1/p}$ instead of $2^{i^2/p}$. The other parameters l_i and d_i are the same as in g . Then the proof of $g_0 \in JN_p(Q_0)$ follows the same steps as the proof of $g \in JN_p(Q_0)$ in [4]. Also we notice that

$$\|g_0\|_{L^p}^p = \int_{Q_0} g_0^p \geq \sum_{i=1}^{\infty} 2^i h_i^p l_i = \sum_{i=1}^{\infty} 2^i \cdot \frac{2^{i^2}}{i} \cdot 2^{-i^2-i-1/4} = 2^{-1/4} \sum_{i=1}^{\infty} \frac{1}{i} = \infty,$$

thus $g_0 \notin L^p(Q_0)$.

To prove that $g_0 \in VJN_p$, let $a > 0$ be a small enough number. Then there is some positive integer k such that $\delta_k \geq a > \delta_{k+1}$. Let \mathcal{J} be a countable collection of disjoint intervals such that $|J| \leq a$ for every $J \in \mathcal{J}$. Define

$$F(J) := |J| \left(\int_J |g_0 - (g_0)_J| \right)^p.$$

Similar to the function u in Section 3, it is clear that if J does not intersect any of the intervals \hat{I} , then $F(J) = 0$. If J does intersect some intervals, then there is a unique widest interval I_J , such that J intersects \hat{I}_J . Also if J does intersect an interval \hat{I} , if it does not intersect the boundary of the interval, then $F(J) = 0$. Therefore we assume that J intersects the boundary of \hat{I}_J . Then there are at most 2 intervals J for every I such that $I = I_J$. Also we can categorize the intervals J as short, medium or long, depending on the length of $J \setminus \hat{I}_J$, the same way as in Definition 3.7.

We have the following bound for $F(J)$:

$$F(J) \leq c \left[|J|^{1-p} \left(\int_{J \setminus \hat{I}} g_0 \right)^p + \min(|J \setminus \hat{I}|, |J \cap \hat{I}|) h_I^p \right], \quad (4.5)$$

where $c = c(p)$ and $I = I_J$. The proof is the same as for [4, Lemma 3.6].

If $I = I_J$ and $|I| = 2^{-i}$ with $i \leq k$, then the fact that $|J \setminus \hat{I}| \leq |J| \leq a \leq \delta_k \leq \delta_i$ implies that J is short. Hence in this case $F(J) \leq c \min(|J \setminus \hat{I}|, |J \cap \hat{I}|) h_I^p \leq c d_k h_k^p$. The total number of dyadic intervals I that are not shorter than 2^{-k} is

$1 + 2 + 4 + \dots + 2^k \leq 2^{k+1}$. Therefore

$$\sum_{\substack{J \in \mathcal{J} \\ |I_J| \geq 2^{-k}}} F(J) \leq 2^{k+1} \cdot 2cd_k h_k^p = c2^k \cdot 2^{-k^2-2k} \frac{2^{k^2}}{k} \leq c2^{-k}.$$

Since $k \rightarrow \infty$ when $a \rightarrow 0$, we notice that the limit of this sum is 0.

Assume from now on that $I = I_J$ and $i > k$. If J is short, then $F(J) \leq cd_i h_i^p \leq c2^{-i^2-2i} 2^{i^2} = c2^{-2i}$. This implies that

$$\sum_{\substack{J \in \mathcal{J} \\ |I_J| < 2^{-k} \\ J \text{ short}}} F(J) \leq \sum_{i=k+1}^{\infty} 2^i \cdot 2c2^{-2i} = c \sum_{i=1}^{\infty} 2^{-k-i} = c2^{-k}$$

which tends to zero when $a \rightarrow 0$.

If J is medium, then for the second term in (4.5) we have essentially the same bound as in the short case. For the first term we have the bound

$$|J|^{1-p} \left(\int_{J \setminus \hat{I}} g_0 \right)^p \leq c(p)2^{-ip}.$$

For the proof we refer to [4, Lemma 3.8], because $g_0 \leq g$ pointwise. This implies that

$$\begin{aligned} \sum_{\substack{J \in \mathcal{J} \\ |I_J| < 2^{-k} \\ J \text{ medium}}} F(J) &\leq \sum_{i=k+1}^{\infty} 2^i \cdot 2c(2^{-2i} + 2^{-ip}) = c \sum_{i=1}^{\infty} 2^{-k-i} + 2^{(1-p)(k+i)} \\ &\leq c \left(2^{-k} + 2^{(1-p)k} \right). \end{aligned}$$

This also tends to zero when $a \rightarrow 0$.

Finally if J is long, we have the same Carleson property as in [4, Lemma 3.9]:

$$\sum_{\substack{I \text{ dyadic} \\ \exists J \text{ long s.t. } I=I_J}} |I| \leq 2.$$

The bound for the first term in (4.5) is the same as in the medium case:

$$|J|^{1-p} \left(\int_{J \setminus \hat{I}} g_0 \right)^p \leq c(p)2^{-ip} = c2^{(1-p)i} |I|.$$

For the second term we have the bound

$$\min(|J \setminus \hat{I}|, |J \cap \hat{I}|) h_I^p \leq l_i h_i^p = 2^{-i^2-i-1/4} \frac{2^{i^2}}{i} \leq \frac{|I|}{i}.$$

This implies that

$$\begin{aligned} \sum_{\substack{J \in \mathcal{J} \\ |I_J| < 2^{-k} \\ J \text{ long}}} F(J) &= \sum_{\substack{I \text{ long} \\ |I| < 2^{-k}}} \sum_{\substack{J \text{ long} \\ I=I_J}} F(J) \leq \sum_{\substack{I \text{ long} \\ |I| < 2^{-k}}} 2c \left(2^{(1-p)i} |I| + \frac{|I|}{i} \right) \\ &\leq c \left(2^{(1-p)k} + \frac{1}{k} \right) \sum_{I \text{ long}} |I| \leq c \left(2^{(1-p)k} + \frac{1}{k} \right). \end{aligned}$$

This also tends to zero.

In conclusion

$$\begin{aligned} \sum_{J \in \mathcal{J}} |J| \left(\int_J |g_0 - (g_0)_J| \right)^p &= \sum_{\substack{J \in \mathcal{J} \\ |I_J| \geq 2^{-k}}} F(J) + \sum_{\substack{J \in \mathcal{J} \\ |I_J| < 2^{-k} \\ J \text{ short}}} F(J) \\ &+ \sum_{\substack{J \in \mathcal{J} \\ |I_J| < 2^{-k} \\ J \text{ medium}}} F(J) + \sum_{\substack{J \in \mathcal{J} \\ |I_J| < 2^{-k} \\ J \text{ long}}} F(J) \\ &\leq c2^{-k} + c2^{-k} + c2^{-k} + c2^{(1-p)k} + c2^{(1-p)k} + \frac{c}{k} \end{aligned}$$

and this holds for any admissible collection \mathcal{J} . This implies that

$$\lim_{a \rightarrow 0} \sup_{\substack{Q_i \subset Q_0 \\ l(Q_i) \leq a}} \sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |g_0 - (g_0)_{Q_i}| \right)^p \leq \lim_{k \rightarrow \infty} c \left(2^{-k} + 2^{(1-p)k} + \frac{1}{k} \right) = 0$$

and so by definition $g_0 \in VJN_p(Q_0)$. \square

Now we would like to extend this result to the multidimensional case. The following proposition is a counterpart of Proposition 2.5.

4.6. Proposition. *Let $Q_0 \subset \mathbb{R}^n$ be a bounded cube, $f \in L^1(Q_0)$, and $\tilde{f}(x, t) := f(x)$ its trivial extension to $(x, t) \in \tilde{Q}_0 := Q_0 \times [0, l(Q_0)) \subset \mathbb{R}^{n+1}$. Then $f \in VJN_p(Q_0)$ if and only if $\tilde{f} \in VJN_p(\tilde{Q}_0)$.*

Proof. The proof is very similar to the proof of [4, Proposition 4.1]. The only difference is that in the partition of Q_0 into subcubes we have the additional criterion that $l(Q_i) \leq a$ for every i and for some $a > 0$. Ultimately we get the result by taking the limit $a \rightarrow 0$. \square

By using Propositions 2.5 and 4.6 multiple times, we can extend both g and g_0 into the multidimensional cube $Q_0 \subset \mathbb{R}^n$ for any n , such that $g \in JN_p(Q_0) \setminus VJN_p(Q_0)$ and $g_0 \in VJN_p(Q_0) \setminus L^p(Q_0)$. In conclusion we have shown that for any bounded cube $Q_0 \subset \mathbb{R}^n$

$$L^p(Q_0) \subsetneq VJN_p(Q_0) \subsetneq JN_p(Q_0) \subsetneq L^{p, \infty}(Q_0).$$

5. THE SPACES JN_p , VJN_p AND CJN_p DEFINED ON \mathbb{R}^n

The definition of the space JN_p can be extended from the bounded cube Q_0 into the whole Euclidian space \mathbb{R}^n . In this section we deal with the entire space \mathbb{R}^n and always assume that $JN_p = JN_p(\mathbb{R}^n)$ and $VJN_p = VJN_p(\mathbb{R}^n)$ etc. unless otherwise specified.

5.1. Definition (JN_p). Let $1 < p < \infty$. Then $f \in JN_p$ if $f \in L^1_{loc}$ and there is a constant $K < \infty$ such that

$$\sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p \leq K^p$$

for all countable collections of pairwise disjoint cubes $(Q_i)_{i=1}^{\infty}$ in \mathbb{R}^n . We denote the smallest such number K by $\|f\|_{JN_p}$.

The embedding $JN_p(Q_0) \subset L^{p,\infty}(Q_0)$ in Theorem 2.4 has been proven in a number of ways. However all the proofs, that we know of, only consider JN_p on bounded sets. They don't address the question of whether this holds if we replace Q_0 with \mathbb{R}^n . Clearly any constant function $f = c$ is in the space $JN_p(\mathbb{R}^n)$ even though it is in $L^{p,\infty}(\mathbb{R}^n)$ only if $c = 0$. So let us define the space

$$L^{p,\infty}/\mathbb{R} := \{f \in L^1_{loc} : \exists c \in \mathbb{R} \text{ s.t. } f - c \in L^{p,\infty}\}.$$

Then for the sake of completeness we prove the following theorem.

5.2. Theorem. *Let $1 < p < \infty$ and $f \in JN_p$. Then $f \in L^{p,\infty}/\mathbb{R}$ and there is a constant b such that*

$$\|f - b\|_{L^{p,\infty}} \leq c\|f\|_{JN_p}$$

where $c = c(n, p)$.

Proof. First let $Q \subset \mathbb{R}^n$ be any bounded cube. It is clear that if we restrict f to this cube, then we have $f \in JN_p(Q)$ and therefore from Theorem 2.4 we get that $\|f - f_Q\|_{L^{p,\infty}(Q)} \leq c\|f\|_{JN_p(Q)}$. Further from Definition 5.1 it immediately follows that

$$\int_Q |f - f_Q| \leq |Q|^{-1/p} \|f\|_{JN_p}.$$

Now let us define a sequence of cubes $(Q_m)_{m=1}^\infty$ such that $|Q_m| = 2^m$ and the center of every cube is the origin. Then clearly we have $Q_1 \subset Q_2 \subset \dots$ and $\bigcup_{m=1}^\infty Q_m = \mathbb{R}^n$. Let us prove that the sequence of integral averages f_{Q_m} is a Cauchy sequence. First we notice that for the difference of two consecutive elements we have the following estimate:

$$\begin{aligned} |f_{Q_{i+1}} - f_{Q_i}| &\leq \frac{|Q_{i+1}|}{|Q_i|} \int_{Q_{i+1}} |f_{Q_{i+1}} - f| + \int_{Q_i} |f - f_{Q_i}| \\ &\leq 2|Q_{i+1}|^{-1/p} \|f\|_{JN_p} + |Q_i|^{-1/p} \|f\|_{JN_p} \\ &= (2^{1-1/p} + 1) \|f\|_{JN_p} 2^{-i/p}. \end{aligned}$$

Now let $\epsilon > 0$ and $N = N(\epsilon)$ an integer that will depend on ϵ . Let $m, k \geq N$ and let us assume that $k \geq m$. Then

$$\begin{aligned} |f_{Q_k} - f_{Q_m}| &\leq \sum_{i=m}^{k-1} |f_{Q_{i+1}} - f_{Q_i}| \leq \sum_{i=N}^{\infty} (2^{1-1/p} + 1) \|f\|_{JN_p} 2^{-i/p} \\ &= c(p) \|f\|_{JN_p} 2^{-N/p} < \epsilon \end{aligned}$$

when N is large enough. Thus f_{Q_m} is a Cauchy sequence and we can set $b := \lim_{m \rightarrow \infty} f_{Q_m}$. Then we have

$$\begin{aligned} |\{x \in \mathbb{R}^n : |f(x) - b| > t\}| &= \lim_{m \rightarrow \infty} |\{x \in Q_m : |f(x) - b| > t\}| \\ &\leq \lim_{m \rightarrow \infty} \left(\left| \left\{ x \in Q_m : |f(x) - f_{Q_m}| > \frac{t}{2} \right\} \right| \right. \\ &\quad \left. + \left| \left\{ x \in Q_m : |f_{Q_m} - b| > \frac{t}{2} \right\} \right| \right) \\ &\leq \lim_{m \rightarrow \infty} c \frac{\|f\|_{JN_p(Q_m)}^p}{\left(\frac{t}{2}\right)^p} \end{aligned}$$

$$\leq c \frac{\|f\|_{JN_p}^p}{t^p}$$

for all $t > 0$. This completes the proof. \square

Notice also that the zero-extension of the function we had in Section 2 in $L^{p,\infty}(Q_0) \setminus JN_p(Q_0)$ is in $L^{p,\infty}(\mathbb{R}^n)/\mathbb{R} \setminus JN_p(\mathbb{R}^n)$. This shows that $JN_p(\mathbb{R}^n) \subsetneq L^{p,\infty}(\mathbb{R}^n)/\mathbb{R}$. It is easy to see, by using a similar method as in the previous proof, that $JN_1(\mathbb{R}^n) = L^1(\mathbb{R}^n)/\mathbb{R}$, if we define JN_1 the same way as in Definition 5.1.

The vanishing subspace VJN_p can be defined in the whole space as well.

5.3. Definition (VJN_p). Let $1 < p < \infty$. Then the vanishing subspace VJN_p of JN_p is defined by setting

$$VJN_p := \overline{D_p(\mathbb{R}^n)} \cap JN_p^{JN_p},$$

where

$$D_p(\mathbb{R}^n) := \{f \in C^\infty(\mathbb{R}^n) : |\nabla f| \in L^p(\mathbb{R}^n)\}.$$

The following characterization of VJN_p is proven in [16, Theorem 3.2].

5.4. Theorem. Let $1 < p < \infty$. Then $f \in VJN_p$ if and only if $f \in JN_p$ and

$$\lim_{a \rightarrow 0} \sup_{\substack{Q_i \subset \mathbb{R}^n \\ l(Q_i) \leq a}} \sum_{i=1}^{\infty} |Q_i| \left(\int_{Q_i} |f - f_{Q_i}| \right)^p = 0$$

where the supremum is taken over all collections of pairwise disjoint cubes Q_i such that the side length of each Q_i is at most a .

We also study $CJN_p(\mathbb{R}^n)$, which is another subspace of $JN_p(\mathbb{R}^n)$. This space is defined analogously to the space of continuous mean oscillation CMO , which is a subspace of BMO . In the case of the bounded cube, $CJN_p(Q_0) = VJN_p(Q_0)$. Hence we only define CJN_p in the whole space.

5.5. Definition (CJN_p). Let $1 < p < \infty$. Then the subspace CJN_p of JN_p is defined by setting

$$CJN_p := \overline{C_c^\infty(\mathbb{R}^n)}^{JN_p},$$

where $C_c^\infty(\mathbb{R}^n)$ denotes the set of smooth functions with compact support in \mathbb{R}^n .

The following characterization of CJN_p is proven in [16, Theorem 4.3].

5.6. Theorem. Let $1 < p < \infty$. Then $f \in CJN_p$ if and only if $f \in VJN_p$ and

$$\lim_{a \rightarrow \infty} \sup_{\substack{Q \subset \mathbb{R}^n \\ l(Q) \geq a}} |Q|^{1/p} \int_Q |f - f_Q| = 0$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ such that the side length of Q is at least a .

It follows directly from the definition of CJN_p that $L^p(\mathbb{R}^n) \subset CJN_p(\mathbb{R}^n)$, by density of C_c^∞ in L^p and by Equation (2.3). Furthermore we notice from Theorems 5.4 and 5.6 and from Definition 5.1 that adding a constant to a function doesn't affect whether the function is in JN_p or in the subspaces VJN_p or CJN_p . Thus if $f \in L^p$, then $f + c \in CJN_p$ for any constant c . If we define L^p/\mathbb{R} the same way as $L^{p,\infty}/\mathbb{R}$, then it is immediate that

$$L^p/\mathbb{R} \subseteq CJN_p \subseteq VJN_p \subseteq JN_p.$$

Again we can ask whether these inclusions are strict. We examine this by extending the previous functions into multidimensional versions in the whole space.

5.7. Lemma. *Let $I \subset \mathbb{R}$ be a bounded interval such that $|I| = L$. Let $f \in L^1(I)$, $n \in \mathbb{Z}_+$ and $1 < p < \infty$. Define the cube $Q_0 := I \times [0, L]^{n-1} \subset \mathbb{R}^n$. Define the function*

$$\hat{f}(x_1, x_2, \dots, x_n) := \begin{cases} f(x_1), & \text{if } x_1 \in I \text{ and } 0 \leq x_i < L \text{ for all } i \geq 2 \\ f_I & \text{elsewhere.} \end{cases}$$

Then $\hat{f} \in JN_p$ if and only if $f \in JN_p(I)$ and

$$2^{\frac{1-n}{p}} \|f\|_{JN_p(I)} L^{\frac{n-1}{p}} \leq \|\hat{f}\|_{JN_p} \leq c(n, p) \|f\|_{JN_p(I)} L^{\frac{n-1}{p}}.$$

Proof. If $\hat{f} \in JN_p$, then clearly if we restrict \hat{f} to Q_0 , we have $\hat{f} \in JN_p(Q_0)$. Then Proposition 2.5 implies that

$$2^{\frac{1-n}{p}} \|f\|_{JN_p(I)} L^{\frac{n-1}{p}} \leq \|\hat{f}\|_{JN_p(Q_0)} \leq \|\hat{f}\|_{JN_p}$$

and so $f \in JN_p(I)$.

Now assume that $f \in JN_p(I)$. Without loss of generality we may assume that $f_I = 0$. Let $Q \subset \mathbb{R}^n$ be a cube such that $Q \cap Q_0 \neq \emptyset$ and $Q \setminus Q_0 \neq \emptyset$. Since $Q \cap Q_0$ is an intersection of two cubes, it is a rectangle. Therefore we can denote $Q \cap Q_0 = J_1 \times J_2 \times \dots \times J_n = J_1 \times K$, where J_i are intervals such that $J_1 \subset I$ and $J_i \subset [0, L[$ for $i \geq 2$. The notation $|\cdot|$ for sets below refers to the Lebesgue measure, which can be either 1-, n - or $n-1$ -dimensional depending on the context. Then

$$\begin{aligned} |Q| \left(\int_Q |\hat{f} - f_Q| \right)^p &\leq 2^p |Q \cap Q_0|^{1-p} \left(\int_{Q \cap Q_0} |\hat{f}| \right)^p \\ &= 2^p |J_1|^{1-p} |K|^{1-p} \left(\int_K \int_{J_1} |f(x_1)| dx_1 dx_2 \dots dx_n \right)^p \\ &= 2^p |K| |J_1|^{1-p} \left(\int_0^\infty |\{x \in J_1 : |f(x)| > t\}| dt \right)^p \\ &\leq 2^p |K| |J_1|^{1-p} \left(\int_0^{|J_1|^{-1/p} \|f\|_{L^{p,\infty}(J_1)}} |J_1| dt \right. \\ &\quad \left. + \int_{|J_1|^{-1/p} \|f\|_{L^{p,\infty}(J_1)}}^\infty \frac{\|f\|_{L^{p,\infty}(J_1)}^p}{t^p} dt \right)^p \\ &= 2^p |K| |J_1|^{1-p} \left(|J_1|^{1-1/p} \|f\|_{L^{p,\infty}(J_1)} \right. \\ &\quad \left. + \|f\|_{L^{p,\infty}(J_1)}^p \frac{1}{p-1} (|J_1|^{-1/p} \|f\|_{L^{p,\infty}(J_1)})^{1-p} \right)^p \\ &\leq \left(\frac{2p}{p-1} \right)^p \|f\|_{L^{p,\infty}(I)}^p |K| \\ &\leq c(p) \|f\|_{JN_p(I)}^p |K|. \end{aligned}$$

Here we used the triangle inequality, Cavalieri's principle and Theorem 2.4.

Let us look at $Q \cap Q_0 = J_1 \times \dots \times J_n$. Clearly we have $|J_i| \leq l(Q)$ for every i . Further we know that $|J_j| < l(Q)$ for at least one j - otherwise we would have $|Q \cap Q_0| = |Q|$.

If $|J_1| < l(Q)$, then the cube Q intersects with the boundary of I . This implies that K , the $n-1$ -dimensional side of $Q \cap Q_0$, is located at the boundary of Q_0 . In other words we can say

$$|K| \leq |\{x \in Q \cap Q_0 : x_1 \in \partial I\}| \leq |\{x \in Q \cap Q_0 : x \in \partial Q_0\}| = |Q \cap \partial Q_0|.$$

Here the first inequality is an equality if Q intersects only one of the boundary points of I . If it intersects the other boundary point, we have $|\{x \in Q \cap Q_0 : x_1 \in \partial I\}| = 2|K|$.

If $|J_1| = l(Q)$, then $|J_i| < l(Q)$ for some $i \geq 2$. By the same reasoning as before we know that Q intersects the boundary of the i :th dimensional interval $[0, L[$ and therefore

$$\begin{aligned} |K| &= |J_2||J_3|\dots|J_n| \leq |J_1||J_2|\dots|J_{i-1}||J_{i+1}|\dots|J_n| \\ &\leq |\{x \in Q \cap Q_0 : x_i \in \partial[0, L[\}| \leq |\{x \in Q \cap Q_0 : x \in \partial Q_0\}| \\ &= |Q \cap \partial Q_0|. \end{aligned}$$

Thus in any case we have $|K| \leq |Q \cap \partial Q_0|$. Now let \mathcal{Q} be a partition of \mathbb{R}^n into disjoint cubes and define

$$F(Q) := |Q| \left(\int_Q |\hat{f} - \hat{f}_Q| \right)^p.$$

Then

$$\begin{aligned} \sum_{Q \in \mathcal{Q}} F(Q) &= \sum_{\substack{Q \in \mathcal{Q} \\ Q \subset Q_0}} F(Q) + \sum_{\substack{Q \in \mathcal{Q} \\ Q \subset \mathbb{R}^n \setminus Q_0}} F(Q) + \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} F(Q) \\ &\leq \|\hat{f}\|_{JN_p(Q_0)}^p + 0 + \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} c(p) \|f\|_{JN_p(I)}^p |K| \\ &\leq \|f\|_{JN_p(I)}^p L^{n-1} + c(p) \|f\|_{JN_p(I)}^p \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} |Q \cap \partial Q_0| \\ &\leq \|f\|_{JN_p(I)}^p (L^{n-1} + c(p) |\partial Q_0|) \\ &= \|f\|_{JN_p(I)}^p (L^{n-1} + c(p) 2nL^{n-1}) \\ &= c(n, p) \|f\|_{JN_p(I)}^p L^{n-1}. \end{aligned}$$

Here we used Proposition 2.5. Finally by taking the supremum this gives us the result

$$\|\hat{f}\|_{JN_p} \leq c(n, p) \|f\|_{JN_p(I)} L^{\frac{n-1}{p}}.$$

□

The moral is that if we have any function in $JN_p(I)$ for a bounded interval I , we can extend the function into multiple dimensions and into the whole space so that we get a function in $JN_p(\mathbb{R}^n)$. It is easy to see that $\hat{f} \in L^p(\mathbb{R}^n)/\mathbb{R}$ if and only if

$f \in L^p(I)$ and $\hat{f} \in L^{p,\infty}(\mathbb{R}^n)/\mathbb{R}$ if and only if $f \in L^{p,\infty}(I)$. The same results hold also if $n = 1$.

5.8. Lemma. *Let $I \subset \mathbb{R}$ be a bounded interval such that $|I| = L$. Let $f \in L^1(I)$, $n \in \mathbb{Z}_+$ and $1 < p < \infty$. Define the cube $Q_0 := I \times [0, L]^{n-1} \subset \mathbb{R}^n$. Define the function*

$$\hat{f}(x_1, x_2, \dots, x_n) := \begin{cases} f(x_1), & \text{if } x_1 \in I \text{ and } 0 \leq x_i < L \text{ for all } i \geq 2 \\ f_I & \text{elsewhere.} \end{cases}$$

Then if $\hat{f} \in CJN_p$, we have $f \in VJN_p(I)$. Especially if $f \notin VJN_p(I)$, then $\hat{f} \notin VJN_p$. Further if we assume that

$$\lim_{a \rightarrow 0} \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p = 0, \quad (5.9)$$

then $\hat{f} \in CJN_p$, if $f \in VJN_p(I)$.

Proof. If $\hat{f} \in CJN_p$, then $\hat{f} \in VJN_p$. Clearly if we restrict \hat{f} to Q_0 , we have $\hat{f} \in VJN_p(Q_0)$. Then Proposition 4.6 implies that $f \in VJN_p(I)$.

If $f \notin VJN_p(I)$ and we restrict \hat{f} to Q_0 , then Proposition 4.6 implies that $\hat{f} \notin VJN_p(Q_0)$ which also implies that $\hat{f} \notin VJN_p$.

Now assume that $f \in VJN_p(I)$ and assumption (5.9) holds. Without loss of generality we may assume that $f_I = 0$. First we notice that then $f \in JN_p(I)$, which means that $\hat{f} \in JN_p$, by Lemma 5.7. Now let $a > 0$. Let $Q \subset \mathbb{R}^n$ be a cube such that $Q \cap Q_0 \neq \emptyset$, $Q \setminus Q_0 \neq \emptyset$ and $l(Q) \leq a$. Similar to the previous proof, we can write $Q \cap Q_0 = J_1 \times J_2 \times \dots \times J_n = J_1 \times K$. Then

$$|Q| \left(\int_Q |\hat{f} - \hat{f}_Q| \right)^p \leq 2^p |K| |J_1|^{1-p} \left(\int_{J_1} |f| \right)^p$$

as in the previous proof. Further we have $|J_1| \leq l(Q) \leq a$, which means that

$$|J_1|^{1-p} \left(\int_{J_1} |f| \right)^p \leq \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p.$$

As in the previous proof, we also have $|K| \leq |Q \cap \partial Q_0|$. Now let \mathcal{Q} be a partition of \mathbb{R}^n into disjoint cubes such that the side length of each cube is at most a . Define

$$F(Q) := |Q| \left(\int_Q |\hat{f} - \hat{f}_Q| \right)^p.$$

Then

$$\begin{aligned} \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} F(Q) &\leq \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} 2^p |K| \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p \\ &\leq 2^p \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} |Q \cap \partial Q_0| \end{aligned}$$

$$\begin{aligned}
&\leq 2^p \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p |\partial Q_0| \\
&= 2^p \cdot 2nL^{n-1} \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p.
\end{aligned}$$

Therefore

$$\begin{aligned}
\sum_{Q \in \mathcal{Q}} F(Q) &= \sum_{\substack{Q \in \mathcal{Q} \\ Q \subset Q_0}} F(Q) + \sum_{\substack{Q \in \mathcal{Q} \\ Q \subset \mathbb{R}^n \setminus Q_0}} F(Q) + \sum_{\substack{Q \in \mathcal{Q} \\ Q \cap Q_0 \neq \emptyset \\ Q \setminus Q_0 \neq \emptyset}} F(Q) \\
&\leq \sup_{\mathcal{Q}} \sum_{\substack{Q \in \mathcal{Q} \\ Q \subset Q_0}} F(Q) + 0 + 2^p \cdot 2nL^{n-1} \sup_{\substack{J \subset I \\ |J| \leq a}} |J|^{1-p} \left(\int_J |f| \right)^p.
\end{aligned}$$

If we restrict \hat{f} to Q_0 , then Proposition 4.6, together with $f \in VJN_p(I)$, implies that $\hat{f} \in VJN_p(Q_0)$. This gives us that

$$\limsup_{a \rightarrow 0} \sup_{\substack{Q \in \mathcal{Q} \\ Q \subset Q_0}} F(Q) = 0.$$

For the other term we have a similar limit by assumption (5.9). Then we have

$$\limsup_{a \rightarrow 0} \sup_{Q \in \mathcal{Q}} F(Q) = 0$$

and so Theorem 5.4 gives us that $\hat{f} \in VJN_p$. Finally we know that $f \in L^1(I)$ and \hat{f} is supported in Q_0 . Therefore

$$\begin{aligned}
\lim_{a \rightarrow \infty} \sup_{\substack{Q \subset \mathbb{R}^n \\ l(Q) \geq a}} |Q|^{1/p} \int_Q |\hat{f} - \hat{f}_Q| &\leq \lim_{a \rightarrow \infty} \sup_{\substack{Q \subset \mathbb{R}^n \\ l(Q) \geq a}} |Q|^{1/p} 2 \int_Q |\hat{f}| \\
&\leq \lim_{a \rightarrow \infty} \sup_{\substack{Q \subset \mathbb{R}^n \\ l(Q) \geq a}} |Q|^{1/p-1} 2 \int_{Q_0} |\hat{f}| \\
&= 2L^{n-1} \int_I |f| \lim_{a \rightarrow \infty} (a^n)^{1/p-1} = 0,
\end{aligned}$$

and so Theorem 5.6 gives us that $\hat{f} \in CJN_p$. This completes the proof. \square

5.10. Lemma. *Let g_0 be the function in Proposition 4.4 such that $g_0 \in VJN_p(Q_0) \setminus L^p(Q_0)$ with $Q_0 \subset \mathbb{R}$. Then we have*

$$\lim_{a \rightarrow 0} \sup_{\substack{J \subset Q_0 \\ |J| \leq a}} |J|^{1-p} \left(\int_J g_0 \right)^p = 0.$$

Proof. Let a be a small positive number such that $\delta_N < a \leq \delta_{N-1}$ for some positive integer N . Let $J \subset Q_0$ be an interval such that $|J| \leq a$. Then there is some integer $m \geq N$ such that $\delta_m < |J| \leq \delta_{m-1}$. Like in earlier cases, if J does not intersect any interval \hat{I} , then

$$|J|^{1-p} \left(\int_J g_0 \right)^p = 0.$$

Therefore we assume that J does intersect some interval \hat{I} . Then there is a unique widest interval I_J such that J intersects \hat{I}_J .

First assume that $|I_J| = |I| = 2^{-i} > 2^{-m}$. Then the fact that $|J \setminus \hat{I}| \leq |J| \leq \delta_{m-1} \leq \delta_i$ implies that \hat{I} is the only interval that J intersects. Then

$$\begin{aligned} |J|^{1-p} \left(\int_J g_0 \right)^p &\leq |J|^{1-p} \left(\int_J h_i \right)^p = |J| h_i^p \leq \delta_{m-1} h_{m-1}^p \leq 2^{-m^2} \cdot 2^{(m-1)^2} \\ &= 2^{1-2m} \leq 2^{-N}. \end{aligned}$$

As $a \rightarrow 0$, we know that $N \rightarrow \infty$. Therefore this upper bound converges to 0.

Assume now that $|I_J| = 2^{-i} \leq 2^{-m}$. Then J intersects at most one interval of length $|\hat{I}| = l_m$, two intervals of length $|\hat{I}| = l_{m+1}$ etc. If I is an interval of length 2^{-m} , then in this case we have

$$\int_J g_0 \leq \min(|J|, l_m) h_m + \sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{\hat{I}'} g_0.$$

For the second term we have the following upper bound.

$$|J|^{1-p} \left(\sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{\hat{I}'} g_0 \right)^p \leq c 2^{-mp}.$$

For the proof we refer to [4, Lemma 3.8]. For the first term we notice that

$$|J|^{1-p} \min(|J|, l_m)^p \leq l_m$$

whether $|J| < l_m$ or $l_m \leq |J|$. This means that

$$\begin{aligned} |J|^{1-p} \left(\int_J g_0 \right)^p &\leq c |J|^{1-p} (\min(|J|, l_m) h_m)^p + c |J|^{1-p} \left(\sum_{\substack{I' \subsetneq I \\ I' \text{ dyadic}}} \int_{\hat{I}'} g_0 \right)^p \\ &\leq c l_m h_m^p + c 2^{-mp} \leq c 2^{-m^2-m} \cdot 2^{m^2} + c 2^{-m} \leq c 2^{-N}. \end{aligned}$$

Hence whenever $|J| \leq a$, we have

$$|J|^{1-p} \left(\int_J g_0 \right)^p \leq c(p) 2^{-N}.$$

This means that

$$\lim_{a \rightarrow 0} \sup_{\substack{J \subset Q_0 \\ |J| \leq a}} |J|^{1-p} \left(\int_J g_0 \right)^p \leq \lim_{N \rightarrow \infty} c(p) 2^{-N} = 0$$

and therefore g_0 satisfies Equation (5.9). \square

Now Lemmas 5.7, 5.8 and 5.10 together with the examples $g \in JN_p(Q_0) \setminus VJN_p(Q_0)$ from Proposition 4.3 and $g_0 \in VJN_p(Q_0) \setminus L^p(Q_0)$ from Proposition 4.4, imply the following corollary.

5.11. Corollary. *For any $n \in \mathbb{Z}_+$ there exists a function $\hat{g} \in JN_p(\mathbb{R}^n) \setminus VJN_p(\mathbb{R}^n)$ and a function $\hat{g}_0 \in CJN_p(\mathbb{R}^n) \setminus L^p(\mathbb{R}^n)/\mathbb{R}$.*

In conclusion we know that

$$L^p/\mathbb{R} \subsetneq CJN_p \subseteq VJN_p \subsetneq JN_p \subsetneq L^{p,\infty}/\mathbb{R}.$$

It still remains an open question whether the spaces $CJN_p(\mathbb{R}^n)$ and $VJN_p(\mathbb{R}^n)$ coincide. It would be interesting to construct an example of a function in $VJN_p \setminus CJN_p$.

Acknowledgements. I would like to thank my supervisor Riikka Korte for many fruitful discussions on the space JN_p , and for having the idea of the function g_0 in Proposition 4.4. I would also like to thank my colleague Kim Myyryläinen for providing the proof of Proposition 2.7.

REFERENCES

- [1] D. Aalto, L. Berkovits, O. E. Kansanen, and H. Yue. John-Nirenberg lemmas for a doubling measure. *Studia Math.* 204(1):21–37, 2011.
- [2] L. Berkovits, J. Kinnunen, and J. M. Martell. Oscillation estimates, self-improving results and good- λ inequalities. *J. Funct. Anal.*, 270(9):3559–3590, 2016.
- [3] A. Brudnyi and Y. Brudnyi. On the Banach structure of multivariate BV spaces. *Dissertationes Math.*, 548:1-52, 2020.
- [4] G. Dafni, T. Hytönen, R. Korte, and H. Yue. The space JN_p : Nontriviality and duality. *J. Funct. Anal.*, 275(3):577–603, 2018.
- [5] Ó. Domínguez and M. Milman. Sparse Brudnyi and John–Nirenberg spaces. arXiv: 2107.05117, 2021.
- [6] R. Hurri-Syrjänen, N. Marola and A. V. Vähäkangas. Aspects of local-to-global results. *Bull. Lond. Math. Soc.*, 46:1032–1042, 2014.
- [7] F. John and L. Nirenberg. On functions of bounded mean oscillation. *Comm. Pure Appl. Math.*, 14:415–426, 1961.
- [8] J. Kinnunen and K. Myyryläinen. Dyadic John–Nirenberg space. arXiv: 2107.00492, 2021.
- [9] N. Marola and O. Saari. Local to global results for spaces of BMO type. *Math. Z.*, 282:473–484, 2016.
- [10] M. Milman. Marcinkiewicz spaces, Garsia-Rodemich spaces and the scale of John-Nirenberg self improving inequalities. *Ann. Acad. Sci. Fenn. Math.*, 41:491-501, 2016.
- [11] K. Myyryläinen. Median-type John–Nirenberg space in metric measure spaces. arXiv: 2104.05380, 2021.
- [12] U. Neri. Fractional integration on the space H^1 and its dual. *Studia Math.*, 53:175-189, 1975.
- [13] D. Sarason. Functions of vanishing mean oscillation. *Trans. Amer. Math. Soc.*, 207:391-405, 1975.
- [14] J. Sun, G. Xie and D. Yang. Localized John–Nirenberg–Campanato spaces. *Anal. Math. Phys.* 11, Paper No. 29, 47 pp, 2021.
- [15] J. Tao, D. Yang and W. Yuan. John-Nirenberg-Campanato spaces. *Nonlinear Anal.* 189, 111584, 36 pp, 2019.
- [16] J. Tao, D. Yang and W. Yuan. Vanishing John–Nirenberg spaces. *Adv. Calc. Var.*, <https://doi.org/10.1515/acv-2020-0061>, 2021.
- [17] J. Tao, D. Yang and W. Yuan. A Survey on Function Spaces of John–Nirenberg Type. arXiv: 2108.01511, 2021.

(T.T.) AALTO UNIVERSITY, DEPARTMENT OF MATHEMATICS AND SYSTEMS ANALYSIS, ESPOO, FINLAND

Email address: timo.i.takala@aalto.fi