

THE MOORE-PENROSE INVERSE OF ACCRETIVE OPERATORS WITH APPLICATION TO QUADRATIC OPERATOR PENCILS

FAIROUZ BOUCHELACHEM¹, MOHAMMED BENHARRAT^{2*}

ABSTRACT. We establish some relationships between an m -accretive operator and its Moore-Penrose inverse. We derive some perturbation result of the Moore-Penrose inverse of a maximal accretive operator. As an application we give a factorization theorem for a quadratic pencil of accretive operators. Also, we study a result of existence, uniqueness, and maximal regularity of the strict solution for complete abstract second order differential equation. Illustrative examples are also given.

1. INTRODUCTION

The Moore-Penrose inverse of a linear operator in Hilbert space is a useful generalization of the ordinary inverse. This generalized inverse is an important theoretical and practical tool in algebra and analysis (Markov chains, singular differential and difference equations, iterative methods...), see [4, 31]. In particular, in [11, 12] (and the references therein) the perturbation analysis for the Moore-Penrose inverse of closed operators has been considered. Also, the expression of the generalized inverse of the perturbed operator has been investigated. In the paper [19] necessary and sufficient conditions for the cone nonnegativity of Moore-Penrose inverses of unbounded Gram operators are derived. These conditions include statements on acuteness of certain closed convex cones in infinite-dimensional real Hilbert spaces. In [3] a complete description of the left quotient and the right quotient of two bounded operators operators is given via the Moore-Penrose inverse. The objectives of this paper are to derive the properties of m -accretive operators via the Moore-Penrose inverse and establish some interesting results, especially for the perturbation analysis of Moore-Penrose inverses as well as of maximal accretive operators. Recall that a linear operator T with domain $\mathcal{D}(T)$ in a complex Hilbert space \mathcal{H} , is called accretive if its numerical range $W(T)$ is contained in the closed right half-plane, and if further has no proper accretive extensions in \mathcal{H} , it called maximal accretive, m -accretive for short. In particular, every m -accretive operator is accretive and closed densely defined, its adjoint is also m -accretive (cf. [16], p. 279). This class is of particular interest and related to the semi-group theory in following sens: an operator T is m -accretive if and only if $-T$ generates a strongly continuous contraction semigroup (Theorem of Lumer-Phillips).

Date: 15/01/2021.

* Corresponding author

This work was supported by the Laboratory of Fundamental and Applicable Mathematics of Oran (LM-FAO) and the Algerian research project: PRFU, no. C00L03ES310120180002.

2010 *Mathematics Subject Classification.* Primary 47A10; 47A56.

Key words and phrases. Moore-Penrose inverse, Accretive operators, Quadratic operator pencil, Perturbation, Semigroup of contractions.

In this paper, we explore the following two questions, the first what can be said about the m -accretivity of the Moore-Penrose of an m -accretive operator and conversely? the second concern the perturbation problem: Let T be m -accretive operator with a bounded Moore-Penrose inverse, what condition on the operator S can guarantee that $T + S$ is m -accretive and its Moore-Penrose inverse exists and it has the simplest expression?

In this work, we give a certain answers to the mentioned problems. This paper is organized as follows: In section 2, we establish some relationships between an m -accretive operator and its Moore-Penrose inverse. In section 3, we consider the perturbation for the m -accretive operator and its Moore-Penrose inverse. We prove that under weaker conditions that considered perturbation does not change the null space and the range space, consequently the perturbed operator is a closed EP operator. Utilizing this result, we study a class of a quadratic operator pencil $\mathcal{Q}(\lambda) = \lambda^2 I - 2\lambda T - S$, ($\lambda \in \mathbb{C}$), where the coefficients of which are accretive operators. Our aim, in Section 4, is to investigate a canonical factorization like $(\lambda I - Z_1)(\lambda I - Z_2)$ for of such pencils based on the perturbation theory of accretive operators. We also obtain a criterion in order that the linear factors, into which the pencil splits, generates an holomorphic semi-group of contraction operators. As an illustration, in section 5, we establish a theorem of existence, uniqueness, and maximal regularity of the strict solution of an abstract second order evolutionary equations generated by such pencils.

2. ACCRETIVE OPERATOR AND THE MOORE-PENROSE INVERSE

Throughout this paper \mathcal{H} is a complex Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. For a closed linear operator T on \mathcal{H} we denote by $\mathcal{D}(T)$, $\mathcal{R}(T)$, $\mathcal{N}(T)$, $\sigma(T)$ and $\rho(T)$ the domain, the range, the kernel, the spectrum and the resolvent set of T , respectively. The space of bounded linear operators on \mathcal{H} is denoted by $\mathcal{B}(\mathcal{H})$. For two possibly unbounded linear operators T, S on \mathcal{H} their product TS is defined on its natural domain $\mathcal{D}(TS) := \{x \in \mathcal{D}(S) : Sx \in \mathcal{D}(T)\}$ and their sum $T + S$ is defined in $\mathcal{D}(T + S) = \mathcal{D}(T) \cap \mathcal{D}(S)$. An inclusion $T \subseteq S$ denotes inclusion of graphs, i.e., it means that S extends T . A possibly unbounded operator T on \mathcal{H} commutes with a bounded operator $S \in \mathcal{B}(\mathcal{H})$ if the graph of T is $S \times S$ -invariant, or equivalently if $ST \subseteq TS$.

Recall that a linear operator T with domain $\mathcal{D}(T)$ in \mathcal{H} is said to be accretive if

$$\operatorname{Re} \langle Tx, x \rangle \geq 0 \quad \text{for all } x \in \mathcal{D}(T)$$

or, equivalently if

$$\|(\lambda + T)x\| \geq \lambda \|x\| \quad \text{for all } x \in \mathcal{D}(T) \text{ and } \lambda > 0.$$

An accretive operator T is called *maximal accretive*, or *m -accretive* for short, if T has no proper accretive extensions in \mathcal{H} . The following conditions are equivalent:

- (1) T is m -accretive.
- (2)

$$(\lambda + T)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \text{and} \quad \|(\lambda + T)^{-1}\| \leq \frac{1}{\lambda} \text{ for } \lambda > 0.$$

- (3) T is accretive densely defined and $\mathcal{R}(\lambda + T) = \mathcal{H}$ for some (and hence for every) $\lambda > 0$;
- (4) T is accretive densely defined and closed, and T^* is accretive;
- (5) $-T$ generates contractive one-parameter semigroup $\mathcal{T}(t) = \exp(-tT)$, $t \geq 0$.

In particular, a bounded accretive operator is m -accretive.

The numerical range of a linear operator $T : \mathcal{D}(T) \rightarrow \mathcal{H}$ it is defined by

$$W(T) := \{ \langle Tx, x \rangle : x \in \mathcal{D}(T), \text{ with } \|x\| = 1 \}, \quad (2.1)$$

It is well-known that $W(T)$ is a convex set of the complex plane (the Toeplitz-Hausdorff theorem), and in general is neither open nor closed, even for a closed operator T . Clearly, an operator T is accretive when $W(T)$ is contained in the closed right half-plane

$$W(T) \subset \overline{\mathbb{C}_+} := \{z \in \mathbb{C} : \operatorname{Re}(z) \geq 0\}.$$

Further, if T is m -accretive operator then $W(T)$ has the so-called spectral inclusion property

$$\sigma(T) \subset \overline{W(T)}. \quad (2.2)$$

Recall that a linear operator T in \mathcal{H} is called *sectorial* with vertex $z = 0$ and semi-angle $\omega \in [0, \pi/2)$, or ω -accretive for short, if its numerical range is contained in a closed sector with semi-angle ω ,

$$W(T) \subset \overline{\mathcal{S}(\omega)} := \{z \in \mathbb{C} : |\arg z| \leq \omega\} \quad (2.3)$$

or, equivalently,

$$|\operatorname{Im} \langle Tx, x \rangle| \leq \tan \omega \operatorname{Re} \langle Tx, x \rangle \quad \text{for all } x \in \mathcal{D}(T).$$

An ω -accretive operator T is called m - ω -accretive, if it is m -accretive. We have T is m - ω -accretive if and only if the operators $e^{\pm i\theta}T$ is m -accretive for $\theta = \frac{\pi}{2} - \omega$, $0 < \omega \leq \pi/2$. The resolvent set of an m - ω -accretive operator T contains the set $\mathbb{C} \setminus \overline{\mathcal{S}(\omega)}$ and

$$\|(T - \lambda I)^{-1}\| \leq \frac{1}{\operatorname{dist}(\lambda, \overline{\mathcal{S}(\omega)})}, \quad \lambda \in \mathbb{C} \setminus \overline{\mathcal{S}(\omega)}.$$

In particular, m - $\pi/2$ -accretivity means m -accretivity. A 0-accretive operator is symmetric. An operator is positive if and only if it is m -0-accretive.

It is known that the C_0 -semigroup $\mathcal{T}(t) = \exp(-tT)$, $t \geq 0$, has contractive and holomorphic continuation into the sector $\overline{\mathcal{S}(\pi/2\omega)}$ if and only if the generator T is m - ω -accretive, see [16, Theorem V-3.35].

Recall that for bounded operator T , we have

$$\operatorname{Re}(T) = \frac{1}{2}(T + T^*) \quad \text{and} \quad \operatorname{Im}(T) = \frac{1}{2i}(T - T^*),$$

where $\operatorname{Re}(T)$ and $\operatorname{Im}(T)$ are self-adjoint operators and called it the real and imaginary parts of T , with

$$T = \operatorname{Re}(T) + i\operatorname{Im}(T),$$

Such decomposition is unique and called the cartesian decomposition of T . In this case, T is accretive if $\operatorname{Re}(T)$ is a nonnegative operator.

The spectral radius and the numerical radius of a bounded operator T are defined, respectively, by

$$r(T) = \sup_{\lambda \in \sigma(T)} |\lambda|$$

and

$$w(T) = \sup_{\|x\|=1} |\langle Tx, x \rangle|.$$

The next results is a generalization of [1, Lemma 2. and Theorem 2.] from matrices to bounded operators.

Lemma 2.1. *Let $T \in \mathcal{B}(\mathcal{H})$ such that $\operatorname{Re}(T) \geq \delta I$, for some $\delta > 0$. Denote by $\mathcal{S} = \operatorname{Im}(T)(\operatorname{Re}(T))^{-1}$. Then there exists $\lambda_0 \in \sigma(\mathcal{S})$, $|\lambda_0| = r(\mathcal{S}) = w(\mathcal{S}) = \|\mathcal{S}\|$ such that T is a sectorial operator with semiangle $\omega = \arctan |\lambda_0|$ and*

$$|\lambda_0| \leq \sqrt{\frac{\|T\|^2}{\delta^2} - 1}. \quad (2.4)$$

Proof. By assumption,

$$\langle \operatorname{Re}(T)x, x \rangle \geq \delta \|x\|^2$$

for all $x \in \mathcal{H}$. Hence we get

$$|\langle \operatorname{Im}(T)x, x \rangle| \leq \|\operatorname{Im}(T)\| \|x\|^2 \leq \frac{1}{\delta} \|\operatorname{Im}(T)\| \langle \operatorname{Re}(T)x, x \rangle.$$

Thus, T is sectorial operator with a vertex at the origin with a semiangle ω . In particular, for $\operatorname{Re}(T)^{-\frac{1}{2}}x$, we have

$$\left| \langle \operatorname{Im}(T)\operatorname{Re}(T)^{-\frac{1}{2}}x, \operatorname{Re}(T)^{-\frac{1}{2}}x \rangle \right| \leq \left\| \operatorname{Re}(T)^{-\frac{1}{2}}\operatorname{Im}(T)\operatorname{Re}(T)^{-\frac{1}{2}} \right\| \|x\|^2$$

this implies that $\tan(\omega) = \|\mathcal{T}\|$, where $\mathcal{T} = \operatorname{Re}(T)^{-\frac{1}{2}}\operatorname{Im}(T)\operatorname{Re}(T)^{-\frac{1}{2}}$. Since \mathcal{T} is a self-adjoint operator we have $r(\mathcal{T}) = w(\mathcal{T}) = \|\mathcal{T}\|$. Thus assert the existence of a $\lambda_0 \in \sigma(\mathcal{T})$ such that $|\lambda_0| = r(\mathcal{T}) = w(\mathcal{T}) = \|\mathcal{T}\|$. Since

$$\mathcal{T} = \operatorname{Re}(T)^{-\frac{1}{2}}\mathcal{S}\operatorname{Re}(T)^{\frac{1}{2}} = \operatorname{Re}(T)^{\frac{1}{2}}\mathcal{S}^*\operatorname{Re}(T)^{-\frac{1}{2}},$$

the self-adjoint operators \mathcal{T} and \mathcal{S} have the same spectrum (which is real) and hence have the same closure of the numerical range. This shows that $\lambda_0 \in \sigma(\mathcal{S})$, with $|\lambda_0| = w(\mathcal{T}) = w(\mathcal{S})$ and hence $\omega = \arctan |\lambda_0|$.

Now, assume that λ_0 is an eigenvalue of \mathcal{T} with $|\lambda_0| = \|\mathcal{T}\|$, there exists $u \in \mathcal{H}$ with $\|u\| = 1$ and $\mathcal{T}u = \lambda_0 u$. We have

$$\operatorname{Re}(T)^{-\frac{1}{2}}T\operatorname{Re}(T)^{-\frac{1}{2}} = (I + i\mathcal{T})$$

and

$$\left\| \operatorname{Re}(T)^{-\frac{1}{2}}T\operatorname{Re}(T)^{-\frac{1}{2}} \right\|^2 \geq \|(I + i\mathcal{T})u\|^2 = 1 + \|\mathcal{T}\|^2 = 1 + |\lambda_0|^2,$$

which implies that

$$|\lambda_0|^2 \leq \frac{\|T\|^2}{\delta^2} - 1.$$

Now we consider the general case. Let $\varepsilon > 0$. It follows from the spectral theorem that there exists a self-adjoint bounded operator P such that $\|P\| \leq \varepsilon$ and the operator $\mathcal{T} + P$ has an eigenvalue such that the modulus equals $\|\mathcal{T} + P\|$. As above, we take the operator $I + i(\mathcal{T} + P)$ instead of $(I + i\mathcal{T})$, we get

$$\frac{\|T\|^2}{\delta^2} + \varepsilon^2 \geq 1 + |\lambda_0|^2.$$

Letting $\varepsilon \rightarrow 0$, we obtain (2.4). □

If we assume, in Lemma 2.1, that the numerical range of T is closed, then $\lambda_0 \in W(T)$, but the extreme points of the numerical range are in the point spectrum, so λ_0 must be an eigenvalue of \mathcal{T} . So, we have the following

Corollary 2.2. *Let $T \in \mathcal{B}(\mathcal{H})$ with closed numerical range such that $\operatorname{Re}(T) \geq \delta I$, for some $\delta > 0$. Denote by $\mathcal{S} = \operatorname{Im}(T)(\operatorname{Re}(T))^{-1}$. Then there exists $\lambda_0 \in \sigma_p(\mathcal{S})$, $|\lambda_0| = w(\mathcal{S})$ such that T is a sectorial operator with semiangle $\omega = \arctan |\lambda_0|$ and λ_0 verifies (2.4).*

Remark 2.3. (1) Since $\operatorname{Re}(T)$ is strongly nonnegative, we know that

$$\sigma(\mathcal{S}) \subseteq \left\{ \frac{\beta}{\alpha} : \alpha \in \overline{W(\operatorname{Re}(T))}, \beta \in \overline{W(\operatorname{Im}(T))} \right\}.$$

So $\lambda_0 = \frac{\beta_0}{\alpha_0}$ for $\alpha_0 \in \overline{W(\operatorname{Re}(T))}$ and $\beta_0 \in \overline{W(\operatorname{Im}(T))}$.

(2) As we can see from the proof above that T can be represented as

$$T = \operatorname{Re}(T)^{\frac{1}{2}}(I + i\mathcal{T})\operatorname{Re}(T)^{\frac{1}{2}}$$

with $\mathcal{T} = \operatorname{Re}(T)^{-\frac{1}{2}}\operatorname{Im}(T)\operatorname{Re}(T)^{-\frac{1}{2}}$ and $\tan(\omega) = \|\mathcal{T}\|$. This is exactly the representation given in [16, Theorem VI-3.2]. In our case the selfadjoint operator is uniquely determined.

Next, in order to give some new results about accretive operator by using the Moore-Penrose inverse, let recall the definition of this generalized inverse for a closed densely defined operator.

Definition 2.4. [4] Let T be a closed densely defined on \mathcal{H} . Then there exists a unique closed densely defined operator T^\dagger , with domain $\mathcal{D}(T^\dagger) = \mathcal{R}(T) \oplus \mathcal{N}(T)^\perp$ such that

$$\begin{aligned} TT^\dagger T &= T \quad \text{on } \mathcal{D}(T), & T^\dagger T T^\dagger &= T^\dagger \quad \text{on } \mathcal{D}(T^\dagger), \\ TT^\dagger &= P_{\overline{\mathcal{R}(T)}} \quad \text{on } \mathcal{D}(T^\dagger), & T^\dagger T &= P_{\mathcal{N}(T)^\perp} \quad \text{on } \mathcal{D}(T), \end{aligned}$$

with $P_{\mathcal{M}}$ denotes the orthogonal projection onto a closed subspace \mathcal{M} .

This unique operator T^\dagger is called the Moore-Penrose inverse of T . (or the Maximal Tseng generalized Inverse in the terminology of [4]). Clearly,

- (1) $\mathcal{N}(T^\dagger) = \mathcal{R}(T)^\perp$,
- (2) $\mathcal{R}(T^\dagger) = \mathcal{N}(T)^\perp \cap \mathcal{D}(T)$.

As a consequence of the closed graph theorem T^\dagger is bounded if and only if $\mathcal{R}(T)$ is closed in \mathcal{H} , see [4].

Now, if we assume that T is an m-accretive operator, then

$$\mathcal{N}(T) = \mathcal{N}(T^*) \quad \text{and} \quad \mathcal{N}(T) \subseteq \mathcal{D}(T) \cap \mathcal{D}(T^*). \quad (2.5)$$

Thus $\overline{\mathcal{R}(T)} = \overline{\mathcal{R}(T^*)}$ and $\mathcal{H} = \overline{\mathcal{R}(T)} \oplus \mathcal{N}(T)$. Consequently, the operator T is written in a matrix form with respect to mutually orthogonal subspaces decomposition as follows

$$T = \begin{bmatrix} T_1 & 0 \\ 0 & 0 \end{bmatrix} : \begin{bmatrix} \overline{\mathcal{R}(T)} \\ \mathcal{N}(T) \end{bmatrix} \longrightarrow \begin{bmatrix} \overline{\mathcal{R}(T)} \\ \mathcal{N}(T) \end{bmatrix};$$

with T_1 is an operator on $\overline{\mathcal{R}(T)} \cap \mathcal{D}(T)$ is injective with dense range in $\overline{\mathcal{R}(T)}$. Also, its Moore-Penrose inverse is given by

$$T^\dagger = \begin{bmatrix} T_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} : \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix} \longrightarrow \begin{bmatrix} \overline{\mathcal{R}(T)} \\ \mathcal{N}(T) \end{bmatrix},$$

with T_1^{-1} from $\mathcal{R}(T)$ to $\overline{\mathcal{R}(T)} \cap \mathcal{D}(T)$ is closed operator densely defined on $\overline{\mathcal{R}(T)}$ and $\mathcal{N}(T^\dagger) = \mathcal{N}(T) = \mathcal{N}(T^*)$. Further, $\mathcal{R}(T)$ is closed if and only if T_1^{-1} is bounded from $\mathcal{R}(T)$ to $\mathcal{R}(T) \cap \mathcal{D}(T)$.

In the next result, we redefine a bounded sectorial operator with semiangle $< \pi/2$ via the Moore-Penrose inverse.

Lemma 2.5. *Let $T \in \mathcal{B}(\mathcal{H})$. T is a sectorial operator with semiangle ω , $0 \leq \omega < \pi/2$, if and only if the following two conditions are fulfilled:*

- (1) $\overline{\text{Re}(T)} \geq 0$.
- (2) $\overline{\mathcal{R}(T)} \subset \overline{\mathcal{R}(\text{Re}(T))}$.

In this case, $\omega = \arctan |\lambda_0|$ for some $\lambda_0 \in \sigma(\mathcal{S})$, $|\lambda_0| = w(\mathcal{S})$ where $\mathcal{S} = \text{Im}(T)(\text{Re}(T))^\dagger$.

Proof. Let $T \in \mathcal{B}(\mathcal{H})$ be a sectorial operator with semiangle ω . Therefore $\text{Re}(T)$ is a nonnegative and $\mathcal{N}(T) = \mathcal{N}(T^*) = \mathcal{N}(\text{Re}(T))$. It follows that $\overline{\mathcal{R}(T)} = \overline{\mathcal{R}(\text{Re}(T))}$.

Conversely, let the two conditions of the theorem hold. Then $\mathcal{N}(T) = \mathcal{N}(T^*) \subset \mathcal{N}(\text{Re}(T))$, indeed, if $Tx = 0$ for some $x \neq 0$, then $\text{Re} \langle Tx, x \rangle = \langle \text{Re}(T)x, x \rangle = 0$. Consequently, $\text{Re}(T)x = 0$ and $T^*x = 2\text{Re}(T)x - Tx = 0$. Since $\mathcal{H} = \mathcal{N}(\text{Re}(T)) \oplus \overline{\mathcal{R}(\text{Re}(T))}$, we get

$$\overline{\mathcal{R}(\text{Re}(T))} \subset \overline{\mathcal{R}(T)} = \overline{\mathcal{R}(T^*)}$$

with $\overline{\mathcal{R}(T)}$ reduces T . By (2) we obtain $\overline{\mathcal{R}(\text{Re}(T))} = \overline{\mathcal{R}(T)} = \overline{\mathcal{R}(T^*)}$. It follows that the subspace $\overline{\mathcal{R}(T)}$ reduces also the operator $\text{Re}(T)$. Moreover, the restriction of $\text{Re}(T)$ to $\overline{\mathcal{R}(T)}$ is strongly nonnegative. So by Lemma 2.1, the restriction of T to $\overline{\mathcal{R}(T)}$ is sectorial operator with semiangle $\omega = \arctan |\lambda_0|$ such that $\lambda_0 \in \sigma(\mathcal{S}_{|\overline{\mathcal{R}(T)}})$ and $|\lambda_0| = \sup_{\|x\|=1} |\langle \mathcal{S}_{|\overline{\mathcal{R}(T)}} x, x \rangle|$, where $\mathcal{S}_{|\overline{\mathcal{R}(T)}} = \text{Im}(T)(\text{Re}(T)|_{\overline{\mathcal{R}(T)}})^{-1} P_{|\overline{\mathcal{R}(T)}}$. In this case $\mathcal{S} = \text{Im}(T)(\text{Re}(T))^\dagger$. □

Now, we consider unbounded operator T .

Proposition 2.6. *If T is m -accretive operator, then T^\dagger is m -accretive.*

Proof. By assumption,

$$\text{Re} \langle Tx, x \rangle \geq 0 \quad \text{for all } x \in \mathcal{D}(T) \cap \mathcal{N}(T)^\perp = \mathcal{R}(T^\dagger).$$

Hence

$$\text{Re} \langle y, T^\dagger y \rangle \geq 0 \quad \text{for all } y \in \mathcal{R}(T).$$

Now let $x \in \mathcal{D}(T^\dagger) = \mathcal{R}(T) \oplus \mathcal{N}(T)$, then $x = x_1 + x_2$, with $x_1 \in \mathcal{R}(T)$ and $x_2 \in \mathcal{N}(T)$. Therefore,

$$\text{Re} \langle x, T^\dagger x \rangle = \text{Re} \langle x_1, T^\dagger x_1 \rangle \geq 0,$$

which implies

$$\text{Re} \langle x, T^\dagger x \rangle \geq 0 \quad \text{for all } x \in \mathcal{D}(T^\dagger).$$

Since T^\dagger is closed densely defined and $(T^\dagger)^*$ is accretive, it follows that T^\dagger is m-accretive. \square

It well known that by [4, Theorem 2; p 341], $T^{\dagger\dagger} = T$, this yields to

Corollary 2.7. T^\dagger is m-accretive operator if and only if T is m-accretive.

Corollary 2.8. T is m-accretive operator with closed range if and only if T^\dagger is bounded and accretive.

Corollary 2.9. If T is m-accretive operator with closed range, then T is an EP (Equal Projections) operator, that is, T^\dagger bounded and $TT^\dagger = T^\dagger T$ on $\mathcal{D}(T)$.

Proposition 2.10. Let T be an accretive bounded operator. If $W(T) \subseteq \overline{\mathbb{D}}$ and $W(T^\dagger) \subseteq \overline{\mathbb{D}}$, then T is unitary on $\mathcal{R}(T)$.

Proof. It well known by [9, Theorem 1.3-1] that the numerical radius is equivalent to the usual operator norm;

$$w(T) \leq \|T\| \leq 2w(T).$$

Hence, the assumption that $w(T^\dagger) \leq 1$ implies that T^\dagger is bounded. Thus $\mathcal{R}(T)$ is closed. Since T is m-accretive, then $\mathcal{R}(T) = \mathcal{R}(T^*) = \mathcal{N}(T)^\perp$. We consider the restriction of T from $\mathcal{R}(T)$ into itself. Since $T_{|\mathcal{R}(T)}^\dagger = (T_{|\mathcal{R}(T)})^{-1}$, $w((T_{|\mathcal{R}(T)})^{-1}) = w(T_{|\mathcal{R}(T)}^\dagger) = w(T^\dagger) \leq 1$. Combining this with $w(T_{|\mathcal{R}(T)}) = w(T) \leq 1$ and applying [30, Corollary 1.] to $(T_{|\mathcal{R}(T)})^{-1}$ and $T_{|\mathcal{R}(T)}$, we conclude that $T_{|\mathcal{R}(T)}$ is unitary on $\mathcal{R}(T)$. \square

3. A PERTURBATION RESULTS

In the following, we shall consider the perturbation of Moore-Penrose inverse of m-accretive operators. This gives an example of EP-operators, see [14, Theorem 3.12]. The part about the Moore-Penrose invertibility of the operator $T + S$ appears to be new.

Theorem 3.1. Let T is m-accretive operator and S is bounded and accretive. We have

- (1) $T + S$ is m-accretive.
- (2) If $\mathcal{R}(T)$ is closed, $\mathcal{R}(S) \subseteq \mathcal{R}(T)$ and $\|T^\dagger S\| < 1$. Then
 - $\mathcal{R}(T + S) = \mathcal{R}(T)$ is closed and $\mathcal{N}(T + S) = \mathcal{N}(T)$.
 - $T + S$ is an EP operator, and

$$(T + S)^\dagger = (I + T^\dagger S)^{-1} T^\dagger = T^\dagger (I + S T^\dagger)^{-1}.$$

In particular,

$$T^\dagger = (T + S)^\dagger (I + S T^\dagger),$$

and

$$\|(T + S)^\dagger - T^\dagger\| \leq \frac{\|S\| \|T^\dagger\|^2}{1 - \|T^\dagger S\|}.$$

- Further, if S is θ -accretive operator with $\theta \in [0, \frac{\pi}{2})$, then

$$\|(T + S)^\dagger\| \leq 2 \|T^\dagger\| + (1 + \tan \theta)^2 \|T^\dagger\|^2.$$

Proof. (1) Clearly, the operator $T + S$, with $\mathcal{D}(T + S) = \mathcal{D}(T)$, is densely defined, closed and accretive. Since also its adjoint operator $(T + S)^* = T^* + S^*$ is accretive, the operator $T + S$ is m-accretive.

(2) If $\mathcal{R}(S) \subseteq \mathcal{R}(T)$, then it is obvious that $\mathcal{R}(T + S) \subseteq \mathcal{R}(T)$ and $TT^\dagger S = P_{\mathcal{R}(T)}S = S$. Conversely, let $y \in \mathcal{R}(T)$, so $y = Tx$ for some $x \in \mathcal{D}(T)$. The condition $\|T^\dagger S\| < 1$ implies that $(I + T^\dagger S)^{-1}$ exists and bounded. Hence, there exists $u \in \mathcal{D}(T)$ such that $x = (I + T^\dagger S)u$. This shows that $y = T(I + T^\dagger S)u = Tu + Su \in \mathcal{R}(T + S)$. Hence $\mathcal{R}(T) \subseteq \mathcal{R}(T + S)$. Consequently, $\mathcal{R}(T + S) = \mathcal{R}(T)$ is closed.

Since T and $T + S$ are m-accretive with closed ranges, then

$$\mathcal{N}(T + S) = \mathcal{R}(T + S)^\perp = \mathcal{R}(T)^\perp = \mathcal{N}(T).$$

Now we prove that $(T + S)^\dagger = (I + T^\dagger S)^{-1}T^\dagger$. Since, $\mathcal{R}(T + S)$ is closed and $\mathcal{N}(T + S) = \mathcal{R}(T + S)^\perp$, by Corollary 2.9, it follows that $T + S$ is an EP operator.

Put $\mathbf{T} = (I + T^\dagger S)^{-1}T^\dagger$. We show that \mathbf{T} satisfies all the axioms of the Definition 2.4. First let us remark that, since $(I + T^\dagger S)^{-1}$ is invertible, $\mathcal{D}(\mathbf{T}) = \mathcal{D}(T^\dagger) = \mathcal{R}(T) \oplus \mathcal{N}(T)$, $\mathcal{N}(\mathbf{T}) = \mathcal{N}(T^\dagger) = \mathcal{R}(T)^\perp = \mathcal{N}(T + S)$.

Let $v \in \mathcal{R}(\mathbf{T})$, then there exists $u \in \mathcal{D}(\mathbf{T})$ such that $v = \mathbf{T}u = (I + T^\dagger S)^{-1}T^\dagger u$. Hence $T^\dagger u = v + T^\dagger S v \in \mathcal{R}(T) \cap \mathcal{D}(T)$. So $v = T^\dagger u - T^\dagger S v \in \mathcal{D}(T)$.

Now for $v \in \mathcal{D}(T)$,

$$\begin{aligned} \mathbf{T}(T + S)v &= (I + T^\dagger S)^{-1}T^\dagger(T + S)v \\ &= (I + T^\dagger S)^{-1}T^\dagger(T + TT^\dagger S)v \quad (\text{since } S = TT^\dagger S) \\ &= (I + T^\dagger S)^{-1}T^\dagger T(I + T^\dagger S)v \\ &= (I + T^\dagger S)^{-1}P_{\mathcal{N}(T)^\perp}(I + T^\dagger S)v \\ &= (I + T^\dagger S)^{-1}P_{\mathcal{R}(T)}(I + T^\dagger S)v \\ &= (I + T^\dagger S)^{-1}(I + T^\dagger S)v \\ &= v = P_{\mathcal{R}(T)}v \\ &= P_{\mathcal{R}(T+S)}v \\ &= P_{\mathcal{N}(T+S)^\perp}v. \end{aligned}$$

and for $u \in \mathcal{D}(\mathbf{T})$,

$$\begin{aligned} (T + S)\mathbf{T}u &= (T + S)(I + T^\dagger S)^{-1}T^\dagger u \\ &= (T + TT^\dagger S)(I + T^\dagger S)^{-1}T^\dagger u \quad (\text{since } S = TT^\dagger S) \\ &= T(I + T^\dagger S)(I + T^\dagger S)^{-1}T^\dagger u \\ &= TT^\dagger u = P_{\mathcal{R}(T)}u \\ &= P_{\mathcal{R}(T+S)}u. \end{aligned}$$

The uniqueness of $(T + S)^\dagger$ follows from Definition 2.4.

Since $\mathcal{R}(S) \subseteq \mathcal{R}(T)$, by Neumann series, we have

$$(I + T^\dagger S)^{-1}T^\dagger = \sum_{n=0}^{\infty} (-T^\dagger S)^n T^\dagger = \sum_{n=0}^{\infty} T^\dagger (-ST^\dagger)^n = T^\dagger (I + ST^\dagger)^{-1}. \quad (3.1)$$

For the last inequality, we can see that

$$\begin{aligned} (T + S)^\dagger - T^\dagger &= (I + ST^\dagger)^{-1}T^\dagger - (I + ST^\dagger)(I + ST^\dagger)^{-1}T^\dagger \\ &= [I - (I + ST^\dagger)](I + ST^\dagger)^{-1}T^\dagger \\ &= (-ST^\dagger)(I + ST^\dagger)^{-1}T^\dagger. \end{aligned}$$

Hence we get the desired inequality.

Now, the fact that $\mathcal{H} = \mathcal{R}(T + S) \oplus \mathcal{N}(T + S) = \mathcal{R}(T) \oplus \mathcal{N}(T)$, and T is invertible from $\mathcal{R}(T)$ to $\mathcal{R}(T)$, we applied the [32, Proposition 3.9.], to obtain the last estimate. \square

Similarly, we have

Theorem 3.2. *Let T is m -accretive operator and S is bounded and accretive, then the Theorem 3.1 hold true, if $\mathcal{R}(T)$ is closed, $\mathcal{N}(T) \subseteq \mathcal{N}(S)$ and $\|ST^\dagger\| < 1$.*

Proof. (1) Since the operator $T + S$ is m -accretive, its adjoint operator $(T + S)^* = T^* + S^*$ is also m -accretive.

(2) If $\mathcal{R}(T^*)$ is closed and $\mathcal{N}(T) \subseteq \mathcal{N}(S)$, then it is obvious that the last inclusion gives $\mathcal{R}(S^*) \subseteq \mathcal{R}(T^*)$. Also, the condition $\|ST^\dagger\| < 1$ implies $\|(ST^\dagger)^*\| = \|(T^*)^\dagger S^*\| < 1$, and conversely. Hence by Theorem 3.1, $\mathcal{R}(T + S) = \mathcal{R}(T^* + S^*) = \mathcal{R}(T^*) = \mathcal{R}(T)$ is closed and $\mathcal{N}(T + S) = \mathcal{N}(T^* + S^*) = \mathcal{N}(T^*) = \mathcal{N}(T)$. Now, we proceed as in the proof of Theorem 3.1. \square

Remark 3.3. Recall that the reduced minimum modulus of a non-zero operator T is defined by

$$\gamma(T) = \inf\{\|Tx\| : x \in N(T)^\perp \cap \mathcal{D}(T), \|x\| = 1\}.$$

If $T = 0$ then we take $\gamma(T) = \infty$. Note that (see [16]), $\mathcal{R}(T)$ is closed if and only if $\gamma(T) > 0$. In that case, $\gamma(T) = \frac{1}{\|T^\dagger\|}$, where T^\dagger is the Moore-Penrose inverse of T . Let us remark that if we assume that $\|S\| < \frac{1}{\gamma(T)}$ instead the condition $\|T^\dagger S\| < 1$, then the Theorem 3.1 hold true.

Proposition 3.4. *Let T be an accretive such that T^2 is m -accretive. Then*

- (i) *T is m -accretive. Further, if T is θ -accretive with $\theta < \pi/4$, then T^2 is m - 2θ -accretive.*
- (ii) *If $\mathcal{R}(T)$ is closed, then $\mathcal{R}(T^2)$ is closed and $\gamma(T^2) \geq \frac{\gamma(T)^2}{2}$.*

Proof. (i) Since T is an accretive operator, by [13, Theorem 1.2], we have ,

$$\|Tx\|^2 \leq \nu \|x\|^2 + \frac{1}{\nu} \|T^2x\|^2, \quad (3.2)$$

for all $x \in \mathcal{D}(T^2)$ and an arbitrary $\nu > 0$. Choosing $\nu > 0$ so large that $\frac{1}{\nu} < 1$, we obtain T is T^2 -bounded with lower bound < 1 . Then $T^2 + T$ with domain $\mathcal{D}(T^2)$ is m -accretive. Now, let us remark that

$$\left(\frac{1}{4}I + T^2 + T\right)x = \left(\frac{1}{2}I + T\right)^2x$$

for all $x \in \mathcal{D}(T^2)$. Since the operator on the left-hand side is invertible, then $(\frac{1}{2}I + T)^2$ is invertible, so $\frac{1}{2}I + T$ is also invertible. It follows that T is m-accretive. Now, we applied [6, Theorem 4.].

(ii) By the Landau-Kolmogorov inequality, [18, Theorem.], applied to T , we have

$$\|Tx\|^2 \leq 2 \|T^2x\| \|x\| ,$$

for all $x \in \mathcal{N}(T^2)^\perp \cap \mathcal{D}(T^2)$. It follows that

$$\|x\|^2 \gamma(T)^2 \leq \|Tx\|^2 \leq 2 \|T^2x\| \|x\| ,$$

and hence

$$\|T^2x\| \geq \frac{\gamma(T)^2}{2} \|x\| ,$$

for all $x \in \mathcal{N}(T^2)^\perp \cap \mathcal{D}(T^2)$. Now by the definition of $\gamma(T^2)$ we obtain $\gamma(T^2) \geq \frac{\gamma(T)^2}{2}$. \square

By Proposition 3.4, Theorem 3.1 and Theorem 3.2,

Corollary 3.5. *Let T^2 be m-accretive, T accretive with closed range and S is bounded and accretive. If $\mathcal{R}(S) \subseteq \mathcal{R}(T)$ and $\|(T^\dagger)^2 S\| < 1$. (or $\mathcal{N}(T) \subseteq \mathcal{N}(S)$ and $\|S(T^2)^\dagger\| < 1$). Then*

- $\mathcal{R}(T^2 + S) = \mathcal{R}(T^2) = \mathcal{R}(T)$ is closed and $\mathcal{N}(T^2 + S) = \mathcal{N}(T^2) = \mathcal{N}(T)$.
- $\mathcal{H} = \mathcal{R}(T^2 + S) \oplus \mathcal{N}(T^2 + S)$.
- $T^2 + S$ is an EP operator, and

$$(T^2 + S)^\dagger = (I + (T^\dagger)^2 S)^{-1} (T^\dagger)^2 = (T^\dagger)^2 (I + S(T^\dagger)^2)^{-1}.$$

If further, T is injective, then T , T^2 and $T^2 + S$ are invertible, and

$$(T^2 + S)^{-1} = (I + T^{-2} S)^{-1} T^{-2} = T^{-2} (I + S T^{-2})^{-1}.$$

Proof. Since T^2 is m-accretive, then T is also m-accretive by Proposition 3.4. Hence $\mathcal{N}(T^2) = \mathcal{N}(T)$ and $\mathcal{R}(T^2) = \mathcal{R}(T)$. By [3, Lemma 5.5], we have $(T^\dagger)^2 = (T^2)^\dagger$. Now the result is obtained by Theorem 3.1 and Theorem 3.2. \square

Remark 3.6. • By Remark 3.3, the Corollary 4.2 hold true if we assume $\|T^\dagger S\| < \gamma(T)$ instead $\|(T^\dagger)^2 S\| < 1$.

- Ôta showed in [27, Theorem 2.1] that, if T is closed and an accretive such that there is a positive integer n with $\mathcal{D}(T^n)$ is dense in \mathcal{H} and $\mathcal{R}(T^n) \subset \mathcal{D}(T)$, then T is bounded . In particular, for a closed and accretive operator T , if $\mathcal{R}(T)$ is contained in $\mathcal{D}(T)$, or in $\mathcal{D}(T^*)$, then T is automatically bounded, see also [27, Theorem 3.3].
- In general, if T^2 is m-accretive; then T fails to be accretive. Take $T = i \frac{d}{dx}$ on $L^2(\mathbb{R})$. The operator T has its spectrum on both sides of the origin. But $T^2 = -\frac{d^2}{dx^2}$ is a nonnegative selfadjoint operator.

4. AN APPLICATION TO QUADRATIC OPERATOR PENCIL

Consider in the Hilbert space \mathcal{H} the following quadratic operator pencil

$$Q(\lambda) = \lambda^2 I - 2\lambda T - S, \quad (4.1)$$

where $\lambda \in \mathbb{C}$ is the spectral parameter and the two operators T and S with domain $\mathcal{D}(T)$ and $\mathcal{D}(S)$, respectively.

One of the approaches to study the spectral properties of quadratic operator pencil (4.1) consists of the reducing them to a first order system in a suitable space, [7, 28]. However, as it was pointed out in [33], this way may be unpractical in the situation when the space is difficult to construct or it is complicated for applications. In addition, as it was mentioned in [34, 35, 36], the direct treatment of higher order problems allows to get more general results. An useful approach to address the study and computation of the spectral structure of quadratic operators pencil is through the use of factorization. This method was developed by Krein and Langer [15], for a quadratic pencils of self-adjoint operators and by Langer [20], for self-adjoint polynomial operator pencils, see also [20, 8, 10]. The main idea of this approach consist to factoring them and studying the spectral properties of the factors. Of particular interest is the separation of spectral values of Q_n between the spectra of the roots. Such separation may be complicated, even in the case of eigenvalues, see [29] and references therein. The problem is of great importance in spectral theory of such general operators. Moreover, its understanding is crucial in the study of performance properties of many systems.

The purpose of this section is to extend some earlier factorization results essentially those given in [15, 10, 22, 24] for the self-adjoint quadratic operators case, to (4.1) based on the perturbation theory of accretive operators together with the uniquely determined fractional powers of the maximal accretive operators. We also obtain a criterion in order that the linear factors, into which the pencil splits, generates a holomorphic semi-group of contraction operators.

We mention that if T is m -accretive, then for each $\alpha \in (0, 1)$ the fractional powers T^α , $0 < \alpha < 1$, are defined by the following Balakrishnan formula, see [2],

$$T^\alpha x = \frac{\sin(\pi\alpha)}{\pi} \int_0^\infty \lambda^{\alpha-1} T(\lambda + T)^{-1} x dt,$$

for all $x \in \mathcal{D}(T)$. The operators T^α are $m - (\alpha\pi)/2$ -accretive and, if $\alpha \in (0, 1/2)$, then $\mathcal{D}(T^\alpha) = \mathcal{D}(T^{*\alpha})$. It was proved in [17, Theorem 5.1] that, if T is m -accretive, then $\mathcal{D}(T^{1/2}) \cap \mathcal{D}(T^{*1/2})$ is a core of both $T^{1/2}$ and $T^{*1/2}$ and the real part $\operatorname{Re} T^{1/2} := (T^{1/2} + T^{*1/2})/2$ defined on $\mathcal{D}(T^{1/2}) \cap \mathcal{D}(T^{*1/2})$ is a selfadjoint operator. Further, by [17, Corollary 2],

$$\mathcal{D}(T) = \mathcal{D}(T^*) \implies \mathcal{D}(T^{1/2}) = \mathcal{D}(T^{*1/2}) = \mathcal{D}(T_R^{1/2}) = \mathcal{D}[\phi], \quad (4.2)$$

where ϕ is the closed form associated with the sectorial operator T via the first representation theorem [16, Sect. VI.2.1] and T_R is the non-negative selfadjoint operator associated with the real part of ϕ given by $\operatorname{Re} \phi := (\phi + \phi^*)/2$, see [2, 16, 17, 23, 25, 26].

Our result of this section read as follows

Theorem 4.1. *Let T^2 be m -accretive, T is accretive and S is accretive bounded operator. Then, we have*

- (1) *The operator $\Upsilon = T^2 + S$ with domain $\mathcal{D}(T^2)$ is m -accretive.*

- (2) Υ admits a fractional powers Υ^α m - $(\alpha\pi/2)$ -accretive for each $0 < \alpha < 1$.
- (3) $Z_1 = T + \Upsilon^{\frac{1}{2}}$ and $Z_2 = T - \Upsilon^{\frac{1}{2}}$ with domain $\mathcal{D}(T) \cap \mathcal{D}(\Upsilon^{\frac{1}{2}})$ are T^2 -bounded with lower bound < 1 and closable operators. Further, Z_1 is accretive densely defined.
If further, $\mathcal{D}(\Upsilon^{\frac{1}{2}}) \subset \mathcal{D}(T)$, then
- (4) Z_1 is m - $\pi/4$ -accretive and for any $\varepsilon > 0$, there exists $r > 0$, such that $-Z_2 - r$ is m - $\pi/4 + \varepsilon$ -accretive.
In particular, $-Z_1$ and $Z_2 + r$ generates holomorphic C_0 -semigroup of contraction operators $\mathcal{T}_1(z)$ and $\mathcal{T}_2(z)$ of angle $\frac{\pi}{4}$ and $\pi/4 - \varepsilon$, respectively.
- (5) The spectra of Z_1 and Z_2 are separated,

$$\sigma(Z_1) \cap \sigma(Z_2) = \emptyset.$$

- (6) If $T(\mathcal{D}(T^2)) \subset \mathcal{D}(T^2)$ and $\Upsilon^{\frac{1}{2}}(\mathcal{D}(T^2)) \subset \mathcal{D}(T^2)$, then Q take the following form,

$$\mathcal{Q}(\lambda)x = \frac{1}{2}(\lambda I - Z_1)(\lambda I - Z_2)x + \frac{1}{2}(\lambda I - Z_2)(\lambda I - Z_1)x, \quad (4.3)$$

for all $x \in \mathcal{D}(T^2)$.

In particular, if $TS = ST$ on $\mathcal{D}(T^2)$, then Q admits the following canonical factorization

$$\mathcal{Q}(\lambda)x = (\lambda I - Z_1)(\lambda I - Z_2)x = (\lambda I - Z_2)(\lambda I - Z_1)x, \quad (4.4)$$

for all $x \in \mathcal{D}(T^2)$.

Proof. (1) An immediate consequence of Theorem 3.1-(1).

(2) Υ admits fractional powers Υ^α m - $(\alpha\pi/2)$ -accretive for each $0 < \alpha < 1$, see [2, 17].

(3) By (2); Υ admits unique root $\Upsilon^{\frac{1}{2}}$ m - $(\pi/4)$ -accretive operator with $\mathcal{D}(T^2)$ is a core of $\Upsilon^{\frac{1}{2}}$. So we define the following operators

$$Z_1 = T + \Upsilon^{\frac{1}{2}}$$

and

$$Z_2 = T - \Upsilon^{\frac{1}{2}}$$

with domain $\mathcal{D}(T) \cap \mathcal{D}(\Upsilon^{\frac{1}{2}})$. Both of Z_1 and Z_2 are densely defined on \mathcal{H} with numerical range is not the whole complex plane, it follows that Z_1 and Z_2 are closable operators. Now, we prove that Z_1 and Z_2 are T^2 -bounded with lower bound < 1 .

By Proposition 3.4, T is also m -accretive. Hence, by [13, Theorem 1.2], we have for an arbitrary $\rho_1 > 0$ and $\rho_2 > 0$,

$$\|Tx\|^2 \leq \rho_1 \|x\|^2 + \frac{1}{\rho_1} \|T^2x\|^2, \quad (4.5)$$

and

$$\left\| \Upsilon^{\frac{1}{2}}x \right\|^2 \leq \rho_2 \|x\|^2 + \frac{1}{\rho_2} \|\Upsilon x\|^2, \quad (4.6)$$

for all $x \in \mathcal{D}(T^2)$ (cf. by [25, Chap. 2, Theorem 6.10]).

By (4.5) and (4.6), it follows that

$$\begin{aligned}
\|Z_i x\|^2 &\leq 2 \|Tx\|^2 + 2 \left\| \Lambda^{\frac{1}{2}} x \right\|^2 \\
&\leq 2(\rho_1 \|x\|^2 + \frac{1}{\rho_1} \|T^2 x\|^2) + 2(\rho_2 \|x\|^2 + \frac{1}{\rho_2} \|\Lambda x\|^2) \\
&\leq 2(\rho_1 \|x\|^2 + \frac{1}{\rho_1} \|T^2 x\|^2) + 2\rho_2 \|x\|^2 + \frac{4}{\rho_2} (\|T^2 x\|^2 + \|Sx\|^2) \\
&\leq 2(\rho_1 + \rho_2 + \frac{2\|S\|^2}{\rho_2}) \|x\|^2 + 2(\frac{1}{\rho_1} + \frac{2}{\rho_2}) \|T^2 x\|^2 \\
&\leq \nu_1 \|x\|^2 + \nu_2 \|T^2 x\|^2,
\end{aligned}$$

for some $\nu_1, \nu_2 > 0$, $i = 1, 2$ and all $x \in \mathcal{D}(A^2)$. Since ρ_1 and ρ_2 are arbitrary, we can choose $\nu_2 < 1$.

(4) Now assume that $\mathcal{D}(\Upsilon^{\frac{1}{2}}) \subset \mathcal{D}(T)$. It follows that

$$\|Tx\| \leq a \|x\| + b \left\| \Upsilon^{\frac{1}{2}} x \right\| \quad (4.7)$$

for all $x \in \mathcal{D}(\Upsilon^{\frac{1}{2}})$ and for some nonnegative constants a and b . By (4.6), we obtain

$$\|Tx\|^2 \leq 2a(1 + \rho_2) \|x\|^2 + \frac{2b}{\rho_2} \|\Upsilon x\|^2,$$

for all $x \in \mathcal{D}(\Upsilon)$ and an arbitrary $\rho_2 > 0$. Thus

$$\left\| T(t + \Upsilon^{\frac{1}{2}})^{-1} x \right\|^2 \leq 2a(1 + \rho_2) \left\| (t + \Upsilon^{\frac{1}{2}})^{-1} x \right\|^2 + \frac{2b}{\rho_2} \left\| \Upsilon(t + \Upsilon^{\frac{1}{2}})^{-1} x \right\|^2,$$

for all $x \in \mathcal{H}$.

Hence

$$\left\| T(t + \Upsilon^{\frac{1}{2}})^{-1} \right\|^2 \leq \frac{2a}{t^2} (1 + \rho_2) + \frac{2b}{\rho_2} \left\| \Upsilon(t + \Upsilon^{\frac{1}{2}})^{-1} \right\|^2.$$

Letting t to $+\infty$, we assert that

$$M = \sup_{t>0} \left\| T(t + \Upsilon^{\frac{1}{2}})^{-1} \right\| < \frac{2b}{\rho_2}.$$

(cf. [37, Proposition 2.12]). Since ρ_2 is arbitrary, we can choose it such that $\frac{2b}{\rho_2} < 1$. Since T is m -accretive and $\Upsilon^{\frac{1}{2}}$ is m - $(\pi/4)$ -accretive, then Z_1 is m -accretive. By [16, Theorem IX-1.24], the factor $-Z_1$ generates holomorphic C_0 -semigroup $\mathcal{T}_1(z)$ of angle $\frac{\pi}{4}$.

On the other hand, since $\Upsilon^{\frac{1}{2}}$ is m - $(\pi/4)$ -accretive and $-T$ satisfy (4.7), by [16, Theorem IX-2.4], for any $\varepsilon > 0$, there exist nonnegative constants r and s such that $a, b < s$ and $(Z_2 + r)$ is the generator of holomorphic C_0 -semigroup $\mathcal{T}_2(z)$ of angle $\frac{\pi}{2} - \frac{\pi}{4} - \varepsilon$. This implies that $-(Z_2 + r)$ is m - ψ -accretive with $\psi = \frac{\pi}{4} + \varepsilon$.

(5) It follows from the item (4), $W(Z_1)$ is contained in the right half complex plan and $W(Z_2)$ in the left side with a non zero distance between their closure.

(6) We have $\mathcal{D}(T^2) \subset \mathcal{D}(Z_1) = \mathcal{D}(Z_2) = \mathcal{D}(\Upsilon^{\frac{1}{2}}) \subset \mathcal{D}(T)$.

The fact that $T(\mathcal{D}(T^2)) \subset \mathcal{D}(T^2)$ and $\Upsilon^{\frac{1}{2}}(\mathcal{D}(T^2)) \subset \mathcal{D}(T^2)$, we have $\mathcal{D}(T^2) \subset \mathcal{D}(T\Upsilon^{\frac{1}{2}})$, $\mathcal{D}(T^2) \subset \mathcal{D}(\Upsilon^{\frac{1}{2}}T)$ and $\mathcal{D}(T^2) \subset \mathcal{D}(Z_1^2)$. Now, by items (1), (2) and (3), we can easily verify that

$$Z_1^2x - TZ_1x - Z_1Tx - Sx = 0,$$

for all $x \in \mathcal{D}(T^2)$, hence on $\mathcal{D}(T^2)$, we have

$$\begin{aligned} Q(\lambda) &= Q(\lambda) - (Z_1^2 - TZ_1 - Z_1T - S) \\ &= \lambda^2I - 2\lambda T - S - Z_1^2 + TZ_1 + Z_1T + S \\ &= \lambda^2I - Z_1^2 - T(\lambda - Z_1) - (\lambda - Z_1)T \\ &= \frac{1}{2}(\lambda - Z_1)(\lambda I + Z_1 - 2T) + \frac{1}{2}(\lambda I + Z_1 - 2T)(\lambda - Z_1) \\ &= \frac{1}{2}(\lambda I - Z_1)(\lambda I - Z_2) + \frac{1}{2}(\lambda I - Z_2)(\lambda I - Z_1). \end{aligned}$$

This gives the form (4.3). Now, if $TS = ST$ on $\mathcal{D}(T^2)$, then $\Upsilon T = T\Upsilon$. Thus $\Upsilon^{\frac{1}{2}}$ commutes with T on $\mathcal{D}(T^2)$, which implies that (4.4). \square

Now the fact that $\mathcal{N}(\Upsilon^{\frac{1}{2}}) = \mathcal{N}(\Upsilon)$ and $\overline{\mathcal{R}(\Upsilon^{\frac{1}{2}})} = \overline{\mathcal{R}(\Upsilon)}$, by Corollary 3.5 and Theorem 3.2, we have

Corollary 4.2. *Let T^2 be m -accretive, T accretive with closed range and S is bounded and accretive. If $\mathcal{R}(S) \subseteq \mathcal{R}(T)$ and $\|(T^\dagger)^2S\| < 1$ (or $\mathcal{N}(T) \subseteq \mathcal{N}(S)$ and $\|S(T^2)^\dagger\| < 1$). Then*

- $\mathcal{R}(Z_1) = \mathcal{R}(Z_2) = \mathcal{R}(\Upsilon^{\frac{1}{2}}) = \mathcal{R}(T)$ is closed and $\mathcal{N}(Z_1) = \mathcal{N}(Z_2) = \mathcal{N}(\Upsilon^{\frac{1}{2}}) = \mathcal{N}(T)$.
- $\mathcal{H} = \mathcal{R}(\Upsilon^{\frac{1}{2}}) \oplus \mathcal{N}(\Upsilon^{\frac{1}{2}})$.
- Z_1, Z_2 and $\Upsilon^{\frac{1}{2}}$ are EP operators.
- If further, T is injective, then Z_1, Z_2 and $\Upsilon^{\frac{1}{2}}$ are invertible.

An immediate consequence of this corollary, the operator Z_1 and Z_2 are written in a matrices form with respect to mutually orthogonal subspaces decomposition as follows

$$Z_1 = \begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix} : \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix} \longrightarrow \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix};$$

and

$$Z_2 = \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} : \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix} \longrightarrow \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix};$$

with A and B on $\mathcal{R}(T) \cap \mathcal{D}(T)$ are injective operators with closed range. In this case, if $AB = BA$, we have

$$Q(\lambda) = \begin{bmatrix} (\lambda - A)(\lambda - B) & 0 \\ 0 & \lambda^2 \end{bmatrix} : \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix} \longrightarrow \begin{bmatrix} \mathcal{R}(T) \\ \mathcal{N}(T) \end{bmatrix}.$$

Also, $Q(0)$ is Moore-penrose invertible and $Q^\dagger(0) = Z_1^\dagger Z_2^\dagger = A^{-1}B^{-1}$. If T is injective, then $Q(0)$ is invertible and $Q^{-1}(0) = Z_1^{-1}Z_2^{-1}$. Consequently, S is also invertible. The

block Vandermonde operator corresponding to Z_1, Z_2 is given by

$$\mathcal{V}(Z_1, Z_2) = \begin{bmatrix} I & I \\ Z_1 & Z_2 \end{bmatrix}.$$

We have,

$$\mathcal{V}(Z_1, Z_2) = \begin{bmatrix} I & 0 \\ Z_2 & I \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & Z_1 - Z_2 \end{bmatrix} \begin{bmatrix} I & I \\ 0 & I \end{bmatrix},$$

where the left and right factors on the right-hand side are invertible. So $\mathcal{V}(Z_1, Z_2)$ is invertible if and only if $2\Upsilon^{\frac{1}{2}} = (Z_1 - Z_2)$ is invertible. Now, we apply [22, Corollary 29.12, Corollary 29.13 and Remark 29.14] taking in account that $\mathcal{N}(\Lambda^{\frac{1}{2}}) = \mathcal{N}((\Lambda^{\frac{1}{2}})^*)$, we obtain,

$$\sigma(Z_1) \cap \sigma(Z_2) = \emptyset \quad \text{and} \quad \sigma(Z_1) \cup \sigma(Z_2) = \sigma(Q(.)).$$

5. AN APPLICATION TO A SECOND ORDER LINEAR BOUNDARY VALUE PROBLEM

Denote $[0, +\infty)$ by \mathbb{R}_+ , and let $C^k(\mathbb{R}_+, \mathcal{D})$ be the set of all k -times (strongly) continuously differentiable functions mapping \mathbb{R}_+ , into $\mathcal{D} \subseteq \mathcal{H}$. In this section we consider the following abstract second order linear boundary value problem,

$$u''(t) - 2Tu'(t) - Su(t) = 0, \quad t \in (0, 1), \quad (5.1)$$

$$u(0) = u_0, \quad u(1) = u_1 \quad (5.2)$$

where $u' = \frac{d}{dt}$.

Theorem 5.1. *Let T^2 be m -accretive, T is accretive and S is bounded and accretive. Assume that*

- (1) $\mathcal{D}(\Upsilon^{\frac{1}{2}}) \subset \mathcal{D}(T)$.
- (2) $T(\mathcal{D}(T^2)) \subset \mathcal{D}(T^2)$ and $\Upsilon^{\frac{1}{2}}(\mathcal{D}(T^2)) \subset \mathcal{D}(T^2)$.
- (3) T commutes with $\Upsilon^{1/2}$ on $\mathcal{D}(T^2)$.

Then of any constant vectors $u_0, u_1 \in \mathcal{D}(T)$ the vector valued function,

$$u(t) = e^{-(1-t)Z_1}x_0 + e^{tZ_2}x_1, \quad t \in (0, 1),$$

with

$$x_0 = (I - e^{-2\Upsilon^{\frac{1}{2}}})^{-1} [-e^{Z_2}u_0 + u_1]$$

and

$$x_1 = (I - e^{-2\Upsilon^{\frac{1}{2}}})^{-1} [u_0 - e^{-Z_1}u_1]$$

is uniquely determined solution of (5.1)-(5.2), with $u(\cdot) \in C^\infty((0, 1), \mathcal{H}) \cap C^1((0, 1), \mathcal{D}(T))$.

Proof. Under the assumptions, by Theorem 4.1, the factors $-Z_1$ and Z_2 generates bounded holomorphic C_0 -semigroups. By [5, Lemma 2.38],

$$x(t) = e^{-(1-t)Z_1}x_0 \in \mathcal{D}(Z_1)^k$$

and

$$y(t) = e^{tZ_2}x_1 \in \mathcal{D}(Z_2)^k,$$

for all $k \in \mathbb{N}$, $x_0, x_1 \in \mathcal{H}$ and $t \in (0, 1)$. This implies that

$$u(t) = x(t) + y(t) \in \mathcal{H},$$

$$u'(t) = Z_1x(t) + Z_2y(t) \in \mathcal{D}(T)$$

and

$$u^{(2)}(t) = Z_1^2x(t) + Z_2^2y(t)$$

for all $t \in (0, 1)$. We can easily see that u verifies (5.1). Since $(I - e^{-2\gamma\frac{1}{2}})^{-1}$ and $(I - e^{-2\gamma\frac{1}{2}})^{-1}$ exist and bounded on \mathcal{H} . Thus, we have

$$u(0) = e^{-Z_1}x_0 + x_1 = u_0,$$

and

$$u(1) = x_0 + e^{-Z_2}x_1 = u_1,$$

This completes the proof. \square

Example 5.2. Let Ω is a smooth bounded domain of \mathbb{R}^n , Γ be the boundary of Ω , and $\xi \in \mathbb{C}$ with $\text{Re}(\xi) \geq 0$, $\eta \geq 0$, $\eta_1 \in \mathbb{R}$. We consider the following initial-boundary value problem in $L^2(\Omega)$,

$$u''(t, x) - 2(\eta\Delta - i\eta_1\Delta^2)u'(t, x) - \xi u(t, x) = 0, \quad (t, x) \in (0, 1) \times \Omega, \quad (5.3)$$

$$u(0, x) = u_0, \quad u(1, x) = u_1, \quad x \in \Omega, \quad (5.4)$$

$$u|_{\Gamma} = \Delta u|_{\Gamma} = 0, \quad (5.5)$$

where Δ denotes the Laplacian. It is known that $-\Delta$ with domain $H_0^1(\Omega) \cap H^2(\Omega) = \{u \in H^2(\Omega); u|_{\Gamma} = 0\}$ is a positive, self-adjoint operator on $L^2(\Omega)$. Its $L^2(\Omega)$ - normalized eigenfunctions are denoted w_j , and its eigenvalues counted with their multiplicities are denoted λ_j :

$$-\Delta w_j = \lambda_j w_j. \quad (5.6)$$

It is well known that $0 < \lambda_1 \leq \dots \leq \lambda_j \rightarrow \infty$. Functional calculus can be defined using the eigenfunction expansion. In particular; if we denote by $(-\Delta)^\alpha$ the fractional powers of the Dirichlet Laplacian, with $0 \leq \alpha \leq 1$, then

$$(-\Delta)^\alpha u = \sum_{j=1}^{\infty} \lambda_j^\alpha u_j w_j$$

with

$$u_j = \int_{\Omega} u(x) w_j(y) dx$$

for $u \in \mathcal{D}((-\Delta)^\alpha) = \{u : (\lambda_j^\alpha u_j) \in \ell^2(\mathbb{N})\}$. Also, if $m\alpha$ is an integer, then

$$\mathcal{D}((-\Delta)^{m\alpha}) = H_0^{m\alpha}(\Omega) \cap H^{2m\alpha}(\Omega), \quad m \geq 1. \quad (5.7)$$

See [21], for more details.

Set

$$T = \eta\Delta - i\eta_1\Delta^2, \quad \mathcal{D}(T) = \mathcal{D}(\Delta^2) = \{u \in H^4(\Omega); u|_{\Gamma} = \Delta u|_{\Gamma} = 0\}.$$

and

$$Su = \xi u, \quad \mathcal{D}(S) = L^2(\Omega)$$

Then the abstract version of problem (5.3)-(5.4) takes the form (5.1)-(5.2). We have,

- S is bounded and accretive.
- T is m-accretive.

- The operator

$$T^2 = \eta^2 \Delta^2 + \eta_1^2 \Delta^4 - 2i\eta\eta_1 \Delta^3$$

with

$$\mathcal{D}(T^2) = \mathcal{D}(\Delta^4) = \{u \in H^8(\Omega); u|_\Gamma = \Delta u|_\Gamma = \Delta^2 u|_\Gamma = \Delta^3 u|_\Gamma = 0\}$$

is m-accretive. Indeed, T^2 is densely defined and closed, and $(T^2)^* = \eta^2 \Delta^2 + \eta_1^2 \Delta^4 + 2i\eta\eta_1 \Delta^3$ is accretive.

- The operator

$$\Upsilon = T^2 + S = \eta^2 \Delta^2 + \eta_1^2 \Delta^4 - 2i\eta\eta_1 \Delta^3 + \xi,$$

with domain $\mathcal{D}(\Delta^4)$, is m-accretive.

- Υ admits a square root $\Upsilon^{1/2}$ m- $(\pi/4)$ -accretive with $\mathcal{D}(\Delta^4)$ is a core of $\Upsilon^{1/2}$.
- The operators factors

$$Z_1 = T + \Upsilon^{\frac{1}{2}} = \eta\Delta - i\eta_1\Delta^2 + (\eta^2\Delta^2 + \eta_1^2\Delta^4 - 2i\eta\eta_1\Delta^3 + \xi)^{\frac{1}{2}}$$

and

$$Z_2 = T - \Upsilon^{\frac{1}{2}} = \eta\Delta - i\eta_1\Delta^2 - (\eta^2\Delta^2 + \eta_1^2\Delta^4 - 2i\eta\eta_1\Delta^3 + \xi)^{\frac{1}{2}}$$

with domain $\mathcal{D}(\Delta^2)$ are closed operators.

- By an argument of functional calculus, we obtain

$$Tu = - \sum_{j=1}^{\infty} (\eta + i\eta_1\lambda_j)\lambda_j u_j w_j$$

for $u \in \mathcal{D}(\Delta^2)$. If $(\eta, \eta_1) \neq (0, 0)$ then T is injective. Thus, is invertible with

$$T^{-1}u = - \sum_{j=1}^{\infty} \frac{1}{(\eta + i\eta_1\lambda_j)\lambda_j} u_j w_j$$

for $u \in \mathcal{D}(\Delta^2)$.

- Assume that $(\eta, \eta_1) \neq (0, 0)$ and $\xi \neq 0$. If $\sum_{j=1}^{\infty} \frac{1}{(\eta + i\eta_1\lambda_j)^2 \lambda_j^2} < \frac{1}{|\xi|}$ then $\|ST^{-2}\| < 1$. In fact,

$$\begin{aligned} \|T^{-1}u\|^2 &= \langle T^{-2}u, u \rangle \\ &= \sum_{j=1}^{\infty} \frac{1}{(\eta + i\eta_1\lambda_j)^2 \lambda_j^2} |\langle u, w_j \rangle|^2 \\ &\leq \sum_{j=1}^{\infty} \frac{1}{(\eta + i\eta_1\lambda_j)^2 \lambda_j^2} \|u\|^2 \\ &< \frac{1}{|\xi|} \|u\|^2. \end{aligned}$$

This implies that

$$\|ST^{-2}\| \leq \|S\| \|T^{-1}\|^2 = |\xi| \|T^{-1}\|^2 < 1.$$

- If we assume that $(\eta, \eta_1) \neq (0, 0)$, $\xi \neq 0$ and $\sum_{j=1}^{\infty} \frac{1}{(\eta + i\eta_1\lambda_j)^2\lambda_j^2} < \frac{1}{|\xi|}$; then by Theorem 3.2 and Corollary 4.2, we conclude that $\Upsilon^{\frac{1}{2}}$, Z_1 and $-Z_2$ are m - $(\pi/4)$ -accretive invertible operators. In particular, Z_1 and $-Z_2$ generates holomorphic C_0 -semigroup of contraction operators $\mathcal{T}_1(z)$ and $\mathcal{T}_2(z)$ of angle $\frac{\pi}{4}$. If we assume further T commutes with $\Upsilon^{\frac{1}{2}}$ on $\mathcal{D}(T^2)$. Then all the statements of Theorem 5.1 hold. Consequently, for any pair of vectors $u_0, u_1 \in \mathcal{D}(T)$ the vector valued function,

$$u(t, x) = e^{-(1-t)Z_1}v_0(x) + e^{tZ_2}v_1(x), \quad t \in (0, 1), \quad x \in \Omega$$

with

$$v_0(x) = (I - e^{-2\Upsilon^{\frac{1}{2}}})^{-1} [-e^{Z_2}u_0(x) + u_1(x)] \quad x \in \Omega$$

and

$$v_1(x) = (I - e^{-2\Upsilon^{\frac{1}{2}}})^{-1} [u_0(x) - e^{-Z_1}u_1(x)] \quad x \in \Omega$$

is the unique solution of (5.3)-(5.5).

REFERENCES

- [1] Yu. M. Arlinskiĭ and A. B. Popov, *On sectorial matrices*, Linear Algebra Appl. 370 (2003), 133–146.
- [2] A. V. Balakrishnan, *Fractional powers of closed operators and the semi-groups generated by them*, Pacific J. Math. 10 (1960), 419–437.
- [3] M. Benharrat, *Right and left quotient of two bounded operators on Hilbert spaces*, Commun. Korean Math. Soc. Vol. 35, No. 2, (2020) 547–563.
- [4] A. Ben-Israel and T. N. E. Greville, *Generalized Inverses: Theory and Applications*. Second edition, Springer, New York, 2003.
- [5] E. B. Davies, *One Parameter Semigroups*. Academic Press, London (1980).
- [6] R. deLaubenfels, *Powers of Generators of Holomorphic Semigroups*, Proc. Amer. Math. Soc. Vol. 99, No. 1 (Jan., 1987), pp. 105–108.
- [7] C. Engstrom, A. Torshage, *On equivalence and linearization of operator matrix functions with unbounded entries*, Integral Equations and Operator Theory, 89(4) (2017), 465–492.
- [8] D. Eschwé M. Langer, *Variational principles for eigenvalues of selfadjoint operator functions*, Integral Equations Operator Theory 49 (2004), 287–321.
- [9] K. Gustafson, D. Rao, *Numerical Range, the Field of Values of Linear Operators and Matrices*, Springer, New York, 1997.
- [10] K. Harbarth, H. Langer, *A factorization theorem for operator pencils*, Integral Equations Operator Theory 2 (1979), 344–364
- [11] Q. Huang and W. Zhai, *Perturbations and expressions for generalized inverses in Banach spaces and Moore-Penrose inverses in Hilbert spaces of closed linear operators*, Linear Algebra Appl. 435 (2011), no. 1, 117–127.
- [12] Q. Huang, L. Zhu, J. Yu, *Some new perturbation results for generalized inverses of closed linear operators in Banach spaces*. Banach J. Math. Anal. 6 (2) (2012), 58–68.
- [13] M. Hayashi and T. Ozawa, *On Landau-Kolmogorov inequalities for dissipative operators*, Proc. Amer. Math. Soc. 145 (2017), 847–852.
- [14] H. Qianglian, Z. Lanping, Y. Jiena, *Some new perturbation results for generalized inverses of closed linear operators in Banach spaces*. Banach J. Math. Anal. 6 no. 2 (2012), 58–68
- [15] M.G. Krein and H. Langer, *On the theory of quadratic pencils of self-adjoint operators*, Dokl. Akad. Nauk SSSR 154 (1964), 1258–1261 (Russian); English transl., Soviet Math. Dokl. 5 (1964), 266–269.
- [16] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, New York (1995).
- [17] T. Kato, *Fractional powers of dissipative operators*, Proc. Japan Acad. 13 (3) (1961), 246–274.
- [18] T. Kato, *On an Inequality of Hardy, Littlewood, and Pólya*, Advances in Math. 7 (1971), 217–218.

- [19] T. Kurmayya, G. Ramesh, *Cone nonnegativity of Moore–Penrose inverses of unbounded Gram operators*. Ann. Funct. Anal. 7 (2016), no. 2, 338–347.
- [20] H. Langer, *Factorization of operator pencils*. Acta. Scient. Math. 38(1976), 83–96.
- [21] J. L. Lions, E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications V II*, Springer (1972)
- [22] A. S. Markus, *Introduction to the spectral theory of polynomial operator pencils*, volume 71 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1988.
- [23] Martinez, C., Sanz, M., Marco, L.: Fractional powers of operators, J. Math. Soc. Japan **40**, 331–347 (1988).
- [24] M. Möller, V. Pivovarchik, *Spectral Theory of Operator Pencils, Hermite-Biehler Functions, and their Applications*, Birkhäuser (2015).
- [25] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Berlin-Heidelberg-New York: Springer 1983.
- [26] R. S. Phillips, *Dissipative Operators and Hyperbolic Systems of Partial Differential Equations*, Transactions of the American Mathematical Society Vol. 90, No. 2 (Feb., 1959), pp. 193-254
- [27] S. Ôta, *Closed linear operators with domain containing their range*, Proc. Edinburgh Math. Soc. 27(1984), 229–233.
- [28] L. Rodman, *An introduction to operator polynomials*, Birkhauser Basel (1989).
- [29] A. A. Shkalikov, *Strongly damped pencils of operators and solvability of the corresponding operator-differential equations*, Math. USSR Sb. 63 (1989), 97-119.
- [30] J. G. Stampfli, *Minimal range theorems for operators with thin spectra*, Pac. J. Math., 23 (1967), 601–612.
- [31] G. Wang, Y. Wei, and S. Qiao, *Generalized Inverses: Theory and Computations*. Developments in Mathematics, vol 53. Springer, Singapore (2018).
- [32] A.F.M. ter Elsta, M. Sauter, H. Vogt, *A generalisation of the form method for accretive forms and operators*, J. Funct. Anal., 269 (3) (2015), 705-744.
- [33] T.-J. Xiao and J. Liang, *The Cauchy Problem for Higher-Order Abstract Differential Equations*, Lecture Notes in Mathematics, 1701. Springer-Verlag, Berlin, 1998.
- [34] T.-J. Xiao and J. Liang, *Differential operators and C-wellposedness of complete second order abstract Cauchy problems*, Pacific J. Math. 186 (1) (1998), 167–200.
- [35] T.-J. Xiao, J. Liang. *Pazy-type characterization for differentiability of propagators of higher order Cauchy problems in Banach spaces*. Discrete Contin. Dyn. Syst. Ser. A 5 (3) (1999), 651-662.
- [36] T.-J. Xiao and J. Liang, *Higher order abstract Cauchy problems: their existence and uniqueness families*, J. London Math. Soc. (2) 67 (1) (2003), 149–164.
- [37] A. Yoshikawa, *On Perturbation of closed operators in a Banach space*, J. Fac. Sci. Hokkaido Univ., 22 (1972), 50–61.

¹ DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ ORAN 1-AHMED BEN BELLA, BP 1524 ORAN-EL M'NAOUAR, 31000 ORAN, ALGÉRIE.

Email address: fairouzbouchelaghem@yahoo.fr

² DÉPARTEMENT DE GÉNIE DES SYSTÈMES, ECOLE NATIONALE POLYTECHNIQUE D'ORAN-AUDIN (EX. ENSET D'ORAN), BP 1523 ORAN-EL M'NAOUAR, 31000 ORAN, ALGÉRIE.

Email address: mohammed.benharrat@enp-oran.dz, mohammed.benharrat@gmail.com