

THE INCIDENCE VARIETY COMPACTIFICATION OF STRATA OF d -DIFFERENTIALS IN GENUS 0

DUC-MANH NGUYEN

WITH AN APPENDIX BY VINCENT KOZIARZ AND DUC-MANH NGUYEN

ABSTRACT. Given $d \in \mathbb{Z}_{\geq 2}$, for every $\underline{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n$ such that $k_i \geq 1 - d$ and $k_1 + \dots + k_n = -2d$, denote by $\Omega^d \mathcal{M}_{0,n}(\underline{k})$ and $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ the corresponding stratum of d -differentials in genus 0 and its projectivization respectively. We show that the incidence variety compactification of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ is isomorphic to the blow-up of $\overline{\mathcal{M}}_{0,n}$ along a specific sheaf of ideals. Along the way we obtain an explicit divisor representing the tautological line bundle on the incidence variety. In the case where none of the k_i is divisible by d , the self-intersection number of this divisor computes the volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$. We prove a recursive formula which allows one to compute the volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ from the volumes of other strata of lower dimension.

1. INTRODUCTION

1.1. Statements of the main results. Given a Riemann surface X , we will denote by K_X its canonical line bundle. Let g be a non-negative integer, and d a positive one. For any vector $\underline{k} = (k_1, \dots, k_n)$ of integers such that $k_i > -d$, and $k_1 + \dots + k_n = d(2g - 2)$, $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is the space of tuples (X, x_1, \dots, x_n, q) , where X is a compact Riemann surface of genus g , $\{x_1, \dots, x_n\}$ are n marked points on X , and q is a non-zero meromorphic d -differential on X (that is a meromorphic section of the bundle $K_X^{\otimes d}$) such that

$$\operatorname{div}(q) = \sum_{i=1}^n k_i x_i.$$

The space $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is called a *stratum* of d -differentials in genus g . There is a natural action of \mathbb{C}^* on $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ which consists in multiplying the differential q by a scalar in \mathbb{C}^* . We denote by $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ the quotient of $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ by this \mathbb{C}^* -action.

It is well known that $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ and $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ are complex algebraic orbifolds (see for instance [30, 52, 53, 5]). It is shown in [46] that $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ and $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ carry some natural volume forms that will be denoted by $d\operatorname{vol}$ and $d\operatorname{vol}_1$ respectively. In the case $d \in \{1, 2, 3, 4, 6\}$, $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ (resp. $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$) carries another particular volume form known as the *Masur-Veech volume*, which will be denoted by $d\operatorname{vol}^{MV}$ (resp. $d\operatorname{vol}_1^{MV}$) (see §6.2 below). The Masur-Veech volume is of interest because it is related to the enumeration of tilings of surfaces by triangles and squares. By construction, $d\operatorname{vol}$ and $d\operatorname{vol}_1$ differ from $d\operatorname{vol}^{MV}$ and $d\operatorname{vol}_1^{MV}$ by a multiplicative constant, which belongs to either \mathbb{Q} , or $\sqrt{3}\mathbb{Q}$ (see [46, Prop.5.9]).

Date: September 21, 2021.

The primary goal of this paper is to compute the volume of the strata of d -differentials in genus 0 where all the orders of the zeros and poles of the d -differentials are not multiples of d . To this purpose, we investigate a natural compactification of $\Omega^d \mathcal{M}_{0,n}(\underline{k})$ known as the *Incidence Variety Compactification* (see [26, 4, 5]). Denote by $\overline{\mathcal{M}}_{0,n}$ the Deligne-Mumford-Knudsen compactification of $\mathcal{M}_{0,n}$. There is a holomorphic vector bundle $\mathcal{H}_{0,n}^{(d)}$ over $\overline{\mathcal{M}}_{0,n}$ whose fiber over a point $(C, x_1, \dots, x_n) \in \overline{\mathcal{M}}_{0,n}$ is the space $H^0(C, d\omega_C + \sum_{i=1}^n (d-1)x_i)$, where ω_C is the dualizing sheaf of C . Note that we have (see §5.2)

$$\dim_{\mathbb{C}} H^0(C, d\omega_C + \sum_{i=1}^n (d-1)x_i) = (d-1)(n-2) - 1.$$

By definition, $\Omega^d \mathcal{M}_{0,n}(\underline{k})$ is contained in the complement of the zero section of $\mathcal{H}_{0,n}^{(d)}$. Hence $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ is a subset of the projective bundle $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ associated to $\mathcal{H}_{0,n}^{(d)}$. The Incidence Variety Compactification of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ is defined to be its closure in $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$. This compactification will be denoted by $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{0,n}(\underline{k})$.

Our first main result is an explicit description of $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ as a complex algebraic variety. Define

$$(1) \quad \mu_i := -\frac{k_i}{d}, \quad i = 1, \dots, n, \quad \text{and} \quad \mu := (\mu_1, \dots, \mu_n).$$

We will call μ_i the weight of the i -th marked point on the pointed curves parametrized by $\overline{\mathcal{M}}_{0,n}$. Observe that we have $\mu_i < 1$, and $\mu_1 + \dots + \mu_n = 2$.

Recall that the boundary $\partial \overline{\mathcal{M}}_{0,n} := \overline{\mathcal{M}}_{0,n} \setminus \mathcal{M}_{0,n}$ of $\overline{\mathcal{M}}_{0,n}$ is a simple normal crossing divisor. The irreducible components of $\partial \overline{\mathcal{M}}_{0,n}$ are in bijection with partitions of $\{1, \dots, n\}$ into two subsets $\{I_0, I_1\}$ such that $\min\{|I_0|, |I_1|\} \geq 2$. The irreducible component of $\partial \overline{\mathcal{M}}_{0,n}$ associated to \mathcal{S} will be denoted by $D_{\mathcal{S}}$. Denote by \mathcal{P} the set of partitions associated to the components of $\partial \overline{\mathcal{M}}_{0,n}$. For any $\mathcal{S} = \{I_0, I_1\} \in \mathcal{P}$, by convention we will always choose the numbering of the subsets I_0, I_1 such that

$$(2) \quad \sum_{i \in I_0} \mu_i \leq 1 \leq \sum_{i \in I_1} \mu_i \quad \Leftrightarrow \quad \sum_{i \in I_0} k_i \geq -d \geq \sum_{i \in I_1} k_i.$$

We will associate to each component $D_{\mathcal{S}}$ of $\partial \overline{\mathcal{M}}_{0,n}$ the weight $\mu_{\mathcal{S}}$ defined by

$$(3) \quad \mu_{\mathcal{S}} := \frac{1}{2} \cdot \left(\sum_{i \in I_1} \mu_i - \sum_{i \in I_0} \mu_i \right) = 1 - \sum_{i \in I_0} \mu_i.$$

Note that $\mu_{\mathcal{S}}$ is always non-negative. We warn the reader that our convention on the numbering of I_0, I_1 , and our definition of $\mu_{\mathcal{S}}$ are different from [34].

Consider now a point \mathbf{x} in $\overline{\mathcal{M}}_{0,n}$. Let $C_{\mathbf{x}}$ denote the stable curve parametrized by \mathbf{x} . Assume that $C_{\mathbf{x}}$ has $(r+1)$ irreducible components, which will be denoted by $C_{\mathbf{x}}^0, \dots, C_{\mathbf{x}}^r$. Let $\mathcal{N}(\mathbf{x})$ be the set of nodes of $C_{\mathbf{x}}$. Note that $|\mathcal{N}(\mathbf{x})| = r$. For any $\alpha \in \mathcal{N}(\mathbf{x})$, let $\mathcal{S}_{\alpha} := \{I_{0,\alpha}, I_{1,\alpha}\}$ be the associated partition of $\{1, \dots, n\}$. Recall that $\dim \overline{\mathcal{M}}_{0,n} = n-3$. Let $\mathcal{U}_{\mathbf{x}}$ be a neighborhood of \mathbf{x} which satisfies the following conditions

- (i) $\mathcal{U}_{\mathbf{x}}$ does not intersect any boundary divisor $D_{\mathcal{S}}$ such that $\mathcal{S} \notin \{\mathcal{S}_{\alpha}, \alpha \in \mathcal{N}(\mathbf{x})\}$,
- (ii) $\mathcal{U}_{\mathbf{x}}$ can be identified with an open subset of \mathbb{C}^{n-3} such that, for each $\alpha \in \mathcal{N}(\mathbf{x})$, there is a coordinate function t_{α} such that $D_{\mathcal{S}_{\alpha}} \cap \mathcal{U}_{\mathbf{x}}$ is defined by the equation $t_{\alpha} = 0$.

As $\overline{\mathcal{M}}_{0,n}$ is a projective variety, we can actually choose $\mathcal{U}_{\mathbf{x}}$ to be an open affine of $\overline{\mathcal{M}}_{0,n}$ such that t_α are elements of the coordinate ring of $\mathcal{U}_{\mathbf{x}}$.

Splitting a node $\alpha \in \mathcal{N}(\mathbf{x})$ into two points, we obtain two subcurves $\hat{C}_{\mathbf{x},\alpha}^0, \hat{C}_{\mathbf{x},\alpha}^1$ of $C_{\mathbf{x}}$, where $\hat{C}_{\mathbf{x},\alpha}^k$ contains the i -th marked points with $i \in I_{k,\alpha}$. Given $j \in \{0, \dots, r\}$, for each $\alpha \in \mathcal{N}(\mathbf{x})$, define

$$(4) \quad \beta_{j,\alpha} = \begin{cases} d\mu_{S_\alpha} & \text{if } C_{\mathbf{x}}^j \subset \hat{C}_{\mathbf{x},\alpha}^0, \\ 0 & \text{otherwise.} \end{cases}$$

Let

$$(5) \quad \beta_j = (\beta_{j,\alpha})_{\alpha \in \mathcal{N}(\mathbf{x})} \in (\mathbb{Z}_{\geq 0})^{\mathcal{N}(\mathbf{x})}, \quad \text{and} \quad t^{\beta_j} = \prod_{\alpha \in \mathcal{N}(\mathbf{x})} t_\alpha^{\beta_{j,\alpha}}.$$

Let $\mathcal{O}_{\overline{\mathcal{M}}_{0,n}}$ be the structure sheaf of $\overline{\mathcal{M}}_{0,n}$, and $\mathcal{I}_{\mathcal{U}_{\mathbf{x}}}$ be the ideal sheaf of $\mathcal{O}_{\overline{\mathcal{M}}_{0,n}|_{\mathcal{U}_{\mathbf{x}}}}$ generated by $\{t^{\beta_0}, \dots, t^{\beta_r}\}$. We will prove

Theorem 1.1.

- (i) The family $\{\mathcal{I}_{\mathcal{U}_{\mathbf{x}}}, \mathbf{x} \in \overline{\mathcal{M}}_{0,n}\}$ defines a sheaf of ideals \mathcal{I} of $\mathcal{O}_{\overline{\mathcal{M}}_{0,n}}$.
- (ii) The incidence variety compactification $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ is isomorphic to the blow-up $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ of $\overline{\mathcal{M}}_{0,n}$ along \mathcal{I} .
- (iii) Let $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$ be the blow-up projection. For every $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$ the fiber $\hat{p}^{-1}(\{\mathbf{x}\})$ is always a projective space, whose dimension is determined by \underline{k} and the stratum of $\overline{\mathcal{M}}_{0,n}$ to which \mathbf{x} belongs.
- (iv) Every irreducible component of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k}) := \widehat{\mathcal{M}}_{0,n}(\underline{k}) \setminus \mathcal{M}_{0,n}$ is a divisor. The set of irreducible components of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$ is in bijection with the set $\hat{\mathcal{P}}(\mu)$ of partitions \mathcal{S} of $\{1, \dots, n\}$ such that either
 - $\mathcal{S} = \{I_0, I_1\}$, and $\sum_{i \in I_0} \mu_i = \sum_{i \in I_1} \mu_i = 1$, or
 - $\mathcal{S} = \{I_0, I_1, \dots, I_r\}$, $\sum_{i \in I_0} \mu_i < 1$ and $\sum_{i \in I_j} \mu_i > 1$ for all $j = 1, \dots, r$.

Remark 1.2.

- The blow-up $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ can be equal to $\overline{\mathcal{M}}_{0,n}$ for instance in the case $k_i < 0$ for all $i = 1, \dots, n$. In §3.3, we specify a necessary and sufficient condition for $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ to be isomorphic to $\overline{\mathcal{M}}_{0,n}$.
- A stratum of $\overline{\mathcal{M}}_{0,n}$ is a subset that parametrizes stable n -pointed curves (of genus 0) having the same dual graph. For each $\mathcal{S} = \{I_0, \dots, I_r\} \in \hat{\mathcal{P}}(\mu)$, there is a distinguished tree $T_{\mathcal{S}}$ associated to \mathcal{S} constructed as follows: the vertices of $T_{\mathcal{S}}$ are labeled by I_0, \dots, I_r , and every edge of $T_{\mathcal{S}}$ joins the vertex labeled by I_0 to one of the vertices labeled by I_1, \dots, I_r . Denote by $D_{\mathcal{S}}^*$ the stratum of $\overline{\mathcal{M}}_{0,n}$ corresponding to $T_{\mathcal{S}}$. Then the irreducible component of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$ associated to \mathcal{S} , denoted by $\hat{D}_{\mathcal{S}}$, is the closure of $\hat{p}^{-1}(D_{\mathcal{S}}^*)$ in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. In the case $\mathcal{S} \in \mathcal{P}$, $\hat{D}_{\mathcal{S}}$ is actually the proper transform of the boundary divisor $D_{\mathcal{S}} \subset \partial \overline{\mathcal{M}}_{0,n}$.

Our second main result shows that for all strata $\Omega^d \mathcal{M}_{0,n}(\underline{k})$ such that d does not divide any of the k_i , $i \in \{1, \dots, n\}$, $\text{vol}_1(\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k}))$ can be computed by the self-intersection number of an explicit

divisor in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. Consider the following \mathbb{Q} -divisor in $\overline{\mathcal{M}}_{0,n}$

$$(6) \quad \mathcal{D}_\mu := \frac{d}{2} \left(\sum_{i=1}^n -\mu_i \psi_i + \sum_{S \in \mathcal{P}} (1 - \mu_S) D_S \right) = \frac{d}{(n-2)(n-1)} \sum_{S \in \mathcal{P}} (|I_0| - 1)(|I_1| - 1 - (n-1)\mu_S) \cdot D_S.$$

This divisor actually represents a line bundle $\bar{\mathcal{L}}_\mu$ over $\overline{\mathcal{M}}_{0,n}$. By construction $\hat{p}^{-1} \mathcal{I} \cdot \mathcal{O}_{\widehat{\mathcal{M}}_{0,n}(\underline{k})}$ is isomorphic to the line bundle associated to a Cartier divisor \mathcal{E} in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ such that $\text{supp}(\mathcal{E}) = \hat{p}^{-1}(\text{supp}(\mathcal{I}))$. We will call \mathcal{E} the exceptional divisor of the blowup $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$. Define

$$(7) \quad \hat{\mathcal{D}}_\mu := \hat{p}^* \mathcal{D}_\mu + \mathcal{E}.$$

Theorem 1.3. *The restriction of the tautological line bundle $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ to $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is isomorphic to the line bundle associated to $\hat{\mathcal{D}}_\mu$. Moreover, if d does not divide k_i , for all $i = 1, \dots, n$, then*

$$(8) \quad \text{vol}_1(\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})) = \frac{(-1)^{n-3}}{d^{n-2}} \cdot \frac{(2\pi)^{n-2}}{2^{n-2}(n-2)!} \cdot \hat{\mathcal{D}}_\mu^{n-3}$$

where $\hat{\mathcal{D}}_\mu^{n-3}$ means the self-intersection number $\underbrace{\hat{\mathcal{D}}_\mu \cdots \hat{\mathcal{D}}_\mu}_{n-3}$ of $\hat{\mathcal{D}}_\mu$ in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$.

Remark 1.4.

- In the cases where $k_i < 0$, for all $i = 1, \dots, n$, we have $\widehat{\mathcal{M}}_{0,n}(\underline{k}) \simeq \overline{\mathcal{M}}_{0,n}$, $\hat{\mathcal{D}}_\mu \simeq \mathcal{D}_\mu$, and (8) is the content of [34, Th. 1.1].
- For general \underline{k} , the fact that $\hat{\mathcal{D}}_\mu^{n-3}$ computes the volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ follows from the results of [10]. The novelty of Theorem 1.3 is that we have an explicit expression of the divisor $\hat{\mathcal{D}}_\mu$.
- In Appendix §A, we provide the self-intersection number $\hat{\mathcal{D}}_\mu^{n-3}$ for all the strata $\Omega^d \mathcal{M}_{0,n}(\underline{k})$, where $n \in \{4, 5\}$, and $d \in \{3, 4, 6\}$. We also compute the ratio $\frac{d \text{vol}^{MV}}{d \text{vol}}$ and the Masur-Veech volumes of those strata, which give the asymptotics of the counting of tilings of the sphere by triangles or squares.

For large values of n , a direct computation of $\hat{\mathcal{D}}_\mu^{n-3}$ by intersection theory seems to be an unrealistic task. To compute $\hat{\mathcal{D}}_\mu^{n-3}$ more efficiently we prove a recursive formula which relates $\hat{\mathcal{D}}_\mu^{n-3}$ to the self-intersection number of the divisors representing the tautological line bundles on strata of lower dimensions. Specifically, for each $k \in \mathbb{Z}_{\geq 3}$, let

$$L_k = \{v = (v_1, \dots, v_k) \in \mathbb{Q}^k, \sum_{i=1}^k v_i = 2, \text{ and } v_i < 1, \text{ for all } i = 1, \dots, k\}.$$

We define a map $\mathcal{J}_k : L_k \rightarrow \mathbb{Z}$ as follows: given $v = (v_1, \dots, v_k) \in L_k$, there is a smallest natural number e such that $ev_i \in \mathbb{Z}$, for all $i = 1, \dots, k$. Set $\ell_i = -ev_i$. We then have $\ell_1 + \dots + \ell_k = -2e$, $\text{gcd}(e, \ell_1, \dots, \ell_k) = 1$, and $\ell_i > -e$, $i = 1, \dots, k$. Let $\underline{\ell} := (\ell_1, \dots, \ell_k)$. Let $\Omega^e \mathcal{M}_{0,k}(\underline{\ell})$ be the stratum of e -differentials associated to the vector $\underline{\ell}$. We can consider the closure $\mathbb{P}\Omega^e \overline{\mathcal{M}}_{0,k}(\underline{\ell})$ of $\mathbb{P}\Omega \mathcal{M}_{0,k}(\underline{\ell})$ in $\mathbb{P}\mathcal{H}_{0,k}^{(e)}$. By Theorem 1.1, we know that $\mathbb{P}\Omega \overline{\mathcal{M}}_{0,k}(\underline{\ell})$ is isomorphic to the blow-up $\widehat{\mathcal{M}}_{0,k}(\underline{\ell})$ of $\overline{\mathcal{M}}_{0,k}$. We then define

$$(9) \quad \mathcal{J}_k(v) := c_1^{k-3}(\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,k}^{(e)}}) \cap [\widehat{\mathcal{M}}_{0,k}(\underline{\ell})],$$

where $[\widehat{\mathcal{M}}_{0,k}(\mathcal{L})]$ is the class of $\widehat{\mathcal{M}}_{0,k}(\mathcal{L})$ in $A_*(\mathbb{P}\mathcal{H}_{0,k}^{(e)})$. Note that by definition $\mathcal{J}_3(\nu) = 1$, for all $\nu \in L_3$.

In what follows, given $I \subset \{1, \dots, n\}$, we define $\mu(I) := \sum_{i \in I} \mu_i$. As usual $|I|$ is the cardinality of I . We consider the following families of partitions of $\{1, \dots, n\}$

- \mathcal{T}_{1a} is the set of partitions $\{I_0, I_1\}$, where $\mu(I_0) < 1 < \mu(I_1)$, $|I_0| = 2$ and $\mu(I_0) \notin \mathbb{Z}$.
- \mathcal{T}_{1b} is the set of partitions $\{I_{00}, I_{01}, I_1\}$ such that
 - $|I_{00}| = 2$, $\mu(I_{00}) = 1$,
 - $|I_{01}| = 1$, $\mu(I_{01}) < 0$,
 - $|I_1| \geq 2$, $\mu(I_1) > 1$, and $\mu(I_1) \notin \mathbb{Z}$.
- \mathcal{T}_{2a} is the set of partitions $\{I_0, I_1, I_2\}$, where I_0, I_1, I_2 satisfy
 - $|I_0| = 1$, $\mu(I_0) < 0$,
 - $|I_j| \geq 2$, $\mu(I_j) > 1$, and $\mu(I_j) \notin \mathbb{Z}$, for $j = 1, 2$.
- \mathcal{T}_{2b} is the set of partitions $\{I_{01}, I_{02}, I_1, I_2\}$ of $\{1, \dots, n\}$ such that
 - $|I_{01}| = |I_{02}| = 1$,
 - $|I_j| \geq 2$, $\mu(I_j) > 1$, and $\mu(I_j) \notin \mathbb{Z}$ for $j = 1, 2$,
 - $\mu(I_1) + \mu(I_{01}) = \mu(I_2) + \mu(I_{02}) = 1$.

We denote by \mathcal{T} the union $\mathcal{T}_{1a} \cup \mathcal{T}_{1b} \cup \mathcal{T}_{2a} \cup \mathcal{T}_{2b}$. Elements of \mathcal{T} will be called *primary partitions* with respect to μ .

- For $\mathcal{S} = \{I_0, I_1\} \in \mathcal{T}_{1a}$ or $\mathcal{S} = \{I_{00}, I_{01}, I_1\} \in \mathcal{T}_{1b}$, we define
 - $e(\mathcal{S})$ is the smallest natural number such that $e(\mathcal{S})\mu_i \in \mathbb{Z}$, for all $i \in I_1$,
 - $m(\mathcal{S}) := d(\mu(I_1) - 1)$,
 - $\nu(\mathcal{S})$ is the vector in $\mathbb{Q}^{|I_1|+1}$ whose $|I_1|$ first coordinates are given by $\{\mu_i, i \in I_1\}$, and the last coordinate is $2 - \mu(I_1)$.
- For $\mathcal{S} = \{I_0, I_1, I_2\} \in \mathcal{T}_{2a}$ or $\mathcal{S} = \{I_{01}, I_{02}, I_1, I_2\} \in \mathcal{T}_{2b}$,
 - $n_1(\mathcal{S}) = |I_1|, n_2(\mathcal{S}) = |I_2|$,
 - $m_1(\mathcal{S}) = d(\mu(I_1) - 1), m_2(\mathcal{S}) = d(\mu(I_2) - 1), m(\mathcal{S}) = m_1(\mathcal{S})m_2(\mathcal{S})$.
 - For $j \in \{1, 2\}$, $e_j(\mathcal{S})$ is the smallest natural number such that $e_j(\mathcal{S})\mu_i \in \mathbb{Z}$ for all $i \in I_j$.
 - $\nu_j(\mathcal{S})$ is the vector in $\mathbb{Q}^{|I_j|+1}$ whose $|I_j|$ first coordinates are given by $\{\mu_i, i \in I_j\}$, and the last coordinate is $2 - \mu(I_j)$.
 - For $\mathcal{S} \in \mathcal{T}_{2b}$, we define

$$\epsilon(\mathcal{S}) := \begin{cases} 1 & \text{if } \mu(I_{01}) \neq \mu(I_{02}) \\ 2 & \text{if } \mu(I_{01}) = \mu(I_{02}). \end{cases}$$

Theorem 1.5. *We have*

$$(10) \quad \begin{aligned} \mathcal{J}_n(\mu) &= \sum_{\mathcal{S} \in \mathcal{T}_{1a}} a_{\mathcal{S}} \cdot \left(\frac{d}{e(\mathcal{S})} \right)^{n-4} \cdot \mathcal{J}_{n-1}(\nu(\mathcal{S})) - \sum_{\mathcal{S} \in \mathcal{T}_{1b}} a_{\mathcal{S}} \cdot \left(\frac{d}{e(\mathcal{S})} \right)^{n-5} \cdot \mathcal{J}_{n-2}(\nu(\mathcal{S})) \\ &\quad - \sum_{\mathcal{S} \in \mathcal{T}_{2a}} a_{\mathcal{S}} \cdot \frac{d^{n-5}}{e_1^{n_1(\mathcal{S})-2}(\mathcal{S})e_2^{n_2(\mathcal{S})-2}(\mathcal{S})} \cdot \mathcal{J}_{n_1(\mathcal{S})+1}(\nu_1(\mathcal{S}))\mathcal{J}_{n_2(\mathcal{S})+1}(\nu_2(\mathcal{S})) \\ &\quad + \sum_{\mathcal{S} \in \mathcal{T}_{2b}} a_{\mathcal{S}} \cdot \frac{d^{n-6}}{e_1^{n_1(\mathcal{S})-2}(\mathcal{S})e_2^{n_2(\mathcal{S})-2}(\mathcal{S})} \cdot \mathcal{J}_{n_1(\mathcal{S})+1}(\nu_1(\mathcal{S}))\mathcal{J}_{n_2(\mathcal{S})+1}(\nu_2(\mathcal{S})), \end{aligned}$$

where the coefficient a_S is given by

$$a_S = \begin{cases} \frac{d(n-3)}{(n-1)(n-2)} - \frac{m}{n-2} & \text{if } S \in \mathcal{T}_{1a} \\ \frac{d(n-3)}{(n-1)(n-2)} \cdot m & \text{if } S \in \mathcal{T}_{1b} \\ \frac{d(n_1 n_2 (m_1 + m_2) - m_1 n_1 - m_2 n_2)}{(n-1)(n-2)} - \frac{m}{n-2} & \text{if } S \in \mathcal{T}_{2a} \\ \epsilon \cdot \frac{d n_1 n_2}{(n-1)(n-2)} \cdot m & \text{if } S \in \mathcal{T}_{2b}. \end{cases}$$

(for simplicity, we write n_i, m_i, m, ϵ in the place of $n_i(S), m_i(S), m(S), \epsilon(S)$ respectively).

As applications, we will use (10) to recover some results of [3] (see §9.3).

1.2. Motivations and related works. Abelian differentials and quadratic differentials on Riemann surfaces are central objects in Teichmüller theory, dynamics in moduli spaces, billiards in rational polygons, interval exchange transformations among others. For some surveys on these fascinating fields of research, we refer to [40, 57, 56]. From the flat metric point of view, d -differentials are natural generalizations of translation surfaces and half-translation surfaces, the objects associated to Abelian and quadratic differentials. For $d \in \{1, 2, 3, 4, 6\}$, d -differentials also arise from the counting of triangulations and quadrangulations of surfaces (see for instance [51, 12, 11, 35]).

In view of applications, it is important to have an adequate compactification of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$. The incidence variety compactification $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{g,n}(\underline{k})$ of strata of Abelian differentials and strata of d -differentials were introduced in [26, 4, 5]. The characterizing properties of points in $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{g,n}(\underline{k})$ are given in [4] and [5]. However, a global description of $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{g,n}(\underline{k})$ has not been known.

Another approach to study $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is to use flat metrics with conical singularities. In [51], Thurston introduced the moduli spaces of flat metrics on the sphere with cone angles prescribed at the singularities, where all the angles are supposed to be smaller than 2π . Those spaces can be interpreted as strata $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$, where all the k_i satisfy $1 - d \leq k_i \leq -1$. Thurston showed that $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ can be endowed with a complex hyperbolic metric which is not complete in general, and the associated metric completions are complex hyperbolic cone manifolds. It follows in particular that the volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ can be computed by the integral of the volume form associated to the complex hyperbolic metric. These volumes have been computed by different methods in [43] and in [34]. Theorem 1.3 is in fact a generalization of the results in [34] to the cases where some of the orders k_i are positive.

The space $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ can also be viewed as the moduli space of n -weighted pointed genus zero smooth curves, where the weight of the i -th marked point is $\mu_i = -\frac{k_i}{d}$. When the weights μ_i satisfy $0 < \mu_i < 1$, moduli compactifications of such spaces have been studied in [29]. In this perspective, it would be interesting to know whether similar moduli compactifications can be constructed in the case some of the weights are negative, and to clarify the connection between the incidence variety compactification $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ and those moduli compactifications of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$.

For $d \in \{1, 2, 3, 4, 6\}$, an important numerical invariant of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is its Masur-Veech volume (see §6.3). The Masur-Veech volumes of strata of Abelian differentials and quadratic differentials (that is $d \in \{1, 2\}$) have been extensively studied in view of applications to billiards and Teichmüller dynamics (see for instance [18, 19, 14, 41, 3, 27, 47, 9]). It has been also known that the volume of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$, when none of the k_i is a multiple of d , is closely related to the self-intersection number of the tautological divisor on $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{g,n}(\underline{k})$ (see [47, 9, 10]). A recursive formula to compute this self-intersection number is proposed in [48, 49]. However, to the author's knowledge, no explicit value of

the Masur-Veech volume of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ was known in the cases where $d \in \{3, 4, 6\}$, and some of the k_i are positive.

Finally, we would like to mention a beautiful formula computing the Masur-Veech volumes of all strata of quadratic differentials in genus zero which was conjectured by Kontsevich and proved by Athreya-Eskin-Zorich in [3]. The proof of this formula in [3] uses in an essential way a result of Eskin-Kontsevich-Zorich [13] on the sum of the Lyapunov exponents of the Teichmüller geodesic flow. The authors of [3] asked whether one can bypass the result of [13] in proving Kontsevich's formula. We are hopeful that the results of the present paper may contribute to an independent proof of this formula providing an affirmative answer to the challenge raised by the authors of [3].

1.3. Plan of the proofs. We now give the plan of the proofs of the main results. We first observe that $\Omega^d \mathcal{M}_{0,n}(\underline{k})$ is the complement of the zero section in the total space of a line bundle \mathcal{L}_μ over $\mathcal{M}_{0,n}$, the fiber $\mathcal{L}_{\mu,\mathbf{x}}$ over a point $\mathbf{x} \sim (\mathbb{P}^1, x_1, \dots, x_n)$ being the line generated by the d -differential $q_{\mathbf{x}} = \prod_{i=1}^n (z - x_i)^{k_i} (dz)^d$. Since the poles of $q_{\mathbf{x}}$ have order at most $d - 1$, \mathcal{L}_μ is actually a line subbundle of $\mathcal{H}_{0,n|\mathcal{M}_{0,n}}^{(d)}$. By taking the projectivization, we get a section $\mathbf{f} : \mathcal{M}_{0,n} \rightarrow \mathbb{P}\mathcal{H}_{0,n|\mathcal{M}_{0,n}}^{(d)}$ of the projective bundle $\mathbb{P}\mathcal{H}_{0,n|\mathcal{M}_{0,n}}^{(d)}$ whose image is precisely $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$.

We would like to extend \mathcal{L}_μ to a line subbundle of $\mathcal{H}_{0,n}^{(d)}$ over $\overline{\mathcal{M}}_{0,n}$. To this end, we consider the Kawamata line bundle \mathcal{K}_μ on the universal curve $\overline{C}_{0,n}$ (see Definition 2.1). In Theorem 2.2, we show that there is a line bundle $\tilde{\mathcal{L}}_\mu$ over $\overline{\mathcal{M}}_{0,n}$ such that $\tilde{\mathcal{L}}_{\mu|\mathcal{M}_{0,n}} \sim \mathcal{L}_\mu$, and $p^* \tilde{\mathcal{L}}_\mu \sim \mathcal{K}_\mu$, where $p : \overline{C}_{0,n} \rightarrow \overline{\mathcal{M}}_{0,n}$ is the natural projection. Since $\tilde{\mathcal{L}}_\mu \sim p_* \mathcal{K}_\mu$, by using the Riemann-Roch-Grothendieck formula, we show that $\tilde{\mathcal{L}}_\mu$ is isomorphic to the line bundle associated to the divisor \mathcal{D}_μ defined in (6).

In fact, to prove Theorem 2.2, we construct for every $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$ and a neighborhood $\mathcal{U}_{\mathbf{x}} \subset \overline{\mathcal{M}}_{0,n}$ as in §1.1 an explicit family of $(r + 1)$ meromorphic sections Φ_0, \dots, Φ_r of $K_{\overline{C}_{0,n}/\overline{\mathcal{M}}_{0,n}}$ such that $t^{\beta_0} \Phi_0, \dots, t^{\beta_r} \Phi_r$, where t^{β_j} are defined in (4), are all trivializing sections of $\mathcal{K}_{\mu|p^{-1}(\mathcal{U}_{\mathbf{x}})}$. Those sections will play a particularly important role in our proof.

By construction, the restriction of $t^{\beta_j} \Phi_j$ to the fiber $C_{\mathbf{x}} \simeq p^{-1}(\{\mathbf{x}\})$ is a collection of d -differentials each of which is defined on an irreducible component of $C_{\mathbf{x}}$. When there is at least one non-zero differential, this collection can be interpreted as an element of the fiber of $\mathcal{H}_{0,n}^{(d)}$ over \mathbf{x} . But it may happen that all of the differentials in this collection vanish identically. This means that $\tilde{\mathcal{L}}_\mu$ is not a line subbundle of $\mathcal{H}_{0,n}^{(d)}$.

In Theorem 3.1 we will show that the data $(\mathcal{U}_{\mathbf{x}}, t^{\beta_0}, \dots, t^{\beta_r})$ patch together to give a well defined sheaf of ideals \mathcal{I} of the structure sheaf of $\overline{\mathcal{M}}_{0,n}$. We then consider the blow-up of $\overline{\mathcal{M}}_{0,n}$ along \mathcal{I} , denoted by $\widehat{\mathcal{M}}_{0,n}(\underline{k})$, which is a projective variety equipped with a birational surjective morphism $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$. In §3, we undertake a study in depth of the geometry of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. In particular, we will show that the fiber of \hat{p} over a point \mathbf{x} is isomorphic to $\mathbb{P}_{\mathbb{C}}^{r_0-1}$, where r_0 is the number of principal subcurves of $C_{\mathbf{x}}$ (see Definition 3.4 and Theorem 3.13). Knowledge about the fiber of \hat{p} allows us to show that the set of irreducible components of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$ is in bijection with the set of partitions in $\hat{\mathcal{P}}(\mu)$.

Once $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is constructed, in Theorem 5.2, we show that there is a natural embedding $\hat{\mathbf{f}} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \mathbb{P}\mathcal{H}_{0,n}^{(d)}$ extending the section \mathbf{f} . The image of $\hat{\mathbf{f}}$ is precisely the closure of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ in $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ showing that $\mathbb{P}\Omega^d \widehat{\mathcal{M}}_{0,n}(\underline{k})$ is isomorphic to $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. This completes the proof of Theorem 1.1.

Let $\hat{\mathcal{L}}_\mu$ denote the pullback of $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ to $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. In Propostion 7.2 we show that $\hat{\mathcal{L}}_\mu \sim \hat{p}^* \tilde{\mathcal{L}}_\mu + \mathcal{E}$, where we abusively denote by \mathcal{E} the line bundle associated to the exceptional divisor of the blow-up $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$. By definition, $\hat{\mathcal{L}}_\mu$ is represented by the divisor $\hat{\mathcal{D}}_\mu$.

There is a natural Hermitian metric on the line bundle \mathcal{L}_μ over $\mathcal{M}_{0,n}$ called the Hodge norm. The curvature form of the Hodge norm is denoted by Θ . Assuming that none of the k_i is a multiple of d , we have (see Lemma 6.3)

$$(11) \quad \int_{\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})} d\text{vol}_1 = \frac{(-1)^{n-3}}{d^{n-2}} \cdot \frac{2\pi}{2^{n-2}(n-2)!} \int_{\mathcal{M}_{0,n}} (i\Theta)^{n-3}.$$

Based on the construction of [6], in [10] Costantini-Möller-Zachhuber construct a compactification $\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ which is a complex algebraic orbifold. There is a birational surjective morphism $\mathcal{P} : \mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \mathbb{P}\Omega^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ which is identity over $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$. Identifying $\mathbb{P}\Omega^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ with $\widehat{\mathcal{M}}_{0,n}(\underline{k})$, we can consider \mathcal{P} as a birational map from $\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ onto $\widehat{\mathcal{M}}_{0,n}(\underline{k})$.

The space $\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ carries a tautological line bundle denoted by $\mathcal{O}(-1)_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})}$ which satisfies $\mathcal{P}^* \hat{\mathcal{L}}_\mu \sim \mathcal{O}(-1)_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})}^{\otimes d}$. It also follows from a result in [10] that $(\frac{1}{2\pi}\Theta)^k$ defines a representative in the sense of currents of $c_1^k(\mathcal{O}(-1)_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})}^{\otimes d})$, for all $k \in \mathbb{N}$. Hence the right hand side of (11) can be computed by the self-intersection number of the divisor $\hat{\mathcal{D}}_\mu$ in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$, which proves Theorem 1.3.

Since $\hat{\mathcal{D}}_\mu$ represents the restriction of the tautological line bundle $\hat{\mathcal{L}}_\mu$ on $\widehat{\mathcal{M}}_{0,n}(\underline{k})$, we have

$$(\hat{\mathcal{D}}_\mu)^N = c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{\mathcal{D}}_\mu]$$

where $[\hat{\mathcal{D}}_\mu]$ denotes the class of $\hat{\mathcal{D}}_\mu$ in $A_*(\widehat{\mathcal{M}}_{0,n}(\underline{k}))$. Using (7), we can express $\hat{\mathcal{D}}_\mu$ as a combination of irreducible components of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$. Thus, to compute $(\hat{\mathcal{D}}_\mu)^N$, it suffices to compute $c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{\mathcal{D}}_S]$, for each $S \in \hat{\mathcal{P}}(\mu)$, where $\hat{\mathcal{D}}_S$ is the irreducible component of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$ associated to S .

We will partition $\hat{\mathcal{P}}(\mu)$ into two subsets: $\hat{\mathcal{P}}_0(\mu)$ is the set of partitions $\mathcal{S} = \{I_0, I_1\}$ such that $\mu(I_0) = \mu(I_1) = 1$, and $\hat{\mathcal{P}}_1(\mu)$ is the set of partions $\mathcal{S} = \{I_0, \dots, I_r\}$ such that $\mu(I_0) < 1$ and $\mu(I_j) > 1$ for $j = 1, \dots, r$. For $\mathcal{S} \in \hat{\mathcal{P}}_1(\mu)$ we will show that $\hat{\mathcal{D}}_S$ is birational to a projective bundle $\mathbb{P}E_S$ over the space

$$\overline{\mathcal{M}}_S = \overline{\mathcal{M}}_{0,n_0+r} \times \widehat{\mathcal{M}}_{0,n_1+1}(\underline{k}^{(1)}) \times \dots \times \widehat{\mathcal{M}}_{0,n_r+1}(\underline{k}^{(r)}),$$

where $n_j = |I_j|$, and the vector $\underline{k}^{(j)}$ is determined by $\{\mu_i, i \in I_j\}$. Moreover, the restriction of $\hat{\mathcal{L}}_\mu$ to $\hat{\mathcal{D}}_S$ can be identified with $\mathcal{O}(-1)_{\mathbb{P}E_S}$. From this, we derive that $c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{\mathcal{D}}_S]$ can be computed from $c_1^{n_j-2}(\hat{\mathcal{L}}_j) \cap [\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})]$, where $\hat{\mathcal{L}}_j$ is the tautological line bundle on $\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$. For a dimensional reason, it turns out that $c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{\mathcal{D}}_S] = 0$ unless $\mathcal{S} \in \mathcal{T}_{1a}$ or $\mathcal{S} \in \mathcal{T}_{2a}$.

For $\mathcal{S} = \{I_0, I_1\} \in \hat{\mathcal{P}}_0(\mu)$, we use a different strategy to compute the divisor representative of the restriction of $\hat{\mathcal{L}}_\mu$ to $\hat{\mathcal{D}}_S$. Let D_S^* be the stratum of codimension 1 in $\overline{\mathcal{M}}_{0,n}$ associated to \mathcal{S} . We can consider D_S^* as an open dense subset of $\hat{\mathcal{D}}_S$. Denote by $\partial \hat{\mathcal{D}}_S$ the complement of D_S^* in $\hat{\mathcal{D}}_S$. Every

irreducible component of $\partial\hat{D}_S$ is a divisor in \hat{D}_S , which is in fact the intersection of \hat{D}_S with another divisor $\hat{D}_{S'}$, for some $S' \in \hat{\mathcal{P}}(\mu) \setminus \{S\}$. There is a natural trivializing section σ of $\hat{\mathcal{L}}_\mu$ over D_S^* . This section extends to a meromorphic section of $\hat{\mathcal{L}}_\mu$ over \hat{D}_S . By computing the vanishing order of σ along each component of $\partial\hat{D}_S$, we show that $\hat{\mathcal{L}}_{\mu|\hat{D}_S}$ is represented by an explicit divisor which is a combination of irreducible components of $\partial\hat{D}_S$.

Let $\hat{\mathcal{P}}(\mu, S)$ be the set of $S \in \hat{\mathcal{P}}(\mu) \setminus \{S\}$ such that $\hat{D}_{S',S} := \hat{D}_{S'} \cap \hat{D}_S \neq \emptyset$. It turns out that $c_1^{N-2}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S',S}]$ is non zero only if the partition of $\{1, \dots, n\}$ associated to $\hat{D}_{S',S}$ belongs to $\mathcal{T}_{1b} \cup \mathcal{T}_{2b}$. We obtain the recursive formula (10) by combining the calculations in two cases $S \in \hat{\mathcal{P}}_0(\mu)$ and $S \in \hat{\mathcal{P}}_1(\mu)$.

Organization. The paper is organized as follows: in §2, we define the Kawamata line bundle \mathcal{K}_μ , and show that \mathcal{K}_μ is the pullback of a line bundle $\tilde{\mathcal{L}}_\mu$ on $\overline{\mathcal{M}}_{0,n}$. That $\tilde{\mathcal{L}}_\mu$ is represented by the divisor \mathcal{D}_μ is deduced from the Grothendieck-Riemann-Roch formula.

In §3, we construct $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ and study the fibers of the blow-up projection $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$. Using the results in §3, in §4 we show that every irreducible component of $\partial\widehat{\mathcal{M}}_{0,n}(\underline{k})$ corresponds uniquely to a partition in $\hat{\mathcal{P}}(\mu)$. We investigate the geometric interpretation of the elements of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ in §5. This allows us to define an embedding $\hat{\mathbf{f}} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \mathbb{P}\mathcal{H}_{0,n}^{(d)}$, which realizes $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ as the closure of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ in $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ proving Theorem 1.1.

In §6, we recall the complex affine manifold (orbifold) structure and the volume forms on $\Omega^d \mathcal{M}_{0,n}(\underline{k})$. The proof of Theorem 1.3 is given in §7. In this proof we make use of the results in [10].

The last part of the paper is devoted to the proof of (10). In §8, we investigate the restriction of the tautological line bundle on $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ to its boundary. Our goal is to determine a divisor representing the tautological line bundle on each boundary component. This enables us to express the self-intersection number of $c_1(\hat{\mathcal{L}}_\mu)$ on $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ as a sum of (products of) self-intersections of the tautological line bundle on the incidence varieties of strata of d -differentials of lower dimension. The proof of Theorem 1.5 is given in §9. As an application of this result, we give an alternative proof of Kontsevich's formula (which was originally proved in [3]) for two families of strata of quadratic differentials. Finally, in Appendix A, which is a joint work with V. Koziarz, we give the explicit values of the self-intersection number of $\hat{\mathcal{D}}_\mu$ and of the Masur-Veech volume of the strata $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$, where $n \in \{4, 5\}$, $d \in \{2, 3, 4, 6\}$, and none of the k_i is divisible by d .

Acknowledgment: the author is indebted to Vincent Koziarz for the helpful and inspiring discussions, which played an important role in the realization of this work. He is grateful to Pascal Autissier, Eric Balandraud, and Yohan Brunearbe for their enlightening remarks and comments. He thanks Martin Möller and Adrien Sauvaget for the stimulating discussions.

The author is partly supported by the French ANR project ANR-19-CE40-0003.

2. THE KAWAMATA LINE BUNDLE

2.1. Definitions. Since holomorphic Abelian differentials on $\mathbb{P}_\mathbb{C}^1$ do not exist, in what follows we always suppose that $d > 1$. Let $p : \overline{\mathcal{C}}_{0,n} \rightarrow \overline{\mathcal{M}}_{0,n}$ be the universal curve over $\overline{\mathcal{M}}_{0,n}$, and $K_{\overline{\mathcal{C}}_{0,n}/\overline{\mathcal{M}}_{0,n}}$

the relative canonical line bundle. Let Γ_i , $i = 1, \dots, n$, be the image of $\overline{\mathcal{M}}_{g,n}$ by the i -th tautological section of p .

Let $\mathcal{S} = \{I_0, I_1\}$ be a partition in \mathcal{P} . A generic point $\mathbf{x} \in D_{\mathcal{S}}$ represents a stable curve $C_{\mathbf{x}}$ with two irreducible components $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^1$, where $C_{\mathbf{x}}^j$ contains the i -th marked point if and only if $i \in I_j$. By definition, $p^{-1}(D_{\mathcal{S}})$ is a divisor of $\overline{\mathcal{C}}_{0,n}$ with two irreducible components denoted by $F_{\mathcal{S}}^0$ and $F_{\mathcal{S}}^1$, where $F_{\mathcal{S}}^j \cap C_{\mathbf{x}} = C_{\mathbf{x}}^j$.

Definition 2.1. *The Kawamata line bundle is the following line bundle on $\overline{\mathcal{C}}_{0,n}$*

$$(12) \quad \mathcal{K}_{\mu} := d \left(K_{\overline{\mathcal{C}}_{0,n}/\overline{\mathcal{M}}_{0,n}} + \sum_{i=1}^n \mu_i \Gamma_i - \sum_{S \in \mathcal{P}} \mu_S F_S^0 \right).$$

The line bundle \mathcal{K}_{μ} was first introduced by Kawamata in [31]. Note also that our convention on the numbering I_0, I_1 is the converse of the one in [31].

For any $\mathbf{x} = (\mathbb{P}^1, x_1, \dots, x_n) \in \mathcal{M}_{0,n}$, $H^0(C_{\mathbf{x}}, \mathcal{K}_{\mu, \mathbf{x}})$ is generated by the d -differential $\prod_{i=1}^n (z - x_i)^{k_i} (dz)^d$. It follows that $p_* \mathcal{K}_{\mu|C_{0,n}}$ gives a line bundle \mathcal{L}_{μ} over $\mathcal{M}_{0,n}$. The complement of the zero section in the total space of \mathcal{L}_{μ} can be identified with $\Omega^d \mathcal{M}_{0,n}(k)$. Our goal in this section is to show the following

Theorem 2.2. *There is a line bundle $\tilde{\mathcal{L}}_{\mu}$ over $\overline{\mathcal{M}}_{0,n}$ such that $\mathcal{K}_{\mu} \sim p^* \tilde{\mathcal{L}}_{\mu}$ and $\tilde{\mathcal{L}}_{\mu| \mathcal{M}_{0,n}} \sim \mathcal{L}_{\mu}$. Moreover, $\tilde{\mathcal{L}}_{\mu}$ is isomorphic to the line bundle associated to the divisor*

$$\mathcal{D}_{\mu} := \frac{d}{2} \left(\sum_{i=1}^n -\mu_i \psi_i + \sum_{S \in \mathcal{P}} (1 - \mu_S) D_S \right).$$

Remark 2.3. *In the case all the weights μ_i are positive, Theorem 2.2 is a consequence of a result by Kawamata (see [31, Th.4]).*

2.2. A construction of the universal curve near the boundary of $\overline{\mathcal{M}}_{0,n}$. To prove Theorem 2.2, we will show that for any $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$ there exists a neighborhood \mathcal{U} of \mathbf{x} such that the restriction $\mathcal{K}_{\mu|p^{-1}(\mathcal{U})}$ is trivial, that is \mathcal{K}_{μ} admits trivializing sections on $p^{-1}(\mathcal{U})$. For $\mathbf{x} \in \mathcal{M}_{0,n}$ this is clear, so we only need to consider the case $\mathbf{x} \in \partial \overline{\mathcal{M}}_{0,n}$.

Let $\mathbf{x} \in \partial \overline{\mathcal{M}}_{0,n}$ be a point in a stratum of codimension r . This means that \mathbf{x} is contained in the intersection of r boundary divisors. As usual the marked points of $C_{\mathbf{x}}$ are denoted by x_i , $i = 1, \dots, n$. The stable curve $C_{\mathbf{x}}$ has $r+1$ components denoted by $C_{\mathbf{x}}^0, \dots, C_{\mathbf{x}}^r$. Let I_j denote the subset of $\{1, \dots, n\}$ that records the marked points contained in $C_{\mathbf{x}}^j$. Note that I_j can be empty.

The topology of $C_{\mathbf{x}}$ is encoded by its dual graph $T_{\mathbf{x}}$, which is a tree since $C_{\mathbf{x}}$ has genus zero. We label the vertices of $T_{\mathbf{x}}$ by $\{v_0, \dots, v_r\}$, where v_j represents the irreducible component $C_{\mathbf{x}}^j$. Each node of $C_{\mathbf{x}}$ corresponds to an edge of $T_{\mathbf{x}}$. Since $T_{\mathbf{x}}$ is a tree, there is at most one edge between any pair of its vertices. Therefore, a node of $C_{\mathbf{x}}$ corresponds uniquely to an unordered pair $\{j, j'\}$, with $j, j' \in \{0, \dots, r\}$ and $j \neq j'$. Let $\mathcal{N}(\mathbf{x})$ denote the set pairs $\{j, j'\}$ associated with the nodes of $C_{\mathbf{x}}$. The point on $C_{\mathbf{x}}^j$ that corresponds to a node $\{j, j'\}$ will be denoted by $y_{jj'}$. For all $j \in \{0, \dots, r\}$, let

$$\mathcal{N}_j(\mathbf{x}) := \{j' \in \{0, \dots, r\}, \{j, j'\} \in \mathcal{N}(\mathbf{x})\}, \quad \text{and} \quad n_j := |I_j| + |\mathcal{N}_j(\mathbf{x})|.$$

The pointed curve $\mathbf{x}_j := (C_{\mathbf{x}}^j, (x_i)_{i \in I_j}, (y_{jj'})_{j' \in \mathcal{N}_j(\mathbf{x})})$ represents an element of \mathcal{M}_{0,n_j} . A neighborhood of \mathbf{x} in its stratum is identified with $U_0 \times \cdots \times U_r$, where U_j is a neighborhood of \mathbf{x}_j in \mathcal{M}_{0,n_j} . Since $\dim \mathcal{M}_{0,n_j} = n_j - 3$, we can identify U_j with a neighborhood of $0 \in \mathbb{C}^{n_j-3}$.

Let $u_j = (u_{j,1}, \dots, u_{j,n_j-3})$ be a system of holomorphic coordinates on U_j . By definition, for all $u_j \in U_j$, the pointed curve parametrized by u_j is isomorphic to $(\mathbb{P}_{\mathbb{C}}^1, (a_{ji})_{i \in I_j}, (b_{jj'})_{j' \in \mathcal{N}_j})$, where a_{ji} is the marked point corresponding to x_i and $b_{jj'}$ is the point corresponding to the node $y_{jj'}$. We consider a_{ji} and $b_{jj'}$ as holomorphic functions of u_j . Note that the points $\{a_{ji}(u_j), i \in I_j\}$ and $\{b_{jj'}(u_j), j' \in \mathcal{N}_j(\mathbf{x})\}$ are pairwise distinct.

Let Δ be a small disc about $0 \in \mathbb{C}$. The set

$$\mathcal{U} := \Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \times \cdots \times U_r$$

can be identified with an open neighborhood of \mathbf{x} in $\overline{\mathcal{M}}_{0,n}$. Our goal now is to construct a family of n -pointed genus zero curves over \mathcal{U} degenerating to $C_{\mathbf{x}}$. Consider

$$\mathcal{X} := \underbrace{\mathbb{P}_{\mathbb{C}}^1 \times \cdots \times \mathbb{P}_{\mathbb{C}}^1}_{r+1} \times \mathcal{U}.$$

Let z_{j-1} be the inhomogeneous coordinate on the j -th $\mathbb{P}_{\mathbb{C}}^1$ factor of \mathcal{X} . For any $\alpha = \{j, j'\} \in \mathcal{N}(\mathbf{x})$, the coordinate on the factor α of $\Delta^{\mathcal{N}(\mathbf{x})}$ will be denoted by t_{α} . Let $\pi_j : \mathcal{X} \rightarrow \mathbb{P}_{\mathbb{C}}^1$ be the projection from \mathcal{X} onto its $(j+1)$ -th $\mathbb{P}_{\mathbb{C}}^1$ factor, for $j = 0, \dots, r$. Let us write $z = (z_0, \dots, z_r)$, $t = (t_{\alpha})_{\alpha \in \mathcal{N}(\mathbf{x})}$, $u = (u_0, \dots, u_r)$.

Let $\overline{C}_{\mathcal{U}}$ denote the set of points $(z, t, u) \in \mathcal{X}$ which satisfy

$$(13) \quad (z_j - b_{jj'})(z_{j'} - b_{j'j}) = t_{\alpha}, \quad \forall \alpha = \{j, j'\} \in \mathcal{N}(\mathbf{x}).$$

Let $p : \overline{C}_{\mathcal{U}} \rightarrow \mathcal{U}$ be the natural projection. Given $t := (t_{\alpha})_{\alpha \in \mathcal{N}(\mathbf{x})} \in \Delta^{\mathcal{N}(\mathbf{x})}$, and $u := (u_j)_{0 \leq j \leq r} \in U_0 \times \cdots \times U_r$, let $C(t, u)$ be the curve in $(\mathbb{P}_{\mathbb{C}}^1)^{r+1}$ defined by the equations (13). By construction, we have $C(0, 0) \simeq C_{\mathbf{x}}$.

Let $Z(t) := \{\alpha \in \mathcal{N}(\mathbf{x}), t_{\alpha} = 0\}$. The elements of $Z(t)$ are in bijection with the nodes of $C(t, u)$. Remove from the tree $T_{\mathbf{x}}$ the edges representing the elements of $Z(t)$, we get $(r(t) + 1)$ subtrees of $T_{\mathbf{x}}$, where $r(t) := |Z(t)|$. Denote those subtrees by $T_{\mathbf{x}}^k(t), k = 0, \dots, r(t)$. Let $J_0(t), \dots, J_r(t)$ denote the partition of $\{0, 1, \dots, r\}$ such that $j \in J_k(t)$ if and only if $v_j \in T_{\mathbf{x}}^k(t)$ (recall that v_0, \dots, v_r are the vertices of $T_{\mathbf{x}}$). The following lemma provides some basic properties of $C(t, u)$, its proof is left to the reader.

Lemma 2.4.

- (i) *The curve $C(t, u)$ has $r(t) + 1$ irreducible components, each of which is associated with a subtree in the family $\{T_{\mathbf{x}}^0(t), \dots, T_{\mathbf{x}}^{r(t)}(t)\}$. The component associated with $T_{\mathbf{x}}^k(t)$ will be denoted by $C^k(t, u)$.*
- (ii) *Given $k \in \{0, \dots, r(t)\}$, for every $j \in J_k(t)$, the restriction of π_j to $C^k(t, u)$ realizes an isomorphism from $C^k(t, u)$ onto $\mathbb{P}_{\mathbb{C}}^1$.*
- (iii) *If $j \notin J_k(t)$, then the projection π_j maps $C^k(t, u)$ to a point that corresponds to a node of $C(t, u)$.*

We now construct the tautological sections of p . For every $i \in \{1, \dots, n\}$, there is a unique $j \in \{0, \dots, r\}$ such that $i \in I_j$. Recall that by construction, we have a holomorphic map $a_{ji} : U_j \rightarrow \mathbb{P}_{\mathbb{C}}^1$

which associates to $u_j \in U_j$ the marked point corresponding to x_i on the curve parametrized by u_j . Let $k \in \{0, \dots, r(t)\}$ be such that $j \in J_k(t)$. By Lemma 2.4(ii) the projection π_j restricts to an isomorphism from $C^k(t, u)$ onto $\mathbb{P}_{\mathbb{C}}^1$. Denote by $\varphi_{ji}(t, u)$ the unique point on $C^k(t, u)$ that is mapped to $a_{ji}(u_j)$ by π_j . Define

$$\begin{aligned} \sigma_i : \mathcal{U} &\rightarrow \overline{C}_{\mathcal{U}} \\ (t, u) &\mapsto (\varphi_{ji}(t, u), t, u). \end{aligned}$$

Then $\{\sigma_i, i = 1, \dots, n\}$ are the tautological sections of the universal curve restricted to \mathcal{U} . This means that the pointed curve parametrized by $(t, u) \in \mathcal{U}$ is isomorphic to $(C(t, u), (\sigma_i(t, u))_{1 \leq i \leq n})$.

For all $i \in \{1, \dots, n\}, \ell \in \{0, \dots, r\}$, $\pi_{\ell} \circ \sigma_i(t, u)$ is the $(\ell + 1)$ -th coordinate of the i -th marked point on $C(t, u) \subset (\mathbb{P}_{\mathbb{C}}^1)^{r+1}$. By definition, if $i \in I_j$, then $a_{ji} = \pi_j \circ \sigma_i(t, u)$. To lighten the notation, we will write $a_{\ell i}(t, u) := \pi_{\ell} \circ \sigma_i(t, u)$ for all $\ell \in \{0, \dots, r\}$.

2.3. Trivializing sections of the Kawamata line bundle. For $j = 0, \dots, r$, consider the following meromorphic d -differential on \mathcal{X}

$$(14) \quad \Phi_j(t, u) = \prod_{i=1}^n (z_j - a_{ji})^{k_i} (dz_j)^d.$$

In this subsection we will prove

Theorem 2.5. *Let $t^{\beta_j}, j = 0, \dots, r$, be the monomials defined in (5). Then for all $j \in \{0, 1, \dots, r\}$, the d -differential $t^{\beta_j} \Phi_j$ gives a trivializing section of the Kawamata line bundle \mathcal{K}_{μ} over $\overline{C}_{\mathcal{U}}$.*

2.3.1. Trivializing section in a neighborhood of a generic point in the boundary. To illustrate the ideas of the proofs of Theorem 2.5, we first prove a special case, where \mathbf{x} is a generic point in the boundary of $\overline{\mathcal{M}}_{0,n}$. Consider a generic point \mathbf{x} in a boundary divisor $D_S \subset \partial \overline{\mathcal{M}}_{0,n}$, where $S = \{I_0, I_1\} \in \mathcal{P}$. Let $n_0 = |I_0|, n_1 = |I_1|$. Without loss of generality, we can assume that $I_0 = \{1, \dots, n_0\}$, and $I_1 = \{n_0 + 1, \dots, n\}$. The fiber $p^{-1}(\mathbf{x})$ is isomorphic to a reducible genus zero curve $C_{\mathbf{x}}$ which has two components denoted by $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^1$. The marked points on $C_{\mathbf{x}}$ will be denoted by $x_i, i = 1, \dots, n$, where $x_i \in C_{\mathbf{x}}^0$ for $1 \leq i \leq n_0$, and $x_i \in C_{\mathbf{x}}^1$ for $n_0 + 1 \leq i \leq n$. The point in $C_{\mathbf{x}}^0$ (resp. in $C_{\mathbf{x}}^1$) that corresponds to the unique node of $C_{\mathbf{x}}$ is denoted by y_{01} , (resp. by y_{10}).

Since $C_{\mathbf{x}}^0$ is a genus zero curve, we can identify $C_{\mathbf{x}}^0$ with $\mathbb{P}_{\mathbb{C}}^1$ such that $y_{01} = 0, x_{n_0-1} = 1, x_{n_0} = \infty$. Similarly, we can identify $C_{\mathbf{x}}^1$ with $\mathbb{P}_{\mathbb{C}}^1$ such that $y_{10} = 0, x_{n-1} = 1, x_n = \infty$. Let U_0 be a neighborhood of the point (x_1, \dots, x_{n_0-2}) in $(\mathbb{C} \setminus \{0, 1\})^{n_0-2}$, and U_1 a neighborhood of $(x_{n_0+1}, \dots, x_{n-2}) \in (\mathbb{C} \setminus \{0, 1\})^{n_1-2}$. Since $x_i \neq x_j$ if $i \neq j$, for all $u_0 = (u_{0,1}, \dots, u_{0,n_0-2}) \in U_0$, we can assume that $u_{0,i} \neq u_{0,j}$ if $i \neq j$. Similarly, for all $u_1 = (u_{1,1}, \dots, u_{1,n_1-2}) \in U_1$, we have $u_{1,i} \neq u_{1,j}$ if $i \neq j$.

Let Δ be a small disc about 0 in \mathbb{C} . Define

$$\mathcal{U} := \Delta \times U_0 \times U_1 \quad \text{and} \quad \overline{C}_{|\mathcal{U}} := \{(z_0, z_1, t, u_0, u_1) \in \mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1 \times \mathcal{U}, z_0 z_1 = t\}$$

(here we use the identification $\mathbb{P}_{\mathbb{C}}^1 \simeq \mathbb{C} \cup \{\infty\}$).

For $i = 0, 1$, let π_i denote the canonical projection from $\mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1$ onto its $(i + 1)$ -th factor. Given $t \in \Delta$, let C_t be the curve in $\mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1$ defined by the equation $z_0 z_1 = t$. For all $t \in \Delta^* := \Delta \setminus \{0\}$, the restrictions of π_0 and π_1 to C_t realize two isomorphisms between C_t and $\mathbb{P}_{\mathbb{C}}^1$. For $t = 0$, C_0 is the

reducible curve $\mathbb{P}_{\mathbb{C}}^1 \times \{0\} \cup \{0\} \times \mathbb{P}_{\mathbb{C}}^1$. Define

$$\begin{aligned} \sigma_0 : \Delta \times \mathbb{C}^* &\rightarrow \mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1 \\ (t, x) &\mapsto (x, t/x) \end{aligned}$$

and

$$\begin{aligned} \sigma_1 : \Delta \times \mathbb{C}^* &\rightarrow \mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1 \\ (t, x) &\mapsto (t/x, x). \end{aligned}$$

By convention, we set $\sigma_0(t, \infty) = (\infty, 0)$ and $\sigma_1(t, \infty) = (0, \infty)$. Note that $\sigma_0(t, x)$ (resp. $\sigma_1(t, x)$) is the unique point in C_t that is mapped to x by π_0 (resp. by π_1).

For $u_0 := (u_{0,1}, \dots, u_{0,n_0-2}) \in U_0$, define

$$\sigma_0(t, u_0) := (\sigma_0(t, u_{0,1}), \dots, \sigma_0(t, u_{0,n_0-2}), \sigma_0(t, 1), \sigma_0(t, \infty)).$$

Similarly, for $u_1 := (u_{1,1}, \dots, u_{1,n_1-2}) \in U_1$, define

$$\sigma_1(t, u_1) := (\sigma_1(t, u_{1,1}), \dots, \sigma_1(t, u_{1,n_1-2}), \sigma_1(t, 1), \sigma_1(t, \infty)).$$

The pointed curve parametrized by $(t, u_0, u_1) \in \Delta \times U_0 \times U_1$ is isomorphic to $(C_t, \sigma_0(t, u_0), \sigma_1(t, u_1))$. Note that if $t \in \Delta^*$ then

$$(C_t, \sigma_0(t, u_0), \sigma_1(t, u_1)) \stackrel{\pi_0}{\sim} (\mathbb{P}_{\mathbb{C}}^1, u_{0,1}, \dots, u_{0,n_0-2}, 1, \infty, \frac{t}{u_{1,1}}, \dots, \frac{t}{u_{1,n_1-2}}, t, 0)$$

and

$$(C_t, \sigma_0(t, u_0), \sigma_1(t, u_1)) \stackrel{\pi_1}{\sim} (\mathbb{P}_{\mathbb{C}}^1, \frac{t}{u_{0,1}}, \dots, \frac{t}{u_{0,n_0-2}}, t, 0, u_{1,1}, \dots, u_{1,n_1-2}, 1, \infty).$$

By convention, set $u_{0,n_0-1} = u_{1,n_1-1} = 1$, and $u_{0,n_0} = u_{1,n_1} = \infty$. We then have

$$a_{0,i}(t, u_0, u_1) = \begin{cases} u_{0,i}, & \text{for } i = 1, \dots, n_0, \\ \frac{t}{u_{1,i-n_0}}, & \text{for } i = n_0 + 1, \dots, n, \end{cases}$$

and,

$$a_{1,i}(t, u_0, u_1) = \begin{cases} \frac{t}{u_{0,i}}, & \text{for } i = 1, \dots, n_0, \\ u_{1,i-n_0}, & \text{for } i = n_0 + 1, \dots, n. \end{cases}$$

Thus

$$\Phi_0 = (z_0 - u_{0,1})^{k_1} \dots (z_0 - u_{0,n_0-2})^{k_{n_0-2}} (z_0 - 1)^{k_{n_0-1}} (z_0 - \frac{t}{u_{1,1}})^{k_{n_0+1}} \dots (z_0 - \frac{t}{u_{1,n_1-2}})^{k_{n-2}} (z_0 - t)^{k_{n-1}} z_0^{k_n} (dz_0)^d,$$

and

$$\Phi_1 = (z_1 - \frac{t}{u_{0,1}})^{k_1} \dots (z_1 - \frac{t}{u_{0,n_0-2}})^{k_{n_0-2}} (z_1 - t)^{k_{n_0-1}} z_1^{k_{n_0}} (z_1 - u_{1,1})^{k_{n_0+1}} \dots (z_1 - u_{1,n_1-2})^{k_{n-2}} (z_1 - 1)^{k_{n-1}} (dz_1)^d.$$

Recall that we have

$$d\mu_S = d + k_1 + \dots + k_{n_0} = -(d + k_{n_0+1} + \dots + k_n) \geq 0.$$

Lemma 2.6. *The differentials $t^{d\mu_S}\Phi_0$ and Φ_1 represent two trivializing sections of the Kawamata line bundle \mathcal{K}_μ on $\overline{C}_{\mathcal{U}}$. Moreover, there is a non-vanishing holomorphic function f on $U_0 \times U_1$ such that*

$$f = \frac{\Phi_1}{t^{d\mu_S}\Phi_0}.$$

Proof. Remark that since \mathbf{x} is a generic point of D_S , it is not contained in any other boundary divisor $D_{S'}$ with $S' \neq S$. This means that for Δ, U_0, U_1 small enough, $p^{-1}(\mathcal{U})$ is disjoint from all the divisors $F_{S'}^0$, where $S' \neq S$.

By definition, we have $F_S^0 = \mathbb{P}_{\mathbb{C}}^1 \times \{0\} \times \{0\} \times U_0 \times U_1$. Note that F_S^0 is defined by the equation $\{z_1 = 0\}$ in the open affine $\{(z_0, z_1, t, u_0, u_1) \in \mathbb{C}^2 \times \mathcal{U}, z_0 z_1 = t\}$ of $\overline{C}_{|\mathcal{U}}$.

We first show that Φ_1 is a trivializing section of \mathcal{K}_μ . In $\overline{C}_{|\mathcal{U}}$, the divisors Γ_i are given by the following equations

$$\Gamma_i = \begin{cases} \{z_0 = u_{0,i}\} & \text{for } i = 1, \dots, n_0 - 1, \\ \{z_0 = \infty, z_1 = 0\} & \text{for } i = n_0, \\ \{z_1 = u_{1,i-n_0}\} & \text{for } i = n_0 + 1, \dots, n - 1, \\ \{z_0 = 0, z_1 = \infty\} & \text{for } i = n. \end{cases}$$

Note that we have $\Gamma_{n_0} = \{(\infty, 0)\} \times \mathcal{U}$ and $\Gamma_n = \{(0, \infty)\} \times \mathcal{U}$. Outside the union $\Gamma_{n_0} \cup \Gamma_n$, since $t = z_0 z_1$, we can write

$$\Phi_1 = \phi_1(z_0, z_1, u_0, u_1) z_1^{d\mu_S} \left(\frac{dz_1}{z_1} \right)^d.$$

where

$$\phi_1(z_0, z_1, u_0, u_1) = \frac{\prod_{i=1}^{n_0-1} (u_{0,i} - z_0)^{k_i} \cdot \prod_{i=1}^{n_1-1} (z_1 - u_{1,i})^{k_{n_0+i}}}{\prod_{i=1}^{n_0-1} u_{0,i}^{k_i}}.$$

Observe that $\frac{dz_1}{z_1}$ is a trivializing section of the relative canonical line bundle K of the projection $\overline{C}_{|\mathcal{U}} \rightarrow \mathcal{U}$, $z_1^{d\mu_S}$ is a trivializing section of $-d\mu_S F_S^0$, and the function ϕ_1 vanishes to the order k_i at the divisor Γ_i , $i \notin \{n_0, n\}$. Thus Φ_1 is a trivializing section of \mathcal{K}_μ on $\overline{C}_{|\mathcal{U}} \setminus (\Gamma_{n_0} \cup \Gamma_n)$.

Near a point in Γ_{n_0} , we use the coordinate $z'_0 = 1/z_0$. The set $\overline{C}_{|\mathcal{U}}$ is then defined by the equation $z_1 = tz'_0$. Hence a trivializing section of K is given by $dz'_0 = \frac{dz_1}{t}$. Note that

$$\phi_1 = (z'_0)^{d+k_{n_0}-d\mu_S} \frac{\prod_{i=1}^{n_0-1} (z'_0 u_{0,i} - 1)^{k_i} \cdot \prod_{i=1}^{n_1-1} (z_1 - u_{1,i})^{k_{n_0+i}}}{\prod_{i=1}^{n_0-1} u_{0,i}^{k_i}}.$$

Thus

$$\Phi_1 = \frac{\prod_{i=1}^{n_0-1} (z'_0 u_{0,i} - 1)^{k_i} \cdot (z'_0)^{k_{n_0}} \cdot \prod_{i=1}^{n_1-1} (z_1 - u_{1,i})^{k_{n_0+i}}}{\prod_{i=1}^{n_0-1} u_{0,i}^{k_i}} t^{d\mu_S} \left(\frac{dz_1}{t} \right)^d.$$

In this local chart F_S^0 is defined by $t = 0$, and Γ_{n_0} is defined by $z'_0 = 0$. Therefore, Φ_1 is a trivializing section of \mathcal{K}_μ .

Finally, near a point in Γ_n , we use the coordinate $z'_1 = 1/z_1$. Then $\overline{C}_{|\mathcal{U}}$ is defined by the equation $z_0 = tz'_1$. Hence a trivializing section of K is given by $dz'_1 = -\frac{dz_1}{z_1^2}$. We can write

$$\phi_1 = \frac{\prod_{i=1}^{n_0-1} (u_{0,i} - z_0)^{k_i} \cdot \prod_{i=1}^{n_1-1} (1 - u_{1,i} z'_1)^{k_{n_0+i}}}{\prod_{i=1}^{n_0-1} u_{0,i}^{k_i}} (z'_1)^{d\mu_S + d + k_n}.$$

Thus

$$\Phi_1 = \frac{\prod_{i=1}^{n_0-1} (u_{0,i} - z_0)^{k_i} \cdot \prod_{i=1}^{n_1-1} (1 - u_{1,i} z'_1)^{k_{n_0+i}} \cdot (z'_1)^{k_n}}{\prod_{i=1}^{n_0-1} u_{0,i}^{k_i}} \left(\frac{dz_1}{z_1^2} \right)^d.$$

Since Γ_n is defined by the equation $z'_1 = 0$, and Γ_n does not intersect the divisor F_S^0 , we can conclude that Φ_1 is a trivializing section of \mathcal{K}_μ .

That $t^{d\mu_S} \Phi_0$ is also a trivializing section of \mathcal{K}_μ follows from similar arguments. Using the relations $t = z_0 z_1$ and $\frac{dz_0}{z_0} = -\frac{dz_1}{z_1}$ (as sections of K) one can easily check that

$$\frac{\Phi_1}{\Phi_0} = t^{d\mu_S} \frac{(-1)^{d+k_{n_0}+k_n} u_{0,1}^{k_1} \cdots u_{0,n_0-2}^{k_{n_0-2}}}{u_{1,1}^{k_{n_0+1}} \cdots u_{1,n_1-2}^{k_{n-2}}}.$$

The lemma is then proved. \square

2.3.2. *Ratios of trivializing sections of \mathcal{K}_μ .* We are now back to the general case where \mathbf{x} is a point in a stratum of codimension r . Our goal is to show

Proposition 2.7. *For all $j = 0, \dots, r$, the d -differential Φ_j induces a meromorphic section of the Kawamata line bundle \mathcal{K}_μ on $\overline{\mathcal{C}}_{\mathcal{U}}$. Let t^{β_j} , $j = 0, \dots, r$, be the monomials in the variables t defined in (4). Then for all $j, j' \in \{0, 1, \dots, r\}$, there is a holomorphic non-vanishing function $f_{jj'}$ on $\Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \times \cdots \times U_r$ such that*

$$\frac{t^{\beta_j} \Phi_j}{t^{\beta_{j'}} \Phi_{j'}} = f_{jj'}.$$

Proof. We first notice that the 1-forms dz_j , $j = 0, \dots, r$, induce $(r+1)$ meromorphic sections of the relative canonical line bundle $K_{\overline{\mathcal{C}}/\mathcal{U}}$. It follows in particular that $t^{\beta_j} \Phi_j$, $j = 0, \dots, r$, are meromorphic sections of $K_{\overline{\mathcal{C}}/\mathcal{U}}$, and hence of \mathcal{K}_μ . We claim that if the components $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^k$ are connected by a node, that is $\alpha = \{j, k\} \in \mathcal{N}(\mathbf{x})$, then there is a holomorphic non-vanishing function f_{jk} on $\Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \times \cdots \times U_r$ such that

$$\frac{\Phi_j}{\Phi_k} = \frac{t^{\beta_k}}{t^{\beta_j}} f_{jk}.$$

Associated to $\alpha \in \mathcal{N}(\mathbf{x})$, we have a partition $\mathcal{S}_\alpha = (I_{0,\alpha}, I_{1,\alpha}) \in \mathcal{P}$. Let $\hat{C}_{\mathbf{x},\alpha}^0$ and $\hat{C}_{\mathbf{x},\alpha}^1$ be the subcurves of $C_{\mathbf{x}}$ that are separated by α . The numbering is chosen such that $x_i \in \hat{C}_{\mathbf{x},\alpha}^0$ if and only if $i \in I_{0,\alpha}$. Without loss of generality, we can assume that $C_{\mathbf{x}}^j \subset \hat{C}_{\mathbf{x},\alpha}^1$ and $C_{\mathbf{x}}^k \subset \hat{C}_{\mathbf{x},\alpha}^0$. In what follows, to lighten the notations, the subscript α will be omitted.

The relations

$$z_j = b_{jk} + \frac{t}{z_k - b_{kj}} \quad \text{and} \quad a_{ji} = b_{jk} + \frac{t}{a_{ki} - b_{kj}}$$

imply

$$(15) \quad z_j - a_{ji} = -t \frac{z_k - a_{ki}}{(z_k - b_{kj})(a_{ki} - b_{kj})}.$$

As a consequence of (13), we have

$$(16) \quad dz_j = \frac{-t}{(z_k - b_{kj})^2} dz_k,$$

as sections of $K_{\overline{C}/\mathcal{U}}$. Hence

$$\begin{aligned} \Phi_j &= \frac{(-t)^{\sum_{i=1}^n k_i} \prod_{i=1}^n (z_k - a_{ki})^{k_i}}{(z_k - b_{kj})^{\sum_{i=1}^n k_i} \prod_{i=1}^n (a_{ki} - b_{kj})^{k_i}} \times \frac{(-t)^d}{(z_k - b_{kj})^{2d}} (dz_k)^d \\ &= (-t)^{-d} \frac{\prod_{i=1}^n (z_k - a_{ki})^{k_i}}{\prod_{i=1}^n (a_{ki} - b_{kj})^{k_i}} (dz_k)^d \\ &= \frac{(-1)^d t^{-d}}{\prod_{i=1}^n (a_{ki} - b_{kj})^{k_i}} \Phi_k. \end{aligned}$$

Here we used the equality $\sum_{i=1}^n k_i = -2d$. Recall that a_{ki} is the $(k+1)$ -th coordinate of the marked point $\sigma_i(t, u) \in C(t, u)$. We have two possibilities

- $i \in I_0$, that is $x_i \in \hat{C}_{\mathbf{x}}^0$: if $x_i \in C_{\mathbf{x}}^k$, then since the marked point x_i and the point corresponding to the node α on $C_{\mathbf{x}}^k$ are distinct, $(a_{ki} - b_{kj})$ is a holomorphic non-vanishing function on U_k . Otherwise, x_i is contained in another component $C_{\mathbf{x}}^{k'}$ of $\hat{C}_{\mathbf{x}}^0$. From Lemma 2.4(iii), we know that the projection π_k maps $\sigma_i(t, u)$ to a point close to a node $b_{kj'}$ which must be different from b_{kj} . Since these two nodes are distinct, it follows that $(a_{ki} - b_{kj})$ is also a holomorphic non-vanishing function on $\Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \times \cdots \times U_r$.
- $i \in I_1$, that is $x_i \in \hat{C}_{\mathbf{x}}^1$: let us write $a_{ki} - b_{kj} = \frac{t}{a_{ji} - b_{jk}}$. By the same argument as above, we conclude that $(a_{ji} - b_{jk})$ is a holomorphic non-vanishing function on $\Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \times \cdots \times U_r$.

We then have

$$\begin{aligned} \frac{\Phi_j}{\Phi_k} &= \frac{(-1)^d t^{-d}}{\prod_{i=1}^n (a_{ki} - b_{kj})^{k_i}} \\ &= (-1)^d t^{-d - \sum_{i \in I_1} k_i} \frac{\prod_{i \in I_1} (a_{ji} - b_{jk})^{k_i}}{\prod_{i \in I_0} (a_{ki} - b_{kj})^{k_i}} \\ &= (-1)^d t^{d\mu_S} \frac{\prod_{i \in I_1} (a_{ji} - b_{jk})^{k_i}}{\prod_{i \in I_0} (a_{ki} - b_{kj})^{k_i}}. \end{aligned}$$

Recall that

$$\mu_S = \frac{1}{2} \left(\sum_{i \in I_1} \mu_i - \sum_{i \in I_0} \mu_i \right) = \sum_{i \in I_1} \mu_i - 1 = \frac{-1}{d} \left(\sum_{i \in I_1} k_i + d \right).$$

It remains to show that $t^{d\mu_S} = t^{\beta_k} / t^{\beta_j}$. Indeed, all the nodes $\alpha' \in \mathcal{N}(\mathbf{x})$ different from α do not separate the components $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^k$. Thus we have $\beta_{j,\alpha'} = \beta_{k,\alpha'}$. By definition, $\beta_{k,\alpha} = d\mu_S$, while $\beta_{j,\alpha} = 0$, and the claim follows. As a consequence, we get $\frac{t^{\beta_j} \Phi_j}{t^{\beta_k} \Phi_k} = f_{jk}$, where

$$f_{jk} := (-1)^d \frac{\prod_{i \in I_1} (a_{ji} - b_{jk})^{k_i}}{\prod_{i \in I_0} (a_{ki} - b_{kj})^{k_i}}$$

is a holomorphic non-vanishing function on $\Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \cdots \times U_r$. Since the curve $C_{\mathbf{x}}$ is connected, the lemma is proved. \square

2.3.3. Proof Theorem 2.5.

Proof. Let $(z^0, t^0, u^0) \in \overline{C_{\mathbf{x}}}$. Assume first that z^0 is a smooth point of $C(t^0, u^0)$, which means that z^0 is contained in a unique irreducible component $C^k(t^0, u^0)$ of $C(t^0, u^0)$. Since the curve $C(t^0, u^0)$ can be obtained from $C_{\mathbf{x}} \simeq C(0, 0)$ by smoothening some nodes, there is a surjective map $\varphi : C(t^0, u^0) \rightarrow C(0, 0) \simeq C_{\mathbf{x}}$, determined up to isotopy, that contracts some simple closed curves on $C(t^0, u^0)$ onto nodes in $C_{\mathbf{x}}$.

By definition, $J_k(t^0)$ is the set of $j \in \{0, \dots, r\}$ such that $C_{\mathbf{x}}^j$ is contained in $\varphi(C^k(t^0, u^0))$. Pick a j in $J_k(t^0)$. By Lemma 2.4, the projection π_j restricts to an isomorphism from $C^k(t^0, u^0)$ onto $\mathbb{P}_{\mathbb{C}}^1$. Thus, Φ_j is a trivializing section of the line bundle $d(K_{\overline{C}/\mathcal{U}} + \sum_{i=1}^n \mu_i \Gamma_i)$ in a neighborhood of (z^0, t^0, u^0) . It remains to check that t^{β_j} is a trivializing section of $\sum_{S \in \mathcal{P}} -d\mu_S F_S^0$ near (z^0, t^0, u^0) . Clearly, we only need to consider family of the divisors $\{F_{S_\alpha}^0, \alpha \in \mathcal{N}(\mathbf{x})\}$ because \mathcal{U} does not intersect D_S if $D_S \notin \{D_{S_\alpha}, \alpha \in \mathcal{N}(\mathbf{x})\}$.

Consider a node $\alpha \in Z(t^0)$, that is $t_\alpha^0 = 0$. Let $p_\alpha(t^0, u^0) \in C(t^0, u^0)$ be the corresponding node in $C(t^0, u^0)$. Concretely, if $\alpha = \{j, j'\}$ then $p_\alpha(t^0, u^0)$ is the point $z \in C(t^0, u^0)$ such that $z_j = b_{jj'}$ and $z_{j'} = b_{j'j}$. Denote by $\hat{C}(t^0, u^0)_\alpha^0$ and $\hat{C}(t^0, u^0)_\alpha^1$ the subcurves of $C(t^0, u^0)$ that are separated by $p_\alpha(t^0, u^0)$. Observe that $C^k(t^0, u^0) \subset \hat{C}(t^0, u^0)_\alpha^0$ if and only if $C_{\mathbf{x}}^j \subset \hat{C}_{\mathbf{x}, \alpha}^0$. This means that $C^k(t^0, u^0) \times \{(t^0, u^0)\}$ is contained in the divisor $F_{S_\alpha}^0$ if and only if $\beta_{j, \alpha} = d\mu_{S_\alpha}$. Since for all $\alpha \in \mathcal{N}(\mathbf{x}) \setminus Z(t^0)$, $t_\alpha \neq 0$ in a neighborhood of (z^0, t^0, u^0) , we conclude that the function

$$t^{\beta_j} = \prod_{\alpha \in \mathcal{N}(\mathbf{x})} t_\alpha^{\beta_{j, \alpha}}$$

gives a trivializing section of $-\sum_\alpha d\mu_{S_\alpha} F_{S_\alpha}^0$ in a neighborhood of (z^0, t^0, u^0) . As a consequence, $t^{\beta_j} \Phi_j$ is a trivializing section of the Kawamata line bundle in a neighborhood of (z^0, t^0, u^0) for all $j \in J_k(t)$.

Assume now that z^0 is a node of $C(t^0, u^0)$ which is contained in two components $C^k(t^0, u^0)$ and $C^{k'}(t^0, u^0)$. Let $\alpha = \{j, j'\} \in Z(t^0)$ be the corresponding node in $C_{\mathbf{x}}$. This means that $t_\alpha^0 = 0$, and we have

$$(z_j^0, z_{j'}^0) = (b_{jj'}(u^0), b_{j'j}(u^0)).$$

We can assume that $j \in J_k(t^0)$, $j' \in J_{k'}(t^0)$, and that $\beta_{j, \alpha} = 0$ and $\beta_{j', \alpha} = d\mu_{S_\alpha}$. This implies that the curve $C^{k'}(t^0, u^0) \times \{(t^0, u^0)\}$ is contained in the divisor $F_{S_\alpha}^0$. Note that near the point (z^0, t^0, u^0) , the divisor $F_{S_\alpha}^0$ is defined by the equation $z_j - b_{jj'} = 0$.

For all $i \in I_{1, \alpha}$, we have $a_{ji}(u^0) \neq b_{jj'}(u^0)$. Thus $z_j - a_{ji}$ is an invertible function near (z^0, t^0, u^0) . However for all $i \in I_{0, \alpha}$, we have $a_{ji}(u^0) = b_{jj'}(u^0) = z_j^0$ (see Lemma 2.4 (iii)), therefore $z_j - a_{ji}$ is not invertible near (z^0, t^0, u^0) . Using (15), we can write

$$z_j - a_{ji} = -t_\alpha \frac{z_{j'} - a_{j'i}}{(z_{j'} - b_{j'j})(a_{j'i} - b_{j'j})} = -(z_j - b_{jj'}) \cdot \frac{z_{j'} - a_{j'i}}{a_{j'i} - b_{j'j}}$$

where the functions $(z_{j'} - a_{j'i})$ and $(a_{j'i} - b_{j'j})$ are invertible near (z^0, t^0, u^0) . It follows that

$$(17) \quad \Phi_j = (-1)^{\sum_{i \in I_{0,\alpha}} k_i} (z_j - b_{jj'})^{d + \sum_{i \in I_{0,\alpha}} k_i} \times \prod_{i \in I_{1,\alpha}} (z_j - a_{ji})^{k_i} \times \prod_{i \in I_{0,\alpha}} \left(\frac{z_{j'} - a_{j'i}}{a_{j'i} - b_{j'j}} \right)^{k_i} \times \left(\frac{dz_j}{z_j - b_{jj'}} \right)^d.$$

In a neighborhood of (z^0, t^0, u^0) , a trivializing section of $K_{\overline{C}/\mathcal{U}}$ is given by $\frac{dz_j}{z_j - b_{jj'}} = -\frac{dz_{j'}}{z_{j'} - b_{j'j}}$. Recall that $(z_j - b_{jj'})^{d + \sum_{i \in I_{0,\alpha}} k_i} = (z_j - b_{jj'})^{d\mu_{S_\alpha}}$ is a trivializing section of $-d\mu_{S_\alpha} F_{S_\alpha}^0$ in a neighborhood of (z^0, t^0, u^0) . Equality (17) shows that Φ_j is a trivializing section of $d(K_{\overline{C}/\mathcal{U}} - \mu_{S_\alpha} F_{S_\alpha}^0)$ in this neighborhood.

Since $\beta_{j,\alpha} = 0$, by the same arguments as the previous case, we see that

$$t^{\beta_j} = \prod_{\alpha' \in \mathcal{N}(\mathbf{x}) \setminus \{\alpha\}} t_{\alpha'}^{\beta_{j,\alpha'}}$$

is a trivializing section of $\sum_{\alpha' \in \mathcal{N}(\mathbf{x}) \setminus \{\alpha\}} -d\mu_{S_{\alpha'}} F_{S_{\alpha'}}^0$. Since (z^0, t^0, u^0) does not belong to any divisor Γ_i , $i = 1, \dots, n$, it follows that $t^{\beta_j} \Phi_j$ is a trivializing section of \mathcal{K}_μ in a neighborhood of (z^0, t^0, u^0) .

We have then showed that for every $(z^0, t^0, u^0) \in \overline{C}\mathcal{U}$ there is some $j \in \{0, \dots, r\}$ such that $t^{\beta_j} \Phi_j$ is a trivializing section of \mathcal{K}_μ in a neighborhood of (z^0, t^0, u^0) . Since all the sections $t^{\beta_j} \Phi_j$, $j = 0, \dots, r$, differ from a fixed one, say $t^{\beta_0} \Phi_0$, by a non-vanishing function depending only on (t, u) by Proposition 2.7, we conclude that all of them are trivializing sections of \mathcal{K}_μ on $\overline{C}\mathcal{U}$. \square

2.4. Proof of Theorem 2.2.

Proof. By Theorem 2.5, every point $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$ has a neighborhood \mathcal{U} such that the line bundle \mathcal{K}_μ is trivial over $p^{-1}(\mathcal{U})$. Every holomorphic section Φ of \mathcal{K}_μ on $p^{-1}(\mathcal{U})$ is then given by a holomorphic function. Since all the fibers of p are compact and connected, Φ must be constant on the fibers. Thus $\Phi = p^*f$, where f is a holomorphic function on \mathcal{U} . This implies that $p_*\mathcal{K}_\mu|_{\mathcal{U}} \sim \mathcal{O}_{\mathcal{U}}$. Thus $p_*\mathcal{K}_\mu$ is a line bundle $\tilde{\mathcal{L}}_\mu$ on $\overline{\mathcal{M}}_{0,n}$ and we have $\mathcal{K}_\mu \sim p^*\tilde{\mathcal{L}}_\mu$. It is clear from the definition that the restriction of $\tilde{\mathcal{L}}_\mu$ to $\mathcal{M}_{0,n}$ is \mathcal{L}_μ .

The expression of the divisor on $\overline{\mathcal{M}}_{0,n}$ corresponding to $\tilde{\mathcal{L}}_\mu$ can be derived from the Grothendieck-Riemann-Roch formula. Details of the calculation can be found in [34, §8.4]. \square

2.5. Example. To close this section, we give some explicit formulas for the trivializing sections of the Kawamata line bundle near a point \mathbf{x} in a stratum of codimension 2 of $\overline{\mathcal{M}}_{0,n}$. In this case, $C_{\mathbf{x}}$ is a stable curve with 3 irreducible components denoted by $C_{\mathbf{x}}^0, C_{\mathbf{x}}^1, C_{\mathbf{x}}^2$. We assume that $C_{\mathbf{x}}^0$ intersects both $C_{\mathbf{x}}^1$ and $C_{\mathbf{x}}^2$. Let α_1 (resp. α_2) denote nodes between $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^1$ (resp. between $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^2$). Let $\{I_0, I_1, I_2\}$ be the partition of $\{1, \dots, n\}$ where I_j records the marked points in the component $C_{\mathbf{x}}^j$ of $C_{\mathbf{x}}$, for $j = 0, 1, 2$. Set $n_j = |I_j|$, $j = 0, 1, 2$. Without loss of generality, we can assume that $I_0 = \{1, \dots, n_0\}$, $I_1 = \{n_0 + 1, \dots, n_0 + n_1\}$, and $I_2 = \{n_0 + n_1 + 1, \dots, n\}$. Since $C_{\mathbf{x}}$ is stable, we must have $n_0 \geq 1$, $n_1 \geq 2$, and $n_2 \geq 2$.

Recall that each component of $C_{\mathbf{x}}$ is isomorphic to $\mathbb{P}_{\mathbb{C}}^1$. Let x, y, z be the inhomogeneous coordinates on $C_{\mathbf{x}}^0, C_{\mathbf{x}}^1$, and $C_{\mathbf{x}}^2$ respectively. For $i \in \{1, \dots, n\}$, let a_i be the (inhomogeneous) coordinate of the i -th marked point of $C_{\mathbf{x}}$ in its component. Using the automorphisms of $\mathbb{P}_{\mathbb{C}}^1$, we can assume that

- In $C_{\mathbf{x}}^0$, 0 corresponds to the node α_1 , 1 corresponds to the node α_2 , and $a_{n_0} = \infty$.

- In $C_{\mathbf{x}}^1$, 0 corresponds to the node α_1 , $a_{n_0+n_1-1} = 1$, $a_{n_0+n_1} = \infty$.
- In $C_{\mathbf{x}}^2$, 0 corresponds to the node α_2 , $a_{n-1} = 1$, $a_n = \infty$.

Let U_0 be an open neighborhood of $(a_i)_{1 \leq i \leq n_0-1}$ in \mathbb{C}^{n_0-1} , U_1 an open neighborhood of $(a_{n_0+i})_{1 \leq i \leq n_1-2}$ in \mathbb{C}^{n_1-2} , and U_2 an open neighborhood of $(a_{n_0+n_1+i})_{1 \leq i \leq n_2-2}$ in \mathbb{C}^{n_2-2} .

Let $D \subset \mathbb{C}$ be a small disk about 0. A neighborhood of \mathbf{x} in $\overline{\mathcal{M}}_{0,n}$ can be identified with $\mathcal{U} = D^2 \times U_0 \times U_1 \times U_2$. Let (t_1, t_2) be the coordinates of D^2 , where t_i corresponds to the node α_i . For all $(t_1, t_2) \in D^2$, define

$$C_{(t_1, t_2)} := \{(x, y, z) \in (\mathbb{P}_{\mathbb{C}}^1)^3, xy = t_1, (x-1)z = t_2\} \subset (\mathbb{P}_{\mathbb{C}}^1)^3.$$

The universal curve over \mathcal{U} is given by

$$\overline{C}_{\mathcal{U}} := \{(x, y, z, t_1, t_2, X, Y, Z) \in (\mathbb{P}_{\mathbb{C}}^1)^3 \times D^2 \times U_0 \times U_1 \times U_2, xy = t_1, (x-1)z = t_2\}$$

where $X = (x_i)_{1 \leq i \leq n_0-1} \in U_0$, $Y = (y_i)_{n_0+1 \leq i \leq n_0+n_1-2} \in U_1$, and $Z = (z_i)_{n_0+n_1 \leq i \leq n-2} \in U_2$. We set

$$x_{n_0} = \infty, y_{n_0+n_1-1} = 1, y_{n_0+n_1} = \infty, z_{n-1} = 1, z_n = \infty,$$

and define

$$\begin{aligned} \sigma_i(t_1, t_2, X, Y, Z) &:= \left(x_i, \frac{t_1}{x_i}, \frac{t_2}{x_i-1}\right) \in C_{(t_1, t_2)}, \quad i = 1, \dots, n_0, \\ \sigma_i(t_1, t_2, X, Y, Z) &:= \left(\frac{t_1}{y_i}, y_i, \frac{t_2 y_i}{t_1 - y_i}\right) \in C_{(t_1, t_2)}, \quad i = n_0 + 1, \dots, n_1, \\ \sigma_i(t_1, t_2, X, Y, Z) &:= \left(\frac{t_2 + z_i}{z_i}, \frac{t_1 z_i}{t_2 + z_i}, z_i\right) \in C_{(t_1, t_2)}, \quad i = n_0 + n_1 + 1, \dots, n. \end{aligned}$$

The pointed curve represented by (t_1, t_2, X, Y, Z) is isomorphic to $(C_{(t_1, t_2)}, (\sigma_i(t_1, t_2, X, Y, Z))_{1 \leq i \leq n})$. Define

$$\begin{aligned} \Phi_0 &:= \prod_{i \in I_0} (x - x_i)^{k_i} \prod_{i \in I_1} \left(x - \frac{t_1}{y_i}\right)^{k_i} \prod_{i \in I_2} \left(x - \frac{t_2 + z_i}{z_i}\right)^{k_i} (dx)^d, \\ \Phi_1 &:= \prod_{i \in I_0} \left(y - \frac{t_1}{x_i}\right)^{k_i} \prod_{i \in I_1} (y - y_i)^{k_i} \prod_{i \in I_2} \left(y - \frac{t_1 z_i}{t_2 + z_i}\right)^{k_i} (dy)^d, \\ \Phi_2 &:= \prod_{i \in I_0} \left(z - \frac{t_2}{x_i - 1}\right)^{k_i} \prod_{i \in I_1} \left(z - \frac{t_2 y_i}{t_1 - y_i}\right)^{k_i} \prod_{i \in I_2} (z - z_i)^{k_i} (dz)^d. \end{aligned}$$

Set $\hat{k}_j := \sum_{i \in I_j} k_i$, $j = 0, 1, 2$, and $\nu_1 := -d - \hat{k}_1 = d + \hat{k}_0 + \hat{k}_2$, $\nu_2 := -d - \hat{k}_2 = d + \hat{k}_0 + \hat{k}_1$. Using the relations $y = \frac{t_1}{x}$, and $z = \frac{t_2}{x-1}$, one can readily check that

$$\begin{aligned} \frac{\Phi_1}{\Phi_0} &= (-1)^d \frac{\prod_{i \in I_1} y_i^{k_i}}{\prod_{i \in I_0} x_i^{k_i}} \cdot \prod_{i \in I_2} \left(\frac{z_i}{t_2 + z_i}\right)^{k_i} t_1^{\nu_1}, \\ \frac{\Phi_2}{\Phi_0} &= (-1)^d \prod_{i \in I_1} \left(\frac{y_i}{t_1 - y_i}\right)^{k_i} \frac{\prod_{i \in I_2} z_i^{k_i}}{\prod_{i \in I_0} (x_i - 1)^{k_i}} t_2^{\nu_2}. \end{aligned}$$

We have the following possibilities:

- (a) $\nu_1 \leq 0$ and $\nu_2 \leq 0$: in this case $\beta_0 = (0, 0), \beta_1 = (-\nu_1, 0), \beta_2 = (0, -\nu_2)$, and $\Phi_0 \sim t_1^{-\nu_1} \Phi_1 \sim t_2^{-\nu_2} \Phi_2$ are trivializing sections of the Kawamata line bundle. The fiber $\tilde{\mathcal{L}}_{\mu, \mathbf{x}}$ is generated by the d -differential which is equal to $\prod_{i \in I_0} (x - a_i)^{k_i} \cdot x^{\hat{k}_1} \cdot (x - 1)^{\hat{k}_2} \cdot (dx)^d$ on $C_{(0,0)}^0$ and vanishes identically on the components $C_{(0,0)}^1, C_{(0,0)}^2$.
- (b) $\nu_1 \geq 0$ and $\nu_2 \leq 0$: in this case $\beta_0 = (\nu_1, 0), \beta_1 = (0, 0), \beta_2 = (\nu_1, -\nu_2)$, and $t_1^{\nu_1} \Phi_0 \sim \Phi_1 \sim t_1^{\nu_1} t_2^{-\nu_2} \Phi_2$ are trivializing sections of \mathcal{K}_μ . The fiber $\tilde{\mathcal{L}}_{\mu, \mathbf{x}}$ is generated by the d -differential which is equal to $\prod_{i \in I_1} (y - a_i)^{k_i} \cdot y^{\hat{k}_0 + \hat{k}_2} \cdot (dy)^d$ on $C_{(0,0)}^1$ and vanishes identically on $C_{(0,0)}^0$ and $C_{(0,0)}^2$.
- (c) $\nu_1 \leq 0$ and $\nu_2 \geq 0$: in this case $\beta_0 = (0, \nu_2), \beta_1 = (-\nu_1, \nu_2), \beta_2 = (0, 0)$, and $t_2^{\nu_2} \Phi_0 \sim t_1^{-\nu_1} t_2^{\nu_2} \Phi_1 \sim \Phi_2$ are trivializing sections of \mathcal{K}_μ . The fiber $\tilde{\mathcal{L}}_{\mu, \mathbf{x}}$ is generated by the d -differential which is equal to $\prod_{i \in I_2} (z - a_i)^{k_i} \cdot z^{\hat{k}_0 + \hat{k}_1} \cdot (dz)^d$ on $C_{(0,0)}^2$ and vanishes identically on $C_{(0,0)}^0$ and $C_{(0,0)}^1$.
- (d) $\nu_1 \geq 0$ and $\nu_2 \geq 0$: in this case, $\beta_0 = (\nu_1, \nu_2), \beta_1 = (0, \nu_2), \beta_2 = (\nu_1, 0)$, and $t_1^{\nu_1} t_2^{\nu_2} \Phi_0 \sim t_2^{\nu_2} \Phi_1 \sim t_1^{\nu_1} \Phi_2$ are all trivializing sections of \mathcal{K}_μ . The d -differentials on $C_{(0,0)}$ associated with $\tilde{\mathcal{L}}_{\mu, \mathbf{x}}$ vanish identically on all three components. In particular, we do not have a geometric interpretation of elements of the fiber $\tilde{\mathcal{L}}_{\mu, \mathbf{x}}$ in this case.

3. BLOW-UP OF $\overline{\mathcal{M}}_{0,n}$

3.1. Construction. Let \mathbf{x} be a point in a stratum of codimension r in $\overline{\mathcal{M}}_{0,n}$. By definition, \mathbf{x} is contained in the intersection $\cap_{\alpha \in \mathcal{N}(\mathbf{x})} D_{S_\alpha}$. Recall that $\mathcal{U}_\mathbf{x}$ is a neighborhood of \mathbf{x} that satisfies the following

- (i) $\mathcal{U}_\mathbf{x}$ does not intersect any boundary divisor $D_S \notin \{D_{S_\alpha}, \alpha \in \mathcal{N}(\mathbf{x})\}$,
- (ii) $\mathcal{U}_\mathbf{x}$ can be identified with an open subset of \mathbb{C}^{n-3} such that, for each $\alpha \in \mathcal{N}(\mathbf{x})$, there is a coordinate function t_α such that $D_{S_\alpha} \cap \mathcal{U}_\mathbf{x}$ is defined by the equation $t_\alpha = 0$.

Denote the coordinates on $\mathcal{U}_\mathbf{x}$ by (t, u) , where $t = (t_\alpha)_{\alpha \in \mathcal{N}(\mathbf{x})}$, and $u = (u_1, \dots, u_{n-3-r})$. Recall that for $j \in \{0, \dots, r\}$, $\beta_j = (\beta_{j,\alpha})_{\alpha \in \mathcal{N}(\mathbf{x})}$, where

$$\beta_{j,\alpha} = \begin{cases} d\mu_{S_\alpha} = d + \sum_{i \in I_{0,\alpha}} k_i & \text{if } C_\mathbf{x}^j \subset \hat{C}_{\mathbf{x},\alpha}^0, \\ 0 & \text{otherwise,} \end{cases}$$

and $t^{\beta_j} = \prod_{\alpha \in \mathcal{N}(\mathbf{x})} t_\alpha^{\beta_{j,\alpha}}$. To simplify the notations, throughout this section, we will write D_α instead of D_{S_α} , and μ_α instead of μ_{S_α} .

Let $\mathcal{O}_{\overline{\mathcal{M}}_{0,n}}$ be the structure sheaf of $\overline{\mathcal{M}}_{0,n}$, and $\mathcal{I}_{\mathcal{U}_\mathbf{x}}$ be the sheaf of ideals of $\mathcal{O}_{\overline{\mathcal{M}}_{0,n}|_{\mathcal{U}_\mathbf{x}}}$ generated by $\{t^{\beta_0}, \dots, t^{\beta_r}\}$. We first show

Theorem 3.1. *The sheaves of ideals $\{\mathcal{I}_{\mathcal{U}_\mathbf{x}}, \mathbf{x} \in \overline{\mathcal{M}}_{0,n}\}$ patch together and give rise to an ideal sheaf \mathcal{I} of $\mathcal{O}_{\overline{\mathcal{M}}_{0,n}}$.*

Proof. We need to show that $\mathcal{I}_{\mathcal{U}_\mathbf{x}}|_{\mathcal{U}_\mathbf{x} \cap \mathcal{U}_{\mathbf{x}'}} \simeq \mathcal{I}_{\mathcal{U}_{\mathbf{x}'}}|_{\mathcal{U}_\mathbf{x} \cap \mathcal{U}_{\mathbf{x}'}}$ for all $\mathbf{x}, \mathbf{x}' \in \overline{\mathcal{M}}_{0,n}$. We first consider the case $\mathbf{x}' \in \mathcal{U}_\mathbf{x}$. In this case, all the boundary divisors that contain \mathbf{x}' contain also \mathbf{x} . Thus $\mathcal{N}(\mathbf{x}')$ is a subset of $\mathcal{N}(\mathbf{x})$. Let $s = |\mathcal{N}(\mathbf{x}')|$. The curve $C_{\mathbf{x}'}$ has $(s + 1)$ irreducible components, which will be denoted by $C_{\mathbf{x}'}^0, \dots, C_{\mathbf{x}'}^s$. For all $\alpha \in \mathcal{N}(\mathbf{x}')$, let t'_α be a function in the affine coordinate ring of $\mathcal{U}_{\mathbf{x}'}$ that defines $D_\alpha \cap \mathcal{U}_{\mathbf{x}'}$. Let $t' = (t'_\alpha)_{\alpha \in \mathcal{N}(\mathbf{x}'})$. For $j' = 0, \dots, s$, define the exponent vectors $\beta_{j'} = (\beta'_{j',\alpha})_{\alpha \in \mathcal{N}(\mathbf{x}'})$, and

the monomial $t'^{\beta'_j}$ in the same way as β_j and t^{β_j} . By a slight abuse of notation, we consider β'_j as a vector in $\mathbb{Z}_{\geq 0}^{\mathcal{N}(\mathbf{x})}$ by setting $\beta'_{j,\alpha} = 0$ for all $\alpha \in \mathcal{N}(\mathbf{x}) \setminus \mathcal{N}(\mathbf{x}')$.

By definition $\mathcal{I}_{\mathcal{U}_{\mathbf{x}'}}$ is generated by $(t'^{\beta'_0}, \dots, t'^{\beta'_s})$. Since the restrictions of t'_α and t_α to $\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}$ define the same divisor, t'_α/t_α is an invertible regular function on $\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}$. It follows that $\mathcal{I}_{\mathcal{U}_{\mathbf{x}'|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}}} = (t^{\beta'_0}, \dots, t^{\beta'_s})$.

Since the curve $C_{\mathbf{x}'}$ can be obtained from $C_{\mathbf{x}}$ by smoothening the nodes in $\mathcal{N}(\mathbf{x}) \setminus \mathcal{N}(\mathbf{x}')$, there is a continuous map $\varphi : C_{\mathbf{x}'} \rightarrow C_{\mathbf{x}}$ that sends each component of $C_{\mathbf{x}'}$ onto a subcurve of $C_{\mathbf{x}}$. Consider an irreducible component $C_{\mathbf{x}}^i$ of $C_{\mathbf{x}}$. There is an irreducible component $C_{\mathbf{x}'}^j$ such that $C_{\mathbf{x}}^i \subset \varphi(C_{\mathbf{x}'}^j)$. By construction, for all $\alpha \in \mathcal{N}(\mathbf{x}')$, we have $\beta_{i,\alpha} = \beta'_{j,\alpha}$. For all $\alpha \in \mathcal{N}(\mathbf{x}) \setminus \mathcal{N}(\mathbf{x}')$, t_α is an invertible regular function on $\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}$ (because \mathbf{x}' is not contained in the divisor D_α). Hence $t^{\beta_i}/t^{\beta'_j}$ is an invertible regular function on $\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}$. It follows that

$$\mathcal{I}_{\mathcal{U}_{\mathbf{x}'|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}}} \simeq (t^{\beta'_0}, \dots, t^{\beta'_s}) \simeq (t^{\beta_0}, \dots, t^{\beta_r}) \simeq \mathcal{I}_{\mathcal{U}_{\mathbf{x}}|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}}.$$

Consider now the general case. For every $\mathbf{x}'' \in \mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}$, we consider the open affine $\mathcal{U}_{\mathbf{x}''}$ of \mathbf{x}'' which satisfies (i) and (ii). The above arguments show that we have

$$\mathcal{I}_{\mathcal{U}_{\mathbf{x}}|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'} \cap \mathcal{U}_{\mathbf{x}''}} \simeq \mathcal{I}_{\mathcal{U}_{\mathbf{x}''}|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'} \cap \mathcal{U}_{\mathbf{x}''}} \simeq \mathcal{I}_{\mathcal{U}_{\mathbf{x}''}|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'} \cap \mathcal{U}_{\mathbf{x}''}}.$$

It follows that $\mathcal{I}_{\mathcal{U}_{\mathbf{x}}|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}} \simeq \mathcal{I}_{\mathcal{U}_{\mathbf{x}'}|\mathcal{U}_{\mathbf{x}} \cap \mathcal{U}_{\mathbf{x}'}}$, and the theorem follows. \square

Definition 3.2. Let $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ be the blow-up of $\overline{\mathcal{M}}_{0,n}$ with respect to the sheaf of ideals \mathcal{I} (see [28, p.163]). By definition $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ comes equipped with a projection $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$. As an analytic space, $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ can be described as follows: let $\mathcal{U}_{\mathbf{x}}^*$ denote the set $\mathcal{U}_{\mathbf{x}} \cap \mathcal{M}_{0,n}$, that is $\mathcal{U}_{\mathbf{x}}^* = \{(t, u) \in \mathcal{U}_{\mathbf{x}}, t_\alpha \neq 0, \forall \alpha \in \mathcal{N}(\mathbf{x})\}$. Assuming that $r > 0$, let

$$\widehat{\mathcal{U}}_{\mathbf{x}}^* := \{(t, u, [t^{\beta_0} : \dots : t^{\beta_r}]), (t, u) \in \mathcal{U}_{\mathbf{x}}^* \subset \mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^r,$$

and

$$\mathcal{U}_{\mathbf{x}}^\# = \{(t, u, [v_0 : \dots : v_r]) \in \mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^r, v_j t^{\beta_k} = v_k t^{\beta_j}, \forall j, k \in \{0, \dots, r\}, j \neq k\} \subset \mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^r.$$

Then $\widehat{\mathcal{U}}_{\mathbf{x}} := \hat{p}^{-1}(\mathcal{U}_{\mathbf{x}})$ is isomorphic to the irreducible component of $\mathcal{U}_{\mathbf{x}}^\#$ that contains $\widehat{\mathcal{U}}_{\mathbf{x}}^*$. Equivalently, one can define $\widehat{\mathcal{U}}_{\mathbf{x}}$ to be the closure of $\widehat{\mathcal{U}}_{\mathbf{x}}^*$ in $\mathcal{U} \times \mathbb{P}_{\mathbb{C}}^r$.

It is worth noticing that $\widehat{\mathcal{U}}_{\mathbf{x}}$ depends only on $\mathcal{I}_{\mathcal{U}_{\mathbf{x}}}$, which means that if in the definition of $\widehat{\mathcal{U}}_{\mathbf{x}}$ we take two different generating sets of $\mathcal{I}_{\mathcal{U}_{\mathbf{x}}}$, then the resulting varieties are isomorphic. We have the following classical result (see for instance [28, Ch.II.§7, Prop. 7.16]).

Proposition 3.3. *The space $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is a complex projective variety, and the natural projection $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$ is a birational, proper, surjective morphism.*

We will spend the remainder of this section studying the geometry of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. In particular, we will be interested in the fibers of the projection $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$.

3.2. Support of the ideal sheaf \mathcal{I} . Consider a point $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$ as in §3.1. Recall that $\mathcal{N}(\mathbf{x})$ is the set of pairs $\{j, j'\}$, where $j, j' \in \{0, \dots, r\}$ and $j \neq j'$, that represent the nodes of $C_{\mathbf{x}}$. The node corresponding to the pair $\alpha = \{j, j'\}$ is obtained by identifying the point $y_{jj'} \in C_{\mathbf{x}}^j$ with the point $y_{j'j} \in C_{\mathbf{x}}^{j'}$. We assign to $y_{jj'}$ and $y_{j'j}$ some weights as follows: to the node $\alpha = \{j, j'\} \in \mathcal{N}(\mathbf{x})$ we have an associated partition $\mathcal{S}_{\alpha} = \{I_{0,\alpha}, I_{1,\alpha}\}$ of $\{1, \dots, n\}$. Let $\hat{C}_{\mathbf{x},\alpha}^0$ and $\hat{C}_{\mathbf{x},\alpha}^1$ be the subcurves of $C_{\mathbf{x}}$ that are separated by α . Without loss of generality, we can assume that $C_{\mathbf{x}}^j \subset \hat{C}_{\mathbf{x},\alpha}^0$, and $C_{\mathbf{x}}^{j'} \subset \hat{C}_{\mathbf{x},\alpha}^1$. We then define

$$(18) \quad \mu(y_{jj'}) = \sum_{i \in I_{1,\alpha}} \mu_i \quad \text{and} \quad \mu(y_{j'j}) = \sum_{i \in I_{0,\alpha}} \mu_i.$$

By definition, we have

$$(19) \quad \mu(y_{jj'}) + \mu(y_{j'j}) = 2, \quad \text{for all } \{j, j'\} \in \mathcal{N}(\mathbf{x}),$$

and

$$(20) \quad \sum_{i \in I_j} \mu_i + \sum_{j' \in \mathcal{N}_j(\mathbf{x})} \mu(y_{jj'}) = 2, \quad \text{for each } j \in \{0, \dots, r\},$$

where $\mathcal{N}_j(\mathbf{x}) = \{j' \in \{0, \dots, r\}, \{j, j'\} \in \mathcal{N}(\mathbf{x})\}$ is the set of nodes of the component $C_{\mathbf{x}}^j$.

Recall that the dual graph $T_{\mathbf{x}}$ of $C_{\mathbf{x}}$ is a tree whose vertices are denoted by $\{v_0, \dots, v_r\}$, with v_j representing the component $C_{\mathbf{x}}^j$ of $C_{\mathbf{x}}$. Each node of $C_{\mathbf{x}}$ corresponds to a geometric edge of $T_{\mathbf{x}}$. Such an edge consists of a pair of oriented edges with opposite orientations. The oriented edge from v_j to $v_{j'}$ will be denoted by $e_{jj'}$. We associate to $e_{jj'}$ the weight $\mu(e_{jj'})$ defined by:

$$\mu(e_{jj'}) := \mu(y_{j'j}) - \mu(y_{jj'}).$$

Note that if $\alpha = \{j, j'\} \in \mathcal{N}(\mathbf{x})$ then

$$(21) \quad \mu_{\alpha} = \frac{1}{2} \left(\sum_{i \in I_{1,\alpha}} \mu_i - \sum_{i \in I_{0,\alpha}} \mu_i \right) = \frac{1}{2} |\mu(e_{jj'})|.$$

We will call μ_{α} the weight of α .

Definition 3.4. Assume that all the nodes in $C_{\mathbf{x}}$ have strictly positive weights. In this case, the component $C_{\mathbf{x}}^j$ of $C_{\mathbf{x}}$ is said to be principal, if we have $\mu(e_{jj'}) > 0$ (or equivalently $\mu(y_{jj'}) < 1$), for every $j' \in \mathcal{N}_j(\mathbf{x})$.

In the case some nodes in $C_{\mathbf{x}}$ have zero weight, we first smoothen those nodes to obtain another curve $C_{\mathbf{x}'}$ parametrized by a point \mathbf{x}' in a neighborhood of \mathbf{x} . Each irreducible component of $C_{\mathbf{x}'}$ corresponds to the union of some components in $C_{\mathbf{x}}$ that are connected by nodes of zero weight. The union corresponding to a principal component of $C_{\mathbf{x}'}$ is called a principal subcurve of $C_{\mathbf{x}}$. A component of $C_{\mathbf{x}}$ is said to be principal if it is contained in a principal subcurve.

Lemma 3.5. If $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^{j'}$ are contained in the same principal subcurve, then $\beta_j = \beta_{j'}$.

Proof. By definition, if the node α does not separate $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^{j'}$, then $\beta_{j,\alpha} = \beta_{j',\alpha}$. Since by definition all the edges that separate $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^{j'}$ have weight zero, the lemma follows. \square

Lemma 3.6. There always exists at least one principal component.

Proof. It suffices to consider the case $C_{\mathbf{x}}$ does not have zero weighted nodes. If there exists no principal component, then for every vertex v_j of $T_{\mathbf{x}}$ there exists an oriented edge $e_{jj'}$ such that $\mu(e_{jj'}) < 0$. Consider a path γ in $T_{\mathbf{x}}$ that is composed uniquely by oriented edges with negative weight. Let $e_{jj'}$ be the last edge of this path. Since there exists j'' such that $\mu(e_{j'j''}) < 0$ (note that we must have $j'' \neq j$), we can always extend γ to a longer path. Thus, there exists such a path γ of length $r + 1$. Since $T_{\mathbf{x}}$ has exactly $r + 1$ vertices, this means that γ must contain a cycle. But $T_{\mathbf{x}}$ is a tree, so this is impossible. \square

Lemma 3.7. *If the component $C_{\mathbf{x}}^j$ of $C_{\mathbf{x}}$ is not principal, then there is a principal component $C_{\mathbf{x}}^k$ such that the unique path in $T_{\mathbf{x}}$ from v_j to v_k consists of edges of weight ≤ 0 , and at least one of those edges has negative weight.*

Proof. Again, it is enough to prove the lemma in the case $C_{\mathbf{x}}$ does not have zero weighted nodes. Assume that $C_{\mathbf{x}}^j$ is not principal. Set $j_0 = j$. By definition, $C_{\mathbf{x}}^{j_0}$ has an adjacent component $C_{\mathbf{x}}^{j_1}$ such that the edge $e_{j_0j_1}$ is negatively weighted. If $C_{\mathbf{x}}^{j_1}$ is not principal then there exists $C_{\mathbf{x}}^{j_2}$ such that $\mu(e_{j_1j_2}) < 0$. Note that $\mu(e_{j_1j_0}) > 0$, therefore $e_{j_1j_2} \neq e_{j_1j_0}$. Continue this procedure we get a sequence of negatively weighted edges $(e_{j_0j_1}, e_{j_1j_2}, \dots, e_{j_{i-1}j_i})$ of $T_{\mathbf{x}}$. The concatenation of those edges gives a path γ in $T_{\mathbf{x}}$. If $C_{\mathbf{x}}^{j_i}$ is not principal, then the path γ can be extended to a longer path. Since $T_{\mathbf{x}}$ is a tree, the length of γ cannot be larger than r , which means that the procedure must terminate after finitely many steps. Let s be the length of γ when the procedure terminates. Then $C_{\mathbf{x}}^{j_s}$ is a principal component, and $\gamma = e_{j_{s-1}j_s} * \dots * e_{j_0j_1}$ is the desired path. \square

As a consequence of Lemma 3.7, we have

Lemma 3.8. *If $C_{\mathbf{x}}^j$ is not a principal component of $C_{\mathbf{x}}$, then there exist $k \in \{1, \dots, r_0\}$ and a non-empty subset $\mathfrak{s}(j, k)$ of $\mathcal{N}(\mathbf{x})$ such that*

$$t^{\beta_j} = \left(\prod_{\alpha \in \mathfrak{s}(j, k)} t_{\alpha}^{d\mu_{\alpha}} \right) \cdot t^{\beta_k}.$$

(here we write μ_{α} instead of $\mu_{S_{\alpha}}$).

Proof. By Lemma 3.7, there is a principal component $C_{\mathbf{x}}^k$ such that the path from v_j to v_k in $T_{\mathbf{x}}$ consists only of negatively weighted edges. This means that for any node α that separates $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^k$, we always have $C_{\mathbf{x}}^j \subset \hat{C}_{\mathbf{x}, \alpha}^0$. Thus, $t^{\beta_j} / t^{\beta_k} = \prod_{\alpha \in \mathfrak{s}(j, k)} t_{\alpha}^{d\mu_{\alpha}}$, where $\mathfrak{s}(j, k)$ is the set of nodes that separate $C_{\mathbf{x}}^j$ and $C_{\mathbf{x}}^k$. \square

Let $\hat{C}_{\mathbf{x}}^1, \dots, \hat{C}_{\mathbf{x}}^{r_0}$ denote the principal subcurves of $C_{\mathbf{x}}$. Let β_m^* be the exponent vector associated with any irreducible component of $\hat{C}_{\mathbf{x}}^m$ (by Lemma 3.5 all the components of $\hat{C}_{\mathbf{x}}^m$ have the same associated exponent vector). An immediate consequence of Lemma 3.8 is

Corollary 3.9. *The ideal sheaf $\mathcal{I}|_{\mathcal{U}_{\mathbf{x}}}$ is generated by the functions $\{t^{\beta_1^*}, \dots, t^{\beta_{r_0}^*}\}$.*

We now show

Proposition 3.10. *A point $\mathbf{x} \in \overline{\mathcal{M}}_{0, n}$ is **not** contained in the support of the sheaf of ideals \mathcal{I} if and only if $C_{\mathbf{x}}$ has a unique principal subcurve, or equivalently, the union of all the principal irreducible components of $C_{\mathbf{x}}$ is connected.*

Proof. Recall that given \mathbf{x} in $\overline{\mathcal{M}}_{0,n}^{(r)}$, the stratum of codimension r in $\overline{\mathcal{M}}_{0,n}$, the sheaf \mathcal{I} is generated by $(r+1)$ functions $\{t^{\beta_0}, \dots, t^{\beta_r}\}$ in a neighborhood of \mathbf{x} . Assume first that $C_{\mathbf{x}}$ has only one principal subcurve. We claim that if $C_{\mathbf{x}}^k$ is a principal irreducible component, then $t^{\beta_k} = 1$, which implies that $\mathcal{I} \simeq \mathcal{O}_{\overline{\mathcal{M}}_{0,n}}$ in a neighborhood of \mathbf{x} . Suppose that $t^{\beta_k} \neq 1$. Then there is a node $\alpha_0 = \{j_0, j'_0\}$ such that $C_{\mathbf{x}}^k \subset \hat{C}_{\mathbf{x},\alpha_0}^0$. Without loss of generality, we can assume that α_0 separates $C_{\mathbf{x}}^{j_0}$ and $C_{\mathbf{x}}^k$. This means that the unique path in $T_{\mathbf{x}}$ from v_{j_0} to v_k contains the edge $e_{j_0 j'_0}$, and $\mu(e_{j_0 j'_0}) > 0$. It follows in particular that $C_{\mathbf{x}}^{j_0}$ is not principal, because otherwise we would have two principal components that are not connected by a sequence of nodes of weight zero contradicting the assumption that $C_{\mathbf{x}}$ has only one principal subcurve.

Since $C_{\mathbf{x}}^{j_0}$ is not principal, by Lemma 3.7, there is a principal component $C_{\mathbf{x}}^j$ such that the path from v_{j_0} to v_j consists only of negatively weighted edges. Since $\mu(e_{j_0 j'_0}) > 0$, the edge $e_{j_0 j'_0}$ is not contained in this path. Since $T_{\mathbf{x}}$ is a tree, the unique path in $T_{\mathbf{x}}$ from v_k to v_j is the concatenation of the path from v_k to v_{j_0} and the path from v_{j_0} to v_j . In particular, the path from v_k to v_j contains $e_{j_0 j'_0}$, and again we have a contradiction to the assumption that $C_{\mathbf{x}}$ has only one principal component. Therefore, we can conclude that $t^{\beta_k} = 1$.

Assume now that $C_{\mathbf{x}}$ has at least two principal subcurves. We claim that $t^{\beta_j} \neq 1$ for all $j \in \{0, \dots, r\}$, which implies that $\mathbf{x} \in \text{supp}(\mathcal{I})$. By Lemma 3.8, we know that if $C_{\mathbf{x}}^j$ is not a principal component, then there exists a principal component $C_{\mathbf{x}}^k$ such that t^{β_j}/t^{β_k} is a non constant monomial in the variables $(t_{\alpha})_{\alpha \in \mathcal{N}(\mathbf{x})}$. Thus $t^{\beta_j} \neq 1$ in this case. If $C_{\mathbf{x}}^j$ is a principal component, since there are more than one principal subcurves, there is a principal component $C_{\mathbf{x}}^k$ and a node $\alpha = \{k, k'\}$ attached to $C_{\mathbf{x}}^k$ which separates $C_{\mathbf{x}}^k$ and $C_{\mathbf{x}}^j$. Since $C_{\mathbf{x}}^k$ is principal, $\mu(y_{kk'}) < 1$, where $y_{kk'}$ is the point in $C_{\mathbf{x}}^k$ corresponding to the node α . This implies that $C_{\mathbf{x}}^j \subset \hat{C}_{\mathbf{x},\alpha}^0$. Consequently, $\beta_{j,\alpha} = d\mu_{\alpha} > 0$, and $t^{\beta_j} \neq 1$, since it contains the factor $t_{\alpha}^{d\mu_{\alpha}}$. The proof of the proposition is now complete. \square

Proposition 3.10 implies

Corollary 3.11. *If for all $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$, the curve $C_{\mathbf{x}}$ has only one principal subcurve, then the blow-up $\widehat{\mathcal{M}}_{0,n}(k)$ is isomorphic to $\overline{\mathcal{M}}_{0,n}$.*

Remark 3.12. *It is shown in [34, §5] that if all the weights $\{\mu_i, i = 1, \dots, n\}$ are positive then $C_{\mathbf{x}}$ has a unique principal subcurve for all $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$. However, this positivity condition is not necessary. For instance, for $\mu = (\frac{1}{2}, \frac{-1}{2})$ or equivalently $\underline{k} = (-1^5, 1)$, we also have $\widehat{\mathcal{M}}_{0,6}(-1^5, 1) \simeq \overline{\mathcal{M}}_{0,6}$.*

3.3. Fiber of the blow-up. Our goal now is to show

Theorem 3.13. *Let r_0 be the number of principal subcurves of $C_{\mathbf{x}}$. Then \mathbf{x} is contained in the support of \mathcal{I} if and only if $r_0 \geq 2$, and we have*

$$\hat{p}^{-1}(\{\mathbf{x}\}) \simeq \mathbb{P}_{\mathbb{C}}^{r_0-1}.$$

For the proof of Theorem 3.13, we will need some preparation. We first observe that if α is zero weighted node in $\mathcal{N}(\mathbf{x})$, then t_{α} is not involved in any of the monomials t^{β_j} . Thus it is enough to consider the case where $\mu_{\alpha} > 0$ for all $\alpha \in \mathcal{N}(\mathbf{x})$. This implies in particular that each principal subcurve has only one irreducible component. Without loss of generality, we can assume that the set of principal components of $C_{\mathbf{x}}$ consists of $C_{\mathbf{x}}^0, C_{\mathbf{x}}^1, \dots, C_{\mathbf{x}}^{r_0-1}$.

By definition $\widehat{\mathcal{U}}_{\mathbf{x}} := \widehat{\rho}^{-1}(\mathcal{U}_{\mathbf{x}})$ is the irreducible component of the algebraic set

$$\mathcal{U}_{\mathbf{x}}^{\#} := \{(t, u, [v_0 : \cdots : v_{r_0-1}]) \in \mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}, v_i t^{\beta_j} = v_j t^{\beta_i}, \forall i, j = 0, \dots, r_0 - 1\}$$

that contains $\widehat{\mathcal{U}}_{\mathbf{x}}^* = \{(t, u, [t^{\beta_0} : \cdots : t^{\beta_{r_0-1}}]), (t, u) \in \mathcal{U}_{\mathbf{x}}, t_{\alpha} \neq 0 \forall \alpha \in \mathcal{N}(\mathbf{x})\}$. It follows in particular that $\widehat{\rho}^{-1}(\{\mathbf{x}\}) \subset \{\mathbf{x}\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}$.

Let us partition the set $\mathcal{N}(\mathbf{x})$ of nodes of $C_{\mathbf{x}}$ into two subsets as follows: \mathfrak{b} is the set of nodes that connect a principal component and a non-principal one, and \mathfrak{c} is the set of nodes that connect two non-principal components. By assumption, each edge in \mathfrak{b} joins a vertex in $\{v_0, \dots, v_{r_0-1}\}$ and a vertex in $\{v_{r_0}, \dots, v_r\}$.

Let us denote the set $\{v_0, \dots, v_{r_0-1}\}$ by Y . Remove Y and all the edges in \mathfrak{b} from $T_{\mathbf{x}}$, we get a disjoint union Z of finitely many subtrees of $T_{\mathbf{x}}$, which will be denoted by Z_1, \dots, Z_{s_0} . By definition, the edges of Z correspond to elements of \mathfrak{c} .

Contracting the edges corresponding to elements of \mathfrak{c} in $T_{\mathbf{x}}$, we get a tree which has $r_0 + s_0$ vertices. A vertex of this tree corresponds to either a vertex in Y , or a connected component of Z . Since the edges of this tree correspond to the elements of \mathfrak{b} , we have

$$(22) \quad |\mathfrak{b}| = r_0 + s_0 - 1.$$

Denote the set of edges that connect Z_j to Y by \mathfrak{b}_j , $j = 1, \dots, s_0$. Let $m_j = |\mathfrak{b}_j|$. Denote by $\alpha_{j,0}$ the edge in \mathfrak{b}_j that is contained in the path from v_0 to Z_j . The remaining edges of \mathfrak{b}_j are labeled by $\alpha_{j,i}$, $i = 1, \dots, m_j - 1$. To simplify the notation, in what follows we will write $v_{j,i} = d\mu_{\alpha_{j,i}}$ for all $\alpha \in \mathfrak{b}$. Define

$$x_{j,i} := \frac{t_{\alpha_{j,0}}^{v_{j,0}}}{t_{\alpha_{j,i}}^{v_{j,i}}}, \quad i = 1, \dots, m_j - 1.$$

Note that the family $\{x_{j,i}, j = 1, \dots, s_0, i = 1, \dots, m_j - 1\}$ contains exactly $|\mathfrak{b}| - s_0 = r_0 - 1$ functions.

For each $m \in \{1, \dots, r_0 - 1\}$, let γ_m denote the unique path in $T_{\mathbf{x}}$ from v_0 to v_m . Let $(Z_{j_1}, \dots, Z_{j_{s(m)}})$ be the sequence of components of Z that meet γ_m . Note that for $k \in \{1, \dots, s(m)\}$, γ_m contains exactly two edges in \mathfrak{b}_{j_k} , namely $\alpha_{j_k,0}$ and another edge α_{j_k,i_k} . Hence, the sequence of edges in \mathfrak{b} that are contained in γ_m is

$$\mathfrak{b}(0, m) := (\alpha_{j_1,0}, \alpha_{j_1,i_1}, \dots, \alpha_{j_{s(m)},0}, \alpha_{j_{s(m)},i_{s(m)}}).$$

Lemma 3.14. *We have*

$$\frac{t^{\beta_m}}{t^{\beta_0}} = h_m \prod_{k=1}^{s(m)} x_{j_k,i_k}$$

where h_m depends only on $(t_{\alpha})_{\alpha \in \mathfrak{c}}$.

Proof. The function t^{β_m}/t^{β_0} involves t_{α} if and only if α separates v_0 and v_m . If $\alpha \in \mathfrak{b}$ then α is contained in the sequence $\mathfrak{b}(0, m)$. Observe that the edge of $T_{\mathbf{x}}$ corresponding to $\alpha_{j_k,0}$ (with the orientation induced by γ_m) is positively weighted, while the edge corresponding to α_{j_k,i_k} is negatively weighted, for all $k = 1, \dots, s(m)$. Therefore we have,

$$\beta_{0,\alpha_{j_k,0}} = 0, \quad \beta_{m,\alpha_{j_k,0}} = v_{j_k,0}, \quad \text{and} \quad \beta_{0,\alpha_{j_k,i_k}} = v_{j_k,i_k}, \quad \beta_{m,\alpha_{j_k,i_k}} = 0.$$

As a consequence,

$$\frac{t^{\beta_m}}{t^{\beta_0}} = h_m \cdot \frac{\prod_{k=1}^{s(m)} t^{\nu_{j_k,0}} \alpha_{j_k,0}^{\nu_{j_k,0}}}{\prod_{k=1}^{s(m)} t^{\nu_{j_k,i_k}} \alpha_{j_k,i_k}^{\nu_{j_k,i_k}}} = h_m \prod_{k=1}^{s(m)} x_{j_k i_k}$$

where h_m depends only on $(t_\alpha)_{\alpha \in \mathfrak{c}}$. \square

Lemma 3.15. Fix $(t_\alpha)_{\alpha \in \mathfrak{c}} \in (\mathbb{C}^*)^{\mathfrak{c}}$, and let $h_m = h_m((t_\alpha)_{\alpha \in \mathfrak{c}})$ be as in Lemma 3.14. Then the map

$$\begin{aligned} \Xi : (\mathbb{C}^*)^{\mathfrak{b}} &\rightarrow \mathbb{C}^{r_0-1} \\ (t_\alpha)_{\alpha \in \mathfrak{b}} &\mapsto \left(h_m \cdot \prod_{k=1}^{s(m)} x_{j_k i_k} \right)_{1 \leq m \leq r_0-1} \end{aligned}$$

is an open map.

Proof. Recall that $\alpha_{j_{s(m)}, i_{s(m)}}$ is the last edge in the path from v_0 to v_m for $m = 1, \dots, r_0 - 1$. Since every edge $\alpha_{j,i}$, with $i \in \{1, \dots, m_j - 1\}$, is the last edge of the path from v_0 to some v_k , the correspondence $v_m \mapsto \alpha_{j_{s(m)}, i_{s(m)}}$ is a bijection between $\{v_1, \dots, v_m\}$ and $\{\alpha_{j,i}, j = 1, \dots, s_0, i = 1, \dots, m_j - 1\}$. In particular, we have $\#\{\alpha_{j,i}, j = 1, \dots, s_0, i = 1, \dots, m_j - 1\} = |\mathfrak{b}| - s_0 = r_0 - 1$. Define

$$\begin{aligned} \Xi_1 : (\mathbb{C}^*)^{\mathfrak{b}} &\rightarrow \mathbb{C}^{r_0-1} \\ (t_\alpha)_{\alpha \in \mathfrak{b}} &\mapsto (x_{ji})_{1 \leq j \leq s_0, 1 \leq i \leq m_j - 1} \end{aligned}$$

Since $t_{\alpha_{j,i}}$ is only involved in x_{ji} , it is clear that Ξ_1 is an open map.

For $m = 1, \dots, r_0 - 1$, define

$$\phi_m := \prod_{k=1}^{s(m)} x_{j_k i_k},$$

and

$$\begin{aligned} \Xi_2 : \mathbb{C}^{r_0-1} &\rightarrow \mathbb{C}^{r_0-1} \\ (x_{ji})_{1 \leq j \leq s_0, 1 \leq i \leq m_j - 1} &\mapsto (\phi_m)_{1 \leq m \leq r_0-1} \end{aligned}$$

Observe that by construction, if $s(m) > 1$, then there is $m' \in \{2, \dots, r_0\}$ such that $\phi_m = \phi_{m'} x_{j_{s(m)}, i_{s(m)}}$. This is because $(\alpha_{j_1,0}, \alpha_{j_1,i_1}, \dots, \alpha_{j_{s-1},0}, \alpha_{j_{s-1},i_{s-1}})$ is the sequence of the edges in \mathfrak{b} that are contained in the path from v_0 to $v_{m'}$. Using this property, by induction over $\max\{s(m), m = 1, \dots, r_0 - 1\}$, one readily shows that Ξ_2 is an open map.

Since $\Xi = D \circ \Xi_2 \circ \Xi_1$, where $D : \mathbb{C}^{r_0-1} \rightarrow \mathbb{C}^{r_0-1}$ is the linear map defined by the diagonal matrix $\text{diag}(h_1, \dots, h_{r_0-1})$, the lemma follows. \square

Recall that $\mathfrak{b} = \{\alpha_{j,i}, j = 1, \dots, s_0, i = 0, \dots, m_j - 1\}$. For each $j \in \{1, \dots, s_0\}$, we choose an integer c_j such that $\nu_{j,i}$ divides c_j for all $i \in \{0, \dots, m_j - 1\}$. Set

$$s_{j,i} := \frac{c_j}{\nu_{j,i}}.$$

Let $\Delta^* = \{\lambda \in \mathbb{C}, 0 < |\lambda| < 1\}$. For every $\lambda \in \Delta^*$, we define an action ρ_λ of λ on $\mathbb{C}^{\mathcal{N}(\mathbf{x})}$ as follows:

$$\begin{cases} \rho_\lambda(t)_\alpha = t_\alpha, & \text{for all } \alpha \in \mathfrak{c} \\ \rho_\lambda(t)_{\alpha_{j,i}} = \lambda^{s_{j,i}} t_{\alpha_{j,i}}, & \text{for all } \alpha_{j,i} \in \mathfrak{b}. \end{cases}$$

Lemma 3.16. Let $t' = \rho_\lambda(t)$. Then for all $m \in \{1, \dots, r_0 - 1\}$, we have

$$\frac{t'^{\beta_m}}{t'^{\beta_0}} = \frac{t^{\beta_m}}{t^{\beta_0}}.$$

Proof. By Lemma 3.14, we have

$$\frac{t^{\beta_m}}{t^{\beta_0}} = h_m(t) \prod_{k=1}^{s(m)} x_{j_k, i_k}(t)$$

where $h_m(t)$ depends only on $(t_\alpha)_{\alpha \in c}$. Since $(t'_\alpha)_{\alpha \in c} = (t_\alpha)_{\alpha \in c}$, it is enough to show that $x_{ji}(t') = x_{ji}(t)$, for all $j = 1, \dots, s_0$, $i = 1, \dots, m_j - 1$. By definition

$$x_{j,i}(t') = \frac{t'^{\nu_{j,0}}}{t'^{\nu_{j,i}}} = \frac{\lambda^{s_{j,0}\nu_{j,0}} t^{\nu_{j,0}}}{\lambda^{s_{j,i}\nu_{j,i}} t^{\nu_{j,i}}} = \frac{t^{\nu_{j,0}}}{t^{\nu_{j,i}}} = x_{j,i}(t).$$

since we have $s_{j,0}\nu_{j,0} = s_{j,i}\nu_{j,i} = c_j$. The lemma is then proved. \square

3.4. Proof of Theorem 3.13.

Proof. By Proposition 3.10, we already know that $\mathbf{x} \in \text{supp}(\mathcal{I})$ if and only if $C_{\mathbf{x}}$ contains at least two principal subcurves, that is $r_0 \geq 2$. It remains to show that $\hat{p}^{-1}(\{\mathbf{x}\}) \simeq \mathbb{P}_{\mathbb{C}}^{r_0-1}$.

As remarked above, we can assume that all the nodes of $C_{\mathbf{x}}$ have strictly positive weight. By Corollary 3.9, $\mathcal{I}_{\mathcal{U}_{\mathbf{x}}}$ is generated by $\{t^{\beta_0}, \dots, t^{\beta_{r_0-1}}\}$. Therefore, $\widehat{\mathcal{U}}_{\mathbf{x}}$ is a subvariety of $\mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}$, which means that $\hat{p}^{-1}(\{\mathbf{x}'\}) \subset \{\mathbf{x}'\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}$ for all $\mathbf{x}' \in \mathcal{U}_{\mathbf{x}}$.

Recall that $\mathcal{U}_{\mathbf{x}}$ can be identified with $\Delta^{\mathcal{N}(\mathbf{x})} \times U$, where $\Delta \subset \mathbb{C}$ is the unit disc, and U is an open neighborhood of 0 in \mathbb{C}^{n-3-r} . In this identification $\mathbf{x} = (0, 0) \in \Delta^{\mathcal{N}(\mathbf{x})} \times U$. Define

$$\mathcal{V}_{\mathbb{C}}^* = \{(t, u) \in \mathcal{U}_{\mathbf{x}}, t_\alpha = 0, \forall \alpha \in \mathfrak{b}, \text{ and } t_\alpha \neq 0, \forall \alpha \in \mathfrak{c}\}.$$

We will show that for all $\mathbf{y}_0 = (t^0, u^0) \in \mathcal{V}_{\mathbb{C}}^*$, we have

$$\{\mathbf{y}_0\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1} \subset \widehat{\mathcal{U}}_{\mathbf{x}} = \hat{p}^{-1}(\mathcal{U}_{\mathbf{x}}).$$

Since $\{\mathbf{y}_0\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}$ intersects $\widehat{\mathcal{U}}_{\mathbf{x}}$ in a subvariety of $\mathbb{P}_{\mathbb{C}}^{r_0-1}$, it is enough to show that $\widehat{\mathcal{U}}_{\mathbf{x}} \cap \{\mathbf{y}_0\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}$ contains an open subset of $\{\mathbf{y}_0\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1}$.

Let Ξ be the function in Lemma 3.15 with $h_m = h_m((t_\alpha)_{\alpha \in c})$. Identifying \mathbb{C}^{r_0-1} with an open subset of $\mathbb{P}_{\mathbb{C}}^{r_0-1}$, we can consider Ξ as a map from $(\mathbb{C}^*)^{\mathfrak{b}}$ to $\mathbb{P}_{\mathbb{C}}^{r_0-1}$. Consider $t \in \Delta^{\mathcal{N}(\mathbf{x})}$ such that

$$\begin{cases} t_\alpha = t_\alpha^0, & \forall \alpha \in \mathfrak{c}, \\ t_\alpha \neq 0, & \forall \alpha \in \mathfrak{b}. \end{cases}$$

Observe that $(t, u^0, \Xi((t_\alpha)_{\alpha \in \mathfrak{b}})) \in \widehat{\mathcal{U}}_{\mathbf{x}}$. By Lemma 3.16, we also have $(\rho_\lambda(t), u^0, \Xi((t_\alpha)_{\alpha \in \mathfrak{b}})) \in \widehat{\mathcal{U}}_{\mathbf{x}}$, for any $\lambda \in \Delta^*$. For all $\alpha \in \mathfrak{b}$, we have $\lim_{\lambda \rightarrow 0} \rho_\lambda(t)_\alpha = 0$. It follows that $\lim_{\lambda \rightarrow 0} \rho_\lambda(t) = t^0$. Consequently, $(t^0, u^0, \Xi((t_\alpha)_{\alpha \in \mathfrak{b}})) \in \widehat{\mathcal{U}}_{\mathbf{x}}$.

Let $W_{\mathbf{y}_0}$ denote the image of $(\Delta^*)^{\mathfrak{b}}$ by Ξ . By Lemma 3.15, $W_{\mathbf{y}_0}$ is an open subset of $\mathbb{C}^{r_0-1} \subset \mathbb{P}_{\mathbb{C}}^{r_0-1}$. Since $(t_\alpha)_{\alpha \in \mathfrak{b}}$ can be chosen arbitrarily in $(\Delta^*)^{\mathfrak{b}}$, we conclude that $\{\mathbf{y}_0\} \times W_{\mathbf{y}_0} \subset \widehat{\mathcal{U}}_{\mathbf{x}}$. Therefore, we have $\{\mathbf{y}_0\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1} \subset \widehat{\mathcal{U}}_{\mathbf{x}}$ for all $\mathbf{y}_0 \in \mathcal{V}_{\mathbb{C}}^*$. Since $\mathbf{x} = (0, 0)$ is a limit of points in $\mathcal{V}_{\mathbb{C}}^*$, we also have $\{\mathbf{x}\} \times \mathbb{P}_{\mathbb{C}}^{r_0-1} \subset \widehat{\mathcal{U}}_{\mathbf{x}}$, and the theorem follows. \square

Remark 3.17. In the case one uses the generator family $\{t^{\beta_0}, \dots, t^{\beta_r}\}$ of $\mathcal{I}_{\mathcal{U}_{\mathbf{x}}}$ instead of $\{t^{\beta_0}, \dots, t^{\beta_{r_0-1}}\}$ in the definition of the blow-up, we have that $\widehat{\mathcal{U}}_{\mathbf{x}}$ is the closure of the set $\{(t, u, [t^{\beta_0} : \dots : t^{\beta_r}], (t, u) \in \mathcal{U}_{\mathbf{x}}^*) \subset \mathcal{U}_{\mathbf{x}}^* \times \mathbb{P}_{\mathbb{C}}^r$ in $\mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^r$. In this case, the fiber $\hat{p}^{-1}(\{\mathbf{x}\})$ can be described as follows: for $i =$

$0, \dots, r_0 - 1$, let ℓ_i be the number of irreducible components of the principal subcurve \hat{C}_x^i . Denote by $\ell = \sum_{i=0}^{r_0-1} \ell_i$. We choose a numbering of the irreducible components of C_x such that for $i = 0, \dots, r_0 - 1$, \hat{C}_x^i is the union $C_x^{\ell_0+\dots+\ell_{i-1}} \cup \dots \cup C_x^{\ell_0+\dots+\ell_i-1}$, and C_x^ℓ, \dots, C_x^r are the non-principal components. In this setting, $\hat{p}^{-1}(\{\mathbf{x}\})$ is the image of $\mathbb{P}_\mathbb{C}^{r_0-1}$ in $\mathbb{P}_\mathbb{C}^r$ under the following embedding

$$\iota : \begin{array}{ccc} \mathbb{P}_\mathbb{C}^{r_0-1} & \rightarrow & \mathbb{P}_\mathbb{C}^r \\ [v_0 : \dots : v_{r_0-1}] & \mapsto & [\underbrace{v_0 : \dots : v_0}_{\ell_0} : \dots : \underbrace{v_{r_0-1} : \dots : v_{r_0-1}}_{\ell_{r_0-1}} : \underbrace{0 : \dots : 0}_{r+1-\ell}] \end{array}$$

We close this section by a combinatorial criterion for $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ to be isomorphic to $\mathcal{M}_{0,n}$.

Proposition 3.18. *The blow-up $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is isomorphic to $\overline{\mathcal{M}}_{0,n}$ if and only if there does not exist a partition $\{I_0, I_1, I_2\}$ of $\{1, \dots, n\}$ such that*

$$(23) \quad \sum_{i \in I_1} \mu_i > 1, \quad \text{and} \quad \sum_{i \in I_2} \mu_i > 1.$$

Proof. Assume first that such a partition exists. Consider the following partitions of $\{1, \dots, n\}$: $\mathcal{S}_1 = \{I_1, I_0 \sqcup I_2\}$ and $\mathcal{S}_2 = \{I_2, I_0 \sqcup I_1\}$. Since $\mu_i < 1$ and $\mu_1 + \dots + \mu_n = 2$, condition (23) implies that $|I_1| \geq 2, |I_2| \geq 2$, and $I_0 \neq \emptyset$. Therefore, both $\mathcal{S}_1, \mathcal{S}_2$ belong to \mathcal{P} . Denote by D_1 and D_2 the boundary divisors of $\overline{\mathcal{M}}_{0,n}$ associated with \mathcal{S}_1 and \mathcal{S}_2 respectively.

Consider a point $\mathbf{x} \in D_1 \cap D_2$. The stable curve (C_x, x_1, \dots, x_n) represented by \mathbf{x} has three components C_x^0, C_x^1, C_x^2 , where $x_i \in C_x^j$ if and only if $i \in I_j$. Note that C_x^0 intersects both C_x^1 and C_x^2 . It is straightforward to check that both C_x^1 and C_x^2 are principal components of C_x . Since they are disjoint, it follows from Theorem 3.13 that $\hat{p}^{-1}(\{\mathbf{x}\}) \simeq \mathbb{P}_\mathbb{C}^1$. Hence $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is not isomorphic to $\overline{\mathcal{M}}_{0,n}$.

Suppose now that $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is not isomorphic to $\overline{\mathcal{M}}_{0,n}$. Then there exists a point $\mathbf{x} \in \overline{\mathcal{M}}_{0,n}$ such that C_x has more than one principal subcurve. Smoothing the nodes of zero weight of C_x , we can assume that each principal subcurve contains a single irreducible component.

Let C_x^1 and C_x^2 be two principal components of C_x . Recall that C_x^1 and C_x^2 correspond to the vertices v_1, v_2 of T_x respectively. Consider the unique path γ in T_x from v_1 to v_2 . The first edge in γ corresponds to a node α_1 between C_x^1 and a non-principal component. Similarly, the last edge in γ corresponds to a node α_2 between C_x^2 and a non-principal component. For $j \in \{1, 2\}$, the node α_j is associated to a partition $\mathcal{S}_j = \{I_{j0}, I_{j1}\} \in \mathcal{P}$, where

$$\mu(I_{j1}) = \sum_{i \in I_{j1}} \mu_i > 1.$$

By assumption $C_x^1 \subset \hat{C}_{x,\alpha_1}^1$ and $C_x^2 \subset \hat{C}_{x,\alpha_2}^1$. Observe that we must have $\hat{C}_{x,\alpha_2}^1 \subset \hat{C}_{x,\alpha_1}^0$, which implies that $I_{21} \subset I_{10}$. Set $I_0 := I_{10} \setminus I_{21}$. Then $\{I_0, I_{11}, I_{21}\}$ is a partition of $\{1, \dots, n\}$ which satisfies (23). \square

4. BOUNDARY COMPONENTS OF THE BLOW-UP

Let $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k}) := \hat{p}^{-1}(\partial \overline{\mathcal{M}}_{0,n})$. We will call $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$ the boundary of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. In this section, we investigate the irreducible components of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$. We start by

Proposition 4.1. *Every irreducible component of $\partial \widehat{\mathcal{M}}_{0,n}(\underline{k})$ is a divisor.*

Proof. Since $\partial\overline{\mathcal{M}}_{0,n}$ is locally defined by a single equation, so is $\partial\widehat{\mathcal{M}}_{0,n}(k)$. \square

The first family of irreducible components of $\partial\widehat{\mathcal{M}}_{0,n}(k)$ arise from the irreducible components of $\partial\overline{\mathcal{M}}_{0,n}$. Consider a boundary divisor $D_S \subset \partial\overline{\mathcal{M}}_{0,n}$ for some $S = \{I_0, I_1\} \in \mathcal{P}$. By Proposition 3.10, we know that D_S is not contained in the support of \mathcal{I} . Let \hat{D}_S denote the *proper transform* of D_S in $\widehat{\mathcal{M}}_{0,n}(k)$. Then \hat{D}_S is an irreducible component of $\partial\widehat{\mathcal{M}}_{0,n}(k)$. Note that \hat{D}_S is isomorphic to the blow-up of D_S along the ideal sheaf $i_S^{-1}\mathcal{I}$, where $i_S : D_S \rightarrow \overline{\mathcal{M}}_{0,n}$ is the natural embedding (see[28, Chap.II, Cor.7.15]).

Our goal now is to look for irreducible components \hat{D} of $\partial\widehat{\mathcal{M}}_{0,n}(k)$ whose projection in $\overline{\mathcal{M}}_{0,n}$ is contained in the support of \mathcal{I} . Recall that a stratum of $\overline{\mathcal{M}}_{0,n}$ is the set of points that parametrize pointed stable curves which have the same dual graph. Each stratum of $\overline{\mathcal{M}}_{0,n}$ is a (connected) submanifold of $\overline{\mathcal{M}}_{0,n}$ whose codimension is equal to the number of edges in the dual graph.

Lemma 4.2. *Let S be a stratum of $\partial\overline{\mathcal{M}}_{0,n}$. If $S \cap \hat{p}(\hat{D}) \neq \emptyset$, then $\overline{S} \subset \hat{p}(\hat{D})$.*

Proof. Let r be the codimension of S . Consider a point $\mathbf{x} \in S \cap \hat{p}(\hat{D})$. Let $\mathcal{U}_{\mathbf{x}}$ be a neighborhood of \mathbf{x} as in §3.1, and $\widehat{\mathcal{U}}_{\mathbf{x}} = \hat{p}^{-1}(\mathcal{U}_{\mathbf{x}})$. Then $\hat{D} \cap \widehat{\mathcal{U}}_{\mathbf{x}}$ is an irreducible component of the algebraic set defined by the equations

$$v_i t^{\beta_j} = v_j t^{\beta_i}, \quad i, j = 0, \dots, r, \quad \text{and} \quad \prod_{\alpha \in \mathcal{N}(\mathbf{x})} t_{\alpha} = 0,$$

in $\mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^r$. By definition, S is defined by the equations $t_{\alpha} = 0$, for all $\alpha \in \mathcal{N}(\mathbf{x})$. Recall that $\mathcal{U}_{\mathbf{x}} \simeq \Delta^{\mathcal{N}(\mathbf{x})} \times U_0 \times \dots \times U_r$. Remark that the equations defining \hat{D} do not involve the parameters of U_0, \dots, U_r . Therefore, if \hat{D} contains a point $(0, u_0, \dots, u_r, [v]) \in \mathcal{U}_{\mathbf{x}} \times \mathbb{P}^r$, then $\{0\} \times U_0 \times \dots \times U_r \times \{[v]\} \subset \hat{D}$, which means that $\hat{p}(\hat{D})$ contains an open subset of S . This implies that $S \subset \hat{p}(\hat{D})$ (because $\hat{p}(\hat{D}) \cap S$ must be a Zariski closed subset of S). Since $\hat{p}(\hat{D})$ must be a closed subset of $\overline{\mathcal{M}}_{0,n}$, we conclude that $\overline{S} \subset \hat{p}(\hat{D})$. \square

Corollary 4.3. *There is a stratum S of $\overline{\mathcal{M}}_{0,n}$ such that \hat{D} is the closure of $\hat{p}^{-1}(S)$ in $\widehat{\mathcal{M}}_{0,n}(k)$.*

Proof. It follows from Lemma 4.2 that there is a unique maximal stratum S of $\overline{\mathcal{M}}_{0,n}$ such that $\overline{S} = \hat{p}(\hat{D})$. By Theorem 3.13, $\hat{p}^{-1}(S)$ is a projective bundle over S . Thus $\hat{p}^{-1}(S)$ is an irreducible subvariety of $\widehat{\mathcal{M}}_{0,n}(k)$. Note that we have $\hat{D} \subset \hat{p}^{-1}(S)$. Since $\hat{p}^{-1}(S) \subset \partial\widehat{\mathcal{M}}_{0,n}(k)$, and \hat{D} is supposed to be an irreducible component of $\partial\widehat{\mathcal{M}}_{0,n}(k)$, we must have $\hat{D} = \hat{p}^{-1}(S)$. \square

Proposition 4.4. *Let S be a stratum in $\overline{\mathcal{M}}_{0,n}$ of codimension $r \geq 2$. Let T_S be the dual graph, and $\{I_0, \dots, I_r\}$ the partition of $\{1, \dots, n\}$ associated to S . The vertices of T_S are denoted by v_0, \dots, v_r , where v_j is the vertex associated to I_j . Assume that $\hat{D} = \hat{p}^{-1}(S)$. Then up to renumbering of the subsets in $\{I_0, \dots, I_r\}$, we have*

$$(24) \quad \begin{cases} \sum_{i \in I_0} \mu_i < 1, \\ \sum_{i \in I_j} \mu_i > 1, \quad \text{for all } j = 1, \dots, r, \end{cases}$$

and for every $j \in \{1, \dots, r\}$ there is an edge in T_S joining v_0 to v_j .

Conversely, assume that $\{I_0, \dots, I_r\}$ satisfies (24), and every edge of T_S joins v_0 to a vertex in $\{v_1, \dots, v_r\}$. Then $\hat{p}^{-1}(S)$ is an irreducible component of $\partial\widehat{\mathcal{M}}_{0,n}(k)$.

Proof. Let \mathbf{x} be a point in S . We claim that the curve $C_{\mathbf{x}}$ parametrized by \mathbf{x} has exactly one non-principal component. By Theorem 3.13, $\hat{p}^{-1}(\{\mathbf{x}\}) \simeq \mathbb{P}_{\mathbb{C}}^{r_0-1}$, where r_0 is the number of principal subcurves of $C_{\mathbf{x}}$. Since \hat{D} has codimension one, we must have $r_0 = r$, that is $C_{\mathbf{x}}$ has r principal subcurves. Since $C_{\mathbf{x}}$ has $r + 1$ irreducible components, this implies that each principal subcurve contains exactly one irreducible component, and there is only one non-principal component. As usual, denote by $C_{\mathbf{x}}^0, \dots, C_{\mathbf{x}}^r$ the irreducible components of $C_{\mathbf{x}}$, where the marked points contained in $C_{\mathbf{x}}^j$ are recorded in I_j . Without loss of generality, we can assume that $C_{\mathbf{x}}^0$ is the unique non-principal component of $C_{\mathbf{x}}$. Since two principal subcurves cannot be adjacent, there must be a node between $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^j$, for all $j \in \{1, \dots, r\}$. It follows that the partition $\{I_0, \dots, I_r\}$ satisfies (24). The proof of the converse statement is straightforward. \square

We now prove

Theorem 4.5. *The set of irreducible components of $\partial\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is in bijection with the set of partitions $\mathcal{S} = \{I_0, I_1, \dots, I_r\}$ of $\{1, \dots, n\}$ with $r \geq 1$ such that either $r = 1$ and $\{I_0, I_1\} \in \mathcal{P}$, or $r \geq 2$ and \mathcal{S} satisfies (24).*

Proof. Let \hat{D} be an irreducible component of $\partial\widehat{\mathcal{M}}_{0,n}(\underline{k})$ and S the stratum of $\overline{\mathcal{M}}_{0,n}$ such that $\hat{D} = \overline{\hat{p}^{-1}(S)}$. Let r be the codimension of S in $\overline{\mathcal{M}}_{0,n}$. If $r = 1$, then S is the set of generic points of a boundary divisor in $\partial\overline{\mathcal{M}}_{0,n}$, which corresponds to a partition $\{I_0, I_1\} \in \mathcal{P}$. In this case \hat{D} is the proper transform of the corresponding boundary divisor of $\overline{\mathcal{M}}_{0,n}$. It is clear that the correspondence $\hat{D} \mapsto \{I_0, I_1\}$ is one-to-one.

If $r > 1$, then it follows from Proposition 4.4 that the partition $\{I_0, I_1, \dots, I_r\}$ associated to S satisfies the required condition. The correspondence $\hat{D} \mapsto \{I_0, \dots, I_r\}$ is also one-to-one because given $\{I_0, \dots, I_r\}$, there is a unique configuration of the dual graph such that the curves parametrized by S has exactly r principle components. \square

Definition 4.6. *We denote by $\hat{\mathcal{P}}(\mu)$ the set of partitions of $\{1, \dots, n\}$ that are associated to the irreducible components of $\partial\widehat{\mathcal{M}}_{0,n}(\underline{k})$. The divisor of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ associated to \mathcal{S} is denoted by $\hat{D}_{\mathcal{S}}$, its projection in $\overline{\mathcal{M}}_{0,n}$ is denoted by $D_{\mathcal{S}}$. Remark that if $\mathcal{S} \in \mathcal{P}$ then $D_{\mathcal{S}}$ is the boundary divisor of $\overline{\mathcal{M}}_{0,n}$ which is associated to \mathcal{S} , and $\hat{D}_{\mathcal{S}}$ is the proper transform of $D_{\mathcal{S}}$.*

5. GEOMETRIC INTERPRETATION

5.1. Differentials associated to points in the blow-up. We now give the geometric interpretation of the points in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. Let \mathbf{x} be a point in $\overline{\mathcal{M}}_{0,n}$, and $\mathcal{U}_{\mathbf{x}}$ a neighborhood of \mathbf{x} satisfying the conditions (i) and (ii) in §3.1. In what follows, to simplify the notation, we will write \mathcal{U} and $\widehat{\mathcal{U}}$ instead of $\mathcal{U}_{\mathbf{x}}$ and $\widehat{\mathcal{U}}_{\mathbf{x}}$. Recall from Theorem 2.5 we have constructed $(r + 1)$ trivializing sections $t^{\beta_0}\Phi_0, \dots, t^{\beta_r}\Phi_r$ of the Kawamata line bundle \mathcal{K}_{μ} over $\overline{\mathcal{C}}_{0,n}|\mathcal{U}$. By multiplying Φ_1, \dots, Φ_r by some non-vanishing holomorphic functions on \mathcal{U} , we can actually write

$$t^{\beta_0}\Phi_0 = \dots = t^{\beta_{r-1}}\Phi_{r-1} = t^{\beta_r}\Phi_r.$$

For every $\mathbf{x}' \in \mathcal{U}$, denote by $\Phi_j(\mathbf{x}')$ the restriction of Φ_j to the curve $C_{\mathbf{x}'}$. Consider a vector $v := (v_0, \dots, v_r) \in \mathbb{C}^{r+1} \setminus \{0\}$ such that $(\mathbf{x}', [v]) \in \hat{p}^{-1}(\{\mathbf{x}'\})$, where $[v] \in \mathbb{P}_{\mathbb{C}}^r$ is the line generated by v .

Assume that $\mathbf{x}' \in \mathcal{U}^* := \mathcal{U} \cap \mathcal{M}_{0,n}$ (that is $t_\alpha(\mathbf{x}') \neq 0$ for all $\alpha \in \mathcal{N}(\mathbf{x})$). In this case, we have

$$\hat{p}^{-1}(\{\mathbf{x}'\}) = (\mathbf{x}', [t^{\beta_0}(\mathbf{x}') : \cdots : t^{\beta_r}(\mathbf{x}')]) \in \mathcal{U} \times \mathbb{P}_{\mathbb{C}}^r.$$

Hence $v = \lambda \cdot (t^{\beta_0}(\mathbf{x}'), \dots, t^{\beta_r}(\mathbf{x}'))$ for some $\lambda \in \mathbb{C}^*$. Therefore, $v_0\Phi_0(\mathbf{x}') = \cdots = v_r\Phi_r(\mathbf{x}')$, which means that all the elements in the family $\{v_0\Phi_0(\mathbf{x}'), \dots, v_r\Phi_r(\mathbf{x}')\}$ give the same d -differential on $C_{\mathbf{x}'}$. Thus to every element $(\mathbf{x}', [v]) \in \hat{p}^{-1}(\mathcal{U}^*)$ we have an associated d -differential on $C_{\mathbf{x}'}$ determined up to multiplication by a scalar in \mathbb{C}^* , that is an element of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(k)$.

More generally, for all $\mathbf{x}' \in \mathcal{U}$, let $C_{\mathbf{x}'}^0, \dots, C_{\mathbf{x}'}^{r'}$ denote the irreducible components of $C_{\mathbf{x}'}$. Associated to $C_{\mathbf{x}'}$ we have a partition $\{J_0, \dots, J_{r'}\}$ of the set $\{0, \dots, r\}$ defined as follows: let $f : C_{\mathbf{x}'} \rightarrow C_{\mathbf{x}}$ be a surjection that pinches some simple closed curves in $C_{\mathbf{x}'}$ into nodes in $C_{\mathbf{x}}$. Then for all $k \in \{0, \dots, r'\}$, J_k is the set of indexes $i \in \{0, \dots, r\}$ such that $f(C_{\mathbf{x}'}^k)$ contains $C_{\mathbf{x}}^i$. In other words, $\{C_{\mathbf{x}}^i, i \in J_k\}$ are the irreducible components of the subcurve $f(C_{\mathbf{x}'}^k) \subset C_{\mathbf{x}}$. Note that the partition $\{J_0, \dots, J_{r'}\}$ depends only on the stratum of \mathbf{x}' in $\overline{\mathcal{M}}_{0,n}$.

For all $i, j \in \{0, \dots, r\}$, by definition, we must have $v_i t^{\beta_j} = v_j t^{\beta_i}$ and $t^{\beta_i}\Phi_i = t^{\beta_j}\Phi_j$. But since $t^{\beta_i}(\mathbf{x}')$ and $t^{\beta_j}(\mathbf{x}')$ can vanish, $v_i\Phi_i(\mathbf{x}')$ may not be equal to $v_j\Phi_j(\mathbf{x}')$ in general. However, we have

Lemma 5.1. *For all $i, j \in J_k$, $v_i\Phi_i(\mathbf{x}')$ and $v_j\Phi_j(\mathbf{x}')$ give the same d -differential on the component $C_{\mathbf{x}'}^k$.*

Proof. We first remark that the function $t^{\beta_i - \beta_j} := t^{\beta_i}/t^{\beta_j}$ involves only t_α where the node α separates $C_{\mathbf{x}}^i$ and $C_{\mathbf{x}}^j$. In $C_{\mathbf{x}'}$ these nodes are smoothed, which means that $t_\alpha(\mathbf{x}') \neq 0$ for all α 's that separate $C_{\mathbf{x}}^i$ and $C_{\mathbf{x}}^j$. Therefore, $t^{\beta_i - \beta_j}(\mathbf{x}') \neq 0$.

Obviously, we only need to consider the case $(v_i, v_j) \neq (0, 0)$. Without loss of generality, we can suppose that $v_j \neq 0$. By definition, there is a sequence $\{(\mathbf{x}'_s, [v^{(s)}])\}_{s \in \mathbb{N}} \subset \hat{p}^{-1}(\mathcal{U}^*)$, where $v^{(s)} = (v_0^{(s)}, \dots, v_r^{(s)})$, converging to $(\mathbf{x}', [v])$. Since $\mathbf{x}'_s \in \mathcal{U}^*$, we have $v_i^{(s)}/v_j^{(s)} = t^{\beta_i}(\mathbf{x}'_s)/t^{\beta_j}(\mathbf{x}'_s) = t^{\beta_i - \beta_j}(\mathbf{x}'_s)$. It follows that

$$\frac{v_i}{v_j} = \lim_{s \rightarrow \infty} \frac{v_i^{(s)}}{v_j^{(s)}} = \lim_{s \rightarrow \infty} t^{\beta_i}(\mathbf{x}'_s)/t^{\beta_j}(\mathbf{x}'_s) = t^{\beta_i - \beta_j}(\mathbf{x}').$$

Remark that $\frac{\Phi_j}{\Phi_i} = t^{\beta_i - \beta_j}$ is a well defined function in a neighborhood of \mathbf{x}' . Thus we have

$$\frac{\Phi_j(\mathbf{x}')}{\Phi_i(\mathbf{x}')} = t^{\beta_i - \beta_j}(\mathbf{x}') = \frac{v_i}{v_j}$$

which implies that $v_i\Phi_i(\mathbf{x}') = v_j\Phi_j(\mathbf{x}')$.

We now observe that the restrictions of $\Phi_i(\mathbf{x}')$ and $\Phi_j(\mathbf{x}')$ to the component $C_{\mathbf{x}'}^k$ give non-zero d -differentials. This is because the restrictions of the projections π_i and π_j to $C_{\mathbf{x}'}^k$ give two isomorphisms from $C_{\mathbf{x}'}^k$ onto $\mathbb{P}_{\mathbb{C}}^1$, and Φ_i and Φ_j are the pullbacks of non-vanishing d -differentials on $\mathbb{P}_{\mathbb{C}}^1$ by π_i and π_j respectively. Therefore, $v_i\Phi_i(\mathbf{x}')$ and $v_j\Phi_j(\mathbf{x}')$ give the same d -differential on $C_{\mathbf{x}'}^k$. \square

Denote by $q_k(\mathbf{x}', v)$ the d -differential given by $v_j\Phi_j(\mathbf{x}')$ on $C_{\mathbf{x}'}^k$. By Lemma 5.1 $q_k(\mathbf{x}', v)$ is independent of the choice of $j \in J_k$. Thus for every $(\mathbf{x}', [v]) \in \widehat{\mathcal{U}}$, we can associate to $(\mathbf{x}', [v])$ the tuple $q(\mathbf{x}', [v]) := (q_0(\mathbf{x}', v), \dots, q_{r'}(\mathbf{x}', v))$ of d -differentials on the components of $C_{\mathbf{x}'}$, which are determined up to simultaneous multiplication by the same scalar in \mathbb{C}^* .

5.2. Embedding into $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$. Let us now have a closer look at the d -differentials in the family $(q_0(\mathbf{x}', v), \dots, q_{r'}(\mathbf{x}', v))$. Lemma 3.8 (see also Remark 3.17) applied to \mathbf{x}' implies that if $C_{\mathbf{x}'}^k$ is not a principal component of $C_{\mathbf{x}'}$, then $v_j = 0$ for all $j \in J_k$, hence $q_k(\mathbf{x}', v) = 0$. Assume now that $C_{\mathbf{x}'}^k$ is a principal component of $C_{\mathbf{x}'}$. Let $\{x'_1, \dots, x'_n\}$ be the marked points of $C_{\mathbf{x}'}$. By construction $q_k(\mathbf{x}', v)$ has a zero of order k_i at every point $x'_i \in C_{\mathbf{x}'}^k$. Recall that by definition, we have $k_i \geq 1 - d$. The remaining zeros and poles of $q_k(\mathbf{x}', v)$ are the points in $C_{\mathbf{x}'}^k$ that correspond to nodes of $C_{\mathbf{x}'}$. The order of the zero at such a point y' is given by $-d\mu(y')$, where $\mu(y')$ is the weight of y' defined in (18). Since $C_{\mathbf{x}'}^k$ is principal, we have $\mu(y') \leq 1$, hence $q_k(\mathbf{x}', v)$ has a pole of order at most d at y' . As a consequence, we see that the tuple $(q_0(\mathbf{x}', v), \dots, q_{r'}(\mathbf{x}', v))$ actually defines a holomorphic section of the line bundle $\mathcal{K}_{\mathbf{x}'}^{(d)} := d\omega_{C_{\mathbf{x}'}} + \sum_{i=1}^n (d-1) \cdot \{x'_i\}$, where $\omega_{C_{\mathbf{x}'}}$ is the dualizing sheaf of $C_{\mathbf{x}'}$.

Let $\varphi : \tilde{C}_{\mathbf{x}'} \rightarrow C_{\mathbf{x}'}$ be the normalization of $C_{\mathbf{x}'}$. For $k = 0, \dots, r'$, the connected component of $\tilde{C}_{\mathbf{x}'}$ corresponding to $C_{\mathbf{x}'}^k$ is denoted by $\tilde{C}_{\mathbf{x}'}^k$. We consider $\{x'_1, \dots, x'_n\}$ as a subset of $\tilde{C}_{\mathbf{x}'}$. For each $k \in \{0, \dots, r'\}$, denote by P_k the set of indices $i \in \{1, \dots, n\}$ such that $x'_i \in \tilde{C}_{\mathbf{x}'}^k$. Let N_k denote the set of points in $\tilde{C}_{\mathbf{x}'}^k$ that correspond to the nodes of $C_{\mathbf{x}'}$. One can readily check that

$$\varphi^*(\omega_{C_{\mathbf{x}'}} - \mathcal{K}_{\mathbf{x}'}^{(d)})|_{\tilde{C}_{\mathbf{x}'}^k} \sim (1-d) \left(\omega_{\tilde{C}_{\mathbf{x}'}^k} + \sum_{y_j \in N_k} \{y_j\} + \sum_{i \in P_k} \{x'_i\} \right).$$

In particular

$$\deg \varphi^*(\omega_{C_{\mathbf{x}'}} - \mathcal{K}_{\mathbf{x}'}^{(d)})|_{\tilde{C}_{\mathbf{x}'}^k} = (1-d)(|N_k| + |P_k| - 2) < 0,$$

which implies that $\dim H^0(\tilde{C}_{\mathbf{x}'}, \varphi^*(\omega_{C_{\mathbf{x}'}} - \mathcal{K}_{\mathbf{x}'}^{(d)})) = 0$. Therefore

$$(25) \quad \dim H^1(C_{\mathbf{x}'}, \mathcal{K}_{\mathbf{x}'}^{(d)}) = \dim H^0(C_{\mathbf{x}'}, \omega_{C_{\mathbf{x}'}} - \mathcal{K}_{\mathbf{x}'}^{(d)}) = 0$$

(see for instance [1, Chap.10, §2]). By Riemann-Roch theorem, (25) implies that

$$(26) \quad \dim H^0(C_{\mathbf{x}'}, \mathcal{K}_{\mathbf{x}'}^{(d)}) = \deg(\mathcal{K}_{\mathbf{x}'}^{(d)}) + 1 = (d-1)(n-2) - 1.$$

In particular, we see that $\dim H^0(C_{\mathbf{x}'}, \mathcal{K}_{\mathbf{x}'}^{(d)})$ does not depend on \mathbf{x}' . Observe that $\mathcal{K}_{\mathbf{x}'}^{(d)}$ is the restriction of the line bundle

$$\mathcal{K}_{0,n}^{(d)} := dK_{\overline{C}_{0,n}/\overline{\mathcal{M}}_{0,n}} + \sum_{i=1}^n (d-1) \cdot \Gamma_i$$

on $\overline{C}_{0,n}$ to $C_{\mathbf{x}'}$ (recall that $K_{\overline{C}_{0,n}/\overline{\mathcal{M}}_{0,n}}$ is the relative canonical line bundle of the projection $p : \overline{C}_{0,n} \rightarrow \overline{\mathcal{M}}_{0,n}$, and Γ_i is the divisor associated with the i -th section of p). It follows from a classical result (see [28, III.12.9]) that $p_*\mathcal{K}_{0,n}^{(d)}$ is a vector bundle $\mathcal{H}_{0,n}^{(d)}$ of rank $(d-1)(n-2) - 1$ over $\overline{\mathcal{M}}_{0,n}$. Denote by $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ the associated projective bundle.

We have a natural section $\mathbf{f} : \mathcal{M}_{0,n} \rightarrow \mathbb{P}\mathcal{H}_{0,n}^{(d)}$ which associates to $\mathbf{x}' = (\mathbb{P}_{\mathbb{C}}^1, x_1, \dots, x_n)$ the line generated by the d -differential $\prod_{i=1}^n (z-x_i)^{k_i} (dz)^d$ viewed as element of $H^0(C_{\mathbf{x}'}, \mathcal{K}_{0,n|C_{\mathbf{x}'}}^{(d)})$. By definition, $\mathbf{f}(\mathcal{M}_{0,n}) = \mathbb{P}\Omega^d \mathcal{M}_{0,n}(k)$.

From what we have seen, the correspondence $(\mathbf{x}', [v]) \mapsto \mathbb{C}^* \cdot (q_0(\mathbf{x}', v), \dots, q_{r'}(\mathbf{x}', v))$ provides us with a map

$$\hat{\mathbf{f}}_{\widehat{\mathcal{U}}} : \widehat{\mathcal{U}} \rightarrow \mathbb{P}\mathcal{H}_{0,n|u}^{(d)}.$$

We now show

Theorem 5.2. *The maps $\hat{\mathbf{f}}_{\widehat{\mathcal{U}}}$ are proper injective morphisms of algebraic varieties and patch together to give an injective morphism $\hat{\mathbf{f}} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \mathbb{P}\mathcal{H}_{0,n}^{(d)}$ extending the section $\mathbf{f} : \mathcal{M}_{0,n} \rightarrow \mathbb{P}\mathcal{H}_{0,n|\mathcal{M}_{0,n}}^{(d)}$.*

Proof. Consider $\widehat{\mathcal{C}}_{0,n|\mathcal{U}} := \widehat{\mathcal{U}} \times_{\mathcal{U}} \overline{\mathcal{C}}_{0,n|\mathcal{U}}$. We have the following commutative diagram

$$\begin{array}{ccc} \widehat{\mathcal{C}}_{0,n|\mathcal{U}} & \xrightarrow{p_2} & \overline{\mathcal{C}}_{0,n|\mathcal{U}} \\ p_1 \downarrow & & \downarrow p \\ \widehat{\mathcal{U}} & \xrightarrow{\hat{p}} & \mathcal{U} \end{array}$$

For $i = 0, \dots, r$, set

$$\widehat{\mathcal{U}}^i := \{(\mathbf{x}', [v_0 : \dots : v_r]) \in \widehat{\mathcal{U}}, v_i \neq 0\}, \text{ and } \widehat{\mathcal{C}}_{0,n|\mathcal{U}}^i := p_1^{-1}(\widehat{\mathcal{U}}^i).$$

We now remark that the family of differentials $\{\frac{v_0}{v_i}\Phi_0, \dots, \frac{v_r}{v_i}\Phi_r\}$ defines a holomorphic section of the line bundle $p_2^*\mathcal{K}_{0,n}^{(d)}$ over $\widehat{\mathcal{C}}_{0,n|\mathcal{U}}^i$. In fact, over $\widehat{\mathcal{C}}_{0,n|\mathcal{U}}^i$, we have $\Phi_i = \frac{v_j}{v_i}\Phi_j$ for all $j \in \{0, \dots, r\}$. By definition, this section gives a section of $p_{1*}p_2^*\mathcal{K}_{0,n}^{(d)}$ over $\widehat{\mathcal{U}}^i$. Now, since $\mathcal{K}_{0,n|\mathbb{C}_x}^{(d)} \sim \mathcal{K}_{\mathbf{x}'}^{(d)}$ for all $\mathbf{x}' \in \mathcal{U}$, it follows from (25), (26) and [28, III.12.9] (see also [1, Chap. 9, Prop.3.3]) that we have

$$p_{1*}p_2^*\mathcal{K}_{0,n}^{(d)} \sim \hat{p}^*p_*\mathcal{K}_{0,n}^{(d)} \sim \hat{p}^*\mathcal{H}_{0,n}^{(d)}.$$

Therefore, the mapping $(\mathbf{x}', [v_0 : \dots : v_n]) \in \widehat{\mathcal{U}}^i \mapsto \Phi_i(\mathbf{x}')$ gives a section of the bundle $\hat{p}^*\mathcal{H}_{0,n}^{(d)}$ over $\widehat{\mathcal{U}}^i$. Taking the image in the associated projective bundle, we get a section $\hat{\Psi}_i$ of $\hat{p}^*\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ over $\widehat{\mathcal{U}}^i$. For all $i, j \in \{0, \dots, r\}$, the sections $\hat{\Psi}_i$ and $\hat{\Psi}_j$ coincide in the intersection $\widehat{\mathcal{U}}^i \cap \widehat{\mathcal{U}}^j$. Thus we obtain a holomorphic section $\hat{\Psi} : \widehat{\mathcal{U}} \rightarrow \hat{p}^*\mathbb{P}\mathcal{H}_{0,n}^{(d)}$. By definition the map $\hat{\mathbf{f}}_{\widehat{\mathcal{U}}}$ is the composition $\hat{q} \circ \hat{\Psi}$, where $\hat{q} : \hat{p}^*\mathbb{P}\mathcal{H}_{0,n}^{(d)} \rightarrow \mathbb{P}\mathcal{H}_{0,n|\mathcal{U}}^{(d)}$ is the natural projection. It follows that $\hat{\mathbf{f}}_{\widehat{\mathcal{U}}}$ is a morphism of algebraic varieties.

It is straightforward to check that $\hat{\mathbf{f}}_{\widehat{\mathcal{U}}}$ is injective. Since $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is the blow-up along the sheaf of ideals \mathcal{I} , the maps $\hat{\mathbf{f}}_{\widehat{\mathcal{U}}}$ patch together to give an embedding of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ into $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ whose restriction to $\mathcal{M}_{0,n}$ equals \mathbf{f} . \square

5.3. Proof of Theorem 1.1.

Proof. The first assertion is the content of Theorem 3.1. For the second assertion, we have $\hat{\mathbf{f}}(\mathcal{M}_{0,n}) = \mathbf{f}(\mathcal{M}_{0,n}) = \mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$. Since $\mathcal{M}_{0,n}$ is dense in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ and $\hat{\mathbf{f}}$ is proper, $\hat{\mathbf{f}}(\widehat{\mathcal{M}}_{0,n}(\underline{k}))$ equals the closure $\mathbb{P}\Omega^d \widehat{\mathcal{M}}_{0,n}(\underline{k})$ of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ in $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$, that is the incidence variety compactification of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$. Since $\hat{\mathbf{f}}$ is an embedding, we conclude that $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is isomorphic to $\mathbb{P}\Omega^d \widehat{\mathcal{M}}_{0,n}(\underline{k})$. The third assertion is the content of Theorem 3.13, and the fourth assertion is the content of Theorem 4.5. \square

6. VOLUME FORM ON MODULI SPACE OF d -DIFFERENTIALS

In this section, we review some relevant properties of the moduli spaces of d -differentials. In what follows, g will be a fixed non-negative integer, $\underline{k} := (k_1, \dots, k_n)$ is a vector of integers such that $k_i \geq 1 - d$ and $k_1 + \dots + k_n = d(2g - 2)$.

6.1. Canonical cyclic cover and local chart. Recall that $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is the space of pairs (X, q) where X is a compact Riemann surface of genus g and q is a meromorphic d -differential on X whose zeros and poles have orders prescribed by \underline{k} (we always suppose that the zeros and poles of q are numbered). The quotient $\Omega^d \mathcal{M}_{g,n}(\underline{k})/\mathbb{C}^*$ is denoted by $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$. A d -differential is said to be *primitive* if it is not a power of some d' -differential, with $d' < d$, on the same Riemann surface. The spaces $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ have been studied by different approaches. In particular, we have

Theorem 6.1 ([30, 53, 5, 50]). *Each stratum $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ of primitive d -differentials with $d > 1$ is a complex orbifold of dimension $2g + n - 2$.*

Remark 6.2. *It follows from Theorem 6.1 that $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is a complex orbifold of dimension $2g + n - 3$.*

Let ζ be a fixed primitive d -th root of unity. Let (X, q) be a primitive d -differential in $\Omega^d \mathcal{M}_{g,n}(\underline{k})$. The *canonical cyclic covering* of (X, q) is a triple $(\hat{X}, \hat{\omega}, \tau)$, where \hat{X} is a compact Riemann surface, $\hat{\omega}$ a holomorphic 1-form on \hat{X} , and τ an automorphism of order d of \hat{X} such that

- (i) X is isomorphic to $\hat{X}/\langle \tau \rangle$,
- (ii) $\hat{\omega}^d = \varpi^* q$, where $\varpi : \hat{X} \rightarrow X \simeq \hat{X}/\langle \tau \rangle$ is the canonical projection,
- (iii) $\tau^* \hat{\omega} = \zeta \hat{\omega}$.

It is well known that such a triple always exists and is unique up to isomorphism (see for instance [21, 5, 46]).

Let \hat{Z} denote the inverse image of the set of zeros and poles of q in \hat{X} . Since τ preserves \hat{Z} , τ induces an isomorphism τ^* of $H^1(\hat{X}, \hat{Z}, \mathbb{C})$. Define

$$V_\zeta := \ker(\tau^* - \zeta \text{id}) \subset H^1(\hat{X}, \hat{Z}, \mathbb{C}).$$

One can then define a *period mapping* $\Phi : \tilde{U} \rightarrow V_\zeta$, where \tilde{U} is a finite cover of a neighborhood U of (X, q) in $\Omega^d \mathcal{M}_{g,n}(\underline{k})$, which is locally biholomorphic. The period mappings form an atlas of orbifold local charts for $\Omega^d \mathcal{M}_{g,n}(\underline{k})$. Since the transition functions of those charts are given by linear isomorphisms, this atlas endows $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ with a complex affine orbifold structure (see [5] or [46] for more details). Note also that Φ induces a local chart $\hat{\Phi}$ of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ which maps a neighborhood of $(X, \mathbb{C}^* \cdot q)$ onto an open subset of the projective space $\mathbb{P}V_\zeta$.

6.2. Volume form and tautological line bundle. Let $\mathbf{p} : H^1(\hat{X}, \hat{Z}; \mathbb{C}) \rightarrow H^1(\hat{X}; \mathbb{C})$ denote the canonical projection, that is for any $\eta \in H^1(\hat{X}, \hat{Z}; \mathbb{C})$, $\mathbf{p}(\eta)$ is the restriction of η to the absolute cycles in $H_1(\hat{X}, \mathbb{Z})$. It is not difficult to see that $\mathbf{p}(V_\zeta) = \ker(\tau^* - \zeta \text{Id}) \subset H^1(\hat{X}, \mathbb{C})$ (see for instance [46, Lem.4.1]). Moreover, we have

$$(27) \quad \dim(\ker \mathbf{p} \cap V_\zeta) = \text{card}(\{k_1, \dots, k_n\} \cap (d\mathbb{Z}))$$

(for a proof see for instance [46, Prop.4.2]). In [46], it is shown that $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ carries a canonical volume form $d\text{vol}$ which is parallel with respect to its affine complex orbifold structure. In the case $\ker \mathbf{p} \cap V_\zeta = \{0\}$, the volume form $d\text{vol}$ is defined as follows: on $H^1(\hat{X}, \mathbb{C})$ we have a Hermitian form (\cdot, \cdot) arising from the intersection form on $H^1(\hat{X}, \mathbb{R})$. It is not difficult to see that the restriction of (\cdot, \cdot) to $\mathbf{p}(V_\zeta)$ is non-degenerate.

By a slight abuse of notation, we also denote the pullback of this Hermitian form to $H^1(\hat{X}, \hat{Z}, \mathbb{C})$ by (\cdot, \cdot) . Since $\mathbf{p}|_{V_\zeta}$ is an isomorphism, the restriction of (\cdot, \cdot) to V_ζ is also non-degenerate. Denote by ϑ the

imaginary part of $(\cdot, \cdot)_{|V_\zeta}$. Then $\frac{1}{N!} \vartheta^N$ is a volume form on V_ζ , where $N = \dim V_\zeta = \dim \Omega^d \mathcal{M}_{g,n}(\underline{k})$. Concretely, if (z_1, \dots, z_N) is a system of coordinates on V_ζ in which $(\cdot, \cdot)_{|V_\zeta}$ is given by the matrix $\text{diag}(\underbrace{1, \dots, 1}_r, \underbrace{-1, \dots, -1}_s)$, where $r + s = N$, then

$$\frac{1}{N!} \vartheta^N = (-1)^s \left(\frac{t}{2}\right)^N dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge dz_N \wedge d\bar{z}_N.$$

Using local charts by period mappings, this volume form gives rise to a well defined volume form $d\text{vol}$ on $\Omega^d \mathcal{M}_{g,n}(\underline{k})$.

By construction, $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is a \mathbb{C}^* -principal bundle over $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$. Denote by \mathcal{L} the associated line bundle. We will call \mathcal{L} the *tautological line bundle* of $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$. The space $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ can be naturally identified with the complement of the zero section in the total space of \mathcal{L} . In the local charts by period mappings, where $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ is locally identified with $\mathbb{P}V_\zeta$, the line bundle \mathcal{L} is identified with $\mathcal{O}(-d)_{\mathbb{P}V_\zeta}$, that is the d -th tensor power of the tautological line bundle $\mathcal{O}(-1)_{\mathbb{P}V_\zeta}$. Recall that we can also view $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ as a subspace of $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$. In this setting, \mathcal{L} is simply the restriction of $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ to $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$.

There is a natural Hermitian metric on \mathcal{L} which is defined as follows: an element ξ of \mathcal{L} outside of the zero section is simply a pair $(X, q) \in \Omega^d \mathcal{M}_{g,n}(\underline{k})$. It is well known that $|q|^{\frac{2}{d}}$ defines an Euclidean metric on X with conical singularities at the zeros and poles of q . We define the norm of ξ to be

$$\|\xi\| := \left(\text{Area}(X, |q|^{\frac{2}{d}})\right)^{\frac{d}{2}} = \left(\int_X q^{\frac{2}{d}}\right)^{\frac{d}{2}}.$$

Equivalently,

$$\|\xi\| = \left(\frac{1}{d} \text{Area}(\hat{X}, |\hat{\omega}|^2)\right)^{\frac{d}{2}} = \left(\frac{t}{2d} \int_{\hat{X}} \hat{\omega} \wedge \bar{\hat{\omega}}\right)^{\frac{d}{2}},$$

where $(\hat{X}, \hat{\omega}, \tau)$ is the canonical cyclic cover of (X, q) . Note that $\left(\frac{t}{2} \int_{\hat{X}} \hat{\omega} \wedge \bar{\hat{\omega}}\right)^{\frac{1}{2}}$ is the Hodge norm of $\hat{\omega}$. We will call $\|(X, q)\|$ the Hodge norm of (X, q) .

Let $\Omega_1^d \mathcal{M}_{g,n}(\underline{k})$ denote the set of $(X, q) \in \Omega^d \mathcal{M}_{g,n}(\underline{k})$ such that $\|(X, q)\| < 1/d$. Let $\text{pr}_1 : \Omega_1^d \mathcal{M}_{g,n}(\underline{k}) \rightarrow \mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ be the natural projection. Define vol_1 to be the pushforward of the restriction of vol to $\Omega_1^d \mathcal{M}_{g,n}(\underline{k})$ onto $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ by pr_1 . Then vol_1 is actually a volume form on $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$. Moreover, we have (see [35, Lem.3.2])

Lemma 6.3. *Assume that none of the k_1, \dots, k_n is a multiple of d . Then*

$$(28) \quad d\text{vol}_1 = \frac{1}{d^N} \cdot \frac{2\pi}{2^N N!} \cdot (-t\Theta)^{N-1}$$

pointwise on $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$, where $\dim \mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k}) = N - 1$, and Θ is the curvature form of the Hodge norm on \mathcal{L} .

Note that our definitions of $d\text{vol}$ and $d\text{vol}_1$ in the present paper and the ones in [35] are slightly different, so that the constant on the right hand side of (28) does not depend on the signature of $(\cdot, \cdot)_{|V_\zeta}$.

6.3. Comparison with Masur-Veech volume. If $d \in \{1, 2, 3, 4, 6\}$, there is another natural volume form on $\Omega^d \mathcal{M}_{g,n}(\underline{k})$ that we now describe. Recall that if A is a \mathbb{Z} -module, then $H^1(\hat{X}, \hat{Z}, A)$ is the space of morphisms of \mathbb{Z} -modules $\eta : H_1(\hat{X}, \hat{Z}, \mathbb{Z}) \rightarrow A$. Let

$$\Lambda_\zeta = \begin{cases} V_\zeta \cap H^1(\hat{X}, \hat{Z}, \mathbb{Z}[t]) & \text{if } d \in \{2, 4\}, \\ V_\zeta \cap H^1(\hat{X}, \hat{Z}, \mathbb{Z}[e^{\frac{2\pi i}{3}}]) & \text{if } d \in \{3, 6\}. \end{cases}$$

Then Λ_ζ is a lattice of V_ζ . There is unique volume form on V_ζ such that the co-volume of Λ_ζ is 1. Since the transition maps of the local charts by the period mappings preserves Λ_ζ , this volume form is well defined on $\Omega^d \mathcal{M}_{g,n}(\underline{k})$. It will be referred to as the *Masur-Veech measure* and denoted by $d\text{vol}^{MV}$. This volume form induces a volume form $d\text{vol}_1^{MV}$ on $\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})$ in the same manner as $d\text{vol}_1$ is induced from $d\text{vol}$. In fact, the Masur-Veech volume arises naturally from the counting problem of tilings of surfaces by triangles and squares (see for instance [18, 12, 11, 35]). From the definition, there is a constant $\lambda \in \mathbb{R}^*$ such that $d\text{vol} = \lambda d\text{vol}^{MV}$ and $d\text{vol}_1 = \lambda d\text{vol}_1^{MV}$. It can be shown that the constant λ always belongs to either \mathbb{Q} or $\sqrt{3}\mathbb{Q}$ (see [46, Prop.5.9]).

7. VOLUME AND SELF-INTERSECTION OF DIVISORS

7.1. Divisors representing the tautological line bundle. Let $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ denote the tautological line bundle over $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$. Define $\hat{\mathcal{L}}_\mu$ to be the pullback of $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ to $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ by $\hat{\mathbf{f}}$. We will call $\hat{\mathcal{L}}_\mu$ the tautological line bundle over $\widehat{\mathcal{M}}_{0,n}(\underline{k})$.

Recall that \mathcal{L}_μ is the line bundle over $\mathcal{M}_{0,n}$ whose fiber over a point $\mathbf{x} = (\mathbb{P}^1, x_1, \dots, x_n)$ is the complex line generated by the d -differential $q_{\mathbf{x}} := \prod (z - x_i)^{k_i} (dz)^d$. By construction, we have

Lemma 7.1. *The restriction of $\hat{\mathcal{L}}_\mu$ to $\mathcal{M}_{0,n}$ is isomorphic to \mathcal{L}_μ .*

Our goal now is to show the following

Proposition 7.2. *Let \mathcal{E} be the exceptional divisor of the blow-up $\hat{p} : \widehat{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \overline{\mathcal{M}}_{0,n}$. We abusively denote also by \mathcal{E} the associated line bundle on $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. Then we have $\hat{\mathcal{L}}_\mu \sim \hat{p}^* \bar{\mathcal{L}}_\mu + \mathcal{E}$, where $\bar{\mathcal{L}}_\mu$ is the line bundle over $\overline{\mathcal{M}}_{0,n}$ in Theorem 2.2. As a consequence, $\hat{\mathcal{L}}_\mu$ is isomorphic to the line bundle associated to the divisor $\hat{\mathcal{D}}_\mu := \hat{p}^* \mathcal{D}_\mu + \mathcal{E}$.*

Proof. By definition, \mathcal{E} is a Cartier divisor in $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ whose defining sheaf is $\hat{p}^{-1} \mathcal{I}$. Consider a point \mathbf{x} in a stratum of codimension $r \geq 1$ of $\overline{\mathcal{M}}_{0,n}$. Let $\mathcal{U}_{\mathbf{x}}, (t, u), \beta_0, \dots, \beta_r$ be as in §3.1. Then $\widehat{\mathcal{U}}_{\mathbf{x}} = \hat{p}^{-1}(\mathcal{U}_{\mathbf{x}})$ is the irreducible component of the algebraic set

$$\{(t, u, [v_0 : \dots : v_r]) \in \mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^{r-1}, v_k t^{\beta_j} = v_j t^{\beta_k}, j, k \in \{0, \dots, r\}\}.$$

which contains the set $\{(t, u, [t^{\beta_0} : \dots : t^{\beta_r}]), (t, u) \in \mathcal{U}_{\mathbf{x}}^*\}$. Observe that $\widehat{\mathcal{U}}_{\mathbf{x}}$ is covered by the family of open subsets

$$\widehat{\mathcal{U}}_{\mathbf{x}}^i := \{(t, u, [v_0 : \dots : v_r]) \in \widehat{\mathcal{U}}_{\mathbf{x}}, v_i \neq 0\}, i = 0, \dots, r.$$

On each $\widehat{\mathcal{U}}_{\mathbf{x}}^i$, we can assume that $v_i = 1$. Hence $t^{\beta_j} = v_j t^{\beta_i}$, for all $j \in \{1, \dots, r\}$. It follows that $\hat{p}^{-1} \mathcal{I}|_{\widehat{\mathcal{U}}_{\mathbf{x}}^i}$ is generated by t^{β_i} . Note that for any $j \in \{0, \dots, r\}$ such that $j \neq i$, we have $\frac{\beta_i}{\beta_j} = \frac{v_i}{v_j}$. Thus

t^{β_i}/t^{β_j} is a non-vanishing regular function on $\widehat{\mathcal{U}}_x^i \cap \widehat{\mathcal{U}}_x^j$. This means that \mathcal{E} is the Cartier divisor whose local equation on $\widehat{\mathcal{U}}_x^i$ is t^{β_i} .

In the proof of Theorem 5.2, we have seen that a trivialization of $\widehat{\mathcal{L}}_\mu|_{\widehat{\mathcal{U}}_x^i} \simeq \mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}|_{\widehat{\mathcal{U}}_x^i}}$ is given by Φ_i . Since we can write $\Phi_i = \frac{t^{\beta_i}\Phi_i}{t^{\beta_i}}$, and $t^{\beta_i}\Phi_i$ is a trivialization of the bundle $\widehat{\mathcal{L}}_\mu$ over \mathcal{U}_x (see Theorem 2.5) and $1/t^{\beta_i}$ is a trivializing section of \mathcal{E} over $\widehat{\mathcal{U}}_x^i$, the proposition follows. \square

7.2. Multiscale differential compactification. In [10], based on the construction in [6], Costantini-Möller-Zachhuber construct a compactification $\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})$ of $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$, which is a complex algebraic orbifold such that the complement of $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ in $\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})$ is a normal crossing divisor. Elements of $\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})$ are called *projectivized multiscale differentials*. The space $\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})$ comes equipped with a tautological line bundle denoted by $\mathcal{O}(-1)_{\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})}$. Moreover, we have a natural morphism $\mathcal{P} : \mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k}) \rightarrow \mathbb{P}\mathcal{H}_{0,n}^{(d)}$ such that the restriction of \mathcal{P} to $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ is the identity, and we have

$$(29) \quad \mathcal{P}^* \mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}} \sim \mathcal{O}(-1)_{\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})}^{\otimes d}.$$

Note that $\mathcal{P}(\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k}))$ must be the closure of $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ in $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$, that is the Incidence Variety Compactification $\mathbb{P}\Omega^d\overline{\mathcal{M}}_{0,n}(\underline{k})$.

Denote the restriction of $\mathcal{O}(-1)_{\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})}$ to $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ by \mathcal{L}' . In local charts by period mappings where $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ is locally identified with $\mathbb{P}V_\zeta^+$, \mathcal{L}' is identified with $\mathcal{O}(-1)_{\mathbb{P}V_\zeta^+}$, that is the restriction of $\mathcal{O}(-1)_{\mathbb{P}V_\zeta}$ to $\mathbb{P}V_\zeta^+$ (see §6.2). By definition, the intersection form (\cdot, \cdot) provides us with a Hermitian metric on $\mathcal{O}(-1)_{\mathbb{P}V_\zeta^+}$. Denote by $\|\cdot\|'$ the induced Hermitian metric on \mathcal{L}' . We will call $\|\cdot\|'$ the Hodge norm of \mathcal{L}' . Let $\Theta' \in \Omega^{1,1}(\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k}))$ denote the curvature form of $\|\cdot\|'$. The following result is proved in [10]

Theorem 7.3 ([10], Theorem 1.4). *The $(1, 1)$ -form $\frac{1}{2\pi}\Theta'$ on $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ defines a closed current on $\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})$ which represents the first Chern class of $\mathcal{O}(-1)_{\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})}$. Moreover, for every $k \in \mathbb{N}$, $(\frac{1}{2\pi} \cdot \Theta')^k$ is also a closed current representing the cohomology class $c_1^k(\mathcal{O}(-1)_{\mathbb{P}\Xi^d\overline{\mathcal{M}}_{0,n}(\underline{k})})$.*

7.3. Proof of Theorem 1.3.

Proof. That $\widehat{\mathcal{D}}_\mu$ represents the restriction of $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ to $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ is a consequence of Proposition 7.2. It remains to prove equality (8).

Denote the restriction of $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ to $\mathbb{P}\Omega^d\mathcal{M}_{0,n}(\underline{k})$ by \mathcal{L} . It follows from (29) that we have $\mathcal{L} \sim \mathcal{L}'^{\otimes d}$. Moreover, the curvature form Θ of the Hodge norm on \mathcal{L} satisfies

$$(30) \quad \Theta = d\Theta'.$$

Assume that d does not divide k_i for all $i = 1, \dots, n$. It follows from Lemma 6.3 and (30) that we have

$$(31) \quad d\text{vol}_1 = \frac{1}{d} \cdot \frac{2\pi}{2^{n-2}(n-2)!} \cdot (-t\Theta')^{n-3}$$

at every point of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$. Therefore

$$\begin{aligned} \text{vol}_1(\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})) &= \frac{1}{d} \cdot \frac{2\pi}{2^{n-2}(n-2)!} \int_{\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})} (-t\Theta')^{n-3} \\ &= \frac{(-1)^{n-3}}{d} \cdot \frac{(2\pi)^{n-2}}{2^{n-2}(n-2)!} \int_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})} c_1^{n-3}(\mathcal{O}(-1)_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})}) \quad (\text{by Theorem 7.3}) \end{aligned}$$

By Theorem 1.1, we can consider \mathcal{P} as a surjective birational morphism from $\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})$ onto $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. Hence

$$\begin{aligned} \text{vol}_1(\mathbb{P}\Omega^d \mathcal{M}_{g,n}(\underline{k})) &= \frac{(-1)^{n-3}}{d^{n-2}} \cdot \frac{(2\pi)^{n-2}}{2^{n-2}(n-2)!} \int_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})} c_1^{n-3}(\mathcal{P}^* \mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(n)}}) \\ &= \frac{(-1)^{n-3}}{d^{n-2}} \cdot \frac{(2\pi)^{n-2}}{2^{n-2}(n-2)!} \int_{\mathbb{P}\Xi^d \overline{\mathcal{M}}_{0,n}(\underline{k})} c_1^{n-3}(\mathcal{P}^* \hat{\mathcal{L}}_\mu) \\ &= \frac{(-1)^{n-3}}{d^{n-2}} \cdot \frac{(2\pi)^{n-2}}{2^{n-2}(n-2)!} (\mathcal{P}^* \hat{\mathcal{D}}_\mu)^{n-3} \quad (\text{since } \hat{\mathcal{L}}_\mu \sim \hat{\mathcal{D}}_\mu \text{ by Proposition 7.2}) \\ &= \frac{(-1)^{n-3}}{d^{n-2}} \cdot \frac{(2\pi)^{n-2}}{2^{n-2}(n-2)!} (\hat{\mathcal{D}}_\mu)^{n-3} \quad (\text{since } \mathcal{P} \text{ is birational}). \end{aligned}$$

Theorem 1.3 is then proved. \square

8. TAUTOLOGICAL LINE BUNDLE ON THE BOUNDARY OF $\widehat{\mathcal{M}}_{0,n}(\underline{k})$

In this section, we study the restriction of the tautological line bundle $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,n}^{(d)}}$ on irreducible components of the boundary of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$. The knowledge about this restriction will allow us to derive the recursive formula in Theorem 1.5.

Recall that by Theorem 4.5 we know that each irreducible component of the boundary of $\widehat{\mathcal{M}}_{0,n}(\underline{k})$ corresponds uniquely to a partition of $\{1, \dots, n\}$ in $\hat{\mathcal{P}}(\mu)$. We divide $\hat{\mathcal{P}}(\mu)$ into two subsets: $\hat{\mathcal{P}}_0(\mu)$ is the set of partitions $\{I_0, I_1\} \in \mathcal{P}$ such that $\mu(I_0) = \mu(I_1) = 1$, and $\hat{\mathcal{P}}_1(\mu) = \hat{\mathcal{P}}(\mu) \setminus \hat{\mathcal{P}}_0(\mu)$. Throughout this section, given $\mathcal{S} \in \hat{\mathcal{P}}(\mu)$, we denote by $D_{\mathcal{S}}^*$ the stratum of $\overline{\mathcal{M}}_{0,n}$ associated to \mathcal{S} .

8.1. Tautological line bundle on boundary divisors in $\hat{\mathcal{P}}_1(\mu)$. Consider a partition $\mathcal{S} = \{I_0, \dots, I_r\} \in \hat{\mathcal{P}}_1(\mu)$. Let $n_j = |I_j|$, $j = 0, \dots, r$. Recall that the stable curves parametrized by $D_{\mathcal{S}}^*$ have $r+1$ components each of which corresponds to a subset I_j in the partition \mathcal{S} . Moreover, there is a node between the component associated to I_0 and the component associated to I_j for all $j \in \{1, \dots, r\}$. Since the component associated to I_0 is adjacent to all the other components, we have

$$D_{\mathcal{S}}^* \simeq \mathcal{M}_{0,n_0+r} \times \mathcal{M}_{0,n_1+1} \times \cdots \times \mathcal{M}_{0,n_r+1}.$$

For each $j = 1, \dots, r$, we define a vector $\underline{k}^{(j)} \in \mathbb{Z}^{n_j+1}$ by setting the n_j first entries to be $\{k_i, i \in I_j\}$ (the ordering is unimportant), and the last entry $k_{n_j+1}^{(j)} := -2d - \sum_{i \in I_j} k_i = \sum_{i \notin I_j} k_i$. By definition, we have $k_{n_j+1}^{(j)} > -d$. Thus $\Omega^d \mathcal{M}_{0,n_j+1}(\underline{k}^{(j)})$ is a subspace of $\mathcal{H}_{0,n_j+1}^{(d)}$. We can then consider the closure of

$\mathbb{P}\Omega^d \mathcal{M}_{0,n_j+1}(\underline{k}^{(j)})$ in $\mathbb{P}\mathcal{H}_{0,n_j+1}^{(d)}$, which is isomorphic to $\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$. Define

$$\mathcal{M}_{\mathcal{S}} := \overline{\mathcal{M}}_{0,n_0+r} \times \prod_{j=1}^r \widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)}).$$

For $j = 1, \dots, r$, let $p_{\mathcal{S},j} : \mathcal{M}_{\mathcal{S}} \rightarrow \widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$ be the canonical projection. Let \mathcal{L}_j denote the tautological line bundle on $\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$. In what follows, for simplicity, we will denote by $\mathcal{O}(-1)$ the tautological line bundle on $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$.

Proposition 8.1. *Let $E_{\mathcal{S}}$ denote the vector bundle $p_{\mathcal{S},1}^* \widehat{\mathcal{L}}_1 \oplus \dots \oplus p_{\mathcal{S},r}^* \widehat{\mathcal{L}}_r$ over $\mathcal{M}_{\mathcal{S}}$. Then there is a birational surjective morphism $i_{\mathcal{S}} : \mathbb{P}E_{\mathcal{S}} \rightarrow \widehat{D}_{\mathcal{S}}$ such that $i_{\mathcal{S}}^* \mathcal{O}(-1) \sim \mathcal{O}(-1)_{\mathbb{P}E_{\mathcal{S}}}$.*

Proof. Locally on $\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$, a trivializing section of $\widehat{\mathcal{L}}_j$ is given by a family of pairs (C^j, q_j) , where C^j is a stable curve in $\overline{\mathcal{M}}_{0,n_j+1}$ and q_j a non-zero d -differential on C^j . Therefore, locally an element of $E_{\mathcal{S}}$ can be identified with a tuple $((C^0, 0), (C^1, q_1), \dots, (C^r, q_r), z)$, where

- $(C^0, 0)$ is a stable curve parametrized by $\overline{\mathcal{M}}_{0,n_0+r}$ equipped with the zero d -differential,
- (C^j, q_j) is a stable curve C^j parametrized by $\overline{\mathcal{M}}_{0,n_j+1}$ equipped with a d -differential $q_j \neq 0$ such that $(C^j, q_j) \in \Omega^d \overline{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$ for $j = 1, \dots, r$,
- $z = (z_1, \dots, z_r) \in \mathbb{C}^r$.

Such a tuple defines an element of $\mathcal{H}_{0,n}^{(d)}$ as follows: let C be the stable curve represented by a point in S , which is obtained by identifying the last marked point on C^j with a marked point on C^0 to form a node for $j = 1, \dots, r$. We consider the C^j 's as subcurves of C . Then the family $\{z_1 q_1, \dots, z_r q_r\}$ gives a d -differential q on C (q vanishes identically on C^0). The pair (C, q) is clearly an element of $\mathcal{H}_{0,n}^{(d)}$.

Thus, by taking the projectivizations, we get a map $i_{\mathcal{S}} : \mathbb{P}E_{\mathcal{S}} \rightarrow \mathbb{P}\mathcal{H}_{0,n}^{(d)}$.

Let $\mathbb{P}E_{\mathcal{S}}^*$ be the restriction of $\mathbb{P}E_{\mathcal{S}}$ to $D_{\mathcal{S}}^* = \mathcal{M}_{0,n_0+r} \times \mathcal{M}_{0,n_1+1} \times \dots \times \mathcal{M}_{0,n_r+1} \subset \mathcal{M}_{\mathcal{S}}$. If C^j is a curve parametrized by a point in \mathcal{M}_{0,n_j+1} , then up to a multiplication by \mathbb{C}^* , there is a unique d -differential q_j on C^j such that $(C^j, q_j) \in \Omega^d \mathcal{M}_{0,n_j+1}(\underline{k}^{(j)})$. Thus the restriction of $i_{\mathcal{S}}$ to $\mathbb{P}E_{\mathcal{S}}^*$ is an isomorphism between $\mathbb{P}E_{\mathcal{S}}^*$ and $\widehat{\mathcal{P}}^{-1}(D_{\mathcal{S}}^*) \subset \widehat{\mathcal{M}}_{0,n}(\underline{k}) \subset \mathbb{P}\mathcal{H}_{0,n}^{(d)}$ (see §5.1). It follows that $i_{\mathcal{S}}(\mathbb{P}E_{\mathcal{S}}) = \widehat{\mathcal{P}}^{-1}(D_{\mathcal{S}}^*) = \widehat{D}_{\mathcal{S}}$, and $i_{\mathcal{S}} : \mathbb{P}E_{\mathcal{S}} \rightarrow \widehat{D}_{\mathcal{S}}$ is birational. It is also clear from the definition that $i_{\mathcal{S}}^* \mathcal{O}(-1) \sim \mathcal{O}(-1)_{\mathbb{P}E_{\mathcal{S}}}$. \square

Remark 8.2. *The map $i_{\mathcal{S}}$ is not one-to-one in general. For instance, let q_1 and q'_1 be two d -differentials on the same curve C^1 such that both (C^1, q_1) and (C^1, q'_1) are elements of $\Omega^d \overline{\mathcal{M}}_{0,n_1+1}(\underline{k}^{(1)})$, but q_1 and q'_1 are not proportional (such d -differentials exist only if C^1 is reducible). Then the tuples $((C^0, 0), (C^1, q_1), \dots, (C^r, q_r), (0, \dots, 0, 1))$ and $((C^0, 0), (C^1, q'_1), \dots, (C^r, q_r), (0, \dots, 0, 1))$ give the same element of $\mathcal{H}_{0,n}^{(d)}$, that is a d -differential on C which is equal to q_r on C^r and vanishes identically on the other subcurves.*

Definition 8.3. *For $\mathcal{S} = \{I_0, \dots, I_r\} \in \widehat{\mathcal{P}}_1(\mu)$, we define the multiplicity of $\widehat{D}_{\mathcal{S}}$ to be*

$$\mathbf{m}(\widehat{D}_{\mathcal{S}}) := d^r \cdot \prod_{j=1}^r \left(\sum_{i \in I_j} \mu_i - 1 \right).$$

Let T_S be the dual graph of the curves parametrized by D_S^* . The vertices of T_S are denoted by v_0, \dots, v_r , where v_j corresponds to the irreducible component that contains the marked points with indices in I_j . By assumption, for each $j \in \{1, \dots, r\}$, there is an edge from v_0 to v_j , which will be denoted by $e_{0,j}$. In §3.2, we have defined the weight of $e_{0,j}$ to be $\mu(e_{0,j}) = \sum_{i \in I_j} \mu_i - 1$. Thus, we have

$$\mathbf{m}(\hat{D}_S) = d^r \cdot \prod_{j=1}^r \mu(e_{0,j}).$$

8.2. Tautological line bundle on boundary divisors in $\hat{\mathcal{P}}_0(\mu)$. Consider now a divisor \hat{D}_S in $\widehat{\mathcal{M}}_{0,n}(k)$ with $\mathcal{S} = \{I_0, I_1\} \in \hat{\mathcal{P}}_0(\mu)$. Recall that \hat{D}_S is the proper transform of the boundary divisor D_S in $\widehat{\mathcal{M}}_{0,n}$ associated to \mathcal{S} . Since D_S is not contained in the support of \mathcal{I} , \hat{D}_S is actually the blow-up of D_S along the ideal sheaf $\mathcal{I}|_{D_S}$ (see [28, Chap.II, Cor.7.15]). Note also that the local generating sets of \mathcal{I} do not involve the defining function of D_S .

8.2.1. Boundary of \hat{D}_S . By definition, D_S is the closure of the stratum $D_S^* \simeq \mathcal{M}_{0,n_0+1} \times \mathcal{M}_{0,n_1+1}$ in $\widehat{\mathcal{M}}_{0,n}$, where $n_j = |I_j|$, $j = 0, 1$. We consider D_S^* as an open dense subset of \hat{D}_S . Denote by $\partial\hat{D}_S$ the complement of D_S^* in \hat{D}_S . The following proposition follows from Propostions 4.1 and 4.4.

Proposition 8.4. *Each irreducible component of $\partial\hat{D}_S$ is a divisor in \hat{D}_S , which corresponds to the intersection $\hat{D}_S \cap \hat{D}_{S'}$, where S' is an element of $\hat{\mathcal{P}}(\mu) \setminus \{\mathcal{S}\}$ such that $D_{S'}$ intersects D_S . Moreover, let $D_{S',S}^*$ be the stratum of $\widehat{\mathcal{M}}_{0,n}$ such that $D_{S'} \cap D_S = \overline{D_{S',S}^*}$. Then we have*

$$\hat{D}_{S'} \cap \hat{D}_S = \overline{\hat{p}^{-1}(D_{S',S}^*)}.$$

Definition 8.5. *Let $\hat{\mathcal{P}}(\mu, \mathcal{S})$ be the set of partitions S' in $\hat{\mathcal{P}}(\mu) \setminus \{\mathcal{S}\}$ such that $D_{S'}$ intersects D_S . For all $S' \in \hat{\mathcal{P}}(\mu, \mathcal{S})$, we denote the intersection $D_{S'} \cap D_S$ by $D_{S',S}$ and the intersection $\hat{D}_{S'} \cap \hat{D}_S$ by $\hat{D}_{S',S}$. The stratum of $\widehat{\mathcal{M}}_{0,n}$ whose closure is equal to $D_{S',S}$ will be denoted by $D_{S',S}^*$.*

8.2.2. Meromorphic section of $\mathcal{O}(-1)$ over \hat{D}_S . Let q be a meromorphic d -differential on $\mathbb{P}_{\mathbb{C}}^1$ which has a pole of order d at $x_0 \in \mathbb{P}_{\mathbb{C}}^1$. The residue of q at x_0 is defined as follows: let $(\hat{C}, \hat{\omega}, \tau)$ be the canonical cyclic cover of $(\mathbb{P}_{\mathbb{C}}^1, q)$. Let \hat{x}_0 be a point in \hat{C} which is mapped to x_0 . The preimage of x_0 in \hat{C} consists of $\{\tau^k(\hat{x}_0), k = 0, \dots, d-1\}$. The Abelian differential $\hat{\omega}$ has a simple pole at \hat{x}_0 . Let c be the residue of $\hat{\omega}$ at \hat{x}_0 . Then the residue of $\hat{\omega}$ at any other point in the preimage of x_0 is equal to $\lambda \cdot c$, where λ is a d -th root of unity. Thus c^d does not depend on the choice of \hat{x}_0 , we will call it the residue of $q_{\mathbf{x}}$ at the node of $C_{\mathbf{x}}$. Concretely, if $z_0 = 0$, and $q = \frac{f(z)(dz)^d}{z^d}$, where $f(z)$ is a holomorphic function in a neighborhood of 0 such that $f(0) \neq 0$, then the residue of q at 0 is simply $f(0)$.

We now construct a trivializing section of $\mathcal{O}(-1)|_{D_S^*}$: for any $\mathbf{x} \in D_S^*$, let $\mathcal{U}_{\mathbf{x}} \subset \widehat{\mathcal{M}}_{0,n}$ be a neighborhood of \mathbf{x} as in §3. Let Φ_0, Φ_1 be the meromorphic sections of the Kawamata line bundle defined in (14). Since the unique node of $C_{\mathbf{x}}$ has weight zero, up to a non-vanishing holomorphic function on $\mathcal{U}_{\mathbf{x}}$, we have $\Phi_0 = \Phi_1$. The curve $C_{\mathbf{x}}$ has two components $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^1$. The restrictions of Φ_0 to $C_{\mathbf{x}}^0$ and the restriction of Φ_1 to $C_{\mathbf{x}}^1$ give a well defined d -differential on $C_{\mathbf{x}}$. This d -differential is actually an element of the fiber of $\mathcal{O}(-1)$ over \mathbf{x} (here we identify $\mathcal{U}_{\mathbf{x}}$ with an open subset of $\widehat{\mathcal{M}}_{0,n}(k)$). Since Φ_0 and Φ_1 have a pole of order d at the node of $C_{\mathbf{x}}$, their residue at this node is non-zero. Thus by multiplying Φ_0 and Φ_1 by a regular function non-vanishing on D_S we can assume that the residue

of Φ_0 and Φ_1 at the node is one. This normalizing condition allows us to patch the local sections (Φ_0, Φ_1) together to obtain a trivializing section σ of $\mathcal{O}(-1)$ over $D_{\mathcal{S}}^*$. The section σ extends naturally to a meromorphic section of $\mathcal{O}(-1)$ over $\hat{D}_{\mathcal{S}}$. We will compute the vanishing order of σ along each of the irreducible components of $\partial\hat{D}_{\mathcal{S}}$.

Let $\hat{\mathbf{x}}$ be a point in $\partial\hat{D}_{\mathcal{S}}$, and \mathbf{x} its image in $\overline{\partial\mathcal{M}_{0,n}}$. We can suppose that $\hat{\mathbf{x}}$ is a generic point in $\hat{D}_{\mathcal{S}'\cdot\mathcal{S}}$, for some $\mathcal{S}' = \{I'_0, \dots, I'_r\} \in \hat{\mathcal{P}}(\mu, \mathcal{S})$. This implies that $\mathbf{x} \in D_{\mathcal{S}'\cdot\mathcal{S}}^*$.

Let us describe a neighborhood of $\hat{\mathbf{x}}$ in $\hat{D}_{\mathcal{S}}$. Let \mathbf{x}' be a point in $\mathcal{U}_{\mathbf{x}} \cap D_{\mathcal{S}'}^*$. By definition, the curve $C_{\mathbf{x}'}$ is obtained from $C_{\mathbf{x}}$ by smoothening the node associated to \mathcal{S} . Denote the irreducible components of $C_{\mathbf{x}'}$ by $C_{\mathbf{x}'}^0, \dots, C_{\mathbf{x}'}^r$, where $C_{\mathbf{x}'}^j$ contains the marked points with index in I'_j . By convention, for each $j \in \{1, \dots, r\}$, there is a node between $C_{\mathbf{x}'}^0$ and $C_{\mathbf{x}'}^j$. Let t_j be the coordinate function on $\mathcal{U}_{\mathbf{x}}$ associated to this node. This means that $D_{\mathcal{S}'} \cap \mathcal{U}_{\mathbf{x}}$ is defined by the equations $\{t_j = 0, j = 1, \dots, r\}$.

Topologically, $C_{\mathbf{x}}$ is obtained from $C_{\mathbf{x}'}$ by pinching a simple closed curve on an irreducible component $C_{\mathbf{x}'}^i$ to a node. We denote the irreducible components of $C_{\mathbf{x}}$ as follows: for $j \in \{0, \dots, r\}$ and $j \neq i$, $C_{\mathbf{x}}^j$ is the component of $C_{\mathbf{x}}$ which corresponds to $C_{\mathbf{x}'}^j$. The component $C_{\mathbf{x}}^i$ corresponds to the union of two components of $C_{\mathbf{x}}$. We denote one of the two components by $C_{\mathbf{x}}^i$ and the other one by $C_{\mathbf{x}}^{r+1}$. Let t_{r+1} be the coordinate function associated with the node between $C_{\mathbf{x}}^i$ and $C_{\mathbf{x}}^{r+1}$. By construction, $\mathcal{U}_{\mathbf{x}} \cap D_{\mathcal{S}}$ is defined by the equation $t_{r+1} = 0$.

For $i = 1, \dots, r$, set $m_j := d(\sum_{i \in I'_j} \mu_i - 1) \in \mathbb{Z}_{\geq 0}$. By assumption, we have

- $t^{\beta_0} = \prod_{j=1}^r t_j^{m_j}$,
- $t^{\beta_k} = \frac{\prod_{j=1}^r t_j^{m_j}}{t_k^{m_k}}$ for $k \in \{1, \dots, r\}$, and
- $t^{\beta_{r+1}} = t^{\beta_i}$.

Since $C_{\mathbf{x}}$ has r principal subcurves, $\widehat{\mathcal{U}}_{\mathbf{x}}$ is covered by the open subsets $\widehat{\mathcal{U}}_{\mathbf{x}}^k$, $k = 1, \dots, r$, where $\widehat{\mathcal{U}}_{\mathbf{x}}^k$ is defined by the equations

$$(32) \quad t^{\beta_j} = v_j t^{\beta_k}, \quad j = 1, \dots, r.$$

Remark that when $t_1 \cdots t_r \neq 0$, (32) is equivalent to

$$(33) \quad t_k^{m_k} = v_j t_j^{m_j}, \quad j = 1, \dots, r.$$

Thus we can use (33) as defining equations of $\widehat{\mathcal{U}}_{\mathbf{x}}^k$.

Note that by definition, $\hat{D}_{\mathcal{S}'}$ is defined by the equations $t_1 = \dots = t_r = 0$ in $\widehat{\mathcal{U}}_{\mathbf{x}}$. In the open $\{(t, u, [v_1 : \dots : v_r]) \in \widehat{\mathcal{U}}_{\mathbf{x}}, v_j \neq 0, j = 1, \dots, r\}$, $\hat{D}_{\mathcal{S}'}$ is defined by any of the equations $t_j = 0$.

Lemma 8.6. *If $C_{\mathbf{x}}^i$ is a principal component of $C_{\mathbf{x}}$, then the order of σ along $\hat{D}_{\mathcal{S}'\cdot\mathcal{S}}$ is zero.*

Proof. We first remark that it suffices to consider the case $i = r$. Indeed, if $\mathcal{S}' \in \hat{\mathcal{P}}_1(\mu)$, then $i \in \{1, \dots, r\}$. In this case, up to a renumbering, we can suppose that $i = r$. In the case, $\mathcal{S}' = \{I'_0, I'_1\} \in \hat{\mathcal{P}}_0(\mu)$, then both $C_{\mathbf{x}'}^0$ and $C_{\mathbf{x}'}^1$ are principal. Hence up to a renumbering of $\{I'_0, I'_1\}$, we can always assume that $i = 1$.

Let $\Phi_0, \Phi_1, \dots, \Phi_{r+1}$ be as in §2. Since $C_{\mathbf{x}}^r$ and $C_{\mathbf{x}}^{r+1}$ are separated by a node of weight zero, we have $\Phi_r = \Phi_{r+1}$. By assumption, the node between $C_{\mathbf{x}}^r$ and $C_{\mathbf{x}}^{r+1}$ is associated to the partition $\mathcal{S} = \{I_0, I_1\}$ of $\{1, \dots, n\}$. Without loss of generality, we can assume that the component $C_{\mathbf{x}}^{r+1}$ contains the i -th marked

point for all $i \in I_1$. Smoothing all the nodes between C_x^0 and the components C_x^j , $j = 1, \dots, r$, we obtain a point \mathbf{y} in the stratum D_S^* . The curve C_y parametrized by \mathbf{y} has two components C_y^0, C_y^1 , where C_y^1 is mapped onto C_x^{r+1} by the degenerating map $\varphi_y : C_y \rightarrow C_x$.

We now observe that the restriction of the section σ to C_y^1 is a d -differential having a zero of order k_i at the i -th marked point for all $i \in I_1$, and a pole of order d at the node associated to \mathcal{S} . Thus, by multiplying Φ_{r+1} by a non-vanishing regular function on $\mathcal{U}_x \cap D_S$, we can write

$$\sigma|_{\widehat{\mathcal{U}}_x^r \cap D_S^*} = \Phi_{r+1}|_{\mathcal{U}_x^r \cap D_S^*}.$$

as sections of $\mathcal{O}(-1)$. By Theorem 5.2, we know that $\Phi_r = \Phi_{r+1}$ is a trivializing section of $\mathcal{O}(-1)|_{\widehat{\mathcal{U}}_x^r}$. Thus, σ extends to a trivializing section of $\mathcal{O}(-1)$ on $\widehat{\mathcal{U}}_x^r \cap \widehat{D}_S$, and the order of σ along $\widehat{D}_{S', \mathcal{S}}$ is zero. \square

Lemma 8.7. *Assume that $\mathcal{S}' \in \widehat{\mathcal{P}}(\mu, \mathcal{S}) \cap \widehat{\mathcal{P}}_1(\mu)$, and $i = 0$. Then the vanishing order of σ along $\widehat{D}_{S', \mathcal{S}}$ is $-\mathbf{m}(\widehat{D}_{S'})$, where $\mathbf{m}(\widehat{D}_{S'})$ is the multiplicity of $\widehat{D}_{S'}$ (see Definition 8.3).*

Proof. Consider a point $\mathbf{y} \in \mathcal{U}_x \cap D_S^*$. The curve C_y parametrized by \mathbf{y} is obtained from C_x by smoothing the nodes associated to the parameters t_1, \dots, t_r . By definition, C_y has two components C_y^0 and C_y^1 separated by the node associated to \mathcal{S} . The degenerating map $\varphi_y : C_y \rightarrow C_x$ maps C_y^0 and C_y^1 onto two subcurves of C_x that are separated by the node between C_x^0 and C_x^{r+1} . Without loss of generality, we can assume that $C_x^0 \subset \varphi_y(C_y^0)$.

Let $\Phi_0, \dots, \Phi_{r+1}$ be the sections of the Kawamata line bundle over $p^{-1}(\mathcal{U}_x)$ defined in §2.3. Since $C_x^0 \subset \varphi_y(C_y^0)$, the restriction of Φ_0 to C_y^0 is a non-zero d -differential with a zero of order k_i at the i -th marked point, for all $i \in I_0$, and a pole of order d at the node of C_y^0 . Multiplying Φ_0 by a regular non-vanishing function on $\mathcal{U}_x \cap D_S$, we can write

$$\sigma = \Phi_0$$

as sections of the Kawamata line bundle over $p^{-1}(\mathcal{U}_x \cap D_S^*)$. This implies that Φ_0 is an extension of σ to $\widehat{\mathcal{U}}_x \cap \widehat{D}_S$.

Consider Φ_0 as a meromorphic section of $\mathcal{O}(-1)$ over $\widehat{\mathcal{U}}_x \cap \widehat{D}_S$. We wish to calculate the vanishing order of this section along the divisor $\widehat{D}_{S', \mathcal{S}}$ of \widehat{D}_S . It is enough to restrict our calculation to the open affine $\widehat{\mathcal{U}}_x^r$. By Theorem 1.1 we know that a trivializing section of $\mathcal{O}(-1)$ on $\widehat{\mathcal{U}}_x^r$ is Φ_r , and we have

$$\frac{\Phi_0}{\Phi_r} = \frac{t^{\beta_r}}{t^{\beta_0}} = t_r^{-m_r}.$$

Thus, the vanishing order of Φ_0 along $\widehat{D}_{S', \mathcal{S}}$ is equal to the vanishing order of $t_r^{-m_r}$ along the divisor $\widehat{D}_{S'}$.

We claim that the order of $t_r^{-m_r}$ along $\widehat{D}_{S'}$ is equal to $\mathbf{m}(\widehat{D}_{S'})$. Let A be localization of the affine coordinate ring of $\widehat{\mathcal{U}}_x^r$ at the ideal generated by $\{t_1, \dots, t_r\}$, that is the defining ideal of $\widehat{D}_{S'}$. By definition (see [25, §1.2]), the order of $t_r^{-m_r}$ along $\widehat{D}_{S'}$ is the length of $A/(t_r^{-m_r})$ as an A -module. It is not difficult to see that

$$A/(t_r^{-m_r}) \simeq \mathbb{C}(v_1, \dots, v_{r-1})[t_1, \dots, t_r]/(t_1^{m_1}, \dots, t_r^{m_r})$$

Since the residue field of A is $\mathbb{C}(v_1, \dots, v_{r-1})$, it follows

$$\ell_A(A/(t_r^{-m_r})) = \dim_{\mathbb{C}(v_1, \dots, v_{r-1})} A/(t_r^{-m_r}) = m_1 \cdots m_r = \mathbf{m}(\widehat{D}_{S'}).$$

(see [25, Ex A.1.1]). Thus the order of σ along $\hat{D}_{S',S} = -\mathbf{m}(\hat{D}_{S'})$. This concludes the proof of the lemma. \square

Combining Lemma 8.6 and Lemma 8.7, we get

Theorem 8.8. *Let $\hat{\mathcal{P}}^*(\mu, \mathcal{S})$ be the set of $S' = \{I'_0, \dots, I'_r\} \in \hat{\mathcal{P}}(\mu, \mathcal{S})$, such that $S' \in \hat{\mathcal{P}}_1(\mu)$ and the curves parametrized by $D_{S',S}^*$ are obtained from the ones parametrized by $D_{S'}^*$, by pinching a loop in the component associated to I'_0 . Then the restriction of the tautological line bundle $\mathcal{O}(-1)$ to \hat{D}_S is represented by the Weil divisor*

$$W_S := - \sum_{S' \in \hat{\mathcal{P}}^*(\mu, \mathcal{S})} \mathbf{m}(\hat{D}_{S'}) \cdot \hat{D}_{S',S}.$$

9. RECURSIVE FORMULA

9.1. Self-intersection of the first Chern class of the tautological line bundle on divisors. For each $k \in \mathbb{Z}_{\geq 3}$, let

$$L_k = \{v = (v_1, \dots, v_k) \in \mathbb{Q}^k, \sum_{i=1}^k v_i = 2, \text{ and } v_i < 1, \text{ for all } i = 1, \dots, k\}.$$

Recall that $\mathcal{J}_k : L_k \rightarrow \mathbb{Z}$ is the map defined as follows: given $v = (v_1, \dots, v_k) \in L_k$, there is a smallest natural number e such that $ev_i \in \mathbb{Z}$, for all $i = 1, \dots, k$. Set $\ell_i = -ev_i$. We then have $\ell_1 + \dots + \ell_k = -2e$, $\gcd(e, \ell_1, \dots, \ell_k) = 1$, and $\ell_i > -e$, $i = 1, \dots, k$. Thus we can embed $\widehat{\mathcal{M}}_{0,k}(\underline{\ell})$ into $\mathbb{P}\mathcal{H}_{0,k}^{(e)}$, where $\underline{\ell} = (\ell_1, \dots, \ell_k)$. We then define

$$\mathcal{J}_k(v) := c_1^{k-3}(\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,k}^{(e)}}) \cap [\widehat{\mathcal{M}}_{0,k}(\underline{\ell})],$$

where $[\widehat{\mathcal{M}}_{0,k}(\underline{\ell})]$ is the class of $\widehat{\mathcal{M}}_{0,k}(\underline{\ell})$ in $A_*(\mathbb{P}\mathcal{H}_{0,k}^{(e)})$. Note that by definition, $\mathcal{J}_3(v) = 1$, for all $v \in L_3$.

Lemma 9.1. *If there exists $i \in \{1, \dots, k\}$ such that $v_i \in \mathbb{Z}$ then $\mathcal{J}_k(v) = 0$.*

Proof. Let $\hat{\mathcal{L}}_v$ denote the restriction of $\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0,k}^{(e)}}$ to $\widehat{\mathcal{M}}_{0,k}(\underline{\ell})$. By Theorem 7.3, $(\frac{1}{2\pi}\Theta_v)^{k-3}$ (where Θ_v is the curvature form of the Hodge norm on $\hat{\mathcal{L}}_v|_{\widehat{\mathcal{M}}_{0,k}(\underline{\ell})}$) is a representative in the sense of currents of $c_1^{k-3}(\hat{\mathcal{L}}_v)$. The condition $v_i \in \mathbb{Z}$ is equivalent to $\ell_i \in e\mathbb{Z}$. Since $\{\ell_1, \dots, \ell_k\} \cap e\mathbb{Z} \neq \emptyset$, we have $\ker \mathbf{p} \cap V_{\underline{\ell}} \neq \{0\}$, where $V_{\underline{\ell}}$ and \mathbf{p} are defined in §6. In this case, a direct computation shows that Θ_v^{k-3} vanishes identically on $\mathbb{P}V_{\underline{\ell}}$. Thus $c_1^{k-3}(\hat{\mathcal{L}}_v) = 0 \in H^{2(k-3)}(\widehat{\mathcal{M}}_{0,k}(\underline{\ell}))$. \square

Consider a divisor \hat{D}_S in $\partial\widehat{\mathcal{M}}_{0,n}(\underline{k})$ which is associated to a partition $\mathcal{S} = \{I_0, I_1, \dots, I_r\} \in \hat{\mathcal{P}}_1(\mu)$. For $j = 1, \dots, r$, we set $(v_1^{(j)}, \dots, v_{n_j}^{(j)}) = (\mu_i, i \in I_j)$ (the ordering of $\{\mu_i, i \in I_j\}$ is unimportant), and define

$$v_{n_j+1}^{(j)} = 2 - \sum_{i \in I_j} \mu_i.$$

Let $v^{(j)} := (v_1^{(j)}, \dots, v_{n_j+1}^{(j)})$. Let e_j be the smallest natural number such that $\ell_k^{(j)} := -e_j v_k^{(j)} \in \mathbb{Z}$, and $\underline{\ell}^{(j)} = (\ell_1^{(j)}, \dots, \ell_{n_j+1}^{(j)})$.

Recall that $N = n - 3 = \dim \widehat{\mathcal{M}}_{0,n}(\underline{k})$, and we must have $n_0 + r \geq 3$.

Proposition 9.2. *If $n_0 + r > 3$ then*

$$c_1^{N-1}(\mathcal{O}(-1)) \cap [\hat{D}_S] = 0.$$

Assume that $n_0 + r = 3$, then we must have $r \in \{1, 2\}$, that is $\mathcal{S} \in \mathcal{T}_{1a} \cup \mathcal{T}_{2a}$, and

$$(34) \quad c_1^{N-1}(\mathcal{O}(-1)) \cap [\hat{D}_S] = (-1)^{r+1} \frac{d^{N-r}}{e_1^{n_1-2} \cdots e_r^{n_r-2}} \prod_{j=1}^r \mathcal{J}_{n_j+1}(v^{(j)}).$$

Proof. For $j = 1, \dots, r$, $i = 1, \dots, n_j + 1$, define $k_i^{(j)} = -dv_i^{(j)}$, and $\underline{k}^{(j)} = (k_1^{(j)}, \dots, k_{n_j+1}^{(j)})$. By Theorem 1.1, we know that $\widehat{\mathcal{M}}_{0, n_j+1}(\underline{k}^{(j)})$ is the closure of the projectivized stratum $\mathbb{P}\Omega^d \mathcal{M}_{0, n_j+1}(\underline{k}^{(j)})$ in $\mathbb{P}\mathcal{H}_{0, n_j+1}^{(d)}$. Let $\hat{\mathcal{L}}_j$ be the tautological line bundle on $\widehat{\mathcal{M}}_{0, n_j+1}(\underline{k}^{(j)})$. By Proposition 8.1, there is a surjective birational morphism

$$i_S : \mathbb{P}E_S \rightarrow \hat{D}_S.$$

where E_S is the rank r vector bundle $\hat{\mathcal{L}}_1 \oplus \cdots \oplus \hat{\mathcal{L}}_r$ over the space $\mathcal{M}_S = \overline{\mathcal{M}}_{0, n_0+r} \times \prod_{j=1}^r \widehat{\mathcal{M}}_{0, n_j+1}(\underline{k}^{(j)})$ (here we abusively denote by $\hat{\mathcal{L}}_j$ the pullback of the tautological line bundle on $\widehat{\mathcal{M}}_{0, n_j+1}(\underline{k}^{(j)})$ to \mathcal{M}_S). Moreover, we have $i_S^* \mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0, n}^{(d)}} \sim \mathcal{O}(-1)_{\mathbb{P}E_S}$. It follows

$$c_1^{N-1}(\mathcal{O}(-1)_{\mathbb{P}\mathcal{H}_{0, n}^{(d)}}) \cap [\hat{D}_S] = c_1^{N-1}(\mathcal{O}(-1)_{\mathbb{P}E_S}) \cap [\mathbb{P}E_S] = (-1)^{N-1} c_1^{N-1}(\mathcal{O}(1)_{\mathbb{P}E_S}) \cap [\mathbb{P}E_S].$$

By definition, we have

$$c_1^{N-1}(\mathcal{O}(1)_{\mathbb{P}E_S}) \cap [\mathbb{P}E_S] = s_{N-r}(E_S) \cap [\mathcal{M}_S]$$

where s_{N-r} is the $(N-r)$ -th Segre class of E_S (see [25, §3.1]). Since E_S is the direct sum of the line bundles $\hat{\mathcal{L}}_1, \dots, \hat{\mathcal{L}}_r$, its total Chern class is therefore

$$c(E_S) = \prod_{j=1}^r (1 + c_1(\hat{\mathcal{L}}_j)).$$

Hence, the total Segre class of E_S is given by

$$s(E_S) = \prod_{j=1}^r (1 + c_1(\hat{\mathcal{L}}_j))^{-1}.$$

It follows that

$$s_{N-r}(E_S) = (-1)^{N-r} \sum_{\alpha_1 + \cdots + \alpha_r = N-r} c_1^{\alpha_1}(\hat{\mathcal{L}}_1) \cdots c_1^{\alpha_r}(\hat{\mathcal{L}}_r).$$

Thus

$$(35) \quad s_{N-r}(E_S) \cap [\mathcal{M}_S] = (-1)^{N-r} \sum_{\alpha_1 + \cdots + \alpha_r = N-r} c_1^{\alpha_1}(\hat{\mathcal{L}}_1) \cap [\widehat{\mathcal{M}}_{0, n_1+1}(\underline{k}^{(1)})] \cdots c_1^{\alpha_r}(\hat{\mathcal{L}}_r) \cap [\widehat{\mathcal{M}}_{0, n_r+1}(\underline{k}^{(r)})].$$

We now remark that

$$\dim \widehat{\mathcal{M}}_{0, n_1+1}(\underline{k}^{(1)}) + \cdots + \dim \widehat{\mathcal{M}}_{0, n_r+1}(\underline{k}^{(r)}) = \sum_{j=1}^r n_j - 2r = n - n_0 - 2r,$$

while $N-r = n-3-r$. Therefore, if $n_0 + r > 3$ then $N-r > n - n_0 - 2r$, hence $s_{N-r}(E_S) \cap [\mathcal{M}_S] = 0$.

Since we must have $n_0 + r \geq 3$, it remains to consider the case $n_0 + r = 3$. By assumption, $r \geq 1$, hence we have $r \in \{1, 2, 3\}$. We first claim that $r \neq 3$. Indeed, if $r = 3$ then $n_0 = 0$, which means that $I_0 = \emptyset$. We have $\sum_{i \in I_j} \mu_i > 1$, for all $j = 1, 2, 3$. But this implies that $\mu_1 + \dots + \mu_n > 3$, which is a contradiction.

Assume now that $r = 2$. Then (35) gives

$$s_{N-2}(E_S) \cap [\mathcal{M}_S] = (-1)^{N-2} c_1^{n_1-2}(\hat{\mathcal{L}}_1) \cap [\widehat{\mathcal{M}}_{0,n_1+1}(\underline{k}^{(1)})] \cdot c_1^{n_2-2}(\hat{\mathcal{L}}_2) \cap [\widehat{\mathcal{M}}_{0,n_2+1}(\underline{k}^{(2)})].$$

Let e_j be the smallest natural number such that $\ell_i^{(j)} := -e_j \mu_i^{(j)} \in \mathbb{Z}$. Note that we have $\underline{k}^{(j)} = (d/e_j) \cdot \underline{\ell}^{(j)}$. The compactification of $\mathbb{P}\Omega^{e_j} \mathcal{M}_{0,n_j+1}(\underline{\ell}^{(j)})$ in $\mathbb{P}\mathcal{H}_{0,n_j+1}^{(e_j)}$ is isomorphic to $\widehat{\mathcal{M}}_{0,n_j+1}(\underline{\ell}^{(j)})$. Let $\hat{\mathcal{L}}_{\mu^{(j)}}$ be the tautological line bundle over $\widehat{\mathcal{M}}_{0,n_j+1}(\underline{\ell}^{(j)})$. We have a natural surjective birational morphism $f_j : \widehat{\mathcal{M}}_{0,n_j+1}(\underline{\ell}^{(j)}) \rightarrow \widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})$ which is induced by the map $(C, q) \mapsto (C, q^{\otimes (d/e_j)})$, for all $(C, q) \in \Omega^{e_j} \mathcal{M}_{0,n_j+1}(\underline{\ell}^{(j)})$. By construction, we have

$$f_j^* \hat{\mathcal{L}}_j \sim \hat{\mathcal{L}}_{\mu^{(j)}}^{\otimes \frac{d}{e_j}}.$$

Thus

$$c_1^{n_j-2}(\hat{\mathcal{L}}_j) \cap [\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})] = \left(\frac{d}{e_j}\right)^{n_j-2} \cdot c_1^{n_j-2}(\hat{\mathcal{L}}_{\mu^{(j)}}) \cap [\widehat{\mathcal{M}}_{0,n_j+1}(\underline{\ell}^{(j)})] = \left(\frac{d}{e_j}\right)^{n_j-2} \mathcal{J}_{n_j+1}(\mu^{(j)}).$$

Hence (34) is proved for this case. The case $n_0 = 2, r = 1$ follows from the same arguments. \square

We now turn to the case \hat{D}_S with $S = \{I_0, I_1\} \in \hat{\mathcal{P}}_0(\mu)$. By Theorem 8.8, the restriction of $\mathcal{O}(-1)$ to \hat{D}_S is represented by a Weil divisor which is supported on the union of divisors $\hat{D}_{S',S}$, with $S' \in \hat{\mathcal{P}}^*(\mu, S)$. Given $S' = \{I'_0, I'_1, \dots, I'_r\} \in \hat{\mathcal{P}}^*(\mu, S)$, we define $S' * S := \{I'_0 \cap I_0, I'_0 \cap I_1, I'_1, \dots, I'_r\}$. Remark that $S' * S$ is the partition associated to the stratum $D_{S',S}^*$ in $\overline{\mathcal{M}}_{0,n}$ (see Definition 8.5). Note however that $D_{S',S}^*$ is not uniquely determined by $S' * S$.

Proposition 9.3. *Let $S' = \{I'_0, \dots, I'_r\} \in \hat{\mathcal{P}}^*(\mu, S)$. If $n_0 + r > 4$ then*

$$c_1^{N-2}(\mathcal{O}(-1)) \cap [\hat{D}_{S',S}] = 0.$$

If $n_0 + r = 4$, then we must have $r \in \{1, 2\}$, and

$$(36) \quad c_1^{N-2}(\mathcal{O}(-1)) \cap [\hat{D}_{S',S}] = (-1)^{r+1} \frac{d^{N-r-1}}{e_1^{n_1-2} \dots e_r^{n_r-2}} \prod_{j=1}^r \mathcal{J}_{n_j+1}(\mu^{(j)}).$$

*In particular, $c_1^{N-2}(\mathcal{O}(-1)) \cap [\hat{D}_{S',S}] = 0$ unless $S' * S \in \mathcal{T}_{1b} \cup \mathcal{T}_{2b}$.*

Proof. Let $I'_{00} = I'_0 \cap I_0$ and $I'_{01} = I'_0 \cap I_1$. The dual graph $T_{S',S}$ associated to the stratum $D_{S',S}^*$ can be constructed as follows: the vertices of $T_{S',S}$ are labeled by $\{I'_{00}, I'_{01}, I'_1, \dots, I'_r\}$. There is an edge between I'_{00} and I'_{01} . For all $j \in \{1, \dots, r\}$, there is an edge between I'_j and I'_{00} if and only if $I'_j \subset I_0$, otherwise there is an edge between I'_j and I_{01} . We set r_0 (resp. r_1) to be the number of vertices among $\{I'_1, \dots, I'_r\}$ that are connected to I'_{00} (resp. to I'_{01}) by an edge. Note that we have $r = r_0 + r_1$.

Let $n_{00} = |I'_{00}|$, $n_{01} = |I'_{01}|$, and $n_j = |I'_j|$, $j = 1, \dots, r$. We define

$$\mathcal{M}_{S',S} = \overline{\mathcal{M}}_{0,n_{00}+r_0+1} \times \overline{\mathcal{M}}_{0,n_{01}+r_1+1} \times \prod_{j=1}^r \widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)}).$$

Note that $\mathcal{M}_{S',S}$ is a divisor in $\mathcal{M}_{S'}$. Let $E_{S',S}$ denote the restriction of the vector bundle $E_{S'}$ to $\mathcal{M}_{S',S}$. By the same arguments as in Proposition 8.1, we see that there is a surjective birational morphism $i_{S',S} : \mathbb{P}E_{S',S} \rightarrow \hat{D}_{S',S}$ such that $i_{S',S}^* \mathcal{O}(-1) \sim \mathcal{O}(-1)_{\mathbb{P}E_{S',S}}$. Thus, by the same arguments as in Proposition 9.2, we have

$$\begin{aligned} c_1^{N-2}(\mathcal{O}(-1)) \cap [\hat{D}_{S',S}] &= c_1^{N-2}(\mathcal{O}(-1)_{\mathbb{P}E_{S',S}}) \cap [\mathbb{P}E_{S',S}] \\ &= (-1)^{N-2} c_1^{N-2}(\mathcal{O}(1)_{\mathbb{P}E_{S',S}}) \cap [\mathbb{P}E_{S',S}] \\ &= (-1)^{N-2} s_{N-r-1}(E_{S',S}) \cap [\mathcal{M}_{S',S}] \\ &= (-1)^{r+1} \sum_{\alpha_1 + \dots + \alpha_r = N-r-1} \prod_{j=1}^r c_1^{\alpha_j}(\hat{\mathcal{L}}_j) \cap [\widehat{\mathcal{M}}_{0,n_j+1}(\underline{k}^{(j)})]. \end{aligned}$$

We now observe that

$$\dim \widehat{\mathcal{M}}_{0,n_1+1}(\underline{k}^{(1)}) + \dots + \dim \widehat{\mathcal{M}}_{0,n_r+1}(\underline{k}^{(r)}) = \sum_{j=1}^r n_j - 2r = n - n_0 - 2r,$$

while $N - r - 1 = n - r - 4$. Recall that we must have $n_{00} + r_0 + 1 \geq 3$ and $n_{01} + r_1 + 1 \geq 3$, which implies that $n_0 + r = n_{00} + n_{01} + r_0 + r_1 \geq 4$. If $n_0 + r > 4$ then $N - r - 1 > n - n_0 - 2r$, hence $c_1^{N-2}(\mathcal{O}(-1)) \cap [\hat{D}_{S',S}] = 0$. Thus it suffices to consider the case $n_0 + r = 4$. Since $r \geq 1$, we have $r \in \{1, 2, 3, 4\}$. We claim that $r \in \{1, 2\}$. Indeed, if $r = 4$ then $n_0 = 0$, which means that $I'_0 = \emptyset$. We then have

$$\sum_{i=1}^n \mu_i = \sum_{j=1}^4 \sum_{i \in I'_j} \mu_i > 4$$

which is impossible. If $r = 3$ then $n_0 = 1$, and we can suppose that $n_{00} = 0$ and $n_{01} = 1$. Since we must have $n_{00} + r_0 + 1 = 3$, it follows that $r_0 = 2$. Without loss of generality, we can assume that $I_0 = I'_1 \cup \dots \cup I'_2$. But we would have

$$1 = \sum_{i \in I_0} \mu_i = \sum_{j=1}^2 \sum_{i \in I'_j} \mu_i > 2$$

which is again impossible. Therefore, we must have $r \in \{1, 2\}$. If $r = 2$, then $n_0 = 2$, and the conditions $n_{00} + r_0 + 1 = n_{01} + r_1 + 1 = 3$ implies that $r_0 = r_1 = 1$. If $r = 1$, then $n_0 = 3$. We can assume that $r_0 = 1$ and $r_1 = 0$, which implies that $n_{00} = 1$ and $n_{01} = 2$. The rest of the proof follows the same lines as Proposition 9.2. \square

9.2. Proof of Theorem 1.5.

Proof. Recall that for any subset $I \subset \{1, \dots, n\}$, we define $\mu(I) := \sum_{i \in I} \mu_i$. By Theorem 1.3, we know that $\hat{\mathcal{L}}_\mu \sim \hat{p}^* \bar{\mathcal{L}}_\mu + \mathcal{E}$, where $\hat{\mathcal{L}}_\mu$ is the restriction of the tautological line bundle on $\mathbb{P}\mathcal{H}_{0,n}^{(d)}$ to $\widehat{\mathcal{M}}_{0,n}(\underline{k})$.

Thus

$$c_1^N(\hat{\mathcal{L}}_\mu) \cap [\widehat{\mathcal{M}}_{0,n}(\underline{k})] = c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{p}^* \bar{\mathcal{L}}_\mu + \mathcal{E}].$$

In [34, §8], it was shown that

$$(37) \quad \bar{\mathcal{L}}_\mu \sim \mathcal{D}_\mu \sim \frac{d}{n-2} \sum_{\mathcal{S}=\{I_0, I_1\} \in \mathcal{P}} (|I_0| - 1)(\mu(I_0) - \frac{|I_0|}{n-1}) D_{\mathcal{S}}.$$

Equality (37) induces an explicit expression of $\hat{p}^* \bar{\mathcal{L}}_\mu + \mathcal{E}$ as a combination of the boundary divisors $\{\hat{D}_{\mathcal{S}}, \mathcal{S} \in \hat{\mathcal{P}}(\mu)\}$. Let us write

$$[\hat{p}^* \mathcal{D}_\mu + \mathcal{E}] = \sum_{\mathcal{S} \in \hat{\mathcal{P}}(\mu)} \lambda_{\mathcal{S}} [\hat{D}_{\mathcal{S}}].$$

For $\mathcal{S} \in \hat{\mathcal{P}}_1(\mu)$, Proposition 9.2 implies that $c_1^{N-1}(\mathcal{O}(-1)) \cap \hat{D}_{\mathcal{S}} = 0$ unless \mathcal{S} belongs to \mathcal{T}_{1a} or \mathcal{T}_{2a} . We will calculate $\lambda_{\mathcal{S}}$ for all $\mathcal{S} \in \mathcal{T}_{1a} \cup \mathcal{T}_{2a}$.

- $\mathcal{S} = \{I_0, I_1\} \in \mathcal{T}_{1a}$. Since $\hat{D}_{\mathcal{S}}$ is not contained in $|\mathcal{E}|$ (that is the support of the exceptional divisor), the coefficient of $\hat{D}_{\mathcal{S}}$ is given by the coefficient of $D_{\mathcal{S}}$ in \mathcal{D}_μ , that is

$$\begin{aligned} \lambda_{\mathcal{S}} &= \frac{d}{n-2} (\mu(I_0) - \frac{2}{n-1}) \\ &= \frac{1}{n-2} \left(\frac{d(n-3)}{n-1} - m(\mathcal{S}) \right) \\ &= a_{\mathcal{S}}. \end{aligned}$$

- $\mathcal{S} = \{I_0, I_1, I_2\} \in \mathcal{T}_{2a}$. In this case, we have $\hat{D}_{\mathcal{S}} \subset |\mathcal{E}|$. In fact, $\hat{D}_{\mathcal{S}}$ is contained in the inverse image of two divisors $D_{\mathcal{S}_1}$ and $D_{\mathcal{S}_2}$ in $\partial \widehat{\mathcal{M}}_{0,n}$, where $\mathcal{S}_1 = \{I_0 \cup I_2, I_1\}$ and $\mathcal{S}_2 = \{I_0 \cup I_1, I_2\}$. Recall that $m_j := d(\mu(I_j) - 1)$, $j = 1, 2$. By assumption, we have $|I_0| = 1$. Therefore, $n_1 + n_2 = n - 1$, and the coefficient of $D_{\mathcal{S}_1}$ in \mathcal{D}_μ is

$$b_{\mathcal{S}_1} = \frac{d}{n-2} \cdot n_2 \cdot \left((2 - \mu(I_1)) - \frac{n_2 + 1}{n-1} \right) = \frac{dn_2(n_1 - 1)}{(n-1)(n-2)} - \frac{n_2 m_1}{n-2}.$$

Similarly, the coefficient of $D_{\mathcal{S}_2}$ in \mathcal{D}_μ is

$$b_{\mathcal{S}_2} = \frac{dn_1(n_2 - 1)}{(n-1)(n-2)} - \frac{n_1 m_2}{n-2}.$$

It follows from Lemma 9.4 that the multiplicity of $\hat{D}_{\mathcal{S}}$ in $\hat{p}^* D_{\mathcal{S}_1}$ is m_2 , and in $\hat{p}^*(D_{\mathcal{S}_2})$ is m_1 . Thus the coefficient of $\hat{D}_{\mathcal{S}}$ in $\hat{p}^* \mathcal{D}_\mu$ is equal to

$$m_2 b_{\mathcal{S}_1} + m_1 b_{\mathcal{S}_2} = \frac{d(m_2 n_2 (n_1 - 1) + m_1 n_1 (n_2 - 1))}{(n-1)(n-2)} - \frac{m_1 m_2 (n-1)}{n-2}.$$

By Lemma 9.5, the coefficient of $\hat{D}_{\mathcal{S}}$ in \mathcal{E} is $m_1 m_2$. Therefore,

$$\begin{aligned} \lambda_{\mathcal{S}} &= \frac{d(m_2 n_2 (n_1 - 1) + m_1 n_1 (n_2 - 1))}{(n-1)(n-2)} - \frac{m_1 m_2 (n-1)}{n-2} + m_1 m_2 \\ &= \frac{d(m_2 n_2 (n_1 - 1) + m_1 n_1 (n_2 - 1))}{(n-1)(n-2)} - \frac{m_1 m_2}{n-2} \\ &= a_{\mathcal{S}}. \end{aligned}$$

In conclusion, we have

$$\sum_{S \in \hat{\mathcal{P}}_1(\mu)} \lambda_S \cdot c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_S] = \sum_{S \in \mathcal{T}_{1a} \cup \mathcal{T}_{2a}} a_S \cdot c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_S].$$

Let us now consider a divisor \hat{D}_{S_0} associated to a partition $S_0 = \{I_0, I_1\} \in \hat{\mathcal{P}}_0(\mu)$. Recall that we have $\mu(I_0) = \mu(I_1) = 1$. This implies that the coefficient of D_{S_0} in \mathcal{D}_μ is $\frac{d(|I_0|-1)(|I_1|-1)}{(n-1)(n-2)}$. Since \hat{D}_{S_0} is not contained in the support of \mathcal{E} , we have

$$\lambda_{S_0} = \frac{d(|I_0|-1)(|I_1|-1)}{(n-1)(n-2)}.$$

From Theorem 8.8, we have

$$c_1(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S_0}] \sim \sum_{S_1 \in \hat{\mathcal{P}}^*(\mu, S_0)} -\mathbf{m}(S_1) [\hat{D}_{S_1 \cdot S_0}].$$

Thus

$$(38) \quad c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap \sum_{S_0 \in \hat{\mathcal{P}}_0(\mu)} \lambda_{S_0} [\hat{D}_{S_0}] = c_1^{N-2}(\hat{\mathcal{L}}_\mu) \cap \sum_{S_0 \in \hat{\mathcal{P}}_0(\mu)} \sum_{S_1 \in \hat{\mathcal{P}}^*(\mu, S_0)} -\lambda_{S_0} \cdot \mathbf{m}(S_1) \cdot [\hat{D}_{S_1 \cdot S_0}].$$

By Proposition 9.3, we know that $c_1^{N-2}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S_1 \cdot S_0}] = 0$ unless the partition $S = S' * S$ associated to $\hat{D}_{S_1 \cdot S_0}$ belongs to either \mathcal{T}_{1b} or \mathcal{T}_{2b} .

- If $S = \{I_{00}, I_{01}, I_1\} \in \mathcal{T}_{1b}$, then there is a unique choice for (S_0, S_1) , namely $S_0 = \{I_{00}, I_{01} \cup I_1\}$, and $S_1 = \{I_{00} \cup I_{01}, I_1\}$. Recall that by definition we have $|I_{00}| = 2$ and $|I_{01}| = 1$. Thus we have

$$\lambda_{S_0} \mathbf{m}(S_1) = \frac{d(n-3)}{(n-1)(n-2)} \cdot m(S) = a_S.$$

- For $S = \{I_{01}, I_{02}, I_1, I_2\} \in \mathcal{T}_{2b}$, we have two cases: if $\mu(I_{01}) \neq \mu(I_{02})$, then S_0 and S_1 are uniquely determined by S , namely $S_0 = \{I_{01} \cup I_1, I_{02} \cup I_2\}$ and $S_1 = \{I_{01} \cup I_{02}, I_1, I_2\}$. If $\mu(I_{01}) = \mu(I_{02})$, then the partition S arises from two pairs (S_0, S_1) and (S'_0, S_1) , where S_0 and S_1 are as above, and $S'_0 = \{I_{01} \cup I_2, I_{02} \cup I_1\}$. In either case, we have

$$\lambda_{S_0} \mathbf{m}(S_1) = \lambda_{S'_0} \mathbf{m}(S_1) = \frac{dn_1 n_2}{(n-1)(n-2)} \cdot \mathbf{m}(S_1) = \frac{dn_1 n_2}{(n-1)(n-2)} \cdot m(S).$$

and

$$c_1^{N-2}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S_1 \cdot S_0}] = c_1^{N-2}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S'_1 \cdot S_0}]$$

Thus

$$\sum_{S_0 \in \hat{\mathcal{P}}_0(\mu)} \lambda_{S_0} \cdot c_1^{N-1}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S_0}] = - \sum_{S \in \mathcal{T}_{1b} \cup \mathcal{T}_{2b}} a_S \cdot c_1^{N-2}(\hat{\mathcal{L}}_\mu) \cap [\hat{D}_{S_1 \cdot S_0}]$$

where S_0 and S_1 are constructed from by S as above. We can now apply Proposition 9.2 and Proposition 9.3 to conclude. \square

Lemma 9.4. *Let $S = \{I_0, I_1, \dots, I_r\}$ be a partition in $\hat{\mathcal{P}}_1(\mu)$. For $j = 1, \dots, r$, let $m_j := d(\mu(I_j) - 1)$. Consider a point $\mathbf{x} \in D_S^*$. Let $C_{\mathbf{x}}^0, \dots, C_{\mathbf{x}}^r$ be the components of $C_{\mathbf{x}}$ where $C_{\mathbf{x}}^j$ is associated to the subset I_j in S . Let $\mathcal{U}_{\mathbf{x}}$ and $\widehat{\mathcal{U}}_{\mathbf{x}}$ be as in §3. Let t_j denote the coordinate function on $\mathcal{U}_{\mathbf{x}}$ that corresponds to the node between $C_{\mathbf{x}}^0$ and $C_{\mathbf{x}}^j$. Then the order of t_j along \hat{D}_S is equal to $\frac{\prod_{i=1}^r m_i}{m_j}$.*

Proof. Recall that $\widehat{\mathcal{U}}_{\mathbf{x}}$ is an irreducible component of the algebraic set

$$\mathcal{U}_{\mathbf{x}}^{\#} = \{(t, u, [v_1 : \cdots : v_r]) \in \mathcal{U}_{\mathbf{x}} \times \mathbb{P}_{\mathbb{C}}^{r-1}, v_k t^{\beta_j} = v_j t^{\beta_k}, j, k = 1, \dots, r\}$$

where

$$t^{\beta_j} = \frac{\prod_{i=1}^r t_i^{m_i}}{t_j^{m_j}}.$$

Over $\mathcal{U}_{\mathbf{x}}^* \times \mathbb{P}_{\mathbb{C}}^{r-1}$, the equations defining $\widehat{\mathcal{U}}_{\mathbf{x}}$ are equivalent to

$$(39) \quad v_1 t_1^{m_1} = \cdots = v_r t_r^{m_r}.$$

Thus we can actually use (39) to define $\widehat{\mathcal{U}}_{\mathbf{x}}$. For concreteness, let us assume that $j = r$. Consider the open affine

$$\widehat{\mathcal{U}}_{\mathbf{x}}^r := \{(t, u, [v_1 : \cdots : v_r]) \in \widehat{\mathcal{U}}_{\mathbf{x}}, v_r \neq 0\}$$

of $\widehat{\mathcal{U}}_{\mathbf{x}}$. Note that $\widehat{\mathcal{U}}_{\mathbf{x}}^r$ is defined by the equations

$$t_r^{m_r} = v_j t_j^{m_j}, \text{ for all } j \in \{1, \dots, r-1\}.$$

In $\widehat{\mathcal{U}}_{\mathbf{x}}^r$, the divisor \widehat{D}_S is defined by the equations $t_1 = \cdots = t_r = 0$.

Let A be the localization of the coordinate ring of $\widehat{\mathcal{U}}_{\mathbf{x}}^r$ at the ideal generated by $\{t_1, \dots, t_r\}$. Since $t_j^{m_j} = \frac{t_r^{m_r}}{v_j}$ in A , every element of A is represented by a quotient P/Q , with $P, Q \in \mathbb{C}(v_1, \dots, v_{r-1})[t_1, \dots, t_r]$, where for $j = 1, \dots, r-1$, the degree of t_j in P and Q is at most $m_j - 1$, and the (t_1, \dots, t_r) -free term in Q is non-zero.

The vanishing order of t_r along \widehat{D}_S is defined to be $\ell(A/(t_r))$, that is the length of $A/(t_r)$ as an A -module (see [25, §1.2]). We have

$$A/(t_r) \simeq \mathbb{C}(v_1, \dots, v_{r-1})[t_1, \dots, t_{r-1}]/(t_1^{m_1}, \dots, t_{r-1}^{m_{r-1}}).$$

Observe that $\mathbb{C}(v_1, \dots, v_{r-1})$ is the residue field of the ideal generated by $\{t_1, \dots, t_r\}$, and we have

$$\dim_{\mathbb{C}(v_1, \dots, v_{r-1})} \mathbb{C}(v_1, \dots, v_{r-1})[t_1, \dots, t_{r-1}]/(t_1^{m_1}, \dots, t_{r-1}^{m_{r-1}}) = \prod_{j=1}^{r-1} m_j.$$

Therefore, we can conclude that $\ell(A/(t_r)) = \prod_{j=1}^{r-1} m_j$ (see [25, Ex. A.1.1]), and the lemma follows. \square

Lemma 9.5. *Let $S = \{I_0, I_1, \dots, I_r\}$ be a partition in $\widehat{\mathcal{P}}_1(\mu)$. Then the coefficient of \widehat{D}_S in \mathcal{E} is $(r-1)\mathbf{m}(S)$.*

Proof. Let \mathbf{x} be a point in D_S^* , and $\mathcal{U}_{\mathbf{x}}, \widehat{\mathcal{U}}_{\mathbf{x}}$ be as in §3. Consider the open affine $\widehat{\mathcal{U}}_{\mathbf{x}}^r \subset \widehat{\mathcal{U}}_{\mathbf{x}}$. On $\widehat{\mathcal{U}}_{\mathbf{x}}^r$, the Cartier divisor \mathcal{E} is defined by the function $t^{\beta_r} = \prod_{j=1}^{r-1} t_j^{m_j}$. By Lemma 9.4, we know that the order of t_j along \widehat{D}_S is $\frac{\prod_{i=1}^r m_i}{m_j}$. Thus the order of $t_j^{m_j}$ is $\prod_{i=1}^r m_i = \mathbf{m}(S)$, and the order of t^{β_r} is $(r-1)\mathbf{m}(S)$. \square

9.3. Applications. It turns out that (10) is particularly efficient even for large values of n for the case of quadratic differentials (that is $d = 2$). Indeed, for $\mu \in (\frac{1}{2}\mathbb{Z})^n$, if n is odd then $\mathcal{J}_n(\mu) = 0$ (by Lemma 9.1) since there must be an integral entry in μ . Therefore we have $\mathcal{T}_{1a} = \mathcal{T}_{2a} = \emptyset$. Making use of this feature, we will show

Proposition 9.6. *We have*

$$(40) \quad \mathcal{J}_{2g+2}\left(\frac{1^{2g+1}}{2}, \frac{3-2g}{2}\right) = (-1)^{g+1} \frac{((2g-1)!!)^2}{2g-1}, \text{ for all } g = 1, 2, \dots$$

$$(41) \quad \mathcal{J}_{2g+2}\left(\frac{1^{g+3}}{2}, \frac{-1^{g-1}}{2}\right) = (-1)^{g-1} \frac{(2g-1)!}{2^{g-1}}, \text{ for all } g = 1, 2, \dots$$

Proof. For (40), we first observe that $\mathcal{J}_4(\frac{1^4}{2}) = 1$, which means that (40) is true for $g = 1$. To simplify the notation, let us write $X_g = \mathcal{J}_{2g+2}(\frac{1^{2g+1}}{2}, \frac{3-2g}{2})$.

We apply (10) to compute X_g . As remarked above, we have $\mathcal{T}_{1a} = \mathcal{T}_{2a} = \emptyset$. We claim that in this case we also have $\mathcal{T}_{2b} = \emptyset$. This is because if $\{I_{01}, I_{02}, I_1, I_2\} \in \mathcal{T}_{2b}$, then each of I_{01} and I_{02} contains a point with negative weight. Since there is only one negative weight in $(\frac{1^{2g+1}}{2}, \frac{3-2g}{2})$, this is impossible.

We now remark that for $g \geq 2$ a partition in \mathcal{T}_{1b} has the form

$$\left\{ \left\{ \frac{1}{2}, \frac{1}{2} \right\}, \left\{ \frac{3-2g}{2} \right\}, \left\{ \frac{1^{2g-1}}{2} \right\} \right\}.$$

In particular, $\#\mathcal{T}_{1b} = \binom{2g+1}{2} = g(2g+1)$. For each $\mathcal{S} \in \mathcal{T}_{1b}$, we have $m(\mathcal{S}) = 2g-3$. Thus (10) implies

$$X_g = -\frac{2(2g-1)}{2g(2g+1)}(2g-3)g(2g+1)X_{g-1} = -(2g-1)(2g-3)X_{g-1}.$$

If $X_{g-1} = (-1)^g \frac{((2g-3)!!)^2}{2^{g-3}}$, then $X_g = (-1)^{g+1} \frac{((2g-1)!!)^2}{2^{g-1}}$ as desired.

We now turn to (41). Let us write $Y_{g-1} = \mathcal{J}_{2g+2}(\frac{1^{g+3}}{2}, \frac{-1^{g-1}}{2})$. Our goal is to show that

$$(42) \quad Y_g = (-1)^g \frac{(2g+1)!}{2^g}.$$

Consider the weight vector $\mu = (\frac{1^{g+4}}{2}, \frac{-1^g}{2})$. A partition in \mathcal{T}_{1a} with respect to μ must have the form

$$\left\{ \left\{ \frac{1}{2}, \frac{1}{2} \right\}, \left\{ \frac{-1}{2} \right\}, \left\{ \frac{1^{g+2}}{2}, \frac{-1^{g-1}}{2} \right\} \right\}$$

Therefore, $\#\mathcal{T}_{1a} = \binom{g+4}{2} \binom{g}{1} = \frac{g(g+3)(g+4)}{2}$. For all $\mathcal{S} \in \mathcal{T}_{1a}$ we have $m(\mathcal{S}) = 1$. Thus the contribution of \mathcal{T}_{1a} in Y_g is

$$-\frac{2(2g+1)}{(2g+2)(2g+3)} \cdot \frac{g(g+3)(g+4)}{2} \cdot Y_{g-1}.$$

A partition in \mathcal{T}_{2a} must have the form

$$\{I_{01}, I_{02}, I_1, I_2\} = \left\{ \left\{ \frac{-1}{2} \right\}, \left\{ \frac{-1}{2} \right\}, \left\{ \frac{1^{k+3}}{2}, \frac{-1^k}{2} \right\}, \left\{ \frac{1^{g-k+1}}{2}, \frac{-1^{g-k-2}}{2} \right\} \right\}.$$

By convention, we will suppose that $n_1 := |I_1| \leq n_2 := |I_2|$. For all $k \in \mathbb{Z}_{\geq 0}$, $k \leq \frac{g}{2} - 1$, let \mathcal{T}_{2b}^k denote the set of partitions $\mathcal{S} = \{I_{01}, I_{02}, I_1, I_2\} \in \mathcal{T}_{2b}$ such that $n_1 = 2k + 3$ and $n_2 = 2g - 2k - 1$. Note that if $n_1 = 2k + 3$ then there are $k \binom{-1}{2}$ -entries in I_1 . We have

$$\#\mathcal{T}_{2b}^k = e_k \binom{g}{2} \binom{g-2}{k} \binom{g+4}{k+3} \quad \text{where } e_k = \begin{cases} 1 & \text{if } k < \frac{g}{2} - 1 \\ 1/2 & \text{if } k = \frac{g}{2} - 1. \end{cases}$$

For each $\mathcal{S} \in \mathcal{T}_{2b}^k$, we have $m(\mathcal{S}) = 1$ and $\epsilon(\mathcal{S}) = 2$. Therefore, the contribution of \mathcal{T}_{2b}^k in Y_g is

$$e_k \cdot 2 \cdot \frac{2(2k+3)(2g-2k-1)}{(2g+3)(2g+2)} \cdot \frac{g(g-1)}{2} \binom{g-2}{k} \binom{g+4}{k+3} Y_k Y_{g-k-2}, \quad \text{for } k = 0, \dots, \lfloor g/2 \rfloor - 1.$$

It follows that the total contribution of \mathcal{T}_{2b} in Y_g is equal to

$$\frac{g(g-1)}{(2g+3)(2g+2)} \sum_{k=0}^{g-2} (2k+3)(2g-2k-1) \binom{g-2}{k} \binom{g+4}{k+3} Y_k Y_{g-k-2}.$$

Thus (10) is equivalent to

$$\begin{aligned} (2g+3)(2g+2)Y_g &= -(2g+1)(g+4)(g+3)g \cdot Y_{g-1} + \\ &\quad + g(g-1) \sum_{k=0}^{g-2} (2k+3)(2g-2k-1) \binom{g-2}{k} \binom{g+4}{k+3} Y_k Y_{g-k-2}. \end{aligned}$$

Assuming (42) holds for $1, 2, \dots, g-1$, we need to show

$$(43) \quad \begin{aligned} (2g+3)! &= 2(2g+1)(g+4)(g+3)g(2g-1)! + \\ &\quad + 4g(g-1) \sum_{k=0}^{g-2} (2k+3)(2g-2k-1) \binom{g-2}{k} \binom{g+4}{k+3} (2k+1)!(2g-2k-3)! \end{aligned}$$

We have

$$\begin{aligned} (2g+1)(g+4)(g+3)g(2g-1)! &= \frac{(g+4)(g+3)}{2} (2g+1)! \\ &= \binom{g}{0} \binom{g+4}{2} (2 \cdot 0 + 1)!(2g - 2 \cdot 0 + 1)! \\ &= \binom{g}{g} \binom{g+4}{g+2} (2 \cdot g + 1)!(2g - 2 \cdot g + 1)! \end{aligned}$$

and

$$4g(g-1)(2k+3)(2g-2k-1) \binom{g-2}{k} (2k+1)!(2g-2k-3)! = \binom{g}{k+1} (2k+3)!(2g-2k-1)!$$

Hence (43) is equivalent to

$$(2g+3)! = \sum_{k=0}^g \binom{g}{k} \binom{g+4}{k+2} (2k+1)!(2g-2k+1)!$$

This identity is the content of Lemma 9.7. The proposition is then proved. \square

Lemma 9.7. *We have*

$$(44) \quad (2g+3)! = \sum_{k=0}^g \binom{g}{k} \binom{g+4}{k+2} (2k+1)!(2g-2k+1)!$$

for all $g \in \mathbb{Z}_{\geq 0}$.

Proof. We first observe

$$\begin{aligned} \binom{g}{k} \binom{g+4}{k+2} (2k+1)!(2g-2k+1)! &= \frac{g!}{k!(g-k)!} \frac{(g+4)!}{(k+2)!(g-k+2)!} (2k+1)!(2g-2k+1)! \\ &= g!(g+4)! \frac{(2k+1)!}{k!(k+2)!} \frac{(2g-2k+1)!}{(g-k)!(g-k+2)!} \\ &= 2^g g!(g+4)! \frac{(2k+1)!!}{(k+2)!} \frac{(2g-2k+1)!!}{(g-k+2)!} \end{aligned}$$

Thus (44) is equivalent to

$$(45) \quad \frac{(2g+3)!}{2^g g!(g+4)!} = \sum_{k=0}^g \frac{(2k+1)!!}{(k+2)!} \frac{(2g-2k+1)!!}{(g-k+2)!}$$

Consider the power series

$$U(x) = \sum_{k=0}^{\infty} \frac{(2k+1)!!}{(k+2)!} x^k$$

Then the right hand side of (45) is the coefficient of x^g in the power series of $U^2(x)$. We remark that

$$\frac{(2k+1)!!}{(k+2)!} = \frac{1}{k+2} \frac{2^{k+1} \prod_{i=0}^k (\frac{1}{2} + i)}{(k+1)!}.$$

Let $V(x) = (1-2x)^{-\frac{1}{2}}$. Then

$$V^{(k)}(0) = 2^k \cdot \frac{1}{2} \cdot \frac{3}{2} \cdots \frac{2k-1}{2} = (2k-1)!!.$$

Thus

$$V(x) = 1 + \sum_{k=1}^{\infty} \frac{(2k-1)!!}{k!} x^k.$$

Let $V_1(x) := \int_0^x V(t) dt = 1 - \sqrt{1-2x}$. The power series of V_1 is given by

$$V_1(x) = x + \sum_{k=1}^{\infty} \frac{(2k-1)!!}{(k+1)!} x^{k+1} = x + \sum_{k=0}^{\infty} \frac{(2k+1)!!}{(k+2)!} x^{k+2}$$

Thus

$$\frac{V_1(x) - x}{x^2} = \sum_{k=0}^{\infty} \frac{(2k+1)!!}{(k+2)!} x^k = U(x).$$

Thus the right hand side of (45) is the coefficient a_g of the x^g term in the power series of the function

$$U^2(x) = \frac{(1-x-\sqrt{1-2x})^2}{x^4} = \frac{2-4x+x^2-2(1-x)\sqrt{1-2x}}{x^4}.$$

A direct calculation shows that

$$a_g = \frac{2(g+1)(2g+3)!!}{(g+4)!} = \frac{2(g+1)(2g+3)!}{2^{g+1}(g+1)!(g+4)!} = \frac{(2g+3)!}{2^g g!(g+4)!}$$

as desired. \square

Remark 9.8. The vectors $(\frac{1}{2}^{2g+1}, \frac{3-2g}{2})$ and $(\frac{1}{2}^{g+3}, -\frac{1}{2}^{g-1})$ correspond respectively to the strata $\Omega^2 \mathcal{M}_{2g+2}(-1^{2g+1}, 2g-3)$ and $\Omega^2 \mathcal{M}_{2g+2}(-1^{g+3}, 1^{g-1})$ of quadratic differentials. In [3], Athreya-Eskin-Zorich study the volumes of strata of quadratic differentials in genus 0 with respect to the Masur-Veech volume form $d\text{vol}^*$. They prove the following formula, which is conjectured by M. Kontsevich,

$$(46) \quad V(k_1, \dots, k_n) = 2\pi^2 \cdot \prod_{i=1}^n v(k_i)$$

where $V(k_1, \dots, k_n)$ is the volume of the stratum $\mathbb{P}\Omega^2 \mathcal{M}_{0,n}(k_1, \dots, k_n)$ (with respect to $d\text{vol}^*$), and

$$v(k) := \frac{k!!}{(k+1)!!} \cdot \pi^k \cdot \begin{cases} \pi & \text{if } k \text{ is odd,} \\ 2 & \text{if } k \text{ is even} \end{cases}$$

for all $k \in \mathbb{Z}_{\geq -1}$. In particular, we have

$$V(-1^{2g+1}, 2g-3) = 2\pi^2 \cdot \frac{(2g-3)!!}{(2g-2)!!} \cdot \pi^{2g-2} = \frac{(2g-3)!!}{2^{g-2}(g-1)!} \pi^{2g}$$

and

$$V(-1^{g+3}, 1^{g-1}) = 2\pi^2 \cdot \left(\frac{\pi^2}{2}\right)^{g-1} = \frac{\pi^{2g}}{2^{g-2}}.$$

When all of the k_i are odd, it can be shown that

$$d\text{vol}^* = (-1)^{\frac{n-2}{2}} \cdot 2^{n-1} \cdot (n-2) \cdot d\text{vol}.$$

Note that in this case n must be even. By Theorem 1.3, we have that

$$V(k_1, \dots, k_n) = (-1)^{\frac{n}{2}} \cdot 2 \cdot \frac{\pi^{n-2}}{(n-3)!} \mathcal{J}_n\left(-\frac{k_1}{2}, \dots, -\frac{k_n}{2}\right).$$

Thus, (40) and (41) allow one to recover Kontsevich's formula for the strata $\Omega^2 \mathcal{M}_{2g+2}(-1^{2g+1}, 2g-3)$ and $\Omega^2 \mathcal{M}_{2g+2}(-1^{g+3}, 1^{g-1})$, $g \in \mathbb{Z}_{\geq 1}$.

APPENDIX A. COMPUTATIONS OF MASUR-VEECH VOLUMES IN LOW DIMENSION BY VINCENT KOZIARZ AND DUC-MANH NGUYEN

By Theorem 1.3, we know that the self-intersection number \hat{D}_μ^{n-3} gives the volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ with respect to $d\text{vol}_1$. In this section, we compute this self-intersection number and the corresponding Masur-Veech volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$, for $n = 4, 5$, $d \in \{3, 4, 6\}$, in the case none of the μ_i is an integer. It is well known that the Masur-Veech volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(\underline{k})$ gives the asymptotics of the number of tilings of the sphere by triangles and squares with prescribed constraints at some vertices (see for instance [51, 12, 11, 35]).

To compute the ratio $\frac{d\text{vol}^{MV}}{d\text{vol}}$, our strategy goes as follows: we first observe that it is enough to compute this ratio locally near any point in $\Omega^d \mathcal{M}_{0,n}(\underline{k})$. Recall that an element $(\mathbb{P}_{\mathbb{C}}^1, x_1, \dots, x_n, q) \in \Omega^d \mathcal{M}_{0,n}(\underline{k})$ defines a flat metric with conical singularities on the sphere, where the cone angle at x_i is

$\alpha_i := 2\pi(1 + \frac{k_i}{d})$. To simplify the discussion, let us assume that there is at most one angle in $\{\alpha_1, \dots, \alpha_n\}$ that is greater than 2π . Thus, we can assume that $0 < \alpha_i < 2\pi$ for $i = 1, \dots, n-1$. We construct a flat surface in $\Omega^d \mathcal{M}_{0,n}(k)$ as follows: let P be a $2(n-1)$ -gon in the plane whose vertices are denoted by $p_1, \dots, p_{2(n-1)}$ in the counterclockwise ordering. We suppose that P satisfies the following

- the sides $\overline{p_{2i-1}p_{2i}}$ and $\overline{p_{2i}p_{2i+1}}$ have the same length, and
- the interior angle at p_{2i} is α_i ,

for $i = 1, \dots, n-1$, (by convention $p_{2n-1} = p_1$). Gluing $\overline{p_{2i-1}p_{2i}}$ and $\overline{p_{2i}p_{2i+1}}$ together, we obtain a flat surface M homeomorphic to the sphere with n conical singularities x_1, \dots, x_n , where x_i corresponds to the vertex p_{2i} of P for $i = 1, \dots, n-1$, and x_n is the identification of $p_1, p_3, \dots, p_{2n-3}$. The cone angle at x_i is clearly α_i for $i = 1, \dots, n-1$. Thus the cone angle at x_n must be α_n . Hence M is an element of $\Omega^d \mathcal{M}_{0,n}(k)$.

Let z_i be the complex number associated to the vector $\overrightarrow{p_{2i}p_{2i-1}}$ for $i = 1, \dots, n-1$. Then the complex number associated to $\overrightarrow{p_{2i}p_{2i+1}}$ is $e^{i\alpha_i} z_i$. It follows that the numbers (z_1, \dots, z_{n-1}) satisfy

$$(47) \quad (1 - e^{i\alpha_1})z_1 + \dots + (1 - e^{i\alpha_{n-1}})z_{n-1} = 0.$$

A neighborhood of M in $\Omega^d \mathcal{M}_{0,n}(k)$ can be identified with an open subset of the hyperplane V defined by (47) in \mathbb{C}^{n-1} . Since $e^{i\alpha_i} \neq 1$ for all $i = 1, \dots, n-1$, we can identify this hyperplane with \mathbb{C}^{n-2} via the natural projection $\mathbb{C}^{n-1} \rightarrow \mathbb{C}^{n-2}$, $(z_1, \dots, z_{n-1}) \mapsto (z_1, \dots, z_{n-2})$. In the coordinates (z_1, \dots, z_{n-2}) , the area of M is given by a Hermitian matrix H , that is

$$\text{Area}(M) = (\bar{z}_1, \dots, \bar{z}_{n-2}) \cdot H \cdot (z_1, \dots, z_{n-2}).$$

Hence $d\text{vol} = \det H \cdot d\lambda_{2(n-2)}$, where $\lambda_{2(n-2)}$ is the Lebesgue measure on \mathbb{C}^{n-2} .

In this setting, the Masur-Veech volume on $\Omega^d \mathcal{M}_{0,n}(k)$ is the unique (real) $(n-2, n-2)$ -form on V such that the lattice $\Lambda := V \cap (\mathbb{Z} \oplus \zeta\mathbb{Z})^{n-1}$ has covolume 1, where $\zeta = i$ if $d = 4$, and $\zeta = e^{\frac{2\pi i}{3}}$ if $d \in \{3, 6\}$. Let Λ' be the projection of Λ in \mathbb{C}^{n-2} . Then Λ' is a sublattice of $(\mathbb{Z} \oplus \zeta\mathbb{Z})^{n-2}$. Let m be the index of Λ' in $(\mathbb{Z} \oplus \zeta\mathbb{Z})^{n-2}$. Since Λ' has covolume $m \cdot \text{Im}(\zeta)^{n-2}$ with respect to the Lebesgue measure, we must have

$$\frac{d\text{vol}^{MV}}{d\lambda_{2(n-2)}} = \frac{1}{m \cdot \text{Im}(\zeta)^{n-2}}.$$

As a consequence, we get

$$(48) \quad \frac{d\text{vol}^{MV}}{d\text{vol}} = \frac{1}{m \cdot \text{Im}(\zeta)^{n-2} \cdot \det H}.$$

In the cases where there are more than one angle in $\{\alpha_1, \dots, \alpha_n\}$ that are greater than 2π , we construct a surface in $\Omega^d \mathcal{M}_{0,n}(k)$ from several polygons, and compute the ratio $d\text{vol}^{MV}/d\text{vol}$ by the same method.

A.1. Case $n = 4$. In this case, $\widehat{\mathcal{M}}_{0,n}(k)$ is isomorphic to $\overline{\mathcal{M}}_{0,4} \simeq \mathbb{P}_{\mathbb{C}}^1$, and $\widehat{\mathcal{D}}_{\mu} = \mathcal{D}_{\mu}$. Thus, the self-intersection number is simply the degree of \mathcal{D}_{μ} . The values of the ratio $d\text{vol}^{MV}/d\text{vol}$ and the Masur-Veech volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(k)$ are recorded in Table 1 (the values on the last column are the product of the ones on the third and the fourth columns with $-\frac{\pi^2}{2d}$).

d	$(-k_i)$	$\frac{1}{d} \deg(\hat{\mathcal{D}}_\mu)$	$d\text{vol}^{MV}/d\text{vol}$	M-V volume
2	(1, 1, 1, 1)	1/2	-1	$\pi^2/8$
3	(2, 2, 1, 1)	1/3	-16/9	$8\pi^2/81$
4	(3, 2, 2, 1)	1/4	-1	$\pi^2/32$
4	(3, 3, 1, 1)	1/4	-2	$\pi^2/16$
4	(3, 3, 3, -1)	-1/4	2	$\pi^2/16$
6	(4, 3, 3, 2)	1/3	-4/9	$\pi^2/81$
6	(4, 4, 3, 1)	1/6	-8/9	$\pi^2/81$
6	(5, 3, 2, 2)	1/6	-8/9	$\pi^2/81$
6	(5, 3, 3, 1)	1/6	-4/3	$\pi^2/54$
6	(5, 4, 2, 1)	1/6	-16/9	$2\pi^2/81$
6	(5, 4, 4, -1)	-1/6	16/9	$2\pi^2/81$
6	(5, 5, 1, 1)	1/6	-16/3	$2\pi^2/27$
6	(5, 5, 3, -1)	-1/6	8/3	$\pi^2/27$
6	(5, 5, 4, -2)	-1/3	16/9	$4\pi^2/81$
6	(5, 5, 5, -3)	-1/2	8/3	$\pi^2/9$

TABLE 1. Volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(k)$ when $n = 4$.

A.2. **Case $n = 5$.** It follows from Theorem 3.13 and Proposition 3.18 that the exceptional divisor \mathcal{E} is non-trivial if and only if there exists a partition $\{I_0, I_1, I_2\}$ of $\{1, \dots, 5\}$ such that $\sum_{i \in I_1} \mu_i > 1$ and $\sum_{i \in I_2} \mu_i > 1$. This implies that $|I_0| = 1$ and $|I_1| = |I_2| = 2$. Thus the stratum of $\overline{\mathcal{M}}_{0,5}$ corresponding to this partition is just one point. Let E be the Weil divisor associated to the Cartier divisor \mathcal{E} . Then each component of E maps to a single point in $\overline{\mathcal{M}}_{0,5}$. Therefore, we have

$$\hat{p}^* \mathcal{D}_\mu \cdot E = \mathcal{D}_\mu \cdot \hat{p}_* E = 0.$$

Hence

$$(49) \quad (\hat{p}^* \mathcal{D}_\mu + \mathcal{E})^2 = \mathcal{D}_\mu^2 + \mathcal{E}^2.$$

Since we have an explicit expression of \mathcal{D}_μ and the intersection of boundary divisors in $\overline{\mathcal{M}}_{0,n}$ is well understood, the computation of \mathcal{D}_μ^2 is straightforward. It remains to compute \mathcal{E}^2 .

To simplify the discussion, let us assume that E has a unique irreducible component associated to a partition $\{I_0, I_1, I_2\}$ as above. Define $r_i := d(\sum_{k \in I_i} \mu_k - 1)$, $i = 1, 2$. By assumption, we have $r_i \in \mathbb{Z}_{>0}$.

By construction, a neighborhood of E in $\overline{\mathcal{M}}_{0,5}(k)$ can be described as $\widehat{\mathcal{U}} = \{(t_1, t_2, [u : v]) \in \mathcal{U} \times \mathbb{P}^1, vt_1^{r_1} = ut_2^{r_2}\}$, where \mathcal{U} is a neighborhood of $(0, 0)$ in \mathbb{C}^2 . Define $U = \{(t_1, t_2, [u : v]) \in \widehat{\mathcal{U}}, v \neq 0\}$ and $V = \{(t_1, t_2, [u : v]) \in \widehat{\mathcal{U}}, u \neq 0\}$. We have

$$U \simeq \{(t_1, t_2, u) \in \mathcal{U} \times \mathbb{C}, t_1^{r_1} = ut_2^{r_2}\} \text{ and } V \simeq \{(t_1, t_2, v) \in \mathcal{U} \times \mathbb{C}, vt_1^{r_1} = t_2^{r_2}\}.$$

Note that $E \cap U$ and $E \cap V$ are both defined by $t_1 = t_2 = 0$.

Lemma A.1. *We have $\mathcal{E} \cdot E = -1$.*

Proof. Since $E \simeq \mathbb{P}^1$, it is enough to check that the restriction of the line bundle associated to \mathcal{E} to E is precisely the tautological line bundle $\mathcal{O}_{\mathbb{P}^1}(-1)$. This is straightforward from the construction. \square

We now show the following formula, which generalizes the usual self-intersection number of the exceptional divisor of the blow-up of a surface at one point.

Lemma A.2. *We have*

$$(50) \quad \mathcal{E}^2 = -r_1 r_2.$$

By Lemma A.1, it is enough to show that the order of \mathcal{E} along E is $r_1 r_2$. This is of course a consequence of Lemma 9.5. We will give here below an alternative proof using complex analytic arguments.

Proof of Lemma A.2 in the analytic setting. Since the order of a function along a Weil divisor is only well defined for normal spaces, we first need consider the normalization of $\widehat{\mathcal{U}}$. Let \tilde{U} (resp. \tilde{V}) be the normalization of U (resp. V). Then the normalization $\tilde{\mathcal{U}}$ of $\widehat{\mathcal{U}}$ is obtained by gluing \tilde{U} and \tilde{V} over $U \cap V$. Let $\tilde{\varphi} : \tilde{\mathcal{U}} \rightarrow \widehat{\mathcal{U}}$ denote the normalizing map.

Let $c := \gcd(r_1, r_2)$ and $m_i := r_i/c$, $i = 1, 2$. Denote by G_{m_i} the cyclic group $\mathbb{Z}/m_i\mathbb{Z}$. It is a well known fact that \tilde{U} is isomorphic to the quotient \mathbb{C}^2/G_{m_1} , where the action of G_{m_1} on \mathbb{C}^2 is generated by $(t, x) \mapsto (\zeta_1 t, \zeta_1^{-m_1} x)$ (here ζ_1 is a primitive m_1 -th root of unity). The normalizing projection $\tilde{\varphi}|_{\tilde{U}} : \tilde{U} \rightarrow U$ is induced by the map $\varphi_U : \mathbb{C}^2 \rightarrow U$, $(t, x) \mapsto (t_1, t_2, u) = (xt^{m_2}, t^{m_1}, x^{r_1})$.

Let \tilde{E} be the pre-image of E in $\tilde{\mathcal{U}}$. The intersection $\tilde{E} \cap \tilde{U}$ is identified with $(\{0\} \times \mathbb{C})/G_{m_1}$. Observe that the restriction of $\tilde{\varphi}$ to \tilde{E} has degree c . This is because $\widehat{\mathcal{U}}$ has c local branches through every point $\{(0, 0)\} \times \mathbb{C}^*$. Therefore, we have

$$\tilde{E} \cdot \tilde{\varphi}^* \mathcal{E} = c \cdot (E \cdot \mathcal{E}) = -c.$$

A neighborhood of a regular point in \tilde{E} can be identified with a neighborhood in \mathbb{C}^2 of a point $(0, x)$ with $x \neq 0$. In this neighborhood \tilde{E} is defined by the equation $t = 0$, while the Cartier divisor $\tilde{\varphi}^* \mathcal{E}$ is defined by $t_2^{r_2} = t_2^{r_2 m_1} = t_2^{c m_1 m_2}$. Thus the order of $\tilde{\varphi}^* \mathcal{E}$ along \tilde{E} is $c m_1 m_2$. It follows that

$$\tilde{\varphi}^* \mathcal{E} \cdot \tilde{\varphi}^* \mathcal{E} = c m_1 m_2 \cdot (\tilde{E} \cdot \tilde{\varphi}^* \mathcal{E}) = -c^2 m_1 m_2 = -r_1 r_2.$$

Since $\deg \tilde{\varphi} = 1$, we get $\mathcal{E}^2 = -r_1 r_2$. □

In Table 2, we record the values of $\hat{\mathcal{D}}_\mu^2$ in the case where $n = 5$ and $d \in \{3, 4, 6\}$, together with the values of the Masur-Veech volume of the corresponding stratum of d -differentials. The values on the last column are the product of the ones in the third and the fourth columns with $\frac{\pi^3}{3!d}$.

REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths: *Geometry of Algebraic Curves. Vol II* (with a contribution by J. Harris) Grundlehren der Mathematischen Wissenschaften 268, *Springer, Heidelberg* (2011).
- [2] A. Avila, A. Eskin, and M. Möller: Symplectic and Isometric $SL(2, \mathbb{R})$ -invariant subbundles of the Hodge bundle, *J. Reine Angew. Math.* **732** (2017), pp. 1–20.
- [3] J.S. Athreya, A. Eskin, and A. Zorich: Right-angled billiards and volumes of moduli spaces of quadratic differentials on $\mathbb{C}P^1$. With an appendix by Jon Chaika. *Ann. Sci. de l'E.N.S.* (4) **49** (2016), no. 6, 1311–1386.
- [4] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, M. Möller: Compactification of strata of Abelian differentials, *Duke Math. Journ.* **167** (2018), no.12, 2347–2416.
- [5] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, M. Möller: Strata of k -differentials, *Algebr. Geom.* **6** (2019), no.2, 196–233.
- [6] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, M. Möller: The moduli spaces of multi-scale differentials, preprint, *arXiv:1910.13492*.

- [7] D. Chen and Q. Gendron: Towards a classification of connected components of the strata of k -differentials, preprint, *arXiv:2101.01650*.
- [8] D. Chen, M. Möller, and A. Sauvaget: Masur-Veech volumes and intersection theory: the principal strata of quadratic differentials (with an appendix by G. Borot, A. Giacchetto, and D. Lewanski), *preprint*, arXiv:1912.02267.
- [9] D. Chen, M. Möller, A. Sauvaget, and D. Zagier: Masur-Veech volumes and intersection theory on moduli spaces of Abelian differentials, *preprint*, arXiv:1901.01785.
- [10] M. Costantini, M. Möller, and J. Zachhuber: The area is a good enough metric, *preprint*, arXiv:1910.14151.
- [11] P. Engel: Hurwitz Theory of elliptic orbifolds I, *Geometry & Topology* 25 (2021), 229–274.
- [12] P. Engel and P. Smillie: The number of convex tilings of the sphere by triangles, squares, or hexagons, *Geometry & Topology* 22 (2018), 2839–2864.
- [13] A. Eskin, M. Kontsevich, and A. Zorich: Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow, *Publ. Math. IHÉS*, **120** (2014) no.1, 207–333.
- [14] A. Eskin, H. Masur, and A. Zorich: Moduli spaces of abelian differentials: The principal boundary, counting problems, and the Siegel-Veech constants, *Publ. Math. I.H.É.S.* **97** (2003), 61–179.
- [15] A. Eskin, M. Mirzakhani, and A. Mohammadi: Isolation, equidistribution, and orbit closures for the $SL(2, \mathbb{R})$ action on moduli space, *Annals of Math.* **182** (2015), no.2, pp. 673–721.
- [16] A. Eskin and M. Mirzakhani: Invariant and stationary measures for the $SL(2, \mathbb{R})$ action on moduli space, *Publ. Math. IHÉS* **127** (2018), no.1, 95–324.
- [17] A. Eskin, C. McMullen, R. Mukamel, and A. Wright: Billiards, quadrilaterals, and moduli spaces, *J. Amer. Math. Soc.* **33** (2020), no. 4, 1039–1086.
- [18] A. Eskin and A. Okounkov: Asymptotics of numbers of branched coverings of a torus and volumes of the moduli spaces of holomorphic differentials, *Invent. Math.* **145** (2001), no. 1, p.59–103.
- [19] A. Eskin and A. Okounkov: Pillowcases and quasimodular forms, *Algebraic geometry and number theory*, 1–15, *Progr. Math.* **253**, Birkhäuser, Boston, MA (2006).
- [20] A. Eskin, A. Okounkov, and R. Pandharipande: The Theta characteristics of a branched covering, *Advances in Math.* **217** (2008), no.3, p.873–888.
- [21] H. Esnault and E. Viehweg: *Lectures on vanishing theorems. Notes grew out of the DMV-seminar on algebraic geometry, held at Reimsburg, October 13-19, 1991*. Basel: Birkhäuser Verlag, 1992.
- [22] B. Farb and D. Margalit: *A primer on Mapping Class Groups* (2012).
- [23] G. Farkas and R. Pandharipande: The moduli space of twisted canonical divisors, *J. Inst. Math. Jussieu*, **17**, no. 3, pp. 615–672.
- [24] S. Filip: Splitting mixed Hodge structures over affine invariant manifolds, *Annals of Math.* **183** (2016), no.2, pp. 681–713.
- [25] W. Fulton: Intersection theory, **Ergebnisse der Mathematik und ihrer Grenzgebiete** **3**, 1984.
- [26] Q. Gendron: The Deligne-Mumford and the Incidence Variety Compactifications of the strata of the moduli space of Abelian differentials, *Annales de l'Institut Fourier*, **68**, no.3 (2018), pp. 1169–1240.
- [27] E. Goujard: Volumes of strata of moduli spaces of quadratic differentials: getting explicit values *Ann. Inst. Fourier* **66** (2016), no. 6, 2203–2251.
- [28] R. Hartshorne: Algebraic Geometry, **Graduate Text in Mathematics** **52**, Springer 1977.
- [29] B. Hassett: Moduli spaces of weighted pointed stable curves, *Adv. Math.* **173** (2003), 316–352
- [30] J. Hubbard and H. Masur: Quadratic differentials and foliations, *Acta Math.* **142**, no.3-4 (1979), p.221–274.
- [31] Y. Kawamata: Subadjunction of log canonical divisors for a subvariety of codimension 2, *Contemporary Math.* **207** (1997), 79–88
- [32] S. Keel: Intersection theory of Moduli Space of Stable N -pointed curves of genus zeros, *Trans. Amer. Math. Soc.* **330** (1992), 545–574.
- [33] M. Kontsevich and A. Zorich: Connected components of the moduli spaces of Abelian differentials, *Invent. Math.* **153** (2003), no.3, 631–678.
- [34] V. Koziarz and D.-M. Nguyen: Complex hyperbolic volume and intersection of boundary divisors in moduli spaces of pointed genus zero curves, *Ann. Sci. de l'É.N.S. (4)* **51** (2018), no.6, 1549–1597.
- [35] V. Koziarz and D.-M. Nguyen: Variation of Hodge structure and enumeration of tilings of surfaces by triangles and squares, *J. École Polytech. Math.* **8** (2021), 831–857.

- [36] F. Labourie: Cyclic surfaces and Hitchin components in rank 2, *Annals of Math. (2)* **185** (2017), no.1, 1–58.
- [37] E. Lanneau: Connected components of the strata of the moduli spaces of quadratic differentials, *Annales Scient. de l'E.N.S. (4)* **41** (2008), no.1, 1–56.
- [38] H. Masur: Interval exchange transformations and measured foliations, *Annals of Math.* **115** (1982), 169–200.
- [39] H. Masur and J. Smillie: Hausdorff dimension of sets of nonergodic measured foliations, *Annals of Math.* **134** (1991), 455–543.
- [40] H. Masur and S. Tabachnikov: Rational billiards and flat structures, *Handbook of dynamical systems, 1A*, North-Holland, Amsterdam (2002), pp. 1015–1089.
- [41] H. Masur and A. Zorich: Multiple saddle connections on flat surfaces and the boundary principle of the moduli space of quadratic differentials, *Geom. Funct. Anal.* **18** (2008), no.3, pp. 919–987.
- [42] A. Mathew: Flatness, semicontinuity, and base change, notes available at <https://math.uchicago.edu/~amathew/semicontinuity.pdf>
- [43] C. McMullen: The Gauss-Bonnet theorem for cone manifolds and volumes of moduli spaces, *Amer. J. Math.*, **139** (2017), no.1, 261–291.
- [44] D. Mumford: Abelian quotient of the Teichmüller modular group, *J. Analyse Math.* **18** (1967), 227–244.
- [45] D. Mumford: Hirzebruch's Proportionality Theorem in the non-compact case, *Invent. Math.*, **42** (1977), 239–272.
- [46] D.-M. Nguyen: Volume forms on moduli spaces of d-differentials, *Geometry & Topology* (to appear), *arXiv:1902.04830*.
- [47] A. Sauvaget: Volumes and Siegel-Veech constants of $\mathcal{H}(2g-2)$ and Hodge integrals *Geom. Funct. Anal.* **28** (2018) no.6, 1756–1779.
- [48] A. Sauvaget: Volume of moduli space of flat surfaces, *preprint*, arXiv:2004.03198.
- [49] A. Sauvaget: Notes on volumes of moduli of k-differentials, *private communication*.
- [50] J. Schmitt: Dimension theory of the moduli space of twisted k-differentials, *Documenta Mathematica* **23** (2018), 871–894.
- [51] W.P. Thurston: Shapes of polyhedra, *Geom. & Top. Monogr.*, Vol. 1: The Epstein birthday schrift, 511–549, Geom. Topol. Publ., Coventry (1998).
- [52] W.A. Veech: Moduli spaces of quadratic differentials, *J. Analyse Math.* **55** (1990), 117 – 171.
- [53] W.A. Veech: Flat surfaces, *Amer. J. Math.* **115** (1993), no.3, pp. 589–689.
- [54] A. Wright: The field of definition of affine invariant submanifolds of the moduli space of abelian differentials. *Geometry & Topology* **18** (2014), no. 3, 1323–1341.
- [55] A. Wright: Cylinder deformations in orbit closures of translation surfaces, *Geometry & Topology* **19** (2015), 413–438.
- [56] A. Wright: From rational billiards to dynamics on moduli spaces, *Bull. Amer. Math. Soc.* (2016).
- [57] A. Zorich: Flat surfaces, *Frontiers in number theory, physics, and geometry*, Springer, Berlin (2006), pp. 437–583.

UNIV. BORDEAUX, IMB, CNRS, UMR 5251, F-33400 TALENCE, FRANCE

Email address, V. Koziarz: vincent.koziarz@math.u-bordeaux.fr

Email address, D.-M. Nguyen: duc-manh.nguyen@math.u-bordeaux.fr

d	$(-k_i)$	$\frac{1}{d^2} \hat{D}_\mu^2$	$d\text{vol}^{MV}/d\text{vol}$	M-V volume
3	(2, 1, 1, 1, 1)	1/9	64/27	$32\pi^3/2187$
3	(2, 2, 2, 1, -1)	-2/9	-64/27	$64\pi^3/2187$
3	(2, 2, 2, 2, -2)	-2/9	-64/27	$64\pi^3/2187$
4	(2, 2, 2, 1, 1)	1/8	1	$\pi^3/192$
4	(3, 2, 1, 1, 1)	1/16	2	$\pi^3/192$
4	(3, 2, 2, 2, -1)	-3/16	-1	$\pi^3/128$
4	(3, 3, 2, 1, -1)	-1/8	-2	$\pi^3/96$
4	(3, 3, 2, 2, -2)	-1/4	-1	$\pi^3/96$
4	(3, 3, 3, 1, -2)	-1/4	-2	$\pi^3/48$
4	(3, 3, 3, 2, -3)	-3/16	-2	$\pi^3/64$
6	(3, 3, 2, 2, 2)	1/6	16/27	$4\pi^3/243$
6	(3, 3, 3, 2, 1)	1/9	8/9	$2\pi^3/729$
6	(4, 3, 2, 2, 1)	1/12	32/27	$2\pi^3/729$
6	(4, 3, 3, 1, 1)	1/18	16/9	$2\pi^3/729$
6	(4, 3, 3, 3, -1)	-5/36	-8/9	$5\pi^3/1458$
6	(4, 4, 2, 1, 1)	1/18	64/27	$8\pi^3/2187$
6	(4, 4, 3, 2, -1)	-1/9	-32/27	$8\pi^3/2187$
6	(4, 4, 3, 3, -2)	-2/9	-16/27	$8\pi^3/2187$
6	(4, 4, 4, 1, -1)	-1/18	-64/27	$8\pi^3/2187$
6	(4, 4, 4, 3, -3)	-1/4	-16/9	$\pi^3/81$
6	(5, 2, 2, 2, 1)	1/36	64/27	$4\pi^3/2187$
6	(5, 3, 2, 1, 1)	1/36	32/9	$2\pi^3/729$
6	(5, 3, 3, 2, -1)	-1/12	-16/9	$\pi^3/243$
6	(5, 3, 3, 3, -2)	-2/9	-8/9	$4\pi^3/729$
6	(5, 4, 1, 1, 1)	1/36	64/9	$4\pi^3/729$
6	(5, 4, 2, 2, -1)	-1/12	-64/27	$4\pi^3/729$
6	(5, 4, 3, 1, -1)	-1/18	-32/9	$4\pi^3/729$
6	(5, 4, 3, 2, -2)	-1/6	-32/27	$4\pi^3/729$
6	(5, 4, 3, 3, -3)	-1/4	-8/9	$\pi^3/162$
6	(5, 4, 4, 1, -2)	-1/9	-64/27	$16\pi^3/2187$
6	(5, 4, 4, 2, -3)	-1/4	-32/27	$2\pi^3/243$
6	(5, 4, 4, 3, -4)	-2/9	-32/27	$16\pi^3/2187$
6	(5, 4, 4, 4, -5)	-5/36	-64/27	$20\pi^3/2187$
6	(5, 5, 2, 1, -1)	-1/18	-64/9	$8\pi^3/729$
6	(5, 5, 2, 2, -2)	-1/9	-64/27	$16\pi^3/2187$
6	(5, 5, 3, 1, -2)	-1/9	-32/9	$8\pi^3/729$
6	(5, 5, 3, 2, -3)	-1/6	-16/9	$2\pi^3/243$
6	(5, 5, 3, 3, -4)	-2/9	-16/9	$8\pi^3/729$
6	(5, 5, 4, -1, -1)	1/18	64/9	$8\pi^3/729$
6	(5, 5, 4, 1, -3)	-1/6	-32/9	$4\pi^3/243$
6	(5, 5, 4, 2, -4)	-2/9	-64/27	$32\pi^3/2187$
6	(5, 5, 4, 3, -5)	-5/36	-32/9	$10\pi^3/729$
6	(5, 5, 5, -1, -2)	1/9	64/9	$16\pi^3/729$
6	(5, 5, 5, 1, -4)	-2/9	-64/9	$32\pi^3/729$
6	(5, 5, 5, 2, -5)	-5/36	-64/9	$20\pi^3/729$
6	(5, 5, 5, 4, -7)	7/36	64/9	$28\pi^3/729$
6	(5, 5, 5, 5, -8)	4/9	64/9	$64\pi^3/729$

TABLE 2. Volume of $\mathbb{P}\Omega^d \mathcal{M}_{0,n}(k)$ when $n = 5$.