

CRITICAL VALUES FOR THE β -TRANSFORMATION WITH A HOLE AT 0

PIETER ALLAART AND DERONG KONG

ABSTRACT. Given $\beta \in (1, 2]$, let T_β be the β -transformation on the unit circle $[0, 1)$ such that $T_\beta(x) = \beta x \pmod{1}$. For each $t \in [0, 1)$ let $K_\beta(t)$ be the survivor set consisting of all $x \in [0, 1)$ whose orbit $\{T_\beta^n(x) : n \geq 0\}$ never hits the open interval $(0, t)$. Kalle et al. proved in [*Ergodic Theory Dynam. Systems*, 40 (9): 2482–2514, 2020] that the Hausdorff dimension function $t \mapsto \dim_H K_\beta(t)$ is a non-increasing Devil's staircase. So there exists a critical value $\tau(\beta)$ such that $\dim_H K_\beta(t) > 0$ if and only if $t < \tau(\beta)$. In this paper we determine the critical value $\tau(\beta)$ for all $\beta \in (1, 2]$, answering a question of Kalle et al. (2020). For example, we find that for the *Komornik-Loreti constant* $\beta \approx 1.78723$ we have $\tau(\beta) = (2 - \beta)/(\beta - 1)$. Furthermore, we show that (i) the function $\tau : \beta \mapsto \tau(\beta)$ is left continuous on $(1, 2]$ with right-hand limits everywhere, but has countably infinitely many discontinuities; (ii) τ has no downward jumps, with $\tau(1+) = 0$ and $\tau(2) = 1/2$; and (iii) there exists an open set $O \subset (1, 2]$, whose complement $(1, 2] \setminus O$ has zero Hausdorff dimension, such that τ is real-analytic, convex and strictly decreasing on each connected component of O . Consequently, the dimension $\dim_H K_\beta(t)$ is not jointly continuous in β and t . Our strategy to find the critical value $\tau(\beta)$ depends on certain substitutions of Farey words and a renormalization scheme from dynamical systems.

CONTENTS

1. Introduction	2
2. Farey words and Farey intervals	8
2.1. Farey words	8
2.2. Quasi-greedy expansions, Farey intervals and Lyndon intervals	10
2.3. Greedy expansions and the symbolic survivor set	12
3. Substitution of Lyndon words	12
3.1. An equivalent definition of the substitution	12
3.2. Properties of the substitution	14
4. Critical values in a basic interval	18
5. Geometrical structure of the basic intervals and bifurcation sets	22
5.1. Tree structure of the Lyndon intervals and relative bifurcation sets	23
5.2. Infinite bifurcation set	27
6. Critical values in the bifurcation sets	29
7. Càdlàg property of the critical value function	34
Acknowledgements	38
References	38

2020 *Mathematics Subject Classification*. Primary: 37B10, 28A78; Secondary: 68R15, 26A30, 37E05.

Key words and phrases. β -transformation, survivor set, Farey word, Lyndon word, Farey interval, bifurcation set, substitution operator, Hausdorff dimension.

1. INTRODUCTION

The mathematical study of dynamical systems with holes, called *open dynamical systems*, was first proposed by Pianigiani and Yorke [27] in 1979. In recent years open dynamical systems have received considerable attention from both theoretical and applied perspectives (cf. [12, 13, 14]). In the general setting, one considers a discrete dynamical system (X, T) , where X is a compact metric space and $T : X \rightarrow X$ is a continuous map having positive topological entropy. Let $H \subset X$ be an open connected set, called the *hole*. It is interesting to study the set of points $x \in X$ whose orbit $\{T^n(x) : n \geq 0\}$ never hits the hole H . In other words, we are interested in the *survivor set*

$$K(H) = \{x \in X : T^n(x) \notin H \forall n \geq 0\} = X \setminus \bigcup_{n=0}^{\infty} T^{-n}(H).$$

It is known that the size of $K(H)$ depends not only on the size but also on the position of the hole H (cf. [7]). In [29, 30] Urbański considered C^2 -expanding, orientation-preserving circle maps with a hole of the form $(0, t)$. In particular, he proved that for the doubling map T_2 on the circle $\mathbb{R}/\mathbb{Z} \sim [0, 1)$, i.e., $T_2 : [0, 1) \rightarrow [0, 1)$; $x \mapsto 2x \pmod{1}$, the Hausdorff dimension of the survivor set $K_2(t) := \{x \in [0, 1) : T_2^n(x) \notin (0, t) \forall n \geq 0\}$ depends continuously on the parameter $t \in [0, 1)$. Furthermore, he showed that the dimension function $\eta_2 : t \mapsto \dim_H K_2(t)$ is a devil's staircase, and studied its bifurcation set. Carminati and Tiozzo [9] showed that the function η_2 has an interesting analytic property: the local Hölder exponent of η_2 at any bifurcation point t is equal to $\eta_2(t)$. For the doubling map T_2 with an arbitrary hole $(a, b) \subset [0, 1)$, Glendinning and Sidorov [18] studied (i) when the survivor set $K_2(a, b) = \{x \in [0, 1) : T_2^n(x) \notin (a, b) \forall n \geq 0\}$ is nonempty; (ii) when $K_2(a, b)$ is infinite; and (iii) when $K_2(a, b)$ has positive Hausdorff dimension. They proved that when the size of the hole (a, b) is strictly smaller than 0.175092, the survivor set $K_2(a, b)$ has positive Hausdorff dimension. The work of Glendinning and Sidorov was partially extended by Clark [10] to the β -dynamical system $([0, 1), T_\beta)$ with a hole (a, b) , where $\beta \in (1, 2]$ and $T_\beta(x) := \beta x \pmod{1}$.

Motivated by the above works, Kalle et al. [20] considered the survivor set in the β -dynamical system $([0, 1), T_\beta)$ with a hole at zero. More precisely, for $t \in [0, 1)$ they determined the Hausdorff dimension of the survivor set

$$K_\beta(t) = \{x \in [0, 1) : T_\beta^n(x) \notin (0, t) \forall n \geq 0\},$$

and showed that the dimension function $\eta_\beta : t \mapsto \dim_H K_\beta(t)$ is a non-increasing Devil's staircase. So there exists a critical value $\tau(\beta) \in [0, 1)$ such that $\dim_H K_\beta(t) > 0$ if and only if $t < \tau(\beta)$. Kalle et al. [20] gave general lower and upper bounds for $\tau(\beta)$. In particular, they showed that $\tau(\beta) \leq 1 - \frac{1}{\beta}$ for all $\beta \in (1, 2]$, and the equality $\tau(\beta) = 1 - \frac{1}{\beta}$ holds for infinitely many $\beta \in (1, 2]$. They left open the interesting question to determine $\tau(\beta)$ for all $\beta \in (1, 2]$. In this paper we give a complete description of the critical value

$$(1.1) \quad \tau(\beta) = \sup \{t : \dim_H K_\beta(t) > 0\} = \inf \{t : \dim_H K_\beta(t) = 0\}$$

for each $\beta \in (1, 2]$. Qualitatively, our main result is:

Theorem 1.

- (i) *The function $\tau : \beta \mapsto \tau(\beta)$ is left continuous on $(1, 2]$ with right-hand limits everywhere (càdlàg), and as a result has only countably many discontinuities;*
- (ii) *τ has no downward jumps;*

- (iii) *There is an open set $O \subset (1, 2]$, whose complement $(1, 2] \setminus O$ has zero Hausdorff dimension, such that τ is real-analytic, convex and strictly decreasing on each connected component of O .*

Quantitatively, the main results are Theorem 2 and Propositions 6.2 and 6.3. Together with Proposition 1.12, they specify the value of $\tau(\beta)$ for all $\beta \in (1, 2]$. In Proposition 1.9 below we give an explicit description of the discontinuities of the map τ , which shows that the dimension $\dim_H K_\beta(t)$ is not jointly continuous in β and t . The closures of the connected components of the set O in Theorem 1 (iii) form a pairwise disjoint collection $\{I_\alpha\}$ of closed intervals which we call *basic intervals* (see Definition 1.5). In the remainder of this introduction we describe these basic intervals by using certain substitutions on Farey words. We then give a formula for $\tau(\beta)$ on each basic interval (see Theorem 2) and decompose the complement $(1, 2] \setminus \bigcup_\alpha I_\alpha$ into countably many disjoint subsets (see Theorem 3), which are of two essentially different types. We then calculate $\tau(\beta)$ on each subset.




FIGURE 1. The graph of the critical value function $\tau(\beta)$ for $\beta \in (1, 2]$. We see that $\tau(\beta) \leq 1 - 1/\beta$ for all $\beta \in (1, 2]$, and the function τ is strictly decreasing in each basic interval $I^{\mathbf{S}}$. For example, the basic interval generated by the Farey word 01 is given by $I^{01} = [\beta_\ell, \beta_*] \approx [1.61803, 1.73867]$ with $((10)^\infty)_{\beta_\ell} = (1100(10)^\infty)_{\beta_*} = 1$. Furthermore, for any $\beta \in I^{01}$ we have $\tau(\beta) = (00(10)^\infty)_\beta = \frac{1}{\beta(\beta^2-1)}$; see Example 1.7 for more details.

To describe the critical value $\tau(\beta)$ we first introduce the Farey words, also called standard words (see [24, Ch. 2.2]). Following a recent paper of Carminati, Isola and Tiozzo [8] we define recursively a sequence of ordered sets $(F_n)_{n=0}^\infty$. Let $F_0 = (0, 1)$, and for $n \geq 0$ the ordered set $F_{n+1} = (v_1, \dots, v_{2^{n+1}+1})$ is obtained from $F_n = (w_1, \dots, w_{2^n+1})$ by inserting for

each $1 \leq j \leq 2^n$ the new word $w_j w_{j+1}$ between the two neighboring words w_j and w_{j+1} . So,

$$(1.2) \quad \begin{aligned} F_1 &= (0, 01, 1), & F_2 &= (0, 001, 01, 011, 1), \\ F_3 &= (0, 0001, 001, 00101, 01, 01011, 011, 0111, 1), \end{aligned}$$

and so on (see Section 2 for more details on Farey words). Set $\Omega_F^* := \bigcup_{n=1}^{\infty} F_n \setminus F_0$. Then each word in Ω_F^* is called a non-degenerate *Farey word*. Note that any word in Ω_F^* has length at least two, and begins with digit 0 and ends with digit 1. We will use the Farey words as basic bricks to construct infinitely many pairwise disjoint closed intervals, so that we can explicitly determine $\tau(\beta)$ for β in each of these intervals. Furthermore, we will show that these closed intervals cover $(1, 2]$ up to a set of zero Hausdorff dimension.

The construction of these basic intervals depends on certain substitutions of Farey words. For this reason we need to introduce a larger class of words, called Lyndon words; see [20, Lemma 3.2].

Definition 1.1. A word $\mathbf{s} = s_1 \dots s_m \in \{0, 1\}^*$ is called *Lyndon* if

$$s_{i+1} \dots s_m \succ s_1 \dots s_{m-i} \quad \forall 0 < i < m.$$

Here and throughout the paper we use lexicographical order \succ between sequences and words; see Section 2. Clearly, the words 0 and 1 are Lyndon. Let Ω_L^* denote the set of all Lyndon words of length at least two. Then by Definition 1.1 it follows that each $\mathbf{s} \in \Omega_L^*$ has a prefix 0 and a suffix 1. It is well-known that each Farey word is Lyndon (cf. [8, Proposition 2.8]). Thus $\Omega_F^* \subset \Omega_L^*$.

Now we define a substitution operator \bullet in Ω_L^* . This requires the following notation. By a *word* we mean a finite string of zeros and ones. For any two words $\mathbf{u} = u_1 \dots u_m, \mathbf{v} = v_1 \dots v_n$ we denote by $\mathbf{uv} = u_1 \dots u_m v_1 \dots v_n$ their concatenation. Furthermore, we write \mathbf{u}^∞ for the periodic sequence with periodic block \mathbf{u} . For a word $\mathbf{w} = w_1 \dots w_n \in \{0, 1\}^n$ we denote $\mathbf{w}^- := w_1 \dots w_{n-1}0$ if $w_n = 1$, and $\mathbf{w}^+ := w_1 \dots w_{n-1}1$ if $w_n = 0$. Furthermore, we denote by $\mathbb{L}(\mathbf{w})$ the lexicographically largest cyclic permutation of \mathbf{w} . Now for two words $\mathbf{s} = s_1 \dots s_m \in \Omega_L^*$ and $\mathbf{r} = r_1 \dots r_\ell \in \{0, 1\}^\ell$ we define

$$(1.3) \quad \mathbf{s} \bullet \mathbf{r} := c_1 \dots c_{\ell m},$$

where

$$c_1 \dots c_m = \begin{cases} \mathbf{s}^- & \text{if } r_1 = 0 \\ \mathbb{L}(\mathbf{s})^+ & \text{if } r_1 = 1, \end{cases}$$

and for $1 \leq j < \ell$,

$$c_{jm+1} \dots c_{(j+1)m} = \begin{cases} \mathbb{L}(\mathbf{s}) & \text{if } r_j r_{j+1} = 00 \\ \mathbb{L}(\mathbf{s})^+ & \text{if } r_j r_{j+1} = 01 \\ \mathbf{s}^- & \text{if } r_j r_{j+1} = 10 \\ \mathbf{s} & \text{if } r_j r_{j+1} = 11. \end{cases}$$

For an equivalent definition of the substitution operator \bullet , see Section 3.

Example 1.2. Let $\mathbf{r} = 01, \mathbf{s} = 001$ and $\mathbf{t} = 011$ be three words in Ω_F^* . Then $\mathbb{L}(\mathbf{r}) = 10$ and $\mathbb{L}(\mathbf{s}) = 100$. So, by (1.3) it follows that

$$\begin{aligned} \mathbf{r} \bullet \mathbf{s} &= \mathbf{r} \bullet 001 = \mathbf{r}^- \mathbb{L}(\mathbf{r}) \mathbb{L}(\mathbf{r})^+ = 001011 \in \Omega_L^*, \\ \mathbf{s} \bullet \mathbf{t} &= \mathbf{s} \bullet 011 = \mathbf{s}^- \mathbb{L}(\mathbf{s})^+ \mathbf{s} = 000101001 \in \Omega_L^*. \end{aligned}$$

Then $\mathbb{L}(\mathbf{r} \bullet \mathbf{s}) = 110010$, and thus

$$(\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t} = (\mathbf{r} \bullet \mathbf{s}) \bullet 011 = (\mathbf{r} \bullet \mathbf{s})^- \mathbb{L}(\mathbf{r} \bullet \mathbf{s})^+ (\mathbf{r} \bullet \mathbf{s}) = 001010110011001011,$$

and

$$\begin{aligned} \mathbf{r} \bullet (\mathbf{s} \bullet \mathbf{t}) &= \mathbf{r} \bullet 000101001 \\ &= \mathbf{r}^- \mathbb{L}(\mathbf{r}) \mathbb{L}(\mathbf{r}) \mathbb{L}(\mathbf{r})^+ \mathbf{r}^- \mathbb{L}(\mathbf{r})^+ \mathbf{r}^- \mathbb{L}(\mathbf{r}) \mathbb{L}(\mathbf{r})^+ = 001010110011001011. \end{aligned}$$

Hence, $(\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t} = \mathbf{r} \bullet (\mathbf{s} \bullet \mathbf{t})$, suggesting that the operator \bullet is associative. On the other hand, observe that $\mathbf{r} \bullet \mathbf{s} = 001011 \neq 000101 = \mathbf{s} \bullet \mathbf{r}$. So \bullet is not commutative.

From Example 1.2 we see that Ω_F^* is not closed under the substitution operator \bullet , since $\mathbf{r} \bullet \mathbf{s} = 001011 \notin \Omega_F^*$. Hence we need the larger collection Ω_L^* . It turns out that Ω_L^* is a non-Abelian semi-group under the substitution operator \bullet .

Proposition 1.3. (Ω_L^*, \bullet) forms a non-Abelian semi-group.

Remark 1.4. The substitution operator \bullet defined in (1.3) is similar to that introduced by Allaart [2], who used it to study the entropy plateaus in unique q -expansions.

Let

$$(1.4) \quad \Lambda := \{\mathbf{S} = \mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \cdots \bullet \mathbf{s}_k : \mathbf{s}_i \in \Omega_F^* \text{ for any } 1 \leq i \leq k; k \in \mathbb{N}\}$$

be the set of all substitutions of Farey words from Ω_F^* . Then by Proposition 1.3 it follows that $\Omega_F^* \subset \Lambda \subset \Omega_L^*$. Moreover, both inclusions are strict. For instance, $001011 = 01 \bullet 001 \in \Lambda \setminus \Omega_F^*$ by Example 1.2 and Proposition 2.4 below, and $0010111 \in \Omega_L^* \setminus \Lambda$.

Given $\beta \in (1, 2]$, for a sequence $(c_i) \in \{0, 1\}^{\mathbb{N}}$ we write

$$((c_i))_\beta := \sum_{i=1}^{\infty} \frac{c_i}{\beta^i}.$$

Now we define the basic intervals.

Definition 1.5. A closed interval $I = [\beta_\ell, \beta_*] \subset (1, 2]$ is called a *basic interval* if there exists a word $\mathbf{S} \in \Lambda$ such that

$$(\mathbb{L}(\mathbf{S})^\infty)_{\beta_\ell} = 1 \quad \text{and} \quad (\mathbb{L}(\mathbf{S})^+ \mathbf{S}^- \mathbb{L}(\mathbf{S})^\infty)_{\beta_*} = 1.$$

The interval $I = I^{\mathbf{S}}$ is also called a basic interval generated by \mathbf{S} .

The subscripts for the endpoints β_ℓ and β_* of a basic interval will be clarified when we define the Lyndon intervals (see Definition 1.8 below). Our second main result gives a formula for $\tau(\beta)$ for any β in a basic interval $I^{\mathbf{S}}$.

Theorem 2.

- (i) *The basic intervals $I^{\mathbf{S}}, \mathbf{S} \in \Lambda$ are pairwise disjoint.*
- (ii) *If $I^{\mathbf{S}}$ is a basic interval generated by $\mathbf{S} \in \Lambda$, then*

$$(1.5) \quad \tau(\beta) = (\mathbf{S}^- \mathbb{L}(\mathbf{S})^\infty)_\beta \quad \text{for every } \beta \in I^{\mathbf{S}}.$$

- (iii) *The function τ is strictly decreasing on $I^{\mathbf{S}}$, and is real-analytic and strictly convex in the interior of $I^{\mathbf{S}}$.*

Remark 1.6. Note that (iii) follows immediately from (ii). For the special case when $\mathbf{S} \in \Omega_F^*$, the formula (1.5) was stated without proof by Kalle et al. [20].

Example 1.7.

(i) Let $\mathbf{s} = 01 \in \Omega_F^*$. Then by Definition 1.5 the basic interval $I^{01} = [\beta_\ell, \beta_*]$ satisfies

$$(\mathbb{L}(01)^\infty)_{\beta_\ell} = ((10)^\infty)_{\beta_\ell} = 1 \quad \text{and} \quad (\mathbb{L}(01)^+(01)^-\mathbb{L}(01)^\infty)_{\beta_*} = (1100(10)^\infty)_{\beta_*} = 1.$$

By numerical calculation we get $I^{01} \approx [1.61803, 1.73867]$ (see Figure 1). In fact, $\beta_\ell = (1 + \sqrt{5})/2$. Theorem 2 yields that

$$\tau(\beta) = (00(10)^\infty)_\beta = \frac{1}{\beta(\beta^2 - 1)} \quad \forall \beta \in I^{01}.$$

(ii) Let $\mathbf{s}_1 = \mathbf{s}_2 = 01 \in \Omega_F^*$. Then $\mathbf{S} = \mathbf{s}_1 \bullet \mathbf{s}_2 = 01 \bullet 01 = 0011$. By Definition 1.5 the basic interval $I^{\mathbf{s}_1 \bullet \mathbf{s}_2} = I^{0011} = [\beta_\ell, \beta_*]$ is given implicitly by

$$\begin{aligned} (\mathbb{L}(0011)^\infty)_{\beta_\ell} &= ((1100)^\infty)_{\beta_\ell} = 1, \\ (\mathbb{L}(0011)^+(0011)^-\mathbb{L}(0011)^\infty)_{\beta_*} &= (11010010(1100)^\infty)_{\beta_*} = 1. \end{aligned}$$

Numerical calculation gives $I^{0011} \approx [1.75488, 1.78431]$ (see Figure 1), and Theorem 2 implies

$$\tau(\beta) = (\mathbf{S}^-\mathbb{L}(\mathbf{S})^\infty)_\beta = (0010(1100)^\infty)_\beta = \frac{1}{\beta^3} + \frac{1 + \beta}{\beta^2(\beta^4 - 1)} \quad \forall \beta \in I^{0011}.$$

Next, we introduce the Lyndon intervals.

Definition 1.8. For each Lyndon word $\mathbf{S} \in \Omega_L^*$ the interval $J^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_r^{\mathbf{S}}] \subset (1, 2]$ is called a *Lyndon interval* generated by \mathbf{S} if

$$(\mathbb{L}(\mathbf{S})^\infty)_{\beta_\ell^{\mathbf{S}}} = 1 \quad \text{and} \quad (\mathbb{L}(\mathbf{S})^+\mathbf{S}^\infty)_{\beta_r^{\mathbf{S}}} = 1.$$

If in particular $\mathbf{S} \in \Omega_F^*$, we call $J^{\mathbf{S}}$ a *Farey interval*.

We remark that the Farey intervals defined in [20, Definition 4.5] are half-open intervals, which is slightly different from our definition. It turns out that the discontinuity points of τ are precisely the right endpoints of the Lyndon intervals $J^{\mathbf{S}}$ with $\mathbf{S} \in \Lambda$:

Proposition 1.9. *The function τ is continuous on $(1, 2] \setminus \{\beta_r^{\mathbf{S}} : \mathbf{S} \in \Lambda\}$. On the other hand, for each $\mathbf{S} \in \Lambda$, we have*

$$(1.6) \quad \lim_{\beta \searrow \beta_r^{\mathbf{S}}} \tau(\beta) = (\mathbf{S}^\infty)_{\beta_r^{\mathbf{S}}} > (\mathbf{S}0^\infty)_{\beta_r^{\mathbf{S}}} = \tau(\beta_r^{\mathbf{S}}).$$

Remark 1.10. Proposition 1.9 implies that, although the dimension $\dim_H K_\beta(t)$ is continuous in t for fixed β , it is not jointly continuous in β and t . In particular, when $t = \tau(\beta_r^{\mathbf{S}})$ for $\mathbf{S} \in \Lambda$, the function $\beta \mapsto \dim_H K_\beta(t)$ has a jump at $\beta_r^{\mathbf{S}}$.

It was shown in [20, Section 4] that the Farey intervals $J^{\mathbf{s}}, \mathbf{s} \in \Omega_F^*$ are pairwise disjoint, and the *bifurcation set*

$$E := (1, 2] \setminus \bigcup_{\mathbf{s} \in \Omega_F^*} J^{\mathbf{s}}$$

has zero Hausdorff dimension. We strengthen this result slightly and show in Proposition 5.6 (i) that E is uncountable and has zero packing dimension.

From Definitions 1.5 and 1.8 it follows that $I^{\mathbf{S}} \subset J^{\mathbf{S}}$ for any $\mathbf{S} \in \Lambda$, and the two intervals $I^{\mathbf{S}}$ and $J^{\mathbf{S}}$ have the same left endpoint (see Proposition 5.1.) In Proposition 5.6 (ii) we show

that for any $\mathbf{S} \in \Lambda$ the Lyndon intervals $J^{\mathbf{S}\bullet\mathbf{r}}$, $\mathbf{r} \in \Omega_F^*$ are pairwise disjoint subsets of $J^{\mathbf{S}} \setminus I^{\mathbf{S}}$, and the *relative bifurcation set*

$$E^{\mathbf{S}} := (J^{\mathbf{S}} \setminus I^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S}\bullet\mathbf{r}}$$

is also uncountable and has zero box-counting dimension. In Proposition 5.1 we prove that the Lyndon intervals $J^{\mathbf{S}}$, $\mathbf{S} \in \Lambda$ have a tree structure. This gives rise to the *infinite bifurcation set*

$$(1.7) \quad E_{\infty} := \bigcap_{k=1}^{\infty} \bigcup_{\mathbf{S} \in \Lambda(k)} J^{\mathbf{S}},$$

where $\Lambda(k) := \{\mathbf{S} = \mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_k : \mathbf{s}_i \in \Omega_F^* \forall 1 \leq i \leq k\}$. It follows at once that E_{∞} is uncountable; we show in Proposition 5.8 that it has zero Hausdorff dimension.

Combining the above results we obtain our last main theorem:

Theorem 3. *The interval $(1, 2]$ can be partitioned as*

$$(1, 2] = E \cup E_{\infty} \cup \bigcup_{\mathbf{S} \in \Lambda} E^{\mathbf{S}} \cup \bigcup_{\mathbf{S} \in \Lambda} I^{\mathbf{S}},$$

and the basic intervals $\{I^{\mathbf{S}} : \mathbf{S} \in \Lambda\}$ cover $(1, 2]$ up to a set of zero Hausdorff dimension.

Remark 1.11. It is worth mentioning that the Lyndon intervals $J^{\mathbf{S}}$ and the relative bifurcation sets $E^{\mathbf{S}}$ constructed in our paper have similar geometrical structure as the relative entropy plateaus and relative bifurcation sets studied in [1], where they were used to describe the local dimension of the set of univoque bases.

The following result was established in the proof of [20, Theorem D].

Proposition 1.12. *For any $\beta \in (1, 2]$ we have $\tau(\beta) \leq 1 - \frac{1}{\beta}$. Furthermore,*

$$\tau(\beta) = 1 - \frac{1}{\beta} \quad \text{for any } \beta \in E.$$

Thus, in view of Theorem 3 it remains to determine $\tau(\beta)$ for $\beta \in E^{\mathbf{S}}$ with $\mathbf{S} \in \Lambda$ and for $\beta \in E_{\infty}$. In Proposition 6.2 we compute $\tau(\beta)$ for $\beta \in E^{\mathbf{S}}$ by relating the relative bifurcation set $E^{\mathbf{S}}$ to the bifurcation set E via a renormalization map $\Psi_{\mathbf{S}}$. Proposition 6.3 gives an expression for $\tau(\beta)$ when $\beta \in E_{\infty}$. As an illustration of the latter, in Proposition 6.4 we construct in each Farey interval $J^{\mathbf{s}}$ a transcendental base $\beta_{\infty}^{\mathbf{s}} \in E_{\infty}$ and give an explicit formula for $\tau(\beta_{\infty}^{\mathbf{s}})$. Here we point out an interesting connection with unique β -expansions: Let $\beta \approx 1.78723$ be the *Komornik-Loreti constant* (cf. [21]); that is, β is the smallest base in which 1 has a unique expansion. Then it follows from Proposition 6.4 that $\beta = \beta_{\infty}^{01} \in E_{\infty}$, and $\tau(\beta) = \frac{2-\beta}{\beta-1} \approx 0.270274$.

The rest of the paper is organized as follows. In Section 2 we recall some properties of Farey words and Farey intervals, as well as greedy and quasi-greedy β -expansions. In Section 3 we give an equivalent definition of the substitution operator \bullet , and prove Proposition 1.3. The proof of Theorem 2 is given in Section 4. At the heart of the argument is Proposition 4.1, which clarifies the role of the special Lyndon words $\mathbf{S} \in \Lambda$ and is used in several settings to derive the upper bound for $\tau(\beta)$. The bifurcation sets $E^{\mathbf{S}}$, $\mathbf{S} \in \Lambda$ and E_{∞} are studied in detail in Section 5, where we show that all of these sets have zero Hausdorff dimension, proving Theorem 3. In Section 6 we determine the critical value $\tau(\beta)$ for β in the relative

bifurcation sets $E^{\mathbf{S}}$ and the infinite bifurcation set E_{∞} . Finally, in Section 7 we show that the function $\beta \mapsto \tau(\beta)$ is càdlàg, and prove Proposition 1.9 and Theorem 1.

2. FAREY WORDS AND FAREY INTERVALS

In this section we recall some properties of Farey words, which are vital in determining the critical value $\tau(\beta)$. We also recall from [20] the Farey intervals, and review basic properties of greedy and quasi-greedy β -expansions.

First we introduce some terminology from symbolic dynamics (cf. [23]). Let $\{0, 1\}^{\mathbb{N}}$ be the set of all infinite sequences of zeros and ones. Denote by σ the left shift map. Then $(\{0, 1\}^{\mathbb{N}}, \sigma)$ is a full shift. By a *word* we mean a finite string of zeros and ones. Let $\{0, 1\}^*$ be the set of all words over the alphabet $\{0, 1\}$ together with the empty word ϵ . For a word $\mathbf{c} \in \{0, 1\}^*$ we denote its length by $|\mathbf{c}|$, and for a digit $a \in \{0, 1\}$ we denote by $|\mathbf{c}|_a$ the number of occurrences of a in the word \mathbf{c} . For two words $\mathbf{c} = c_1 \dots c_m$ and $\mathbf{d} = d_1 \dots d_n$ in $\{0, 1\}^*$ we write $\mathbf{cd} = c_1 \dots c_m d_1 \dots d_n$ for their concatenation. For $n \in \mathbb{N}$ we denote by \mathbf{c}^n the n -fold concatenation of \mathbf{c} with itself, and by \mathbf{c}^{∞} the periodic sequence with period block \mathbf{c} .

Throughout the paper we will use the lexicographical order ' \prec, \preceq, \succ ' or ' \succcurlyeq ' between sequences and words. For example, for two sequences $(c_i), (d_i) \in \{0, 1\}^{\mathbb{N}}$, we say $(c_i) \prec (d_i)$ if $c_1 < d_1$, or there exists $n \in \mathbb{N}$ such that $c_1 \dots c_n = d_1 \dots d_n$ and $c_{n+1} < d_{n+1}$. For two words \mathbf{c}, \mathbf{d} , we say $\mathbf{c} \prec \mathbf{d}$ if $\mathbf{c}0^{\infty} \prec \mathbf{d}0^{\infty}$. We also recall from Section 1 that if $\mathbf{c} = c_1 \dots c_m$ with $c_m = 0$, then $\mathbf{c}^+ = c_1 \dots c_{m-1}1$; and if $\mathbf{c} = c_1 \dots c_m$ with $c_m = 1$, then $\mathbf{c}^- = c_1 \dots c_{m-1}0$. Finally, for a word $\mathbf{c} = c_1 c_2 \dots c_n$ we denote its *reflection* by $\bar{\mathbf{c}} := (1 - c_1)(1 - c_2) \dots (1 - c_n)$.

2.1. Farey words. Farey words have attracted much attention in the literature due to their intimate connection with rational rotations on the circle (see [24, Chapter 2]) and their one-to-one correspondence with the rational numbers in $[0, 1]$ (see (2.1) below). In the following we adopt the definition from a recent paper of Carminati, Isola and Tiozzo [8].

First we recursively define a sequence of ordered sets $F_n, n = 0, 1, 2, \dots$. Let $F_0 = (0, 1)$; and for $n \geq 0$ the ordered set $F_{n+1} = (v_1, \dots, v_{2^{n+1}+1})$ is obtained from $F_n = (w_1, \dots, w_{2^n+1})$ by

$$\begin{cases} v_{2i-1} = w_i & \text{for } 1 \leq i \leq 2^n + 1, \\ v_{2i} = w_i w_{i+1} & \text{for } 1 \leq i \leq 2^n. \end{cases}$$

In other words, F_{n+1} is obtained from F_n by inserting for each $1 \leq j \leq 2^n$ the new word $w_j w_{j+1}$ between the two neighboring words w_j and w_{j+1} . See (1.2) for examples. Note that for each $n \geq 0$ the ordered set F_n consists of $2^n + 1$ words which are listed from the left to the right in lexicographically increasing order. We call $w \in \{0, 1\}^*$ a *Farey word* if $w \in F_n$ for some $n \geq 0$, and we denote by $\Omega_F := \bigcup_{n=1}^{\infty} F_n$ the set of all Farey words. As shown in [8, Proposition 2.3], the set Ω_F can be bijectively mapped to $\mathbb{Q} \cap [0, 1]$ via the map

$$(2.1) \quad \xi : \Omega_F \rightarrow \mathbb{Q} \cap [0, 1]; \quad \mathbf{s} \mapsto \frac{|\mathbf{s}|_1}{|\mathbf{s}|}.$$

So, $\xi(\mathbf{s})$ is the frequency of the digit 1 in \mathbf{s} .

For each $n \geq 1$ set

$$F_n^* := F_n \setminus \{0, 1\},$$

and

$$F_n^0 := \{w \in F_n^* : |w|_0 > |w|_1\}, \quad F_n^1 := \{w \in F_n^* : |w|_1 > |w|_0\}.$$

For example, $F_1^* = (01), F_2^* = (001, 01, 011)$, and $F_2^0 = (001), F_2^1 = (011)$. The following decomposition can be deduced from [8, Proposition 2.3].

Lemma 2.1. *For any $n \geq 2$ we have $F_n^* = F_n^0 \cup F_1^* \cup F_n^1$.*

The ordered sets $F_n^*, n \geq 1$ can also be obtained via substitutions. We define the two substitution operators by

$$(2.2) \quad U_0 : \begin{cases} 0 & \mapsto 0 \\ 1 & \mapsto 01, \end{cases} \quad \text{and} \quad U_1 : \begin{cases} 0 & \mapsto 01 \\ 1 & \mapsto 1. \end{cases}$$

Then U_0 and U_1 naturally induce a map on $\{0, 1\}^*$ or $\{0, 1\}^{\mathbb{N}}$. For example,

$$U_0 : \{0, 1\}^* \rightarrow \{0, 1\}^*; \quad c_1 \dots c_n \mapsto U_0(c_1) \dots U_0(c_n).$$

The following result was proven in [8, Proposition 2.9].

Lemma 2.2. *For each $a \in \{0, 1\}$ the map $U_a : F_n^* \rightarrow F_{n+1}^a$ is bijective.*

By Lemmas 2.1 and 2.2 it follows that the ordered sets F_n^* can be obtained by the substitution operators U_0 and U_1 on the set $F_1^* = (01)$. We will clarify this in the next proposition. Let Ω_F^* be the set of all non-degenerate Farey words, i.e.,

$$\Omega_F^* = \bigcup_{n=1}^{\infty} F_n^*.$$

For a word $\mathbf{c} = c_1 \dots c_m \in \{0, 1\}^*$ let $\mathbb{S}(\mathbf{c})$ and $\mathbb{L}(\mathbf{c})$ be the lexicographically smallest and largest cyclic permutations of \mathbf{c} , respectively. In other words, $\mathbb{S}(\mathbf{c})$ is the lexicographically smallest word among

$$c_1 c_2 \dots c_m, \quad c_2 \dots c_m c_1, \quad c_3 \dots c_m c_1 c_2, \quad \dots, \quad c_m c_1 \dots c_{m-1};$$

and $\mathbb{L}(\mathbf{c})$ is the lexicographically largest word in the above list. The following properties of Farey words are well known (see, e.g., [8, Proposition 2.5]).

Lemma 2.3. *Let $\mathbf{s} = s_1 \dots s_m \in \Omega_F^*$. Then*

- (i) $\mathbb{S}(\mathbf{s}) = \mathbf{s}$ and $\mathbb{L}(\mathbf{s}) = s_m s_{m-1} \dots s_1$.
- (ii) \mathbf{s}_- is a palindrome; that is $s_1 \dots s_{m-1} (s_m - 1) = (s_m - 1) s_{m-1} s_{m-2} \dots s_1$.
- (iii) The word \mathbf{s} has a conjugate $\tilde{\mathbf{s}} \in \Omega_F^*$, given by

$$(2.3) \quad \tilde{\mathbf{s}} := \overline{\mathbb{L}(\mathbf{s})} = 0 \overline{s_2 \dots s_{m-1}} 1.$$

The last equality in (2.3) follows from statements (i) and (ii). In terms of the correspondence (2.1), if $\xi(\mathbf{s}) = r \in \mathbb{Q} \cap [0, 1]$, then $\xi(\tilde{\mathbf{s}}) = 1 - r$. Note also that the conjugate of $\tilde{\mathbf{s}}$ is simply \mathbf{s} itself.

The following explicit description of Ω_F^* will be useful in Section 4 to prove the upper bound for $\tau(\beta)$.

Proposition 2.4. *Ω_F^* consists of all words in one of the following forms:*

- (i) 01^p or $0^p 1$ for some $p \in \mathbb{N}$;
- (ii) $01^p 01^{p+t_1} \dots 01^{p+t_N} 01^{p+1}$ for some $p \in \mathbb{N}$ and Farey word $0t_1 \dots t_N 1 \in \Omega_F^*$.
- (iii) $0^{p+1} 10^{p+t_1} 1 \dots 0^{p+t_N} 10^p 1$ for some $p \in \mathbb{N}$ and Farey word $0t_1 \dots t_N 1 \in \Omega_F^*$;

Proof. Note that $01 = U_1(0) = U_0(1) \in F_1^* \subset \Omega_F^*$. Furthermore, for $p \in \mathbb{N}$ and $0t_1 \dots t_N 1 \in \Omega_F^*$ we have

$$\begin{aligned} 01^p &= U_1(01^{p-1}) = U_1^{p-1}(U_0(1)), \\ 0^p 1 &= U_0(0^{p-1} 1) = U_0^{p-1}(U_1(0)), \\ 01^p 01^{p+t_1} \dots 01^{p+t_N} 01^{p+1} &= U_1^p(U_0(0t_1 \dots t_N 1)), \\ 0^{p+1} 10^{p+t_1} 1 \dots 0^{p+t_N} 10^p 1 &= U_0^p(U_1(0\overline{t_1 \dots t_N} 1)). \end{aligned}$$

By Lemma 2.3 (iii), if $0t_1 \dots t_N 1 \in \Omega_F^*$ then $0\overline{t_1 \dots t_N} 1 \in \Omega_F^*$ as well. Hence by Lemma 2.2, all the above words lie in Ω_F^* .

To prove the converse, it suffices to show that each word in Ω_F^* is of the form $U_0^p(U_1(\mathbf{t}))$ or $U_1^p(U_0(\mathbf{t}))$ for some $p \geq 0$ and Farey word $\mathbf{t} \in \Omega_F$. This is clearly true for $01 = U_0^0(U_1(0))$, where U_0^0 denotes the identity map. Let $n \geq 1$ and suppose the statement is true for all Farey words in F_n^* . Take $\mathbf{s} \in F_{n+1}^*$ with $\mathbf{s} \neq 01$. By Lemmas 2.1 and 2.2, $\mathbf{s} = U_0(\mathbf{t})$ or $\mathbf{s} = U_1(\mathbf{t})$ for some Farey word $\mathbf{t} \in F_n^*$. We assume the former, as the argument for the second case is similar. By the induction hypothesis, either $\mathbf{t} = U_0^p(U_1(\mathbf{u}))$ for some $\mathbf{u} \in \Omega_F$ and $p \geq 0$, in which case $\mathbf{s} = U_0^{p+1}(U_1(\mathbf{u}))$; or $\mathbf{t} = U_1^p(U_0(\mathbf{u}))$ for some $\mathbf{u} \in \Omega_F$ and $p \geq 1$, in which case $\mathbf{s} = U_0(U_1(\mathbf{v}))$ where $\mathbf{v} = U_1^{p-1}(U_0(\mathbf{u})) \in \Omega_F$. In both cases, \mathbf{s} is of the required form. \square

Observe that the two types of words in Proposition 2.4 (i) are each others conjugates, and the conjugate of a Farey word of type (ii) is a Farey word of type (iii), and vice versa. For more properties of Farey words we refer to the book of Lothaire [24] and the references therein.

2.2. Quasi-greedy expansions, Farey intervals and Lyndon intervals. Given $\beta \in (1, 2]$, let $\delta(\beta) = \delta_1(\beta)\delta_2(\beta) \dots \in \{0, 1\}^{\mathbb{N}}$ be the *quasi-greedy* β -expansion of 1 (cf. [11]), i.e., $\delta(\beta)$ is the lexicographically largest sequence not ending with 0^∞ such that $(\delta_i(\beta))_\beta = 1$. The following property of $\delta(\beta)$ is well known (cf. [6]).

Lemma 2.5.

- (i) *The map $\beta \mapsto \delta(\beta)$ is an increasing bijection from $\beta \in (1, 2]$ to the set of sequences $(a_i) \in \{0, 1\}^{\mathbb{N}}$ not ending with 0^∞ and satisfying*

$$\sigma^n((a_i)) \preceq (a_i) \quad \forall n \geq 0.$$

- (ii) *The map $\beta \mapsto \delta(\beta)$ is left continuous everywhere on $(1, 2]$ with respect to the order topology, and it is right continuous at $\beta_0 \in (1, 2)$ if and only if $\delta(\beta_0)$ is not periodic. Furthermore, if $\delta(\beta_0) = (a_1 \dots a_m)^\infty$ with minimal period m , then $\delta(\beta) \searrow a_1 \dots a_m^+ 0^\infty$ as $\beta \searrow \beta_0$.*

Recall from Definition 1.1 that for a word $\mathbf{s} = s_1 \dots s_m \in \Omega_L^*$ we have $s_{i+1} \dots s_m \succ s_1 \dots s_{m-i}$ for all $1 \leq i < m$. The following basic fact can be found in [5, Theorem 1.5.3]:

Lemma 2.6. *Let $\mathbf{c} = c_1 \dots c_m \in \{0, 1\}^*$, and suppose two cyclic permutations of \mathbf{c} are equal (that is, $c_{i+1} \dots c_m c_1 \dots c_i = c_{j+1} \dots c_m c_1 \dots c_j$ where $i \neq j$). Then \mathbf{c} is periodic; in other words, $\mathbf{c} = \mathbf{b}^k$ for some word \mathbf{b} and $k \geq 2$.*

In fact, the length of \mathbf{b} in Lemma 2.6 can be taken to equal $\gcd(|i - j|, m)$.

Lemma 2.7. *Let $\mathbf{s} \in \Omega_L^*$ and $\mathbf{a} = \mathbb{L}(\mathbf{s}) = a_1 \dots a_m$. Then*

$$(2.4) \quad a_{i+1} \dots a_m \prec a_1 \dots a_{m-i} \quad \forall 1 \leq i < m.$$

Furthermore,

$$(2.5) \quad \sigma^n(\mathbf{a}^+ \mathbf{s} \mathbf{a}^\infty) \preccurlyeq \mathbf{a}^+ \mathbf{s} \mathbf{a}^\infty \quad \forall n \geq 0.$$

Proof. First we prove (2.4). Since \mathbf{s} is Lyndon, it is not periodic. Hence $\mathbf{a} = \mathbb{L}(\mathbf{s})$ is not periodic, because any cyclic permutation of a periodic word is periodic. Since $\mathbf{a} = \mathbb{L}(\mathbf{s})$, we have

$$a_{i+1} \dots a_m \preccurlyeq a_1 \dots a_{m-i} \quad \forall 1 \leq i < m.$$

Suppose equality holds for some i . Then

$$a_{i+1} \dots a_m a_1 \dots a_i = a_1 \dots a_{m-i} a_1 \dots a_i \succcurlyeq a_1 \dots a_{m-i} a_{m-i+1} \dots a_m = \mathbf{a},$$

so $a_{i+1} \dots a_m a_1 \dots a_i = \mathbb{L}(\mathbf{s}) = \mathbf{a}$ by definition of $\mathbb{L}(\mathbf{s})$. By Lemma 2.6, this cannot happen, since \mathbf{a} is not periodic.

Next we prove (2.5). Since $\mathbf{s} = s_1 \dots s_m$ is a Lyndon word, any word of length $k \in \{1, \dots, m-1\}$ occurring in $\mathbf{a} = \mathbb{L}(\mathbf{s})$ is lexicographically larger than or equal to $s_1 \dots s_k$. By (2.4) it follows that

$$(2.6) \quad a_{k+1} \dots a_m^+ s_1 \dots s_k \preccurlyeq a_1 \dots a_{m-k} a_{m-k+1} \dots a_m \prec a_1 \dots a_m^+$$

for all $0 < k < m$. Hence, by (2.6) and (2.4) we conclude that $\sigma^n(\mathbf{a}^+ \mathbf{s} \mathbf{a}^\infty) \prec \mathbf{a}^+ \mathbf{s} \mathbf{a}^\infty$ for all $n \geq 1$. This completes the proof. \square

Lemma 2.8. *Let $\beta \in (1, 2)$. Then $\delta(\beta)$ is periodic if and only if $\delta(\beta) = \mathbb{L}(\mathbf{s})^\infty$ for some Lyndon word \mathbf{s} of length at least two.*

Proof. Suppose $\delta(\beta) = (a_1 \dots a_m)^\infty$ with minimal period block $\mathbf{a} = a_1 \dots a_m$. Then $m \geq 2$ since $\beta < 2$. Take $\mathbf{s} := \mathbb{S}(\mathbf{a})$. Then $\mathbf{a} = \mathbb{L}(\mathbf{s})$, and

$$(2.7) \quad s_{i+1} \dots s_m \succcurlyeq s_1 \dots s_{m-i} \quad \forall 1 \leq i < m.$$

If equality holds for some i , then we deduce just as in the proof of Lemma 2.7 that \mathbf{s} is periodic. But then \mathbf{a} is also periodic, contradicting that m is the minimal period of $\delta(\beta)$. Hence, strict inequality holds in (2.7), and \mathbf{s} is Lyndon. The converse is trivial. \square

Recall the Farey intervals and Lyndon intervals from Definition 1.8. The following properties of Lyndon intervals and Farey intervals were established in the proof of [20, Theorem C].

Lemma 2.9.

- (i) *The Farey intervals $J^{\mathbf{s}}$, $\mathbf{s} \in \Omega_F^*$ are pairwise disjoint, and their union is dense in $(1, 2]$.*
- (ii) *Any two Lyndon intervals are either disjoint, or one is contained in the other.*
- (iii) *For any Lyndon interval $J^{\mathbf{S}}$, $\mathbf{S} \in \Omega_L^*$ there exists a unique Farey interval $J^{\mathbf{s}}$ such that $J^{\mathbf{S}} \subset J^{\mathbf{s}}$.*

Note that (iii) follows immediately from (i) and (ii).

2.3. Greedy expansions and the symbolic survivor set. Given $\beta \in (1, 2]$ and $t \in [0, 1)$, we call the sequence $(d_i) \in \{0, 1\}^{\mathbb{N}}$ a β -*expansion* of t if $((d_i))_{\beta} = t$. Note that a point $t \in [0, 1)$ may have multiple β -expansions. We denote by $b(t, \beta) = (b_i(t, \beta)) \in \{0, 1\}^{\mathbb{N}}$ the *greedy* β -expansion of t , which is the lexicographically largest expansion of t in base β . Since $T_{\beta}(t) = \beta t \pmod{1}$, it follows that $b(T_{\beta}^n(t), \beta) = \sigma^n(b(t, \beta)) = b_{n+1}(t, \beta)b_{n+2}(t, \beta) \dots$. The following result was established by Parry [26] and de Vries and Komornik [15, Lemma 2.5 and Proposition 2.6].

Lemma 2.10. *Let $\beta \in (1, 2]$. The map $t \mapsto b(t, \beta)$ is an increasing bijection from $[0, 1)$ to*

$$\{(d_i) \in \{0, 1\}^{\mathbb{N}} : \sigma^n((d_i)) \prec \delta(\beta) \forall n \geq 0\}.$$

Furthermore,

- (i) *The map $t \mapsto b(t, \beta)$ is right-continuous everywhere in $[0, 1)$ with respect to the order topology in $\{0, 1\}^{\mathbb{N}}$;*
- (ii) *If $b(t_0, \beta)$ does not end with 0^{∞} , then the map $t \mapsto b(t, \beta)$ is continuous at t_0 ;*
- (iii) *If $b(t_0, \beta) = b_1 \dots b_m 0^{\infty}$ with $b_m = 1$, then $b(t, \beta) \nearrow b_1 \dots b_m^- \delta(\beta)$ as $t \nearrow t_0$.*

Recall that the survivor set $K_{\beta}(t)$ consists of all $x \in [0, 1)$ whose orbit $\{T_{\beta}^n(x) : n \geq 0\}$ avoids the hole $(0, t)$. To describe the dimension of $K_{\beta}(t)$ we introduce the topological entropy of a symbolic set. For a subset $X \subset \{0, 1\}^{\mathbb{N}}$ its *topological entropy* $h_{top}(X)$ is defined by

$$h_{top}(X) := \liminf_{n \rightarrow \infty} \frac{\log \#B_n(X)}{n},$$

where $\#B_n(X)$ denotes the number of all length n words occurring in sequences of X . The following result for the Hausdorff dimension of $K_{\beta}(t)$ can be essentially deduced from Raith [28] (see also [20]).

Lemma 2.11. *Given $\beta \in (1, 2]$ and $t \in [0, 1)$, the Hausdorff dimension of $K_{\beta}(t)$ is given by*

$$\dim_H K_{\beta}(t) = \frac{h_{top}(\mathbf{K}_{\beta}(t))}{\log \beta},$$

where

$$\mathbf{K}_{\beta}(t) = \{(d_i) \in \{0, 1\}^{\mathbb{N}} : b(t, \beta) \preccurlyeq \sigma^n((d_i)) \prec \delta(\beta) \forall n \geq 0\}.$$

In order to determine the critical value $\tau(\beta)$ for β inside any Farey interval J^s , we first need to develop some properties of the substitution operator \bullet from (1.3). We do this in the next section.

3. SUBSTITUTION OF LYNDON WORDS

In this section we give an equivalent definition of the substitution operator in Ω_L^* introduced in (1.3), and prove that Ω_L^* forms a semi-group under this substitution operator. This will play a crucial role in the rest of the paper.

3.1. An equivalent definition of the substitution. Given a Lyndon word $\mathbf{s} \in \Omega_L^*$ with $\mathbf{a} = \mathbb{L}(\mathbf{s})$, we construct a directed graph $G = (V, E)$ as in Figure 2. The directed graph G has two starting vertices ‘Start-0’ and ‘Start-1’. The directed edges in the graph G take labels from $\{0, 1\}$, and the vertices in G take labels from $\{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$. Denote by \mathcal{L}^E the edge labeling and by \mathcal{L}^V the vertex labeling. Then for each directed edge $e \in E(G)$ we have $\mathcal{L}^E(e) \in \{0, 1\}$, and for each vertex $v \in V(G)$ we have $\mathcal{L}^V(v) \in \{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$. The labeling

maps \mathcal{L}^E and \mathcal{L}^V naturally induce the maps on the infinite edge paths and infinite vertex paths in G , respectively. For example, for an infinite edge path $e_1e_2\dots$ we have

$$\mathcal{L}^E(e_1e_2\dots) = \mathcal{L}^E(e_1)\mathcal{L}^E(e_2)\dots \in \{0, 1\}^{\mathbb{N}}.$$

Here we call $e_1e_2\dots$ an *infinite edge path* in G , if the initial vertex of e_1 is one of the starting vertices and for any $i \geq 1$ the *terminal vertex* $t(e_i)$ equals the *initial vertex* $i(e_{i+1})$. Similarly, for an infinite vertex path $v = v_1v_2\dots$ we have

$$\mathcal{L}^V(v_1v_2\dots) = \mathcal{L}^V(v_1)\mathcal{L}^V(v_2)\dots \in \{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}^{\mathbb{N}},$$

where we call $v_1v_2\dots$ an *infinite vertex path* in G , if v_1 is one of the starting vertices and for any $i \geq 1$ there exists a directed edge $e \in E(G)$ such that $i(e) = v_i$ and $t(e) = v_{i+1}$.




FIGURE 2. The directed graph $G = (V, E)$ with the edge labels from $\{0, 1\}$ and vertex labels from $\{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$, where $\mathbf{s} \in \Omega_L^*$ and $\mathbf{a} = \mathbb{L}(\mathbf{s})$.

Let $X_E = X_E(G)$ be the *edge shift* consisting of all labeling of infinite edge paths in G , i.e.,

$$X_E := \{\mathcal{L}^E(e_1e_2\dots) : e_1e_2\dots \text{ is an infinite edge path in } G\}.$$

One can verify easily that $X_E = \{0, 1\}^{\mathbb{N}}$. Also, let $X_V = X_V(G)$ be the *vertex shift* which consists of all labeling of infinite vertex paths in G , i.e.,

$$X_V := \{\mathcal{L}^V(v_1v_2\dots) : v_1v_2\dots \text{ is an infinite vertex path in } G\}.$$

Then any sequence in X_V is an infinite concatenation of words from $\{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$. Observe that the edge shift X_E is *right-resolving*, which means that out-going edges from the same vertex have different labels (cf. [23]). Moreover, different vertices have different labels. So for each $(d_i) \in X_E$ there is a unique infinite edge path $e_1e_2\dots$ in G such that $d_1d_2\dots = \mathcal{L}^E(e_1e_2\dots)$.

Definition 3.1. The substitution map $\Phi_{\mathbf{s}}$ from X_E to X_V is defined by

$$\Phi_{\mathbf{s}} : X_E \rightarrow X_V; \quad \mathcal{L}^E(e_1e_2\dots) \mapsto \mathcal{L}^V(t(e_1)t(e_2)\dots),$$

where $t(e_i)$ denotes the terminal vertex of the directed edge e_i .

We can extend the substitution map $\Phi_{\mathbf{s}}$ to a map from $B_*(X_E)$ to $B_*(X_V)$ by

$$(3.1) \quad \Phi_{\mathbf{s}} : B_*(X_E) \rightarrow B_*(X_V); \quad \mathcal{L}^E(e_1\dots e_n) \mapsto \mathcal{L}^V(t(e_1)\dots t(e_n)),$$

where $B_*(X_E)$ consists of all labelings of finite edge paths in G , and $B_*(X_V)$ consists of all labelings of finite vertex paths in G . So, by (1.3) and (3.1) it follows that for any two words $\mathbf{s} \in \Omega_L^*$ and $\mathbf{r} \in \{0, 1\}^*$ we have

$$(3.2) \quad \mathbf{s} \bullet \mathbf{r} = \Phi_{\mathbf{s}}(\mathbf{r}).$$

Example 3.2. Let $\mathbf{s} = 01$ and $\mathbf{r} = 001011$. Then $\mathbf{s} \in \Omega_F^*$ and $\mathbf{r} \in \Omega_L^* \setminus \Omega_F^*$. Furthermore, $\mathbf{s}_- = 00$, $\mathbf{a} = \mathbb{L}(\mathbf{s}) = 10$, $\mathbf{a}^+ = 11$. So by the definition of $\Phi_{\mathbf{s}}$ it follows that

$$\begin{aligned} \Phi_{\mathbf{s}}(\mathbf{r}) &= \Phi_{\mathbf{s}}(001011) = \mathbf{s}_- \mathbf{a} \mathbf{a}^+ \mathbf{s}_- \mathbf{a}^+ \mathbf{s} = 001011001101, \\ \Phi_{\mathbf{s}}(\mathbf{r}^-) &= \Phi_{\mathbf{s}}(001010) = \mathbf{s}_- \mathbf{a} \mathbf{a}^+ \mathbf{s}_- \mathbf{a}^+ \mathbf{s}_- = 001011001100. \end{aligned}$$

Observe that $\Phi_{\mathbf{s}}(\mathbf{r}^-) = \Phi_{\mathbf{s}}(\mathbf{r})^-$. By Definition 1.1 one can check that $\Phi_{\mathbf{s}}(\mathbf{r}) \in \Omega_L^*$. Furthermore,

$$\Phi_{\mathbf{s}}(\mathbb{L}(\mathbf{r})) = \Phi_{\mathbf{s}}(110010) = \mathbf{a}^+ \mathbf{s} \mathbf{s}_- \mathbf{a} \mathbf{a}^+ \mathbf{s}_- = 110100101100 = \mathbb{L}(\Phi_{\mathbf{s}}(\mathbf{r})),$$

and

$$\Phi_{\mathbf{s}}(\mathbf{r}^\infty) = \Phi_{\mathbf{s}}((001011)^\infty) = (\mathbf{s}_- \mathbf{a} \mathbf{a}^+ \mathbf{s}_- \mathbf{a}^+ \mathbf{s})^\infty = (001011001101)^\infty = \Phi_{\mathbf{s}}(\mathbf{r})^\infty.$$

3.2. Properties of the substitution. Motivated by Examples 1.2 and 3.2 we study the properties of the substitution $\Phi_{\mathbf{s}}$. We will show that Ω_L^* forms a semi-group under the substitution operator defined in Definition 3.1. First we prove the monotonicity of $\Phi_{\mathbf{s}}$.

Lemma 3.3. *Let $\mathbf{s} \in \Omega_L^*$. Then the map $\Phi_{\mathbf{s}}$ is strictly increasing in $X_E = \{0, 1\}^{\mathbb{N}}$.*

Proof. Let (d_i) and (d'_i) be two sequences in X_E , and let $(e_i), (e'_i)$ be their corresponding edge paths; thus, $(d_i) = \mathcal{L}^E((e_i))$ and $(d'_i) = \mathcal{L}^E((e'_i))$. Suppose $(d_i) \prec (d'_i)$. Then there exists $k \in \mathbb{N}$ such that $d_1 \dots d_{k-1} = d'_1 \dots d'_{k-1}$ and $d_k < d'_k$. If $k = 1$, then $d_1 = 0$ and $d'_1 = 1$. So, $\mathcal{L}^V(t(e_1)) = \mathbf{s}_-$ and $\mathcal{L}^V(t(e'_1)) = \mathbf{a}^+$. By Definition 3.1 it follows that $\Phi_{\mathbf{s}}((d_i)) \prec \Phi_{\mathbf{s}}((d'_i))$.

If $k > 1$, then $e_1 \dots e_{k-1} = e'_1 \dots e'_{k-1}$, which implies that the initial vertices of e_k and e'_k coincide. Since $d_k < d'_k$, by the definition of \mathcal{L}^V it follows that (see Figure 2)

$$\mathcal{L}^V(t(e_k)) \prec \mathcal{L}^V(t(e'_k)).$$

By Definition 3.1 we also have $\Phi_{\mathbf{s}}((d_i)) \prec \Phi_{\mathbf{s}}((d'_i))$. This completes the proof. \square

Lemma 3.4. *Let $\mathbf{s} \in \Omega_L^*$. Then for any word $\mathbf{d} = d_1 \dots d_k \in B_*(X_E)$ with $k \geq 2$, we have*

$$\begin{cases} \Phi_{\mathbf{s}}(\mathbf{d}^-) = \Phi_{\mathbf{s}}(\mathbf{d})^- & \text{if } d_k = 1, \\ \Phi_{\mathbf{s}}(\mathbf{d}^+) = \Phi_{\mathbf{s}}(\mathbf{d})^+ & \text{if } d_k = 0. \end{cases}$$

Proof. Since $\mathbf{d} = d_1 \dots d_k \in B_*(X_E)$, there exists a unique finite edge path $e_1 \dots e_k$ such that $\mathcal{L}^E(e_1 \dots e_k) = \mathbf{d}$. If $d_k = 1$, then $\mathbf{d}^- = d_1 \dots d_{k-1}0$ can be represented by a unique finite edge path $e'_1 \dots e'_k$ with $e'_1 \dots e'_{k-1} = e_1 \dots e_{k-1}$. By the definition of \mathcal{L}^V it follows that $\mathcal{L}^V(t(e'_k)) = \mathcal{L}^V(t(e_k))^-$. Therefore, by Definition 3.1 it follows that

$$\begin{aligned} \Phi_{\mathbf{s}}(\mathbf{d}^-) &= \Phi_{\mathbf{s}}(\mathcal{L}^E(e'_1 \dots e'_k)) = \Phi_{\mathbf{s}}(\mathcal{L}^E(e_1 \dots e_{k-1} e'_k)) \\ &= \mathcal{L}^V(t(e_1) \dots t(e_{k-1}) t(e'_k)) \\ &= \mathcal{L}^V(t(e_1) \dots t(e_k))^- = \Phi_{\mathbf{s}}(\mathbf{d})^-. \end{aligned}$$

This proves the first equality of the lemma. The second equality follows analogously. \square

Lemma 3.5. *Let $\mathbf{s}, \mathbf{r} \in \Omega_L^*$. Then $\Phi_{\mathbf{s}}(\mathbf{r}^\infty) = \Phi_{\mathbf{s}}(\mathbf{r})^\infty$ and $\Phi_{\mathbf{s}}(\mathbb{L}(\mathbf{r})^\infty) = \Phi_{\mathbf{s}}(\mathbb{L}(\mathbf{r}))^\infty$.*

Proof. Let $\mathbf{r} = r_1 \dots r_\ell \in \Omega_L^*$. Then $r_1 = 0$ and $r_\ell = 1$. Let $e_1 \dots e_\ell$ be the unique edge path representing \mathbf{r} , i.e., $\mathbf{r} = \mathcal{L}^E(e_1 \dots e_\ell)$. Since $r_\ell r_1 = 10$, by the definition of \mathcal{L}^E it follows that the edge path representing $r_1 \dots r_\ell r_1$ ends at the vertex with the label $\mathbf{s}_- = \mathcal{L}^V(r_1)$ (see Figure 2). Since the vertices of G have different labels, by the definition of $\Phi_{\mathbf{s}}$ it follows that $\Phi_{\mathbf{s}}(\mathbf{r}^\infty) = \Phi_{\mathbf{s}}(\mathbf{r})^\infty$. The second statement follows analogously. \square

Recall the operator \bullet from (3.2). In the following we prove Proposition 1.3 by showing that Ω_L^* is closed under \bullet and that \bullet is associative. The proof will be split into a sequence of lemmas. First we prove that Ω_L^* is closed under \bullet .

Lemma 3.6. *For any $\mathbf{s}, \mathbf{r} \in \Omega_L^*$ we have $\mathbf{s} \bullet \mathbf{r} \in \Omega_L^*$.*

Proof. Let $\mathbf{s} = s_1 \dots s_m \in \Omega_L^*$ and $\mathbf{a} = \mathbb{L}(\mathbf{s})$. Then there exists $j \in \{1, \dots, m-1\}$ such that

$$(3.3) \quad \mathbf{a} = s_{j+1} \dots s_m s_1 \dots s_j.$$

Let $\mathbf{r} = r_1 \dots r_\ell \in \Omega_L^*$. Then we can write $\mathbf{s} \bullet \mathbf{r} = \Phi_{\mathbf{s}}(\mathbf{r}) = b_1 \dots b_{m\ell}$. Furthermore, there exists a finite edge path $e_1 \dots e_\ell$ representing \mathbf{r} such that

$$\Phi_{\mathbf{s}}(\mathbf{r}) = \mathcal{L}^V(t(e_1) \dots t(e_\ell)) =: \mathbf{b}_1 \dots \mathbf{b}_\ell,$$

where each block $\mathbf{b}_i \in \{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$. Note that $\mathbf{b}_1 = \mathbf{s}_-$ since the block \mathbf{r} begins with $r_1 = 0$. By Definition 1.1 it suffices to prove

$$(3.4) \quad b_{i+1} \dots b_{m\ell} \succ b_1 \dots b_{m\ell-i} \quad \text{for any } 0 < i < m\ell.$$

We split the proof of (3.4) into two cases.

Case I. $i = km$ for some $k \in \{1, 2, \dots, \ell-1\}$. Then $b_{i+1} \dots b_{m\ell} = \mathbf{b}_{k+1} \dots \mathbf{b}_\ell$. Since \mathbf{r} is a Lyndon word, we have $r_{k+1} \dots r_\ell \succ r_1 \dots r_{\ell-k}$. So, (3.4) follows directly by Lemma 3.3.

Case II. $i = km + p$ for some $k \in \{0, 1, \dots, \ell-1\}$ and $p \in \{1, \dots, m-1\}$. Then $b_{i+1} \dots b_{m\ell} = b_{i+1} \dots b_{i+m-p} \mathbf{b}_{k+2} \dots \mathbf{b}_\ell$. In the following we prove (3.4) by considering the four possible choices of $\mathbf{b}_{k+1} \in \{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$. If $\mathbf{b}_{k+1} = \mathbf{s}$, then by using that $\mathbf{s} \in \Omega_L^*$ we conclude that

$$b_{i+1} \dots b_{i+m-p} = s_{p+1} \dots s_m \succ s_1 \dots s_{m-p} = b_1 \dots b_{m-p},$$

proving (3.4). Similarly, if $\mathbf{b}_{k+1} = \mathbf{a}^+$, then by (3.3) one can also prove that $b_{i+1} \dots b_{i+m-p} \succ b_1 \dots b_{m-p}$. Now we assume $\mathbf{b}_{k+1} = \mathbf{s}_-$. Then by using $\mathbf{s} \in \Omega_L^*$ it follows that

$$(3.5) \quad b_{i+1} \dots b_{i+m-p} = s_{p+1} \dots s_m^- \succ s_1 \dots s_{m-p} = b_1 \dots b_{m-p}.$$

Observe that the word \mathbf{s}_- can only be followed by \mathbf{a} or \mathbf{a}^+ in G (see Figure 2). So $\mathbf{b}_{k+2} \in \{\mathbf{a}, \mathbf{a}^+\}$. Since $\mathbf{a} = \mathbb{L}(\mathbf{s})$, we obtain that

$$(3.6) \quad b_{i+m-p+1} \dots b_{i+m} = a_1 \dots a_p \succ s_{m-p+1} \dots s_m \succ s_{m-p+1} \dots s_m^- = b_{m-p+1} \dots b_m.$$

Thus, by (3.5) and (3.6) we conclude that $b_{i+1} \dots b_{i+m} \succ b_1 \dots b_m$, proving (3.4). Finally, suppose $\mathbf{b}_{k+1} = \mathbf{a}$. Note that the word \mathbf{a} can only be followed by \mathbf{a} or \mathbf{a}^+ in G . Then by (3.3) and using $\mathbf{s} \in \Omega_L^*$ we have

$$b_{i+1} \dots b_{i+m} \succ s_1 \dots s_m \succ b_1 \dots b_m.$$

This completes the proof. \square

To prove that \bullet is associative we need the following result, which says that the two operators \bullet and \mathbb{L} are commute.

Lemma 3.7. *For any $\mathbf{s}, \mathbf{r} \in \Omega_L^*$ we have $\mathbb{L}(\mathbf{s} \bullet \mathbf{r}) = \mathbf{s} \bullet \mathbb{L}(\mathbf{r})$.*

Proof. The proof is similar to that of Lemma 3.6. Let $\mathbf{r} = r_1 \dots r_\ell \in \Omega_L^*$. First we claim that $\mathbf{s} \bullet \mathbb{L}(\mathbf{r})$ is a cyclic permutation of $\mathbf{s} \bullet \mathbf{r}$. Note that $\mathbb{L}(\mathbf{r}) = r_{j+1} \dots r_\ell r_1 \dots r_j$ for some $1 < j < \ell$. Then $r_j = 0$ and $r_{j+1} = 1$. By the definition of $\Phi_{\mathbf{s}}$ it follows that

$$(3.7) \quad \mathbf{s} \bullet \mathbf{r} = \Phi_{\mathbf{s}}(r_1 \dots r_\ell) = \Phi_{\mathbf{s}}(r_1 \dots r_j) \Phi_{\mathbf{s}}(r_{j+1} \dots r_\ell).$$

On the other hand, since $\mathbf{r} = r_1 \dots r_\ell \in \Omega_L^*$, we have $r_\ell = 1$ and $r_1 = 0$. So, by the definition of $\Phi_{\mathbf{s}}$ we obtain that

$$\mathbf{s} \bullet \mathbb{L}(\mathbf{r}) = \Phi_{\mathbf{s}}(r_{j+1} \dots r_\ell r_1 \dots r_j) = \Phi_{\mathbf{s}}(r_{j+1} \dots r_\ell) \Phi_{\mathbf{s}}(r_1 \dots r_j).$$

This, together with (3.7), proves that $\mathbf{s} \bullet \mathbb{L}(\mathbf{r})$ is indeed a cyclic permutation of $\mathbf{s} \bullet \mathbf{r}$. So, in the following it suffices to prove that $\mathbf{s} \bullet \mathbb{L}(\mathbf{r})$ is the lexicographically largest cyclic permutation of itself.

Write $\mathbf{s} = s_1 \dots s_m \in \Omega_L^*$ with $\mathbf{a} = \mathbb{L}(\mathbf{s}) = a_1 \dots a_m$, and write $\mathbb{L}(\mathbf{r}) = c_1 \dots c_\ell$. Then by Lemma 2.7 it follows that

$$(3.8) \quad \begin{aligned} a_{i+1} \dots a_m &\prec a_1 \dots a_{m-i} \quad \forall 0 < i < m; \\ c_{i+1} \dots c_\ell &\prec c_1 \dots c_{\ell-i} \quad \forall 0 < i < \ell. \end{aligned}$$

Write $\mathbf{s} \bullet \mathbb{L}(\mathbf{r}) = \mathbf{b}_1 \dots \mathbf{b}_\ell = b_1 \dots b_{m\ell}$, where each $\mathbf{b}_i \in \{\mathbf{s}_-, \mathbf{s}, \mathbf{a}, \mathbf{a}^+\}$. Then it suffices to prove that

$$(3.9) \quad b_{i+1} \dots b_{m\ell} \prec b_1 \dots b_{m\ell-i} \quad \forall 0 < i < m\ell.$$

Since $\mathbb{L}(\mathbf{r})$ has a prefix $c_1 = 1$, we see that $b_1 \dots b_m = \mathbf{b}_1 = \mathbf{a}^+$. So, by using (3.8) and the same argument as in the proof of Lemma 3.6 we can prove (3.9). \square

The next lemma will be used in the proof of Lemma 5.3 and Proposition 6.2.

Lemma 3.8. *Let $\mathbf{s} \in \Omega_L^*$ and take two sequences $(c_i), (d_i) \in \{0, 1\}^{\mathbb{N}}$.*

(i) *If $d_1 = 1$, then*

$$\sigma^n((c_i)) \prec (d_i) \quad \forall n \geq 0 \quad \implies \quad \sigma^n(\Phi_{\mathbf{s}}((c_i))) \prec \Phi_{\mathbf{s}}((d_i)) \quad \forall n \geq 0.$$

(ii) *If $d_1 = 0$, then*

$$\sigma^n((c_i)) \succ (d_i) \quad \forall n \geq 0 \quad \implies \quad \sigma^n(\Phi_{\mathbf{s}}((c_i))) \succ \Phi_{\mathbf{s}}((d_i)) \quad \forall n \geq 0.$$

Proof. First we prove (i). Suppose $d_1 = 1$ and $\sigma^n((c_i)) \prec (d_i)$ for all $n \geq 0$. Then $\Phi_{\mathbf{s}}((d_i))$ begins with $\mathbb{L}(\mathbf{s})^+$. If $n \equiv 0 \pmod{|\mathbf{s}|}$, then by Lemma 3.3 it follows that $\sigma^n(\Phi_{\mathbf{s}}((c_i))) \prec \Phi_{\mathbf{s}}((d_i))$. If $n \not\equiv 0 \pmod{|\mathbf{s}|}$, then by using $\Phi_{\mathbf{s}}(d_1) = \mathbb{L}(\mathbf{s})^+$ and the same argument as in the proof of Lemma 3.7 one can verify that $\sigma^n(\Phi_{\mathbf{s}}((c_i))) \prec \Phi_{\mathbf{s}}((d_i))$.

The proof of (ii) is similar. Suppose $d_1 = 0$ and $\sigma^n((c_i)) \succ (d_i)$ for all $n \geq 0$. Then $\Phi_{\mathbf{s}}((d_i))$ begins with \mathbf{s}_- . If $n \equiv 0 \pmod{|\mathbf{s}|}$, then by Lemma 3.3 it follows that $\sigma^n(\Phi_{\mathbf{s}}((c_i))) \succ \Phi_{\mathbf{s}}((d_i))$. If $n \not\equiv 0 \pmod{|\mathbf{s}|}$, then by using $\Phi_{\mathbf{s}}(d_1) = \mathbf{s}_-$ and the same argument as in the proof of Lemma 3.6 one can verify that $\sigma^n(\Phi_{\mathbf{s}}((c_i))) \succ \Phi_{\mathbf{s}}((d_i))$. \square

Finally, we show that \bullet is associative.

Lemma 3.9. *For any three words $\mathbf{r}, \mathbf{s}, \mathbf{t} \in \Omega_L^*$ we have $(\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t} = \mathbf{r} \bullet (\mathbf{s} \bullet \mathbf{t})$.*

Proof. Let $\mathbf{r} = r_1 \dots r_m, \mathbf{s} = s_1 \dots s_n$ and $\mathbf{t} = t_1 \dots t_\ell$. Then we can write $(\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t}$ as

$$(3.10) \quad (\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t} = \mathbf{B}_1 \mathbf{B}_2 \dots \mathbf{B}_\ell,$$

where each $\mathbf{B}_i \in \{(\mathbf{r} \bullet \mathbf{s})^-, \mathbf{r} \bullet \mathbf{s}, \mathbb{L}(\mathbf{r} \bullet \mathbf{s}), \mathbb{L}(\mathbf{r} \bullet \mathbf{s})^+\}$. Since $t_1 = 0$, we have $\mathbf{B}_1 = (\mathbf{r} \bullet \mathbf{s})^-$. Furthermore, by the definition of $\Phi_{\mathbf{r} \bullet \mathbf{s}}$ it follows that for $1 < i \leq \ell$,

$$(3.11) \quad \mathbf{B}_i = \begin{cases} \mathbb{L}(\mathbf{r} \bullet \mathbf{s}) & \text{if } t_{i-1}t_i = 00, \\ \mathbb{L}(\mathbf{r} \bullet \mathbf{s})^+ & \text{if } t_{i-1}t_i = 01, \\ (\mathbf{r} \bullet \mathbf{s})^- & \text{if } t_{i-1}t_i = 10, \\ \mathbf{r} \bullet \mathbf{s} & \text{if } t_{i-1}t_i = 11. \end{cases}$$

On the other hand, we write

$$\mathbf{s} \bullet \mathbf{t} = \mathbf{b}_1 \mathbf{b}_2 \dots \mathbf{b}_\ell,$$

where each $\mathbf{b}_i \in \{\mathbf{s}_-, \mathbf{s}, \mathbb{L}(\mathbf{s}), \mathbb{L}(\mathbf{s})^+\}$. Similar to (3.11) we also have by the definition of $\Phi_{\mathbf{s}}$ that for $1 < i \leq \ell$,

$$(3.12) \quad \mathbf{b}_i = \begin{cases} \mathbb{L}(\mathbf{s}) & \text{if } t_{i-1}t_i = 00, \\ \mathbb{L}(\mathbf{s})^+ & \text{if } t_{i-1}t_i = 01, \\ \mathbf{s}_- & \text{if } t_{i-1}t_i = 10, \\ \mathbf{s} & \text{if } t_{i-1}t_i = 11. \end{cases}$$

Then

$$(3.13) \quad \mathbf{r} \bullet (\mathbf{s} \bullet \mathbf{t}) = \Phi_{\mathbf{r}}(\mathbf{b}_1 \mathbf{b}_2 \dots \mathbf{b}_\ell) = \Phi_{\mathbf{r}}(\mathbf{b}_1) \Phi_{\mathbf{r}}^{\mathbf{b}_1}(\mathbf{b}_2) \Phi_{\mathbf{r}}^{\mathbf{b}_2}(\mathbf{b}_3) \dots \Phi_{\mathbf{r}}^{\mathbf{b}_{\ell-1}}(\mathbf{b}_\ell),$$

where $\Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i)$ is the labeling of the vertex path of \mathbf{b}_i with the starting vertex determined by the terminal vertex of the vertex path representing \mathbf{b}_{i-1} . Let $t(\mathbf{b}_{i-1})$ be the last digit of \mathbf{b}_{i-1} and let $i(\mathbf{b}_i)$ be the first digit of \mathbf{b}_i . Then by the definition of $\Phi_{\mathbf{r}}$ it follows that

$$(3.14) \quad \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i) = \Phi_{\mathbf{r}}(\mathbf{b}_i) \iff t(\mathbf{b}_{i-1}) \neq i(\mathbf{b}_i).$$

Comparing (3.10) and (3.13) it suffices to prove that

$$\mathbf{B}_1 = \Phi_{\mathbf{r}}(\mathbf{b}_1) \quad \text{and} \quad \mathbf{B}_i = \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i) \quad \text{for any } 2 \leq i \leq \ell.$$

Since \mathbf{t} is a Lyndon word, $t_1 = 0$. This implies that $\mathbf{b}_1 = \mathbf{s}_-$, so Lemma 3.4 yields that

$$\mathbf{B}_1 = (\mathbf{r} \bullet \mathbf{s})^- = \Phi_{\mathbf{r}}(\mathbf{s})^- = \Phi_{\mathbf{r}}(\mathbf{s}_-) = \Phi_{\mathbf{r}}(\mathbf{b}_1).$$

Now suppose $2 \leq i \leq \ell$, and we prove $\mathbf{B}_i = \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i)$ by considering the following four cases.

Case I. $t_{i-1}t_i = 00$. Then by (3.11) and (3.12) we have $\mathbf{B}_i = \mathbb{L}(\mathbf{r} \bullet \mathbf{s})$ and $\mathbf{b}_i = \mathbb{L}(\mathbf{s})$. Observe by (3.12) that in this case $t(\mathbf{b}_{i-1}) = 0$ and $i(\mathbf{b}_i) = 1$. So by (3.14) and Lemma 3.7 it follows that

$$\mathbf{B}_i = \mathbb{L}(\mathbf{r} \bullet \mathbf{s}) = \mathbf{r} \bullet \mathbb{L}(\mathbf{s}) = \mathbf{r} \bullet \mathbf{b}_i = \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i).$$

Case II. $t_{i-1}t_i = 01$. Then by (3.11) and (3.12) we have $\mathbf{B}_i = \mathbb{L}(\mathbf{r} \bullet \mathbf{s})^+$ and $\mathbf{b}_i = \mathbb{L}(\mathbf{s})^+$. Note by (3.12) that $t(\mathbf{b}_{i-1}) = 0$ and $i(\mathbf{b}_i) = 1$. So by (3.14), Lemmas 3.4 and 3.7 we still have

$$\mathbf{B}_i = \mathbb{L}(\mathbf{r} \bullet \mathbf{s})^+ = (\mathbf{r} \bullet \mathbb{L}(\mathbf{s}))^+ = \mathbf{r} \bullet (\mathbb{L}(\mathbf{s})^+) = \mathbf{r} \bullet \mathbf{b}_i = \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i).$$

Case III. $t_{i-1}t_i = 10$. Then by (3.11) and (3.12) we have $\mathbf{B}_i = (\mathbf{r} \bullet \mathbf{s})^-$ and $\mathbf{b}_i = \mathbf{s}_-$. Note by (3.12) that $t(\mathbf{b}_{i-1}) = 1$ and $i(\mathbf{b}_i) = 0$. So by (3.14) and Lemma 3.4 we obtain that

$$\mathbf{B}_i = (\mathbf{r} \bullet \mathbf{s})^- = \mathbf{r} \bullet (\mathbf{s}_-) = \mathbf{r} \bullet \mathbf{b}_i = \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i).$$

Case IV. $t_{i-1}t_i = 11$. Then by (3.11) and (3.12) we have $\mathbf{B}_i = \mathbf{r} \bullet \mathbf{s}$ and $\mathbf{b}_i = \mathbf{s}$. Note by (3.12) that $t(\mathbf{b}_{i-1}) = 1$ and $i(\mathbf{b}_i) = 0$. So by (3.14) it follows that

$$\mathbf{B}_i = \mathbf{r} \bullet \mathbf{s} = \mathbf{r} \bullet \mathbf{b}_i = \Phi_{\mathbf{r}}^{\mathbf{b}_i}(\mathbf{b}_i)$$

as desired.

Therefore, by Cases I-IV, (3.10) and (3.13) we conclude that $(\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t} = \mathbf{r} \bullet (\mathbf{s} \bullet \mathbf{t})$. \square

Remark 3.10. The proof of Lemma 3.9 suggests that for any $\mathbf{r}, \mathbf{s} \in \Omega_L^*$ and for any $\mathbf{t} \in \{0, 1\}^*$ or $\mathbf{t} \in \{0, 1\}^{\mathbb{N}}$ we also have $(\mathbf{r} \bullet \mathbf{s}) \bullet \mathbf{t} = \mathbf{r} \bullet (\mathbf{s} \bullet \mathbf{t})$.

Proof of Proposition 1.3. The proposition follows by Lemmas 3.6 and 3.9 and Example 1.2, which shows that \bullet is not commutative. \square

4. CRITICAL VALUES IN A BASIC INTERVAL

In this section we will prove Theorem 2. Recall from (1.4) that Λ consists of all words \mathbf{S} of the form

$$\mathbf{S} = \mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \cdots \bullet \mathbf{s}_k, \quad k \in \mathbb{N},$$

where each $\mathbf{s}_i \in \Omega_F^*$. By Proposition 1.3 it follows that $\Lambda \subset \Omega_L^*$, and each $\mathbf{S} \in \Lambda$ can be uniquely represented in the above form. Take $\mathbf{S} \in \Lambda$. As in Definition 1.5 we let $I^{\mathbf{S}} := [\beta_\ell^{\mathbf{S}}, \beta_*^{\mathbf{S}}]$ be the basic interval generated by \mathbf{S} . Then by Lemmas 2.5 and 2.7 it follows that

$$(4.1) \quad \delta(\beta_\ell^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^\infty \quad \text{and} \quad \delta(\beta_*^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^- \mathbb{L}(\mathbf{S})^\infty.$$

To prove Theorem 2 we first prove the following proposition, which provides one of the key tools in this paper and will be used again in Section 6.

Proposition 4.1. *For any $\mathbf{S} \in \Lambda$ the set*

$$(4.2) \quad \Gamma(\mathbf{S}) := \{(x_i) : \mathbf{S}^\infty \preceq \sigma^n((x_i)) \preceq \mathbb{L}(\mathbf{S})^\infty \forall n \geq 0\}$$

is countable.

We point out that the specific form of \mathbf{S} is essential in this proposition: it is not enough to merely assume that \mathbf{S} is a Lyndon word. For instance, take $\mathbf{S} = 0010111 \in \Omega_L^*$. Then $\mathbb{L}(\mathbf{S}) = 1110010$, and it is easy to see that $\Gamma(\mathbf{S}) \supset \{10, 110\}^{\mathbb{N}}$. For $\mathbf{S} = \mathbf{s} \in \Omega_F^*$, Proposition 4.1 follows from the following stronger result, proved in [20, Proposition 4.4].

Lemma 4.2. *For any $\mathbf{s} = s_1 \dots s_m \in \Omega_F^*$, the set*

$$\Gamma(\mathbf{s}) := \{(x_i) : \mathbf{s}_\infty \preceq \sigma^n((x_i)) \preceq \mathbb{L}(\mathbf{s})^\infty \forall n \geq 0\}$$

consists of exactly m different elements.

In order to reduce the technicalities in the proof of Proposition 4.1, we extend the definition from Lemma 2.3 (iii) and define the *conjugate* of any word $\mathbf{S} \in \Lambda$ by

$$\varphi(\mathbf{S}) := \overline{\mathbb{L}(\mathbf{S})}.$$

Lemma 4.3. *The function $\varphi : \Lambda \rightarrow \{0, 1\}^*$; $\mathbf{S} \mapsto \varphi(\mathbf{S})$ is a semigroup automorphism on (Λ, \bullet) . That is, φ maps Λ bijectively onto itself, and*

$$(4.3) \quad \varphi(\mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_k) = \varphi(\mathbf{s}_1) \bullet \cdots \bullet \varphi(\mathbf{s}_k) \quad \forall \mathbf{s}_1, \dots, \mathbf{s}_k \in \Omega_F^*.$$

Furthermore, φ is its own inverse:

$$(4.4) \quad \varphi(\varphi(\mathbf{S})) = \mathbf{S} \quad \forall \mathbf{S} \in \Lambda.$$

Proof. We prove (4.3) and (4.4) simultaneously by induction on the degree k of $\mathbf{S} = \mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_k$. For $k = 1$, (4.3) is trivial and (4.4) follows from Lemma 2.3 (iii). Now suppose (4.3) and (4.4) both hold for any word $\mathbf{S} = \mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_k$ of degree k , and consider $\mathbf{S} \bullet \mathbf{r}$ with $\mathbf{r} \in \Omega_F^*$. Set $\tilde{\mathbf{S}} := \varphi(\mathbf{S})$. We claim first that, for any word $\mathbf{t} \in \{0, 1\}^*$,

$$(4.5) \quad \Phi_{\mathbf{S}}(\bar{\mathbf{t}}) = \overline{\Phi_{\tilde{\mathbf{S}}}(\mathbf{t})}.$$

The expression on the right is well defined since, by (4.3), $\tilde{\mathbf{S}} = \varphi(\mathbf{s}_1) \bullet \cdots \bullet \varphi(\mathbf{s}_k) \in \Omega_L^*$.

Write $\mathbf{A} := \mathbb{L}(\mathbf{S})$ and $\tilde{\mathbf{A}} := \mathbb{L}(\tilde{\mathbf{S}})$. By (4.4), $\varphi(\tilde{\mathbf{S}}) = \mathbf{S}$, so we have

$$(4.6) \quad \overline{\mathbf{A}} = \tilde{\mathbf{S}} \quad \text{and} \quad \tilde{\mathbf{A}} = \overline{\mathbf{S}}.$$

Now note the rotational skew-symmetry in the edge labels of the directed graph in Figure 2: The edge path corresponding to the word $\bar{\mathbf{t}}$ is just the 180° rotation about the center of the figure of the edge path corresponding to \mathbf{t} . On the other hand, replacing the vertex labels $\mathbf{S}, \mathbf{S}^-, \mathbf{A}$ and \mathbf{A}^+ by $\tilde{\mathbf{S}} = \overline{\mathbf{A}}, \tilde{\mathbf{S}}^- = \overline{\mathbf{A}^+}, \tilde{\mathbf{A}} = \overline{\mathbf{S}}$ and $\tilde{\mathbf{A}}^+ = \overline{\mathbf{S}^-}$ respectively and rotating the whole graph by 180° , we get the original graph back except that all the vertex labels and edge labels are reflected. This implies (4.5).

Now we can apply (4.5) to $\mathbf{t} = \overline{\mathbb{L}(\mathbf{r})}$ and obtain:

$$(4.7) \quad \varphi(\mathbf{S}) \bullet \varphi(\mathbf{r}) = \tilde{\mathbf{S}} \bullet \varphi(\mathbf{r}) = \Phi_{\tilde{\mathbf{S}}}(\overline{\mathbb{L}(\mathbf{r})}) = \overline{\Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r}))} = \overline{\mathbf{S} \bullet \mathbb{L}(\mathbf{r})} = \overline{\mathbb{L}(\mathbf{S} \bullet \mathbf{r})} = \varphi(\mathbf{S} \bullet \mathbf{r}).$$

Since $\mathbf{r} \in \Omega_F^*$ was arbitrary, the induction hypothesis (4.3) and (4.7) give

$$(4.8) \quad \varphi(\mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_k \bullet \mathbf{s}_{k+1}) = \varphi(\mathbf{s}_1) \bullet \cdots \bullet \varphi(\mathbf{s}_k) \bullet \varphi(\mathbf{s}_{k+1}) \quad \forall \mathbf{s}_1, \dots, \mathbf{s}_{k+1} \in \Omega_F^*.$$

Thus, (4.3) holds for $k+1$ in place of k . Next, by Lemma 2.3 (iii), $\varphi(\mathbf{s}_i) \in \Omega_F^*$ and $\varphi(\varphi(\mathbf{s}_i)) = \mathbf{s}_i$ for each i , so applying (4.8) with $\varphi(\mathbf{s}_i)$ in place of \mathbf{s}_i for each i , we conclude that $\varphi(\varphi(\mathbf{S}')) = \mathbf{S}'$ for every $\mathbf{S}' \in \Lambda$ of degree $k+1$ also.

Thus, we have proved (4.3) and (4.4) by induction. Now the remaining statements of the lemma follow immediately: by (4.3), Lemma 2.3 (iii) and Proposition 1.3 it follows that $\varphi(\mathbf{S}) \in \Lambda$ for every $\mathbf{S} \in \Lambda$, whereas (4.4) implies that $\varphi : \Lambda \rightarrow \Lambda$ is bijective. Therefore, φ is an automorphism of (Λ, \bullet) . \square

Define

$$\overline{\Gamma(\mathbf{S})} := \{(\overline{x_i}) : (x_i) \in \Gamma(\mathbf{S})\}, \quad \mathbf{S} \in \Lambda.$$

It is clear that $\overline{\Gamma(\mathbf{S})}$ has the same cardinality as $\Gamma(\mathbf{S})$. Observe also by (4.6) that

$$(4.9) \quad \begin{aligned} \overline{\Gamma(\mathbf{S})} &= \{(y_i) : \mathbf{S}^\infty \preceq \sigma^n(\overline{(y_i)}) \preceq \mathbb{L}(\mathbf{S})^\infty \forall n \geq 0\} \\ &= \{(y_i) : \overline{\mathbf{S}}^\infty \succeq \sigma^n((y_i)) \succeq \overline{\mathbb{L}(\mathbf{S})}^\infty \forall n \geq 0\} \\ &= \{(y_i) : \mathbb{L}(\varphi(\mathbf{S}))^\infty \succeq \sigma^n((y_i)) \succeq \varphi(\mathbf{S})^\infty \forall n \geq 0\} \\ &= \Gamma(\varphi(\mathbf{S})). \end{aligned}$$

Proof of Proposition 4.1. For $\mathbf{S} = \mathbf{s} \in \Omega_F^*$ the proposition follows from Lemma 4.2. So it suffices to prove that if $\Gamma(\mathbf{S})$ is countable for an $\mathbf{S} \in \Lambda$, then $\Gamma(\mathbf{S} \bullet \mathbf{r})$ is also countable for any $\mathbf{r} \in \Omega_F^*$.

Fix $\mathbf{S} \in \Lambda$ with $\Gamma(\mathbf{S})$ countable; fix $\mathbf{r} \in \Omega_F^*$, and note that \mathbf{r} begins with 0 and $\mathbb{L}(\mathbf{r})$ begins with 1. Therefore $\mathbf{S} \bullet \mathbf{r}$ begins with \mathbf{S}^- and $\mathbb{L}(\mathbf{S} \bullet \mathbf{r}) = \mathbf{S} \bullet \mathbb{L}(\mathbf{r})$ begins with $\mathbb{L}(\mathbf{S})^+$. So

$$(4.10) \quad (\mathbf{S} \bullet \mathbf{r})^\infty \prec \mathbf{S}^\infty \quad \text{and} \quad \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^\infty \succ \mathbb{L}(\mathbf{S})^\infty.$$

By (4.2) and (4.10) it follows that

$$\Gamma(\mathbf{S}) \subseteq \{(x_i) : (\mathbf{S} \bullet \mathbf{r})^\infty \preceq \sigma^n((x_i)) \preceq \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^\infty \forall n \geq 0\} = \Gamma(\mathbf{S} \bullet \mathbf{r}).$$

Since $\Gamma(\mathbf{S})$ is countable, it suffices to prove that the difference set $\Gamma(\mathbf{S} \bullet \mathbf{r}) \setminus \Gamma(\mathbf{S})$ is countable. By Proposition 2.4 the word \mathbf{r} must be of one of the following four types:

- (I) $\mathbf{r} = 01^p$ for some $p \in \mathbb{N}$;
- (II) $\mathbf{r} = 0^p 1$ for some $p \in \mathbb{N}$;
- (III) $\mathbf{r} = 01^p 01^{p+t_1} \dots 01^{p+t_N} 01^{p+1}$ for some $p \in \mathbb{N}$ and $0t_1 \dots t_N 1 \in \Omega_F^*$.
- (IV) $\mathbf{r} = 0^{p+1} 10^{p+t_1} 1 \dots 0^{p+t_N} 10^p 1$ for some $p \in \mathbb{N}$ and $0t_1 \dots t_N 1 \in \Omega_F^*$;

Since the words in (II) and (IV) are the conjugates of the words in (I) and (III), respectively, it suffices by Lemma 4.3 and the relationship (4.9) to consider cases (I) and (III). Let $\mathbf{A} := \mathbb{L}(\mathbf{S})$.

Case I. $\mathbf{r} = 01^p$ for some $p \in \mathbb{N}$. Note that $\mathbf{S} \bullet \mathbf{r} = \Phi_{\mathbf{S}}(01^p) = \mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1}$ and $\mathbb{L}(\mathbf{S} \bullet \mathbf{r}) = \mathbf{S} \bullet \mathbb{L}(\mathbf{r}) = \Phi_{\mathbf{S}}(1^p 0) = \mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^-$. Then $\Gamma(\mathbf{S} \bullet \mathbf{r})$ consists of all sequences $(x_i) \in \{0, 1\}^{\mathbb{N}}$ satisfying

$$(4.11) \quad (\mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1})^\infty \preceq \sigma^n((x_i)) \preceq (\mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^-)^\infty \quad \forall n \geq 0.$$

Take a sequence $(x_i) \in \Gamma(\mathbf{S} \bullet \mathbf{r}) \setminus \Gamma(\mathbf{S})$. Then by (4.2) and (4.11) it follows that $x_{k+1} \dots x_{k+m} = \mathbf{S}^-$ or \mathbf{A}^+ for some $k \geq 0$. If $x_{k+1} \dots x_{k+m} = \mathbf{S}^-$, then by taking $n = k$ in (4.11) we obtain

$$x_{k+m+1} x_{k+m+2} \dots \succ (\mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^-)^\infty.$$

On the other hand, by taking $n = k + m$ in (4.11) we see that the above inequality is indeed an equality. So, $x_{k+1} x_{k+2} \dots = (\mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1})^\infty$.

If $x_{k+1} \dots x_{k+m} = \mathbf{A}^+$, then by taking $n = k$ in (4.11) we have

$$(4.12) \quad x_{k+m+1} x_{k+m+2} \dots \preceq (\mathbf{S}^{p-1} \mathbf{S}^- \mathbf{A}^+)^\infty.$$

Note by (4.11) that $x_{i+1} \dots x_{i+m} \succ \mathbf{S}^-$ for all $i \geq 0$. So by (4.12) there must exist a $j \in \{k+m, k+2m, \dots, k+pm\}$ such that $x_{j+1} \dots x_{j+m} = \mathbf{S}^-$. Then by the same argument as above we conclude that $x_{j+1} x_{j+2} \dots = (\mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1})^\infty$. So, $\Gamma(\mathbf{S} \bullet \mathbf{r}) \setminus \Gamma(\mathbf{S})$ is at most countable.

Case III. $\mathbf{r} = 01^p 01^{p+t_1} \dots 01^{p+t_N} 01^{p+1}$, where $p \in \mathbb{N}$ and $\hat{\mathbf{r}} := 0t_1 \dots t_N 1 \in \Omega_F^*$. Consider the substitution

$$\eta_p := U_1^p \circ U_0 : 0 \mapsto 01^p; \quad 1 \mapsto 01^{p+1}.$$

Then $\mathbf{r} = \eta_p(\hat{\mathbf{r}})$ as shown in the proof of Proposition 2.4. Note by Lemma 2.3 that $\mathbb{L}(\mathbf{r}) = 1^{p+1} 01^{p+t_1} 01^{p+t_2} \dots 01^{p+t_N} 01^p 0$ and $\mathbb{L}(\hat{\mathbf{r}}) = 1t_1 \dots t_N 0$. Then

$$(4.13) \quad \mathbb{L}(\mathbf{r})^\infty = \sigma((01^{p+1} 01^{p+t_1} \dots 01^{p+t_N} 01^p)^\infty) = \sigma(\eta_p((1t_1 \dots t_N 0)^\infty)) = \sigma(\eta_p(\mathbb{L}(\hat{\mathbf{r}})^\infty)).$$

Claim: If $(x_i) \in \Gamma(\mathbf{S} \bullet \mathbf{r})$ begins with $x_1 \dots x_m = \mathbf{S}^-$, then there exists a unique sequence $(z_i) \in \{0, 1\}^{\mathbb{N}}$ such that $(x_i) = \eta_p(z_1 z_2 \dots)$.

Note that \mathbf{r} begins with $01^p 0$ and $\mathbb{L}(\mathbf{r})$ begins with $1^{p+1} 0$. Then $\mathbf{S} \bullet \mathbf{r}$ begins with $\Phi_{\mathbf{S}}(01^p 0) = \mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^-$ and $\mathbb{L}(\mathbf{S} \bullet \mathbf{r}) = \mathbf{S} \bullet \mathbb{L}(\mathbf{r})$ begins with $\Phi_{\mathbf{S}}(1^{p+1} 0) = \mathbf{A}^+ \mathbf{S}^p \mathbf{S}^-$. Let $(x_i) \in \Gamma(\mathbf{S} \bullet \mathbf{r})$ with $x_1 \dots x_m = \mathbf{S}^-$. Then

$$(4.14) \quad \mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^- \preceq x_{n+1} \dots x_{n+m(p+2)} \preceq \mathbf{A}^+ \mathbf{S}^p \mathbf{S}^- \quad \forall n \geq 0.$$

By taking $n = 0$ in (4.14) it follows that

$$(4.15) \quad x_{m+1} \dots x_{m(p+2)} \succ \mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^-.$$

On the other hand, by taking $n = m$ in (4.14) we have

$$(4.16) \quad x_{m+1} \cdots x_{m(p+3)} \preceq \mathbf{A}^+ \mathbf{S}^p \mathbf{S}^-.$$

By (4.14)–(4.16) it follows that

$$\text{either } x_{m+1} \cdots x_{m(p+2)} = \mathbf{A}^+ \mathbf{S}^{p-1} \mathbf{S}^- \quad \text{or} \quad x_{m+1} \cdots x_{m(p+3)} = \mathbf{A}^+ \mathbf{S}^p \mathbf{S}^-.$$

Observe that in both cases we obtain a block ending with \mathbf{S}^- . Then we can repeat the above argument indefinitely, and conclude that

$$(4.17) \quad (x_i) \in \{\mathbf{S}^- \mathbf{A}^+ \mathbf{S}^{p-1}, \mathbf{S}^- \mathbf{A}^+ \mathbf{S}^p\}^{\mathbb{N}} = \{\Phi_{\mathbf{S}}(\eta_p(0)), \Phi_{\mathbf{S}}(\eta_p(1))\}^{\mathbb{N}}.$$

Since $\eta_p(0) = 01^p$ and $\eta_p(1) = 01^{p+1}$, it follows from (4.17) and the definition of $\Phi_{\mathbf{S}}$ that

$$(4.18) \quad (x_i) = \Phi_{\mathbf{S}}(\eta_p(z_1)) \Phi_{\mathbf{S}}(\eta_p(z_2)) \cdots = \Phi_{\mathbf{S}}(\eta_p(z_1) \eta_p(z_2) \cdots) = \Phi_{\mathbf{S}}(\eta_p(z_1 z_2 \cdots))$$

for some sequence $(z_i) \in \{0, 1\}^{\mathbb{N}}$. The uniqueness of (z_i) follows by the definition of the substitutions η_p and $\Phi_{\mathbf{S}}$. This proves the claim.

Now take a sequence $(x_i) \in \Gamma(\mathbf{S} \bullet \mathbf{r}) \setminus \Gamma(\mathbf{S})$. Then by (4.2) and (4.14) we can find an $n_0 \geq 0$ such that $x_{n_0+1} \cdots x_{n_0+m} = \mathbf{S}^-$ or \mathbf{A}^+ . If $x_{n_0+1} \cdots x_{n_0+m} = \mathbf{A}^+$, then by (4.14) there must exist $n_1 > n_0$ such that $x_{n_1+1} \cdots x_{n_1+m} = \mathbf{S}^-$. So, without loss of generality, we may assume $x_{n_0+1} \cdots x_{n_0+m} = \mathbf{S}^-$. Then by the claim there exists a unique sequence $(z_i) \in \{0, 1\}^{\mathbb{N}}$ such that $x_{n_0+1} x_{n_0+2} \cdots = \Phi_{\mathbf{S}}(\eta_p(z_1 z_2 \cdots)) \in \Gamma(\mathbf{S} \bullet \mathbf{r})$. So, by the definition of $\Gamma(\mathbf{S} \bullet \mathbf{r})$ it follows that

$$(4.19) \quad (\mathbf{S} \bullet \mathbf{r})^\infty \preceq \sigma^n(\Phi_{\mathbf{S}}(\eta_p(z_1 z_2 \cdots))) \preceq \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^\infty \quad \forall n \geq 0$$

Note by $\mathbf{r} = \eta(\hat{\mathbf{r}})$ and Lemma 3.5 that $(\mathbf{S} \bullet \mathbf{r})^\infty = \Phi_{\mathbf{S}}(\mathbf{r}^\infty) = \Phi_{\mathbf{S}}(\eta_p(\hat{\mathbf{r}}^\infty))$. Also, by Lemmas 3.5, 3.7 and (4.13) it follows that $\mathbb{L}(\mathbf{S} \bullet \mathbf{r})^\infty = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^\infty) = \Phi_{\mathbf{S}}(\sigma(\eta_p(\mathbb{L}(\hat{\mathbf{r}})^\infty)))$. So, (4.19) can be rewritten as

$$\Phi_{\mathbf{S}}(\eta_p(\hat{\mathbf{r}}^\infty)) \preceq \sigma^n(\Phi_{\mathbf{S}}(\eta_p(z_1 z_2 \cdots))) \preceq \Phi_{\mathbf{S}}(\sigma(\eta_p(\mathbb{L}(\hat{\mathbf{r}})^\infty))) \quad \forall n \geq 0.$$

By Lemma 3.3 this implies that

$$(4.20) \quad \eta_p(\hat{\mathbf{r}}^\infty) \preceq \sigma^n(\eta_p(z_1 z_2 \cdots)) \preceq \sigma(\eta_p(\mathbb{L}(\hat{\mathbf{r}})^\infty)) \quad \forall n \geq 0.$$

Note $(c_i) \prec (d_i)$ is equivalent to $\eta_p((c_i)) \prec \eta_p((d_i))$. So, by (4.20) and the definition of η_p it follows that

$$\hat{\mathbf{r}}^\infty \preceq \sigma^n(z_1 z_2 \cdots) \preceq \mathbb{L}(\hat{\mathbf{r}})^\infty \quad \forall n \geq 0,$$

which implies that $(z_i) \in \Gamma(\hat{\mathbf{r}})$. Since $\hat{\mathbf{r}} \in \Omega_F^*$, we know that $\Gamma(\hat{\mathbf{r}})$ is finite by Lemma 4.2. Hence there are only countably many choices for the sequence (z_i) , and thus by the claim there are only countably many choices for the tail sequence of (x_i) . Therefore, $\Gamma(\mathbf{S} \bullet \mathbf{r}) \setminus \Gamma(\mathbf{S})$ is at most countable. \square

Recall from Lemma 2.11 the symbolic survivor set $\mathbf{K}_\beta(t)$. To prove Theorem 2 we also recall the following result from [20, Lemma 3.7].

Lemma 4.4. *Let $\beta \in (1, 2]$ and $t \in [0, 1)$. If $\sigma^m(\delta(\beta)) \preceq b(t, \beta)$, then*

$$\mathbf{K}_\beta(t) = \{(d_i) : b(t, \beta) \preceq \sigma^n((d_i)) \preceq (\delta_1(\beta) \cdots \delta_m(\beta)^-)^{\infty} \forall n \geq 0\}.$$

Proof of Theorem 2. That the basic intervals $I^{\mathbf{S}}, \mathbf{S} \in \Lambda$ are pairwise disjoint will be shown in Proposition 5.1 below. In what follows, we fix a basic interval $I^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_*^{\mathbf{S}}]$. Take $\beta \in I^{\mathbf{S}}$, and let $t^* = (\Phi_{\mathbf{S}}(0^\infty))_\beta = (\mathbf{S}^- \mathbf{A}^\infty)_\beta$, where $\mathbf{A} = \mathbb{L}(\mathbf{S}) = A_1 \dots A_m$. Then by (4.1) and Lemma 2.5 it follows that

$$(4.21) \quad \mathbf{A}^\infty = \delta(\beta_\ell^{\mathbf{S}}) \preceq \delta(\beta) \preceq \delta(\beta_*^{\mathbf{S}}) = \mathbf{A}^+ \mathbf{S}^- \mathbf{A}^\infty.$$

Since $\mathbf{S} = \mathbf{s}_1 \bullet \dots \bullet \mathbf{s}_k$ with each $\mathbf{s}_i \in \Omega_F^*$, by Proposition 1.3 we have $\mathbf{S} \in \Omega_L^*$. If $\beta \in (\beta_\ell^{\mathbf{S}}, \beta_*^{\mathbf{S}}]$, then by Lemma 2.7 we have

$$\sigma^n(\mathbf{S}^- \mathbf{A}^\infty) \preceq \mathbf{A}^\infty \prec \delta(\beta) \quad \forall n \geq 0;$$

and by Lemma 2.10 it follows that $\mathbf{S}^- \mathbf{A}^\infty$ is the greedy β -expansion of t^* , i.e., $b(t^*, \beta) = \mathbf{S}^- \mathbf{A}^\infty$. If $\beta = \beta_\ell^{\mathbf{S}}$, then $\sigma^n(\mathbf{S}^- \mathbf{A}^\infty) \preceq \mathbf{A}^\infty = \delta(\beta)$ for all $n \geq 0$; and in this case one can verify that the greedy β -expansion of t^* is given by $b(t^*, \beta) = \mathbf{S}0^\infty$.

First we prove $\tau(\beta) \geq t^*$. Note that $\mathbf{A} = A_1 \dots A_m$. Let $t_N := ((\mathbf{S}^- \mathbf{A}^N A_1 \dots A_j)^\infty)_\beta$, where the index $j \in \{1, \dots, m\}$ satisfies $\mathbf{S} = \mathbb{S}(\mathbf{A}) = A_{j+1} \dots A_m A_1 \dots A_j$. Note by Lemma 2.7 that $A_{i+1} \dots A_m \prec A_1 \dots A_{m-i}$ for any $0 < i < m$. Then by (4.21) one can verify that

$$\begin{aligned} \sigma^n((\mathbf{S}^- \mathbf{A}^N A_1 \dots A_j)^\infty) &= \sigma^n(\mathbf{S}^- \mathbf{A}^{N+1} (A_1 \dots A_j^- \mathbf{A}^{N+1})^\infty) \\ &\prec \mathbf{A}^\infty \preceq \delta(\beta) \quad \forall n \geq 0. \end{aligned}$$

So, $b(t_N, \beta) = (\mathbf{S}^- \mathbf{A}^N A_1 \dots A_j)^\infty$. This implies that any sequence (x_i) constructed by arbitrarily concatenating blocks of the form

$$\mathbf{S}^- \mathbf{A}^k A_1 \dots A_j, \quad k > N$$

satisfies $(\mathbf{S}^- \mathbf{A}^N A_1 \dots A_j)^\infty \preceq \sigma^n((x_i)) \prec \delta(\beta)$ for all $n \geq 0$. So,

$$\{\mathbf{S}^- \mathbf{A}^{N+1} A_1 \dots A_j, \mathbf{S}^- \mathbf{A}^{N+2} A_1 \dots A_j\}^{\mathbb{N}} \subset \mathbf{K}_\beta(t_N).$$

By Lemma 2.11 this implies that $\dim_H K_\beta(t_N) > 0$ for all $N \geq 1$. Thus $\tau(\beta) \geq t_N$ for all $N \geq 1$. Note that $t_N \nearrow t^*$ as $N \rightarrow \infty$. We then conclude that $\tau(\beta) \geq t^*$.

Next we prove $\tau(\beta) \leq t^*$. By (4.21) and Lemma 4.4 it follows that

$$(4.22) \quad \begin{aligned} \mathbf{K}_\beta(t^*) &\subset \{(x_i) : \mathbf{S}^- \mathbf{A}^\infty \preceq \sigma^n((x_i)) \prec \mathbf{A}^+ \mathbf{S}^- \mathbf{A}^\infty \forall n \geq 0\} \\ &= \{(x_i) : \mathbf{S}^- \mathbf{A}^\infty \preceq \sigma^n((x_i)) \preceq \mathbf{A}^\infty \forall n \geq 0\} =: \Gamma. \end{aligned}$$

Note by Proposition 4.1 that $\Gamma(\mathbf{S}) = \{(x_i) : \mathbf{S}^\infty \preceq \sigma^n((x_i)) \preceq \mathbf{A}^\infty \forall n \geq 0\}$ is a countable subset of Γ . Furthermore, any sequence in the difference set $\Gamma \setminus \Gamma(\mathbf{S})$ must end with $\mathbf{S}^- \mathbf{A}^\infty$. So, Γ is also countable. By (4.22) this implies that $\dim_H K_\beta(t^*) = 0$, and thus $\tau(\beta) \leq t^*$. This completes the proof. \square

5. GEOMETRICAL STRUCTURE OF THE BASIC INTERVALS AND BIFURCATION SETS

In this section we will prove Theorem 3. The proof will be split into two subsections. In Subsection 5.1 we demonstrate the tree structure of the Lyndon intervals $J^{\mathbf{S}}, \mathbf{S} \in \Lambda$ and the relative bifurcation sets $E^{\mathbf{S}}, \mathbf{S} \in \Lambda$, from which it follows that the basic intervals $I^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_*^{\mathbf{S}}]$, $\mathbf{S} \in \Lambda$ are pairwise disjoint. We show that each relative bifurcation set $E^{\mathbf{S}}$ has zero box-counting dimension, and the bifurcation set E has zero packing dimension. In Subsection 5.2 we prove that the infinite bifurcation set E_∞ has zero Hausdorff dimension.

5.1. Tree structure of the Lyndon intervals and relative bifurcation sets. Given $\mathbf{S} \in \Lambda$, recall from Definitions 1.5 and 1.8 the basic interval $I^{\mathbf{S}} = [\beta_{\ell}^{\mathbf{S}}, \beta_{*}^{\mathbf{S}}]$ and the Lyndon interval $J^{\mathbf{S}} = [\beta_{\ell}^{\mathbf{S}}, \beta_r^{\mathbf{S}}]$ generated by \mathbf{S} , respectively. Then by Lemmas 2.5 and 2.7 it follows that

$$(5.1) \quad \delta(\beta_{\ell}^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^{\infty}, \quad \delta(\beta_{*}^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^{+}\mathbf{S}^{-}\mathbb{L}(\mathbf{S})^{\infty} \quad \text{and} \quad \delta(\beta_r^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^{+}\mathbf{S}^{\infty}.$$

First we show that the Lyndon intervals $J^{\mathbf{S}}, \mathbf{S} \in \Lambda$ have a tree structure.

Proposition 5.1. *Let $\mathbf{S} \in \Lambda$. Then $I^{\mathbf{S}} \subset J^{\mathbf{S}}$. Furthermore,*

- (i) *for any $\mathbf{r} \in \Omega_F^*$ we have $J^{\mathbf{S}\bullet\mathbf{r}} \subset J^{\mathbf{S}} \setminus I^{\mathbf{S}}$;*
- (ii) *for any two different words $\mathbf{r}, \mathbf{r}' \in \Omega_F^*$ we have $J^{\mathbf{S}\bullet\mathbf{r}} \cap J^{\mathbf{S}\bullet\mathbf{r}'} = \emptyset$.*

Proof. Let $I^{\mathbf{S}} = [\beta_{\ell}^{\mathbf{S}}, \beta_{*}^{\mathbf{S}}]$ and $J^{\mathbf{S}} = [\beta_{\ell}^{\mathbf{S}}, \beta_r^{\mathbf{S}}]$. Then by (5.1) it follows that

$$\delta(\beta_{*}^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^{+}\mathbf{S}^{-}\mathbb{L}(\mathbf{S})^{\infty} \prec \mathbb{L}(\mathbf{S})^{+}\mathbf{S}^{\infty} = \delta(\beta_r^{\mathbf{S}}),$$

which implies $\beta_{*}^{\mathbf{S}} < \beta_r^{\mathbf{S}}$ by Lemma 2.5. So $I^{\mathbf{S}} \subset J^{\mathbf{S}}$.

For (i), let $\mathbf{r} \in \Omega_F^*$. Then \mathbf{r} begins with digit 0 and ends with digit 1. By Lemmas 3.5 and 3.7 this implies that

$$\begin{aligned} \delta(\beta_{\ell}^{\mathbf{S}\bullet\mathbf{r}}) &= \mathbb{L}(\mathbf{S}\bullet\mathbf{r})^{\infty} = (\mathbf{S}\bullet\mathbb{L}(\mathbf{r}))^{\infty} = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^{\infty}) \\ &\succ \Phi_{\mathbf{S}}(10^{\infty}) = \mathbb{L}(\mathbf{S})^{+}\mathbf{S}^{-}\mathbb{L}(\mathbf{S})^{\infty} = \delta(\beta_{*}^{\mathbf{S}}). \end{aligned}$$

So, $\beta_{\ell}^{\mathbf{S}\bullet\mathbf{r}} > \beta_{*}^{\mathbf{S}}$. Furthermore, by Lemmas 3.4, 3.5 and 3.7 it follows that

$$\begin{aligned} \delta(\beta_r^{\mathbf{S}\bullet\mathbf{r}}) &= \mathbb{L}(\mathbf{S}\bullet\mathbf{r})^{+}(\mathbf{S}\bullet\mathbf{r})^{\infty} \\ &= \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^{+})\Phi_{\mathbf{S}}(\mathbf{r}^{\infty}) = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^{+}\mathbf{r}^{\infty}) \\ &\prec \Phi_{\mathbf{S}}(1^{\infty}) = \mathbb{L}(\mathbf{S})^{+}\mathbf{S}^{\infty} = \delta(\beta_r^{\mathbf{S}}). \end{aligned}$$

This proves $\beta_r^{\mathbf{S}\bullet\mathbf{r}} < \beta_r^{\mathbf{S}}$. Hence, $J^{\mathbf{S}\bullet\mathbf{r}} = [\beta_{\ell}^{\mathbf{S}\bullet\mathbf{r}}, \beta_r^{\mathbf{S}\bullet\mathbf{r}}] \subset (\beta_{*}^{\mathbf{S}}, \beta_r^{\mathbf{S}}] = J^{\mathbf{S}} \setminus I^{\mathbf{S}}$.

Next we prove (ii). Let \mathbf{r}, \mathbf{r}' be two different Farey words in Ω_F^* . By Lemma 2.9 (i) it follows that $J^{\mathbf{r}} \cap J^{\mathbf{r}'} = \emptyset$. Write $J^{\mathbf{r}} = [\beta_{\ell}^{\mathbf{r}}, \beta_r^{\mathbf{r}}]$ and $J^{\mathbf{r}'} = [\beta_{\ell}^{\mathbf{r}'}, \beta_r^{\mathbf{r}'}]$. Since $J^{\mathbf{r}}$ and $J^{\mathbf{r}'}$ are disjoint, we may assume $\beta_r^{\mathbf{r}} < \beta_{\ell}^{\mathbf{r}'}$. By (5.1) and Lemma 2.5 it follows that

$$(5.2) \quad \mathbb{L}(\mathbf{r})^{+}\mathbf{r}^{\infty} = \delta(\beta_r^{\mathbf{r}}) \prec \delta(\beta_{\ell}^{\mathbf{r}'}) = \mathbb{L}(\mathbf{r}')^{\infty}.$$

Then by (5.1), (5.2) and Lemma 3.3 we obtain that

$$\begin{aligned} \delta(\beta_r^{\mathbf{S}\bullet\mathbf{r}}) &= \mathbb{L}(\mathbf{S}\bullet\mathbf{r})^{+}(\mathbf{S}\bullet\mathbf{r})^{\infty} = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^{+}\mathbf{r}^{\infty}) \\ &\prec \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r}')^{\infty}) = \mathbb{L}(\mathbf{S}\bullet\mathbf{r}')^{\infty} = \delta(\beta_{\ell}^{\mathbf{S}\bullet\mathbf{r}'}). \end{aligned}$$

It follows that $\beta_r^{\mathbf{S}\bullet\mathbf{r}} < \beta_{\ell}^{\mathbf{S}\bullet\mathbf{r}'}$, and hence $J^{\mathbf{S}\bullet\mathbf{r}} \cap J^{\mathbf{S}\bullet\mathbf{r}'} = \emptyset$. □

Remark 5.2. Proposition 5.1 implies that the Lyndon intervals $J^{\mathbf{S}}, \mathbf{S} \in \Lambda$ have a tree structure. More precisely, we say $J^{\mathbf{R}}$ is an *offspring* of $J^{\mathbf{S}}$ if there exists a word $\mathbf{T} \in \Lambda$ such that $\mathbf{R} = \mathbf{S}\bullet\mathbf{T}$. Then any offspring of $J^{\mathbf{S}}$ is a subset of $J^{\mathbf{S}}$. Furthermore, if $J^{\mathbf{S}'}$ is not an offspring of $J^{\mathbf{S}}$ and $J^{\mathbf{S}}$ is not an offspring of $J^{\mathbf{S}'}$, then Proposition 5.1 implies that $J^{\mathbf{S}'} \cap J^{\mathbf{S}} = \emptyset$. Consequently, the basic intervals $I^{\mathbf{S}}, \mathbf{S} \in \Lambda$ are pairwise disjoint.

Recall from Section 1 the bifurcation set $E = (1, 2] \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{r}}$ and the relative bifurcation sets $E^{\mathbf{S}} = (J^{\mathbf{S}} \setminus I^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S} \bullet \mathbf{r}}$ with $\mathbf{S} \in \Lambda$. Next we will show that E is bijectively mapped to $E^{\mathbf{S}}$ via the map

$$(5.3) \quad \Psi_{\mathbf{S}} : (1, 2] \rightarrow J^{\mathbf{S}} \setminus I^{\mathbf{S}} = (\beta_*^{\mathbf{S}}, \beta_r^{\mathbf{S}}]; \quad \beta \mapsto \delta^{-1} \circ \Phi_{\mathbf{S}} \circ \delta(\beta),$$

where $\delta(\beta)$ is the quasi-greedy β -expansion of 1.

We mention that $\Psi_{\mathbf{S}}$ is not surjective, which somewhat complicates the proof of Proposition 5.4 below. For example, let $\mathbf{S} = 011$. Then $\delta(\beta_*^{\mathbf{S}}) = 111010(110)^\infty$ and $\delta(\beta_r^{\mathbf{S}}) = 111(011)^\infty$. Take $\beta \in (\beta_*^{\mathbf{S}}, \beta_r^{\mathbf{S}}]$ such that $\delta(\beta) = 1110110^\infty$. One can verify that $\beta \notin \Psi_{\mathbf{S}}((1, 2])$, since $\delta(\beta)$ cannot be written as a concatenation of words from $\{\mathbf{S}, \mathbf{S}^-, \mathbb{L}(\mathbf{S}), \mathbb{L}(\mathbf{S})^+\}$.

Lemma 5.3. *For any $\mathbf{S} \in \Lambda$ the map $\Psi_{\mathbf{S}}$ is well-defined and strictly increasing.*

Proof. Let $\mathbf{S} \in \Lambda$ with $\mathbf{A} = \mathbb{L}(\mathbf{S})$. First we show that the map $\Psi_{\mathbf{S}} : (1, 2] \rightarrow J^{\mathbf{S}} \setminus I^{\mathbf{S}} = (\beta_*^{\mathbf{S}}, \beta_r^{\mathbf{S}}]$ is well-defined. Note that

$$(5.4) \quad \delta(\beta_*^{\mathbf{S}}) = \mathbf{A}^+ \mathbf{S}^- \mathbf{A}^\infty = \Phi_{\mathbf{S}}(10^\infty) \quad \text{and} \quad \delta(\beta_r^{\mathbf{S}}) = \mathbf{A}^+ \mathbf{S}^\infty = \Phi_{\mathbf{S}}(1^\infty).$$

Take $\beta \in (1, 2]$. Then $10^\infty \prec \delta(\beta) \preceq 1^\infty$. By Lemma 3.3 and (5.4) it follows that

$$\delta(\beta_*^{\mathbf{S}}) \prec \Phi_{\mathbf{S}}(\delta(\beta)) \preceq \delta(\beta_r^{\mathbf{S}}).$$

Thus, by Lemma 2.5 it suffices to prove that

$$(5.5) \quad \sigma^n(\Phi_{\mathbf{S}}(\delta(\beta))) \preceq \Phi_{\mathbf{S}}(\delta(\beta)) \quad \forall n \geq 0.$$

Note by Lemma 2.5 that $\sigma^n(\delta(\beta)) \preceq \delta(\beta)$ for all $n \geq 0$, and $\delta(\beta)$ begins with digit 1. Thus, (5.5) follows by Lemma 3.8 (i), and we conclude that the map $\Psi_{\mathbf{S}}$ is well-defined.

The monotonicity of $\Psi_{\mathbf{S}} = \delta^{-1} \circ \Phi_{\mathbf{S}} \circ \delta$ follows since both maps δ and $\Phi_{\mathbf{S}}$ are strictly increasing by Lemmas 2.5 and 3.3, respectively. This completes the proof. \square

Proposition 5.4. *For any $\mathbf{S} \in \Lambda$ we have $\Psi_{\mathbf{S}}(E) = E^{\mathbf{S}}$.*

Proof. We first prove that

$$(5.6) \quad \Psi_{\mathbf{S}}(\beta_\ell^{\mathbf{r}}) = \beta_\ell^{\mathbf{S} \bullet \mathbf{r}} \quad \text{and} \quad \Psi_{\mathbf{S}}(\beta_r^{\mathbf{r}}) = \beta_r^{\mathbf{S} \bullet \mathbf{r}}, \quad \forall \mathbf{r} \in \Omega_L^*.$$

Observe that $\delta(\beta_\ell^{\mathbf{r}}) = \mathbb{L}(\mathbf{r})^\infty$. Then by Lemmas 3.5 and 3.7 it follows that

$$\Phi_{\mathbf{S}}(\delta(\beta_\ell^{\mathbf{r}})) = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^\infty) = (\mathbf{S} \bullet \mathbb{L}(\mathbf{r}))^\infty = \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^\infty = \delta(\beta_\ell^{\mathbf{S} \bullet \mathbf{r}}),$$

so $\Psi_{\mathbf{S}}(\beta_\ell^{\mathbf{r}}) = \beta_\ell^{\mathbf{S} \bullet \mathbf{r}}$. Similarly, since $\delta(\beta_r^{\mathbf{r}}) = \mathbb{L}(\mathbf{r})^+ \mathbf{r}^\infty$, Lemmas 3.4, 3.5 and 3.7 imply that

$$\begin{aligned} \Phi_{\mathbf{S}}(\delta(\beta_r^{\mathbf{r}})) &= \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^+ \mathbf{r}^\infty) = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^+) \Phi_{\mathbf{S}}(\mathbf{r}^\infty) \\ &= (\mathbf{S} \bullet \mathbb{L}(\mathbf{r}))^+ \Phi_{\mathbf{S}}(\mathbf{r}^\infty) = \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^+ (\mathbf{S} \bullet \mathbf{r})^\infty = \delta(\beta_r^{\mathbf{S} \bullet \mathbf{r}}). \end{aligned}$$

We conclude that $\Psi_{\mathbf{S}}(\beta_r^{\mathbf{r}}) = \beta_r^{\mathbf{S} \bullet \mathbf{r}}$. This proves (5.6).

Note by Lemma 2.5 (ii) that the map $\beta \mapsto \delta(\beta)$ is left continuous in $(1, 2]$, and is right continuous at a point β_0 if and only if $\delta(\beta_0)$ is *not* periodic. Hence by Lemma 2.8 it follows that the map $\beta \mapsto \delta(\beta)$ is continuous at β_0 if and only if $\delta(\beta_0)$ is not of the form $\mathbb{L}(\mathbf{r})^\infty$ for a Lyndon word \mathbf{r} . Since $\Phi_{\mathbf{S}}$ is clearly continuous with respect to the order topology and the map δ^{-1} is continuous, it follows that $\Psi_{\mathbf{S}}$ is continuous at β_0 if and only if $\delta(\beta_0)$ is not of the form $\mathbb{L}(\mathbf{r})^\infty$ for a Lyndon word \mathbf{r} . Moreover, $\Psi_{\mathbf{S}}$ is left continuous everywhere.

Note that if $\delta(\beta_0) = \mathbb{L}(\mathbf{r})^\infty$, then by Lemma 2.5 (ii) it follows that as β decreases to β_0 the sequence $\delta(\beta)$ converges to $\mathbb{L}(\mathbf{r})^+0^\infty$ with respect to the order topology, i.e., $\lim_{\beta \searrow \beta_0} \delta(\beta) = \mathbb{L}(\mathbf{r})^+0^\infty$. Since $\Psi_{\mathbf{S}}$ is increasing, it follows that

$$\Psi_{\mathbf{S}}((1, 2]) = (J^{\mathbf{S}} \setminus I^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_{\mathbf{L}}^*} (p^{\mathbf{r}}, q^{\mathbf{r}}],$$

where $\delta(p^{\mathbf{r}}) = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^\infty)$ and $\delta(q^{\mathbf{r}}) = \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r})^+0^\infty)$. Note that $(p^{\mathbf{r}}, q^{\mathbf{r}}] \subset J^{\mathbf{S}\bullet\mathbf{r}}$. By Lemma 2.9 (iii) there is a (unique) Farey word $\hat{\mathbf{r}}$ such that $J^{\mathbf{r}} \subset J^{\hat{\mathbf{r}}}$. Applying (5.6) to both \mathbf{r} and $\hat{\mathbf{r}}$ and using Lemma 5.3 we conclude that $J^{\mathbf{S}\bullet\mathbf{r}} \subset J^{\mathbf{S}\bullet\hat{\mathbf{r}}}$. Hence, $(p^{\mathbf{r}}, q^{\mathbf{r}}] \subset J^{\mathbf{S}\bullet\hat{\mathbf{r}}}$. Therefore, if $\beta \in E^{\mathbf{S}} = (J^{\mathbf{S}} \setminus I^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S}\bullet\mathbf{r}}$, then β lies in the range of $\Psi_{\mathbf{S}}$. This implies that

$$(5.7) \quad E^{\mathbf{S}} \subset \Psi_{\mathbf{S}}((1, 2]).$$

Now assume first that $\beta \in \Psi_{\mathbf{S}}(E)$. Then $\beta = \Psi_{\mathbf{S}}(\hat{\beta})$, where $\hat{\beta} \notin J^{\mathbf{r}}$ for any $\mathbf{r} \in \Omega_F^*$. Hence $\beta \notin J^{\mathbf{S}\bullet\mathbf{r}}$ for any $\mathbf{r} \in \Omega_F^*$ by (5.6) and since $\Psi_{\mathbf{S}}$ is increasing. Therefore, $\beta \in E^{\mathbf{S}}$.

Conversely, suppose $\beta \in E^{\mathbf{S}}$. By (5.7), $\beta = \Psi_{\mathbf{S}}(\hat{\beta})$ for some $\hat{\beta} \in (1, 2]$. If $\hat{\beta} \in J^{\mathbf{r}}$ for some $\mathbf{r} \in \Omega_F^*$, then $\beta \in \Psi_{\mathbf{S}}(J^{\mathbf{r}}) \subset J^{\mathbf{S}\bullet\mathbf{r}}$ by (5.6), contradicting that $\beta \in E^{\mathbf{S}}$. Hence, $\hat{\beta} \in E$ and then $\beta \in \Psi_{\mathbf{S}}(E)$. This completes the proof. \square

Kalle et al. proved in [20, Theorem C] that the Farey intervals $J^{\mathbf{r}}$, $\mathbf{r} \in \Omega_F^*$ cover the whole interval $(1, 2]$ up to a set of zero Hausdorff dimension. Here we strengthen this result and show that the bifurcation set E is uncountable and has zero packing dimension. Furthermore, we show that each relative bifurcation set $E^{\mathbf{S}}$ is uncountable and has zero box-counting dimension. The proof uses the following simple lemma.

Lemma 5.5. *Let $J^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_r^{\mathbf{S}}] =: [p, q]$ be any Lyndon interval. Then the length of $J^{\mathbf{S}}$ satisfies*

$$|J^{\mathbf{S}}| \leq \frac{q}{q-1} q^{-|\mathbf{S}|}.$$

Proof. Since $\delta(p) = \mathbb{L}(\mathbf{S})^\infty$ and $\delta(q) = \mathbb{L}(\mathbf{S})^+\mathbf{S}^\infty$, we have

$$(\mathbb{L}(\mathbf{S})^+0^\infty)_p = 1 = (\mathbb{L}(\mathbf{S})^+\mathbf{S}^\infty)_q =: ((c_i))_q.$$

It follows that

$$|J^{\mathbf{S}}| = q - p = \sum_{i=1}^{|\mathbf{S}|} \frac{c_i}{q^{i-1}} + \sum_{i=|\mathbf{S}|+1}^{\infty} \frac{c_i}{q^{i-1}} - \sum_{i=1}^{|\mathbf{S}|} \frac{c_i}{p^{i-1}} \leq \sum_{i=|\mathbf{S}|+1}^{\infty} \frac{1}{q^{i-1}} = \frac{q}{q-1} q^{-|\mathbf{S}|},$$

as required. \square

Proposition 5.6.

(i) *The bifurcation set*

$$E = (1, 2] \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{r}}$$

is uncountable and has zero packing dimension.

(ii) *For any $\mathbf{S} \in \Lambda$ the relative bifurcation set*

$$E^{\mathbf{S}} = (J^{\mathbf{S}} \setminus I^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S}\bullet\mathbf{r}}$$

is uncountable and has zero box-counting dimension.

Proof. (i) First we prove $\dim_P E = 0$. Let $\rho_N \in (1, 2]$ such that $\delta(\rho_N) = (10^{N-1})^\infty$. Then by Lemma 2.5 it follows that $\rho_N \searrow 1$ as $N \rightarrow \infty$. Thus $E = \bigcup_{N=1}^\infty (E \cap [\rho_N, 2])$. By the countable stability of packing dimension (cf. [16]) it suffices to prove that

$$(5.8) \quad \dim_B(E \cap [\rho_N, 2]) = 0 \quad \text{for all } N \in \mathbb{N}.$$

Let $N \in \mathbb{N}$. Take a Farey interval $J^{\mathbf{s}} := [p, q] \subset [\rho_N, 2]$ with $\mathbf{s} = s_1 \dots s_m \in \Omega_F^*$ such that

$$(5.9) \quad m > N + 2 - 3 \log_2(\rho_N - 1).$$

Write $\mathbb{L}(\mathbf{s}) = a_1 \dots a_m$. Then $\delta(p) = (a_1 \dots a_m)^\infty$. Since $p \geq \rho_N$, by Lemma 2.5 we have $(a_1 \dots a_m)^\infty = \delta(p) \succcurlyeq \delta(\rho_N) = (10^{N-1})^\infty$, which implies that $a_1 \dots a_{N+1} \succcurlyeq 10^{N-1}$. Then by Proposition 2.4 we conclude that

$$(5.10) \quad s_1 \dots s_{N+1} \succcurlyeq 0^N 1.$$

Note that

$$(\mathbb{L}(\mathbf{s})^+ 0^\infty)_p = 1 = (\mathbb{L}(\mathbf{s})^+ \mathbf{s}_\infty)_q =: ((c_i))_q.$$

So, by (5.10) it follows that

$$\sum_{i=1}^m \frac{c_i}{p^i} = 1 = \sum_{i=1}^m \frac{c_i}{q^i} + \sum_{i=m+1}^\infty \frac{c_i}{q^i} > \sum_{i=1}^m \frac{c_i}{q^i} + \frac{1}{q^{m+N+1}},$$

which implies

$$\frac{1}{q^{m+N+1}} < \sum_{i=1}^\infty \left(\frac{1}{p^i} - \frac{1}{q^i} \right) = \frac{q-p}{(p-1)(q-1)}.$$

Whence,

$$(5.11) \quad |J^{\mathbf{s}}| = q-p > \frac{(p-1)(q-1)}{q^{N+1}} q^{-m} \geq \frac{(\rho_N-1)^2}{2^{N+1}} q^{-m}.$$

On the other hand, by Lemma 5.5 it follows that

$$(5.12) \quad |J^{\mathbf{s}}| \leq \frac{q}{q-1} q^{-m} \leq \frac{2}{\rho_N-1} q^{-m} \leq \frac{2}{\rho_N-1} \rho_N^{-m}.$$

Now we list all of the Farey intervals in $[\rho_N, 2]$ in a decreasing order according to their length, say $J^{\mathbf{s}^1}, J^{\mathbf{s}^2}, \dots$. In other words, $|J^{\mathbf{s}^i}| \geq |J^{\mathbf{s}^j}|$ for any $i < j$. For a Farey interval $J^{\mathbf{s}}$, if $J^{\mathbf{s}} = J^{\mathbf{s}^k}$ we then define its *order index* as $o(J^{\mathbf{s}}) = k$.

Set $C_N := 2 \log 2 / \log \rho_N$. Let $J^{\mathbf{s}'}$ be a Farey interval with $|\mathbf{s}'| > C_N m$. Then by (5.9), (5.11) and (5.12) it follows that

$$|J^{\mathbf{s}'}| \leq \frac{2}{\rho_N-1} \rho_N^{-C_N m} = \frac{2}{\rho_N-1} 2^{-2m} < \frac{(\rho_N-1)^2}{2^{N+1}} 2^{-m} \leq |J^{\mathbf{s}}|.$$

This implies that

$$(5.13) \quad o(J^{\mathbf{s}}) \leq \sum_{k=1}^{\lfloor C_N m \rfloor} \# \{ \mathbf{s}' \in \Omega_F^* : |\mathbf{s}'| = k \} \leq \sum_{k=1}^{\lfloor C_N m \rfloor} (k-1) < C_N^2 m^2,$$

where the second inequality follows by (2.1) since the number of non-degenerate Farey words of length k is at most $k-1$ (see [8, Proposition 2.3]). Together with (5.12), (5.13) implies that

$$\liminf_{i \rightarrow \infty} \frac{-\log |J^{\mathbf{s}^i}|}{o(J^{\mathbf{s}^i})} = +\infty.$$

Note that $[\rho_N, 2] \cap E = [\rho_N, 2] \setminus \bigcup_{\mathbf{s} \in \Omega_F^*} J^{\mathbf{s}}$. So, by [17, Proposition 3.6] we conclude (5.8). This proves $\dim_P E = 0$.

Next we prove that E is uncountable. For $\mathbf{s} \in \Omega_F^*$ let $\hat{J}^{\mathbf{s}} = (\beta_\ell^{\mathbf{s}}, \beta_r^{\mathbf{s}})$ be the interior of the Farey interval $J^{\mathbf{s}} = [\beta_\ell^{\mathbf{s}}, \beta_r^{\mathbf{s}}]$. By Lemma 2.9 (i) it follows that the compact set

$$\hat{E} := [1, 2] \setminus \bigcup_{\mathbf{s} \in \Omega_F^*} \hat{J}^{\mathbf{s}}$$

is nonempty and has no isolated points. Hence, \hat{E} is a perfect set and is therefore uncountable. Since $\hat{E} \setminus E$ is countable, it follows that E is uncountable as well.

(ii) In a similar way we prove $\dim_B E^{\mathbf{S}} = 0$. Note that $E^{\mathbf{S}} = (\beta_*^{\mathbf{S}}, \beta_r^{\mathbf{S}}] \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S} \bullet \mathbf{r}}$. Fix a Farey word $\mathbf{r} = r_1 \dots r_m$. Then the Lyndon interval $J^{\mathbf{S} \bullet \mathbf{r}} = [\beta_\ell^{\mathbf{S} \bullet \mathbf{r}}, \beta_r^{\mathbf{S} \bullet \mathbf{r}}] =: [p_{\mathbf{r}}, q_{\mathbf{r}}]$ satisfies

$$(\mathbb{L}(\mathbf{S} \bullet \mathbf{r})^+ 0^\infty)_{p_{\mathbf{r}}} = 1 = (\mathbb{L}(\mathbf{S} \bullet \mathbf{r})^+ (\mathbf{S} \bullet \mathbf{r})^\infty)_{q_{\mathbf{r}}} =: ((d_i))_{q_{\mathbf{r}}}.$$

So,

$$\sum_{i=1}^{m|\mathbf{S}|} \frac{d_i}{p_{\mathbf{r}}^i} = 1 = \sum_{i=1}^{m|\mathbf{S}|} \frac{d_i}{q_{\mathbf{r}}^i} + \sum_{i=m|\mathbf{S}|+1}^{\infty} \frac{d_i}{q_{\mathbf{r}}^i} > \sum_{i=1}^{m|\mathbf{S}|} \frac{d_i}{q_{\mathbf{r}}^i} + \frac{1}{q_{\mathbf{r}}^{(m+1)|\mathbf{S}|+1}},$$

where the inequality follows by observing that $\mathbf{S} \in \Omega_L^*$ and thus $\mathbf{S} \bullet \mathbf{r}^\infty \succcurlyeq 0^{|\mathbf{S}|} 10^\infty$. Therefore,

$$\frac{1}{q_{\mathbf{r}}^{(m+1)|\mathbf{S}|+1}} \leq \sum_{i=1}^{\infty} \left(\frac{1}{p_{\mathbf{r}}^i} - \frac{1}{q_{\mathbf{r}}^i} \right) = \frac{q_{\mathbf{r}} - p_{\mathbf{r}}}{(p_{\mathbf{r}} - 1)(q_{\mathbf{r}} - 1)},$$

which implies

$$|J^{\mathbf{S} \bullet \mathbf{r}}| = q_{\mathbf{r}} - p_{\mathbf{r}} \geq \frac{(p_{\mathbf{r}} - 1)(q_{\mathbf{r}} - 1)}{q_{\mathbf{r}}^{|\mathbf{S}|+1}} q_{\mathbf{r}}^{-|\mathbf{S} \bullet \mathbf{r}|} > \frac{(\beta_*^{\mathbf{S}} - 1)^2}{2^{|\mathbf{S}|+1}} q_{\mathbf{r}}^{-|\mathbf{S} \bullet \mathbf{r}|}.$$

On the other hand, by Lemma 5.5 it follows that

$$|J^{\mathbf{S} \bullet \mathbf{r}}| \leq \frac{q_{\mathbf{r}}}{q_{\mathbf{r}} - 1} q_{\mathbf{r}}^{-m|\mathbf{S}|} \leq \frac{2}{\beta_*^{\mathbf{S}} - 1} q_{\mathbf{r}}^{-|\mathbf{S} \bullet \mathbf{r}|}.$$

Now let $(J^{\mathbf{S} \bullet \mathbf{r}_i})$ be an enumeration of the intervals $J^{\mathbf{S} \bullet \mathbf{r}}$, $\mathbf{r} \in \Omega_F^*$, arranged in order by decreasing length. Then by a similar argument as in (i) above, we obtain

$$\liminf_{i \rightarrow \infty} \frac{-\log |J^{\mathbf{S} \bullet \mathbf{r}_i}|}{\log o(J^{\mathbf{S} \bullet \mathbf{r}_i})} = +\infty.$$

Thus, $\dim_B E^{\mathbf{S}} = 0$.

Finally, since we showed in (i) that E is uncountable, we conclude by Lemma 5.3 and Proposition 5.4 that $E^{\mathbf{S}} = \Psi_{\mathbf{S}}(E)$ is also uncountable. This completes the proof. \square

5.2. Infinite bifurcation set. Recall from (1.7) that

$$E_\infty = \bigcap_{n=1}^{\infty} \bigcup_{\mathbf{S} \in \Lambda(n)} J^{\mathbf{S}},$$

where

$$\Lambda(n) = \{\mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \dots \bullet \mathbf{s}_n : \mathbf{s}_i \in \Omega_F^* \text{ for all } 1 \leq i \leq n\}.$$

In particular, $\Lambda(1) = \Omega_F^*$ and $\Lambda = \bigcup_{n=1}^{\infty} \Lambda(n)$. Note that $(1, 2] = E \cup \bigcup_{\mathbf{s} \in \Omega_F^*} J^{\mathbf{s}}$. Furthermore, for each word $\mathbf{S} \in \Lambda$ we have

$$J^{\mathbf{S}} \setminus I^{\mathbf{S}} = E^{\mathbf{S}} \cup \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S}\mathbf{r}}.$$

By iteration of the above equation we obtain the following partition of the interval $(1, 2]$.

Lemma 5.7. *The interval $(1, 2]$ can be partitioned as*

$$(1, 2] = E \cup E_{\infty} \cup \bigcup_{\mathbf{S} \in \Lambda} E^{\mathbf{S}} \cup \bigcup_{\mathbf{S} \in \Lambda} I^{\mathbf{S}}.$$

To complete the proof of Theorem 3 we still need the following dimension result for E_{∞} .

Proposition 5.8. *We have $\dim_H E_{\infty} = 0$.*

Proof. Note by (1.7) that

$$(5.14) \quad E_{\infty} = \bigcap_{n=1}^{\infty} \bigcup_{\mathbf{S} \in \Lambda(n)} J^{\mathbf{S}} \subset \bigcap_{n=1}^{\infty} \bigcup_{\mathbf{S} \in \Lambda: |\mathbf{S}| \geq n} J^{\mathbf{S}}.$$

This suggests covering E_{∞} by the intervals $J^{\mathbf{S}}$ for $\mathbf{S} \in \Lambda$ with $|\mathbf{S}| \geq n$ for a sufficiently large n . To this end, we first estimate the diameter of $J^{\mathbf{S}}$. Take $\mathbf{S} \in \Lambda$ with $|\mathbf{S}| = m$, and write $J^{\mathbf{S}} = [p, q]$. Then $\delta(p) = \mathbb{L}(\mathbf{S})^{\infty}$ and $\delta(q) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^{\infty}$, and it follows from Lemma 5.5 that

$$(5.15) \quad |J^{\mathbf{S}}| \leq \frac{q}{q-1} \cdot q^{-m}.$$

Let (β_n) be an arbitrary sequence in $(1, 2)$ decreasing to 1. We will use (5.15) to show that $\dim_H(E_{\infty} \cap (\beta_n, 2]) = 0$ for all $n \in \mathbb{N}$, so the result will follow from the countable stability of Hausdorff dimension (cf. [16]). Fix $n \in \mathbb{N}$. Observe that, if $J^{\mathbf{S}} = [p, q]$ intersects $(\beta_n, 2]$, then $q > \beta_n$ and so by (5.15),

$$(5.16) \quad |J^{\mathbf{S}}| \leq \frac{2}{\beta_n - 1} \beta_n^{-m} =: C_n \beta_n^{-m}.$$

Next, we count how many words $\mathbf{S} \in \Lambda$ there are with $|\mathbf{S}| = m$. Call this number N_m . Observe that if $\mathbf{S} = \mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \cdots \bullet \mathbf{s}_k$ and $|\mathbf{s}_i| = l_i$ for $i = 1, \dots, k$, then $|\mathbf{S}| = l_1 l_2 \cdots l_k$. Note by [8, Proposition 2.3] that $\#\{\mathbf{r} \in \Omega_F^* : |\mathbf{r}| = l\} \leq l - 1$ for any $l \geq 2$. Thus, for any given tuple (l_1, \dots, l_k) , the number of possible choices for the words $\mathbf{s}_1, \dots, \mathbf{s}_k$ is at most $l_1 l_2 \cdots l_k = |\mathbf{S}| = m$. It remains to estimate how many *ordered factorizations* of m there are, that is to estimate the number

$$f_m := \#\{(l_1, \dots, l_k) : k \in \mathbb{N}, \quad l_i \in \mathbb{N}_{\geq 2} \forall i \quad \text{and} \quad l_1 l_2 \cdots l_k = m\}.$$

By considering the possible values of l_1 , it is easy to see that f_m satisfies the recursion

$$f_m = \sum_{d|m, d>1} f_{m/d},$$

where we set $f_1 := 1$. (See [19].) We claim that $f_m \leq m^2$. This is trivial for $m = 1$, so let $m \geq 2$ and assume $f_n \leq n^2$ for all $n < m$; then

$$f_m = \sum_{d|m, d>1} f_{m/d} \leq \sum_{d|m, d>1} \left(\frac{m}{d}\right)^2 \leq m^2 \sum_{d=2}^{\infty} \frac{1}{d^2} = m^2 \left(\frac{\pi^2}{6} - 1\right) < m^2.$$

This proves the claim, and we thus conclude that $N_m \leq m^3$. Now, given $\varepsilon > 0$ and $\delta > 0$, choose N large enough so that $C_n(\beta_n)^{-N} < \delta$. Using (5.14) and (5.16), we obtain

$$\mathcal{H}_\delta^\varepsilon(E_\infty \cap (\beta_n, 2]) \leq \sum_{\mathbf{S} \in \Lambda: |\mathbf{S}| \geq N, J^{\mathbf{S}} \cap (\beta_n, 2] \neq \emptyset} |J^{\mathbf{S}}|^\varepsilon \leq \sum_{m=N}^{\infty} m^3 C_n^\varepsilon \beta_n^{-m\varepsilon} \rightarrow 0$$

as $N \rightarrow \infty$. This shows that $\dim_H(E_\infty \cap (\beta_n, 2]) = 0$, as desired. \square

Proof of Theorem 3. The theorem follows by Proposition 5.6, Lemma 5.7 and Proposition 5.8. \square

6. CRITICAL VALUES IN THE BIFURCATION SETS

By Proposition 1.12, Theorem 2 and Theorem 3 it suffices to determine the critical value $\tau(\beta)$ for

$$\beta \in \bigcup_{\mathbf{S} \in \Lambda} E^{\mathbf{S}} \cup E_\infty.$$

First we compute $\tau(\beta)$ for $\beta \in \bigcup_{\mathbf{S} \in \Lambda} E^{\mathbf{S}}$. Recall from Lemma 5.3 and Proposition 5.4 that for each $\mathbf{S} \in \Lambda$ the map $\Psi_{\mathbf{S}}$ bijectively maps the bifurcation set $E = (1, 2] \setminus \bigcup_{\mathbf{s} \in \Omega_F^*} J^{\mathbf{s}}$ to the relative bifurcation set $E^{\mathbf{S}} = (J^{\mathbf{S}} \setminus I^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S}\mathbf{r}}$.

Lemma 6.1. *Let $\hat{\beta} \in E \setminus \{2\}$ with $\delta(\hat{\beta}) = \delta_1 \delta_2 \dots$. Also let $\mathbf{S} \in \Lambda$, and set $\beta := \Psi_{\mathbf{S}}(\hat{\beta})$. Then*

- (i) $b(\tau(\hat{\beta}), \hat{\beta}) = 0\delta_2\delta_3\dots$; and
- (ii) *the map $\hat{t} \mapsto (\Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta})))_{\beta}$ is continuous at $\tau(\hat{\beta})$.*

Proof. First we prove (i). Note by Proposition 1.12 that $\tau(\hat{\beta}) = 1 - 1/\hat{\beta} = (0\delta_2\delta_3\dots)_{\hat{\beta}}$. So by Lemma 2.10 it suffices to verify that

$$(6.1) \quad \sigma^n(0\delta_2\delta_3\dots) \prec \delta_1\delta_2\dots \quad \forall n \geq 0.$$

By Lemma 2.5, it is immediate that $\sigma^n(0\delta_2\delta_3\dots) \prec (\delta_i)$ for all $n \geq 0$. If equality holds for some n , then $\delta(\hat{\beta}) = (\delta_i)$ is periodic with period $m \geq 2$ (since $\hat{\beta} \neq 2$), so by Lemma 2.8, $\delta(\hat{\beta}) = \mathbb{L}(\mathbf{r})^\infty$ for some Lyndon word \mathbf{r} . This implies $\hat{\beta} = \beta_{\hat{t}}^{\mathbf{r}} \in J^{\mathbf{r}}$. But then by Lemma 2.9, $\hat{\beta} \in J^{\mathbf{s}}$ for some Farey word \mathbf{s} , and so $\hat{\beta} \notin E$, a contradiction. This proves (6.1), and then yields statement (i).

For (ii), note by Lemma 2.10 (ii) that the map $\hat{t} \mapsto b(\hat{t}, \hat{\beta})$ is continuous at all points \hat{t} for which $b(\hat{t}, \hat{\beta})$ does not end with 0^∞ . Furthermore, the map $\Phi_{\mathbf{S}}$ is continuous with respect to the order topology. So, by statement (i) it follows that the map $\hat{t} \mapsto (\Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta})))_{\beta}$ is continuous at $\tau(\hat{\beta})$, completing the proof. \square

Proposition 6.2. *Let $\mathbf{S} \in \Lambda$. Then for any $\beta \in E^{\mathbf{S}}$ we have*

$$\tau(\beta) = \left(\Phi_{\mathbf{S}}(0\delta_2\delta_3\dots) \right)_{\beta},$$

where $1\delta_2\delta_3\dots$ is the quasi-greedy expansion of 1 in base $\hat{\beta} := \Psi_{\mathbf{S}}^{-1}(\beta)$.

Proof. Let $\beta \in E^{\mathbf{S}}$. Then by Lemma 5.3 and Proposition 5.4 there exists a unique $\hat{\beta} \in E$ such that $\beta = \Psi_{\mathbf{S}}^{-1}(\beta) \in E$, in other words, $\delta(\beta) = \Phi_{\mathbf{S}}(\delta(\hat{\beta}))$. Write $\delta(\hat{\beta}) = \delta_1\delta_2\dots$, and set $t^* := (\Phi_{\mathbf{S}}(0\delta_2\delta_3\dots))_{\beta}$. We will show that $\tau(\beta) = t^*$, by proving that $h_{top}(\mathbf{K}_{\beta}(t)) > 0$ for

$t < t^*$, and $\mathbf{K}_\beta(t)$ is countable for $t > t^*$. We consider separately the two cases: (i) $\hat{\beta} < 2$ and (ii) $\hat{\beta} = 2$.

Case I. $\hat{\beta} < 2$. First, for notational convenience, we define the map

$$\Theta_{\mathbf{S}, \hat{\beta}} : (0, 1) \rightarrow (0, 1); \quad \hat{t} \mapsto (\Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta})))_{\beta}.$$

Since by Lemma 6.1 the map $\Theta_{\mathbf{S}, \hat{\beta}}$ is continuous at $\tau(\hat{\beta})$ and $t^* = \Theta_{\mathbf{S}, \hat{\beta}}(\tau(\hat{\beta})) = (\Phi_{\mathbf{S}}(0\delta_2\delta_3\dots))_{\beta}$, it is, by the monotonicity of the set-valued map $t \mapsto \mathbf{K}_\beta(t)$, sufficient to prove the following two things:

$$(6.2) \quad \hat{t} < \tau(\hat{\beta}) \quad \implies \quad h_{top}(\mathbf{K}_\beta(\Theta_{\mathbf{S}, \hat{\beta}}(\hat{t}))) > 0,$$

and

$$(6.3) \quad \hat{t} > \tau(\hat{\beta}) \quad \implies \quad \mathbf{K}_\beta(\Theta_{\mathbf{S}, \hat{\beta}}(\hat{t})) \text{ is countable.}$$

First, take $\hat{t} < \tau(\hat{\beta})$ and set $t := \Theta_{\mathbf{S}, \hat{\beta}}(\hat{t}) = (\Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta})))_{\beta}$. Since $\sigma^n(b(\hat{t}, \hat{\beta})) \prec \delta(\hat{\beta})$ for all $n \geq 0$, Lemma 3.8 implies that

$$\sigma^n(\Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta}))) \prec \Phi_{\mathbf{S}}(\delta(\hat{\beta})) = \delta(\beta) \quad \forall n \geq 0.$$

Hence, $b(t, \beta) = \Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta}))$. Now

$$\begin{aligned} \Phi_{\mathbf{S}}(\mathbf{K}_{\hat{\beta}}(\hat{t})) &= \{\Phi_{\mathbf{S}}((x_i)) : b(\hat{t}, \hat{\beta}) \preceq \sigma^n((x_i)) \prec \delta(\hat{\beta}) \quad \forall n \geq 0\} \\ &\subset \{\Phi_{\mathbf{S}}((x_i)) : \Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta})) \preceq \sigma^n(\Phi_{\mathbf{S}}((x_i))) \prec \Phi_{\mathbf{S}}(\delta(\hat{\beta})) \quad \forall n \geq 0\} \\ &= \{\Phi_{\mathbf{S}}((x_i)) : b(t, \beta) \preceq \sigma^n(\Phi_{\mathbf{S}}((x_i))) \prec \delta(\beta) \quad \forall n \geq 0\} \\ &\subset \{(y_i) : b(t, \beta) \preceq \sigma^n((y_i)) \prec \delta(\beta) \quad \forall n \geq 0\} = \mathbf{K}_\beta(t), \end{aligned}$$

where the first inclusion again follows by Lemma 3.8. We deduce that

$$h_{top}(\mathbf{K}_\beta(t)) \geq h_{top}(\Phi_{\mathbf{S}}(\mathbf{K}_{\hat{\beta}}(\hat{t}))) = |\mathbf{S}|^{-1} h_{top}(\mathbf{K}_{\hat{\beta}}(\hat{t})) > 0,$$

where the last inequality follows since $\hat{t} < \tau(\hat{\beta})$. This gives (6.2).

Next, let $\hat{t} > \tau(\hat{\beta})$ and set $t := \Theta_{\mathbf{S}, \hat{\beta}}(\hat{t})$. Then by the same argument as above we have $b(t, \beta) = \Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta}))$. Since $\hat{\beta} \in E$, there exists a sequence of Farey intervals $J^{\mathbf{r}_k} = [\beta_{\ell}^{\mathbf{r}_k}, \beta_r^{\mathbf{r}_k}]$ with $\mathbf{r}_k \in \Omega_F^*$ such that $\hat{q}_k := \beta_{\ell}^{\mathbf{r}_k} \searrow \hat{\beta}$ as $k \rightarrow \infty$.

We claim that $b(\hat{t}, \hat{\beta}) \succ (\mathbf{r}_k)^{\infty}$ for all sufficiently large k . This can be seen as follows. As explained in the proof of Lemma 6.1, $\delta(\hat{\beta})$ is not periodic, and therefore by Lemma 2.5 (ii) the map $\beta' \mapsto \delta(\beta')$ is continuous at $\hat{\beta}$ (where we use β' to denote a generic base). This implies $\delta(\hat{q}_k) \searrow \delta(\hat{\beta}) = 1\delta_2\delta_3\dots$ as $k \rightarrow \infty$. But $\delta(\hat{q}_k) = \delta(\beta_{\ell}^{\mathbf{r}_k}) = \mathbb{L}(\mathbf{r}_k)^{\infty}$, and by Lemma 2.3 $\mathbb{L}(\mathbf{r}_k)$ is the word obtained from \mathbf{r}_k by flipping the first and last digits. Thus, $(\mathbf{r}_k)^{\infty}$ converges to $0\delta_2\delta_3\dots$ in the order topology. Since $\hat{t} > \tau(\hat{\beta})$ implies $b(\hat{t}, \hat{\beta}) \succ b(\tau(\hat{\beta}), \hat{\beta}) = 0\delta_2\delta_3\dots$, the claim follows.

We can now deduce that, for all sufficiently large k ,

$$\begin{aligned} \mathbf{K}_\beta(t) &= \{(y_i) : b(t, \beta) \preceq \sigma^n((y_i)) \prec \delta(\beta) \quad \forall n \geq 0\} \\ &= \{(y_i) : \Phi_{\mathbf{S}}(b(\hat{t}, \hat{\beta})) \preceq \sigma^n((y_i)) \prec \Phi_{\mathbf{S}}(\delta(\hat{\beta})) \quad \forall n \geq 0\} \\ &\subset \{(y_i) : \Phi_{\mathbf{S}}((\mathbf{r}_k)^{\infty}) \preceq \sigma^n((y_i)) \prec \Phi_{\mathbf{S}}(\mathbb{L}(\mathbf{r}_k)^{\infty}) \quad \forall n \geq 0\} \\ &= \{(y_i) : (\mathbf{S} \bullet \mathbf{r}_k)^{\infty} \preceq \sigma^n((y_i)) \prec \mathbb{L}(\mathbf{S} \bullet \mathbf{r}_k)^{\infty} \quad \forall n \geq 0\}, \end{aligned}$$

where the inclusion follows using the claim and $\delta(\hat{\beta}) \prec \delta(\hat{q}_k) = \mathbb{L}(\mathbf{r}_k)^\infty$. Hence, $\mathbf{K}_\beta(t)$ is countable by Proposition 4.1. This establishes (6.3).

Case II. $\hat{\beta} = 2$. In this case, $\delta(\hat{\beta}) = 1^\infty$, so $\delta(\beta) = \Phi_{\mathbf{S}}(\delta(\hat{\beta})) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty$ and $t^* = (\Phi_{\mathbf{S}}(01^\infty))_\beta = (\mathbf{S}^- \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty)_\beta = (\mathbf{S}0^\infty)_\beta$. Recall that $\beta = \beta_r^{\mathbf{S}}$ is the right endpoint of the Lyndon interval $J^{\mathbf{S}}$.

If $t < t^*$, then $b(t, \beta) \prec b(t^*, \beta) = \mathbf{S}0^\infty$, so by Lemma 2.10 (iii) there exists $k \in \mathbb{N}$ such that $b(t, \beta) \preceq \mathbf{S}^- \mathbb{L}(\mathbf{S})^+ \mathbf{S}^k 0^\infty$. It follows that

$$\begin{aligned} \mathbf{K}_\beta(t) &= \{(x_i) : b(t, \beta) \preceq \sigma^n((x_i)) \prec \delta(\beta) \forall n \geq 0\} \\ &\supset \{\mathbb{L}(\mathbf{S})^+ \mathbf{S}^k \mathbf{S}^-, \mathbb{L}(\mathbf{S})^+ \mathbf{S}^{k+1} \mathbf{S}^-\}^{\mathbb{N}}, \end{aligned}$$

and hence $h_{\text{top}}(\mathbf{K}_\beta(t)) > 0$.

Now suppose $t > t^*$. Then $b(t, \beta) \succ b(t^*, \beta) = \mathbf{S}0^\infty$, so by Lemma 4.4,

$$\begin{aligned} \mathbf{K}_\beta(t) &= \{(x_i) : b(t, \beta) \preceq \sigma^n((x_i)) \prec \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty \forall n \geq 0\} \\ &\subset \{(x_i) : \mathbf{S}0^\infty \preceq \sigma^n((x_i)) \prec \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty \forall n \geq 0\} \\ &= \{(x_i) : \mathbf{S}^\infty \preceq \sigma^n((x_i)) \preceq \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty \forall n \geq 0\} \\ &= \{(x_i) : \mathbf{S}^\infty \preceq \sigma^n((x_i)) \preceq \mathbb{L}(\mathbf{S})^\infty \forall n \geq 0\}, \end{aligned}$$

where the second equality follows by using $\mathbf{S} \in \Omega_L^*$, so $\sigma^n((x_i)) \succ \mathbf{S}0^\infty$ for all $n \geq 0$ if and only if $\sigma^n((x_i)) \succ \mathbf{S}^\infty$ for all $n \geq 0$. Therefore, $\mathbf{K}_\beta(t)$ is countable by Proposition 4.1. This completes the proof. \square

Next we will determine the critical value $\tau(\beta)$ for $\beta \in E_\infty$. Recall from (1.7) that

$$E_\infty = \bigcap_{n=1}^{\infty} \bigcup_{\mathbf{S} \in \Lambda(n)} J^{\mathbf{S}},$$

where for each $n \in \mathbb{N}$ the Lyndon intervals $J^{\mathbf{S}}, \mathbf{S} \in \Lambda(n)$ are pairwise disjoint. Thus, for any $\beta \in E_\infty$ there exists a unique sequence of words (\mathbf{s}_k) with each $\mathbf{s}_k \in \Omega_F^*$ such that

$$\{\beta\} = \bigcap_{n=1}^{\infty} J^{\mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_n}.$$

We call (\mathbf{s}_k) the *coding* of β .

Proposition 6.3. *For any $\beta \in E_\infty$ with its coding (\mathbf{s}_k) we have*

$$\tau(\beta) = \lim_{n \rightarrow \infty} (\mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_n 0^\infty)_\beta.$$

Proof. Take $\beta \in E_\infty$. For $k \geq 1$ let $\mathbf{S}_k := \mathbf{s}_1 \bullet \cdots \bullet \mathbf{s}_k$, and write $t_k := (\mathbf{S}_k 0^\infty)_\beta$. Note that $\beta \in J^{\mathbf{S}_k} = [\beta_\ell^{\mathbf{S}_k}, \beta_r^{\mathbf{S}_k}]$ for all $k \geq 1$. Hence

$$(6.4) \quad \delta(\beta) \succ \delta(\beta_\ell^{\mathbf{S}_k}) = \mathbb{L}(\mathbf{S}_k)^\infty,$$

which implies that $b(t_k, \beta) = \mathbf{S}_k 0^\infty$ for all $k \geq 1$. Observe that $\mathbf{S}_{k+1} = \mathbf{S}_k \bullet \mathbf{s}_{k+1}$ begins with \mathbf{S}_k^- . Therefore,

$$t_{k+1} = (\mathbf{S}_{k+1} 0^\infty)_\beta < (\mathbf{S}_k 0^\infty)_\beta = t_k,$$

so the sequence (t_k) is decreasing. Since $t_k \geq 0$ for all $k \geq 1$, the limit $t^* := \lim_{k \rightarrow \infty} t_k$ exists. We will now show that $\tau(\beta) = t^*$.

First we prove $\tau(\beta) \leq t^*$. Since t_k decreases to t^* as $k \rightarrow \infty$, it suffices to prove that $\tau(\beta) \leq t_k$ for all $k \geq 1$. Let $q_k := \beta_r^{\mathbf{S}^k}$ for all $k \geq 1$. Then $q_k > \beta$ since $\beta \in J^{\mathbf{S}^k}$, and $q_k \searrow \beta$ as $k \rightarrow \infty$. Set $t'_k := (\mathbf{S}_k 0^\infty)_{q_k}$. Since $q_k > \beta$, one can verify that $b(t'_k, q_k) = \mathbf{S}_k 0^\infty = b(t_k, \beta)$. So,

$$(6.5) \quad \begin{aligned} \mathbf{K}_\beta(t_k) &= \{(x_i) : b(t_k, \beta) \preceq \sigma^n((x_i)) \prec \delta(\beta) \forall n \geq 0\} \\ &\subset \{(x_i) : \mathbf{S}_k 0^\infty \preceq \sigma^n((x_i)) \prec \delta(q_k) \forall n \geq 0\} = \mathbf{K}_{q_k}(t'_k). \end{aligned}$$

Note by Proposition 6.2 and Case II of its proof that $\tau(q_k) = (\mathbf{S}_k 0^\infty)_{q_k} = t'_k$. This implies that $\dim_H K_{q_k}(t'_k) = 0$, and thus by (6.5) we have $\dim_H K_\beta(t_k) = 0$. Hence $\tau(\beta) \leq t_k$ for any $k \geq 1$. Letting $k \rightarrow \infty$ we obtain that $\tau(\beta) \leq t^*$.

Next we prove $\tau(\beta) \geq t^*$. Note that $\beta = \Psi_{\mathbf{S}_k}(\beta_k)$, where $\beta_k \in E_\infty$ has coding $(\mathbf{s}_{k+1}, \mathbf{s}_{k+2}, \dots)$. Let $a(\hat{t}, \beta_k)$ denote the quasi-greedy expansion of \hat{t} in base β_k (cf. [15, Lemma 2.3]). Observe that the map $\hat{t} \mapsto a(\hat{t}, \beta_k)$ is strictly increasing and left continuous everywhere in $(0, 1)$, and thus the map $\hat{t} \mapsto (\Phi_{\mathbf{S}_k}(a(\hat{t}, \beta_k)))_\beta$ is also left continuous in $(0, 1)$. So, by the same argument as in the proof of (6.2) it follows that

$$\tau(\beta) \geq (\Phi_{\mathbf{S}_k}(a(\tau(\beta_k), \beta_k)))_\beta \geq (\Phi_{\mathbf{S}_k}(0^\infty))_\beta > (\mathbf{S}_k^- 0^\infty)_\beta$$

for every $k \in \mathbb{N}$, and letting $k \rightarrow \infty$ gives $\tau(\beta) \geq t^*$. \square

To illustrate Proposition 6.3 we construct in each Farey interval $J^{\mathbf{s}}$ a transcendental base $\beta \in E_\infty$ and give an explicit formula for the critical value $\tau(\beta)$. Recall from [4] that the classical *Thue-Morse sequence* $(\theta_i)_{i=0}^\infty = 01101001\dots$ is defined recursively as follows. Let $\theta_0 = 0$; and if $\theta_0 \dots \theta_{2^n-1}$ is defined for some $n \geq 0$, then

$$(6.6) \quad \theta_{2^n} \dots \theta_{2^{n+1}-1} = \overline{\theta_0 \dots \theta_{2^n-1}}.$$

By the definition of (θ_i) it follows that

$$(6.7) \quad \theta_{2k+1} = 1 - \theta_k, \quad \theta_{2k} = \theta_k \quad \text{for any } k \geq 0.$$

Komornik and Loreti [21] showed that

$$(6.8) \quad \theta_{i+1}\theta_{i+2}\dots \prec \theta_1\theta_2\dots \quad \forall i \geq 1.$$

Proposition 6.4. *Given $\mathbf{s} = s_1 \dots s_m = 0\mathbf{c}1 \in \Omega_F^*$, let $\beta := \beta_\infty^{\mathbf{s}} \in (1, 2]$ such that*

$$(\theta_1\mathbf{c}\theta_2\theta_3\mathbf{c}\theta_4 \dots \theta_{2k+1}\mathbf{c}\theta_{2k+2}\dots)_\beta = 1.$$

Then $\beta \in E_\infty \cap J^{\mathbf{s}}$ is transcendental, and

$$\tau(\beta) = \frac{2 \sum_{j=2}^m s_j \beta^{m-j} + \beta^{m-1} - \beta^m}{\beta^m - 1}.$$

We point out that in the above proposition, \mathbf{c} may be the empty word. To prove the transcendence of β we recall the following result due to Mahler [25].

Lemma 6.5 (Mahler, 1976). *If z is an algebraic number in the open unit disc, then the number $Z := \sum_{i=1}^\infty \theta_i z^i$ is transcendental.*

Proof of Proposition 6.4. Let $\mathbf{s} = s_1 \dots s_m = 0\mathbf{c}1 \in \Omega_F^*$. First we prove that

$$(6.9) \quad \delta(\beta) = \theta_1\mathbf{c}\theta_2\theta_3\mathbf{c}\theta_4 \dots \theta_{2k+1}\mathbf{c}\theta_{2k+2}\dots =: (\delta_i).$$

By Lemma 2.5 it suffices to prove that $\sigma^n((\delta_i)) \preceq (\delta_i)$ for all $n \geq 1$. Note by Lemma 2.3 that $\delta_1 \dots \delta_m = 1\mathbf{c}1 = \mathbb{L}(\mathbf{s})^+ =: a_1 \dots a_m^+$. Take $n \in \mathbb{N}$, and write $n = mk + j$ with $k \in \mathbb{N} \cup \{0\}$ and $j \in \{1, 2, \dots, m\}$. We will prove $\sigma^n((\delta_i)) \prec (\delta_i)$ in the following three cases.

Case I. $j \in \{1, 2, \dots, m-2\}$. Note by (6.7) that $\theta_{2k} = 1 - \theta_{2k+1}$. This implies that $\sigma^n((\delta_i))$ begins with either $a_{j+1} \dots a_m$ or $a_{j+1} \dots a_m^+ s_1 \dots s_{m-1}$. By Lemma 2.7 it follows that $\sigma^n((\delta_i)) \prec (\delta_i)$.

Case II. $j = m-1$. Then $\sigma^n((\delta_i))$ begins with $\theta_{2k} \theta_{2k+1} \mathbf{c}$ for some $k \in \mathbb{N}$. If $\theta_{2k} = 0$, then it is clear that $\sigma^n((\delta_i)) \prec (\delta_i)$ since $\delta_1 = 1$. Otherwise, by (6.7) it follows that $\theta_{2k} \theta_{2k+1} \mathbf{c} = 10\mathbf{c} = 1s_1 \dots s_{m-1}$. Hence,

$$\sigma^n((\delta_i)) = 1s_1 \dots s_{m-1} \delta_{n+m+1} \delta_{n+m+2} \dots \prec 1a_2 \dots a_m^+ \delta_{m+1} \delta_{m+2} = (\delta_i),$$

where the strict inequality follows since $s_1 \dots s_{m-1} \preccurlyeq a_2 \dots a_m$.

Case III. $j = m$. Then

$$\sigma^n((\delta_i)) = \theta_{2k+1} \mathbf{c} \theta_{2k+2} \theta_{2k+3} \mathbf{c} \theta_{2k+4} \dots \prec \theta_1 \mathbf{c} \theta_2 \theta_3 \mathbf{c} \theta_4 \dots = (\delta_i),$$

where the strict inequality follows by (6.8).

Therefore, by Cases I–III we establish (6.9). Next we show that $\beta \in E_\infty$. For $k \in \mathbb{N}$ let $\mathbf{S}_k := \mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \dots \bullet \mathbf{s}_k$ with $\mathbf{s}_1 = \mathbf{s}$ and $\mathbf{s}_i = 01$ for all $2 \leq i \leq k$. Then $\mathbf{S}_k \in \Lambda$ for all $k \in \mathbb{N}$. So it suffices to show that $\beta \in J^{\mathbf{S}_k}$ for all $k \geq 1$. First we claim that

$$(6.10) \quad \mathbf{S}_k = \overline{\theta_1 \mathbf{c} \theta_2 \theta_3 \mathbf{c} \theta_4} \dots \overline{\theta_{2^{k-1}} \mathbf{c} \theta_{2^k}^+}, \quad \mathbb{L}(\mathbf{S}_k) = \theta_1 \mathbf{c} \theta_2 \theta_3 \mathbf{c} \theta_4 \dots \theta_{2^{k-1}} \mathbf{c} \theta_{2^k}^-$$

for all $k \geq 1$.

Since $\mathbf{S}_1 = \mathbf{s} = 0\mathbf{c}1 = \overline{\theta_1 \mathbf{c} \theta_2^+}$ and $\mathbb{L}(\mathbf{S}_1) = 1\mathbf{c}0 = \theta_1 \mathbf{c} \theta_2^-$, (6.10) holds for $k = 1$. Now suppose (6.10) holds for a given $k \in \mathbb{N}$. Then

$$\begin{aligned} \mathbf{S}_{k+1} &= \mathbf{S}_k \bullet (01) = \mathbf{S}_k^- \mathbb{L}(\mathbf{S}_k)^+ = \overline{\theta_1 \mathbf{c} \theta_2} \dots \overline{\theta_{2^{k-1}} \mathbf{c} \theta_{2^k}^-} \theta_1 \mathbf{c} \theta_2 \dots \theta_{2^{k-1}} \mathbf{c} \theta_{2^k} \\ &= \overline{\theta_1 \mathbf{c} \theta_2} \dots \overline{\theta_{2^{k+1}-1} \mathbf{c} \theta_{2^{k+1}}^+}, \end{aligned}$$

where the last equality follows since, by the definition of (θ_i) in (6.6), $\theta_{2^{k+1}} \dots \theta_{2^{k+1}} = \overline{\theta_1 \dots \theta_{2^k}^+}$. Similarly, by the induction hypothesis and Lemma 3.7 it follows that

$$\begin{aligned} \mathbb{L}(\mathbf{S}_{k+1}) &= \mathbb{L}(\mathbf{S}_k \bullet (01)) = \mathbf{S}_k \bullet \mathbb{L}(01) = \mathbf{S}_k \bullet (10) = \mathbb{L}(\mathbf{S}_k)^+ \mathbf{S}_k^- \\ &= \theta_1 \mathbf{c} \theta_2 \dots \theta_{2^{k-1}} \mathbf{c} \theta_{2^k} \overline{\theta_1 \mathbf{c} \theta_2} \dots \overline{\theta_{2^k-1} \mathbf{c} \theta_{2^k}^-} \\ &= \theta_1 \mathbf{c} \theta_2 \dots \theta_{2^{k+1}-1} \mathbf{c} \theta_{2^{k+1}}^-. \end{aligned}$$

Hence, by induction (6.10) holds for all $k \geq 1$.

Next, recall that the Lyndon interval $J^{\mathbf{S}_k} = [\beta_\ell^{\mathbf{S}_k}, \beta_r^{\mathbf{S}_k}]$ satisfies

$$(6.11) \quad \delta(\beta_\ell^{\mathbf{S}_k}) = \mathbb{L}(\mathbf{S}_k)^\infty \quad \text{and} \quad \delta(\beta_r^{\mathbf{S}_k}) = \mathbb{L}(\mathbf{S}_k)^+ \mathbf{S}_k^\infty.$$

By (6.9) and (6.10) it follows that $\delta(\beta)$ begins with $\mathbb{L}(\mathbf{S}_k)^+$, so by (6.11), $\delta(\beta) \succ \mathbb{L}(\mathbf{S}_k)^\infty = \delta(\beta_\ell^{\mathbf{S}_k})$. Thus, $\beta > \beta_\ell^{\mathbf{S}_k}$ for all $k \geq 1$. On the other hand, by (6.9), (6.10), and Lemmas 3.7 and 3.9 it follows that $\delta(\beta)$ also begins with

$$\begin{aligned} \mathbb{L}(\mathbf{S}_{k+2})^+ &= \mathbb{L}(\mathbf{S}_k \bullet (01 \bullet 01))^+ = \mathbb{L}(\mathbf{S}_k \bullet (0011))^+ \\ &= (\mathbf{S}_k \bullet \mathbb{L}(0011))^+ = (\mathbf{S}_k \bullet (1100))^+ = \mathbb{L}(\mathbf{S}_k)^+ \mathbf{S}_k^- \mathbb{L}(\mathbf{S}_k)^+, \end{aligned}$$

which is strictly smaller than a prefix of $\delta(\beta_r^{\mathbf{S}_k})^+ = \mathbb{L}(\mathbf{S}_k)^+ \mathbf{S}_k^\infty$. This implies that $\beta < \beta_r^{\mathbf{S}_k}$ for all $k \geq 1$. Hence, $\beta \in J^{\mathbf{S}_k}$ for all $k \geq 1$, and thus $\beta \in E_\infty \cap J^{\mathbf{s}}$. Furthermore, by (6.9),

(6.10) and Proposition 6.3 it follows that

$$\begin{aligned}
\tau(\beta) &= \lim_{k \rightarrow \infty} (\mathbf{S}_k 0^\infty)_\beta = (\overline{\theta_1 \mathbf{c} \theta_2 \theta_3 \mathbf{c} \theta_4 \dots})_\beta \\
&= \sum_{k=0}^{\infty} \left(\frac{1}{\beta^{mk+1}} + 2 \sum_{j=2}^{m-1} \frac{s_j}{\beta^{mk+j}} + \frac{1}{\beta^{mk+m}} \right) - (\theta_1 \mathbf{c} \theta_2 \theta_3 \mathbf{c} \theta_4 \dots)_\beta \\
&= \frac{\beta^{m-1} + 2 \sum_{j=2}^{m-1} s_j \beta^{m-j} + 1}{\beta^m - 1} - 1 \\
&= \frac{2 \sum_{j=2}^m s_j \beta^{m-j} + \beta^{m-1} - \beta^m}{\beta^m - 1},
\end{aligned}$$

where we recall that $\mathbf{c} = s_2 \dots s_{m-1}$, and the last equality follows by using $s_m = 1$.

Finally, the transcendence of β follows by using (6.9), Lemma 6.5 and a similar argument as in the proof of [22, Proposition 5.2]. \square

Remark 6.6.

- When $\mathbf{s} = 01$, the base $\beta_\infty^{01} \approx 1.78723$ given in Proposition 6.4 is the *Komornik-Loreti constant* (cf. [21]), whose transcendence was first proved by Allouche and Cosnard [3]. In this case we obtain $\tau(\beta_\infty^{01}) = (2 - \beta_\infty^{01})/(\beta_\infty^{01} - 1) \approx .270274$.
- When $\mathbf{s} = 001$, the base $\beta_\infty^{001} \approx 1.55356$ is a critical value for the fat Sierpinski gaskets studied by Li and the second author in [22]. In this case we have $\tau(\beta_\infty^{001}) \approx .241471$.

By Proposition 6.4 and numerical calculation we give in Table 1 the triples $(\mathbf{s}, \beta_\infty^{\mathbf{s}}, \tau(\beta_\infty^{\mathbf{s}}))$ for all $\mathbf{s} \in F_3^* \subset \Omega_F^*$. Based on Proposition 6.4 we conjecture that each base $\beta \in E_\infty$ is transcendental.

$\mathbf{s} =$	0001	001	00101	01	01011	011	0111
$\beta_\infty^{\mathbf{s}} \approx$	1.43577	1.55356	1.59998	1.78723	1.83502	1.91988	1.96452
$\tau(\beta_\infty^{\mathbf{s}}) \approx$	0.218562	0.241471	0.336114	0.270274	0.432175	0.40305	0.455933

TABLE 1. The triples $(\mathbf{s}, \beta_\infty^{\mathbf{s}}, \tau(\beta_\infty^{\mathbf{s}}))$ with $\mathbf{s} \in F_3^* \subset \Omega_F^*$.

7. CÀDLÀG PROPERTY OF THE CRITICAL VALUE FUNCTION

In this section we prove Proposition 1.9 and Theorem 1. Recall by Lemma 5.7 that the interval $(1, 2]$ can be partitioned as

$$(7.1) \quad (1, 2] = E \cup \bigcup_{\mathbf{s} \in \Omega_F^*} J^{\mathbf{s}} = E \cup E_\infty \cup \bigcup_{\mathbf{S} \in \Lambda} E^{\mathbf{S}} \cup \bigcup_{\mathbf{S} \in \Lambda} I^{\mathbf{S}}.$$

Here we emphasize that the bifurcation set E , the relative bifurcation sets $E^{\mathbf{S}}$ and the infinite bifurcation sets E_∞ featuring in (7.1) all have Lebesgue measure zero in view of Propositions 5.6 and 5.8. Hence the basic intervals $I^{\mathbf{S}}$, $\mathbf{S} \in \Lambda$, and then certainly the Lyndon intervals $J^{\mathbf{S}}$, $\mathbf{S} \in \Lambda$, are dense in $(1, 2]$. This allows for approximation of points in E , $E^{\mathbf{S}}$ and E_∞ by left and/or right endpoints of such Lyndon intervals. We also recall from Theorem 2 that

for any basic interval $I^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_*^{\mathbf{S}}]$ with $\delta(\beta_\ell^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^\infty$ and $\delta(\beta_*^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^- \mathbb{L}(\mathbf{S})^\infty$, the critical value is given by

$$(7.2) \quad \tau(\beta) = (\Phi_{\mathbf{S}}(0^\infty))_\beta = (\mathbf{S}^- \mathbb{L}(\mathbf{S})^\infty)_\beta \quad \text{for any } \beta \in I^{\mathbf{S}}.$$

And, by Proposition 6.2 it follows that for each $\beta \in E^{\mathbf{S}}$ we have

$$(7.3) \quad \tau(\beta) = (\Phi_{\mathbf{S}}(0\delta_2\delta_3\dots))_\beta,$$

where $1\delta_2\delta_3\dots = \delta(\hat{\beta})$ with $\hat{\beta} = \Psi_{\mathbf{S}}^{-1}(\beta) \in E$. In particular, when $\beta \in E$ we have $\tau(\beta) = 1 - \frac{1}{\beta}$ (see Proposition 1.12). When $\beta \in E_\infty$, it follows by Proposition 6.3 that

$$(7.4) \quad \tau(\beta) = \lim_{n \rightarrow \infty} (\mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \dots \bullet \mathbf{s}_n 0^\infty)_\beta,$$

where (\mathbf{s}_k) is the unique coding of β (that is $\beta \in J^{\mathbf{s}_1 \bullet \dots \bullet \mathbf{s}_k}$ for all $k \in \mathbb{N}$).

From (7.2) it is clear that the critical value function τ is continuous inside each basic interval $I^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_*^{\mathbf{S}}]$. So, in view of (7.1), we still need to consider the continuity of τ for $\beta \in E \cup E_\infty \cup \bigcup_{\mathbf{S} \in \Lambda} (E^{\mathbf{S}} \setminus \{\beta_\ell^{\mathbf{S}}\})$, the left continuity of τ at $\beta = \beta_\ell^{\mathbf{S}}$ and $\beta = \beta_r^{\mathbf{S}}$, and the right continuity of τ at $\beta = \beta_*^{\mathbf{S}}$. We need the following lemma:

Lemma 7.1. *If $\beta \in (1, 2)$ and $\delta(\beta)$ is periodic, then $\beta \in \bigcup_{\mathbf{S} \in \Lambda} I^{\mathbf{S}}$.*

Proof. Assume $\delta(\beta)$ is periodic. In view of (7.1) it suffices to prove

$$\beta \notin E \cup E_\infty \cup \bigcup_{\mathbf{S} \in \Lambda} E^{\mathbf{S}}.$$

First we prove $\beta \notin E$. By Lemma 2.8, $\delta(\beta) = \mathbb{L}(\mathbf{S}')^\infty$ for some Lyndon word \mathbf{S}' . This means β is the left endpoint of a Lyndon interval, so by Lemma 2.9, $\beta \in J^{\mathbf{S}}$ for some Farey word \mathbf{s} . Hence $\beta \notin E$.

Next, suppose $\beta \in E^{\mathbf{S}}$ for some $\mathbf{S} \in \Lambda$. Clearly, $\beta \neq \beta_r^{\mathbf{S}}$ since $\delta(\beta_r^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty$ is not periodic. Thus $\beta \in (\beta_*^{\mathbf{S}}, \beta_r^{\mathbf{S}}) \setminus \bigcup_{r \in \Omega_F^*} J^{\mathbf{S} \bullet \mathbf{r}}$. So, by Proposition 5.6 there is a sequence (\mathbf{r}_k) of Farey words such that $\beta_\ell^{\mathbf{S} \bullet \mathbf{r}_k} \searrow \beta$. Write $\delta(\beta) = (a_1 \dots a_n)^\infty$ with minimal period n . Then by Lemma 2.5 it follows that $\delta(\beta_\ell^{\mathbf{S} \bullet \mathbf{r}_k}) \searrow a_1 \dots a_n^+ 0^\infty$, so for all sufficiently large k , $\delta(\beta_\ell^{\mathbf{S} \bullet \mathbf{r}_k})$ contains a block of more than $2m$ zeros, where $m := |\mathbf{S}|$. But this is impossible, since $\delta(\beta_\ell^{\mathbf{S} \bullet \mathbf{r}_k}) = \mathbb{L}(\mathbf{S} \bullet \mathbf{r}_k)^\infty$ is a concatenation of blocks from $\mathbf{S}, \mathbf{S}^-, \mathbb{L}(\mathbf{S})$ and $\mathbb{L}(\mathbf{S})^+$. These blocks all have length m , and only \mathbf{S}^- could possibly consist of all zeros, while \mathbf{S}^- can only be followed by $\mathbb{L}(\mathbf{S})$ or $\mathbb{L}(\mathbf{S})^+$. Thus, $\delta(\beta_\ell^{\mathbf{S} \bullet \mathbf{r}_k})$ cannot contain a block of $2m$ zeros. This contradiction shows that $\beta \notin E^{\mathbf{S}}$.

Finally, suppose $\beta \in E_\infty$. Then there is a sequence (\mathbf{S}_k) of words in Λ such that β lies in the interior of $J^{\mathbf{S}_k}$ for each k . Note that $\beta = \beta_\ell^{\mathbf{S}'}$ is the left endpoint of $J^{\mathbf{S}'}$. Thus, $J^{\mathbf{S}'} \cap J^{\mathbf{S}_k} \neq \emptyset$, and therefore by Lemma 2.9, it must be the case that $J^{\mathbf{S}'} \subset J^{\mathbf{S}_k}$ for all k . But this is impossible, since $|J^{\mathbf{S}_k}| \rightarrow 0$. Hence, $\beta \notin E_\infty$. \square

Proof of Proposition 1.9. First fix $\beta_0 \in (1, 2] \setminus \{\beta_r^{\mathbf{S}} : \mathbf{S} \in \Lambda\}$. It is sufficient to prove that

- (*) for each $N \in \mathbb{N}$ there exists $r > 0$ such that if $\beta \in (1, 2]$ satisfies $|\beta - \beta_0| < r$, then there is a word $s_1 \dots s_N$ such that $\tau(\beta)$ has a β -expansion beginning with $s_1 \dots s_N$, and $\tau(\beta_0)$ has a β_0 -expansion beginning with $s_1 \dots s_N$.

For, if $\tau(\beta) = (s_1 \dots s_N c_1 c_2 \dots)_\beta$ and $\tau(\beta_0) = (s_1 \dots s_N d_1 d_2 \dots)_{\beta_0}$, then

$$\begin{aligned} |\tau(\beta) - \tau(\beta_0)| &\leq |(s_1 \dots s_N c_1 c_2 \dots)_\beta - (s_1 \dots s_N c_1 c_2 \dots)_{\beta_0}| \\ &\quad + |(s_1 \dots s_N c_1 c_2 \dots)_{\beta_0} - (s_1 \dots s_N d_1 d_2 \dots)_{\beta_0}| \\ &\leq \sum_{i=1}^{\infty} \left| \frac{1}{\beta^i} - \frac{1}{\beta_0^i} \right| + \sum_{i=N+1}^{\infty} \frac{1}{\beta_0^i} \\ &= \frac{|\beta - \beta_0|}{(\beta - 1)(\beta_0 - 1)} + \frac{1}{(\beta_0 - 1)\beta_0^N} \\ &< \frac{r}{(\beta - 1)(\beta_0 - 1)} + \frac{1}{(\beta_0 - 1)\beta_0^N}, \end{aligned}$$

and this can be made as small as desired by choosing N sufficiently large and r sufficiently small. In view of (7.1) we prove (*) by considering several cases.

Case 1. $\beta_0 \in (\beta_\ell^{\mathbf{S}}, \beta_r^{\mathbf{S}})$ for some basic interval $I^{\mathbf{S}} = [\beta_\ell^{\mathbf{S}}, \beta_r^{\mathbf{S}}]$ with $\mathbf{S} \in \Lambda$. It is clear from Theorem 2 that (*) holds in this case.

Case 2. $\beta_0 \in E$. Then by Proposition 5.6 there exists a sequence of Farey intervals $J^{\mathbf{s}_k} = [\beta_\ell^{\mathbf{s}_k}, \beta_r^{\mathbf{s}_k}]$ such that $\beta_\ell^{\mathbf{s}_k} \rightarrow \beta_0$ as $k \rightarrow \infty$. Furthermore, $|J^{\mathbf{s}_k}| \rightarrow 0$ as $k \rightarrow \infty$. This implies that the length $|\mathbf{s}_k|$ of the Farey word \mathbf{s}_k goes to infinity as $k \rightarrow \infty$. Let $N \in \mathbb{N}$ be given. We can choose $r > 0$ small enough so that if a Farey interval $J^{\mathbf{s}}$ intersects $(\beta_0 - r, \beta_0 + r)$, then $|\mathbf{s}| > N$ and

$$(7.5) \quad \delta_1(\beta) \dots \delta_N(\beta) = \delta_1(\beta_0) \dots \delta_N(\beta_0).$$

We can guarantee (7.5) because for $\beta_0 \in E \setminus \{2\}$ the expansion $\delta(\beta_0)$ is not periodic by Lemma 7.1, so by Lemma 2.5 the map $\beta \mapsto \delta(\beta)$ is continuous at β_0 . Furthermore, for $\beta_0 = 2$ the map $\beta \mapsto \delta(\beta)$ is left continuous at β_0 .

Now take $\beta \in (\beta_0 - r, \beta_0 + r)$. By (7.1) we have either $\beta \in E$ or $\beta \in J^{\mathbf{s}}$ for some $\mathbf{s} \in \Omega_F^*$. If $\beta \in E$, then by Proposition 1.12 it follows that $\tau(\beta) = 1 - \frac{1}{\beta} = (0\delta_2(\beta)\delta_3(\beta) \dots)_\beta$ and

$$(7.6) \quad \tau(\beta_0) = 1 - \frac{1}{\beta_0} = (0\delta_2(\beta_0)\delta_3(\beta_0) \dots)_{\beta_0},$$

so (*) holds by (7.5).

Next we assume $\beta \in J^{\mathbf{s}}$ with $\mathbf{s} = s_1 \dots s_m \in \Omega_F^*$. By our choice of r it follows that $m = |\mathbf{s}| > N$, and (7.5) holds. Since $\beta \in J^{\mathbf{s}} = [\beta_\ell^{\mathbf{s}}, \beta_r^{\mathbf{s}}]$, we have $\mathbb{L}(\mathbf{s})^\infty \preccurlyeq \delta(\beta) \preccurlyeq \mathbb{L}(\mathbf{s})^+ \mathbf{s}_\infty$. By (7.5) it follows that

$$(7.7) \quad \delta_1(\beta) \dots \delta_N(\beta) = \delta_1(\beta_0) \dots \delta_N(\beta_0) = a_1 \dots a_N,$$

where $a_1 \dots a_m = \mathbb{L}(\mathbf{s})$. Observe by (7.1)–(7.4) that $\tau(\beta)$ has a β expansion beginning with $\mathbf{s}_- = 0a_2 \dots a_m$, where the equality follows by Lemma 2.3. So, by (7.7) it follows that $\tau(\beta)$ has a β -expansion with prefix

$$(7.8) \quad s_1 \dots s_N = 0a_2 \dots a_N = 0\delta_2(\beta_0) \dots \delta_N(\beta_0).$$

This, together with (7.6), gives (*).

Case 3. $\beta_0 \in E^{\mathbf{S}} \setminus \{\beta_r^{\mathbf{S}}\}$ for some $\mathbf{S} \in \Lambda$. The proof is similar to that of Case 2. By Proposition 5.6 it follows that

$$(7.9) \quad \beta \in (\beta_*^{\mathbf{S}}, \beta_r^{\mathbf{S}}) \setminus \bigcup_{\mathbf{r} \in \Omega_F^*} J^{\mathbf{S}\bullet\mathbf{r}},$$

and there exists a sequence (\mathbf{r}_k) of Farey words such that $\beta_{\ell}^{\mathbf{S}\bullet\mathbf{r}_k} \rightarrow \beta$ as $k \rightarrow \infty$. This implies that $|\mathbf{r}_k| \rightarrow \infty$ as $k \rightarrow \infty$.

Let $N \in \mathbb{N}$ be given; without loss of generality we may assume that $N = N'|\mathbf{S}|$ for some integer N' . By Lemma 7.1 we can choose $r > 0$ sufficiently small so that if a Lyndon interval $J^{\mathbf{S}\bullet\mathbf{r}}$ intersects $(\beta_0 - r, \beta_0 + r)$, then $|\mathbf{r}| > N'$ and

$$(7.10) \quad \delta_1(\beta) \dots \delta_{N'|\mathbf{S}|}(\beta) = \delta_1(\beta_0) \dots \delta_{N'|\mathbf{S}|}(\beta_0) = \mathbf{S} \bullet (\delta_1(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0))$$

for any $\beta \in (\beta_0 - r, \beta_0 + r)$, where $\hat{\beta}_0 = \Psi_{\mathbf{S}}^{-1}(\beta_0) \in E$.

Now take $\beta \in (\beta_0 - r, \beta_0 + r)$. By (7.9) it follows that either $\beta \in E^{\mathbf{S}}$ or $\beta \in J^{\mathbf{S}\bullet\mathbf{r}}$ for some $\mathbf{r} \in \Omega_F^*$. If $\beta \in E^{\mathbf{S}}$, we let $\hat{\beta} = \Psi_{\mathbf{S}}^{-1}(\beta)$. Then by (7.10) it follows that

$$\mathbf{S} \bullet (\delta_1(\hat{\beta}) \dots \delta_{N'}(\hat{\beta})) = \delta_1(\beta) \dots \delta_{N'|\mathbf{S}|}(\beta) = \mathbf{S} \bullet (\delta_1(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0)),$$

which implies $\delta_1(\hat{\beta}) \dots \delta_{N'}(\hat{\beta}) = \delta_1(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0)$. So, by (7.3) it follows that $\tau(\beta)$ has a β -expansion with a prefix

$$\mathbf{S} \bullet (0\delta_2(\hat{\beta}) \dots \delta_{N'}(\hat{\beta})) = \mathbf{S} \bullet (0\delta_2(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0)),$$

which coincides with a prefix of a β_0 -expansion of $\tau(\beta_0)$. This yields (*).

If $\beta \in J^{\mathbf{S}\bullet\mathbf{r}}$ for some $\mathbf{r} = r_1 \dots r_m \in \Omega_F^*$, then by our assumption on r we have $m > N'$. Furthermore,

$$(\mathbf{S} \bullet \mathbb{L}(\mathbf{r}))^\infty = \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^\infty \preceq \delta(\beta) \preceq \mathbb{L}(\mathbf{S} \bullet \mathbf{r})^+(\mathbf{S} \bullet \mathbf{r})^\infty = (\mathbf{S} \bullet \mathbb{L}(\mathbf{r}))^+(\mathbf{S} \bullet \mathbf{r})^\infty.$$

So, if we write $\mathbb{L}(\mathbf{r}) = b_1 \dots b_m$, then by (7.10) it follows that

$$(7.11) \quad \mathbf{S} \bullet (b_1 \dots b_{N'}) = \delta_1(\beta) \dots \delta_{N'|\mathbf{S}|}(\beta) = \mathbf{S} \bullet (\delta_1(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0)).$$

From (7.11) we obtain that $b_1 \dots b_{N'} = \delta_1(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0)$. Observe by (7.1)–(7.4) and Lemma 2.3 that $\tau(\beta)$ has a β -expansion beginning with $(\mathbf{S} \bullet \mathbf{r})^- = \mathbf{S} \bullet (0b_2 \dots b_m)$. So, by (7.3) and (7.11) it follows that $\tau(\beta)$ has a β -expansion with a prefix

$$\mathbf{S} \bullet (0b_2 \dots b_{N'}) = \mathbf{S} \bullet (0\delta_2(\hat{\beta}_0) \dots \delta_{N'}(\hat{\beta}_0))$$

of length N , which is also a prefix of a β_0 -expansion of $\tau(\beta_0)$. This again gives (*).

Case 4. $\beta_0 = \beta_{\ell}^{\mathbf{S}}$ for some $\mathbf{S} \in \Lambda$. Here the right continuity of τ at β_0 follows from Theorem 2. The left-continuity follows from the arguments in Case 2 and Case 3 and the left-continuity of the map $\beta \mapsto \delta(\beta)$ at β_0 : If $\mathbf{S} \in \Omega_F^*$, then $\tau(\beta_{\ell}^{\mathbf{S}})$ has a $\beta_{\ell}^{\mathbf{S}}$ -expansion $0\delta_2(\beta_{\ell}^{\mathbf{S}})\delta_3(\beta_{\ell}^{\mathbf{S}})\dots$. Otherwise, $\mathbf{S} = \mathbf{S}' \bullet \mathbf{r}$ for some $\mathbf{S}' \in \Lambda$ and $\mathbf{r} \in \Omega_F^*$, and $\tau(\beta_{\ell}^{\mathbf{S}})$ has a $\beta_{\ell}^{\mathbf{S}}$ -expansion $\Phi_{\mathbf{S}'}(0\delta_2(\beta_{\ell}^{\mathbf{r}})\delta_3(\beta_{\ell}^{\mathbf{r}})\dots)$.

Case 5. $\beta_0 = \beta_*^{\mathbf{S}}$ for some $\mathbf{S} \in \Lambda$. Here the left continuity at β_0 follows from Theorem 2. The right-continuity can be seen as follows. First note that $\delta(\beta_*^{\mathbf{S}}) = \mathbb{L}(\mathbf{S})^+\mathbf{S}^-\mathbb{L}(\mathbf{S})^\infty$, and $\tau(\beta_*^{\mathbf{S}}) = (\mathbf{S}^-\mathbb{L}(\mathbf{S})^\infty)_{\beta_*^{\mathbf{S}}}$. Observe also that $\beta_*^{\mathbf{S}} = \lim_{\hat{\beta} \searrow 1} \Psi_{\mathbf{S}}(\hat{\beta})$, so by Lemma 2.5, $\Phi_{\mathbf{S}}(\delta(\hat{\beta})) \searrow \delta(\beta_*^{\mathbf{S}})$ as $\hat{\beta} \searrow 1$.

Now let $N \in \mathbb{N}$ be given, and assume without loss of generality that $N = N'|\mathbf{S}|$ for some integer N' . We choose $r > 0$ small enough so that, if $\mathbf{r} \in \Omega_F^*$ and $J^{\mathbf{S}\bullet\mathbf{r}}$ intersects $(\beta_0, \beta_0 + r)$,

then $|\mathbf{r}| > N'$. Now take $\beta \in (\beta_0, \beta_0 + r)$. If $\beta \in E^{\mathbf{S}}$, then $\beta = \Psi_{\mathbf{S}}(\hat{\beta})$ for some $\hat{\beta} \in E$, and $\delta(\beta) = \Phi_{\mathbf{S}}(\delta(\hat{\beta}))$. Note that, if $\beta \searrow \beta_0$, then $\hat{\beta} \searrow 1$ and so $\delta(\hat{\beta}) \searrow 10^\infty$. Thus, we may also assume r is small enough so that $\delta(\hat{\beta})$ begins with $10^{N'-1}$. Then $\tau(\beta)$ has a β -expansion beginning with $\Phi_{\mathbf{S}}(0^{N'}) = \mathbf{S}^{-\mathbb{L}(\mathbf{S})^{N'-1}}$, which is also a prefix of length N of a β_0 -expansion of $\tau(\beta_0)$.

Similarly, if $\beta \in J^{\mathbf{S} \bullet \mathbf{r}}$ then we may assume r is small enough so that \mathbf{r} begins with $0^{N'-1}$. Then $\tau(\beta)$ has a β -expansion beginning with $(\mathbf{S} \bullet \mathbf{r})^-$, and therefore beginning with $\mathbf{S}^{-\mathbb{L}(\mathbf{S})^{N'-1}}$, and we conclude as above.

Case 6. $\beta_0 \in E_\infty$. Then there exists a sequence (\mathbf{s}_k) of Farey words such that

$$\{\beta\} = \bigcap_{k=1}^{\infty} J^{\mathbf{S}_k},$$

where $\mathbf{S}_k := \mathbf{s}_1 \bullet \mathbf{s}_2 \bullet \dots \bullet \mathbf{s}_k$. Note that $J^{\mathbf{S}_k} \supset J^{\mathbf{S}_{k+1}}$ for all $k \geq 1$, and $|\mathbf{S}_k| \rightarrow \infty$ as $k \rightarrow \infty$.

Let $N \in \mathbb{N}$ be given, and choose k so large that $|\mathbf{S}_k| > N$. Choose $r > 0$ sufficiently small so that $(\beta_0 - r, \beta_0 + r) \subset J^{\mathbf{S}_k}$. Take $\beta \in (\beta_0 - r, \beta_0 + r)$. Then $\beta \in J^{\mathbf{S}_k}$, so by (7.1)–(7.4) it follows that $\tau(\beta)$ has a β -expansion beginning with \mathbf{S}_k^- , which is also a prefix (of length at least N) of a β_0 -expansion of $\tau(\beta_0)$. Hence, we obtain (*).

Finally, we consider $\beta_0 = \beta_r^{\mathbf{S}}$ for $\mathbf{S} \in \Lambda$. The left continuity at β_0 (i.e. the analog of (*) for $\beta \in (\beta_0 - r, \beta_0)$) follows just as in Case 3. The jump at β_0 (i.e. (1.6)) can be seen as follows. Note by Proposition 6.2 that $\tau(\beta_0) = (\mathbf{S}0^\infty)_{\beta_0} < (\mathbf{S}^\infty)_{\beta_0}$. Thus it suffices to prove

$$(7.12) \quad \lim_{\beta \searrow \beta_0} \tau(\beta) = (\mathbf{S}^\infty)_{\beta_0}.$$

First we assume $\mathbf{S} \in \Omega_F^*$. Then $\delta(\beta_0) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty$, and by Lemma 2.3 it follows that $\mathbf{S}^\infty = 0\delta_2(\beta_0)\delta_3(\beta_0)\dots$. So, by the same argument as in Case 2 we obtain (7.12).

Next suppose $\mathbf{S} = \mathbf{S}' \bullet \mathbf{r}$ for some $\mathbf{S}' \in \Lambda$ and $\mathbf{r} \in \Omega_F^*$. Then

$$\delta(\beta_0) = \mathbb{L}(\mathbf{S})^+ \mathbf{S}^\infty = (\mathbf{S}' \bullet \mathbb{L}(\mathbf{r}))^+ (\mathbf{S}' \bullet \mathbf{r})^\infty = \Phi_{\mathbf{S}'}(\mathbb{L}(\mathbf{r})^+ \mathbf{r}_\infty) = \Phi_{\mathbf{S}'}(\delta(\hat{\beta}_0)),$$

where $\hat{\beta}_0 = \Psi_{\mathbf{S}'}^{-1}(\beta_0) \in E$. This implies that

$$(7.13) \quad \Phi_{\mathbf{S}'}(0\delta_2(\hat{\beta}_0)\delta_3(\hat{\beta}_0)\dots) = \Phi_{\mathbf{S}'}(\mathbf{r}_\infty) = (\mathbf{S}' \bullet \mathbf{r})^\infty = \mathbf{S}^\infty.$$

By (7.13) and the same argument as in Case 3 we again obtain (7.12). \square

Proof of Theorem 1. The theorem follows by Proposition 1.9, Theorem 2 and Theorem 3. \square

ACKNOWLEDGEMENTS

The first author is partially supported by Simons Foundation grant # 709869. The second author thanks Charlene Kalle for some useful discussions when she visited Chongqing University in November 2018. He is supported by NSFC No. 11971079 and the Fundamental and Frontier Research Project of Chongqing No. cstc2019jcyj-msxmX0338 and No. cx2019067.

REFERENCES

- [1] P. Allaart and D. Kong. Relative bifurcation sets and the local dimension of univoque bases. *Ergodic Theory Dynam. Systems*, 41: 2241–2273, 2021.
- [2] P. C. Allaart. An algebraic approach to entropy plateaus in non-integer base expansions. *Discrete Contin. Dyn. Syst.*, 39(11):6507–6522, 2019.

- [3] J.-P. Allouche and M. Cosnard. The Komornik-Loreti constant is transcendental. *Amer. Math. Monthly*, 107(5):448–449, 2000.
- [4] J.-P. Allouche and J. Shallit. The ubiquitous Prouhet-Thue-Morse sequence. In *Sequences and their applications (Singapore, 1998)*, Springer Ser. Discrete Math. Theor. Comput. Sci., pages 1–16. Springer, London, 1999.
- [5] J.-P. Allouche and J. Shallit. *Automatic sequences. Theory, applications, generalizations*. Cambridge University Press, Cambridge, 2003.
- [6] C. Baiocchi and V. Komornik. Greedy and quasi-greedy expansions in non-integer bases. *arXiv:0710.3001v1*, 2007.
- [7] L. A. Bunimovich and A. Yurchenko. Where to place a hole to achieve a maximal escape rate. *Israel J. Math.*, 182:229–252, 2011.
- [8] C. Carminati, S. Isola, and G. Tiozzo. Continued fractions with $SL(2, Z)$ -branches: combinatorics and entropy. *Trans. Amer. Math. Soc.*, 370(7):4927–4973, 2018.
- [9] C. Carminati and G. Tiozzo. The local Hölder exponent for the dimension of invariant subsets of the circle. *Ergodic Theory Dynam. Systems*, 37(6):1825–1840, 2017.
- [10] L. Clark. The β -transformation with a hole. *Discrete Contin. Dyn. Syst.*, 36(3):1249–1269, 2016.
- [11] Z. Daróczy and I. Kátai. Univoque sequences. *Publ. Math. Debrecen*, 42(3-4):397–407, 1993.
- [12] M. Demers, P. Wright, and L.-S. Young. Escape rates and physically relevant measures for billiards with small holes. *Comm. Math. Phys.*, 294(2):353–388, 2010.
- [13] M. F. Demers. Markov extensions for dynamical systems with holes: an application to expanding maps of the interval. *Israel J. Math.*, 146:189–221, 2005.
- [14] M. F. Demers and L.-S. Young. Escape rates and conditionally invariant measures. *Nonlinearity*, 19(2):377–397, 2006.
- [15] M. de Vries and V. Komornik. A two-dimensional univoque set. *Fund. Math.*, 212(2):175–189, 2011.
- [16] K. Falconer. *Fractal geometry: Mathematical foundations and applications*. John Wiley & Sons Ltd., Chichester, 1990.
- [17] K. Falconer. *Techniques in fractal geometry*. John Wiley & Sons Ltd., Chichester, 1997.
- [18] P. Glendinning and N. Sidorov. The doubling map with asymmetrical holes. *Ergodic Theory Dynam. Systems*, 35(4):1208–1228, 2015.
- [19] E. Hille. A Problem in “Factorisatio Numerorum.” *Acta Arith.*, 2:134–144, 1936.
- [20] C. Kalle, D. Kong, N. Langeveld, and W. Li. The β -transformation with a hole at 0. *Ergodic Theory Dynam. Systems*, 40(9):2482–2514, 2020.
- [21] V. Komornik and P. Loreti, Unique developments in noninteger bases, *Amer. Math. Monthly*, 105 (1998), 636–639.
- [22] D. Kong and W. Li, Critical base for the unique codings of fat Sierpinski gasket, *Nonlinearity*, 33(9): 4484–4511, 2020.
- [23] D. Lind and B. Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995.
- [24] M. Lothaire. *Algebraic combinatorics on words*, volume 90 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2002.
- [25] K. Mahler. *Lectures on transcendental numbers*. Lecture Notes in Mathematics, Vol. 546. Springer-Verlag, Berlin, 1976.
- [26] W. Parry. On the β -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [27] G. Pianigiani and J. A. Yorke. Expanding maps on sets which are almost invariant. Decay and chaos. *Trans. Amer. Math. Soc.*, 252:351–366, 1979.
- [28] P. Raith. Continuity of the Hausdorff dimension for invariant subsets of interval maps. *Acta Math. Univ. Comenian. (N.S.)*, 63(1):39–53, 1994.
- [29] M. Urbański. On Hausdorff dimension of invariant sets for expanding maps of a circle. *Ergodic Theory Dynam. Systems*, 6(2):295–309, 1986.
- [30] M. Urbański. Invariant subsets of expanding mappings of the circle. *Ergodic Theory Dynam. Systems*, 7(4):627–645, 1987.

(P. Allaart) MATHEMATICS DEPARTMENT, UNIVERSITY OF NORTH TEXAS, 1155 UNION CIR #311430,
DENTON, TX 76203-5017, U.S.A.

Email address: `allaart@unt.edu`

(D. Kong) COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, CHONGQING 401331,
PEOPLE'S REPUBLIC OF CHINA.

Email address: `derongkong@126.com`