

ON A CLASS OF SELF-SIMILAR SETS WHICH CONTAIN FINITELY MANY COMMON POINTS

ZHIQIANG WANG, KAN JIANG, DERONG KONG, AND WENXIA LI

ABSTRACT. For $\lambda \in (0, 1/2]$ let $K_\lambda \subset \mathbb{R}$ be a self-similar set generated by the iterated function system $\{\lambda x, \lambda x + 1 - \lambda\}$. Given $x \in (0, 1/2)$, let $\Lambda(x)$ be the set of $\lambda \in (0, 1/2]$ such that $x \in K_\lambda$. In this paper we show that $\Lambda(x)$ is a topological Cantor set having zero Lebesgue measure and full Hausdorff dimension. Furthermore, we show that for any $y_1, \dots, y_p \in (0, 1/2)$ there exists a full Hausdorff dimensional set of $\lambda \in (0, 1/2]$ such that y_1, \dots, y_p are common points of K_λ .

1. INTRODUCTION

For $\lambda \in (0, 1/2]$ let K_λ be the *self-similar set* generated by the *iterated function system* (simply called, *IFS*) $\{f_{\lambda,d}(x) = \lambda x + d(1 - \lambda) : d = 0, 1\}$. Then K_λ is the unique nonempty compact set satisfying (cf. [3])

$$(1.1) \quad K_\lambda = f_{\lambda,0}(K_\lambda) \cup f_{\lambda,1}(K_\lambda) = \left\{ (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1} : i_n \in \{0, 1\} \ \forall n \geq 1 \right\}.$$

It is obvious that the convex hull of K_λ is the unit interval $[0, 1]$ for all $\lambda \in (0, 1/2]$. Then 0 and 1 are common points of K_λ for all $\lambda \in (0, 1/2]$. For other $x \in (0, 1)$ it is natural to ask how likely the self-similar sets $K_\lambda, \lambda \in (0, 1/2]$ contain the common point x ? Or even ask how likely the self-similar sets $K_\lambda, \lambda \in (0, 1/2]$ contain any given points $y_1, \dots, y_p \in (0, 1)$? These questions are motivated by the work of Boes, Darst and Erdős [2], in which they considered a class of fat Cantor sets C_λ with positive Lebesgue measure. They showed that for a given point $x \in (0, 1)$ the set of parameters $\lambda \in (0, 1/2)$ such that $x \in C_\lambda$ is of first category.

Given $x \in [0, 1]$, let

$$(1.2) \quad \Lambda(x) := \{\lambda \in (0, 1/2] : x \in K_\lambda\}.$$

Then $\Lambda(x)$ consists of all $\lambda \in (0, 1/2]$ such that x is the common point of K_λ . Note that K_λ is symmetric, i.e., $x \in K_\lambda$ if and only if $1 - x \in K_\lambda$. Then $\Lambda(x) = \Lambda(1 - x)$ for any $x \in [0, 1]$. So, we only need to consider $x \in [0, 1/2]$. Note that $\Lambda(0) = (0, 1/2]$; and $\Lambda(1/2) = \{1/2\}$, because $1/2 \in K_\lambda$ if and only if $\lambda = 1/2$. So, it is interesting to study $\Lambda(x)$ for $x \in (0, 1/2)$.

Recall that a set $F \subset \mathbb{R}$ is called a *Cantor set* if it is a non-empty compact set containing neither interior nor isolated points. Our first result considers the topology of $\Lambda(x)$.

2010 *Mathematics Subject Classification*. Primary: 28A78; Secondary: 28A80, 37B10.

Key words and phrases. Hausdorff dimension, thickness, self-similar set, Cantor set, intersection.

Theorem 1.1. *For any $x \in (0, 1/2)$ the set $\Lambda(x)$ is a Cantor set with $\min \Lambda(x) = x$ and $\max \Lambda(x) = 1/2$.*

By Theorem 1.1 it follows that $\Lambda(x)$ is a fractal set for any $x \in (0, 1/2)$. Our next result considers $\Lambda(x)$ from the Lebesgue measure and fractal dimension perspectives.

Theorem 1.2. *For any $x \in (0, 1/2)$ the set $\Lambda(x)$ is a Lebesgue null set of full Hausdorff dimension. Furthermore,*

$$\lim_{\delta \rightarrow 0^+} \dim_H(\Lambda(x) \cap (\lambda - \delta, \lambda + \delta)) = \frac{\log 2}{-\log \lambda} \quad \forall \lambda \in \Lambda(x),$$

where \dim_H denotes the Hausdorff dimension.

Since $\Lambda(1/2) = \{1/2\}$ and $\Lambda(1-x) = \Lambda(x)$ for any $x \in (0, 1/2)$, by Theorem 1.2 it follows that $\Lambda(x)$ has zero Lebesgue measure for any $x \in (0, 1)$. Then the union $\bigcup_{x \in \mathbb{Q} \cap (0, 1)} \Lambda(x)$ also has zero Lebesgue measure. Taking the complement we obtain the following result which is also interesting in number theory.

Corollary 1.3. *For Lebesgue almost every $\lambda \in (0, 1/2]$ the set $K_\lambda \setminus \{0, 1\}$ contains only irrational numbers.*

Given $y_1, \dots, y_p \in (0, 1/2)$, by Theorems 1.1 and 1.2 it follows that the intersection $\bigcap_{i=1}^p \Lambda(y_i)$ is small from the topological and Lebesgue measure perspectives. Furthermore, by using the *thickness* method introduced by Newhouse [8] we show that $\bigcap_{i=1}^p \Lambda(y_i)$ contains a sequence of Cantor sets whose thickness can be arbitrarily large. From this we conclude that there exists a full Hausdorff dimensional set of $\lambda \in (0, 1/2]$ such that $y_1, \dots, y_p \in K_\lambda$.

Theorem 1.4. *For any points $y_1, y_2, \dots, y_p \in (0, 1/2)$ we have*

$$\dim_H \bigcap_{i=1}^p \Lambda(y_i) = 1.$$

Recently, the last three authors studied in [6] analogous objects but with different family of self-similar sets (their self-similar sets have different convex hulls). We indicate that our method is different from theirs. Furthermore, Theorem 1.4 shows that the intersection of any finitely many $\Lambda(y_i)$ has full Hausdorff dimension, while in [6, Theorem 1.5] their method can only prove that the intersection of two associated sets has full Hausdorff dimension.

The rest of the paper is organized as follows. In Section 2 we prove Theorem 1.1 for the topology of $\Lambda(x)$; and in Section 3 we investigate the local dimension of $\Lambda(x)$ and prove Theorem 1.2. In Section 4 we consider the intersection $\bigcap_{i=1}^p \Lambda(y_i)$ and prove Theorem 1.4; and in the final section we make some further remarks.

2. TOPOLOGICAL PROPERTIES OF $\Lambda(x)$

In this section we will investigate the topology of $\Lambda(x)$, and prove Theorem 1.1. First we recall some terminology from symbolic dynamics (cf. [7]). Let $\{0, 1\}^{\mathbb{N}}$ be the set of all infinite

sequences of zeros and ones. For a *word* we mean a finite string of zeros and ones. Let $\{0, 1\}^*$ be the set of all words over the alphabet $\{0, 1\}$ together with the empty word ϵ . For two words $\mathbf{c} = c_1 \dots c_m, \mathbf{d} = d_1 \dots d_n$ from $\{0, 1\}^*$ we write $\mathbf{cd} = c_1 \dots c_m d_1 \dots d_n$ for their concatenation. In particular, for $n \in \mathbb{N}$ we denote by \mathbf{c}^n the n -fold concatenation of \mathbf{c} with itself, and by \mathbf{c}^∞ the periodic sequence with period block \mathbf{c} . Throughout the paper we will use lexicographical order ' \prec, \preceq, \succ ' or ' \succcurlyeq ' between sequences and words. For example, for two sequences $(c_i), (d_i) \in \{0, 1\}^\mathbb{N}$, we say $(c_i) \prec (d_i)$ if $c_1 < d_1$, or there exists $n \in \mathbb{N}$ such that $c_1 \dots c_n = d_1 \dots d_n$ and $c_{n+1} < d_{n+1}$. For two words \mathbf{c}, \mathbf{d} , we say $\mathbf{c} \prec \mathbf{d}$ if $\mathbf{c}0^\infty \prec \mathbf{d}0^\infty$.

Let $\lambda \in (0, 1/2]$. We define the coding map $\pi_\lambda : \{0, 1\}^\mathbb{N} \rightarrow K_\lambda$ by

$$(2.1) \quad \pi_\lambda((i_n)) = \lim_{n \rightarrow \infty} f_{\lambda, i_1} \circ f_{\lambda, i_2} \circ \dots \circ f_{\lambda, i_n}(0) = (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1}.$$

If $\lambda \in (0, 1/2)$, then the IFS $\{f_{\lambda, d}(x) = \lambda x + d(1 - \lambda) : d = 0, 1\}$ satisfies the strong separation condition, and thus the map π_λ is bijective. If $\lambda = 1/2$, then $\pi_{1/2}$ is bijective up to a countable set. The map π_λ defined in (2.1) naturally induces a function with two parameters:

$$(2.2) \quad \Pi : \{0, 1\}^\mathbb{N} \times (0, 1/2] \rightarrow [0, 1]; \quad ((i_n), \lambda) \mapsto \pi_\lambda((i_n)).$$

Equipped with $\{0, 1\}^\mathbb{N}$ the order topology given by the metric

$$(2.3) \quad \rho((i_n), (j_n)) = 2^{-\inf\{n \geq 1 : i_n \neq j_n\}},$$

and with $\{0, 1\}^\mathbb{N} \times (0, 1/2]$ the product topology we show that the function Π is continuous.

Lemma 2.1. *The function Π is continuous. Furthermore,*

- (i) *for $\lambda \in (0, 1/2]$ the function $\Pi(\cdot, \lambda)$ is increasing with respect to the lexicographical order, and is strictly increasing if $\lambda \in (0, 1/2)$;*
- (ii) *if $0^\infty \prec (i_n) \preceq 01^\infty$, then $\Pi((i_n), \cdot)$ has positive derivative in $(0, 1/2)$.*

Proof. First we prove the continuity of Π . For any two points $((i_n), \lambda_1), ((j_n), \lambda_2) \in \{0, 1\}^\mathbb{N} \times (0, 1/2]$ we have

$$(2.4) \quad |\Pi((j_n), \lambda_2) - \Pi((i_n), \lambda_1)| \leq |\Pi((j_n), \lambda_2) - \Pi((i_n), \lambda_2)| + |\Pi((i_n), \lambda_2) - \Pi((i_n), \lambda_1)|.$$

Note that if $\rho((j_n), (i_n)) \leq 2^{-m}$, then $|\Pi((j_n), \lambda_2) - \Pi((i_n), \lambda_2)| \leq \lambda_2^{m-1} \leq 2^{1-m}$. So the first term in (2.4) converges to zero as $\rho((j_n), (i_n)) \rightarrow 0$. Moreover, since the series $\Pi((i_n), \lambda) = (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1}$ with parameter λ converges uniformly in $(0, 1/2]$, the second term in (2.4) also converges to zero as $|\lambda_2 - \lambda_1| \rightarrow 0$. Therefore, Π is continuous.

For (i) let $\lambda \in (0, 1/2]$ and take two sequences $(i_n), (j_n) \in \{0, 1\}^\mathbb{N}$. Suppose $(i_n) \prec (j_n)$. Then there exists a $m \in \mathbb{N}$ such that $i_1 \dots i_{m-1} = j_1 \dots j_{m-1}$ and $i_m < j_m$. This implies that

$$\begin{aligned} \Pi((i_n), \lambda) &= (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1} \leq (1 - \lambda) \left(\sum_{n=1}^m i_n \lambda^{n-1} + \sum_{n=m+1}^{\infty} \lambda^{n-1} \right) \\ &\leq (1 - \lambda) \sum_{n=1}^m j_n \lambda^{n-1} \\ &\leq \Pi((j_n), \lambda), \end{aligned}$$

where the second inequality follows by $\lambda \in (0, 1/2]$, and this inequality is strict if $\lambda \in (0, 1/2)$.

For (ii) let $(i_n) \in \{0, 1\}^{\mathbb{N}}$ with $0^\infty \prec (i_n) \preceq 01^\infty$. Then $i_1 = 0$. So for any $\lambda \in (0, 1/2)$ we have $\Pi((i_n), \lambda) = (1 - \lambda) \sum_{n=2}^{\infty} i_n \lambda^{n-1}$. This implies that

$$(2.5) \quad \frac{d\Pi((i_n), \lambda)}{d\lambda} = \sum_{n=2}^{\infty} n \left(\frac{n-1}{n} - \lambda \right) i_n \lambda^{n-2} > 0,$$

where the inequality follows since $\lambda < 1/2$ and $(i_n) \succ 0^\infty$. This completes the proof. \square

Note that the map Π defined in (2.2) is surjective but not injective. Given $x \in [0, 1]$, for $\lambda \in (0, 1/2]$ we consider the horizontal fiber

$$\Gamma_x(\lambda) := \Pi^{-1}(x) \cap (\{0, 1\}^{\mathbb{N}} \times \{\lambda\}) = \left\{ ((i_n), \lambda) \in \{0, 1\}^{\mathbb{N}} \times (0, 1/2] : (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1} = x \right\}.$$

Then $\Gamma_x(\lambda) \neq \emptyset$ if and only if $\lambda \in \Lambda(x)$, where $\Lambda(x)$ is defined in (1.2). Furthermore, by Lemma 2.1 (i) it follows that for any $\lambda \in \Lambda(x) \cap (0, 1/2)$ the fiber set $\Gamma_x(\lambda)$ contains only one sequence; and for $\lambda = 1/2 \in \Lambda(x)$ the set $\Gamma_x(1/2)$ contains at most two sequences. This defines a map

$$\Psi_x : \Lambda(x) \rightarrow \{0, 1\}^{\mathbb{N}}; \quad \lambda \mapsto \Psi_x(\lambda),$$

where $\Psi_x(\lambda)$ denotes the lexicographically largest sequence in $\Gamma_x(\lambda)$. The sequence $\Psi_x(\lambda)$ is also called the *greedy coding* of x in base λ .

Given $x \in (0, 1/2)$, we reserve the notation $\Psi_x(1/2) = (x_n)$ for the greedy coding of x in base $1/2$. Then (x_n) begins with 0 and does not end with 1^∞ .

Lemma 2.2. *For any $x \in (0, 1/2)$ the map $\Psi_x : \Lambda(x) \rightarrow \Omega(x)$ is a decreasing homeomorphism, where*

$$\Omega(x) := \left\{ (i_n) \in \{0, 1\}^{\mathbb{N}} : (x_n) \preceq (i_n) \preceq 01^\infty \right\}.$$

Proof. Let $x \in (0, 1/2)$. By Lemma 2.1 it follows that Ψ_x is strictly decreasing. Observe that $x \notin K_\lambda$ for any $\lambda < x$. Then $\Lambda(x) \subset [x, 1/2]$. Note that $\Psi_x(x) = 01^\infty$ and $\Psi_x(1/2) = (x_n)$. Since Ψ_x is monotone decreasing, we have

$$(x_n) \preceq \Psi_x(\lambda) \preceq 01^\infty \quad \forall \lambda \in [x, 1/2].$$

So, $\Psi_x(\Lambda(x)) \subset \Omega(x)$.

Next we show that $\Psi_x(\Lambda(x)) = \Omega(x)$. Let $(i_n) \in \Omega(x)$. Then by Lemma 2.1 it follows that

$$\Pi\left((i_n), \frac{1}{2}\right) \geq \Pi\left((x_n), \frac{1}{2}\right) = x \quad \text{and} \quad \Pi((i_n), \lambda) \searrow 0 < x \quad \text{as} \quad \lambda \searrow 0.$$

So, by the continuity of Π in Lemma 2.1 there must exist a $\lambda \in (0, 1/2]$ such that

$$(2.6) \quad \Pi((i_n), \lambda) = x.$$

If $\lambda \in (0, 1/2)$, then (2.6) gives that $\Psi_x(\lambda) = (i_n)$. If $\lambda = 1/2$, then by (2.6) and using $(i_n) \succ (x_n)$ we still have $\Psi_x(\lambda) = (i_n)$. This proves $\Psi_x(\Lambda(x)) = \Omega(x)$. Hence, $\Psi_x : \Lambda(x) \rightarrow \Omega(x)$ is a decreasing bijection.

To complete the proof it remains to prove the continuity of Ψ_x and its inverse Ψ_x^{-1} . Since the proof for the continuity of Ψ_x^{-1} is similar to that of Ψ_x , we only prove it for Ψ_x . Take $\lambda_* \in \Lambda(x)$. Suppose Ψ_x is not continuous at λ_* . Then there exists a $N \in \mathbb{N}$ such that for any $\delta > 0$ we can find $\lambda \in \Lambda(x) \cap (\lambda_* - \delta, \lambda_* + \delta)$ such that $|\Psi_x(\lambda) - \Psi_x(\lambda_*)| \geq 2^{-N}$. Letting $\delta = 1/k$ with $k = 1, 2, \dots$, we can find a sequence $(\lambda_k) \subset \Lambda(x)$ such that

$$(2.7) \quad \lim_{k \rightarrow \infty} \lambda_k = \lambda_* \quad \text{and} \quad |\Psi_x(\lambda_k) - \Psi_x(\lambda_*)| \geq 2^{-N} \quad \forall k \geq 1.$$

Write $\Psi_x(\lambda_k) = (i_n^{(k)})$ and $\Psi_x(\lambda_*) = (i_n^*)$. Then by (2.7) we have $i_1^{(k)} \dots i_N^{(k)} \neq i_1^* \dots i_N^*$ for all $k \geq 1$. Note that $(\{0, 1\}^{\mathbb{N}}, \rho)$ is a compact metric space, where ρ is defined in (2.3). So we can find a subsequence $(k_j) \subset \mathbb{N}$ such that the limit $\lim_{j \rightarrow \infty} (i_n^{(k_j)})$ exists, say (i'_n) . Then $i'_1 \dots i'_N \neq i_1^* \dots i_N^*$. Observe that

$$(2.8) \quad \Pi((i_n^{(k_j)}), \lambda_{k_j}) = x = \Pi((i_n^*), \lambda_*) \quad \forall j \geq 1.$$

Letting $j \rightarrow \infty$ in (2.8), by (2.7) and Lemma 2.1 it follows that

$$(2.9) \quad \Pi((i'_n), \lambda_*) = x = \Pi((i_n^*), \lambda_*).$$

If $\lambda_* = 1/2$, then by Lemma 2.1 (i) it follows that $\Psi_x(\lambda_{k_j}) \succ \Psi_x(\lambda_*) = (i_n^*)$ for all $j \geq 1$, and thus $(i'_n) \succ (i_n^*)$. Note that (i_n^*) is the greedy coding of x in base λ_* . Then by (2.9) it follows that $(i'_n) = (i_n^*)$, leading to a contradiction with $i'_1 \dots i'_N \neq i_1^* \dots i_N^*$. If $\lambda_* < 1/2$, then (2.9) gives that $(i'_n) = (i_n^*)$. This again leads to a contradiction. Therefore, Ψ_x is continuous at λ_* . Since $\lambda_* \in \Lambda(x)$ is arbitrary, Ψ_x is continuous in $\Lambda(x)$. This completes the proof. \square

Proof of Theorem 1.1. Let $x \in (0, 1/2)$. By Lemma 2.2 it follows that $\min \Lambda(x) = \Psi_x^{-1}(01^\infty) = x$ and $\max \Lambda(x) = \Psi_x^{-1}((x_n)) = 1/2$. Observe that $(\Omega(x), \rho)$ is a Cantor set. Then by Lemma 2.2 we conclude that $\Lambda(x)$ is also a Cantor set. \square

3. LEBESGUE MEASURE AND HAUSDORFF DIMENSION OF $\Lambda(x)$

In this section we will prove Theorem 1.2, which states that for any $x \in (0, 1/2)$ the set $\Lambda(x)$ is a Lebesgue null set of full Hausdorff dimension. First we consider the local dimension of $\Lambda(x)$.

Proposition 3.1. *Let $x \in (0, 1/2)$. Then for any $\lambda \in \Lambda(x)$ we have*

$$(3.1) \quad \lim_{\delta \rightarrow 0^+} \dim_H(\Lambda(x) \cap (\lambda - \delta, \lambda + \delta)) = \dim_H K_\lambda = -\frac{\log 2}{\log \lambda}.$$

The second equality in (3.1) is obvious, since for any $\lambda \in \Lambda(x)$ the self-similar set K_λ is generated by the IFS $\{\lambda x, \lambda x + (1 - \lambda)\}$ satisfying the open set condition (cf. [5]). So it suffices to prove the first equality in (3.1).

Lemma 3.2. *Let $x \in (0, 1/2)$. Then for any $\lambda \in (x, 1/2)$ we have*

$$\dim_H(\Lambda(x) \cap [x, \lambda]) \leq \dim_H K_\lambda.$$

Proof. Let $\lambda \in (x, 1/2)$. Note by Lemma 2.2 that $\pi_\lambda \circ \Psi_x : \Lambda(x) \cap [x, \lambda] \rightarrow K_\lambda$ is injective. Since Hausdorff dimension is preserved by Lipschitz mappings, it suffices to prove that for any $\lambda_1, \lambda_2 \in \Lambda(x) \cap [x, \lambda]$ we have

$$(3.2) \quad |\pi_\lambda(\Psi_x(\lambda_1)) - \pi_\lambda(\Psi_x(\lambda_2))| \geq C|\lambda_1 - \lambda_2|,$$

where $C > 0$ is a constant independent of λ_1 and λ_2 .

Take $\lambda_1, \lambda_2 \in \Lambda(x) \cap [x, \lambda]$ with $\lambda_1 < \lambda_2$, and write $\Psi_x(\lambda_1) = (i_n)$, $\Psi_x(\lambda_2) = (j_n)$. By Lemma 2.2 we have $i_1 = j_1 = 0$ and $(i_n) \succ (j_n)$. Then there exists $m \geq 2$ such that $i_1 \dots i_{m-1} = j_1 \dots j_{m-1}$ and $i_m > j_m$. Note that

$$(1 - \lambda_1) \sum_{n=2}^{\infty} i_n \lambda_1^{n-1} = x = (1 - \lambda_2) \sum_{n=2}^{\infty} j_n \lambda_2^{n-1}.$$

Then

$$\begin{aligned} \frac{x(1 - \lambda_1 - \lambda_2)}{\lambda_1 \lambda_2 (1 - \lambda_1)(1 - \lambda_2)} (\lambda_2 - \lambda_1) &= \frac{x}{\lambda_1(1 - \lambda_1)} - \frac{x}{\lambda_2(1 - \lambda_2)} \\ &= \sum_{n=2}^{\infty} i_n \lambda_1^{n-2} - \sum_{n=2}^{\infty} j_n \lambda_2^{n-2} \\ &\leq \sum_{n=2}^{m-1} i_n \lambda_1^{n-2} + \sum_{n=m}^{\infty} \lambda_1^{n-2} - \sum_{n=2}^{m-1} i_n \lambda_2^{n-2} \\ &\leq \sum_{n=m}^{\infty} \lambda_1^{n-2} = \frac{\lambda_1^{m-2}}{1 - \lambda_1}, \end{aligned}$$

where the first inequality follows by $i_1 \dots i_{m-1} = j_1 \dots j_{m-1}$, and the second inequality follows by $\lambda_1 < \lambda_2$. This, together with $\lambda_1 < \lambda_2 \leq \lambda$, implies that

$$\lambda^m \geq \lambda_1^{m-1} \lambda_2 \geq \frac{x(1 - \lambda_1 - \lambda_2)}{1 - \lambda_2} (\lambda_2 - \lambda_1) \geq x(1 - 2\lambda)(\lambda_2 - \lambda_1).$$

Therefore,

$$\begin{aligned} |\pi_\lambda(\Psi_x(\lambda_1)) - \pi_\lambda(\Psi_x(\lambda_2))| &= (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1} - (1 - \lambda) \sum_{n=1}^{\infty} j_n \lambda^{n-1} \\ &\geq (1 - \lambda) \left(\lambda^{m-1} - \sum_{n=m+1}^{\infty} \lambda^{n-1} \right) \\ &= (1 - 2\lambda) \lambda^{m-1} \geq C|\lambda_2 - \lambda_1|, \end{aligned}$$

where $C = \frac{x(1-2\lambda)^2}{\lambda} > 0$ (since $\lambda < 1/2$). This proves (3.2), and then completes the proof. \square

Lemma 3.3. *Let $x \in (0, 1/2)$. If $\lambda \in \Lambda(x) \setminus \{1/2\}$ such that $\Psi_x(\lambda)$ does not end with 0^∞ , then for any $\delta \in (0, 1/2 - \lambda)$,*

$$\dim_H(\Lambda(x) \cap [\lambda, \lambda + \delta]) \geq \dim_H K_\lambda.$$

Proof. Let $\lambda \in \Lambda(x) \setminus \{1/2\}$ such that $\Psi_x(\lambda) = (c_n)$ does not end with 0^∞ . Take $\delta \in (0, 1/2 - \lambda)$. We will construct a sequence of subsets in $\Lambda(x) \cap [\lambda, \lambda + \delta]$ whose Hausdorff dimension can be arbitrarily close to $\dim_H K_\lambda$.

Since $\lambda < 1/2$, by Lemma 2.2 we have $(c_n) \succ \Psi_x(1/2) = (x_n)$. Then there exists $n_0 \geq 2$ such that $c_1 \dots c_{n_0-1} = x_1 \dots x_{n_0-1}$ and $c_{n_0} > x_{n_0}$. Since (c_n) does not end with 0^∞ , we can find an increasing sequence $(n_k) \subset \mathbb{N}$ such that $n_0 < n_1 < n_2 < \dots$, and $c_{n_k} = 1$ for all $k \geq 1$. Now for $k \geq 1$, we define

$$(3.3) \quad \Omega_{\lambda,k} := \left\{ c_1 \dots c_{n_k-1} 0 i_1 i_2 \dots : i_{n+1} \dots i_{n+k} \neq 0^k \ \forall n \geq 0 \right\}.$$

Note by Lemma 2.2 that $\Psi_x(\Lambda(x) \cap [\lambda, 1/2]) = \{(i_n) : (x_n) \preceq (i_n) \preceq (c_n)\}$. Then by using $c_{n_0} > x_{n_0}$ and $c_{n_k} = 1$ it follows that

$$(3.4) \quad \Omega_{\lambda,k} \subset \Psi_x(\Lambda(x) \cap [\lambda, 1/2]) \quad \text{for all } k \geq 1.$$

Since $\delta \in (0, 1/2 - \lambda)$, by (3.3), (3.4) and Lemma 2.2 there exists a $N \in \mathbb{N}$ such that

$$\Lambda(x) \cap [\lambda, \lambda + \delta] \supset \Psi_x^{-1}(\Omega_{\lambda,k}) \quad \forall k \geq N.$$

So, to finish the proof it suffices to prove that

$$(3.5) \quad \lim_{k \rightarrow \infty} \dim_H \Psi_x^{-1}(\Omega_{\lambda,k}) \geq \dim_H K_\lambda.$$

Take $k \geq N$, and consider the map $\pi_\lambda \circ \Psi_x : \Psi_x^{-1}(\Omega_{\lambda,k}) \rightarrow \pi_\lambda(\Omega_{\lambda,k})$. Let $\lambda_1, \lambda_2 \in \Psi_x^{-1}(\Omega_{\lambda,k})$ with $\lambda_1 < \lambda_2$, and write $\Psi_x(\lambda_1) = (i_n), \Psi_x(\lambda_2) = (j_n)$. Then $(i_n), (j_n) \in \Omega_{\lambda,k}$. Since $\lambda_1 < \lambda_2$, by Lemma 2.2 we have $(i_n) \succ (j_n)$. So there exists $m > n_k$ such that $i_1 \dots i_{m-1} = j_1 \dots j_{m-1}$ and $i_m > j_m$. Note that $i_m i_{m+1} \dots$ does not contain k consecutive zeros. Then

$$(3.6) \quad x = (1 - \lambda_1) \sum_{n=1}^{\infty} i_n \lambda_1^{n-1} > (1 - \lambda_2) \sum_{n=1}^{\infty} i_n \lambda_1^{n-1} > (1 - \lambda_2) \left(\sum_{n=1}^m i_n \lambda_1^{n-1} + \lambda_1^{m+k-1} \right).$$

On the other hand,

$$(3.7) \quad x = (1 - \lambda_2) \sum_{n=1}^{\infty} j_n \lambda_2^{n-1} \leq (1 - \lambda_2) \sum_{n=1}^m i_n \lambda_2^{n-1}.$$

Note that $\lambda_i \in \Psi_x^{-1}(\Omega_{\lambda,k}) \subset \Lambda(x) \cap [\lambda, \lambda + \delta]$. Then by (3.6) and (3.7) it follows that

$$\begin{aligned} \lambda^{m+k-1} &\leq \lambda_1^{m+k-1} < \sum_{n=1}^m i_n (\lambda_2^{n-1} - \lambda_1^{n-1}) \\ &< \sum_{n=1}^{\infty} (\lambda_2^{n-1} - \lambda_1^{n-1}) = \frac{1}{1 - \lambda_2} - \frac{1}{1 - \lambda_1} = \frac{\lambda_2 - \lambda_1}{(1 - \lambda_1)(1 - \lambda_2)}. \end{aligned}$$

This implies that

$$\begin{aligned}
|\pi_\lambda(\Psi_x(\lambda_1)) - \pi_\lambda(\Psi_x(\lambda_2))| &= (1 - \lambda) \sum_{n=1}^{\infty} i_n \lambda^{n-1} - (1 - \lambda) \sum_{n=1}^{\infty} j_n \lambda^{n-1} \\
(3.8) \quad &\leq (1 - \lambda) \sum_{n=m}^{\infty} \lambda^{n-1} \\
&= \lambda^{m-1} < \frac{\lambda_2 - \lambda_1}{\lambda^k (1 - \lambda_1)(1 - \lambda_2)} < \frac{4}{\lambda^k} (\lambda_2 - \lambda_1),
\end{aligned}$$

where the last inequality follows by $\lambda_i < 1/2$.

So, by (3.3) and (3.8) it follows that

$$\begin{aligned}
\dim_H \Psi_x^{-1}(\Omega_{\lambda,k}) &\geq \dim_H \pi_\lambda(\Omega_{\lambda,k}) \\
&= \dim_H \pi_\lambda(\{(i_n) : i_{n+1} \dots i_{n+k} \neq 0^k \ \forall n \geq 0\}) \\
&\geq \dim_H \pi_\lambda(\{(i_n) \in \{0, 1\}^{\mathbb{N}} : i_n = 1 \text{ for all } n \equiv 0 \pmod{k}\}) \\
&= -\frac{(k-1) \log 2}{k \log \lambda} \rightarrow \frac{\log 2}{-\log \lambda} = \dim_H K_\lambda,
\end{aligned}$$

as $k \rightarrow \infty$. This proves (3.5), and then completes the proof. \square

Proof of Proposition 3.1. Take $\lambda \in \Lambda(x)$. Note that $\Lambda(x) \subset [x, 1/2]$ and $x, 1/2 \in \Lambda(x)$. We will prove (3.1) in the following two cases.

Case I. $\lambda \in \Lambda(x) \cap [x, 1/2)$. Then by Lemma 3.2 it follows that for any $\delta \in (0, 1/2 - \lambda)$,

$$\dim_H(\Lambda(x) \cap (\lambda - \delta, \lambda + \delta)) \leq \dim_H(\Lambda(x) \cap [x, \lambda + \delta]) \leq \dim_H K_{\lambda+\delta} = \frac{\log 2}{-\log(\lambda + \delta)}.$$

This implies that

$$(3.9) \quad \lim_{\delta \rightarrow 0^+} \dim_H(\Lambda(x) \cap (\lambda - \delta, \lambda + \delta)) \leq \frac{\log 2}{-\log \lambda} = \dim_H K_\lambda.$$

On the other hand, take $\delta \in (0, 1/2 - \lambda)$. Note by Theorem 1.1 that $\Lambda(x)$ is a Cantor set, and $\lambda \in \Lambda(x)$. Then we can find a sequence (λ_k) in $\Lambda(x) \cap (\lambda - \delta, \lambda + \delta)$ such that each $\Psi_x(\lambda_k)$ does not end with 0^∞ , and $\lambda_k \rightarrow \lambda$ as $k \rightarrow \infty$. Therefore, by Lemma 3.3 it follows that

$$\begin{aligned}
\dim_H(\Lambda(x) \cap (\lambda - \delta, \lambda + \delta)) &\geq \dim_H(\Lambda(x) \cap [\lambda_k, \lambda + \delta]) \\
&\geq \dim_H K_{\lambda_k} = \frac{\log 2}{-\log \lambda_k} \rightarrow \frac{\log 2}{-\log \lambda} = \dim_H K_\lambda,
\end{aligned}$$

as $k \rightarrow \infty$. This, together with (3.9), proves (3.1).

Case II. $\lambda = 1/2$. The proof is similar to that for the second part of Case I. Let $\delta > 0$. Since $\Lambda(x)$ is a Cantor set and $\max \Lambda(x) = 1/2$, there exists a sequence (λ_k) in $\Lambda(x) \cap (1/2 - \delta, 1/2)$ such that each $\Psi_x(\lambda_k)$ does not end with 0^∞ , and $\lambda_k \nearrow 1/2$ as $k \rightarrow \infty$. Then by Lemma 3.3

it follows that

$$\begin{aligned} \dim_H(\Lambda(x) \cap (1/2 - \delta, 1/2 + \delta)) &\geq \dim_H(\Lambda(x) \cap [\lambda_k, \lambda_{k+1}]) \\ &\geq \dim_H K_{\lambda_k} = \frac{\log 2}{-\log \lambda_k} \rightarrow 1 = \dim_H K_{1/2}, \end{aligned}$$

proving (3.1). □

As a direct consequence of Proposition 3.1 we have the following dimensional result of $\Lambda(x)$.

Corollary 3.4. *Let $x \in (0, 1/2)$. Then for any open interval $I \subset \mathbb{R}$ with $\Lambda(x) \cap I \neq \emptyset$ we have*

$$\dim_H(\Lambda(x) \cap I) = \sup_{\lambda \in \Lambda(x) \cap I} \dim_H K_\lambda.$$

Proof of Theorem 1.2. By Corollary 3.4 it follows that

$$\dim_H \Lambda(x) = \dim_H(\Lambda(x) \cap (x, 1/2)) = \sup_{\lambda \in (x, 1/2)} \dim_H K_\lambda = 1.$$

Furthermore, for any $n \in \mathbb{N}$ the Hausdorff dimension of $\Lambda_n(x) := \Lambda(x) \cap [x, 1/2 - 1/n]$ is strictly smaller than one, and thus each $\Lambda_n(x)$ has zero Lebesgue measure. Since $\Lambda(x) \setminus \{1/2\} = \bigcup_{n=1}^{\infty} \Lambda_n(x)$, the set $\Lambda(x)$ also has zero Lebesgue measure. This together with Proposition 3.1 completes the proof. □

4. HAUSDORFF DIMENSION OF THE INTERSECTION $\bigcap_{i=1}^p \Lambda(y_i)$

Given finitely many numbers $y_1, y_2, \dots, y_p \in (0, 1/2)$, we will show in this section that the intersection $\bigcap_{i=1}^p \Lambda(y_i)$ has full Hausdorff dimension (Theorem 1.4). Note by Theorem 1.1 that each set $\Lambda(y_i)$ is a Cantor set. We will construct in each $\Lambda(y_i)$ a sequence of Cantor subsets $C_\ell(y_i), \ell \geq 1$ such that each $C_\ell(y_i)$ has the same maximum point $1/2$, and the thickness of $C_\ell(y_i)$ tends to infinity as $\ell \rightarrow \infty$. Then by using a result from Hunt, Kan and Yorke [4] (see Lemma 4.2 below) we conclude that the intersection $\bigcap_{i=1}^p \Lambda(y_i)$ contains a sequence of Cantor subsets whose thickness tends to infinity. This, together with Lemma 4.1 (see below), implies that $\bigcap_{i=1}^p \Lambda(y_i)$ has full Hausdorff dimension.

4.1. Thickness of a Cantor set in \mathbb{R} . First we recall the *thickness* of a Cantor set in \mathbb{R} introduced by Newhouse [8] (see [1] for some recent progress). Let E be a Cantor set in \mathbb{R} with its convex hull $\text{conv}(E) = E_0$. Then the complement $E_0 \setminus E = \bigcup_{n=1}^{\infty} V_n$ is the union of countably many disjoint open intervals. The sequence $\mathcal{V} = (V_1, V_2, \dots)$ is called a *defining sequence* for E . If moreover $|V_1| \geq |V_2| \geq |V_3| \geq \dots$, where $|V|$ denotes the diameter of a set $V \subset \mathbb{R}$, then we call \mathcal{V} an *ordered defining sequence* for E . Let $E_n := E_0 \setminus \bigcup_{k=1}^n V_k$. Then E_n is the union of finitely many closed intervals. So, for any $n \geq 1$, the open interval V_n is contained in some connected component of E_{n-1} , say E_{n-1}^* . Then the set $E_{n-1}^* \setminus V_n$ is the union of two closed intervals $L_{\mathcal{V}}(V_n)$ and $R_{\mathcal{V}}(V_n)$, where we always assume that $L_{\mathcal{V}}(V_n)$ lies to the left of $R_{\mathcal{V}}(V_n)$. We emphasize that both intervals $L_{\mathcal{V}}(V_n)$ and $R_{\mathcal{V}}(V_n)$ have positive

length, since otherwise E will contain isolated points which is impossible. Then the thickness of E with respect to the defining sequence \mathcal{V} is defined by

$$(4.1) \quad \tau_{\mathcal{V}}(E) := \inf_{n \geq 1} \min \left\{ \frac{|L_{\mathcal{V}}(V_n)|}{|V_n|}, \frac{|R_{\mathcal{V}}(V_n)|}{|V_n|} \right\}.$$

And the *thickness* of E is defined by

$$(4.2) \quad \tau(E) := \sup_{\mathcal{V}} \tau_{\mathcal{V}}(E),$$

where the supremum is taken over all defining sequences \mathcal{V} for E . It was shown in [1] that $\tau(E) = \tau_{\mathcal{V}}(E)$ for every ordered defining sequence \mathcal{V} for E .

The following result on a lower bound for the Hausdorff dimension of a Cantor set in \mathbb{R} by using thickness was proven by Newhouse [9] (see also, [10]).

Lemma 4.1. [9] *If E is a Cantor set in \mathbb{R} , then*

$$\dim_H E \geq \frac{\log 2}{\log \left(2 + \frac{1}{\tau(E)} \right)}.$$

Two Cantor sets in \mathbb{R} are called *interleaved* if neither set lies in a gap of the other. The following result for the intersection of two interleaved Cantor sets was shown by Hunt, Kan and Yorke [4].

Lemma 4.2. [4, Theorem 1] *There exists a function $\varphi : (0, \infty) \rightarrow (0, \infty)$ such that for all interleaved Cantor sets E and F in \mathbb{R} with $\tau(E) \geq t$ and $\tau(F) \geq t$, there exists a Cantor subset $K \subset E \cap F$ with $\tau(K) \geq \varphi(t)$.*

Remark 4.3. (i) In [4, P. 882] the authors pointed out that when t is sufficiently large, $\varphi(t)$ is of order \sqrt{t} . So, Lemma 4.2 implies that if the thicknesses of two interleaved Cantor sets E and F in \mathbb{R} are sufficiently large, then the thickness of the resulting Cantor set $K \subset E \cap F$ is also very large.

(ii) It is clear that if two Cantor sets E and F in \mathbb{R} have the same maximum point ξ , then they are interleaved. Furthermore, if the maximum point ξ is also an accumulation point of $E \cap F$, then from the proof of [4, Theorem 1] (see also [4, Page 887]) it follows that the resulting Cantor set $K \subset E \cap F$ in Lemma 4.2 can be required to have the maximum point ξ .

4.2. Construction of Cantor subsets of $\Lambda(x)$. Let $x \in (0, 1/2)$. We will construct a sequence of Cantor subsets $\{C_\ell(x)\}_{\ell=1}^\infty$ of $\Lambda(x)$ such that each $C_\ell(x)$ has the maximum point $1/2$, and $\tau(C_\ell(x)) \rightarrow +\infty$ as $\ell \rightarrow \infty$. Recall that $\Psi_x(1/2) = (x_n)$ satisfies that $x_1 = 0$ and (x_n) does not end with 1^∞ . Denote by (n_k) the set of all indices such that $x_{n_k} = 0$. Then $x_{n_k} = 0$ for any $k \geq 1$; and $x_n = 1$ for any $n_k < n < n_{k+1}$. Recall from Lemma 2.2 that $\Omega(x) = \Psi_x(\Lambda(x)) = \{(i_n) : (x_i) \preceq (i_n) \preceq 01^\infty\}$. For $k \geq 1$, we define

$$(4.3) \quad \Omega_k(x) := \{(i_n) : x_1 \cdots x_{n_k-1} 10^\infty \preceq (i_n) \preceq x_1 \cdots x_{n_k-1} 1^\infty\}.$$

Then $\Omega_k(x) \subset \Omega(x)$, which implies that

$$(4.4) \quad F_k(x) := \Psi_x^{-1}(\Omega_k(x)) \subset \Lambda(x).$$

Note by (4.3) that for each $k \geq 1$ the set $(\Omega_k(x), \rho)$ is a topological Cantor set, where ρ is the metric defined in (2.3). By Lemma 2.2 it follows that each $F_k(x)$ is a Cantor subset of $\Lambda(x)$. Now for any $\ell \geq 1$, we define

$$(4.5) \quad C_\ell(x) := \{1/2\} \cup \bigcup_{k=\ell}^{\infty} F_k(x).$$

Then $C_\ell(x) \subset \Lambda(x)$ for all $\ell \geq 1$. Observe by (4.3) that the largest sequence in $\Omega_{k+1}(x)$ is $x_1 \dots x_{n_{k+1}-1} 1^\infty = x_1 \dots x_{n_k-1} 0 1^\infty$, which is strictly smaller than the smallest sequence $x_1 \dots x_{n_k-1} 1 0^\infty$ in $\Omega_k(x)$. So, by (4.4) and Lemma 2.2 it follows that $\max F_k(x) < \min F_{k+1}(x)$ for all $k \geq 1$. Furthermore, since the maximum sequence in $\Omega_k(x)$ decreases to $(x_n) = \Psi_x(1/2)$ as $k \rightarrow \infty$, again by Lemma 2.2 we conclude that $\min F_k(x) \nearrow 1/2$ as $k \rightarrow \infty$. Therefore, each $C_\ell(x)$ defined in (4.5) is a Cantor subset of $\Lambda(x)$, and $\min C_\ell(x) = \min F_\ell(x) \nearrow 1/2$ as $\ell \rightarrow \infty$. Furthermore, all of these Cantor subsets $C_\ell(x)$, $\ell \geq 1$ have the same maximum point $1/2$.

Let $\ell \geq 1$, and take $k \geq \ell$. First we describe a defining sequence for the Cantor set $F_k(x)$. Let $[\alpha_k, \beta_k]$ be the convex hull of $F_k(x)$. Then

$$\alpha_k = \Psi_x^{-1}(x_1 x_2 \dots x_{n_k-1} 1^\infty), \quad \beta_k = \Psi_x^{-1}(x_1 x_2 \dots x_{n_k-1} 1 0^\infty).$$

Clearly, α_k and β_k depend on x . For simplicity we will suppress this dependence in our notation if no confusion arises. By Lemma 2.2 it follows that

$$[\alpha_k, \beta_k] \setminus F_k(x) = \bigcup_{\omega \in \{0,1\}^*} V_{k,\omega},$$

where

$$V_{k,\omega} := (\Psi_x^{-1}(x_1 x_2 \dots x_{n_k-1} 1 \omega 10^\infty), \Psi_x^{-1}(x_1 x_2 \dots x_{n_k-1} 1 \omega 01^\infty)).$$

We enumerate these open intervals $V_{k,\omega}$, $\omega \in \{0,1\}^*$ according first to the length of ω and then to the lexicographical order of ω :

$$(4.6) \quad \begin{aligned} &V_{k,\epsilon}; \quad V_{k,0}, V_{k,1}; \quad V_{k,00}, V_{k,01}, V_{k,10}, V_{k,11}; \\ &V_{k,000}, V_{k,001}, V_{k,010}, V_{k,011}, V_{k,100}, V_{k,101}, V_{k,110}, V_{k,111}; \quad \dots, \end{aligned}$$

where ϵ is the empty word. Then for $\omega = i_1 i_2 \dots i_q \in \{0,1\}^*$ the open interval $V_{k,\omega}$ is indeed at the $N(\omega)$ -th position in (4.6), where

$$N(\omega) = 2^q + \sum_{n=1}^q i_n 2^{q-n}.$$

Note that for $\omega = \epsilon$ we have $N(\epsilon) = 1$. So, we can also write $V_{k,\omega} = V_{k,N(\omega)}$. Hence, we obtain a defining sequence $\mathcal{V}_k = \{V_{k,j}\}_{j=1}^{\infty}$ for $F_k(x)$ (see Figure 1):

$$(4.7) \quad [\alpha_k, \beta_k] \setminus F_k(x) = \bigcup_{j=1}^{\infty} V_{k,j}.$$




FIGURE 1. A defining sequence $\mathscr{W}_\ell = \{(\beta_k, \alpha_{k+1}), V_{k,j} : k \geq \ell, j \geq 1\}$ for the Cantor set $C_\ell(x) = \{1/2\} \cup \bigcup_{k=\ell}^\infty F_k(x)$, and for each $k \geq \ell$ a defining sequence $\mathscr{V}_k = \{V_{k,j}\}_{j=1}^\infty$ for the Cantor set $F_k(x)$; see (4.7) and (4.8) for more explanation.

Now we turn to describe a defining sequence for $C_\ell(x)$ based on the defining sequence \mathscr{V}_k for $F_k(x)$ with $k \geq \ell$. Note that $\text{conv}(C_\ell(x)) = [\alpha_\ell, 1/2]$ and $\beta_k = \max F_k(x) < \min F_{k+1}(x) = \alpha_{k+1}$ for all $k \geq \ell$. Then by (4.5) and (4.7) it follows that (see Figure 1)

$$[\alpha_\ell, 1/2] \setminus C_\ell(x) = \bigcup_{k=\ell}^\infty (\beta_k, \alpha_{k+1}) \cup \bigcup_{k=\ell}^\infty \bigcup_{j=1}^\infty V_{k,j}.$$

We enumerate the open intervals (β_k, α_{k+1}) and $V_{j,k}$ with $k \geq \ell, j \geq 1$ in the following way:

$$(4.8) \quad \begin{array}{cccccc} (\beta_\ell, \alpha_{\ell+1}) & (\beta_{\ell+1}, \alpha_{\ell+2}) & (\beta_{\ell+2}, \alpha_{\ell+3}) & (\beta_{\ell+3}, \alpha_{\ell+4}) & \cdots \\ \downarrow & \downarrow & \downarrow & \downarrow & \\ V_{\ell,1} & V_{\ell,2} & V_{\ell,3} & V_{\ell,4} & \cdots \\ \swarrow & \swarrow & \swarrow & \swarrow & \\ V_{\ell+1,1} & V_{\ell+1,2} & V_{\ell+1,3} & V_{\ell+1,4} & \cdots \\ \swarrow & \swarrow & \swarrow & \swarrow & \\ V_{\ell+2,1} & V_{\ell+2,2} & V_{\ell+2,3} & V_{\ell+2,4} & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \end{array}$$

This means that we first remove from $[\alpha_\ell, 1/2]$ the open interval $(\beta_\ell, \alpha_{\ell+1})$, and next remove $V_{\ell,1}$, and then $(\beta_{\ell+1}, \alpha_{\ell+2}), V_{\ell+1,1}, V_{\ell,2}, (\beta_{\ell+2}, \alpha_{\ell+3})$, and so on. Thus, (4.8) gives a defining sequence $\mathscr{W}_\ell = \{(\beta_k, \alpha_{k+1}), V_{k,j} : k \geq \ell, j \geq 1\}$ for $C_\ell(x)$.

4.3. Hausdorff dimension of $\bigcap_{i=1}^p \Lambda(y_i)$. Based on the defining sequence \mathscr{W}_ℓ for $C_\ell(x)$ we will show that the thickness of $C_\ell(x)$ with respect to \mathscr{W}_ℓ goes to infinity as $\ell \rightarrow \infty$.

Proposition 4.4. *For any $x \in (0, 1/2)$ we have*

$$\lim_{\ell \rightarrow \infty} \tau_{\mathscr{W}_\ell}(C_\ell(x)) = +\infty.$$

Note by (4.8) that

$$L_{\mathscr{W}_\ell}((\beta_k, \alpha_{k+1})) = [\alpha_k, \beta_k], \quad R_{\mathscr{W}_\ell}((\beta_k, \alpha_{k+1})) = [\alpha_{k+1}, 1/2] \quad \text{for any } k \geq \ell;$$

and

$$L_{\mathcal{W}_\ell}(V_{k,j}) = L_{\mathcal{Y}_k}(V_{k,j}), \quad R_{\mathcal{W}_\ell}(V_{k,j}) = R_{\mathcal{Y}_k}(V_{k,j}) \quad \text{for any } k \geq \ell, j \geq 1.$$

Therefore, by (4.1) it follows that

$$(4.9) \quad \begin{aligned} \tau_{\mathcal{W}_\ell}(C_\ell(x)) &= \inf_{k \geq \ell, j \geq 1} \min \left\{ \frac{\beta_k - \alpha_k}{\alpha_{k+1} - \beta_k}, \frac{1/2 - \alpha_{k+1}}{\alpha_{k+1} - \beta_k}, \frac{|L_{\mathcal{Y}_k}(V_{k,j})|}{|V_{k,j}|}, \frac{|R_{\mathcal{Y}_k}(V_{k,j})|}{|V_{k,j}|} \right\} \\ &= \inf_{k \geq \ell} \min \left\{ \tau_{\mathcal{Y}_k}(F_k(x)), \frac{\beta_k - \alpha_k}{\alpha_{k+1} - \beta_k}, \frac{1/2 - \alpha_{k+1}}{\alpha_{k+1} - \beta_k} \right\}. \end{aligned}$$

Now we prove Proposition 4.4 in the following two cases: (A) $x \in (0, 1/2) \setminus \{1/4\}$; (B) $x = 1/4$.

Case A. $x \in (0, 1/2) \setminus \{1/4\}$. Write $\Psi_x(1/2) = (x_n)$. Since $\Psi_{1/4}(1/2) = 010^\infty$, there exists $m \geq 3$ such that $x_m = 1$. First we need the following bounds.

Lemma 4.5. *Let $x \in (0, 1/2) \setminus \{1/4\}$ with $(x_n) = \Psi_x(1/2)$, and let $m \geq 3$ such that $x_m = 1$.*

(i) *If $\lambda_1, \lambda_2 \in \Lambda(x)$ satisfy $\Psi_x(\lambda_1) = j_1 j_2 \cdots j_q 1^\infty$ and $\Psi_x(\lambda_2) = j_1 j_2 \cdots j_q 0^\infty$, then*

$$\lambda_2 - \lambda_1 \geq \frac{1}{4} \lambda_2^q.$$

(ii) *If $\lambda_3, \lambda_4 \in \Lambda(x)$ satisfy $\Psi_x(\lambda_3) = x_1 \cdots x_m j_1 \cdots j_q 10^\infty$ and $\Psi_x(\lambda_4) = x_1 \cdots x_m j_1 \cdots j_q 01^\infty$, then*

$$\lambda_4 - \lambda_3 \leq \min \left\{ 2(1 - 2\lambda_3) \lambda_3^{q+2}, 2(1 - 2\lambda_4) \frac{\lambda_4^{m+q}}{\lambda_3^{m-2}} \right\}.$$

Proof. For (i), let $\lambda_1, \lambda_2 \in \Lambda(x)$ with $\Psi_x(\lambda_1) = j_1 j_2 \cdots j_q 1^\infty$ and $\Psi_x(\lambda_2) = j_1 j_2 \cdots j_q 0^\infty$. Then by Lemma 2.2 we have $\lambda_1 < \lambda_2$. Note that $x = \pi_{\lambda_1}(j_1 j_2 \cdots j_q 1^\infty) = \pi_{\lambda_2}(j_1 j_2 \cdots j_q 0^\infty)$. Then

$$\begin{aligned} \lambda_2^q &= \pi_{\lambda_2}(0^q 1^\infty) = \pi_{\lambda_2}(j_1 j_2 \cdots j_q 1^\infty) - \pi_{\lambda_2}(j_1 j_2 \cdots j_q 0^\infty) \\ &= \pi_{\lambda_2}(j_1 j_2 \cdots j_q 1^\infty) - \pi_{\lambda_1}(j_1 j_2 \cdots j_q 1^\infty) \\ &\leq 4(\lambda_2 - \lambda_1), \end{aligned}$$

where the inequality follows by Lemma 2.1 (ii) since for any $(i_n) \in \Omega(x)$ we have

$$\frac{d\pi_\lambda((i_n))}{d\lambda} = \sum_{n=2}^{\infty} n \left(\frac{n-1}{n} - \lambda \right) i_n \lambda^{n-2} \leq \sum_{n=2}^{\infty} \frac{n-1}{2^{n-2}} = 4.$$

This proves (i).

For (ii) let $\lambda_3, \lambda_4 \in \Lambda(x)$ such that

$$\Psi_x(\lambda_3) = x_1 \cdots x_m j_1 \cdots j_q 10^\infty \quad \text{and} \quad \Psi_x(\lambda_4) = x_1 \cdots x_m j_1 \cdots j_q 01^\infty,$$

where $x_m = 1$ with $m \geq 3$. Then by Lemma 2.2 we have $\lambda_3 < \lambda_4$. Note that $x = \pi_{\lambda_3}(x_1 \cdots x_m j_1 j_2 \cdots j_q 10^\infty) = \pi_{\lambda_4}(x_1 \cdots x_m j_1 j_2 \cdots j_q 01^\infty)$. Then

$$(4.10) \quad \begin{aligned} &\pi_{\lambda_3}(0^{m+q} 10^\infty) - \pi_{\lambda_4}(0^{m+q+1} 1^\infty) \\ &= \pi_{\lambda_4}(x_1 \cdots x_m j_1 j_2 \cdots j_q 0^\infty) - \pi_{\lambda_3}(x_1 \cdots x_m j_1 j_2 \cdots j_q 0^\infty) \\ &\geq \frac{1}{2} \lambda_3^{m-2} (\lambda_4 - \lambda_3), \end{aligned}$$

where the last inequality follows by Lemma 2.1 (ii) since by using $x_m = 1$ with $m \geq 3$ and (2.5) we have

$$\frac{d\pi_\lambda(x_1 \cdots x_m j_1 j_2 \cdots j_q 0^\infty)}{d\lambda} \geq m \left(\frac{m-1}{m} - \lambda \right) x_m \lambda^{m-2} \geq \frac{1}{2} \lambda^{m-2}.$$

Therefore, by (4.10) and using $\lambda_3 < \lambda_4$ it follows that

$$\begin{aligned} \lambda_4 - \lambda_3 &\leq \frac{2}{\lambda_3^{m-2}} \left((1 - \lambda_3) \lambda_3^{m+q} - \lambda_4^{m+q+1} \right) \\ &\leq \frac{2}{\lambda_3^{m-2}} \left((1 - \lambda_3) \lambda_3^{m+q} - \lambda_3^{m+q+1} \right) = 2(1 - 2\lambda_3) \lambda_3^{q+2}, \end{aligned}$$

and

$$\begin{aligned} \lambda_4 - \lambda_3 &\leq \frac{2}{\lambda_3^{m-2}} \left((1 - \lambda_3) \lambda_3^{m+q} - \lambda_4^{m+q+1} \right) \\ &\leq \frac{2}{\lambda_3^{m-2}} \left((1 - \lambda_4) \lambda_4^{m+q} - \lambda_4^{m+q+1} \right) = 2(1 - 2\lambda_4) \frac{\lambda_4^{m+q}}{\lambda_3^{m-2}}. \end{aligned}$$

This completes the proof. \square

Proof of Proposition 4.4 with $x \in (0, 1/2) \setminus \{1/4\}$. Write $(x_n) = \Psi_x(1/2)$, and let $m \geq 3$ such that $x_m = 1$. Now take ℓ sufficiently large such that $n_\ell > m$, where the subsequence (n_k) is an enumeration of all indices n such that $x_n = 0$. Take $k \geq \ell$. By (4.9) we need to estimate lower bounds for $\tau_{\gamma_k}(F_k(x))$, $(\beta_k - \alpha_k)/(\alpha_{k+1} - \beta_k)$ and $(1/2 - \alpha_{k+1})/(\alpha_{k+1} - \beta_k)$, respectively.

(A1) A lower bound for $\tau_{\gamma_k}(F_k(x))$. Note by (4.6) that for any $j \in \mathbb{N}$ there exists a unique word $\omega = i_1 \dots i_q \in \{0, 1\}^*$ such that $V_{k,j} = V_{k,\omega}$. Then

$$(4.11) \quad \begin{aligned} L_{\gamma_k}(V_{k,j}) &= [\Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 1^\infty), \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 10^\infty)] =: [\gamma_1, \gamma_2], \\ R_{\gamma_k}(V_{k,j}) &= [\Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 01^\infty), \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 0^\infty)] =: [\gamma_3, \gamma_4], \end{aligned}$$

and

$$V_{k,j} = (\Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 10^\infty), \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 01^\infty)) = (\gamma_2, \gamma_3).$$

Note that $n_k > m$. Then by Lemma 4.5 it follows that

$$\gamma_2 - \gamma_1 \geq \frac{1}{4} \gamma_2^{n_k+q+1}, \quad \gamma_4 - \gamma_3 \geq \frac{1}{4} \gamma_4^{n_k+q+1},$$

and

$$\gamma_3 - \gamma_2 \leq 2(1 - 2\gamma_2) \gamma_2^{n_k+q-m+2}.$$

Observe by Lemma 2.2 and (4.11) that $\alpha_k \leq \gamma_1 < \gamma_2 < \gamma_3 < \gamma_4$, where $\alpha_k = \min F_k(x)$. So, for any $j \geq 1$ we obtain that

$$\begin{aligned} \frac{|L_{\gamma_k}(V_{k,j})|}{|V_{k,j}|} &= \frac{\gamma_2 - \gamma_1}{\gamma_3 - \gamma_2} \geq \frac{\gamma_2^{m-1}}{8(1 - 2\gamma_2)} \geq \frac{\alpha_k^{m-1}}{8(1 - 2\alpha_k)}, \\ \frac{|R_{\gamma_k}(V_{k,j})|}{|V_{k,j}|} &= \frac{\gamma_4 - \gamma_3}{\gamma_3 - \gamma_2} \geq \frac{\gamma_4^{m-1}}{8(1 - 2\gamma_2)} \geq \frac{\alpha_k^{m-1}}{8(1 - 2\alpha_k)}. \end{aligned}$$

Therefore, for any $k \geq \ell$ we have

$$(4.12) \quad \tau_{\mathcal{Y}_k}(F_k(x)) \geq \frac{\alpha_k^{m-1}}{8(1-2\alpha_k)} \geq \frac{x^{m-1}}{8(1-2\alpha_k)},$$

where the last inequality holds since $\alpha_k \in \Lambda(x) \subset [x, 1/2]$.

(A2) A lower bound for $(\beta_k - \alpha_k)/(\alpha_{k+1} - \beta_k)$. Note that

$$(4.13) \quad \alpha_k = \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1^\infty), \quad \beta_k = \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 10^\infty).$$

Furthermore, by the definition of (n_k) it follows that

$$(4.14) \quad \alpha_{k+1} = \Psi_x^{-1}(x_1 \cdots x_{n_k-1} x_{n_k} \cdots x_{n_{k+1}-1} 1^\infty) = \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 01^\infty).$$

So, by (4.13), (4.14) and Lemma 4.5 we obtain that

$$\beta_k - \alpha_k \geq \frac{1}{4} \beta_k^{n_k}, \quad \alpha_{k+1} - \beta_k \leq 2(1 - 2\beta_k) \beta_k^{n_k - m + 1}.$$

This implies

$$(4.15) \quad \frac{\beta_k - \alpha_k}{\alpha_{k+1} - \beta_k} \geq \frac{\beta_k^{m-1}}{8(1-2\beta_k)} \geq \frac{x^{m-1}}{8(1-2\beta_k)}.$$

(A3) A lower bound for $(1/2 - \alpha_{k+1})/(\alpha_{k+1} - \beta_k)$. Note by (4.13), (4.14) and Lemma 4.5 that

$$\alpha_{k+1} - \beta_k \leq 2(1 - 2\alpha_{k+1}) \frac{\alpha_{k+1}^{n_k-1}}{\beta_k^{m-2}}.$$

This implies that

$$(4.16) \quad \frac{1/2 - \alpha_{k+1}}{\alpha_{k+1} - \beta_k} \geq \frac{\beta_k^{m-2}}{4\alpha_{k+1}^{n_k-1}} > \frac{x^{m-2}}{4 \cdot 2^{1-n_k}},$$

where the last inequality follows by $x < \beta_k < \alpha_{k+1} < 1/2$.

Note that $m \geq 3$ is a constant depending on x , and $\alpha_k \nearrow 1/2$, $\beta_k \nearrow 1/2$ as $k \rightarrow \infty$. Letting $k \rightarrow \infty$ in (4.12), (4.15) and (4.16) we conclude by (4.9) that

$$\tau_{\mathcal{Y}_\ell}(C_\ell(x)) \rightarrow +\infty \quad \text{as } \ell \rightarrow \infty.$$

This completes the proof. \square

Case B. $x = 1/4$. Then $(x_n) = \Phi_x(1/2) = 010^\infty$. We first need the following estimation.

Lemma 4.6. *Let $x = 1/4$.*

(i) *If $\lambda_1, \lambda_2 \in \Lambda(x)$ satisfy $\Psi_x(\lambda_1) = 010^m j_1 j_2 \cdots j_q 1^\infty$ and $\Psi_x(\lambda_2) = 010^m j_1 j_2 \cdots j_q 0^\infty$, then*

$$\lambda_2 - \lambda_1 \geq \frac{\lambda_2^{m+2+q}}{1 - 2\lambda_1 + (m+3)2^{-m}}.$$

(ii) *If $\lambda_3, \lambda_4 \in \Lambda(x)$ satisfy $\Psi_x(\lambda_3) = 01j_1 j_2 \cdots j_q 10^\infty$ and $\Psi_x(\lambda_4) = 01j_1 j_2 \cdots j_q 01^\infty$, then*

$$\lambda_4 - \lambda_3 \leq \lambda_3^{2+q}.$$

Proof. For (i) we note by Lemma 2.2 that $\lambda_1 < \lambda_2$. Since $x = \pi_{\lambda_1}(010^m j_1 j_2 \cdots j_q 1^\infty) = \pi_{\lambda_2}(010^m j_1 j_2 \cdots j_q 0^\infty)$, we have

$$\begin{aligned} \lambda_2^{m+2+q} &= \pi_{\lambda_2}(0^{m+2+q} 1^\infty) = \pi_{\lambda_2}(010^m j_1 j_2 \cdots j_q 1^\infty) - \pi_{\lambda_2}(010^m j_1 j_2 \cdots j_q 0^\infty) \\ &= \pi_{\lambda_2}(010^m j_1 j_2 \cdots j_q 1^\infty) - \pi_{\lambda_1}(010^m j_1 j_2 \cdots j_q 1^\infty) \\ &\leq (1 - 2\lambda_1 + (m+3)2^{-m})(\lambda_2 - \lambda_1), \end{aligned}$$

where the inequality follows by Lemma 2.1 (ii) since by (2.5) we have

$$\begin{aligned} \frac{d\pi_\lambda(010^m j_1 j_2 \cdots j_q 1^\infty)}{d\lambda} &\leq (1 - 2\lambda) + \sum_{n=m+3}^{\infty} n \left(\frac{n-1}{n} - \lambda \right) \lambda^{n-2} \\ &\leq 1 - 2\lambda + \sum_{n=m+3}^{\infty} \frac{n-1}{2^{n-2}} = 1 - 2\lambda + (m+3)2^{-m}. \end{aligned}$$

This proves (i).

For (ii), note by Lemma 2.2 that $\lambda_3 < \lambda_4$. Then by using $x = \pi_{\lambda_3}(01j_1 j_2 \cdots j_q 10^\infty) = \pi_{\lambda_4}(01j_1 j_2 \cdots j_q 01^\infty)$ it follows that we have

$$\begin{aligned} \pi_{\lambda_3}(0^{2+q} 10^\infty) - \pi_{\lambda_4}(0^{3+q} 1^\infty) &= \pi_{\lambda_4}(01j_1 j_2 \cdots j_q 0^\infty) - \pi_{\lambda_3}(01j_1 j_2 \cdots j_q 0^\infty) \\ &\geq \pi_{\lambda_4}(010^\infty) - \pi_{\lambda_3}(010^\infty) \\ &= (1 - \lambda_3 - \lambda_4)(\lambda_4 - \lambda_3). \end{aligned}$$

This implies that

$$\begin{aligned} \lambda_4 - \lambda_3 &\leq \frac{1}{1 - \lambda_3 - \lambda_4} \left((1 - \lambda_3)\lambda_3^{2+q} - \lambda_4^{3+q} \right) \\ &\leq \frac{1}{1 - \lambda_3 - \lambda_4} \left((1 - \lambda_3)\lambda_3^{2+q} - \lambda_4\lambda_3^{2+q} \right) = \lambda_3^{2+q}, \end{aligned}$$

as desired. \square

Proof of Proposition 4.4 with $x = 1/4$. Let $x = 1/4$. Then $(x_n) = \Psi_x(1/2) = 010^\infty$. Let $\ell \geq 1$ such that $n_\ell > 2$. Take $k \geq \ell$. Similar to Case A we consider lower bounds for $\tau_{\gamma_k}(F_k(x))$, $(\beta_k - \alpha_k)/(\alpha_{k+1} - \beta_k)$ and $(1/2 - \alpha_{k+1})/(\alpha_{k+1} - \beta_k)$, respectively.

(B1) A lower bound for $\tau_{\gamma_k}(F_k(x))$. For $j \in \mathbb{N}$ let $\omega = i_1 i_2 \cdots i_q$ be the unique word such that $V_{k,j} = V_{k,\omega}$. Then

$$\begin{aligned} L\gamma_k(V_{k,j}) &= [\Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 1^\infty), \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 10^\infty)] =: [\gamma_1, \gamma_2], \\ R\gamma_k(V_{k,j}) &= [\Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 01^\infty), \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 0^\infty)] =: [\gamma_3, \gamma_4], \end{aligned}$$

and

$$V_{k,j} = (\Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 10^\infty), \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1 \omega 01^\infty)) = (\gamma_2, \gamma_3).$$

Then $\alpha_k \leq \gamma_1 < \gamma_2 < \gamma_3 < \gamma_4$. Note that $x_1 \dots x_{n_k-1} = 010^{n_k-3}$. Then by Lemma 4.6 it follows that

$$\gamma_2 - \gamma_1 \geq \frac{\gamma_2^{n_k+q+1}}{1 - 2\gamma_1 + n_k 2^{3-n_k}}, \quad \gamma_4 - \gamma_3 \geq \frac{\gamma_4^{n_k+q+1}}{1 - 2\gamma_3 + n_k 2^{3-n_k}},$$

and

$$\gamma_3 - \gamma_2 \leq \gamma_2^{n_k+q}.$$

This implies that for all $j \geq 1$,

$$\begin{aligned} \frac{|L_{\gamma_k}(V_{k,j})|}{|V_{k,j}|} &= \frac{\gamma_2 - \gamma_1}{\gamma_3 - \gamma_2} \geq \frac{\gamma_2}{1 - 2\gamma_1 + n_k 2^{3-n_k}} \geq \frac{\alpha_k}{1 - 2\alpha_k + n_k 2^{3-n_k}}, \\ \frac{|R_{\gamma_k}(V_{k,j})|}{|V_{k,j}|} &= \frac{\gamma_4 - \gamma_3}{\gamma_3 - \gamma_2} \geq \frac{\gamma_4}{1 - 2\gamma_3 + n_k 2^{3-n_k}} \geq \frac{\alpha_k}{1 - 2\alpha_k + n_k 2^{3-n_k}}. \end{aligned}$$

Therefore, for any $k \geq \ell$ we have

$$(4.17) \quad \tau_{\gamma_k}(F_k(x)) \geq \frac{\alpha_k}{1 - 2\alpha_k + n_k 2^{3-n_k}}.$$

(B2) A lower bound for $(\beta_k - \alpha_k)/(\alpha_{k+1} - \beta_k)$. Note that

$$\alpha_k = \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 1^\infty), \quad \beta_k = \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 10^\infty),$$

and

$$\alpha_{k+1} = \Psi_x^{-1}(x_1 \cdots x_{n_k-1} x_{n_k} \cdots x_{n_{k+1}-1} 1^\infty) = \Psi_x^{-1}(x_1 x_2 \cdots x_{n_k-1} 01^\infty).$$

Then by Lemma 4.6 it follows that

$$(4.18) \quad \beta_k - \alpha_k \geq \frac{\beta_k^{n_k}}{1 - 2\alpha_k + n_k 2^{3-n_k}}, \quad \alpha_{k+1} - \beta_k \leq \beta_k^{n_k-1}.$$

This implies

$$(4.19) \quad \frac{\beta_k - \alpha_k}{\alpha_{k+1} - \beta_k} \geq \frac{\beta_k}{1 - 2\alpha_k + n_k 2^{3-n_k}}.$$

(B3) A lower bound for $(1/2 - \alpha_{k+1})/(\alpha_{k+1} - \beta_k)$. Note that

$$\frac{1}{4} = x = \pi_{\alpha_{k+1}}(x_1 x_2 \cdots x_{n_k-1} 01^\infty) = \pi_{\alpha_{k+1}}(010^{n_k-2} 1^\infty) = (1 - \alpha_{k+1})\alpha_{k+1} + \alpha_{k+1}^{n_k}.$$

This implies that

$$(4.20) \quad \left(\frac{1}{2} - \alpha_{k+1}\right)^2 = \alpha_{k+1}^{n_k}, \quad \text{and thus} \quad \frac{1}{2} - \alpha_{k+1} = \alpha_{k+1}^{n_k/2}.$$

Note by (4.18) that $\alpha_{k+1} - \beta_k \leq \beta_k^{n_k-1} < \alpha_{k+1}^{n_k-1}$. This, together with (4.20), implies that

$$(4.21) \quad \frac{1/2 - \alpha_{k+1}}{\alpha_{k+1} - \beta_k} \geq \frac{1}{\alpha_{k+1}^{n_k/2-1}}.$$

Since $\lim_{k \rightarrow \infty} \alpha_k = \lim_{k \rightarrow \infty} \beta_k = 1/2$, letting $k \rightarrow \infty$ in (4.17), (4.19) and (4.21) we conclude by (4.9) that $\tau_{\mathcal{W}_\ell}(C_\ell(x)) \rightarrow +\infty$ as $\ell \rightarrow \infty$. This completes the proof. \square

Proof of Theorem 1.4. Let $y_1, \dots, y_p \in (0, 1)$. Then by (4.2), (4.5) and Proposition 4.4 it follows that each $\Lambda(y_i)$ contains a sequence of Cantor subsets $C_\ell(y_i)$, $\ell \geq 1$ such that $\max C_\ell(y_i) = 1/2$ for all $\ell \geq 1$, and the thickness $\tau(C_\ell(y_i)) \rightarrow +\infty$ as $\ell \rightarrow \infty$. So, by Lemma 4.2 and Remark 4.3 (i) it follows that for each $\ell \geq 1$ and for any $i, j \in \{1, 2, \dots, p\}$ the intersection $C_\ell(y_i) \cap C_\ell(y_j)$ contains a Cantor subset $C_\ell(y_i, y_j)$ such that

$$(4.22) \quad \tau(C_\ell(y_i, y_j)) \rightarrow +\infty \quad \text{as } \ell \rightarrow \infty.$$

Note that for any $k \in \{1, 2, \dots, p\}$ we have $\min C_\ell(y_k) \nearrow 1/2 = \max C_\ell(y_k)$ as $\ell \rightarrow \infty$. Then by (4.22) it follows that the maximum point $1/2$ is an accumulation point of $C_\ell(y_i) \cap C_\ell(y_j)$ for any $\ell \geq 1$. So, by Remark 4.3 (ii) we can require that the resulting Cantor set $C_\ell(y_i, y_j) \subset C_\ell(y_i) \cap C_\ell(y_j)$ has the maximum point $1/2$ for any $i, j \in \{1, 2, \dots, p\}$ and $\ell \geq 1$.

Proceeding this argument for all y_1, y_2, \dots, y_p we obtain that for any $\ell \geq 1$ the intersection $\bigcap_{i=1}^p C_\ell(y_i)$ contains a Cantor subset $C_\ell(y_1, \dots, y_p)$ such that $\max C_\ell(y_1, \dots, y_p) = 1/2$, and the thickness $\tau(C_\ell(y_1, \dots, y_p)) \rightarrow +\infty$ as $\ell \rightarrow \infty$. Therefore, by Lemma 4.1 it follows that

$$\dim_H \bigcap_{i=1}^p \Lambda(y_i) \geq \dim_H \bigcap_{i=1}^p C_\ell(y_i) \geq \dim_H C_\ell(y_1, \dots, y_p) \geq \frac{\log 2}{\log(2 + \frac{1}{\tau(C_\ell(y_1, \dots, y_p))})} \rightarrow 1,$$

as $\ell \rightarrow \infty$. This completes the proof. \square

At the end of this section we remark that in the proof of Theorem 1.4 we construct a sequence of Cantor subsets $C_\ell(y_1, \dots, y_p), \ell \geq 1$ in the intersection $\bigcap_{i=1}^p \Lambda(y_i)$ such that the thickness $\tau(C_\ell(y_1, \dots, y_p)) \rightarrow +\infty$ as $\ell \rightarrow \infty$. By a recent work of Yavicoli [12, remark of Theorem 4] it follows that the intersection $\bigcap_{i=1}^p \Lambda(y_i)$ contains arbitrarily long arithmetic progression.

5. FINAL REMARKS

At the end of this paper we point out that our results Theorem 1.1–1.4 can be extended to higher dimensions. To illustrate this we give two examples.

Example 5.1. For $\lambda \in (0, 1/2]$ let K_λ be the self-similar set defined in (1.1). Then for $n \in \mathbb{N}$ the product set $\bigotimes_{i=1}^n K_\lambda$ is a also self-similar set in \mathbb{R}^n . For $\mathbf{a} = (a_1, \dots, a_n) \in (0, 1/2)^n$ let

$$\Lambda(\mathbf{a}) := \{\lambda \in (0, 1/2] : \mathbf{a} \in \bigotimes_{i=1}^n K_\lambda\}$$

be the set of parameters $\lambda \in (0, 1/2]$ such that the n -dimensional self-similar set $\bigotimes_{i=1}^n K_\lambda$ contains the given point \mathbf{a} . It is clear that $\Lambda(\mathbf{a}) = \bigcap_{i=1}^n \Lambda(a_i)$, where $\Lambda(x)$ is defined as in (1.2). So, by Theorems 1.2 and 1.4 it follows that $\Lambda(\mathbf{a})$ has zero Lebesgue measure and full Hausdorff dimension for any $\mathbf{a} \in (0, 1/2)^n$.

Example 5.2. Let $n \in \mathbb{N}$ and let $\lambda_i \in (0, 1/2]$ for all $1 \leq i \leq n$. Then $\bigotimes_{i=1}^n K_{\lambda_i} \subset \mathbb{R}^n$ is a self-affine set generated by the IFS $\{(\lambda_1 x_1, \lambda_2 x_2, \dots, \lambda_n x_n) + \mathbf{i} : \mathbf{i} \in \bigotimes_{i=1}^n \{0, 1 - \lambda_i\}\}$. For any $\mathbf{b} = (b_1, \dots, b_n) \in (0, 1/2)^n$ let

$$\Lambda'(\mathbf{b}) := \{(\lambda_1, \lambda_2, \dots, \lambda_n) \in (0, 1/2]^n : \mathbf{b} \in \bigotimes_{i=1}^n K_{\lambda_i}\}.$$

Then $(\lambda_1, \lambda_2, \dots, \lambda_n) \in \Lambda'(\mathbf{b})$ if and only if $b_i \in K_{\lambda_i}$ for all $1 \leq i \leq n$, which is also equivalent to $\lambda_i \in \Lambda(b_i)$ for all $1 \leq i \leq n$. So, $\Lambda'(\mathbf{b}) = \bigotimes_{i=1}^n \Lambda(b_i)$. By Theorem 1.1 it follows that for any $\mathbf{b} \in (0, 1/2)^n$ the set $\Lambda'(\mathbf{b})$ is a Cantor set in \mathbb{R}^n , i.e., it is a non-empty compact, totally disconnected and perfect set in \mathbb{R}^n . Furthermore, by [3, product formula 7.2] and

Theorem 1.2 we obtain that $\Lambda'(\mathbf{b})$ has Lebesgue measure zero and $\dim_H \Lambda'(\mathbf{b}) = n$ for any $\mathbf{b} \in (0, 1/2)^n$.

ACKNOWLEDGEMENTS

K. Jiang was supported by NSFC No. 11701302, Zhejiang Provincial NSF No. LY20A010009, and the K.C. Wong Magna Fund in Ningbo University. D. Kong was supported by NSFC No. 11971079 and the Fundamental and Frontier Research Project of Chongqing No. cstc2019jcyj-msxmX0338 and No. cx2019067. W. Li was supported by NSFC No. 12071148 and Science and Technology Commission of Shanghai Municipality (STCSM) No. 18dz2271000.

REFERENCES

- [1] S. Astels. Cantor sets and numbers with restricted partial quotients. *Trans. Amer. Math. Soc.*, 352(1):133–170, 2000.
- [2] D. Boes, R. Darst, and P. Erdős. Fat, symmetric, irrational Cantor sets. *Amer. Math. Monthly*, 88(5):340–341, 1981.
- [3] K. Falconer. *Fractal geometry: Mathematical foundations and applications*. John Wiley & Sons Ltd., Chichester, 1990.
- [4] B. R. Hunt, I. Kan, and J. A. Yorke. When Cantor sets intersect thickly. *Trans. Amer. Math. Soc.*, 339(2):869–888, 1993.
- [5] J. E. Hutchinson. Fractals and self-similarity. *Indiana Univ. Math. J.*, 30(5):713–747, 1981.
- [6] K. Jiang, D. Kong, and W. Li. How likely can a point be in different cantor sets. *arxiv.org/abs/2102.13264*, 2021.
- [7] D. Lind and B. Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995.
- [8] S. E. Newhouse. Nondensity of axiom A(a) on S^2 . In *Global Analysis (Proc. Sympos. Pure Math., Vol. XIV, Berkeley, Calif., 1968)*, pages 191–202. Amer. Math. Soc., Providence, R.I., 1970.
- [9] S. E. Newhouse. The abundance of wild hyperbolic sets and non-smooth stable sets for diffeomorphisms. *Inst. Hautes Etudes Sci. Publ. Math.*, 50:101–151, 1979.
- [10] J. Palis and F. Takens. *Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations. Fractal dimensions and infinitely many attractors. Cambridge Studies in Advanced Mathematics, 35*. Cambridge University Press, Cambridge, 1993.
- [11] G. Pianigiani and J. A. Yorke. Expanding maps on sets which are almost invariant. Decay and chaos. *Trans. Amer. Math. Soc.*, 252:351–366, 1979.
- [12] A. Yvicoli. Patterns in thick comapct sets. *Israel J. Math.*, pages 1–32, 2021.

(Z. Wang) SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, PEOPLE'S REPUBLIC OF CHINA

Email address: zhiqiangwzy@163.com

(K. Jiang) DEPARTMENT OF MATHEMATICS, NINGBO UNIVERSITY, PEOPLE'S REPUBLIC OF CHINA

Email address: jiangkan@nbu.edu.cn

(D. Kong) COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, 401331, CHONGQING, P.R.CHINA

Email address: derongkong@126.com

(W. Li) SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, PEOPLE'S REPUBLIC OF CHINA

Email address: wxli@math.ecnu.edu.cn