

A Spectral Approach to Off-Policy Evaluation for POMDPs

Yash Nair^{1*}, Nan Jiang²

¹Harvard University

²University of Illinois Urbana-Champaign

yashnair@college.harvard.edu, nanjiang@illinois.edu

Abstract

We consider off-policy evaluation (OPE) in Partially Observable Markov Decision Processes, where the evaluation policy depends only on observable variables but the behavior policy depends on latent states (Tennenholtz *et al.*, 2020a). Prior work on this problem uses a causal identification strategy based on one-step observable proxies of the hidden state, which relies on the invertibility of certain one-step moment matrices. In this work, we relax this requirement by using spectral methods and extending one-step proxies both into the past and future. We empirically compare our OPE methods to existing ones and demonstrate their improved prediction accuracy and greater generality. Lastly, we derive a separate Importance Sampling (IS) algorithm which relies on rank, distinctness, and positivity conditions, and not on the strict sufficiency conditions of observable trajectories with respect to the reward and hidden-state structure required by Tennenholtz *et al.* (2020a).

1 Introduction

We consider the problem of estimating the value of a policy—called the *evaluation policy*—given access only to a batch of trajectories obtained from a different policy—called the *behavior policy*—known as the *off-policy evaluation* (OPE) problem. The OPE problem is well-motivated by its application to real-world scenarios in which the deployment of the evaluation policy is potentially too costly for its performance to be determined by directly intervening with it, but rather must be inferred from previously existing data under a different policy. As an example, in healthcare settings, it may be unsafe to directly test a new experimental treatment with potentially unknown harms and side-effects on patients, and thus reasoning based on previous treatment strategies from electronic health records (EHRs) to infer the new treatment’s efficacy is necessary. Alternatively, when designing a new personalized educational curriculum, which could potentially have harmful effects on students’ learning (Mandel *et al.*, 2016), the use of historical curriculum data is required.

OPE has received much attention in the setting of Markov Decision Processes (MDPs) in which the underlying state is fully observable, and both evaluation and behavior policies are (potentially non-deterministic and/or non-stationary) functions of state (Sutton and Barto, 2018; Jiang and Li, 2016; Precup *et al.*, 2000). We, however, consider the more challenging regime of Partially Observable MDPs (POMDPs), in which one has access only to *observable* trajectories formed under the behavior policy, which do not include underlying latent states (i.e. hidden confounders), and must infer the value of the evaluation policy given only these observables. Following Tennenholtz *et al.* (2020a), we consider the setting in which the behavior policy depends on the latent state, whereas the evaluation policy at any given time step depends only on the observed history up to that time step. This choice is motivated, for example, by the medical treatment problem and autonomous driving scenarios described by Tennenholtz *et al.* (2020a).

In this setup, Tennenholtz *et al.* (2020a) builds on the work of Miao *et al.* (2018) and gives a causal effect identification scheme, given two conditionally independent views of the confounder. Their estimator uses one-step observable proxies of the hidden state, and crucially relies on the invertibility of certain *one-step* moment matrices, which is a strong assumption and limits the applicability of their method. Furthermore, they use moment matrices whose entries are *conditional* probabilities, which can be difficult to estimate when the condition event is rare and impossible if measure zero.

In this work we extend Tennenholtz *et al.* (2020a) and address the above issues, inspired by techniques for solving similar problems in the literature of spectral learning of Predictive State Representations (PSRs) and Hidden Markov Models (HMMs) (Hsu *et al.*, 2012; Boots *et al.*, 2011; Kulesza *et al.*, 2015). Our work extends and improves upon Tennenholtz *et al.* (2020a) in several aspects:

- Our estimators work for a broader family of POMDPs, specifically those in which the observation space is larger than hidden state space, as is the case in many real-world settings.
- Our estimators estimate matrices of *joint* probabilities rather than *conditionals*, and thus improve generality by not having to condition on probability zero events.
- We are able to further relax a crucial assumption in their

*Contact Author

work regarding the invertibility of one-step moment matrices by working with *multi-step proxies* for the hidden state; for example, our estimator uses the entire extended history preceding the confounder as one of the proxies, rather than just the previous time step.

While we are able to successfully use multi-step histories as the past proxy, we show that there is difficulty in directly using extended futures as the future proxy (which would further relax rank assumptions). However, by employing the eigendecomposition technique of Kuroki and Pearl (2014), we are able—with minimal additional assumptions—to give an estimator which also incorporates extended futures and thus applies even more generally.

We then empirically validate our estimators, comparing them to prior estimators (Tennenholtz *et al.*, 2020a) and demonstrate their superior predictive accuracy. Finally, by again employing the eigendecomposition technique of Kuroki and Pearl (2014), we provide a separate Importance Sampling (IS) algorithm which requires only rank, positivity, and distinctness conditions on certain probability matrices, and not the sufficiency assumptions required by the IS algorithm of Tennenholtz *et al.* (2020a).

2 Related Work

OPE in POMDPs Most closely related to our work is that of Tennenholtz *et al.* (2020a), who consider the finite-horizon OPE problem in tabular POMDPs with equal-sized hidden state and observation spaces by extending Miao *et al.* (2018) and writing the probability of seeing reward r_t , in terms of observable moment matrices under the behavior measure. These matrices have dimensions which scale with the size of the observation space, \mathcal{Z} , and are required to be invertible. They further develop a Decoupled POMDP model which factors the state space, \mathcal{U} , into both observed and unobserved variables, allowing for a modified OPE algorithm whose moment matrices scale with $|\mathcal{U}|$, which is often much smaller than $|\mathcal{Z}|$. Our work mitigates difficulty in the more-general POMDP setting by allowing for rich observation spaces with greater cardinality than state space, requiring that the observable matrices have rank $|\mathcal{U}|$, rather than $|\mathcal{Z}|$, and further relaxes assumptions and attains superior performance by extending both histories and futures. Additionally, Oberst and Sontag (2019) consider counterfactuals, asking about the outcome of a counterfactual trajectory under the evaluation policy, given the observable trajectory’s outcome under the behavior policy. Their work differs from ours in that, just as in Tennenholtz *et al.* (2020a), we do not ask about the evaluation policy’s counterfactual performance compared to the behavior policy’s, but rather are concerned with determining the value of the evaluation policy given observable trajectories under the behavior.

OPE with unobserved confounding The OPE problem has also been studied in the setting of the MDP with Unmeasured Confounding (MDPUC) (Zhang and Bareinboim, 2016). The MDPUC framework is orthogonal to POMDPs, with the local confounders assumed to be i.i.d. across time steps. The OPE problem is considered in an infinite-horizon, non-tabular MDPUC by Bennett *et al.* (2020). Critically,

however, their work assumes access to a latent variable model for confounders, which we do not assume. Specifically, in the spirit of Tennenholtz *et al.* (2020a) and Miao *et al.* (2018), our initial relaxations of rank assumptions do not even depend on non-parametric identification of the latent variable model. Our extended-future estimator and IS results, however, do rest on the identification of aspects of the latent variable model, but obtain estimates from observable data instead of assuming access to a model.

Causal effect identification Our work is closely related to unobserved confounding in the causal inference literature, and, specifically, the use of negative controls to minimize confounding bias. Of particular relevance are Miao *et al.* (2018), Kuroki and Pearl (2014) and Greenland and Lash (2012), which all consider the static problem of identifying causal effects, assuming multiple observable proxies of the latent confounder. Kuroki and Pearl (2014) first identify the confounder’s error mechanism via eigenvalue analysis and then use the matrix adjustment method (Greenland and Lash, 2012) to determine the causal effect, while Miao *et al.* (2018) directly determine the causal effect without identifying any of the aspects of the confounding model, allowing for weaker assumptions. Our work extends the latter’s result to the setting of POMDPs just as in Tennenholtz *et al.* (2020a); our spectral result in the setting of POMDPs also applies to the static setting and hence relaxes rank assumptions in Miao *et al.* (2018). Additionally, we extend Kuroki and Pearl (2014)’s analysis to identify causal effects of actions on futures in our extended-future estimator as well as to recover the behavior policy and reward distribution in our IS algorithm.

3 Preliminaries

POMDPs A POMDP is a 7-tuple $\langle \mathcal{U}, \mathcal{Z}, \mathcal{A}, O, T, R, H \rangle$, where \mathcal{U} and \mathcal{Z} denote the hidden state and observation spaces, respectively, \mathcal{A} denotes the set of actions, $O : \mathcal{U} \times \mathcal{Z} \rightarrow [0, 1]$ gives observation emission probabilities, with $O(z|u) := \Pr[z|u]$, $T : \mathcal{U} \times \mathcal{U} \times \mathcal{A}$ governs the hidden state transition dynamics, with $T(u'|u, a) := \Pr[u'|u, a]$, R is a (potentially non-deterministic) function of the hidden state and action giving the reward which is supported in the set \mathcal{R} , and H denotes the horizon. At each time step, the agent selects an action, a , transitions under T , to a new hidden state, u' , based on the current hidden state, u , and a and then observes an observation, z , drawn according to $O(\cdot|u')$, and reward, r , drawn from its conditional distribution over the hidden state and action. In this work, we restrict ourselves to finite \mathcal{U} , \mathcal{Z} , and \mathcal{A} . Furthermore, we assume that $|\mathcal{U}| \leq |\mathcal{Z}|$, reflecting the fact that in most real-world settings, the hidden state space is often much smaller than the rich observation space.

Using the notation of Tennenholtz *et al.* (2020a), a trajectory τ of length t denotes a sequence $(u_0, z_0, a_0, \dots, u_t, z_t, a_t)$ while τ^o denotes the observable trajectory $(z_0, a_0, \dots, z_t, a_t)$. We use \mathcal{T}_t to denote the set of trajectories of length t , and, correspondingly, \mathcal{T}_t^o to denote the set of observable trajectories of length t . Additionally, an observable history of length t is given by a sequence of the form $(z_0, a_0, \dots, a_{t-1}, z_t)$, and is denoted

by h_t^o . Finally, we assume the existence of an observation z_{-1} preceding the initial time step which is conditionally independent of z_0 and a_0 given u_0 .

Policies Let π_b and π_e denote the behavior and evaluation policies, respectively. We consider the setting in which the behavior policy at time step t , $\pi_b^{(t)}$, depends only on the hidden state, u , whereas the evaluation policy at time t , $\pi_e^{(t)}$, depends on the observable history, h_t^o . In particular, $\pi_b^{(t)}(a_t|u_t)$ denotes the probability that an agent (with access to hidden state) following π_b selects action a_t at time step t given that it is at u_t , and $\pi_e^{(t)}(a_t|h_t^o)$ denotes the probability that an agent (without access to hidden state) following π_e selects action a_t at time step t given that it has seen the observable history h_t^o . Additionally, we consider the finite-horizon undiscounted setting so that for any policy π (either depending on observable histories or hidden state), its value $v_H(\pi)$ is given by $\mathbb{E}_\pi \left[\sum_{t=0}^H r_t \right]$, the expected sum of rewards gotten by following π . Finally, let P^b and P^e denote measures over trajectories induced by the behavior and evaluation policies, respectively.

In addition to the above, we use a double vertical bar notation to indicate intervening (Boots *et al.*, 2011). For example, $P(z_0, \dots, z_t | a_0, \dots, a_{t-1})$ denotes the probability of seeing observations (z_0, \dots, z_t) , given that we intervened with actions a_0, \dots, a_{t-1} . Given a policy, π , $P^\pi(z_0, \dots, z_t | a_0, \dots, a_{t-1})$ is in general not equal to $P(z_0, \dots, z_t | a_0, \dots, a_{t-1})$, as in the former, actions are random variables, which may convey information about hidden state, whereas in the latter, they are non-random interventions. This can also be written using the *do* operator of Pearl (2009).

We use matrix notations similar to Tennenholtz *et al.* (2020a). Let random variables $\mathbf{x}, \mathbf{y}, \mathbf{z}$ be supported in $X = \{x_1, \dots, x_{n_1}\}, Y = \{y_1, \dots, y_{n_2}\}, Z = \{z_1, \dots, z_{n_3}\}$. Then $P(y|X), P(X), P(Y, x, Z)$ denote a row vector, column vector, and matrix, with $(P(y|X))_i = \Pr[y=x_i], (P(X))_i = \Pr[x_i], (P(Y, x, Z))_{i,j} = \Pr[y_i, x, z_j]$. For joint probabilities, the first (second) set listed denotes vectorization across rows (columns); for conditionals, the set listed before (after) the conditioning bar indicates vectorization across rows (columns).

Problem Formally, the OPE problem asks, given a batch of observable trajectories collected under the behavior policy, to estimate the value of the evaluation policy. In other words, we must estimate $v_H(\pi_e)$ given only observable trajectories τ^o drawn from the behavior measure P^b .

3.1 One-Step Proxies

We now briefly review the OPE algorithm of Tennenholtz *et al.* (2020a), who approach the problem by estimating $P^e(r_t)$, which can then be used to approximate $v_H(\pi_e)$ by estimating the expected reward at each timestep. (In Sections 4 and 5 we also focus on estimating $P^e(r_t)$.) They make the following assumptions on observable probability matrices.

Assumption 1. $P^b(Z_i|a_i, Z_{i-1})$ is invertible $\forall i \leq H, a_i \in \mathcal{A}$.

Defining $\Pi_e^e(\tau^o) = \prod_{i=0}^t \pi_e^{(i)}(a_i|h_i^o)$, they give the following estimator:

Theorem 1. Under Assumption 1, $P^e(r_t)$ can be written as

$$\sum_{\tau^o \in \mathcal{T}_t^o} \Pi_e^e(\tau^o) P^b(r_t, z_t | a_t, Z_{t-1}) \cdot \prod_{i=t-1}^0 (P^b(Z_{i+1}|a_{i+1}, Z_i)^{-1} P^b(Z_{i+1}, z_i | a_i, Z_{i-1})) \cdot P^b(Z_0|a_0, Z_{-1})^{-1} P^b(Z_0). \quad (1)$$

We give a sketch of their proof (for details, see Tennenholtz *et al.* (2020a)).

Proof Sketch. Note that

$$P^e(r_t) = \sum_{\tau^o \in \mathcal{T}_t^o} \Pi_e^e(\tau^o) P^b(r_t, z_t, z_{t-1}, \dots, z_0 | a_0, \dots, a_t). \quad (2)$$

The analysis rests on non-parametric identification of the causal effect of a_i on the outcome (u_{i+1}, z_i) given confounded proxies z_{i-1} and z_i at each time step. This is done by first noting that

$$P(r_t, z_t, z_{t-1}, \dots, z_0 | a_0, \dots, a_t) = P^b(r_t, z_t | a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i | a_i, U_i) \right) P^b(U_0), \quad (3)$$

and extending off of the static causal identification setting, to show that

$$P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_{i-1}, U_{i-1}) = P^b(U_{i+1}, z_i | a_i, Z_{i-1}) P^b(Z_i | a_i, Z_{i-1})^{-1} \cdot P^b(Z_i, z_{i-1} | a_{i-1}, Z_{i-2}) P^b(Z_{i-1} | a_{i-1}, Z_{i-2})^{-1} \cdot P^b(Z_{i-1} | a_{i-1}, U_{i-1}) \quad (4)$$

and

$$P^b(r_t, z_t | a_t, U_t) P^b(U_t, z_{t-1} | a_{t-1}, U_{t-1}) = P^b(r_t, z_t | a_t, Z_{t-1}) P^b(Z_t | a_t, Z_{t-1})^{-1} \cdot P^b(Z_t, z_{t-1} | a_{t-1}, Z_{t-2}) P^b(Z_{t-1} | a_{t-1}, Z_{t-2})^{-1} \cdot P^b(Z_{t-1} | a_{t-1}, U_{t-1}) \quad (5)$$

while also observing that

$$P^b(Z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_{i-1}, Z_{i-2}) = P^b(Z_i, z_{i-1} | a_{i-1}, Z_{i-2}). \quad (6)$$

The result then follows from these four core identities and induction. \square

In the language of causal inference, the proof is made possible by, at each time step i , statically viewing z_{i-1} as an observable proxy emitted by the latent state u_i , giving the causal diagram in Figure 1(a) illustrating that the proxies z_i and z_{i-1} are conditionally independent given u_i , allowing for the application of the identification scheme of Miao *et al.* (2018).




Figure 1: Causal diagrams for proxies of the confounder u_i : one-step proxies (a), multi-step histories (b), multi-step histories, multi-step futures (c), and multi-step futures with reward-observation outcome (d).

4 Relaxing Rank Assumptions

The proof of Theorem 1 requires that $|\mathcal{U}| = |\mathcal{Z}|$ which is unrealistic in most real-world settings, where the hidden state space is often much smaller than the rich observation space. Inspired by spectral learning of PSRs and POMDPs (Boots *et al.*, 2011; Kulesza *et al.*, 2015; Hsu *et al.*, 2012), we show that we can allow for the more general assumption that $|\mathcal{U}| \leq |\mathcal{Z}|$ by requiring that $\text{rank}(P^b(Z_i|a_i, Z_{i-1})) = |\mathcal{U}|$. We then further relax rank assumptions by using the entire observable history as a proxy, rather than just the observation from the previous time step. The results in the following two sections hold using matrices of conditional probabilities, however we state them using joint probabilities as doing so is more general by allowing for probability zero events which would otherwise be conditioned on. Additionally, all matrix rank assumptions (2–4) are stated with equality rather than inequality (i.e., rank equal to $|\mathcal{U}|$ rather than at least $|\mathcal{U}|$) as all such matrices have rank upper bounded by $|\mathcal{U}|$; assuming rank $|\mathcal{U}|$ should be thought of as a “full rank” assumption.

4.1 Spectral Relaxation of One-Step Proxies

Assumption 2. $\text{rank}(P^b(Z_i, a_i, Z_{i-1})) = |\mathcal{U}|, \forall i \leq H, a_i \in \mathcal{A}$.

In contrast to Theorem 1 which requires $|\mathcal{Z}| = |\mathcal{U}|$ and $P^b(Z_i|a_i, Z_{i-1})$ to be invertible, here we only require $|\mathcal{Z}| \geq |\mathcal{U}|$ and the matrix to have the maximal possible rank $|\mathcal{U}|$. Moreover, we place the maximum rank condition on the joint probability matrix $P^b(Z_i, a_i, Z_{i-1})$ instead of its conditional version in Assumption 1, which is strictly more general: Assumption 2 automatically implies that $\text{rank}(P^b(Z_i|a_i, Z_{i-1})) = |\mathcal{U}|$ as $P^b(Z_i, a_i, Z_{i-1}) = P^b(Z_i|a_i, Z_{i-1})\text{diag}(P^b(Z_{i-1}, a_i))$ (for matrices, A, B , $\text{rank}(AB) \leq \min(\text{rank}(A), \text{rank}(B))$).

With this Assumption, we are able to derive the following estimator of $P^e(r_t)$.

Theorem 2. *Under Assumption 2, there exists matrices $M_{i,a_i} \in \mathbb{R}^{|\mathcal{U}| \times |\mathcal{Z}|}$ and $M'_{i,a_i} := (M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}))^+$*

such that $P^e(r_t)$ is equal to

$$\sum_{\tau^o \in \mathcal{T}_t^o} \Pi_e(\tau^o) P^b(r_t, z_t, a_t, Z_{t-1}) M'_{t,a_t} \cdot \prod_{i=t-1}^0 (M_{i+1,a_{i+1}} P^b(Z_{i+1}, z_i, a_i, Z_{i-1}) M'_{i,a_i}) M_{0,a_0} P^b(Z_0) \quad (7)$$

Remark 1 (Choice of M_{i,a_i}). *The matrices M_{i,a_i} must be chosen so as to satisfy the non-degenerate conditions that $\text{rank}(M_{i,a_i}) = |\mathcal{U}|$ and $\text{rank}(M_{i,a_i} P^b(Z_i, a_i, Z_{i-1})) = |\mathcal{U}|$. One such choice is that M_{i,a_i} is the top $|\mathcal{U}|$ left singular vectors of $P^b(Z_i, a_i, Z_{i-1})$. Alternatively, such a non-degenerate M_{i,a_i} may be drawn from any continuous probability distribution over matrices of the appropriate dimension, almost surely.*

The proof of Theorem 2 relies on deriving the following analogs to equations (4) – (5) and using equations (3) and (6) to similarly proceed by induction (see Appendix A.1 for proof).

Lemma 1. *Given Assumption 2, define M_{i,a_i} and M'_{i,a_i} as above. Then*

$$P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_{i-1}, U_{i-1}) = P^b(U_{i+1}, z_i, a_i, Z_{i-1}) M'_{i,a_i} M_{i,a_i} P^b(Z_i, z_{i-1}, a_{i-1}, Z_{i-2}) \cdot M'_{i-1,a_{i-1}} M_{i-1,a_{i-1}} P^b(Z_{i-1} | a_{i-1}, U_{i-1}) \quad (8)$$

and

$$P^b(r_t, z_t | a_t, U_t) P^b(U_t, z_{t-1} | a_{t-1}, U_{t-1}) = P^b(r_t, z_t, a_t, Z_{t-1}) M'_{t,a_t} M_{t,a_t} P^b(Z_t, z_{t-1}, a_{t-1}, Z_{t-2}) \cdot M'_{t-1,a_{t-1}} M_{t-1,a_{t-1}} P^b(Z_{t-1} | a_{t-1}, U_{t-1}) \quad (9)$$

4.2 Multi-Step Histories

We now show that, by extending histories, even weaker rank assumptions can be made. Specifically, letting \mathcal{H}_t^o denote the set of observable histories of length t , the matrices, $P^b(Z_i, a_i, Z_{i-1})$ and $P^b(Z_{i+1}, z_i, a_i, Z_{i-1})$, in Lemma 1 and Theorem 2 can be replaced with $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)$ and $P^b(Z_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o)$, respectively. We make the following rank assumption:

Assumption 3. $\text{rank}(P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)) = |\mathcal{U}|, \forall i \leq H, a_i \in \mathcal{A}$.

This assumption is strictly weaker than our previous Assumption 2. This is because $P^b(Z_i, a_i, Z_{i-1})$ can be written as $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) J_i$, where $J_i \in \mathbb{R}^{|\mathcal{H}_{i-1}^o| \times |\mathcal{Z}_{i-1}|}$ marginalizes out all parts of the history prior to timestep $i-1$ for each z_{i-1} . Since $P^b(Z_i, a_i, Z_{i-1})$ is a projection of $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)$, the rank of the latter cannot be smaller than that of the former. We then obtain the following:

Theorem 3. *Under Assumption 3, there exists matrices M_{i,a_i} and $M'_{i,a_i} := (M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o))^+$, such that the probabil-*

ity $P^e(r_t)$ is equal to

$$\sum_{\tau^o \in \mathcal{T}_i^o} \Pi_e(\tau^o) P^b(r_t, z_t, a_t, \mathcal{H}_{i-1}^o) M'_{t, a_t} \cdot \prod_{i=t-1}^0 (M_{i+1, a_{i+1}} P^b(Z_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i}) M_{0, a_0} P^b(Z_0) \quad (10)$$

Remark 2 (Choice of M_{i, a_i}). *Similar to Remark 1, the matrices M_{i, a_i} must be chosen so that $\text{rank}(M_{i, a_i}) = \text{rank}(M_{i, a_i} P(Z_i, a_i, \mathcal{H}_{i-1}^o)) = |\mathcal{U}|$. Again, two viable choices are for M_{i, a_i} to be the top $|\mathcal{U}|$ left singular vectors of $P(Z_i, a_i, \mathcal{H}_{i-1}^o)$ or to draw M_{i, a_i} from a continuous probability distribution. Furthermore, M_{i, a_i} can be chosen to first marginalize out multi-step histories \mathcal{H}_{i-1}^o into one-step pasts \mathcal{Z}_{i-1} , and then act as a rank $|\mathcal{U}|$ projection matrix, as in Theorem 2. This choice of M_{i, a_i} illustrates that Theorem 3 is a strict generalization of Theorem 2 as the latter is recoverable with this alternate choice of M_{i, a_i} , which first performs marginalization.*

We give a proof of Theorem 3 in Appendix A.2, but, at a high level, Figures 1(a) and (b) show the static causal structure is unchanged when extending the length of observable histories, allowing for essentially the same analysis as in the one-step case. We also remark that Theorem 3 holds using conditional probability matrices $P^b(Z_i | a_i, \mathcal{H}_{i-1}^o)$ and $P^b(Z_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$ in place of $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)$ and $P^b(Z_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o)$, respectively. An analogous statement applies to Theorem 2.

5 Multi-Step Futures

In the previous section we relaxed the rank assumptions in prior work by replacing the one-step observation with *histories* of observations as proxies. It is then very natural to ask: can we make a similar extension to the future? Indeed, in the PSR literature that inspire our work, it is common to use moment matrices about multi-step histories and futures (Boots *et al.*, 2011) to relax the rank assumptions of earlier works that estimate one-step moment matrices (Hsu *et al.*, 2012).

To this end, our goal in this section is to extend futures and replace the matrices $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)$ and $P^b(Z_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o)$ with matrices of the form $P^b(\mathcal{F}_i^o, a_i, \mathcal{H}_{i-1}^o)$ and $P^b(\mathcal{F}_{i+1}^o, z_i, a_i, \mathcal{H}_{i-1}^o)$, respectively (\mathcal{F}_i^o denotes a set of futures at timestep i , containing subtrajectories f_i of the form $(z_i, z_{i+1}, a_{i+1}, \dots, z_H, a_H)$), so as to further relax rank assumptions by only assuming that $\text{rank}(P^b(\mathcal{F}_i^o, a_i, \mathcal{H}_{i-1}^o)) = |\mathcal{U}|$. Such an assumption would imply Assumption 3 and thus be even weaker than that made in the previous section.

Unfortunately, such a goal is not immediately possible to achieve via identification of $P^b(r_t, z_t, z_{t-1}, \dots, z_0 | a_0, \dots, a_t)$ using previous techniques. The primary difficulty, as indicated by the red edge in Figure 1(c), is that, when extending futures, the treatment, a_i , *does* have a causal effect on the future f_i , rendering it a post-treatment variable rather than a negative control as was

the case with z_i , in Figures 1(a) and (b), on which a_i has no causal effect.

While our approach cannot use extended futures as a proxy for the confounder under rank $|\mathcal{U}|$ assumptions on $P^b(\mathcal{F}_i^o, a_i, \mathcal{H}_{i-1}^o)$ alone, we show that it is in fact possible to perform OPE by working with conditional moment matrices $P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o)$ and making relatively weak positivity and distinctness assumptions about the reward distributions:

Assumption 4. $\text{rank}(P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o)) = |\mathcal{U}|, \forall i \leq H, a_i \in \mathcal{A}$.

Assumption 5. *For each $a_i \in \mathcal{A}$, there exists $r_i \in \mathcal{R}$ such that the probabilities $P^b(r_i | a_i, u_i), u_i \in \mathcal{U}$ are all distinct.*

Intuitively, Assumption 5 requires that the reward at each timestep—another post-treatment variable—contains enough information about hidden state so as to enable its use, in tandem with futures, to perform causal effect identification (see Appendix A.3 for proof):

Theorem 4. *Under Assumptions 4–5, there exist M_{i, a_i} and M'_{i, a_i} (chosen to be non-degenerate w.r.t. $P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o)$, analogously to Remarks 1–2), such that $P^e(r_t)$ equals*

$$\sum_{\tau^o \in \mathcal{T}_i^o} \Pi_e(\tau^o) P^b(r_t, z_t | a_t, \mathcal{H}_{i-1}^o) M'_{t, a_t} \cdot \prod_{i=t-1}^0 (M_{i+1, a_{i+1}} P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1}) M'_{i, a_i}) M_{0, a_0} P^b(Z_0), \quad (11)$$

where the matrices $P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1})$ are identifiable.

The proof of equation (11) in Theorem 4 is similar to those of Theorems 2 and 3. The key difference with the previous approaches, however, is the identification of the matrices $P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1})$, which relies on the diagonalization strategy of Kuroki and Pearl (2014). Essentially, by diagonalizing products of observable moment matrices, we are able to identify the unobserved matrices $P^b(\mathcal{F}_{i+1}^o | a_{i+1}, U_{i+1})$ and $P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$ which allows for identification of

$$P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1}) = P^b(\mathcal{F}_{i+1}^o | a_{i+1}, U_{i+1}) P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$$

6 Experiments

We now empirically show that our methods extend the range of environments in which we can successfully perform accurate OPE. In particular, we show that: our history-extension approach of Theorem 3 is more accurate than that of Theorem 1, the baseline estimator of Tennenholtz *et al.* (2020a), even in environments satisfying Assumption 1; using our extended-history estimator significantly outperforms the baseline in environments under which Assumption 1 does not hold; and that while Theorem 3’s estimator *still* performs well on environments in which Assumption 3 doesn’t hold, the extended-future estimator from Theorem 4 can produce even more accurate results.




Figure 2: Average absolute residuals $|\hat{P}^e(r_t) - P^e(r_t)|$ over randomly generated POMDP environments which satisfy Assumption 1.




Figure 3: Estimates of $P^e(r_t)$ in POMDP environment which violates Assumption 1.

Setup For each of our experiments, we conduct 50 independent trials in POMDPs with $|\mathcal{U}| = |\mathcal{Z}| = |\mathcal{A}| = 2$, $\mathcal{R} = \{0, 1\}$, and $H = 4$. For the first experiment, comparing the estimators of Theorems 1 and 3, the probability parameters of the POMDP and policies are drawn uniformly at random from $[0, 1]$ each trial, whereas in the two latter experiments, the parameters are chosen specifically, so as to violate Assumptions 1 and 3, and remain constant across trials. We note however, the extended-future estimator of Theorem 4 can be quite sensitive to ill-conditioned matrices, and so the environment for the third experiment was chosen somewhat carefully, through adaptive search, so as to yield an environment with well-conditioned matrices. For the first experiment, during each trial for estimator E , we compute, at each timestep, the residual $|\hat{P}^e(r_t = 1) - P^e(r_t = 1)|$, where $\hat{P}^e(r_t = 1)$ is E 's estimate of $P^e(r_t = 1)$ after seeing n trajectories for various values of n . For the latter experiments, the results of a given trial for estimator E consist of E 's probability estimates of $P^e(r_0 = 1), \dots, P^e(r_H = 1)$ after seeing n trajectories, for various values of n . The final results for each experiment are then averaged over the 50 trials and presented with standard error bars.




Figure 4: Estimates of $P^e(r_t)$ in POMDP environment which violates Assumption 3.

Results Figure 2 shows that even in environments in which Assumption 1 *does* hold, the quality of the baseline estimator of Theorem 1 degrades substantially, on average, as t grows for both 5000 and 10,000 trajectories. On the other hand, our history-extended estimator of Theorem 3 maintains accurate estimates of $P^e(r_t = 1)$ even for larger values of t , and is in general, much more accurate than the baseline at all timesteps.

Figure 3 extends these results to the second experiment in which we compare our estimator from Theorem 3 to the baseline of Theorem 1 when its Assumption 1 is violated. At 5000 trajectories, both estimators incur a noticeable degree of error at multiple timesteps, although the errors of the baseline estimator are significantly worse. However, at 6000 samples, the extended-history estimator of Theorem 3 incurs negligible error in its estimates at *all* timesteps, especially as compared to the baseline which still performs extremely poorly, often producing estimates far outside of $[0, 1]$.

Lastly, the robustness of the extended-history estimator of Theorem 3 to an environment in which its Assumption 3 is violated is illustrated in Figure 4. While Theorem 3's estimator is somewhat biased for both 500,000 and 1,000,000 sampled trajectories, the estimates are, in absolute terms, relatively accurate, especially when compared to the errors incurred by the baseline estimator of Theorem 1 when its key Assumption 1 was violated in the second experiment. In Appendix B, we further show that, in fact, Theorem 3's estimator is still highly accurate on average across environments that violate its Assumption 3. Figure 4 also shows that the extended-future estimator of Theorem 4—which does not require Assumption 3—produces more accurate results.

7 Importance Sampling

Importance Sampling (Precup *et al.*, 2000) is another family of standard OPE methods, which complements the so-called “direct methods” that either estimate a model or approximate the Bellman equations (Dudík *et al.*, 2011; Jiang and Li, 2016). (Our approach in previous sections can be roughly categorized as a direct method.) Tennenholtz *et al.* (2020a) illustrate the difficulty of IS in POMDPs by showing that its immediate extension to POMDPs is neither unbiased nor con-

sistent unless strong sufficiency assumptions are made. By extending the eigendecomposition technique of Kuroki and Pearl (2014), we show, that OPE in POMDPs is amenable to a consistent IS estimator under much weaker assumptions involving only rank $|\mathcal{U}|$, distinctness, and positivity conditions.

In order to satisfy certain rank assumptions of our analysis we require the following assumptions on the relative sizes of hidden state, action and reward spaces:

Assumption 6. $|\mathcal{A}| \geq |\mathcal{U}|$ and $|\mathcal{R}| \geq |\mathcal{U}|$

Under rank $|\mathcal{U}|$ conditions on certain moment matrices, we are then able to identify the behavior policy probabilities $\pi_b(a_i|U_i)$ via eigendecompositions. We, additionally, make a weak assumption on the reward distribution so that the causal structure of rewards and observations are identical:

Assumption 7. *The reward function R depends only latent state and not on action.*

Assumption 7 allows for the reward distribution $P^b(r_i|U_i)$ to also be written in terms of observable probabilities similar to $\pi_b^{(i)}(a_i|U_i)$, under analogous rank $|\mathcal{U}|$ assumptions. Additionally, in order to ensure uniqueness in the eigendecompositions, we further require certain sets of probabilities to be distinct. As these conditions are in terms of probability matrices which lack succinct representation, we relegate the assumptions and proof to Appendix C.1 but summarize here:

Theorem 5. *Under Assumptions 6–7, rank $|\mathcal{U}|$ and distinctness conditions, the vectors $\pi_b^{(i)}(a_i|U_i)$, and $P^b(r_i|U_i)$ are identifiable under the behavior measure w.r.t. an unknown, but consistent, ordering on U_i for all $i \leq H$.*

Defining $\Pi_b(u_{0:t}) = \prod_{i=0}^t \pi_b^{(i)}(a_i|u_i)$ and letting v denote value, we then obtain an IS procedure under the two following additional rank and positivity assumptions, as well as positivity and distinctness assumptions (see Appendix C):

Assumption 8. $\text{rank}(P^b(R_i|U_i)) = |\mathcal{U}|$

Assumption 9. $P^b(v|\tau^o) > 0, \forall v, \tau^o$ and $\pi_b^{(i)}(a_i|u_i) > 0, \forall i \leq H, a_i \in \mathcal{H}, u_i \in \mathcal{U}$.

Theorem 6. *Under Assumptions 6–9, as well as additional rank, distinctness, and positivity conditions, we have that $v_H(\pi_e) = \mathbb{E}[vW_{e,b}(v, \tau^o)]$ with*

$$W_{e,b}(v, \tau^o) := \frac{1}{P^b(v|\tau^o)} \Pi_e(\tau^o) \Gamma_b(v, \tau^o)$$

and

$$\Gamma_b(v, \tau^o) = \sum_{r_{0:H}: \sum r_i = v} \sum_{u_{0:H}} \frac{\prod_{i=0}^H P^b(r_i|u_i) P^b(u_{0:H}|\tau^o)}{\Pi_b(u_{0:H})},$$

where $W_{e,b}(v, \tau^o)$ are identifiable in terms of observable probabilities.

We give a sketch of the proof (for details, see Appendix C).

Proof sketch. We have that

$$\begin{aligned} v_H(\pi_e) &= \sum_{\tau \in \mathcal{T}_H} \sum_v v P^e(v|\tau) P^e(\tau) \\ &= \sum_{\tau \in \mathcal{T}_H} \sum_v v P^b(v|u_{0:H}) P^b(\tau) \left(\frac{\Pi_e(\tau^o)}{\Pi_b(u_{0:H})} \right) \\ &= \sum_v v \sum_{\tau^o \in \mathcal{T}_t^o} P^b(\tau^o) \Pi_e(\tau^o) \sum_{u_{0:H}} \frac{P^b(v|u_{0:H}) P^b(u_{0:H}|\tau^o)}{\Pi_b(u_{0:H})} \\ &= \sum_v v P^b(v|\tau^o) \sum_{\tau^o \in \mathcal{T}_t^o} W_{e,b}(v, \tau^o) P^b(\tau^o) = \mathbb{E}[vW_{e,b}(v, \tau^o)]. \end{aligned}$$

Identification of $W_{e,b}(v, \tau^o)$ is made possible by Theorem 5 and writing $P^b(u_{0:H}|\tau^o)$ in terms observable and identifiable probabilities via eigenvalue analysis and pseudoinversion. \square

We also conducted experiments comparing our IS estimator to the baseline (biased) IS estimator of Tennenholtz *et al.* (2020a) and found that, in environments which meet the required conditions of Theorem 5 and with sufficiently well-conditioned moment matrices that our estimator must pseudoinvert, it outperforms the baseline for horizon comprising two timesteps. However, for longer horizons our IS estimator often produces less accurate value estimates than the baseline, indicating that, while our IS estimator, in contrast to the baseline, is statistically consistent, the number of sampled trajectories required to guarantee good performance may often be quite high for longer horizons.

8 Conclusion and Future Work

In this paper we consider OPE in POMDPs. We extend prior work, relaxing assumptions on observable probability matrices to only have rank $|\mathcal{U}|$ rather than $|\mathcal{Z}|$, simultaneously generalizing to POMDPs in which $|\mathcal{Z}| \geq |\mathcal{U}|$. By additionally extending one of the proxy variables into the past, we further relax assumptions by assuming the matrices $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)$ have rank $|\mathcal{U}|$, allowing for our estimator to apply to POMDPs in which the previous observation may not contain enough information about state, but the entire observable history does. We show that futures, however, cannot be immediately extended using this strategy, and thus make relatively weak assumptions on the reward distribution so as to allow for identification using rank $|\mathcal{U}|$ moment matrices involving both histories and futures. We then experimentally compare our main estimators and show that they both outperform the baseline given in Tennenholtz *et al.* (2020a) and perform well in a variety of different environments. Finally, we give a consistent IS algorithm for OPE in POMDPs which only depends on rank, distinctness, and positivity conditions on certain probability matrices and not on sufficiency assumptions.

References

Andrew Bennett, Nathan Kallus, Lihong Li, and Ali Mousavi. Off-policy evaluation in infinite-horizon reinforcement learning with latent confounders, 2020.

- Byron Boots, Sajid M Siddiqi, and Geoffrey J Gordon. Closing the learning-planning loop with predictive state representations. *The International Journal of Robotics Research*, 30(7):954–966, 2011.
- Miroslav Dudík, John Langford, and Lihong Li. Doubly robust policy evaluation and learning. In *Proceedings of the 28th International Conference on International Conference on Machine Learning*, pages 1097–1104, 2011.
- Sander Greenland and Timothy Lash. Bias Analysis. *Modern Epidemiology*, 3:423–437, 2012.
- Daniel Hsu, Sham M. Kakade, and Tong Zhang. A spectral algorithm for learning hidden markov models. *Journal of Computer and System Sciences*, 78(5):1460 – 1480, 2012. JCSS Special Issue: Cloud Computing 2011.
- Nan Jiang and Lihong Li. Doubly robust off-policy value evaluation for reinforcement learning. In Maria Florina Balcan and Kilian Q. Weinberger, editors, *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 652–661, New York, New York, USA, 20–22 Jun 2016. PMLR.
- Alex Kulesza, Nan Jiang, and Satinder Singh. Spectral learning of predictive state representations with insufficient statistics. In *Proceedings of the Twenty-Ninth AAAI Conference on Artificial Intelligence*, AAAI’15, page 2715–2721. AAAI Press, 2015.
- Manabu Kuroki and Judea Pearl. Measurement bias and effect restoration in causal inference. *Biometrika*, 101(2):423–437, 03 2014.
- Travis Mandel, Yun-En Liu, Emma Brunskill, and Zoran Popović. Offline evaluation of online reinforcement learning algorithms. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 30, 2016.
- Wang Miao, Zhi Geng, and Eric J Tchetgen Tchetgen. Identifying causal effects with proxy variables of an unmeasured confounder. *Biometrika*, 105(4):987–993, 2018.
- Michael Oberst and David Sontag. Counterfactual off-policy evaluation with Gumbel-max structural causal models. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 4881–4890, Long Beach, California, USA, 09–15 Jun 2019. PMLR.
- Judea Pearl. *Causality: Models, Reasoning and Inference*. Cambridge University Press, USA, 2nd edition, 2009.
- Doina Precup, Richard S. Sutton, and Satinder P. Singh. Eligibility traces for off-policy policy evaluation. In *Proceedings of the Seventeenth International Conference on Machine Learning*, ICML ’00, page 759–766, San Francisco, CA, USA, 2000. Morgan Kaufmann Publishers Inc.
- G. W. Stewart. *Matrix Algorithms*. Society for Industrial and Applied Mathematics, 2001.
- Richard S Sutton and Andrew G Barto. *Reinforcement learning: An introduction*. MIT press, 2018.
- Guy Tennenholtz, Uri Shalit, and Shie Mannor. Off-policy evaluation in partially observable environments. *Proceedings of the AAAI Conference on Artificial Intelligence*, 34(06):10276–10283, Apr. 2020.
- Guy Tennenholtz, Uri Shalit, and Shie Mannor. Off-policy evaluation in partially observable environments. *Proceedings of the AAAI Conference on Artificial Intelligence*, 34(06):10276–10283, Apr. 2020.
- Philip Thomas and Emma Brunskill. Data-efficient off-policy policy evaluation for reinforcement learning. In Maria Florina Balcan and Kilian Q. Weinberger, editors, *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 2139–2148, New York, New York, USA, 20–22 Jun 2016. PMLR.
- J. Zhang and Elias Bareinboim. Markov decision processes with unobserved confounders : A causal approach. 2016.

A OPE via Observable Proxies

We extend the identification scheme of Tennenholtz *et al.* (2020a) and Miao *et al.* (2018) to make weaker rank assumptions. The proofs in this section use the same general strategy as in Tennenholtz *et al.* (2020a), but use spectral methods to relax their assumptions.

A.1 One-Step Proxies

Proof of Lemma 1

Proof. First, notice that

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, a_i, Z_{i-1}) = P^b(U_{i+1}, z_i, a_i, Z_{i-1}) \\ \implies & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, a_i, Z_{i-1}) M'_{i,a_i} = P^b(U_{i+1}, z_i, a_i, Z_{i-1}) M'_{i,a_i}. \end{aligned} \quad (12)$$

Additionally,

$$P^b(Z_i, a_i, Z_{i-1}) = P^b(Z_i | a_i, U_i) P^b(U_i, a_i, Z_{i-1}) \quad (13)$$

$$\implies P^b(Z_i, a_i, Z_{i-1}) M'_{i,a_i} = P^b(Z_i | a_i, U_i) P^b(U_i, a_i, Z_{i-1}) M'_{i,a_i}. \quad (14)$$

From the definition of M'_{i,a_i} , we have that

$$\begin{aligned} & P^b(Z_i, a_i, Z_{i-1}) M'_{i,a_i} \\ &= P^b(Z_i, a_i, Z_{i-1}) (M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}))^+ = P^b(Z_i, a_i, Z_{i-1}) V_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+ \\ &= U_{i,a_i} \Sigma_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+, \end{aligned}$$

which has rank $|\mathcal{U}|$, thus implying, by equation (14), that $\text{rank}(P^b(U_i, a_i, Z_{i-1}) M'_{i,a_i}) = |\mathcal{U}|$, and so equation (12) implies that

$$P^b(U_{i+1}, z_i | a_i, U_i) = P^b(U_{i+1}, z_i, a_i, Z_{i-1}) M'_{i,a_i} (P^b(U_i, a_i, Z_{i-1}) M'_{i,a_i})^{-1}. \quad (15)$$

Now, notice that equation (13) implies that

$$M_{i,a_i} P^b(Z_i | a_i, U_i) P^b(U_i, a_i, Z_{i-1}) = M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}). \quad (16)$$

Similarly notice that $\text{rank}(M_{i,a_i} P^b(Z_i, a_i, Z_{i-1})) = |\mathcal{U}|$ as, from the definition of M_{i,a_i} , we have $M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}) = I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i} V_{i,a_i}^T$. This, in turn, implies that $\text{rank}(M_{i,a_i} P^b(Z_i | a_i, U_i)) = |\mathcal{U}|$ by equation (16), and so equation (16) implies that

$$P^b(U_i, a_i, Z_{i-1}) = (M_{i,a_i} P^b(Z_i | a_i, U_i))^{-1} M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}),$$

implying that

$$P^b(U_i, a_i, Z_{i-1}) M'_{i,a_i} = (M_{i,a_i} P^b(Z_i | a_i, U_i))^{-1} M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}) M'_{i,a_i}. \quad (17)$$

Now, we claim that $M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}) M'_{i,a_i} = I_{|\mathcal{U}| \times |\mathcal{U}|}$ since

$$\begin{aligned} M_{i,a_i} P^b(Z_i, a_i, Z_{i-1}) M'_{i,a_i} &= I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T (I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T)^+ \\ &= I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+ \end{aligned}$$

and thus equation (17) becomes

$$P^b(U_i, a_i, Z_{i-1}) M'_{i,a_i} = (M_{i,a_i} P^b(Z_i | a_i, U_i))^{-1}. \quad (18)$$

Finally, substituting (18) into (15) gives

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) \\ &= P^b(U_{i+1}, z_i, a_i, Z_{i-1}) M'_{i,a_i} M_{i,a_i} P^b(Z_i | a_i, U_i). \end{aligned}$$

This, in combination with the fact that

$$P^b(Z_i | a_i, U_i) P^b(U_i, z_{i-1}, a_{i-1}, Z_{i-2}) = P^b(Z_i, z_{i-1}, a_{i-1}, Z_{i-2}) \quad (19)$$

immediately implies that

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_i, U_i) = P^b(U_{i+1}, z_i, a_i, Z_{i-1}) M'_{i,a_i} \\ & \cdot M_{i,a_i} P^b(Z_i, z_{i-1}, a_{i-1}, Z_{i-2}) M'_{a_{i-1}} M_{a_{i-1}} P^b(Z_{i-1} | a_{i-1}, U_{i-1}), \end{aligned}$$

as desired. Replacing U_{i+1} with r_i gives the second part of the Lemma. \square

The proof of Theorem 2 then follows first from writing $P(r_t, z_t, \dots, z_0 | a_0, \dots, a_t)$ as

$$P^b(r_t, z_t | a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i | a_i, U_i) \right) P^b(U_0),$$

using Lemma 1 and equation (19) above to inductively show that

$$\begin{aligned} & P^b(r_t, z_t | a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i | a_i, U_i) \right) P^b(U_0) \\ &= P^b(r_t, z_t, a_t, Z_{t-1}) M'_{t, a_t} \prod_{i=t-1}^0 (M_{i+1, a_{i+1}} P^b(Z_{i+1}, z_i, a_i, Z_{i-1}) M'_{i, a_i}) M_{0, a_0} P^b(Z_0), \end{aligned} \quad (20)$$

and then using the fact that

$$P^e(r_t) = \sum_{\tau^o \in \mathcal{T}_t^o} \Pi_e(\tau^o) P(r_t, z_t, \dots, z_0 | a_0, \dots, a_t)$$

to obtain the final result.

A.2 Multi-Step Histories

Proof of Theorem 3

The proof of Theorem 3 is essentially the same as that of Theorem 2, except that we replace the one-step history Z_{i-1} with \mathcal{H}_{i-1}^o . For completeness, we include it below, first proving a Lemma analogous to Lemma 1:

Lemma 2. *Given Assumption 3, let $U_{i, a_i} \Sigma_{i, a_i} V_{i, a_i}^T$ be an SVD of $P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)$ with the diagonal entries of Σ_{i, a_i} in descending order and define $M_{i, a_i} := I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i, a_i}^T$ and $M'_{i, a_i} := (M_{a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o))^+$. Then*

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_{i-1}, U_{i-1}) \\ &= P^b(U_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i} M_{i, a_i} P^b(Z_i, z_{i-1}, a_{i-1}, \mathcal{H}_{i-2}^o) \\ &\quad \cdot M'_{i-1, a_{i-1}} M_{i-1, a_{i-1}} P^b(Z_{i-1} | a_{i-1}, U_{i-1}) \end{aligned}$$

and

$$\begin{aligned} & P^b(r_t, z_t | a_t, U_t) P^b(U_t, z_{t-1} | a_{t-1}, U_{t-1}) \\ &= P^b(r_t, z_t, a_t, \mathcal{H}_{i-1}^o) M'_{t, a_t} M_{t, a_t} P^b(Z_t, z_{t-1}, a_{t-1}, \mathcal{H}_{t-2}^o) \\ &\quad \cdot M'_{t-1, a_{t-1}} M_{t-1, a_{t-1}} P^b(Z_{t-1} | a_{t-1}, U_{t-1}) \end{aligned}$$

Proof. First, notice that

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, a_i, \mathcal{H}_{i-1}^o) = P^b(U_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) \\ \implies & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i} = P^b(U_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i}. \end{aligned} \quad (21)$$

Additionally,

$$P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) = P^b(Z_i | a_i, U_i) P^b(U_i, a_i, \mathcal{H}_{i-1}^o) \quad (22)$$

$$\implies P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i} = P^b(Z_i | a_i, U_i) P^b(U_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i}. \quad (23)$$

From the definition of M'_{i, a_i} , we have that

$$\begin{aligned} & P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i} \\ &= P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) (M_{i, a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o))^+ = P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) V_{i, a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i, a_i})^+ \\ &= U_{i, a_i} \Sigma_{i, a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i, a_i})^+ = U_{i, a_i} I_{|\mathcal{Z}| \times |\mathcal{U}|} = U_{i, a_i} I_{|\mathcal{Z}| \times |\mathcal{U}|}, \end{aligned}$$

which has rank $|\mathcal{U}|$, thus implying, by equation (23), that $\text{rank}(P^b(U_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i}) = |\mathcal{U}|$, and so equation (21) implies that

$$P^b(U_{i+1}, z_i | a_i, U_i) = P^b(U_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i} (P^b(U_i, a_i, \mathcal{H}_{i-1}^o) M'_{i, a_i})^{-1}. \quad (24)$$

Now, notice that equation (22) implies that

$$M_{i,a_i} P^b(Z_i|a_i, U_i) P^b(U_i, a_i, \mathcal{H}_{i-1}^o) = M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o). \quad (25)$$

Similarly notice that $\text{rank}(M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o)) = |U|$ as, from the definition of M_{i,a_i} , we have $M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) = I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i} V_{i,a_i}^T$. This, in turn, implies that $\text{rank}(M_{i,a_i} P^b(Z_i|a_i, U_i)) = |U|$ by equation (25), and so equation (25) implies that

$$P^b(U_i, a_i, \mathcal{H}_{i-1}^o) = (M_{i,a_i} P^b(Z_i|a_i, U_i))^{-1} M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o),$$

implying that

$$P^b(U_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = (M_{i,a_i} P^b(Z_i|a_i, U_i))^{-1} M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i}. \quad (26)$$

Now, we claim that $M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = I_{|\mathcal{U}| \times |\mathcal{U}|}$ since

$$\begin{aligned} M_{i,a_i} P^b(Z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} &= I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T (I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T)^+ \\ &= I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+ \end{aligned}$$

and thus equation (26) becomes

$$P^b(U_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = (M_{i,a_i} P^b(Z_i|a_i, U_i))^{-1}. \quad (27)$$

Finally, substituting (27) into (24) gives

$$\begin{aligned} &P^b(U_{i+1}, z_i|a_i, U_i) \\ &= P^b(U_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} M_{i,a_i} P^b(Z_i|a_i, U_i). \end{aligned}$$

This, in combination with the fact that

$$P^b(Z_i|a_i, U_i) P^b(U_i, z_{i-1}, a_{i-1}, \mathcal{H}_{i-2}^o) = P^b(Z_i, z_{i-1}, a_{i-1}, \mathcal{H}_{i-2}^o) \quad (28)$$

immediately implies that

$$\begin{aligned} &P^b(U_{i+1}, z_i|a_i, U_i) P^b(U_i, z_{i-1}|a_i, U_i) = P^b(U_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} \\ &\quad \cdot M_{i,a_i} P^b(Z_i, z_{i-1}, a_{i-1}, \mathcal{H}_{i-2}^o) M'_{a_{i-1}} M_{a_{i-1}} P^b(Z_{i-1}|a_{i-1} U_{i-1}), \end{aligned}$$

as desired. Replacing U_{i+1} with r_i and i with gives the second part of the Lemma. \square

Again, the proof of Theorem 3 follows first from writing $P(r_t, z_t, \dots, z_0|a_0, \dots, a_t)$ as

$$P^b(r_t, z_t|a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i|a_i, U_i) \right) P^b(U_0),$$

using Lemma 2 and equation (28) above to inductively show that

$$\begin{aligned} &P^b(r_t, z_t|a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i|a_i, U_i) \right) P^b(U_0) \\ &= P^b(r_t, z_t, a_t, \mathcal{H}_{t-1}^o) M'_{t,a_t} \prod_{i=t-1}^0 (M_{i+1,a_{i+1}} P^b(Z_{i+1}, z_i, a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i}) M_{0,a_0} P^b(Z_0), \quad (29) \end{aligned}$$

and again using the fact that

$$P^e(r_t) = \sum_{\tau^o \in \mathcal{T}_t^o} \Pi_e(\tau^o) P(r_t, z_t, \dots, z_0|a_0, \dots, a_t)$$

to obtain the final result.

A.3 Multi-Step Futures

Proof of equation (11) of Theorem 4

The proof of equation (11) of Theorem 4 is, again, essentially the same as those of Theorems 2 and 3, except that we replace the one-step history Z_{i-1} with \mathcal{H}_{i-1}^o and the one-step future Z_i with \mathcal{F}_i^o and here, we use moment matrices of conditional probabilities rather than joints. Again, for completeness, we include it below, first proving a Lemma analogous to Lemmas 1 and 2:

Lemma 3. *Given Assumption 4, let $U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T$ be an SVD of $P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o)$ with the diagonal entries of Σ_{i,a_i} in descending order and define $M_{i,a_i} := I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T$ and $M'_{i,a_i} := (M_{a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o))^+$. Then*

$$\begin{aligned} P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_{i-1}, U_{i-1}) \\ = P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} M_{i,a_i} P^b(\mathcal{F}_i^o, z_{i-1} | a_{i-1}, \mathcal{H}_{i-2}^o | a_i) \\ \cdot M'_{i-1,a_{i-1}} M_{i-1,a_{i-1}} P^b(\mathcal{F}_{i-1}^o | a_{i-1}, U_{i-1}) \end{aligned}$$

and

$$\begin{aligned} P^b(r_t, z_t | a_t, U_t) P^b(U_t, z_{t-1} | a_{t-1}, U_{t-1}) \\ = P^b(r_t, z_t | a_t, \mathcal{H}_{t-1}^o) M'_{t,a_t} M_{t,a_t} P^b(\mathcal{F}_t^o, z_{t-1} | a_{t-1}, \mathcal{H}_{t-2}^o | a_t) \\ \cdot M'_{t-1,a_{t-1}} M_{t-1,a_{t-1}} P^b(\mathcal{F}_{t-1}^o | a_{t-1}, U_{t-1}) \end{aligned}$$

Proof. First, notice that

$$\begin{aligned} P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i | a_i, \mathcal{H}_{i-1}^o) &= P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) \\ \implies P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} &= P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i}. \end{aligned} \quad (30)$$

Additionally,

$$P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) = P^b(\mathcal{F}_i^o | a_i, U_i) P^b(U_i | a_i, \mathcal{H}_{i-1}^o) \quad (31)$$

$$\implies P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = P^b(\mathcal{F}_i^o | a_i, U_i) P^b(U_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i}. \quad (32)$$

From the definition of M'_{i,a_i} , we have that

$$\begin{aligned} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} \\ = P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) (M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o))^+ = P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) V_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+ \\ = U_{i,a_i} \Sigma_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+ = U_{i,a_i} I_{\mathcal{Z} \times |\mathcal{U}|} = U_{i,a_i} I_{\mathcal{Z} \times |\mathcal{U}|}, \end{aligned}$$

which has rank $|\mathcal{U}|$, thus implying, by equation (32), that $\text{rank}(P^b(U_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i}) = |\mathcal{U}|$, and so equation (30) implies that

$$P^b(U_{i+1}, z_i | a_i, U_i) = P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} (P^b(U_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i})^{-1}. \quad (33)$$

Now, notice that equation (31) implies that

$$M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, U_i) P^b(U_i | a_i, \mathcal{H}_{i-1}^o) = M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o). \quad (34)$$

Similarly notice that $\text{rank}(M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o)) = |\mathcal{U}|$ as, from the definition of M_{i,a_i} , we have $M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) = I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i} V_{i,a_i}^T$. This, in turn, implies that $\text{rank}(M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, U_i)) = |\mathcal{U}|$ by equation (34), and so equation (34) implies that

$$P^b(U_i | a_i, \mathcal{H}_{i-1}^o) = (M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, U_i))^{-1} M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o),$$

implying that

$$P^b(U_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = (M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, U_i))^{-1} M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i}. \quad (35)$$

Now, we claim that $M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = I_{|\mathcal{U}| \times |\mathcal{U}|}$ since

$$\begin{aligned} M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} &= I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T (I_{|\mathcal{U}| \times |\mathcal{Z}|} U_{i,a_i}^T U_{i,a_i} \Sigma_{i,a_i} V_{i,a_i}^T)^+ \\ &= I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i} (I_{|\mathcal{U}| \times |\mathcal{Z}|} \Sigma_{i,a_i})^+ \end{aligned}$$

and thus equation (35) becomes

$$P^b(U_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} = (M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, U_i))^{-1}. \quad (36)$$

Finally, substituting (36) into (33) gives

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) \\ &= P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} M_{i,a_i} P^b(\mathcal{F}_i^o | a_i, U_i). \end{aligned}$$

This, in combination with the fact that

$$P^b(\mathcal{F}_i^o | a_i, U_i) P^b(U_i, z_{i-1} | a_{i-1}, \mathcal{H}_{i-2}^o) = P^b(\mathcal{F}_i^o, z_{i-1} | a_{i-1}, \mathcal{H}_{i-2}^o | a_i) \quad (37)$$

immediately implies that

$$\begin{aligned} & P^b(U_{i+1}, z_i | a_i, U_i) P^b(U_i, z_{i-1} | a_i, U_i) = P(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) M'_{i,a_i} \\ & \cdot M_{i,a_i} P^b(\mathcal{F}_i^o, z_{i-1} | a_{i-1}, \mathcal{H}_{i-2}^o | a_i) M'_{a_{i-1}} M_{a_{i-1}} P^b(Z_{i-1} | a_{i-1} U_{i-1}), \end{aligned}$$

as desired. Replacing U_{i+1} with r_i and i with gives the second part of the Lemma. \square

Again, the proof of Theorem 4 follows first from writing $P(r_t, z_t, \dots, z_0 | a_0, \dots, a_t)$ as

$$P^b(r_t, z_t | a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i | a_i, U_i) \right) P^b(U_0),$$

using Lemma 3 and equation (37) above to inductively show that

$$\begin{aligned} & P^b(r_t, z_t | a_t, U_t) \left(\prod_{i=t-1}^0 P^b(U_{i+1}, z_i | a_i, U_i) \right) P^b(U_0) \\ &= P^b(r_t, z_t | a_t, \mathcal{H}_{t-1}^o) M'_{t,a_t} \prod_{i=t-1}^0 (M_{i+1,a_{i+1}} P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1}) M'_{i,a_i}) M_{0,a_0} P^b(Z_0), \quad (38) \end{aligned}$$

and again using the fact that

$$P^e(r_t) = \sum_{\tau^o \in \mathcal{T}_t^o} \Pi_e(\tau^o) P(r_t, z_t, \dots, z_0 | a_0, \dots, a_t)$$

to obtain equation (11).

Identification of $P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1})$

Our identification strategy for $P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1})$ in Theorem 4 extends that of Kuroki and Pearl (2014). Define the following matrices

$$\begin{aligned} P_{i,a_i}^\tau &:= \begin{pmatrix} 1 & P^b(\mathcal{F}_i^o | a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o | a_i) \\ P^b(\mathcal{H}_{i-1}^o | a_i) & P^b(\mathcal{F}_i^o, \mathcal{H}_{i-1}^o | a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o, \mathcal{H}_{i-1}^o | a_i) \\ \vdots & \vdots & \ddots & \vdots \\ P^b(\mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o | a_i) & P^b(\mathcal{F}_i^o, \mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o | a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o, \mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o | a_i) \end{pmatrix}, \\ Q_{i,a_i,r_i}^\tau &:= \begin{pmatrix} P^b(r_i | a_i) & P^b(\mathcal{F}_i^o, r_i | a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o, r_i | a_i) \\ P^b(\mathcal{H}_{i-1}^o, r_i | a_i) & P^b(\mathcal{F}_i^o, \mathcal{H}_{i-1}^o, r_i | a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o, \mathcal{H}_{i-1}^o, r_i | a_i) \\ \vdots & \vdots & \ddots & \vdots \\ P^b(\mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o, r_i | a_i) & P^b(\mathcal{F}_i^o, \mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o, r_i | a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o, \mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o, r_i | a_i) \end{pmatrix}, \\ U_{i,a_i}^\tau &:= \begin{pmatrix} 1 & P^b(\mathcal{F}_i^o | u_i^{(1)}, a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o | u_i^{(1)}, a_i) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & P^b(\mathcal{F}_i^o | u_i^{(\kappa)}, a_i) & \dots & P^b(\mathcal{F}_i^o | \mathcal{F}_i^o | u_i^{(\kappa)}, a_i) \end{pmatrix}, \\ R_{i,a_i}^\tau &:= \begin{pmatrix} 1 & P^b(\mathcal{H}_{i-1}^o | a_i, u_i^{(1)}) & \dots & P^b(\mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o | a_i, u_i^{(1)}) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & P^b(\mathcal{H}_{i-1}^o | a_i, u_i^{(\kappa)}) & \dots & P^b(\mathcal{H}_{i-1}^o | \mathcal{H}_{i-1}^o | a_i, u_i^{(\kappa)}) \end{pmatrix}, \\ M_{i,a_i}^\tau &:= \text{diag}(P^b(u_i^{(1)} | a_i), \dots, P^b(u_i^{(\kappa)} | a_i)). \end{aligned}$$

Lastly, define

$$\Delta_{i,a_i,r_i} := \text{diag}(P^b(r_i|a_i, u_i^{(1)}), \dots, P^b(r_i|a_i, u_i^{(\kappa)})).$$

The ordering $u_i^{(j)}$ is such that the elements on the diagonal of $\Delta_{i,z_{i+1}}$ are in non-decreasing order. That is, $P^b(r_i|a_i, u_i^{(1)}) \geq \dots \geq P^b(r_i|a_i, u_i^{(\kappa)})$, where $\kappa = |\mathcal{U}|$. Furthermore, S^j denotes the j th element of set S .

Now, we claim that Assumptions 4 and 5 imply the following

Lemma 4.

$$\text{rank}(P_{i,a_i}^\tau) = \text{rank}(Q_{i,a_i,r_i}^\tau) = |\mathcal{U}| \quad (39)$$

Proof. Define

$$E_d = \begin{pmatrix} 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & -1 \\ 0 & 1 & \cdots & 0 & -1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -1 \end{pmatrix} \in \mathbb{R}^{d \times d}$$

It is clear that

$$E_{|\mathcal{H}_{i-1}|}^T P_{i,a_i}^\tau E_{|\mathcal{F}_i|} = P^b(\mathcal{H}_{i-1}, \mathcal{F}_i|a_i) \text{ and } E_{|\mathcal{H}_{i-1}|}^T Q_{i,a_i,r_i}^\tau E_{|\mathcal{F}_i|} = P^b(\mathcal{H}_{i-1}, \mathcal{F}_i, r_i|a_i).$$

Hence, it suffices to show that

$$\text{rank}(P^b(\mathcal{H}_{i-1}, \mathcal{F}_i|a_i)) = \text{rank}(P^b(\mathcal{H}_{i-1}, \mathcal{F}_i, r_i|a_i)) = |\mathcal{U}|,$$

as E_d is clearly invertible for all d .

To see why

$$\text{rank}(P^b(\mathcal{H}_{i-1}, \mathcal{F}_i|a_i)) = \text{rank}(P^b(\mathcal{H}_{i-1}, \mathcal{F}_i, r_i|a_i)) = |\mathcal{U}|,$$

note that we have

$$\begin{aligned} P^b(\mathcal{H}_{i-1}, \mathcal{F}_i|a_i)^T &= P^b(\mathcal{F}_i, \mathcal{H}_{i-1}|a_i) \\ &= P^b(\mathcal{F}_i|a_i, U_i) P^b(U_i, \mathcal{H}_{i-1}|a_i) \\ &= P^b(\mathcal{F}_i|a_i, U_i) P^b(U_i|a_i, \mathcal{H}_{i-1}) \text{diag}(P^b(\mathcal{H}_{i-1}|a_i)) = P^b(\mathcal{F}_i|a_i, \mathcal{H}_{i-1}) \text{diag}(P^b(\mathcal{H}_{i-1}|a_i)). \end{aligned}$$

We are guaranteed that $\text{diag}(P^b(\mathcal{H}_{i-1}|a_i))$ is invertible (otherwise we would be conditioning on measure 0 events), and thus invertibility of P_{i,a_i}^τ follows from Assumption 4. As for Q_{i,a_i,r_i}^τ , note that

$$P^b(\mathcal{H}_{i-1}, \mathcal{F}_i, r_i|a_i)^T = P^b(\mathcal{F}_i|a_i, U_i) \text{diag}(P^b(r_i|a_i, U_i)) P^b(U_i|a_i, \mathcal{H}_{i-1}) \text{diag}(P^b(\mathcal{H}_{i-1}|a_i)).$$

As $\text{diag}(P^b(r_i|a_i, U_i))$ is invertible (the positivity of Assumption 5), we have that Q_{i,a_i,r_i}^τ will have rank $|\mathcal{U}|$. \square

Now, notice that

$$P_{i,a_i}^\tau = (R_{i,a_i}^\tau)^T M_{i,a_i}^\tau U_{i,a_i}^\tau \quad (40)$$

and

$$Q_{i,a_i,r_i}^\tau = (R_{i,a_i}^\tau)^T M_{i,a_i}^\tau \Delta_{i,a_i,r_i} U_{i,a_i}^\tau \quad (41)$$

Now, for any subset Λ of $\{2, \dots, |\mathcal{F}_i^o|\}$ of size $|\mathcal{U}| - 1$, define $U_{i,a_i}^\tau(\Lambda)$ to be the $|\mathcal{U}| \times |\mathcal{U}|$ matrix whose first column is the first column of U_{i,a_i}^τ and whose j th column is the Λ_{j-1} th column of U_{i,a_i}^τ for all $2 \leq j \leq |\mathcal{U}|$, where Λ_j is the j th smallest element of Λ . Equivalently, $U_{i,a_i}^\tau(\Lambda)$ can be written as $U_{i,a_i}^\tau L(\Lambda)$ where $L(\Lambda)$ is the $|\mathcal{F}_i^o| \times |\mathcal{U}|$ matrix whose first column is e_1 and whose j th column is $e_{\Lambda_{j-1}}$, where e_k denotes the k th standard basis vector of $\mathbb{R}^{|\mathcal{F}_i^o|}$, for $2 \leq j \leq |\mathcal{U}|$.

Note that we can identify a maximal set of $|\mathcal{U}|$ linearly independent columns of P_{i,a_i}^τ , containing the first column, via Gaussian elimination. Let Λ be the set of indices of these columns. Right-multiplying equations (40) and (41) by $L(\Lambda)$ thus gives,

$$P_{i,a_i}^\tau L(\Lambda) = (R_{i,a_i}^\tau)^T M_{i,a_i}^\tau U_{i,a_i}^\tau(\Lambda) \quad (42)$$

and

$$Q_{i,a_i,r_i}^\tau L(\Lambda) = (R_{i,a_i}^\tau)^T M_{i,a_i}^\tau \Delta_{i,a_i,r_i} U_{i,a_i}^\tau(\Lambda) \quad (43)$$

By assumption, $U_{i,a_i}^\tau(\Lambda)$ is invertible. Additionally, Lemma 4 implies that M_{i,a_i}^τ is invertible and $\text{rank}(R_{i,a_i}^\tau) = |\mathcal{U}|$, via equation (40), and so, from equation (42), we have $(P_{i,a_i}^\tau L(\Lambda))^+ = (U_{i,a_i}^\tau(\Lambda))^{-1} (M_{i,a_i}^\tau)^{-1} ((R_{i,a_i}^\tau)^T)^+$. This along with equation (43) gives the following:

Lemma 5. *Under Assumption 4, we have that*

$$(P_{i,a_i}^\tau L(\Lambda))^+ Q_{i,a_i,r_i}^\tau L(\Lambda) = (U_{i,a_i}^\tau(\Lambda))^{-1} \Delta_{i,a_i,r_i} U_{i,a_i}^\tau(\Lambda).$$

Lemma 5 then allows us to diagonalize the *observable* product $(P_{i,a_i}^\tau L(\Lambda))^+ Q_{i,a_i,r_i}^\tau L(\Lambda)$, as in Kuroki and Pearl (2014), to obtain the left eigenvalue matrix $T_{i,a_i}^\tau(\Lambda)$. Note that each row of $T_{i,a_i}^\tau(\Lambda)$ differs from the corresponding row of $U_{i,a_i}^\tau(\Lambda)$ only by a constant factor and that the first entry of each row of $U_{i,a_i}^\tau(\Lambda)$ is 1. This is guaranteed by the distinctness of Assumption 5, which guarantees distinct eigenvalues, Δ_{i,a_i,r_i} , and thus a unique ordering on the eigenvectors of $T_{i,a_i}^\tau(\Lambda)$. Hence, we have that $U_{i,a_i}^\tau(\Lambda) = \text{diag}((T_{i,a_i}^\tau)_1)^{-1} T_{i,a_i}^\tau(\Lambda)$, where $(T_{i,a_i}^\tau)_1$ denotes the first column of T_{i,a_i}^τ , thereby allowing for the successful identification of $U_{i,a_i}^\tau(\Lambda)$.

Finally, the remaining columns of U_{i,a_i}^τ outside of those indexed in Λ are identifiable by simply determining the linear dependence among the columns of P_{i,a_i}^τ and recognizing that they are the same for U_{i,a_i}^τ . That is, if, for $j \notin \{1\} \cup \Lambda$

$$(P_{i,a_i}^\tau)_j = \sum_{j' \in \{1\} \cup \Lambda} c_{j'}^j (P_{i,a_i}^\tau)_{j'},$$

for constants $c_{j'}$, then

$$(U_{i,a_i}^\tau)_j = \sum_{j' \in \{1\} \cup \Lambda} c_{j'}^j (U_{i,a_i}^\tau)_{j'},$$

thus allowing for the identification of all columns of U_{i,a_i}^τ . Again, the coefficients $c_{j'}^j$, for each $j \notin \{1\} \cup \Lambda, j' \in \{1\} \cup \Lambda$ are identifiable via Gaussian elimination on P_{i,a_i}^τ .

Remark 3. *We remark that, in practice, our estimates of the matrices P_{i,a_i}^τ will be noisy and thus often have rank greater than $|\mathcal{U}|$ (even if the true moment matrix has rank $|\mathcal{U}|$). In such cases, instead of identifying $|\mathcal{U}|$ linearly independent columns which also contain the first, one may consider all matrices $P_{i,a_i}^\tau L(\Lambda)$ for all choices of $\Lambda \subseteq \{2, \dots, |\mathcal{F}_i^o|\}$ of size $|\mathcal{U}| - 1$ and select the choice of Λ corresponding to the matrix of lowest condition number. Alternatively, this can be mitigated by making the support of f_i^o more coarse—by combining elements of the support—so that $P_{i,a_i}^\tau, Q_{i,a_i,r_i}^\tau$ and U_{i,a_i}^τ all have second dimension equal to $|\mathcal{U}|$ instead of $|\mathcal{F}_i^o|$.*

The last remaining step is to use the identification of $U_{i+1,a_{i+1}}^\tau$ to identify

$$P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1}) = P^b(\mathcal{F}_{i+1} | a_{i+1}, U_{i+1}) P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o).$$

Note that

$$P^b(\mathcal{F}_{i+1} | a_{i+1}, U_{i+1}) P^b(U_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) = P^b(\mathcal{F}_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o).$$

The fact that $P^b(\mathcal{F}_{i+1} | a_{i+1}, U_{i+1})$ has full column rank and is identifiable then immediately implies the identifiability of $P^b(U_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$ as

$$P^b(U_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o) = (P^b(\mathcal{F}_{i+1} | a_{i+1}, U_{i+1}))^+ P^b(\mathcal{F}_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o),$$

and because $P^b(\mathcal{F}_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$ is observable. Then, $P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$ is readily obtained via marginalization across a_{i+1} in $P^b(U_{i+1}, a_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$. Finally, as we have identified both $P^b(\mathcal{F}_{i+1} | a_{i+1}, U_{i+1})$ and $P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o)$, the identification of $P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1})$ follows from the fact that

$$P^b(\mathcal{F}_{i+1}^o, z_i | a_i, \mathcal{H}_{i-1}^o | a_{i+1}) = P^b(\mathcal{F}_{i+1} | a_{i+1}, U_{i+1}) P^b(U_{i+1}, z_i | a_i, \mathcal{H}_{i-1}^o).$$

B Experimental Details

B.1 Choice of Parameters in Experiment 2

In these experiments, we give a choice of parameters violating Assumption 1 by making $\text{rank}(P^b(U_i | a_i, Z_{i-1})) = 1 < |\mathcal{U}|$.

Define

$$P^b(Z_i | U_i) = \begin{pmatrix} \rho_0 & \rho_1 \\ 1 - \rho_0 & 1 - \rho_1 \end{pmatrix}, P^b(U_{i+1} | a_i = 0, U_i) = \begin{pmatrix} a & c \\ 1 - a & 1 - c \end{pmatrix}, P^b(U_{i+1} | a_i = 1, U_i) = \begin{pmatrix} b & d \\ 1 - b & 1 - d \end{pmatrix},$$

$$\pi_b^{(i)}(A_i | U_i) = \begin{pmatrix} \epsilon & \delta \\ 1 - \epsilon & 1 - \delta \end{pmatrix}, \text{ and } P^b(U_i) = \begin{pmatrix} s_i \\ 1 - s_i \end{pmatrix}.$$

Note that s_i for $i > 0$ are not actually free to choose (but rather determined by the other parameters). Also, define

$$P^b(Z_{i-1} | U_i) = \begin{pmatrix} \zeta_0^i & \zeta_1^i \\ 1 - \zeta_0^i & 1 - \zeta_1^i \end{pmatrix}.$$

Noting that

$$P^b(u_i|a_i, z_{i-1}) = \frac{P^b(a_i, z_{i-1}|u_i)P^b(u_i)}{\sum_{u'_i} P^b(a_i, z_{i-1}|u'_i)P^b(u'_i)},$$

we have that

$$P^b(U_i|a_i = 0, Z_{i-1}) = \left(\begin{array}{c} \frac{\epsilon\zeta_0 s_i}{\epsilon\zeta_0 s_i + \delta\zeta_1(1-s_i)} \quad \frac{\epsilon(1-\zeta_0)s_i}{\delta(1-\zeta_1)(1-s_i)} \\ \frac{\delta\zeta_1(1-s_i)}{\epsilon\zeta_0 s_i + \delta\zeta_1(1-s_i)} \quad \frac{\epsilon(1-\zeta_0)s_i + \delta(1-\zeta_1)(1-s_i)}{\epsilon(1-\zeta_0)s_i + \delta(1-\zeta_1)(1-s_i)} \end{array} \right)$$

and

$$P^b(U_i|a_i = 1, Z_{i-1}) = \left(\begin{array}{c} \frac{(1-\epsilon)\zeta_0 s_i}{(1-\epsilon)\zeta_0 s_i + (1-\delta)\zeta_1(1-s_i)} \quad \frac{(1-\epsilon)(1-\zeta_0)s_i}{(1-\epsilon)(1-\zeta_0)s_i + (1-\delta)(1-\zeta_1)(1-s_i)} \\ \frac{(1-\delta)\zeta_1(1-s_i)}{(1-\epsilon)\zeta_0 s_i + (1-\delta)\zeta_1(1-s_i)} \quad \frac{(1-\delta)(1-\zeta_1)(1-s_i)}{(1-\epsilon)(1-\zeta_0)s_i + (1-\delta)(1-\zeta_1)(1-s_i)} \end{array} \right).$$

A sufficient condition for $\text{rank}(P^b(U_i|a_i = 0, Z_{i-1})) = \text{rank}(P^b(U_i|a_i = 1, Z_{i-1})) = 1 < 2$ is that $\zeta_0 = \zeta_1$. We now write $P^b(Z_{i-1}|U_i)$ in terms of the other parameters.

Using the fact that

$$P^b(u_{i-1}, a_{i-1}|u_i) = \frac{P^b(u_i|u_{i-1}, a_{i-1})\pi_b^{(i-1)}(a_{i-1}|u_{i-1})P^b(u_{i-1})}{\sum_{a'_{i-1}, u'_{i-1}} P^b(u_i|u'_{i-1}, a'_{i-1})\pi_b^{(i-1)}(a'_{i-1}|u'_{i-1})P^b(u'_{i-1})}$$

we have that

$$P^b(U_{i-1}, a_{i-1} = 0|U_i) = \left(\begin{array}{c} \frac{a\epsilon s_{i-1}}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})} \quad \frac{(1-a)\epsilon s_{i-1}}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))} \\ \frac{c\delta(1-s_{i-1})}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})} \quad \frac{(1-c)\delta(1-s_{i-1})}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))} \end{array} \right)$$

and

$$P^b(U_{i-1}, a_{i-1} = 1|U_i) = \left(\begin{array}{c} \frac{b(1-\epsilon)s_{i-1}}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})} \quad \frac{(1-b)(1-\epsilon)s_{i-1}}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))} \\ \frac{d(1-\delta)(1-s_{i-1})}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})} \quad \frac{(1-d)(1-\delta)(1-s_{i-1})}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))} \end{array} \right).$$

Marginalizing across a_{i-1} gives that

$$P^b(U_{i-1}|U_i) = \left(\begin{array}{c} \frac{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1}}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})} \quad \frac{(1-a)\epsilon s_{i-1} + (1-b)(1-\epsilon)s_{i-1}}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))} \\ \frac{c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})} \quad \frac{(1-c)\delta(1-s_{i-1}) + (1-d)(1-\delta)(1-s_{i-1})}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))} \end{array} \right)$$

Finally, using the fact that

$$P^b(z_{i-1}|u_i) = \sum_{u'_{i-1}} P^b(z_{i-1}|u'_{i-1})P^b(u'_{i-1}|u_i),$$

we have that

$$\zeta_0 = \frac{\rho_0(a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1}) + \rho_1(c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))}{a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})}$$

and

$$\zeta_1 = \frac{\rho_0((1-a)\epsilon s_{i-1} + (1-b)(1-\epsilon)s_{i-1}) + \rho_1((1-c)\delta(1-s_{i-1}) + (1-d)(1-\delta)(1-s_{i-1}))}{1 - (a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1}))}.$$

Finally,

$$\begin{aligned} \zeta_0 = \zeta_1 &\iff (\rho_0 s_{i-1} + \rho_1(1-s_{i-1}))(a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1} + c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})) \\ &= \rho_0(a\epsilon s_{i-1} + b(1-\epsilon)s_{i-1}) + \rho_1(c\delta(1-s_{i-1}) + d(1-\delta)(1-s_{i-1})). \end{aligned} \quad (44)$$

Thus, for Experiment 2, we chose a setting of parameters so as to satisfy equation (44), thus violating Assumption 1.

B.2 Coarsening the Support in Experiment 3

As mentioned in Remark 3, one way of dealing with noisy estimates is by coarsening the support of the future distribution. This is done in our third experiment by taking the future, f_i , of u_i to consist only of a_{i+1} . That is, the set \mathcal{F}_i^o is simply \mathcal{A}_{i+1} . This has the benefit of having the same cardinality as $|\mathcal{U}|$ and thus allowing us to bypass identification of a strict subset of $|\mathcal{U}|$ (nearly) linearly independent columns. Additionally, the violation of Assumption 3 in Experiment 3—via way of making $\text{rank}(P^b(Z_i|U_i)) = 1$ as mentioned in Appendix B.5—does not render $\text{rank}(P^b(\mathcal{A}_{i+1}|U_i)) = 1$, thus allowing for the use of this choice of future in Experiment 3 to guarantee that $\text{rank}(P^b(\mathcal{F}_i^o|a_i, \mathcal{H}_{i-1}^o)) = |\mathcal{U}|$.




Figure 5: Average absolute residuals $|\hat{P}^e(r_t) - P^e(r_t)|$ over randomly generated POMDP environments which violate Assumption 3.

B.3 Adaptive Search in Experiment 3

In the adaptive search conducted in Experiment 3, we first generate the POMDP parameters uniformly at random until the matrices P_{i,a_i}^T have sufficiently low condition number and the observable product, $(P_{i,a_i}^T)^+ Q_{i,a_i,r_i}^T$, to be diagonalized has sufficiently low eigenvector condition number, which, in the case of 2×2 matrices is simply the inverse of the spectral gap (Stewart, 2001). After such a setting of parameters has been identified, we then again randomly generate parameters from a uniform distribution biased in the direction of what appears to be the optimal setting, with stricter stopping conditions on the condition numbers. For example, if, after the first round, one of the probability parameter settings is 0.9, we then draw that parameter from $\text{Unif}[0.9, 1]$ in the next round so as to find a setting of parameters which yields even better conditioned matrices.

B.4 Robustness of Theorem 3 in Environments Violating Assumption 3

The setup for this experiment is essentially the same as that of Experiment 1 described in Section 6, wherein the residual of each estimate of $P^e(r_t = 1)$ is averaged across 50 trials. Figure 5 shows results for the estimators of Theorems 3 and 4, where the average is over environments violating Theorem 3’s Assumption 3. In particular, note that, even though the environments violate Assumption 3, the average residuals produced by Theorem 3’s estimators are very low, indicating the high-quality estimates produced by Theorem 3 even in environments which violate its assumptions. Theorem 4, however, produces, on average, much larger residuals. This is in contrast to Experiment 3 in Figure 4, wherein Theorem 4’s estimator performs much better—in fact, even better than Theorem 3’s. The reason for the diminished quality of Theorem 4 in Experiment 4 is due to the fact that, on average, the matrices that need to be inverted and diagonalized have extremely poor matrix and eigenvector condition numbers, resulting in severe degradation of Theorem 4, as mentioned in Section 6, even though its assumptions are not explicitly violated.

B.5 Choice of Parameters in Experiments 3 and 4

In Experiments 3 and 4, in order to violate Assumption 3, we simply chose POMDPs in which $\text{rank}(P^b(Z_i|U_i)) < |\mathcal{U}|$. As $|\mathcal{Z}| = |\mathcal{U}| = 2$ in our experiments, this meant that we chose

$$P^b(Z_i|U_i) = \begin{pmatrix} \rho & \rho \\ 1 - \rho & 1 - \rho \end{pmatrix},$$

for $\rho \in [0, 1]$.

B.6 Importance Sampling Experiments and Results

Number of trajectories	True value (\pm standard error)	Theorem 6 estimated value	Baseline IS estimated value
1,000,000	1.119 \pm 0.002	157.639	1.430
5,000,000	1.119 \pm 0.002	1.116	1.445

Figure 6: Estimated values compared to true value of estimators of Theorem 6 and the baseline for horizon $H = 1$.

We also conducted experiments comparing our IS algorithm to that of Tennenholtz *et al.* (2020a) in an environment—again, chosen through adaptive search, similar to that described in Appendix B.3, so as to produce well-conditioned matrices—meeting the assumptions of Theorem 6 but not the sufficiency conditions of the baseline IS estimator of Tennenholtz *et al.*

Number of trajectories	True value (\pm standard error)	Theorem 6 estimated value	Baseline IS estimated value
1,000,000	1.863 \pm 0.002	309.309	2.314
5,000,000	1.863 \pm 0.002	2.878	2.166

Figure 7: Estimated values compared to true value of estimators of Theorem 6 and the baseline for horizon $H = 2$.

(2020a). Figures 6 and 7 shows the results of the two estimators compared at both 1,000,000 and 5,000,000 number of sampled trajectories at horizons $H = 1$ and $H = 2$. In Figure 6, we see that at horizon 1 (i.e., two time-steps), our estimator produces very inaccurate results for 1,000,000 sampled trajectories, but is much more accurate with 5,000,000 trajectories than even the baseline, which is known to be biased and inconsistent.

However, for $H = 2$, it appears that even 5,000,000 number of trajectories is not enough to beat the baseline IS estimator of Tennenholtz *et al.* (2020a). While the quality of Theorem 6’s estimator improves greatly from 1,000,000 to 5,000,000 number of trajectories, Figure 7 illustrates that our estimator requires even more trajectories in order to produce more accurate results than the biased and inconsistent baseline.

C OPE via Importance Sampling

We now give a derivation for the IS estimator given in Section 6. As shown in the proof sketch of Theorem 6, we can write

$$v_H(\pi_e) = \mathbb{E}[vW(v, \tau^o)] \quad (45)$$

with

$$W(v, \tau^o) := \frac{1}{P^b(v|\tau^o)} \Pi_e(\tau^o) \Gamma_b(v, \tau^o) \quad (46)$$

and

$$\Gamma_b(v, \tau^o) = \sum_{r_{0:H}: \sum r_i = v} \sum_{u_{0:H}} \frac{\prod_{i=0}^H P^b(r_i|u_i) P^b(u_{0:H}|\tau^o)}{\Pi_b(u_{0:H})}. \quad (47)$$

Hence, all that must be shown is the identifiability of $\Gamma_b(v, \tau^o)$, as the terms of $\frac{1}{P^b(v|\tau^o)} \Pi_e(\tau^o)$ are all either given or directly estimable from observable data.

C.1 Identifiability of $\Gamma_b(v, \tau^o)$

First, we show that under certain rank and distinctness assumptions, the vectors $\pi_b^{(i)}(a_i|U_i)$ and $P^b(r_i|U_i)$ are identifiable by extending the diagonalization method of Kuroki and Pearl (2014). We will treat both of the above vectors individually. Again, letting $\kappa = |\mathcal{U}|$, we first define a matrix which will be used in both derivations:

$$\Delta_{i, z_{i+1}} := \text{diag}(P^b(z_{i+1}|u_i^{(1)}), \dots, P^b(z_{i+1}|u_i^{(\kappa)})).$$

The ordering $u_i^{(j)}$ is such that the elements on the diagonal of $\Delta_{i, z_{i+1}}$ are in non-decreasing order. That is, $P^b(z_{i+1}|u_i^{(1)}) \geq \dots \geq P^b(z_{i+1}|u_i^{(\kappa)})$. Furthermore, we will let z^j , r^j , and a^j denote the j th elements of the sets \mathcal{Z} , \mathcal{R} , and \mathcal{A} , respectively. Henceforth, the vectors $\pi_b^{(i)}(a_i|U_i)$ and $P^b(z_i|U_i)$ are defined such that their j th entries are $\pi_b^{(i)}(a_i|u_i^{(j)})$ and $P^b(z_i|u_i^{(j)})$, respectively. Throughout our identification procedure for both of these vectors, we make the following crucial distinctness assumption, which will allow for unique eigendecomposition (up to constants):

Assumption 10 (Distinctness). *For each $z_{i+1} \in \mathcal{Z}$, the probabilities $\{P^b(z_{i+1}|u_i) : u_i \in \mathcal{U}\}$ are all distinct.*

Identifying $\pi_b^{(i)}(a_i|U_i)$

Again, similar to those defined by Kuroki and Pearl (2014), we first define the following matrices which will be key to the analysis, explicitly state all assumptions in terms of them, and then show the result.

Define the matrix following matrices

$$P_{i, z_i}^a := \begin{pmatrix} 1 & P^b(a_i^1|z_i) & \dots & P^b(a_i^{|\mathcal{A}|-1}|z_i) \\ P^b(z_{i-1}^1|z_i) & P^b(a_i^1, z_{i-1}^1|z_i) & \dots & P^b(a_i^{|\mathcal{A}|-1}, z_{i-1}^1|z_i) \\ \vdots & \vdots & \ddots & \vdots \\ P^b(z_{i-1}^{|\mathcal{Z}|-1}|z_i) & P^b(z_{i-1}^{|\mathcal{Z}|-1}, a_i^1|z_i) & \dots & P^b(a_i^{|\mathcal{A}|-1}, z_{i-1}^{|\mathcal{Z}|-1}|z_i) \end{pmatrix},$$

$$\begin{aligned}
Q_{i,z_i,z_{i+1}}^a &:= \begin{pmatrix} P^b(z_{i+1}|z_i) & P^b(a_i^1, z_{i+1}|z_i) & \cdots & P^b(a_i^{|\mathcal{A}|-1}, z_{i+1}|z_i) \\ P^b(z_{i-1}^1, z_{i+1}|z_i) & P^b(a_i^1, z_{i-1}^1, z_{i+1}|z_i) & \cdots & P^b(a_i^{|\mathcal{A}|-1}, z_{i-1}^1, z_{i+1}|z_i) \\ \vdots & \vdots & \ddots & \vdots \\ P^b(z_{i-1}^{|\mathcal{Z}|-1}, z_{i+1}|z_i) & P^b(a_i^1, z_{i-1}^{|\mathcal{Z}|-1}, z_{i+1}|z_i) & \cdots & P^b(a_i^{|\mathcal{A}|-1}, z_{i-1}^{|\mathcal{Z}|-1}, z_{i+1}|z_i) \end{pmatrix}, \\
U_i^a &:= \begin{pmatrix} 1 & \pi_b^{(i)}(a_i^1|u_i^{(1)}) & \cdots & \pi_b^{(i)}(a_i^{|\mathcal{A}|-1}|u_i^{(1)}) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \pi_b^{(i)}(a_i^1|u_i^{(\kappa)}) & \cdots & \pi_b^{(i)}(a_i^{|\mathcal{A}|-1}|u_i^{(\kappa)}) \end{pmatrix}, \\
R_{i,z_i}^a &= \begin{pmatrix} 1 & P^b(z_{i-1}^1|z_i, u_i^{(1)}) & \cdots & P^b(z_{i-1}^{|\mathcal{Z}|-1}|z_i, u_i^{(1)}) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & P^b(z_{i-1}^1|z_i, u_i^{(\kappa)}) & \cdots & P^b(z_{i-1}^{|\mathcal{Z}|-1}|z_i, u_i^{(\kappa)}) \end{pmatrix}, \\
M_{i,z_i}^a &:= \text{diag}(P^b(u_i^{(1)}|z_i), \dots, P^b(u_i^{(\kappa)}|z_i)).
\end{aligned}$$

In addition to Assumptions 6, 7, and 10, we also make the following rank assumption:

Assumption 11 (Rank). *The matrices P_{i,z_i}^a and Q_{i,z_i}^a have rank at least $|\mathcal{U}|$.*

With these conditions, we now show how to obtain the matrix U_i^a , which contains the desired probabilities, via diagonalization of observable matrices.

Again, we have that

$$P_{i,z_i}^a = R_{i,z_i}^a T M_{i,z_i}^a U_i^a \quad (48)$$

and

$$Q_{i,z_i,z_{i+1}}^a = R_{i,z_i}^a T M_{i,z_i}^a \Delta_{i,z_{i+1}} U_i^a \quad (49)$$

The identification of U_i^a from equations (48) and (49) is then essentially the same as that of $P^b(\mathcal{F}_{i+1}^o, z_i|a_i, \mathcal{H}_{i-1}^o||a_{i+1})$ in Appendix A.3, wherein we can either identify $|\mathcal{U}|$ independent columns of P_{i,z_i}^a (by looking at condition numbers of the noisy estimates of submatrices of P_{i,z_i}^a) or coarsen the support of $|\mathcal{A}|$ to make P_{i,z_i}^a , $Q_{i,z_i,z_{i+1}}^a$, and U_i^a have second dimension equal to $|\mathcal{U}|$ and then employ the eigendecomposition strategy of Kuroki and Pearl (2014) by diagonalizing $(P_{i,z_i}^a L(\Lambda))^+ Q_{i,z_i,z_{i+1}}^a L(\Lambda)$ in the former case or $(P_{i,z_i}^a)^+ Q_{i,z_i,z_{i+1}}^a$ in the latter, and renormalizing by the appropriate constants to make the first column of the left eigenvector matrix equal to $\mathbf{1}$.

Identifying $P^b(r_i|U_i)$

The identification procedure for $P^b(r_i|U_i)$ is essentially the same as that in the previous section, except that we use the following matrices:

$$\begin{aligned}
P_{i,a_i,z_i}^r &= \begin{pmatrix} 1 & P^b(r_i^1|z_i) & \cdots & P^b(r_i^{|\mathcal{R}|-1}|z_i) \\ P^b(z_{i-1}^1, a_i|z_i) & P^b(r_i^1, z_{i-1}^1, a_i|z_i) & \cdots & P^b(r_i^{|\mathcal{R}|-1}, z_{i-1}^1, a_i|z_i) \\ \vdots & \vdots & \ddots & \vdots \\ P^b(z_{i-1}^{|\mathcal{Z}|-1}, a_i|z_i) & P^b(r_i^1, z_{i-1}^{|\mathcal{Z}|-1}, a_i|z_i) & \cdots & P^b(r_i^{|\mathcal{R}|-1}, z_{i-1}^{|\mathcal{Z}|-1}, a_i|z_i) \end{pmatrix}, \\
Q_{i,z_i,z_{i+1},a_i}^r &= \begin{pmatrix} P^b(z_{i+1}|z_i) & P^b(r_i^1, z_{i+1}|z_i) & \cdots & P^b(r_i^{|\mathcal{R}|-1}, z_{i+1}|z_i) \\ P^b(z_{i-1}^1, z_{i+1}, a_i|z_i) & P^b(r_i^1, z_{i-1}^1, z_{i+1}, a_i|z_i) & \cdots & P^b(r_i^{|\mathcal{R}|-1}, z_{i-1}^1, z_{i+1}, a_i|z_i) \\ \vdots & \vdots & \ddots & \vdots \\ P^b(z_{i-1}^{|\mathcal{Z}|-1}, z_{i+1}, a_i|z_i) & P^b(r_i^1, z_{i-1}^{|\mathcal{Z}|-1}, z_{i+1}, a_i|z_i) & \cdots & P^b(r_i^{|\mathcal{R}|-1}, z_{i-1}^{|\mathcal{Z}|-1}, z_{i+1}, a_i|z_i) \end{pmatrix}, \\
U_i^r &= \begin{pmatrix} 1 & P^b(r_i^1|u_i^{(1)}) & \cdots & P^b(r_i^{|\mathcal{R}|-1}|u_i^{(1)}) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & P^b(r_i^1|u_i^{(k)}) & \cdots & P^b(r_i^{|\mathcal{R}|-1}|u_i^{(k)}) \end{pmatrix}, \\
R_{i,z_i,a_i}^r &= \begin{pmatrix} 1 & P^b(a_i, z_{i-1}^1|z_i, u_i^{(1)}) & \cdots & P^b(a_i, z_{i-1}^{|\mathcal{Z}|-1}|z_i, u_i^{(1)}) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & P^b(a_i, z_{i-1}^1|z_i, u_i^{(k)}) & \cdots & P^b(a_i, z_{i-1}^{|\mathcal{Z}|-1}|z_i, u_i^{(k)}) \end{pmatrix},
\end{aligned}$$

and

$$M_{i,z_i}^r = \text{diag}(P^b(u_i^{(1)}|z_i), \dots, P^b(u_i^{(k)}|z_i)).$$

Again, analogously to the previous section, in addition to Assumptions 6, 7, and 10, we also make the following rank assumption:

Assumption 12 (Rank). *The matrices P_{i,z_i}^r and Q_{i,z_i}^r have rank at least $|\mathcal{U}|$.*

Importantly, we have the same identities on the above matrices:

$$P_{i,z_i}^r = R_{i,z_i}^r{}^T M_{i,z_i}^r U_i^a \quad (50)$$

and

$$Q_{i,z_i,z_{i+1}}^r = R_{i,z_i}^r{}^T M_{i,z_i}^r \Delta_{i,z_{i+1}} U_i^r \quad (51)$$

The procedure for identifying U_i^r is then precisely the same as that described in the previous section and Appendix A.3. Importantly, because the diagonal matrix is the *same* matrix $\Delta_{i,z_{i+1}}$ used to identify $\pi_b^{(i)}(a_i|U_i)$ and because of the distinctness prescribed by Assumption 10, the ordering $u_i^{(1)}, \dots, u_i^{(k)}$ is the same, so that when we identify the vector $P^b(r_i|U_i)$, the ordering of entries is indeed in corresponding order to our identification of $\pi_b^{(i)}(a_i|U_i)$.

We summarize the results from the previous 2 sections here before we proceed with the rest of the identification procedure for $\Gamma_b(v, \tau^\circ)$.

Theorem 7. *Under Assumptions 6, 7, 10, 11, and 12, the vectors $\pi_b^{(i)}(a_i|U_i)$ and $P^b(r_i|U_i)$ are identifiable, under the consistent ordering that their j th entries are $\pi_b^{(i)}(a_i|u_i^{(j)})$ and $P^b(r_i|u_i^{(j)})$, respectively.*

We now proceed with the identification procedure for $\Gamma_b(v, \tau^\circ)$. The sole remaining ingredient is the identification of $P^b(U_{0:H}|\mathcal{T}_H^\circ)$, which we now derive.

Identifying $P^b(U_{0:H}|\mathcal{T}_H^\circ)$

Lemma 6. *Under Assumptions 6, 7, and 8, the matrix $P^b(U_{0:H}|\mathcal{T}_H^\circ)$ is identifiable.*

Proof. Note that $P^b(R_{0:H}|U_{0:H}) = \bigotimes_{h=0}^H P^b(R_h|U_h)$. Thus $\text{rank}(P^b(R_{0:H}|U_{0:H})) = |\mathcal{U}|^{H+1}$ by Assumption 8, and thus $P^b(R_{0:H}|U_{0:H})$ has full column rank. The identifiability of $P^b(R_h|U_h)$ (up to the unknown ordering) implies the identifiability of $P^b(R_{0:H}|U_{0:H})$ (also up to the unknown ordering). Finally, note that

$$\begin{aligned} P^b(R_{0:H}|U_{0:H})P^b(U_{0:H}|\mathcal{T}_H^\circ) &= P^b(R_{0:H}|\mathcal{T}_H^\circ) \\ \implies P^b(U_{0:H}|\mathcal{T}_H^\circ) &= P^b(R_{0:H}|U_{0:H})^+ P^b(R_{0:H}|\mathcal{T}_H^\circ), \end{aligned}$$

thus rendering $P^b(U_{0:H}|\mathcal{T}_H^\circ)$ identifiable, as $P^b(R_{0:H}|\mathcal{T}_H^\circ)$ is observable. \square

Combining Identifications

Finally, recalling that

$$\begin{aligned} \Gamma_b(v, \tau^\circ) &= \sum_{r_{0:H}:\sum r_i=v} \sum_{u_{0:H}} \frac{\prod_{i=0}^H P^b(r_i|u_i) P^b(u_{0:H}|\tau^\circ)}{\Pi_b(u_{0:H})}, \\ &= \sum_{r_{0:H}:\sum r_i=v} \sum_{1 \leq j_0, \dots, j_H \leq \kappa} \frac{\prod_{i=0}^H P^b(r_i|u_i^{(j_i)}) P^b(u_0^{(j_0)}, \dots, u_H^{(j_H)}|\tau^\circ)}{\Pi_b(u_0^{(j_0)}, \dots, u_H^{(j_H)})}, \end{aligned}$$

the above derivations allow for identification for *each* of the terms $P^b(r_i|u_i^{(j_i)})$, $P^b(u_0^{(j_0)}, \dots, u_H^{(j_H)}|\tau^\circ)$, and $\Pi_b(u_0^{(j_0)}, \dots, u_H^{(j_H)})$, thus implying the desired identifiability for the entire IS estimation procedure, as desired.

We summarize this result and all conditions below:

Theorem 8. *Under Assumptions 6, 7, 8, 9, 10, 11, 12, the quantity $\Gamma_b(v, \tau^\circ)$ is identifiable for all values v and $\tau^\circ \in \mathcal{T}_H^\circ$. Hence, the importance weights*

$$W(v, \tau^\circ) := \frac{1}{P^b(v|\tau^\circ)} \Pi_e(\tau^\circ) \Gamma_b(v, \tau^\circ)$$

are all identifiable, allowing for the IS procedure given by $v_H(\pi_e) = \mathbb{E}[vW(v, \tau^\circ)]$