

NUMERICAL CONTINUED FRACTION INTERPOLATION

OLIVER SALAZAR CELIS*

Abstract. We show that highly accurate approximations can often be obtained from constructing Thiele interpolating continued fractions by a Greedy selection of the interpolation points together with an early termination condition. The obtained results are comparable with the outcome from state-of-the-art rational interpolation techniques based on the barycentric form.

Key words. Thiele continued fractions, univariate rational interpolation, best approximations

AMS subject classifications. 65D05, 65D15, 41A20, 41A50

1. Introduction. An alternative title to this note could be *The unreasonable effectiveness of Thiele continued fractions*. Indeed, it is known [1, 3] and further detailed in Section 3.5 that the construction of Thiele interpolating continued fractions can suffer from numerical instabilities. As argued in [3], a careful selection of the ordering of the interpolation points is therefore needed. Here we propose a Greedy strategy, one which takes the next point where the error is maximal. It is shown in Theorem 3.2 that, at least from an existence point of view, this strategy is well motivated. We cannot prove that it is the overall *best* strategy, but we rather give ample numerical evidence that the approach can lead to results that are competitive with other well-known approaches [10, 7, 2] for univariate rational interpolation and that the observed behavior is typical.

2. Thiele continued fraction rational interpolation. Consider a sequence of distinct complex points $(x_i)_{i \in \mathbb{N}}$ together with function valuations $f(x_i) = f_i$ of the complex function $f(x)$. The continued fraction

$$(2.1) \quad d_0 + \cfrac{x - x_0}{d_1} + \cfrac{x - x_1}{d_2} + \cfrac{x - x_2}{\dots}$$

with $d_i = \varphi_i[x_0, \dots, x_i]$ where

$$(2.2) \quad \begin{cases} \varphi_0[x_k] = f_k & k \geq 0 \\ \varphi_{i+1}[x_0, \dots, x_i, x_k] = \frac{x_k - x_i}{\varphi_i[x_0, \dots, x_{i-1}, x_k] - \varphi_i[x_0, \dots, x_i]} & k > i \end{cases}$$

generates rational interpolants when considering its successive convergents. The n th convergent

$$(2.3) \quad C_n(x) = \varphi_0[x_0] + \cfrac{x - x_0}{\varphi_1[x_0, x_1]} + \cfrac{x - x_1}{\varphi_2[x_0, x_1, x_2]} + \cfrac{x - x_2}{\dots} + \cfrac{x - x_{n-1}}{\varphi_n[x_0, \dots, x_n]}$$

has the property that $C_n(x_i) = f_i$ for $i = 0, \dots, n$, provided that the inverse differences $\varphi_i[x_0, \dots, x_i] \neq \infty$ exist and none of the tails

$$(2.4) \quad \begin{cases} T_{i,n}(x) = \varphi_i[x_i] + \cfrac{x - x_i}{\dots} + \cfrac{x - x_{n-1}}{\varphi_n[x_0, \dots, x_n]} & 0 \leq i < n \\ T_{n,n}(x) = \varphi_n[x_0, \dots, x_n] \end{cases}$$

*Department of Mathematics and Computer Science. University of Antwerp, Antwerp, Belgium. (oliver.salazarcelis@uantwerpen.be), ING Belgium, Brussels, Belgium

vanish at x_{i+1} . When $T_{i,n}(x_{i+1}) = 0$, then x_{i+1} is *unattainable*, meaning that it is a common zero of the numerator and denominator of $C_n(x)$ and $\lim_{x \rightarrow x_{i+1}} C_n(x) \neq f_{i+1}$.

3. Existence, sampling and numerical stability.

3.1. Off-diagonal interpolants. First, it is not difficult to show that the numerator and denominator degrees of the convergents $C_n(x)$ are at most $\lceil n/2 \rceil$ and $\lfloor n/2 \rfloor$ respectively. If desired, more off-diagonal interpolants can be obtained by considering $f(x) - p_\ell(x)$ instead of $f(x)$, with $p_\ell(x)$ an interpolating polynomial of degree at most ℓ . The introduction of such a polynomial could be useful to *dampen* functions $f(x)$ with a wide varying range. However, in what follows our interest is purely in Thiele continued fractions, hence we do not explore further potential benefits of non-diagonal interpolants.

3.2. Unattainability. While unattainability is undesirable, it is rarely encountered in practice except for pathological cases which are mostly related to symmetry of $f(x)$. Therefore we only mention that for $i = 1, \dots, n$, it can be checked a posteriori if desired: either by checking for zeros of the tails $T_{i-1,n}(x)$ at x_i or from direct evaluation of the convergents $C_n(x)$ at x_i . The next theorem says that a Thiele continued fraction is irreducible, up to possible unattainable points.

THEOREM 3.1. *Provided that the convergents $C_n(x)$ exist, its only common factors are of the form $(x - x_i)$ with $x_i \in \{x_0, \dots, x_{n-1}\}$ an interpolation point. In such a case the x_i are unattainable and are characterized by vanishing of the tails $T_{i+1,n}(x_i) = 0$.*

Proof. Elaborate proofs can for instance be found in [15, 3]. \square

3.3. Existence. Non-existence of inverse differences on the other hand is more inconvenient as it renders contributions from subsequent tails meaningless. The existence problem depends entirely on the ordering of the $(x_i)_{i \in \mathbb{N}}$. Specifically, it is required for two consecutive convergents $C_{i-1}(x)$ and $C_i(x)$ to be different ($i > 0$) in order for the inverse differences $\varphi_i[x_0, \dots, x_i]$ to exist.

THEOREM 3.2. *If the points $(x_i)_{i \in \mathbb{N}}$ are ordered such that every two consecutive convergents of the continued fraction (2.1) are different, then $\varphi_i[x_0, \dots, x_i] \neq \infty$.*

The proof is detailed in Appendix A, where it is shown that the inverse differences can essentially be interpreted as the ratio of two (linearized) residuals of successive convergents. This interpretation is used next to motivate a Greedy selection of the interpolation points.

3.4. Adaptive Greedy selection. Given a finite sequence of points $(x_i)_{0 \leq i \leq n}$, the condition of Theorem 3.2 where successive convergents should be different gives us a heuristic way to choose the next point x_{i+1} in the construction of $C_{i+1}(x)$ with $0 < i < n$. Given $C_i(x)$ and (x_0, \dots, x_i) , reorder the remaining points (x_{i+1}, \dots, x_n) to determine $C_{i+1}(x)$ such that $|C_i(x_{i+1}) - f(x_{i+1})|$ is maximal. In this way, $C_n(x)$ is ultimately constructed in an adaptive Greedy way by choosing in each step the point where for $0 < i < n$ the error between $C_i(x)$ and $f(x)$ is maximal. For the first point x_0 one can take a point where $|f(x_0)|$ is minimum. As such, at least one zero of $f(x)$ is accurately represented when present in the data.

This Greedy selection ensures the existence of the inverse differences (see Appendix A). A similar strategy is adopted by the AAA approach [7], although there the choice is motivated by numerical considerations rather than existence. We cannot

prove that the Greedy strategy is best for numerical stability in the context of Thiele continued fractions (discussed next), but the examples in this note suggest that the observed stable behaviour is typical.

3.5. Numerical stability. It is known [1] that the computation of the inverse differences and successive convergents in a Thiele interpolating continued fraction can suffer from numerical instabilities and worst case exponential loss of precision may occur in the calculation of the inverse differences [3]. Nevertheless, the backward evaluation of the continued fraction often leads to near machine precision (roughly $2\text{e-}16$) magnitudes for the residuals $f(x_i) - C_n(x_i)$ in practice [1]. We do not attempt an analysis like in [6] because any error analysis would necessarily depend on both the function $f(x)$ and the chosen interpolation points x_i . Instead, we provide compelling numerical evidence in Sections 4 and 5 that accurate approximations are often obtained, even on hard numerical problems, when the Greedy selection strategy of Section 3.4 is employed.

3.6. Early termination. From the discussion in the previous section, it is important to obtain the Thiele interpolating continued fractions in as few steps as possible. Besides the greedy selection, we add a stopping criterion when constructing $C_n(x)$. If the maximum absolute error in the remaining points is below a prescribed tolerance, say $\text{tol}=5\text{e-}15$, then we stop the recursion. The precise condition used is

$$\max |C_i(x_k) - f(x_k)| < \text{tol} \times \max_{i < j \leq n} (|f(x_j)|) \quad \text{for } i < k \leq n.$$

It essentially tells us that within numerical tolerance, the function to be approximated is rational. This can also be understood from the deeper connection to the shape of the blocks in the Walsh table [14], a topic we do not address further.

4. Example: interpolation of $|x|$ and \sqrt{x} . To illustrate that the Greedy selection strategy works well in practice, we start with the example of interpolation of $f(x) = |x|$ in Newman [9] points for $x \in [-1, 1]$. This problem demonstrates the approximation power of rational functions as compared to polynomials which can only achieve $\mathcal{O}(n^{-1})$ accuracy. However, it has proven quite challenging numerically and has been used to assess the performance of several rational approximations schemes based on for instance the barycentric form [10, 7, 5].

Newman approximations are rational interpolants in the $2n + 1$ points

$$(4.1) \quad (-1, -\eta, \dots, -\eta^{n-1}, 0, \eta^{n-1}, \dots, 1), \quad \text{with } \eta = e^{-1/\sqrt{n}}.$$

These points cluster around the origin, approaching it at an exponential rate as n increases. It is shown in [16] that the asymptotic rate of convergence of rational interpolants to $|x|$ is root-exponential $\mathcal{O}(n^{-1/2}e^{-\sqrt{n}})$.

Figure 1 shows the results of Thiele interpolation in Newman points and $f(x) = |x|$ with n up to 50. All obtained interpolants use all interpolation points in their construction, meaning that for $n = 50$ we have $2n + 1 = 101$ interpolation points and we construct $C_{2n}(x) = C_{100}(x)$. Mind that this is a challenging example because $C_{100}(x)$ is a rational function of degree $[n] = 50$ and $[n] = 50$ in numerator and denominator respectively. For instance the robust approach [10, see Fig. 5.6, p. 75] on the same example already breaks down at $n \approx 11$ for reasons related to rounding errors. The Thiele interpolants on the other hand provide accurate results and follow the theoretical root-exponential convergence. These results are in line with those observed in the more recent AAA approach [7, see Fig. 6.10, p. 1511]




Fig. 1: Adaptive Thiele interpolation of $|x|$ in Newman points for $n = 6, \dots, 50$. Left: maximum error on $[-1, 1]$ in approximation of $|x|$ by Thiele interpolation. Odd values of n give approximations with poles in $[-1, 1]$, while the even n approximations are pole-free. The asymptotic bound [16] is superimposed (dotted line). Right: 2-norm of the interpolation error in the Newman points (4.1) from backward evaluation of the Thiele interpolant. The evaluations remain highly accurate around 15 decimal digits precision and errors grow linearly (dotted line) as n increases.




Fig. 2: Adaptive Thiele interpolation of \sqrt{x} in squared Newman points (4.2) for $n = 5, \dots, 400$. Left: maximum error on $[0, 1]$ in approximation of \sqrt{x} by Thiele interpolation. The asymptotic bound [16] is superimposed (dotted line). Right: 2-norm of the interpolation error in the chosen interpolation points from backward evaluation of the Thiele interpolant. The evaluations remain highly accurate around 15 decimal digits precision and errors grow logarithmically (dotted line) as n increases.

An equivalent, but computationally easier, problem is to approximate $f(x) = \sqrt{x}$ on $x \in [0, 1]$. For this interpolation problem we take the square of the Newman-

points (4.1) leading to $n + 1$ unique points in $[0, 1]$:

$$(4.2) \quad (0, \eta^{2(n-1)}, \eta^{2(n-2)}, \dots, 1), \quad \text{with } \eta = e^{-1/\sqrt{n}}.$$

Note that these points cluster even closer near the origin than the original Newman points (4.1).

Figure 2 shows the results of Thiele interpolation in squared Newman points and $f(x) = \sqrt{x}$ with n up to 400. The error in the interpolation points seems to decay logarithmically with n . However, this is merely a consequence of the fact that for $n > 60$ the interpolants are no longer of full degree. Beyond that point, not all given interpolation points are used in the continued fraction construction and their degree only rises slowly. In fact, for $n = 400$ only 116 points are used after which the early termination kicks in. Nevertheless, the overall approximation quality improves in a root-exponential manner. We note that our implementation of the AAA method [7] fails for this example when $n > 50$ as it introduces real poles inside the interpolation interval.

5. Best approximations. While interpolation can deliver good approximations, interpolants are hardly ever *best* approximations. For the remainder of this section, let $f(x) \in \mathcal{C}[a, b]$. If $C_n^*(x)$ is a rational function, then it is called a best approximation to $f(x)$ on $x \in [a, b]$ when

$$C_n^*(x) = \operatorname{argmin}_{C_n(x), x \in [a, b]} |f(x) - C_n(x)|.$$

Recent advances in best rational approximation have been made in [2, 8, 4]. We refer the interested reader to those references for further details on best approximations.

Provided that $C_n^*(x)$ is of full degree, it is known that there exists a so-called *alternant set* consisting of $n + 2$ ordered nodes

$$(5.1) \quad a \leq \tilde{x}_0 < \tilde{x}_1 < \dots < \tilde{x}_{n+1} \leq b,$$

where $|f(x) - C_n^*(x)|$ attains its global extremum over all $x \in [a, b]$ with alternating signs:

$$f(\tilde{x}_j) - C_n^*(\tilde{x}_j) = (-1)^{\delta+j} \lambda, \quad j = 0, \dots, n + 1.$$

Here $\delta \in \{0, 1\}$ and $\lambda = \max_{x \in [a, b]} |f(x) - C_n^*(x)|$. The alternating set is usually the starting point for the iterative Remez algorithm. From an interpolation point of view, the alternant set is not immediately useful because the construction of $C_n(x)$ requires only $n + 1$ points and the value λ is unknown. However, we can use an alternative approach as discussed next.

Due to continuity, the error $f(x) - C_n^*(x)$ must attain zero between each pair $(\tilde{x}_j, \tilde{x}_{j+1})$ of neighboring points of the alternant set. This means that there must exist at least $n + 1$ points x_0, \dots, x_n

$$(5.2) \quad a \leq \tilde{x}_0 < x_0 < \tilde{x}_1 < x_1 < \dots < x_n < \tilde{x}_{n+1} \leq b,$$

such that $f(x_i) = C_n^*(x_i)$ for $i = 0, \dots, n$. This idea is exploited in the BRASIL [4] algorithm which, given an initial guess of $n + 1$ interpolation points, iteratively rescales the interval widths between successive interpolation points with the goal of equilibrating the local errors. In each step, a new set of interpolation points is determined.

Figure 3 illustrates the best rational approximation for $f(x) = \sin(20x)/(1 + 25x^2)$ on $x \in [-1, 2]$. To obtain the initial interpolation points, a low accuracy continued

fraction $C_{49}(x)$ is constructed from 100 Chebychev points of the first kind between -1 and 2 . Then the BRASIL iteration is ran, which relocates the interpolation points in each step. The interpolation itself is done using the adaptive Thiele continued fractions approach rather than using the barycentric form as in [4]. Other parameters, such as step size are put equal to 0.01 and convergence acceleration was not implemented. The final error $f(x) - C_{49}(x)$ equioscillates $n + 2 = 51$ times on $x \in [-1, 2]$ between the maximum error $\approx 1.76\text{e-}08$.




Fig. 3: Best approximation of $f(x) = \sin(20x)/(1 + 25x^2)$ on $x \in [-1, 2]$. The interpolation points are shown with dots. Left: Best approximation $C_{49}(x)$ (dashed line), the function $f(x)$ is shown in gray. Right: equioscillating error between $f(x)$ and $C_{49}(x)$.

For the last illustration, we turn to the example of [12]. In Figure 4, the results for $f(x) = \sqrt{x}$ on $x \in [0, 1]$ are shown. We found that a convenient starting point was obtained from a low(er) accuracy approximation based on 1000 linearly spaced points between 0 and 1 raised to the power 6 where the endpoints are removed. The iteration is started with $n + 1 = 81$ points adaptively chosen from the aforementioned set leading to the full degree interpolant $C_{80}^*(x)$ with maximum leveled error $\approx 4.39\text{e-}12$. Compared to the interpolants of similar size (i.e. constructed with the same number of interpolation points) in Figure 2a, the best approximation is roughly twice as accurate.

It is worth noting that one easily obtains an approximation of the same quality for $f(x) = |x|$ on $x \in [-1, 1]$ by taking $C_{80}^*(x^2)$. Such an approximation would be equivalent to the one obtained in [2, Fig. 7.1] of degree 80 in numerator and denominator. Mind that here the interpolation points range over more than 20 order of magnitudes and some of them are even below machine precision. A similar phenomenon is observed in [4, Fig. 3]. To be fair, the direct best approximation of $f(x) = |x|$ on $x \in [-1, 1]$ more often fails using the current approach, because the continued fraction representation struggles to maintain the exact symmetry.

6. Conclusions. We have shown that highly accurate approximations can often be obtained with Thiele continued fractions when incorporating a Greedy selection of the interpolation points together with an early termination condition. The obtained results are comparable with state-of-the art rational interpolation techniques




Fig. 4: Best approximation to \sqrt{x} in $n + 1 = 81$ points for $x \in [0, 1]$. The error from the final Thiele interpolant $C_{80}^*(x)$ equioscillates $n + 2 = 82$ times between the maximum error (shown in dashed). The location of the interpolation points is shown with dots. Left: Log scale x-axis plot (base 10) of the equioscillating error to visualize the area close to the origin. Right: regular plot of the equioscillating error to visualize the area closer to 1.

based on the barycentric form such as AAA [7] and can be used for rational minimax approximation [4].

The main advantage of the Thiele continued fraction is the simplicity of its construction. It does not require specialized linear algebra implementations such as SVD, but rather relies on an elementary recursion (see Appendix B for an implementation). Of course accumulation of rounding errors does occur, hence inevitably one can find examples where the approach will break down. The possibility of numerical breakdown is applicable for all rational interpolation approaches, including AAA [8, see §6 p.3172–3173].

Interpolation naturally assumes that the given data is noise free. Currently we do not see how a continued fraction approach could be used in such a context. However, we believe that for interpolation purposes a continued fraction method can often be competitive with other approaches.

Appendix A. Proof of Theorem 3.2.

For the proof of Theorem 3.2 we first relate the construction of the inverse difference to the (linearized) residuals of the convergents $C_n(x)$ of the continued fraction (2.1). To that end, let $C_n(x) = A_n(x)/B_n(x)$ where the n th numerator $A_n(x)$ and n th denominator $B_n(x)$ satisfy the recurrence relation [13]

$$(A.1) \quad \begin{pmatrix} A_n(x) \\ B_n(x) \end{pmatrix} = \begin{pmatrix} \varphi_n[x_0, \dots, x_n]A_{n-1}(x) + (x - x_n)A_{n-2}(x) \\ \varphi_n[x_0, \dots, x_n]B_{n-1}(x) + (x - x_n)B_{n-2}(x) \end{pmatrix},$$

with

$$\begin{cases} A_{-2}(x) = 0, & B_{-2}(x) = 1 \\ A_{-1}(x) = 1, & B_{-1}(x) = 0. \\ A_0(x) = \varphi_n[x_0] = f(x_0), & B_0(x) = 1 \end{cases}$$

THEOREM A.1. *If the points $(x_i)_{i \in \mathbb{N}}$ are ordered such that every two consecutive convergents of the continued fraction (2.1) are different then for $i = -1, 0, 1, \dots$*

$$\begin{cases} \varphi_0[x_k] = -\frac{R_{-2}(x_k)}{R_{-1}(x_k)}, & k > i = -1 \\ \varphi_{i+1}[x_0, \dots, x_i, x_k] = -(x_k - x_i) \frac{R_{i-1}(x_k)}{R_i(x_k)}, & k > i \geq 0 \end{cases},$$

where the linearized residuals $R_i(x)$ are defined as

$$R_i(x) = f(x)B_i(x) - A_i(x).$$

Proof. The proof is by induction. For $i = -1$

$$-\frac{R_{-2}(x_k)}{R_{-1}(x_k)} = -\frac{f(x_k)B_{-2}(x_k) - A_{-2}(x_k)}{f(x_k)B_{-1}(x_k) - A_{-1}(x_k)} = -\frac{f(x_k)}{-1} = \varphi_0[x_k].$$

And for $i = 0$

$$\begin{aligned} -(x_k - x_0) \frac{R_{-1}(x_k)}{R_0(x_k)} &= -(x_k - x_0) \frac{f(x_k)B_{-1}(x_k) - A_{-1}(x_k)}{f(x_k)B_0(x_k) - A_0(x_k)} \\ &= -(x_k - x_0) \frac{-1}{f(x_k) - f(x_0)} = \varphi_1[x_0, x_k]. \end{aligned}$$

Assume that the hypothesis $\varphi_i[x_0, \dots, x_{i-1}, x_k] = -(x_k - x_{i-1})R_{i-2}(x_k)/R_{i-1}(x_k)$ holds. First note that from application of (A.1) we have

$$R_i(x_k) = (x_k - x_{i-1})R_{i-2}(x_k) + \varphi_i[x_0, \dots, x_i]R_{i-1}(x_k).$$

Hence

$$-(x_k - x_i) \frac{R_{i-1}(x_k)}{R_i(x_k)} = \frac{-(x_k - x_i)R_{i-1}(x_k)}{(x_k - x_{i-1})R_{i-2}(x_k) + \varphi_i[x_0, \dots, x_i]R_{i-1}(x_k)}$$

Provided that $R_{i-1}(x_k) \neq 0$ and application of the induction hypothesis gives

$$-(x_k - x_i) \frac{R_{i-1}(x_k)}{R_i(x_k)} = \frac{-(x_k - x_i)}{-\varphi_i[x_0, \dots, x_{i-1}, x_k] + \varphi_i[x_0, \dots, x_i]} = \varphi_{i+1}[x_0, \dots, x_i, x_k].$$

For the last step, the presumed ordering of the points is important. If two consecutive convergents $C_{i-1}(x)$ and $C_i(x)$ are different, then the polynomial

$$(A.2) \quad A_{i-1}(x)B_i(x) - A_i(x)B_{i-1}(x) \neq 0$$

is of degree at most $\max\{[(i-1)/2] + [i/2], [i/2] + [(i-1)/2]\} \leq i$ which already vanishes at i points $(x_j)_{0 \leq j \leq i-1}$

$$R_{i-1}(x_j)B_i(x_j) - R_i(x_j)B_{i-1}(x_j) = 0, \quad j = 0, \dots, i-1.$$

Due to the condition (A.2) it cannot be that both $R_{i-1}(x)$ and $R_i(x)$ simultaneously vanish at other points. \square

Proof of Theorem 3.2. From Theorem A.1, we know that the inverse differences can be expressed as

$$\varphi_{i+1}[x_0, \dots, x_i, x_{i+1}] = -(x_{i+1} - x_i) \frac{R_{i-1}(x_{i+1})}{R_i(x_{i+1})}, \quad i \geq 0$$

and the presumed ordering of the $(x_i)_{i \in \mathbb{N}}$ ensures that the ratio $R_{i-1}(x_{i+1})/R_i(x_{i+1})$ is well-defined because its numerator and denominator cannot vanish simultaneously. In addition, $\varphi_{i+1}[x_0, \dots, x_i, x_{i+1}] = \infty$ would imply $R_i(x_{i+1}) = 0$ which contradicts that consecutive convergents $C_{i+1}(x)$ and $C_i(x)$ are different. \square

Appendix B. R Code. All codes have been developed in the open source R language [11]. The basic code for performing Thiele interpolation together with an example is given below.

```
evalcfrac <- function(aa,zz,x){
  #backward evaluation of continued fraction aa1 + (x-zz1)/aa2+...
  j = max(which(is.finite(aa))); res=rep(0,length(x))
  if(j>1){for(i in j:2){res=(x-zz[i-1])/(aa[i]+res)}}
  return(aa[1]+res)}
cfrac_interpolate <-function(xx,ff,tol=5e-15,NN=0){
  if(NN==0){NN=length(xx);} #use tol criteria to stop
  aa=rep(NA,NN); zz=rep(NA,NN); rr=rep(NA,NN)
  for(k in 1:NN){ #main loop
    if(k==1){ #init
      rr=ff #inverse differences
      i=which.min(abs(ff)) #smallest value
    }else{ i=which.max(abs(evalcfrac(aa,zz,xx)-ff)) #adaptive choice
      rr=(xx-zz[k-1])/(rr-aa[k-1])} #inverse differences
    aa[k]=rr[i];zz[k]=xx[i] #store cfrac coef
    ff=ff[xx!=xx[i]];rr=rr[xx!=xx[i]];xx=xx[xx!=xx[i]]; #reduce data
    if(k<NN){
      if(max(abs(evalcfrac(aa,zz,xx)-ff)<tol*max(abs(ff)))){
        print(paste("target precision reached earlier at n=",k))
        break();}}
  };return(list( "a"=aa[is.finite(aa)],"z"=zz[is.finite(zz)]))}
#example
f <- function(x) cos(exp(x))
x=seq(-1,1,length.out=100); y=f(x)
cfr=cfrac_interpolate(x,y)
plot(x,abs(f(x)-evalcfrac(cfr$a,cfr$z,x)))
The basic implementation of the BRASIL [4] iteration is given below.
findExtrema <- function(xk,a,b,cfr,f){
  xk = sort(unique(c(a,xk,b))) #add endpoints
  sk = numeric(0);resids = numeric(0);residabs = numeric(0)
  for(i in 2:length(xk)){ ## get extrema
    obj_func <- function(x) (f(x)-evalcfrac(cfr$a,cfr$z,x))
    opt = optimize(f = function(x) abs(obj_func(x)),
      lower = xk[i-1], upper = xk[i],maximum = TRUE, tol = 1e-30)
    resids = c(resids, f(opt$maximum)-evalcfrac(cfr$a,cfr$z,opt$maximum))
    sk = c(sk, opt$maximum)
  };res <- list("sk"= sk,"resids"= resids,"residabs" = abs(resids)) }
brasil <- function(xk,a,b,cfr,f,smax=1e-1,tt=1e-1, tol=1e-3){
  stop = FALSE
  while(!stop){
```

```

oldx = cfr$z[is.finite(cfr$z)]
res = findExtrema(oldx,a=a,b=b,cfr,f)
eps = max(res$residabs)/min(res$residabs)-1
if(eps<tol){ stop =TRUE
}else{
  h = mean(res$residabs) #mean err
  g = max(abs(res$residabs-h)) #max deviation from mean err
  gk = (res$residabs-h)/g
  s = min(c(smax, tt*g/h)) #stepzise
  ck = (1-s)^gk
  xkk = unique(sort(c(a,oldx,b)))
  lk = ck*diff(xkk) #rescaled length
  w = sum(sort(lk)) #normalisation
  newx = rep(NA, length(oldx))
  for(i in 1:length(newx)){
    newx[i] = (a*w + (b-a)*sum(lk[1:i]))/w
  };cfr = cfrac_interpolate(newx,f(newx),tol=0)
}
};return(cfr)}
#example sqrt
f <- function(x) sqrt((x))
x=unique(seq(0,1,length.out=1000)^6); x = sort(x)[2:(length(x)-1)]
cfr = cfrac_interpolate(x,f(x),tol=5e-15, NN=81) #init
cfrstar = brasil(cfr$z,0,1,cfr,f,smax=1e-1,tt=1e-1, tol=2e-4)
plot(x,abs(f(x)- evalcfrac(cfrstar$a, cfrstar$z,x)),type="l")

```

REFERENCES

- [1] A. A. CUYT, L. JACOBSEN, AND B. M. VERDONK, *Instability and modification of thiele interpolating continued fractions*, Applied Numerical Mathematics, 4 (1988), pp. 253–262.
- [2] S.-I. FILIP, Y. NAKATSUKASA, L. N. TREFETHEN, AND B. BECKERMANN, *Rational minimax approximation via adaptive barycentric representations*, SIAM Journal on Scientific Computing, 40 (2018), pp. A2427–A2455.
- [3] P. R. GRAVES-MORRIS, *Practical, Reliable, Rational Interpolation*, IMA Journal of Applied Mathematics, 25 (1980), pp. 267–286.
- [4] C. HOFREITHER, *An algorithm for best rational approximation based on barycentric rational interpolation*, Numerical Algorithms, 88 (2021), pp. 365–388.
- [5] G. ION VICTOR AND A. ATHANASIOS C., *Rational approximation of the absolute value function from measurements: a numerical study of recent methods*, arXiv, (2020).
- [6] W. B. JONES AND W. J. THRON, *Numerical stability in evaluating continued fractions*, Mathematics of Computation, 28 (1974), pp. 795–810.
- [7] Y. NAKATSUKASA, O. SÈTE, AND L. TREFETHEN, *The AAA algorithm for rational approximation*, SIAM Journal on Scientific Computing, 40 (2018), pp. 1494–1522.
- [8] Y. NAKATSUKASA AND L. N. TREFETHEN, *An algorithm for real and complex rational minimax approximation*, SIAM Journal on Scientific Computing, 42 (2020), pp. A3157–A3179.
- [9] D. J. NEWMAN, *Rational approximation to $|x|$* , Michigan Mathematical Journal, 11 (1964), pp. 11–14.
- [10] R. PACHÓN, *Algorithms for polynomial and rational approximation*, PhD thesis, University of Oxford, 2010.
- [11] R CORE TEAM, *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, 2020, <https://www.R-project.org/>.
- [12] R. VARGA, A. RUTTAN, AND A. KARPENTER, *Numerical results on best numerical approximation of $|x|$ on $[-1, +1]$* , Mathematics of the USSR-Sbornik, 74 (2007), p. 271.
- [13] J. WALLIS, *Arithmetica Infinitorum*, 1655.
- [14] J. L. WALSH, *On approximation to an analytic function by rational functions of best approximation*, Mathematische Zeitschrift, 38 (1934), pp. 163–176.
- [15] H. WERNER, *A reliable method for rational interpolation*, in Padé Approximation and its Applications, L. Wuytack, ed., Berlin, Heidelberg, 1979, Springer Berlin Heidelberg, pp. 257–277.

- [16] T. F. XIE AND S. P. ZHOU, *The asymptotic property of approximation to $|x|$ by Newman's rational operators*, Acta Mathematica Hungarica, 103 (2004), pp. 313–319.