

**SPECTRAL FLOW INSIDE ESSENTIAL SPECTRUM IV:
 F^*F IS A REGULAR DIRECTION**

NURULLA AZAMOV

ABSTRACT. Let H_0 and V be self-adjoint operators such that V admits a factorisation $V = F^*JF$ with bounded self-adjoint J and $|H_0|^{1/2}$ compact F . Flow of singular spectrum of the path of self-adjoint operators $H_0 + rV$, $r \in \mathbb{R}$, — also called spectral flow, through a point λ outside the essential spectrum of H_0 is well studied, and appears in such diverse areas as differential geometry and condensed matter physics.

Inside the essential spectrum the spectral flow through λ for such a path is well-defined if the norm limit

$$\lim_{y \rightarrow 0^+} F(H_0 + rV - \lambda - iy)^{-1} F^*$$

exists for at least one value of the coupling variable $r \in \mathbb{R}$. This raises the question: given a self-adjoint operator H_0 and $|H_0|^{1/2}$ compact operator F , for which real numbers λ there exists a bounded self-adjoint operator J such that the limit above exists? Real numbers λ for which this statement is true we call essentially regular or semi-regular and the operator $V = F^*JF$ we call a regular direction for H_0 at λ .

In this paper we prove that λ is semi-regular for H_0 if and only if the direction F^*F is regular.

1. INTRODUCTION

Flow of eigenvalues of a norm-continuous path of self-adjoint operators,

$$H_r = H_0 + rV,$$

which share the common essential spectrum, σ_{ess} , through a point λ outside the essential spectrum is well studied. The resulting integer number is also called *spectral flow*, which has independent origins in operator theory [Kr] and differential geometry [APS] and since then appeared in such areas as index theory and condensed matter physics, see e.g. [C].

For λ inside the essential spectrum the spectral flow, whether it is flow of eigenvalues or more generally flow of singular spectrum, is not well-defined, due to well-known extreme volatility of singular spectrum embedded in the essential spectrum. The spectral shift function (SSF) $\xi(\lambda)$ could have been considered as an analogue of spectral flow, however it is not integer-valued in σ_{ess} . It was found in [Az] and [AD] that SSF admits a natural decomposition into the sum of so-called absolutely continuous (ACSSF), $\xi^{(a)}$, and singular (SSSF), $\xi^{(s)}$, spectral shift functions, with $\xi^{(s)}(\lambda)$ being integer-valued for a.e. $\lambda \in \sigma_{ess}$, provided H_0 and V obey a certain trace class condition. Later $\xi^{(s)}(\lambda)$ was given other descriptions, most notably, as

2000 *Mathematics Subject Classification.* Primary 47A40.

Key words and phrases. Spectral flow, essential spectrum, resonance index, limiting absorption principle.

total resonance index (TRI), see [Az2] and [AD]. TRI is integer-valued for a.e. λ , and coincides with the classical spectral flow outside σ_{ess} , see [Az3].

In order to define TRI one does not need a trace class condition, — it suffices to assume the limiting absorption principle (LAP), see e.g. [AMG] and [Y] for more information on LAP. LAP admits many interpretations. We shall outline one which we will use.

Let H_0 be a self-adjoint operator on a Hilbert space \mathcal{H} and $F: \mathcal{H} \rightarrow \mathcal{K}$ be a closed $|H_0|^{1/2}$ -compact operator, which we call a *rigging* in \mathcal{H} . Then the pair H_0 and F is said to obey LAP if there exists the norm limit, $T_{\lambda+i0}(H_0)$, of the operator

$$T_{\lambda+iy}(H_0) := FR_{\lambda+iy}(H_0)F^* := F(H_0 - \lambda - iy)^{-1}F^*$$

for a.e. $\lambda \in \mathbb{R}$. TRI is well-defined at λ for a pair of operators H_0 and $V = F^*JF$, where $J \in \mathcal{B}_{sa}(\mathcal{K})$, if the norm limit $T_{\lambda+i0}(H_r)$ exists for at least one value of the coupling variable r , in which case it automatically exists for all r except a discrete set. We say that a real number λ is *essentially regular* or *semi-regular* for H_0 if there exists at least one $J \in \mathcal{B}_{sa}(\mathcal{K})$ such that the norm limit $T_{\lambda+i0}(H_0 + rF^*JF)$ exists for at least one r . In this case we also say that $V = F^*JF$ is a *regular direction* for H_0 at λ .

For more motivation for this paper I refer to papers [Az2, Az3, AD, AD2] and their introductions.

The main result of this paper is the following theorem.

Theorem 1.1. *Let H_0 be a self-adjoint operator and F be a closed $|H_0|^{1/2}$ -compact operator. If λ is semi-regular for H_0 , then F^*F is a regular direction.*

Theorem 1.1 is simple but important, as it gives a natural choice of a regular direction. It also allows to simplify the definition of a semi-simple point λ of a s.-a. operator H_0 as follows: if F^*F is a regular direction for H_0 at λ then λ is semi-simple. Finally, Theorem 1.1 shows that semi-simplicity of a point λ depends on a relationship between a self-adjoint operator H_0 and a rigging F only.

2. PROOF OF THEOREM 1.1

The premise means by definition that there exists a regular direction $V = F^*JF$ at H_0 , that is, $T_{\lambda+i0}(H_0 + rV)$ exists for all real numbers r except a discrete set. We need to show that for some real number r the norm limit

$$T_{\lambda+i0}(H_0 + rF^*F) =: T_{\lambda+i0}(\tilde{H}_r)$$

also exists. The second resolvent identity applied to the operator

$$\tilde{H}_{sr} = H_r + r(sF^*F - V),$$

where $s \in \mathbb{R}$, gives

$$T_z(\tilde{H}_{sr}) = T_z(H_r + rF^*(s - J)F) = \left[1 + rT_z(H_r)(s - J)\right]^{-1}T_z(H_r).$$

Thus, for some real number s the norm limit $T_{\lambda+i0}(\tilde{H}_{sr})$ exists if and only if the operator

$$1 + rT_{\lambda+i0}(H_r)(s - J)$$

is invertible and this is what we will prove. Assume the contrary. Then, since $T_{\lambda+i0}(H_r)(s - J)$ is compact, by Fredholm alternative for some non-zero analytic vector-valued function φ_s we have for all real numbers s

$$[1 + rT_{\lambda+i0}(H_r)(s - J)]\varphi_s = 0.$$

We can assume that $s > \|J\|$. Thus, the equality above means that -1 is an eigenvalue of $r\sqrt{s - J}T_{\lambda+i0}(H_r)\sqrt{s - J}$ for all real $s > \|J\|$, so, for non-zero vector function ψ_s we have

$$(1) \quad r\sqrt{s - J}T_{\lambda+i0}(H_r)\sqrt{s - J}\psi_s = -\psi_s.$$

Taking the scalar product of both sides of this equality by ψ_s and then taking the imaginary part of both sides we get

$$\langle \psi_s, \sqrt{s - J} \Im T_{\lambda+i0}(H_r) \sqrt{s - J} \psi_s \rangle = 0,$$

and since $\Im T_{\lambda+i0}(H_r) \geq 0$ from this we find

$$\Im T_{\lambda+i0}(H_r) \sqrt{s - J} \psi_s = 0.$$

Therefore, the equality (1) turns into

$$(2) \quad r\sqrt{s - J} \Re T_{\lambda+i0}(H_r) \sqrt{s - J} \psi_s = -\psi_s.$$

Recall that s is large enough for the operator $s - J$ to be invertible. Now we use a well-known lemma: for an analytic path of self-adjoint operators N_s the eigenvalue equation

$$N_s \varphi_s = \lambda(s) \varphi_s$$

implies

$$(N'_s \varphi_s, \varphi_s) = \lambda'(s) (\varphi_s, \varphi_s).$$

Applying this lemma to (2) gives

$$(\sqrt{s - J}^{-1} \Re T_{\lambda+i0}(H_r) \sqrt{s - J} \psi_s, \psi_s) + (\sqrt{s - J} \Re T_{\lambda+i0}(H_r) \sqrt{s - J}^{-1} \psi_s, \psi_s) = 0.$$

Combining this with (2) implies

$$(\psi_s, (s - J)^{-1} \psi_s) = 0.$$

Now since for large s the operator $(s - J)^{-1}$ is positive definite, it follows that for such s we have $\psi_s = 0$. Which is clearly impossible. This contradiction completes the proof.

Corollary 2.1. *If $V = F^*JF$ is a regular direction then so is $F^*|J|F$.*

Proof. This corollary is a consequence of the proof of Theorem 1.1. In the proof we need to replace $s - J$ by $s|J| - J$ and assume that $s > 1$. There is one slight difficulty, as the operator $\sqrt{s|J| - J}$ is not necessarily invertible, but it can be easily overcome: clearly ψ_s belongs to the closure of the range of $|J|$ and so we can restrict the eigenvalue equation to this subspace on which the operator $\sqrt{s|J| - J}$ is invertible. \square

An argument used in the proof of Theorem 1.1 also allows to prove

Corollary 2.2. *Suppose λ is a semi-regular point for H_0 . If $J \geq 0$ and the direction $V = F^*JF$ is λ -regular at H_0 then so is the direction $F^*\tilde{J}F$ for any $\tilde{J} \geq J$.*

Proof. Proof follows verbatim that of Theorem 1.1, with some obvious changes such as replacing $s - J$ by $s\tilde{J} - J$. \square

Acknowledgements. The author thanks his wife, Feruza, for financially supporting him during the work on this paper.

REFERENCES

- [AMG] W. O. Amrein, A. B. de Monvel, V. Georgescu, *C_0 -groups, commutator methods and spectral theory of N -body Hamiltonians*, Birkhäuser, 1996.
- [APS] M. Atiyah, V. Patodi, I. M. Singer, *Spectral Asymmetry and Riemannian Geometry. III*, Math. Proc. Camb. Phil. Soc. **79** (1976), 71–99.
- [Az] N. A. Azamov, *Absolutely continuous and singular spectral shift functions*, Dissertationes Math. **480** (2011), 1–102.
- [Az2] N. A. Azamov, *Spectral flow inside essential spectrum*, Dissertationes Math. **518** (2016), 1–156.
- [Az3] N. A. Azamov, *Spectral flow and resonance index*, Dissertationes Math. **528** (2017), 1–91.
- [AD] N. A. Azamov, T. W. Daniels, *Singular spectral shift function for resolvent comparable operators*, Math. Nachrichten (2019), 292, 1911–1930.
- [AD2] N. A. Azamov, T. W. Daniels, *Coupling resonances and spectral properties of the product of resolvent and perturbation*, in preparation.
- [C] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, 1994.
- [Kr] M. G. Kreĭn, *On the trace formula in perturbation theory*, Mat. Sb., **33** 75 (1953), 597–626.
- [Y] D. R. Yafaev, *Mathematical scattering theory: general theory*, Providence, R. I., AMS, 1992.

INDEPENDENT SCHOLAR, ADELAIDE, SA, AUSTRALIA
 Email address: azamovnurulla@gmail.com