

LARGE DEVIATIONS, MOMENT ESTIMATES AND ALMOST SURE INVARIANCE PRINCIPLES FOR SKEW PRODUCTS WITH MIXING BASE MAPS AND EXPANDING ON THE AVERAGE FIBERS

YEOR HAFOUTA

DEPARTMENT OF MATHEMATICS
THE OHIO STATE UNIVERSITY

ABSTRACT. In this paper we show how to apply classical probabilistic tools for partial sums $\sum_{j=0}^{n-1} \varphi \circ \tau^j$ generated by a skew product τ , built over a sufficiently well mixing base map and a random expanding dynamical system. Under certain regularity assumptions on the observable φ , we obtain a central limit theorem (CLT) with rates, a functional CLT, an almost sure invariance principle (ASIP), a moderate deviations principle, several exponential concentration inequalities and Rosenthal type moment estimates for skew products with α , ϕ or ψ mixing base maps and expanding on the average random fiber maps. All of the results are new even in the uniformly expanding case. The main novelty here (contrary to [2]) is that the random maps are not independent, they do not preserve the same measure and the observable φ depends also on the base space. For stretched exponentially α -mixing base maps our proofs are based on multiple correlation estimates, which make the classical method of cumulants applicable. For ϕ or ψ mixing base maps, we obtain an ASIP and maximal and concentration inequalities by establishing an L^∞ convergence of the iterates \mathcal{K}^n of a certain transfer operator \mathcal{K} with respect to a certain sub- σ -algebra, which yields an appropriate (reverse) martingale-coboundary decomposition.

1. INTRODUCTION AND A PREVIEW OF THE MAIN RESULTS

1.1. Quenched limit theorems for random dynamical systems. Let (X, \mathcal{B}, m) be a probability space and let $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$ be an invertible ergodic probability preserving system. Let $T_\omega : X \rightarrow X$, $\omega \in \Omega$ be a family of non-singular maps (i.e. $m \circ T_\omega^{-1} \ll m$) so that the corresponding skew product τ given by $\tau(\omega, x) = (\sigma\omega, T_\omega x)$ is measurable. A random dynamical system is formed by the sequence of compositions

$$T_\omega^n x, n \geq 0 \text{ where } T_\omega^n = T_{\sigma^{n-1}\omega} \circ \dots \circ T_{\sigma\omega} \circ T_\omega$$

taken along the orbit of a “random” point ω . The system $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$ is often referred to as the *driving system*, and the map σ is often referred to as the *base map*.

Let $\varphi : \Omega \times X \rightarrow \mathbb{R}$ be a measurable function (“an observable”) and let μ be a τ -invariant probability measure on $\Omega \times X$. Then μ can be decomposed as $\mu = \int \mu_\omega d\mathbb{P}(\omega)$, where μ_ω is a family of probability measures on X so that $(T_\omega)_* \mu_\omega = \mu_{\sigma\omega}$ for \mathbb{P} -a.e. ω . Set $S_n \varphi = \sum_{j=0}^{n-1} \varphi \circ \tau^j$. Then

$$S_n \varphi(\omega, x) := S_n^\omega \varphi(x) = \sum_{j=0}^{n-1} \varphi_{\sigma^j \omega} \circ T_\omega^j,$$

where $\varphi_\omega(\cdot) = \varphi(\omega, \cdot)$. For \mathbb{P} almost every ω we can consider the sequence of functions $S_n^\omega \varphi(\cdot)$ on the probability space $(X, \mathcal{B}, \mu_\omega)$ as random variables. Limit theorems for such sequences are called *quenched limit theorems*. Among the first papers dealing with quenched limit theorems for random dynamical systems are [35, 36], where in [35] a quenched large deviations principle was obtained, and in [36] a central limit theorem and a law of iterated logarithm were established. Since then quenched limit theorems for random dynamical systems were extensively studied. For instance, in [16, 22, 23, 20] almost sure invariance principle (almost sure approximation by a sum of independent Gaussians) was established for random expanding or hyperbolic maps T_ω , in [28, 21] Berry-Esseen theorems (optimal rates in the CLT) were obtained for similar classes of maps and in [28, 17, 18, 19] local central limit theorems were

achieved. In addition, in [30] several limit theorems were extended to random non-uniformly hyperbolic or expanding maps. We would also like to refer to [6] for related results concerning mixing rates for random non-uniformly hyperbolic maps and to [31] for related results concerning sequential dynamical systems, where an almost sure invariance principle (ASIP) was obtained. We note that in many of the examples these results are obtained for the unique measure μ such that μ_ω is absolutely continuous with respect to m . However, some results hold true even for maps $T_\omega : \mathcal{E}_\omega \rightarrow \mathcal{E}_{\sigma\omega} \subset X$ which are defined on random subsets of X (see [37]), where in this case the most notable choices of μ_ω are the, so called, random Gibbs measures (see [28, 42]).

1.2. Limit theorems skew-products. Let us consider the sums $S_n\varphi = \sum_{j=0}^{n-1} \varphi \circ \tau^j$ as random variables on the probability space $(\Omega \times X, \mathcal{F} \times \mathcal{B}, \mu)$. In this paper we will focus on limit theorems for such sequences of random variables. In order to demonstrate the difference between such limit theorems and the quenched ones, let us focus on the CLT. The quenched CLT means that for \mathbb{P} -a.e. ω , for all real t we have

$$\lim_{n \rightarrow \infty} \mu_\omega (\{x : S_n^\omega \varphi(x) - \mu_\omega(S_n^\omega \varphi) \leq t\sqrt{n}\}) = \frac{1}{\sqrt{2\pi\sigma}} \int_{-\infty}^t e^{-\frac{s^2}{2\sigma^2}} ds$$

where $\sigma \geq 0$ is the number which satisfies that $\sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_{\mu_\omega}(S_n^\omega \varphi)$ for \mathbb{P} -a.e. ω (see [36]). On the other hand, the CLT for the skew product means that for all real t we have

$$\lim_{n \rightarrow \infty} \mu (\{x : S_n \varphi(x) - \mu(S_n \varphi) \leq t\sqrt{n}\}) = \frac{1}{\sqrt{2\pi\Sigma}} \int_{-\infty}^t e^{-\frac{s^2}{2\Sigma^2}} ds$$

where $\Sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_\mu(S_n \varphi)$. Note that, in contrast with the quenched case, the summands $X_j = \varphi \circ \tau^j$ form a stationary sequence and, in applications, the existence of the limit Σ^2 follows from a sufficiently fast decay of $\text{Cov}(X_0, X_n)$ as $n \rightarrow \infty$. We also remark that both CLT's above are formulated when σ and Σ are positive, and when one of them vanishes then the convergence is towards the constant function 0.

When $\mu_\omega(\varphi_\omega)$ does not depend on ω then $\mu_\omega(\varphi_\omega) = \mu(\varphi)$ and $\sigma^2 = \Sigma^2$. In this case the quenched CLT implies the CLT for $S_n\varphi$ by integrating $\mu_\omega(\{x : S_n^\omega \varphi(x) - \mu_\omega(S_n^\omega \varphi) \leq t\sqrt{n}\})$ with respect to \mathbb{P} (and similarly other distributive limit theorems for the skew product follow from the quenched ones). However, it is less likely to be true when $\mu_\omega(\varphi_\omega)$ depends on ω . Remark that even when $\mu_\omega(\varphi_\omega)$ does not depend on ω other finer results like the ASIP do not follow by integration. Indeed the ASIP concerns an almost sure approximation of the partial sums at question by a sum of independent Gaussian random variables, but the quenched ASIP provides a construction of such a Gaussian process which depends on the fiber ω .

1.2.1. Annealed limit theorems: iid maps. A particular well studied case is when the maps $T_{\sigma^j\omega}$ are independent. That is, $\Omega = \mathcal{Y}^{\mathbb{Z}}$ is a product space, the coordinates ω_j of $\omega = (\omega_j)$ are independent (with σ being the left shift) and $T_\omega = T_{\omega_0}$ depends only on the 0-th coordinate. In this case the statistical behavior of the skew product τ can be investigated using the, so called, annealed transfer operator, given by (see [9, 8, 34]),

$$\mathcal{A}g(x) = \int \mathcal{L}_\omega g(x) d\mathbb{P}(\omega)$$

where \mathcal{L}_ω is the transfer operator corresponding to T_ω and the underlying reference measure m . In [2] it was shown that for several classes of random expanding maps, the operator \mathcal{A} is quasi compact. Using that, a variety of limit theorems were obtained¹ for random variables of the form

$$S_n \varphi(\omega, x) = \sum_{j=0}^{n-1} \varphi(T_{\omega_{j-1}} \circ \dots \circ T_{\omega_0} x)$$

where (ω, x) are distributed according to a τ -invariant measure μ of the form $\mathbb{P} \times (h dm)$ for some continuous function h , which satisfies $\mathcal{A}h = h$. The latter assumption means that the maps T_ω preserve

¹Such as a central limit theorem, a Berry-Esseen theorem, a local central limit theorem, local large deviations principle and an almost sure invariance principle.

the same measure $\nu = h dm$. The point is that once quasi compactness is achieved the classical Nagaev-Guivarch method (see [32]) can be applied. This method was applied successfully to obtain limit theorems for deterministic dynamical systems, i.e. when $T_\omega = T$ does not depend on ω , and in [2] (see also [5]) this method was applied to obtain annealed limit theorems. We note that since both the function φ and the measure $h dm$ do not depend on ω , and all the maps T_ω preserve the measure $h dm$ the fiberwise centering constant $\mu_\omega(S_n^\omega \varphi)$ and the usual centering constant $\mu(S_n \varphi)$ are both equal to $n \int \varphi(x) h(x) dm$. Hence, as discussed in the previous section, in this setup some annealed results like the CLT already follow from the quenched ones.

Independence here is crucial, since it yields that the iterates on the annealed transfer operator can be written as

$$(1.1) \quad \mathcal{A}^n g = \int \mathcal{L}_\omega^n g d\mathbb{P}(\omega),$$

where $\mathcal{L}_\omega^n = \mathcal{L}_{\sigma^{n-1}\omega} \circ \dots \circ \mathcal{L}_{\sigma\omega} \circ \mathcal{L}_\omega$, which is the transfer operator of T_ω^n . Hence, the statistical behavior of the iterates τ^n of the skew product can be described by the iterates of \mathcal{A} . Note that in this iid setup this approach works only when $\varphi(\omega, x) = \varphi(x)$ does not depend on ω since it requires substituting φ (and appropriate functions of φ) into the annealed operator.

1.2.2. The motivation behind the present paper: non iid maps and random functions. The starting point of this paper is the observation that when the coordinates (ω_j) are not independent² there is no apparent relation between the iterates τ^n of τ and the iterates of the annealed operator \mathcal{A} defined above. Thus, a natural question arising from [2, 5] is which limit theorems hold true for mixing base maps with non-independent coordinates, and functions φ which depend on ω . Moreover, the assumptions in [2] require all the maps T_ω to preserve the same absolutely continuous measure $\nu = h dm$, and it is also desirable to prove limit theorems without such assumptions. We note that without the above assumptions even the CLT was not obtained before for the skew products considered in this paper, which will be our first result.

The question described above was also one of the main motivations in [29], where a CLT, a local CLT and a renewal theorem were obtained for several classes of skew products with mixing base maps such as Markov shifts and non-uniform Young towers, together with uniformly expanding random maps. These results were obtained by a certain type of integration argument, however the method of [29] does not involve the iterates of an annealed transfer operator, and instead we studied directly integrals of the form $\int \mathcal{L}_\omega^n g_\omega d\mathbb{P}(\omega)$, and their complex perturbations (relying on the fiberwise ‘‘spectral’’ properties and a certain type of periodic point approach which was introduced in [28]). While [29] was the first paper to discuss limit theorem for skew products with non independent fiber maps and random observables, all the results there were obtained for fiberwise centered observables φ (i.e. $\mu_\omega(\varphi_\omega) = 0$). Moreover, the maps T_ω in [29] were uniformly expanding, the base map had a periodic point and the random transfer operator satisfied certain regularity assumptions as functions of ω around the periodic orbit. From this point of view, a second motivation for the present paper is to prove limit theorem for skew products with non-independent fiber maps $T_{\sigma^j\omega}$ without the fiberwise centralization assumption and without additional topological assumptions like the behavior around a periodic orbit. We note that apart from the CLT we did not consider in [29] any of the limit theorems obtained in the present paper, and so almost all the results in the present paper are new even under the fiberwise centering assumption.

1.3. Our new results and the method of the proofs. As explained in the previous section, the goal of this paper is to obtain limit theorems with deterministic centering conditions for skew products τ built over mixing base maps and non-uniformly expanding maps T_ω . More precisely, we still consider a product space $\Omega = \mathcal{Y}^{\mathbb{Z}}$, but with ‘‘weakly-dependent’’ coordinates ω_j instead of independent ones. We consider a family of non-uniformly expanding map $T_\omega = T_{\omega_0}$ and observables of the form $\varphi(\omega, x) = \varphi_{\omega_0}(x)$ and prove limit theorems for sequences of the form $Z_n = S_n \varphi - n \int \varphi d\mu$, where

$$S_n \varphi(\omega, x) = \sum_{j=0}^{n-1} \varphi_{\omega_j}(T_{\omega_{j-1}} \circ \dots \circ T_{\omega_0} x) = \sum_{j=0}^{n-1} \varphi_{\sigma^j \omega}(T_\omega^j(x))$$

²That is, that maps $T_{\sigma^j\omega}$ are not iid.

considered as a random variables on the probability space $(\Omega \times X, \mathcal{F} \times \mathcal{B}, \mu)$, where μ is the unique τ -invariant measure with μ_ω being absolutely continuous with respect to m (or when μ_ω is a random Gibbs measure). These results are obtained for a certain type of observables φ so that $\varphi_\omega(\cdot)$ has bounded variation, uniformly in ω . When the maps T_ω are expanding on the average we will also have a certain scaling assumption³, which was shown in [20] to be necessary for quenched limit theorems, and which is similarly necessary for obtaining limit theorems for the skew product. In what follows we will always assume that $\int \varphi d\mu = 0$, which is not really a restriction since we can always replace φ with $\varphi - \int \varphi d\mu$.

We obtain our results using two different methods, as described below.

1.3.1. A (functional) CLT, moment estimates, moderate deviations and exponential concentration inequalities for α -mixing driving systems via the method of cumulants. We assume first that the coordinates ω_n are α -mixing, with the n -th α mixing coefficient α_n (defined in (2.17)) satisfy $\alpha_n = O(e^{-c n^\eta})$ for some $c, \eta > 0$ (i.e. it is stretched exponential). The first step towards limit theorems is standard for stationary processes: we show that under the weaker condition $\sum_n n \alpha_n < \infty$, the limit

$$s^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_\mu(S_n), \quad S_n = S_n \varphi$$

exists and that it vanishes if and only if φ admits a certain co-boundary representation. When $s^2 > 0$ we show that $n^{-1/2} S_n$ converges in distribution towards a centered normal random variable with variance s^2 . More precisely, we obtain the convergence rate

$$\sup_{t \in \mathbb{R}} \left| \mu(S_n \leq t s \sqrt{n}) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-\frac{1}{2} x^2} dx \right| \leq C n^{-\frac{1}{2+4\gamma}}, \quad \gamma = 1/\eta.$$

An annealed CLT (that is for independent maps) was obtained in [5] for random toral automorphism and in [2] for more general maps. When the base map is only mixing (and φ depend on ω) it was obtained in [29] for fiberwise centered potentials (i.e. $\mu_\omega(\varphi_\omega) = 0$). One of the results in this paper is the CLT for stretched exponentially α -mixing base maps but without the fiberwise centering assumption (in fact, we will obtain a functional CLT, see Theorem 2.19 and the last paragraph of this section).

We also obtain certain type of large deviations results, which is often refereed to as a moderate deviations principle (see [14]). These results yield, for instance, that for every closed interval $[a, b]$ we have

$$\lim_{n \rightarrow \infty} \frac{1}{a_n^2} \ln \mu \left\{ (\omega, x) : \frac{S_n(\omega, x)}{a_n s n^{1/2}} \in [a, b] \right\} = -\frac{1}{2} \inf_{x \in [a, b]} x^2$$

where a_n is a sequence so that $a_n \rightarrow \infty$ and $a_n = o(n^{\frac{1}{2+4\gamma}})$. We also obtain several types of ‘‘stretched’’ exponential concentration inequalities (2.19), (2.20) and Gaussian moment estimates of Rosenthal type (2.21). These result are obtained using the method of cumulants. More precisely, we first obtain a certain type of multiple correlation estimates (see Proposition 3.4), and then by applying general theorem we conclude that the k -th cumulant of the sum S_n is at most of order $n(k!)^{1+\gamma} (c_0)^{k-2}$ for $k \geq 3$, where c_0 is some constant (see Theorem 3.1). Then we can apply the method of cumulants [46, 15]. In the annealed setup, using the quasi compactness of the annealed transfer operator large deviations principles and exponential concentration inequalities were obtained in [2], and the above results show that there is a similar behavior when the maps are not independent and the function φ depends on ω (see also the results in the next section where better exponential concentration inequalities are described).

The above multiple correlation estimates together with the method of cumulants and the Rosenthal type moment estimates also yield a functional CLT. Let us consider the random function $S_n(t) = n^{-1/2} S_{[nt]}$ on $[0, 1]$. Then we show that it converges in distribution in the Skorokhod space $D[0, 1]$ to sW , where W is a standard Brownian motion and $s^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_\mu(S_n)$.

1.3.2. Limit theorems ϕ or ψ mixing driving systems via martingale methods: almost sure invariance principle, concentration inequalities and maximal moment estimates. One of the strongest methods to prove central limit theorems and related results in probability theory and dynamical systems is the, so called, martingale-coboundary representation (Gordin’s method). For a sufficiently chaotic dynamical system (Y, \mathcal{G}, μ, T) and an observable $\varphi : Y \rightarrow \mathbb{R}$ it means that φ can be represented

³That is $\text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) < \infty$ for some tempered random variable K .

as $\varphi = u + \chi - \chi \circ T$ for some sufficiently regular function χ , and $(u \circ T^n)$ forms a reverse martingale difference. Such results are well known for deterministic expanding (or hyperbolic) dynamical systems, and we refer to [16, 41, 20] for a quenched and sequential versions of such martingale methods. When the base map σ of the skew product τ is either (sufficiently fast) ϕ or ψ mixing (see (2.22) and (2.23) for the relevant definitions) we obtain a certain type of L^∞ martingale-coboundary representation (i.e. $\chi \in L^\infty$) for the underlying class of observables φ with respect to the skew product τ . This was already established in [2] in the annealed setup, and here using different arguments we obtain such a representation for skew products with mixing base maps.

Once an L^∞ martingale-coboundary decomposition is achieved, as usual, we can apply the Azuma-Hoeffding inequality together with Chernoff's bounding method and obtain exponential concentration inequalities of the form

$$\mathbb{P}(|S_n - \mathbb{E}[S_n]| \geq tn + c_1) \leq c_2 e^{-c_3 nt^2}, t > 0$$

where c_1, c_2, c_3 are positive constants. These concentration inequities are better than the ones we obtain using the method of cumulants, although they involve the stronger notions of ϕ or ψ mixing instead of α -mixing⁴. Another immediate consequence is moment estimates of the form

$$\left\| \max_{1 \leq k \leq n} |S_k - \mathbb{E}[S_k]| \right\|_{L^p} = O(n^{1/2})$$

which hold for every $p \geq 1$. Such results are known in the annealed case [2], and we extend them to the skew products considered in this paper.

The idea behind the martingale-coboundary representation is as follows. Consider the sub- σ -algebra \mathcal{F}_0 of $\Omega \times X$ generated by the projection $\pi_0(\omega, x) = ((\omega_j)_{j \geq 0}, x)$, where $\omega = (\omega_j)_{j \in \mathbb{Z}}$. Then τ preserves \mathcal{F}_0 since $T_\omega = T_{\omega_0}$ depends only on ω_0 , and \mathcal{F}_0 can be viewed as a sub-system (or a factor) given by $(\Omega \times X, \mathcal{F}_0, \mu, \tau)$. Our main argument is that, under quite mild ϕ or ψ mixing rates for the coordinates ω_j , the iterates $\mathcal{K}^n \varphi$ of the transfer operator \mathcal{K} corresponding to this system converge fast enough in $L^\infty(\mu)$ towards $\mu(\varphi)\mathbf{1}$, where $\mathbf{1}$ is the function taking the constant value 1, and φ is our given observable. This convergence can be established for every function φ so that $\|\varphi\|_{K,2} = \text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi(\omega, \cdot)\|_{BV}) < \infty$ for an appropriate tempered random variable $K(\omega)$, or for any observable with $\text{esssup}_{\omega \in \Omega} \|\varphi(\omega, \cdot)\|_{BV} < \infty$ when the maps T_ω are uniformly expanding. We stress that in any case this is not a spectral result (even under exponential mixing), since the convergence of \mathcal{K}^n is not in an operator norm, and, in general, it does not have exponential rate. Indeed we only prove that

$$(1.2) \quad \|\mathcal{K}^n \varphi - \mu(\varphi)\|_{L^\infty} \leq C \|\varphi\|_{K,2} \cdot \gamma_n$$

where $\gamma_n = \delta^n + \phi_R([n/2])$ or $\gamma_n = \delta^n + \psi([n/2])$, and $\delta \in (0, 1)$ and $\phi_R(\cdot)$ and $\psi(\cdot)$ are the reverse ϕ -mixing coefficients and ψ -mixing coefficients defined in (2.22) and (2.23), respectively.

Another consequence of the martingale-coboundary representation is the almost sure invariance principle (ASIP). In [13] the authors proved that under certain assumptions, a reverse martingale M_n can be approximated almost surely by a sum of independent Gaussians. One consequence of the methods in [13] is for sums of the form $W_n = \sum_{j=0}^{n-1} \varphi \circ \tau^j$. For such sums, the conditions of [13, Theorem 3.2] shows that there is a coupling with a sequence of iid centered normal random variables Z_j with variance $s^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}(W_n)$ so that

$$\sup_{1 \leq k \leq n} \left| W_k - \sum_{j=1}^k Z_j \right| = O(n^{1/4} (\log n)^{1/2} (\log \log n)^{1/4}), \text{ almost surely.}$$

In our notations, the first and second conditions of [13, Theorem 3.2] about \mathcal{K} can be verified using (1.2). In order to show that the third (and last condition) about \mathcal{K} in [13, Theorem 3.2] is in force we will also need to provide more general estimates on expression of the form

$$\left\| \mathcal{K}^i(\bar{\varphi} \mathcal{K}^j \bar{\varphi}) - \mu(\mathcal{K}^i(\bar{\varphi} \mathcal{K}^j \bar{\varphi})) \right\|_{L^\infty}$$

for $1 \leq i, j \leq n$, where $\bar{\varphi} = \varphi - \mu(\varphi)$.

⁴However, they only require summable ϕ or ψ mixing coefficients and not stretched exponential ones.

We note that in [2] the annealed ASIP was obtained using Gouezel's approach [24] and not the martingale-coboundary approach. Gouezel's approach was also used in [3] to obtain an ASIP for non-independent maps with mixing base maps, but as indicated in [3] the results are mostly applicable for Gordin-Denker maps.

Finally, we also prove a vector-valued almost sure invariance principle for skew products with uniformly expanding random maps and exponentially fast α -mixing base maps via the method of Gouezel [24]. As we have mentioned above, this method was applied in [2] in the annealed setting, while in [3] it was applied for Gordin-Denker systems. In a final section we also discuss a few extensions such as different types of mixing base maps like Young towers or Gibbs-Markov maps, application of the method of cumulants for nonconventional sums of the form $S_n = \sum_{m=1}^n \prod_{j=1}^{\ell} \varphi_j \circ \tau^{q_j(m)}$, for polynomial $q_j(m)$, as well as extension of the results for different class of random expanding maps (the ones in [42]).

2. PRELIMINARIES AND MAIN RESULTS

2.1. The random maps. We begin by recalling the setup from [12]. Let (X, \mathcal{G}) be a measurable space endowed with a probability measure m and a notion of a variation $v: L^1(X, m) \rightarrow [0, \infty]$ which satisfies the following conditions:

- (V1) $v(th) = |t|v(h)$;
- (V2) $v(g+h) \leq v(g) + v(h)$;
- (V3) $\|h\|_{L^\infty} \leq C_v(\|h\|_1 + v(h))$ for some constant $1 \leq C_v < \infty$;
- (V4) for any $C > 0$, the set $\{h: X \rightarrow \mathbb{R} : \|h\|_1 + v(h) \leq C\}$ is $L^1(m)$ -compact;
- (V5) $v(\mathbf{1}) = 0$, where $\mathbf{1}$ denotes the function equal to 1 on X ;
- (V6) $\{h: X \rightarrow \mathbb{R}_+ : \|h\|_1 = 1 \text{ and } v(h) < \infty\}$ is $L^1(m)$ -dense in $\{h: X \rightarrow \mathbb{R}_+ : \|h\|_1 = 1\}$;
- (V7) for any $f \in L^1(X, m)$ such that $\text{essinf} f > 0$, we have

$$v(1/f) \leq \frac{v(f)}{(\text{essinf} f)^2}.$$

- (V8) $v(fg) \leq \|f\|_{L^\infty} \cdot v(g) + \|g\|_{L^\infty} \cdot v(f)$;
- (V9) for $M > 0$, $f: X \rightarrow [-M, M]$ measurable and every C^1 function $h: [-M, M] \rightarrow \mathbb{C}$, we have $v(h \circ f) \leq \|h'\|_{L^\infty} \cdot v(f)$.

We define

$$BV = BV(X, m) = \{g \in L^1(X, m) : v(g) < \infty\}.$$

Then, BV is a Banach space with respect to the norm

$$\|g\|_{BV} = \|g\| + v(g).$$

2.1. Remark. Observe that (V3) and (V8) imply that

$$(2.1) \quad \|fg\|_{BV} \leq C_v \|f\|_{BV} \cdot \|g\|_{BV} \quad \text{for } f, g \in BV.$$

2.2. Remark. We observe that in [12], assumption (V5) is replaced by the weaker $v(\mathbf{1}) < +\infty$. However, for the examples we have in mind, our stronger version is satisfied. In particular, (V5) implies that $\|\mathbf{1}\|_{BV} = 1$.

The rest of our setup is almost identical to [20], with a single additional requirement which will be indicated in what follows. Let $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$ be a probability space and $\sigma: \Omega \rightarrow \Omega$ an invertible ergodic measure-preserving transformation. Let $T_\omega: X \rightarrow X$, $\omega \in \Omega$ be a collection of non-singular transformations (i.e. $m \circ T_\omega^{-1} \ll m$ for each ω) acting on X . Each transformation T_ω induces the corresponding transfer operator \mathcal{L}_ω acting on $L^1(X, m)$ and defined by the following duality relation

$$(2.2) \quad \int_X (\mathcal{L}_\omega \phi) \varphi dm = \int_X \phi(\varphi \circ T_\omega) dm, \quad \phi \in L^1(X, m), \quad \varphi \in L^\infty(X, m).$$

Thus, we obtain a cocycle of transfer operators $(\Omega, \mathcal{F}, \mathbb{P}, \sigma, L^1(X, m), \mathcal{L})$ that we denote by $\mathcal{L} = (\mathcal{L}_\omega)_{\omega \in \Omega}$. For $\omega \in \Omega$ and $n \in \mathbb{N}$, set

$$\mathcal{L}_\omega^n := \mathcal{L}_{\sigma^{n-1}\omega} \circ \dots \circ \mathcal{L}_{\sigma\omega} \circ \mathcal{L}_\omega.$$

We recall the notion of a tempered random variable.

2.3. Definition. We say that a measurable map $K : \Omega \rightarrow (0, +\infty)$ is *tempered* if

$$\lim_{n \rightarrow \pm\infty} \frac{1}{n} \log K(\sigma^n \omega) = 0, \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega.$$

In this paper we will consider the following assumptions on the random transfer operators.

2.4. Definition. A cocycle $\mathcal{L} = (\mathcal{L}_\omega)_{\omega \in \Omega}$ of transfer operators is said to be *good* if the following conditions hold:

- Ω is a Borel subset of a separable, complete metric space and σ is a homeomorphism. Moreover, \mathcal{L} is \mathbb{P} -continuous, i.e. Ω can be written as a countable union of measurable sets such that $\omega \mapsto \mathcal{L}_\omega$ is continuous on each of those sets;
- There is a tempered random variable $N(\omega)$ so that

$$(2.3) \quad v(g \circ T_\omega) \leq N(\omega)v(g), \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega \text{ and } g \in BV.$$

- there exists a random variable $C : \Omega \rightarrow (0, +\infty)$ such that $\log C \in L^1(\Omega, \mathbb{P})$ and

$$\|\mathcal{L}_\omega h\|_{BV} \leq C(\omega)\|h\|_{BV}, \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega \text{ and } h \in BV;$$

- there exist $N \in \mathbb{N}$ and random variables $\alpha^N, K^N : \Omega \rightarrow (0, +\infty)$ such that

$$\int_{\Omega} \log \alpha^N d\mathbb{P} < 0, \quad \log K^N \in L^1(\Omega, \mathbb{P})$$

and, for \mathbb{P} -a.e. $\omega \in \Omega$ and $h \in BV$,

$$v(\mathcal{L}_\omega^N h) \leq \alpha^N(\omega)v(h) + K^N(\omega)\|h\|_1;$$

- for each $a > 0$ and \mathbb{P} -a.e. $\omega \in \Omega$, there exist random numbers $n_c(\omega) < +\infty$ and $\alpha_0(\omega), \alpha_1(\omega), \dots$ such that for every $h \in \mathcal{C}_a$,

$$(2.4) \quad \text{essinf}_x(\mathcal{L}_\omega^n h)(x) \geq \alpha_n\|h\|_1 \quad \text{for } n \geq n_c,$$

where

$$(2.5) \quad \mathcal{C}_a := \{h \in L^\infty(X, m) : h \geq 0 \text{ and } v(h) \leq a\|h\|_1\};$$

- $\log(\text{essinf}_{x \in X}(\mathcal{L}_\omega \mathbf{1})(x)) \in L^1(\Omega, \mathbb{P})$.

Finally, we say that the cocycle \mathcal{L} is *uniformly random* if the random variables C, α^N, K^N and n_c are constants and $\alpha_n(\omega)$ does not depend on n and ω .

2.5. Remark.

- Definition 2.4 almost coincides with [20, Definition 3], the only difference being the addition of (2.3) (which was considered in [20, Section 3].)
- The log-integrability assumption specified at the end of Definition 2.4 may easily be checked on explicit examples (see e.g. the discussion in [4, Remark 2.12]).
- Furthermore, this assumption implies a certain version of the “random covering” similar to (2.4), see [20, Remark 4].

Let us now give examples of systems satisfying our requirements. Our first example is essentially taken from [12].

2.6. Example (Lasota-Yorke cocycles). Consider $X = [0, 1]$, endowed with Lebesgue measure m and the classical notion of variation v . We say that $T : X \rightarrow X$ is a piecewise monotonic non-singular map (p.m.n.s map for short) if the following conditions hold:

- T is piecewise monotonic, i.e. there exists a subdivision $0 = a_0 < a_1 < \dots < a_N = 1$ such that for each $i \in \{0, \dots, N-1\}$, the restriction $T_i = T|_{(a_i, a_{i+1})}$ is monotonic (in particular it is a homeomorphism on its image).
- T is non-singular, i.e. there exists $|T'| : [0, 1] \rightarrow \mathbb{R}_+$ such that for any measurable $E \subset (a_i, a_{i+1})$, $m(T(E)) = \int_E |T'| dm$.

The intervals $(a_i, a_{i+1})_{i \in \{0, \dots, N-1\}}$ are called the intervals of T . We also set $N(T) := N$ and $\lambda(T) := \text{essinf}_{[0,1]} |T'|$.

We consider a family $(T_\omega)_{\omega \in \Omega}$ of random p.m.n.s as above, and such that $T : \Omega \times [0, 1] \rightarrow [0, 1]$, $(\omega, x) \mapsto T_\omega(x)$ is measurable. Denoting $N_\omega = N(T_\omega)$ and $\lambda_\omega = \lambda(T_\omega)$, we assume that

- The map $\omega \mapsto \left(v \left(\frac{1}{|T'_\omega|} \right), N_\omega, \lambda_\omega, a_1, \dots, a_{N_\omega-1} \right)$ is measurable.
- We have the following expanding-on-average property:

$$\lim_{K \rightarrow \infty} \int_{\Omega} \log \min(\lambda_\omega, K) d\mathbb{P}(\omega) \in (0, +\infty]$$

- The maps $\log(N_\omega)$ and $\log^+ \left(\frac{N_\omega}{\lambda_\omega} \right)$ are integrable.
- The map $\log^+ \left(v \left(\frac{1}{|T'_\omega|} \right) \right)$ is integrable.
- T_ω is covering, i.e. for any interval $I \subset [0, 1]$, there exists a random number $n_c(\omega) > 0$ such that for any $n \geq n_c$, one has

$$(2.6) \quad \text{essinf}_{[0,1]} \mathcal{L}_\omega^n(\mathbf{1}_I) > 0.$$

- $\log(\text{essinf}_{x \in X} (\mathcal{L}_\omega \mathbf{1})(x)) \in L^1(\Omega, \mathbb{P})$.

We will call a cocycle satisfying the previous assumptions an *expanding on average Lasota-Yorke cocycle*. For a countably-valued measurable family $(T_\omega)_{\omega \in \Omega}$ of expanding on average Lasota-Yorke cocycle, the associated cocycle of transfer operators $(\mathcal{L}_\omega)_{\omega \in \Omega}$ is good (see [19]).

The following example can be fruitfully compared to a similar one by Kifer [38].

2.7. Example. We consider $X = \mathbb{S}^1$, endowed with the Lebesgue measure m and the notion of variation given by $v(\phi) := \int_X |\phi'| dm = \|\phi'\|_{L^1}$. We consider a measurable map $T : \Omega \times X \rightarrow X$ such that $T_\omega := T(\omega, \cdot)$ is C^r , $r \geq 2$. In addition, we make the following assumptions:

- There exists a tempered random variable $N(\omega)$ so that (2.3) holds true;
- The map $\omega \in \Omega \mapsto \left(\int_X \frac{|T''_\omega|}{(T'_\omega)^2} dm, \lambda_\omega \right)$ is measurable, where $\lambda_\omega = \inf_{[0,1]} |T'_\omega|$.
- The following expanding on average property holds:

$$(2.7) \quad \int_{\Omega} \log(\lambda_\omega) d\mathbb{P}(\omega) > 0.$$

- The map $\log \left(\int_X \frac{|T''_\omega|}{(T'_\omega)^2} dm \right)$ is \mathbb{P} -integrable.
- $\log(\text{essinf}_{x \in X} (\mathcal{L}_\omega \mathbf{1})(x)) \in L^1(\Omega, \mathbb{P})$.

We call a family $(T_\omega)_{\omega \in \Omega}$ satisfying the previous assumptions a *smooth expanding on average cocycle*. For a family $(T_\omega)_{\omega \in \Omega}$, countably-valued and measurable, of smooth expanding on average cocycle which satisfy (2.3), the associated cocycle of transfer operators $(\mathcal{L}_\omega)_{\omega \in \Omega}$ is good (see [19, Example 16]). We note that our expansion on average condition (2.7) implies that \mathbb{P} -a.s, T_ω has non-vanishing derivative, hence is a local diffeomorphism and a monotonic map of the circle. As noted in [20, Example 6], smooth expanding on average cocycles satisfy a stronger version of the random covering property (which by [12, Remark 0.1] implies the one formulated in (2.6)): for each non-trivial interval $I \subset X$, for \mathbb{P} -a.e $\omega \in \Omega$, there is a $n_c := n_c(\omega, I) < \infty$ such that for all $n \geq n_c$,

$$T_\omega^n(I) = X.$$

2.2. The one dimensionality of the top Oseledets space: a summary of known results. In this section we recall two results from [20] that will be in constant use in the course of the proofs of all of our results.

2.8. Theorem ([20] Theorem 12). *Let $\mathcal{L} = (\mathcal{L}_\omega)_{\omega \in \Omega}$ be a good cocycle of transfer operators. Then, the following holds:*

- *there exists an essentially unique measurable family $(h_\omega)_{\omega \in \Omega} \subset BV$ such that $h_\omega \geq 0$, $\int_X h_\omega dm = 1$ and*

$$\mathcal{L}_\omega h_\omega = h_{\sigma\omega}, \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega;$$

- there is a random variable $\ell : \Omega \rightarrow (0, +\infty)$ such that for \mathbb{P} -a.e. $\omega \in \Omega$,

$$(2.8) \quad h_\omega \geq \ell(\omega) \quad m\text{-a.e.};$$

- for \mathbb{P} -a.e. $\omega \in \Omega$,

$$(2.9) \quad BV = \text{span}\{h_\omega\} \oplus BV^0,$$

where

$$BV^0 = \left\{ h \in BV : \int_X h \, dm = 0 \right\};$$

- $\omega \mapsto \|h_\omega\|_{BV}$ is tempered;
- there exist $\lambda > 0$ and for each $\epsilon > 0$, a tempered random variable $D = D_\epsilon : \Omega \rightarrow (0, +\infty)$ such that for \mathbb{P} -a.e. $\omega \in \Omega$ and $n \in \mathbb{N}$,

$$(2.10) \quad \|\mathcal{L}_\omega^n \Pi(\omega)\|_{BV} \leq D(\omega)e^{-\lambda n}$$

and

$$(2.11) \quad \|\mathcal{L}_\omega^n (Id - \Pi(\omega))\|_{BV} \leq D(\omega)e^{\epsilon n},$$

where $\Pi(\omega) : BV \rightarrow BV^0$ is a projection associated to the splitting (2.9).

Finally, for uniformly random cocycles the random variables $\ell(\omega)$ and $D(\omega)$ can be replaced with positive constants and $\omega \rightarrow \|h_\omega\|_{BV}$ is a bounded random variable.

2.9. Corollary ([20] Corollary 13). *Let $\mathcal{L} = (\mathcal{L}_\omega)_{\omega \in \Omega}$ be a good cocycle of transfer operators.*

Then, the following holds:

- If $(h_\omega)_{\omega \in \Omega} \subset BV$ is given by Theorem 2.8, then

$$(2.12) \quad \omega \mapsto \|1/h_\omega\|_{BV} \text{ is tempered.}$$

- For \mathbb{P} -a.e. $\omega \in \Omega$,

$$(2.13) \quad BV = \text{span}\{\mathbb{1}\} \oplus BV_\omega^0,$$

where

$$BV_\omega^0 = \left\{ h \in BV : \int_X h \, d\mu_\omega = 0 \right\},$$

and $d\mu_\omega = h_\omega dm$, $\omega \in \Omega$;

- there exist $\lambda' > 0$ and a tempered random variable $\tilde{D} : \Omega \rightarrow (0, +\infty)$ such that for \mathbb{P} -a.e. $\omega \in \Omega$ and $n \in \mathbb{N}$,

$$(2.14) \quad \|L_\omega^n \tilde{\Pi}(\omega)\|_{BV} \leq \tilde{D}(\omega)e^{-\lambda' n}$$

$$(2.15) \quad \|L_\omega^n (Id - \tilde{\Pi}(\omega))\|_{BV} \leq \tilde{D}(\omega),$$

where $\tilde{\Pi}(\omega) : BV \rightarrow BV_\omega^0$ is a projection associated to the splitting (2.13), and

$$L_\omega^n g = \mathcal{L}_\omega^n(gh_\omega)/h_{\sigma^n \omega}, \quad g \in BV, \quad n \in \mathbb{N}.$$

Finally, for uniformly random cocycles the random variable $\tilde{D}(\omega)$ can be replaced with a positive constant.

Since $\mathcal{L}_\omega h_\omega = h_{\sigma\omega}$ and \mathcal{L}_ω satisfy the duality relation (2.2) the measure μ_ω satisfies that for \mathbb{P} -a.e. ω we have $(T_\omega)_* \mu_\omega = \mu_{\sigma\omega}$. Thus μ_ω gives raise to a T -invariant probability measure μ on $\Omega \times X$ so that

$$\mu(A \times B) = \int_A \mu_\omega(B) d\mathbb{P}(\omega) = \int_{A \times B} h(\omega, x) d\mathbb{P}(\omega) dm(x)$$

for every measurable sets A in Ω and B in X , where $h(\omega, x) = h_\omega(x)$.

2.3. Main results: limit theorems for mixing base maps.

2.4. The observable. Let us take a measurable $\varphi : \Omega \times X \rightarrow \mathbb{R}$ so that $\int \varphi d\mu = 0$. Let $\tilde{K}(\omega)$ be the tempered random variable defined by

$$\tilde{K}(\omega) = \max \left(D(\omega), \tilde{D}(\omega), N(\omega), \|1/h_\omega\|_{BV} \right)$$

where $D(\omega)$, $\tilde{D}(\omega)$ and $N(\omega)$ are specified in the definition of a good cocycles and in Theorem 2.8 and Corollary 2.9. In order to describe our assumptions on the observable φ , we will need the following classical result (see [7, Proposition 4.3.3.]).

2.10. Proposition. *Let $\tilde{K} : \Omega \rightarrow (0, +\infty)$ be a tempered random variable. For each $\epsilon > 0$, there exists a tempered random variable $\tilde{K}_\epsilon : \Omega \rightarrow (1, +\infty)$ such that*

$$\frac{1}{\tilde{K}_\epsilon(\omega)} \leq \tilde{K}(\omega) \leq \tilde{K}_\epsilon(\omega) \quad \text{and} \quad \tilde{K}_\epsilon(\omega)e^{-\epsilon|n|} \leq \tilde{K}_\epsilon(\sigma^n \omega) \leq \tilde{K}_\epsilon(\omega)e^{\epsilon|n|},$$

for \mathbb{P} -a.e. $\omega \in \Omega$ and $n \in \mathbb{Z}$.

Next, using the notations of Proposition 2.10 let $K(\omega) = \tilde{K}_\epsilon(\omega)$ for some $\epsilon < \lambda''/3$, where $\lambda'' = \min(\lambda, \lambda')$, and λ and λ' care specified in Theorem 2.8 and Corollary 2.9, respectively.

2.11. Remark. From now on we will replace both λ and λ' by their minimum, which for notational convenience will be denoted by λ .

In what follows we will consider an observable $\varphi : \Omega \times X \rightarrow \mathbb{R}$ satisfying the scaling condition

$$(2.16) \quad \text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) < \infty$$

which was first introduced in [19]. In the uniformly random case $\tilde{K}(\omega)$ (and hence $K(\omega)$) can be replace by a positive constant, and so the scaling condition reads

$$\text{esssup}_{\omega \in \Omega} \|\varphi_\omega\|_{BV} < \infty.$$

The main goal in this paper is to obtain limit theorems for the sequence of functions

$$S_n = S_n \varphi = \sum_{j=0}^{n-1} \varphi \circ \tau^j$$

under certain mixing assumptions on the driving system $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$ and the above assumptions on the observable φ .

2.12. Remark. For expanding on the average maps the scaling condition (2.16) is necessary for limit theorems, see [20, Appendix]. In any case, our results are also new in the uniformly random case, and the readers who would prefer can just consider this case together with the assumption that $\text{esssup}_{\omega \in \Omega} \|\varphi_\omega\|_{BV} < \infty$.

2.5. Limit theorems. Let us first introduce our assumptions on the base map. Let (ξ_n) be a two sided stationary sequence taking values on some measurable space \mathcal{Y} . We assume here that $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$ is the corresponding shift system. Namely, $\Omega = \mathcal{Y}^{\mathbb{Z}}$, $\sigma((\omega_j)_j) = (\omega_{j+1})_j$ is the left shift and if $\pi_0 : \Omega \rightarrow \mathcal{Y}$ denotes the 0-th coordinate projection, then (ξ_n) has the same distribution as $(\pi_0 \circ \sigma^n)$. We also assume that $T_\omega = T_{\omega_0}$ and $\varphi(\omega, \cdot) = \varphi(\omega_0, \cdot)$ depend only on 0-th coordinate ω_0 of ω .

2.5.1. Limit theorems for stretched exponentially fast α -mixing driving processes.

Let $(\Omega_0, \mathcal{F}, \mathbf{P})$ be the probability space on which (ξ_n) is defined. We recall that the α -mixing (dependence) coefficient between two sub- σ -algebras \mathcal{G}, \mathcal{H} of \mathcal{F} is given by

$$\alpha(\mathcal{G}, \mathcal{H}) = \sup \{ |\mathbf{P}(A \cap B) - \mathbf{P}(A)\mathbf{P}(B)| : A \in \mathcal{G}, B \in \mathcal{H} \}.$$

The α -dependence coefficients of (ξ_n) are defined by

$$(2.17) \quad \alpha_n = \sup_k \alpha(\mathcal{F}_{-\infty, k}, \mathcal{F}_{k+n, \infty}) = \alpha(\mathcal{F}_{-\infty, 0}, \mathcal{F}_{n, \infty})$$

where $\mathcal{F}_{-\infty, k}$ is the σ -algebra generated by $\xi_j, j \leq k$ and $\mathcal{F}_{k+n, \infty}$ is generated by $\xi_j, j \geq k+n$. The last equality holds true due to stationarity. Let us consider the following class of mixing assumptions on the base map:

2.13. Assumption (Stretched exponential α mixing rates). There exist positive constants c_1, c_2 and η so that $\alpha_n \leq c_1 e^{-c_2 n^\eta}$ for every n .

Our first result concerns the variance of S_n and the central limit theorem (with rates).

2.14. Theorem. *Suppose that the cocycle \mathcal{L} is good. Let φ be an observable so that $\|\varphi\|_K := \text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) < \infty$, where $\varphi_\omega = \varphi(\omega, \cdot)$. Suppose that $\sum_n n \alpha_n < \infty$. Then the limit*

$$s = \lim_{n \rightarrow \infty} n^{-1/2} \|S_n - \mathbb{E}[S_n]\|_{L^2(\mu)}$$

exists and it vanishes if and only if $\varphi = r \circ \tau - r$ for some $r \in L^2(\mu)$. If in addition Assumption 2.13 is satisfied then $n^{-1/2} S_n$ converges in distribution to sZ , where Z is a standard normal random variable. Moreover, there is a constant $C > 0$ so that for all $n \in \mathbb{N}$,

$$(2.18) \quad \sup_{t \in \mathbb{R}} |\mu(S_n - \mathbb{E}[S_n] \leq ts\sqrt{n}) - \Phi(t)| \leq Cn^{-\frac{1}{2+4\gamma}}$$

where $\gamma = 1/\eta$ and Φ is the standard normal distribution function. The constant C depends only on $c_1, c_2, \eta, \|\varphi\|_K$ and the constant C_v (from the definition of the variation $v(\cdot)$), and an explicit formula for C can be recovered from the proof.

The proof of Theorem 2.14 appears in Section 3.2.1. As discussed in Sections 1.2 and 1.3, when the quenched CLT holds true with a deterministic centering, then the CLT for the skew product follows by integration. This was the approach for the CLT in [2], but in the setup of this paper the function φ and the measure μ_ω depend on ω , and so the quenched CLT only holds with fiberwise centering. Thus, the novelty of Theorem 2.14 is that the CLT is obtained for the skew product beyond the annealed case considered in [2]. Moreover, Theorem 2.18 also strengthens the CLT in [29], since our maps T_ω are not uniformly expanding, and the observable φ is not fiberwise centered.

Next, let us discuss our results concerning moderate deviations and exponential concentration inequalities.

2.15. Theorem. *Suppose that the cocycle \mathcal{L} is good, and let φ be an observable so that $\|\varphi\|_K = \text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) < \infty$. Let Assumption 2.13 hold and set $\gamma = \frac{1}{\eta}$. Then there exist constants $a_1, a_2 > 0$ so that for every $x > 0$ and $n \in \mathbb{N}$,*

$$(2.19) \quad P(S_n - \mathbb{E}[S_n] \geq x) \leq \exp\left(-\frac{x^2}{2(a_1 + a_2 x n^{-\frac{1}{2+4\gamma}})^{\frac{1+2\gamma}{1+\gamma}}}\right).$$

All the constants depend only on $c_1, c_2, \eta, \|\varphi\|_K$ and C_v from the definition of the variation $v(\cdot)$, and an explicit formula for them can be recovered from the proof.

We will also prove the following:

2.16. Theorem. *Suppose that the cocycle \mathcal{L} is good, and let φ be an observable so that $\|\varphi\|_K = \text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) < \infty$. Let Assumption 2.13 hold and set $\gamma = \frac{1}{\eta}$. Let us also assume that the asymptotic variance s^2 is positive.*

(i) Set $v_n = \sqrt{\text{Var}(S_n)}$ and when $v_n > 0$ also set $Z_n = \frac{S_n - \mathbb{E}[S_n]}{v_n}$. Let Φ be the standard normal distribution function. Then there exist constants $s_3, s_4, s_5 > 0$ so that for every $n \geq a_3$ we have $v_n > 0$ and for every $0 \leq x < a_4 n^{\frac{1}{2+4\gamma}}$,

$$(2.20) \quad \left| \ln \frac{P(Z_n \geq x)}{1 - \Phi(x)} \right| \leq a_5 (1 + x^3) n^{-\frac{1}{2+4\gamma}} \quad \text{and} \\ \left| \ln \frac{P(Z_n \leq -x)}{\Phi(-x)} \right| \leq a_5 (1 + x^3) n^{-\frac{1}{2+4\gamma}}.$$

The constants a_4, a_5 depend only on $c_1, c_2, \eta, \|\varphi\|_K$ and C_v , and an explicit formula for them can be recovered from the proof.

(ii) Let $a_n, n \geq 1$ be a sequence of real numbers so that

$$\lim_{n \rightarrow \infty} a_n = \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n n^{-\frac{1}{2+4\gamma}} = 0.$$

Then the sequence $W_n = (sn^{\frac{1}{2}}a_n)^{-1}S_n$, $n \geq 1$ satisfies the moderate deviations principle with speed $s_n = a_n^2$ and the rate function $I(x) = \frac{x^2}{2}$. Namely, for every Borel measurable set $\Gamma \subset \mathbb{R}$,

$$-\inf_{x \in \Gamma^\circ} I(x) \leq \liminf_{n \rightarrow \infty} \frac{1}{a_n^2} \ln \mu(W_n \in \Gamma) \leq \limsup_{n \rightarrow \infty} \frac{1}{a_n^2} \ln \mu(W_n \in \Gamma) \leq -\inf_{x \in \bar{\Gamma}} I(x)$$

where Γ° is the interior of Γ and $\bar{\Gamma}$ is its closure.

We also obtain the following Roesenthal type moments estimates.

2.17. Theorem. *Suppose that \mathcal{L} is a good cocycle. If $\|\varphi\|_K = \text{esssup}_{\omega \in \Omega} (K(\omega)\|\varphi_\omega\|_{BV}) < \infty$, then under Assumption 2.13, there exist a constant c_0 so that with $\gamma = 1/\eta$ for every integer $p \geq 1$ we have*

$$(2.21) \quad |\mathbb{E}[(S_n - \mathbb{E}[S_n])^p] - (\text{Var}(S_n))^{\frac{p}{2}} \mathbb{E}[Z^p]| \leq (c_0)^p (p!)^{1+\gamma} \sum_{1 \leq u \leq \frac{p-1}{2}} n^u \frac{p^u}{(u!)^2} = O(n^{(p-1)/2})$$

where Z be a standard normal random variable. In particular, $\|S_n - \mathbb{E}[S_n]\|_{L^p} = O(\sqrt{n})$ for every p . As in the previous theorems, the constant c_0 depends (explicitly) only on $c_1, c_2, \eta, \|\varphi\|_K$ and C_v .

We remark that Theorem 2.17 provides another proof of the CLT by the method of moments. Indeed, if $s^2 > 0$ then it follows that for every integer $p \geq 1$ the p -th moment of $(S_n - \mathbb{E}[S_n])n^{-1/2}s^{-1}$ converges to $\mathbb{E}[Z^p]$, where s^2 is the asymptotic variance. In fact, for even p 's we get the convergence rate $O(n^{-1/2})$, while for odd p 's we get the rate $O(n^{-1})$.

2.18. Remark. The proofs of Theorems 2.15, 2.16 and 2.17 appear in Section 3.2.2.

Theorems 2.15, 2.16 and 2.17 are well established for sufficiently fast mixing (in the probabilistic sense) sequences of random variables, where one of the most notable methods of proof is the, so-called, method of cumulants (see [46]). For random dynamical systems, a moderate deviations principle was obtained in [21], using a random complex Perron-Frobenius theorem. In the setup of [2], annealed (local) large deviations principles and exponential concentration inequalities were obtained for iid maps, and we expect that for independent maps the methods in [2] will yield results like Theorems 2.15, 2.16 and 2.17 as well. The novelty in Theorems 2.15, 2.16 and 2.17 is that we show how to apply the method of cumulants in the context of skew products with non independent fiber maps, which results in concentration inequalities, moderate deviations principles and Gaussian moment estimates beyond the annealed setup [2].

Finally, let us consider the random function $\mathcal{S}_n(t) = n^{-1/2}(S_{[nt]} - \mathbb{E}[S_{nt}])$ on $[0, 1]$. We also obtain a functional CLT.

2.19. Theorem. *Let \mathcal{L} be a good cocycle. Suppose that $\text{esssup}_{\omega \in \Omega} (K(\omega)\|\varphi_\omega\|_{BV}) < \infty$ and that Assumption 2.13 holds true. Then the random function \mathcal{S}_n converges in distribution towards the distribution of $\{sW_t\}$ where W is a standard Brownian motion (restricted to $[0, 1]$) and s^2 is the asymptotic variance.*

2.20. Remark. The proof of Theorem 2.19 appears in Section 3.3. In [2] an almost sure invariance principle (ASIP) was obtained, which yields the functional CLT. In the next section, using different mixing coefficients for the base map we will obtain an ASIP for the more general skew products considered in this paper. However Theorem 2.19 shows that the functional CLT already holds true for stretched exponential α mixing base maps.

2.5.2. An almost sure invariance principle and exponential concentration inequalities for ϕ and ψ -mixing driving processes (via martingale methods).

Let $(\Omega_0, \mathcal{F}, \mathbf{P})$ be the probability space on which (ξ_n) is defined. We recall that the ϕ -mixing and ψ (dependence) coefficient between two sub- σ -algebras \mathcal{G}, \mathcal{H} of \mathcal{F} is given by

$$\phi(\mathcal{G}, \mathcal{H}) = \sup \{ |\mathbf{P}(B|A) - \mathbf{P}(B)| : A \in \mathcal{G}, B \in \mathcal{H}, \mathbf{P}(A) > 0 \}$$

and

$$\psi(\mathcal{G}, \mathcal{H}) = \sup \left\{ \left| \frac{\mathbf{P}(A \cap B)}{\mathbf{P}(A)\mathbf{P}(B)} - 1 \right| : A \in \mathcal{G}, B \in \mathcal{H}, \mathbf{P}(A)\mathbf{P}(B) > 0 \right\}.$$

The reverse ϕ -mixing coefficients of (ξ_n) are defined by

$$(2.22) \quad \phi_{n,R} = \sup_k \phi(\mathcal{F}_{k+n,\infty}, \mathcal{F}_{-\infty,k}) = \phi(\mathcal{F}_{n,\infty}, \mathcal{F}_{-\infty,0})$$

while the ψ -mixing coefficients of (ξ_n) are defined by

$$(2.23) \quad \psi_n = \sup_k \psi(\mathcal{F}_{-\infty, k}, \mathcal{F}_{k+n, \infty}) = \psi(\mathcal{F}_{-\infty, 0}, \mathcal{F}_{n, \infty})$$

where $\mathcal{F}_{-\infty, k}$ is the σ -algebra generated by $\xi_j, j \leq k$ and $\mathcal{F}_{k+n, \infty}$ is generated by $\xi_j, j \geq k+n$. It is clear from the definitions of the mixing coefficients that

$$\alpha_n \leq \phi_{n, R} \leq \psi_n.$$

2.21. Theorem (Exponential concentration and maximal inequalities). *Let \mathcal{L} be a good cocycle. Suppose the observable satisfies that $\text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi_\omega\|_{BV}) < \infty$.*

Let \mathcal{F}_0 be the σ algebra generated by the map $\pi(\omega, x) = ((\omega_j)_{j \geq 0}, x)$, namely the one generated by \mathcal{B} and the coordinates with non-negative indexes in the ω direction. If either $\text{ess-inf}_x h_\omega(x) > 0$ and $\sum_n \phi_{n, R} < \infty$ or $\sum_n \psi_n < \infty$ then there is an \mathcal{F}_0 -measurable function $\chi \in L^\infty(\mu)$ so that if we set $u = \varphi + \chi \circ \tau - \chi$ then $(u \circ \tau^n)$ is a reverse martingale difference with respect to the reverse filtration $\{\tau^{-n} \mathcal{F}_0\}$. As a consequence:

(i) *There are constants $a_1, a_2, a_3 > 0$ so that the following exponential concentration inequality holds true: for every $t > 0$ we have*

$$(2.24) \quad \mathbb{P}(|S_n - \mathbb{E}[S_n]| \geq tn + a_1) \leq a_2 e^{-a_3 n t^2}.$$

The constants a_1, a_2, a_3 depend only on $\tilde{\Phi} = \sum_n \phi_{n, R} < \infty$ and c (or $\tilde{\Psi} = \sum_n \psi_n < \infty$), the constant C_v and $\|\varphi\|_{K, 2} = \text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi_\omega\|_{BV})$, and an explicit formula for them can be recovered from the proof.

(ii) *For every $p \geq 2$ we have*

$$(2.25) \quad \left\| \max_{1 \leq k \leq n} |S_k - \mathbb{E}[S_k]| \right\|_{L^p} \leq C_p n^{1/2}$$

where $C_p > 0$ is a constant (which can be recovered from the proof and depends only on p and the above constants).

The proof of Theorem 2.21 appears in Section 4. Let us note that once the martingale coboundary representation $\varphi = u + \chi - \chi \circ \tau$ is established, Theorem 2.21 (i) follows from the Azuma–Hoeffding inequality together with Chernoff’s bounding method and Theorem 2.21 (ii) follows from, the so called, Rio’s inequality [45] (see [43, Proposition 7]).

To obtain the martingale coboundary representation we show that if \mathcal{K} is the transfer operator⁵ corresponding to the system $(\Omega \times X, \mathcal{F}_0, \mu, \tau)$ then

$$(2.26) \quad \|\mathcal{K}^n \varphi - \mu(\varphi)\|_{L^\infty} = O_{n \rightarrow \infty}(\delta^n + \gamma_{\lfloor n/2 \rfloor})$$

where γ_n is either ψ_n or $\phi_{n, R}$, depending on the case, and $\delta \in (0, 1)$. Once this is established we can take

$$\chi = \sum_{n \geq 1} \mathcal{K}^n \varphi.$$

The proof of (2.26) is given in Proposition 4.3 (i).

Our next result is an almost sure invariance principle.

2.22. Theorem (ASIP). *Let \mathcal{L} be a good cocycle, and suppose that the observable satisfies that $\text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi_\omega\|_{BV}) < \infty$.*

When $\text{ess-inf}_x h_\omega(x) > 0$ we set $\gamma_n = \phi_{R, n}$, while otherwise we set $\gamma_n = \psi_n$. In both cases, assume that

$$\sum_{n \geq 2} n^{5/2} (\log n)^3 \gamma_n^4 < \infty \quad \text{and} \quad \sum_{n \geq 2} n (\log n)^3 \gamma_n^2 < \infty$$

⁵Namely the one satisfying the duality relation

$$\int (\mathcal{K}g) \cdot f d\mu = \int g \cdot (f \circ \tau), \quad g \in L^1(\Omega \times X, \mathcal{F}_0, \mu), \quad f \in L^\infty(\Omega \times X, \mathcal{F}_0, \mu).$$

and

$$\sum_{n \geq 2} \frac{(\log n)^3}{n^2} \left(\sum_{k=0}^n (k+1)\gamma_k \right)^2 < \infty.$$

Then the limit

$$s^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \left[(S_n - \mathbb{E}[S_n])^2 \right]$$

exists and the following version of the almost sure invariance principle holds true: there is a coupling of $(\varphi \circ \tau^n)$ with a sequence of iid Gaussian random variables Z_j with zero mean and variance s^2 so that

$$\sup_{1 \leq k \leq n} \left| (S_k - \mathbb{E}[S_k]) - \sum_{j=1}^k Z_j \right| = O(n^{1/4}(\log n)^{1/2}(\log \log n)^{1/4}), \text{ almost surely.}$$

2.23. Remark. The ASIP implies the functional CLT, see [44]. Thus, Theorem 2.22 yields better results than Theorem 2.19 for ϕ_R or ψ mixing driving sequences (which are not necessarily stretched-exponentially mixing).

The proof of Theorem 2.22 appears in Section 4, and it relies on an application of [13, Theorem 3.2]. In addition to (2.26), in order to apply [13, Theorem 3.2] we will show that for all $1 \leq i, j \leq n$ we have

$$(2.27) \quad \left\| \mathcal{K}^i(\bar{\varphi} \mathcal{K}^j \bar{\varphi}) - \mu(\mathcal{K}^i(\bar{\varphi} \mathcal{K}^j \bar{\varphi})) \right\|_{L^\infty} = O_{n \rightarrow \infty}(\delta^n + \gamma_n)$$

where $\bar{\varphi} = \varphi - \mu(\varphi)$ and δ and γ_n are as in (2.26). The proof of (2.27) is given in Proposition 4.3 (ii).

2.24. Remark. As discussed in Section 1.3.2, the martingale-coboundary decomposition in Theorem 2.21 (and its consequences) is comparable with the annealed case [2], and the main novelty is that we obtain it for more general skew products and functions φ which depend on ω . Moreover, we do not assume that all T_ω preserve the same absolutely continuous probability measure. The ASIP's we obtain are comparable to ASIP's in [2] and [3] (see the discussion in Section 1.3.2).

2.5.3. A vector valued almost sure invariance principle in the uniformly random case for exponentially fast α -mixing base maps.

Let us take a vector-valued measurable function $\varphi = (\varphi_1, \dots, \varphi_d) : \Omega \times X \rightarrow \mathbb{R}^d$ so that $\varphi_\omega = \varphi(\omega, \cdot)$ depend on ω only through ω_0 and $\text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_{\omega, i}\|_{BV}) < \infty$ for all $1 \leq i \leq d$. Let us also assume that $\mu(\varphi_i) = 0$ for every i . Set $S_n = \sum_{j=0}^{n-1} \varphi \circ \tau^j$.

2.25. Theorem. *Suppose that $\alpha_n = O(\alpha^n)$ for some $\alpha \in (0, 1)$. Then there is a positive semi definite matrix Σ^2 so that*

$$\Sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Cov}(S_n).$$

Moreover, Σ^2 is positive definite if and only if $\varphi \cdot v \neq r - r \circ \tau$ for all unit vectors v and all $r \in L^2$.

Assume now that there are constants $C > 0$ and $\delta \in (0, 1)$ so that

$$(2.28) \quad \|\mathcal{L}_\omega^n \mathbf{1} - h_{\sigma^n \omega}\|_{BV} \leq C\delta^n$$

namely, that $K(\omega)$ is a bounded random variable. Then there is a coupling of $(\varphi \circ \tau^n)$ with a sequence of independent Gaussian centered random vectors (Z_n) so that $\text{Cov}(Z_n) = \Sigma^2$ and for every $\varepsilon > 0$,

$$\left| (S_n - \mathbb{E}[S_n]) - \sum_{j=1}^n Z_j \right| = o(n^{1/4+\varepsilon}), \text{ almost surely.}$$

3. LIMIT THEOREMS VIA THE METHOD OF CUMULANTS FOR α -MIXING DRIVING PROCESSES

We recall next that the k -th cumulant of a random variable W with finite moments of all orders is given by

$$\Gamma_k(W) = \frac{1}{i^k} \frac{d^k}{dt^k} (\ln \mathbb{E}[e^{itW}]) \Big|_{t=0}.$$

Note that $\Gamma_1(W) = \mathbb{E}[W]$, $\Gamma_2(W) = \text{Var}(W)$ and that $\Gamma_k(aW) = a^k \Gamma_k(W)$ for any $a \in \mathbb{R}$ and $k \geq 1$.

From now on we will assume that $\mathbb{E}[S_n] = 0$ for all n , namely we will replace φ by $\varphi - \mu(\varphi)$. The main result in this section is the following.

3.1. Theorem. *Let \mathcal{L} be a good cocycle, and suppose that Assumption 2.13 holds true and that $\|\varphi\|_K = \text{esssup}_{\omega \in \Omega} (K(\omega)\|\varphi_\omega\|_{BV}) < \infty$. Then, with $\gamma = 1/\eta$, there exists a constant c_0 which depends only on $\|\varphi\|_K$ and the constants from Assumption 2.13 so that for any $k \geq 3$,*

$$|\Gamma_k(S_n)| \leq n(k!)^{1+\gamma}(c_0)^{k-2}.$$

We will prove Theorem 3.1 by applying the following Proposition 3.3, which appears in [25] as Corollary 3.2.

Let us start with a few preparations. Let V be a finite set and $\rho : V \times V \rightarrow [0, \infty)$ be so that $\rho(v, v) = 0$ and $\rho(u, v) = \rho(v, u)$ for all $u, v \in V$. For every $A, B \subset V$ set

$$\rho(A, B) = \min\{\rho(a, b) : a \in A, b \in B\}.$$

We assume here that there exist $c_0 \geq 1$ and $u_0 \geq 0$ so that

$$(3.1) \quad |\{u \in V : \rho(u, v) \leq s\}| \leq c_0 s^{u_0}$$

for all $v \in V$ and $s \geq 1$.

Next, let $X_v, v \in V$ be a collection of centered random variables with finite moments of all orders, and for each $v \in V$ and $t \in (0, \infty]$ let $\varrho_{v,t} \in (0, \infty]$ be so that $\|X_v\|_t \leq \varrho_{v,t}$.

3.2. Assumption. For some $0 < \delta \leq \infty$ and all $k \geq 1$, $b > 0$ and a finite collection $A_j, j \in \mathcal{J}$ of (nonempty) subsets of V so that $\min_{i \neq j} \rho(A_i, A_j) \geq b$ and $r := \sum_{j \in \mathcal{J}} |A_j| \leq k$ we have

$$(3.2) \quad \left| \mathbb{E} \left[\prod_{j \in \mathcal{J}} \prod_{i \in A_j} X_i \right] - \prod_{j \in \mathcal{J}} \mathbb{E} \left[\prod_{i \in A_j} X_i \right] \right| \leq (r-1) \left(\prod_{j \in \mathcal{J}} \prod_{i \in A_j} \varrho_{i, (1+\delta)k} \right) \gamma_\delta(b, k)$$

where $\gamma_\delta(b, r)$ is some nonnegative number which depends only on δ, b and r , and $|\Delta|$ stands for the cardinality of a finite set Δ .

Set $W = \sum_{v \in V} X_v$. In the course of the proof of Theorems 2.14, 2.15, 2.16 and 2.19 we will need the following general result.

3.3. Proposition (Corollary 3.2, [25]). *Let (3.1) be and Assumption 3.2 be in force. Assume also that*

$$\tilde{\gamma}_\delta(m, k) := \max\{\gamma_\delta(m, r)/r : 1 \leq r \leq k\} \leq de^{-am^\eta}$$

for some $a, \eta > 0$, $d \geq 1$ and all $k, m \geq 1$. Then there exists a constant c which depends only on c_0, a, u_0 and η so that for every $k \geq 2$,

$$(3.3) \quad |\Gamma_k(W)| \leq d^k |V| c^k (k!)^{1+\frac{u_0}{\eta}} (M_k^k + M_{(1+\delta)k}^k)$$

where for all $q > 0$,

$$M_q = \max\{\varrho_{v,q} : v \in V\} \quad \text{and} \quad M_q^k = (M_q)^k.$$

When the X_v 's are bounded and (3.2) holds true with $\delta = \infty$ we can always take $\varrho_{v,t} = \varrho_{v,\infty}$, $t > 0$ and then for any $k \geq 2$,

$$(3.4) \quad |\Gamma_k(W)| \leq 2d^k |V| M_\infty^k c^k (k!)^{1+\frac{u_0}{\eta}}.$$

When $\delta < \infty$ and there exist $\theta \geq 0$ and $M > 0$ so that

$$(3.5) \quad (\varrho_{v,k})^k \leq M^k (k!)^\theta$$

for any $v \in V$ and $k \geq 1$, then for any $k \geq 2$,

$$(3.6) \quad |\Gamma_k(W)| \leq 3C^{\frac{\theta}{1+\theta}} d^k |V| c^k (1+\delta)^k M^k (k!)^{1+\frac{u_0}{\eta}+\theta}$$

where C is some absolute constant.

Theorem 3.1 will follow from the following result, which is proved in the next section.

3.4. Proposition. *For a good cocycle \mathcal{L} and an observable φ satisfying (2.16) we have the following. Fix some n and set $V = \{0, 1, \dots, n-1\}$ and $X_v = \varphi \circ \tau^v$. Set also $\rho(x, y) = |x - y|$, and let $t = \delta = \infty$, $\gamma_\infty(b, k) = \gamma_b = e^{-(\lambda-\varepsilon)b/3} + \alpha_{\lfloor b/3 \rfloor}$. Then condition (3.2) holds true with the above choices and with*

$$\varrho_{v, \infty} = A_0 \max(\text{esssup}_{\omega \in \Omega}(K(\omega)\|\varphi_\omega\|_{BV}), \|\varphi\|_{L^\infty})$$

where A_0 is a constant which depends only on $\lambda - 3\varepsilon$ and on the constant C so that $\sup |g| \leq C\|g\|_{BV}$ for every function $g : X \rightarrow \mathbb{C}$ (and the dependence can be easily recovered from the proof).

If in addition Assumption 2.13 holds then the conditions of Proposition 3.3 hold true with $u_0 = 1$, $c_0 = 2$ and $\gamma = 1/\eta$.

3.1. Multiple correlation estimates: proof of Proposition 3.4. Our goal is to show that (3.2) holds true with the desired upper bounds. We first need the following result.

3.5. Lemma. *For every two measurable functions g, h on $\mathcal{Y}^{\mathbb{N}}$ with $g, h \in L^\infty$ (w.r.t to the law of (ξ_n)) and all $k \in \mathbb{Z}$ and $n \in \mathbb{N}$ we have*

$$(3.7) \quad \begin{aligned} & |\mathbb{E}[g(\dots, \xi_{k-1}, \xi_k)h(\xi_{k+n}, \xi_{k+n+1}, \dots)] - \mathbb{E}[g(\dots, \xi_{k-1}, \xi_k)] \cdot \mathbb{E}[h(\xi_{k+n}, \xi_{k+n+1}, \dots)]| \\ & \leq \frac{1}{4} \|g(\dots, \xi_{k-1}, \xi_k)\|_{L^\infty} \|h(\xi_{k+n}, \xi_{k+n+1}, \dots)\|_{L^\infty} \alpha_n. \end{aligned}$$

Proof. By [11, Ch.4], we have

$$\alpha(\mathcal{G}, \mathcal{H}) = \frac{1}{4} \sup\{\|\mathbb{E}[h|\mathcal{G}] - \mathbb{E}[h]\|_{L^1} : h \in L^\infty(\Omega, \mathcal{G}, \mathbf{P}), \|h\|_{L^\infty} \leq 1\}.$$

Taking $g = g(\dots, \xi_{k-1}, \xi_k)$ and $h = h(\xi_{k+n}, \xi_{k+n+1}, \dots)$, $\mathcal{G} = \mathcal{F}_{-\infty, k}$ and $\mathcal{H} = \mathcal{F}_{k+n, \infty}$ we get

$$|\mathbb{E}[hg] - \mathbb{E}[g]\mathbb{E}[h]| = |\mathbb{E}[(\mathbb{E}[h|\mathcal{G}] - \mathbb{E}[h])g]| \leq \frac{1}{4} \alpha(\mathcal{G}, \mathcal{H}) \|g\|_{L^\infty} \|h\|_{L^\infty}.$$

□

Next, is it clearly enough to prove Proposition 3.4 when $\|\varphi\|_{L^\infty}$ and $\text{esssup}_{\omega \in \Omega}(K(\omega)\|\varphi_\omega\|_{BV})$ do not exceed 1, for otherwise we can just divide φ by the maximum between the two. Recall also our assumption that $K(\omega)e^{-\varepsilon|m|} \leq K(\sigma^m \omega) \leq K(\omega)e^{\varepsilon|m|}$ for some $\varepsilon < \lambda/3$ (recall Remark 2.11).

The first step in the proof of Proposition 3.4 is the following result.

3.6. Lemma (Fiberwise multiple correlation estimates). *Let B_1, B_2, \dots, B_m be nonempty intervals in the nonnegative integers so that B_i is to the left of B_{i+1} and B_1 contains 0. Let us denote by d_i the gap between B_i to B_{i+1} (namely the distance). Let us fix some ω and let f_i be a family of functions so that $K(\sigma^i \omega)\|f_i\|_{BV} \leq 1$ and $\|f_i\|_{L^\infty} \leq 1$. Let us define $F_j = F_{B_j, \omega} = \prod_{i \in B_j} f_i \circ T_\omega^i$. Then*

$$\left| \left(\int \left(\prod_{j=1}^m F_j \right) d\mu_\omega \right) - \left(\prod_{j=1}^m \int F_j d\mu_\omega \right) \right| \leq A \sum_{j=1}^{m-1} e^{-(\lambda-\varepsilon)d_j},$$

where $A = C^2 \sup_{d \in \mathbb{N}} 2de^{-(\lambda-\varepsilon)d}$ and λ comes from (2.10) and (2.14) (recall Remark 2.11).

Proof. The proof will be carried out by induction on m . Let us first prove the lemma in the case $m = 2$. We first note that for all functions g_0, g_1, \dots, g_q we have

$$\mathfrak{v} \left(\prod_{k=0}^q g_k \circ T_\omega^k \right) \leq \sum_{k=0}^q \left(\prod_{0 \leq s < k} \|g_s\|_\infty \right) \cdot (\mathfrak{v}(g_k \circ T_\omega^k)) \cdot \left(\prod_{k < s \leq q} \|g_s\|_\infty \right)$$

where $\|g_i\|_\infty = \sup \|g_i\|_{L^\infty}$, and hence

$$(3.8) \quad \left\| \prod_{k=0}^q g_k \circ T_\omega^k \right\|_{BV} \leq \prod_{k=0}^q \|g_k\|_\infty + \sum_{k=0}^q \left(\prod_{0 \leq s < k} \|g_s\|_\infty \right) \cdot \left(\prod_{s=0}^{k-1} K(\sigma^s \omega) \mathfrak{v}(g_k) \right) \left(\prod_{k < s \leq q} \|g_s\|_\infty \right)$$

where we have used (2.3), that $N(\omega) \leq K(\omega)$ and that

$$\left\| \prod_{k=0}^q g_k \circ T_\omega^k \right\|_{L^1} \leq \left\| \prod_{k=0}^q g_k \circ T_\omega^k \right\|_{L^\infty} \leq \prod_{k=0}^q \|g_k\|_\infty.$$

Let us write $B_1 = \{0, 1, \dots, d\}$. Taking $g_k = f_k$ for $0 \leq k \leq d = q$ and noting that $K(\sigma^s \omega) \|g_s\|_\infty \leq C$ for some constant C which depends⁶ only the space X we conclude that

$$\|F_1\|_{BV} \leq C(d+1) \leq 2Cd.$$

Now, if we write $B_2 = \{d+n, d+n+1, \dots, d+n+L\}$ then

$$\mu_\omega(F_1 F_2) = \mu_\omega(F_1 \cdot G_2 \circ T_\omega^{d+n}) = \mu_{\sigma^{n+d}\omega}(G_2 L_\omega^{n+d} F_1)$$

where

$$G_2 = \prod_{u \in B_2} f_u \circ T_{\sigma^u \omega}^{u-n-d}.$$

By (2.14) we have,

$$\|L_\omega^{n+d} F_1 - \mu_\omega(F_1)\|_{BV} \leq K(\omega) \|F_1\|_{BV} e^{-\lambda(d+n)} \leq 2dCK(\omega) e^{-\lambda(d+n)}.$$

Therefore, using also that μ_ω is an equivariant family and that (since $n+d \in B_2$)

$$\|G_2\|_{L^\infty} \leq \|f_{n+d}\|_{L^\infty} \leq CK(\sigma^{n+d}\omega)^{-1}$$

we get that

$$\begin{aligned} & |\mu_\omega(F_1 F_2) - \mu_\omega(F_1) \mu_\omega(F_2)| = |\mu_{\sigma^{n+d}\omega}(G_2 L_\omega^{n+d} F_1) - \mu_\omega(F_1) \mu_{\sigma^{n+d}\omega}(G_2)| \\ &= \left| \int (L_\omega^{d+n} F_1 - \mu_\omega(F_1)) G_2 d\mu_{\sigma^{n+d}\omega} \right| \leq 2dCK(\omega) e^{-\lambda(d+n)} \|G_2\|_{L^\infty} \\ &\leq 2dCK(\omega) e^{-\lambda(d+n)} K(\sigma^{n+d}\omega)^{-1} \leq 2dC^2 e^{-(\lambda-\varepsilon)(d+n)} = (2C^2 d e^{-(\lambda-\varepsilon)d}) e^{-(\lambda-\varepsilon)n}. \end{aligned}$$

This proves the lemma for $m = 2$.

Next, let us complete the induction step. Let d be the right end point of B_{m-1} . Then $d + d_m$ is the left end point of B_m and we can write

$$\mu_\omega\left(\prod_k F_k\right) = \mu_\omega\left(\prod_{k < m} F_k \cdot (G_m \circ T_\omega^{d+d_m})\right) = \mu_{\sigma^{d+d_m}\omega}(L_\omega^{d+d_m}(\prod_{k < m} F_k) \cdot G_m)$$

where G_m is some function. Now we observe that

$$\left\| \prod_{k < m} F_k \right\|_{BV} \leq C(d+1) \leq 2Cd$$

which is proved exactly as in the previous case (even though there are gaps between the blocks B_j , we can set $g_i = 1$ when i does not belong to one of the B_j 's, and then $v(g_i) = 0$). Thus, as in the case $m = 2$, we have

$$\left| \mu_\omega\left(\prod_k F_k\right) - \mu_\omega(F_m) \mu_\omega\left(\prod_{k < m} F_k\right) \right| \leq (2C^2 d e^{-(\lambda-\varepsilon)d}) e^{-(\lambda-\varepsilon)d_m}.$$

The induction is completed by the above inequality, taking into account that $|\mu_\omega(F_m)| \leq 1$. \square

Integrating over ω yields the following corollary of Lemma 3.6.

3.7. Corollary. *Let τ be the skew product. Let $B_j, 1 \leq j \leq m$ be blocks as in Lemma 3.6. Set $G_j = \prod_{i \in B_j} \varphi \circ \tau^i$. Let us denote by b_j the left end point of B_j . Then*

$$(3.9) \quad \left| \int \prod_{j=1}^m G_j d\mu - \int \left(\prod_{j=1}^m \int \left(\prod_{i \in B_j} \varphi_{\sigma^i \omega} \circ T_{\sigma^{b_j} \omega}^{i-b_j} \right) d\mu_{\sigma^{b_j} \omega} \right) d\mathbb{P}(\omega) \right| \leq A \sum_{j=1}^m e^{-\lambda d_j}.$$

⁶ C is a constant which satisfies that $\|g\|_\infty = \sup |g| \leq C \|g\|_{BV}$ for every complex function on X .

The next step of the proof is to estimate the second term inside the absolute value on the left hand side of (3.9). To obtain appropriate estimates, we first need the following lemma:

3.8. Lemma. *Let us fix some $k \in \mathbb{N}$ and set*

$$F_\omega = \prod_{j=0}^k \varphi_{\sigma^k \omega} \circ T_\omega^k.$$

Then for every $n \in \mathbb{N}$ and for \mathbb{P} a.e. ω we have

$$|\mu_\omega(F_\omega) - m(F_\omega \mathcal{L}_{\sigma^{-n}\omega}^n \mathbf{1})| \leq C e^{-n(\lambda-\varepsilon)}$$

where C is such that $\|g\|_{L^\infty} \leq C\|g\|_{BV}$ for every function g on X with bounded variation (recall that such a constant C exists by our assumption on the variation $v(\cdot)$).

Proof. Using (2.10), that $K(\sigma^{-n}\omega) \leq e^{\varepsilon n} K(\omega)$ and that $\|F_\omega\|_{L^\infty} \leq \|\varphi_\omega\|_{L^\infty} \leq C\|\varphi_\omega\|_{BV} \leq CK(\omega)^{-1}$ we obtain that

$$\begin{aligned} & |\mu_\omega(F_\omega) - m(F_\omega \mathcal{L}_{\sigma^{-n}\omega}^n \mathbf{1})| = \\ & \left| \int (h_\omega - \mathcal{L}_{\sigma^{-n}\omega}^n \mathbf{1}) F_\omega dm \right| \leq CK(\omega)^{-1} \int |h_\omega - \mathcal{L}_{\sigma^{-n}\omega}^n \mathbf{1}| dm \\ & \leq K(\omega)^{-1} e^{-\lambda n} K(\sigma^{-n}\omega) \leq C e^{-n(\lambda-\varepsilon)}. \end{aligned}$$

□

Taking into account that $|\mu_\omega(F_\omega)| \leq 1$, that $|m(F_\omega \mathcal{L}_{\sigma^{-n}\omega}^n \mathbf{1})| = |m(F_\omega \circ T_{\sigma^{-n}\omega}^n)| \leq 1$ and that $|\prod_j \alpha_j - \prod_j \beta_j| \leq \sum_j |\alpha_j - \beta_j|$ for all numbers α_j, β_j so that $|\alpha_j|, |\beta_j| \leq 1$ we get the following result directly from Corollary 3.7 and Lemma 3.8.

3.9. Corollary. *Let b_j be the left end point of the block B_j . Let us also set $r_j = d_j/3$ and $r_0 = r_1$. Then there exists a constant $A_1 > 0$ which does not depend on ω or on the blocks so that in the notations of Corollary 3.7 and Lemma 3.8 we have*

$$\left| \int \prod_{j=1}^m G_j d\mu - \int \left(\prod_{j=0}^d m(\varphi_{\omega,j} \mathcal{L}_{\sigma^{b_j-d_j}\omega}^{d_j} \mathbf{1}) \right) d\mathbb{P}(\omega) \right| \leq A_1 \sum_{j=1}^{m-1} e^{-(\lambda-\varepsilon)r_j}$$

where

$$\varphi_{\omega,j} = \prod_{i \in B_j} \varphi_{\sigma^i \omega} \circ T_{\sigma^{b_j}\omega}^{i-b_j}.$$

Now, we observe that $m(\varphi_{\omega,j} \mathcal{L}_{\sigma^{n_j-d_j}\omega}^{d_j} \mathbf{1})$ is a function of $\xi_{b_j-r_j}, \dots, \xi_{b_{j+1}-r_j}$ (i.e. of the coordinates $\omega_{b_j-r_j}, \dots, \omega_{b_{j+1}-r_j}$). Namely, in distribution it can be written as

$$m(\varphi_{\omega,j} \mathcal{L}_{\sigma^{n_j-d_j}\omega}^{d_j} \mathbf{1}) = f_j(\xi_{b_j-r_j}, \dots, \xi_{b_{j+1}-r_j})$$

for some measurable function f_j . Since $m(\varphi_{\omega,j} \mathcal{L}_{\sigma^{n_j-d_j}\omega}^{d_j} \mathbf{1}) = m(\varphi_{\omega,j} \circ T_{\sigma^{n_j-d_j}\omega}^{d_j})$ and $|\varphi_{\omega,j}| \leq 1$, we can insure that $|f_j| \leq 1$. Using [25, (2.20)] and Corollary 3.9 we conclude that:

3.10. Corollary. *Let $G_j, 1 \leq j \leq m$ be as in Corollary 3.7 (defined by some blocks B_j with gaps d_j). There are constants $A > 1$ and $\delta_0 \in (0, 1)$ which do not depend on the blocks so that*

$$\left| \int \left(\prod_{j=1}^m G_j \right) d\mu - \left(\prod_{j=1}^m \int G_j d\mu \right) \right| \leq A \sum_{j=1}^m (\delta_0^{r_j} + \alpha([r_j])).$$

All that is left is to notice that Corollary 3.10 is a reformulation of Proposition 3.4, using the notations of this section.

3.2. Limit theorems via the method of cumulants.

3.2.1. *The CLT: proof of Theorem 2.14.* First, by Proposition 3.4 we have that (3.2) holds true with the numbers $\varrho_{i,(1+\delta)k}$ and $\gamma_\delta(b, k)$ specified in Proposition 3.4. By taking $r = 2$, $A_1 = \{0\}$ and $A_2 = \{n\}$ in (3.2) we see that

$$|\mathbb{E}_\mu[\varphi \cdot \varphi^n]| = O(\delta^n + \alpha_{[n/3]})$$

for some $\delta \in (0, 1)$. Hence, if $\sum n\alpha_n < \infty$ then $\sum_n n|\mathbb{E}_\mu[\varphi \cdot \varphi^n]| < \infty$ and the results concerning the asymptotic variance s^2 follow from the general theory of (weakly) stationary processes (see [33] and Lemma 3.11 below).

Now, suppose that $s^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_\mu(S_n) > 0$, where $S_n = S_n \varphi$. To prove the CLT and the convergence rate (2.18), by applying [46, Corollary 2.1], taking into account Theorem 3.1, we get the CLT and the rate (2.18) for $S_n / \sqrt{\text{Var}(S_n)}$. To get the same rate for S_n / \sqrt{n} we need the following general fact from the theory of stationary real-valued sequences, which for the sake of convenience is stated as a lemma.

3.11. Lemma. *Let Y_n be a centered weakly stationary sequence of square integrable random variables. Set $b_n = \mathbb{E}[Y_0 Y_n]$ and $S_n = \sum_{j=1}^n Y_j$. Suppose that $\sum_k k|b_k| < \infty$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[S_n^2] = b_0 + 2 \sum_{n \geq 1} b_n := s^2$$

and

$$\left| \frac{1}{n} \mathbb{E}[S_n^2] - s^2 \right| \leq 2n^{-1} \sum_{k=1}^{\infty} k|b_k|.$$

Let us give a reminder of the short proof. We have $\frac{1}{n} \mathbb{E}[S_n^2] = \sum_{k=1}^{n-1} (1 - k/n)b_k + b_0$ and so

$$\left| \frac{1}{n} \mathbb{E}[S_n^2] - s^2 \right| = \left| 2 \sum_{k=n}^{\infty} b_k + 2n^{-1} \sum_{k=1}^{n-1} kb_k \right| \leq 2n^{-1} \left(\sum_{k=n}^{\infty} k|b_k| + \sum_{k=1}^{n-1} k|b_k| \right) \leq 2n^{-1} \sum_{k \geq 1} k|b_k|.$$

Using this lemma together with [26, Lemma 3.3] with $a = 2$ and that

$$\left\| \frac{S_n}{\sqrt{\text{Var}(S_n)}} - \frac{S_n}{s\sqrt{n}} \right\|_{L^2} = \|S_n\|_{L^2} \left| \frac{1}{\sqrt{\text{Var}(S_n)}} - \frac{1}{s\sqrt{n}} \right| = O(n^{1/2}) \cdot O(n^{-3/2}) = O(n^{-1})$$

we obtain (2.18).

3.2.2. *A moderate deviations principle, stretched exponential concentration inequalities and Rosenthal type estimates: proof of Theorems 2.15, 2.16 and 2.17.*

First, Theorem 2.15 follows from Theorem 3.1 and [46, Lemma 2.3]. The estimates (2.20) stated in Theorem 2.16 follow from Theorem 3.1 and [15, Lemma 2.3] (which is a consequence of [46, Lemma 2.3]). The moderate deviations principle stated in Theorem 2.16 follows from Theorem 3.1 and [15, Theorem 1.1]. We note that the conditions of [46, Lemma 2.3], [15, Lemma 2.3] and [15, Theorem 1.1] are certain estimates on the growth rates (in k) of the cumulants $\Gamma_k(S_n)$, and the role of Theorem 3.1 is that it shows that the conditions of all of these results are in force in the setup of this paper.

3.3. A functional CLT via the method of cumulants: proof of Theorem 2.19. Let us first show that the sequence \mathcal{S}_n is tight. By Theorem 2.17 we have that

$$\|S_n\|_4 = O(\sqrt{n})$$

where $\|\cdot\|_4 = \|\cdot\|_{L^4}$, and therefore, using also stationarity and the Hölder inequality we get that for all $t_1 < t_2 \leq r_1 < r_2$,

$$\begin{aligned} \mathbb{E} \left[(\mathcal{S}_n(r_2) - \mathcal{S}_n(r_1))^2 (\mathcal{S}_n(t_2) - \mathcal{S}_n(t_1))^2 \right] &\leq \|\mathcal{S}_n(r_2) - \mathcal{S}_n(r_1)\|_4^2 \|\mathcal{S}_n(t_2) - \mathcal{S}_n(t_1)\|_4^2 \\ &\leq C \left(\frac{[r_2 n] - [t_1 n]}{n} \right)^2. \end{aligned}$$

Thus, by [10, Ch.15], $\mathcal{S}_n(\cdot)$ is a tight sequence in the Skorokhod space $D[0, 1]$.

Now let us show that the finite-dimensional distributions converge. Let us fix some $t_1 < t_2 < \dots < t_d$. Set $X_k = \varphi \circ \tau^k$. Next, let us recall the following general fact. Given a vector valued sequence of random variables $Y_n = (Y_{1,n}, \dots, Y_{d,n})$, by the multidimensional version of Levi's theorem, in order to show that Y_n converges in distribution as $n \rightarrow \infty$ towards a given random variable \mathcal{Z} , it is enough to show that for every $a \in \mathbb{R}^d$ we have

$$\lim_{n \rightarrow \infty} \mathbb{E}[e^{i(a \cdot Y_n)}] = \mathbb{E}[e^{i(a \cdot \mathcal{Z})}].$$

Therefore, it is enough to show that any linear combination of $Y_{j,n}$, $j = 1, 2, \dots, d$ converges in distribution towards the corresponding linear combination of the coordinates of \mathcal{Z} . Returning to our problem, to obtain the appropriate convergence of the distribution of $(\mathcal{S}_n(it_j))_{j=1}^d$ it is enough to show that any linear combination of $\mathcal{S}_n(t_j)$ converges towards a centered normal random variable with an appropriate variance. More precisely, let $a_1, \dots, a_d \in \mathbb{R}$. Then we need to show that $\sum_{j=1}^d a_j \mathcal{S}_n(t_j)$ converges in distribution towards a centered normal random variable with variance

$$s^2 \sum_{j=1}^d (a_j + \dots + a_d)^2 (t_j - t_{j-1})$$

where $t_0 = 0$ and $s^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[S_n^2]$. We first notice that

$$\sum_{j=1}^d a_j \mathcal{S}_n(t_j) = n^{-1/2} \sum_{j=1}^d (a_j + \dots + a_d) (S_{[nt_j]} - S_{[nt_{j-1}]})$$

where we set $t_0 = 0$ and $S_0 = 0$. Thus, using stationarity, we have

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{j=1}^d a_j \mathcal{S}_n(t_j) \right)^2 \right] &= n^{-1} \sum_{j=1}^d (a_j + \dots + a_d)^2 \mathbb{E}[S_{[nt_j] - [nt_{j-1}]}^2] \\ &+ 2n^{-1} \sum_{1 \leq j_1 < j_2 \leq d} (a_{j_1} + \dots + a_{j_d})(a_{j_2} + \dots + a_{j_d}) \mathbb{E} \left[(S_{[nt_{j_2}]} - S_{[nt_{j_2-1}]}) (S_{[nt_{j_1}]} - S_{[nt_{j_1-1}]}) \right]. \end{aligned}$$

Now, the first summand on the above right hand side converges to

$$s^2 \sum_{j=1}^d (a_j + \dots + a_d)^2 (t_j - t_{j-1}),$$

while the second summand (the double sum) converges to 0 because $|\mathbb{E}[\varphi \cdot \varphi \circ \tau^n]|$ converges to 0 stretched exponentially fast. Therefore, the asymptotic variance of $\sum_{j=1}^d a_j \mathcal{S}_n(t_j)$ has the desired form. Now, let us consider the following array of random variables. Set

$$Y_k = Y_k^{(n, a_1, \dots, a_d, t_1, \dots, t_d)} = (a_1 + \dots + a_j) \varphi \circ \tau^k \text{ if } [nt_{j-1}] \leq k < [nt_j].$$

Then,

$$\begin{aligned} \sum_{j=1}^d (a_j + \dots + a_d) (S_{[nt_j]} - S_{[nt_{j-1}]}) &= \sum_{j=1}^d (a_j + \dots + a_d) \sum_{s=[nt_{j-1}]}^{[nt_j]-1} \varphi \circ \tau^s \\ &= \sum_{j=1}^d (a_j + \dots + a_d) \sum_{s=0}^{[nt_d]-1} \mathbb{I}([nt_{j-1}] \leq s < [nt_j]) \varphi \circ \tau^s = \sum_{s=0}^{[nt_d]-1} Y_s. \end{aligned}$$

On the other other hand, arguing as in the proof of Theorem 3.1 (replacing each appearance of $\varphi \circ \tau^k$ by Y_k) we get the same kind of estimates on the cumulants of

$$\tilde{S}_n := \sum_{s=0}^{[nt_d]-1} Y_s,$$

that is, there exists a constant c_0 which might depend on t_j and a_j so that for every k we have

$$|\Gamma_k(\tilde{S}_n)| \leq n(k!)^{1+\gamma} (c_0)^{k-2}.$$

Thus, by applying [46, Corollary 2.1] we get that

$$\sum_{s=0}^{[nt_d]-1} Y_s^{(n, a_1, \dots, a_d)} / w_n$$

converges towards the standard normal distribution, where w_n is the standard deviation of the numerator. Note that, as we have shown, $w_n^2/n \rightarrow s^2 \sum_{j=1}^d (a_1 + \dots + a_d)^2 (t_j - t_{j-1})$, which is positive unless either $s = 0$ or $a_1 = \dots = a_d = 0$, which are both trivial cases. Thus, in any case we obtain the desired convergence of the linear combination $\sum_{j=1}^d a_j \mathcal{S}_n(t_j)$ and the proof of Theorem 2.19 is complete.

4. LIMIT THEOREMS VIA MARTINGALE APPROXIMATION FOR ϕ AND ψ MIXING DRIVING PROCESSES

4.1. Some expectation estimates using mixing coefficients. In the course of the proof of Theorem 2.22 we will need the following two relatively simple lemmas.

4.1. Lemma. *Let \mathcal{G}, \mathcal{H} be two sub- σ -algebras of a given σ -algebra on some space measure space. Let g be a real-valued bounded \mathcal{G} -measurable function and h be an \mathcal{H} -measurable real-valued integrable function. Then*

$$|\mathbb{E}[hg] - \mathbb{E}[h]\mathbb{E}[g]| \leq \frac{1}{2} \|h\|_{L^\infty} \|g\|_{L^1} \phi(\mathcal{G}, \mathcal{H})$$

Proof. By [11, Ch. 4] we have

$$\|\mathbb{E}[h|\mathcal{G}] - \mathbb{E}[h]\|_{L^\infty} \leq \frac{1}{2} \|h\|_{L^\infty} \phi(\mathcal{G}, \mathcal{H})$$

which clearly implies the lemma. \square

The next result is:

4.2. Lemma. *Let \mathcal{G}, \mathcal{H} be two sub- σ -algebras of a given σ -algebra on some space measure space. Let g a real-valued bounded \mathcal{G} -measurable function and h be an \mathcal{H} -measurable real-valued integrable function. Suppose also that $\psi = \psi(\mathcal{G}, \mathcal{H}) < 1$. Then*

$$|\mathbb{E}[hg] - \mathbb{E}[h]\mathbb{E}[g]| \leq 4 \|hg\|_{L^1} C_\psi \psi$$

where $C_\psi = (1 - \psi)^{-1}$.

Proof. By [11, Ch.4] we have,

$$\|\mathbb{E}[h|\mathcal{G}] - \mathbb{E}[h]\|_{L^\infty} \leq \|h\|_{L^1} \psi(\mathcal{G}, \mathcal{H}).$$

Hence

$$|\mathbb{E}[hg] - \mathbb{E}[h]\mathbb{E}[g]| \leq \|h\|_{L^1} \|g\|_{L^1} \psi.$$

Taking $h, g \geq 0$ we get that

$$|\mathbb{E}[hg] - \mathbb{E}[h]\mathbb{E}[g]| \leq \mathbb{E}[h]\mathbb{E}[g]\psi.$$

Thus,

$$\mathbb{E}[h]\mathbb{E}[g] \leq (1 - \psi)^{-1} \mathbb{E}[hg] = C_\psi \mathbb{E}[hg].$$

Therefore, for nonnegative functions we have

$$|\mathbb{E}[hg] - \mathbb{E}[h]\mathbb{E}[g]| \leq C_\psi \psi \mathbb{E}[hg].$$

Now the general result follows by writing $h = h^+ - h^-$ and $g = g^+ - g^-$ where h^\pm and g^\pm are nonnegative functions so that $h^+ + h^- = |h|$ and $g^+ + g^- = |g|$, and using that both $(g, h) \rightarrow \mathbb{E}[g]\mathbb{E}[h]$ and $(g, h) \rightarrow \mathbb{E}[hg]$ are bilinear in (g, h) . \square

4.2. Convergence of the iterates of the transfer operator with respect to a sub- σ -algebra.

Let \mathcal{F}_0 be the σ -algebra generated by the map $\pi(\omega, x) = ((\omega_j)_{j \geq 0}, x)$, namely the one generated by \mathcal{B} and the coordinates with non-negative indexes in the ω direction. Then $(\tau^{-k}\mathcal{F}_0)_{k \geq 0}$ is a decreasing sequence of σ -algebras and $\tau^{-k}\mathcal{F}_0$ is generated by τ^k and the coordinates ω_j for $j \geq k$. In particular τ preserves \mathcal{F}_0 .

Next, let us define a transfer operator with respect to \mathcal{F}_0 . For each function $g \in L^1(\mu)$ there is a unique \mathcal{F}_0 -measurable function G so that

$$\mathbb{E}[g|\tau^{-1}\mathcal{F}_0] = G \circ \tau.$$

Let us define $\mathcal{K}g = G$, where we formally set G to be 0 outside the image of τ (if τ is not onto). Then

$$\mathbb{E}[g|\tau^{-1}\mathcal{F}_0] = \mathcal{K}g \circ \tau.$$

Notice that for $g \in L^1(\Omega \times X, \mathcal{F}_0, \mu)$, $f \in L^\infty(\Omega \times X, \mathcal{F}_0, \mu)$ we have

$$\int (\mathcal{K}g)f d\mu = \int (\mathcal{K}g \circ \tau)f \circ \tau d\mu = \int \mathbb{E}[g|\tau^{-1}\mathcal{F}_0] \cdot f \circ \tau d\mu = \int g \cdot f \circ \tau d\mu$$

and therefore \mathcal{K} can also be defined using the usual duality relation. That is, it is the transfer operator of τ with respect to $(\Omega \times X, \mathcal{F}_0, \mu)$.

The proof of Theorems 2.21 and 2.22 is based on the following result.

4.3. Proposition. *Under the assumptions of Theorems 2.21 and 2.22, and when $\mu(\varphi) = 0$ we have the following:*

(i) *We have*

$$(4.1) \quad \|\mathcal{K}^n \varphi\|_{L^\infty} \leq C(e^{-(\lambda-2\varepsilon)n/2} + \psi_{[n/2]}) := C\gamma_{2,n}.$$

Moreover, if $h_\omega \geq c^{-1} > 0$ for some constant $c > 1$ then

$$(4.2) \quad \|\mathcal{K}^n \varphi\|_{L^\infty} \leq Cc(e^{-(\lambda-2\varepsilon)n/2} + \phi_{[n/2],R}) := C\gamma_{1,n}.$$

Here $C = C_\varphi$ is a constant having the form $C_\varphi = AC_v \text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi_\omega\|_{BV})$ where A is an absolute constant and C_0 is any constant satisfying $\|g\|_{L^\infty} \leq C_0 \|g\|_{BV}$ and $\|fg\|_{BV} \leq C_0 \|g\|_{BV} \|f\|_{BV}$ for all functions $g, f : X \rightarrow \mathbb{C}$.

(ii) *We have*

$$\|\mathcal{K}^i(\varphi \mathcal{K}^j \varphi) - \mu(\mathcal{K}^i(\varphi \mathcal{K}^j \varphi))\|_{L^\infty} \leq C\gamma_{2, \max(i,j)}.$$

If $h_\omega \geq c^{-1} > 0$ for some constant $c > 1$ then

$$\|\mathcal{K}^i(\varphi \mathcal{K}^j \varphi) - \mu(\mathcal{K}^i(\varphi \mathcal{K}^j \varphi))\|_{L^\infty} \leq Cc\gamma_{1, \max(i,j)}.$$

Proof of Theorems 2.21 and 2.22 based on Proposition 4.3. First, theorem 2.21 (i) follows since if we set $\chi = \sum_{n=1}^\infty \mathcal{K}^n \varphi$ and $u = \varphi + \chi \circ \tau - \chi$, then $\|\chi\|_{L^\infty} < \infty$ and $(u \circ \tau^n)$ is a reverse martingale difference with respect to the reverse filtration $\{\tau^{-n}\mathcal{F}_0\}$. Moreover, the differences $u \circ \tau^n$ are uniformly bounded (as χ and φ are in L^∞). Thus by the Azuma-Hoeffding inequality for every $\beta > 0$ we have,

$$\mathbb{E}_\mu[e^{\lambda \sum_{j=0}^{n-1} u \circ \tau^j}] \leq e^{\beta^2 n \|u\|_{L^\infty}^2}.$$

Now the proof proceeds by using the Chernoff bounding method: by the Markov inequality for all $t > 0$ we have

$$\mu \left\{ \sum_{j=0}^{n-1} u \circ \tau^j \geq tn \right\} \leq e^{-\beta tn} e^{\beta^2 n \|u\|_{L^\infty}^2}.$$

Taking $\beta = \beta_t = \frac{t}{2\|u\|_{L^\infty}}$ and replacing u with $-u$ we get that

$$\mu \left\{ \pm \sum_{j=0}^{n-1} u \circ \tau^j \geq tn \right\} \leq e^{-\frac{nt^2}{4\|u\|_{L^\infty}^2}}.$$

The proof of Theorem 2.21 (i) is completed now by noticing that

$$(4.3) \quad \left\| S_n \varphi - \sum_{j=0}^{n-1} u \circ \tau^j \right\|_{L^\infty} = \|\chi - \chi \circ \tau^n\|_{L^\infty} \leq 2\|\chi\|_{L^\infty}.$$

Next, the proof of Theorem 2.21 (ii) is completed by applying [43, Proposition 7] with the reverse martingale $(u \circ \tau^n)$ and using (4.3).

In order to prove Theorem 2.22, we apply [13, Theorem 3.2] with the bounded function φ and the probability preserving system $(\Omega \times X, \mathcal{F}_0, \mu, \tau)$, whose transfer operator is \mathcal{K} . Now, since we have assumed that $\mu(\varphi) = 0$, in order for the conditions of [13, Theorem 3.2] to be in force we need the following three estimates to hold

$$\sum_{n \geq 2} n^{5/2} (\log n)^3 \|\mathcal{K}\varphi\|_{L^4(\mu)}^4 < \infty \quad \text{and} \quad \sum_{n \geq 2} n (\log n)^3 \|\mathcal{K}\varphi\|_{L^2(\mu)}^2 < \infty$$

and

$$\sum_{n \geq 2} \frac{(\log n)^3}{n^2} \left(\sum_{i=1}^n \sum_{j=0}^{n-i} \|\mathcal{K}^i(\varphi \mathcal{K}^j(\varphi)) - \mu(\varphi \mathcal{K}^j(\varphi))\|_{L^2(\mu)} \right)^2 < \infty.$$

The above three conditions are verified by Proposition 4.3 and the mixing rates specified in the formulation of Theorem 2.22, and the proof of Theorem 2.22 is complete. \square

Proof of Proposition 4.3. (i) Since $L^\infty(\mu)$ is the dual of $L^1(\mu)$ and φ and $\mathcal{K}^n \varphi$ are \mathcal{F}_0 -measurable, it is enough to show that for every $g \in L^1(\Omega \times X, \mathcal{F}_0, \mu)$ so that $\|g\|_{L^1} \leq 1$ we have

$$\left| \int g \cdot (\mathcal{K}^n \varphi) d\mu \right| \leq \gamma_n \|g\|_{L^1(\mu)}$$

where γ_n is one of the desired upper bounds. To achieve that let us first note that \mathcal{K}^n is the dual of the restriction of the Koopman operator $f \rightarrow f \circ \tau^n$ acting on \mathcal{F}_0 -measurable functions. Thus,

$$(4.4) \quad \begin{aligned} \int g \cdot (\mathcal{K}^n \varphi) d\mu &= \int \varphi \cdot (g \circ \tau^n) d\mu = \int \left(\int \varphi_\omega \cdot (g_{\sigma^n \omega} \circ T_\omega^n) d\mu_\omega \right) d\mathbb{P}(\omega) \\ &= \int \left(\int (L_\omega^n \varphi_\omega) \cdot g_{\sigma^n \omega} d\mu_{\sigma^n \omega} \right) d\mathbb{P}(\omega). \end{aligned}$$

Now, using (2.14) and that $\|\varphi\|_K = \text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) < \infty$ we get that

$$\|L_\omega^n \varphi_\omega - \mu_\omega(\varphi_\omega)\|_{L^\infty} \leq C_0 \|\varphi\|_K e^{-\lambda n}.$$

Hence, using also the σ -invariance of \mathbb{P} ,

$$\int g \cdot (\mathcal{K}^n \varphi) d\mu = \int \mu_\omega(\varphi_\omega) \mu_{\sigma^n \omega}(g_{\sigma^n \omega}) d\mathbb{P}(\omega) + I$$

where $|I| \leq C e^{-\lambda n} \|g\|_{L^1(\mu)}$. Next, let us write

$$\mu_{\sigma^n \omega}(g_{\sigma^n \omega}) = m(g_{\sigma^n \omega} h_{\sigma^n \omega}).$$

By (2.10) we have

$$\left\| h_{\sigma^n \omega} - \mathcal{L}_{\sigma^{[n/2]} \omega}^{n - [n/2]} \mathbf{1} \right\|_{L^\infty} \leq C_0 K(\sigma^{[n/2]} \omega) e^{-\lambda n/2} \leq C_0 K(\omega) e^{-(\lambda - \varepsilon)n/2}.$$

Observe next that since $\|1/h_\omega\|_{BV} \leq K(\omega)$ we have

$$m(|g|) = \mu_{\sigma^n \omega}(|g|/h_{\sigma^n \omega}) \leq C_0 K(\sigma^n \omega) \mu_\omega(|g|)$$

for every function g and recall that $K(\sigma^n \omega) \leq K(\omega) e^{\varepsilon n}$. Combining this with the previous estimates we get that

$$(4.5) \quad \begin{aligned} \left| m(g_{\sigma^n \omega} h_{\sigma^n \omega}) - m(g_{\sigma^n \omega} \mathcal{L}_{\sigma^{[n/2]} \omega}^{n - [n/2]} \mathbf{1}) \right| C_0 &\leq K(\omega) e^{-(\lambda - \varepsilon)n/2} m(|g_{\sigma^n \omega}|) \\ &\leq C K(\omega)^2 \mu_{\sigma^n \omega}(|g_{\sigma^n \omega}|) e^{-(\lambda - 3\varepsilon)n/2}. \end{aligned}$$

Therefore,

$$(4.6) \quad \int g \cdot (\mathcal{K}^n \varphi) d\mu = \int \mu_\omega(\varphi_\omega) m(g_{\sigma^n \omega} \mathcal{L}_{\sigma^{[n/2]} \omega}^{n-[n/2]} \mathbf{1}) d\mathbb{P}(\omega) + I + J$$

where $|I| \leq C e^{-\lambda n} \|g\|_{L^1(\mu)}$ and $|J| \leq C' e^{-(\lambda-3\varepsilon)n/2} \|g\|_{L^1(\mu)}$ and we have used that $K(\omega)^2 \|\varphi_\omega\|_{BV}$ is bounded.

Next, using (2.10) and that $K(\omega)$ is tempered we have $h_\omega = \lim_{n \rightarrow \infty} \mathcal{L}_{\sigma^{-n} \omega}^n \mathbf{1}$, and therefore h_ω depends only on the coordinates ω_j for $j \leq 0$. Thus

$$\mu_\omega(\varphi_\omega) = F(\omega_j; j \leq 0)$$

for some measurable function F so that $|F| \leq \|\varphi\|_{L^1(\mu)}$. Observe also that the random variable

$$G_n(\omega) = m(g_{\sigma^n \omega} \mathcal{L}_{\sigma^{[n/2]} \omega}^{n-[n/2]} \mathbf{1})$$

depends only on $\omega_j, j \geq [n/2]$ since $g_\omega(x)$ is a function of x and $\omega_j, j \geq 0$ (i.e. it factors through π_0). In the case when $h_\omega \geq c^{-1} > 0$ for some constant $c > 0$ we have

$$|G_n(\omega)| = \left| \mu_{\sigma^n \omega} (g_{\sigma^n \omega} \mathcal{L}_{\sigma^{[n/2]} \omega}^{n-[n/2]} (1/h_{\sigma^{[n/2]} \omega})) \right| \leq c \mu_{\sigma^n \omega} (|g_{\sigma^n \omega}|).$$

Thus, using also lemma 4.1 we see that there is a constant $C > 0$ so that

$$\left| \int \mu_\omega(\varphi_\omega) m(g_{\sigma^n \omega} \mathcal{L}_{\sigma^{[n/2]} \omega}^{n-[n/2]} \mathbf{1}) d\mathbb{P}(\omega) \right| \leq C \phi_{[n/2], R} \int |G_n(\omega)| d\mathbb{P}(\omega) \leq c C \phi_{[n/2], R} \|g\|_{L^1(\mu)}$$

where we have taken into account that $\int \mu_\omega(\varphi_\omega) d\mathbb{P}(\omega) = \mu(\varphi) = 0$. This, together with (4.6) and the previous estimates on I and J , proves (4.2).

To prove (4.1), we first use (4.5) in order to obtain that

$$(4.7) \quad |G_n(\omega)| \leq C \mu_{\sigma^n \omega} (|g_{\sigma^n \omega}|) (1 + CK^2(\omega) e^{-(\lambda-3\varepsilon)n/2}) \leq C' \mu_{\sigma^n \omega} (|g_{\sigma^n \omega}|) K(\omega)^2.$$

Taking into account that

$$\text{esssup}_{\omega \in \Omega} (\|\varphi_\omega\|_{L^\infty} K(\omega)^2) \leq C \text{esssup}_{\omega \in \Omega} (\|\varphi_\omega\|_{BV} K(\omega)^2) < \infty$$

we conclude that $G_n(\omega) \mu_\omega(\varphi_\omega)$ is integrable. Now, we would like to apply Lemma 4.2, but the problem is that G_n is not bounded. To overcome that, for each $M > 0$ set $G_n^{(M)}(\omega) = G_n(\omega) \mathbb{I}(|G_n(\omega)| \leq M)$. Then, since $G_n(\omega) \mu_\omega(\varphi_\omega)$ is integrable, by the dominated convergence theorem we have

$$\int \mu_\omega(\varphi_\omega) G_n(\omega) d\mathbb{P}(\omega) = \lim_{M \rightarrow \infty} \int \mu_\omega(\varphi_\omega) G_n^{(M)}(\omega) d\mathbb{P}(\omega).$$

Now, taking n so that $\psi_{[n/2]} \leq 1/2$ and using that $\mu(\varphi) = 0$ we get from Lemma 4.2 that

$$\begin{aligned} \left| \int \mu_\omega(\varphi_\omega) G_n^{(M)}(\omega) d\mathbb{P}(\omega) \right| &\leq 2 \left(\int |G_n^{(M)}(\omega) \mu_\omega(\varphi_\omega)| d\mathbb{P}(\omega) \right) \psi_{[n/2]} \leq \\ &2 \left(\int |G_n(\omega) \mu_\omega(\varphi_\omega)| d\mathbb{P}(\omega) \right) \psi_{[n/2]}. \end{aligned}$$

Using also (4.7) and that $\text{esssup}_{\omega \in \Omega} (\|\varphi_\omega\|_{BV} K(\omega)^2) < \infty$, we conclude that

$$\left| \int \mu_\omega(\varphi_\omega) G_n(\omega) d\mathbb{P}(\omega) \right| \leq 2 (\text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi_\omega\|_{BV})) C' \|g\|_{L^1} \psi_{[n/2]}$$

and (4.1) follows (using also (4.6)).

(ii) First, since \mathcal{K} weakly contracts the L^∞ norm (being defined through conditional expectations) and φ is bounded we have

$$\|\mathcal{K}^i(\varphi \mathcal{K}^j \varphi) - \mu(\mathcal{K}^i(\varphi \mathcal{K}^j \varphi))\|_{L^\infty} \leq 2 \|\varphi\|_{L^\infty} \|\mathcal{K}^j \varphi\|_{L^\infty}.$$

This together with Proposition 4.3 (i) provides the desired estimate when $j \geq i$. The estimate in the case $i > j$ is carried out similarly to the proof of (i). Let $g \in L^1(\Omega \times X, \mu, \mathcal{F}_0)$. Let us first show that

$$(4.8) \quad \int \mathcal{K}^i(\varphi \mathcal{K}^j \varphi) g d\mu = \int \mu_\omega(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)) \mu_{\sigma^{i+j} \omega}(g_{\sigma^{i+j} \omega}) d\mathbb{P}(\omega) + I$$

where $|I| \leq C_2 e^{-\lambda i}$, and C_2 is some constant.

In order to prove (4.8), using that \mathcal{K} satisfies the duality relation and the disintegration $\mu = \int \mu_\omega d\mathbb{P}(\omega)$ we first have

$$\begin{aligned}
\int \mathcal{K}^i(\varphi \mathcal{K}^j \varphi) g d\mu &= \int (\varphi \mathcal{K}^j \varphi) \cdot g \circ \tau^i d\mu = \int \mathcal{K}^j \varphi \cdot (\varphi \cdot (g \circ \tau^i)) d\mu = \int (\varphi \cdot (\varphi \circ \tau^j)) \cdot g \circ \tau^{i+j} d\mu \\
(4.9) \qquad &= \int \left(\int \varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j) \cdot (g_{\sigma^{i+j} \omega} \circ T_\omega^{i+j}) d\mu_\omega \right) d\mathbb{P}(\omega) \\
&= \int \left(\int L_\omega^{i+j}(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)) g_{\sigma^{i+j} \omega} d\mu_{\sigma^{i+j} \omega} \right) d\mathbb{P}(\omega).
\end{aligned}$$

Next, since $L_\omega^n(f \circ T_\omega^n) = f$ for every function f and n , we have

$$L_\omega^{i+j}(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)) = L_{\sigma^j \omega}^i(\varphi_{\sigma^j \omega} L_\omega^j \varphi_\omega).$$

By (2.14) we have

$$\|L_\omega^j \varphi_\omega - \mu_\omega(\varphi_\omega)\|_{BV} \leq K(\omega) \|\varphi_\omega\|_{BV} e^{-\lambda j}.$$

In particular,

$$\|L_\omega^j \varphi_\omega\|_{BV} \leq CK(\omega) \|\varphi_\omega\|_{BV}$$

for some constant C . Since $\|uv\|_{BV} \leq C_0 \|u\|_{BV} \|v\|_{BV}$ for every two functions u, v we have

$$\|\varphi_{\sigma^j \omega} L_\omega^j \varphi_\omega\|_{BV} \leq C_0 CK(\omega) \|\varphi_\omega\|_{BV} \|\varphi_{\sigma^j \omega}\|_{BV}.$$

Thus by (2.14),

$$\begin{aligned}
&\|L_{\sigma^j \omega}^i(\varphi_{\sigma^j \omega} L_\omega^j \varphi_\omega) - \mu_{\sigma^j \omega}(\varphi_{\sigma^j \omega} L_\omega^j \varphi_\omega)\|_{BV} \\
&\leq C_0 CK(\omega) K(\sigma^j \omega) \|\varphi_\omega\|_{BV} \|\varphi_{\sigma^j \omega}\|_{BV} e^{-\lambda i} \leq C_0 C \|\varphi\|_K^2 e^{-\lambda i},
\end{aligned}$$

where $\|\varphi\|_K = \text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV})$. Observe next that

$$\mu_{\sigma^j \omega}(\varphi_{\sigma^j \omega} L_\omega^j \varphi_\omega) = \mu_\omega(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)).$$

The desired inequality (4.8) follows from the above estimates.

Observe that the function $\mu_\omega(\varphi_\omega \cdot \varphi_{\sigma^j \omega} \circ T_\omega^j)$ depends only on ω_k for $k \leq j$ and that it is bounded by $CK^{-2}(\omega)$ for some constant $C > 0$ (since $\text{esssup}_{\omega \in \Omega} (K(\omega)^2 \|\varphi_\omega\|_{BV}) < \infty$). Therefore, the same arguments in the proof of (i) yield that

$$\int \mu_\omega(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)) \mu_{\sigma^{i+j} \omega}(g_{\sigma^{i+j} \omega}) d\mathbb{P}(\omega) = \int \mu_\omega(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)) d\mathbb{P}(\omega) \cdot \int \mu_{\sigma^{i+j} \omega}(g_{\sigma^{i+j} \omega}) d\mathbb{P}(\omega) + J$$

where $|J| \leq \gamma_i \|g\|_{L^1}$ and γ_i is one of the right hand sides on the upper bounds in (i) (depending on the case) with n replaced by i . Notice next that

$$\int \mu_\omega(\varphi_\omega \cdot (\varphi_{\sigma^j \omega} \circ T_\omega^j)) d\mathbb{P}(\omega) = \int \mathcal{K}^i(\varphi \mathcal{K}^j \varphi) d\mu$$

(this can be seen by taking $g = 1$ in (4.9)). Hence,

$$\left| \int (\mathcal{K}^i(\varphi \mathcal{K}^j \varphi) - \mu(\mathcal{K}^i(\varphi \mathcal{K}^j \varphi)) g d\mu \right| \leq C(e^{-\lambda i} + \gamma_i) \|g\|_{L^1}$$

and the desired estimate follows again since L^∞ is the dual of L^1 . \square

5. A VECTOR VALUED ASIP FOR SKEW PRODUCTS WITH UNIFORMLY EXPANDING FIBER MAPS AND EXPONENTIALLY FAST α -MIXING BASE MAPS

Let us first explain why the matrix Σ^2 exists. For a fixed vector v the limit $s_v^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[(S_n \cdot v)^2]$ exists, by considering the real-valued observable $\varphi \cdot v$. Then the matrix Σ^2 from Theorem 2.25 is given by $(\Sigma^2)_{i,j} = \frac{1}{2}(s_{e_i+e_j}^2 - s_{e_i}^2 - s_{e_j}^2)$. This matrix satisfies $\Sigma^2 v \cdot v = s_v^2$ and so it is not positive definite if and only if $\varphi \cdot v$ is a coboundary for some unit vector v . Note that this part does not require T_ω to be uniformly expanding.

We assume next that there exist constants $C > 0$ and $\delta \in (0, 1)$ so that for \mathbb{P} a.e. ω we have

$$(5.1) \quad \|\mathcal{L}_\omega^n \mathbf{1} - h_{\sigma^n \omega}\|_{BV} \leq C\delta^n$$

(this is the uniform expansion assumption).

The proof of Theorem 2.25 relies on an application of [24, Theorem 1.2]. The main condition of [24, Theorem 1.2] is the content of the following lemma. Once the lemma is proven Theorem 2.25 follows from [24, Theorem 1.2] applied with an arbitrary large p .

5.1. Lemma. *There exists $\varepsilon_0 > 0$, $c, C > 0$ such that for any $n, m > 0$, $b_1 < b_2 < \dots < b_{n+m+1}$, $k > 0$ and $t_1, \dots, t_{n+m} \in \mathbb{R}^d$ with $|t_j| \leq \varepsilon_0$ we have*

$$(5.2) \quad \left| \mathbb{E}_\mu \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} B_\ell) + i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} B_\ell)} \right) - \mathbb{E}_\mu \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} B_\ell)} \right) \cdot \mathbb{E}_\mu \left(e^{i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} B_\ell)} \right) \right| \leq C^{n+m} e^{-ck},$$

where $B_\ell = \varphi \circ \tau^\ell$.

Proof. First, denoting by \mathbb{E}_ω the expectation with respect to μ_ω , by [20, Lemma 24] there are $\varepsilon_0 > 0$, $c, C > 0$ with the property that for every $n, m > 0$, $b_1 < b_2 < \dots < b_{n+m+1}$, $k > 0$ and $t_1, \dots, t_{n+m} \in \mathbb{R}^d$ such that $|t_j| \leq \varepsilon_0$,

$$(5.3) \quad \left| \mathbb{E}_\omega \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} A_\ell) + i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} A_\ell)} \right) - \mathbb{E}_\omega \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} A_\ell)} \right) \cdot \mathbb{E}_\omega \left(e^{i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} A_\ell)} \right) \right| \leq C^{n+m} e^{-ck},$$

where $\mathbb{E}_\omega(g) = \int gh_\omega dm$ and

$$A_\ell := \varphi_{\sigma^\ell \omega} \circ T_\omega^\ell, \quad \ell \in \mathbb{N}.$$

Let

$$G(\omega) = \mathbb{E}_\omega \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} A_\ell)} \right)$$

and

$$F(\omega) = \mathbb{E}_\omega \left(e^{i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} A_\ell)} \right).$$

Then with $B_\ell = \varphi \circ \tau^\ell$ we have

$$(5.4) \quad \left| \mathbb{E}_\mu \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} B_\ell) + i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} B_\ell)} \right) - \mathbb{E}_\mu \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} B_\ell)} \right) \cdot \mathbb{E}_\mu \left(e^{i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} B_\ell)} \right) \right| \leq C^{n+m} e^{-ck} + |\text{Cov}_{\mathbb{P}}(G, F)|.$$

Using (5.1) and that $(T_\omega)_* \mu_\omega = \mu_{\sigma \omega}$ we get that there are $k_0 \in \mathbb{Z}$ and functions G_1 and F_1 so that

$$\|G(\omega) - G_1(\dots, \omega_{k_0-1}, \omega_{k_0+[k/4]})\|_{L^\infty} \leq C' \delta^{k/4}$$

and

$$\|G(\omega) - G_1(\omega_{k_0+k-[k/4]}, \omega_{k_0+k-[k/4]+1}, \dots)\|_{L^\infty} \leq C' \delta^{k/4}.$$

Thus,

$$|\text{Cov}_{\mathbb{P}}(G, F)| \leq |\text{Cov}_{\mathbb{P}}(G_1, F_1)| + C'' \delta^{k/4}$$

where we have used that G_1, G_2, G and F are uniformly bounded (so the above constants C', C'' do not depend on the choice of b_j, t_j etc.). On the other hand, by (3.7),

$$|\text{Cov}_{\mathbb{P}}(G_1, F_1)| \leq C''' \alpha^{k/2}.$$

Thus,

$$(5.5) \quad \left| \mathbb{E}_{\mu} \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} B_{\ell}) + i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} B_{\ell})} \right) - \mathbb{E}_{\mu} \left(e^{i \sum_{j=1}^n t_j \cdot (\sum_{\ell=b_j}^{b_{j+1}-1} B_{\ell})} \right) \cdot \mathbb{E}_{\mu} \left(e^{i \sum_{j=n+1}^{n+m} t_j \cdot (\sum_{\ell=b_j+k}^{b_{j+1}+k-1} B_{\ell})} \right) \right| \leq C^{m+m} e^{-ck} + C'' \delta^{k/4} + C''' \alpha^{k/2}.$$

□

6. EXTENSIONS AND GENERALIZATIONS AND ADDITIONAL RESULTS, A SHORT DISCUSSION

In this section we will describe a few additional results which can also be obtained using the methods of the current paper. In order not to overload the paper the section is presented in a form of a discussion rather than explicit formulations of theorems.

6.1. More general mixing base maps for continuous in ω transfer operators, a short discussion. Let $(\xi_n)_{n \in \mathbb{Z}}$ be a stationary process taking values on a metric space (\mathcal{Y}, d) satisfying the following approximation and mixing conditions:

There are sub- σ -algebras $\mathcal{G}_{n,m}$ on the underlying probability space so that $\mathcal{G}_{n,m} \subset \mathcal{G}_{n_1, m_1}$ if $[n, m] \subset [n_1, m_1]$ and for each r and n there is an $\mathcal{G}_{n-r, n+r}$ measurable random variable $\xi_{n,r}$ so that:

- (1) **approximation:** $\|d(\xi_n, \xi_{n,r})\|_{L^\infty} \leq A_1 \beta^r$, $\beta \in (0, 1)$
- (2) **mixing:** the sequences $(\xi_{2nr, r})_{n \in \mathbb{Z}}$ are α (or ϕ_R or ψ) mixing uniformly in r .

We note that the above uniform approximation by α -mixing sequences applies to Young towers, when $\alpha_n = O(n^{-(p-2)})$ is the tails of the tower are $O(n^{-p})$ for some $p \geq 3$. We can also take several classes of smooth maps on the interval or Gibbs-Markov maps [1] for which such an approximation holds with $\psi_n = O(\delta^n)$ for some $\delta \in (0, 1)$.

Let $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$ be the shift system constructed as before. Then all the results stated in the paper hold true when $\omega \rightarrow \mathcal{L}_\omega$ and $\omega \rightarrow \varphi_\omega$ are Hölder continuous in ω (on a set with probability 1). The main point is that Lemma 3.8 and the similar approximations used in the construction of the martingale (i.e. in the proof of Proposition 4.3) can be obtained by first approximating (taking $r = r_n = \varepsilon_0 n$ for some small ε_0) and using the mixing conditions on the approximating sequences. The main reason we did not include such results in the body of the paper is that it would make the notations more complicated, and that the additional essentially global regularity assumptions on the transfer operators are somehow less natural.

6.2. Extension to random Gibbs measures. Let us consider now the random expanding maps T_ω as in [42]. Let $\mu_\omega = h_\omega \nu_\omega$ be a random Gibbs measure corresponding to a given random logarithmically α -Hölder continuous potential, and let λ_ω be the exponent of the random pressure. Namely, if \mathcal{L}_ω is the transfer operator corresponding to the random potential, then

$$\mathcal{L}_\omega h_\omega = \lambda_\omega h_{\sigma\omega}, (\mathcal{L}_\omega)^* \nu_{\sigma\omega} = \lambda_\omega \nu_\omega.$$

Next, for the sake of simplicity let us consider here random expanding maps as in [28, Chapter 5]. Then there is a constant $K > 0$ so that with $\tilde{\mathcal{L}}_\omega = \mathcal{L}_\omega / \lambda_\omega$ we have

$$\|\tilde{\mathcal{L}}_\omega^n - \nu_\omega \otimes h_{\sigma^n \omega}\|_{Holder} \leq K e^{-\lambda n}$$

where $\|\cdot\|_{Holder}$ is the usual Hölder norm corresponding to the exponent α and $\nu \otimes h(g) = \nu(g)h$. Plugging in $g = \mathbf{1}$ we get similar estimates to the ones we had in (2.10):

$$\|\tilde{\mathcal{L}}_\omega^n - h_{\sigma^n \omega}\|_{Holder} \leq K e^{-\lambda n}.$$

Remark also that $h_\omega \geq c > 0$ for some constant $c > 0$ (see [28]).

The main additional difficulty here is to estimate expressions of the form $\mu_\omega(F_\omega)$ (as in Lemma 3.8) by functions of the coordinates in places j for $|j| \leq n$. Once this is achieved, we can use the approximation argument (similarly to Lemma 3.8) which was essential in the proofs of all of the results stated in the body of the paper. The main difference in comparison with the case when $\nu_\omega = m$ does not depend on ω is that now we need to approximate ν_ω by functions of the first n coordinates (exponentially fast in n). For uniformly expanding maps, this follows from the construction of ν_ω as a certain uniform limit (see [28, Ch. 4-5]).

6.3. Extension to nonconventional sums (multiple recurrences). Let us consider partial “non-conventional” sums of the form

$$S_n \varphi = \sum_{m=1}^n \prod_{j=1}^{\ell} \varphi \circ \tau^{q_j(m)}$$

where ℓ is an integer and $q_j(n)$ are positive integer-valued sequences. The statistical properties of such sums were studied for several classes of expanding or hyperbolic maps (in particular), see [39, 40, 25] and references therein. When all q_j 's are polynomials, we believe that all the results obtained using the method of cumulants (i.e. Theorems 2.14, 2.15, 2.16, 2.17 and an appropriate version of Theorem 2.19) can be obtained for such sums exactly as in [25], relying on a version of Proposition 3.4 applied with $\rho(n, m) = \max_{1 \leq i, j \leq \ell} |q_i(m) - q_j(n)|$. The main idea is that by induction on the number of blocks we can show that the conditions of Proposition 3.3 with that $\rho = \rho_\ell$ hold true for

$$X_m = \prod_{j=1}^{\ell} \varphi \circ \tau^{q_j(m)}.$$

That is, by an inductive argument similar to the one in [28, Corollary 1.3.11], we can prove the following result.

6.1. Lemma. *Let $r \in \mathbb{N}$ and let B_1, B_2, \dots, B_k be finite subsets of \mathbb{N} so that the distance between B_j and B_{j+1} is d_j . Set $r_j = \lfloor d_j/3 \rfloor$. Let $\mathcal{C} = \{C_j : 1 \leq j \leq s\}$ be a partition of $\{1, 2, \dots, k\}$ and set $Y_j = \prod_{k \in C_j} \prod_{u \in B_k} \varphi \circ \tau^u$. Then, assuming that $\|\varphi\|_{L^\infty} \leq 1$ and that $\text{esssup}_{\omega \in \Omega} (K(\omega) \|\varphi_\omega\|_{BV}) \leq 1$, there is an absolute constant $A > 1$ so that*

$$\left| \mathbb{E}_\mu \left[\prod_{j=1}^s Y_j \right] - \prod_{j=1}^s \mathbb{E}_\mu[Y_j] \right| \leq A^m \sum_{j=1}^m (\delta^{r_j} + \alpha([r_j]))$$

where $\delta = e^{-(\lambda-3\varepsilon)/2} \in (0, 1)$.

We note that in order to prove a version of the functional CLT for the sums above we first need to use the arguments in [40, 27] to compute the variance of the limiting Gaussian, which for general polynomials might differ from a Brownian motion, and this can also be done by using the above lemma.

REFERENCES

- [1] J. Aaronson, M. Denker, *Local Limit Theorems for Gibbs-Markov Maps*, Stoch. Dyn. 1 (2001), 193-237.
- [2] R. Aimino, M. Nicol and S. Vaienti, *Annealed and quenched limit theorems for random expanding dynamical systems*, Probab. Th. Rel. Fields 162, 233-274, (2015).
- [3] J. Atnip, *An Almost Sure Invariance Principle for Several Classes of Random Dynamical Systems*, preprint, arXiv:1702.07691.
- [4] J. Atnip, G. Froyland, C. González-Tokman and S. Vaienti, *Thermodynamic Formalism for Random Weighted Covering Systems*, Comm. Math. Phys. **386** (2021), 819-902.
- [5] A. Ayer, C. Liverani, M. Stenlund, *Quenched CLT for random toral automorphism*. Discret. Contin. Dyn. Syst. 24, 331-348 (2009)
- [6] J. F. Alves, W. Bahsoun and R. Ruziboev, *Almost sure rates of mixing for partially hyperbolic attractors*, Journal of Differential Equations, Volume 311, 2022, Pages 98-157.
- [7] L. Arnold, *Random dynamical systems*, Springer Monogr. Math., Springer, Berlin, 1998.
- [8] Baladi, V.: *Correlation spectrum of quenched and annealed equilibrium states for random expanding maps*. Commun. Math. Phys. 186, 671-700 (1997)

- [9] Baladi, V., Young, L.-S.: On the spectra of randomly perturbed expanding maps. *Commun. Math. Phys.* 156, 355–385 (1993)
- [10] P. Billingsley, *Convergence of Probability Measures*, Wiley, New York (1968)
- [11] R.C. Bradley, *Introduction to Strong Mixing Conditions*, Volume 1, Kendrick Press, Heber City, 2007.
- [12] J. Buzzi, *Exponential decay of correlations for random Lasota-Yorke maps*, *Commun. Math. Phys.* **208** (1999), 25–54.
- [13] C. Cuny, F. Merlevede, *Strong invariance principles with rate for “reverse” martingale differences and applications*, *J. Theor. Prob.* 28, pages137–183 (2015).
- [14] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications*, 2nd edn. Applications of Mathematics, vol. 38. Springer, New York (1998).
- [15] H. Döring and P. Eichelsbacher, Moderate deviations via cumulants, *J. Theor. Probab.* 26 (2013), 360–385.
- [16] D. Dragičević, G. Froyland, C. González-Tokman, and S. Vaienti. Almost Sure Invariance Principle for random piecewise expanding maps, *Nonlinearity* 31 (2018), 2252–2280.
- [17] D. Dragičević, G. Froyland, C. González-Tokman, and S. Vaienti. A Spectral Approach for Quenched Limit Theorems for Random Expanding Dynamical Systems, *Comm. Math. Phys.* 360 (2018), 1121–1187.
- [18] D. Dragičević, G. Froyland, C. González-Tokman, and S. Vaienti. A Spectral Approach for Quenched Limit Theorems for Random Expanding Dynamical Systems, *Tran. Amer. Math. Soc.* 360 (2018), 1121–1187.
- [19] D. Dragičević, J. Sedro *Quenched limit theorems for expanding on average cocycles*, preprint, arXiv 2105.00548.
- [20] D. Dragičević, Y. Hafouta, J. Sedro *A vector-valued almost sure invariance principle for random expanding on average cocycles*, <https://arxiv.org/abs/2108.08714>, 41 pages.
- [21] D. Dragičević and Y. Hafouta *Limit theorems for random expanding or Anosov dynamical systems and vector-valued observables*, *Ann. Henri Poincaré* **21** (2020), 3869–3917.
- [22] D. Dragičević and Y. Hafouta, *Almost sure invariance principle for random distance expanding maps with a nonuniform decay of correlations*, *Thermodynamic Formalism*, CIRM Jean-Morlet Chair Subseries, Springer-Verlag, to appear.
- [23] D. Dragičević and Y. Hafouta *Almost sure invariance principle for random dynamical systems via Gouëzel’s approach*, *Nonlinearity*, 34 6773 (2021).
- [24] S. Gouëzel, *Almost sure invariance principle for dynamical systems by spectral methods*, *Annals of Probability* **38** (2010), 1639–1671.
- [25] Y. Hafouta, *Nonconventional moderate deviations and exponential concentration inequalities*, *Ann. Inst. H. Poincaré Probab. Statist.*, Vol. 56, No. 1, 428–448 (2020).
- [26] Y. Hafouta and Yu. Kifer, *Berry-Esseen type estimates for nonconventional sums*, *Stoch. Proc. Appl.* 126 (2016), 2430–2464.
- [27] Y. Hafouta and Yu. Kifer, *Nonconventional polynomial CLT*, *Stochastics*, 89 (2017), 550–591.
- [28] Y. Hafouta and Yu. Kifer, *Nonconventional limit theorems and random dynamics*, World Scientific, Singapore, 2018.
- [29] Y. Hafouta, *Limit theorems for some skew products with mixing base maps*, *Erg. Theor. Dyn. Syst.*, Volume 41 , Issue 1 , January 2021 , pp. 241 - 271
- [30] Y. Hafouta, *Limit theorems for random non-uniformly expanding or hyperbolic maps*, *Ann. Henri Poincaré* 23, 293–332 (2022).
- [31] N. Haydn, M. Nicol, A. Török and S. Vaienti, *Almost sure invariance principle for sequential and non-stationary dynamical systems*, *Trans. Amer. Math. Soc.* **369** (2017), 5293–5316.
- [32] H. Hennion and L. Hervé, *Limit Theorems for Markov Chains and Stochastic Properties of Dynamical Systems by Quasi-Compactness*, *Lecture Notes in Mathematics* vol. 1766, Springer, Berlin, 2001.
- [33] I.A. Ibragimov and Yu.V. Linnik, *Independent and Stationary Sequences of Random Variables*, Wolters-Noordhoff, Groningen, 1971.
- [34] Ishitani, H.: Central limit theorems for the random iterations of 1-dimensional transformations (dynamics of complex systems). *RIMS Kokyuroku*, Kyoto Univ. 1404, 21–31 (2004)
- [35] Yu. Kifer, *Perron-Frobenius theorem, large deviations, and random perturbations in random environments*, *Math. Z.* 222(4) (1996), 677–698.
- [36] Yu. Kifer, *Limit theorems for random transformations and processes in random environments*, *Trans. Amer. Math. Soc.* 350 (1998), 1481–1518.
- [37] Y. Kifer, PD. Lui, *Random Dynamics*, in *Handbook of Dynamical Systems 1B*, edited by B. Hasselblatt and A. Katok, Elsevier, 2006, pp. 379–499.
- [38] Yu. Kifer, *Thermodynamic formalism for random transformations revisited*, *Stoch. Dyn.* 8 (2008), 77–102.
- [39] Yu. Kifer, *Nonconventional limit theorems*, *Probab. Th. Rel. Fields*, 148 (2010), 71–106.
- [40] Yu. Kifer and S.R.S Varadhan, *Nonconventional limit theorems in discrete and continuous time via martingales*, *Ann. Probab.* 42 (2014), 649–688.

- [41] A Korepanov, Z Kosloff, I Melbourne *Martingale-coboundary decomposition for families of dynamical systems*, Annales Inst. H. Poincaré Analyse Non Linéaire **35** (2018), 859-885
- [42] V. Mayer, B. Skorulski and M. Urbański, *Distance expanding random mappings, thermodynamical formalism, Gibbs measures and fractal geometry*, Lecture Notes in Mathematics, vol. 2036 (2011), Springer.
- [43] F. Merlevéde, M. Peligrad and S. Utev, *Recent advances in invariance principles for stationary sequences*, Probab. Surveys 3 , 1-36 (2006).
- [44] W. Philipp and W.F. Stout, *Almost sure invariance principles for partial sums of weakly dependent random variables*, Mem. Amer. Math. Soc. 161 (1975).
- [45] E. Rio. *Théorie asymptotique des processus aléatoires faiblement dépendants. Mathématiques & Applications* (Berlin) [Mathematics & Applications] 31, Springer Verlag, Berlin, 2000.
- [46] L. Saulis and V.A. Statulevicius, *Limit Theorems for Large Deviations*, Kluwer Academic, Dordrecht, Boston, 1991.

Email address: yeor.hafouta@mail.huji.ac.il, hafuta.1@osu.edu