

Lie Theory for Asymptotic Symmetries in General Relativity: The NU Group

David Prinz* and Alexander Schmeding†

September 26, 2021

We study the Newman–Unti (NU) group from the viewpoint of infinite-dimensional geometry. The NU group is a topological group in a natural coarse topology, but it does not become a manifold and hence a Lie group in this topology. To obtain a manifold structure we consider a finer Whitney-type topology. This turns the unit component of the NU group into an infinite-dimensional Lie group. We then study the Lie theoretic properties of this group. Surprisingly, the group operations of the full NU group become discontinuous, whence the NU group does not support a Lie group structure. Recall that the NU group contains the Bondi–Metzner–Sachs (BMS) group as a subgroup and a Lie group structure of the BMS was constructed in a previous article. The lack of a Lie group structure on the NU group implies that, despite the known splitting of the NU Lie algebra into a direct sum of the BMS Lie algebra and conformal rescalings, the BMS group cannot be embedded as a Lie subgroup into the NU group.

Keywords: Bondi–Metzner–Sachs group, Newman–Unti group, asymptotically flat spacetimes, infinite-dimensional Lie group, analytic Lie group, smooth representation, Trotter product formula

MSC2020: 22E66 (primary mathematics), 22E65 (secondary mathematics); 83C30 (primary physics), 83C35 (secondary physics)

Contents

1. Introduction and statement of results	2
1.1. Physical motivation	3
1.2. Mathematical motivation	3

*Humboldt-Universität zu Berlin, Germany: prinz@math.hu-berlin.de

†Nord universitetet i Levanger, Norway: alexander.schmeding@nord.no

1.3. Original results of the present article	4
1.4. A recap of the Bondi–Metzner–Sachs group	5
2. Asymptotically flat spacetimes	6
2.1. Penrose’s conformal extension	6
2.2. Newman and Unti’s coordinate-wise definition	8
3. Lie theory for the NU group	9
3.1. General constructions	9
3.2. The Newman–Unti group	12
3.3. The group \mathcal{N} in the fine very strong topology	16
3.4. Lie group structure of the Newman–Unti group	22
4. Conclusion	27
A. Essentials from infinite-dimensional calculus	27
B. Auxiliary results for Section 3.3	29
References	35

1. Introduction and statement of results

In [PS21] we have studied the Lie group structure of the Bondi–Metzner–Sachs (BMS) group [BBM62, Sac62] from the viewpoint of infinite-dimensional geometry. The BMS group is defined as the subgroup of spacetime-diffeomorphisms that preserve asymptotic flatness in the sense of Bondi, Van der Burg, Metzner and Sachs. In this article we extend our investigations to the Newman–Unti (NU) group [NU62], which turns out to be much more involved. On the level of Lie algebras it is a well-known fact that the corresponding Lie algebras of the BMS and NU groups are related via the following direct sum decomposition [BT10b, BT10a]

$$\mathfrak{nu}_4 \cong \mathfrak{bms}_4 \oplus \mathfrak{conf}_{\mathcal{S}}, \tag{1}$$

where \mathfrak{nu}_4 and \mathfrak{bms}_4 denote the respective Lie algebras for the NU and BMS groups in 4 dimensions of spacetime and $\mathfrak{conf}_{\mathcal{S}}$ denotes the abelian Lie algebra of conformal rescalings on $\text{scri } \mathcal{S}$, the corresponding null boundary of the spacetime. Interestingly, this decomposition is not compatible with the smooth structure and does therefore not carry over to the level global Lie group level. More precisely, there is a topological obstruction which is similar to the situation for diffeomorphism groups of non-compact manifolds. There the Lie algebra of the diffeomorphism group needs to be defined as the Lie algebra of compactly supported vector fields and is not equal to the Lie algebra of all vector fields. Unfortunately, this restriction is not compatible with the decomposition of equation (1). In the following, we will briefly compare the BMS and NU groups from a physical and from a mathematical perspective. Furthermore, we

refer to our previous article [PS21] for a general introduction to asymptotic symmetry groups in General Relativity and infinite-dimensional Lie groups.

Acknowledgments: The authors wish to thank G. Barnich and H. Glöckner for helpful discussions on the subject of this work. Furthermore, we thank H. Glöckner for access to his preprint [Glö21]. DP is additionally supported by the University of Potsdam.

1.1. Physical motivation

Asymptotic symmetry groups capture the symmetries of physical models in General Relativity with a finite matter content. Such models are given e.g. by isolated stars, but also complete universes with a finite matter distribution. Therefore, asymptotically flat spacetimes are ideally suited to study gravitational waves - a topic that has regained a lot of interest after the experimental verification of gravitational waves at LIGO in 2016 [A⁺16]. However, there are different approaches to asymptotically flat spacetimes, each of which leading to different asymptotic symmetry groups, see e.g. [Fri18, Ruz20] for overviews and [PS21] for a general introduction. Specifically, there are coordinate based approaches, such as the original BMS and NU approach [BBM62, Sac62, NU62], as well as the geometric approach by Penrose [Pen63, Pen64, Pen65, Pen68]. We refer to [SSW75] for a characterization of the BMS and NU groups in terms of the geometric approach of Penrose. Physically, the BMS group and the NU group differ in the definition of the corresponding ‘supertranslations’, given for 4-dimensional spacetimes with vanishing cosmological constant via the function spaces $\mathcal{S} := C^\infty(\mathbb{S}^2)$ for the BMS group and $\mathcal{N} \subset C^\infty(\mathbb{R} \times \mathbb{S}^2)$ for the NU group, where the subset is characterized in Definition 3.8. Geometrically, the difference lies in the action of the corresponding groups on scri $\mathcal{I} := \mathbb{R} \times \mathbb{S}^2$ (cf. Definition 2.2 and Proposition 2.4), the null-boundary of the aforementioned spacetimes, and its induced (degenerate) metric. More precisely, the NU group allows for conformal rescalings of the induced metric, whereas the BMS group is additionally subjected to the invariance of a specific (2,2)-tensor, defining a ‘strong conformal geometry’, [Pen74, SSW75].

1.2. Mathematical motivation

The NU group is known to be an “infinite-dimensional group”. This directly motivates the question: Is it an infinite-dimensional Lie group? The answer to this depends on the definition of infinite-dimensional Lie group. The present articles concept of an infinite-dimensional Lie group is a group with a manifold structure modeled on a locally convex space, such that the group operations are smooth. As ordinary calculus breaks down beyond the realm of Banach spaces (and the NU group can not be modeled on a Banach space), smoothness here refers to smoothness in the sense of Bastiani calculus (see Appendix A for more information). We refer to [PS21] for an overview on infinite-dimensional Lie groups. Specifically, we have discussed that while the Lie group exponential provides a local diffeomorphism (exponential coordinates) between finite-dimensional Lie groups and their Lie algebras, this correspondence breaks down in general in the infinite-dimensional case. As an example, the Lie group exponential of

the diffeomorphism group fails to be locally surjective. In this article we encounter another complication: For the BMS group it was sufficient to consider function spaces on compact manifolds. The NU group is modeled on spaces of functions on non-compact manifolds. This might seem like a minor issue, but it leads to major complications stemming from the function space topologies. Again this complication is well known for the diffeomorphism group of a non-compact manifold [Sch15]. To illustrate this, recall that for a compact manifold K the Lie algebra of the Lie group $\text{Diff}(K)$ is the Lie algebra of vector fields on K with the negative of the usual bracket of vector fields. The natural topology on this Lie algebra is the compact open C^∞ -topology. This topology allows one to simultaneously control a function and up to finitely many of its derivatives on any given compact set. The Lie group exponential of $\text{Diff}(K)$ is the map which sends a vector field to its time 1 flow and it is not difficult to see that it is smooth with respect to the compact open C^∞ -topology.

Switching now to a non-compact manifold N , consider the Lie algebra of vector fields on N with the compact-open C^∞ -topology. Since N is non-compact, this topology does not control the behavior of vector fields at infinity. Indeed, it is too coarse in the non-compact case. Extrapolating from the compact case, also the Lie group exponential is problematic: On a non-compact manifold there are vector fields whose flow explodes in arbitrarily short time, so the flow map is simply not available on the algebra of all vector fields. All of these problems can be solved by switching to the Lie algebra of compactly supported vector fields on N with a finer topology. This topology, called the fine-very strong topology, yields enough control to work with the restricted algebra. Moreover, flows exist now and give rise to a smooth Lie group exponential turning the group $\text{Diff}(N)$ into a Lie group.

Returning to the NU group, the problems we face stem from the well known difficulties for diffeomorphism groups which were just outlined. The BMS group is a product of an infinite-dimensional abelian group (indeed a vector space of functions on a compact manifold) with a finite-dimensional Lie group. For the NU group the abelian part is replaced by a parametrized version of the diffeomorphism group of a non-compact manifold. So one should expect that all problems associated to diffeomorphism groups in the passage from compact to non-compact domains are again present.

1.3. Original results of the present article

For the present article we investigate the NU group as infinite-dimensional Lie groups in the setting of Bastiani calculus (see Appendix A for more information on the calculus). Let us first recall the (algebraic structure of the NU group. We denote by $\text{SO}^+(3, 1)$ the orthochronous Lorentz group and note that it is a finite dimensional Lie group which acts on the function space $C^\infty(\mathbb{R} \times \mathbb{S}^2)$. This action will be recalled later, for now we only mention that the action restricts to a certain (non-commutative) group $\mathcal{N} \subseteq C^\infty(\mathbb{R} \times \mathbb{S}^2)$ and the NU group is then the semidirect product

$$\text{NU} = \mathcal{N} \rtimes \text{SO}^+(3, 1) . \tag{2}$$

There is a canonical identification $I: \text{BMS} \rightarrow \text{NU}$ identifying the BMS group as a subgroup of the NU group. Note that the semidirect product (2) is topologically much

more involved compared to the BMS case. The reason is that the functions in \mathcal{N} are defined on a non-compact manifold, while functions in \mathcal{S} are defined on a compact domain. The non-compactness of $\mathbb{R} \times \mathbb{S}^2$ implies that $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ admits at least three qualitatively different choices of function space topologies (the compact open C^∞ -topology, the Whitney topologies and the fine very strong topology). Only two of these topologies provide additionally either a topological vector space structure (compact open topology) or a manifold structure (fine very strong topology) on $C^\infty(\mathbb{R} \times \mathbb{S}^2)$.

As our first result, we prove that \mathcal{N} and also the NU group becomes a topological group in the compact open C^∞ -topology. The inclusion of the BMS group becomes then a morphism of topological groups. The compact open C^∞ -topology is too coarse to construct a manifold structure on \mathcal{N} , whence there is no Lie group structure to be gained here.

Passing to the fine very strong topology, we construct a manifold structure on \mathcal{N} which turns the group \mathcal{N} into a Lie group. Note that the model space of \mathcal{N} is neither a Banach nor a Fréchet space, whence Bastiani calculus is needed to make sense of this. We then identify the Lie algebra of \mathcal{N} and establish that \mathcal{N} is a regular Lie group. Recall that a Lie group is regular if a certain kind of ordinary differential equations can be solved on the Lie group and the solution depends smoothly on parameters, cf. e.g. the introduction in [PS21]. Regularity is a prerequisite for advanced tools in (infinite-dimensional) Lie theory. Surprisingly, the action of the Lorentz group on \mathcal{N} is discontinuous with respect to the fine topology, so the NU group can not be turned into a Lie group if \mathcal{N} carries this topology. However, we show that the action on the unit component \mathcal{N}_0 of \mathcal{N} is smooth and thus the unit component NU_0 of the NU group becomes a Lie group if we endow \mathcal{N}_0 with the fine topology. As a consequence of the semidirect product structure, also NU_0 is a regular Lie group. Moreover, we establish for \mathcal{N} and NU_0 that they are not real analytic Lie groups (whence the Baker–Campbell–Hausdorff series does not yield a model for the group structure). However, our analysis shows that the strong Trotter and the strong commutator properties hold for both groups.

1.4. A recap of the Bondi–Metzner–Sachs group

We have established in [PS21] that the BMS group is an infinite-dimensional Lie group modeled on a Fréchet space. This follows directly from its semidirect product form [Sac62, McC72]

$$\text{BMS} = \mathcal{S} \rtimes \text{SO}^+(3, 1), \quad (3)$$

where $\mathcal{S} = C^\infty(\mathbb{S}^2)$ is a function space (viewed as an abelian Lie group) and $\text{SO}^+(3, 1)$ denotes the Lorentz group (which is in particular a finite-dimensional Lie group) together with the smoothness of the group action on the infinite-dimensional space of supertranslations \mathcal{S} [PS21]. We found that the BMS group is regular in the sense of Milnor [Mil84]. As a consequence, we obtain the validity of the Trotter product formula and the commutator formula on the BMS group. We remark that these formulae are important tools in the representation theory of Lie groups. Additionally, we found a more surprising results of our investigation: The BMS group is **not** an analytic Lie

group. In a nutshell, the reason for this is that the group product incorporates function evaluations of smooth (but necessarily analytic) mappings. In particular, this entails that

- the well known Baker–Campbell–Hausdorff series does not provide a local model for the Lie group multiplication, and
- there can not be a complexification of the BMS group which continues the real BMS group multiplication as a complex (infinite-dimensional) Lie group. (See also [McC92] on complexifications of the BMS group.)

Moreover, we remark that due to this defect either the Lie group exponential does not provide a local diffeomorphism onto an identity neighborhood or the BCH-series can not converge on any neighborhood of 0 in the Lie algebra.

Finally, we have also discuss the case of the generalized BMS group (or gBMS), which can be identified with the semidirect product $\mathcal{S} \rtimes \text{Diff}(\mathbb{S}^2)$, and behaves quite similarly (when it comes to the Lie theory) to the BMS group.

2. Asymptotically flat spacetimes

We start this article with an introduction to different definitions of asymptotic flatness, where we reproduce important material from [PS21] in order to be self-contained. Opposed to the historical development, we start with the geometric approach due to Penrose and then discuss the coordinate-based approaches, started by Bondi et al. and Sachs, and then continued by Newman and Unti. The reason for this is that the conformal extension due to Penrose is more fundamental, as it directly addresses the spacetime-geometry, whereas the coordinate-based constructions started by Bondi et al. and Sachs is more restrictive, by using specific coordinate representations (which can be interpreted as an additional choice of gauge). We refer to [Fri18] for an excellent overview of the different approaches.

2.1. Penrose’s conformal extension

A very geometrical approach to this subject is given via Penrose’s conformal extension [Pen63, Pen64, Pen65, Pen68]. Basically the idea is that, given a spacetime (M, g) , we call it asymptotically simple if there exists a conformal embedding into a so-called ‘extended spacetime’ $(\widehat{M}, \widehat{g})$, which is a manifold with boundary that represents the points ‘at infinity’. To this end, we start with the definition of a spacetime:

2.1 Definition (Spacetime) Let (M, g) be a Lorentzian manifold. We call (M, g) a spacetime, if it is smooth, connected, 4-dimensional and time-orientable.¹

¹We fix the spacetime-dimension solely for a cleaner presentation. Our main results are actually valid in arbitrary spacetime-dimensions, i.e. for any compact manifold K and conformal group $\text{Conf}(K)$.

With the following construction, we want to unify spacetimes that approach Minkowski spacetime ‘at infinity’. For this, we construct the boundary

$$\mathcal{I} := i_+ \sqcup \mathcal{I}_+ \sqcup i_0 \sqcup \mathcal{I}_- \sqcup i_-, \quad (4)$$

where i_+ and i_- correspond to timelike infinity, \mathcal{I}_+ and \mathcal{I}_- to lightlike infinity and finally i_0 to spacelike infinity. This is then set as the boundary of a so-called ‘extended spacetime’ (or sometimes also ‘unphysical spacetime’) $(\widehat{M}, \widehat{g})$, and then related to the spacetime (M, g) via a conformal embedding $\iota: M \hookrightarrow \widehat{M}$, as follows:

2.2 Definition (Asymptotically simple (and empty) spacetime) Let (M, g) be an oriented and causal spacetime. We call (M, g) an asymptotically simple spacetime, if it admits a conformal extension $(\widehat{M}, \widehat{g})$ in the sense of Penrose: That is, if there exists an embedding $\iota: M \hookrightarrow \widehat{M}$ and a smooth function $\varsigma \in C^\infty(\widehat{M})$, such that:

1. \widehat{M} is a manifold with interior $\iota(M)$ and boundary \mathcal{I} , i.e. $\widehat{M} \cong \iota(M) \sqcup \mathcal{I}$
2. $\varsigma|_{\iota(M)} > 0$, $\varsigma|_{\mathcal{I}} \equiv 0$ and $d\varsigma|_{\mathcal{I}} \neq 0$; additionally $\iota_*g \equiv \varsigma^2\widehat{g}$
3. Each null geodesic of $(\widehat{M}, \widehat{g})$ has two distinct endpoints on \mathcal{I}

We call (M, g) an asymptotically simple and empty spacetime, if additionally:²

4. $(R_{\mu\nu})|_{\iota^{-1}(\bar{\mathcal{O}})} \equiv 0$, where $\bar{\mathcal{O}} \subset \widehat{M}$ is an open neighborhood of $\mathcal{I} \subset \widehat{M}$

Using this construction, the BMS and Newman–Unti groups can be seen as diffeomorphisms acting on \mathcal{I} , cf. [SSW75]. Furthermore, the geometry of asymptotically simple and empty spacetimes can be characterized as follows:

2.3 Proposition *Let (M, g) be an asymptotically simple and empty spacetime. Then (M, g) is globally hyperbolic and thus parallelizable.*

Proof. A classical result due to Ellis and Hawking [HE73, Proposition 6.9.2] shows that (M, g) is globally hyperbolic. For parallelizability of (M, g) we note that the spacetimes is assumed to be four-dimensional. Hence, being globally hyperbolic, the space-submanifold is three-dimensional and thus parallelizable, as it is orientable. \square

Moreover, in the case of a vanishing cosmological constant $\Lambda = 0$, the topology of \mathcal{I} can be characterized as follows:

2.4 Proposition ([New89, Corollary 2]) *Let (M, g) be an asymptotically simple spacetime with vanishing cosmological constant $\Lambda = 0$. Then the two components representing lightlike infinity $\mathcal{I}_\pm \supset \mathcal{I}$ are both homeomorphic to $\mathbb{R} \times \mathbb{S}^2$.*

²This condition can be modified to allow electromagnetic radiation near \mathcal{I} . We remark that asymptotically simple and empty spacetimes are also called asymptotically flat.

2.2. Newman and Unti's coordinate-wise definition

The original approach to this subject was due to the pioneering works of Bondi, van der Burg and Metzner [BBM62] and Sachs [Sac62] via a coordinate-wise definition, which are described in [PS21]. In this article we focus on the coordinate approach of Newman and Unti [NU62] in the form of [BL12], where also an explicit relation between the BMS and NU gauges is discussed. As it turned out later using Penrose's conformal extension, asymptotically flat spacetimes are parallelizable (Proposition 2.3). Thus, the following coordinate functions can be defined globally, modulo possible singularities. Furthermore, we remark that they are constructed for spherically symmetric situations, which motivates the use of spherical coordinates for the spatial submanifold:

2.5 Definition (NU coordinate functions) Let (M, g) be an asymptotically simple spacetime with globally defined coordinate functions $x^\alpha: M \rightarrow \mathbb{R}^4$, denoted via $x^\alpha := (t, x, y, z)$. Then we introduce the so-called NU coordinate functions $y^\alpha: M \rightarrow \mathbb{R} \times \mathcal{I} \times \mathbb{S}^2$, where $\mathcal{I} \subseteq \mathbb{R}$ is an open subinterval, denoted via $y^\alpha = (u, \varrho, \vartheta, \varphi)$, as follows: The first coordinate u is fixed and serves as a label for null surfaces, the second coordinate $\varrho(u)$ is an affine parameter for null geodesics and the two remaining ones $z^a := (\vartheta, \varphi)$ are angular coordinates. Given these coordinates, the metric in the Newman and Unti approach can be expressed as follows:

$$g_{\mu\nu} dx^\mu \otimes dx^\nu \equiv W du \otimes du - (du \otimes d\varrho + d\varrho \otimes du) \quad (5a)$$

$$+ \varrho^2 h_{ab} (dz^a - U^a du) \otimes (dz^b - U^b du), \quad (5b)$$

where h_{ab} is the metric on the (deformed) unit sphere, which we decompose as follows

$$h_{ab} dz^a \otimes dz^b \equiv h_{ab}^{(1)} dz^a \otimes dz^b + \frac{1}{\varrho} h_{ab}^{(2)} dz^a \otimes dz^b + \mathcal{O}(\varrho^{-1}), \quad (5c)$$

where $h_{ab}^{(1)}$ is conformally flat and $h_{ab}^{(2)}$ is traceless with respect to $h_{ab}^{(1)}$, i.e. $h^{(1)ab} h_{ab}^{(2)} \equiv 0$. Here, we have expressed the metric degrees of freedom via a real function on the spacetime $W \in C^\infty(M, \mathbb{R})$, a vector field on the unit sphere $U \in \mathfrak{X}(\mathbb{S}^2)$ and a metric on the unit sphere $h \in \text{Met}(\mathbb{S}^2)$.

2.6 Definition (NU asymptotic flatness) Given the coordinate functions from Definition 2.5 with the angular functions $z^a := (\vartheta, \varphi)$ transformed into stereographic coordinates $\zeta := \cot(\vartheta/2) \exp(i\varphi)$ and $\bar{\zeta} := \cot(\vartheta/2) \exp(-i\varphi)$, the spacetime (M, g) is called asymptotically flat in the sense of NU, if the following relations are satisfied [BL12]:

$$W = -2\varrho \partial_u k + 4 \exp(-2k) \partial_\zeta \partial_{\bar{\zeta}} k + \mathcal{O}(\varrho^{-1}), \quad U^a = \mathcal{O}(\varrho^{-2}) \quad (6)$$

and Equation (5c), where $k(u, \vartheta, \varphi)$ is the conformal factor such that $h_{ab}^{(1)} dz^a \otimes dz^b \equiv \exp(2k) d\zeta \otimes d\bar{\zeta}$.

2.7 Remark We emphasize the explicit relation between the BMS and the NU gauges that is worked out in [BL12, Section 4].

3. Lie theory for the NU group

Asymptotic symmetry groups are subgroups of the diffeomorphism group that preserve the chosen boundary condition and gauge fixing. We focus in this article on the NU group. Our aim is to establish (infinite-dimensional) Lie group structures on the NU group. For readers who are not familiar with calculus beyond Banach spaces, we have compiled the basic definitions in Appendix A (and we suggest to review them before continuing). Furthermore, as the NU group is an extension of the (perhaps more well known) Bondi–Metzner–Sachs (BMS) group, let us recall some facts and notation on the BMS_v group from [PS21].

Let $\mathcal{S} := C^\infty(\mathbb{S}^2) := C^\infty(\mathbb{S}^2, \mathbb{R})$ be the abelian group of supertranslations and $\text{SO}^+(3, 1)$ the orthochronous Lorentz group $\text{SO}^+(3, 1)$. Then $\text{SO}^+(3, 1)$ acts by conformal transformations on the sphere \mathbb{S}^2 . Furthermore, we can identify elements in $\text{SO}^+(3, 1)$ with Möbius transformations. Recall that a Möbius transformation admits a matrix representation $\Lambda_f = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for $f \in \text{SO}^+(3, 1)$ which acts conformally on the Riemann sphere $\hat{\mathbb{C}} \cong \mathbb{S}^2$. The conformal factor of the transformation f can thus be expressed as

$$K: \text{SO}^+(3, 1) \times \hat{\mathbb{C}} \rightarrow]0, \infty[, \quad K_f(\zeta) := \frac{1 + \|\zeta\|^2}{\|a\zeta + b\|^2 + \|c\zeta + d\|^2}, \Lambda_f = \begin{bmatrix} a & b \\ c & d \end{bmatrix}. \quad (7)$$

Identifying the sphere \mathbb{S}^2 with the Riemann sphere, we obtain a smooth group action

$$\sigma: \mathcal{S} \times \text{SO}^+(3, 1) \rightarrow \mathcal{S}, (f, \alpha) \mapsto K_f(\cdot)^{-1} \cdot \alpha \circ f$$

and the BMS group is the semidirect product $\text{BMS} = \mathcal{S} \rtimes \text{SO}^+(3, 1)$ with respect to this action. To spell it out explicitly, the group product of the BMS group is

$$(F, \phi)(G, \psi) = (F + \sigma(G, \phi), \phi \circ \psi) = (F + K_\phi^{-1} \cdot G \circ \phi, \phi \circ \psi).$$

In [PS21] it was shown that this structure turns the BMS group into an infinite-dimensional regular Lie group.

3.1. General constructions

Before we begin, let us recall several general constructions which will be used throughout the following sections. We will encounter spaces of differentiable mappings as infinite-dimensional manifolds. Let us repeat some important definitions and properties of these manifolds. In the following M, N will always denote smooth paracompact manifolds and $C^k(M, N)$ the space of k -times continuously differentiable mappings from M to N where $k \in \mathbb{N}_0 \cup \{\infty\}$.

3.1 (Compact open C^k -topology) If nothing else is said, we topologize $C^k(M, N)$ with the compact open C^k -topology. This is the topology turning the mapping

$$C^k(M, N) \rightarrow \prod_{\ell \in \mathbb{N}_0, \ell \leq k} C(T^\ell M, T^\ell N), \quad f \mapsto (T^\ell f)_{0 \leq \ell \leq k}$$

into a topological embedding. Here the spaces on the right hand side are topologized with the compact open topology and T^ℓ denotes the ℓ -fold iterated tangent functor. Recall from [AGS20, Appendix A] that this topology turns $C^k(M, N)$ into a Banach (resp. Fréchet) manifold for $k \in \mathbb{N}_0$ (resp. $k = \infty$) if M is compact and N is a finite dimensional manifold. Moreover, one can prove that if N is a locally convex vector space, then also $C^k(M, N)$ is a locally convex vector space with the pointwise operations.

If M is non-compact, the compact open C^k -topology does not control the behaviour of mappings at infinity. Moreover, the compact open C^k -topology does not turn $C^k(M, N)$ into a manifold if N is not a vector space. For this reason one introduces the so called fine very strong topology on $C^k(M, N)$. We recall its definition now and refer to [HS17] for more information.

3.2 (the fine very strong topology) We now endow $C^\infty(M, N)$ with the so called \mathcal{FD} -topology or *fine very strong topology* and write $C_{\text{FS}}^\infty(M, N)$ for the space endowed with this topology. This is a Whitney type topology controlling functions and their derivatives on locally finite families of compact sets. Before we describe a basis of the fine very strong topology, we have to construct a basis for the strong topology which we will then refine. To this end, we recall the construction of the so called basic neighborhoods (see [HS17]). Consider f smooth, A compact, $\varepsilon > 0$ together with a pair of charts (U, ψ) and (V, φ) such that $A \subseteq V$ and $\psi \circ f \circ \varphi^{-1}$ makes sense. Then we use standard multiindex notation to define an *elementary f -neighborhood*

$$\mathcal{N}^r(f; A, \varphi, \psi, \varepsilon) := \left\{ \begin{array}{l} g \in C^\infty(M, N), \quad \psi \circ g|_A \text{ makes sense,} \\ \sup_{\alpha \in \mathbb{N}_0^d, |\alpha| < r} \sup_{x \in \varphi(A)} \|\partial^\alpha \psi \circ f \circ \varphi^{-1}(x) - \partial^\alpha \psi \circ g \circ \varphi^{-1}(x)\| < \varepsilon \end{array} \right\}.$$

A basic neighborhood of f arises now as the intersection of (possibly countably many) elementary neighborhoods $\mathcal{N}^r(f; A_i, \varphi_i, \psi_i, \varepsilon_i)$ where the family $(V_i, \varphi_i)_{i \in I}$ is locally finite. We remark that basic neighborhoods form the basis of the very strong topology. To obtain the fine very strong topology, one declares the sets

$$\{g \in C^\infty(M, N) \mid \exists K \subseteq M \text{ compact such that } \forall x \in M \setminus K, g(x) = f(x)\} \quad (\star)$$

to be open and constructs a subbase of the fine very strong topology as the collection of sets (\star) (where $f \in C^\infty(M, N)$ and the basic neighborhoods of the very strong topology. If M is compact, the fine very strong topology coincides with the compact open C^∞ -topology.

3.3 The fine-very strong topology turns $C^\infty(M, N)$ into an infinite-dimensional manifold (cf. [Mic80] and [HS17]). If $N = F$ is a locally convex space, the pointwise operations turn $C_{\text{FS}}^\infty(M, \mathbb{R}^n)$ into a vector space. However, for non-compact M , $C_{\text{FS}}^\infty(M, F)$ is disconnected, whence it is a manifold but not a locally convex space. The largest

locally convex space contained in $C_{\text{fs}}^\infty(M, F)$ is the space of compactly supported maps

$$C_c^\infty(M, F) := \{f \in C^\infty(M, F) \mid \exists K \subseteq M \text{ compact, s.t. } f|_{M \setminus K} \equiv 0\}$$

We shall always topologize $C_c^\infty(M, F)$ with the fine very strong topology.

Recall from [AGS20, Lemma A.10] that for M compact and N a finite dimensional manifold the manifold $C^k(M, N)$ is canonical, i.e. a mapping

$$h: A \rightarrow C^k(M, N) \text{ for any smooth manifold } A$$

is of class C^ℓ if and only if the adjoint map $h^\wedge: A \times M \rightarrow N, (a, m) \mapsto h(a)(m)$ is a $C^{\ell, k}$ -map. This means that h^\wedge is ℓ -times continuously differentiable with respect to the A -component of the product and each of these differentials is then k -times differentiable with respect to the M -component (cf. A.6). This is an extremely useful property of the compact open C^k -topology. We warn the reader that the corresponding statement is false for the fine very strong topology if M is non-compact.

Finally, we recall from [HN12, Lemma 2.2.3] the concept of a semidirect product (all asymptotic symmetry groups in this article will turn out to be semidirect products).

3.4 (Semidirect product of groups) Let N and H be groups and $\text{Aut}(N)$ the group of automorphisms of N . Assume that $\delta: H \rightarrow \text{Aut}(N)$ is a group homomorphism. Then we define a multiplication on $N \times H$ by

$$(n, h)(m, g) := (n\delta(h)(m), hg) \tag{8}$$

This multiplication turns $N \times H$ into a group denoted by $N \rtimes_\delta H$, where $N \cong N \times \{e\}$ is a normal subgroup and $H \cong \{e\} \times H$ is a subgroup. Furthermore, each element $x \in N \rtimes_\delta H$ has a unique representation as $x = nh, n \in N, h \in H$.

If H, N are Lie groups (or analytic Lie groups) and $\delta^\wedge: H \times N \rightarrow H, (h, n) \mapsto \delta(h)(n)$ is smooth (analytic)³, then $N \rtimes_\delta H$ is a Lie group (analytic Lie group). Its Lie algebra is a semi-direct product of Lie algebras.

Almost local mappings

To establish smoothness of certain mappings with respect to the function space topologies just defined we need Glöckner's concept of almost local mappings, see [Glö05]. We present here a version of this technique which allows for parameter dependent almost local mappings. These results were communicated to us by H. Glöckner and will appear in [Glö21]. We remark here that the proofs for these results are variants of the proofs for the statements without parameter in [Glö05].

³If N is finite-dimensional, it suffices to require that δ is a Lie group morphism. In general, there is no Lie group structure on $\text{Aut}(N)$ which guarantees smoothness of the group operation in $N \rtimes_\delta H$.

3.5 Let M, N be finite-dimensional smooth manifolds and E, F be locally convex spaces. Fix an open set $\Omega \subseteq C_c^\infty(M, E)$. Furthermore, consider a smooth (possibly infinite-dimensional) manifold P and a map

$$f: P \times \Omega \rightarrow C_c^\infty(N, F).$$

The map f is called an *almost local* if for each $p \in P$ there exist open p -neighborhood $Q \subseteq P$, a locally finite cover $(U_n)_{n \in A}$ of M by relatively compact open subsets and a locally finite cover $(V_n)_{n \in A} \subseteq N$ of open relatively compact sets (where both families are indexed by the same set A) such that the following condition holds

$$\forall n \in A, \forall q \in Q, \forall \sigma, \tau \in \Omega, \quad \sigma|_{U_n} = \tau|_{U_n} \Rightarrow f(q, \sigma)|_{V_n} = f(q, \tau)|_{V_n}.$$

If each $(p, \sigma) \in P \times \Omega$ has an open neighborhood $P_0 \times \Omega_0$ such that $f|_{P_0 \times \Omega_0}$ is almost local, f is called *locally almost local*.

This notion is relevant due to Glöckner's smoothness proposition for locally almost local maps (which is a parameter dependent version of [Glö05, Theorem 3.2]).

3.6 Proposition *In the situation of 3.5 assume that the map*

$$f: P \times \Omega \rightarrow C_c^\infty(N, F)$$

has the following properties

1. *The restriction of f to a mapping*

$$P \times (\Omega \cap C_K^\infty(M, E)) \rightarrow C_c^\infty(N, F)$$

is smooth for each compact subset $K \subseteq M$; and

2. *f is an almost local map (or locally almost local).*

Then f is smooth.

3.7 Remark The statement of Proposition 3.6 simplifies several assumptions and the result obtained. We mention that [Glö21] establishes in particular a version for finite orders of differentiability and works not only with the spaces $C_c^\infty(M, E)$ but more generally with C^k -sections of locally convex vector bundles.

3.2. The Newman–Unti group

Let us first recall the definition of the Newman–Unti group as a certain semidirect product. let $\text{Diff}^+(\mathbb{R})$ be the group of all smooth orientation preserving diffeomorphisms of \mathbb{R} (recall that a diffeomorphism of \mathbb{R} is orientation preserving if and only if it has positive derivative everywhere).

3.8 Definition Define $\mathcal{N} := \left\{ F \in C^\infty(\mathbb{R} \times \mathbb{S}^2) \mid F(\cdot, \mathbf{z}) \in \text{Diff}^+(\mathbb{R}), \quad \forall \mathbf{z} \in \mathbb{S}^2 \right\}$. Then \mathcal{N} becomes a group with respect to the product

$$F \cdot G(u, \mathbf{z}) := F(G(u, \mathbf{z}), \mathbf{z}).$$

The unit of the product is the map $p: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}, (t, \mathbf{z}) \mapsto t$ and the inverse $F^{-1}: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}$ is for $\mathbf{z} \in \mathbb{S}^2$ given by $F(\cdot, \mathbf{z})^{-1}$, where the inverse is computed in $\text{Diff}^+(\mathbb{R})$. Note that the inverse is the unique smooth map⁴ satisfying the implicit equation

$$t = F(F^{-1}(t, \mathbf{z}), \mathbf{z}) \quad (t, \mathbf{z}) \in \mathbb{R} \times \mathbb{S}^2. \quad (9)$$

Furthermore, using the conformal factor from (7), the mapping

$$\tau: \mathcal{N} \times \text{SO}^+(3, 1) \rightarrow \mathcal{N}, (F, \phi) \mapsto \left((t, \mathbf{z}) \mapsto K_\phi^{-1}(\mathbf{z})F(K_\phi(\mathbf{z})t, \phi(\mathbf{z})) \right),$$

makes sense. A quick calculation shows that it is a group action which induces a group morphism $\hat{\tau}: \text{SO}^+(3, 1) \rightarrow \text{Aut}(\mathcal{S}), \phi \mapsto \tau(\cdot, \phi)$ and we can form the semidirect product of groups. The *Newman–Unti (NU) group* is now the semidirect product

$$\text{NU} := \mathcal{N} \rtimes_\tau \text{SO}^+(3, 1).$$

Explicitly, the multiplication $(F, \phi) \cdot (G, \psi)$ of the NU group is given by

$$(F(\tau(G, \phi)(t, \mathbf{z}), \mathbf{z}), \phi \circ \psi(\mathbf{z})) = (F(K_\phi^{-1}(\mathbf{z})G(K_\phi(\mathbf{z})t, \phi(\mathbf{z})), \mathbf{z}), \phi(\psi(\mathbf{z}))).$$

By construction of the group structures we obtain an injective group morphism:

$$I: \text{BMS} \rightarrow \text{NU}, (F, \phi) \mapsto ((t, \mathbf{z}) \mapsto t + F(\mathbf{z}), \phi) \quad (10)$$

3.9 Remark Our definition of the NU-group might look odd to the reader used to the usual presentations in the physics literature. In [AE18, Definition 5.1] the definition of the space \mathcal{N} seems only to require that $F \in C^\infty(\mathbb{R} \times \mathbb{S}^2)$ satisfies

$$\frac{\partial F}{\partial t}(t, \mathbf{z}) > 0 \quad \forall (t, \mathbf{z}) \in \mathbb{R} \times \mathbb{S}^2 \quad (11)$$

(an impression one could also have from the formulation in [SSW75], though a second glance shows that the maps need to induce diffeomorphisms of \mathcal{S}) we note that this definition would **not** lead to a group structure on \mathcal{N} . As an example, we could consider the mapping $F: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}, F(t, \mathbf{z}) := \arctan(t)$ which would satisfy (11) but its inverse \tan can not be extended to a smooth map on \mathbb{R} .

Compared to the BMS group, one replaces the abelian group of supertranslations by the non-abelian group \mathcal{N} . Beyond the non-abelian structure there is another significant difference between the two groups. Whereas the supertranslations are smooth

⁴Note that smoothness of the inverse in all variables is guaranteed by the implicit function theorem.

mappings on a compact domain, the group $\mathcal{N} = C^\infty(\mathbb{R} \times \mathbb{S}^2)$ consists of smooth mappings on a non-compact domain. Thus for \mathcal{S} there is in principal only one choice for the function space topology, while for \mathcal{N} the different topologies do not coincide.⁵ Before we investigate this behaviour let us note for later use that elements in \mathcal{N} are proper maps:

3.10 Lemma *If $F \in \mathcal{N}$, then for every $K \subseteq \mathbb{R}$ compact, $F^{-1}(K)$ is compact, i.e. F is a proper map.*

Proof. Let $K \subseteq \mathbb{R}$ be compact and let us show that the preimage $F^{-1}(K)$ is compact. To this end, pick a sequence $(r_n, \mathbf{z}_n) \in F^{-1}(K), n \in \mathbb{N}$. We need to show that this sequence has a convergent subsequence. Due to the compactness of K , we see that (by passing to a subsequence) we may assume that $F(r_n, \mathbf{z}_n)$ converges in K towards a limit, say z . Now \mathbb{S}^2 is compact, whence we can pass again to a subsequence and assume that \mathbf{z}_n converges towards some $\mathbf{z}_* \in \mathbb{S}^2$. Now $F(\cdot, \mathbf{z})$ is a diffeomorphism, whence $r_* := F(\cdot, \mathbf{z}_*)^{-1}(z)$ exists. We claim that now $r_n \rightarrow r_*$. To see this, consider the diffeomorphisms

$$G_n := F(\cdot, \mathbf{z}_*)^{-1} \circ F(\cdot, \mathbf{z}_n), \quad n \in \mathbb{N}.$$

By construction this sequence of diffeomorphisms converges (pointwise) towards the identity and satisfies $G_n(r_n) \rightarrow r_*$ as $n \rightarrow \infty$. We see that $G_n^{-1}(r_*) \rightarrow r_*$ and $r_n - G_n(r_*) \rightarrow 0$, whence also r_n converges towards r_* . We conclude that (r_n, \mathbf{z}_n) has a convergent subsequence and thus $F^{-1}(K)$ is compact. Since K was arbitrary, F is a proper map. \square

We now topologize the Newman–Unti group with respect to the compact open C^∞ topology and consider whether it becomes a topological or Lie group in that way. This is the coarsest function space topology and the NU group with this topology can be analyzed similarly to the BMS group.

3.11 Proposition *Endow \mathcal{N} with the subspace topology induced by the compact open C^∞ -topology on $C^\infty(\mathbb{R} \times \mathbb{S}^2)$. Then the NU-group becomes a topological group.*

Proof. From 3.3 we see that a map $f: M \rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2)$ is smooth if and only if the associated map $f^\wedge: M \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}, (m, (t, \mathbf{z})) \mapsto f(m)(t, \mathbf{z})$ is smooth. This entails that the evaluation map $\text{ev}: C^\infty(\mathbb{R} \times \mathbb{S}^2) \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}, (f, t, \mathbf{z}) \mapsto f(t, \mathbf{z})$ is smooth, as it is the adjoint map to the identity $C^\infty(\mathbb{R} \times \mathbb{S}^2) \rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2)$. In the following we prove directly that group product and inversion in the NU-group are continuous (it will be clear from the proof that this entails that also \mathcal{N} is a topological group with the subspace topology and the group action of $\text{SO}^+(3, 1)$ on \mathcal{N} is continuous).

⁵The space $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ supports a variety of inequivalent function space topologies beyond the compact open C^∞ and the fine very strong topology. For example one can define several Whitney-type topologies, see [Mic80, Section 4]. As these additional topologies do not turn \mathcal{N} into a manifold we will ignore them.

Step 1: *The group product of the NU-group is continuous.* Let us first extend the group product to the locally convex space $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ containing \mathcal{N} . To write it in terms of composition of mappings, consider $\text{pr}_2: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{S}^2, (t, \mathbf{z}) \mapsto \mathbf{z}$. Then we can write the product as:

$$P: (C^\infty(\mathbb{R} \times \mathbb{S}^2) \times \text{SO}^+(3, 1)(\mathbb{C}))^2 \rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2) \times \text{SO}^+(3, 1),$$

$$((F, \phi), (G, \psi)) \mapsto (F \circ (\hat{\tau}(\phi)(G), \text{pr}_2), \phi \circ \psi) \quad (12)$$

As $\text{SO}^+(3, 1)$ is a Lie group, it suffices to establish continuity of the first component of P . Note that ψ is irrelevant to the first component of P , whence due to the exponential law it suffices to establish smoothness of the mapping

$$C^\infty(\mathbb{R} \times \mathbb{S}^2) \times \text{SO}^+(3, 1) \times C^\infty(\mathbb{R} \times \mathbb{S}^2) \times \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}$$

$$(F, \phi, G, t, \mathbf{z}) \mapsto F(K_\phi^{-1}(\mathbf{z})G(K_\phi(\mathbf{z}) \cdot t, \phi(\mathbf{z})), \mathbf{z})$$

However, this mapping is easily seen to be smooth as it can be written as a composition of the smooth mappings K, δ and the smooth action of $\text{SO}^+(3, 1)$ on \mathbb{S}^2 and the smooth evaluation map ev . We deduce from the exponential law that the map P on $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ is smooth. In particular, it is continuous and restricts to the group product of the NU-group on \mathcal{N} . As \mathcal{N} carries the subspace topology, this establishes continuity of the group product.

Step 2: *Inversion in the NU-group is continuous.* Recall that in a semidirect product the inversion is given by the formula

$$\iota: \mathcal{N} \times \text{SO}^+(3, 1) \rightarrow \mathcal{N} \times \text{SO}^+(3, 1), (F, \Lambda) \mapsto (\tau(F^{-1}, \Lambda^{-1}), \Lambda^{-1})$$

where F^{-1} is the inverse in the group \mathcal{N} . Since $\text{SO}^+(3, 1)$ is a Lie group, inversion is smooth and we may suppress it in the following computations. Having endowed \mathcal{N} with the subspace topology of $C^\infty(\mathbb{R} \times \mathbb{S}^2)$, continuity of ι will follow if we can establish the continuity of the mapping

$$i_1: \mathcal{N} \times \text{SO}^+(3, 1) \rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2), (F, \phi) \mapsto \tau(F^{-1}, \phi).$$

We apply now the exponential law [AS15, Theorem B] in its strong form: The map i_1 is continuous if and only if the adjoint map

$$i_1^\wedge: (\mathcal{N} \times \text{SO}^+(3, 1)) \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}, ((F, \phi), (t, \mathbf{z})) \mapsto K_\phi^{-1}(\mathbf{z})F^{-1}(t, \phi(\mathbf{z}))$$

is a $C^{0, \infty}$ -mapping. Recall that a mapping is of class $C^{0, \infty}$ if it is continuous and infinitely often continuously differentiable with respect to the second component (i.e. in the present case wrt. (t, \mathbf{z})). Rewriting the latter formula we see that

$$i_1^\wedge(F, \phi)(t, \mathbf{z}) = \text{ev}(\delta(\phi), \mathbf{z}) \cdot \text{ev}(F^{-1}, (t, \phi(\mathbf{z}))).$$

Now δ and the evaluation maps are smooth and we know that $\text{SO}^+(3, 1)$ acts smoothly on \mathbb{S}^2 . From the chain rule for $C^{r, s}$ -mappings [AS15, Lemma 3.17 and 3.18] we deduce that i_1^\wedge is a $C^{0, \infty}$ -map if the map $H: \mathcal{N} \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}, (F, (t, \mathbf{z})) \mapsto F^{-1}(t, \mathbf{z})$ is a

$C^{0,\infty}$ -map. By construction we see that $H(F, \cdot)$ solves the implicit equation (9) for the function $F \in \mathcal{N}$. Hence we can treat the whole equation as an implicit equation with parameter F . It is well known that the implicit function theorem with (continuous) parameter yields a smooth solution which depends continuously on the parameter. In other words the map H will be of class $C^{0,\infty}$ as a result of the implicit function theorem. Thus we can deduce continuity of the inversion in the NU-group from an implicit function theorem with parameter in a locally convex space (a suitable version of the theorem is recorded in [Gl606, Proposition 2.1.]). \square

3.12 Lemma *The canonical inclusion $I(F, \phi) = ((t, \mathbf{z}) \mapsto (t + F(\mathbf{z}), \phi))$, (10) of the BMS group into the NU-group is a topological group morphism.*

Proof. From the definition of I it is clear that it will be continuous if its first component

$$\mathcal{S} \rightarrow \mathcal{N}, \quad F \mapsto p + F \circ \text{pr}_2$$

is continuous, where $\text{pr}_2: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{S}^2$ denotes the projection onto the second component. We exploit that \mathcal{N} carries the subspace topology of $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ and the exponential law [AS15, Theorem B]. It suffices to prove that the mapping

$$C^\infty(\mathbb{S}^2) \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}, \quad (F, (t, \mathbf{z})) \mapsto t + F(\mathbf{z}) \quad (13)$$

is a $C^{0,\infty}$ -map. Indeed this map is even smooth as [AS15, Proposition 3.20] shows that the evaluation map ev on $C^\infty(\mathbb{S}^2)$ is smooth. Hence $(t, \mathbf{z}) \mapsto t + \text{ev}(F, \mathbf{z})$ and thus also I are smooth. \square

3.13 Remark Note that the compact open C^∞ -topology there does not seem to be any sensible way to turn \mathcal{N} into a submanifold. Indeed in the compact open topology $\text{Diff}^+(\mathbb{R})$ is not a submanifold of $C^\infty(\mathbb{R})$ in any suitable sense. Thus the compact open topology does not seem to provide a suitable structure to turn \mathcal{N} into an infinite-dimensional Lie group.

While the compact open C^∞ -topology is too coarse to turn \mathcal{N} into a manifold. Switching to a finer topology this problem can be remedied. We will investigate the resulting structure in the next section.

3.3. The group \mathcal{N} in the fine very strong topology

In this section we endow $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ with the fine very strong topology. We will see that \mathcal{N} is an open subset of $C_{\text{fs}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ and this endows \mathcal{N} with a manifold structure as $C^\infty(\mathbb{R} \times \mathbb{S}^2)$ is a manifold (cf. [Mic80]). The following result seems to be new (and while the related set $\text{Diff}^+(\mathbb{R}) \subseteq C_{\text{fs}}^\infty(\mathbb{R})$ is open (cf. [Mic80, Section 10]) this is not immediately useful to establish the following).

3.14 Proposition *The set \mathcal{N} is an open subset of $C_{\text{fs}}^\infty(\mathbb{R} \times \mathbb{S}^2)$.*

The proof of Proposition 3.14 is postponed to Appendix B. Having now a manifold structure at our disposal, the group \mathcal{N} turns out to be a Lie group.

3.15 Proposition *The submanifold structure turns $\mathcal{N} \subseteq C_{\text{fS}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ into a Lie group.*

Proof. Recall from [HS17] that in the fine very strong topology $C_{\text{prop}}^\infty(\mathbb{R} \times \mathbb{S}^2)$, the subset of proper mappings, is an open subset. By Lemma 3.10 and Proposition 3.14 the group \mathcal{N} is an open subset of $C_{\text{prop}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ whence it is an open submanifold of $C_{\text{fS}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ (see [Mic80, Theorem 10.4] for a construction of the manifold structure on the function space).

Step 1: *The multiplication is smooth.* We rewrite the group product of \mathcal{N} as follows

$$F \cdot G = F \circ (G \times \text{id}_{\mathbb{S}^2}) \circ (\text{id}_{\mathbb{R}} \times \Delta) = (\text{id}_{\mathbb{R}} \times \Delta)^*(\text{Comp}(F, G \times \text{id}_{\mathbb{S}^2})), \quad (14)$$

where $\Delta: \mathbb{S}^2 \rightarrow \mathbb{S}^2 \times \mathbb{S}^2$, $\mathbf{z} \mapsto (\mathbf{z}, \mathbf{z})$ is the diagonal map, and we have

$$\begin{aligned} (\text{id}_{\mathbb{R}} \times \Delta)^*: C^\infty(\mathbb{R} \times \mathbb{S}^2, \mathbb{R} \times \mathbb{S}^2) &\rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2 \times \mathbb{S}^2, \mathbb{R} \times \mathbb{S}^2), \phi \mapsto \phi \circ (\text{id}_{\mathbb{R}} \times \Delta) \\ \text{Comp}: C^\infty(\mathbb{R} \times \mathbb{S}^2) \times C_{\text{prop}}^\infty(\mathbb{R} \times \mathbb{S}^2 \times \mathbb{S}^2, \mathbb{R} \times \mathbb{S}^2) &, (F, \phi) \mapsto F \circ \phi \end{aligned}$$

(where the subscript prop denotes the open subset of proper mappings). From [Mic80, Corollary 10.14 and Theorem 11.4] we deduce that $(\text{id}_{\mathbb{R}} \times \Delta)^*$ and Comp are smooth. Furthermore, the map $C^\infty(\mathbb{R} \times \mathbb{S}^2) \rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2 \times \mathbb{S}^2, \mathbb{R} \times \mathbb{S}^2)$, $G \mapsto G \times \text{id}_{\mathbb{S}^2}$ is smooth by compactness of \mathbb{S}^2 and [Mic80, Corollary 11.10 1.]. We deduce from (14) that the group product is smooth as a composition of smooth mappings.

Step 2: *\mathcal{N} is a Lie group.* Since we already know that the group product is smooth and inversion $I: \mathcal{N} \rightarrow \mathcal{N}$ is a group anti-morphism, the formula $I(F) = G^{-1} \cdot I(F \cdot G)$ shows that it suffices to prove that I is smooth in an open neighborhood of the identity element $p: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}$, $(x, \mathbf{z}) \mapsto x$. In B.5 we construct a local model ι for the pointwise inversion I on $\Phi(G_0) \subseteq C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ where Φ is a chart and G_0 a suitable p -neighborhood. We will now use the auxiliary results from Appendix B to establish smoothness of ι .

Then Lemma B.7 shows that ι restricts to a smooth map on the closed subspaces $C_L^\infty(\mathbb{R} \times K)$. Define now for $R > 0$ the set $O_R := \{\gamma \in \Phi(G_0) \mid \sup_{(x,k)} |\gamma(x,k)| < R\}$. By construction O_R is an open subset of $\Phi(G_0) \subseteq C_c^\infty(\mathbb{R} \times K)$ and we see that $\Phi(G_0) = \bigcup_{R>0} O_R$. Moreover, Lemma B.8 implies that for every $R > 0$ the restriction of ι to $O_R \cap \Phi(G_0)$ is almost local. Hence, Glöckner's smoothness theorem [Glö05, Theorem 3.2] shows that ι is smooth, whence I is smooth on G_0 .

In conclusion the manifold structure turns the group operations into smooth maps, whence \mathcal{N} is an infinite-dimensional Lie group. \square

In essence the group \mathcal{N} is a version of the group $\text{Diff}^+(\mathbb{R})$ with an added parameter which is not directly visible in the composition of the group. hence the Lie algebra should be given by a Lie algebra of (compactly supported) vector fields where every vector fields depends on a parameter which is not relevant for the Lie bracket. The next proposition shows that this is indeed the case. To formulate it, we recall from

B.4 that the set of all compactly supported mappings $C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ becomes a locally convex space if we endow it with the subspace topology induced by the strong very fine topology.

3.16 Proposition *The Lie algebra of \mathcal{N} can be identified as $\mathbf{L}(\mathcal{N}) = C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ with the Lie bracket given by*

$$[X, Y](u, \theta) := -[X(\cdot, \theta), Y(\cdot, \theta)](u), \quad (u, \theta) \in \mathbb{R} \times \mathbb{S}^2$$

and the Lie bracket on the right is the usual Lie bracket of vector fields (on \mathbb{R}).

Proof. The unit in \mathcal{N} is the projection $p(u, \theta) = u$ and we use [Mic80, 10.12] to identify

$$T_p\mathcal{N} = \{(p, X) \in C^\infty(\mathbb{R} \times \mathbb{S}^2, T\mathbb{R}^2) \mid X \in C_c^\infty(\mathbb{R} \times \mathbb{S}^2)\}.$$

This identifies the Lie algebra as a locally convex space. Now we need to compute the Lie bracket. Again the proof is a variant of the classical argument by which the Lie algebra of the diffeomorphism group can be identified, [Mil84, 6.]. Take $X \in \mathbf{L}(\mathcal{N}) = C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$. Since multiplication in \mathcal{N} is given by composition in the u -component, we deduce from [Mic80, Corollary 11.6] that the extension of X to a right invariant vector field R_X on \mathcal{N} is given by the formula

$$R^X(F) = X \circ (F \times \text{pr}_2), \quad F \in \mathcal{N}$$

where $\text{pr}_2: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{S}^2$ is the projection onto the second component. Consider the vector field $R^X \times 0 \in \mathcal{V}(\mathcal{N} \times (\mathbb{R} \times \mathbb{S}^2))$. We shall now prove that $R^X \times 0$ is related to $X_0: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R} \times T\mathbb{S}^2, (u, \mathbf{z}) \mapsto (X(u, \mathbf{z}), 0(\mathbf{z}))$, where $0(\mathbf{z})$ is the zero-vector field on \mathbb{S}^2 . To this end consider the left-action

$$\alpha: \mathcal{N} \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R} \times \mathbb{S}^2, \quad (F, (u, \mathbf{z})) \mapsto (F(u, \mathbf{z}), \mathbf{z}).$$

As the evaluation $\text{ev}: C_{fS}^\infty(\mathbb{R} \times \mathbb{S}^2) \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}, (F, (u, \mathbf{z})) \mapsto F(u, \mathbf{z})$ is smooth by [Mic80, Corollary 11.6], it is easy to see that α is a Lie group action. Moreover, [Mic80, Corollary 11.6] implies that $T_{(F, (u, \mathbf{z}))} \text{ev}(X, Y) = X(u, \mathbf{z}) + dF(u, \mathbf{z}; Y)$. Plugging the vector field $R^X \times 0$ into this formula we deduce that

$$T\alpha \circ (R^X \times 0)(F, (u, \mathbf{z})) = (X(F(u, \mathbf{z})), 0_{\mathbf{z}}) = X_0(\alpha(F, (u, \mathbf{z}))).$$

As relatedness is inherited by the Lie bracket of vector fields, we find

$$\begin{aligned} \left([R^X, R^Y](p)(u, \mathbf{z}), 0(\mathbf{z}) \right) &= T\alpha \left([R^X \times 0, R^Y \times 0] \right) (p, (u, \mathbf{z})) \\ &= \left([X(\cdot, \mathbf{z}), Y(\cdot, \mathbf{z})](u), 0(\mathbf{z}) \right). \end{aligned}$$

where the bracket on the right hand side is the usual bracket of vector fields. By definition the Lie bracket on $\mathbf{L}(\mathcal{N})$ is given as $[X, Y] = -[R^X, R^Y](p)$ with the sign shift arising due to the computation with right invariant fields. This proves the claim. \square

Having identified the Lie algebra, the next step to develop the Lie theory of \mathcal{N} is to establish the regularity of this Lie group. Let us briefly recall the concept of a regular Lie group. A Lie group G is C^r -semiregular, $r \in \mathbb{N}_0 \cup \{\infty\}$ if for every C^r -curve $\gamma: [0, 1] \rightarrow \mathbf{L}(G)$ the initial value problem

$$\begin{cases} \eta'(t) = T_{\mathbf{1}}\rho_{\eta(t)}(\gamma(t)) & \rho_g(h) := hg \\ \eta(0) = \mathbf{1} \end{cases} \quad (15)$$

has a unique C^{r+1} -solution $\text{Evol}(\gamma) := \eta: [0, 1] \rightarrow G$. Moreover, if in addition the evolution map $\text{evol}: C^r([0, 1], \mathbf{L}(G)) \rightarrow G, \gamma \mapsto \text{Evol}(\gamma)(1)$ is smooth, we say G is C^r -regular. If G is C^∞ -regular (the weakest of the regularity conditions), G is called *regular (in the sense of Milnor)*. To employ advanced techniques in infinite-dimensional Lie theory, one needs to require regularity of the Lie groups involved, cf. [Glö15b]. Note that for a constant curve $\eta(t) \equiv v \in \mathbf{L}(G)$ we simply recover the Lie group exponential $\text{evol}(\eta(t)) = \exp(v)$. Thus every regular infinite-dimensional Lie group admits a Lie group exponential.

Again, since the regularity of the group $\text{Diff}^+(\mathbb{R})$ is a well known fact (see e.g. [KM97, Sch15] for proofs in the convenient and in the Bastiani setting), it is not hard to imagine that these proofs carry over to \mathcal{N} (as we have in principle just added another parameter to the construction). As the modification is again not trivial we supply the necessary details now.

Let us first take a look at the differential equation we need to solve. Consider first the initial condition given by a C^k -curve $\gamma: [0, 1] \rightarrow \mathbf{L}(\mathcal{N})$. The Lie algebra $\mathbf{L}(\mathcal{N})$ has been identified as $C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$, so we can think of elements in the Lie algebra as parameter dependent vector fields on \mathbb{R} (with smooth dependence on a parameter $\mathbf{z} \in \mathbb{S}^2$). To understand the derivative of the right translation $\rho_G(F) = F \cdot G$ where the product is the group product of \mathcal{N} , we apply [Mic80, Corollary 11.6] twice to (14). This shows that if we identify $X \in \mathbf{L}(\mathcal{N}) = C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ then

$$T\rho_F(X) = X \circ (F, \text{pr}_2), \text{ i.e. } T\rho_F(X)(t, \mathbf{z}) = X(F(t, \mathbf{z}), \mathbf{z}).$$

In other words a C^{k+1} -curve $\eta: [0, 1] \rightarrow \mathcal{N}$ solves the differential equation (15) if the adjoint map $\eta^\wedge: [0, 1] \times \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{S}^2$ solves the time and parameter dependent flow equation

$$\frac{\partial}{\partial s} \eta^\wedge(s, t, \mathbf{z}) = \gamma^\wedge(s, \eta^\wedge(s, t, \mathbf{z}), \mathbf{z}). \quad (16)$$

We recall now from [AS15, Proposition 3.20] that the evaluation map

$$\text{ev}_k: C^k([0, 1], C_c^\infty(\mathbb{R} \times \mathbb{S}^2)) \times [0, 1] \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$$

is a $C^{\infty, k}$ -map. As the evaluation ev of smooth functions is a smooth map, the chain rule [AS15, Lemma 3.18] for $C^{r, s}$ -maps implies that

$$\begin{aligned} H: [0, 1] \times \left(C^k([0, 1], C_c^\infty(\mathbb{R} \times \mathbb{S}^2)) \times (\mathbb{R} \times \mathbb{S}^2) \right) &\rightarrow \mathbb{R}, \\ (s, (\gamma, (u, \mathbf{z}))) &\mapsto \text{ev}(\text{ev}_k(\gamma, s), (u, \mathbf{z})) = \gamma(s)(u, \mathbf{z}) \end{aligned}$$

is a mapping of class $C^{k,\infty}$. Hence the right hand side of (16) is a $C^{k,\infty}$ mapping in its entries and we can apply the solution theory for differential equation whose right hand side is of class $C^{r,s}$.

3.17 Lemma *Let $k \in \mathbb{N}_0$. For every $\gamma \in C^k([0, 1], C^\infty(\mathbb{R} \times \mathbb{S}^2))$ the differential equation (16) admits a unique solution $\eta_\gamma^\wedge: [0, 1] \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}$ which is of class $C^{k+1,\infty}$. Moreover, η_γ^\wedge depends smoothly on γ .*

Proof. Instead of the differential equation (16) we consider directly the differential equation

$$\frac{\partial}{\partial s} \eta(s, u, \mathbf{z}) = H(s, \gamma, \eta(s, u, \mathbf{z}), \mathbf{z}) = \gamma(s)(\eta(s, u, \mathbf{z}), \mathbf{z}) \quad (17)$$

where we regard γ and \mathbf{z} as parameters on which the right hand side H depends smoothly. We have already seen that H is a $C^{k,\infty}$ -map (where the k is with respect to the time variable s). We can now apply [AS15, Theorem 5.6] which ensures that for every parameter γ, \mathbf{z} there exists a unique solution $\eta_\gamma^\wedge(s, u, \mathbf{z})$ on some time interval (a priori depending on the parameters) around 0. Note that this is exactly the differentiable dependence on parameters and time we claimed for the solution in the statement of the lemma. Since $\gamma(s)(\cdot, \mathbf{z})$ is a compactly supported time dependent vector field on \mathbb{R} for every choice of the parameters γ, \mathbf{z} , the usual argument (see e.g. [Lee13, Theorem 9.16]) shows that the solution exists on all of $[0, 1]$. \square

We have now constructed candidates for the solution of the regularity problem for \mathcal{N} . Now these candidates need to be identified with smooth mappings taking values in the manifold of mappings.

3.18 Proposition *The Lie group \mathcal{N} is C^r -regular for all $r \in \mathbb{N}_0$.*

Proof. Fix $k \in \mathbb{N}_0$ and $\gamma \in C^k([0, 1], C_c^\infty(\mathbb{R} \times \mathbb{S}^2))$. From Lemma 3.17 we obtain a solution η_γ^\wedge of (16). By construction we have for fixed $s \in [0, 1]$ that $\eta_\gamma^\wedge(s, \cdot) \in C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$, whence we can define $\eta_\gamma: [0, 1] \rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^2)$, $\eta_\gamma(s) := \eta_\gamma^\wedge(s, \cdot)$. Note now that $C_c^\infty(\mathbb{R} \times \mathbb{S}^2) = \lim_{\rightarrow} C_L^\infty(\mathbb{R} \times \mathbb{S}^2)$, where L runs through the compact subsets of $\mathbb{R} \times \mathbb{S}^2$. The inductive limit is compactly regular by [Mic80, 4.7.8]. This means that since $\gamma: [0, 1] \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ is continuous with compact image, the image of γ is already contained in a step of the limit. Thus for γ there exists a compact set $L_1(\gamma) \subseteq \mathbb{R}$ such that $\gamma(s)|_{\mathbb{R} \times \mathbb{S}^2 \setminus L_1(\gamma) \times \mathbb{S}^2} \equiv 0$ for all $s \in [0, 1]$. Note that since the initial condition for the differential equation (17) is $\gamma(0)(u, \mathbf{z}) = u = p(u)$, we see that

$$|\eta_\gamma(s)(u, \mathbf{z}) - u| = \left| \int_0^s \frac{d}{ds} \eta_\gamma(s)(u, \mathbf{z}) ds \right| \leq \int_0^s |\gamma(s)(\eta_\gamma(s)(u, \mathbf{z}))| ds,$$

is bounded by $\sup_{s,u,\mathbf{z}} |\gamma(s)(u, \mathbf{z})| < \infty$. So if $R := \sup_{s,u,\mathbf{z}} |\gamma(s)(u, \mathbf{z})|$ and $L_1 = [a, b]$ then outside of the compact set $L_R := [a - R, b + R]$ the map $\eta_\gamma(s)|_{\mathbb{R} \times \mathbb{S}^2 \setminus (L_R \times \mathbb{S}^2)}$ coincides with p for all $s \in [0, 1]$.

Step 1: η_γ is a C^{k+1} -curve to $C^\infty(\mathbb{R} \times \mathbb{S}^2)$. From the preliminary considerations we see that $\eta_\gamma - p$ takes its image in $C_{L_R \times \mathbb{S}^2}^\infty(\mathbb{R} \times \mathbb{S}^2)$. As the mapping

$$C_{fS}^\infty(\mathbb{R} \times \mathbb{S}^2) \rightarrow C_{fS}^\infty(\mathbb{R} \times \mathbb{S}^2), F \mapsto F + p$$

is smooth (see [Mic80, Remark 4.11] and note that this even restricts on the component of p to a manifold chart), it suffices to prove the C^{k+1} -property of $\eta_\gamma - p$ as a mapping into $C_{L_R \times \mathbb{S}^2}^\infty(\mathbb{R} \times \mathbb{S}^2)$. However, $\eta_\gamma^\wedge: [0, 1] \times (\mathbb{R} \times \mathbb{S}^2) \rightarrow \mathbb{R}$ is a $C^{k+1, \infty}$ -mapping due to Lemma 3.17. Subtracting p , we can exploit the exponential law [AS15, Theorem B] to see that the mapping $\eta_\gamma - p: [0, 1] \rightarrow C_{L_R \times \mathbb{S}^2}^\infty(\mathbb{R} \times \mathbb{S}^2)$ is a C^{k+1} -map. The crucial point here is that the space on the right hand side is endowed with the compact open C^∞ -topology which coincides on the subspace with the fine very strong topology (see e.g. [HS17, Remark 4.5]). Thus η_γ is a C^{k+1} -map.

Step 2: η_γ is a C^{k+1} -map with image in \mathcal{N} . By construction we have $\eta_\gamma(0) = p$ and for every fixed $\mathbf{z} \in \mathbb{S}^2$ the map $\eta_\gamma^\wedge(s, \cdot, \mathbf{z}) \in \text{Diff}(\mathbb{R})$ by the flow property. Now since $p(\cdot, \mathbf{z}) \in \text{Diff}^+(\mathbb{R})$ and $s \mapsto \gamma_\eta(s)(\cdot, \mathbf{z}) \in C_c^\infty(\mathbb{R})$ is continuous for every \mathbf{z} , we see that $\gamma_\eta(s)(\cdot, \mathbf{z}) \in \text{Diff}^+(\mathbb{R})$ for every s, \mathbf{z} as $\text{Diff}^+(\mathbb{R}) \cap C_c^\infty(\mathbb{R})$ is the connected component of the identity in the diffeomorphism group.

Step 3: C^0 -semiregularity and the map evol From our discussion of the differential equation governing regularity of \mathcal{N} , we see that η_γ is a solution for (15) for the initial value γ . Thus \mathcal{N} is C^k -semiregular. To establish C^0 -regularity of the group \mathcal{N} we consider the map

$$\text{evol}: C([0, 1], C_c^\infty(\mathbb{R} \times \mathbb{S}^2)) \rightarrow \mathcal{N} \subseteq C^\infty(\mathbb{R} \times \mathbb{S}^2), \quad \gamma \mapsto ((u, \mathbf{z}) \mapsto \eta_\gamma^\wedge(1, u, \mathbf{z})),$$

where again η_γ^\wedge solves (16). In view of [Glö15b, Lemma 3.1] the group \mathcal{N} will be C^0 -regular if evol is smooth.

Step 4: $e_K: C([0, 1], C_K^\infty(\mathbb{R} \times \mathbb{S}^2)) \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2), \gamma \mapsto \text{evol}(\gamma) - p$ is smooth for every $K \subseteq \mathbb{R} \times \mathbb{S}^2$ compact. Consider for $R > 0$ the open set

$$O_R := \{\gamma \in C([0, 1], C_L^\infty(\mathbb{R} \times \mathbb{S}^2)) \mid \sup_{(s, u, \mathbf{z}) \in [0, 1] \times \mathbb{R} \times \mathbb{S}^2} |\gamma(s)(u, \mathbf{z})| < R\}.$$

As the O_R exhaust $C([0, 1], C_L^\infty(\mathbb{R} \times \mathbb{S}^2))$, it suffices to prove smoothness of e_K on every O_R . To this end, we recall from Lemma 3.17 that the adjoint map

$$e_K^\wedge: O_R \cap C([0, 1], C_K^\infty(\mathbb{R} \times \mathbb{S}^2)) \times \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}, (\gamma, u, \mathbf{z}) \mapsto \eta_\gamma(1)(u, \mathbf{z})$$

is smooth. We now need to create a situation where the exponential law can be applied. As the compact set K is contained in some compact set $L \times \mathbb{S}^2$ we proceed as in Step 1: For every $\gamma \in O_R$, the solution η_γ^\wedge to (16) takes its image in $C_{L_R \times \mathbb{S}^2}^\infty(\mathbb{R} \times \mathbb{S}^2)$ for a compact set L_R only depending on R . Hence we deduce that there is a compact subset $K_R \subseteq \mathbb{R} \times \mathbb{S}^2$ such that $\text{evol}(\gamma) - p$ takes its image in $C_{K_R}^\infty(\mathbb{R} \times \mathbb{S}^2)$. Applying now the exponential law [AS15, Theorem B], the smoothness of e_K follows.

Step 5: \mathcal{N} is C^0 -regular To prove that evol is smooth, we exploit that $C_c^\infty(\mathbb{R} \times \mathbb{S}^2) = \lim_{\rightarrow} C_L^\infty(\mathbb{R} \times \mathbb{S}^2)$ is a compactly regular inductive limit. Thus Mujica's theorem

[Muj83] yields an isomorphism

$$C([0, 1], C_c^\infty(\mathbb{R} \times \mathbb{S}^2)) = C([0, 1], \varinjlim C_L^\infty(\mathbb{R} \times \mathbb{S}^2)) \cong \varinjlim C([0, 1], C_L^\infty(\mathbb{R} \times \mathbb{S}^2)).$$

On each step $C_L^\infty(\mathbb{R} \times \mathbb{S}^2)$ the topology coincides with the compact open C^∞ -topology and we can thus apply the exponential law [AS15, Theorem A] for C^0, ∞ -mappings. Thus $C([0, 1], C_L^\infty(\mathbb{R} \times \mathbb{S}^2)) \cong C_L^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R}))$ for every L . Passing to the limit, we deduce that the mapping

$$\Theta: C([0, 1], C_c^\infty(\mathbb{R} \times \mathbb{S}^2)) \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R})), \quad \gamma \mapsto ((u, \mathbf{z}) \mapsto \gamma(\cdot)(u, \mathbf{z}))$$

is an isomorphism of locally convex spaces. We combine this with the fact that the image of evol is contained in the component of the unit $p \in \mathcal{N}$. Thus it suffices to establish smoothness of the map

$$E: C_c^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R})) \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2), \quad h \mapsto \text{evol}(\Theta^{-1}(h)) - p.$$

Pick now $K \subseteq \mathbb{R} \times \mathbb{S}^2$ compact and consider the restriction of E to $C_K^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R}))$. As $\Theta^{-1}(C_K^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R}))) = C([0, 1], C_L^\infty(\mathbb{R} \times \mathbb{S}^2))$, we have $E|_{C_K^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R}))} = e_K \circ \Theta^{-1}|_{C_K^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R}))}$, whence the restriction is smooth by Step 4.

Let us now show that E is a locally almost local map. For this we work locally on the open sets $\Theta(O_R)$ which by construction exhaust $C_c^\infty(\mathbb{R} \times \mathbb{S}^2, C([0, 1], \mathbb{R}))$. Hence we fix $R > 1$ and may assume that $\sup_s \sup_{(u, \mathbf{z})} |F(u, \mathbf{z})(s)| < R$ for every F we consider. From the definition of E we see that $E(F)|_K = E(G)|_K$ on some compact subset K if and only if $\eta_{\Theta^{-1}(F)}$ and $\eta_{\Theta^{-1}(G)}$ coincide on K . To obtain such sets we define for $n \in \mathbb{Z}$ the open relatively compact sets $U_n^R :=]n - R, n + 2 + R[\times \mathbb{S}^2$ and $V_n :=]n, n + 2[\times \mathbb{S}^2$. Clearly the resulting families are locally finite and cover $\mathbb{R} \times \mathbb{S}^2$. Assume now that F and G coincide on the open set U_n^R . By definition this implies that $\gamma := \Theta(F)$ and $\gamma' := \Theta(G)$ satisfy $\gamma(s)(u, \mathbf{z}) = \gamma'(s)(u, \mathbf{z})$ for all $s \in [0, 1]$, $(u, \mathbf{z}) \in U_n^R$. Moreover, η_γ is the unique solution of the initial value problem

$$\begin{cases} \frac{d}{ds} \eta_\gamma(s)(u, \mathbf{z}) = \gamma(s)(\eta_\gamma(s)(u, \mathbf{z}), \mathbf{z}) & \forall s \in [0, 1], (u, \mathbf{z}) \in \mathbb{R} \times \mathbb{S}^2, \\ \eta_\gamma(0)(u, \mathbf{z}) = u. \end{cases} \quad (18)$$

Now if $(u, \mathbf{z}) \in V_n$, we deduce from $\sup_{(s, u, \mathbf{z})} |\gamma^\wedge(s, u, \mathbf{z})| < R$ that the flow of (18) starting at (u, \mathbf{z}) stays inside of U_n^R . The same observations hold for γ' . Hence $\eta_\gamma(s)(u, \mathbf{z}) = \eta_{\gamma'}(s)(u, \mathbf{z})$ for all $(u, \mathbf{z}) \in V_n$ by uniqueness of solutions to the differential equations (18). We conclude that E restricts on every $\Theta(O_R)$ to an almost local map, whence is locally almost local.

In conclusion E satisfies the prerequisites of Proposition 3.6 and is thus smooth. This concludes the proof. \square

3.4. Lie group structure of the Newman–Unti group

We have seen in the last section that the component \mathcal{N} of the semidirect product $\mathcal{N} \rtimes_\tau \text{SO}^+(3, 1)$ comprising the Newman–Unti group is a Lie group with respect to the

fine very strong topology. Thus we can ask whether the Newman–Unti group can be turned into a Lie group, as it is the semidirect product of two Lie groups. The key is of course the action

$$\tau: \mathcal{N} \times \mathrm{SO}^+(3, 1) \rightarrow \mathcal{N}, (F, f) \mapsto (t, \mathbf{z}) \mapsto K_f^{-1}(\mathbf{z}) \cdot F(K_f(\mathbf{z}), f(\mathbf{z}))$$

which needs to be smooth (with respect to the manifold structure we just constructed on \mathcal{N}). However, we obtain first the following negative result,

3.19 Proposition *The group action $\tau: \mathcal{N} \times \mathrm{SO}^+(3, 1) \rightarrow \mathcal{N}$ is **not** smooth in the sense of convenient analysis. So in particular it is not smooth in the Bastiani sense and thus the Newman–Unti group does not become a Lie group (neither in the Bastiani nor in the convenient setting) if we endow \mathcal{N} with the Lie group structure from Proposition 3.15.*

Proof. We shall construct a smooth curve with values in $\mathcal{N} \times \mathrm{SO}^+(3, 1)$ which is mapped by the action map τ to a non-smooth curve. This not only shows that the action can not be smooth in the Bastiani setting but also in the setting of convenient calculus.

Identify elements in $\mathrm{SO}^+(3, 1)$ with complex 2×2 matrices and consider the smooth curve:

$$\Lambda: \mathbb{R} \rightarrow \mathrm{SO}^+(3, 1), \quad s \mapsto \Lambda_s := \begin{bmatrix} 1+s & 1 \\ s & 1 \end{bmatrix}.$$

Identifying \mathbb{S}^2 with the extended complex plane, via the stereographical projection map $\kappa: \mathbb{S}^2 \rightarrow \hat{\mathbb{C}}$, we define $\phi_s := \kappa^{-1} \circ \Lambda(s) \circ \kappa$. This yields for $\zeta = \kappa(\mathbf{z})$ the relation $K_{\phi_s}(\mathbf{z}) = \frac{1+\|\zeta\|^2}{\|(1+s)\zeta+1\|^2 + \|s\zeta+1\|^2}$. To construct the desired curve with values in the product manifold $\mathcal{N} \times \mathrm{SO}^+(3, 1)$ we consider the element $n: \mathbb{R} \times \mathbb{S}^2, (t, \mathbf{z}) \mapsto t+1$ of \mathcal{N} and define the smooth curve $c(s) := (n, \phi_s)$. We will now show that the curve $\tau \circ c$ with values in \mathcal{N} is not smooth. A trivial computation yields $\tau \circ c(s)(t, \mathbf{z}) = t + K_{\phi_s}(\mathbf{z})^{-1}$ and for $\mathbf{z} = \kappa^{-1}(1)$ we obtain

$$\tau \circ c(s)(t, \kappa^{-1}(1)) = t + \frac{(2+s)^2 + (1+s)^2}{2}.$$

Observe now that for every compact interval $[a, b] \subseteq \mathbb{R}$ containing more than one point it is impossible to find $K \subseteq \mathbb{R} \times \mathbb{S}^2$ compact such that $\tau \circ c(s)|_{(\mathbb{R} \times \mathbb{S}^2) \setminus K}$ is constant in s as no compact subset of $\mathbb{R} \times \mathbb{S}^2$ contains the set $\mathbb{R} \times \{\kappa^{-1}(1)\}$. However, a curve with values in $C_{\mathbb{R}\mathbb{S}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ which violates this condition can not be smooth by [KM97, Lemma 42.5]. Since \mathcal{N} is an open submanifold of $C_{\mathbb{R}\mathbb{S}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ the curve $\tau \circ c$ can not be smooth. \square

The problem identified in the proof of Proposition 3.19 is that the action does not respect the convergence in \mathcal{N} . Namely, we picked an element n which is not contained in the same connected component as the identity element $p \in \mathcal{N}$ and thus did not coincide with p outside of any compact subset of $\mathbb{R} \times \mathbb{S}^2$. We will now show that this is

the only defect of this action. In other words, if we restrict from \mathcal{N} to the connected component

$$\mathcal{N}_0 := \{F \in \mathcal{N} \mid \exists K \subseteq \mathbb{R} \times \mathbb{S}^2 \text{ compact, such that } (F - p)(u, \mathbf{z}) = 0, \forall (u, \mathbf{z}) \notin K\},$$

then we observe the following.

3.20 Lemma *The restriction of τ to the connected component $\mathcal{N}_0 \subseteq \mathcal{N}$ yields a group action*

$$\tau_0: \mathcal{N}_0 \times \text{SO}^+(3, 1) \rightarrow \mathcal{N}_0.$$

Proof. We have to show that $\tau_0(F, f) \in \mathcal{N}_0$ if $F \in \mathcal{N}_0$. To this end, pick $T > 0$ such that outside of the compact set $K := [-T, T] \times \mathbb{S}^2$ we have $F(t, \mathbf{z}) = t = p(t, \mathbf{z})$ if $|t| > T$. Since \mathbb{S}^2 is compact, there is $M := \inf_{\mathbf{z}} K_f(\mathbf{z}) > 0$. Now if $|t| > T/M$ we have $F(K_f(\mathbf{z})t, \mathbf{z}) = K_f(\mathbf{z})t$, whence $\tau_0(F, f)(t, \mathbf{z}) = K_f^{-1}(\mathbf{z})F(K_f(\mathbf{z})t, f(\mathbf{z})) = t = p(t, \mathbf{z})$ for all such t and $\tau_0(F, f) \in \mathcal{N}_0$. \square

For the restricted action we have ruled out the pathology exploited in Proposition 3.19. We will show now that the restricted action is smooth, whence it yields a Lie group structure on the restricted semidirect product.

3.21 Theorem *The unit component $\text{NU}_0 := \mathcal{N}_0 \rtimes_{\tau_0} \text{SO}^+(3, 1)$ of the Newman–Unti group becomes a Lie group with respect to the submanifold structure $\mathcal{N}_0 \subseteq C_{fS}^\infty(\mathbb{R} \times \mathbb{S}^2)$.*

Proof. It suffices to prove that the action τ_0 is smooth. Note that the map

$$\Phi: C^\infty(\mathbb{R} \times \mathbb{S}^2) \supseteq \mathcal{N}_0 \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2), \quad F \mapsto F - p$$

is a chart when restricted to the open set \mathcal{N}_0 . We write $\Omega := \Phi(\mathcal{N}_0)$ for the open image of Φ in $C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ and see that it suffices to show that

$$\tilde{\tau}: \Omega \times \text{SO}^+(3, 1) \rightarrow C_c^\infty(\mathbb{R} \times \mathbb{S}^2), \quad (\alpha, f) \mapsto \left((t, \mathbf{z}) \mapsto K_f^{-1}(\mathbf{z})\alpha(K_f(\mathbf{z})t, f(\mathbf{z})) \right)$$

is smooth. We will proceed in two steps and verify the prerequisites of Proposition 3.6. Before we proceed, it is useful to construct neighborhoods in $\text{SO}^+(3, 1)$ bounding the maximal conformal factor: Since \mathbb{S}^2 is compact, for every $f \in \text{SO}^+(3, 1)$ the constant $\sup_{\mathbf{z} \in \mathbb{S}^2} K_f(\mathbf{z})$ is finite. As the conformal factor $K_f(\mathbf{z})$ is continuous (even smooth) in $(f, \mathbf{z}) \in \text{SO}^+(3, 1) \times \mathbb{S}^2$, the set $O_R = \{g \in \text{SO}^+(3, 1) \mid \sup_{\mathbf{z} \in \mathbb{S}^2} K_g(\mathbf{z}) < R\} \subseteq \text{SO}^+(3, 1)$ is open. Moreover $\text{SO}^+(3, 1) = \bigcup_{R \geq 1} O_R$. If $L \subseteq \mathbb{R} \times \mathbb{S}^2$ is a compact set, we can find $T_L > 0$ such that the compact set $[-T_L, T_L] \times \mathbb{S}^2$ contains L . Enlarging the set even further, we define the compact set $L_R := [-RT_L, RT_L] \times \mathbb{S}^2$ which again contains L for all $R \geq 1$.

Step 1: $\tilde{\tau}$ restricts to a smooth map on $\Phi(\mathcal{N}_0) \cap C_L^\infty(\mathbb{R} \times \mathbb{S}^2) \times \text{SO}^+(3, 1)$. Clearly it suffices to prove the claim for every open set $\Omega \cap C_L^\infty(\mathbb{R} \times \mathbb{S}^2) \times O_R$ where $R \geq 1$. Consider now the adjoint map of the restriction $\tilde{\tau}_R$ of $\tilde{\tau}$ given by

$$\tilde{\tau}_R^\vee: (\Omega \cap C_L^\infty(\mathbb{R} \times \mathbb{S}^2)) \times O_R \times \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}, \quad (\alpha, f, u, \mathbf{z}) \mapsto K_f^{-1}(\mathbf{z}) \cdot \alpha(K_f(\mathbf{z})u, f(\mathbf{z})).$$

By construction we see that $\tilde{\tau}_R^\vee$ vanishes outside of $(\Omega \cap C_L^\infty(\mathbb{R} \times \mathbb{S}^2)) \times O_R \times L_R$. Since L_R is compact, we can apply [AS19, Lemma C.3] to see that $\tilde{\tau}_R$ will be smooth if the adjoint map $\tilde{\tau}_R^\vee$ is smooth. However, $K_f(\mathbf{z}), K_f^{-1}(\mathbf{z})$ and the canonical action $\text{SO}^+(3, 1) \times \mathbb{S}^2 \rightarrow \mathbb{S}^2$ are all smooth. Thus the smoothness of $\tilde{\tau}_R^\vee$ follows directly from the smoothness of the evaluation $\text{ev}: C_L^\infty(\mathbb{R} \times \mathbb{S}^2) \times \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}$ (cf. [AS15, Proposition 3.20]).

Step 2: *The mapping $\tilde{\tau}$ is almost local.* Let $R \in \mathbb{N}$ and define the open set $Q_R := \{f \in O_R \mid \forall \mathbf{z} \in \mathbb{S}^2, 1/R < K_f(\mathbf{z}) < R\}$. Then $\text{SO}^+(3, 1) = \bigcup_{R \in \mathbb{N}} Q_R$. We fix $R \in \mathbb{N}$ and define families of relatively compact open sets as follows: Let A be the set of all integers and define $V_a :=]a - 2, a + 2[\times \mathbb{S}^2$. Then the V_a form a locally finite family of relatively compact sets covering $\mathbb{R} \times \mathbb{S}^2$. To define the sets U_a we need to distinguish several cases: For the first case, assume that $(0, \mathbf{z}) \in V_a$, then we define $U_s :=]\min(R(a - 2), -2R), \max(R(a + 2), 2R)[\times \mathbb{S}^2$. Then by construction we have that if $(u, \mathbf{z}) \in V_a$, then for every $f \in Q_R$ we have $|K_f(\mathbf{z})u| \leq R|u|$, whence $(K_f(\mathbf{z})u, f(\mathbf{z})) \in U_a$ for all $f \in Q_R$. For the other cases, assume that $(0, \mathbf{z}) \notin V_a$. Then we define

$$U_n := \begin{cases}](a - 2)/R, R(a + 2)[\times \mathbb{S}^2 & a - 2 \geq 0 \\]R(a - 2), (a + 2)/R[\times \mathbb{S}^2 & a + 2 \leq 0 \end{cases}$$

Thus for $a - 2 \geq 0$, $(u, \mathbf{z}) \in V_a$ and $f \in Q_R$ we obtain the inequalities

$$\frac{a - 2}{R} < \frac{u}{R} \leq K_f(\mathbf{z})u \leq Ru < R(a + 2)$$

and thus $(K_f(\mathbf{z})u, f(\mathbf{z})) \in U_a$ by construction. Similarly we obtain the same in the case $a + 2 \leq 0$. In conclusion, we have constructed a locally finite family $(U_a)_a$ of relatively compact sets which covers $\mathbb{R} \times \mathbb{S}^2$. If now $f \in Q_R$ and $\alpha|_{U_a} \equiv \beta|_{U_a}$ then we deduce from the construction of the locally finite families that for $(u, \mathbf{z}) \in V_a$ we have

$$\tilde{\tau}(\alpha, f)(u, \mathbf{z}) = K_f^{-1}(\mathbf{z})\alpha(K_f(\mathbf{z})u, \mathbf{z}) = K_f^{-1}(\mathbf{z})\beta(K_f(\mathbf{z})u, \mathbf{z}) = \tilde{\tau}(\beta, f)(u, \mathbf{z}).a$$

This shows that $\tilde{\tau}$ is almost local.

Combining Step 1 and 2, the smoothness of $\tilde{\tau}$ follows from Proposition 3.6. \square

While this indeed constitutes a Lie group structure on the unit component of the NU-group (seen as a submanifold of $C_{fS}^\infty(\mathbb{R} \times \mathbb{S}^2) \times \text{SO}^+(3, 1)$), we note that this subgroup does not accommodate the image of the canonical inclusion of the BMS group. Recall that the inclusion was given by

$$I: \text{BMS} \rightarrow \text{NU}, \quad (F, \phi) \mapsto (p + F \circ \text{pr}_2, \phi).$$

Then the formula for the first component shows that the only element which gets mapped by I into the subgroup $\mathcal{N}_0 \rtimes \text{SO}^+(3, 1)$ is the identity supertranslation $F \equiv 0$.

Note that since \mathcal{N}_0 is the unit component of the Lie group \mathcal{N} we have $\mathbf{L}(\mathcal{N}_0) = \mathbf{L}(\mathcal{N}) = C_c^\infty(\mathbb{R} \times \mathbb{S}^2)$ with the bracket computed in Proposition 3.16. Now NU_0 is a semidirect product of \mathcal{N}_0 and $\text{SO}^+(3, 1)$, whence its Lie algebra is given as the semidirect product

$$\mathbf{L}(\text{NU}_0) = C_c^\infty(\mathbb{R} \times \mathbb{S}^2) \rtimes_{d\tau} \mathbf{L}(\text{SO}^+(3, 1)),$$

where $d\tau$ is the derived action of τ . Moreover, as \mathcal{N} is C^0 -regular by Proposition 3.18, $\mathrm{SO}^+(3, 1)$ is C^0 -regular as a finite dimensional Lie group and C^0 -regularity is an extension property we obtain:

3.22 Corollary *The Lie group NU_0 is C^0 -regular.*

As a consequence of C^0 -regularity we obtain the following properties of NU_0 :

3.23 Lemma *The strong Trotter and the strong commutator property hold for the Lie groups \mathcal{N} and NU_0 .*

Proof. For any C^0 -regular Lie group the strong Trotter property holds due to [Han20, Theorem 1]. Further, it is known that the strong Trotter property implies the (strong) commutator property [Glö15a, Theorem H]. \square

On the other hand we have the following negative results:

3.24 Proposition *The Lie groups \mathcal{N} and NU_0 are not analytic Lie groups.*

Proof. It is well known that the group $\mathrm{Diff}^+(\mathbb{R})$ is not convenient real analytic by [KM97, 43.3 Remarks]. As \mathcal{N} is essentially a parametrized version of $\mathrm{Diff}^+(\mathbb{R})$ these results carry over to \mathcal{N} . So we only need to notice that if a Lie group is not convenient real analytic, it can not be real analytic in the Bastiani sense.

For the group NU_0 the corresponding results follow now at once from the semidirect product structure. Note that as an alternative, we could also have deduced the lack of an analytic structure from the failure of τ to be analytic (this works exactly as the corresponding proof for the BMS group, see [PS21, Proposition 3.9]). \square

Note that as a consequence of Proposition 3.24 for both Lie groups the Baker–Campbell–Hausdorff series does not provide a local model for the multiplication on the Lie algebra. It is yet unclear as to whether the Lie group \mathcal{N} and NU_0 are locally exponential, i.e. that their Lie group exponential induces a local diffeomorphism between a 0-neighborhood in the Lie algebra and a unit neighborhood in the Lie group. Let us note that if \mathcal{N} is not locally exponential this property carries over to NU_0 due to the semidirect product structure (however, if \mathcal{N} was locally exponential it is not easy to deduce local exponentiality of NU_0). Since $\mathrm{Diff}^+(\mathbb{R})$ is not locally exponential (see e.g. [KM97]) we strongly suspect that also \mathcal{N} is not locally exponential. A proof of this statement would require a detailed analysis and adaption of the arguments for the non local exponentiality of $\mathrm{Diff}(\mathbb{R})$ in [Gra88]. This is beyond the scope of the current paper, but we pose the following

3.25 Conjecture *The Lie groups \mathcal{N} and NU_0 are not locally exponential.*

4. Conclusion

We have studied the NU group from the viewpoint of infinite-dimensional Lie group theory. In particular, we have discussed several possible topologies, which turn the supertranslation part \mathcal{N} of the NU group either into a topological group (Proposition 3.11) or into a Lie group (Proposition 3.15). However, we have also shown that only the unit component NU_0 of the complete NU group becomes a Lie group, as the group action is not smooth on the full NU group (Proposition 3.19 and Theorem 3.21). Furthermore, we have shown that both, the Newman–Unti supertranslation group \mathcal{N} as well as the unit component of the Newman–Unti group NU_0 , are regular in the sense of Milnor (Proposition 3.18 and Corollary 3.22). Moreover, we have shown that both these Lie groups are not analytic (Proposition 3.24). Additionally, we have also shown that while the BMS group can be embedded into the NU group as a topological group (Lemma 3.12), this inclusion is not possible on the level of Lie groups, contrary to their Lie algebras, which split as a direct sum (cf. Equation (1)). Finally, we remark our previous article [PS21] in which we have studied the BMS group from a Lie theoretic perspective.

A. Essentials from infinite-dimensional calculus

In this appendix we collect background on infinite-dimensional calculus and Lie groups. We follow [Nee06]. In the following we will always pick a field $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ to be either the real or complex numbers. Then \mathbb{K}^\times denotes the invertible elements in the field \mathbb{K} .

A.1 Definition Let E, F be locally convex spaces over a field \mathbb{K} , $U \subseteq E$ be an open subset, $f: U \rightarrow F$ a map and $r \in \mathbb{N}_0 \cup \{\infty\}$. If it exists, we define for $(x, h) \in U \times E$ the directional derivative

$$df(x, h) := D_h f(x) := \lim_{\mathbb{K}^\times \ni t \rightarrow 0} t^{-1}(f(x + th) - f(x)).$$

We say that f is C^r if the iterated directional derivatives

$$d^{(k)}f(x, y_1, \dots, y_k) := (D_{y_k} D_{y_{k-1}} \cdots D_{y_1} f)(x)$$

exist for all $k \in \mathbb{N}_0$ such that $k \leq r$, $x \in U$ and $y_1, \dots, y_k \in E$ and define continuous maps $d^{(k)}f: U \times E^k \rightarrow F$. If f is $C_{\mathbb{R}}^\infty$, it is called *smooth*. If f is $C_{\mathbb{C}}^\infty$, it is said to be *complex analytic* or *holomorphic* and that f is of class $C_{\mathbb{C}}^\omega$.⁶

A.2 Definition (Complexification of a locally convex space) Let E be a real locally convex topological vector space. Endow the locally convex product $E_{\mathbb{C}} := E \times E$ with the following operation

$$(x + iy).(u, v) := (xu - yv, xv + yu), \quad \forall x, y \in \mathbb{R}, u, v \in E.$$

⁶Recall from [Dah11, Proposition 1.1.16] that $C_{\mathbb{C}}^\infty$ functions are locally given by series of continuous homogeneous polynomials (cf. [BS71b, BS71a]). This justifies the abuse of notation.

The complex vector space $E_{\mathbb{C}}$ is called the *complexification* of E . Identify E with the closed real subspace $E \times \{0\}$ of $E_{\mathbb{C}}$.

A.3 Definition Let E, F be real locally convex spaces and $f: U \rightarrow F$ defined on an open subset $U \subseteq E$. f is called *real analytic* (or $C_{\mathbb{R}}^{\omega}$) if f extends to a $C_{\mathbb{C}}^{\infty}$ -map $\tilde{f}: \tilde{U} \rightarrow F_{\mathbb{C}}$ on an open neighborhood \tilde{U} of U in the complexification $E_{\mathbb{C}}$.

For $r \in \mathbb{N}_0 \cup \{\infty, \omega\}$, being of class $C_{\mathbb{K}}^r$ is a local condition, i.e. if $f|_{U_{\alpha}}$ is $C_{\mathbb{K}}^r$ for every member of an open cover $(U_{\alpha})_{\alpha}$ of its domain, then f is $C_{\mathbb{K}}^r$. (See [Glö02a, pp. 51-52] for the case of $C_{\mathbb{R}}^{\omega}$, the other cases are clear by definition.) In addition, the composition of $C_{\mathbb{K}}^r$ -maps (if possible) is again a $C_{\mathbb{K}}^r$ -map (cf. [Glö02a, Propositions 2.7 and 2.9]).

A.4 (C^r -Manifolds and C^r -mappings between them) For $r \in \mathbb{N}_0 \cup \{\infty, \omega\}$, manifolds modeled on a fixed locally convex space can be defined as usual. The model space of a locally convex manifold and the manifold as a topological space will always be assumed to be Hausdorff spaces. However, we will not assume that infinite-dimensional manifolds are second countable or paracompact. We say M is a *Banach* (or *Fréchet*) manifold if all its modeling spaces are Banach (or Fréchet) spaces.

A.5 Direct products of locally convex manifolds, tangent spaces and tangent bundles as well as C^r -maps between manifolds may be defined as in the finite-dimensional setting. For C^r -manifolds M, N we use the notation $C^r(M, N)$ for the set of all C^r -maps from M to N . Furthermore, we define *locally convex Lie groups* as groups with a C^{∞} -manifold structure turning the group operations into C^{∞} -maps.

A.6 Let E_1, E_2 and F be locally convex spaces and $f: U \times V \rightarrow F$ be a mapping on a product of locally convex subsets $U \subseteq E_1$ and $V \subseteq E_2$ be open. Given $k, \ell \in \mathbb{N}_0 \cup \{\infty\}$, we say that f is $C^{k, \ell}$ if f is continuous and there exist continuous mappings $d^{(i, j)} f: U \times V \times E_1^i \times E_2^j \rightarrow F$ for all $i, j \in \mathbb{N}_0$ such that $i \leq k$ and $j \leq \ell$ such that

$$d^{(i, j)} f(x, y, v_1, \dots, v_i, w_1, \dots, w_j) = (D_{(v_i, 0)} \cdots D_{(v_1, 0)} D_{(0, w_j)} \cdots D_{(0, w_1)} f)(x, y)$$

for all $x \in U^0, y \in V^0$ and $v_1, \dots, v_i \in E_1, w_1, \dots, w_j \in E_2$.

One can also define $C^{k, \ell}$ -maps $M_1 \times M_2 \rightarrow N$ if M_1 is a C^k -manifold, M_2 a C^{ℓ} -manifold and N a $C^{k+\ell}$ -manifold, checking the property in local charts. Moreover, one can also consider mappings which are complex differentiable in one component and only real differentiable in the other. We refer to [AS15] for more information.

A.7 Remark The calculus used in the present article is also known as “Bastiani-calculus”. It is not the only valid choice of calculus beyond the realm of Banach spaces. An alternative approach is the so called “convenient calculus” (see [KM97]).

In general both calculi do not coincide outside of Fréchet spaces (where the Bastiani calculus is stronger as it requires continuity of the underlying mappings). However, in the present paper we are exclusively working with Fréchet spaces and manifolds modeled on Fréchet spaces. Thus there is no difference between our calculus and the convenient calculus in this setting.

B. Auxiliary results for Section 3.3

In this appendix we compile some auxiliary results used in the construction of the Lie group structure for the Newman–Unti group. These results are easy extensions of well known results. However, we were not able to find them in citable form in the literature, whence they are compiled here for the readers convenience. We start with a technical lemma:

B.1 Lemma *If $F \in C_{fS}^\infty(\mathbb{R} \times K)$ for K a compact manifold such that $\frac{\partial}{\partial u}F(u, \mathbf{z}) > 0$, $\forall (u, \mathbf{z}) \in \mathbb{R} \times \mathbb{S}^2$, then there is an open F -neighborhood $O \subseteq C_{fS}^\infty(\mathbb{R} \times K)$ of mappings with this property.*

Proof. Pick a finite atlas $(\psi_j, U_j)_{j=1, \dots, k}$ of K together with compact sets $A_j \subseteq U_j$ such that the A_j cover K . Choose a locally finite family $(K_i)_{i \in \mathbb{N}}$ of compact subsets which cover \mathbb{R} . For every $i \in \mathbb{N}, j = 1, \dots, k$ there is $\varepsilon_{i,j} > 0$ such that for $G \in C^\infty(\mathbb{R} \times K)$ the condition on the directional derivative $d(F - G)$

$$\sup_{(u, \mathbf{z}) \in K_i \times A_j} \left| d \left((F - G) \circ (\text{id}_R \times \psi_j^{-1}) \right) (u, \mathbf{z}; 1, 0) \right| < \varepsilon_{i,j} \quad (19)$$

implies that $G(\cdot, \mathbf{z})$ satisfies $\frac{\partial}{\partial u}G(u, \mathbf{z}) > 0$ for all $(u, \mathbf{z}) \in K_i \times A_j$. By construction, the family of compact sets $(K_i \times A_j)_{i,j}$ is locally finite and we see that there is a basic open neighborhood O_F of F in $C_{fS}^\infty(\mathbb{R} \times K)$ consisting only of mappings which satisfy (19) (compare [HS17, Definition 1.6]). \square

The next result is a parametrized version of [Hir76, Lemma 2.1.3] (the proof is completely analogous apart from the presence of another parameter).

B.2 Lemma *Let K be a compact manifold and $F \in C^\infty(\mathbb{R} \times K)$ be a mapping such that for every $k \in K$ the partial map $F(\cdot, k)$ is an embedding. Furthermore, we fix $W \subseteq \mathbb{R}$ open and relatively compact. Then there exists an open neighborhood $N_F(W)$ of F in the fine very strong topology such that for every $G \in N_F(W)$ the partial maps $G(\cdot, k): \mathbb{R} \rightarrow \mathbb{R}, k \in K$ satisfy*

1. $G(\cdot, k)$ is an immersion,
2. $G(\cdot, k)|_W$ is an embedding.

Proof. Lemma B.1 provides an open neighborhood O of F in the fine very strong topology consisting only of mappings G such that the partial maps $G(\cdot, k)$ are immersions (note that in the proof we assumed the stronger condition that the derivative of the partial maps $F(\cdot, k)$ is everywhere positive. It is easy to see that the same argument holds for mappings whose partial mappings have everywhere negative derivative). We will that we can shrink O such that every $G \in O$ restricts to an embedding $G(\cdot, k)$ on W for all $k \in K$. Assume to the contrary that this is not possible to construct such a neighborhood. Then there must be a sequence $G_N \rightarrow F_N$ of mappings $G_N \in C^\infty(\mathbb{R} \times K)$, where convergence of the sequence is in the sense of uniform convergence of the function and its first derivative on the compact set $\overline{W} \times K$ (actually convergence here is even stronger, but we only need the convergence on the compact set). In other words

$$\lim_{n \rightarrow \infty} \sup_{i=0,1} \sup_{(x,k) \in \overline{W} \times K} \left| \frac{\partial^i}{\partial x} G_n(x, k) - \frac{\partial^i}{\partial x} F(x, k) \right| = 0 \quad (20)$$

By assumption there exist distinct $a_{n,k}, b_{n,k} \in W$ such that $G_n(a_{n,k}, k) = G_n(b_{n,k}, k)$ for every $n \in \mathbb{N}, k \in K$. Now \overline{W} is compact whence $a_{n,k} \rightarrow a_k$ and $b_{n,k} \rightarrow b_k$ for some $a_k, b_k \in \overline{W}$. Evaluating (20) for $i = 0$ this entails $F(a_k, k) = F(b_k, k)$ for every $k \in K$. This entails $a_k = b_k$. Choosing subsequences if necessary, we may assume that the sequence of unit vectors

$$v_{n,k} := \frac{a_{n,k} - b_{n,k}}{|a_{n,k} - b_{n,k}|}$$

converges (either to 1 or -1). Now we take a Taylor expansion in the x -variable while we treat the k -variable as a parameter on which the function in x and its derivative in x depend uniformly continuous (using compactness of K). Now by uniformity of the Taylor expansion (with remainder in integral form) we see that

$$\lim_{n \rightarrow \infty} \sup_{(x,k) \in \overline{W} \times K} \frac{|G_n(a_{n,k}, k) - G_n(b_{n,k}, k) - \frac{\partial}{\partial x} F(b_{n,k}, k)(a_n - b_n)|}{|a_n - b_n|} = 0.$$

Thus $\frac{\partial}{\partial x} F(b_{n,k}, k)v_{n,k} \rightarrow 0$ for every $k \in K$. Note that the absolute values of this sequence converge to $|\frac{\partial}{\partial x} F(b_k, k)|$ and this contradicts that $F(\cdot, k)$ is an immersion for every $k \in K$. \square

B.3 Remark From the proof of Lemma B.2 it is clear that the condition that $G(\cdot, k)$ restricts to an embedding on W for every $k \in K$ requires us only to control the function and its derivative on the compact set $\overline{W} \times K$. Hence if we have a locally finite family $(K_i)_{i \in \mathbb{N}}$ of compact sets covering \mathbb{R} , we can intersect countably many of the neighborhoods constructed in Lemma B.2 and still retain an open set in the fine very strong topology (this is due to the fact that the fine very strong topology admits to control a function and its derivatives on any locally finite family of compacta simultaneously, cf. [HS17] and see also [Hir76, Proof of Theorem 2.1.4]).

This enables us to prove Proposition 3.14, i.e. the set \mathcal{N} is an open subset of $C_{\text{fs}}^\infty(\mathbb{R} \times \mathbb{S}^2)$. While the proof is similar to the classical proof, it requires us to

Proof. Proof of Proposition 3.14 Adapting [Hir76, Theorem 1.7] let us recall that a mapping $f \in C^\infty(\mathbb{R})$ is an element of $\text{Diff}^+(\mathbb{R})$ if and only if f satisfies the following conditions

1. $\frac{d}{du}f(u) > 0, \forall u \in \mathbb{R}$ (i.e. f is an orientation preserving local diffeomorphism)
2. f is proper,
3. f is injective.

To see that this is true, we need only notice that these conditions entail that f is a diffeomorphism: Local diffeomorphisms are open maps, i.e. the image of f is open. Proper maps have closed image, whence by connectedness of \mathbb{R} , f must be surjective. Hence f is a bijective local diffeomorphism and thus a diffeomorphism.

Let us now show that for each $F \in \mathcal{N}$ there is an open neighborhood which contains only mappings $G: \mathbb{R} \times \mathbb{S}^2 \rightarrow \mathbb{R}$ whose partial mappings $G(\cdot, \mathbf{z})$ satisfy 1.-3. Indeed we know already from Lemma B.1 that F admits an open neighborhood O such that every partial map $G(\cdot, \mathbf{z})$ for $G \in O$ satisfies 1.

Step 1: *A neighborhood of proper maps.* The set of proper mappings $\mathcal{P} \subseteq C_{\text{fs}}^\infty(\mathbb{R} \times \mathbb{S}^2)$ is open by [Hir76, Theorem 1.5]. For every $\mathbf{z} \in \mathbb{S}^2$ the inclusion $i_{\mathbf{z}}: \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{S}^2, u \mapsto (u, \mathbf{z})$ is a smooth and proper map. So for every $G \in \mathcal{P}$ the partial map $G(\cdot, \mathbf{z}) = G \circ i_{\mathbf{z}}$ is proper. Since $\mathcal{N} \subseteq \mathcal{P}$ by Lemma 3.10, we intersect O and \mathcal{P} to obtain an open neighborhood of mappings satisfying 1.-2.

Step 2: *An F -neighborhood $O \subseteq C^\infty(\mathbb{R} \times \mathbb{S}^2)$ such that $G(\cdot, \mathbf{z})$ is injective for $G \in O$.* The argument is a variant of [Hir76, Theorem 2.1.4]: Fix the locally finite family $K_i = [i - 1, i + 1], i \in \mathbb{Z}$ of compact sets covering \mathbb{R} . Then the open sets $U_i =]i - 2, i + 2[, i \in \mathbb{Z}$ contain the compact sets $K_i \subseteq U_i$ and form a locally finite family. As $F(\cdot, \mathbf{z})$ is a proper embedding (so in particular a closed injective map), we have for each $\mathbf{z} \in \mathbb{S}^2$ two closed disjoint sets $F(K_i, \mathbf{z})$ and $F(\mathbb{R} \setminus U_i, \mathbf{z})$. By construction, the first set is compact and connected, while the second set is closed and consists of two connected components. The minimal distance $3r_{i,\mathbf{z}}$ (for some $r_{i,\mathbf{z}} > 0$ between the closed sets is realized by one of the pairs $(F(i - 2, \mathbf{z}), F(i - 1, \mathbf{z}))$ or $(F(i + 1, \mathbf{z}), F(i + 2, \mathbf{z}))$).

We define now two disjoint open sets

$$A_{i,\mathbf{z}} := \{x \in \mathbb{R} \mid \sup_{y \in F(K_i, \mathbf{z})} |x - y| < r_{i,\mathbf{z}}\} \text{ and } B_{i,\mathbf{z}} := \{x \in \mathbb{R} \mid \sup_{y \in F(K_i, \mathbf{z})} |x - y| > 2r_{i,\mathbf{z}}\}$$

By construction $F(K_i, \mathbf{z}) \subseteq A_{i,\mathbf{z}}$ and $F(\mathbb{R} \setminus U_i, \mathbf{z}) \subseteq B_{i,\mathbf{z}}$. Using continuity of F and [Eng77, 3.2.10 The Wallace theorem], there is a compact neighborhood $L_{i,\mathbf{z}}$ of \mathbf{z} such that $F(K_i \times L_{i,\mathbf{z}}) \subseteq A_{i,\mathbf{z}}$. We will now shrink $L_{i,\mathbf{z}}$ such that also $F((\mathbb{R} \setminus U_i) \times L_{i,\mathbf{z}}) \subseteq B_{i,\mathbf{z}}$ holds. By construction $\frac{\partial}{\partial u}F(u, s) > 0$ holds for all $s \in \mathbb{S}^2$, hence

$$F(i - 2, s), F(i + 2, s) \in B_{i,\mathbf{z}} \text{ implies } F(\mathbb{R} \setminus U_i, s) \subseteq B_{i,\mathbf{z}}. \quad (21)$$

Applying again continuity of F and Wallace theorem, we can shrink $L_{i,\mathbf{z}}$ such that every $s \in L_{i,\mathbf{z}}$ satisfies (21). Define now the open F -neighborhood

$$N_{F,i,\mathbf{z}} := \{G \in C^\infty(\mathbb{R} \times \mathbb{S}^2) \mid \forall s \in L_{i,\mathbf{z}}, G(K_i, s) \subseteq A_{i,\mathbf{z}}, G(i-1, s), G(i+2, s) \in B_{i,\mathbf{z}}\} \cap O,$$

where O is the open F -neighborhood from Lemma B.1. By construction, (21) implies that every $G \in N_{F,i,\mathbf{z}}$ satisfies $G(\mathbb{R} \setminus U_i, s) \subseteq B_{i,\mathbf{z}}$.

Now we exploit compactness of \mathbb{S}^2 to find finitely many $\mathbf{z}_{i,1}, \dots, \mathbf{z}_{i,j_i}$ such that $\mathbb{S}^2 = \bigcup_{k=1}^{j_i} L_{i,\mathbf{z}_k}$. Intersecting the open sets N_{F,i,\mathbf{z}_k} , we find an open F -neighborhood of functions which satisfy the condition (21) for all $s \in \mathbb{S}^2$, but with different sets $A_{i,\mathbf{z}}, B_{i,\mathbf{z}}$. Note in addition that this neighborhood only controls functions on the compact set $K_i \cup \{i-2, i+2\} \times \mathbb{S}^2$.

Repeat the construction for all i to obtain an open F -neighborhood of functions which satisfy $G(K_i, s) \subseteq A_{i,\mathbf{z}}$ and $G(\mathbb{R} \setminus U_i, s) \subseteq B_{i,\mathbf{z}}$ for all $i \in \mathbb{Z}$, some $\mathbf{z} \in \mathbb{S}^2$ and disjoint open subsets $A_{i,\mathbf{z}}, B_{i,\mathbf{z}}$. Applying Lemma B.2 and Remark B.3 we can shrink the open neighborhood to obtain an open neighborhood Ω of F such that every $G(\cdot, s)|_{U_i}, i \in \mathbb{Z}, s \in \mathbb{S}^2$ is an embedding. Let us show now that the partial maps for every element in Ω are injective. If $x \in K_i, y \in \mathbb{R}$ are two distinct points and $s \in \mathbb{S}^2$, we distinguish two cases: If $y \in U_i$, then $G(x, s) \neq G(y, s)$ as $G(\cdot, s)$ is an embedding on U_i . If $y \in \mathbb{R} \setminus U_i$, we then have $G(x, s) \in A_{i,\mathbf{z}}$ and $G(y, s) \in B_{i,\mathbf{z}}$ for some \mathbf{z} . Since these sets are disjoint, we see that also $G(x, s) \neq G(y, s)$. This concludes the proof. \square

It is possible to extend the proof of Proposition 3.14 to groups of smooth maps defined on $\mathbb{R}^n \times K$ (replacing \mathbb{R} with a multidimensional space and \mathbb{S}^2 with a compact manifold). While it is apparent from the proof that we have only exploited the compactness of \mathbb{S}^2 , the argument for \mathbb{R}^n is slightly more involved. While groups with a compact manifold K replacing \mathbb{S}^2 have been considered in the literature (corresponding to different spacetime dimensions), the extension to \mathbb{R}^n does not seem to be physically relevant. We thus omit the discussion of any details to this extension.

Auxiliary results for the inversion in \mathcal{N}

In this subsection we provide the necessary details to establish smoothness of the inversion mapping for the \mathcal{N} -component of the Newman–Unti group. By construction this is just a parametrized version of the inversion map in the Lie group $\text{Diff}^+(\mathbb{R})$. However, since manifolds of mappings on non-compact manifolds are somewhat delicate, there seems to be no quick way of leveraging the fact that we already know that inversion in $\text{Diff}^+(\mathbb{R})$ is smooth. Instead we have to mimic the proof for the smoothness of inversion in the diffeomorphism group as outlined in [Glö05] but with an additional parameter inserted. We start with some preparation:

B.4 Let K be a compact manifold and $p: \mathbb{R} \times K \rightarrow \mathbb{R}, (x, k) \mapsto x$. Consider the subset

$$G = \left\{ F \in C^\infty(\mathbb{R} \times K) \mid \begin{array}{l} F(\cdot, k) \in \text{Diff}^+(\mathbb{R}) \forall k \in K \text{ and} \\ \exists L \subseteq \mathbb{R} \text{ compact with } (F-p)|_{(\mathbb{R} \setminus L) \times K} = 0 \end{array} \right\}$$

Then G is an open subset of $C_{fS}^\infty(\mathbb{R} \times K)$. To see this, note that Lemma B.1 entails that the set $\{F \in C^\infty(\mathbb{R} \times K) \mid F(\cdot, k) \in \text{Diff}^+(\mathbb{R}), \forall k \in K\}$ is open in the fine very strong topology. In addition, the set

$$\Omega_p := \{F \in C^\infty(\mathbb{R} \times K) \mid \exists L \subseteq \mathbb{R} \text{ compact with } (F - p)|_{(\mathbb{R} \setminus L) \times K} = 0\}$$

is open in this topology. Thus G is open as the intersection of two open sets.

Let us fix some subspaces of $C^\infty(\mathbb{R} \times K)$: First of all we consider the open subset

$$C_c^\infty(\mathbb{R} \times K) := \{F \in C^\infty(\mathbb{R} \times K) \mid \exists L \subseteq \mathbb{R} \text{ with } F|_{(\mathbb{R} \setminus L) \times K} \equiv 0\}$$

and note that the fine very strong topology turns it into a locally convex vector space. Furthermore, we define for every $L \subseteq \mathbb{R}$ compact the subset

$$C_L^\infty(\mathbb{R} \times K) := \{F \in C_c^\infty(\mathbb{R} \times K) \mid F|_{(\mathbb{R} \setminus L) \times K} \equiv 0\}.$$

Then $C_L^\infty(\mathbb{R} \times K)$ is a closed locally convex subspace of $C_c^\infty(\mathbb{R} \times K)$ and one can prove that $C_c^\infty(\mathbb{R} \times K)$ is the locally convex inductive limit of the spaces $C_L^\infty(\mathbb{R} \times K)$ (partially ordered by obvious inclusion as L runs through all compact subset of \mathbb{R}). Moreover, the subspace topology on $C_L^\infty(\mathbb{R} \times K)$ is a Fréchet topology, i.e. a the space is a complete metrizable space. See [Glö02b] for more information.

B.5 Arguing as in Lemma B.1 we define another open subset of $C_{fS}^\infty(\mathbb{R} \times K)$:

$$U_0 := \left\{ F \in C_c^\infty(\mathbb{R} \times K) \mid \sup_{(x,k) \in \mathbb{R} \times K} \left| \frac{\partial}{\partial x} F(x, k) \right| < 1 \right\}.$$

Observe that the map

$$\Phi: \Omega_p \rightarrow C_c^\infty(\mathbb{R} \times K), \quad F \mapsto F - p$$

is a homeomorphism, mapping G onto an open subset of the locally convex space. Indeed the restriction of Φ to G is a chart for the manifold G . We define the open subset $G_0 = \Phi^{-1}(U_0) \cap G$ of G and observe that it is a neighborhood of p .

We now build a local model for the pointwise inversion map $F^{-1}(x, k) := (F(\cdot, k)^{-1})(x)$, by defining

$$U_0 \cap \Phi(G) \rightarrow C_c^\infty(\mathbb{R} \times K), \gamma \mapsto \gamma^* := \Phi((\Phi^{-1}(\gamma))^{-1})$$

Obviously inversion in the group G will be smooth in a neighborhood of p if and only if the local inversion map is smooth on $U_0 \cap \Phi(G) = \Phi(G_0)$.

B.6 A quick computation yields the validity of the following formulae for $\gamma, \eta \in \Phi(G_0)$:

$$\Phi^{-1}(\Phi(\gamma) \circ (\Phi(\eta) \times \text{id}_K)) = \eta + \gamma \circ ((p + \eta) \times \text{id}_K) \quad (22)$$

$$\gamma^* + \gamma \circ ((p + \gamma^*) \times \text{id}_K) = 0 \text{ and } \gamma + \gamma^* \circ ((p + \gamma) \times \text{id}_K) = 0, \quad (23)$$

where (23) is a direct consequence of (22).

B.7 Lemma For every $L \subseteq \mathbb{R}$ compact, the mapping

$$\iota_L: \Phi(G_0) \cap C_L^\infty(\mathbb{R} \times K) \rightarrow C_L^\infty(\mathbb{R} \times K), \quad \gamma \mapsto \gamma^*$$

makes sense and is smooth.

Proof. Let us show that $\iota_L(\gamma)|_{(\mathbb{R} \setminus L) \times K} \equiv 0$. If we fix $k \in K$ then we need to prove that the map $\gamma^*(\cdot, k)$ vanishes outside of L if γ vanishes outside of L . Due to our definition of U_0 this pointwise property is a consequence of [Glö05, Step 3 in the proof of Lemma 5.1.]. Thus ι_L makes sense and takes its image in $C_L^\infty(\mathbb{R} \times K)$. To establish smoothness of ι_L we consider the adjoint map

$$\iota_L^\vee: \Phi(G_0) \cap C_L^\infty(\mathbb{R} \times K) \times (\mathbb{R} \times K) \rightarrow \mathbb{R}, \quad (\gamma, x, k) \mapsto \gamma^*(x, k).$$

By construction every $\gamma \in C_L^\infty(\mathbb{R} \times K)$ vanishes outside of the compact set $L \times K \subseteq \mathbb{R} \times K$. Thus [AS19, Lemma C.3] implies that ι_L will be smooth if ι_L^\vee is smooth. However, for $\iota_L^\vee(\gamma) = \gamma^*$ we obtain from (23) the implicit equation

$$\gamma^*(x, k) + \gamma(x + \gamma^*(x, k), k) = 0, \quad (24)$$

which depends on the parameter $\gamma \in \Phi(G_0) \subseteq C_L^\infty(\mathbb{R} \times K)$. Now the topology of $C_L^\infty(\mathbb{R} \times K)$ coincides with the subspace topology induced by the compact open C^∞ -topology on $C^\infty(\mathbb{R} \times K)$ (cf. [HS17, Remark 4.5]). In particular, [AS15] entails that the evaluation map $\text{ev}: C_L^\infty(\mathbb{R} \times K) \times \mathbb{R} \times K \rightarrow \mathbb{R}$, $(\gamma, x, k) \mapsto \gamma(x, k)$ is smooth. Hence the left hand side is given by the smooth function

$$H: \Phi(G_0) \times \mathbb{R} \times K \times \mathbb{R} \rightarrow \mathbb{R}, \quad H(\eta, x, k, Z) = Z + \text{ev}(\eta, (x + Z, k))$$

Taking the derivative with respect to the Z variable we obtain the continuous linear map

$$d_4 H(\gamma, x, k, Z; \bullet) = \bullet \cdot \left(1 + \frac{\partial}{\partial x} \gamma(x + Z, k) \right).$$

Since $\gamma \in \Phi(G_0) \subseteq U_0$, we see that $\|\text{id}_{\mathbb{R}} - d_4 H(\eta, x, k, Z; \bullet)\|_{\text{op}} = \left| \frac{\partial}{\partial x} \gamma(x + Z, k) \right| < 1$. In other words, $d_4 H(\gamma, x, k, Z)$ is invertible. We can thus again apply the implicit function theorem with parameters in locally convex spaces [Glö06, Theorem 2.3] to deduce that ι_L^\vee is smooth. \square

To extend the smoothness assertion now from the closed subspace $C_L^\infty(\mathbb{R} \times K)$ to all of $C_c^\infty(\mathbb{R} \times K)$ we employ almost local mappings, 3.5 (here without parameter).

B.8 Lemma Let $R > 0$ and O_R be as in Step 2 of the proof of Proposition 3.15. The restriction of the inversion map

$$\Phi(G_0) \cap O_R \rightarrow C_c^\infty(\mathbb{R} \times K), \quad \gamma \mapsto \gamma^*$$

is an almost local map.

Proof. Denote for $x \in \mathbb{R}$ by $B_r(x)$ the r -ball around x in \mathbb{R}

Step 1: For all $r > 0$, $x \in \mathbb{R}$ and $\gamma \in O_R$ we have $B_r(x) \subseteq (p + \gamma)(B_{r+R}(x) \times K)$. Let $y \in B_r(x)$ and recall that $p + \gamma = \Phi^{-1}(\gamma) \in G_0$, whence for every fixed $k \in K$ the map $\Phi^{-1}(\gamma)(\cdot, k)$ is a diffeomorphism of \mathbb{R} there is a unique element y_k such that

$$y = p(y_k, k) + \gamma(y_k, k) = y_k + \gamma(y_k, k), \text{ and } |y_k - x| = |y - \gamma(y_k, k) - x| \leq r + R,$$

where we exploited that $\gamma \in O_R$. In particular, $y_k \in B_{r+R}(x)$ for all $k \in K$.

Step 2: If $R, r > 0$ and $x \in \mathbb{R}$, then for all $\gamma, \eta \in O_R$ such that $\gamma|_{B_{r+R}(x) \times K} = \eta|_{B_{r+R}(x) \times K}$ we have $\gamma^*|_{B_r(x) \times K} = \eta^*|_{B_{r+R}(x) \times K}$. Let $y \in B_r(x)$. By Step 1 we find for every $k \in K$ a unique element $y_k \in B_{r+R}(x)$ such that $y_k + \gamma(y_k, k) = y$. Then from (23) we deduce that

$$\gamma^*(y, k) = \gamma^*(y_k + \gamma(y_k, k), k) = -\gamma(y_k, k) = -\eta(y_k, k) = \eta^*(y_k + \eta(y_k, k), k) = \eta^*(y, k)$$

Step 3: *Inversion is almost local.* Define for $z \in \mathbb{Z}$ the open sets $V_z := B_2(z) \times K$, $W_z := B_3(z) \times K$, $U_z := B_{2+R}(z) \times K$ and $X_z := B_{3+R}(z) \times K$. We deduce from Step 2 that the restriction of the inversion map to $O_R \cap \Phi(G_0)$ together with the families $\{U_z\}_{z \in \mathbb{Z}}$, $\{X_z\}_{z \in \mathbb{Z}}$, $\{V_z\}_{z \in \mathbb{Z}}$ and $\{W_z\}_{z \in \mathbb{Z}}$ satisfy the requirements in the definition of an almost local mapping. \square

References

- [A⁺16] Abbott, B. P. et al. *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116** (2016)(6):061102. doi:10.1103/PhysRevLett.116.061102. ArXiv:1602.03837.
- [AE18] Alessio, F. and Esposito, G. *On the structure and applications of the Bondi–Metzner–Sachs group*. Int. J. Geom. Meth. Mod. Phys. **15** (2018)(02):1830002. doi:10.1142/S0219887818300027. ArXiv:1709.05134.
- [AGS20] Amiri, H., Glöckner, H. and Schmeding, A. *Lie groupoids of mappings taking values in a Lie groupoid*. Arch. Math., Brno **56** (2020)(5):307–356. ArXiv:1811.02888.
- [AS15] Alzaareer, H. and Schmeding, A. *Differentiable mappings on products with different degrees of differentiability in the two factors*. Expo. Math. **33** (2015)(2):184–222. doi:10.1016/j.exmath.2014.07.002. ArXiv:1208.6510.
- [AS19] Amiri, H. and Schmeding, A. *A differentiable monoid of smooth maps on Lie groupoids*. J. Lie Theory **29** (2019)(4):1167–1192
- [BBM62] Bondi, H., van der Burg, M. G. J. and Metzner, A. W. K. *Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems*. Proc. Roy. Soc. Lond. A **269** (1962):21–52. doi:10.1098/rspa.1962.0161

- [BL12] Barnich, G. and Lambert, P.-H. *A Note on the Newman-Unti group and the BMS charge algebra in terms of Newman-Penrose coefficients.* Adv. Math. Phys. **2012** (2012):197385. ArXiv:1102.0589 [gr-qc]
- [BS71a] Bochnak, J. and Siciak, J. *Analytic functions in topological vector spaces.* Studia Math. **39** (1971):77–112
- [BS71b] Bochnak, J. and Siciak, J. *Polynomials and multilinear mappings in topological vector spaces.* Studia Math. **39** (1971):59–76
- [BT10a] Barnich, G. and Troessaert, C. *Aspects of the BMS/CFT correspondence.* JHEP **05** (2010):062. doi:10.1007/JHEP05(2010)062. ArXiv:1001.1541.
- [BT10b] Barnich, G. and Troessaert, C. *Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited.* Phys. Rev. Lett. **105** (2010):111103. doi:10.1103/PhysRevLett.105.111103. ArXiv:0909.2617.
- [Dah11] Dahmen, R. *Direct limit constructions in infinite dimensional Lie theory.* Ph.D. thesis, University of Paderborn 2011. Urn:nbn:de:hbz:466:2-239
- [Eng77] Engelking, R. *General topology. A revised and enlarged translation. Translated by the author,* vol. 60 (PWN - Panstwowe Wydawnictwo Naukowe, Warszawa, 1977)
- [Fri18] Friedrich, H. *Peeling or not peeling — is that the question?* Class. Quant. Grav. **35** (2018)(8):083001. doi:10.1088/1361-6382/aaafdb. ArXiv:1709.07709.
- [Glö02a] Glöckner, H. *Infinite-dimensional Lie groups without completeness restrictions.* In *Geometry and analysis on finite- and infinite-dimensional Lie groups (Będlewo, 2000), Banach Center Publ.*, vol. 55, pp. 43–59 (Polish Acad. Sci., Warsaw, 2002)
- [Glö02b] Glöckner, H. *Lie group structures on quotient groups and universal complexifications for infinite-dimensional Lie groups.* J. Funct. Anal. **194** (2002)(2):347–409
- [Glö05] Glöckner, H. *Diff(\mathbb{R}^n) as a Milnor-Lie group.* Math. Nachr. **278** (2005)(9):1025–1032
- [Glö06] Glöckner, H. *Implicit functions from topological vector spaces to Banach spaces.* Isr. J. Math. **155** (2006):205–252
- [Glö15a] Glöckner, H. *Measurable regularity properties of infinite-dimensional lie groups* 2015. ArXiv:1601.02568.
- [Glö15b] Glöckner, H. *Regularity properties of infinite-dimensional Lie groups, and semiregularity* 2015. ArXiv:1208.0715.

- [Glö21] Glöckner, H. *Semidirect products involving lie groups of compactly supported diffeomorphisms* 2021. In preparation
- [Gra88] Grabowski, J. *Free subgroups of diffeomorphism groups*. Fundam. Math. **131** (1988)(2):103–121. doi:10.4064/fm-131-2-103-121
- [Han20] Hanusch, M. *The strong Trotter property for locally μ -convex Lie groups*. J. Lie Theory **30** (2020)(1):25–32. ArXiv:1802.08923.
- [HE73] Hawking, S. W. and Ellis, G. F. R. *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics (Cambridge University Press, 1973). doi:10.1017/CBO9780511524646
- [Hir76] Hirsch, M. W. *Differential topology*, vol. 33 (Springer, New York, NY, 1976)
- [HN12] Hilgert, J. and Neeb, K.-H. *Structure and Geometry of Lie Groups*. Springer Monographs in Mathematics (Springer, 2012). doi:10.1007/978-0-387-84794-8
- [HS17] Hjelle, E. O. and Schmeding, A. *Strong topologies for spaces of smooth maps with infinite-dimensional target*. Expositiones Mathematicae **35** (2017)(1):13–53. doi:10.1016/j.exmath.2016.07.004
- [KM97] Kriegl, A. and Michor, P. W. *The convenient setting of global analysis*, vol. 53 (Providence, RI: American Mathematical Society, 1997)
- [Lee13] Lee, J. M. *Introduction to smooth manifolds*, vol. 218 (New York, NY: Springer, 2013). doi:10.1007/978-1-4419-9982-5
- [McC72] McCarthy, P. J. *Structure of the Bondi-Metzner-Sachs group*. J. Math. Phys. **13** (1972):1837–1842
- [McC92] McCarthy, P. J. *Real and complex asymptotic symmetries in quantum gravity, irreducible representations, polygons, polyhedra, and the A, D, E series*. Philos. Trans. R. Soc. Lond., Ser. A **338** (1992)(1650):271–299
- [Mic80] Michor, P. W. *Manifolds of differentiable mappings, Shiva Mathematics Series*, vol. 3 (Shiva Publishing Ltd., Nantwich, 1980)
- [Mil84] Milnor, J. *Remarks on infinite dimensional Lie groups*. Relativity, Groups, and Topology II, Les Houches, 1983, B.S. DeWitt, R. Stora, Eds., Elsevier, Amsterdam (1984)
- [Muj83] Mujica, J. *Spaces of continuous functions with values in an inductive limit*. Functional analysis, holomorphy, and approximation theory, Proc. Semin., Rio de Janeiro 1979, Lect. Notes Pure Appl. Math. 83, 359–367 (1983). 1983
- [Nee06] Neeb, K.-H. *Towards a Lie theory of locally convex groups*. Jpn. J. Math. **1** (2006)(2):291–468. doi:10.1007/s11537-006-0606-y. ArXiv:1501.06269.

- [New89] Newman, R. *The global structure of simple space-times*. Commun. Math. Phys. **123** (1989):17–52. doi:https://doi.org/10.1007/BF01244016
- [NU62] Newman, E. T. and Unti, T. W. J. *Behavior of Asymptotically Flat Empty Spaces*. J. Math. Phys. **3** (1962)(5):891. doi:10.1063/1.1724303
- [Pen63] Penrose, R. *Asymptotic properties of fields and space-times*. Phys. Rev. Lett. **10** (1963):66–68
- [Pen64] Penrose, R. *Conformal treatment of infinity*. Relativity, groups and topology: the 1963 Les Houches lectures (1964). Republished in Gen. Relativ. Gravit. 43:901–922, 2011.
- [Pen65] Penrose, R. *Zero rest mass fields including gravitation: Asymptotic behavior*. Proc. Roy. Soc. Lond. A **284** (1965):159
- [Pen68] Penrose, R. *Structure of space-time*. In *Battelle Rencontres* (1968)
- [Pen74] Penrose, R. *Relativistic Symmetry Groups*, pp. 1–58 (Springer Netherlands, Dordrecht, 1974). doi:10.1007/978-94-010-2144-9_1
- [PS21] Prinz, D. and Schmeding, A. *Lie theory for asymptotic symmetries in general relativity: The BMS group* 2021. ArXiv:2106.12513 [gr-qc]
- [Ruz20] Ruzziconi, R. *On the Various Extensions of the BMS Group*. Ph.D. thesis, Brussels U. and Intl. Solvay Inst., Brussels 2020. ArXiv:2009.01926.
- [Sac62] Sachs, R. K. *Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times*. Proc. Roy. Soc. Lond. A **270** (1962):103–126. doi:10.1098/rspa.1962.0206
- [Sch15] Schmeding, A. *The diffeomorphism group of a non-compact orbifold*. Dissertationes Math. **507** (2015):179. doi:10.4064/dm507-0-1. ArXiv:1301.5551.
- [SSW75] Schmidt, B., Sommers, P. and Walker, M. *A characterization of the bondi-metzner-sachs group*. General Relativity and Gravitation **6** (1975):489–497. doi:10.1007/BF00762453