

From the AKNS system to the matrix Schrödinger equation with vanishing potentials: Direct and inverse problems*

Francesco Demontis[†], Cornelis van der Mee*

Abstract

We relate the scattering theory of the focusing AKNS system with vanishing boundary conditions to that of the matrix Schrödinger equation. The corresponding Miura transformation which allows this connection, converts the focusing matrix nonlinear Schrödinger (NLS) equation into a new nonlocal integrable equation. We apply the matrix triplet method to derive the multisoliton solutions of the nonlocal integrable equation, thus proposing a new method to solve the matrix NLS equation.

1 Introduction

For nearly 50 years nonlinear Schrödinger (NLS) equations have served as the basic models for describing surface waves on deep waters [1, 5, 59], signals along optical fibres [38, 37, 54], particle states in Bose-Einstein condensates [50, 51, 40], plasma oscillations [57], and magnetic spin waves [21, 58]. NLS equations with solutions decaying at infinity have been studied in detail by means of the inverse scattering transform [3, 5, 19, 29, 4], where a time invariant canonical transformation converts the initial-value problem of the matrix NLS equation into the (elementary) time evolution of the scattering data of the so-called AKNS system.

*LaTeX compilation date and time: 09/02/2022 at 19:22:00

[†]Dipartimento di Matematica e Informatica, Università di Cagliari, Via Ospedale 72, 09124 Cagliari, Italy. Email: fdemontis@unica.it, cornelis110553@gmail.com

In this article we present an apparently new method for solving matrix NLS equations inspired by the Miura transformation [5, 2] which traditionally allows one to derive Korteweg-de Vries (KdV) solutions from modified Korteweg-de Vries (mKdV) solutions (but not necessarily vice versa). By a similar transformation the direct and inverse scattering theory of the focusing $n = m_1 + m_2$ AKNS system can be related to the direct and inverse scattering theory of a matrix Schrödinger equation whose $n \times n$ matrix potential \mathcal{Q} satisfies the adjoint symmetry relation

$$\mathcal{Q}^\dagger = \sigma_3 \mathcal{Q} \sigma_3, \quad (1.1)$$

where $\sigma_3 = I_{m_1} \oplus (-I_{m_2})$ and the dagger denotes the matrix conjugate transpose. The traditional application of the matrix Schrödinger equation to quantum graphs, quantum wires, and quantum mechanical scattering of particles with internal structure [14, 15, 16, 17, 18, 27, 31, 32, 33, 34, 35, 36, 42, 43, 44, 45, 46, 47, 48] has led to the almost exclusive development of matrix Schrödinger scattering theory for selfadjoint potentials for which $\mathcal{Q}^\dagger = \mathcal{Q}$. After the seminal papers on the scalar Schrödinger scattering theory by Faddeev [28], Deift and Trubowitz [22], and others (see [20]), in the matrix case the half-line theory has been described in detail by Agranovich and Marchenko [7] and by Aktosun and Weder [12, 13] and the full-line theory by Wadati and Kamijo [56]; the most general small energy asymptotics of the scattering data, crucial to developing the direct and inverse scattering theory rigorously, is due to Klaus [41] in the scalar case and to Aktosun et al. [10] in the matrix case.

Energy losses in quantum graphs, quantum wires, and particles with internal structure naturally lead to matrix Schrödinger potentials whose imaginary part $[\mathcal{Q} - \mathcal{Q}^\dagger]/2i$ has constant sign. In this article we thus require a modified direct and inverse scattering theory when solving the matrix NLS equation by using the matrix Schrödinger equation for potentials satisfying

$$\mathcal{Q}^\dagger = \sigma_3 \mathcal{Q} \sigma_3. \quad (1.2)$$

In this article we depart from the direct and inverse scattering theory of the focusing AKNS system [3, 4, 29]

$$v_x = (-ik\sigma_3 + \mathcal{Q})v, \quad (1.3)$$

where $v = v(x, k)$ is a vector function with $n = m_1 + m_2$ components and the potential \mathcal{Q} anticommutes with $\sigma_3 = I_{m_1} \oplus (-I_{m_2})$. Letting $L =$

$i\sigma_3[(d/dx)I_n - \mathcal{Q}]$ stand for the AKNS Hamiltonian, we easily verify that L^2 is the matrix Schrödinger Hamiltonian given by

$$\begin{aligned} L^2v &= -\sigma_3[(d/dx)I_n - \mathcal{Q}]\sigma_3[(d/dx)I_n - \mathcal{Q}]v \\ &= -[(d/dx)I_n + \mathcal{Q}][(d/dx)I_n - \mathcal{Q}]v \\ &= -v_{xx} + \mathcal{Q}^2v - \mathcal{Q}v_x + (\mathcal{Q}v)_x = -v_{xx} + \mathbf{Q}v, \end{aligned}$$

where

$$\mathbf{Q} = \mathcal{Q}^2 + \mathcal{Q}_x \quad (1.4)$$

is a matrix Schrödinger potential obtained from \mathcal{Q} by the Miura transform (1.4). Assuming that the entries of \mathcal{Q} and \mathcal{Q}_x belong to the weighted L^1 -space $L^1(\mathbb{R}; (1 + |x|)dx)$, we arrive from (1.3) at the matrix Schrödinger equation

$$-v_{xx} + \mathbf{Q}v = k^2v, \quad (1.5)$$

where \mathbf{Q} is a so-called Faddeev class potential (i.e., its entries belong to $L^1(\mathbb{R}; (1 + |x|)dx)$) satisfying the adjoint symmetry relation (1.2).

It is well-known [59, 3, 4, 29] that for a suitable time evolution of the AKNS scattering data the potential $\mathcal{Q} = \mathcal{Q}(x; t)$ satisfies the matrix NLS equation

$$i\mathcal{Q}_t + \mathcal{Q}_{xx} - 2\mathcal{Q}^3 = 0_{n \times n}, \quad (1.6)$$

where \mathcal{Q} anticommutes with σ_3 . Direct AKNS scattering theory involves Jost solutions and certain scattering coefficients which are analytic in k in either upper half complex plane \mathbb{C}^+ or the lower half complex plane \mathbb{C}^- , with continuous extensions up to the real k -line [3, 4, 29]. On the other hand, traditionally direct matrix Schrödinger scattering theory involves Jost solutions and certain scattering coefficients which are continuous in $k \in \mathbb{C}^+ \cup \mathbb{R}$ and analytic in $k \in \mathbb{C}^+$ ([7, 12, 13] for the half-line theory and [56, 10] for the full line theory). In fact, the values $\pm k$ of the AKNS spectral variable can be made to correspond to the matrix Schrödinger spectral variable k , which greatly simplifies direct and inverse scattering theory. It is therefore natural to wonder which nonlinear evolution equation $\mathbf{Q}(x; t)$ is satisfied when $\mathcal{Q}(x; t)$ is a matrix NLS solution. We are thus lead to a nonlocal integrable equation for $\mathbf{Q}(x; t)$ to be derived from a Lax pair.

The principal objective of this article is to derive the multisoliton solutions of the focusing NLS system by applying the Miura transform (1.4) to the matrix Schrödinger equation which is then solved explicitly by the so-called

matrix triplet method (see [55] and references therein). These solutions are obtained by solving the corresponding Marchenko integral equations. Once obtained, these solutions are verified by direct substitution into the nonlocal integrable equation satisfied by $\mathbf{Q}(x;t)$.

Let us discuss the contents of this article. In Sec. 2 we derive the non-local integrable equation for \mathbf{Q} from a Lax pair. Next, in Sections 3-4 we develop the direct and inverse scattering theory of the matrix Schrödinger equation (1.5) for Faddeev class potentials \mathbf{Q} satisfying (1.2). In particular, we introduce the Jost solutions and the scattering coefficients, write them as Fourier transforms of L^1 -functions, and derive the Marchenko integral equations to solve the inverse scattering problem. The rather technical proof of the adjoint symmetry relations of the Marchenko integral kernels is deferred to Appendix B. The relationship between our Jost solutions and scattering coefficients and those prevailing in AKNS scattering theory will be worked out in Sec. 5. We then go on to derive the time evolution of the scattering data in Sec. 6. In Sec. 7 we apply the so-called matrix triplet method to derive the multisoliton solutions. Three appendices have been attached. In Appendix A we derive the nonlocal integrable equation for \mathbf{Q} directly from the NLS equation for \mathcal{Q} without using Lax pairs. In Appendix B we derive the adjoint symmetry relations for the Marchenko integral kernels. Finally, in Appendix C we prove by direct substitution that the multisoliton solutions satisfy the nonlocal integrable equation. Moreover, the emphasis on the methods used in this article allows one to construct the reflectionless solutions of the hierarchy of equations with Lax pairs $\{(L^j, A)\}_{j=1}^{\infty}$, where L is the AKNS Hamiltonian and A describes the time evolution (see Sec. 2). In fact, we present the details only for the first two equations of the hierarchy.

We adopt blackboard boldface symbols for many of the quantities pertaining to the AKNS system, thus deviating from the notational system adopted in [4]. Boldface symbols are reserved for many of the quantities pertaining to the matrix Schrödinger equation. We also deviate from the praxis of [3, 4] in allowing right and left to correspond to the real line endpoints of transmission, both in the AKNS case and in the (matrix) Schrödinger case. Thus we deviate from the right vs. left conventions adopted in [3, 4].

2 Lax Pair for the new integrable model

As to be indicated shortly, the matrix NLS system is governed by a Lax pair $\{L, A\}$ of linear operators [49, 5, 26], where $Lv = kv$ is the AKNS eigenvalue problem and $v_t = Av$ describes the time evolution. Then the matrix NLS system is equivalent to the zero curvature condition

$$L_t + LA - AL = \mathbf{0},$$

where $\mathbf{0}$ denotes the zero operator on a suitable function space. We now observe that $\{L^2, A\}$ is also a Lax pair of linear operators, where $L^2v = k^2v$ is the matrix Schrödinger eigenvalue problem and $v_t = Av$ describes the time evolution. Then the accompanying nonlinear evolution equation is equivalent to the zero curvature condition

$$(L^2)_t + L^2A - AL^2 = \mathbf{0}.$$

Putting

$$L = i\sigma_3(\partial_x I_n - \mathcal{Q}), \quad (2.1a)$$

$$A = i\sigma_3(2\partial_x^2 I_n - 2\mathcal{Q}\partial_x - \mathcal{B}_0), \quad (2.1b)$$

where $n = m_1 + m_2$ and \mathcal{B}_0 is to be determined, we compute

$$\begin{aligned} \mathbf{0} &= -(L_t + LA - AL) \\ &= i\sigma_3 \mathcal{Q}_t + \sigma_3(\partial_x I_n - \mathcal{Q})\sigma_3(2\partial_x^2 I_n - 2\mathcal{Q}\partial_x - \mathcal{B}_0) \\ &\quad - \sigma_3(2\partial_x^2 I_n - 2\mathcal{Q}\partial_x - \mathcal{B}_0)\sigma_3(\partial_x I_n - \mathcal{Q}) \\ &= i\sigma_3 \mathcal{Q}_t + (\partial_x I_n + \mathcal{Q})(2\partial_x^2 I_n - 2\mathcal{Q}\partial_x - \mathcal{B}_0) \\ &\quad - (2\partial_x^2 I_n + 2\mathcal{Q}\partial_x - \sigma_3 \mathcal{B}_0 \sigma_3)(\partial_x I_n - \mathcal{Q}) \\ &= i\sigma_3 \mathcal{Q}_t + 2\partial_x^3 I_n - 2\mathcal{Q}_x \partial_x - 2\mathcal{Q}\partial_x^2 - (\mathcal{B}_0)_x - \mathcal{B}_0 \partial_x + 2\mathcal{Q}\partial_x^2 - 2\mathcal{Q}^2 \partial_x - \mathcal{Q}\mathcal{B}_0 - 2\partial_x^3 I_n \\ &\quad + 2\mathcal{Q}_{xx} + 4\mathcal{Q}_x \partial_x + 2\mathcal{Q}\partial_x^2 - 2\mathcal{Q}\partial_x^2 + 2\mathcal{Q}\mathcal{Q}_x + 2\mathcal{Q}^2 \partial_x + \sigma_3 \mathcal{B}_0 \sigma_3 \partial_x - \sigma_3 \mathcal{B}_0 \sigma_3 \mathcal{Q} \\ &= i\sigma_3 \mathcal{Q}_t - 2(-\mathcal{Q}_x + \frac{1}{2}[\mathcal{B}_0 - \sigma_3 \mathcal{B}_0 \sigma_3])\partial_x \\ &\quad - (\mathcal{B}_0)_x - \mathcal{Q}\mathcal{B}_0 + 2\mathcal{Q}_{xx} + 2\mathcal{Q}\mathcal{Q}_x - \sigma_3 \mathcal{B}_0 \sigma_3 \mathcal{Q}. \end{aligned}$$

To cancel the ∂_x terms, \mathcal{B}_0 should be the sum of \mathcal{Q}_x and a matrix commuting with σ_3 . Thus taking $\mathcal{B}_0 = \mathcal{Q}^2 + \mathcal{Q}_x = \mathcal{Q}$, we get the matrix NLS equation

$$i\sigma_3 \mathcal{Q}_t + \mathcal{Q}_{xx} - 2\mathcal{Q}^3 = 0_{n \times n}. \quad (2.2)$$

Putting $\mathcal{L} = L^2 = -\partial_x^2 + \mathcal{Q}$, we compute

$$\begin{aligned}
i\sigma_3[\mathcal{L}_t + \mathcal{L}A - A\mathcal{L}] &= i\sigma_3\mathcal{Q}_t \\
&- (-\partial_x^2 + \sigma_3\mathcal{Q}\sigma_3) [2\partial_x^2 - 2\mathcal{Q}\partial_x - \mathcal{Q}] + [2\partial_x^2 - 2\mathcal{Q}\partial_x - \mathcal{Q}] (-\partial_x^2 + \mathcal{Q}) \\
&= i\sigma_3\mathcal{Q}_t + 4(-\mathcal{Q}_x + \frac{1}{2}[\mathcal{Q} - \sigma_3\mathcal{Q}\sigma_3])\partial_x^2 \\
&+ 2(-\mathcal{Q}_{xx} + \mathcal{Q}_x + \sigma_3\mathcal{Q}\sigma_3\mathcal{Q} - \mathcal{Q}\mathcal{Q})\partial_x \\
&+ \mathcal{Q}_{xx} + \sigma_3\mathcal{Q}\sigma_3\mathcal{Q} - 2\mathcal{Q}\mathcal{Q}_x - \mathcal{Q}^2.
\end{aligned}$$

Then (1.4) and $\sigma_3\mathcal{Q} = -\mathcal{Q}\sigma_3$ imply that the terms in the third member involving ∂_x^2 and ∂_x vanish. Therefore, we have arrived at the nonlinear evolution equation

$$i\sigma_3\mathcal{Q}_t + \mathcal{Q}_{xx} - \mathcal{Q}^2 + \sigma_3\mathcal{Q}\sigma_3\mathcal{Q} - 2\mathcal{Q}\mathcal{Q}_x = 0_{n \times n}, \quad (2.3)$$

where

$$\mathcal{Q}(x; t) = - \int_x^\infty dy \frac{1}{2} (\mathcal{Q} - \sigma_3\mathcal{Q}\sigma_3) = \int_{-\infty}^x dy \frac{1}{2} (\mathcal{Q} - \sigma_3\mathcal{Q}\sigma_3). \quad (2.4)$$

A proof of the integrability of (2.3) based directly on (1.4) and (1.6) will be given in Appendix A. Finally, using (1.4) in (2.3) and singling out the block off-diagonal component we easily get the derivative of (2.2) with respect to x and hence (2.2) itself; the block diagonal component yields the commutator of (2.2) and \mathcal{Q} .

3 Direct Scattering Problem

In this section we introduce the Jost solutions and scattering coefficients for the matrix Schrödinger equation (1.5) with Faddeev class potential \mathcal{Q} satisfying (1.2). For the scalar Schrödinger equation with real Faddeev class potential the direct scattering theory is well documented [28, 22, 19, 20, 2]. For selfadjoint potentials, the matrix theory on the half-line is discussed at length in [7, 12, 13], whereas that on the full-line can be found in [56, 10]. Here we give the modifications caused by the adjoint symmetry (1.2).

3.1 Jost solutions of the matrix Schrödinger equation

Let us define the *Jost solution from the left* $F_l(x, k)$ and the *Jost solution from the right* $F_r(x, k)$ as those solutions of the matrix Schrödinger equation

(1.5) which satisfy the asymptotic conditions

$$F_l(x, k) = e^{ikx} [I_n + o(1)], \quad x \rightarrow +\infty, \quad (3.1a)$$

$$F_r(x, k) = e^{-ikx} [I_n + o(1)], \quad x \rightarrow -\infty, \quad (3.1b)$$

where $n = m_1 + m_2$. Calling $m_l(x, k) = e^{-ikx} F_l(x, k)$ and $m_r(x, k) = e^{ikx} F_r(x, k)$ *Faddeev functions*, we easily define them as the unique solutions of the Volterra integral equations

$$m_l(x, k) = I_n + \int_x^\infty dy \frac{e^{2ik(y-x)} - 1}{2ik} \mathbf{Q}(y) m_l(y, k), \quad (3.2a)$$

$$m_r(x, k) = I_n + \int_{-\infty}^x dy \frac{e^{2ik(x-y)} - 1}{2ik} \mathbf{Q}(y) m_r(y, k). \quad (3.2b)$$

Then, for each $x \in \mathbb{R}$, $m_l(x, k)$ and $m_r(x, k)$ are continuous in $k \in \mathbb{C}^+ \cup \mathbb{R}$, are analytic in $k \in \mathbb{C}^+$, and tend to I_n as $k \rightarrow \infty$ from within $\mathbb{C}^+ \cup \mathbb{R}$. For $0 \neq k \in \mathbb{R}$ we can reshuffle (3.2) and arrive at the asymptotic relations

$$F_l(x, k) = e^{ikx} A_l(k) + e^{-ikx} B_l(k) + o(1), \quad x \rightarrow -\infty, \quad (3.3a)$$

$$F_r(x, k) = e^{-ikx} A_r(k) + e^{ikx} B_r(k) + o(1), \quad x \rightarrow +\infty, \quad (3.3b)$$

where

$$A_{r,l}(k) = I_n - \frac{1}{2ik} \int_{-\infty}^\infty dy \mathbf{Q}(y) m_{r,l}(y, k), \quad (3.4a)$$

$$B_{r,l}(k) = \frac{1}{2ik} \int_{-\infty}^\infty dy e^{\mp 2iky} \mathbf{Q}(y) m_{r,l}(y, k). \quad (3.4b)$$

Thus $A_{r,l}(k)$ is continuous in $0 \neq k \in \mathbb{C}^+ \cup \mathbb{R}$, is analytic in $k \in \mathbb{C}^+$, and tends to I_n as $k \rightarrow \infty$ from within $\mathbb{C}^+ \cup \mathbb{R}$, while $2ik[I_n - A_{r,l}(k)]$ has the finite limit $-\Delta_{r,l} = \int_{-\infty}^\infty dy \mathbf{Q}(y) m_{r,l}(y, k)$ as $k \rightarrow 0$ from within $\mathbb{C}^+ \cup \mathbb{R}$. By the same token, $B_{r,l}(k)$ is continuous in $0 \neq k \in \mathbb{R}$, vanishes as $k \rightarrow \pm\infty$, and satisfies $2ikB_{r,l}(k) \rightarrow -\Delta_{r,l}$ as $k \rightarrow 0$ along the real k -axis.

Putting

$$\mathbf{F}_l(x, k) = \begin{pmatrix} F_l(x, -k) & F_l(x, k) \\ F'_l(x, -k) & F'_l(x, k) \end{pmatrix}, \quad \mathbf{F}_r(x, k) = \begin{pmatrix} F_r(x, k) & F_r(x, -k) \\ F'_r(x, k) & F'_r(x, -k) \end{pmatrix},$$

where the prime denotes differentiation with respect to x , we obtain

$$\mathbf{F}_r(x, k) = \mathbf{F}_l(x, k) \begin{pmatrix} A_r(k) & B_r(-k) \\ B_r(k) & A_r(-k) \end{pmatrix}, \quad (3.5a)$$

$$\mathbf{F}_l(x, k) = \mathbf{F}_r(x, k) \begin{pmatrix} A_l(-k) & B_l(k) \\ B_l(-k) & A_l(k) \end{pmatrix}, \quad (3.5b)$$

where $0 \neq k \in \mathbb{R}$. Using that $\mathbf{F}_{r,l}(x, k)$ satisfies the linear first order system

$$\begin{pmatrix} V \\ V' \end{pmatrix}' = \begin{pmatrix} 0_{n \times n} & I_n \\ \mathbf{Q}(x) - k^2 I_n & 0_{n \times n} \end{pmatrix} \begin{pmatrix} V \\ V' \end{pmatrix} \quad (3.6)$$

with traceless system matrix, we see that, for $0 \neq k \in \mathbb{R}$, $\mathbf{F}_{r,l}(x, k)$ has a determinant not depending on $x \in \mathbb{R}$. Using (3.1) we easily verify that $\det \mathbf{F}_{r,l}(x, k) = (2ik)^n$ for $0 \neq k \in \mathbb{R}$.

Putting

$$\begin{aligned} \Phi(x, k) &= \begin{pmatrix} F_r(x, k) & F_l(x, k) \\ F_r'(x, k) & F_l'(x, k) \end{pmatrix} \\ &= \mathbf{F}_r(x, k) \begin{pmatrix} I_n & B_l(k) \\ 0_{n \times n} & A_l(k) \end{pmatrix} = \mathbf{F}_l(x, k) \begin{pmatrix} A_r(k) & 0_{n \times n} \\ B_r(k) & I_n \end{pmatrix}, \end{aligned} \quad (3.7)$$

we easily see that, for $0 \neq k \in \mathbb{R}$, $\det A_r(k) = \det A_l(k)$. By analytic continuation we get $\det A_r(k) = \det A_l(k)$ for $0 \neq k \in \mathbb{C}^+ \cup \mathbb{R}$.

Letting $\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ denote the second Pauli matrix, we define

$$\sigma_2 \otimes \sigma_3 = \begin{pmatrix} 0_{n \times n} & -i\sigma_3 \\ i\sigma_3 & 0_{n \times n} \end{pmatrix}$$

as the Kronecker product of σ_2 and σ_3 (cf. [39]).

Proposition 3.1 *For $0 \neq k \in \mathbb{C}^+ \cup \mathbb{R}$, let $\mathbf{V}(x, k)$ and $\mathbf{W}(x, k)$ be two size compatible matrix solutions of the linear first order system (3.6). Then*

$$\mathbf{W}(x, -k^*)^\dagger (\sigma_2 \otimes \sigma_3) \mathbf{V}(x, k)$$

is independent of $x \in \mathbb{R}$. Similarly, for $0 \neq k \in \mathbb{R}$ the matrix

$$\mathbf{W}(x, k)^\dagger (\sigma_2 \otimes \sigma_3) \mathbf{V}(x, k)$$

is independent of $x \in \mathbb{R}$.

Proof. It is easily verified that

$$(\sigma_2 \otimes \sigma_3) \begin{pmatrix} 0_{n \times n} & I_n \\ \mathbf{Q}(x) - k^2 I_n & 0_{n \times n} \end{pmatrix} (\sigma_2 \otimes \sigma_3) = - \begin{pmatrix} 0_{n \times n} & I_n \\ \mathbf{Q}(x) - (-k^*)^2 I_n & 0_{n \times n} \end{pmatrix}^\dagger.$$

Then

$$\begin{aligned} & \frac{\partial}{\partial x} [\mathbf{W}(x, -k^*)^\dagger (\sigma_2 \otimes \sigma_3) \mathbf{V}(x, k)] \\ &= \mathbf{W}(x, -k^*)^\dagger \begin{pmatrix} 0_{n \times n} & I_n \\ \mathbf{Q}(x) - (-k^*)^2 I_n & 0_{n \times n} \end{pmatrix}^\dagger (\sigma_2 \otimes \sigma_3) \mathbf{V}(x, k) \\ &+ \mathbf{W}(x, -k^*)^\dagger (\sigma_2 \otimes \sigma_3) \begin{pmatrix} 0_{n \times n} & I_n \\ \mathbf{Q}(x) - k^2 I_n & 0_{n \times n} \end{pmatrix} \mathbf{V}(x, k) = 0_{2n \times 2n}, \end{aligned}$$

as claimed. The second part follows in the same way. \blacksquare

Let us now apply either part of Proposition 3.1 to derive identities for the A and B coefficients by equating the asymptotics as $x \rightarrow +\infty$ to the asymptotics as $x \rightarrow -\infty$. Using the second part for $\mathbf{V} = \mathbf{W} = \Phi$ we get

$$A_{r,l}(k)^\dagger \sigma_3 A_{r,l}(k) - B_{r,l}(k)^\dagger \sigma_3 B_{r,l}(k) = \sigma_3, \quad (3.8a)$$

$$B_{r,l}(k)^\dagger = -\sigma_3 B_{l,r}(k) \sigma_3, \quad (3.8b)$$

where $0 \neq k \in \mathbb{R}$. Using the second part for $\mathbf{V} = \mathbf{W} = \mathbf{F}_{r,l}$, we get

$$A_{r,l}(k)^\dagger \sigma_3 B_{r,l}(-k) = B_{r,l}(k)^\dagger \sigma_3 A_{r,l}(-k), \quad (3.9)$$

where $0 \neq k \in \mathbb{R}$. Using the second part for $\mathbf{V} = \mathbf{F}_l$ and $\mathbf{W} = \mathbf{F}_r$ we obtain

$$A_r(k)^\dagger = \sigma_3 A_l(-k) \sigma_3, \quad B_r(k)^\dagger = -\sigma_3 B_l(k) \sigma_3, \quad (3.10)$$

where $0 \neq k \in \mathbb{R}$. Finally, using the first part for $\mathbf{V} = \mathbf{W} = \Phi$ we get

$$A_{r,l}(-k^*)^\dagger = \sigma_3 A_{l,r}(k) \sigma_3, \quad (3.11)$$

where $0 \neq k \in \mathbb{C}^+ \cup \mathbb{R}$.

Introducing the *reflection coefficients*

$$R_{r,l}(k) = B_{r,l}(k) A_{r,l}(k)^{-1} = -A_{l,r}(k)^{-1} B_{l,r}(-k) \quad (3.12)$$

and the *transmission coefficients* $A_{r,l}(k)^{-1}$, we obtain the *Riemann-Hilbert problem*

$$(F_l(x, -k) \quad F_r(x, -k)) = (F_r(x, k) \quad F_l(x, k)) \begin{pmatrix} A_r(k)^{-1} & -R_l(k) \\ -R_r(k) & A_l(k)^{-1} \end{pmatrix}, \quad (3.13)$$

where the matrix $\mathbf{S}(k)$ containing the A and R quantities is called the *scattering matrix* and the discussion of the nonsingularity of $A_{r,l}(k)$ will be presented shortly. Then it is easily verified that

$$R_{r,l}(k)^\dagger = \sigma_3 R_{r,l}(-k) \sigma_3, \quad (3.14)$$

and

$$\mathbf{S}(k)^\dagger (\sigma_3 \oplus \sigma_3) \mathbf{S}(k) = \sigma_3 \oplus \sigma_3, \quad (3.15)$$

provided $0 \neq k \in \mathbb{R}$ and $\det A_{r,l}(k) \neq 0$.

Above we have defined $\Delta_{r,l}$ as follows:

$$\Delta_{r,l} = \lim_{k \rightarrow 0} 2ik A_{r,l}(k) = - \lim_{k \rightarrow 0^\pm} 2ik B_{r,l}(k),$$

where the first limit may be taken from the upper half-plane. Then the matrices $\Delta_{r,l}$ have the same determinant. If $\Delta_{r,l}$ is nonsingular, we say that we are in the *generic case*; if instead $\Delta_{r,l}$ is singular, we are in the *exceptional case*. We are said to be in the *superexceptional case* if $\Delta_{r,l} = 0_{n \times n}$ and $A_{r,l}(k)$ tends to a nonsingular matrix, $A_{r,l}(0)$ say, as $k \rightarrow 0$ from within $\mathbb{C}^+ \cup \mathbb{R}$.

Throughout this article, we assume the absence of *spectral singularities*, i.e., the absence of nonzero real k for which $\det A_{r,l}(k) = 0$. Under this condition the reflection coefficients $R_{r,l}(k)$ are continuous in $0 \neq k \in \mathbb{R}$.

3.2 Triangular representations and Wiener algebras

The Jost solutions allow the triangular representations

$$F_l(x, k) = e^{ikx} I_n + \int_x^\infty dy e^{iky} K(x, y), \quad (3.16a)$$

$$F_r(x, k) = e^{-ikx} I_n + \int_{-\infty}^x dy e^{-iky} J(x, y), \quad (3.16b)$$

where for every $x \in \mathbb{R}$

$$\int_x^\infty dy \|K(x, y)\| + \int_{-\infty}^x dy \|J(x, y)\| < +\infty. \quad (3.17)$$

To derive (3.16), we introduce the auxiliary matrix functions $K(x, y)$ and $J(x, y)$ as the unique solutions of the Volterra integral equations

$$\begin{aligned} K(x, y) &= \frac{1}{2} \int_{\frac{1}{2}[x+y]}^{\infty} dz \mathbf{Q}(z) + \frac{1}{2} \int_x^{\infty} dz \mathbf{Q}(z) \int_{\max(z, x+y-z)}^{z+y-x} d\hat{y} K(z, \hat{y}) \\ &= \frac{1}{2} \int_{\frac{1}{2}[x+y]}^{\infty} dz \mathbf{Q}(z) + \frac{1}{2} \int_x^y dz \int_{x+\frac{1}{2}(y-z)}^{\infty} dw \mathbf{Q}(w) K(w, w+z-x), \end{aligned} \quad (3.18a)$$

$$\begin{aligned} J(x, y) &= \frac{1}{2} \int_{-\infty}^{\frac{1}{2}[x+y]} dz \mathbf{Q}(z) + \frac{1}{2} \int_{-\infty}^x dz \mathbf{Q}(z) \int_{z+y-x}^{\min(z, x+y-z)} d\hat{y} J(z, \hat{y}) \\ &= \frac{1}{2} \int_{-\infty}^{\frac{1}{2}[x+y]} dz \mathbf{Q}(z) + \frac{1}{2} \int_y^x dz \int_{-\infty}^{x-\frac{1}{2}[z-y]} dw \mathbf{Q}(w) J(w, w+z-x). \end{aligned} \quad (3.18b)$$

Solving (3.18) by iteration we easily derive (3.16) and (3.17). Moreover,

$$K(x, x) = \frac{1}{2} \int_x^{\infty} dy \mathbf{Q}(y), \quad J(x, x) = \frac{1}{2} \int_{-\infty}^x dy \mathbf{Q}(y). \quad (3.19)$$

Let us now introduce some necessary terminology and well-known results on Fourier transforms of L^1 -functions. By the (continuous) Wiener algebra \mathcal{W} we mean the complex vector space of constants plus Fourier transforms of L^1 -functions

$$\mathcal{W} = \{c + \hat{h} : c \in \mathbb{C}, h \in L^1(\mathbb{R})\}$$

endowed with the norm $|c| + \|h\|_1$. Here we define the Fourier transform as follows: $(\mathcal{F}h)(k) = \hat{h}(k) = \int_{-\infty}^{\infty} dy e^{iky} h(y)$. The invertible elements of the commutative Banach algebra \mathcal{W} with unit element are exactly those $c + \hat{h} \in \mathcal{W}$ for which $c \neq 0$ and $c + \hat{h}(k) \neq 0$ for each $k \in \mathbb{R}$ [30].

The algebra \mathcal{W} has the two closed subalgebras \mathcal{W}^+ and \mathcal{W}^- consisting of those $c + \hat{h} \in \mathcal{W}$ for which h is supported on \mathbb{R}^+ and \mathbb{R}^- , respectively. The invertible elements of \mathcal{W}^{\pm} are exactly those $c + \hat{h} \in \mathcal{W}^{\pm}$ for which $c \neq 0$ and $c + \hat{h}(k) \neq 0$ for each $k \in \mathbb{C}^{\pm} \cup \mathbb{R}$ [30]. Letting \mathcal{W}_0^{\pm} and \mathcal{W}_0 stand for the (nonunital) closed subalgebras of \mathcal{W}^{\pm} and \mathcal{W} consisting of those $c + \hat{h}$ for which $c = 0$, we obtain the direct sum decompositions

$$\mathcal{W} = \mathbb{C} \oplus \mathcal{W}_0^+ \oplus \mathcal{W}_0^-, \quad \mathcal{W}_0 = \mathcal{W}_0^+ \oplus \mathcal{W}_0^-.$$

By Π_{\pm} we now denote the (bounded) projections of \mathcal{W} onto \mathcal{W}_0^{\pm} along $\mathbb{C} \oplus \mathcal{W}_0^{\mp}$. Then Π_+ and Π_- are complementary projections. In fact,

$$(\Pi_{\pm}f)(k) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} d\zeta \frac{f(\zeta)}{\zeta - (k \pm i0^+)}, \quad (3.20)$$

where $f \in \mathcal{W}_0 \cap L^p(\mathbb{R})$ for some $p \in (1, +\infty)$. These direct sum decompositions can be coupled by the Fourier transform to the natural direct sum decomposition of $L^1(\mathbb{R})$ as follows:

$$\begin{array}{ccc} L^1(\mathbb{R}) & = & L^1(\mathbb{R}^-) \oplus L^1(\mathbb{R}^+) \\ \downarrow \mathcal{F} & & \downarrow \mathcal{F} \quad \oplus \quad \downarrow \mathcal{F} \\ \mathcal{W}_0 & = & \mathcal{W}_0^- \oplus \mathcal{W}_0^+ \end{array}$$

Throughout this article we denote the vector spaces of $n \times m$ matrices with entries in \mathcal{W} , \mathcal{W}^{\pm} , and \mathcal{W}_0^{\pm} by $\mathcal{W}^{n \times m}$, $\mathcal{W}^{\pm n \times m}$, and $\mathcal{W}_0^{\pm n \times m}$, respectively. We write $L^1(\mathbb{R})^{n \times m}$ and $L^1(\mathbb{R}^{\pm})^{n \times m}$ for the vector spaces of $n \times m$ matrices with entries in $L^1(\mathbb{R})$ and $L^1(\mathbb{R}^{\pm})$, respectively. Using a submultiplicative matrix norm, we can turn all of these vector spaces into Banach spaces. It is then clear that $\mathcal{W}^{n \times n}$ and $\mathcal{W}^{\pm n \times n}$ are noncommutative Banach algebras with unit element and $\mathcal{W}_0^{\pm n \times n}$ are (nonunital) noncommutative Banach algebras. The projections Π^{\pm} can be extended in a natural way to act on matrices of Wiener algebra elements. Hence, according to (3.16) and (3.17), for each $x \in \mathbb{R}$ the Faddeev functions $m_{r,l}(x, \cdot) \in \mathcal{W}_+^{n \times n}$.

For $n = 1$ the following result is most easily proved using the Gelfand theory of commutative Banach algebras [30]. The extension to arbitrary n is elementary, because the determinant and the cofactor matrix of an element of $\mathcal{W}^{n \times n}$ belong to \mathcal{W} and $\mathcal{W}^{n \times n}$, respectively.

Theorem 3.2 *If for some complex $n \times n$ matrix H_{∞} and some $H \in L^1(\mathbb{R})^{n \times n}$ the Fourier transform $H_{\infty} + \int_{-\infty}^{\infty} dz e^{ikz} H(z)$ is a nonsingular matrix for every $k \in \mathbb{R}$ and if $\det H_{\infty} \neq 0$, then there exists $K \in L^1(\mathbb{R})^{n \times n}$ such that*

$$\left[H_{\infty} + \int_{-\infty}^{\infty} dz e^{ikz} H(z) \right]^{-1} = (H_{\infty})^{-1} + \int_{-\infty}^{\infty} dz e^{ikz} K(z)$$

for every $k \in \mathbb{R}$.

Using that $m_{r,l}(x, \cdot) \in \mathcal{W}_+^{n \times n}$, we easily prove with the help of (3.4) that $2ik[I_n - A_{r,l}(k)]$ belongs to $\mathcal{W}_+^{n \times n}$ and $2ikB_{r,l}(k)$ belongs to $\mathcal{W}^{n \times n}$. Assuming the absence of spectral singularities and to be in the generic case, we

easily prove that the reflection coefficients $R_{r,l}(k)$ belong to $\mathcal{W}_0^{n \times n}$ and the transmission coefficients $A_{r,l}(k)^{-1}$ to $\mathcal{W}_+^{n \times n}$. Indeed,

$$\frac{k}{k+1}A_{r,l}(k) = \frac{k}{k+i}I_n - \frac{2ik[I_n - A_{r,l}(k)]}{2i(k+i)}$$

belongs to $\mathcal{W}_+^{n \times n}$ and therefore, by Theorem 3.2 and $k/(k+i) \in \mathcal{W}$,

$$A_{r,l}(k)^{-1} = \frac{k}{k+i} \left[\frac{k}{k+i}A_{r,l}(k) \right]^{-1}$$

belongs to $\mathcal{W}^{n \times n}$. Analogously, since

$$\frac{k}{k+i}B_{r,l}(k) = \frac{2ikB_{r,l}(k)}{2i(k+i)}$$

belongs to $\mathcal{W}^{n \times n}$, we easily see that

$$R_{r,l}(k) = \left[\frac{k}{k+i}B_{r,l}(k) \right] \left[\frac{k}{k+i}A_{r,l}(k) \right]^{-1}$$

belongs to $\mathcal{W}_0^{n \times n}$.

The exceptional case requires more attention. Assuming that the potential $\mathbf{Q} \in L^1(\mathbb{R}; (1+|x|)^2 dx)$, we can differentiate (3.2) with respect to k and solve the resulting integral equations by iteration. Next, with the help of (3.4) we obtain

$$A_{r,l}(k) + B_{r,l}(k) = I_n + \int_{-\infty}^{\infty} dy \frac{e^{\mp 2iky} - 1}{2ik} \mathbf{Q}(y) m_{r,l}(y, k).$$

This expression satisfies

$$\lim_{k \rightarrow 0^\pm} [A_{r,l}(k) + B_{r,l}(k)] = I_n \mp \int_{-\infty}^{\infty} dy y \mathbf{Q}(y) m_{r,l}(y, 0) = I_n \mp E_{r,l},$$

provided $\mathbf{Q} \in L^1(\mathbb{R}; (1+|x|)^2 dx)$. Taking the limit as $k \rightarrow 0^\pm$ we get

$$\begin{aligned} \lim_{\substack{k \rightarrow 0 \\ k \in \mathbb{C}^+ \cup \mathbb{R}}} \frac{2ikA_{r,l}(k) - \Delta_{r,l}}{2ik} &= I_n - \frac{1}{2i} \int_{-\infty}^{\infty} dy \mathbf{Q}(y) \dot{m}_{r,l}(y, 0) \\ &= I_n - \frac{1}{2i} G_{r,l}, \end{aligned} \tag{3.21a}$$

$$\begin{aligned} \lim_{k \rightarrow 0^\pm} \frac{2ikB_{r,l}(k) + \Delta_{r,l}}{2ik} &= \int_{-\infty}^{\infty} dy \mathbf{Q}(y) [\mp y m_{r,l}(y, 0) + \frac{1}{2i} \dot{m}_{r,l}(y, 0)] \\ &= \mp E_{r,l} + \frac{1}{2i} G_{r,l}, \end{aligned} \tag{3.21b}$$

where the overdot indicates differentiation with respect to k .

In the superexceptional case, where $\Delta_{r,l} = 0_{n \times n}$, it is clear that $A_{r,l} \in \mathcal{W}_+^{n \times n}$, provided $\mathbf{Q} \in L^1(\mathbb{R}; (1 + |x|)^2 dx)$. Assuming the absence of spectral singularities, using the nonsingularity of $A_{r,l}(0)$, and applying Theorem 3.2, we get the transmission coefficients $A_{r,l}(k)^{-1}$ to belong to $\mathcal{W}^{n \times n}$. Since $A_{r,l}(k) + B_{r,l}(k)$ belongs to $\mathcal{W}^{n \times n}$, we conclude that

$$R_{r,l}(k) = [A_{r,l}(k) + B_{r,l}(k)]A_{r,l}(k)^{-1} - I_n \in \mathcal{W}_0^{n \times n}.$$

At present it is not known if, under the absence of spectral singularities, the reflection and transmission coefficients belong to $\mathcal{W}^{n \times n}$ in any other exceptional case and for general $\mathbf{Q} \in L^1(\mathbb{R}; (1 + |x|)dx)$. Under the condition $\mathbf{Q} \in L^1(\mathbb{R}; (1 + |x|)dx)$, the continuity of the reflection and transmission coefficients at $k = 0$ is known for $n = 1$ [41] and for selfadjoint potentials [10].

4 Inverse scattering problem

In this section we introduce the Marchenko integral equations for the matrix Schrödinger equation (1.5) with Faddeev class potential \mathbf{Q} satisfying (1.2). We make use of the hypothesis that the reflection coefficients $R_{r,l} \in \mathcal{W}_0^{n \times n}$, something that we have not been able to prove in the most general exceptional case. For the sake of simplicity we assume that the poles of $A_{r,l}(k)^{-1}$ in \mathbb{C}^+ are simple. The extension to multiple pole situations is rather technical but straightforward [24]. Inverse scattering theory is well documented in the scalar case [28, 22, 19, 20, 2], as it is in the matrix half-line case [7, 12, 13]. Without any details, the matrix full-line case is given in [56].

Let us first write the *transmission coefficients* in the form

$$A_r(k)^{-1} = A_{r0}(k) + \sum_{s=1}^N \frac{\tau_{r;s}}{k - k_s}, \quad A_l(k)^{-1} = A_{l0}(k) + \sum_{s=1}^N \frac{\tau_{l;s}}{k - k_s}, \quad (4.1)$$

where k_1, \dots, k_N are the distinct simple poles of $A_{r,l}(k)^{-1}$ in \mathbb{C}^+ , $\tau_{r;s}$ and $\tau_{l;s}$ are the residues of $A_r(k)^{-1}$ and $A_l(k)^{-1}$ at $k = k_s$ ($s = 1, \dots, N$), and $A_{r0}(k)$ and $A_{l0}(k)$ are continuous in $k \in \mathbb{C}^+ \cup \mathbb{R}$, are analytic in $k \in \mathbb{C}^+$, and tend to I_n as $k \rightarrow \infty$ from within $\mathbb{C}^+ \cup \mathbb{R}$.

Letting $\tau_{r;s}$ and $\tau_{r;\bar{s}}$ (or: $\tau_{l;s}$ and $\tau_{l;\bar{s}}$) stand for the residues of $A_r(k)^{-1}$ (or: $A_l(k)^{-1}$) at k_s and $-k_s^*$, we get with the help of (3.11)

$$\begin{aligned}\tau_{r;\bar{s}} &= \lim_{k \rightarrow -k_s^*} (k + k_s^*) A_r(k)^{-1} = \sigma_3 \lim_{k \rightarrow -k_s^*} (k + k_s^*) A_l(-k^*)^{\dagger -1} \sigma_3 \\ &= -\sigma_3 \left(\lim_{\zeta \rightarrow k_s} (\zeta - k_s) A_l(\zeta)^{-1} \right)^{\dagger} \sigma_3 = -\sigma_3 \tau_{l;s}^{\dagger} \sigma_3.\end{aligned}\quad (4.2)$$

Similarly, we get $\tau_{l;\bar{s}} = -\sigma_3 \tau_{r;s}^{\dagger} \sigma_3$.

Next, let us consider the Riemann-Hilbert problem

$$\begin{pmatrix} F_l(x, -k) & F_r(x, -k) \end{pmatrix} = \begin{pmatrix} F_r(x, k) & F_l(x, k) \end{pmatrix} \begin{pmatrix} A_r(k)^{-1} & -R_l(k) \\ -R_r(k) & A_l(k)^{-1} \end{pmatrix}, \quad (4.3)$$

where $\det A_{r,l}(k) \neq 0$ for $0 \neq k \in \mathbb{R}$. In the generic and superexceptional cases while assuming the absence of spectral singularities, we write

$$R_r(k) = \int_{-\infty}^{\infty} d\alpha e^{-ik\alpha} \hat{R}_r(\alpha), \quad R_l(k) = \int_{-\infty}^{\infty} d\alpha e^{ik\alpha} \hat{R}_l(\alpha), \quad (4.4)$$

where the entries of the $n \times n$ matrix functions $\hat{R}_{r,l}(\alpha)$ belong to $L^1(\mathbb{R})$. In fact, we only need to assume the absence of spectral singularities as well as $R_{r,l} \in \mathcal{W}_0^{n \times n}$ for the derivations below to be valid. Rewriting the left half of (4.3) we obtain

$$\begin{aligned}e^{ikx} F_l(x, -k) &= e^{ikx} F_r(x, k) A_r(k)^{-1} - e^{-ikx} F_l(x, k) e^{2ikx} R_r(k) \\ &= e^{ikx} F_r(x, k) A_{r0}(k) + \sum_{s=1}^N \frac{e^{ikx} F_r(x, k) - e^{ik_s x} F_r(x, k_s)}{k - k_s} \tau_{r;s} \\ &\quad + i \sum_{s=1}^N e^{2ik_s x} \frac{e^{-ik_s x} F_l(x, k_s)}{k - k_s} N_{r;s} - e^{-ikx} F_l(x, k) e^{2ikx} R_r(k),\end{aligned}\quad (4.5)$$

where the *norming constants* $N_{r;s}$ ($s = 1, \dots, N$) are defined by

$$F_r(x, k_s) \tau_{r;s} = i F_l(x, k_s) N_{r;s}, \quad s = 1, \dots, N. \quad (4.6)$$

By the same token, rewriting the right half of (4.3) we get

$$\begin{aligned}
e^{-ikx} F_r(x, -k) &= e^{-ikx} F_l(x, k) A_l(k)^{-1} - e^{ikx} F_r(x, k) e^{-2ikx} R_l(k) \\
&= e^{-ikx} F_l(x, k) A_{l0}(k) + \sum_{s=1}^N \frac{e^{-ikx} F_l(x, k) - e^{-ik_s x} F_l(x, k_s)}{k - k_s} \tau_{l;s} \\
&\quad + i \sum_{s=1}^N e^{-2ik_s x} \frac{e^{ik_s x} F_r(x, k_s)}{k - k_s} N_{l;s} - e^{ikx} F_r(x, k) e^{-2ikx} R_l(k), \quad (4.7)
\end{aligned}$$

where the norming constants $N_{l;s}$ ($s = 1, \dots, N$) are defined by

$$F_l(x, k_s) \tau_{l;s} = i F_r(x, k_s) N_{l;s}, \quad s = 1, \dots, N. \quad (4.8)$$

For every $x \in \mathbb{R}$ the identities (4.5) and (4.7) are equations in $\mathcal{W}^{n \times n}$. Using Π_- to project these two equations onto $\mathcal{W}_-^{n \times n}$ along $\mathcal{W}_{+,0}^{n \times n}$, we obtain

$$\begin{aligned}
e^{ikx} F_l(x, -k) &= I_n + i \sum_{s=1}^N e^{2ik_s x} \frac{e^{-ik_s x} F_l(x, k_s)}{k - k_s} N_{r;s} \\
&\quad - \Pi_- [e^{-ikx} F_l(x, k) e^{2ikx} R_r(k)], \\
e^{-ikx} F_r(x, -k) &= I_n + i \sum_{s=1}^N e^{-2ik_s x} \frac{e^{ik_s x} F_r(x, k_s)}{k - k_s} N_{l;s} \\
&\quad - \Pi_- [e^{ikx} F_r(x, k) e^{-2ikx} R_l(k)],
\end{aligned}$$

respectively. Using (3.16), (4.4), and the identity

$$\int_0^\infty dw e^{-ikw} e^{ik_s w} = \frac{-i}{k - k_s}, \quad s = 1, 2, \dots, N, \quad (4.9)$$

we strip off the Fourier transforms and arrive at the Marchenko integral equations

$$K(x, y) + \Omega_r(x + y) + \int_x^\infty dz K(x, z) \Omega_r(z + y) = 0_{n \times n}, \quad (4.10a)$$

$$J(x, y) + \Omega_l(x + y) + \int_{-\infty}^x dz J(x, z) \Omega_l(z + y) = 0_{n \times n}, \quad (4.10b)$$

where the *Marchenko integral kernels* are given by

$$\Omega_r(w) = \hat{R}_r(w) + \sum_{s=1}^N e^{ik_s w} N_{r;s}, \quad (4.11a)$$

$$\Omega_l(w) = \hat{R}_l(w) + \sum_{s=1}^N e^{-ik_s w} N_{l;s}. \quad (4.11b)$$

Indeed, to prove (4.10a) and (4.11a) we compute

$$\begin{aligned} & \int_0^\infty d\alpha e^{-ik\alpha} K(x, x + \alpha) \\ &= - \int_0^\infty d\alpha e^{-ik\alpha} \sum_{s=1}^N \left[e^{ik_s(\alpha+2x)} N_{r;s} + \int_0^\infty d\beta e^{ik_s(\alpha+\beta+2x)} K(x, x + \beta) N_{r;s} \right] \\ & - \int_0^\infty d\alpha e^{-ik\alpha} \left[\hat{R}_r(\alpha + 2x) + \int_0^\infty d\beta K(x, x + \beta) \hat{R}_r(\alpha + \beta + 2x) \right]. \end{aligned}$$

Stripping off the Fourier transform and putting $y = x + \alpha$ and $z = x + \beta$, we get (4.10a) and (4.11a). Analogously, to prove (4.10b) and (4.11b) we compute

$$\begin{aligned} & \int_0^\infty d\alpha e^{-ik\alpha} J(x, x - \alpha) \\ &= - \int_0^\infty d\alpha e^{-ik\alpha} \sum_{s=1}^N \left[e^{-ik_s(2x-\alpha)} N_{l;s} + \int_0^\infty d\beta e^{-ik_s(2x-\alpha-\beta)} J(x, x - \beta) N_{l;s} \right] \\ & - \int_0^\infty d\alpha e^{-ik\alpha} \left[\hat{R}_l(2x - \alpha) + \int_0^\infty d\beta J(x, x - \beta) \hat{R}_l(2x - \alpha - \beta) \right]. \end{aligned}$$

Stripping off the Fourier transform and putting $y = x - \alpha$ and $z = x - \beta$, we get (4.10b).

As in [24], we can prove the adjoint symmetry relations

$$\Omega_{r,l}(w) = \sigma_3 \Omega_{r,l}(w)^\dagger \sigma_3, \quad (4.12)$$

thus implying the following symmetry relations for the norming constants:

$$N_{r;s} = \sigma_3 N_{r;s}^\dagger \sigma_3, \quad N_{l;s} = \sigma_3 N_{l;s}^\dagger \sigma_3. \quad (4.13)$$

For the rather tedious details we refer to Appendix B.

5 AKNS vs. matrix Schrödinger data

Let us define the Jost solutions of the AKNS system (1.3) as those solutions of (1.3) that for $k \in \mathbb{R}$ satisfy the asymptotic relations

$$\mathbb{F}_l(x, k) = e^{-ikx\sigma_3}[I_n + o(1)], \quad x \rightarrow +\infty, \quad (5.1a)$$

$$\mathbb{F}_r(x, k) = e^{-ikx\sigma_3}[I_n + o(1)], \quad x \rightarrow -\infty. \quad (5.1b)$$

Since we are dealing with a first order system, there exist nonsingular matrices $\mathbb{A}_l(k)$ and $\mathbb{A}_r(k)$ such that

$$\mathbb{F}_r(x, k) = \mathbb{F}_l(x, k)\mathbb{A}_r(k), \quad \mathbb{F}_l(x, k) = \mathbb{F}_r(x, k)\mathbb{A}_l(k), \quad (5.2)$$

and hence $\mathbb{A}_l(k)$ and $\mathbb{A}_r(k)$ are each other's inverses. Equations (5.1) and (5.2) imply the asymptotic relations

$$\mathbb{F}_l(x, k) = e^{-ikx\sigma_3}\mathbb{A}_l(k)[I_n + o(1)], \quad x \rightarrow -\infty, \quad (5.3a)$$

$$\mathbb{F}_r(x, k) = e^{-ikx\sigma_3}\mathbb{A}_r(k)[I_n + o(1)], \quad x \rightarrow +\infty. \quad (5.3b)$$

Since the matrix $-ik\sigma_3 + \mathcal{Q}$ in (1.3) has trace $-ik(m_1 - m_2)$, the determinants of $\mathbb{F}_l(x, k)$ and $\mathbb{F}_r(x, k)$ equal $e^{-ik(m_1 - m_2)x}$. As a result, $\det \mathbb{A}_{r,l}(k) = 1$.

Let us relate the AKNS and matrix Schrödinger Jost solutions. Putting

$$\mathbf{e}_1 = I_{m_1} \oplus 0_{m_2 \times m_2}, \quad \mathbf{e}_2 = 0_{m_1 \times m_1} \oplus I_{m_2},$$

we obtain from the principal asymptotic relations

$$\mathbb{F}_l(x, k) = F_l(x, -k)\mathbf{e}_1 + F_l(x, k)\mathbf{e}_2, \quad (5.4a)$$

$$\mathbb{F}_r(x, k) = F_r(x, k)\mathbf{e}_1 + F_r(x, -k)\mathbf{e}_2, \quad (5.4b)$$

as well as

$$F_l(x, k) = \mathbb{F}_l(x, -k)\mathbf{e}_1 + \mathbb{F}_l(x, k)\mathbf{e}_2, \quad (5.5a)$$

$$F_r(x, k) = \mathbb{F}_r(x, k)\mathbf{e}_1 + \mathbb{F}_r(x, -k)\mathbf{e}_2. \quad (5.5b)$$

Using $\mathbf{e}_1\mathbf{e}_2 = \mathbf{e}_2\mathbf{e}_1 = 0_{n \times n}$ and $\mathbf{e}_1^2 + \mathbf{e}_2^2 = I_n$ it is easy to derive (5.5) from (5.4) and vice versa. Using (5.4) we obtain the triangular representations for the AKNS system

$$\mathbb{F}_l(x, k) = e^{-ikx\sigma_3} + \int_x^\infty dy K(x, y)e^{-iky\sigma_3}, \quad (5.6a)$$

$$\mathbb{F}_r(x, k) = e^{-ikx\sigma_3} + \int_{-\infty}^x dy J(x, y)e^{-iky\sigma_3}, \quad (5.6b)$$

where the kernel functions K and J coincide with those for the matrix Schrödinger system.

Let us now relate the AKNS and matrix Schrödinger scattering coefficients. Let us introduce the standard partitioning

$$H = \begin{pmatrix} H_1 & H_2 \\ H_3 & H_4 \end{pmatrix} \quad (5.7)$$

of an $n \times n$ matrix H into the $m_1 \times m_1$ matrix H_1 , the $m_1 \times m_2$ matrix H_2 , the $m_2 \times m_1$ matrix H_3 , and the $m_2 \times m_2$ matrix H_4 . Using the asymptotic relations (3.3) and (5.3) we obtain for $k \in \mathbb{R}$

$$\begin{aligned} e^{ikx} A_l(k) + e^{-ikx} B_l(k) &= \begin{pmatrix} e^{ikx} \mathbb{A}_{l1}(-k) & e^{-ikx} \mathbb{A}_{l2}(k) \\ e^{-ikx} \mathbb{A}_{l3}(-k) & e^{ikx} \mathbb{A}_{l4}(k) \end{pmatrix}, \\ e^{-ikx} A_r(k) + e^{ikx} B_r(k) &= \begin{pmatrix} e^{-ikx} \mathbb{A}_{r1}(k) & e^{ikx} \mathbb{A}_{r2}(-k) \\ e^{ikx} \mathbb{A}_{r3}(k) & e^{-ikx} \mathbb{A}_{r4}(-k) \end{pmatrix}. \end{aligned}$$

Consequently, for $0 \neq k \in \mathbb{R}$ we have

$$A_l(k) = \begin{pmatrix} \mathbb{A}_{l1}(-k) & 0_{m_1 \times m_2} \\ 0_{m_2 \times m_1} & \mathbb{A}_{l4}(k) \end{pmatrix}, \quad A_r(k) = \begin{pmatrix} \mathbb{A}_{r1}(k) & 0_{m_1 \times m_2} \\ 0_{m_2 \times m_1} & \mathbb{A}_{r4}(-k) \end{pmatrix}, \quad (5.8a)$$

$$B_l(k) = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{A}_{l2}(k) \\ \mathbb{A}_{l3}(-k) & 0_{m_2 \times m_2} \end{pmatrix}, \quad B_r(k) = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{A}_{r2}(-k) \\ \mathbb{A}_{r3}(k) & 0_{m_2 \times m_2} \end{pmatrix}. \quad (5.8b)$$

Conversely, for $0 \neq k \in \mathbb{R}$ we get in terms of blocks of $A_{r,l}(k)$ and $B_{r,l}(k)$

$$\mathbb{A}_l(k) = \begin{pmatrix} A_{l1}(k) & B_{l2}(k) \\ B_{l3}(-k) & A_{l4}(k) \end{pmatrix}, \quad (5.9a)$$

$$\mathbb{A}_r(k) = \begin{pmatrix} A_{r1}(k) & B_{r2}(-k) \\ B_{r3}(k) & A_{r4}(-k) \end{pmatrix}. \quad (5.9b)$$

Since $\mathbb{A}_l(k)$ and $\mathbb{A}_r(k)$ are continuous in $k \in \mathbb{R}$ and tend to I_n as $k \rightarrow \pm\infty$, we see that $A_l(k)$, $B_l(k)$, $A_r(k)$, and $B_r(k)$ are continuous in $k \in \mathbb{R}$ and have finite limits as $k \rightarrow \pm\infty$. Of course, the block entries of $\mathbb{A}_{l,r}(k)$ have all the usual continuity and analyticity properties, also at $k = 0$. Thus, we are always in the exceptional case. We are in the superexceptional case iff $k = 0$ is not a spectral singularity of the AKNS system (1.3). Further, the matrix

Schrödinger reflection coefficients are given by

$$R_l(k) = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{R}_{l1}(k) \\ \mathbb{R}_{l4}(k) & 0_{m_2 \times m_2} \end{pmatrix} = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{A}_{l2}(k)\mathbb{A}_{l4}(k)^{-1} \\ \mathbb{A}_{l3}(-k)\mathbb{A}_{l1}(-k)^{-1} & 0_{m_2 \times m_2} \end{pmatrix}, \quad (5.10a)$$

$$R_r(k) = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{R}_{r4}(k) \\ \mathbb{R}_{r1}(k) & 0_{m_2 \times m_2} \end{pmatrix} = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{A}_{r2}(-k)\mathbb{A}_{r4}(-k)^{-1} \\ \mathbb{A}_{r3}(k)\mathbb{A}_{r1}(k)^{-1} & 0_{m_2 \times m_2} \end{pmatrix}. \quad (5.10b)$$

Finally, the AKNS system is reflectionless iff the matrix Schrödinger system is reflectionless.

Let $\tau_{r;s}$ and $\tau_{l;s}$ be the residues of $A_r(k)^{-1}$ and $A_l(k)^{-1}$ at the simple pole $k_s \in \mathbb{C}^+$. Now let $\tau_{l4;s}$ and $\tau_{r1;s}$ be the residues of $\mathbb{A}_{l4}(k)^{-1}$ and $\mathbb{A}_{r1}(k)^{-1}$ at $k = k_s$ and $\tau_{l1;s}$ and $\tau_{r4;s}$ the residues of $\mathbb{A}_{l1}(k)^{-1}$ and $\mathbb{A}_{r4}(k)^{-1}$ at the simple pole $k = -k_s$. Then (5.8a) implies that

$$\tau_{r;s} = \tau_{r1;s} \oplus (-\tau_{r4;s}), \quad \tau_{l;s} = (-\tau_{l1;s}) \oplus \tau_{l4;s}.$$

Let us define the AKNS norming constants as follows:

$$\mathbb{F}_r(x, k_s) \mathbf{e}_1 \tau_{r1;s} = i \mathbb{F}_l(x, k_s) \mathbf{e}_2 \mathbb{N}_{r1;s}, \quad (5.11a)$$

$$\mathbb{F}_r(x, -k_s) \mathbf{e}_2 \tau_{r4;s} = -i \mathbb{F}_l(x, -k_s) \mathbf{e}_1 \mathbb{N}_{r4;s}, \quad (5.11b)$$

$$\mathbb{F}_l(x, k_s) \mathbf{e}_2 \tau_{l4;s} = i \mathbb{F}_r(x, k_s) \mathbf{e}_1 \mathbb{N}_{l4;s}, \quad (5.11c)$$

$$\mathbb{F}_l(x, -k_s) \mathbf{e}_1 \tau_{l1;s} = -i \mathbb{F}_r(x, -k_s) \mathbf{e}_2 \mathbb{N}_{l1;s}. \quad (5.11d)$$

We then obtain

$$N_{r;s} = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{N}_{r4;s} \\ \mathbb{N}_{r1;s} & 0_{m_2 \times m_2} \end{pmatrix}, \quad N_{l;s} = \begin{pmatrix} 0_{m_1 \times m_1} & \mathbb{N}_{l1;s} \\ \mathbb{N}_{l4;s} & 0_{m_2 \times m_2} \end{pmatrix}. \quad (5.12)$$

6 Time evolution

In this section we establish the time evolution of the scattering data of the matrix Schrödinger equation and the accompanying AKNS system. We then go on to derive, in either case, the Marchenko integral kernels as a function of time. These results allow us, in the next section, to derive the reflectionless solutions of the nonlocal integrable equation (2.3) and hence of the focusing matrix NLS equation.

6.1 Matrix Schrödinger time evolution

The Lax pair $\{\mathcal{L}, A\}$ for the matrix Schrödinger equation (2.3) is given by (2.1), where $\mathcal{B}_0 = \mathbf{Q}$. Equation (2.3) is compatible with the linear system

$$\mathcal{L}v = k^2v, \quad v_t = Av,$$

where $\mathcal{L} = -\partial_x^2 + \mathbf{Q}$. We may therefore write

$$\begin{aligned} v_t &= Av = 2i\sigma_3 v_{xx} - 2i\sigma_3 \mathcal{Q}v_x - i\sigma_3 \mathbf{Q}v \\ &= 2i\sigma_3(\mathbf{Q} - k^2\mathbf{1})v - 2i\sigma_3 \mathcal{Q}v_x - i\sigma_3 \mathbf{Q}v \\ &= i\sigma_3 \{(\mathbf{Q} - 2k^2\mathbf{1})v - 2\mathcal{Q}v_x\}, \end{aligned}$$

where $\mathbf{1}$ stands for the identity operator on a suitable function space. Next, we compute

$$\begin{aligned} (v_x)_t &= (Av)_x = i\sigma_3 ((\mathbf{Q} - 2k^2\mathbf{1})v_x + \mathbf{Q}_x v - 2\mathcal{Q}_x v_x - 2\mathcal{Q}(\mathbf{Q} - k^2\mathbf{1})v) \\ &= i\sigma_3 (\mathbf{Q}_x - 2\mathcal{Q}\mathbf{Q} + 2k^2\mathcal{Q})v + i\sigma_3 (\mathbf{Q} - 2k^2\mathbf{1} - 2\mathcal{Q}_x)v_x. \end{aligned}$$

Hence, putting $\mathbf{V} = \begin{pmatrix} v \\ v_x \end{pmatrix}$ we get the linear system

$$\mathbf{V}_x = \mathbf{X}(x, k; t)\mathbf{V}, \quad \mathbf{V}_t = \mathbf{T}(x, k; t)\mathbf{V}, \quad (6.1)$$

where

$$\mathbf{X}(x, k; t) = \begin{pmatrix} 0_{n \times n} & I_n \\ \mathbf{Q}(x; t) - k^2 I_n & 0_{n \times n} \end{pmatrix}, \quad (6.2a)$$

$$\mathbf{T}(x, k; t) = \begin{pmatrix} i\sigma_3(\mathbf{Q} - 2k^2 I_n) & -2i\sigma_3 \mathcal{Q} \\ i\sigma_3(\mathbf{Q}_x - 2\mathcal{Q}\mathbf{Q} + 2k^2 \mathcal{Q}) & i\sigma_3(\mathbf{Q} - 2k^2 I_n - 2\mathcal{Q}_x) \end{pmatrix}. \quad (6.2b)$$

Then we easily compute

$$i(\sigma_3 \oplus \sigma_3)(\mathbf{X}_t - \mathbf{T}_x + \mathbf{X}\mathbf{T} - \mathbf{T}\mathbf{X}) = \begin{pmatrix} 0_{n \times n} & 0_{n \times n} \\ \mathbf{E}_{21} & 0_{n \times n} \end{pmatrix},$$

where

$$\begin{aligned} \mathbf{E}_{21} &= i\sigma_3 \mathbf{Q}_t + \mathbf{Q}_{xx} - 2(\mathcal{Q}\mathbf{Q})_x + 2k^2 \mathcal{Q}_x \\ &\quad - \sigma_3(\mathbf{Q} - k^2 I_n)\sigma_3(\mathbf{Q} - 2k^2 I_n) + (\mathbf{Q} - 2k^2 I_n - 2\mathcal{Q}_x)(\mathbf{Q} - k^2 I_n) \\ &= i\sigma_3 \mathbf{Q}_t + \mathbf{Q}_{xx} + \mathbf{Q}^2 - \sigma_3 \mathbf{Q} \sigma_3 \mathbf{Q} - 2\mathcal{Q}\mathbf{Q}_x - 4\mathcal{Q}_x \mathbf{Q} \\ &\quad + k^2(2\mathcal{Q}_x + 2\sigma_3 \mathbf{Q} \sigma_3 + \mathbf{Q} - \mathbf{Q} - 2\mathbf{Q} + 2\mathcal{Q}_x) \\ &= i\sigma_3 \mathbf{Q}_t + \mathbf{Q}_{xx} + \mathbf{Q}^2 - \sigma_3 \mathbf{Q} \sigma_3 \mathbf{Q} - 2\mathcal{Q}\mathbf{Q}_x - 2(\mathbf{Q}^2 - \sigma_3 \mathbf{Q} \sigma_3 \mathbf{Q}) \\ &= i\sigma_3 \mathbf{Q}_t + \mathbf{Q}_{xx} - \mathbf{Q}^2 + \sigma_3 \mathbf{Q} \sigma_3 \mathbf{Q} - 2\mathcal{Q}\mathbf{Q}_x, \end{aligned}$$

as claimed. Thus the zero curvature condition for the AKNS pair $\{\mathbf{X}, \mathbf{T}\}$ is equivalent to the nonlinear evolution equation (2.3).

There exist nonsingular matrices $C_{\mathbf{F}_l}(k; t)$ and $C_{\mathbf{F}_r}(k; t)$ not depending on $k \in \mathbb{R}$ such that

$$\mathbf{F}_l(x, k; t) = \mathbf{V}(x, k; t)C_{\mathbf{F}_l}(k; t)^{-1}, \quad \mathbf{F}_r(x, k; t) = \mathbf{V}(x, k; t)C_{\mathbf{F}_r}(k; t)^{-1}.$$

Then a simple differentiation yields

$$\begin{aligned} [C_{\mathbf{F}_l}(k; t)]_t C_{\mathbf{F}_l}(k; t)^{-1} &= \mathbf{F}_l^{-1} \mathbf{T} \mathbf{F}_l - \mathbf{F}_l^{-1} [\mathbf{F}_l]_t, \\ [C_{\mathbf{F}_r}(k; t)]_t C_{\mathbf{F}_r}(k; t)^{-1} &= \mathbf{F}_r^{-1} \mathbf{T} \mathbf{F}_r - \mathbf{F}_r^{-1} [\mathbf{F}_r]_t, \end{aligned}$$

where the two left-hand sides do not depend on $x \in \mathbb{R}$. We may therefore compute the $x \rightarrow \pm\infty$ limits of the two right-hand sides and obtain

$$[C_{\mathbf{F}_l}(k; t)]_t C_{\mathbf{F}_l}(k; t)^{-1} = [C_{\mathbf{F}_r}(k; t)]_t C_{\mathbf{F}_r}(k; t)^{-1} = -2ik^2(\boldsymbol{\sigma}_3 \oplus \boldsymbol{\sigma}_3).$$

Consequently,

$$C_{\mathbf{F}_l}(k; t) = \left[e^{-2ik^2 t \boldsymbol{\sigma}_3} \oplus e^{-2ik^2 t \boldsymbol{\sigma}_3} \right] C_{\mathbf{F}_l}(k; 0), \quad (6.3a)$$

$$C_{\mathbf{F}_r}(k; t) = \left[e^{-2ik^2 t \boldsymbol{\sigma}_3} \oplus e^{-2ik^2 t \boldsymbol{\sigma}_3} \right] C_{\mathbf{F}_r}(k; 0). \quad (6.3b)$$

Relating $\mathbf{F}_{r,l}(x, k; t)$ by means of the equalities

$$\mathbf{F}_r(x, k; t) = \mathbf{F}_l(x, k; t) \mathbf{A}_r(k; t), \quad \mathbf{F}_l(x, k; t) = \mathbf{F}_r(x, k; t) \mathbf{A}_l(k; t),$$

where the factors $\mathbf{A}_{r,l}(k; t)$ are given by the matrices [cf. (3.5)]

$$\mathbf{A}_r(k; t) = \begin{pmatrix} A_r(k; t) & B_r(-k; t) \\ B_r(k; t) & A_r(-k; t) \end{pmatrix}, \quad \mathbf{A}_l(k; t) = \begin{pmatrix} A_l(-k; t) & B_l(k; t) \\ B_l(-k; t) & A_l(k; t) \end{pmatrix},$$

we compute

$$\begin{aligned} [\mathbf{A}_r]_t &= -\mathbf{F}_l^{-1} [\mathbf{F}_l]_t \mathbf{F}_l^{-1} \mathbf{F}_r + \mathbf{F}_l^{-1} [\mathbf{F}_r]_t \\ &= -\mathbf{F}_l^{-1} (\mathbf{T} \mathbf{F}_l - \mathbf{F}_l [C_{\mathbf{F}_l}(k; t)]_t C_{\mathbf{F}_l}(k; t)^{-1}) \mathbf{A}_r \\ &\quad + \mathbf{F}_l^{-1} (\mathbf{T} \mathbf{F}_r - \mathbf{F}_r [C_{\mathbf{F}_r}(k; t)]_t C_{\mathbf{F}_r}(k; t)^{-1}) \\ &= [C_{\mathbf{F}_l}(k; t)]_t C_{\mathbf{F}_l}(k; t)^{-1} \mathbf{A}_r - \mathbf{A}_r [C_{\mathbf{F}_l}(k; t)]_t C_{\mathbf{F}_l}(k; t)^{-1} \\ &= 2ik^2 (\mathbf{A}_r (\boldsymbol{\sigma}_3 \oplus \boldsymbol{\sigma}_3) - (\boldsymbol{\sigma}_3 \oplus \boldsymbol{\sigma}_3) \mathbf{A}_r). \end{aligned}$$

Using that $\mathbf{A}_{r,l}(k;t)$ are each other's inverses, we get

$$\mathbf{A}_{r,l}(k;t) = \left[e^{-2ik^2t\sigma_3} \oplus e^{-2ik^2t\sigma_3} \right] \mathbf{A}_{r,l}(k;0) \left[e^{2ik^2t\sigma_3} \oplus e^{2ik^2t\sigma_3} \right].$$

Therefore,

$$A_{r,l}(k;t) = e^{-2ik^2t\sigma_3} A_{r,l}(k;0) e^{2ik^2t\sigma_3}, \quad (6.4a)$$

$$B_{r,l}(k;t) = e^{-2ik^2t\sigma_3} B_{r,l}(k;0) e^{2ik^2t\sigma_3}. \quad (6.4b)$$

For the reflection coefficients we get

$$R_{r,l}(k;t) = e^{-2ik^2t\sigma_3} R_{r,l}(k;0) e^{2ik^2t\sigma_3}. \quad (6.5)$$

Moreover, if k_s is a simple pole of $A_{r,l}(k;t)$ in \mathbb{C}^+ , (6.4a) implies that

$$\tau_{r;s}(t) = e^{-2ik_s^2t\sigma_3} \tau_{r;s}(0) e^{2ik_s^2t\sigma_3}, \quad \tau_{l;s}(t) = e^{-2ik_s^2t\sigma_3} \tau_{l;s}(0) e^{2ik_s^2t\sigma_3}.$$

As a result,

$$[\tau_{r;s}]_t = 2ik_s^2 (\tau_{r;s}\sigma_3 - \sigma_3\tau_{r;s}), \quad [\tau_{l;s}]_t = 2ik_s^2 (\tau_{l;s}\sigma_3 - \sigma_3\tau_{l;s}). \quad (6.6)$$

Let us now study the time evolution of the norming constants corresponding to a simple pole k_s of $A_r(k;t)^{-1}$ in \mathbb{C}^+ . Equations (6.3) imply

$$[\mathbf{F}_{r,l}]_t = \mathbf{T}\mathbf{F}_{r,l} + 2ik^2\mathbf{F}_{r,l}(\sigma_3 \oplus \sigma_3). \quad (6.7)$$

Now recall (4.6) and (4.8) in the form

$$F_r(x, k_s; t) \tau_{r;s} = iF_l(x, k_s; t) N_{r;s}(t), \quad (6.8a)$$

$$F_l(x, k_s; t) \tau_{l;s} = iF_r(x, k_s; t) N_{l;s}(t). \quad (6.8b)$$

Using the right/left upper block of (6.7) for $k = k_s$ to differentiate (6.8) with respect to t for $k = k_s$ and then substituting (6.6) and (6.8), we get

$$\begin{aligned} F_l(x, k_s; t) \{ [N_{r;s}]_t - 2ik_s^2 (N_{r;s}\sigma_3 - \sigma_3 N_{r;s}) \} &= 0_{n \times n}, \\ F_r(x, k_s; t) \{ [N_{l;s}]_t - 2ik_s^2 (N_{l;s}\sigma_3 - \sigma_3 N_{l;s}) \} &= 0_{n \times n}. \end{aligned}$$

Using (3.1) we obtain

$$\begin{aligned} [N_{r;s}]_t &= 2ik_s^2 (N_{r;s}(t)\sigma_3 - \sigma_3 N_{r;s}(t)), \\ [N_{l;s}]_t &= 2ik_s^2 (N_{l;s}(t)\sigma_3 - \sigma_3 N_{l;s}(t)). \end{aligned}$$

Consequently,

$$N_{r;s}(t) = e^{-2ik_s^2t\sigma_3} N_{r;s}(0) e^{2ik_s^2t\sigma_3}, \quad (6.9a)$$

$$N_{l;s}(t) = e^{-2ik_s^2t\sigma_3} N_{l;s}(0) e^{2ik_s^2t\sigma_3}. \quad (6.9b)$$

6.2 AKNS time evolution

Equations (6.4a) and (5.8a), applied for $k \in \mathbb{R}$ and then extended by analytic continuation, imply that $\mathbb{A}_{l4}(k; t)$ and $\mathbb{A}_{r1}(k; t)$ for $k \in \mathbb{C}^+ \cup \mathbb{R}$ and $\mathbb{A}_{l1}(k; t)$ and $\mathbb{A}_{r4}(k; t)$ for $k \in \mathbb{C}^- \cup \mathbb{R}$ are time independent. Equations (6.4b) and (5.8b) imply

$$\mathbb{A}_{l2}(k; t) = e^{-4ik^2t} \mathbb{A}_{l2}(k; 0), \quad \mathbb{A}_{l3}(k; t) = e^{4ik^2t} \mathbb{A}_{l3}(k; 0), \quad (6.10a)$$

$$\mathbb{A}_{r2}(k; t) = e^{-4ik^2t} \mathbb{A}_{r2}(k; 0), \quad \mathbb{A}_{r3}(k; t) = e^{4ik^2t} \mathbb{A}_{r3}(k; 0), \quad (6.10b)$$

where $k \in \mathbb{R}$. Equations (6.5) and (5.10) imply

$$\mathbb{R}_{l1}(k; t) = e^{-4ik^2t} \mathbb{R}_{l1}(k; 0), \quad \mathbb{R}_{l4}(k; t) = e^{4ik^2t} \mathbb{R}_{l4}(k; 0), \quad (6.11a)$$

$$\mathbb{R}_{r1}(k; t) = e^{4ik^2t} \mathbb{R}_{r1}(k; 0), \quad \mathbb{R}_{r4}(k; t) = e^{-4ik^2t} \mathbb{R}_{r4}(k; 0), \quad (6.11b)$$

where $k \in \mathbb{R}$. Finally, (6.9) and (5.12) imply

$$\mathbb{N}_{l1;s}(t) = e^{-4ik_s^2t} \mathbb{N}_{l1;s}(0), \quad \mathbb{N}_{l4;s}(t) = e^{4ik_s^2t} \mathbb{N}_{l4;s}(0), \quad (6.12a)$$

$$\mathbb{N}_{r1;s}(t) = e^{4ik_s^2t} \mathbb{N}_{r1;s}(0), \quad \mathbb{N}_{r4;s}(t) = e^{-4ik_s^2t} \mathbb{N}_{r4;s}(0), \quad (6.12b)$$

where k_s is a simple pole of $A_{r,l}(k; t)^{-1}$ in \mathbb{C}^+ .

6.3 Marchenko kernels and time evolution

The Marchenko integral kernels of the matrix Schrödinger system can be expressed in terms of those of the AKNS system as follows:

$$\Omega_r(w; t) = \begin{pmatrix} 0_{m_1 \times m_1} & \Omega_{r4}(w; t) \\ \Omega_{r1}(w; t) & 0_{m_2 \times m_2} \end{pmatrix}, \quad \Omega_l(w; t) = \begin{pmatrix} 0_{m_1 \times m_1} & \Omega_{l1}(w; t) \\ \Omega_{l4}(w; t) & 0_{m_2 \times m_2} \end{pmatrix}, \quad (6.13)$$

where in the case of simple poles

$$\Omega_{r1}(w; t) = \hat{R}_{r1}(w; t) + \sum_{s=1}^N e^{ik_s w} e^{4ik_s^2 t} N_{r1;s}(0), \quad (6.14a)$$

$$\Omega_{r4}(w; t) = \hat{R}_{r4}(w; t) + \sum_{s=1}^N e^{ik_s w} e^{-4ik_s^2 t} N_{r4;s}(0), \quad (6.14b)$$

$$\Omega_{l1}(w; t) = \hat{R}_{l1}(w; t) + \sum_{s=1}^N e^{-ik_s w} e^{-4ik_s^2 t} N_{l1;s}(0), \quad (6.14c)$$

$$\Omega_{l4}(w; t) = \hat{R}_{l4}(w; t) + \sum_{s=1}^N e^{-ik_s w} e^{4ik_s^2 t} N_{l4;s}(0). \quad (6.14d)$$

Consequently, for simple poles we obtain

$$\Omega_r(w; t) = \hat{R}_r(w; t) + \sum_{s=1}^N e^{ik_s w} e^{-4ik_s^2 t} N_{r;s}(0), \quad (6.15a)$$

$$\Omega_l(w; t) = \hat{R}_l(w; t) + \sum_{s=1}^N e^{-ik_s w} e^{-4ik_s^2 t} N_{l;s}(0), \quad (6.15b)$$

which we could also have obtained directly from (4.11) and (6.13). As a result, we have arrived at the linear PDEs

$$[\Omega_r]_t = 4i\sigma_3[\Omega_r]_{\alpha\alpha}, \quad [\Omega_l]_t = 4i\sigma_3[\Omega_l]_{\alpha\alpha}, \quad (6.16)$$

in accordance with (6.6).

Let us discuss the adjoint symmetry relations of the Marchenko integral kernels. According to Corollary B.4, those of the matrix Schrödinger equation read

$$\Omega_r(w; t)^\dagger = \sigma_3 \Omega_r(w; t) \sigma_3, \quad \Omega_l(w; t)^\dagger = \sigma_3 \Omega_l(w; t) \sigma_3.$$

Using (6.13) this yields the adjoint symmetry relations for the AKNS system

$$\Omega_{r1}(w; t)^\dagger = -\Omega_{r4}(w; t), \quad \Omega_{r4}(w; t)^\dagger = -\Omega_{r1}(w; t), \quad (6.17a)$$

$$\Omega_{l1}(w; t)^\dagger = -\Omega_{l4}(w; t), \quad \Omega_{l4}(w; t)^\dagger = -\Omega_{l1}(w; t). \quad (6.17b)$$

7 Multisoliton solutions

In this section we apply the matrix triplet method to write the Marchenko integral kernels in separated form and solve them by separation of variables. This method has been successfully applied to the Korteweg-de Vries (KdV) equation [6, 11], the NLS equation [8, 25], the sine-Gordon equation [53, 9], the modified Korteweg-de Vries (mKdV) equation [23], and the Toda lattice equation [52]. An introduction to this method can be found in [55].

In the reflectionless case we can write the Marchenko integral kernels of the AKNS system as follows:

$$\Omega_{r1}(w; t) = \mathcal{C}_r e^{-w\mathcal{A}_r} e^{-4it\mathcal{A}_r^2} \mathcal{B}_r, \quad \Omega_{r4}(w; t) = -\mathcal{B}_r^\dagger e^{-w\mathcal{A}_r^\dagger} e^{4it\mathcal{A}_r^{\dagger 2}} \mathcal{C}_r^\dagger, \quad (7.1a)$$

$$\Omega_{l1}(w; t) = \mathcal{C}_l e^{w\mathcal{A}_l} e^{4it\mathcal{A}_l^2} \mathcal{B}_l, \quad \Omega_{l4}(w; t) = -\mathcal{B}_l^\dagger e^{w\mathcal{A}_l^\dagger} e^{-4it\mathcal{A}_l^{\dagger 2}} \mathcal{C}_l^\dagger, \quad (7.1b)$$

where we have taken into account the adjoint symmetry relations (6.17). Here $(\mathcal{A}_r, \mathcal{B}_r, \mathcal{C}_r)$ and $(\mathcal{A}_l, \mathcal{B}_l, \mathcal{C}_l)$ are matrix triplets, where all of the eigenvalues of \mathcal{A}_r and \mathcal{A}_l have a positive real part. Let us now define

$$\mathbf{A}_r = \begin{pmatrix} \mathcal{A}_r^\dagger & 0_{p \times p} \\ 0_{p \times p} & \mathcal{A}_r \end{pmatrix}, \quad \mathbf{B}_r = \begin{pmatrix} 0_{p \times m_1} & -\mathcal{C}_r^\dagger \\ \mathcal{B}_r & 0_{p \times m_2} \end{pmatrix}, \quad \mathbf{C}_r = \begin{pmatrix} \mathcal{B}_r^\dagger & 0_{m_1 \times p} \\ 0_{m_2 \times p} & \mathcal{C}_r \end{pmatrix},$$

$$\mathbf{A}_l = \begin{pmatrix} \mathcal{A}_l & 0_{p \times p} \\ 0_{p \times p} & \mathcal{A}_l^\dagger \end{pmatrix}, \quad \mathbf{B}_l = \begin{pmatrix} 0_{p \times m_1} & \mathcal{B}_l \\ -\mathcal{C}_l^\dagger & 0_{p \times m_2} \end{pmatrix}, \quad \mathbf{C}_l = \begin{pmatrix} \mathcal{C}_l & 0_{m_1 \times p} \\ 0_{m_2 \times p} & \mathcal{B}_l^\dagger \end{pmatrix},$$

where \mathcal{A}_r and \mathcal{A}_l are both matrices of order p whose eigenvalues have a positive real part. Putting $\tilde{\sigma}_3 = \sigma_3 \otimes I_p = I_p \oplus (-I_p)$, we obtain

$$\Omega_r(w; t) = \mathbf{C}_r e^{-w\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{B}_r, \quad (7.2a)$$

$$\Omega_l(w; t) = \mathbf{C}_l e^{w\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{B}_l, \quad (7.2b)$$

where

$$\tilde{\sigma}_3 \mathbf{A}_{r,l} = \mathbf{A}_{r,l} \tilde{\sigma}_3, \quad \mathbf{B}_{r,l} \sigma_3 = -\tilde{\sigma}_3 \mathbf{B}_{r,l}, \quad \sigma_3 \mathbf{C}_{r,l} = \mathbf{C}_{r,l} \tilde{\sigma}_3. \quad (7.3)$$

Solving the Marchenko integral equations we get

$$K(x, y, t) = -\mathbf{W}_r(x; t) e^{-y\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{B}_r, \quad (7.4a)$$

$$J(x, y, t) = -\mathbf{W}_l(x; t) e^{y\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{B}_l, \quad (7.4b)$$

where

$$\mathbf{W}_r(x; t) = \mathbf{C}_r e^{-x\mathbf{A}_r} + \int_x^\infty dz K(x, z, t) \mathbf{C}_r e^{-z\mathbf{A}_r},$$

$$\mathbf{W}_l(x; t) = \mathbf{C}_l e^{x\mathbf{A}_l} + \int_{-\infty}^x dz J(x, z, t) \mathbf{C}_l e^{z\mathbf{A}_l}.$$

Substituting (7.4) into (4.10) and solving (4.10) for $\mathbf{W}_{r,l}(x; t)$ we get

$$\mathbf{W}_r(x; t) = \mathbf{C}_r e^{-x\mathbf{A}_r} \left[I_{2p} + e^{-x\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{P}_r e^{-x\mathbf{A}_r} \right]^{-1},$$

$$\mathbf{W}_l(x; t) = \mathbf{C}_l e^{x\mathbf{A}_l} \left[I_{2p} + e^{x\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{P}_l e^{x\mathbf{A}_l} \right]^{-1},$$

provided the inverse matrices exist. Here

$$\begin{aligned}\mathbf{P}_r &= \int_0^\infty dz e^{-z\mathbf{A}_r} \mathbf{B}_r \mathbf{C}_r e^{-z\mathbf{A}_r} = \begin{pmatrix} 0_{p \times p} & \mathcal{Q}_{r,\Sigma} \\ \mathcal{N}_{r,\Xi} & 0_{p \times p} \end{pmatrix}, \\ \mathbf{P}_l &= \int_0^\infty dz e^{-z\mathbf{A}_l} \mathbf{B}_l \mathbf{C}_l e^{-z\mathbf{A}_l} = \begin{pmatrix} 0_{p \times p} & \mathcal{N}_{l,\Xi} \\ \mathcal{Q}_{l,\Sigma} & 0_{p \times p} \end{pmatrix},\end{aligned}$$

are the unique solutions of the Sylvester equations

$$\mathbf{A}_r \mathbf{P}_r + \mathbf{P}_r \mathbf{A}_r = \mathbf{B}_r \mathbf{C}_r, \quad \mathbf{A}_l \mathbf{P}_l + \mathbf{P}_l \mathbf{A}_l = \mathbf{B}_l \mathbf{C}_l. \quad (7.5)$$

More precisely, given $(x, t) \in \mathbb{R}^2$, the Marchenko integral equations (4.10) are uniquely solvable (in an L^1 -setting) iff the algebraic equations for $\mathbf{W}_{r,l}(x; t)$ are uniquely solvable. Consequently,

$$\begin{aligned}K(x, y, t) &= -\mathbf{C}_r e^{-x\mathbf{A}_r} \left[I_{2p} + e^{-x\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{P}_r e^{-x\mathbf{A}_r} \right]^{-1} e^{-y\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{B}_r \\ &= -\mathbf{C}_r \left[I_{2p} + e^{-2x\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{P}_r \right]^{-1} e^{-(x+y)\mathbf{A}_r} e^{4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{B}_r \\ &= -\mathbf{C}_r \mathbf{\Pi}_r(x; t)^{-1} e^{-(y-x)\mathbf{A}_r} \mathbf{B}_r,\end{aligned} \quad (7.6a)$$

$$\begin{aligned}J(x, y, t) &= -\mathbf{C}_l e^{x\mathbf{A}_l} \left[I_{2p} + e^{x\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{P}_l e^{x\mathbf{A}_l} \right]^{-1} e^{y\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{B}_l \\ &= -\mathbf{C}_l \left[I_{2p} + e^{2x\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{P}_l \right]^{-1} e^{(x+y)\mathbf{A}_l} e^{4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{B}_l \\ &= -\mathbf{C}_l \mathbf{\Pi}_l(x; t)^{-1} e^{-(x-y)\mathbf{A}_l} \mathbf{B}_l,\end{aligned} \quad (7.6b)$$

where

$$\mathbf{\Pi}_r(x; t) = e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} + \mathbf{P}_r, \quad \mathbf{\Pi}_l(x; t) = e^{-2x\mathbf{A}_l} e^{-4it\tilde{\sigma}_3 \mathbf{A}_l^2} + \mathbf{P}_l. \quad (7.7)$$

Observe that, for $(x, t) \in \mathbb{R}^2$, the matrices $\mathbf{\Pi}_{r,l}(x; t)$ are invertible iff the Marchenko integral equations (4.10) are uniquely solvable. Using (3.5) [with $\mathbf{Q} = \mathbf{Q}^2 + \mathbf{Q}_x$ instead of \mathbf{Q}] and (3.19), we get

$$-\mathcal{Q}(x; t) + \int_x^\infty dy \mathcal{Q}(y; t)^2 = 2K(x, x; t) = -2\mathbf{C}_r \mathbf{\Pi}_r(x; t)^{-1} \mathbf{B}_r, \quad (7.8a)$$

$$\mathcal{Q}(x; t) + \int_{-\infty}^x dy \mathcal{Q}(y; t)^2 = 2J(x, x; t) = -2\mathbf{C}_l \mathbf{\Pi}_l(x; t)^{-1} \mathbf{B}_l, \quad (7.8b)$$

where $\mathbf{\Pi}_r(x; t)$ and $\mathbf{\Pi}_l(x; t)$ are defined by (7.7). The matrix NLS solution then equals the \mp block off-diagonal parts of (7.8), where we recover the

explicit (matrix) NLS solutions obtained before by the matrix triplet method [8, 25, 55]. Differentiating (7.8) with respect to x and using (1.4) we obtain the following solutions of the nonlocal integrable equation (2.2):

$$\begin{aligned} \mathbf{Q}(x; t) &= -\frac{d}{dx} \left(-2\mathbf{C}_r \mathbf{\Pi}_r(x; t)^{-1} \mathbf{B}_r \right) \\ &= -4\mathbf{C}_r \mathbf{\Pi}_r(x; t)^{-1} \mathbf{A}_r e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{\Pi}_r(x; t)^{-1} \mathbf{B}_r, \end{aligned} \quad (7.9a)$$

$$\begin{aligned} \mathbf{Q}(x; t) &= +\frac{d}{dx} \left(-2\mathbf{C}_l \mathbf{\Pi}_l(x; t)^{-1} \mathbf{B}_l \right) \\ &= -4\mathbf{C}_l \mathbf{\Pi}_l(x; t)^{-1} \mathbf{A}_l e^{-2x\mathbf{A}_l} e^{-4it\tilde{\sigma}_3 \mathbf{A}_l^2} \mathbf{\Pi}_l(x; t)^{-1} \mathbf{B}_l. \end{aligned} \quad (7.9b)$$

Acknowledgments

The authors have been partially supported by the Regione Autonoma della Sardegna research project *Algorithms and Models for Imaging Science* [AMIS] (RASSR57257, intervento finanziato con risorse FSC 2014-2020 – Patto per lo Sviluppo della Regione Sardegna), and by INdAM-GNFM.

A Deriving the nonlocal integrable equation

In this appendix we present an alternative proof of (2.3). Indeed, let us start from the matrix NLS equation

$$i\sigma_3 \mathcal{Q}_t + \mathcal{Q}_{xx} - 2\mathcal{Q}^3 = 0_{n \times n},$$

where we seek solutions \mathcal{Q} that anticommute with σ_3 . Putting

$$\mathcal{Q} = \mathcal{Q}^2 + \mathcal{Q}_x,$$

so that

$$\begin{aligned} \mathcal{Q}_x &= \mathcal{Q}_x \mathcal{Q} + \mathcal{Q} \mathcal{Q}_x + \mathcal{Q}_{xx}, \\ \mathcal{Q}_{xx} &= \mathcal{Q}_{xx} \mathcal{Q} + 2\mathcal{Q}_x^2 + \mathcal{Q} \mathcal{Q}_{xx} + \mathcal{Q}_{xxx}, \end{aligned}$$

we obtain

$$\begin{aligned}
-i\sigma_3\mathcal{Q}_t &= -(i\sigma_3\mathcal{Q}_t)\mathcal{Q} + \mathcal{Q}(i\sigma_3\mathcal{Q}_t) - (i\sigma_3\mathcal{Q}_t)_x \\
&= (\mathcal{Q}_{xx} - 2\mathcal{Q}^3)\mathcal{Q} - \mathcal{Q}(\mathcal{Q}_{xx} - 2\mathcal{Q}^3) + \mathcal{Q}_{xxx} - 2(\mathcal{Q}^3)_x \\
&= \mathcal{Q}_{xx}\mathcal{Q} - \mathcal{Q}\mathcal{Q}_{xx} + \mathcal{Q}_{xxx} - 2\mathcal{Q}_x\mathcal{Q}^2 - 2\mathcal{Q}\mathcal{Q}_x\mathcal{Q} - 2\mathcal{Q}^2\mathcal{Q}_x \\
&= (\mathcal{Q}_x - \mathcal{Q}_x\mathcal{Q} - \mathcal{Q}\mathcal{Q}_x)\mathcal{Q} - \mathcal{Q}(\mathcal{Q}_x - \mathcal{Q}_x\mathcal{Q} - \mathcal{Q}\mathcal{Q}_x) \\
&+ (\mathcal{Q}_{xx} - \mathcal{Q}_{xx}\mathcal{Q} - 2\mathcal{Q}_x^2 - \mathcal{Q}\mathcal{Q}_{xx}) - 2\mathcal{Q}_x\mathcal{Q}^2 - 2\mathcal{Q}\mathcal{Q}_x\mathcal{Q} - 2\mathcal{Q}^2\mathcal{Q}_x \\
&= \mathcal{Q}_x\mathcal{Q} - \mathcal{Q}\mathcal{Q}_x + \mathcal{Q}_{xx} \\
&- 3\mathcal{Q}_x\mathcal{Q}^2 - 2\mathcal{Q}\mathcal{Q}_x\mathcal{Q} - \mathcal{Q}^2\mathcal{Q}_x - \mathcal{Q}_{xx}\mathcal{Q} - 2\mathcal{Q}_x^2 - \mathcal{Q}\mathcal{Q}_{xx} \\
&= \mathcal{Q}_x\mathcal{Q} - \mathcal{Q}\mathcal{Q}_x + \mathcal{Q}_{xx} - \mathcal{Q}_x\mathcal{Q} - \mathcal{Q}\mathcal{Q}_x - 2\mathcal{Q}_x\mathcal{Q}^2 - 2\mathcal{Q}_x^2 \\
&= \mathcal{Q}_{xx} - 2\mathcal{Q}\mathcal{Q}_x - 2\mathcal{Q}_x\mathcal{Q}^2 - 2\mathcal{Q}_x^2 \\
&= \mathcal{Q}_{xx} - 2\mathcal{Q}\mathcal{Q}_x - 2\mathcal{Q}_x\mathcal{Q},
\end{aligned}$$

where $\mathcal{Q}_x = \frac{1}{2}(\mathcal{Q} - \sigma_3\mathcal{Q}\sigma_3)$ and hence

$$\mathcal{Q}(x; t) = - \int_x^\infty dy \frac{1}{2} (\mathcal{Q} - \sigma_3\mathcal{Q}\sigma_3) = \int_{-\infty}^x dy \frac{1}{2} (\mathcal{Q} - \sigma_3\mathcal{Q}\sigma_3).$$

Here the integrand depends on $(y, t) \in \mathbb{R}^2$. We have thus reproduced (2.3) by a more direct method.

B Adjoint symmetry of Marchenko kernels

In this appendix we prove the adjoint symmetry relations (4.12) for the Marchenko integral kernels and (4.13) for the norming constants. Even though these relations are easily established for scalar Schrödinger equations [28, 22, 19, 20] or for matrix Schrödinger equations without discrete eigenvalues (where they are immediate from (3.14)), they require considerably more effort in the general matrix Schrödinger case.

Let $G_l(x, k)$ and $G_r(x, k)$ be the $n \times n$ matrix solutions of the dual matrix Schrödinger equation

$$-\phi''(x, k) + \phi(x, k)\mathcal{Q}(x) = k^2\phi(x, k) \quad (\text{B.1})$$

under the asymptotic conditions

$$G_l(x, k) = e^{ikx} [I_n + o(1)], \quad x \rightarrow +\infty, \quad (\text{B.2a})$$

$$G_r(x, k) = e^{-ikx} [I_n + o(1)], \quad x \rightarrow -\infty. \quad (\text{B.2b})$$

Then the dual Faddeev functions satisfy the Volterra integral equations

$$e^{-ikx}G_l(x, k) = I_n + \int_x^\infty dy \frac{e^{2ik(y-x)} - 1}{2ik} e^{-iky}G_l(y, k)\mathbf{Q}(y), \quad (\text{B.3a})$$

$$e^{ikx}G_r(x, k) = I_n + \int_{-\infty}^x dy \frac{e^{2ik(x-y)} - 1}{2ik} e^{iky}G_r(y, k)\mathbf{Q}(y). \quad (\text{B.3b})$$

In analogy with Proposition 3.1 we prove

Proposition B.1 *The dual Faddeev functions $e^{-ikx}G_l(x, k)$ and $e^{ikx}G_r(x, k)$ are continuous in $k \in \mathbb{C}^+ \cup \mathbb{R}$, are analytic in $k \in \mathbb{C}^+$, and tend to I_n as $k \rightarrow \infty$ from within $\mathbb{C}^+ \cup \mathbb{R}$, irrespective of the choice of $x \in \mathbb{R}$.*

Using the symmetry relation (1.2) in the matrix Schrödinger equation (1.5) with k replaced by $-k^*$, we see that $\sigma_3\Psi(x, -k^*)^\dagger\sigma_3$ satisfies the dual matrix Schrödinger equation (B.1) whenever $\Psi(x, k)$ satisfies the original matrix Schrödinger equation (1.5). Equations (3.1) and (B.2) then imply the following adjoint symmetry relations:

$$G_{r,l}(x, k) = \sigma_3 F_{r,l}(x, -k^*)^\dagger \sigma_3. \quad (\text{B.4})$$

These relations can also be derived directly from the Volterra integral equations (3.2) and (B.3). With the help of (3.3), (3.10), and (B.4) we obtain for $0 \neq k \in \mathbb{R}$ the asymptotic relations

$$\begin{aligned} G_l(x, k) &= e^{ikx}A_r(k) - e^{-ikx}B_r(-k) + o(1), & x \rightarrow -\infty, \\ G_r(x, k) &= e^{-ikx}A_l(k) - e^{ikx}B_l(-k) + o(1), & x \rightarrow +\infty. \end{aligned}$$

Putting

$$\begin{aligned} \mathbf{G}_r(x, k) &= \begin{pmatrix} G'_r(x, -k) & -G_r(x, -k) \\ -G'_r(x, k) & G_r(x, k) \end{pmatrix} = (\sigma_2 \otimes \sigma_3)\mathbf{F}_r(x, -k)^\dagger(\sigma_2 \otimes \sigma_3), \\ \mathbf{G}_l(x, k) &= \begin{pmatrix} G'_l(x, k) & -G_l(x, k) \\ -G'_l(x, -k) & G_l(x, -k) \end{pmatrix} = (\sigma_2 \otimes \sigma_3)\mathbf{F}_l(x, -k)^\dagger(\sigma_2 \otimes \sigma_3), \end{aligned}$$

where $\sigma_2 \otimes \sigma_3 = \begin{bmatrix} 0_{n \times n} & -i\sigma_3 \\ i\sigma_3 & 0_{n \times n} \end{bmatrix}$, we get for $0 \neq k \in \mathbb{R}$

$$\mathbf{G}_r(x, k) = \begin{pmatrix} A_l(-k) & B_l(k) \\ B_l(-k) & A_l(k) \end{pmatrix} \mathbf{G}_l(x, k), \quad (\text{B.5a})$$

$$\mathbf{G}_l(x, k) = \begin{pmatrix} A_r(k) & B_r(-k) \\ B_r(k) & A_r(-k) \end{pmatrix} \mathbf{G}_r(x, k), \quad (\text{B.5b})$$

where the matrices containing the A and B quantities are each other's inverses. By analytic continuation we obtain for $k \in \mathbb{C}^+ \cup \mathbb{R}$

$$\mathbf{G}_{r,l}(x, k) = (\sigma_2 \otimes \sigma_3) \mathbf{F}_{r,l}(x, -k^*)^\dagger (\sigma_2 \otimes \sigma_3).$$

Using (B.5) and (3.12) we easily derive the Riemann-Hilbert problem

$$\begin{pmatrix} G_r(x, -k) \\ G_l(x, -k) \end{pmatrix} = \begin{pmatrix} A_r(k)^{-1} & -R_l(k) \\ -R_r(k) & A_l(k)^{-1} \end{pmatrix} \begin{pmatrix} G_l(x, k) \\ G_r(x, k) \end{pmatrix}, \quad (\text{B.6})$$

where we are either in the generic case or in the superexceptional case and there are no spectral singularities.

Assume that the poles k_s of $A_r(k)^{-1}$ are simple and hence (4.1) is true. Rewriting the bottom half of (B.6) we get

$$\begin{aligned} e^{ikx} G_l(x, -k) &= A_{l0}(k)^{-1} e^{ikx} G_r(x, k) - e^{2ikx} R_r(k) e^{-ikx} G_l(x, k) \\ &= A_{l0}(k) e^{ikx} G_r(x, k) + \sum_{s=1}^N \tau_{l;s} \frac{e^{ikx} G_r(x, k) - e^{ik_s x} G_r(x, k_s)}{k - k_s} \\ &\quad + i \sum_{s=1}^N \frac{e^{2ik_s x} \mathbb{N}_{r;s}}{k - k_s} e^{-ikx} G_l(x, k) - e^{2ikx} R_r(k) e^{-ikx} G_l(x, k), \end{aligned} \quad (\text{B.7})$$

where

$$\tau_{l;s} G_r(x, k_s) = i \mathbb{N}_{r;s} G_l(x, k_s) \quad (\text{B.8})$$

for the ‘‘dual’’ norming constants $\mathbb{N}_{r;s}$. By the same token,

$$\begin{aligned} e^{-ikx} G_r(x, -k) &= A_r(k)^{-1} e^{-ikx} G_l(x, k) - e^{-2ikx} R_l(k) e^{ikx} G_r(x, k) \\ &= A_{r0}(k) e^{-ikx} G_l(x, k) + \sum_{s=1}^N \tau_{r;s} \frac{e^{-ikx} G_l(x, k) - e^{-ik_s x} G_l(x, k_s)}{k - k_s} \\ &\quad + i \sum_{s=1}^N \frac{e^{-2ik_s x} \mathbb{N}_{l;s}}{k - k_s} e^{ikx} G_r(x, k) - e^{-2ikx} R_l(k) e^{ikx} G_r(x, k), \end{aligned} \quad (\text{B.9})$$

where

$$\tau_{r;s} G_l(x, k_s) = i \mathbb{N}_{l;s} G_r(x, k_s) \quad (\text{B.10})$$

for another set of ‘‘dual’’ norming constants $\mathbb{N}_{l;s}$. For every $x \in \mathbb{R}$ the identities (B.7) and (B.9) are equations in $\mathcal{W}^{n \times n}$. Using Π_- to project these

two equations onto $\mathcal{W}_-^{n \times n}$ along $\mathcal{W}_{+,0}^{n \times n}$, we obtain

$$\begin{aligned} e^{ikx} G_l(x, -k) &= I_n + i \sum_{s=1}^N \frac{e^{2ik_s x} \mathbb{N}_{r;s}}{k - k_s} e^{-ikx} G_l(x, k) \\ &\quad - \Pi_- [e^{2ikx} R_r(k) e^{-ikx} G_l(x, k)], \end{aligned} \quad (\text{B.11a})$$

$$\begin{aligned} e^{-ikx} G_r(x, -k) &= I_n + i \sum_{s=1}^N \frac{e^{-2ik_s x} \mathbb{N}_{l;s}}{k - k_s} e^{ikx} G_r(x, k) \\ &\quad - \Pi_- [e^{-2ikx} R_l(k) e^{ikx} G_r(x, k)]. \end{aligned} \quad (\text{B.11b})$$

Using the triangular representations

$$e^{-ikx} G_l(x, k) = I_n + \int_x^\infty dy e^{ik(y-x)} \mathbb{K}(y, x), \quad (\text{B.12a})$$

$$e^{ikx} G_r(x, k) = I_n + \int_{-\infty}^x dy e^{ik(x-y)} \mathbb{J}(y, x), \quad (\text{B.12b})$$

in (B.11) and stripping off the Fourier transforms we get the dual Marchenko equations

$$\mathbb{K}(y, x) + \omega_r(y + x) + \int_x^\infty dz \omega_r(y + z) \mathbb{K}(z, x) = 0_{n \times n}, \quad (\text{B.13a})$$

$$\mathbb{J}(y, x) + \omega_l(y + x) + \int_{-\infty}^x dz \omega_l(y + z) \mathbb{J}(z, x) = 0_{n \times n}, \quad (\text{B.13b})$$

where the dual Marchenko kernels are given by

$$\begin{aligned} \omega_r(w) &= \hat{R}_r(w) + \sum_{s=1}^N e^{ik_s w} \mathbb{N}_{r;s}, \\ \omega_l(w) &= \hat{R}_l(w) + \sum_{s=1}^N e^{-ik_s w} \mathbb{N}_{l;s}. \end{aligned}$$

Theorem B.2 *Let the generic case or the superexceptional case be satisfied and assume there do not exist any spectral singularities. Then the dual Marchenko kernels are related to the Marchenko kernels as follows:*

$$\omega_r(w) = \sigma_3 \Omega_r(w)^\dagger \sigma_3, \quad \omega_l(w) = \sigma_3 \Omega_l(w)^\dagger \sigma_3. \quad (\text{B.14})$$

Proof. Taking the adjoint in (4.10a), we see that the matrix functions $\sigma_3 \Omega_{l,r}(w)^\dagger \sigma_3$ satisfy the dual Marchenko equations (B.13), where

$$\mathbb{K}(y, x) = \sigma_3 K(x, y)^\dagger \sigma_3, \quad \mathbb{J}(y, x) = \sigma_3 J(x, y)^\dagger \sigma_3.$$

Since the Marchenko equation (B.13a) is uniquely solvable in $L^1(x, +\infty)^{n \times n}$ for large enough x and (B.13b) is uniquely solvable in $L^1(-\infty, x)^{n \times n}$ for large enough $-x$, we obtain the symmetry relations

$$\omega_r(w) = \sigma_3 \Omega_r(w)^\dagger \sigma_3, \quad \omega_l(w) = \sigma_3 \Omega_l(w)^\dagger \sigma_3, \quad (\text{B.15})$$

where $w \geq x_0 > -\infty$ in the first identity and $w \leq x_0 < +\infty$ in the second identity. Since the Fourier transformed reflection coefficients satisfy

$$\hat{R}_r(\alpha)^\dagger = \sigma_3 \hat{R}_r(\alpha) \sigma_3, \quad \hat{R}_l(\alpha)^\dagger = \sigma_3 \hat{R}_l(\alpha) \sigma_3,$$

we see that $\Omega_r(w) - \hat{R}_r(w)$ is an entire analytic matrix function of w that satisfies (B.15) for $w \geq x_0 > -\infty$. Similarly, $\Omega_l(w) - \hat{R}_l(w)$ is an entire analytic matrix function that satisfies (B.15) for $w \leq x_0 < +\infty$. By analytic continuation, (B.14) holds true. \blacksquare

We immediately have the symmetry relations for the norming constants

$$N_{r;s} = \sigma_3 \mathbb{N}_{r;s}^\dagger \sigma_3, \quad N_{l;s} = \sigma_3 \mathbb{N}_{l;s}^\dagger \sigma_3,$$

provided k_s is a simple pole of $A_r(k)^{-1}$.

Theorem B.3 *Let the generic case or the superexceptional case be satisfied and assume there do not exist any spectral singularities. Then the dual Marchenko kernels satisfy*

$$\omega_r(w) = \Omega_r(w), \quad \omega_l(w) = \Omega_l(w).$$

Proof. Using (3.16) and (B.12) we easily verify that for $0 \neq k \in \mathbb{R}$

$$\begin{aligned} & F_r(x, k)G_r(x, -k) - F_r(x, -k)G_r(x, k) \\ &= F_r(x, k) [A_r(k)^{-1}G_l(x, k) - R_l(k)G_r(x, k)] \\ &= [-F_r(x, k)R_l(k) + F_l(x, k)A_l(k)^{-1}] G_r(x, k) \\ &= F_r(x, k)A_r(k)^{-1}G_l(x, k) - F_l(x, k)A_l(k)^{-1}G_r(x, k), \end{aligned}$$

which can be meromorphically extended to the upper half-plane. In the same way we get

$$\begin{aligned}
F_l(x, k)G_l(x, -k) - F_l(x, -k)G_l(x, k) \\
&= F_l(x, k) [-R_r(k)G_l(x, k) + A_l(k)^{-1}G_r(x, k)] \\
&\quad - [F_r(x, k)A_r(k)^{-1} - F_l(x, k)R_r(k)] G_l(x, k) \\
&= F_l(x, k)A_l(k)^{-1}G_r(x, k) - F_r(x, k)A_r(k)^{-1}G_l(x, k),
\end{aligned}$$

which can be meromorphically extended to the upper half-plane. If k_s is a simple pole of $A_r(k)^{-1}$ and hence of $A_l(k)^{-1}$, we get by taking the residues

$$F_r(x, k_s)\tau_{r;s}G_l(x, k_s) - F_l(x, k_s)\tau_{l;s}G_r(x, k_s) \quad (\text{B.16})$$

as well as the negative of this expression. Note that either term in the expression (B.16) is exponentially decaying as $x \rightarrow \pm\infty$. Using (4.6), (4.8), (B.8), and (B.10), we can write the latter expression in the two equivalent forms

$$iF_l(x, k_s) [N_{r;s} - \mathbb{N}_{r;s}] G_l(x, k_s) = iF_r(x, k_s) [\mathbb{N}_{l;s} - N_{l;s}] G_r(x, k_s).$$

Utilizing the asymptotic behavior of the Jost and dual Jost matrices as $x \rightarrow \pm\infty$, we obtain $e^{2ik_s x} [N_{r;s} - \mathbb{N}_{r;s}]$ as $x \rightarrow +\infty$ and $e^{-2ik_s x} [\mathbb{N}_{l;s} - N_{l;s}]$ as $x \rightarrow -\infty$. Therefore,

$$\mathbb{N}_{r;s} = N_{r;s}, \quad \mathbb{N}_{l;s} = N_{l;s}.$$

Consequently, Theorem B.3 is true if the poles of $A_{r,l}(k)^{-1}$ are simple. \blacksquare

Corollary B.4 *Let the generic case or the superexceptional case be satisfied and assume there do not exist any spectral singularities. Then the Marchenko kernels satisfy*

$$\Omega_r(w) = \sigma_3 \Omega_r(w)^\dagger \sigma_3, \quad \Omega_l(w) = \sigma_3 \Omega_l(w)^\dagger \sigma_3.$$

In other words, the Marchenko kernels are σ_3 -hermitian.

C Direct substitution

In this appendix we derive the multisoliton solution $\mathcal{Q}(x; t)$ of the nonlocal nonlinear evolution equation (2.3) in the reflectionless case by substituting

the derivatives of the expressions

$$\int_x^\infty dy \mathbf{Q}(y; t) = 2K(x, x; t) = -2\mathbf{C}_r \mathbf{\Pi}_r(x; t)^{-1} \mathbf{B}_r, \quad (\text{C.1a})$$

$$\int_{-\infty}^x dy \mathbf{Q}(y; t) = 2J(x, x; t) = -2\mathbf{C}_l \mathbf{\Pi}_l(x; t)^{-1} \mathbf{B}_l, \quad (\text{C.1b})$$

with respect to $x \in \mathbb{R}$ directly into (2.3), where $\mathbf{\Pi}_{r,l}(x; t)$ are defined by (7.7).

Recall that $\boldsymbol{\sigma}_3 = I_{m_1} \oplus (-I_{m_2})$ and $\tilde{\sigma}_3 = I_p \oplus (-I_p)$, where p is the order of the matrices $A_{r,l}$ in (C.1) and (7.3) is satisfied. Then (7.3) implies the intertwining relations

$$\tilde{\sigma}_3 \mathbf{P}_{r,l} = -\mathbf{P}_{r,l} \tilde{\sigma}_3, \quad (\text{C.2a})$$

$$\tilde{\sigma}_3 \mathbf{\Pi}_{r,l}(x; t) = \tilde{\mathbf{\Pi}}_{r,l}(x; t) \tilde{\sigma}_3, \quad (\text{C.2b})$$

where

$$\tilde{\mathbf{\Pi}}_r(x; t) = e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} - \mathbf{P}_r, \quad \tilde{\mathbf{\Pi}}_l(x; t) = e^{-2x\mathbf{A}_l} e^{-4it\tilde{\sigma}_3 \mathbf{A}_l^2} - \mathbf{P}_l.$$

Theorem C.1 *Let $(\mathbf{A}_r, \mathbf{B}_r, \mathbf{C}_r)$ be a matrix triplet satisfying (7.3) for which the Sylvester equation (7.5) has a unique solution \mathbf{P}_r . Then the matrix function $\mathbf{Q}(x; t)$ defined by*

$$\mathbf{Q}(x; t) = -4\mathbf{C}_r \mathbf{\Pi}_r(x; t)^{-1} \mathbf{A}_r e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{\Pi}_r(x; t)^{-1} \mathbf{B}_r$$

satisfies the nonlinear evolution equation (2.3) in those $(x, t) \in \mathbb{R}^2$ for which the matrix $\mathbf{\Pi}_r(x; t)$ is nonsingular.

Proof. Let us introduce the abbreviations

$$\mathbf{E} = e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2}, \quad [\dots] = \mathbf{\Pi}(x; t) = \mathbf{E} + \mathbf{P}_r = e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} + \mathbf{P}_r,$$

so that

$$[\bullet] = \tilde{\mathbf{\Pi}}(x; t) = \tilde{\sigma}_3 [\dots] \tilde{\sigma}_3 = \mathbf{E} - \mathbf{P}_r = e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} - \mathbf{P}_r.$$

Then $\mathbf{E}_x = 2\mathbf{A}_r \mathbf{E} = 2\mathbf{E} \mathbf{A}_r$. In terms of these abbreviations we now define the product

$$\Gamma_{,j_1, j_2, \dots, j_r} = [\dots]^{-1} \mathbf{A}_r^{j_1} \mathbf{E} [\dots]^{-1} \mathbf{A}_r^{j_2} \mathbf{E} [\dots]^{-1} \dots [\dots]^{-1} \mathbf{A}_r^{j_r} \mathbf{E} [\dots]^{-1}$$

consisting of $r+1$ factors $[\dots]^{-1}$ interrupted by the consecutive factors $\mathbf{A}_r^{j_s} \mathbf{E}$, where j_s is a nonnegative integer ($s = 1, 2, \dots, r$). Then (7.9a) implies that

$$\mathbf{Q}(x; t) = -4\mathbf{C}_r [\dots]^{-1} \mathbf{A}_r \mathbf{E} [\dots]^{-1} \mathbf{B}_r = -4\mathbf{C}_r \Gamma_{,1} \mathbf{B}_r. \quad (\text{C.3})$$

In the subscript string $, j_1, \dots, j_r$, we replace a comma by a semicolon if the factor $[\bullet]^{-1}$ takes the place of $[\dots]^{-1}$. Using (7.3) we get from (C.3)

$$\sigma_3 \mathbf{Q}(x; t) \sigma_3 = 4\mathbf{C}_r [\bullet]^{-1} \mathbf{A}_r \mathbf{E} [\bullet]^{-1} \mathbf{B}_r = 4\mathbf{C}_r \Gamma_{;1} \mathbf{B}_r. \quad (\text{C.4})$$

Using (C.3), we easily derive the identity

$$\begin{aligned} [\Gamma_{,j_1, j_2, \dots, j_r}]_x &= -2\Gamma_{,1, j_1, \dots, j_r} - 2\Gamma_{,j_1, 1, j_2, \dots, j_r} - \dots - 2\Gamma_{,j_1, \dots, j_r, 1} \\ &\quad + 2\Gamma_{,j_1+1, j_2, \dots, j_r} + 2\Gamma_{,j_1, j_2+1, j_3, \dots, j_r} + \dots + 2\Gamma_{,j_1, \dots, j_r-1, j_r+1}, \end{aligned}$$

where we have first differentiated the $r+1$ factors $[\dots]^{-1}$ and then the factors $\mathbf{A}_r^{j_s} \mathbf{E}$ ($s = 1, 2, \dots, r$). A similar differentiation rule holds if some (or all) of the subscripted commas are replaced by semicolons.

Using the above x -differentiation rule we get

$$\mathbf{Q}_x = 8\mathbf{C}_r (2\Gamma_{,1,1} - \Gamma_{,2}) \mathbf{B}_r. \quad (\text{C.5})$$

Using the above x -differentiation rule again we get

$$\mathbf{Q}_{xx} = 16\mathbf{C}_r (-6\Gamma_{,1,1,1} + 3\Gamma_{,1,2} + 3\Gamma_{,2,1} - \Gamma_{,3}) \mathbf{B}_r. \quad (\text{C.6})$$

Next, using (C.3) and (C.4) we compute

$$\begin{aligned} \mathcal{Q}_x &= \frac{1}{2}(\mathbf{Q} - \sigma_3 \mathbf{Q} \sigma_3) = -2\mathbf{C}_r \Gamma_{,1} \mathbf{B}_r - 2\mathbf{C}_r \Gamma_{;1} \mathbf{B}_r \\ &= \frac{\partial}{\partial x} \mathbf{C}_r \left(\left[e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} + \mathbf{P}_r \right]^{-1} + \left[e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} - \mathbf{P}_r \right]^{-1} \right) \mathbf{B}_r, \end{aligned}$$

implying that

$$\begin{aligned} \mathcal{Q}(x; t) &= 2\mathbf{C}_r \left[e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} \pm \mathbf{P}_r \right]^{-1} \times \\ &\quad \times e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} \left[e^{2x\mathbf{A}_r} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mp \mathbf{P}_r \right]^{-1} \mathbf{B}_r \\ &= 2\mathbf{C}_r \Gamma_{,0} \mathbf{B}_r = 2\mathbf{C}_r \Gamma_{;0} \mathbf{B}_r, \end{aligned} \quad (\text{C.7})$$

where we have used the identity

$$[\dots]^{-1} \mathbf{E}[\bullet]^{-1} = [\bullet]^{-1} \mathbf{E}[\dots]^{-1} = \frac{1}{2} ([\dots]^{-1} + [\bullet]^{-1}). \quad (\text{C.8})$$

Using (7.5) we get

$$\mathbf{B}_r \mathbf{C}_r = \mathbf{A}_r[\dots] + [\dots] \mathbf{A}_r - 2\mathbf{A}_r \mathbf{E}, \quad (\text{C.9a})$$

$$\mathbf{B}_r \mathbf{C}_r = \mathbf{A}_r[\dots] - [\bullet] \mathbf{A}_r. \quad (\text{C.9b})$$

Equations (C.3) and (C.9a) imply the identity

$$\mathbf{Q}(x; t)^2 = 16\mathbf{C}_r (\Gamma_{,1,2} + \Gamma_{,2,1} - 2\Gamma_{,1,1,1}) \mathbf{B}_r. \quad (\text{C.10})$$

Equations (C.9b) and (7.3) imply the identity

$$\begin{aligned} \sigma_3 \mathbf{Q} \sigma_3 \mathbf{Q} &= -16\mathbf{C}_r \Gamma_{,1}; \mathbf{B}_r \mathbf{C}_r \Gamma_{,1}, \mathbf{B}_r \\ &= -16\mathbf{C}_r \Gamma_{,1}; (\mathbf{A}_r[\dots] - [\bullet] \mathbf{A}_r) \Gamma_{,1}, \mathbf{B}_r \\ &= -16\mathbf{C}_r [\bullet]^{-1} \mathbf{A}_r \mathbf{E}[\bullet]^{-1} (\mathbf{A}_r[\dots] - [\bullet] \mathbf{A}_r) [\dots]^{-1} \mathbf{A}_r \mathbf{E}[\dots]^{-1} \mathbf{B}_r \\ &= -16\mathbf{C}_r [\bullet]^{-1} (\mathbf{A}_r \mathbf{E}[\bullet]^{-1} \mathbf{A}_r^2 \mathbf{E} - \mathbf{A}_r^2 \mathbf{E}[\dots]^{-1} \mathbf{A}_r \mathbf{E}) [\dots]^{-1} \mathbf{B}_r \\ &= -16\mathbf{C}_r \Gamma_{,1;2}, \mathbf{B}_r + 16\mathbf{C}_r \Gamma_{,2;1}, \mathbf{B}_r. \end{aligned} \quad (\text{C.11})$$

Equations (C.5), (C.7), and (C.9b) imply the identity

$$\begin{aligned} \mathbf{Q} \mathbf{Q}_x &= 16\mathbf{C}_r [\dots]^{-1} \mathbf{E}[\bullet]^{-1} \mathbf{B}_r \mathbf{C}_r (2\Gamma_{,1,1} - \Gamma_{,2}) \mathbf{B}_r \\ &= 32\mathbf{C}_r [\dots]^{-1} \mathbf{E}[\bullet]^{-1} (\mathbf{A}_r[\dots] - [\bullet] \mathbf{A}_r) [\dots]^{-1} \mathbf{A}_r \mathbf{E}[\dots]^{-1} \mathbf{A}_r \mathbf{E}[\dots]^{-1} \mathbf{B}_r \\ &\quad - 16\mathbf{C}_r [\dots]^{-1} \mathbf{E}[\bullet]^{-1} (\mathbf{A}_r[\dots] - [\bullet] \mathbf{A}_r) [\dots]^{-1} \mathbf{A}_r^2 \mathbf{E}[\dots]^{-1} \mathbf{B}_r \\ &= 32\mathbf{C}_r \Gamma_{,0;2,1}, \mathbf{B}_r - 32\mathbf{C}_r \Gamma_{,1,1,1}, \mathbf{B}_r - 16\mathbf{C}_r \Gamma_{,0;3}, \mathbf{B}_r + 16\mathbf{C}_r \Gamma_{,1,2}, \mathbf{B}_r. \end{aligned} \quad (\text{C.12})$$

Finally, using $\tilde{\sigma}_3[\dots] \tilde{\sigma} = [\bullet]$ as well as

$$\mathbf{E}_t = -4i\tilde{\sigma}_3 \mathbf{A}_r^2 \mathbf{E},$$

$$([\dots]^{-1})_t = -[\dots]^{-1} (-4i\tilde{\sigma}_3 \mathbf{A}_r^2 \mathbf{E}) [\dots]^{-1} = 4i\tilde{\sigma}_3 [\bullet]^{-1} \mathbf{A}_r^2 \mathbf{E} [\dots]^{-1} = 4i\tilde{\sigma}_3 \Gamma_{,2},$$

$$([\bullet]^{-1})_t = -[\bullet]^{-1} (-4i\tilde{\sigma}_3 \mathbf{A}_r^2 \mathbf{E}) [\bullet]^{-1} = 4i\tilde{\sigma}_3 [\dots]^{-1} \mathbf{A}_r^2 \mathbf{E} [\bullet]^{-1} = 4i\tilde{\sigma}_3 \Gamma_{,2},$$

we compute with the help of (7.3) and (C.3)

$$\begin{aligned} i\sigma_3 \mathbf{Q}_t &= -4i\sigma_3 \mathbf{C}_r (\Gamma_{,1})_t \mathbf{B}_r = -4i\mathbf{C}_r \tilde{\sigma}_3 (\Gamma_{,1})_t \mathbf{B}_r \\ &= -4i\mathbf{C}_r \tilde{\sigma}_3 ([\dots]^{-1} \mathbf{A}_r \mathbf{E} [\dots]^{-1})_t \mathbf{B}_r \\ &= 16\mathbf{C}_r \tilde{\sigma}_3 (\tilde{\sigma}_3 \Gamma_{,2}, \mathbf{A}_r \mathbf{E} [\dots]^{-1} - [\dots]^{-1} \tilde{\sigma}_3 \mathbf{A}_r^3 \mathbf{E} [\dots]^{-1} \\ &\quad + [\dots]^{-1} \mathbf{A}_r \mathbf{E} \tilde{\sigma}_3 \Gamma_{,2}) \mathbf{B}_r \\ &= 16\mathbf{C}_r \Gamma_{,2;1}, \mathbf{B}_r - 16\mathbf{C}_r \Gamma_{,3}, \mathbf{B}_r + 16\mathbf{C}_r \Gamma_{,1;2}, \mathbf{B}_r. \end{aligned} \quad (\text{C.13})$$

To prove (2.3) we now employ (C.13), (C.6), (C.10), (C.11), and (C.12) to compute

$$\begin{aligned}
& i\sigma_3 Q_t + Q_{xx} - Q^2 + \sigma_3 Q \sigma_3 Q - 2QQ_x \\
&= 16C_r \Gamma_{;2,1}, \mathbf{B}_r - 16C_r \Gamma_{;3}, \mathbf{B}_r + 16C_r \Gamma_{;1,2}, \mathbf{B}_r \\
&\quad - 96C_r \Gamma_{;1,1,1}, \mathbf{B}_r + 48C_r \Gamma_{;1,2}, \mathbf{B}_r + 48C_r \Gamma_{;2,1}, \mathbf{B}_r - 16C_r \Gamma_{;3}, \mathbf{B}_r \\
&\quad - 16C_r \Gamma_{;1,2}, \mathbf{B}_r - 16C_r \Gamma_{;2,1}, \mathbf{B}_r + 32C_r \Gamma_{;1,1,1}, \mathbf{B}_r \\
&\quad - 16C_r \Gamma_{;1,2}, \mathbf{B}_r + 16C_r \Gamma_{;2,1}, \mathbf{B}_r \\
&\quad - 64C_r \Gamma_{;0,2,1}, \mathbf{B}_r + 64C_r \Gamma_{;1,1,1}, \mathbf{B}_r + 32C_r \Gamma_{;0,3}, \mathbf{B}_r - 32C_r \Gamma_{;1,2}, \mathbf{B}_r \\
&= 32C_r \Gamma_{;2,1}, \mathbf{B}_r - 16C_r \Gamma_{;3}, \mathbf{B}_r + 32C_r \Gamma_{;2,1}, \mathbf{B}_r - 16C_r \Gamma_{;3}, \mathbf{B}_r \\
&\quad - 64C_r \Gamma_{;0,2,1}, \mathbf{B}_r + 32C_r \Gamma_{;0,3}, \mathbf{B}_r. \tag{C.14}
\end{aligned}$$

We now regroup the terms in the last member of (C.14) as follows:

$$\begin{aligned}
& i\sigma_3 Q_t + Q_{xx} - Q^2 + \sigma_3 Q \sigma_3 Q - 2QQ_x \\
&= 32C_r \Gamma_{;2,1}, \mathbf{B}_r + 32C_r \Gamma_{;2,1}, \mathbf{B}_r - 64C_r \Gamma_{;0,2,1}, \mathbf{B}_r \\
&\quad - 16C_r \Gamma_{;3}, \mathbf{B}_r - 16C_r \Gamma_{;3}, \mathbf{B}_r + 32C_r \Gamma_{;0,3}, \mathbf{B}_r. \tag{C.15}
\end{aligned}$$

Using (C.8) to derive the identities

$$\Gamma_{;2,1} + \Gamma_{;2,1} - 2\Gamma_{;0,2,1} = 0_{2p \times 2p}, \quad \Gamma_{;3} + \Gamma_{;3} - 2\Gamma_{;0,3} = 0_{2p \times 2p},$$

it appears that the right-hand side of (C.15) equals the zero matrix. Thus the expression for $\mathbf{Q}(x; t)$ in the statement of Theorem C.1 satisfies (2.3), as claimed. \blacksquare

By applying Theorem C.1 to the matrix triplet $(-\mathbf{A}_l, \mathbf{B}_l, -\mathbf{C}_l)$ we prove

Theorem C.2 *Let $(\mathbf{A}_l, \mathbf{B}_l, \mathbf{C}_l)$ be a matrix triplet satisfying (7.3) for which the Sylvester equation (7.5) has a unique solution \mathbf{P}_l . Then the matrix function $\mathbf{Q}(x; t)$ defined by*

$$\mathbf{Q}(x; t) = -4\mathbf{C}_l \mathbf{\Pi}_l(x; t)^{-1} \mathbf{A}_l e^{-2x\mathbf{A}_l} e^{-4it\tilde{\sigma}_3 \mathbf{A}_r^2} \mathbf{\Pi}_l(x; t)^{-1} \mathbf{B}_l$$

satisfies the nonlinear evolution equation (2.3) in those $(x, t) \in \mathbb{R}^2$ for which the matrix $\mathbf{\Pi}_l(x; t)$ is nonsingular.

References

- [1] M.J. Ablowitz, *Nonlinear Dispersive Waves. Asymptotic Analysis and Solitons*, Cambridge Texts in Applied Mathematics **47**, Cambridge University Press, Cambridge, 2011.
- [2] M.J. Ablowitz and P.A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.
- [3] M.J. Ablowitz, D.J. Kaup, A.C. Newell, and H. Segur, *The inverse scattering transform – Fourier analysis for nonlinear problems*, Stud. Appl. Math. **53**, 249–315 (1974).
- [4] M.J. Ablowitz, B. Prinari, and A.D. Trubatch, *Discrete and Continuous Nonlinear Schrödinger Systems*, Cambridge University Press, Cambridge, 2003.
- [5] M.J. Ablowitz and H. Segur, *Solitons and the Inverse Scattering Transform*, SIAM, Philadelphia, 1981.
- [6] H. Aden and B. Carl, *On realizations of solutions of the KdV equation by determinants on operator ideals*, J. Math. Phys. **37**, 1833–1857 (1996).
- [7] Z.S. Agranovich and V.A. Marchenko, *The Inverse Problem of Scattering Theory*, Gordon and Breach, New York, 1963.
- [8] T. Aktosun, F. Demontis, and C. van der Mee, *Exact solutions to the focusing nonlinear Schrödinger equation*, Inverse Problems **23**, 2171–2195 (2007).
- [9] T. Aktosun, F. Demontis, and C. van der Mee, *Exact solutions to the sine-Gordon equation*, J. Math. Phys. **51** (2010), 123521, 27 pp.
- [10] T. Aktosun, M. Klaus, and C. van der Mee, *Small-energy asymptotics of the scattering matrix for the self-adjoint matrix Schrödinger operator on the line*, J. Math. Phys. **42**, 4627–4652 (2001).
- [11] T. Aktosun and C. van der Mee, *Explicit solutions to the Korteweg-de Vries equation on the half-line*, Inverse Problems **22**, 2165–2174 (2006).

- [12] T. Aktosun and R. Weder, *Inverse scattering on the half line for the matrix Schrödinger equation*, Journal of Mathematical Physics, Analysis, Geometry **14**, 237–269 (2018).
- [13] T. Aktosun and R. Weder, *Direct and Inverse Scattering for the Matrix Schrödinger Equation*, Applied Mathematical Sciences **203**, Springer, New York, 2020.
- [14] G. Berkolaiko, *An elementary introduction to quantum graphs*. In: *Geometric and Computational Spectral Theory*, Contemporary Mathematics **700**, Amer. Math. Soc., Providence RI, 2017, pp. 41–72; also: arXiv:1603.07356v2 (2016).
- [15] G. Berkolaiko, R. Carlson, S.A. Fulling, and P. Kuchment (eds.), *Quantum Graphs and their Applications*, Contemporary Mathematics **415**, Amer. Math. Soc., Providence RI, 2006.
- [16] G. Berkolaiko and P. Kuchment, *Introduction to Quantum Graphs*, Mathematical Surveys and Monographs **186**, Amer. Math. Soc., Providence RI, 2013.
- [17] G. Berkolaiko and W. Liu, *Simplicity of eigenvalues and non-vanishing of eigenfunctions of a quantum graph*, J. Math. Anal. Appl. **445**, 803–818 (2017).
- [18] J. Boman and P. Kurasov, *Symmetries of quantum graphs and the inverse scattering problem*, Adv. Appl. Math. **35**, 58–70 (2005).
- [19] F. Calogero and A. Degasperis, *Spectral Transform and Solitons, I*, Studies in Mathematics and its Applications **13**, Elsevier, Amsterdam and New York, 1982.
- [20] K. Chadan and P. Sabatier, *Inverse Problems in Quantum Scattering Theory*, 2nd ed., Springer, New York, 1989.
- [21] M. Chen, M.A. Tsankov, J.M. Nash, and C.E. Patton, *Backward-volume-water microwave-envelope solitons in yttrium iron garnet films*, Phys. Rev. B **49**, 12773–12790 (1994).
- [22] P. Deift and E. Trubowitz, *Inverse scattering on the line*, Commun. Pure Appl. Math. **32**, 121–251 (1979).

- [23] F. Demontis, *Exact solutions to the modified Korteweg-de Vries equation*, *Theor. Math. Phys.* **168**, 886–897 (2011).
- [24] F. Demontis and C. van der Mee, *Marchenko equations and norming constants of the matrix Zakharov-Shabat system*, *Operators and Matrices* **2**, 79–113 (2008).
- [25] F. Demontis and C. van der Mee, *Explicit solutions of the cubic matrix nonlinear Schrödinger equation*, *Inverse Problems* **24**, 02520 (2008), 16 pp.
- [26] W. Eckhaus and A. van Harten, *The Inverse Scattering Transformation and the Theory of Solitons*, North-Holland, Amsterdam, 1981.
- [27] P. Exner, J.P. Keating, P. Kuchment, T. Sunada, and A. Teplyaev (eds.), *Analysis on Graphs and its Applications*, *Proceedings of Symposia in Pure Mathematics* **77**, Amer. Math. Soc., Providence RI, 2008.
- [28] L.D. Faddeev, *Properties of the S-matrix of the one-dimensional Schrödinger equation*, *Amer. Math. Soc. Transl. Series* **2**, **65**, 139–166 (1964).
- [29] L.D. Faddeev and L.A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons*, *Classics in Mathematics*, Springer, New York, 1987.
- [30] I.M. Gelfand, D.A. Raikov, and G.E. Shilov, *Commutative Normed Rings*, Chelsea Publ., New York, 1964.
- [31] N.I. Gerasimenko, *The inverse scattering problem on a noncompact graph*, *Theor. Math. Phys.* **75**, 460–470 (1988).
- [32] I. Gerasimenko and B.S. Pavlov, *A scattering problem on a noncompact graphs*, *Theor. Math. Phys.* **74**, 230–240 (1988).
- [33] B. Gutkin and U. Smilansky, *Can one hear the shape of a graph?*, *J. Phys. A: Math. Gen.* **34**, 6061–6068 (2001).
- [34] M.S. Harmer, *Inverse scattering for the matrix Schrödinger operator and Schrödinger operator on graphs with general self-adjoint boundary conditions*, *ANZIAM J.* **44**, 161–168 (2002).

- [35] M.S. Harmer, *The matrix Schrödinger operator and Schrödinger operator on graphs*, Ph.D. thesis, University of Auckland, New Zealand, 2004.
- [36] M.S. Harmer, *Inverse scattering on matrices with boundary conditions*, J. Phys. A: Math. Gen. **38**, 4875–4885 (2005).
- [37] A. Hasegawa, *Optical Solitons in Fibers*, Springer Series in Photonics **9**, Springer, New York, 2002.
- [38] A. Hasegawa and F. Tappert, *Transmission of stationary nonlinear optical pulses in dispersive dielectric fibers. I. Anomalous dispersion*, Appl. Phys. Lett. **23**(3), 142–144 (1973); *II. Normal dispersion*, Appl. Phys. Lett. **23**(4), 171–172 (1973).
- [39] R.A. Horn and C.J. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1994.
- [40] P.G. Kevrekidis, D.J. Frantzeskakis, R. Carretero-González, *Emergent Non-linear Phenomena in Bose-Einstein Condensates*, Springer, Berlin, 2008.
- [41] M. Klaus, *Low-energy behaviour of the scattering matrix for the Schrödinger equation on the line*, Inverse Problems **4**, 505–512 (1988).
- [42] V. Kostykin and R. Schrader, *Kirchhoff's rule for quantum wires*, J. Phys. A: Math. Gen. **32**, 595–630 (1999).
- [43] V. Kostykin and R. Schrader, *Kirchhoff's rule for quantum wires. II. The inverse problem with possible applications to quantum computers*, Fortschr. Phys. **48**, 703–716 (2000).
- [44] P. Kuchment, *Quantum graphs. I. Some basic structures*, Waves Random Media **14**, S107–S128 (2004).
- [45] P. Kuchment, *Quantum graphs. II. Some spectral properties of quantum and combinatorial graphs*, J. Phys. A: Math. Gen. **38**, 4887–4900 (2005).
- [46] P. Kurasov and M. Nowaczyk, *Geometric properties of quantum graphs and vertex scattering matrices*, Opusc. Math. **30**, 295–309 (2010).

- [47] P. Kurasov and M. Nowaczyk, *Inverse spectral problem for quantum graphs*, J. Phys. A: Math. Gen. **38**, 4901–4915 (2005).
- [48] P. Kurasov and F. Stenberg, *On the inverse scattering problem on branching graphs*, J. Phys. A: Math. Gen. **35**, 101–121 (2002).
- [49] P. Lax, *Integrals of nonlinear equations of evolution and solitary waves*, Commun. Pure Appl. Math. **21**, 467–490 (1968).
- [50] C.J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, 2nd ed., Cambridge University Press, Cambridge, 2008.
- [51] L.P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superconductivity*, Oxford University Press, Oxford, 2016.
- [52] C. Schiebold, *An operator theoretic approach to the Toda lattice equation*, Physica D **122**, 37–61 (1998).
- [53] C. Schiebold, *Solutions of the sine-Gordon equation coming in clusters*, Revista Matemática Complutense **15**, 265–325 (2002).
- [54] J.K. Shaw, *Mathematical Principles of Optical Fiber Communications*, CBMS-NSF Regional Conference Series in Applied Mathematics **76**, SIAM, Philadelphia, 2004.
- [55] C. van der Mee, *Nonlinear Evolution Models of Integrable Type*, SIMAI e-Lecture Notes **11**, SIMAI, Torino, 2013.
- [56] M. Wadati and T. Kamijo, *On the extension of inverse scattering method*, Prog. Theor. Phys. **52**, 397–414 (1974).
- [57] V.E. Zakharov, *Hamilton formalism for hydrodynamic plasma models*, Sov. Phys. JETP **33**, 927–932 (1971).
- [58] V.E. Zakharov and A.F. Popkov, *Contribution to the nonlinear theory of magnetostatic spin waves*, Sov. Phys. JETP **57**, 350–355 (1983).
- [59] V.E. Zakharov and A.B. Shabat, *Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media*, Sov. Phys. JETP **34**, 62–69 (1972).