

SHEARING IN SOME SIMPLE RANK ONE THEORIES

M. MALLIARIS AND S. SHELAH

ABSTRACT. Dividing asks about inconsistency along indiscernible sequences. In order to study the finer structure of simple theories without much dividing, the authors recently introduced *shearing*, which essentially asks about inconsistency along generalized indiscernible sequences. Here we characterize the shearing of the random graph. We then use shearing to distinguish between the random graph and the theories $T_{n,k}$, the higher-order analogues of the triangle-free random graph. It follows that shearing is distinct from dividing in simple unstable theories, and distinguishes meaningfully between classes of simple unstable rank one theories. The paper begins with an overview of shearing, and includes open questions.

This paper is dedicated to Moshe Jarden, in honor of his many contributions to field arithmetic.

One of the central points of contact between fields and model theory is the definition of forking/dividing, developed in the second author's book [19], which significantly abstracts the notions of algebraic independence in algebraically closed fields, or linear independence in vector spaces. Informally, dividing asks about inconsistency along indiscernible sequences. This definition has substantial explanatory power within stable theories, but outside of stable theories, it appears that new ideas are also needed. Thus, in order to study the finer structure of simple theories without much dividing, the authors recently introduced *shearing*, which essentially asks about inconsistency along generalized indiscernible sequences. The aim of this paper is to further develop this very interesting definition.

The paper [11] proved a first separation theorem showing shearing is a priori useful for detecting differences in complexity in simple unstable theories. The proof that this notion is strictly weaker than dividing, and that it can be found in the random graph and indeed in any theory with the independence property, was deferred to the present manuscript. Below, we carry out the characterization of shearing in the random graph announced there. We then show that shearing can distinguish between the random graph and the theories $T_{n,k}$, the higher-order analogues of the triangle-free random graph studied by Hrushovski. Perhaps this may open the door for a more careful structural analysis of simple rank one theories (whose structure is not visible to dividing) via shearing. Along the way we review what is known and record many natural questions.

1. PRELIMINARIES

In this section we recall the central definition of *shearing*, Definition 1.2. The reader may prefer to look ahead, or to read on for the motivated development.

Thanks: Research partially supported by NSF CAREER award 1553653, by an NSF-BSF award (NSF 2051825, BSF 3013005232), and ISF grant 1838/19. Paper 1221 in Shelah's list.

The basic idea will be that whereas dividing corresponds to inconsistency along an indiscernible sequence, shearing corresponds to inconsistency along a generalized indiscernible sequence. One initial reason it might be hoped this would give some power is that generalized indiscernible sequences arise from the skeletons of GEM-models.¹ So from consistency or inconsistency along such sequences in a given GEM-model one might hope to produce larger GEM-models in which a given type was realized, or stayed omitted. Going further, one might hope to show a difference in complexity between theories in this way, say, by showing that if T_2 has recurrent shearing for a certain kind of generalized indiscernible sequence and T_1 does not, then it would be possible to build a model of a theory interpreting both of them whose reduct to T_1 is quite saturated and whose reduct to T_2 is (say) not even \aleph_1 -saturated. This was done in [11], using GEM-models, and working through those proofs allowed us to arrive at the present definition of shearing. Once identified, the definition makes sense in any context, not only that of GEM-models.

We shall use three ideas from the model theory of Ehrenfeucht-Mostowski models: that of an index model class \mathcal{K} , that of a context \mathbf{c} , and that of a generalized indiscernible sequence for such a class, which we call \mathcal{K} -indiscernible. In various guises, these have long histories in model theory (a partial list might include [1], [16], [19], [17], [3], [10], [11]). To balance the demands of keeping the paper short but also reasonably self-contained, we point the reader to where these are clearly written down, and give here an English summary.

Index model class: See [11] Definitions 2.3, p. 4 and 2.9, p. 5. Briefly, \mathcal{K} is a class of linearly ordered models, closed under isomorphism but not necessarily an elementary class, in a signature expanding $\{<\}$. We require that \mathcal{K} is universal, and that for every $I \in \mathcal{K}$ there is some \aleph_0 -saturated $J \in \mathcal{K}$ extending it. (Since the class is not necessarily elementary, “ \aleph_0 -saturated” always abbreviates “ \aleph_0 -universal and \aleph_0 -homogeneous”.) Finally, the class has to be Ramsey, in the usual sense of GEM-models (or equivalently, by a theorem of Scow, in the sense of Nešetřil [14] and of Kechris-Pestov-Todorčević [7]; see [17] Theorem 4.31). Simple examples include: the class of linear orders, or the class of linear orders partitioned by countably many unary predicates $\{P_n : n < \omega\}$, where note that requiring the countably many predicates to partition the domain (every element has a color; some colors may be empty) makes it not an elementary class.

Context: See [11] Definition 2.12. Briefly, a context $\mathbf{c} = (I, \mathcal{K})$ is a choice of index model class \mathcal{K} along with a choice of $I \in \mathcal{K}$ which is not obviously trivial. For example, taking \mathcal{K} to be the class of linear orders and asking for universality means \mathcal{K} contains lots of finite linear orders, but it is not so useful to choose one of them when asking about uniform inconsistency. The various conditions essentially rule out analogues of this: I is closed under functions, if any; I is not generated by any of its finite subsets; closure in the sense of the function symbols in the language corresponds reasonably to algebraicity (things which only appear finitely many times in saturated extensions of I do so for this reason).

A *countable context* just means that, in addition, I is countably infinite.

\mathcal{K} -indiscernible: See [11] Section 3, or Definitions 3.2-3.5. In a usual indiscernible sequence in a model M , indexed by a linear order $(I, <)$, if the finite sequences \bar{t}, \bar{s}

¹Generalized Ehrenfeucht-Mostowski models – informally, like Ehrenfeucht-Mostowski models except that we allow the index models to be reasonable expansions of linear orders, varying within an index model class \mathcal{K} ; for a recent exposition, see the early sections of [10], [11].

from I have the same order-type [i.e., the same quantifier-free type in the language of order] then the corresponding tuples of elements they index have the same type in M . In general, if I is no longer just a linear order but belongs to some \mathcal{K} , then we may ask that if \bar{t}, \bar{s} have the same quantifier-free type in I then the corresponding tuples of elements they index have the same type in M .

The subtlety is that in the full definition, one more level of generality is implicit, as follows. To specify a \mathcal{K} -indiscernible sequence, we choose a map f from finite sequences from I to finite sequences from M such that if \bar{t}, \bar{s} have the same length and quantifier-free type in I then their images have the same length and type in M . Write $\bar{a}_{\bar{t}}$ for $f(\bar{t})$. Observe that requirements on coherence of these maps between arities are conspicuously absent. For instance, $f(t_1 t_2)$ isn't required to be the concatenation of $f(t_1)$ with $f(t_2)$. Indeed, here is a very useful construction which fits the definition. Choose a particular quantifier-free k -type τ of some sequence \bar{t} of elements of I . Choose some \aleph_0 -saturated $J \in \mathcal{K}$ with $I \subseteq J$. Consider the set $Q_\tau^J = \{\bar{s} : \text{tp}_{\text{qf}}(\bar{s}, \emptyset, J) = \text{tp}_{\text{qf}}(\bar{t}, \emptyset, I)\}$. Consider a map f taking each \bar{s} in Q_τ^J to some $\bar{a}_{\bar{s}}$ in M , so that the types of any two such sequences in M are the same. Extend f to $\omega^> J$ by sending anything outside Q_τ^J to the empty set (or to a single fixed element). Then the image of f is a very reasonable \mathcal{K} -indiscernible sequence.

Notation 1.1. *When $I_0 \subseteq J \in \mathcal{K}$ is a set, writing $J[I_0]$ means J expanded by constants for the elements of I_0 , and likewise for $J[\bar{s}]$ when $\bar{s} \subseteq J$ is a sequence.*

We now arrive at our central definition.

Definition 1.2 (Shearing, [11] Definition 5.2). *Suppose we are given a context \mathbf{c} , a theory T , $M \models T$, $A \subseteq M$, and a formula $\varphi(\bar{x}, \bar{c})$ of the language of T with $\bar{c} \in \omega^> M$. We say that*

the formula $\varphi(\bar{x}, \bar{c})$ shears over A in M for (I_0, I_1, \mathbf{c})

when there exist a model N , a sequence \mathbf{b} in N , enumerations \bar{s}_0 of I_0 and \bar{t} of I_1 , and an \aleph_0 -saturated $J \supseteq I$ such that:

- (1) $I_0 \subseteq I_1$ are finite subsets of I
- (2) $M \preceq N$
- (3) $\mathbf{b} = \langle \bar{b}_{\bar{s}} : \bar{s} \in \omega^>(J[I_0]) \rangle$ is \mathcal{K} -indiscernible in N over A
- (4) $\bar{c} = \bar{b}_{\bar{t}}$, and
- (5) the set of formulas

$$\{\varphi(\bar{x}, \bar{b}_{\bar{t}'}') : \bar{t}' \in {}^{\text{lg}(\bar{t})}(J), \text{tp}_{\text{qf}}(\bar{t}', \bar{s}_0, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}_0, I)\}$$

is contradictory.

Let us look at this definition a little more closely. From the second paragraph of the section, the reader may anticipate that it will be useful to be able to iterate any inconsistency we may find. Perhaps we start by observing that in some larger $J \supseteq I$, the tuples satisfying the same quantifier-free type as our given \bar{t} can form indices for a generalized indiscernible sequence along which some φ is inconsistent. If we fix one of these instances, say \bar{s}_0 , can this happen again over \bar{s}_0 ? That is, can we find some \bar{t}_1 from I so that in some larger $J \supseteq I$, the tuples satisfying the quantifier-free type of \bar{t}_1 over \bar{s}_0 form indices for a generalized indiscernible sequence along which our φ is again inconsistent? Does this stop after finitely many steps? This motivates the definition of \mathbf{c} -*superstability*, which was key to [11], and explains the appearance of a finite “ I_0 ” in Definition 1.2.

We state here the local definition of “ \mathbf{c} -superstable” just for relational languages, which suffices for the present paper. Note that countability of the context \mathbf{c} is used in an essential way, as I is written as the union of an increasing chain of finite sets.

Definition 1.3 ([11] Definition 6.2, local version). *Let \mathbf{c} be a countable context and Δ a set of formulas. We say (T, Δ) is \mathbf{c} -unsuperstable for \mathbf{c} when there are:*

- (a) *an increasing sequence of nonempty finite sets $\langle I_n : n < \omega \rangle$ with $I_m \subseteq I_n \subseteq I$ for $m < n < \omega$ and $\bigcup_n I_n = I$, which are given along with a choice of enumeration \bar{s}_n for each I_n where $\bar{s}_n \trianglelefteq \bar{s}_{n+1}$ for each n*
- (b) *an increasing sequence of nonempty, possibly infinite, sets $B_n \subseteq B_{n+1} \subseteq \mathfrak{C}_T$ in the monster model for T , with $B := \bigcup_n B_n$*
- (c) *and a partial type p over B ,*

such that for each n , for some formula $\varphi(\bar{x}, \bar{c})$ from $p \upharpoonright B_{n+1}$ where $\varphi(\bar{x}, \bar{y}) \in \Delta$, we have that

$$\varphi(\bar{x}, \bar{c}) \quad (I_n, I_{n+1})\text{-shears over } B_n.$$

Definition 1.4. *Continuing Definition 1.3, for a countable context \mathbf{c} ,*

- (1) *we may write (T, φ) to mean $(T, \{\varphi\})$.*
- (2) *We say T is \mathbf{c} -superstable if (T, Δ) is \mathbf{c} -superstable where Δ is the set of all formulas in the language.*
- (3) *We say T is \mathbf{c} -stable if (T, φ) is \mathbf{c} -superstable for every φ in the language.*

Discussion 1.5. These investigations into the fine structure of forking highlight a longstanding terminological point, which hopefully should not cause confusion if explicitly pointed out. Namely, in the classical case, both “superstable” and “supersimple” correspond naturally to “ $\kappa(T) = \aleph_0$.” In Definition 1.3, we would also have been justified in using “supersimple”. It seems to us that stable is the right one to use for various reasons, so hopefully this does not confuse the reader in a sentence like “a theory is simple if there is *some* context \mathbf{c} for which it is \mathbf{c} -stable.”

For context and easy quotation, the next local summary theorem includes some already known facts, and some new results proved below. We assume our theories T are all complete.

Theorem 1.6 (Local shearing). *For now all contexts are countable.*

- (1) *Dividing implies shearing.*
- (2) *Shearing does not imply dividing, i.e., it is strictly weaker than dividing.*
- (3) *If φ is a stable formula in the theory T , then (T, φ) is \mathbf{c} -superstable for every countable context \mathbf{c} .*
- (4) *If φ is an unstable formula in the theory T , then (T, φ) is \mathbf{c} -unsuperstable for some countable context \mathbf{c} . (So the previous item is a characterization of stable formulas.)*
- (5) *In the previous item, we can take \mathbf{c} to be a countable context from the class \mathcal{K} of linear orders, that is, (T, φ) is unstable if and only if it is \mathbf{c} -unsuperstable for a countable context chosen from the class of infinite linear orders.*
- (6) *If (T, φ) is \mathbf{c} -superstable for some countable context \mathbf{c} , then φ is simple, i.e. it does not have the tree property in the theory T . (So the natural focus of shearing is in some sense within simplicity.)*

- (7) More precisely, in the previous item, the following are equivalent: (a) φ is simple in T and² $\kappa_{\text{loc}}(T, \varphi) = \aleph_0$, and (b) there is some countable context \mathbf{c} for which (T, φ) is \mathbf{c} -superstable.

Proof. (1) See [11] Claim 5.8.

(2) It was stated in [11] (see the end of §5) that this would be proved in the present paper, which it is indeed: we shall show that there is nontrivial shearing for the random graph and also for the theories $T_{n,k}$ for $n > k \geq 2$, theories which have dividing only for equality.

(3)-(4) See Conclusion 3.10 below.

(5) Conclusion 3.10 and Example 3.8 below. But this should be read with caution, see 1.7.

(6)-(7) See [11] Lemma 9.5 and Theorem 9.6. \square

Discussion 1.7. Caution: just because dividing implies shearing, one should not jump to conclusions about linear orders. As noted in 3.10(5), the random graph is \mathbf{c} -unsuperstable for contexts coming from \mathcal{K} the class of linear orders, see 3.8 below. (This uses an indexing by pairs.) However the random graph is \mathbf{c} -superstable for some contexts coming from expansions of linear orders by predicates, see 3.9.

Discussion 1.8. In 3.10(7), we might want a characterization of “ (T, φ) is simple” and wonder why the apparently extra assumption “ $\kappa_{\text{loc}}(T, \varphi) = \aleph_0$ ” appears. The reason is that in 3.10, we are using countable contexts while making no assumptions on cardinality of the theory, so there is an extra point about the length of the shearing chain. To fully capture simplicity, it would be natural to consider larger contexts to deal with the analogue of arbitrarily large $\kappa(T)$. If the reader is inspired by this remark, [11] §10 is a beginning.

2. BACKGROUND

In later sections we will use shearing to distinguish between various theories which have no dividing in the usual sense (other than that coming from equality). In this expository section we review why this is so. For background on simple theories, see the survey [2].

Definition 2.1. For each $n \geq 2$, let $T_{n,1}$ be the theory of the generic K_{n+1} -free graph. For $n > k \geq 2$, let $T_{n,k}$ be the theory of the generic $(n+1)$ -free $(k+1)$ -hypergraph.

So, in this notation, $T_{3,2}$ is the theory of the generic tetrahedron-free three-hypergraph. For $k \geq 2$, i.e. for the case where the edge is really a hyperedge, these theories are simple unstable with trivial forking, as shown by Hrushovski [4]. For context, we begin by reviewing why the graph versions of these theories (triangle-free graph and its relatives) *do* have a lot of dividing.

Fact 2.2. $T_{n,1}$ is not simple.

Proof sketch. First consider the triangle-free random graph, in our notation $T_{2,1}$. Let $\varphi(x, y, z) = R(x, y) \wedge R(x, z)$. Consider in some sufficiently saturated model of $T_{2,1}$ an infinite sequence of pairs $\mathbf{a} = \langle \bar{a}_i : i < \omega \rangle$ where each $\bar{a}_i = a_i^0 a_i^1$, and such

²Recall that $\kappa_{\text{loc}}(T, \varphi) = \aleph_0$ means that every φ -type does not fork over a finite set.

that $R(a_i^0, a_j^1)$ for all $i \neq j$ but for all i, j $\neg R(a_i^0, a_j^0)$ and $\neg R(a_i^1, a_j^1)$. Informally, the sequence \mathbf{a} is a bipartite graph which is the complement of a matching. Then the sequence

$$\{\varphi(x, a_i^0, a_i^1) : i < \omega\}$$

has the property that each formula is individually consistent, but the sequence is 2-inconsistent, since if $i \neq j$ then any element satisfying $\{R(x, a_i^0), R(x, a_j^1)\}$ would form a triangle. Moreover, it is easy to see that for any \bar{a}_i in \mathbf{a} , we can construct a sequence isomorphic to \mathbf{a} which is indiscernible over \bar{a}_i , and continuing in this way we may construct the tree property for φ , showing that φ is not simple.

It is easy to extend this idea to $n > 2$ using $\varphi = R(x, y_0) \wedge \cdots \wedge R(x, y_{n-1})$, replacing \mathbf{a} by a sequence of n -tuples $\bar{a}_i = a_i^0 \cdots a_i^{n-1}$ and specifying that $R(a_i^s, a_j^t)$ holds there if and only if $(s \neq t) \wedge (i \neq j)$. \square

When we consider hypergraphs instead of graphs the situation is quite different.

Fact 2.3 (Hrushovski c. 2002, see [4]). *For $n > k \geq 2$, $T_{n,k}$ is simple unstable with only trivial dividing.*

Proof sketch. We sketch the proof for $T = T_{3,2}$, the tetrahedron-free 3-hypergraph, since this extends naturally to larger arities but at a notational cost. Suppose there were some formula $\varphi(x, a^0, \dots, a^{\ell-1})$, some $m > 1$, and some indiscernible sequence $\mathbf{a} = \langle \bar{a}_i : i < \omega \rangle$ such that $\bar{a}_0 = \langle a_0^0, \dots, a_0^{\ell-1} \rangle = \langle a^0, \dots, a^{\ell-1} \rangle$ and

$$(1) \quad \{\varphi(x, a_i^0, \dots, a_i^{\ell-1}) : i < \omega\}$$

is m -inconsistent. Let's consider \mathbf{a} as being arranged so that each \bar{a}_i is a column, and each $\langle a_i^s : i < \omega \rangle$ is a row. We don't assume anything about how the edges hold on \mathbf{a} , but there are some constraints, e.g. because \mathbf{a} is indiscernible and exists in a model of T , the edge $R(x, y, z)$ cannot hold on any three distinct elements in any row of \mathbf{a} , otherwise (by indiscernibility) the row would contain a tetrahedron. By quantifier elimination, and without loss of generality ignoring the trivial forking coming from equality, we may assume φ is a boolean combination of instances of $R(x; y, z)$.

Now if (1) is inconsistent, there must be some tetrahedron which appears. In particular, there must be elements b, c, d which occur in \mathbf{a} with the following three properties: first, the quantifier-free type of \mathbf{a} implies $R(b, c, d)$; second, $\{\varphi(x, a_i^0, \dots, a_i^{\ell-1}) : i < \omega\} \vdash \{R(x, b, c), R(x, b, d), R(x, c, d)\}$; and third, because each individual formula in (1) is consistent, b, c, d are not all in one column of \mathbf{a} . But (1) can only imply instances of formulas all of whose parameters occur in the same column – look at the definition of (1) and notice that none of its formulas include parameters from distinct columns. [This is the crucial difference in having an edge of higher arity than 2.] So this contradiction can never arise, and this completes the proof. \square

3. ANALYSIS OF THE RANDOM GRAPH

Convention 3.1. *Reminder: all contexts are countable.*

The theory of the random graph, $T_{\mathbf{rg}}$, is central to the picture. In this section we analyze it carefully and give a complete characterization of countable contexts \mathbf{c} for which $T_{\mathbf{rg}}$ is \mathbf{c} -superstable, Theorem 3.7. This result was announced in [11],

along with Definition 3.12 and the motivating example §3.1 preceding it; although it isn't strictly necessary, we repeat the example here for clarity.

The direct route to Theorem 3.7 is via Definition 3.2, Claim 3.5 and Claim 3.6. There are two discussions and a claim which are mainly explanatory: Discussion 3.1, Discussion 3.2 and Claim 3.5.

3.1. A motivating example. Fix for awhile $\mathbf{c} = (I, \mathcal{K}) = (I_{\mathbf{c}}, \mathcal{K}_{\mathbf{c}})$ some countable context and we shall investigate how \mathbf{c} -shearing may arise for $T_{\mathbf{rg}}$ inside a GEM-model and try to find a characteristic property of a countable context which explains how such shearing occurs. Consider $M = \text{GEM}(I, \Phi)$, where $\Phi \in \Upsilon[T_{\mathbf{rg}}]$ thus $(M, R^M) \models T_{\mathbf{rg}}$, and let $p \in \mathbf{S}(M \upharpoonright \tau(T_{\mathbf{rg}}))$ be a nonalgebraic type. Fix J such that $I \subseteq J \in \mathcal{K}$ and J is \aleph_0 -saturated. Let $N = \text{GEM}(J, \Phi)$. Since all the contexts we will consider are well behaved (see [11] Convention 2.7, or take the following as a provisional definition of good behavior) we have $M \preceq N$, so we will identify the sequence $\langle \bar{a}_t : t \in I \rangle$ which generates M with a subsequence of $\langle \bar{a}_t : t \in J \rangle$.

By quantifier elimination, we may suppose p is equivalent to $\{R(x, b_\alpha)^{i_\alpha} \wedge x \neq b_\alpha : \alpha < \kappa\}$ for some infinite κ , where each $i_\alpha \in \{0, 1\}$. As M is generated by $\{\bar{a}_t : t \in I\}$, each b_α may be written as $\sigma_\alpha^M(\bar{a}_{\bar{t}_\alpha})$ for some $\tau(\Phi)$ -term σ_α and some $\bar{t}_\alpha \in \text{inc}(I)$. This representation may not be unique; there is no harm in choosing our enumeration to include all representations. So without loss of generality, for some $\kappa = \kappa + |\tau(\Phi)|$,

$$(2) \quad p(x) \equiv \{R(x, \sigma_\alpha^M(\bar{a}_{\bar{t}_\alpha}))^{i_\alpha} \wedge x \neq \sigma_\alpha^M(\bar{a}_{\bar{t}_\alpha}) : \alpha < \kappa\}.$$

where for each $\alpha < \kappa$ and $\bar{t} \in {}^\omega I$ if $\sigma_\alpha^M(\bar{a}_{\bar{t}}) = \sigma_\alpha^M(\bar{a}_{\bar{t}_\alpha})$, then

$$(3) \quad \text{for some } \beta < \kappa, \sigma_\beta = \sigma \text{ and } \bar{t}_\beta = \bar{t}.$$

This may increase the length of the enumeration, but will not change the size of the type in $\tau(T_{\mathbf{rg}})$. [From the point of view of $M \upharpoonright \tau(T_{\mathbf{rg}})$, we may appear to list some (say) $R(x, b_\alpha)^{i_\alpha} \wedge x \neq b_\alpha$ many times, because we have listed an instance for each way of writing b_α in M in terms of the skeleton. From the point of view of our enumeration, which has access to $\tau(\Phi)$, for each b_α there are potentially $|I| + |\tau(\Phi)|$ such representations.]

Recalling our fixed \aleph_0 -saturated J extending I and its associated $N = \text{GEM}(J, \Phi)$, we ask about potential \mathbf{c} -dividing. Working in N , consider the set of formulas

$$q(x) = q_{I_0}(x) = \{R(x, \sigma_\alpha^N(\bar{a}_{\bar{u}}))^{i_\alpha} \wedge x \neq \sigma_\alpha^N(\bar{a}_{\bar{u}}) : \alpha < \kappa, \\ \text{tp}_{\text{qf}}(\bar{u}, I_0, J) = \text{tp}_{\text{qf}}(\bar{t}_\alpha, I_0, I)\}.$$

To show $q(x)$ is consistent, it would suffice to check that whenever

$$(4) \quad R(x, \sigma_\alpha^N(\bar{a}_{\bar{v}}))^{i_\alpha} \in q \text{ and } R(x, \sigma_\beta^N(\bar{a}_{\bar{w}}))^{i_\beta} \in q$$

we have that if $\sigma_\alpha^N(\bar{a}_{\bar{v}}) = \sigma_\beta^N(\bar{a}_{\bar{w}})$ then $i_\alpha = i_\beta$. Suppose this fails, i.e., for some suitable $\alpha, \beta, \bar{v}_*, \bar{w}_*$ which we fix for awhile, q contains the contradictory formulas

$$(5) \quad R(x, \sigma_\alpha^N(\bar{a}_{\bar{v}_*})) \text{ and } \neg R(x, \sigma_\beta^N(\bar{a}_{\bar{w}_*})).$$

In other words,

$$(6) \quad \sigma_\alpha^N(\bar{a}_{\bar{v}_*}) = \sigma_\beta^N(\bar{a}_{\bar{w}_*}) = b \text{ but } i_\alpha \neq i_\beta \text{ (here w.l.o.g. } i_\alpha = 1, i_\beta = 0).$$

Informally, what has happened is that in N , there is a ‘‘positive line’’

$$\text{Pos} = \{\sigma_\alpha^N(\bar{a}_{\bar{v}}) : \text{tp}_{\text{qf}}(\bar{v}, I_0, J) = \text{tp}_{\text{qf}}(\bar{t}_\alpha, I_0, I)\} \subseteq \text{Dom}(N)$$

and a “negative line”

$$\text{Neg} = \{\sigma_\beta^N(\bar{a}_{\bar{w}}) : \text{tp}_{\text{qf}}(\bar{w}, I_0, J) = \text{tp}_{\text{qf}}(\bar{t}_\beta, I_0, I)\} \subseteq \text{Dom}(N)$$

and the problem is that

$$(7) \quad \text{Pos} \cap \text{Neg} \neq \emptyset, \text{ witnessed by } b = \sigma_\alpha^N(\bar{a}_{\bar{v}_*}) = \sigma_\beta^N(\bar{a}_{\bar{w}_*}).$$

However, it is also important to notice that both “lines” have “points from I ”, and that these are not the point(s) of intersection:³

$$(8) \quad \sigma_\alpha^N(\bar{a}_{\bar{t}_\alpha}) \neq \sigma_\beta^N(\bar{a}_{\bar{t}_\beta})$$

else our original p would be inconsistent. (So both Pos and Neg have size ≥ 2 .)

Definition 3.2 abstracts the key property of \mathbf{c} behind this picture. Towards this, observe that writing $\mathfrak{r}_\alpha = \text{tp}_{\text{qf}}(\bar{t}_\alpha, I_0, I)$, we have that “ $\sigma_\alpha^N(\bar{a}_{\bar{v}}) = \sigma_\alpha^N(\bar{a}_{\bar{u}})$ ” is an equivalence relation on $Q_{\mathfrak{r}_\alpha}^J$ (asserting that \bar{v}, \bar{u} are equivalent), and similarly for σ_β^N and \mathfrak{r}_β . Note that the fact that \bar{t}_α and \bar{t}_β may have different types is not important (as will be explained). The third formula, F , will give the analogue of equation (6) “points of intersection.” After giving the definition, we will work towards proving the characterization in Theorem 3.7.⁴

Definition 3.2. *The context $\mathbf{c} = (I, \mathcal{K})$ has property \odot when:*

For every finite $I_0 \subseteq I$ there is a finite I_1 with $I_0 \subseteq I_1 \subseteq I$, letting \bar{s}, \bar{t} be I_0, I_1 respectively, such that for any \aleph_0 -saturated $J \supseteq I$ there exist quantifier-free (possibly infinitary) formulas of $\tau(\mathcal{K})$ called $F(\bar{x}_1, \bar{x}_2; \bar{y})$, $E_1(\bar{x}_1, \bar{x}_2; \bar{y})$, $E_2(\bar{x}_1, \bar{x}_2; \bar{y})$, such that $\ell(\bar{x}_1) = \ell(\bar{x}_2) = \text{lg}(\bar{t})$, $\text{lg}(\bar{y}) = \text{lg}(\bar{s})$, and:

- (i) *for $\mathbf{i} = 1, 2$ $E_{\mathbf{i}}(\bar{x}_1, \bar{x}_2; \bar{s})$ defines an equivalence relation on $Y_{\bar{s}} = \{\bar{t}' \in \text{lg}(\bar{t})J : \text{tp}_{\text{qf}}(\bar{t}', \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, J)\}$.*
- (ii) *$F(\bar{x}_1, \bar{x}_2; \bar{y})$ defines a nonempty one to one partial function from $Y_{\bar{s}}/E_1(-, -; \bar{s})$ to $Y_{\bar{s}}/E_2(-, -; \bar{s})$, and*
- (iii) *F has no fixed points, in other words for no $\bar{t} \in Y_{\bar{s}}$ is it the case that $F(\bar{t}, \bar{t}; \bar{s})$.*

We now work towards a characterization, Theorem 3.7. Note that the intent is $\odot_{\mathbf{c}}$ means \mathbf{c} is too expressive, $\neg\odot_{\mathbf{c}}$ means \mathbf{c} is reasonable. For a high-level view of this property, see §4.

3.2. Discussion. As a warm-up to Claim 3.5, let us verify that indeed this property has captured \mathbf{c} -shearing, by reversing the abstraction above. Suppose that \mathbf{c} has property \odot . Fix a finite subset I_0 of $I_{\mathbf{c}}$ and $\Psi \in \Upsilon[T_{\mathbf{rg}}]$. Let $I_1, \bar{s}, \bar{t}, E_1, E_2, F$ witness property \odot for I_0 . Fix an \aleph_0 -saturated $J \in \mathcal{K}_{\mathbf{c}}$ with $I \subseteq J$, and we will find a formula of $\tau(T_{\mathbf{rg}})$ which (I_0, I_1) -shears for \mathbf{c} as follows. Let $N = \text{GEM}(J, \Psi)$ have skeleton $\langle \bar{a}_t : t \in J \rangle$. Without loss of generality, $\|N\| \geq |J|$. Let σ_1, σ_2 be two new $(\ell(\bar{a}_t) + \ell(\bar{a}_{\bar{s}}))$ -place function symbols not already in $\tau(\Psi)$. Let $\mathfrak{r} = \text{tp}_{\text{qf}}(\bar{t}, I_0, I)$. Let $Y_{\bar{s}} = Q_{\mathfrak{r}}^J$ be the set of realizations of \mathfrak{r} in J .

Expand N to N^+ by interpreting σ_1, σ_2 as follows.

³If p is a complete type in M , then we will have the stronger statement that “the restrictions of Pos and Neg to I have no intersection,” i.e. $\{\sigma_\alpha^N(\bar{a}_{\bar{v}}) : \text{tp}_{\text{qf}}(\bar{v}, I_0, I) = \text{tp}_{\text{qf}}(\bar{t}_\alpha, I_0, I)\} \cap \{\sigma_\beta^N(\bar{a}_{\bar{w}}) : \text{tp}_{\text{qf}}(\bar{w}, I_0, I) = \text{tp}_{\text{qf}}(\bar{t}_\beta, I_0, I)\} = \emptyset$, but we do not need this here.

⁴In Definition 3.2, note that \odot , “the circle property” which abstracts the above analysis will be the indicator of complexity ($T_{\mathbf{rg}}$ is \mathbf{c} -unsuperstable), whereas its negation $\neg\odot$ will be the indicator of non-complexity ($T_{\mathbf{rg}}$ is \mathbf{c} -superstable).

- *Formal description:* We require the expansion to satisfy: for any $\bar{t}', \bar{t}'' \in Y_{\bar{s}}$,
 - (a) for $\mathbf{i} = 1, 2$,

$$N^+ \models \sigma_{\mathbf{i}}(\bar{a}_{\bar{t}'}, \bar{a}_{\bar{s}}) = \sigma_{\mathbf{i}}(\bar{a}_{\bar{t}''}, \bar{a}_{\bar{s}}).$$

if and only if $J \models E_{\mathbf{i}}(\bar{t}', \bar{t}''; \bar{s})$.

- (b) $N^+ \models \sigma_1(\bar{a}_{\bar{t}'}, \bar{a}_{\bar{s}}) = \sigma_2(\bar{a}_{\bar{t}'}, \bar{a}_{\bar{s}})$ if and only if $J \models F(\bar{t}', \bar{t}''; \bar{s})$.

Once we have done this, since J is \aleph_0 -homogeneous, we can also ensure that for any \bar{u} realizing the same quantifier-free type as \bar{s} in J that the analogues of (a), (b) hold with \bar{u} in place of \bar{s} , remembering to then replace \bar{t} by some \bar{v} such that $\text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{u}, \emptyset, J) = \text{tp}_{\text{qf}}(\bar{t} \hat{\wedge} \bar{s}, \emptyset, J)$.

- *Informal description:* Take any set of distinct elements of $\text{Dom}(N)$ of size $|J|$, thus $\geq |Y_{\bar{s}}|$, and interpret the functions to take values in this set according to the following informal heuristic. Given $\bar{t}' \in Y_{\bar{s}}$, let us say “the image of \bar{t}' ” to mean $\bar{a}_{\bar{t}'} \hat{\wedge} \bar{a}_{\bar{s}}$. Then: images of elements of $Y_{\bar{s}}$ are sent to the same $b \in \text{Dom}(N^+)$ by σ_1 if and only if they are in the same E_1 -class; they are sent to the same $b \in \text{Dom}(N^+)$ by σ_2 if and only if they are in the same E_2 -class; and the values of σ_1 and σ_2 should coincide if and only if the E_1 - and E_2 -classes of the respective elements were matched by F . Ensure that the parallel conditions hold replacing \bar{s} by any other \bar{u} from J with the same quantifier-free type.

Let $\Phi' \geq \Phi$ be given by applying the Ramsey property to N^+ , so in Φ' (and any template extending it) (a) and (b) will remain true, as will their analogues for $\bar{u} \equiv_{\text{qf}} \bar{s}$.

Now in the model $N' = \text{GEM}(I, \Phi')$, consider the formula

$$R(x, \sigma_1(\bar{a}_{\bar{t}})) \wedge \neg R(x, \sigma_2(\bar{a}_{\bar{t}})).$$

Property (b) and the assumption 3.2(iii) that F has no fixed points ensure that in $N' \models \sigma_1(\bar{a}_{\bar{t}}) \neq \sigma_2(\bar{a}_{\bar{t}})$, so this is a consistent formula. However, F is a partial function and is nonempty, where non-emptiness is witnessed say by $N' \models F(\bar{w}', \bar{w}''; \bar{s})$. Since J is \aleph_0 -homogeneous and $\bar{w}' \in Y_{\bar{s}}$, for any other $\bar{t}' \in Y_{\bar{s}}$ there is $\bar{t}'' \in Y_{\bar{s}}$ such that

$$\text{tp}_{\text{qf}}(\bar{t}' \hat{\wedge} \bar{t}'', \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{w}' \hat{\wedge} \bar{w}'', \bar{s}, J).$$

Since F is an invariant of the quantifier-free type, this means $J \models F(\bar{t}', \bar{t}''; \bar{s})$. In short, the homogeneity of J tells us that if F is a partial one to one function it must be a bijection. It follows that

$$\{R(x, \sigma_1(\bar{a}_{\bar{t}})) \wedge \neg R(x, \sigma_2(\bar{a}_{\bar{t}})) : \bar{t} \in Y_{\bar{s}}\}$$

is inconsistent in the following strong sense: for every $\bar{t}_* \in Y_{\bar{s}}$, there is some $\bar{t}_{**} \in Y_{\bar{s}}$ such that

$$\{R(x, \sigma_1(\bar{a}_{\bar{t}_*})) \wedge \neg R(x, \sigma_2(\bar{a}_{\bar{t}_*})), R(x, \sigma_1(\bar{a}_{\bar{t}_{**}})) \wedge \neg R(x, \sigma_2(\bar{a}_{\bar{t}_{**}}))\}$$

is inconsistent. Finally, observe this clearly satisfies the definition of shearing, since the range of the map $f : Y_{\bar{s}} \rightarrow {}^2N'$ given by

$$\bar{t}' \mapsto \langle \sigma_1(\bar{a}_{\bar{t}'}) \rangle \hat{\wedge} \langle \sigma_2(\bar{a}_{\bar{t}'}) \rangle$$

is a $\mathcal{K}_{\mathbf{c}}$ -indiscernible sequence. This completes the discussion.

Claim 3.5 now repeats this move in the context of an inductive argument, which gives the a priori stronger conclusion of \mathbf{c} -unsuperstability; the minor but important new points to notice in the proof of 3.5 are the conditions there labelled (c), (e) which ensure that the images of the new Skolem functions at each inductive step are disjoint from those at earlier stages and allow us to “continue along the independence property” (in the random graph, disjointness is enough to ensure that the union of the formulas built at each step is indeed a type).

Remark 3.4. *Given a complete theory T and a formula $\varphi(x, y)$, let $p(x)$ be a partial φ -type in some $M \models T$. Let $\Gamma(x)$ be the infinite set of formulas expressing that $\varphi(x, y)$ has the independence property. Observe that if $p(x) \cup \Gamma(x)$ is consistent, for any κ there is $N \models T$, $M \preceq N$ containing a sequence $\langle b_i : i < \kappa \rangle$ such that φ has the independence property over this sequence [i.e., for any two finite disjoint $\sigma, \tau \subseteq \kappa$, $\{\varphi(x, b_i) : i \in \sigma\} \cup \{\neg\varphi(x, b_j) : j \in \tau\}$ is consistent] and moreover $p(x) \cup \{\varphi(x, b_i) \wedge \neg\varphi(x, b_j)\}$ is consistent for any $i \neq j$.*

Claim 3.5. *Let \mathbf{c} be a countable context and suppose \mathbf{c} has property \odot . Then:*

- (1) $T_{\mathbf{rg}}$ is \mathbf{c} -unsuperstable; moreover,
- (2) (T, φ) is \mathbf{c} -unsuperstable, for any T , φ with the independence property.

Proof. Let $\langle s_\ell : \ell < \omega \rangle$ list $I_{\mathbf{c}}$. Define $I_n = \{s_\ell : \ell < n\}$, so each $I_n \in [I_{\mathbf{c}}]^{<\aleph_0}$. By induction on n , we shall define an increasing sequence $\Phi_n \in \Upsilon[T_{\mathbf{rg}}]$ (so $n < m$ implies $\Phi_n \leq \Phi_m$) and an increasing sequence p_n of partial φ -types such that $p_n(x) \cup \Gamma(x)$ is consistent for each n , where Γ is from 3.4. (In the case of the random graph, take $\varphi(x, y) = R(x, y)$. The length of x need not be 1, but we will drop overlines for simplicity.)

For $n = 0$, let $M_0 = \text{GEM}(I, \Phi_0)$ for some $\Phi_0 \in \Upsilon[T_{\mathbf{rg}}]$, and let $p_0 = \{x = x\}$.

For $n + 1$, suppose we have defined Φ_n and $M_n = \text{GEM}(I, \Phi_n)$. Let \bar{s}_n list $I_{n,0} := I_n$. Apply \odot with $I_{n,0}$ in place of I_0 . Let $I_{n,1}$, $E_{n,0}$, $E_{n,1}$, $E_{n,2}$ be as returned by the definition of \odot , and let \bar{s}_n, \bar{t}_n list $I_{n,0}$ and $I_{n,1}$ respectively.

Next we define N_n . In the case of the random graph, let $N_n = \text{GEM}(J, \Phi_n)$. As Φ is nice, $M_n \preceq N_n$ and $N_n \upharpoonright \tau(T_{\mathbf{rg}}) \models T_{\mathbf{rg}}$. For an arbitrary theory, let N_n be an elementary extension of $\text{GEM}(J, \Phi_n)$ such that $N_n \setminus M_n$ contains a sequence $\langle b_i : i < |J| \rangle$ as in 3.4 for φ and the partial type p_n . In this case it follows additionally that $M_n \preceq N_n$ and $N_n \upharpoonright \tau(T) \models T$.

Let $\mathbf{r}_n = \text{tp}_{\text{qf}}(\bar{t}_n, \bar{s}_n, J)$, so as usual $Q_{\mathbf{r}_n}^J$ is the set of realizations of \mathbf{r}_n in J .

We may expand N_n to N_n^+ by adding two new Skolem functions $\sigma_{n,1}$ and $\sigma_{n,2}$, interpreted so that:

- (a) for $\mathbf{i} = 1, 2$, whenever $\bar{t}', \bar{t}'' \in Q_{\mathbf{r}_n}^J$, $N_n^+ \models \sigma_{n,\mathbf{i}}(\bar{a}_{\bar{t}'}, \bar{a}_{\bar{s}_n}) = \sigma_{n,\mathbf{i}}(\bar{a}_{\bar{t}''}, \bar{a}_{\bar{s}_n})$ if and only if $J \models E_{n,\mathbf{i}}(\bar{t}', \bar{t}''; \bar{s}_n)$.
- (b) $N_n^+ \models \sigma_{n,1}(\bar{a}_{\bar{t}'}, \bar{a}_{\bar{s}_n}) = \sigma_{n,2}(\bar{a}_{\bar{t}'}, \bar{a}_{\bar{s}_n})$ if and only if $J \models E_{n,0}(\bar{t}', \bar{t}''; \bar{s}_n)$.
- (c) for $m < n$ and $\mathbf{i}, \ell = 1, 2$, $N_n^+ \models \sigma_{n,\mathbf{i}}(\bar{a}_{\bar{t}_m}, \bar{a}_{\bar{s}_n}) \neq \sigma_{m,\ell}(\bar{a}_{\bar{t}_m}, \bar{a}_{\bar{s}_m})$.
- (d) For any \bar{u} realizing the same quantifier-free type as \bar{s}_n in J , the analogues of (a), (b), (c) hold with \bar{u} in place of \bar{s}_n (remembering to then replace \bar{t}_n by some \bar{v} such that $\text{tp}_{\text{qf}}(\bar{v} \hat{\ } \bar{u}, \emptyset, J) = \text{tp}_{\text{qf}}(\bar{t}_n \hat{\ } \bar{s}_n, \emptyset, J)$).
- (e) $\sigma_n(\bar{a}_{\bar{t}_i}, \bar{a}_{\bar{s}_n}) = b_i$ for some $i < \kappa$, i.e. the functions choose elements from our independent sequence. [For the random graph, essentially the whole model is an independent sequence, so it suffices to ask that $\sigma_n(\bar{a}_{\bar{t}_i}, \bar{a}_{\bar{s}_n}) \notin M_n$.]

Let $\Phi_{n+1} \in \Upsilon[T_{\mathbf{rg}}]$ extending Φ_n be given by applying the Ramsey property ([11] Corollary 2.10) to N_n^+ , and then (a), (b), (c), (d) will remain true in any template

extending Ψ_{n+1} . Moreover,

$$p_{n+1} := p_n \cup \{\varphi(x, \sigma_{n,1}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n})) \wedge \neg\varphi(x, \sigma_{n,2}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n}))\}$$

is consistent, and remains consistent with Γ from 3.4.

Let $\Phi_\omega = \bigcup_n \Phi_n$, so $\Phi_\omega \in \Upsilon[T]$. Consider any $\Psi \geq \Phi_\omega$ and let $N = \text{GEM}_{\tau(T)}(I_{\mathbf{c}}, \Psi)$. For each n , apply the non-fixed-point clause of 3.2(iii) to observe that

$$\neg E_0(\bar{t}_n, \bar{t}_n; \bar{s}_n)$$

i.e. in the model N ,

$$\sigma_{n,1}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n}) \neq \sigma_{n,2}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n}).$$

Remembering (c) and (e) above, this ensures the following set of formulas is a type:

$$p(x) = \{\varphi(x, \sigma_{n,1}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n})) \wedge \neg\varphi(x, \sigma_{n,2}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n})) : n < \omega\}.$$

So p is a partial type in $N \upharpoonright T_{\mathbf{rg}}$, and let us show that for every n it \mathbf{c} -shears over I_n . Why? First notice that $\bar{t}_n \subseteq I$ by construction. Second, notice that by the positive part of condition 3.2(iii), for each n there exists $\bar{u}_n \in Q_{\tau_n}^{J_n}$ such that

$$N \models \sigma_{n,1}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n}) = \sigma_{n,2}(\bar{a}_{\bar{u}_n}, \bar{a}_{\bar{s}_n}).$$

This means that

$$\{\varphi(x, \sigma_{n,1}(\bar{a}_{\bar{t}_n}, \bar{a}_{\bar{s}_n})) \wedge \neg\varphi(x, \sigma_{n,2}(\bar{a}_{\bar{w}'_n}, \bar{a}_{\bar{s}_n})) : \text{tp}_{\text{qf}}(\bar{t}'_n \hat{\ } \bar{w}'_n, \bar{s}_n, J_n = \bar{t}_n \hat{\ } \bar{w}_n, \bar{s}_n, J_n)\}$$

will be inconsistent. Thus T is not \mathbf{c} -superstable, as desired. \square

For the random graph, we also have a converse.

Claim 3.6. *Let \mathbf{c} be a countable context and suppose $T_{\mathbf{rg}}$ is not \mathbf{c} -superstable. Then \mathbf{c} has property \odot .*

Proof. In this proof we appeal directly to the definition of \mathbf{c} -shearing from Definition 1.2, and \mathbf{c} -unsuperstability, Definition 1.3. Let some finite $I_0 \subseteq I$ be given, and we will show how to find the rest of the data so that \odot is satisfied. Recall from Definition 1.3 that:

we say T is *unsuperstable* for \mathbf{c} when there are:

- (a) an increasing sequence of nonempty finite sets $\langle I_n : n < \omega \rangle$ with $I_m \subseteq I_n \subseteq I$ for $m < n < \omega$ and $\bigcup_n I_n = I$, which are given along with a choice of enumeration \bar{s}_n for each I_n where $\bar{s}_n \trianglelefteq \bar{s}_{n+1}$ for each n
- (b) an increasing sequence of nonempty, possibly infinite, sets $B_n \subseteq B_{n+1} \subseteq \mathfrak{C}_T$ in the monster model for T , with $B := \bigcup_n B_n$
- (c) and a partial type p over (B) , such that

$$p \upharpoonright B_{n+1} (I_n, I_{n+1})\text{-shears over } B_n.$$

Let I_n from Definition 1.3 be a finite subset of I containing I_0 . Without loss of generality, $I_0 = I_n$. Let \bar{s} be an enumeration of I_n , and let \bar{t} be an enumeration of I_{n+1} . Let N be the monster model for $T_{\mathbf{rg}}$. Let p be the type given by 1.3 which (I_n, I_{n+1}) -shears. [We can ignore the B_{n+1}, B_n from that definition.] In the remainder of the proof, we use I_0, I_1 instead of I_n, I_{n+1} . Let $\varphi(x, \bar{b}_{\bar{t}}) \in p$, \mathbf{b} be the formula and \mathcal{K} -indiscernible sequence in N which witness this instance of shearing. Recall from Definition 1.2 that:

we say *the formula* $\varphi(\bar{x}, \bar{c})$ *shears* over A in M for (I_0, I_1, \mathbf{c}) when there exist a model N , a sequence \mathbf{b} in N , enumerations \bar{s} of I_0 and \bar{t} of I_1 , and an \aleph_0 -saturated $J \supseteq I$ such that:

- (1) $I_0 \subseteq I_1$ are finite subsets of I
- (2) $M \preceq N$
- (3) $\mathbf{b} = \langle \bar{b}_{\bar{t}'} : \bar{t}' \in {}^\omega(J[I_0]) \rangle$ is \mathcal{K} -indiscernible in N over A
- (4) $\bar{c} = \bar{b}_{\bar{t}}$, and
- (5) the set of formulas

$$\{\varphi(\bar{x}, \bar{b}_{\bar{t}'}) : \bar{t}' \in \text{lg}(\bar{t})(J), \text{tp}_{\text{qf}}(\bar{t}', \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, I)\}$$

is contradictory.

Note that by the conditions on p in 1.3, this formula will be nonalgebraic, and we may assume $\ell(x) = 1$.

Let J be \aleph_0 -saturated such that $I \subseteq J \in \mathcal{K}$. Let $n = \text{lg}(\bar{b}_{\bar{t}})$. Fixing notation, let $\bar{b}_{\bar{t}} = \langle \bar{b}_{\bar{t}, i} : i < n \rangle$. Since $\varphi(x, \bar{b}_{\bar{t}})$ is a consistent nonalgebraic formula in the random graph, we can write n as the disjoint union of sets A, B, C so that

$$\varphi(x, \bar{b}_{\bar{t}}) \equiv \bigwedge_{i \in A} R(x, \bar{b}_{\bar{t}, i}) \wedge \bigwedge_{j \in B} \neg R(x, \bar{b}_{\bar{t}, j}) \wedge \bigwedge_{k \in C} x \neq \bar{b}_{\bar{t}, k}.$$

Define $Y_{\bar{s}} = \{\bar{t}' \in \text{lg}(\bar{t})(J) : \text{tp}_{\text{qf}}(\bar{t}', \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, J)\}$. In order to define E_1, E_2 , observe that since the set of formulas

$$\{\varphi(\bar{x}, \bar{b}_{\bar{t}'}) : \bar{t}' \in \text{lg}(\bar{t})(J), \text{tp}_{\text{qf}}(\bar{t}', \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, I)\}$$

is contradictory, there must be some $\bar{t}_\alpha, \bar{t}_\beta \in Y_{\bar{s}}$ and $i_\alpha \in A, j_\beta \in B$ such that

$$\bar{b}_{\bar{t}_\alpha, i_\alpha} = \bar{b}_{\bar{t}_\beta, j_\beta}.$$

Let $E_1(\bar{x}_1, \bar{x}_2; \bar{s})$ be the following two-place relation on $Y_{\bar{s}}$:

$$E_1(\bar{t}_1, \bar{t}_2; \bar{s}) \text{ iff } N \models \bar{b}_{\bar{t}_1, i_\alpha} = \bar{b}_{\bar{t}_2, i_\alpha}.$$

Let $E_2(\bar{x}_1, \bar{x}_2; \bar{s})$ be the following two-place relation on $Y_{\bar{s}}$:

$$E_2(\bar{t}_1, \bar{t}_2; \bar{s}) \text{ iff } N \models \bar{b}_{\bar{t}_1, j_\beta} = \bar{b}_{\bar{t}_2, j_\beta}.$$

Then for $\ell = 1, 2$, clearly:

- $E_\ell = E_\ell(\bar{x}_1, \bar{x}_2; \bar{s})$ is an equivalence relation on $Y_{\bar{s}}$.
- The truth value of “ $E_\ell(\bar{t}_1, \bar{t}_2; \bar{s})$ ” is determined by $\text{tp}_{\text{qf}}(\bar{t}_1 \wedge \bar{t}_2, \bar{s}, J)$, for any $\bar{t}_1, \bar{t}_2 \in Y_{\bar{s}}$, in other words, it is an invariant of the quantifier-free type (thus, said to be definable by a possibly infinitary quantifier-free formula).

Finally, define $F(\bar{x}_1, \bar{x}_2; \bar{s})$, a two place relation on $Y_{\bar{s}}$, by: $F(\bar{t}_1, \bar{t}_2; \bar{s})$ if and only if

$$N \models \bar{b}_{\bar{t}_1, i_\alpha} = \bar{b}_{\bar{t}_2, j_\beta}.$$

Then $F(\bar{x}_1, \bar{x}_2; \bar{s})$ defines a nonempty one to one partial function from $Y_{\bar{s}}/E_1(-, -; \bar{s})$ to $Y_{\bar{s}}/E_2(-, -; \bar{s})$.

It remains to check that F has no fixed points. Recall that for our original tuple \bar{t} , we know that

$$\{\bar{b}_{\bar{t}, i} : i \in A\} \cap \{\bar{b}_{\bar{t}, j} : j \in B\} = \emptyset$$

because $\varphi(x, \bar{b}_{\bar{t}})$ is a consistent $\tau(T_{\text{rg}})$ -formula in N . Thus, $J \models \neg F(\bar{t}, \bar{t}; \bar{s})$. Since F is definable by a possibly infinitary quantifier-free formula, this remains true for all tuples from $Y_{\bar{s}}$. This completes the verification of \odot . \square

3.3. Discussion: obtaining \odot directly from the example in 3.1. For completeness, we work out the analysis of 3.1. Recall equations (5) and (6) there. Recall that $\bar{t}_\alpha, \bar{t}_\beta$ come from equation (2) for our given α, β (so \bar{v}_* and \bar{w}_* share their respective quantifier-free types over \bar{s}). As J is \aleph_0 -homogeneous, there are \bar{v}, \bar{w} from J such that

$$\text{tp}_{\text{qf}}(\bar{v}_* \hat{\wedge} \bar{w}, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{w}_*, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}_\alpha \hat{\wedge} \bar{t}_\beta, \bar{s}, I).$$

(It may be that $\bar{v} = \bar{v}_*$ or $\bar{w} = \bar{w}_*$, and if so, no problem.)

Let $\bar{t} = \bar{s} \hat{\wedge} \bar{t}_\alpha \hat{\wedge} \bar{t}_\beta$. Without loss of generality, we may simultaneously replace \bar{v}_* and \bar{w}_* respectively by

$$\bar{s} \hat{\wedge} \bar{v}_* \hat{\wedge} \bar{w}, \quad \bar{s} \hat{\wedge} \bar{v} \hat{\wedge} \bar{w}_*.$$

So for us,

$$Y_{\bar{s}} = \{\bar{t}' \in {}^{\text{lg}(\bar{t})}J : \text{tp}_{\text{qf}}(\bar{t}', \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, J)\}$$

and this set includes \bar{v}_* and \bar{w}_* . After possibly adding dummy variables to $\sigma_\alpha, \sigma_\beta$ we may assume their operation is unchanged. Let $E_1(\bar{x}_1, \bar{x}_2; \bar{s})$ be the following two-place relation on $Y_{\bar{s}}$:

$$E_1(\bar{t}_1, \bar{t}_2; \bar{s}) \text{ iff } N \models \text{“}\sigma_\alpha(\bar{a}_{\bar{t}_1}) = \sigma_\alpha(\bar{a}_{\bar{t}_2})\text{”}.$$

Let $E_2(\bar{x}_1, \bar{x}_2; \bar{s})$ be the following two-place relation on $Y_{\bar{s}}$:

$$E_2(\bar{t}_1, \bar{t}_2; \bar{s}) \text{ iff } N \models \text{“}\sigma_\beta(\bar{a}_{\bar{t}_1}) = \sigma_\beta(\bar{a}_{\bar{t}_2})\text{”}.$$

Then for $\ell = 1, 2$, clearly:

- $E_\ell = E_\ell(\bar{x}_1, \bar{x}_2; \bar{s})$ is an equivalence relation on $Y_{\bar{s}}$.
- The truth value of “ $E_\ell(\bar{t}_1, \bar{t}_2; \bar{s})$ ” is determined by $\text{tp}_{\text{qf}}(\bar{t}_1 \hat{\wedge} \bar{t}_2, \bar{s}, J)$, for any $\bar{t}_1, \bar{t}_2 \in Y_{\bar{s}}$, in other words, it is an invariant of the quantifier-free type (thus, said to be definable by a possibly infinitary quantifier-free formula).

Finally, define $F(\bar{x}, \bar{x}_2; \bar{s})$, a two place relation on $Y_{\bar{s}}$, by: $F(\bar{t}_1, \bar{t}_2; \bar{s})$ if and only if

$$N \models \sigma_\alpha(\bar{a}_{\bar{t}_1}) = \sigma_\beta(\bar{a}_{\bar{t}_2}).$$

Then F naturally defines a subset $X_{\bar{s}}$ of $Y_{\bar{s}} \times Y_{\bar{s}}$ i.e. if $\bar{t}_1, \bar{t}_2 \in Y_{\bar{s}}$ then the truth value of $F(\bar{t}_1, \bar{t}_2; \bar{s})$ is the same for any $(\bar{t}'_1, \bar{t}'_2) \in (\bar{t}_1/E_1) \times (\bar{t}_2/E_2)$. F is not empty because of equation (6), and respects the equivalence relations. (If $(\bar{t}'_\ell, \bar{t}''_\ell) \in F$ for $\ell = 1, 2$ then $\bar{t}'_1 E_1 \bar{t}'_2$ iff $\bar{t}''_1 E_2 \bar{t}''_2$. Together we get that F is a 1-to-1 partial function from Y/E_1 into Y/E_2 , so by homogeneity of J it is a full function onto.)

It remains to show that F has no fixed points. Now, whether or not $F(\bar{u}, \bar{u}; \bar{s})$ is an invariant of the quantifier-free type $\text{tp}_{\text{qf}}(\bar{u}, \bar{s}, J)$. We know that

$$N \models \sigma_\alpha(\bar{a}_{\bar{t}}) \neq \sigma_\beta(\bar{a}_{\bar{t}})$$

by our definition of \bar{t} , because the original type p was consistent. It follows that

$$\neg F(\bar{t}, \bar{t}; \bar{s})$$

and that this is an invariant of $\text{tp}_{\text{qf}}(\bar{t}, \bar{s}, I)$. So it will remain be true for any tuple from $Y_{\bar{s}}$. This proves F has no fixed points, which completes the verification of \odot and the discussion.

Summarizing, we arrive at:

Theorem 3.7. *Let \mathbf{c} be a countable context. T_{rg} is \mathbf{c} -unsuperstable if and only if \mathbf{c} has property \odot .*

Proof. Claim 3.5 and Claim 3.6. □

We now give a positive and a negative example of \odot . First, we verify that $T_{\mathbf{rg}}$ is \mathbf{c} -unsuperstable for contexts coming from linear orders. In the following example, the context need not be countable.

Example 3.8. *Let \mathcal{K} be the class of infinite linear orders, Let $I \in \mathcal{K}$. Then $\mathbf{c} = (I, \mathcal{K})$ has \odot .*

Proof. Let $M \models T_{\mathbf{rg}}$ and let $\langle a_t : t \in I \rangle$ be any infinite indiscernible sequence. Recall that if $\bar{t} = t_0 t_1$ we let $\bar{a}_{\bar{t}}$ denote $a_{t_0} \hat{\ } a_{t_1}$. Choose $t_0^*, t_1^* \in I$ with $t_0^* < t_1^*$. Let $\mathbf{r} = \text{tp}_{\text{qf}}(t_0^* \hat{\ } t_1^*, \emptyset, I)$. (In this example, $\bar{s} = \langle \rangle$, $\bar{t} = \langle t_0^*, t_1^* \rangle$ are the corresponding values from \odot .) Recall that $Q_{\mathbf{r}}^I$ is the set of realizations of \mathbf{r} in I . Then

$$\{\bar{a}_{\bar{t}} : \bar{t} \in {}^2 I, \bar{t} \in Q_{\mathbf{r}}^I\}$$

is a \mathcal{K} -indiscernible sequence. Define E_0, E_1, E_2 on $Q_{\mathbf{r}}^I$ as follows:

$$E_1(t_1 t_2, t'_1 t'_2) \equiv t_1 = t'_1$$

Now

$$\begin{aligned} E_2(t_1 t_2, t'_1 t'_2) &\equiv t_2 = t'_2 \\ E_0(t_1 t_2, t'_1 t'_2) &\equiv t_1 = t'_2. \end{aligned}$$

We can also see the \mathbf{c} -dividing directly:

$$\{\varphi(x, \bar{a}_{\bar{t}}) : \bar{t} \in Q_{\mathbf{r}}^I\}$$

is inconsistent since for every $t_1 t_2 \in Q_{\mathbf{r}}^I$, there is $t_0 \in I$ such that $t_0 t_1 \in Q_{\mathbf{r}}^I$. \square

There are also natural examples which do not have \odot . In the following example, I also need not be countable.

Example 3.9. *Let \mathcal{K}_μ be the class of linear orders expanded by μ unary predicates which partition the domain. Let $I \in \mathcal{K}_\mu$ and suppose that for each predicate P_α , $|P_\alpha^I| \leq 1$. Let $\mathbf{c} = (I, \mathcal{K}_\mu)$. Then \mathbf{c} does not have \odot .*

Proof. See [10] Claim 5.10, where it is shown that for any such \mathbf{c} and $M = \text{GEM}(I, \Phi)$, if p is a type of $M \upharpoonright \tau(T_{\mathbf{rg}})$ then there is $\Psi \geq \Phi$ in which p is realized. (It follows that $T_{\mathbf{rg}}$ is \mathbf{c} -superstable.) *For the interested reader, the main idea of that proof is quite close to that of Claim 5.5 below; just leave out the relations R on J .* \square

Conclusion 3.10. *The following are equivalent:*

- (a) *The formula φ is stable (with respect to the complete theory T).*
- (b) *(T, φ) is \mathbf{c} -superstable for every countable context \mathbf{c} .*

Proof. For (a) implies (b): This is because stable formulas have definitions, thus a fortiori weak definitions. We assemble some facts from [11] for a more detailed proof. Suppose for a contradiction that (T, φ) were \mathbf{c} -unsuperstable for some countable context $\mathbf{c} = (I, \mathcal{K})$. Let $M = \text{GEM}(I, \Phi)$. Then [11] Corollary 6.16 would build a larger template Φ_* so that for any $\Psi \geq \Phi_*$, the model $\text{GEM}(I, \Psi)$ is not \aleph_1 -saturated for φ -types. In particular, that construction builds a φ -type p which does not have a weak \bar{t}_* -definition for any finite \bar{t}_* in I , and moreover cannot have one in any $\text{GEM}(I, \Psi)$ for any $\Psi \geq \Phi_*$, see [11] Remark 6.15 or Corollary 4.25. However, by [11] Claim 4.15, in any $\text{GEM}(I, \Psi)$ any type in a stable formula has a definition, thus a weak definition. Contradiction.

For (b) implies (a): Suppose first that φ is not simple (has the tree property). Then it has long dividing chains, and since dividing implies shearing ([11], Claim 5.8 and Remark 5.9), φ is not \mathbf{c} -superstable. If φ is simple unstable, φ has the independence property, and we can apply Claim 3.5(b). \square

4. INTERLUDE ON “eq”

In this section, we observe that it isn’t an accident that our negative examples of \odot have a certain form. There is a nice further explanation of this, and of \odot , once we define the analogue of M^{eq} in this context.

Definition 4.1. *We say that the countable context $\mathbf{c} = (I, \mathcal{K})$ is essentially separated when there is a finite $I_0 \subseteq I$ such that $s \neq t \in I$ implies that $\text{tp}_{\text{qf}}(s, I_0, I) \neq \text{tp}_{\text{qf}}(t, I_0, I)$.*

The idea is that “ $\neg\odot$ ” is for all intents and purposes essentially separated (one might say, is essentially *essentially separated*). In this section, we outline a proof of this, which involves defining the analogue of M^{eq} for contexts, defining for any context \mathbf{c} a so-called “eq-extension” and then pointing out that the property $\neg\odot$ for \mathbf{c} is really saying that some such “ \mathbf{c}^{eq} ” is well behaved. Since the section is primarily explanatory, we will be brief.

Definition 4.2. *We say the context \mathbf{d} is an eq-extension of our context \mathbf{c} when there is \bar{E} such that:*

- (1) $\bar{E} = \langle E_i : i < i_* \rangle$, $\bar{\varphi} = \langle \varphi_j : j < j_* \rangle$
- (2) $E_i = E_i(\bar{x}_i, \bar{y}_i)$ is a possibly infinitary quantifier-free formula in the vocabulary $\tau(\mathcal{K})$, with $\text{lg}(\bar{x}_i) = \text{lg}(\bar{y}_i) =: n_i$, such that for every $I' \in \mathcal{K}$ it defines an equivalence relation on tuples of I' of length n_i .
We stipulate $E_0 = “x = y”$. (Alternately, list the defining formulas φ_i for E_i separately.)
- (3) For $I \in \mathcal{K}$, let I^+ be the analogue of \mathfrak{C}^{eq} using the E_i ’s for $i < i_*$. (Why not all possible E s? as then we may lose countability.) That is,
 - (a) the universe of I^+ is $I \cup \{\bar{t}/E_i^I : i < i_*, \bar{t} \in {}^{n_i}I\}$,
but we identify t/E_0 with t for $t \in I$.
 - (b) $P^{I^+} = P^I$ for $P \in \tau(\mathcal{K})$ a predicate.
 - (c) $P_*^{I^+} = I$.
 - (d) for $F \in \tau(\mathcal{K})$ a function symbol, if any, $F^{I^+} = F$, so its domain is defined.
 - (e) $F_i^{I^+} = \{(\bar{t}, \bar{t}/E_i^I) : \bar{t} \in {}^{n_i}I\}$.
 - (f) $P_{\varphi_j}^{I^+} = \{(\bar{t}_{i_0}/E_{i_0}, \dots, \bar{t}_{i_{m(j)-1}}/E_{i_{m(j)-1}}) : m(j) < \omega, i_\ell < i_*, \text{ and } I \models \varphi[\bar{t}_{i_0}, \bar{t}_{i_1}, \dots]\}$.
 - (g) $\mathcal{K}_{\mathbf{d}} = \{I^+ : I \in \mathcal{K}_{\mathbf{c}}\}$, and $I_{\mathbf{d}} = (I_{\mathbf{c}})^+$, so we have defined $\mathbf{d} = (I_{\mathbf{d}}, \mathcal{K}_{\mathbf{d}})$.

Definition 4.3. *We say $\mathbf{d} = \mathbf{c}^{\text{eq}}$ when we use all possible E_i, φ_i up to equivalence. (Since we have required these sequences to be countable, this of course puts some restrictions on the contexts \mathbf{c} for which \mathbf{c}^{eq} is presently defined. We will not at present require such a canonical extension to exist, but it is reasonable to define it.)*

Claim 4.4. *For any countable context \mathbf{c} , any \mathbf{d} defined from it as in 4.2 is also a context. Moreover, if $\lambda \geq |I_{\mathbf{c}}| + |\{\text{tp}(\bar{s}, \emptyset, I) : \bar{s} \in {}^{\omega}I, I \in \mathcal{K}_{\mathbf{c}}\}|$ then also $\lambda \geq |I_{\mathbf{d}}| + |\{\text{tp}(\bar{s}, \emptyset, I) : \bar{s} \in {}^{\omega}I, I \in \mathcal{K}_{\mathbf{d}}\}|$.*

Definition 4.5. For $I \in \mathcal{K}_{\mathbf{c}}$ and finite $\bar{s} \in \omega^{>}I$:

- (1) let $\text{dcl}(\bar{s}, I) = \{t \in I : \text{we cannot find } J \in \mathcal{K}_{\mathbf{c}}, \bar{w} \in \text{lg } \bar{s} J \text{ and } t_1 \neq t_2 \in J \text{ such that for } \ell = 1, 2, \text{tp}_{\text{qf}}(\bar{w} \hat{\ } \langle t_\ell \rangle, \emptyset, J) = \text{tp}_{\text{qf}}(\bar{s} \hat{\ } \langle t \rangle, \emptyset, J)\}$.
- (2) let $\text{acl}(\bar{s}, I)$ be defined similarly, replacing “ $t_1 \neq t_2$ ” by “ $\langle t_\ell : \ell < \omega \rangle$ ”.

Claim 4.6. For a countable context \mathbf{c} the following are equivalent.

- (a) \mathbf{c} satisfies \odot .
- (b) for some eq-extension \mathbf{d} of \mathbf{c} , for every finite \bar{s} from $I_{\mathbf{d}}$, there are $r_0 \neq r_1 \in I_{\mathbf{d}}$ realizing the same complete quantifier-free type over \bar{s} .
- (c) for every eq-extension \mathbf{d} of \mathbf{c} , there is a finite $I_0 \subseteq I_{\mathbf{d}}$ such that $\text{dcl}(I_0, I_{\mathbf{d}}) \neq I_{\mathbf{d}}$.

Proof. We prove the nontrivial arrow, (c) implies (a). Fixing \mathbf{d} and $I_0 \subseteq I_{\mathbf{d}}$, we can choose $\bar{s} \in \omega^{>}(I_{\mathbf{c}})$ such that $I_0 \subseteq \text{dcl}(\bar{s}, I_{\mathbf{d}})$, and we can find $r_1 \in I_{\mathbf{d}}$ such that $r_1 \notin \text{dcl}(\bar{s}, I_{\mathbf{d}})$. Let \bar{t}_1 and E_1 be such that $r_1 = \bar{s} \hat{\ } \bar{t}_1 / E_1$. Likewise, there is $r_2 \in I_{\mathbf{d}}$ realizing $\text{tp}_{\text{qf}}(r_1, \bar{s}, I_{\mathbf{d}})$ and so there are $\bar{t}_2 \in \omega^{>}I_{\mathbf{d}}$ and E_2 such that $r_2 = \bar{s} \hat{\ } \bar{t}_2 / E_2$. Without loss of generality, $\bar{t}_1 = \bar{t}_2$, call it \bar{t} , and without loss of generality $\bar{s} \hat{\ } \bar{t}$ has no repetitions. Let $J \in \mathcal{K}_{\mathbf{c}}$ be \aleph_0 -saturated extending $I_{\mathbf{c}}$, hence $J^+ \in \mathcal{K}_{\mathbf{d}}$ is \aleph_0 -saturated extending $I_{\mathbf{d}}$. As $\text{tp}_{\text{qf}}(r_1, \bar{s}, I_{\mathbf{d}}) = \text{tp}_{\text{qf}}(r_2, \bar{s}, I_{\mathbf{d}})$, it remains to define F , as follows. In J^+ we let $F = \{\bar{u} \hat{\ } \bar{v} / E_1, \bar{u} \hat{\ } \bar{v} / E_2 : \bar{u} \hat{\ } \bar{v} \subseteq J^+ \text{ realizes } \text{tp}_{\text{qf}}(\bar{s} \hat{\ } \bar{t}, \emptyset, I_{\mathbf{d}} = I_{\mathbf{d}}^+)\}$. \square

Remark 4.7. In the previous claim, note that as we can use \aleph_0 -saturated $I \in \mathcal{K}_{\mathbf{c}}$, $\text{dcl}(-, I) = \text{acl}(-, I)$.

5. SEPARATING THE RANDOM GRAPH AND $T_{n,k}$

We now show one can separate T_{rg} and $T_{3,2}$ using countable contexts. That is, for each $n > k \geq 2$, we prove that there are countable contexts \mathbf{c} for which T_{rg} is \mathbf{c} -superstable but $T_{n,k}$ is \mathbf{c} -unsuperstable, Theorem 5.8.

Definition 5.1. Let $\mathcal{K}_{n,k}$ be the following index model class. In $\tau(\mathcal{K}_{n,k})$ we have a binary relation $<$, unary predicates $\{P_q : q \in \mathbb{Q}\}$, and a $(k+1)$ -place relation R , and on $I \in \mathcal{K}$:

- $<$ is a linear order
- the P_q are disjoint unary predicates which partition I
- R is symmetric irreflexive (a hyperedge), and has no cliques of size $n+1$, i.e. for any distinct i_0, \dots, i_n from I it is not the case that R^I holds on all $k+1$ -element subsets of $\{i_0, \dots, i_n\}$. [Note this is a universal statement.]

Claim 5.2. $\mathcal{K}_{n,k}$ is a Ramsey class and (satisfies our hypotheses).

Proof. The key point is being a Ramsey class. By a theorem of Scow, it suffices to check this for the class of finite substructures of members of \mathcal{K} , so we may cite the general Nešetřil-Rödl theorem for relational structures, see [5] Theorem 3.21. \square

For the next few claims we shall use:

Definition 5.3. Given $n > k \geq 2$, let $\mathbf{c}_{n,k}$ be the context (I, \mathcal{K}) where $\mathcal{K} = \mathcal{K}_{n,k}$ and I has domain \mathbb{Q} , $<^I$ is the usual linear order on \mathbb{Q} , $P_q^I = \{q\}$, and $R^I = \emptyset$.

Discussion 5.4. Clearly each $\mathbf{c}_{n,k}$ is a countable context. Recall the intent of n, k : the edge relation R has arity $k + 1$, and the forbidden configuration is a ‘large clique’ where large means size $n + 1$. Note that since we make no demands on R in 5.1 other than having no large cliques, there is no problem choosing an I in which there are no instances of R at all. However, the point will be that when we consider \aleph_0 -saturated $J \supseteq I$, instances of R will appear.

Claim 5.5. $T_{\mathbf{rg}}$ is $\mathbf{c}_{n,k}$ -superstable, for any $n > k \geq 2$.

Proof. Fix n, k for the course of the proof and let $\mathbf{c} = \mathbf{c}_{n,k}$, $I = I_{\mathbf{c}}, \mathcal{K} = \mathcal{K}_{\mathbf{c}}$. It will suffice to show there is no instance of \mathbf{c} -dividing for non-algebraic formulas $\varphi(x, \bar{a})$ in the monster model for the random graph.

Step 1: Analysis of the problem. Suppose for a contradiction that there were some:

- finite \bar{s}, \bar{t} from I (without loss of generality $\bar{s} \trianglelefteq \bar{t}$ and without loss of generality \bar{t} is strictly increasing without repetition)
- $\mathbf{r} = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, I)$
- J which is \aleph_0 -saturated with $I \subseteq J \in \mathcal{K}$,
- a \mathcal{K} -indiscernible sequence $\{\bar{b}_{\bar{t}'} : \bar{t}' \in Q_{\mathbf{r}}^J\}$ in the monster model for $T_{\mathbf{rg}}$,
- and a formula $\varphi(x, \bar{y})$ with $\ell(\bar{y}) = \ell(\bar{b}_{\bar{t}})$,

such that

$$(9) \quad \varphi(x, \bar{b}_{\bar{t}}) \text{ is consistent and non-algebraic}$$

however

$$(10) \quad \{\varphi(x, \bar{b}_{\bar{t}'}) : \bar{t}' \in Q_{\mathbf{r}}^J\} \text{ is inconsistent.}$$

As before, it suffices to consider the case where $\ell(x) = 1$, and we may partition $m = \ell(\bar{b}_{\bar{t}})$ into disjoint sets A, B, C such that

$$\varphi(x, \bar{b}_{\bar{t}}) \equiv \bigwedge_{i \in A} R(x, \bar{b}_{\bar{t}, i}) \wedge \bigwedge_{j \in B} \neg R(x, \bar{b}_{\bar{t}, j}) \wedge \bigwedge_{k \in C} x \neq \bar{b}_{\bar{t}, k}.$$

As we have excluded dividing because of equality, the contradiction in (10) must be because we have $\bar{v}, \bar{w} \in Q_{\mathbf{r}}^J$ and i, j such that $i \in A, j \in B$, and

$$(11) \quad \bar{a}_{\bar{v}, i} = \bar{a}_{\bar{w}, j}.$$

Recall here that $\bar{a}_{\bar{v}, i}$ denotes the i -th element of the tuple $\bar{a}_{\bar{v}}$.

Step 2: A property that we would like \bar{v}, \bar{w} , our witnesses to collision, to have. Let \bar{u} enumerate, in increasing order, the set $\text{Range}(\bar{v}) \cap \text{Range}(\bar{w})$ of elements common to both sequences. (By the second line of Step 1, this will include the elements of \bar{s} .) Here we will follow an idea from the proof of [10], Claim 5.10. By ‘‘an interval of consecutive elements of \bar{u} ’’ we shall mean a set of elements which are all less than u_0 , or all greater than $u_{\text{lg}(\bar{u})-1}$, or all strictly between u_i, u_{i+1} for some $0 \leq i < \text{lg}(\bar{u}) - 1$.

Now consider the following potential property of \bar{v}, \bar{w} :

- (\star) within each interval of consecutive elements of \bar{u} , all elements of \bar{v} falling in this interval are strictly below all elements of \bar{w} falling in the same interval.

In this step and the next, let us show that we may assume our \bar{v}, \bar{w} (which were chosen to witness the contradiction) also satisfy (\star).

Suppose not, that is, suppose we chose our \bar{v}, \bar{w} so that⁵

$$\{(i, j) : v_i, w_j \text{ fall in the same interval of } \bar{u} \text{ but } v_i \leq w_j \}$$

is minimized, but we were not able to choose this number to be zero. That is, within at least one interval, say (u_i, u_{i+1}) of \bar{u} , we have elements v_k, w_j such that the following holds. ((If one of the endpoints is $+\infty$ or $-\infty$, the same argument applies substituting this notation throughout.)

$$\begin{aligned} u_i < \{v \in \bar{v} : u_i < v < v_k\} \cup \{w \in \bar{w} : u_i < w < w_j\} \\ &< w_j < v_k < \\ &\{v \in \bar{v} : v_k < v < u_{i+1}\} \cup \{w \in \bar{w} : w_j < w < u_{i+1}\} < u_{i+1}. \end{aligned}$$

where some or all of the sets in the first and third lines may be empty. Recalling that J is \aleph_0 -saturated, we will justify in the next step that we may choose w'_j, v'_k so that

- $w_j < v'_k < w'_j < v_k$ and
- writing \bar{w}' for the result of substituting w'_j for w_j in \bar{w} , and writing \bar{v}' for the result of substituting v'_k for v_k in \bar{v} , we have that

$$\text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{w}, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{v}' \hat{\wedge} \bar{w}, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{w}', \bar{s}, J).$$

Why is this sufficient? Recalling that (11) is an invariant of $\text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{w}, \bar{s}, J)$, we will then have that

$$\begin{aligned} \bar{a}_{\bar{v}, i} &= \bar{a}_{\bar{w}, j} \\ \bar{a}_{\bar{v}, i} &= \bar{a}_{\bar{w}', j} \\ \bar{a}_{\bar{v}', i} &= \bar{a}_{\bar{w}, j} \end{aligned}$$

so by transitivity of equality, $\bar{a}_{\bar{v}', i} = \bar{a}_{\bar{w}', j}$. But now \bar{v}', \bar{w}' are witnesses to the collision which have a strictly lower number of $w - v$ crossings than \bar{v}, \bar{w} , whose number of crossings we had assumed to be minimal. This contradiction shows that, modulo the next step, we may indeed choose \bar{v}, \bar{w} to have property (\star) .

Step 3: There is no nontrivial algebraicity in J . Suppose we are given a finite set $c_0 < \dots < c_p$ from J , and $d \in J$ such that $c_r < d < c_{r+1}$. Then we claim the realizations of $\text{tp}_{\text{qf}}(d, \bar{c}, J)$ are dense in the interval $(c_r, c_{r+1})_J$. Why? This quantifier-free type is determined by the ordering $<$, the predicate which holds of d , and a partition of $[\{c_0, \dots, c_p\}]^k$ into $X \cup Y$ so that the quantifier-free type specifies $\{R(d, \tau) : \tau \in X\}$ and $\{-R(d, \tau) : \tau \in Y\}$. Since the type already has a realization d , X cannot contain any $(n+1)$ -cliques. So this collection of conditions is consistent, and is realized densely in the interval $(c_r, c_{r+1})_J$ by \aleph_0 -saturation.

Step 4: Using property (\star) to contradict inconsistency. At this point in the proof, we have \bar{v}, \bar{w}, i, j witnessing the contradiction, where \bar{v}, \bar{w} satisfy property (\star) from Step 2. We claim we can choose $\bar{z} \in Q_{\mathfrak{r}}^J$ such that

$$(12) \quad \text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{w}, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{v} \hat{\wedge} \bar{z}, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{z} \hat{\wedge} \bar{w}, \bar{s}, J).$$

Why? It suffices to show we can do this within each interval of \bar{u} . Within each interval of \bar{u} , all the elements of v are below all the elements of w in the interval.

⁵informally, the sum over all intervals of \bar{u} the number of elements of \bar{w} less than elements of \bar{v} in each given interval

Suppose the number of elements of w in the interval is n_w . Let v_a, w_b be the maximal element of v in the interval and the minimal element of w in the interval, respectively. (Since \bar{v}, \bar{w} were assumed to be in strictly increasing order, v_{a-1}, w_{b+1} are the next largest and next smallest, respectively; the elements of w in the interval are, in order, $w_b < \dots < w_{b+n_w-1}$.) By induction on $\ell < n_w$, we apply the claim of Step 3 with \bar{c} an enumeration of $\text{Range}(\bar{v}) \cup \text{Range}(\bar{w}) \setminus w_{b+\ell}$, and $d = w_{b+\ell}$, to choose a realization $w'_{b+\ell}$ which is $< w_b$ (and necessarily strictly above v_a). Informally, one by one, we copy the elements of w to the left. Note that equation (12) implies a fortiori that $\text{tp}_{\text{qf}}(\bar{z}, \bar{s}, J) = \text{tp}_{\text{qf}}(\bar{t}, \bar{s}, J)$, which we called $\bar{\tau}$.

The result is \bar{z} with the desired property, and note that within each interval of \bar{u} , all the elements of \bar{v} there are below all the elements of \bar{z} there which are in turn below all the elements of \bar{w} there.

For the final contradiction, recall our original assumption (11) which was an invariant of the quantifier-free type $\text{tp}_{\text{qf}}(\bar{v} \wedge \bar{w}, \bar{s}, J)$. Thus

$$\begin{aligned} \bar{a}_{\bar{v},i} &= \bar{a}_{\bar{w},j} \\ \bar{a}_{\bar{z},i} &= \bar{a}_{\bar{w},j} \\ \bar{a}_{\bar{v},i} &= \bar{a}_{\bar{z},j} \end{aligned}$$

so by transitivity of equality, $\bar{a}_{\bar{z},i} = \bar{a}_{\bar{z},j}$. This must be an invariant of $\text{tp}_{\text{qf}}(\bar{z}, \bar{s}, J)$ but since this equals $\text{tp}_{\text{qf}}(\bar{t}, \bar{s}, J)$, our original formula must have been inconsistent, contradicting (9). This completes the proof. \square

For the converse collection of claims, showing $\mathbf{c}_{n,k}$ -unsuperstability for the theories $T_{n,k}$, we begin with a warm-up, deriving a single instance of dividing for the case of $n = 3, k = 2$. We shall then upgrade this result to unsuperstability and to general n, k . However, in this warm-up case the notation is a bit simpler, and all the main ideas are represented.

Claim 5.6. *The theory $T_{3,2}$ contains nontrivial $\mathbf{c}_{3,2}$ -dividing, coming from a formula which is a conjunction of positive instances of the edge relation.*

Proof. We start with $I = I_{\mathbf{c}_{3,2}}$, which is countable, linearly ordered, each element is named by a different predicate, and there are no instances of R . Choose $\mathbf{t}_0, \mathbf{t}_1, \mathbf{t}_2 \in I$ such that $I \models P_0(\mathbf{t}_0) \wedge P_1(\mathbf{t}_1) \wedge P_2(\mathbf{t}_2)$, so it follows that $\mathbf{t}_0, \mathbf{t}_1, \mathbf{t}_2$ are distinct and (without loss of generality) $\mathbf{t}_0 < \mathbf{t}_1 < \mathbf{t}_2$. Because these elements are from I , note that their quantifier-free type specifies there are no instances of R . We will write $\bar{\mathbf{t}} = \langle \mathbf{t}_0, \mathbf{t}_1, \mathbf{t}_2 \rangle$.

Consider some \aleph_0 -saturated $J \in \mathcal{K}$ with $I \subseteq J$. Let $M \models T_{n,k}$ be fairly saturated. Choose a \mathcal{K} -indiscernible sequence $\langle a_t : t \in J \rangle$ so that for any distinct $t_{i_0}, t_{i_1}, t_{i_2}$ from J ,

$$M \models R(a_{t_{i_0}}, a_{t_{i_1}}, a_{t_{i_2}}) \iff J \models R(t_{i_0}, t_{i_1}, t_{i_2}).$$

(The point is that the forbidden configuration never occurs on J , so we can build the map indexing the indiscernible sequence, say, by induction.) Now recalling $\bar{\mathbf{t}}$ from the first paragraph, consider in M the formula

$$\varphi(x, \bar{a}_{\bar{\mathbf{t}}}) = R(x, a_{\mathbf{t}_0}, a_{\mathbf{t}_1}) \wedge R(x, a_{\mathbf{t}_0}, a_{\mathbf{t}_2}) \wedge R(x, a_{\mathbf{t}_1}, a_{\mathbf{t}_2}).$$

By our choice of $\bar{\mathbf{t}}$ and our choice of the indiscernible sequence, $M \models \neg R(a_{\mathbf{t}_0}, a_{\mathbf{t}_1}, a_{\mathbf{t}_2})$ (because the triple of indices come from I , which has no instances of R) so $\varphi(x, \bar{a}_{\bar{\mathbf{t}}})$

is a consistent formula of M . Since J is \aleph_0 -saturated for \mathcal{K} , we may find v_0, v_1, v_2 in J such that:⁶

$$\begin{aligned} \text{tp}_{\text{qf}}(\mathbf{t}_0 \wedge \mathbf{t}_1 \wedge \mathbf{t}_2, \emptyset, I) &= \text{tp}_{\text{qf}}(v_0 \wedge v_1 \wedge \mathbf{t}_2, \emptyset, J) \\ &= \text{tp}_{\text{qf}}(v_0 \wedge \mathbf{t}_1 \wedge v_2, \emptyset, J) \\ &= \text{tp}_{\text{qf}}(\mathbf{t}_0 \wedge v_1 \wedge v_2, \emptyset, J) \end{aligned}$$

and $J \models R(v_0, v_1, v_2)$. Then writing

$$\begin{aligned} \bar{t}_{\{1,2\}} &= \mathbf{t}_0 \wedge v_1 \wedge v_2 \\ \bar{t}_{\{0,2\}} &= v_0 \wedge \mathbf{t}_1 \wedge v_2 \\ \bar{t}_{\{0,1\}} &= v_0 \wedge v_1 \wedge \mathbf{t}_2 \end{aligned}$$

we see that writing $\mathfrak{r} = \text{tp}_{\text{qf}}(\bar{\mathbf{t}}, \emptyset, I)$ we have that

$$\{\varphi(x, \bar{a}_{\bar{t}}) : \bar{t} \in Q_{\mathfrak{r}}^J\}$$

is inconsistent, because the set of formulas

$$\{\varphi(x, \bar{a}_{\bar{t}_{\{1,2\}}}), \varphi(x, \bar{a}_{\bar{t}_{\{0,2\}}}), \varphi(x, \bar{a}_{\bar{t}_{\{0,1\}}})\}$$

is already inconsistent, recalling that $M \models R(a_{v_0}, a_{v_1}, a_{v_2})$ and

$$\begin{aligned} \varphi(x, \bar{a}_{\bar{t}_{\{1,2\}}}) &\vdash R(x, a_{v_1}, a_{v_2}) \\ \varphi(x, \bar{a}_{\bar{t}_{\{0,2\}}}) &\vdash R(x, a_{v_0}, a_{v_2}) \\ \varphi(x, \bar{a}_{\bar{t}_{\{0,1\}}}) &\vdash R(x, a_{v_0}, a_{v_1}). \end{aligned}$$

This completes the warm-up. \square

Claim 5.7. *For any $n > k \geq 2$, $T_{n,k}$ is $\mathbf{c}_{n,k}$ -unsuperstable.*

Proof. Now we fix $n > k \geq 2$ and let us prove that $T_{n,k}$ is $\mathbf{c}_{n,k}$ -unsuperstable. For this proof, $I = I_{\mathbf{c}_{n,k}}$ and $\mathcal{K} = \mathcal{K}_{\mathbf{c}_{n,k}}$. Recall the meaning of n, k : in $T_{n,k}$ the edge relation R has arity $k+1$, and the forbidden configuration has arity $n+1$.

To satisfy Definition 1.3 in this case, it would suffice to show that whenever we are given a finite $m < \omega$ and sets I_m, B_m, p_m such that I_m is a finite subset of I , B_m is a subset of the monster model of $T_{n,k}$, and p_m is a partial non-algebraic type over B_m , we can find $I_{m+1}, B_{m+1}, p_{m+1}$ such that I_{m+1} is finite and $I_m \subseteq I_{m+1} \subseteq I$, $B_{m+1} \supseteq B_m$ is a set of parameters in the monster model for $T_{n,k}$, and p_{m+1} is a partial non-algebraic type over B_{m+1} which extends p_m , and such that p_{m+1} $\mathbf{c}_{n,k}$ -divides over B_m .

Suppose then that I_m, B_m, p_m are given.

Choose distinct t_0, \dots, t_{n-1} from $I \setminus I_m$, so that $0 \leq i < j < n$ implies $t_i < t_j$ in I . Since we are in I , the element t_i is named by the predicate P_{t_i} , and since $t_i \neq t_j$ their corresponding predicates are necessarily distinct. Also, since we are in I , there are necessarily no instances of R on $\{t_0, \dots, t_{n-1}\}$ in I . Let $\bar{t} = \langle t_0, \dots, t_{n-1} \rangle$. Let $\mathfrak{r} = \text{tp}_{\text{qf}}(\bar{t}, \bar{I}_m, I)$.

Fix $J \supseteq I$ such that $J \in \mathcal{K}$ and J is \aleph_0 -saturated. Choose a sequence $\langle a_t : t \in J \rangle$ which is \mathcal{K} -indiscernible and such that

$$(13) \quad M \models R(a_{t_{i_0}}, \dots, a_{t_{i_k}}) \iff J \models R(t_{i_0}, \dots, t_{i_k}).$$

⁶recalling that tp_{qf} in J is determined by ordering, the predicates P_q , and the instances of R .

Now we will need some notation for two different but related things. First, recalling \bar{t} from the previous paragraph, $\bar{a}_{\bar{t}}$ is now well defined (it is the tuple $\langle a_{t_i} : i < n \rangle$). For $u \in [n]^\ell$, let $\bar{t} \upharpoonright u$ denote the ℓ -tuple of elements of J given by $\langle t_i : i \in u \rangle$, thus we let

$$\bar{a}_{\bar{t} \upharpoonright u} \text{ denote the } \ell\text{-tuple of elements of } M \text{ given by } \langle a_{t_i} : i \in u \rangle.$$

Informally, we select an appropriate ℓ -tuple from the sequence of length n . (We will mostly use $\ell = k$ or $\ell = k + 1$.) Second, if $\bar{v} = \langle v_0, \dots, v_{n-1} \rangle$ is any sequence from Q_τ^J , and $u \in [n]^k$, let

$$\begin{aligned} \bar{v}_u \text{ denote the } n\text{-tuple of elements of } J \text{ given by } \langle t'_i : i < n \rangle \\ \text{where } t'_i = v_i \text{ if } i \in u \text{ and } t'_i = t_i \text{ if } i \notin u. \end{aligned}$$

Informally, this is \bar{v} where the i th element is replaced by t_i if $i \notin u$.

The next step is to choose $\bar{v} \in Q_\tau^J$ (and \bar{v} will be fixed as this choice for the rest of the proof) so that:⁷

- (i) $\text{tp}_{\text{qf}}(\bar{t}, I_m, I) = \text{tp}_{\text{qf}}(\bar{v}_u, I_m, J)$ for all $u \in [n]^k$
- and (ii) $J \models R(\bar{v} \upharpoonright w)$ for all $w \in [n]^{k+1}$.

For example, it would suffice to choose $\langle v_i : i < n \rangle$ by induction on i as follows. Remember we assumed that $J \models t_0 < \dots < t_n$, with $J \models P_{t_i}(t_i)$. Choose v_i to be any element of J satisfying the following two conditions:

- (a) $J \models t_i < v_i < t_{i+1}$ and $P_{t_i}(v_i)$.
- (b) partition $[\{t_0, \dots, t_n\} \cup \{v_j : j < i\}]^k$ into two sets: $Y = [\{v_j : j < i\}]^k$, and X is the complement of Y (so members of X contain at least one t_ℓ). Ask that t_i satisfies: $\tau \cup \{t_i\} \in R^J$ for $\tau \in Y$, and $\tau \cup \{t_i\} \notin R^J$ for $\tau \in X$.

At each step, this choice is possible because these conditions will not produce an $(n+1)$ -clique, and J is \aleph_0 -saturated. Since quantifier-free type in J is determined by the ordering $<$, the predicates P_ℓ , and the edge relation R , this suffices.

Now in M , consider the formula

$$\varphi(x, \bar{a}_{\bar{t}}) = \bigwedge_{u \in [n]^k} R(x, \bar{a}_{\bar{t} \upharpoonright u}).$$

Since there are no instances of R on \bar{t} in J , there are correspondingly no instances of R on $\{a_{t_i} : i < n\}$ in M by equation (13), so this formula is consistent. However, the set of formulas

$$\{\varphi(x, \bar{a}_{\bar{t}'} : \bar{t}' \in Q_\tau^J\}$$

is inconsistent, because by condition (a) above, it includes the set of formulas

$$\{\varphi(x, \bar{a}_{\bar{v}_u} : u \in [n]^k\}$$

which is inconsistent as for each $u \in [n]^k$, $\varphi(x, \bar{a}_{\bar{v}_u}) \vdash R(x, \bar{a}_{\bar{v} \upharpoonright u})$, and also $M \models R(\bar{a}_{\bar{v} \upharpoonright w})$ for each $w \in [n]^{k+1}$ by condition (ii) and equation (13); together, these

⁷Notice that condition (ii) is legal as the forbidden configuration has size $n + 1$. Notice also that u has size k , and the edge relation has arity $k + 1$. Each \bar{v}_u has ‘most,’ i.e. all but k , of its elements from \bar{t} ; this will be more noticeable when $n \gg k$. So a priori, substituting in a few elements from \bar{v} into the sequence \bar{t} should not cause edges to appear.

would give us an $(n + 1)$ -clique which is forbidden. This is our desired instance of $\mathbf{c}_{n,k}$ -dividing.

To finish, let $I_{m+1} = I_m \cup \{t_i : i < n\}$. Let B_{m+1} be an elementary submodel of M which contains $B_m \cup \{a_t : t \in J\}$. Let $p_{m+1} = p_m \cup \{\varphi(x, \bar{a}_t)\}$. (Note that in our theory $T_{n,k}$, p_{m+1} is consistent simply because the parameters from $\{a_{t_i} : i < n\}$ are disjoint to B_m .)

This completes the proof. \square

Theorem 5.8. *For each $n > k \geq 2$, there is a countable context \mathbf{c} such that $T_{\mathbf{rg}}$ is \mathbf{c} -superstable but $T_{n,k}$ is \mathbf{c} -unsuperstable.*

Proof. Use $\mathbf{c}_{n,k}$ from Definition 5.3. By Claim 5.5, $T_{\mathbf{rg}}$ is \mathbf{c} -superstable. By Claim 5.7, $T_{n,k}$ is $\mathbf{c}_{n,k}$ -unsuperstable. \square

6. SOME QUESTIONS AND FUTURE DIRECTIONS

Problem 6.1. *Characterize the theories which have the same shearing as the random graph, that is, the theories T which satisfy: for every countable context \mathbf{c} , T is \mathbf{c} -unsuperstable if and only if \mathbf{c} has property \odot .*

Remark 6.2. *By inspection, the theories $T_{\mathbf{m}}$ from [12] appear to have the same shearing as the random graph. This suggests the class from Problem 6.1 is interesting, spanning classes in Keisler's order and the interpretability order.*

Problem 6.3. *Investigate the relation of shearing and dividing in "basic" non-simple theories such as T_{feq} , for example, investigate whether it is possible to characterize there the shearing which does not come from dividing.*

Problem 6.4. *The equivalence of shearing and dividing in stable theories is explained by the phenomenon of "weak definitions," see explanation and references in the proof of 3.10 above. The existence of weak definitions is a property of (T, φ) and of a context. Thus the methods of this paper open up the possibility of developing a theory of the relative strength of weak definitions, as contexts vary, in simple unstable theories.*

Remark 6.5. *The methods of this paper also represent an interesting interaction between elementary classes (the theories T) and a priori non-elementary classes (the index model classes \mathcal{K}).*

Remark 6.6. *Since, by the proofs above, shearing is strictly weaker than dividing, it necessarily fails some of the axioms for independence relations in simple theories [8]; it may be useful to sort out which hold and which do not.*

Question 6.7. *Is there an analogous axiomatic characterization of shearing?*

Remark 6.8. *The statement and proof of 3.10 implicitly raise the analogous question for NIP theories, i.e., it may be worth while to explicitly sort out what can be said about shearing for formulas without the independence property.*

REFERENCES

- [1] A. Ehrenfeucht and A. Mostowski. “Models of axiomatic theories admitting automorphisms.” *Fund. Math.* 43 (1956), 50–68.
- [2] R. Grossberg, J. Iovino, O. Lessmann, “A primer of simple theories.” *Arch. Math. Logic* 41 (2002), no. 6, 541–580.
- [3] V. Guingona, C. D. Hill, and L. Scow. “Characterizing model-theoretic dividing lines via collapse of generalized indiscernibles.” *Ann. Pure Appl. Logic* 168 (2017), no. 5, 1091–1111.
- [4] E. Hrushovski, “Pseudofinite fields and related structures.” *Model Theory and Applications, Quaderni di Matematica*, vol. 11, Aracne, Rome, 2002, pp. 151–212.
- [5] J. Hubička and J. Nešetřil. “All those Ramsey classes (Ramsey classes with closures and forbidden homomorphisms).” arXiv:1606.07979v4, to appear, *Advances in Math.*
- [6] I. Kaplan and N. Ramsey. “Transitivity of Kim-independence.” *Advances in Math.* 379 (2021).
- [7] A. S. Kechris, V. G. Pestov and S. Todorčević, “Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups.” *Geom. Funct. Anal.* 15 (2005), no. 1, 106–189.
- [8] B. Kim and A. Pillay, “Simple theories.” *APAL* 99 (1997) 149–164.
- [9] M. Malliaris, “Hypergraph sequences as a tool for saturation of ultrapowers.” *Journal of Symbolic Logic*, 77, 1 (2012) 195–223.
- [10] M. Malliaris and S. Shelah, “A new look at interpretability and saturation.” *Ann Pure Appl Logic* 170, 5 (2019) 642–671.
- [11] M. Malliaris and S. Shelah, “A separation theorem for simple theories.” *Trans. AMS*, to appear.
- [12] M. Malliaris and S. Shelah, “Keisler’s order is not simple (and simple theories may not be either).” *Advances in Math.*, to appear.
- [13] M. Malliaris and S. Shelah, “New simple theories from hypergraph sequences.” *Manuscript* 1206, 2021.
- [14] J. Nešetřil. “Ramsey classes and homogeneous structures.” *Combin. Probab. Comput.* 14 (2005), no. 1-2, 171–189.
- [15] J. Nešetřil and V. Rödl. “Ramsey classes of set systems.” *J. Comb. Th. A* 34 (1983), 183–201.
- [16] M. O. Rabin, “Universal groups of automorphisms of models.” In *Symposium on the Theory of Models*, proceedings of the 1963 International Symposium at Berkeley, J. W. Addison, L. Henkin, A. Tarski, eds. North-Holland Publ. Co., Amsterdam (1965).
- [17] L. Scow. “Characterization of NIP theories by ordered graph-indiscernibles,” *Annals of Pure and Applied Logic* 163 (2012), pp. 1624–1641.
- [18] S. Shelah, “Simple unstable theories.” *Ann. Math. Logic*, 19, 177–203 (1980).
- [19] S. Shelah, *Classification Theory and the number of non-isomorphic models*, North-Holland, rev. ed. 1990 (first edition, *Classification Theory*, published 1978).
- [20] Shelah, “General non structure theory.” Paper E59, version of January 2016. <http://shelah.logic.at/files/E59.pdf>.
- [21] S. Shelah, “Toward classifying unstable theories.” *Annals of Pure and Applied Logic* 80 (1996) 229–255.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVENUE, CHICAGO, IL 60637, USA

Email address: mem@math.uchicago.edu

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

Email address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>