

NORMALIZERS AND CENTRALIZERS OF SUBNORMAL SUBSYSTEMS OF FUSION SYSTEMS

ELLEN HENKE

ABSTRACT. Every saturated fusion system corresponds to a group-like structure called a regular locality. In this paper we introduce and study (regular) normalizers and centralizers of partial subnormal subgroups of regular localities. This is then used to show that there is a reasonable notion of normalizers and centralizers of subnormal subsystems of fusion systems.

1. INTRODUCTION

In this paper we introduce and study normalizers and centralizers of subnormal subsystems of saturated fusion systems. In doing so, we generalize and unify some concepts which were known before: Aschbacher defined centralizers of normal subsystems as well as normalizers and centralizers of components of fusion systems (cf. [2, Chapter 6] and [3, Sections 2.1 and 2.2]). The latter construction was generalized by Oliver [17] to define normalizers of products of components. All these concepts are special cases of the normalizers and centralizers introduced in this paper. However, the existence of centralizers of normal subsystems (and the concrete construction of such centralizers given in [11]) is used in the proof of our results.

Our approach uses the fact that there is an (essentially unique) regular locality associated to every saturated fusion system. Here regular localities are group-like structures introduced by Chermak. Recently, Chermak and the author of this paper [8] established a kind of dictionary which allows us to translate between concepts in saturated fusion systems and in regular localities. In particular, subnormal subsystems of saturated fusion systems correspond to partial subnormal subgroups of regular localities.

In the present paper, we investigate first “regular normalizers” and centralizers of partial subnormal subgroups of regular localities. Only afterwards, this is used to define and study normalizers and centralizers of subnormal subsystems. Along the way, it becomes automatically transparent how one can translate between regular localities and fusion systems with regards to the concepts we will be concerned with. Generally, some flexibility in moving between fusion systems and localities may lead to simplifications in Aschbacher’s program to give a new treatment of the classification of finite simple groups. Before we state our results on regular localities, we collect some definitions, notation and background results. The reader who is only interested in the fusion-theoretic results is invited to go immediately to Subsection 1.3.

1.1. Some background. Throughout this paper we write homomorphisms on the right hand side of the argument. The reader is referred e.g. to [8, Section 3] for a detailed summary of the required definitions and results concerning partial groups and localities. As in this reference and in Chermak’s original papers [5, 7], we write $\mathbf{W}(\mathcal{L})$ for the set of words in a set \mathcal{L} , \emptyset for the empty word, and $v_1 \circ v_2 \circ \dots \circ v_n$ for the concatenation of words $v_1, \dots, v_n \in \mathbf{W}(\mathcal{L})$. Recall that a partial group consists of a set \mathcal{L} , a “product” $\Pi: \mathbf{D} \rightarrow \mathcal{L}$ defined only on the words in some subset \mathbf{D} of $\mathbf{W}(\mathcal{L})$, and an involutory bijection $\mathcal{L} \rightarrow \mathcal{L}, f \mapsto f^{-1}$ called an “inversion map”, subject to certain group-like axioms. If \mathcal{L} is a partial group with product $\Pi: \mathbf{D} \rightarrow \mathcal{L}$ then, given $(x_1, \dots, x_n) \in \mathbf{D}$, we write also $x_1 x_2 \cdots x_n$ for $\Pi(x_1, \dots, x_n)$. Moreover, we have the following important definitions.

- A subset $\mathcal{H} \subseteq \mathcal{L}$ is called a *partial subgroup* of \mathcal{L} if $\Pi(w) \in \mathcal{H}$ for every $w \in \mathbf{W}(\mathcal{H}) \cap \mathbf{D}$ and $h^{-1} \in \mathcal{H}$ for every $h \in \mathcal{H}$.
- A partial subgroup \mathcal{H} of \mathcal{L} is called a *subgroup* of \mathcal{L} if $\mathbf{W}(\mathcal{H}) \subseteq \mathbf{D}$. Observe that every subgroup of \mathcal{L} forms an actual group. We call a subgroup \mathcal{H} of \mathcal{L} a *p*-subgroup if it is a *p*-group.
- Given $f \in \mathcal{L}$, we write $\mathbf{D}(f) := \{x \in \mathcal{L} : (f^{-1}, x, f) \in \mathbf{D}\}$ for the set of elements $x \in \mathcal{L}$ for which the *conjugate* $x^f := \Pi(f^{-1}, x, f)$ is defined. This gives us a conjugation map $c_f: \mathbf{D}(f) \rightarrow \mathcal{L}$ defined by $x \mapsto x^f$.
- We say that a partial subgroup \mathcal{N} of \mathcal{L} is a *partial normal subgroup* of \mathcal{L} (and write $\mathcal{N} \trianglelefteq \mathcal{L}$) if $x^f \in \mathcal{N}$ for every $f \in \mathcal{L}$ and every $x \in \mathbf{D}(f) \cap \mathcal{N}$.
- We call a partial subgroup \mathcal{H} of \mathcal{L} a *partial subnormal subgroup* of \mathcal{L} (and write $\mathcal{H} \trianglelefteq \trianglelefteq \mathcal{L}$) if there is a sequence $\mathcal{H}_0, \mathcal{H}_1, \dots, \mathcal{H}_k$ of partial subgroups of \mathcal{L} such that

$$\mathcal{H} = \mathcal{H}_0 \trianglelefteq \mathcal{H}_1 \trianglelefteq \dots \trianglelefteq \mathcal{H}_{k-1} \trianglelefteq \mathcal{H}_k = \mathcal{L}.$$

- If $f \in \mathcal{L}$ and $\mathcal{X} \subseteq \mathbf{D}(f)$ set $\mathcal{X}^f := \{x^f : x \in \mathcal{X}\}$. For every $\mathcal{X} \subseteq \mathcal{L}$, call

$$N_{\mathcal{L}}(\mathcal{X}) := \{f \in \mathcal{L} : \mathcal{X} \subseteq \mathbf{D}(f) \text{ and } \mathcal{X}^f = \mathcal{X}\}$$

the *normalizer* of \mathcal{X} in \mathcal{L} .

- For $\mathcal{X} \subseteq \mathcal{L}$, we call

$$C_{\mathcal{L}}(\mathcal{X}) := \{f \in \mathcal{L} : \mathcal{X} \subseteq \mathbf{D}(f), x^f = x \text{ for all } x \in \mathcal{X}\}$$

the *centralizer* of \mathcal{X} in \mathcal{L} .

- For any two subsets $\mathcal{X}, \mathcal{Y} \subseteq \mathcal{L}$, their *product* can be naturally defined by

$$\mathcal{X}\mathcal{Y} := \{\Pi(x, y) : x \in \mathcal{X}, y \in \mathcal{Y}, (x, y) \in \mathbf{D}\}.$$

A *locality* is a triple (\mathcal{L}, Δ, S) such that \mathcal{L} is a partial group, S is maximal among the *p*-subgroups of \mathcal{L} , and Δ is a set of subgroups of S subject to certain axioms, which imply in particular that $N_{\mathcal{L}}(P)$ is a subgroup of \mathcal{L} for every $P \in \Delta$. If $R \leq S$ and $f \in \mathcal{L}$ with $R \subseteq \mathbf{D}(f)$ and $R^f \subseteq S$, then R^f is a subgroup of S and $c_f|_R$ turns out to be an injective group homomorphism from R to R^f . Moreover, for every $f \in \mathcal{L}$, the subset

$$S_f := \{x \in S : x \in \mathbf{D}(f), x^f \in S\}$$

is a subgroup of S and indeed an element of Δ . In particular, $c_f|_{S_f}$ is an injective group homomorphism $S_f \rightarrow S$.

Definition 1.1. If \mathcal{H} is a partial subgroup of \mathcal{L} and $T = S \cap \mathcal{H}$, then $\mathcal{F}_T(\mathcal{H})$ denotes the fusion system over T generated by the maps of the form $c_h: R \rightarrow R^h$ where $h \in \mathcal{H}$ and $R \leq T$ with $R^h \leq T$. Equivalently, $\mathcal{F}_T(\mathcal{H})$ is generated by all the maps $c_h|_{S_h \cap \mathcal{H}}$ with $h \in \mathcal{H}$.

Notice that we have in particular $\mathcal{F}_S(\mathcal{L})$ defined for every locality (\mathcal{L}, Δ, S) . We say that (\mathcal{L}, Δ, S) is a locality *over* \mathcal{F} if $\mathcal{F} = \mathcal{F}_S(\mathcal{L})$.

Below we will consider situations in which we are given a partial subgroup \mathcal{X} of \mathcal{L} such that, for some *p*-subgroup S_0 and some set Γ of subgroups of S_0 , the triple $(\mathcal{X}, \Gamma, S_0)$ is a locality. Notice, that we may then also form the fusion system $\mathcal{F}_{S_0}(\mathcal{X})$ according to the definition above, even if S_0 is not a subgroup of S .

Chermak [6] defined a locality (\mathcal{L}, Δ, S) over a saturated fusion system \mathcal{F} to be *regular* if the set Δ equals a certain set $\delta(\mathcal{F})$ of subgroups of S and $N_{\mathcal{L}}(P)$ is a group of characteristic *p* for every $P \in \Delta$. Here a finite group G is *of characteristic p* if $C_G(O_p(G)) \leq O_p(G)$.

We will list now some theorems which will be crucial in our approach. The first one follows essentially from the existence and uniqueness of centric linking systems. For a detailed proof see e.g. [12, Theorem A] and [13, Lemma 10.4].

Background Theorem 1. *Let \mathcal{F} be a saturated fusion system. Then there exists a regular locality (\mathcal{L}, Δ, S) over \mathcal{F} , which is unique up to an isomorphism which restricts to the identity on S .*

The next theorem was first observed by Chermak [6, Theorem C]. A proof can also be found in [13, Theorem 10.16(e), Corollary 10.19].

Background Theorem 2. *Let (\mathcal{L}, Δ, S) be a regular locality and $\mathcal{H} \trianglelefteq \mathcal{L}$. Then $\mathcal{E} := \mathcal{F}_{S \cap \mathcal{H}}(\mathcal{H})$ is saturated and $(\mathcal{H}, \delta(\mathcal{E}), S \cap \mathcal{H})$ is a regular locality over \mathcal{E} . If $\mathcal{H} \trianglelefteq \mathcal{L}$, then $C_{\mathcal{L}}(\mathcal{H}) \trianglelefteq \mathcal{L}$.*

As a consequence of the above theorem, for every partial normal subgroup \mathcal{H} of \mathcal{L} , the centralizer $C_{\mathcal{L}}(\mathcal{H})$ can be given the structure of a regular locality.

Building on Background Theorem 2, it is easy to define a notion of components of regular localities. If (\mathcal{L}, Δ, S) is regular, then we write $\text{Comp}(\mathcal{L})$ for the set of components of \mathcal{L} . The product $E(\mathcal{L})$ of components of \mathcal{L} forms a partial normal subgroups of \mathcal{L} . Similarly, there is a natural notion of components of fusion systems introduced by Aschbacher [2, Chapter 9] such that there is a normal subsystem $E(\mathcal{F})$ of \mathcal{F} which is a central product of the components. Again, we write $\text{Comp}(\mathcal{F})$ for the set of components of \mathcal{F} .

Background Theorem 3 ([8, Proposition 7.1(a), Lemma 7.3(a),(b)]). *The map*

$$\hat{\Psi}: \{\mathcal{H}: \mathcal{H} \trianglelefteq \mathcal{L}\} \rightarrow \{\mathcal{E}: \mathcal{E} \trianglelefteq \mathcal{F}\}, \mathcal{H} \mapsto \mathcal{F}_{S \cap \mathcal{H}}(\mathcal{H})$$

is well-defined and a bijection. It restricts to a bijection from the set of partial normal subgroups of \mathcal{L} to the set of normal subsystems of \mathcal{F} , and to a bijection from $\text{Comp}(\mathcal{L})$ to $\text{Comp}(\mathcal{F})$. Moreover,

$$\hat{\Psi}(E(\mathcal{L})) = E(\mathcal{F}).$$

1.2. Normalizers and centralizers in regular localities. Suppose (\mathcal{L}, Δ, S) is a regular locality and $\mathcal{H} \trianglelefteq \mathcal{L}$. The normalizer $N_{\mathcal{L}}(\mathcal{H})$ as defined above is not the right subset to consider if we want to find a partial subgroup of \mathcal{L} in which \mathcal{H} is normal. Indeed, \mathcal{H} is in many cases not even contained in $N_{\mathcal{L}}(\mathcal{H})$. We will however find that there is another partial subgroup $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ of \mathcal{L} which one can also reasonably think of as a “normalizer of \mathcal{H} ”.

One can observe relatively easily that $\tilde{T} := S \cap E(\mathcal{L})$ is an element of Δ . In particular, $G := N_{\mathcal{L}}(\tilde{T})$ is a finite group of characteristic p . We show in Lemma 2.13 that G acts on \mathcal{L} via conjugation. More precisely, for every $g \in G$, we have $\mathcal{L} \subseteq \mathbf{D}(g)$ (i.e. conjugation by g is defined on all elements of \mathcal{L}), and this gives an action of the group G on the set \mathcal{L} . It turns also out that the conjugation map $c_g: \mathcal{L} \rightarrow \mathcal{L}$ is for every element $g \in G$ an automorphism of \mathcal{L} . As G acts on \mathcal{L} , the set-wise stabilizer

$$N_G(\mathcal{X}) := \{g \in G: \mathcal{X}^g = \mathcal{X}\}$$

of \mathcal{X} in the group G is for every subset \mathcal{X} of \mathcal{L} a subgroup of G . Set $N_S(\mathcal{X}) := S \cap N_G(\mathcal{X}) = S \cap N_{\mathcal{L}}(\mathcal{X})$.

If we consider now a partial subnormal subgroup \mathcal{H} of \mathcal{L} as above then, similarly as in finite groups, every component of \mathcal{L} is either contained in \mathcal{H} or in the centralizer of \mathcal{H} . Hence, it is reasonable to think that the product $E(\mathcal{L})$ of components should be contained in $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$. There is moreover a Frattini Argument for localities (cf. [7, Corollary 3.11]) which implies that

$$\mathcal{L} = E(\mathcal{L})N_{\mathcal{L}}(\tilde{T}) = E(\mathcal{L})G.$$

Therefore, the following definition seems natural.

Definition 1. *Let (\mathcal{L}, Δ, S) be a regular locality over \mathcal{F} , set $\tilde{T} := E(\mathcal{L}) \cap S$ and $G := N_{\mathcal{L}}(\tilde{T})$. For every partial subnormal subgroup \mathcal{H} of \mathcal{L} , the subset*

$$\mathbb{N}_{\mathcal{L}}(\mathcal{H}) := E(\mathcal{L})N_G(\mathcal{H})$$

is called the regular normalizer of \mathcal{H} in \mathcal{L} .

If $\mathcal{H} \trianglelefteq \mathcal{L}$, then $\mathcal{H}^g = \mathcal{H}$ for every $g \in G$. So we have in this case that $N_G(\mathcal{H}) = G$ and thus $\mathbb{N}_{\mathcal{L}}(\mathcal{H}) = E(\mathcal{L})G = \mathcal{L}$ as one would expect. In the general case, the properties of $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ are summarized in the following theorem.

Theorem A. *Let (\mathcal{L}, Δ, S) be a regular locality and $\mathcal{H} \trianglelefteq \mathcal{L}$. Then the regular normalizer $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ is a partial subgroup of \mathcal{L} . Moreover, setting*

$$\tilde{T} := E(\mathcal{L}) \cap S, \quad G := N_{\mathcal{L}}(\tilde{T}), \quad T := \mathcal{H} \cap S, \quad \text{and } \mathcal{E} := \mathcal{F}_T(\mathcal{H}),$$

the following hold for every $S_0 \in \text{Syl}_p(N_G(\mathcal{H}))$:

- (a) *There exists a set Γ of subgroups of S_0 such that $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \Gamma, S_0)$ is a regular locality. In particular, the fusion system $\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ is well-defined and saturated.*
- (b) *$\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. If S_0 is chosen such that $T \leq S_0$, then $T = \mathcal{H} \cap S_0$ and $\mathcal{E} \trianglelefteq \mathcal{F}_0$.*
- (c) *$\text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) = \text{Comp}(\mathcal{L})$, $E(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) = E(\mathcal{L})$ and $\tilde{T} = E(\mathcal{L}) \cap S_0$. In particular, $\text{Comp}(\mathcal{F}) = \text{Comp}(\mathcal{F}_0)$ and $E(\mathcal{F}) = E(\mathcal{F}_0)$.*
- (d) *$S \cap \mathbb{N}_{\mathcal{L}}(\mathcal{H}) = N_S(\mathcal{H})$.*
- (e) *Suppose \mathcal{X} is a partial subgroup of \mathcal{L} such that $\mathcal{H} \trianglelefteq \mathcal{X}$ and $(\mathcal{X}, \Gamma_{\mathcal{X}}, S_{\mathcal{X}})$ is a locality for some p -subgroup $S_{\mathcal{X}}$ of \mathcal{X} with $T \leq S_{\mathcal{X}}$, and some set $\Gamma_{\mathcal{X}}$ of subgroups of $S_{\mathcal{X}}$. Then $\mathcal{X} \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.*
- (f) *$C_{\mathcal{L}}(T) \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ and*

$$C_{\mathcal{L}}(\mathcal{H}) = C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\mathcal{H}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H}).$$

With the notation as above, Theorem A(a) says that $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ gives rise to a saturated fusion system, which is however not necessarily a subsystem of $\mathcal{F} := \mathcal{F}_S(\mathcal{L})$ (as S_0 might not be a subgroup of S). It follows from Background Theorem 2 and parts (a),(b),(f) of Theorem A that $C_{\mathcal{L}}(\mathcal{H})$ can be given the structure of a regular locality. Thus, $C_{\mathcal{L}}(\mathcal{H})$ gives also rise to a saturated fusion system, which again is not necessarily a subsystem of \mathcal{F} . One can see here that it might be an advantage to work with regular localities rather than with fusion systems in some contexts. Nevertheless, we want to have some more fusion-theoretic results in place as well. We therefore define normalizers and centralizers of subnormal subsystems of saturated fusion systems now.

Definition 2. *Let \mathcal{F} be a saturated fusion system over S and let \mathcal{E} be a subnormal subsystem of \mathcal{F} over $T \leq S$. Fix a regular locality (\mathcal{L}, Δ, S) over \mathcal{F} and $\mathcal{H} \trianglelefteq \mathcal{L}$ with $T = S \cap \mathcal{H}$ and $\mathcal{E} = \mathcal{F}_T(\mathcal{H})$ (which exist by Background Theorems 1 and 3). We call then*

$$N_{\mathcal{F}}(\mathcal{E}) := \mathcal{F}_{N_S(\mathcal{H})}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$$

the normalizer of \mathcal{E} in \mathcal{F} and

$$C_{\mathcal{F}}(\mathcal{E}) := \mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H})).$$

the centralizer of \mathcal{E} in \mathcal{F} .

Recall that a regular locality (\mathcal{L}, Δ, S) over \mathcal{F} is by Background Theorem 1 essentially unique. Moreover, for a given regular locality (\mathcal{L}, Δ, S) over \mathcal{F} , a partial subnormal subgroup \mathcal{H} as above is also unique by Background Theorem 3. This can be used to show that $N_{\mathcal{F}}(\mathcal{E})$ and $C_{\mathcal{F}}(\mathcal{E})$ do not depend on the choice of (\mathcal{L}, Δ, S) and \mathcal{H} . Alternatively, we prove some characterizations of $N_{\mathcal{F}}(\mathcal{E})$ and $C_{\mathcal{F}}(\mathcal{E})$ in terms of fusion systems, which imply that $N_{\mathcal{F}}(\mathcal{E})$ and $C_{\mathcal{F}}(\mathcal{E})$ are well-defined.

1.3. Normalizers and centralizers of subnormal subsystems. We will now investigate the properties of $N_{\mathcal{F}}(\mathcal{E})$ and $C_{\mathcal{F}}(\mathcal{E})$ as defined above. The reader is referred to [4, Chapter I] for an introduction to the theory of fusion systems. Throughout this subsection we assume the following hypothesis:

\mathcal{F} is a saturated fusion system over S and \mathcal{E} is a subnormal subsystem of \mathcal{F} over $T \leq S$.

We will need the following definitions.

Definition 3.

- If Q, R are p -groups, \mathcal{D} is a fusion system over R and $\varphi: R \rightarrow Q$ is an injective group homomorphism, then write \mathcal{D}^φ for the fusion system over $R\varphi$ with

$$\mathrm{Hom}_{\mathcal{D}^\varphi}(P\varphi, Q\varphi) = \{\varphi^{-1}\psi\varphi: \psi \in \mathrm{Hom}_{\mathcal{D}}(P, Q)\} \text{ for all } P, Q \leq R.$$

- By $\mathrm{Aut}(\mathcal{F})$ we denote the set of automorphisms α of S with $\mathcal{F}^\alpha = \mathcal{F}$.
- Set $N_S(\mathcal{E}) := \{s \in N_S(T): c_s|_T \in \mathrm{Aut}(\mathcal{E})\}$.
- Every element of $\mathcal{E}^\mathcal{F} := \{\mathcal{E}^\varphi: \varphi \in \mathrm{Hom}_{\mathcal{F}}(T, S)\}$ is called an \mathcal{F} -conjugate of \mathcal{E} .
- The subsystem \mathcal{E} is said to be fully \mathcal{F} -normalized, if

$$|N_S(\mathcal{E})| \geq |N_S(\tilde{\mathcal{E}})| \text{ for all } \tilde{\mathcal{E}} \in \mathcal{E}^\mathcal{F}.$$

- Similarly, \mathcal{E} is called fully \mathcal{F} -centralized if

$$|C_S(\mathcal{E})| \geq |C_S(\tilde{\mathcal{E}})| \text{ for all } \tilde{\mathcal{E}} \in \mathcal{E}^\mathcal{F}.$$

It is shown in Lemma 3.27 below that every \mathcal{F} -conjugate of \mathcal{E} is again a subnormal subsystem of \mathcal{F} . Note that there is always a fully \mathcal{F} -normalized \mathcal{F} -conjugate and a fully \mathcal{F} -centralized \mathcal{F} -conjugate of \mathcal{E} .

If \mathcal{F} is constrained, then \mathcal{F} has a model, i.e. a group of characteristic p realizing \mathcal{F} (cf. [4, Theorem III.5.10]). Moreover, if \mathcal{F} is constrained and G is a model for \mathcal{F} , then it is shown in [8, Lemma 2.9] that there exists a unique subnormal subgroup H of G with $T = H \cap S$ and $\mathcal{E} = \mathcal{F}_T(H)$. This is used to formulate the following proposition. It gives a characterization of $N_{\mathcal{F}}(\mathcal{E})$, which one could take alternatively as a definition of this subsystem.

Theorem B.

- $N_{\mathcal{F}}(\mathcal{E})$ is a subsystem of \mathcal{F} over $N_S(\mathcal{E})$.
- Fix $T_0, \tilde{T} \leq S$ such that

$$E(\mathcal{E}) \text{ is a fusion system over } T_0 \text{ and } E(\mathcal{F}) \text{ is a fusion system over } \tilde{T}.$$

Then $N_{\mathcal{F}}(\tilde{T})$ is constrained and $N_{\mathcal{E}}(T_0)$ is subnormal in $N_{\mathcal{F}}(\tilde{T})$.

- Choose a model G for $N_{\mathcal{F}}(\tilde{T})$ and $H \trianglelefteq G$ with $H \cap S = T$ and $\mathcal{F}_T(H) = N_{\mathcal{E}}(T_0)$. Then $N_S(H) = N_S(\mathcal{E})$ and

$$N_{\mathcal{F}}(\mathcal{E}) = \langle E(\mathcal{F})N_S(\mathcal{E}), \mathcal{F}_{N_S(H)}(N_G(H)) \rangle.$$

For the definition of $E(\mathcal{F})N_S(\mathcal{E}) := (E(\mathcal{F})N_S(\mathcal{E}))_{\mathcal{F}}$, which is used in part (c) of the theorem above, the reader is referred to Definition 2.7. Some important properties of $N_{\mathcal{F}}(\mathcal{E})$ are summarized in the following theorem.

Theorem C.

- \mathcal{E} is a subsystem of $N_{\mathcal{F}}(\mathcal{E})$, which is $N_{\mathcal{F}}(\mathcal{E})$ -invariant.
- Every saturated subsystem \mathcal{D} of \mathcal{F} with $\mathcal{E} \trianglelefteq \mathcal{D}$ is contained in $N_{\mathcal{F}}(\mathcal{E})$.
- If \mathcal{E} is fully \mathcal{F} -normalized, then $N_{\mathcal{F}}(\mathcal{E})$ is saturated, \mathcal{E} is normal in $N_{\mathcal{F}}(\mathcal{E})$, and

$$E(N_{\mathcal{F}}(\mathcal{E})) = E(\mathcal{F}).$$

- If $X, Y \leq N_S(\mathcal{E})$ and $\alpha \in \mathrm{Hom}_{\mathcal{F}}(X, Y)$ with $T \leq X \cap Y$ and $\varphi|_T \in \mathrm{Aut}(\mathcal{E})$, then φ is a morphism in $N_{\mathcal{F}}(\mathcal{E})$. In particular, $C_{\mathcal{F}}(T) = C_{N_{\mathcal{F}}(\mathcal{E})}(T)$.
- If \mathcal{F} is constrained, G is a model for \mathcal{F} and $H \trianglelefteq G$ with $T = S \cap H$ and $\mathcal{F}_T(H) = \mathcal{E}$, then $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{N_S(H)}(N_G(H))$.

If \mathcal{E} is fully \mathcal{F} -normalized, then notice that $N_{\mathcal{F}}(\mathcal{E})$ is by parts (b) and (c) of Theorem C the largest saturated subsystem of \mathcal{F} containing \mathcal{E} as a normal subsystem. In Lemma 3.31 we give some equivalent conditions for \mathcal{E} to be fully \mathcal{F} -normalized. In particular, given a regular locality (\mathcal{L}, Δ, S) and $\mathcal{H} \trianglelefteq \mathcal{L}$ with $\mathcal{H} \cap S = T$ and $\mathcal{F}_S(\mathcal{H}) = \mathcal{E}$, we show that $N_S(\mathcal{E}) = N_S(\mathcal{H})$; moreover,

setting $G = N_{\mathcal{L}}(E(\mathcal{L}) \cap S)$ as before, \mathcal{E} is fully \mathcal{F} -normalized if and only if $N_S(\mathcal{E}) = N_S(\mathcal{H})$ is a Sylow p -subgroup of $N_G(\mathcal{H})$.

When proving Theorem C, we obtain along the way a nice notion of a product subsystem $\mathcal{E}R = (\mathcal{E}R)_{\mathcal{F}}$ for every subgroup R of $N_S(\mathcal{E})$. The existence of such a product was known before in the case that \mathcal{E} is normal (cf. [2, Chapter 8] or [10]). For arbitrary \mathcal{E} , a precise description of the product subsystem $\mathcal{E}R = (\mathcal{E}R)_{\mathcal{F}}$ is given in Definition 2.7. We have the following theorem.

Theorem D. *Let $R \leq N_S(\mathcal{E})$. Then the following hold.*

- (a) *The subsystem $\mathcal{E}R = (\mathcal{E}R)_{\mathcal{F}}$ is saturated and the unique saturated subsystem \mathcal{D} of \mathcal{F} over TR with $O^p(\mathcal{D}) = O^p(\mathcal{E})$. Moreover, $\mathcal{E} \trianglelefteq \mathcal{E}R$.*
- (b) *If (\mathcal{L}, Δ, S) is a regular locality over \mathcal{F} and $\mathcal{H} \trianglelefteq \trianglelefteq \mathcal{L}$ with $T = S \cap \mathcal{H}$ and $\mathcal{F}_T(\mathcal{H}) = \mathcal{E}$, then $\mathcal{E}R = \mathcal{F}_{TR}(\mathcal{H}R)$.*

With the above results about the normalizer $N_{\mathcal{F}}(\mathcal{E})$ in place, it is also relatively easy to study the centralizer $C_{\mathcal{F}}(\mathcal{E})$. First of all, we can show that there is a reasonable notion of a centralizer of \mathcal{E} in S .

Theorem E. *The set $\{R \leq S : \mathcal{E} \subseteq C_{\mathcal{F}}(R)\}$ has (with respect to inclusion) a unique maximal member $C_S(\mathcal{E})$. Moreover, $C_{\mathcal{F}}(\mathcal{E})$ is a fusion system over $C_S(\mathcal{E})$.*

Next we want to list some of the properties of the centralizer of \mathcal{E} in \mathcal{F} . For that we will use the following definition.

Definition 4. *Given two saturated subsystems \mathcal{F}_1 and \mathcal{F}_2 of \mathcal{F} over S_1 and S_2 respectively, we say that \mathcal{F}_1 commutes with \mathcal{F}_2 , if $\mathcal{F}_i \subseteq C_{\mathcal{F}}(S_{3-i})$ for each $i = 1, 2$.*

If (\mathcal{L}, Δ, S) is a regular locality over \mathcal{F} and $\mathcal{H} \trianglelefteq \trianglelefteq \mathcal{L}$ with $S \cap \mathcal{H} = T$ and $\mathcal{E} = \mathcal{F}_T(\mathcal{H})$, then the subgroup $C_S(\mathcal{E})$ from above is actually equal to $C_S(\mathcal{H})$. Moreover, we show in Lemma 3.38 that \mathcal{E} is fully centralized if and only if $C_S(\mathcal{H})$ is a maximal p -subgroup of $C_{\mathcal{L}}(\mathcal{H})$.

Centralizers of normal subsystems were defined by Aschbacher [2, Chapter 6]. We show that the centralizer $C_{\mathcal{F}}(\mathcal{E})$ defined above coincides with the centralizer subsystem defined by Aschbacher if \mathcal{E} is normal in \mathcal{F} . An alternative construction of centralizers of normal subsystems was given in [11]. It allows us to give an easy description of $C_{\mathcal{F}}(\mathcal{E})$ if \mathcal{E} is fully centralized (cf. Theorem F(c) below).

Theorem F. (a) *$C_{\mathcal{F}}(\mathcal{E})$ is a subsystem of $N_{\mathcal{F}}(\mathcal{E})$ over $C_S(\mathcal{E})$. Moreover, $C_{\mathcal{F}}(\mathcal{E})$ commutes with \mathcal{E} , and every saturated subsystem of \mathcal{F} which commutes with \mathcal{E} is contained in $C_{\mathcal{F}}(\mathcal{E})$.*
 (b) *$C_{\mathcal{F}}(\mathcal{E})$ is weakly $N_{\mathcal{F}}(\mathcal{E})$ -invariant in the sense of Definition 2.2 below. If \mathcal{E} is fully \mathcal{F} -normalized, then $C_{\mathcal{F}}(\mathcal{E}) = C_{N_{\mathcal{F}}(\mathcal{E})}(\mathcal{E})$ is normal in $N_{\mathcal{F}}(\mathcal{E})$.*
 (c) *Suppose \mathcal{E} is fully centralized. Then $C_{\mathcal{F}}(\mathcal{E})$ is saturated and thus the largest saturated subsystem of \mathcal{F} that commutes with \mathcal{E} . Moreover,*

$$C_{\mathcal{F}}(\mathcal{E}) = \langle O^p(\text{Aut}_{C_{\mathcal{F}}(T)}(P)) : P \leq C_S(\mathcal{E}) \rangle_{C_S(\mathcal{E})}.$$

For more information on the centralizer $C_{\mathcal{F}}(\mathcal{E})$, the reader is referred to Subsection 3.5. It seems difficult to obtain a neat description of this subsystem if \mathcal{E} is not fully centralized. However, the subsystem $C_{\mathcal{F}}(\mathcal{E})$ is uniquely determined by the properties stated in Theorem F(c) and Lemma 3.43.

2. PRELIMINARIES

Throughout this section let \mathcal{F} be a fusion system over a p -group S .

We adapt the terminology and notation from [4, Chapter I], except that we write homomorphisms on the right hand side. We use moreover the following notation.

Notation 2.1.

- Given $T \leq S$, we write $\mathcal{F}|_T$ for the full subcategory of \mathcal{F} whose objects are all the subgroups of T .
- If T is any p -group and Φ a set of injective group homomorphisms between subgroups of T , then $\langle \Phi \rangle_T$ denotes the fusion system over T generated by Φ , i.e. the smallest fusion system over T containing every morphism in Φ .

2.1. Invariant subsystems. As in [4, Definition I.6.1], we say that a subsystem \mathcal{E} of \mathcal{F} over T is \mathcal{F} -invariant if T is strongly closed, $\mathcal{E}^\alpha = \mathcal{E}$ for every $\alpha \in \text{Aut}_{\mathcal{F}}(T)$ and the Frattini condition holds, i.e. for every $P \leq T$ and every $\varphi \in \text{Hom}_{\mathcal{F}}(P, T)$, there exists $\varphi_0 \in \text{Hom}_{\mathcal{E}}(P, T)$ and $\alpha \in \text{Aut}_{\mathcal{F}}(T)$ with $\varphi = \varphi_0 \alpha$. There is also a different notion of \mathcal{F} -invariant subsystems in the literature. To distinguish the two, we introduce the following definition.

Definition 2.2. We call a subsystem \mathcal{E} of \mathcal{F} over T *weakly \mathcal{F} -invariant*, if T is strongly closed and, for all $P \leq Q \leq T$, all $\varphi \in \text{Hom}_{\mathcal{E}}(P, Q)$ and all $\alpha \in \text{Hom}_{\mathcal{F}}(Q, T)$, we have

$$\varphi^\alpha := (\alpha|_P)^{-1} \varphi \alpha \in \text{Hom}_{\mathcal{E}}(P\alpha, Q\alpha).$$

If \mathcal{F} is saturated, it is shown in [4, Proposition I.6.4] that \mathcal{E} is \mathcal{F} -invariant if and only if it is weakly \mathcal{F} -invariant. In general, the two notions are different, but we have the following lemma.

Lemma 2.3. *Let \mathcal{F} be a fusion system over S and \mathcal{E} a subsystem over \mathcal{F} over $T \leq S$.*

- If \mathcal{E} is \mathcal{F} -invariant, then \mathcal{E} is weakly \mathcal{F} -invariant.*
- Suppose T is strongly \mathcal{F} -closed and $\mathcal{E}^\alpha = \mathcal{E}$ for every $\alpha \in \text{Aut}_{\mathcal{F}}(T)$. Then T is \mathcal{F} -invariant if and only if $\mathcal{F}|_T = \langle \mathcal{E}, \text{Aut}_{\mathcal{F}}(T) \rangle_T$.*

Proof. This is shown in the proof of [4, Proposition I.6.4] if \mathcal{F} is saturated. The arguments do not use the assumption that \mathcal{F} is saturated. \square

2.2. Commuting Subsystems. As in Definition 4, we say that two saturated subsystems \mathcal{F}_i over S_i ($i = 1, 2$) commute in \mathcal{F} if $\mathcal{F}_i \subseteq C_{\mathcal{F}}(S_{3-i})$. Notice that this definition makes also sense if \mathcal{F} is not saturated. If \mathcal{F}_1 and \mathcal{F}_2 commute then $[S_1, S_2] = 1$ and thus $S_1 S_2$ is a subgroup of S . We set in this case

$$\mathcal{F}_1 * \mathcal{F}_2 := \langle \varphi \in \text{Hom}_{\mathcal{F}}(P_1 P_2, S_1 S_2) : P_i \leq S_i, \varphi|_{P_i} \in \text{Hom}_{\mathcal{F}_i}(P_i, S_i) \text{ for each } i = 1, 2 \rangle_{S_1 S_2}.$$

The following lemma could be concluded from the results in [8], but for simplicity we write out part of the argument.

Lemma 2.4. *Let \mathcal{F}_1 and \mathcal{F}_2 be saturated subsystems of \mathcal{F} over S_1 and S_2 respectively such that \mathcal{F}_1 and \mathcal{F}_2 commute in \mathcal{F} . Then $\mathcal{D} := \mathcal{F}_1 * \mathcal{F}_2$ is a saturated subsystem of \mathcal{F} with $\mathcal{D}|_{S_i} = \mathcal{F}_i$, $\mathcal{F}_i \trianglelefteq \mathcal{D}$ and $\mathcal{F}_i \subseteq C_{\mathcal{F}}(S_{3-i})$ for each $i = 1, 2$.*

Proof. By [11, Proposition 3.3], $\mathcal{D} = \mathcal{F}_1 * \mathcal{F}_2$ is a saturated subsystem of \mathcal{F} . Moreover, by [11, Lemma 3.2], $S_1 \cap S_2 \leq Z(\mathcal{F}_1) \cap Z(\mathcal{F}_2)$.

Let $P_i \leq S_i$ and $\varphi \in \text{Hom}_{\mathcal{F}}(P_1 P_2, S_1 S_2)$ with $\varphi|_{P_i} \in \text{Hom}_{\mathcal{F}_i}(P_i, S_i)$ for $i = 1, 2$. Fix now $i \in \{1, 2\}$. Notice that $\hat{P}_i := (P_1 P_2) \cap S_i = P_i Z$, where $Z := P_{3-i} \cap S_i \leq S_1 \cap S_2 \leq Z(\mathcal{F}_1) \cap Z(\mathcal{F}_2)$. As $Z \leq P_{3-i} \cap Z(\mathcal{F}_{3-i})$ and $\varphi|_{P_{3-i}}$ is a morphism in \mathcal{F}_{3-i} , we have in particular $\varphi|_Z = \text{id}_Z$. As $Z \leq Z(\mathcal{F}_i)$ and $\varphi|_{P_i}$ is a morphism in \mathcal{F}_i , it follows that $\varphi|_{\hat{P}_i}$ is a morphism in \mathcal{F}_i . This property implies that S_i is strongly closed in \mathcal{D} and

$$\mathcal{D}|_{S_i} = \mathcal{F}_i.$$

In particular, \mathcal{F}_i is \mathcal{F} -invariant. By assumption, \mathcal{F}_i is saturated and thus weakly normal in \mathcal{D} .

As $\mathcal{F}_i \subseteq C_{\mathcal{F}}(S_{3-i})$, every morphism $\varphi_i \in \text{Hom}_{\mathcal{F}_i}(P_i, S_i)$ extends to a morphism

$$\varphi \in \text{Hom}_{\mathcal{F}}(P_i S_{3-i}, S_1 S_2)$$

with $\varphi|_{S_{3-i}} = \text{id}_{S_{3-i}}$. By definition of \mathcal{D} , φ is then a morphism in \mathcal{D} . This shows that $\mathcal{F}_i \subseteq C_{\mathcal{D}}(S_{3-i})$. Using this property for $P_i := S_i$, it follows moreover that the pair $(\mathcal{F}_i, \mathcal{D})$ satisfies the extension condition for normal subsystems (cf. [4, Definition I.6.1]). Hence, \mathcal{F}_i is normal in \mathcal{F} . \square

2.3. Two lemmas on fusion systems.

Lemma 2.5. *If \mathcal{E} is a subnormal subsystem of \mathcal{F} over $T \leq S$ and $R \in \mathcal{F}^{cr}$, then $R \cap T \in \mathcal{E}^{cr}$.*

Proof. If \mathcal{E} is normal, this is shown in [1, Lemma 1.20(d)] (or alternatively in [9, Lemma 4.6]). The result follows thus by induction on the subnormal length of \mathcal{E} . \square

Lemma 2.6. *Let \mathcal{E} be a normal subsystem of \mathcal{F} over $T \leq S$ with $E(\mathcal{E}) = E(\mathcal{F})$. If $P \leq S$ with $P \cap T \in \delta(\mathcal{E})$, then $P \in \delta(\mathcal{F})$.*

Proof. Since the set $\delta(\mathcal{E})$ is overgroup-closed in T , we have $(PO_p(\mathcal{F})) \cap T \in \delta(\mathcal{E})$. Hence, by [9, Lemma 4.9(a)], $PO_p(\mathcal{F}) \in \delta(\mathcal{F})$. Using Background Theorem 1, it follows now from [13, Lemma 10.6] that $P \in \delta(\mathcal{F})$. \square

2.4. Products of subnormal subsystems with p -groups. If \mathcal{F} is saturated and \mathcal{E} is a normal subsystem of \mathcal{F} over $T \leq S$, then a product subsystem $\mathcal{E}R$ was defined by Aschbacher [2, Chapter 8] for every subgroup $R \leq S$. A new construction of this product subsystems was given in [10], which we generalize in the following definition.

Definition 2.7. Let \mathcal{E} be a subsystem of \mathcal{F} over $T \leq S$.

- For $P \leq S$ set

$$A_{\mathcal{F}, \mathcal{E}}^{\circ}(P) := \langle \varphi \in \text{Aut}_{\mathcal{F}}(P) : \varphi \text{ } p'\text{-element, } [P, \varphi] \leq P \cap T, \varphi|_{P \cap T} \in \text{Aut}_{\mathcal{E}}(P \cap T) \rangle.$$

- If \mathcal{F} is saturated and \mathcal{E} is subnormal in \mathcal{F} , then for every $R \leq N_S(\mathcal{E})$ set

$$\mathcal{E}R := (\mathcal{E}R)_{\mathcal{F}} := \langle A_{\mathcal{F}, \mathcal{E}}^{\circ}(P) : P \leq TR, P \cap T \in \mathcal{E}^{cr} \rangle_{TR}.$$

For the remainder of this section let \mathcal{F} be a saturated fusion system over a p -group S .

Proposition 2.8. *Suppose \mathcal{E} is a normal subsystem of \mathcal{F} over $T \leq S$. Then $N_S(\mathcal{E}) = S$. Moreover, for every $R \leq S$, the subsystem $\mathcal{E}R = (\mathcal{E}R)_{\mathcal{F}}$ is saturated and*

$$\mathcal{E}R = \langle A_{\mathcal{F}, \mathcal{E}}^{\circ}(P) : P \leq TR, P \cap T \in \mathcal{E}^c \rangle_{TR}.$$

Moreover, $\mathcal{E} \trianglelefteq \mathcal{E}R$ and $\mathcal{E}R$ is the unique saturated subsystem \mathcal{D} of \mathcal{F} over TR with $O^p(\mathcal{D}) = O^p(\mathcal{E})$.

Proof. As $\mathcal{E} \trianglelefteq \mathcal{F}$, we have $T \trianglelefteq S$ and $\text{Aut}_S(T) \leq \text{Aut}_{\mathcal{F}}(T) \leq \text{Aut}(\mathcal{E})$. Hence, $N_S(\mathcal{E}) = S$. Set now

$$\mathcal{G} := \langle A_{\mathcal{F}, \mathcal{E}}^{\circ}(P) : P \leq TR, P \cap T \in \mathcal{E}^c \rangle_{TR}.$$

Then by [10, Theorem 1], \mathcal{G} is saturated and the unique saturated subsystem \mathcal{D} of \mathcal{F} over TR with $O^p(\mathcal{D}) = O^p(\mathcal{E})$. By [8, Lemma 2.33], $\mathcal{E} \trianglelefteq \mathcal{G}$. Hence, it remains to argue that \mathcal{G} equals $\mathcal{E}R$. Clearly, $\mathcal{E}R$ is contained in \mathcal{G} . As \mathcal{G} is saturated, it follows from Alperin's Fusion Theorem (cf. [4, Theorem I.3.6]) that \mathcal{G} is generated by all the automorphism groups $\text{Aut}_{\mathcal{G}}(P)$, where $P \in \mathcal{G}^{cr}$ is fully \mathcal{G} -normalized. For every $P \in \mathcal{G}^{cr}$, Lemma 2.5 gives $P \cap T \in \mathcal{E}^{cr}$, and [10, Lemma 4.7] implies therefore that $O^p(\text{Aut}_{\mathcal{G}}(P)) = A_{\mathcal{F}, \mathcal{E}}^{\circ}(P)$ is contained in $\mathcal{E}R$. If $P \in \mathcal{G}^{cr}$ is fully \mathcal{G} -normalized, then P is fully \mathcal{G} -automized and hence $\text{Aut}_{\mathcal{G}}(P) = \text{Aut}_{TR}(P)O^p(\text{Aut}_{\mathcal{G}}(P))$ is contained in $\mathcal{E}R$. This shows that $\mathcal{G} \subseteq \mathcal{E}R$ and thus $\mathcal{G} = \mathcal{E}R$ as required. \square

The following lemma can be seen as a Frattini argument for fusion systems.

Lemma 2.9. *Let \mathcal{E} a normal subsystem of \mathcal{F} over $T \leq S$. Then*

$$\mathcal{F} = \langle \mathcal{E}S, N_{\mathcal{F}}(T) \rangle.$$

More precisely, for every subgroup $P \leq S$ and every morphism $\varphi \in \text{Hom}_{\mathcal{F}}(P, S)$, there exists $\psi \in \text{Hom}_{\mathcal{E}P}(P, TP)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(T)}(P\psi, S)$ such that $\varphi = \psi\alpha$.

Proof. We use throughout that, for every subgroup $R \leq S$, the subsystem $\mathcal{E}R$ defined above coincides by Proposition 2.8 with the equally denoted subsystem defined in [10]. It was first shown by Aschbacher [3, 1.3.2] that $\mathcal{F} = \langle \mathcal{E}S, N_{\mathcal{F}}(T) \rangle$; a proof using localities is given in [9, Lemma 4.10]. We use the property $\mathcal{F} = \langle \mathcal{E}S, N_{\mathcal{F}}(T) \rangle$ now to prove the second part of the claim. Fix $P \leq S$ and $\varphi \in \text{Hom}_{\mathcal{F}}(P, S)$.

By the definition of $\mathcal{E}S = (\mathcal{E}S)_{\mathcal{F}}$, every morphism in $\mathcal{E}S$ is a product of morphisms each of which is either a restriction of a morphisms in $A_{\mathcal{F}, \mathcal{E}}^{\circ}(Q)$ for some $Q \leq S$ with $Q \cap T \in \mathcal{E}^{cr}$, or a morphism induced by conjugation with an element of S . Notice that morphisms induced by conjugation with elements of S are in particular morphisms in $N_{\mathcal{F}}(T)$. Since $\mathcal{F} = \langle \mathcal{E}S, N_{\mathcal{F}}(T) \rangle$, it follows that φ can be factorized as the product of morphisms, each of which is either in $N_{\mathcal{F}}(T)$ or a restriction of an automorphism in $A_{\mathcal{F}, \mathcal{E}}^{\circ}(Q)$ for some $Q \leq S$ with $Q \cap T \in \mathcal{E}^{cr}$. By induction on the length of such a product, we may assume that φ can be factorized as

$$\varphi = \psi\alpha\gamma_0$$

where $\psi \in \text{Hom}_{\mathcal{E}P}(P, TP)$, $\alpha \in \text{Hom}_{N_{\mathcal{F}}(T)}(P\psi, S)$ and either

- (i) γ_0 is a morphism in $N_{\mathcal{F}}(T)$; or
- (ii) γ_0 is a restriction of a morphism $\gamma \in A_{\mathcal{F}, \mathcal{E}}^{\circ}(Q)$ for some $Q \leq S$ with $Q \cap T \in \mathcal{E}^{cr}$ and $P\psi\alpha \leq Q$.

In case (i), $\alpha\gamma_0$ is a morphism in $N_{\mathcal{F}}(T)$ and the claim holds. Hence, we may assume that (ii) holds. It follows from the definition of $\mathcal{E}P = (\mathcal{E}P)_{\mathcal{F}}$ that $TP = T(P\psi)$. Moreover, α extends to $\hat{\alpha} \in \text{Hom}_{\mathcal{F}}(T(P\psi), T(P\psi\alpha))$ with $T\hat{\alpha} = T$. Replacing Q by $Q \cap T(P\psi\alpha)$ and γ by $\gamma|_{Q \cap T(P\psi\alpha)}$, we may assume that $Q \leq T(P\psi\alpha)$. Notice that

$$\delta := \hat{\alpha}\gamma\hat{\alpha}^{-1} \in A_{\mathcal{F}, \mathcal{E}}^{\circ}(Q\hat{\alpha}^{-1})$$

with

$$\varphi = \psi\delta|_{P\psi}\hat{\alpha}.$$

By [8, Lemma 2.6], we have $(Q\hat{\alpha}^{-1}) \cap T = (Q \cap T)\hat{\alpha}^{-1} \in \mathcal{E}^{cr}$. Moreover, $Q\hat{\alpha}^{-1} \leq T(P\psi) = TP$. Hence, δ is a morphism in $\mathcal{E}P$ and thus $\psi\delta|_{P\psi}$ is a morphism in $\mathcal{E}P$. As $\hat{\alpha}$ is a morphism in $N_{\mathcal{F}}(T)$ the assertion follows. \square

2.5. Some lemmas on localities.

Lemma 2.10. *Let (\mathcal{L}, Δ, S) be a locality, $\mathcal{N} \trianglelefteq \mathcal{L}$ and $T := \mathcal{N} \cap S$. Then for every $g \in \mathcal{L}$ there exist $n \in \mathcal{N}$ and $f \in N_{\mathcal{L}}(T)$ such that $(n, f) \in \mathbf{D}$, $g = nf$ and $S_g = S_{(n, f)}$.*

Proof. This is [9, Lemma 3.13]. \square

Lemma 2.11. *Let (\mathcal{L}, Δ, S) and $(\tilde{\mathcal{L}}, \tilde{\Delta}, \tilde{S})$ be localities, and suppose $\alpha: \mathcal{L} \rightarrow \tilde{\mathcal{L}}$ is an isomorphism of partial groups. Let moreover \mathcal{H} be a partial subgroup of \mathcal{L} . Then the following hold:*

- (a) *If $S\alpha = \tilde{S}$, then $(S \cap \mathcal{H})\alpha = \tilde{S} \cap (\mathcal{H}\alpha)$.*
- (b) *If $(S \cap \mathcal{H})\alpha = \tilde{S} \cap (\mathcal{H}\alpha)$, then $\alpha|_{S \cap \mathcal{H}}$ induces an isomorphism from $\mathcal{F}_{S \cap \mathcal{H}}(\mathcal{H})$ to $\mathcal{F}_{\tilde{S} \cap (\mathcal{H}\alpha)}(\mathcal{H}\alpha)$.*

Proof. As α is injective, $S\alpha = \tilde{S}$ implies $(S \cap \mathcal{H})\alpha = (S\alpha) \cap (\mathcal{H}\alpha) = \tilde{S} \cap (\mathcal{H}\alpha)$. So assume from now on just that $(S \cap \mathcal{H})\alpha = \tilde{S} \cap (\mathcal{H}\alpha)$. Let $h \in \mathcal{H}$. It is sufficient to show that

$$\alpha^{-1}(c_h|_{S_h \cap \mathcal{H}})\alpha = c_{h\alpha}|_{\tilde{S}_{h\alpha} \cap (\mathcal{H}\alpha)} \quad (2.1)$$

To prove this let $x \in S_h \cap \mathcal{H}$, i.e. $x \in \mathbf{D}(h)$ and $x^h \in S$. As α is a homomorphism of partial groups, we have then $x\alpha \in \tilde{\mathbf{D}}(h\alpha)$ (where $\tilde{\mathbf{D}}$ denotes the domain of the product on $\tilde{\mathcal{L}}$). Moreover, since $x^h \in S \cap \mathcal{H}$, we have

$$(x\alpha)^{h\alpha} = (x^h)\alpha \in (S \cap \mathcal{H})\alpha = \tilde{S} \cap (\mathcal{H}\alpha).$$

In particular, as $x\alpha \in (S \cap \mathcal{H})\alpha = \tilde{S} \cap (\mathcal{H}\alpha)$, it follows $x\alpha \in \tilde{S}_{h\alpha}$, which proves

$$(S_h \cap \mathcal{H})\alpha \leq \tilde{S}_{h\alpha} \cap (\mathcal{H}\alpha).$$

It follows from [14, Lemma 2.18] that α^{-1} is also an isomorphism of partial groups. Hence, using the property we just proved with α^{-1} in place of α , we can conclude that $(\tilde{S}_{h\alpha} \cap (\mathcal{H}\alpha))\alpha^{-1} \leq S_h \cap \mathcal{H}$ and thus $\tilde{S}_{h\alpha} \cap (\mathcal{H}\alpha) = (S_h \cap \mathcal{H})\alpha$. As $(x\alpha)^{h\alpha} = (x^h)\alpha = (x\alpha)(\alpha^{-1}c_h\alpha)$, property (2.1) follows. \square

Corollary 2.12. *Let (\mathcal{L}, Δ, S) be a regular locality, $\mathcal{H} \trianglelefteq \mathcal{L}$ and $\alpha \in \text{Aut}(\mathcal{L})$ with $(\mathcal{H} \cap S)\alpha \leq S$. Then $(\mathcal{H} \cap S)\alpha = (\mathcal{H}\alpha) \cap S$ and α induces an isomorphism from $\mathcal{F}_{S \cap \mathcal{H}}(\mathcal{H})$ to $\mathcal{F}_{S \cap (\mathcal{H}\alpha)}(\mathcal{H}\alpha)$.*

Proof. It is a consequence of Background Theorem 2 and the definition of a locality that $S \cap \mathcal{H}$ is a maximal p -subgroup of \mathcal{H} . Notice that $\alpha \in \text{Aut}(\mathcal{L})$ induces an isomorphism from \mathcal{H} to $\mathcal{H}\alpha$. Hence, $(S \cap \mathcal{H})\alpha$ is a maximal p -subgroup of $\mathcal{H}\alpha$. As $(S \cap \mathcal{H})\alpha \leq S \cap (\mathcal{H}\alpha)$ by assumption, it follows $(S \cap \mathcal{H})\alpha = S \cap (\mathcal{H}\alpha)$. The assertion follows now from Lemma 2.11(b). \square

A linking locality is a locality (\mathcal{L}, Δ, S) such that $\mathcal{F}_S(\mathcal{L})$ is saturated, $\mathcal{F}_S(\mathcal{L})^{cr} \subseteq \Delta$ and $N_{\mathcal{L}}(P)$ is of characteristic p for every $P \in \Delta$. If \mathcal{F} is saturated, then the set \mathcal{F}^s of *subcentric subgroups* is the largest possible object set of a linking locality. It consists of the subgroups $P \leq S$ such that, for every fully \mathcal{F} -normalized \mathcal{F} -conjugate Q of P , the normalizer $N_{\mathcal{F}}(Q)$ is constrained (cf. [12, Proposition 3.1]).

For every saturated fusion system \mathcal{F} , the set $\delta(\mathcal{F})$ contains \mathcal{F}^{cr} (see e.g. [13, Lemma 10.4]). Hence, every regular locality is a linking locality. For every linking locality (\mathcal{L}, Δ, S) , there is the *generalized Fitting subgroup* $F^*(\mathcal{L})$ defined (cf. [13, Definition 3]).

We set $Z(\mathcal{L}) = C_{\mathcal{L}}(\mathcal{L})$ for every partial group \mathcal{L} . A linking locality (\mathcal{L}, Δ, S) is called *quasisimple* if $\mathcal{L} = O^p(\mathcal{L})$ (i.e. $\mathcal{L} \neq \mathcal{N}S$ for every proper partial normal subgroup \mathcal{N} of \mathcal{L}) and $\mathcal{L}/Z(\mathcal{L})$ is simple (i.e. $\mathcal{L}/Z(\mathcal{L})$ has precisely two partial normal subgroups).

Suppose now that (\mathcal{L}, Δ, S) is a regular locality. By Background Theorem 2, a partial subnormal subgroup \mathcal{K} of \mathcal{L} can be regarded as a regular locality. It is called a *component* if it is quasisimple. As mentioned already in the introduction, the product $E(\mathcal{L})$ of components of \mathcal{L} forms a partial normal subgroup of \mathcal{L} . The *generalized Fitting subgroup* $F^*(\mathcal{L})$ equals $E(\mathcal{L})O_p(\mathcal{L})$ (cf. [13, Theorem 4(a),(b),(c)]).

Lemma 2.13. *Let (\mathcal{L}, Δ, S) be a regular locality. Set*

$$\tilde{T} := E(\mathcal{L}) \cap S \text{ and } T^* := F^*(\mathcal{L}) \cap S.$$

Then \tilde{T} and T^ are elements of Δ , and $G = N_{\mathcal{L}}(\tilde{T}) = N_{\mathcal{L}}(T^*)$. Moreover, $\mathcal{L} \subseteq \mathbf{D}(f)$ for every $f \in G$, and G acts on \mathcal{L} via conjugation.*

Proof. By [13, Lemma 11.9], we have $E(\mathcal{L}) \trianglelefteq \mathcal{L}$ and $F^*(\mathcal{L}) = \tilde{T}O_p(\mathcal{L})$. Therefore $G = N_{\mathcal{L}}(\tilde{T}) = N_{\mathcal{L}}(T^*)$. By [13, Theorem 2], $F^*(\mathcal{L})$ is centric in \mathcal{L} . Hence, by [13, Lemma 10.11(c)], we have

$$\mathcal{L} \subseteq \mathbf{D}(f) \text{ and } c_f \in \text{Aut}(\mathcal{L})$$

for all $f \in N_{\mathcal{L}}(T^*)$. Notice that $x^1 = x$ for all $x \in \mathcal{L}$. Moreover, if $f, g \in N_{\mathcal{L}}(T^*)$, we have $u = (g^{-1}, f^{-1}, x, f, g) \in \mathbf{D}$ via $(S_x \cap T^*)^{fg}$. Hence, $x^{fg} = \Pi((fg)^{-1}, x, (fg)) = \Pi(u) = (x^f)^g$. Thus, $N_{\mathcal{L}}(T^*)$ acts on \mathcal{L} via conjugation. \square

2.6. Some group theoretic lemmas. Throughout this section let G be a finite group.

Lemma 2.14. *If $H \trianglelefteq G$, then $O_p(G) \leq N_G(O^p(H))$.*

Proof. Let G be a minimal counterexample. Notice that $H \neq G$ as otherwise the assertion clearly holds. As $H \trianglelefteq G$, there exists thus $N \trianglelefteq G$ with $H \trianglelefteq N$ and $N \neq G$. Then N is not a counterexample and hence $[O^p(H), O_p(N)] \leq O^p(H)$. Notice that

$$[O_p(G), H] \leq [O_p(G), N] \leq O_p(G) \cap N = O_p(N).$$

Hence, by a property of coprime action (cf. [15, 8.2.7(b)]), we have

$$[O_p(G), O^p(H)] = [O_p(G), O^p(H), O^p(H)] \leq [O_p(N), O^p(H)] \leq O^p(H)$$

and so the assertion holds. \square

Recall from the introduction that G is of *characteristic p* if $C_G(O_p(G)) \leq O_p(G)$.

Lemma 2.15. *Let G be a group of characteristic p . Then the following hold:*

- (a) $N_G(P)$ is of characteristic p for all non-trivial p -subgroups $P \leq G$.
- (b) Every subnormal subgroup of G is of characteristic p .
- (c) If $H \leq G$ with $O_p(G) \leq H$, then H is of characteristic p .

Proof. See [16, Lemma 1.2(a),(c)] for parts (a) and (b). If H is as in (c), then $O_p(G) \leq O_p(H)$ and hence $C_H(O_p(H)) \leq C_G(O_p(G)) \leq O_p(G) \leq O_p(H)$. \square

Lemma 2.16. *Let G be of characteristic p and $H \trianglelefteq G$. Then $N_G(H)$ is of characteristic p .*

Proof. As $H \trianglelefteq G$, it follows from Lemma 2.15(b) that H is of characteristic p . By Lemma 2.14, we have $O_p(G) \leq N_G(O^p(H))$. Hence, $N_G(O^p(H))$ has characteristic p by Lemma 2.15(c). Notice that $N_G(H) \leq N_G(O^p(H))$ as $O^p(H)$ is characteristic in H . Moreover, $H \trianglelefteq N_G(O^p(H))$. Hence, replacing G by $N_G(O^p(H))$, we can and will assume from now on that $O^p(H) \trianglelefteq G$.

Let $T \in \text{Syl}_p(H)$. Since $H \trianglelefteq N_G(H)$, it is a consequence of [9, Lemma 4.3] that $N_G(H)$ is of characteristic p if and only if $N_{N_G(H)}(T)$ is of characteristic p . However, as $O^p(H) \trianglelefteq G$ and $H = O^p(H)T$, we have $N_{N_G(H)}(T) = N_G(T)$. By Lemma 2.15(a), $N_G(T)$ has characteristic p . \square

Lemma 2.17. *Let \mathcal{F} be a constrained fusion system. Suppose G and \tilde{G} are models for \mathcal{F} . Let moreover $H \trianglelefteq G$ and $\tilde{H} \trianglelefteq \tilde{G}$ such that $\mathcal{F}_{S \cap H}(H) = \mathcal{F}_{S \cap \tilde{H}}(\tilde{H})$. Then*

$$\mathcal{F}_{N_S(H)}(N_G(H)) = \mathcal{F}_{N_S(\tilde{H})}(N_G(\tilde{H})).$$

Proof. As G and \tilde{G} are models, it follows from the [4, Theorem III.5.10] that there exists an isomorphism $\alpha: G \rightarrow \tilde{G}$ which restricts to the identity on S . If Δ is the set of all subgroups of S , then (G, Δ, S) and (\tilde{G}, Δ, S) are localities. Hence, it is a special case of Lemma 2.11 that

$$S \cap X = S \cap (X\alpha) \text{ and } \mathcal{F}_{S \cap X}(X) = \mathcal{F}_{S \cap X}(X)^\alpha = \mathcal{F}_{S \cap (X\alpha)}(X\alpha) \text{ for all } X \leq S. \quad (2.2)$$

Notice that $H\alpha \trianglelefteq \tilde{G}$. Moreover (2.2) together with our assumption gives $\mathcal{F}_{S \cap \tilde{H}}(\tilde{H}) = \mathcal{F}_{S \cap H}(H) = \mathcal{F}_{S \cap (H\alpha)}(H\alpha)$. By [8, Lemma 2.9], \tilde{H} is the unique subnormal subgroup of \tilde{G} realizing $\mathcal{F}_{S \cap \tilde{H}}(\tilde{H})$. Hence, $\tilde{H} = H\alpha$. It follows $N_G(H)\alpha = N_{\tilde{G}}(\tilde{H})$. Hence, using (2.2) now for $X = N_G(H)$, we obtain $\mathcal{F}_{N_S(H)}(N_G(H)) = \mathcal{F}_{N_S(\tilde{H})}(N_G(\tilde{H}))$. \square

3. NORMALIZERS AND CENTRALIZERS

Throughout this section we assume the following hypothesis.

Hypothesis 3.1. *Let (\mathcal{L}, Δ, S) be a regular locality and $\mathcal{H} \trianglelefteq \trianglelefteq \mathcal{L}$. Set*

$$\begin{aligned} \tilde{T} &:= E(\mathcal{L}) \cap S, \quad G := N_{\mathcal{L}}(\tilde{T}), \\ T &:= \mathcal{H} \cap S, \quad \mathcal{E} := \mathcal{F}_T(\mathcal{H}), \quad T_0 := E(\mathcal{H}) \cap S, \\ \mathcal{M} &:= \left(\prod_{\mathcal{K} \in \text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H})} \mathcal{K} \right) \text{ and } T_0^\perp := \mathcal{M} \cap S. \end{aligned}$$

We use above and throughout this section that, by Background Theorem 2, \mathcal{E} is saturated and $(\mathcal{H}, \delta(\mathcal{E}), T)$ is a regular locality. In particular $E(\mathcal{H}) \trianglelefteq \mathcal{H}$ and thus T_0 is well-defined.

3.1. Some basic results.

- Lemma 3.2.** (a) *$E(\mathcal{H})$ and \mathcal{M} are partial normal subgroups of $E(\mathcal{L})$ with $E(\mathcal{L}) = E(\mathcal{H})\mathcal{M}$. Moreover, $\tilde{T} = T_0 T_0^\perp$, $\mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{M})$ and $\mathcal{M} \subseteq C_{\mathcal{L}}(\mathcal{H})$.*
(b) *We have $\text{Comp}(\mathcal{M}) = \text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H})$, i.e. $\text{Comp}(\mathcal{L})$ is the disjoint union of $\text{Comp}(\mathcal{M})$ and $\text{Comp}(\mathcal{H})$.*
(c) *$\mathcal{H}\mathcal{M}$ is a partial subnormal subgroup of \mathcal{L} with $\mathcal{H} \trianglelefteq \mathcal{H}\mathcal{M}$ and $\mathcal{M} \trianglelefteq \mathcal{H}\mathcal{M}$.*

Proof. (a,c) We refer in this proof to the notion of an (internal) central product, which was defined in [13, Definition 4.1]. However, we will not need to go into the details of the definition.

By [13, Remark 11.2(b)], we have $\text{Comp}(\mathcal{H}) \subseteq \text{Comp}(\mathcal{L})$. By [13, Lemma 4.5], forming the internal central product is associative. Hence, it follows from [13, Theorem 11.18(a)] that $E(\mathcal{L})$ is a central product of $E(\mathcal{H})$ and \mathcal{M} , and from [13, Theorem 11.18(c)] that $\mathcal{H}\mathcal{M}$ is subnormal in \mathcal{L} and a central product of \mathcal{H} and \mathcal{M} . In particular, by [13, Lemma 4.9], $E(\mathcal{H})$ and \mathcal{M} are normal in $E(\mathcal{L}) = E(\mathcal{H})\mathcal{M}$, and \mathcal{H} and \mathcal{M} are normal in $\mathcal{H}\mathcal{M}$. It is moreover a consequence of [13, Lemma 4.8] that $\mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{M})$ and $\mathcal{M} \subseteq C_{\mathcal{L}}(\mathcal{H})$.

By Background Theorem 2, $E(\mathcal{L})$ can be regarded as a regular locality with Sylow subgroup \tilde{T} . Hence, it follows from [7, Theorem 5.1] (applied with $E(\mathcal{L})$ in place of \mathcal{L}) that $\tilde{T} = (E(\mathcal{H}) \cap \tilde{T})(\mathcal{M} \cap \tilde{T}) = T_0 T_0^\perp$.

(b) Part (a) yields that $\mathcal{M} \trianglelefteq E(\mathcal{L}) \trianglelefteq \mathcal{L}$ is subnormal in \mathcal{L} . In particular, $\text{Comp}(\mathcal{M})$ is well-defined by Background Theorem 2. Using [13, Remark 11.2], one sees that $\text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H}) \subseteq \text{Comp}(\mathcal{M})$ and $\text{Comp}(\mathcal{M}) \subseteq \text{Comp}(\mathcal{L})$. If $\mathcal{K} \in \text{Comp}(\mathcal{M}) \cap \text{Comp}(\mathcal{H})$, then part (a) yields that $\mathcal{K} = Z(\mathcal{K})$, which contradicts \mathcal{K} being quasisimple. Hence, $\text{Comp}(\mathcal{M}) \subseteq \text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H})$ and the assertion holds. \square

By Lemma 2.13, G is a subgroup of \mathcal{L} which acts on \mathcal{L} via conjugation. Thus, for every subset \mathcal{X} of \mathcal{L} , the set-wise stabilizer

$$N_G(\mathcal{X}) := \{f \in G : \mathcal{X}^f = \mathcal{X}\}$$

is a subgroup of G . In particular, $N_G(\mathcal{H})$ is a subgroup of G . As $S \leq G$, it follows that $N_S(\mathcal{H}) := S \cap N_G(\mathcal{H})$ is a subgroup of S .

Lemma 3.3. *The following hold:*

- (a) $N_{\mathcal{H}}(\tilde{T}) = N_{\mathcal{H}}(T_0) \trianglelefteq \trianglelefteq G$.
(b) $N_{\mathcal{F}}(\tilde{T})$ is constrained and G is a model for $N_{\mathcal{F}}(\tilde{T})$. Moreover, $N_{\mathcal{H}}(\tilde{T}) \trianglelefteq \trianglelefteq G$ is a model for $N_{\mathcal{E}}(T_0)$. In particular,

$$N_{\mathcal{E}}(T_0) \trianglelefteq \trianglelefteq N_{\mathcal{F}}(\tilde{T}).$$

- (c) $N_G(\mathcal{H}) = N_G(N_{\mathcal{H}}(\tilde{T})) = N_G(N_{\mathcal{H}}(T_0))$ is of characteristic p .
(d) $N_{E(\mathcal{L})}(\tilde{T}) \subseteq N_G(\mathcal{H})$.
(e) $N_G(\mathcal{H}) \subseteq N_G(E(\mathcal{H})) = N_G(T_0) = N_G(\mathcal{M}) = N_G(T_0^\perp)$.
(f) If $g \in G$ with $T_0^g \leq T$, then $g \in N_G(E(\mathcal{H}))$.

Proof. (a) Using Lemma 3.2(a), one can observe that $\mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{M}) \subseteq C_{\mathcal{L}}(T_0^\perp)$ and then $N_{\mathcal{H}}(T_0) \subseteq N_{\mathcal{H}}(T_0 T_0^\perp) = N_{\mathcal{H}}(\tilde{T})$. As $T_0 \leq \tilde{T}$, we have $T_0 = E(\mathcal{H}) \cap \tilde{T}$. Thus, $E(\mathcal{H}) \trianglelefteq \mathcal{H}$ implies that T_0 is normalized by $N_{\mathcal{H}}(\tilde{T})$. This proves (a).

(b) As $\tilde{T} \in \Delta$ by Lemma 2.13, the group G is of characteristic p . By [9, Lemma 3.10(b)], we have $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$, i.e. $N_{\mathcal{F}}(\tilde{T})$ is constrained and G is a model for $N_{\mathcal{F}}(\tilde{T})$. The same argument with $(\mathcal{H}, \delta(\mathcal{E}), T)$ and \mathcal{E} in place of (\mathcal{L}, Δ, S) and \mathcal{F} yields that $N_{\mathcal{H}}(T_0)$ is a model for $N_{\mathcal{E}}(T_0)$. By (a), $N_{\mathcal{H}}(T_0) = N_{\mathcal{H}}(\tilde{T})$. If $\mathcal{H} = \mathcal{H}_0 \trianglelefteq \mathcal{H}_1 \trianglelefteq \cdots \trianglelefteq \mathcal{H}_n = \mathcal{L}$ is a subnormal series of \mathcal{H} in \mathcal{L} , then $N_{\mathcal{H}}(\tilde{T}) = N_{\mathcal{H}_0}(\tilde{T}) \trianglelefteq N_{\mathcal{H}_1}(\tilde{T}) \trianglelefteq \cdots \trianglelefteq N_{\mathcal{H}_n}(\tilde{T}) = G$ is a subnormal series of $N_{\mathcal{H}}(\tilde{T})$ in G , i.e. $N_{\mathcal{H}}(\tilde{T}) \trianglelefteq G$. In particular, $N_{\mathcal{E}}(T_0) \trianglelefteq N_{\mathcal{F}}(\tilde{T})$ by [4, Proposition I.6.2]. This proves (b).

(f) Let $g \in G$ with $T_0^g \leq T$ and $\mathcal{K} \in \text{Comp}(\mathcal{H})$. Recall that $c_g \in \text{Aut}(\mathcal{L})$ by Lemma 2.13 and so $\mathcal{K}^g \in \text{Comp}(\mathcal{L})$ by [13, Lemma 11.12]. Notice moreover that $\mathcal{K} \cap S \leq T_0$ and thus $(\mathcal{K} \cap S)^g \leq T_0^g \leq T$. If \mathcal{K}^g were not a component of \mathcal{H} , then Lemma 3.2(a) would imply that $\mathcal{K}^g \subseteq \mathcal{M} \subseteq C_{\mathcal{L}}(\mathcal{H}) \subseteq C_{\mathcal{L}}(T)$ and thus $(\mathcal{K} \cap S)^g$ were abelian. As \mathcal{K} is quasisimple and $\mathcal{K} \cap S \cong (\mathcal{K} \cap S)^g$, this would contradict [13, Lemma 7.10]. Hence, $\mathcal{K}^g \in \text{Comp}(\mathcal{H})$. As \mathcal{K} was arbitrary, this yields $E(\mathcal{H})^g = E(\mathcal{H})$.

(c) Using parts (a) and (b), Lemma 2.16 yields that $N_G(N_{\mathcal{H}}(\tilde{T})) = N_G(N_{\mathcal{H}}(T_0))$ is of characteristic p . Hence, it remains only to show that $N_G(\mathcal{H}) = N_G(N_{\mathcal{H}}(\tilde{T}))$.

Clearly, $N_G(\mathcal{H}) \leq N_G(\mathcal{H} \cap G) = N_G(N_{\mathcal{H}}(\tilde{T}))$. Fix now $g \in N_G(N_{\mathcal{H}}(\tilde{T}))$. Then $T_0^g \leq N_{\mathcal{H}}(\tilde{T}) \cap \tilde{T} \leq \mathcal{H} \cap S = T$. Hence, by (f), we have $E(\mathcal{H})^g = E(\mathcal{H})$. It follows from the Frattini Lemma [7, Corollary 3.11] applied with $(\mathcal{H}, \delta(\mathcal{E}), T)$ in place of (\mathcal{L}, Δ, S) that $\mathcal{H} = E(\mathcal{H})N_{\mathcal{H}}(T_0) = E(\mathcal{H})N_{\mathcal{H}}(\tilde{T})$. Using again that $c_g \in \text{Aut}(\mathcal{L})$ by Lemma 2.13, we can conclude that $\mathcal{H}^g = E(\mathcal{H})^g N_{\mathcal{H}}(\tilde{T})^g = E(\mathcal{H})N_{\mathcal{H}}(\tilde{T}) = \mathcal{H}$. Hence, $N_G(N_{\mathcal{H}}(\tilde{T})) \leq N_G(\mathcal{H})$ and so equality holds.

(d) By Lemma 3.2(a), $E(\mathcal{H}) \trianglelefteq E(\mathcal{L}) = E(\mathcal{H})\mathcal{M}$ and $\mathcal{M} \subseteq C_{\mathcal{L}}(\mathcal{H}) \subseteq C_{\mathcal{L}}(T_0) \subseteq N_{\mathcal{L}}(T_0)$. In particular, $N_{E(\mathcal{L})}(\tilde{T})$ normalizes $T_0 = E(\mathcal{H}) \cap \tilde{T}$. Using the Dedekind argument for localities [7, Lemma 1.10] and part (a), we conclude that $N_{E(\mathcal{L})}(\tilde{T}) \subseteq N_{E(\mathcal{L})}(T_0) = N_{E(\mathcal{H})}(T_0)\mathcal{M} = N_{E(\mathcal{H})}(\tilde{T})\mathcal{M}$ and thus $N_{E(\mathcal{L})}(\tilde{T}) = N_{E(\mathcal{H})}(\tilde{T})N_{\mathcal{M}}(\tilde{T})$. As $N_{E(\mathcal{H})}(\tilde{T}) \subseteq N_{\mathcal{H}}(\tilde{T}) \subseteq N_G(N_{\mathcal{H}}(\tilde{T}))$ and $\mathcal{M} \subseteq C_{\mathcal{L}}(\mathcal{H})$, this implies $N_{E(\mathcal{L})}(\tilde{T}) \subseteq N_G(N_{\mathcal{H}}(\tilde{T}))$. Now (d) follows from (c).

(e) By Lemma 2.13, we have $c_g \in \text{Aut}(\mathcal{L})$ for every $g \in G$. Hence, for $g \in N_G(\mathcal{H})$, c_g restricts to an automorphism of \mathcal{H} . It follows now from [13, Lemma 11.12] that G acts on $\text{Comp}(\mathcal{L})$, and that $N_G(\mathcal{H})$ acts on $\text{Comp}(\mathcal{H})$. In particular, $N_G(\mathcal{H}) \leq N_G(E(\mathcal{H}))$. Similarly, $N_G(E(\mathcal{H}))$ acts on $\text{Comp}(\mathcal{H}) = \text{Comp}(E(\mathcal{H}))$ (cf. [13, Remark 11.2]). Hence, $N_G(E(\mathcal{H}))$ acts also on $\text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H})$, which implies $N_G(E(\mathcal{H})) \leq N_G(\mathcal{M})$. Notice that $N_G(E(\mathcal{H})) \leq N_G(E(\mathcal{H}) \cap \tilde{T}) = N_G(T_0)$. By (f), we have $N_G(T_0) \leq N_G(E(\mathcal{H}))$. We know thus that

$$N_G(E(\mathcal{H})) = N_G(T_0) \leq N_G(\mathcal{M}).$$

By Lemma 3.2(a),(b), $\mathcal{M} \trianglelefteq E(\mathcal{L}) \trianglelefteq \mathcal{L}$ is subnormal in \mathcal{L} , $\text{Comp}(\mathcal{H}) = \text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{M})$ and $\mathcal{M} = E(\mathcal{M})$. Hence, applying the same property with \mathcal{M} in place of \mathcal{H} , we obtain $N_G(\mathcal{M}) = N_G(T_0^\perp) \leq N_G(E(\mathcal{H}))$. This shows (e). \square

Lemma 3.4. *Let $f \in N_{\mathcal{L}}(T)$. Then $\mathcal{H} \subseteq \mathbf{D}(f)$. Moreover, there exist $n \in E(\mathcal{L})$ and $g \in G$ such that $(n, g) \in \mathbf{D}$, $f = ng$ and $\mathcal{H}^f = \mathcal{H}^g$.*

Proof. By Lemma 2.10, there exist $n \in E(\mathcal{L})$ and $g \in G$ such that $(n, g) \in \mathbf{D}$, $f = ng$ and $S_f = S_{(n,g)}$. As $f \in N_{\mathcal{L}}(T)$, we have then $T \leq S_f \leq S_n$. By Lemma 3.2(c), $\mathcal{H}\mathcal{M}$ is a partial subnormal subgroup of \mathcal{L} with $\mathcal{H} \trianglelefteq \mathcal{H}\mathcal{M}$. Notice that $E(\mathcal{L}) = E(\mathcal{H})\mathcal{M} \subseteq \mathcal{H}\mathcal{M}$ by Lemma 3.2(a). So $T^n \subseteq S \cap \mathcal{H} = T$ and thus $n \in N_{E(\mathcal{L})}(T) \subseteq N_{\mathcal{H}\mathcal{M}}(T)$. As $\mathcal{H}\mathcal{M} \trianglelefteq \mathcal{L}$, Background Theorem 2 yields that $\mathcal{H}\mathcal{M}$ can be given the structure of a regular locality with Sylow subgroup $S \cap (\mathcal{H}\mathcal{M})$. Hence, by [13, Theorem 10.16(f)] applied with $\mathcal{H}\mathcal{M}$ in place of \mathcal{L} , we have $N_{\mathcal{H}\mathcal{M}}(T) \subseteq N_{\mathcal{H}\mathcal{M}}(\mathcal{H})$.

In particular, $n \in N_{\mathcal{L}}(\mathcal{H})$. Fixing $h \in \mathcal{H}$, it follows $(n^{-1}, h, n) \in \mathbf{D}$ and thus $Q := S_{(n^{-1}, h, n)} \in \Delta$ by [7, Corollary 2.7]. As (\mathcal{L}, Δ, S) is regular, we have $\Delta = \delta(\mathcal{F})$ and [13, Lemma 10.2] yields $Q \cap F^*(\mathcal{L}) \in \Delta$. By Lemma 2.13, $g \in N_{\mathcal{L}}(F^*(\mathcal{L}) \cap S)$ and then $v = (g^{-1}, n^{-1}, h, n, g) \in \mathbf{D}$ via $(Q \cap F^*(\mathcal{L}))^g$. By [7, Lemma 1.4(f)], $g^{-1}n^{-1} = f^{-1}$. Hence, using the ‘‘associativity axiom’’ in partial groups, we can conclude that $(f^{-1}, h, f) \in \mathbf{D}$ and $h^f = \Pi(v) = (h^n)^g$. Since this holds for an arbitrary element $h \in \mathcal{H}$, this shows $\mathcal{H} \subseteq \mathbf{D}(f)$ and $\mathcal{H}^f = (\mathcal{H}^n)^g = \mathcal{H}^g$, where the last equality uses $n \in N_{\mathcal{L}}(\mathcal{H})$. This proves the assertion. \square

3.2. The regular normalizer and the centralizer in \mathcal{L} . Recall from the introduction that

$$\mathbb{N}_{\mathcal{L}}(\mathcal{H}) := E(\mathcal{L})N_G(\mathcal{H})$$

is called the *regular normalizer* of \mathcal{H} in \mathcal{L} .

In this subsection we will prove Theorem A and some related results. Before we do so let us consider the special case that \mathcal{F} is constrained.

Remark 3.5. Let \mathcal{F} be constrained. Then by [13, Lemma 11.6], $E(\mathcal{L}) = 1$ and \mathcal{L} is a model for \mathcal{F} . In particular, \mathcal{H} is a subnormal subgroup of \mathcal{L} , $\tilde{T} = 1$ and $\mathbb{N}_{\mathcal{L}}(\mathcal{H}) = N_{\mathcal{L}}(\mathcal{H})$.

Theorem 3.6. *The regular normalizer $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ is a partial subgroup of \mathcal{L} . Moreover, fixing $S_0 \in \text{Syl}_p(N_G(\mathcal{H}))$ and setting*

$$\Delta_0 := \{P \leq S_0 : P \cap \tilde{T} \in E(\mathcal{F})^s\},$$

the following hold:

- (a) *The triple $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \Delta_0, S_0)$ is a linking locality and the fusion system $\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ is saturated. Moreover, $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 and*

$$\delta(\mathcal{F}_0) := \{P \leq S_0 : PO_p(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) \in \Delta_0\}.$$

- (b) $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. *If $T \leq S_0$, then $T = \mathcal{H} \cap S_0$ and $\mathcal{E} \trianglelefteq \mathcal{F}_0$.*
(c) $\text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) = \text{Comp}(\mathcal{L})$, $E(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) = E(\mathcal{L})$ and $\tilde{T} = E(\mathcal{L}) \cap S_0$. *In particular, $\text{Comp}(\mathcal{F}) = \text{Comp}(\mathcal{F}_0)$ and $E(\mathcal{F}) = E(\mathcal{F}_0)$.*
(d) $E(\mathcal{H}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, and $\mathcal{M} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. *In particular, $E(\mathcal{E}) \trianglelefteq \mathcal{F}_0$ and $\mathcal{F}_{T_0^\perp}(\mathcal{M}) \trianglelefteq \mathcal{F}_0$.*
(e) $N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}) = N_G(\mathcal{H})$. *In particular, $S \cap \mathbb{N}_{\mathcal{L}}(\mathcal{H}) = N_S(\mathcal{H})$.*

Proof. (a) Set $T^* := F^*(\mathcal{L}) \cap S$ and recall from Lemma 2.13 that $G = N_{\mathcal{L}}(T^*)$. So $H := N_G(\mathcal{H}) \leq N_{\mathcal{L}}(T^*)$. By Lemma 3.3(d), we have

$$N_{E(\mathcal{L})}(T^*) = N_{E(\mathcal{L})}(\tilde{T}) \leq H.$$

In particular, \tilde{T} is contained in H and then a normal p -subgroup of H . Therefore $\tilde{T} \leq S_0$. By Lemma 3.3(c), H is of characteristic p . By Background Theorem 3, we have $\mathcal{F}_{S \cap E(\mathcal{L})}(E(\mathcal{L})) = E(\mathcal{F})$ and by [8, Lemma 7.22], $\delta(E(\mathcal{F})) = E(\mathcal{F})^s$. Hence,

$$\Delta_0 = \{P \leq S_0 : P \cap E(\mathcal{L}) \in \delta(\mathcal{F}_{S \cap E(\mathcal{L})}(E(\mathcal{L})))\}.$$

It follows now from [9, Theorem 6.1] (applied with $E(\mathcal{L})$ in place of \mathcal{N}) that $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ is a partial subgroups of \mathcal{L} , $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \Delta_0, S_0)$ is a linking locality, $\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ is saturated and $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 , where

$$\delta(\mathcal{F}_0) = \{P \leq S_0 : PO_p(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) \in \Delta_0\}^1.$$

This shows (a).

We will use now throughout that \mathcal{F}_0 is well-defined and saturated, and that $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 .

¹The reader might want to observe that the arguments to prove [9, Theorem 6.1] become actually easier in the case that \mathcal{N} equals $E(\mathcal{L})$.

(e) By the Dedekind Lemma, $N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}) = N_{E(\mathcal{L})}(\tilde{T})N_G(\mathcal{H})$. Hence, Lemma 3.3(d) implies that $N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}) = N_G(\mathcal{H})$. As $\tilde{T} \trianglelefteq S$, it follows in particular that $S \cap \mathbb{N}_{\mathcal{L}}(\mathcal{H}) = S \cap N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}) = S \cap N_G(\mathcal{H}) = N_S(\mathcal{H})$.

(b) If \mathcal{H} is a partial normal subgroup of \mathcal{L} , then $\mathcal{E} = \mathcal{F}_T(\mathcal{H}) \trianglelefteq \mathcal{F}_0$ by (a) and Background Theorem 2. Thus, it is sufficient to show $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

We prove first that \mathcal{H} is contained in $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$. It follows from the Frattini Lemma [7, Corollary 3.11] applied with $(\mathcal{H}, \delta(\mathcal{E}), T)$ in place of (\mathcal{L}, Δ, S) that $\mathcal{H} = E(\mathcal{H})N_{\mathcal{H}}(T_0)$. As $E(\mathcal{H}) \subseteq E(\mathcal{L})$ by Lemma 3.2(a) and $N_{\mathcal{H}}(T_0) = N_{\mathcal{H}}(\tilde{T}) \subseteq N_G(\mathcal{H})$ by Lemma 3.3(a), it follows $\mathcal{H} \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

As \mathcal{H} is a partial subgroup of \mathcal{L} , it is also a partial subgroup of $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$. Let $f \in \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ and $h \in \mathcal{H}$ such that $h \in \mathbf{D}(f)$. We need to show that $h^f \in \mathcal{H}$.

Clearly $E(\mathcal{L}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ as $E(\mathcal{L}) \trianglelefteq \mathcal{L}$ is contained in $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$. Hence, by Lemma 2.10 applied with $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ in place of (\mathcal{L}, Δ, S) , there exists $x \in E(\mathcal{L})$ and $g \in N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T})$ such that $(x, g) \in \mathbf{D}$, $f = xg$ and

$$(S_0)f = (S_0)_{(x,g)}.$$

It follows from [7, Lemma 1.4(f), Proposition 2.6(b)] that $(g^{-1}, x^{-1}) \in \mathbf{D} \cap \mathbf{W}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ via $(S_0)_{f^{-1}} = (S_0)_{f^{-1}}^f = (S_0)_{(x,g)}^{xg}$ and $f^{-1} = g^{-1}x^{-1}$. So $u := (g^{-1}, x^{-1}, h, x, g) \in \mathbf{D}$ via $(S_0)_{(f^{-1}, x, f)}$ and, by the axioms of a partial group, $h^f = \Pi(f^{-1}, h, f) = \Pi(u) = (h^x)^g$.

By Lemma 3.2(a),(b), $E(\mathcal{L}) = E(\mathcal{H})\mathcal{M} \subseteq \mathcal{H}\mathcal{M}$, and $\mathcal{H}\mathcal{M}$ is a partial subgroup of \mathcal{L} with $\mathcal{H} \trianglelefteq \mathcal{H}\mathcal{M}$. In particular, $h^x \in \mathcal{H}$. By (e), $g \in N_G(\mathcal{H})$. Thus, it follows $h^f = (h^x)^g \in \mathcal{H}$ as required.

(c) As every component of \mathcal{L} is subnormal in \mathcal{L} and contained in the partial subgroup $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ of \mathcal{L} , it is straightforward to see that every component of \mathcal{L} is subnormal in $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ (cf. [13, Lemma 3.7(a)]). Thus $\text{Comp}(\mathcal{L}) \subseteq \text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$.

Suppose now that there is a component $\mathcal{K} \in \text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) \setminus \text{Comp}(\mathcal{L})$. As $E(\mathcal{L}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, it follows from [13, Lemma 11.16] (applied with $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), E(\mathcal{L}))$ in place of $(\mathcal{L}, \mathcal{N})$ and combined with [13, Theorem 10.16(e)]) that

$$\mathcal{K} \subseteq C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(E(\mathcal{L})) \subseteq C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}) \subseteq N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}).$$

By (e) and Lemma 3.3(c), $N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T}) = N_G(\mathcal{H})$ is a group of characteristic p . As \mathcal{K} is a partial subnormal subgroup of $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$, it follows that \mathcal{K} is a partial subnormal subgroup of $N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T})$ and thus by Lemma 2.15(b) of characteristic p . However, by [13, Lemma 7.10], the fact that \mathcal{K} is a quasisimple locality implies $O_p(\mathcal{K}) = Z(\mathcal{K})$. This contradicts \mathcal{K} being of characteristic p and shows thus that

$$\text{Comp}(\mathcal{L}) = \text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})).$$

In particular, $E(\mathcal{L}) = E(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$. Using (a), it follows thus from Background Theorem 3 that $\text{Comp}(\mathcal{F}) = \text{Comp}(\mathcal{F}_0)$ and $E(\mathcal{F}) = E(\mathcal{F}_0)$.

(d) As $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, it follows from [13, Lemma 11.13] that $E(\mathcal{H}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. By Lemma 3.2(a), $\mathcal{M} \trianglelefteq E(\mathcal{L})$. Moreover, by (e) and Lemma 3.3(e), $N_{\mathbb{N}_{\mathcal{L}}(\tilde{T})}(\tilde{T}) = N_G(\mathcal{H})$ normalizes \mathcal{M} . As $E(\mathcal{L}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ with $S_0 \cap E(\mathcal{L}) = \tilde{T}$, it follow now from [7, Corollary 3.13] that $\mathcal{M} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

It is a consequence of Background Theorem 3 (applied with $(\mathcal{N}, \delta(\mathcal{E}), T)$ in place of (\mathcal{L}, Δ, S)) that $E(\mathcal{E}) = \mathcal{F}_{T_0}(E(\mathcal{H}))$. As $T_0 = E(\mathcal{H}) \cap \tilde{T} = E(\mathcal{H}) \cap S_0$ and $T_0^\perp = \mathcal{M} \cap \tilde{T} = \mathcal{M} \cap S_0$, Background Theorem 3 (applied with $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ in place of (\mathcal{L}, Δ, S)) gives now also $E(\mathcal{E}) \trianglelefteq \mathcal{F}_0$ and $\mathcal{F}_{T_0^\perp}(\mathcal{M}) \trianglelefteq \mathcal{F}_0$. \square

Lemma 3.7. *Let \mathcal{X} be a partial subgroup of \mathcal{L} such that $\mathcal{H} \trianglelefteq \mathcal{X}$ and $(\mathcal{X}, S_{\mathcal{X}}, \Gamma_{\mathcal{X}})$ is a locality for some p -subgroup $S_{\mathcal{X}}$ of \mathcal{X} with $T \leq S_{\mathcal{X}}$ and some set $\Gamma_{\mathcal{X}}$ of subgroups of $S_{\mathcal{X}}$. Then $\mathcal{X} \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.*

Proof. It is a consequence of Background Theorem 2 (and the definition of a locality) that T is a maximal p -subgroup of \mathcal{H} . As $T \leq S_{\mathcal{X}} \cap \mathcal{H}$, it follows thus that $T = S_{\mathcal{X}} \cap \mathcal{H}$. The Frattini argument for localities [7, Corollary 3.11] (applied with $(\mathcal{X}, S_{\mathcal{X}}, \Delta_{\mathcal{X}})$ in place of (\mathcal{L}, Δ, S)) gives therefore $\mathcal{X} = \mathcal{H}N_{\mathcal{X}}(T)$. By Theorem 3.6 $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ is a partial subgroup of \mathcal{L} with $\mathcal{H} \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. Thus it remains to show that $N_{\mathcal{X}}(T) \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

Let $f \in N_{\mathcal{X}}(T)$. By Lemma 3.4, we have $\mathcal{H} \subseteq \mathbf{D}(f)$ and there exist $n \in E(\mathcal{L})$ and $g \in G$ such that $(n, g) \in \mathbf{D}$, $f = ng$ and $\mathcal{H}^f = \mathcal{H}^g$. As $\mathcal{H} \trianglelefteq \mathcal{X}$, it follows $\mathcal{H}^g = \mathcal{H}^f = \mathcal{H}$ and thus $g \in N_G(\mathcal{H})$. Hence, $f = ng \in E(\mathcal{L})N_G(\mathcal{H}) = \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. This completes the proof. \square

Lemma 3.8. *Let $f \in N_G(T)$. Then $f \in N_G(\mathcal{H})$ if and only if $c_f|_T \in \text{Aut}(\mathcal{E})$. In particular,*

$$N_S(\mathcal{E}) = N_S(\mathcal{H}) = N_S(N_{\mathcal{H}}(\tilde{T})) = N_S(N_{\mathcal{E}}(T_0)).$$

Proof. Recall that $c_f \in \text{Aut}(\mathcal{L})$ by Lemma 2.13. In particular, as $\mathcal{H} \trianglelefteq \mathcal{L}$, we have $\mathcal{H}^f \trianglelefteq \mathcal{L}$. Moreover, setting $\alpha := c_f|_T$, it follows from Lemma 2.12 that $S \cap \mathcal{H}^f = T^f = T$ and

$$\mathcal{F}_T(\mathcal{H}^f) = \mathcal{E}^{\alpha}.$$

In particular, if $\mathcal{H}^f = \mathcal{H}$, then $\alpha \in \text{Aut}(\mathcal{E})$. On the other hand, if $\alpha \in \text{Aut}(\mathcal{E})$, then $\mathcal{F}_T(\mathcal{H}^f) = \mathcal{E}^{\alpha} = \mathcal{E} = \mathcal{F}_T(\mathcal{H})$. Since the map $\hat{\Psi}$ from Background Theorem 3 is injective, we see thus that $\alpha \in \text{Aut}(\mathcal{E})$ implies $\mathcal{H}^f = \mathcal{H}$. This shows $f \in N_G(\mathcal{H})$ if and only if $c_f|_T \in \text{Aut}(\mathcal{E})$. In particular,

$$N_S(\mathcal{E}) = N_S(\mathcal{H}). \quad (3.1)$$

By Lemma 3.3(b), G is a model for $N_{\mathcal{F}}(\tilde{T})$ and $N_{\mathcal{H}}(\tilde{T}) \trianglelefteq G$ is a model for $N_{\mathcal{E}}(T_0)$. It follows from Lemma 2.15(a) that $(G, \delta(N_{\mathcal{F}}(\tilde{T})), S)$ is a regular locality over $N_{\mathcal{F}}(\tilde{T})$. Thus, applying (3.1) with $(G, N_{\mathcal{H}}(\tilde{T}), N_{\mathcal{E}}(T_0))$ in place of $(\mathcal{L}, \mathcal{H}, \mathcal{E})$, we obtain $N_S(N_{\mathcal{H}}(\tilde{T})) = N_S(N_{\mathcal{E}}(T_0))$. Lemma 3.3(c) implies that $N_S(\mathcal{H}) = N_S(N_{\mathcal{H}}(\tilde{T}))$. Hence the assertion holds. \square

Lemma 3.9. (a) $C_{\mathcal{L}}(T) \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

(b) $C_{\mathcal{L}}(\mathcal{H}) = C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\mathcal{H}) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

Proof. (a) As T is a p -subgroup of $N_G(\mathcal{H})$, we may choose $S_0 \in \text{Syl}_p(N_G(\mathcal{H}))$ with $T \leq S_0$. Set $\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$. By Theorem 3.6(a),(b), \mathcal{F}_0 is saturated and $\mathcal{E} \trianglelefteq \mathcal{F}_0$. Let $f \in C_{\mathcal{L}}(T)$. By Lemma 2.10, there exists $x \in E(\mathcal{L})$ and $g \in G$ such that $(x, g) \in \mathbf{D}$, $f = xg$ and $S_{(x,g)} = S_f$. Notice that $E(\mathcal{L}) \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, so $c_x|_T \in \text{Aut}_{\mathcal{F}_0}(T) \leq \text{Aut}(\mathcal{E})$. As $(c_x|_T) \circ (c_g|_T) = c_f|_T = \text{id} \in \text{Aut}(\mathcal{E})$, it follows $c_g|_T \in \text{Aut}(\mathcal{E})$. Hence, Lemma 3.8 yields $g \in N_G(\mathcal{H})$ and thus $f = xg \in E(\mathcal{L})N_G(\mathcal{H}) = \mathbb{N}_{\mathcal{L}}(\mathcal{H})$.

(b) Using (a), we see that $C_{\mathcal{L}}(\mathcal{H}) \subseteq C_{\mathcal{L}}(T) \subseteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ and thus $C_{\mathcal{L}}(\mathcal{H}) = C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\mathcal{H})$. Recall that $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ forms a regular locality by Theorem 3.6(a). Hence, part (b) follows from Background Theorem 2 applied with $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ in place of \mathcal{L} . \square

Proof of Theorem A. Parts (a),(b),(c) follows from Theorem 3.6(a),(b),(c), and part (d) is stated in Theorem 3.6(e). Part (e) is stated in Lemma 3.7, and part (f) in Lemma 3.9. \square

3.3. The normalizer of \mathcal{E} . We continue to assume Hypothesis 3.1 and assume in addition for the remainder of this section the following hypothesis.

Hypothesis 3.10. *Let S_0 be a Sylow p -subgroup of $N_G(\mathcal{H})$ with $T \leq S_0$. Set*

$$\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})).$$

We will use throughout that, by Theorem 3.6, \mathcal{F}_0 is well-defined and saturated, $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 , $E(\mathcal{L}) = E(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, $\tilde{T} = E(\mathcal{L}) \cap S_0$, $E(\mathcal{F}) = E(\mathcal{F}_0) \trianglelefteq \mathcal{F}_0$, $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, $T = S_0 \cap \mathcal{H}$ and $\mathcal{E} \trianglelefteq \mathcal{F}_0$. We use moreover throughout that

$$N_S(\mathcal{E}) = N_S(\mathcal{H})$$

by Lemma 3.8. Recall that

$$N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{N_S(\mathcal{H})}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})),$$

so in particular, $N_{\mathcal{F}}(\mathcal{E})$ is a fusion system over $N_S(\mathcal{H}) = N_S(\mathcal{E})$.

In this subsection, we will prove Theorem B, Theorem C(a),(b),(d) and Theorem D, as well as some related results concerning in particular the connection between $N_{\mathcal{F}}(\mathcal{E})$ and \mathcal{F}_0 .

Lemma 3.11. *For every $R \leq N_S(\mathcal{E}) \cap S_0$, we have*

$$E(\mathcal{F})R := (E(\mathcal{F})R)_{\mathcal{F}} = \mathcal{F}_{\tilde{T}R}(E(\mathcal{L})R) = (E(\mathcal{F})R)_{\mathcal{F}_0}$$

and $(E(\mathcal{L})R, \delta(E(\mathcal{F})R), \tilde{T}R)$ is a regular locality over $E(\mathcal{F})R$. In particular, $E(\mathcal{F})R \subseteq N_{\mathcal{F}}(\mathcal{E})$.

Proof. By Background Theorem 3, we have $E(\mathcal{F}) = \mathcal{F}_{S \cap E(\mathcal{L})}(E(\mathcal{L})) = \mathcal{F}_{\tilde{T}}(E(\mathcal{L})) = \mathcal{F}_{S_0 \cap E(\mathcal{L})}(E(\mathcal{L}))$. Hence, by [8, Theorem D(a)] applied twice (once with \mathcal{L} and once with $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ in place of \mathcal{L}), we have

$$(E(\mathcal{F})R)_{\mathcal{F}} = \mathcal{F}_{\tilde{T}R}(E(\mathcal{L})R) = (E(\mathcal{F})R)_{\mathcal{F}_0}$$

and $(E(\mathcal{L})R, \delta(E(\mathcal{F})R), \tilde{T}R)$ is a regular locality over $E(\mathcal{F})R$. As $\tilde{T}R \leq N_S(\mathcal{E})$ and $E(\mathcal{L})R \leq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, it follows that $E(\mathcal{F})R \subseteq N_{\mathcal{F}}(\mathcal{E})$. \square

For every $R \leq N_S(\mathcal{E})$ we write from now on usually $E(\mathcal{F})R$ for $(E(\mathcal{F})R)_{\mathcal{F}}$.

Lemma 3.12. *Let $X \leq N_S(\mathcal{E})$ and $\varphi \in \text{Hom}_{\mathcal{F}}(X, S)$. Then there exist*

$$\psi \in \text{Hom}_{E(\mathcal{F})X}(X, X\tilde{T}) \subseteq \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(X, X\tilde{T})$$

and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(X\psi, S)$ such that $\varphi = \psi\alpha$. If $X \leq S_0$, then ψ is a morphism in \mathcal{F}_0 .

Proof. The existence of $\psi \in \text{Hom}_{E(\mathcal{F})X}(X, \tilde{T}X)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(X\psi, S)$ with $\varphi = \psi\alpha$ follows from Lemma 2.9 applied with $E(\mathcal{F})$ in place of \mathcal{E} . The assertion follows now from Lemma 3.11. \square

Lemma 3.13. *Set $\mathcal{B} := \mathcal{F}_{N_S(\mathcal{H})}(N_G(\mathcal{H}))$. Then the following hold:*

(a) *We have*

$$N_{\mathcal{F}}(\mathcal{E}) = \langle E(\mathcal{F})N_S(\mathcal{E}), \mathcal{B} \rangle.$$

If $N_S(\mathcal{E}) \leq S_0$, then also

$$N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \cap \mathcal{F}_0 = \mathcal{F}_0|_{N_S(\mathcal{E})}.$$

(b) *If $P \leq N_S(\mathcal{E})$ and $\varphi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(P, N_S(\mathcal{E}))$, then there exist*

$$\psi \in \text{Hom}_{E(\mathcal{F})P}(P, \tilde{T}P) \text{ and } \alpha \in \text{Hom}_{\mathcal{B}}(P\psi, N_S(\mathcal{E}))$$

such that $\varphi = \psi\alpha$.

(c) *\mathcal{E} is a $N_{\mathcal{F}}(\mathcal{E})$ -invariant subsystem of $N_{\mathcal{F}}(\mathcal{E})$.*

Proof. As $N_S(\mathcal{E}) = N_S(\mathcal{H})$ is a p -subgroup of $N_G(\mathcal{H})$, there exists always a Sylow p -subgroup of $N_G(\mathcal{H})$ containing $N_S(\mathcal{E})$. Hence, we may assume throughout the proof that $N_S(\mathcal{E}) \leq S_0$.

(a,b) As $N_S(\mathcal{E}) \leq S_0$ and $\mathcal{E} \trianglelefteq \mathcal{F}_0$, we have $S_0 \cap S = N_S(\mathcal{E}) = N_S(\mathcal{H})$. Notice that $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{N_S(\mathcal{H})}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ is contained in $\mathcal{F} = \mathcal{F}_S(\mathcal{L})$ and in $\mathcal{F}_0 = \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$. Hence,

$$N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F} \cap \mathcal{F}_0 \subseteq \mathcal{F}_0|_{N_S(\mathcal{E})}.$$

It follows moreover from Lemma 3.11 that $E(\mathcal{F})N_S(\mathcal{E}) = \mathcal{F}_{N_S(\mathcal{E})}(E(\mathcal{L})N_S(\mathcal{E})) \subseteq \mathcal{F}_0$. Clearly, \mathcal{B} is contained in \mathcal{F}_0 . Hence,

$$\mathcal{G} := \langle E(\mathcal{F})N_S(\mathcal{E}), \mathcal{B} \rangle \subseteq \mathcal{F}_0 \cap \mathcal{F} \subseteq \mathcal{F}_0|_{N_S(\mathcal{E})}.$$

Let now $P, Q \leq N_S(\mathcal{E})$ and $\varphi \in \text{Hom}_{\mathcal{F}_0}(P, Q)$. Notice that any morphism in $\mathcal{F}_0|_{N_S(\mathcal{E})}$ is of this form. As $E(\mathcal{F}) \trianglelefteq \mathcal{F}_0$, it follows from Lemma 2.9 that there exists $\psi \in \text{Hom}_{(E(\mathcal{F})P)_{\mathcal{F}_0}}(P, \tilde{T}P)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}_0}(\tilde{T})}(P\psi, N_S(\mathcal{E}))$ such that $\varphi = \psi\alpha$. By Lemma 3.11,

$$(E(\mathcal{F})P)_{\mathcal{F}_0} = (E(\mathcal{F})P)_{\mathcal{F}} \subseteq (E(\mathcal{F})N_S(\mathcal{E}))_{\mathcal{F}} \subseteq \mathcal{G}$$

and

$$(E(\mathcal{F})P)_{\mathcal{F}_0} \subseteq N_{\mathcal{F}}(\mathcal{E}).$$

By Lemma 2.13 applied with $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ in place of \mathcal{L} , we have $\tilde{T} \in \delta(\mathcal{F}_0)$. Hence, by [9, Lemma 3.10(b)] and Lemma 3.6(d),

$$N_{\mathcal{F}_0}(\tilde{T}) = \mathcal{F}_{S_0}(N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\tilde{T})) = \mathcal{F}_{S_0}(N_G(\mathcal{H})).$$

As $P\psi$ and $P\psi\alpha = P\varphi$ are contained in $N_S(\mathcal{E}) = N_S(\mathcal{H})$, it follows that α is a morphism in $\mathcal{B} = \mathcal{F}_{N_S(\mathcal{H})}(N_G(\mathcal{H}))$. Since $N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F}_0|_{N_S(\mathcal{E})}$, this proves (b). Observe that $\mathcal{B} \subseteq \mathcal{G} \cap N_{\mathcal{F}}(\mathcal{E})$ and so the argument shows also that a morphism φ as above is a morphism in \mathcal{G} and in $N_{\mathcal{F}}(\mathcal{E})$. Hence, $\mathcal{F}_0|_{N_S(\mathcal{E})}$ is contained in \mathcal{G} and in $N_{\mathcal{F}}(\mathcal{E})$. This completes the proof of (a).

(c) Using (a) one sees that $\mathcal{E} \subseteq \mathcal{F} \cap \mathcal{F}_0 = N_{\mathcal{F}}(\mathcal{E})$. As \mathcal{E} is normal in \mathcal{F}_0 , T is strongly $N_{\mathcal{F}}(\mathcal{E})$ -closed. Moreover, $\mathcal{E}^\alpha = \mathcal{E}$ for every $\alpha \in \text{Aut}_{\mathcal{F}_0}(T)$ and thus also for every $\alpha \in \text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(T)$. By Lemma 2.3(b) remains to show that every $N_{\mathcal{F}}(\mathcal{E})$ -morphism between subgroups of T is in $\langle \mathcal{E}, \text{Aut}_{\mathcal{F}}(T) \rangle_T$. Fix $f \in \mathbb{N}_{\mathcal{L}}(\mathcal{H})$. We need to show that $c_f|_{T \cap S_f} \in \langle \mathcal{E}, \text{Aut}_{\mathcal{F}}(T) \rangle_T$.

By Lemma 2.10 applied with $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ and \mathcal{H} in place of (\mathcal{L}, Δ, S) and \mathcal{N} , there exist $h \in \mathcal{H}$ and $g \in N_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(T)$ such that $(h, g) \in \mathbf{D}$, $f = hg$ and $(S_0)_{(h,g)} = (S_0)_f$. As $T \leq S_0$ and T is strongly closed in \mathcal{F}_0 , we have in particular $T \cap S_{(h,g)} = T \cap S_f$. Hence, $c_f|_{T \cap S_f}$ is the composition of $c_h|_{T \cap S_f}$ and $c_g|_{(T \cap S_f)^h}$. Here $c_h|_{T \cap S_f}$ is a morphism in \mathcal{E} , and $c_g|_{(T \cap S_f)^h}$ extends to $c_g|_T$, which is an element of $\text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(T)$. This proves (c). \square

Proof of Theorem B. As already observed above, $N_{\mathcal{F}}(\mathcal{E})$ is by definition a fusion system over $N_S(\mathcal{H}) = N_S(\mathcal{E})$, i.e. (a) holds. As throughout, T_0 and \tilde{T} are defined as in Hypothesis 3.1. Background Theorem 3 gives then however that $E(\mathcal{E})$ is a fusion system over T_0 and $E(\mathcal{F})$ is a fusion system over \tilde{T} . By Lemma 3.3(b), $N_{\mathcal{F}}(\tilde{T})$ is constrained, G is a model for $N_{\mathcal{F}}(\tilde{T})$, and $N_{\mathcal{H}}(\tilde{T}) \trianglelefteq \trianglelefteq G$ is a model for $N_{\mathcal{E}}(T_0) \trianglelefteq \trianglelefteq N_{\mathcal{F}}(\tilde{T})$. In particular, part (b) holds.

If \tilde{G} is now an arbitrary a model for $N_{\mathcal{F}}(\tilde{T})$ and $\tilde{H} \trianglelefteq \trianglelefteq \tilde{G}$ with $\mathcal{F}_{S \cap \tilde{H}}(\tilde{H}) = N_{\mathcal{E}}(T_0)$, then it follows from Lemma 2.17 that $\mathcal{F}_{N_S(\tilde{H})}(N_{\tilde{G}}(\tilde{H})) = \mathcal{F}_{N_S(N_{\mathcal{H}}(\tilde{T}))}(N_G(N_{\mathcal{H}}(\tilde{T})))$. As $N_G(\mathcal{H}) = N_G(N_{\mathcal{H}}(\tilde{T}))$ by Lemma 3.3(c), we can conclude for such \tilde{G} and \tilde{H} that $N_S(\tilde{H}) = N_S(N_{\mathcal{H}}(\tilde{T})) = N_S(\mathcal{H}) = N_S(\mathcal{E})$ and

$$\mathcal{F}_{N_S(\tilde{H})}(N_{\tilde{G}}(\tilde{H})) = \mathcal{F}_{N_S(\mathcal{H})}(N_G(\mathcal{H})).$$

Hence, part (c) follows from the first statement in Lemma 3.13(a). \square

Lemma 3.14. *Let $X, Y \leq N_S(\mathcal{E})$ with $T \leq X \cap Y$ and $\varphi \in \text{Hom}_{\mathcal{F}}(X, Y)$ with $\varphi|_T \in \text{Aut}(\mathcal{E})$. Then φ is a morphism in $N_{\mathcal{F}}(\mathcal{E})$.*

Proof. By Lemma 3.12, there exists $\psi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(X, \tilde{T}X)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(X\psi, S)$ such that $\varphi = \psi\alpha$. As \mathcal{E} is $N_{\mathcal{F}}(\mathcal{E})$ -invariant by Lemma 3.13(c), we have $T\psi = T$ and $\mathcal{E}^\psi = \mathcal{E}$. Hence, $T = T\psi \leq X\psi$, $T\alpha = T\psi\alpha = T\varphi = T$ and $\mathcal{E}^\alpha = (\mathcal{E}^\psi)^\alpha = \mathcal{E}^\varphi = \mathcal{E}$. By Lemma 3.3(b), G is a model for $N_{\mathcal{F}}(\tilde{T})$, i.e. there exists $g \in G$ with $\alpha = c_g|_{X\psi}$. As $c_g|_T = \alpha|_T \in \text{Aut}(\mathcal{E})$, it follows from Lemma 3.8 that $g \in N_G(\mathcal{H})$. As $X\psi \leq \tilde{T}X \leq N_S(\mathcal{E}) = N_S(\mathcal{H})$, it follows $X\psi\alpha = (X\psi)^g \leq S \cap N_G(\mathcal{H}) = N_S(\mathcal{H})$. So α is a morphism in $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{N_S(\mathcal{H})}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$. Hence, $\varphi = \psi\alpha$ is in $N_{\mathcal{F}}(\mathcal{E})$. \square

Corollary 3.15. *We have $C_{\mathcal{F}}(T) = C_{N_{\mathcal{F}}(\mathcal{E})}(T)$.*

Proof. Clearly $C_{N_{\mathcal{F}}(\mathcal{E})}(T) \subseteq C_{\mathcal{F}}(T)$. If $P, Q \leq C_S(T)$ and $\varphi \in \text{Hom}_{C_{\mathcal{F}}(T)}(P, Q)$, the φ extends to $\hat{\varphi} \in \text{Hom}_{\mathcal{F}}(PT, QT)$ with $\hat{\varphi}|_T = \text{id}_T$. As $\text{id}_T \in \text{Aut}(\mathcal{E})$, Lemma 3.14 yields that $\hat{\varphi}$ is a morphism in $N_{\mathcal{F}}(\mathcal{E})$. Hence, $\varphi = \hat{\varphi}|_P$ is a morphism in $C_{N_{\mathcal{F}}(\mathcal{E})}(T)$. \square

Corollary 3.16. *If $N_S(\mathcal{E}) \leq S_0$, then $C_{\mathcal{F}}(T) = C_{\mathcal{F}_0}(T)|_{C_S(T)}$.*

Proof. As $N_S(\mathcal{E}) \leq S_0$, Lemma 3.13(a) gives that $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_0|_{N_S(\mathcal{E})}$. In particular, $N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F}_0$ and so Corollary 3.15 implies

$$C_{\mathcal{F}}(T) = C_{N_{\mathcal{F}}(\mathcal{E})}(T) \subseteq C_{\mathcal{F}_0}(T)|_{C_S(T)}.$$

On the other hand, if $P, Q \leq C_S(T)$ and $\varphi \in \text{Hom}_{C_{\mathcal{F}_0}(T)}(P, Q)$, then φ extends to $\hat{\varphi} \in \text{Hom}_{\mathcal{F}_0}(PT, QT)$ with $\hat{\varphi}|_T = \text{id}_T$. Using Lemma 3.8, one observes that PT and QT are contained in $TC_S(T) \leq N_S(\mathcal{H}) = N_S(\mathcal{E})$. Hence, it follows that $\hat{\varphi}$ is a morphism in $\mathcal{F}_0|_{N_S(\mathcal{E})} = N_{\mathcal{F}}(\mathcal{E})$. This yields that φ is a morphism in $C_{N_{\mathcal{F}}(\mathcal{E})}(T) = C_{\mathcal{F}}(T)$ and proves thus that $C_{\mathcal{F}_0}(T)|_{C_S(T)} \subseteq C_{\mathcal{F}}(T)$. \square

Our next goal is to show that every saturated subsystem \mathcal{D} of \mathcal{F} with $\mathcal{E} \trianglelefteq \mathcal{D}$ is contained in $N_{\mathcal{F}}(\mathcal{E})$. We first state some technical lemmas which are used in the proof.

Lemma 3.17. *Let $T_0^\perp \leq Q \leq N_S(\mathcal{E})$ and $\varphi \in \text{Hom}_{\mathcal{F}}(Q, S)$ such that $T_0^\perp \varphi = T_0^\perp$. Then*

$$(Q \cap T_0)\varphi \leq T_0.$$

Proof. We may choose S_0 such that $N_S(\mathcal{E}) \leq S_0$. By Lemma 3.12, there exist then $\psi \in \text{Hom}_{\mathcal{F} \cap \mathcal{F}_0}(Q, \tilde{T}Q)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(Q\tilde{T}, S)$ with $\varphi = \psi\alpha$. Theorem 3.6(d) implies that T_0 and T_0^\perp are strongly closed in \mathcal{F}_0 . Thus,

$$(Q \cap T_0)\psi \leq T_0$$

and $T_0^\perp \psi = T_0^\perp$. This yields $T_0^\perp \alpha = T_0^\perp \psi\alpha = T_0^\perp \varphi = T_0^\perp$. By Lemma 3.3(b), G realizes $N_{\mathcal{F}}(\tilde{T})$. Hence, α is realized as a conjugation homomorphism by an element of G which then lies in $N_G(T_0^\perp)$ as $T_0^\perp \alpha = T_0^\perp$. Recall from Lemma 3.3(e) that $N_G(T_0^\perp) = N_G(T_0)$. Hence, it follows that $T_0\alpha = T_0$ and thus $(Q \cap T_0)\varphi = (Q \cap T_0)\psi\alpha \leq T_0\alpha = T_0$. \square

Lemma 3.18. *Let $Q \in \mathcal{E}^{cr}$ and $\varphi \in \text{Hom}_{\mathcal{F}}(Q\tilde{T}, S)$ with $Q\varphi = Q$. Then $T_0\varphi = T_0$ and $T_0^\perp \varphi = T_0^\perp$.*

Proof. If $\mathcal{K} \in \text{Comp}(\mathcal{H})$, then $\mathcal{F}_{S \cap \mathcal{K}}(\mathcal{K}) \trianglelefteq \mathcal{E}$ by Background Theorem 3. Hence, by Lemma 2.5, $Q \cap \mathcal{K} \in \mathcal{F}_{S \cap \mathcal{K}}(\mathcal{K})^{cr}$. In particular, $Q \cap \mathcal{K} \not\leq Z(\mathcal{K})$ as $Z(\mathcal{K}) \neq \mathcal{K} \cap S$ by [13, Lemma 7.10]. On the other hand, if $\mathcal{K} \in \text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H})$, then $\mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{K})$ by [13, Lemma 3.5, Lemma 11.17]. Hence, as $Q \leq T \subseteq \mathcal{H}$, we have in this case $Q \cap \mathcal{K} \subseteq \mathcal{H} \cap \mathcal{K} \subseteq C_{\mathcal{K}}(\mathcal{K}) = Z(\mathcal{K})$. This shows

$$\text{Comp}(\mathcal{H}) = \{\mathcal{K} \in \text{Comp}(\mathcal{L}) : \mathcal{K} \cap Q \not\leq Z(\mathcal{K})\}.$$

As \tilde{T} is strongly \mathcal{F} -closed, φ is a morphism in $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$. Hence, there exists $f \in G$ with $\varphi = c_f|_{Q\tilde{T}}$. Then conjugation by f induces by Lemma 2.13 an automorphism of \mathcal{L} and thus permutes by [13, Lemma 11.12] the components of \mathcal{L} . Moreover, for every $\mathcal{K} \in \text{Comp}(\mathcal{L})$, c_f induces an isomorphism from \mathcal{K} to \mathcal{K}^f so that $Z(\mathcal{K})^f = Z(\mathcal{K}^f)$. As $Q^f = Q\varphi = Q$, we have also $(\mathcal{K} \cap Q)^f = \mathcal{K}^f \cap Q$. So for every $\mathcal{K} \in \text{Comp}(\mathcal{L})$, we have $\mathcal{K} \cap Q \not\leq Z(\mathcal{K})$ if and only if $\mathcal{K}^f \cap Q \not\leq Z(\mathcal{K}^f)$. Therefore, f permutes the elements of $\text{Comp}(\mathcal{H})$ via conjugation and thus also the elements of $\text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H})$. This implies $E(\mathcal{H})^f = E(\mathcal{H})$ and $\mathcal{M}^f = \mathcal{M}$. As $\varphi = c_f|_{Q\tilde{T}}$, the assertion follows. \square

Lemma 3.19. *Let $R \leq S_0$ and set*

$$\mathcal{G} := (\mathcal{E}R)_{\mathcal{F}_0}$$

Then the following hold:

- (a) \mathcal{G} is saturated, $\mathcal{E} \trianglelefteq \mathcal{F}$, $Op(\mathcal{G}) = Op(\mathcal{E})$ and $(\mathcal{H}R, \delta(\mathcal{G}), TR)$ is a regular locality over \mathcal{G} with $E(\mathcal{H}R) = E(\mathcal{H})$. In particular, $E(\mathcal{G}) = E(\mathcal{E})$. Moreover, if $R \leq N_S(\mathcal{E})$, then $\mathcal{G} = \mathcal{F}_{TR}(\mathcal{H}R) \subseteq \mathcal{F}$.

- (b) If $P \leq TR$ with $P \cap T \in \delta(\mathcal{E})$, then $P \in \delta(\mathcal{G})$.
(c) Suppose $P \in \delta(\mathcal{G})$. Then $N_{\mathcal{H}R}(P)$ and $N_{\mathcal{H}}(P)$ are subgroups of \mathcal{L} . In particular,

$$\text{Aut}_{\mathcal{H}}(P) := \{c_h : h \in N_{\mathcal{H}}(P)\}$$

is a subgroup of $\text{Aut}(P)$. Moreover, $O^p(\text{Aut}_{\mathcal{G}}(P)) = O^p(\text{Aut}_{\mathcal{H}}(P))$.

- (d) If $P \leq TR \cap N_S(\mathcal{E})$ such that $P \cap T \in \mathcal{E}^{cr}$, then $P \cap T \in \delta(\mathcal{E})$ and thus $P \in \delta(\mathcal{G})$. Moreover,

$$O^p(\text{Aut}_{\mathcal{G}}(P)) = O^p(\text{Aut}_{\mathcal{H}}(P)) = A_{\mathcal{F}_0, \mathcal{E}}^\circ(P) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P).$$

Proof. (a) We use as before that \mathcal{F}_0 is well-defined and saturated, $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 with $T = S_0 \cap \mathcal{H}$, $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ and $\mathcal{E} = \mathcal{F}_T(\mathcal{H}) \trianglelefteq \mathcal{F}_0$. By Proposition 2.8, \mathcal{G} is saturated, $\mathcal{E} \trianglelefteq \mathcal{G}$, $O^p(\mathcal{G}) = O^p(\mathcal{E})$ and our notion of a product of a normal subsystem with a p -subgroup coincides with the one used in [8]. Thus, it follows from [8, Theorem D(a)] that \mathcal{G} is saturated and $(\mathcal{H}R, \delta(\mathcal{G}), TR)$ is a regular locality over \mathcal{G} . In particular, if $R \leq N_S(\mathcal{E})$, then

$$\mathcal{G} = \mathcal{F}_{TP}(\mathcal{H}P) \subseteq \mathcal{F}_S(\mathcal{L}) = \mathcal{F}.$$

For an arbitrary subgroup $R \leq S_0$, \mathcal{H} is a partial normal subgroup of $\mathcal{H}P$ of p -power index. Thus, by [13, Lemma 11.14], $E(\mathcal{H}P) = E(\mathcal{H})$. It follows now from Background Theorem 3 applied twice (once with $(\mathcal{H}, T, \mathcal{E})$ and once with $(\mathcal{H}P, TP, \mathcal{G})$ in place of $(\mathcal{L}, S, \mathcal{F})$) that $E(\mathcal{E}) = \mathcal{F}_{T_0}(E(\mathcal{H})) = E(\mathcal{G})$. This proves (a).

(b) As $E(\mathcal{E}) = E(\mathcal{G})$, part (b) follows from Lemma 2.6 applied with \mathcal{G} in place of \mathcal{F} .

(c) Let $P \in \delta(\mathcal{G})$. As $(\mathcal{H}R, \delta(\mathcal{G}), TR)$ is a regular locality over \mathcal{G} , it follows that $N_{\mathcal{H}R}(P)$ is a subgroup of $\mathcal{H}R$ and thus of \mathcal{L} . Moreover,

$$\text{Aut}_{\mathcal{G}}(P) = \{c_g : g \in N_{\mathcal{H}R}(P)\}.$$

Since $\mathcal{H} \trianglelefteq \mathcal{H}R$, $N_{\mathcal{H}}(P) = \mathcal{H} \cap N_{\mathcal{H}R}(P)$ is a normal subgroup of $N_{\mathcal{H}R}(P)$. In particular, $\text{Aut}_{\mathcal{H}}(P)$ is a subgroup of $\text{Aut}(P)$. By [14, Lemma 6.1] (applied with $(\mathcal{H}P, \delta(\mathcal{G}), TP)$ in place of (\mathcal{L}, Δ, S)), we have $O^p(N_{\mathcal{H}P}(P)) = O^p(N_{\mathcal{H}}(P))$. Observe that every p' -element of $\text{Aut}_{\mathcal{H}}(P)$ can be realized by conjugation with a p' -element of $N_{\mathcal{H}}(P)$, and similarly every p' -automorphism of $\text{Aut}_{\mathcal{G}}(P)$ can be realized by conjugation with a p' -element of $N_{\mathcal{H}P}(P)$. This implies

$$O^p(\text{Aut}_{\mathcal{H}}(P)) = \{c_h : h \in O^p(N_{\mathcal{H}}(P))\} = \{c_h : h \in O^p(N_{\mathcal{H}R}(P))\} = O^p(\text{Aut}_{\mathcal{G}}(P)).$$

Hence (c) holds.

(d) By [13, Lemma 10.4], $\mathcal{E}^{cr} \subseteq \delta(\mathcal{E})$. Moreover, if $P \leq TR$ with $P \cap T = \mathcal{E}^c$, then [10, Lemma 4.7] gives $O^p(\text{Aut}_{\mathcal{G}}(P)) = A_{\mathcal{F}_0, \mathcal{E}}^\circ(P)$. Hence, supposing P is a subgroup of TR with $P \cap T \in \mathcal{E}^{cr}$, it follows from (b) and (c) that $O^p(\text{Aut}_{\mathcal{H}}(P)) = O^p(\text{Aut}_{\mathcal{G}}(P)) = A_{\mathcal{F}_0, \mathcal{E}}^\circ(P)$. Assume now in addition that $P \leq N_S(\mathcal{E})$. Then $A_{\mathcal{F}_0, \mathcal{E}}^\circ(P) \leq \text{Aut}_{\mathcal{H}}(P) \leq \text{Aut}_{\mathcal{F}}(P)$. As $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \cap \mathcal{F}_0$ by Lemma 3.13(b), it follows $A_{\mathcal{F}_0, \mathcal{E}}^\circ(P) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P)$. This proves (d). \square

We use from now on without further reference that, for all $P \leq N_S(\mathcal{E})$ with $P \cap T \in \mathcal{E}^{cr}$, the normalizer $N_{\mathcal{H}}(P)$ is a subgroup of \mathcal{L} and

$$\text{Aut}_{\mathcal{H}}(P) := \{c_h : h \in N_{\mathcal{H}}(P)\}$$

is a subgroup of $\text{Aut}(P)$. Notice that this is true by Lemma 3.19(b),(c) (applied e.g. for a Sylow subgroup S_0 of $N_G(\mathcal{H})$ containing $N_S(\mathcal{H}) = N_S(\mathcal{E})$, and with $R = N_S(\mathcal{E})$).

Lemma 3.20. *Let $P \leq N_S(\mathcal{E})$ such that $P \cap T \in \mathcal{E}^{cr}$. Then $\text{Aut}_{\mathcal{H}}(P) \trianglelefteq \trianglelefteq \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P)$.*

Proof. By Lemma 3.2(a), $\mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{M}) \subseteq C_{\mathcal{L}}(T_0^\perp)$. In particular,

$$\text{Aut}_{\mathcal{H}}(P) \leq \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P).$$

As $\mathcal{F}_{T_0}(E(\mathcal{H})) = E(\mathcal{E}) \trianglelefteq \mathcal{E}$ by Background Theorem 3, it follows from Lemma 2.5 and [13, Lemma 10.4] that

$$P_0 := P \cap T_0 \in E(\mathcal{E})^{cr} \subseteq \delta(E(\mathcal{E})).$$

By Lemma 3.2(a), $E(\mathcal{H})$ and \mathcal{M} are partial normal subgroups of $E(\mathcal{L})$ with $E(\mathcal{L}) = E(\mathcal{H})\mathcal{M}$ and $E(\mathcal{H}) \subseteq \mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{M})$. The latter property implies by [13, Lemma 3.5] that $E(\mathcal{H})$ commutes with \mathcal{M} in the sense of [13, Definition 2]. Thus, it follows from [13, Lemma 10.15(d)] that

$$X := P_0 T_0^\perp \in \delta(\mathcal{F}_{S \cap E(\mathcal{L})}(E(\mathcal{L}))).$$

By Background Theorem 3, we have $\mathcal{F}_{S \cap E(\mathcal{L})}(E(\mathcal{L})) = E(\mathcal{F})$ and by [8, 7.22], $\delta(E(\mathcal{F})) \subseteq \delta(\mathcal{F})$.² We can conclude that

$$X \in \Delta = \delta(\mathcal{F}).$$

In particular, $N_{\mathcal{L}}(X)$ is a group.

It follows from Lemma 3.3(e) that $P \leq N_S(\mathcal{E}) = N_S(\mathcal{H}) \leq N_S(T_0^\perp) = N_S(T_0)$. Thus, $P \leq N_S(P_0)$ and

$$P \leq N_S(X).$$

Notice moreover that

$$N_{\mathcal{H}}(P) \leq N_{\mathcal{H}}(X)$$

as $P_0 = P \cap E(\mathcal{H})$ is normalized by $N_{\mathcal{H}}(P)$ and $\mathcal{H} \subseteq C_{\mathcal{L}}(T_0^\perp)$.

Let now $\mathcal{H} = \mathcal{H}_0 \trianglelefteq \mathcal{H}_1 \trianglelefteq \cdots \trianglelefteq \mathcal{H}_n = \mathcal{L}$ be a subnormal series of \mathcal{H} in \mathcal{L} . Then

$$N_{\mathcal{H}}(P) = N_{N_{\mathcal{H}}(X)}(P) = N_{N_{\mathcal{H}_0}(X)}(P) \trianglelefteq N_{N_{\mathcal{H}_1}(X)}(P) \trianglelefteq \cdots \trianglelefteq N_{N_{\mathcal{H}_n}(X)}(P) = N_{N_{\mathcal{L}}(X)}(P)$$

is a subnormal series of $N_{\mathcal{H}}(P)$ in $N_{N_{\mathcal{L}}(X)}(P)$. This leads to a subnormal series of $\text{Aut}_{\mathcal{H}}(P)$ in

$$\text{Aut}_{N_{\mathcal{L}}(X)}(P) = \{c_g|_P : g \in N_{N_{\mathcal{L}}(X)}(P)\}.$$

Thus, it remains only to argue that $\text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P) \leq \text{Aut}_{N_{\mathcal{L}}(X)}(P)$. By [9, Lemma 3.10(b)], $N_{\mathcal{F}}(X) = \mathcal{F}_{N_S(X)}(N_{\mathcal{L}}(X))$ and thus

$$\text{Aut}_{N_{\mathcal{L}}(X)}(P) = \text{Aut}_{N_{\mathcal{F}}(X)}(P).$$

Let $\varphi \in \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P)$. Then φ extends to $\hat{\varphi} \in \text{Hom}_{\mathcal{F}}(PT_0^\perp, S)$ with $T_0^\perp \hat{\varphi} = T_0^\perp$. By Lemma 3.17, we have $P_0 \hat{\varphi} \leq T_0$ and thus $P_0 \hat{\varphi} \leq P \cap T_0 = P_0$. This implies $X \hat{\varphi} = X$ and proves thus $\text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P) \leq \text{Aut}_{N_{\mathcal{F}}(X)}(P) = \text{Aut}_{N_{\mathcal{L}}(X)}(P)$ as required. \square

Lemma 3.21. *Let $P \leq N_S(\mathcal{E})$ and $\chi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(P, N_S(\mathcal{E}))$. Then*

$$\chi^{-1} A_{\mathcal{F}, \mathcal{E}}^\circ(P)\chi = A_{\mathcal{F}, \mathcal{E}}^\circ(P\chi) \text{ and } \chi^{-1} A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P)\chi = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P\chi).$$

In particular, $A_{\mathcal{F}, \mathcal{E}}^\circ(P) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P)$ if and only if $A_{\mathcal{F}, \mathcal{E}}^\circ(P\chi) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P\chi)$.

Proof. Recall that \mathcal{E} is $N_{\mathcal{F}}(\mathcal{E})$ -invariant by Lemma 3.13(c). In particular, \mathcal{E} is weakly $N_{\mathcal{F}}(\mathcal{E})$ -invariant by Lemma 2.3(a). We can conclude that $(P \cap T)\chi = (P\chi) \cap T$ and

$$\chi^{-1} \text{Aut}_{\mathcal{E}}(P \cap T)\chi = \text{Aut}_{\mathcal{E}}((P\chi) \cap T).$$

Since the map $\text{Aut}_{\mathcal{F}}(P) \rightarrow \text{Aut}_{\mathcal{F}}(P\chi), \delta \mapsto \chi^{-1}\delta\chi$ is an isomorphism which restricts to an isomorphism from $\text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(P) \rightarrow \text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(P)$, the assertion is now easy to check. \square

Lemma 3.22. *Let $P \leq N_S(\mathcal{E}) \cap S_0$ such that $P \cap T \in \mathcal{E}^{cr}$. Then there exists $\chi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(P, N_S(\mathcal{E}))$ such that $P\chi \leq S_0$ and the following hold:*

- (i) $O_p(N_{\text{Aut}_{\mathcal{F}}(P\chi)}((P\chi) \cap T)) \leq \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P\chi)$;
- (ii) $(P\chi) \cap T$ is fully normalized in $(\mathcal{E}(P\chi))_{\mathcal{F}_0}$ and an element of \mathcal{E}^{cr} .

²Alternatively, a direct argument involving Lemma 10.2, Lemma 10.6, Lemma 10.15(d) and Lemma 11.5 in [13] could be given to show $\delta(\mathcal{F}_{S \cap E(\mathcal{L})}(E(\mathcal{L}))) \subseteq \delta(\mathcal{F})$.

Proof. To ease notation set $Q := P \cap T$. We show first

$$\text{There exists } \varphi \in \text{Hom}_{\mathcal{F}}(P, S) \text{ such that } N_{\text{Aut}_S(P\varphi)}(Q\varphi) \in \text{Syl}_p(N_{\text{Aut}_{\mathcal{F}}(P\varphi)}(Q\varphi)). \quad (3.2)$$

For the proof fix $\beta \in \text{Hom}_{\mathcal{F}}(P, S)$ such that $P\beta$ is fully \mathcal{F} -normalized. Then by the Sylow axiom [4, Proposition I.2.5], we have $\text{Aut}_S(P\beta) \in \text{Syl}_p(\text{Aut}_{\mathcal{F}}(P\beta))$. As $H := N_{\text{Aut}_{\mathcal{F}}(P\beta)}(Q\beta)$ is a subgroup of $\text{Aut}_{\mathcal{F}}(P\beta)$, there exists $\gamma \in \text{Aut}_{\mathcal{F}}(P\beta)$ such that $H^\gamma \cap \text{Aut}_S(P\beta) \in \text{Syl}_p(H^\gamma)$. Note that $P\beta\gamma = P\beta$ and $H^\gamma = N_{\text{Aut}_{\mathcal{F}}(P\beta)}(Q\beta\gamma)$. Hence (3.2) holds for $\varphi := \beta\gamma$.

Let now φ be as in (3.2). Then in particular,

$$O_p(N_{\text{Aut}_{\mathcal{F}}(P\varphi)}(Q\varphi)) \leq \text{Aut}_S(P\varphi) \leq \text{Aut}_{N_{\mathcal{F}}(\tilde{T})}(P\varphi).$$

By Lemma 3.12, there exist $\psi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(P, \tilde{T}P) \cap \text{Hom}_{\mathcal{F}_0}(P, \tilde{T}P)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(P\psi, S)$ such that $\varphi = \psi\alpha$. Notice that we have an isomorphism from $N_{\text{Aut}_{\mathcal{F}}(P\varphi)}(Q\varphi)$ to $N_{\text{Aut}_{\mathcal{F}}(P\psi)}(Q\psi)$ defined by $\delta \mapsto \alpha\delta\alpha^{-1}$. Hence,

$$O_p(N_{\text{Aut}_{\mathcal{F}}(P\psi)}(Q\psi)) = \alpha O_p(N_{\text{Aut}_{\mathcal{F}}(P\varphi)}(Q\varphi))\alpha^{-1} \leq \text{Aut}_{N_{\mathcal{F}}(\tilde{T})}(P\psi). \quad (3.3)$$

Since $\mathcal{E} \trianglelefteq \mathcal{F}_0$ and ψ is a morphism in \mathcal{F}_0 , it follows from $Q \in \mathcal{E}^{cr}$ and [8, Lemma 2.6] that $Q\psi \in \mathcal{E}^{cr}$. Hence, using (3.3) and applying Lemma 3.18 with $Q\psi$ in place of Q , we can conclude that

$$O_p(N_{\text{Aut}_{\mathcal{F}}(P\psi)}(Q\psi)) \leq \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P\psi). \quad (3.4)$$

Observe now that $P\psi \leq N_S(\mathcal{E}) = N_S(\mathcal{H}) \leq N_S(T_0^\perp)$ by Lemma 3.8 and Lemma 3.3(e). Moreover, by Lemma 3.2(a), $\mathcal{H} \subseteq C_{\mathcal{L}}(\mathcal{M}) \subseteq C_{\mathcal{L}}(T_0^\perp)$. As $N_{\mathcal{L}}(T_0^\perp)$ is by [12, Lemma 5.5] a partial subgroup of \mathcal{L} , it follows $\mathcal{H}(P\psi) \subseteq N_{\mathcal{L}}(T_0^\perp)$. Using Lemma 3.19(a), we see now that

$$\mathcal{G} := (\mathcal{E}(P\psi))_{\mathcal{F}_0} = \mathcal{F}_{T(P\psi)}(\mathcal{H}(P\psi)) \subseteq N_{\mathcal{F}}(T_0^\perp)$$

is saturated. Thus, by [4, Lemma I.2.6(c)], there exists $\rho \in \text{Hom}_{\mathcal{G}}(N_{T(P\psi)}(Q\psi), T(P\psi))$ such that $Q\psi\rho$ is fully \mathcal{G} -normalized. Notice that $Q\psi \trianglelefteq P\psi$ and thus $P\psi\rho$ is defined. Moreover, by definition of \mathcal{G} , we have $T(P\psi) = T(P\psi\rho)$ and thus $\mathcal{G} = (\mathcal{E}(P\psi\rho))_{\mathcal{F}_0}$. Since $\mathcal{E} \trianglelefteq \mathcal{F}_0$ and ψ and ρ are morphisms in \mathcal{F}_0 , we have that $(P\psi\rho) \cap T = Q\psi\rho$ is fully \mathcal{G} -normalized and moreover an element of \mathcal{E}^{cr} by [8, Lemma 2.6]. Since ρ is a morphism in \mathcal{G} and $\mathcal{G} \subseteq N_{\mathcal{F}}(T_0^\perp)$, it follows from (3.4) that

$$O_p(N_{\text{Aut}_{\mathcal{F}}(P\psi\rho)}((P\psi\rho) \cap T)) = \rho^{-1} O_p(N_{\text{Aut}_{\mathcal{F}}(P\psi)}(Q\psi))\rho \leq \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P\psi\rho).$$

Hence, the assertion follows for $\chi = \psi\rho$. \square

Lemma 3.23. *Let $P \leq N_S(\mathcal{E}) \cap S_0$ such that $P \cap T \in \mathcal{E}^{cr}$. Then*

$$A_{\mathcal{F}, \mathcal{E}}^\circ(P) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P) = A_{\mathcal{F}_0, \mathcal{E}}^\circ(P) = O^p(\text{Aut}_{\mathcal{H}}(P)).$$

Proof. Set $\mathcal{G} := (\mathcal{E}P)_{\mathcal{F}_0}$. By Lemma 3.19, \mathcal{G} is saturated, $(\mathcal{H}P, \delta(\mathcal{G}), TP)$ is a regular locality over \mathcal{G} , $P \in \delta(\mathcal{G})$ and

$$O^p(\text{Aut}_{\mathcal{G}}(P)) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P) = A_{\mathcal{F}_0, \mathcal{E}}^\circ(P) = O^p(\text{Aut}_{\mathcal{H}}(P)). \quad (3.5)$$

Thus, it remains to show $\text{Aut}_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}(P) = A_{\mathcal{F}, \mathcal{E}}^\circ(P)$. Set

$$A := N_{\text{Aut}_{\mathcal{F}}(P)}(P \cap T).$$

Replacing P by a suitable $N_{\mathcal{F}}(\mathcal{E})$ -conjugate, Lemma 3.21 and Lemma 3.22 allow us to assume that

$$P \cap T \text{ is fully } \mathcal{G}\text{-normalized and } O_p(A) \leq \text{Aut}_{N_{\mathcal{F}}(T_0^\perp)}(P). \quad (3.6)$$

Clearly, $A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^\circ(P) \leq A_{\mathcal{F}, \mathcal{E}}^\circ(P)$, so it remains to show the converse inclusion. For the proof let $\varphi \in A_{\mathcal{F}, \mathcal{E}}^\circ(P)$ be a p' -automorphism. By (3.5), it remains to show that $\varphi \in O^p(\text{Aut}_{\mathcal{H}}(P))$.

Observe that $\varphi_0 := \varphi|_{P \cap T} \in \text{Aut}_{\mathcal{E}}(P \cap T)$ is a p' -automorphism. Set $\hat{P} := PC_{TP}(P \cap T)$. We show first that

$$\text{there exists } \beta \in \text{Aut}_{\mathcal{G}}(\hat{P}) \text{ such that } \beta|_{P \cap T} = \varphi_0. \quad (3.7)$$

Notice that φ_0 is a morphism in \mathcal{G} and $C_{TP}(P \cap T) \leq N_{\varphi_0}^{\mathcal{G}}$ (where we use the notation introduced in [4, Definition I.2.2]). Using e.g. [10, Remark 2.1]), observe furthermore that

$$\varphi_0^{-1} \text{Aut}_P(P \cap T) \varphi_0 = \text{Aut}_{P\varphi_0}(P \cap T) = \text{Aut}_P(P \cap T)$$

and hence $P \leq N_{\varphi_0}^{\mathcal{G}}$. Therefore, $\hat{P} \leq N_{\varphi_0}^{\mathcal{G}}$. Since \mathcal{G} is saturated and $P \cap T$ is by (3.6) fully \mathcal{G} -normalized, it follows that φ_0 extends to $\beta \in \text{Hom}_{\mathcal{G}}(\hat{P}, TP)$ (cf. [4, Proposition I.2.5]). As φ is a p' -automorphism and $[P, \varphi] \leq P \cap T$, using a property of coprime action (see e.g. [15, 8.2.7(a)]), we can conclude that $P = (P \cap T)C_P(\varphi)$. As $\varphi|_{P \cap T} = \varphi_0 = \beta|_{P \cap T}$ and $P \cap T \trianglelefteq P$, it is a consequence of [10, Lemma 2.4] that

$$[C_P(\varphi), \beta] := \langle x^{-1}(x\beta) : x \in C_P(\varphi) \rangle \leq C_{TP}((P \cap T)\beta) = C_{TP}(P \cap T) \leq \hat{P}.$$

Notice that β normalizes $P \cap T$ and thus also $C_{TP}(P \cap T)$. As $\hat{P} = (P \cap T)C_P(\varphi)C_{TP}(P \cap T)$, we can conclude that $\hat{P}\beta = \hat{P}$. Hence, (3.7) holds.

We choose now β as in (3.7). Since φ_0 is a p' -element, replacing β by a suitable power of itself, we may assume that β is a p' -element of $\text{Aut}_{\mathcal{G}}(\hat{P})$. As $\mathcal{G} = (\mathcal{E}P)_{\mathcal{F}_0}$, it follows from [10, Lemma 4.7] that $\beta \in O^p(\text{Aut}_{\mathcal{G}}(\hat{P})) \leq A_{\mathcal{F}_0, \mathcal{E}}^{\circ}(\hat{P})$. In particular,

$$[\hat{P}, \beta] \leq \hat{P} \cap T.$$

As above, it follows from [10, Lemma 2.4] that $[C_P(\varphi), \beta] \leq C_{TP}(P \cap T)$. Hence,

$$[C_P(\varphi), \beta] \leq C_T(P \cap T) \leq P \cap T$$

since $P \cap T \in \mathcal{E}^c$. As $P = (P \cap T)C_P(\varphi)$ and $(P \cap T)\beta = P \cap T$, this yields $P\beta = P$ and thus, by (3.5),

$$\psi := \beta|_P \in O^p(\text{Aut}_{\mathcal{G}}(P)) = O^p(\text{Aut}_{\mathcal{H}}(P)) = A_{N_{\mathcal{F}}(\mathcal{E}), \mathcal{E}}^{\circ}(P).$$

Notice that $\psi|_{P \cap T} = \varphi_0 = \varphi|_{P \cap T}$ and thus $(\varphi\psi^{-1})|_{P \cap T} = \text{id}_{P \cap T}$. Moreover, $[P, \psi] \leq P \cap T$ and $[P, \varphi] \leq P \cap T$, so $[P, \varphi\psi^{-1}] \leq P \cap T$. By a property of coprime action (see e.g. [15, 8.2.7(b)]), $C_A(P/P \cap T) \cap C_A(P \cap T)$ is a normal p -subgroup of A and thus contained in $O_p(A)$. Hence, $\varphi\psi^{-1} \in O_p(A)$ and thus

$$\varphi \in X := O_p(A)O^p(\text{Aut}_{\mathcal{H}}(P)).$$

Notice that $\text{Aut}_{\mathcal{H}}(P) \leq A$ normalizes $O_p(A)$ and therefore X is a group with $O_p(A) \leq O_p(X)$. By Lemma 3.20, $\text{Aut}_{\mathcal{H}}(P) \trianglelefteq \trianglelefteq \text{Aut}_{N_{\mathcal{F}}(T_0^{\perp})}(P)$ and thus $O^p(\text{Aut}_{\mathcal{H}}(P)) \trianglelefteq \trianglelefteq \text{Aut}_{N_{\mathcal{F}}(T_0^{\perp})}(P)$. By assumption (3.6), we have $O_p(A) \leq \text{Aut}_{N_{\mathcal{F}}(T_0^{\perp})}(P)$. Hence, $O^p(\text{Aut}_{\mathcal{H}}(P))$ is subnormal in X . By Lemma 2.14, this implies that $O_p(A) \leq O_p(X)$ normalizes $O^p(\text{Aut}_{\mathcal{H}}(P))$. In particular, $O^p(\text{Aut}_{\mathcal{H}}(P))$ is normal in X . As $O^p(\text{Aut}_{\mathcal{H}}(P))$ has p -power index in X , this implies $O^p(X) = O^p(\text{Aut}_{\mathcal{H}}(P))$. As φ is a p' -automorphism of P , it follows $\varphi \in O^p(X) = O^p(\text{Aut}_{\mathcal{H}}(P))$ as required. \square

Proposition 3.24. *Let \mathcal{D} be a saturated subsystem of \mathcal{F} with $\mathcal{E} \trianglelefteq \mathcal{D}$. Then $\mathcal{D} \subseteq N_{\mathcal{F}}(\mathcal{E})$.*

Proof. Let $R \leq S$ such that \mathcal{D} is a fusion system over R . As $\mathcal{E} \trianglelefteq \mathcal{D}$, we have $R \leq N_S(\mathcal{E})$. Lemma 2.9 gives

$$\mathcal{D} = \langle (\mathcal{E}R)_{\mathcal{D}}, N_{\mathcal{D}}(T) \rangle.$$

Notice that we may choose S_0 always such that $N_S(\mathcal{E}) \leq S_0$. For $P \leq TR$ with $P \cap T \in \mathcal{E}^{cr}$, it follows thus from Lemma 3.23 that $A_{\mathcal{D}, \mathcal{E}}^{\circ}(P) \leq A_{\mathcal{F}, \mathcal{E}}^{\circ}(P) \leq \text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(P)$. Hence, by definition of $(\mathcal{E}R)_{\mathcal{D}}$, we have $(\mathcal{E}R)_{\mathcal{D}} \subseteq N_{\mathcal{F}}(\mathcal{E})$.

A morphism in $N_{\mathcal{D}}(T)$ is the restriction of a morphism $\varphi \in \text{Hom}_{\mathcal{D}}(X, Y)$ where $X, Y \leq R$ with $T \leq X \cap Y$ and $T\varphi = T$. As $\mathcal{E} \trianglelefteq \mathcal{D}$, for any such φ , we have $\varphi|_T \in \text{Aut}(\mathcal{E})$. Thus φ is a morphism in $N_{\mathcal{F}}(\mathcal{E})$ by Lemma 3.14. This shows $N_{\mathcal{D}}(T) \subseteq N_{\mathcal{F}}(\mathcal{E})$ and therefore $\mathcal{D} \subseteq N_{\mathcal{F}}(\mathcal{E})$. \square

Proof of Theorem D. Choose S_0 such that $N_S(\mathcal{E}) = N_S(\mathcal{H}) \leq S_0$. It follows then from Lemma 3.23 and the definitions of $\mathcal{E}R := (\mathcal{E}R)_{\mathcal{F}}$ and $(\mathcal{E}R)_{\mathcal{F}_0}$ that $(\mathcal{E}R)_{\mathcal{F}} = (\mathcal{E}R)_{\mathcal{F}_0}$. Hence, by Lemma 3.19(a), $\mathcal{E}R$ is saturated, $\mathcal{E} \trianglelefteq \mathcal{E}R$, $O^p(\mathcal{E}R) = O^p(\mathcal{E})$ and $(\mathcal{H}R, \delta(\mathcal{E}R), TR)$ is a regular locality over $\mathcal{E}R$.

Fix now a saturated subsystem \mathcal{D} of \mathcal{F} over TR with $O^p(\mathcal{D}) = O^p(\mathcal{E})$. If $\mathcal{E} \trianglelefteq \mathcal{D}$, then it follows from Proposition 3.24 that \mathcal{D} is contained in $N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F}_0$ and is thus equal to $(\mathcal{E}R)_{\mathcal{F}_0} = \mathcal{E}R$ by Proposition 2.8 applied with \mathcal{F}_0 in place of \mathcal{F} . Hence,

$$\mathcal{D} = \mathcal{E}R \text{ if } \mathcal{E} \trianglelefteq \mathcal{D}. \quad (3.8)$$

Using this property with $(O^p(\mathcal{E}), T)$ in place of (\mathcal{E}, R) (along with the fact that $O^p(\mathcal{E}) = O^p(O^p(\mathcal{E}))$ by [8, Lemma 2.29]) we can conclude the following statement.

$$\text{Every saturated subsystem } \mathcal{D}' \text{ of } \mathcal{F} \text{ over } T \text{ with } O^p(\mathcal{D}') = O^p(\mathcal{E}) \text{ equals } (O^p(\mathcal{E})T)_{\mathcal{F}}. \quad (3.9)$$

As $T \trianglelefteq TR$, by [4, Theorem I.7.4], there is a unique normal subsystem \mathcal{E}' of \mathcal{D} over T of p -power index. Then $O^p(\mathcal{E}') = O^p(\mathcal{D}) = O^p(\mathcal{E})$ by [8, Lemma 2.29] and assumption. Hence, applying (3.9) twice, we obtain $\mathcal{E} = (O^p(\mathcal{E})T)_{\mathcal{F}} = \mathcal{E}' \trianglelefteq \mathcal{D}$. By (3.8), this implies $\mathcal{D} = \mathcal{E}R$. \square

3.4. Conjugates of \mathcal{E} . In this subsection we will look at properties of \mathcal{F} -conjugates of \mathcal{E} and complete the proof of Theorem C. Many results we prove below are not needed in the proofs of the main theorems, but seem interesting on their own.

We continue to assume Hypotheses 3.1 and 3.10. Recall from the introduction that

$$\mathcal{E}^{\mathcal{F}} := \{\mathcal{E}^{\varphi} : \varphi \in \text{Hom}_{\mathcal{F}}(T, S)\}$$

denotes the set of \mathcal{F} -conjugates of \mathcal{E} .

Lemma 3.25. *We have $\mathcal{E}^{\mathcal{F}} = \mathcal{E}^{N_{\mathcal{F}}(\tilde{T})}$ and $T^{\mathcal{F}} = T^{N_{\mathcal{F}}(\tilde{T})}$.*

Proof. As $T \leq N_S(\mathcal{E})$, it follows from Lemma 3.12 that there exists $\psi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(T, \tilde{T}T)$ and $\alpha \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(T\psi, S)$ such that $\varphi = \psi\alpha$. By Lemma 3.13(c), \mathcal{E} is $N_{\mathcal{F}}(\mathcal{E})$ -invariant. Hence, $T\psi = T$ and $\mathcal{E}^{\psi} = \mathcal{E}$, which yields $T\varphi = T\psi\alpha = T\alpha \in T^{N_{\mathcal{F}}(\tilde{T})}$ and similarly $\mathcal{E}^{\varphi} = (\mathcal{E}^{\psi})^{\alpha} = \mathcal{E}^{\alpha} \in \mathcal{E}^{N_{\mathcal{F}}(\tilde{T})}$. The assertion follows. \square

Lemma 3.26.

- (a) *Let $g \in G$ such that $T^g \leq S$. Then $\mathcal{H}^g \trianglelefteq \trianglelefteq \mathcal{L}$ with $\mathcal{H}^g \cap S = T^g$. Moreover, setting $\varphi := c_g|_T$, we have $\mathcal{E}^{\varphi} = \mathcal{F}_{T^g}(\mathcal{H}^g) \trianglelefteq \trianglelefteq \mathcal{F}$.*
- (b) *For every $g \in G$, the conjugation automorphism $c_g \in \text{Aut}(\mathcal{L})$ restricts to isomorphisms from $N_G(\mathcal{H})$ to $N_G(\mathcal{H}^g)$ and from $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ to $\mathbb{N}_{\mathcal{L}}(\mathcal{H}^g)$. For $S_0 \in \text{Syl}_p(N_G(\mathcal{H}))$ we have $S_0^g \in \text{Syl}_p(N_G(\mathcal{H}^g))$.*

Proof. (a) By Lemma 2.13, $c_g \in \text{Aut}(\mathcal{L})$ and thus $\mathcal{H}^g \trianglelefteq \trianglelefteq \mathcal{L}$. It follows moreover from Corollary 2.12 that $\mathcal{E}^{\varphi} = \mathcal{F}_{T^g}(\mathcal{H}^g)$. By Background Theorem 3, we have $\mathcal{F}_{T^g}(\mathcal{H}^g) \trianglelefteq \trianglelefteq \mathcal{F}$.

(b) Using again $c_g \in \text{Aut}(\mathcal{L})$, one observes easily that $N_G(\mathcal{H})c_g = N_G(\mathcal{H}^g)$, i.e. c_g induces an isomorphism from $N_G(\mathcal{H})$ to $N_G(\mathcal{H}^g)$, which then takes a Sylow p -subgroup of $N_G(\mathcal{H})$ to a Sylow p -subgroup of $N_G(\mathcal{H}^g)$. By [13, Lemma 11.12], we have also $E(\mathcal{L})c_g = E(\mathcal{L})$. As $\mathbb{N}_{\mathcal{L}}(\mathcal{H}) = E(\mathcal{L})N_G(\mathcal{H})$ and $\mathbb{N}_{\mathcal{L}}(\mathcal{H}^g) = E(\mathcal{L})N_G(\mathcal{H}^g)$ the assertion follows. \square

Lemma 3.27. *Every \mathcal{F} -conjugate of \mathcal{E} is a subnormal subsystem of \mathcal{F} .*

Proof. As $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$, this follows from Lemma 3.25 and Lemma 3.26. \square

Recall that we call \mathcal{E} *fully \mathcal{F} -normalized*, if

$$|N_S(\mathcal{E})| \geq |N_S(\tilde{\mathcal{E}})| \text{ for all } \tilde{\mathcal{E}} \in \mathcal{E}^{\mathcal{F}}.$$

Notice that $\mathcal{E}^{\mathcal{F}} = (\mathcal{E}')^{\mathcal{F}}$ for every $\mathcal{E}' \in \mathcal{E}^{\mathcal{F}}$. Hence, we can always choose a fully \mathcal{F} -normalized \mathcal{F} -conjugate of \mathcal{E} .

Definition 3.28. The subsystem \mathcal{E} is called *fully \mathcal{F} -automized* if

$$\text{Aut}_S(T) \cap \text{Aut}(\mathcal{E}) \in \text{Syl}_p(\text{Aut}_{\mathcal{F}}(T) \cap \text{Aut}(\mathcal{E})).$$

We will prove below that \mathcal{E} is fully \mathcal{F} -normalized if and only if T is fully \mathcal{F} -centralized and \mathcal{E} is fully \mathcal{F} -automized.

Lemma 3.29. *Suppose $N_S(\mathcal{E}) \leq S_0$. Then the following hold.*

(a) $\text{Aut}_{S_0}(T)$ is a Sylow p -subgroup of $\text{Aut}_{\mathcal{F}_0}(T)$ and of

$$\text{Aut}_{\mathcal{F}_0}(T) \cap \text{Aut}_{\mathcal{F}}(T) = \text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(T) = \text{Aut}_{\mathcal{F}}(T) \cap \text{Aut}(\mathcal{E}).$$

(b) *The following properties are equivalent.*

- (i) \mathcal{E} is fully automized;
- (ii) T is fully $N_{\mathcal{F}}(\mathcal{E})$ -automized.
- (iii) $\text{Aut}_{N_S(\mathcal{E})}(T) = \text{Aut}_{S_0}(T)$.

Proof. (a) As $\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ is saturated and $T \trianglelefteq S_0$, we have $\text{Aut}_{S_0}(T) \in \text{Syl}_p(\text{Aut}_{\mathcal{F}_0}(T))$. Notice moreover that $\text{Aut}_{S_0}(T) \leq \text{Aut}_{\mathcal{F}}(T)$ since $\mathcal{F} = \mathcal{F}_S(\mathcal{L})$. Hence, $\text{Aut}_{S_0}(T)$ is also a Sylow p -subgroup of $\text{Aut}_{\mathcal{F}_0}(T) \cap \text{Aut}_{\mathcal{F}}(T)$. It follows from Lemma 3.13(b) that

$$\text{Aut}_{\mathcal{F}}(T) \cap \text{Aut}_{\mathcal{F}_0}(T) = \text{Aut}_{\mathcal{F} \cap \mathcal{F}_0}(T) = \text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(T).$$

By Lemma 3.13(c), \mathcal{E} is $N_{\mathcal{F}}(\mathcal{E})$ invariant and so $\text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(T) \leq \text{Aut}_{\mathcal{F}}(T) \cap \text{Aut}(\mathcal{E})$. The converse inclusion follows from Lemma 3.14. So (a) holds.

(b) Notice that $\text{Aut}_S(T) \cap \text{Aut}(\mathcal{E}) = \text{Aut}_{N_S(\mathcal{E})}(T)$. Hence, by (a), property (i) is equivalent to $\text{Aut}_{N_S(\mathcal{E})}(T) \in \text{Syl}_p(\text{Aut}_{N_{\mathcal{F}}(\mathcal{E})}(T))$, i.e. to (ii). Assume now $N_S(\mathcal{E}) \leq S_0$. Then $\text{Aut}_{N_S(\mathcal{E})}(T) \leq \text{Aut}_{S_0}(T)$. So (ii) is by (a) equivalent to $\text{Aut}_{N_S(\mathcal{E})}(T) = \text{Aut}_{S_0}(T)$. \square

Lemma 3.30.

- (a) *Suppose $C_S(T) \leq S_0$. Then T is fully \mathcal{F} -centralized if and only if $C_S(T) = C_{S_0}(T)$.*
- (b) *If T is fully centralized and $N_S(\mathcal{E}) \leq S_0$, then $C_{\mathcal{F}}(T) = C_{\mathcal{F}_0}(T)$.*

Proof. (a) It follows from Lemma 3.8 that $C_G(T) \leq X := N_G(\mathcal{H})$ and thus $C_G(T) = C_X(T) \trianglelefteq N_X(T)$. Notice that S_0 is a Sylow p -subgroup of X which is contained in $N_X(T)$. Thus, S_0 is also a Sylow p -subgroup of $N_G(T)$. As $C_G(T) \trianglelefteq N_X(T)$, it follows that $C_{S_0}(T) = S_0 \cap C_G(T) \in \text{Syl}_p(C_G(T))$. Since $C_S(T) \leq C_{S_0}(T)$ by assumption, we can conclude that $C_S(T) \in \text{Syl}_p(C_G(T))$ if and only if $C_S(T) = C_{S_0}(T)$. Since, $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$, the condition $C_S(T) \in \text{Syl}_p(C_G(T))$ is by [4, Proposition I.5.4] equivalent to T being fully $N_{\mathcal{F}}(\tilde{T})$ -centralized. It follows from Lemma 3.25 that this is in turn equivalent to T being fully \mathcal{F} -centralized.

(b) Suppose now that T is fully centralized or equivalently that $C_S(T) = C_{S_0}(T)$. Assume furthermore $N_S(\mathcal{E}) \leq S_0$. By Lemma 3.13(a), we have $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_0|_{N_S(\mathcal{E})}$. Using Corollary 3.15, we can in particular conclude that $C_{\mathcal{F}}(T) = C_{N_{\mathcal{F}}(\mathcal{E})}(T) \subseteq C_{\mathcal{F}_0}(T)$. Moreover, if $P, Q \leq C_{S_0}(T) = C_S(T)$ and $\varphi \in \text{Hom}_{C_{\mathcal{F}_0}(T)}(P, Q)$, then φ extends to $\hat{\varphi} \in \text{Hom}_{\mathcal{F}_0}(PT, QT)$ with $\hat{\varphi}|_T = \text{id}_T$. Notice that PT and QT are contained in $N_S(\mathcal{E})$. Hence, $\hat{\varphi}$ is a morphism in $\mathcal{F}_0|_{N_S(\mathcal{E})} = N_{\mathcal{F}}(\mathcal{E})$. Thus, $\varphi = \hat{\varphi}|_P$ is a morphism in $N_{N_{\mathcal{F}}(\mathcal{E})}(T) = C_{\mathcal{F}}(T)$. This proves (b). \square

Lemma 3.31. *The following properties are equivalent:*

- (i) \mathcal{E} is fully \mathcal{F} -normalized;
- (ii) $N_S(\mathcal{H}) \in \text{Syl}_p(N_G(\mathcal{H}))$;

(iii) \mathcal{E} is fully \mathcal{F} -automized and T is fully centralized.

If so, then $N_{\mathcal{F}}(\mathcal{E})$ is saturated, $\mathcal{E} \trianglelefteq N_{\mathcal{F}}(\mathcal{E})$, $E(N_{\mathcal{F}}(\mathcal{E})) = E(\mathcal{F})$ and $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(N_{\mathcal{F}}(\mathcal{E})), N_S(\mathcal{E}))$ is a regular locality over $N_{\mathcal{F}}(\mathcal{E})$.

Proof. Recall that $N_S(\mathcal{E}) = N_S(\mathcal{H})$ and $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{N_S(\mathcal{H})}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$. In particular, if (ii) holds, then Theorem 3.6(a),(b),(c) applied with $S_0 = N_S(\mathcal{H})$ gives that $N_{\mathcal{F}}(\mathcal{E})$ is saturated, $\mathcal{E} \trianglelefteq N_{\mathcal{F}}(\mathcal{E})$, $E(N_{\mathcal{F}}(\mathcal{E})) = E(\mathcal{F})$ and $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(N_{\mathcal{F}}(\mathcal{E})), N_S(\mathcal{E}))$ is a regular locality over $N_{\mathcal{F}}(\mathcal{E})$. Thus, it remains only to prove the equivalence of properties (i)-(iii). For the proof choose $S_0 \in \text{Syl}_p(N_G(\mathcal{H}))$ such that $N_S(\mathcal{E}) = N_S(\mathcal{H}) \leq S_0$.

(ii) \iff (iii) Observe that

$$|\text{Aut}_{N_S(\mathcal{E})}(T)| \cdot |C_S(T)| = |N_S(\mathcal{E})| \leq |S_0| = |\text{Aut}_{S_0}(T)| \cdot |C_{S_0}(T)|. \quad (3.10)$$

The inequality here is an equality if and only if $N_S(\mathcal{E}) = S_0$, i.e. if and only if (ii) holds. We have moreover $\text{Aut}_{N_S(\mathcal{E})}(T) \leq \text{Aut}_{S_0}(T)$ with equality if and only if \mathcal{E} is fully automized (see Lemma 3.29(b)). As $C_S(T) \leq N_S(\mathcal{E}) \leq S_0$, we have moreover $C_S(T) \leq C_{S_0}(T)$. Lemma 3.30(a) gives that $C_S(T) = C_{S_0}(T)$ if and only if T is fully centralized. Hence, the inequality in (3.10) is also an equality if and only if (iii) holds.

(i) \implies (ii) To prove this implication by contraposition, assume (ii) is false, i.e. $N_S(\mathcal{E}) < S_0$. As $S \in \text{Syl}_p(G)$ and S_0 is a p -subgroup of G , it follows from Sylow's Theorem that there exists $g \in G$ with $S_0^g \leq S$. Notice that $T \leq N_S(\mathcal{E}) \leq S_0$. So in particular, $T^g \leq S$ and $\varphi = c_g|_T$ is a morphism in \mathcal{F} . By Lemma 3.26(a), we have $\mathcal{E}^\varphi = \mathcal{F}_{T^g}(\mathcal{H}^g)$. By Lemma 3.26(b), $S_0^g \leq N_G(\mathcal{H}^g)$ and thus $S_0^g \leq N_S(\mathcal{H}^g) = N_S(\mathcal{E}^\varphi)$ by Lemma 3.8. Hence, $|N_S(\mathcal{E})| < |S_0| = |S_0^g| \leq |N_S(\mathcal{E}^\varphi)|$ and therefore \mathcal{E} is not fully \mathcal{F} -normalized.

(ii) \implies (i) Assume (ii) and fix $\mathcal{E}' \in \mathcal{E}^{\mathcal{F}}$ such that \mathcal{E}' is fully \mathcal{F} -normalized. It is sufficient to show that $|N_S(\mathcal{E}')| \leq |N_S(\mathcal{E})|$. By Lemma 3.25 we have $\mathcal{E}' \in \mathcal{E}^{N_{\mathcal{F}}(\tilde{T})}$. As $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$, there exists thus $g \in G$ with $T^g \leq S$ such that, setting $\varphi = c_g|_T$, we have $\mathcal{E}' = \mathcal{E}^\varphi$. Then Lemma 3.26(a) gives $\mathcal{E}' = \mathcal{F}_{T^g}(\mathcal{H}^g)$. By Lemma 3.26(b), c_g induces an isomorphism from $N_G(\mathcal{H})$ to $N_G(\mathcal{H}^g)$. By Lemma 3.8, $N_S(\mathcal{E}') = N_S(\mathcal{H}^g)$ is a p -subgroup of $N_G(\mathcal{H}^g)$. So (ii) implies $|N_S(\mathcal{E}')| \leq |N_S(\mathcal{H})| = |N_S(\mathcal{E})|$ as required, \square

Proof of Theorem C. Part (a) is a restatement of Lemma 3.13(c), part (b) was shown as Proposition 3.24, part (c) follows from Lemma 3.31, and part (d) was proved as Lemma 3.14 and Corollary 3.15. If \mathcal{F} is constrained, then by Remark 3.5, \mathcal{L} is a model for \mathcal{F} , \mathcal{H} is a subnormal subgroup of \mathcal{L} realizing \mathcal{E} , $\mathbb{N}_{\mathcal{L}}(\mathcal{H}) = N_{\mathcal{L}}(\mathcal{H})$ and thus $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{N_S(\mathcal{H})}(N_{\mathcal{L}}(\mathcal{H}))$. Part (e) follows thus from Lemma 2.17. \square

We conclude this section with two lemmas which make it easier to move to fully \mathcal{F} -normalized conjugates of \mathcal{E} .

Lemma 3.32. *Let $\beta \in \text{Hom}_{\mathcal{F}}(N_S(\mathcal{E}), S)$. Then $N_S(\mathcal{E})\beta \leq N_S(\mathcal{E}^\beta)$.*

Proof. If $s \in N_S(\mathcal{E})$, then $c_s|_T \in \text{Aut}(\mathcal{E})$. So using [10, Remark 2.1], one can observe that

$$c_{s\beta}|_{T\beta} = \beta^{-1}(c_s|_T)\beta \in \text{Aut}(\mathcal{E}^\beta)$$

and thus $s\beta \in N_S(\mathcal{E}^\beta)$. \square

Lemma 3.33. (a) *Let $\hat{\mathcal{E}} \in \mathcal{E}^{\mathcal{F}}$ such that $\hat{\mathcal{E}}$ is fully \mathcal{F} -normalized. Then there exists $\alpha \in \text{Hom}_{\mathcal{F}}(N_S(\mathcal{E}), S)$ such that $\mathcal{E}^\alpha = \hat{\mathcal{E}}$.*

(b) *Fix $\alpha \in \text{Hom}_{\mathcal{F}}(N_S(\mathcal{E}), S)$ such that \mathcal{E}^α is fully normalized. Then for all $P, Q \leq N_S(\mathcal{E})$,*

$$\text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(P, Q) = \{\varphi \in \text{Hom}_{\mathcal{F}}(P, Q) : \alpha^{-1}\varphi\alpha \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E}^\alpha)}(P\alpha, Q\alpha)\}.$$

In particular, if \mathcal{E} is also fully normalized, then $N_S(\mathcal{E})\alpha = N_S(\mathcal{E}^\alpha)$ and α induces an isomorphism from $N_{\mathcal{F}}(\mathcal{E})$ to $N_{\mathcal{F}}(\mathcal{E}^\alpha)$.

Proof. (a) Let $\hat{\mathcal{E}}$ be a subsystem over \hat{T} . As $\hat{\mathcal{E}} \in \mathcal{E}^{\mathcal{F}}$, there exists $\varphi \in \text{Hom}_{\mathcal{F}}(T, S)$ such that $\hat{T} = T\varphi$ and $\hat{\mathcal{E}} = \mathcal{E}^{\varphi}$. Notice that $(\text{Aut}_{\mathcal{F}}(T) \cap \text{Aut}(\mathcal{E}))^{\varphi} = \text{Aut}_{\mathcal{F}}(\hat{T}) \cap \text{Aut}(\hat{\mathcal{E}})$. In particular, $\text{Aut}_{N_S(\mathcal{E})}(T)^{\varphi}$ is a p -subgroup of $\text{Aut}_{\mathcal{F}}(\hat{T}) \cap \text{Aut}(\hat{\mathcal{E}})$. As $\hat{\mathcal{E}}$ is fully normalized, Lemma 3.31 gives that \hat{T} is fully centralized and $\text{Aut}_S(\hat{T}) \cap \text{Aut}(\mathcal{E}) \in \text{Syl}_p(\text{Aut}_{\mathcal{F}}(\hat{T}) \cap \text{Aut}(\hat{\mathcal{E}}))$. Hence, by Sylow's Theorem there exists $\gamma \in \text{Aut}_{\mathcal{F}}(\hat{T}) \cap \text{Aut}(\hat{\mathcal{E}})$ such that $\text{Aut}_{N_S(\mathcal{E})}(T)^{\varphi\gamma} \leq \text{Aut}_S(\hat{T})$. This means that $N_S(\mathcal{E}) \leq N_{\varphi\gamma}$. Hence, by the extension axiom [4, Proposition I.2.5], $\varphi\gamma$ extends to $\alpha \in \text{Hom}_{\mathcal{F}}(N_S(\mathcal{E}), S)$. As $\gamma \in \text{Aut}(\hat{\mathcal{E}})$, we have $\mathcal{E}^{\alpha} = \mathcal{E}^{\varphi\gamma} = \hat{\mathcal{E}}^{\gamma} = \hat{\mathcal{E}}$. This proves (a).

(b) For the proof of (b) let $\alpha \in \text{Hom}_{\mathcal{F}}(N_S(\mathcal{E}), S)$ be arbitrary such that \mathcal{E}^{α} is fully normalized. By Lemma 3.32, $N_S(\mathcal{E})\alpha \leq N_S(\mathcal{E}^{\alpha})$. If \mathcal{E} is fully normalized, then $|N_S(\mathcal{E})\alpha| = |N_S(\mathcal{E})| \geq |N_S(\mathcal{E}^{\alpha})|$ and so $N_S(\mathcal{E})\alpha = N_S(\mathcal{E}^{\alpha})$.

By Lemma 3.2(a), we have $\tilde{T} = T_0T_0^{\perp} \subseteq TC_S(\mathcal{H}) \leq N_S(\mathcal{H}) = N_S(\mathcal{E})$. As $\tilde{T} = E(\mathcal{L}) \cap S$ is strongly \mathcal{F} -closed, it follows that $\tilde{T}\alpha = \tilde{T}$ and α is a morphism in $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$. Thus, we may pick $g \in G$ such that $\alpha = c_g|_{N_S(\mathcal{E})}$.

Recall that $N_S(\mathcal{E}) = N_S(\mathcal{H})$. By Lemma 3.26(a), we have $\mathcal{E}^{\alpha} = \mathcal{F}_{T^g}(\mathcal{H}^g)$ and $T^g = \mathcal{H}^g \cap S$. In particular, $N_S(\mathcal{E}^{\alpha}) = N_S(\mathcal{H}^g)$ by Lemma 3.8. As stated in Lemma 3.26(b), c_g induces an isomorphism from $N_G(\mathcal{H})$ to $N_G(\mathcal{H}^g)$. As \mathcal{E}^{α} is fully normalized, it follows from Lemma 3.31 that $N_S(\mathcal{E}^{\alpha}) = N_S(\mathcal{H}^g) \in \text{Syl}_p(N_G(\mathcal{H}^g))$. Hence, $N_S(\mathcal{E}^{\alpha})^{g^{-1}}$ is a Sylow p -subgroup of $N_G(\mathcal{H})$. This allows us to choose S_0 such that

$$S_0 = N_S(\mathcal{E}^{\alpha})^{g^{-1}}.$$

By Lemma 3.32, we have $N_S(\mathcal{E})^g = N_S(\mathcal{E})\alpha \leq N_S(\mathcal{E}^{\alpha})$ and thus $N_S(\mathcal{E}) \leq S_0$. Now by Lemma 3.26(b), c_g induces an isomorphism from $\mathbb{N}_{\mathcal{L}}(\mathcal{H})$ to $\mathbb{N}_{\mathcal{L}}(\mathcal{H}^g)$ with $S_0c_g = N_S(\mathcal{E}^{\alpha}) = N_S(\mathcal{H}^g)$. Hence, using Theorem 3.6(a), it follows from Lemma 2.11 that c_g induces an isomorphism from $\mathcal{F}_0 := \mathcal{F}_{S_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$ to $N_{\mathcal{F}}(\mathcal{E}^{\alpha}) = \mathcal{F}_{N_S(\mathcal{H}^g)}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}^g))$.

If \mathcal{E} is fully normalized, then Lemma 3.31 implies $N_S(\mathcal{E}) = S_0$ and thus $\mathcal{F}_0 = N_{\mathcal{F}}(\mathcal{E})$, i.e. α induces an isomorphism from $N_{\mathcal{F}}(\mathcal{E})$ to $N_{\mathcal{F}}(\mathcal{E}^{\alpha})$. In the general case, Lemma 3.13(a) gives $\mathcal{F}_0 \cap \mathcal{F} = N_{\mathcal{F}}(\mathcal{E})$. Given $\varphi \in \text{Hom}_{\mathcal{F}}(P, Q)$, we have thus $\varphi \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E})}(P, Q)$ if and only if $\varphi \in \text{Hom}_{\mathcal{F}_0}(P, Q)$, and this is the case if and only if $\alpha^{-1}\varphi\alpha = (c_g|_{S_0})^{-1}\varphi(c_g|_{S_0}) \in \text{Hom}_{N_{\mathcal{F}}(\mathcal{E}^{\alpha})}(P\alpha, Q\alpha)$. This proves the assertion. \square

3.5. The centralizer of \mathcal{E} and fully centralized conjugates of \mathcal{E} . We continue to assume Hypotheses 3.1 and 3.10. In particular, S_0 is a Sylow p -subgroup of $N_G(\mathcal{H})$ containing T .

If \mathcal{E} is normal in \mathcal{F} , then it was first shown by Aschbacher that there is a unique largest element of the set

$$\{R \leq S : \mathcal{E} \subseteq C_{\mathcal{F}}(R)\},$$

which we denote by $C_S(\mathcal{E}) = C_S(\mathcal{E})_{\mathcal{F}}$ (cf. [2, Chapter 6] or [11, Theorem 1(a)]). We extend this result now to the case that \mathcal{E} is subnormal in \mathcal{F} . As \mathcal{F}_0 is saturated and $\mathcal{E} \trianglelefteq \mathcal{F}_0$, we can use that the subgroup $C_{S_0}(\mathcal{E})_{\mathcal{F}_0}$ is defined as the largest element of the set $\{R \leq S_0 : \mathcal{E} \subseteq C_{\mathcal{F}_0}(R)\}$. Notice that the following proposition implies Theorem E.

Proposition 3.34. *The subgroup $C_S(\mathcal{H})$ is with respect to inclusion a maximal element of the set $\mathcal{X} := \{R \leq S : \mathcal{E} \subseteq C_{\mathcal{F}}(R)\}$. If $C_S(\mathcal{H}) \leq S_0$, then $C_S(\mathcal{H}) = S \cap C_{S_0}(\mathcal{E})_{\mathcal{F}_0}$.*

Proof. By [13, Lemma 3.5], we have $\mathcal{H} \subseteq C_{\mathcal{L}}(C_S(\mathcal{H}))$, which implies easily that $\mathcal{E} = \mathcal{F}_T(\mathcal{H}) \subseteq C_{\mathcal{F}}(C_S(\mathcal{H}))$. Hence, $C_S(\mathcal{H}) \in \mathcal{X}$.

Choose now S_0 such that $C_S(\mathcal{H}) \leq S_0$. Recall from Theorem 3.6(a),(b) that \mathcal{F}_0 is saturated and $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 with $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$, $T = S_0 \cap \mathcal{H}$ and $\mathcal{E} \trianglelefteq \mathcal{F}_0$. Hence, it follows from [12, Proposition 4] that $C_{S_0}(\mathcal{E})_{\mathcal{F}_0} = C_{S_0}(\mathcal{H})$. As $C_S(\mathcal{H}) \leq S_0$ by assumption, we can conclude that $C_S(\mathcal{H}) = S \cap C_{S_0}(\mathcal{H}) = S \cap C_{S_0}(\mathcal{E})_{\mathcal{F}_0}$.

Choose now S_0 such that $N_S(\mathcal{E}) \leq S_0$. We have then $N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F}_0$ by Lemma 3.13(a). Let $R \in \mathcal{X}$. Then \mathcal{E} and $\mathcal{F}_R(R)$ commute in the sense of Definition 4. Hence, by Lemma 2.4, $\mathcal{D} := \mathcal{E} * \mathcal{F}_R(R)$ is a saturated subsystem of \mathcal{F} over TR with $\mathcal{E} \trianglelefteq \mathcal{D}$ and $\mathcal{E} \subseteq C_{\mathcal{D}}(R)$. Proposition 3.24 yields now $\mathcal{D} \subseteq N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F}_0$. In particular, $R \leq N_S(\mathcal{E}) \leq S_0$. As $\mathcal{E} \subseteq C_{\mathcal{D}}(R) \subseteq C_{\mathcal{F}_0}(R)$, it follows $R \leq C_{S_0}(\mathcal{E})_{\mathcal{F}_0}$. Hence, $R \leq S \cap C_{S_0}(\mathcal{E})_{\mathcal{F}_0} = C_S(\mathcal{H})$. This proves the assertion. \square

Definition 3.35. We write $C_S(\mathcal{E})$ or $C_S(\mathcal{E})_{\mathcal{F}}$ for $C_S(\mathcal{H})$, i.e. for the maximal element of the set

$$\{R \leq S : \mathcal{E} \subseteq C_{\mathcal{F}}(R)\}.$$

We call $C_S(\mathcal{E}) = C_S(\mathcal{E})_{\mathcal{F}}$ the *centralizer of \mathcal{E} in S* .

The reader might want to observe that

$$C_S(\mathcal{E}) = C_S(\mathcal{H}) \leq N_S(\mathcal{H}) = N_S(\mathcal{E})$$

by Lemma 3.8.

Recall from the introduction that $C_{\mathcal{F}}(\mathcal{E})$ is defined as

$$C_{\mathcal{F}}(\mathcal{E}) := \mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H})).$$

We call $C_{\mathcal{F}}(\mathcal{E})$ the *centralizer of \mathcal{E} in \mathcal{F}* .

Note that $C_{\mathcal{F}}(\mathcal{E})$ depends a priori not only on \mathcal{F} and \mathcal{E} , but also on \mathcal{L} . However, the properties we show below imply that $C_{\mathcal{F}}(\mathcal{E})$ is indeed independent of the choice of \mathcal{L} . Alternatively, this can be seen more directly using Background Theorems 1 and 3 combined with Lemma 2.11.

The reader is referred to [4, Definition I.7.1] for the definition of the focal subgroup $\text{foc}(\mathcal{F})$ and the hyperfocal subgroup $\text{hnp}(\mathcal{F})$. We extend the definition of the focal subgroup to fusion systems which are not saturated. Namely, for every fusion system \mathcal{G} over a p -group X , we set

$$\text{foc}(\mathcal{G}) := \langle x^{-1}(x\varphi) : x \in X, \varphi \in \text{Hom}_{\mathcal{F}}(\langle x \rangle, X) \rangle.$$

It is always true that $\text{hnp}(\mathcal{F}) \leq \text{foc}(\mathcal{F})$. For every subgroup $R \leq S$ with $\text{hnp}(\mathcal{F}) \leq R$, it is shown in [4, Theorem I.7.4] that

$$\mathcal{F}_R := \langle O^p(\text{Aut}_{\mathcal{F}}(P)) : P \leq R \rangle_R \quad (3.11)$$

is the unique saturated subsystem of \mathcal{F} over R of p -power index. This concept will play a role in obtaining a description of $C_{\mathcal{F}}(\mathcal{E})$ if \mathcal{E} is fully centralized.

If \mathcal{E} is normal in \mathcal{F} , then a subsystem $C_{\mathcal{F}}(\mathcal{E})$ was first defined by Aschbacher [2, Chapter 6], and a new construction of $C_{\mathcal{F}}(\mathcal{E})$ was given in [11]. However, if \mathcal{E} is normal, then by Background Theorem 2, we have $\mathcal{H} \trianglelefteq \mathcal{L}$. Hence, [8, Proposition 6.7] states in this case that the subsystem $C_{\mathcal{F}}(\mathcal{E})$ defined in [2, 11] equals $\mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))$ and coincides thus with the centralizer $C_{\mathcal{F}}(\mathcal{E})$ defined in this paper. Using this, the following theorem follows from [11, Proposition 1, Theorem 2] and the construction of $C_{\mathcal{F}}(\mathcal{E})$ given in [11].

Theorem 3.36. *Suppose $\mathcal{E} \trianglelefteq \mathcal{F}$. Then $\text{hnp}(C_{\mathcal{F}}(T)) \leq \text{foc}(C_{\mathcal{F}}(T)) \leq C_S(\mathcal{E})$. Moreover,*

$$C_{\mathcal{F}}(\mathcal{E}) = C_{\mathcal{F}}(T)_{C_S(\mathcal{E})}$$

is normal in \mathcal{F} . Every saturated subsystem of \mathcal{F} which commutes with \mathcal{E} is contained in $C_{\mathcal{F}}(\mathcal{E})$. \square

We use here the notation introduced in (3.11) for $C_{\mathcal{F}}(T)$ in place of \mathcal{F} . Notice that this is possible, as T is in the situation above fully centralized and thus $C_{\mathcal{F}}(T)$ is saturated.

Proof. By Lemma 3.31, $N_{\mathcal{F}}(\mathcal{E})$ is saturated, $\mathcal{E} \trianglelefteq N_{\mathcal{F}}(\mathcal{E})$ and $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(N_{\mathcal{F}}(\mathcal{E})), N_S(\mathcal{E}))$ is a regular locality over $N_{\mathcal{F}}(\mathcal{E})$. By Lemma 3.9, we have $C_{\mathcal{L}}(\mathcal{H}) = C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\mathcal{H})$ and thus $C_{\mathcal{F}}(\mathcal{E}) = C_{N_{\mathcal{F}}(\mathcal{E})}(\mathcal{E})$ by definition. It follows from Lemma 3.14 that $C_{\mathcal{F}}(T) = C_{N_{\mathcal{F}}(\mathcal{E})}(T)$. Hence, the assertion follows from Theorem 3.36. \square

Lemma 3.37. *The centralizer $C_{\mathcal{F}_0}(\mathcal{E})$ is normal in \mathcal{F}_0 , $\text{foc}(C_{\mathcal{F}_0}(T)) \leq C_{S_0}(\mathcal{E}) := C_{S_0}(\mathcal{E})_{\mathcal{F}_0}$ and*

$$C_{\mathcal{F}_0}(\mathcal{E}) = \mathcal{F}_{C_{S_0}(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H})) = C_{\mathcal{F}_0}(T)C_{S_0}(\mathcal{E}).$$

Moreover, $(C_{\mathcal{L}}(\mathcal{H}), \delta(C_{\mathcal{F}_0}(\mathcal{E})), C_{S_0}(\mathcal{H}))$ is a regular locality over $C_{\mathcal{F}_0}(\mathcal{E})$. In particular, $C_{S_0}(\mathcal{H}) = C_{S_0}(\mathcal{E})_{\mathcal{F}_0}$ is a maximal p -subgroup of $C_{\mathcal{L}}(\mathcal{H})$.

Proof. As \mathcal{F}_0 is saturated and $\mathcal{E} \trianglelefteq \mathcal{F}_0$, it follows from Theorem 3.36 that $C_{\mathcal{F}_0}(\mathcal{E}) = C_{\mathcal{F}_0}(T)C_{S_0}(\mathcal{E})$ is normal in \mathcal{F}_0 . Recall that $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 and $C_{\mathcal{L}}(\mathcal{H}) = C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\mathcal{H})$ by Lemma 3.9. Hence, the remaining part of the assertion follows from Background Theorem 3 and the definition $C_{\mathcal{F}_0}(\mathcal{E})$. \square

Recall that \mathcal{E} is called *fully \mathcal{F} -centralized* if

$$|C_S(\mathcal{E})| \geq |C_S(\tilde{\mathcal{E}})| \text{ for all } \tilde{\mathcal{E}} \in \mathcal{E}^{\mathcal{F}}.$$

Lemma 3.38.

- (a) *Choose S_0 such that $C_S(\mathcal{H}) = C_S(\mathcal{E}) \leq S_0$. Then the following are equivalent:*
 - (i) \mathcal{E} is fully \mathcal{F} -centralized;
 - (ii) $C_S(\mathcal{H}) = C_{S_0}(\mathcal{H})$;
 - (iii) $C_S(\mathcal{H})$ is a maximal p -subgroup of $C_{\mathcal{L}}(\mathcal{H})$.
- (b) *Let $\hat{\mathcal{E}} \in \mathcal{E}^{\mathcal{F}}$ such that $\hat{\mathcal{E}}$ is fully \mathcal{F} -centralized. Then there exists $\alpha \in \text{Hom}_{\mathcal{F}}(TC_S(\mathcal{E}), S)$ such that $\mathcal{E}^\alpha = \hat{\mathcal{E}}$.*
- (c) *If $\alpha \in \text{Hom}_{\mathcal{F}}(TC_S(\mathcal{E}), S)$, then α is a morphism in $N_{\mathcal{F}}(\tilde{T})$ and $C_S(\mathcal{E})\alpha \leq C_S(\mathcal{E}^\alpha)$.*

Proof. **(a,b)** Choose S_0 such that $C_S(\mathcal{H}) \leq S_0$. By Lemma 3.37, $C_{S_0}(\mathcal{H}) \geq C_S(\mathcal{H})$ is a maximal p -subgroup of $C_{\mathcal{L}}(\mathcal{H})$. Hence, (ii) and (iii) are equivalent. Thus, it remains only to show that (i) and (ii) are equivalent and that (c) holds.

Step 1: We show that (i) implies (ii). Notice that $TC_{S_0}(\mathcal{H})$ is a p -subgroup of $N_G(\mathcal{H})$. As $S \in \text{Syl}_p(G)$, there exists thus $g \in G$ with $(TC_{S_0}(\mathcal{H}))^g \leq S$. As $T^g \leq S$, $\varphi = c_g|_T$ is an \mathcal{F} -morphism and $T\varphi = T^g \in T^{\mathcal{F}}$. By Lemma 3.26(a), we have $T^g = \mathcal{H}^g \cap S$ and $\mathcal{E}^\varphi = \mathcal{F}_{T^g}(\mathcal{H}^g)$. In particular, $C_S(\mathcal{E}^\varphi) = C_S(\mathcal{H}^g)$. Using Lemma 3.26(b), one observes that $C_{S_0}(\mathcal{H})^g \leq C_S(\mathcal{H}^g)$. Thus, if $C_S(\mathcal{H}) < C_{S_0}(\mathcal{H})$, then $|C_S(\mathcal{E})| = |C_S(\mathcal{H})| < |C_S(\mathcal{H}^g)| = |C_S(\mathcal{E}^\varphi)|$ and \mathcal{E} is thus not fully centralized. So Step 1 is complete.

Step 2: We show that (b) holds and (ii) implies (i). For the proof fix $\hat{\mathcal{E}} \in \mathcal{E}^{\mathcal{F}}$ fully centralized (and notice that such $\hat{\mathcal{E}}$ always exists). By Lemma 3.25, there is $\varphi \in \text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(T, S)$ such that $\mathcal{E}^\varphi = \hat{\mathcal{E}}$. As $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$, there exists $g \in G$ with $\varphi = c_g|_T$. By Lemma 3.26(a), we have $\mathcal{F}_{T^g}(\mathcal{H}^g) = \hat{\mathcal{E}}$ and thus $C_S(\hat{\mathcal{E}}) = C_S(\mathcal{H}^g)$. Fix $\hat{S}_0 \in \text{Syl}_p(N_G(\mathcal{H}^g))$ such that

$$T^g C_S(\mathcal{H}^g) \leq \hat{S}_0.$$

As T is a maximal p -subgroup of \mathcal{H} , the conjugate $T^g = T\varphi$ is a maximal p -subgroup of \mathcal{H}^g and thus equal to $S \cap \mathcal{H}^g$. By Theorem 3.6(b) applied with \mathcal{H}^g in place of \mathcal{H} , we have moreover $\mathcal{H}^g \cap \hat{S}_0 = T^g = S \cap \mathcal{H}^g$. Step 1 applied with $(\hat{\mathcal{E}}, \mathcal{H}^g)$ in place of $(\mathcal{E}, \mathcal{H})$ gives

$$C_S(\mathcal{H}^g) = C_{\hat{S}_0}(\mathcal{H}^g).$$

Notice that S_0^g is a Sylow p -subgroup of $N_G(\mathcal{H}^g)$ as c_g induces an isomorphism from $N_G(\mathcal{H})$ to $N_G(\mathcal{H}^g)$ by Lemma 3.26(b). Hence, there exists $x \in N_G(\mathcal{H}^g)$ such that $S_0^{gx} = \hat{S}_0$. Observe that $\mathcal{H}^g = \mathcal{H}^{gx}$. Moreover, $T^{gx} \leq S_0^{gx} \cap \mathcal{H}^g = \hat{S}_0 \cap \mathcal{H}^g = T^g$ and thus $T^{gx} = T^g \leq S$. Thus, replacing φ and g by c_{gx} and gx , we may assume that $S_0^g = \hat{S}_0$. Hence, using $c_g \in \text{Aut}(\mathcal{L})$, we may conclude that

$$C_S(\hat{\mathcal{E}}) = C_S(\mathcal{H}^g) = C_{\hat{S}_0}(\mathcal{H}^g) = C_{S_0}(\mathcal{H})^g \geq C_S(\mathcal{H})^g = C_S(\mathcal{E})^g. \quad (3.12)$$

So $\alpha := c_g|_{TC_S(\mathcal{E})}$ is an element of $\text{Hom}_{N_{\mathcal{F}}(\tilde{T})}(TC_S(\mathcal{E}), S)$ with $\mathcal{E}^\alpha = \hat{\mathcal{E}}$. This proves (b). If we assume in addition that $C_S(\mathcal{H}) = C_{S_0}(\mathcal{H})$, then we obtain from (3.12) that $C_S(\mathcal{E})^g = C_S(\mathcal{H})^g = C_{S_0}(\mathcal{H})^g = C_S(\mathcal{H}^g) = C_S(\hat{\mathcal{E}})$. As $\hat{\mathcal{E}}$ was chosen to be fully centralized, it follows then that \mathcal{E} is fully centralized. This completes Step 2 and thus the proof of (a) and (b).

(c) By Lemma 3.2(a), we have $\tilde{T} = T_0 T_0^\perp \subseteq TC_S(\mathcal{H}) = TC_S(\mathcal{E})$. As \tilde{T} is strongly \mathcal{F} -closed, it follows that a morphism $\alpha \in \text{Hom}_{\mathcal{F}}(TC_S(\mathcal{E}), S)$ is a morphism in $N_{\mathcal{F}}(\tilde{T})$. As $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$, there exists thus $g \in G$ with $\alpha = c_g|_{TC_S(\mathcal{E})}$. By Lemma 3.26(a), we have $S \cap \mathcal{H}^g = T^g$ and $\mathcal{E}^\alpha = \mathcal{F}_{T^g}(\mathcal{H}^g)$. In particular, $C_S(\mathcal{E}^\alpha) = C_S(\mathcal{H}^g)$. Using Lemma 3.26(b), we can conclude that $C_S(\mathcal{E})\alpha = C_S(\mathcal{E})^g = C_S(\mathcal{H})^g \leq S \cap C_G(\mathcal{H}^g) = C_S(\mathcal{H}^g) = C_S(\mathcal{E}^\alpha)$. \square

Proposition 3.39. *If \mathcal{E} is fully centralized, then $C_{\mathcal{F}}(\mathcal{E})$ is saturated, $\text{foc}(C_{\mathcal{F}}(T)) \leq C_S(\mathcal{E})$ and*

$$C_{\mathcal{F}}(\mathcal{E}) = \langle O^p(\text{Aut}_{C_{\mathcal{F}}(T)}(P)) : P \leq C_S(\mathcal{E}) \rangle_{C_S(\mathcal{E})}.$$

If in addition S_0 is chosen such that $C_S(\mathcal{E}) \leq S_0$, then $C_{\mathcal{F}}(\mathcal{E}) = C_{\mathcal{F}_0}(\mathcal{E})$ is normal in \mathcal{F}_0 .

Proof. Choose first S_0 such that $C_S(\mathcal{E}) \leq S_0$. Assuming that \mathcal{E} is fully \mathcal{F} -centralized, part (a) gives $C_{S_0}(\mathcal{E})_{\mathcal{F}_0} = C_{S_0}(\mathcal{H}) = C_S(\mathcal{H}) = C_S(\mathcal{E})$. Hence, using the definition of $C_{\mathcal{F}}(\mathcal{E})$ and Lemma 3.37, we can conclude that

$$\text{foc}(C_{\mathcal{F}_0}(T)) \leq C_S(\mathcal{E}), \quad (3.13)$$

and that

$$C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H})) = \mathcal{F}_{C_{S_0}(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H})) = C_{\mathcal{F}_0}(\mathcal{E}) = C_{\mathcal{F}_0}(T)_{C_S(\mathcal{E})} \quad (3.14)$$

is saturated and normal in \mathcal{F}_0 . It remains to show that $\text{foc}(C_{\mathcal{F}}(T)) \leq C_S(\mathcal{E})$ and $C_{\mathcal{F}}(\mathcal{E}) = \langle O^p(\text{Aut}_{C_{\mathcal{F}}(T)}(P)) : P \leq C_S(\mathcal{E}) \rangle_{C_S(\mathcal{E})}$. For the proof we may choose S_0 such that $N_S(\mathcal{E}) \leq S_0$. By Corollary 3.16, we have then

$$C_{\mathcal{F}}(T) = C_{\mathcal{F}_0}(T)|_{C_S(T)}.$$

In particular, if $x \leq C_S(T)$ and $\varphi \in \text{Hom}_{C_{\mathcal{F}}(T)}(x, C_S(T))$, then φ is a morphism in $C_{\mathcal{F}_0}(T)$ and $x^{-1}(x\varphi) \in \text{foc}(C_{\mathcal{F}_0}(T)) \leq C_S(\mathcal{E})$ by (3.13). This shows $\text{foc}(C_{\mathcal{F}}(T)) \leq C_S(\mathcal{E})$.

As $C_S(\mathcal{E}) \leq C_S(T)$, it follows moreover that $\text{Aut}_{C_{\mathcal{F}_0}(T)}(P) = \text{Aut}_{C_{\mathcal{F}}(T)}(P)$ for every $P \leq C_S(\mathcal{E})$. Hence, using (3.11) and (3.14), we can conclude that

$$\begin{aligned} C_{\mathcal{F}}(\mathcal{E}) &= C_{\mathcal{F}_0}(T)_{C_S(\mathcal{E})} = \langle O^p(\text{Aut}_{C_{\mathcal{F}_0}(T)}(P)) : P \leq C_S(\mathcal{E}) \rangle_{C_S(\mathcal{E})} \\ &= \langle O^p(\text{Aut}_{C_{\mathcal{F}}(T)}(P)) : P \leq C_S(\mathcal{E}) \rangle_{C_S(\mathcal{E})}. \end{aligned}$$

This proves the assertion. \square

Lemma 3.40.

- (a) *Suppose T is fully \mathcal{F} -centralized. Then \mathcal{E} is fully \mathcal{F} -centralized, $\text{foc}(C_{\mathcal{F}}(T)) \leq C_S(\mathcal{E})$ and $C_{\mathcal{F}}(\mathcal{E}) = C_{\mathcal{F}}(T)_{C_S(\mathcal{E})}$.*
- (b) *If \mathcal{E} is fully \mathcal{F} -normalized, then $C_{\mathcal{F}}(\mathcal{E}) = C_{N_{\mathcal{F}}(\mathcal{E})}(\mathcal{E}) \trianglelefteq N_{\mathcal{F}}(\mathcal{E})$.*

Proof. (a) Assume T is fully \mathcal{F} -centralized and choose S_0 such that $C_S(T) \leq S_0$. Lemma 3.30(a) gives then that $C_S(T) = C_{S_0}(T)$. In particular, $C_S(\mathcal{H}) = C_{C_S(T)}(\mathcal{H}) = C_{C_{S_0}(T)}(\mathcal{H}) = C_{S_0}(\mathcal{H})$. Hence, by Lemma 3.38(a), \mathcal{E} is fully centralized. Using that $C_{\mathcal{F}}(T)$ is saturated, part (a) follows now from Proposition 3.39 and (3.11).

(b) If \mathcal{E} is fully normalized, then Lemma 3.31 yields $N_S(\mathcal{E}) = N_S(\mathcal{H}) \in \text{Syl}_p(N_G(\mathcal{H}))$. Hence, we may assume that $S_0 = N_S(\mathcal{E})$ and thus $\mathcal{F}_0 = N_{\mathcal{F}}(\mathcal{E})$. The assertion follows then from Lemma 3.37. \square

We now want to consider the properties of $C_{\mathcal{F}}(\mathcal{E})$ if \mathcal{E} is not necessarily fully centralized or fully normalized. The reader might want to recall Definition 2.2.

Lemma 3.41. *Suppose $C_S(\mathcal{H}) = C_S(\mathcal{E}) \leq S_0$. Then*

$$C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \cap C_{\mathcal{F}_0}(\mathcal{E}) = C_{\mathcal{F}_0}(\mathcal{E})|_{C_S(\mathcal{E})}.$$

In particular, $C_{\mathcal{F}}(\mathcal{E})$ is weakly $N_{\mathcal{F}}(\mathcal{E})$ -invariant.

Proof. By Lemma 3.37, $\mathcal{C} := C_{\mathcal{F}_0}(\mathcal{E}) = \mathcal{F}_{C_{S_0}(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))$ is a normal subsystem of \mathcal{F}_0 . In particular, \mathcal{C} is weakly \mathcal{F}_0 -invariant, which implies that $\mathcal{F} \cap \mathcal{C}$ is weakly $(\mathcal{F} \cap \mathcal{F}_0)$ -invariant. By Lemma 3.13(a), we have $N_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \cap \mathcal{F}_0$ if S_0 is chosen such that $N_S(\mathcal{E}) \leq S_0$. Hence, it is sufficient to prove that $C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \cap \mathcal{C} = \mathcal{C}|_{C_S(\mathcal{E})}$.

By Proposition 3.34, both $C_{\mathcal{F}}(\mathcal{E})$ and $\mathcal{F} \cap \mathcal{C}$ are fusion systems over

$$C_S(\mathcal{H}) = C_S(\mathcal{E}) = S \cap C_{S_0}(\mathcal{E})_{\mathcal{F}_0} = S \cap C_{S_0}(\mathcal{H}).$$

Notice that $C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))$ is contained in $\mathcal{C} = \mathcal{F}_{C_{S_0}(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))$ and thus in $\mathcal{F} \cap \mathcal{C}$. Hence

$$C_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{C} \cap \mathcal{F} \subseteq \mathcal{C}|_{C_S(\mathcal{E})}$$

and it remains to show that $\mathcal{C}|_{C_S(\mathcal{E})} \subseteq C_{\mathcal{F}}(\mathcal{E})$.

Recall from Lemma 3.6(a),(b),(c) that $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \delta(\mathcal{F}_0), S_0)$ is a regular locality over \mathcal{F}_0 with $\mathcal{H} \trianglelefteq \mathbb{N}_{\mathcal{L}}(\mathcal{H})$ and $\text{Comp}(\mathcal{L}) = \text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}))$. Since $C_{\mathcal{L}}(\mathcal{H}) = C_{\mathbb{N}_{\mathcal{L}}(\mathcal{H})}(\mathcal{H})$ by Lemma 3.9, it follows from [13, Lemma 11.16, Theorem 10.16(e)] (applied with $(\mathbb{N}_{\mathcal{L}}(\mathcal{H}), \mathcal{H})$ in place of $(\mathcal{L}, \mathcal{N})$) that

$$\text{Comp}(C_{\mathcal{L}}(\mathcal{H})) = \text{Comp}(\mathbb{N}_{\mathcal{L}}(\mathcal{H})) \setminus \text{Comp}(\mathcal{H}) = \text{Comp}(\mathcal{L}) \setminus \text{Comp}(\mathcal{H}).$$

Thus, $E(C_{\mathcal{L}}(\mathcal{H})) = \mathcal{M}$ by definition of \mathcal{M} . Here \mathcal{M} is subnormal in \mathcal{L} as $\mathcal{M} \trianglelefteq E(\mathcal{L}) \trianglelefteq \mathcal{L}$ by Lemma 3.2(a). Thus, it follows from Background Theorem 2 and the definition of a locality that $T_0^\perp = \mathcal{M} \cap S$ is a maximal p -subgroup of \mathcal{M} . As $T_0^\perp \leq C_S(\mathcal{H}) \leq C_{S_0}(\mathcal{H})$ by Lemma 3.2(a), it follows that $T_0^\perp = C_{S_0}(\mathcal{H}) \cap \mathcal{M}$. Hence, $E(\mathcal{C}) = \mathcal{F}_{T_0^\perp}(\mathcal{M})$ by Background Theorem 3 applied with $(C_{\mathcal{L}}(\mathcal{H}), C_{S_0}(\mathcal{H}), \mathcal{C})$ in place of $(\mathcal{L}, S, \mathcal{F})$.

Let now $P, Q \leq C_S(\mathcal{E}) = C_S(\mathcal{H})$ and $\varphi \in \text{Hom}_{\mathcal{C}}(P, Q)$. We need to show that φ is a morphism in $N_{\mathcal{F}}(\mathcal{E})$. By Lemma 2.9 (applied with \mathcal{C} and $E(\mathcal{C})$ in place of \mathcal{F} and \mathcal{E}), there exist $\psi \in \text{Hom}_{E(\mathcal{C})P}(P, PT_0^\perp)$ and a morphism $\alpha: P\psi \rightarrow Q$ in $N_{\mathcal{C}}(T_0^\perp)$ such that $\varphi = \psi\alpha$. By [8, Theorem D(a)], we have $E(\mathcal{C})P = \mathcal{F}_{T_0^\perp P}(\mathcal{M}P)$. Notice that $T_0^\perp P \leq C_S(\mathcal{H})$ and $\mathcal{M}P \subseteq C_{\mathcal{L}}(\mathcal{H})$. Hence, $E(\mathcal{C})P$ is contained in $C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))$ and ψ is a morphism in $C_{\mathcal{F}}(\mathcal{E})$. In particular, $P\psi \leq C_S(\mathcal{H})$.

By Lemma 2.13 applied with $C_{\mathcal{L}}(\mathcal{H})$ in place of \mathcal{L} , we have $T_0^\perp = \mathcal{M} \cap S_0 = E(C_{\mathcal{L}}(\mathcal{H})) \cap S_0 \in \delta(\mathcal{C})$. Hence, by [9, Lemma 3.10(b)], $N_{\mathcal{C}}(T_0^\perp)$ is realized by the group $N_{C_{\mathcal{L}}(\mathcal{H})}(T_0^\perp)$. In particular, $\alpha = c_g|_{P\psi}$ for some $g \in N_{C_{\mathcal{L}}(\mathcal{H})}(T_0^\perp) \subseteq C_{\mathcal{L}}(\mathcal{H})$. As $P\psi$ and Q are contained in $C_S(\mathcal{H}) = C_S(\mathcal{E})$, it follows that α is a morphism in $C_{\mathcal{F}}(\mathcal{E})$. Hence, $\varphi = \psi\alpha$ is a morphism in $C_{\mathcal{F}}(\mathcal{E})$. This shows $\mathcal{C}|_{C_S(\mathcal{E})} \subseteq C_{\mathcal{F}}(\mathcal{E})$ and thus $C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \cap \mathcal{C} = \mathcal{C}|_{C_S(\mathcal{E})}$ as required. \square

Lemma 3.42. *Every saturated subsystem of \mathcal{F} which commutes with \mathcal{E} is contained in $C_{\mathcal{F}}(\mathcal{E})$.*

Proof. Let \mathcal{G} be a saturated subsystem of \mathcal{F} which commutes with \mathcal{E} . Then by Lemma 2.4, $\mathcal{D} := \mathcal{E} * \mathcal{G}$ is a saturated subsystem of \mathcal{F} such that $\mathcal{E} \trianglelefteq \mathcal{D}$, and moreover \mathcal{E} and \mathcal{G} commute in \mathcal{D} . Hence, by Proposition 3.24, $\mathcal{D} \subseteq N_{\mathcal{F}}(\mathcal{E})$. Choose now S_0 such that $N_S(\mathcal{E}) \leq S_0$. Then $\mathcal{D} \subseteq N_{\mathcal{F}}(\mathcal{E}) \subseteq \mathcal{F}_0$. As \mathcal{E} and \mathcal{G} commute in \mathcal{D} , they commute also in \mathcal{F}_0 . Since $\mathcal{E} \trianglelefteq \mathcal{F}_0$, it follows now from Theorem 3.36 that $\mathcal{G} \subseteq C_{\mathcal{F}_0}(\mathcal{E})$. Using Lemma 3.41, we can conclude that $\mathcal{G} \subseteq \mathcal{F} \cap C_{\mathcal{F}_0}(\mathcal{E}) = C_{\mathcal{F}}(\mathcal{E})$. \square

Proof of Theorem F. We have $C_{\mathcal{L}}(\mathcal{H}) \subseteq C_{\mathcal{L}}(T)$ and, by [13, Lemma 3.5], $\mathcal{H} \subseteq C_{\mathcal{L}}(C_S(\mathcal{H}))$. As $\mathcal{E} = \mathcal{F}_T(\mathcal{H})$ and $C_{\mathcal{F}}(\mathcal{E}) = \mathcal{F}_{C_S(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))$, it follows that $C_{\mathcal{F}}(\mathcal{E}) \subseteq C_{\mathcal{F}}(T)$ and $\mathcal{E} \subseteq C_{\mathcal{F}}(C_S(\mathcal{H}))$, i.e. \mathcal{E} and $C_{\mathcal{F}}(\mathcal{E})$ commute in \mathcal{F} . Lemma 3.9 implies that $C_{\mathcal{F}}(\mathcal{E})$ is contained in $N_{\mathcal{F}}(\mathcal{E})$, and clearly $C_S(\mathcal{E})$ is a fusion system over $C_S(\mathcal{H}) = C_S(\mathcal{E})$. Now part (a) follows from Lemma 3.42.

Part (b) follows from Lemma 3.40(b) and Lemma 3.41. Part (c) is a consequence of (a) and Proposition 3.39. \square

The following lemma together with Lemma 3.38(b),(c) yields a description of $C_{\mathcal{F}}(\mathcal{E})$ in terms of the fusion systems \mathcal{E} and \mathcal{F} .

Lemma 3.43. *Fix $\alpha \in \text{Hom}_{\mathcal{F}}(TC_S(\mathcal{E}), S)$ such that \mathcal{E}^α is fully \mathcal{F} -centralized. Then for all $P, Q \leq C_S(\mathcal{E})$,*

$$\text{Hom}_{C_{\mathcal{F}}(\mathcal{E})}(P, Q) = \{\varphi \in \text{Hom}_{\mathcal{F}}(P, Q) : \alpha^{-1}\varphi\alpha \in \text{Hom}_{C_{\mathcal{F}}(\mathcal{E}^\alpha)}(P\alpha, Q\alpha)\}.$$

In particular, if \mathcal{E} is also fully \mathcal{F} -centralized, then $C_S(\mathcal{E})\alpha = C_S(\mathcal{E}^\alpha)$ and α induces an isomorphism from $C_{\mathcal{F}}(\mathcal{E})$ to $C_{\mathcal{F}}(\mathcal{E}^\alpha)$.

Proof. Let $\alpha \in \text{Hom}_{\mathcal{F}}(TC_S(\mathcal{E}), S)$ such that \mathcal{E}^α is fully centralized. By Lemma 3.38(c), α is a morphism in $N_{\mathcal{F}}(\tilde{T}) = \mathcal{F}_S(G)$ and $C_S(\mathcal{E})\alpha \leq C_S(\mathcal{E}^\alpha)$. In particular, there exists $g \in G$ such that $\alpha = c_g|_{TC_S(\mathcal{E})}$.

By Lemma 3.26(a), we have $\mathcal{E}^\alpha = \mathcal{F}_{T^g}(\mathcal{H}^g)$ and $T^g = \mathcal{H}^g \cap S$. Fix $\hat{S}_0 \in \text{Syl}_p(N_G(\mathcal{H}^g))$ such that $T^g C_S(\mathcal{H}^g) \leq N_S(\mathcal{H}^g) = N_S(\mathcal{E}^\alpha) \leq \hat{S}_0$. As \mathcal{E}^α is fully centralized, it follows from Lemma 3.38(a) that $C_S(\mathcal{H}^g) = C_{\hat{S}_0}(\mathcal{H}^g)$. As G acts on \mathcal{L} , one can observe that c_g induces an isomorphism from $N_G(\mathcal{H})$ to $N_G(\mathcal{H}^g)$. Thus, $\hat{S}_0^{g^{-1}}$ is a Sylow p -subgroup of $N_G(\mathcal{H})$ containing T . Hence, we may choose S_0 such that

$$S_0 := \hat{S}_0^{g^{-1}}.$$

Notice that $C_S(\mathcal{H})^g = C_S(\mathcal{E})\alpha \leq S \cap N_G(\mathcal{H}^g) = N_S(\mathcal{H}^g) \leq \hat{S}_0$ and thus $C_S(\mathcal{H}) \leq S_0$. Moreover,

$$C_{S_0}(\mathcal{H})^g = C_{S_0^g}(\mathcal{H}^g) = C_{\hat{S}_0}(\mathcal{H}^g) = C_S(\mathcal{H}^g).$$

Set $\hat{\mathcal{F}}_0 := \mathcal{F}_{\hat{S}_0}(\mathbb{N}_{\mathcal{L}}(\mathcal{H}^g))$. By Proposition 3.39, $C_{\hat{\mathcal{F}}_0}(\mathcal{E}^\alpha) = C_{\mathcal{F}}(\mathcal{E}^\alpha)$. Lemma 3.37 gives now that $(C_{\mathcal{L}}(\mathcal{H}), \delta(C_{\mathcal{F}_0}(\mathcal{E})), C_{S_0}(\mathcal{H}))$ is a regular locality over $C_{\mathcal{F}_0}(\mathcal{E})$, and $(C_{\mathcal{L}}(\mathcal{H}^g), \delta(C_{\mathcal{F}}(\mathcal{E}^\alpha)), C_S(\mathcal{H}^g))$ is a regular locality over $C_{\mathcal{F}}(\mathcal{E}^\alpha)$. Moreover, $\tilde{\alpha} := c_g \in \text{Aut}(\mathcal{L})$ induces an isomorphism from $C_{\mathcal{L}}(\mathcal{H})$ to $C_{\mathcal{L}}(\mathcal{H}^g)$ which takes $C_{S_0}(\mathcal{H})$ to $C_S(\mathcal{H}^g)$. Thus, it follows from Lemma 2.11 that

$$C_{\mathcal{F}_0}(\mathcal{E})^{\tilde{\alpha}} = \mathcal{F}_{C_{S_0}(\mathcal{H})}(C_{\mathcal{L}}(\mathcal{H}))^{\tilde{\alpha}} = \mathcal{F}_{C_{S_0}(\mathcal{H})^g}(C_{\mathcal{L}}(\mathcal{H})^g) = \mathcal{F}_{C_S(\mathcal{H}^g)}(C_{\mathcal{L}}(\mathcal{H}^g)) = C_{\mathcal{F}}(\mathcal{E}^\alpha). \quad (3.15)$$

Let now $P, Q \leq C_S(\mathcal{H}) = C_S(\mathcal{E})$ and $\varphi \in \text{Hom}_{\mathcal{F}}(P, Q)$. As φ is a morphism in \mathcal{F} , Lemma 3.41 yields that φ is a morphism in $C_{\mathcal{F}}(\mathcal{E})$ if and only if it is a morphism in $C_{\mathcal{F}_0}(\mathcal{E})$. By (3.15), this is the case if and only if $\alpha^{-1}\varphi\alpha = \tilde{\alpha}^{-1}\varphi\tilde{\alpha} \in \text{Hom}_{C_{\mathcal{F}}(\mathcal{E}^\alpha)}(P\alpha, Q\alpha)$. This yields the first part of the assertion.

If \mathcal{E} is fully centralized, then $|C_S(\mathcal{E})| \geq |C_S(\mathcal{E}^\alpha)|$ and hence $C_S(\mathcal{E})\alpha = C_S(\mathcal{E}^\alpha)$. Moreover, $C_{\mathcal{F}_0}(\mathcal{E}) = C_{\mathcal{F}}(\mathcal{E})$ by Proposition 3.39. Thus, by (3.15), α induces an isomorphism from $C_{\mathcal{F}}(\mathcal{E})$ to $C_{\mathcal{F}}(\mathcal{E}^\alpha)$. \square

REFERENCES

1. K. K. S. Andersen, B. Oliver, and J. Ventura, *Reduced, tame and exotic fusion systems*, Proc. Lond. Math. Soc. (3) **105** (2012), no. 1, 87–152.
2. M. Aschbacher, *The generalized Fitting subsystem of a fusion system*, Mem. Amer. Math. Soc. **209** (2011), no. 986, vi+110.
3. ———, *On fusion systems of component type*, Mem. Amer. Math. Soc. **257** (2019), no. 1236, v+182.
4. M. Aschbacher, R. Kessar, and B. Oliver, *Fusion systems in algebra and topology*, London Math. Soc. Lecture Note Series, vol. 391, Cambridge University Press, 2011.
5. A. Chermak, *Fusion systems and localities*, Acta Math. **211** (2013), no. 1, 47–139.
6. ———, *Finite Localities III*, arXiv:1505.06161 (2016).
7. ———, *Finite Localities I*, <https://homepages.abdn.ac.uk/d.j.benson/pages/html/archive/chermak.html>, older version on arXiv:1505.07786v3 (first version 2015, corrected version 2020).
8. A. Chermak and E. Henke, *Fusion systems and localities – a dictionary*, arXiv:1706.05343v2 (2017).

9. V. Grazian and E. Henke, *Kernels of localities*, arXiv:2109.10962 (2021).
10. E. Henke, *Products in fusion systems*, J. Algebra **376** (2013), 300–319.
11. ———, *Centralizers of normal subsystems revisited*, J. Algebra **511** (2018), 364–387.
12. ———, *Subcentric linking systems*, Trans. Amer. Math. Soc. **371** (2019), no. 5, 3325–3373.
13. ———, *Commuting partial normal subgroups and regular localities*, arXiv:2103.00955v2 (2021).
14. ———, *Extensions of homomorphisms between localities*, Forum Math. Sigma **9** (2021), Paper No. e63.
15. H. Kurzweil and B. Stellmacher, *The theory of finite groups*, Universitext, Springer-Verlag, New York, 2004.
16. U. Meierfrankenfeld and B. Stellmacher, *Applications of the FF-Module Theorem and related results*, J. Algebra **351** (2012), 64–106.
17. B. Oliver, *Normalizers of sets of components in fusion systems*, preprint on <https://www.math.univ-paris13.fr/~bobol/>.

INSTITUT FÜR ALGEBRA, FAKULTÄT MATHEMATIK, TECHNISCHE UNIVERSITÄT DRESDEN, 01062 DRESDEN, GERMANY

Email address: `ellen.henke@tu-dresden.de`