

SPECTRAL FLOW INSIDE ESSENTIAL SPECTRUM V: ON ABSORBING POINTS OF COUPLING RESONANCES

NURULLA AZAMOV

ABSTRACT. Let H_0 and V be self-adjoint operators, such that V admits a factorisation $V = F^*JF$ with bounded self-adjoint J and $|H_0|^{1/2}$ -compact F . Coupling resonance functions, $r_j(z)$, of the pair H_0 and V can be defined as $r_j(z) = -\sigma_j(z)^{-1}$, where $\sigma_j(z)$ are eigenvalues of the compact-operator valued holomorphic function $F(H_0 - z)^{-1}F^*J$. Taken together, the functions $r_j(z)$ form an infinite-valued holomorphic function on the resolvent set of H_0 . These functions contain a lot of information about the pair H_0, V (this is well-known in the case of rank one V). A point z_0 of the resolvent set we call *absorbing* if some $r_j(z)$ goes to ∞ as $z \rightarrow z_0$ along some half-interval.

In this note I present some partial results concerning absorbing points of coupling resonances.

Let H_0 and V be self-adjoint operators on a (complex separable) Hilbert space \mathcal{H} , such that V admits a factorisation $V = F^*JF$ with bounded self-adjoint J and $|H_0|^{1/2}$ -compact F . *Coupling resonance functions*, $r_j(z)$, of the pair H_0 and V can be defined as

$$r_j(z) = -\sigma_j(z)^{-1},$$

where $\sigma_j(z)$ are eigenvalues of the compact-operator valued holomorphic function $F(H_0 - z)^{-1}F^*J$. Taken together, the functions $r_j(z)$ form an infinite-valued holomorphic function, which we denote $\mathcal{R}(z) = \mathcal{R}(z; H_0, V)$, on the resolvent set of H_0 . These functions contain a lot of information about the pair H_0 and V , — this is well-known in the case of a rank one perturbation $V = \langle v, \cdot \rangle v$, which in essence is equivalent to the theory of Herglotz-Nevanlinna-Pick functions, see e.g. [7, 11], since in this case the function

$$\mathbb{C}_+ \ni z \mapsto \langle v, (H_0 - z)^{-1}v \rangle$$

is Herglotz-Nevanlinna-Pick and it allows to recover the pair H_0 and V up to a unitary equivalence, — assuming the pair is irreducible, using the same unitary operator.

More information about coupling resonance functions, their applications and why they are so-called can be found in the introductions of papers [1, 3, 4].

The general case is complicated by the fact that the function $\mathcal{R}(z)$ is infinite-valued, and can have potentially quite an erratic behaviour, typical for such functions. However, as it turns out, there are only two types of singularities given in the following theorem, see [2, Theorem 3.1]:

Theorem 1. *Coupling resonance points can have only two types of singularities:*

- (1) *continuous branching points of finite period, or*
- (2) *absorbing points, whether isolated or not.*

Moreover, isolated absorbing points, if they exist, must have infinite period of branching.

Recall, [2], that a point z_0 of the resolvent set is called *absorbing* if some coupling resonance function goes to ∞ as $z \rightarrow z_0$ along some half-interval, γ , ending at z_0 . In this case it is not difficult to see that the point z_0 has a small enough neighbourhood such that if another half-interval γ' ending at z_0 is homotopic to γ in the domain of holomorphy of $\mathcal{R}(z)$ then the coupling resonance function goes to infinity along γ' too.

Clearly, Theorem 1 significantly reduces the possible types of singularities which coupling resonance functions can have. I believe that more is true, as specified in the following

2000 *Mathematics Subject Classification.* Primary 47A40;
Key words and phrases. coupling resonances.

Conjecture 1. *Assume that H_0 and V is as above. Then the pair H_0 and V does not have absorbing points, whether isolated or not.*

It would be nice to have this proved, since then the analytic function $\mathcal{R}(z)$ could have only one type of singularity: continuous branching points of finite period, which are quite manageable. Concerning these branching points, it is not difficult to demonstrate their existence. What is their meaning is currently being investigated, see [4].

In this paper I present a solution to a very special case of Conjecture 1.

First I present some preliminaries. Let $s_1(T), s_2(T), \dots$ be s -numbers of a compact operator T , see e.g. [9, 11] for details, and let $\lambda_1(T), \lambda_2(T), \dots$ be eigenvalues of T written in the order of decreasing magnitudes. The following holds for any $n = 1, 2, \dots$ and $p \in (0, \infty)$, see [9]:

$$(1) \quad \sum_{j=1}^n |\lambda_j(T)|^p \leq \sum_{j=1}^n [s_j(T)]^p.$$

By $\|T\|_1$ we denote the trace class norm of T . We denote $R_z(H_0) = (H_0 - z)^{-1}$.

Theorem 2. *Let H_0 and V be self-adjoint operators. If V admits a factorisation $V = F^* J F$ with bounded self-adjoint J and closed $|H_0|^{1/2}$ -compact F such that $FR_z(H_0)F^*J$ is trace-class then all coupling resonance functions of the pair (H_0, V) do not have isolated absorbing points.*

Proof. Assume that z_0 is an isolated absorbing point of the necessarily infinite-valued in a neighbourhood of z_0 (see Theorem 1) resonance function with values $r_j(z)$, $j \in \mathbb{Z}$, in a neighbourhood of z_0 . Without loss of generality, we can assume that r_j take values in the upper half-plane. For a real s , the numbers $(s - r_j(z))^{-1}$ are eigenvalues of the trace class operator $FR_z(H_s)F^*J$ and thus the series

$$f(z) := \sum_{j \in \mathbb{Z}} (s - r_j(z))^{-1}$$

converges absolutely and uniformly on compact subsets of a deleted neighbourhood of z_0 . The function $f(z)$ is single-valued in the deleted neighbourhood and takes values in \mathbb{C}_+ since all summands of the series take values in \mathbb{C}_+ . Also, since z_0 is an absorbing point, the function f admits analytic continuation to z_0 where it takes zero value. Indeed, by (1) for any $z \in O$ the series

$$f(z) := \sum_{j \in \mathbb{Z}} |s - r_j(z)|^{-1}$$

is bounded by $\sup_{z \in O} \|FR_z(H_s)F^*J\|_1$, where O is a neighbourhood of z_0 . But this contradicts the openness principle, since $f|_O$ does not take values in \mathbb{C}_- . \square

The following theorem of T. Radó gives a positive answer to a very weak version of the famous Painlevé problem about removable sets, see e.g. [8]. Its proof can be found in [10, §§49].

Theorem 3. *Let G be an open set in \mathbb{C} . If a continuous map $u: G \rightarrow \mathbb{C}$ is holomorphic outside of its set of zeros then it is holomorphic in G .*

In my opinion, it seems to be highly unlikely that an absorbing point, if it exists, can be non-isolated. In any case, the set of absorbing points is small in the following sense: the set of rays emanating from a point $z_0 \in \mathbb{C}_+$ which hit an absorbing point has Lebesgue measure zero. This easily follows from the above mentioned properties of absorbing points, the Riemann Mapping Theorem and the Nevanlinna-Luzin-Privalov theorem. I present a proof of absence of non-isolated absorbing points in a very special case.

Theorem 4. *Let $FR_z(H_0)F^*J$ be trace class, O an open set in \mathbb{C}_+ , K a compact subset of O , and $r(z)$ a possibly multi-valued coupling resonance function which is either regular in $O \setminus K$ or may have continuous branching points. If $r(z)$ goes to ∞ , whenever z goes to a point of K along any half-interval in $O \setminus K$, then the set K is empty.*

Proof. We define a function

$$f(z) := \sum_j (s - r_j(z))^{-1},$$

where the sum is taken over all branches of the multi-valued resonance function in $O \setminus K$. By the argument of the proof of Theorem 2, the function $f(z)$ is continuous in O , is holomorphic outside K and vanishes in K . Hence, by Rado's Theorem 3, $f(z)$ is holomorphic in O . So, the set of zeros of this function is discrete, and so is K . Hence, by Theorem 2 the set K is empty. \square

It is possible that Conjecture 1 does not hold in full generality. If so, it would be interesting to find sufficient conditions on the pair H_0 and V under which the conjecture holds. In particular, the case of $FR_z(H_0)F^* \in \mathcal{L}_{1,\infty}$, where $\mathcal{L}_{1,\infty}$ is the Dixmier ideal, see e.g. [6], seems to be especially interesting, see [5].

Acknowledgements. I thank my wife for financially supporting me during the work on this paper.

REFERENCES

- [1] N. A. Azamov, *Spectral flow inside essential spectrum*, Dissertationes Math. **518**, 1-156 (2016)
- [2] N. A. Azamov, *Spectral flow inside essential spectrum III: coupling resonances near essential spectrum*, arXiv: 2109.04675
- [3] N. A. Azamov and T. W. Daniels, *Singular spectral shift function for resolvent comparable operators*, Math. Nachr. **292**, 1911-1930 (2019)
- [4] N. A. Azamov and T. W. Daniels, *Coupling resonances and spectral properties of the product of resolvent and perturbation*, in preparation
- [5] N. Azamov, E. McDonald, F. Sukochev, D. Zanin, *A Dixmier trace formula for the density of states*, Commun. Math. Phys. **377**, 2597–2628 (2020)
- [6] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, 1994
- [7] W. F. Donoghue, *Monotone Matrix Functions and Analytic Continuation*, Springer, Berlin, Heidelberg, New York, 1974
- [8] J. J. Dudziak, *Vitushkin's conjecture for removable sets*, Universitext, Springer Science+Business Media, LLC 2010
- [9] I. C. Gohberg, M. G. Krein, *Introduction to the Theory of Linear Non-selfadjoint Operators*, Providence, R. I., AMS, Trans. Math. Monographs, vol. 18, 1969.
- [10] B. V. Shabat, *Introduction to complex analysis, part II*, Providence, R. I., AMS, 1992
- [11] B. Simon, *Trace Ideals and their Applications*, Second edition, Math. Surveys Monogr. (Amer. Math. Soc., 2005)

INDEPENDENT SCHOLAR, ADELAIDE, SA, AUSTRALIA
 Email address: azamovnurulla@gmail.com